



(11) **EP 2 949 724 A1**

(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 153(4) EPC

(43) Date of publication:  
**02.12.2015 Bulletin 2015/49**

(51) Int Cl.:  
**C09K 11/06<sup>(2006.01)</sup> C07D 403/14<sup>(2006.01)</sup>**  
**H01L 51/50<sup>(2006.01)</sup>**

(21) Application number: **14742808.0**

(86) International application number:  
**PCT/JP2014/051184**

(22) Date of filing: **22.01.2014**

(87) International publication number:  
**WO 2014/115743 (31.07.2014 Gazette 2014/31)**

(84) Designated Contracting States:  
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB**  
**GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO**  
**PL PT RO RS SE SI SK SM TR**  
Designated Extension States:  
**BA ME**

(30) Priority: **23.01.2013 JP 2013010153**  
**23.01.2013 JP 2013010154**

(71) Applicant: **Kyushu University,**  
**National University Corporation**  
**Fukuoka-shi, Fukuoka 812-8581 (JP)**

(72) Inventors:  
• **HIRATA Shuzo**  
**Fukuoka-shi**  
**Fukuoka 812-8581 (JP)**

- **SAKAI Yumi**  
**Kurume-shi**  
**Fukuoka 830-8511 (JP)**
- **MASUI Kensuke**  
**Ashigarakami-gun**  
**Kanagawa 258-8577 (JP)**
- **SHIZU Katsuyuki**  
**Fukuoka-shi**  
**Fukuoka 812-8581 (JP)**
- **TANAKA Hiroyuki**  
**Fukuoka-shi**  
**Fukuoka 812-8581 (JP)**
- **ADACHI Chihaya**  
**Fukuoka-shi**  
**Fukuoka 812-8581 (JP)**

(74) Representative: **Zimmermann & Partner**  
**Patentanwälte mbB**  
**Postfach 330 920**  
**80069 München (DE)**

(54) **LIGHT EMITTING MATERIAL AND ORGANIC LIGHT EMITTING ELEMENT USING SAME**

(57) An organic light emitting device containing a compound represented by the following general formula in a light emitting layer have a high light emission efficiency. In the general formula (1), R<sup>1</sup> and R<sup>2</sup> represent carbazolyl, aryl, heteroaryl, alkyl or cycloalkyl; R<sup>5</sup> and R<sup>6</sup> represent alkyl; R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> represent aryl, alkyl or carbazolyl; R<sup>10</sup> represents carbazolyl which may be substituted with carbazolyl, aryl, heteroaryl, alkyl or cycloalkyl; n<sub>1</sub>, n<sub>2</sub>, n<sub>6</sub> and n<sub>7</sub> represent an integer of from 0 to 4; n<sub>5</sub> represents an integer of from 0 to 3; n<sub>8</sub> and n<sub>9</sub> represent an integer of from 0 to 5; and n<sub>10</sub> represents 0 or 1.

**EP 2 949 724 A1**





Fig. 1



## Description

## Technical Field


5 **[0001]** The present invention relates to a light emitting material having a high light emission efficiency, and an organic light emitting device, such as an organic electroluminescent device (organic EL device), using the same.

## Background Art

10 **[0002]** An organic light emitting device, such as an organic electroluminescent device, has been actively studied for enhancing the light emission efficiency thereof. In particular, various studies for enhancing the light emitting efficiency have been made by newly developing and combining an electron transporting material, a hole transporting material, a light emitting material and the like constituting an organic electroluminescent device. There are studies relating to an organic electroluminescent device utilizing a compound containing a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure, which are found among them, and some proposals have been made hitherto.


15 **[0003]** For example, Patent Document 1 describes the use of the compound represented by the following general formula as an electron transporting material of an electron transporting layer of an organic electroluminescent device. In the following general formula, n represents 1 or 2, Ar represents an arylene group or a heteroarylene group, R<sub>3</sub> and R<sub>4</sub> each represent a hydrogen atom or an aryl group, X<sub>1</sub> to X<sub>3</sub> each represent =CR- or =N-, R represents a hydrogen atom or a substituent, and Cz represents a carbazolyl group.

[chem 1]



35 **[0004]** Patent Document 1 describes the compound A having the following structure as the compound represented by the general formula, and also describes the example of an organic electroluminescent device using the compound in an electron transporting layer. However, a compound obtained by introducing a substituent to the carbazolyl group of the compound A is not investigated. Furthermore, Patent Document 1 does not consider the usefulness of the compound A as a light emitting material.

[chem 2]




55 **[0005]** Patent Document 2 describes that the compound having the structure represented by the following general formula emits blue fluorescent light, and the compound is useful as a light emitting device material. In the following general formula, R<sup>11</sup> and R<sup>12</sup> each represent a hydrogen atom, an aliphatic hydrocarbon group, an aryl group or a heterocyclic group, R<sup>1</sup> and R<sup>2</sup> each represent a hydrogen atom or a substituent that does not contain an amino group, and L represents a linking group. Patent Document 2 describes that an organic light emitting device using the compound A as a light emitting material emits blue fluorescent light. However, Patent Document 2 also does not investigate a

compound obtained by introducing a substituent to the carbazolyl group of the compound A.

[chem 3]

5

10



15

**[0006]** Patent Document 3 describes the compound having a partial structure containing three carbazole structures connected to each other, and describes specific examples using the compound having the partial structure as a host material of a light emitting layer of an organic light emitting device. As examples of the compound having a partial structure containing three carbazole structures connected to each other, the compound having the following structure is exemplified among the many example structures. However, there is no example that uses the compound, and the usefulness of the compound as a light emitting material is not mentioned.


20

[chem 4]

25

30

35



40

Citation List

Patent Documents

45

**[0007]**

Patent Document 1: JP-A-2009-21336

Patent Document 2: JP-A-2002-193952

Patent Document 3: WO 2012/077902

50

Summary of Invention

Technical Problem

55

**[0008]** As described above, there have been some studies on a compound containing a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure, and proposals relating to application thereof to an organic electroluminescent device have been slightly made. However, like Patent Documents 1 to 3 described above, the literatures referring to the compound containing a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure discuss only the general use-

fulness of the group of compounds represented by the extensive general formulae encompassing the numerous other compounds, but do not focus the compound containing a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure and do not investigate the details of the compound. Accordingly, the relationship between the chemical structure of the group of compounds containing a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure and the usefulness of the compounds as a light emitting material has not been sufficiently clarified, and it is the current situation that it is difficult to estimate the usefulness as a light emitting material based on the chemical structure. In particular, Patent Document 1 and 2 do not evaluate the usefulness of the compounds as a light emitting material, and it is difficult to obtain any suggestion relating to the usefulness as a light emitting material based on the literatures. Furthermore, Patent Document 3 specifically describes the usefulness of the compound A as a light emitting material, but the compound A does not emit delayed fluorescent light and is not sufficiently satisfactory in the light emission efficiency.

**[0009]** The present inventors have considered these problems and have made investigations for evaluating in detail the usefulness of the compounds containing a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure as a light emitting material of an organic light emitting device. The inventors also have made investigations for providing a general formula of compounds that are particularly useful as a light emitting material and generalizing the structure of an organic light emitting device having a high light emission efficiency.

#### Solution to Problem


**[0010]** As a result of earnest investigations for achieving the objects, the inventors have clarified that compounds that contain a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure and satisfy the particular structural condition are particularly useful as a light emitting material. In particular, the inventors have found compounds that are useful as a delayed fluorescent material in the compounds that contain a carbazole structure and a 2,4,6-triphenyl-1,3,5-triazine structure, and have clarified that an organic light emitting device having a high light emission efficiency may be provided inexpensively. Based on the knowledge, the inventors have provided the following inventions as measures for solving the problems.

#### **[0011]**

(1) A light emitting material containing a compound represented by the following general formula (1):

[chem 5]

General Formula (1)



wherein in the general formula (1),

R<sup>1</sup> and R<sup>2</sup> each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl

group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$R^{10}$  represents a carbazolyl group, provided that the carbazolyl group may be substituted with a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$n_1$ ,  $n_2$ ,  $n_6$  and  $n_7$  each independently represent an integer of from 0 to 4;

$n_5$  represents an integer of from 0 to 3;

$n_8$  and  $n_9$  each independently represent an integer of from 0 to 5; and


$n_{10}$  represents 0 or 1,

provided that when  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$ ,  $R^2$  and  $R^5$  to  $R^9$  corresponding to  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  respectively each may be the same as or different from each other.

(2) The light emitting material according to the item (1), wherein the compound represented by the general formula (1) is a compound represented by the following general formula (2) :

[chem 6]

General Formula (2)



wherein in the general formula (2),

$R^1$  and  $R^2$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$n_1$ ,  $n_2$ ,  $n_6$  and  $n_7$  each independently represent an integer of from 0 to 4;

$n_5$  represents an integer of from 0 to 3; and


$n_8$  and  $n_9$  each independently represent an integer of from 0 to 5,

provided that when  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$ ,  $R^2$  and  $R^5$  to  $R^9$  corresponding to  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  respectively each may be the same as or different from each other.

(3) The light emitting material according to the item (1), wherein the compound represented by the general formula (1) is a compound represented by the following general formula (3) :

[chem 7]

General Formula (3)



wherein in the general formula (3),

$R^1$  to  $R^4$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$n1$  to  $n4$  and  $n7$  each independently represent an integer of from 0 to 4;

$n5$  and  $n6$  each represent an integer of from 0 to 3; and

$n8$  and  $n9$  each independently represent an integer of from 0 to 5,

provided that when  $n1$  to  $n9$  each represent an integer of 2 or more, plural groups represented by  $R^1$  to  $R^9$  corresponding to  $n1$  to  $n9$  respectively each may be the same as or different from each other.

(4) The light emitting material according to any one of the items (1) to (3), wherein the light emitting material emits delayed fluorescent light.

(5) The light emitting material according to the item (2) or (4), wherein in the general formula (2),  $n6$  represents 0.

(6) The light emitting material according to any one of the items (2), (4) and (5), wherein in the general formula (2),  $n1$  represents an integer of from 1 to 4.

(7) The light emitting material according to the item (6), wherein in the general formula (2),  $R^1$  is bonded to the 3-position of the carbazolyl group.

(8) The light emitting material according to the item (6) or (7), wherein in the general formula (2),  $n2$  represents an integer of from 1 to 4.

(9) The light emitting material according to the item (8), wherein in the general formula (2),  $R^2$  is bonded to the 6-position of the carbazolyl group.

(10) The light emitting material according to any one of the items (1), (2) and (4) to (9), wherein in the general formulae (1) and (2),  $R^1$  and  $R^2$  each independently represent a substituted or unsubstituted 9-carbazolyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted pyridyl group, an alkyl group having from 1 to 6 carbon atoms or a cycloalkyl group having from 5 to 7 carbon atoms.

(11) The light emitting material according to any one of the items (1), (2) and (4) to (9), wherein in the general formulae (1) and (2), R<sup>1</sup> and R<sup>2</sup> each independently represent a 9-carbazolyl group, a phenyl group, a tolyl group, a dimethylphenyl group, a trimethylphenyl group, a biphenyl group, a pyridyl group, a pyrrolyl group, a tert-butyl group or a cyclohexyl group.

(12) The light emitting material according to any one of the items (2), (4) and (5), wherein in the general formula (2), both n<sub>1</sub> and n<sub>2</sub> represent 0.

(13) The light emitting material according to the item (3) or (4), wherein in the general formula (3), at least one of n<sub>1</sub> to n<sub>4</sub> represents an integer of from 1 to 4.

(14) The light emitting material according to the item (3) or (4), wherein in the general formula (3), n<sub>1</sub> to n<sub>4</sub> each independently represent an integer of from 1 to 4.

(15) The light emitting material according to any one of the items (3), (4), (13) and (14), wherein in the general formula (3), R<sup>1</sup> to R<sup>4</sup> each independently represent a substituted or unsubstituted 9-carbazolyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted pyridyl group, an alkyl group having from 1 to 6 carbon atoms or a cycloalkyl group having from 5 to 7 carbon atoms.

(16) The light emitting material according to any one of the items (3), (4), (13) and (14), wherein in the general formula (3), R<sup>1</sup> to R<sup>4</sup> each independently represent a 9-carbazolyl group, a phenyl group, a tolyl group, a dimethylphenyl group, a trimethylphenyl group, a biphenyl group, a pyridyl group, a pyrrolyl group, a tert-butyl group or a cyclohexyl group.

(17) The light emitting material according to any one of the items (3), (4) and (13) to (16), wherein in the general formula (3), both n<sub>5</sub> and n<sub>6</sub> represent 0.

(18) The light emitting material according to any one of the items (1) to (17), wherein in the general formulae (1) to (3), all n<sub>7</sub>, n<sub>8</sub> and n<sub>9</sub> represent 0.


(19) A delayed fluorescence emitter having a structure represented by the general formula (1).

(20) A compound represented by the general formula (2).

(21) A compound represented by the following general formula (4):

[chem 8]

General Formula (4)



wherein in the general formula (4),

R<sup>11</sup> to R<sup>14</sup> each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

R<sup>15</sup> and R<sup>16</sup> each independently represent an alkyl group;

R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;  
 n<sup>11</sup> to n<sup>14</sup> and n<sup>17</sup> each independently represent an integer of from 0 to 4;  
 n<sup>15</sup> and n<sup>16</sup> each represent an integer of from 0 to 3; and  
 n<sup>18</sup> and n<sup>19</sup> each independently represent an integer of from 0 to 5,  
 provided that  
 at least one of n<sup>11</sup> to n<sup>14</sup> represents an integer of from 1 to 4, and  
 when n<sup>11</sup> to n<sup>19</sup> each represent an integer of 2 or more, plural groups represented by R<sup>11</sup> to R<sup>19</sup> corresponding to n<sup>11</sup> to n<sup>19</sup> respectively each may be the same as or different from each other.

(22) An organic light emitting device containing a substrate having thereon a light emitting layer containing the light emitting material according to any one of the items (1) to (18).

(23) The organic light emitting device according to the item (22), wherein the organic light emitting device emits delayed fluorescent light.

(24) The organic light emitting device according to the item (22) or (23), wherein the organic light emitting device is an organic electroluminescent device.

#### Advantageous Effects of Invention

**[0012]** The organic light emitting device of the invention has such a feature that the device has a high light emission efficiency. The compound and the light emitting material of the invention may be effectively used for producing the organic light emitting device. In particular, the delayed fluorescence emitter of the invention has such a feature that when the material is used in a light emitting layer of an organic light emitting device, the organic light emitting device emits delayed fluorescent light with a light emission efficiency that is drastically enhanced.

#### Brief Description of Drawings

##### **[0013]**

[Fig. 1] Fig. 1 is a schematic cross sectional view showing an example of a layer structure of an organic electroluminescent device.

[Fig. 2] Fig. 2 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 1.

[Fig. 3] Fig. 3 is the transient decay curves of the organic electroluminescent device using the compound 1.

[Fig. 4] Fig. 4 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 1.

[Fig. 5] Fig. 5 is a graph showing the electric current density-external quantum efficiency characteristics of the organic electroluminescent device using the compound 1.

[Fig. 6] Fig. 6 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 2.

[Fig. 7] Fig. 7 is the transient decay curves of the organic electroluminescent device using the compound 2.

[Fig. 8] Fig. 8 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 2.

[Fig. 9] Fig. 9 is a graph showing the electric current density-external quantum efficiency characteristics of the organic electroluminescent device using the compound 2.

[Fig. 10] Fig. 10 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 3.

[Fig. 11] Fig. 11 is the transient decay curves of the organic electroluminescent device using the compound 3.

[Fig. 12] Fig. 12 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 3.

[Fig. 13] Fig. 13 is a graph showing the electric current density-external quantum efficiency characteristics of the organic electroluminescent device using the compound 3.

[Fig. 14] Fig. 14 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 4.

[Fig. 15] Fig. 15 is the transient decay curves of the organic electroluminescent device using the compound 4.

[Fig. 16] Fig. 16 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 4.

[Fig. 17] Fig. 17 is a graph showing the electric current density-external quantum efficiency characteristics of the

organic electroluminescent device using the compound 4.

[Fig. 18] Fig. 18 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 27.

[Fig. 19] Fig. 19 is the transient decay curves of the organic electroluminescent device using the compound 27.

[Fig. 20] Fig. 20 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 27.

[Fig. 21] Fig. 21 is a graph showing the electric current density-external quantum efficiency characteristics of the organic electroluminescent device using the compound 27.

[Fig. 22] Fig. 22 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 28.

[Fig. 23] Fig. 23 is the transient decay curves of the organic electroluminescent device using the compound 28.

[Fig. 24] Fig. 24 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 28.

[Fig. 25] Fig. 25 is a graph showing the electric current density-external quantum efficiency characteristics of the organic electroluminescent device using the compound 28.

[Fig. 26] Fig. 26 is the light emission spectra of the organic photoluminescent device and the organic electroluminescent device using the compound 29.

[Fig. 27] Fig. 27 is the transient decay curves of the organic electroluminescent device using the compound 29.

[Fig. 28] Fig. 28 is a graph showing the voltage-electric current density characteristics of the organic electroluminescent device using the compound 29.

[Fig. 29] Fig. 29 is a graph showing the electric current density-external quantum efficiency characteristics of the organic electroluminescent device using the compound 29.

[Fig. 30] Fig. 30 is the light emission spectrum of the toluene solution of the compound A.

[Fig. 31] Fig. 31 is the transient decay curves of the toluene solution of the compound A.

#### Description of Embodiments


**[0014]** The contents of the invention will be described in detail below. The constitutional elements may be described below with reference to representative embodiments and specific examples of the invention, but the invention is not limited to the embodiments and the examples. In the description, a numerical range expressed with reference to an upper limit and/or a lower limit means a range that includes the upper limit and/or the lower limit. In the invention, the hydrogen atom that is present in the molecule in the compound used in the invention is not particularly limited in isotope species, and for example, all the hydrogen atoms in the molecule may be  $^1\text{H}$ , and all or a part of them may be  $^2\text{H}$  (deuterium (D)).

Compound represented by General Formula (1)

**[0015]** The light emitting material of the invention contains the compound represented by the following general formula (1). The organic light emitting device of the invention contains the compound represented by the following general formula (1) as a light emitting material of a light emitting layer. The compound represented by the following general formula (1) will be described.

[chem 9]

## General Formula (1)



25 **[0016]** In the general formula (1),  $R^1$  and  $R^2$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group.

**[0017]** The bonding position of the carbazolyl group herein is not limited, and is preferably a 9-carbazolyl group or a 3-carbazolyl group, and more preferably a 9-carbazolyl group.

30 **[0018]** The aryl group may be a monocyclic ring or a fused ring, and preferably has from 6 to 14 carbon atoms, and more preferably from 6 to 10 carbon atoms. Preferred examples thereof include a phenyl group.

**[0019]** The heteroaryl group may be a monocyclic ring or a fused ring, and preferably has from 2 to 12 carbon atoms, more preferably from 3 to 10 carbon atoms, and further preferably from 3 to 6 carbon atoms. Specific examples thereof include a pyridyl group and a pyrrolyl group, and preferred examples thereof include a 1-pyridyl group, a 2-pyridyl group, a 3-pyridyl group, a 1-pyrrolyl group, a 2-pyrrolyl group and a 3-pyrrolyl group.

35 **[0020]** The alkyl group may be linear or branched, and preferably has from 1 to 12 carbon atoms, more preferably from 1 to 6 carbon atoms, and further preferably from 1 to 4 carbon atoms. Specific examples thereof include a methyl group, an ethyl group, a n-propyl group, an isopropyl group, a n-butyl group, an isobutyl group and a tert-butyl group, and preferred examples thereof include a tert-butyl group.

40 **[0021]** The cycloalkyl group may be a monocyclic ring or a fused ring, and preferably has from 5 to 12 carbon atoms, and more preferably from 5 to 7 carbon atoms. Specific examples thereof include a cyclopentyl group, a cyclohexyl group and a cycloheptyl group, and preferred examples thereof include a cyclohexyl group.

45 **[0022]** The carbazolyl group, the aryl group, the heteroaryl group, the alkyl group and the cycloalkyl group capable of being represented by  $R^1$  and  $R^2$  each may have a substituent. In the case where the group has a substituent, the substitution position and the number of the substituent are not particularly limited. The number of the substituent on the group is preferably from 0 to 6, and more preferably from 0 to 4, and for example, may be preferably from 0 to 2. In the case where the group has plural substituents, the substituents may be the same as or different from each other, and preferably the same as each other. Examples of the substituent include a hydroxyl group, a halogen atom, a cyano group, an alkyl group having from 1 to 12 carbon atoms, an alkoxy group having from 1 to 12 carbon atoms, an alkylthio group having from 1 to 12 carbon atoms, an alkyl-substituted amino group having from 1 to 12 carbon atoms, an acyl group having from 2 to 12 carbon atoms, an aryl group having from 6 to 14 carbon atoms, a heteroaryl group having from 3 to 13 carbon atoms, a diarylamino group having from 12 to 20 carbon atoms, a substituted or unsubstituted carbazolyl group having from 12 to 20 carbon atoms, an alkenyl group having from 2 to 10 carbon atoms, an alkynyl group having from 2 to 10 carbon atoms, an alkoxy carbonyl group having from 2 to 10 carbon atoms, an alkylsulfonyl group having from 1 to 10 carbon atoms, a haloalkyl group having from 1 to 10 carbon atoms, an amide group, an alkylamide group having from 2 to 10 carbon atoms, a trialkylsilyl group having from 3 to 12 carbon atoms, a trialkylsilylalkyl group having from 4 to 12 carbon atoms, a trialkylsilylalkenyl group having from 5 to 14 carbon atoms, a trialkylsilylalkynyl group having from 5 to 14 carbon atoms, and a nitro group. In these specific examples, the substituent that is capable of being further substituted with a substituent may be substituted. The carbazolyl group, the aryl group, the heteroaryl

50

55

group, the alkyl group and the cycloalkyl group capable of being represented by  $R^1$  and  $R^2$  in the general formula (1) that are unsubstituted are also preferred. An alkyl-substituted aryl group is also preferred, and examples thereof include a tolyl group, a dialkylphenyl group and a trialkylphenyl group, specific examples of which include a 1-tolyl group, a 2-tolyl group, a 3-tolyl group, a 2,6-dimethylphenyl group, a 2,4-dimethylphenyl group and a 2,4,6-trimethylphenyl group.

5 **[0023]** In the general formula (1),  $R^1$  and  $R^2$  each preferably independently represent a substituted or unsubstituted 9-carbazolyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted pyridyl group, an alkyl group having from 1 to 6 carbon atoms or a cycloalkyl group having from 5 to 7 carbon atoms.  $R^1$  and  $R^2$  each more preferably independently represent a 9-carbazolyl group, a phenyl group, a tolyl group, a dimethylphenyl group, a trimethylphenyl group, a biphenyl group, a pyridyl group, a pyrrolyl group, a tert-butyl group or a cyclohexyl group.

10 **[0024]** In the general formula (1),  $n_1$  and  $n_2$  each independently represent an integer of from 0 to 4, preferably an integer of from 0 to 3, and more preferably an integer of from 0 to 2. In the case where  $n_1$  is 2 or more, plural groups represented by  $R^1$  may be the same or different, and in the case where  $n_2$  is 2 or more, plural groups represented by  $R^2$  may be the same or different.  $n_1$  and  $n_2$  may be the same as or different from each other. Examples of the case where  $n_1$  and  $n_2$  are the same as each other include the case where both of them are 0, the case where both of them are 1, and the case where both of them are 2. Examples of the case where  $n_1$  and  $n_2$  are different from each other include the case where  $n_1$  is 1 and  $n_2$  is 0.

15 **[0025]** In the general formula (1),  $R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group. For the description and the preferred range of the alkyl group, reference may be made to the description for the alkyl group capable of being represented by  $R^1$  and  $R^2$ . For example, an unsubstituted alkyl group may be used as  $R^5$  and  $R^6$ , and a methyl group may be preferably used as  $R^5$  and  $R^6$ .

20 **[0026]**  $n_5$  represents an integer of from 0 to 3, and  $n_4$  represents an integer of from 0 to 4.  $n_5$  and  $n_5$  each preferably independently represent an integer of from 0 to 2, and more preferably 0 or 1, and the case where both of them are 0 is also preferred. In particular, the case where  $n_6$  is 0 is preferably used. In the case where  $n_5$  is 2 or more, plural groups represented by  $R^5$  may be the same or different, and in the case where  $n_6$  is 2 or more, plural groups represented by  $R^6$  may be the same or different.  $n_5$  and  $n_6$  may be the same as or different from each other. Preferred examples of the case where  $n_5$  and  $n_6$  are the same as each other include the case where both of them are 0 and the case where both of them are 1.

25 **[0027]** In the general formula (1),  $R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group. For the descriptions and the preferred ranges of the aryl group, the alkyl group, the carbazolyl group and the substituent, reference may be made to the corresponding descriptions for  $R^1$  and  $R^2$ .

30 **[0028]**  $n_7$  represents an integer of from 0 to 4, and  $n_8$  and  $n_9$  each independently represent an integer of from 0 to 5.  $n_7$  preferably represents an integer of from 0 to 2, and more preferably 0 or 1, and  $n_7$  also preferably represents 0.  $n_8$  and  $n_9$  each preferably represent an integer of from 0 to 2, and more preferably 0 or 1. In the case where  $n_7$  is 2 or more, plural groups represented by  $R^7$  may be the same or different, in the case where  $n_8$  is 2 or more, plural groups represented by  $R^8$  may be the same or different, and in the case where  $n_9$  is 2 or more, plural groups represented by  $R^9$  may be the same or different.  $n_8$  and  $n_9$  may be the same as or different from each other. Preferred examples of the case where  $n_8$  and  $n_9$  are the same as each other include the case where both of them are 0 and the case where both of them are 1. Examples of the case where  $n_8$  and  $n_9$  are different from each other include the case where  $n_8$  is 1 and  $n_9$  is 0.

35 **[0029]** Preferred examples of  $R^8$  and  $R^9$  in the general formula (1) include a phenyl group, a tolyl group, a dimethylphenyl group and a trimethylphenyl group, and more specific examples thereof include a phenyl group, a 1-tolyl group, a 2-tolyl group, a 3-tolyl group, a 2,6-dimethylphenyl group, a 2,4-dimethylphenyl group and a 2,4,6-trimethylphenyl group. The case where  $R^8$  and  $R^9$  in the general formula (1) each represent a substituted or unsubstituted carbazolyl group is also preferred. In this case, the substituted or unsubstituted carbazolyl group is preferably bonded to the 4-position of the phenyl group bonded to the triazine ring. Examples of the substituted carbazolyl group herein include a carbazolyl group substituted with a carbazolyl group. Preferred examples thereof include a structure represented by the following general formula (5).

[chem 10]


General Formula (5)

5

10

15

20



25

**[0030]** In the general formula (5), D1 to D3 each independently have a structure represented by the following general formula (6), provided that D3 may represent a hydrogen atom, a substituted or unsubstituted aryl group or a substituted or unsubstituted alkyl group;  $R^{7'}$ ,  $R^{8'}$  and  $R^{9'}$  each independently represent a substituted or unsubstituted aryl group or a substituted or unsubstituted alkyl group; and  $n7'$ ,  $n8'$  and  $n9'$  each independently represent an integer of from 0.4.

[chem 11]


General Formula (6)

30

35

40

45



50

**[0031]** In the general formula (6), for the definitions and the preferred ranges of  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^6$ ,  $n1$ ,  $n2$ ,  $n5$  and  $n6$ , reference may be made to the corresponding descriptions therefor in the general formula (1).

55

**[0032]** In the general formula (5), D1 and D2 may be the same as or different from each other. In the case where D6 has the structure represented by the general formula (6), all D1 to D3 may be the same as each other, only two of them may be the same as each other, or all of them may be different from each other. The compound having D1 to D3 that are the same as each other has such an advantage that the compound may be easily synthesized.


**[0033]** In the general formulae (1) and (6),  $R^{10}$  represents a carbazolyl group, provided that the carbazolyl group may be substituted with a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl

group. For the descriptions and the preferred ranges of the carbazolyl group, the aryl group, the heteroaryl group, the alkyl group, the cycloalkyl group and the substituent, reference may be made to the corresponding descriptions for R<sup>1</sup> and R<sup>2</sup>.

[0034] n<sub>10</sub> represents 0 or 1. The compound represented by the general formula (1), in which n<sub>10</sub> represents 0, may be represented by the following general formula (2), and the compound represented by the general formula (1), in which n<sub>10</sub> represents 1, may be represented by the following general formula (3).

[chem 12]

General Formula (2)



[0035] In the general formula (2), for the definitions and the preferred ranges of R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup> to R<sup>9</sup>, n<sub>1</sub>, n<sub>2</sub>, and n<sub>5</sub> to n<sub>9</sub>, reference may be made to the corresponding descriptions therefor in the general formula (1).


[0036] In the general formula (2), in the case where at least one of R<sup>1</sup> and R<sup>2</sup> is present, the bonding position thereof may be any of the 1- to 8-positions of the carbazole ring, and is preferably any of the 2- to 7-positions, more preferably any of the 3-, 4-, 6- and 7-positions, and further preferably any of the 3- and 6-positions. Preferred examples of the case include the case where n<sub>1</sub> represents an integer of from 1 to 4, and R<sup>1</sup> is bonded to the 3-position of the carbazole ring, the case where n<sub>2</sub> represents an integer of from 1 to 4, and R<sup>2</sup> is bonded to the 6-position of the carbazole ring, and the case where both n<sub>1</sub> and n<sub>2</sub> each represent an integer of from 1 to 4, and R<sup>1</sup> is bonded to the 3-position of the carbazole ring, whereas R<sup>2</sup> is bonded to the 6-position thereof. In the general formula (2), in the case where both R<sup>1</sup> and R<sup>2</sup> are present, R<sup>1</sup> and R<sup>2</sup> may be the same as or different from each other.

[0037] In the general formula (2), in the case where R<sup>5</sup> is present, the substitution position thereof may be any of the positions among the 1- to 4-positions of the carbazole ring that do not have the carbazolyl group substituted thereon. The carbazolyl group is preferably substituted on the 3-position of the carbazole ring. In the case where R<sup>6</sup> is present, the substitution position thereof is preferably any of the 5-, 6- and 7-positions among the 5- to 8-positions of the carbazole ring, more preferably any of the 6- and 7-positions thereof, and further preferably the 6-position thereof.

[0038] In the case where R<sup>7</sup> is present, the substitution position thereof may be any position on the benzene ring. In the case where R<sup>8</sup> and R<sup>9</sup> are present, the substitution positions thereof may be any of the 2- to 6-positions.

[chem 13]

General Formula (3)



**[0039]** In the general formula (3), for the definitions and the preferred ranges of  $R^1$ ,  $R^2$ ,  $R^5$  to  $R^9$ ,  $n_1$ ,  $n_2$ , and  $n_5$  to  $n_9$ , reference may be made to the corresponding descriptions therefor in the general formula (1).

**[0040]** In the general formula (3),  $R^3$  and  $R^4$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group. For the descriptions and the preferred ranges of the carbazolyl group, the aryl group, the heteroaryl group, the alkyl group, the cycloalkyl group and the substituent, reference may be made to the corresponding descriptions for  $R^1$  and  $R^2$  in the general formula (1).

**[0041]** In the general formula (3),  $n_3$  and  $n_4$  each independently represent an integer of from 0 to 4, preferably an integer of from 0 to 3, and more preferably an integer of from 0 to 2. In the case where  $n_3$  is 2 or more, plural groups represented by  $R^3$  may be the same or different, and in the case where  $n_4$  is 2 or more, plural groups represented by  $R^4$  may be the same or different.

**[0042]**  $n_1$  to  $n_4$  may be the same as or different from each other. Examples of the case where  $n_1$  to  $n_4$  are the same as each other include the case where all of them are 0, the case where all of them are 1, and the case where all of them are 2. Examples of the case where  $n_1$  to  $n_4$  are different from each other include the case where  $n_1$  and  $n_3$  are 1, and  $n_2$  and  $n_4$  are 0, and the case where  $n_1$  and  $n_2$  are 1, and  $n_3$  and  $n_4$  are 0. In the general formula (3), at least one of  $n_1$  to  $n_4$  is preferably an integer of from 1 to 4, and  $n_1$  to  $n_4$  each preferably independently represent an integer of from 1 to 4.

**[0043]** In the general formula (3), in the case where at least one of  $R^1$  to  $R^4$  is present, the bonding position thereof may be any of the 1- to 8-positions of the carbazole ring, and is preferably any of the 2- to 7-positions, more preferably any of the 3-, 4-, 6- and 7-positions, and further preferably any of the 3- and 6-positions. Examples of the case include the case where  $R^1$  is bonded only to the 3-position, the case where  $R^1$  is bonded to the 3-position, and  $R^2$  is bonded to the 6-position, the case where  $R^1$  and  $R^3$  are bonded to the 3-position, and the case where  $R^1$  and  $R^3$  are bonded to the 3-position, and  $R^2$  and  $R^4$  are bonded to the 6-position. In the general formula (3), in the case where two or more of  $R^1$  to  $R^4$  are present, they may be the same as or different from each other.

**[0044]** In the general formula (3), in the case where  $R^5$  is present, the substitution position thereof may be any of the positions among the 1- to 4-positions of the carbazole ring that do not have the carbazolyl group substituted thereon. In the case where  $R^6$  is present, the substitution position thereof may be any of the positions among the 5- to 8-positions of the carbazole ring that do not have the carbazolyl group substituted thereon. The carbazolyl group is preferably substituted on the 3- and 6-positions of the carbazole ring.

**[0045]** In the case where  $R^7$  is present, the substitution position thereof may be any position on the benzene ring. In the case where  $R^8$  and  $R^9$  are present, the substitution positions thereof may be any of the 2- to 6-positions.

**[0046]** Specific examples of the compound represented by the general formula (1) are shown below, but the compound


represented by the general formula (1) capable of being used in the invention is not construed as being limited to the specific examples.

[chem 14]


5

10

15



Compound 1




Compound 2


20

25

30



Compound 3



Compound 4

35

40


45

50

55


[chem 15]

5



Compound 5


10



Compound 6


15

20



Compound 7

25



Compound 8

30

35

40


45

50

55




5



Compound 15

10




Compound 16

15


[chem 18]

20



Compound 17


25



Compound 18


30

35



Compound 19

40



Compound 20

45

50


55

[chem 19]


5

10

15



Compound 21




Compound 22

20


25

30

35



Compound 23



Compound 24


[chem 20]

40


45

50

55



Compound 25



Compound 26


[chem 21]

5


10

15

20



Compound 27




Compound 28

25

30

35

40



Compound 29

45

50

55


[chem 22]

5


10

15

20



Compound 30




Compound 31

25


30

35

40



Compound 32



Compound 33

45

50


55

[chem 23]


5

10

15



Compound 34



Compound 35


20

25

30

35

40



Compound 36

45

50

55


[chem 24]

5


10

15

20



Compound 37




Compound 38

25


30

35

40



Compound 39



Compound 40

45

50

55


[chem25]

5


10

15

20



Compound 41




Compound 42

25


30

35

40



Compound 43



Compound 44

45

50

55


[chem 26]

5


10

15

20



Compound 45




Compound 46

25


30

35

40



Compound 47




Compound 48

45

50

55

[chem 27]



Compound 49

**[0047]** The molecular weight of the compound represented by the general formula (1) is preferably 1,500 or less, more preferably 1,200 or less, further preferably 1,000 or less, and still further preferably 800 or less, for example, in the case where an organic layer containing the compound represented by the general formula (1) is intended to be formed as a film by a vapor deposition method. The lower limit of the molecular weight is 639 or more for the compound represented by the general formula (2), and 804 or more for the compound represented by the general formula (3).

**[0048]** The compound represented by the general formula (1) may be formed into a film by a coating method irrespective of the molecular weight thereof. The compound that has a relatively large molecular weight may be formed into a film by a coating method.

**[0049]** As an application of the invention, it may be considered that a compound that contains plural structures each represented by the general formula (1) in the molecule is used in a light emitting layer of an organic light emitting device.



**[0050]** For example, it may be considered that a polymerizable monomer having a structure represented by the general formula (1) is polymerized to form a polymer, and the polymer is used in a light emitting layer of an organic light emitting device. Specifically, it may be considered that a monomer that has a polymerizable functional group at any of  $R^1$ ,  $R^2$  and  $R^5$  to  $R^{10}$ , preferably any of  $R^1$ ,  $R^2$ ,  $R^8$ ,  $R^9$  and  $R^{10}$ , in the general formula (1) is prepared, and is homopolymerized or copolymerized with another monomer to prepare a polymer containing repeating units, and the polymer is used in a light emitting layer of an organic light emitting device. In alternative, it may be considered that the compounds represented by the general formula (1) are coupled to form a dimer or a trimer, and the dimer or the trimer is used in a light emitting layer of an organic light emitting device.

**[0051]** Examples of the structure of the repeating unit constituting the polymer containing the structure represented by the general formula (1) include a structure represented by the general formula (1), in which any of  $R^1$ ,  $R^2$  and  $R^5$  to  $R^{10}$ , preferably any of  $R^1$ ,  $R^2$ ,  $R^8$ ,  $R^9$  and  $R^{10}$ , in the general formula (1) has a structure represented by the following general formula (17) or (18).

[chem 28]

General Formula (17)

General Formula (18)




[0052] In the general formulae (17) and (18),  $L^1$  and  $L^2$  each represent a linking group. The linking group preferably has from 0 to 20 carbon atoms, more preferably from 1 to 15 carbon atoms, and further preferably from 2 to 10 carbon atoms. The linking group preferably has a structure represented by  $-X^{11}-L^{11}-$ , wherein  $X^{11}$  represents an oxygen atom or a sulfur atom, and preferably an oxygen atom, and  $L^{11}$  represents a linking group, preferably a substituted or unsubstituted alkylene group having from 1 to 10 carbon atoms or a substituted or unsubstituted arylene group, and more preferably a substituted or unsubstituted alkylene group having from 1 to 10 carbon atoms or a substituted or unsubstituted phenylene group.

[0053] In the general formulae (17) and (18),  $R^{101}$ ,  $R^{102}$ ,  $R^{103}$  and  $R^{104}$  each independently represent a substituent, preferably a substituted or unsubstituted alkyl group having from 1 to 6 carbon atoms, a substituted or unsubstituted alkoxy group having from 1 to 6 carbon atoms, or a halogen atom, more preferably a substituted or unsubstituted alkyl group having from 1 to 3 carbon atoms, a substituted or unsubstituted alkoxy group having from 1 to 3 carbon atoms, a fluorine atom or a chlorine atom, and further preferably a substituted or unsubstituted alkyl group having from 1 to 3 carbon atoms or a substituted or unsubstituted alkoxy group having from 1 to 3 carbon atoms.


[0054] Specific examples of the structure of the repeating unit include a structure, in which any of  $R^1$ ,  $R^2$  and  $R^5$  to  $R^{10}$ , preferably any of  $R^1$ ,  $R^2$ ,  $R^8$ ,  $R^9$  and  $R^{10}$ , in the general formula (1) has a structure represented by any of the following formulae (21) to (24). Two or more of  $R^1$ ,  $R^2$  and  $R^5$  to  $R^{10}$  may have a structure represented by any of the following formulae (21) to (24), and the case where only one of  $R^1$ ,  $R^2$  and  $R^5$  to  $R^{10}$  has a structure represented by any of the following formulae (21) to (24) is preferred.

[chem 29]


Formula (21)




Formula (22)



Formula (23)




Formula (24)



**[0055]** The polymer having the repeating unit containing the structure represented by any of the formulae (21) to (24) may be synthesized in such a manner that a hydroxyl group is introduced to least one of R<sup>1</sup>, R<sup>2</sup> and R<sup>5</sup> to R<sup>10</sup>, preferably at least one of R<sup>1</sup>, R<sup>2</sup>, R<sup>8</sup>, R<sup>9</sup> and R<sup>10</sup>, in the general formula (1), and the hydroxyl group as a linker is reacted with the following compound to introduce a polymerizable group thereto, followed by polymerizing the polymerizable group.

[chem 30]




**[0056]** The polymer containing the structure represented by the general formula (1) may be a polymer containing only a repeating unit having the structure represented by the general formula (1), or a polymer further containing a repeating unit having another structure. The repeating unit having the structure represented by the general formula (1) contained in the polymer may be only one kind or two or more kinds. Examples of the repeating unit that does not have the structure represented by the general formula (1) include a repeating unit derived from a monomer that is used for ordinary copolymerization. Examples of the repeating unit include a repeating unit derived from a monomer having an ethylenic unsaturated bond, such as ethylene and styrene.

Synthesis Method of Compound represented by General Formula (2)

**[0057]** In the compound represented by the general formula (1), the compound represented by the general formula (2) is a novel compound. The synthesis method of the compound represented by the general formula (2) is not particularly limited, and the compound represented by the general formula (2) may be synthesized by appropriately combining the known synthesis methods and conditions. For example, the compound represented by the general formula (2) may be synthesized through the following reaction.

[chem 31]




[0058] In the above formulae,  $R^1$ ,  $R^2$ ,  $R^5$  to  $R^9$ ,  $n_1$ ,  $n_2$ , and  $n_5$  to  $n_9$  have the same definition as in the general formula (2), and X represents a halogen atom, preferably a chlorine atom, a bromine atom or an iodine atom, and more preferably a bromine atom. The aforementioned reaction may be performed by appropriately optimizing the conditions that are used for the ordinary coupling reaction of an amine and a halide. The amine and the halide as reaction raw materials may be synthesized by utilizing the known synthesis methods. For example, for the synthesis method of a carbazoly-carbazole, reference may be made to Appl. Phys. Lett., 101, 093306 (2012), and the like. For the details of the reaction, reference may be made to the synthesis examples described later. The compound represented by the general formula (2) may also be synthesized by combining the other known synthesis reactions.

Compound represented by General Formula (4)

[0059] In the compound represented by the general formula (1), the compound represented by the following general formula (4) is a novel compound.

[chem 32]  
General Formula (4)




[0060] In the general formula (4),  $R^{11}$  to  $R^{14}$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;  $R^{15}$  and  $R^{16}$  each independently represent an alkyl group;  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;  $n11$  to  $n14$  and  $n17$  each independently represent an integer of from 0 to 4;  $n15$  and  $n16$  each represent an integer of from 0 to 3; and  $n18$  and  $n19$  each independently represent an integer of from 0 to 5, provided that at least one of  $n11$  to  $n14$  represents an integer of from 1 to 4, and when  $n11$  to  $n19$  each represent an integer of 2 or more, plural groups represented by  $R^{11}$  to  $R^{19}$  corresponding to  $n11$  to  $n19$  respectively each may be the same as or different from each other.

[0061] In the general formula (4), the preferred ranges of  $R^{11}$  to  $R^{19}$  and  $n11$  and  $n19$ , reference may be made to the descriptions for  $R^1$  to  $R^9$  and  $n1$  to  $n9$  in the general formula (1), provided that  $n11$  to  $n14$  in the general formula (4) are different from  $n1$  to  $n4$  in the general formula (1) in such a point that that at least one of  $n11$  to  $n14$  represents an integer of from 1 to 4. In the general formula (4), at least two of  $n11$  to  $n14$  each preferably represent an integer of from 1 to 4. For example, all  $n11$  to  $n14$  each preferably represent an integer of from 1 to 4, and examples of the case include the case where all  $n11$  to  $n14$  represent 1, the case where  $n11$  and  $n12$  represent 1, and  $n13$  and  $n14$  represent 0, and the case where  $n11$  and  $n13$  represent 1, and  $n12$  and  $n14$  represent 0. In the case where  $R^{11}$  to  $R^{14}$  are present,  $R^{11}$  and  $R^{13}$  are preferably bonded to the 3-position of the carbazole ring, and  $R^{12}$  and  $R^{14}$  are preferably bonded to the 6-position of the carbazole ring.

Synthesis Method of Compound represented by General Formula (4)

[0062] The synthesis method of the compound represented by the general formula (4) is not particularly limited, and the compound represented by the general formula (4) may be synthesized by appropriately combining the known synthesis methods and conditions. For example, the compound represented by the general formula (4) may be synthesized through the following reaction.

[chem 33]



[0063] In the above formulae,  $R^1$  to  $R^9$ ,  $n1$  to  $n9$  have the same definition as in the general formula (3), and  $X$  represents a halogen atom, preferably a chlorine atom, a bromine atom or an iodine atom, and more preferably a bromine atom. The aforementioned reaction may be performed by appropriately optimizing the conditions that are used for the ordinary coupling reaction of an amine and a halide. The amine and the halide as reaction raw materials may be synthesized by utilizing the known synthesis methods. For example, for the synthesis method of a carbazolylicarbazole, reference may be made to Appl. Phys. Lett., 101, 093306 (2012), and the like. For the details of the reaction, reference may be made to the synthesis examples described later. The compound represented by the general formula (4) may also be synthesized by combining the other known synthesis reactions.

#### Organic Light Emitting Device

[0064] The compound represented by the general formula (1) of the invention is useful as a light emitting material of an organic light emitting device. Accordingly, the compound represented by the general formula (1) of the invention may be effectively used as a light emitting material in a light emitting layer of an organic light emitting device. The compound represented by the general formula (1) includes a delayed fluorescent material emitting delayed fluorescent light, i.e. a delayed fluorescence emitter. Thus, the invention provides an invention relating to a delayed fluorescence emitter having the structure represented by the general formula (1), an invention relating to the use of the compound represented by the general formula (1) as a fluorescence emitter, and an invention relating to a method for emitting delayed fluorescent light with the compound represented by the general formula (1). An organic light emitting device that uses the compound as a light emitting material has features that the device emits delayed fluorescent light and has a high light emission efficiency. The principle of the features may be described as follows for an organic electroluminescent device as an example.

**[0065]** In an organic electroluminescent device, carriers are injected from an anode and a cathode to a light emitting material to form an excited state for the light emitting material, with which light is emitted. In the case of a carrier injection type organic electroluminescent device, in general, excitons that are excited to the excited singlet state are 25% of the total excitons generated, and the remaining 75% thereof are excited to the excited triplet state. Accordingly, the use of phosphorescence, which is light emission from the excited triplet state, provides a high energy utilization. However, the excited triplet state has a long lifetime and thus causes saturation of the excited state and deactivation of energy through mutual action with the excitons in the excited triplet state, and therefore the quantum yield of phosphorescence may generally be often not high. A delayed fluorescent material emits fluorescent light through the mechanism that the energy of excitons transits to the excited triplet state through intersystem crossing or the like, and then transits to the excited singlet state through reverse intersystem crossing due to triplet-triplet annihilation or absorption of thermal energy, thereby emitting fluorescent light. It is considered that among the materials, a thermal activation type delayed fluorescent material emitting light through absorption of thermal energy is particularly useful for an organic electroluminescent device. In the case where a delayed fluorescent material is used in an organic electroluminescent device, the excitons in the excited singlet state normally emit fluorescent light. On the other hand, the excitons in the excited triplet state emit fluorescent light through intersystem crossing to the excited singlet state by absorbing the heat generated by the device. At this time, the light emitted through reverse intersystem crossing from the excited triplet state to the excited single state has the same wavelength as fluorescent light since it is light emission from the excited single state, but has a longer lifetime (light emission lifetime) than the normal fluorescent light and phosphorescent light, and thus the light is observed as fluorescent light that is delayed from the normal fluorescent light and phosphorescent light. The light may be defined as delayed fluorescent light. The use of the thermal activation type exciton transition mechanism may raise the proportion of the compound in the excited single state, which is generally formed in a proportion only of 25%, to 25% or more through the absorption of the thermal energy after the carrier injection. A compound that emits strong fluorescent light and delayed fluorescent light at a low temperature of lower than 100°C undergoes the intersystem crossing from the excited triplet state to the excited singlet state sufficiently with the heat of the device, thereby emitting delayed fluorescent light, and thus the use of the compound may drastically enhance the light emission efficiency.

**[0066]** The use of the compound represented by the general formula (1) of the invention as a light emitting material of a light emitting layer may provide an excellent organic light emitting device, such as an organic photoluminescent device (organic PL device) and an organic electroluminescent device (organic EL device). At this time, the compound represented by the general formula (1) of the invention may have a function of assisting light emission of another light emitting material contained in the light emitting layer, i.e., as a so-called assist dopant. Specifically, the compound represented by the general formula (1) of the invention contained in the light emitting layer may have a lowest excited singlet energy that is between the lowest excited singlet energy of the host material contained in the light emitting layer and the lowest excited singlet energy of the another light emitting material contained in the light emitting layer.

**[0067]** The organic photoluminescent device has a structure containing a substrate having formed thereon at least a light emitting layer. The organic electroluminescent device has a structure containing at least an anode, a cathode and an organic layer formed between the anode and the cathode. The organic layer contains at least a light emitting layer, and may be formed only of a light emitting layer, or may have one or more organic layer in addition to the light emitting layer. Examples of the organic layer include a hole transporting layer, a hole injection layer, an electron barrier layer, a hole barrier layer, an electron injection layer, an electron transporting layer and an exciton barrier layer. The hole transporting layer may be a hole injection and transporting layer having a hole injection function, and the electron transporting layer may be an electron injection and transporting layer having an electron injection function. A specific structural example of an organic electroluminescent device is shown in Fig. 1. In Fig. 1, the numeral 1 denotes a substrate, 2 denotes an anode, 3 denotes a hole injection layer, 4 denotes a hole transporting layer, 5 denotes a light emitting layer, 6 denotes an electron transporting layer, and 7 denotes a cathode.

**[0068]** The members and the layers of the organic electroluminescent device will be described below. The descriptions for the substrate and the light emitting layer may also be applied to the substrate and the light emitting layer of the organic photoluminescent device.

Substrate

**[0069]** The organic electroluminescent device of the invention is preferably supported by a substrate. The substrate is not particularly limited and may be those that have been commonly used in an organic electroluminescent device, and examples thereof used include those formed of glass, transparent plastics, quartz and silicon.

Anode

**[0070]** The anode of the organic electroluminescent device used is preferably formed of as an electrode material a metal, an alloy or an electroconductive compound each having a large work function (4 eV or more), or a mixture thereof.

Specific examples of the electrode material include a metal, such as Au, and an electroconductive transparent material, such as CuI, indium tin oxide (ITO), SnO<sub>2</sub> and ZnO. A material that is amorphous and is capable of forming a transparent electroconductive film, such as IDIXO (In<sub>2</sub>O<sub>3</sub>-ZnO), may also be used. The anode may be formed in such a manner that the electrode material is formed into a thin film by such a method as vapor deposition or sputtering, and the film is patterned into a desired pattern by a photolithography method, or in the case where the pattern may not require high accuracy (for example, approximately 100 μm or more), the pattern may be formed with a mask having a desired shape on vapor deposition or sputtering of the electrode material. In alternative, in the case where a material capable of being applied as a coating, such as an organic electroconductive compound, is used, a wet film forming method, such as a printing method and a coating method, may be used. In the case where emitted light is to be taken out through the anode, the anode preferably has a transmittance of more than 10%, and the anode preferably has a sheet resistance of several hundred Ohm per square or less. The thickness thereof may be generally selected from a range of from 10 to 1,000 nm, and preferably from 10 to 200 nm, while depending on the material used.

#### Cathode

**[0071]** The cathode is preferably formed of as an electrode material a metal (referred to as an electron injection metal), an alloy or an electroconductive compound each having a small work function (4 eV or less), or a mixture thereof. Specific examples of the electrode material include sodium, a sodium-potassium alloy, magnesium, lithium, a magnesium-copper mixture, a magnesium-silver mixture, a magnesium-aluminum mixture, a magnesium-indium mixture, an aluminum-aluminum oxide (Al<sub>2</sub>O<sub>3</sub>) mixture, indium, a lithium-aluminum mixture, and a rare earth metal. Among these, a mixture of an electron injection metal and a second metal that is a stable metal having a larger work function than the electron injection metal, for example, a magnesium-silver mixture, a magnesium-aluminum mixture, a magnesium-indium mixture, an aluminum-aluminum oxide (Al<sub>2</sub>O<sub>3</sub>) mixture, a lithium-aluminum mixture, and aluminum, are preferred from the standpoint of the electron injection property and the durability against oxidation and the like. The cathode may be produced by forming the electrode material into a thin film by such a method as vapor deposition or sputtering. The cathode preferably has a sheet resistance of several hundred Ohm per square or less, and the thickness thereof may be generally selected from a range of from 10 nm to 5 μm, and preferably from 50 to 200 nm. For transmitting the emitted light, any one of the anode and the cathode of the organic electroluminescent device is preferably transparent or translucent, thereby enhancing the light emission luminance.

**[0072]** The cathode may be formed with the electroconductive transparent materials described for the anode, thereby forming a transparent or translucent cathode, and by applying the cathode, a device having an anode and a cathode, both of which have transmittance, may be produced.

#### Light Emitting Layer

**[0073]** The light emitting layer is a layer, in which holes and electrons injected from the anode and the cathode, respectively, are recombined to form excitons, and then the layer emits light. A light emitting material may be solely used as the light emitting layer, but the light emitting layer preferably contains a light emitting material and a host material. The light emitting material used may be one kind or two or more kinds selected from the group of compounds represented by the general formula (1) of the invention. In order that the organic electroluminescent device and the organic photoluminescent device of the invention exhibit a high light emission efficiency, it is important that the singlet excitons and the triplet excitons generated in the light emitting material are confined in the light emitting material. Accordingly, a host material is preferably used in addition to the light emitting material in the light emitting layer. The host material used may be an organic compound that has excited singlet energy and excited triplet energy, at least one of which is higher than those of the light emitting material of the invention. As a result, the singlet excitons and the triplet excitons generated in the light emitting material of the invention are capable of being confined in the molecules of the light emitting material of the invention, thereby eliciting the light emission efficiency thereof sufficiently. Even though the singlet excitons and the triplet excitons are not confined sufficiently, a high light emission efficiency may be obtained in some cases, and thus a host material that is capable of achieving a high light emission efficiency may be used in the invention without any particular limitation. In the organic light emitting device and the organic electroluminescent device of the invention, the light emission occurs in the light emitting material of the invention contained in the light emitting layer. The emitted light contains both fluorescent light and delayed fluorescent light. However, a part of the emitted light may contain emitted light from the host material, or the emitted light may partially contain emitted light from the host material.

**[0074]** In the case where the host material is used, the amount of the compound of the invention as the light emitting material contained in the light emitting layer is preferably 0.1% by weight or more, and more preferably 1% by weight or more, and is preferably 50% by weight or less, more preferably 20% by weight or less, and further preferably 10% by weight or less.

**[0075]** The host material in the light emitting layer is preferably an organic compound that has a hole transporting

function and an electron transporting function, prevents the emitted light from being increased in wavelength, and has a high glass transition temperature.

#### Injection Layer

5  
[0076] The injection layer is a layer that is provided between the electrode and the organic layer, for decreasing the driving voltage and enhancing the light emission luminance, and includes a hole injection layer and an electron injection layer, which may be provided between the anode and the light emitting layer or the hole transporting layer and between the cathode and the light emitting layer or the electron transporting layer. The injection layer may be provided depending on necessity.

#### Barrier Layer

15  
[0077] The barrier layer is a layer that is capable of inhibiting charges (electrons or holes) and/or excitons present in the light emitting layer from being diffused outside the light emitting layer. The electron barrier layer may be disposed between the light emitting layer and the hole transporting layer, and inhibits electrons from passing through the light emitting layer toward the hole transporting layer. Similarly, the hole barrier layer may be disposed between the light emitting layer and the electron transporting layer, and inhibits holes from passing through the light emitting layer toward the electron transporting layer. The barrier layer may also be used for inhibiting excitons from being diffused outside the light emitting layer. Thus, the electron barrier layer and the hole barrier layer each may also have a function as an exciton barrier layer. The term "the electron barrier layer" or "the exciton barrier layer" referred herein is intended to include a layer that has both the functions of an electron barrier layer and an exciton barrier layer by one layer.

#### Hole Barrier Layer

25  
[0078] The hole barrier layer has the function of an electron transporting layer in a broad sense. The hole barrier layer has a function of inhibiting holes from reaching the electron transporting layer while transporting electrons, and thereby enhances the recombination probability of electrons and holes in the light emitting layer. As the material for the hole barrier layer, the materials for the electron transporting layer described later may be used depending on necessity.

#### Electron Barrier Layer

30  
[0079] The electron barrier layer has the function of transporting holes in a broad sense. The electron barrier layer has a function of inhibiting electrons from reaching the hole transporting layer while transporting holes, and thereby enhances the recombination probability of electrons and holes in the light emitting layer.

#### Exciton Barrier Layer

40  
[0080] The exciton barrier layer is a layer for inhibiting excitons generated through recombination of holes and electrons in the light emitting layer from being diffused to the charge transporting layer, and the use of the layer inserted enables effective confinement of excitons in the light emitting layer, and thereby enhances the light emission efficiency of the device. The exciton barrier layer may be inserted adjacent to the light emitting layer on any of the side of the anode and the side of the cathode, and on both the sides. Specifically, in the case where the exciton barrier layer is present on the side of the anode, the layer may be inserted between the hole transporting layer and the light emitting layer and adjacent to the light emitting layer, and in the case where the layer is inserted on the side of the cathode, the layer may be inserted between the light emitting layer and the cathode and adjacent to the light emitting layer. Between the anode and the exciton barrier layer that is adjacent to the light emitting layer on the side of the anode, a hole injection layer, an electron barrier layer and the like may be provided, and between the cathode and the exciton barrier layer that is adjacent to the light emitting layer on the side of the cathode, an electron injection layer, an electron transporting layer, a hole barrier layer and the like may be provided. In the case where the barrier layer is provided, the material used for the barrier layer preferably has excited singlet energy and excited triplet energy, at least one of which is higher than the excited singlet energy and the excited triplet energy of the light emitting layer, respectively.

#### Hole Transporting Layer

55  
[0081] The hole transporting layer is formed of a hole transporting material having a function of transporting holes, and the hole transporting layer may be provided as a single layer or plural layers.

[0082] The hole transporting material has one of injection or transporting property of holes and barrier property of

electrons, and may be any of an organic material and an inorganic material. Examples of known hole transporting materials that may be used herein include a triazole derivative, an oxadiazole derivative, an imidazole derivative, a carbazole derivative, an indolocarbazole derivative, a polyaryllkane derivative, a pyrazoline derivative, a pyrazolone derivative, a phenylenediamine derivative, an arylamine derivative, an amino-substituted chalcone derivative, an oxazole derivative, a styrylanthracene derivative, a fluorenone derivative, a hydrazone derivative, a stilbene derivative, a silazane derivative, an aniline copolymer and an electroconductive polymer, particularly a thiophene oligomer. Among these, a porphyrin compound, an aromatic tertiary amine compound and a styrylamine compound are preferably used, and an aromatic tertiary amine compound is more preferably used.

#### 10 Electron Transporting Layer

**[0083]** The electron transporting layer is formed of a material having a function of transporting electrons, and the electron transporting layer may be provided as a single layer or plural layers.

**[0084]** The electron transporting material (which may also function as a hole barrier material in some cases) needs only to have a function of transporting electrons, which are injected from the cathode, to the light emitting layer. Examples of the electron transporting layer that may be used herein include a nitro-substituted fluorene derivative, a diphenylquinone derivative, a thiopyran dioxide derivative, carbodiimide, a fluorenylidene methane derivative, anthraquinodimethane and anthrone derivatives, and an oxadiazole derivative. The electron transporting material used may be a thiaziazole derivative obtained by replacing the oxygen atom of the oxadiazole ring of the oxadiazole derivative by a sulfur atom, or a quinoxaline derivative having a quinoxaline ring, which is known as an electron attracting group. Furthermore, polymer materials having these materials introduced to the polymer chain or having these materials used as the main chain of the polymer may also be used.


**[0085]** In the production of the organic electroluminescent device, the compound represented by the general formula (1) may be used not only in the light emitting layer but also in the other layers than the light emitting layer. In this case, the compound represented by the general formula (1) used in the light emitting layer and the compound represented by the general formula (1) used in the other layers than the light emitting layer may be the same as or different from each other. For example, the compound represented by the general formula (1) may be used in the injection layer, the barrier layer, the hole barrier layer, the electron barrier layer, the exciton barrier layer, the hole transporting layer, the electron transporting layer and the like described above. The film forming method of the layers are not particularly limited, and the layers may be produced by any of a dry process and a wet process.

**[0086]** Specific examples of preferred materials that may be used in the organic electroluminescent device are shown below, but the materials that may be used in the invention are not construed as being limited to the example compounds. The compound that is shown as a material having a particular function may also be used as a material having another function. In the structural formulae of the example compounds, R and R<sub>1</sub> to R<sub>10</sub> each independently represent a hydrogen atom or a substituent, and n represents an integer of from 3 to 5.

**[0087]** Preferred examples of a compound that may also be used as the host material of the light emitting layer are shown below.


[chem 34]

5




10

15




20

25



30

35



40



45

50

55



[chem 35]

5





10

15






20

25



30

35




40


45

50

55




[chem 36]




[chem 37]


5



10




15




20

25


30



35



40




45

50

55


[chem 38]

5




10

15



20

25




30

**[0088]** Preferred examples of a compound that may be used as the hole injection material are shown below.

[chem 39]

35


40



45


50

55



[0089] Preferred examples of a compound that may be used as the hole transporting material are shown below.


[chem 40]



50

55



5



10

15


20



25


[chem 41]

30



35

40




45

50


55

5



10


15



20

25


30



35

[chem 42]

40





45

50



55

5



10

15





20

25

30



[chem 43]

35



40


45




50

55

5




10



15

[chem 44]


20



25

30


35



40


45

50



55

[chem 45]



[0090] Preferred examples of a compound that may be used as the electron barrier material are shown below.


50

55

[chem 46]

5

10



**[0091]** Preferred examples of a compound that may be used as the hole barrier material are shown below.

15

[chem 47]

20

25

30


35

40



45

50

55





5



10

15




20

25

30

35



40

45

**[0092]** Preferred examples of a compound that may be used as the electron transporting material are shown below.

50


55

[chem 48]

5


10

15




20

25




30

35



40

45




50

55

[chem 49]


5

10



15

20




25

30


35

40




45

50




55

[chem 50]



[0093] Preferred examples of a compound that may be used as the electron injection material are shown below.

[chem 51]






[0094] Preferred examples of a compound as a material that may be added are shown below. For example, the compound may be added as a stabilizing material.

55



[chem 52]

5



10

15



20

25

**[0095]** The organic electroluminescent device thus produced by the aforementioned method emits light on application of an electric field between the anode and the cathode of the device. In this case, when the light emission is caused by the excited single energy, light having a wavelength that corresponds to the energy level thereof may be confirmed as fluorescent light and delayed fluorescent light. When the light emission is caused by the excited triplet energy, light having a wavelength that corresponds to the energy level thereof may be confirmed as phosphorescent light. The normal fluorescent light has a shorter light emission lifetime than the delayed fluorescent light, and thus the light emission lifetime may be distinguished between the fluorescent light and the delayed fluorescent light.

30

**[0096]** The phosphorescent light may substantially not be observed with a normal organic compound, such as the compound of the invention, at room temperature since the excited triplet energy is converted to heat or the like due to the instability thereof, and is immediately deactivated with a short lifetime. The excited triplet energy of the normal organic compound may be measured by observing light emission under an extremely low temperature condition.

35

**[0097]** The organic electroluminescent device of the invention may be applied to any of a single device, a structure with plural devices disposed in an array, and a structure having anodes and cathodes disposed in an X-Y matrix. According to the invention, an organic light emitting device that is largely improved in light emission efficiency may be obtained by adding the compound represented by the general formula (1) in the light emitting layer. The organic light emitting device, such as the organic electroluminescent device, of the invention may be applied to a further wide range of purposes. For example, an organic electroluminescent display apparatus may be produced with the organic electroluminescent device of the invention, and for the details thereof, reference may be made to S. Tokito, C. Adachi and H. Murata, "Yuki EL Display" (Organic EL Display) (Ohmsha, Ltd.). In particular, the organic electroluminescent device of the invention may be applied to organic electroluminescent illumination and backlight which are highly demanded.

40

Example

**[0098]** The features of the invention will be described more specifically with reference to synthesis examples and working examples below. The materials, processes, procedures and the like shown below may be appropriately modified unless they deviate from the substance of the invention. Accordingly, the scope of the invention is not construed as being limited to the specific examples shown below.


50

Synthesis Example 1

55

**[0099]** In this synthesis example, a compound 1 was synthesized according to the following scheme.

[chem 53]



15 **[0100]** 3-Carbazolylcarbazole (0.25 g, 0.75 mmol), 2-bromo-4,6-diphenyl-1,3,5-triazine (0.29 g, 0.75 mmol), copper iodide (0.021 g, 0.11 mmol), 18-crown-6-ether (0.030 g, 0.11 mmol) and potassium carbonate (0.62 g, 4.5 mmol) were added to dodecylbenzene (2.0 mL), and the mixture was heated to 220°C under a nitrogen gas environment for 2 days. After completing the reaction, the product was extracted with dichloromethane, dried over sodium carbonate, and then purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 80/20), thereby providing the compound 1 (yield amount: 0.25 g (4.0 mmol), yield: 53%).

20  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ ) :  $\delta$  (ppm) 9.08 (d,  $J = 9.0$  Hz, 2H), 8.84 (d,  $J = 6.5$  Hz, 4H), 8.32 (d,  $J = 2.0$  Hz, 1H), 8.20 (d,  $J = 8.0$  Hz, 2H), 8.15 (d,  $J = 7.5$  Hz, 1H), 7.91 (d,  $J = 8.5$  Hz, 2H), 7.75 (d,  $J = 9.0$  Hz, 1H), 7.67-7.59 (m, 8H), 7.55-7.51 (m, 1H), 7.45-7.41 (m, 4H), 7.38-7.35 (m, 1H), 7.33-7.29 (m, 2H)

#### Synthesis Example 2

25 **[0101]** In this synthesis example, a compound 2 was synthesized according to the following schemes.

30

35


40

45


50

55


[chem 54]



Scheme 1



Scheme 2



Scheme 3

## Scheme 1

40

## Synthesis of 3,6-Di-tert-butyl-9-(9-tosyl-9H-carbazol-3-yl)-9H-carbazole

45

**[0102]** 3-Bromo-9-tosyl-9H-carbazole (4.47 g, 11.5 mmol), 3,6-di-tert-butyl-9H-carbazole (3.04 g, 10.9 mmol), and copper (I) oxide (3.96g, 27.6 mmol) were added to dodecylbenzene (2.0 mL), and the mixture was heated to 220°C under a nitrogen gas environment for 20 hours. After completing the reaction, copper (I) oxide was removed by filtration, and then the product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 70/30) (yield amount: 2.21 g (11.5 mmol), yield: 32.2%).

50

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ (ppm) 8.38 (d, J = 8.5 Hz, 1H), 8.16 (d, J = 1.5 Hz, 1H), 8.07 (d, J = 2.0 Hz, 1H), 7.87 (d, J = 7.5 Hz, 1H), 7.80 (d, J = 8.5 Hz, 2H), 7.66 (dd, J = 2.5 Hz, J = 2.0 Hz, 1H), 7.57-7.53 (m, 1H), 7.46 (dd, J = 2.0 Hz, J = 1.5 Hz, 2H), 7.40-7.37 (m, 1H), 7.32 (d, J = 9.0 Hz, 2H), 7.20 (d, J = 8.0 Hz, 2H), 2.33 (s, 3H), 1.47 (s, 18H)

## Scheme 2

55

## Synthesis of 3,6-Di-tert-butyl-9-(9H-carbazol-3-yl)-9H-carbazole

**[0103]** 3,6-di-tert-butyl-9-(9-tosyl-9H-carbazol-3-yl)-9H-carbazole (2.08g, 3.47 mmol) and potassium hydroxide (22.2 g, 39.5 mmol) were added to a mixed solvent of tetrahydrofuran (4.6 mL), dimethylsulfoxide (2.3 mL) and water (0.7 mL), and the mixture was stirred at 70°C for 5 hours. After completing the reaction, the reaction mixture was neutralized

## EP 2 949 724 A1

with sulfuric acid, and the product was extracted with toluene, which was then dried over sodium sulfate. After removing the solvent with an evaporator, the product was purified by recrystallization (isopropanol/methanol = 50/50) (yield amount: 1.17 g (2.51 mmol), yield: 72.5%).

5  $^1\text{H-NMR}$  (500 MHz, DMSO) :  $\delta$  (ppm) 11.54 (s, 1H), 8.34 (d, J = 2.0 Hz, 1H), 8.30 (d, J = 1.5 Hz, 2H), 8.20 (d, J = 8.0 Hz, 1H), 7.72 (d, J = 8.5 Hz, 1H), 7.56 (d, J = 8.5 Hz, 1H), 7.53 (dd, J = 2.0 Hz, J = 2.0 Hz, 1H), 7.48-7.44 (m, 3H), 7.25 (d, J = 8.5 Hz, 2H), 7.19-7.16 (m, 1H), 1.43 (s, 18H)

### Scheme 3

10 Synthesis of 9'-(4-(4,6-Diphenyl-1,3,5-triazin-2-yl)phenyl)-9'H-(3',6'-d i-tert-butylcarbazolyl)carbazole

15 **[0104]** Copper iodide (0.01 g, 0.05 mmol), 18-crown-5-ether (0.027 g, 0.10 mmol), potassium carbonate (0.138 g, 1.00 mmol) and dodecylbenzene (0.9 mL) were added to 3-(3,6-di-tert-butylcarbazolyl)carbazole (0.34 g, 0.77 mmol) and 4-bromophenyldiphenyltriazine (0.38 g, 0.97 mmol), and the mixture was reacted at 220°C overnight. After completing the reaction, dichloromethane was added thereto, and the mixture was filtered with Celite. The filtrate was concentrated and then extracted with dichloromethane. The organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: cyclohexane/dichloromethane = 70/30), thereby providing the compound 2 (yield amount: 0.3 g, crude yield: 51.9%).

MALDI-TOF-MS  $m/z$  = 751 ( $[\text{M}]^+$ )

20  $^1\text{H-NMR}$  (500 MHz, DMSO) :  $\delta$  (ppm) 9.08 (d, J = 8.5 Hz, 2H), 8.82 (d, J = 7.5 Hz, 4H), 8.57 (d, J = 2.0 Hz, 1H), 8.41 (d, J = 7.5 Hz, 1H), 8.33 (d, J = 2.0 Hz, 2H), 8.08 (d, J = 8.5 Hz, 2H), 7.82 (d, 9.0 Hz, 1H), 7.76-7.65 (m, 8H), 7.58-7.55 (m, 1H), 7.50 (dd, J = 1.5 Hz, J = 1.5 Hz, 2H), 7.39-7.36 (m, 1H), 7.33 (d, 9.0 Hz, 2H), 1.43 (s, 18H)

### Synthesis Example 3

25

**[0105]** In this synthesis example, a compound 3 was synthesized according to the following schemes.

30

35





40

45

50

55

[chem 55]



## Scheme 1

45

## Synthesis of 3,6-Diphenylcarbazole

**[0106]** Toluene (200 mL), ethanol (100 mL) and water (53 mL) were added to 3,6-dibromocarbazole (13 g, 40 mmol), phenylboronic acid (11.7 g, 96 mmol) and sodium carbonate (21.9 g, 207 mmol), and after deaeration, the mixture was stirred. Pd(PPh<sub>3</sub>)<sub>4</sub> (4.16 g, 3.99 mmol) was further added thereto, and after deaeration, the mixture was reacted at 80°C for 12 hours. After completing the reaction, the product was extracted with dichloromethane. The organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: cyclohexane/dichloromethane = 50/50), and recrystallized from hexane (yield amount: 4.32 g, yield: 33.8%).

55 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ (ppm) 8.34 (d, J = 1.5 Hz, 2H), 8.11 (s, 1H), 7.72 (dd, J = 1.0 Hz, J = 1.0 Hz, 4H), 7.69 (dd, J = 1.5 Hz, J = 2.0 Hz, 2H), 7.51-7.46 (m, 6H), 7.36-7.33 (m, 2H)

## Scheme 2

## Synthesis of 3,6-Diphenyl-9-tosyl-9H-carbazole

- 5 **[0107]** 3-Bromo-9-tosyl-9H-carbazole (4.20 g, 13.2 mmol), 3,6-diphenyl-9H-carbazole (4.87 g, 12.5 mmol) and copper(I) oxide (4.54 g, 31.8 mmol) were added to dodecylbenzene (12.3 mL), and the mixture was heated to 220°C under a nitrogen gas environment for 20 hours. After completing the reaction, copper (I) oxide was removed by filtration, and then the filtrate was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 50/50) (yield amount: 3.24 g (5.07 mmol), yield: 38.5%).
- 10 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ (ppm) 8.58 (d, J = 8.5 Hz, 1H), 8.43 (d, J = 1.5 Hz, 2H), 8.41 (d, J = 8.5 Hz, 1H), 8.14 (d, J = 2.0 Hz, 1H), 7.92 (d, J = 7.5 Hz, 1H), 7.84 (d, J = 8.5 Hz, 2H), 7.75-7.72 (m, 6H), 7.68 (dd, J = 1.5 Hz, J=2.0Hz, 2H), 7.59-7.56 (m, 1H), 7.52-7.45 (m, 5H), 7.42-7.39 (m, 1H), 7.37-7.35 (m, 2H), 2.34 (s, 3H)

## Scheme 3

- 15 Synthesis of 9-(9H-Carbazol-3-yl)-3,6-diphenyl-9H-carbazole

- 20 **[0108]** 3,6-Diphenyl-9-tosyl-9H-carbazole (3.22 g, 5.03 mmol) and potassium hydroxide (3.22 g, 57.3 mmol) were added to a mixed solvent of tetrahydrofuran (6.7mL), dimethylsulfoxide (3.3mL) and water (1.0 mL), and the mixture was stirred at 70°C for 5 hours. After completing the reaction, the reaction mixture was neutralized with sulfuric acid, and the product was extracted with toluene, which was then dried over sodium sulfate. After removing the solvent with an evaporator, the product was purified by recrystallization (chloroform/hexane = 50/50) (yield amount: 1.20 g (2.48 mmol), yield: 49.2%).
- 25 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ (ppm) 8.44 (d, J = 2.0 Hz, 2H), 8.29 (s, 1H), 8.27 (s, 1H), 8.08 (d, J = 8.0 Hz, 1H), 7.75 (d, J = 7.0 Hz, 4H), 7.68 (dd, J=1.5 Hz, J = 1.5 Hz, 2H), 7.60 (dd, J = 1.5 Hz, J = 2.0 Hz, 1H), 7.50-7.47 (m, 8H), 7.37-7.34 (m, 2H), 7.30-7.27 (m, 1H)

## Scheme 4

- 30 Synthesis of 9'-(4-(4,6-Diphenyl-1,3,5-triazin-2-yl)phenyl)-9'H-(3',6'-d iphenylcarbazolyl)carbazole

- 35 **[0109]** Copper iodide (0.010 g, 0.04 mmol), 18-crown-5-ether (0.038 g, 0.14 mmol), potassium carbonate (0.10 g, 0.72 mmol) and dodecylbenzene (0.72 mL) were added to 3-(3,6-diphenylcarbazolyl)carbazole (0.35 g, 0.72 mmol) and 4-bromophenyldiphenyltriazine (0.31 g, 0.80 mmol), and the mixture was reacted at 220°C overnight. After completing the reaction, dichloromethane was added thereto, and the mixture was filtered with Celite. The filtrate was concentrated and then extracted with dichloromethane. The organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 70/30), thereby providing the compound 3 (yield amount: 0.10 g, crude yield: 17.5%).
- 40 <sup>1</sup>H-NMR (500 MHz, DMSO): δ (ppm) 9.09 (d, J = 8.5 Hz, 2H), 8.82 (d, J = 7.0 Hz, 4H), 8.78 (d, J = 1.5 Hz, 2H), 8.68 (d, J = 2.0 Hz, 1H), 8.43 (d, J = 8.0 Hz, 1H), 8.09 (d, J = 8.5 Hz, 2H), 7.85 (d, 7.5 Hz, 5H), 7.81 (dd, J = 1.5 Hz, J = 1.5 Hz, 2H), 7.76-7.67 (m, 9H), 7.53-7.49 (m, 6H), 7.42-7.35 (m, 3H) MALDI-TOF-MS: m/z = 791 ([M]<sup>+</sup>)


## Synthesis Example 4

- 45 **[0110]** In this synthesis example, a compound 4 was synthesized according to the following schemes.


50

55


[chem 56]



Scheme 1



Scheme 2



Scheme 3

## Scheme 1

## 35 Synthesis of 9-(9-(9-Tosyl-9H-carbazolyl-3-yl)-9H-carbazolyl-3-yl)-9H-carbazole

**[0111]** 3-Bromo-9-tosyl-9H-carbazole (1.70 g, 4.36 mmol), 3-carbazolylcarbazole (1.52 g, 4.59 mmol) and copper(I) oxide (2.22 g, 15.5 mmol) were added to dodecylbenzene (4.3 mL), and the mixture was reacted at 220°C under an inert atmosphere for 20 hours. After completing the reaction, copper(I) oxide was removed by filtration, and the product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 50/50) (yield amount: 1.14 g (1.75 mmol), yield: 38.1%).

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): 8.60 (d, J = 9.0 Hz, 1H), 8.41 (d, J = 8.5 Hz, 1H), 8.31 (d, J = 1.0 Hz, 1H), 8.20-8.17 (m, 3H), 8.14 (d, J = 8.0 Hz, 1H), 7.94 (d, J = 7.5 Hz, 1H), 7.84 (d, J = 8.5 Hz, 2H), 7.75 (dd, J = 2.0 Hz, J = 2.0 Hz, 1H), 7.58-7.54 (m, 3H), 7.48-7.40 (m, 7H), 7.35-7.29 (m, 3H), 7.22 (d, J = 8.0 Hz, 2H), 2.34 (s, 3H)

## Scheme 2

## Synthesis of 9-(9-(9H-carbazolyl-3-yl)-9H-carbazolyl-3-yl)-9H-carbazole

**[0112]** 9-(9-(9-Tosyl-9H-carbazolyl-3-yl)-9H-carbazolyl-3-yl)-9H-carbazole (1.05g, 2.16 mmol) and potassium hydroxide (1.38 g, 24.6 mmol) were added to a mixed solvent of tetrahydrofuran (2.9 mL), dimethylsulfoxide (1.4 mL) and water (0.4 mL), and the mixture was stirred at 70°C for 5 hours. After completing the reaction, the reaction mixture was neutralized with sulfuric acid, and the product was extracted with toluene, which was then dried over sodium sulfate. After removing the solvent with an evaporator, the product was purified by recrystallization (chloroform/hexane = 20/80) (yield amount: 0.62 g (1.3 mmol), yield: 59%).

<sup>1</sup>H-NMR (500 MHz, DMSO): 11.62 (s, 1H), 8.56 (d, J = 2.0 Hz, 1H), 8.50 (d, J = 2.0 Hz, 1H), 8.37 (d, J = 7.5 Hz, 1H), 8.29 (d, J = 8.0 Hz, 2H), 8.25 (d, J = 8.0 Hz, 1H), 7.80 (d, J = 8.5 Hz, 1H), 7.66 (dd, J = 2.0 Hz, J = 2.0 Hz, 1H), 7.61-7.7.59 (m, 3H), 7.50-7.43 (m, 4H), 7.41-7.38 (m, 3H), 7.32-7.29 (m, 3H), 7.22-7.19 (m, 1H)

## Scheme 3

Synthesis of 9'-(4-(4,6-Diphenyl-1,3,5-triazin-2-yl)phenyl)-9'H-3,9':3', 9"-tercarbazole

5 **[0113]** Dodecylbenzene (0.6 mL) was added to 3-(3-(3-carbazolyl)carbazolyl)carbazole (0.26 g, 0.5 mmol), 3,6-dibromo-9-tosylcarbazole (0.23 g, 0.6 mmol), copper iodide (0.005 g, 0.02 mmol), 18-crown-6-ether (0.025 g, 0.09 mmol) and potassium carbonate (0.09 g, 0.65 mmol), and the mixture was reacted at 220°C overnight. After completing the reaction, dichloromethane was added thereto, and the mixture was filtered with Celite. After concentrating, the organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 70/30), thereby providing the compound 4 (yield amount: 0.11 g, crude yield: 23.1%).

10 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): 9.10 (d, J = 8.5 Hz, 2H), 8.84 (d, J = 7.0 Hz, 4H), 8.41 (d, J = 2.0 Hz, 1H), 8.33 (d, J = 1.5 Hz, 1H), 8.20-8.15 (m, 4H), 7.92 (d, J = 8.5 Hz, 2H), 7.80 (d, J = 8.5 Hz, 1H), 7.69-7.55 (m, 12H), 7.48 (s, 2H), 7.43 (s, 4H), 7.35-7.29 (m, 3H)

15 MALDI-TOF-MS: m/z = 803 ([M-1]<sup>+</sup>)

## Synthesis Example 5

20 **[0114]** In this synthesis example, a compound 28 was synthesized according to the following schemes.

25

30

35

40

45

50


55

[chem 57]

5

10


15



20

25


30



35

40

45



Scheme 1

50

Synthesis of 9-Tosyl-3,3',6,6''-tetraphenyl-3',6'-dicarbazolylicarbazole

55

**[0115]** Dodecylbenzene (3.5 mL) was added to 3,6-dibromo-9-tosylcarbazole (1.92 g, 4.0 mmol), 3,6-diphenylcarbazole (2.58 g, 8.08 mmol) and copper oxide (1.39 g, 9.7 mmol), and the mixture was reacted at 220°C overnight. After completing the reaction, the product was extracted with dichloromethane. The organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 70/30) (yield amount: 1.42 g, yield: 19.0%).

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ (ppm) 8.66 (d, J = 9.0 Hz, 2H), 8.40 (d, J = 2.0 Hz, 4H), 8.18 (d, J = 2.0 Hz, 2H), 7.97 (d, J = 8.50 Hz, 2H), 7.82 (dd, J = 2.0 Hz, J = 2.0 Hz, 2H), 7.72 (d, J = 7.0 Hz, 8H), 7.67 (dd, J = 2.0 Hz, J = 2.0 Hz, 4H),

7.48-7.46 (m, 12H), 7.36-7.33 (m, 6H), 2.42 (s, 3H)

#### Scheme 2

#### 5 Synthesis of 3,3'',6,6''-Tetraphenyl-3',6'-dicarbazoly carbazole

10 **[0116]** THF (2.3 mL), DMSO (1.4 mL) and water (0.3 mL) were added to 9-tosyl-3,3'',6,6''-tetraphenyl-3',6'-dicarbazoly carbazole (0.25 g, 0.25 mmol) and potassium hydroxide (0.14 g, 2.5 mmol), and the mixture was refluxed under heating for 4 hours. After completing the reaction, the reaction mixture was neutralized with 1N HCl. The product was extracted with water and dichloromethane, and the organic layer was dried over sodium sulfate and concentrated with an evaporator. The product was recrystallized from hexane and dichloromethane (yield amount: 0.16 g, yield: 80.0%).  
15 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ (ppm) 8.52 (s, 1H), 8.42 (d, J = 2.0 Hz, 4H), 8.30 (d, J = 2.0 Hz, 2H), 7.77 (d, J = 8.5 Hz, 2H), 7.73 (d, J = 8.0 Hz, 8H), 7.70 (d, J = 2.0 Hz, 1H), 7.67 (dd, J = 2.0 Hz, J = 1.5 Hz, 5H), 7.49-7.46 (m, 12H), 7.35-7.33 (m, 4H)

#### Scheme 3

#### Synthesis of 9'-(4-(4,6-diphenyl-1,3,5-triazin-2-yl)phenyl)-9'H-(3',6'-diphenylcarbazolyl)carbazole

20 **[0117]** Dodecylbenzene (1.2 mL) was added to 3,3'',6,6''-tetraphenyl-3',6'-dicarbazoly carbazole (0.97 g, 1.2 mmol), diphenyltriazine (0.47 g, 1.21 mmol), copper iodide (0.004 g, 0.02 mmol), 18-crown-6-ether (0.035 g, 0.13 mmol) and potassium carbonate (0.17 g, 1.43 mmol), and the mixture was reacted at 220°C for 2 days. After completing the reaction, the product was extracted with dichloromethane. The organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 70/30), thereby providing the compound 28 (yield amount: 0.20 g, yield: 15.3%).  
25 <sup>1</sup>H-NMR (500 MHz, d<sub>6</sub>-DMSO) : δ (ppm) 9.16 (d, J = 9.0 Hz, 2H), 8.87-8.83 (m, 6H), 8.77 (d, J = 8.5 Hz, 4H), 8.23 (d, J = 8.5 Hz, 2H), 7.96 (d, J = 8.5 Hz, 2H), 7.86-7.80 (m, 16H), 7.76-7.71 (m, 4H), 7.55 (d, J = 8.5 Hz, 4H), 7.52-7.49 (m, 8H), 7.37-7.34 (m, 4H)

#### 30 Synthesis Example 6

**[0118]** In this synthesis example, a compound 29 was synthesized according to the following schemes.

35

40

45

50


55

[chem 58]

5

10


15



20


25

30



35

40



45

Scheme 1

Synthesis of 3,3',6,6''-Tetra-tert-butyl-9'-tosyl-3',6'-dicarbazolylicarbazole

50

55

**[0119]** Dodecylbenzene (4 mL) and NMP (1 mL) were added to 3,6-di-tert-butylcarbazole (2.69 g, 9.59 mmol), 3,6-dibromocarbazole (2.40 g, 5.00 mmol) and copper oxide (1.71 g, 12.3 mmol), and the mixture was reacted at 220°C overnight. After completing the reaction, copper oxide was removed by filtration, and the filtrate was concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: cyclohexane/dichloromethane = 50/50), thereby providing a matter that was considered to be the target product in an amount of approximately 0.44 g (yield: approximately 10%). Copper (II) oxide (0.85 g, 5.94 mmol) and dodecylbenzene (2 mL) were added to the product containing the mono-substituted compound and the raw materials, and the mixture was heated under refluxing at 220°C overnight (yield amount: 1.02 g, yield: 23.3%).

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ (ppm) 8.56 (d, J = 8.5 Hz, 2H), 8.14 (d, J = 1.5 Hz, 4H), 8.05 (d, J = 2.0 Hz, 2H), 7.91 (d,

J= 8.5 Hz, 2H), 7.73 (dd, J = 2.0 Hz, J = 2.0 Hz, 2H), 7.44 (dd, J = 2.0 Hz, J = 2.0 Hz, 4H), 7.33-7.2.8 (m, 6H), 2.39 (s, 3H), 1.45

(s, 36H)

5

Scheme 2

Synthesis of (3,6-Di(3,6-di-tert-butylcarbazolyl))carbazole

10 **[0120]** THF (49 mL), DMSO (24 mL) and water (5 mL) were added to 3,3",6,6"-tetra-tert-butyl-9'-tosyl-3',6'-dicarbazolylcarbazole (4.78 g, 5.45 mmol) and potassium hydroxide (2.41 g, 56 mmol), and the mixture was stirred under heating for 3 hours. After completing the reaction, the reaction mixture was neutralized with 1N HCl and extracted with dichloromethane. The organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: cyclohexane/dichloromethane = 70/30) and recrystallized from hexane (yield amount: 2.95 g, yield: 75%).

15 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ (ppm) 8.42 (s, 1H), 8.17-8.15 (m, 6H), 7.69-7.67 (m, 1H), 7.61 (d, J= 8.5 Hz, 2H), 7.65-7.55 (m, 1H), 7.47-7.43 (m, 4H), 7.32-7.30 (m, 4H), 1.48-1.46 (m, 36H)

Scheme 3

20

Synthesis of 9'-(4-(4,6-diphenyl-1,3,5-triazin-2-yl)phenyl)-9'H-(3',6'-di-tert-butylcarbazolyl)carbazole

25 **[0121]** Dodecylbenzene (0.6 mL) was added to (3,6-di(3,6-di-tert-butylcarbazolyl))carbazole (0.51 g, 0.7 mmol), 3,6-dibromo-9-tosylcarbazole (0.31 g, 0.81 mmol), copper iodide (0.007 g, 0.04 mmol), 18-crown-6-ether (0.02 g, 0.08 mmol) and potassium carbonate (0.09 g, 0.65 mmol), and the mixture was refluxed under heating overnight. After completing the reaction, the mixture was filtered with Celite. After concentrating, the organic layer was dried over magnesium sulfate and concentrated with an evaporator. The product was purified by silica gel column chromatography (developing solvent: hexane/dichloromethane = 50/50), thereby providing the compound 29 (yield amount: 0.16 g, crude yield: 22.5%).

MALDI-TOF-MS: m/z = 1030 ([M+1]<sup>+</sup>)

30

Example 1

35 **[0122]** In this example, toluene solutions of the compounds 1 to 4 and 27 to 29 (concentration: 10<sup>-5</sup> mol/L) were prepared and irradiated with light at 300 K under nitrogen bubbling, and thus light emission was observed. The absorption wavelength, the light emission wavelength and the quantum yield are shown in Table 1. The toluene solutions exhibited delayed fluorescent light.

Table 1

40

Compound	Absorption wavelength (nm)	Light emission wavelength (nm)	Quantum yield (%)
Compound 1	360	450	71.6
Compound 2	370	461	56.9
Compound 3	357	451	68.1
45 Compound 4	360	442	82.5
Compound 27	361	448	57.1
Compound 28	360	450	74.2
50 Compound 29	370	460	68.5

Example 2

**[0123]** In this example, organic photoluminescent devices were produced and evaluated for the characteristics.

55 **[0124]** The compound 1 and DPEPO were vapor-deposited on a silicon substrate from separate vapor deposition sources under condition of a vacuum degree of  $5.0 \times 10^{-4}$  Pa to form a thin film having a thickness of 100 nm having a concentration of the compound 1 of 6% by weight at a rate of 0.3 nm/sec, and thus an organic photoluminescent device was produced.

[0125] Organic photoluminescent devices were produced in the same manner except that the compounds 2 to 4 and 27 to 29 were used instead of the compound 1.

[0126] The organic photoluminescent devices thus produced were irradiated with light having a wavelength of 337 nm with a N<sub>2</sub> laser, and the light emission spectrum from the thin film on irradiation was evaluated for the characteristics at 300 K.

[0127] The light emission spectrum of the organic photoluminescent device using the compound 1 is shown in Fig. 2, the light emission spectrum of the organic photoluminescent device using the compound 2 is shown in Fig. 6, the light emission spectrum of the organic photoluminescent device using the compound 3 is shown in Fig. 10, and the light emission spectrum of the organic photoluminescent device using the compound 4 is shown in Fig. 14. The light emission spectrum of the organic photoluminescent device using the compound 27 is shown in Fig. 18, the light emission spectrum of the organic photoluminescent device using the compound 28 is shown in Fig. 22, and the light emission spectrum of the organic photoluminescent device using the compound 29 is shown in Fig. 26.

### Example 3

[0128] In this example, organic electroluminescent devices were produced and evaluated for the characteristics.

[0129] Thin films were laminated on a glass substrate having formed thereon an anode formed of indium tin oxide (ITO) having a thickness of 100 nm, by a vacuum vapor deposition method at a vacuum degree of  $5.0 \times 10^{-4}$  Pa. Firstly,  $\alpha$ -NPD was formed to a thickness of 35 nm on ITO, and thereon CBP was formed to a thickness of 10 nm. Further thereon, the compound 1 and DPEPO were vapor-deposited from separate vapor deposition sources to form a thin film having a thickness of 15 nm having a concentration of the compound 1 of 6% by weight. Further thereon, TPBi was formed to a thickness of 40 nm, further lithium fluoride (LiF) was vacuum vapor-deposited to a thickness of 0.8 nm, and then aluminum (Al) was vapor-deposited to a thickness of 80 nm to form a cathode, thereby producing an organic electroluminescent device.

[0130] Organic electroluminescent devices were produced in the same manner except that the compounds 2 to 4 and 27 to 29 were used instead of the compound 1.

[0131] The organic electroluminescent devices thus produced were measured with Semiconductor Parameter Analyzer (E5273A, produced by Agilent Technologies, Inc.), Optical Power Meter (1930C, produced by Newport Corporation), Fiber Optic Spectrometer (USB2000, produced by Ocean Optics, Inc.) and Streak Camera (Model C4334, produced by Hamamatsu Photonics K.K.).

[0132] The light emission spectrum of the organic electroluminescent device using the compound 1 is shown in Fig. 2, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 3, the voltage-electric current density characteristics thereof are shown in Fig. 4, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 5. The light emission spectrum of the organic electroluminescent device using the compound 2 is shown in Fig. 6, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 7, the voltage-electric current density characteristics thereof are shown in Fig. 8, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 9. The light emission spectrum of the organic electroluminescent device using the compound 3 is shown in Fig. 10, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 11, the voltage-electric current density characteristics thereof are shown in Fig. 12, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 13. The light emission spectrum of the organic electroluminescent device using the compound 4 is shown in Fig. 14, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 15, the voltage-electric current density characteristics thereof are shown in Fig. 16, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 17. The light emission spectrum of the organic electroluminescent device using the compound 27 is shown in Fig. 18, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 19, the voltage-electric current density characteristics thereof are shown in Fig. 20, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 21. The light emission spectrum of the organic electroluminescent device using the compound 28 is shown in Fig. 22, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 23, the voltage-electric current density characteristics thereof are shown in Fig. 24, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 25. The light emission spectrum of the organic electroluminescent device using the compound 29 is shown in Fig. 26, the transient decay curves under ordinary pressure and in vacuum thereof are shown in Fig. 27, the voltage-electric current density characteristics thereof are shown in Fig. 28, and the electric current density-external quantum efficiency characteristics thereof are shown in Fig. 29.


[0133] The transient decay curve shows the measurement result of the light emission lifetime obtained by measuring the process where the light emission intensity is deactivated on irradiating the compound with excitation light. In ordinary one-component light emission (fluorescent light or phosphorescent light), the light emission intensity decays monoexponentially. This means that the light emission intensity decays linearly on a graph with the semilogarithm as the ordinate.

In a transient decay curve of a delayed fluorescence emitter, while a linear component (fluorescent light) is observed in the initial stage of observation, a component that deviates from the linearity appears after several microseconds. The later component is light emission of the delayed component, and the signal thereof added to the initial component appears as a long tail curve on the longer time side. Thus, the measurement of the light emission lifetime revealed that the compound of the invention is a light emitting material that contained a delayed component in addition to a fluorescent component.

#### Comparative Example 1

**[0134]** In this comparative example, a toluene solution of the compound A shown below described in JP-A-2009-21336 and JP-A-2002-193952 (concentration:  $10^{-5}$  mol/L) was prepared and irradiated with light at 300 K under nitrogen bubbling, and thus the light emission spectrum shown in Fig. 30 was obtained. The solution exhibited the transient decay curve shown in Fig. 31, in which no delayed fluorescent component was observed.

[chem 59]



#### Industrial Applicability

**[0135]** The organic light emitting device of the invention is capable of achieving a high light emission efficiency. The compound represented by the general formula (1) is useful as a light emitting material of such an organic light emitting device. Accordingly, the invention has high industrial applicability.

#### Reference Signs List


**[0136]**

- 1 substrate  
 2 anode  
 3 hole injection layer  
 4 hole transporting layer  
 5 light emitting layer  
 6 electron transporting layer  
 7 cathode

## 10 Claims

1. A light emitting material containing a compound represented by the following general formula (1):

15 General Formula (1)



wherein in the general formula (1),

$R^1$  and  $R^2$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$R^{10}$  represents a carbazolyl group which may be substituted with a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$n_1$ ,  $n_2$ ,  $n_6$  and  $n_7$  each independently represent an integer of from 0 to 4;

$n_5$  represents an integer of from 0 to 3;


$n_8$  and  $n_9$  each independently represent an integer of from 0 to 5; and

$n_{10}$  represents 0 or 1,

provided that when  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$ ,  $R^2$  and  $R^5$  to  $R^9$  corresponding to  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  respectively each may be the same as or different from each other.

2. The light emitting material according to Claim 1, wherein the compound represented by the general formula (1) is a compound represented by the following general formula (2):

General Formula (2)



wherein in the general formula (2),

$R^1$  and  $R^2$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$n_1$ ,  $n_2$ ,  $n_6$  and  $n_7$  each independently represent an integer of from 0 to 4;


$n_5$  represents an integer of from 0 to 3; and

$n_8$  and  $n_9$  each independently represent an integer of from 0 to 5,

provided that when  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$ ,  $R^2$  and  $R^5$  to  $R^9$  corresponding to  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  respectively each may be the same as or different from each other.

3. The light emitting material according to Claim 1, wherein the compound represented by the general formula (1) is a compound represented by the following general formula (3):

General Formula (3)



wherein in the general formula (3),

30  $R^1$  to  $R^4$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$n_1$  to  $n_4$  and  $n_7$  each independently represent an integer of from 0 to 4;

35  $n_5$  and  $n_6$  each represent an integer of from 0 to 3; and  $n_8$  and  $n_9$  each independently represent an integer of from 0 to 5,


provided that when  $n_1$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$  to  $R^9$  corresponding to  $n_1$  to  $n_9$  respectively each may be the same as or different from each other.

- 40
- 45
- 50
- 55
4. The light emitting material according to any one of Claims 1 to 3, wherein the light emitting material emits delayed fluorescent light.
  5. The light emitting material according to any one of Claims 2 to 4, wherein in the general formula (2),  $n_6$  represents 0.
  6. The light emitting material according to any one of Claims 2, 4 and 5, wherein in the general formula (2),  $n_1$  represents an integer of from 1 to 4.
  7. The light emitting material according to Claim 6, wherein in the general formula (2),  $R^1$  is bonded to the 3-position of the carbazolyl group.
  8. The light emitting material according to Claim 6 or 7, wherein in the general formula (2),  $n_2$  represents an integer of from 1 to 4.
  9. The light emitting material according to Claim 8, wherein in the general formula (2),  $R^2$  is bonded to the 6-position of the carbazolyl group.
  10. The light emitting material according to any one of Claims 1, 2 and 4 to 9, wherein in the general formulae (1) and (2),  $R^1$  and  $R^2$  each independently represent a substituted or unsubstituted 9-carbazolyl group, a substituted or

unsubstituted phenyl group, a substituted or unsubstituted pyridyl group, an alkyl group having from 1 to 6 carbon atoms or a cycloalkyl group having from 5 to 7 carbon atoms.

- 5 11. The light emitting material according to any one of Claims 1, 2 and 4 to 9, wherein in the general formulae (1) and (2), R<sup>1</sup> and R<sup>2</sup> each independently represent a 9-carbazolyl group, a phenyl group, a tolyl group, a dimethylphenyl group, a trimethylphenyl group, a biphenyl group, a pyridyl group, a pyrrolyl group, a tert-butyl group or a cyclohexyl group.
- 10 12. The light emitting material according to any one of Claims 2, 4 and 5, wherein in the general formula (2), both n<sub>1</sub> and n<sub>2</sub> represent 0.
13. The light emitting material according to Claim 3 or 4, wherein in the general formula (3), at least one of n<sub>1</sub> to n<sub>4</sub> represents an integer of from 1 to 4.
- 15 14. The light emitting material according to Claim 3 or 4, wherein in the general formula (3), n<sub>1</sub> to n<sub>4</sub> each independently represent an integer of from 1 to 4.
16. The light emitting material according to any one of Claims 3, 4, 13 and 14, wherein in the general formula (3), R<sup>1</sup> to R<sup>4</sup> each independently represent a substituted or unsubstituted 9-carbazolyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted pyridyl group, an alkyl group having from 1 to 6 carbon atoms or a cycloalkyl group having from 5 to 7 carbon atoms.
- 20 15. The light emitting material according to any one of Claims 3, 4, 13 and 14, wherein in the general formula (3), R<sup>1</sup> to R<sup>4</sup> each independently represent a substituted or unsubstituted 9-carbazolyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted pyridyl group, an alkyl group having from 1 to 6 carbon atoms or a cycloalkyl group having from 5 to 7 carbon atoms.
- 25 16. The light emitting material according to any one of Claims 3, 4, 13 and 14, wherein in the general formula (3), R<sup>1</sup> to R<sup>4</sup> each independently represent a 9-carbazolyl group, a phenyl group, a tolyl group, a dimethylphenyl group, a trimethylphenyl group, a biphenyl group, a pyridyl group, a pyrrolyl group, a tert-butyl group or a cyclohexyl group.
17. The light emitting material according to any one of Claims 3, 4 and 13 to 16, wherein in the general formula (3), both n<sub>5</sub> and n<sub>6</sub> represent 0.
- 30 18. The light emitting material according to any one of Claims 1 to 17, wherein in the general formulae (1) to (3), all n<sub>7</sub>, n<sub>8</sub> and n<sub>9</sub> represent 0.
19. A delayed fluorescence emitter having a structure represented by the following general formula (1):

General Formula (1)



wherein in the general formula (1),

$R^1$  and  $R^2$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$R^{10}$  represents a carbazolyl group which may be substituted with a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$n_1$ ,  $n_2$ ,  $n_6$  and  $n_7$  each independently represent an integer of from 0 to 4;

$n_5$  represents an integer of from 0 to 3;


$n_8$  and  $n_9$  each independently represent an integer of from 0 to 5; and

$n_{10}$  represents 0 or 1,

provided that when  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$ ,  $R^2$  and  $R^5$  to  $R^9$  corresponding to  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  respectively each may be the same as or different from each other.

20. A compound represented by the following general formula (2):

General Formula (2)



wherein in the general formula (2),

$R^1$  and  $R^2$  each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

$R^5$  and  $R^6$  each independently represent a substituted or unsubstituted alkyl group;

$R^7$ ,  $R^8$  and  $R^9$  each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

$n_1$ ,  $n_2$ ,  $n_6$  and  $n_7$  each independently represent an integer of from 0 to 4;

$n_5$  represents an integer of from 0 to 3; and


$n_8$  and  $n_9$  each independently represent an integer of from 0 to 5,

provided that when  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  each represent an integer of 2 or more, plural groups represented by  $R^1$ ,  $R^2$  and  $R^5$  to  $R^9$  corresponding to  $n_1$ ,  $n_2$  and  $n_5$  to  $n_9$  respectively each may be the same as or different

from each other.

21. A compound represented by the following general formula (4):

General Formula (4)



wherein in the general formula (4),

R<sup>11</sup> to R<sup>14</sup> each independently represent a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted aryl group, a substituted or unsubstituted heteroaryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted cycloalkyl group;

R<sup>15</sup> and R<sup>16</sup> each independently represent an alkyl group;

R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> each independently represent a substituted or unsubstituted aryl group, a substituted or unsubstituted alkyl group or a substituted or unsubstituted carbazolyl group;

n<sup>11</sup> to n<sup>14</sup> and n<sup>17</sup> each independently represent an integer of from 0 to 4;

n<sup>15</sup> and n<sup>16</sup> each represent an integer of from 0 to 3; and

n<sup>18</sup> and n<sup>19</sup> each independently represent an integer of from 0 to 5,

provided that

at least one of n<sup>11</sup> to n<sup>14</sup> represents an integer of from 1 to 4, and

when n<sup>11</sup> to n<sup>19</sup> each represent an integer of 2 or more, plural groups represented by R<sup>11</sup> to R<sup>19</sup> corresponding to n<sup>11</sup> to n<sup>19</sup> respectively each may be the same as or different from each other.

22. An organic light emitting device containing a substrate having thereon a light emitting layer containing the light emitting material according to any one of Claims 1 to 18.

23. The organic light emitting device according to Claim 22, wherein the organic light emitting device emits delayed fluorescent light.

24. The organic light emitting device according to Claim 22 or 23, wherein the organic light emitting device is an organic electroluminescent device.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30





Fig. 31



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2014/051184

5	A. CLASSIFICATION OF SUBJECT MATTER C09K11/06(2006.01)i, C07D403/14(2006.01)i, H01L51/50(2006.01)i	
	According to International Patent Classification (IPC) or to both national classification and IPC	
10	B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C09K11/06, C07D403/14, H01L51/50	
15	Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Jitsuyo Shinan Koho 1922-1996 Jitsuyo Shinan Toroku Koho 1996-2014 Kokai Jitsuyo Shinan Koho 1971-2014 Toroku Jitsuyo Shinan Koho 1994-2014	
	Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAplus/REGISTRY(STN), Scopus, JSTPlus/JMEDPlus/JST7580(JDreamIII)	
20	C. DOCUMENTS CONSIDERED TO BE RELEVANT	
	Category*	Citation of document, with indication, where appropriate, of the relevant passages
25	X A	WO 2011/049325 A2 (CHEIL INDUSTRIES INC.), 28 April 2011 (28.04.2011), [105], chemical formula 25 & US 2012/0205636 A1 & KR 10-2011-0043342 A & TW 201114743 A
30	X A	WO 2012/077902 A2 (CHEIL INDUSTRIES INC.), 14 June 2012 (14.06.2012), [73], compound 28a & KR 10-2012-0013173 A
35	A	Serevicius T et al, "Enhanced electroluminescence based on thermally activated delayed fluorescence from a carbazole -triazine derivative" Physical Chemistry Chemical Physics, 15, 38, pp.15850-15855, 2013
40	<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.	
45	* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
50	Date of the actual completion of the international search 22 April, 2014 (22.04.14)	Date of mailing of the international search report 28 April, 2014 (28.04.14)
55	Name and mailing address of the ISA/ Japanese Patent Office	Authorized officer
	Facsimile No.	Telephone No.

Form PCT/ISA/210 (second sheet) (July 2009)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2014/051184

5

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

10

15

20

25

30

35

40

45

50

55

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Ayataka Endo et al, "Efficient up-conversion of triplet excitons into a singlet state and its application for organic light emitting diodes" Applied Physics Letters 98,083302(2011)	1-24
A	Hiroki Uoyama et al, "Highly efficient organic light-emitting diodes from delayed fluorescence", Nature, Vol. 492, pp.234-238 (2012)	1-24
A	Kenichi Goushi et al, "Organic light-emitting diodes employing efficient reverse intersystem crossing for triplet-to-singlet state conversion", Nature Photonics 6,253-258(2012)	1-24
A	Hironori KAJI, "Fabrication and Analysis of Organic LEDs and Solar Cells", IEICE Technical Report 2013, vol.113, no.18, pages 43 to 48	1-24

Form PCT/ISA/210 (continuation of second sheet) (July 2009)

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- JP 2009021336 A [0007] [0134]
- JP 2002193952 A [0007] [0134]
- WO 2012077902 A [0007]

**Non-patent literature cited in the description**

- *Appl. Phys. Lett.*, 2002, vol. 101, 093306 [0058]
- *Appl. Phys. Lett.*, 2012, vol. 101, 093306 [0063]

专利名称(译)	发光材料和使用其的有机发光元件		
公开(公告)号	<a href="#">EP2949724A4</a>	公开(公告)日	2016-06-29
申请号	EP2014742808	申请日	2014-01-22
[标]申请(专利权)人(译)	国立大学法人九州大学		
申请(专利权)人(译)	KYUSHU大学, 国立大学公司		
当前申请(专利权)人(译)	KYUSHU大学, 国立大学公司		
[标]发明人	HIRATA SHUZO SAKAI YUMI MASUI KENSUKE SHIZU KATSUYUKI TANAKA HIROYUKI ADACHI CHIHAYA		
发明人	HIRATA SHUZO SAKAI YUMI MASUI KENSUKE SHIZU KATSUYUKI TANAKA HIROYUKI ADACHI CHIHAYA		
IPC分类号	C09K11/06 C07D403/14 H01L51/50		
CPC分类号	C07D403/14 C09K2211/1029 C09K2211/1059 H01L51/0067 H01L51/0072 H01L51/5012 C09K11/06 C09K2211/1007		
优先权	2013010153 2013-01-23 JP 2013010154 2013-01-23 JP		
其他公开文献	EP2949724A1 EP2949724B1		
外部链接	<a href="#">Espacenet</a>		

#### 摘要(译)

在发光层中含有由以下通式表示的化合物的有机发光器件具有高发光效率。在通式(1)中, R1和R2代表咪唑基, 芳基, 杂芳基, 烷基或环烷基; R5和R6代表烷基; R7, R8和R9代表芳基, 烷基或咪唑基; R10代表咪唑基, 其可以被咪唑基, 芳基, 杂芳基, 烷基或环烷基取代; n1, n2, n6和n7表示0至4的整数; n5表示0至3的整数; n8和n9表示0至5的整数; n10代表0或1。