

(19)



(11)

EP 2 626 399 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
03.12.2014 Bulletin 2014/49

(51) Int Cl.:
C09K 11/06 ^(2006.01) **C07C 15/28** ^(2006.01)
H01L 51/00 ^(2006.01) **H05B 33/10** ^(2006.01)

(21) Application number: **13154326.6**

(22) Date of filing: **07.02.2013**

(54) Amine-based compound and organic light-emitting diode including the same

Aminbasierte Verbindung und organische lichtemittierende Diodenvorrichtung damit

Composé à base d'amine et diode électroluminescente organique le comprenant

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **07.02.2012 KR 20120012532**
08.06.2012 KR 20120061676

(43) Date of publication of application:
14.08.2013 Bulletin 2013/33

(73) Proprietor: **Samsung Display Co., Ltd.**
Yongin-City, Gyeonggi-Do, 446-711 (KR)

(72) Inventors:

- **Kim, Young-Kook**
Gyeonggi-Do 446-711 (KR)
- **Hwang, Seok-Hwan**
Gyeonggi-Do 446-711 (KR)
- **Jung, Hye-Jin**
Gyeonggi-Do 446-711 (KR)
- **Lim, Jin-O**
Gyeonggi-Do 446-711 (KR)
- **Han, Sang-Hyun**
Gyeonggi-Do 446-711 (KR)
- **Jeong, Eun-Jae**
Gyeonggi-Do 446-711 (KR)

- **Kim, Soo-Yon**
Gyeonggi-Do 446-711 (KR)
- **Park, Jun-Ha**
Gyeonggi-Do 446-711 (KR)
- **Lee, Eun-Young**
Gyeonggi-Do 446-711 (KR)
- **Lee, Jong-Hyuk**
Gyeonggi-Do 446-711 (KR)

(74) Representative: **Russell, Tim et al**
Venner Shipley LLP
200 Aldersgate
London EC1A 4HD (GB)

(56) References cited:
US-A- 5 935 721

- **KUM HEE LEE ET AL: "Synthesis and Electroluminescent Properties of Blue Fluorescent Triphenylamine Substituted Anthracene Derivatives for OLEDs", MOLECULAR CRYSTALS AND LIQUID CRYSTALS, vol. 530, no. 1, 11 October 2010 (2010-10-11), pages 48/[204]-55/[211], XP55060418, ISSN: 1542-1406, DOI: 10.1080/15421406.2010.495882**

EP 2 626 399 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to an amine-based compound for organic light-emitting diodes, and an organic light-emitting diode including the compound.

[0002] Organic light-emitting diodes (OLEDs), which are self-emitting diodes, have advantages such as wide viewing angles, excellent contrast, quick response, high brightness, excellent driving voltage characteristics, and can provide multicoloured images.

[0003] A typical OLED has a structure including a substrate, and an anode, a hole transport layer (HTL), an emission layer (EML), an electron transport layer (ETL), and a cathode which are sequentially stacked on the substrate. In this regard, the HTL, the EML, and the ETL are organic thin-films comprising organic compounds.

[0004] An operating principle of an OLED having the above-described structure is as follows.


[0005] When a voltage is applied between the anode and the cathode, holes injected from the anode move to the EML via the HTL, and electrons injected from the cathode move to the EML via the ETL. The holes and electrons recombine in the EML to generate excitons. When the excitons drop from an excited state to a ground state, light is emitted.

[0006] Kum Hee Lee et.al. disclose blue fluorescent triphenylamine substituted anthracenes derivatives, especially 9-(naphthalen-2-yl)-10-(4-triphenylamine)anthracene, comprised in the organic layer of organic light-emitting devices ("Synthesis and electroluminescent properties of blue fluorescent triphenylamine substituted anthracene derivatives for OLEDs", Molecular Crystals and Liquid Crystals, Vol. 530, No.1, 11 October 2010)

[0007] The present invention provides an amine-based compound having a novel structure and an organic light-emitting diode including the amine-based compound.

[0008] According to a first aspect of the present invention there is provided an amine-based compound represented by Formula 1 below:

Formula 1



wherein, in Formula 1, Ar_1 and Ar_2 are each independently a substituted or unsubstituted C_6-C_{60} aryl group or a substituted or unsubstituted C_2-C_{60} heteroaryl group;

X_1 is a substituted or unsubstituted C_6-C_{60} arylene group or a substituted or unsubstituted C_2-C_{60} heteroarylene group;

m is an integer from 1 to 5; and

at least one substituent of each of the substituted C_6-C_{60} aryl group, the substituted C_2-C_{60} heteroaryl group, the substituted C_6-C_{60} arylene group, and the substituted C_2-C_{60} heteroarylene group is one of a deuterium atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; $-NO_2$; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a tri(C_6-C_{60} aryl)silyl group; a C_1-C_{60} alkyl group, a C_1-C_{60} alkoxy group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group; a C_1-C_{60} alkyl group, a C_1-C_{60} alkoxy group, a C_2-C_{60} alkenyl group and a C_2-C_{60} alkynyl group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, $-NO_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof; a C_3-C_{60} cycloalkyl group, a C_3-C_{60} cycloalkenyl group, a C_6-C_{60} aryl group, a C_2-C_{60} heteroaryl group, a C_6-C_{60} aralkyl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthiol group; a C_3-C_{60} cycloalkyl group, a C_3-C_{60} cycloalkenyl group, a C_6-C_{60} aryl group, a C_2-C_{60} heteroaryl group, a C_6-C_{60} aralkyl group, a C_6-C_{60} aryloxy group, and a C_6-C_{60} arylthiol group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, $-NO_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic

acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one fluorine (F), a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group and a C₂-C₆₀ heteroaryl group,

wherein at least one of Ar₁ and Ar₂ is a C₆-C₆₀ aryl group substituted with at least one electron withdrawing group selected from the group consisting of -F; -CN; -NO₂; a C₁-C₆₀ alkyl group substituted with at least one -F; a C₂-C₆₀ heteroaryl group; and a C₂-C₆₀ heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.

[0009] According to another aspect of the present invention, there is provided an organic light-emitting diode comprising a first electrode, a second electrode disposed opposite to the first electrode, and an organic layer disposed between the first electrode and the second electrode, the organic layer comprising at least one of the amine-based compounds according to the invention in its first aspect.


[0010] A more complete appreciation of the present invention, and many of the attendant advantages thereof, will be readily apparent as the same becomes better understood by reference to the following detailed description when considered in conjunction with the accompanying drawings, wherein:

Figure. 1 schematically illustrates the structure of an organic light-emitting diode according to an embodiment of the invention.

[0011] As used herein, the term "and/or" includes any and all combinations of one or more of the associated listed items.

[0012] According to an aspect of the present invention, there is provided an amine-based compound represented by Formula 1:

Formula 1



[0013] In Formula 1 above, Ar₁ and Ar₂ are each independently a substituted or unsubstituted C₆-C₆₀ aryl group, or a substituted or unsubstituted C₂-C₆₀ heteroaryl group; and X₁ is a substituted or unsubstituted C₆-C₆₀ arylene group, or a substituted or unsubstituted C₂-C₆₀ heteroarylene group and m is an integer selected from 1, 2, 3, 4 and 5.

[0014] At least one substituent of each of the substituted C₆-C₆₀ aryl group, the substituted C₂-C₆₀ heteroaryl group, the substituted C₆-C₆₀ arylene group, and the substituted C₂-C₆₀ heteroarylene group is one of a deuterium atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; -NO₂; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a tri(C₆-C₆₀aryl)silyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, and a C₂-C₆₀ alkynyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group and a C₂-C₆₀ alkynyl group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof; a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthiol group; a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, and a C₆-C₆₀ arylthiol group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one fluorine (F), a

C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group and a C₂-C₆₀ heteroaryl group.


[0015] In Formula 1 above, at least one of Ar₁ and Ar₂ is a C₆-C₆₀ aryl group substituted with at least one electron withdrawing group selected from the group consisting of -F; -CN; -NO₂; a C₁-C₆₀ alkyl group substituted with at least one -F; a C₂-C₆₀ heteroaryl group; and a C₂-C₆₀ heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.

[0016] For example, the at least one electron withdrawing group may be selected from the group consisting of: -F; -CN; -NO₂; a C₁-C₂₀ alkyl group substituted with at least one -F; a C₂-C₂₀ heteroaryl group including a ring which contains a N atom; and a C₂-C₂₀ heteroaryl group that includes a ring which contains a N atom and is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₂-C₂₀ heteroaryl group.


[0017] In an embodiment of the present invention, the at least one electron withdrawing group in Formula 1 is selected from the group consisting of -F; -CN; a C₁-C₂₀ alkyl group substituted with at least one -F; a pyrrolyl group, a pyrazolyl group, an imidazolyl group, an imidazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a benzoimidazolyl group, an indolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a phthalazinyl group, an indolizynyl group, a quinazolynyl group, a cinnolinyl group, an indazolyl group, a carbazolyl group, a phenazinyl group, a phenanthridinyl group, a triazinyl group, a pyridazinyl group, a triazolyl group, and a tetrazolyl; and a pyrrolyl group, a pyrazolyl group, an imidazolyl group, an imidazolyl group, an imidazopyridinyl group, imidazopyrimidinyl, pyridinyl, pyrazinyl, pyrimidinyl, benzoimidazolyl, indolyl, purinyl, quinolinyl, isoquinolinyl, phthalazinyl, indolizynyl, quinazolynyl, cinnolinyl, indazolyl, carbazolyl, phenazinyl, phenanthridinyl, triazinyl, pyridazinyl, triazolyl, and a tetrazolyl group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group and a carbazolyl group.

[0018] For example, the at least one electron withdrawing group in Formula 1 above may be selected from, but is not limited to, the group consisting of: -F; -CN; a C₁-C₂₀ alkyl group substituted with at least one -F; a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolynyl group, a triazinyl group, a benzoimidazolyl group; a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolynyl group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group.


[0019] In an embodiment of the invention, the at least one electron withdrawing group in Formula 1 above is selected from the group consisting of -F; -CN; -CH₂F; -CHF₂; -CF₃; and groups represented by Formulae 2(1) to 2(14) below:




Formula 2(1)




Formula 2(2)




Formula 2(3)




Formula 2(4)




Formula 2(5)




Formula 2(6)




Formula 2(7)




Formula 2(8)




Formula 2(9)




Formula 2(10)




Formula 2(11)



Formula 2(12)



Formula 2(13)



Formula 2(14)

[0020] In Formulae 2(1) to 2(14) above, Z₁₁ to Z₁₈ may each independently be a hydrogen atom, a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, or a carbazolyl group.


[0021] For example, Z₁₁ to Z₁₈ in Formulae 2(1) to 2(14) above may each independently be a hydrogen atom, a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonate group or a salt thereof, a phosphoric acid or a salt thereof, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, or a carbazolyl group, but are not limited thereto.

[0022] In Formula 1 above, the at least one of Ar₁ and Ar₂ may be a C₆-C₆₀ aryl group substituted with at least two electron withdrawing groups.


[0023] In an embodiment of the present invention, the at least one of Ar₁ and Ar₂ is a phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl group that is substituted with at least two electron withdrawing groups. The electron withdrawing groups may each independently be selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group.

[0024] At least one of Ar₁ and Ar₂ in the amine-based compound of Formula 1 above is a C₆-C₆₀ aryl group substituted with at least one of the above electron withdrawing groups. Accordingly, the amine-based compound may be represented by Formula 1(1) or Formula 1(2) below:

Formula 1(1)



Formula 1(2)



[0025] In Formula 1(1) above, Ar_2 is a substituted or unsubstituted C_6 - C_{20} aryl group, or a substituted or unsubstituted C_2 - C_{20} heteroaryl group. In Formulae 1(1) and 1(2) above, the rings A and B are each independently a substituted C_6 - C_{20} aryl group; R_1 and R_2 are each independently an electron withdrawing group selected from the group consisting of: -F; -CN; -NO₂; a C_1 - C_{60} alkyl group substituted with at least one -F; a C_2 - C_{60} heteroaryl group; and a C_2 - C_{60} heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_1 - C_{60} alkoxy group substituted with at least one -F, a C_1 - C_{60} alkoxy group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_6 - C_{60} aryl group, and a C_2 - C_{60} heteroaryl group; and p and q are each independently an integer selected from 1, 2, 3, 4, 5, 6, 7, 8 and 9.

[0026] The electron withdrawing group in Formulae 1(1) and 1(2) are as described above, and thus a detailed description thereof will not be repeated herein.

[0027] For example, the amine-based compound may be represented by Formula 1(1) above, wherein at least one of R_1 in Formula 1(1) may be -CN.

[0028] In an embodiment of the present invention, the amine-based compound is represented by Formula 1(2) above, wherein at least one of R_1 and R_2 in Formula 1(2) is -CN.

[0029] In an embodiment of the present invention, the amine-based compound is represented by Formula 1(1) above, wherein the ring A in Formula 1(1) is a substituted phenyl group, a substituted biphenyl group, a substituted naphthyl group, a substituted anthryl group, a substituted phenanthrenyl group, a substituted pyrenyl group, or a substituted fluorenyl group, but is not limited thereto.

[0030] In another embodiment of the present invention, the amine-based compound is represented by Formula 1(2) above, wherein the rings A and B in Formula 1(2) may be each independently a substituted phenyl group, a substituted biphenyl group, a substituted naphthyl group, a substituted anthryl group, a substituted phenanthrenyl group, a substituted pyrenyl group, or a substituted fluorenyl group.

[0031] The amine-based compound may be represented by Formula 1(1), wherein the A ring may be a substituted phenyl group, a substituted biphenyl group, a substituted naphthyl group, a substituted anthryl group, a substituted phenanthrenyl group, a substituted pyrenyl group, or a substituted fluorenyl group; R_1 may be at least one electron withdrawing group selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkyl group substituted with at least one -F, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group; and p may be 2, 3, or 4, for example, may be 2.

[0032] In an embodiment of the present invention, the amine-based compound is represented by Formula 1(2), wherein the ring A and the ring B are each independently a substituted phenyl group, a substituted biphenyl group, a substituted naphthyl group, a substituted anthryl group, a substituted phenanthrenyl group, a substituted pyrenyl group, or a substituted fluorenyl group; R_1 and R_2 are each independently at least one electron withdrawing group selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkyl group substituted with at least one -F, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group; and p and q are each

independently an integer of 2, 3, or 4, for example, p and q may both be an integer of 2.

[0033] At least one of Ar₁ and Ar₂ in Formula 1 above may be a phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl group that are substituted with at least one of the above-listed electron withdrawing groups.


[0034] In Formula 1 above, Ar₁ and Ar₂ may each independently be a substituted or unsubstituted phenyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted azulenyl group, a substituted or unsubstituted heptalenyl group, a substituted or unsubstituted indacenyl group, a substituted or unsubstituted acenaphthyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted phenalenyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluoranthenyl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted naphthacenyl group, a substituted or unsubstituted picenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted pentaphenyl group, a substituted or unsubstituted hexacenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted imidazolynyl group, a substituted or unsubstituted imidazopyridinyl group, a substituted or unsubstituted imidazopyrimidinyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted benzoimidazolyl group, a substituted or unsubstituted indolyl group, a substituted or unsubstituted purinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted phthalazinyl group, a substituted or unsubstituted indolizynyl group, a substituted or unsubstituted naphthyridinyl group, a substituted or unsubstituted quina-
zolinyl group, a substituted or unsubstituted cinnolinyl group, a substituted or unsubstituted indazolyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted phenazinylene group, a substituted or unsubstituted phenanthridinyl group, a substituted or unsubstituted pyranyl group, a substituted or unsubstituted chromenyl group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranyl group, a substituted or unsubstituted thiophenyl group, a substituted or unsubstituted benzothiophenyl group, a substituted or unsubstituted isothiazolyl group, a substituted or unsubstituted benzoimidazolyl group, a substituted or unsubstituted isoxazolyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted oxadiazolyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted triazolyl group, a substituted or unsubstituted tetrazolyl group, or a substituted or unsubstituted phenanthrolinyl group. At least one of Ar₁ and Ar₂ may be a phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl group that are substituted with at least one of the above-listed electron withdrawing groups.

[0035] For example, Ar₁ and Ar₂ in Formula 1 above may each independently be a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, or a substituted or unsubstituted phenanthrolinyl group. At least one of Ar₁ and Ar₂ may be a phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl group that is substituted with at least one electron withdrawing group selected from the group consisting of -F; -CN; a C₁-C₂₀ alkyl group substituted with at least one -F; a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazolinyl group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group.


[0036] In Formula 1, Ar₁ and Ar₂ may be linked together by a single bond.

[0037] The amine-based compound may be represented by any one of Formulae 1A to 1J below:


Formula 1A




Formula 1B




Formula 1C




Formula 1D




Formula 1E




Formula 1F




Formula 1G




Formula 1H



Formula 1I



Formula 1J



15 Ar_2 in Formulae 1A to 1I may be the same as described above in conjunction with other formulae.

16 **[0038]** At least one of R_{11} to R_{15} in Formulae 1A and 1B, at least one of R_{11} to R_{17} in Formulae 1C and 1D, at least
 17 one of R_{11} to R_{18} in Formulae 1E and 1J, and at least one of R_{11} to R_{19} in Formulae 1F, 1G, 1H and 1I may each
 18 independently be an electron withdrawing group selected from the group consisting of -F; -CN; -NO₂; a C₁-C₆₀ alkyl
 19 group substituted with at least one -F; a C₂-C₆₀ heteroaryl group; and a C₂-C₆₀ heteroaryl group substituted with at least
 20 one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine,
 21 hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof,
 22 a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl
 23 group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.

24 **[0039]** For example, at least one of R_{11} to R_{15} of Formulae 1A and 1B, at least one of R_{11} to R_{17} in Formulae 1C and
 25 1D, at least one of R_{11} to R_{18} of Formulae 1E and 1J, and at least one of R_{11} to R_{19} of Formulae 1F, 1G, 1H and 1I may
 26 each independently be, but are not limited to, an electron withdrawing group selected from the group consisting of: -F;
 27 -CN; a C₁-C₂₀ alkyl group substituted with at least one -F; a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a
 28 quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl
 29 group; a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny
 30 group, a phthalazinyl group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with
 31 at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group,
 32 hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a
 33 salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl
 34 group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group,
 35 and a carbazolyl group.


36 **[0040]** In an embodiment of the present invention, Ar_2 in Formulae 1A to 1I may be, but are not limited thereto, a
 37 phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl
 38 group that are substituted with at least one electron withdrawing group selected from the group consisting of -F; -CN; a
 39 C₁-C₂₀ alkyl group substituted with at least one -F; a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl
 40 group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group;
 41 and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny
 42 group, a phthalazinyl group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with
 43 at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group,
 44 hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a
 45 salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl
 46 group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group,
 47 and a carbazolyl group.

48 **[0041]** In an embodiment of the invention, the amine-based compound may be represented by Formula 1A-(1) or
 49 1A-(2) below:


50

55

Formula 1A-(1)



Formula 1A-(2)



[0042] In Formulae 1A-(1) and 1A-(2), R_{12} , R_{14} , R_{22} , and R_{24} may each independently be an electron withdrawing group selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group; and Ar_2 may be a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, or a substituted or unsubstituted fluorenyl group.

[0043] In Formulae 1A-(1) and 1A-(2), R_{12} , R_{14} , R_{22} and R_{24} may each independently be selected from the group consisting of -F; -CN; -CH₂F; -CHF₂; -CF₃; and groups represented by Formulae 2(1) to 2(14) above.

[0044] In Formulae 1A-(1) and 1A-(2) above, R_{12} , R_{14} , R_{22} and R_{24} may each independently be one of the groups represented by Formulae 2(1) to 2(8) above.

[0045] In Formulae 1A-(1) and 1A-(2) above, R_{12} , R_{14} , R_{22} and R_{24} may each independently be the group represented by Formula 2(2) above, but are not limited thereto.


[0046] In Formula 1A-(1), Ar_2 may be a phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl group, but are not limited thereto.

[0047] In Formula 1, X_1 may be a substituted or unsubstituted phenylene group, a substituted or unsubstituted pentalenylene group, a substituted or unsubstituted indenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted azulenylenylene group, a substituted or unsubstituted heptalenylene group, a substituted or unsubstituted indacenylene group, a substituted or unsubstituted acenaphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted phenalenylenylene group, a substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted fluoranthenylenylene group, a substituted or unsubstituted triphenylenylene group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted naphthacenylenylene group, a substituted or unsubstituted pincenylenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted pentaphenylenylene group, a substituted or unsubstituted hexacenylenylene group, a substituted or unsubstituted pyrrolylene group, a substituted or unsubstituted pyrazolylenylene group, a substituted or unsubstituted imidazolylene group, a substituted or unsubstituted imidazolinylenylene group, a substituted or unsubstituted imidazopyridinylenylene group, a substituted or unsubstituted imidazopyrimidinylenylene group, a substituted or unsubstituted pyridinylenylene group, a substituted or unsubstituted pyrazinylenylene group, a substituted or unsubstituted pyrimidinylenylene group, a substituted or unsubstituted indolylenylene group, a substituted or unsubstituted purinylenylene group, a substituted or unsubstituted quinolinylenylene group, a substituted or unsubstituted phthalazinylenylene group, a substituted or unsubstituted indolizinylenylene group, a substituted or unsubstituted naphthyridinylenylene group, a substituted or unsubstituted quinazolinylenylene group, a substituted or unsubstituted cinnolinylenylene group, a substituted or unsubstituted indazolylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted phenazinylenylene group, a substituted or unsubstituted phenanthridinylenylene group, a substituted or unsubstituted pyranylene group, a substituted or unsubstituted chromenylenylene group, a substituted or unsubstituted furanylenylene group, a substituted or unsubstituted benzofuranylenylene group, a substituted or unsubstituted thiophenylenylene group, a substituted or unsubstituted benzothiophenylenylene group, a substituted or unsubstituted isothiazolylenylene group, a substituted or unsubstituted benzoimidazolylene group, a substituted or unsubstituted isoxazolylene group, a substituted or unsubstituted dibenzothiophenylenylene group, a substituted or unsubstituted dibenzofuranylenylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted oxadiazolylenylene group, a substituted or unsubstituted


pyridazinylene group, a substituted or unsubstituted triazolylene group, or a substituted or unsubstituted tetrazolylene group. X₁ in Formula 1 may have at least one substituent, which may be selected from among the substituents described above.

[0048] In an embodiment, X₁ in Formula 1 is, but is not limited to, a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted triphenylenylene group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted pyridinylene group, a substituted or unsubstituted pyrazinylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted quinolinylene group, a substituted or unsubstituted quinazolinylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted dibenzothioophenylylene group, a substituted or unsubstituted dibenzofuranylylene group, a substituted or unsubstituted triazinylylene group, a substituted or unsubstituted pyridazinylylene group, a substituted or unsubstituted triazolylene group, or a substituted or unsubstituted tetrazolylylene group.


[0049] For example, X₁ in Formula 1 above may be a group represented by one of Formulae 5(1) to 5(16) below:




Formula 5(1)




Formula 5(2)




Formula 5(3)




Formula 5(4)




Formula 5(5)




Formula 5(6)




Formula 5(7)




Formula 5(8)




Formula 5(9)




Formula 5(10)




Formula 5(11)




Formula 5(12)




Formula 5(13)



Formula 5(14)



Formula 5(15)



Formula 5(16)


[0050] In Formulae 5(1) to 5(16), Z_1 to Z_8 may each independently be one of a hydrogen atom; a deuterium atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; -NO₂; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; phosphoric acid or a salt thereof; a C₁-C₂₀ alkyl group; a C₁-C₂₀ alkoxy group; a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy groups that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, sulfonic acid group or a salt thereof, and phosphoric acid or a salt thereof; a C₆-C₂₀ aryl group; a C₂-C₂₀ heteroaryl group; and a C₆-C₂₀ aryl group and a C₂-C₂₀ heteroaryl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a C₆-C₂₀ aryl group and a C₂-C₂₀ heteroaryl group.


[0051] In Formulae 5(1) to 5(16), * indicates a binding site to anthracene in Formula 1, and *, indicates a binding site to N in Formula 1.

[0052] For example, Z_1 to Z_8 in Formulae 5(1) to 5(16) may each independently be, but are not limited to, a hydrogen atom; a deuterium atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; -NO₂; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a methyl group, an ethyl group, a propyl group, a butyl group, and a pentyl group; a methoxy group, an ethoxy group, a propoxy group, a butoxy group, and a pentoxy group; a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, and a pentoxy group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid or a salt thereof; a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, and a fluorenyl group; a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolyl group, and a carbazolyl group; a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a fluorenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a quinolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group and a pyridinyl group.


[0053] In Formula 1, m is an integer from 1 to 5. In an embodiment of the present invention, m is 1, 2, or 3, but is not limited thereto.

[0054] The amine-based compound of Formula 1 may be, for example, any one of Compounds 1 to 109 below, but is not limited thereto:







16




17




18




19




20




21




22




23




24




25




26




27



28




29




30

55


5



31




32




33


10



34



35




36


15

20


25



37




38




39

30


35



40




41




42

40


45



43



44




45

50


55

5




46

10




47




48

15




49

20




50




51

25




52

30




53




54

35




55

40




56




57

45



58


50



59


55

5




60

10




61

15




62

20




63

25




64

30




65

35




66

40




67

45



68











50




69

55


5
10
15
20
25
30
35
40
45
50
55



5




80




81

10

15




82




83

20

25




84




85

30

35




86




87

40

45



88




89


50

55


5
10
15
20
25
30
35
40
45
50
55




90




91




92




93




94




95




96




97



98




99



[0055] Since at least one of Ar_1 and Ar_2 in the amine-based compound of Formula 1 above is a C_6-C_{60} aryl group substituted with at least one electron withdrawing group selected from the group consisting of -F; -CN; $-NO_2$; a C_1-C_{60} alkyl group substituted with at least one -F; a C_2-C_{60} heteroaryl group; and a C_2-C_{60} heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, $-NO_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1-C_{60} alkyl group, a C_1-C_{60} alkoxy group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_6-C_{60} aryl group, and a C_2-C_{60} heteroaryl group, a moiety represented by "A" in Formula 1' below may be able to withdraw electrons.

Formula 1'



[0056] The amine-based compound of Formula 1 has a naphthyl-anthracene core abundant in electrons and a moiety A with electron withdrawing ability, and thus the amine-based compound may have good electron transport characteristics. When the electron withdrawing group is a substituted or unsubstituted C₂-C₆₀ heteroaryl group, the substituted or unsubstituted C₂-C₆₀ heteroaryl group is linked to N of Formula 1, not directly, but via a C₆-C₆₀ aryl group. Therefore, an organic light-emitting diode including the amine-based compound of Formula 1 may have improved efficiency characteristics. When the electron withdrawing group is -CN, an organic light-emitting diode including the amine-based compound of Formula 1 may have improved lifetime characteristics.

[0057] Not wishing to be bound by a particular theory, in either of i) an amine-based compound including a naphthyl-anthracene core that lacks the above-described electron withdrawing group or ii) an amine-based compound including a naphthyl-anthracene core with pyridine directly linked to N, the highest occupied molecular orbital (HOMO) electron density may be focused near the anthracene moiety. However, in the amine-based compound of Formula 1 above, the HOMO electron density may be dispersed near the amine moiety, so that the lowest unoccupied molecular orbital (LUMO) electron density may be relatively fixed near anthracene. This may result in the amine-based compound of Formula 1 having improved dipole characteristics. Thus, the electron transport characteristics of the amine-based compound of Formula 1 may be improved.

[0058] Therefore, an organic light-emitting diode including any of the amine-based compounds represented by Formula 1 above may have a low driving voltage, a high luminance, a high efficiency, and a long lifetime.

[0059] The amine-based compound of Formula 1 may be synthesized by any known organic synthesis method. A synthesis method of the amine-based compound of Formula 1 may be understood by those of ordinary skill in the art from the examples that will be described below.

[0060] At least one of the amine-based compounds according to the invention in its first aspect may be used between a pair of electrodes of an organic light-emitting diode. For example, at least one of the amine-based compounds may be in an EML and/or between a cathode and the EML (for example, in the ETL, the EIL, or a functional layer having both electron transport and electron injection capabilities).

[0061] According to another aspect of the present invention, an organic light-emitting diode includes a first electrode, a second electrode disposed opposite to the first electrode, and an organic layer disposed between the first electrode and the second electrode, wherein the organic layer includes at least one of the amine-based compounds of Formula 1 according to the invention in its first aspect.

[0062] As used herein, "(for example, the organic layer) including at least one amine-based compound" means (the organic layer) including one of the amine-based compounds of Formula 1 according to the invention in its first aspect, or at least two different amine-based compounds of Formula 1 according to the invention in its first aspect.

[0063] The organic layer may include only Compound 1 as the amine-based compound. Compound 1 may be in the EML or ETL of the organic light-emitting diode. The organic layer may include Compounds 1 and 3 as the amine-based compound. Compounds 1 and 3 may be in the same layer (for example, in the ETL) or may be in different layers (for example, in the EML and ETL, respectively).

[0064] The organic layer may include at least one layer selected from among a hole injection layer (HIL), a hole transport layer (HTL), a functional layer having both hole injection and hole transport capabilities (hereinafter, "H-functional layer"), a buffer layer, an electron blocking layer (EBL), an emission layer (EML), a hole blocking layer (HBL), an electron transport layer (ETL), an electron injection layer (EIL), and a functional layer having both electron injection and electron transport capabilities (hereinafter, "E-functional layer").

[0065] The term "organic layer" as used herein refers to a single layer and/or a plurality of layers disposed between the first and second electrodes of the organic light-emitting diode.

[0066] The organic layer may include an EML, wherein at least one of the amine-based compounds may be included in the EML.

[0067] The amine-based compound in the EML may serve as a host. When the amine-based compound in the EML

serves as a host, the EML may further include a fluorescent dopant. The fluorescent dopant may be a blue fluorescent dopant. The amine-based compound in the EML may serve as a dopant. When the amine-based compound in the EML serves as a dopant, the amine-based compound may be a blue fluorescent dopant.

[0068] The organic layer may include an ETL, wherein at least one of the amine-based compounds may be included in the ETL.

[0069] Figure 1 is a schematic sectional view of an organic light-emitting diode 10 according to an embodiment of the invention. Hereinafter, a structure of an organic light-emitting diode according to an embodiment of the invention and a method of manufacturing the same will now be described with reference to Figure 1.

[0070] The substrate 11 may be any substrate that is used in existing organic light-emitting diodes. The substrate 11 may be a glass substrate or a transparent plastic substrate with strong mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0071] The first electrode 13 may be formed by depositing or sputtering a first electrode-forming material on the substrate 11. When the first electrode 13 constitutes an anode, a material having a high work function may be used as the first electrode-forming material to facilitate hole injection. The first electrode 13 may be a reflective electrode or a transmission electrode. Suitable first electrode-forming materials include transparent and conductive materials such as ITO, IZO, SnO₂, and ZnO. The first electrode 13 may be formed as a reflective electrode using magnesium (Mg), aluminium (Al), aluminium-lithium (Al-Li), calcium (Ca), magnesium-indium (Mg-In), magnesium-silver (Mg-Ag), or the like.

[0072] The first electrode 13 may have a single-layer structure or a multi-layer structure including at least two layers. For example, the first electrode 13 may have a three-layered structure of ITO/Ag/ITO, but is not limited thereto.

[0073] The organic layer 15 may be disposed on the first electrode 13.


[0074] The organic layer 15 may include a HIL, a HTL, a buffer layer, an EML, an ETL, and an EIL.

[0075] The HIL may be formed on the first electrode 13 by vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like.


[0076] When the HIL is formed using vacuum deposition, vacuum deposition conditions may vary according to the compound that is used to form the HIL, and the desired structure and thermal properties of the HIL to be formed. For example, vacuum deposition may be performed at a temperature of about 100°C to about 500°C, a pressure of about 0.133 x 10⁻⁶ Pa (10⁻⁸ torr) to about 0.133 Pa (10⁻³ torr), and a deposition rate of about 0.001 to about 10 nm/sec (about 0.01 to about 100 Å/sec). However, the deposition conditions are not limited thereto.

[0077] When the HIL is formed using spin coating, the coating conditions may vary according to the compound that is used to form the HIL, and the desired structure and thermal properties of the HIL to be formed. For example, the coating rate may be in the range of about 2000 rpm to about 5000 rpm, and a temperature at which heat treatment is performed to remove a solvent after coating may be in the range of about 80°C to about 200°C. However, the coating conditions are not limited thereto.

[0078] The HIL 130 may comprise any material that is commonly used to form a HIL. Examples of the material that can be used to form the HIL are, but are not limited to, N,N'-diphenyl-N,N'-bis-[4-(phenyl-m-tolyl-amino)-phenyl]-biphenyl-4,4'-diamine, (DNTPD), a phthalocyanine compound such as copper phthalocyanine, 4,4',4"-tris (3-methylphenylphenylamino)triphenylamine (m-MTDATA), N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB), 4,4'4"-Tris(N,N-diphenylamino)triphenylamine (TDATA), 4,4'4"-Tris-(N-(naphthyl-2-yl)-N-phenylamine)triphenylamine (2-TNATA), poly-aniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), and polyaniline/poly(4-styrenesulfonate) (PANI/PSS).



5



10

2-TNATA

15

[0079] The thickness of the HIL may be about 10 nm (100 Å) to about 1000 nm (10000 Å), and preferably may be about 10 nm (100 Å) to about 100 nm (1000 Å). When the thickness of the HIL is within these ranges, the HIL may have good hole injecting ability without a substantial increase in driving voltage.


20

[0080] Then, a HTL may be formed on the HIL by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like. When the HTL is formed using vacuum deposition or spin coating, the conditions for deposition and coating may be similar to those for the formation of the HIL, though the conditions for the deposition and coating may vary according to the material that is used to form the HTL.

25

[0081] Non-limiting examples of suitable known hole transport materials are carbazole derivatives, such as N-phenylcarbazole or polyvinylcarbazole, N,N'-bis(3-methylphenyl)-N,N'-diphenyl-[1,1'-biphenyl]-4,4'-diamine (TPD), 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), and N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB).

30



35

TPD**NPB**

40

[0082] The thickness of the HTL may be about 5 nm (50 Å) to about 200 nm (2000 Å), and preferably may be about 10 nm (100 Å) to about 150 nm (1500 Å). When the thickness of the HTL is within these ranges, the HTL may have good hole transporting ability without a substantial increase in driving voltage.

45


[0083] The H-functional layer (having both hole injection and hole transport capabilities) may contain at least one material from each group of the hole injection layer materials and hole transport layer materials. The thickness of the H-functional layer may be from about 10 nm (100Å) to about 1000 nm (10,000Å), and preferably may be from about 10 nm (100Å) to about 100 nm (1,000Å). When the thickness of the H-functional layer is within these ranges, the H-functional layer may have good hole injection and transport capabilities without a substantial increase in driving voltage.

50


[0084] At least one of the HIL, HTL, and H-functional layer may include at least one of a compound of Formula 300 below and a compound of Formula 350 below:

55

Formula 300



Formula 350



[0085] In each of Formulae 300 and 350, Ar_{11} and Ar_{12} may each independently be a substituted or unsubstituted C_6 - C_{60} arylene group and Ar_{21} and Ar_{22} may each independently be a substituted or unsubstituted C_6 - C_{60} aryl group. Ar_{11} and Ar_{12} may each independently be one of a phenylene group, a naphthylene group, a phenanthrenylene group, and a pyrenylene group; and a phenylene group, a naphthylene group, a phenanthrenylene group, a fluorenylene group, and a pyrenylene group that are substituted with at least one of a phenyl group, a naphthyl group, and an anthryl group. Ar_{21} and Ar_{22} may each independently be one of a phenyl group, a naphthyl group, a phenanthrenyl group, and a pyrenyl group; and a phenyl group, a naphthyl group, a phenanthrenyl group, a fluorenyl group, and a pyrenyl group that are substituted with at least one of a phenyl group, a naphthyl group, and an anthryl group.

[0086] In Formula 300, e and f may each independently be an integer from 0 to 5, for example, may be 0, 1, or 2. In a non-limiting embodiment, e may be 1, and f may be 0.


[0087] In each of Formulae 300 and 350, R_{51} to R_{58} , R_{61} to R_{69} , R_{71} , and R_{72} may each independently be a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, $-NO_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{60} cycloalkyl group, a substituted or unsubstituted C_6 - C_{60a} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, or a substituted or unsubstituted C_6 - C_{60} arylthiol group. In some non-limiting embodiments, R_5 , to R_{58} , R_{61} to R_{69} , R_{71} , and R_{72} may each independently be one of a hydrogen atom; a deuterium atom; a halogen atom; a hydroxyl group; a cyano group; $-NO_2$; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a C_1 - C_{10} alkyl group (for example, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, or the like); a C_1 - C_{10} alkoxy group (for example, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, or the like); a C_1 - C_{10} alkyl group and a C_1 - C_{10} alkoxy group that are substituted with at least one of a deuterium

atom, a halogen atom, a hydroxyl group, a cyano group, $-\text{NO}_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid or a salt thereof; a phenyl group; a naphthyl group; an anthryl group; a fluorenyl group; a pyrenyl group; and a phenyl group, a naphthyl group, an anthryl group, a fluorenyl group, and a pyrenyl group that are substituted with at least one of a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, $-\text{NO}_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{10} alkyl group, and a C_1 - C_{10} alkoxy group.

[0088] In Formula 300, R_{59} may be one of a phenyl group; a naphthyl group; an anthryl group; a biphenyl group; a pyridyl group; and a phenyl group, a naphthyl group, an anthryl group, a biphenyl group, and a pyridyl group that are substituted with at least one of a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, $-\text{NO}_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{20} alkyl group, and a substituted or unsubstituted C_1 - C_{20} alkoxy group.


[0089] In an embodiment the compound of Formula 300 is a compound represented by Formula 300A below:

Formula 300A





[0090] R_{51} , R_{60} , R_{61} and R_{59} in Formula 300A are as defined above, and thus a detailed description thereof will not be provided herein.

[0091] At least one of the HIL, HTL, and H-functional layer may include at least one of compounds represented by Formulae 301 to 320 below:





5



10



15



20




25

30



35


40



45

50


55




[0092] At least one of the HIL, HTL, and H-functional layer may further include a charge-generating material for improved layer conductivity, in addition to a known hole injecting material, hole transport material, and/or material having both hole injection and hole transport capabilities as described above.

[0093] The charge-generating material may be, for example, a p-dopant. The p-dopant may be one of quinine derivatives, metal oxides, and compounds with a cyano group, but are not limited thereto. Non-limiting examples of the p-dopant include quinone derivatives such as tetracyanoquinonodimethane (TCNQ), 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F₄-TCNQ), and the like; metal oxides such as tungsten oxide, molybdenum oxide, and the like; and cyano-containing compounds such as Compound 200 below.

Compound 200



F4-TCNQ



[0094] When the hole injection layer, hole transport layer, or H-functional layer further includes a charge-generating material, the charge-generating material may be homogeneously dispersed or inhomogeneously distributed in the layer.


[0095] A buffer layer may be disposed between at least one of the HIL, HTL, and H-functional layer, and the EML. The buffer layer may compensate for an optical resonance distance of light according to a wavelength of the light emitted from the EML, and thus may increase efficiency. The buffer layer may include any hole injecting material or hole transporting material that is widely known. The buffer layer may include the same material as one of the materials included in the HIL, HTL, and H-functional layer that lies beneath the buffer layer.

[0096] Then, an EML may be formed on the HTL, H-functional layer, or buffer layer by vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like. When the EML is formed using vacuum deposition or spin coating, the deposition and coating conditions may be similar to those for the formation of the HIL, though the conditions for deposition and coating may vary according to the material that is used to form the EML.


[0097] The EML may include at least one of the amine-based compounds of Formula 1.

[0098] The amine-based compound in the EML may serve as a dopant, for example, as a blue fluorescent dopant. The EML may further include a host, in addition to the amine-based compound.

[0099] Example of the host are Alq₃, 4,4'-N,N'-dicarbazole-biphenyl (CBP), poly(n-vinylcarbazole) (PVK), 9,10-di(naphthalene-2-yl)anthracene (ADN), tris(4-carbazoyl-9-ylphenyl)amine (TCTA), 1,3,5-tris(N-phenylbenzimidazole-2-yl)benzene (TPBI), 3-tert-butyl-9,10-di-2-naphthylanthracene (TBADN), E3, distyrylarylene (DSA), dmCBP (see formula below), and Compounds 501 to 509 below, but are not limited thereto.



TPBI




TBADN




E3

5




PVK



ADN

10


15



CBP

20


25




dmCBP

30


35



501




502




503

40


45



504



505




506


55

5


10



507



508



509


15 **[0100]** An anthracene-based compound represented by Formula 400 below may be used as the host.

Formula 400

20

25

30



35 **[0101]** In Formula 400, Ar_{112} and Ar_{112} may each independently be a substituted or unsubstituted C_6-C_{60} arylene group; Ar_{113} to Ar_{116} may each independently be a substituted or unsubstituted C_1-C_{10} alkyl group or a substituted or unsubstituted C_6-C_{60} aryl group; and g , h , i , and j may each independently be an integer selected from 0, 1, 2, 3 and 4.


[0102] Ar_{112} and Ar_{112} in Formula 400 may each independently be a phenylene group, a naphthylene group, a phenanthrenylene group, or a pyrenylene group; or a phenylene group, a naphthylene group, a phenanthrenylene group, a fluorenyl group, or a pyrenylene group that are substituted with at least one of a phenyl group, a naphthyl group, and an anthryl group.

40 **[0103]** In Formula 400 above, g , h , i , and j may each independently be 0, 1, or 2.

45 **[0104]** Ar_{113} to Ar_{116} in Formula 400 may each independently be one of a C_1-C_{10} alkyl group substituted with at least one of a phenyl group, a naphthyl group, and an anthryl group; a phenyl group; a naphthyl group; an anthryl group; a pyrenyl group; a phenanthrenyl group; a fluorenyl group; a phenyl group, a naphthyl group, an anthryl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group that are substituted with at least one of a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, $-NO_2$, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a pyrenyl group, a phenanthrenyl group, and a fluorenyl group; and


50

55




[0105] For example, the anthracene compound of Formula 400 above may be one of the compounds represented by the following formulae, but is not limited thereto:


5




10




15



20




25




30

35




40


45



50


55





[0106] An anthracene-based compound represented by Formula 401 below may be used as the host.

Formula 401



[0107] Ar_{122} to Ar_{125} in Formula 401 above may be defined as described above in conjunction with Ar_{113} of Formula 400, and thus a detailed description thereof will not be provided here.

[0108] Ar₁₂₆ and Ar₁₂₇ in Formula 401 above may each independently be a C₁-C₁₀ alkyl group, for example, a methyl group, an ethyl group, or a propyl group.

[0109] In Formula 401, k and l may each independently be an integer from 0 to 4, for example, 0, 1, or 2.

[0110] For example, the anthracene compound of Formula 401 above may be one of the compounds represented by the following formulae, but is not limited thereto:

5





10

15

20

25

30




[0111] The amine-based compound in the EML may serve as a host. The EML may further include a dopant, for example, a blue dopant, a green dopant, or a red dopant, in addition to the amine-based compound.

[0112] Non-limiting examples of the blue dopant are compounds represented by the following formulae.


35

40


45



F₂Irpic






(F₂ppy)₂Ir(tmd)




Ir(dfppz)₃

50



55




5





10




15



20




25




DPVBi

30



DPAVBi

40



TBPe


45

50

[0113] Non-limiting examples of the red dopant are compounds represented by the following formulae. The red dopant may be DCM or DCJTb, which will be described later.


55

5




PtOEP

10





Ir(piq)₃




Btp₂Ir(acac)

15




Ir(pq)₂(acac)




Ir(2-phq)₃

20


25



Ir(BT)₂(acac)



Ir(flq)₂(acac)




Ir(fliq)₂(acac)


30

35

40



DCM



DCJTb


45

50

[0114] Non-limiting examples of the green dopant are compounds represented by the following formulae. In an embodiment, the green dopant may be C₅₄₅T represented below.


55

5




Ir(ppy)_3

10




$\text{Ir(ppy)}_2(\text{acac})$



Ir(mppy)_3

15




20

C545T

25


[0115] Non-limiting examples of the dopant that may be used in the EML are Pt complexes represented by the following formulae.

30




D1

35




D2




D3

40




D4

45



D5


50




D6

55


5



D7




D8




D9


10



D10



D11




D12


15

20


25



D13




D14




D15

30


35



D16



D17




D18


40

45


50



D19




D20




D21

55


5
10
15
20
25
30
35
40
45
50
55




D22




D23




D24




D25




D26




D27




D28




D29




D30




D31




D32




D33




D34




D35




D36




D37




D38




D39




D40




D41




D42




D43




D44




D45




D46




D47



D48








D49



D50

15 **[0116]** Non-limiting examples of the dopant that may be used in the EML are Os complexes represented by the following formulae.


Os(fppz)₂(CO)₂Os(fppz)₂(PPh₂Me)₂Os(bppz)₂(PPh₃)₂Os(fptz)₂(PPh₂Me)₂Os(lptz)₂(PPhMe₂)₂

40 **[0117]** When the EML includes both a host and a dopant, the amount of the dopant may be from about 0.01 to about 15 parts by weight based on 100 parts by weight of the host. However, the amount of the dopant is not limited to this range.

45 **[0118]** The thickness of the EML may be about 10 nm (100 Å) to about 100 nm (1000 Å), and preferably may be about 20 nm (200 Å) to about 60 nm (600 Å). When the thickness of the EML is within these ranges, the EML may have good light emitting ability without a substantial increase in driving voltage.


50 **[0119]** Then, an ETL may be formed on the EML by vacuum deposition, spin coating, casting, or the like. When the ETL is formed using vacuum deposition or spin coating, the deposition and coating conditions may be similar to those for the formation of the HIL, though the deposition and coating conditions may vary according to a compound that is used to form the ETL. A material for forming the ETL may be any known material that can stably transport electrons injected from an electron injecting electrode (cathode). Examples of materials for forming the ETL are a quinoline derivative, such as tris(8-quinolinolate)aluminum (Alq₃), TAZ, BAlq, beryllium bis(benzoquinolin-10-olate (Bebq₂), 9,10-di(naphthalene-2-yl)anthracene (ADN), Compound 201, and Compound 202, but are not limited thereto.

55






TAZ

Compound 201



BAiq

Compound 202



BCP

[0120] The ETL may include at least one of the amine-based compounds described above.

[0121] When the amine-based material of Formula 1 is used as a material for forming the ETL, efficiency and/or lifetime of the organic light-emitting diode may be improved. The ETL including the amine-based compound of Formula 1 may further include a metal complex, for example, lithium quinolate.



[0122] The thickness of the ETL may be in a range of about 10 nm (100 Å) to about 100 nm (1,000 Å), and preferably, may be about 15 nm (150 Å) to about 50 nm (500 Å). When the thickness of the ETL is within these ranges, the ETL

may have satisfactory electron transporting ability without a substantial increase in driving voltage.

[0123] The ETL may further include a metal-containing material, in addition to any known electron-transporting organic compound.

[0124] The metal-containing material may be a lithium (Li) complex. Non-limiting examples of the Li complex are lithium quinolate (Liq) and Compound 203 below:

Compound 203




[0125] Then, an EIL, which facilitates injection of electrons from the cathode, may be formed on the ETL. Any suitable electron-injecting material may be used to form the EIL.

[0126] Examples of materials for forming the EIL are LiF, NaCl, CsF, Li₂O, and BaO, which are known in the art. The deposition and coating conditions for forming the EIL 18 may be similar to those for the formation of the HIL, though the deposition and coating conditions may vary according to the material that is used to form the EIL 18.

[0127] The thickness of the EIL may be about 0.1 nm (1 Å) to about 10 nm (100 Å), and preferably, may be about 0.3 nm (3 Å) to about 9 nm (90 Å). When the thickness of the EIL is within these ranges, the EIL may have satisfactory electron injection ability without a substantial increase in driving voltage.

[0128] The second electrode 17 is disposed on the organic layer 15. The second electrode 17 may be a cathode that is an electron injection electrode. A material for forming the second electrode 17 may be a metal, an alloy, an electroconductive compound, which have a low work function, or a mixture thereof. In this regard, the second electrode 17 may comprise lithium (Li), magnesium (Mg), aluminium (Al), aluminium (Al)-lithium (Li), calcium (Ca), magnesium (Mg)-indium (In), magnesium (Mg)-silver (Ag), or the like, and may be formed as a thin film type transmission electrode. To manufacture a top-emission light-emitting diode, the transmission electrode may comprise indium tin oxide (ITO) or indium zinc oxide (IZO).

[0129] When a phosphorescent dopant is used in the EML, a HBL may be formed between the HTL and the EML or between the H-functional layer and the EML by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like, in order to prevent diffusion of triplet excitons or holes into the ETL. When the HBL is formed using vacuum deposition or spin coating, the conditions for deposition and coating may be similar to those for the formation of the HIL, although the conditions for deposition and coating may vary according to the material that is used to form the HBL. Any known hole-blocking material may be used. Non-limiting examples of hole-blocking materials are oxadiazole derivatives, triazole derivatives, and phenanthroline derivatives. For example, BCP represented by the following formula may be used as a material for forming the HBL.

**BCP**

[0130] The thickness of the HBL may be from about 2 nm (20 Å) to about 100 nm (1000 Å), and preferably, may be

from about 3 nm (30 Å) to about 30 nm (300 Å). When the thickness of the HBL is within these ranges, the HBL may have a good hole blocking ability without a substantial increase in driving voltage.

[0131] Examples of the unsubstituted C₁-C₆₀ alkyl group (or C₁-C₆₀ alkyl group) as used herein are C₁-C₆₀ linear or branched alkyl groups, such as methyl, ethyl group, propyl, ethyl, propyl, isobutyl, sec-butyl, pentyl, iso-amyl, and hexyl.

5 Examples of the substituted C₁-C₆₀ alkyl group are a C₁-C₆₀ alkyl group of which at least one hydrogen atom is substituted with at least one of a deuterium atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; -NO₂; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a tri(C₆-C₆₀aryl)silyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, and a C₂-C₆₀ alkynyl group; an a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, and a C₂-C₆₀ alkynyl group of which at least one hydrogen atom are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, and a phosphoric acid or a salt thereof,; a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, and a C₆-C₆₀ arylthiol group; and a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, and a C₆-C₆₀ arylthiol group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.

20 **[0132]** The unsubstituted C₁-C₆₀ alkoxy group (or C₁-C₆₀ alkoxy group) may be a group represented by -OA, wherein A is an unsubstituted C₁-C₆₀ alkyl group described above. Examples of the unsubstituted C₁-C₆₀ alkoxy group are a methoxy group, an ethoxy group, and an isopropoxy group. At least one of the hydrogen atoms in the alkoxy group may be substituted with the substituents described above in conjunction with the substituted C₁-C₆₀ alkyl group.

25 **[0133]** The unsubstituted C₂-C₆₀ alkenyl group (or C₂-C₆₀ alkenyl group) is a hydrocarbon chain having a carbon-carbon double bond in the center or at a terminal of the unsubstituted C₂-C₆₀ alkyl group. Examples of the alkenyl group are an ethenyl group, a propenyl group, a butenyl group, and the like. At least one hydrogen atom in the unsubstituted C₂-C₆₀ alkenyl group may be substituted with those substituents described in conjunction with the substituted C₁-C₆₀ alkyl group.

30 **[0134]** The unsubstituted C₂-C₆₀ alkynyl group (C₂-C₆₀ alkynyl group) is a C₂-C₆₀ alkyl group having at least one carbon-carbon triple bond in the center or at a terminal thereof. Examples of the unsubstituted C₂-C₆₀ alkynyl group (or C₂-C₆₀ alkynyl group) are an ethenyl group, a propynyl group, and the like. At least one hydrogen atom in the alkynyl group may be substituted with the substituents described above in conjunction with the C₁-C₆₀ alkyl group.

35 **[0135]** As used herein, the unsubstituted C₃-C₆₀ cycloalkyl group indicates a cyclic, monovalent C₃-C₆₀ saturated hydrocarbon group. Non-limiting examples of the unsubstituted C₃-C₆₀ cycloalkyl group are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cyclooctyl. At least one hydrogen atom in the cycloalkynyl group may be substituted with those substituents described above in conjunction with the C₁-C₆₀ alkyl group.

40 **[0136]** As used herein, the unsubstituted C₃-C₆₀ cycloalkenyl group indicates a non-aromatic, cyclic unsaturated hydrocarbon group with at least one carbon-carbon double bond. Examples of the unsubstituted C₃-C₆₀ cycloalkenyl group are cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, 1,3-cyclohexadienyl, 1,4-cyclohexadienyl, 2,4-cycloheptadienyl, and 1,5-cyclooctadienyl. At least one hydrogen atom in the cycloalkynyl group may be substituted with those substituents described above in conjunction with the substituted C₁-C₆₀ alkyl group.

45 **[0137]** The unsubstituted C₆-C₆₀ aryl group is a monovalent group having a carbocyclic system having 6 to 60 carbon atoms including at least one aromatic ring. The unsubstituted C₆-C₆₀ arylene group is a bivalent group having a carbocyclic system having 6 to 60 carbon atoms including at least one aromatic ring. When the aryl group and the arylene group have at least two rings, they may be fused to each other via a single bond. At least one hydrogen atom in the aryl group and the arylene group may be substituted with the substituents described above in conjunction with the C₁-C₆₀ alkyl group.

50 **[0138]** Examples of the substituted or unsubstituted C₆-C₆₀ aryl group are a phenyl group, a C₁-C₁₀ alkylphenyl group (e.g., an ethylphenyl group), a C₁-C₁₀ alkylbiphenyl group (e.g., an ethylbiphenyl group), a halophenyl group (e.g., an o-, m- or p-fluorophenyl group and a dichlorophenyl group), a dicyanophenyl group, a trifluoromethoxyphenyl group, an o-, m- or p-tolyl group, an o-, m- or p-cumenyl group, a mesityl group, a phenoxyphenyl group, a (α,α-dimethylbenzene)phenyl group, a (N,N'-dimethyl)aminophenyl group, a (N,N'-diphenyl)aminophenyl group, a pentalenyl group, an indenyl group, a naphthyl group, a halonaphthyl group (e.g., a fluoronaphthyl group), a C₁-C₁₀ alkylnaphthyl group (e.g., a methylnaphthyl group), a C₁-C₁₀ alkoxynaphthyl group (e.g., a methoxynaphthyl group), an anthracenyl group, an azulenyl group, a heptalenyl group, an acenaphthylenyl group, a phenalenyl group, a fluorenyl group, an anthraquinolyl group, a methylanthryl group, a phenanthryl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, an ethylchrysenyl group, a picenyl group, a perylenyl group, a chloroperylene group, a pentaphenyl group, a pentacenyl group, a tetraphenylenyl group, a hexaphenyl group, hexacenyl group, a rubicenyl group, a coronenyl group, a trinaphthylenyl group, a heptaphenyl group, a heptacenyl group, a pyranthrenyl group, and an ovalenyl group. Examples of the substituted

C₆-C₆₀ aryl group may be inferred based on those of the unsubstituted C₆-C₆₀ aryl group and the substituted C₁-C₆₀ alkyl group described above. Examples of the substituted or unsubstituted C₆-C₆₀ arylene group may be inferred based on those examples of the substituted or unsubstituted C₆-C₆₀ aryl group described above.

[0139] The unsubstituted C₂-C₆₀ heteroaryl group is a monovalent group having at least one aromatic ring having at least one heteroatom selected from among N, O, P, and S as a ring-forming atom. The unsubstituted C₂-C₆₀ heteroarylene group is a bivalent group having at least one aromatic ring having at least one heteroatom selected from among N, O, P, and S. In this regard, when the heteroaryl group and the heteroarylene group have at least two rings, they may be fused to each other via a single bond. At least one hydrogen atom in the heteroaryl group and the heteroarylene group may be substituted with those substituents described above in conjunction with the C₁-C₆₀ alkyl group.

[0140] Examples of the unsubstituted C₂-C₆₀ heteroaryl group are a pyrazolyl group, an imidazolyl group, an oxazolyl group, a thiazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a pyridinyl group, a pyridazinyl group, a pyrimidinyl group, a triazinyl group, a carbazolyl group, an indolyl group, a quinolinyl group, an isoquinolinyl group, a benzimidazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a dibenzothiophenyl group, a dibenzofuranyl group and a phenanthrolyl group. Examples of the unsubstituted C₂-C₆₀ heteroarylene group may be inferred based on those examples of the substituted or unsubstituted C₂-C₆₀ arylene group described above.

[0141] The substituted or unsubstituted C₆-C₆₀ aryloxy group indicates -OA₂ (wherein A₂ is a substituted or unsubstituted C₆-C₆₀ aryl group described above). The substituted or unsubstituted C₆-C₆₀ arylthiol group indicates -SA₃ (wherein A₃ is a substituted or unsubstituted C₆-C₆₀ aryl group described above).


[0142] Hereinafter, the present invention will be described in detail with reference to the following synthesis examples and other examples. However, these examples are for illustrative purposes only and are not intended to limit the scope of the present invention.

EXAMPLES

Synthesis Example 1: Synthesis of Compound 1

[0143] Compound 1 was synthesized according to Reaction Scheme 1 below:

Reaction Scheme 1



Synthesis of Intermediate 3-1

[0144] 8.60g (20.0 mmol) of Intermediate 1-1, 5.66g (20.0 mmol) of Intermediate 2-1, 1.15g (1.0 mmol) of tetrakis(triphenylphosphine)palladium(0) (Pd(PPh₃)₄), and 8.29g (60.0 mmol) of K₂CO₃ were dissolved in 50 mL of a mixed solution of THF/H₂O (2:1), and then the resultant solution was stirred at about 70°C for about 5 hours. The resultant mixture was cooled to room temperature, followed by three times of extraction with 50 mL of water and 50 mL of diethylether. An organic layer was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified by silica gel column chromatography to obtain 7.33 g (80% Yield) of Intermediate 3-1.

Synthesis of Compound 1

[0145] 4.58 g (10.0 mmol) of Intermediate 3-1, 2.85 g (12.0 mmol) of Intermediate 4-1, 0.18 g (0.2 mmol) of Pd₂(dba)₃ (tris(dibenzylideneacetone)dipalladium(o)), 0.04 g (0.4 mmol) of tri-tert-butylphosphine (P(t-Bu)₃), and 1.44 g (15.0 mmol) of NaOtBu were dissolved in 50 mL of toluene, and the resultant solution was then refluxed for about 3 hours. The resultant mixture was cooled to room temperature, followed by three times of extraction with 40 mL of water and 40 mL of diethylether. An organic layer was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified by silica gel column chromatography to obtain 4.80 g (78% Yield) of Compound 1. This compound was identified using mass spectroscopy/fast atom bombardment (MS/FAB) and ¹H NMR.

C₄₆H₃₀FN: calc. 615.24, and found; 615.22

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.13-8.11 (dd, 1H), 7.87-7.85 (m, 1H), 7.84-7.80 (m, 3H), 7.72-7.69 (m, 2H), 7.67 (d, 1H), 7.65 (d, 1H), 7.59-7.56 (m, 2H), 7.54-7.51 (dd, 1H), 7.48-7.41 (m, 4H), 7.37-7.23 (m, 6H), 7.18-7.14 (m, 2H), 7.09-7.06 (m, 1H), 6.98-6.94 (m, 2H), 6.85-6.83 (dd, 1H), 6.79-6.75 (m, 2H)

Synthesis Example 2: Synthesis of Compound 3

[0146] 4.98 g of Compound 3 (Yield 80%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-3 instead of Intermediate 4-1 was used. Compound 3 was identified using MS/FAB and ¹H NMR.

C₄₇H₃₀N₂: calc. 622.24, and found; 622.23

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.84-7.82 (m, 2H), 7.81 (d, 1H), 7.78-7.76 (m, 1H), 7.72-7.68 (m, 3H), 7.66 (d, 1H), 7.65 (d, 1H), 7.63-7.59 (m, 2H), 7.57-7.52 (m, 3H), 7.47-7.43 (m, 3H), 7.41-7.38 (m, 2H), 7.37-7.27 (m, 6H), 7.17 (dd, 1H), 7.13-7.09 (m, 2H), 6.99-6.95 (m, 1H), 6.88-6.85 (m, 1H)

Synthesis Example 3: Synthesis of Compound 4

[0147] 4.86 g of Compound 4 (Yield 75%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-4 instead of Intermediate 4-1 was used. Compound 4 was identified using MS/FAB and ¹H NMR.

C₄₉H₃₂N₂: calc. 648.26, and found; 648.27

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.86-7.82 (m, 2H), 7.81 (d, 1H), 7.73-7.68 (m, 2H), 7.66 (d, 1H), 7.65-7.52 (m, 3H), 7.60-7.58 (m, 2H), 7.54-7.49 (m, 3H), 7.46-7.42 (m, 3H), 7.40-7.28 (m, 8H), 6.99-6.95 (m, 1H), 6.90-6.84 (m, 4H), 6.74-6.70 (m, 2H)

Synthesis Example 4: Synthesis of Compound 5

[0148] 4.77g of Compound 5 (Yield 70%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-5 instead of Intermediate 4-1 was used. Compound 5 was identified using MS/FAB and ¹H NMR.

C₅₁H₃₆FN: calc. 681.28, and found; 681.27

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.84-7.82 (m, 2H), 7.80 (d, 1H), 7.78-7.75 (m, 1H), 7.72-7.68 (m, 2H), 7.67 (d, 1H), 7.65 (d, 1H), 7.62-7.58 (m, 2H), 7.56-7.52 (m, 2H), 7.47-7.44 (m, 1H), 7.38-7.27 (m, 6H), 7.14-7.08 (m, 2H), 6.98-6.96 (m, 1H), 6.94-6.89 (m, 2H), 6.85-6.83 (dd, 1H), 6.79-6.77 (m, 2H), 6.75 (d, 1H), 6.73-6.70 (m, 2H), 1.66 (s, 6H)

Synthesis Example 5: Synthesis of Compound 6

[0149] 4.87 g of Compound 6 (Yield 66%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-6 instead of Intermediate 4-1 was used. Compound 6 was identified using MS/FAB and ¹H NMR.

C₅₅H₃₅N₃: calc. 737.28, and found; 737.29

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.07-8.05 (m, 1H), 7.86-7.83 (m, 2H), 7.81 (d, 1H), 7.73-7.69 (m, 2H), 7.68-7.64 (m, 2H), 7.62-7.58 (m, 2H), 7.50-7.44 (m, 6H), 7.42-7.28 (m, 11H), 7.26-7.23 (m, 2H), 6.99-6.96 (m, 1H), 6.90-6.86 (m, 2H), 6.81-6.78 (dd, 1H), 6.73-6.69 (m, 2H)

Synthesis Example 6: Synthesis of Compound 7

[0150] 4.82 g of Compound 7 (Yield 71%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-7 instead of Intermediate 4-1 was used. Compound 7 was identified

using MS/FAB and ^1H NMR.

$\text{C}_{49}\text{H}_{30}\text{N}_2\text{S}$: calc. 678.21; and found 678.22

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.15-8.13 (m, 1H), 8.09-8.06 (m, 1H), 7.84-7.80 (m, 4H), 7.72-7.69 (m, 2H), 7.68-7.64 (m, 2H), 7.61-7.58 (m, 2H), 7.54-7.51 (m, 1H), 7.47-7.41 (m, 2H), 7.39-7.27 (m, 8H), 7.16 (d, 1H), 7.13-7.10 (dd, 1H), 7.04-7.01 (m, 1H), 6.93-6.90 (m, 2H), 6.88-6.84 (m, 2H)

Synthesis Example 7: Synthesis of Compound 11

[0151] 4.17 g of Compound 11 (Yield 62%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-11 instead of Intermediate 4-1 was used. Compound 11 was identified using MS/FAB and ^1H NMR.

$\text{C}_{48}\text{H}_{29}\text{F}_2\text{NO}$: calc. 673.22, and found; 673.21

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 7.87-7.85 (m, 1H), 7.84-7.81 (m, 3H), 7.76-7.70 (m, 3H), 7.68-7.66 (dd, 1H), 7.65-7.64 (m, 1H), 7.62-7.58 (m, 3H), 7.55-7.50 (m, 3H), 7.46-7.40 (m, 2H), 7.37-7.29 (m, 5H), 7.16-7.13 (dd, 1H), 7.10-7.03 (m, 2H), 6.98-6.93 (m, 2H), 6.86-6.82 (m, 2H)

Synthesis Example 8: Synthesis of Compound 13

[0152] 4.52 g of Compound 13 (Yield 79%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-13 instead of Intermediate 4-1 was used. Compound 13 was identified using MS/FAB and ^1H NMR.

$\text{C}_{43}\text{H}_{28}\text{N}_2$: calc. 572.23, and found; 572.23

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 7.84-7.82 (m, 2H), 7.80 (d, 1H), 7.73-7.70 (m, 2H), 7.67 (d, 1H), 7.65 (d, 1H), 7.62-7.58 (m, 2H), 7.54-7.52 (dd, 1H), 7.47-7.43 (m, 2H), 7.38-7.29 (m, 7H), 7.22-7.14 (m, 4H), 7.11-7.06 (m, 1H), 6.97-6.95 (m, 1H), 6.89-6.86 (m, 1H), 6.84-6.81 (m, 2H)

Synthesis Example Q: Synthesis of Compound 14

[0153] 4.29 g of Compound 14 (Yield 75%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-14 instead of Intermediate 4-1 was used. Compound 14 was identified using MS/FAB and ^1H NMR.

$\text{C}_{43}\text{H}_{28}\text{N}_2$: calc. 572.23, and found; 572.24

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 7.86-7.82 (m, 2H), 7.81 (d, 1H), 7.73-7.68 (m, 2H), 7.66 (d, 1H), 7.65 (d, 1H), 7.61-7.57 (m, 2H), 7.54-7.49 (m, 1H), 7.46-7.42 (m, 1H), 7.40-7.28 (m, 7H), 7.22-7.17 (m, 4H), 7.10-7.06 (m, 1H), 6.98-6.95 (m, 2H), 6.88-6.84 (m, 1H), 6.80-6.76 (m, 2H)

Synthesis Example 10: Synthesis of Compound 17

[0154] 4.49 g of Compound 17 (Yield 73%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-17 instead of Intermediate 4-1 was used. Compound 17 was identified using MS/FAB and ^1H NMR.

$\text{C}_{43}\text{H}_{28}\text{F}_3\text{N}$: calc. 615.22, and found; 615.23

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 7.85-7.81 (m, 2H), 7.79 (d, 1H), 7.74-7.69 (m, 2H), 7.67 (d, 1H), 7.66 (d, 1H), 7.62-7.59 (m, 2H), 7.56-7.48 (m, 3H), 7.43-7.41 (m, 1H), 7.34-7.23 (m, 5H), 7.18-7.15 (m, 4H), 7.06-7.03 (m, 1H), 6.97-6.95 (m, 2H), 6.86-6.83 (m, 1H), 6.78-6.74 (m, 2H)

Synthesis Example 11: Synthesis of Compound 18

[0155] 5.81 g of Compound 18 (Yield 70%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-18 instead of Intermediate 4-1 was used. Compound 18 was identified using MS/FAB and ^1H NMR.

$\text{C}_{61}\text{H}_{42}\text{N}_2\text{Si}$: calc. 830.31, and found; 830.30

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 7.82-7.80 (m, 2H), 7.79 (d, 1H), 7.73-7.68 (m, 2H), 7.66 (d, 1H), 7.65 (d, 1H), 7.60-7.55 (m, 8H), 7.52-7.49 (m, 1H), 7.45-7.42 (m, 1H), 7.37-7.26 (m, 15H), 7.24-7.20 (m, 3H), 7.16-7.14 (m, 1H), 7.06-7.02 (m, 2H), 6.96-6.94 (m, 2H), 6.80-6.76 (m, 2H)

Synthesis Example 12: Synthesis of Compound 19

[0156] 3.70 g of Compound 19 (Yield 58%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-19 instead of Intermediate 4-1 was used. Compound 19 was identified using MS/FAB and ¹H NMR.

C₄₂H₂₄F₅N: calc. 637.18, and found; 637.19

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.85-7.83 (m, 2H), 7.81 (d, 1H), 7.75-7.71 (m, 2H), 7.68 (d, 1H), 7.66-6.62 (m, 3H), 7.56-7.54 (dd, 1H), 7.50-7.46 (m, 1H), 7.40-7.31 (m, 5H), 7.25-7.20 (m, 2H), 7.12-7.09 (m, 1H), 7.02-6.99 (m, 1H), 6.92-6.88 (m, 4H)

Synthesis Example 13: Synthesis of Compound 20

[0157] 4.54 g of Compound 20 (Yield 76%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-20 instead of Intermediate 4-1 was used. Compound 20 was identified using MS/FAB and ¹H NMR.

C₄₄H₂₇N₃: calc. 597.22, and found; 597.23

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.83-7.81 (m, 2H), 7.80 (d, 1H), 7.72-7.69 (m, 2H), 7.67 (d, 1H), 7.66 (d, 1H), 7.62-7.58 (m, 2H), 7.54-7.51 (m, 1H), 7.47-7.43 (m, 1H), 7.40-7.28 (m, 9H), 7.13-7.10 (m, 1H), 7.02-6.99 (m, 4H), 6.89-6.85 (m, 2H)

Synthesis Example 14: Synthesis of Compound 21

[0158] 3.44 g of Compound 21 (Yield 59%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-21 instead of Intermediate 4-1 was used. Compound 21 was identified using MS/FAB and ¹H NMR.

C₄₂H₂₇F₂N: calc. 583.21, and found; 583.22

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.84-7.82 (m, 2H), 7.81 (d, 1H), 7.74-7.70 (m, 2H), 7.68 (d, 1H), 7.67 (d, 1H), 7.64-7.61 (m, 2H), 7.55-7.52 (dd, 1H), 7.48-7.44 (m, 1H), 7.37-7.30 (m, 5H), 7.23-7.20 (m, 4H), 7.15-7.09 (m, 5H), 7.04-7.00 (m, 2H)

Synthesis Example 15: Synthesis of Compound 22

[0159] 4.03 g of Compound 22 (Yield 56%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-22 instead of Intermediate 4-1 was used. Compound 22 was identified using MS/FAB and ¹H NMR.

C₅₁H₃₃FN₄: calc. 720.27, and found; 720.28

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.52-8.47 (m, 4H), 7.87-7.82 (m, 5H), 7.76-7.73 (m, 2H), 7.70 (d, 1H), 7.68 (d, 1H), 7.62-7.56 (m, 5H), 7.50-7.46 (m, 1H), 7.44-7.31 (m, 7H), 7.24-7.18 (m, 2H), 7.13-7.09 (m, 2H), 7.00-6.98 (m, 1H), 6.85-6.82 (m, 2H)

Synthesis Example 16: Synthesis of Compound 23

[0160] 5.35 g of Compound 23 (Yield 70%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-23 instead of Intermediate 4-1 was used. Compound 23 was identified using MS/FAB and ¹H NMR.

C₅₆H₃₆N₄: calc. 764.29, and found; 764.28

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.85-7.82 (m, 3H), 7.81-7.77 (m, 3H), 7.74-7.70 (m, 2H), 7.69-7.67 (m, 1H), 7.66-7.65 (m, 2H), 7.61-7.52 (m, 5H), 7.47-7.30 (m, 12H), 7.27-7.23 (m, 1H), 7.12-7.10 (m, 1H), 7.02-6.98 (m, 4H), 6.93-6.91 (m, 2H)

Synthesis Example 17: Synthesis of Compound 24

[0161] 4.74 g of Compound 24 (Yield 76%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-24 instead of Intermediate 4-1 was used. Compound 24 was identified using MS/FAB and ¹H NMR.

C₄₇H₃₂N₂: calc. 624.26, and found; 624.25

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.49-8.46 (m, 1H), 7.84-7.78 (m, 5H), 7.76-7.68 (m, 3H), 7.67-7.64 (m, 3H), 7.62-7.59 (m, 2H), 7.55-7.53 (m, 1H), 7.48-7.45 (m, 1H), 7.39-7.28 (m, 6H), 7.23-7.19 (m, 4H), 7.11-7.09 (m, 1H), 7.02-6.98 (m, 3H), 6.90-6.86 (m, 2H)

Synthesis Example 18: Synthesis of Compound 25

[0162] 4.93 g of Compound 25 (Yield 79%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-25 instead of Intermediate 4-1 was used. Compound 25 was identified using MS/FAB and ¹H NMR.

C₄₇H₃₂N₂: calc. 624.26, and found; 624.25

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.69 (d, 1H), 8.46-8.42 (m, 1H), 7.91-7.97 (m, 1H), 7.86-7.81 (m, 3H), 7.75-7.70 (m, 2H), 7.69-7.66 (m, 2H), 7.63-7.60 (m, 2H), 7.57-7.54 (m, 1H), 7.51-7.47 (m, 2H), 7.40-7.29 (m, 7H), 7.15-7.10 (m, 4H), 6.98-6.91 (m, 3H), 6.87-6.85 (m, 1H), 6.82-6.78 (m, 2H)

Synthesis Example 19: Synthesis of Compound 26

[0163] 4.99 g of Compound 26 (Yield 80%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-26 instead of Intermediate 4-1 was used. Compound 26 was identified using MS/FAB and ¹H NMR.

C₄₇H₃₂N₂: calc. 624.26, and found; 624.25

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.56-8.53 (m, 2H), 7.85-7.82 (m, 2H), 7.80 (d, 1H), 7.73-7.69 (m, 2H), 7.68-7.66 (m, 1H), 7.65 (d, 1H), 7.61-7.51 (m, 7H), 7.47-7.44 (m, 1H), 7.38-7.26 (m, 5H), 7.18-7.13 (m, 4H), 7.09-7.04 (m, 1H), 6.93-6.88 (m, 3H), 6.84-6.80 (m, 2H)

Synthesis Example 20: Synthesis of Compound 27

[0164] 4.29 g of Compound 27 (Yield 66%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-27 instead of Intermediate 4-1 was used. Compound 27 was identified using MS/FAB and ¹H NMR.

C₄₈H₃₁N₃: calc. 649.25, and found; 649.26

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.72 (d, 1H), 8.42-8.39 (m, 1H), 7.95-7.92 (m, 1H), 7.84-7.81 (m, 2H), 7.79 (d, 1H), 7.71-7.68 (m, 2H), 7.66 (d, 1H), 7.64 (d, 1H), 7.62-7.57 (m, 2H), 7.55-7.52 (m, 1H), 7.49-7.44 (m, 2H), 7.40-7.26 (m, 9H), 7.11-7.06 (m, 1H), 7.02-7.00 (m, 2H), 6.93-6.89 (m, 2H), 6.79-6.77 (m, 2H)

Synthesis Example 21: Synthesis of Compound 30

[0165] 4.99 g of Compound 30 (Yield 74%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-30 instead of Intermediate 4-1 was used. Compound 30 was identified using MS/FAB and ¹H NMR.

C₅₁H₃₄N₂: calc. 674.27, and found; 674.26

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.74 (d, 1H), 8.44-8.31 (m, 1H), 8.12-8.08 (dd, 1H), 7.94-7.90 (m, 1H), 7.87-7.80 (m, 4H), 7.75-7.71 (m, 2H), 7.68 (d, 1H), 7.65 (d, 1H), 7.60-7.56 (m, 2H), 7.54-7.52 (m, 1H), 7.49-7.41 (m, 5H), 7.38-7.21 (m, 8H), 7.11-7.07 (m, 2H), 7.03-7.00 (m, 3H), 6.87-6.85 (dd, 1H)

Synthesis Example 22: Synthesis of Compound 31

[0166] 4.98 g of Compound 31 (Yield 80%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-31 instead of Intermediate 4-1 was used. Compound 31 was identified using MS/FAB and ¹H NMR.

C₄₇H₃₀N₂: calc. 622.24, and found; 622.23

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.05-8.01 (dd, 1H), 7.89-7.86 (m, 1H), 7.84-7.80 (m, 3H), 7.74-7.69 (m, 2H), 7.67 (d, 1H), 7.65 (d, 1H), 7.59-7.56 (m, 2H), 7.53-7.50 (m, 1H), 7.48-7.41 (m, 4H), 7.40-7.22 (m, 8H), 7.13-7.09 (m, 2H), 7.05-7.03 (m, 1H), 6.96-6.93 (dd, 1H), 6.88-6.84 (m, 2H)

Synthesis Example 23: Synthesis of Compound 32

[0167] 5.40 g of Compound 32 (Yield 73%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-32 instead of Intermediate 4-1 was used. Compound 32 was identified using MS/FAB and ¹H NMR.

C₅₆H₄₀N₂: calc. 740.32, and found; 740.31

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.76 (d, 1H), 8.51-8.47 (dd, 1H), 7.91-7.88 (m, 1H), 7.83-7.79 (m, 3H), 7.78-7.76 (m, 1H), 7.73-7.69 (m, 2H), 7.68 (d, 1H), 7.66 (d, 1H), 7.62-7.58 (m, 2H), 7.56-7.51 (m, 2H), 7.47-7.43 (m, 2H), 7.39-7.26

EP 2 626 399 B1

(m, 8H), 7.15-7.10 (m, 2H), 7.06-7.04 (m, 1H), 6.94-6.90 (m, 3H), 6.87 (d, 1H), 6.83-6.81 (m, 2H), 1.67 (s, 6H)

Synthesis Example 24: Synthesis of Compound 35

5 **[0168]** 5.05 g of Compound 35 (Yield 67%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-35 instead of Intermediate 4-1 was used. Compound 35 was identified using MS/FAB and ¹H NMR.

C₅₈H₃₈N₂O: calc. 778.30, and found; 778.29

10 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.84-7.80 (m, 4H), 7.78-7.76 (m, 1H), 7.74-7.65 (m, 6H), 7.62-7.59 (m, 3H), 7.55-7.49 (m, 4H), 7.46-7.40 (m, 3H), 7.38-7.27 (m, 5H), 7.19-7.11 (m, 2H), 7.05 (d, 1H), 6.87 (d, 1H), 6.79-6.74 (m, 2H), 1.64 (s, 6H)

Synthesis Example 25: Synthesis of Compound 36

15 **[0169]** 5.80 g of Compound 36 (Yield 70%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-36 instead of Intermediate 4-1 was used. Compound 36 was identified using MS/FAB and ¹H NMR.

C₆₃H₄₄N₂: calc. 828.35, and found; 828.34

20 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.08-8.04 (m, 2H), 7.85-7.82 (m, 2H), 7.81 (d, 1H), 7.77-7.75 (m, 1H), 7.72-7.68 (m, 2H), 7.68 (d, 1H), 7.66 (d, 1H), 7.62-7.58 (m, 2H), 7.56-7.52 (m, 2H), 7.47-7.43 (m, 1H), 7.36-7.26 (m, 12H), 7.15-7.10 (m, 4H), 7.04-7.02 (m, 1H), 6.99-6.91 (m, 3H), 6.89 (d, 1H), 6.85-6.81 (m, 2H), 1.65 (s, 6H)

Synthesis Example 26: Synthesis of Compound 18

25 **[0170]** 5.04 g of Compound 38 (Yield 75%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate 4-38 instead of Intermediate 4-1 was used. Compound 38 was identified using MS/FAB and ¹H NMR.

C₅₁H₃₂N₂: calc. 672.26, and found; 672.25

30 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.50-8.47 (m, 1H), 8.16-8.12 (m, 1H), 7.96-7.92 (m, 1H), 7.86-7.82 (m, 2H), 7.80 (d, 1H), 7.73-7.69 (m, 4H), 7.67 (d, 1H), 7.65 (d, 1H), 7.60-7.52 (m, 5H), 7.47-7.41 (m, 2H), 7.39-7.28 (m, 8H), 7.22-7.18 (m, 2H), 7.07-7.03 (m, 1H), 6.97-6.95 (m, 2H)

Synthesis Example 27: Synthesis of Compound 42

Synthesis of Intermediate 3-42

35 **[0171]** Intermediate 3-42 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-42 instead of Intermediate 2-1 was used.

Synthesis of Compound 42

40 **[0172]** 4.54 g of Compound 42 (Yield 65%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-42 and 4-27, instead of Intermediates 3-1 and 4-1, were used. Compound 42 was identified using MS/FAB and ¹H NMR.

C₅₂H₃₃N₃: calc. 699.27, and found; 699.28

45 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.78 (d, 1H), 8.45-8.42 (m, 1H), 8.11-8.08 (m, 1H), 7.94-7.91 (m, 1H), 7.89-7.86 (m, 1H), 7.83-7.80 (m, 2H), 7.78 (d, 1H), 7.72-7.67 (m, 3H), 7.65-7.62 (dd, 1H), 7.58-7.55 (m, 3H), 7.54-7.52 (m, 1H), 7.48-7.43 (m, 2H), 7.40-7.33 (m, 5H), 7.31-7.27 (m, 4H), 7.18-7.15 (dd, 1H), 6.99-6.95 (m, 1H), 6.82-6.79 (m, 2H), 6.77-6.73 (m, 2H)

Synthesis Example 28: Synthesis of Compound 43

[0173] 5.24 g of Compound 43 (Yield 71%) was prepared in the same manner as in the method of preparing Compound 42 of Synthesis Example 27, except that Intermediate 4-43 instead of Intermediate 4-27 was used. Compound 43 was identified using MS/FAB and ¹H NMR.

55 C₅₆H₃₈N₂: calc. 738.30, and found; 738.31

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.10-8.08 (m, 1H), 7.90-7.87 (m, 1H), 7.85-7.82 (m, 2H), 7.80 (d, 1H), 7.78-7.75 (m, 1H), 7.71-7.65 (m, 4H), 7.59-7.57 (m, 1H), 7.55-7.51 (m, 4H), 7.46-7.42 (m, 1H), 7.39-7.27 (m, 8H), 7.16-7.10 (m, 3H), 6.97-6.93 (m, 1H), 6.89-6.86 (m, 2H), 6.83-6.81 (dd, 1H), 6.79 (d, 1H), 1.66 (s, 6H)

Synthesis Example 29: Synthesis of Compound 45Synthesis of Intermediate 3-45

5 **[0174]** Intermediate 3-45 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-45 instead of Intermediate 2-1 was used.

Synthesis of Compound 45

10 **[0175]** 5.38 g of Compound 45 (Yield 80%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-45 and 4-31, instead of Intermediates 3-1 and 4-1, were used. Compound 45 was identified using MS/FAB and ¹H NMR.

C₅₁H₃₂N₂: calc. 672.26, and found; 672.27

15 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.03-8.00 (m, 1H), 7.88-7.85 (m, 1H), 7.84-7.79 (m, 5H), 7.73-7.68 (m, 2H), 7.65-7.63 (m, 1H), 7.60 (d, 1H), 7.55-7.37 (m, 9H), 7.35-7.28 (m, 5H), 7.20-7.15 (m, 2H), 7.10-7.06 (m, 1H), 6.97-6.92 (m, 1H), 6.86-6.85 (dd, 1H), 6.80-6.77 (m, 2H)

Synthesis Example 30: Synthesis of Compound 48

20 **[0176]** 5.50 g of Compound 48 (Yield 76%) was prepared in the same manner as in the method of preparing Compound 45 of Synthesis Example 29, except that Intermediate 4-48 instead of Intermediate 4-31 was used. Compound 48 was identified using MS/FAB and ¹H NMR.

C₅₅H₃₆N₂: calc. 724.29, and found; 724.30

25 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.56-8.53 (dd, 1H), 8.06-8.03 (m, 1H), 7.85-7.78 (m, 6H), 7.76-7.69 (m, 5H), 7.66-7.60 (m, 3H), 7.55-7.40 (m, 7H), 7.36-7.32 (m, 5H), 7.29-7.25 (m, 1H), 7.21-7.17 (m, 2H), 7.11-7.07 (m, 1H), 6.99-6.94 (m, 1H), 6.87-6.85 (dd, 1H), 6.82-6.79 (m, 2H)

Synthesis Example 31: Synthesis of Compound 51Synthesis of Intermediate 3-51

[0177] Intermediate 3-51 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-51 instead of Intermediate 2-1 was used.

Synthesis of Compound 51

[0178] 4.29 g of Compound 51 (Yield 66%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-51 and 4-4, instead of Intermediates 3-1 and 4-1, were used. Compound 51 was identified using MS/FAB and ¹H NMR.

40 C₄₈H₃₁N₃: calc. 649.25, and found; 649.26

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.50 (d, 1H), 7.93-7.90 (m, 2H), 7.89-7.87 (m, 2H), 7.85 (d, 1H), 7.84-7.81 (m, 1H), 7.74-7.69 (m, 2H), 7.65-7.61 (m, 2H), 7.56-7.49 (m, 5H), 7.46-7.37 (m, 6H), 7.36-7.34 (m, 1H), 7.32-7.27 (m, 2H), 7.14-7.10 (m, 2H), 6.99-6.94 (m, 2H), 6.83-6.79 (m, 2H)

Synthesis Example 32: Synthesis of Compound 54

[0179] 4.23 g of Compound 54 (Yield 65%) was prepared in the same manner as in the method of preparing Compound 51 of Synthesis Example 31, except that Intermediate 4-27 instead of Intermediate 4-4 was used. Compound 54 was identified using MS/FAB and ¹H NMR.

50 C₄₇H₃₀N₄: calc. 650.25, and found; 650.24

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.73 (m, 1H), 8.55 (d, 1H), 8.52-8.47 (m, 1H), 7.96-7.88 (m, 5H), 7.86 (d, 1H), 7.84-7.81 (m, 1H), 7.73-7.68 (m, 2H), 7.56-7.53 (m, 1H), 7.49-7.38 (m, 6H), 7.36-7.28 (m, 5H), 7.16-7.13 (m, 2H), 7.00-6.95 (m, 2H), 6.89-6.85 (m, 2H)

55

Synthesis Example 33: Synthesis of Compound 57Synthesis of Intermediate 3-57

5 **[0180]** Intermediate 3-57 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-57 instead of Intermediate 2-1 was used.

Synthesis of Compound 57

10 **[0181]** 4.97 g of Compound 57 (Yield 67%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-57 and 4-32, instead of Intermediates 3-1 and 4-1, were used. Compound 57 was identified using MS/FAB and ¹H NMR.

C₅₅H₃₉N₃: calc. 741.31, and found; 741.32

15 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.77 (dd, 1H), 8.52 (d, 1H), 8.23-8.20 (m, 1H), 7.93-7.90 (m, 1H), 7.89 (d, 1H), 7.87 (d, 1H), 7.85-7.82 (m, 1H), 7.78-7.75 (m, 1H), 7.72-7.68 (m, 2H), 7.53-7.44 (m, 6H), 7.42-7.23 (m, 9H), 7.14-7.09 (m, 2H), 6.99-6.95 (m, 1H), 6.87-6.83 (m, 4H), 6.78 (d, 1H), 1.63 (s, 6H)

Synthesis Example 34: Synthesis of Compound 58Synthesis of Intermediate 3-58

[0182] Intermediate 3-58 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-58 instead of Intermediate 2-1 was used.

Synthesis of Compound 58

[0183] 5.67 g of Compound 58 (Yield 81%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-58 and 4-25, instead of Intermediates 3-1 and 4-1, were used. Compound 58 was identified using MS/FAB and ¹H NMR.

30 C₅₃H₃₆N₂: calc. 700.29, and found; 700.30

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.74 (dd, 1H), 8.51-8.48 (m, 1H), 7.94-7.90 (m, 1H), 7.83-7.77 (m, 5H), 7.74-7.65 (m, 6H), 7.55-7.52 (m, 1H), 7.48-7.44 (m, 4H), 7.38-7.25 (m, 7H), 7.19-7.15 (m, 2H), 7.08-7.03 (m, 3H), 6.96-6.92 (m, 2H), 6.86-6.83 (m, 1H), 6.80-6.79 (m, 2H)

Synthesis Example 35: Synthesis of Compound 59

[0184] 5.46 g of Compound 59 (Yield 78%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-24 instead of Intermediate 4-25 was used. Compound 59 was identified using MS/FAB and ¹H NMR.

40 C₅₃H₃₆N₂: calc. 700.29, and found; 700.29

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.58-8.54 (m, 1H), 7.86-7.80 (m, 7H), 7.78-7.70 (m, 5H), 7.68-7.65 (m, 3H), 7.54-7.51 (m, 1H), 7.47-7.44 (m, 3H), 7.37-7.26 (m, 6H), 7.17-7.13 (m, 2H), 7.09-7.07 (m, 1H), 7.01-6.97 (m, 2H), 6.98-6.93 (m, 3H), 6.84-6.81 (m, 2H)

Synthesis Example 36: Synthesis of Compound 60

[0185] 5.08 g of Compound 60 (Yield 70%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-27 instead of Intermediate 4-25 was used. Compound 60 was identified using MS/FAB and ¹H NMR.

50 C₅₄H₃₅N₃: calc. 725.28, and found; 725.27

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.71-8.69 (m, 1H), 8.44-8.41 (m, 1H), 7.93-7.91 (m, 1H), 7.84-7.78 (m, 5H), 7.74-7.69 (m, 4H), 7.68-7.65 (m, 2H), 7.55-7.52 (m, 1H), 7.48-7.43 (m, 4H), 7.39-7.24 (m, 9H), 7.12-7.11 (m, 1H), 7.03-6.99 (m, 2H), 6.93-6.89 (m, 2H), 6.82-6.80 (m, 2H)

Synthesis Example 37: Synthesis of Compound 61

[0186] 6.04 g of Compound 61 (Yield 74%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-32 instead of Intermediate 4-25 was used. Compound 61 was

EP 2 626 399 B1

identified using MS/FAB and ^1H NMR.

$\text{C}_{62}\text{H}_{44}\text{N}_2$: calc. 816.35, and found; 816.34

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.69-8.67 (dd, 1H), 8.41-8.38 (m, 1H), 7.91-7.88 (m, 1H), 7.83-7.76 (m, 6H), 7.74-7.68 (m, 4H), 7.67-7.64 (m, 2H), 7.56-7.51 (m, 2H), 7.47-7.44 (m, 4H), 7.37-7.25 (m, 8H), 7.15-7.09 (m, 2H), 6.99-6.94 (m, 1H), 6.90-6.86 (m, 3H), 6.83-6.79 (m, 2H), 6.77-6.76 (m, 1H), 1.64 (s, 6H)

Synthesis Example 38: Synthesis of Compound 62

[0187] 5.93 g of Compound 62 (Yield 79%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-30 instead of Intermediate 4-25 was used. Compound 62 was identified using MS/FAB and ^1H NMR.

$\text{C}_{57}\text{H}_{38}\text{N}_2$: calc. 750.30, and found; 750.29

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.51-8.48 (m, 2H), 8.17-8.15 (dd, 1H), 7.87-7.78 (m, 6H), 7.74-7.65 (m, 6H), 7.55-7.51 (m, 5H), 7.48-7.40 (m, 6H), 7.38-7.28 (m, 5H), 7.23 (t, 1H), 7.14-7.13 (m, 1H), 7.06-7.02 (m, 2H), 6.95-6.93 (dd, 1H), 6.88-6.85 (m, 2H)

Synthesis Example 39: Synthesis of Compound 63

[0188] 5.85 g of Compound 63 (Yield 78%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-63 instead of Intermediate 4-25 was used. Compound 63 was identified using MS/FAB and ^1H NMR.

$\text{C}_{57}\text{H}_{38}\text{N}_2$: calc. 750.30, and found; 750.31

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.50-8.47 (m, 2H), 8.15-8.13 (dd, 1H), 7.86-7.77 (m, 6H), 7.73-7.69 (m, 4H), 7.68-7.65 (m, 2H), 7.56-7.52 (m, 5H), 7.49-7.39 (m, 6H), 7.37-7.29 (m, 5H), 7.22 (t, 1H), 7.18-7.16 (m, 1H), 7.11-7.07 (m, 2H), 6.96-6.94 (dd, 1H), 6.89-6.86 (m, 2H)

Synthesis Example 40: Synthesis of Compound 64

[0189] 5.40 g of Compound 64 (Yield 72%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-48 instead of Intermediate 4-25 was used. Compound 64 was identified using MS/FAB and ^1H NMR.

$\text{C}_{57}\text{H}_{38}\text{N}_2$: calc. 750.30, and found; 750.31

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.48-8.45 (m, 1H), 8.16-8.14 (dd, 1H), 7.87-7.82 (m, 3H), 7.81-7.68 (m, 10H), 7.67-7.64 (m, 3H), 7.54-7.51 (m, 1H), 7.48-7.40 (m, 6H), 7.38-7.21 (m, 7H), 7.09-7.07 (m, 1H), 7.01-6.99 (m, 2H), 6.95-6.94 (dd, 1H), 6.87-6.83 (m, 2H)

Synthesis Example 41: Synthesis of Compound 66

[0190] 5.38 g of Compound 66 (Yield 77%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-31 instead of Intermediate 4-25 was used. Compound 66 was identified using MS/FAB and ^1H NMR.

$\text{C}_{53}\text{H}_{34}\text{N}_2$: calc. 698.27, and found; 698.28

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.06-8.04 (dd, 1H), 7.88-7.85 (m, 1H), 7.84-7.79 (m, 4H), 7.78-7.76 (m, 1H), 7.74-7.69 (m, 4H), 7.67-7.64 (m, 2H), 7.56-7.52 (m, 1H), 7.48-7.29 (m, 13H), 7.22 (t, 1H), 7.13-7.11 (m, 1H), 7.05-7.01 (m, 2H), 6.96-6.94 (dd, 1H), 6.87-6.84 (m, 2H)

Synthesis Example 42: Synthesis of Compound 67

[0191] 5.36 g of Compound 67 (Yield 74%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-67 instead of Intermediate 4-25 was used. Compound 67 was identified using MS/FAB and ^1H NMR.

$\text{C}_{55}\text{H}_{36}\text{N}_2$: calc. 724.29, and found; 724.30

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 7.85-7.82 (m, 2H), 7.81-7.79 (m, 2H), 7.78-7.76 (m, 1H), 7.73-7.68 (m, 4H), 7.67 (d, 1H), 7.65 (d, 1H), 7.60-7.57 (m, 2H), 7.54-7.52 (dd, 1H), 7.48-7.28 (m, 13H), 7.26-7.22 (m, 2H), 7.16-7.15 (m, 1H), 7.06-7.01 (m, 2H), 6.94-6.90 (m, 2H), 6.87-6.84 (m, 2H)

Synthesis Example 43: Synthesis of Compound 68

[0192] 5.45 g of Compound 68 (Yield 78%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-68 instead of Intermediate 4-25 was used. Compound 68 was identified using MS/FAB and ¹H NMR.

C₅₃H₃₄N₂: calc. 698.27, and found; 698.28

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.86-7.83 (m, 2H), 7.82-7.79 (m, 2H), 7.78-7.76 (m, 2H), 7.74-7.68 (m, 4H), 7.66-7.63 (m, 3H), 7.57-7.54 (m, 2H), 7.53-7.50 (m, 2H), 7.47-7.43 (m, 3H), 7.41-7.28 (m, 8H), 7.18-7.16 (m, 1H), 7.06-7.05 (dd, 1H), 6.92-6.88 (m, 2H), 6.85-6.82 (m, 2H)

Synthesis Example 44: Synthesis of Compound 70

[0193] 3.56 g of Compound 70 (Yield 53%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-70 instead of Intermediate 4-25 was used. Compound 70 was identified using MS/FAB and ¹H NMR.

C₅₀H₂₉N₃: calc. 671.24, and found; 671.23

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.22-8.20 (m, 2H), 7.92-7.88 (m, 2H), 7.85-7.82 (m, 2H), 7.80 (d, 1H), 7.75-7.69 (m, 4H), 7.67-7.66 (dd, 1H), 7.65-7.63 (m, 2H), 7.61 (d, 1H), 7.55-7.53 (dd, 1H), 7.49-7.42 (m, 7H), 7.38-7.29 (m, 5H), 7.02-6.97 (m, 1H)

Synthesis Example 45: Synthesis of Compound 73

[0194] 4.97g of Compound 73 (Yield 65%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-43 instead of Intermediate 4-25 was used. Compound 73 was identified using MS/FAB and ¹H NMR.

C₅₈H₄₀N₂: calc. 764.32, and found; 764.33

¹H NMR (CDCl₃, 400MHz) δ (ppm) 7.85-7.82 (m, 2H), 7.81-7.79 (m, 2H), 7.78-7.75 (m, 2H), 7.73-7.68 (m, 4H), 7.67-7.64 (m, 2H), 7.56-7.52 (m, 2H), 7.47-7.44 (m, 3H), 7.39-7.27 (m, 8H), 7.13-7.09 (m, 2H), 7.02-7.00 (m, 1H), 6.96-6.92 (m, 2H), 6.86-6.84 (dd, 1H), 6.82-6.79 (m, 2H), 6.77 (d, 1H), 1.65 (s, 6H)

Synthesis Example 46: Synthesis of Compound 76

[0195] 6.06 g of Compound 76 (Yield 78%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-76 instead of Intermediate 4-25 was used. Compound 76 was identified using MS/FAB and ¹H NMR.

C₅₉H₄₀N₂: calc. 776.32, and found; 776.32

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.68-8.67 (m, 1H), 8.39-8.36 (m, 1H), 7.94-7.91 (m, 1H), 7.84-7.80 (m, 3H), 7.79-7.78 (m, 1H), 7.73-7.69 (m, 4H), 7.67-7.61 (m, 4H), 7.55-7.38 (m, 10H), 7.37-7.24 (m, 7H), 7.18-7.16 (m, 1H), 7.06-7.02 (m, 2H), 6.91-6.88 (m, 2H), 6.82-6.78 (m, 2H)

Synthesis Example 47: Synthesis of Compound 77

[0196] 5.22 g of Compound 77 (Yield 72%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-77 instead of Intermediate 4-25 was used. Compound 77 was identified using MS/FAB and ¹H NMR.

C₅₄H₃₅N₃: calc. 725.28, and found; 725.27

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.55-8.52 (m, 2H), 7.85-7.82 (m, 2H), 7.81-7.77 (m, 3H), 7.74-7.70 (m, 4H), 7.67 (d, 1H), 7.65 (d, 1H), 7.58-7.52 (m, 5H), 7.47-7.44 (m, 3H), 7.39-7.28 (m, 7H), 7.11-7.09 (m, 1H), 7.02-6.99 (m, 2H), 6.93-6.89 (m, 2H), 6.81-6.77 (m, 2H)

Synthesis Example 48: Synthesis of Compound 78

[0197] 5.63 g of Compound 78 (Yield 75%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-78 instead of Intermediate 4-25 was used. Compound 78 was identified using MS/FAB and ¹H NMR.

C₅₇H₃₈N₂: calc. 750.30, and found; 750.29

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.67-8.65 (dd, 1H), 8.40-8.37 (m, 1H), 7.95-7.92 (m, 1H), 7.85-7.82 (m, 2H), 7.81-7.79 (m, 2H), 7.78-7.76 (m, 2H), 7.74-7.68 (m, 4H), 7.66-7.63 (m, 3H), 7.58-7.52 (m, 4H), 7.48-7.44 (m, 4H), 7.37-7.25 (m,

EP 2 626 399 B1

8H), 7.16-7.15 (m, 1H), 7.02-7.00 (dd, 1H), 6.95-6.91 (m, 2H), 6.84-6.81 (m, 2H)

Synthesis Example 49: Synthesis of Compound 79

5 **[0198]** 5.00 g of Compound 79 (Yield 69%) was prepared in the same manner as in the method of preparing Compound 58 of Synthesis Example 34, except that Intermediate 4-79 instead of Intermediate 4-25 was used. Compound 79 was identified using MS/FAB and ¹H NMR.

C₅₄H₃₅N₃: calc. 725.28, and found; 725.27

10 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.49-8.46 (m, 1H), 7.84-7.77 (m, 7H), 7.74-7.68 (m, 5H), 7.67-7.63 (m, 3H), 7.56-7.54 (dd, 1H), 7.47-7.43 (m, 3H), 7.39-7.26 (m, 8H), 7.19-7.17 (m, 1H), 7.03-6.99 (m, 2H), 6.92-6.88 (m, 2H), 6.81-6.78 (m, 2H)

Synthesis Example 50: Synthesis of Compound 82

Synthesis of Intermediate 3-82

15 **[0199]** Intermediate 3-82 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-82 instead of Intermediate 2-1 was used.

Synthesis of Compound 82

20 **[0200]** 5.20 g of Compound 82 (Yield 67%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-82 and 4-25, instead of Intermediates 3-1 and 4-1, were used. Compound 82 was identified using MS/FAB and ¹H NMR.

C₅₉H₄₀N₂: calc. 776.32, and found; 776.33

25 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.68 (d, 1H), 8.41-8.38 (m, 1H), 7.94-7.91 (m, 1H), 7.86-7.80 (m, 5H), 7.74-7.64 (m, 10H), 7.56-7.53 (m, 1H), 7.49-7.44 (m, 4H), 7.38-7.24 (m, 7H), 7.13-7.09 (m, 2H), 7.04-6.99 (m, 3H), 6.96-6.92 (m, 2H), 6.87-6.83 (m, 1H), 6.81-6.78 (m, 2H)

Synthesis Example 51: Synthesis of Compound 83

30 **[0201]** 5.58 g of Compound 83 (Yield 68%) was prepared in the same manner as in the method of preparing Compound 82 of Synthesis Example 50, except that Intermediate 4-48 instead of Intermediate 4-25 was used. Compound 83 was identified using MS/FAB and ¹H NMR.

C₆₃H₄₂N₂: calc. 826.33, and found; 826.32

35 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.46-8.43 (m, 1H), 8.17-8.14 (m, 1H), 7.87-7.80 (m, 6H), 7.79-7.75 (m, 2H), 7.74-7.63 (m, 12H), 7.56-7.54 (dd, 1H), 7.49-7.40 (m, 6H), 7.38-7.22 (m, 7H), 7.16-7.14 (m, 1H), 7.05-7.00 (m, 2H), 6.95-6.93 (dd, 1H), 6.85-6.82 (m, 2H)

Synthesis Example 52: Synthesis of Compound 84

Synthesis of Intermediate 3-84

40 **[0202]** Intermediate 3-84 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-84 instead of Intermediate 2-1 was used.

Synthesis of Compound 84

45 **[0203]** 4.90 g of Compound 84 (Yield 62%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-84 and 4-30, instead of Intermediates 3-1 and 4-1, were used. Compound 84 was identified using MS/FAB and ¹H NMR.

C₆₀H₄N₂: calc. 790.33, and found; 790.32

50 ¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.69 (d, 1H), 8.40-8.37 (m, 1H), 8.11-8.08 (m, 1H), 7.93-7.90 (m, 2H), 7.89-7.85 (m, 2H), 7.84-7.79 (m, 4H), 7.72-7.68 (m, 2H), 7.62-7.59 (m, 2H), 7.53-7.51 (m, 1H), 7.49-7.42 (m, 6H), 7.40-7.25 (m, 8H), 7.13-7.11 (m, 1H), 7.07-7.05 (dd, 1H), 6.98-6.96 (dd, 1H), 6.90-6.86 (m, 2H), 6.82 (d, 1H), 1.62 (s, 6H)

Synthesis Example 53 : Synthesis of Compound 85

55 **[0204]** 4.90 g of Compound 84 (Yield 64%) was prepared in the same manner as in the method of preparing Compound

EP 2 626 399 B1

84 of Synthesis Example 52, except that Intermediate 4-27 instead of Intermediate 4-30 was used. Compound 85 was identified using MS/FAB and ^1H NMR.

$\text{C}_{57}\text{H}_{39}\text{N}_3$: calc. 765.31, and found; 765.30

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.68 (d, 1H), 8.38-8.35 (m, 1H), 7.95-7.92 (m, 2H), 7.90-7.88 (m, 1H), 7.85-7.80 (m, 4H), 7.73-7.69 (m, 2H), 7.63-7.60 (m, 2H), 7.55-7.52 (m, 2H), 7.48-7.44 (m, 2H), 7.41-7.33 (m, 5H), 7.32-7.25 (m, 4H), 7.12-7.09 (m, 1H), 7.03-6.96 (m, 2H), 6.93-6.90 (dd, 1H), 6.88-6.84 (m, 2H), 6.80 (d, 1H), 1.61 (s, 6H)

Synthesis Example 54: Synthesis of Compound 86

[0205] 4.73 g of Compound 86 (Yield 64%) was prepared in the same manner as in the method of preparing Compound 84 of Synthesis Example 52, except that Intermediate 4-31 instead of Intermediate 4-30 was used. Compound 86 was identified using MS/FAB and ^1H NMR.

$\text{C}_{56}\text{H}_{38}\text{N}_2$: calc. 738.30, and found; 738.31

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.11-8.08 (m, 1H), 7.94-7.91 (m, 1H), 7.89-7.85 (m, 2H), 7.84-7.79 (m, 4H), 7.72-7.69 (m, 2H), 7.62-7.59 (m, 2H), 7.54-7.51 (m, 1H), 7.49-7.41 (m, 5H), 7.40-7.34 (m, 5H), 7.32-7.25 (m, 3H), 7.14-7.12 (m, 1H), 7.01-6.99 (m, 1H), 6.93-6.87 (m, 3H), 6.85 (d, 1H), 1.62 (s, 6H)

Synthesis Example 55: Synthesis of Compound 89

Synthesis of Intermediate 3-89

[0206] Intermediate 3-89 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-89 instead of Intermediate 2-1 was used.

Synthesis of Compound 89

[0207] 5.61 g of Compound 89 (Yield 75%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-89 and 4-4, instead of Intermediates 3-1 and 4-1, were used. Compound 89 was identified using MS/FAB and ^1H NMR.

$\text{C}_{57}\text{H}_{36}\text{N}_2$: calc. 748.29, and found; 748.30

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.25-8.19 (m, 2H), 8.11-8.09 (m, 1H), 8.06-8.04 (m, 1H), 7.85-7.82 (m, 2H), 7.81-7.80 (m, 1H), 7.78-7.77 (m, 1H), 7.72-7.68 (m, 3H), 7.64-7.61 (m, 2H), 7.59-7.55 (m, 3H), 7.54-7.48 (m, 3H), 7.47-7.43 (m, 3H), 7.42-7.35 (m, 5H), 7.34-7.27 (m, 3H), 7.12-7.08 (m, 2H), 7.02-6.98 (m, 2H), 6.90-6.86 (m, 2H)

Synthesis Example 56: Synthesis of Compound 92

Synthesis of Intermediate 3-92

[0208] Intermediate 3-92 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-92 instead of Intermediate 2-1 was used.

Synthesis of Compound 92

[0209] 4.61 g of Compound 92 (Yield 61%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-92 and 4-27, instead of Intermediates 3-1 and 4-1, were used. Compound 92 was identified using MS/FAB and ^1H NMR.

$\text{C}_{54}\text{H}_{33}\text{N}_3\text{S}$: calc. 755.24, and found; 755.25

^1H NMR (CDCl_3 , 400MHz) δ (ppm) 8.70-8.68 (m, 1H), 8.40-8.37 (m, 2H), 8.01 (d, 1H), 7.93-7.91 (m, 1H), 7.88-7.86 (dd, 1H), 7.85-7.82 (m, 2H), 7.81-7.80 (m, 1H), 7.78-7.76 (m, 1H), 7.73-7.69 (m, 2H), 7.64 (d, 1H), 7.63-7.62 (m, 1H), 7.54-7.51 (m, 1H), 7.48-7.43 (m, 2H), 7.40-7.35 (m, 4H), 7.34-7.24 (m, 5H), 7.13-7.12 (m, 1H), 7.06-7.03 (dd, 1H), 6.98-6.93 (m, 1H), 6.92-6.84 (m, 4H)

Synthesis Example 57: Synthesis of Compound 95

Synthesis of Intermediate 3-95

[0210] Intermediate 3-95 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-95 instead of Intermediate 2-1 was used.

Synthesis of Compound 95

[0211] 4.74 g of Compound 95 (Yield 62%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-95 and 4-30, instead of Intermediates 3-1 and 4-1, were used.

Compound 95 was identified using MS/FAB and ¹H NMR.

C₅₇H₃₆N₂O: calc. 764.28, and found; 764.29

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.69-8.67 (m, 1H), 8.41-8.38 (m, 1H), 8.11-8.08 (m, 1H), 8.04-8.02 (m, 1H), 7.99-7.96 (m, 2H), 7.94-7.91 (m, 1H), 7.87-7.85 (m, 1H), 7.84-7.77 (m, 6H), 7.73-7.69 (m, 2H), 7.55-7.52 (m, 1H), 7.48-7.41 (m, 5H), 7.40-7.36 (m, 3H), 7.35-7.25 (m, 5H), 7.10-7.04 (m, 2H), 6.98-6.96 (m, 1H), 6.94-6.92 (dd, 1H), 6.88-6.85 (m, 2H)

Synthesis Example 58: Synthesis of Compound 97Synthesis of Intermediate 3-97

[0212] Intermediate 3-97 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-97 instead of Intermediate 2-1 was used.

Synthesis of Compound 97

[0213] 4.96 g of Compound 97 (Yield 61%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-97 and 4-4, instead of Intermediates 3-1 and 4-1, were used. Compound 97 was identified using MS/FAB and ¹H NMR.

C₆₁H₃₉N₃: calc. 813.31, and found; 813.32

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.31-8.29 (m, 1H), 7.84-7.82 (m, 2H), 7.81-7.80 (m, 1H), 7.77-7.74 (m, 1H), 7.72-7.68 (m, 4H), 7.64-7.61 (m, 2H), 7.60-7.58 (m, 1H), 7.54-7.47 (m, 7H), 7.46-7.41 (m, 4H), 7.40-7.36 (m, 5H), 7.34-7.28 (m, 4H), 7.24-7.22 (m, 1H), 7.16-7.14 (m, 1H), 7.06-7.00 (m, 2H), 6.86-6.80 (m, 3H)

Synthesis Example 59: Synthesis of Compound 98

[0214] 4.65 g of Compound 98 (Yield 59%) was prepared in the same manner as in the method of preparing Compound 97 of Synthesis Example 58, except that Intermediate 4-31 instead of Intermediate 4-4 was used. Compound 98 was identified using MS/FAB and ¹H NMR.

C₅₉H₃₇N₃: calc. 787.30, and found; 787.29

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.32-8.30 (m, 1H), 8.10-8.07 (dd, 1H), 7.87-7.85 (m, 1H), 7.84-7.80 (m, 2H), 7.80-7.79 (m, 1H), 7.76-7.74 (m, 1H), 7.73-7.68 (m, 4H), 7.61-7.58 (m, 1H), 7.55-7.51 (m, 1H), 7.50-7.43 (m, 7H), 7.42-7.35 (m, 6H), 7.34-7.23 (m, 6H), 7.13-7.10 (m, 1H), 7.04-7.03 (dd, 1H), 6.94-6.91 (dd, 1H), 6.87-6.84 (m, 2H)

Synthesis Example 60: Synthesis of Compound 99Synthesis of Intermediate 3-99

[0215] Intermediate 3-99 was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 2-98 instead of Intermediate 2-1 was used.

Synthesis of Compound 99

[0216] 5.03 g of Compound 99 (Yield 55%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-99 and 4-30, instead of Intermediates 3-1 and 4-1, were used. Compound 99 was identified using MS/FAB and ¹H NMR.

C₆₉H₄₅N₃: calc. 915.36, and found; 915.35

¹H NMR (CDCl₃, 400MHz) δ (ppm) 8.90 (s, 1H), 8.60-8.58 (dd, 1H), 8.45 (s, 1H), 8.13-8.11 (dd, 1H), 7.94-7.91 (m, 1H), 7.87-7.75 (m, 5H), 7.72-7.58 (m, 7H), 7.54-7.21 (m, 23), 7.16-7.14 (m, 1H), 7.03-6.99 (m, 2H), 6.89-6.84 (m, 2H)

Synthesis Example 61: Synthesis of Compound 103

[0217] 5.71 g of Compound 103 (Yield 76%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate B37 instead of Intermediate 4-1 was used. Compound 103 was identified using MS/FAB and ¹H NMR.

$C_{56}H_{37}N_3$: calc. 751.30, found 751.28

1H NMR ($CDCl_3$, 400MHz) δ (ppm) 8.76 (s, 2H), 8.57 (d, 2H), 8.12 (d, 1H), 7.93 (d, 2H), 7.87-7.65 (m, 9H), 7.61-7.27 (m, 15H), 7.14-7.10 (m, 2H), 7.08-7.05 (m, 3H), 6.98 (d, 1H)

5 Synthesis Example 62: Synthesis of Compound 104

[0218] 5.23 g of Compound 104 (Yield 72%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediate B38 instead of Intermediate 4-1 was used. Compound 104 was identified using MS/FAB and 1H NMR.

10 $C_{53}H_{34}N_4$: calc. 726.28, found 726.27

1H NMR ($CDCl_3$, 400MHz) δ (ppm) 8.78 (s, 2H), 8.56 (d, 2H), 7.94 (d, 2H), 7.84-7.80 (m, 3H), 7.72-7.60 (m, 7H), 7.54-7.25 (m, 11H), 7.12-7.08 (m, 5H), 7.02-6.99 (m, 2H)

15 Synthesis Example 63: Synthesis of Compound 107

[0219] 6.21 g of Compound 107 (yield 75%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-58 and B37, instead of Intermediates 3-1 and 4-1, were used. Compound 107 was identified using MS/FAB and 1H NMR.

20 Synthesis of Compound 107


[0220] 6.21 g of Compound 107 (Yield 75%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that Intermediates 3-107 and B37, instead of Intermediates 3-1 and 4-1, were used. Compound 107 was identified using MS/FAB and 1H NMR.

25 $C_{62}H_{41}N_3$: calc. 827.33, found 827.31


1H NMR ($CDCl_3$, 400MHz) δ (ppm) 8.76 (s, 2H), 8.57 (d, 2H), 8.11 (d, 1H), 7.94 (d, 2H), 7.87-7.62 (m, 13H), 7.54-7.26 (m, 15H), 7.12-7.10 (m, 3H), 7.06-7.03 (m, 1H), 6.96-6.92 (m, 2H)

Intermediates


30 **[0221]**




5




3-1



3-42




3-45




3-51


10



3-57




3-58




3-82

15


20



3-84




3-89




3-92

25


30



3-95




3-97




3-99

35


40




4-1



4-3




4-4




4-5


45




4-6



4-7




4-11




4-13

50


55




4-14



4-17




4-18




4-19


5




4-20



4-21




4-22




4-23


10




4-24



4-25




4-26




4-27


15




4-30



4-31




4-32




4-35

20


25




4-36



4-38




4-43




4-48

30


35




4-63



4-67




4-68




4-70

40


45




4-76



4-77




4-78




4-79

50

55



B37



B38

Example 1


[0222] To manufacture an anode, a coming $15 \Omega/\text{cm}^2$ (1200 \AA) ITO glass substrate was cut to a size of 50 mm x 50 mm x 0.7 mm and then sonicated in isopropyl alcohol and pure water each for five minutes, and then cleaned by irradiation of ultraviolet rays for 30 minutes and exposure to ozone. The resulting glass substrate was loaded into a vacuum deposition diode.

[0223] 2-TNATA was deposited on the ITO glass substrate to form an HIL having a thickness of 600 \AA on the anode, and then 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPS) was deposited on the HIL to form a HTL having a thickness of 300 \AA .

[0224] Subsequently, 9,10-di-naphthalene-2-yl-anthracene (ADN) and 4,4'-bis[2-(4-(N,N-diphenylamino)phenyl)vinyl]biphenyl (DPAVBi) were co-deposited on the HTL in a weight ratio of 98:2 to form an EML having a thickness of about 300 \AA .

[0225] Then, Compound 1 was deposited on the EML to form an ETL having a thickness of about 300 \AA , and then LiF was deposited on the ETL to form an EIL having a thickness of about 10 \AA . Then, Al was deposited on the EIL to form a second electrode (cathode) having a thickness of about 3000 \AA , thereby completing the manufacture of an organic light-emitting diode.

<DPAVBi>

Example 2

[0226] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 4, instead of Compound 1, was used to form the ETL.

Example 1

[0227] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 14, instead of Compound 1, was used to form the ETL.

Example 4

[0228] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 23, instead of Compound 1, was used to form the ETL.

Example 5

[0229] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 25, instead of Compound 1, was used to form the ETL.

Example 6

[0230] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 27, instead of Compound 1, was used to form the ETL.

Example 7

[0231] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 31, instead of Compound 1, was used to form the ETL.

Example 8

[0232] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 32, instead of Compound 1, was used to form the ETL.

5

Example 9

[0233] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 42, instead of Compound 1, was used to form the ETL.

10

Example 10

[0234] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 48, instead of Compound 1, was used to form the ETL.

15

Example 11

[0235] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 58, instead of Compound 1, was used to form the ETL.

20

Example 12

[0236] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 60, instead of Compound 1, was used to form the ETL.

25

Example 13

[0237] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 62, instead of Compound 1, was used to form the ETL.

30

Example 14

[0238] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 66, instead of Compound 1, was used to form the ETL.

35

Example 15

[0239] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 70, instead of Compound 1, was used to form the ETL.

40

Example 16

[0240] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 77, instead of Compound 1, was used to form the ETL.

45

Example 17

[0241] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 82, instead of Compound 1, was used to form the EML.

50

Example 18

[0242] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 86, instead of Compound 1, was used to form the ETL.

55

Example 19

[0243] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound

97, instead of Compound 1, was used to form the EML.

Example 20

5 **[0244]** An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 25, instead of DPAVBi, was used to form the EML, and Alq₃, instead of Compound 1, was used to form the ETL.

Example 21

10 **[0245]** An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound 86, instead of DPAVBi, was used to form the EML, and Alq₃, instead of Compound 1, was used to form the ETL.

Example 22

15 **[0246]** An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 103, instead of Compound 1, was used to form the ETL.

Example 23


20 **[0247]** An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 104, instead of Compound 1, was used to form the ETL.

Example 24

25 **[0248]** An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 107, instead of Compound 1, was used to form the ETL.

Comparative Example 1

30 **[0249]** An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Alq₃, instead of Compound 1, was used to form the ETL.




Comparative Example 2

45 **[0250]** Compound A was synthesized according to Reaction Scheme A below:

50

55

Reaction Scheme A

Synthesis of Intermediate 3-A

[0251] Intermediate 3-A was prepared in the same manner as in the method of preparing Intermediate 3-1 of Synthesis Example 1, except that Intermediate 1-A, instead of Intermediate 1, was used.

Synthesis of Compound A

[0252] 3.23 g of Compound A (Yield 63%) was prepared in the same manner as in the method of preparing Compound 1 of Synthesis Example 1, except that 4.24 g (10 mmol) of Intermediate 3-A and 2.23 g (12.0 mmol) of Intermediate 4-A, instead of Intermediates 3-1 and 4-1, were used.


Manufacture of organic light-emitting diode

[0253] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound A, instead of Compound 1, was used to form the ETL.

Comparative Example 3

[0254] Compound B was synthesized according to Reaction Scheme B below:

Reaction Scheme B

Synthesis of Compound B


[0255] Intermediate B was prepared in the same manner as in the method of preparing Compound A of Comparative Example 2, except that Intermediate 4-B, instead of Intermediate 4-A, was used.

Manufacture of organic light-emitting diode

[0256] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound B, instead of Compound 1, was used to form the ETL.

Comparative Example 4

[0257] Compound C was synthesized according to Reaction Scheme C below:

Synthesis of Compound C


[0258] Compound C was prepared in the same manner as in the method of preparing Compound A of Comparative Example 2, except that Intermediate 4-C, instead of Intermediate 4-A, was used.

Manufacture of organic light-emitting diode

[0259] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound C, instead of Compound 1, was used to form the ETL.

Comparative Example 5

[0260] Compound D was synthesized according to Reaction Scheme D below:

Synthesis of Compound D

[0261] Compound D was prepared in the same manner as in the method of preparing Compound A of Comparative Example 2, except that Intermediate 4-D, instead of Intermediate 4-A, was used.

Manufacture of organic light-emitting diode

[0262] An organic light-emitting diode was manufactured in the same manner as in Example 1, except that Compound D, instead of Compound 1, was used to form the ETL.

Evaluation Example 1

[0263] Driving voltages, luminances, emitting-light colours, efficiencies (at current density of 50mA/cm², and half-life spans (at 100mA/cm²) of the organic light-emitting diodes of Examples 1 to 24 and Comparative Examples 1 to 5 were measured using a PR650 (Spectroscan) Source Measurement Unit (available from Photo Research, Inc.). The results are shown in Table 1 below.

Table 1

	EML host	EML dopant	ETL	Driving voltage (V)	Luminance (cd/m ²)	Efficiency (cd/A)	Light-emitting colour	Half-life span (hr)
Example 1	ADN	DPVABi	Compound 1	5.36	3,335	6.67	blue	492
Example 2	ADN	DPVABi	Compound 4	5.32	3,390	6.78	blue	596
Example 3	ADN	DPVABi	Compound 14	5.29	3,380	6.76	blue	542

EP 2 626 399 B1

(continued)

	EML host	EML dopant	ETL	Driving voltage (V)	Luminance (cd/m ²)	Efficiency (cd/A)	Light-emitting colour	Half-life span (hr)	
5	Example 4	ADN	DPVABi	Compound 23	5.24	3,470	6.94	blue	589
10	Example 5	ADN	DPVABi	Compound 25	5.26	3,465	6.93	blue	536
	Example 6	ADN	DPVABi	Compound 27	5.21	3,505	7.01	blue	564
15	Example 7	ADN	DPVABi	Compound 31	5.26	3,280	6.56	blue	632
	Example 8	ADN	DPVABi	Compound 32	5.30	3,440	6.88	blue	559
20	Example 9	ADN	DPVABi	Compound 42	5.23	3,565	7.13	blue	658
	Example 10	ADN	DPVABi	Compound 48	5.31	3,435	6.87	blue	524
25	Example 11	ADN	DPVABi	Compound 58	5.26	3,580	7.16	blue	532
	Example 12	ADN	DPVABi	Compound 60	5.24	3,515	7.03	blue	582
30	Example 13	ADN	DPVABi	Compound 62	5.31	3,605	7.21	blue	529
	Example 14	ADN	DPVABi	Compound 66	5.27	3,345	6.69	blue	608
35	Example 15	ADN	DPVABi	Compound 70	5.16	3,015	6.03	blue	583
	Example 16	ADN	DPVABi	Compound 77	5.21	3,560	7.12	blue	637
40	Example 17	ADN	DPVABi	Compound 82	5.32	3,460	6.92	blue	543
	Example 18	ADN	DPVABi	Compound 86	5.43	3,445	6.89	blue	469
45	Example 19	ADN	DPVABi	Compound 97	5.62	3,070	6.14	blue	486
	Example 20	ADN	Compound 25	Alq ₃	6.86	2,190	4.38	blue	216
50	Example 21	ADN	Compound 86	Alq ₃	6.97	2,260	4.52	blue	238
	Example 22	ADN	DPVABi	Compound 103	5.35	3,640	7.82	blue	659
55	Example 23	ADN	DPVABi	Compound 104	5.12	3,445	6.89	blue	618
	Example 24	ADN	DPVABi	Compound 107	5.63	3,590	7.18	blue	689

(continued)

	EML host	EML dopant	ETL	Driving voltage (V)	Luminance (cd/m ²)	Efficiency (cd/A)	Light-emitting colour	Half-life span (hr)	
5	Comparative Example 1	ADN	DPVABi	Alq ₃	7.35	2,065	4.13	blue	145
10	Comparative Example 2	ADN	DPVABi	Compound A	5.71	2,865	5.73	blue	311
	Comparative Example 3	ADN	DPVABi	Compound B	5.75	2,730	5.46	blue	320
15	Comparative Example 4	ADN	DPVABi	Compound C	5.73	2,845	5.69	blue	297
	Comparative Example 5	ADN	DPVABi	Compound D	5.73	2,830	5.66	blue	213

[0264] Referring to Table 1, the organic light-emitting devices of Examples 1 to 19 and 22 to 24 are found to have lower driving voltages, higher luminances, higher efficiencies, and better lifetime characteristics as compared to the organic light-emitting diodes of Comparative Examples 1 to 5. The organic light-emitting devices of Examples 20 and 21 are found to have lower driving voltages and better lifetime characteristics as compared to the organic light-emitting diode of Comparative Example 1.


[0265] As described above, an organic light-emitting diode including any of the amine-based compounds according to the invention in its first aspect may have a low driving voltage, a high luminance, a high efficiency, and a long lifetime.

[0266] While the present invention has been described in connection with certain exemplary embodiments thereof, it is to be understood that the invention is not limited to the disclosed embodiments but, on the contrary, is intended to cover various modifications and equivalent arrangements included within the scope of the appended claims.

Claims

1. An amine-based compound represented by Formula 1 below:

Formula 1



wherein, in Formula 1, Ar₁ and Ar₂ are each independently a substituted or unsubstituted C₆-C₆₀ aryl group or a substituted or unsubstituted C₂-C₆₀ heteroaryl group; and wherein Ar₁ and Ar₂ are optionally linked by a single bond;


X₁ is a substituted or unsubstituted C₆-C₆₀ arylene group or a substituted or unsubstituted C₂-C₆₀ heteroarylene group;

m is an integer from 1 to 5; and


at least one substituent of each of the substituted C₆-C₆₀ aryl group, the substituted C₂-C₆₀ heteroaryl group, the substituted C₆-C₆₀ arylene group, and the substituted C₂-C₆₀ heteroarylene group is one of a deuterium

atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; -NO₂; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; a phosphoric acid or a salt thereof; a tri(C₆-C₆₀aryl)silyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group and a C₂-C₆₀ alkynyl group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof; a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthiol group; a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, and a C₆-C₆₀ arylthiol group that is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one fluorine (F), a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group and a C₂-C₆₀ heteroaryl group, wherein at least one of Ar₁ and Ar₂ is a C₆-C₆₀ aryl group substituted with at least one electron withdrawing group selected from the group consisting of -F; -CN; -NO₂; a C₁-C₆₀ alkyl group substituted with at least one -F; a C₂-C₆₀ heteroaryl group; and a C₂-C₆₀ heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.


2. The amine-based compound of claim 1, wherein the at least one electron withdrawing group is selected from the group consisting of: -F; -CN; -NO₂; a C₁-C₂₀ alkyl group substituted with at least one -F; a C₂-C₂₀ heteroaryl group including a ring-containing a N atom; and a C₂-C₂₀ heteroaryl group that includes a ring-containing a N atom and is substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₂-C₂₀ heteroaryl group.
3. The amine-based compound of claim 1 or claim 2, wherein the at least one electron withdrawing group is selected from the group consisting of -F; -CN; -CH₂F; -CHF₂; -CF₃; and groups represented by Formulae 2(1) to 2(14) below:




Formula 2(1)




Formula 2(2)




Formula 2(3)




Formula 2(4)




Formula 2(5)




Formula 2(6)




Formula 2(7)




Formula 2(8)




Formula 2(9)




Formula 2(10)




Formula 2(11)



Formula 2(12)



Formula 2(13)




Formula 2(14)

wherein, in Formulae 2(1) to 2(14), Z₁₁ to Z₁₈ are each independently a hydrogen atom, a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, or a carbazolyl group.


4. The amine-based compound of any preceding claim, wherein Ar₁ and Ar₂ are each independently a C₆-C₆₀ aryl group selected from a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted quinolyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, or a substituted or unsubstituted phenanthrolinyl group.
5. The amine-based compound of any preceding claim, wherein Ar₁ and Ar₂ are linked by a single bond.
6. The amine-based compound of any preceding claim, wherein the at least one of Ar₁ and Ar₂ is a C₆-C₆₀ aryl group substituted with at least two electron withdrawing groups.
7. The amine-based compound of any preceding claim, wherein the at least one of Ar₁ and Ar₂ is a phenyl group, a biphenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, or a fluorenyl group that is substituted with at least two electron withdrawing groups; and the electron withdrawing groups are each independently selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group.
8. The amine-based compound of claim 1, wherein the amine-based compound is represented by Formula 1(1) or

1(2) below:

Formula 1(1)



Formula 1(2)



wherein, in Formula 1(1), Ar_2 is a substituted or unsubstituted C_6 - C_{20} aryl group or a substituted or unsubstituted C_2 - C_{20} heteroaryl group; and

in Formulae 1(1) and 1(2), the ring groups A and B are each independently a substituted C_6 - C_{20} aryl group; R_1 and R_2 are each independently an electron withdrawing group selected from the group consisting of -F; -CN; - NO_2 ; a C_1 - C_{60} alkyl group substituted with at least one -F; a C_2 - C_{60} heteroaryl group; and a C_2 - C_{60} heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, - NO_2 , an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_1 - C_{60} alkyl group substituted with at least one -F, a C_1 - C_{60} alkoxy group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_6 - C_{60} aryl group, and a C_2 - C_{60} heteroaryl group; and

p and q are each independently an integer from 1 to 9.


9. The amine-based compound of claim 8, wherein the amine-based compound is represented by Formula 1(1) in which at least one of R_1 is -CN; or is represented by Formula 1(2) in which at least one of R_1 and one of R_2 is -CN.

10. The amine-based compound of claim 7, wherein the amine-based compound is represented by Formula 1(1), wherein the ring group A is a substituted phenyl group, a substituted biphenyl group, a substituted naphthyl group, a substituted anthryl group, a substituted phenanthrenyl group, a substituted pyrenyl group, or a substituted fluorenyl group; R_1 is at least one electron withdrawing group selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, - NO_2 , an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkyl group substituted with at least one -F, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group; and p is 2, 3, or 4; or., wherein the amine-based compound is represented by Formula 1(2), wherein the ring group A and the ring group B are each independently a substituted phenyl group, a substituted biphenyl group, a substituted naphthyl group, a substituted anthryl group, a substituted phenanthrenyl group, a substituted pyrenyl group, or a substituted fluorenyl group; R_1 and R_2 are each independently at least one electron withdrawing group selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a ben-


zoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group; and p and q are each independently 2, 3, or 4.

11. The amine-based compound of claim 1, wherein the amine-based compound is represented by one of Formulae 1A to 1J below:


Formula 1A




Formula 1B




Formula 1C




Formula 1D



Formula 1E



Formula 1F



Formula 1G


Formula 1H

5

10

15

20



25


Formula 1I

Formula 1J

30

35

40



45


50

55


wherein, in Formulae 1A to 1J, Ar_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenylene group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted azulenyl group, a substituted or unsubstituted heptalenyl group, a substituted or unsubstituted indacenyl group, a substituted or unsubstituted acenaphthyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted phenalenyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluoranthrenyl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted naphthacenyl group, a substituted or unsubstituted picenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted pentaphenyl group, a substituted or unsubstituted hexacenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted imidazolynyl group, a substituted or unsubstituted imidazopyridinyl group, a substituted or unsubstituted imidazopyrimidinyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted benzimidazolyl group, a substituted or unsubstituted indolyl group, a

- substituted or unsubstituted purinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted phthalazinyl group, a substituted or unsubstituted indolizinyl group, a substituted or unsubstituted naphthyridinyl group, a substituted or unsubstituted quinazolinyl group, a substituted or unsubstituted cinolinyl group, a substituted or unsubstituted indazolyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted phenazinyl group, a substituted or unsubstituted phenanthridinyl group, a substituted or unsubstituted pyranyl group, a substituted or unsubstituted chromenyl group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranyl group, a substituted or unsubstituted thiophenyl group, a substituted or unsubstituted benzothiophenyl group, a substituted or unsubstituted isothiazolyl group, a substituted or unsubstituted benzoimidazolyl group, a substituted or unsubstituted isoxazolyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted oxadiazolyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted triazolyl group, a substituted or unsubstituted tetrazolyl group, or a substituted or unsubstituted phenanthrolinyl group,
- substituents of the substituted phenyl group, the substituted pentalenyl group, the substituted indenyl group, the substituted naphthyl group, the substituted azulenyl group, the substituted heptalenyl group, the substituted indacenyl group, the substituted acenaphthyl group, the substituted fluorenyl group, the substituted phenalenyl group, the substituted phenanthrenyl group, the substituted anthryl group, the substituted fluoranthenyl group, the substituted triphenylenyl group, the substituted pyrenyl group, the substituted chrysenyl group, the substituted naphthacenyl group, the substituted picenyl, the substituted perylenyl group, the substituted pentaphenyl group, the substituted hexacenyl group, the substituted pyrrolyl group, the substituted pyrazolyl group, the substituted imidazolyl group, the substituted imidazolyl group, the substituted imidazopyridinyl group, the substituted imidazopyrimidinyl group, the substituted pyridinyl group, the substituted pyrazinyl group, the substituted pyrimidinyl group, the substituted benzoimidazolyl group, the substituted indolyl group, the substituted purinyl group, the substituted quinolinyl group, the substituted phthalazinyl group, the substituted indolizinyl group, the substituted naphthyridinyl group, the substituted quinazolinyl group, the substituted cinolinyl group, the substituted indazolyl group, the substituted carbazolyl group, the substituted phenazinyl group, the substituted phenanthridinyl group, the substituted pyranyl group, the substituted chromenyl group, the substituted furanyl group, the substituted benzofuranyl group, the substituted thiophenyl group, the substituted benzothiophenyl group, the substituted isothiazolyl group, the substituted benzoimidazolyl group, the substituted isoxazolyl group, the substituted dibenzothiophenyl group, the substituted dibenzofuranyl group, the substituted triazinyl group, the substituted oxadiazolyl group, the substituted pyridazinyl group, the substituted triazolyl group, the substituted tetrazolyl group, and the substituted phenanthrolinyl group, and R₁₁ to R₁₉ are each independently a hydrogen atom; a deuterium atom; -F; -Cl; -Br; -I; -CN; a hydroxyl group; -NO₂; an amino group; an amidino group; hydrazine; hydrazone; a carboxyl group or a salt thereof; a sulfonic acid group or a salt thereof; phosphoric acid or a salt thereof; a tri(C₆-C₆₀aryl)silyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, and a C₂-C₆₀ alkynyl group; a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group and a C₂-C₆₀ alkynyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, and phosphoric acid or a salt thereof; a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, and a C₆-C₆₀ arylthiol group; and a C₃-C₆₀ cycloalkyl group, a C₃-C₆₀ cycloalkenyl group, a C₆-C₆₀ aryl group, a C₂-C₆₀ heteroaryl group, a C₆-C₆₀ aralkyl group, a C₆-C₆₀ aryloxy group, and a C₆-C₆₀ arylthiol group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group and a C₂-C₆₀ heteroaryl group, wherein at least one of R₁₁ to R₁₅ in Formulae 1A and 1B, at least one of R₁₁ to R₁₇ in Formulae 1C and 1D, at least one of R₁₁ to R₁₈ in Formulae 1E and 1J, and at least one of R₁₁ to R₁₉ in Formula 1F, 1G, 1H and 1I are each independently an electron withdrawing group selected from the group consisting of -F; -CN; -NO₂; a C₁-C₆₀ alkyl group substituted with at least one -F; a C₂-C₆₀ heteroaryl group; and a C₂-C₆₀ heteroaryl group substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one -F, a C₁-C₆₀ alkoxy group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.
12. The amine-based compound of claim 11, wherein the amine-based compound is represented by Formula 1A-(1) or 1A-(2) below:

Formula 1A-(1)




Formula 1A-(2)




wherein, in Formulae 1A-(1) and 1A-(2), R₁₂, R₁₄, R₂₂, and R₂₄ are each independently an electron withdrawing group selected from the group consisting of a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a triazinyl group, a benzoimidazolyl group, and a carbazolyl group; and a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a quinazoliny group, a phthalazinyl group, a benzoimidazolyl group, and a carbazolyl group that are substituted with at least one of a deuterium atom, -F, -Cl, -Br, -I, -CN; a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, and a carbazolyl group; and in Formula 1A-(1) Ar₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, or a substituted or unsubstituted fluorenyl group.


13. The amine-based compound of claim 12, wherein R₁₂, R₁₄, R₂₂ and R₂₄ are each independently selected from the group consisting of -F; -CN; -CH₂F; -CHF₂; -CF₃; and groups represented by Formulae 2(1) to 2(14) below:




Formula 2(1)




Formula 2(2)




Formula 2(3)




Formula 2(4)




Formula 2(5)




Formula 2(6)




Formula 2(7)




Formula 2(8)




Formula 2(9)




Formula 2(10)




Formula 2(11)



Formula 2(12)



Formula 2(13)




Formula 2(14)

wherein, in Formulae 2(1) to 2(14), Z₁₁ to Z₁₈ are each independently a hydrogen atom, a deuterium atom, -F, -Cl, -Br, -I, -CN, a hydroxyl group, -NO₂, an amino group, an amidino group, hydrazine, hydrazone, a carboxyl group or a salt thereof, a sulfonic acid group or a salt thereof, phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkyl group substituted with at least one -F, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthryl group, a phenanthrenyl group, a pyrenyl group, a pyridinyl group, a triazinyl group, or a carbazolyl group.


14. The amine-based compound of any preceding claim, wherein X₁ is a substituted or unsubstituted phenylene group, a substituted or unsubstituted pentalenylene group, a substituted or unsubstituted indenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted azulenylene group, a substituted or unsubstituted heptalenylene group, a substituted or unsubstituted indacenylene group, a substituted or unsubstituted acenaphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted phenalenylene group, a substituted or unsubstituted phenanthrenylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted fluoranthrenylene group, a substituted or unsubstituted triphenylenylene group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted naphthacenylene group, a substituted or unsubstituted picenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted pentaphenylene group, a substituted or unsubstituted hexacenylenylene group, a substituted or unsubstituted pyrrolylene group, a substituted or unsubstituted pyrazolylenylene group, a substituted or unsubstituted imidazolylene group, a substituted or unsubstituted imidazolinylenylene group, a substituted or unsubstituted imidazopyridinylenylene group, a substituted or unsubstituted imidazopyrimidinylenylene group, a substituted or unsubstituted pyridinylenylene group, a substituted or unsubstituted pyrazinylenylene group, a substituted or unsubstituted pyrimidinylenylene group, a substituted or unsubstituted indolylenylene group, a substituted or unsubstituted purinylenylene group, a substituted or unsubstituted quinolinylenylene group, a substituted or unsubstituted phthalazinylenylene group, a substituted or unsubstituted indolizinylenylene group, a substituted or unsubstituted naphthyridinylenylene group, a substituted or unsubstituted quinazolinylenylene group, a substituted or unsubstituted cinnolinylenylene group, a substituted or unsubstituted indazolylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted phenazinylenylene group, a substituted or unsubstituted phenanthridinylenylene group, a substituted or unsubstituted pyranylene group, a substituted or unsubstituted chromenylenylene group, a substituted or unsubstituted furanylenylene group, a substituted or unsubstituted benzofuranylenylene group, a substituted or unsubstituted thiophenylenylene group, a substituted or unsubstituted benzothiophenylenylene group, a substituted or unsubstituted isothiazolylenylene group, a substituted or unsubstituted benzoimidazolylene group, a substituted or unsubstituted isoxazolylene group, a substituted or unsubstituted dibenzothiophenylenylene group, a substituted or unsubstituted benzofuranylenylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted oxadiazolylenylene group, a substituted or unsubstituted pyridazinylenylene group, a substituted or unsubstituted triazolylene group, or a substituted or unsubstituted tetrazolylenylene group.

15. The amine-based compound of claim 14, wherein X₁ is a group represented by one of Formulae 5(1) to 5(16) below:


5




Formula 5(1)



Formula 5(2)




Formula 5(3)




Formula 5(4)


15




Formula 5(5)



Formula 5(6)



Formula 5(7)




Formula 5(8)


20

25


30



Formula 5(9)




Formula 5(10)




Formula 5(11)

35


40



Formula 5(12)



Formula 5(13)




Formula 5(14)

45

50


55

5




4

10




5




6

15




7

20




8




9

25




10

30




11




12

35




13

40




14




15

45




16

50




17




18

55


5



19




20




21


10



22



23




24

15


20

25


30



25




26




27

35


40



28



29




30

45


50

55


5



31




32




33


10



34



35




36

15


20

25


30



37




38




39

35


40



40




41




42

45


50



43




44




45

55


5



46




47




48


10



49




50




51

15


20



52




53




54

25


30



55



56




57


35

40

45



58




59


50

55


5




10




15




20




25




30




35




40



45




50




55

5




70




71

10




72




73

15

20



74




75


25

30

35




76




77

40

45



78




79


50

55

5




80




81

10

15




82




83

20

25




84




85

30

35




86




87

40

45



88




89


50

55

5




90




91

10

15




92




93

20

25




94




95

30

35




96




97

40

45




98



99

50

55



19. An organic light-emitting diode comprising a first electrode, a second electrode disposed opposite to the first electrode, and an organic layer disposed between the first electrode and the second electrode, the organic layer comprising at least one of the amine-based compounds of claims 1 to 18.

20. The organic light-emitting diode of claim 19, wherein the organic layer comprises at least one of a hole injection layer, a hole transport layer, a functional layer having both hole injection and hole transport capabilities, a buffer layer, an electro blocking layer, an emission layer, a hole blocking layer, an electron transport layer, an electron injection layer, and a functional layer having both electron injection and electron transport capabilities.

21. The organic light-emitting diode of claim 20, wherein the organic layer comprises an electron transport layer, and the amine-based compound is included in the electron transport layer, preferably wherein the electron transport layer further comprises a metal complex.


22. The organic light-emitting diode of claim 20, wherein the organic layer comprises an emission layer, and the amine-

based compound is included in the emission layer, preferably.


wherein the amine-based compound in the emission layer serves as a host, and the emission layer further comprises a blue fluorescent dopant.

23. The organic light-emitting diode of claim 22, wherein the amine-based compound in the emission layer serves as a dopant, and the emission layer further comprises at least one of an anthracene-based compound represented by Formula 400 below and an anthracene-based compound represented by Formula 401 below:

Formula 400



Formula 401




wherein, in Formulae 400 and 401, Ar_{112} and Ar_{112} are each independently a substituted or unsubstituted C_6-C_{60} arylene group; Ar_{113} to Ar_{116} , and Ar_{122} to Ar_{125} are each independently a substituted or unsubstituted C_1-C_{10} alkyl group, or a substituted or unsubstituted C_6-C_{60} aryl group; Ar_{126} and Ar_{127} are each independently a C_1-C_{10} alkyl group; and g , h , i , j , k , and l are each independently an integer from 0 to 4.

24. The organic light-emitting diode of claim 20, wherein the organic layer comprises at least one of a hole injection layer, a hole transport layer, and a functional layer having both hole injection and hole transport capabilities, and the at least one of the hole injection layer, the hole transport layer, and the functional layer having both hole injection and hole transport capabilities comprises a p-dopant.

Patentansprüche

1. Aminbasierte Verbindung, wiedergegeben durch nachstehende Formel 1 :

Formel 1



wobei, in Formel 1, Ar₁ und Ar₂ jeweils unabhängig voneinander ein substituierter oder unsubstituierter C₆-C₆₀-Arylrest oder ein substituierter oder unsubstituierter C₂-C₆₀-Heteroarylrest sind; und wobei Ar₁ und Ar₂ gegebenenfalls über eine Einfachbindung verbunden sind;

X₁ ein substituierter oder unsubstituierter C₆-C₆₀-Arylenrest oder ein substituierter oder unsubstituierter C₂-C₆₀-Heteroarylenrest ist;


m eine ganze Zahl von 1 bis 5 ist; und

mindestens ein Substituent von jedem des substituierten C₆-C₆₀-Arylrestes, des substituierten C₂-C₆₀-Heteroarylrestes, des substituierten C₆-C₆₀-Arylenrestes und des substituierten C₂-C₆₀-Heteroarylenrestes eines ist von einem Deuteriumatom; -F; -Cl; -Br; -I; -CN; einer Hydroxylgruppe; -NO₂; einer Aminogruppe; einer Amidinogruppe; Hydrazin; Hydrazon; einer Carboxylgruppe oder einem Salz davon; einer Sulfonsäuregruppe oder einem Salz davon; einer Phosphorsäure oder einem Salz davon; einem Tri(C₆-C₆₀-aryl)silylrest; einem C₁-C₆₀-Alkylrest, einem C₁-C₆₀-Alkoxyrest, einem C₂-C₆₀-Alkenylrest, einem C₂-C₆₀-Alkinylrest; einem C₁-C₆₀-Alkylrest, einem C₁-C₆₀-Alkoxyrest, einem C₂-C₆₀-Alkenylrest und einem C₂-C₆₀-Alkinylrest, der substituiert ist mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon; einem C₃-C₆₀-Cycloalkylrest, einem C₃-C₆₀-Cycloalkenylrest, einem C₆-C₆₀-Arylrest, einem C₂-C₆₀-Heteroarylrest, einem C₆-C₆₀-Aralkylrest, einem C₆-C₆₀-Aryloxyrest, einem C₆-C₆₀-Arylthiolrest; einem C₃-C₆₀-Cycloalkylrest, einem C₃-C₆₀-Cycloalkenylrest, einem C₆-C₆₀-Arylrest, einem C₂-C₆₀-Heteroarylrest, einem C₆-C₆₀-Aralkylrest, einem C₆-C₆₀-Aryloxyrest und einem C₆-C₆₀-Arylthiolrest, der substituiert ist mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einem C₁-C₆₀-Alkylrest, einem mit mindestens einem Fluoratom (F) substituierten C₁-C₆₀-Alkylrest, einem C₁-C₆₀-Alkoxyrest; einem C₂-C₆₀-Alkenylrest, einem C₂-C₆₀-Alkinylrest; einem C₆-C₆₀-Arylrest und einem C₂-C₆₀-Heteroarylrest, wobei mindestens eines von Ar₁ und Ar₂ ein C₆-C₆₀-Arylrest ist, substituiert mit mindestens einer elektronenziehenden Gruppe, ausgewählt aus -F; -CN; -NO₂; einem mit mindestens einem -F substituierten C₁-C₆₀-Alkylrest; einem C₂-C₆₀-Heteroarylrest; und einem C₂-C₆₀-Heteroarylrest, substituiert mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C₁-C₆₀-Alkylrest, einem mit mindestens einem -F substituierten C₁-C₆₀-Alkylrest, einem C₁-C₆₀-Alkoxyrest, einem C₂-C₆₀-Alkenylrest, einem C₂-C₆₀-Alkinylrest, einem C₆-C₆₀-Arylrest und einem C₂-C₆₀-Heteroarylrest.


2. Aminbasierte Verbindung nach Anspruch 1, wobei mindestens eine elektronenziehende Gruppe ausgewählt ist aus: -F; -CN; -NO₂, einem mit mindestens einem -F substituierten C₁-C₂₀-Alkylrest; einem C₂-C₂₀-Heteroarylrest, umfassend einen ein N-Atom enthaltenden Ring; und einem C₂-C₂₀-Heteroarylrest, der einen ein N-Atom enthaltenden

Ring umfasst und substituiert ist mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, Phosphorsäure oder einem Salz davon, einem C₁-C₂₀-Alkylrest, einem mit mindestens einem -F substituierten C₁-C₂₀-Alkylrest, einem C₁-C₂₀-Alkoxyrest, einem C₆-C₂₀-Arylrest und einem C₂-C₂₀-Heteroarylrest.


3. Aminbasierte Verbindung nach Anspruch 1 oder Anspruch 2, wobei mindestens eine elektronenziehende Gruppe ausgewählt ist aus -F; -CN; -CH₂F; -CHF₂; -CF₃ und den durch die nachstehenden Formeln 2(1) bis 2(14) dargestellten Resten:




Formel 2(1)




Formel 2(2)




Formel 2(3)




Formel 2(4)




Formel 2(5)




Formel 2(6)




Formel 2(7)




Formel 2(8)




Formel 2(9)




Formel 2(10)




Formel 2(11)



Formel 2(12)



Formel 2(13)




Formel 2(14)

wobei, in den Formeln 2(1) bis 2(14), Z₁₁ bis Z₁₈ jeweils unabhängig voneinander ein Wasserstoffatom, ein Deuteriumatom, -F, -Cl, -Br, -I, -CN, eine Hydroxylgruppe, -NO₂, eine Aminogruppe, eine Amidinogruppe, Hydrazin, Hydrazon, eine Carboxylgruppe oder ein Salz davon, eine Sulfonsäuregruppe oder ein Salz davon, Phosphorsäure


oder ein Salz davon, ein C₁-C₂₀-Alkylrest, ein mit mindestens einem -F substituierter C₁-C₂₀-Alkylrest, ein C₁-C₂₀-Alkoxyrest, eine Phenylgruppe, eine Naphthylgruppe, eine Anthrylgruppe, eine Phenanthrenylgruppe, eine Pyrenylgruppe, eine Pyridinylgruppe, eine Triazinylgruppe oder eine Carbazolylgruppe sind.

- 5 4. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei Ar₁ und Ar₂ jeweils unabhängig voneinander ein C₆-C₆₀-Arylrest sind, ausgewählt aus einer substituierten oder unsubstituierten Phenylgruppe, einer substituierten oder unsubstituierten Naphthylgruppe, einer substituierten oder unsubstituierten Fluorenylgruppe, einer substituierten oder unsubstituierten Phenanthrenylgruppe, einer substituierten oder unsubstituierten Anthrylgruppe, einer substituierten oder unsubstituierten Triphenylenylgruppe, einer substituierten oder unsubstituierten Pyrenylgruppe, einer substituierten oder unsubstituierten Chrysenylgruppe, einer substituierten oder unsubstituierten Pyridinylgruppe, einer substituierten oder unsubstituierten Pyrazinylgruppe, einer substituierten oder unsubstituierten Pyrimidinylgruppe, einer substituierten oder unsubstituierten Chinolylgruppe, einer substituierten oder unsubstituierten Carbazolylgruppe, einer substituierten oder unsubstituierten Triazinylgruppe, einer substituierten oder unsubstituierten Dibenzothiophenylgruppe, einer substituierten oder unsubstituierten Dibenzofuranylgruppe oder einer substituierten oder unsubstituierten Phenanthrolinylgruppe.
- 10 5. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei Ar₁ und Ar₂ über eine Einfachbindung verbunden sind.
- 15 6. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei mindestens eines von Ar₁ und Ar₂ ein mit mindestens zwei elektronenziehenden Gruppen substituierter C₆-C₆₀-Arylrest ist.
- 20 7. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei mindestens eines von Ar₁ und Ar₂ eine Phenylgruppe, eine Biphenylgruppe, eine Naphthylgruppe, eine Anthrylgruppe, eine Phenanthrenylgruppe, eine Pyrenylgruppe oder eine Fluorenylgruppe ist, die mit mindestens zwei elektronenziehenden Gruppen substituiert ist; und
- 25 die elektronenziehenden Gruppen jeweils unabhängig voneinander ausgewählt sind aus einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolylgruppe, einer Isochinolylgruppe, einer Chinazolylgruppe, einer Triazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe; und einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolylgruppe, einer Isochinolylgruppe, einer Chinazolylgruppe, einer Phthalazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN; einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C₁-C₂₀-Alkylrest, einem mit mindestens einem -F substituierten C₁-C₂₀-Alkylrest, einem C₁-C₂₀-Alkoxyrest, einer Phenylgruppe, einer Naphthylgruppe, einer Anthrylgruppe, einer Phenanthrenylgruppe, einer Pyrenylgruppe, einer Pyridinylgruppe, einer Triazinylgruppe und einer Carbazolylgruppe.
- 30 8. Aminbasierte Verbindung nach Anspruch 1, wobei die aminbasierte Verbindung durch die nachstehende Formel 1(1) oder 1 (2) wiedergegeben wird:
- 35 40

Formel 1(1)



Formel 1(2)



wobei, in Formel 1(1), Ar₂ ein substituierter oder unsubstituierter C₆-C₂₀-Arylrest oder ein substituierter oder unsubstituierter C₂-C₂₀-Heteroarylrest ist; und in den Formeln 1(1) und 1(2) die Ringgruppen A und B jeweils unabhängig voneinander ein substituierter C₆-C₂₀-Arylrest sind;

R₁ und R₂ jeweils unabhängig voneinander eine elektronenziehende Gruppe sind, ausgewählt aus -F, -CN, -NO₂; einem mit mindestens einem -F substituierten C₁-C₆₀-Alkylrest; einem C₂-C₆₀-Heteroarylrest und einem C₂-C₆₀-Heteroarylrest, substituiert mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C₁-C₆₀-Alkylrest, einem mit mindestens einem -F substituierten C₁-C₆₀-Alkylrest, einem C₁-C₆₀-Alkoxyrest, einem C₂-C₆₀-Alkylrest, einem C₂-C₆₀-Alkylrest; einem C₆-C₆₀-Arylrest und einem C₂-C₆₀-Heteroarylrest; und p und q jeweils unabhängig voneinander eine ganze Zahl von 1 bis 9 sind.

9. Aminbasierte Verbindung nach Anspruch 8, wobei die aminbasierte Verbindung durch die Formel 1 (1) wiedergegeben wird, wobei mindestens eines der R₁ -CN ist; oder durch die Formel 1(2) wiedergegeben wird, wobei mindestens eines der R₁ und eines der R₂ -CN ist.

10. Aminbasierte Verbindung nach Anspruch 7, wobei die aminbasierte Verbindung durch die Formel 1(1) wiedergegeben wird, wobei die Ringgruppe A eine substituierte Phenylgruppe, eine substituierte Biphenylgruppe, eine substituierte Naphthylgruppe, eine substituierte Anthrylgruppe, eine substituierte Phenanthrenylgruppe, eine substituierte Pyrenylgruppe oder eine substituierte Fluorenylgruppe ist;


R₁ mindestens eine elektronenziehende Gruppe ist, ausgewählt aus einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolinylgruppe, einer Isochinolinylgruppe, einer Chinazolinylgruppe, einer Triazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe; und einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolinylgruppe, einer Isochinolinylgruppe, einer Chinazolinylgruppe, einer Phthalazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN; einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C₁-C₂₀-Alkylrest, einem mit mindestens einem -F substituierten C₁-C₂₀-Alkylrest, einem C₁-C₂₀-Alkoxyrest, einer Phenylgruppe, einer Naphthylgruppe, einer Anthrylgruppe, einer Phenanthrenylgruppe, einer Pyrenylgruppe, einer Pyridinylgruppe, einer Triazinylgruppe und einer Carbazolylgruppe; und p 2, 3 oder 4 ist; oder wobei die aminbasierte Verbindung durch die Formel 1(2) wiedergegeben wird, wobei die Ringgruppe A und die Ringgruppe B jeweils unabhängig voneinander eine substituierte Phenylgruppe, eine substituierte Biphenylgruppe, eine substituierte Naphthylgruppe, eine substituierte Anthrylgruppe, eine substituierte Phenanthrenylgruppe, eine substituierte Pyrenylgruppe oder eine substituierte Fluorenylgruppe sind;

R₁ und R₂ jeweils unabhängig voneinander mindestens eine elektronenziehende Gruppe sind, ausgewählt aus einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolinylgruppe, einer Isochinolinylgruppe, einer Chinazolinylgruppe, einer Triazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe; und einer


Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolinylgruppe, einer Isochinolinylgruppe, einer Chinazolinylgruppe, einer Phthalazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN; einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C₁-C₂₀-Alkylrest, einem mit mindestens einem -F substituierten C₁-C₂₀-Alkylrest, einem C₁-C₂₀-Alkoxyrest, einer Phenylgruppe, einer Naphthylgruppe, einer Anthrylgruppe, einer Phenanthrenylgruppe, einer Pyrenylgruppe, einer Pyridinylgruppe, einer Triazinylgruppe und einer Carbazolylgruppe; und p und q jeweils unabhängig voneinander 2, 3 oder 4 sind.

11. Aminbasierte Verbindung nach Anspruch 1, wobei die aminbasierte Verbindung durch eine der nachstehenden Formeln 1A bis 1J wiedergegeben wird:


Formel 1A




Formel 1B



Formel 1C



Formel 1D




Formel 1E

Formel 1F

5

10

15



20

Formel 1G

Formel 1H


25

30

35

40

45




50

55

Formel 1I

Formel 1J



wobei, in den Formeln 1A bis 1J, Ar_2 eine substituierte oder unsubstituierte Phenylgruppe, eine substituierte oder unsubstituierte Pentalenylgruppe, eine substituierte oder unsubstituierte Indenylengruppe, eine substituierte oder unsubstituierte Naphthylgruppe, eine substituierte oder unsubstituierte Azulenylgruppe, eine substituierte oder unsubstituierte Heptalenylgruppe, eine substituierte oder unsubstituierte Indacenylgruppe, eine substituierte oder unsubstituierte Acenaphthylgruppe, eine substituierte oder unsubstituierte Fluorenylgruppe, eine substituierte oder unsubstituierte Phenalenylgruppe, eine substituierte oder unsubstituierte Phenanthrenylgruppe, eine substituierte oder unsubstituierte Anthrylgruppe, eine substituierte oder unsubstituierte Fluoranthenylgruppe, eine substituierte oder unsubstituierte Triphenylenylgruppe, eine substituierte oder unsubstituierte Pyrenylgruppe, eine substituierte oder unsubstituierte Chrysenylgruppe, eine substituierte oder unsubstituierte Naphthacenylgruppe, eine substituierte oder unsubstituierte Picenylgruppe, eine substituierte oder unsubstituierte Perylenylgruppe, eine substituierte oder unsubstituierte Pentaphenylgruppe, eine substituierte oder unsubstituierte Hexacenylgruppe, eine substituierte oder unsubstituierte Pyrrolylgruppe, eine substituierte oder unsubstituierte Pyrazolygruppe, eine substituierte oder unsubstituierte Imidazolylgruppe, eine substituierte oder unsubstituierte Imidazolinygruppe, eine substituierte oder unsubstituierte Imidazopyridinygruppe, eine substituierte oder unsubstituierte Imidazopyrimidinygruppe, eine substituierte oder unsubstituierte Pyridinygruppe, eine substituierte oder unsubstituierte Pyrazinygruppe, eine substituierte oder unsubstituierte Pyrimidinygruppe, eine substituierte oder unsubstituierte Benzimidazolylgruppe, eine substituierte oder unsubstituierte Indolygruppe, eine substituierte oder unsubstituierte Purinygruppe, eine substituierte oder unsubstituierte Chinolinygruppe, eine substituierte oder unsubstituierte Phthalazinygruppe, eine substituierte oder unsubstituierte Indolizinygruppe, eine substituierte oder unsubstituierte Naphthyridinygruppe, eine substituierte oder unsubstituierte Chinazolinygruppe, eine substituierte oder unsubstituierte Cinnolinygruppe, eine substituierte oder unsubstituierte Indazolylgruppe, eine substituierte oder unsubstituierte Carbazolylgruppe, eine substituierte oder unsubstituierte Phenazinygruppe, eine substituierte oder unsubstituierte Phenanthridinygruppe, eine substituierte oder unsubstituierte Pyranlygruppe, eine substituierte oder unsubstituierte Chromenylgruppe, eine substituierte oder unsubstituierte Furanylgruppe, eine substituierte oder unsubstituierte Benzofuranylgruppe, eine substituierte oder unsubstituierte Thiophenylgruppe, eine substituierte oder unsubstituierte Benzothiophenylgruppe, eine substituierte oder unsubstituierte Isothiazolylgruppe, eine substituierte oder unsubstituierte Benzimidazolylgruppe, eine substituierte oder unsubstituierte Isoxazolylgruppe, eine substituierte oder unsubstituierte Dibenzothiophenylgruppe, eine substituierte oder unsubstituierte Dibenzofuranylgruppe, eine substituierte oder unsubstituierte Triazinylgruppe, eine substituierte oder unsubstituierte Oxadiazolylgruppe, eine substituierte oder unsubstituierte Pyridazinylgruppe, eine substituierte oder unsubstituierte Triazolylgruppe, eine substituierte oder unsubstituierte Tetrazolylgruppe oder eine substituierte oder unsubstituierte Phenanthrolinygruppe ist, Substituenten der substituierten Phenylgruppe, der substituierten Pentalenylgruppe, der substituierten Indenylgruppe, der substituierten Naphthylgruppe, der substituierten Azulenylgruppe, der substituierten Heptalenylgruppe, der

substituierten Indacenylgruppe, der substituierten Acenaphthylgruppe, der substituierten Fluorenylgruppe, der substituierten Phenalenylgruppe, der substituierten Phenanthrenylgruppe, der substituierten Anthrylgruppe, der substituierten Fluoranthenylgruppe, der substituierten Triphenylenylgruppe, der substituierten Pyrenylgruppe, der substituierten Chrysenylgruppe, der substituierten Naphthacenylgruppe, der substituierten Picenylgruppe, der substituierten Perylenylgruppe, der substituierten Pentaphenylgruppe, der substituierten Hexacenylgruppe, der substituierten Pyrrolylgruppe, der substituierten Pyrazolylgruppe, der substituierten Imidazolylgruppe, der substituierten Imidazolylgruppe, der substituierten Imidazopyridinylgruppe, der substituierten Imidazopyrimidinylgruppe, der substituierten Pyridinylgruppe, der substituierten Pyrazinylgruppe, der substituierten Pyrimidinylgruppe, der substituierten Benzimidazolylgruppe, der substituierten Indolylgruppe, der substituierten Purinylgruppe, der substituierten Chinolinylgruppe, der substituierten Phthalazinylgruppe, der substituierten Indolizinylgruppe, der substituierten Naphthyridinylgruppe, der substituierten Chinazolinylgruppe, der substituierten Cinnolinylgruppe, der substituierten Indazolylgruppe, der substituierten Carbazolylgruppe, der substituierten Phenazinylgruppe, der substituierten Phenanthridinylgruppe, der substituierten Pyranylgruppe, der substituierten Chromenylgruppe, der substituierten Furanylgruppe, der substituierten Benzofuranylgruppe, der substituierten Thiophenylgruppe, der substituierten Benzothiophenylgruppe, der substituierten Isothiazolylgruppe, der substituierten Benzimidazolylgruppe, der substituierten Isoxazolylgruppe, der substituierten Dibenzothiophenylgruppe, der substituierten Dibenzofuranylgruppe, der substituierten Triazinylgruppe, der substituierten Oxadiazolylgruppe, der substituierten Pyridazinylgruppe, der substituierten Triazolylgruppe, der substituierten Tetrazolylgruppe und der substituierten Phenanthrolinylgruppe, und R_{11} bis R_{19} jeweils unabhängig voneinander ein Wasserstoffatom; ein Deuteriumatom; -F; -Cl; -Br; -I; -CN; eine Hydroxylgruppe; $-NO_2$; eine Aminogruppe; eine Amidinogruppe; Hydrazin; Hydrazon; eine Carboxylgruppe oder ein Salz davon; eine Sulfonsäuregruppe oder ein Salz davon; Phosphorsäure oder ein Salz davon; ein Tri(C_6-C_{60} -aryl)silylrest; ein C_1-C_{60} -Alkylrest, ein C_1-C_{60} -Alkoxyrest, ein C_2-C_{60} -Alkenylrest und ein C_2-C_{60} -Alkinylrest; ein C_1-C_{60} -Alkylrest, ein C_1-C_{60} -Alkoxyrest, ein C_2-C_{60} -Alkenylrest und ein C_2-C_{60} -Alkinylrest, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, $-NO_2$, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon und Phosphorsäure oder einem Salz davon; ein C_3-C_{60} -Cycloalkylrest, ein C_3-C_{60} -Cycloalkenylrest, ein C_6-C_{60} -Arylrest, ein C_2-C_{60} -Heteroarylrest, ein C_6-C_{60} -Aralkylrest, ein C_6-C_{60} -Aryloxyrest und ein C_6-C_{60} -Arylthiolrest; und ein C_3-C_{60} -Cycloalkylrest, ein C_3-C_{60} -Cycloalkenylrest, ein C_6-C_{60} -Arylrest, ein C_2-C_{60} -Heteroarylrest, ein C_6-C_{60} -Aralkylrest, ein C_6-C_{60} -Aryloxyrest und ein C_6-C_{60} -Arylthiolrest, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, $-NO_2$, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, Phosphorsäure oder einem Salz davon, einem C_1-C_{60} -Alkylrest, einem mit mindestens einem -F substituierten C_1-C_{60} -Alkylrest, einem C_1-C_{60} -Alkoxyrest, einem C_2-C_{60} -Alkenylrest, einem C_2-C_{60} -Alkinylrest; einem C_6-C_{60} -Arylrest und einem C_2-C_{60} -Heteroarylrest, sind, wobei mindestens eines von R_{11} bis R_{15} in den Formeln 1A und 1B, mindestens eines von R_{11} bis R_{17} in den Formeln 1C und 1D, mindestens eines von R_{11} bis R_{18} in den Formeln 1E und 1J und mindestens eines von R_{11} bis R_{19} in Formel 1F, 1G, 1H und 1I jeweils unabhängig voneinander eine elektronenziehende Gruppe sind, ausgewählt aus -F; -CN; $-NO_2$, einem mit mindestens einem -F substituierten C_1-C_{60} -Alkylrest; einem C_2-C_{60} -Heteroarylrest; und einem C_2-C_{60} -Heteroarylrest, substituiert mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, $-NO_2$, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C_1-C_{60} -Alkylrest, einem mit mindestens einem -F substituierten C_1-C_{60} -Alkylrest, einem C_1-C_{60} -Alkoxyrest, einem C_2-C_{60} -Alkenylrest, einem C_2-C_{60} -Alkinylrest; einem C_6-C_{60} -Arylrest und einem C_2-C_{60} -Heteroarylrest.

12. Aminbasierte Verbindung nach Anspruch 11, wobei die aminbasierte Verbindung durch die nachstehende Formel 1A-(1) oder 1A-(2) wiedergegeben wird:


Formel 1A-(1)

Formel 1A-(2)

5

10


15



20

25


30




wobei, in den Formeln 1A-(1) oder 1A-(2), R_{12} , R_{14} , R_{22} und R_{24} jeweils unabhängig voneinander eine elektronenziehende Gruppe sind, ausgewählt aus einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolinylgruppe, einer Isochinolinylgruppe, einer Chinazolinylgruppe, einer Triazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe; und einer Pyridinylgruppe, einer Pyrazinylgruppe, einer Pyrimidinylgruppe, einer Chinolinylgruppe, einer Isochinolinylgruppe, einer Chinazolinylgruppe, einer Phthalazinylgruppe, einer Benzimidazolylgruppe und einer Carbazolylgruppe, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN; einer Hydroxylgruppe, $-NO_2$, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C_1 - C_{20} -Alkylrest, einem mit mindestens einem -F substituierten C_1 - C_{20} -Alkylrest, einem C_1 - C_{20} -Alkoxyrest, einer Phenylgruppe, einer Naphthylgruppe, einer Anthrylgruppe, einer Phenanthrenylgruppe, einer Pyrenylgruppe, einer Pyridinylgruppe, einer Triazinylgruppe und einer Carbazolylgruppe; und in Formel 1A-(1) Ar_2 eine substituierte oder unsubstituierte Phenylgruppe, eine substituierte oder unsubstituierte Biphenylgruppe, eine substituierte oder unsubstituierte Naphthylgruppe, eine substituierte oder unsubstituierte Anthrylgruppe, eine substituierte oder unsubstituierte Phenanthrenylgruppe, eine substituierte oder unsubstituierte Pyrenylgruppe oder eine substituierte oder unsubstituierte Fluorenylgruppe ist.

13. Aminbasierte Verbindung nach Anspruch 12, wobei R_{12} , R_{14} , R_{22} und R_{24} jeweils unabhängig voneinander ausgewählt sind aus F; -CN; $-CH_2F$, $-CHF_2$; $-CF_3$ und den durch die nachstehenden Formeln 2(1) bis 2(14) dargestellten Resten:


55




Formel 2(1)




Formel 2(2)




Formel 2(3)




Formel 2(4)




Formel 2(5)




Formel 2(6)




Formel 2(7)




Formel 2(8)




Formel 2(9)




Formel 2(10)




Formel 2(11)



Formel 2(12)



Formel 2(13)




Formel 2(14)

wobei, in den Formeln 2(1) bis 2(14), Z_{11} bis Z_{18} jeweils unabhängig voneinander ein Wasserstoffatom, ein Deuteriumatom, -F, -Cl, -Br, -I, -CN, eine Hydroxylgruppe, -NO₂, eine Aminogruppe, eine Amidinogruppe, Hydrazin, Hydrazon, eine Carboxylgruppe oder ein Salz davon, eine Sulfonsäuregruppe oder ein Salz davon, Phosphorsäure oder ein Salz davon, ein C₁-C₂₀-Alkylrest, ein mit mindestens einem -F substituiertes C₁-C₂₀-Alkylrest, ein C₁-C₂₀-Alkoxyrest, eine Phenylgruppe, eine Naphthylgruppe, eine Anthrylgruppe, eine Phenanthrenylgruppe, eine Pyrenylgruppe, eine Pyridinylgruppe, eine Triazinylgruppe oder eine Carbazolylgruppe sind.


14. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei X_1 eine substituierte oder unsubstituierte Phenylengruppe, eine substituierte oder unsubstituierte Pentalenylengruppe, eine substituierte oder unsubstituierte Indenylengruppe, eine substituierte oder unsubstituierte Naphthylengruppe, eine substituierte oder unsubstituierte Azulenylengruppe, eine substituierte oder unsubstituierte Heptalenylengruppe, eine substituierte oder unsubstituierte Indacenylengruppe, eine substituierte oder unsubstituierte Acenaphthylengruppe, eine substituierte oder un-

substituierte Fluorenylengruppe, eine substituierte oder unsubstituierte Phenalenylengruppe, eine substituierte oder unsubstituierte Phenanthrenylengruppe, eine substituierte oder unsubstituierte Anthrylengruppe, eine substituierte oder unsubstituierte Fluoranthenylengruppe, eine substituierte oder unsubstituierte Triphenylenylengruppe, eine substituierte oder unsubstituierte Pyrenylengruppe, eine substituierte oder unsubstituierte Chrysenylengruppe, eine substituierte oder unsubstituierte Naphthacenylengruppe eine substituierte oder unsubstituierte Picenylengruppe, eine substituierte oder unsubstituierte Perylenylengruppe, eine substituierte oder unsubstituierte Pentaphenylengruppe, eine substituierte oder unsubstituierte Hexacenylengruppe, eine substituierte oder unsubstituierte Pyrrolylengruppe, eine substituierte oder unsubstituierte Pyrazolylengruppe, eine substituierte oder unsubstituierte Imidazolylengruppe, eine substituierte oder unsubstituierte Imidazolinylenylengruppe, eine substituierte oder unsubstituierte Imidazopyridinylenylengruppe, eine substituierte oder unsubstituierte Imidazopyrimidinylenylengruppe, eine substituierte oder unsubstituierte Pyridinylenylengruppe, eine substituierte oder unsubstituierte Pyrazinylenylengruppe, eine substituierte oder unsubstituierte Pyrimidinylenylengruppe, eine substituierte oder unsubstituierte Indolylenylengruppe, eine substituierte oder unsubstituierte Purinylenylengruppe, eine substituierte oder unsubstituierte Chinolinylenylengruppe, eine substituierte oder unsubstituierte Phthalazinylenylengruppe, eine substituierte oder unsubstituierte Indolizinylenylengruppe, eine substituierte oder unsubstituierte Naphthyridinylenylengruppe, eine substituierte oder unsubstituierte Chinazolinylenylengruppe, eine substituierte oder unsubstituierte Cinnolinylenylengruppe, eine substituierte oder unsubstituierte Indazolylengruppe, eine substituierte oder unsubstituierte Carbazolylengruppe, eine substituierte oder unsubstituierte Phenazinylenylengruppe, eine substituierte oder unsubstituierte Phenanthridinylenylengruppe, eine substituierte oder unsubstituierte Pyranlylenylengruppe, eine substituierte oder unsubstituierte Chromenylengruppe, eine substituierte oder unsubstituierte Furanylenylengruppe, eine substituierte oder unsubstituierte Benzofuranylenylengruppe, eine substituierte oder unsubstituierte Thiophenylengruppe, eine substituierte oder unsubstituierte Benzothiophenylengruppe, eine substituierte oder unsubstituierte Isothiazolylengruppe, eine substituierte oder unsubstituierte Benzimidazolylengruppe, eine substituierte oder unsubstituierte Isoxazolylengruppe, eine substituierte oder unsubstituierte Dibenzothiophenylengruppe, eine substituierte oder unsubstituierte Dibenzofuranylenylengruppe, eine substituierte oder unsubstituierte Triazinylenylengruppe, eine substituierte oder unsubstituierte Oxadiazolylengruppe, eine substituierte oder unsubstituierte Pyridazinylenylengruppe, eine substituierte oder unsubstituierte Triazolylengruppe oder eine substituierte oder unsubstituierte Tetrazolylengruppe ist.


15. Aminbasierte Verbindung nach Anspruch 14, wobei X₁ ein durch eine der nachstehenden Formeln 5(1) bis 5(16) dargestellter Rest ist:




Formel 5(1)




Formel 5(2)




Formel 5(3)




Formel 5(4)




Formel 5(5)




Formel 5(6)




Formel 5(7)




Formel 5(8)




Formel 5(9)




Formel 5(10)




Formel 5(11)




Formel 5(12)




Formel 5(13)



Formel 5(14)



Formel 5(15)



Formel 5(16)

wobei, in den Formeln 5(1) bis 5(16), Z₁ bis Z₈ jeweils unabhängig voneinander eines von einem Wasserstoffatom; einem Deuteriumatom; -F; -Cl; -Br; -I; -CN; einer Hydroxylgruppe; -NO₂; einer Aminogruppe; einer Amidinogruppe; Hydrazin; Hydrazon; einer Carboxylgruppe oder einem Salz davon; einer Sulfonsäuregruppe oder einem Salz davon; Phosphorsäure oder einem Salz davon; einem C₁-C₂₀-Alkylrest; einem C₁-C₂₀-Alkoxyrest; einem C₁-C₂₀-Alkylrest und einem C₁-C₂₀-Alkoxyrest, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon und Phosphorsäure oder einem Salz davon; einem C₆-C₂₀-Arylrest; einem C₂-C₂₀-Heteroarylrest; und einem C₆-C₂₀-Arylrest und einem C₂-C₂₀-Heteroarylrest, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, Phosphorsäure oder einem Salz davon, einem C₁-C₂₀-Alkylrest, einem C₁-C₂₀-Alkoxyrest, einem C₆-C₂₀-Arylrest und einem C₂-C₂₀-Heteroarylrest, sind; wobei * eine Bindungsstelle an Anthracen in Formel 1 kennzeichnet; und wobei ** eine Bindungsstelle an N in Formel 1 kennzeichnet.

16. Aminbasierte Verbindung nach Anspruch 15, wobei Z₁ bis Z₈ jeweils unabhängig voneinander eines von einem Wasserstoffatom; einem Deuteriumatom; -F; -Cl; -Br; -I; -CN; einer Hydroxylgruppe; -NO₂; einer Aminogruppe; einer Amidinogruppe; Hydrazin; Hydrazon; einer Carboxylgruppe oder einem Salz davon; einer Sulfonsäuregruppe oder einem Salz davon; einer Phosphorsäure oder einem Salz davon; einer Methylgruppe, einer Ethylgruppe, einer Propylgruppe, einer Butylgruppe und einer Pentylgruppe; einer Methoxygruppe, einer Ethoxygruppe, einer Propoxygruppe, einer Butoxygruppe und einer Pentoxygruppe; einer Methylgruppe, einer Ethylgruppe, einer Propylgruppe, einer Butylgruppe, einer Pentylgruppe, einer Methoxygruppe, einer Ethoxygruppe, einer Propoxygruppe, einer Butoxygruppe und einer Pentoxygruppe, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon und einer Phosphorsäure oder einem Salz davon; einer Phenylgruppe, einer Naphthylgruppe, einer Anthrylgruppe, einer Phenanthrenylgrup-

pe, einer Pyrenylgruppe und einer Fluorenylgruppe; einer Pyridinylgruppe, einer Pyrimidinylgruppe, einer Triazinylgruppe, einer Chinolylgruppe und einer Carbazolylgruppe; einer Phenylgruppe, einer Naphthylgruppe, einer Anthrylgruppe, einer Phenanthrenylgruppe, einer Pyrenylgruppe, einer Fluorenylgruppe; einer Pyridinylgruppe, einer Pyrimidinylgruppe, einer Triazinylgruppe, einer Chinolylgruppe und einer Carbazolylgruppe, die substituiert sind mit mindestens einem von einem Deuteriumatom, -F, -Cl, -Br, -I, -CN, einer Hydroxylgruppe, -NO₂, einer Aminogruppe, einer Amidinogruppe, Hydrazin, Hydrazon, einer Carboxylgruppe oder einem Salz davon, einer Sulfonsäuregruppe oder einem Salz davon, einer Phosphorsäure oder einem Salz davon, einem C₁-C₂₀-Alkylrest und einem C₁-C₂₀-Alkoxyrest, sind.

5


10

17. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei m 1, 2 oder 3 ist.


18. Aminbasierte Verbindung nach einem vorangehenden Anspruch, wobei die aminbasierte Verbindung eine der nachstehenden Verbindungen 1 bis 109 ist:

15


20



1




2




3

25


30



4




5




6

35


40



7




8




9

45


50



10




11




12

55


5



13




14




15


10



16




17




18

15


20



19




20




21

25


30



22



23




24


35

40


45



25



26




27


50

55


5



28




29




30

10


15



31




32




33

20


25



34




35




36

30


35



37




38




39

40


45



40



41




42


50

55


5



43




44




45


10



46



47




48

15


20

25


30



49




50




51

35


40



52




53




54

45


50



55




56




57

55


5
10
15
20
25
30
35
40
45
50
55




58




59




60




61




62




63




64



65




66




67

5




68




69

10

15




70




71

20

25




72




73

30

35




74




75

40

45



76




77

50


55

5




88

10




89

15




90

20




91

25




92

30




93

35




94

40




95

45



96


50




97

55


5
10
15
20
25
30
35
40
45
50
55




98




99




100




101




102




103




104




105




106



107



108




109

19. Organische lichtemittierende Diodenvorrichtung, umfassend eine erste Elektrode, eine gegenüber der ersten Elektrode angeordnete zweite Elektrode und eine zwischen der ersten Elektrode und der zweiten Elektrode angeordnete organische Schicht, wobei die organische Schicht mindestens eine der aminbasierten Verbindungen nach den


Ansprüchen 1 bis 18 umfasst.

- 5
20. Organische lichtemittierende Diodenvorrichtung nach Anspruch 19, wobei die organische Schicht mindestens eine von einer Lochinjektionsschicht, einer Lochtransportschicht, einer funktionellen Schicht mit sowohl Lochinjektions- als auch Lochtransportvermögen, einer Pufferschicht, einer Elektronensperrschicht, einer Emissionsschicht, einer Lochsperrschicht, einer Elektronentransportschicht, einer Elektroneninjectionsschicht und einer funktionellen Schicht mit sowohl Elektroneninjektions- als auch Elektronentransportvermögen umfasst.
- 10
21. Organische lichtemittierende Diodenvorrichtung nach Anspruch 20, wobei die organische Schicht eine Elektronentransportschicht umfasst und die aminbasierte Verbindung in der Elektronentransportschicht enthalten ist, vorzugsweise wobei die Elektronentransportschicht weiterhin einen Metallkomplex umfasst.
- 15
22. Organische lichtemittierende Diodenvorrichtung nach Anspruch 20, wobei die organische Schicht eine Emissionsschicht umfasst und die aminbasierte Verbindung in der Emissionsschicht enthalten ist, vorzugsweise wobei die aminbasierte Verbindung in der Emissionsschicht als Hostmaterial fungiert und die Emissionsschicht weiterhin einen blau fluoreszierenden Dotanden umfasst.
- 20
23. Organische lichtemittierende Diodenvorrichtung nach Anspruch 22, wobei die aminbasierte Verbindung in der Emissionsschicht als Dotand fungiert und die Emissionsschicht weiterhin mindestens eine einer durch nachstehende Formel 400 dargestellten Verbindung auf Anthracenbasis und einer durch nachstehende Formel 401 dargestellten Verbindung auf Anthracenbasis umfasst:

Formel 400



Formel 401



wobei, in den Formeln 400 und 401, Ar_{111} und Ar_{112} jeweils unabhängig voneinander ein substituierter oder unsubstituierter C_6-C_{60} -Arylenrest sind; Ar_{113} bis Ar_{116} und Ar_{122} bis Ar_{125} jeweils unabhängig voneinander ein substituierter oder unsubstituierter C_1-C_{10} -Alkylrest oder ein substituierter oder unsubstituierter C_6-C_{60} -Arylrest sind; Ar_{126}


und Ar₁₂₇ jeweils unabhängig voneinander ein C₁-C₁₀-Alkylrest sind; und g, h, i, j, k und l jeweils unabhängig voneinander eine ganze Zahl von 0 bis 4 sind.

24. Organische lichtemittierende Diodenvorrichtung nach Anspruch 20, wobei die organische Schicht mindestens eine von einer Lochinjektionsschicht, einer Lochtransportschicht und einer funktionellen Schicht mit sowohl Lochinjektions- als auch Lochtransportvermögen umfasst und mindestens eine von der Lochinjektionsschicht, der Lochtransportschicht und der funktionellen Schicht mit sowohl Lochinjektions- als auch Lochtransportvermögen eine p-Dotierung umfasst.

Revendications

1. Composé à base d'amine représenté par la Formule 1 ci-dessous :

Formule 1



dans lequel, dans la Formule 1, Ar₁ et Ar₂ sont chacun indépendamment un groupe aryle en C₆ à C₆₀ substitué ou non ou un groupe hétéroaryle en C₂ à C₆₀ substitué ou non ; et dans lequel Ar₁ et Ar₂ sont éventuellement liés par une liaison simple ;

X₁ est un groupe arylène en C₆ à C₆₀ substitué ou non ou un groupe hétéroarylène en C₂ à C₆₀ substitué ou non ; m est un nombre entier de 1 à 5 ; et


au moins un substituant de chacun du groupe aryle en C₆ à C₆₀ substitué, du groupe hétéroaryle en C₂ à C₆₀ substitué, du groupe arylène en C₆ à C₆₀ substitué, et du groupe hétéroarylène en C₂ à C₆₀ substitué est l'un parmi un atome de deutérium ; -F ; -Cl ; -Br ; -I ; -CN ; un groupe hydroxyle ; -NO₂ ; un groupe amino ; un groupe amidino ; une hydrazine ; une hydrazone ; un groupe carboxyle ou un de ses sels ; un groupe acide sulfonique ou un de ses sels ; un acide phosphorique ou un de ses sels ; un groupe triaryl (C₆ à C₆₀) silyle ; un groupe alkyle en C₁ à C₆₀, un groupe alkoxy en C₁ à C₆₀, un groupe alcényle en C₂ à C₆₀, un groupe alcynyle en C₂ à C₆₀ ; un groupe alkyle en C₁ à C₆₀, un groupe alkoxy en C₁ à C₆₀, un groupe alcényle en C₂ à C₆₀ et un groupe alcynyle en C₂ à C₆₀ qui est substitué par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels ; un groupe cycloalkyle en C₃ à C₆₀, un groupe cycloalcényle en C₃ à C₆₀, un groupe aryle en C₆ à C₆₀, un groupe hétéroaryle en C₂ à C₆₀, un groupe aralkyle en C₆ à C₆₀, un groupe aryloxy en C₆ à C₆₀, un groupe arylthio en C₆ à C₆₀ ; un groupe cycloalkyle en C₃ à C₆₀, un groupe cycloalcényle en C₃ à C₆₀, un groupe aryle en C₆ à C₆₀, un groupe hétéroaryle en C₂ à C₆₀, un groupe aralkyle en C₆ à C₆₀, un groupe aryloxy en C₆ à C₆₀ et un groupe arylthio en C₆ à C₆₀ qui est substitué par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₆₀, un groupe alkyle en C₁ à C₆₀ substitué par au moins un atome de fluor (F), un groupe alkoxy en C₁ à C₆₀, un groupe alcényle en C₂ à C₆₀, un groupe alcynyle en C₂ à C₆₀, un groupe aryle en C₆ à C₆₀ et un groupe hétéroaryle en C₂ à C₆₀,

dans lequel au moins un des Ar₁ et Ar₂ est un groupe aryle en C₆ à C₆₀ substitué par au moins un groupe attracteur d'électrons choisi dans le groupe constitué par -F ; -CN ; -NO₂ ; un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F ; un groupe hétéroaryle en C₂ à C₆₀ ; et un groupe hétéroaryle en C₂ à C₆₀ substitué par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino,


un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels ; un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₆₀, un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₆₀, un groupe alcényle en C₂ à C₆₀, un groupe alcyne en C₂ à C₆₀, un groupe aryle en C₆ à C₆₀, et un groupe hétéroaryle en C₂ à C₆₀.

2. Composé à base d'amine selon la revendication 1, dans lequel ledit au moins groupe attracteur d'électrons est choisi dans le groupe constitué par : -F ; -CN ; -NO₂ ; un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F ; un groupe hétéroaryle en C₂ à C₂₀ comprenant un noyau contenant un atome N ; et un groupe hétéroaryle en C₂ à C₂₀ qui comprend un noyau contenant un atome N et est substitué par un moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe aryle en C₆ à C₂₀, et un groupe hétéroaryle en C₂ à C₂.


3. Composé à base d'amine selon la revendication 1 ou la revendication 2, dans lequel ledit au moins un groupe attracteur d'électrons est choisi dans le groupe constitué par -F ; -CN ; -CH₂F ; -CHF₂ ; -CF₃ ; et les groupes représentés par les Formules 2(1) à 2(14) ci-dessous :




Formule 2(1)




Formule 2(2)




Formule 2(3)




Formule 2(4)




Formule 2(5)




Formule 2(6)




Formule 2(7)




Formule 2(8)






Formule 2(9)



Formule 2(10)



Formule 2(11)



15 dans lequel, dans les Formules 2(1) à 2(14), Z₁₁ à Z₁₈ sont chacun indépendamment un atome d'hydrogène, un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, un groupe pyridinyle, un groupe triazinyle, ou un groupe carbazolyle.

20 4. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel Ar₁ et Ar₂ sont chacun indépendamment un groupe aryle en C₆ à C₆₀ choisi parmi un groupe phényle substitué ou non, un groupe naphthyle substitué ou non, un groupe fluorényle substitué ou non, un groupe phénanthrényle substitué ou non, un groupe anthryle substitué ou non, un groupe triphénylényle substitué ou non, un groupe pyrényle substitué ou non, un groupe chrysényle substitué ou non, un groupe pyridinyle substitué ou non, un groupe pyrazinyle substitué ou non, un groupe pyrimidinyle substitué ou non, un groupe quinolyle substitué ou non, un groupe carbazolyle substitué ou non, un groupe triazinyle substitué ou non, un groupe dibenzothiophényle substitué ou non, un groupe dibenzofurannyle substitué ou non, ou un groupe phénanthrolinyle substitué ou non.

25 5. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel Ar₁ et Ar₂ sont liés par une liaison simple.

30 6. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel ledit au moins Ar₁ et Ar₂ est un groupe aryle en C₆ à C₆₀ substitué par au moins deux groupes attracteurs d'électrons.


35 7. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel ledit au moins Ar₁ et Ar₂ est un groupe phényle, un groupe biphényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, ou un groupe fluorényle qui est substitué par au moins deux groupes attracteurs d'électrons ; et

40 les groupes attracteurs d'électrons sont chacun indépendamment choisis dans le groupe constitué par un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinoxalinyle, un groupe triazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle ; et un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinoxalinyle, un groupe phtalazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN ; un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, un groupe pyridinyle, un groupe triazinyle, et un groupe carbazolyle.


45 8. Composé à base d'amine selon la revendication 1, dans lequel le composé à base d'amine est représenté par la Formule 1(1) ou 1(2) ci-dessous :

50

Formule 1(1)



Formule 1(2)



dans lequel, dans la Formule 1(1), Ar₂ est un groupe aryle en C₆ à C₂₀ substitué ou non ou un groupe hétéroaryle en C₂ à C₂₀ substitué ou non ; et

dans les Formules 1(1) et 1(2), les groupes à noyau A et B sont chacun indépendamment un groupe aryle en C₆ à C₂₀ substitué ;

R₁ et R₂ sont chacun indépendamment un groupe attracteur d'électrons choisi dans le groupe constitué par -F ; -CN ; -NO₂ ; un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F ; un groupe hétéroaryle en C₂ à C₆₀ ; et un groupe hétéroaryle en C₂ à C₆₀ substitué par un moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₆₀, un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₆₀, un groupe alcényle en C₂ à C₆₀, un groupe alcynyle en C₂ à C₆₀, un groupe aryle en C₆ à C₆₀, et un groupe hétéroaryle en C₂ à C₆₀ ; et

p et q sont chacun indépendamment un nombre entier de 1 à 9.

9. Composé à base d'amine selon la revendication 8, dans lequel le composé à base d'amine est représenté par la Formule 1(1) dans laquelle au moins un de R₁ est -CN ; ou est représenté par la Formule 1(2) dans laquelle au moins un de R₁ et un de R₂ est -CN.

10. Composé à base d'amine selon la revendication 7, dans lequel le composé à base d'amine est représenté par la Formule 1(1), dans laquelle le groupe à noyau A est un groupe phényle substitué, un groupe biphenyle substitué, un groupe naphthyle substitué, un groupe anthryle substitué, un groupe phénanthrényle substitué, un groupe pyrényle substitué, ou un groupe florényle substitué ;


R₁ est au moins un groupe attracteur d'électrons choisi dans le groupe constitué par un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinazolinyne, un groupe triazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle ; et un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinazolinyne, un groupe phtalazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN ; un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, un groupe pyridinyle, un groupe triazinyle, et un groupe carbazolyle ; et p est 2, 3, ou 4 ; ou, dans lequel le composé à base d'amine est représenté par la Formule 1(2), dans laquelle le groupe à noyau A et le groupe à noyau B sont chacun indépendamment un groupe phényle substitué, un groupe biphenyle substitué, un groupe naphthyle substitué, un groupe anthryle substitué, un groupe phénanthrényle substitué, un groupe pyrényle substitué, ou un groupe florényle substitué ;

R₁ et R₂ sont chacun indépendamment au moins un groupe attracteur d'électrons choisi dans le groupe constitué


par un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinazolinyne, un groupe triazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle ; et un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinazolinyne, un groupe phtalazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN ; un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, un groupe pyridinyle, un groupe triazinyle, et un groupe carbazolyle ; et p et q sont chacun indépendamment 2, 3, ou 4.

11. Composé à base d'amine selon la revendication 1, dans lequel le composé à base d'amine est représenté par une des Formules 1A à 1J ci-dessous :


Formule 1A




Formule 1B




Formule 1C




Formule 1D




Formule 1E




Formule 1F




Formule 1G




Formule 1H



Formule 1I



Formule 1J



30

35

40

45

50

dans lequel, dans les Formules 1A à 1J, Ar₂ est un groupe phényle substitué ou non, un groupe pentalényle substitué ou non, un groupe indényle substitué ou non, un groupe naphthyle substitué ou non, un groupe azulényle substitué ou non, un groupe heptalényle substitué ou non, un groupe indacényle substitué ou non, un groupe acénaphthyle substitué ou non, un groupe fluorényle substitué ou non, un groupe phénalényle substitué ou non, un groupe phénanthrényle substitué ou non, un groupe anthryle substitué ou non, un groupe fluoranthényle substitué ou non, un groupe triphénylényle substitué ou non, un groupe pyrényle substitué ou non, un groupe chrysényle substitué ou non, un groupe naphtacényle substitué ou non, un groupe picényle substitué ou non, un groupe pérylényle substitué ou non, un groupe pentaphényle substitué ou non, un groupe hexacényle substitué ou non, un groupe pyrrolyle substitué ou non, un groupe pyrazolyle substitué ou non, un groupe imidazolyle substitué ou non, un groupe imidazolynyle substitué ou non, un groupe imidazopyridinyle substitué ou non, un groupe imidazopyrimidinyle substitué ou non, un groupe pyridinyle substitué ou non, un groupe pyrazinyle substitué ou non, un groupe pyrimidinyle substitué ou non, un groupe benzoimidazolyle substitué ou non, un groupe indolyle substitué ou non, un groupe purinyle substitué ou non, un groupe quinolinyle substitué ou non, un groupe phtalazinyle substitué ou non, un groupe indolizynyle substitué ou non, un groupe naphtyridinyle substitué ou non, un groupe quinazolynyle substitué ou non, un groupe cinolinyle substitué ou non, un groupe indazolyle substitué ou non, un groupe carbazolyle substitué ou non, un groupe phénazinyle substitué ou non, un groupe phénanthridinyle substitué ou non, un groupe pyranyle substitué ou non, un groupe chroményle substitué ou non, un groupe furannyle substitué ou non, un groupe benzofurannyle substitué ou non, un groupe thiophényle substitué ou non, un groupe benzothiophényle substitué ou non, un groupe isothiazolyle substitué ou non, un groupe benzoimidazolyle substitué ou non, un groupe isoxazolyle substitué ou non, un groupe dibenzothiophényle substitué ou non, un groupe dibenzofurannyle substitué ou non, un groupe triazinyle substitué ou non, un groupe oxadiazolyle substitué ou non, un groupe pyridazinyle substitué ou non, un groupe triazolyle substitué ou non, un groupe tétrazolyle substitué ou non, ou un groupe phénanthrolinyle substitué ou non,

55


les substituants du groupe phényle substitué, du groupe pentalényle substitué, du groupe indényle substitué, du groupe naphthyle substitué, du groupe azulényle substitué, du groupe heptalényle substitué, du groupe indacényle substitué, du groupe acénaphthyle substitué, du groupe fluorényle substitué, du groupe phénalényle substitué, du groupe phénanthrényle substitué, du groupe anthryle substitué, du groupe fluoranthényle substitué, du groupe triphénylényle substitué, du groupe pyrényle substitué, du groupe chrysényle substitué, du groupe naphtacényle substitué, du groupe picényle substitué, du groupe pérylényle substitué, du groupe pentaphényle substitué, du groupe hexacényle substitué, du groupe pyrrolyle substitué, du groupe pyrazolyle substitué, du groupe imidazolyle substitué, du groupe imidazolynyle substitué, du groupe imidazopyridinyle substitué, du groupe imidazopyrimidinyle

substitué, du groupe pyridinyle substitué, du groupe pyrazinyle substitué, du groupe pyrimidinyle substitué, du groupe benzoimidazole substitué, du groupe indolyle substitué, du groupe purinyle substitué, du groupe quinolinyle substitué, du groupe phtalazinyle substitué, du groupe indolizinyne substitué, du groupe naphthyridinyle substitué, du groupe quinazolinyne substitué, du groupe cinolinyle substitué, du groupe indazolyle substitué, du groupe carbazolyle substitué, du groupe phénazinyle substitué, du groupe phénanthridinyle substitué, du groupe pyranyle substitué, du groupe chroményne substitué, du groupe furannyle substitué, du groupe benzofurannyle substitué, du groupe thiophényne substitué, du groupe benzothiophényne substitué, du groupe isothiazolyle substitué, du groupe benzoimidazolyle substitué, du groupe isoxazolyle substitué, du groupe dibenzothiophényne substitué, du groupe dibenzofurannyle substitué, du groupe triazinyle substitué, du groupe oxadiazolyle substitué, du groupe pyridazinyle substitué, du groupe triazolyle substitué, du groupe tétrazolyle substitué, et du groupe phénanthrolinyle substitué, et R_{11} à R_{19} sont chacun indépendamment un atome d'hydrogène ; un atome de deutérium ; -F ; -Cl ; -Br ; -I ; -CN ; un groupe hydroxyle ; -NO₂ ; un groupe amino ; un groupe amidino ; une hydrazine ; une hydrazone ; un groupe carboxyle ou un de ses sels ; un groupe acide sulfonique ou un de ses sels ; un acide phosphorique ou un de ses sels ; un groupe triaryl (C₆ à C₆₀) silyle ; un groupe alkyle en C₁ à C₆₀, un groupe alkoxy en C₁ à C₆₀, un groupe alcényne en C₂ à C₆₀, et un groupe alcynyle en C₂ à C₆₀ ; un groupe alkyle en C₁ à C₆₀, un groupe alkoxy en C₁ à C₆₀, un groupe alcényne en C₂ à C₆₀ et un groupe alcynyle en C₂ à C₆₀ qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, et un acide phosphorique ou un de ses sels ; un groupe cycloalkyle en C₃ à C₆₀, un groupe cycloalcényne en C₃ à C₆₀, un groupe aryle en C₆ à C₆₀, un groupe hétéroaryle en C₂ à C₆₀, un groupe aralkyle en C₆ à C₆₀, un groupe aryloxy en C₆ à C₆₀, un groupe arylthio en C₆ à C₆₀ ; un groupe cycloalkyle en C₃ à C₆₀, un groupe cycloalcényne en C₃ à C₆₀, un groupe aryle en C₆ à C₆₀, un groupe hétéroaryle en C₂ à C₆₀, un groupe aralkyle en C₆ à C₆₀, un groupe aryloxy en C₆ à C₆₀, et un groupe arylthio en C₆ à C₆₀ qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₆₀, un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₆₀, un groupe alcényne en C₂ à C₆₀, un groupe alcynyle en C₂ à C₆₀, un groupe aryle en C₆ à C₆₀ et un groupe hétéroaryle en C₂ à C₆₀.


dans lequel au moins un des R_{11} à R_{15} dans les Formules 1A et 1B, au moins un des R_{11} à R_{17} dans les Formules 1C et 1D, au moins un des R_{11} à R_{18} dans les Formules 1E et 1J, et au moins un des R_{11} à R_{19} dans la Formule 1F, 1G, 1H et 1I est indépendamment un groupe attracteur d'électrons choisi dans le groupe constitué par -F ; -CN ; -NO₂ ; un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F ; un groupe hétéroaryle en C₂ à C₆₀ ; et un groupe hétéroaryle en C₂ à C₆₀ substitué par un moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₆₀, un groupe alkyle en C₁ à C₆₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₆₀, un groupe alcényne en C₂ à C₆₀, un groupe alcynyle en C₂ à C₆₀, un groupe aryle en C₆ à C₆₀, et un groupe hétéroaryle en C₂ à C₆₀.

12. Composé à base d'amine selon la revendication 11, dans lequel le composé à base d'amine est représenté par la Formule 1A-(1) ou 1A-(2) ci-dessous :

Formule 1A-(1)




Formule 1A-(2)




dans lequel, dans les Formules 1A-(1) et 1A-(2), R_{12} , R_{14} , R_{22} , et R_{24} sont chacun indépendamment un groupe attracteur d'électrons choisi dans le groupe constitué par un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinazolinyne, un groupe triazinyle, un

groupe benzoimidazolyle, et un groupe carbazolyle ; et un groupe pyridinyle, un groupe pyrazinyle, un groupe pyrimidinyle, un groupe quinolinyle, un groupe isoquinolinyle, un groupe quinazolinyle, un groupe phtalazinyle, un groupe benzoimidazolyle, et un groupe carbazolyle qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN ; un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, un groupe pyridinyle, un groupe triazinyle, et un groupe carbazolyle ; et dans la Formule 1A-(1), Ar₂ est un groupe phényle substitué ou non, un groupe biphenyle substitué ou non, un groupe naphthyle substitué ou non, un groupe anthryle substitué ou non, un groupe phénanthrényle substitué ou non, un groupe pyrényle substitué ou non, ou un groupe fluorényle substitué ou non.


13. Composé à base d'amine selon la revendication 12, dans lequel R₁₂, R₁₄, R₂₂ et R₂₄ sont chacun indépendamment choisis dans le groupe constitué par -F ; -CN ; -CH₂F ; -CHF₂ ; -CF₃ ; et les groupes représentés par les Formules 2(1) à 2(14) ci-dessous :




Formule 2(1)




Formule 2(2)




Formule 2(3)




Formule 2(4)




Formule 2(5)




Formule 2(6)




Formule 2(7)




Formule 2(8)






Formule 2(9)



Formule 2(10)



Formule 2(11)



15 dans lequel, dans les Formules 2(1) à 2(14), Z₁₁ à Z₁₈ sont chacun indépendamment un atome d'hydrogène, un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkyle en C₁ à C₂₀ substitué par au moins un -F, un groupe alkoxy en C₁ à C₂₀, un groupe phényle, un groupe naphyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, un groupe pyridinyle, un groupe triazinyle, ou un groupe carbazolyne.

20

14. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel X₁ est un groupe phénylène substitué ou non, un groupe pentalénylène substitué ou non, un groupe indénylène substitué ou non, un groupe naphtylène substitué ou non, un groupe azulénylène substitué ou non, un groupe heptalénylène substitué ou non, un groupe indacénylène substitué ou non, un groupe acénaphtylène substitué ou non, un groupe fluorénylène substitué ou non, un groupe phénalénylène substitué ou non, un groupe phénanthrényle substitué ou non, un groupe anthrylène substitué ou non, un groupe fluoranthénylène substitué ou non, un groupe triphénylényle substitué ou non, un groupe pyrényle substitué ou non, un groupe chrysénylène substitué ou non, un groupe naphtacénylène substitué ou non, un groupe picénylène substitué ou non, un groupe pérylényle substitué ou non, un groupe pentaphénylène substitué ou non, un groupe hexacénylène substitué ou non, un groupe pyrrolylène substitué ou non, un groupe pyrazolylène substitué ou non, un groupe imidazolylène substitué ou non, un groupe imidazolinyène substitué ou non, un groupe imidazopyridinyène substitué ou non, un groupe imidazopyrimidinyène substitué ou non, un groupe pyridinyène substitué ou non, un groupe pyrazinyène substitué ou non, un groupe pyrimidinyène substitué ou non, un groupe indolylène substitué ou non, un groupe purinyène substitué ou non, un groupe quinolinyène substitué ou non, un groupe phtalazinyène substitué ou non, un groupe indolizinyène substitué ou non, un groupe naphtyridinyène substitué ou non, un groupe quinazolinyène substitué ou non, un groupe cinnolinyène substitué ou non, un groupe indazolylène substitué ou non, un groupe carbazolylène substitué ou non, un groupe phénazinyène substitué ou non, un groupe phénanthridinyène substitué ou non, un groupe pyranlyène substitué ou non, un groupe chroménylène substitué ou non, un groupe furanylène substitué ou non, un groupe benzofuranylène substitué ou non, un groupe thiophénylène substitué ou non, un groupe benzothiophénylène substitué ou non, un groupe isothiazolylène substitué ou non, un groupe benzoimidazolylène substitué ou non, un groupe isoxazolylène substitué ou non, un groupe dibenzothiophénylène substitué ou non, un groupe dibenzofuranylène substitué ou non, un groupe triazinylène substitué ou non, un groupe oxadiazolylène substitué ou non, un groupe pyridazinyène substitué ou non, un groupe triazolylène substitué ou non, ou un groupe tétrazolylène substitué ou non.

25

30

35

40


45

15. Composé à base d'amine selon la revendication 14, dans lequel X₁ est un groupe représenté par une des Formules 5(1) à 5(16) ci-dessous :


50

55


5




Formule 5(1)



Formule 5(2)




Formule 5(3)




Formule 5(4)

10


15




Formule 5(5)



Formule 5(6)



Formule 5(7)




Formule 5(8)


20

25


30



Formule 5(9)




Formule 5(10)




Formule 5(11)

35


40



Formule 5(12)



Formule 5(13)




Formule 5(14)

45


50

55

5



Formule 5(15)



Formule 5(16)

10

15

20

25

dans lequel, dans les Formules 5(1) à 5(16), Z_1 à Z_8 sont chacun indépendamment l'un parmi un atome d'hydrogène ; un atome de deutérium ; -F ; -Cl ; -Br ; -I ; -CN ; un groupe hydroxyle ; -NO₂ ; un groupe amino ; un groupe amidino ; une hydrazine ; une hydrazone ; un groupe carboxyle ou un de ses sels ; un groupe acide sulfonique ou un de ses sels ; un acide phosphorique ou un de ses sels ; un groupe alkyle en C₁ à C₂₀ ; un groupe alkoxy en C₁ à C₂₀ ; un groupe aryle en C₆ à C₂₀ et un groupe hétéroaryle en C₂ à C₂₀ qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, et un acide phosphorique ou un de ses sels ; un groupe aryle en C₆ à C₂₀ ; un groupe hétéroaryle en C₂ à C₂₀ ; et un groupe aryle en C₆ à C₂₀ et un groupe hétéroaryle en C₂ à C₂₀ qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, un groupe alkoxy en C₁ à C₂₀, un groupe aryle en C₆ à C₆₀ et un groupe hétéroaryle en C₂ à C₂₀ ;

dans lequel * indique un site de liaison à l'antracène dans la Formule 1 ; et

dans lequel *† indique un site de liaison à N dans la Formule 1.

30

16. Composé à base d'amine selon la revendication 15, dans lequel Z_1 à Z_8 sont chacun indépendamment l'un parmi un atome d'hydrogène ; un atome de deutérium ; -F ; -Cl ; -Br ; -I ; -CN ; un groupe hydroxyle ; -NO₂ ; un groupe amino ; un groupe amidino ; une hydrazine ; une hydrazone ; un groupe carboxyle ou un de ses sels ; un groupe acide sulfonique ou un de ses sels ; un acide phosphorique ou un de ses sels ; un groupe méthyle, un groupe éthyle, un groupe propyle, un groupe butyle, et un groupe pentyle ; un groupe méthoxy, un groupe éthoxy, un groupe propoxy, un groupe butoxy, et un groupe pentoxy ; un groupe méthyle, un groupe éthyle, un groupe propyle, un groupe butyle, et un groupe pentyle, un groupe méthoxy, un groupe éthoxy, un groupe propoxy, un groupe butoxy, et un groupe pentoxy qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, et un acide phosphorique ou un de ses sels ; un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, et un groupe fluorényle ; un groupe pyridinyle, un groupe pyrimidinyle, un groupe triazinyle, un groupe quinolyne, et un groupe carbazolyle ; un groupe phényle, un groupe naphthyle, un groupe anthryle, un groupe phénanthrényle, un groupe pyrényle, et un groupe fluorényle ; un groupe pyridinyle, un groupe pyrimidinyle, un groupe triazinyle, un groupe quinolyne, et un groupe carbazolyle qui sont substitués par au moins l'un parmi un atome de deutérium, -F, -Cl, -Br, -I, -CN, un groupe hydroxyle, -NO₂, un groupe amino, un groupe amidino, une hydrazine, une hydrazone, un groupe carboxyle ou un de ses sels, un groupe acide sulfonique ou un de ses sels, un acide phosphorique ou un de ses sels, un groupe alkyle en C₁ à C₂₀, et un groupe alkoxy en C₁ à C₂₀.

40

45


50

17. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel m est 1, 2, ou 3.


18. Composé à base d'amine selon l'une quelconque des revendications précédentes, dans lequel le composé à base d'amine est un des composés 1 à 109 ci-dessous :

55


5



1




2




3


10



4




5




6


20



7




8




9


30



10




11




12


40



13




14




15

45


50



16




17




18

55


5
10
15
20
25
30
35
40
45
50
55




19




20




21




22




23




24




25




26




27



28




29




30


5



31




32




33


10



34




35




36

15


20



37




38




39

25


30



40




41




42

35


40



43



44




45

45


50

55


5



46




47




48


10



49




50




51

15


20



52




53




54

25


30



55




56




57

35

40



58




59

45


50

55


5




10




15




20




25




30




35




40



45




50




55

5




70




71

10




72




73

15

20




74




75

25

30



76




77


35

40

45



78




79

50


55

5




80

10




81

15




82

20




83

25




84

30




85

35




86

40




87

45



88


50




89

55


5
10
15
20
25
30
35
40
45
50
55




90




91




92




93




94




95




96




97



98



99



19. Diode électroluminescente organique comprenant une première électrode, une seconde électrode placée à l'opposé de la première, et une couche organique placée entre la première et la seconde électrode, la couche organique comprenant au moins un des composés à base d'amine selon les revendications 1 à 18.

20. Diode électroluminescente organique selon la revendication 19, dans laquelle la couche organique comprend au moins l'une parmi une couche d'injection de trous, une couche de transport de trous, une couche fonctionnelle ayant à la fois des possibilités d'injection de trous et de transport de trous, une couche tampon, une couche de blocage d'électrons, une couche d'émission, une couche de blocage de trous, une couche de transport d'électrons, une couche d'injection d'électrons, et une couche fonctionnelle ayant à la fois des possibilités d'injection d'électrons et de transport d'électrons.


21. Diode électroluminescente organique selon la revendication 20, dans laquelle la couche organique comprend une couche de transport d'électrons, et le composé à base d'amine est compris dans la couche de transport d'électrons, de préférence, dans laquelle la couche de transport d'électrons comprend en outre un complexe métallique.

22. Diode électroluminescente organique selon la revendication 20, dans laquelle la couche organique comprend une


couche d'émission, et le composé à base d'amine est compris dans la couche d'émission, de préférence, dans laquelle le composé à base d'amine dans la couche d'émission joue le rôle d'hôte, et la couche d'émission comprend en outre un dopant fluorescent bleu.

- 5 **23.** Diode électroluminescente organique selon la revendication 22, dans laquelle le composé à base d'amine dans la couche d'émission joue le rôle de dopant, et la couche d'émission comprend en outre au moins l'un parmi un composé à base d'anthracène représenté par la Formule 400 ci-dessous et un composé à base d'anthracène représenté par la Formule 401 ci-dessous :

10 Formule 400



25 Formule 401



40 dans laquelle, dans les Formules 400 et 401, Ar₁₁₁ et Ar₁₁₂ sont chacun indépendamment un groupe arylène en C₆ à C₆₀ substitué ou non ; Ar₁₁₃ à Ar₁₁₆ et Ar₁₂₂ à Ar₁₂₅ sont chacun indépendamment un groupe alkyle en C₁ à C₁₀ substitué ou non, ou un groupe aryle en C₆ à C₆₀ substitué ou non ; Ar₁₂₆ et Ar₁₂₇ sont chacun indépendamment un groupe alkyle en C₁ à C₁₀ ; et g, h, i, j, k, et l sont chacun indépendamment un nombre entier de 0 à 4.

- 45 **24.** Diode électroluminescente organique selon la revendication 20, dans laquelle la couche organique comprend au moins l'une parmi une couche d'injection de trous, une couche de transport de trous, et une couche fonctionnelle ayant à la fois des possibilités d'injection de trous et de transport de trous, et l'une au moins parmi la couche d'injection de trous, la couche de transport de trous, et la couche fonctionnelle ayant à la fois des possibilités d'injection de trous et de transport de trous comprend un dopant p.

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- Synthesis and electroluminescent properties of blue fluorescent triphenylamine substituted anthracene derivatives for OLEDs. *Molecular Crystals and Liquid Crystals*, 11 October 2010, vol. 530 (1 [0006])

专利名称(译)	胺类化合物和包括其的有机发光二极管		
公开(公告)号	EP2626399B1	公开(公告)日	2014-12-03
申请号	EP2013154326	申请日	2013-02-07
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	KIM YOUNG KOOK HWANG SEOK HWAN JUNG HYE JIN LIM JIN O HAN SANG HYUN JEONG EUN JAE KIM SOO YON PARK JUN HA LEE EUN YOUNG LEE JONG HYUK		
发明人	KIM, YOUNG-KOOK HWANG, SEOK-HWAN JUNG, HYE-JIN LIM, JIN-O HAN, SANG-HYUN JEONG, EUN-JAE KIM, SOO-YON PARK, JUN-HA LEE, EUN-YOUNG LEE, JONG-HYUK		
IPC分类号	C09K11/06 C07C15/28 H01L51/00 H05B33/10		
CPC分类号	C09K11/06 C07C211/56 C07C211/59 C07C211/61 C07C217/92 C07C255/58 C07C2603/18 C07C2603/24 C07C2603/26 C07C2603/50 C07D209/88 C07D213/36 C07D213/74 C07D307/91 C07D333/76 C07F7/0805 C09K2211/1007 C09K2211/1011 C09K2211/1014 H01L51/0058 H01L51/006 H01L51/0081 H05B33/10		
代理机构(译)	RUSSELL , TIM		
优先权	1020120012532 2012-02-07 KR 1020120061676 2012-06-08 KR		
其他公开文献	EP2626399A1		
外部链接	Espacenet		

摘要(译)

胺类化合物和包含胺类化合物的有机发光二极管。

Formula 1

