

(19)



(11)

EP 2 966 706 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
04.09.2019 Bulletin 2019/36

(51) Int Cl.:
H01L 51/54^(2006.01)

(21) Application number: **15175686.3**

(22) Date of filing: **07.07.2015**

(54) ORGANIC ELECTROLUMINESCENT MATERIALS AND DEVICES

ORGANISCHE ELEKTROLUMINESZENDE MATERIALIEN UND VORRICHTUNGEN

MATÉRIAUX ET DISPOSITIFS ÉLECTROLUMINESCENTS ORGANIQUES

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **09.07.2014 US 201462022300 P**
19.08.2014 US 201462038925 P
06.10.2014 US 201462060192 P
24.11.2014 US 201462083490 P
09.06.2015 US 201514734712

(43) Date of publication of application:
13.01.2016 Bulletin 2016/02

(73) Proprietor: **Universal Display Corporation**
Ewing, NJ 08618 (US)

(72) Inventors:

- **ZENG, Lichang**
Ewing, NJ 08618 (US)
- **LAYEK, Suman**
Ewing, NJ 08618 (US)
- **BOUDREAU, Pierre-Luc T.**
Ewing, NJ 08618 (US)
- **ELSHENAWY, Zeinab**
Ewing, NJ 08618 (US)
- **KOTTAS, Gregg**
Ewing, NJ 08618 (US)
- **DYATKIN, Alexey Borisovich**
Ewing, NJ 08618 (US)

- **JOSEPH, Scott**
Ewing, NJ 08618 (US)
- **ADAMOVICH, Vadim**
Ewing, NJ 08618 (US)
- **XIA, Chuanjun**
Ewing, NJ 08618 (US)
- **WANG, Ting-Chih**
Ewing, NJ 08618 (US)
- **YEAGER, Walter**
Ewing, NJ 08618 (US)

(74) Representative: **Hansen, Norbert**
Maiwald Patentanwalts- und
Rechtsanwaltsgesellschaft mbH
Elisenhof
Elisenstraße 3
80335 München (DE)

(56) References cited:
WO-A1-2013/032297 WO-A1-2013/077362
CN-A- 102 850 329 JP-A- 2004 022 334
JP-A- 2014 125 449 US-A1- 2012 126 208
US-A1- 2014 001 456

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

EP 2 966 706 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

FIELD OF THE INVENTION

5 **[0001]** The present invention relates to compounds for use as hosts, blocking materials, and electron transporting materials, and devices, such as organic light emitting diodes, including the same.


BACKGROUND

10 **[0002]** Opto-electronic devices that make use of organic materials are becoming increasingly desirable for a number of reasons. Many of the materials used to make such devices are relatively inexpensive, so organic opto-electronic devices have the potential for cost advantages over inorganic devices. In addition, the inherent properties of organic materials, such as their flexibility, may make them well suited for particular applications such as fabrication on a flexible substrate. Examples of organic opto-electronic devices include organic light emitting devices (OLEDs), organic photo-transistors, organic photovoltaic cells, and organic photodetectors. For OLEDs, the organic materials may have performance advantages over conventional materials. For example, the wavelength at which an organic emissive layer emits light may generally be readily tuned with appropriate dopants.

15 **[0003]** OLEDs make use of thin organic films that emit light when voltage is applied across the device. OLEDs are becoming an increasingly interesting technology for use in applications such as flat panel displays, illumination, and backlighting. Several OLED materials and configurations are described in U.S. Pat. Nos. 5,844,363, 6,303,238, and 5,707,745.

20 **[0004]** One application for phosphorescent emissive molecules is a full color display. Industry standards for such a display call for pixels adapted to emit particular colors, referred to as "saturated" colors. In particular, these standards call for saturated red, green, and blue pixels. Color may be measured using CIE coordinates, which are well known to the art.

25 **[0005]** One example of a green emissive molecule is tris(2-phenylpyridine) iridium, denoted Ir(ppy)₃, which has the following structure:



40 **[0006]** In this, and later figures herein, we depict the dative bond from nitrogen to metal (here, Ir) as a straight line.

[0007] As used herein, the term "organic" includes polymeric materials as well as small molecule organic materials that may be used to fabricate organic opto-electronic devices. "Small molecule" refers to any organic material that is not a polymer, and "small molecules" may actually be quite large. Small molecules may include repeat units in some circumstances. For example, using a long chain alkyl group as a substituent does not remove a molecule from the "small molecule" class. Small molecules may also be incorporated into polymers, for example as a pendent group on a polymer backbone or as a part of the backbone. Small molecules may also serve as the core moiety of a dendrimer, which consists of a series of chemical shells built on the core moiety. The core moiety of a dendrimer may be a fluorescent or phosphorescent small molecule emitter. A dendrimer may be a "small molecule," and it is believed that all dendrimers currently used in the field of OLEDs are small molecules.

45 **[0008]** As used herein, "top" means furthest away from the substrate, while "bottom" means closest to the substrate. Where a first layer is described as "disposed over" a second layer, the first layer is disposed further away from substrate. There may be other layers between the first and second layer, unless it is specified that the first layer is "in contact with" the second layer. For example, a cathode may be described as "disposed over" an anode, even though there are various organic layers in between.

50 **[0009]** As used herein, "solution processible" means capable of being dissolved, dispersed, or transported in and/or deposited from a liquid medium, either in solution or suspension form.

[0010] A ligand may be referred to as "photoactive" when it is believed that the ligand directly contributes to the photoactive properties of an emissive material. A ligand may be referred to as "ancillary" when it is believed that the

ligand does not contribute to the photoactive properties of an emissive material, although an ancillary ligand may alter the properties of a photoactive ligand.

[0011] As used herein, and as would be generally understood by one skilled in the art, a first "Highest Occupied Molecular Orbital" (HOMO) or "Lowest Unoccupied Molecular Orbital" (LUMO) energy level is "greater than" or "higher than" a second HOMO or LUMO energy level if the first energy level is closer to the vacuum energy level. Since ionization potentials (IP) are measured as a negative energy relative to a vacuum level, a higher HOMO energy level corresponds to an IP having a smaller absolute value (an IP that is less negative). Similarly, a higher LUMO energy level corresponds to an electron affinity (EA) having a smaller absolute value (an EA that is less negative). On a conventional energy level diagram, with the vacuum level at the top, the LUMO energy level of a material is higher than the HOMO energy level of the same material. A "higher" HOMO or LUMO energy level appears closer to the top of such a diagram than a "lower" HOMO or LUMO energy level.

[0012] As used herein, and as would be generally understood by one skilled in the art, a first work function is "greater than" or "higher than" a second work function if the first work function has a higher absolute value. Because work functions are generally measured as negative numbers relative to vacuum level, this means that a "higher" work function is more negative. On a conventional energy level diagram, with the vacuum level at the top, a "higher" work function is illustrated as further away from the vacuum level in the downward direction. Thus, the definitions of HOMO and LUMO energy levels follow a different convention than work functions.

[0013] More details on OLEDs, and the definitions described above, can be found in US Pat. No. 7,279,704.


[0014] US 2014/0001456 A1 describes OLEDs comprising aromatic heterocyclic compounds in the electron transporting zone of an emitting layer.

[0015] CN 102850329 A describes triazinyl derivatives for application in OLEDs.

SUMMARY OF THE INVENTION

[0016] The present invention is directed to a composition of materials as defined in claim 1.

[0017] A composition of materials comprising a first compound and a second compound is provided. The first compound has a structure of formula:



, Formula I.

In Formula I:

G¹ is selected from the group consisting of dibenzofuran, dibenzothiophene, dibenzoselenophene, and fluorene; L¹, L² and L³ are each independently selected from the group consisting of direct bond, phenyl, biphenyl, terphenyl, pyridine, pyrimidine, and combinations thereof;

G⁴ is selected from the group consisting of phenyl, biphenyl, terphenyl, naphthalene, phenanthrene, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phenanthroline, and combinations thereof;

G², G³, and G⁵ are each independently selected from the group consisting of phenyl, biphenyl, terphenyl, fluorene, naphthalene, phenanthrene, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phenanthroline, aza-fluorene, and combinations thereof;

G², G³, G⁴, and G⁵ are each optionally further substituted with one or more unfused substituents selected from the group consisting of deuterium, alkyl, alkoxy, cycloalkyl, cycloalkoxy, halogen, nitro, nitrile, silyl, phenyl, biphenyl, terphenyl, pyridine, and combinations thereof;

m is an integer from 0 to 7,


n is an integer from 0 to 4;

when m or n is larger than 1, then each G⁴ or G⁵ can be same or different;

when n is 0, and m is equal to or greater than 1, then each G⁴ is selected from the group consisting of phenyl, and biphenyl;

when n is equal to or greater than 1, then L¹ is not a direct bond; and
when m and n are both 0, then L¹ is biphenyl.

[0018] The second compound has a structure of formula II:



In the structure of Formula II:

Ar¹ is selected from the group consisting of triphenylene, and aza-triphenylene;

Ar² is selected from the group consisting of a direct bond, phenyl, biphenyl, terphenyl, naphthalene, pyridine, dibenzofuran, dibenzothiophene, dibenzoselenophene, aza-dibenzofuran, aza-dibenzothiophene, aza-dibenzoselenophene, and combinations thereof;

Ar³ is selected from the group consisting of benzene, biphenyl, terphenyl, naphthalene, pyridine, dibenzofuran, dibenzothiophene, dibenzoselenophene, aza-dibenzofuran, aza-dibenzothiophene, aza-dibenzoselenophene, carbazole, aza-carbazole, and combinations thereof; and

Ar¹, Ar² and Ar³ are each, independently, optionally further substituted with one or more substitutions selected from the group consisting of deuterium, halogen, alkyl, aryl, heteroaryl, and combinations thereof.

[0019] According to another embodiment of the present disclosure, a device that includes one or more organic light emitting devices is also provided. At least one of the one or more organic light emitting devices can include an anode, a cathode, and an organic layer, disposed between the anode and the cathode. The organic layer can include a composition of materials as defined in any one of the claims.

[0020] In yet another aspect of the present disclosure, a method for fabricating an organic light emitting device is provided. The organic light emitting device can include a first electrode, a second electrode, and a first organic layer disposed between the first electrode and the second electrode, where the first organic layer comprises a first composition comprising a mixture of a first compound and a second compound.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021]

FIG. 1 shows an organic light emitting device.

FIG. 2 shows an inverted organic light emitting device that does not have a separate electron transport layer.

FIG. 3 shows Formula I as disclosed herein.

FIG. 4 shows Formula II as disclosed herein.

DETAILED DESCRIPTION

[0022] Generally, an OLED comprises at least one organic layer disposed between and electrically connected to an anode and a cathode. When a current is applied, the anode injects holes and the cathode injects electrons into the organic layer(s). The injected holes and electrons each migrate toward the oppositely charged electrode. When an electron and hole localize on the same molecule, an "exciton," which is a localized electron-hole pair having an excited energy state, is formed. Light is emitted when the exciton relaxes via a photoemissive mechanism. In some cases, the exciton may be localized on an excimer or an exciplex. Non-radiative mechanisms, such as thermal relaxation, may also occur, but are generally considered undesirable.

[0023] The initial OLEDs used emissive molecules that emitted light from their singlet states ("fluorescence") as disclosed, for example, in U.S. Pat. No. 4,769,292. Fluorescent emission generally occurs in a time frame of less than 10 nanoseconds.

[0024] More recently, OLEDs having emissive materials that emit light from triplet states ("phosphorescence") have been demonstrated. Baldo et al., "Highly Efficient Phosphorescent Emission from Organic Electroluminescent Devices," Nature, vol. 395, 151-154, 1998; ("Baldo-I") and Baldo et al., "Very high-efficiency green organic light-emitting devices

based on electrophosphorescence," Appl. Phys. Lett., vol. 75, No. 3, 4-6 (1999) ("Baldo-II"). Phosphorescence is described in more detail in US Pat. No. 7,279,704 at cols. 5-6.

5 [0025] FIG. 1 shows an organic light emitting device 100. The figures are not necessarily drawn to scale. Device 100 may include a substrate 110, an anode 115, a hole injection layer 120, a hole transport layer 125, an electron blocking layer 130, an emissive layer 135, a hole blocking layer 140, an electron transport layer 145, an electron injection layer 150, a protective layer 155, a cathode 160, and a barrier layer 170. Cathode 160 is a compound cathode having a first conductive layer 162 and a second conductive layer 164. Device 100 may be fabricated by depositing the layers described, in order. The properties and functions of these various layers, as well as example materials, are described in more detail in US 7,279,704 at cols. 6-10.

10 [0026] More examples for each of these layers are available. For example, a flexible and transparent substrate-anode combination is disclosed in U.S. Pat. No. 5,844,363. An example of a p-doped hole transport layer is m-MTDATA doped with F₄-TCNQ at a molar ratio of 50:1, as disclosed in U.S. Patent Application Publication No. 2003/0230980. Examples of emissive and host materials are disclosed in U.S. Pat. No. 6,303,238 to Thompson et al. An example of an n-doped electron transport layer is BPhen doped with Li at a molar ratio of 1:1, as disclosed in U.S. Patent Application Publication No. 2003/0230980. U.S. Pat. Nos. 5,703,436 and 5,707,745 disclose examples of cathodes including compound cathodes having a thin layer of metal such as Mg:Ag with an overlying transparent, electrically-conductive, sputter-deposited ITO layer. The theory and use of blocking layers is described in more detail in U.S. Pat. No. 6,097,147 and U.S. Patent Application Publication No. 2003/0230980. Examples of injection layers are provided in U.S. Patent Application Publication No. 2004/0174116. A description of protective layers may be found in U.S. Patent Application Publication No. 2004/0174116.

20 [0027] FIG. 2 shows an inverted OLED 200. The device includes a substrate 210, a cathode 215, an emissive layer 220, a hole transport layer 225, and an anode 230. Device 200 may be fabricated by depositing the layers described, in order. Because the most common OLED configuration has a cathode disposed over the anode, and device 200 has cathode 215 disposed under anode 230, device 200 may be referred to as an "inverted" OLED. Materials similar to those described with respect to device 100 may be used in the corresponding layers of device 200. FIG. 2 provides one example of how some layers may be omitted from the structure of device 100.

25 [0028] The simple layered structure illustrated in FIGS. 1 and 2 is provided by way of nonlimiting example, and it is understood that embodiments of the invention may be used in connection with a wide variety of other structures. The specific materials and structures described are exemplary in nature, and other materials and structures may be used. Functional OLEDs may be achieved by combining the various layers described in different ways, or layers may be omitted entirely, based on design, performance, and cost factors. Other layers not specifically described may also be included. Materials other than those specifically described may be used. Although many of the examples provided herein describe various layers as comprising a single material, it is understood that combinations of materials, such as a mixture of host and dopant, or more generally a mixture, may be used. Also, the layers may have various sublayers. The names given to the various layers herein are not intended to be strictly limiting. For example, in device 200, hole transport layer 225 transports holes and injects holes into emissive layer 220, and may be described as a hole transport layer or a hole injection layer. In one embodiment, an OLED may be described as having an "organic layer" disposed between a cathode and an anode. This organic layer may comprise a single layer, or may further comprise multiple layers of different organic materials as described, for example, with respect to FIGS. 1 and 2.

30 [0029] Structures and materials not specifically described may also be used, such as OLEDs comprised of polymeric materials (PLEDs) such as disclosed in U.S. Pat. No. 5,247,190 to Friend et al. By way of further example, OLEDs having a single organic layer may be used. OLEDs may be stacked, for example as described in U.S. Pat. No. 5,707,745 to Forrest et al. The OLED structure may deviate from the simple layered structure illustrated in FIGS. 1 and 2. For example, the substrate may include an angled reflective surface to improve out-coupling, such as a mesa structure as described in U.S. Pat. No. 6,091,195 to Forrest et al., and/or a pit structure as described in U.S. Pat. No. 5,834,893 to Bulovic et al.

35 [0030] Unless otherwise specified, any of the layers of the various embodiments may be deposited by any suitable method. For the organic layers, preferred methods include thermal evaporation, ink-jet, such as described in U.S. Pat. Nos. 6,013,982 and 6,087,196, organic vapor phase deposition (OVPD), such as described in U.S. Pat. No. 6,337,102 to Forrest et al., and deposition by organic vapor jet printing (OVJP), such as described in U.S. Pat. No. 7,431,968. Other suitable deposition methods include spin coating and other solution based processes. Solution based processes are preferably carried out in nitrogen or an inert atmosphere. For the other layers, preferred methods include thermal evaporation. Preferred patterning methods include deposition through a mask, cold welding such as described in U.S. Pat. Nos. 6,294,398 and 6,468,819, and patterning associated with some of the deposition methods such as ink-jet and OVJD. Other methods may also be used. The materials to be deposited may be modified to make them compatible with a particular deposition method. For example, substituents such as alkyl and aryl groups, branched or unbranched, and preferably containing at least 3 carbons, may be used in small molecules to enhance their ability to undergo solution processing. Substituents having 20 carbons or more may be used, and 3-20 carbons is a preferred range. Materials

with asymmetric structures may have better solution processibility than those having symmetric structures, because asymmetric materials may have a lower tendency to recrystallize. Dendrimer substituents may be used to enhance the ability of small molecules to undergo solution processing.

[0031] Devices fabricated in accordance with the present invention may

5 further optionally comprise a barrier layer. One purpose of the barrier layer is to protect the electrodes and organic layers from damaging exposure to harmful species in the environment including moisture, vapor and/or gases, etc. The barrier layer may be deposited over, under or next to a substrate, an electrode, or over any other parts of a device including an edge. The barrier layer may comprise a single layer, or multiple layers. The barrier layer may be formed by various known chemical vapor deposition techniques and may include compositions having a single phase as well as compositions having multiple phases. Any suitable material or combination of materials may be used for the barrier layer. The barrier layer may incorporate an inorganic or an organic compound or both. The preferred barrier layer comprises a mixture of a polymeric material and a non-polymeric material as described in U.S. Pat. No. 7,968,146, PCT Pat. Application Nos. PCT/US2007/023098 and PCT/US2009/042829. To be considered a "mixture", the aforesaid polymeric and non-polymeric materials comprising the barrier layer should be deposited under the same reaction conditions and/or at the same time. The weight ratio of polymeric to non-polymeric material may be in the range of 95:5 to 5:95. The polymeric material and the non-polymeric material may be created from the same precursor material. In one example, the mixture of a polymeric material and a non-polymeric material consists essentially of polymeric silicon and inorganic silicon.

[0032] Devices fabricated in accordance with the invention can be incorporated into a wide variety of electronic component modules (or units) that can be incorporated into a variety of electronic products or intermediate components.

20 Examples of such electronic products or intermediate components include display screens, lighting devices such as discrete light source devices or lighting panels, etc. that can be utilized by the end-user product manufacturers. Such electronic component modules can optionally include the driving electronics and/or power source(s). Devices fabricated in accordance with embodiments of the invention can be incorporated into a wide variety of consumer products that have one or more of the electronic component modules (or units) incorporated therein. Such consumer products would include any kind of products that include one or more light source(s) and/or one or more of some type of visual displays. Some examples of such consumer products include flat panel displays, computer monitors, medical monitors, televisions, billboards, lights for interior or exterior illumination and/or signaling, heads-up displays, fully or partially transparent displays, flexible displays, laser printers, telephones, cell phones, tablets, phablets, personal digital assistants (PDAs), laptop computers, digital cameras, camcorders, viewfinders, micro-displays, 3-D displays, vehicles, a large area wall, theater or stadium screen, or a sign. Various control mechanisms may be used to control devices fabricated in accordance with the present invention, including passive matrix and active matrix. Many of the devices are intended for use in a temperature range comfortable to humans, such as 18 degrees C. to 30 degrees C., and more preferably at room temperature (20-25 degrees C), but could be used outside this temperature range, for example, from -40 degree C to + 80 degree C.

35 **[0033]** The materials and structures described herein may have applications in devices other than OLEDs. For example, other optoelectronic devices such as organic solar cells and organic photodetectors may employ the materials and structures. More generally, organic devices, such as organic transistors, may employ the materials and structures.

[0034] The term "halo," "halogen," or "halide" as used herein includes fluorine, chlorine, bromine, and iodine.

40 **[0035]** The term "alkyl" as used herein contemplates both straight and branched chain alkyl radicals. Preferred alkyl groups are those containing from one to fifteen carbon atoms and includes methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, and the like. Additionally, the alkyl group may be optionally substituted.

[0036] The term "cycloalkyl" as used herein contemplates cyclic alkyl radicals. Preferred cycloalkyl groups are those containing 3 to 7 carbon atoms and includes cyclopropyl, cyclopentyl, cyclohexyl, and the like. Additionally, the cycloalkyl group may be optionally substituted.

45 **[0037]** The term "alkenyl" as used herein contemplates both straight and branched chain alkene radicals. Preferred alkenyl groups are those containing two to fifteen carbon atoms. Additionally, the alkenyl group may be optionally substituted.

[0038] The term "alkynyl" as used herein contemplates both straight and branched chain alkyne radicals. Preferred alkynyl groups are those containing two to fifteen carbon atoms. Additionally, the alkynyl group may be optionally substituted.

50 **[0039]** The terms "aralkyl" or "arylalkyl" as used herein are used interchangeably and contemplate an alkyl group that has as a substituent an aromatic group. Additionally, the aralkyl group may be optionally substituted.

[0040] The term "heterocyclic group" as used herein contemplates aromatic and non-aromatic cyclic radicals. Hetero-aromatic cyclic radicals also means heteroaryl. Preferred hetero-non-aromatic cyclic groups are those containing 3 or 7 ring atoms which includes at least one hetero atom, and includes cyclic amines such as morpholino, piperidino, pyrrolidino, and the like, and cyclic ethers, such as tetrahydrofuran, tetrahydropyran, and the like. Additionally, the heterocyclic group may be optionally substituted.

55 **[0041]** The term "aryl" or "aromatic group" as used herein contemplates single-ring groups and polycyclic ring systems.

The polycyclic rings may have two or more rings in which two carbons are common to two adjoining rings (the rings are "fused") wherein at least one of the rings is aromatic, e.g., the other rings can be cycloalkyls, cycloalkenyls, aryl, heterocycles, and/or heteroaryls. Additionally, the aryl group may be optionally substituted.

5 [0042] The term "heteroaryl" as used herein contemplates single-ring hetero-aromatic groups that may include from one to three heteroatoms, for example, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, triazole, pyrazole, pyridine, pyrazine and pyrimidine, and the like. The term heteroaryl also includes polycyclic hetero-aromatic systems having two or more rings in which two atoms are common to two adjoining rings (the rings are "fused") wherein at least one of the rings is a heteroaryl, e.g., the other rings can be cycloalkyls, cycloalkenyls, aryl, heterocycles, and/or heteroaryls. Additionally, the heteroaryl group may be optionally substituted.

10 [0043] The alkyl, cycloalkyl, alkenyl, alkynyl, aralkyl, heterocyclic group, aryl, and heteroaryl may be optionally substituted with one or more substituents selected from the group consisting of hydrogen, deuterium, halogen, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, cyclic amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acid, ether, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof.

15 [0044] As used herein, "substituted" indicates that a substituent other than H is bonded to the relevant position, such as carbon. Thus, for example, where R¹ is mono-substituted, then one R¹ must be other than H. Similarly, where R¹ is di-substituted, then two of R¹ must be other than H. Similarly, where R¹ is unsubstituted, R¹ is hydrogen for all available positions.

20 [0045] The "aza" designation in the fragments described herein, i.e. aza-dibenzofuran, aza-dibenzothiophene, etc. means that one or more of the C-H groups in the respective fragment can be replaced by a nitrogen atom, for example, and without any limitation, azatriphenylene encompasses both dibenzo[*f,h*]quinoxaline and dibenzo[*f,h*]quinoline. One of ordinary skill in the art can readily envision other nitrogen analogs of the aza-derivatives described above, and all such analogs are intended to be encompassed by the terms as set forth herein.

25 [0046] It is to be understood that when a molecular fragment is described as being a substituent or otherwise attached to another moiety, its name may be written as if it were a fragment (e.g. phenyl, phenylene, naphthyl, dibenzofuryl) or as if it were the whole molecule (e.g. benzene, naphthalene, dibenzofuran). As used herein, these different ways of designating a substituent or attached fragment are considered to be equivalent.

30 [0047] Often, the emissive layer (EML) of OLED devices exhibiting good lifetime and efficiency requires more than two components (e.g. 3 or 4 components). Fabricating such EMLs using vacuum thermal evaporation (VTE) process then requires evaporating 3 or 4 evaporation source materials in separate VTE sublimation crucibles, which is very complicated and costly compared to a standard two-component EML with a single host and an emitter, which requires only two evaporation sources.

35 [0048] Premixing two or more materials and evaporating them from one VTE sublimation crucible can reduce the complexity of the fabrication process. However, the co-evaporation must be stable and produce an evaporated film having a composition that remains constant through the evaporation process. Variations in the film's composition may adversely affect the device performance. In order to obtain a stable co-evaporation from a mixture of compounds under vacuum, one would assume that the materials must have the same evaporation temperature under the same condition. However, this may not be the only parameter one has to consider. When two compounds are mixed together, they may interact with each other and the evaporation property of the mixture may differ from their individual properties. On the other hand, materials with slightly different evaporation temperatures may form a stable co-evaporation mixture. Therefore, it is extremely difficult to achieve a stable co-evaporation mixture. So far, there have been very few stable co-evaporation mixture examples. "Evaporation temperature" of a material is measured in a vacuum deposition tool at a constant pressure, normally between 1,33322x10⁻⁵ Pa to 1,33322x10⁻⁶ Pa (1x10⁻⁷ Torr to 1x10⁻⁸ Torr), at a 2Å/sec deposition rate on a surface positioned at a set distance away from the evaporation source of the material being evaporated, e.g. sublimation crucible in a VTE tool. The various measured values such as temperature, pressure, deposition rate, etc. disclosed herein are expected to have nominal variations because of the expected tolerances in the measurements that produced these quantitative values as understood by one of ordinary skill in the art.


40 [0049] Many factors other than temperature can contribute to the ability to achieve stable co-evaporation, such as the miscibility of the different materials and the phase transition temperatures of the different materials. The inventors found that when two materials have similar evaporation temperatures, and similar mass loss rate or similar vapor pressures, the two materials can co-evaporate consistently. "Mass loss rate" of a material is defined as the percentage of mass lost over time ("percentage/minute" or "%/min") and is determined by measuring the time it takes to lose the first 10% of the mass of a sample of the material as measured by thermal gravity analysis (TGA) under a given experimental condition at a given constant temperature for a given material after the a steady evaporation state is reached. The given constant temperature is one temperature point that is chosen so that the value of mass loss rate is between about 0.05 to 0.50 %/min. A skilled person in this field should appreciate that in order to compare two parameters, the experimental condition should be consistent. The method of measuring mass loss rate and vapor pressure is well known in the art and can be found, for example, in Bull. et al. Mater. Sci. 2011, 34, 7.

[0050] In the state of the art OLED devices, the EML may consist of three or more components. In one example, the EML can consist of two host-type compounds and an emitter combination (e.g. a hole transporting cohost (h-host), an electron transporting cohost (e-host), and a compound capable of functioning as an emitter in an OLED at room temperature). In another example, the EML can consist of one host-type compound and two emitter-type compounds (e.g., a host compound and two compounds each capable of functioning as an emitter in an OLED at room temperature). Conventionally, in order to fabricate such EMLs having three or more components using VTE process, three or more evaporation sources are required, one for each of the components. Because the concentration of the components are important for the device performance, typically, the rate of deposition of each component is measured individually during the deposition process. This makes the VTE process complicated and costly. Thus, it is desired to premix at least two of the components of such EMLs to reduce the number of VTE evaporation sources.

[0051] If any two of the three or more components of the EMLs can be premixed and form a stable mixture of co-evaporation source, then the number of evaporation sources required for EML layer fabrication would be reduced. In order for materials to be premixable into an evaporation source, they should co-evaporate and deposit uniformly without changing the ratio. The ratio of the components in the mixture should be the same as the ratio of the components in the evaporation deposited films from these premixed materials. Therefore, the concentration of the two components in the deposited film is controlled by their concentration in the premixed evaporation source.

[0052] This disclosure describes a new class of h- and e-hosts that can be premixed and stably co-evaporated from a single source.

[0053] A composition of materials comprising a first compound and a second compound is disclosed. The first compound has a structure of formula:



, Formula I.

In Formula I:

G^1 is selected from the group consisting of dibenzofuran, dibenzothiophene, dibenzoselenophene, and fluorene; L^1 , L^2 and L^3 are each independently selected from the group consisting of direct bond, phenyl, biphenyl, terphenyl, pyridine, pyrimidine, and combinations thereof;

G^4 is selected from the group consisting of phenyl, biphenyl, terphenyl, naphthalene, phenanthrene, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phenanthroline, and combinations thereof;

G^2 , G^3 , and G^5 are each independently selected from the group consisting of phenyl, biphenyl, terphenyl, fluorene, naphthalene, phenanthrene, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phenanthroline, aza-fluorene, and combinations thereof;

G^2 , G^3 , G^4 , and G^5 are each optionally further substituted with one or more unfused substituents selected from the group consisting of deuterium, alkyl, alkoxy, cycloalkyl, cycloalkoxy, halogen, nitro, nitrile, silyl, phenyl, biphenyl, terphenyl, pyridine, and combinations thereof;

m is an integer from 0 to 7,

n is an integer from 0 to 4;

when m or n is larger than 1, then each G^4 or G^5 can be same or different;

when n is 0, and m is equal to or greater than 1, then each G^4 is selected from the group consisting of phenyl, and biphenyl;


when n is equal to or greater than 1, then L^1 is not a direct bond; and

when m and n are both 0, then L^1 is biphenyl.


[0054] In some embodiments, one or more of L^1 , L^2 and L^3 can be a direct bond, and the direct bond can be a single bond or a double bond. When L^1 is a direct bond, $n = 0$.

[0055] In some embodiments, n is 0, while n is equal to or greater than 1 in other embodiments. In some embodiments, m and n are both 0. In some embodiments, m is equal to or greater than 1.

[0056] In some embodiments, G⁴ has the structure selected from the group consisting of:




[0057] In some embodiments, G¹ has the structure selected from the group consisting of:




wherein:


20 X is selected from a group consisting of O, S and Se;
 RB¹ and RB² are independently selected from a group consisting of hydrogen, deuterium, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, halogen, and combinations thereof; and
 RB¹ and RB² are optionally joined to form a ring.

[0058] In some embodiments, L¹ is selected from the group consisting of:



[0059] In some embodiments, G², G³ and G⁵ are independently selected from the group consisting of:






15 wherein

R^{B1} and R^{B2} are independently selected from a group consisting of hydrogen, deuterium, alkyl, cycloalkyl, alkoxy, aryl, heteroaryl, halogen, and combinations thereof; and

20 R^{B1} and R^{B2} are optionally joined to form a ring.

[0060] In some embodiments, at least one of G^2 , G^3 , G^4 and G^5 is substituted with at least one fluorine atom.


[0061] In some embodiments, the first compound has the formula:



35 where X is selected from a group consisting of O, S and Se.


[0062] In some embodiments, the first compound is selected from the group consisting of:

40 Compound A1 through A3, each represented by the formula



45 wherein in Compound A1: X=O,
in Compound A2: X=S,
in Compound A3: X=Se

Compound A4 through A6, each represented by the formula



50 wherein in Compound A4: X=O,
in Compound A5: X=S,
in Compound A6: X=Se



55

EP 2 966 706 B1

Compound A7 through A9, each represented by the formula

Compound A10 through A12, each represented by the formula

5



10

wherein in Compound A7: X=O,
in Compound A8: X=S,
in Compound A9: X=Se



wherein in Compound A10: X=O,
in Compound A11: X=S,
in Compound A12: X=Se

15

Compound A13 through A15, each represented by the formula

Compound A16 through A18, each represented by the formula

20



25

wherein in Compound A13: X=O,
in Compound A14: X=S,
in Compound A15: X=Se



wherein in Compound A16: X=O,
in Compound A17: X=S,
in Compound A18: X=Se

30

Compound A19 through A21, each represented by the formula

Compound A22 through A24, each represented by the formula

35



40

wherein in Compound A19: X=O,
in Compound A20: X=S,
in Compound A21: X=Se



wherein in Compound A22: X=O,
in Compound A23: X=S,
in Compound A24: X=Se

45

Compound A25 through A27, each represented by the formula

Compound A28 through A30, each represented by the formula

50




55

wherein in Compound A25: X=O,
in Compound A26: X=S,
in Compound A27: X=Se

wherein in Compound A28: X=O,
in Compound A29: X=S,
in Compound A30: X=Se


EP 2 966 706 B1

Compound A31 through A33, each represented by the formula




wherein in Compound A31: X=O,
in Compound A32: X=S,
in Compound A33: X=Se

Compound A34 through A36, each represented by the formula




wherein in Compound A34: X=O,
in Compound A35: X=S,
in Compound A36: X=Se

Compound A37 through A39, each represented by the formula




wherein in Compound A37: X=O,
in Compound A38: X=S,
in Compound A39: X=Se

Compound A40 through A42, each represented by the formula




wherein in Compound A40: X=O,
in Compound A41: X=S,
in Compound A42: X=Se

Compound A43 through A45, each represented by the formula




wherein in Compound A43: X=O,
in Compound A44: X=S,
in Compound A45: X=Se

Compound A46 through A48, each represented by the formula




wherein in Compound A46: X=O,
in Compound A47: X=S,
in Compound A48: X=Se

Compound A49 through A51, each represented by the formula



wherein in Compound A49: X=O,
in Compound A50: X=S,
in Compound A51: X=Se

Compound A52 through A54, each represented by the formula




wherein in Compound A52: X=O,
in Compound A53: X=S,
in Compound A54: X=Se

EP 2 966 706 B1

Compound A55 through A57, each represented by the formula


5



10

wherein in Compound A55: X=O,
in Compound A56: X=S,
in Compound A57: X=Se


Compound A58 through A60, each represented by the formula



wherein in Compound A58: X=O,
in Compound A59: X=S,
in Compound A60: X=Se

15


Compound A61 through A63, each represented by the formula



20

wherein in Compound A61: X=O,
in Compound A62: X=S,
in Compound A63: X=Se

Compound A64 through A66, each represented by the formula




25

wherein in Compound A64: X=O,
in Compound A65: X=S,
in Compound A66: X=Se

30


Compound A67 through A69, each represented by the formula



35

wherein in Compound A67: X=O,
in Compound A68: X=S,
in Compound A69: X=Se

Compound A70 through A72, each represented by the formula




40

wherein in Compound A70: X=O,
in Compound A71: X=S,
in Compound A72: X=Se

45


Compound A73 through A75, each represented by the formula



50

wherein in Compound A73: X=O,
in Compound A74: X=S,
in Compound A75: X=Se

Compound A76 through A78, each represented by the formula



55

wherein in Compound A76: X=O,
in Compound A77: X=S,
in Compound A78: X=Se



EP 2 966 706 B1

Compound A82 through A84, each represented by the formula

5

Compound A79 through A81, each represented by the formula

10



15

wherein in Compound A79: X=O,
in Compound A80: X=S,
in Compound A81: X=Se



wherein in Compound A82: X=O,
in Compound A83: X=S,
in Compound A84: X=Se

20

Compound A85 through A87, each represented by the formula

Compound A88 through A90, each represented by the formula

25



30

wherein in Compound A85: X=O,
in Compound A86: X=S,
in Compound A87: X=Se



wherein in Compound A88: X=O,
in Compound A89: X=S,
in Compound A90: X=Se

35

Compound A91 through A93, each represented by the formula

Compound A94 through A96, each represented by the formula

40



45

wherein in Compound A91: X=O,
in Compound A92: X=S,
in Compound A93: X=Se


wherein in Compound A94: X=O,
in Compound A95: X=S,
in Compound A96: X=Se

50

55


EP 2 966 706 B1

Compound A97 through A99, each represented by the formula




wherein in Compound A97: X=O,
in Compound A98: X=S,
in Compound A99: X=Se

Compound A100 through A102, each represented by the formula




wherein in Compound A100: X=O,
in Compound A101: X=S,
in Compound A102: X=Se

Compound A103 through A105, each represented by the formula




wherein in Compound A103: X=O,
in Compound A104: X=S,
in Compound A105: X=Se

Compound A106 through A108, each represented by the formula




wherein in Compound A106: X=O,
in Compound A107: X=S,
in Compound A108: X=Se

Compound A109 through A111, each represented by the formula




wherein in Compound A109: X=O,
in Compound A110: X=S,
in Compound A111: X=Se

Compound A112 through A114, each represented by the formula




wherein in Compound A112: X=O,
in Compound A113: X=S,
in Compound A114: X=Se

Compound A115 through A117, each represented by the formula




wherein in Compound A115: X=O,
in Compound A116: X=S,
in Compound A117: X=Se

5




Compound B1




Compound B2

10


15



Compound B3



Compound B4




Compound B5


20

25


30



Compound B6



Compound B7






Compound B8

35

40


[0063] In some embodiments, n is 0, m is 1, and G⁴-G¹ has a structure selected from the group consisting of:

45



50

55



and

[0064] In some embodiments, the first compound is selected from the group consisting of:

30

35

40

45

50



55

EP 2 966 706 B1

Compound C1 through C3, each represented by the formula

Compound C4 through C6, each represented by the formula

5



10

wherein in Compound C1: X=O,
in Compound C2: X=S,
in Compound C3: X=Se



wherein in Compound C4: X=O,
in Compound C5: X=S,
in Compound C6: X=Se

15

Compound C7 through C9, each represented by the formula

Compound C10 through C12, each represented by the formula

20



25

wherein in Compound C7: X=O,
in Compound C8: X=S,
in Compound C9: X=Se

wherein in Compound C10: X=O,
in Compound C11: X=S,
in Compound C12: X=Se

30

35

40

45


50

55

EP 2 966 706 B1

Compound C13 through C15, each represented by the formula


5



10

wherein in Compound C13: X=O,
in Compound C14: X=S,
in Compound C15: X=Se

Compound C16 through C18, each represented by the formula




wherein in Compound C16: X=O,
in Compound C17: X=S,
in Compound C18: X=Se

15

Compound C19 through C21, each represented by the formula


20



25

wherein in Compound C19: X=O,
in Compound C20: X=S,
in Compound C21: X=Se

Compound C22 through C24, each represented by the formula




wherein in Compound C22: X=O,
in Compound C23: X=S,
in Compound C24: X=Se

30

Compound C25 through C27, each represented by the formula


35



40

wherein in Compound C25: X=O,
in Compound C26: X=S,
in Compound C27: X=Se

Compound C28 through C30, each represented by the formula




wherein in Compound C28: X=O,
in Compound C29: X=S,
in Compound C30: X=Se

45

Compound C31 through C33, each represented by the formula


50



55

wherein in Compound C31: X=O,
in Compound C32: X=S,
in Compound C33: X=Se

Compound C34 through C36, each represented by the formula




wherein in Compound C34: X=O,
in Compound C35: X=S,
in Compound C36: X=Se

EP 2 966 706 B1

Compound C37 through C39, each represented by the formula


5



10

wherein in Compound C37: X=O,
in Compound C38: X=S,
in Compound C39: X=Se

Compound C40 through C42, each represented by the formula




wherein in Compound C40: X=O,
in Compound C41: X=S,
in Compound C42: X=Se

15

Compound C43 through C45, each represented by the formula


20



wherein in Compound C43: X=O,
in Compound C44: X=S,
in Compound C45: X=Se

25


Compound C46 through C48, each represented by the formula



wherein in Compound C46: X=O,
in Compound C47: X=S,
in Compound C48: X=Se

Compound C49 through C51, each represented by the formula


30



wherein in Compound C49: X=O,
in Compound C50: X=S,
in Compound C51: X=Se

35


Compound C52 through C54, each represented by the formula



wherein in Compound C52: X=O,
in Compound C53: X=S,
in Compound C54: X=Se

Compound C55 through C57, each represented by the formula


45



wherein in Compound C55: X=O,
in Compound C56: X=S,
in Compound C57: X=Se

50

Compound C58 through C60, each represented by the formula



wherein in Compound C58: X=O,
in Compound C59: X=S,
in Compound C60: X=Se



55

EP 2 966 706 B1

Compound C61 through C63, each represented by the formula

Compound C64 through C66, each represented by the formula

5



10

wherein in Compound C61: X=O,
in Compound C62: X=S,
in Compound C63: X=Se



wherein in Compound C64: X=O,
in Compound C65: X=S,
in Compound C66: X=Se

15

Compound C67 through C69, each represented by the formula

Compound C70 through C72, each represented by the formula

20



25

wherein in Compound C67: X=O,
in Compound C68: X=S,
in Compound C69: X=Se



wherein in Compound C70: X=O,
in Compound C71: X=S,
in Compound C72: X=Se

30

Compound C73 through C75, each represented by the formula

Compound C76 through C78, each represented by the formula

35



40

wherein in Compound C73: X=O,
in Compound C74: X=S,
in Compound C75: X=Se



wherein in Compound C76: X=O,
in Compound C77: X=S,
in Compound C78: X=Se

45

Compound C79 through C81, each represented by the formula

Compound C82 through C84, each represented by the formula

50



55


wherein in Compound C79: X=O,
in Compound C80: X=S,
in Compound C81: X=Se

wherein in Compound C82: X=O,
in Compound C83: X=S,
in Compound C84: X=Se

EP 2 966 706 B1

Compound C85 through C87, each represented by the formula


5



10

wherein in Compound C85: X=O,
in Compound C86: X=S,
in Compound C87: X=Se

Compound C88 through C90, each represented by the formula




wherein in Compound C88: X=O,
in Compound C89: X=S,
in Compound C90: X=Se

15

Compound C91 through C93, each represented by the formula


20



wherein in Compound C91: X=O,
in Compound C92: X=S,
in Compound C93: X=Se

25

Compound C94 through C96, each represented by the formula




wherein in Compound C94: X=O,
in Compound C95: X=S,
in Compound C96: X=Se

30

Compound C97 through C99, each represented by the formula


35



wherein in Compound C97: X=O,
in Compound C98: X=S,
in Compound C99: X=Se

40

Compound C100 through C102, each represented by the formula




wherein in Compound C100: X=O,
in Compound C101: X=S,
in Compound C102: X=Se

45

Compound C103 through C105, each represented by the formula


50



wherein in Compound C103: X=O,
in Compound C104: X=S,
in Compound C105: X=Se

55

Compound C106 through C108, each represented by the formula




wherein in Compound C106: X=O,
in Compound C107: X=S,
in Compound C108: X=Se

EP 2 966 706 B1

Compound C109 through C111, each represented by the formula

5




10

wherein in Compound C109: X=O,
in Compound C110: X=S,
in Compound C111: X=Se

15


Compound C112 through C114, each represented by the formula



wherein in Compound C112: X=O,
in Compound C113: X=S,
in Compound C114: X=Se

Compound C115 through C117, each represented by the formula

20




25

wherein in Compound C115: X=O,
in Compound C116: X=S,
in Compound C117: X=Se

30


Compound C118 through C120, each represented by the formula



wherein in Compound C118: X=O,
in Compound C119: X=S,
in Compound C120: X=Se

Compound C121 through C123, each represented by the formula

35




40

wherein in Compound C121: X=O,
in Compound C122: X=S,
in Compound C123: X=Se

45

Compound C124 through C126, each represented by the formula



wherein in Compound C124: X=O,
in Compound C125: X=S,
in Compound C126: X=Se


50

55

EP 2 966 706 B1

Compound C127 through C129, each represented by the formula

5




10

wherein in Compound C127: X=O,
in Compound C128: X=S,
in Compound C129: X=Se

15


Compound C130 through C132, each represented by the formula



wherein in Compound C130: X=O,
in Compound C131: X=S,
in Compound C132: X=Se

Compound C133 through C135, each represented by the formula


20



25

wherein in Compound C133: X=O,
in Compound C134: X=S,
in Compound C135: X=Se


30



wherein in Compound C136: X=O,
in Compound C137: X=S,
in Compound C138: X=Se

Compound C139 through C141, each represented by the formula

35




40

wherein in Compound C139: X=O,
in Compound C140: X=S,
in Compound C141: X=Se

45

Compound C142 through C144, each represented by the formula



wherein in Compound C142: X=O,
in Compound C143: X=S,
in Compound C144: X=Se

50



55

EP 2 966 706 B1

Compound C145 through C147, each represented by the formula

Compound C148 through C150, each represented by the formula

5



10

wherein in Compound C145: X=O,
in Compound C146: X=S,
in Compound C147: X=Se



wherein in Compound C148: X=O,
in Compound C149: X=S,
in Compound C150: X=Se

15

Compound C151 through C153, each represented by the formula

Compound C154 through C156, each represented by the formula

20



25

wherein in Compound C151: X=O,
in Compound C152: X=S,
in Compound C153: X=Se



wherein in Compound C154: X=O,
in Compound C155: X=S,
in Compound C156: X=Se

30

Compound C157 through C159, each represented by the formula

Compound C160 through C162, each represented by the formula

35



40

wherein in Compound C157: X=O,
in Compound C158: X=S,
in Compound C159: X=Se

wherein in Compound C160: X=O,
in Compound C161: X=S,
in Compound C162: X=Se

45


50

55

EP 2 966 706 B1

Compound C163 through C165, each represented by the formula

5




10

wherein in Compound C163: X=O,
in Compound C164: X=S,
in Compound C165: X=Se

15


Compound C166 through C168, each represented by the formula



wherein in Compound C166: X=O,
in Compound C167: X=S,
in Compound C168: X=Se

Compound C169 through C171, each represented by the formula

20




25

wherein in Compound C169: X=O,
in Compound C170: X=S,
in Compound C171: X=Se

30


Compound C172 through C174, each represented by the formula



wherein in Compound C172: X=O,
in Compound C173: X=S,
in Compound C174: X=Se

Compound C175 through C177, each represented by the formula

35




40

wherein in Compound C175: X=O,
in Compound C176: X=S,
in Compound C177: X=Se

45

Compound C178 through C180, each represented by the formula



wherein in Compound C178: X=O,
in Compound C179: X=S,
in Compound C180: X=Se

50

55



EP 2 966 706 B1

Compound C181 through C183, each represented by the formula

Compound C184 through C186, each represented by the formula

5

10



15

wherein in Compound C181: X=O,
in Compound C182: X=S,
in Compound C183: X=Se



wherein in Compound C184: X=O,
in Compound C185: X=S,
in Compound C186: X=Se

Compound C187 through C189, each represented by the formula

Compound C190 through C192, each represented by the formula

20

25



30

wherein in Compound C187: X=O,
in Compound C188: X=S,
in Compound C189: X=Se



wherein in Compound C190: X=O,
in Compound C191: X=S,
in Compound C192: X=Se

Compound C193 through C195, each represented by the formula

Compound C196 through C198, each represented by the formula

35

40



45

wherein in Compound C193: X=O,
in Compound C194: X=S,
in Compound C195: X=Se

wherein in Compound C196: X=O,
in Compound C197: X=S,
in Compound C198: X=Se

50



55

EP 2 966 706 B1

Compound C199 through C201, each represented by the formula

Compound C202 through C204, each represented by the formula

5



10

wherein in Compound C199: X=O,
in Compound C200: X=S,
in Compound C201: X=Se



wherein in Compound C202: X=O,
in Compound C203: X=S,
in Compound C204: X=Se

15

Compound C205 through C207, each represented by the formula

Compound C208 through C210, each represented by the formula

20



25

wherein in Compound C205: X=O,
in Compound C206: X=S,
in Compound C207: X=Se



wherein in Compound C208: X=O,
in Compound C209: X=S,
in Compound C210: X=Se

30

Compound C211 through C213, each represented by the formula

Compound C214 through C216, each represented by the formula

35



40

wherein in Compound C211: X=O,
in Compound C212: X=S,
in Compound C213: X=Se

wherein in Compound C214: X=O,
in Compound C215: X=S,
in Compound C216: X=Se

45


50

55

EP 2 966 706 B1

Compound C217 through C219, each represented by the formula

5




10

15

wherein in Compound C217: X=O,
in Compound C218: X=S,
in Compound C219: X=Se


Compound C220 through C222, each represented by the formula



wherein in Compound C220: X=O,
in Compound C221: X=S,
in Compound C222: X=Se

Compound C223 through C225, each represented by the formula

20




25

30

wherein in Compound C223: X=O,
in Compound C224: X=S,
in Compound C225: X=Se


Compound C226 through C228, each represented by the formula



wherein in Compound C226: X=O,
in Compound C227: X=S,
in Compound C228: X=Se

Compound C229 through C231, each represented by the formula

35




40

45

wherein in Compound C229: X=O,
in Compound C230: X=S,
in Compound C231: X=Se

Compound C232 through C234, each represented by the formula



wherein in Compound C232: X=O,
in Compound C233: X=S,
in Compound C234: X=Se


50

55

EP 2 966 706 B1

Compound C235 through C237, each represented by the formula

5




10

wherein in Compound C235: X=O,
in Compound C236: X=S,
in Compound C237: X=Se

15

Compound C238 through C240, each represented by the formula




wherein in Compound C238: X=O,
in Compound C239: X=S,
in Compound C240: X=Se

,

,

Compound C241 through C243, each represented by the formula

20




25

wherein in Compound C241: X=O,
in Compound C242: X=S,
in Compound C243: X=Se

30

Compound C244 through C246, each represented by the formula




wherein in Compound C244: X=O,
in Compound C245: X=S,
in Compound C246: X=Se

,

,

Compound C247 through C249, each represented by the formula

35




40

wherein in Compound C247: X=O,
in Compound C248: X=S,
in Compound C249: X=Se

45

Compound C250 through C252, each represented by the formula



wherein in Compound C250: X=O,
in Compound C251: X=S,
in Compound C252: X=Se

,

, and


50

55

Compound C253 through C255, each represented by the formula

5

10




15

wherein in Compound C253: X=O,
in Compound C254: X=S,
in Compound C255: X=Se

[0065] In some embodiments, the first compound has a formula:

20

25




30

where L¹ is biphenyl.

[0066] In some embodiments, the first compound is selected from the group consisting of:

35

40



Compound D1


Compound D2

45

and


50

55



Compound D3

[0067] The composition as claimed comprises a second compound having a structure of formula II:



Formula II. In the structure of Formula II:


Ar¹ is selected from the group consisting of triphenylene, and aza-triphenylene;

Ar² is selected from the group consisting of a direct bond, phenyl, biphenyl, terphenyl, naphthalene, pyridine, dibenzofuran, dibenzothiophene, dibenzoselenophene, aza-dibenzofuran, aza-dibenzothiophene, aza-dibenzoselenophene, and combinations thereof;

Ar³ is selected from the group consisting of benzene, biphenyl, terphenyl, naphthalene, pyridine, dibenzofuran, dibenzothiophene, dibenzoselenophene, aza-dibenzofuran, aza-dibenzothiophene, aza-dibenzoselenophene, carbazole, aza-carbazole, and combinations thereof; and

Ar¹, Ar² and Ar³ are each, independently, optionally further substituted with one or more substitutions selected from the group consisting of deuterium, halogen, alkyl, aryl, heteroaryl, and combinations thereof.

[0068] In some embodiments, the second compound is selected from the group consisting of



where:

X is selected from the group consisting of O, S and Se;

R¹ and R⁴ each independently represents mono, di, or tri, substitution, or no substitution;

R², R³, R⁵, and R⁶ each independently represents mono, di, tri, or tetra substitution, or no substitution; and

R¹ to R⁶ are each independently selected from the group consisting of hydrogen, deuterium, benzene, biphenyl, terphenyl, naphthalene, fluorene, triphenylene, phenanthrene, dibenzofuran, dibenzothiophene, carbazole and combinations thereof.


[0069] In some embodiments, the second compound is selected from the group consisting of

EP 2 966 706 B1

Compound E1 through E3, each represented by the formula

Compound E4 through E6, each represented by the formula


5



10

wherein in Compound E1: X=O,
in Compound E2: X=S,
in Compound E3: X=Se

15



wherein in Compound E4: X=O,
in Compound E5: X=S,
in Compound E6: X=Se

20

25

30

35

40


45

50

55

Compound E7 through E9, each represented by the formula

5




10

wherein in Compound E7: X=O,
in Compound E8: X=S,
in Compound E9: X=Se

15


Compound E10 through E12, each represented by the formula



wherein in Compound E10: X=O,
in Compound E11: X=S,
in Compound E12: X=Se

Compound E13 through E15, each represented by the formula

20




25

wherein in Compound E13: X=O,
in Compound E14: X=S,
in Compound E15: X=Se

30


Compound E16 through E18, each represented by the formula



wherein in Compound E16: X=O,
in Compound E17: X=S,
in Compound E18: X=Se

Compound E19 through E21, each represented by the formula

35




40

wherein in Compound E19: X=O,
in Compound E20: X=S,
in Compound E21: X=Se

45

Compound E22 through E24, each represented by the formula




wherein in Compound E22: X=O,
in Compound E23: X=S,
in Compound E24: X=Se

50

55

Compound E25 through E27, each represented by the formula

5




10


wherein in Compound E25: X=O,
in Compound E26: X=S,
in Compound E27: X=Se

15


20



Compound E28



Compound E29




Compound E30

25


[0070] In some embodiments, the mixture of the first compound and the second compound is selected from the group consisting of:

30



(Compound E1 , Compound C1); (

40



Compound E2


Compound C2

Compound E5


50

55


5



Compound C65



Compound E8




Compound C74


10

); (, (); (


15



Compound E11



Compound C74




Compound E17


20

, (); (,


25



Compound C74



Compound E8




Compound A5

30


); (, (); (

35


40



Compound E8



Compound C17




Compound E17


45

, (); (,


50



Compound A5






Compound E26





Compound C74


55

); (, (); (



[0071] In some embodiments, the mixture of the first compound and the second compound is selected from the group consisting of:





15 **[0072]** In some embodiments, the first compound has an evaporation temperature T1 of 150 to 350 °C; the second compound has an evaporation temperature T2 of 150 to 350 °C; an absolute value of T1-T2 is less than 20 °C; the first compound has a concentration C1 in said mixture and a concentration C2 in a film formed by evaporating the mixture in a vacuum deposition tool at a constant pressure between $1,33322 \times 10^{-4}$ Pa to $1,33322 \times 10^{-7}$ Pa (1×10^{-6} Torr to 1×10^{-9} Torr), at a $2 \text{ \AA}/\text{sec}$ deposition rate on a surface position at a predefined distance away from the mixture being evaporated; and the absolute value of (C1-C2)/C1 is less than 5%.

20 **[0073]** In some embodiments, the first compound has a vapor pressure of P1 at T1 at 101,325 kPa (1 atm), the second compound has a vapor pressure of P2 at T2 at 101,325 kPa (1 atm); and the ratio of P1/P2 is within the range of 0.90 to 1.10.

25 **[0074]** In some embodiments, the first compound has a first mass loss rate and the second compound has a second mass loss rate, wherein the ratio between the first mass loss rate and the second mass loss rate is within the range of 0.90 to 1.10.

30 **[0075]** In some embodiments, the first compound and the second compound each has a purity in excess of 99 % as determined by high pressure liquid chromatography.


35 **[0076]** In some embodiments, the composition also comprises a third compound. In some embodiments, the third compound has a different chemical structure than the first and second compounds. In some embodiments, the third compound has a third mass loss rate and the ratio between the first mass loss rate and third mass loss rate is within the range of 0.90 to 1.10. In some embodiments, the third compound has an evaporation temperature T3 of 150 to 350 °C, and the absolute value of T1-T3 is less than 20 °C.

40 **[0077]** In some embodiments, the composition is in liquid form at a temperature less than T1 and T2.

45 **[0078]** According to another embodiment of the present disclosure, a device that includes one or more organic light emitting devices is also provided. At least one of the one or more organic light emitting devices can include an anode, a cathode, and an organic layer, disposed between the anode and the cathode. The organic layer can include a composition of materials as defined in the claims.

50 **[0079]** In some embodiments, the organic layer is an emissive layer and the composition comprises a host.


55 **[0080]** In some embodiments, the organic layer also includes a phosphorescent emissive dopant. In some embodiments, the phosphorescent emissive dopant is a transition metal complex having at least one ligand or part of the ligand if the ligand is more than bidentate selected from the group consisting of:



5

10

15




20

25

30


35




40

45

50



55



where:

- 50 each X¹ to X¹³ are independently selected from the group consisting of carbon and nitrogen;
 X is selected from the group consisting of BR', NR', PR', O, S, Se, C=O, S=O, SO₂, CR'R'', SiR'R'', and GeR'R'';
 R' and R'' are optionally fused or joined to form a ring;
 each R_a, R_b, R_c, and R_d may represent from mono substitution to the possible maximum number of substitution, or
 no substitution;
- 55 R', R'', R_a, R_b, R_c, and R_d are each independently selected from the group consisting of hydrogen, deuterium, halide,
 alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl,
 heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof; and

any two adjacent substituents of R_a , R_b , R_c , and R_d are optionally fused or joined to form a ring or form a multidentate ligand.

[0081] In some embodiments, the organic layer is a blocking layer and the composition is a blocking material in the organic layer. In some embodiments, the organic layer is an electron transporting layer and the composition is an electron transporting material in the organic layer.

[0082] In some embodiments, the first device is selected from the group consisting of a consumer product, an electronic component module, an organic light-emitting device, and a lighting panel.

[0083] In some embodiments, at least one of R_a , R_b , R_c , and R_d is selected from the group consisting of alkyl, cycloalkyl, partially or fully deuterated variants thereof, partially or fully fluorinated variants thereof, and combinations thereof.

[0084] In yet another aspect of the present disclosure, a method for fabricating an organic light emitting device is provided. The organic light emitting device can include a first electrode, a second electrode, and a first organic layer disposed between the first electrode and the second electrode, where the first organic layer comprises a first composition comprising a mixture of a first compound and a second compound.

[0085] In another aspect, the method includes providing a substrate having the first electrode disposed thereon; depositing the first composition over the first electrode; and depositing the second electrode over the first organic layer. In another aspect, the first composition comprises a first compound of Formula I and a second compound of Formula II.


COMBINATION WITH OTHER MATERIALS


[0086] The materials described herein as useful for a particular layer in an organic light emitting device may be used in combination with a wide variety of other materials present in the device. For example, emissive dopants disclosed herein may be used in conjunction with a wide variety of hosts, transport layers, blocking layers, injection layers, electrodes and other layers that may be present. The materials described or referred to below are non-limiting examples of materials that may be useful in combination with the compounds disclosed herein, and one of skill in the art can readily consult the literature to identify other materials that may be useful in combination.

HIL/HTL:

[0087] A hole injecting/transporting material to be used in the present invention is not particularly limited, and any compound may be used as long as the compound is typically used as a hole injecting/transporting material. Examples of the material include, but are not limited to: a phthalocyanine or porphyrin derivative; an aromatic amine derivative; an indolocarbazole derivative; a polymer containing fluorohydrocarbon; a polymer with conductivity dopants; a conducting polymer, such as PEDOT/PSS; a self-assembly monomer derived from compounds such as phosphonic acid and silane derivatives; a metal oxide derivative, such as MoO_x ; a p-type semiconducting organic compound, such as 1,4,5,8,9,12-Hexaazatriphenylenehexacarbonitrile; a metal complex, and a cross-linkable compound.


[0088] Examples of aromatic amine derivatives used in HIL or HTL include, but are not limited to the following general structures:





wherein Met is a metal; (Y¹⁰³-Y¹⁰⁴) is a bidentate ligand, Y¹⁰³ and Y¹⁰⁴ are independently selected from C, N, O, P, and S; L¹⁰¹ is another ligand; k' is an integer value from 1 to the maximum number of ligands that may be attached to the metal; and k'+k'' is the maximum number of ligands that may be attached to the metal.

[0095] In one aspect, the metal complexes are:





wherein (O-N) is a bidentate ligand, having metal coordinated to atoms O and N.

[0096] In another aspect, Met is selected from Ir and Pt. In a further aspect, (Y¹⁰³-Y¹⁰⁴) is a carbene ligand.

[0097] Examples of organic compounds used as host are selected from the group consisting aromatic hydrocarbon cyclic compounds such as benzene, biphenyl, triphenyl, triphenylene, naphthalene, anthracene, phenalene, phenanthrene, fluorene, pyrene, chrysene, perylene, azulene; group consisting aromatic heterocyclic compounds such as dibenzothiophene, dibenzofuran, dibenzoselenophene, furan, thiophene, benzofuran, benzothiophene, benzoselenophene, carbazole, indolocarbazole, pyridylindole, pyrrolodipyridine, pyrazole, imidazole, triazole, oxazole, thiazole, oxadiazole, oxatriazole, dioxazole, thiadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, oxazine, oxathiazine, oxadiazine, indole, benzimidazole, indazole, indoxazine, benzoxazole, benzisoxazole, benzothiazole, quinoline, isoquinoline, cinoline, quinazoline, quinoxaline, naphthyridine, phthalazine, pteridine, xanthene, acridine, phenazine, phenothiazine, phenoxazine, benzofuropyridine, furodipyridine, benzothienopyridine, thienodipyridine, benzoselenophenopyridine, and selenophenodipyridine; and group consisting 2 to 10 cyclic structural units which are groups of the same type or different types selected from the aromatic hydrocarbon cyclic group and the aromatic heterocyclic group and are bonded to each other directly or via at least one of oxygen atom, nitrogen atom, sulfur atom, silicon atom, phosphorus atom, boron atom, chain structural unit and the aliphatic cyclic group. Wherein each group is further substituted by a substituent selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof.

[0098] In one aspect, host compound contains at least one of the following groups in the molecule:





20 wherein R^{101} to R^{107} is independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, when it is aryl or heteroaryl, it has the similar definition as Ar's mentioned above. k is an integer from 0 to 20 or 1 to 20; k' is an integer from 0 to 20. X^{101} to X^{108} is selected from C (including CH) or N.


25 Z^{101} and Z^{102} is selected from NR^{101} , O, or S.

HBL:

30 **[0099]** A hole blocking layer (HBL) may be used to reduce the number of holes and/or excitons that leave the emissive layer. The presence of such a blocking layer in a device may result in substantially higher efficiencies as compared to a similar device lacking a blocking layer. Also, a blocking layer may be used to confine emission to a desired region of an OLED.

[0100] In one aspect, compound used in HBL contains the same molecule or the same functional groups used as host described above.

35 **[0101]** In another aspect, compound used in HBL contains at least one of the following groups in the molecule:




45 wherein k is an integer from 1 to 20; L^{101} is an another ligand, k' is an integer from 1 to 3.

ETL:


50 **[0102]** Electron transport layer (ETL) may include a material capable of transporting electrons. Electron transport layer may be intrinsic (undoped), or doped. Doping may be used to enhance conductivity. Examples of the ETL material are not particularly limited, and any metal complexes or organic compounds may be used as long as they are typically used to transport electrons.

[0103] In one aspect, compound used in ETL contains at least one of the following groups in the molecule:



wherein R^{101} is selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, when it is aryl or heteroaryl, it has the similar definition as Ar's mentioned above. Ar^1 to Ar^3 has the similar definition as Ar's mentioned above. k is an integer from 1 to 20. X^{101} to X^{108} is selected from C (including CH) or N.

[0104] In another aspect, the metal complexes used in ETL include, but are not limited to the following general formula:



wherein (O-N) or (N-N) is a bidentate ligand, having metal coordinated to atoms O, N or N, N; L^{101} is another ligand; k' is an integer value from 1 to the maximum number of ligands that may be attached to the metal.

[0105] In any above-mentioned compounds used in each layer of the OLED device, the hydrogen atoms can be partially or fully deuterated. Thus, any specifically listed substituent, such as, without limitation, methyl, phenyl, pyridyl, etc. encompasses undeuterated, partially deuterated, and fully deuterated versions thereof. Similarly, classes of substituents such as, without limitation, alkyl, aryl, cycloalkyl, heteroaryl, etc. also encompass undeuterated, partially deuterated, and fully deuterated versions thereof.





[0106] In addition to and / or in combination with the materials disclosed herein, many hole injection materials, hole transporting materials, host materials, dopant materials, exciton/hole blocking layer materials, electron transporting and electron injecting materials may be used in an OLED. Non-limiting examples of the materials that may be used in an OLED in combination with materials disclosed herein are listed in Table A below. Table A lists non-limiting classes of materials, non-limiting examples of compounds for each class, and references that disclose the materials.

TABLE A

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Hole injection materials		
Phthalocyanine and porphyrin compounds		Appl. Phys. Lett. 69, 2160 (1996)






EP 2 966 706 B1

(continued)

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Hole injection materials		
Starburst triarylamines		J. Lumin. 72-74, 985 (1997)
CF _x Fluorohydrocarbon polymer	$\left[\text{CH}_x\text{F}_y \right]_n$	Appl. Phys. Lett. 78, 673 (2001)
Conducting polymers (e.g., PEDOT:PSS, polyaniline, polythiophene)		Synth. Met. 87, 171 (1997) WO2007002683
Phosphonic acid and silane SAMs		US20030162053
Triaryamine or polythiophene polymers with conductivity dopants	 <p style="text-align: right;">and</p>	EP1725079A1

EP 2 966 706 B1

(continued)






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Hole injection materials		
<p>Organic compounds with conductive inorganic compounds, such as molybdenum and tungsten oxides</p>		<p>US20050123751 SID Symposium Digest, 37, 923 (2006) WO2009018009</p>
<p>n-type semiconducting organic complexes</p>		<p>US20020158242</p>
<p>Metal organometallic complexes</p>		<p>US20060240279</p>
<p>Cross-linkable compounds</p>		<p>US20080220265</p>
<p>Polythiophene based polymers and copolymers</p>		<p>WO 2011075644 EP2350216</p>

EP 2 966 706 B1

(continued)

Hole transporting materials			
5	Triarylamines (e.g., TPD, α -NPD)		Appl. Phys. Lett. 51, 913 (1987)
10			US5061569
15			EP650955
20			
25			J. Mater. Chem. 3, 319 (1993)
30			
35			
40			Appl. Phys. Lett. 90, 183503 (2007)
45			
50			Appl. Phys. Lett. 90, 183503 (2007)
55			









(continued)

Hole transporting materials		
5 Triarylamine on spirofluorene core		Synth. Met. 91, 209 (1997)
10 Arylamine carbazole compounds		Adv. Mater. 6, 677 (1994), US20080124572
25 Triarylamine with (di) benzothiophene/(di)ben zofuran		US20070278938, US20080106190 US20110163302
30 Indolocarbazoles		Synth. Met. 111, 421 (2000)
40 Isoindole compounds		Chem. Mater. 15, 3148 (2003)







50

55

(continued)

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
5 Metal carbene complexes		US20080018221
15 Phosphorescent OLED host materials		
15 Red hosts		
20 Arylcarbazoles		Appl. Phys. Lett. 78, 1622 (2001)
25 Metal 8-hydroxyquinolates (e.g., Alq ₃ , BALq)		Nature 395, 151 (1998)
30		US20060202194
35		WO2005014551
40		WO2006072002
45 50 Metal phenoxybenzothiazole compounds		Appl. Phys. Lett. 90, 123509 (2007)
55 Conjugated oligomers and polymers (e.g., polyfluorene)		Org. Electron. 1, 15 (2000)

(continued)








Phosphorescent OLED host materials		
Red hosts		
5 10	Aromatic fused rings 	WO2009066779, WO2009066778, WO2009063833, US20090045731, US20090045730, WO2009008311, US20090008605, US20090009065
15 20	Zinc complexes 	WO2010056066
25	Chrysene based compounds 	WO2011086863
Green hosts		
30 35	Arylcarbazoles 	Appl. Phys. Lett. 78, 1622 (2001)
40		US20030175553
45		WO2001039234

50








55

EP 2 966 706 B1







(continued)

Green hosts			
5	Aryltriphenylene compounds		US20060280965
10			US20060280965
15			WO2009021126
20			WO2009021126
25	Poly-fused heteroaryl compounds		US20090309488 US20090302743 US20100012931
30			
35			
40	Donor acceptor type molecules		WO2008056746
45			WO2010107244
50			
55			

(continued)

Green hosts		
5	<p>Aza-carbazole/DBT/DBF</p> 	JP2008074939
10		US20100187984
15		US20100187984
20		US20100187984
25	<p>Polymers (e.g., PVK)</p> 	Appl. Phys. Lett. 77, 2280 (2000)
30	<p>Spirofluorene compounds</p> 	WO2004093207
35	<p>Metal phenoxybenzoxazole compounds</p> 	WO2005089025
40		WO2005089025
45		WO2006132173
50		WO2006132173
55		JP200511610







(continued)

Green hosts			
5	Spiro fluorene-carbazole compounds		JP2007254297
10			JP2007254297
15	Indolocarbazoles		WO2007063796
20			WO2007063754
25	5-member ring electron deficient heterocycles (e.g., triazole, oxadiazole)		J. Appl. Phys. 90, 5048 (2001)
30			WO2004107822
35			
40			
45			







50

55

(continued)







Green hosts		
5 10 15	Tetraphenylene complexes 	US20050112407
20	Metal phenoxy pyridine compounds 	WO2005030900
25 30	Metal coordination complexes (e.g., Zn, Al with N^N ligands) 	US20040137268, US20040137267
Blue hosts		
35 40	Arylcarbazoles 	Appl. Phys. Lett, 82, 2422 (2003)
45		US20070 190359
50	Dibenzothiophene/ Dibenzofuran-carbazole compounds 	WO2006114966, US20090167162

(continued)

Blue hosts		
5 10		US20090 167162
15 20		WO2009086028
25		US20090030202, US20090017330
30 35		US20100084966
40	Silicon aryl compounds 	US20050238919
45 50		WO2009003898







55

(continued)

Blue hosts		
5 10	Silicon/Germanium aryl compounds 	EP2034538A
15	Aryl benzoyl ester 	WO2006100298
20 25	Carbazole linked by nonconjugated groups 	US20040115476
30	Aza-carbazoles 	US20060121308
35 40	High triplet metal organometallic complex 	US7154114
Phosphorescent dopants		
Red dopants		
45 50	Heavy metal porphyrins (e.g., PtOEP) 	Nature 395, 151 (1998)







55

(continued)






Phosphorescent dopants		
Red dopants		
5 10		Appl. Phys. Lett. 78, 1622 (2001)
15		US20030072964
20		US20030072964
25		US20060202194
30		US20060202194
35		US20070087321

55

(continued)

Phosphorescent dopants		
Red dopants		
5 10		<p>US20080261 076 US20100090591</p>
15 20		<p>US20070087321</p>
25	 <p>H₁₇C₈</p>	<p>Adv. Mater. 19, 739 (2007)</p>
30 35 40		<p>WO2009100991</p>
45 50		<p>WO2008101842</p>
55		<p>US7232618</p>

(continued)






Phosphorescent dopants		
Red dopants		
5 Platinum(II) organometallic complexes		WO2003040257
10 15 20		US20070103060
25 Osmium(III) complexes		Chem. Mater. 17, 3532 (2005)
30 Ruthenium(II) complexes		Adv. Mater. 17, 1059 (2005)
35 40 Rhenium (I), (II), and (III) complexes		US20050244673

45

50

55







(continued)

Green dopants		
5 10	Iridium(III) organometallic complexes  and its derivatives	Inorg. Chem. 40, 1704 (2001)
15 20		US20020034656
25 30		US7332232
35 40		US20090108737
45 50		WO2010028151








55

EP 2 966 706 B1







(continued)

Green dopants		
5		EP1841834B
10		
15		US20060127696
20		
25		US20090039776
30		
35		US6921915
40		
45		US20100244004
50		
55		US6687266






(continued)

Green dopants		
5		Chem. Mater. 16, 2480 (2004)
10		
15		US20070190359
20		US 20060008670 JP2007123392
25		WO2010086089, WO2011044988
30		
35		Adv. Mater. 16, 2003 (2004)
40		Angew. Chem. Int. Ed. 2006, 45, 7800
45		
50		WO2009050290

(continued)

Green dopants		
5		US20090165846
10		US20080015355
15		
20		US20010015432
25		
30		US20100295032
35		
40	Monomer for polymeric metal organometallic compounds	US7250226, US7396598
45		
50	Pt(II) organometallic complexes, including polydentate ligands	Appl. Phys. Lett. 86, 153505 (2005)
55		






(continued)

Green dopants		
5		Appl. Phys. Lett. 86, 153505 (2005)
10		
15		Chem. Lett. 34, 592 (2005)
20		
25		WO2002015645
30		
35		US20060263635
40		
45		US20060182992 US20070103060







50

55

(continued)

Green dopants			
5 10 15	Cu complexes		WO2009000673
20 25			US20070111026
30 35	Gold complexes		Chem. Commun. 2906 (2005)
40	Rhenium(III) complexes		Inorg. Chem. 42, 1248 (2003)
45 50 55	Osmium(II) complexes		US7279704

(continued)

Green dopants		
5 Deuterated organometallic complexes		US20030138657
15 Organometallic complexes with two or more metal centers		US20030152802
30		US7090928
Blue dopants		
40 Iridium(III) organometallic complexes		WO2002002714
45		WO2006009024
50		US20060251923 US20110057559 US20110204333
55		







EP 2 966 706 B1

(continued)

Blue dopants		
5 10	<p>The structure shows an iridium (Ir) atom coordinated to three 2-imidazolylphenyl ligands. Each ligand consists of a phenyl ring attached to the 2-position of an imidazole ring. The entire complex is enclosed in large square brackets with a subscript '3' at the bottom right.</p>	US7393599, WO2006056418, US20050260441, WO2005019373
15 20	<p>The structure shows an iridium (Ir) atom coordinated to three 2-(1H-benzofuran-2-yl)imidazole ligands. Each ligand features a benzofuran moiety attached to the 2-position of an imidazole ring. The complex is shown within large square brackets with a subscript '3' at the bottom right.</p>	US7534505
25 30	<p>The structure shows an iridium (Ir) atom coordinated to two 2-(1H-benzofuran-2-yl)imidazole ligands and one 2-(1H-benzofuran-2-yl)imidazole ligand. The two 2-(1H-benzofuran-2-yl)imidazole ligands are enclosed in large square brackets with a subscript '2' at the bottom right, and the single 2-(1H-benzofuran-2-yl)imidazole ligand is also enclosed in large square brackets. The entire complex is enclosed in a larger set of square brackets.</p>	WO2011051404
35 40	<p>The structure shows an iridium (Ir) atom coordinated to two 2-(1H-benzofuran-2-yl)imidazole ligands. The complex is shown within large square brackets with a subscript '2' at the bottom right. The iridium atom has a positive charge (+).</p>	US7445855
45 50	<p>The structure shows an iridium (Ir) atom coordinated to three 2-(1H-benzofuran-2-yl)imidazole ligands. Each ligand has a benzofuran moiety attached to the 2-position of an imidazole ring. The complex is shown within large square brackets with a subscript '3' at the bottom right.</p>	US20070190359, US20080297033 US20100148663

55







(continued)

Blue dopants		
5		US7338722
10		
15		US20020134984
20		
25		Angew. Chem. Int. Ed. 47, 4542 (2008)
30		
35		Chem. Mater. 18, 5119 (2006)
40		
45		Inorg. Chem. 46, 4308 (2007)
50		
55		WO2005123873





(continued)

Blue dopants			
5		WO2005123873	
10		WO2007004380	
15		WO2006082742	
20		WO2006082742	
25		WO2006082742	
30	Osmium(II) complexes		US7279704
35		US7279704	
40		Organometallics 23, 3745 (2004)	
45	Gold complexes		Appl. Phys. Lett. 74, 1361 (1999)
50	Platinum(II) complexes		WO2006098120, WO2006103874
55			

(continued)

Blue dopants			
5 10	Pt tetradentate complexes with at least one metal-carbene bond		US7655323
Exciton/hole blocking layer materials			
15 20	Bathocuprine compounds (e.g., BCP, BPhen)		Appl. Phys. Lett. 75, 4 (1999)
25			Appl. Phys. Lett. 79, 449 (2001)
30	Metal 8-hydroxyquinolates (e.g., BAlq)		Appl. Phys. Lett. 81, 162 (2002)
35 40	5-member ring electron deficient heterocycles such as triazole, oxadiazole, imidazole, benzimidazole		Appl. Phys. Lett. 81, 162 (2002)
45 50	Triphenylene compounds		US20050025993

(continued)







Exciton/hole blocking layer materials			
5 10 15	Fluorinated aromatic compounds		Appl. Phys. Lett. 79, 156 (2001)
20 25	Phenothiazine-S-oxide		WO2008132085
30	Silylated five-membered nitrogen, oxygen, sulfur or phosphorus dibenzoheterocycles		WO2010079051
35 40	Aza-carbazoles		US20060121308

45

50

55

(continued)




Electron transporting materials		
5 10	Anthracene-benzimidazole compounds 	WO2003060956
15 20		US20090179554
25 30	Aza triphenylene derivatives 	US20090115316
35	Anthracene-benzothiazole compounds 	Appl. Phys. Lett. 89, 063504 (2006)
40	Metal 8-hydroxyquinolates (e.g., Alq ₃ , Zrq ₄) 	Appl. Phys. Lett. 51, 913 (1987) US7230107
45 50	Metal hydroxybenzoquinolates 	Chem. Lett. 5, 905 (1993)

55

(continued)

Electron transporting materials		
5	Bathocuprine compounds such as BCP, BPhen, etc.	Appl. Phys. Lett. 91, 263503 (2007)
10		
15		Appl. Phys. Lett. 79, 449 (2001)
20	5-member ring electron deficient heterocycles (e.g., triazole, oxadiazole, imidazole, benzoimidazole)	Appl. Phys. Lett. 74, 865 (1999)
25		Appl. Phys. Lett. 55, 1489 (1989)
30		Jpn. J. Apply. Phys. 32, L917 (1993)
35		
40	Silole compounds	Org. Electron. 4, 113 (2003)
45	Arylborane compounds	J. Am. Chem. Soc. 120, 9714 (1998)
50		
55	Fluorinated aromatic compounds	J. Am. Chem. Soc. 122, 1832 (2000)

(continued)

Electron transporting materials			
5 10	Fullerene (e.g., C ₆₀)		US20090101870
15 20	Triazine complexes		US20040036077
25 30	Zn (N^N) complexes		US6528187

EXPERIMENTAL

Synthesis Examples

[0107] Chemical abbreviations used throughout this document are as follows:

SPhos is dicyclohexyl(2',6'-dimethoxy-[1,1'-biphenyl]-2-yl)phosphine,

Pd₂(dba)₃ is tri(dibenzylideneacetone)dipalladium(0),


Pd(PPh₃)₄ is tetrakis(triphenylphosphine) palladium (0),

DCM is dichloromethane,

EtOAc is ethyl acetate,


DME is dimethoxyethane, and

THF is tetrahydrofuran.

Synthesis of Compound A5Synthesis of 4-(3-bromo-5-chlorophenyl)dibenzo[*b,d*]thiophene5 **[0108]**

15 **[0109]** Dibenzo[*b,d*]thiophen-4-ylboronic acid (3.0 g, 13.15 mmol) and 1,3-dibromo-5-chlorobenzene (10.67 g, 39.5 mmol) were dissolved in toluene (150 ml) under a nitrogen atmosphere in a nitrogen-flushed 250 mL two-necked round-bottomed flask to give a colorless solution. K_2CO_3 (7.27 g, 52.6 mmol) in water (50 ml) was added to the reaction mixture, followed by $Pd(PPh_3)_4$ (0.304 g, 0.263 mmol). The reaction mixture was then heated to reflux under nitrogen overnight (~12 hours). After cooling to room temperature (~22 °C), the organic phase was isolated, the solvent was evaporated, and the unreacted 1,3-dibromo-5-chlorobenzene was distilled off under reduced pressure. The residue was subjected to column chromatography on the silica gel with heptanes/DCM (9/1, v/v) as the eluent, to obtain 4-(3-bromo-5-chlorophenyl)dibenzo[*b,d*] thiophene (3.5 g, 71.2 %) as a white solid.

20


25 Synthesis of 4-(5-chloro-[1,1':4',1''-terphenyl]-3-yl)dibenzo[*b,d*]thiophene**[0110]**

40 **[0111]** A solution of 4-(3-bromo-5-chlorophenyl)dibenzo[*b,d*]thiophene (4.0 g, 10.70 mmol), [1,1'-biphenyl]-4-ylboronic acid (2.120 g, 10.70 mmol), K_2CO_3 (3.0 g, 21.4 mmol) and $Pd(PPh_3)_4$ (0.37 g, 0.32 mmol) in toluene (150 ml) and water (50 ml) was refluxed under nitrogen overnight (~12 hours). After cooling to room temperature (~22 °C), the organic layer was isolated and the solvent was evaporated. The residue was purified by column chromatography on silica gel with heptane/DCM (4/1, v/v) as the eluent to isolate 4-(5-chloro-[1,1':4',1''-terphenyl]-3-yl)dibenzo[*b,d*]thiophene (1.4 g 29%) as a white solid.

45

Synthesis of 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane50 **[0112]**

55




15 **[0113]** A mixture of 4-(5-chloro-[1,1':4',1''-terphenyl]-3-yl)dibenzo[*b,d*]thiophene (1.40 g, 3.13 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (1.59 g, 6.26 mmol), potassium acetate (0.92 g, 9.40 mmol), SPhos (0.25 g, 0.61 mmol) and Pd₂(dba)₃ (0.11 g, 0.12 mmol) in dioxane (150 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the reaction mixture was diluted with EtOAc, washed with brine and water, and dried over Na₂SO₄. Upon evaporation of the solvent, the residue was purified by column chromatography on silica gel with heptanes/EtOAc (9/1, v/v) as the eluent to yield 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.1 g, 65%) as a white solid.

20

Synthesis of Compound A5

[0114]




40 **[0115]** A solution of 2-chloro-4,6-diphenyl-1,3,5-triazine (1.84 g, 6.87 mmol), 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.70 g, 6.87 mmol), Pd(PPh₃)₄ (0.16 g, 0.137 mmol), and K₂CO₃ (2.85 g, 20.61 mmol) in DME (150 ml), toluene (100 ml) and water (50 ml) was refluxed under nitrogen overnight (~12 hours). After cooling to room temperature (~ 22 °C), the solid was isolated by filtration, and washed successively with water, methanol, and EtOAc. The crude product was dissolved in hot toluene, filtered through a short plug of silica gel and recrystallized from toluene to yield **Compound A5** (3.1 g, 70%) as a white solid.

45 **Synthesis of Compound A11**

Synthesis of 4-(5-chloro-[1,1'-biphenyl]-3-yl)-6-phenyldibenzo[*b,d*]thiophene


[0116]



[0117] A solution of 3-bromo-5-chloro-1,1'-biphenyl (10 g, 37.4 mmol), (6-phenyldibenzo[*b,d*]thiophen-4-yl)boronic acid (11.37 g, 37.4 mmol), Pd(PPh₃)₄ (0.432 g, 0.374 mmol), and K₂CO₃ (10.33 g, 74.8 mmol) in toluene (150 ml) and water (30 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the organic phase was isolated and the solvent was evaporated. The residue was purified by column chromatography on silica gel with heptane/DCM (4/1, v/v) as the eluent to yield 4-(5-chloro-[1,1'-biphenyl]-3-yl)-6-phenyldibenzo[*b,d*]thiophene (12.1 g, 72.4 %) as a white solid.

Synthesis of 4,4,5,5-tetramethyl-2-(5-(6-phenyldibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-1,3,2-dioxaborolane


[0118]



[0119] A solution of 4-(5-chloro-[1,1'-biphenyl]-3-yl)-6-phenyldibenzo[*b,d*]thiophene (13.0 g, 29.1 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (14.77 g, 58.2 mmol), Pd₂(dba)₃ (0.20 g, 0.22 mmol), SPhos (0.35 g, 0.85 mmol), and potassium acetate (8.56 g, 87 mmol) in dioxane (200 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the reaction solution was quenched with water and extracted with EtOAc. The combined organic extracts were dried over Na₂SO₄ and the solvent was evaporated. The residue was purified by column chromatography on silica gel with heptane/EtOAc (9/1, v/v) as the eluent to yield 4,4,5,5-tetramethyl-2-(5-(6-phenyldibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-1,3,2-dioxaborolane (13.2 g, 84%) as a white solid.

Synthesis of Compound A11

[0120]




[0121] A solution of 4,4,5,5-tetramethyl-2-(5-(6-phenyldibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-1,3,2-dioxaborolane (3.55 g, 6.59 mmol), 2-chloro-4,6-diphenyl-1,3,5-triazine (1.765 g, 6.59 mmol), Pd(PPh₃)₄ (0.152 g, 0.132 mmol), and K₂CO₃ (1.822 g, 13.18 mmol) in toluene (100 ml), DME (100 ml) and water (50 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the organic layer was isolated, filtered through a short plug of silica gel and concentrated. The precipitate was collected, washed successively with heptane, ethanol, and heptane to yield **Compound A11** (3.9 g, 92%) as a white solid.

Synthesis of Compound A14

Synthesis of Compound A14

[0122]




15

[0123] A solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.44 g, 7.44 mmol), 2-(5-chloro-[1,1'-biphenyl]-3-yl)-4,6-diphenyl-1,3,5-triazine (2.92 g, 6.95 mmol), Pd₂(dba)₃ (0.159 g, 0.174 mmol), SPhos (0.4 g, 0.976 mmol), and K₃PO₄ (4.80 g, 20.9 mmol) in toluene (125 ml), DME (100 ml) and water (25 ml) was refluxed under nitrogen for 18 h. After cooling to room temperature (~22°C), the solid was collected by filtration, dissolved in boiling toluene (800 ml), and filtered through a short plug of silica gel. Upon evaporation off the solvent, **Compound A14** (3.50 g, 70%) was recrystallized from toluene to yield a white solid.

Synthesis of Compound A17

20 Synthesis of Compound A17

[0124]




35

[0125] A solution of 2-(6-([1,1'-biphenyl]-3-yl)dibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.51 g, 7.60 mmol), 2-(5-chloro-[1,1'-biphenyl]-3-yl)-4,6-diphenyl-1,3,5-triazine (2.9 g, 6.91 mmol), Pd₂(dba)₃ (0.190 g, 0.207 mmol), SPhos (.5 g, 1.220 mmol), and K₃PO₄ (4.77 g, 20.7 mmol) in toluene (125 ml), DME (100 ml) and water (20 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the solid was collected by filtration, dissolved in boiling toluene (800 ml) and filtered through a short plug of silica gel. Upon evaporation off the solvent, **Compound A17** (3.75 g, 76%) was recrystallized from toluene to yield a white solid.

Synthesis of Compound A32

Synthesis of 4-(5-chloro-[1,1'-biphenyl]-3-yl)dibenzo[*b,d*]thiophene

[0126]




55

[0127] A solution of 3-bromo-5-chloro-1,1'-biphenyl (14.8 g, 55.3 mmol), dibenzo[*b,d*]thiophen-4-ylboronic acid (12.62 g, 55.3 mmol), Pd(PPh₃)₄ (0.639 g, 0.553 mmol) and K₂CO₃ (15.29 g, 111 mmol) in toluene (150 ml) and water (30 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the organic phase was isolated. After evaporating off the solvent, the residue was purified by column chromatography on silica gel with hep-

tane/DCM (85/15,v/v) as the eluent to yield 4-(5-chloro-[1,1'-biphenyl]-3-yl)dibenzo[*b,d*]thiophene (15.4 g, 70%) as a white solid.

Synthesis of 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane


[0128]



[0129] A solution of 4-(5-Chloro-[1,1'-biphenyl]-3-yl)dibenzo[*b,d*]thiophene (11.88 g, 32.0 mmol), 4,4,4',4'',5,5,5',5''-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (16.27 g, 64.1 mmol), Pd₂(dba)₃ (280 mg), SPhos (0.32 g, 0.78 mmol), and potassium acetate (9.43 g, 96 mmol) in dioxane (200 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the reaction mixture was quenched with water and extracted with EtOAc. After evaporating off the solvent, the residue was purified by column chromatography on silica gel with heptane/EtOAc (9/1, v/v) as the eluent and recrystallization from heptane to yield 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (11.1 g, 74.9 %) as a white solid.

Synthesis of Compound A32

[0130]




[0131] A solution of 2-([1,1'-biphenyl]-4-yl)-4-chloro-6-phenyl-1,3,5-triazine (2.24 g, 6.52 mmol), 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.01 g, 6.52 mmol), Pd(PPh₃)₄ (0.151 g, 0.130 mmol) and K₂CO₃ (1.801 g, 13.03 mmol) in toluene (180 ml), DME (30 ml), and water (30 ml) was refluxed under nitrogen for 12 h. After cooling to room temperature (~ 22 °C), the solid was collected by filtration and washed successively with ethanol, water, and ethanol. The crude product was recrystallized from toluene to yield **Compound A32** (2.7 g, 64 %) as a white solid.

Synthesis of Compound A35

Synthesis of Compound A35

[0132]


EP 2 966 706 B1



15 **[0133]** A solution of 2,4-di([1,1'-biphenyl]-4-yl)-6-chloro-1,3,5-triazine (3.0 g, 7.14 mmol) and 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.30 g, 7.14 mmol), Pd(PPh₃)₄ (0.164 g, 0.14 mmol), and K₂CO₃ (1.975 g, 14.29 mmol) in toluene (100 ml), DME (100 ml), and water (50 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22 °C), the solid was collected by filtration and washed successively with ethanol, water, ethanol, and heptane. The crude product was dissolved in hot toluene, filtered through a short plug of silica gel and recrystallized from toluene to yield **Compound A35** (3.7 g, 72% yield) as a white solid.

20 **Synthesis of Compound A38**


Synthesis of 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine



35 **[0135]** A Grignard reagent solution prepared by refluxing 2-bromo-9,9-dimethyl-9H-fluorene (19.33 g, 70.8 mmol) and Mg (2.58 g, 106 mmol) in dry THF (100 ml) under nitrogen for 2 h was transferred dropwise into a solution of 2,4-dichloro-6-phenyl-1,3,5-triazine (8.0 g, 35.4 mmol) in dry THF (50 ml) at room temperature (~ 22°C). The reaction mixture was stirred under nitrogen overnight (~ 12 hours), quenched with concentrated HCl solution and extracted with EtOAc. The organic phase was isolated and the solvent was evaporated. The residue was purified by column chromatography on silica gel with heptane/DCM (9/1, v/v) as the eluent and recrystallization from heptane to yield 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (11 g, 81%) as yellow crystals.

40

Synthesis of Compound A38




[0137] A solution of 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (3.0 g, 7.82 mmol), 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.61 g, 7.82 mmol), Pd(PPh₃)₄ (0.18 g, 0.156 mmol) and K₂CO₃ (2.16 g, 15.63 mmol) in toluene (100 ml), DME (100 ml) and water (50 ml) was refluxed under nitrogen for 12 h. After cooling to room temperature (~ 22°C), the organic phase was isolated and the solvent was evaporated. The residue was purified by column chromatography on silica gel with heptane/DCM (6/4, v/v) as the eluent and trituration with heptane to yield **Compound A38** (4.3 g, 80%) as a white solid.

Synthesis of Compound A41

Synthesis of Compound A41

[0138]




[0139] A solution of 4,4,5,5-tetramethyl-2-(5-(6-phenyldibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-1,3,2-dioxaborolane (1.98 g, 3.68 mmol), 2-([1,1'-biphenyl]-3-yl)-4-chloro-6-phenyl-1,3,5-triazine (1.264 g, 3.68 mmol), Pd(PPh₃)₄ (0.085 g, 0.074 mmol), and K₂CO₃ (1.016 g, 7.35 mmol) in DME (150 ml) and water (5 ml) was refluxed under nitrogen for 12 h. After cooling to room temperature (~ 22°C), the solid was collected by filtration, washed successively with ethanol, water, ethanol, and heptane, then dissolved in boiling toluene and filtered through a short plug of silica gel. Upon evaporation off the solvent, **Compound A41** (2.3 g, 87%) was recrystallized from toluene to yield a white solid.

Synthesis of Compound A47

Synthesis of Compound A47


[0140]



[0141] A solution of 4,4,5,5-tetramethyl-2-(5-(6-phenyldibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-1,3,2-dioxaborolane (3.18 g, 5.91 mmol), 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (2.267 g, 5.91 mmol) Pd(PPh₃)₄ (0.136 g, 0.118 mmol) and potassium carbonate (1.632 g, 11.81 mmol) in toluene (30 ml), DME (100 ml) and water (20 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22°C), the organic layer was isolated and the solvent was evaporated. The residue was purified by column chromatography on silica gel with heptane/DCM(1/1, v/v) as the eluent to yield **Compound A47** (2.1 g, 47%) as a white solid.

Synthesis of Compound A110Synthesis of Compound A110

5 [0142]




15 [0143] A solution of 2-(5-(dibenzo[*b,d*]thiophen-4-yl)-[1,1'-biphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.70 g, 3.68 mmol), 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (1.43 g, 3.68 mmol), Pd(PPh₃)₄ (0.085 g, 0.074 mmol), and K₂CO₃ (1.02 g, 7.35 mmol) in DME (120 ml) and water (20 ml) was refluxed under nitrogen for 14 h. After cooling to room temperature (~22°C), the precipitate was collected by filtration, washed successively with ethanol, water, ethanol and heptane to yield **Compound A110** (2.1 g, 89% yield), as a white solid.

20

Synthesis of Compound A113Synthesis of Compound A113

25 [0144]




40 [0145] A solution of 2-(5-chloro-[1,1'-biphenyl]-3-yl)-4,6-diphenyl-1,3,5-triazine (3.4 g, 8.10 mmol), 2-(4-(dibenzo[*b,d*]thiophen-4-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.28 g, 8.50 mmol), Pd₂(dba)₃ (0.222 g, 0.243 mmol), SPhos (0.199 g, 0.486 mmol), and K₂CO₃ (3.36 g, 24.29 mmol) in toluene (16 ml), DME (48 ml), and water (16 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the solid was collected by filtration and triturated with ethanol. The crude product was dissolved in boiling toluene, then filtered through a short plug of silica gel, and recrystallized from toluene to yield **Compound A113** (4.25 g, 82%) as a white solid.

45

Synthesis of Compound A116Synthesis of 2-(3-bromo-5-chlorophenyl)-9,9-dimethyl-9H-fluorene

50

[0146]




EP 2 966 706 B1

[0147] A solution of (9,9-Dimethyl-9*H*-fluoren-2-yl)boronic acid (5.0 g, 21.0 mmol), 1,3-dibromo-5-chlorobenzene (14.19 g, 52.5 mmol), Pd(PPh₃)₄ (0.49 g, 0.42 mmol), and K₂CO₃ (5.80 g, 42.0 mmol) in toluene (200 ml) and water (50 ml) was refluxed under nitrogen for 18 h. After cooling to room temperature (~22°C), the organic layer was isolated and the excess of 1,3-dibromo-5-chlorobenzene was distilled off. The residue was purified by column chromatography on silica gel with heptane/DCM (9/1, v/v) as the eluent to yield 2-(3-bromo-5-chlorophenyl)-9,9-dimethyl-9*H*-fluorene (6.2 g, 77%) as a colorless crystalline solid.

Synthesis of 4-(3-chloro-5-(9,9-dimethyl-9*H*-fluoren-2-yl)phenyl)dibenzo[*b,d*]thiophene


[0148]



[0149] A solution of 2-(3-bromo-5-chlorophenyl)-9,9-dimethyl-9*H*-fluorene (7.7 g, 20.07 mmol), dibenzo[*b,d*]thiophen-4-ylboronic acid (4.58 g, 20.07 mmol), Pd(PPh₃)₄ (0.464 g, 0.401 mmol), and K₂CO₃ (5.55 g, 40.1 mmol) in DME (150 ml) and water (20 ml) was refluxed under nitrogen for 12 h. After cooling to room temperature (~22°C), the organic phase was isolated and the solvent was evaporated. The crude product was purified by column chromatography on silica gel with heptane/DCM (9/1 to 4/1, v/v) as the eluent to yield 4-(3-chloro-5-(9,9-dimethyl-9*H*-fluoren-2-yl)phenyl)dibenzo[*b,d*]thiophene (9.0 g, 92%) as a white crystalline solid.

Synthesis of 2-(3-(dibenzo[*b,d*]thiophen-4-yl)-5-(9,9-dimethyl-9*H*-fluoren-2-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane


[0150]



[0151] A solution of 4-(3-chloro-5-(9,9-dimethyl-9*H*-fluoren-2-yl)phenyl)dibenzo[*b,d*]thiophene (9.5 g, 19.51 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (9.91 g, 39.0 mmol), Pd₂(dba)₃ (0.268 g, 0.293 mmol), SPhos (0.240 g, 0.585 mmol), and potassium acetate (5.74 g, 58.5 mmol) in dioxane was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the reaction mixture was diluted with water and extracted with ethyl acetate. The combined organic extracts were filtered and evaporated. The crude product was purified by column chromatography on silica gel with heptane/DCM (1/1, v/v) as the eluent to yield 2-(3-(dibenzo[*b,d*]thiophen-4-yl)-5-(9,9-dimethyl-9*H*-fluoren-2-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (7.8 g, 69.1%) as a white crystalline solid.

Synthesis of Compound A116

[0152]



[0153] A solution of 2-(3-(dibenzo[*b,d*]thiophen-4-yl)-5-(9,9-dimethyl-9*H*-fluoren-2-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5.84 g, 10.09 mmol), 2-chloro-4,6-diphenyl-1,3,5-triazine (2.70 g, 10.09 mmol), Pd(PPh₃)₄ (0.233 g, 0.202 mmol), and K₂CO₃ (3.49 g, 25.2 mmol) in DME (100 ml), toluene (100 ml), and water (50 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the precipitate was collected by filtration, then washed successively with water, ethanol, and heptane to yield Compound A116 (5.5 g, 80%) as a white solid.

Synthesis of Compound B3

20

Synthesis of Compound B3

[0154]

25

30

35

[0155] A solution of 2-(5-(9,9-dimethyl-9*H*-fluoren-2-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3 g, 5.47 mmol), 2-chloro-4,6-diphenyl-1,3,5-triazine (1.464 g, 5.47 mmol), Pd(PPh₃)₄ (0.126 g, 0.109 mmol), and K₂CO₃ (1.512 g, 10.94 mmol) in toluene (75 ml), DME (75 ml), and water (20 ml) was refluxed under nitrogen for 6 h. After cooling to room temperature (~22°C), the precipitate was collected by filtration, then washed successively with water, ethanol, heptanes, and ethanol to yield **Compound B3** (2.8 g, 78%) as a white solid.

40

Synthesis of Compound B6

Synthesis of 2-(5-chloro-[1,1':4',1''-terphenyl]-3-yl)-9,9-dimethyl-9*H*-fluorene

45

[0156]

50

55


[0157] A solution of 2-(3-bromo-5-chlorophenyl)-9,9-dimethyl-9*H*-fluorene (5 g, 13.03 mmol), [1,1'-biphenyl]-4-ylbo-

EP 2 966 706 B1

ronic acid (2.58 g, 13.03 mmol), Pd(PPh₃)₄ (0.301 g, 0.261 mmol), and K₂CO₃ (5.40 g, 39.1 mmol) in DME (150 ml) and water (25 ml) was refluxed under nitrogen for 12 h. After cooling to room temperature (~22°C), the organic phase was isolated and the solvent was evaporated. The crude product was purified by column chromatography on silica gel with heptane/DCM (1/1, v/v) as the eluent and recrystallized from heptane to yield 2-(5-chloro-[1,1':4',1''-terphenyl]-3-yl)-9,9-dimethyl-9H-fluorene (3.6 g, 60.5%) as colorless crystals.

Synthesis of 2-(5-(9,9-dimethyl-9H-fluoren-2-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane


[0158]



[0159] A solution of 2-(5-chloro-[1,1':4',1''-terphenyl]-3-yl)-9,9-dimethyl-9H-fluorene (6.8 g, 14.88 mmol), 4,4,4',4',5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (7.56 g, 29.8 mmol), Pd₂(dba)₃ (0.273 g, 0.298 mmol), SPhos (0.244 g, 0.595 mmol), and potassium acetate (2.92 g, 29.8 mmol) in dioxane (100 ml) and DME (100 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the solid was filtered off. Upon evaporating off the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (1/1, v/v) as the eluent to yield 2-(5-(9,9-dimethyl-9H-fluoren-2-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5.0 g, 61.3%) as a white solid.

Synthesis of Compound B6

[0160]




[0161] A solution of 2-(5-(9,9-Dimethyl-9H-fluoren-2-yl)-[1,1':4',1''-terphenyl]-3-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5.17 g, 9.43 mmol), 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (3.62 g, 9.43 mmol), Pd(PPh₃)₄ (0.218 g, 0.189 mmol), and potassium carbonate (2.61 g, 18.85 mmol) in DME (75 ml), toluene (75 ml), and water (10 ml) was refluxed under nitrogen for 15 h. After cooling to room temperature (~22°C), the solid was collected by filtration, then washed successively with ethanol, water, ethanol, and heptane to yield **Compound B6** (4.2 g, 58%) as a white crystalline solid.

Synthesis of Compound B7

Synthesis of Compound B7

[0162]




15 **[0163]** A solution of 2-(3,5-bis(9,9-dimethyl-9H-fluoren-2-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (4 g, 6.80 mmol), 2-chloro-4,6-diphenyl-1,3,5-triazine (1.819 g, 6.80 mmol), Pd(PPh₃)₄ (0.079 g, 0.068 mmol), and K₂CO₃ (1.878 g, 13.59 mmol) in DME (75 ml), toluene (75 ml), and water (10 ml) was refluxed under nitrogen for 18 h. After cooling to room temperature (~22°C), the solid was collected by filtration, then washed successively with ethanol, water, ethanol and heptane to yield **Compound B7** (3.5 g, 74%) as a white crystalline solid.

20 **Synthesis of Compound C21**

Synthesis of 2-(dibenzo[*b,d*]selenophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

25 **[0164]**

30



35 **[0165]** Into a solution of dibenzo[*b,d*]selenophene (7 g, 30.3 mmol) in anhydrous THF (151 ml) a solution of sec-butyllithium (23.79 ml, 33.3 mmol) was added dropwise at -78 °C. The resulting mixture was stirred at this temperature for 2 h and warmed to room temperature (~22°C). After cooling the mixture to -78 °C, the mixture was quenched with 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (7.72 ml, 37.9 mmol) by syringe over ~1 minute, then gradually warmed to room temperature (~22°C) and stirred overnight (~12 hours). The resulting mixture was quenched with methanol and the solvent was removed *in vacuo*. The crude product was purified by column chromatography on silica gel with heptane/DCM (4/1 to 1/1, v/v) as the eluent to yield 2-(dibenzo[*b,d*]selenophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (7 g, 65%) as a yellow oil.


40

45 Synthesis of 4-phenyldibenzo[*b,d*]selenophene

50

[0166]

55




[0167] A solution of 2-(dibenzo[*b,d*]selenophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (7.0 g, 19.60 mmol), iodobenzene (2.62 ml, 23.52 mmol), Pd(PPh₃)₄ (0.453 g, 0.392 mmol), and K₂CO₃ (8.13 g, 58.8 mmol) in THF (78 ml)

EP 2 966 706 B1

and water (19.60 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the reaction mixture was partitioned with ethyl acetate and water. The organic phase was isolated, then washed with brine and dried over Na₂SO₄. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (9/1, v/v) as the eluent to yield 4-phenyldibenzo[*b,d*]selenophene (5.3 g, 88%) as a colorless oil.

Synthesis of 4,4,5,5-tetramethyl-2-(6-phenyldibenzo[*b,d*]selenophen-4-yl)-1,3,2-dioxaborolane


[0168]



[0169] A solution of 4-phenyldibenzo[*b,d*]selenophene (5.3 g, 17.25 mmol) in THF (108 ml) was cooled to -78 °C and treated slowly with a solution of *sec*-butyllithium 1.4 M (16.63 ml, 23.29 mmol) in cyclohexane. The resulting mixture was stirred at this -78 °C for 1h before being allowed to warm to room temperature (~22°C). The dark red solution was cooled to -78 °C and quenched with 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5.28 ml, 25.9 mmol) by syringe. The reaction mixture was allowed to warm gradually to room temperature (~22°C) and stirred for 16 h. The resulting mixture was quenched with methanol, then the solvent was removed *in vacuo*. The residue was dissolved in DCM, washed with water and brine, and then dried over Na₂SO₄. After evaporating the solvent, the crude product was recrystallized from heptane to yield 4,4,5,5-tetramethyl-2-(6-phenyldibenzo[*b,d*]selenophen-4-yl)-1,3,2-dioxaborolane (5 g, 67%) as a pale yellow solid.

Synthesis of Compound C21

[0170]




[0171] A solution of 4,4,5,5-tetramethyl-2-(6-phenyldibenzo[*b,d*]selenophen-4-yl)-1,3,2-dioxaborolane (2.0 g, 4.62 mmol), 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (1.882 g, 4.85 mmol), Pd(PPh₃)₄ (0.160 g, 0.139 mmol), and K₂CO₃ (1.914 g, 13.85 mmol) in DME (28 ml), toluene (9 ml), and water (9 ml) was refluxed under nitrogen for 8 h. After cooling to room temperature (~22°C), the solids were collected by filtration, then washed with water and ethanol, dissolved in boiling toluene, and finally filtered through a short plug of silica gel. After evaporating the solvent, **Compound C21** (2.6 g, 74%) recrystallized from toluene as a white solid.

Synthesis of Compound C23

Synthesis of Compound C23

[0172]




[0173] A solution of 4,4,5,5-tetramethyl-2-(3-(6-phenyldibenzo[b,d]thiophen-4-yl)phenyl)-1,3,2-dioxaborolane (3.0 g, 6.49 mmol), 2-([1,1'-biphenyl]-4-yl)-4-chloro-6-phenyl-1,3,5-triazine (2.454 g, 7.14 mmol), Pd(PPh₃)₄ (0.375 g, 0.324 mmol), and K₂CO₃ (2.69 g, 19.46 mmol) in toluene (13 ml), DME (39 ml), and water (13 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the solids were collected by filtration, then triturated with ethanol, dissolved in boiling toluene, and filtered through a short plug of silica gel. After evaporating the solvent, **Compound C23** (3.78 g, 91%) recrystallized from toluene as a white solid.

15

Synthesis of Compound C29

Synthesis of Compound C29

[0174]



[0175] A solution of 2-(4-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (2.75g, 8.00 mmol), 6-phenyl-dibenzo[b,d]thiophen-4-yl boronic acid (2.68 g, 8.80 mmol), Pd₂(dba)₃ (0.20 g, 0.22 mmol) and SPhos (0.40 g, 0.98 mmol), and K₃PO₄ (5.52 g, 24.00 mmol) in toluene (150 ml), DME (125 ml) and water (30 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the precipitate was collected by filtration and washed with water and DCM, before being dissolved in boiling toluene and filtered through a short plug of silica gel. After evaporating the solvent, **Compound C29** (2.42 g, 53%) was recrystallized from toluene to give a white solid.


35

40

Synthesis of Compound C47

Synthesis of Compound C47

[0176]




[0177] A suspension of 4,4,5,5-tetramethyl-2-(3-(6-phenyldibenzo[b,d]thiophen-4-yl)phenyl)-1,3,2-dioxaborolane (3.09 g, 6.69 mmol), 2-(4-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (2.3 g, 6.69 mmol), Pd₂(dba)₃ (0.123 g, 0.134 mmol),

and SPhos (0.110 g, 0.268 mmol) and K_3PO_4 (4.26 g, 20.07 mmol) in toluene (20ml), DME (30 ml), and water (10 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature ($\sim 22^\circ C$), the solid was collected by filtration, then dissolved in boiling toluene, filtered through a short plug of silica, and recrystallized from toluene to yield **Compound C47** (3.51g, 81%) as a white solid.

Synthesis of Compound C56

Synthesis of Compound C56

[0178]




[0179] A solution of 4,4,5,5-tetramethyl-2-(4-(6-phenyldibenzo[*b,d*]thiophen-4-yl)phenyl)-1,3,2-dioxaborolane (3.5 g, 7.57 mmol), 2-(3-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (2.169 g, 6.31 mmol), $Pd_2(dba)_3$ (0.17 g, 0.19 mmol), SPhos (0.23 g, 0.57 mmol), and K_3PO_4 (4.02 g, 18.9 mmol) in toluene (100 ml), DME (100 ml), and water (10 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature ($\sim 22^\circ C$), the reaction mixture was filtered through a plug of silica gel. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM(9/1 to 4/1, v/v) as eluent and recrystallization from DCM to yield **Compound C56** (2.2 g, 54%) as a white solid.

Synthesis of Compound C65

Synthesis of Compound C65

[0180]




[0181] A mixture solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.75 g, 8.11 mmol), 2-chloro-4,6-diphenyl-1,3,5-triazine (2.61 g, 9.73 mmol), $Pd_2(dba)_3$ (0.149 g, 0.162 mmol), SPhos (0.266 g, 0.649 mmol), and potassium phosphate hydrate (3.74 g, 16.22 mmol) in toluene (90 mL) and water (10 mL) was refluxed under nitrogen overnight (~ 12 hours). Upon completion, the toluene was evaporated and the mixture was extracted with dichloromethane (not completely soluble) and washed with brine and water. The organic layers were combined, dried over Na_2SO_4 , and concentrated under vacuum. The crude materials was triturated with ethanol and then with toluene to yield **Compound C65** (3.0 g, 65%) as a light-yellow solid.

Synthesis of Compound C68

Synthesis of Compound C68

5 [0182]




20 [0183] A solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (4 g, 8.65 mmol), 2-([1,1'-biphenyl]-4-yl)-4-chloro-6-phenyl-1,3,5-triazine (2.75 g, 8.00 mmol), Pd(PPh₃)₄ (0.28 g, 0.24 mmol), and K₂CO₃ (3.31 g, 24 mmol) in toluene (125 ml), DME (100 ml) and water (25 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the precipitate was collected by filtration then rinsed with toluene. The crude product was triturated successively with toluene and methanol, then sublimed under vacuum to yield **Compound C68** (4.25 g, 83%) as a white solid.

25

Synthesis of Compound C71

Synthesis of Compound C71

30 [0184]




45 [0185] A suspension of 2-([1,1'-biphenyl]-3-yl)-4-chloro-6-phenyl-1,3,5-triazine (2.1 g, 6.11 mmol), 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.82 g, 6.11 mmol), Pd(PPh₃)₄ (0.141 g, 0.122 mmol), and K₂CO₃ (2.53 g, 18.32 mmol) in DME (150 ml) and water (20 ml) was refluxed under nitrogen for 3 h. After cooling to room temperature (~22°C), the solid was collected by filtration, then washed with water and ethanol. The solid was dissolved in boiling toluene, filtered through a short plug of silica gel. After evaporating the solvent, the crude product was recrystallized from toluene to yield **Compound C71** (2.9 g, 74%) as a light-yellow solid.

50 **Synthesis of Compound C73**

Synthesis of Compound C73

55 [0186]




[0187] A solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[b,d]furan-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (4.2 g, 9.41 mmol), 2-(3-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (3.46 g, 10.07 mmol), Pd₂(dba)₃ (0.258 g, 0.282 mmol), SPhos (0.463 g, 1.129 mmol), and K₃PO₄ (6.49 g, 28.2 mmol) in toluene (125 ml), DME (100 ml), and water (25 ml) was refluxed under nitrogen for 18 h. After cooling to room temperature (~22°C), the solid was collected by filtration, dissolved in boiling toluene, filtered through a short plug of silica gel, and recrystallized from toluene to yield Compound C73 (4.5 g, 76%) as a white solid.

Synthesis of Compound C74

Synthesis of Compound C74

[0188]




[0189] A solution of 2,4-diphenyl-6-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1,3,5-triazine (2.64 g, 6.07 mmol), 4-([1,1'-biphenyl]-4-yl)-6-bromodibenzo[b,d]thiophene (2.53 g, 6.10 mmol), Pd₂(dba)₃ (0.139 g, 0.152 mmol), SPhos (0.187 g, 0.455 mmol), and K₃PO₄ (2.146 g, 10.11 mmol) in toluene (75 ml), DME (75 ml), and water (7.50 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22°C), the solid was collected by filtration, purified by column chromatography on silica gel with heptane/DCM (4/1 to 7/3, v/v) as eluent and recrystallization from heptane to yield **Compound C74** (2.0 g, 61%) as a white solid.

Synthesis of Compound C75

Synthesis of 4-([1,1'-biphenyl]-4-yl)dibenzo[b,d]selenophene


[0190]



[0191] A solution of 4-iodo-dibenzo[*b,d*]selenophene (10 g, 28.0 mmol), [1,1'-biphenyl]-4-ylboronic acid (8.32 g, 42.0 mmol), Pd(PPh₃)₄ (1.624 g, 1.400 mmol), and K₂CO₃ (7.74 g, 56.0 mmol) in DME (200 ml) and water (40 ml) was refluxed under nitrogen for 24 h. After cooling to room temperature (~22°C), the solid was collected by filtration, washed with water and heptane, then dissolved in boiling toluene and filtered through a short plug of silica gel. After evaporating the solvent, 4-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]selenophene (8.0 g, 74%) was recrystallized from toluene as a white solid.

Synthesis of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]selenophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane


[0192]



[0193] Into a solution of 4-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]selenophene (5.5 g, 14.35 mmol) in THF (150 ml) a solution of *sec*-butyl lithium (18.45 ml, 25.8 mmol) was added dropwise at -78 °C. The resulting mixture was stirred at -78 °C for 5 h before 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5.12 ml, 25.1 mmol) was added in one portion. The reaction mixture was gradually warmed to room temperature (~22°C) and stirred for 16 h before quenching with water. The resulting mixture was extracted with ethyl acetate, then dried over Na₂SO₄. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (4/1 to 3/2, v/v) as the eluent to yield 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]selenophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.6 g, 36%) as a white solid.

Synthesis of Compound C75

[0194]




[0195] A solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]selenophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.62 g, 5.15 mmol), 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (2 g, 5.15 mmol), Pd(PPh₃)₄ (0.179 g, 0.155 mmol), and K₂CO₃ (1.424 g, 10.30 mmol) in DME (150 ml), toluene (50 ml), and water (40 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the solid was collected by filtration, washed successively with water and heptane, then dissolved in boiling toluene and filtered through a short plug of silica gel. The crude product further purified by recrystallized successively from heptane and toluene to yield **Compound C75** (2.1 g, 59%) as white crystals.

Synthesis of Compound C83

Synthesis of Compound C83

[0196]




15 **[0197]** A suspension of 2-(4-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (2.409 g, 7.01 mmol), 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.7 g, 5.84 mmol), Pd₂(dba)₃ (0.107 g, 0.117 mmol), SPhos (0.096 g, 0.234 mmol), and K₂CO₃ (2.421 g, 17.52 mmol) in toluene (20 ml), DME (65 ml), and water (15 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~ 22°C), the reaction mixture was diluted with water. The solid was collected by filtration, washed with water and ethanol, redissolved in hot toluene, and filtered through a short plug of silica gel. After evaporating the solvent, the residue was recrystallized from EtOAc to yield **Compound C83** (3.2 g, 85%) as a white solid.

20

Synthesis of Compound C101

Synthesis of Compound C101

[0198]



40 **[0199]** A solution of 2-(6-([1,1'-biphenyl]-3-yl)dibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (4.03 g, 8.73 mmol), 2-(3-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (2.50 g, 7.27 mmol), Pd₂(dba)₃ (0.20 g, 0.22 mmol), SPhos (0.27 g, 0.65 mmol), and K₃PO₄ (4.63 g, 21.8 mmol) in toluene (100 ml), DME (100 ml), and water (10 ml) was refluxed under nitrogen overnight (~ 12 hours). After cooling to room temperature (~22°C), the reaction mixture was diluted with DCM and filtered through a plug of silica gel. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (4/1 to 3/2, v/v) as the eluent and recrystallization from DCM to yield **Compound C101** (1.6 g, 43%) as a white solid.

45


Synthesis of Compound C110

Synthesis of Compound C110

[0200]

50

55




15

[0201] A suspension of 2-(6-([1,1'-biphenyl]-3-yl)dibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.1 g, 6.70 mmol), 2-(4-chlorophenyl)-4,6-diphenyl-1,3,5-triazine (2.54 g, 7.37 mmol), Pd₂(dba)₃ (0.123 g, 0.134 mmol), and SPhos (0.110 g, 0.268 mmol), and K₃PO₄ (4.27 g, 20.11 mmol) in toluene (20.00 ml), DME (30.0 ml) and water (10 ml) was refluxed under nitrogen overnight. After cooling to room temperature, it was diluted with water and the solid was collected by filtration and washed with ethanol. The crude product was dissolved in boiling toluene and filtered through a short plug of silica gel. Upon evaporation off the solvent, **Compound C110** (4.2 g, 97 %) was recrystallized from toluene as a white solid.

Synthesis of Compound C119

Synthesis of Compound C119

[0202]




35

[0203] A solution of (6-phenyldibenzo[b,d]thiophen-4-yl)boronic acid (3.14 g, 10.32 mmol), 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (3.6 g, 9.38 mmol), Pd(PPh₃)₄ (0.217 g, 0.188 mmol), and K₂CO₃ (3.89 g, 28.1 mmol) in DME (180 ml) and water (50 ml) was refluxed under nitrogen for 14 h. After cooling to room temperature (~22°C), the solid was collected by filtration, washed successively with methanol, water, ethanol, ethyl acetate and heptane, then dissolved in dichloromethane and filtered through a short plug of silica gel. After evaporating the solvent, the crude product was triturated with ethanol and heptane to yield **Compound C119** (4.0 g, 70%) as a white solid.

Synthesis of Compound C131

Synthesis of 2-(4-chloro-2-heptyl)-4,6-bis(9,9-dimethyl-9H-fluoren-2-yl)-1,3,5-triazine

[0204]




[0205] A solution of 2,4-dichloro-6-(4-chlorophenyl)-1,3,5-triazine (5 g, 19.19 mmol), (9,9-dimethyl-9H-fluoren-2-yl)bo-

ronic acid (9.14 g, 38.4 mmol), Pd(PPh₃)₄ (0.444 g, 0.384 mmol), and K₂CO₃ (7.96 g, 57.6 mmol) in DME (150 ml) and water (15 ml) was refluxed under nitrogen for 13 h. After cooling to room temperature (~22°C), the organic phase was isolated. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (4/1, v/v) as the eluent to yield 2-(4-chlorophenyl)-4,6-bis(9,9-dimethyl-9H-fluoren-2-yl)-1,3,5-triazine (5.43 g, 49.1 %) as a white solid.

Synthesis of Compound C131

[0206]




[0207] A solution of 2-(4-chlorophenyl)-4,6-bis(9,9-dimethyl-9H-fluoren-2-yl)-1,3,5-triazine (5.43 g, 9.42 mmol), (6-phenyldibenzo[*b,d*]thiophen-4-yl)boronic acid (2.87 g, 9.42 mmol), Pd₂(dba)₃ (0.129 g, 0.141 mmol), SPhos (0.116 g, 0.283 mmol), and K₃PO₄ (4.34 g, 18.85 mmol) in DME (200 ml) and water (25 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the organic phase was isolated and purified by column chromatography on silica gel with heptane/DCM (1/1, v/v) as the eluent to yield **Compound C131** as a white solid.

Synthesis of Compound C134

Synthesis of Compound C134

[0208]




[0209] A solution of (6-([1,1'-biphenyl]-4-yl)dibenzo[*b,d*]thiophen-4-yl)boronic acid (2.25 g, 5.92 mmol), 2-chloro-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (2.4 g, 6.25 mmol), Pd(PPh₃)₄ (0.137 g, 0.118 mmol), and K₂CO₃ (2.453 g, 17.75 mmol) in DME (200 ml) and water (50 ml) was refluxed under nitrogen for 14 h. After cooling to room temperature (~22°C), the solid was collected by filtration, washed successively with methanol, water, ethanol, ethyl acetate and heptane, then dissolved in boiling toluene and filtered through a short plug of silica gel. After evaporating the solvent, the crude product was triturated with ethanol and heptane to yield **Compound C134** (3.0 g, 75%)

Synthesis of Compound C139

Synthesis of Compound C139

[0210]




15 **[0211]** A solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[b,d]furan-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (4.41 g, 9.89 mmol), 2-(3-chlorophenyl)-4-(9,9-dimethyl-9H-fluoren-2-yl)-6-phenyl-1,3,5-triazine (4.25 g, 9.24 mmol), Pd₂(dba)₃ (0.211 g, 0.231 mmol), SPhos (0.379 g, 0.924 mmol), and K₃PO₄ (6.38 g, 27.7 mmol) in toluene (125 ml), DME (100 ml), and water (30 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the reaction mixture was extracted with toluene. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/toluene (4/1 to 1/1, v/v) as eluent to yield **Compound C139** (4.1 g, 59.7%) as a white solid.

Synthesis of Compound C173

20 Synthesis of 2-(2,8-diphenyldibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

25 **[0212]**




35 **[0213]** A solution of *sec*-butyllithium in cyclohexane (28.5 ml, 39.9 mmol) was added dropwise into a solution of 2,8-diphenyldibenzo[b,d]thiophene (7.45 g, 22.14 mmol) in anhydrous THF at -78 °C. The reaction mixture was stirred at -78 °C for 2 h, while 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (7.21 g, 38.8 mmol) was added at a rate of 1 mL/min. The reaction mixture was gradually warmed to room temperature (~22°C) and stirred for 16 h before quenching with a 10% NH₄Cl aqueous solution. The resulting mixture was extracted with ethyl acetate. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (1/1, v/v) as the eluent and then recrystallized from heptane to yield 2-(2,8-diphenyldibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5.5 g, 53.7%) as white crystals.

40

Synthesis of Compound 173

45 **[0214]**



[0215] A solution of 2-(2,8-diphenyldibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.03 g, 6.55


EP 2 966 706 B1

mmol), 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (2.54 g, 6.55 mmol), Pd₂(dba)₃ (0.090 g, 0.098 mmol), SPhos (0.081 g, 0.197 mmol), and K₃PO₄ (3.02 g, 13.11 mmol) in DME (100 ml), toluene (100 ml), and water (10 ml) was refluxed under nitrogen for 16 h. After cooling to room temperature (~22°C), the solid was collected by filtration, washed successively with ethanol, water, ethanol and heptane, and then triturated with boiling toluene to yield **Compound 173** (4.0 g, 95%) as a white solid.

Synthesis of Compound C185

Synthesis of 2-(6,8-diphenyldibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane


[0216]



[0217] Sec-butyllithium in cyclohexane (38.2 ml, 53.5 mmol) was added dropwise into a solution of 2,4-diphenyldibenzo[*b,d*]thiophene (10 g, 29.7 mmol) in anhydrous THF at -78 °C. The reaction mixture was stirred at -78 °C for 2 h, while 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (10.61 ml, 52.0 mmol) was added at a rate of 1 mL/min. The reaction mixture was gradually warmed to room temperature (~22°C) and stirred for 16 h before being quenched with a 10% NH₄Cl aqueous solution. The resulting mixture was extracted with ethyl acetate. After evaporating the solvent, the residue was purified by column chromatography on silica gel with heptane/DCM (4/1 to 0/1, v/v) as the eluent, then recrystallized from heptane to yield 2-(6,8-diphenyldibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (9.5 g, 69%) as white crystals.

Synthesis of Compound C185

[0218]




[0219] A solution of 2-(6,8-diphenyldibenzo[*b,d*]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (4.29 g, 9.27 mmol), 2-(3-bromophenyl)-4,6-diphenyl-1,3,5-triazine (3 g, 7.73 mmol), Pd(PPh₃)₄ (0.269 g, 0.232 mmol), and K₂CO₃ (2.14 g, 15.45 mmol) in DME (200 ml) and water (40.0 ml) was refluxed under nitrogen for 5 h. After cooling to room temperature (~22°C), the solid was collected by filtration, dissolved in boiling xylene, then filtered through a short plug of silica gel and recrystallized from xylene to yield **Compound C185** (2.6 g, 52.3%) as a white solid.

Synthesis of Compound C251

Synthesis of Compound C251

[0220]




15

[0221] A solution of 2-(4'-(4-chloro-[1,1'-biphenyl]-3-yl)-4,6-diphenyl-1,3,5-triazine (3.0 g, 7.14 mmol), 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.53 g, 7.64 mmol), Pd₂(dba)₃ (0.196 g, 0.214 mmol), SPhos (0.40 g, 0.976 mmol), and K₃PO₄ (4.93 g, 21.43 mmol) in DME (80 ml), toluene (160 ml), and water (25 ml) was refluxed for 18 h. After cooling to room temperature (~22°C), the solid was collected by filtration, washed with water, then dissolved in boiling toluene and filtered through a short plug of silica gel, and recrystallized from toluene to yield **Compound C251** (3.43 g, 67%) as a white solid.

Synthesis of Compound C254

20 Synthesis of Compound C254

[0222]



35


[0223] A solution of 2-(6-([1,1'-biphenyl]-4-yl)dibenzo[b,d]thiophen-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.72 g, 5.89 mmol), 2-(3-bromophenyl)-4,6-bis(4-fluorophenyl)-1,3,5-triazine (2.5 g, 5.89 mmol), Pd₂(dba)₃ (0.081 g, 0.088 mmol), SPhos (0.073 g, 0.177 mmol), and K₃PO₄ (2.71 g, 11.79 mmol) in DME (200 ml) and water (50 ml) was refluxed under nitrogen for 18 h. After cooling to room temperature (~22°C), the solid was collected by filtration and purified by column chromatography on silica gel with heptane/DCM (1/1, v/v) as the eluent and recrystallized from heptane to yield **Compound C254** (2.5 g, 62%) as white crystals.

40 Device Examples


[0224] Application in OLED. All devices were fabricated by high vacuum (~1,33322x10⁻⁵ Pa (~10⁻⁷ Torr)) thermal evaporation. The anode electrode was 80 nm of indium tin oxide (ITO). The cathode electrode consisted of 1 nm of LiF followed by 100 nm of Al. All devices were encapsulated with a glass lid sealed with an epoxy resin in a nitrogen glove box (<1 ppm of H₂O and O₂) immediately after fabrication, and a moisture getter was incorporated inside the package.

[0225] Device Examples - Set 1. A first set of device examples have organic stacks consisting of, sequentially, from the ITO surface, 10 nm of LG101 (from LG Chem) as the hole injection layer (HIL), 45 nm of 4,4'-bis[N-(1-naphthyl)-N-phenylaminobiphenyl (NPD) as the hole-transport layer (HTL), and 30 nm of Compound E8 with 20 wt% of inventive compound (Compound C65) or comparative compound (CC-1) and 10 wt% of emitter GD as the emissive layer (EML). On top of the EML, 50 nm of Compound C65 or CC-1 was deposited as the hole blocking layer (HBL), followed by 40 nm of tris(8-hydroxyquinolino)aluminum (Alq₃) as the electron-transport layer (ETL). The structures of the compounds used are shown below.


5



NPD




Compound E8




GD

10


15



Compound C65



CC-1

Alq₃

20

[0226] Table D1, below, is a summary of the device data, voltage (V), external efficiency (EQE) and power efficiency (PE), recorded at 9000 nits for the Device Example 1.

Table D1

25

Device	EML	HBL	Color	V(V)	EQE (%)	PE (lm/W)
Device C-1	E8: CC-1: GD	CC-1	Green	8.1	14.5	20.5
Device 1	E8: C65: GD	C65	Green	7.6	15.1	22.5


30

[0227] The data in Table D1 shows that Device 1 using inventive compound Compound C65 as the co-host and HBL achieves higher efficiency at a lower driving voltage than Device C-1 using comparative compound CC-1 as the co-host and HBL.


[0228] Device Examples - Set 2. A second set of device examples have the same structure as that in Device Example 1 except that an inventive Compound C101 or a comparative compound CC-2 doped with 15% of GD is used as a two-component EML. The chemical structures of the inventive and comparative compounds used are presented below.

35

40



CC-2



Compound C101

45

[0229] Table D2, below, is a summary of the relative device data recorded at 9000 nits for the Device Example 2. Device lifetime LT97 is defined as the time it takes for devices to decay to 97% of their original luminance under a constant current density with an initial luminance of 9000 nits, and the values are normalized to that of Device C-2.

50

Table D2

55

Device	EML	HBL	Color	V (V)	EQE (%)	PE (lm/W)	LT97
Device C-2	CC-2: GD	CC-2	Green	7.5	14.7	22	100
Device 2	C101: GD	C101	Green	7.6	14.7	23	130


[0230] The data in Table D2 shows that Device 2 using inventive compound Compound C101 as the host and HBL achieves higher efficiency and longer lifetime than Device C-2 using comparative compound CC-2 as the host and HBL.

[0231] **Device Examples - Set 3.** A third set of device examples have the same structure as that in Device Example 1. The chemical structures of the inventive and comparative compounds used are presented below.

5

10

15



[0232] **Table D3**, below, is a summary of the relative device data recorded at 1000 nits for the Device Examples 3. Device lifetime LT95 defined as the time it takes for devices to decay to 95% of their original luminance under a constant current density with an initial luminance of 1000 nits, was calculated, with an acceleration factor of 1.8, from the values measured at a current density of 50 mA/cm², and normalized to that of Device C-3.

20

Table D3

25

Device	EML	HBL	Color	LT95
Device C-3	E8:CC-3:GD	CC-3	Green	100
Device 3	E8:A5:GD	A5	Green	694
Device 4	E8:A116:GD	A116	Green	529
Device 5	E8:C74:GD	C74	Green	537

30

[0233] The data in Table D3 show that, using as cohost with E8 in EML and as EBL, inventive compounds having substitutions on dibenzothiophene or bridging phenyl demonstrate longer lifetime than comparative compound that does not have these substitutions.


35

[0234] **Device Examples - Set 4.** A fourth set of devices have the same structure as that in Device Example 2. The chemical structures of the inventive and comparative compounds used are presented below.

40


45

50




55


5



,




,




,

10


15



,




,




,

20


25



,




,




,

30


35



,



,






,

45

50




55

5






15

20






25

30



35

40




45

50

55

5

10



[0235] Table D4, below, is a summary of the relative device data recorded at 1000 nits for the Device Example 4. Device lifetime LT95 was normalized to that of Device C-4.

15

Table D4

20

25

30

35

40

45

50

55

Device	EML	HBL	Color	LT95
Device C-4	CC-4:GD	CC-4	Green	100
Device C-5	CC-5:GD	CC-5	Green	3
Device C-6	CC-6:GD	CC-6	Green	3
Device C-7	CC-7:GD	CC-7	Green	147
Device C-8	CC-8:GD	CC-8	Green	93
Device C-9	CC-9:GD	CC-9	Green	62
Device C-10	CC-10:GD	CC-10	Green	88
Device 6	A10:GD	A10	Green	899
Device 7	A11:GD	A11	Green	602
Device 8	A14:GD	A14	Green	1029
Device 9	A17:GD	A17	Green	848
Device 10	A32:GD	A32	Green	969
Device 11	A38:GD	A38	Green	721
Device 12	A47:GD	A47	Green	859
Device 13	A110:GD	A110	Green	902
Device 14	A113:GD	A113	Green	1248
Device 15	B3:GD	B3	Green	1136
Device 16	B7:GD	B7	Green	849
Device 17	C23:GD	C23	Green	1466
Device 18	C29:GD	C29	Green	344
Device 19	C47:GD	C47	Green	915
Device 20	C56:GD	C56	Green	287
Device 21	C71:GD	C71	Green	181
Device 22	C83:GD	C83	Green	444
Device 23	C110:GD	C110	Green	477
Device 24	C119:GD	C119	Green	285
Device 25	C134:GD	C134	Green	192
Device 26	C140:GD	C140	Green	824

(continued)

Device	EML	HBL	Color	LT95
Device 27	C173:GD	C173	Green	760
Device 28	C251:GD	C251	Green	1830

[0236] The data in Table D4 show that OLED devices using inventive compounds as hosts and HBL have longer lifetime than that using comparative compounds.

[0237] The device data in Tables D1 through D4 together show that the inventive compounds having unique chemical structures are superior to comparative compounds when they are used as hosts or co-host and EBL in OLEDs. It is well accepted that OLED device performance is highly dependent on the materials properties, which are attributable to materials chemical structure.

Premixture Examples


[0238] The compatibility of selected h- and e-hosts was evaluated by compositional analysis of films fabricated by single-source co-evaporation of the premixture of these two components. A first set of potential premixtures of selected h- and e-hosts are presented in Table PM1.

Table PM1: Potential premixtures comprising selected h- and e-hosts


Premixtures	h-hosts	e-hosts
PM-A1	Compound E1	Compound C1
PM-A2	Compound E2	Compound C2
PM-A3	Compound E5	Compound C65
PM-A4	Compound E8	Compound C74
PM-A5	Compound E11	Compound C74
PM-A6	Compound E17	Compound C74
PM-A7	Compound E8	CC-1
PM-A8	Compound E8	Compound A5
PM-A9	Compound E8	Compound C17
PM-A10	Compound E17	Compound A5
PM-A11	Compound E25	CC-1
PM-A12	Compound E26	Compound C74
PM-A13	Compound E26	Compound C248
PM-A14	Compound E28	Compound C74
PM-A15	Compound E29	Compound C74
PM-A16	Compound E30	Compound C74

[0239] **Premixture Example - Set 1:** For premixture PM-A4, Compound E8 and Compound C74 were provided at a weight ratio of 7:3, then they were physically mixed, grinded and loaded into an evaporation source. The premixed compositions were thermally co-evaporated at a rate of 2 Å/s in a vacuum chamber under a pressure less than $1,33322 \times 10^{-5}$ Pa (10^{-7} Torr), and deposited onto glass substrates. The substrates were replaced continuously after deposition of 500 Å of film without stopping the deposition and cooling the source. The compositions of films were analyzed by high-performance liquid chromatography (HPLC) and the results are shown in Table 2.

5



Compound E8



Compound C74

10

15

Table PM2: HPLC composition (%) of sequentially deposited films from premixture (PM-A4) comprising Compound E8 and Compound C74 with weight ratio 7:3. (HPLC Conditions C18, 100 45 min, Detected wavelength 254 nm)
(Due to different absorption coefficients, the HPLC composition may or may not agree with the weight ratio.)

20

Films	Compound E8	Compound C74
Plate1	69.5	30.5
Plate2	68.4	31.6
Plate3	68.2	31.8
Plate4	68.2	31.8
Plate5	68.4	31.6
Plate6	69.3	30.7
Plate7	70.6	29.4
Plate8	71.7	28.3
Plate9	73.0	27.0

30


35

[0240] As shown in Table PM2, the composition of the components Compound E8 and Compound C74 did not change significantly from plate 1 through plate 9. The minor fluctuations in the concentrations do not reveal any trend and can be explained by the accuracy of HPLC analysis. Normally, the change of the concentration before and after depositions within 5% throughout the process is considered to be good and useful for commercial OLED application.


40

[0241] Premixture Example - Set 2: For premixture PM-A12, Compound E26 and Compound C74 were provided at a weight ratio of 3:2, then they were physically mixed, grinded, and loaded into an evaporation source. The premixed compositions were thermally co-evaporated at a rate of 2 Å/s in a vacuum chamber under a pressure less than 1.33322×10^{-5} Pa (10^{-7} Torr), and deposited onto glass substrates. The substrates were replaced continuously after deposition of 500 Å of film without stopping the deposition and cooling the source. The compositions of films were analyzed by high-performance liquid chromatography (HPLC) and the results are shown in Table PM3.

45



Compound E26



Compound C74

50

55

Table PM3: HPLC composition (%) of sequentially deposited films from premixture (PM-A12) comprising Compound E26 and Compound C74 with weight ratio 3:2. (HPLC Conditions C18, 100 45 min, Detected wavelength 254 nm)
(Due to different absorption coefficients, the HPLC composition may or may not agree with the weight ratio.)

Films	Compound E26	Compound C74
Plate1	67.1	32.9
Plate2	67.3	32.7
Plate3	68.4	31.6
Plate4	69.6	30.4
Plate5	70.8	29.2
Plate6	71.9	28.1
Plate7	72.9	27.1
Plate8	73.9	26.1

[0242] Again, the results of plates 1 to 8 show only minor variations and would be considered to be good and useful for commercial OLED application.

[0243] **Premixture Example - Set 3.** For premixture PM-A16, Compound E30 and Compound C74 were provided at a weight ratio of 1:1, then they were physically mixed, grinded and loaded into an evaporation source. The premixed compositions were thermally co-evaporated at a rate of 2 Å/s in a vacuum chamber under a pressure less than $1,33322 \times 10^{-5}$ Pa (10^{-7} Torr), and deposited onto glass substrates. The substrates were replaced continuously after deposition of 500 Å of film without stopping the deposition and cooling the source. The compositions of films were analyzed by high-performance liquid chromatography (HPLC) and the results are shown in Table PM4.




Table PM4: HPLC composition (%) of sequentially deposited films from premixture (PM-A16) comprising Compound E30 and Compound C74 with weight ratio 1:1. (HPLC Conditions C18, 100 45 min, Detected wavelength 254 nm)
(Due to different absorption coefficients, the HPLC composition may or may not agree with the weight ratio.)

Films	Compound E30	Compound C74
Plate1	52.3	47.7
Plate2	51.6	48.4
Plate3	52.1	47.9
Plate4	52.9	47.1
Plate5	53.9	46.1
Plate6	54.6	45.4
Plate7	51.8	48.2

[0244] The data in Tables PM2, PM3 and PM4 show that the ratio of the two components in premixtures PM-A4, PM-A12 and PM-A16 does not change significantly over a continuous single-source coevaporation. The minor fluctuations in the concentrations do not reveal any trend and can be explained by the accuracy of HPLC analysis. Normally, the

change of the concentration before and after depositions within 5% throughout the process is considered to be good and useful for commercial OLED application. These experiments conclude that PM-A4, PM-A12 and PM-A16 are stable premixtures for coevaporation. The coevaporation stability of these premixtures is believed to trackable to the unique chemical structures associated with these two classes of materials.

5


Claims

1. A composition of materials comprising a first compound and a second compound, wherein the first compound has the formula:

10

15

20



, Formula I;

25

wherein G^1 is selected from the group consisting of dibenzofuran, dibenzothiophene, dibenzoselenophene, and fluorene;

wherein L^1 , L^2 and L^3 are each independently selected from the group consisting of direct bond, phenyl, biphenyl, terphenyl, pyridine, pyrimidine, and combinations thereof;

30

wherein G^4 is selected from the group consisting of phenyl, biphenyl, terphenyl, naphthalene, phenanthrene, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phenanthroline, and combinations thereof;

wherein G^2 , G^3 , and G^5 are each independently selected from the group consisting of phenyl, biphenyl, terphenyl, fluorene, naphthalene, phenanthrene, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phenanthroline, aza-fluorene, and combinations thereof;

35

wherein G^2 , G^3 , G^4 , and G^5 are each optionally further substituted with one or more unfused substituents selected from the group consisting of deuterium, alkyl, alkoxy, cycloalkyl, cycloalkoxy, halogen, nitro, nitrile, silyl, phenyl, biphenyl, terphenyl, pyridine, and combinations thereof;

wherein m is an integer from 0 to 7,

wherein n is an integer from 0 to 4;

wherein, when m or n is larger than 1, then each G^4 or G^5 can be same or different;

40


wherein, when n is 0, and m is equal to or greater than 1, then each G^4 is selected from the group consisting of phenyl, and biphenyl;

wherein, when n is equal to or greater than 1, then L^1 is not a direct bond; and

wherein, when m and n are both 0, then L^1 is biphenyl,

and wherein the second compound has the formula II:

45



, Formula II;

50

wherein Ar^1 is selected from the group consisting of triphenylene, and aza-triphenylene;

wherein Ar^2 is selected from the group consisting of a direct bond, phenyl, biphenyl, terphenyl, naphthalene, pyridine, dibenzofuran, dibenzothiophene, dibenzoselenophene, aza-dibenzofuran, aza-dibenzothiophene, aza-dibenzoselenophene, and combinations thereof;

55

wherein Ar^3 is selected from the group consisting of benzene, biphenyl, terphenyl, naphthalene, pyridine, dibenzofuran, dibenzothiophene, dibenzoselenophene, aza-dibenzofuran, aza-dibenzothiophene, aza-dibenzoselenophene, carbazole, aza-carbazole, and combinations thereof; and

wherein Ar^1 , Ar^2 and Ar^3 are each, independently, optionally further substituted with one or more substitutions


selected from the group consisting of deuterium, halogen, alkyl, aryl, heteroaryl, and combinations thereof.

2. The composition of Claim 1, wherein n is 0.

5 3. The composition of Claim 1, wherein n is equal to or greater than 1.

4. The composition of Claim 1, wherein m and n are both 0.

10 5. The composition of Claim 1, wherein G¹ has the structure selected from the group consisting of:




wherein X is selected from a group consisting of O, S and Se;

wherein R^{B1} and R^{B2} are independently selected from a group consisting of hydrogen, deuterium, alkyl, cycloalkyl, alkoxy, aryl, heteraryl, halogen, and combinations thereof; and

wherein R^{B1} and R^{B2} are optionally joined to form a ring.

20 6. The composition of Claim 1, wherein at least one of G², G³, G⁴ and G⁵ is substituted with at least one fluorine atom.

7. The composition of Claim 1, wherein the first compound has the formula:



wherein X is selected from a group consisting of O, S and Se.

8. The composition of Claim 1, wherein the first compound is selected from the group consisting of:


40

45

50


55

Compound A1 through A3, each represented by the formula




wherein in Compound A1: X=O,
in Compound A2: X=S,
in Compound A3: X=Se

Compound A4 through A6, each represented by the formula




wherein in Compound A4: X=O,
in Compound A5: X=S,
in Compound A6: X=Se

Compound A7 through A9, each represented by the formula




wherein in Compound A7: X=O,
in Compound A8: X=S,
in Compound A9: X=Se

Compound A10 through A12, each represented by the formula




wherein in Compound A10: X=O,
in Compound A11: X=S,
in Compound A12: X=Se

Compound A13 through A15, each represented by the formula



wherein in Compound A13: X=O,
in Compound A14: X=S,
in Compound A15: X=Se


Compound A16 through A18, each represented by the formula



wherein in Compound A16: X=O,
in Compound A17: X=S,
in Compound A18: X=Se

Compound A19 through A21, each represented by the formula


5



10


wherein in Compound A19: X=O,
in Compound A20: X=S,
in Compound A21: X=Se

Compound A22 through A24, each represented by the formula



wherein in Compound A22: X=O,
in Compound A23: X=S,
in Compound A24: X=Se


Compound A25 through A27, each represented by the formula



wherein in Compound A25: X=O,
in Compound A26: X=S,
in Compound A27: X=Se

Compound A28 through A30, each represented by the formula


15



20


wherein in Compound A28: X=O,
in Compound A29: X=S,
in Compound A30: X=Se

Compound A31 through A33, each represented by the formula



wherein in Compound A31: X=O,
in Compound A32: X=S,
in Compound A33: X=Se


Compound A34 through A36, each represented by the formula



wherein in Compound A34: X=O,
in Compound A35: X=S,
in Compound A36: X=Se

Compound A37 through A39, each represented by the formula


30



35


wherein in Compound A37: X=O,
in Compound A38: X=S,
in Compound A39: X=Se

Compound A40 through A42, each represented by the formula



wherein in Compound A40: X=O,
in Compound A41: X=S,
in Compound A42: X=Se


Compound A43 through A45, each represented by the formula



wherein in Compound A43: X=O,
in Compound A44: X=S,
in Compound A45: X=Se

Compound A46 through A48, each represented by the formula


40



45


wherein in Compound A46: X=O,
in Compound A47: X=S,
in Compound A48: X=Se

Compound A49 through A51, each represented by the formula



wherein in Compound A49: X=O,
in Compound A50: X=S,
in Compound A51: X=Se

Compound A52 through A54, each represented by the formula




wherein in Compound A52: X=O,
in Compound A53: X=S,
in Compound A54: X=Se

50


55

Compound A55 through A57, each represented by the formula




wherein in Compound A55: X=O,
in Compound A56: X=S,
in Compound A57: X=Se

Compound A58 through A60, each represented by the formula




wherein in Compound A58: X=O,
in Compound A59: X=S,
in Compound A60: X=Se

Compound A61 through A63, each represented by the formula




wherein in Compound A61: X=O,
in Compound A62: X=S,
in Compound A63: X=Se

Compound A64 through A66, each represented by the formula




wherein in Compound A64: X=O,
in Compound A65: X=S,
in Compound A66: X=Se

Compound A67 through A69, each represented by the formula




wherein in Compound A67: X=O,
in Compound A68: X=S,
in Compound A69: X=Se

Compound A70 through A72, each represented by the formula




wherein in Compound A70: X=O,
in Compound A71: X=S,
in Compound A72: X=Se

Compound A73 through A75, each represented by the formula




wherein in Compound A73: X=O,
in Compound A74: X=S,
in Compound A75: X=Se

Compound A76 through A78, each represented by the formula




wherein in Compound A76: X=O,
in Compound A77: X=S,
in Compound A78: X=Se

Compound A79 through A81, each represented by the formula




wherein in Compound A79: X=O,
in Compound A80: X=S,
in Compound A81: X=Se

Compound A82 through A84, each represented by the formula




wherein in Compound A82: X=O,
in Compound A83: X=S,
in Compound A84: X=Se

Compound A85 through A87, each represented by the formula



wherein in Compound A85: X=O,
in Compound A86: X=S,
in Compound A87: X=Se


Compound A88 through A90, each represented by the formula



wherein in Compound A88: X=O,
in Compound A89: X=S,
in Compound A90: X=Se


EP 2 966 706 B1

Compound A91 through A93, each represented by the formula




wherein in Compound A91: X=O,
in Compound A92: X=S,
in Compound A93: X=Se

Compound A94 through A96, each represented by the formula




wherein in Compound A94: X=O,
in Compound A95: X=S,
in Compound A96: X=Se

Compound A97 through A99, each represented by the formula




wherein in Compound A97: X=O,
in Compound A98: X=S,
in Compound A99: X=Se

Compound A100 through A102, each represented by the formula




wherein in Compound A100: X=O,
in Compound A101: X=S,
in Compound A102: X=Se

Compound A103 through A105, each represented by the formula




wherein in Compound A103: X=O,
in Compound A104: X=S,
in Compound A105: X=Se

Compound A106 through A108, each represented by the formula




wherein in Compound A106: X=O,
in Compound A107: X=S,
in Compound A108: X=Se

Compound A109 through A111, each represented by the formula




wherein in Compound A109: X=O,
in Compound A110: X=S,
in Compound A111: X=Se

Compound A112 through A114, each represented by the formula




wherein in Compound A112: X=O,
in Compound A113: X=S,
in Compound A114: X=Se


Compound A115 through A117, each represented by the formula




wherein in Compound A115: X=O,
in Compound A116: X=S,
in Compound A117: X=Se




Compound B1



Compound B2




Compound B3




9. The composition of Claim 1, wherein the first compound is selected from the group consisting of:

30 Compound C1 through C3, each represented by the formula Compound C4 through C6, each represented by the formula Compound C7 through C9, each represented by the formula




40 wherein in Compound C1: X=O,
in Compound C2: X=S,
in Compound C3: X=Se , wherein in Compound C4: X=O,
in Compound C5: X=S,
in Compound C6: X=Se , wherein in Compound C7: X=O,
in Compound C8: X=S,
in Compound C9: X=Se ,

45 Compound C10 through C12, each represented by the formula Compound C13 through C15, each represented by the formula Compound C16 through C18, each represented by the formula




55 wherein in Compound C10: X=O,
in Compound C11: X=S,
in Compound C12: X=Se , wherein in Compound C13: X=O,
in Compound C14: X=S,
in Compound C15: X=Se , wherein in Compound C16: X=O,
in Compound C17: X=S,
in Compound C18: X=Se ,

Compound C19 through C21, each represented by the formula




wherein in Compound C19: X=O,
in Compound C20: X=S,
in Compound C21: X=Se

Compound C22 through C24, each represented by the formula




wherein in Compound C22: X=O,
in Compound C23: X=S,
in Compound C24: X=Se

Compound C25 through C27, each represented by the formula




wherein in Compound C25: X=O,
in Compound C26: X=S,
in Compound C27: X=Se

Compound C28 through C30, each represented by the formula




wherein in Compound C28: X=O,
in Compound C29: X=S,
in Compound C30: X=Se

Compound C31 through C33, each represented by the formula




wherein in Compound C31: X=O,
in Compound C32: X=S,
in Compound C33: X=Se

Compound C34 through C36, each represented by the formula




wherein in Compound C34: X=O,
in Compound C35: X=S,
in Compound C36: X=Se

Compound C37 through C39, each represented by the formula




wherein in Compound C37: X=O,
in Compound C38: X=S,
in Compound C39: X=Se

Compound C40 through C42, each represented by the formula




wherein in Compound C40: X=O,
in Compound C41: X=S,
in Compound C42: X=Se

Compound C43 through C45, each represented by the formula




wherein in Compound C43: X=O,
in Compound C44: X=S,
in Compound C45: X=Se

Compound C46 through C48, each represented by the formula




wherein in Compound C46: X=O,
in Compound C47: X=S,
in Compound C48: X=Se

Compound C49 through C51, each represented by the formula



wherein in Compound C49: X=O,
in Compound C50: X=S,
in Compound C51: X=Se


Compound C52 through C54, each represented by the formula



wherein in Compound C52: X=O,
in Compound C53: X=S,
in Compound C54: X=Se


Compound C55 through C57, each represented by the formula

5




wherein in Compound C55: X=O,
in Compound C56: X=S,
in Compound C57: X=Se

Compound C58 through C60, each represented by the formula



wherein in Compound C58: X=O,
in Compound C59: X=S,
in Compound C60: X=Se


Compound C61 through C63, each represented by the formula



wherein in Compound C61: X=O,
in Compound C62: X=S,
in Compound C63: X=Se


Compound C64 through C66, each represented by the formula

15




wherein in Compound C64: X=O,
in Compound C65: X=S,
in Compound C66: X=Se

Compound C67 through C69, each represented by the formula



wherein in Compound C67: X=O,
in Compound C68: X=S,
in Compound C69: X=Se


Compound C70 through C72, each represented by the formula



wherein in Compound C70: X=O,
in Compound C71: X=S,
in Compound C72: X=Se


Compound C73 through C75, each represented by the formula

30




wherein in Compound C73: X=O,
in Compound C74: X=S,
in Compound C75: X=Se

Compound C76 through C78, each represented by the formula



wherein in Compound C76: X=O,
in Compound C77: X=S,
in Compound C78: X=Se


Compound C79 through C81, each represented by the formula



wherein in Compound C79: X=O,
in Compound C80: X=S,
in Compound C81: X=Se


Compound C82 through C84, each represented by the formula

40




wherein in Compound C82: X=O,
in Compound C83: X=S,
in Compound C84: X=Se

Compound C85 through C87, each represented by the formula



wherein in Compound C85: X=O,
in Compound C86: X=S,
in Compound C87: X=Se

Compound C88 through C90, each represented by the formula




wherein in Compound C88: X=O,
in Compound C89: X=S,
in Compound C90: X=Se

50


55

Compound C91 through C93, each represented by the formula




wherein in Compound C91: X=O,
in Compound C92: X=S,
in Compound C93: X=Se

Compound C94 through C96, each represented by the formula




wherein in Compound C94: X=O,
in Compound C95: X=S,
in Compound C96: X=Se

Compound C97 through C99, each represented by the formula




wherein in Compound C97: X=O,
in Compound C98: X=S,
in Compound C99: X=Se

Compound C100 through C102, each represented by the formula




wherein in Compound C100: X=O,
in Compound C101: X=S,
in Compound C102: X=Se

Compound C103 through C105, each represented by the formula




wherein in Compound C103: X=O,
in Compound C104: X=S,
in Compound C105: X=Se

Compound C106 through C108, each represented by the formula




wherein in Compound C106: X=O,
in Compound C107: X=S,
in Compound C108: X=Se

Compound C109 through C111, each represented by the formula




wherein in Compound C109: X=O,
in Compound C110: X=S,
in Compound C111: X=Se

Compound C112 through C114, each represented by the formula




wherein in Compound C112: X=O,
in Compound C113: X=S,
in Compound C114: X=Se

Compound C115 through C117, each represented by the formula




wherein in Compound C115: X=O,
in Compound C116: X=S,
in Compound C117: X=Se

Compound C118 through C120, each represented by the formula




wherein in Compound C118: X=O,
in Compound C119: X=S,
in Compound C120: X=Se

Compound C121 through C123, each represented by the formula



wherein in Compound C121: X=O,
in Compound C122: X=S,
in Compound C123: X=Se


Compound C124 through C126, each represented by the formula



wherein in Compound C124: X=O,
in Compound C125: X=S,
in Compound C126: X=Se


EP 2 966 706 B1

Compound C127 through C129, each represented by the formula




wherein in Compound C127: X=O,
in Compound C128: X=S,
in Compound C129: X=Se

Compound C130 through C132, each represented by the formula




wherein in Compound C130: X=O,
in Compound C131: X=S,
in Compound C132: X=Se

Compound C133 through C135, each represented by the formula




wherein in Compound C133: X=O,
in Compound C134: X=S,
in Compound C135: X=Se

Compound C136 through C138, each represented by the formula




wherein in Compound C136: X=O,
in Compound C137: X=S,
in Compound C138: X=Se

Compound C139 through C141, each represented by the formula




wherein in Compound C139: X=O,
in Compound C140: X=S,
in Compound C141: X=Se

Compound C142 through C144, each represented by the formula




wherein in Compound C142: X=O,
in Compound C143: X=S,
in Compound C144: X=Se

Compound C145 through C147, each represented by the formula




wherein in Compound C145: X=O,
in Compound C146: X=S,
in Compound C147: X=Se

Compound C148 through C150, each represented by the formula




wherein in Compound C148: X=O,
in Compound C149: X=S,
in Compound C150: X=Se

Compound C151 through C153, each represented by the formula




wherein in Compound C151: X=O,
in Compound C152: X=S,
in Compound C153: X=Se

Compound C154 through C156, each represented by the formula




wherein in Compound C154: X=O,
in Compound C155: X=S,
in Compound C156: X=Se

Compound C157 through C159, each represented by the formula



wherein in Compound C157: X=O,
in Compound C158: X=S,
in Compound C159: X=Se

Compound C160 through C162, each represented by the formula




wherein in Compound C160: X=O,
in Compound C161: X=S,
in Compound C162: X=Se

EP 2 966 706 B1

Compound C163 through C165, each represented by the formula

5




10

wherein in Compound C163: X=O,
in Compound C164: X=S,
in Compound C165: X=Se

15


Compound C166 through C168, each represented by the formula



wherein in Compound C166: X=O,
in Compound C167: X=S,
in Compound C168: X=Se

Compound C175 through C177, each represented by the formula

Compound C169 through C171, each represented by the formula




wherein in Compound C169: X=O,
in Compound C170: X=S,
in Compound C171: X=Se

Compound C178 through C180, each represented by the formula


Compound C172 through C174, each represented by the formula

20




25

wherein in Compound C172: X=O,
in Compound C173: X=S,
in Compound C174: X=Se




wherein in Compound C175: X=O,
in Compound C176: X=S,
in Compound C177: X=Se



wherein in Compound C178: X=O,
in Compound C179: X=S,
in Compound C180: X=Se

Compound C181 through C183, each represented by the formula

30




35

wherein in Compound C181: X=O,
in Compound C182: X=S,
in Compound C183: X=Se


40

Compound C184 through C186, each represented by the formula



wherein in Compound C184: X=O,
in Compound C185: X=S,
in Compound C186: X=Se


Compound C187 through C189, each represented by the formula



wherein in Compound C187: X=O,
in Compound C188: X=S,
in Compound C189: X=Se

Compound C190 through C192, each represented by the formula


45



50


wherein in Compound C190: X=O,
in Compound C191: X=S,
in Compound C192: X=Se

Compound C193 through C195, each represented by the formula



wherein in Compound C193: X=O,
in Compound C194: X=S,
in Compound C195: X=Se

Compound C196 through C198, each represented by the formula




wherein in Compound C196: X=O,
in Compound C197: X=S,
in Compound C198: X=Se

55


EP 2 966 706 B1

Compound C199 through C201, each represented by the formula




wherein in Compound C199: X=O,
in Compound C200: X=S,
in Compound C201: X=Se

Compound C202 through C204, each represented by the formula




wherein in Compound C202: X=O,
in Compound C203: X=S,
in Compound C204: X=Se

Compound C205 through C207, each represented by the formula




wherein in Compound C205: X=O,
in Compound C206: X=S,
in Compound C207: X=Se

Compound C208 through C210, each represented by the formula




wherein in Compound C208: X=O,
in Compound C209: X=S,
in Compound C210: X=Se

Compound C211 through C213, each represented by the formula




wherein in Compound C211: X=O,
in Compound C212: X=S,
in Compound C213: X=Se

Compound C214 through C216, each represented by the formula




wherein in Compound C214: X=O,
in Compound C215: X=S,
in Compound C216: X=Se

Compound C217 through C219, each represented by the formula




wherein in Compound C217: X=O,
in Compound C218: X=S,
in Compound C219: X=Se

Compound C220 through C222, each represented by the formula




wherein in Compound C220: X=O,
in Compound C221: X=S,
in Compound C222: X=Se

Compound C223 through C225, each represented by the formula




wherein in Compound C223: X=O,
in Compound C224: X=S,
in Compound C225: X=Se

Compound C226 through C228, each represented by the formula




wherein in Compound C226: X=O,
in Compound C227: X=S,
in Compound C228: X=Se

Compound C229 through C231, each represented by the formula




wherein in Compound C229: X=O,
in Compound C230: X=S,
in Compound C231: X=Se

Compound C232 through C234, each represented by the formula




wherein in Compound C232: X=O,
in Compound C233: X=S,
in Compound C234: X=Se

Compound C235 through C237, each represented by the formula




wherein in Compound C235: X=O,
in Compound C236: X=S,
in Compound C237: X=Se

Compound C238 through C240, each represented by the formula




wherein in Compound C238: X=O,
in Compound C239: X=S,
in Compound C240: X=Se

Compound C241 through C243, each represented by the formula




wherein in Compound C241: X=O,
in Compound C242: X=S,
in Compound C243: X=Se

Compound C244 through C246, each represented by the formula




wherein in Compound C244: X=O,
in Compound C245: X=S,
in Compound C246: X=Se

Compound C247 through C249, each represented by the formula




wherein in Compound C247: X=O,
in Compound C248: X=S,
in Compound C249: X=Se

Compound C250 through C252, each represented by the formula



wherein in Compound C250: X=O,
in Compound C251: X=S,
in Compound C252: X=Se



Compound C253 through C255, each represented by the formula



wherein in Compound C253: X=O,
in Compound C254: X=S,
in Compound C255: X=Se


and

10. The composition of Claim 1, wherein the first compound is selected from the group consisting of:



and

5




10

Compound D3

11. The composition of Claim 1, wherein the second compound is selected from the group consisting of

15

20



25

where:

X is selected from the group consisting of O, S and Se;

R¹ and R⁴ each independently represents mono, di, or tri, substitution, or no substitution;

30

R², R³, R⁵, and R⁶ each independently represents mono, di, tri, or tetra substitution, or no substitution; and

R¹ to R⁶ are each independently selected from the group consisting of hydrogen, deuterium, benzene, biphenyl, terphenyl, naphthalene, fluorene, triphenylene, phenanthrene, dibenzofuran, dibenzothiophene, carbazole and combinations thereof.

35

12. The composition of Claim 1, wherein the second compound is selected from the group consisting of

40

45

50




55

Compound E1 through E3, each represented by the formula

Compound E4 through E6, each represented by the formula

Compound E7 through E9, each represented by the formula

5



10

wherein in Compound E1: X=O,
in Compound E2: X=S,
in Compound E3: X=Se

wherein in Compound E4: X=O,
in Compound E5: X=S,
in Compound E6: X=Se




wherein in Compound E7: X=O,
in Compound E8: X=S,
in Compound E9: X=Se

Compound E10 through E12, each represented by the formula

Compound E13 through E15, each represented by the formula

Compound E16 through E18, each represented by the formula

20



25

wherein in Compound E10: X=O,
in Compound E11: X=S,
in Compound E12: X=Se

wherein in Compound E13: X=O,
in Compound E14: X=S,
in Compound E15: X=Se




wherein in Compound E16: X=O,
in Compound E17: X=S,
in Compound E18: X=Se

Compound E19 through E21, each represented by the formula

Compound E22 through E24, each represented by the formula

Compound E25 through E27, each represented by the formula

30




35

wherein in Compound E19: X=O,
in Compound E20: X=S,
in Compound E21: X=Se


wherein in Compound E22: X=O,
in Compound E23: X=S,
in Compound E24: X=Se

wherein in Compound E25: X=O,
in Compound E26: X=S,
in Compound E27: X=Se


40



Compound E28



Compound E29




Compound E30


45

13. The composition of Claim 1, wherein the mixture of the first compound and the second compound is selected from the group consisting of:


50




Compound E1



Compound C1



Compound E2




Compound C2


55

(Compound E1 , Compound C1); (Compound E2 , Compound C2);


5
10
15
20
25
30
35
40
45
50
55




(**Compound E5** , **Compound C65**); (**Compound E8** ,




Compound C74); (**Compound E11** , **Compound C74**);



(**Compound E17** , **Compound C74**); (**Compound E8** ,




Compound A5); (**Compound E8** , **Compound C17**);




(**Compound E17** , **Compound A5**); (**Compound E26** , **Compound C74**);

5




10



15

20



25

30

(Compound E8, Compound C65); (Compound E8, Compound A116); and (Compound E8, Compound D2).

14. A first device comprising a first organic light emitting device, the first organic light emitting device comprising:

35


- an anode;
- a cathode; and
- an organic layer, disposed between the anode and the cathode, comprising a composition of materials as defined in any one of the previous claims.

40

Patentansprüche

1. Eine Zusammensetzung von Materialien umfassend eine erste Verbindung und eine zweite Verbindung, wobei die erste Verbindung die folgende Formel aufweist:

45



50

55

, Formel I;

wobei G¹ ausgewählt ist aus der Gruppe bestehend aus Dibenzofuran, Dibenzothiophen, Dibenzoselenophen, und Fluoren;

wobei L¹, L² und L³ jeweils unabhängig ausgewählt sind aus der Gruppe bestehend aus direkter Bindung, Phenyl, Biphenyl, Terphenyl, Pyridin, Pyrimidin, und Kombinationen davon;

wobei G⁴ ausgewählt ist aus der Gruppe bestehend aus Phenyl, Biphenyl, Terphenyl, Naphthalin, Phenanthren, Pyridin, Pyrimidin, Pyrazin, Chinolin, Isochinolin, Phenanthrolin, und Kombinationen davon;

wobei G², G³, und G⁵ jeweils unabhängig ausgewählt sind aus der Gruppe bestehend aus Phenyl, Biphenyl, Terphenyl, Fluoren, Naphthalin, Phenanthren, Pyridin, Pyrimidin, Pyrazin, Chinolin, Isochinolin, Phenanthrolin, Aza-fluoren, und Kombinationen davon;

wobei G², G³, G⁴, und G⁵ jeweils gegebenenfalls weiter substituiert sind mit einem oder mehreren nicht fusionierten Substituenten ausgewählt aus der Gruppe bestehend aus Deuterium, Alkyl, Alkoxy, Cycloalkyl, Cycloalkoxyl, Halogen, Nitro, Nitril, Silyl, Phenyl, Biphenyl, Terphenyl, Pyridin, und Kombinationen davon;

wobei m eine ganze Zahl von 0 bis 7 ist,

wobei n eine ganze Zahl von 0 bis 4 ist;


wobei, wenn m oder n größer als 1 ist, dann können jedes G⁴ oder G⁵ gleich oder verschieden sein;

wobei, wenn n gleich 0 ist, und m gleich oder größer als 1 ist, dann ist jedes G⁴ ausgewählt aus der Gruppe bestehend aus Phenyl und Biphenyl;

wobei, wenn n gleich oder größer als 1 ist, dann ist L¹ nicht eine direkte Bindung; und

wobei, wenn m und n beide 0 sind, dann ist L¹ Biphenyl,

und wobei die zweite Verbindung die Formel II aufweist:



wobei Ar¹ ausgewählt ist aus der Gruppe bestehend aus Triphenylen und Azatriphenylen;

wobei Ar² ausgewählt ist aus der Gruppe bestehend aus einer direkten Bindung, Phenyl, Biphenyl, Terphenyl, Naphthalin, Pyridin, Dibenzofuran, Dibenzothiophen, Dibenzoselenophen, Aza-dibenzofuran, Aza-dibenzothiophen, Aza-dibenzoselenophen und Kombinationen davon;

wobei Ar³ ausgewählt ist aus der Gruppe bestehend aus Benzol, Biphenyl, Terphenyl, Naphthalin, Pyridin, Dibenzofuran, Dibenzothiophen, Dibenzoselenophen, Aza-dibenzofuran, Aza-dibenzothiophen, Aza-dibenzoselenophen, Carbazol, Aza-carbazol und Kombinationen davon; und


wobei Ar¹, Ar² und Ar³ jeweils unabhängig gegebenenfalls ferner substituiert sind mit einem oder mehreren Substitutionen ausgewählt aus der Gruppe bestehend aus Deuterium, Halogen, Alkyl, Aryl, Heteroaryl und Kombinationen davon.

2. Die Zusammensetzung nach Anspruch 1, wobei n gleich 0 ist.

3. Die Zusammensetzung nach Anspruch 1, wobei n gleich oder größer als 1 ist.

4. Die Zusammensetzung nach Anspruch 1, wobei m und n beide 0 sind.

5. Die Zusammensetzung nach Anspruch 1, wobei G¹ die Struktur ausgewählt aus der Gruppe bestehend aus:



aufweist,

wobei X ausgewählt ist aus einer Gruppe bestehend aus O, S und Se;


wobei R^{B1} und R^{B2} unabhängig ausgewählt sind aus einer Gruppe bestehend aus Wasserstoff, Deuterium, Alkyl, Cycloalkyl, Alkoxy, Aryl, Heteraryl, Halogen und Kombinationen davon; und

wobei R^{B1} und R^{B2} gegebenenfalls verknüpft sind um einen Ring zu bilden.

6. Die Zusammensetzung nach Anspruch 1, wobei mindestens einer von G², G³, G⁴ und G⁵ mit mindestens einem Fluoratom substituiert ist.

EP 2 966 706 B1

7. Die Zusammensetzung nach Anspruch 1, wobei die erste Verbindung die Formel aufweist:



wobei X ausgewählt ist aus einer Gruppe bestehend aus O, S und Se.

15 8. Die Zusammensetzung nach Anspruch 1, wobei die erste Verbindung ausgewählt ist aus der Gruppe bestehend aus:

20

25

30

35

40

45

50




55

Verbindung A1 bis A3, jeweils wiedergegeben durch die Formel

Verbindung A4 bis A6, jeweils wiedergegeben durch die Formel

Verbindung A7 bis A9, jeweils wiedergegeben durch die Formel

5



10

wobei in Verbindung A1: X=O,
in Verbindung A2: X=S,
in Verbindung A3: X=Se

wobei in Verbindung A4: X=O,
in Verbindung A5: X=S,
in Verbindung A6: X=Se

wobei in Verbindung A7: X=O,
in Verbindung A8: X=S,
in Verbindung A9: X=Se




15

Verbindung A10 bis A12, jeweils wiedergegeben durch die Formel

Verbindung A13 bis A15, jeweils wiedergegeben durch die Formel

Verbindung A16 bis A18, jeweils wiedergegeben durch die Formel

20



25

wobei in Verbindung A10: X=O,
in Verbindung A11: X=S,
in Verbindung A12: X=Se

wobei in Verbindung A13: X=O,
in Verbindung A14: X=S,
in Verbindung A15: X=Se




wobei in Verbindung A16: X=O,
in Verbindung A17: X=S,
in Verbindung A18: X=Se

Verbindung A19 bis A21, jeweils wiedergegeben durch die Formel

Verbindung A22 bis A24, jeweils wiedergegeben durch die Formel

Verbindung A25 bis A27, jeweils wiedergegeben durch die Formel

30



35

wobei in Verbindung A19: X=O,
in Verbindung A20: X=S,
in Verbindung A21: X=Se

wobei in Verbindung A22: X=O,
in Verbindung A23: X=S,
in Verbindung A24: X=Se




wobei in Verbindung A25: X=O,
in Verbindung A26: X=S,
in Verbindung A27: X=Se

Verbindung A28 bis A30, jeweils wiedergegeben durch die Formel

Verbindung A31 bis A33, jeweils wiedergegeben durch die Formel

Verbindung A34 bis A36, jeweils wiedergegeben durch die Formel

40



45

wobei in Verbindung A28: X=O,
in Verbindung A29: X=S,
in Verbindung A30: X=Se

wobei in Verbindung A31: X=O,
in Verbindung A32: X=S,
in Verbindung A33: X=Se


wobei in Verbindung A34: X=O,
in Verbindung A35: X=S,
in Verbindung A36: X=Se

50

55

Verbindung A37 bis A39, jeweils wiedergegeben durch die Formel


5



10


wobei in Verbindung A37: X=O,
in Verbindung A38: X=S,
in Verbindung A39: X=Se

Verbindung A40 bis A42, jeweils wiedergegeben durch die Formel



wobei in Verbindung A40: X=O,
in Verbindung A41: X=S,
in Verbindung A42: X=Se


Verbindung A43 bis A45, jeweils wiedergegeben durch die Formel



wobei in Verbindung A43: X=O,
in Verbindung A44: X=S,
in Verbindung A45: X=Se

15

Verbindung A46 bis A48, jeweils wiedergegeben durch die Formel




20

wobei in Verbindung A46: X=O,
in Verbindung A47: X=S,
in Verbindung A48: X=Se


25

Verbindung A49 bis A51, jeweils wiedergegeben durch die Formel



wobei in Verbindung A49: X=O,
in Verbindung A50: X=S,
in Verbindung A51: X=Se


Verbindung A52 bis A54, jeweils wiedergegeben durch die Formel



wobei in Verbindung A52: X=O,
in Verbindung A53: X=S,
in Verbindung A54: X=Se

Verbindung A55 bis A57, jeweils wiedergegeben durch die Formel


30



wobei in Verbindung A55: X=O,
in Verbindung A56: X=S,
in Verbindung A57: X=Se


35

Verbindung A58 bis A60, jeweils wiedergegeben durch die Formel



wobei in Verbindung A58: X=O,
in Verbindung A59: X=S,
in Verbindung A60: X=Se


Verbindung A61 bis A63, jeweils wiedergegeben durch die Formel



wobei in Verbindung A61: X=O,
in Verbindung A62: X=S,
in Verbindung A63: X=Se

Verbindung A64 bis A66, jeweils wiedergegeben durch die Formel


40



wobei in Verbindung A64: X=O,
in Verbindung A65: X=S,
in Verbindung A66: X=Se

45


Verbindung A67 bis A69, jeweils wiedergegeben durch die Formel



wobei in Verbindung A67: X=O,
in Verbindung A68: X=S,
in Verbindung A69: X=Se

50

Verbindung A70 bis A72, jeweils wiedergegeben durch die Formel




wobei in Verbindung A70: X=O,
in Verbindung A71: X=S,
in Verbindung A72: X=Se

55

EP 2 966 706 B1

Verbindung A73 bis A75, jeweils wiedergegeben durch die Formel


5



10


wobei in Verbindung A73: X=O,
in Verbindung A74: X=S,
in Verbindung A75: X=Se

Verbindung A76 bis A78, jeweils wiedergegeben durch die Formel



wobei in Verbindung A76: X=O,
in Verbindung A77: X=S,
in Verbindung A78: X=Se


Verbindung A79 bis A81, jeweils wiedergegeben durch die Formel



wobei in Verbindung A79: X=O,
in Verbindung A80: X=S,
in Verbindung A81: X=Se

Verbindung A82 bis A84, jeweils wiedergegeben durch die Formel


15



20


wobei in Verbindung A82: X=O,
in Verbindung A83: X=S,
in Verbindung A84: X=Se

Verbindung A85 bis A87, jeweils wiedergegeben durch die Formel



wobei in Verbindung A85: X=O,
in Verbindung A86: X=S,
in Verbindung A87: X=Se

Verbindung A88 bis A90, jeweils wiedergegeben durch die Formel




25

wobei in Verbindung A88: X=O,
in Verbindung A89: X=S,
in Verbindung A90: X=Se

Verbindung A91 bis A93, jeweils wiedergegeben durch die Formel


30



35


wobei in Verbindung A91: X=O,
in Verbindung A92: X=S,
in Verbindung A93: X=Se

Verbindung A94 bis A96, jeweils wiedergegeben durch die Formel



wobei in Verbindung A94: X=O,
in Verbindung A95: X=S,
in Verbindung A96: X=Se


Verbindung A97 bis A99, jeweils wiedergegeben durch die Formel



wobei in Verbindung A97: X=O,
in Verbindung A98: X=S,
in Verbindung A99: X=Se

Verbindung A100 bis A102, jeweils wiedergegeben durch die Formel


40



45


wobei in Verbindung A100: X=O,
in Verbindung A101: X=S,
in Verbindung A102: X=Se

Verbindung A103 bis A105, jeweils wiedergegeben durch die Formel



wobei in Verbindung A103: X=O,
in Verbindung A104: X=S,
in Verbindung A105: X=Se

Verbindung A106 bis A108, jeweils wiedergegeben durch die Formel




wobei in Verbindung A106: X=O,
in Verbindung A107: X=S,
in Verbindung A108: X=Se

50


55

Verbindung A109 bis A111, jeweils wiedergegeben durch die Formel




wobei in Verbindung A109: X=O,
in Verbindung A110: X=S,
in Verbindung A111: X=Se

Verbindung A112 bis A114, jeweils wiedergegeben durch die Formel



wobei in Verbindung A112: X=O,
in Verbindung A113: X=S,
in Verbindung A114: X=Se

Verbindung A115 bis A117, wiedergegeben durch die Formel



wobei in Verbindung A115: X=O,
in Verbindung A116: X=S,
in Verbindung A117: X=Se


5

10


15

20


25



Verbindung B1




Verbindung B2




Verbindung B3


30



Verbindung B4




Verbindung B5




Verbindung B6

40

45



Verbindung B7



Verbindung B8

, und

55




9. Die Zusammensetzung nach Anspruch 1, wobei die erste Verbindung ausgewählt ist aus der Gruppe bestehend aus:

Verbindung C1 bis C3, jeweils wiedergegeben durch die Formel

Verbindung C4 bis C6, jeweils wiedergegeben durch die Formel

Verbindung C7 bis C9, jeweils wiedergegeben durch die Formel

5



10

wobei in Verbindung C1: X=O,
in Verbindung C2: X=S,
in Verbindung C3: X=Se

wobei in Verbindung C4: X=O,
in Verbindung C5: X=S,
in Verbindung C6: X=Se

wobei in Verbindung C7: X=O,
in Verbindung C8: X=S,
in Verbindung C9: X=Se




15

Verbindung C10 bis C12, jeweils wiedergegeben durch die Formel

Verbindung C13 bis C15, jeweils wiedergegeben durch die Formel

Verbindung C16 bis C18, jeweils wiedergegeben durch die Formel

20



25

wobei in Verbindung C10: X=O,
in Verbindung C11: X=S,
in Verbindung C12: X=Se

wobei in Verbindung C13: X=O,
in Verbindung C14: X=S,
in Verbindung C15: X=Se




wobei in Verbindung C16: X=O,
in Verbindung C17: X=S,
in Verbindung C18: X=Se

Verbindung C19 bis C21, jeweils wiedergegeben durch die Formel

Verbindung C22 bis C24, jeweils wiedergegeben durch die Formel

Verbindung C25 bis C27, jeweils wiedergegeben durch die Formel

30



35

wobei in Verbindung C19: X=O,
in Verbindung C20: X=S,
in Verbindung C21: X=Se

wobei in Verbindung C22: X=O,
in Verbindung C23: X=S,
in Verbindung C24: X=Se




wobei in Verbindung C25: X=O,
in Verbindung C26: X=S,
in Verbindung C27: X=Se

Verbindung C28 bis C30, jeweils wiedergegeben durch die Formel

Verbindung C31 bis C33, jeweils wiedergegeben durch die Formel

Verbindung C34 bis C36, jeweils wiedergegeben durch die Formel

40



45

wobei in Verbindung C28: X=O,
in Verbindung C29: X=S,
in Verbindung C30: X=Se

wobei in Verbindung C31: X=O,
in Verbindung C32: X=S,
in Verbindung C33: X=Se

wobei in Verbindung C34: X=O,
in Verbindung C35: X=S,
in Verbindung C36: X=Se

50




55

Verbindung C37 bis C39, jeweils wiedergegeben durch die Formel

Verbindung C40 bis C42, jeweils wiedergegeben durch die Formel

Verbindung C43 bis C45, jeweils wiedergegeben durch die Formel

5



10

wobei in Verbindung C37: X=O,
in Verbindung C38: X=S,
in Verbindung C39: X=Se

wobei in Verbindung C40: X=O,
in Verbindung C41: X=S,
in Verbindung C42: X=Se

wobei in Verbindung C43: X=O,
in Verbindung C44: X=S,
in Verbindung C45: X=Se




15

Verbindung C46 bis C48, jeweils wiedergegeben durch die Formel

Verbindung C49 bis C51, jeweils wiedergegeben durch die Formel

Verbindung C52 bis C54, jeweils wiedergegeben durch die Formel

20



wobei in Verbindung C46: X=O,
in Verbindung C47: X=S,
in Verbindung C48: X=Se

wobei in Verbindung C49: X=O,
in Verbindung C50: X=S,
in Verbindung C51: X=Se

wobei in Verbindung C52: X=O,
in Verbindung C53: X=S,
in Verbindung C54: X=Se




25

Verbindung C55 bis C57, jeweils wiedergegeben durch die Formel

Verbindung C58 bis C60, jeweils wiedergegeben durch die Formel

Verbindung C61 bis C63, jeweils wiedergegeben durch die Formel

30



35

in Verbindung C56: X=S,
in Verbindung C57: X=Se

wobei in Verbindung C58: X=O,
in Verbindung C59: X=S,
in Verbindung C60: X=Se

wobei in Verbindung C61: X=O,
in Verbindung C62: X=S,
in Verbindung C63: X=Se




40

Verbindung C64 bis C66, jeweils wiedergegeben durch die Formel

Verbindung C67 bis C69, jeweils wiedergegeben durch die Formel

Verbindung C70 bis C72, jeweils wiedergegeben durch die Formel

45



wobei in Verbindung C64: X=O,
in Verbindung C65: X=S,
in Verbindung C66: X=Se

wobei in Verbindung C67: X=O,
in Verbindung C68: X=S,
in Verbindung C69: X=Se


wobei in Verbindung C70: X=O,
in Verbindung C71: X=S,
in Verbindung C72: X=Se

50

55

Verbindung C73 bis C75, jeweils wiedergegeben durch die Formel


5



10


wobei in Verbindung C73: X=O,
in Verbindung C74: X=S,
in Verbindung C75: X=Se

Verbindung C76 bis C78, jeweils wiedergegeben durch die Formel



wobei in Verbindung C76: X=O,
in Verbindung C77: X=S,
in Verbindung C78: X=Se


Verbindung C79 bis C81, jeweils wiedergegeben durch die Formel



wobei in Verbindung C79: X=O,
in Verbindung C80: X=S,
in Verbindung C81: X=Se

Verbindung C82 bis C84, jeweils wiedergegeben durch die Formel

15




20

wobei in Verbindung C82: X=O,
in Verbindung C83: X=S,
in Verbindung C84: X=Se


25

Verbindung C85 bis C87, jeweils wiedergegeben durch die Formel



wobei in Verbindung C85: X=O,
in Verbindung C86: X=S,
in Verbindung C87: X=Se


Verbindung C88 bis C90, jeweils wiedergegeben durch die Formel



wobei in Verbindung C88: X=O,
in Verbindung C89: X=S,
in Verbindung C90: X=Se

Verbindung C91 bis C93, jeweils wiedergegeben durch die Formel


30



wobei in Verbindung C91: X=O,
in Verbindung C92: X=S,
in Verbindung C93: X=Se


35

Verbindung C94 bis C96, jeweils wiedergegeben durch die Formel



wobei in Verbindung C94: X=O,
in Verbindung C95: X=S,
in Verbindung C96: X=Se


Verbindung C97 bis C99, jeweils wiedergegeben durch die Formel



wobei in Verbindung C97: X=O,
in Verbindung C98: X=S,
in Verbindung C99: X=Se

Verbindung C100 bis C102, jeweils wiedergegeben durch die Formel


40



wobei in Verbindung C100: X=O,
in Verbindung C101: X=S,
in Verbindung C102: X=Se


45

Verbindung C103 bis C105, jeweils wiedergegeben durch die Formel



wobei in Verbindung C103: X=O,
in Verbindung C104: X=S,
in Verbindung C105: X=Se

Verbindung C106 bis C108, jeweils wiedergegeben durch die Formel



wobei in Verbindung C106: X=O,
in Verbindung C107: X=S,
in Verbindung C108: X=Se

50




55

Verbindung C109 bis C111, jeweils wiedergegeben durch die Formel

Verbindung C112 through C114, jeweils wiedergegeben durch die Formel

Verbindung C115 bis C117, jeweils wiedergegeben durch die Formel

5



10

wobei in Verbindung C109: X=O,
in Verbindung C110: X=S,
in Verbindung C111: X=Se

wobei in Verbindung C112: X=O,
in Verbindung C113: X=S,
in Verbindung C114: X=Se

wobei in Verbindung C115: X=O,
in Verbindung C116: X=S,
in Verbindung C117: X=Se




15

Verbindung C118 bis C120, jeweils wiedergegeben durch die Formel

Verbindung C121 bis C123, jeweils wiedergegeben durch die Formel

Verbindung C124 bis C126, jeweils wiedergegeben durch die Formel

20



25

wobei in Verbindung C118: X=O,
in Verbindung C119: X=S,
in Verbindung C120: X=Se

wobei in Verbindung C121: X=O,
in Verbindung C122: X=S,
in Verbindung C123: X=Se

wobei in Verbindung C124: X=O,
in Verbindung C125: X=S,
in Verbindung C126: X=Se




30

Verbindung C127 bis C129, jeweils wiedergegeben durch die Formel

Verbindung C130 bis C132, jeweils wiedergegeben durch die Formel

Verbindung C133 bis C135, jeweils wiedergegeben durch die Formel

35



40

wobei in Verbindung C127: X=O,
in Verbindung C128: X=S,
in Verbindung C129: X=Se

wobei in Verbindung C130: X=O,
in Verbindung C131: X=S,
in Verbindung C132: X=Se

wobei in Verbindung C133: X=O,
in Verbindung C134: X=S,
in Verbindung C135: X=Se




45

Verbindung C136 bis C138, jeweils wiedergegeben durch die Formel

Verbindung C139 bis C141, jeweils wiedergegeben durch die Formel

Verbindung C142 bis C144, jeweils wiedergegeben durch die Formel

50



wobei in Verbindung C136: X=O,
in Verbindung C137: X=S,
in Verbindung C138: X=Se

wobei in Verbindung C139: X=O,
in Verbindung C140: X=S,
in Verbindung C141: X=Se

wobei in Verbindung C142: X=O,
in Verbindung C143: X=S,
in Verbindung C144: X=Se

55




EP 2 966 706 B1

Verbindung C145 bis C147, jeweils wieder-
gegeben durch die Formel

Verbindung C148 bis C150, jeweils wieder-
gegeben durch die Formel

Verbindung C151 bis C153, jeweils wieder-
gegeben durch die Formel

5



10

wobei in Verbindung C145: X=O,
in Verbindung C146: X=S,
in Verbindung C147: X=Se

wobei in Verbindung C148: X=O,
in Verbindung C149: X=S,
in Verbindung C150: X=Se

wobei in Verbindung C151: X=O,
in Verbindung C152: X=S,
in Verbindung C153: X=Se




15

Verbindung C154 bis C156, jeweils wieder-
gegeben durch die Formel

Verbindung C157 bis C159, jeweils wieder-
gegeben durch die Formel

Verbindung C160 bis C162, jeweils wieder-
gegeben durch die Formel

20



25

wobei in Verbindung C154: X=O,
in Verbindung C155: X=S,
in Verbindung C156: X=Se

wobei in Verbindung C157: X=O,
in Verbindung C158: X=S,
in Verbindung C159: X=Se

wobei in Verbindung C160: X=O,
in Verbindung C161: X=S,
in Verbindung C162: X=Se




30

Verbindung C163 bis C165, jeweils wieder-
gegeben durch die Formel

Verbindung C166 bis C168, jeweils wieder-
gegeben durch die Formel

Verbindung C169 bis C171, jeweils wieder-
gegeben durch die Formel

35



40

wobei in Verbindung C163: X=O,
in Verbindung C164: X=S,
in Verbindung C165: X=Se

wobei in Verbindung C166: X=O,
in Verbindung C167: X=S,
in Verbindung C168: X=Se




wobei in Verbindung C169: X=O,
in Verbindung C170: X=S,
in Verbindung C171: X=Se

Verbindung C175 bis C177, jeweils wieder-
gegeben durch die Formel

Verbindung C178 bis C180, jeweils wieder-
gegeben durch die Formel

45

Verbindung C172 bis C174, jeweils wieder-
gegeben durch die Formel



50

wobei in Verbindung C172: X=O,
in Verbindung C173: X=S,
in Verbindung C174: X=Se

wobei in Verbindung C175: X=O,
in Verbindung C176: X=S,
in Verbindung C177: X=Se

wobei in Verbindung C178: X=O,
in Verbindung C179: X=S,
in Verbindung C180: X=Se




55

Verbindung C181 bis C183, jeweils wieder-
gegeben durch die Formel

Verbindung C184 bis C186, jeweils wieder-
gegeben durch die Formel

Verbindung C187 bis C189, jeweils wieder-
gegeben durch die Formel

5



10

wobei in Verbindung C181: X=O,
in Verbindung C182: X=S,
in Verbindung C183: X=Se

wobei in Verbindung C184: X=O,
in Verbindung C185: X=S,
in Verbindung C186: X=Se

wobei in Verbindung C187: X=O,
in Verbindung C188: X=S,
in Verbindung C189: X=Se




15

Verbindung C190 bis C192, jeweils wieder-
gegeben durch die Formel

Verbindung C193 bis C195, jeweils wieder-
gegeben durch die Formel

Verbindung C196 bis C198, jeweils wieder-
gegeben durch die Formel

20



25

wobei in Verbindung C190: X=O,
in Verbindung C191: X=S,
in Verbindung C192: X=Se

wobei in Verbindung C193: X=O,
in Verbindung C194: X=S,
in Verbindung C195: X=Se

wobei in Verbindung C196: X=O,
in Verbindung C197: X=S,
in Verbindung C198: X=Se




30

Verbindung C199 bis C201, jeweils wieder-
gegeben durch die Formel

Verbindung C202 bis C204, jeweils wieder-
gegeben durch die Formel

Verbindung C205 bis C207, jeweils wieder-
gegeben durch die Formel

35



40

wobei in Verbindung C199: X=O,
in Verbindung C200: X=S,
in Verbindung C201: X=Se

wobei in Verbindung C202: X=O,
in Verbindung C203: X=S,
in Verbindung C204: X=Se




wobei in Verbindung C205: X=O,
in Verbindung C206: X=S,
in Verbindung C207: X=Se

Verbindung C208 bis C210, jeweils wieder-
gegeben durch die Formel

Verbindung C211 bis C213, jeweils wieder-
gegeben durch die Formel

Verbindung C214 bis C216, jeweils wieder-
gegeben durch die Formel

45



50

wobei in Formel C208: X=O,
in Verbindung C209: X=S,
in Verbindung C210: X=Se

wobei in Verbindung C211: X=O,
in Verbindung C212: X=S,
in Verbindung C213: X=Se

wobei in Verbindung C214: X=O,
in Verbindung C215: X=S,
in Verbindung C216: X=Se




55

Verbindung C217 bis C219, jeweils wieder-
gegeben durch die Formel

Verbindung C220 bis C222, jeweils wieder-
gegeben durch die Formel

Verbindung C223 bis C225, jeweils wieder-
gegeben durch die Formel

5



10

wobei in Verbindung C217: X=O,
in Verbindung C218: X=S,
in Verbindung C219: X=Se

wobei in Verbindung C220: X=O,
in Verbindung C221: X=S,
in Verbindung C222: X=Se

wobei in Verbindung C223: X=O,
in Verbindung C224: X=S,
in Verbindung C225: X=Se




15

Verbindung C226 bis C228, jeweils wieder-
gegeben durch die Formel

Verbindung C229 bis C231, jeweils wieder-
gegeben durch die Formel

Verbindung C232 bis C234, jeweils wieder-
gegeben durch die Formel

20



25

wobei in Verbindung C226: X=O,
in Verbindung C227: X=S,
in Verbindung C228: X=Se

wobei in Verbindung C229: X=O,
in Verbindung C230: X=S,
in Verbindung C231: X=Se

wobei in Verbindung C232: X=O,
in Verbindung C233: X=S,
in Verbindung C234: X=Se




30

Verbindung C235 bis C237, jeweils wieder-
gegeben durch die Formel

Verbindung C238 bis C240, jeweils wieder-
gegeben durch die Formel

Verbindung C241 bis C243, jeweils wiedergegeben
durch die Formel

35



wobei in Verbindung C235: X=O,
in Verbindung C236: X=S,
in Verbindung C237: X=Se

wobei in Verbindung C238: X=O,
in Verbindung C239: X=S,
in Verbindung C240: X=Se

wobei in Verbindung C241: X=O,
in Verbindung C242: X=S,
in Verbindung C243: X=Se

40


45

50

55


EP 2 966 706 B1

Verbindung C244 bis C246, jeweils wieder-
gegeben durch die Formel




wobei in Verbindung C244: X=O,
in Verbindung C245: X=S,
in Verbindung C246: X=Se

Verbindung C247 bis C249, jeweils wieder-
gegeben durch die Formel




wobei in Verbindung C247: X=O,
in Verbindung C248: X=S,
in Verbindung C249: X=Se

Verbindung C250 bis C252, jeweils wieder-
gegeben durch die Formel





wobei in Verbindung C250: X=O,
in Verbindung C251: X=S,
in Verbindung C252: X=Se

Verbindung C253 bis C255, jeweils wieder-




wobei in Verbindung C253: X=O,
in Verbindung C254: X=S,
in Verbindung C255: X=Se

10. Die Zusammensetzung nach Anspruch 1, wobei die erste Verbindung ausgewählt ist aus der Gruppe bestehend aus:




und



11. Die Zusammensetzung nach Anspruch 1, wobei die zweite Verbindung ausgewählt ist aus der Gruppe bestehend aus

5

10



, und

,

wobei:

15

20

X ist ausgewählt aus der Gruppe bestehend aus O, S und Se;

R¹ und R⁴ unabhängig für mono-, di-, oder tri-Substitution, oder keine Substitution;

R², R³, R⁵, und R⁶ stehen unabhängig für mono-, di-, tri-, oder tetra-Substitution, oder keine Substitution; und

R¹ bis R⁶ sind jeweils unabhängig ausgewählt aus der Gruppe bestehend aus Wasserstoff, Deuterium, Benzol, Biphenyl, Terphenyl, Naphthalin, Fluorene, Triphenylen, Phenanthren, Dibenzofuran, Dibenzothiophen, Carbazol und Kombinationen davon.

12. Die Zusammensetzung nach Anspruch 1, wobei die zweite Verbindung ausgewählt ist aus der Gruppe bestehend aus:

25

30

35

40

45

50



55

EP 2 966 706 B1

Verbindung E1 bis E3, jeweils wiedergegeben durch die Formel

Verbindung E4 bis E6, jeweils wiedergegeben durch die Formel

5



10

wobei in Verbindung E1: X=O,
in Verbindung E2: X=S,
in Verbindung E3: X=Se



wobei in Verbindung E4: X=O,
in Verbindung E5: X=S,
in Verbindung E6: X=Se

15

Verbindung E7 bis E9, jeweils wiedergegeben durch die Formel

Verbindung E10 bis E12, jeweils wiedergegeben durch die Formel

20



wobei in Verbindung E7: X=O,
in Verbindung E8: X=S,
in Verbindung E9: X=Se

wobei in Verbindung E10: X=O,
in Verbindung E11: X=S,
in Verbindung E12: X=Se




25

Verbindung E13 bis E15, jeweils wiedergegeben durch die Formel

Verbindung E16 bis E18, jeweils wiedergegeben

Verbindung E19 bis E21, jeweils wiedergegeben durch die Formel

30



wobei in Verbindung E13: X=O,
in Verbindung E14: X=S,
in Verbindung E15: X=Se

wobei in Verbindung E16: X=O,
in Verbindung E17: X=S,
in Verbindung E18: X=Se

wobei in Verbindung E19: X=O,
in Verbindung E20: X=S,
in Verbindung E21: X=Se



35

40

Verbindung E22 bis E24, jeweils wiedergegeben durch die Formel

Verbindung E25 bis E27, jeweils wiedergegeben durch die Formel

45




wobei in Verbindung E22: X=O,
in Verbindung E23: X=S,
in Verbindung E24: X=Se

wobei in Verbindung E25: X=O,
in Verbindung E26: X=S,
in Verbindung E27: X=Se

50


55

5




Verbindung E28

10



Verbindung E29

, und




Verbindung E30


15

13. Die Zusammensetzung nach Anspruch 1, wobei die Mischung aus der ersten Verbindung und der zweiten Verbindung ausgewählt ist aus der Gruppe bestehend aus:


20




Verbindung E1



Verbindung C1



Verbindung E2




Verbindung C2


25

(Verbindung E1 , Verbindung C1); (Verbindung E2 , Verbindung C2);


30



Verbindung E5



Verbindung C65




Verbindung E8


35

(Verbindung E5 , Verbindung C65); (Verbindung E8 ,


40



Verbindung C74



Verbindung E11



Verbindung C74


45

); (Verbindung E11 , Verbindung C74);

50

55


5



(**Verbindung E17** , **Verbindung C74**); (**Verbindung E8** ,

10


15



Verbindung A5); (**Verbindung E8** , **Verbindung C17**);

20


25



(**Verbindung E17** , **Verbindung A5**); (**Verbindung E26** , **Verbindung C74**);

30


35



(**Verbindung E26** , **Verbindung C248**); (**Verbindung E28** ,

40


45



Verbindung C74); (**Verbindung E29** , **Verbindung C74**);

50

55



(Verbindung E8, Verbindung C65); (Verbindung E8, Verbindung A116); und (Verbindung E8, Verbindung D2); .


14. Eine erste Vorrichtung umfassend eine erste organische Licht emittierende Vorrichtung, die erste organische Licht emittierende Vorrichtung umfassend:

15
 eine Anode;
 eine Kathode; und
 eine organische Schicht, angeordnet zwischen der Anode und der Kathode, umfassend eine Zusammensetzung von Materialien wie in einem der vorhergehenden Ansprüche definiert.

20

Revendications

1. Une composition de matières comprenant un premier composé et un second composé, où le premier composé est représenté par la formule :



où G¹ est sélectionné parmi le groupe constitué de dibenzofurane, dibenzothiophène, dibenzosélénophène, et fluorène ;

où L¹, L² et L³ sont chacun indépendamment sélectionnés parmi le groupe constitué d'une liaison directe, de phényle, biphényle, terphényle, pyridine, pyrimidine, et de combinaisons de ceux-ci ;

où G⁴ est sélectionné parmi le groupe constitué de phényle, biphényle, terphényle, naphthalène, phénanthrène, pyridine, pyrimidine, pyrazine, quinoléine, isoquinoléine, phénanthroline, et de combinaisons de ceux-ci ;

où G², G³, et G⁵ sont chacun indépendamment sélectionnés parmi le groupe constitué de phényle, biphényle, terphényle, fluorène, naphthalène, phénanthrène, pyridine, pyrimidine, pyrazine, quinoléine, isoquinoléine, phénanthroline, azáfluorène, et de combinaisons de ceux-ci ;

où G², G³, G⁴, et G⁵ sont chacun éventuellement en outre substitués avec un ou plusieurs substituants non-fusionnés sélectionnés parmi le groupe constitué de deutérium, alkyle, alkoxy, cycloalkyle, cycloalkoxy, halogène, nitro, nitrile, silyle, phényle, biphényle, terphényle, pyridine, et de combinaisons de ceux-ci ;

où m est un nombre entier de 0 à 7,

où n est un nombre entier de 0 à 4 ;


où, lorsque m ou n est supérieur à 1, alors chaque G⁴ ou G⁵ peut être identique ou différent ;

où, lorsque n est égal à 0, et m est égal ou supérieur à 1, alors chaque G⁴ est sélectionné parmi le groupe constitué de phényle et biphényle ;

où, lorsque n est égal ou supérieur à 1, alors L¹ n'est pas une liaison directe ; et

où, lorsque m et n sont tous les deux égaux à 0, alors L¹ est biphényle,

et où le second composé est représenté par la formule II :



5

où Ar¹ est sélectionné parmi le groupe constitué de triphénylène et azatriphénylène ;
 où Ar² est sélectionné parmi le groupe constitué d'une liaison directe, de phényle, biphényle, terphényle, naphthalène, pyridine, dibenzofurane, dibenzothiophène, dibenzosélénophène, azadibenzofurane, azadibenzothiophène, azadibenzosélénophène, et de combinaisons de ceux-ci ;

10 où Ar³ est sélectionné parmi le groupe constitué de benzène, biphényle, terphényle, naphthalène, pyridine, dibenzofurane, dibenzothiophène, dibenzosélénophène, azadibenzofurane, azadibenzothiophène, azadibenzosélénophène, carbazole, azacarbazole, et de combinaisons de ceux-ci ; et

15 où Ar¹, Ar² et Ar³ sont chacun, indépendamment, éventuellement en outre substitués avec une ou plusieurs substitutions sélectionnées parmi le groupe constitué de deutérium, halogène, alkyle, aryle, hétéroaryle, et de combinaisons de ceux-ci.


2. La composition selon la revendication 1, où n est égal à 0.

20 3. La composition selon la revendication 1, où n est égal ou supérieur à 1.

4. La composition selon la revendication 1, où m et n sont tous les deux égaux à 0.

25 5. La composition selon la revendication 1, où G¹ a la structure sélectionnée parmi le groupe constitué de :

25



30


où X est sélectionné parmi un groupe constitué de O, S et Se ;
 où R^{B1} et R^{B2} sont indépendamment sélectionnés parmi un groupe constitué d'hydrogène, deutérium, alkyle, cycloalkyle, alkoxy, aryle, hétéroaryle, halogène, et de combinaisons de ceux-ci ; et
 35 où R^{B1} et R^{B2} sont éventuellement reliés pour former un cycle.

35

6. La composition selon la revendication 1, où au moins un parmi G², G³, G⁴ et G⁵ est substitué avec au moins un atome de fluor.

40 7. Le composé selon la revendication 1, où le premier composé est représenté par la formule :

40



50


où X est sélectionné parmi un groupe constitué de O, S et Se.

8. La composition selon la revendication 1, où le premier composé est sélectionné parmi le groupe constitué de :

55


5

Composé A1 à A3, chacun représenté par la formule




10

Composé A4 à A6, chacun représenté par la formule



Composé A7 à A9, chacun représenté par la formule



15


où dans le Composé A1 : X=O,
dans le Composé A2 : X=S,
dans le Composé A3 : X=Se

où dans le Composé A4 :
X=O,
dans le Composé A5 :
X=S,
dans le Composé A6 :
X=Se

où dans le Composé A7 : X=O,
dans le Composé A8 : X=S,
dans le Composé A9 : X=Se


20

Composé A10 à A12, chacun représenté par la formule




25

Composé A13 à A15, chacun représenté par la formule



30

Composé A16 à A18, chacun représenté par la formule




où dans le Composé A10 : X=O,
dans le Composé A11 : X=S,
dans le Composé A12 : X=Se

où dans le Composé A13 : X=O,
dans le Composé A14 : X=S,
dans le Composé A15 : X=Se

où dans le Composé A16 : X=O,
dans le Composé A17 : X=S,
dans le Composé A18 : X=Se


35

Composé A19 à A21, chacun représenté par la formule




40

Composé A22 à A24, chacun représenté par la formule



Composé A25 à A27, chacun représenté par la formule



45

où dans le Composé A19 : X=O,
dans le Composé A20 : X=S,
dans le Composé A21 : X=Se

où dans le Composé A22 : X=O,
dans le Composé A23 : X=S,
dans le Composé A24 : X=Se


où dans le Composé A25 : X=O,
dans le Composé A26 : X=S,
dans le Composé A27 : X=Se

50

55

Composé A28 à A30, chacun représenté par la formule

5




10

où dans le Composé A28 : X=O,
dans le Composé A29 : X=S,
dans le Composé A30 : X=Se


15

Composé A31 à A33, chacun représenté par la formule



où dans le Composé A31 : X=O,
dans le Composé A32 : X=S,
dans le Composé A33 : X=Se


Composé A34 à A36, chacun représenté par la formule



où dans le Composé A34 : X=O,
dans le Composé A35 : X=S,
dans le Composé A36 : X=Se

Composé A37 à A39, chacun représenté par la formule

20




25

où dans le Composé A37 : X=O,
dans le Composé A38 : X=S,
dans le Composé A39 : X=Se


30

Composé A40 à A42, chacun représenté par la formule



où dans le Composé A40 : X=O,
dans le Composé A41 : X=S,
dans le Composé A42 : X=Se


Composé A43 à A45, chacun représenté par la formule



où dans le Composé A43 : X=O,
dans le Composé A44 : X=S,
dans le Composé A45 : X=Se

Composé A46 à A48, chacun représenté par la formule

35




40

où dans le Composé A46 : X=O,
dans le Composé A47 : X=S,
dans le Composé A48 : X=Se


45

Composé A49 à A51, chacun représenté par la formule



où dans le Composé A49 : X=O,
dans le Composé A50 : X=S,
dans le Composé A51 : X=Se

Composé A52 à A54, chacun représenté par la formule




où dans le Composé A52 : X=O,
dans le Composé A53 : X=S,
dans le Composé A54 : X=Se

50

55

Composé A55 à A57, chacun représenté par la formule

5




10

où dans le Composé A55 : X=O,
dans le Composé A56 : X=S,
dans le Composé A57 : X=Se


15

Composé A58 à A60, chacun représenté par la formule



où dans le Composé A58 : X=O,
dans le Composé A59 : X=S,
dans le Composé A60 : X=Se


Composé A61 à A63, chacun représenté par la formule



où dans le Composé A61 : X=O,
dans le Composé A62 : X=S,
dans le Composé A63 : X=Se

Composé A64 à A66, chacun représenté par la formule

20




25

où dans le Composé A64 : X=O,
dans le Composé A65 : X=S,
dans le Composé A66 : X=Se


30

Composé A67 à A69, chacun représenté par la formule



où dans le Composé A67 : X=O,
dans le Composé A68 : X=S,
dans le Composé A69 : X=Se


Composé A70 à A72, chacun représenté par la formule



où dans le Composé A70 : X=O,
dans le Composé A71 : X=S,
dans le Composé A72 : X=Se

Composé A73 à A75, chacun représenté par la formule

35




40

où dans le Composé A73 : X=O,
dans le Composé A74 : X=S,
dans le Composé A75 : X=Se


45

Composé A76 à A78, chacun représenté par la formule



où dans le Composé A76 : X=O,
dans le Composé A77 : X=S,
dans le Composé A78 : X=Se

Composé A79 à A81, chacun représenté par la formule



où dans le Composé A79 : X=O,
dans le Composé A80 : X=S,
dans le Composé A81 : X=Se

Composé A82 à A84, chacun

Composé A85 à A87, chacun


Composé A88 à A90, chacun

50

55

représenté par la formule

5




10

15


où dans le Composé A82 : X=O,
dans le Composé A83 : X=S,
dans le Composé A84 : X=Se

représenté par la formule



où dans le Composé A85 : X=O,
dans le Composé A86 : X=S,
dans le Composé A87 : X=Se


représenté par la formule



où dans le Composé A88 : X=O,
dans le Composé A89 : X=S,
dans le Composé A90 : X=Se

Composé A91 à A93, chacun
représenté par la formule

20




25

30


où dans le Composé A91 : X=O,
dans le Composé A92 : X=S,
dans le Composé A93 : X=Se

Composé A94 à A96, chacun
représenté par la formule



où dans le Composé A94 : X=O,
dans le Composé A95 : X=S,
dans le Composé A96 : X=Se


Composé A97 à A99, chacun
représenté par la formule



où dans le Composé A97 : X=O,
dans le Composé A98 : X=S,
dans le Composé A99 : X=Se

Composé A100 à A102, chacun
représenté par la formule

35




40

45


où dans le Composé A100 : X=O,
dans le Composé A101 : X=S,
dans le Composé A102 : X=Se

Composé A103 à A105, chacun
représenté par la formule



où dans le Composé A103 : X=O,
dans le Composé A104 : X=S,
dans le Composé A105 : X=Se

Composé A106 à A108, chacun
représenté par la formule



où dans le Composé A106 : X=O,
dans le Composé A107 : X=S,
dans le Composé A108 : X=Se

Composé A109 à A111, chacun
représenté par la formule


50

Composé A112 à A114, chacun
représenté par la formule

Composé A115 à A117, chacun
représenté par la formule


55

5




où dans le Composé A109 : X=O,
dans le Composé A110 : X=S,
dans le Composé A111 : X=Se

10




où dans le Composé A112 : X=O,
dans le Composé A113 : X=S,
dans le Composé A114 : X=Se




où dans le Composé A115 : X=O,
dans le Composé A116 : X=S,
dans le Composé A117 : X=Se

15


20



Composé B1,




Composé B2,




Composé B3,

30


35



Composé B4,



Composé B5,




Composé B6,

40


45

50



Composé B7,

et




Composé B8.

55

9. La composition selon la revendication 1, où le premier composé est sélectionné parmi le groupe constitué de :

5 Composé C28 à C30, chacun représenté par la formule




10

où dans le Composé C28 : X=O,
dans le Composé C29 : X=S,
dans le Composé C30 : X=Se


15

Composé C31 à C33, chacun représenté par la formule




où dans le Composé C31 : X=O,
dans le Composé C32 : X=S,
dans le Composé C33 : X=Se

Composé C34 à C36, chacun représenté par la formule



où dans le Composé C34 : X=O,
dans le Composé C35 : X=S,
dans le Composé C36 : X=Se

Composé C37 à C39, chacun représenté par la formule




20

où dans le Composé C37 : X=O,
dans le Composé C38 : X=S,
dans le Composé C39 : X=Se


25

Composé C40 à C42, chacun représenté par la formule



où dans le Composé C40 : X=O,
dans le Composé C41 : X=S,
dans le Composé C42 : X=Se


Composé C43 à C45, chacun représenté par la formule



où dans le Composé C43 : X=O,
dans le Composé C44 : X=S,
dans le Composé C45 : X=Se

30

Composé C46 à C48, chacun représenté par la formule




35

où dans le Composé C46 : X=O,
dans le Composé C47 : X=S,
dans le Composé C48 : X=Se


40

Composé C49 à C51, chacun représenté par la formule



où dans le Composé C49 : X=O,
dans le Composé C50 : X=S,
dans le Composé C51 : X=Se

Composé C52 à C54, chacun représenté par la formule



où dans le Composé C52 : X=O,
dans le Composé C53 : X=S,
dans le Composé C54 : X=Se


45

50

55

Composé C55 à C57, chacun représenté par la formule

5




10

où dans le Composé C55 : X=O,
dans le Composé C56 : X=S,
dans le Composé C57 : X=Se

15

Composé C64 à C66, chacun représenté par la formule

20




25

où dans le Composé C64 : X=O,
dans le Composé C65 : X=S,
dans le Composé C66 : X=Se

30

Composé C73 à C75, chacun représenté par la formule

35



40


où dans le Composé C73 : X=O,
dans le Composé C74 : X=S,
dans le Composé C75 : X=Se

45

50


55

Composé C58 à C60, chacun représenté par la formule




où dans le Composé C58 : X=O,
dans le Composé C59 : X=S,
dans le Composé C60 : X=Se

Composé C67 à C69, chacun représenté par la formule




où dans le Composé C67 : X=O,
dans le Composé C68 : X=S,
dans le Composé C69 : X=Se

Composé C61 à C63, chacun représenté par la formule




où dans le Composé C61 : X=O,
dans le Composé C62 : X=S,
dans le Composé C63 : X=Se

Composé C70 à C72, chacun représenté par la formule




où dans le Composé C70 : X=O,
dans le Composé C71 : X=S,
dans le Composé C72 : X=Se

Composé C76 à C78, chacun représenté par la formule



où dans le Composé C76 : X=O,
dans le Composé C77 : X=S,
dans le Composé C78 : X=Se


Composé C79 à C81, chacun représenté par la formule



où dans le Composé C79 : X=O,
dans le Composé C80 : X=S,
dans le Composé C81 : X=Se

Composé C82 à C84, chacun représenté par la formule

5




10

où dans le Composé C82 : X=O,
dans le Composé C83 : X=S,
dans le Composé C84 : X=Se


15

Composé C85 à C87, chacun représenté par la formule



où dans le Composé C85 : X=O,
dans le Composé C86 : X=S,
dans le Composé C87 : X=Se


Composé C88 à C90, chacun représenté par la formule



où dans le Composé C88 : X=O,
dans le Composé C89 : X=S,
dans le Composé C90 : X=Se

Composé C91 à C93, chacun représenté par la formule


20



où dans le Composé C91 : X=O,
dans le Composé C92 : X=S,
dans le Composé C93 : X=Se


25

Composé C94 à C96, chacun représenté par la formule



où dans le Composé C94 : X=O,
dans le Composé C95 : X=S,
dans le Composé C96 : X=Se


Composé C97 à C99, chacun représenté par la formule



où dans le Composé C97 : X=O,
dans le Composé C98 : X=S,
dans le Composé C99 : X=Se

Composé C100 à C102, chacun représenté par la formule


30



où dans le Composé C100 : X=O,
dans le Composé C101 : X=S,
dans le Composé C102 : X=Se


35

Composé C103 à C105, chacun représenté par la formule



où dans le Composé C103 : X=O,
dans le Composé C104 : X=S,
dans le Composé C105 : X=Se

Composé C106 à C108, chacun représenté par la formule



où dans le Composé C106 : X=O,
dans le Composé C107 : X=S,
dans le Composé C108 : X=Se

40


45

50

55

Composé C109 à C111, chacun représenté par la formule

5




10

où dans le Composé C109 : X=O,
dans le Composé C110 : X=S,
dans le Composé C111 : X=Se

15

Composé C118 à C120, chacun représenté par la formule

20




25

où dans le Composé C118 : X=O,
dans le Composé C119 : X=S,
dans le Composé C120 : X=Se

30

Composé C127 à C129, chacun représenté par la formule

35




40

où dans le Composé C127 : X=O,
dans le Composé C128 : X=S,
dans le Composé C129 : X=Se

Composé C136 à C138, chacun représenté par la formule


45

Composé C112 à C114, chacun représenté par la formule




où dans le Composé C112 : X=O,
dans le Composé C113 : X=S,
dans le Composé C114 : X=Se

Composé C121 à C123, chacun représenté par la formule



où dans le Composé C121 : X=O,
dans le Composé C122 : X=S,
dans le Composé C123 : X=Se


Composé C130 à C132, chacun représenté par la formule



où dans le Composé C130 : X=O,
dans le Composé C131 : X=S,
dans le Composé C132 : X=Se


Composé C139 à C141, chacun représenté par la formule

Composé C115 à C117, chacun représenté par la formule




où dans le Composé C115 : X=O,
dans le Composé C116 : X=S,
dans le Composé C117 : X=Se

Composé C124 à C126, chacun représenté par la formule



où dans le Composé C124 : X=O,
dans le Composé C125 : X=S,
dans le Composé C126 : X=Se

Composé C133 à C135, chacun représenté par la formule




où dans le Composé C133 : X=O,
dans le Composé C134 : X=S,
dans le Composé C135 : X=Se

Composé C142 à C144, chacun représenté par la formule

50


55

5




où dans le Composé C136 : X=O,
dans le Composé C137 : X=S,
dans le Composé C138 : X=Se

Composé C145 à C147, chacun
représenté par la formule



où dans le Composé C139 : X=O,
dans le Composé C140 : X=S,
dans le Composé C141 : X=Se

Composé C148 à C150, chacun
représenté par la formule




où dans le Composé C142 : X=O,
dans le Composé C143 : X=S,
dans le Composé C144 : X=Se


Composé C151 à C153, chacun
représenté par la formule

10


15



où dans le Composé C145 : X=O,
dans le Composé C146 : X=S,
dans le Composé C147 : X=Se



où dans le Composé C148 : X=O,
dans le Composé C149 : X=S,
dans le Composé C150 : X=Se



où dans le Composé C151 : X=O,
dans le Composé C152 : X=S,
dans le Composé C153 : X=Se


20

25


Composé C154 à C156, chacun
représenté par la formule

Composé C157 à C159, chacun
représenté par la formule


Composé C160 à C162, chacun
représenté par la formule



où dans le Composé C154 : X=O,
dans le Composé C155 : X=S,
dans le Composé C156 : X=Se



où dans le Composé C157 : X=O,
dans le Composé C158 : X=S,
dans le Composé C159 : X=Se



où dans le Composé C160 : X=O,
dans le Composé C161 : X=S,
dans le Composé C162 : X=Se


30

35


Composé C163 à C165, chacun
représenté par la formule

Composé C166 à C168, chacun
représenté par la formule


Composé C169 à C171, chacun
représenté par la formule



où dans le Composé C163 : X=O,
dans le Composé C164 : X=S,



où dans le Composé C166 : X=O,
dans le Composé C167 : X=S,



où dans le Composé C169 : X=O,
dans le Composé C170 : X=S,


45

50

55

dans le Composé C165 : X=Se


Composé C172 à C174, chacun représenté par la formule



où dans le Composé C172 : X=O,
dans le Composé C173 : X=S,
dans le Composé C174 : X=Se

dans le Composé C168 : X=Se


Composé C175 à C177, chacun représenté par la formule



où dans le Composé C175 : X=O,
dans le Composé C176 : X=S,
dans le Composé C177 : X=Se


dans le Composé C171 : X=Se

Composé C178 à C180, chacun représenté par la formule




où dans le Composé C178 : X=O,
dans le Composé C179 : X=S,
dans le Composé C180 : X=Se

Composé C181 à C183, chacun représenté par la formule




où dans le Composé C181 : X=O,
dans le Composé C182 : X=S,
dans le Composé C183 : X=Se

Composé C184 à C186, chacun représenté par la formule




où dans le Composé C184 : X=O,
dans le Composé C185 : X=S,
dans le Composé C186 : X=Se

Composé C187 à C189, chacun représenté par la formule




où dans le Composé C187 : X=O,
dans le Composé C188 : X=S,
dans le Composé C189 : X=Se

Composé C190 à C192, chacun représenté par la formule




où dans le Composé C190 : X=O,
dans le Composé C191 : X=S,
dans le Composé C192 : X=Se

Composé C193 à C195, chacun représenté par la formule



où dans le Composé C193 : X=O,
dans le Composé C194 : X=S,
dans le Composé C195 : X=Se

Composé C196 à C198, chacun représenté par la formule




où dans le Composé C196 : X=O,
dans le Composé C197 : X=S,
dans le Composé C198 : X=Se

Composé C199 à C201, chacun représenté par la formule

Composé C202 à C204, chacun représenté par la formule


Composé C205 à C207, chacun représenté par la formule

5




où dans le Composé C199 : X=O,
dans le Composé C200 : X=S,
dans le Composé C201 : X=Se

10



où dans le Composé C202 : X=O,
dans le Composé C203 : X=S,
dans le Composé C204 : X=Se




où dans le Composé C205 : X=O,
dans le Composé C206 : X=S,
dans le Composé C207 : X=Se

Composé C208 à C210, chacun
représenté par la formule


Composé C211 à C213, chacun
représenté par la formule

Composé C214 à C216, chacun
représenté par la formule


15



où dans le Composé C208 : X=O,
dans le Composé C209 : X=S,
dans le Composé C210 : X=Se



où dans le Composé C211 : X=O,
dans le Composé C212 : X=S,
dans le Composé C213 : X=Se



où dans le Composé C214 : X=O,
dans le Composé C215 : X=S,
dans le Composé C216 : X=Se


20

Composé C217 à C219, chacun
représenté par la formule


Composé C220 à C222, chacun
représenté par la formule

Composé C223 à C225, chacun
représenté par la formule


25



où dans le Composé C217 : X=O,
dans le Composé C218 : X=S,
dans le Composé C219 : X=Se



où dans le Composé C220 : X=O,
dans le Composé C221 : X=S,
dans le Composé C222 : X=Se



où dans le Composé C223 : X=O,
dans le Composé C224 : X=S,
dans le Composé C225 : X=Se


30

Composé C226 à C228, chacun
représenté par la formule


Composé C229 à C231, chacun
représenté par la formule

Composé C232 à C234, chacun
représenté par la formule


40



où dans le Composé C226 : X=O,



où dans le Composé C229 : X=O,



où dans le Composé C232 : X=O,

50

55

dans le Composé C227 : X=S,
dans le Composé C228 : X=Se

dans le Composé C230 : X=S,
dans le Composé C231 : X=Se




dans le Composé C233 : X=S,
dans le Composé C234 : X=Se

5 Composé C235 à C237, chacun
représenté par la formule

Composé C238 à C240, chacun
représenté par la formule

Composé C241 à C243, chacun
représenté par la formule

10



15

où dans le Composé C235 : X=O,
dans le Composé C236 : X=S,
dans le Composé C237 : X=Se

où dans le Composé C238 : X=O,
dans le Composé C239 : X=S,
dans le Composé C240 : X=Se




où dans le Composé C241 : X=O,
dans le Composé C242 : X=S,
dans le Composé C243 : X=Se

Composé C244 à C246, chacun
représenté par la formule

Composé C247 à C249, chacun
représenté par la formule

Composé C250 à C252, chacun
représenté par la formule

20



25

où dans le Composé C244 : X=O,
dans le Composé C245 : X=S,
dans le Composé C246 : X=Se


où dans le Composé C247 : X=O,
dans le Composé C248 : X=S,
dans le Composé C249 : X=Se

où dans le Composé C250 : X=O,
dans le Composé C251 : X=S,
dans le Composé C252 : X=Se, et

30

Composé C253 à C255, chacun
représenté par la formule

35



40

où dans le Composé C253 : X=O,
dans le Composé C254 : X=S,
dans le Composé C255 : X=Se


45

10. La composition selon la revendication 1, où le premier composé est sélectionné parmi le groupe constitué de :

50


55

5



10

15




20

25

Composé D1,
Composé D2, et

30




35

40

Composé D3,

11. La composition selon la revendication 1, où le second composé est sélectionné parmi le groupe constitué de

45




50

55

et

5

10



où:

15

X est sélectionné parmi le groupe constitué de O, S et Se ;

R¹ et R⁴ représentent chacun indépendamment une mono, di, ou tri, substitution, ou pas de substitution ;

R², R³, R⁵, et R⁶ représentent chacun indépendamment une mono, di, tri, ou tétra substitution, ou pas de substitution ; et

20

R¹ à R⁶ sont chacun indépendamment sélectionnés parmi le groupe constitué d'hydrogène, deutérium, benzène, biphenyle, terphenyle, naphthalène, fluorène, triphenylène, phénanthrène, dibenzofurane, dibenzothiophène, carbazole et de combinaisons de ceux-ci.

12. La composition selon la revendication 1, où le second composé est sélectionné parmi le groupe constitué de

25

30

35

40


45

50

55

Composé E1 à E3, chacun représenté par la formule


5



10


où dans le Composé E1 : X=O,
dans le Composé E2 : X=S,
dans le Composé E3 : X=Se

Composé E4 à E6, chacun représenté par la formule



où dans le Composé E4 : X=O,
dans le Composé E5 : X=S,
dans le Composé E6 : X=Se


Composé E7 à E9, chacun représenté par la formule



où dans le Composé E7 : X=O,
dans le Composé E8 : X=S,
dans le Composé E9 : X=Se

Composé E10 à E12, chacun représenté par la formule

15




20

où dans le Composé E10 : X=O,
dans le Composé E11 : X=S,
dans le Composé E12 : X=Se


25

Composé E13 à E15, chacun représenté par la formule



où dans le Composé E13 : X=O,
dans le Composé E14 : X=S,
dans le Composé E15 : X=Se


Composé E16 à E18, chacun représenté par la formule



où dans le Composé E16 : X=O,
dans le Composé E17 : X=S,
dans le Composé E18 : X=Se

Composé E19 à E21, chacun représenté par la formule

30




35

où dans le Composé E19 : X=O,
dans le Composé E20 : X=S,
dans le Composé E21 : X=Se

40

Composé E22 à E24, chacun représenté par la formule



où dans le Composé E22 : X=O,
dans le Composé E23 : X=S,
dans le Composé E24 : X=Se


45

50

55

Composé E25 à E27, chacun représenté par la formule

5




10


où dans le Composé E25 : X=O,
dans le Composé E26 : X=S,
dans le Composé E27 : X=Se

15

20



Composé E28,




Composé E29,

et

25

30




Composé E30


35

13. La composition selon la revendication 1, où le mélange du premier composé et du second composé est sélectionné parmi le groupe constitué de :


40




(Composé E1,



Composé C1) ;



(Composé E2,






Composé C2) ;

45

50

55

5






(Composé E5,

Composé C65) ;

(Composé E8,

15






20

Composé C74) ;

(Composé E11,

Composé C74) ;

25






30

(Composé CE17,

Composé C74) ;

(Composé E8,

35







40

Composé A5) ;

(Composé E8,

Composé C17) ;

45



55


(Composé E17,

Composé A5) ;

(Composé E26,


Composé C74) ;

5




(Composé E26,

10




Composé C248) ;




(Composé E28,

15




Composé C74) ;

20




(Composé E29,




Composé C74) ;

25



(Composé E30,

30



Composé C74) ;

(Composé E8, Composé C65) ;

(Composé E8, Composé A116) ; et (Composé E8, Composé D2).

35 **14.** Un premier dispositif comprenant un premier dispositif électroluminescent organique, le premier dispositif électroluminescent organique comprenant :

une anode ;

une cathode ; et

40 une couche organique, située entre l'anode et la cathode, comprenant une composition de matières telle que défini dans l'une quelconque des revendications précédentes.

45

50

55




FIGURE 1





FIGURE 2



Formula I



Formula II

FIGURE 3

FIGURE 4

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5844363 A [0003] [0026]
- US 6303238 B [0003] [0026]
- US 5707745 A [0003] [0026] [0029]
- US 7279704 B [0013] [0024] [0025] [0106]
- US 20140001456 A1 [0014]
- CN 102850329 A [0015]
- US 4769292 A [0023]
- US 20030230980 A [0026]
- US 5703436 A [0026]
- US 6097147 A [0026]
- US 20040174116 A [0026]
- US 5247190 A, Friend [0029]
- US 6091195 A, Forrest [0029]
- US 5834893 A, Bulovic [0029]
- US 6013982 A [0030]
- US 6087196 A [0030]
- US 6337102 B, Forrest [0030]
- US 7431968 B [0030]
- US 6294398 B [0030]
- US 6468819 B [0030]
- US 7968146 B [0031]
- US 2007023098 W [0031]
- US 2009042829 W [0031]
- WO 2007002683 A [0106]
- US 20030162053 A [0106]
- EP 1725079 A1 [0106]
- US 20050123751 A [0106]
- WO 2009018009 A [0106]
- US 20020158242 A [0106]
- US 20060240279 A [0106]
- US 20080220265 A [0106]
- WO 2011075644 A [0106]
- EP 2350216 A [0106]
- US 5061569 A [0106]
- EP 650955 A [0106]
- US 20080124572 A [0106]
- US 20070278938 A [0106]
- US 20080106190 A [0106]
- US 20110163302 A [0106]
- US 20080018221 A [0106]
- US 20060202194 A [0106]
- WO 2005014551 A [0106]
- WO 2006072002 A [0106]
- WO 2009066779 A [0106]
- WO 2009066778 A [0106]
- WO 2009063833 A [0106]
- US 20090045731 A [0106]
- US 20090045730 A [0106]
- WO 2009008311 A [0106]
- US 20090008605 A [0106]
- US 20090009065 A [0106]
- WO 2010056066 A [0106]
- WO 2011086863 A [0106]
- US 20030175553 A [0106]
- WO 2001039234 A [0106]
- US 20060280965 A [0106]
- WO 2009021126 A [0106]
- US 20090309488 A [0106]
- US 20090302743 A [0106]
- US 20100012931 A [0106]
- WO 2008056746 A [0106]
- WO 2010107244 A [0106]
- JP 2008074939 B [0106]
- US 20100187984 A [0106]
- WO 2004093207 A [0106]
- WO 2005089025 A [0106]
- WO 2006132173 A [0106]
- JP 200511610 B [0106]
- JP 2007254297 B [0106]
- WO 2007063796 A [0106]
- WO 2007063754 A [0106]
- WO 2004107822 A [0106]
- US 20050112407 A [0106]
- WO 2005030900 A [0106]
- US 20040137268 A [0106]
- US 20040137267 A [0106]
- US 20070190359 A [0106]
- WO 2006114966 A [0106]
- US 20090167162 A [0106]
- WO 2009086028 A [0106]
- US 20090030202 A [0106]
- US 20090017330 A [0106]
- US 20100084966 A [0106]
- US 20050238919 A [0106]
- WO 2009003898 A [0106]
- EP 2034538 A [0106]
- WO 2006100298 A [0106]
- US 20040115476 A [0106]
- US 20060121308 A [0106]
- US 7154114 B [0106]
- US 20030072964 A [0106]
- US 20070087321 A [0106]
- US 20080261076 A [0106]
- US 20100090591 A [0106]
- WO 2009100991 A [0106]
- WO 2008101842 A [0106]
- US 7232618 B [0106]
- WO 2003040257 A [0106]

- US 20070103060 A [0106]
- US 20050244673 A [0106]
- US 20020034656 A [0106]
- US 7332232 B [0106]
- US 20090108737 A [0106]
- WO 2010028151 A [0106]
- EP 1841834 B [0106]
- US 20060127696 A [0106]
- US 20090039776 A [0106]
- US 6921915 B [0106]
- US 20100244004 A [0106]
- US 6687266 B [0106]
- US 20060008670 A [0106]
- JP 2007123392 B [0106]
- WO 2010086089 A [0106]
- WO 2011044988 A [0106]
- WO 2009050290 A [0106]
- US 20090165846 A [0106]
- US 20080015355 A [0106]
- US 20010015432 A [0106]
- US 20100295032 A [0106]
- US 7250226 B [0106]
- US 7396598 B [0106]
- WO 2002015645 A [0106]
- US 20060263635 A [0106]
- US 20060182992 A [0106]
- WO 2009000673 A [0106]
- US 20070111026 A [0106]
- US 20030138657 A [0106]
- US 20030152802 A [0106]
- US 7090928 B [0106]
- WO 2002002714 A [0106]
- WO 2006009024 A [0106]
- US 20060251923 A [0106]
- US 20110057559 A [0106]
- US 20110204333 A [0106]
- US 7393599 B [0106]
- WO 2006056418 A [0106]
- US 20050260441 A [0106]
- WO 2005019373 A [0106]
- US 7534505 B [0106]
- WO 2011051404 A [0106]
- US 7445855 B [0106]
- US 20080297033 A [0106]
- US 20100148663 A [0106]
- US 7338722 B [0106]
- US 20020134984 A [0106]
- WO 2005123873 A [0106]
- WO 2007004380 A [0106]
- WO 2006082742 A [0106]
- WO 2006098120 A [0106]
- WO 2006103874 A [0106]
- US 7655323 B [0106]
- US 20050025993 A [0106]
- WO 2008132085 A [0106]
- WO 2010079051 A [0106]
- WO 2003060956 A [0106]
- US 20090179554 A [0106]
- US 20090115316 A [0106]
- US 7230107 B [0106]
- US 20090101870 A [0106]
- US 20040036077 A [0106]
- US 6528187 B [0106]

Non-patent literature cited in the description

- **BALDO et al.** Highly Efficient Phosphorescent Emission from Organic Electroluminescent Devices. *Nature*, 1998, vol. 395, 151-154 [0024]
- **BALDO et al.** Very high-efficiency green organic light-emitting devices based on electrophosphorescence. *Appl. Phys. Lett.*, 1999, vol. 75 (3), 4-6 [0024]
- **BULL et al.** *Mater. Sci.*, 2011, vol. 34, 7 [0049]
- *Appl. Phys. Lett.*, 1996, vol. 69, 2160 [0106]
- *J. Lumin.*, 1997, vol. 72-74, 985 [0106]
- *Appl. Phys. Lett.*, 2001, vol. 78, 673 [0106]
- *Synth. Met.*, 1997, vol. 87, 171 [0106]
- *SID Symposium Digest*, 2006, vol. 37, 923 [0106]
- *Appl. Phys. Lett.*, 1987, vol. 51, 913 [0106]
- *J. Mater. Chem.*, 1993, vol. 3, 319 [0106]
- *Appl. Phys. Lett.*, 2007, vol. 90, 183503 [0106]
- *Synth. Met.*, 1997, vol. 91, 209 [0106]
- *Adv. Mater.*, 1994, vol. 6, 677 [0106]
- *Synth. Met.*, 2000, vol. 111, 421 [0106]
- *Chem. Mater.*, 2003, vol. 15, 3148 [0106]
- *Appl. Phys. Lett.*, 2001, vol. 78, 1622 [0106]
- *Nature*, 1998, vol. 395, 151 [0106]
- *Appl. Phys. Lett.*, 2007, vol. 90, 123509 [0106]
- *Org. Electron.*, 2000, vol. 1, 15 [0106]
- *Appl. Phys. Lett.*, 2000, vol. 77, 2280 [0106]
- *J. Appl. Phys.*, 2001, vol. 90, 5048 [0106]
- *Appl. Phys. Lett.*, 2003, vol. 82, 2422 [0106]
- *Adv. Mater.*, 2007, vol. 19, 739 [0106]
- *Chem. Mater.*, 2005, vol. 17, 3532 [0106]
- *Adv. Mater.*, 2005, vol. 17, 1059 [0106]
- *Inorg. Chem.*, 2001, vol. 40, 1704 [0106]
- *Chem. Mater.*, 2004, vol. 16, 2480 [0106]
- *Adv. Mater.*, 2004, vol. 16, 2003 [0106]
- *Angew. Chem. Int. Ed.*, 2006, vol. 45, 7800 [0106]
- *Appl. Phys. Lett.*, 2005, vol. 86, 153505 [0106]
- *Chem. Lett.*, 2005, vol. 34, 592 [0106]
- *Chem. Commun.*, 2005, 2906 [0106]
- *Inorg. Chem.*, 2003, vol. 42, 1248 [0106]
- *Angew. Chem. Int. Ed.*, 2008, vol. 47, 4542 [0106]
- *Chem. Mater.*, 2006, vol. 18, 5119 [0106]
- *Inorg. Chem.*, 2007, vol. 46, 4308 [0106]
- *Organometallics*, 2004, vol. 23, 3745 [0106]
- *Appl. Phys. Lett.*, 1999, vol. 74, 1361 [0106]
- *Appl. Phys. Lett.*, 1999, vol. 75, 4 [0106]
- *Appl. Phys. Lett.*, 2001, vol. 79, 449 [0106]

EP 2 966 706 B1

- *Appl. Phys. Lett.*, 2002, vol. 81, 162 [0106]
- *Appl. Phys. Lett.*, 2001, vol. 79, 156 [0106]
- *Appl. Phys. Lett.*, 2006, vol. 89, 063504 [0106]
- *Chem. Lett.*, 1993, vol. 5, 905 [0106]
- *Appl. Phys. Lett.*, 2007, vol. 91, 263503 [0106]
- *Appl. Phys. Lett.*, 1999, vol. 74, 865 [0106]
- *Appl. Phys. Lett.*, 1989, vol. 55, 1489 [0106]
- *Jpn. J. Apply. Phys.*, 1993, vol. 32, L917 [0106]
- *Org. Electron.*, 2003, vol. 4, 113 [0106]
- *J. Am. Chem. Soc.*, 1998, vol. 120, 9714 [0106]
- *J. Am. Chem. Soc.*, 2000, vol. 122, 1832 [0106]

专利名称(译)	有机电致发光材料和器件		
公开(公告)号	EP2966706B1	公开(公告)日	2019-09-04
申请号	EP2015175686	申请日	2015-07-07
[标]申请(专利权)人(译)	环球展览公司		
申请(专利权)人(译)	通用显示器公司		
当前申请(专利权)人(译)	通用显示器公司		
[标]发明人	ZENG LICHANG LAYEK SUMAN BOUDREAULT PIERRE LUC T ELSHENAWY ZEINAB KOTTAS GREGG DYATKIN ALEXEY BORISOVICH JOSEPH SCOTT ADAMOVICH VADIM XIA CHUANJUN WANG TING CHIH YEAGER WALTER		
发明人	ZENG, LICHANG LAYEK, SUMAN BOUDREAULT, PIERRE-LUC T. ELSHENAWY, ZEINAB KOTTAS, GREGG DYATKIN, ALEXEY BORISOVICH JOSEPH, SCOTT ADAMOVICH, VADIM XIA, CHUANJUN WANG, TING-CHIH YEAGER, WALTER		
IPC分类号	H01L51/54		
CPC分类号	C07D251/24 C07D405/10 C07D409/04 C07D409/10 C07D421/10 H01L51/0052 H01L51/0067 H01L51/0071 H01L51/0073 H01L51/0074 C09K11/025 C09K11/06 C09K2211/1007 C09K2211/1059 C09K2211/1092 C09K2211/1096 C09K2211/185 H01L51/0085 H01L51/5016 H01L51/0058 H01L51/5072		
代理机构(译)	汉森NORBERT		
优先权	62/022300 2014-07-09 US 62/038925 2014-08-19 US 62/060192 2014-10-06 US 62/083490 2014-11-24 US 14/734712 2015-06-09 US		
其他公开文献	EP2966706A2 EP2966706A3		
外部链接	Espacenet		
摘要(译)			

描述了包含第一化合物和第二化合物的组合物。该组合物可以是具有根据式I的结构的第一化合物和具有根据式II的结构第二化合物的混合物。该组合物还可以是具有根据式III的结构的第一化合物的混合物，和还描述了具有根据式IV的结构第二化合物，包括用于式I和II或式III和IV的组合物的装置，例如OLED，以及制备它们的方法。

