



US 20150255721A1

(19) **United States**(12) **Patent Application Publication**
Lim et al.(10) **Pub. No.: US 2015/0255721 A1**
(43) **Pub. Date: Sep. 10, 2015**(54) **COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE INCLUDING THE
SAME***255/58* (2013.01); *C07D 307/91* (2013.01);
H01L 51/0061 (2013.01); *C07C 2103/54*
(2013.01); *H01L 51/0056* (2013.01); *H01L*
51/0073 (2013.01); *H01L 51/0058* (2013.01)(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-City (KR)(57) **ABSTRACT**


A compound is represented by Formula 1:

(72) Inventors: **Jin-O Lim**, Yongin-City (KR);
Young-Kook Kim, Yongin-City (KR);
Jong-Woo Kim, Yongin-City (KR);
Seok-Hwan Hwang, Yongin-City (KR)


Formula 1

(21) Appl. No.: **14/444,881**(22) Filed: **Jul. 28, 2014**(30) **Foreign Application Priority Data**

Mar. 10, 2014 (KR) 10-2014-0027950

Publication Classification(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 407/12 (2006.01)
C07D 307/91 (2006.01)
H01L 27/32 (2006.01)
C07C 255/58 (2006.01)
C07C 211/61 (2006.01)
H01L 51/50 (2006.01)(52) **U.S. Cl.**
CPC *H01L 51/006* (2013.01); *C07C 211/61*
(2013.01); *C07D 407/12* (2013.01); *H01L*
51/5096 (2013.01); *H01L 51/5056* (2013.01);
H01L 51/5072 (2013.01); *H01L 51/0003*
(2013.01); *H01L 27/3248* (2013.01); *C07C*

The substituents R_1 to R_{14} , Ar_1 to Ar_4 , X , and Y of Formula 1 are described herein. An organic light emitting device includes at least one of the compound represented by Formula 1. The organic light-emitting device including the compound may have a low driving voltage, high efficiency, high brightness, and long lifespan.



**COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE INCLUDING THE
SAME**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] This application claims priority to and the benefit of Korean Patent Application No. 10-2014-0027950, filed on Mar. 10, 2014, in the Korean Intellectual Property Office, the entire content of which is incorporated herein by reference.

BACKGROUND

[0002] 1. Field

[0003] One or more embodiments relate to a compound and an organic light-emitting device including the same.

[0004] 2. Description of the Related Art

[0005] Organic light emitting devices are self-emission devices that have wide viewing angles, high contrast ratios, short response time, and excellent brightness, driving voltage, and response speed characteristics, and produce full-color images.

[0006] An organic light-emitting device may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, are recombined in the emission layer to produce excitons. These excitons change from an excited state to a ground state, thereby generating light.


SUMMARY

[0007] Aspects of one or more embodiments are directed toward a novel compound and an organic light-emitting device including the same.

[0008] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0009] According to one or more embodiments of the present disclosure, a compound is represented by Formula 1 below:

Formula 1



[0010] where, in the above Formula 1,

[0011] R_1 to R_{14} may be each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group;

[0012] Ar_1 to Ar_4 may be each independently selected from a hydrogen, a deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, $-N(Q_1)(Q_2)$, $-Si(Q_3)(Q_4)(Q_5)$, and $-B(Q_6)(Q_7)$;

[0013] X and Y may be each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene

group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group;

[0014] m and n may be each independently an integer of 0 to 5; and

[0015] at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₂-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₂-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic hetero-condensed polycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic hetero-condensed polycyclic group may be selected from:

[0016] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0017] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇),

[0018] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

[0019] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid

or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅) and —B(Q₂₆)(Q₂₇); and

[0020] —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), where:

[0021] Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇ and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

[0022] According to one or more embodiments of the present disclosure, an organic light-emitting device includes: a first electrode; a second electrode facing the first electrode; and an organic layer including an emission layer between the first electrode and the second electrode, where the organic layer includes at least one of the compound described above.

[0023] According to one or more embodiments of the present disclosure, a flat panel display apparatus includes the organic light-emitting device, where the first electrode of the organic light-emitting device is electrically coupled to a source electrode or a drain electrode of a thin film transistor.

BRIEF DESCRIPTION OF THE DRAWINGS


[0024] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with the accompanying drawing, which is a schematic view of an organic light-emitting device according to an embodiment of the present disclosure.

DETAILED DESCRIPTION

[0025] Reference will now be made in detail to certain embodiments, examples of which are illustrated in the accompanying drawing, where like reference numerals refer to like elements throughout. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the accompanying drawing, to explain aspects of the present description. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list. Also, in the context of the present application, when a first element is referred to as being “on” a second element, it can be directly on the second element or be indirectly on the second element with one or more intervening elements interposed therebetween. Further, the use of “may” when describ-

ing embodiments of the present disclosure refers to “one or more embodiments of the present disclosure.”

[0026] A compound according to an embodiment is represented by Formula 1 below:



Formula 1

[0027] where, in Formula 1,

[0028] R₁ to R₁₄ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group;

[0029] Ar₁ to Ar₄ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), and —B(Q₆)(Q₇);

[0030] X and Y may be each independently selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkylene

group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group;

[0031] m and n may be each independently an integer of 0 to 5; and

[0032] at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₂-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₂-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic hetero-condensed polycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic hetero-condensed polycyclic group may be selected from:

[0033] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0034] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from:

[0035] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

[0036] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

[0037] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic

group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅) and —B(Q₂₆)(Q₂₇); and

[0038] —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), where:


[0039] Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇ and Q₃₁ to Q₃₇ may be each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

[0040] The compound of Formula 1 according to an embodiment includes a chrysene derivative core, and due to the inclusion of the chrysene derivative, the compound has a high glass transition temperature (T_g) or a high melting point. Accordingly, during emission (e.g., light emission), the compound of Formula 1 may exhibit increased durability at high temperature due to a heat resistance against Joule's heat (e.g., Joule heating, ohmic heating or resistive heating) that may occur in an organic layer, between organic layers, or between an organic layer and a metal electrode during emission (e.g., light emission). An organic light-emitting device manufactured by using (utilizing) the compound of Formula 1 according to an embodiment of the present disclosure has high durability during preservation and driving (e.g., during storage and operation).

[0041] Substituents of Formula 1 will now be described in more detail.

[0042] According to an embodiment, R₁ to R₁₄ in Formula 1 may be each independently selected from a hydrogen, a deuterium, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group.

[0043] According to another embodiment, R₁ and R₂ in Formula 1 may be each independently a substituted or unsubstituted C₁-C₂₀ alkyl group, or a group represented by Formula 2a below:



2a

[0044] Z₁ in Formula 2a may be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁ to C₂₀ alkyl group, a substituted or unsubstituted C₆ to C₂₀ aryl group, a substituted or unsubstituted C₁ to C₂₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group;

[0045] p is an integer of 1 to 5; and when p is two or more, a plurality of Z₁ may be identical to, or different from, each other; and * indicates a binding site to a neighboring atom.

[0046] According to another embodiment, R₃ to R₁₄ in Formula 1 may be each independently a hydrogen or a deuterium.

[0047] According to another embodiment of the present disclosure, X and Y in Formula 1 may be each independently selected from:


[0048] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group (an isoxazolylene group), a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzoimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group (a benzoxazolylene group), an isobenzooxazolylene group (an isobenzoxazolylene group), a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group;

[0049] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphth-


ylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylylylene group, a pyrenylene group, a chrysenylene group, a naphthacenylylene group, a picenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a pyrrolylylene group, a thiophenylylene group, a furanylylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group (an isoxazolylene group), a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, an isoindolylylene group, an indolylylene group, an indazolylene group, a purinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzoimidazolylene group, a benzofuranylylene group, a benzothiofenylene group, an isobenzothiazolylene group, a benzooxazolylene group (a benzoxazolylene group), an isobenzooxazolylene group (an isobenzoxazolylene group), a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylylene group, and an imidazopyrimidinylylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylylyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl

group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

[0050] According to another embodiment of the present disclosure, X and Y in Formula 1 may be each independently a group represented by Formulae 3a or 3b below:



3a



3b

[0051] where, in Formulae 3a and 3b,

[0052] Z_1 may be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 to C_{20} alkyl group, a substituted or unsubstituted C_6 to C_{20} aryl group, a substituted or unsubstituted C_1 to C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group;

[0053] p is an integer of 1 to 7; and when p is two or more, a plurality of Z_1 may be identical to, or different from, each other; and * indicates a binding site to a neighboring atom.

[0054] According to another embodiment of the present disclosure, Ar_1 to Ar_4 in Formula 1 may be each independently selected from:

[0055] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylylyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl


an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0056] a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, —Si(Q_3)(Q_4)(Q_5), a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzoox-

azolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group and a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, and

[0057] Q_3 to Q_5 may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

[0058] According to another embodiment, Ar_1 to Ar_4 in Formula 1 may be each independently represented by one of Formulae 4a to 4e below:



[0059] where, in Formulae 4a to 4e,

[0060] H_1 may be $CR_{21}R_{22}$, O, NR_{23} , or S;

[0061] R_{21} , R_{22} , R_{23} , and Z_1 may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_6 - C_{20} aryl group, a substituted or unsubstituted C_1 - C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a

substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, and $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$;


[0062] Q_3 to Q_5 may be each independently selected from a hydrogen, a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

[0063] p is an integer of 1 to 9; and when p is two or more, a plurality of Z_1 may be identical to, or different from, each other; and * indicates a binding site to a neighboring atom.


[0064] According to another embodiment, in Formula 1, Ar_1 or Ar_2 may be identical to Ar_3 or Ar_4 . For example, Ar_1 may be identical to Ar_3 or Ar_4 , or Ar_2 may be identical to Ar_3 or Ar_4 , or Ar_3 may be identical to Ar_1 or Ar_2 , or Ar_4 may be identical to Ar_1 or Ar_2 .

[0065] According to another embodiment, m and n in Formula 1 may be each independently 0 or 1.


[0066] According to an embodiment, Formula 1 may be represented by Formulae 2, 3, or 4 below.



Formula 2



Formula 3







Formula 4


[0067] Ar_1 to Ar_4 , X , Y , m , and n in Formulae 2, 3, and 4 are the same as those described above with respect Formula 1.

[0068] According to another embodiment, the compound of Formula 1 may be any one of compounds 1-62 illustrated below:


-continued




-continued




-continued




6






8




-continued



-continued




-continued




13

-continued




15


14



16




-continued




17

-continued




19


18



20




-continued




21


-continued



23




22




24

-continued




29


-continued



31




30




32

-continued




33


-continued



35




34




36

-continued




37

-continued




39


38




40




-continued




-continued





42



44




-continued




-continued

45




47

46




48

49




-continued




50


-continued



52




51




53

-continued




54


-continued



56




55



57


-continued

58




-continued


60




59



61



-continued



62

[0069] The compound represented by Formula 1 may be synthesized by using (utilizing) an organic synthetic method generally used in the art, and should be apparent to those of ordinary skill in the art. A synthesis method for the compound represented by Formula 1 should be apparent to one of ordinary skill in the art in view of the following embodiments.

[0070] The compound of Formula 1 may be used (utilized) between a pair of electrodes of an organic light-emitting device. For example, the compound may be included an electron transport region or an emission layer. Accordingly, an organic light-emitting device according to an embodiment includes: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode, where the organic layer includes an emission layer, and the organic layer includes at least one of the compounds described above.

[0071] The expression “(an organic layer) includes at least one of the compounds” used herein may include a case in which “(an organic layer)” includes one compound of Formula 1 and a case in which two or more different compounds of Formula 1 are included.

[0072] For example, the organic layer may include, as the compound, only Compound 1. In this regard, Compound 1 may exist in (or be present in) an electron transport layer of the organic light-emitting device. In another embodiment, the organic layer may include, as the compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may exist in (or be present in) one layer, such as a same or an identical layer (for example, Compound 1 and Compound 2 may all exist in, or be present in, one electron transport layer), or different layers (for example, Compound 1 may exist in, or be present in, an emission layer and Compound 2 may exist in, or be present in, an electron transport layer).

[0073] In one embodiment, the organic layer includes: i) a hole transport region that is disposed between the first electrode (anode) and the emission layer, and the hole transport region includes at least one selected from a hole injection layer, a hole transport layer, a buffer layer, and an electron

blocking layer, and ii) an electron transport region that is disposed between the emission layer and the second electrode (cathode), and the electron transport region includes at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer. For example, the emission layer may include the compound represented by Formula 1.

[0074] The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0075] The accompanying drawing is a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190. Here, in one embodiment, the first electrode 110 is connected (e.g., coupled) or configured to be connected (e.g., coupled) to a source electrode or drain electrode of a thin film transistor.

[0076] Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with the accompanying drawing.

[0077] In the accompanying drawing, a substrate may be additionally disposed under the first electrode 110 and/or above the second electrode 190. The substrate may include a glass substrate or transparent plastic substrate, each having excellent (or suitable) mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water repellency.

[0078] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode on the substrate. When the first electrode 110 is an anode, the material for the first electrode 110 may be selected from materials having a high work function to promote ease of hole supplying (or hole injection). The first electrode 110 may be a reflective electrode or a transmissive electrode. The material for the first electrode 120 may include a transparent and highly conductive material, and examples of such a material include indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), and zinc oxide (ZnO), but the present disclosure is not limited thereto. When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode, at least one of magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag) may be used (utilized), but the present disclosure is not limited thereto.

[0079] The first electrode 110 may have a single-layer structure, or a multi-layer structure including two or more layers. For example, the first electrode 110 may have a three-layer/ed structure of ITO/Ag/ITO, but the structure of the first electrode 110 is not limited thereto.

[0080] The organic layer 150 is disposed on the first electrode 110. The organic layer 150 may include an emission layer.

[0081] The organic layer 150 may further include a hole transport region disposed between the first electrode and the emission layer, and an electron transport region disposed between the emission layer and the second electrode.

[0082] The hole transport region may include at least one selected from a hole injection layer (HIL), a hole transport

layer (HTL), a buffer layer, and an electron blocking layer (EBL), and the electron transport region may include at least one selected from a hole blocking layer (HBL), an electron transport layer (ETL), and an electron injection layer (EIL), but the present disclosure is not limited thereto.

[0083] The hole transport region may have a single-layer/ed structure including (or formed of) a single material, a single-layer/ed structure including (or formed of) a plurality of different materials, or a multi-layer/ed structure having a plurality of layers including (or formed of) a plurality of different materials.

[0084] For example, the hole transport region may have a single-layer/ed structure including (or formed of) a plurality of different materials, or a structure of hole injection layer/hole transport layer, a structure of hole injection layer/hole transport layer/buffer layer, a structure of hole injection layer/buffer layer, a structure of hole transport layer/buffer layer, or a structure of hole injection layer/hole transport layer/electron blocking layer, where layers of each structure are sequentially stacked from the first electrode **110** in the stated order, but the present disclosure is not limited thereto.



[0085] When the hole transport region includes a hole injection layer, the hole injection layer may be formed on the first electrode **110** by using (utilizing) various suitable methods, such as vacuum deposition, spin coating, casting, a Langmuir-Blodgett (LB) method, ink-jet printing, laser-printing, or laser-induced thermal imaging, but the present disclosure is not limited thereto.

[0086] When a hole injection layer is formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec in consideration of a composition of a compound to be deposited for the hole injection layer, and the structure of the hole injection layer to be formed.


[0087] When a hole injection layer is formed by spin coating, the spin coating may be performed at a coating rate of about 2000 rpm to about 5000 rpm, and at a temperature of about 80° C. to 200° C. in consideration of a composition of a compound to be deposited for the hole injection layer, and the structure of the hole injection layer to be formed.

[0088] When the hole transport region includes a hole transport layer, the hole transport layer may be formed on either the first electrode **110** or the hole injection layer by using (utilizing) various suitable methods, such as vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When the hole transport layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole transport layer may be the same as those described above with respect to the deposition and coating conditions for the hole injection layer.

[0089] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β-NPB, TPD, Spiro-TPD, Spiro-NPB, α-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyanilinedodecylbenzenesulfonic acid (PaniDBSA), poly(3,4-ethylenedioxythiophene)poly(4-styrenesulfonate) (PEDOTPSS), polyanilinecamphor sulfonic acid (PaniCSA), (polyaniline)poly(4-styrenesulfonate) (PANIPSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below:




-continued




2-TNATA


-continued




Spiro-TPD




Spiro-NPB




NPB




α -NPB




β -NPB




TAPC



TPD



HMTPD



[0090] where, in Formulae 201 and 202,

[0091] L_{201} to L_{205} may be understood by referring to the description provided herein in connection with X and Y;

[0092] $\text{xa}1$ to $\text{xa}4$ may be each independently selected from 0, 1, 2, and 3;

[0093] $\text{xa}5$ may be selected from 1, 2, 3, 4, and 5; and

[0094] R_{201} to R_{204} may be each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group.

[0095] Where, in Formulae 201 and 202,

[0096] L_{201} to L_{205} may be each independently selected from:

[0097] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a carbazolylene group, and a triazinylylene group; and

[0098] a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-fluorenylylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a carbazolylene group, and a triazinylylene group, each substituted with at least one of a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinolinylyl group, an isoquinolinylyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group;

[0099] $\text{xa}1$ to $\text{xa}4$ may be each independently 0, 1, or 2;


[0100] $\text{xa}5$ may be 1, 2, or 3; and

[0101] R_{201} to R_{204} may be each independently selected from:


[0102] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinylyl group, an isoquinolinylyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group; and

[0103] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinylyl group, an isoquinolinylyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, an azulenylyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinylyl group, an isoquinolinylyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolyl group, and a triazinyl group.


[0104] The compound represented by Formula 201 may be represented by Formula 201A, but the present disclosure is not limited thereto:



[0105] For example, the compound represented by Formula 201 may be represented by Formula 201A-1 below, but the present disclosure is not limited thereto:



[0106] For example, the compound represented by Formula 202 may be represented by Formula 202A below, but the present disclosure is not limited thereto:



[0107] L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} in Formulae 201A, 201A-1 and 202A may be the same as described above with respect to Formulae 201 and 202, and R_{211} may be understood by referring to the description provided herein in connection with R_{203} (e.g., R_{211} may be the same as R_{203} as described above with respect to Formulae 201 and 202), and R_{213} to R_{216} may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic heterocondensed polycyclic group.

[0108] For example, in Formulae 201A, 201A-1, and 202A,

[0109] L_{201} to L_{203} may be each independently selected from:

[0110] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a carbazolylylene group, and a triazinylylene group; and

[0111] a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-fluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a carbazolylylene group, and a triazinylylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl

group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group;

[0112] $xa1$ to $xa3$ may be each independently 0 or 1;

[0113] R_{203} , R_{211} , and R_{212} may be each independently selected from:

[0114] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group; and

[0115] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, carboxylic acid a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group;

[0116] R_{213} and R_{214} may be each independently selected from:

[0117] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group;

[0118] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group;

[0119] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group; and

[0120] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group,

a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

[0121] R₂₁₅ and R₂₁₆ may be each independently selected from:

[0122] a hydrogen atom, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0123] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

[0124] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, and a triazinyl group; and

[0125] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phe-


nyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxaliny group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and

[0126] xa5 is 1 or 2.


[0127] R₂₁₃ and R₂₁₄ in Formulae 201A, and 201A-1 may, optionally, bind to each other (e.g., combine together) to form a saturated or unsaturated ring.

[0128] The compound represented by Formula 201 and the compound represented by Formula 202 may each include any one of compounds HT1 to HT20 illustrated below, but the present disclosure is not limited thereto.

HT1




HT2




-continued

HT3




-continued


HT5



HT4




HT6




-continued


HT7



HT8




HT9




-continued


HT10




HT11



HT12




-continued




HT13

-continued




HT17


HT18




HT14




HT19




HT15



HT20



HT16




[0129] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1000 Å. When the hole transport region includes both a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 10000 Å, for example, about 100 Å to

about 1000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2000 Å, for example about 100 Å to about 1500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within any of the foregoing ranges, suitable (or satisfactory) hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0130] The hole transport region may further include, in addition to the above-described materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0131] The charge-generation material may include, for example, a p-dopant. The p-dopant may be one of a quinone derivative, a metal oxide, and a cyano group-containing compound, but the present disclosure is not limited thereto. For example, non-limiting examples of the p-dopant include a quinone derivative, such as tetracyanoquinonodimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide, and Compound HT-D1 illustrated below, but the present disclosure is not limited thereto.



Compound HT-D1

F4-TCNQ

[0132] The hole transport region may further include, in addition to the hole injection layer and the hole transport layer, at least one of a buffer layer and an electron blocking layer. Because the buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, light-emission efficiency of a formed organic light-emitting device may be improved. Materials that are included in the hole transport region may be used (utilized) as a material included in the buffer layer. The electron blocking layer prevents (or reduces) injection of electrons from the electron transport region.


[0133] An emission layer is formed on the first electrode 110 or the hole transport region by using (utilizing) various suitable methods, such as vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When the emission layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the emission layer may be the same as those


described above with respect to the deposition and coating conditions for the hole injection layer.

[0134] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub pixel. In some embodiments, the emission layer may have a stacked structure of a red emission layer, a green emission layer, and a blue emission layer, or may include a red-light emission material, a green-light emission material, and a blue-light emission material, which are mixed with each other in a single layer, to emit white light.

[0135] The emission layer may include a host and a dopant.

[0136] The host may include at least one selected from TPBi, TBADN, ADN (also referred to as "DNA"), CBR, CDBP, and TCP:





[0137] According to another embodiment, the host may include a compound represented by Formula 301 below.



[0138] where, in Formula 301,

[0139] Ar_{301} may be selected from:

[0140] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

[0141] a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a

C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic hetero-condensed polycyclic group, and —Si(Q_{301})(Q_{302})(Q_{303}) (Q_{301} to Q_{303} may be each independently selected from a hydrogen, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_6 - C_{60} aryl group, and a C_1 - C_{60} heteroaryl group);

[0142] L_{301} may be understood by referring to the description provided in connection with L_{201} (e.g., L_{301} may be the same as L_{201} as described above with respect to Formulae 201 and 202);

[0143] R_{301} may be selected from:

[0144] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group;

[0145] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

[0146] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and

[0147] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

[0148] $\text{xb}1$ may be selected from 0, 1, 2, and 3; and

[0149] $\text{xb}2$ may be selected from 1, 2, 3, and 4.

[0150] Where, in Formula 301,

[0151] L_{301} may be selected from:

[0152] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, and a chrysenylene group; and

[0153] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, and a chrysenylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

[0154] R₃₀₁ may be selected from:


[0155] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group;

[0156] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

[0157] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group; and






[0158] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group, but the present disclosure is not limited thereto.

[0159] For example, the host may include a compound represented by Formula 301A below:




[0160] Substituents of Formula 301A may be understood by corresponding descriptions provided herein (e.g., in Formula 301A, L₃₀₁, x_{b1}, R₃₀₁, and x_{b2} may be the same as those described above with respect to Formula 301).

[0161] The compound represented by Formula 301 may include at least one of Compounds H1 to H42, but Formula 301 is not limited thereto:



Formula 301A

-continued



H6


H7

H8

H9

H10

-continued



H11

H12





H13

H14






H15

H16


-continued



-continued




-continued




H26


-continued




H32




H27




H33




H28




H29




H30



H31




H34




H35

-continued




H36


-continued




H39




H37



H40




H38



H41


-continued

H42




-continued

H45




[0162] According to another embodiment, the host may include at least one of Compounds H43 to H49 below, but the host is not limited thereto:


H43





H46




H47




H44



H48



-continued



[0163] The dopant may include the compound of Formula 1 according to an embodiment of the present disclosure.

[0164] An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but the dopant is not limited thereto.

[0165] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within any of the foregoing ranges, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.


[0166] Then, an electron transport region may be disposed on the emission layer.

[0167] The electron transport region may include at least one selected from a hole blocking layer, an electron transport layer (ETL), and an electron injection layer, but the electron transport region is not limited thereto.

[0168] For example, the electron transport region may have a structure of electron transport layer/electron injection layer or a structure of hole blocking layer/electron transport layer/electron injection layer, where layers of each structure are sequentially stacked from the emission layer in the stated order, but the electron transport region is not limited thereto.



[0169] According to an embodiment, the organic layer 150 of the organic light-emitting device may include an electron transport region disposed between the emission layer and the second electrode 190. The electron transport region may include at least one selected from an electron transport layer and an electron injection layer.

[0170] The electron transport layer may include at least one selected from BCP, Bphen, Alq₃, Balq, TAZ, and NTAZ.




H49

-continued




TAZ



NTAZ

[0171] According to another embodiment, the electron transport layer may include at least one compound selected from a compound represented by Formula 601 and a compound represented by Formula 602 illustrated below:



Formula 601

[0172] where, in Formula 601,

[0173] Ar₆₀₁ may be selected from:

[0174] a naphthalene, a heptalene, a fluorene, a spirofluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene;

[0175] a naphthalene, a heptalene, a fluorene, a spirofluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₃-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic hetero-condensed polycyclic group, and —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃) (Q₃₀₁

to Q₃₀₃ may be each independently selected from a hydrogen, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₆-C₆₀ aryl group, and a

[0176] C₁-C₆₀ heteroaryl group);

[0177] L₆₀₁ may be understood by referring to the description provided in connection with L₂₀₁ (e.g., L₆₀₁ may be the same as L₂₀₁ as described above with respect to Formulae 201 and 202); and

[0178] E₆₀₁ may be selected from:

[0179] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group; and


[0180] a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl

group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0181] xe1 may be selected from 0, 1, 2, and 3; and

[0182] xe2 may be selected from 1, 2, 3, and 4.

Formula 602



[0183] Where, in Formula 602,

[0184] X₆₁₁ may be N or C-(L₆₁₁)_{xe611}-R₆₁₁, X₆₁₂ may be N or C-(L₆₁₂)_{xe612}-R₆₁₂, X₆₁₃ may be N or C-(L₆₁₃)_{xe613}-R₆₁₃ and at least one of X₆₁₁ to X₆₁₃ may be N;

[0185] L₆₁₁ to L₆₁₆ may be understood by referring to the description provided herein in connection with L₂₀₁ (e.g., each of L₆₁₁ to L₆₁₆ may be the same L₂₀₁ as described above with respect to Formulae 201 and 202);

[0186] R₆₁₁ to R₆₁₆ may be each independently selected from:



[0187] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

[0188] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt

thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and


[0189] xe611 to xe616 may be each independently selected from 0, 1, 2, and 3.

[0190] The compound represented by Formula 601 and the compound represented by Formula 602 may each be selected from Compounds ET1 to ET15 illustrated below.




-continued


ET7



ET8




ET9




-continued


ET10



ET11




ET12




-continued


ET13



ET14




ET15




[0193] The metal-containing material may include a Li complex, but the present disclosure is not limited thereto. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

ET-D1



ET-D2




[0194] The electron transport region may include a hole blocking layer. The hole blocking layer may be formed (or included), when the emission layer includes a phosphorescent dopant, to prevent (or reduce) diffusion of excitons or holes into an electron transport layer.

[0195] When the electron transport region includes a hole blocking layer, the hole blocking layer may be formed on the emission layer by using (utilizing) various suitable methods, such as vacuum deposition, spin coating, casting, a Langmuir-Blodgett (LB) method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When the hole blocking layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole blocking layer may be the same as those described above with respect to the deposition and coating conditions for the hole injection layer.


[0196] The hole blocking layer may include, for example, at least one of BCP and Bphen, but the hole blocking layer is not limited thereto.

[0191] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within any of the foregoing ranges, the electron transport layer may have suitable (or satisfactory) electron transport characteristics without a substantial increase in driving voltage.

[0192] Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.



BCP



[0197] A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within any of the foregoing ranges, the hole blocking layer may have suitable (or excellent) hole blocking characteristics without a substantial increase in driving voltage.

[0198] The electron transport layer may be formed on the emission layer or the hole blocking layer by using (utilizing) various suitable methods, such as vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When an electron transport layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the electron transport layer may be the same as those described above with respect to the deposition and coating conditions for the hole injection layer.

[0199] According to an embodiment, the organic layer 150 of the organic light-emitting device includes an electron transport region disposed between the emission layer and the second electrode 190, wherein the electron transport region includes an electron transport layer.

[0200] The electron transport region may include an electron injection layer that allows electrons to be easily provided from the second electrode 190.

[0201] The electron injection layer may be formed on the electron transport layer by using (utilizing) various suitable methods, such as vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When an electron injection layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the electron injection layer may be the same as those described above with respect to the deposition and coating conditions for the hole injection layer.

[0202] The electron injection layer may include at least one selected from, LiF, NaCl, CsF, Li₂O, BaO, and LiQ, but the electron injection layer is not limited thereto.

[0203] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within any of the foregoing ranges, the electron injection layer may have suitable (or satisfactory) electron injection characteristics without a substantial increase in driving voltage.

[0204] The second electrode 190 is disposed on the organic layer 130 having such a structure. The second electrode 190 may be a cathode which is an electron injection electrode, and in this regard, a material for the second electrode 190 may be metal, an alloy, an electrically conductive compound, and a mixture thereof, which have a relatively low work function, but the second electrode is not limited thereto. Detailed examples of the material for the second electrode 190 include lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—

In), or magnesium-silver (Mg—Ag). According to another embodiment, the material for forming the second electrode 190 may be ITO or IZO. The second electrode 190 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode.

[0205] Also, an organic layer according to an embodiment may be formed by depositing the compound according to an embodiment, or may be formed by using (utilizing) a wet method in which the compound according to an embodiment is prepared in the form of solution and then the solution including the compound is used (utilized) for coating.

[0206] An organic light-emitting device according to an embodiment may be used (utilized) in various flat panel display apparatuses, such as a passive matrix organic light-emitting display apparatus or an active matrix organic light-emitting display apparatus. For example, when the organic light-emitting device is included in an active matrix organic light-emitting display apparatus, a first electrode disposed on a substrate acts as a pixel and may be electrically coupled to (e.g. electrically connected to) a source electrode or a drain electrode of a thin film transistor. In addition, the organic light-emitting device may be included in a flat panel display apparatus that emits light in opposite directions.

[0207] Hereinbefore, the organic light-emitting device has been described with reference to the accompanying drawing, but the present disclosure is not limited thereto.

[0208] Hereinafter, definitions of substituents used herein are described (the number of carbon numbers used to describe (or restrict) a substituent is not limited, and does not limit properties of the substituent, and unless defined otherwise, the definition of the substituent is consistent with a general definition thereof).

[0209] As used herein, a C₁-C₆₀ alkyl group refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and detailed examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group, but the present disclosure is not limited thereto. A C₁-C₆₀ alkyne group used herein refers to a divalent group having the same structure as that described above with respect to the C₁-C₆₀ alkyl group.

[0210] As used herein, a C₁-C₆₀ alkoxy group refers to a monovalent group represented by —OA₁₀₁ (where A₁₀₁ is the above-described C₁-C₆₀ alkyl group), and detailed examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group, but the present disclosure is not limited thereto.

[0211] As used herein, a C₂-C₆₀ alkenyl group refers to a hydrocarbon group having at least one carbon double bond at a center portion (e.g., in the middle) or terminal end of a C₂-C₆₀ alkyl group corresponding to the above-described C₁-C₆₀ alkyl group, and detailed examples thereof include an ethenyl group, a propenyl group (or a propenyl group), and a butenyl group, but the present disclosure is not limited thereto. As used herein, a C₂-C₆₀ alkenylene group refers to a divalent group having the same structure as that described above with respect to the C₂-C₆₀ alkenyl group.

[0212] As used herein, a C₂-C₆₀ alkynyl group refers to a hydrocarbon group having one carbon triple bond at a center portion (e.g., in the middle) or terminal end of the above-described C₂-C₆₀ alkyl group, and detailed examples thereof include an ethynyl group, and a propynyl group, but the present disclosure is not limited thereto. As used herein, a

C_2 - C_{60} alkynylene group refers to a divalent group having the same structure as that described above with respect to the C_2 - C_{60} alkynyl group.

[0213] As used herein, a C_3 - C_{10} cycloalkyl group refers to a monovalent hydrocarbon monocyclic group having 3 to 10 carbon atoms, and detailed examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group, but the present disclosure is not limited thereto. As used herein, a C_3 - C_{10} cycloalkylene group refers to a divalent group having the same structure as that described above with respect to the C_3 - C_{10} cycloalkyl group.

[0214] As used herein, a C_2 - C_{10} heterocycloalkyl group refers to a monovalent monocyclic group having at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 2 to 10 carbon atoms, and detailed examples thereof include a tetrahydrofuranyl group, and a tetrahydrothiophenyl group, but the present disclosure is not limited thereto. As used herein, a C_2 - C_{10} heterocycloalkylene group refers to a divalent group having the same structure as that described above with respect to the C_2 - C_{10} heterocycloalkyl group.

[0215] As used herein, a C_3 - C_{10} cycloalkenyl group refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in the ring thereof and does not have aromaticity, and detailed examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group, but the present disclosure is not limited thereto. As used herein, a C_3 - C_{10} cycloalkenylene group refers to a divalent group having the same structure as that described above with respect to the C_3 - C_{10} cycloalkenyl group.

[0216] As used herein, a C_2 - C_{10} heterocycloalkenyl group refers to a monovalent monocyclic group that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, 2 to 10 carbon atoms, and at least one double bond in its ring. Detailed examples of the C_2 - C_{10} heterocycloalkenyl group include a 2,3-hydrofuranyl group and a 2,3-hydrothiophenyl group, but the present disclosure is not limited thereto. As used herein, a C_2 - C_{10} heterocycloalkenylene group refers to a divalent group having the same structure as that described above with respect to the C_2 - C_{10} heterocycloalkenyl group.

[0217] As used herein, a C_6 - C_{60} aryl group refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C_6 - C_{60} arylene group, as used herein, refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the C_6 - C_{60} aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group, but the present disclosure is not limited thereto. When the C_6 - C_{60} aryl group and the C_6 - C_{60} arylene group each include two or more rings, the rings may, optionally, be fused to each other (e.g., combined).

[0218] As used herein, a C_1 - C_{60} heteroaryl group refers to a monovalent group having a carbocyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 1 to 60 carbon atoms. As used herein, a C_1 - C_{60} heteroarylene group refers to a divalent group having a carbocyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 1 to 60 carbon atoms. Examples of the C_1 - C_{60} heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group, but the

present disclosure is not limited thereto. When the C_1 - C_{60} heteroaryl group and the C_2 - C_{60} heteroarylene group each include two or more rings, the rings may, optionally, be fused to each other (e.g., combined).

[0219] As used herein, a C_6 - C_{60} aryloxy group used herein indicates $—OA_{102}$ (where A_{102} is the C_6 - C_{60} aryl group), and a C_6 - C_{60} arylthio group indicates $—SA_{103}$ (where A_{103} is the C_6 - C_{60} aryl group).

[0220] As used herein, a monovalent non-aromatic condensed polycyclic group refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other, only carbon atoms as a ring forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group, but the present disclosure is not limited thereto. As used herein, a divalent non-aromatic condensed polycyclic group refers to a divalent group having the same structure as that described above with respect to the monovalent non-aromatic condensed polycyclic group.

[0221] As used herein, a monovalent non-aromatic condensed heteropolycyclic group refers to a monovalent group (for example, having 2 to 60 carbon atoms) that has two or more rings condensed to each other (e.g., combined with each other), has a heteroatom selected from N, O, P, and S, other than carbon atoms, as a ring forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group, but the present disclosure is not limited thereto. As used herein, a divalent non-aromatic condensed heteropolycyclic group refers to a divalent group having the same structure as that described above with respect to the monovalent non-aromatic condensed heteropolycyclic group.

[0222] In the present specification, at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_2 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_2 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic hetero-condensed polycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_2 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_2 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic hetero-condensed polycyclic group may be selected from:

[0223] a deuterium, $—F$, $—Cl$, $—Br$, $—I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0224] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, $—F$,

—Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

[0225] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

[0226] a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅) and —B(Q₂₆)(Q₂₇); and

[0227] —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), where:

[0228] Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇ and Q₃₁ to Q₃₇ may be each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

[0229] For example, at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₂-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₂-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic hetero-condensed polycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl

group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic hetero-condensed polycyclic group may be selected from:

[0230] a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0231] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quiazolinyl group, a cinolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

[0232] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a

picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0233] a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyly group, a heptalenyl group, an indacenyly group, an acenaphthyl group, a fluorenyly group, a spirofluorenyly group, a benzofluorenyly group, a dibenzofluorenyly group, a phenalenyl group, a phenanthrenyl group, an anthracenyly group, a fluoranthenyly group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyly group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphonic acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyly group, a heptalenyl group, an indacenyly group, an acenaphthyl group, a fluorenyly group, a spirofluorenyly group, a ben-

zofluorenyly group, a dibenzofluorenyly group, a phenalenyl group, a phenanthrenyl group, an anthracenyly group, a fluoranthenyly group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyly group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and

[0234] —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), where:

[0235] Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyly group, a heptalenyl group, an indacenyly group, an acenaphthyl group, a fluorenyly group, a spirofluorenyly group, a benzofluorenyly group, a dibenzofluorenyly group, a phenalenyl group, a phenanthrenyl group, an anthracenyly group, a fluoranthenyly group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyly group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyly group, a pentacenyly group, a rubicenyly group, a coronenyly group, an ovalenyly group, a pyrrolyly group, a thiophenyly group, a furanyly group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group (an isoxazolyl group), a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group (a benzoxazolyl group), an isobenzooxazolyl group (an isobenzoxazolyl group), a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a

dibenzofuranyl group, a dibenzothiophenyl group, a benzo-carbazolyl group, a dibenzocarbazolyl group, a thiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

[0236] As used herein, the term “Ph” refers to a phenyl group, the term “Me”, as used herein, refers to a methyl group, the term “Et”, as used herein, refers to an ethyl group, and the term “tert-Bu” or “Bu”, as used herein, refers to a tert-butyl group.




[0237] Hereinafter, an organic light-emitting device according to an embodiment will be described with reference to Synthesis Examples and Examples. In describing the Synthesis Examples, the wording “B was used instead of A” means that a molar equivalent of A was identical to a molar equivalent of B.

SYNTHESIS EXAMPLE

Synthesis Example 1

Synthesis of Compound 1

[0238]



Synthesis of Intermediate I-1

[0239] 6.6 g (20 mmol) of Compound I was dissolved in 200 mL of ethylether, and then, 3.2 g (24 mmol) of AlCl_3 was slowly added dropwise thereto, and the resultant was stirred for 30 minutes. At a temperature of 0°C ., 1.1 g (30 mmol) of lithium aluminum hydride was slowly added dropwise thereto, and then, the resultant was stirred for 1 hour while refluxing. The reaction solution was cooled to room temperature, and then, 6 M HCl was added thereto. 40 mL of water was added thereto, and then, the resultant was extracted three times with 40 mL of ethylether to obtain an organic layer, which was then dried with magnesium sulfate. The residual was obtained by removing a solvent used herein by evaporation, and the residual was then separation-purified by silicagel column chromatography to obtain 5.4 g (yield of 85%) of Intermediate I-1. The obtained compound was identified by LC-MS. $\text{C}_{25}\text{H}_{16}$ M^+ 317.1

Synthesis of Intermediate I-2

[0240] 5.4 g (17 mmol) of Intermediate I-1 was dissolved in 150 mL of dimethyl sulfoxide (DMSO), and then, 14 g (127 mmol) of potassium t-butoxide was added thereto to form a mixture. Then, the mixture was stirred at a temperature of 70°C . for 30 minutes. 10 mL (136 mmol) of iodomethane was slowly added dropwise thereto to form a mixture, and then the

mixture was stirred for 1 hour. When the reaction stopped, distilled water was added thereto to produce a solid product, and then, recrystallization was performed thereon to produce 3.5 g (yield of 60%) of Intermediate I-2. The obtained compound was identified by LC-MS. $C_{27}H_{20} M^+$ 345.2

Synthesis of Intermediate I-3

[0241] 3.4 g (10 mmol) of Intermediate I-2 was dissolved in 60 mL of tetrahydrofuran (THF), and then, 1.1 mL (21 mmol) of Br_2 was slowly added dropwise thereto to form a mixture, and then, the mixture was stirred at room temperature for 20 hours. 10 mL (136 mmol) of iodomethane was slowly added dropwise thereto, and then the mixture was stirred for 1 hour. 40 mL of water was added thereto, and then, the resultant was extracted three times with 30 mL of ethylether to obtain an organic layer, which was then dried with magnesium sulfate. The residual was obtained by removing a solvent used herein by evaporation, and the residual was then separation-purified by silicagel column chromatography to obtain 2.5 g (yield of 50%) of Intermediate I-3. The obtained compound was identified by LC-MS. $C_{27}H_{18}Br_2 M^+$ 501.0


Synthesis of Compound 1

[0242] 2.5 g (5.0 mmol) of Intermediate I-3, 1.7 g (10 mmol) of diphenylamine, 0.09 g (0.1 mmol) of $Pd_2(dba)_3$, 0.02 g (0.1 mmol) of $P(tBu)_3$, and 0.72 g (7.5 mmol) of NaOtBu were dissolved in 40 mL of toluene to form a mixture, and then, the mixture was stirred at a temperature of 80° C. for 3 hours. The reaction solution was cooled to room temperature, and then, 30 mL of water was added thereto, and an extraction process was performed thereon three times with 30 mL of ethylether. An organic layer was collected and then dried by using (utilizing) magnesium sulfate. Then, the residual was obtained by evaporating a solvent therefrom, and the residual was separation-purified by silica gel column chromatography to obtain 2.78 g (yield of 82%) of Compound 1. An obtained compound was confirmed by 1H NMR and MS/FAB. $C_{51}H_{38}N_2$; calc. 678.30, found 678.29

Synthesis Example 2

Synthesis of Compound 4

[0243]




[0244] 3.35 g (yield of 78%) of Compound 4 was synthesized as in the synthesis of Compound 1, except that Intermediate A-1 was used (utilized) instead of diphenylamine. An obtained compound was confirmed by ^1H NMR and MS/FAB. $\text{C}_{63}\text{H}_{42}\text{N}_2\text{O}_2$: calc. 858.32. found 858.34.

Synthesis Example 3

Synthesis of Compound 15

[0245]



Synthesis of Intermediate I-4

[0246] 10 g (20 mmol) of Intermediate I-3, 2.2 g (10 mmol) of Intermediate A-2, 0.18 g (0.2 mmol) of $\text{Pd}_2(\text{dba})_3$, 0.04 g (0.2 mmol) of $\text{P}(\text{tBu})_3$, and 1.4 g (15 mmol) of NaOtBu were dissolved in 50 mL of toluene to form a mixture, and then, the

mixture was stirred at a temperature of 80°C . for 3 hours. The reaction solution was cooled to room temperature, and then, 30 mL of water was added thereto, and an extraction process was performed thereon three times with 40 mL of ethylether. An organic layer was collected and then dried by using (utilizing) magnesium sulfate. Then, the residual was obtained by

removing a solvent therefrom by evaporation, and the residual was subjected to silica gel column chromatography to obtain 3.2 g (yield of 50%) of Intermediate I-4. The obtained compound was identified by LC-MS. $C_{42}H_{31}BrN_2$ M^+ 643.2

Synthesis of Intermediate I-5

[0247] 3.2 g (5.0 mmol) of Intermediate I-4, 1.0 g (5.0 mmol) of 4-bromophenylboronic acid, 0.29 g (0.25 mmol) of $Pd(PPh_3)_4$, and 1.6 g (15 mmol) of Na_2CO_3 were dissolved in 40 mL of a toluene/ethanol (a volumetric ratio of 21) mixed solution, and then, the resultant solution was stirred at a temperature of 90° C. for 10 hours. The reaction solution was cooled to room temperature, and then, 30 mL of water was added thereto, and an extraction process was performed thereon three times with 30 mL of ethylether. An organic layer was collected and then dried by using (utilizing) magnesium sulfate. Then, the residual was obtained by evaporating a solvent therefrom, and the residual was separation-purified by silica gel column chromatography to obtain 2.9 g (yield of 81%) of Intermediate I-5. The obtained compound was identified by LC-MS. $C_{48}H_{35}BrN_2$ M^+ 719.2


Synthesis of Compound 15

[0248] 2.93 g (yield of 68%) of Compound 15 was synthesized as in the synthesis of Intermediate I-4, except that Intermediate I-5 was used (utilized) instead of Intermediate I-3. An obtained compound was confirmed by 1H NMR and MS/FAB. $C_{63}H_{48}N_4$: calc. 860.39. found 860.40.

Synthesis Example 4

Synthesis of Compound 27

[0249]




[0250] 3.23 g (yield of 83%) of Compound 27 was synthesized as in the synthesis of Compound 1, except that Intermediate A-3 was used (utilized) instead of diphenylamine. An obtained compound was confirmed by 1H NMR and MS/FAB. $C_{59}H_{42}N_2$: calc. 778.33. found 778.32.


Synthesis Example 5

Synthesis of Compound 49

[0251]



-continued



Synthesis of Intermediate I-6

[0252] 3.54 g (yield of 52%) of Intermediate I-6 was synthesized as in the synthesis of Intermediate I-4, except that Intermediate A-1 was used (utilized) instead of Intermediate A-2. The obtained compound was identified by LC-MS. $C_{45}H_{30}BrNO$ M^+ 680.1

Synthesis of Compound 49

[0253] 2.88 g (yield of 75%) of Compound 49 was synthesized as in the synthesis of Intermediate I-4, except that Intermediate I-6 was used (utilized) instead of Intermediate I-3,

and diphenylamine was used instead of Intermediate A-2. An obtained compound was confirmed by 1H NMR and MS/FAB. $C_{57}H_{40}N_2O$: calc. 768.31. found 768.30.

[0254] Additional compounds were synthesized by using (utilizing) the synthesis methods described above and appropriate intermediate materials. 1H NMR and MS/FAB results of the additional synthetic compounds are shown in Table 1 below.

[0255] Methods of synthesizing compounds other than the compounds shown in Table 1 should be apparent to one of ordinary skill in the art by referring to the synthesis methods (or path) and source materials described above.

TABLE 1

Compound	1H NMR ($CDCl_3$, 400 MHz)	MS/FAB	
		found	calc.
1	$\delta = 8.92$ (s, 1H), 8.51 (d, 1H), 8.02 (d, 1H), 7.90 (s, 1H), 7.82-7.80 (m, 1H), 7.71-7.63 (m, 2H), 7.59-7.49 (m, 2H), 7.42-7.38 (m, 1H), 7.08-7.03 (m, 8H), 6.79-6.78 (m, 1H), 6.66-6.60 (m, 5H), 6.30-6.24 (m, 8H), 1.63 (s, 6H)	678.29	678.30
3	$\delta = 8.91$ (s, 1H), 8.50 (d, 1H), 8.01 (d, 1H), 7.89 (s, 1H), 7.82-7.80 (m, 1H), 7.71-7.55 (m, 3H), 7.50-7.48 (m, 1H), 7.41-7.37 (m, 5H), 7.08-7.04 (m, 4H), 6.81-6.80 (m, 1H), 6.68-6.62 (m, 3H), 6.55-6.51 (m, 4H), 6.27-6.20 (m, 4H), 1.63 (s, 6H), 0.24 (s, 18H)	822.40	822.38
4	$\delta = 8.92$ (s, 1H), 8.51 (d, 1H), 8.49 (s, 1H), 8.01 (d, 1H), 7.95-7.93 (m, 1H), 7.83-7.81 (m, 2H), 7.75-7.64 (m, 6H), 7.59-7.55 (m, 1H), 7.50-7.46 (m, 3H), 7.42-7.38 (m, 3H), 7.10-6.95 (m, 7H), 6.81-6.80 (m, 1H), 6.67-6.63 (m, 3H), 6.44 (d, 1H), 6.26-6.24 (m, 4H), 1.64 (s, 6H)	858.34	858.32
8	$\delta = 8.91$ (s, 1H), 8.51 (d, 1H), 8.49 (s, 1H), 8.01 (d, 1H), 7.97-7.95 (m, 1H), 7.83-7.81 (m, 1H), 7.77-7.62 (m, 4H), 7.60-7.55 (m, 3H), 7.50-7.38 (m, 7H), 7.22-7.14 (m, 2H), 7.07-6.93 (m, 7H), 6.72 (d, 1H), 6.66-6.62 (m, 2H), 6.47-6.45 (dd, 1H), 6.25-6.23 (m, 2H), 6.14 (d, 1H), 6.09-6.07 (m, 2H), 1.63 (s, 6H)	844.36	844.35
12	$\delta = 8.91$ (s, 1H), 8.51 (d, 1H), 8.29 (s, 1H), 8.02 (d, 1H), 7.83-7.81 (m, 2H), 7.77-7.73 (m, 2H), 7.69-7.64 (m, 3H), 7.59-7.55 (m, 1H), 7.50-7.46 (m, 2H), 7.42-7.40 (m, 3H), 7.35-7.31 (m, 1H), 7.12-6.99 (m, 7H), 6.80-6.79 (m, 1H), 6.69-6.64 (m, 4H), 6.48-6.46 (m, 2H), 6.27-6.24 (m, 2H), 6.19-6.17 (m, 2H), 1.63 (s, 6H), 1.61 (s, 6H)	884.39	884.38
15	$\delta = 8.99$ (s, 1H), 8.77 (d, 1H), 8.53 (s, 1H), 8.19 (d, 1H), 7.84-7.82 (m, 1H), 7.72-7.65 (m, 2H), 7.60-7.57 (m, 1H), 7.50-7.48 (m, 3H), 7.43-7.38 (m, 1H), 7.31-7.30 (m, 4H), 7.10-7.04 (m, 4H), 6.95-6.93 (m, 2H), 6.70-6.64 (m, 3H), 6.59-6.56 (dd, 1H), 6.26-6.24 (m, 2H), 6.10-6.08 (m, 2H), 2.25 (d, 12H), 1.63 (s, 6H)	860.40	860.39
19	$\delta = 8.96$ (s, 1H), 8.51 (d, 1H), 8.09-8.07 (m, 1H), 8.02 (d, 1H), 7.90 (s, 1H), 7.82-7.80 (m, 1H), 7.69-7.49 (m, 14H), 7.42-7.38 (m, 4H), 7.33-7.32 (m, 1H), 7.15-7.04 (m, 7H),	924.40	924.39

TABLE 1-continued

Compound	¹ H NMR (CDCl ₃ , 400 MHz)	MS/FAB	
		found	calc.
22	6.86-6.83 (m, 1H), 6.69-6.64 (m, 3H), 6.20-6.15 (m, 7H), 1.67 (s, 6H) δ = 8.92 (s, 1H), 8.51 (d, 1H), 8.02 (d, 1H), 7.90 (s, 1H), 7.82-7.80 (m, 1H), 7.71-7.64 (m, 2H), 7.59-7.55 (m, 5H), 7.49-7.35 (m, 12H), 7.11-7.08 (m, 4H), 6.85-6.84 (m, 1H), 6.71-6.67 (m, 3H), 6.50-6.46 (m, 4H), 6.28-6.21 (m, 4H), 1.63 (s, 6H)	830.36	830.37
24	δ = 8.91 (s, 1H), 8.53 (d, 1H), 8.04-8.01 (m, 2H), 7.69-7.50 (m, 23H), 7.41-7.38 (m, 3H), 7.34-7.31 (m, 2H), 7.13-7.04 (m, 6H), 6.75-6.74 (m, 1H), 6.68-6.62 (m, 2H), 6.50-6.47 (dd, 1H), 6.22-6.19 (m, 2H), 6.13-6.11 (m, 2H), 1.63 (s, 6H)	1018.40	1018.41
27	δ = 8.92 (s, 1H), 8.51 (d, 1H), 8.02 (m, 1H), 7.91-7.83 (m, 4H), 7.74 (d, 1H), 7.68-7.64 (m, 1H), 7.60-7.39 (m, 11H), 7.34 (d, 1H), 7.30 (d, 1H), 7.12-7.04 (m, 6H), 6.84 (d, 1H), 6.66-6.61 (m, 3H), 6.26-6.24 (m, 2H), 6.20-6.18 (m, 2H), 1.62 (s, 6H)	778.32	778.33
30	δ = 8.92 (s, 1H), 8.52 (d, 1H), 8.36 (d, 2H), 8.20-8.17 (m, 2H), 8.05-8.02 (m, 2H), 7.98-7.85 (m, 1H), 7.85-7.82 (dd, 1H), 7.73-7.62 (m, 9H), 7.59-7.55 (m, 3H), 7.51-7.49 (m, 1H), 7.43-7.38 (m, 3H), 7.07-7.02 (m, 5H), 6.76-6.75 (m, 1H), 6.65-6.59 (m, 3H), 6.46-6.43 (dd, 1H), 6.16-6.14 (m, 2H), 6.08-6.06 (m, 2H), 1.63 (s, 6H)	878.38	878.37
35	δ = 8.92 (s, 1H), 8.51 (d, 1H), 8.02 (d, 1H), 7.90 (s, 1H), 7.83-7.80 (m, 1H), 7.70 (d, 1H), 7.68-7.64 (m, 1H), 7.59-7.55 (m, 1H), 7.50-7.48 (m, 1H), 7.42-7.38 (m, 1H), 7.08-7.03 (m, 4H), 6.79-6.78 (m, 1H), 6.67-6.59 (m, 3H), 6.20-6.15 (m, 4H), 1.63 (s, 6H)	688.36	688.37
38	δ = 8.92 (s, 1H), 8.52 (d, 1H), 8.03 (d, 1H), 7.88 (s, 1H), 7.84-7.79 (m, 5H), 7.74 (d, 1H), 7.68-7.64 (m, 1H), 7.59-7.48 (m, 4H), 7.42-7.37 (m, 3H), 7.26-7.21 (m, 2H), 7.10-7.04 (m, 4H), 6.98-6.87 (m, 6H), 6.67-6.61 (m, 2H), 6.37-6.31 (m, 4H), 1.62 (s, 6H)	858.34	858.32
45	δ = 8.92 (s, 1H), 8.51 (d, 1H), 8.21 (d, 1H), 8.02 (d, 1H), 7.90 (s, 1H), 7.83-7.80 (m, 1H), 7.75-7.63 (m, 2H), 7.59-7.55 (m, 2H), 7.52-7.48 (m, 5H), 7.42-7.23 (m, 6H), 7.10-7.03 (m, 6H), 6.88-6.85 (m, 2H), 6.70-6.61 (m, 4H), 6.35-6.33 (m, 2H), 6.16-6.13 (m, 4H), 1.63 (s, 6H)	843.37	843.36
48	δ = 8.91 (s, 1H), 8.52 (d, 1H), 8.06-8.02 (m, 2H), 7.90-7.87 (m, 2H), 7.83-7.80 (m, 2H), 7.75-7.71 (m, 2H), 7.68-7.55 (m, 2H), 7.51-7.38 (m, 4H), 7.09-7.03 (m, 6H), 6.90-6.85 (m, 2H), 6.70-6.61 (m, 4H), 6.34-6.32 (m, 2H), 6.15-6.12 (m, 4H), 1.63 (s, 6H)	784.30	784.29
54	δ = 8.92 (s, 1H), 8.52 (d, 1H), 8.02 (d, 1H), 7.90 (s, 1H), 7.82-7.80 (m, 1H), 7.71-7.64 (m, 2H), 7.59-7.56 (m, 3H), 7.51-7.35 (m, 7H), 7.08-7.03 (m, 6H), 6.79-6.78 (m, 1H), 6.68-6.59 (m, 4H), 6.44-6.42 (m, 2H), 6.22-6.16 (m, 6H), 1.63 (s, 6H)	754.34	754.33
57	δ = 8.81 (s, 1H), 8.51 (d, 1H), 8.40 (d, 1H), 7.83-7.80 (m, 2H), 7.68-7.64 (m, 1H), 7.59-7.55 (m, 1H), 7.42 (d, 1H), 7.32-7.15 (m, 11H), 7.09-7.03 (m, 8H), 6.96-6.95 (m, 1H), 6.66-6.61 (m, 4H), 6.52-6.48 (m, 1H), 6.42-6.40 (dd, 1H), 6.21-6.15 (m, 8H)	802.34	802.33
59	δ = 8.82 (s, 1H), 8.51 (d, 1H), 8.39 (d, 1H), 7.83-7.80 (m, 2H), 7.69-7.65 (m, 1H), 7.59-7.55 (m, 1H), 7.43-7.37 (m, 5H), 7.32-7.29 (dd, 1H), 7.26-7.14 (m, 10H), 7.09-7.03 (m, 4H), 6.98-6.97 (m, 1H), 6.66-6.61 (m, 2H), 6.55-6.50 (m, 6H), 6.27-6.20 (m, 4H), 0.24 (s, 18H)	946.42	946.41
62	δ = 8.89 (s, 1H), 8.52 (d, 1H), 8.17 (d, 1H), 7.96-7.94 (m, 1H), 7.90 (s, 1H), 7.83-7.81 (m, 2H), 7.70-7.64 (m, 5H), 7.59-7.55 (m, 1H), 7.49-7.45 (m, 3H), 7.42-7.31 (m, 4H), 7.13-7.12 (m, 1H), 7.06-6.94 (m, 8H), 6.76-6.74 (dd, 1H), 6.69-6.65 (m, 2H), 6.28-6.25 (m, 4H), 2.33-2.24 (m, 2H), 2.12-2.04 (m, 2H), 1.58 (s, 6H)	886.37	886.36

EXAMPLES


Example 1

[0256] An anode was prepared by cutting a Corning 15 Ωcm^2 (1200 Å) ITO glass substrate to a size of 50 mm×50 mm×0.7 mm, ultrasonically cleaning the glass substrate by using (utilizing) isopropyl alcohol and pure water for 5 minutes each, and then irradiating UV light for 30 minutes thereto and exposing the ITO glass substrate to ozone to clean. Then, the anode was loaded into a vacuum deposition apparatus.


[0257] 2-TNATA was deposited on the ITO layer (anode) to form a hole injection layer having a thickness of 600 Å, and then, NPB was deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.

[0258] Subsequently, 9,10-di-naphthalene-2-yl-anthracene (DNA, host) and Compound 1(dopant) were co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.

[0259] Thereafter, Alq3 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, and LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å. Al was deposited on the electron injection layer to form a second electrode (cathode) having a thickness of 3000 Å, thereby completing the manufacture of an organic light-emitting device.




2-TNATA



NPB

-continued



DNA

Example 2

[0260] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound 4 was used (utilized) instead of Compound 1.

Example 3

[0261] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound 15 was used (utilized) instead of Compound 1.

Example 4

[0262] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound 27 was used (utilized) instead of Compound 1.

Example 5

[0263] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound 49 was used (utilized) instead of Compound 1.

Example 6

[0264] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound 57 was used (utilized) instead of Compound 1.


Example 7

[0265] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound 59 was used (utilized) instead of Compound 1.

Comparative Example 1

[0266] An organic light-emitting device was manufactured as in Example 1, except that in forming an emission layer, Compound A illustrated below was used (utilized) instead of Compound 1.

Compound A



[0267] In Examples 1 to 7, compounds, represented by Formulae 1 to 3 were used (utilized) as a dopant material for a blue emission layer. As a result, compared with Comparative Example 1 that included N,N,N',N'-tetraphenyl-pyrene-1,6-diamine (Compound A), which is a material used in organic light-emitting devices, the compounds represented by Formulae 1 to 3 had excellent I-V-L characteristics, for example, low driving voltage and high efficiency, and, for example, high lifespan improvement effects, leading to substantially prolonged lifespan. Representative characteristics and lifespan results were summarized and results thereof are shown in Table 2 below.

TABLE 2

Material	Driving voltage (V)	Current density (mA/cm ²)	Brightness (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr@100 mA/cm ²)	
Example 1	Compound 1	6.11	50	3210	6.42	Blue	380
Example 2	Compound 4	6.18	50	3388	6.78	Blue	358
Example 3	Compound 15	6.17	50	3515	7.03	Blue	362
Example 4	Compound 27	6.22	50	3328	6.66	Blue	351
Example 5	Compound 49	6.23	50	3540	7.08	Blue	345
Example 6	Compound 57	6.09	50	3452	6.90	Blue	354
Example 7	Compound 59	6.32	50	3580	7.16	Blue	325
Comparative Example 1	Compound A	7.01	50	2645	5.29	Blue	258

[0268] When compounds having a structure represented by Formula 1 according to an embodiment of the present disclosure are used (utilized) as a dopant for an emission layer of a blue light-emitting device, compared to known compounds, high efficiency and long lifespan may be obtained.

[0269] An organic light-emitting device including the compound according to an embodiment of the present disclosure may have a low driving voltage, high efficiency, high brightness, and long lifespan.


[0270] It should be understood that the example embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0271] While one or more embodiments of the present invention have been described with reference to the accompanying drawing, it will be understood by those of ordinary skill in the art that various changes in form and details may be made herein without departing from the spirit and scope of the present invention as defined by the following claims, and equivalents thereof.

What is claimed is:

1. A compound represented by Formula 1 below:

Formula 1



wherein, in Formula 1,

R₁ to R₁₄ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group;

Ar₁ to Ar₄ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or

unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), and —B(Q₆)(Q₇);

X and Y are each independently selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic hetero-condensed polycyclic group;

m and n are each independently an integer of 0 to 5; and at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₂-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₂-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic hetero-condensed polycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₂-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic hetero-condensed polycyclic group is selected from:

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy

group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic hetero-condensed polycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and

—N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), wherein:


Q₁ to Q₇, Q₁₁ to Q₁₇, Q₂₁ to Q₂₇ and Q₃₁ to Q₃₇ are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group.

2. The compound of claim 1, wherein:

R₁ to R₁₄ in Formula 1 are each independently selected from a hydrogen, a deuterium, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group.

3. The compound of claim 1, wherein:

R_1 and R_2 in Formula 1 are each independently a substituted or unsubstituted C_1 - C_{20} alkyl group, or a group represented by Formula 2a below:



2a

wherein, in Formula 2a,

Z_1 is selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 to C_{20} alkyl group, a substituted or unsubstituted C_6 to C_{20} aryl group, a substituted or unsubstituted C_1 to C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic heterocondensed polycyclic group;

p is an integer of 1 to 5;

when p is 2 or more, a plurality of Z_1 are identical to, or different from, each other; and

* indicates a binding site to a neighboring atom.

4. The compound of claim 1, wherein:

R_3 to R_{14} in Formula 1 are each independently a hydrogen or a deuterium.

5. The compound of claim 1, wherein:

X and Y in Formula 1 are each independently selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spirofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylene group, an isoxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylenylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylenylene group, a benzothiofenylene group, an isobenzothiazolylenylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothiofenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spirofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an


group, a triazolylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothiofenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spirofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylene group, an isoxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylenylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylenylene group, a benzothiofenylene group, an isobenzothiazolylenylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothiofenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spirofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an


isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

6. The compound of claim 1, wherein:

X and Y in Formula 1 are each independently a group represented by Formulae 3a or 3b below:



3a



3b

wherein, in Formulae 3a and 3b,

Z_1 is selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 to C_{20} alkyl group, a substituted or unsubstituted C_6 to C_{20} aryl group, a substituted or unsubstituted C_1 to C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic heterocondensed polycyclic group;

p is an integer of 1 to 7, and when p is 2 or more, a plurality of Z_1 are identical to, or different from, each other; and * indicates a binding site to a neighboring atom.

7. The compound of claim 1, wherein:

Ar_1 to Ar_4 in Formula 1 are each independently selected from:

a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group,

a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and


a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, C_1 - C_{20} alkoxy group, —Si(Q_3)(Q_4)(Q_5), a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a ben-

zofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a pice-nyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothia-zolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinox-alinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenaziny group, a benzimidazolyl group, a benzofuranyl group, a ben-zothiophenyl group, an isobenzothiazolyl group, a ben-zoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazi-nyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group and a dibenzocarpa-zolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group,


wherein Q_3 to Q_5 are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed poly-cyclic group.

8. The compound of claim 1, wherein:

Ar_1 to Ar_4 in Formula 1 are each independently represented by one of Formulae 4a to 4e below:




4a




4b


-continued



4c



4d



4e

wherein, in Formulae 4a to 4e,

H_1 is $CR_{21}R_{22}$, O, NR_{23} , or S;

R_{21} , R_{22} , R_{23} , and Z_1 are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_6 - C_{20} aryl group, a substituted or unsubstituted C_1 - C_{20} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic hetero-condensed polycyclic group, and —Si(Q_3)(Q_4)(Q_5);

Q_3 to Q_5 are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic hetero-condensed polycyclic group;

p is an integer of 1 to 9, when p is 2 or more, a plurality of Z_1 are identical to, or different from, each other; and

* indicates a binding site to a neighboring atom.

9. The compound of claim 1, wherein:


in Formula 1, Ar_1 or Ar_2 are identical to Ar_3 or Ar_4 .

10. The compound of claim 1, wherein:

m and n in Formula 1 are each independently 0 or 1.

11. The compound of claim 1, wherein:


the compound represented by Formula 1 is a compound represented by Formula 2:



Formula 2

13. The compound of claim 1, wherein:


the compound represented by Formula 1 is a compound represented by Formula 4 below:



Formula 4

12. The compound of claim 1, wherein:


the compound represented by Formula 1 is a compound represented by Formula 3 below:



Formula 3

14. The compound of claim 1, wherein:


the compound of Formula 1 is any one of compounds 1 to 62:



1


-continued

2




-continued


4




3



5




-continued




6

-continued




8


7




9




-continued




-continued




11



13




-continued




14

-continued




16




15

17




-continued




18


-continued



20




19




21

-continued




22


-continued



24




23




25

-continued




26

-continued




28


27



29




-continued




30


-continued



32




31




33

-continued




34


-continued



36




35



37


-continued

38




-continued


40




39




41




-continued




-continued




43



45




-continued




51


-continued



53




52




54

-continued




55

-continued




57




56

58




-continued



59



-continued



60

61

62




15. An organic light-emitting device comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer comprising an emission layer between the first electrode and the second electrode,
 wherein the organic layer comprises at least one of the compound of claim 1.

16. The organic light-emitting device of claim 15, wherein:
 the first electrode is an anode,
 the second electrode is a cathode, and
 the organic layer comprises:
 a hole transport region comprising at least one selected from a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer between the first electrode and the emission layer, and
 an electron transport region comprising at least one selected from a hole blocking layer, an electron trans-

port layer, and an electron injection layer between the emission layer and the second electrode.

17. The organic light-emitting device of claim 16, wherein: the emission layer comprises the compound.

18. The organic light-emitting device of claim 16, wherein: the hole transport region comprises at least one of a compound represented by Formula 201A and a compound represented by Formula 202A:



wherein, in Formulae 201A and 202A,

L_{201} to L_{203} are each independently selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a carbazolylylene group, and a triazinylylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a pyrenylene group, a chrysenylene group, a pyridinylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a carbazolylylene group, and a triazinylylene group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group,

an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group;

x_{a1} to x_{a3} are each independently 0 or 1;

R_{203} , R_{211} , and R_{212} are each independently selected from:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group;

R_{213} and R_{214} are each independently selected from:

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolylyl group, and a triazinylyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

R₂₁₅ and R₂₁₆ are each independently selected from:

a hydrogen atom, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

nyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, and a triazinyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group, each substituted with at least one of a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, and a triazinyl group; and

xa5 is 1 or 2.

19. The organic light-emitting device of claim **15**, wherein the organic layer is formed by utilizing a wet process.

20. A flat panel display apparatus comprising the organic light-emitting device of claim **15**, wherein the first electrode of the organic light-emitting device is configured to be electrically coupled to a source electrode or a drain electrode of a thin film transistor.

* * * * *

专利名称(译)	包括其的化合物和有机发光装置		
公开(公告)号	US20150255721A1	公开(公告)日	2015-09-10
申请号	US14/444881	申请日	2014-07-28
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO., LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO., LTD.		
[标]发明人	LIM JIN O KIM YOUNG KOOK KIM JONG WOO HWANG SEOK HWAN		
发明人	LIM, JIN-O KIM, YOUNG-KOOK KIM, JONG-WOO HWANG, SEOK-HWAN		
IPC分类号	H01L51/00 C07D407/12 C07D307/91 H01L27/32 C07C255/58 C07C211/61 H01L51/50		
CPC分类号	H01L51/006 H01L51/0058 C07D407/12 H01L51/5096 H01L51/5056 H01L51/5072 H01L51/0003 H01L27/3248 C07C255/58 C07D307/91 H01L51/0061 C07C2103/54 H01L51/0056 H01L51/0073 C07C211/61 C07B2200/05 C07C217/92 C07C2603/18 C07C2603/26 C07C2603/54 C07F7/081 H01L51/0081 H01L51/5016		
优先权	1020140027950 2014-03-10 KR		
其他公开文献	US9337431		
外部链接	Espacenet USPTO		

摘要(译)

化合物由式1表示：取代基R 1 至R 14 ， Ar 1 至Ar 4 ， X和Y均为式1这里描述的。有机发光装置包括由式1表示的化合物中的至少一种。包含该化合物的有机发光装置可以具有低驱动电压，高效率，高亮度和长寿命。

