



US 20130113367A1

(19) **United States**(12) **Patent Application Publication**
Jung et al.(10) **Pub. No.: US 2013/0113367 A1**(43) **Pub. Date: May 9, 2013**(54) **HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**(75) Inventors: **Hye-Jin Jung**, Yongin-City (KR);
Seok-Hwan Hwang, Yongin-City (KR);
Young-Kook Kim, Yongin-City (KR);
Jin-O Lim, Yongin-City (KR);
Sang-Hyun Han, Yongin-City (KR);
Soo-Yon Kim, Yongin-City (KR);
Dae-Yup Shin, Yongin-City (KR)(73) Assignee: **Samsung Mobile Display Co., Ltd.**,
Yongin-City (KR)(21) Appl. No.: **13/444,938**(22) Filed: **Apr. 12, 2012**(30) **Foreign Application Priority Data**


Nov. 3, 2011 (KR) 10-2011-0114117

Publication Classification(51) **Int. Cl.**
H05B 33/14 (2006.01)
C07D 401/04 (2006.01)
C07D 403/04 (2006.01)
C07D 413/10 (2006.01)
C07D 401/12 (2006.01)
C07D 405/12 (2006.01)
C07F 7/02 (2006.01)
C07D 409/10 (2006.01)
C07D 209/94 (2006.01)
C07D 401/14 (2006.01)(52) **U.S. Cl.**USPC **313/504**; 548/418; 546/276.7; 544/331;
544/212; 546/256; 548/406; 548/143


(57)

ABSTRACT

A heterocyclic compound represented by Formula 1A and an organic light-emitting device including the heterocyclic compound:



Formula 1A

at least one of R₁ to R₁₃ is a group represented by Formula 1B below:

Formula 1B



wherein R₁ to R₁₅, Ar₁, Ar₂, A, B, a, and b are defined as in the specification. The organic light-emitting device may include an organic layer containing the heterocyclic compound, and thus may have a low driving voltage, a high-emission efficiency, and long lifespan characteristics.

FIG. 1



**HETEROCYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED PATENT
APPLICATION

[0001] This application makes reference to, incorporates the same herein, and claims all benefits accruing under 35 U.S.C. §119 from an application earlier filed in the Korean Intellectual Property Office on 3 Nov. 2011 and there duly assigned Serial No. 10-2011-0114117.

BACKGROUND OF THE INVENTION

[0002] 1. Field of the Invention

[0003] The present invention relates to a heterocyclic compound and an organic light-emitting device including the heterocyclic compound.

[0004] 2. Description of the Related Art

[0005] Organic light-emitting diode (OLEDs), which are self-emitting devices, have advantages such as wide viewing angles, excellent contrast, quick response, high brightness, excellent driving voltage characteristics, and can provide multicolored images.


[0006] A typical OLED has a structure including a substrate, and an anode, a hole transport layer (HTL), an emission layer (EML), an electron transport layer (ETL), and a cathode which are sequentially stacked on the substrate. In this regard, the HTL, the EML, and the ETL are organic thin films formed of organic compounds.

[0007] An operating principle of an OLED having the above-described structure is as follows. When a voltage is applied between the anode and the cathode, holes injected from the anode move to the EML via the HTL, and electrons injected from the cathode move to the EML via the ETL. The holes and electrons recombine in the EML to generate excitons. When the excitons drop from an excited state to a ground state, light is emitted.

SUMMARY

[0008] The present invention provides a novel heterocyclic compound for an organic light-emitting device with a low voltage, a high luminance, a high efficiency, and a long lifespan.


[0009] According to an aspect of the present invention, there is provided a heterocyclic compound represented by Formula 1A below:



Formula 1A

wherein, in Formula 1A, R₁ to R₁₃ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl

group, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₂-C₃₀ alkynyl group, a substituted or unsubstituted C₁-C₃₀ alkoxy group, a substituted or unsubstituted C₃-C₃₀ cycloalkyl group, a substituted or unsubstituted C₃-C₃₀ cycloalkenyl group, a substituted or unsubstituted C₅-C₃₀ aryl group, a substituted or unsubstituted C₂-C₃₀ heteroaryl group, a substituted or unsubstituted C₅-C₃₀ aryloxy group, a substituted or unsubstituted C₅-C₃₀ arylthio group, a —N(Q₁)(Q₂) group, and a group represented by Formula 1B below, wherein Q₁ and Q₂ are each independently one of a hydrogen atom, a deuterium atom, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₂-C₃₀ alkynyl group, a substituted or unsubstituted C₁-C₃₀ alkoxy group, a substituted or unsubstituted C₃-C₃₀ cycloalkyl group, a substituted or unsubstituted C₃-C₃₀ cycloalkenyl group, a substituted or unsubstituted C₆-C₃₀ aryl group, a substituted or unsubstituted C₆-C₃₀ aryloxy group, a substituted or unsubstituted C₆-C₃₀ arylthio group, and a substituted or unsubstituted C₃-C₃₀ heteroaryl group:



Formula 1B

wherein, in Formula 1B, R₁₄ and R₁₅ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₂-C₃₀ alkynyl group, a substituted or unsubstituted C₁-C₃₀ alkoxy group, a substituted or unsubstituted C₃-C₃₀ cycloalkyl group, a substituted or unsubstituted C₃-C₃₀ cycloalkenyl group, a substituted or unsubstituted C₅-C₃₀ aryl group, a substituted or unsubstituted C₂-C₃₀ heteroaryl group, a substituted or unsubstituted C₅-C₃₀ aryloxy group, and a substituted or unsubstituted C₅-C₃₀ arylthio group;

at least one of R₁ to R₁₃ is a group represented by Formula 1B above;

Ar₁ and Ar₂ are each independently a substituted or unsubstituted C₅-C₃₀ aryl group or a substituted or unsubstituted C₃-C₃₀ heteroaryl group, wherein Ar₁ and Ar₂ are optionally linked to each other;

A and B are a divalent linker, and are each independently one of a substituted or unsubstituted C₅-C₃₀ arylene group or a substituted or unsubstituted C₃-C₃₀ heteroarylene group; and a is an integer from 0 to 3, and b is an integer from 0 to 3, wherein if a is 2 or greater, the two or more A are identical to or different from each other, and if b is 2 or greater, the two or more B are identical to or different from each other.

[0010] According to another aspect of the present invention, there is provided an organic light-emitting device including: a first electrode; a second electrode disposed opposite to the first electrode; and an organic layer disposed between the first electrode and the second electrode, wherein the organic layer includes at least one layer, and includes at least one of the heterocyclic compounds represented by Formula 1A above.

BRIEF DESCRIPTION OF THE DRAWING


[0011] The above and other features and advantages of the present invention will become more apparent by describing in detail exemplary embodiments thereof with reference to the attached drawing in which:

[0012] FIG. 1 schematically illustrates the structure of an organic light-emitting device according to an embodiment of the present invention.

DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS


[0013] As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0014] According to an aspect of the present invention, there is provided a heterocyclic compound represented by Formula 1A:



Formula 1A

wherein, in Formula 1A, R_1 to R_{13} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted C_2 - C_{30} alkenyl group, a substituted or unsubstituted C_2 - C_{30} alkynyl group, a substituted or unsubstituted C_1 - C_{30} alkoxy group, a substituted or unsubstituted C_3 - C_{30} cycloalkyl group, a substituted or unsubstituted C_3 - C_{30} cycloalkenyl group, a substituted or unsubstituted C_5 - C_{30} aryl group, a substituted or unsubstituted C_2 - C_{30} heteroaryl group, a substituted or unsubstituted C_5 - C_{30} aryloxy group, a substituted or unsubstituted C_5 - C_{30} arylthio group, a $-N(Q_1)(Q_2)$ group, and a group represented by Formula 1B below, wherein Q_1 and Q_2 are each independently one of a hydrogen atom, a deuterium atom, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted C_2 - C_{30} alkenyl group, a substituted or unsubstituted C_2 - C_{30} alkynyl group, a substituted or unsubstituted C_1 - C_{30} alkoxy group, a substituted or unsubstituted C_3 - C_{30} cycloalkyl group, a substituted or unsubstituted C_3 - C_{30} cycloalkenyl group, a substituted or unsubstituted C_6 - C_{30} aryl group, a substituted or unsubstituted C_6 - C_{30} aryloxy group, a substituted or unsubstituted C_6 - C_{30} arylthio group, and a substituted or unsubstituted C_3 - C_{30} heteroaryl group; at least one of R_1 to R_{13} is a group represented by Formula 1B below;



Formula 1B

wherein, in Formula 1B, R_{14} and R_{15} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted C_2 - C_{30} alkenyl group, a substituted or unsubstituted C_2 - C_{30} alkynyl group, a substituted or unsubstituted C_1 - C_{30} alkoxy group, a substituted or unsubstituted C_3 - C_{30} cycloalkyl group, a substituted or unsubstituted C_3 - C_{30} cycloalkenyl group, a substituted or unsubstituted C_5 - C_{30} aryl group, a substituted or unsubstituted C_2 - C_{30} heteroaryl group, a substituted or unsubstituted C_5 - C_{30} aryloxy group, and a substituted or unsubstituted C_5 - C_{30} arylthio group;

[0015] Ar_1 and Ar_2 are each independently a substituted or unsubstituted C_5 - C_{30} aryl group or a substituted or unsubstituted C_3 - C_{30} heteroaryl group, wherein Ar_1 and Ar_2 may be optionally linked to each other;

[0016] A and B are a divalent linker, and are each independently one of a substituted or unsubstituted C_5 - C_{30} arylene group or a substituted or unsubstituted C_3 - C_{30} heteroarylene group;


[0017] a is an integer from 0 to 3, and b is an integer from 0 to 3, wherein if a is 2 or greater, the two or more A are identical to or different from each other, and if b is 2 or greater, the two or more B are identical to or different from each other; and

[0018] * indicates a binding site.


[0019] For example, Ar_1 and Ar_2 may be each independently one of a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenoxy group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyranil group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted diazinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzimidazolyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted azulenyl group, a substituted or unsubstituted heptalenyl group, a substituted or unsubstituted indacenyl group, a substituted or unsubstituted acenaphthyl group, a substituted or unsubstituted phenalenyl group, a substituted or unsubstituted phenanthridinyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluoranthenyl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted naphthacenyl group, a substituted or unsubstituted picenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted pentaphenyl group, a substituted or unsubstituted hexacenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted imidazopyridinyl group, a substituted or unsubstituted imidazopyrimidinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted imidazopyrimidinyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted indolyl group, a substituted or unsubstituted indoliziny

group, a substituted or unsubstituted isoindolyl group, a substituted or unsubstituted pyridoindolyl group, a substituted or unsubstituted indazolyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted purinyl group, a substituted or unsubstituted benzoquinolinyl group, a substituted or unsubstituted phthalazinyl group, a substituted or unsubstituted naphthyridinyl group, a substituted or unsubstituted quinoxalinyl group, a substituted or unsubstituted quinazoliny group, a substituted or unsubstituted cinnolinyl group, a substituted or unsubstituted phenazinyl group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted thiophenyl group, a substituted or unsubstituted benzothiophenyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted thiazolyl group, a substituted or unsubstituted isothiazolyl group, a substituted or unsubstituted benzothiazolyl group, a substituted or unsubstituted oxazolyl group, a substituted or unsubstituted isoxazolyl group, a substituted or unsubstituted oxadiazolyl group, a substituted or unsubstituted triazolyl group, and a substituted or unsubstituted tetrazolyl group.


[0020] For example, Ar₁ and Ar₂ may be each independently one of the groups represented by Formulae 2A to 2J below, but are not limited thereto:




Formula 2A




Formula 2B




Formula 2C




Formula 2D




Formula 2E




Formula 2F




Formula 2G



Formula 2H




Formula 2I




Formula 2J

[0021] In Formulae 2A to 2J, Z₁₁, Z₁₂, Z₁₃, and Z₁₄ may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyridinyl group, and a substituted or unsubstituted quinolyl group; a plurality of Z₁₁, Z₁₂, Z₁₃, and Z₁₄ may be identical to or different from each other; r is an integer from 1 to 9; s is an integer from 1 to 5; t is an integer from 1 to 4; u is an integer from 1 to 4; and * indicates a binding site.


[0022] For example, Ar₁ and Ar₂ may be each independently one of the groups represented by Formulae 3A to 3S below, but are not limited thereto:




-continued



Formula 3Q



Formula 3R




Formula 3S


[0023] In Formulae 3A to 3S, * indicates a binding site.

[0024] A and B may be each independently one of a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted carbazolyene group, a substituted or unsubstituted dibenzothiophenylene group, a substituted or unsubstituted pyridinylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted spiro-fluorenylene group, a substituted or unsubstituted furanylene group, a substituted or unsubstituted thiophenylene group, and a substituted or unsubstituted oxadiazolylene group.


[0025] For example, A and B may be each independently one of the groups represented by Formulae 4A to 4L below, but are not limited thereto:



Formula 4A




Formula 4B




Formula 4C


-continued




Formula 4D




Formula 4E




Formula 4F




Formula 4G




Formula 4H



Formula 4I



Formula 4J



Formula 4K

Formula 4L


[0026] In Formulae 4A to 4L, Z_{21} , Z_{22} , and Z_{23} may be each independently one of a hydrogen atom, a deuterium atom, a

halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, and a substituted or unsubstituted pyridinyl group, wherein a plurality of Z_{21} , Z_{22} , and Z_{23} may be identical to or different from each other;


v, w, and x are an integer from 1 to 4; and

* and *' indicate binding sites.


[0027] For example, A and B may be each independently one of the groups represented by Formulae 5A to 5R below, but are not limited thereto:




Formula 5A




Formula 5B




Formula 5C



Formula 5D




Formula 5E




Formula 5F


-continued




Formula 5G




Formula 5H




Formula 5I




Formula 5J



Formula 5K




Formula 5L




Formula 5M

-continued


Formula 5N




Formula 5O




Formula 5P



Formula 5Q



Formula 5R




[0028] In Formulae 5A to 5R, * and *' indicate binding sites.

[0029] R_1 to R_{13} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyranyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzoimidazolyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted

imidazolyl group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted pyrazolyl group, a $N(Q_1)(Q_2)$ group, and a group represented by Formula 1B above, wherein Q_1 and Q_2 may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group.

Formula 1B



[0030] In Formula 1B, R_{14} and R_{15} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, and a substituted or unsubstituted pyridinyl group, and a substituted or unsubstituted phenanthrenyl group;

[0031] Ar_1 and Ar_2 may be each independently one of a substituted or unsubstituted phenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted spiro-fluorenyl group, and a substituted or unsubstituted oxadiazolyl group;


[0032] A and B may be each independently one of a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted dibenzothiophenylene group, a substituted or unsubstituted pyridinylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted spiro-fluo-

renylene group, a substituted or unsubstituted furanylene group, a substituted or unsubstituted thiophenylylene group, and a substituted or unsubstituted oxadiazolylylene group; and


a is an integer from 0 to 2, and b is an integer from 0 to 2, wherein if a is 2, the two A are identical to or different from each other, and if b is 2, the two B are identical to or different from each other.

[0033] For example, R_1 to R_{13} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a group represented by Formula 1B above, and groups represented by Formulae 6A to 6G below; and


R_{14} and R_{15} may be each independently one of a hydrogen atom, a deuterium atom, and the groups represented by Formulae 6A to 6L below, but are not limited thereto:




Formula 6A




Formula 6B




Formula 6C



Formula 6D




Formula 6E




Formula 6F


-continued




Formula 6G




Formula 6H




Formula 6I



Formula 6J



Formula 6K



Formula 6L

[0034] In Formulae 6A to 6L, Z_{31} , Z_{32} , and Z_{33} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted quinolyl group, and a $N(Q_{11})(Q_{12})$ group, wherein Q_{11} and Q_{12} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a

hydroxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group;

a plurality of Z_{31} and Z_{32} may be identical to or different from each other;

p may be an integer from 1 to 9;


q may be an integer from 1 to 5; and

* indicates a binding site.


For example, R_1 to R_{13} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted tert-butyl group, a cyano group, $-\text{CD}_3$, $-\text{CF}_3$, a group represented by Formula 1B above, and groups represented by Formulae 7A to 7S below, but are not limited thereto.

[0035] R_{14} and R_{15} may be each independently one of a hydrogen atom, a deuterium atom, and the groups represented by Formulae 7A to 7H below, but are not limited thereto:


-continued



Formula 7F




Formula 7G




Formula 7H


Formula 7A




Formula 7B





Formula 7C




Formula 7D




Formula 7E



Formula 7I




Formula 7J




Formula 7K

-continued



Formula 7L

-continued




Formula 7S


[0036] In Formulae 7A to 7S, * indicates a binding site.

[0037] The heterocyclic compound of Formula 1A above may be a compound represented by Formula 1C, 1D, or 1E below:

Formula 7M





Formula 1C




Formula 1D

Formula 7N





Formula 1E


Formula 7O




Formula 7P



Formula 7Q



Formula 7R



[0038] In Formulae 1C, 1D and 1E, Ar₁ and Ar₂ may be each independently one of a substituted or unsubstituted phenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted spiro-fluorenyl group, and a substituted or unsubstituted oxadiazolyl group;

A and B may be each independently one of a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted dibenzothiophenylene group, a substituted or unsubstituted pyridinylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted perylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted spiro-fluorenylene group, a substituted or unsubstituted furanylene group, a substituted or unsubstituted thiophenylene group, and a substituted or unsubstituted oxadiazolylene group;

[0039] a is an integer from 0 to 2, and b is an integer from 0 to 2, wherein if a is 2, the two A are identical to or different from each other, and if b is 2, the two B are identical to or different from each other;

[0040] R_1 to R_{13} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyranlyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzimidazolyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted pyrazolyl group, and a $N(Q_1)(Q_2)$ group, wherein Q_1 and Q_2 may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or

unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group; and



[0041] R_{14} and R_{15} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, and a substituted or unsubstituted pyridinyl group, and a substituted or unsubstituted phenanthrenyl group.

[0042] In some embodiments, in Formulae 1C, 1D and 1E, Ar_1 and Ar_2 may be each independently one of the groups represented by Formulae 2A to 2J above. In some other embodiments, in Formulae 1C, 1D and 1E, Ar_1 and Ar_2 may be each independently one of the groups represented by Formulae 3A to 3S above.











[0043] In still other embodiments, in Formulae 1C, 1D and 1E, A and B may be each independently one of the groups represented by Formulae 4A to 4L above. In yet still other embodiments, in Formulae 1C, 1D and 1E, A and B may be each independently one of the groups represented by Formulae 5A to 5R above.

[0044] In some embodiments, in Formulae 1C, 1D and 1E, R_1 to R_{13} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, and the groups represented by Formulae 6A to 6L above; and R_{14} and R_{15} may be each independently one of a hydrogen atom, a deuterium atom, and the groups represented by Formulae 6A to 6L above. In some other embodiments, in Formulae 1C, 1D and 1E, R_1 to R_{13} may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted tert-butyl group, a cyano group, $-CD_3$, $-CF_3$, and groups represented by Formulae 7A to 7S below; and R_{14} and R_{15} may be each independently one of a hydrogen atom, a deuterium atom, and the groups represented by Formulae 7A to 7H above.



[0045] The heterocyclic compound represented by Formula 1A above may be one of the Compounds 1 to 75 below, but is not limited thereto:




-continued




-continued
13




14




15




16




17




18




19



20




21




22


-continued
23




24




25




26




27




28




29




30




31




32




-continued




33




34




35




36




37



38




39




40

-continued


47




48




49



50




51




-continued

52





53




54


55



56




57




-continued


58




59







60



61




-continued




-continued


66




67




68




69



70




71




-continued


72




73



74



75



[0046] The heterocyclic compound of Formula 1A above may be used as an emitting material, a hole injecting material, and/or a hole transporting material of an organic light-emitting device. The heterocyclic compound of Formula 1 above, which has a heterocyclic group in the molecules thereof, has a high glass transition temperature (T_g) or a high melting point due to the inclusion of the heterocyclic group. Thus, the heterocyclic compound has high heat resistance against Joule's heat generated in an organic layer, between organic layers, or between an organic layer and a metal electrode when light emission occurs, and has high durability in high-temperature environments. When a substituent such as a fluorenyl group is introduced, the heterocyclic compound of Formula 1A above may improve morphology of the organic layer, and thus improving characteristics of an organic light-emitting device. In the compound of Formula 1A indenocarbazole moieties with fused aromatic rings possess abundant π -electrons. This is advantageous for $\pi \rightarrow \pi^*$ transition, and appropriate arrangement of orbitals involved in light emission may lead to high light-emission efficiency. Double bonds are linked to indenocarbazole in a smooth resonance form, thus facilitating delocalization of π -electrons. This may also improve emission efficiency. Introduction of arylamine groups that may lead to a high light-emission efficiency via $n \rightarrow \pi^*$ transition in the compound of Formula 1A may further improve the emission efficiency of the compound of Formula 1A.

[0047] As used herein, the term "substituted A" of the "substituted or unsubstituted A (wherein A is a certain substituent)" refers to a group A of which at least one hydrogen atom is substituted with one of a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, an amino group, hydrazine, hydrazone, a carboxyl group or a salt derivative thereof, a sulfonic acid group or a salt derivative thereof, a phosphoric acid group or a salt derivative thereof, a C₁-C₃₀ alkyl group, a C₂-C₃₀ alkenyl group, a C₂-C₃₀ alkynyl group, a C₁-C₃₀ alkoxy group, a C₃-C₃₀ cycloalkyl group, a C₃-C₃₀ cycloalkenyl group, a C₅-C₃₀ aryl group, a C₅-C₃₀ aryloxy group, a C₅-C₃₀ arylthio group, a C₃-C₃₀ heteroaryl group, a group represented by N(Q₁₀₁)(Q₁₀₂), and a group represented by Si(Q₁₀₃)(Q₁₀₄)(Q₁₀₅), wherein Q₁₀₁ to Q₁₀₅ may be each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, an amino group, a nitro group, a carboxyl group, a C₁-C₃₀ alkyl group, a C₂-C₃₀ alkenyl group, a C₂-C₃₀ alkynyl group, a C₁-C₃₀ alkoxy group, a C₃-C₃₀ cycloalkyl group, a C₃-C₃₀ cycloalkenyl group, a C₅-C₃₀ aryl group, a C₅-C₃₀ aryloxy group, a C₅-C₃₀ arylthio group, and a C₂₀-C₃₀ heteroaryl group.

[0048] For example, the term "substituted A" may refer to a group A of which at least one hydrogen atom is substituted with one of a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a methoxy group, an ethoxy group, a phenyl group, a biphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spirofluorenyl group, a phenalenyl group, a phenanthrenyl group, a phenanthridinyl group, a phenanthrolinyl group, an anthryl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrycenyl group, a naphthacenylyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pyrrolyl group, an imidazolyl

group, a benzoimidazolyl group, a pyrazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, an imidazopyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, a pyridoindolyl group, an indazolyl group, a furinyl group, a quinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a phthalazinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenazinyl group, a furanyl group, a benzofuranyl group, a dibenzofuranyl group, a thiophenyl group, a benzothiophenyl group, a dibenzothiophenyl group, a thiazolyl group, an isothiazolyl group, a benzothiazolyl group, an oxazolyl group, a benzooxazolyl group, an isoxazolyl group, an oxadiazolyl group, a triazolyl group, a triazinyl group, a tetrazolyl group, a group represented by N(Q₁₀₁)(Q₁₀₂), and a group represented by Si(Q₁₀₃)(Q₁₀₄)(Q₁₀₅).

[0049] As used herein, the unsubstituted C₁-C₃₀ alkyl group refers to a linear or branched saturated hydrocarbon that lacks one hydrogen atom from alkane. Non-limiting examples of the unsubstituted C₁-C₃₀ alkyl group include methyl, ethyl, propyl, isobutyl, sec-butyl, pentyl, isoamyl, and hexyl. Substituents of the substituted C₁-C₃₀ alkyl group are the same as described above in conjunction with the "substituted A".

[0050] As used herein, the unsubstituted C₂-C₃₀ alkenyl group indicates a hydrocarbon chain having at least one carbon-carbon double bond in the center or at a terminal of the unsubstituted C₂-C₃₀ alkyl group. Non-limiting examples of the unsubstituted C₂-C₃₀ alkenyl group include ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, propadienyl, isoprenyl, and allyl. Substituents of the substituted C₂-C₃₀ alkenyl group are the same as described above in conjunction with the "substituted A".

[0051] As used herein, the unsubstituted C₂-C₃₀ alkynyl group indicates a hydrocarbon chain having at least one carbon-carbon triple bond in the center or at a terminal of the unsubstituted C₂-C₆₀ alkyl group. Non-limiting examples of the unsubstituted C₂-C₃₀ alkynyl group include ethynyl, propynyl, and acetylenyl. Substituents of the substituted C₂-C₃₀ alkynyl group are the same as described above in conjunction with the "substituted A".

[0052] As used herein, the unsubstituted C₁-C₃₀ alkoxy group refers to a group represented by —OY, wherein Y is an unsubstituted C₁-C₃₀ alkyl group as described above. Non-limiting examples of the unsubstituted C₁-C₃₀ alkoxy group are a methoxy group, an ethoxy group, an isopropoxy group, a butoxy group, and a pentoxy group. Substituents of the substituted C₁-C₃₀ alkoxy group are the same as described above in conjunction with the "substituted A".

[0053] As used herein, the unsubstituted C₃-C₃₀ cycloalkyl group indicates a cyclic saturated hydrocarbon group. Non-limiting examples of the unsubstituted C₃-C₃₀ cycloalkyl group are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cyclooctyl. Substituents of the substituted C₃-C₃₀ cycloalkyl group are the same as described above in conjunction with the "substituted A".

[0054] As used herein, the unsubstituted C₃-C₃₀ cycloalkenyl group refers to a non-aromatic, cyclic unsaturated hydrocarbon group with at least one carbon-carbon double bond. Non-limiting examples of the unsubstituted C₃-C₃₀ cycloalkenyl group are cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, 1,3-cyclohexadienyl, 1,4-cyclohexadienyl, 2,4-cycloheptadienyl, and 1,5-cyclooctadi-

enylactenyl. Substituents of the substituted C_5-C_{60} cycloalkenyl group are the same as described above in conjunction with the "substituted A".

[0055] As used herein, the unsubstituted C_5-C_{30} aryl group indicates a monovalent group including a C_5-C_{30} carbocyclic aromatic system, which may be monocyclic or polycyclic. In a polycyclic group, at least two rings may be fused to each other. Non-limiting examples of the unsubstituted C_5-C_{30} aryl group are phenyl, pentalenyl, indenyl, naphthyl, azulenyl, heptalenyl, indacenyl, acenaphthyl, fluorenyl, spiro-fluorenyl, phenalenyl, phenanthrenyl, anthryl, fluoranthenyl, triphenylenyl, pyrenyl, chrysenyl, naphthacenyl, picenyl, perylenyl, pentaphenyl, and hexacacenyl. Substituents of the substituted C_5-C_{30} aryl group are the same as described above in conjunction with the "substituted A".

[0056] As used herein, the unsubstituted C_5-C_{30} aryloxy group refers to a monovalent group with a carbon atom of the C_5-C_{30} aryl group attached via an oxygen linker ($-O-$). Substituents of the substituted C_5-C_{30} aryloxy group are the same as described above in conjunction with the "substituted A".

[0057] As used herein, the unsubstituted C_5-C_{30} arylthio group refers to a monovalent group with a carbon atom of the C_5-C_{30} aryl group attached via a sulfur linker ($-S-$). Non-limiting examples of the unsubstituted C_5-C_{30} arylthio group are a phenylthio group, a naphthylthio group, an indanylthio group, and an indenylthio group. Substituents of the substituted C_5-C_{30} arylthio group are the same as described above in conjunction with the "substituted A".

[0058] As used herein, the unsubstituted C_3-C_{30} heteroaryl group indicates a monovalent group with at least one ring including at least one heteroatom selected from among N, O, P, and S, which may be a monocyclic or polycyclic group. In a polycyclic group, at least two rings may be fused to each other. Non-limiting examples of the unsubstituted C_3-C_{30} heteroaryl group are pyrrolyl, imidazolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolinyl, benzoquinolinyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, carbazolyl, phenanthridinyl, acridinyl, phenanthrolinyl, phenazinyl, benzooxazolyl, benzoimidazolyl, furanyl, benzofuranyl, thiophenyl, benzothiophenyl, thiazolyl, isothiazolyl, benzothiazolyl, isoxazolyl, oxazolyl, triazolyl, tetrazolyl, oxadiazolyl, triazinyl, and benzooxazolyl. Substituents of the substituted C_3-C_{30} heteroaryl group are the same as described above in conjunction with the "substituted A".

[0059] In the specification, the unsubstituted C_1-C_{30} alkylene group refers to a linear or branched divalent group that lacks two hydrogen atoms from the unsubstituted C_1-C_{30} alkylene group. Examples of the unsubstituted C_1-C_{30} alkylene group may be inferred based on those of the unsubstituted C_1-C_{30} alkyl group described above. Substituents of the substituted C_1-C_{30} alkylene group are the same as those described above in conjunction with the "substituted A".

[0060] As used herein, the unsubstituted C_5-C_{30} arylene group indicates a divalent group including a C_5-C_{30} carbocyclic aromatic system, which may be monocyclic or polycyclic. Examples of the unsubstituted C_5-C_{30} arylene group may be inferred based on those of the unsubstituted C_5-C_{30} alkyl group described above. Substituents of the substituted C_5-C_{30} arylene group are the same as those described above in conjunction with the "substituted A".

[0061] The heterocyclic compound of Formula 1A may be synthesized using a known organic synthesis method. A syn-

thesis method of the heterocyclic compound of Formula 1A may be understood by one of ordinary skill in the art from the examples that will be described below.

[0062] The heterocyclic compound of Formula 1A above may be used in an organic light-emitting device.

[0063] According to another aspect of the present invention, an organic light-emitting device includes: a first electrode; a second electrode disposed opposite to the first electrode; and an organic layer disposed between the first electrode and the second electrode. The organic layer, including at least one layer, may contain at least one of the heterocyclic compounds of Formula 1A described above.

[0064] As used herein, the term "organic layer" refers to a layer containing an organic compound and including at least one layer. For example, the organic layer may include at least one layer selected from among a hole injection layer, a hole transport layer, a hole injection and transport layer having both hole injection and hole transport capabilities, an electron blocking layer, an emission layer, a hole blocking layer, an electron injection layer, an electron transport layer, and an electron injection and transport layer having both electron injection and electron transport capabilities. The organic layer may further include an inorganic compound or an inorganic material such as an organometallic complex. In one embodiment, the organic layer may include both an organic compound and an inorganic compound or may include one layer including an organic compound and another layer including an inorganic compound or material. For example, the organic layer may include both an organic compound and an organometallic complex in one layer. In another embodiment, the organic layer may include a layer containing an organic compound and a layer containing an inorganic compound or an inorganic material.

[0065] At least one of the heterocyclic compounds listed above may be included in one layer of the organic layer, and in some other embodiments, at least one of the heterocyclic compounds listed above may be included in different layers of the organic layer. For example, the organic layer may include one of the heterocyclic compounds as an emitting dopant in an emission layer, and another heterocyclic compound as a hole transport material in a hole transport layer. In another embodiment, the organic layer may include one of the heterocyclic compounds as an emitting dopant and another heterocyclic compound as an emitting host in an emission layer. In another embodiment, the organic layer may include one of the heterocyclic compounds as an emitting dopant and another heterocyclic compound as an emitting host in an emission layer, and still another heterocyclic compound as a hole transport material in a hole transport layer.

[0066] The organic layer may include at least one of an emission layer, a hole injection layer, a hole transport layer, and a hole injection and transport layer having both hole injection and hole transport capabilities, and at least one of the emission layer, the hole injection layer, the hole transport layer, and the hole injection and transport layer may include the heterocyclic compound.

[0067] In some embodiments, the organic light-emitting device may have a first electrode/hole injection layer/hole transport layer/emission layer/electron transport layer/second electrode structure, wherein the emission layer, the hole transport layer, or the hole injection layer may include the heterocyclic compound. In some embodiments, at least two of the emission layer, the hole transport layer, and the hole injection layer may include the heterocyclic compound. In

these embodiments, each of the at least two layers may include a different heterocyclic compound. As described above, a layer of the organic layer may include a mixture of at least two of the heterocyclic compounds listed above, or a mixture of one of the heterocyclic compounds and a non-heterocyclic compound.

[0068] In some embodiments, the organic layer may include an emission layer, which may include a host and a dopant, and the heterocyclic compound may be a fluorescent host, a phosphorescent host, or a fluorescent dopant of the emission layer.

[0069] In some embodiments, the organic layer may include an emission layer, which may further include an anthracene compound, an arylamine compound, or a styryl compound. The emission layer may or may not include the heterocyclic compound.

[0070] The organic layer may include an emission layer, which may include a host and a dopant. The emission layer may further include a phosphorescent dopant. For example, the phosphorescent dopant may be, but is not limited to, an organometallic complex including at least one of iridium (Ir), platinum (Pt), osmium (Os), rhenium (Re), titanium (Ti), zirconium (Zr), hafnium (Hf), and a combination of at least two thereof. The emission layer may or may not include the heterocyclic compound.

[0071] At least one of the hole injection layer, the hole transport layer, and the hole injection and transport layer may further include a charge generating material, in addition to the heterocyclic compound. The charge generating material may be, for example, a p-dopant. The hole injection layer, the hole transport layer, and the hole injection and transport layer may or may not include the heterocyclic compound.

[0072] The organic layer may further include an electron transport layer, which may include an electron transporting organic compound and a metal-containing material. The metal-containing material may include a lithium (Li) complex. The electron transport layer may or may not include the heterocyclic compound.

[0073] At least one organic layer disposed between the first electrode and the second electrode may be formed using deposition or a wet process.

[0074] As used herein, the terms "wet process" refers to a process involving applying a mixture of a specific material and a solvent to a predetermined substrate, and drying and/or thermally treating to remove at least part of the solvent, thereby forming a layer including the specific material on the substrate.

[0075] For example, the organic layer may be formed using a general vacuum deposition method. In some other embodiments, the organic layer may be formed by applying the mixture of the heterocyclic compound and the solvent to a region to form the organic layer (for example, on the hole transport layer) using spin coating, spraying, inkjet printing, dipping, casting, gravure coating, bar coating, roll coating, wire bar coating, screen coating, flexo coating, offset coating, laser transferring, or the like, and drying and/or thermally treating the mixture coating the region to form the organic layer to remove at least part of the solvent.

[0076] In another embodiment, the organic layer may be formed using a laser transfer method by which an organic layer is formed on a base film using vacuum deposition or a wet process described above, and then transferred to a region

to form the organic layer of an organic light-emitting device (for example, on the hole transport layer of the organic light-emitting device) using laser.

[0077] FIG. 1 is a schematic sectional view of an organic light-emitting device 10 according to an embodiment of the present invention. Hereinafter, a structure of an organic light-emitting device according to an embodiment of the present invention and a method of manufacturing the same will now be described with reference to FIG. 1.

[0078] Referring to FIG. 1, the organic light-emitting device 10 according to the present embodiment includes a first electrode 13, an organic layer 15, and a second electrode 17, which are sequentially stacked on a substrate 11 in this order.

[0079] The substrate 11 may be any substrate that is used in conventional organic light-emitting devices. In some embodiments the substrate 11 may be a glass substrate or a transparent plastic substrate with strong mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0080] The first electrode 13 may be formed by depositing or sputtering a first electrode-forming material on the substrate 11. When the first electrode 13 constitutes an anode, a material having a high work function may be used as the first electrode-forming material to facilitate hole injection. The first electrode 13 may be a reflective electrode or a transmission electrode. Suitable first electrode-forming materials are transparent and conductive materials such as ITO, IZO, SnO₂, and ZnO. The first electrode 13 may be formed as a reflective electrode using magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), or the like.

[0081] The organic layer 15 may be disposed on the first electrode 13. As described above, the organic layer 15 indicates any layer interposed between the first electrode 13 and the second electrode 17. The organic layer 15 may further include an inorganic compound or material such as a metal complex.

[0082] The organic layer 15 may include a hole injection layer (HIL), a hole transport layer (HTL), an emission layer (EML), an electron transport layer (ETL), and an electron injection layer (EIL).


[0083] The HIL may be formed on the first electrode 13 by vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like.

[0084] When the HIL is formed using vacuum deposition, vacuum deposition conditions may vary according to the compound that is used to form the HIL, and the desired structure and thermal properties of the HIL to be formed. For example, vacuum deposition may be performed at a temperature of about 100° C. to about 500° C., a pressure of about 10⁻⁸ torr to about 10⁻³ torr, and a deposition rate of about 0.01 to about 100 Å/sec. However, the deposition conditions are not limited thereto.


[0085] When the HIL is formed using a wet process such as spin coating, the coating conditions may vary according to the compound that is used to form the HIL, and the desired structure and thermal properties of the HIL to be formed. For example, the coating rate may be from about 2,000 rpm to about 5,000 rpm, and a temperature at which heat treatment is performed to remove a solvent after coating may be from about 80° C. to about 200° C. However, the coating conditions are not limited thereto.

[0086] For example, as a HIL material, the heterocyclic compound of Formula 1A or any known hole injection materials may be used. Non-limiting examples of known hole injection materials include N,N'-diphenyl-N,N'-bis-[4-(phenyl-m-tolyl-amino)-phenyl]-biphenyl-4,4'-diamine, (DNTPD), a phthalocyanine compound such as copperphthalocyanine, 4,4',4''-tris(3-methylphenylphenylamino)triphenylamine (m-MTDATA), N,N'-di(1-naphthyl)-N,N'-diphenylbenzidine (NPB), 4,4',4''-Tris(N,N-diphenylamino)triphenylamine (TDATA), 4,4',4''-tris {N,-(2-naphthyl)-N-phenylamino}-triphenylamine (2T-NATA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), and polyaniline/poly(4-styrenesulfonate) (PANI/PSS).

m-MTDATA




TDATA






-continued

2T-NATA



PEDOT/PSS






[0087] The thickness of the HIL may be about 100 Å to about 10,000 Å, and in some embodiments, may be about 100 Å to about 1,000 Å. When the thickness of the HIL is within these ranges, the HIL may have improved hole injecting ability without a substantial increase in driving voltage.

[0088] Then, a HTL may be formed on the HIL by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like. When the HTL is formed using vacuum deposition or spin coating, the conditions for deposition and coating may be similar to those for the formation of the HIL, though the conditions for the deposition and coating may vary according to the material that is used to form the HTL.

[0089] As a HTL material, the heterocyclic compound of Formula 1A or any known hole transporting materials may be used. Examples of known hole transporting materials include carbazole derivatives such as N-phenylcarbazole, polyvinylcarbazole, and the like; triphenylamine materials such as TPD (N,N'-bis(3-methylphenyl)-N,N'-diphenyl-[1,1'-biphenyl]-4,4'-diamine); and amine derivatives having an aromatic condensed ring, such as α -NPD (4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl, TCTA (4,4',4''-tris(N-carbazolyl)triphenylamine), and the like.




[0090] The thickness of the HTL may be about 50 Å to about 1,000 Å, and in some embodiments, may be about 100 Å to about 800 Å. When the thickness of the HTL is within these ranges, the HTL may have improved hole transporting ability without a substantial increase in driving voltage.

[0091] In some embodiments, instead of the HIL and the HTL, a hole injection and transport layer having both hole injection and hole transport capabilities may be formed. As a material for the hole injection and transport layer, the heterocyclic compound of Formula 1 or Formula 2, or any known materials may be used.

[0092] At least one of the hole injection layer, the hole transport layer, and the hole injection and transport layer may further include a charge generating material for improved layer conductivity, in addition to a known hole injection material and a known hole transport material.

[0093] The charge generating material may be, for example, a p-dopant. Non-limiting examples of the p-dopant include quinone derivatives such as tetracyanoquinodimethane (TCNQ), 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinodimethane (F4TCNQ), and the like; metal oxides such as tungsten oxide, molybdenum oxide, and the like; and cyano-containing compounds such as Compound 100 below.




TCNQ


F4TCNQ

Compound 100


zene), TBADN ((3-tert-butyl-9,10-di(naphth-2-yl)anthracene), DSA (distyrylarylene), and E3.




CBP




PVK




DSA



AND



TPBI



TBADN


[0094] When the hole injection layer, the hole transport layer, or the hole injection and transport layer having both hole injection and hole transport capabilities further includes a charge generating material, the charge generating material may be, but is not limited to, homogeneous dispersed or inhomogeneously distributed in the layer.

[0095] Then, an EML may be formed on the HTL or the hole injection and transport layer having both hole injection and hole transport capabilities by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like. When the EML is formed using vacuum deposition or spin coating, the deposition and coating conditions may be similar to those for the formation of the HIL, though the conditions for deposition and coating may vary according to the material that is used to form the EML.

[0096] As an EML material, at least one of the heterocyclic compound of Formula 1A, and known light-emitting materials (including hosts and dopants) may be used. When including the heterocyclic compound of Formula 1A, the EML may further include a known phosphorescent host, fluorescent host, phosphorescent dopant, or fluorescent dopant. The heterocyclic compound may also serve as a phosphorescent host, a fluorescent host, or a fluorescent dopant.

[0097] The heterocyclic compound of Formula 1A may be used as a host. In another embodiment a known dopant may be used. Non-limiting examples of known hosts include Alq3, CBP (4,4'-N,N'-dicabazole-biphenyl), PVK (poly(n-vinylcabazole), ADN (9,10-di(naphthalene-2-yl)anthracene), TCTA, TPBI ((1,3,5-tris(N-phenylbenzimidazole-2-yl)ben-


-continued





E3

[0098] The heterocyclic compound of Formula 1A may be used as a dopant. In another embodiment a known dopant may be used. For example, at least one of a fluorescent dopant and a phosphorescent dopant may be used. For example, the phosphorescent dopant may include, but is not limited to, an organometallic complex including at least one selected from the group consisting of iridium (Ir), platinum (Pt), osmium (Os), rhenium (Re), titanium (Ti), zirconium (Zr), hafnium (Hf), and a combination of at least two thereof.





[0099] Non-limiting examples of known red dopants include PtOEP (Pt(II) octaethylporphine), Ir(piq)₃ (tris(2-phenylisoquinoline)iridium), and Btp₂Ir(acac) (bis(2-(2'-benzothienyl)-pyridinato-N,C3')iridium(acetylacetonate)).



PtOEP


Ir(piq)₃Btp₂Ir(acac)

[0100] Non-limiting examples of known green dopants include Ir(ppy)₃ (tris(2-phenylpyridine) iridium), Ir(ppy)₂(acac) (bis(2-phenylpyridine)(acetylacetonato)iridium(III)), Ir(mppy)₃ (tris(2-(4-tolyl)phenylpyridine)iridium), and C545T (10-(2-benzothiazolyl)-1,1,7,7-tetramethyl-2,3,6,7-tetrahydro-1H,5H,11H-[1]benzopyrano[6,7,8-ij]-quinoxalin-11-one).

Ir(ppy)₃Ir(mppy)₃Ir(ppy)₂(acac)

C545T

[0101] Non-limiting examples of known blue dopants include F₂Irpic (bis[3,5-difluoro-2-(2-pyridyl)phenyl](picolinato)iridium(III)), (F₂ppy)₂Ir(tmd), Ir(dfppz)₃, DPVBi (4,4'-bis(2,2'-diphenylethen-1-yl)biphenyl), DPAVBi (4,4'-bis[4-(diphenylamino)styryl]biphenyl), and TBPe (2,5,8,11-tetra-tert-butyl perylene).



[0102] When the EML includes both a host and a dopant, the amount of the dopant may be from about 0.01 to about 15 parts by weight based on 100 parts by weight of the host. However, the amount of the dopant is not limited to this range.

[0103] The thickness of the EML may be from about 100 Å to about 1,000 Å, and in some embodiments, may be from about 200 Å to about 600 Å. When the thickness of the EML is within these ranges, the EML may have improved light emitting ability without a substantial increase in driving voltage.

[0104] When a phosphorescent dopant is also used to form the EML, a HBL may be formed between the ETL and the EML by using vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, or the like, in order to prevent diffusion of triplet excitons or holes into an ETL. When the HBL is formed using vacuum deposition or spin coating, the conditions for deposition and coating may be similar to those for the formation of the HIL, although the conditions for deposition and coating may vary according to the material that is used to form the HBL. Any known hole-blocking material may be used. Non-limiting examples of hole-blocking materials include an oxadiazole derivative, a


triazole derivative, and a phenanthroline derivative. For example, BCP may be used as a material for forming the HBL.

[0105] The thickness of the HBL may be from about 50 Å to about 1,000 Å, and in some embodiments, may be from about 100 Å to about 300 Å. When the thickness of the HBL is within these ranges, the HBL may have improved hole blocking ability without a substantial increase in driving voltage.


[0106] Then, an ETL may be formed on the HBL or EML by vacuum deposition, spin coating, casting, or the like. When the ETL is formed using vacuum deposition or spin coating, the deposition and coating conditions may be similar to those for the formation of the HIL, though the deposition and coating conditions may vary according to the material that is used to form the ETL.

[0107] The ETL may be formed of any known hole transporting material. Non-limiting examples of known ETL materials include quinoline derivatives, and in particular, Alq₃ (tris(8-quinolinolate)aluminum), BCP (2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline), Bphen (4,7-diphenyl-1,10-phenanthroline; 4,7-diphenyl-1,10-phenanthroline), TAZ


(3-(4-biphenyl)-4-phenyl-5-tert-butylphenyl-1,2,4-triazole), NTAZ (4-(Naphthalen-1-yl)-3,5-diphenyl-4H-1,2,4-triazole), tBu-PBD (2-(4-biphenyl)-5-(4-tert-butylphenyl)-1,3,4-oxadiazole), BAiq (see the following formula), Bebq₂ (beryllium bis(benzoquinolin-10-olate), AND (9,10-di(naphthalene-2-yl)anthracene), Compound 101, and Compound 102.




BAiq




BCP



NTAZ




Bphen




TAZ

-continued




tBu-PBD

Compound 101




Compound 102



[0108] The thickness of the ETL may be from about 100 Å to about 1,000 Å, and in some other embodiments, may be from about 150 Å to about 500 Å. When the thickness of the ETL is within these ranges, the ETL may have satisfactory electron transporting ability without a substantial increase in driving voltage.

[0109] In some embodiments the ETL may include an electron-transporting organic compound and a metal-containing material. The metal-containing material may include a lithium (Li) complex. Non-limiting examples of the Li complex include lithium quinolate (LiQ) and Compound 103 below:



Compound 103

[0110] Then, an EIL, which facilitates injection of electrons from the cathode, may be formed on the ETL. Any suitable electron-injecting material may be used to form the EIL.

[0111] Examples of materials for forming the EIL include LiF, NaCl, CsF, Li₂O, and BaO, which are known in the art. The deposition and coating conditions for forming the EIL may be similar to those for the formation of the HIL, though the deposition and coating conditions may vary according to the material that is used to form the EIL.

[0112] The thickness of the EIL may be about 1 Å to about 100 Å, and in some embodiments, may be from about 3 Å to about 90 Å. When the thickness of the EIL is within these ranges, the EIL may have satisfactory electron injection ability without a substantial increase in driving voltage.




[0113] The second electrode 17 is disposed on the organic layer 15. The second electrode 17 may be a cathode that is an electron injection electrode. A material for forming the second electrode 17 may be a metal, an alloy, an electro-conductive compound, which have a low work function, or a mixture thereof. In this regard, the second electrode 17 may be formed of lithium (Li), magnesium (Mg), aluminum (Al), aluminum (Al)-lithium (Li), calcium (Ca), magnesium (Mg)-indium (In), magnesium (Mg)-silver (Ag), or the like, and may be formed as a thin film type transmission electrode. To manufacture a top-emission light-emitting device, the transmission electrode may be formed of indium tin oxide (ITO) or indium zinc oxide (IZO).


[0114] Hereinafter, the present invention will be described in detail with reference to the following synthesis examples and other examples. However, these examples are for illustrative purposes only and are not intended to limit the scope of the present invention.

Synthesis Example 1

Synthesis of Compound 8

[0115]





Synthesis of Intermediate I-1

[0116] 6.34 g (20.0 mmol) of 2-bromo-9,9-dimethyl-7-fluoreneboronic acid, 4.04 g (20.0 mmol) of 2-bromonitrobenzene, 1.15 g (1.0 mmol) of Pd(PPh₃)₄ (tetrakis(triphenylphosphine)palladium), and 8.29 g (60.0 mmol) of K₂CO₃ were dissolved in 60 mL of a mixed tetrahydrofuran (THF) and H₂O (2:1 by volume) solution to obtain a solution, which was then stirred at about 70° C. for about 5 hours. The reaction solution was cooled to room temperature, and 40 mL of water was added thereto, followed by three times of extraction with 50 mL of ethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 6.86 g of Intermediate I-1 (Yield: 87%). This compound was identified using mass spectroscopy/fast atom bombardment (MS/FAB). C₂₁H₁₆BrNO₂; calc. 393.04. found 393.14

Synthesis of Intermediate I-2

[0117] 3.94 g (10.0 mmol) of Intermediate I-1 and 5.77 g (22 mmol) of triphenylphosphine (PPh₃) were dissolved in 30 mL of 1,2-dichlorobenzene to obtain a solution, which was then stirred at about 220° C. for about 12 hours. The reaction solution was cooled to room temperature, and the solvent was removed therefrom under vacuum conditions, followed by three times of extraction with 50 mL of water and 50 mL of dichloromethane. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 2.83 g of Intermediate I-2 (Yield: 78%). This compound was identified using MS/FAB. C₂₁H₁₆BrN; calc. 361.05. found 361.26

Synthesis of Intermediate I-3

[0118] 3.62 g (10.0 mmol) of Intermediate I-2, 4.80 g (15.0 mmol) of 9,9-dimethyl-2-iodofluorene, 0.19 g (1.0 mmol) of CuI, 0.05 g (0.2 mmol) of 18-Crown-6, and 4.15 g (30.0

mmol) of K₂CO₃ were dissolved in 30 mL of 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU) to obtain a solution, which was then stirred at about 170° C. for about 12 hours. The reaction solution was cooled to room temperature, followed by three times of extraction with 50 mL of water and 50 mL of dichloromethane. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 4.03 g of Intermediate I-3 (Yield: 92%). This compound was identified using MS/FAB. C₃₆H₂₈BrN; calc. 553.14. found 553.25

Synthesis of Intermediate I-4

[0119] 8.461 g (50 mmol) of diphenylamine, 10.983 g (60 mmol) of bromostyrene (Compound E), 0.915 g (1 mmol) of Pd₂(dba)₃, 0.202 g (1 mmol) of PtBu₃, and 69.611 g (100 mmol) of KOtBu were dissolved in 300 mL of toluene to obtain a solution, which was then stirred at about 85° C. for about 4 hours. The reaction solution was cooled to room temperature, followed by three times of extraction with 100 mL of water and 100 mL of diethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 7.33 g of Intermediate I-4 (Yield: 54%). This compound was identified using MS/FAB. C₂₀H₁₇N; calc. 271.13. found 271.36

Synthesis of Compound 8


[0120] 2.76 g (5 mmol) of Intermediate I-3, 1.36 g (5 mmol) of Intermediate I-4, 0.056 g (0.25 mmol) of Pd(OAc)₂, 0.76 g (0.25 mmol) of tri(o-tolyl)phosphine ((p-tolyl)₃P), and 1.019 g (10 mmol) of triethylamine were dissolved in 100 mL of dimethylacetamide (DMAc) to obtain a solution, which was then stirred at about 85° C. for about 4 hours. The reaction solution was cooled to room temperature, followed by three times of extraction with 100 mL of water and 100 mL of diethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 1.71 g of Compound 8 (Yield: 46%). This compound was identified using MS/FAB. C₅₆H₄₄N₂; calc. 744.35. found 744.53


[0121] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.18-8.14 (m, 1H), 7.93 (d, 1H), 7.88-7.85 (m, 1H), 7.76 (d, 1H), 7.71-7.69 (m, 1H), 7.64-7.60 (m, 2H), 7.48-7.45 (m, 3H), 7.40-7.38 (m, 1H), 7.35-7.30 (m, 3H), 7.23-7.03 (m, 9H), 6.70-6.63 (m, 5H), 6.17-6.13 (m, 4H), 1.76 (s, 6H), 1.64 (s, 6H)

Synthesis Example 2

Synthesis of Compound 16

[0122]








[0123] Intermediate I-5 was synthesized using iodobenzene, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Intermediate I-3 in Synthesis Example 1. Intermediate I-6 was synthesized using Intermediate I-5, instead of Intermediate I-3, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1.

[0124] 1.76 g (2.8 mmol) of Intermediate I-6, 0.081 g (0.08 mmol) of (carbonyl)chloro(hydrido)tris(triphenylphosphine) ruthenium(II), and 0.56 g (28 mmol) of D₂O were dissolved in 30 mL of 1,4-dioxane to obtain a solution, which was then stirred at about 80° C. for about 12 hours. The reaction solution was cooled to room temperature, followed by solvent removal, three times of extraction with 50 mL of water and 50 mL of dichloromethane. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 1.33 g of Compound 16 (Yield: 76%). This compound was identified using MS/FAB. C₄₇H₃₄D₂N₂: calc. 630.81. found 630.92

[0125] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.73-7.71 (m, 1H), 7.61 (d, 1H), 7.56-7.51 (m, 2H), 7.48-7.30 (m, 7H), 7.24-7.19 (m, 1H), 7.08-7.02 (m, 5H), 6.79-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)





Synthesis of Intermediate I-7

[0127] 1.96 ml (11.0 mmol) of diisobutylaluminumhydride was added to 10 mL of tetrahydrofuran (THF) to obtain a solution, followed by temperature control to about 0° C., a slow dropwise addition of 1.78 mL (10.0 mmol) of diphenylacetylene, and stirring at room temperature for about 6 hours. The reaction solution was cooled to about -78° C., and 0.77 mL (15.0 mmol) of a bromine solution was slowly added thereto over about 10 minutes. The temperature of the reaction solution was then raised to room temperature and stirred for about 1 hour. 10.0 mL of sodium potassium tartrate was slowly dropwise added to the reaction solution at about 0° C., followed by stirring at room temperature for about 30 minutes, and three times of extraction with 20 mL of water and 20 mL of diethylether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 1.74 g of Intermediate I-7 (Yield: 67%). This compound was identified using MS/FAB. $C_{56}H_{44}N_2$; calc. 258.00. found 258.13

Synthesis of Intermediate I-8

[0128] 5.18 g (20.0 mmol) of Intermediate I-7, 4.02 g (20.0 mmol) of 4-bromophenyl boronic acid, 1.15 g (1.0 mmol) of tetrakis(triphenylphosphine)palladium ($Pd(PPh_3)_4$), and 8.29 g (60.0 mmol) of K_2CO_3 were dissolved in 60 mL of a mixed tetrahydrofuran (THF) and H_2O (2:1 by volume) solution to obtain a solution, which was then stirred at about 70° C. for about 5 hours. The reaction solution was cooled to room temperature, and 40 mL of water was added thereto, followed by three times of extraction with 50 mL of ethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to

obtain 5.49 g of Intermediate I-8 (Yield: 82%). This compound was identified using MS/FAB. $C_{20}H_{15}Br$; calc. 334.04. found 334.14

Synthesis of Intermediate I-9

[0129] Intermediate I-9 was synthesized using Intermediate I-8, instead of 4-bromostyrene, in the same manner as in the synthesis of Intermediate I-4 in Synthesis Example 1.

Synthesis of Intermediate I-10

[0130] Intermediate I-10 was synthesized using 2-bromopyridine, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Intermediate I-3 in Synthesis Example 1.

Synthesis of Compound 19



[0131] Compound 19 was synthesized using Intermediates I-10 and I-9, instead of Intermediates I-3 and I-4, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{58}H_{43}N_3$; calc. 781.34. found 781.53


[0132] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.73-8.71 (m, 1H), 8.12-8.05 (m, 1H), 7.84-7.79 (m, 1H), 7.74-7.73 (m, 1H), 7.69-7.56 (m, 4H), 7.48-7.32 (m, 13H), 7.27-7.25 (m, 1H), 7.11-7.03 (m, 5H), 6.91-6.88 (m, 2H), 6.72-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 4

Synthesis of Compound 29

[0133]





Synthesis of Intermediate I-11

[0134] 5.37 g (20.0 mmol) of 2,4-dibromo-6-fluoro-phenylamine, 4.88 g (40.0 mmol) of phenylboronic acid, 1.15 g (1.0 mmol) of tetrakis(triphenylphosphine)palladium ($\text{Pd}(\text{PPh}_3)_4$), and 8.29 g (60.0 mmol) of K_2CO_3 were dissolved in 60 mL of a mixed tetrahydrofuran (THF) and H_2O (2:1 by volume) solution to obtain a solution, which was then stirred at about 70° C. for about 5 hours. The reaction solution was cooled to room temperature, and 40 mL of water was added thereto, followed by three times of extraction with 50 mL of ethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 3.95 g of Intermediate I-11 (Yield: 75%). This compound was identified using MS/FAB. $\text{C}_{18}\text{H}_{14}\text{FN}$: calc. 263.11. found 263.25

Synthesis of Intermediate I-12

[0135] 2.63 g (10 mmol) of Intermediate I-11, 2.45 g (12 mmol) of iodobenzene, 0.83 g (0.2 mmol) of $\text{Pd}_2(\text{dba})_3$, 0.040 g (0.2 mmol) of PtBu_3 , and 13.9 g (20 mmol) of KOtBu were dissolved in 60 mL of toluene to obtain a solution, which was then stirred at about 85° C. for about 4 hours. The reaction solution was cooled to room temperature, followed by three times of extraction with 100 mL of water and 100 mL of diethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 2.14 g of Intermediate I-12 (Yield: 63%). This compound was identified using MS/FAB. $\text{C}_{24}\text{H}_{18}\text{FN}$: calc. 339.14. found 339.29

Synthesis of Intermediate I-13

[0136] Intermediate I-13 was synthesized using Intermediate I-12, instead of diphenylamine, in the same manner as in

the synthesis of Intermediate I-4 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{32}H_{24}FN$: calc. 441.53. found 441.71

Synthesis of Compound 29



[0137] Compound 29 was synthesized using Intermediates I-5 and I-13, instead of Intermediates I-3 and I-4, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{59}H_{43}FN_2$: calc. 798.34. found 798.53

[0138] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.11 (m, 1H), 7.93 (d, 1H), 7.77-7.76 (m, 1H), 7.72-7.70 (m, 2H), 7.65-7.59 (m, 3H), 7.55-7.29 (m, 18H), 7.24-7.19 (m, 1H), 7.14-7.04 (m, 5H), 6.66-6.60 (m, 3H), 6.15-6.12 (m, 2H), 1.78 (s, 6H)



Synthesis Example 5

Synthesis of Compound 39

[0139]



-continued



Synthesis of Intermediate I-14

[0140] 20.1 g (60 mmol) of 9,10-dibromoanthracene, 8.461 g (50 mmol) of diphenylamine, 0.915 g (1 mmol) of $Pd_2(dba)_3$, 0.202 g (1 mmol) of $PtBu_3$, and 69.611 g (100 mmol) of $KOtBu$ were dissolved in 300 mL of toluene to obtain a solution, which was then stirred at about 85° C. for about 4 hours. The reaction solution was cooled to room temperature, followed by three times of extraction with 100 mL of water and 100 mL of diethyl ether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 12.9 g of Intermediate I-14 (Yield: 61%). This compound was identified using MS/FAB. $C_{26}H_{18}BrN$: calc. 423.06. found 423.13

Synthesis of Intermediate I-15

[0141] 4.24 g (10.0 mmol) of Intermediate I-14, 2.54 g (10.0 mmol) of bis(pinacolato)diborane, 0.36 g (0.5 mmol) of [1,1'-bis(diphenylphosphino)ferrocene]dichloro palladium (II) ($PdCl_2(dppf)_2$), and 2.94 g (30.0 mmol) of $KOAc$ were

dissolved in 40 mL of DMSO to obtain a solution, which was then stirred at about 80° C. for about 6 hours. The reaction solution was cooled to room temperature, followed by three times of extraction with 50 mL of water and 50 mL of diethylether. The organic phase was collected and was dried using magnesium sulfate to evaporate the solvent. The residue was separated and purified using silica gel column chromatography to obtain 1.88 g of Intermediate I-15 (Yield: 74%). This compound was identified using MS/FAB. $C_{32}H_{30}BNO_2$; calc. 471.23. found 471.33

Synthesis of Intermediate I-16

[0142] Intermediate I-16 was synthesized using Intermediate I-15 and vinyl bromide, instead of 4-bromophenyl boronic acid and Intermediate I-7, in the same manner as in the synthesis of Intermediate I-8 in Synthesis Example 3. This compound was identified using MS/FAB. $C_{28}H_{21}N$; calc. 371.47. found 371.61

Synthesis of Compound 39


[0143] Compound 39 was synthesized using Intermediates I-5 and I-16, instead of Intermediates I-3 and I-4, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{55}H_{40}N_2$; calc. 728.31. found 728.54

[0144] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.93 (d, 1H), 7.89-7.86 (d, 1H), 7.75-7.72 (m, 2H), 7.62-7.58 (m, 3H), 7.56-7.48 (m, 3H), 7.44-7.41 (m, 2H), 7.38-7.30 (m, 4H), 7.24-7.19 (m, 1H), 7.15 (s, 1H), 7.11 (s, 1H), 7.04-6.98 (m, 6H), 6.88-6.84 (m, 2H), 6.63-6.60 (m, 2H), 5.96-5.94 (m, 4H), 1.77 (s, 6H)


Synthesis Example 6

Synthesis of Compound 47

[0145]



-continued



Synthesis of Intermediate I-17

[0146] Intermediate I-17 was synthesized using Intermediate I-5, instead of Intermediate I-14, in the same manner as in the synthesis of Intermediate I-15 in Synthesis Example 5. This compound was identified using MS/FAB. $C_{33}H_{32}BNO_2$: calc. 485.25. found 485.33

Synthesis of Intermediate I-18

[0147] Intermediate I-18 was synthesized using Intermediate I-17 and 2,5-dibromopyridine, instead of 2-bromonitrobenzene and 2-bromo-9,9-dimethyl-7-fluoreneboronic acid, in the same manner as in the synthesis of Intermediate I-1 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{32}H_{23}BrN_2$: calc. 515.44. found 515.62

Synthesis of Compound 47

[0148] Compound 47 was synthesized using Intermediate I-18, instead of Intermediate I-3, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{52}H_{39}N_3$: calc. 705.31. found 705.43

[0149] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.87-8.85 (m, 1H), 8.16-8.12 (m, 1H), 8.07-8.04 (m, 1H), 7.92 (d, 1H), 7.86-7.83 (m, 1H), 7.78 (d, 1H), 7.69-7.67 (m, 1H), 7.62-7.57 (m, 2H), 7.56-7.48 (m, 4H), 7.44-7.41 (m, 2H), 7.38-7.30 (m, 4H), 7.26-7.19 (m, 2H), 7.08-7.03 (m, 4H), 6.85-6.82 (m, 2H), 6.66-6.63 (m, 2H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 7

Synthesis of Compound 1

[0150] Compound 1 was synthesized using ethylodide (C_2H_5I), instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{43}H_{36}N_2$: calc. 580.28. found 580.42

[0151] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.17 (d, 1H), 8.00 (d, 1H), 7.75 (d, 1H), 7.69-7.65 (m, 2H), 7.49-7.36 (m, 6H), 7.24-7.20 (m, 1H), 7.08-7.00 (m, 5H), 6.69-6.63 (m, 4H), 6.17-6.12 (m, 4H), 4.71 (q, 2H), 1.77 (s, 6H), 1.31 (t, 3H)

Synthesis Example 8

Synthesis of Compound 3

[0152] Compound 3 was synthesized using iodobenzene (C_6H_5I), instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis

Example 1. This compound was identified using MS/FAB. $C_{47}H_{36}N_2$: calc. 628.28. found 628.41

[0153] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.73-7.71 (m, 1H), 7.61 (d, 1H), 7.56-7.51 (m, 3H), 7.48-7.30 (m, 9H), 7.24-7.19 (m, 1H), 7.08-7.02 (m, 4H), 6.79-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 9

Synthesis of Compound 4

[0154] Compound 4 was synthesized using 2-iodonaphthalene, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{51}H_{38}N_2$: calc. 678.30. found 678.61

[0155] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 2H), 7.84 (d, 1H), 7.75 (d, 1H), 7.72-7.70 (m, 1H), 7.65-7.61 (m, 2H), 7.57-7.52 (m, 2H), 7.48-7.30 (m, 9H), 7.22-7.18 (m, 1H), 7.08-7.03 (m, 5H), 6.7-6.63 (m, 4H), 6.17-6.13 (m, 3H), 1.76 (s, 6H)

Synthesis Example 10

Synthesis of Compound 5

[0156] Compound 5 was synthesized using 9-bromophenanthrene, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{55}H_{40}N_2$: calc. 728.31. found 728.62

[0157] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.66-8.64 (m, 1H), 8.31-8.29 (m, 1H), 8.17-8.12 (m, 2H), 7.98 (d, 1H), 7.76 (d, 1H), 7.71-7.64 (m, 3H), 7.59-7.54 (m, 1H), 7.51-7.30 (m, 9H), 7.35-7.30 (m, 2H), 7.08-7.03 (m, 5H), 6.70-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 11

Synthesis of Compound 7

[0158] Compound 7 was synthesized using 4-bromobiphenyl, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{53}H_{40}N_2$: calc. 704.31. found 704.65

[0159] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 1H), 7.76 (d, 1H), 7.71-7.69 (m, 1H), 7.64-7.59 (m, 3H), 7.53-7.38 (m, 9H), 7.35-7.29 (m, 3H), 7.20-7.16 (m, 2H), 7.08-7.03 (m, 5H), 6.70-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 12

Synthesis of Compound 9

[0160] Compound 9 was synthesized using iodobenzene-d5, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{47}H_{31}D_5N_2$: calc. 633.31. found 633.59

[0161] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 1H), 7.76 (d, 1H), 7.71-7.69 (m, 1H), 7.62-7.59 (m, 1H), 7.48-7.44 (m, 2H), 7.35-7.30 (m, 3H), 7.23-7.19 (m, 1H), 7.08-7.03 (m, 6H), 6.70-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 13

Synthesis of Compound 12

[0162] Compound 9 was synthesized using Intermediate I-19, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{56}H_{41}N_5$: calc. 783.33. found 783.54

[0163] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.83 (d, 1H), 8.78-8.75 (m, 3H), 8.15 (m, 1H), 7.92-7.88 (m, 1H), 7.77 (d, 1H), 7.71-7.57 (m, 7H), 7.48-7.35 (m, 8H), 7.08-7.03 (m, 5H), 6.68-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 14

Synthesis of Compound 14

[0164] Intermediate I-20 was synthesized using 2-bromo-9,9-diphenyl-7-fluoreneboronic acid and iodobenzene, instead of 2-bromo-9,9-dimethyl-7-fluoreneboronic acid and 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1.

[0165] Intermediate 14 was synthesized using Intermediate I-20, instead of Intermediate I-6, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0166] $C_{57}H_{38}D_2N_2$: calc. 754.33. found 754.61

[0167] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 2H), 7.74 (d, 1H), 7.56-7.51 (m, 2H), 7.47-7.31 (m, 12H), 7.26-7.22 (m, 1H), 7.15-7.11 (m, 1H), 7.08-6.98 (m, 13H), 6.69-6.63 (m, 3H), 6.16-6.13 (m, 3H),

Synthesis Example 15

Synthesis of Compound 17

[0168] Compound 17 was synthesized using Intermediate I-5, instead of Intermediate I-10, in the same manner as in the synthesis of Compound 19 in Synthesis Example 3. This compound was identified using MS/FAB.

[0169] $C_{59}H_{44}N_2$: calc. 780.35. found 780.58

[0170] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.77 (d, 1H), 7.69-7.66 (m, 2H), 7.61-7.59 (m, 1H), 7.56-7.51 (m, 2H), 7.48-7.30 (m, 14H), 7.24-7.19 (m, 1H), 7.11-7.03 (m, 6H), 6.91-6.88 (m, 2H), 6.72-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 16

Synthesis of Compound 20

[0171] Intermediate I-22 was synthesized using Intermediate I-21, instead of 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1.

[0172] Compound 20 was synthesized using Intermediate I-22, instead of Intermediate I-6, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0173] $C_{55}H_{42}D_2N_2$: calc. 782.36. found 782.59

[0174] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.26-8.21 (m, 1H), 7.93 (d, 1H), 7.81-7.76 (m, 5H), 7.72-7.70 (m, 1H), 7.66-7.64 (m, 1H), 7.55-7.50 (m, 5H), 7.47-7.29 (m, 9H), 7.08-7.03 (m, 4H), 6.69-6.63 (m, 5H), 6.16-6.13 (m, 4H), 1.78 (s, 6H)

Synthesis Example 17

Synthesis of Compound 21

[0175] Compound 21 was synthesized using tert-butyl iodide and Intermediates I-23 in the synthesis of Intermediate I-3, instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{47}H_{44}N_2$: calc. 636.35. found 636.59

[0176] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 7.97-7.94 (m, 1H), 7.92 (d, 1H), 7.77-7.74 (m, 1H), 7.70 (d, 1H), 7.65-7.63 (m, 1H), 7.48-7.39 (m, 5H), 7.35-7.19 (m, 5H), 7.07-7.03 (m, 4H), 6.99-6.94 (m, 3H), 6.82-6.78 (m, 3H), 6.63-6.61 (m, 2H), 6.53-6.50 (m, 2H), 1.87 (s, 3H), 1.80 (s, 6H), 1.79 (s, 6H)

Synthesis Example 18

Synthesis of Compound 22

[0177] Compound 22 was synthesized using iodobenzene and Intermediate I-24 in the synthesis of Intermediate I-3, instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{48}H_{37}FN_2$: 22. calc. 660.29. found 660.42

[0178] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.72-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.56-7.51 (m, 2H), 7.48-7.31 (m, 9H), 7.24-7.19 (m, 1H), 7.10-7.03 (m, 3H), 7.00-6.92 (m, 3H), 6.82-6.78 (m, 1H), 6.70-6.67 (m, 1H), 6.63-6.57 (m, 3H), 2.01 (s, 3H), 1.76 (s, 6H)

Synthesis Example 19

Synthesis of Compound 23

[0179] Compound 23 was synthesized using iodobenzene and Intermediate I-25 in the synthesis of Intermediate I-3, instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB.

[0180] $C_{48}H_{35}N_3$: calc. 653.28. found 653.49

[0181] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.10 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.72-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.56-7.51 (m, 2H), 7.48-7.30 (m, 11H), 7.24-7.19

(m, 1H), 7.10-7.03 (m, 3H), 6.78-6.73 (m, 4H), 6.66-6.63 (m, 1H), 6.22-6.20 (m, 2H), 1.73 (s, 6H)

Synthesis Example 20

Synthesis of Compound 26

[0182] Compound 26 was synthesized using Intermediate I-26, instead of Intermediate I-4, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0183] $C_{57}H_{38}D_2N_2$: calc. 754.33. found 754.51

[0184] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.12 (m, 3H), 8.04-8.00 (m, 2H), 7.93 (d, 1H), 7.82-7.76 (m, 4H), 7.71-7.69 (m, 1H), 7.62-7.51 (m, 5H), 7.46-7.31 (m, 9H), 7.24-7.19 (m, 1H), 7.06-7.01 (m, 2H), 6.73-6.70 (m, 2H), 6.65-6.61 (m, 1H), 6.22-6.17 (m, 2H), 1.75 (s, 6H)

Synthesis Example 21

Synthesis of Compound 27

[0185] Compound 27 was synthesized using Intermediate I-27, instead of Intermediate I-4, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0186] $C_{56}H_{42}D_2N_2$: calc. 746.36. found 746.41

[0187] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.93 (d, 1H), 7.77-7.71 (m, 2H), 7.72-7.71 (m, 1H), 7.61 (d, 1H), 7.56-7.51 (m, 3H), 7.48-7.30 (m, 7H), 7.23-7.19 (m, 1H), 7.14-7.04 (m, 5H), 6.73-6.63 (m, 5H), 6.39-6.37 (m, 1H), 6.23-6.20 (m, 2H), 1.76 (s, 6H), 1.61 (s, 6H)

Synthesis Example 22

Synthesis of Compound 28

[0188] Compound 28 was synthesized using Intermediate I-28, instead of Intermediate I-4, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0189] $C_{59}H_{41}D_2N_3$: calc. 795.35. found 795.53

[0190] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.22-8.20 (m, 1H), 8.16-8.11 (m, 1H), 7.93 (d, 1H), 7.77-7.76 (m, 1H), 7.72-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.56-7.19 (m, 21H), 7.09-7.04 (m, 2H), 6.87-6.81 (m, 3H), 6.66-6.63 (m, 1H), 6.31-6.29 (m, 2H), 1.77 (s, 6H)

Synthesis Example 23

Synthesis of Compound 30

[0191] Compound 30 was synthesized using Intermediate I-30, instead of Intermediate I-4, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0192] $C_{47}H_{24}D_{12}N_2$: calc. 640.36. found 640.58

[0193] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.11 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.72-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.54-7.51 (m, 2H), 7.46-7.29 (m, 8H), 7.23-7.19 (m, 1H), 6.69-6.66 (m, 2H), 1.76 (s, 6H)

Synthesis Example 24

Synthesis of Compound 31

[0194] Compound 31 was synthesized using iodobenzene and Intermediate I-31 in the synthesis of Intermediate I-3,

instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB.

[0195] $C_{45}H_{34}N_4$: calc. 630.27. found 630.39

[0196] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.53-8.50 (m, 2H), 8.16-8.11 (m, 1H), 8.08-8.06 (m, 2H), 7.93 (d, 1H), 7.77-7.76 (m, 1H), 7.72-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.55-7.30 (m, 12H), 7.24-7.20 (m, 1H), 7.17-7.14 (m, 2H), 7.07 (s, 1H), 7.03 (s, 1H), 6.97-6.93 (m, 2H), 1.75 (s, 6H)

Synthesis Example 25

Synthesis of Compound 35

[0197] Compound 35 was synthesized using iodobenzene and Intermediate I-32 in the synthesis of Intermediate I-3, instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{53}H_{38}N_2S$: calc. 734.27. found 734.59

[0198] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.12 (m, 1H), 8.06-8.04 (m, 1H), 7.93 (d, 1H), 7.82 (d, 1H), 7.75 (d, 1H), 7.72-7.70 (m, 2H), 7.68-7.66 (m, 1H), 7.62-7.57 (m, 2H), 7.56-7.29 (m, 12H), 7.24-7.19 (m, 1H), 7.09-7.03 (m, 3H), 6.87-6.81 (m, 3H), 6.66-6.63 (m, 1H), 6.32-6.29 (m, 2H), 1.78 (s, 6H)

Synthesis Example 26

Synthesis of Compound 36

[0199] Compound 36 was synthesized using iodobenzene and Intermediate I-33 in the synthesis of Intermediate I-3, instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{53}H_{38}N_{20}$: calc. 718.29. found 718.59

[0200] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.17-8.12 (m, 1H), 7.93 (d, 1H), 7.83 (d, 1H), 7.77-7.71 (m, 4H), 7.62-7.56 (m, 3H), 7.55-7.30 (m, 12H), 7.24-7.19 (m, 1H), 7.09-7.03 (m, 3H), 6.92-6.88 (m, 1H), 6.84-6.81 (m, 2H), 6.66-6.63 (m, 1H), 6.32-6.29 (m, 2H), 1.78 (s, 6H)

Synthesis Example 27

Synthesis of Compound 37

[0201] Compound 37 was synthesized using iodobenzene and Intermediate I-34 in the synthesis of Intermediate I-3, instead of 9,9-dimethyl-2-iodofluorene and diphenylamine, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{53}H_{52}N_2Si_2$: calc. 772.36. found 772.55

[0202] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.93 (d, 1H), 7.83 (d, 1H), 7.77-7.75 (m, 1H), 7.72-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.56-7.51 (m, 2H), 7.48-7.41 (m, 4H), 7.39-7.30 (m, 7H), 7.22-7.19 (m, 1H), 7.07 (s, 1H), 7.03 (s, 1H), 6.83-6.80 (m, 2H), 6.60-6.56 (m, 4H), 1.76 (s, 6H), 0.24 (s, 18H)

Synthesis Example 28

Synthesis of Compound 40

[0203] Compound 40 was synthesized using 1,4-dibromobenzene, instead of 9,10-dibromoanthracene, in the same

manner as in the synthesis of Compound 39 in Synthesis Example 5. This compound was identified using MS/FAB.

[0204] C₅₃H₄₀N₂: calc. 704.31. found 704.54

[0205] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.71-7.70 (m, 1H), 7.62-7.59 (m, 1H), 7.56-7.41 (m, 11H), 7.38-7.30 (m, 3H), 7.24-7.19 (m, 1H), 7.08-7.01 (m, 6H), 6.86-6.82 (m, 2H), 6.66-6.63 (m, 2H), 6.16-6.13 (m, 4H), 1.78 (s, 6H)

Synthesis Example 29

Synthesis of Compound 41

[0206] Intermediate I-35 was synthesized using 2,7-dibromo-9,9-dimethylfluorene, instead of 9,10-dibromoanthracene, in the same manner as in the synthesis of Compound 39 in Synthesis Example 5.

[0207] Compound 41 was synthesized using Intermediate I-35, instead of Intermediate I-6, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0208] C₅₆H₄₂D₂N₂: calc. 746.36. found 746.62

[0209] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.93 (d, 1H), 7.77-7.59 (m, 6H), 7.56-7.52 (m, 2H), 7.44-7.30 (m, 7H), 7.24-7.19 (m, 1H), 7.09-7.04 (m, 4H), 6.67-6.63 (m, 3H), 6.46-6.45 (m, 1H), 6.16-6.13 (m, 4H), 1.78 (s, 6H), 1.61 (s, 6H)

Synthesis Example 30

Synthesis of Compound 44

[0210] Compound 44 was synthesized using 1,4-dibromobenzene in the synthesis of Intermediate I-18, instead of 2,5-dibromopyridine, in the same manner as in the synthesis of Compound 47 in Synthesis Example 6. This compound was identified using MS/FAB. C₅₃H₄₀N₂: calc. 704.31. found 704.43

[0211] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.83-7.77 (m, 3H), 7.66-7.59 (m, 2H), 7.54-7.50 (m, 2H), 7.48-7.30 (m, 6H), 7.38-7.30 (m, 4H), 7.26 (s, 1H), 7.22-7.19 (m, 1H), 7.08-7.03 (m, 4H), 6.91 (s, 1H), 6.87 (s, 1H), 6.70-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.78 (s, 6H)

Synthesis Example 31

Synthesis of Compound 48

[0212] Compound 48 was synthesized using 2,5-dibromofuran, instead of 2,5-dibromopyridine, in the same manner as in the synthesis of Compound 47 in Synthesis Example 6. This compound was identified using MS/FAB. C₅₁H₃₈N₂O: calc. 694.29. found 694.51

[0213] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.96-7.93 (m, 1H), 7.79-7.76 (m, 2H), 7.62-7.51 (m, 5H), 7.46-7.41 (m, 3H), 7.38-7.30 (m, 3H), 7.24-7.19 (m, 2H), 7.08-7.00 (m, 5H), 6.93-6.91 (m, 1H), 6.70-6.63 (m, 5H), 6.16-6.13 (m, 4H), 1.79 (s, 6H)

Synthesis Example 32

Synthesis of Compound 49

[0214] Compound 49 was synthesized using 2,5-dibromothiophene, instead of 2,5-dibromopyridine, in the same manner as in the synthesis of Compound 47 in Synthesis

Example 6. This compound was identified using MS/FAB. C₅₁H₃₈N₂S: calc. 710.27. found 710.46

[0215] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.85-7.83 (m, 1H), 7.78 (d, 1H), 7.62-7.59 (m, 1H), 7.56-7.50 (m, 4H), 7.44-7.41 (m, 2H), 7.38-7.30 (m, 5H), 7.28-7.19 (m, 2H), 7.12-7.02 (m, 7H), 6.76-6.73 (m, 2H), 6.66-6.63 (m, 2H), 6.16-6.13 (m, 4H), 1.79 (s, 6H)

Synthesis Example 33

Synthesis of Compound 51

[0216] Compound 51 was synthesized using 2,7-dibromo-9,9-dimethyl-9H-fluorene, instead of 2,5-dibromopyridine, in the same manner as in the synthesis of Compound 47 in Synthesis Example 6. This compound was identified using MS/FAB.

[0217] C₆₂H₄₈N₂: calc. 820.38. found 820.46 ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.17-8.11 (m, 1H), 7.82-7.77 (m, 2H), 7.69-7.64 (m, 4H), 7.62-7.59 (m, 1H), 7.57-7.30 (m, 14H), 7.24-7.19 (m, 1H), 7.08-7.03 (m, 5H), 6.70-6.63 (m, 4H), 6.16-6.13 (m, 4H), 1.70 (s, 6H), 1.60 (s, 6H)

Synthesis Example 34

Synthesis of Compound 54

[0218] Intermediate I-36 was synthesized using 2-bromo-4-(trifluoromethyl)-1-nitrobenzene in the synthesis of Intermediate I-1, instead of 2-bromonitrobenzene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1.

[0219] Intermediate 54 was synthesized using Intermediate I-36, instead of Intermediate I-6, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0220] C₄₈H₃₃D₂F₃N₂: calc. 698.28. found 698.41

[0221] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.12-8.10 (m, 1H), 7.90 (d, 1H), 7.77-7.74 (m, 2H), 7.71-7.70 (m, 1H), 7.63 (d, 1H), 7.56-7.50 (m, 2H), 7.47-7.34 (m, 6H), 7.08-7.04 (m, 5H), 6.69-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 35

Synthesis of Compound 56

[0222] Intermediate I-38 was synthesized using Intermediate I-37 in the synthesis of Intermediate I-1, instead of 2-bromonitrobenzene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 2.

[0223] Intermediate 56 was synthesized using Intermediate I-38, instead of Intermediate I-6, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.

[0224] C₅₃H₃₈D₂N₂: calc. 706.33. found 706.57

[0225] ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 8.00-7.99 (m, 1H), 7.91 (d, 1H), 7.78-7.76 (m, 1H), 7.73-7.69 (m, 2H), 7.66-7.61 (m, 2H), 7.56-7.34 (m, 13H), 7.08-7.03 (m, 4H), 6.70-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.78 (s, 6H)

Synthesis Example 36

Synthesis of Compound 61

[0226] Compound 61 was synthesized using Intermediate I-39 and iodobenzene in the synthesis of Intermediate I-1, instead of 2-bromonitrobenzene and 9,9-dimethyl-2-iodof-

luorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{59}H_{45}N_3$: calc. 795.36. found 795.61

[0227] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 7.96 (d, 1H), 7.77 (d, 1H), 7.72-7.70 (m, 1H), 7.65-7.64 (m, 1H), 7.56-7.51 (m, 2H), 7.49-7.42 (m, 5H), 7.39-7.34 (m, 3H), 7.15-7.13 (m, 1H), 7.09-7.03 (m, 9H), 6.76-6.73 (m, 1H), 6.70-6.62 (m, 6H), 6.25-6.21 (m, 4H), 6.17-6.12 (m, 4H), 1.80 (s, 6H)

Synthesis Example 37

Synthesis of Compound 62

[0228] Compound 62 was synthesized using Intermediate I-40 and iodobenzene in the synthesis of Intermediate I-1, instead of 2-bromonitrobenzene and 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{65}H_{49}N_3$: calc. 871.39. found 871.51

[0229] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.12-8.10 (m, 1H), 7.91 (d, 1H), 7.77-7.76 (m, 1H), 7.71-7.69 (m, 2H), 7.63-7.51 (m, 6H), 7.48-7.42 (m, 6H), 7.39-7.34 (m, 2H), 7.08-7.02 (m, 8H), 6.86-6.82 (m, 2H), 6.70-6.63 (m, 6H), 6.16-6.13 (m, 8H), 1.77 (s, 6H)

Synthesis Example 38

Synthesis of Compound 64







[0230] Compound 64 was synthesized using Intermediate I-41 and iodobenzene in the synthesis of Intermediate I-1, instead of 2-bromonitrobenzene and 9,9-dimethyl-2-iodofluorene, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{62}H_{45}N_5$: calc. 859.36. found 859.47

[0231] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 9.07-9.06 (m, 1H), 8.80-8.76 (m, 4H), 8.56-8.54 (m, 1H), 7.95-7.89 (m, 3H), 7.77-7.76 (m, 1H), 7.71-7.70 (m, 1H), 7.63-7.60 (m, 4H), 7.55-7.34 (m, 11H), 7.08-7.03 (m, 5H), 6.69-6.63 (m, 4H), 6.16-6.13 (m, 4H), 1.75 (s, 6H)


Synthesis Example 39

Synthesis of Compound 66

[0232]



-continued



66

[0233] Intermediate I-42, instead of Intermediate I-2, was synthesized using Intermediate 9,9-dimethyl-7-fluoreneboronic acid, instead of 2-bromo-9,9-dimethyl-7-fluoreneboronic acid, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. Subsequently, Intermediate I-43, instead of Intermediate I-3, was synthesized using 1-bromo-4-iodobenzene, instead of 9,9-dimethyl-2-iodofluorene. Next, Intermediate I-44, instead of Compound 8, was synthesized by reacting Intermediate I-43, instead of Intermediate I-3, with Intermediate I-4.

[0234] Compound 66 was synthesized using Intermediate I-44, instead of Intermediate I-6, in the same manner as in the synthesis of Compound 16 in Synthesis Example 2. This compound was identified using MS/FAB.


[0235] $C_{47}H_{34}D_2N_2$; calc. 630.30. found 630.41

[0236] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.11 (m, 1H), 7.86-7.84 (m, 1H), 7.77 (d, 1H), 7.72-7.68 (m, 2H), 7.63-7.61 (m, 1H), 7.44-7.40 (m, 4H), 7.35-7.29 (m, 2H), 7.22-7.16 (m, 3H), 7.09-7.03 (m, 5H), 6.69-6.63 (m, 4H), 6.16-6.13 (m, 4H), 1.74 (s, 6H)

Synthesis Example 40

Synthesis of Compound 68

[0237]



Synthesis of Intermediate I-46

[0238] As illustrated in a synthesis flow diagram above, 2-bromo-7-iodo-9,9-dimethyl-9H-fluorene was reacted with styrene to obtain Intermediate I-45, which was then reacted with bis(pinacolato)diborane to obtain Intermediate I-46.

Synthesis of Compound 68

[0239] Intermediate 68 was synthesized using Intermediate I-46, instead of 9,9-dimethyl-7-fluoreneboronic acid, in the same manner as in the synthesis of Intermediate I-44 in Synthesis Example 39. This compound was identified using MS/FAB.


[0240] $C_{55}H_{42}N_2$; calc. 730.33. found 730.42


[0241] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.16-8.11 (m, 1H), 7.93 (d, 1H), 7.77 (d, 1H), 7.73-7.69 (m, 2H), 7.62-7.59 (m, 1H), 7.52-7.46 (m, 2H), 7.45-7.26 (m, 13H), 7.20-7.15 (m, 2H), 7.08-7.03 (m, 5H), 6.70-6.63 (m, 4H), 6.17-6.13 (m, 4H), 1.76 (s, 6H)

Synthesis Example 41

Synthesis of Compound 71

[0242]









synthesis of Intermediate I-3 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{35}H_{27}N$: calc. 461.21. found 461.28

Synthesis of Compound 71

[0248] Compound 71 was synthesized using 4-bromotriphenylamine and Intermediate I-51, instead of Intermediates I-3 and I-4, in the same manner as in the synthesis of Compound 8 in Synthesis Example 1. This compound was identified using MS/FAB. $C_{53}H_{40}N_2$: calc. 704.31. found 704.40

[0249] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.04-8.01 (m, 1H), 7.85-7.83 (m, 1H), 7.78 (d, 1H), 7.71-7.69 (m, 1H), 7.63-7.60 (m, 4H), 7.56-7.41 (m, 8H), 7.38-7.34 (m, 1H), 7.30-7.26 (m, 1H), 7.22-7.17 (m, 2H), 7.09-7.03 (m, 5H), 6.91-6.87 (m, 1H), 6.70-6.63 (m, 4H), 6.16-6.11 (m, 4H), 1.74 (s, 6H)



Synthesis Example 42


Synthesis of Compound 73

[0250] Compound 73 was synthesized using Intermediate I-46 of Synthesis Example in the synthesis of Intermediate I-49, instead of 9,9-dimethyl-7-fluoreneboronic acid, in the same manner as in the synthesis of Compound 71 in Synthesis Example 41. This compound was identified using MS/FAB. $C_{73}H_{55}N_3$: calc. 973.43. found 973.52

[0251] 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 8.04-8.01 (m, 1H), 7.91 (d, 1H), 7.77 (d, 1H), 7.71-7.69 (m, 2H), 7.64-7.61 (m, 4H), 7.56-7.51 (m, 2H), 7.49-7.26 (m, 14H), 7.08-7.03 (m, 8H), 6.91-6.87 (m, 1H), 6.70-6.63 (m, 8H), 6.16-6.11 (m, 7H), 1.76 (s, 6H)


[0252] Intermediates I-19 and I-44 used in Synthesis Examples 7-42 are represented by the following formulae.

I-19




I-20

I-21




I-22

I-23



I-24



I-25



I-26



-continued

I-27





I-28

I-30




I-31


I-32



I-33




I-34





I-35

-continued




I-36

I-37





I-38





I-39

I-40




I-41

I-42



I-43

I-44



Example 1

[0253] A 15 Ω/cm^2 (1200 Å) ITO glass substrate (available from Corning Co.) was cut to a size of 50 mm×50 mm×0.7 mm, ultrasonically washed with isopropyl alcohol for 5 minutes and then with pure water for 5 minutes, and washed again with UV ozone for 30 minutes. 2-TNATA was vacuum-deposited on the ITO glass substrate to form an HIL having a thickness of 600 Å on the anode, and then 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPB) was vacuum-deposited on the HIL to form a HTL having a thickness of 300 Å. 98 wt % of ADN as a blue fluorescent host and 2 wt % of Compound 8 above as a fluorescent dopant were deposited on the HTL to form an EML having a thickness of 300 Å. Alq₃ was vacuum-deposited on the EML to form an ETL having a thickness of 300 Å. LiF was vacuum-deposited on the ETL to form an EIL having a thickness of 10 Å and Al was vacuum-deposited on the EIL to form a cathode having a thickness of 3,000 Å, thereby completing the manufacture of an organic light-emitting device.

Example 2

[0254] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 16, instead of Compound 8, was used as a dopant to form the EML.

Example 3

[0255] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 19, instead of Compound 8, was used as a dopant to form the EML.

Example 4

[0256] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 25, instead of Compound 8, was used as a dopant to form the EML.

Example 5

[0257] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 29, instead of Compound 8, was used as a dopant to form the EML.

Example 6

[0258] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 39, instead of Compound 8, was used as a dopant to form the EML.

Example 7

[0259] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 47, instead of Compound 8, was used as a dopant to form the EML.

Example 8

[0260] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 56, instead of Compound 8, was used as a dopant to form the EML.

Example 9

[0261] An organic light-emitting device was manufactured in the same manner as in Example 1, except that Compound 62, instead of Compound 8, was used as a dopant to form the EML.

Comparative Example 1

[0262] An organic light-emitting device was manufactured in the same manner as in Example 1, except that DPAVB_i, instead of Compound 8, was used as a dopant to form the EML.

Evaluation Example

[0263] Driving voltages, current densities, luminance, efficiencies, emitting-light colors, half-life spans of the organic light-emitting devices of Examples 1 to 9 and Comparative Example 1 were measured using a PR650 (Spectroscan) Source Measurement Unit. (available from Photo Research, Inc.). The results are shown in Table 1 below.

TABLE 1

	EML host	EML dopant	Driving voltage (V)	Current Density	Luminance (cd/m ²)	Efficiency (cd/A)	Luminescent Color	Half-life span (hr @100 mA)
Example 1	ADN	Compound 8	6.22	50	3,132	6.26	blue	251
Example 2	ADN	Compound 16	6.27	50	3,055	6.11	blue	315
Example 3	ADN	Compound 19	6.20	50	3,065	6.13	blue	264
Example 4	ADN	Compound 25	6.27	50	3,185	6.37	blue	298
Example 5	ADN	Compound 29	6.26	50	2,985	5.97	blue	253
Example 6	ADN	Compound 39	6.29	50	3,115	6.23	bluish green	241
Example 7	ADN	Compound 47	6.18	50	3,019	6.03	bluish green	263
Example 8	ADN	Compound 56	6.20	50	3,022	6.04	bluish green	244
Example 9	ADN	Compound 62	6.23	50	3,215	6.43	bluish green	249
Comparative Example 1	ADN	DPAVB _i	7.85	50	2,065	4.13	blue	145

¹A half life-span at a current density of 100 mA/cm²


[0264] Referring to Table 2, the organic light-emitting devices of Examples 1 to 9 were found to have better performance in terms of driving voltage, luminance, efficiency, and lifetime, as compared with the organic light-emitting device of Comparative Example 1.

[0265] As described above, an organic light-emitting device including the heterocyclic compound according to embodiments of the present invention may have a high performance, for example, a low driving voltage, a high luminance, a high efficiency, and a long lifetime.


[0266] While the present invention has been particularly shown and described with reference to exemplary embodiments thereof, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present invention as defined by the following claims

What is claimed is:

1. A heterocyclic compound represented by Formula 1A below:



wherein, in Formula 1A, R₁ to R₁₃ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₁-C₃₀ alkoxy group, a substituted or unsubstituted C₃-C₃₀ cycloalkyl group, a substituted or unsubstituted C₃-C₃₀ cycloalkenyl group, a substituted or unsubstituted C₅-C₃₀ aryl group, a substituted or unsubstituted C₂-C₃₀ heteroaryl group, a substituted or unsubstituted C₅-C₃₀ aryloxy group, a substituted or unsubstituted C₅-C₃₀ arylthio group, a—N(Q₁)(Q₂) group, and a group represented by Formula 1B below, wherein Q₁ and Q₂ are each independently one of a hydrogen atom, a deuterium atom, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₁-C₃₀ alkoxy group, a substituted or unsubstituted C₃-C₃₀ cycloalkyl group, a substituted or unsubstituted C₃-C₃₀ cycloalkenyl group, a substituted or unsubstituted C₆-C₃₀ aryl group, a substituted or unsubstituted C₆-C₃₀ aryloxy group, a substituted or unsubstituted C₆-C₃₀ arylthio group, and a substituted or unsubstituted C₃-C₃₀ heteroaryl group:



wherein, in Formula 1B, R₁₄ and R₁₅ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₂-C₃₀ alkenyl group, a substituted or unsubstituted C₁-C₃₀ alkoxy group, a substituted or unsubstituted C₃-C₃₀ cycloalkyl group, a substituted or unsubstituted C₃-C₃₀ cycloalkenyl group, a substituted or unsubstituted C₅-C₃₀ aryl group, a substituted or unsubstituted C₂-C₃₀ heteroaryl group, a substituted or unsubstituted C₅-C₃₀ aryloxy group, and a substituted or unsubstituted C₅-C₃₀ arylthio group;

at least one of R₁ to R₁₃ is a group represented by Formula 1B above;

Ar₁ and Ar₂ are each independently a substituted or unsubstituted C₅-C₃₀ aryl group or a substituted or unsubstituted C₃-C₃₀ heteroaryl group, wherein Ar₁ and Ar₂ are optionally linked to each other;


A and B are a divalent linker, and are each independently one of a substituted or unsubstituted C₅-C₃₀ arylene group or a substituted or unsubstituted C₃-C₃₀ heteroarylene group; and

a is an integer from 0 to 3, and b is an integer from 0 to 3, wherein if a is 2 or greater, the two or more A are identical to or different from each other, and if b is 2 or greater, the two or more B are identical to or different from each other.


2. The heterocyclic compound of claim 1, wherein Ar₁ and Ar₂ are each independently one of a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenoxy group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted carbozoyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyranlyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted diazinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzimidazolyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted azulenyl group, a substituted or unsubstituted heptalenyl group, a substituted or unsubstituted indacenyl group, a substituted or unsubstituted acenaphthyl group, a substituted or unsubstituted phenalenyl group, a substituted or unsubstituted phenanthridinyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluoranthenyl group, a substituted or unsubstituted triphenylenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted naphthacenyl group, a substituted or unsubstituted picenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted pentaphenyl group, a substituted or unsubstituted hexacenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolinyll group, a substituted or unsubstituted pyrazolyl group, a substituted or unsubstituted imidazopyridinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted imidazopyrimidinyl

group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted indolyl group, a substituted or unsubstituted indolizynyl group, a substituted or unsubstituted isoindolyl group, a substituted or unsubstituted pyridoindolyl group, a substituted or unsubstituted indazolyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted purinyl group, a substituted or unsubstituted benzoquinolinyl group, a substituted or unsubstituted phthalazinyl group, a substituted or unsubstituted naphthyridinyl group, a substituted or unsubstituted quinoxalinyl group, a substituted or unsubstituted quinazolinyl group, a substituted or unsubstituted cinnolinyl group, a substituted or unsubstituted phenazinyl group, a substituted or unsubstituted furanyl group, a substituted or unsubstituted benzofuranyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted thiophenyl group, a substituted or unsubstituted benzothiophenyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted thiazolyl group, a substituted or unsubstituted isothiazolyl group, a substituted or unsubstituted benzothiazolyl group, a substituted or unsubstituted oxazolyl group, a substituted or unsubstituted isoxazolyl group, a substituted or unsubstituted oxadiazolyl group, a substituted or unsubstituted triazolyl group, and a substituted or unsubstituted tetrazolyl group.


3. The heterocyclic compound of claim 1, wherein Ar_1 and Ar_2 are each independently one of the groups represented by Formulae 2A to 2J below;




Formula 2A




Formula 2B




Formula 2C




Formula 2D




Formula 2E




Formula 2F




Formula 2G



Formula 2H



Formula 2I



Formula 2J

wherein, in Formulae 2A to 2J, Z_{11} , Z_{12} , Z_{13} , and Z_{14} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyridinyl group, and a substituted or unsubstituted quinolyl group;


a plurality of Z_{11} , Z_{12} , Z_{13} , and Z_{14} are identical to or different from each other;

r , is an integer from 1 to 9;

s is an integer from 1 to 5;

* indicates a binding site.

4. The heterocyclic compound of claim 3, wherein Ar_1 and Ar_2 are each independently one of the groups represented by Formulae 3A to 3S below:



Formula 3A

Formula 3B

Formula 3C

Formula 3D

Formula 3E

Formula 3F


Formula 3G

Formula 3H

Formula 3I

Formula 3J

-continued



Formula 3K

Formula 3L

Formula 3M

Formula 3N


Formula 3O

Formula 3P

Formula 3Q

Formula 3R

-continued




Formula 3S


wherein, in Formulae 3A to 3S, * indicates a binding site.

5. The heterocyclic compound of claim 1, wherein A and B are each independently one of a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted dibenzothiophenylylene group, a substituted or unsubstituted pyridinylylene group, a substituted or unsubstituted pyrimidinylylene group, a substituted or unsubstituted triazininylylene group, a substituted or unsubstituted anthrylylene group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenylylene group, a substituted or unsubstituted chrysenylylene group, a substituted or unsubstituted perylenylylene group, a substituted or unsubstituted spiro-fluorenylylene group, a substituted or unsubstituted furanylylene group, a substituted or unsubstituted thiophenylylene group, and a substituted or unsubstituted oxadiazolylylene group.


6. The heterocyclic compound of claim 1, wherein Ar₁ and Ar₂ are each independently one of the groups represented by Formulae 4A to 4L below;




Formula 4A




Formula 4B



Formula 4C




Formula 4D




Formula 4E


-continued




Formula 4F




Formula 4G




Formula 4H




Formula 4I



Formula 4J



Formula 4K




Formula 4L

wherein, in Formulae 4A to 4L, Z₂₁, Z₂₂, and Z₂₃ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, and a substituted or unsubstituted pyridinyl group, wherein a plurality of Z₂₁, Z₂₂, and Z₂₃ are identical to or different from each other;


v, w, and x are an integer from 1 to 4; and

* and *' indicate binding sites.

7. The heterocyclic compound of claim 6, wherein A and B are each independently one of the groups represented by Formulae 5A to 5R below:




-continued



Formula 5R

wherein, in Formulae 5A to 5R, * and *' indicate binding sites.

8. The heterocyclic compound of claim 1, wherein R_1 to R_{13} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyranyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzoimidazolyl group, a substituted or unsubstituted benzooxazolyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted pyrazolyl group, a $N(Q_1)(Q_2)$ group, and a group represented by Formula 1B below, wherein Q_1 and Q_2 are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group:



Formula 1B

R_{14} and R_{15} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a

substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, and a substituted or unsubstituted pyridinyl group, and a substituted or unsubstituted phenanthrenyl group;


Ar_1 and Ar_2 are each independently one of a substituted or unsubstituted phenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted perylenyl group, a substituted or unsubstituted spiro-fluorenyl group, and a substituted or unsubstituted oxadiazolyl group;

A and B are each independently one of a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted carbazolylene group, a substituted or unsubstituted dibenzothiophenylene group, a substituted or unsubstituted pyridinylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted spiro-fluorenylene group, a substituted or unsubstituted furanylene group, a substituted or unsubstituted thiophenylene group, and a substituted or unsubstituted oxadiazolylene group; and

a is an integer from 0 to 2, and b is an integer from 0 to 2, wherein if a is 2, the two A are identical to or different from each other, and if b is 2, the two B are identical to or different from each other.










9. The heterocyclic compound of claim 8, wherein R_1 to R_{13} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a group represented by Formula 1B above, and groups represented by Formulae 6A to 6L below; and

R_{14} and R_{15} are each independently one of a hydrogen atom, a deuterium atom, and groups represented by Formulae 6A to 6L below:



Formula 6A

-continued



Formula 6B

Formula 6C

Formula 6D

Formula 6E

Formula 6F



Formula 6G

Formula 6H

Formula 6I

Formula 6J

-continued



Formula 6K

Formula 6L

wherein, in Formulae 6A to 6L, Z_{31} , Z_{32} , and Z_{33} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted quinolyl group, and a $N(Q_{11})(Q_{12})$ group;

Q_{11} and Q_{12} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group;

a plurality of Z_{31} and Z_{32} are identical to or different from each other;


p is an integer from 1 to 9;

q is an integer from 1 to 5; and

* indicates a binding site.

10. The heterocyclic compound of claim 8, wherein R_1 to R_{13} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted tert-butyl group, a cyano group, $-CD_3$, $-CF_3$, a group represented by Formula 1B above, and groups represented by Formulae 7A to 7S below; and

R_{14} and R_{15} are each independently one of a hydrogen atom, a deuterium atom, and groups represented by Formulae 7A to 7H below:



Formula 7A

Formula 7B

Formula 7C

Formula 7D

Formula 7E


Formula 7F

Formula 7G

Formula 7H

Formula 7I

-continued



Formula 7J

Formula 7K

Formula 7L


Formula 7M

Formula 7N


Formula 7O

Formula 7P


-continued



Formula 7Q




Formula 7R




Formula 7S

wherein, in Formulae 7A to 7S, * indicates a binding site.


11. A heterocyclic compound represented by Formula 1C, 1D, or 1E below:



Formula 1C



Formula 1D



Formula 1E

wherein, in Formula 1C, 1D, and 1E, R₁ to R₁₃ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a nitro group, a carboxyl group, a substituted or unsub-

stituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted biphenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyranlyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted pyrazinyl group, a substituted or unsubstituted pyridazinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted quinolinyl group, a substituted or unsubstituted benzimidazolyl group, a substituted or unsubstituted benzoxazolyl group, a substituted or unsubstituted pentalenyl group, a substituted or unsubstituted indenyl group, a substituted or unsubstituted pyrrolyl group, a substituted or unsubstituted imidazolyl group, a substituted or unsubstituted pyrazolyl group, and a N(Q₁)(Q₂) group, wherein Q₁ and Q₂ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a carboxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group;

R₁₄ and R₁₅ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, and a substituted or unsubstituted phenanthrenyl group;


Ar₂ are each independently one of a substituted or unsubstituted phenyl group, a substituted or unsubstituted terphenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted spiro-fluorenyl group, a substituted or unsubstituted dibenzothiophenyl group, a substituted or unsubstituted dibenzofuranyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted pyrimidinyl group, a substituted or unsubstituted triazinyl group, a substituted or unsubstituted chrysenyl group, a substituted or unsubstituted perylenyl group, a

substituted or unsubstituted spiro-fluorenyl group, and a substituted or unsubstituted oxadiazolyl group;


A and B are each independently one of a substituted or unsubstituted phenylene group, a substituted or unsubstituted naphthylene group, a substituted or unsubstituted fluorenylene group, a substituted or unsubstituted carbazolyene group, a substituted or unsubstituted dibenzothiophenylene group, a substituted or unsubstituted pyridinylene group, a substituted or unsubstituted pyrimidinylene group, a substituted or unsubstituted triazinylene group, a substituted or unsubstituted anthrylene group, a substituted or unsubstituted phenanthrenyl group, a substituted or unsubstituted pyrenylene group, a substituted or unsubstituted chrysenylene group, a substituted or unsubstituted perylenylene group, a substituted or unsubstituted spiro-fluorenylene group, a substituted or unsubstituted furanylene group, a substituted or unsubstituted thiophenylene group, and a substituted or unsubstituted oxadiazolylene group; and

a is an integer from 0 to 2, and b is an integer from 0 to 2, wherein if a is 2, the two A are identical to or different from each other, and if b is 2, the two B are identical to or different from each other.


12. The heterocyclic compound of claim 11, wherein Ar_1 and Ar_2 are each independently one of the groups represented by Formulae 2A to 2J below:




Formula 2A




Formula 2B




Formula 2C




Formula 2D




Formula 2E




Formula 2F




Formula 2G



Formula 2H



Formula 2I



Formula 2J

wherein, in Formulae 2A to 2J, Z_{11} , Z_{12} , Z_{13} , and Z_{14} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyridinyl group, and a substituted or unsubstituted quinolyl group;


a plurality of Z_{11} , Z_{12} , Z_{13} , and Z_{14} are identical to or different from each other;

r is an integer from 1 to 9;

s, t and u are an integer from 1 to 4; and

* indicates a binding site.

13. The heterocyclic compound of claim 11, wherein Ar₁ and Ar₂ are each independently one of the groups represented by Formulae 3A to 3S below:



Formula 3A

Formula 3B

Formula 3C

Formula 3D

Formula 3E

Formula 3F


Formula 3G

Formula 3H

Formula 3I

Formula 3J

-continued



Formula 3K

Formula 3L

Formula 3M

Formula 3N


Formula 3O

Formula 3P

Formula 3Q

Formula 3R


-continued




Formula 3S

wherein, in Formulae 3A to 3S, * indicates a binding site.


14. The heterocyclic compound of claim 11, wherein A and B are each independently one of the groups represented by Formulae 4A to 4L below:




Formula 4A




Formula 4B




Formula 4C




Formula 4D



Formula 4E




Formula 4F




Formula 4G

-continued



Formula 4H




Formula 4I



Formula 4J



Formula 4K




Formula 4L

wherein, in Formulae 4A to 4L, Z_{21} , Z_{22} , and Z_{23} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, and a substituted or unsubstituted pyridinyl group, wherein a plurality of Z_{21} , Z_{22} , and Z_{23} are identical to or different from each other;


v , w , and x are an integer from 1 to 4; and

* and *f indicate binding sites.

15. The heterocyclic compound of claim 11, wherein A and B are each independently one of the groups represented by Formulae 5A to 5R below:




Formula 5A




Formula 5B

pyl group, a substituted or unsubstituted butyl group, and groups represented by Formulae 6A to 6L below; and


R_{14} and R_{15} are each independently one of a hydrogen atom, a deuterium atom, and groups represented by Formulae 6A to 6L below:




Formula 6A




Formula 6B




Formula 6C




Formula 6D




Formula 6E



Formula 6F




Formula 6G




Formula 6H


-continued




Formula 6I



Formula 6J



Formula 6K



Formula 6L

wherein, in Formulae 6A to 6L, Z_{31} , Z_{32} , and Z_{33} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a cyano group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted methoxy group, a substituted or unsubstituted ethoxy group, a substituted or unsubstituted ethenyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted pyridinyl group, a substituted or unsubstituted carbazolyl group, a substituted or unsubstituted quinolyl group, and a $N(Q_{11})(Q_{12})$ group;

Q_{11} and Q_{12} are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a hydroxyl group, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted butyl group, a substituted or unsubstituted pentyl group, a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthryl group, a substituted or unsubstituted fluorenyl group, a substituted or unsubstituted carbazolyl group, and a substituted or unsubstituted pyridinyl group;

a plurality of Z_{31} and Z_{32} are identical to or different from each other;


p is an integer from 1 to 9;

q is an integer from 1 to 5; and


* indicates a binding site.

17. The heterocyclic compound of claim 11, wherein R₁ to R₁₃ are each independently one of a hydrogen atom, a deuterium atom, a halogen atom, a substituted or unsubstituted methyl group, a substituted or unsubstituted ethyl group, a substituted or unsubstituted propyl group, a substituted or unsubstituted tert-butyl group, a cyano group, —CD₃, —CF₃, and groups represented by Formulae 7A to 7S below; and


R₁₄ and R₁₅ are each independently one of a hydrogen atom, a deuterium atom, and groups represented by Formulae 7A to 7H below:




Formula 7A




Formula 7B




Formula 7C




Formula 7D



Formula 7E




Formula 7F




Formula 7G


-continued




Formula 7H




Formula 7I




Formula 7J



Formula 7K




Formula 7L




Formula 7M


-continued



Formula 7N




Formula 7O




Formula 7P


-continued



Formula 7Q




Formula 7R




Formula 7S

wherein, in Formulae 7A to 7S, * indicates a binding site.


18. The heterocyclic compound of claim 1, wherein the heterocyclic compound of Formula 1A above is one of the compounds represented by Formulae 1 to 7S below:




1




2




3



4













5




6











-continued










-continued



-continued




-continued




-continued


44




45




46



47




48




-continued


49




50



51




52





-continued

53




54


55




56



57




58




-continued


59




60



61




62




-continued


63



64





65




66

67




-continued


68




69




70




71



72




73




-continued

74




75




19. The heterocyclic compound of claim 1, wherein the heterocyclic compound of Formula 1A above is one of the compounds represented by Formulae 8, 16, 19, 29, 39, 47, 56, and 62 below:


Compound 8




Compound 16



Compound 19




Compound 29




-continued


Compound 39




Compound 47



Compound 56



Compound 62



20. An organic light-emitting device comprising:
 a first electrode;
 a second electrode disposed opposite to the first electrode;
 and
 an organic layer disposed between the first electrode and the second electrode,
 wherein the organic layer comprises at least one layer, and comprises at least one of the heterocyclic compound of claim 1.

21. The organic light-emitting device of claim 20, wherein the organic layer comprises at least one of a hole injection layer, a hole transport layer, a hole injection and transport layer having both hole injection and hole transport capabilities, an emission layer, an electron injection layer, an electron transport layer, and an electron injection and transport layer having both electron injection and electron transport capabilities.

22. The organic light-emitting device of claim 21, wherein at least one of the emission layer, the hole injection layer, the

hole transport layer, and the hole injection and transport layer comprises the heterocyclic compound.

23. The organic light-emitting device of claim **21**, wherein the organic layer comprises the emission layer, the emission layer comprises a host and a dopant, and the heterocyclic compound is a fluorescent host, a phosphorescent host, or a fluorescent dopant of the emission layer.

24. The organic light-emitting device of claim **22**, wherein the emission layer is a blue emission layer.

25. The organic light-emitting device of claim **21**, wherein the organic layer comprises the emission layer, the emission layer comprises a host and a phosphorescent dopant.

26. The organic light-emitting device of claim **21**, wherein at least one of the hole injection layer, the hole transport layer, and the hole injection and transport layer further comprises a charge generating material.

27. The organic light-emitting device of claim **21**, wherein the charge generating material is a p-type dopant.

28. The organic light-emitting device of claim **21**, wherein the organic layer comprises the electron transport layer, and the electron transport layer comprises an electron transporting organic compound and a metal-containing material.

29. The organic light-emitting device of claim **28**, wherein the metal-containing material comprises a lithium (Li) complex.

30. The organic light-emitting device of claim **20**, wherein at least one layer of the organic layer is formed using a wet process.

* * * * *

专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	US20130113367A1	公开(公告)日	2013-05-09
申请号	US13/444938	申请日	2012-04-12
[标]申请(专利权)人(译)	JUNG HYE JIN HWANG SEOK HWAN 金荣KOOK LIM JINØ 韩相HYUN 金秀YON SHIN DAE YUP		
申请(专利权)人(译)	JUNG , HYE-JIN HWANG , SEOK-HWAN KIM , YOUNG-KOOK LIM , JIN-O HAN , SANG-HYUN KIM , SOO-YON SHIN , DAE-YUP		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	JUNG HYE JIN HWANG SEOK HWAN KIM YOUNG KOOK LIM JIN O HAN SANG HYUN KIM SOO YON SHIN DAE YUP		
发明人	JUNG, HYE-JIN HWANG, SEOK-HWAN KIM, YOUNG-KOOK LIM, JIN-O HAN, SANG-HYUN KIM, SOO-YON SHIN, DAE-YUP		
IPC分类号	H05B33/14 C07D401/04 C07D403/04 C07D413/10 C07D401/12 C07D405/12 C07F7/02 C07D409/10 C07D209/94 C07D401/14		
CPC分类号	C07D401/14 C07D401/04 C07D401/12 C07D403/04 C07D209/94 C07D405/12 C07D409/04 C07D409 /12 C07D413/04 C07D405/04		
优先权	1020110114117 2011-11-03 KR		
其他公开文献	US8828559		
外部链接	Espacenet USPTO		

摘要(译)

由式1A表示的杂环化合物和包含该杂环化合物的有机发光装置：R1至R13中的至少一个是由下式1B表示的基团：其中R1至R15，Ar1，Ar2，A，B，a和b如说明书中所定义。有机发光装置可以包括含有杂环化合物的有机层，因此可以具有低驱动电压，高发光效率和长寿命特性。

