



US010115907B2

(12) **United States Patent**
Shin et al.

(10) **Patent No.: US 10,115,907 B2**
 (45) **Date of Patent: Oct. 30, 2018**

(54) **COMPOUND AND ORGANIC
 LIGHT-EMITTING DEVICE INCLUDING
 THE SAME**

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
 Yongin, Gyeonggi-Do (KR)

(72) Inventors: **Daeyup Shin**, Yongin (KR);
Youngkook Kim, Yongin (KR); **Jihwan**
Yoon, Yongin (KR)

(73) Assignee: **SAMSUNG DISPLAY CO., LTD.**,
 Yongin, Gyeonggi-do (KR)

(*) Notice: Subject to any disclaimer, the term of this
 patent is extended or adjusted under 35
 U.S.C. 154(b) by 417 days.

(21) Appl. No.: **14/736,375**

(22) Filed: **Jun. 11, 2015**

(65) **Prior Publication Data**

US 2016/0190465 A1 Jun. 30, 2016

(30) **Foreign Application Priority Data**

Dec. 31, 2014 (KR) 10-2014-0195956

(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 487/04 (2006.01)
 (Continued)

(52) **U.S. Cl.**
 CPC **H01L 51/0061** (2013.01); **C07D 487/04**
 (2013.01); **C07D 495/04** (2013.01);
 (Continued)

(58) **Field of Classification Search**
 CPC H01L 51/0061; H01L 51/0072; H01L
 51/0074; H01L 51/0065–51/0068;
 (Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

2007/0134515 A1 6/2007 Fukuoka et al.
 2009/0085468 A1* 4/2009 Funahashi C07C 211/54
 313/504

(Continued)

FOREIGN PATENT DOCUMENTS

KR 10-2007-0087028 A 8/2007
 KR 10-2008-0096733 A 11/2008

(Continued)

OTHER PUBLICATIONS

“Naphtho[1,2-b:5,6-b']dithiophene”, <http://www.sigmaaldrich.com/catalog/search?interfact=All&term=naththo%5B1%2C2-b%3A5%2C6>.

(Continued)


Primary Examiner — Ruiyun Zhang

(74) Attorney, Agent, or Firm — Lee & Morse, P.C.

(57) **ABSTRACT**

A compound, an organic light-emitting device, and a flat display apparatus, the compound being represented by any one selected from the following Formula 1, Formula 2, and Formula 3:

<Formula 1>




(Continued)

10


190
150
110

-continued

<Formula 2>



<Formula 3>



20 Claims, 1 Drawing Sheet

(51) **Int. Cl.**

C07D 495/04 (2006.01)
C09K 11/06 (2006.01)
H01L 51/50 (2006.01)

(52) **U.S. Cl.**

CPC **C09K 11/06** (2013.01); **H01L 51/0072**
 (2013.01); **H01L 51/0074** (2013.01); **C09K**
2211/1007 (2013.01); **C09K 2211/1014**
 (2013.01); **C09K 2211/1029** (2013.01); **C09K**

2211/1037 (2013.01); **C09K 2211/1044**
 (2013.01); **C09K 2211/1088** (2013.01); **C09K**
2211/1092 (2013.01); **C09K 2211/181**
 (2013.01); **H01L 51/5012** (2013.01); **H01L**
51/5076 (2013.01)

(58) **Field of Classification Search**

CPC . H01L 51/5012; C07D 487/04; C07D 495/04;
 C09K 11/06; C09K 2211/1007; C09K
 2211/1018; C09K 2211/1029; C09K
 2211/1033; C09K 2211/1037; C09K
 2211/1044; C09K 2211/1088; C09K
 2211/1092

USPC 428/690, 917; 257/E51.049, E51.051;
 313/504; 549/42

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

2011/0040107	A1	2/2011	Goto et al.	
2011/0068683	A1	3/2011	Kawamura et al.	
2014/0051865	A1*	2/2014	Takimiya	C07D 495/04 549/42
2015/0011780	A1	1/2015	Takimiya et al.	
2017/0229654	A1*	8/2017	Gao	H01L 51/0035

FOREIGN PATENT DOCUMENTS

KR	10-2010-0123735	A	11/2010
WO	WO 2009/128559	A1	10/2009
WO	WO 2013/121664	A1	8/2013

OTHER PUBLICATIONS

Yoshihito Kunugi, et al., "Single Crystal Organic Field-effect Transistors Based on Dinaphthothienothiophene Isomers", Journal of Photopolymer Science and Technology, vol. 23, No. 3(2010) 323-326.

* cited by examiner

10

190
150
110

1

COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

Korean Patent Application No. 10-2014-0195956, filed on Dec. 31, 2014, in the Korean Intellectual Property Office, and entitled: "Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

Embodiments relate to a compound and an organic light-emitting device including the same.

2. Description of the Related Art



Organic light-emitting devices are self-emission devices that have wide viewing angles, high contrast ratios, short response times, and excellent brightness, driving voltage, and response speed characteristics, and produce full-color images.

The organic light-emitting device may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, are recombined in the emission layer to produce excitons. These excitons change from an excited state to a ground state, thereby generating light.

SUMMARY

Embodiments are directed to a compound and an organic light-emitting device including the same.


The embodiments may be realized by providing a compound represented by any one selected from the following Formula 1, Formula 2, and Formula 3:



2

-continued

<Formula 3>




wherein in Formulae 1, 2, and 3, X and Y are each independently S, O, NR₁, or CR₂R₃, A to F are each independently hydrogen, deuterium, or a substituted or unsubstituted C₆-C₆₀ aryl moiety fused to a corresponding backbone, Ar₁ to Ar₄ and R₁ to R₃ are each independently selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, at least one substituent of the substituted C₁-C₆₀ alkyl group, substituted C₆-C₆₀ aryl group, substituted C₂-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group; a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇); a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₂-C₆₀ alkyl group, a


3

C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), and —B(Q₂₆)(Q₂₇); and —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇), wherein Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ are each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.


The compound represented by one of Formulae 1 to 3 may be represented by one of the following Formulae 4 to 6:



<Formula 4>



<Formula 5>




<Formula 6>

wherein, in Formulae 4 to 6, Ar₁ to Ar₄, X, and Y are defined the same as Ar₁ to Ar₄, X, and Y of Formulae 1 to 3.


The compound represented by one of Formulae 1 to 3 may be represented by one of the following Formulae 7 to 9:

4


<Formula 7>



<Formula 8>




<Formula 9>




wherein, in Formulae 7 to 9, Ar₁ to Ar₄, X, and Y are defined the same as Ar₁ to Ar₄, X, and Y of Formulae 1 to 3.

The compound represented by one of Formulae 1 to 3 may be represented by one of the following 10 to 12:

<Formula 10>




<Formula 11>



5

-continued

<Formula 12>





wherein, in Formulae 10 to 12 Ar₁ to Ar₄, X, and Y are defined the same as Ar₁ to Ar₄, X, and Y of Formulae 1 to 3.

R₁ to R₃ may each independently be a substituted or unsubstituted C₁-C₂₀ alkyl group or a substituted or unsubstituted C₆-C₆₀ aryl group.

R₁ to R₃ may each independently be a methyl group or a phenyl group.

Ar₁ to Ar₄ may each independently be a substituted or unsubstituted C₆-C₂₀ aryl group or a substituted or unsubstituted C₁-C₂₀ heteroaryl group.


Ar₁ to Ar₄ may each independently be a group represented by one of the following Formulae 2a and 2b:



wherein, in Formulae 2a and 2b, Z₁ may be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, a substituted or unsubstituted a monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group; p may be an integer of 1 to 7, * indicates a binding site to a neighboring atom, when p is 2 or more, each Z₁ may be identical or different from one another.



Ar₁ to Ar₄ may each independently be a group represented by one of the following Formulae 3a to 3k, in which * indicates a binding site to a neighboring atom:

6





7

-continued



The compound represented by one of Formula 1, Formula 2, and Formula 3 may be one of the following compounds:

**8**

-continued

3j

5

10

15

3k

20

25

30

35

40




45

50

55


60

65








9

-continued




10

-continued




11

-continued



12

-continued



13

The embodiments may be realized by providing an organic light-emitting device including a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes the compound according to an embodiment.

The organic layer may be formed by a wet process.

The first electrode may be an anode, the second electrode may be a cathode, and the organic layer may include a hole transport region between the first electrode and the emission layer, the hole transport region including at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and an electron transport region between the emission layer and the second electrode, the electron transport region including at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

The compound may be included in the emission layer.


The hole transport region may include a charge-generating material.

The electron transport region may include a metal complex.

The metal complex may be a lithium complex.

The metal complex may be a lithium quinolate.

The metal complex may be represented by the following Formula ET-D2:



The embodiments may be realized by providing a flat display apparatus including a thin film transistor, the thin film transistor including a source electrode and a drain electrode; and the organic light-emitting device according to an embodiment, wherein the first electrode of the organic light-emitting device is electrically connected to the source electrode or the drain electrode of the thin film transistor.

BRIEF DESCRIPTION OF THE DRAWING

Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawing in which:

FIG. 1 illustrates a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION


Example embodiments will now be described more fully hereinafter with reference to the accompanying drawing; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.

14


In the drawing FIGURES, the dimensions of layers and regions may be exaggerated for clarity of illustration. Like reference numerals refer to like elements throughout.

According to an aspect, provided is a compound represented by any one selected from the following Formula 1, Formula 2, and Formula 3:


<Formula 1>



<Formula 2>



<Formula 3>



In Formulae 1, 2, and 3,

X and Y may each independently be, e.g., S, O, NR₁, or CR₂R₃,

A to F may each independently be or include, e.g., hydrogen, deuterium, or a substituted or unsubstituted C₆-C₆₀ aryl moiety fused to a corresponding backbone, e.g., fused to the core moiety of Formulae 1 to 3. For example, A to F may not necessarily be a ring, and may represent hydrogen bound to the core moiety of the Formulae at the connection points with A to F.

Ar₁ to Ar₄ and R₁ to R₃ may each independently be selected from or include, e.g., a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

At least one substituent of the substituted C₁-C₆₀ alkyl group, substituted C₆-C₆₀ aryl group, substituted C₂-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic

acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₂-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅) and —B(Q₂₆)(Q₂₇); and

—N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇),

wherein Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may each independently be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

Compounds represented by Formulae 1 to 3 according to an embodiment may have a rigid main luminescent structure, which may be enriched with electrons due to a non-covalent electron pair of a heteroatom. Due to effects of a heavy atom, spin-orbit coupling effects may be improved,

leading to high luminescent efficiency. Also, luminescent wavelengths of the compounds may be controllable by adding or removing ring structures corresponding to A to F.

The compounds represented by Formulae 1 to 3 according to an embodiment may have a high glass transition temperature (T_g) or a high melting point. Accordingly, a heat resistance against Joule heat that may occur within an organic layer, or between the organic layer and a metal electrode during emission, and durability at high temperature increase may be improved.

An organic light-emitting device manufactured by using a compound according to an embodiment may have high durability during preservation and driving.


Hereinafter, substituents of Formulae 1, 2, and 3 will be described in detail.

In an implementation, A to F in Formulae 1, 2 and 3 may each independently be or include, e.g., a substituted or unsubstituted C₆-C₆₀ aryl moiety fused to a corresponding backbone.


In an implementation, A to F in Formulae 1, 2 and 3 may not be the aryl moiety fused to a corresponding backbone. For example, A to F may each independently be hydrogen or deuterium.

For example, the compound represented by one of Formulae 1 to 3 may be represented by one of the following Formulae 4 to 6.


<Formula 4>



<Formula 5>



<Formula 6>




In an implementation, when A to F in Formulae 1 to 3 include the fused C₆-C₆₀ aryl moiety, A to F may each independently be, e.g., a phenyl moiety, or a naphthyl moiety.


In an implementation, the compound represented by one of Formulae 1 to 3 may be represented by one of the following Formulae 7 to 9.

17


<Formula 7>



<Formula 8>




<Formula 9>




In an implementation, the compound represented by one of Formulae 1 to 3 may be represented by one of the following Formulae 10 to 12.

<Formula 10>




<Formula 11>



18

-continued

<Formula 12>




In an implementation, R_1 to R_3 may each independently be or include, e.g., a substituted or unsubstituted C_1 - C_{20} alkyl group or a substituted or unsubstituted C_6 - C_{60} aryl group. For example, R_1 to R_3 may each independently be a methyl group or a phenyl group.


In an implementation, Ar_1 to Ar_4 may each independently be or include, e.g., a substituted or unsubstituted C_6 - C_{20} aryl group or a substituted or unsubstituted C_1 - C_{20} heteroaryl group.

In an implementation, Ar_1 to Ar_4 may each independently be a group represented by one of the following Formulae 2a and 2b.

2a



2b



In Formulae 2a and 2b,










Z_1 may be selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_2 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_2 - C_{60} heteroaryl group, a substituted or unsubstituted a monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group;

p may be an integer selected from 1 to 7; and * indicates a binding site;

when p is 2 or more, each Z_1 may be identical or different from one another.

In an implementation, Ar_1 to Ar_4 may each independently be a group represented by one of the following Formulae 3a to 3k, in which * represents a binding site to a neighboring atom.

19



20

-continued

3a

3j

5

3b

10

15

3c

3k

20

3d

25

3e

30

In an implementation, the compound represented by one of Formulae 1 to 3 may be one of the following compounds.

3f

35

3g

40

45

3h



50

3i

55


60

65



21

-continued



5


10

15

20

22

-continued




25

30

35

40

45





50

55


60

65



23

-continued

**24**


-continued

5

10

15

20



25

30

35

40






45

50

55






60

65



25

-continued



26

The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of an organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1.

In FIG. 1, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or transparent plastic substrate, each with excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode on the substrate. When the first electrode 110 is an anode, the material for the first electrode 110 may be selected from materials with a high work function to make holes be easily injected. The first electrode 110 may be a reflective electrode or a transmissive electrode. The material for the first electrode 110 may be a transparent and highly conductive material, and examples of such a material are indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), and zinc oxide (ZnO). When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode, at least one of magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag) may be used.

The first electrode 110 may have a single-layer structure, or a multi-layer structure including two or more layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

An organic layer 150 may be disposed on the first electrode 110. The organic layer 150 may include an emission layer.

The organic layer 150 may further include a hole transport region between the first electrode and the emission layer, and/or an electron transport region between the emission layer and the second electrode.

The hole transport region may include at least one selected from a hole transport layer (HTL), a hole injection layer (HIL), a buffer layer, and an electron blocking layer. The electron transport region may include at least one selected from a hole blocking layer (HBL), an electron transport layer (ETL), and an electron injection layer (EIL).

The hole transport region may have a single-layered structure formed of a single material, a single-layered structure formed of a plurality of different materials, or a multi-layered structure having a plurality of layers formed of a plurality of different materials.

For example, the hole transport region may have a single-layered structure formed of a plurality of different materials, or a structure of hole injection layer/hole transport layer, a structure of hole injection layer/hole transport layer/buffer layer, a structure of hole injection layer/buffer layer, a structure of hole transport layer/buffer layer, or a structure of hole injection layer/hole transport layer/electron blocking layer, wherein layers of each structure are sequentially stacked from the first electrode 110 in this stated order.

27


When the hole transport region includes a hole injection layer, the hole injection layer may be formed on the first electrode **110** by using various methods, such as vacuum deposition, spin coating casting, a Langmuir-Blodgett (LB) method, ink-jet printing, laser-printing, or laser-induced thermal imaging.

When a hole injection layer is formed by vacuum deposition, for example, the vacuum deposition may be performed at a temperature of a deposition temperature of about 100° C. to about 500° C., at a vacuum degree of about 10^{-8} to about 10^{-3} torr, and/or at a deposition rate of about 0.01 to about 100 Å/sec in consideration of a compound for a hole injection layer to be deposited, and the structure of a hole injection layer to be formed.

When a hole injection layer is formed by spin coating, the spin coating may be performed at a coating rate of about 2,000 rpm to about 5,000 rpm, and at a temperature of about 80° C. to 200° C. in consideration of a compound for a hole injection layer to be deposited, and the structure of a hole injection layer to be formed.





When the hole transport region includes a hole transport layer, the hole transport layer may be formed on the first electrode **110** or the hole injection layer by using various methods, e.g., vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When the hole transport layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole transport layer may be the same as the deposition and coating conditions for the hole injection layer.

The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB, β -NPB, TPD, Spiro-TPD, Spiro-NPB, α -NPB, TAPC, HMTDPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), (polyaniline)/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, and a compound represented by Formula 202 below.




28

-continued




29


-continued





TPD



Spiro-TPD




Spiro-NPB

 α -NPB


TAPC

30


-continued



HMTPD



<Formula 201>



<Formula 202>

²⁵ In Formulae 201 and 202,

xa1 to xa4 may each independently be selected from 0, 1, 2, and 3;

xa5 may be selected from 1, 2, 3, 4, and 5; and

R₂₀₁ to R₂₀₄ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

In an implementation, in Formulae 201 and 202,

L_{201} to L_{205} may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylenylene group, a pyrenylene group, a chrysenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinoxalinylene group, a carbazolylenylene group, and a triazinylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorene group, a dibenzofluorene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolyene group, and a triazinylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl

31

group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

xa1 to xa4 may be each independently 0, 1, or 2;


xa5 may be 1, 2, or 3;

R₂₀₁ to R₂₀₄ may each independently be selected from:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an azulenyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group.


The compound represented by Formula 201 may be represented by Formula 201A:



For example, the compound represented by Formula 201 may be represented by Formula 201A-1 below.


32

<Formula 201A-1>



For example, the compound represented by Formula 202 may be represented by Formula 202A below.

<Formula 202A>



L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ in Formulae 201A, 201A-1, and 202A are already described in detail above, and R₂₁₁ may be understood by referring to the description provided herein in connection with R₂₀₃, and R₂₁₃ to R₂₁₆ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

For example, in Formulae 201A, 201A-1, and 202A,

L₂₀₁ to L₂₀₃ may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a quinolinylene group, an isoquinolinylene group, a quinoxalinylene group, a quinazolinylene group, a carbazolylenylene group, and a triazinylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, a chrysenylene group, a pyridinylene group, a pyrazinylene


35

carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and


xa5 may be 1 or 2.

R_{213} and R_{214} in Formulae 201A, and 201A-1 may bind to each other to form a saturated or unsaturated ring.


The compound represented by Formula 201, and/or the compound represented by Formula 202 may include one of compounds HT1 to HT20 illustrated below.




HT1



HT2



HT3



HT4


36

-continued

37

-continued

HT5



38

-continued

HT7

5

10

15

20

25

30

35

40

HT6


45

50


55

60

65




HT8



39

-continued

HT9



40

-continued

HT11


5

10

15

20

25



30

HT12


35

40

HT10

45

50





55

HT13





60

65






41

-continued



42

-continued




A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes both a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or heterogeneously dispersed in the hole transport region.

The charge-generation material may be, e.g., a p-dopant. The p-dopant may be one selected from, e.g., a quinone

43

derivative, a metal oxide, and a cyano group-containing compound. Examples of the p-dopant may include a quinone derivative, such as tetracyanoquinonodimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide, and Compound HT-D1 illustrated below.



<Compound HT-D1>

<F4-TCNQ>

The hole transport region may further include a buffer layer, in addition to an electron blocking layer, a hole injection layer, and a hole transport layer. Since the buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, light-emission efficiency of a formed organic light-emitting device may be improved. For use as a material included in the buffer layer, materials that are included in the hole transport region may be used. The electron blocking layer prevents injection of electrons from the electron transport region.


An emission layer may be formed on the first electrode 110 or the hole transport region by using various methods, e.g., vacuum deposition, spin coating, casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When an emission layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the emission may be the same as those for the hole injection layer.

When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub pixel. In some embodiments, the emission layer may have a stacked structure of a red emission layer, a green emission layer, and a blue emission layer, or may include a red-light emission material, a green-light emission material, and a blue-light emission material, which are mixed with each other in a single layer, to emit white light.


The emission layer may include a host and a dopant.

The host may include, e.g., at least one selected from TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, and TCP.


44




TPBi




TBADN



ADN




CBP




CDBP

45

-continued



In an implementation, the host may include a compound represented by Formula 301 below.



<Formula 301>

In Formula 301,

Ar_{301} may be selected from:

a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q_{301})(Q_{302})(Q_{303}) (Q_{301} to Q_{303} may be each independently selected from a hydrogen, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_6 - C_{60} aryl group, and a C_1 - C_{60} heteroaryl group);

a description of L_{301} may be understood by referring to the description provided in connection with L_{201} .

R_{301} may be selected from:

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group;
a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl

46

group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

xb1 may be selected from 0, 1, 2, and 3;

xb2 may be selected from 1, 2, 3, and 4.

In Formula 301,

L_{301} may be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, and a chrysenylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a pyrenylene group, and a chrysenylene group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

R_{301} may be selected from:

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-


47

fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group;

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group; and

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, and a chrysenyl group.



For example, the host may include a compound represented by Formula 301A below:



[(L₃₀₁)_{xb1}—R₃₀₁]_{xb2}







Descriptions of substituents of Formula 301A may be understood by referring to the descriptions provided herein.

In an implementation, the compound represented by Formula 301 may include at least one of Compounds H1 to H42.




48

-continued




49

-continued




50

-continued




51

-continued



52


-continued



53

-continued

H31




5

10

15


H32



20

25


H33



35

40

H34



50

55


60

65


54

-continued


H35



H36




H37



55


-continued




H38

56


-continued




H41



H39



H42



H40

The dopant may include the compound represented by one of Formulae 1 to 3 according to an embodiment.

An amount of the dopant in the emission layer may be, e.g., about 0.01 to about 15 parts by weight, based on 100 parts by weight of the host.

A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.



Then, an electron transport region may be disposed on the emission layer.

The electron transport region may include, e.g., at least one selected from a hole blocking layer, an electron transport layer (ETL), and an electron injection layer.

When the electron transport region includes a hole blocking layer, the hole blocking layer may be formed on the emission layer by using various methods, e.g., vacuum deposition, spin coating casting, a Langmuir-Blodgett (LB) method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When the hole blocking layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the hole blocking layer may be determined by referring to the deposition and coating conditions for the hole injection layer.

The hole blocking layer may include, e.g., at least one of BCP and Bphen.

57




A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, e.g., about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within these ranges, the hole blocking layer may have excellent hole blocking characteristics without a substantial increase in driving voltage.

For example, the electron transport region may have a structure of electron transport layer/electron injection layer or a structure of hole blocking layer/electron transport layer/electron injection layer, wherein layers of each structure are sequentially stacked from the emission layer in the stated order.




According to an embodiment, the organic layer 150 of the organic light-emitting device may include an electron transport region disposed between the emission layer and the second electrode 190, and the electron transport region may include an electron transport layer. The electron transport layer may include a plurality of layers. For example, the electron transport layer may include a first electron transport layer and a second electron transport layer.

The electron transport layer may further include at least one selected from BCP, Bphen, Alq₃, Balq, TAZ, and NTAZ.




58

-continued



In an implementation, the electron transport layer may include at least one compound selected from a compound represented by Formula 601 and a compound represented by Formula 602 illustrated below.



<Formula 601>

In Formula 601,

Ar₆₀₁ may be selected from:

a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene; and

a naphthalene, a heptalene, a fluorene, a spiro-fluorene, a benzofluorene, a dibenzofluorene, a phenalene, a phenanthrene, an anthracene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a naphthacene, a picene, a perylene, a pentaphene, and an indenoanthracene, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₃-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, monovalent non-aromatic condensed heteropolycyclic group, and —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃) (Q₃₀₁ to Q₃₀₃ may be each independently selected from a hydrogen,

59

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group).

Descriptions of L₆₀₁ may be understood by referring to the description provided in connection with L₂₀₃.

E₆₀₁ may be selected from:

a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthroli-
n-1-yl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-1-yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran-1-yl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocar-
bazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group; and

a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthroli-
n-1-yl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-1-yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran-1-yl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocar-
bazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenale-
n-1-yl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl


60

group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthroli-1-yl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-1-yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran-1-yl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.

xe1 may be selected from 0, 1, 2, and 3; and

xe2 may be selected from 1, 2, 3, and 4.

<Formula 602>



In Formula 602,

X₆₁₁ may be N or C-(L₆₁₁)_{xe611}-R₆₁₁, X₆₁₂ may be N or C-(L₆₁₂)_{xe612}-R₆₁₂, X₆₁₃ may be N or C-(L₆₁₃)_{xe613}-R₆₁₃, and at least one selected from X₆₁₁ to X₆₁₃ may be N;

L₆₁₁ to L₆₁₆ may be understood by referring to the description provided herein in connection with L₂₀₃;

R₆₁₁ and R₆₁₆ may each independently be selected from:

a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolynyl group, an isoquinolynyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and


a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolynyl group, an isoquinolynyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolynyl group, an isoquinolynyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, and a triazinyl group; and

xe611 to xe616 may each independently be selected from 0, 1, 2, and 3.

The compound represented by Formula 601 and/or the compound represented by Formula 602 may be selected from Compounds ET1 to ET15 illustrated below.

61

ET1



5


10

15

20

ET2

25



30


35

40

45

ET3

50



55


60

65

62

-continued

ET4




ET5

30

35

40

45




ET6

55




60

65






63

-continued


**64**

-continued




65

-continued




ET13

ET14



ET15




A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.


The metal-containing material may include a lithium (Li) complex. The Li complex may include, e.g., Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.

66

ET-D1



ET-D2



The electron transport region may include an electron injection layer that allows electrons to be easily provided from the second electrode **190**.

The electron injection layer may be formed on the electron transport layer by using various methods, e.g., vacuum deposition, spin coating casting, a LB method, ink-jet printing, laser-printing, or laser-induced thermal imaging. When an electron injection layer is formed by vacuum deposition or spin coating, deposition and coating conditions for the electron injection layer may be the same as those for the hole injection layer.

The electron injection layer may include at least one selected from, LiF, NaCl, CsF, Li₂O, BaO, and LiQ.

A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

The second electrode **190** may be disposed on the organic layer **150** having such a structure. The second electrode **190** may be a cathode which is an electron injection electrode, and in this regard, a material for the second electrode **190** may be selected from metal, an alloy, an electrically conductive compound, and a mixture thereof, which have a relatively low work function. Examples of the material for the second electrode **190** may include lithium (Li), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag). In an implementation, the material for forming the second electrode **190** may be ITO or IZO. The second electrode **190** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode.

In an implementation, an organic layer according to an embodiment may be formed by depositing the compound according to an embodiment, or may be formed by using a wet method in which the compound according to an embodiment is prepared in the form of solution and then the solution of the compound is used for coating.

An organic light-emitting device according to an embodiment may be used in various flat panel display apparatuses, such as a passive matrix organic light-emitting display apparatus or an active matrix organic light-emitting display apparatus. For example, when the organic light-emitting device is included in an active matrix organic light-emitting display apparatus, a first electrode disposed on a substrate

acts as a pixel and may be electrically connected to a source electrode or a drain electrode of a thin film transistor. In addition, the organic light-emitting device may be included in a flat panel display apparatus that emits light in opposite directions.

Hereinbefore, the organic light-emitting device has been described with reference to FIG. 1.

Hereinafter, definitions of substituents used herein will be presented (the number of carbon numbers used to restrict a substituent is not limited, and does not limit properties of the substituent, and unless defined otherwise, the definition of the substituent is consistent with a general definition thereof).

A C_1 - C_{60} alkyl group used herein refers to a linear or branched aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and detailed examples thereof are a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A C_1 - C_{60} alkylene group used herein refers to a divalent group having the same structure as the C_1 - C_{60} alkyl group.

A C_1 - C_{60} alkoxy group used herein refers to a monovalent group represented by $-OA_{101}$ (wherein A_{101} is the C_1 - C_{60} alkyl group), and detailed examples thereof are a methoxy group, an ethoxy group, and an isopropoxy group.

A C_2 - C_{60} alkenyl group used herein refers to a hydrocarbon group having at least one carbon double bond in the middle or at the terminal of the C_2 - C_{60} alkyl group, and detailed examples thereof are an ethenyl group, a propenyl group, and a butenyl group. A C_2 - C_{60} alkenylene group used herein refers to a divalent group having the same structure as the C_2 - C_{60} alkenyl group.

A C_2 - C_{60} alkynyl group used herein refers to a hydrocarbon group having one carbon triple bond in the middle or at the terminal of the C_2 - C_{60} alkyl group, and detailed examples thereof are an ethynyl group, and a propynyl group. A C_2 - C_{60} alkynylene group used herein refers to a divalent group having the same structure as the C_2 - C_{60} alkynyl group.

A C_3 - C_{10} cycloalkyl group used herein refers to a monovalent hydrocarbon monocyclic group having 3 to 10 carbon atoms, and detailed examples thereof are a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C_3 - C_{10} cycloalkylene group used herein refers to a divalent group having the same structure as the C_3 - C_{10} cycloalkyl group.

A C_1 - C_{10} heterocycloalkyl group used herein refers to a monovalent monocyclic group having at least one hetero atom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and detailed examples thereof are a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group. A C_2 - C_{10} heterocycloalkylene group used herein refers to a divalent group having the same structure as the C_2 - C_{10} heterocycloalkyl group.

A C_3 - C_{10} cycloalkenyl group used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one double bond in the ring thereof and does not have aromaticity, and detailed examples thereof are a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. A C_3 - C_{10} cycloalkenylene group used herein refers to a divalent group having the same structure as the C_3 - C_{10} cycloalkenyl group.

A C_2 - C_{10} heterocycloalkenyl group used herein refers to a monovalent monocyclic group that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, 2 to 10 carbon atoms, and at least one double bond in its ring. Detailed examples of the C_2 - C_{10} heterocycloalkenyl group

are a 2,3-hydrofuranlyl group and a 2,3-hydrothiophenyl group. A C_2 - C_{10} heterocycloalkenylene group used herein refers to a divalent group having the same structure as the C_2 - C_{10} heterocycloalkenyl group.

A C_6 - C_{60} aryl group used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C_6 - C_{60} arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Detailed examples of the C_6 - C_{60} aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C_6 - C_{60} aryl group and the C_6 - C_{60} arylene group each include two or more rings, the rings may be fused to each other.

A C_1 - C_{60} heteroaryl group used herein refers to a monovalent group having a carboncyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 2 to 60 carbon atoms. A C_1 - C_{60} heteroarylene group used herein refers to a divalent group having a carbocyclic aromatic system that has at least one hetero atom selected from N, O, P, and S as a ring-forming atom, and 2 to 60 carbon atoms. Examples of the C_1 - C_{60} heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C_1 - C_{60} heteroaryl group and the C_1 - C_{60} heteroarylene group each include two or more rings, the rings may be fused to each other.

A C_6 - C_{60} aryloxy group used herein indicates $-OA_{102}$ (wherein A_{102} is the C_6 - C_{60} aryl group), and a C_6 - C_{60} arylthio group indicates $-SA_{103}$ (wherein A_{103} is the C_6 - C_{60} aryl group).

A monovalent non-aromatic condensed polycyclic group used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other, only carbon atoms as a ring forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. A divalent non-aromatic condensed polycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

A monovalent non-aromatic condensed heteropolycyclic group used herein refers to a monovalent group (for example, having 2 to 60 carbon atoms) that has two or more rings condensed to each other, has a heteroatom selected from N, O, P, and S, other than carbon atoms, as a ring forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. A divalent non-aromatic condensed heteropolycyclic group used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

In the present specification, at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_2 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_2 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_2 - C_{10} heterocycloalkyl

group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_2 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), and —B(Q₁₆)(Q₁₇);

a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_2 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_2 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅) and —B(Q₂₆)(Q₂₇); and

—N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), and —B(Q₃₆)(Q₃₇).

Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ may be each independently selected from a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a

C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_2 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_2 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

For example, at least one substituent of the substituted C_3 - C_{10} cycloalkylene group, the substituted C_2 - C_{10} heterocycloalkylene group, the substituted C_3 - C_{10} cycloalkenylene group, the substituted C_2 - C_{10} heterocycloalkenylene group, the substituted C_6 - C_{60} arylene group, the substituted C_1 - C_{60} heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_2 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_2 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophe-nyl group, a benzocarbazolyl group, a dibenzocarbazolyl

group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{N}(\text{Q}_{11})(\text{Q}_{12})$, $-\text{Si}(\text{Q}_{13})(\text{Q}_{14})(\text{Q}_{15})$, and $-\text{B}(\text{Q}_{16})(\text{Q}_{17})$;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-
 10 yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran-
 15 yl group, a dibenzothiophenyl group, a dibenzocarbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-
 25 yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran-
 30 yl group, a dibenzothiophenyl group, a dibenzocarbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or

a salt thereof, a phosphoric acid or a salt thereof, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-
 35 yl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran-
 40 yl group, a dibenzothiophenyl group, a dibenzocarbazolyl group, a benzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{Si}(\text{Q}_{23})(\text{Q}_{24})(\text{Q}_{25})$, and $-\text{B}(\text{Q}_{26})(\text{Q}_{27})$; and
 $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{Si}(\text{Q}_{33})(\text{Q}_{34})(\text{Q}_{35})$, and $-\text{B}(\text{Q}_{36})(\text{Q}_{37})$;

Q_{11} to Q_{17} , Q_{21} to Q_{27} , and Q_{31} to Q_{37} may be each independently selected from a hydrogen, a deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a furinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolynyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzoimidazolyl group, a benzofuran-
 60 yl group, a benzothiophenyl group, an isobenzothiazolyl

73

group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group.


The term "Ph" used herein refers to a phenyl group, the term "Me" used herein refers to a methyl group, the term "Et" used herein refers to an ethyl group, and the term "ter-Bu" or "Bu" used herein refers to a tert-butyl group.

Hereinafter, an organic light-emitting device according to an embodiment will be described in detail with reference to Synthesis Examples and Examples.

The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.


Synthesis Example

Synthesis of Compound A



74

-continued



Synthesis of Intermediate A-1

13.8 g (100 mmol) of 3-nitroaniline was dissolved in 300 mL of DMSO, and then, at ambient temperature, 23 g (101 mmol) of N-Iodosuccinimide was slowly added dropwise thereto. The result was stirred for 5 hours at ambient temperature, and then 200 mL of water was added thereto, generating a solid. The solid was filtered therefrom. The solid was dissolved in 200 mL of dichloromethane, and then, an extraction process was performed thereon three times by using 100 mL of water. A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified

75

by silica gel column chromatography to obtain 16.6 g (yield of 63%) of Intermediate A-1. The obtained compound was confirmed by LC-MS. $C_6H_5IN_2O_2$: M+1 265.1

Synthesis of Intermediate A-2

16.6 g (63 mmol) of Intermediate A-1, 6.8 g (69.2 mmol) of trimethylsilylacetylene, 1.38 g (3.2 mmol) of $PdCl_2(PPh_3)_2$, 1.2 g (6.4 mmol) of CuI were dissolved in 150 mL of THF, and then, 17.4 mL (126 mmol) of triethylamine was slowly added thereto and heated at a temperature of 70° C. Five hours after, the heated solution was cooled to ambient temperature, and then 100 mL of water was added thereto, and an extraction process was performed thereon three times. A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 8.8 g (yield of 60%) of Intermediate A-2. The obtained compound was confirmed by LC-MS. $C_{11}H_{14}N_2O_2Si$: M+1 235.3

Synthesis of Intermediate A-3

8.8 g (37.5 mmol) of Intermediate A-2 was dissolved in 100 mL of a mixed solution of MeOH/THF/H₂O (3/5/2 volume ratio), and then, 15.6 g (113 mmol) of K_2CO_3 was added thereto, and the result was stirred at ambient temperature for 3 hours. After the mixed solution was removed therefrom by evaporation, an extraction process was performed thereon three times by using 60 mL of dichloromethane and 50 mL of water. A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 5.96 g (yield of 98%) of Intermediate A-3. The obtained compound was confirmed by LC-MS. $C_8H_6N_2O_2$: M+1 163.1

Synthesis of Intermediate A-4

5.96 g (36.7 mmol) of Intermediate A-3, 9.2 g (34.8 mmol) of Intermediate A-1, 0.8 g (1.8 mmol) of $PdCl_2(PPh_3)_2$, 0.68 g (3.6 mmol) of CuI were dissolved in 80 mL of THF, and then, 15 mL (110 mmol) of triethylamine was slowly added thereto and heated at a temperature of 70° C. After the mixed solution was removed therefrom by evaporation, an extraction process was performed thereon three times by using 60 mL of dichloromethane and 50 mL of water. A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 9.63 g (yield of 88%) of Intermediate A-4. The obtained compound was confirmed by LC-MS. $C_{14}H_{10}N_4O_4$: M+1 299.3

Synthesis of Intermediate A-5

6.6 g (22.1 mmol) of Intermediate A-4 was dissolved in a mixed solution of 100 mL of CH_3CN and 50 mL of sulfone, and then, 11.2 mL (d=1.12 g/mL, 98%, 86.6 mmol) of $BF_3 \cdot Et_2O$ was slowly added thereto at a temperature of -30° C., and at the same temperature, the result was stirred for 10 minutes. 14.8 mL (111 mmol) of tert-butyl nitride was slowly added thereto, and the temperature was raised to ambient temperature, and the result was stirred for 1 hour. After a solvent was evaporated, 60 mL of DMSO was added thereto, and 20 g (89 mmol) of $CuBr_2$ were added thereto, and the result was stirred at ambient temperature for 3 hours.

76

100 mL of cold water was added to the resulting solution, and then, a generated solid was filtered therefrom, and then an extraction process was performed thereon three times by using 60 mL of dichloromethane and 50 mL of water. A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 3.76 g (yield of 40%) of Intermediate A-5. The obtained compound was confirmed by LC-MS. $C_{14}H_6Br_2N_2O_4$: M+1 424.9

Synthesis of Intermediate A-6

3.2 g (7.47 mmol) of Intermediate A-5 was dissolved in 100 mL of acetone and 1 mL of an acetic acid, and then, 2.61 g (16.5 mmol) of $KMnO_4$ was added thereto, and the result was refluxed while heating and stirred for 2 hours. The solution was cooled to ambient temperature, and then, 300 mL of acetone was added thereto and the resultant solution was subjected to sonication for 10 minutes. The generated MnO_2 was removed therefrom by filtering, and the resultant solution was evaporated and the residual was separation-purified by silica gel column chromatography to produce 3.37 g (yield of 99%) of Intermediate A-6. The obtained compound was confirmed by LC-MS. $C_{14}H_6Br_2N_2O_6$: M+1 456.9

Synthesis of Intermediate A-7

2.88 g (7 mmol) of Intermediate A-6 was dissolved in 100 mL of acetic acid, and then, the result was strongly stirred at a temperature of 80° C. for 5 minutes, and then, 14.2 g (63 mmol) of $SnCl_2 \cdot H_2O$ and 12 mL of 1N HCl were added thereto, and then stirred at a temperature of 80° C. for 5 hours. Once the reaction was complete, a generated solid was washed with an acetic acid and methanol to produce 0.93 g (yield of 37%) of Intermediate A-7. The obtained compound was confirmed by LC-MS. $C_{14}H_8Br_2N_2$: M+1 362.9

Synthesis of Intermediate A-8

0.93 g (2.59 mmol) of Intermediate A-7, 0.1 g (0.05 mmol) of CuI, 2.14 g (15.5 mmol) of K_2CO_3 , and 6.34 g (31 mmol) of iodobenzene were dissolved in 30 mL of DMPU, and then, the resultant solution was heated at a temperature of 120° C. for 4 hours. The solution was cooled to ambient temperature, and then, an extraction process was performed thereon three times by using 50 mL of water and 50 mL of Et_2O . A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 0.86 g (yield of 64%) of Intermediate A-8. The obtained compound was confirmed by LC-MS. $C_{26}H_{16}Br_2N_2$: M+1 517.2

Synthesis of Compound A


0.86 g (1.66 mmol) of Intermediate A-8, 0.7 g (4.14 mmol) of diphenylamine, 0.2 g (0.1 mmol) of $Pd_2(dba)_3$, 0.02 g (0.1 mmol) of $P(tBu)_3$, and 0.64 g (6.6 mmol) of $NaOtBu$ were dissolved in 20 mL of toluene, and then, the mixture was stirred at a temperature of 80° C. for 4 hours. The solution was cooled to ambient temperature, and then, an extraction process was performed thereon three times by using 50 mL of water and 50 mL of Et_2O . A collected organic layer was dried by using magnesium sulfate, and

77


then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 0.89 g (yield of 78%) of Compound A. The obtained compound was confirmed by LC-MS and ^1H NMR. $\text{C}_{50}\text{H}_{36}\text{N}_4$; $\text{M}+1$ 693.9

^1H NMR (CDCl_3 , 400 MHz) $\square$ 7.69 (d, 2H), 7.59-7.44 (m, 8H), 7.39-7.34 (m, 2H), 7.21-7.14 (m, 8H), 7.07 (s, 2H), 6.97-6.92 (m, 4H), 6.88-6.80 (m, 10H)

Synthesis of Compound B



78



Synthesis of Intermediate B-1

Intermediate B-1 was synthesized from naphthalene-2-thiol as described herein.

Synthesis of Intermediate B-2

3.4 g (10 mmol) of Intermediate B-1 was dissolved in 30 mL of dichloromethane, and then, at a temperature of -10°C ., 1 mL (20 mmol) of Br_2 was slowly added thereto. The solution was stirred at ambient temperature for 2 hours, and then, an extraction process was performed thereon three times by using 30 mL of water and 10 mL of 10% sodium thiosulfate aqueous solution. A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 4.58 g (yield of 92%) of Compound B-2. The obtained compound was identified by LC-MS. $\text{C}_{22}\text{H}_{10}\text{Br}_2\text{S}_2$; $\text{M}+1$ 499.2

Synthesis of Compound B

2.49 g (5 mmol) of Intermediate B-2, 2.12 g (12.5 mmol) of diphenylamine, 0.35 g (0.3 mmol) of $\text{Pd}_2(\text{dba})_3$, 0.06 g (0.1 mmol) of $\text{P}(\text{tBu})_3$, and 1.44 g (15 mmol) of NaOtBu were dissolved in 20 mL of toluene, and then, the mixture was stirred at a temperature of 80°C . for 4 hours. The solution was cooled to ambient temperature, and then, an extraction process was performed thereon three times by using 50 mL of water and 50 mL of Et_2O . A collected organic layer was dried by using magnesium sulfate, and then, the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 2.73 g (yield of 81%) of Compound A. The obtained compound was confirmed by LC-MS and ^1H NMR. $\text{C}_{46}\text{H}_{30}\text{N}_2\text{S}_2$; $\text{M}+1$ 675.9

^1H NMR (CDCl_3 , 400 MHz) $\square$ 8.39 (dd, 2H), 7.60-7.55 (m, 2H), 7.44-7.40 (m, 2H), 7.32 (s, 2H), 7.24-7.19 (m, 8H), 7.09-7.05 (m, 2H), 6.95-6.92 (m, 4H), 6.92-6.78 (m, 8H)

EXAMPLE 1

A top-type resonance device was manufactured by using Compound A.

A doping amount in a blue emission layer was 3%, and the top-type resonance device had the following structure.

ITO/HT-D1/HT13/HT3/H23+(Compound A)/ET1+ET-D1(5:5)/LiF/AgMg

EXAMPLE 2

A top-type resonance device was manufactured by using Compound B.


79

A doping amount in a blue emission layer was 3%, and the top-type resonance device and the following structure.

ITO/HT-D1/HT13/HT3/H23+(Compound B)/ET1+ET-D1(5:5)/LiF/AgMg

Comparative Example 1

As a Comparative Example 1, a doping amount in a blue emission layer was 3%, and the top-type resonance device and the following structure ITO/HT-D1/HT13/HT3/H23+(Compound 1000)/ET1+ET-D1(5:5)/LiF/AgMg.



Compound 1000

Results of Comparative Example 1 and Examples 1 and 2 are shown in Table 1 below.

TABLE 1

	Emission layer dopant	Driving voltage (V)	Current density (mA/cm ²)	y color coordinate	Efficiency (cd/A)	Half lifespan (hr @ 100 mA/cm ²)
Example 1	Compound A	3.8	11.8	0.043	4.8	100
Example 2	Compound B	3.9	12.9	0.041	4.1	110
Comparative Example 1	Comparative Example 1	3.9	14.5	0.039	3.5	85

Referring to Table 1, it may be seen that the organic light-emitting devices manufactured according to Examples 1 and 2 had better luminescent characteristics than that manufactured according to Comparative Example 1.

A compound according to an embodiment is a luminescent material with high efficiency and a luminescent wavelength thereof is controllable. An organic light-emitting device has a low driving voltage and a high current efficiency.

The embodiments may provide a luminescent material with high efficiency.

Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with


80

other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.


What is claimed is:

1. A compound represented by any one selected from the following Formula 1, Formula 2, and Formula 3:

<Formula 1>




<Formula 2>



-continued

<Formula 3>



wherein, in Formula 1, X and Y are each independently S, O, or NR₁,

wherein, in Formulae 2 and 3,

X and Y are each independently S, O, NR₁, or CR₂R₃,

wherein, in Formulae 1, 2, and 3,

A to F are each independently hydrogen, deuterium, or a substituted or unsubstituted C₆-C₆₀ aryl moiety fused to a corresponding backbone, at least one of A or B being

81

a substituted or unsubstituted C₆-C₆₀ aryl moiety fused to the corresponding backbone,

Ar₁ to Ar₄ and R₁ to R₃ are each independently a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

at least one substituent of the substituted C₁-C₆₀ alkyl group, substituted C₆-C₆₀ aryl group, substituted C₂-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is:

a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), or —B(Q₁₆)(Q₁₇);

a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group;


a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₂-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group, each substituted with a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₂-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), or —B(Q₂₆)(Q₂₇); or

82


—N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), or —B(Q₃₆)(Q₃₇), wherein Q₁₁ to Q₁₇, Q₂₁ to Q₂₇, and Q₃₁ to Q₃₇ are each independently a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₂-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group.

2. The compound as claimed in claim 1, wherein the compound represented by one of Formulae 1 to 3 is represented by one of the following Formulae 5 or 6:

<Formula 5>




<Formula 6>



wherein, in Formulae 5 and 6, Ar₁ to Ar₄, X, and Y are defined the same as Ar₁ to Ar₄, X, and Y of Formulae 1 to 3.

3. The compound as claimed in claim 1, wherein the compound represented by one of Formulae 1 to 3 is represented by one of the following Formulae 7 to 9:


<Formula 7>




83

-continued

<Formula 8>




<Formula 9>




wherein, in Formulae 7 to 7, Ar₁ to Ar₄, X, and Y are defined the same as Ar₁ to Ar₄, X, and Y of Formulae 1 to 3.

4. The compound as claimed in claim 1, wherein the compound represented by one of Formulae 1 to 3 is represented by one of the following Formulae 10 to 12:


<Formula 10>



<Formula 11>



<Formula 12>



84

wherein, in Formulae 10 to 12 Ar₁ to Ar₄, X, and Y are defined the same as Ar₁ to Ar₄, X, and Y of Formulae 1 to 3.

5. The compound as claimed in claim 1, wherein R₁ to R₃ are each independently a substituted or unsubstituted C₁-C₂₀ alkyl group or a substituted or unsubstituted C₆-C₆₀ aryl group.


6. The compound as claimed in claim 1, wherein R₁ to R₃ are each independently a methyl group or a phenyl group.

7. The compound as claimed in claim 1, wherein Ar₁ to Ar₄ are each independently a substituted or unsubstituted C₆-C₂₀ aryl group or a substituted or unsubstituted C₁-C₂₀ heteroaryl group.

8. The compound as claimed in claim 1, wherein Ar₁ to Ar₄ are each independently a group represented by one of the following Formulae 2a and 2b:


2a

20



2b

25



wherein, in Formulae 2a and 2b,

Z₁ is a hydrogen, a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₂-C₆₀ heteroaryl group, a substituted or unsubstituted a monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group;

p is an integer of 1 to 7,


* indicates a binding site to a neighboring atom,

when p is 2 or more, each Z₁ is identical or different from one another.

9. The compound as claimed in claim 1, wherein Ar₁ to Ar₄ are each independently a group represented by one of the following Formulae 3a to 3k, in which * indicates a binding site to a neighboring atom:









3a

65



85

-continued

**86**

-continued

3b

5

10

3c

15

20

3d

25

3e

30

3f

35

3g

40

45

3h

50

3i

55


60

65

3j




3k

10. The compound as claimed in claim 1, wherein the compound represented by one of Formula 1, Formula 2, and Formula 3 is one of the following compounds:



87

-continued

**88**

-continued

5

10

15

20

25

30

35

40





45

50

55





60

65




89

-continued



90

-continued



11. An organic light-emitting device, comprising

a first electrode;

a second electrode facing the first electrode; and

an organic layer between the first electrode and the second electrode, the organic layer including an emission layer,

wherein the organic layer includes the compound as claimed in claim 1.

12. The organic light-emitting device as claimed in claim 11, wherein the organic layer is formed by a wet process.

13. The organic light-emitting device as claimed in claim 11, wherein

the first electrode is an anode,

the second electrode is a cathode, and

the organic layer includes:

a hole transport region between the first electrode and the emission layer, the hole transport region including at least one of a hole injection layer, a hole transport layer, a buffer layer, and an electron blocking layer, and

an electron transport region between the emission layer and the second electrode, the electron transport region including at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

14. The organic light-emitting device as claimed in claim 13, wherein the compound is included in the emission layer.

15. The organic light-emitting device as claimed in claim 13, wherein the hole transport region includes a charge-generating material.

16. The organic light-emitting device as claimed in claim 13, wherein the electron transport region includes a metal complex.

17. The organic light-emitting device as claimed in claim 16, wherein the metal complex is a lithium complex.


18. The organic light-emitting device as claimed in claim 16, wherein the metal complex is a lithium quinolate.

19. The organic light-emitting device as claimed in claim 16, wherein the metal complex is represented by the following Formula ET-D2:

91

92

<ET-D2>



5

10

20. A flat display apparatus, comprising:
a thin film transistor, the thin film transistor including a
source electrode and a drain electrode; and
the organic light-emitting device as claimed in claim 11,
wherein the first electrode of the organic light-emitting
device is electrically connected to the source electrode
or the drain electrode of the thin film transistor.

15

20

* * * * *

专利名称(译)	包括其的化合物和有机发光装置		
公开(公告)号	US10115907	公开(公告)日	2018-10-30
申请号	US14/736375	申请日	2015-06-11
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	SHIN DAEYUP KIM YOUNGKOOK YOON JIHWAN		
发明人	SHIN, DAEYUP KIM, YOUNGKOOK YOON, JIHWAN		
IPC分类号	H01L51/00 C07D487/04 C07D495/04 C09K11/06 H01L51/50		
CPC分类号	H01L51/0061 C07D495/04 C09K11/06 H01L51/0072 H01L51/0074 C07D487/04 H01L51/5076 C09K2211/1007 C09K2211/1014 C09K2211/1029 C09K2211/1037 C09K2211/1044 C09K2211/1088 C09K2211/1092 C09K2211/181 H01L51/5012		
审查员(译)	张 , RUIYUN		
优先权	1020140195956 2014-12-31 KR		
其他公开文献	US20160190465A1		
外部链接	Espacenet		

摘要(译)

化合物，有机发光装置和平板显示装置，该化合物由选自下式1，式2和式3中的任何一种表示：

