



US 20180301638A1

(19) **United States**(12) **Patent Application Publication**
AHN et al.(10) **Pub. No.: US 2018/0301638 A1**(43) **Pub. Date: Oct. 18, 2018**(54) **HETEROCYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME***C07F 9/64* (2006.01)*C09K 11/02* (2006.01)*C09K 11/06* (2006.01)*C07F 9/6596* (2006.01)*C07F 9/572* (2006.01)(71) Applicant: **Samsung Display Co., Ltd.**, Yongin-si
(KR)(52) **U.S. Cl.**CPC *H01L 51/0072* (2013.01); *H01L 51/5004*(2013.01); *C07F 9/64* (2013.01); *C09K 11/025*(2013.01); *C09K 11/06* (2013.01); *H01L**51/0094* (2013.01); *C09K 2211/1018*(2013.01); *H01L 51/0067* (2013.01); *C07F**9/5728* (2013.01); *H01L 51/5012* (2013.01);*H01L 51/5016* (2013.01); *H01L 2251/5376*(2013.01); *H01L 2251/552* (2013.01); *C07F**9/6596* (2013.01)(72) Inventors: **Heechoon AHN**, Yongin-si (KR);
Soobyung KO, Yongin-si (KR); **Mieun**
JUN, Yongin-si (KR); **Sungbum KIM**,
Yongin-si (KR); **Mina JEON**, Yongin-si
(KR); **Youngkook KIM**, Yongin-si
(KR); **Seokhwan HWANG**, Yongin-si
(KR)(21) Appl. No.: **15/788,605**

(57)

ABSTRACT(22) Filed: **Oct. 19, 2017**

An organic light-emitting device includes a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer. The emission layer includes at least one heterocyclic compound of Formula 1. The heterocyclic compound may be a host or a delayed fluorescent dopant. The organic light-emitting device including the heterocyclic compound may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

(30) **Foreign Application Priority Data**

Apr. 14, 2017 (KR) 10-2017-0048532

Publication Classification(51) **Int. Cl.***H01L 51/00* (2006.01)*H01L 51/50* (2006.01)**10****190****150****110**

10

190
150
110

HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims priority to and the benefit of Korean Patent Application No. 10-2017-0048532, filed on Apr. 14, 2017, in the Korean Intellectual Property Office, the entire content of which is incorporated herein by reference.

BACKGROUND

1. Field

[0002] One or more aspects of one or more embodiments of the present disclosure relate to a heterocyclic compound and an organic light-emitting device including the same.

2. Description of the Related Art

[0003] Organic light-emitting devices are self-emission devices that produce full-color images, have wide viewing angles, high contrast ratios, short response times, and excellent characteristics in terms of brightness, driving voltage, and response speed, compared to devices in the pertinent art.


[0004] An example of such organic light-emitting device may include a first electrode disposed (e.g., positioned) on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode, for example, may move toward the emission layer through the hole transport region, and electrons provided from the second electrode, for example, may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may then recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.

SUMMARY

[0005] One or more aspects of one or more embodiments of the present disclosure are directed to a novel heterocyclic compound and an organic light-emitting device including the same.

[0006] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.


[0007] An aspect of one or more embodiments of the present disclosure is directed to a heterocyclic compound represented by Formula 1:




Formula 1

-continued


Formula 2-1




Formula 2-2



Formula 3-1



Formula 3-2



[0008] In Formulae 1, 2-1, 2-2, 3-1, and 3-2,

[0009] X_1 may be O or S,

[0010] Ar_1 and Ar_2 may each independently be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_2 - C_{60} heterocyclic group,

[0011] a_1 and a_2 may each independently be an integer from 0 to 3,

[0012] R_1 and R_2 may each independently be a group represented by Formula 2-1,

[0013] R_3 may be a group represented by Formula 2-2,

[0014] b_1 and b_2 may each independently be 0 or 1,

[0015] a_1 , a_2 , b_1 , and b_2 may satisfy $a_1 + b_1 \geq 1$ and $a_2 + b_2 \leq 1$,

[0016] A_1 to A_4 may each independently be a benzene group, a naphthalene group, a phenanthrene group, a fluo-ranthen group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, or a tetrazine group,

[0017] A_5 to A_8 may each independently be a benzene group, a naphthalene group, a phenanthrene group, a fluo-ranthen group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, or a tetrazine group,

[0018] Y_1 may be a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, $Si(R_{19})(R_{20})$, $S(=O)(R_{19})$, $S(=O)_2(R_{19})$, or $P(=O)(R_{19})$,

[0019] Y_2 may be a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, $Si(R_{19})(R_{20})$, $S(=O)(R_{19})$, $S(=O)_2(R_{19})$, or $P(=O)(R_{19})$,

[0020] Y_3 may be a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, $Si(R_{19})(R_{20})$, $S(=O)(R_{19})$, $S(=O)_2(R_{19})$, or $P(=O)(R_{19})$,

[0021] Y_4 may be a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, $Si(R_{19})(R_{20})$, $S(=O)(R_{19})$, $S(=O)_2(R_{19})$, or $P(=O)(R_{19})$,

[0022] Y_5 may be a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, $Si(R_{19})(R_{20})$, $S(=O)(R_{19})$, $S(=O)_2(R_{19})$, or $P(=O)(R_{19})$,

[0023] when Y_1 is O, a group represented by Formula 2-1 may be different from a group represented by Formula 2-2,

[0024] R_{11} to R_{14} may each independently be a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1-C_{60} alkyl group, a substituted or unsubstituted C_2-C_{60} alkenyl group, a substituted or unsubstituted C_2-C_{60} alkynyl group, a substituted or unsubstituted C_1-C_{60} alkoxy group, a substituted or unsubstituted C_3-C_{10} cycloalkyl group, a substituted or unsubstituted C_3-C_{10} cycloalkenyl group, a substituted or unsubstituted C_1-C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3-C_{10} cycloalkenyl group, a substituted or unsubstituted C_1-C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6-C_{60} aryl group, a substituted or unsubstituted C_6-C_{60} aryloxy group, a substituted or unsubstituted C_6-C_{60} arylthio group, a substituted or unsubstituted C_1-C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-S(=O)(Q_1)$, or $-P(=O)(Q_1)(Q_2)$, two or more $R_{11}(s)$ may optionally be linked to form a substituted or unsubstituted C_5-C_{60} carbocyclic group or a substituted or unsubstituted C_2-C_{60} heterocyclic group, two or more $R_{12}(s)$ may optionally be linked to form a substituted or unsubstituted C_5-C_{60} carbocyclic group or a substituted or unsubstituted C_2-C_{60} heterocyclic group, two or more $R_{13}(s)$ may optionally be linked to form a substituted or unsubstituted C_5-C_{60} carbocyclic group or a substituted or unsubstituted C_2-C_{60} heterocyclic group, and two or more $R_{14}(s)$ may optionally be linked to form a substituted or unsubstituted C_5-C_{60} carbocyclic group, or a substituted or unsubstituted C_2-C_{60} heterocyclic group,

[0025] when Y_2 is a single bond, at least one of R_{13} and R_{14} may be a group represented by Formula 3-1 or a group represented by Formula 3-2,

[0026] R_{15} to R_{20} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1-C_{60} alkyl group, a substituted or unsubstituted C_2-C_{60} alkenyl group, a substituted or unsubstituted C_2-C_{60} alkynyl group, a substituted or unsubstituted C_1-C_{60} alkoxy group, a substituted or unsubstituted C_3-C_{10} cycloalkyl group, a substituted or unsubstituted C_3-C_{10} cycloalkenyl group, a substituted or unsubstituted C_1-C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3-C_{10} cycloalkenyl group, a substituted or unsubstituted C_1-C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6-C_{60} aryl group, a substituted or unsubstituted C_6-C_{60} aryloxy group, a substituted or unsubstituted C_6-C_{60} arylthio group, a substituted or unsubstituted C_1-C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic, $-S(=O)(Q_1)$, and $-P(=O)(Q_1)(Q_2)$,

[0027] b_{11} to b_{18} may each independently be an integer from 1 to 8,

[0028] at least one substituent of the substituted C_5-C_{60} carbocyclic group, the substituted C_2-C_{60} heterocyclic group, the substituted C_1-C_{60} alkyl group, the substituted

C_2-C_{60} alkenyl group, the substituted C_2-C_{60} alkynyl group, the substituted C_1-C_{60} alkoxy group, the substituted C_3-C_{10} cycloalkyl group, the substituted C_1-C_{10} heterocycloalkyl group, the substituted C_3-C_{10} cycloalkenyl group, the substituted C_1-C_{10} heterocycloalkenyl group, the substituted C_6-C_{60} aryl group, the substituted C_6-C_{60} aryloxy group, the substituted C_6-C_{60} arylthio group, the substituted C_1-C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0029] deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group;

[0030] a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{11})(Q_{12})(Q_{13})$, $-N(Q_{11})(Q_{12})$, $-B(Q_{11})(Q_{12})$, $-C(=O)(Q_{11})$, $-S(=O)_2(Q_{11})$, $-P(=O)(Q_{11})(Q_{12})$;

[0031] a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0032] a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{21})(Q_{22})(Q_{23})$, $-N(Q_{21})(Q_{22})$, $-B(Q_{21})(Q_{22})$, $-C(=O)(Q_{21})$, $-S(=O)_2(Q_{21})$, and $-P(=O)(Q_{21})(Q_{22})$; and

[0033] $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0034] where Q_1 , Q_2 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group,

a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

[0035] * indicates a binding site to a neighboring atom.

[0036] Another aspect of one or more embodiments of the present disclosure is directed to an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes at least one of the heterocyclic compounds described above.


BRIEF DESCRIPTION OF THE DRAWING

[0037] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with the drawing, which is a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

[0038] Reference will now be made in more detail to embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the drawing, to explain aspects of the present description. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” “one of,” and “selected from,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list. Further, the use of “may” when describing embodiments of the present invention refers to “one or more embodiments of the present invention.”

[0039] A heterocyclic compound according to an embodiment is represented by Formula 1 below:



Formula 1

[0040] X₁ in Formula 1 may be O or S.

[0041] Ar₁ and Ar₂ in Formula 1 may each independently be a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group.

[0042] In one embodiment, Ar₁ and Ar₂ may each independently be selected from:

[0043] a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine

group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, and a quinazoline group; and

[0044] a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, and a quinazoline group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂), and

[0045] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0046] In one or more embodiments, Ar₁ and Ar₂ may each independently be a substituted or unsubstituted aromatic C₆-C₃₀ carbocyclic group, or a substituted or unsubstituted π electron-depleted nitrogen-containing C₂-C₃₀ heterocyclic group.

[0047] In one or more embodiments, Ar₁ and Ar₂ may each independently be selected from:

[0048] a benzene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, and a triazine group; and

[0049] a benzene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, and a triazine group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a pyridinyl group, a pyrimidinyl group, a triazinyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothio-

phenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, and $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, and

[0050] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0051] In one embodiment, Ar_1 and Ar_2 may each independently be a benzene group or a pyridine group.

[0052] In one embodiment, Ar_1 and Ar_2 may each be a benzene group, but embodiments of the present disclosure are not limited thereto.


[0053] In Formula 1, a_1 and a_2 may each independently be an integer from 0 to 3, b_1 and b_2 may each independently be 0 or 1, and a_1 , a_2 , b_1 , and b_2 may satisfy $a_1+b_1 \geq 1$ and $a_2+b_2 \geq 1$.

[0054] In one embodiment, a_1 and a_2 may each independently be 0 or 1, but embodiments of the present disclosure are not limited thereto.

[0055] In one embodiment, a_1 , a_2 , b_1 , and b_2 may satisfy $a_1+b_1=1$ and $a_2+b_2=1$.

[0056] In one embodiment, b_1 and b_2 may satisfy $b_1+b_2=0$ or $b_1+b_2=1$.

[0057] In Formula 1, R_1 and R_2 may each be a group represented by Formula 2-1, and R_3 may be a group represented by Formula 2-2:



[0058] A_1 to A_4 in Formulae 2-1 and 2-2 may each independently be a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, or a tetrazine group.

[0059] In one embodiment, A_1 to A_4 may each independently be a benzene group, a pyridine group, or a pyrimidine group.

[0060] In one embodiment, A_1 to A_4 may each be a benzene group, but embodiments of the present disclosure are not limited thereto.

[0061] In one embodiment, A_5 to A_8 in Formulae 3-1 and 3-2 (provided below) may each independently be a benzene group, a pyridine group, or a pyrimidine group.

[0062] In one embodiment, A_5 to A_8 in Formulae 3-1 and 3-2 (provided below) may each be a benzene group, but embodiments of the present disclosure are not limited thereto.

[0063] In Formulae 2-1 and 2-2, Y_1 may be a single bond, O, S, SO_2 , $\text{N}(\text{R}_{19})$, $\text{C}(\text{R}_{19})(\text{R}_{20})$, $\text{Si}(\text{R}_{19})(\text{R}_{20})$, $\text{S}(\text{=O})(\text{R}_{19})$, $\text{S}(\text{=O})_2(\text{R}_{19})$, or $\text{P}(\text{=O})(\text{R}_{19})$,

[0064] Y_2 may be a single bond, O, S, SO_2 , $\text{N}(\text{R}_{19})$, $\text{C}(\text{R}_{19})(\text{R}_{20})$, $\text{Si}(\text{R}_{19})(\text{R}_{20})$, $\text{S}(\text{=O})(\text{R}_{19})$, $\text{S}(\text{=O})_2(\text{R}_{19})$, or $\text{P}(\text{=O})(\text{R}_{19})$,

[0065] when Y_2 is O, a group represented by Formula 2-1 and a group represented by Formula 2-2 may be different from each other, and

[0066] when Y_2 is a single bond, at least one of R_{13} and R_{14} may be a group represented by Formula 3-1 or a group represented by Formula 3-2 (provided below).

[0067] In one embodiment, when Y_2 is O, Y_1 may be a single bond, S, SO_2 , $\text{N}(\text{R}_{19})$, $\text{C}(\text{R}_{19})(\text{R}_{20})$, or $\text{Si}(\text{R}_{19})(\text{R}_{20})$.

[0068] In one embodiment, Y_1 may be a single bond, O, S, SO_2 , $\text{N}(\text{R}_{19})$, $\text{C}(\text{R}_{19})(\text{R}_{20})$, or $\text{Si}(\text{R}_{19})(\text{R}_{20})$.

[0069] In one embodiment, Y_2 may be a single bond, O, S, SO_2 , $\text{N}(\text{R}_{19})$, $\text{C}(\text{R}_{19})(\text{R}_{20})$, or $\text{Si}(\text{R}_{19})(\text{R}_{20})$.

[0070] b_{11} to b_{14} in Formulae 2-1 and 2-2 may each independently be an integer from 1 to 8. In one embodiment, b_{11} to b_{14} may each independently be an integer from 1 to 4, but embodiments of the present disclosure are not limited thereto.

[0071] In Formulae 2-1 and 2-2, R_{11} to R_{14} may each independently be a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{S}(\text{=O})(\text{Q}_1)$, or $-\text{P}(\text{=O})(\text{Q}_1)(\text{Q}_2)$,

[0072] where two or more $\text{R}_{11}(\text{s})$ may optionally be linked to form a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_2 - C_{60} heterocyclic group; two or more $\text{R}_{12}(\text{s})$ may optionally be linked to form a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_2 - C_{60} heterocyclic group; two or more $\text{R}_{13}(\text{s})$ may optionally be linked to form a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_2 - C_{60} heterocyclic group; and two or more $\text{R}_{14}(\text{s})$ may optionally be linked to form a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_2 - C_{60} heterocyclic group.

[0073] In one embodiment, R_{11} to R_{14} may each independently be selected from:

[0074] a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl

group, a quinoxaliny group, a quinazolinyl group, a carbazoyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

[0075] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazolinyl group, a carbazoyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a phenyl group, a naphthyl group, a fluorenyl group, a carbazoyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group.


[0076] In one or more embodiments, R₁₁ to R₁₄ may each independently be selected from:

[0077] a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a phenyl group, an acridyl group, and a carbazoyl group; and


[0078] a phenyl group, an acridyl group, and a carbazoyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, and a C₁-C₂₀ alkoxy group.

[0079] In one embodiment, R₁₁ to R₁₄ may each independently be a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, a phenyl group, an acridyl group substituted with a methyl group, or a carbazoyl group.

Formula 3-1



Formula 3-2



[0080] Y₃ in Formulae 3-1 and 3-2 may be a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉). In one embodiment, Y₃ may be a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), or Si(R₁₉)(R₂₀), but embodiments of the present disclosure are not limited thereto.

[0081] Y₄ in Formulae 3-1 and 3-2 may be a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉),

S(=O)₂(R₁₉), or P(=O)(R₁₉). In one embodiment, Y₄ may be a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), or Si(R₁₉)(R₂₀), but embodiments of the present disclosure are not limited thereto.

[0082] Y₅ in Formulae 3-1 and 3-2 may be a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉). In one embodiment, Y₅ may be a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), or Si(R₁₉)(R₂₀), but embodiments of the present disclosure are not limited thereto.

[0083] R₁₅ to R₂₀ in Formulae 3-1 and 3-2 may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted

[0084] C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic, —S(=O)(Q₁), and —P(=O)(Q₁)(Q₂) group.

[0085] In one embodiment, R₁₅ to R₂₀ may each independently be selected from:

[0086] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazolinyl group, a carbazoyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

[0087] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazolinyl group, a carbazoyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a phenyl group, a naphthyl group, a fluorenyl group, a carbazoyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group.

[0088] In one or more embodiments, R_{15} to R_{20} may each independently be selected from:


[0089] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C_1 - C_{20} alkoxy group, and a phenyl group; and

[0090] a phenyl group substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, and a C_1 - C_{20} alkoxy group.

[0091] In one or more embodiments, R_{13} to R_{18} may each independently be hydrogen, deuterium, a methyl group, or a phenyl group, but embodiments of the present disclosure are not limited thereto.

[0092] b_{15} to b_{18} in Formulae 3-1 and 3-2 may each independently be an integer from 1 to 8. In one embodiment, b_{15} to b_{18} may each independently be an integer from 1 to 4, but embodiments of the present disclosure are not limited thereto.

[0093] In one embodiment, the group represented by Formula 3-1 may be a group represented by Formula 3-1-1:




[0094] In Formula 3-1-1,

[0095] Y_3 , R_{15} , and R_{16} are the same as described herein, and

[0096] * indicates a binding site to a neighboring atom.

[0097] * in Formulae 2-1, 2-2, 3-1, and 3-2 indicates a binding site to a neighboring atom.

[0098] In one embodiment, when Y_1 is a single bond, the group represented by Formula 2-1 may be a group represented by Formula 2-2-1:



[0099] In Formula 2-2-1,

[0100] R_{28} may be a group represented by Formula 3-1 or a group represented by Formula 3-2,

[0101] definitions of R_{21} to R_{27} are each independently the same as those provided in connection with R_{11} to R_{14} .

[0102] In one embodiment, R_{21} to R_{27} may each independently be selected from:

[0103] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl

group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

[0104] a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group.


[0105] In one embodiment, R_{23} may be selected from:

[0106] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and


[0107] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a phenyl group, a naphthyl group, a fluorenyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, and [0108] R₂₁, R₂₂, and R₂₄ to R₂₇ may each be hydrogen, but embodiments of the present disclosure are not limited thereto.

[0109] In one embodiment, the heterocyclic compound represented by Formula 1 may be represented by Formula 1-1 or 1-2:

Formula 1-1



Formula 1-2



[0110] In Formulae 1-1 and 1-2,

[0111] definitions of A₁₁, A₁₂, A₂₁, A₂₂, A₃₁, and A₃₂ are each independently the same as those provided in connection with A₁ to A₄,

[0112] definitions of Y₁₁ and Y₁₃ are the same as that provided in connection with Y₁,

[0113] definition of Y₁₂ is the same as that provided in connection with Y₂,

[0114] and when one of Y₁₁ and Y₁₂ is oxygen (O), the other one of Y₁₁ and Y₁₂ is not O,

[0115] definitions of R₁₀₁, R₁₀₂, R₁₁₁, R₁₁₂, R₁₂₁, and R₁₂₂ are each independently the same as those provided in connection with R₁₁ to R₁₄, and

[0116] X₁, Ar₁, and Ar₂ are respectively the same as described in connection with Formula 1.

[0117] At least one substituent of the substituted C₅-C₆₀ carbocyclic group, the substituted C₂-C₆₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0118] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0119] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0120] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;


[0121] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and


[0122] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0123] where Q₁, Q₂, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.


[0124] In one embodiment, the heterocyclic compound may be one of Compounds 1 to 30 illustrated below, but embodiments of the present disclosure are not limited thereto:




1




3




4



5




2




6

-continued




7



-continued



8




9




10

11




12


-continued



13




14




15


-continued




16



17




18




19

-continued




20

-continued




23


21




24



22




25




-continued

26




-continued


30




27



28



29



[0125] Since the heterocyclic compound represented by Formula 1 includes phosphine oxide (or phosphine sulfide) and has a structure in which the P atom of the phosphine oxide is substituted with N-containing hetero ring to form a P—N bond, the heterocyclic compound may have excellent stability, as compared with a compound having no P—N bond. Also, when the heterocyclic compound is applied in an organic light-emitting device, the lifespan and efficiency of the organic light-emitting device may increase.

[0126] Therefore, an electronic device, for example, an organic light-emitting device, which includes the heterocyclic compound represented by Formula 1, may have low driving voltage, high current density, and high efficiency.

[0127] Also, the heterocyclic compound represented by Formula 1 is a compound that may emit delayed fluorescence, and triplet state excitons, as well as singlet state excitons, may be used for light emission according to an appropriate combination of substituents. Therefore, an organic light-emitting device including the heterocyclic compound may have excellent luminescent efficiency.

[0128] In one embodiment, an organic light-emitting device may include: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed (e.g., positioned) between the first electrode and the second electrode, the organic layer including an emission layer and at least one heterocyclic compound of Formula 1.

[0129] In one or more embodiments, the first electrode may be an anode.

[0130] In one or more embodiments, the second electrode may be a cathode.

[0131] In one or more embodiments, the first electrode may be an anode, and the second electrode may be a cathode.

[0132] In one embodiment, the organic layer may further include a hole transport region between the first electrode and the emission layer, and an electron transport region between the emission layer and the second electrode.

[0133] In one embodiment, the hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.

[0134] In one embodiment, the electron transport region may include at least one selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

[0135] In one or more embodiments, the emission layer may include the condensed heterocyclic compound represented by Formula 1. In one embodiment, the emission layer may emit phosphorescence or delayed fluorescence.

[0136] In one embodiment, the heterocyclic compound included in the emission layer may be a host, and the emission layer may further include a phosphorescent dopant.

[0137] In one or more embodiments, the heterocyclic compound included in the emission layer may be a host, and the emission layer may further include a delayed fluorescent dopant satisfying Equation 1 below:

$$|E_{D, S1} - E_{D, T1}| \leq 0.3 \text{ eV.} \quad \text{Equation 1}$$

[0138] In Equation 1,

[0139] $E_{D, S1}$ is a singlet energy level (eV) of the delayed fluorescent dopant, and

[0140] $E_{D, T1}$ is a triplet energy level (eV) of the delayed fluorescent dopant.

[0141] Since a difference between the singlet energy level ($E_{D, S1}$) and the triplet energy level ($E_{D, T1}$) of the delayed fluorescent dopant is 0.3 eV or less, singlet state excitons and triplet state excitons generated by an electric field may easily transit to an intermediate state.

[0142] In one or more embodiments, the heterocyclic compound included in the emission layer may be a delayed fluorescent dopant satisfying Equation 1 below.

$$|E_{D, S1} - E_{D, T1}| \leq 0.3 \text{ eV.} \quad \text{Equation 1}$$

[0143] In Equation 1,

[0144] $E_{D, S1}$ is a singlet energy level (eV) of the delayed fluorescent dopant, and

[0145] $E_{D, T1}$ is a triplet energy level (eV) of the delayed fluorescent dopant.

[0146] Since a difference between the singlet energy ($E_{D, S1}$) and the triplet energy ($E_{D, T1}$) of the delayed fluorescent dopant is 0.3 eV or less, singlet state excitons and triplet state excitons generated by an electric field may easily transit to an intermediate state.

[0147] In one or more embodiments, the heterocyclic compound included in the emission layer may be a delayed fluorescent dopant, the emission layer may further include a host, and the delayed fluorescent dopant and the host may satisfy Equation 2 or 3 below:

$$|E_{H, HOMO} - E_{D, HOMO}| \leq 0.5 \text{ eV} \quad \text{Equation 2}$$

$$|E_{H, LUMO} - E_{D, LUMO}| \leq 0.5 \text{ eV} \quad \text{Equation 3}$$

[0148] In Equations 2 and 3,

[0149] $E_{H, HOMO}$ is a highest occupied molecular orbital level of the host,

[0150] $E_{D, HOMO}$ is a highest occupied molecular orbital level of the dopant,





[0151] $E_{H, LUMO}$ is a lowest unoccupied molecular orbital level of the host, and

[0152] $E_{D, LUMO}$ is a lowest unoccupied molecular orbital level of the dopant.

[0153] Since a difference in highest occupied molecular orbital levels or lowest unoccupied molecular orbital levels of the delayed fluorescent dopant (e.g., the heterocyclic compound) and the host is 0.5 eV or less, charge transfer from the host to the delayed fluorescent dopant may be facilitated. Therefore, an organic light-emitting device including the heterocyclic compound as the delayed fluorescent dopant may have excellent luminescent efficiency.


[0154] For example, the host may include a silicon-based compound, a carbazole-based compound, or a phosphine oxide-based compound:

[0155] According to an embodiment, the host may be one of Compounds H101 to H106 below:




-continued

H105



H106



[0156] At least one layer selected from the electron transport layer and the electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof.

[0157] The organic light-emitting device including the heterocyclic compound may have low driving voltage, high current density, and high efficiency.

[0158] A synthesis method for the heterocyclic compound represented by Formula 1 should be apparent to those of ordinary skill in the art by referring to the following examples (provided below).

[0159] At least one of the heterocyclic compounds of Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the heterocyclic compound may be included in at least one layer selected from a hole transport region and an emission layer. In one or more embodiments, the heterocyclic compound of Formula 1 may be used as a material for a capping layer located outside a pair of electrodes of an organic light-emitting device.

[0160] For example, an organic light-emitting device may include: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes at least one heterocyclic compound represented by Formula 1.

[0161] The reference to a layer including at least one heterocyclic compound, as used herein, may refer to a layer including one or more of the same heterocyclic compound represented by Formula 1 and/or a layer including two or more different heterocyclic compounds represented by Formula 1.

[0162] For example, the organic layer may include, as the heterocyclic compound, only Compound 1. In this embodiment, Compound 1 may exist in an emission layer of the organic light-emitting device. In another example, the

organic layer may include, as the heterocyclic compound, Compound 1 and Compound 2. In this embodiment, Compound 1 and Compound 2 may both exist in the same layer (e.g., Compound 1 and Compound 2 may both exist in an emission layer), or in different layers (e.g., Compound 1 may exist in an emission layer and Compound 2 may exist in an electron transport layer).

[0163] In one embodiment,

[0164] the first electrode of the organic light-emitting device may be an anode,

[0165] the second electrode of the organic light-emitting device may be a cathode,

[0166] the organic layer of the organic light-emitting device may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,

[0167] the hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, a buffer layer, an emission auxiliary layer, and an electron blocking layer, and

[0168] the electron transport region may include at least one layer selected from a hole blocking layer, an electron transport layer, and an electron injection layer. The electron transport region may include the heterocyclic compound represented by Formula 1.

[0169] In one embodiment, the electron transport region of the organic light-emitting device may include an electron transport layer and an electron injection layer, and at least one of the electron transport layer and the electron injection layer may include the heterocyclic compound represented by Formula 1. However, embodiments of the present disclosure are not limited thereto.

[0170] In the organic light-emitting device, at least one layer selected from the electron transport layer and the electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof.

[0171] In the organic light-emitting device, the emission layer may include a dopant and a host, and the host may include at least one compound selected from an anthracene-based compound, a pyrene-based compound, and a spiro-bifluorene-based compound

[0172] The term “organic layer” used herein may refer to a single layer and/or a plurality of layers disposed (e.g., positioned) between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0173] The drawing is a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0174] Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing the organic light-emitting device 10 will be described in connection with the drawing.

[0175] In the drawing, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent (or suitable)

mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and/or water resistance.

[0176] The first electrode **110** may be formed by depositing or sputtering a material for forming the first electrode **110** on the substrate. When the first electrode **110** is an anode, the material for forming the first electrode may be selected from materials with a high work function so as to facilitate hole injection.

[0177] The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode **110** is a transmissive electrode, a material for forming the first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and a combination thereof, but embodiments of the present disclosure are not limited thereto.

[0178] In one or more embodiments, when the first electrode **110** is a semi-transmissive electrode or a reflective electrode, a material for forming the first electrode may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and a combination thereof, but embodiments of the present disclosure are not limited thereto.

[0179] The first electrode **110** may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **110** is not limited thereto.

[0180] The organic layer **150** may be disposed on the first electrode **110**. The organic layer **150** may include an emission layer.



[0181] The organic layer **150** may further include a hole transport region between the first electrode **110** and the emission layer, and an electron transport region between the emission layer and the second electrode **190**.

[0182] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.


[0183] The hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.

[0184] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein for each structure, constituting layers are sequentially stacked from the first electrode **110** in this stated order, but the structure of the hole transport region is not limited thereto.

[0185] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β -NPB, TPD, spiro-TPD, spiro-NPB, methylated NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANT/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




-continued




2-TNATA


-continued




Spiro-TPD





Spiro-NPB




NPB




methylated-NPB

 β -NPB

TAPC




TPD




HMTPD

Formula 201



Formula 202



[0195] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a

[10196] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazoly group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoly group, a dibenzocarbazoly group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂), and

[0197] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0198] In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

[0199] In one or more embodiments, xa5 may be 1, 2, 3, or 4.

[0200] In one or more embodiments, R_{201} to R_{204} and Q_{201} may each independently be selected from:

[0201] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0202] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl

group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), and —N(Q_{31})(Q_{32}), and

[0203] Q_{31} to Q_{33} may be the same as described herein.

[0204] In one or more embodiments, at least one of R_{201} to R_{203} in Formula 201 may be selected from:

[0205] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0206] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,

[0207] but embodiments of the present disclosure are not limited thereto.

[0208] In one or more embodiments, in Formula 202, i) R_{201} and R_{202} may be linked via a single bond, and/or ii) R_{203} and R_{204} may be linked via a single bond.


[0209] In one or more embodiments, at least one of R_{201} to R_{204} in Formula 202 may be selected from:

[0210] a carbazolyl group; and


[0211] a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,

[0212] but embodiments of the present disclosure are not limited thereto.


[0213] The compound represented by Formula 201 may be represented by Formula 201A:




[0214] In one embodiment, the compound represented by Formula 201 may be represented by Formula 201A(1) below, but embodiments of the present disclosure are not limited thereto:




[0215] In one embodiment, the compound represented by Formula 201 may be represented by Formula 201A-1 below, but embodiments of the present disclosure are not limited thereto:



[0216] In one embodiment, the compound represented by Formula 202 may be represented by Formula 202A:



[0217] In one embodiment, the compound represented by Formula 202 may be represented by Formula 202A-1:



[0218] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0219] definitions for L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} are the same as those provided herein,

[0220] definitions of R_{211} and R_{212} may each independently be the same as that provided in connection with R_{203} ,


[0221] R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0222] The hole transport region may include at least one compound selected from Compounds HT1 to HT39, but embodiments of the present disclosure are not limited thereto:


HT1




HT2



HT3




HT4




-continued


HT5




HT6



HT7




HT8




-continued


HT9




HT10




HT11




HT12



HT13




HT14




-continued


HT15




HT16




HT17




HT18



HT19




HT20




-continued


HT21




HT22




HT23



HT24




HT25




-continued


HT26




HT27




HT28




HT29




HT30




HT31




HT32




HT33




-continued
HT34


HT35




HT36




HT37



HT38



HT39



[0223] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer

may be in a range of about 100 Å to about 9,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, for example about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole

injection layer, and the hole transport layer are within any of these ranges, satisfactory (or suitable) hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0224] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer; and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include any of the suitable materials as described above.

[0225] The hole transport region may further include, in addition to the materials described above, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0226] The charge-generation material may be, for example, a p-dopant.

[0227] In one embodiment, the p-dopant may have a LUMO of -3.5 eV or less.

[0228] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments of the present disclosure are not limited thereto.

[0229] For example, the p-dopant may include at least one selected from:


[0230] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and/or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

[0231] a metal oxide, such as tungsten oxide and/or molybdenum oxide;

[0232] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and


[0233] a compound represented by Formula 221 below,

[0234] but embodiments of the present disclosure are not limited thereto:



-continued

Formula 221



[0235] In Formula 221,

[0236] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, provided that at least one of R_{221} to R_{223} has at least one substituent selected from a cyano group, $-F$, $-Cl$, $-Br$, $-I$, a C_1 - C_{20} alkyl group substituted with $-F$, a C_1 - C_{20} alkyl group substituted with $-Cl$, a C_1 - C_{20} alkyl group substituted with Br , and a C_1 - C_{20} alkyl group substituted with $-I$.

[0237] When the organic light-emitting device **10** is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red light-emitting material, a green light-emitting material, and a blue light-emitting material, in which the two or more materials are mixed with each other in a single layer to emit white light.

[0238] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.


[0239] An amount of the dopant in the emission layer may be in a range of about 0.01 parts to about 15 parts by weight, based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.

[0240] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.


[0241] The host may include the heterocyclic compound represented by Formula 1.

[0242] In one or more embodiments, the host may include a silicon-based compound, a carbazole-based compound, or a phosphine oxide-based compound.


[0243] For example, the host may be one of Compounds H101 to H106 below:




H101



H102




H103




H104

-continued


H105



H106



[0244] In one or more embodiments, the host may include a compound represented by Formula 301 below:



Formula 301

In Formula 301,

[0245] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0246] $xb11$ may be 1, 2, or 3,

[0247] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0248] $xb1$ may be an integer from 0 to 5,

[0249] R_{301} may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{301})(Q_{302})(Q_{303}), —N(Q_{301})

(Q₃₀₂), —B(Q₃₀₁)(Q₃₀₂), —C(=O)(Q₃₀₁), —S(=O)₂(Q₃₀₁), and —P(=O)(Q₃₀₁)(Q₃₀₂),

[0250] xb21 may be an integer from 1 to 5, and

[0251] Q₃₀₁ to Q₃₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0252] In one embodiment, Ar₃₀₁ in Formula 301 may be selected from:

[0253] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and

[0254] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0255] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0256] When xb11 in Formula 301 is two or more, two or more Ar₃₀₁(s) may be linked via a single bond.

[0257] In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or Formula 301-2:

[0258] In Formulae 301-1 and 301-2,

[0259] A₃₀₁ to A₃₀₄ may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group,

[0260] X₃₀₁ may be O, S, or N—[(L₃₀₄)_{xb4}—R₃₀₄],

[0261] R₃₁₁ to R₃₁₄ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0262] xb22 and xb23 may each independently be 0, 1, or 2,

[0263] L₃₀₁, xb1, R₃₀₁, and Q₃₁ to Q₃₃ may be the same as described herein,

[0264] definitions of L₃₀₂ to L₃₀₄ may each independently be the same as that provided in connection with L₃₀₁,


[0265] definitions of xb2 to xb4 may each independently be the same as that provided in connection with xb1, and

[0266] definitions of R₃₀₂ to R₃₀₄ may each independently be the same as that provided in connection with R₃₀₁.


[0267] For example, L₃₀₁ to L₃₀₄ in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0268] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene

Formula 301-1



Formula 301-2



group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0269] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazoyl group, a thiazoyl group, an isothiazoyl group, an oxazolyl group, an isoxazolyl group, a thiadiazoyl group, an oxadiazoyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazoyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group; and

isothiazoyl group, an oxazolyl group, an isoxazolyl group, a thiadiazoyl group, an oxadiazoyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazoyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazoyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}), and

[0270] Q_{31} to Q_{33} are the same as described herein.

[0271] In one embodiment, R_{301} to R_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0272] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazoyl group, a thiazoyl group, an isothiazoyl group, an oxazolyl group, an isoxazolyl group, a thiadiazoyl group, an oxadiazoyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazoyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group; and

[0273] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazoyl group, a thiazoyl group, an isothiazoyl group, an oxazolyl group, an isoxazolyl group, a thiadiazoyl group, an oxadiazoyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazoyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group; and

group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}) and —P(=O)(Q_{31})(Q_{32}), and


[0274] Q_{31} to Q_{33} are the same as described herein.

[0275] In one embodiment, the host may include an alkaline earth-metal complex.


[0276] For example, the host may be selected from a Be complex (e.g., Compound H55), a Mg complex, and a Zn complex.

[0277] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, but embodiments of the present disclosure are not limited thereto:


-continued




H2




H3




H4




H5



H6




H7




H1

-continued




H8

-continued




HT13


HT9




HT14




HT10




HT15




HT11




HT16



HT12




HT17




-continued

HT18




-continued


HT22




HT19




HT23




HT20



HT21




HT24




-continued

HT25




-continued


HT27




HT28




HT26



HT29




-continued




HT30

-continued




HT35


HT31




HT32




HT36




HT33




HT37



HT34




HT38




-continued

HT39




-continued


HT41




HT40




HT42



HT43




-continued




HT44


-continued




HT48




HT45




HT49




HT50




HT46




HT51



HT47




HT52




HT53

-continued

HT54



HT55




[0278] The phosphorescent dopant may include an organometallic complex represented by Formula 401:

Formula 401

m(L₄₀₁)_{xc1}(L₄₀₂)_{xc2}.

Formula 402



[0279] In Formulae 401 and 402,

[0280] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0281] L₄₀₁ may be selected from ligands represented by Formula 402, and xc1 may be 1, 2, or 3, wherein, when xc1 is two or more, two or more L₄₀₁(S) may be identical to or different from each other,

[0282] L₄₀₂ may be an organic ligand, and xc2 may be an integer from 0 to 4, wherein, when xc2 is two or more, two or more L₄₀₂(s) may be identical to or different from each other,

[0283] X₄₀₁ to X₄₀₄ may each independently be nitrogen or carbon,

[0284] X₄₀₁ and X₄₀₃ may be linked via a single bond or a double bond, and X₄₀₂ and X₄₀₄ may be linked via a single bond or a double bond,

[0285] A₄₀₁ and A₄₀₂ may each independently be a C₅-C₆₀ carbocyclic group or a C₁-C₆₀ heterocyclic group,

[0286] X₄₀₅ may be a single bond, *-O-*, *-S-*, *-C(=O)-*, *-N(Q₄₁₁)-*, *-C(Q₄₁₁)(Q₄₁₂)-, *-C(Q₄₁₁)=C(Q₄₁₂)-, *-C(Q₄₁₁)=*, or *-C(Q₄₁₁)=*, wherein Q₄₁₁ and Q₄₁₂ may each independently be hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0287] X₄₀₆ may be a single bond, O, or S,

[0288] R₄₀₁ and R₄₀₂ may each independently be selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₂₀ alkyl group, a substituted or unsubstituted C₁-C₂₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), -N(Q₄₀₁)(Q₄₀₂), -B(Q₄₀₁)(Q₄₀₂), -C(=O)(Q₄₀₁), -S(=O)₂(Q₄₀₁), and -P(=O)(Q₄₀₁)(Q₄₀₂), wherein Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a C₆-C₂₀ aryl group, and a C₁-C₂₀ heteroaryl group,

[0289] xc11 and xc12 may each independently be an integer from 0 to 10, and

[0290] * and *' in Formula 402 each indicate a binding site to M in Formula 401.

[0291] In one embodiment, A₄₀₁ and A₄₀₂ in Formula 402 may each independently be a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0292] In one or more embodiments, in Formula 402, i) X₄₀₁ may be nitrogen, and X₄₀₂ may be carbon, or ii) X₄₀₁ and X₄₀₂ may both be nitrogen.

[0293] In one or more embodiments, R₄₀₁ and R₄₀₂ in Formula 402 may each independently be selected from:

[0294] hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino

group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0295] a C₁-C₂₀ alkyl group and a C₁-C₂₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0296] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0297] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0298] —Si(Q₄₀₁)(Q₄₀₂)(Q₄₀₃), —N(Q₄₀₁)(Q₄₀₂), —B(Q₄₀₁)(Q₄₀₂), —C(=O)(Q₄₀₁), —

[0299] S(=O)₂(Q₄₀₁), and —P(=O)(Q₄₀₁)(Q₄₀₂), and





[0300] Q₄₀₁ to Q₄₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0301] In one or more embodiments, when xc1 in Formula 401 is two or more, two or more L₄₀₁(s) may optionally be linked via a linking group X₄₀₇, or two A₄₀₂(s) in two or more L₄₀₁(S) may optionally be linked via a linking group X₄₀₈ (see e.g., Compounds PD1 to PD4 and PD7). X₄₀₇ and X₄₀₈ may each independently be a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q₄₁₃)*, *—C(Q₄₁₃)(Q₄₁₄)*, or *—C(Q₄₁₃)=C(Q₄₁₄)* (wherein Q₄₁₃ and Q₄₁₄ may each independently be hydrogen, deuterium, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), but embodiments of the present disclosure are not limited thereto.







[0302] L₄₀₂ in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L₄₀₂ may be selected from halogen, diketone (e.g., acetylacetonate), carboxylic acid (e.g., picolinate), —C(=O), isonitrile, —CN,

and phosphorus (e.g., phosphine and/or phosphite), but embodiments of the present disclosure are not limited thereto.

[0303] In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, but embodiments of the present disclosure are not limited thereto:




-continued




-continued

PD5




PD11

PD6




PD12

PD7



PD13


PD8



PD14






PD9

PD10




PD15

-continued




-continued

PD16




PD21

PD17




PD22

PD18




PD23

PD19




PD24

PD20



[0304] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0305] The fluorescent dopant may include a compound represented by Formula 501 below:



[0306] In Formula 501,

[0307] Ar₅₀₁ may be a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

[0308] L₅₀₁ to L₅₀₃ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0309] xd1 to xd3 may each independently be an integer from 0 to 3,

[0310] R₅₀₁ and R₅₀₂ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0311] xd4 may be an integer from 1 to 6.

[0312] In one embodiment, Ar₅₀₁ in Formula 501 may be selected from:

[0313] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene

group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

[0314] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0315] In one or more embodiments, L₅₀₁ to L₅₀₃ in Formula 501 may each independently be selected from:

[0316] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

[0317] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chryseryl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0318] In one or more embodiments, R₅₀₁ and R₅₀₂ in Formula 501 may each independently be selected from:


[0319] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0320] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}), and


[0321] Q_{31} to Q_{33} may be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0322] In one or more embodiments, xd4 in Formula 501 may be 2, but embodiments of the present disclosure are not limited thereto.

[0323] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:




-continued




FD2

FD3




-continued




FD4

-continued





FD6

FD7




FD8

FD5




-continued

FD9




-continued


FD13




FD10




FD14




FD11




FD15




FD12



FD16




-continued




FD17

-continued




FD20

FD18





FD19


FD21



FD22




[0324] In one or more embodiments, the fluorescent dopant may be selected from the following compounds, but embodiments of the present disclosure are not limited thereto:




DPVBi


-continued




DPAVBi



TBPe





DCM



DCJTb

Coumarin 6

C545T



[0325] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure including a plurality of layers including a plurality of different materials.

[0326] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

[0327] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein for each structure, constituting layers are sequentially stacked from an emission layer. However, embodiments of the structure of the electron transport region are not limited thereto.

[0328] The electron transport region (e.g., a buffer layer, a hole blocking layer, an electron control layer, and/or an electron transport layer in the electron transport region) may

include a metal-free compound containing at least one 7 electron-depleted nitrogen-containing ring.

[0329] As used herein, the “7 electron-depleted nitrogen-containing ring” may refer to a C_1 - C_{60} heterocyclic group including at least one $*-N=*$ moiety as a ring-forming moiety.

[0330] For example, the “7 electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered heteromonocyclic group including at least one $*-N=*$ moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each including at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropolycyclic group in which at least one 5-membered to 7-membered heteromonocyclic group including at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0331] Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an

isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazole, an imidazopyridine, an imidazopyrimidine, and an azacarbazole, but embodiments of the present disclosure are not limited thereto.

[0332] For example, the electron transport region may include a compound represented by Formula 601:



[0333] In Formula 601,

[0334] Ar_{601} may be a substituted or unsubstituted $\text{C}_5\text{-C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heterocyclic group,

[0335] xe11 may be 1, 2, or 3,

[0336] L_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkylene group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenylene group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylene group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0337] xe1 may be an integer from 0 to 5,

[0338] R_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a substituted or unsubstituted $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a substituted or unsubstituted $\text{C}_1\text{-C}_{10}$ heterocycloalkenyl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryl group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ aryloxy group, a substituted or unsubstituted $\text{C}_6\text{-C}_{60}$ arylthio group, a substituted or unsubstituted $\text{C}_1\text{-C}_{60}$ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $-\text{C}(=\text{O})(\text{Q}_{601})$, $-\text{S}(=\text{O})_2(\text{Q}_{601})$, and $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$,

[0339] Q_{601} to Q_{603} may each independently be a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0340] xe21 may be an integer from 1 to 5.

[0341] In one embodiment, at least one selected from xe11 number of $\text{Ar}_{601}(\text{s})$ and xe21 number of $\text{R}_{601}(\text{S})$ may include the 7 electron-depleted nitrogen-containing ring.

[0342] In one embodiment, Ar_{601} in Formula 601 may be selected from:

[0343] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzox-

azole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0344] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_1\text{-C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$, and


[0345] Q_{31} to Q_{33} may each independently be selected from a $\text{C}_1\text{-C}_{10}$ alkyl group, a $\text{C}_1\text{-C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0346] When xe11 in Formula 601 is two or more, two or more $\text{Ar}_{601}(\text{s})$ may be linked via a single bond.

[0347] In one or more embodiments, Ar_{601} in Formula 601 may be an anthracene group.

[0348] In one or more embodiments, the compound represented by Formula 601 may be represented by Formula 601-1:

Formula 601-1



[0349] In Formula 601-1,

[0350] X_{614} may be N or C(R_{614}), X_{615} may be N or C(R_{615}), X_{616} may be N or C(R_{616}), and at least one of X_{614} to X_{616} may be N,

[0351] definitions of L_{611} to L_{613} may each independently be the same as that provided in connection with L_{601} ,

[0352] definitions of xe611 to xe613 may each independently be the same as that provided in connection with xe1 ,

[0353] definitions of R_{611} to R_{613} may each independently be the same as that provided in connection with R_{601} ,

[0354] R_{614} to R_{616} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0355] In one embodiment, L_{601} and L_{611} to L_{613} in Formulae 601 and 601-1 may each independently be selected from:

[0356] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group;

[0357] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an

azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group,

[0358] but embodiments of the present disclosure are not limited thereto.

[0359] In one or more embodiments, $xe1$ and $xe611$ to $xe613$ in Formulae 601 and 601-1 may each independently be 0, 1, or 2.


[0360] In one or more embodiments, R_{601} and R_{611} to R_{613} in Formulae 601 and 601-1 may each independently be selected from:

[0361] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;


[0362] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0363] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂), and
 [0364] Q₆₀₁ and Q₆₀₂ are the same as described herein.
 [0365] The electron transport region may include at least one compound selected from Compounds ET1 to ET36, but embodiments of the present disclosure are not limited thereto:


ET1



ET2




ET3




-continued

ET4




-continued


ET7




ET5




ET8



ET6




-continued




ET9

-continued





ET11

ET12




ET10

ET13




-continued




ET14

-continued




ET17

ET15





ET18


ET16



ET19




-continued



ET20



-continued



ET23



ET24

ET21




ET22

ET25




-continued

ET26




-continued


ET29




ET27




ET30




ET28



ET31




-continued




ET32

-continued




ET35

ET33






ET36



[0366] In one or more embodiments, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(bi-phenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:

ET34

Alq₃



[0367] A thickness of each of the buffer layer, the hole blocking layer, and/or the electron control layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of each of the buffer layer, the hole blocking layer, and/or the electron control layer is within the range described above, the buffer layer, the hole blocking layer, and/or the electron control layer may have satisfactory (or suitable) hole blocking or electron controlling characteristics without a substantial increase in driving voltage.


[0368] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory (or suitable) electron transport characteristics without a substantial increase in driving voltage.

[0369] The electron transport region (e.g., the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.


[0370] The metal-containing material may include at least one selected from an alkali metal complex and an alkaline earth metal complex. The alkali metal complex may include a metal ion selected from a Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion; and the alkaline earth metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, a Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline

earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenyloxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazoole, a hydroxy diphenylthiadiazole, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0371] For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or Compound ET-D2:



ET-D1



ET-D2

[0372] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0373] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0374] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof.

[0375] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs, but embodiments of the present disclosure are not limited thereto.

[0376] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0377] The rare earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

[0378] The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be an oxide and/or a halide (e.g., fluoride, chloride, bromide, and/or iodide) of the alkali metal, the alkaline earth metal, and the rare earth metal, respectively.

[0379] For example, the alkali metal compound may be selected from alkali metal oxides (such as Li_2O , Cs_2O and/or K_2O) and alkali metal halides (such as LiF , NaF , CsF , KF , LiI , NaI , CsI , KI and/or RbI). In one embodiment, the alkali metal compound may be selected from LiF , Li_2O , NaF , LiI , NaI , CsI , and KI , but embodiments of the present disclosure are not limited thereto.

[0380] The alkaline earth-metal compound may be selected from alkaline earth-metal oxides, such as BaO , SrO , CaO , $\text{Ba}_{1-x}\text{Sr}_x\text{O}$ ($0 < x < 1$) and/or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$). In one embodiment, the alkaline earth-metal compound may be selected from BaO , SrO , and CaO , but embodiments of the present disclosure are not limited thereto.

[0381] The rare earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In one embodiment, the rare earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 , but embodiments of the present disclosure are not limited thereto.

[0382] The alkali metal complex, the alkaline earth metal complex, and the rare earth metal complex may respectively include an ion of an alkali metal, an alkaline earth metal, and a rare earth metal as described above. A ligand coordinated with the metal ion of the alkali metal complex, the alkaline earth metal complex, or the rare earth metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenyloxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazole, a hydroxy diphenylthiadiazole, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0383] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combination thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, the alkali metal, the alkaline earth metal, the rare earth metal, the alkali metal compound, the alkaline earth-metal compound, the rare earth metal compound, the alkali metal complex, the alkaline earth-metal complex, the rare earth metal complex, or a combination thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0384] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory (or suitable) electron injection characteristics without a substantial increase in driving voltage.

[0385] The second electrode 190 may be disposed on the organic layer 150 having the structure according to embodiments of the present disclosure. The second electrode 190 may be a cathode, which is an electron injection electrode, and in this regard, a material for forming the second electrode 190 may be selected from metals, alloys, electrically

conductive compounds, and combinations thereof, which have a relatively low work function.

[0386] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al-Li), calcium (Ca), magnesium-indium (Mg-In), magnesium-silver (Mg-Ag), ITO, and IZO, but embodiments of the present disclosure are not limited thereto. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0387] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[0388] Hereinbefore, the organic light-emitting device has been described with reference to the drawing, but embodiments of the present disclosure are not limited thereto.

[0389] Layers constituting the hole transport region, the emission layer, and layers constituting the electron transport region may each independently be formed in the respective region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0390] When any of the layers constituting the hole transport region, the emission layer, and layers constituting the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100° C. to about 500° C., at a vacuum degree of about 10^{-8} torr to about 10^{-3} torr, and at a deposition rate of about 0.01 Å/sec to about 100 Å/sec, taking into account the characteristics of the material to be included in the layer to be formed, and the desired structure of the layer to be formed.

[0391] When any of the layers constituting the hole transport region, the emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C., taking into account the characteristics of the material to be included in the layer to be formed, and the desired structure of the layer to be formed.

[0392] The term “ $\text{C}_1\text{-C}_{60}$ alkyl group” as used herein may refer to a linear or branched aliphatic saturated hydrocarbon monovalent group having 1 to 60 carbon atoms, and non-limiting examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “ $\text{C}_1\text{-C}_{60}$ alkylene group” as used herein may refer to a divalent group having the same structure as the $\text{C}_1\text{-C}_{60}$ alkyl group.

[0393] The term “ $\text{C}_2\text{-C}_{60}$ alkenyl group” as used herein may refer to a hydrocarbon group having at least one carbon-carbon double bond in the middle and/or at either terminus of the $\text{C}_2\text{-C}_{60}$ alkyl group, and non-limiting examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “ $\text{C}_2\text{-C}_{60}$ alkenylene group” as used herein may refer to a divalent group having the same structure as the $\text{C}_2\text{-C}_{60}$ alkenyl group.

[0394] The term “ $\text{C}_2\text{-C}_{60}$ alkynyl group” as used herein may refer to a hydrocarbon group having at least one carbon-carbon triple bond in the middle and/or at either terminus of the $\text{C}_2\text{-C}_{60}$ alkyl group, and non-limiting examples thereof include an ethynyl group and a propynyl

group. The term “C₂-C₆₀ alkynylene group” as used herein may refer to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0395] The term “C₁-C₆₀ alkoxy group” as used herein may refer to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and non-limiting examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0396] The term “C₃-C₁₀ cycloalkyl group” as used herein may refer to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and non-limiting examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein may refer to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0397] The term “C₁-C₁₀ heterocycloalkyl group” as used herein may refer to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and non-limiting examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein may refer to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0398] The term “C₃-C₁₀ cycloalkenyl group” as used herein may refer to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof, but does not have aromaticity; and non-limiting examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein may refer to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0399] The term “C₁-C₁₀ heterocycloalkenyl group” as used herein may refer to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring, but does not have aromaticity. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranyl group, and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein may refer to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0400] The term “C₆-C₆₀ aryl group” as used herein may refer to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein may refer to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each independently include two or more rings, the respective rings may be fused (condensed) to each other.

[0401] The term “C₁-C₆₀ heteroaryl group” as used herein may refer to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein may refer to a divalent group having a carbocyclic aromatic system that has at least one heteroatom

selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each independently include two or more rings, the respective rings may be condensed (fused) with each other.

[0402] The term “C₆-C₆₀ aryloxy group” as used herein may refer to a group represented by —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group used herein may refer to a group represented by —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0403] The term “monovalent non-aromatic condensed polycyclic group” as used herein may refer to a monovalent group having two or more rings condensed (fused) with each other, only carbon atoms as ring-forming atoms (e.g., having 8 to 60 carbon atoms), and no overall aromaticity in its entire molecular structure. A non-limiting example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” used herein, may refer to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0404] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein may refer to a monovalent group having two or more rings condensed (fused) to each other, at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, and carbon atoms as the remaining ring-forming atoms (e.g., having 1 to 60 carbon atoms), and no overall aromaticity in its entire molecular structure. A non-limiting example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein may refer to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0405] The term “C₅-C₆₀ carbocyclic group” as used herein may refer to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atoms are carbon atoms only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring (such as benzene), a monovalent group (such as a phenyl group), or a divalent group (such as a phenylene group). In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0406] The term “C₁-C₆₀ heterocyclic group” as used herein may refer to a group having the same structure as a C₁-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used, in addition to carbon atoms (e.g., the number of carbon atoms may be in a range of 1 to 60).

[0407] At least one substituent of the substituted C₅-C₆₀ carbocyclic group, the substituted C₁-C₆₀ heterocyclic group, the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group,

the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0408] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

[0409] a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{11})(Q_{12})(Q_{13}), —N(Q_{11})(Q_{12}), —B(Q_{11})(Q_{12}), —C(=O)(Q_{11}), —S(=O)₂(Q_{11}), and —P(=O)(Q_{11})(Q_{12});

[0410] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0411] a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{21})(Q_{22})(Q_{23}), —N(Q_{21})(Q_{22}), —B(Q_{21})(Q_{22}), —C(=O)(Q_{21}), —S(=O)₂(Q_{21}), and —P(=O)(Q_{21})(Q_{22}); and

[0412] —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}), and

[0413] Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} may each independently be selected from hydrogen, deuterium, —F,

—Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0414] The term “Ph” as used herein may refer to a phenyl group, the term “Me” as used herein may refer to a methyl group, the term “Et” as used herein may refer to an ethyl group, the term “ter-Bu” or “Bu^t,” as used herein, may refer to a tert-butyl group, the term “OMe” as used herein may refer to a methoxy group, and “D” as used herein may refer to deuterium.

[0415] The term “biphenyl group” as used herein may refer to a “phenyl group substituted with a phenyl group. For example, the “biphenyl group” may be referred to as a “substituted phenyl group” having a “ C_6 - C_{60} aryl group” as a substituent.

[0416] The term “terphenyl group” as used herein may refer to a “phenyl group substituted with a biphenyl group. For example, the “terphenyl group” may be referred to as a “phenyl group” having, as a substituent, a “ C_6 - C_{60} aryl group substituted with a C_6 - C_{60} aryl group.”

[0417] * and *¹ as used herein, unless defined otherwise, each refers to a binding site to a neighboring atom in a corresponding formula.


[0418] Hereinafter, a compound according to one or more embodiments of the present disclosure, and an organic light-emitting device including the compound, will be described in more detail with reference to Synthesis Examples and Examples. The expression “B was used instead of A” used in describing Synthesis Examples and Examples may refer to an identical number of molar equivalents of A being used in place of molar equivalents of B.

SYNTHESIS EXAMPLES


Synthesis Example 1

Synthesis of Compound 1

[0419]



6267-02-3




1) Synthesis of Intermediate 1-1

[0420] 9,9-dimethyl-9,10-dihydroacridine (CAS# 6267-02-3) was reacted with nBuLi, and then reacted with dichlorophenylphosphine, to obtain Intermediate 1-1. Intermediate 1-1 was identified by liquid chromatography-mass spectrometry (LC-MS). $C_{36}H_{33}N_2P$: M+1 525.4.

2) Synthesis of Compound 1

[0421] 3.2 g of Intermediate 1-1 was dissolved in 30 mL of methylene chloride (MC), and 10 mL of peroxide was added thereto dropwise. Then, the resultant mixture was stirred at room temperature for 2 hours. The obtained reaction mixture was extracted by using MC. Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 2.6 g (yield: 79%) of Compound 1. Compound 1 was identified by LC-MS and 1H -NMR.



Synthesis of Intermediate 2-1

[0423] 9,9-diphenyl-9,10-dihydroacridine (CAS# 20474-15-1) was reacted with nBuLi, and then reacted with dichlorophenylphosphine, to obtain Intermediate 2-1. Intermediate 2-1 was identified by LC-MS.

[0424] $C_{56}H_{41}N_2P$: M+1 773.4.

Synthesis of Compound 2



[0425] 2.6 g of Intermediate 2-1 was dissolved in 30 mL of MC, and 10 mL of peroxide was added thereto dropwise. Then, the resultant mixture was stirred at room temperature

for 2 hours. The obtained reaction mixture was extracted by using MC. Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 2.2 g (yield: 81%) of Compound 2. Compound 2 was identified by LC-MS and ^1H -NMR.


Synthesis Example 3

Synthesis of Compound 3

[0426]



-continued



Synthesis of Intermediate 3-1

[0427] 10,10-diphenyl-5,10-dihydrobenzo [b,e][1,4]azaciline (CAS# 3508-62-1) was reacted with nBuLi, and then reacted with dichlorophenylphosphine, to obtain Intermediate 3-1. Intermediate 3-1 was identified by LC-MS.

[0428] C₅₄H₄₁N₂PSi₂; M+1 805.1.


Synthesis of Compound 3

[0429] 1.9 g of Intermediate 3-1 was dissolved in 30 mL of MC, and 10 mL of peroxide was added thereto dropwise. Then, the resultant mixture was stirred at room temperature for 2 hours. The obtained reaction mixture was extracted by using MC. Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 1.4 g (yield: 69%) of Compound 3. Compound 3 was identified by LC-MS and ^1H -NMR.

Synthesis Example 4

Synthesis of Compound 7

[0430]



Synthesis of Intermediate 7-1

[0431] 3-bromopyridine was reacted with isopropylmagnesium chloride, and then reacted with dichlorophenylphosphine, to obtain phenyl(pyridine-3-yl)phosphinic chloride (Intermediate A). 10-(9H-carbazole-3-yl)-9,9-dimethyl-9,10-dihydroacridine (CAS# 1821228-18-5) was reacted with nBuLi to obtain lithium 3-(9,9-dimethylacridine-10(9H)-yl)carbazolide (Intermediate B). Intermediate A was reacted with Intermediate B to obtain Intermediate 7-1. Intermediate 7-1 was identified by LC-MS.

[0432] $C_{38}H_{30}N_3P$: M+1 560.7.

Synthesis of Compound 7




[0433] 3.3 g of Intermediate 7-1 was dissolved in 30 mL of MC, and 10 mL of peroxide was added thereto dropwise. Then, the resultant mixture was stirred at room temperature for 2 hours. The obtained reaction mixture was extracted by using MC.

[0434] Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 2.2 g (yield: 66%) of Compound 7. Compound 7 was identified by LC-MS and 1H -NMR.

Synthesis Example 5

Synthesis of Compound 8

[0435]



Synthesis of Intermediate 8-1

[0436] 10-(9H-carbazole-3-yl)-9,9-dimethyl-9,10-dihydroacridine (CAS# 1821228-18-5) was reacted with nBuLi, and then reacted with dichlorophenylphosphine, to obtain Intermediate 8-1. Intermediate 8-1 was identified by LC-MS.

[0437] $C_{39}H_{31}N_2P$: M+1 559.3.


Synthesis of Compound 8

[0438] 2.9 g of Intermediate 8-1 was dissolved in 30 mL of MC, and 10 mL of peroxide was added thereto dropwise. Then, the resultant mixture was stirred at room temperature for 2 hours. The obtained reaction mixture was extracted by using MC. Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 2.6 g (yield: 87%) of Compound 8. Compound 8 was identified by LC-MS and 1H -NMR.

Synthesis Example 6

Synthesis of Compound 11

[0439]



[0440] Synthesis of Intermediate 11-1

[0441] 9H-3,9'-bicarbazole (CAS# 18628-07-4) was reacted with nBuLi, and then reacted with dichlorophenylphosphine, to obtain Intermediate 11-1. Intermediate 11-1 was identified by LC-MS.

[0442] C₃₆H₂₅N₂P: M+1 517.33.


Synthesis of Compound 11

[0443] 3.6 g of Intermediate 11-1 was dissolved in 30 mL of MC, and 10 mL of peroxide was added thereto dropwise. Then, the resultant mixture was stirred at room temperature for 2 hours. Then, the obtained reaction mixture was extracted by using MC. Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 3.1 g (yield: 83%) of Compound 11. Compound 11 was identified by LC-MS and ¹H-NMR.

Synthesis Example 7

Synthesis of Compound 27

[0444]



Synthesis of Compound 27

[0445] 2.7 g of Intermediate 3-1 was dissolved in 30 mL of MC, and then reacted with dichlorophenylphosphine.

Then, 0.3 g of sulfur was added thereto dropwise. Then, the resultant mixture was stirred at room temperature for 2 hours. Then, the obtained reaction mixture was extracted by using MC. Then, an organic layer collected therefrom was dried by using magnesium sulfate, and a solvent was evaporated therefrom. Then, a residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 2.2 g (yield: 76%) of Compound 27. Compound 27 was identified by LC-MS and ^1H -NMR.

[0446] ^1H NMR and MS/FAB results of compounds synthesized in Synthesis Examples 1 to 8 are shown in Table 1 below.

[0447] Synthesis methods of compounds other than the compounds shown in Table 1 below should become apparent those of ordinary skill in the art by referring to the synthesis mechanisms and source materials described above.

TABLE 1

Compound	^1H NMR (CDCl_3 , 400 MHz)	MS/FAB	
		Found [M + 1]	calc.
1	7.31 (d, 2H), 7.27-7.23 (m, 3H), 7.18-7.14 (m, 12H), 6.92 (t, 4H), 1.65 (s, 12H)	541.33	540.23
2	7.33 (d, 2H), 7.28-7.22 (m, 19H), 7.19-7.13 (m, 16H), 7.11 (d, 8H), 6.92 (t, 4H)	789.33	788.30
3	7.47 (d, 8H), 7.41-7.36 (m, 20H), 7.30-7.20 (m, 9H), 7.03 (t, 4H)	821.27	820.25
7	9.13 (s, 1H), 8.77 (d, 1H), 8.55 (d, 11H), 8.26 (d, 1H), 7.93 (d, 1H), 7.76 (d, 2H), 7.55-7.51 (m, 5H), 7.36-7.32 (m, 3H), 7.20-7.13 (m, 7H), 6.99 (t, 2H), 1.69 (s, 6H)	576.44	575.21
8	8.54 (d, 1H), 7.93 (d, 1H), 7.76 (d, 4H), 7.52 (d, 1H), 7.36-7.31 (m, 3H), 7.17-7.12 (m, 7H), 6.94 (t, 2H), 1.67 (s, 6H)	575.11	574.22
11	8.56-8.54 (m, 2H), 8.20 (d, 1H), 7.95 (d, 1H), 7.78-7.70 (m, 5H), 7.67 (s, 1H), 7.57 (d, 1H), 7.52-7.48 (m, 7H), 7.39-7.32 (m, 3H), 7.2-7.15 (m, 3H)	533.21	532.17
27	7.47-7.40 (m, 12H), 7.37-7.32 (m, 16H), 7.29-7.25 (m, 3H), 7.01 (t, 4H)	837.41	836.23

EXAMPLES

Example 1


[0448] As an anode, a Corning 15 Ω/cm^2 (500 Å) ITO glass substrate was cut to a size of 50 mm×50 mm×0.5 mm, sonicated with isopropyl alcohol and pure water, for 10 minutes in each, and then cleaned by exposure to ultraviolet rays and ozone for 10 minutes. Then, the resultant ITO glass substrate was placed in a vacuum deposition apparatus.

[0449] NPD was vacuum-deposited on the ITO glass substrate to form a hole injection layer having a thickness of 300 Å, and CzSi was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 200 Å.


[0450] Compound 1 and $\text{Ir}(\text{pmp})_3$ were co-deposited on the hole transport layer at a weight ratio of 92:8 to form an emission layer having a thickness of 250 Å.

[0451] Then, TSP01 was vacuum-deposited on the emission layer to form an electron transport layer having a thickness of 200 Å.



[0452] LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form a cathode having a thickness of 100 Å, thereby completing the manufacture of an organic light-emitting device.



NPD



CzSi

 $\text{Ir}(\text{pmp})_3$ 

TSP01

Example 2

[0453] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in

Example 1, except that Compound 2 was used instead of Compound 1 in forming an emission layer.

Example 3

[0454] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in Example 1, except that Compound 3 was used instead of Compound 1 in forming an emission layer.

Example 4

[0455] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in Example 1, except that Compound 7 was used instead of Compound 1 in forming an emission layer.

Example 5

[0456] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in Example 1, except that Compound 8 was used instead of Compound 1 in forming an emission layer.

Example 6


[0457] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in Example 1, except that Compound 11 was used instead of Compound 1 in forming an emission layer.

Example 7

[0458] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in Example 1, except that Compound 27 was used instead of Compound 1 in forming an emission layer.

Comparative Example 1

[0459] An organic light-emitting device was manufactured in the same (or substantially the same) manner as in Example 1, except that BCPO was used instead of Compound 1 in forming an emission layer.




BCPO

Comparative Examples 2 to 4

[0460] Organic light-emitting devices were manufactured in the same (or substantially the same) manner as in Example 1, except that Compounds 1 to 3 were respectively used instead of Compound 1 in forming an emission layer.


Comparative Compound 1

[0461]




Comparative Compound 2

[0462]



Comparative Compound 3

[0463]



Evaluation Example 1

[0464] The driving voltage, luminescent efficiency, and color coordinates of each of the organic light-emitting devices manufactured according to Examples 1 to 7 and Comparative Examples 1 to 4 were measured by using a Keithley SMU 236 and a luminance meter PR650 at 600 nit. Results thereof are shown in Table 2 below.

TABLE 2

	Emission layer	Driving voltage (V)	Current density (mA/cm ²)	Maximum quantum efficiency (%)	Emission color
Example 1	Compound 1	4.2	5.5	20.5	blue
Example 2	Compound 2	4.3	5.5	19.9	blue
Example 3	Compound 3	4.4	5.5	18.8	blue
Example 4	Compound 7	3.8	5.5	21	blue
Example 5	Compound 8	3.7	5.5	22.1	blue
Example 6	Compound 11	3.8	5.5	19.9	blue
Example 7	Compound 27	4.4	5.5	18.9	blue
Comparative Example 1	BCPO	4.9	5.5	14.9	blue
Comparative Example 2	Comparative Compound 1	4.7	5.5	4.4	blue
Comparative Example 3	Comparative Compound 2	4.9	5.5	14.4	blue
Comparative Example 4	Comparative Compound 3	5.5	5.5	13.3	blue




[0465] Referring to Table 2, it can be seen that the organic light-emitting devices of Examples 1 to 7 exhibited lower driving voltage, as compared with the organic light-emitting devices of Comparative Examples 1 to 4. Also, it can be seen that the organic light-emitting devices of Examples 1 to 7 exhibited excellent characteristics in terms of luminescent efficiency, as compared with the organic light-emitting devices of Comparative Examples 1 to 4.

Example 8



[0466] As an anode, a Corning 15 Ω/cm^2 (1,200 Å) ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm, sonicated with isopropyl alcohol and pure water, for 5 minutes in each, and then cleaned by exposure to ultraviolet rays and ozone for 30 minutes. Then, the resultant ITO glass substrate was placed in a vacuum deposition apparatus.

[0467] α -NPD was vacuum-deposited on the ITO glass substrate to form a hole injection layer having a thickness of 300 Å, and TCTA as a hole transport compound was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 200 Å. CzSi as a hole transport layer compound was vacuum-deposited on the hole transport layer to form a layer having a thickness of 100 Å. Compound H101 and Compound 3 were co-deposited on the CzSi layer at a weight ratio of 90:10 to form an emission layer having a thickness of 200 Å. Then, Compound H101 as an electron transport layer compound was formed to a thickness of 200 Å, and TPBI as an electron injection layer compound was deposited to a thickness of 300 Å.


[0468] LiF, which is an alkali metal halide, was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited to form a cathode electrode having a thickness of 3,000 Å, thereby forming a LiF/Al electrode. In this manner, an organic light-emitting device was manufactured.




-continued

 α -NPD

CBP



H101



DPS

Examples 9 and 10 and Comparative Example 5

[0469] Organic light-emitting devices were manufactured in the same (or substantially the same) manner as in Example 8, except that the compounds shown in Table 3 were respectively used instead of Compound 3 in forming an emission layer.

Evaluation Example 2

[0470] The driving voltage, luminescent efficiency, and color coordinates of each of the organic light-emitting devices manufactured according to Examples 8 to 11 and Comparative Example 5 were measured by using a Keithley SMU 236 and a luminance meter PR650 at 600 nit. Results thereof are shown in Table 3 below.

TABLE 3

	Emission layer	Driving voltage (V)	Current density (mA/cm ²)	Maximum quantum efficiency (%)	Emission color
Example 8	Compound 3	6.4	2	22.5	blue
Example 9	Compound 7	6.2	2	23.9	blue
Example 10	Compound 8	6.3	2	19.3	blue
Example 11	Compound 11	6.2	2	22	blue
Comparative Example 5	DPS	7.4	2	16.7	blue

[0471] Referring to Table 3, it can be seen that the organic light-emitting devices of Examples 8 to 11 exhibited lower driving voltage, as compared with the organic light-emitting device of Comparative Example 5. Also, it can be seen that the organic light-emitting devices of Examples 8 to 11 exhibited excellent maximum quantum efficiency, as compared with the organic light-emitting device of Comparative Example 5.

[0472] An organic light-emitting device including the heterocyclic compound according to embodiments of the present disclosure may have a low driving voltage, a high efficiency, a high luminance, and a long lifespan.

[0473] It should be understood that embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0474] As used herein, the terms “use,” “using,” and “used” may be considered synonymous with the terms “utilize,” “utilizing,” and “utilized,” respectively.

[0475] In addition, the terms “substantially,” “about,” and similar terms are used as terms of approximation and not as terms of degree, and are intended to account for the inherent deviations in measured or calculated values that would be recognized by those of ordinary skill in the art.

[0476] Also, any numerical range recited herein is intended to include all sub-ranges of the same numerical precision subsumed within the recited range. For example, a range of “1.0 to 10.0” is intended to include all subranges between (and including) the recited minimum value of 1.0 and the recited maximum value of 10.0, that is, having a minimum value equal to or greater than 1.0 and a maximum value equal to or less than 10.0, such as, for example, 2.4 to 7.6. Any maximum numerical limitation recited herein is intended to include all lower numerical limitations subsumed therein and any minimum numerical limitation recited in this specification is intended to include all higher numerical limitations subsumed therein. Accordingly, Applicant reserves the right to amend this specification, including the claims, to expressly recite any sub-range subsumed within the ranges expressly recited herein.


[0477] It will be understood that when an element is referred to as being “on,” “connected to,” or “coupled to” another element, it may be directly on, connected, or coupled to the other element or one or more intervening elements may also be present. When an element is referred to as being “directly on,” “directly connected to,” or “directly coupled to” another element, there are no intervening elements present. For example, when a first element is described as being “coupled” or “connected” to a second element, the first element may be directly coupled or con-

nected to the second element or the first element may be indirectly coupled or connected to the second element via one or more intervening elements.


[0478] While one or more embodiments have been described with reference to the drawing, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present disclosure as defined by the following claims and equivalents thereof.

What is claimed is:


1. A heterocyclic compound represented by Formula 1:




Formula 1




Formula 2-1



Formula 2-2



Formula 3-1



Formula 3-2

wherein, in Formulae 1, 2-1, 2-2, 3-1, and 3-2,

X₁ is O or S,

Ar₁ and Ar₂ are each independently a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group,

a₁ and a₂ are each independently an integer from 0 to 3, b₁ and b₂ are each independently 0 or 1,

a₁, a₂, b₁, and b₂ satisfy a₁+b₁≤1 and a₂+b₂≤1,

R₁ and R₂ are each independently a group represented by Formula 2-1,

R₃ is a group represented by Formula 2-2,

A₁ to A₄ are each independently selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, and a tetrazine group,

A₅ to A₈ are each independently selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, and a tetrazine group,

Y₁ is a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉),

Y₂ is a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉),

Y₃ is a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉),

Y₄ is a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉),

Y₅ is a single bond, O, S, SO₂, N(R₁₉), C(R₁₉)(R₂₀), Si(R₁₉)(R₂₀), S(=O)(R₁₉), S(=O)₂(R₁₉), or P(=O)(R₁₉),

wherein when Y₂ is O, a group represented by Formula 2-1 and a group represented by Formula 2-2 are different from each other,

R₁₁ to R₁₄ are each independently selected from a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —S(=O)(Q₁), and —P(=O)(Q₁)(Q₂), wherein two or more R₁₁(s) are optionally linked to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, two or more R₁₂(s) are optionally linked to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, two or more R₁₃(s) are optionally linked to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group, and two or more R₁₄(s) are optionally linked to form a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group,

b₁₁ to b₁₄ are each independently an integer from 1 to 8, when Y₂ is a single bond, at least one of R₁₃ and R₁₄ is a group represented by Formula 3-1 or a group represented by Formula 3-2,

R₁₅ to R₂₀ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group,

a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1-C_{60} alkyl group, a substituted or unsubstituted C_2-C_{60} alkenyl group, a substituted or unsubstituted C_2-C_{60} alkynyl group, a substituted or unsubstituted C_1-C_{60} alkoxy group, a substituted or unsubstituted C_3-C_{10} cycloalkyl group, a substituted or unsubstituted C_1-C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3-C_{10} cycloalkenyl group, a substituted or unsubstituted C_1-C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6-C_{60} aryl group, a substituted or unsubstituted C_6-C_{60} aryloxy group, a substituted or unsubstituted C_6-C_{60} arylthio group, a substituted or unsubstituted C_1-C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-S(=O)(Q_1)$, and $-P(=O)(Q_1)(Q_2)$,

b15 to b18 are each independently an integer from 1 to 8, at least one substituent of the substituted C_5-C_{60} carbocyclic group, the substituted C_2-C_{60} heterocyclic group, the substituted C_1-C_{60} alkyl group, the substituted C_2-C_{60} alkenyl group, the substituted C_2-C_{60} alkynyl group, the substituted C_1-C_{60} alkoxy group, the substituted C_3-C_{10} cycloalkyl group, the substituted C_1-C_{10} heterocycloalkyl group, the substituted C_3-C_{10} cycloalkenyl group, the substituted C_1-C_{10} heterocycloalkenyl group, the substituted C_6-C_{60} aryl group, the substituted C_6-C_{60} aryloxy group, the substituted C_6-C_{60} arylthio group, the substituted C_1-C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group;

a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, and a C_1-C_{60} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{11})(Q_{12})(Q_{13})$, $-N(Q_{11})(Q_{12})$, $-B(Q_{11})(Q_{12})$, $-C(=O)(Q_{11})$, $-S(=O)_2(Q_{11})$, and $-P(=O)(Q_{11})(Q_{12})$;

a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

eroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{21})(Q_{22})(Q_{23})$, $-N(Q_{21})(Q_{22})$, $-B(Q_{21})(Q_{22})$, $-C(=O)(Q_{21})$, $-S(=O)_2(Q_{21})$, and $-P(=O)(Q_{21})(Q_{22})$; and

$-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

Q_1 , Q_2 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1-C_{60} alkyl group, a C_2-C_{60} alkenyl group, a C_2-C_{60} alkynyl group, a C_1-C_{60} alkoxy group, a C_3-C_{10} cycloalkyl group, a C_1-C_{10} heterocycloalkyl group, a C_3-C_{10} cycloalkenyl group, a C_1-C_{10} heterocycloalkenyl group, a C_6-C_{60} aryl group, a C_6-C_{60} aryloxy group, a C_6-C_{60} arylthio group, a C_1-C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

* indicates a binding site to a neighboring atom.

2. The heterocyclic compound of claim 1, wherein

a1, a2, b1, and b2 satisfy $a1+b1=1$ and $a2+b2=1$.

3. The heterocyclic compound of claim 1, wherein

b1 and b2 satisfy $b1+b2=0$ or $b1+b2=1$.

4. The heterocyclic compound of claim 1, wherein,

when Y_2 is O, Y_1 is a single bond, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, or $Si(R_{19})(R_{20})$.

5. The heterocyclic compound of claim 1, wherein

Ar_1 and Ar_2 are each independently a substituted or unsubstituted aromatic C_6-C_{30} carbocyclic group or a substituted or unsubstituted π electron-depleted nitrogen-containing C_2-C_{30} heterocyclic group.

6. The heterocyclic compound of claim 1, wherein

A_1 to A_4 are each independently a benzene group, a pyridine group, or a pyrimidine group, and

A_5 to A_8 are each independently a benzene group, a pyridine group, or a pyrimidine group.

7. The heterocyclic compound of claim 1, wherein


Y_1 is a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, or $Si(R_{19})(R_{20})$,

Y_2 is a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, or $Si(R_{19})(R_{20})$, and


Y_3 is a single bond, O, S, SO_2 , $N(R_{19})$, $C(R_{19})(R_{20})$, or $Si(R_{19})(R_{20})$.

8. The heterocyclic compound of claim 1, wherein the heterocyclic compound represented by Formula 1 is represented by Formula 1-1 or Formula 1-2:

Formula 1-1



Formula 1-2



wherein, in Formulae 1-1 and 1-2,

descriptions of A₁₁, A₁₂, A₂₁, A₂₂, A₃₁, and A₃₂ are each independently the same as the one provided in connection with A₁ to A₄ in Formula 1,

descriptions of Y₁₁ and Y₁₃ are each independently the same as the one provided in connection with Y₁ in Formula 1,

description of Y₁₂ is the same as the one provided in connection with Y₂ in Formula 1,

when one of Y₁₁ and Y₁₂ is O, the other one of Y₁₁ and Y₁₂ is not O,

descriptions of R₁₀₁, R₁₀₂, R₁₁₁, R₁₁₂, R₁₂₁, and R₁₂₂ are each independently the same as the one provided in connection with R₁₁ to R₁₄ in Formula 1, and

descriptions of X₁, Ar₁, and Ar₂ are respectively the same as those provided in connection with Formula 1.

9. The heterocyclic compound of claim 1, wherein

R₁₁ to R₁₄ are each independently selected from:

a group represented by Formula 3-1, a group represented by Formula 3-2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a phenyl group, an acridyl group, and a carbazolyl group; and


a phenyl group, an acridyl group, and a carbazolyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an

ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, and a C₁-C₂₀ alkoxy group.

10. The heterocyclic compound of claim 1, wherein

when Y₁ is a single bond, the group represented by Formula 2-1 is a group represented by Formula 2-2-1:

Formula 2-2-1



wherein, in Formula 2-2-1,

R₂₈ is a group represented by Formula 3-1 or a group represented by Formula 3-2, and

descriptions of R₂₁ to R₂₇ are each independently the same as the one provided in connection with R₁₁ to R₁₄ in Formula 1.

11. The heterocyclic compound of claim 10, wherein


R₂₃ is selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a phenyl group, a naphthyl group, a fluorenyl group, a carbazolyl group, a dibenzofuran group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, and

R₂₁, R₂₂, and R₂₄ to R₂₇ are each hydrogen.

12. The heterocyclic compound of claim 1, wherein the group represented by Formula 3-1 is represented by Formula 3-1-1:



Formula 3-1-1

wherein, in Formula 3-1-1,

descriptions of Y_3 , R_{15} , and R_{16} are respectively the same as those provided in connection with Formula 1, and * indicates a binding site to a neighboring atom.

13. The heterocyclic compound of claim 1, wherein R_{15} to R_{20} are each independently selected from: hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a C_1 - C_{20} alkoxy group, and a phenyl group; and

a phenyl group substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, and a C_1 - C_{20} alkoxy group.

14. An organic light-emitting device comprising:
a first electrode;
a second electrode facing the first electrode; and
an organic layer between the first electrode and the second electrode, the organic layer comprising an emission layer and at least one heterocyclic compound of claim 1.

15. The organic light-emitting device of claim 14, wherein the first electrode is an anode,
the second electrode is a cathode,
the organic layer further comprises a hole transport region, between the first electrode and the emission layer, and an electron transport region, between the emission layer and the second electrode,
the hole transport region comprises at least one layer selected from a hole injection layer, a hole transport layer, a buffer layer, an emission auxiliary layer, and an electron blocking layer, and
the electron transport region comprises at least one layer selected from a hole blocking layer, an electron transport layer, and an electron injection layer.

16. The organic light-emitting device of claim 14, wherein the emission layer comprises the heterocyclic compound.

17. The organic light-emitting device of claim 16, wherein the heterocyclic compound comprised in the emission layer is a host, and
the emission layer further comprises a phosphorescent dopant.

18. The organic light-emitting device of claim 16, wherein the heterocyclic compound comprised in the emission layer is a delayed fluorescent dopant satisfying Equation 1:

$$|E_{D, S1} - E_{D, T1}| \leq 0.3 \text{ eV},$$

Equation 1

wherein, in Equation 1,

$E_{D, S1}$ is a singlet energy level (eV) of the delayed fluorescent dopant, and

$E_{D, T1}$ is a triplet energy level (eV) of the delayed fluorescent dopant.

19. The organic light-emitting device of claim 16, wherein the heterocyclic compound comprised in the emission layer is a delayed fluorescent dopant,

the emission layer further comprises a host, and

the delayed fluorescent dopant and the host satisfy Equation 2 or Equation 3:

$$|E_{H, HOMO} - E_{D, HOMO}| \leq 0.5 \text{ eV}$$

Equation 2

$$|E_{H, HOMO} - E_{D, HOMO}| \leq 0.5 \text{ eV}$$

Equation 3

wherein, in Equations 2 and 3,

$E_{H, HOMO}$ is a highest occupied molecular orbital level of the host,


$E_{D, HOMO}$ is a highest occupied molecular orbital level of the dopant,

$E_{H, LUMO}$ is a lowest unoccupied molecular orbital level of the host, and


$E_{D, LUMO}$ is a lowest unoccupied molecular orbital level of the dopant.

20. The organic light-emitting device of claim 19, wherein the host is one of Compounds H101 to H106:

H101




H102




-continued

H103




H104




-continued

H105



H106




* * * * *


专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	US20180301638A1	公开(公告)日	2018-10-18
申请号	US15/788605	申请日	2017-10-19
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	AHN HEECHOON KO SOOBYUNG JUN MIEUN KIM SUNGBUM JEON MINA KIM YOUNGKOOK HWANG SEOKHWAN		
发明人	AHN, HEECHOON KO, SOOBYUNG JUN, MIEUN KIM, SUNGBUM JEON, MINA KIM, YOUNGKOOK HWANG, SEOKHWAN		
IPC分类号	H01L51/00 H01L51/50 C07F9/64 C09K11/02 C09K11/06 C07F9/6596 C07F9/572		
CPC分类号	H01L51/0072 H01L51/5004 C07F9/64 C09K11/025 C09K11/06 H01L51/0094 C07F9/6596 H01L51/0067 C07F9/5728 H01L51/5012 H01L51/5016 H01L2251/5376 H01L2251/552 C09K2211/1018 C07F9/65616 H01L51/0073 H01L51/0074 H01L51/50		
优先权	1020170048532 2017-04-14 KR		
外部链接	Espacenet USPTO		

摘要(译)


有机发光装置包括第一电极;面向第一电极的第二电极;在第一电极和第二电极之间的有机层中,有机层包括发光层。发光层包括至少一种式1的杂环化合物。杂环化合物可以是主体或延迟荧光掺杂剂。包含杂环化合物的有机发光装置可以具有低驱动电压,高效率,高亮度和长寿命。




Formula 1




Formula 2-1



Formula 2-2



Formula 3-1



Formula 3-2