



US 20190036044A1

(19) **United States**

(12) **Patent Application Publication**
JEON et al.

(10) **Pub. No.: US 2019/0036044 A1**
(43) **Pub. Date: Jan. 31, 2019**

(54) **ORGANOMETALLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

Publication Classification

(71) Applicant: **Samsung Display Co., Ltd., Yongin-si (KR)**

(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07F 15/00 (2006.01)
C09K 11/06 (2006.01)

(72) Inventors: **Mina JEON, Yongin-si (KR); Soobyung KO, Yongin-si (KR); Sungbum KIM, Yongin-si (KR); Heechoon AHN, Yongin-si (KR); Mieun JUN, Yongin-si (KR); Youngkook KIM, Yongin-si (KR); Seokhwan HWANG, Yongin-si (KR)**

(52) **U.S. Cl.**
CPC *H01L 51/0087* (2013.01); *H01L 51/5012* (2013.01); *C09K 11/06* (2013.01); *C07F 15/0086* (2013.01)

(21) Appl. No.: **15/952,587**

(57) **ABSTRACT**

(22) Filed: **Apr. 13, 2018**

An organometallic compound is represented by Formula 1, $M(L_1)_{n1}(L_2)_{n2}$, where M is Ir, Pt, Os, Ru, Pd, Rh, Fe, Co, Ni, Au, Ag, or Cu, L_1 is a ligand, L_2 is a ligand, $n1$ is 1 or 2, and $n2$ is 1 or 2. An organic light-emitting device includes: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and including an emission layer, where the organic layer includes at least one of the organometallic compound.

(30) **Foreign Application Priority Data**

Jul. 28, 2017 (KR) 10-2017-0096378

10

190
150
110

10

190
150
110

**ORGANOMETALLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] This application claims priority to and the benefit of Korean Patent Application No. 10-2017-0096378, filed on Jul. 28, 2017, in the Korean Intellectual Property Office, the entire content of which is incorporated herein by reference.

BACKGROUND

1. Field

[0002] One or more embodiments relate to an organometallic compound and an organic light-emitting device including the same.

2. Description of the Related Art

[0003] Organic light-emitting devices are self-emission devices that produce full-color images, and also have wide viewing angles, high contrast ratios, short response times, and excellent characteristics in terms of brightness, driving voltage, and response speed, as compared to other light-emitting devices.


[0004] An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. These excitons transit (e.g., transition or relax) from an excited state to a ground state, thereby generating light.

SUMMARY


[0005] Aspects of embodiments of the present disclosure provide a novel organometallic compound and an organic light-emitting device including the same.

[0006] Additional aspects of embodiments will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0007] An aspect of an embodiment provides an organometallic compound represented by Formula 1 below:




Formula 1



Formula 2A

-continued

Formula 2B



[0008] In Formulae 1, 2A, and 2B,

[0009] M may be Ir, Pt, Os, Ru, Pd, Rh, Fe, Co, Ni, Au, Ag, or Cu,

[0010] L₁ may be a ligand represented by Formula 2A,

[0011] n₁ may be 1 or 2, wherein, when n₁ is 2, two L₁(s) may be identical to or different from each other,

[0012] L₂ may be a ligand represented by Formula 2B,

[0013] n₂ may be 1 or 2, wherein, when n₂ is 2, two L₂(s) may be identical to or different from each other,

[0014] the sum of n₁ and n₂ may be 2 or 3,

[0015] ring Ar₁ may be a C₅-C₃₀ carbocyclic group or a C₂-C₆₀ heterocyclic group,

[0016] ring Ar₂ may be an N-containing C₂-C₆₀ heterocyclic group,

[0017] X₃ may be N or C(R₃),

[0018] X₄ may be N or C(R₄),

[0019] X₅ may be N, O, or S,

[0020] R₁ to R₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

[0021] R₃ and R₄ may optionally be linked to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group,

[0022] a₁ and a₂ may each independently be an integer from 0 to 10, wherein, when a₁ is two or more, two or more R₁(s) may be identical to or different from each other, and when a₂ is two or more, two or more R₂(s) may be identical to or different from each other,

[0023] * and *' each indicate a binding site to M in Formula 1,

[0024] at least one substituent of the substituted C₄-C₆₀ carbocyclic group, the substituted C₂-C₆₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0025] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0026] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0027] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0028] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

[0029] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0030] Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ alkyl group substituted with at least one selected from deuterium, —F, and a cyano group, a C₆-C₆₀ aryl group substituted with at least one selected from deuterium, —F, and a cyano group, a biphenyl group, and a terphenyl group.

[0031] Another aspect of an embodiment provides an organic light-emitting device that includes: a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes at least one of the organometallic compound described above.


BRIEF DESCRIPTION OF THE DRAWING

[0032] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with the accompanying drawing which is a schematic view of an organic light-emitting device according to an embodiment.

DETAILED DESCRIPTION

[0033] Reference will now be made in more detail to embodiments, examples of which are illustrated in the accompanying drawing, wherein like reference numerals refer to like elements throughout. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the accompanying drawing, to explain aspects of embodiments of the present description. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0034] An organometallic compound according to an embodiment is represented by Formula 1 below:




Formula 1


[0035] M in Formula 1 may be iridium (Ir), platinum (Pt), osmium (Os), ruthenium (Ru), palladium (Pd), rhenium (Rh), iron (Fe), cobalt (Co), nickel (Ni), gold (Au), silver (Ag), or copper (Cu).

[0036] In one embodiment, M in Formula 1 may be iridium or platinum, but embodiments of the present disclosure are not limited thereto.

[0037] In Formula 1, L₁ may be a ligand represented by Formula 2A, and n₁ may be 1, 2, or 3, wherein, when n₁ is two or more, two or more L₁(s) may be identical to or different from each other, L₂ may be a ligand represented by Formula 2B, and n₂ may be 1, 2, or 3, wherein, when n₂ is two or more, two or more L₂(s) may be identical to or different from each other:



Formula 2A



Formula 2B

[0038] The sum of n_1 and n_2 in Formula 1 may be 2 or 3.

[0039] The organometallic compound represented by Formula 1 does not have a salt form consisting of a cation and an anion and is neutral.

[0040] In one embodiment, in Formula 1, M may be platinum and the sum of n_1 and n_2 may be 2, but embodiments of the present disclosure are not limited thereto. For example, M in Formula 1 may be platinum, n_1 may be 1, and n_2 may be 1.

[0041] In one or more embodiments, in Formula 1, M may be iridium and the sum of n_1 and n_2 may be 3, but embodiments of the present disclosure are not limited thereto. For example, in Formula 1, M may be iridium, n_1 may be 2, and n_2 may be 1. For example, in Formula 1, M may be iridium, n_1 may be 1, and n_2 may be 2.

[0042] Ring Ar_1 in Formula 2A may be a C_5 - C_{30} carbocyclic group or a C_2 - C_{60} heterocyclic group.

[0043] For example, ring Ar_1 may be a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, or a quinazoline group.

[0044] In one embodiment, ring Ar_1 may be a benzene group, a pyridine group, a pyrimidine group, or a triazine group.

[0045] Ring Ar_2 in Formula 2A may be an N-containing C_2 - C_{60} heterocyclic group.

[0046] For example, ring Ar_2 may be a π electron-depleted nitrogen-containing C_2 - C_{60} heterocyclic group.

[0047] The term " π electron-depleted nitrogen-containing ring," as used herein, indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0048] For example, the " π electron-depleted nitrogen-containing ring" may be i) a 5-membered to 7-membered heteromonocyclic group having at least one $*-N=*$ moiety, ii) a heteropolycyclic group in which two or more


5-membered to 7-membered heteromonocyclic groups each having at least one $*-N=*$ moiety are condensed with each other (e.g., combined together), or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one $*-N=*$ moiety, is condensed with (e.g., combined together with) at least one C_5 - C_{60} carbocyclic group.

[0049] Examples of the π electron-depleted nitrogen-containing ring may include an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, but embodiments of the present disclosure are not limited thereto.

[0050] In one embodiment, ring Ar_2 may be a pyridine group, a pyrimidine group, a triazine group, a pyrazinyl group, a pyridazine group, an oxazine group, a thiazine group, an imidazole group, a pyrazole group, an oxazole group, an isoxazole group, a thiazole group, an isothiazole group, a triazole group, a furazanyl group, an oxadiazole group, a thiadiazole group, a diathiazole group, a tetrazole group, a quinoline group, an isoquinoline group, a quinazoline group, a quinoxaline group, a naphthyridine group, a diazaphthalene group, an acridine group, a phenanthridine group, a phenanthroline group, a triazindene group, a benzimidazole group, a benzoxazole group, or a benzothiazole group.

[0051] In one or more embodiments, ring Ar_2 may be a pyridine group, a pyrimidine group, or a triazine group, but embodiments of the present disclosure are not limited thereto.

[0052] In one embodiment, ring Ar_2 may be a C_2 - C_{60} heterocyclic group containing N at 2-position. For example, ring Ar_2 may be an N-containing C_2 - C_{60} heterocyclic group in which 2-position is N with respect to a moiety represented by



in Formula 2A, and $*$ indicates a binding site to ring Ar_2 .

[0053] X_3 in Formula 2A may be N or C(R_3).

[0054] X_4 in Formula 2A may be N or C(R_4).

[0055] In one embodiment, X_3 may be $C(R_3)$, and X_4 may be $C(R_4)$, but embodiments of the present disclosure are not limited thereto.

[0056] In one or more embodiments, X_3 may be N, and X_4 may be $C(R_4)$; or X_3 may be $C(R_3)$, and X_4 may be N.

[0057] In one or more embodiment, X_3 and X_4 may each be N at the same time.

[0058] X_5 in Formula 2B may be N, O, or S.

[0059] For example, X_5 may be O.

[0060] In one embodiment, X_5 may be N.

[0061] In one embodiment, X_5 may be S.

[0062] R_1 to R_7 in Formulae 2A and 2B may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), —B(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2), and

[0063] R_3 and R_4 may optionally be linked to form a substituted or unsubstituted C_5 - C_{30} carbocyclic group or a substituted or unsubstituted C_2 - C_{60} heterocyclic group.

[0064] In one embodiment, R_1 and R_2 in Formula 2A may each independently be selected from:

[0065] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0066] a C_1 - C_{60} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, and a C_1 - C_{60} heteroaryl group.

[0067] For example, R_1 and R_2 in Formula 2A may each independently be selected from:

[0068] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a

sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

[0069] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, and a C_1 - C_{60} heteroaryl group, but embodiments of the present disclosure are not limited thereto.

[0070] In one embodiment, R_1 and R_2 in Formula 2A may each independently be hydrogen, deuterium, a methyl group, or a phenyl group, but embodiments of the present disclosure are not limited thereto.

[0071] In one embodiment, R_3 and R_4 in Formula 2A may each independently be selected from:

[0072] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl

group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

[0073] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenaleny group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, and a C₁-C₆₀ heteroaryl group, and

[0074] R₃ and R₄ may optionally be linked to form a group selected from:

[0075] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group; and

[0076] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano

group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, and a C₁-C₆₀ heteroaryl group.

[0077] For example, R₃ and R₄ in Formula 2A may optionally be selected from hydrogen, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C₂-C₂₀ alkenyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a phenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, and

[0078] R₃ and R₄ may optionally be linked to form a group selected from:

[0079] a benzene group and a pyrazine group; and

[0080] a benzene group and a pyrazine group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, and a C₁-C₆₀ heteroaryl group.


[0081] In one embodiment, R₃ and R₄ in Formula 2A may each independently be hydrogen, deuterium, a methyl group, or a phenyl group, but embodiments of the present disclosure are not limited thereto.

[0082] In one or more embodiments, R₃ and R₄ in Formula 2A may be linked to form a benzene group or a pyrazine group, but embodiments of the present disclosure are not limited thereto.


[0083] a1 and a2 in Formula 2A may each independently be an integer from 0 to 10, wherein, when a1 is two or more, two or more R₁(s) may be identical to or different from each other, and when a2 is two or more, two or more R₂(s) may be identical to or different from each other.

[0084] * and *' in Formulae 2A and 2B each indicate a binding site to M in Formula 1.

[0085] In one embodiment, a moiety represented by



in Formula 2A may be a moiety represented by Formula 2A-1:



Formula 2A-1

[0086] R_1 and a_1 in Formula 2A-1 are the same as described in connection with R_1 and a_1 in Formula 2A.

[0087] In Formula 2A-1, Z_1 to Z_4 may each independently be C or N, and at least one of Z_1 to Z_4 may be N.

[0088] For example, Z_1 to Z_4 in Formula 2A-1 may each be C.

[0089] For example, in Formula 2A-1, Z_1 may be N, and Z_2 , Z_3 , and Z_4 may each be C.

[0090] In one or more embodiments, in Formula 2A-1, Z_2 may be N, and Z_1 , Z_3 , and Z_4 may each be C.

[0091] In one or more embodiments, in Formula 2A-1, Z_3 may be N, and Z_1 , Z_2 , and Z_4 may each be C.

[0092] In one or more embodiments, in Formula 2A-1, Z_4 may be N, and Z_1 , Z_2 , and Z_3 may each be C.

[0093] For example, in Formula 2A-1, Z_2 and Z_4 may each be N, and Z_1 and Z_3 may each be C.

[0094] In one or more embodiments, in Formula 2A-1, Z_1 and Z_4 may each be N, and Z_2 and Z_3 may each be C.

[0095] In one or more embodiments, in Formula 2A-1, Z_1 and Z_3 may each be N, and Z_2 and Z_4 may each be C.

[0096] In one or more embodiments, in Formula 2A-1, Z_3 and Z_4 may each be N, and Z_1 and Z_2 may each be C.

[0097] For example, in Formula 2A-1, Z_1 may be C, and Z_2 , Z_3 , and Z_4 may each be N.

[0098] In one or more embodiments, in Formula 2A-1, Z_2 may be C, and Z_1 , Z_3 , and Z_4 may each be N.

[0099] In one or more embodiments, in Formula 2A-1, Z_3 may be C, and Z_1 , Z_2 , and Z_4 may each be N.

[0100] In one or more embodiments, in Formula 2A-1, Z_4 may be C, and Z_1 , Z_2 , and Z_3 may each be N.

[0101] For example, Z_1 to Z_4 in Formula 2A-1 may each be N.

[0102] * in Formula 2A-1 indicates a binding site to a neighboring atom.


[0103] * in Formula 2A-1 indicates a binding site to M in Formula 1.

[0104] In one embodiment, a moiety represented by



Formula 2A-2

in Formula 2A may be a moiety represented by Formula 2A-2:



[0105] R_2 and a_2 in Formula 2A-2 are the same as described in connection with R_2 and a_2 in Formula 2A.

[0106] Z_5 to Z_8 in Formula 2A-2 may each independently be C or N.

[0107] For example, Z_5 to Z_8 in Formula 2A-2 may each be C at the same time.

[0108] For example, in Formula 2A-2, Z_5 may be N, and Z_6 , Z_7 , and Z_8 may each be C.

[0109] In one or more embodiments, in Formula 2A-2, Z_6 may be N, and Z_5 , Z_7 , and Z_8 may each be C.

[0110] In one or more embodiments, in Formula 2A-2, Z_7 may be N, and Z_5 , Z_6 , and Z_8 may each be C.

[0111] In one or more embodiments, in Formula 2A-2, Z_8 may be N, and Z_5 , Z_6 , and Z_7 may each be C.

[0112] For example, in Formula 2A-2, Z_6 and Z_8 may each be N, and Z_5 and Z_7 may each be C.

[0113] In one or more embodiments, in Formula 2A-2, Z_5 and Z_8 may each be N, and Z_6 and Z_7 may each be C.

[0114] In one or more embodiments, in Formula 2A-2, Z_5 and Z_7 may each be N, and Z_6 and Z_8 may each be C.

[0115] In one or more embodiments, in Formula 2A-2, Z_7 and Z_8 may each be N, and Z_5 and Z_6 may each be C.

[0116] For example, in Formula 2A-2, Z_5 may be C, and Z_6 , Z_7 , and Z_8 may each be N.

[0117] In one or more embodiments, in Formula 2A-2, Z_6 may be C, and Z_5 , Z_7 , and Z_8 may each be N.

[0118] In one or more embodiments, in Formula 2A-2, Z_7 may be C, and Z_5 , Z_6 , and Z_8 may each be N.

[0119] In one or more embodiments, in Formula 2A-2, Z_8 may be C, and Z_5 , Z_6 , and Z_7 may each be N.

[0120] For example, in Formula 2A-2, Z_5 to Z_8 may each be N at the same time.

[0121] Formula 2A-2, * indicates a binding site to a neighboring atom.

[0122] In one embodiment, R_5 and R_7 in Formula 2B may be identical to each other, but embodiments of the present disclosure are not limited thereto.

[0123] In one embodiment, R_5 and R_7 in Formula 2B may be different from each other, but embodiments of the present disclosure are not limited thereto.

[0124] In one embodiment, R_5 to R_7 in Formula 2B may each independently be selected from:

[0125] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

[0126] a C_1 - C_{60} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic con-

densed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, and a C₁-C₆₀ heteroaryl group.

[0127] In one or more embodiments, R₅ to R₇ in Formula 2B may each independently be selected from:

[0128] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₃-C₁₀ cycloalkyl group, and a C₆-C₆₀ aryl group; and

[0129] deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₃-C₁₀ cycloalkyl group, and a C₆-C₆₀ aryl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, and a C₁-C₆₀ heteroaryl group.

[0130] For example, R₅ to R₇ in Formula 2B may each independently be selected from:


[0131] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a triphenylenyl group, a fluoranthenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group; and

[0132] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a triphenylenyl group, a fluoranthenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group,

a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, and a C₁-C₆₀ heteroaryl group.

[0133] In one or more embodiments, R₅ to R₇ in Formula 2B may each independently be hydrogen, a methyl group, a phenyl group, or a phenyl group substituted with a methyl group, but embodiments of the present disclosure are not limited thereto.

[0134] In one embodiment, the organometallic compound represented by Formula 1 may be an organometallic compound represented by Formula 10:



Formula 10

[0135] M, X₃ to X₅, R₁ to R₇, a₁, and a₂ in Formula 10 may each independently be the same as described in connection with M, X₃ to X₅, R₁ to R₇, a₁, and a₂ in Formulae 1, 2A, and 2B.

[0136] In Formula 10, Z₁ to Z₈ may each independently be C or N, and at least one of Z₁ to Z₄ may be N.





[0137] Z₁ to Z₈ in Formula 10 may each independently be the same as described in connection with Z₁ to Z₈ in Formulae 2A-1 and 2A-2.

[0138] n in Formula 10 may be 1 or 2.





[0139] In one embodiment, in Formula 10, M may be platinum, and n may be 1.

[0140] In one or more embodiments, in Formula 10, M may be iridium, and n may be 2.

[0141] In one or more embodiments, the organometallic compound may be selected from Compounds 1 to 140, but embodiments of the present disclosure are not limited thereto:



-continued



5


6

7

8


-continued

-continued





9

13





10

14





11

15




12


16




-continued




17




21




18




22




19



23




20




24

-continued


-continued




25




29




26




30




27



31







28




32

-continued




-continued

33




37

34




38

35




39

36




40

-continued




41


-continued




45




42




46




43



47




44




48

-continued




49


-continued




53




50




54




51



55




52




56

-continued




57


-continued




61




58




62




59



63




60



64


-continued

-continued




65

69




66

70







67

71




68


72




-continued




73



74




75




76


-continued




77



78




79




80

-continued




81


-continued




85




82




86




83



87




84




88

-continued


-continued




89




93




90




94




91



95




92




96

-continued




97


-continued




101




98




102




99



103




100




104

-continued




105


-continued




109




106




110




107



111




108




112

-continued




113


-continued




117




114




118




115



119







116




120

-continued




-continued




133


-continued




137




134




138




135



139



136



140

[0142] The organometallic compound represented by Formula 1 includes at least one ligand represented by Formula 2A or 2B. The ligand represented by Formula 2A may include a carbene moiety substituted with an N-containing heterocyclic group, and the ligand represented by Formula 2B may include an —NH group.

[0143] Therefore, N in the ligand represented by Formula 2A, for example, N included in an =N— moiety forming a part of the N-containing heterocyclic group, and —NH in the ligand represented by Formula 2B are linked via an intermolecular hydrogen bond, thereby stabilizing a molecular structure and improving the efficiency and lifespan of a material in applying to an organic light-emitting device.

[0144] Also, due to the intermolecular hydrogen bond described above, it is possible to obtain an effect that increases selectivity to a structural isomer that may be generated by an asymmetric auxiliary ligand in synthesizing an organometallic compound.

[0145] The organometallic compound may emit blue light. For example, the organometallic compound may emit blue light (bottom-emission CIE_{x,y}, color coordinates 0.14, 0.06 to 0.08) having an upper (e.g., a maximum) emission wavelength of greater than or equal to about 440 nm and less than about 495 nm, for example, greater than or equal to about 450 nm and less than or equal to about 471 nm, but embodiments of the present disclosure are not limited thereto. Therefore, the organometallic compound represented by Formula 1 may be usefully used in manufacturing an organic light-emitting device that emits deep blue light.

[0146] A synthesis method for the organometallic compound represented by Formula 1 will become apparent to those of ordinary skill in the art by referring to the following examples.

[0147] At least one of the organometallic compound of Formula 1 may be used between a pair of electrodes of an organic light-emitting device. For example, the organometallic compound may be included in an emission layer. The organometallic compound may act as a dopant in the emission layer.

[0148] Accordingly, provided is an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes at least one organometallic cyclic compound represented by Formula 1.

[0149] As used herein, the expression “(an organic layer) includes at least one organometallic compound” used herein may include a case in which “(an organic layer) includes identical compounds represented by Formula 1” and a case in which “(an organic layer) includes two or more different organometallic compounds represented by Formula 1.”

[0150] For example, the organic layer may include, as the organometallic compound, only Compound 1. In this regard, Compound 1 may exist in an emission layer of the organic light-emitting device. In one or more embodiments, the organic layer may include, as the organometallic compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may exist in an identical layer (for example,

[0151] Compound 1 and Compound 2 may all exist in an emission layer), or different layers (for example, Compound 1 may exist in an emission layer and Compound 2 may exist in an electron transport region).

[0152] In an embodiment,

[0153] the first electrode of the organic light-emitting device may be an anode,

[0154] the second electrode of the organic light-emitting device may be a cathode,

[0155] the organic layer may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,

[0156] the hole transport region may include a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof, and

[0157] the electron transport region may include a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

[0158] The term “organic layer,” as used herein, refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0159] In one or more embodiments, the emission layer may include the organometallic compound represented by Formula 1, and the emission layer may further include a host, wherein an amount of the host in the emission layer may be greater than that of the organometallic compound in the emission layer.

[0160] For example, the host may include at least one of a silicon-containing compound and a phosphine oxide-containing compound.

[0161] In one embodiment, the hole transport region may include an electron blocking layer, and the electron blocking layer may include the organometallic compound; or

[0162] the electron transport region may include a hole blocking layer, and the hole blocking layer may include the organometallic compound.

[0163] In one embodiment, the hole transport region may include a p-dopant, wherein a low unoccupied molecular orbital (LUMO) energy level of the p-dopant is lower than about -3.5 eV.

[0164] In one or more embodiments, the electron transport region may include a hole blocking layer, and the hole blocking layer may include at least one of a silicon-containing compound and a phosphine oxide-containing compound.

DESCRIPTION OF THE ACCOMPANYING DRAWING

[0165] The accompanying drawing is a schematic view of an organic light-emitting device **10** according to an embodiment. The organic light-emitting device **10** includes a first electrode **110**, an organic layer **150**, and a second electrode **190**.

[0166] Hereinafter, the structure of the organic light-emitting device **10** according to an embodiment and a method of manufacturing the organic light-emitting device **10** will be described in connection with the accompanying drawing.

[0167] [First Electrode **110**]

[0168] In the accompanying drawing, a substrate may be additionally disposed under the first electrode **110** or above the second electrode **190**. The substrate may be a glass

substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0169] The first electrode **110** may be formed by depositing or sputtering a material for forming the first electrode **110** on the substrate. When the first electrode **110** is an anode, the material for a first electrode may be selected from materials with a high work function to facilitate hole injection.

[0170] The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode **110** is a transmissive electrode, a material for forming a first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combinations thereof, but embodiments of the present disclosure are not limited thereto. In one or more embodiments, when the first electrode **110** is a semi-transmissive electrode or a reflectable electrode, a material for forming a first electrode may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combinations thereof, but embodiments of the present disclosure are not limited thereto.

[0171] The first electrode **110** may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **110** is not limited thereto.

[0172] [Organic Layer **150**]

[0173] The organic layer **150** is disposed on the first electrode **110**. The organic layer **150** may include an emission layer.

[0174] The organic layer **150** may further include a hole transport region between the first electrode **110** and the emission layer, and an electron transport region between the emission layer and the second electrode **190**.

[0175] [Hole Transport Region in Organic Layer **150**]



[0176] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0177] The hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.


[0178] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein for each structure, constituting layers are sequentially stacked from the first

electrode **110** in this stated order, but the structure of the hole transport region is not limited thereto.









[0179] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANT/CSA), polyaniline/poly(4-styrenesulfonate) (PANT/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




-continued



-continued





[0180] In Formulae 201 and 202,

[0181] L_{201} to L_{204} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0182] L_{205} may be selected from $*-O-*$, $*-S-*$, $*-N(Q_{201})-*$, a substituted or unsubstituted C_1 - C_{20} alkylene group, a substituted or unsubstituted C_2 - C_{20} alkenylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0183] xa_1 to xa_4 may each independently be an integer from 0 to 3,

[0184] xa_5 may be an integer from 1 to 10,

[0185] R_{201} to R_{204} and Q_{201} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0186] In one embodiment, in Formula 202, R_{201} and R_{202} may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R_{203} and R_{204} may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

[0187] In one or more embodiments, in Formulae 201 and 202,

[0188] L_{201} to L_{205} may each independently be selected from:

[0189] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a

heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

[0190] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenyl group, a pentacenylenyl group, a rubicenylenyl group, a coronenylenyl group, an ovalenylenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a benzofuranyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, and $-N(Q_{31})(Q_{32})$, and

[0191] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0192] In one or more embodiments, xa_1 to xa_4 may each independently be 0, 1, or 2.

[0193] In one or more embodiments, xa5 may be 1, 2, 3, or 4.

[0194] In one or more embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from:

[0195] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0196] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂), and

[0197] Q₃₁ to Q₃₃ are the same as described above.

[0198] In one or more embodiments, at least one of R₂₀₁ to R₂₀₃ in Formula 201 may each independently be selected from:

[0199] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0200] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,

[0201] but embodiments of the present disclosure are not limited thereto.

[0202] In one or more embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be linked via a single bond, and/or ii) R₂₀₃ and R₂₀₄ may be linked via a single bond.

[0203] In one or more embodiments, at least one of R₂₀₁ to R₂₀₄ in Formula 202 may be selected from:


[0204] a carbazolyl group; and

[0205] a carbazolyl group, substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,


[0206] but embodiments of the present disclosure are not limited thereto.

[0207] The compound represented by Formula 201 may be represented by Formula 201A:


Formula 201A




[0208] In one embodiment, the compound represented by Formula 201 may be represented by Formula 201A(1) below, but embodiments of the present disclosure are not limited thereto:




[0209] In one embodiment, the compound represented by Formula 201 may be represented by Formula 201A-1 below, but embodiments of the present disclosure are not limited thereto:



[0210] In one embodiment, the compound represented by Formula 202 may be represented by Formula 202A:



[0211] In one embodiment, the compound represented by Formula 202 may be represented by Formula 202A-1:



[0212] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0213] L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} are the same as described above,

[0214] R_{211} and R_{212} may be understood by referring to the description provided herein in connection with R_{203} , and


[0215] R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0216] The hole transport region may include at least one compound selected from Compounds HT1 to HT39, but embodiments of the present disclosure are not limited thereto:


HT1




HT2



HT3




HT4




-continued


HT5




HT6



HT7




HT8




-continued


HT9




HT10



HT11





HT12



-continued



HT13

HT14





HT15

HT16




HT17

HT18




-continued





-continued

HT23





HT24

HT25





HT26

HT27





-continued
HT28

HT29





HT30

HT31





HT32


HT33





HT34

HT35




-continued
HT36

HT38

-continued
HT37

HT37

HT39



[0217] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, for example about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, suitable or satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0218] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[0219] [P-Dopant]

[0220] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0221] The charge-generation material may be, for example, a p-dopant.

[0222] In one embodiment, the LUMO of the p-dopant may be lower than -3.5 eV.

[0223] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments of the present disclosure are not limited thereto.


[0224] For example, the p-dopant may include at least one selected from:

[0225] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);


[0226] a metal oxide, such as tungsten oxide or molybdenum oxide;

[0227] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and


[0228] a compound represented by Formula 221 below:
 [0229] but embodiments of the present disclosure are not limited thereto:



HAT-CN



F4-TCNQ



Formula 221

[0230] In Formula 221,

[0231] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one selected from R_{221} to R_{223} may have at least one substituent selected from a cyano group, $-F$, $-Cl$, $-Br$, $-I$, a C_1 - C_{20} alkyl group substituted with $-F$, a C_1 - C_{20} alkyl group substituted with $-Cl$, a C_1 - C_{20} alkyl group substituted with $-Br$, and a C_1 - C_{20} alkyl group substituted with $-I$.

[0232] [Emission Layer in Organic Layer 150]

[0233] When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red light-emitting material, a green light-

emitting material, and a blue light-emitting material, in which the two or more materials are mixed with each other in a single layer to emit white light.


[0234] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0235] An amount of the dopant in the emission layer may be in a range of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.

[0236] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0237] [Host in Emission Layer]

[0238] In one or more embodiments, the host may include a compound represented by Formula 301 below.



Formula 301

[0239] In Formula 301,

[0240] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0241] $xb11$ may be 1, 2, or 3,

[0242] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0243] $xb1$ may be an integer from 0 to 5,

[0244] R_{301} may be selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{301})(Q_{302})(Q_{303})$, $-N(Q_{301})(Q_{302})$, $-B(Q_{301})(Q_{302})$, $-C(=O)(Q_{301})$, $-S(=O)_2(Q_{301})$, and $-P(=O)(Q_{301})(Q_{302})$,

[0245] $xb21$ may be an integer from 1 to 5,

[0246] Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0247] In one embodiment, Ar₃₀₁ in Formula 301 may be selected from:


[0248] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and

[0249] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and


[0250] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0251] When xb11 in Formula 301 is two or more, two or more Ar₃₀₁(s) may be linked via a single bond.

[0252] In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:



Formula 301-1



Formula 301-2

[0253] In Formulae 301-1 and 301-2,

[0254] A₃₀₁ to A₃₀₄ may each independently be selected from a benzene, a naphthalene, a phenanthrene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine, an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a dibenzofluorene, an indole, a carbazole, a benzocarbazole, a dibenzocarbazole, a furan, a benzofuran,

a dibenzofuran, a naphthofuran, a benzonaphthofuran, a dinaphthofuran, a thiophene, a benzothiophene, a dibenzothiophene, a naphthothiophene, a benzonaphthothiophene group, and a dinaphthothiophene group,

[0255] X₃₀₁ may be O, S, or N-[(L₃₀₄)_{xb4}-R₃₀₄],

[0256] R₃₁₁ to R₃₁₄ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0257] xb22 and xb23 may each independently be 0, 1, or 2,

[0258] L₃₀₁, xb1, R₃₀₁, and Q₃₁ to Q₃₃ may each independently be the same as described above,

[0259] L₃₀₂ to L₃₀₄ may each independently be the same as described in connection with L₃₀₁,

[0260] xb2 to xb4 may each independently be the same as described in connection with xb1, and

[0261] R₃₀₂ to R₃₀₄ may each independently be the same as described in connection with R₃₀₁.

[0262] For example, L₃₀₁ to L₃₀₄ in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0263] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a diben-


zofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a

group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$, and


[0269] Q_{31} to Q_{33} are the same as described above.

[0270] In one or more embodiments, the host may include an alkaline earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), a Mg complex, and a Zn complex.

[0271] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, but embodiments of the present disclosure are not limited thereto:




H1




H2


-continued




H3




H4




H5



H6




H7




H8

-continued


H9




H10




H11



H12




H13




-continued


H14




H15




H16







H17






H18





-continued



-continued




-continued




-continued

H28

H26




H29




H30


H27




H31




-continued




H32



H33




H34




H35


-continued




H36



H37




H38




H39

-continued




H40


-continued




H43




H41



H44




H42



H45


-continued

H46




-continued

H52




H53


H47




H48





H49




H50




H51



H54



H55



[0272] In one embodiment, the host may include at least one selected from a silicon-containing compound (for example, BCPDS used in the following examples or the like) and a phosphine oxide-containing compound (for example, POPCPA used in the following examples or the like).

[0273] However, embodiments of the present disclosure are not limited thereto. In one embodiment, the host may include only one compound, or two or more different

compounds (for example, a host used in the following examples includes BCPDS and POPCPA).

[0274] [Phosphorescent Dopant Included in Emission Layer in Organic Layer 150]

[0275] The phosphorescent dopant may include an organometallic compound represented by Formula 1 below:

[0276] [Electron Transport Region in Organic Layer 150]

[0277] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0278] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

[0279] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein for each structure, constituting layers are sequentially stacked from an emission layer. However, embodiments of the structure of the electron transport region are not limited thereto.


[0280] The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

[0281] As used herein, the term “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0282] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered heteromonocyclic group having at least one $*-N=*$ moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one $*-N=*$ moiety are condensed with each other (e.g., combined together), or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one $*-N=*$ moiety, is condensed with (e.g., combined together with) at least one C_5 - C_{60} carbocyclic group.

[0283] Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole, but are not limited thereto.

[0284] For example, the electron transport region may include a compound represented by Formula 601:



Formula 601

[0285] In Formula 601,

[0286] Ar_{601} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0287] $xe11$ may be 1, 2, or 3,

[0288] L_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0289] $xe1$ may be an integer from 0 to 5,

[0290] R_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{601})(Q_{602})(Q_{603})$, $-C(=O)(Q_{601})$, $-S(=O)_2(Q_{601})$, and $-P(=O)(Q_{601})(Q_{602})$,

[0291] Q_{601} to Q_{603} may each independently be a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0292] $xe21$ may be an integer from 1 to 5.

[0293] In one embodiment, at least one of $Ar_{601}(s)$ in the number of $xe11$ and $R_{601}(s)$ in the number of $xe21$ may include the π electron-depleted nitrogen-containing ring described above.

[0294] In one embodiment, ring Ar_{601} in Formula 601 may be selected from:

[0295] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0296] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triph-

enylene group, a pyrene group, a chrysene group, a naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and


[0297] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0298] When xe11 in Formula 601 is two or more, two or more Ar_{60i}(s) may be linked via a single bond.

[0299] In one or more embodiments, Ar_{60i} in Formula 601 may be an anthracene group.

[0300] In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:

Formula 601-1



[0301] In Formula 601-1,

[0302] X₆₁₄ may be N or C(R₆₁₄), X₆₁₅ may be N or C(R₆₁₅), X₆₁₆ may be N or C(R₆₁₆), and at least one of X₆₁₄ to X₆₁₆ may be N,

[0303] L₆₁₁ to L₆₁₃ may each independently be the same as described in connection with L₆₀₁,

[0304] xe611 to xe613 may each independently be the same as described in connection with xe1,

[0305] R₆₁₁ to R₆₁₃ may each independently be the same as described in connection with R₆₀₁, and

[0306] R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0307] In one embodiment, L₆₀₁ and L₆₁₁ to L₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0308] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylene group, a dibenzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0309] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylene group, a dibenzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a

spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group,

[0310] but embodiments of the present disclosure are not limited thereto.

[0311] In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0312] In one or more embodiments, R₆₀₁ and R₆₁₁ to R₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0313] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

[0314] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl

group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


[0315] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂), and

[0316] Q₆₀₁ and Q₆₀₂ are the same as described above.


[0317] The electron transport region may include at least one compound selected from Compounds ET1 to ET36, but embodiments of the present disclosure are not limited thereto:

-continued


ET1




ET4




ET2




ET5



ET3




ET6




-continued


ET7



ET8




ET9




-continued


ET10






ET11






ET12






-continued



-continued




-continued




-continued

ET19




ET22

ET20




ET23

ET21




ET24


-continued




-continued




ET26




ET29




ET27



ET30




-continued




ET31

-continued




ET34


ET32




ET35




ET33



ET36



[0318] In one or more embodiments, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BALq, 3-(bi-phenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:



[0319] In one embodiment, the electron transport region may include a phosphine oxide-containing compound (for example, TSPO1 used in the following examples or the like), but embodiments of the present disclosure are not limited thereto. In one embodiment, the phosphine oxide-containing compound may be used in a hole blocking layer in the electron transport region, but embodiments of the present disclosure are not limited thereto.

[0320] A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.


[0321] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example,

about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have suitable or satisfactory electron transport characteristics without a substantial increase in driving voltage.


[0322] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0323] The metal-containing material may include at least one selected from alkali metal complex and alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from a Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, a Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenyloxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazaole, a hydroxy diphenylthiadiazole, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0324] For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2



ET-D1



ET-D2

[0325] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0326] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0327] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare

earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof.

[0328] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs, but embodiments of the present disclosure are not limited thereto.

[0329] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0330] The rare earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

[0331] The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare earth metal.

[0332] The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, or KI. In one embodiment, the alkali metal compound may be selected from LiF, Li_2O , NaF, LiI, NaI, CsI, and KI, but embodiments of the present disclosure are not limited thereto.

[0333] The alkaline earth-metal compound may be selected from alkaline earth-metal oxides, such as BaO, SrO, CaO , $\text{Ba}_x\text{Sr}_{1-x}\text{O}$ ($0 < x < 1$), or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$). In one embodiment, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO, but embodiments of the present disclosure are not limited thereto.

[0334] The rare earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In one embodiment, the rare earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 , but embodiments of the present disclosure are not limited thereto.

[0335] The alkali metal complex, the alkaline earth-metal complex, and the rare earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, or the rare earth metal complex may be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenyloxazole, hydroxy phenylthiazole, hydroxy diphenyloxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0336] The electron injection layer may consist of an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any

combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0337] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have suitable or satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0338] [Second Electrode 190]

[0339] The second electrode 190 may be disposed on the organic layer 150 having such a structure. The second electrode 190 may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode 190 may be selected from metal, an alloy, an electrically conductive compound, and a combination thereof, which have a relatively low work function.

[0340] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, but embodiments of the present disclosure are not limited thereto. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0341] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[0342] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0343] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100° C. to about 500° C., at a vacuum degree of about 10^{-8} torr to about 10^{-3} torr, and at a deposition rate of about 0.01 Å/sec to about 100 Å/sec by taking into account a material to be included in a layer to be formed, and the structure of a layer to be formed.

[0344] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to about 200° C. by taking into account a material to be included in a layer to be formed, and the structure of a layer to be formed.

General Definitions of Some of the Substituents

[0345] The term “ $\text{C}_1\text{-C}_{60}$ alkyl group,” as used herein, refers to a linear or branched aliphatic saturated hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “ $\text{C}_1\text{-C}_{60}$ alkylene group,” as used herein, refers to a divalent group having substantially the same

structure as the C₁-C₆₀ alkyl group, except that the C₁-C₆₀ alkylene group is divalent instead of monovalent.

[0346] The term “C₂-C₆₀ alkenyl group,” as used herein, refers to a hydrocarbon group having at least one carbon-carbon double bond at a main chain (e.g., in the middle) or at the terminus (e.g., an end) of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group,” as used herein refers, to a divalent group having substantially the same structure as the C₂-C₆₀ alkenyl group, except that the C₂-C₆₀ alkenylene group is divalent instead of monovalent.

[0347] The term “C₂-C₆₀ alkynyl group,” as used herein, refers to a hydrocarbon group having at least one carbon-carbon triple bond at a main chain (e.g., in the middle) or at the terminus (e.g., an end) of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group, and a propynyl group. The term “C₂-C₆₀ alkynylene group,” as used herein, refers to a divalent group having substantially the same structure as the C₂-C₆₀ alkynyl group, except that the C₂-C₆₀ alkynylene group is divalent instead of monovalent.

[0348] The term “C₁-C₆₀ alkoxy group,” as used herein, refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0349] The term “C₃-C₁₀ cycloalkyl group,” as used herein, refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group,” as used herein refers to a divalent group having substantially the same structure as the C₃-C₁₀ cycloalkyl group, except that the C₃-C₁₀ cycloalkylene group is divalent instead of monovalent.

[0350] The term “C₁-C₁₀ heterocycloalkyl group,” as used herein, refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidiny group, a tetrahydrofuran group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group,” as used herein refers to a divalent group having substantially the same structure as the C₁-C₁₀ heterocycloalkyl group, except that the C₁-C₁₀ heterocycloalkylene group is divalent instead of monovalent.

[0351] The term “C₃-C₁₀ cycloalkenyl group,” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity (e.g., the ring is not aromatic), and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group,” as used herein, refers to a divalent group having substantially the same structure as the C₃-C₁₀ cycloalkenyl group, except that the C₃-C₁₀ cycloalkenylene group is divalent instead of monovalent.

[0352] The term “C₁-C₁₀ heterocycloalkenyl group,” as used herein, refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group include a

4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuran group, and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group,” as used herein, refers to a divalent group having substantially the same structure as the C₁-C₁₀ heterocycloalkenyl group, except that the C₁-C₁₀ heterocycloalkenylene group is divalent instead of monovalent.

[0353] The term “C₆-C₆₀ aryl group,” as used herein, refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other (e.g., combined together).

[0354] The term “C₁-C₆₀ heteroaryl group,” as used herein, refers to a monovalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group,” as used herein, refers to a divalent group having a carbocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be condensed with each other (e.g., combined together).

[0355] The term “C₆-C₆₀ aryloxy group,” as used herein, refers to —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and the term “C₆-C₆₀ arylthio group,” as used herein, indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0356] The term “monovalent non-aromatic condensed polycyclic group,” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) having two or more rings condensed with each other (e.g., combined together), only carbon atoms as ring-forming atoms, and no aromaticity in its entire molecular structure (e.g., the entire group is not aromatic). An example of the monovalent non-aromatic condensed polycyclic group includes a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” as used herein, refers to a divalent group having substantially the same structure as the monovalent non-aromatic condensed polycyclic group, except that the divalent non-aromatic condensed polycyclic group is divalent instead of monovalent.

[0357] The term “monovalent non-aromatic condensed heteropolycyclic group,” as used herein, refers to a monovalent group (for example, having 1 to 60 carbon atoms) having two or more rings condensed to each other (e.g., combined together), at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and no aromaticity in its entire molecular structure (e.g., the entire group is not aromatic). An example of the monovalent non-aromatic condensed heteropolycyclic group includes a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group,” as used herein, refers to a divalent group having substantially the

same structure as the monovalent non-aromatic condensed heteropolycyclic group, except that the divalent non-aromatic condensed heteropolycyclic group is divalent instead of monovalent.

[0358] The term “C₄-C₆₀ carbocyclic group,” as used herein, refers to a monocyclic or polycyclic group having 4 to 60 carbon atoms in which a ring-forming atom is a carbon atom only (e.g., the ring is formed only from carbon atoms). The C₄-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₄-C₆₀ carbocyclic group may be a ring, such as benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C₄-C₆₀ carbocyclic group, the C₄-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0359] The term “C₂-C₆₀ heterocyclic group,” as used herein, refers to a group having substantially the same structure as the C₄-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 2 to 60).

[0360] At least one substituent of the substituted C₄-C₆₀ carbocyclic group, the substituted C₂-C₆₀ heterocyclic group, the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0361] deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0362] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0363] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀

heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0364] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

[0365] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0366] Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ alkyl group substituted with at least one selected from deuterium, —F, and a cyano group, a C₆-C₆₀ aryl group substituted with at least one selected from deuterium, —F, and a cyano group, a biphenyl group, and a terphenyl group.

[0367] The term “Ph,” as used herein, represents a phenyl group, the term “Me,” as used herein, represents a methyl group, the term “Et,” as used herein, represents an ethyl group, the term “ter-Bu” or “Bu^t,” as used herein, represents a tert-butyl group, and the term “OMe,” as used herein, represents a methoxy group.

[0368] The term “biphenyl group,” as used herein, refers to a “phenyl group substituted with a phenyl group. The “biphenyl group” is a “substituted phenyl group” having a “C₆-C₆₀ aryl group” as a substituent.

[0369] The term “terphenyl group,” as used herein, refers to a “phenyl group substituted with a biphenyl group. The “terphenyl group” is a “phenyl group” having, as a substituent, a “C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group.”

[0370] * and *¹ used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.


[0371] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in more detail with reference

to Synthesis Examples and Examples. The expression "B was used instead of A" used in describing Synthesis Examples means that an identical (or substantially identical) number of molar equivalents of A was used in place of molar equivalents of B.

EXAMPLES

Synthesis Example 1: Synthesis of Compound 1

[0372] 1) Synthesis of Intermediate [A-1]




[0373] 1.0 equivalents (eq) of iodobenzene, 1.2 eq of imidazole, 0.01 eq of copper iodide, 2.0 eq of potassium carbonate, and 0.02 eq of L-Proline were dissolved in 0.1 M dimethylsulfonate, and then stirred at a temperature of 130° C. for 24 hours. The reaction mixture was cooled to room temperature, and an organic layer was extracted therefrom three times by using dichloromethane and water. The extracted organic layer was dried by using magnesium sulfate, concentrated, and then purified by column chromatography. The product obtained therefrom was dissolved in 1.0 M tetrahydrofuran (THF) together with 1.2 eq of 2-iodopyridine, and then stirred under reflux at a temperature of 50° C. for 3 hours. The reaction product was cooled to room temperature, and a precipitate obtained therefrom was cooled to room temperature, filtered and separated under reduced pressure, and washed by using diethyl ether to synthesize solid Intermediate [A-1] (yield: 60%).

[0374] 2) Synthesis of Intermediate [B-1]



[0375] 1.0 eq of acetylacetone and 1.05 eq of ammonium carbamate were dissolved in dichloromethane, and then stirred at room temperature for 12 hours. The reaction product was concentrated under reduced pressure to synthesize Intermediate [B-1] (yield: 99%).


[0376] 3) Synthesis of Compound 1



[0377] Intermediate [A-1] and 1.2 eq of silver(I) oxide were dissolved in 1.0 M dichloromethane, and stirred at room temperature for 6 hours. A solid generated after the reaction was filtered, and a solvent was removed from a filtrate to obtain silver-carbene Intermediate. The silver-carbene Intermediate and 1.1 eq of cyclooctadienylplatinum chloride were heated and stirred in a 2-ethoxyethanol solvent at a temperature of 120° C. for 24 hours, and 1.1 eq of Intermediate [B-1] and 2.0 eq of potassium tert-butoxide were added thereto and heated and stirred at a temperature of 120° C. for 12 hours. The reaction mixture was cooled to room temperature, and an organic layer was extracted therefrom three times by using dichloromethane and water. The extracted organic layer was dried by using magnesium sulfate, concentrated, and then purified by column chromatography. The product obtained therefrom was recrystallized by using diethyl ether to synthesize Compound 1 (yield: 23%).


Synthesis Example 2: Synthesis of Compound 5

[0378]




[0379] Compound 5 (yield: 19%) was synthesized in substantially the same manner as in Synthesis Example 1, except that 1,3-diphenyl-1,3-propanedione was used instead of acetyl acetone in synthesizing Intermediate [B-2].

Synthesis Example 3: Synthesis of Compound 41
[0380]



[0381] Compound 41 (yield: 21%) was synthesized in substantially the same manner as in Synthesis Example 1, except that 3-(2,3,5,6-tetramethylphenyl)pentane-2,4-dione was used instead of acetyl acetone in synthesizing Intermediate [B-3].


Synthesis Example 4: Synthesis of Compound 45
[0382]



[0383] Compound 45 (yield: 18%) was synthesized in substantially the same manner as in Synthesis Example 1, except that benzimidazole was used instead of imidazole in synthesizing Intermediate [A-4].

Synthesis Example 5: Synthesis of Compound 69

[0384]



[0385] Compound 69 (yield: 22%) was synthesized in substantially the same manner as in Synthesis Example 1, except that 1H-imidazo[4,5-b]pyrazine was used instead of imidazole in synthesizing Intermediate [A-5].

[0386] ¹H NMR and FAB-MS of Compounds synthesized according to Synthesis Examples 1 to 5 are shown in Table 1.

[0387] Methods of synthesizing compounds other than the compounds shown in Table 1 are recognizable by those of ordinary skill in the art by referring to the synthesis mechanisms and source materials described above.

TABLE 1

Compound	¹ H NMR (CDCl ₃ , 500 MHz)	FAB-MS [M ⁺]	
		found	calc.
1	9.54 (d, 1H), 7.66 (t, 1H), 7.23 (m, 2H), 7.11 (t, 1H), 7.01 (d, 1H), 6.77 (t, 1H), 6.67 (d, 1H), 5.43 (m, 2H), 4.98 (s, 1H), 2.26 (s, 3H), 1.93 (s, 3H), 1.62 (s, 1H) ppm.	513.11	513.11
5	9.73 (d, 1H), 7.80-7.70 (m, 4H), 7.58-7.48 (m, 6H), 7.25-7.20 (m, 2H), 6.78 (t, 1H), 6.65 (d, 1H), 6.35 (s, 1H), 5.42 (m, 2H), 3.80 (s, 1H), 1.54 (s, 3H) ppm.	638.15	638.15
41	9.59 (d, 1H), 7.86-7.82 (m, 3H), 7.24-7.21 (m, 2H), 6.81-6.66 (m, 3H), 5.43 (m, 2H), 3.86 (s, 1H), 2.26-2.07 (m, 6H), 1.46 (s, 1H) ppm.	644.22	644.21
45	9.68 (d, 1H), 7.83-7.80 (m, 3H), 7.30-7.27 (m, 2H), 7.20-7.13 (m, 3H), 6.94-7.02 (m, 3H), 4.98 (s, 1H), 3.84 (s, 1H), 2.23 (s, 3H), 1.97 (s, 3H), 1.41 (s, 1H) ppm.	564.14	564.14
69	9.71 (d, 1H), 8.41-8.34 (m, 3H), 7.83-7.80 (m, 3H), 7.30-7.05 (m, 3H), 4.94 (s, 1H), 3.86 (s, 1H), 2.28 (s, 3H), 1.95 (s, 3H), 1.71 (s, 1H) ppm.	566.13	566.13


Example 1

[0388] As a substrate and an anode, a Corning 15 Ω/cm^2 (1,200 Å) ITO glass substrate was cut to a size of 50 mm \times 50 mm \times 0.7 mm, sonicated with isopropyl alcohol and pure water each for 5 minutes, and then cleaned by exposure to ultraviolet rays and ozone for 30 minutes. Then, the resultant ITO glass substrate was provided to a vacuum deposition apparatus.





[0389] Compound 2-TNATA was vacuum-deposited on the ITO glass substrate to form a hole injection layer having a thickness of 600 Å, and 4,4'-bis[N-(1-naphthyl)-N-phenylaminobiphenyl (NPB) as a hole transport compound was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.

[0390] bis(4-(9H-carbazol-9-yl)phenyl)diphenylsilane (BCPDS) and (4-(1-(4-(diphenylamino)phenyl)cyclohexyl)phenyl)diphenyl-phosphine oxide (POPCPA) as a co-host (weight ratio=1:1) and Compound 1 as a dopant were co-deposited on the hole transport layer, so that a weight ratio of the co-host to the dopant was 90:10, to form a blue fluorescent emission layer having a thickness of 300 Å.

[0391] Then, diphenyl(4-(triphenylsilyl)phenyl)-phosphine oxide (TSPO1) was vacuum-deposited on the blue fluorescent emission layer to form a hole blocking layer having a thickness of 50 Å. Then, Alq₃ was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, LiF, which is an alkali metal halide, was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form a cathode electrode having a thickness of 3,000 Å, thereby forming a LiF/Al electrode. In this manner, an organic light-emitting device was manufactured.



-continued



Examples 2 to 5 and Comparative Examples 1 to 4


[0392] Organic light-emitting devices were manufactured in substantially the same manner as in Example 1, except that Compounds shown in Table 1 were each used instead of Compound 1 as a dopant in forming an emission layer.

Evaluation Example 1


[0393] The driving voltage, current density, luminance, luminescent efficiency, and upper (e.g., maximum) emission wavelength of the organic light-emitting devices manufactured according to Examples 1 to 5 and Comparative Examples 1 to 4 were measured by using Keithley SMU 236 and a luminance meter PR650, and results thereof are shown in Table 2.

TABLE 2


	Dopant Compound	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Luminescent efficiency (cd/A)	Emission color	Maximum emission wavelength (nm)
Example 1	1	5.32	50	4125	8.25	Blue	470
Example 2	5	5.42	50	4025	8.05	Blue	462
Example 3	41	5.48	50	4033	8.07	Blue	463
Example 4	45	5.46	50	4024	8.05	Blue	458
Example 5	69	5.42	50	4055	8.11	Blue	460
Comparative Example 1	Flrpic	6.56	50	3870	7.74	Blue	475
Comparative Example 2	C	6.27	50	4021	8.04	Blue	464
Comparative Example 3	A	6.44	50	3805	7.61	Blue	462
Comparative Example 4	B	5.83	50	3720	7.44	Blue	473




Flrpic



Compound C



Compound A



Compound B

[0394] Referring to Table 1, it is confirmed that the organic light-emitting devices of Examples 1 to 5 have a low driving voltage, high luminance, and high luminescent efficiency and are suitable for deep blue light emission, as compared with those of the organic light-emitting devices of Comparative Examples 1 to 4.

[0395] An organic light-emitting device including the organometallic compound may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

[0396] It should be understood that embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0397] It will be understood that, although the terms “first,” “second,” “third,” etc., may be used herein to

describe various elements, components, regions, layers and/or sections, these elements, components, regions, layers and/or sections should not be limited by these terms. These terms are used to distinguish one element, component, region, layer or section from another element, component, region, layer or section. Thus, a first element, component, region, layer or section described below could be termed a second element, component, region, layer or section, without departing from the spirit and scope of the present disclosure.

[0398] Spatially relative terms, such as “beneath,” “below,” “lower,” “under,” “above,” “upper,” and the like, may be used herein for ease of explanation to describe one element or feature’s relationship to another element(s) or feature(s) as illustrated in the figures. It will be understood that the spatially relative terms are intended to encompass different orientations of the device in use or in operation, in

stituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

R₃ and R₄ are optionally linked to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₂-C₆₀ heterocyclic group,

a1 and a2 are each independently an integer from 0 to 10, wherein, when a1 is two or more, two or more R₁(s) are identical to or different from each other, and when a2 is two or more, two or more R₂(s) are identical to or different from each other,

and **i* each indicate a binding site to M in Formula 1,

at least one substituent of the substituted C₄-C₆₀ carbocyclic group, the substituted C₂-C₆₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic con-

densed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ alkyl group substituted with at least one selected from deuterium, —F, and a cyano group, a C₆-C₆₀ aryl group substituted with at least one selected from deuterium, —F, and a cyano group, a biphenyl group, and a terphenyl group.

2. The organometallic compound of claim 1, wherein:

M is Pt or Ir.

3. The organometallic compound of claim 1, wherein:

ring Ar₁ is a benzene group, a naphthalene group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, or a quinazoline group.

4. The organometallic compound of claim 1, wherein:

ring Ar₂ is a π electron-depleted nitrogen-containing C₂-C₆₀ heterocyclic group.

5. The organometallic compound of claim 1, wherein:

ring Ar₂ is a pyridine group, a pyrimidine group, a triazine group, a pyrazinyl group, a pyridazine group, an oxazine group, a thiazine group, an imidazole group, a pyrazole group, an oxazole group, an isoxazole group, a thiazole group, an isothiazole group, a triazole group, a furazanyl group, an oxadiazole group, a thiadiazole group, a diathiazole group, a tetrazole group, a quinoline group, an isoquinoline group, a quinazoline group, a quinoxaline group, a naphthyridine group, a diaz-anaphthalene group, an acridine group, a phenanthridine group, a phenanthroline group, a triazindene group, a benzimidazole group, a benzoxazole group, or a benzothiazole group.

6. The organometallic compound of claim 1, wherein:

X₃ is C(R₃), and X₄ is C(R₄).

7. The organometallic compound of claim 1, wherein:

R_1 and R_2 are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group; and

a C_1 - C_{60} alkyl group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, and a C_1 - C_{60} heteroaryl group.

8. The organometallic compound of claim 1, wherein:

R_3 and R_4 are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group; and

a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a pyrenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyridinyl group, a pyrimidyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group,


a phenazinyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a biphenyl group, and a terphenyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, and a C_1 - C_{60} heteroaryl group, and

R_3 and R_4 are optionally linked to form a group selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group; and


a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, and a C_1 - C_{60} heteroaryl group.

9. The organometallic compound of claim 1, wherein:

a moiety represented by



in Formula 2A is a moiety represented by Formula 2A-1:



Formula 2A-1

wherein, in Formula 2A-1,

R_1 and a_1 are the same as described in Formula 2A,

Z_1 to Z_4 are each independently C or N,

at least one of Z_1 to Z_4 is N,


* indicates a binding site to a neighboring atom, and

* \dagger indicates a binding site to M in Formula 1.

10. The organometallic compound of claim 1, wherein: a moiety represented by



in Formula 2A is a moiety represented by Formula 2A-2:



Formula 2A-2

wherein, in Formula 2A-2,

R_2 and a_2 are the same as described in Formula 2A,

Z_5 to Z_8 are each independently C or N, and

* indicates a binding site to a neighboring atom.

11. The organometallic compound of claim 1, wherein:

R_5 to R_7 are each independently selected from:


hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a triphenylenyl group, a fluoranthenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl

group, a picenyl group, a perylenyl group, a pentaphe-nyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group; and

- a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a triphenylenyl group, a fluoranthenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphe-nyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group, each substituted with at least one substituent selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, and a C_1 - C_{60} heteroaryl group.

- 12.** The organometallic compound of claim 1, wherein: the organometallic compound represented by Formula 1 is represented by Formula 10:

Formula 10



and

wherein, in Formula 10,





M, X_3 to X_5 , R_1 to R_7 , a_1 , and a_2 are the same as described in claim 1,

Z_1 to Z_8 are each independently C or N,





at least one of Z_1 to Z_4 is N, and

n is 1 or 2.

13. The organometallic compound of claim 1, wherein:
the organometallic compound represented by Formula 1 is
selected from Compounds 1 to 140:



-continued



5


6

7

8


-continued

-continued




9

13




10

14







11

15




12

16




-continued

-continued





17

21





18

22





19

23




20

24




-continued

-continued




25

29




26

30






27

31




28


32




-continued




33




37




34




38




35



39




36



40



-continued

-continued





41

45





42

46




43

47




44

48




-continued




49


-continued




53




50




54




51



55




52




56

-continued




57


-continued




61




58




62




59



63




60




64

-continued




65


-continued




69




66




70




67



71




68




72

-continued




73


-continued




77




74




78




75



79







76



80


-continued



-continued


85

81




86

82




87

83




88

84




-continued




89


-continued




93




90




94




91



95




92




96

-continued




97


-continued




101




98




102




99



103




100




104


-continued




105




109




106




110




107



111




108










112


-continued




-continued




-continued




-continued




122




130




123




131




124










132



-continued



-continued



- 14.** An organic light-emitting device comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer that is disposed between the first electrode and the second electrode and comprises an emission layer,
 wherein the organic layer comprises at least one of the organometallic compound of claim 1.

15. The organic light-emitting device of claim **14**, wherein:

the first electrode is an anode,

the second electrode is a cathode,

the organic layer further comprises a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,

the hole transport region comprises a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof, and

the electron transport region comprises a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

16. The organic light-emitting device of claim **14**, wherein:

the emission layer comprises the organometallic compound.

17. The organic light-emitting device of claim **16**, wherein:

the emission layer further comprises a host, and an amount of the host in the emission layer is larger than an amount of the organometallic compound in the emission layer.

18. The organic light-emitting device of claim **17**, wherein:

the host comprises at least one of a silicon-containing compound and a phosphine oxide-containing compound.

19. The organic light-emitting device of claim **15**, wherein:

the hole transport region comprises an electron blocking layer, and the electron blocking layer comprises the organometallic compound; or

the electron transport region comprises a hole blocking layer, and the hole blocking layer comprises the organometallic compound.

20. The organic light-emitting device of claim **15**, wherein:

the electron transport region comprises a hole blocking layer, and the hole blocking layer comprises a silicon-containing compound and a phosphine oxide-containing compound.

* * * * *

专利名称(译)	有机化合物和有机发光装置，包括相同的装置		
公开(公告)号	US20190036044A1	公开(公告)日	2019-01-31
申请号	US15/952587	申请日	2018-04-13
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	JEON MINA KO SOOBYUNG KIM SUNGBUM AHN HEECHOON JUN MIEUN KIM YOUNGKOOK HWANG SEOKHWAN		
发明人	JEON, MINA KO, SOOBYUNG KIM, SUNGBUM AHN, HEECHOON JUN, MIEUN KIM, YOUNGKOOK HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C07F15/00 C09K11/06		
CPC分类号	H01L51/0087 C07F15/0086 C09K11/06 H01L51/5012 H01L51/5096 H01L51/5016 H01L51/001 H01L2251/558 H01L51/5088 H01L51/506 H01L51/5072 H01L51/5092 H01L51/56 H01L51/5206 H01L51/5221 C09K2211/185 C09K2211/1044 C09K2211/1074 H01L51/0081 H01L51/0094 H01L2251 /308		
优先权	1020170096378 2017-07-28 KR		
外部链接	Espacenet USPTO		

摘要(译)

有机金属化合物由式1表示， $M(L_1)_{n1}(L_2)_{n2}$ ，其中M是Ir，Pt，Os，Ru，Pd，Rh，Fe，Co，Ni，Au，Ag或Cu，L₁是配体，L₂是配体，n₁为1或2，n₂为1或2.有机发光装置包括：第一电极;面向第一电极的第二电极;第一电极和第二电极之间包括发光层的有机层，其中有机层包括至少一种有机金属化合物。

10

190

150

110