



(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2019/0267553 A1**  
YE et al. (43) **Pub. Date: Aug. 29, 2019**

(54) **HETEROCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

51/0068 (2013.01); C09K 11/06 (2013.01); H01L 51/0069 (2013.01)

(71) Applicant: **Samsung Display Co., Ltd.**, Yongin-si (KR)

(72) Inventors: **Jimyoung YE**, Yongin-si (KR); **Myeongsuk KIM**, Yongin-si (KR); **Byeongwook YOO**, Yongin-si (KR); **Soobyung KO**, Yongin-si (KR); **Jihwan YOON**, Yongin-si (KR); **Jaehoon HWANG**, Yongin-si (KR)

(57) **ABSTRACT**

An organic light-emitting device and a heterocyclic compound, the device including a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and including an emission layer, wherein the emission layer includes a compound represented by Formula 1A or a compound represented by Formula 1B:

(21) Appl. No.: **16/181,407**

(22) Filed: **Nov. 6, 2018**

(30) **Foreign Application Priority Data**


Feb. 28, 2018 (KR) ..... 10-2018-0024720

**Publication Classification**


(51) **Int. Cl.**  
**H01L 51/00** (2006.01)  
**C07D 417/14** (2006.01)  
**C09K 11/06** (2006.01)

(52) **U.S. Cl.**  
CPC ..... **H01L 51/0072** (2013.01); **C07D 417/14** (2013.01); **H01L 51/5016** (2013.01); **H01L**

<Formula 1A>



<Formula 1B>



**10**

<b>190</b>
<b>150</b>
<b>110</b>

**FIG. 1**

**10**

<b>190</b>
<b>150</b>
<b>110</b>

**FIG. 2**

**20**

<b>190</b>
<b>150</b>
<b>110</b>
<b>210</b>

# FIG. 3

30

220
190
150
110

# FIG. 4

40

<b>220</b>
<b>190</b>
<b>150</b>
<b>110</b>
<b>210</b>

**HETEROCYCLIC COMPOUND AND  
ORGANIC LIGHT-EMITTING DEVICE  
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED  
APPLICATION

**[0001]** Korean Patent Application No. 10-2018-0024720, filed on Feb. 28, 2018, in the Korean Intellectual Property Office, and entitled: "Heterocyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

**[0002]** Embodiments relate to a heterocyclic compound and an organic light-emitting device including the same.



2. Description of the Related Art

**[0003]** Organic light-emitting devices (OLEDs) are self-emission devices that, as compared with other devices, have wide viewing angles, high contrast ratios, short response times, and excellent characteristics in terms of brightness, driving voltage, and response speed, and produce full-color images.

**[0004]** The organic light-emitting device may include a first electrode on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. Then, the excitons are transitioned from an excited state to a ground state, thereby generating light.

SUMMARY


**[0005]** The embodiments may be realized by providing an organic light-emitting device including a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and including an emission layer, wherein the emission layer includes a compound represented by Formula 1A or a compound represented by Formula 1B:



**[0006]** wherein, in Formulae 1A and 1B, ring  $A_1$  is selected from a  $C_5$ - $C_{60}$  carbocyclic group and a  $C_2$ - $C_{60}$  heterocyclic group,  $X_1$  is S or Se,  $L_1$  and  $L_2$  are each independently selected from a substituted or unsubstituted  $C_5$ - $C_{60}$  carbocyclic group and a substituted or unsubstituted  $C_2$ - $C_{60}$  heterocyclic group,  $a_1$  and  $a_2$  are each independently an integer from 0 to 5, when  $a_1$  is 2, 3, 4, or 5, the 2, 3, 4, or 5  $L_1$ (s) are identical to or different from each other, and when  $a_2$  is 2, 3, 4, or 5, the 2, 3, 4, or 5  $L_2$ (s) are identical to or different from each other, when  $a_1$  is 0,  $*(L_1)_{a_1}$ - $*$  is a single bond, and when  $a_2$  is 0,  $*(L_2)_{a_2}$ - $*$  is a single bond,  $R_1$  and  $R_2$  are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryloxy group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si( $Q_1$ )( $Q_2$ )( $Q_3$ ), —B( $Q_1$ )( $Q_2$ ), —C(=O)( $Q_1$ ), —N( $Q_1$ )( $Q_2$ ), —P(=O)( $Q_1$ )( $Q_2$ ), —P(=S)( $Q_1$ )( $Q_2$ ), —S(=O)( $Q_1$ )( $Q_2$ ), and —S(=O)<sub>2</sub>( $Q_1$ )( $Q_2$ ), at least one of  $R_1$  and  $R_2$  is an electron donating group,  $b_1$  and  $b_2$  are each independently an integer from 1 to 10, when  $b_1$  is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10  $R_1$ (s) are identical to or different from each other, and when  $b_2$  is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10  $R_2$ (s) are identical to or different from each other,  $c_1$  and  $c_2$  are each independently an integer from 1 to 10, at least one substituent of the substituted  $C_5$ - $C_{60}$  carbocyclic group, the substituted  $C_2$ - $C_{60}$  heterocyclic group, the substituted  $C_1$ - $C_{60}$  alkyl group, the substituted  $C_2$ - $C_{60}$  alkenyl group, the substituted  $C_2$ - $C_{60}$  alkynyl group, the substituted  $C_1$ - $C_{60}$  alkoxy group, the substituted  $C_3$ - $C_{10}$  cycloalkyl group, the substituted  $C_1$ - $C_{10}$  heterocycloalkyl group, the substituted  $C_3$ - $C_{10}$  cycloalkenyl group, the substituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, the substituted  $C_6$ - $C_{60}$  aryl group, the substituted  $C_6$ - $C_{60}$  aryloxy group, the substituted  $C_6$ - $C_{60}$  arylthio group, the substituted  $C_1$ - $C_{60}$  heteroaryl group, the substituted  $C_1$ - $C_{60}$  heteroaryloxy group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, and a  $C_1$ - $C_{60}$  alkoxy group; a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, and a  $C_1$ - $C_{60}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a  $C_3$ - $C_{10}$  cycloalkyl group, a  $C_1$ - $C_{10}$

heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>); a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group; a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>), Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted with a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and \* and \*' each indicate a binding site to a neighboring atom.


**[0007]** The embodiments may be realized by providing a heterocyclic compound represented by Formula 1A or Formula 1B:



&lt;Formula 1A&gt;

-continued

&lt;Formula 1B&gt;














**[0008]** wherein, in Formulae 1A and 1B, ring A<sub>1</sub> is selected from a C<sub>5</sub>-C<sub>60</sub> carbocyclic group and a C<sub>2</sub>-C<sub>60</sub> heterocyclic group, X<sub>1</sub> is S or Se, L<sub>1</sub> and L<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group and a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, a<sub>1</sub> and a<sub>2</sub> are each independently an integer from 0 to 5, when a<sub>1</sub> is 2, 3, 4, or 5, the 2, 3, 4, or 5 L<sub>1</sub>(s) are identical to or different from each other, and when a<sub>2</sub> is 2, 3, 4, or 5, the 2, 3, 4, or 5 L<sub>2</sub>(s) are identical to or different from each other, when a<sub>1</sub> is 0, \*-(L<sub>1</sub>)<sub>a1</sub>-\*' is a single bond, and when a<sub>2</sub> is 0, \*-(L<sub>2</sub>)<sub>a2</sub>-\*' is a single bond, R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryloxy group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>), at least one of R<sub>1</sub> and R<sub>2</sub> is an electron donating group, in Formula 1A, (R<sub>1</sub>)<sub>b1</sub>-(L<sub>1</sub>)<sub>a1</sub>-\*' and (R<sub>2</sub>)<sub>b2</sub>-(L<sub>2</sub>)<sub>a2</sub>-\*' are not halogen, a phenoxy group, a phenylthio group, an alkylthio group, or an alkoxy group, b<sub>1</sub> and b<sub>2</sub> are each independently an integer from 1 to 10, when b<sub>1</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>1</sub>(s) are identical to or different from each other, and when b<sub>2</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>2</sub>(s) are identical to or different from each other, c<sub>1</sub> and c<sub>2</sub> are each independently an integer from 1 to 10, at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub>



[0018] In Formula 1B, ring  $A_1$  may be selected from or include, e.g., a  $C_5$ - $C_{60}$  carbocyclic group and a  $C_2$ - $C_{60}$  heterocyclic group.

[0019] In an implementation, ring  $A_1$  may be selected from or include, e.g., a cyclohexadiene group, a dihydronaphthalene group, a dihydroanthracene group, a dihydrobenzoanthracene group, a dihydrophenanthrene group, a dihydrobenzophenanthrene group, a dihydrotriphenylene group, a dihydronaphthacene group, a dihydropyrene group, a dihydrochrysene group, a dihydropicene group, and a dihydrobenzopentaphene group.

[0020] In an implementation, ring  $A_1$  may be a group of, e.g., one of the following Formulae  $A_{1-1}$  to  $A_{1-12}$ . For example, one of the following Formulae  $A_{1-1}$  to  $A_{1-12}$  may share a C—C bond with the 5-membered ring of Formula 1B

 $A_{1-1}$  $A_{1-2}$  $A_{1-3}$  $A_{1-4}$  $A_{1-5}$  $A_{1-6}$  $A_{1-7}$  $A_{1-8}$  $A_{1-9}$  $A_{1-10}$  $A_{1-11}$  $A_{1-12}$ 

-continued

[0021] In Formulae 1A and 1B,  $X_1$  may be, e.g., S or Se. In an implementation,  $X_1$  may be S.

[0022] In Formulae 1A and 1B,  $L_1$  and  $L_2$  may each independently be selected from or include, e.g., a substituted or unsubstituted  $C_5$ - $C_{60}$  carbocyclic group, and a substituted or unsubstituted  $C_2$ - $C_{60}$  heterocyclic group.

[0023] In an implementation,  $L_1$  and  $L_2$  may each independently be selected or include, e.g., a phenylene group, a


pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a (quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, and a dibenzocarbazolylene group; and

**[0024]** a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-fluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, and a dibenzocarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano


group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isooxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzooxazolyl group, an isobenzooxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, and an imidazopyridinyl group.

**[0025]** In an implementation, L<sub>1</sub> to L<sub>2</sub> may each independently be or include, e.g., a phenylene group or a thiophenylene group.


**[0026]** In an implementation, L<sub>1</sub> and L<sub>2</sub> may each independently be selected from or include, e.g., a group represented by one of the following Formulae 3-1 to 3-22.




3-1



3-2









3-3




3-4

-continued




-continued

3-5




3-11

3-6




3-12

3-7




3-13

3-8




3-14

3-9




3-15


3-10



3-16




3-17




3-18


-continued




3-19



3-20



3-21



3-22

[0027] In Formulae 3-1 to 3-22,

[0028]  $Y_1$  may be, e.g., O, S,  $C(Z_3)(Z_4)$ ,  $N(Z_5)$ , or  $Si(Z_6)(Z_7)$ ,

[0029]  $Z_1$  to  $Z_7$  may each independently be selected from or include, e.g., hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, a silolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indolyl group, an isoindolyl group, an

indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzosilolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthyridinyl group, an azafuorenyl group, an azaspiro-bifluorenyl group, an azacarbazolyl group, an azadibenzofuranyl group, an azadibenzothiophenyl group, an azadibenzosilolyl group,  $-Si(Q_{31})(Q_{32})(Q_{33})$ ,  $-N(Q_{31})(Q_{32})$ ,  $-B(Q_{31})(Q_{32})$ ,  $-C(=O)(Q_{31})$ ,  $-S(=O)_2(Q_{31})$ , and  $-P(=O)(Q_{31})(Q_{32})$ .

[0030]  $Q_{31}$  to  $Q_{33}$  may each independently be selected from or include, e.g.,

[0031] a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazoliny group; and

[0032] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, and a quinazoliny group, each substituted with at least one selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, and a phenyl group,

[0033]  $d_2$  may be, e.g., an integer from 0 to 2,

[0034]  $d_3$  may be, e.g., an integer from 0 to 3,

[0035]  $d_4$  may be, e.g., an integer from 0 to 4,

[0036]  $d_5$  may be, e.g., an integer from 0 to 5,

[0037]  $d_6$  may be, e.g., an integer from 0 to 6,

[0038]  $d_8$  may be, e.g., an integer from 0 to 8, and

[0039] \* and \*' each indicate a binding site to a neighboring atom.

[0040] In Formulae 1A and 1B,  $a_1$  and  $a_2$  may each independently be, e.g., an integer from 0 to 5.  $a_1$  indicates the number of  $L_1(s)$ , wherein, when  $a_1$  is 2, 3, 4, or 5, the 2, 3, 4, or 5  $L_1(s)$  may be identical to or different from each other.  $a_2$  may be described by referring to the description of  $a_1$ .

[0041] When  $a_1$  is 0,  $*(L_1)_{a_1}$  may be a single bond, and when  $a_2$  is 0,  $*(L_2)_{a_2}$  may be a single bond.

[0042] In an implementation,  $a_1$  and  $a_2$  may each independently be, e.g., 0, 1, or 2.

[0043] In an implementation, in Formula 1A, at least one of  $a_1$  and  $a_2$  may not be 0.

[0044] In Formulae 1A and 1B,  $R_1$  and  $R_2$  may each independently be selected from or include, e.g., hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$

heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryloxy group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>). In an implementation, at least one selected from R<sub>1</sub> and R<sub>2</sub> may include an electron donating group.

**[0045]** Here, Q<sub>1</sub> to Q<sub>3</sub> are respectively defined the same as described above.

**[0046]** In an implementation, in Formula 1A, (R<sub>1</sub>)<sub>b1</sub>-(L<sub>1</sub>)<sub>a1</sub>-\* may not be a halogen, a phenoxy group, a phenylthio group, an alkylthio group, or an alkoxy group and (R<sub>2</sub>)<sub>b2</sub>-(L<sub>2</sub>)<sub>a2</sub>-\* may not be a halogen, a phenoxy group, a phenylthio group, an alkylthio group, or an alkoxy group

**[0047]** In an implementation, R<sub>1</sub> and R<sub>2</sub> may each independently be selected from or include, e.g.,

**[0048]** hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

**[0049]** a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD<sub>3</sub>, —CD<sub>2</sub>H, —CDH<sub>2</sub>, —CF<sub>3</sub>, —CF<sub>2</sub>H, —CFH<sub>2</sub>, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>);

**[0050]** a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

**[0051]** a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD<sub>3</sub>, —CD<sub>2</sub>H, —CDH<sub>2</sub>, —CF<sub>3</sub>, —CF<sub>2</sub>H, —CFH<sub>2</sub>, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>) and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>); and

**[0052]** —N(Q<sub>1</sub>)(Q<sub>2</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —P(Q<sub>1</sub>)(Q<sub>2</sub>), and —C(=O)(Q<sub>1</sub>).

**[0053]** In an implementation, at least one selected from R<sub>1</sub> and R<sub>2</sub> may include an electron donating group.

**[0054]** Here, Q<sub>1</sub> to Q<sub>3</sub> and Q<sub>31</sub> to Q<sub>33</sub> are respectively defined the same as described above.


**[0055]** In an implementation, R<sub>1</sub> and R<sub>2</sub> may each independently be selected from or include, e.g.,

**[0056]** hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino


group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and

**[0057]** a groups represented by one of the following Formulae 5-1 to 5-6.


**[0058]** In an implementation, at least one selected from R<sub>1</sub> and R<sub>2</sub> may include an electron donating group.




&lt;Formula 5-1&gt;




&lt;Formula 5-2&gt;




&lt;Formula 5-3&gt;



&lt;Formula 5-4&gt;



&lt;Formula 5-5&gt;



&lt;Formula 5-6&gt;

**[0059]** In Formulae 5-1 to 5-6,

**[0060]** K<sub>1</sub> may be selected from, e.g., \*—C(R<sub>2</sub>)(R<sub>30</sub>)—\*, \*—N(R<sub>20</sub>)—\*, and \*—Si(R<sub>20</sub>)(R<sub>30</sub>)—\*,

**[0061]** R<sub>20</sub>, R<sub>30</sub>, R<sub>40</sub>, and R<sub>50</sub> may be defined the same as R<sub>1</sub> and R<sub>2</sub> of Formulae 1A and 1B,

**[0062]** b<sub>40</sub> and b<sub>50</sub> may each independently be, e.g., an integer from 1 to 4, and

**[0063]** \* indicates a binding site to a neighboring atom.


**[0064]** b<sub>1</sub> and b<sub>2</sub> in Formulae 1A and 1B may each independently be an integer from 1 to 10. When b<sub>1</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>1</sub>(s) may be identical to or different from each other, and when b<sub>2</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>2</sub>(s) may be identical to or different from each other.

**[0065]** In an implementation, b<sub>1</sub> and b<sub>2</sub> may each independently be, e.g., 1 or 2.


**[0066]** c<sub>1</sub> and c<sub>2</sub> in Formula 1B may each independently be, e.g., an integer from 1 to 10.

**[0067]** In an implementation, c<sub>1</sub> and c<sub>2</sub> may each independently be, e.g., 1 or 2.


**[0068]** In an implementation, the compound represented by Formula 1B may be represented by, e.g., one of the following Formulae 1B-1 to 1B-4.




&lt;Formula 1B-1&gt;



&lt;Formula 1B-2&gt;



&lt;Formula 1B-3&gt;




&lt;Formula 1B-4&gt;

**[0069]** In Formulae 1B-1 to 1B-4, X<sub>1</sub> may be defined the same as in Formula 1B, L<sub>11</sub> and L<sub>12</sub> may be defined the same as L<sub>1</sub> and L<sub>2</sub> of Formula 1B, a<sub>11</sub> and a<sub>12</sub> may be defined the



same as a1 and a2 of Formula 1B, and  $R_{11}$  and  $R_{12}$  may be defined the same as  $R_1$  and  $R_2$  of Formula 1B.

[0070] In an implementation, the compound represented by Formula 1B may be represented by, e.g., one of the following Formulae 1B-a to 1B-d.







-continued

<Formula 1B-c>








[0071] In Formulae 1B-a to 1B-d,  $L_{11}$  and  $L_{12}$  may be defined the same as  $L_1$  and  $L_2$  of Formula 1B, a11 and a12 may be defined the same as a1 and a2 of Formula 1B, and  $R_{11}$  and  $R_{12}$  may be defined the same as  $R_1$  and  $R_2$  of Formula 1B.

[0072] In an implementation, the compound represented by Formula 1A or 1B may be, e.g., one of the following Compounds 1 to 37.




-continued




-continued

11




12


13




14



15





16



-continued



17

18




19


20



21





22




23

24




-continued


25




26




27




28




29




30



31




32





-continued

33




34


35



36



37



**[0073]** The heterocyclic compound represented by Formula 1A or 1B may include an oxidized thiadiazole-based core. Light in an infrared ray (IR) wavelength range may be emitted by employing a molecular structure including a strong electron donating substituent (e.g., according to Formula 1A or Formula 1B), and the luminescent efficiency may be improved by a thermally activated delayed fluorescent (TADF) mechanism.

**[0074]** An electronic device, e.g., an organic light-emitting device, which includes the heterocyclic compound, may have high external quantum efficiency may enable a long wavelength shift of a maximum emission wavelength, the

electronic device may be suitable for near infrared ray (NIR) emission, and may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

**[0075]** A synthesis method of the heterocyclic compound represented by Formula 1A or 1B may be recognized by those of ordinary skill in the art by referring to Examples provided below.

**[0076]** At least one of the heterocyclic compounds represented by Formula 1A or 1B may be used or included between a pair of electrodes of the organic light-emitting device. For example, the heterocyclic compound may be included in a hole transport region, an electron transport

region, and/or an emission layer. In an implementation, the heterocyclic compound represented by Formula 1A or 1B may be used as a material for forming a capping layer disposed outside the pair of electrodes of the organic light-emitting device.

**[0077]** Another aspect provides an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and including an emission layer. In an implementation, the emission layer may include the heterocyclic compound represented by Formula 1A or 1B.

**[0078]** The expression “(an organic layer) includes at least one of the heterocyclic compound” may mean that “(an organic layer) may include one heterocyclic compound belonging to the category of Formula 1A or 1B or two or more heterocyclic compounds belonging to the category of Formula 1A or 1B.


**[0079]** For example, the organic layer may include only Compound 1 as the heterocyclic compound. In this regard,

Compound 1 may exist only in the emission layer of the organic light-emitting device. In an implementation, the organic layer may include Compound 1 and Compound 2 as the heterocyclic compound. In this case, the Compound 1 and Compound 2 may exist in the same layer (for example, Compound 1 and Compound 2 may exist in the emission layer), or may exist in different layers (for example, Compound 1 may exist in the emission layer and Compound 2 may exist in the electron transport layer).

**[0080]** In an implementation, the emission layer may include a host and a dopant.

**[0081]** In an implementation, the dopant may include a compound represented by Formula 1A or Formula 1B.

**[0082]** In an implementation, the host may include, e.g., at least one selected from a compound represented by Formula 11-HT, a compound represented by Formula 21-HT, a compound represented by Formula 31-HT, a compound represented by Formula 41-HT, a compound represented by Formula 51-HT, a compound represented by Formula 61-ET, and a compound represented by Formula 71-ET:



**[0083]** In Formulae 11-HT, 21-HT, 31-HT, 41-HT, and 51-HT,

**[0084]**  $X_{21}$  may be, e.g., O, S, N-[(L<sub>24</sub>)<sub>a24</sub>-R<sub>24</sub>], B-[(L<sub>24</sub>)<sub>a24</sub>-R<sub>24</sub>], P-[(L<sub>24</sub>)<sub>a24</sub>-R<sub>24</sub>], C(R<sub>24</sub>)(R<sub>25</sub>), or Si(R<sub>24</sub>)(R<sub>25</sub>),

**[0085]** A<sub>11</sub>, A<sub>12</sub>, and A<sub>21</sub> to A<sub>24</sub> may each independently be or include, e.g., a C<sub>5</sub>-C<sub>30</sub> carbocyclic group or a  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group,

**[0086]** A<sub>31</sub> may be or may include, e.g., a substituted or unsubstituted C<sub>5</sub>-C<sub>30</sub> carbocyclic group in which two or more rings are condensed with each other or a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group in which two or more rings are condensed with each other, wherein A<sub>31</sub> may not be a substituted or unsubstituted spirobifluorene group and a substituted or unsubstituted anthracene group,

**[0087]** L<sub>11</sub> to L<sub>13</sub>, L<sub>21</sub> to L<sub>25</sub>, L<sub>31</sub>, L<sub>41</sub> to L<sub>43</sub>, and L<sub>51</sub> to L<sub>55</sub> may each independently be selected from or include, e.g.,

**[0088]** a single bond, a C<sub>5</sub>-C<sub>30</sub> carbocyclic group, and a  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group; and

**[0089]** a C<sub>5</sub>-C<sub>30</sub> carbocyclic group and a  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group, each substituted with at least one selected from hydrogen, deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>51</sub>)(Q<sub>52</sub>)(Q<sub>53</sub>), —N(Q<sub>51</sub>)(Q<sub>52</sub>), and —B(Q<sub>51</sub>)(Q<sub>52</sub>),

**[0090]** a11 to a13, a21 to a25, a31, a41 to a43, and a51 to a55 may each independently be, e.g., an integer from 1 to 5,

**[0091]** R<sub>11</sub>, R<sub>12</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>21</sub>, R<sub>22</sub>, and R<sub>24</sub> to R<sub>29</sub> may each independently be selected from or include, e.g.,

**[0092]** hydrogen, deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group;

**[0093]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub>

cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>51</sub>)(Q<sub>52</sub>)(Q<sub>53</sub>), —N(Q<sub>51</sub>)(Q<sub>52</sub>), and —B(Q<sub>51</sub>)(Q<sub>52</sub>); and

**[0094]** —Si(Q<sub>41</sub>)(Q<sub>42</sub>)(Q<sub>43</sub>), —N(Q<sub>41</sub>)(Q<sub>42</sub>), and —B(Q<sub>41</sub>)(Q<sub>42</sub>),

**[0095]** R<sub>13</sub>, R<sub>23</sub>, R<sub>31</sub>, R<sub>41</sub> to R<sub>43</sub>, and R<sub>51</sub> to R<sub>54</sub> may each independently be selected from or include, e.g.,

**[0096]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group;

**[0097]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>51</sub>)(Q<sub>52</sub>)(Q<sub>53</sub>), —N(Q<sub>51</sub>)(Q<sub>52</sub>) and —B(Q<sub>51</sub>)(Q<sub>52</sub>); and

**[0098]** —Si(Q<sub>41</sub>)(Q<sub>42</sub>)(Q<sub>43</sub>), —N(Q<sub>41</sub>)(Q<sub>42</sub>) and —B(Q<sub>41</sub>)(Q<sub>42</sub>).

**[0099]** In an implementation, R<sub>51</sub> and R<sub>52</sub> may be separate or may optionally be linked to form a saturated or unsaturated ring, and R<sub>53</sub> and R<sub>54</sub> may be separate or may optionally be linked to form a saturated or unsaturated ring,

**[0100]** b31 may be, e.g., an integer from 1 to 10, and

**[0101]** Q<sub>41</sub> to Q<sub>43</sub> and Q<sub>51</sub> to Q<sub>53</sub> may each independently be selected from or include, e.g., hydrogen, deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

**[0102]** In Formulae 61-ET and 71-ET,

**[0103]**  $X_{71}$  may be, e.g., O, S, N-[( $L_{74}$ )<sub>a74</sub>-R<sub>74</sub>], B-[( $L_{74}$ )<sub>a74</sub>-R<sub>74</sub>], P-[( $L_{74}$ )<sub>a74</sub>-R<sub>74</sub>], C(R<sub>74</sub>)(R<sub>75</sub>), or Si(R<sub>74</sub>)(R<sub>75</sub>),

**[0104]**  $A_{61}$ ,  $A_{62}$  and  $A_{71}$  to  $A_{74}$  may each independently be or include, e.g., a C<sub>5</sub>-C<sub>30</sub> carbocyclic group or a C<sub>2</sub>-C<sub>30</sub> heterocyclic group,

**[0105]**  $L_{61}$  to  $L_{63}$  and  $L_{71}$  to  $L_{75}$  may each independently be selected from or include, e.g., a single bond, a substituted or unsubstituted C<sub>5</sub>-C<sub>30</sub> carbocyclic group, and a substituted or unsubstituted C<sub>2</sub>-C<sub>30</sub> heterocyclic group,

**[0106]**  $a_{61}$  to  $a_{63}$  and  $a_{71}$  to  $a_{75}$  may each independently be, e.g., an integer from 1 to 5, and

**[0107]**  $R_{61}$  to  $R_{65}$  and  $R_{71}$  to  $R_{79}$  may each independently be selected from or include, e.g., hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)<sub>2</sub>(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>).

**[0108]** In an implementation, in Formula 61-ET, i) at least one selected from  $A_{61}$  and  $A_{62}$  may include, e.g., a  $\pi$  electron-depleted nitrogen-containing C<sub>2</sub>-C<sub>30</sub> heterocyclic group, or ii) at least one selected from  $R_{61}$  to  $R_{65}$  may each independently be selected from or include, e.g., a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>60</sub> heteroaryl group, and a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-containing monovalent non-aromatic condensed heteropolycyclic group.

**[0109]** In an implementation, in Formula 71-ET, i) at least one selected from  $A_{71}$  to  $A_{74}$  may include, e.g., a  $\pi$  electron-depleted nitrogen-containing C<sub>2</sub>-C<sub>30</sub> heterocyclic group, or ii) at least one selected from  $R_{71}$  to  $R_{73}$  and  $R_{76}$  to  $R_{79}$  may each independently be selected from or include, e.g., a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>60</sub> heteroaryl group, and a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-containing monovalent non-aromatic condensed heteropolycyclic group.

**[0110]** In an implementation, at least one substituent of the substituted C<sub>5</sub>-C<sub>30</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>30</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0111]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)<sub>2</sub>(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

**[0112]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)<sub>2</sub>(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

**[0113]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

**[0114]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)<sub>2</sub>(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>) and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

**[0115]** —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)<sub>2</sub>(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>), and

**[0116]** Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted with a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted


with a C<sub>6</sub>-C<sub>60</sub> aryl group, a terphenyl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group substituted with a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group substituted with a C<sub>6</sub>-C<sub>60</sub> aryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.

[0117] In an implementation, A<sub>11</sub>, A<sub>12</sub> and A<sub>21</sub> to A<sub>24</sub> may each independently be selected from or include, e.g., a benzene group, a naphthalene group, a pyrrole group, a cyclopentene group, a furan group, a thiophene group, an indole group, an indene group, a benzofuran group, a benzothiophene group, a carbazole group, a fluorene group, a dibenzofuran group, and a dibenzothiophene group,


[0118] A<sub>31</sub> may be selected from or include, e.g., a group of Formulae A31-1 to A31-38,

[0119] A<sub>61</sub>, A<sub>62</sub>, and A<sub>71</sub> to A<sub>74</sub> may each independently be selected from or include, e.g., a benzene group, a naphthalene group, a pyrrole group, a cyclopentene group, a furan group, a thiophene group, an indole group, an indene group, a benzofuran group, a benzothiophene group, a carbazole group, a fluorene group, a dibenzofuran group, a dibenzothiophene group, a pyridine group, a pyrimidine group, a pyrazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a benzoisoquinoline group, a quinoxaline group and a benzoquinoxaline group.


-continued




A31-1




A31-2




A31-3




A31-4




A31-5




A31-6




A31-7




A31-8




A31-9



A31-10




A31-11




A31-12


-continued




A31-13




A31-14




A31-15



A31-16




A31-17




A31-18


-continued




A31-19




A31-20




A31-21




A31-22



A31-23




A31-24




A31-25

-continued



-continued



**[0120]** In an implementation, the host may include at least one selected from, e.g., a compound represented by Formula 11-HT, a compound represented by Formula 21-HT, a compound represented by Formula 31-HT, a compound represented by Formula 41-HT, and a compound represented by Formula 51-HT.

**[0121]** In an implementation, the host may include at least one selected from, e.g., a compound represented by Formula 61-ET and a compound represented by Formula 71-ET.

**[0122]** In an implementation, the compound included in the emission layer and represented by Formula 1A or 1B may be, e.g., an NIR-emitting compound having a maximum emission wavelength of about 700 nm or more.

[0123] In an implementation, the compound included in the emission layer and represented by Formula 1A or 1B may be, e.g., a TADF emitter, and the emission layer may emit delayed fluorescence.

[0124] In an implementation, the first electrode of the organic light-emitting device may be an anode, the second electrode of the organic light-emitting device may be a cathode, the organic layer may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode, the hole transport region may include a hole injection layer, a hole transport layer, a buffer layer, an electron blocking layer, or any combination thereof, and the electron transport region may include a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

[0125] In an implementation, the hole transport region may include a charge-generation material. For example, the hole transport region may include a p-dopant, and the p-dopant may have a lowest unoccupied molecular orbital (LUMO) level of about  $-3.5$  eV or less. For example, the p-dopant may include a cyano group-containing compound.

[0126] In an implementation, the electron transport region may include a triazole-containing compound or a benzotriazole-containing compound, and may further include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth metal complex, a rare earth metal complex, or a combination thereof.

[0127] The term “an organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of an organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0128] [Description of FIG. 1]

[0129] FIG. 1 illustrates a schematic cross-sectional view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0130] Hereinafter, the structure of an organic light-emitting device 10 according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIG. 1.

[0131] [First Electrode 110]

[0132] In an implementation, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. For use as the substrate, the substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

[0133] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for forming the first electrode 110 may be selected from materials with a high work function to facilitate hole injection.

[0134] The first electrode 110 may be a reflective electrode, a semi-reflective electrode, or a transmissive electrode. In an implementation, when the first electrode 110 is a transmissive electrode, a material for forming a first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide ( $\text{SnO}_2$ ), zinc oxide (ZnO), and any combinations thereof. In an implementation, when the first electrode 110 is a semi-transmissive electrode or a reflectable electrode, a material for forming a first electrode may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and a combinations thereof.

[0135] In an implementation, the first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. In an implementation, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

[0136] [Organic Layer 150]

[0137] The organic layer 150 is disposed on the first electrode 110. The organic layer 150 may include an emission layer.

[0138] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer and an electron transport region between the emission layer and the second electrode 190.

[0139] In an implementation, the organic layer 150 may include the heterocyclic compound according to an embodiment.

[0140] [Hole Transport Region in Organic Layer 150]

[0141] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.






[0142] The hole transport region may include at least one layer selected from a hole injection layer, a first hole transport layer, a second hole transport layer, an emission auxiliary layer, and an electron blocking layer.

[0143] In an implementation, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein for each structure, constituting layers are sequentially stacked from the first electrode 110 in this stated order.

[0144] In an implementation, the hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (or NPD), (3-NPB, TPD, spiro-TPD, spiro-NPB, methylated-NPB, TAPC, HMTDP, 4,4',

4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202.

-continued





ylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylylene group, a phenanthrenylylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylylylene group, a pyrenylylene group, a chrysenylylene group, a naphthacenylylene group, a picenylylene group, a perylylylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group; and

**[0155]** a phenylene group, a pentalenylylene group, an indenylene group, a naphthylene group, an azulenylylene group, a heptalenylylene group, an indacenylylene group, an acenaphthylene group, a fluorenylylene group, a spiro-bifluorenylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenylylene group, a phenanthrenylylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylylylene group, a pyrenylylene group, a chrysenylylene group, a naphthacenylylene group, a picenylylene group, a perylylylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a  $C_1$ - $C_{10}$  alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylylene group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylylylene group, a pyrenyl group, a chrysenyl group, a naphthacenylylene group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ), and —N( $Q_{31}$ )( $Q_{32}$ ), and

**[0156]**  $Q_{31}$  to  $Q_{33}$  may each independently be selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

**[0157]** In an implementation, xa1 to xa4 may each independently be 0, 1, or 2.

**[0158]** In an implementation, xa5 may be 1, 2, 3, or 4.

**[0159]** In an implementation,  $R_{201}$  to  $R_{204}$  and  $Q_{201}$  may each independently be selected from or include, e.g.,

**[0160]** a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylylene group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylylylene group, a pyrenyl group, a chrysenyl group, a naphthacenylylene group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

**[0161]** a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenylylene group, a phenalenylylene group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylylylene group, a pyrenyl group, a chrysenyl group, a naphthacenylylene group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a  $C_1$ - $C_{10}$  alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenylylene group, a phenalenylylene group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylylylene group, a pyrenyl group, a chrysenyl group, a naphthacenylylene group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ), and —N( $Q_{31}$ )( $Q_{32}$ ), and

**[0162]**  $Q_{31}$  to  $Q_{33}$  are respectively defined the same as described above.

**[0163]** In an implementation, in Formula 201, at least one selected from  $R_{201}$  to  $R_{203}$  may each independently be selected from or include, e.g.,

**[0164]** a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

**[0165]** a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected

from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0166] In an implementation, in Formula 202, i) R<sub>201</sub> and R<sub>202</sub> may be linked via a single bond, and/or ii) R<sub>203</sub> and R<sub>204</sub> may be linked via a single bond.

[0167] In an implementation, in Formula 202, at least one selected from R<sub>201</sub> to R<sub>204</sub> may be selected from or include, e.g.,


[0168] a carbazolyl group; and

[0169] a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0170] The compound represented by Formula 201 may be represented by Formula 201A:




[0171] In an implementation, the compound represented by Formula 201 may be represented by Formula 201A(1).




[0172] In an implementation, the compound represented by Formula 201 may be represented by Formula 201A-1.



[0173] In an implementation, the compound represented by Formula 202 may be represented by Formula 202A.



[0174] In an implementation, the compound represented by Formula 202 may be represented by Formula 202A-1.



[0175] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,

[0176] L<sub>201</sub> to L<sub>203</sub>, xa1 to xa3, xa5, and R<sub>202</sub> to R<sub>204</sub> may respectively be defined the same as described above,

[0177] R<sub>211</sub> and R<sub>212</sub> may respectively be defined the same as described in connection with R<sub>203</sub>,


[0178] R<sub>213</sub> to R<sub>217</sub> may each independently be selected from or include, e.g., hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substi-

tuted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl


group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0179] In an implementation, the hole transport region may include at least one compound selected from Compounds HT1 to HT39.


HT1




HT2



HT3




HT4




-continued


HT5




HT6



HT7




HT8




-continued


HT9




HT10



HT11





HT12





-continued  
HT13

HT14





HT15

HT16




HT17

HT18




-continued


HT19




HT20



HT21




HT22




-continued


HT23




HT24




HT25




-continued  
HT26




HT27




HT28




HT29



HT30




HT31




-continued


HT32




HT33




HT34




HT35



HT36





HT37



-continued

HT38

HT39



[0180] A thickness of the hole transport region may be from about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes at least one selected from a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, e.g., about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å.

[0181] In an implementation, a thickness of the first hole transport layer or the second hole transport layer may be from about 200 Å to about 400 Å.

[0182] When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within any of these ranges, excellent hole transport characteristics may be obtained without a substantial increase in driving voltage.

[0183] The emission auxiliary layer may help increase light emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer. The electron blocking layer may reduce or eliminate the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the aforementioned materials.

[0184] [P-Dopant]

[0185] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0186] The charge-generation material may be, for example, a p-dopant.

[0187] In an implementation, the p-dopant may have a lowest unoccupied molecular orbital (LUMO) level of  $-3.5$  eV or less.

[0188] In an implementation, the p-dopant may include, e.g., at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound.

[0189] In an implementation, the p-dopant may include at least one selected from, e.g.,


[0190] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

[0191] a metal oxide, such as tungsten oxide or molybdenum oxide;


[0192] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0193] a compound represented by Formula 221.


&lt;HAT-CN&gt;



&lt;F4-TCNQ&gt;



-continued



&lt;Formula 221&gt;

[0194] In Formula 221,

[0195]  $R_{221}$  to  $R_{223}$  may each independently be selected from or include, e.g., a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one selected from  $R_{221}$  to  $R_{223}$  may include at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a  $C_1$ - $C_{20}$  alkyl group substituted with —F, a  $C_1$ - $C_{20}$  alkyl group substituted with —C<sub>1</sub>, a  $C_1$ - $C_{20}$  alkyl group substituted with —Br, and a  $C_1$ - $C_{20}$  alkyl group substituted with —I.

[0196] [Emission Layer in Organic Layer 150]

[0197] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure, and the stacked structure may include two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer. The two or more layers may be in direct contact with each other. In some embodiments, the two or more layers may be separated from each other. In one or more embodiments, the emission layer may include two or more materials. The two or more materials may include a red light-emitting material, a green light-emitting material, or a blue light-emitting material. The two or more materials may be mixed with each other in a single layer. The two or more materials mixed with each other in the single layer may emit white light.

[0198] In an implementation, the emission layer may include a host and a dopant. For example, the heterocyclic compound included in the emission layer may be a dopant, and the emission layer may further include a host. The dopant may further include at least one selected from a phosphorescent dopant and fluorescent dopant.

[0199] In an implementation, the heterocyclic compound included in the emission layer may be a near infrared ray (NIR)-emitting compound having a maximum emission wavelength of 700 nm.

[0200] In an implementation, the heterocyclic compound included in the emission layer may be a thermally activated delayed fluorescent (TADF) emitter, and in this regard, the emission layer may emit delayed fluorescence.

[0201] In an implementation, an amount of the dopant in the emission layer may be in a range of, e.g., about 0.01 parts by weight to about 30 parts by weight based on 100 parts by weight of the host.

[0202] In an implementation, a thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent emission characteristics may be obtained without a substantial increase in driving voltage.

[0203] [Host in Emission Layer]

[0204] The emission layer may include a compound represented by Formula 301:



[0205] In an implementation, in Formula 301,

[0206]  $Ar_{301}$  may be or may include, e.g., a substituted or unsubstituted  $C_5$ - $C_{60}$  carbocyclic group or a substituted or unsubstituted  $C_1$ - $C_{60}$  heterocyclic group,

[0207]  $xb11$  may be 1, 2, or 3,

[0208]  $L_{301}$  may be selected from or include, e.g., a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0209]  $xb1$  may be an integer from 0 to 5,

[0210]  $R_{301}$  may be selected from or include, e.g., deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si( $Q_{301}$ )( $Q_{302}$ )( $Q_{303}$ ), —N( $Q_{301}$ )( $Q_{302}$ ), —B( $Q_{301}$ )( $Q_{302}$ ), —C(=O)( $Q_{301}$ ), —S(=O)<sub>2</sub>( $Q_{301}$ ), and —P(=O)( $Q_{301}$ )( $Q_{302}$ ),

[0211]  $xb21$  may be an integer from 1 to 5, and

[0212]  $Q_{301}$  to  $Q_{303}$  may each independently be selected from or include, e.g., a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0213] In one embodiment, in Formula 301,  $Ar_{301}$  may be selected from:


[0214] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and

[0215] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>), and


[0216] Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0217] In Formula 301, when xb11 is two or more, two or more Ar<sub>301</sub>(s) may be linked via a single bond.

[0218] In an implementation, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:



&lt;Formula 301-1&gt;



&lt;Formula 301-2&gt;

[0219] In Formulae 301-1 to 301-2,

[0220] A<sub>301</sub> to A<sub>304</sub> may each independently be selected from or include, e.g., a benzene, a naphthalene, a phenanthrene, a fluoranthene, a triphenylene, a pyrene, a chrysene, a pyridine, a pyrimidine, an indene, a fluorene, a spiro-bifluorene, a benzofluorene, a dibenzofluorene, an indole, a carbazole, benzocarbazole, dibenzocarbazole, a furan, a benzofuran, a dibenzofuran, a naphthofuran, a benzonaphthofuran, dinaphthofuran, a thiophene, a benzothiophene, a dibenzothiophene, a naphthothiophene, a benzonaphthothiophene, and a dinaphthothiophene,

[0221] X<sub>301</sub> may be O, S, or N-[(L<sub>304</sub>)xb4-R<sub>304</sub>],

[0222] R<sub>311</sub> to R<sub>314</sub> may each independently be selected from or include, e.g., hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group,

a biphenyl group, a terphenyl group, a naphthyl group —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),

[0223] xb22 and xb23 may each independently be 0, 1, or 2,

[0224] L<sub>301</sub>, xb1, R<sub>301</sub>, and Q<sub>31</sub> to Q<sub>33</sub> may respectively be defined the same as described above,

[0225] L<sub>302</sub> to L<sub>304</sub> may each independently be defined the same as described in connection with L<sub>301</sub>,

[0226] xb2 to xb4 may each independently be defined the same as described in connection with xb1, and

[0227] R<sub>302</sub> to R<sub>304</sub> may each independently be defined the same as described in connection with R<sub>301</sub>.

[0228] For example, in Formulae 301, 301-1, and 301-2, L<sub>301</sub> to L<sub>304</sub> may each independently be selected from or include, e.g.,

[0229] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene

group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylene group, a quinolylylene group, an isoquinolylylene group, a benzoquinolylylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzimidazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a




group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group,  $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{C}(=\text{O})(\text{Q}_{31})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$ , and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ , and

**[0235]**  $\text{Q}_{31}$  to  $\text{Q}_{33}$  may respectively be defined the same as described above.


**[0236]** In an implementation, the host may include an alkaline earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), a Mg complex, and a Zn complex.

**[0237]** In an implementation, the host may include, e.g., at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55 and H-1a to H-15a.


-continued




H4




H5




H6




H7




H1




H2



H3




H8




H9

-continued


-continued




H10




H15




H11




H16




H12




H17




H13



H18




H14




H19

-continued




H20


H21




H22



H23




-continued




H24


H25



H26




-continued




H27

-continued




H30




H31


H28




H29




H32



H33




H34




-continued

-continued


H35




H38




H36




H39



H37










H40




-continued

-continued




-continued


H48




H49




H50




H51



H52




H53




-continued


H54




H55



H-1a




H-2a




-continued


-continued




H-3a




H-6a




H-4a



H-7a




H-5a




H-8a


-continued




H-9a



H-10a




H-11a




H-12a


-continued



H-13a




H-14a



H-15a

[0238] [Phosphorescent Dopant Included in Emission Layer in Organic Layer 150]

[0239] The phosphorescent dopant may include an organometallic compound represented by Formula 401:



[0240] In Formulae 401 and 402,

[0241] M may be selected from, e.g., iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0242] L<sub>401</sub> may be selected from ligands represented by Formula 402, xc1 may be 1, 2, or 3, and when xc1 is two or more, two or more L<sub>401</sub>(s) may be identical to or different from each other,

[0243] L<sub>402</sub> may be an organic ligand, xc2 may be an integer from 0 to 4, and when xc2 is two or more, two or more L<sub>402</sub>(s) may be identical to or different from each other,

[0244] X<sub>401</sub> to X<sub>404</sub> may each independently be nitrogen or carbon,

[0245] X<sub>401</sub> and X<sub>403</sub> may be linked via a single bond or a double bond, and X<sub>402</sub> and X<sub>404</sub> may be linked via a single bond or a double bond,

[0246] A<sub>401</sub> and A<sub>402</sub> may each independently be a C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a C<sub>1</sub>-C<sub>60</sub> heterocyclic group,

[0247] X<sub>405</sub> may be a single bond, \*—O—\*, \*—S—\*, \*—C(=O)—\*, \*—N(Q<sub>411</sub>)—\*, \*—C(Q<sub>411</sub>)(Q<sub>412</sub>)—\*, \*—C(Q<sub>411</sub>)=C(Q<sub>412</sub>)—\*, \*—C(Q<sub>411</sub>)=\*, or \*=C(Q<sub>411</sub>)=\*, wherein Q<sub>411</sub> and Q<sub>412</sub> may each independently be hydrogen, deuterium, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0248] X<sub>406</sub> may be a single bond, O, or S,

[0249] R<sub>401</sub> and R<sub>402</sub> may each independently be selected from or include, e.g., hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>20</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or

unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>401</sub>)(Q<sub>402</sub>)(Q<sub>403</sub>), —N(Q<sub>401</sub>)(Q<sub>402</sub>), —B(Q<sub>401</sub>)(Q<sub>402</sub>), —C(=O)(Q<sub>401</sub>), —S(=O)<sub>2</sub>(Q<sub>401</sub>), and —P(=O)(Q<sub>401</sub>)(Q<sub>402</sub>), wherein Q<sub>401</sub> to Q<sub>403</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, and a C<sub>1</sub>-C<sub>20</sub> heteroaryl group,

[0250] xc11 and xc12 may each independently be an integer from 0 to 10, and

[0251] \* and \*' in Formula 402 each indicate a binding site to M of Formula 401.

[0252] In an implementation, in Formula 402, A<sub>401</sub> and A<sub>402</sub> may each independently be selected from or include, e.g., a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan(furan) group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiothiophene group, an isobenzothiothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiothiophene group.

[0253] In an implementation, in Formula 402, i) X<sub>401</sub> may be nitrogen, and X<sub>402</sub> may be carbon, or ii) X<sub>401</sub> and X<sub>402</sub> may both be nitrogen.

[0254] In an implementation, in Formula 402, R<sub>401</sub> and R<sub>402</sub> may each independently be selected from or include, e.g.,

[0255] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

[0256] a C<sub>1</sub>-C<sub>20</sub> alkyl group and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0257] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuran group, and a dibenzothiothiophenyl group;

[0258] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

**[0259]** —Si(Q<sub>401</sub>)(Q<sub>402</sub>)(Q<sub>403</sub>), —N(Q<sub>401</sub>)(Q<sub>402</sub>), —B(Q<sub>401</sub>)(Q<sub>402</sub>), —C(=O)(Q<sub>401</sub>), —S(=O)<sub>2</sub>(Q<sub>401</sub>), and —P(=O)(Q<sub>401</sub>)(Q<sub>402</sub>), and

**[0260]** Q<sub>401</sub> to Q<sub>403</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.


**[0261]** In an implementation, in Formula 401, when xc1 is two or more, two A<sub>401</sub>(s) among a plurality of L<sub>401</sub>(s) may optionally be linked via a linking group, X<sub>407</sub>, or two A<sub>402</sub>(s) may optionally be linked via a linking group, X<sub>408</sub> (see Compounds PD1 to PD4 and PD7). X<sub>407</sub> and X<sub>408</sub> may each independently be a single bond, \*—O—\*, \*—S—\*, \*—C(=O)—\*, \*—N(Q<sub>413</sub>)—\*, \*—C(Q<sub>413</sub>)(Q<sub>414</sub>)—\*, or \*—C(Q<sub>413</sub>)=C(Q<sub>414</sub>)—\* (wherein Q<sub>413</sub> and Q<sub>414</sub> may each independently be hydrogen, deuterium, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group).

**[0262]** In an implementation, in Formula 401, L<sub>402</sub> may be a monovalent, divalent, or trivalent organic ligand. For example, L<sub>402</sub> may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), —C(=O), isonitrile, —CN, and phosphorus (for example, phosphine and phosphite).


**[0263]** In an implementation, the phosphorescent dopant may be, for example, selected from Compounds PD1 to PD25.

-continued


PD2




PD3




PD4








PD1



PD5




-continued




-continued

PD6




PD11

PD7




PD12

PD8




PD13

PD9








PD14

PD10




PD15

-continued




-continued

PD16




PD21

PD17




PD22

PD18




PD23

PD19



PD24

PD20




**[0264]** [Fluorescent Dopant in Emission Layer]

**[0265]** The fluorescent dopant may include an arylamine compound or a styrylamine compound.

**[0266]** The fluorescent dopant may include a compound represented by Formula 501:

**[0267]** The fluorescent dopant may include a compound represented by Formula 501:



**[0268]** In Formula 501,

**[0269]** Ar<sub>501</sub> may be or may include, e.g., a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group,

**[0270]** L<sub>501</sub> to L<sub>503</sub> may each independently be selected from or include, e.g., a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

**[0271]** xd1 to xd3 may each independently be an integer from 0 to 3,

**[0272]** R<sub>501</sub> and R<sub>502</sub> may each independently be selected from or include, e.g., a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> het-

eroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

**[0273]** xd4 may be an integer from 1 to 6.

**[0274]** In an implementation, in Formula 501, Ar<sub>501</sub> may be selected from or include, e.g.,

**[0275]** a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

**[0276]** a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

**[0277]** In an implementation, in Formula 501, L<sub>501</sub> to L<sub>503</sub> may each independently be selected from or include, e.g.,

**[0278]** a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group; and

**[0279]** a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro

group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0280] In an implementation, in Formula 501,  $R_{501}$  and  $R_{502}$  may each independently be selected from or include, e.g.,

[0281] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and


[0282] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ), and

[0283]  $Q_{31}$  to  $Q_{33}$  may each independently be selected from or include, e.g., a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


[0284] In an implementation, in Formula 501, xd4 may be 2.

[0285] In an implementation, the fluorescent dopant may be selected from Compounds FD1 to FD22.


FD1



FD2




FD3




-continued

FD4




-continued


FD7




FD5




FD8



FD6




FD9




-continued

-continued


FD10




FD11




FD12




FD13




FD14




FD15




FD16



FD17




FD18




-continued

FD19





-continued

FD21




FD22


FD20




[0286] In an implementation, the fluorescent dopant may be selected from compounds below.



DPVBi



DPAVBi



**[0287]** [Electron Transport Region in Organic Layer 150]

**[0288]** In an implementation, the electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

**[0289]** In an implementation, the electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer.

**[0290]** In an implementation, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein for each structure, constituting layers are sequentially stacked from an emission layer.

**[0291]** The electron transport region (e.g., a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include

a metal-free compound containing at least one  $\pi$  electron-depleted nitrogen-containing ring.


**[0292]** In an implementation, the “ $\pi$  electron-depleted nitrogen-containing ring” indicates a  $C_1$ - $C_{60}$  heterocyclic group having at least one  $*-N=*'$  moiety as a ring-forming moiety.

**[0293]** In an implementation, the “ $\pi$  electron-depleted nitrogen-containing ring” may be or may include, e.g., i) a 5-membered to 7-membered heteromonocyclic group having at least one  $*-N=*'$  moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one  $*-N=*'$  moiety are condensed with each other, or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one  $*-N=*'$  moiety, is condensed with at least one  $C_5$ - $C_{60}$  carbocyclic group.

**[0294]** Examples of the  $\pi$  electron-depleted nitrogen-containing ring may include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quiazoline, a cin-

noline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a thiadiazol, an imidazopyridine, an imidazopyrimidine, and an azacarbazole.

[0295] In an implementation, the electron transport region may include a compound represented by Formula 601:



[0296] In Formula 601,

[0297]  $\text{Ar}_{601}$  may be or may include, e.g., a substituted or unsubstituted  $\text{C}_5$ - $\text{C}_{60}$  carbocyclic group or a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  heterocyclic group,

[0298]  $\text{xe}11$  may be 1, 2, or 3,

[0299]  $\text{L}_{601}$  may be selected from or may include, e.g., a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkylene group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenylene group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  arylene group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0300]  $\text{xe}1$  may be an integer from 0 to 5,

[0301]  $\text{R}_{601}$  may be selected from or may include, e.g., a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  aryloxy group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  arylthio group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$ ,  $-\text{C}(=\text{O})(\text{Q}_{601})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{601})$ , and  $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$ ,

[0302]  $\text{Q}_{601}$  to  $\text{Q}_{603}$  may each independently be a  $\text{C}_1$ - $\text{C}_{10}$  alkyl group, a  $\text{C}_1$ - $\text{C}_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0303]  $\text{xe}21$  may be an integer from 1 to 5.

[0304] In an implementation, at least one of  $\text{Ar}_{601}(\text{s})$  in the number of  $\text{xe}11$  and  $\text{R}_{601}(\text{s})$  in the number of  $\text{xe}21$  may include the  $\pi$  electron-depleted nitrogen-containing ring.

[0305] In an implementation, ring  $\text{Ar}_{601}$  in Formula 601 may be selected from or may include, e.g.,

[0306] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a

phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0307] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1$ - $\text{C}_{20}$  alkyl group, a  $\text{C}_1$ - $\text{C}_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group,  $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$ , and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ , and


[0308]  $\text{Q}_{31}$  to  $\text{Q}_{33}$  may each independently be selected from  $\text{C}_1$ - $\text{C}_{10}$  alkyl group, a  $\text{C}_1$ - $\text{C}_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0309] When  $\text{xe}11$  in Formula 601 is two or more, two or more  $\text{Ar}_{601}(\text{s})$  may be linked via a single bond.

[0310] In an implementation,  $\text{Ar}_{601}$  in Formula 601 may be an anthracene group.

[0311] In an implementation, the compound represented by Formula 601 may be represented by Formula 601-1.

$\langle \text{Formula 601-1} \rangle$



[0312] In Formula 601-1,

[0313]  $\text{X}_{614}$  may be N or  $\text{C}(\text{R}_{614})$ ,  $\text{X}_{615}$  may be N or  $\text{C}(\text{R}_{615})$ ,  $\text{X}_{616}$  may be N or  $\text{C}(\text{R}_{616})$ , and at least one selected from  $\text{X}_{614}$  to  $\text{X}_{616}$  may be N,

[0314]  $\text{L}_{611}$  to  $\text{L}_{613}$  may each independently be defined the same as described in connection with  $\text{L}_{601}$ ,

[0315]  $\text{xe}611$  to  $\text{xe}613$  may each independently be defined the same as described in connection with  $\text{xe}1$ ,

**[0316]**  $R_{611}$  to  $R_{613}$  may each independently be defined the same as described in connection with  $R_{601}$ , and

**[0317]**  $R_{614}$  to  $R_{616}$  may each independently be selected from or include, e.g., hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

**[0318]** In an implementation, the  $L_{601}$  and  $L_{611}$  to  $L_{613}$  in Formulae 601 and 601-1 may each independently be selected from or include, e.g.,

**[0319]** a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazololylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazololylylene group, an isoxazololylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylene group, and an azacarbazolylylene group; and

**[0320]** a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazololylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazololylylene group, an isoxazololylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylylene group, a benzoxazolylene group, an isobenzoxazolylene group, a

triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylene group, and an azacarbazolylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group.

**[0321]** In an implementation, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

**[0322]** In an implementation, in Formulae 601 and 601-1,  $R_{601}$  and  $R_{611}$  to  $R_{613}$  may each independently be selected from or include, e.g.,

**[0323]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

**[0324]** a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluo-


renyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0325] —S(=O)<sub>2</sub>(Q<sub>601</sub>) and —P(=O)(Q<sub>601</sub>)(Q<sub>602</sub>), and


[0326] Q<sub>601</sub> and Q<sub>602</sub> may respectively be defined the same as described above.

[0327] In an implementation, the electron transport region may include at least one compound selected from Compounds ET1 to ET36.


ET1




ET2








ET3



-continued




-continued




-continued

ET10




-continued


ET13




ET11




ET14





ET12



ET15




-continued




-continued

ET16




ET18

ET17




ET19

ET20




-continued




ET21


-continued




ET24




ET22



ET25




ET23




ET26

-continued


ET27



ET28




ET29




-continued


ET30




ET31




ET32







-continued




-continued



[0328] In an implementation, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq<sub>3</sub>, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:






[0329] Thicknesses of the buffer layer, the hole blocking layer, and the electron control layer may each be in a range of about 20 Å to about 1,000 Å, e.g., about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

[0330] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

[0331] The electron transport region (e.g., the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0332] In an implementation, the metal-containing material may include at least one selected from an alkali metal complex and an alkaline earth metal complex. The alkali metal complex may include a metal ion selected from a Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, a Sr ion, and a Ba ion. In an implementation, a ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenyloxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.

[0333] In an implementation, the metal-containing material may include a Li complex. In an implementation, the Li complex may include, e.g., Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



[0334] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0335] In an implementation, the electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0336] In an implementation, the electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or a combinations thereof.

[0337] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In an implementation, the alkali metal may be Li or Cs.

[0338] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0339] The rare earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

[0340] The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare earth metal.

[0341] The alkali metal compound may be selected from alkali metal oxides, such as  $\text{Li}_2\text{O}$ ,  $\text{Cs}_2\text{O}$ , or  $\text{K}_2\text{O}$ , and alkali metal halides, such as  $\text{LiF}$ ,  $\text{NaF}$ ,  $\text{CsF}$ ,  $\text{KF}$ ,  $\text{LiI}$ ,  $\text{NaI}$ ,  $\text{CsI}$ , or  $\text{KI}$ . In an implementation, the alkali metal compound may be selected from  $\text{LiF}$ ,  $\text{Li}_2\text{O}$ ,  $\text{NaF}$ ,  $\text{LiI}$ ,  $\text{NaI}$ ,  $\text{CsI}$ , and  $\text{KI}$ .

[0342] The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as  $\text{BaO}$ ,  $\text{SrO}$ ,  $\text{CaO}$ ,  $\text{Ba}_x\text{Sr}_{1-x}\text{O}$  ( $0 < x < 1$ ), or  $\text{Ba}_x\text{Ca}_{1-x}\text{O}$  ( $0 < x < 1$ ). In an implementation, the alkaline earth-metal compound may be selected from  $\text{BaO}$ ,  $\text{SrO}$ , and  $\text{CaO}$ .

[0343] The rare earth metal compound may be selected from  $\text{YbF}_3$ ,  $\text{ScF}_3$ ,  $\text{ScO}_3$ ,  $\text{Y}_2\text{O}_3$ ,  $\text{Ce}_2\text{O}_3$ ,  $\text{GdF}_3$ , and  $\text{TbF}_3$ . In an implementation, the rare earth metal compound may be selected from  $\text{YbF}_3$ ,  $\text{ScF}_3$ ,  $\text{TbF}_3$ ,  $\text{YbI}_3$ ,  $\text{ScI}_3$ , and  $\text{Tb}_3$ .

[0344] In an implementation, the alkali metal complex, the alkaline earth-metal complex, and the rare earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, or the rare earth metal complex may be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenyloxazole, hydroxy phenylthiazole, hydroxy diphenyloxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene.

[0345] The electron injection layer may consist of or include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof, as described above. In an implementation, the electron injection layer may further include an organic material. When the electron

injection layer further includes an organic material, an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0346] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0347] [Second Electrode 190]

[0348] The second electrode 190 may be disposed on the organic layer 150 having such a structure. The second electrode 190 may be a cathode which is an electron injection electrode, and in this regard, a material for forming the second electrode 190 may be selected from metal, an alloy, an electrically conductive compound, and a combination thereof, which have a relatively low work function.

[0349] In an implementation, the second electrode 190 may include, e.g., at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO. The second electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0350] The second electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[0351] [Description of FIGS. 2 to 4]

[0352] An organic light-emitting device 20 of FIG. 2 may include a first capping layer 210, a first electrode 110, an organic layer 150, and a second electrode 190 which are sequentially stacked in this stated order, an organic light-emitting device 30 of FIG. 3 may include a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220 which are sequentially stacked in this stated order, and an organic light-emitting device 40 of FIG. 4 may include a first capping layer 210, a first electrode 110, an organic layer 150, a second electrode 190, and a second capping layer 220.

[0353] Regarding FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may be understood by referring to the description presented in connection with FIG. 1.

[0354] In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110, which is a semi-transmissive electrode or a transmissive electrode, and the first capping layer 210 toward the outside, and in the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190, which is a semi-transmissive electrode or a transmissive electrode, and the second capping layer 220 toward the outside.

[0355] The first capping layer 210 and the second capping layer 220 may help increase external luminescent efficiency according to the principle of constructive interference.


[0356] In an implementation, the first capping layer 210 and the second capping layer 220 may each independently

be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.


[0357] At least one selected from the first capping layer 210 and the second capping layer 220 may include at least one material selected from, e.g., carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrine derivatives, phthalocyanine derivatives, a naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one embodiment, at least one selected from the first capping layer 210 and the second capping layer 220 may each independently include an amine-based compound.

[0358] In an implementation, at least one selected from the first capping layer 210 and the second capping layer 220 may include the compound represented by Formula 201 or the compound represented by Formula 202.

[0359] In an implementation, the at least one selected from the first capping layer 210 and the second capping layer 220 may include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5



-continued



[0360] Hereinbefore, the organic light-emitting device according to an embodiment has been described in connection with FIGS. 1 to 4.

[0361] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0362] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by vacuum deposition, the deposition may be performed at, e.g., a deposition temperature of about 100° C. to about 500° C., a vacuum degree of about 10<sup>-8</sup> torr to about 10<sup>-3</sup> torr, and/or a deposition speed of about 0.01 Å/sec to about 100 Å/sec by taking into account a material to be included in a layer to be formed, and the structure of a layer to be formed.

[0363] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at, e.g., a coating speed of about 2,000 rpm to about 5,000 rpm and/or at a heat treatment temperature of about 80° C. to about 200° C. by taking into account a material to be included in a layer to be formed, and the structure of a layer to be formed.

[0364] [General Definition of Substituents]

[0365] The term “C<sub>1</sub>-C<sub>60</sub> alkyl group” as used herein refers to a linear or branched aliphatic saturated hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an isoamyl group, and a hexyl

group. The term “C<sub>1</sub>-C<sub>60</sub> alkylene group” as used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>60</sub> alkyl group.

[0366] The term “C<sub>2</sub>-C<sub>60</sub> alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>60</sub> alkenyl group.

[0367] The term “C<sub>2</sub>-C<sub>60</sub> alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and examples thereof include an ethynyl group, and a propynyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkynylene group” as used herein refers to a divalent group having the same structure as the C<sub>2</sub>-C<sub>60</sub> alkynyl group.

[0368] The term “C<sub>1</sub>-C<sub>60</sub> alkoxy group” as used herein refers to a monovalent group represented by —OA<sub>101</sub> (wherein A<sub>101</sub> is the C<sub>1</sub>-C<sub>60</sub> alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0369] The term “C<sub>3</sub>-C<sub>10</sub> cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkylene group” as used herein refers to a divalent group having the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkyl group.

[0370] The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group” as used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group. The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group.

[0371] The term C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group.

[0372] The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Non-limiting examples of the C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranlyl group, and a 2,3-dihydrothiophenyl group. The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group.

[0373] The term “C<sub>6</sub>-C<sub>60</sub> aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C<sub>6</sub>-C<sub>60</sub> arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting

examples of the C<sub>6</sub>-C<sub>60</sub> aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C<sub>6</sub>-C<sub>60</sub> aryl group and the C<sub>6</sub>-C<sub>60</sub> arylene group each include two or more rings, the rings may be fused to each other.

**[0374]** The term “C<sub>1</sub>-C<sub>60</sub> heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C<sub>1</sub>-C<sub>60</sub> heteroarylene group” as used herein refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C<sub>1</sub>-C<sub>60</sub> heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C<sub>1</sub>-C<sub>60</sub> heteroaryl group and the C<sub>1</sub>-C<sub>60</sub> heteroarylene group each include two or more rings, the rings may be condensed with each other.

**[0375]** The term “C<sub>6</sub>-C<sub>60</sub> aryloxy group” as used herein refers to —OA<sub>102</sub> (wherein A<sub>102</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group), and a C<sub>6</sub>-C<sub>60</sub> arylthio group used herein indicates —SA<sub>103</sub> (wherein A<sub>103</sub> is the C<sub>6</sub>-C<sub>60</sub> aryl group).

**[0376]** The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group having two or more rings condensed with each other, only carbon atoms as ring-forming atoms (for example, having 8 to 60 carbon atoms), and no aromaticity in its entire molecular structure. An example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

**[0377]** The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group having two or more rings condensed to each other, at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms (for example, having 1 to 60 carbon atoms), as a ring-forming atom, and no aromaticity in its entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

**[0378]** The term “C<sub>5</sub>-C<sub>60</sub> carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be a ring, such as benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be a trivalent group or a quadrivalent group.

**[0379]** The term “C<sub>1</sub>-C<sub>60</sub> heterocyclic group” as used herein refers to a group having the same structure as the C<sub>1</sub>-C<sub>60</sub> carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and

S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

**[0380]** At least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group, the substituted C<sub>6</sub>-C<sub>60</sub> arylene group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

**[0381]** deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

**[0382]** a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-Coo heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

**[0383]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

**[0384]** a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a

monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group,  $-\text{Si}(\text{Q}_{21})(\text{Q}_{22})(\text{Q}_{23})$ ,  $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$ ,  $-\text{B}(\text{Q}_{21})(\text{Q}_{22})$ ,  $-\text{C}(=\text{O})(\text{Q}_{21})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{21})$ , and  $-\text{P}(=\text{O})(\text{Q}_{21})(\text{Q}_{22})$ ; and

[0385]  $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{C}(=\text{O})(\text{Q}_{31})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$ , and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ ; and

[0386]  $\text{Q}_{11}$  to  $\text{Q}_{13}$ ,  $\text{Q}_{21}$  to  $\text{Q}_{23}$ , and  $\text{Q}_{31}$  to  $\text{Q}_{33}$  may each independently be selected from hydrogen, deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, a  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkyl group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkenyl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0387] The term "Ph" or "ph" used herein refers to a phenyl group, the term "Me" used herein refers to a methyl group, the term "Et" used herein refers to an ethyl group, the term "ter-Bu" or "Bu" used herein refers to a tert-butyl group, and the term "OMe" used herein refers to a methoxy group.

[0388] The term "biphenyl group" as used herein refers to "a phenyl group substituted with a phenyl group." In other words, the "biphenyl group" is a substituted phenyl group having a  $\text{C}_6$ - $\text{C}_{60}$  aryl group as a substituent.

[0389] The term "terphenyl group" as used herein refers to "a phenyl group substituted with a biphenyl group." In other words, the "terphenyl group" is a phenyl group having, as a substituent, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group substituted with a  $\text{C}_6$ - $\text{C}_{60}$  aryl group.

[0390] \* and \*<sup>1</sup> used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.






[0391] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording "B was used instead of A" used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.

[0392] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

#### SYNTHESIS EXAMPLES

##### Synthesis Example 1: Synthesis of Compound 1

[0393]



##### 1) Synthesis of Compound 1-1

[0394] 4-bromobenzaldehyde (1 eq.),  $\text{Pd}(\text{OAc})_2$  (0.05 eq.),  $\text{P}(\text{t-Bu})_3$  (0.5 eq.), and  $\text{NaOt-Bu}$  (1 eq.) were dissolved in a toluene solvent in a nitrogen atmosphere, and 9,9-dimethyl-9,10-dihydroacridine (1 eq.) was added thereto. Then, the reaction mixture was stirred under reflux for 12

hours. After the reaction mixture was cooled to ambient temperature, distilled water was slowly added thereto, and an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography to obtain Compound 1-1 (yield: 87%).

## 2) Synthesis of Compound 1-2

[0395]  $\text{Cp}_2\text{TiCl}_2$  (1.2 eq.) was dissolved in a tetrahydrofuran (THF) solvent under a nitrogen atmosphere at a temperature of  $-78^\circ\text{C}$ . and  $\text{sec-BuMgCl}$  (1.2 eq.) was added to the mixture. The mixture was stirred at a temperature of  $-78^\circ\text{C}$ . for 2 hours, slowly heated to ambient temperature, and stirred for 30 minutes. The mixture was cooled to a temperature of  $-78^\circ\text{C}$ . and Compound 1-1 (1 eq.) was added thereto and stirred for 3 hours. After the reaction was completed, distilled water was slowly added to the mixture, and a solid obtained therefrom was filtered and washed using an ethyl ether solvent. Then, an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 1-2 (yield: 66%).

## 3) Synthesis of Compound 1-3






[0396] Compound 1-2 (1 eq.) and N-bromosuccinimide (2.2 eq.) were dissolved in a  $\text{CCl}_4$  solvent and the mixture was stirred under reflux for 5 hours. A solid obtained therefrom was filtered and washed by using an ethyl ether solvent. Then, an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 1-3 (yield: 42%).

## 4) Synthesis of Compound 1

[0397] Compound 1-3 (1 eq.) and sulfamide (5 eq.) were dissolved in anhydrous ethanol, and HCl gas was added thereto while stirring the mixture under reflux for 8 hours. After the reaction was completed, distilled water was slowly added to the mixture, and an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 1 (yield: 31%).

## Synthesis Example 2: Synthesis of Compound 5

### [0398]



5

### 1) Synthesis of Compound 5-1

[0399] 5-bromothiophene-2-carbaldehyde (1 eq.) and  $\text{Pd}(\text{OAc})_2$  (0.05 eq.),  $\text{P}(\text{t-Bu})_3$  (0.5 eq.), and  $\text{NaOt-Bu}$  (1 eq.) were dissolved in a toluene solvent under a nitrogen atmosphere, and 9,9-dimethyl-9,10-dihydroacridine (1 eq.) was added thereto. Then, the reaction mixture was stirred under reflux for 12 hours. The reaction mixture was cooled to ambient temperature, and distilled water was added thereto. Then, an organic layer was extracted therefrom by using

dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography to obtain Compound 5-1 (yield: 72%).

## 2) Synthesis of Compound 5-2

[0400]  $\text{Cp}_2\text{TiCl}_2$  (1.2 eq.) was dissolved in a THF solvent at a temperature of  $-78^\circ\text{C}$ . under a nitrogen atmosphere, and  $\text{sec-BuMgCl}$  (1.2 eq.) was added to the mixture. The mixture was stirred at a temperature of  $-78^\circ\text{C}$ . for 2 hours, slowly heated to ambient temperature, and stirred for 30 minutes. The mixture was cooled to a temperature of  $-78^\circ\text{C}$ . and Compound 5-1 (1 eq.) was added thereto and stirred for 3 hours. After the reaction was completed, distilled water was slowly added to the mixture, and a solid obtained therefrom was filtered and washed by using an ethyl ether solvent. Then, an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. The result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 5-2 (yield: 53%).

## 3) Synthesis of Compound 5-3



[0401] Compound 5-2 (1 eq.) and N-bromosuccinimide (2.2 eq.) were dissolved in a  $\text{CCl}_4$  solvent, and the mixture was stirred under reflux for 5 hours. A solid obtained therefrom was filtered and washed by using an ethyl ether solvent. Then, an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 5-3 (yield: 40%).

## 4) Synthesis of Compound 5

[0402] Compound 5-3 (1 eq.) and sulfamide (5 eq.) were dissolved in anhydrous ethanol, and HCl gas was added thereto while stirring the mixture under reflux for 8 hours. After the reaction was completed, distilled water was slowly added to the mixture, and an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 5 (yield: 33%).

## Synthesis Example 3: Synthesis of Compound 16

### [0403]



## 1) Synthesis of Compound 16-1

[0404]  $\text{Cp}_2\text{TiCl}_2$  (1.2 eq.) was dissolved in a THF solvent under a nitrogen atmosphere at a temperature of  $-78^\circ\text{C}$ . and  $\text{sec-BuMgCl}$  (1.2 eq.) was added to the mixture. The mixture was stirred at a temperature of  $-78^\circ\text{C}$ . for 2 hours, slowly heated to ambient temperature, and stirred for 30 minutes. The mixture was cooled to a temperature of  $-78^\circ\text{C}$ . and benzaldehyde (1 eq.) was added thereto and stirred for 3 hours. After the reaction was completed, distilled water was slowly added to the mixture, and a solid obtained therefrom was filtered and washed by using an ethyl ether solvent. Then, an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the

result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 16-1 (yield: 80%).

#### 2) Synthesis of Compound 16-2

[0405] Compound 16-1 (1 eq.) and N-bromosuccinimide (2.2 eq.) were dissolved in a  $\text{CCl}_4$  solvent and the mixture was stirred under reflux for 5 hours. A solid obtained therefrom was filtered and washed by using an ethyl ether solvent. Then, an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 16-2 (yield: 49%).

#### 3) Synthesis of Compound 16-3

[0406] Compound 16-2 (1 eq.) and sulfamide (5 eq.) were dissolved in anhydrous ethanol, and HCl gas was added thereto while stirring the mixture under reflux for 8 hours. After the reaction was completed, distilled water was slowly added to the mixture, and an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 16-3 (yield: 49%).

#### 4) Synthesis of Compound 16-4

[0407] Compound 16-3 was dissolved in anhydrous dichloromethane at a temperature of  $0^\circ\text{C}$ . under a nitrogen atmosphere, and anhydrous aluminium chloride (4 eq.) was added thereto and stirred for 3 hours at ambient temperature. Ice water was added to the mixture, and washing was performed thereon by using dichloromethane. Then, a solid obtained therefrom was filtered and dried (yield: 47%).

#### 5) Synthesis of Compound 16-5

[0408] Compound 16-4 (1 eq.) was dissolved in a THF solvent under a nitrogen atmosphere. N-bromosuccinimide (2.2 eq.) dissolved in a small amount of THF was slowly added to the reaction mixture at a temperature of  $0^\circ\text{C}$ . without light and stirred at ambient temperature for 8 hours. After the reaction was completed, distilled water was slowly added thereto, and an organic layer was extracted therefrom by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 16-5 (yield: 72%).

#### 6) Synthesis of Compound 16-6

[0409] Compound 16-5 (1 eq.),  $\text{Pd}(\text{Ph}_3)_4$  (0.05 eq.), and  $\text{K}_2\text{CO}_3$  (1.2 eq.) were dissolved in toluene under a nitrogen atmosphere, and (4-bromophenyl)boronic acid (2.3 eq.) dissolved in a small amount of toluene was added thereto. The reaction mixture was stirred under reflux for 15 hours. After the reaction was completed, the reaction mixture was cooled to ambient temperature, and distilled water was slowly added thereto. Then, an organic layer was extracted there-


from by using dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 16-6 (yield: 46%).

#### 7) Synthesis of Compound 16

[0410] Compound 16-6 (1 eq.), 9,9-dimethyl-9,10-dihydroacridine (2 eq.),  $\text{Pd}(\text{OAc})_2$  (0.05 eq.),  $\text{P}(\text{t-Bu})_3$  (50 wt %, 0.05 eq.), and NaOt-Bu (sodium tert-butoxide, 2 eq.) were stirred with a toluene solvent under a nitrogen atmosphere. Then, the mixture was stirred under reflux for 12 hours at a temperature of  $120^\circ\text{C}$ . After the reaction was completed, the mixture was cooled to ambient temperature, and an organic layer was extracted therefrom by using water and dichloromethane. The extracted organic layer was dried by using  $\text{MgSO}_4$  and concentrated under vacuum. Then, the result obtained therefrom was purified by column chromatography using ethyl acetate/hexane to obtain Compound 16 (yield: 59%).

#### Synthesis Example 4: Synthesis of Compound 17

##### [0411]



## 1) Synthesis of Compound 17-1

[0412] Compound 17-1 (yield: 52%) was synthesized in the same manner as in Synthesis of Compound 16-6, except that (5-bromothiophen-2-yl)boronic acid was used instead of (4-bromophenyl)boronic acid.

## 2) Synthesis of Compound 17

[0413] Compound 17 (yield: 50%) was synthesized in the same manner as in Synthesis of Compound 16, except that Compound 17-1 was used instead of Compound 16-6.

[0414] <sup>1</sup>H NMR and MS/FAB of Compounds synthesized in Synthesis Examples 1 to 4 are shown in Table 1.

[0415] Synthesis methods of compounds other than Compounds shown in Table 1 may also be recognized by those of ordinary skill in the art by referring to the synthesis mechanisms and source materials described above.

TABLE 1

Compound	<sup>1</sup> H NMR (CDCl <sub>3</sub> , 400 MHz)	MS/FAB	
		found	calc.
1	δ = 7.44(d, 4H), 7.26(d, 4H), 7.07(d, 4H), 7.14(t, 4H), 7.11(d, 4H) 6.89(t, 4H), 1.69(s, 12H)	684.26	684.86
5	δ = 7.14(t, 4H), 7.11(d, 4H), 7.07(d, 4H), 6.89(t, 4H), 6.70(d, 2H) 5.98(d, 2H), 1.69(s, 12H)	696.17	696.90
16	δ = 8.17(s, 2H), 8.04(d, 4H), 7.14(t, 4H), 7.11(d, 4H), 7.07(d, 4H), 6.85(d, 2H) 5.98(d, 2H), 1.69(s, 12H)	834.30	835.04
17	δ = 8.20(s, 2H), 8.14(m, 4H), 7.55(d, 4H), 7.27(d, 4H), 7.14(t, 4H), 7.11(d, 4H), 6.89(t, 4H), 1.69(s, 12H)	846.22	847.08

## EXAMPLES

## Example 1

[0416] An ITO film (anode) having a thickness of 120 nm was formed on a glass substrate, and ultrasonic cleaning and pre-treatment (UV—O<sub>3</sub> treatment, thermal treatment) were performed thereon.



[0417] Compound HT3 and HAT-CN were co-deposited on the anode at a weight ratio of 99:1 to form a hole transport layer having a thickness of 120 nm.

[0418] Then, a host (Compound H-1a) and a dopant (Compound 1, 1 wt %) were co-deposited on the hole transport layer to form an emission layer having a thickness of 30 nm.


[0419] Then, ET1 was deposited on the emission layer to form a buffer layer having a thickness of 5 nm, ET2 was deposited on the buffer layer to form an electron transport layer having a thickness of 25 nm, LiF was deposited on the electron transport layer to form an electron injection layer

having a thickness of 0.5 nm, and A1 was vacuum-deposited on the electron injection layer to form a cathode having a thickness of 150 nm, thereby completing the manufacture of an organic light-emitting device.


HT3



H-1a



-continued




1

[0422] Referring to Table 2, it may be seen that the organic light-emitting devices of Examples 1 to 4 had high external quantum efficiency, as compared with the organic light-emitting device of Comparative Example 1 (that employed Pt phthalocyanine (PtPc) as a dopant compound), and may enable a long wavelength shift of a maximum emission wavelength, and thus may be suitable for NIR emission.


[0423] Organic light-emitting devices including the heterocyclic compound according to an embodiment may have a low driving voltage, high efficiency, high luminance, and a long lifespan, and high-quality organic light-emitting devices and organic light-emitting apparatuses may be implemented.

[0424] The embodiments may provide a heterocyclic compound and an organic light-emitting device including the same.



ET1

[0425] Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.



ET2

What is claimed is:

1. An organic light-emitting device, comprising: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode and including an emission layer, wherein the emission layer includes a compound represented by Formula 1A or a compound represented by Formula 1B:

Examples 2 to 4 and Comparative Example 1

[0420] Organic light-emitting devices were manufactured in the same manner as in Example 1, except that Compounds shown in Table 2 were each used instead of Compound 1 as a dopant in forming the emission layer.

Evaluation Example 1

[0421] The driving voltage, current density, luminance, external quantum efficiency (EQE), and maximum emission wavelength of the organic light-emitting devices manufactured according to Examples 1 to 4 and Comparative Example 1 were measured by using Keithley SMU 236 and a luminance meter PR745, and results thereof are shown in Table 2.

<Formula 1A>





TABLE 2

	Dopant compound	HOMO (eV)	LUMO (eV)	S <sub>1</sub> (eV)	ΔE <sub>ST</sub> (eV)	Driving voltage (V)	EQE (%)	Maximum emission wavelength (nm)
Example 1	Compound 1	-5.46	-3.54	1.51	0.01	6.7	2.3	819
Example 2	Compound 5	-5.66	-3.51	1.70	0.01	6.5	2.7	730
Example 3	Compound 16	-5.40	-3.79	1.35	0.00	7.1	0.9	917
Example 4	Compound 17	-5.57	-3.75	1.55	0.01	7.1	1.2	801
Comparative Example 1	PtPc	-5.08	-2.74	1.32 (T <sub>1</sub> )	0.86	7.9	0.5	968

-continued

&lt;Formula 1B&gt;



wherein, in Formulae 1A and 1B, ring A<sub>1</sub> is selected from a C<sub>5</sub>-C<sub>60</sub> carbocyclic group and a C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

X<sub>1</sub> is S or Se,

L<sub>1</sub> and L<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group and a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, a<sub>1</sub> and a<sub>2</sub> are each independently an integer from 0 to 5, when a<sub>1</sub> is 2, 3, 4, or 5, the 2, 3, 4, or 5 L<sub>1</sub>(s) are identical to or different from each other, and when a<sub>2</sub> is 2, 3, 4, or 5, the 2, 3, 4, or 5 L<sub>2</sub>(s) are identical to or different from each other,

when a<sub>1</sub> is 0, \*-(L<sub>1</sub>)<sub>a1</sub>-\* is a single bond, and when a<sub>2</sub> is 0, \*-(L<sub>2</sub>)<sub>a2</sub>-\* is a single bond,

R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>60</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryloxy group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

at least one of R<sub>1</sub> and R<sub>2</sub> is an electron donating group, b<sub>1</sub> and b<sub>2</sub> are each independently an integer from 1 to 10, when b<sub>1</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>1</sub>(s) are identical to or different from each other, and when b<sub>2</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>2</sub>(s) are identical to or different from each other,

c<sub>1</sub> and c<sub>2</sub> are each independently an integer from 1 to 10, at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substi-

tuted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryloxy group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

—Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>),  
—C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),

Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted with a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a


monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and and \*<sup>1</sup> each indicate a binding site to a neighboring atom.

2. The organic light-emitting device as claimed in claim 1, wherein:


the emission layer includes a host and a dopant,

the dopant includes the compound represented by Formula 1A or Formula 1B, and


the host includes at least one selected from a compound represented by Formula 11-HT, a compound represented by Formula 21-HT, a compound represented by Formula 31-HT, a compound represented by Formula 41-HT, a compound represented by Formula 51-HT, a compound represented by Formula 61-ET, and a compound represented by Formula 71-ET:




&lt;Formula 11-HT&gt;




&lt;Formula 21-HT&gt;




&lt;Formula 31-HT&gt;




&lt;Formula 41-HT&gt;



&lt;Formula 51-HT&gt;



&lt;Formula 61-HT&gt;



&lt;Formula 71-HT&gt;

wherein, in Formulae 11-HT, 21-HT, 31-HT, 41-HT, and 51-HT,

$X_{21}$  is O, S, N-[( $L_{24}$ )<sub>a24</sub>-R<sub>24</sub>], B-[( $L_{24}$ )<sub>a24</sub>-R<sub>24</sub>], P-[( $L_{24}$ )<sub>a24</sub>-R<sub>24</sub>], C(R<sub>24</sub>)(R<sub>25</sub>), or Si(R<sub>24</sub>)(R<sub>25</sub>),

A<sub>11</sub>, A<sub>12</sub>, and A<sub>21</sub> to A<sub>24</sub> are each independently a C<sub>5</sub>-C<sub>30</sub> carbocyclic group or a  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group,

A<sub>31</sub> is a substituted or unsubstituted C<sub>5</sub>-C<sub>30</sub> carbocyclic group in which two or more rings are condensed with each other or a substituted or unsubstituted  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group in which two or more rings are condensed with each other, wherein A<sub>31</sub> is not a substituted or unsubstituted spirobifluorene group and a substituted or unsubstituted anthracene group,

L<sub>11</sub> to L<sub>13</sub>, L<sub>21</sub> to L<sub>25</sub>, L<sub>31</sub>, L<sub>41</sub> to L<sub>43</sub>, and L<sub>51</sub> to L<sub>55</sub> are each independently selected from:

a single bond, a C<sub>5</sub>-C<sub>30</sub> carbocyclic group, and a  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group; and

a C<sub>5</sub>-C<sub>30</sub> carbocyclic group and a  $\pi$  electron-depleted nitrogen-free C<sub>2</sub>-C<sub>30</sub> heterocyclic group, each substituted with at least one selected from hydrogen, deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>0</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>5</sub>)(Q<sub>52</sub>)(Q<sub>53</sub>), —N(Q<sub>51</sub>)(Q<sub>52</sub>), and —B(Q<sub>5</sub>)(Q<sub>52</sub>),

a111 to a13, a21 to a25, a31, a41 to a43, and a51 to a55 are each independently an integer from 1 to 5,

R<sub>11</sub>, R<sub>12</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>21</sub>, R<sub>22</sub>, and R<sub>24</sub> to R<sub>29</sub> are each independently selected from:

hydrogen, deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>0</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, a

C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>51</sub>)(Q<sub>52</sub>)(Q<sub>53</sub>), —N(Q<sub>51</sub>)(Q<sub>52</sub>), and —B(Q<sub>51</sub>)(Q<sub>52</sub>); and —Si(Q<sub>41</sub>)(Q<sub>42</sub>)(Q<sub>43</sub>), —N(Q<sub>41</sub>)(Q<sub>42</sub>), and —B(Q<sub>41</sub>)(Q<sub>42</sub>),

R<sub>13</sub>, R<sub>23</sub>, R<sub>31</sub>, R<sub>41</sub> to R<sub>43</sub>, and R<sub>51</sub> to R<sub>54</sub> are each independently selected from:

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>51</sub>)(Q<sub>52</sub>)(Q<sub>53</sub>), —N(Q<sub>51</sub>)(Q<sub>52</sub>), and —B(Q<sub>51</sub>)(Q<sub>52</sub>); and —Si(Q<sub>41</sub>)(Q<sub>42</sub>)(Q<sub>43</sub>), —N(Q<sub>41</sub>)(Q<sub>42</sub>), and —B(Q<sub>41</sub>)(Q<sub>42</sub>),

R<sub>51</sub> and R<sub>52</sub> are separate or are linked to form a saturated or unsaturated ring, and R<sub>53</sub> and R<sub>54</sub> are separate or are linked to form a saturated or unsaturated ring,

b31 is an integer from 1 to 10,

Q<sub>41</sub> to Q<sub>43</sub> and Q<sub>51</sub> to Q<sub>53</sub> are each independently selected from hydrogen, deuterium, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a  $\pi$  electron-depleted nitrogen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a  $\pi$  electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group,

- gen-free C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a π electron-depleted nitrogen-free monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group,
- in Formulae 61-ET and 71-ET,
- X<sub>71</sub> is O, S, N-[(L<sub>74</sub>)<sub>a74</sub>-R<sub>74</sub>], B-[(L<sub>74</sub>)<sub>a74</sub>-R<sub>74</sub>], P-[(L<sub>74</sub>)<sub>a74</sub>-R<sub>74</sub>], C(R<sub>74</sub>)(R<sub>75</sub>), or Si(R<sub>74</sub>)(R<sub>75</sub>),
- A<sub>61</sub>, A<sub>62</sub>, and A<sub>71</sub> to A<sub>74</sub> are each independently a C<sub>5</sub>-C<sub>30</sub> carbocyclic group or a C<sub>2</sub>-C<sub>30</sub> heterocyclic group,
- L<sub>61</sub> to L<sub>63</sub> and L<sub>71</sub> to L<sub>75</sub> are each independently selected from a single bond, a substituted or unsubstituted C<sub>5</sub>-C<sub>30</sub> carbocyclic group, and a substituted or unsubstituted C<sub>2</sub>-C<sub>30</sub> heterocyclic group,
- a<sub>61</sub> to a<sub>63</sub> and a<sub>71</sub> to a<sub>75</sub> are each independently an integer from 1 to 5,
- R<sub>61</sub> to R<sub>65</sub> and R<sub>71</sub> to R<sub>79</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)<sub>2</sub>(Q<sub>1</sub>), —S(=O)<sub>2</sub>(Q<sub>1</sub>), and —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>),
- in Formula 61-ET, i) at least one of A<sub>61</sub> and A<sub>62</sub> is a π electron-depleted nitrogen-containing C<sub>2</sub>-C<sub>30</sub> heterocyclic group, or ii) at least one of R<sub>61</sub> to R<sub>65</sub> is a substituted or unsubstituted π electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted π electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>60</sub> heteroaryl group, and a substituted or unsubstituted π electron-depleted nitrogen-containing monovalent non-aromatic condensed heteropolycyclic group,
- in Formula 71-ET, i) at least one of A<sub>71</sub> to A<sub>74</sub> is a π electron-depleted nitrogen-containing C<sub>2</sub>-C<sub>30</sub> heterocyclic group, or ii) at least one of R<sub>71</sub> to R<sub>73</sub> and R<sub>76</sub> to R<sub>79</sub> is a substituted or unsubstituted π electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted π electron-depleted nitrogen-containing C<sub>1</sub>-C<sub>60</sub> heteroaryl group, and a substituted or unsubstituted π electron-depleted nitrogen-containing monovalent non-aromatic condensed heteropolycyclic group,
- at least one substituent of the substituted C<sub>5</sub>-C<sub>30</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>30</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:
- deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;
- a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)<sub>2</sub>(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);
- a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;
- a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)<sub>2</sub>(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and
- Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)<sub>2</sub>(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>), and
- Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a

hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted with a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group substituted with a C<sub>6</sub>-C<sub>60</sub> aryl group, a terphenyl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group substituted with a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group substituted with a C<sub>6</sub>-C<sub>60</sub> aryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group.


3. The organic light-emitting device as claimed in claim 2, wherein:

A<sub>11</sub>, A<sub>12</sub>, and A<sub>21</sub> to A<sub>24</sub> are each independently selected from a benzene group, a naphthalene group, a pyrrole group, a cyclopentene group, a furan group, a thiophene group, an indole group, an indene group, a benzofuran group, a benzothiophene group, a carbazole group, a fluorene group, a dibenzofuran group, and a dibenzothiophene group,


A<sub>31</sub> is a group of one of the following Formulae A<sub>31</sub>-1 to A<sub>31</sub>-38, and

A<sub>61</sub>, A<sub>62</sub>, and A<sub>71</sub> to A<sub>74</sub> are each independently selected from a benzene group, a naphthalene group, a pyrrole group, a cyclopentene group, a furan group, a thiophene group, an indole group, an indene group, a benzofuran group, a benzothiophene group, a carbazole group, a fluorene group, a dibenzofuran group, a dibenzothiophene group, a pyridine group, a pyrimidine group, a pyrazine group, a triazine group, a pyrazole group, a pyridazine group, a triazole group, a quinoline group, an isoquinoline group, a benzoquinoline group, a benzoisoquinoline group, a quinoxaline group, and a benzoquinoxaline group,


-continued




A31-1




A31-2




A31-3




A31-4




A31-5




A31-6




A31-7




A31-8



A31-9




A31-10




A31-11


-continued




A31-12




A31-13




A31-14



A31-15




A31-16




A31-17


-continued




A31-18




A31-19




A31-20




A31-21



A31-22




A31-23




A31-24


-continued




A31-25




A31-26




A31-27




A31-28



A31-29




A31-30




A31-31


-continued




A31-32




A31-33




A31-34




A31-35



A31-36



A31-37



A31-38

4. The organic light-emitting device as claimed in claim 1, wherein the compound included in the emission layer and represented by Formula 1A or Formula 1B is a near infrared ray (NIR)-emitting compound having a maximum emission wavelength of about 700 nm or more.

5. The organic light-emitting device as claimed in claim 1, wherein:

the compound included in the emission layer and represented by Formula 1A or Formula 1B is a thermally activated delayed fluorescent (TADF) emitter, and the emission layer emits delayed fluorescence.

6. The organic light-emitting device as claimed in claim 1, wherein the organic layer further includes:

a hole transport region between the first electrode and the emission layer; and

an electronic transport region between the emission layer and the second electrode.

7. The organic light-emitting device as claimed in claim 6, wherein:

the hole transport region includes a hole injection layer, a hole transport layer, a buffer layer, an electron blocking layer, or any combination thereof, and

the electron transport region comprises a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

8. The organic light-emitting device as claimed in claim 6, wherein:

the hole transport region includes a p-dopant, and the p-dopant has a lowest unoccupied molecular orbital (LUMO) level of about -3.5 eV or less.



9. The organic light-emitting device as claimed in claim 8, wherein the p-dopant includes a cyano group-containing compound.

10. The organic light-emitting device as claimed in claim 6, wherein:

the electron transport region includes a triazole-containing compound or a benzotriazole-containing compound, and

the electron transport region further includes an alkali metal, alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth metal compound, a rare earth metal compound, an alkali metal complex, alkaline earth metal complex, a rare earth metal complex, or a combination thereof.

11. A heterocyclic compound represented by Formula 1A or Formula 1B:



wherein, in Formulae 1A and 1B,

ring A<sub>1</sub> is selected from a C<sub>5</sub>-C<sub>60</sub> carbocyclic group and a C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

X<sub>1</sub> is S or Se,

L<sub>1</sub> and L<sub>2</sub> are each independently selected from a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group and a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, a<sub>1</sub> and a<sub>2</sub> are each independently an integer from 0 to 5, when a<sub>1</sub> is 2, 3, 4, or 5, the 2, 3, 4, or 5 L<sub>1</sub>(s) are identical to or different from each other, and when a<sub>2</sub> is 2, 3, 4, or 5, the 2, 3, 4, or 5 L<sub>2</sub>(s) are identical to or different from each other,

when a<sub>1</sub> is 0, \*-(L<sub>1</sub>)<sub>a1</sub>-\* is a single bond, and when a<sub>2</sub> is 0, \*-(L<sub>2</sub>)<sub>a2</sub>-\* is a single bond,

R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryloxy group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —N(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

at least one of R<sub>1</sub> and R<sub>2</sub> is an electron donating group, in Formula 1A, (R<sub>1</sub>)<sub>b1</sub>-(L<sub>1</sub>)<sub>a1</sub>-\* and (R<sub>2</sub>)<sub>b2</sub>-(L<sub>2</sub>)<sub>a2</sub>-\* are not halogen, a phenoxy group, a phenylthio group, an alkylthio group, or an alkoxy group,

b<sub>1</sub> and b<sub>2</sub> are each independently an integer from 1 to 10, when b<sub>1</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>1</sub>(s) are identical to or different from each other, and when b<sub>2</sub> is 2, 3, 4, 5, 6, 7, 8, 9, or 10, the 2, 3, 4, 5, 6, 7, 8, 9, or 10 R<sub>2</sub>(s) are identical to or different from each other,

c<sub>1</sub> and c<sub>2</sub> are each independently an integer from 1 to 10, at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryloxy group, the substituted monovalent non-aromatic con-

densed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —N(Q<sub>11</sub>)(Q<sub>12</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —N(Q<sub>21</sub>)(Q<sub>22</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

—Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),





Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>6</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a

C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and and \*<sup>1</sup> each indicate a binding site to a neighboring atom.








12. The heterocyclic compound as claimed in claim 11, wherein ring A<sub>1</sub> is selected from a cyclohexadiene group, a dihydronaphthalene group, a dihydroanthracene group, a dihydrobenzoanthracene group, a dihydrophenanthrene group, a dihydrobenzophenanthrene group, a dihydrotriphenylene group, a dihydronaphthacene group, a dihydropyrene group, a dihydrochrysene group, a dihydropicene group, and a dihydrobenzopentaphene group.

13. The heterocyclic compound as claimed in claim 11, wherein:

the compound is represented by Formula 1B, and ring A<sub>1</sub> is a group of one of the following Formulae A<sub>1-1</sub> to A<sub>1-12</sub>:

A<sub>1-1</sub>A<sub>1-2</sub>A<sub>1-3</sub>A<sub>1-4</sub>A<sub>1-5</sub>

-continued

A<sub>1-6</sub>A<sub>1-7</sub>A<sub>1-8</sub>A<sub>1-9</sub>A<sub>1-10</sub>A<sub>1-11</sub>A<sub>1-12</sub>

fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylylene group, a picenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a pyrrolylylene group, a thiophenylylene group, a furanylylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylylene group, a cinolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylylene group, a benzothiophenylylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylene group, and a dibenzocarbazolylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylylene group, a heptalenylene group, an indacenylylene group, an acenaphthylene group, a fluorenylylene group, a spiro-fluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylylene group, a picenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a pyrrolylylene group, a thiophenylylene group, a furanylylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isooxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylylene group, a cinolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylylene group, a benzothiophenylylene group, an isobenzothiazolylene group, a benzooxazolylene group, an isobenzooxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylene group, and a dibenzocarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a

14. The heterocyclic compound as claimed in claim 11, wherein L<sub>1</sub> and L<sub>2</sub> are each independently selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylylene group, a heptalenylene group, an indacenylylene group, an acenaphthylene group, a fluorenylylene group, a spiro-fluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylylene group, a

sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-fluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a thiadiazolyl group, and an imidazopyridinyl group.

15. The heterocyclic compound as claimed in claim 11, wherein:

i) the heterocyclic compound is represented by Formula 1A,

a1 and a2 are each independently 0, 1, or 2, and

at least one of a1 and a2 in Formula 1A is not 0; or

ii) the heterocyclic compound is represented by Formula 1B, and

a1 and a2 are each independently 0, 1, or 2.

16. The heterocyclic compound as claimed in claim 11, wherein:

R<sub>1</sub> and R<sub>2</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>2</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group;

a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD<sub>3</sub>, —CD<sub>2</sub>H, —CDH<sub>2</sub>, —CF<sub>3</sub>, —CF<sub>2</sub>H, —CFH<sub>2</sub>, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>);

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, —CD<sub>3</sub>, —CD<sub>2</sub>H, —CDH<sub>2</sub>, —CF<sub>3</sub>, —CF<sub>2</sub>H, —CFH<sub>2</sub>, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridi-

nyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenanthroliny group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>); and

—N(Q<sub>1</sub>)(Q<sub>2</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —P(Q<sub>1</sub>)(Q<sub>2</sub>), and —C(=O)(Q<sub>1</sub>),

at least one of R<sub>1</sub> and R<sub>2</sub> is an electron donating group, and

Q<sub>1</sub> to Q<sub>3</sub> and Q<sub>31</sub> to Q<sub>33</sub> are defined the same as those of Formulae 1A and 1B.

17. The heterocyclic compound as claimed in claim 11, wherein:


R<sub>1</sub> and R<sub>2</sub> are each independently selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>2</sub>-C<sub>20</sub> alkenyl group, a C<sub>2</sub>-C<sub>20</sub> alkynyl group, and a C<sub>1</sub>-C<sub>20</sub> alkoxy group; and


a group represented by one of the following Formulae 5-1 to 5-6, and

at least one of R<sub>1</sub> and R<sub>2</sub> is an electron donating group:


<Formula 5-1>



<Formula 5-2>




<Formula 5-3>




-continued


<Formula 5-4>



<Formula 5-5>



<Formula 5-6>



wherein, in Formulae 5-1 to 5-6,

K<sub>1</sub> is selected from \*—C(R<sub>20</sub>)(R<sub>30</sub>)—\*, \*—N(R<sub>20</sub>)—\*, and \*—Si(R<sub>20</sub>)(R<sub>30</sub>)—\*,

R<sub>20</sub>, R<sub>30</sub>, R<sub>40</sub>, and R<sub>50</sub> are defined the same as R<sub>1</sub> and R<sub>2</sub> of Formulae 1A and 1B,

b<sub>40</sub> and b<sub>50</sub> are each independently an integer from 1 to 4, and


indicates a binding site to a neighboring atom.

18. The heterocyclic compound as claimed in claim 11, wherein


the compound is represented by Formula 1B, and

the compound represented by Formula 1B is represented by one of Formulae 1B-1 to 1B-4:

<Formula 1B-1>




<Formula 1B-2>




-continued

<Formula 1B-3>




<Formula 1B-4>




-continued


<Formula 1B-b>



<Formula 1B-c>



<Formula 1B-d>




wherein, in Formulae 1B-1 to 1B-4,  
 $X_1$  is defined the same as that of Formula 1B,  
 $L_{11}$  and  $L_{12}$  are defined the same as  $L_1$  and  $L_2$  of Formula 1B,  
 $a_{11}$  and  $a_{12}$  are defined the same as  $a_1$  and  $a_2$  of Formula 1B, and  
 $R_{11}$  and  $R_{12}$  are defined the same as  $R_1$  and  $R_2$  of Formula 1B.

19. The heterocyclic compound as claimed in claim 11, wherein:

the compound is represented by Formula 1B, and  
 the compound represented by Formula 1B is represented by one of Formulae 1B-a to 1B-d:

<Formula 1B-a>



wherein, in Formulae 1B- to 1B-d,


$L_{11}$  and  $L_{12}$  are defined the same as  $L_1$  and  $L_2$  of Formula 1B,

$a_{11}$  and  $a_{12}$  are defined the same as  $a_1$  and  $a_2$  of Formula 1B, and


$R_{11}$  and  $R_{12}$  are defined the same as  $R_1$  and  $R_2$  of Formula B.

20. The heterocyclic compound as claimed in claim 11, wherein the compound represented by Formula 1A or Formula 1B is one of the following Compounds 1 to 37:


1




2




-continued




3




4




5




6



7




8




9

-continued


10




11




12




13




14



15





16



-continued



17

18




19


20



21





22




23

24




-continued


25




26




27




28




29




30



31




32





-continued

33




34


35



36



37



\* \* \* \* \*

专利名称(译)	杂环化合物和包括其的有机发光装置		
公开(公告)号	<a href="#">US20190267553A1</a>	公开(公告)日	2019-08-29
申请号	US16/181407	申请日	2018-11-06
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	YE JIMYOUNG KIM MYEONGSUK YOO BYEONGWOOK KO SOOBYUNG YOON JIHWAN HWANG JAEHOON		
发明人	YE, JIMYOUNG KIM, MYEONGSUK YOO, BYEONGWOOK KO, SOOBYUNG YOON, JIHWAN HWANG, JAEHOON		
IPC分类号	H01L51/00 C07D417/14 C09K11/06		
CPC分类号	H01L51/0072 C07D417/14 H01L51/0069 H01L51/0068 C09K11/06 H01L51/5016 C09K2211/1018 H01L51/5072 H01L51/5096 H01L51/5088 H01L51/5092 H01L2251/552 H01L51/5056 H01L51/0071 H01L51/5012		
优先权	1020180024720 2018-02-28 KR		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

有机发光器件和杂环化合物，该器件包括第一电极;面向第一电极的第二电极;第一电极和第二电极之间的有机层，包括发光层，其中发光层包括由式1A表示的化合物或由式1B表示的化合物：

**10**

<b>190</b>
<b>150</b>
<b>110</b>