




US 20170179407A1

(19) **United States**(12) **Patent Application Publication**
PARK et al.(10) **Pub. No.: US 2017/0179407 A1**(43) **Pub. Date: Jun. 22, 2017**(54) **CONDENSED CYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**(52) **U.S. Cl.**
CPC **H01L 51/0072** (2013.01); **C07D 221/18**
(2013.01); **H01L 51/0052** (2013.01); **C07D**
405/10 (2013.01); **H01L 51/0073** (2013.01);
C07F 9/5765 (2013.01); **H01L 51/0058**
(2013.01); **C07D 401/10** (2013.01); **C07D**
471/04 (2013.01); **H01L 51/0067** (2013.01);
H01L 51/5012 (2013.01)(71) Applicant: **Samsung Display Co., Ltd**, Yongin-si
(KR)(72) Inventors: **Junha PARK**, Yongin-si (KR);
Youngkook KIM, Yongin-si (KR);
Munki SIM, Yongin-si (KR);
Eunyoung LEE, Yongin-si (KR);
Hyoyoung LEE, Yongin-si (KR);
Eunjae JEONG, Yongin-si (KR);
Seokhwan HWANG, Yongin-si (KR)(57) **ABSTRACT**Provided are a condensed cyclic compound and an organic
light-emitting device. The condensed cyclic compound is
represented by Formula 1:(21) Appl. No.: **15/371,066**(22) Filed: **Dec. 6, 2016**

<Formula 1>

(30) **Foreign Application Priority Data**Dec. 21, 2015 (KR) 10-2015-0182792
Apr. 15, 2016 (KR) 10-2016-0046499**Publication Classification**(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 471/04 (2006.01)
C07F 9/576 (2006.01)
C07D 401/10 (2006.01)
C07D 221/18 (2006.01)
C07D 405/10 (2006.01)

Details about the constituents of Formula 1 is disclosed.

10**190****150****110**

FIG. 1

10

190
150
110

FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

CONDENSED CYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of Korean Patent Application Nos. 10-2015-0182792 filed on Dec. 21, 2015, and 10-2016-0046499 filed on Apr. 15, 2016 in the Korean Intellectual Property Office, the disclosure of which are incorporated herein in their entirety by reference.

BACKGROUND

1. Field

[0002] One or more embodiments relate to a condensed cyclic compound and an organic light-emitting device including the same.

SUMMARY

[0003] Organic light-emitting devices are self-emission devices that offer advantages such as wide viewing angles, high contrast ratios, short response times, and excellent luminance, driving voltage, and response speed characteristics, and produce full-color images.


[0004] The organic light-emitting device may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. Then, the excitons are transitioned from an excited state to a ground state, thereby generating light.

SUMMARY

[0005] One or more embodiments include a novel condensed cyclic compound and an organic light-emitting device including the same.

[0006] Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

[0007] According to one or more embodiments, a condensed cyclic compound is represented by Formula 1:







[0008] In Formula 1,

[0009] ring A may be selected from a C₅-C₃₀ carbocyclic ring and a C₂-C₃₀ heterocyclic ring,

[0010] X₁ may be C(R₁) or N, X₂ may be C(R₂) or N, at least one of X₁ and X₂ may be N,

[0011] R₁ to R₁₁ may each independently be a group represented by one of Formulae 2-1 to 2-5 group, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, substituted or unsubstituted C₁-C₆₀ alkyl group, substituted or unsubstituted C₂-C₆₀ alkenyl group, substituted or unsubstituted C₂-C₆₀ alkynyl group, substituted or unsubstituted C₁-C₆₀ alkoxy group, substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₆-C₆₀ aryloxy group, substituted or unsubstituted C₆-C₆₀ arylthio group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂):



[0012] L₁ to L₁₁ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, substituted or unsubstituted C₆-C₆₀ arylene group, substituted or unsubstituted C₁-C₆₀ heteroarylene group, substituted or unsubstituted divalent non-aromatic

condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0013] a1 to a11 may each independently be 0, 1, 2, 3, 4, or 5,

[0014] Ar₁ to Ar₇ may be a substituted or unsubstituted C₁-C₆₀ alkyl group, substituted or unsubstituted C₂-C₆₀ alkenyl group, substituted or unsubstituted C₂-C₆₀ alkynyl group, substituted or unsubstituted C₁-C₆₀ alkoxy group, substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

[0015] b1 to b4 may each independently be 1, 2, or 3,

[0016] c3 may be 0, 1, 2, 3, 4, 5, or 6,

[0017] at least one of substituent or substituents of the substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0018] deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0019] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₁₁)(Q₁₂)(Q₁₃), -N(Q₁₁)(Q₁₂), -B(Q₁₁)(Q₁₂), -C(=O)(Q₁₁), -S(=O)₂(Q₁₁), and -P(=O)(Q₁₁)(Q₁₂);

[0020] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0021] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₂₁)(Q₂₂)(Q₂₃), -N(Q₂₁)(Q₂₂), -B(Q₂₁)(Q₂₂), -C(=O)(Q₂₁), -S(=O)₂(Q₂₁), and -P(=O)(Q₂₁)(Q₂₂); and

[0022] -Si(Q₃₁)(Q₃₂)(Q₃₃), -N(Q₃₁)(Q₃₂), -B(Q₃₁)(Q₃₂), -C(=O)(Q₃₁), -S(=O)₂(Q₃₁) and -P(=O)(Q₃₁)(Q₃₂),

[0023] wherein Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ each independently hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

[0024] * indicates a binding site to a neighboring atom.

[0025] According to another aspect, an organic light-emitting device includes a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes at least one of condensed cyclic compounds represented by Formula 1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0026] These and/or other aspects will become apparent and more readily appreciated from the following description of the embodiments, taken in conjunction with the accompanying drawings in which:

[0027] FIG. 1 is a schematic cross-sectional view of an organic light-emitting device according to an embodiment;

[0028] FIG. 2 is a schematic cross-sectional view of an organic light-emitting device according to an embodiment;

[0029] FIG. 3 is a schematic cross-sectional view of an organic light-emitting device according to an embodiment; and


[0030] FIG. 4 is a schematic cross-sectional view of an organic light-emitting device according to an embodiment;

DETAILED DESCRIPTION

[0031] Reference will now be made in detail to embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like

elements throughout. In this regard, the present embodiments may have different forms and should not be construed as being limited to the descriptions set forth herein. Accordingly, the embodiments are merely described below, by referring to the figures, to explain aspects of the present description. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

[0032] A condensed cyclic compound according to an embodiment of the present disclosure is represented by Formula 1:








[0033] Ring A in Formula 1 may be selected from a C₅-C₃₀ carbocyclic ring and a C₂-C₃₀ heterocyclic ring. For example, ring A may be selected from a benzene group, a pyridine group, a pyridazine group, a pyrimidine group, a pyrazine group, a naphthalene group, a quinoline group, an isoquinoline group, a cinnoline group, a quinazoline group, a quinoxaline group, a naphthyridine group, an anthracene group, a phenanthrene group, a benzoquinoline group, a phenanthridine group, an acridine group, a phenanthroline group, and a phenazine group. In one embodiment, ring A may be selected from a benzene group, a pyridine group, a naphthalene group, a quinoline group, and an isoquinoline group, but embodiments of the present disclosure are not limited thereto.

[0034] In Formula 1, X₁ may be C(R₁) or N, X₂ may be C(R₂) or N, and at least one of X₁ and X₂ may be N.

[0035] R₁ to R₁₁ in Formula 1 may each independently be selected from the group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, substituted or unsubstituted C₁-C₆₀ alkyl group, substituted or unsubstituted C₂-C₆₀ alkenyl group, substituted or unsubstituted C₂-C₆₀ alkynyl group, substituted or unsubstituted C₁-C₆₀ alkoxy group, substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₆-C₆₀ aryloxy group, substituted or unsubstituted C₆-C₆₀ arylthio group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂),

wherein Q₁ to Q₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group:



[0036] * in Formulae 2-1 to 2-5 indicates a binding site to a neighboring atom.

[0037] L₁ to L₁₁ in Formulae 2-1 to 2-5 may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, substituted or unsubstituted C₆-C₆₀ arylene group, substituted or unsubstituted C₁-C₆₀ heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group. For example, L₁ to L₁₁ may each independently be selected from:

[0038] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene


group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazininylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzosilolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group; and

[0039] a phenylene group, a pentalenylenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-bifluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylenylene group, an indolylenylene group, an indazolylenylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazininylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzosilolylenylene group, a dibenzocarbazolylenylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cyclo-


heptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinylyl group, a cinnolinylyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

[0040] wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.


[0041] In one embodiment, L_1 to L_{11} may each independently be selected from groups represented by Formulae 3-1 to 3-25:



Formular 3-1












Formular 3-2



Formular 3-3

-continued



Formular 3-4

Formular 3-5

Formular 3-6

Formular 3-7

Formular 3-8







Formular 3-9

Formular 3-10

Formular 3-11

Formular 3-12

-continued



Formular 3-13

Formular 3-14


Formular 3-15

Formular 3-16


Formular 3-17

Formular 3-18


-continued




Formular 3-19




Formular 3-20




Formular 3-21




Formular 3-22



Formular 3-23



Formular 3-24



Formular 3-25


[0042] In Formulae 3-1 to 3-25,

[0043] Y, may be O, S, C(Z₅)(Z₆), N(Z₇), or Si(Z₅)(Z₉),


[0044] Z₁ to Z₉ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃), wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

[0045] d₃ may be an integer from 1 to 3,[0046] d₄ may be an integer from 1 to 4,[0047] d₆ may be an integer from 1 to 6,[0048] d₈ may be an integer from 1 to 8, and


[0049] * and *¹ each indicate a binding site to a neighboring atom. For example, L₁ to L₁₁ may each independently be selected from groups represented by Formulae 4-1 to 4-19, but embodiments of the present disclosure are not limited thereto:




Formular 4-1




Formular 4-2




Formular 4-3




Formular 4-4




Formular 4-5




Formular 4-6



Formular 4-7




Formular 4-8




Formular 4-9


-continued




Formular 4-10




Formular 4-11




Formular 4-12




Formular 4-13



Formular 4-14




Formular 4-15




Formular 4-16


-continued



Formular 4-17



Formular 4-18



Formular 4-19

[0050] * and *' in Formulae 4-1 to 4-19 each indicate a binding site to a neighboring atom.

[0051] a₁ to a₁₁ in Formulae 2-1 to 2-5 may each independently be 0, 1, 2, 3, 4, or 5.

[0052] a₁ indicates the number of L₁(s), wherein, when a₁ is 0, *(L₁)_{a₁}-*' indicates a single bond, and when a₁ is two or more, two or more L₁(s) may be identical to or different from each other. a₂ to a₁₁ may be understood by referring to the description of a₁ and the structures of Formulae 2-1 to 2-5. For example, a₁ to a₁₀ may each independently be 0, 1, 2, or 3, but embodiments of the present disclosure are not limited thereto.

[0053] Ar₁ to Ar₇ in Formulae 2-1 to 2-5 may be selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, substituted or unsubstituted C₂-C₆₀ alkenyl group, substituted or unsubstituted C₂-C₆₀ alkynyl group, substituted or unsubstituted C₁-C₆₀ alkoxy group, substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group. For example, Ar₁ to Ar₇ may each independently be selected from:

[0054] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluore-


nyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzonaphthothiophenyl group, a dinaphthothiophenyl group, a benzonaphthosilolyl group, and a dinaphthosilolyl group; and

[0055] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzonaphthothiophenyl group, a dinaphthothiophenyl group, a benzonaphthosilolyl group, and a dinaphthosilolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy


group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzosilolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzonaphthothiophenyl group, a dinaphthothiophenyl group, a benzonaphthosilolyl group, a dinaphthosilolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0056] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.


[0057] In one embodiment, Ar₁ to Ar₇ may each independently be selected from groups represented by Formulae 5-1 to 5-25:



Formula 5-1














Formula 5-2




Formula 5-3

-continued




-continued

Formula 5-4




Formula 5-5

Formula 5-6




Formula 5-7

Formula 5-8




Formula 5-9

Formula 5-10





Formula 5-11

Formula 5-12



Formula 5-13

Formula 5-14



Formula 5-15

Formula 5-16

Formula 5-17


Formula 5-18

Formula 5-19


Formula 5-20

Formula 5-21


-continued




Formula 5-22



Formula 5-23



Formula 5-24



Formula 5-25


[0058] In Formulae 5-1 to 5-25,

[0059] Y_{31} may be O, S, $C(Z_{33})(Z_{34})$, $N(Z_{35})$, or $Si(Z_{36})(Z_{37})$,


[0060] Z_{31} to Z_{37} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and $-Si(Q_{31})(Q_{32})(Q_{33})$, wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0061] $e3$ may be an integer from 1 to 3,[0062] $e4$ may be an integer from 1 to 4,[0063] $e5$ may be an integer from 1 to 5,[0064] $e6$ may be an integer from 1 to 6,[0065] $e7$ may be an integer from 1 to 7,[0066] $e9$ may be an integer from 1 to 9, and


[0067] * indicates a binding site to a neighboring atom. For example, Ar_1 to Ar_7 may each independently be selected from groups represented by Formulae 6-1 to 6-18, but embodiments of the present disclosure are not limited thereto:




Formula 6-1




Formula 6-2




Formula 6-3



Formula 6-4




Formula 6-5




Formula 6-6



Formula 6-7










Formula 6-8



Formula 6-9

-continued



Formula 6-10

Formula 6-11

Formula 6-12

Formula 6-13



Formual 6-14

Formula 6-15

Formula 6-16

-continued

Formula 6-17



Formula 6-18

[0068] * in Formulae 6-1 to 6-18 indicates a binding site to a neighboring atom.

[0069] b1 to b4 in Formulae 2-1 to 2-4 may each independently be 1, 2, or 3. For example, b1 to b4 may be 1, but embodiments of the present disclosure are not limited thereto.

[0070] In one embodiment, R₁ to R₁₁ may each independently be selected from:

[0071] the group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group;

[0072] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

[0073] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a

dibenzofuranyl group, a dibenzothiophenyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a thienyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}); and

[0074] —Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), —B(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1) and —P(=O)(Q_1)(Q_2),

[0075] wherein Q_1 to Q_3 and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.

[0076] In one embodiment, R_1 to R_{11} may each independently be selected from:

[0077] the group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, —Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2),

[0078] wherein Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0079] c_3 in Formula 1 may be 0, 1, 2, 3, 4, 5, or 6. c_3 indicates the number of R_3 (s), wherein, when c_3 is two or more, two or more R_3 (s) may be identical to or different from each other. For example, c_3 may be 0, 1, 2, or 3, but embodiments of the present disclosure are not limited thereto.



[0080] In one embodiment, in Formula 1, X_1 may be N, X_2 may be C(R_2), R_2 may be selected from groups represented by Formulae 2-1 to 2-5, and R_3 to R_{11} may each independently be selected from hydrogen, deuterium, —F, —Cl,

—Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.


[0081] In one embodiment, in Formula 1, X_1 may be C(R_1), X_2 may be N, R_1 may be selected from groups represented by Formulae 2-1 to 2-5, and R_3 to R_{11} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.

[0082] In one embodiment, in Formula 1, R_3 may be selected from groups represented by Formulae 2-1 to 2-5, R_1 , R_2 and R_4 to R_{11} may each independently be selected from the group represented by Formula 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.

[0083] In one embodiment, Formula 1 may be represented by one of Formulae 1A to 1S:




-continued




Formula 1C


-continued




Formula 1H




Formula 1D




Formula 1I




Formula 1E




Formula 1J




Formula 1F



Formula 1K




Formula 1G



Formula 1L


-continued

Formula 1M




-continued


Formula 1R




Formula 1S




Formula 1N




Formula 1O



Formula 1P



Formula 1Q



[0084] In Formulae 1A to 1S,

[0085] R_{31} to R_{36} may be the same as described in connection with R_3 , and

[0086] X_1 , X_2 , and R_3 to R_{11} are the same as described above.


[0087] In one embodiment, in Formulae 1A to 1S, X_1 may be N, X_2 may be $C(R_2)$, and R_2 may be selected from groups represented by Formulae 2-1 to 2-5. For example, in Formulae 1A to 1S, X_1 may be N, X_2 may be $C(R_2)$, R_2 may be selected from groups represented by Formulae 2-1 to 2-5, R_{31} to R_{36} and R_4 to R_{11} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.

[0088] In one embodiment, in Formulae 1A to 1S, X_1 may be $C(R_1)$, X_2 may be N, and R_1 may be selected from groups represented by Formulae 2-1 to 2-5. For example, in Formulae 1A to 1S, X_1 may be $C(R_1)$, X_2 may be N, R_1 may be selected from groups represented by Formulae 2-1 to 2-5, and R_{31} to R_{36} and R_4 to R_{11} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.

[0089] In one embodiment, in Formulae 1A, 1B, 1D to 1G, 1I to 1N, and 1P to 1S, X_1 may be N, X_2 may be $C(R_2)$, and R_{32} may be selected from groups represented by Formulae 2-1 to 2-5. For example, in Formulae 1A, 1B, 1D to 1G, 1I to 1N, and 1P to 1S, X_1 may be N, X_2 may be $C(R_2)$, R_{32} may be selected from groups represented by Formulae 2-1 to 2-5, and R_2 , R_{31} , R_{33} to R_{36} and R_4 to R_{11} may each independently be selected from the group represented by Formula 2-5, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.

[0090] In one embodiment, in Formulae 1A, 1B, 1D to 1G, 1I to 1N, and 1P to 1S, X_1 may be $C(R_1)$, X_2 may be N, and R_{32} may be selected from groups represented by Formulae 2-1 to 2-5. For example, in Formulae 1A, 1B, 1D to 1G, 1I to 1N, and 1P to 1S, X_1 may be $C(R_1)$, X_2 may be N, R_{32} may be selected from groups represented by Formulae 2-1 to 2-5, R_1 , R_{31} , R_{33} to R_{36} , and R_4 to R_{11} may each independently be selected from the group represented by Formula 2-5, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzo-fluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group.


[0091] The condensed cyclic compound represented by Formula 1 may be one of Compounds 1 to 192, but embodiments of the present disclosure are not limited thereto:




1

-continued


2




3







4



-continued




-continued




-continued


10




11



12




13




-continued


14




15



16




17




-continued


18




19



20




21




-continued


22




23



24




-continued






25

-continued



26




27

28

29

-continued


30




31

-continued



34




35




32




33




36



37




-continued




38


39




40




41




-continued



42




43




44

-continued




45


-continued




47



46




48




49

-continued




50

-continued




51


53






52



54




-continued



-continued


55

58




59

56




57





60




-continued



-continued




-continued




66

-continued





67

69






68

70




-continued



-continued


71

74




75

72





73

76




-continued



-continued


77

79




80


78



81




-continued



-continued


82

85




83

86




84

-continued



87


-continued




90

91


88





92




89




93




94




95




96




97




98




99



100




101




-continued


102




103



104




105




-continued


106




107



108




109




-continued


110




111



112




113




-continued


114




115



116




117




-continued


118




119



120




121




-continued


122




123



124




125




-continued


126




127



128




129




-continued


130




131




132



133




-continued




134


-continued




138




135




139




136



140







137







141

-continued




-continued




-continued


150




151



152




153




-continued


154




155




156










157



-continued




-continued




-continued

166




-continued


169




167




170




168



171




-continued




172


-continued




175




173



176




174




177

-continued




178


-continued




181




179



182




180




183

-continued




184


-continued




187




185



188






186



189

-continued



[0092] The condensed cyclic compound represented by Formula 1 has excellent charge transport characteristics and thermal stability. Accordingly, an organic light-emitting device including the condensed cyclic compound represented by Formula 1 may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

[0093] A method of synthesizing the condensed cyclic compound represented by Formula 1 may be understood by referring to the following examples.

[0094] At least one of the condensed cyclic compounds represented by Formula 1 may be used between a pair of electrodes of an organic light-emitting device. In one or more embodiments, the condensed cyclic compound may be included in an electron transport region.

[0095] According to another aspect, an organic light-emitting device includes a first electrode; a second electrode facing the first electrode; and an organic layer that is disposed between the first electrode and the second electrode and includes an emission layer, wherein the organic layer includes at least one of condensed cyclic compounds represented by Formula 1.

[0096] The expression “(an organic layer) includes at least one of the condensed cyclic compounds” used herein may include a case in which “(an organic layer) includes identical condensed cyclic compounds represented by Formula 1” and a case in which “(an organic layer) includes two or more different condensed cyclic compounds represented by Formula 1.”

[0097] For example, the organic layer may include, as the condensed cyclic compound, only Compound 1. In this regard, Compound 1 may exist only in an electron transport layer of the organic light-emitting device. In one or more embodiments, the organic layer may include, as the condensed cyclic compound, Compound 1 and Compound 2. In one or more embodiments, Compound 1 and Compound 2 may both exist in an identical layer (for example, Compound 1 and Compound 2 may both exist in an electron transport layer), or may exist in different layers (for example, Compound 1 may exist in an electron transport layer and Compound 2 may exist in an electron injection layer).

[0098] The organic layer includes i) a hole transport region that is disposed between the first electrode (anode) and the emission layer and includes at least one of a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer; and ii) an electron transport region that is disposed between the emission layer and the second electrode (cathode) and includes at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer. The electron transport region may include at least one condensed cyclic compound represented by Formula 1. For example, the electron transport region of the organic light-emitting device may include the electron transport layer, wherein the electron transport layer may include at least one of the condensed cyclic compound represented by Formula 1. For example, the electron transport region of the organic light-emitting device may include an electron injection layer, and the electron injection layer may include at least one of the organometallic compounds represented by Formula 1.

[0099] The organic light-emitting device may further include at least one selected from a first capping layer disposed in a pathway along which light generated in an emission layer travels toward outside through the first electrode and a second capping layer disposed in a pathway along which light generated in an emission layer travels toward outside through the second electrode, and the at least one selected from the first capping layer and the second capping layer may include at least one of the condensed cyclic compounds represented by Formula 1.

[0100] For example, the organic light-emitting device may have i) a stack structure including a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order, ii) a stack structure including a first capping layer, a first electrode, an organic layer, and a second electrode which are sequentially stacked in this stated order, or iii) a stack structure including a first capping layer, a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order, and at least one selected from the first capping layer and the second capping layer may include the condensed cyclic compound.

[0101] The term “organic layer” used herein refers to a single layer and/or a plurality of layers disposed between the first electrode and the second electrode of the organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[Description of FIG. 1]

[0102] FIG. 1 is a schematic cross-sectional view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0103] Hereinafter, the structure of the organic light-emitting device 10 according to the present embodiment and a method of manufacturing the same will be described in connection with FIG. 1.

[First Electrode 110]

[0104] In FIG. 1, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

[0105] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for forming the first electrode 110 may be selected from materials with a high work function to facilitate hole injection.

[0106] The first electrode 110 may be a reflective electrode, a semi-reflective electrode, or a transmissive electrode. When the first electrode 110 is a transmissible electrode, a material for forming a first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO_2), zinc oxide (ZnO), and any combinations thereof, but embodiments of the present disclosure are not limited thereto. In one or more embodiments, when the first electrode 110 is a semi-transmissible electrode or a reflectable electrode, a material for forming a first electrode may be selected from magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), and any combinations thereof, but embodiments of the present disclosure are not limited thereto.

[0107] The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode 110 is not limited thereto.

[Organic Layer 150]

[0108] The organic layer 150 is disposed on the first electrode 110. The organic layer 150 may include an emission layer.

[0109] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.


[Hole Transport Region in Organic Layer 150]

[0110] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including different materials, or iii) a multi-layered structure having a plurality of layers including different materials.

[0111] The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, and an electron blocking layer (EBL).


[0112] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein for each structure, constituting layers are sequentially stacked from the first electrode 110 in this stated order, but the structure of the hole transport region is not limited thereto.

[0113] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPB), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




m-MTDATA


-continued



TDATA





2-TNATA




NPB


-continued

 β -NPB


TPD






Spiro-TPD



Spiro-NPB



methylated NPB



[0114] In Formulae 201 and 202, L_{201} to L_{204} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, substituted or unsubstituted C_6 - C_{60} arylene group, substituted or unsubstituted C_1 - C_{60} heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0115] L_{205} may be selected from $^*\text{—O—}^*$, $^*\text{—S—}^*$, $^*\text{—N}(\text{Q}_{201})\text{—}^*$, a substituted or unsubstituted C_1 - C_{20} alkylene group, substituted or unsubstituted C_2 - C_{20} alkenylene group, substituted or unsubstituted C_3 - C_{10} cycloalkylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, substituted or unsubstituted C_6 - C_{60} arylene group, substituted or unsubstituted C_1 - C_{60} heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0116] xa1 to xa4 may each independently be an integer from 0 to 3,

[0117] xa5 may be an integer from 1 to 10,

[0118] R_{201} to R_{204} and Q_{201} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, substituted or unsubstituted C_6 - C_{60} aryl group, substituted or unsubstituted C_6 - C_{60} aryloxy group, substituted or unsubstituted C_6 - C_{60} arylthio group, substituted or unsubstituted C_1 - C_{60} heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0119] For example, in Formula 202, R_{201} and R_{202} may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R_{203} and R_{204} may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

[0120] In one embodiment, in Formulae 201 and 202,

[0121] L_{201} to L_{205} may each independently be selected from:

[0122] a phenylene group, a pentalenylene group, an indenylene group, a naphthylenylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-bifluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a benzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylene group, and a pyridinylenylene group; and

[0123] a phenylene group, a pentalenylene group, an indenylene group, a naphthylenylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylenylene group, a fluorenylenylene group, a spiro-bifluorenylenylene group, a benzofluorenylenylene group, a dibenzofluorenylenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a benzothiophenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F , —Cl , —Br , —I , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a

cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0124] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0125] In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

[0126] In one or more embodiments, xa5 may be 1, 2, 3, or 4.

[0127] In one or more embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0128] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group,

an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0129] wherein Q₃₁ to Q₃₃ may be the same as described above.

[0130] In one or more embodiments, at least one of R₂₀₁ to R₂₀₃ in Formula 201 may each independently be selected from:

[0131] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0132] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, but embodiments of the present disclosure are not limited thereto.


[0133] In one or more embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be linked via a single bond, and/or ii) R₂₀₃ and R₂₀₄ may be linked via a single bond.

[0134] In one or more embodiments, at least one of R₂₀₁ to R₂₀₄ in Formula 202 may be selected from:


[0135] a carbazolyl group; and

[0136] a carbazolyl group, substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group and a dibenzothiophenyl group, but embodiments of the present disclosure are not limited thereto.


[0137] The compound represented by Formula 201 may be represented by Formula 201A:




[0138] For example, the compound represented by Formula 201 may be represented by Formula 201A(1), but embodiments of the present disclosure are not limited thereto:




[0139] In one embodiment, the compound represented by Formula 201 may be represented by Formula 201A-1, but embodiments of the present disclosure are not limited thereto:



[0140] The compound represented by Formula 202 may be represented by Formula 202A:



[0141] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:



[0142] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0143] L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} may be the same as described above,

[0144] R_{211} and R_{212} may be the same as described in connection with R_{203} ,


[0145] R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuran group, a benzothiophenyl group, a benzocarbonyl group, a dibenzothiophenyl group, a benzocarbonyl group, a dibenzocarbonyl group, a dibenzosilolyl group, and a pyridinyl group.

[0146] The hole transport region may include at least one compound selected from Compounds HT1 to HT39, but embodiments of the present disclosure are not limited thereto:


HT1




HT2



HT3





HT4





-continued
HT5

HT6



HT7



HT8



-continued



HT9

HT10



HT11



HT12



-continued



HT13

HT14





HT15

HT16




HT17

HT18




-continued


HT19




HT20



HT21




HT22





-continued

HT23





HT24

HT25




HT26

HT27




-continued


HT28




HT29




HT30




HT31




HT32




HT33




HT34




HT35




-continued
HT36


HT37



HT38



HT39



[0147] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one selected from a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, for example about 100 Å to about 1,500 Å. When the thickness of each of the hole transport region, the hole injection layer, and the hole transport layer is within these ranges, satisfactory hole transport characteristics may be obtained without a substantial increase in driving voltage.

[0148] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may each include the materials as described above.

[p-Dopant]

[0149] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-

generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0150] The charge-generation material may be, for example, a p-dopant.

[0151] In one embodiment, a lowest unoccupied molecular orbital (LUMO) of the p-dopant may be -3.5 eV or less.

[0152] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments are not limited thereto.


[0153] In one embodiment, the p-dopant may include at least one selected from:

[0154] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and F4-2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (TCNQ);


[0155] metal oxide, such as tungsten oxide and molybdenum oxide;

[0156] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and


[0157] a compound represented by Formula 221, but embodiments of the present disclosure are not limited thereto:



<HAT-CN>



<F4-TCNQ>



<Formula 211>

[0158] In Formula 221,

[0159] R₂₂₁ to R₂₂₃ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and at least one of R₂₂₁ to R₂₂₃ may have at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C₁-C₂₀ alkyl group substituted with —F, a C₁-C₂₀ alkyl group substituted with —Cl, a C₁-C₂₀ alkyl group substituted with —Br, and a C₁-C₂₀ alkyl group substituted with —I.

[Emission Layer in Organic Layer 150]

[0160] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials are mixed with each other in a single layer to emit white light.


[0161] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0162] An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host, but is not limited thereto.

[0163] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent emission characteristics may be obtained without a substantial increase in driving voltage.

[Host in Emission Layer]

[0164] The host may include a compound represented by Formula 301.



[0165] In Formula 301,

[0166] Ar₃₀₁ may be selected from a substituted or unsubstituted C₅-C₆₀ carbocyclic group or a substituted or unsubstituted C₁-C₆₀ heterocyclic group,

[0167] xb11 may be 1, 2, or 3,

[0168] L₃₀₁ may be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, substituted or unsubstituted C₆-C₆₀ arylene group, substituted or unsubstituted C₁-C₆₀ heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0169] xb1 may be an integer from 0 to 5,

[0170] R₃₀₁ may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₃₀₁)(Q₃₀₂)(Q₃₀₃), —N(Q₃₀₁)(Q₃₀₂), —B(Q₃₀₁)(Q₃₀₂), —C(=O)(Q₃₀₁), —S(=O)₂(Q₃₀₁), and —P(=O)(Q₃₀₁)(Q₃₀₂), and

[0171] xb21 may be an integer from 1 to 5, wherein

[0172] Q₃₀₁ to Q₃₀₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0173] In one embodiment, Ar₃₀₁ in Formula 301 may be selected from

[0174] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene



group, a phenylene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and

[0175] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenylene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0176] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

[0177] When xb11 in Formula 301 is two or more, two or more Ar₃₀₁(s) may be linked via a single bond.

[0178] In one or more embodiments, the compound represented by Formula 301 may be represented by one of Formula 301-1 or 301-2:



[0179] In Formulae 301-1 to 301-2,

[0180] A₃₀₁ to A₃₀₄ may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group,

a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group,

[0181] X₃₀₁ may be O, S, or N-[(L₃₀₄)_{xb4}-R₃₀₄],

[0182] R₃₁₁ to R₃₁₄ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0183] xb22 and xb23 may each independently be 0, 1, or 2,

[0184] L₃₀₁, xb1, R₃₀₁, and Q₃₁ to Q₃₃ may be the same as described above,

[0185] L₃₀₂ to L₃₀₄ may each independently be the same as described in connection with L₃₀₁.

[0186] Xb2 to xb4 may each independently be the same as described in connection with xb1, and

[0187] R₃₀₂ to R₃₀₄ may each independently be the same as described in connection with R₃₀₁.

[0188] For example, L₃₀₁ to L₃₀₄ in Formulae 301, 301-1 and 301-2 may each independently be selected from:

[0189] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a

thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group,

group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group; and

[0190] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolyene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolyene group, a dibenzocarbazolyene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylene group, a pyrazolyene group, a thiazolyene group, an isothiazolylenylene group, an oxazolyene group, an isoxazolyene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylenylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolyene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolyene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrim-

idinyl group, an aza carbazolyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

[0191] wherein Q₃₁ to Q₃₃ may be the same as described above.

[0192] In one embodiment, R₃₀₁ to R₃₀₄ in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0193] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group; and


[0194] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazoyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazoyl group, a dibenzocarbazoyl group, a dibenzosilolyl group, a pyridinyl group, an imidazoyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazoyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁–C₂₀ alkyl group, a C₁–C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an

anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an aza carbazolyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$.


[0195] wherein Q_{31} to Q_{33} may be the same as described above.

[0196] In one or more embodiments, the host may include an alkaline-earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), an Mg complex, and a Zn complex.

[0197] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, but is not limited thereto:




H1




H2


-continued




H3




H4




H5



H6




H7




H8

-continued


H9




H10




H11



H12




H13




-continued


H14




H15




H16



H17




H18




-continued


H19




H20



H21




H22




-continued


H23




H24



H25




-continued




H26


-continued




H29




H30




H31



H27




H28




H32

-continued


H33




H34



H35




H36




-continued


H37




H38



H39




-continued




H40


-continued




H43




H41



H44




H42




H45

-continued


H46




H47




H48




H49



H50




H51




-continued


H52




H53



H54





H55



[Phosphorescent Dopant Included in Emission Layer in Organic Layer **150**]

[0198] In one or more embodiments, the phosphorescent dopant may include an organometallic complex represented by Formula 401:



<Formula 401>

<Formula 402>

[0199] In Formulae 401 and 402,

[0200] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0201] L_{401} may be selected from ligands represented by Formula 402, and xc1 may be 1, 2, or 3, wherein, when xc1 is two or more, two or more $L_{401}(S)$ may be identical to or different from each other,

[0202] L_{402} may be an organic ligand, and xc2 may be an integer from 0 to 4, wherein, when xc2 is two or more, two or more $L_{402}(s)$ may be identical to or different from each other,

[0203] X_{401} to X_{404} may each independently be nitrogen or carbon;

[0204] X_{401} and X_{403} may be linked via a single bond or a double bond, and X_{402} and X_{404} may be linked via a single bond or a double bond,

[0205] A_{401} and A_{402} may each independently be selected from a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

[0206] X_{405} may be a single bond, $^*O^*$, $^*S^*$, $^*C(=O)^*$, $^*N(Q_{411})^*$, $^*C(Q_{411})(Q_{412})^*$, $^*C(Q_{411})=C(Q_{412})^*$, $^*C(Q_{411})^*$, or $^*C(Q_{411})=^*$, Q_{411} and Q_{412} may be hydrogen, deuterium, C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0207] X_{406} may be a single bond, O or S,

[0208] R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, substituted or unsubstituted C_1 - C_{20} alkyl group, substituted or unsubstituted C_1 - C_{20} alkoxy group, substituted or unsubstituted C_3 - C_{10} cycloalkyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, substituted or unsubstituted C_6 - C_{60} aryl group, substituted or unsubstituted C_6 - C_{60} aryloxy group, substituted or unsubstituted C_6 - C_{60} arylthio group, substituted or unsubstituted C_1 - C_{60} heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$ and $-P(=O)(Q_{401})(Q_{402})$, wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_6 - C_{20} aryl group, and a C_1 - C_{20} heteroaryl group,

[0209] xc11 and xc12 may each independently be an integer from 0 to 10,

[0210] $*$ and * in Formula 402 each indicate a binding site to M in Formula 401.

[0211] In one embodiment, A_{401} and A_{402} in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0212] In one or more embodiments, in Formula 402, i) X_{401} may be nitrogen, and X_{402} may be carbon, or ii) X_{401} and X_{402} may all be nitrogen.

[0213] In one or more embodiments, R_{401} and R_{402} in Formula 402 may each independently be selected from:

[0214] hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0215] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0216] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0217] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group and a dibenzothiophenyl group; and


[0218] $-\text{Si}(\text{Q}_{401})(\text{Q}_{402})(\text{Q}_{403})$, $-\text{N}(\text{Q}_{401})(\text{Q}_{402})$, $-\text{B}(\text{Q}_{401})(\text{Q}_{402})$, $-\text{C}(=\text{O})(\text{Q}_{401})$, $\text{S}(=\text{O})_2(\text{Q}_{401})$, and $-\text{P}(=\text{O})(\text{Q}_{401})(\text{Q}_{402})$,

[0219] wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.


[0220] In one or more embodiments, when xc1 in Formula 401 is two or more, two $\text{A}_{401}(\text{s})$ in two or more $\text{L}_{401}(\text{s})$ may optionally be linked via X_{407} , which is a linking group, or two $\text{A}_{402}(\text{s})$ in two or more $\text{L}_{401}(\text{s})$ may optionally be linked via X_{408} , which is a linking group (see Compounds PD1 to PD4 and PD7). X_{407} and X_{408} may each independently be a single bond, $^*\text{—O—}^*$, $^*\text{—S—}^*$, $^*\text{—C(=O)—}^*$, $^*\text{—N(Q}_{413}\text{)—}^*$, $^*\text{—C(Q}_{413}\text{)(Q}_{414}\text{)—}^*$, or $^*\text{—C(Q}_{413}\text{)=C(Q}_{414}\text{)—}^*$ (herein, Q_{413} and Q_{414} may each independently be selected from hydrogen, deuterium, C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), but embodiments of the present disclosure are not limited thereto.

[0221] L_{402} in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L_{402} may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), $-\text{C}(=\text{O})$, isonitrile, $-\text{CN}$, and phosphorus (for example, phosphine, or phosphite), but embodiments of the present disclosure are not limited thereto.

[0222] In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, but embodiments of the present disclosure are not limited thereto:




PD1




PD2


-continued




PD3




PD4




PD5



PD6




PD7




PD8

-continued




PD9

-continued




PD14





PD10

PD15





PD11

PD16





PD12

PD17







PD13

PD18




-continued




-continued

PD19




PD23

PD20



PD24

PD21



PD25


[Fluorescent Dopant in Emission Layer]

PD22

[0223] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0224] The fluorescent dopant may include a compound represented by Formula 501:

<Formula 501>



In Formula 501,

[0225] Ar_{501} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0226] L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, substituted or unsubstituted C_6 - C_{60} arylene group, substituted or unsubstituted C_1 - C_{60} heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0227] $xd1$ to $xd3$ may each independently be an integer from 0 to 3,

[0228] R_{501} and R_{502} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, substituted or unsubstituted C_6 - C_{60} aryl group, substituted or unsubstituted C_6 - C_{60} aryloxy group, substituted or unsubstituted C_6 - C_{60} arylthio group, substituted or unsubstituted C_1 - C_{60} heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0229] $xd4$ may be an integer from 1 to 6.

[0230] In one embodiment, Ar_{501} in Formula 501 may be selected from:

[0231] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, and an indenophenanthrene group; and

[0232] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

[0233] In one or more embodiments, L_{50} to L_{503} in Formula 501 may each independently be selected from:

[0234] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group; and

[0235] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group and a pyridinyl group.

[0236] In one or more embodiments, R_{501} and R_{502} in Formula 501 may each independently be selected from:

[0237] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group and a pyridinyl group; and


[0238] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, and $\text{—Si(Q}_{31}\text{)(Q}_{32}\text{)(Q}_{33}\text{)}$.


[0239] wherein Q_{31} to Q_{33} may be selected from a $\text{C}_1\text{—C}_{10}$ alkyl group, a $\text{C}_1\text{—C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0240] In one or more embodiments, xd4 in Formula 501 may be two, but embodiments of the present disclosure are not limited thereto.


[0241] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:




FD1



FD2



FD3




FD4

-continued


-continued

FD5




-continued


FD8




FD6




FD9




FD7




FD10




-continued




FD12




FD13




FD14




-continued




FD16



FD17




FD18




-continued

FD19





-continued

FD21




FD22


FD20




[0242] In one embodiment, the fluorescent dopant may be selected from the following compounds, but embodiments of the present disclosure are not limited thereto:



DPVBi



DPAVB



[Electron Transport Region in Organic Layer 150]

[0243] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0244] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

[0245] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein for each structure, constituting layers are sequentially stacked from an emission layer. However, embodiments of the structure of the electron transport region are not limited thereto.

[0246] For example, the electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a condensed cyclic compound represented by Formula 1.

[0247] In one embodiment, the electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

[0248] The “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*'$ moiety as a ring-forming moiety.

[0249] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered heteromonocyclic group having at least one $*-N=*'$ moiety, ii) a heteropoly cyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one $*-N=*'$ moiety are condensed with each other, or iii) a heteropoly cyclic group in which at least one of 5-membered to 7-membered heteromonocyclic

groups, each having at least one *—N=* moiety, is condensed with at least one $\text{C}_5\text{—C}_{60}$ carbocyclic group.

[0250] Examples of the π electron-depleted nitrogen-containing ring include an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, but are not limited thereto.

[0251] For example, the electron transport region may include a compound represented by Formula 601:



[0252] In Formula 601,

[0253] Ar_{601} may be a substituted or unsubstituted $\text{C}_5\text{—C}_{60}$ carbocyclic group or a substituted or unsubstituted $\text{C}_1\text{—C}_{60}$ heterocyclic group,

[0254] xe11 may be 1, 2, or 3,

[0255] L_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{—C}_{10}$ cycloalkylene group, substituted or unsubstituted $\text{C}_1\text{—C}_{10}$ heterocycloalkylene group, substituted or unsubstituted $\text{C}_3\text{—C}_{10}$ cycloalkenylene group, substituted or unsubstituted $\text{C}_1\text{—C}_{10}$ heterocycloalkenylene group, substituted or unsubstituted $\text{C}_6\text{—C}_{60}$ arylene group, substituted or unsubstituted $\text{C}_1\text{—C}_{60}$ heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0256] xe1 may be an integer from 0 to 5,

[0257] R_{601} may be selected from a substituted or unsubstituted $\text{C}_3\text{—C}_{10}$ cycloalkyl group, substituted or unsubstituted $\text{C}_1\text{—C}_{10}$ heterocycloalkyl group, substituted or unsubstituted $\text{C}_3\text{—C}_{10}$ cycloalkenyl group, substituted or unsubstituted $\text{C}_1\text{—C}_{10}$ heterocycloalkenyl group, substituted or unsubstituted $\text{C}_6\text{—C}_{60}$ aryl group, substituted or unsubstituted $\text{C}_6\text{—C}_{60}$ aryloxy group, substituted or unsubstituted $\text{C}_6\text{—C}_{60}$ arylthio group, substituted or unsubstituted $\text{C}_1\text{—C}_{60}$ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $\text{—Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$, $\text{—C(=O)}(\text{Q}_{601})$, $\text{—S(=O)}_2(\text{Q}_{601})$, and $\text{—P(=O)}(\text{Q}_{601})(\text{Q}_{602})$, wherein Q_{601} to Q_{603} may each independently be a $\text{C}_1\text{—C}_{10}$ alkyl group, a $\text{C}_1\text{—C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0258] xe21 may be an integer from 1 to 5.

[0259] In one embodiment, at least one selected from Ar_{601} in the number of xe11 and R_{601} in the number of xe21 may include the π electron-depleted nitrogen-containing ring described above.

[0260] In one embodiment, ring Ar_{601} in Formula 601 may be selected from:

[0261] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triph-

enylene group, a pyrene group, a chrysene group, a naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0262] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, phenanthroline group, phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F , —Cl , —Br , —I , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a $\text{C}_1\text{—C}_{20}$ alkyl group, a $\text{C}_1\text{—C}_{20}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $\text{—Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $\text{—S(=O)}_2(\text{Q}_{31})$, and $\text{—P(=O)}(\text{Q}_{31})(\text{Q}_{32})$.


[0263] wherein Q_{31} to Q_{33} may each independently be selected from a $\text{C}_1\text{—C}_{10}$ alkyl group, a $\text{C}_1\text{—C}_{10}$ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0264] When xe11 in Formula 601 is two or more, two or more $\text{Ar}_{601}(\text{S})$ may be linked via a single bond.

[0265] In one or more embodiments, Ar_{601} in Formula 601 may be an anthracene group.

[0266] In one or more embodiments, the compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



[0267] In Formula 601-1,

[0268] X_{614} may be N or C(R_{614}), X_{615} may be N or C(R_{615}), X_{616} may be N or C(R_{616}), at least one of X_{614} to X_{616} may be N,

[0269] L_{611} to L_{613} may each independently be the same as described in connection with L_{601} ,

[0270] xe611 to xe613 may each independently be the same as described in connection with xe1,

[0271] R_{611} to R_{613} may each independently be the same as described in connection with R_{601} , and

[0272] R_{614} to R_{616} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0273] In one embodiment, L_{601} and L_{611} to L_{613} in Formulae 601 and 601-1 may each independently be selected from:

[0274] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0275] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, but embodiments of the present disclosure are not limited thereto.

thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, but embodiments of the present disclosure are not limited thereto.

[0276] In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0277] In one or more embodiments, R_{601} and R_{611} to R_{613} in Formulae 601 and 601-1 may each independently be selected from

[0278] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an

isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

[0279] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a


cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0280] —S(=O)₂(Q₆₀₁) and —P(=O)(Q₆₀₁)(Q₆₀₂),


[0281] wherein Q₆₀₁ and Q₆₀₂ are the same as described above.

[0282] The electron transport region may include at least one compound represented by Compounds ET1 to ET36, but embodiments of the present disclosure are not limited thereto:

ET1




ET2




-continued

ET3




-continued


ET6



ET4





ET7




ET8

ET5




-continued




ET9

-continued




ET11

ET12




ET10

ET13




-continued



ET14



-continued

ET17





ET15

ET18




ET16

ET19




-continued




ET20

-continued




ET23


ET21




ET24



ET22




ET25




-continued

ET26




-continued


ET29




ET27




ET28



ET30




ET31




-continued

ET32




-continued


ET35




ET33




ET36




ET34



[0283] In one embodiment, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BOP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ.

Alq₃



[0284] A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.



[0285] A thickness of the electron transport layer may be from about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the ranges described above, excellent electron transport characteristics may be obtained without a substantial increase in driving voltage.

[0286] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0287] The metal-containing material may include at least one selected from alkali metal complex and alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from an Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, an Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy

phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenylan oxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0288] In one embodiment, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



[0289] The electron transport region may include an electron injection layer that facilitates electron injection from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0290] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including different materials, or iii) a multi-layered structure having a plurality of layers including different materials.

[0291] The electron injection layer may include alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof.

[0292] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one embodiment, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs, but embodiments of the present disclosure are not limited thereto.

[0293] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0294] The rare-earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

[0295] The alkali metal compound, the alkaline earth-metal compound, and the rare-earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal and rare-earth metal.

[0296] The alkali metal compound may be selected from alkali metal oxides, such as Li₂O, Cs₂O, or K₂O, and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, KI,

or RbI. In one embodiment, the alkali metal compound may be selected from LiF, Li₂O, NaF, LiI, NaI, CsI, and KI, but embodiments of the present disclosure are not limited thereto.

[0297] The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as BaO, SrO, CaO, Ba_xSr_{1-x}O (0<x<1), or Ba_xCa_{1-x}O (0<x<1). In one embodiment, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO, but embodiments of the present disclosure are not limited thereto.

[0298] The rare-earth metal compound may be selected from YbF₃, ScF₃, ScO₃, Y₂O₃, Ce₂O₃, GdF₃, and TbF₃. In one embodiment, the rare-earth metal compound may be selected from YbF₃, ScF₃, TbF₃, YbI₃, ScI₃, and TbI₃, but embodiments of the present disclosure are not limited thereto.

[0299] The alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare-earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may each independently be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenyl oxazole, hydroxy phenylthiazole, hydroxy diphenyl oxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

[0300] The electron injection layer may consist of alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, alkali metal, alkaline earth metal, rare-earth metal, alkali metal compound, alkaline earth-metal compound, rare-earth metal compound, alkali metal complex, alkaline earth-metal complex, rare-earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0301] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. If the thickness of the electron injection layer is within the ranges described above, excellent electron injection characteristics may be obtained without a substantial increase in driving voltage.

[Second Electrode 190]

[0302] The second electrode 190 may be disposed on the organic layer 150 having such a structure. The second electrode 190 may be a cathode that is an electron injection electrode, and in this regard, as a metal for forming the second electrode 190, metal, alloy, an electrically conductive compound, and a combination thereof, each having a low work function, may be used.

[0303] The second electrode 190 may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca),

magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, but embodiments of the present disclosure are not limited thereto. The first electrode 190 may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0304] The first electrode 190 may have a single-layered structure, or a multi-layered structure including two or more layers.

[Description of FIGS. 2 to 4]

[0305] FIG. 2 is a schematic view of an organic light-emitting device 20 according to an embodiment. The organic light-emitting device 20 includes a first capping layer 210, the first electrode 110, the organic layer 150, and the second electrode 190, which are sequentially stacked in this stated order. FIG. 3 is a schematic view of an organic light-emitting device 30 according to an embodiment. The organic light-emitting device 30 includes the first electrode 110, the organic layer 150, the second electrode 190, and a second capping layer 220, which are sequentially stacked in this stated order. FIG. 4 is a schematic view of an organic light-emitting device 40 according to an embodiment. The organic light-emitting device 40 includes the first capping layer 210, the first electrode 110, the organic layer 150, the second electrode 190, and the second capping layer 220, which are sequentially stacked in this stated order.

[0306] Regarding FIGS. 2 to 4, the first electrode 110, the organic layer 150, and the second electrode 190 may be understood by referring to the description presented in connection with FIG. 1.

[0307] In the organic layer 150 of each of the organic light-emitting devices 20 and 40, light generated in an emission layer may pass through the first electrode 110 and the first capping layer 210 toward the outside, wherein the first electrode 110 may be a semi-transmissive electrode or a transmissive electrode. In the organic layer 150 of each of the organic light-emitting devices 30 and 40, light generated in an emission layer may pass through the second electrode 190 and the second capping layer 220 toward the outside, wherein the second electrode 190 may be a semi-transmissive electrode or a transmissive electrode.

[0308] The first capping layer 210 and the second capping layer 220 may increase external luminescent efficiency according to the principle of constructive interference.

[0309] The first capping layer 210 and the second capping layer 220 may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.




[0310] At least one selected from the first capping layer 210 and the second capping layer 220 may include the condensed cyclic compound represented by Formula 1.

[0311] At least one selected from the first capping layer 210 and the second capping layer 220 may each independently include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrine derivatives, phthalocyanine derivatives, naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In one embodiment, at least one selected

from the first capping layer **210** and the second capping layer **220** may each independently include an amine-based compound.


[0312] In one embodiment, at least one selected from the first capping layer **210** and the second capping layer **220** may each independently include the compound represented by Formula 201 or the compound represented by Formula 202.

[0313] In one or more embodiments, at least one selected from the first capping layer **210** and the second capping layer **220** may each independently include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5, but embodiments of the present disclosure are not limited thereto.




-continued

CP4



CP5



[0314] Hereinbefore, the organic light-emitting device according to an embodiment has been described in connection with FIGS. 1 to 4, but embodiments of the present disclosure are not limited thereto.

[0315] Respective layers constituting the hole transport region, an emission layer, and respective layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, langmuir-blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0316] When the respective layers of the hole transport region, the emission layer, and the respective layers of the electron transport region are formed by deposition, the deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec by taking into account a material for forming a layer to be deposited, and the structure of a layer to be formed.

[0317] When respective layers constituting the hole transport region, an emission layer, and respective layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C., depending on a material to be included in a layer and the structure of each layer to be formed.

[General Definition of Substituents]

[0318] The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and

examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0319] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0320] The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0321] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0322] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0323] The term “C₁-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent saturated monocyclic group having at least one heteroatom selected from N, O, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuran group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0324] The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and does not have aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0325] The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuran group and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0326] The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and the term “C₆-C₆₀ arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be condensed to each other.

[0327] The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group,” as used herein, refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be condensed to each other.

[0328] The term “C₆-C₆₀ aryloxy group” as used herein refers to —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and the term “C₆-C₆₀ arylthio group” as used herein refers to —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0329] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed to each other, only carbon atoms as a ring-forming atom, and non-aromaticity in the entire molecular structure. Examples of the monovalent non-aromatic condensed polycyclic group include a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” as used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0330] The term “monovalent non-aromatic condensed heteropolycyclic group,” as used herein, refers to a monovalent group (for example, having 2 to 60 carbon atoms) that has two or more rings condensed to each other, has at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and has non-aromaticity in the entire molecular structure. Examples of the monovalent non-aromatic condensed heteropolycyclic group include a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0331] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a monovalent group, such as a cyclic phenyl group, such as benzene, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents

connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0332] The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having the same structure as the C₁-C₆₀ carbocyclic group, except that, as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

[0333] At least one of substituent or substituents of the substituted C₅-C₆₀ carbocyclic group, substituted C₁-C₆₀ heterocyclic group, substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0334] deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0335] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₁₁)(Q₁₂)(Q₁₃), -N(Q₁₁)(Q₁₂), -B(Q₁₁)(Q₁₂), -C(=O)(Q₁₁), -S(=O)₂(Q₁₁), and -P(=O)(Q₁₁)(Q₁₂);

[0336] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0337] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino

group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₂₁)(Q₂₂)(Q₂₃), -N(Q₂₁)(Q₂₂), -B(Q₂₁)(Q₂₂), -C(=O)(Q₂₁), -S(=O)₂(Q₂₁), and -P(=O)(Q₂₁)(Q₂₂); and

[0338] -Si(Q₃₁)(Q₃₂)(Q₃₃), -N(Q₃₁)(Q₃₂), -B(Q₃₁)(Q₃₂), -C(=O)(Q₃₁), -S(=O)₂(Q₃₁), and -P(=O)(Q₃₁)(Q₃₂);

[0339] wherein Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0340] The term “Ph” used herein represents a phenyl group, the term “Me” used herein represents a methyl group, the term “Et” used herein represents an ethyl group, the term “ter-Bu” or “Bu” used herein represents a tert-butyl group, and “OMe” used herein represents a methoxy group.

[0341] The term “biphenyl group” as used therein refers to “a phenyl group substituted with a phenyl group.” In other words, a “biphenyl group” is a substituted phenyl group having a C₆-C₆₀ aryl group as a substituent.

[0342] The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group.” The “terphenyl group” is a “substituted phenyl group” having a “C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group” as a substituent.

[0343] * and *¹ used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.


[0344] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples indicates that a molar equivalent of A was identical to a molar equivalent of B.


EXAMPLE

Synthesis Example 1: Synthesis of Compound 5

[0345]

<Reaction Scheme 1>





Synthesis of Intermediate A-1

[0346] Phenylboronic acid (2.44 g, 20.0 mmol), 9-bromo-10-nitrophenanthrene (6.04 g, 20.0 mmol), $\text{Pd}(\text{PPh}_3)_4$ (1.16 g, 1.00 mmol), tetrabutylammonium bromide (TBAB) (0.322 g, 1.00 mmol), and Na_2CO_3 (6.28 g, 60.0 mmol) were dissolved in 80 mL of a mixed solution of toluene/ethanol/

H_2O (3/3/1), and then, stirred at a temperature of 80° C. for 16 hours. The reaction solution was cooled to room temperature, and then, an extraction process was performed three times thereon by using 60 mL of water and 60 mL of diethylether. An organic layer was dried by using MgSO_4 , and the residue obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 4.91 g of Intermediate A-1 (yield of 82%). The obtained compound was identified by LC-MS. $\text{C}_{20}\text{H}_{13}\text{NO}_2$: $\text{M}+1$ 299.1

Synthesis of Intermediate A-2

[0347] Intermediate A-1 (4.91 g, 16.4 mmol), Tin (5.84 g, 49.2 mmol), and 8.2 mL (82 mmol, conc. 36.5%) of HCl were dissolved in 60 mL of ethanol, and then, stirred at a temperature of 100° C. for 8 hours. The reaction solution was cooled to room temperature, and then, 5 g of sodium hydroxide dissolved in 10 mL of water was added to a filtrate obtained by filtering under reduced pressure, and then, the result was subjected three times to an extraction process using 60 mL of water and 60 mL of dichloromethane. An organic layer obtained therefrom was dried by using MgSO_4 , and the residue obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 3.98 g (yield of 90%) of Intermediate A-2. The obtained compound was identified by LC-MS. $\text{C}_{20}\text{H}_{15}\text{N}$: $\text{M}+1$ 269.1

Synthesis of Intermediate A-3

[0348] Intermediate A-2 (3.98 g, 14.7 mmol) and 3-bromobenzaldehyde (5.38 g, 29.4 mmol) were dissolved in 30 mL of trifluoroacetic acid, and then, the mixture was stirred in a seal tube at a temperature of 130° C. for 3 days. The reaction solution was cooled to room temperature, and then, the reaction was quenched by using NaHCO_3 . The result was subjected to an extraction process using 60 mL of water and 60 mL of dichloromethane. An organic layer obtained therefrom was dried by using MgSO_4 , and the residue obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 3.83 g (yield of 60%) of Intermediate A-3. The obtained compound was identified by LC-MS. $\text{C}_{27}\text{H}_{16}\text{BrN}$: $\text{M}+1$ 433.1

Synthesis of Compound 5

[0349] Intermediate A-3 (3.83 g, 8.82 mmol), 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile (3.05 g, 10 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.51 g, 0.44 mmol), and K_2CO_3 (3.66 g, 2.65 mmol) were dissolved in 60 mL of a mixed solution including THF/ H_2O (a volumetric ratio of 2/1), and then, the mixture was stirred at a temperature of 80° C. for 16 hours. The reaction solution was cooled to room temperature, and then, 40 mL of water was added thereto, and an extraction process was performed three times on the result by using 50 mL of ethyl ether. An organic layer obtained therefrom was dried by using MgSO_4 , and the residue obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 3.63 g of Compound 5 (yield of 72%). The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{43}\text{H}_{28}\text{N}_2$ cal. 572.23, found 572.22.




Synthesis Example 2: Synthesis of Compound 7

[0350] 3.83 g of Compound 7 (yield of 70%) was obtained in the same manner as used to synthesize Compound 5 in Synthesis Example 1, except that 9-phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-carbazole-2-carbonitrile was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{46}\text{H}_{27}\text{N}_3\text{O}$ cal. 621.22, found 621.23.


Synthesis Example 3: Synthesis of Compound 12

[0351]

<Reaction Scheme 2>



-continued



12

Synthesis of Intermediate B-1

[0352] 6.05 g of Intermediate B-1 (yield of 80%) was obtained in the same manner as used to synthesize Intermediate A-1, except that 4-bromophenylboronic acid was used instead of phenylboronic acid. The obtained compound was identified by LC-MS. $\text{C}_{20}\text{H}_{12}\text{NBrO}_2$: $\text{M}+1$ 377.0

Synthesis of Intermediate B-2

[0353] 5.01 g of Intermediate B-2 (yield of 90%) was obtained in the same manner as used to synthesize Intermediate A-2, except that Intermediate B-1 was used instead of Intermediate A-1. The obtained compound was identified by LC-MS. $\text{C}_{20}\text{H}_{14}\text{BrN}$: $\text{M}+1$ 347.0

Synthesis of Intermediate B-3

[0354] 4.38 g of Intermediate B-3 (yield of 70%) was obtained in the same manner as used to synthesize Intermediate A-3, except that Intermediate B-2 was used instead of Intermediate A-2. The obtained compound was identified by LC-MS. $\text{C}_{27}\text{H}_{16}\text{BrN}$: $\text{M}+1$ 433.1

Synthesis of Compound 12

[0355] 4.57 g of Compound 12 (yield of 65%) was obtained in the same manner as used to synthesize Compound 5, except that 2-(3-(dinaphtho[2,1-b:1',2'-d]furan-6-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{53}\text{H}_{31}\text{NO}$ cal. 697.24, found 697.23.

Synthesis Example 4: Synthesis of Compound 19

[0356] 3.98 g of Compound 19 (yield of 60%) was obtained in the same manner as used to synthesize Compound 12 in Synthesis Example 3, except that 4,4,5,5-tetramethyl-2-(10-(naphthalen-1-yl)anthracen-9-yl)-1,3,2-dioxaborolane was used instead of 2-(3-(dinaphtho[2,1-b:1',2'-d]furan-6-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{51}\text{H}_{31}\text{N}$ cal. 657.25, found 657.26.

Synthesis Example 5: Synthesis of Compound 30

[0357] 4.27 g of Compound 30 (yield of 68%) was obtained in the same manner as in Synthesis Example 3, except that, in synthesizing Intermediate B-1, (4-bromonaphthalen-1-yl)boronic acid was used instead of 4-bromophenylboronic acid, and, in synthesizing Compound 12, 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile was used instead of 2-(3-(dinaphtho[2,1-b:1',2'-d]furan-6-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{47}\text{H}_{30}\text{N}_2$ cal. 622.24, found 622.25.

Synthesis Example 6: Synthesis of Compound 47

[0358] 4.23 g (yield of 63%) of Compound 47 was obtained in the same manner as in Synthesis Example 1, except that, in synthesizing Intermediate A-1, 2-naphthaleneboronic acid was used instead of a phenylboronic acid, and, in synthesizing Compound 5, 9-phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-carbazole-2-carbonitrile was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{50}\text{H}_{29}\text{N}_3$ cal. 671.24, found 671.23.

Synthesis Example 7: Synthesis of Compound 56

[0359] 4.36 g of Compound 56 (yield of 70%) was obtained in the same manner as used to synthesize Synthesis Example 1, except that, in synthesizing Intermediate A-1, 2-pyridineboronic acid was used instead of a phenylboronic acid, and, in synthesizing Compound 5, 2-(dinaphtho[2,1-b:1',2'-d]furan-6-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{46}\text{H}_{26}\text{N}_2\text{O}$ cal. 622.20, found 622.19.

Synthesis Example 8: Synthesis of Compound 67

[0360] 3.52 g of Compound 67 (yield of 60%) was obtained in the same manner as in Synthesis Example 1, except that, in synthesizing Intermediate A-1, 4-pyridineboronic acid was used instead of a phenylboronic acid, in synthesizing Intermediate A-3, 4-bromobenzaldehyde was used instead of 3-bromobenzaldehyde, and, in synthesizing Compound 5, 3,3'-(5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-phenylene)dipyridine was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{42}\text{H}_{26}\text{N}_4$ cal. 586.22, found 586.21.

Synthesis Example 9: Synthesis of Compound 79

[0361] 4.51 g of Compound 79 (yield of 67%) was obtained in the same manner as in Synthesis Example 1, except that, in synthesizing Intermediate A-1, 8-quinolineboronic acid was used instead of a phenylboronic acid, and, in synthesizing Compound 5, 9-phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-carbazole-2-carbonitrile was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{49}\text{H}_{28}\text{N}_4$ cal. 672.23, found 672.22.

Synthesis Example 10: Synthesis of Compound 88

[0362] 4.64 g of Compound 88 (yield of 69%) was obtained in the same manner as in Synthesis Example 1, except that, in synthesizing Intermediate A-1, 3-isoquinolineboronic acid was used instead of phenylboronic acid, and, in synthesizing Compound 5, 2-(dinaphtho[2,1-b:1',2'-d]furan-6-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane was used instead of 9,9-dimethyl-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-9H-fluorene-2-carbonitrile. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{50}\text{H}_{28}\text{N}_2\text{O}$ cal. 672.22, found 672.20.

Synthesis Example 11: Synthesis of Compound 89

[0363] Intermediate B-3 (3.91 g, 9.00 mmol) prepared according to Synthesis Example 3 was dissolved in 30 mL of THF, and then, at a temperature of -80°C ., n-butyllithium (3.6 mL, 2.5 M in hexane) was added thereto. After one hour, at the same temperature, chlorodiphenylphosphine (1.99 g, 9 mmol) was added to the resultant mixture. The mixture was stirred at room temperature for 3 hours, followed by adding water thereto, and then, washed three times with diethylether (30 mL). A diethylether layer obtained after the washing was dried by using MgSO_4 , and then, dried under reduced pressure to obtain a product. Then, 30 mL of H_2O_2 was added to the product and stirred for 1 hour. Water was added to the resultant product, which was then washed three times with diethylether (30 mL). A diethylether layer obtained after the washing was dried by using MgSO_4 , and then, dried under reduced pressure to obtain a product, which was then separation-purified by silica gel column chromatography to obtain 3.00 g of Compound 89 (yield of 60%), which was white solid. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{39}\text{H}_{26}\text{NOP}$ cal. 555.18, found 555.19.

Synthesis Example 12: Synthesis of Compound 91

[0364] Intermediate B-3 (4.35 g, 10.0 mmol) prepared according to Synthesis Example 3, diphenyl(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide (4.04 g, 10.0 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.58 g, 0.50 mmol), and K_2CO_3 (4.15 g, 30.0 mmol) were dissolved in 60 mL of a mixed solution including THF/ H_2O (a volumetric ratio of 2/1), and then, stirred at a temperature of 80°C . for 16 hours. The reaction solution was cooled to room temperature, followed by adding with 40 mL of water thereto, and then, washed three times with 50 mL of diethylether. A collected organic layer was dried by using MgSO_4 , and the residue obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 4.29 g (yield of 68%) of Compound 91. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{45}\text{H}_{30}\text{NOP}$ cal. 631.21, found 631.22.

Synthesis Example 13: Synthesis of Compound 103


[0365] 3.45 g of Compound 103 (yield of 67%) was obtained in the same manner as in Synthesis Example 11, except that, in synthesizing Intermediate B-1, (4-bromonaphthalen-1-yl)boronic acid was used instead of 4-bromophenylboronic acid. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{43}\text{H}_{28}\text{NOP}$ cal. 605.19, found 605.18.

Synthesis Example 14: Synthesis of Compound 119


[0366] 4.29 g of Compound 119 (yield of 68%) was obtained in the same manner as in Synthesis Example 12, except that, in synthesizing Compound 91, 4,4,5,5-tetramethyl-2-(10-(phenylsulfonyl)anthracen-9-yl)-1,3,2-dioxaborolane was used instead of diphenyl(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{47}\text{H}_{29}\text{NO}_2\text{S}$ cal. 671.19, found 671.18.

Synthesis Example 15: Synthesis of Compound 129

[0367]



-continued



Synthesis of Intermediate C-1

[0368] 9-bromophenanthrene (2.57 g, 10.0 mmol) was dissolved in 30 mL of THF, and then, at a temperature of -80°C , *n*-butyllithium (4.0 mL, 2.5 M in Hexane) was added thereto. After one hour, at the same temperature, 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.24 mL, 11 mmol) was added thereto. The resultant mixture was stirred at room temperature for 3 hours, followed by adding water thereto, and then, washed three times with 30 mL of diethylether. A diethylether layer obtained after the washing was dried by using MgSO_4 , and then, dried under reduced pressure to obtain a product, which was then separation-purified by silica gel column chromatography to obtain 1.88 g (yield of 62%) of Intermediate C-1. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{20}\text{H}_{31}\text{BO}_2$; $\text{M}+1$ 378.0

Synthesis of Intermediate C-2

[0369] Intermediate C-1 (3.04 g, 10.0 mmol), 1,4-dibromo-2-nitrobenzene (2.81 g, 10.0 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.58 g, 0.50 mmol), tetrabutylammonium bromide (TBAB) (0.161 g, 0.50 mmol), and Na_2CO_3 (3.14 g, 30.0 mmol) were dissolved in 40 mL of a mixed solution including toluene/ethanol/ H_2O (3/3/1), and then, stirred at a temperature of 80°C for 16 hours. The reaction solution was cooled to room temperature, and then, subjected to an extraction process three times by using 60 mL of water and 60 mL of diethylether. An organic layer obtained therefrom was dried by using MgSO_4 , and the residue obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 3.06 g of Intermediate C-2 (yield of 81%). The obtained compound was confirmed by LC-MS. $\text{C}_{20}\text{H}_{12}\text{NBrO}_2$; $\text{M}+1$ 378.0

Synthesis of Intermediate C-3

[0370] Intermediate C-2 (3.78 g, 10.0 mmol), Tin (3.56 g, 30.0 mmol), and HCl 5.0 mL (50 mmol, conc. 36.5%) were dissolved in 40 mL of ethanol, and then, the mixture was stirred at a temperature of 100°C for 8 hours. The reaction solution was cooled to room temperature, and then, 5 g of sodium hydroxide dissolved in 10 mL of water was added to a filtrate obtained by filtering the reaction solution under reduced pressure. The resultant product was subjected to an extraction process three times by using 60 mL of water and 60 mL of dichloromethane. An organic layer obtained there-

from was dried by using MgSO_4 , and the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 2.99 g (yield of 86%) of Intermediate C-3. The obtained compound was confirmed by LC-MS. $\text{C}_{20}\text{H}_{14}\text{BrN}$: $\text{M}+1$ 348.0

Synthesis of Intermediate C-4

[0371] Intermediate C-3 (3.48 g, 10.0 mmol) and 3-bromobenzaldehyde (3.70 g, 20.0 mmol) were dissolved in 20 mL of trifluoroacetic acid, and then, the mixture was stirred in a seal tube at a temperature of 130°C . for 3 days. The reaction solution was cooled to room temperature, followed by quenching using NaHCO_3 . The resultant solution was subjected to an extraction process three times by using 40 mL of water and 60 mL of dichloromethane. An organic layer obtained therefrom was dried by using MgSO_4 , and the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 2.99 g of Intermediate C-4 (yield of 69%). The obtained compound was confirmed by LC-MS. $\text{C}_{27}\text{H}_{16}\text{BrN}$: $\text{M}+1$ 434.1

Synthesis of Compound 129

[0372] Intermediate C-4 (4.34 g, 10.00 mmol) was dissolved in 30 mL of THF, and then, at a temperature of -80°C ., n-butyllithium (4.0 mL, 2.5 M in hexane) was added thereto. After one hour, at the same temperature, chlorodiphenylphosphine (2.20 g, 10 mmol) was added to the mixture. The resultant mixture was stirred at room temperature for 3 hours, followed by adding water thereto, and then, washed three times with diethylether (30 mL). A diethylether layer was dried by using MgSO_4 , and then, dried under reduced pressure to obtain a product. Then, 30 mL of H_2O_2 was added thereto and stirred for 1 hour. After the stirring, water was added thereto, and the resultant product was washed three times with diethylether (30 mL). A diethylether layer obtained after the washing was dried by using MgSO_4 , and then, dried under reduced pressure to obtain a product, which was then separation-purified by silica gel column chromatography to obtain 3.27 g of Compound 129 (yield of 59%). The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{39}\text{H}_{26}\text{NOP}$ cal. 555.18, found 555.19.

Synthesis Example 16: Synthesis of Compound 134

[0373] Intermediate C-4 (4.34 g, 10.0 mmol) prepared according to Synthesis Example 15, diphenyl(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)phosphine oxide (5.04 g, 10.0 mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.58 g, 0.50 mmol), and K_2CO_3 (4.15 g, 30.0 mmol) were dissolved in 60 mL of a mixed solution including THF/ H_2O (a volumetric ratio of 2/1), and then, the mixture was stirred at a temperature of 80°C . for 16 hours. The reaction solution was cooled to room temperature, followed by adding 40 mL of water thereto, and then, subjected to an extraction process three times by using 50 mL of diethylether. A collected organic layer was dried by using MgSO_4 , and the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography, thereby completing the preparation of 4.29 g (yield of 68%) of Compound 134. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{53}\text{H}_{34}\text{NOP}$ cal. 731.24, found 731.23.

Synthesis Example 17: Synthesis of Compound 138

[0374] 4.37 g (yield of 64%) of Compound 138 was obtained in the same manner as in Synthesis Example 16, except that, in synthesizing Intermediate C-2, 3,6-dibromo-2-nitropyridine was used instead of 1,4-dibromo-2-nitrobenzene, and, in synthesizing Compound 134, diphenyl(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)naphthalen-1-yl)phosphine oxide was used instead of diphenyl(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{48}\text{H}_{31}\text{N}_2\text{OP}$ cal. 682.22, found 682.21.

Synthesis Example 18: Synthesis of Compound 143

[0375] 3.88 g (yield of 64%) of Compound 143 was obtained in the same manner as in Synthesis Example 15, except that, in synthesizing Intermediate C-2, 1,4-dibromo-2-nitronaphthalene was used instead of 1,4-dibromo-2-nitrobenzene. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{43}\text{H}_{28}\text{NOP}$ cal. 605.19, found 605.18.

Synthesis Example 19: Synthesis of Compound 146

[0376] 4.43 g of Compound 146 (yield of 65%) was obtained in the same manner as used to synthesize Synthesis Example 16, except that, in synthesizing Intermediate C-2, 1,4-dibromo-2-nitronaphthalene was used instead of 1,4-dibromo-2-nitrobenzene, and, in synthesizing Compound 134, diphenyl(5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-3-yl)phosphine oxide was used instead of diphenyl(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{48}\text{H}_{31}\text{N}_2\text{OP}$ cal. 682.22, found 682.21.

Synthesis Example 20: Synthesis of Compound 153

[0377] 4.17 g of Compound 153 (yield of 61%) was obtained in the same manner as in Synthesis Example 16, except that, in synthesizing Intermediate C-2, 5,8-dibromo-6-nitroquinoline was used instead of 1,4-dibromo-2-nitrobenzene, and, in synthesizing Compound 134, diphenyl(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2-yl)phosphine oxide was used instead of diphenyl(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)phosphine oxide. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{47}\text{H}_{30}\text{N}_3\text{OP}$ cal. 685.22, found 685.23.

Synthesis Example 21: Synthesis of Compound 154

[0378] 4.43 g (yield of 65%) of Compound 154 was obtained in the same manner as in Synthesis Example 16, except that, in synthesizing Intermediate C-2, 5,8-dibromo-7-nitroquinoline was used instead of 1,4-dibromo-2-nitrobenzene, and, in synthesizing Compound 134, diphenyl(3-(4,4,5-trimethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide was used instead of diphenyl(10-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)

phosphine oxide. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{48}\text{H}_{31}\text{N}_2\text{OP}$ cal. 682.22, found 682.21.

Synthesis Example 22: Synthesis of Compound 168

[0379] 4.12 g of Compound 168 (yield of 66%) was obtained in the same manner as in Synthesis Example 16, except that, in synthesizing Intermediate C-2, 5,8-dibromo-6-nitroquinoline was used instead of 1,4-dibromo-2-nitrobenzene, and, in synthesizing Compound 134, 2-(phenylsulfonyl)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine was used instead of diphenyl(10-(4,4,5,5-

tetramethyl-1,3,2-dioxaborolan-2-yl)anthracen-9-yl)phosphine oxide. The obtained compound was identified by MS/FAB and ^1H NMR. $\text{C}_{41}\text{H}_{25}\text{N}_3\text{O}_2\text{S}$ cal. 623.17, found 623.18.

[0380] Additional compounds were synthesized by using the same synthesis paths and methods as described above and appropriate intermediate materials. Table 1 shows ^1H NMR and MS/FAB of these compounds are shown in Table 1.

[0381] Synthesis methods for other compounds than the compounds shown in Table 1 may be understood by one of ordinary skill in the art by referring to the synthesis paths and source materials described above.

TABLE 1

Compound	^1H NMR (CDCl_3 , 400 MHz) δ	MS/FAB	
		found	calc.
5	$\delta = 9.23$ (d, 1H), 8.87 (d, 1H), 8.69-8.65 (m, 2H), 8.36-8.34 (m, 1H), 8.22 (d, 1H), 7.92-7.76 (m, 7H), 7.70-7.64 (m, 3H), 7.60-7.45 (m, 4H), 7.36 (t, 1H), 7.12-7.08 (m, 1H), 1.65 (s, 6H)	572.22	572.23
7	$\delta = 9.22$ (d, 1H), 8.86 (d, 1H), 8.69-8.66 (m, 2H), 8.57-8.55 (m, 1H), 8.23-8.21 (m, 2H), 8.08 (d, 1H), 8.03 (d, 1H), 7.92-7.76 (m, 7H), 7.70-7.48 (m, 8H), 7.41 (t, 1H), 7.32-7.28 (m, 1H), 7.13-7.09 (m, 1H)	621.23	621.22
12	$\delta = 9.23$ -9.19 (m, 1H), 8.83 (t, 2H), 8.69-8.63 (m, 4H), 8.33 (s, 1H), 8.06-7.90 (m, 7H), 7.86-7.48 (m, 15H), 7.09 (t, 1H)	697.23	697.24
19	$\delta = 9.22$ -9.20 (m, 1H), 8.82, 8.73-8.65 (m, 4H), 8.18 (d, 1H), 7.98-7.96 (m, 2H), 7.86-7.79 (m, 6H), 7.72-7.44 (m, 10H), 7.40-7.27 (m, 5H), 7.09 (t, 1H), 6.96 (t, 1H)	657.26	657.25
30	$\delta = 9.27$ (d, 1H), 8.86 (d, 1H), 8.70 (d, 1H), 8.67 (d, 1H), 8.28 (s, 1H), 8.20-8.18 (m, 2H), 8.06-8.02 (m, 2H), 7.95-7.79 (m, 5H), 7.74-7.45 (m, 8H), 7.23 (t, 1H), 7.07 (t, 1H), 1.66 (s, 6H)	622.25	622.24
47	$\delta = 9.29$ -9.26 (m, 1H), 8.82-8.76 (m, 2H), 8.70-8.64 (m, 3H), 8.23, 8.16-8.03 (m, 5H), 7.94 (d, 1H), 7.89-7.79 (m, 6H), 7.70-7.45 (m, 8H), 7.42 (t, 1H), 7.32-7.28 (m, 1H), 7.13 (t, 1H)	671.23	671.24
56	$\delta = 9.31$ -9.28 (m, 1H), 9.01 (d, 1H), 8.89-8.76 (m, 5H), 8.60-8.56 (m, 2H), 8.39 (s, 1H), 7.96-7.87 (m, 6H), 7.83-7.78 (m, 3H), 7.72-7.48 (m, 6H), 7.37 (t, 1H)	622.19	622.20
67	$\delta = 9.52$ -9.50 (m, 1H), 9.23-9.21 (m, 1H), 8.94 (d, 2H), 8.81 (d, 1H), 8.69-8.65 (m, 4H), 8.46 (d, 1H), 8.38-8.35 (m, 2H), 8.06-8.00 (m, 3H), 7.95-7.79 (m, 8H), 7.54-7.46 (m, 3H)	586.21	586.22
79	$\delta = 9.53$ -9.51 (m, 1H), 9.18 (d, 1H), 8.80-8.70 (m, 3H), 8.46 (d, 1H), 8.27 (d, 1H), 8.23 (d, 1H), 8.08 (d, 1H), 8.04-7.94 (m, 3H), 7.86-7.76 (m, 5H), 7.70-7.48 (m, 8H), 7.41 (t, 1H), 7.31-7.28 (m, 1H), 7.16 (t, 1H)	672.22	672.23
88	$\delta = 9.55$ (d, 1H), 9.37-8.34 (m, 1H), 8.92 (d, 1H), 8.85-8.76 (m, 3H), 8.67-8.63 (m, 2H), 8.39 (d, 1H), 8.29 (dd, 1H), 8.19 (d, 1H), 8.01-7.76 (m, 11H), 7.72-7.48 (m, 6H)	672.20	672.22
89	$\delta = 9.23$ -9.16(m, 2H), 8.74-8.66(m, 3H), 7.95-7.93(m, 1H), 7.85-7.81(m, 5H), 7.69-7.61(m, 7H), 7.55-7.47(m, 3H), 7.45-7.39(m, 4H), 7.12-7.07(m, 1H)	555.19	555.18
91	$\delta = 9.23$ -9.21(m, 1H), 8.71-8.65(m, 2H), 8.52-8.50(m, 1H), 8.37-8.36(m, 1H), 8.13-8.11(m, 1H), 7.98-7.97(m, 2H), 7.87-7.81(m, 4H), 7.68-7.61(m, 8H), 7.57-7.48(m, 5H), 7.44-7.41(m, 4H), 7.12-7.08(m, 1H)	631.22	631.21
103	$\delta = 9.29$ -9.24(m, 2H), 8.96-8.94(m, 1H), 8.78-8.71(m, 2H), 8.67-8.66(m, 1H), 8.01-8.00(m, 2H), 7.85-7.81(m, 2H), 7.74-7.72(m, 4H), 7.68-7.66(m, 2H), 7.63-7.53(m, 5H), 7.51-7.49(m, 1H), 7.41-7.39(m, 4H), 7.08-7.06(m, 1H)	605.18	605.19
119	$\delta = 9.76$ -9.74(m, 1H), 9.26-9.22(m, 2H), 8.85-8.68(m, 4H), 8.06-8.01(m, 2H), 7.96-7.93(m, 3H), 7.80-7.63(m, 12H), 7.54-7.52(m, 2H), 7.44-7.42(m, 4H)	671.18	671.19
129	$\delta = 8.84$ -8.83(m, 2H), 8.71-8.61(m, 4H), 8.00-7.99(m, 2H), 7.84-7.74(m, 3H), 7.70-7.68(m, 5H), 7.64-7.62(m, 4H), 7.49-7.42(m, 5H)	555.19	555.18
134	$\delta = 8.84$ -8.83(m, 2H), 8.71-8.62(m, 3H), 8.20-8.18(m, 2H), 8.04-8.00(m, 3H), 7.83-7.68(m, 4H), 7.64-7.62(m, 8H), 7.54-7.45(m, 6H), 7.39-7.37(m, 4H), 7.19-7.16(m, 2H)	731.23	731.24

TABLE 1-continued

Compound	¹ H NMR (CDCl ₃ , 400 MHz) δ	MS/FAB	
		found	calc.
138	δ = 8.93-8.85(m, 4H), 8.73-8.71(m, 1H), 8.59-8.55(m, 2H), 8.01-7.92(m, 4H), 7.79-7.72(m, 10H), 7.64-7.63(m, 2H), 7.53-7.51(m, 3H), 7.45-7.39(m, 5H)	682.21	682.22
143	δ = 8.92-8.90(m, 2H), 8.85-8.69(m, 5H), 8.02-8.00(m, 2H), 7.80-7.69(m, 8H), 7.65-7.63(m, 4H), 7.55-7.53(m, 3H), 7.40-7.38(m, 4H)	605.18	605.19
146	δ = 8.93-8.85(m, 4H), 8.79-8.77(m, 2H), 8.66-8.59(m, 2H), 8.40-8.39(m, 1H), 8.02-7.97(m, 3H), 7.83-7.80(m, 6H), 7.74-7.69(m, 2H), 7.64-7.63(m, 4H), 7.50-7.48(m, 3H), 7.41-7.39(m, 4H)	682.21	682.22
153	δ = 9.76-9.74(m, 1H), 9.26-9.22(m, 2H), 8.85-8.68(m, 4H), 8.06-8.01(m, 2H), 7.96-7.93(m, 3H), 7.80-7.63(m, 12H), 7.54-7.52(m, 2H), 7.44-7.42(m, 4H)	685.23	685.22
154	δ = 9.06-9.02(m, 2H), 8.94-8.79(m, 3H), 8.65-8.63(m, 1H), 8.39-8.38(m, 1H), 8.22-8.21(m, 1H), 8.02-7.93(m, 3H), 7.80-7.74(m, 3H), 7.65-7.63(m, 9H), 7.57-7.56(m, 1H), 7.50-7.48(m, 3H), 7.44-7.41(m, 4H)	682.21	682.22
168	δ = 9.76-9.74(m, 2H), 9.26-9.25(m, 1H), 8.85-8.77(m, 3H), 8.70-8.69(m, 1H), 8.40-8.38(m, 1H), 8.15-8.14(m, 1H), 8.05-8.02(m, 4H), 7.89-7.80(m, 3H), 7.74-7.69(m, 2H), 7.65-7.55(m, 7H)	623.18	623.17

Example 1


[0382] As a substrate and an anode, a Corning 150/cm² (1200 Å) ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm, and then, sonicated with isopropyl alcohol and pure water, each for 5 minutes, and then, cleaned by exposure to ultraviolet light for 30 minutes, and then, to ozone. The resultant glass substrate was mounted on a vacuum deposition apparatus.

[0383] 2-TNATA was vacuum-deposited on the ITO anode on the glass substrate to form a hole injection layer having a thickness of 600 Å, and then, 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPB) was vacuum-deposited on the


hole injection layer to form a hole transport layer having a thickness of 300 Å.

[0384] 9,10-di-naphthalene-2-yl-anthracene (ADN), which is a host, and 4,4'-bis[2-(4-(N,N-diphenylamino)phenyl)vinyl]biphenyl (DPAVBi), which is a dopant, were co-deposited at a weight ratio of 98:2 on the hole transport layer to form an emission layer having a thickness of 300 Å.

[0385] Compound 5 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, LiF was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form a cathode having a thickness of 3,000 Å, thereby completing manufacture of an organic light-emitting device.



2-TNATA



Examples 2 to 22 and Comparative Examples 1 and 2

[0386] Organic light-emitting devices were manufactured in the same manner as in Example 1, except that, in forming the electron transport layer, compounds shown in Table 2 were used instead of Compound 1.

Evaluation Example 1

[0387] The driving voltage, current density, luminance, efficiency, emission color, and half lifespan of the organic light-emitting devices manufactured according to Examples 1 to 22 and Comparative Examples 1 and 2 were measured by using Keithley SMU 236 and luminance meter PR650. Results thereof are shown in Table 2.

TABLE 2

	Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
Example 1	Compound 5	3.60	50	3,425	7.20	Blue	618
Example 2	Compound 7	3.42	50	3,755	8.12	Blue	625
Example 3	Compound 12	3.54	50	3,500	7.53	Blue	656
Example 4	Compound 19	3.38	50	3,615	8.07	Blue	602
Example 5	Compound 30	3.47	50	3,830	8.27	Blue	635
Example 6	Compound 47	3.33	50	3,550	7.75	Blue	620
Example 7	Compound 56	3.46	50	3,745	8.03	Blue	635
Example 8	Compound 67	3.50	50	3,570	7.55	Blue	662
Example 9	Compound 79	3.47	50	3,600	8.10	Blue	627
Example 10	Compound 88	3.52	50	3,815	8.16	Blue	640
Example 11	Compound 89	3.60	50	3,425	6.85	Blue	618
Example 12	Compound 91	3.42	50	3,755	7.51	Blue	625
Example 13	Compound 103	3.54	50	3,500	7.00	Blue	656
Example 14	Compound 119	3.38	50	3,615	7.23	Blue	602
Example 15	Compound 129	3.47	50	3,830	7.66	Blue	635

TABLE 2-continued

	Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
Example 16	Compound 134	3.34	50	3,727	7.45	Blue	627
Example 17	Compound 138	3.41	50	3,750	7.50	Blue	637
Example 18	Compound 143	3.37	50	3,627	7.25	Blue	608
Example 19	Compound 146	3.52	50	3,635	7.27	Blue	650
Example 20	Compound 153	3.59	50	3,487	6.97	Blue	591
Example 21	Compound 154	3.42	50	3,562	7.12	Blue	613
Example 22	Compound 168	3.48	50	3,612	7.22	Blue	647
Comparative Example 1	Compound A	5.06	50	3,010	6.52	Blue	325
Comparative Example 2	Compound B	4.88	50	2,948	5.89	Blue	411






TABLE 2-continued

Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
----------	---------------------	---------------------------------------	--------------------------------	-------------------	----------------	---






TABLE 2-continued




Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
						
67						
						
79						
						
88						

TABLE 2-continued





Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
						
89						
						
91						
						
103						
						
119						

TABLE 2-continued





Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
						
129						
						
134						
						
138						
						
143						

TABLE 2-continued








Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
						
146						
						
153						
						
154						
						
168						

TABLE 2-continued

Material	Driving voltage (V)	Current density (mA/cm ²)	Luminance (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm ²)
						
						
						

[0388] Referring to Table 2, it was confirmed that the organic light-emitting devices of Examples 1 to 22 had better characteristics than those of Comparative Examples 1 and 2, in terms of a driving voltage, luminance, efficiency, and a half lifespan.


[0389] An organic light-emitting device including the condensed cyclic compound according to an embodiment may have a low driving voltage, high luminance, high efficiency, and a long lifespan.

[0390] It should be understood that embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments.

[0391] While one or more embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims.

What is claimed is:


1. A condensed cyclic compound represented by Formula 1:




wherein, in Formula 1,
ring A is selected from a C₅-C₃₀ carbocyclic ring and a C₂-C₃₀ heterocyclic ring,

X₁ is C(R₁) or N, X₂ is C(R₂) or N, at least one of X₁ and X₂ is N,


R₁ to R₁₁ are each independently a group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, substituted or unsubstituted C₁-C₆₀ alkyl group, substituted or unsubstituted C₂-C₆₀ alkenyl group, substituted or unsubstituted C₂-C₆₀ alkynyl group, substituted or unsubstituted C₁-C₆₀ alkoxy group, substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₆-C₆₀ aryloxy group, substituted or unsubstituted C₆-C₆₀ arylthio group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂):




-continued




<Formula 2-2>



<Formula 2-3>



<Formula 2-4>



<Formula 2-5>

L_1 to L_{11} are each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, substituted or unsubstituted C_6 - C_{60} arylene group, substituted or unsubstituted C_1 - C_{60} heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$a1$ to $a11$ are each independently 0, 1, 2, 3, 4, or 5,

Ar_1 to Ar_7 are selected from a substituted or unsubstituted C_1 - C_{60} alkyl group, substituted or unsubstituted C_2 - C_{60} alkenyl group, substituted or unsubstituted C_2 - C_{60} alkynyl group, substituted or unsubstituted C_1 - C_{60} alkoxy group, substituted or unsubstituted C_3 - C_{10} cycloalkyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, substituted or unsubstituted C_6 - C_{60} aryl group, substituted or unsubstituted C_1 - C_{60} heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

$b1$ to $b4$ are each independently 1, 2, or 3,

$c3$ is 0, 1, 2, 3, 4, 5, or 6,

at least one of substituents of the substituted C_3 - C_{10} cycloalkylene group, substituted C_1 - C_{10} heterocycloalkylene group, substituted C_3 - C_{10} cycloalkenylene group, substituted C_1 - C_{10} heterocycloalkenylene group, substituted C_6 - C_{60} arylene group, substituted C_1 - C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substi-

tuted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q_{11})(Q_{12})(Q_{13}), -N(Q_{11})(Q_{12}), -B(Q_{11})(Q_{12}), -C(=O)(Q_{11}), -S(=O)₂(Q_{11}), and -P(=O)(Q_{11})(Q_{12});

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q_{21})(Q_{22})(Q_{23}), -N(Q_{21})(Q_{22}), -B(Q_{21})(Q_{22}), -C(=O)(Q_{21}), -S(=O)₂(Q_{21}), and -P(=O)(Q_{21})(Q_{22}); and

-Si(Q_{31})(Q_{32})(Q_{33}), -N(Q_{31})(Q_{32}), -B(Q_{31})(Q_{32}), -C(=O)(Q_{31}), -S(=O)₂(Q_{31}), and -P(=O)(Q_{31})(Q_{32}), wherein Q_1 to Q_3 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60}

aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

* indicates a binding site to a neighboring atom.

2. The condensed cyclic compound of claim 1, wherein ring A is selected from a benzene group, a pyridine group, a pyridazine group, a pyrimidine group, a pyrazine group, a naphthalene group, a quinoline group, an isoquinoline group, a cinnoline group, a quinazoline group, a quinoxaline group, naphthyridine group, an anthracene group, a phenanthrene group, a benzoquinoline group, a phenanthridine group, an acridine group, phenanthroline group, and a phenazine group.

3. The condensed cyclic compound of claim 1, wherein L_1 to L_{11} are each independently selected from

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiofenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiofenylene group, a benzocarbazolylene group, a dibenzosilolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylene group,


a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiofenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiofenylene group, a benzocarbazolylene group, a dibenzosilolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiofenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrim-

idiny1 group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$.


wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.

4. The condensed cyclic compound of claim 1, wherein


L_1 to L_{11} are each independently selected from groups represented by Formulae 3-1 to 3-25:




Formular 3-1




Formular 3-2




Formular 3-3




Formular 3-4




Formular 3-5



Formular 3-6




Formular 3-7




Formular 3-8


-continued




Formular 3-9




Formular 3-10




Formular 3-11




Formular 3-12



Formular 3-13











Formular 3-14



Formular 3-15

-continued



Formular 3-16

Formular 3-17

Formular 3-18

Formular 3-19



Formular 3-20

Formular 3-21

Formular 3-22

Formular 3-23

-continued



Formular 3-24

Formular 3-25

wherein, in Formulae 3-1 to 3-25,

Y_1 is O, S, $C(Z_5)(Z_6)$, $N(Z_7)$, or $Si(Z_8)(Z_9)$,

Z_1 to Z_9 are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzo-fluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}), wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

d_3 is an integer from 1 to 3,

d_4 is an integer from 1 to 4,


d_6 is an integer from 1 to 6,

d_8 is an integer from 1 to 8, and


* and *' each indicate a binding site to a neighboring atom.

5. The condensed cyclic compound of claim 1, wherein L_1 to L_{11} are each independently selected from groups represented by Formulae 4-1 to 4-19:


Formular 4-1













Formular 4-2



Formular 4-3



-continued



Formular 4-4

Formular 4-5

Formular 4-6

Formular 4-7

Formular 4-8

Formular 4-9







Formular 4-10

Formular 4-11

Formular 4-12

Formular 4-13

-continued



Formular 4-14

Formular 4-15

Formular 4-16

Formular 4-17

Formular 4-18

Formular 4-19

wherein, in Formulae 4-1 to 4-19, * and *' each indicate a binding site to a neighboring atom.


6. The condensed cyclic compound of claim 1, wherein Ar_1 to Ar_7 are each independently selected from:

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzonaphthothiophenyl group, a dinaphthothiophenyl group, a benzonaphthosilolyl group, and a dinaphthosilolyl group; and


a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group,

a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzonaphthothiophenyl group, a dinaphthothiophenyl group, a benzonaphthosilolyl group, and a dinaphthosilolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzosilolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, a benzonaphthofuranyl group, a dinaphthofuranyl group, a benzonaphthothiophenyl group, a dinaphthothiophenyl group, a benzonaphthosilolyl group, a dinaphthosilolyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}) and —P(=O)(Q_{31})(Q_{32}), wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.


7. The condensed cyclic compound of claim 1, wherein Ar_1 to Ar_7 are each independently selected from groups represented by Formulae 5-1 to 5-25:




Formular 5-1




Formular 5-2




Formular 5-3




Formular 5-4




Formular 5-5




Formular 5-6




Formular 5-7




Formular 5-8



Formular 5-9




Formular 5-10




Formular 5-11


-continued




Formular 5-12




Formular 5-13




Formular 5-14




Formular 5-15




Formular 5-16



Formular 5-17




Formular 5-18




Formular 5-19


-continued




Formular 5-20




Formular 5-21




Formular 5-22



Formular 5-23



Formular 5-24



Formular 5-25

wherein, in Formulae 5-1 to 5-25,
 Y_{31} is O, S, C(Z_{33})(Z_{34}), N(Z_{35}), or Si(Z_{36})(Z_{37}),
 Z_{31} to Z_{37} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a

pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}), wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

e_3 is an integer from 1 to 3,

e_4 is an integer from 1 to 4,

e_5 is an integer from 1 to 5,


e_6 is an integer from 1 to 6,

e_7 is an integer from 1 to 7,


e_9 is an integer from 1 to 9, and

* indicates a binding site to a neighboring atom.


8. The condensed cyclic compound of claim 1, wherein Ar_1 to Ar_7 are each independently selected from groups represented by Formulae 6-1 to 6-18:




Formular 6-1




Formular 6-2




Formular 6-3



Formular 6-4




Formular 6-5



Formular 6-6










Formular 6-7



Formular 6-8

-continued



Formular 6-9

Formular 6-10

Formular 6-11




Formular 6-12

Formular 6-13

Formular 6-14

Formular 6-15

-continued



Formular 6-16

Formular 6-17

Formular 6-18

wherein, in Formulae 6-1 to 6-18, * indicates a binding site to a neighboring atom.

9. The condensed cyclic compound of claim 1, wherein R_1 to R_{11} are each independently selected from

a group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thi-

azolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}); and

—Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), —B(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2),

wherein Q_1 to Q_3 and Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.


10. The condensed cyclic compound of claim 1, wherein R_1 to R_{11} are each independently selected from:

a group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, —Si(Q_1)(Q_2)(Q_3), —N(Q_1)(Q_2), —C(=O)(Q_1), —S(=O)₂(Q_1), and —P(=O)(Q_1)(Q_2),


wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

11. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by one of Formulae 1A to 1 S:


Formula 1A




Formula 1B




Formula 1C



Formula 1D




-continued




Formula 1E


-continued




Formula 1J




Formula 1F




Formula 1K




Formula 1G




Formula 1L




Formula 1H



Formula 1M




Formula 1I




Formula 1N

-continued


Formula 1O




Formula 1P




Formula 1Q



Formula 1R



Formula 1S



wherein, in Formulae 1A to 1S,

R_{31} to R_{36} are the same as described in connection with R_3 , and

X_1 , X_2 , and R_3 to R_{11} are the same as described in claim 1.

12. The condensed cyclic compound of claim 11, wherein, in Formulae 1A to 1S,

X_1 is N, and X_2 is $C(R_2)$, and

R_2 is selected from groups represented by Formulae 2-1 to 2-5.

13. The condensed cyclic compound of claim 11, wherein, in Formulae 1A to 1S,

X_1 is $C(R_1)$, X_2 is N, and

R_1 is selected from groups represented by Formulae 2-1 to 2-5.


14. The condensed cyclic compound of claim 11, wherein, in Formulae 1A, 1B, 1D to 1G, 1I to 1N, and 1P to 1S, X_1 is N, X_2 is $C(R_2)$, and

R_{32} is selected from groups represented by Formulae 2-1 to 2-5.


15. The condensed cyclic compound of claim 11, wherein, in Formulae 1A, 1B, 1D to 1G, 1I to 1N, and 1P to 1S, X_1 is $C(R_1)$, X_2 is N, and

R_{32} is selected from groups represented by Formulae 2-1 to 2-5.

16. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is one of the following Compounds 1 to 192:




-continued



-continued


2

5






3

6







4

-continued




-continued




-continued


14




15



16




17




-continued


18




19




20






21







-continued



-continued




-continued




-continued


34




35



36




37




-continued


38




39



40




-continued




41


-continued




44



42




43




45

-continued




46


-continued




48



49




50




51

-continued




51

-continued




53




52



54




-continued



-continued


55

58




56

59




57

60




-continued




61


-continued




62



63




64



65

-continued




66

67

68




-continued



69

70


-continued



-continued


71

74




75

72




73

76




-continued




77

-continued




79


80




78



81




-continued




82

-continued




83


85




84



86




-continued




87

-continued




90

91





88

92




93

89




-continued


94




95



96




97




-continued


98




99



100




101




-continued


102




103



104




105




-continued


106




107



108




109




-continued


110




111



112




113




-continued


114




115



116




117




-continued


118




119



120




121




-continued


122




123



124




125




-continued


126




127



128




129




-continued


130




131




132



133




-continued




134


135




136



137




-continued




138


139







140







141



-continued




-continued




-continued


150




151



152




153




-continued


154




155




156



157




-continued




158


159




160



161




-continued




162

163




164



165


-continued

166




-continued


169




167




170



168




171




145

-continued




172


-continued




175




173



176




174




177

-continued




178


-continued




181




179



182




180






183

-continued




-continued



17. An organic light-emitting device comprising: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer is at least one condensed cyclic compound of Formula 1:

190



<Formula 1>

ring A is selected from a C₅-C₃₀ carbocyclic ring and a C₂-C₃₀ heterocyclic ring,


X₁ is C(R₁) or N, X₂ is C(R₂) or N, at least one of X₁ and X₂ is N,

191


R₁ to R₁₁ are each independently a group represented by one of Formulae 2-1 to 2-5, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, substituted or unsubstituted C₁-C₆₀ alkyl group, substituted or unsubstituted C₂-C₆₀ alkenyl group, substituted or unsubstituted C₂-C₆₀ alkynyl group, substituted or unsubstituted C₁-C₆₀ alkoxy group, substituted or unsubstituted C₃-C₁₀ cycloalkyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, substituted or unsubstituted C₆-C₆₀ aryl group, substituted or unsubstituted C₆-C₆₀ aryloxy group, substituted or unsubstituted C₆-C₆₀ arylthio group, substituted or unsubstituted C₁-C₆₀ heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂):

192


<Formula 2-1>




<Formula 2-2>



<Formula 2-3>



-continued



L_1 to L_{11} are each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, substituted or unsubstituted C_6 - C_{60} arylene group, substituted or unsubstituted C_1 - C_{60} heteroarylene group, substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$a1$ to $a11$ are each independently 0, 1, 2, 3, 4, or 5,

Ar_1 to Ar_7 are selected from a substituted or unsubstituted C_1 - C_{60} alkyl group, substituted or unsubstituted C_2 - C_{60} alkenyl group, substituted or unsubstituted C_2 - C_{60} alkynyl group, substituted or unsubstituted C_1 - C_{60} alkoxy group, substituted or unsubstituted C_3 - C_{10} cycloalkyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, substituted or unsubstituted C_6 - C_{60} aryl group, substituted or unsubstituted C_1 - C_{60} heteroaryl group, substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

$b1$ to $b4$ are each independently 1, 2, or 3,

$c3$ is 0, 1, 2, 3, 4, 5, or 6,

at least one of substituents of the substituted C_3 - C_{10} cycloalkylene group, substituted C_1 - C_{10} heterocycloalkylene group, substituted C_3 - C_{10} cycloalkenylene group, substituted C_1 - C_{10} heterocycloalkenylene group, substituted C_6 - C_{60} arylene group, substituted C_1 - C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group, and substituted monovalent non-aromatic condensed heteropolycyclic group is selected from

deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group;

a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, and a C_1 - C_{60} alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q_{11})(Q_{12})(Q_{13}), -N(Q_{11})(Q_{12}), -B(Q_{11})(Q_{12}), -C(=O)(Q_{11}), -S(=O)₂(Q_{11}), and -P(=O)(Q_{11})(Q_{12});

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q_{21})(Q_{22})(Q_{23}), -N(Q_{21})(Q_{22}), -B(Q_{21})(Q_{22}), -C(=O)(Q_{21}), -S(=O)₂(Q_{21}), and -P(=O)(Q_{21})(Q_{22}); and

-Si(Q_{31})(Q_{32})(Q_{33}), -N(Q_{31})(Q_{32}), -B(Q_{31})(Q_{32}), -C(=O)(Q_{31}), -S(=O)₂(Q_{31}), and -P(=O)(Q_{31})(Q_{32}),

wherein Q_1 to Q_3 , Q_{11} to Q_{13} , Q_{21} to Q_{23} , and Q_{31} to Q_{33} are each independently selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

* indicates a binding site to a neighboring atom.

- 18.** The organic light-emitting device of claim **17**, wherein
the first electrode is an anode,
the second electrode is a cathode,
the organic layer includes a hole transport region between
the first electrode and the emission layer and an electron
transport region between the emission layer and
the second electrode,
the hole transport region includes at least one selected
from a hole injection layer, a hole transport layer, an
emission auxiliary layer, and an electron blocking
layer, and
the electron transport region includes at least one selected
from a buffer layer, a hole blocking layer, an electron
control layer, an electron transport layer, and an electron
injection layer.
- 19.** The organic light-emitting device of claim **18**, wherein
the electron transport region includes an electron transport
layer, and the electron transport layer includes the at
least one condensed cyclic compound.
- 20.** The organic light-emitting device of claim **18**, wherein
the electron transport region includes an electron injection
layer, and the electron injection layer includes the at
least one condensed cyclic compound.

* * * * *

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	US20170179407A1	公开(公告)日	2017-06-22
申请号	US15/371066	申请日	2016-12-06
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	PARK JUNHA KIM YOUNGKOOK SIM MUNKI LEE EUNYOUNG LEE HYOYOUNG JEONG EUNJAE HWANG SEOKHWAN		
发明人	PARK, JUNHA KIM, YOUNGKOOK SIM, MUNKI LEE, EUNYOUNG LEE, HYOYOUNG JEONG, EUNJAE HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C07D471/04 C07F9/576 C07D401/10 C07D221/18 C07D405/10		
CPC分类号	H01L51/0072 C07D221/18 H01L51/0052 C07D405/10 H01L51/0073 C07F9/5765 H01L51/5092 C07D401/10 C07D471/04 H01L51/0067 H01L51/5012 H01L51/5016 H01L51/5072 H01L51/0058 C07F9/65583 C07F9/6561 H01L51/5096		
优先权	1020150182792 2015-12-21 KR 1020160046499 2016-04-15 KR		
外部链接	Espacenet USPTO		

摘要(译)

提供缩合环状化合物和有机发光器件。缩合环状化合物由式1表示：公开了关于式1的组分的细节。

10

190

150

110