



(19) **United States**

(12) **Patent Application Publication**
LEE

(10) **Pub. No.: US 2018/0033974 A1**
(43) **Pub. Date: Feb. 1, 2018**

(54) **POLYCYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(52) **U.S. Cl.**
CPC *H01L 51/0067* (2013.01); *C09K 11/06* (2013.01); *C09K 11/025* (2013.01); *H01L 51/0085* (2013.01); *C07D 401/10* (2013.01); *H01L 51/0072* (2013.01); *H01L 51/0052* (2013.01); *C07D 491/06* (2013.01); *H01L 51/0071* (2013.01); *C07D 495/06* (2013.01); *H01L 51/5072* (2013.01)

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-si (KR)

(72) Inventor: **Jungsub LEE**, Yongin-si (KR)

(21) Appl. No.: **15/437,504**

(22) Filed: **Feb. 21, 2017**

(30) **Foreign Application Priority Data**

Jul. 27, 2016 (KR) 10-2016-0095488

Publication Classification

(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 495/06 (2006.01)
C07D 401/10 (2006.01)
C07D 491/06 (2006.01)
C09K 11/06 (2006.01)
C09K 11/02 (2006.01)

(57) **ABSTRACT**
A polycyclic compound and an organic light-emitting device, the compound being represented by Formula 1:

[Formula 1]






FIG. 1



**POLYCYCLIC COMPOUND AND ORGANIC
LIGHT-EMITTING DEVICE INCLUDING
THE SAME**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] Korean Patent Application No. 10-2016-0095488, filed on Jul. 27, 2016, in the Korean Intellectual Property Office, and entitled: "Polycyclic Compound and Organic Light-Emitting Device Including the Same," is incorporated by reference herein in its entirety.

BACKGROUND

1. Field

[0002] Embodiments relate to a polycyclic compound and an organic light-emitting device including the same.

2. Description of the Related Art


[0003] Organic light-emitting devices are self-emission devices, have wide viewing angles, high contrast ratios, short response times, and excellent brightness, driving voltage, and response speed characteristics, compared to conventional devices, and produce full-color images.

[0004] An example of such organic light-emitting devices may include a first electrode disposed on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially disposed on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, recombine in the emission layer to produce excitons. These excitons transition from an excited state to a ground state, thereby generating light.

SUMMARY

[0005] Embodiments are directed to a polycyclic compound and an organic light-emitting device including the same.


[0006] The embodiments may be realized by providing a polycyclic compound represented by Formula 1:



[Formula 1]

[0007] wherein, in Formula 1, X₁ is C(R₁₁)(R₁₂), N(R₁₃), O or S, and R₁ to R₁₃ are each independently a structure represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group,

a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), or —P(=O)(Q₁)(Q₂), at least one of R₁ to R₁₀ having a structure represented by Formula 2,



[Formula 2]

[0008] wherein, in Formulae 1 and 2, L₁ is a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group, Ar₁ is a substituted or unsubstituted C₁-C₆₀ heterocyclic group having at least one *—N=* moiety, a₁ is an integer from 1 to 5, b₁ is an integer from 1 to 5, when a₁ is two or more, two or more L₁(s) are identical to or different from each other, when b₁ is two or more, two or more Ar₁(s) are identical to or different from each other, Ar₂ is a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, or a substituted or unsubstituted C₁-C₆₀ heteroaryl group, and R₁₃ and Ar₂ are separate or are linked to a neighboring substituent to form a substituted or unsubstituted ring, * and *' each indicate a binding site to a neighboring atom, and at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substi-

tuted C₁-C₆₀ heterocyclic group, or the substituted ring is: deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a diazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group; a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, —Si(Q_{ii})(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), or —P(=O)(Q₁₁)(Q₁₂); a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, a biphenyl group, or a terphenyl group; a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), or —P(=O)(Q₂₁)(Q₂₂); or —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), or —P(=O)(Q₃₁)(Q₃₂), wherein Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryl group substituted with a C₁-C₆₀ alkyl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, a biphenyl group, or a terphenyl group.

[0009] The embodiments may be realized by providing an organic light-emitting device including a first electrode; a second electrode facing the first electrode; and an organic

layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes the polycyclic compound according to an embodiment.

BRIEF DESCRIPTION OF THE DRAWING

[0010] Features will be apparent to those of skill in the art by describing in detail exemplary embodiments with reference to the attached drawings in which:

[0011] FIG. 1 illustrates a schematic view of an organic light-emitting device according to an example embodiment.


DETAILED DESCRIPTION

[0012] Example embodiments will now be described more fully hereinafter with reference to the accompanying drawing; however, they may be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey exemplary implementations to those skilled in the art.


[0013] In the drawing figure, the dimensions of layers and regions may be exaggerated for clarity of illustration. It will also be understood that when a layer or element is referred to as being “on” another layer or element, it can be directly on the other layer or element, or intervening layers may also be present. In addition, it will also be understood that when a layer is referred to as being “between” two layers, it can be the only layer between the two layers, or one or more intervening layers may also be present. Like reference numerals refer to like elements throughout.

[0014] A polycyclic compound according to an embodiment may be represented by Formula 1.

<Formula 1>



[Formula 2]



[0015] X₁ in Formula 1 may be one selected from, e.g., C(R₁₁)(R₁₂), N(R₁₃), O, and S.

[0016] In one or more embodiments, X₁ may be C(R₁₁)(R₁₂).

[0017] In one or more embodiments, X_i may be N(R₁₃).

[0018] In one or more embodiments, X₁ may be O or S.

[0019] R₁ to R₁₃ in Formula 1 may each independently be selected from, e.g., a structure represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy

group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), and —P(=O)(Q₁)(Q₂). In an implementation, at least one selected from R₁ to R₁₀ may have a structure represented by Formula 2.

[0020] In one or more embodiments, R₁₁ and R₁₂ in Formula 1 may each independently be selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, and a substituted or unsubstituted C₁-C₆₀ heteroaryl group.

[0021] In one or more embodiments, R₁₁ and R₁₂ in Formula 1 may each independently be selected from:

[0022] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, and a hexyl group; and

[0023] a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, and a hexyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, a C₁-C₆₀ alkyl group, a C₆-C₆₀ aryl group, and a C₂-C₆₀ heteroaryl group.

[0024] In one or more embodiments, R₁₁ and R₁₂ in Formula 1 may each independently be a methyl group or a phenyl group.

[0025] Ar₂ in Formula 1 may be selected from a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, and a substituted or unsubstituted C₁-C₆₀ heteroaryl group.

[0026] In one or more embodiments, Ar₂ and R₁₃ in Formula 1 may each independently be selected from:

[0027] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group; and

[0028] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group, each substituted with at least one selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an

amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group.

[0029] In an implementation, Ar₂ and R₁₃ in Formula 1 may be separate or linked to a neighboring substituent to form a substituted or unsubstituted benzene group, a substituted or unsubstituted naphthalene group, a substituted or unsubstituted fluorene group, a substituted or unsubstituted spiro-fluorene group, a substituted or unsubstituted indene group, a substituted or unsubstituted pyrrole group, a substituted or unsubstituted thiophene group, a substituted or unsubstituted furan group, a substituted or unsubstituted imidazole group, a substituted or unsubstituted pyrazole group, a substituted or unsubstituted thiazole group, a substituted or unsubstituted isothiazole group, a substituted or unsubstituted oxazole group, a substituted or unsubstituted isoxazole group, a substituted or unsubstituted pyridine group, a substituted or unsubstituted pyrazine group, a substituted or unsubstituted pyrimidine group, a substituted or unsubstituted pyridazine group, a substituted or unsubstituted quinoline group, a substituted or unsubstituted isoquinoline group, a substituted or unsubstituted benzoquinoline group, a substituted or unsubstituted quinoxaline group, a substituted or unsubstituted quinazoline group, a substituted or unsubstituted carbazole group, a substituted or unsubstituted benzimidazole group, a substituted or unsubstituted benzofuran group, a substituted or unsubstituted benzothiophene group, a substituted or unsubstituted isobenzothiophene group, a substituted or unsubstituted benzoxazole group, a substituted or unsubstituted isobenzoxazole group, a substituted or unsubstituted triazole group, a substituted or unsubstituted oxadiazole group, a substituted or unsubstituted triazine group, a substituted or unsubstituted dibenzofuran group, or a substituted or unsubstituted dibenzothiophene group.

[0030] At least one substituent selected from a substituent (s) of the substituted benzene group, the substituted naphthalene group, the substituted fluorene group, the substituted spiro-fluorene group, the substituted indene group, the substituted pyrrole group, the substituted thiophene group, the substituted furan group, the substituted imidazole group, the substituted pyrazole group, the substituted thiazole group, the substituted isothiazole group, the substituted oxazole group, the substituted isoxazole group, the substituted pyridine group, the substituted pyrazine group, the substituted pyrimidine group, the substituted pyridazine group, the substituted quinoline group, the substituted isoquinoline group, the substituted benzoquinoline group, the substituted quinoxaline group, the substituted quinazoline group, the substituted carbazole group, the substituted benzimidazole group, the substituted benzofuran group, the substituted benzothiophene group, the substituted isobenzothiophene group, the substituted benzoxazole group, the substituted isobenzoxazole group, the substituted triazole group, the

substituted oxadiazole group, the substituted triazine group, the substituted dibenzofuran group, and the substituted dibenzothiophene group may be selected from deuterium, —F, —Cl, —Br, —I, a cyano group, and a C₁-C₆₀ alkyl group.

[0031] In one or more embodiments, Ar₂ and R₁₃ in Formula 1 may each independently be selected from:

[0032] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, and a pyrenyl group; and

[0033] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, and a pyrenyl group, each substituted with deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C_m alkyl group, or a C₁-C₂₀ alkoxy group, or

[0034] Ar₂ and R₁₃ in Formula 1 may be separate or may be linked to a neighboring substituent to form a substituted or unsubstituted benzene, a substituted or unsubstituted naphthalene, a substituted or unsubstituted fluorene, a substituted or unsubstituted spiro-fluorene, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine, a substituted or unsubstituted pyridazine, a substituted or unsubstituted quinoline, a substituted or unsubstituted isoquinoline, a substituted or unsubstituted quinoxaline, a substituted or unsubstituted quinazoline, or a substituted or unsubstituted triazine.

[0035] At least one substituent selected from a substituent (s) of the substituted benzene, the substituted naphthalene, the substituted fluorene, the substituted spiro-fluorene, the substituted pyridine, the substituted pyrazine, the substituted pyrimidine, the substituted pyridazine, the substituted quinoline, the substituted isoquinoline, the substituted quinoxaline, the substituted quinazoline, and the substituted triazine may be selected from deuterium, —F, —Cl, —Br, —I, a cyano group, and a C₁-C₆₀ alkyl group.


[0036] In one or more embodiments, Ar_e in Formula 1 may be selected from:

[0037] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, and a chrysenyl group;


[0038] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a Spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, and a chrysenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a Spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzo-

fluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, and a chrysenyl group.


[0039] In one or more embodiments, Ar₂ in Formula 1 may be one selected from groups represented by Formulae 5-1 to 5-52:




Formula 5-1




Formula 5-2




Formula 5-3




Formula 5-4




Formula 5-5



Formula 5-6










Formula 5-7




Formula 5-8

-continued




-continued

Formula 5-9




Formula 5-16

Formula 5-10




Formula 5-17

Formula 5-11



Formula 5-18

Formula 5-12




Formula 5-19

Formula 5-13


Formula 5-14

Formula 5-15




Formula 5-20


-continued




Formula 5-21




Formula 5-22




Formula 5-23



Formula 5-24




Formula 5-25




Formula 5-26


-continued




Formula 5-27




Formula 5-28



Formula 5-29




Formula 5-30




Formula 5-31


-continued




Formula 5-32




Formula 5-33




Formula 5-34



Formula 5-35




Formula 5-36




Formula 5-37


-continued




Formula 5-38




Formula 5-39



Formula 5-40




Formula 5-41




Formula 5-42


-continued




Formula 5-43




Formula 5-44




Formula 5-45



Formula 5-46




Formula 5-47




Formula 5-48


-continued




Formula 5-49



Formula 5-50



Formula 5-51



Formula 5-52

[0040] In Formulae 5-1 to 5-52,

[0041] Z_{31} to Z_{35} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkenyl group, a C_{20} alkynyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a

pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a triazinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, and $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$.

[0042] Q_1 to Q_3 and Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a dibenzofuranyl group.

[0043] e2 may be 1 or 2,

[0044] e3 may be an integer from 1 to 3,

[0045] e4 may be an integer from 1 to 4,

[0046] e5 may be an integer from 1 to 5,


[0047] e6 may be an integer from 1 to 6,

[0048] e7 may be an integer from 1 to 7,


[0049] e9 may be an integer from 1 to 9, and

[0050] * indicates a binding site to a neighboring atom.


[0051] In one or more embodiments, Ar_e in Formula 1 may be represented by one of Formulae 6-1 to 6-4:




Formula 6-1



Formula 6-2



Formula 6-3



Formula 6-4

[0052] wherein * indicates a binding site to a neighboring atom.

[0053] L_1 in Formula 2 may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic con-

densed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group.

[0054] In one or more embodiments, L_1 in Formula 2 may be selected from:

[0055] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzo-fluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylene group, an isoxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylene group, an isoindolylenylene group, an indolylenylene group, an indazolylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylene group, a benzofuranylenylene group, a benzothioiphenylene group, an isobenzothiazolylenylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylenylene group, an oxadiazolylenylene group, a triazinylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a dibenzosilyloxylenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylenylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group; and


[0056] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzo-fluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a phenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylene group, an isoxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylene group, an isoindolylenylene group, an indolylenylene group, an indazolylene group, a purinylenylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylene group, a benzofuranylenylene group, a benzoth-

iophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a dibenzosilolylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzocarbazolyl group, a benzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).


[0057] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, and a dibenzofuranyl group.

[0058] In one or more embodiments, L₁ in Formula 2 may be represented by one of Formulae 3-1 to 3-49 (e.g., Formula 3-1 to 3-43, 3-43', and 3-44 to 3-49).


-continued




Formula 3-3




Formula 3-4




Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-1



Formula 3-2
















Formula 3-10




Formula 3-11


-continued



Formula 3-12




Formula 3-13




Formula 3-14


Formula 3-15




Formula 3-16



Formula 3-17




Formula 3-18




Formula 3-19

Formula 3-20




Formula 3-21

Formula 3-22



Formula 3-23



Formula 3-24

Formula 3-25

Formula 3-26

Formula 3-27

Formula 3-28







Formula 3-29

Formula 3-30

Formula 3-31


Formula 3-32

-continued




-continued

Formula 3-33




Formula 3-34


Formula 3-35





Formula 3-36



Formula 3-37



Formula 3-38



Formula 3-39

Formula 3-40


Formula 3-41

Formula 3-42


Formula 3-43

Formula 3-43'


-continued




Formula 3-44




Formula 3-45




Formula 3-46



Formula 3-47



Formula 3-48



Formula 3-49

[0059] In Formulae 3-1 to 3-49,

[0060] Y_1 may be O, S, $C(Z_3)(Z_4)$, $N(Z_5)$, or $Si(Z_6)(Z_7)$,

[0061] Z_1 to Z_7 may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a triazinyl group, a benzimidazolyl group, a phenanthrolinyl group, and $-Si(Q_{33})(Q_{34})(Q_{35})$,

[0062] wherein Q_{33} to Q_{35} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

[0063] d_2 may be an integer from 0 to 2,

[0064] d_3 may be an integer from 0 to 3,

[0065] d_4 may be an integer from 0 to 4,


[0066] d_5 may be an integer from 0 to 5,

[0067] d_6 may be an integer from 0 to 6,


[0068] d_8 may be an integer from 0 to 8, and

[0069] * and *' each indicate a binding site to a neighboring atom.


[0070] In one or more embodiments, L_1 in Formula 1 may be selected from groups represented by Formulae 4-1 to 4-6:




Formula 4-1



Formula 4-2




Formula 4-3




Formula 4-4

-continued



Formula 4-5



Formula 4-6

[0071] wherein * and *' in Formulae 4-1 and 4-6 each indicate a binding site to a neighboring atom.

[0072] a₁ in Formula 1 indicates the number of L₁(s). a₁ may be 0, 1, 2, or 3. When a₁ is two or three, the two or three L₁(s) may be identical to or different from each other. When a₁ is zero, *-(L₁)_{a1}-*' may be a single bond.

[0073] In one or more embodiments, a₁ in Formula 1 may be 0, 1, or 2.

[0074] In one or more embodiments, a₁ in Formula 1 may be 0 or 1.

[0075] In one or more embodiments, a₁ in Formula 1 may be 1 or 2.

[0076] Ar₁ in Formula 2 may be a substituted or unsubstituted C₁-C₆₀ heterocyclic group having at least one *-N=* moiety.


[0077] For example, An in Formula 1 may be selected from:

[0078] an imidazolyl group, a pyrazole group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


[0079] an imidazolyl group, a pyrazole group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each

substituted with deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group.


[0080] In one or more embodiments, Ar₁ in Formula 1 may be selected from groups represented by Formulae 7-1 to 7-59:




Formula 7-1




Formula 7-2




Formula 7-3




Formula 7-4




Formula 7-5




Formula 7-6




Formula 7-7



Formula 7-8















Formula 7-9




Formula 7-10


-continued




Formula 7-11




Formula 7-12




Formula 7-13




Formula 7-14




Formula 7-15




Formula 7-16




Formula 7-17




Formula 7-18




Formula 7-19




Formula 7-20



Formula 7-21



Formula 7-22



-continued

Formula 7-23

Formula 7-24

Formula 7-25

Formula 7-26

Formula 7-27

Formula 7-28

Formula 7-29

Formula 7-30

Formula 7-31





Formula 7-32

Formula 7-33

Formula 7-34

Formula 7-35

-continued



[0081] wherein, in Formulae 7-1 to 7-59,

[0082] Z_{51} and Z_{52} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, an imidazolyl group, a pyrazole group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group,

[0083] g_2 may be an integer from 0 to 2,

[0084] g_3 may be an integer from 0 to 3,

[0085] g_4 may be an integer from 0 to 4,

[0086] g_5 may be an integer from 0 to 5,

[0087] g_6 may be an integer from 0 to 6, and

[0088] * and *' each indicate a binding site to a neighboring atom.


Formula 7-56

Formula 7-57


Formula 7-58

Formula 7-59


[0089] For example, the polycyclic compound may be represented by one of Formulae 1-1 to 1-4:




[Formula 1-1]



[Formula 1-2]



[Formula 1-3]




[Formula 1-4]


[0090] wherein, in Formulae 1-1 to 1-4,

[0091] L_1 , al , R_1 to R_{10} , and Ar_2 are the same as described above.

[0092] In one or more embodiments, when X_1 in Formula 1 is $N(R_{13})$, the compound represented by Formula 1 may be represented by Formula 11 or 12.



[Formula 11]



[Formula 12]

[0093] R₁ to R₁₀ in Formulae 11 and 12 are the same as described above.

[0094] Ring A₁ in Formula 11 may be selected from a substituted or unsubstituted benzene, a substituted or unsubstituted naphthalene, a substituted or unsubstituted fluorene, a substituted or unsubstituted spiro-fluorene, a substituted or unsubstituted indene, a substituted or unsubstituted pyrrole, a substituted or unsubstituted thiophene, a substituted or unsubstituted furan, a substituted or unsubstituted imidazole, a substituted or unsubstituted pyrazole, a substituted or unsubstituted thiazole, a substituted or unsubstituted isothiazole, a substituted or unsubstituted oxazole, a substituted or unsubstituted isoxazole, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine, a substituted or unsubstituted pyridazine, a substituted or unsubstituted quinoline, a substituted or unsubstituted isoquinoline, a substituted or unsubstituted benzoquinoline, a substituted or unsubstituted quinoxaline, a substituted or unsubstituted quinazoline, a substituted or unsubstituted carbazole, a substituted or unsubstituted benzimidazole, a substituted or unsubstituted benzofuran, a substituted or unsubstituted benzothiophene, a substituted or unsubstituted isobenzothiophene, a substituted or unsubstituted benzoxazole, a substituted or unsubstituted isobenzoxazole, a substituted or unsubstituted triazole, a substituted or unsubstituted oxadiazole, a substituted or unsubstituted triazine, a substituted or unsubstituted dibenzofuran, and a substituted or unsubstituted dibenzothiophene.

[0095] In one or more embodiments, ring A₁ in Formula 11 may be selected from a substituted or unsubstituted benzene, a substituted or unsubstituted naphthalene, a substituted or unsubstituted fluorene, and a substituted or unsubstituted spiro-fluorene.


[0096] R₂₁ and R₂₂ in Formula 12 may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0097] In one or more embodiments, R₂₁ and R₂₂ may each independently be selected from:

[0098] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group; and

[0099] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, and an ovalenyl group.

[0100] In one or more embodiments, the compound represented by Formula 11 may be represented by Formula 11-1.



[0101] In Formula 11-1,


[0102] R_1 to R_{10} are the same as described above.

[0103] R_{23} in Formula 11-1 may be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, or may be linked to two or more neighboring R_{23} (s) to form a substituted or unsubstituted ring.


[0104] r_1 in Formula 11-1 may be an integer from 0 to 4,

[0105] wherein, when r_1 is two or more, two or more R_{23} (s) may be identical to or different from each other.

[0106] In one or more embodiments, the polycyclic compound represented by Formula 1 may be represented by one of Formulae 21 to 24.




-continued




[Formula 22]

[Formula 23]



[Formula 24]




[0107] In Formulae 21 to 24,


[0108] X_1 , L_1 , Ar_1 , Ar_2 , R_1 to R_{10} , a_1 , and b_1 are the same as described above.

[0109] In one or more embodiments, the polycyclic compound may be one selected from Compounds A-1 to A-51.






A-1



-continued




-continued




-continued

A-8





-continued

A-12





A-13

A-9





A-14

A-10




A-15

A-11




-continued

A-16




-continued

A-20




A-21

A-17





A-22

A-18




A-23

A-19




-continued


A-24




A-25



A-26




A-27




-continued


A-28




A-29



A-30



A-31





-continued

-continued



A-35

A-32





A-36

A-33




A-37

A-34




-continued

A-38





-continued

A-41





A-42

A-39




A-43


A-40




A-44



-continued




-continued




A-48


A-46



A-49





A-50



A-51

A-47



[0110] The polycyclic compound represented by Formula 1 may have a structure in which some ring-forming atoms in a phenalene ring condensed with a benzothiophene or a benzofuran are modified with nitrogen atoms. Thus, it is possible to help prevent a reduction in efficiency under a low current condition by increasing electron mobility through molecular energy adjustment. Also, charge balance in a

device may be appropriately adjusted under a high current condition, thereby obtaining the effects of increasing the lifespan and efficiency of the device.

[0111] Therefore, an electronic device (e.g., an organic light-emitting device) including the polycyclic compound represented by Formula 1 may advantageously have a low driving voltage, a high current density, and high efficiency.

[0112] A synthesis method for the polycyclic compound represented by Formula 1 may be understood by referring to the following examples.

[0113] At least one polycyclic compound of Formula 1 may be used or included between a pair of electrodes of an organic light-emitting device. For example, the polycyclic compound may be included in at least one layer selected from a hole transport region and an emission layer. In one or more embodiments, the polycyclic compound represented by Formula 1 may be used as a material for forming a capping layer positioned outside the pair of electrodes of the organic light-emitting device.

[0114] Accordingly, provided is an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes at least one polycyclic compound represented by Formula 1.

[0115] The expression “(an organic layer) includes at least one of the polycyclic compounds” used herein may include a case in which “(an organic layer) includes identical polycyclic compounds represented by Formula 1” and a case in which “(an organic layer) includes two or more different polycyclic compounds represented by Formula 1.”

[0116] For example, the organic layer may include, as the polycyclic compound, only Compound 1. In this regard, Compound 1 may exist in an emission layer of the organic light-emitting device. In one or more embodiments, the organic layer may include, as the polycyclic compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may exist in an identical layer (for example, Compound 1 and Compound 2 may all exist in an emission layer), or different layers (for example, Compound 1 may exist in an emission layer and Compound 2 may exist in an electron transport layer).

[0117] According to one or more embodiments,

[0118] the first electrode of the organic light-emitting device may be an anode,

[0119] the second electrode of the organic light-emitting device may be a cathode,

[0120] the organic layer of the organic light-emitting device may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode,

[0121] the hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, a buffer layer, an emission auxiliary layer, and an electron blocking layer, and

[0122] the electron transport region may include at least one layer selected from a hole blocking layer, an electron transport layer, and an electron injection layer. The electron transport region may include the polycyclic compound represented by Formula 1.

[0123] In one or more embodiments, the electron transport region of the organic light-emitting device may include an electron transport layer and an electron injection layer, and

at least one of the electron transport layer and the electron injection layer may include the polycyclic compound represented by Formula 1.

[0124] In the organic light-emitting device, at least one layer selected from the electron transport layer and the electron injection layer may further include alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof.

[0125] The emission layer of the organic light-emitting device includes a dopant and a host, and the host may include at least one selected from an anthracene-based compound, a pyrene-based compound, and a spiro-bifluorene-based compound.

[0126] The term “organic layer” as used herein refers to a single layer and/or a plurality of layers between the first electrode and the second electrode of an organic light-emitting device. A material included in the “organic layer” is not limited to an organic material.

[0127] FIG. 1 illustrates a schematic view of an organic light-emitting device 10 according to an embodiment. The organic light-emitting device 10 includes a first electrode 110, an organic layer 150, and a second electrode 190.

[0128] Hereinafter, the structure of the organic light-emitting device 10 according to an embodiment and a method of manufacturing the organic light-emitting device 10 will be described in connection with FIG. 1.

[0129] In an implementation, a substrate may be additionally disposed under the first electrode 110 or above the second electrode 190. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water-resistance.

[0130] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for forming the first electrode 110 may be selected from materials with a high work function to facilitate hole injection.

[0131] The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissible electrode, a material for forming the first electrode 110 may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combinations thereof. When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode 110, magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), or any combination thereof may be used.

[0132] The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO.

[0133] The organic layer 150 is disposed on the first electrode 110. The organic layer 150 may include an emission layer.


[0134] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.

[0135] The hole transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.




[0136] The hole transport region may include at least one layer selected from a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer.

[0137] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/emission auxiliary layer structure, wherein, for each structure, constituting layers are sequentially stacked from the first electrode 110 in this stated order.


[0138] The hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β -NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (Pani/DBSA), PEDOT/PSS (poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate)), polyaniline/camphor sulfonic acid (Pani/CSA), polyaniline/poly(4-styrenesulfonate) (Pani/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:



-continued




-continued




β -NPB


-continued




TAPC




TPD




HMTPD



Spiro-TPD




Spiro-NPB




methylated NPB

<Formula 201>



<Formula 202>



[0139] In Formulae 201 and 202,

[0140] L_{201} to L_{204} are each independently selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0141] L_{205} may be selected from $*-O-*$, $*-S-*$, $*-N(Q_{201})-*$, a substituted or unsubstituted C_1 - C_{20} alkylene group, a substituted or unsubstituted C_2 - C_{20} alkenylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0142] xa1 to xa4 may each independently be an integer from 0 to 3,

[0143] xa5 may be an integer from 1 to 10, and

[0144] R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0145] For example, in Formula 202, R₂₀₁ and R₂₀₂ may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R₂₀₃ and R₂₀₄ may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

[0146] In one or more embodiments, in Formulae 201 and 202,

[0147] L₂₀₁ to L₂₀₅ may each independently be selected from:

[0148] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, and a pyridinylene group; and

[0149] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an

indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, and a pyridinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0150] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0151] In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

[0152] In one or more embodiments, xa5 may be 1, 2, 3, or 4.

[0153] In one or more embodiments, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranlyl group, a benzothiophenyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0154] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂),

[0155] wherein Q₃₁ to Q₃₃ are the same as described above.

[0156] In one or more embodiments, at least one of R₂₀₁ to R₂₀₃ in Formula 201 may each independently be selected from:

[0157] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

[0158] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl

group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0159] In one or more embodiments, in Formula 202, i) R₂₀₁ and R₂₀₂ may be linked via a single bond and/or ii) R₂₀₃ and R₂₀₄ may be linked via a single bond.

[0160] In one or more embodiments, at least one of R₂₀₁ to R₂₀₄ in Formula 202 may be selected from:


[0161] a carbazolyl group; and

[0162] a carbazolyl group, substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group.


[0163] The compound represented by Formula 201 may be represented by Formula 201A:




[0164] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1) below:




[0165] In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1 below:



[0166] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A:



[0167] In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:





[0168] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,

[0169] L_{201} to L_{203} , $xa1$ to $xa3$, $xa5$, and R_{202} to R_{204} may be the same as described above,

[0170] R_{211} and R_{212} may be understood by referring to the description provided herein in connection with R_{203} .

[0171] R_{213} to R_{217} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a phenyl group substituted with $-F$, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a Spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.



[0172] The hole transport region may include at least one compound selected from Compounds HT1 to HT39:



-continued



HT3

HT4





HT5

HT6





HT7

HT8





-continued
HT9

HT10





HT11

HT12




HT13


HT14




-continued




HT15




HT16




HT17



HT18




HT19




HT20


-continued
HT21




HT22




HT23









HT24



HT25




-continued




-continued


HT32




HT33




HT34




HT35



HT36





HT37



-continued
HT38

HT39



[0173] A thickness of the hole transport region may be in a range of about 100 Å to about 10,000 Å, e.g., about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, the thickness of the hole injection layer may be in a range of about 100 Å to about 9,000 Å, e.g., about 100 Å to about 1,000 Å, and the thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, e.g., about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0174] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[0175] The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0176] The charge-generation material may be, e.g., a p-dopant.

[0177] In one or more embodiments, a lowest unoccupied molecular orbital (LUMO) level of the p-dopant may be -3.5 eV or less.

[0178] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound.



[0179] For example, the p-dopant may include at least one selected from:


[0180] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

[0181] a metal oxide, such as tungsten oxide or molybdenum oxide;

[0182] 1,4,5,8,9,11-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0183] a compound represented by Formula 221 below:





[0184] In Formula 221,

[0185] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one of R_{221} to R_{223} may have at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C_1 - C_{20} alkyl group substituted with —F, a C_1 - C_{20} alkyl group substituted with —Cl, a C_1 - C_{20} alkyl group substituted with —Br, and a C_1 - C_{20} alkyl group substituted with —I.

[0186] When the organic light-emitting device 10 is a full color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers contact each other or are separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red-light emission material, a green-light emission material, and a blue-light emission material, in which the two or more materials are mixed with each other in a single layer to emit white light.


[0187] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0188] An amount of the dopant in the emission layer may be, e.g., in a range of about 0.01 to about 15 parts by weight based on 100 parts by weight of the host.

[0189] A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0190] In one or more embodiments, the host may include a compound represented by Formula 301 below.

[0191] <Formula 301>



[0192] In Formula 301,

[0193] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0194] x_{b11} may be 1, 2, or 3,

[0195] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsub-

stituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0196] x_{b1} may be an integer from 0 to 5,

[0197] R_{301} may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_{301})(Q_{302})(Q_{303}), —N(Q_{301})(Q_{302}), —B(Q_{301})(Q_{302}), —C(=O)(Q_{301}), —S(=O)₂(Q_{301}), and —P(=O)(Q_{301})(Q_{302}),

[0198] x_{b21} may be an integer from 1 to 5,

[0199] Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0200] In one or more embodiments, Ar_{301} in Formula 301 may be selected from:

[0201] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and


[0202] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —B(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}),

[0203] wherein Q_3 to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.


[0204] When x_{b11} in Formula 301 is two or more, two or more $Ar_{301}(s)$ may be linked via a single bond.

[0205] In one or more embodiments, the compound represented by Formula 301 may be represented by Formulae 301-1 or 301-2:

<Formula 301-1>



<Formula 301-2>



[0206] In Formulae 301-1 to 301-2,

[0207] A_{301} to A_{304} may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group,

[0208] X_{301} may be O, S, or N- $[(L_{304})_{xb4}-R_{304}]$,

[0209] R_{311} to R_{314} may each independently be selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group -Si(Q_{31})(Q_{32})(Q_{33}), -N(Q_{31})(Q_{32}), -B(Q_{31})(Q_{32}), -C(=O)(Q_{31}), -S(=O)₂(Q_{31}), and -P(=O)(Q_{31})(Q_{32}),

[0210] $xb22$ and $xb23$ may each independently be 0, 1, or 2,

[0211] L_{301} , $xb1$, R_{301} , and Q_{31} to Q_{33} may be the same as described above,

[0212] L_{302} to L_{304} may each independently be the same as described in connection with

[0213] L_{301} ,

[0214] $xb2$ to $xb4$ may each independently be the same as described in connection with $xb1$, and

[0215] R_{302} to R_{304} may each independently be the same as described in connection with R_{301} .

[0216] For example, L_{301} to L_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0217] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a

thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylenylene group, an isobenzothiazolylenylene group, a benzoxazolylenylene group, an isobenzoxazolylenylene group, a triazolylenylene group, a tetrazolylenylene group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, and an azacarbazolylenylene group; and

[0218] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylenylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylenylene group, a dibenzocarbazolylenylene group, a dibenzosilolylenylene group, a pyridinylenylene group, an imidazolylenylene group, a pyrazolylenylene group, a thiazolylenylene group, an isothiazolylenylene group, an oxazolylenylene group, an isoxazolylenylene group, a thiadiazolylenylene group, an oxadiazolylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, a triazinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a phenanthridinylenylene group, an acridinylenylene

group, a phenanthrolylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an aza carbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁) and —P(=O)(Q₃₁)(Q₃₂),

[0219] wherein Q₃₁ to Q₃₃ may be the same as described above.

[0220] In one or more embodiments, R₃₀₁ to R₃₀₄ in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0221] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a Spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetra-

zolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and


[0222] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetra-

[0223] Q₃₁ to Q₃₃ may be the same as described above.


[0224] In one or more embodiments, the host may include an alkaline-earth metal complex. For example, the host may

be selected from a Be complex (for example, Compound H55), an Mg complex, and a Zn complex.


[0225] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), a 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55:




H1




H2



H3




H4




H5


-continued




H6




H7




H8



H9




H10











H11

-continued




-continued




-continued


H21




H22



H23





H24




-continued

H25




H26



H27


-continued

H28




-continued


H32




H29




H33




H30




H34




H31



H35




-continued




H36

-continued



H40



H37




H38



H39







H41



H42


-continued



-continued


H43

H47




H48

H44




H49

H45




H50


H46




H51




H52




-continued



H53




H54




H55

[0226] The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:



<Formula 401>



<Formula 402>

[0227] In Formulae 401 and 402,

[0228] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

[0229] L_{401} may be selected from ligands represented by Formula 402, and $xc1$ may be 1, 2, or 3, wherein, when $xc1$ is two or more, two or more $L_{401}(s)$ may be identical to or different from each other,

[0230] L_{402} may be an organic ligand, and $xc2$ may be an integer from 0 to 4, wherein, when $xc2$ is two or more, two or more $L_{402}(s)$ may be identical to or different from each other,

[0231] X_{401} to X_{404} may each independently be nitrogen or carbon,

[0232] X_{401} and X_{403} may be linked via a single bond or a double bond, and X_{402} and X_{404} may be linked via a single bond or a double bond,

[0233] A_{401} and A_{402} may each independently be selected from a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

[0234] X_{405} may be a single bond, $*-O-*$, $*-C(=O)-*I$, $*-N(Q_{411})-*$, $*-C(Q_{411})(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, or $*=C(Q_{411})-*$, wherein Q_{411} and Q_{412} may be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0235] X_{406} may be a single bond, O, or S,

[0236] R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$, wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_6 - C_{20} aryl group, and a C_1 - C_{20} heteroaryl group,

[0237] $xc11$ and $xc12$ may each independently be an integer from 0 to 10, and

[0238] $*$ and * in Formula 402 each indicate a binding site to M in Formula 401.

[0239] In one or more embodiments, A_{401} and A_{402} in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole

group, a benzimidazole group, a benzofuran group, a benzothiophene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0240] In one or more embodiments, in Formula 402, i) X_{401} may be nitrogen, and X_{402} may be carbon, or ii) X_{401} and X_{402} may each be nitrogen at the same time.

[0241] In one or more embodiments, R_{401} and R_{402} in Formula 402 may each independently be selected from:

[0242] hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0243] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0244] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofurananyl group, and a dibenzothiophenyl group;

[0245] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofurananyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofurananyl group, and a dibenzothiophenyl group; and

[0246] —Si(Q_{401})(Q_{402})(Q_{403}), —N(Q_{401})(Q_{402}), —B(Q_{401})(Q_{402}), —C(=O)(Q_{401}), —S(=O)₂(Q_{401}), and —P(=O)(Q_{401})(Q_{402}),

[0247] wherein Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group.


[0248] In one or more embodiments, when xcl in Formula 401 is two or more, two A_{401} (s) in two or more L_{401} (s) may optionally be linked to each other via X_{407} , which is a linking group, or two A_{402} (s) in two or more L_{401} (s) may optionally be linked to each other via X_{408} , which is a linking group (see Compounds PD1 to PD4 and PD7). X_{407} and X_{408} may each independently be a single bond, *—O—*, *—S—*, *—C(=O)—*, *—N(Q_{413})—*, *—C(Q_{413})(Q_{414})—*, or *—C(Q_{413})=C(Q_{414})—* (wherein Q_{413} and Q_{414} may each independently be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group).

[0249] L_{402} in Formula 401 may be a monovalent, divalent, or trivalent organic ligand.


[0250] For example, L_{402} may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), —C(=O), isonitrile, —CN, and phosphorus (for example, phosphine, or phosphite).

[0251] In one or more embodiments, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25:


PD1









PD2



PD3




-continued




-continued

PD4




PD10

PD5



PD11


PD6



PD12


PD7

PD8








PD13

PD9




PD14

-continued




-continued

PD15




PD20

PD16




PD21

PD17



PD22



PD18



PD23


PD19

-continued



[0252] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0253] The fluorescent dopant may include a compound represented by Formula 501 below.



[0254] In Formula 501,

[0255] Ar_{501} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0256] L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0257] $xd1$ to $xd3$ may each independently be an integer of 0 to 3;

[0258] R_{501} and R_{502} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalk-

enyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0259] $xd4$ may be an integer of 1 to 6.

[0260] In one or more embodiments, Ar_{501} in Formula 501 may be selected from:

[0261] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

[0262] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0263] In one or more embodiments, L_{501} to L_{503} in Formula 501 may each independently be selected from:

[0264] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, dibenzosilolylylene group, and a pyridinylylene group; and

[0265] a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naph-

thyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0266] In one or more embodiments, R_{501} and R_{501} in Formula 502 may each independently be selected from:


[0267] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0268] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and —Si(Q_{31})(Q_{32})(Q_{33}),


[0269] wherein Q_{31} to Q_{33} may each be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0270] In one or more embodiments, $xd4$ in Formula 501 may be 2.


[0271] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:



FD1




FD2




FD3

-continued




FD4

-continued




FD7

FD5





FD6


FD8




FD9




-continued




-continued




FD11




FD16




FD12




FD17




FD13



FD18




FD14




-continued

FD19





-continued

FD21




FD22


FD20




[0272] In one or more embodiments, the fluorescent dopant may be selected from the following compounds.



DPVBi



DPAVBi



[0273] The electron transport region may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0274] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer.

[0275] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein, for each structure, constituting layers are sequentially stacked from an emission layer.

[0276] The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one π electron-depleted nitrogen-containing ring.

[0277] The “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0278] For example, the “ π electron-depleted nitrogen-containing ring” may be i) a 5-membered to 7-membered hetero monocyclic group having at least one $*-N=*$ moiety, ii) a heteropoly cyclic group in which two or more 5-membered to 7-membered hetero monocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or iii) a heteropoly cyclic group in which at least one of 5-membered to 7-membered hetero monocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0279] Examples of the 7C electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a thiadiazole, an imidazopyridine, an imidazopyrimidine, and an azacarbazole.

[0280] For example, the electron transport region may include a compound represented by Formula 601:



[0281] In Formula 601,

[0282] Ar_{601} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0283] $xe11$ may be 1, 2, or 3,

[0284] L_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0285] $xe1$ may be an integer from 0 to 5,

[0286] R_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substi-

tuted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₆₀₁)(Q₆₀₂)(Q₆₀₃), —C(=O)(Q₆₀₁), —S(=O)₂(Q₆₀₁), and —P(=O)(Q₆₀₁)(Q₆₀₂).

[0287] Q₆₀₁ to Q₆₀₃ may each independently be a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and [0288] xe21 may be an integer from 1 to 5.

[0289] In one or more embodiments, at least one of Ar₆₀₁(s) in the number of xe11 and/or at least one of R₆₀₁(s) in the number of xe21 may include the it electron-depleted nitrogen-containing ring.

[0290] In one or more embodiments, ring Ar₆₀₁ in Formula 601 may be selected from:

[0291] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0292] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, thiadiazol group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydra-

zono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂).


[0293] wherein Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0294] When xe11 in Formula 601 is two or more, two or more Ar₆₀₁(s) may be linked to each other via a single bond.

[0295] In one or more embodiments, Ar₆₀₁ in Formula 601 may be an anthracene group.

[0296] In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



[0297] In Formula 601-1,

[0298] X₆₁₄ may be N or C(R₆₁₄), X₆₁₅ may be N or C(R₆₁₅), X₆₁₆ may be N or C(R₆₁₆), and at least one selected from X₆₁₄ to X₆₁₆ may be N,

[0299] L₆₁₁ to L₆₁₃ may be each independently substantially the same as described in connection with L₆₀₁,

[0300] xe611 to xe613 may be each independently substantially the same as described in connection with xe1,

[0301] R₆₁₁ to R₆₁₃ may be each independently substantially the same as described in connection with R₆₀₁, and

[0302] R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0303] In one or more embodiments, L₆₀₁ and L₆₁₁ to L₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0304] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a benzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene

group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group; and

[0305] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

azolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group.

[0306] In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

[0307] In one or more embodiments, R₆₀₁ and R₆₁₁ to R₆₁₃ in Formula 601 and 601-1 may each independently be selected from:

[0308] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and





[0309] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

idinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenaziny group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and




[0310] —S(=O)₂(Q₆₀₁), and —P(=O)(Q₆₀₁)(Q₆₀₂),

[0311] wherein Q₆₀₁ and Q₆₀₂ are substantially the same as described above.



[0312] The electron transport region may include at least one compound selected from Compounds ET1 to ET36:



-continued




-continued




-continued

ET10




-continued


ET13




ET11




ET14






ET12



ET15




-continued




-continued

ET16




ET19

ET17



ET20

ET18





ET21

-continued

-continued



ET22

ET25





ET23

ET26




ET24

ET27




-continued




ET28

-continued




ET31


ET29




ET32






ET30




ET33






-continued



ET34

Alq₃

ET35



NTAZ

[0313] In one or more embodiments, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq₃, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:


[0314] The thickness of the buffer layer, the hole blocking layer, or the electron controlling layer may each independently be in a range of about 20 Å to about 1,000 Å, e.g., about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

[0315] A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, e.g., about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the range described above, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.


[0316] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

[0317] The metal-containing material may include at least one selected from an alkali metal complex and an alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from an Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, an Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenylan oxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazole, a hydroxy diphenylthiadiazole, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene.

[0318] For example, the metal-containing material may include a Li complex. The Li complex may include, e.g., Compound ET-D1 (lithium quinolate, LiQ) or ET-D2:



ET-D1



ET-D2

[0319] The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode 190. The electron injection layer may directly contact the second electrode 190.

[0320] The electron injection layer may have i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

[0321] The electron injection layer may include alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof.

[0322] The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one or more embodiments, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs.

[0323] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0324] The rare-earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

[0325] The alkali metal compound, the alkaline earth-metal compound, and the rare-earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare-earth metal.

[0326] The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, KI, or RbI. In one or more embodiments, the alkali metal compound may be selected from LiF, Li_2O , NaF, LiI, NaI, CsI, and KI.

[0327] The alkaline earth-metal compound may be selected from alkaline earth-metal compounds, such as BaO, SrO, CaO, $\text{Ba}_x\text{Sr}_{1-x}\text{O}$ ($0 < x < 1$), or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$). In one or more embodiments, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO.

[0328] The rare-earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In one or more embodiments, the rare-earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 .

[0329] The alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare-earth metal as described above, and a ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, and the rare-earth metal complex may each independently be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenylan oxazole, hydroxy phenylthiazole, hydroxy diphenylan oxadiazole, hydroxy diphenylthiadiazole, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene.

[0330] The electron injection layer may consist of alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, alkali metal, alkaline earth metal, rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare-earth metal complex, or any combinations thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0331] A thickness of the electron injection layer may be in a range of about 1 Å to about 100 Å, e.g., about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

[0332] The second electrode 190 may be disposed on the organic layer 150 having such a structure. The second electrode 190 may be a cathode that is an electron injection electrode, and in this regard, a material for forming the

second electrode **190** may be a material having a low work function, and such a material may be metal, alloy, an electrically conductive compound, or a combination thereof.

[0333] The second electrode **190** may include at least one selected from lithium (Li), silver (Si), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO. The second electrode **190** may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0334] The second electrode **190** may have a single-layered structure, or a multi-layered structure including two or more layers.

[0335] Hereinbefore, the organic light-emitting device has been described with reference to FIG. 1.

[0336] Layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region may be formed in a certain region by using one or more suitable methods selected from vacuum deposition, spin coating, casting, langmuir-blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0337] When the respective layers of the hole transport region, the emission layer, and the respective layers of the electron transport region are formed by deposition, the deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10⁻⁸ to about 10⁻³ torr, and at a deposition rate of about 0.01 to about 100 Å/sec by taking into account a material for forming a layer to be deposited, and the structure of a layer to be formed.

[0338] When layers constituting the hole transport region, an emission layer, and layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to 200° C., depending on a material to be included in a layer and the structure of each layer to be formed.

[0339] The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and non-limiting examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C₁-C₆₀ alkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₆₀ alkyl group.

[0340] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and non-limiting examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkenyl group.

[0341] The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and non-limiting examples thereof include an ethynyl group and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as the C₂-C₆₀ alkynyl group.

[0342] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and non-limiting examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0343] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and non-limiting examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkyl group.

[0344] The term “C₁₀-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent saturated monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and non-limiting examples thereof include a 1,2,3,4-oxatriazolidiny group, a tetrahydrofuranly group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0345] The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and does not have aromaticity, and non-limiting examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0346] The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Non-limiting examples of the C₁-C₁₀ heterocycloalkenyl group include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranly group and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0347] The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and the term “C₆-C₆₀ arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other.

[0348] The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C₁-C₆₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a

triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₁-C₆₀ heteroaryl group and the C₁-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other.

[0349] The term “C₆-C₆₀ aryloxy group” as used herein refers to —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and a C₆-C₆₀ arylthio group used herein indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

[0350] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) that has two or more rings condensed with each other, only carbon atoms as a ring-forming atom, and non-aromaticity in the entire molecular structure. A detailed example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0351] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) that has two or more rings condensed to each other, has at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and has non-aromaticity in the entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group,” used herein, refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0352] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as a benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0353] The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having the same structure as the C₁-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

[0354] At least one substituent of the substituted C₅-C₆₀ carbocyclic group, substituted C₁-C₆₀ heterocyclic group, the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the sub-

stituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0355] deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0356] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0357] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0358] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂); and

[0359] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂),

[0360] wherein Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a

C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0361] The term “Ph”, as used herein, may refer to a phenyl group; the term “Me”, as used herein, may refer to a methyl group; the term “Et”, as used herein, may refer to an ethyl group; the terms “ter-Bu” or “But”, as used herein, may refer to a tert-butyl group; and the term “OMe” as used herein may refer to a methoxy group.

[0362] The term “biphenyl group” as used therein refers to “a phenyl group substituted with a phenyl group.” For example, a “biphenyl group” is a substituted phenyl group having a C_6 - C_{60} aryl group as a substituent.

[0363] The term “terphenyl group” as used herein refers to “a phenyl group substituted with a biphenyl group.” For example, a “terphenyl group” is a substituted phenyl group having a C_6 - C_{60} aryl group substituted with a C_6 - C_{60} aryl group as a substituent.

[0364] * and *’ used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.


[0365] The term “or” is not an exclusive term. For example, a composition including A or B would encompass a composition including A, a composition including B, and a composition including A and B.

[0366] Hereinafter, a compound according to embodiments and an organic light-emitting device according to embodiments will be described in detail with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.

[0367] The following Examples and Comparative Examples are provided in order to highlight characteristics of one or more embodiments, but it will be understood that the Examples and Comparative Examples are not to be construed as limiting the scope of the embodiments, nor are the Comparative Examples to be construed as being outside the scope of the embodiments. Further, it will be understood that the embodiments are not limited to the particular details described in the Examples and Comparative Examples.

SYNTHESIS EXAMPLES


[0368] The structures of compounds used in the Synthesis Examples were as follows:




S

-continued


Int-1




Int-2




Int-3








Int-4



Int-5



-continued




Synthesis Example 1

Int-6

Synthesis of Compound A-1


[0369] Synthesis of Intermediate Int-1

T-1



Int-1

T-2



T-3

[0370] As a starting material, 1 equivalent of Compound S was dissolved in 100 ml of ether and cooled to a temperature of -78°C . Then, 1.2 equivalents of 1.6 M hex-methyl lithium was added thereto and stirred for 1 hour. Then, the resulting mixture was gradually heated to ambient temperature and stirred for 4 hours. After the reaction was completed, an extraction process was performed thereon by using 200 mL of water and an organic layer was dried under reduced pressure. The result was recrystallized with a mixture of ethanol and acetone (volume ratio of 1:1), thereby completing the preparation of Intermediate Int-1 (67%).


[0371] HRMS ($[M]^+$) calcd for $\text{C}_{27}\text{H}_{22}\text{BrNO}$: 456.38, Found: 455

[0372] Calculated values through atom analysis of $\text{C}_{27}\text{H}_{22}\text{BrNO}$: C, 71.06; H, 4.86; Br, 17.51; N, 3.07; O, 3.51


T-4

[0373] Synthesis of Intermediate Int-2

Int-1



-continued




Int-2

[0374] Intermediate Int-1 was dissolved in 150 mL of strong H_3PO_4 and stirred for 5.5 hours. An extraction process was performed thereon by using 200 mL of water and 200 mL of ethyl acetate, and an organic layer was dried under reduced pressure. The result was recrystallized with 150 mL of ethanol, thereby completing the preparation of Intermediate Int-2(72%).

[0375] HRMS ($[M]^+$) calcd for $C_{27}H_{20}BrN$: 438.37, Found: 437

[0376] Synthesis of Compound A-1



[0377] 1 equivalent of Intermediate Int-2 and 1.2 equivalents of Compound T-1 were added to a flask, and 0.03 equivalents of $Pd(dba)_3$, 0.06 equivalents of $(t-Bu)_3P$, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with $MgSO_4$ and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-1(71.4%).


[0378] HRMS ($[M]^+$) calcd for $C_{48}H_{34}N_4$: 666.83, Found: 665

[0379] Calculated values through atom analysis of $C_{48}H_{34}N_4$: C, 86.46; H, 5.14; N, 8.40

Synthesis Example 2

Synthesis of Compound A-2

[0380]



[0381] 1 equivalent of Intermediate Int-2, which was synthesized in the same manner as in Synthesis Example 1, and 1.2 equivalent of Compound T-2 were added to a flask, and 0.03 equivalents of Pd(dba)₃, 0.06 equivalents of (t-Bu)₃P, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO₄ and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-2(73.7%).


[0382] HRMS ([M]⁺) calcd for C₄₈H₃₄N₄: 666.83, Found: 665

[0383] Calculated values through atom analysis of C₄₈H₃₄N₄: C, 86.46; H, 5.14; N, 8.40

Synthesis Example 3

Synthesis of Compound A-10

[0384]



[0385] 1 equivalent of Intermediate Int-2, which was synthesized in the same manner as in Synthesis Example 1, and 1.2 equivalent of Compound T-3 were added to a flask, and 0.03 equivalents of Pd(dba)₃, 0.06 equivalents of (t-Bu)₃P, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO₄ and distilled under reduced

pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-10 (75.9%).


[0386] HRMS ([M]⁺) calcd for C₅₂H₃₆N₄: 716.89, Found: 715

[0387] Calculated values through atom analysis of C₅₂H₃₆N₄: C, 87.12; H, 5.06; N, 7.82

Synthesis Example 4

Synthesis of Compound A-15

[0388]



[0389] 1 equivalent of Intermediate Int-2, which was synthesized in the same manner as in Synthesis Example 1, and 1.2 equivalent of Compound T-4 were added to a flask, and 0.03 equivalents of Pd(dba)₃, 0.06 equivalents of (t-Bu)₃P, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO₄ and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-15 (75.9%).


[0390] HRMS ([M]⁺) calcd for C₅₂H₃₆N₄: 716.89, Found: 715

[0391] Calculated values through atom analysis of C₅₂H₃₆N₄: C, 87.12; H, 5.06; N, 7.82

Synthesis Example 5


Synthesis of Compound A-32

[0392] Synthesis of Intermediate Int-4



Int-3


Int-4



[0393] As a starting material, Intermediate Int-3 and trifluoromethanesulfonic acid ($\text{CF}_3\text{SO}_3\text{H}$) were added to a flask and stirred at ambient temperature for 24 hours. Then, a water and pyridine mixed solvent (volume ratio of 8:1) was added thereto and refluxed and stirred for 30 minutes. Then, the resulting mixture was cooled to ambient temperature, and an extraction process was performed thereon by using methylene chloride. The obtained material was dried by using MgSO_4 and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Intermediate Int-4 (41%).


[0394] HRMS ($[\text{M}]^+$) calcd for $\text{C}_{24}\text{H}_{14}\text{BrNO}$: 412.29, Found: 411

[0395] Synthesis of Compound A-32



Int-4

-continued



A-32

[0396] 1 equivalent of Intermediate Int-4 and 1.2 equivalents of Compound T-2 were added to a flask, and 0.03 equivalents of $\text{Pd}(\text{dba})_3$, 0.06 equivalents of $(\text{t-Bu})_3\text{P}$, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO_4 and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-32 (67%).


[0397] HRMS ($[\text{M}]^+$) calcd for $\text{C}_{45}\text{H}_{28}\text{N}_4\text{O}$: 640.75, Found: 639

[0398] Calculated values through atom analysis of $\text{C}_{45}\text{H}_{28}\text{N}_4\text{O}$: C, 84.35; H, 4.40; N, 8.74; O, 2.50

Synthesis Example 6


Synthesis of Compound A-43

[0399]



Int-4

-continued



A-43

[0400] 1 equivalent of Intermediate Int-4 and 1.2 equivalents of Compound T-3 were added to a flask, and 0.03 equivalents of Pd(dba)₃, 0.06 equivalents of (t-Bu)₃P, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO₄ and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-43 (68%).


[0401] HRMS calcd for C₄₉H₃₀N₄O: 690.81, Found: 689

[0402] Calculated values through atom analysis of C₄₉H₃₀N₄O: C, 85.20; H, 4.38; N, 8.11; O, 2.32


Synthesis Example 7

Synthesis of Compound A-35


[0403] (1) Synthesis of Intermediate Int-6




Int-5



Int-6



Int-6



A-35

[0407] 1 equivalent of Intermediate Int-6 and 1.2 equivalents of Compound T-2 were added to a flask, and 0.03 equivalents of Pd(dba)₃, 0.06 equivalents of (t-Bu)₃P, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO₄ and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-35 (67%).

[0408] HRMS ([M]⁺) calcd for C₄₅H₂₈N₄S: 656.81, Found: 655

[0409] Calculated values through atom analysis of C₄₅H₂₈N₄S: C, 82.29; H, 4.30; N, 8.53; S, 4.88

[0404] As a starting material, Intermediate Int-5 and trifluoromethanesulfonic acid (CF₃SO₃H) were added to a flask and stirred at ambient temperature for 24 hours. Then, a

water and pyridine mixed solvent (volume ratio of 8:1) was added thereto and refluxed and stirred for 30 minutes. Then, the resulting mixture was cooled to ambient temperature, and an extraction process was performed thereon by using methylene chloride. The obtained material was dried by using MgSO₄ and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Intermediate Int-6 (39.7%).


[0405] HRMS ([M]⁺) calcd for C₂₄H₁₄BrNS: 428.35, Found: 427

[0406] (2) Synthesis of Compound A-35

Synthesis Example 8

Synthesis of Compound A-49

[0410]



[0411] 1 equivalent of Intermediate Int-6 and 1.2 equivalents of Compound T-3 were added to a flask, and 0.03 equivalents of $\text{Pd}(\text{dba})_3$, 0.06 equivalents of $(\text{t-Bu})_3\text{P}$, and toluene (based on 0.1 M 1 eq. reagent) were added thereto. Then, the resulting mixture was refluxed and stirred for 5 hours and cooled to ambient temperature. An extraction process was performed thereon by using methylene chloride and a washing process was performed thereon. The obtained material was dried with MgSO_4 and distilled under reduced pressure. Then, the residue obtained therefrom was separated by a column, thereby completing the preparation of Compound A-49 (57%).

[0412] HRMS ($[\text{M}]^+$) calcd for $\text{C}_{49}\text{H}_{30}\text{N}_4\text{S}$: 706.87, Found: 705

[0413] Calculated values through atom analysis of $\text{C}_{49}\text{H}_{30}\text{N}_4\text{S}$: C, 83.26; H, 4.28; N, 7.93; S, 4.54

Example

Example 1




[0414] A glass substrate, in which an ITO/Ag/ITO anode was formed, was ultrasonically cleaned by using isopropyl alcohol and pure water each for 5 minutes and then cleaned by exposure to UV irradiation for 10 minutes and ozone. Then, the glass substrate was loaded into a vacuum deposition apparatus.

[0415] 2-TNATA was vacuum-deposited on the ITO/Ag/ITO anode to form a hole injection layer having a thickness

of 600 Å, and NPB was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.


[0416] Bis-(1-phenylisoquinolyl)acetylacetonate iridium (III) ($\text{Ir}(\text{piq})_2(\text{acac})$) and CBP were co-deposited on the hole transport layer at a weight ratio of 15:85 to form an emission layer having a thickness of 300 Å.

[0417] Compound A-1 was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å, and Al was deposited on the electron transport layer to form a cathode having a thickness of 1,200 Å, thereby completing the manufacture of an organic light-emitting device.



-continued


A-1



Examples 2 to 10 and Comparative Example 1 and 2

[0418] Organic light-emitting devices of Examples 2 to 10 and Comparative Examples 1 and 2 were manufactured in the same manner as in Example 1, except that Compounds shown in Table 1 were each used instead of Compound A-1 in forming an electron transport layer.

[HT-A]



Evaluation Example 1

[0419] The efficiency and lifespan of the organic light-emitting devices manufactured according to Examples 1 to 10 and Comparative Examples 1 and 2 were evaluated by using a Keithley SMU 236 and a luminance meter PR650, and results thereof are shown in Table 1. The lifespan (T_{95}) indicates an amount of time that lapsed when luminance was 95% of initial luminance (100%).

TABLE 1

	Electron transport layer	Efficiency (cd/A)	Lifespan (T_{95}) (hr) (@3,700 nit)
Example 1	A-1	51	923
Example 2	A-2	48.8	911
Example 3	A-10	47.7	897
Example 4	A-15	47.9	921
Example 5	A-23	49.2	943

TABLE 1-continued

	Electron transport layer	Efficiency (cd/A)	Lifespan (T_{95}) (hr) (@3,700 nit)
Example 6	A-29	48.0	937
Example 7	A-32	47.9	956
Example 8	A-35	42	942
Example 9	A-43	46.4	965
Example 10	A-49	42.7	970
Comparative Example 1	Alq3	38.4	789
Comparative Example 2	HT-A	47.4	884

[0420] Referring to Table 1, it may be seen that the organic light-emitting devices of


[0421] Examples 1 to 10 had excellent lifespan characteristics, compared to those of Comparative Examples 1 and 2. Also, it may be seen that the organic light-emitting devices of Examples 1 to 7 had excellent efficiency characteristics, compared to those of Comparative Examples 1 and 2.

[0422] According to one or more example embodiments, an organic light-emitting device including the polycyclic compound may have a low driving voltage, high efficiency, high luminance, and a long lifespan.

[0423] Example embodiments have been disclosed herein, and although specific terms are employed, they are used and are to be interpreted in a generic and descriptive sense only and not for purpose of limitation. In some instances, as would be apparent to one of ordinary skill in the art as of the filing of the present application, features, characteristics, and/or elements described in connection with a particular embodiment may be used singly or in combination with features, characteristics, and/or elements described in connection with other embodiments unless otherwise specifically indicated. Accordingly, it will be understood by those of skill in the art that various changes in form and details may be made without departing from the spirit and scope of the present invention as set forth in the following claims.

What is claimed is:

1. A polycyclic compound represented by Formula 1:




wherein, in Formula 1,

X_1 is $C(R_{11})(R_{12})$, $N(R_{13})$, O, or S, and

R_1 to R_{13} are each independently a structure represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl

group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q_i)(Q₂)(Q₃), —N(Q₁)(Q₂), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), or —P(=O)(Q₁)(Q₂), at least one of R₁ to R₁₀ having a structure represented by Formula 2,

[Formula 2]



wherein, in Formulae 1 and 2,

L₁ is a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

Ar₁ is a substituted or unsubstituted C₁-C₆₀ heterocyclic group having at least one *—N=* moiety,

a₁ is an integer from 1 to 5,

b₁ is an integer from 1 to 5,

when a₁ is two or more, two or more L₁(s) are identical to or different from each other,

when b₁ is two or more, two or more Ar₁(s) are identical to or different from each other,

Ar₂ is a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, or a substituted or unsubstituted C₁-C₆₀ heteroaryl group, and

R₁₃ and Ar₂ are separate or are linked to a neighboring substituent to form a substituted or unsubstituted ring,

* and *' each indicate a binding site to a neighboring atom, and

at least one substituent of the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy

group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ heterocyclic group, or the substituted ring is:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), or —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, a biphenyl group, or a terphenyl group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, a biphenyl group, and a terphenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, a terphenyl group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), or —P(=O)(Q₂₁)(Q₂₂); or —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), or —P(=O)(Q₃₁)(Q₃₂),

wherein Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a

hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryl group substituted with a C₁-C₆₀ alkyl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a C₁-C₆₀ heterocyclic group, a biphenyl group, or a terphenyl group.

2. The polycyclic compound as claimed in claim 1, wherein L₁ is:


a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophenylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a dibenzosilolylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, or an imidazopyrimidinylene group; or

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a spiro-fluorene-benzofluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a rubicenylene group, a coronenylene group, an ovalenylene group, a pyrrolylene group, a thiophe-


nylene group, a furanylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylene group, a benzothiophenylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a dibenzosilolylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, or an imidazopyrimidinylene group, each substituted with deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolylene group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), or —P(=O)(Q₃₁)(Q₃₂),

wherein Q_{31} to Q_3 are each independently a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, or a dibenzofuranyl group.


3. The polycyclic compound as claimed in claim 1, wherein L_1 is a group represented by one of Formula 3-1 to 3-49:




Formula 3-1




Formula 3-2




Formula 3-3




Formula 3-4



Formula 3-5




Formula 3-6




Formula 3-7


-continued




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11




Formula 3-12




Formula 3-13




Formula 3-14



Formula 3-15













Formula 3-16




Formula 3-17

-continued




-continued

Formula 3-18




Formula 3-19




Formula 3-20

Formula 3-21




Formula 3-22


Formula 3-23



Formula 3-24




Formula 3-25



Formula 3-26

Formula 3-27



Formula 3-28

Formula 3-29

Formula 3-30


Formula 3-31

Formula 3-32

Formula 3-33


Formula 3-34

-continued




Formula 3-35

-continued




Formula 3-40


Formula 3-36




Formula 3-41




Formula 3-37




Formula 3-42




Formula 3-38




Formula 3-43




Formula 3-39




Formula 3-43'




-continued




Formula 3-44




Formula 3-45




Formula 3-46



Formula 3-47



Formula 3-48



Formula 3-49

wherein, in Formulae 3-1 to 3-49,

Y_1 is O, S, C(Z_3)(Z_4), N(Z_5), or Si(Z_6)(Z_7),

Z_1 to Z_7 are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a triazinyl group, a benzimidazolyl group, a phenanthrolinyl group, or —Si(Q_{33})(Q_{34})(Q_{35}),

wherein Q_3 to Q_{35} are each independently a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

d_2 is an integer from 0 to 2,

d_3 is an integer from 0 to 3,

d_4 is an integer from 0 to 4,


d_5 is an integer from 0 to 5,

d_6 is an integer from 0 to 6,


d_8 is an integer from 0 to 8, and

* and *' each indicate a binding site to a neighboring atom.


4. The polycyclic compound as claimed in claim 1, wherein L_1 is a group represented by one of Formula 4-1 to 4-6:




Formula 4-1



Formula 4-2




Formula 4-3




Formula 4-4

-continued



Formula 4-5



Formula 4-6

wherein * and ** in Formulae 4-1 to 4-6 each indicate a binding site to a neighboring atom.

5. The polycyclic compound as claimed in claim 1, wherein:

X_1 is $C(R_{11})(R_{12})$, and

R_{11} and R_{12} are each independently:

a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, or a hexyl group; or

a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, or a hexyl group, each substituted with deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, a C_1 - C_{60} alkyl group, a C_6 - C_{60} aryl group, or a C_2 - C_{60} heteroaryl group.

6. The polycyclic compound as claimed in claim 1, wherein:

X_1 is $N(R_{13})$, and

Ar_2 and R_{13} are each independently:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, or a pyrenyl group;


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, or a pyrenyl group, each substituted with deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid group or a salt thereof, a sulfonic acid group or a salt thereof, a phosphoric acid group or a salt thereof, a C_1 - C_{20} alkyl group, or a C_1 - C_{20} alkoxy group, or

Ar_2 and R_{13} are linked to a neighboring substituent to form a substituted or unsubstituted benzene; a substituted or unsubstituted naphthalene; a substituted or unsubstituted fluorene, a substituted or unsubstituted spiro-fluorene, a substituted or unsubstituted pyridine, a substituted or unsubstituted pyrazine, a substituted or unsubstituted pyrimidine, a substituted or unsubstituted pyridazine, a substituted or unsubstituted quinoline, a substituted or unsubstituted isoquinoline, a substituted or unsubstituted quinoxaline, a substituted or unsubstituted quinazoline, or a substituted or unsubstituted triazine, and


at least one substituent of the substituted benzene, the substituted naphthalene, the substituted fluorene, the substituted spiro-fluorene, the substituted pyridine, the substituted pyrazine, the substituted pyrimidine, the substituted pyridazine, the substituted quinoline, the substituted isoquinoline, the substituted quinoxaline, the substituted quinazoline, and the substituted triazine is deuterium, —F, —Cl, —Br, —I, a cyano group, or a C_1 - C_{60} alkyl group.

7. The polycyclic compound as claimed in claim 1, wherein: X_1 is $N(R_{13})$, and

the compound represented by Formula 1 is represented by Formula 11 or 12:



[Formula 11]



[Formula 12]

wherein, in Formulae 11 and 12,

R_1 to R_{10} are defined the same as R_1 to R_{10} of Formula 1, ring A_1 is a substituted or unsubstituted benzene group, a substituted or unsubstituted naphthalene group, a substituted or unsubstituted fluorene group, a substituted or unsubstituted spiro-fluorene group, a substituted or unsubstituted indene group, a substituted or unsubstituted pyrrole group, a substituted or unsubstituted thiophene group, a substituted or unsubstituted furan group, a substituted or unsubstituted imidazole group, a substituted or unsubstituted pyrazole group, a substituted or unsubstituted thiazole group, a substituted or unsubstituted isothiazole group, a substituted or unsubstituted oxazole group, a substituted or unsubstituted isoxazole group, a substituted or unsubstituted pyridine group, a substituted or unsubstituted pyrazine group, a substituted or unsubstituted pyrimidine group, a substituted or unsubstituted pyridazine group, a substituted or unsubstituted quinoline group, a substituted or unsubstituted isoquinoline group, a substituted or unsubstituted benzoquinoline group, a substituted or unsubstituted quinoxaline group, a substituted or unsubstituted quinazoline group, a substituted or unsubstituted carbazole group, a substituted or unsub-

stituted benzimidazole group, a substituted or unsubstituted benzofuran group, a substituted or unsubstituted benzothiophene group, a substituted or unsubstituted isobenzothiophene group, a substituted or unsubstituted benzoxazole group, a substituted or unsubstituted isobenzoxazole group, a substituted or unsubstituted triazole group, a substituted or unsubstituted oxadiazole group, a substituted or unsubstituted triazine group, a substituted or unsubstituted dibenzofuran group, or a substituted or unsubstituted dibenzothiophene group,

R_{21} and R_{22} are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,


at least one substituent of the substituted benzene group, the substituted naphthalene group, the substituted fluorene group, the substituted spiro-fluorene group, the substituted indene group, the substituted pyrrole group, the substituted thiophene group, the substituted furan group, the substituted imidazole group, the substituted pyrazole group, the substituted thiazole group, the substituted isothiazole group, the substituted oxazole group, the substituted isoxazole group, the substituted pyridine group, the substituted pyrazine group, the substituted pyrimidine group, the substituted pyridazine group, the substituted quinoline group, the substituted isoquinoline group, the substituted benzoquinoline group, the substituted quinoxaline group, the substituted quinazoline group, the substituted carbazole group, the substituted benzimidazole group, the substituted benzofuran group, the substituted benzothiophene group, the substituted isobenzothiophene group, the substituted benzoxazole group, the substituted isobenzoxazole group, the substituted triazole group, the substituted oxadiazole group, the substituted triazine group, the substituted dibenzofuran group, the substituted dibenzothiophene group, the substituted 0 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60} alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted mon-

ovalent non-aromatic condensed polycyclic group, or the substituted monovalent non-aromatic condensed heteropolycyclic group is:

- deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, or a C_1 - C_{60} alkoxy group; or
- a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, or a terphenyl group.

8. The polycyclic compound as claimed in claim 7, wherein:

the compound is represented by Formula 11, and the compound represented by Formula 11 is represented by Formula 11-1:



wherein, in Formula 11-1,

R_1 to R_{10} are defined the same as R_1 to R_{10} of Formula 1, R_{23} is hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,

r_1 is an integer from 0 to 4,

when r_1 is two or more, two or more R_{23} (s) are identical to or different from each other and two or more neighboring R_{23} (s) are separate or are linked to form a substituted or unsubstituted ring,

at least one substituent of the substituted C_1 - C_{60} alkyl group, the substituted C_2 - C_{60} alkenyl group, the substituted C_2 - C_{60} alkynyl group, the substituted C_1 - C_{60}

alkoxy group, the substituted C_3 - C_{10} cycloalkyl group, the substituted C_1 - C_{10} heterocycloalkyl group, the substituted C_3 - C_{10} cycloalkenyl group, the substituted C_1 - C_{10} heterocycloalkenyl group, the substituted C_6 - C_{60} aryl group, the substituted C_6 - C_{60} aryloxy group, the substituted C_6 - C_{60} arylthio group, the substituted C_1 - C_{60} heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, the substituted monovalent non-aromatic condensed heteropolycyclic group, or the substituted ring is:

deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, or a C_1 - C_{60} alkoxy group; or


a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_6 - C_{60} aryloxy group, a C_6 - C_{60} arylthio group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, or a terphenyl group.

9. The polycyclic compound as claimed in claim 1, wherein Ar_2 is:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, or a chrysenyl group; or


a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a Spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, or a chrysenyl group, each substituted with deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, or a chrysenyl group.

10. The polycyclic compound as claimed in claim 1, wherein Ar_2 is a group represented by one of Formula 5-1 to 5-52:




Formula 5-1


-continued




Formula 5-2




Formula 5-3




Formula 5-4




Formula 5-5




Formula 5-6



Formula 5-7










Formula 5-8




Formula 5-9

-continued




-continued

Formula 5-10




Formula 5-17

Formula 5-11




Formula 5-18

Formula 5-12




Formula 5-19

Formula 5-13




Formula 5-14




Formula 5-20

Formula 5-15




Formula 5-21

Formula 5-16




-continued




Formula 5-22

-continued




Formula 5-28


Formula 5-23




Formula 5-29




Formula 5-24




Formula 5-30




Formula 5-25



Formula 5-31





Formula 5-26




Formula 5-32


Formula 5-27




-continued




Formula 5-33




Formula 5-34




Formula 5-35



Formula 5-36




Formula 5-37




Formula 5-38


-continued




Formula 5-39




Formula 5-40



Formula 5-41




Formula 5-42




Formula 5-43

-continued




Formula 5-44


-continued




Formula 5-49




Formula 5-45




Formula 5-50




Formula 5-46




Formula 5-51



Formula 5-47



Formula 5-52



Formula 5-48

wherein, in Formulae 5-1 to 5-52, Z_{31} to Z_{35} are each independently hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkenyl group, a C_1 - C_{20} alkynyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a pyrenyl group, a chrysenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a

pyridazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinylyl group, a quinazolinylyl group, a carbazolyl group, a triazinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, or $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

wherein Q_1 to Q_3 and Q_{31} to Q_{33} are each independently a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a phenyl group substituted with a C_1 - C_{10} alkyl group, a biphenyl group, a terphenyl group, a naphthyl group, or a dibenzofuranyl group,

e2 is 1 or 2,

e3 is an integer from 1 to 3,

e4 is an integer from 1 to 4,

e5 is an integer from 1 to 5,

e6 is an integer from 1 to 6,

e7 is an integer from 1 to 7,

e9 is an integer from 1 to 9, and


* indicates a binding site to a neighboring atom.

11. The polycyclic compound as claimed in claim 1, wherein Ar_1 is:


an imidazolyl group, a pyrazole group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylyl group, a quinazolinylyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, or an azacarbazolyl group; or

an imidazolyl group, a pyrazole group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylyl group, a quinazolinylyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, or an azacarbazolyl group, each substituted with deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, or a C_1 - C_{20} alkoxy group.


12. The polycyclic compound as claimed in claim 1, wherein Ar_1 is a group represented by one of Formula 7-1 to 7-59:




Formula 7-1




Formula 7-2




Formula 7-3




Formula 7-4




Formula 7-5




Formula 7-6




Formula 7-7




Formula 7-8



Formula 7-9













Formula 7-10




Formula 7-11


-continued




Formula 7-12




Formula 7-13




Formula 7-14




Formula 7-15




Formula 7-16




Formula 7-17




Formula 7-18




Formula 7-19



Formula 7-20



Formula 7-21



-continued

Formula 7-22

Formula 7-23

Formula 7-24

Formula 7-25

Formula 7-26

Formula 7-27

Formula 7-28

Formula 7-29


Formula 7-30

Formula 7-31


Formula 7-32

Formula 7-33


-continued




Formula 7-52




Formula 7-53




Formula 7-54




Formula 7-55




Formula 7-56



Formula 7-57



Formula 7-58



Formula 7-59

wherein, in Formulae 7-1 to 7-59,

Z_{51} and Z_{52} are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20}

alkoxy group, an imidazolyl group, a pyrazole group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quiazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, or an azacarbazolyl group,

g_2 is an integer from 0 to 2,

g_3 is an integer from 0 to 3,

g_4 is an integer from 0 to 4,


g_5 is an integer from 0 to 5,

g_6 is an integer from 0 to 6, and


* and *' each indicate a binding site to a neighboring atom.

13. The polycyclic compound as claimed in claim 1, wherein the compound represented by Formula 1 is represented by one of Formula 21 to 24:

[Formula 21]




[Formula 22]




-continued

[Formula 23]



[Formula 24]




wherein, in Formulae 21 to 24, X_1 , L_1 , Ar_1 , Ar_2 , R_1 to R_{10} , a_1 , and b_1 are defined the same as X_1 , L_1 , Ar_1 , Ar_2 , R_1 to R_{10} , a_1 , and b_1 of Formula 1.


14. The polycyclic compound as claimed in claim 1, wherein the polycyclic compound is one of Compound A-1 to A-51:

-continued

A-2





A-3




A-4

A-1




-continued

A-5




-continued


A-8



A-6





A-9




A-10

A-7




A-11




-continued


A-12




A-13



A-14




A-15




-continued


A-16




A-17



A-18




A-19




-continued


A-20




A-21



A-22




A-23




-continued


A-24




A-25



A-26




A-27




-continued

A-28




-continued


A-32




A-29



A-33





A-30




A-34

A-31




-continued

A-35





-continued

A-38





A-39

A-36



A-40

A-37





-continued

-continued



A-41

A-45





A-42

A-46




A-43

A-47




A-44




-continued


A-48




A-49



A-50



A-51



15. An organic light-emitting device, comprising:

a first electrode;

a second electrode facing the first electrode; and

an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes the polycyclic compound as claimed in claim 1.

16. The organic light-emitting device as claimed in claim 15, wherein:

the first electrode is an anode,

the second electrode is a cathode, and

the organic layer further includes:

a hole transport region between the first electrode and the emission layer, and

an electron transport region between the emission layer and the second electrode,

the hole transport region includes a hole injection layer, a hole transport layer, a buffer layer, an emission auxiliary layer, or an electron blocking layer, and

the electron transport region includes a hole blocking layer, an electron transport layer, or an electron injection layer.

17. The organic light-emitting device as claimed in claim 16, wherein the electron transport region includes the polycyclic compound.

18. The organic light-emitting device as claimed in claim 17, wherein:

the electron transport region includes an electron transport layer and an electron injection layer, and

the electron transport layer or the electron injection layer includes the polycyclic compound.

19. The organic light-emitting device as claimed in claim 18, wherein the electron transport layer or the electron injection layer further includes an alkali metal, an alkaline earth metal, a rare-earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare-earth metal compound, an alkali metal complex, an alkaline earth-metal complex, or a rare-earth metal complex.

20. The organic light-emitting device as claimed in claim 15, wherein:

the emission layer includes a dopant and a host, and

the host includes an anthracene-based compound, a pyrene-based compound, or a spiro-bifluorene-based compound.

* * * * *

专利名称(译)	多环化合物和包括其的有机发光装置		
公开(公告)号	US20180033974A1	公开(公告)日	2018-02-01
申请号	US15/437504	申请日	2017-02-21
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	LEE JUNGSUB		
发明人	LEE, JUNGSUB		
IPC分类号	H01L51/00 C07D495/06 C07D401/10 C07D491/06 C09K11/06 C09K11/02		
CPC分类号	H01L51/0067 C09K2211/185 C09K11/025 H01L51/0085 C07D401/10 H01L51/0072 H01L51/0052 C07D491/06 H01L51/0071 C07D495/06 H01L51/5072 H01L51/5092 H01L51/5012 H01L51/5016 C09K2211/1007 C09K2211/1029 C09K11/06		
优先权	1020160095488 2016-07-27 KR		
其他公开文献	US10205104		
外部链接	Espacenet USPTO		

摘要(译)

多环化合物和有机发光器件，该化合物由式1表示：

10

190
150
110