



US 20190140184A1

(19) **United States**

(12) **Patent Application Publication**
HAN et al.

(10) **Pub. No.: US 2019/0140184 A1**
(43) **Pub. Date: May 9, 2019**

(54) **CONDENSED-CYCLIC COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

C07D 407/14 (2006.01)
C07D 409/14 (2006.01)

(52) **U.S. Cl.**
CPC *H01L 51/0061* (2013.01); *C07D 319/14* (2013.01); *H01L 51/5012* (2013.01); *C07D 409/14* (2013.01); *C07D 407/14* (2013.01)

(71) Applicant: **SAMSUNG DISPLAY CO., LTD.**,
Yongin-Si (KR)

(72) Inventors: **Sanghyun HAN**, Yongin-si (KR);
Sooyon KIM, Yongin-si (KR);
Youngkook KIM, Yongin-si (KR);
Jongwoo KIM, Yongin-si (KR); **Hyejin JUNG**,
Yongin-si (KR); **Seokhwan HWANG**,
Yongin-si (KR)

(57) **ABSTRACT**

A condensed cyclic compound and an organic light-emitting device including the same. The a condensed cyclic compound represented by Formula 1:

(21) Appl. No.: **16/182,827**

(22) Filed: **Nov. 7, 2018**


(30) **Foreign Application Priority Data**

Nov. 7, 2017 (KR) 10-2017-0147603

Publication Classification

(51) **Int. Cl.**
H01L 51/00 (2006.01)
C07D 319/14 (2006.01)

<Formula 1>



10

190
150
110

FIG. 1

10

190
150
110

FIG. 2

20

190
150
110
210

FIG. 3

30

220
190
150
110

FIG. 4

40

220
190
150
110
210

**CONDENSED-CYCLIC COMPOUND AND
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] This application claims priority under 35 U.S.C. § 119 to Korean Patent Application No. 10-2017-0147603, filed on Nov. 7, 2017, in the Korean Intellectual Property Office, the disclosure of which is incorporated by reference herein in its entirety.

1. TECHNICAL FIELD

[0002] Exemplary embodiments of the present invention relate to a condensed cyclic compound, and more particularly to an organic light-emitting device including the same.

2. DISCUSSION OF RELATED ART


[0003] Organic light-emitting devices may be self-emission devices that produce full-color images, and also have relatively wide viewing angles, relatively high contrast ratios, relatively short response times, relatively high brightness, and relatively fast response speeds.

[0004] An example of such organic light-emitting devices may include a first electrode located on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode, which are sequentially stacked on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. Carriers, such as holes and electrons, may recombine in the emission layer to produce excitons. These excitons may fall from an excited state to a ground state, thus generating light.

SUMMARY

[0005] An exemplary embodiment of the present invention provides a condensed cyclic compound and an organic light-emitting device including the same.

[0006] According to an exemplary embodiment of the present invention, a condensed cyclic compound is represented by Formula 1:



[0007] In Formulae 1 and 2,

[0008] X_1 and X_2 may each independently be O or S,

[0009] L_1 , L_2 , and L_{11} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene, a substituted or unsubstituted C_6 - C_{60} arylene, a substituted or unsubstituted C_1 - C_{60} heteroarylene, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0010] a_1 , a_2 , and a_{11} may each independently be 0, 1, 2, 3, 4, or 5,

[0011] Ar_1 and Ar_2 may each independently be selected from a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0012] R_1 to R_{10} may each independently be selected from a group represented by Formula 2, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group (aryloxy), a substituted or unsubstituted C_6 - C_{60} arylthio group (arylthio), a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, $-B(Q_1)(Q_2)$, $-C(=O)(Q_1)$, $-S(=O)_2(Q_1)$, and $-P(=O)(Q_1)(Q_2)$, and at least two selected from R_1 to R_{10} may each be a group represented by Formula 2,

[0013] at least one selected from the substituted C_3 - C_{10} cycloalkylene group, substituted C_1 - C_{10} heterocycloalkylene group, substituted C_3 - C_{10} cycloalkenylene group, substituted C_1 - C_{10} heterocycloalkenylene group, substituted C_6 - C_{60} arylene group, substituted C_1 - C_{60} heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C_1 - C_{60} alkyl group, substituted C_2 - C_{60} alkenyl group, substituted C_2 - C_{60} alkynyl group, substituted C_1 - C_{60} alkoxy group, substituted C_3 - C_{10} cycloalkyl group, substituted C_1 - C_{10} heterocycloalkyl group, substituted C_3 - C_{10} cycloalkenyl group, substituted C_1 - C_{10} heterocycloalkenyl group, substituted C_6 - C_{60} aryl group, substituted C_6 - C_{60} aryloxy group, substituted C_6 - C_{60} arylthio group, substituted C_1 - C_{60} heteroaryl group, substituted monovalent non-aromatic condensed polycyclic

group, and substituted monovalent non-aromatic condensed heteropolycyclic group may be: selected from:

[0014] deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0015] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₁₁)(Q₁₂)(Q₁₃), -N(Q₁₁)(Q₁₂), -B(Q₁₁)(Q₁₂), -C(=O)(Q₁₁), -S(=O)₂(Q₁₁), and -P(=O)(Q₁₁)(Q₁₂);

[0016] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0017] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q₂₁)(Q₂₂)(Q₂₃), -N(Q₂₁)(Q₂₂), -B(Q₂₁)(Q₂₂), -C(=O)(Q₂₁), -S(=O)₂(Q₂₁), and -P(=O)(Q₂₁)(Q₂₂); or

[0018] -Si(Q₃₁)(Q₃₂)(Q₃₃), -N(Q₃₁)(Q₃₂), -B(Q₃₁)(Q₃₂), -C(=O)(Q₃₁), -S(=O)₂(Q₃₁), or -P(=O)(Q₃₁)(Q₃₂),

[0019] Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, or a terphenyl group.

[0020] According to an exemplary embodiment of the present invention, an organic light-emitting device includes a first electrode. A second electrode faces the first electrode. An organic layer is between the first electrode and the second electrode. The organic layer includes an emission layer and at least one of the condensed cyclic compounds represented by Formula 1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] The above and other features of the present invention will become more apparent by describing in detail exemplary embodiments thereof with reference to the accompanying drawings, in which:

[0022] FIG. 1 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention;

[0023] FIG. 2 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention;


[0024] FIG. 3 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention; and

[0025] FIG. 4 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention.

DETAILED DESCRIPTION


[0026] Exemplary embodiments of the present invention will be described below in more detail with reference to the accompanying drawings. In this regard, the exemplary embodiments may have different forms and should not be construed as being limited to the exemplary embodiments of the present invention described herein.

[0027] A condensed cyclic compound according to an exemplary embodiment of the present invention is represented by Formula 1 below:



[0028] X₁ and X₂ in Formula 1 may each independently be O or S. In an exemplary embodiment of the present invention, X₁ may be O, and X₂ may be S. In an exemplary embodiment of the present invention, X₁ and X₂ may each be O, but exemplary embodiments of the present invention are not limited thereto.

[0029] Two or more selected from R₁ to R₁₀ in Formula 1 may each be a group represented by Formula 2:



[0030] L_1 , L_2 , and L_{11} in Formula 2 may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group. In an exemplary embodiment of the present invention, L_1 , L_2 , and L_{11} may each independently be selected from:

[0031] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzoisothiazolylene group, a benzoxazolylene group, a benzoisoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylene group, a dibenzosilolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group; and


[0032] a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a pyrrolylenylene group, a thiophenylenylene group, a furanylenylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylenylene group, a pyrazinylenylene group, a pyrimidinylenylene group, a pyridazinylenylene group, an

isoindolylene group, an indolylene group, an indazolylene group, a purinylenylene group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolylene group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a benzoisothiazolylene group, a benzoxazolylene group, a benzoisoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylene group, a dibenzosilolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylenylene group, and an imidazopyrimidinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinylenylene group, an isoquinolinylenylene group, a benzoquinolinylenylene group, a phthalazinylenylene group, a naphthyridinylenylene group, a quinoxalinylenylene group, a quinazolinylenylene group, a cinnolinylenylene group, a carbazolyl group, a phenanthridinylenylene group, an acridinylenylene group, a phenanthrolinylenylene group, a phenazinylenylene group, a benzimidazolyl group, a benzofuranylenylene group, a benzothiophenyl group, a benzoisothiazolyl group, a benzoxazolyl group, a benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranylenylene group, a dibenzothiophenylene group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinylenylene group, an imidazopyrimidinylenylene group, —Si(Q_{31})(Q_{32})(Q_{33}), —N(Q_{31})(Q_{32}), —C(=O)(Q_{31}), —S(=O)₂(Q_{31}), and —P(=O)(Q_{31})(Q_{32}); and


[0033] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, or a carbazolyl group.

[0034] In an exemplary embodiment of the present invention, L_1 , L_2 , and L_{11} may each independently be selected from groups represented by Formulae 3-1 to 3-28:


-continued




Formula 3-1




Formula 3-2




Formula 3-3




Formula 3-4




Formula 3-5




Formula 3-6




Formula 3-7




Formula 3-8




Formula 3-9




Formula 3-10




Formula 3-11




Formula 3-12




Formula 3-13




Formula 3-14




Formula 3-15




Formula 3-16




Formula 3-17



Formula 3-18




Formula 3-19




Formula 3-20


-continued




Formula 3-21




Formula 3-22




Formula 3-23




Formula 3-24




Formula 3-25



Formula 3-26



Formula 3-27



Formula 3-28

[0035] In Formulae 3-1 to 3-28,
 [0036] Y₁ may be O, S, C(Z₃)(Z₄), N(Z₅), or Si(Z₆)(Z₇),
 [0037] Z₁ to Z₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a

spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃),

[0038] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

[0039] d₂ may be an integer selected from 1 or 2,

[0040] d₃ may be an integer selected from 1 to 3,


[0041] d₄ may be an integer selected from 1 to 4,

[0042] d₅ may be an integer selected from 1 to 5,


[0043] d₆ may be an integer selected from 1 to 6,

[0044] d₈ may be an integer selected from 1 to 8, and


[0045] * and *' each indicate a binding site to a neighboring atom. In an exemplary embodiment of the present invention, L₁, L₂, and L₁₁ may each independently be selected from groups represented by Formulae 4-1 to 4-15, but exemplary embodiments of the present invention are not limited thereto:




Formular 4-1




Formular 4-2




Formular 4-3



Formular 4-4











Formular 4-5



Formular 4-6

-continued



Formular 4-7

Formular 4-8

Formular 4-9

Formular 4-10

Formular 4-11


Formular 4-12

Formular 4-13

Formular 4-14

-continued

Formular 4-15



[0046] * and * in Formulae 4-1 to 4-15 each indicate a binding site to a neighboring atom.

[0047] a₁, a₂, and a₁₁ in Formula 2 may each independently be an integer selected from 0, 1, 2, 3, 4, or 5. In an exemplary embodiment of the present invention, a₁ indicates the number of L₁(s) in Formula 2, and when a₁ is 0, *(L₁)_{a₁}* is a single bond, and when a₁ is two or more, two or more of L₁(s) may be identical to or different from each other. a₂ and a₁₁ may each be the same as a₁ described above (see, e.g., the description of a₁ and Formula 2 above).

[0048] In an exemplary embodiment of the present invention, a₁, a₂, and a₁₁ may each independently be an integer selected from 0, 1, or 2. In an exemplary embodiment of the present invention, a₁, a₂, and a₁₁ may each independently be an integer selected from 0 or 1, but exemplary embodiments of the present invention are not limited thereto.

[0049] Ar₁ and Ar₂ in Formula 2 may each independently be selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

[0050] In an exemplary embodiment of the present invention, Ar₁ and Ar₂ may each independently be selected from:

[0051] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spirobifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinoliny group, an isoquinoliny group, a benzoquinoliny group, a phthalazinyl group, a naphthyridinyl group, a quinoxaliny group, a quinazoliny group, a cinnoliny group, a carbazolyl group, a phenan-


thridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothiophenyl group, an benzisothiazolyl group, a benzoxazolyl group, an benzisoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group; and

[0052] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl


group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranlyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzosilolyl group, a dibenzofuranlyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0053] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, and a carbazolyl group.


[0054] In an exemplary embodiment of the present invention, Ar₁ and Ar₂ may each independently be selected from groups represented by Formulae 5-1 to 5-24:




Formular 5-1




Formular 5-2




Formular 5-3




Formular 5-4



Formular 5-5













Formular 5-6



Formular 5-7

-continued



Formular 5-8

Formular 5-9

Formular 5-10

Formular 5-11

Formular 5-12

Formular 5-13








Formular 5-14

Formular 5-15

Formular 5-16

Formular 5-17

-continued



Formular 5-18

Formular 5-19

Formular 5-20

Formular 5-21

Formular 5-22

Formular 5-23

Formular 5-24

[0055] In Formulae 5-1 to 5-24,

[0056] Y_{31} may be O, S, C(Z_{33})(Z_{34}), N(Z_{35}), Or Si(Z_{36})(Z_{37}),

[0057] Z_{31} to Z_{37} may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and —Si(Q_{31})(Q_{32})(Q_{33}).

[0058] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group,

[0059] e2 may be an integer selected from 1 or 2,

[0060] e3 may be an integer selected from 1 to 3,


[0061] e4 may be an integer selected from 1 to 4,

[0062] e5 may be an integer selected from 1 to 5,


[0063] e6 may be an integer selected from 1 to 6,

[0064] e7 may be an integer selected from 1 to 7, and


[0065] * indicate a binding site to a neighboring atom. In an exemplary embodiment of the present invention, Ar_1 and Ar_2 may each independently be selected from groups represented by Formulae 6-1 to 6-25, but exemplary embodiments of the present invention are not limited thereto:




Formular 6-1




Formular 6-2




Formular 6-3




Formular 6-4




Formular 6-5



Formular 6-6




Formular 6-7




-continued


Formular 6-8




Formular 6-9




Formular 6-10




Formular 6-11




Formular 6-12




Formular 6-13



Formular 6-14




Formular 6-15




Formular 6-16


-continued




Formular 6-17




Formular 6-18




Formular 6-19




Formular 6-20




Formular 6-21



Formular 6-22




Formular 6-23



Formular 6-24

-continued



Formular 6-25

[0066] * in Formulae 6-1 to 6-25 indicates a binding site to a neighboring atom.

[0067] R_1 to R_{10} in Formula 1 may each independently be selected from a group represented by Formula 2, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_1)(Q_2)(Q_3)$, $-B(Q_1)(Q_2)$, $-C(=O)(Q_1)$, $-S(=O)_2(Q_1)$, and $-P(=O)(Q_1)(Q_2)$, and at least two selected from R_1 to R_{10} may be the group represented by Formula 2, and

[0068] Q_1 to Q_3 may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{60} alkyl group, a C_2 - C_{60} alkenyl group, a C_2 - C_{60} alkynyl group, a C_1 - C_{60} alkoxy group, a C_3 - C_{10} cycloalkyl group, a C_1 - C_{10} heterocycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a C_1 - C_{10} heterocycloalkenyl group, a C_6 - C_{60} aryl group, a C_1 - C_{60} heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0069] In an exemplary embodiment of the present invention, R_1 to R_{10} may each independently be selected from:

[0070] the group represented by Formula 2, hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0071] a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, and $-Si(Q_{31})(Q_{32})(Q_{33})$;

[0072] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a

triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group;

[0073] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C₁-C₂₀ alkyl group, and a C₁-C₂₀ alkoxy group, and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

[0074] —Si(Q₁)(Q₂)(Q₃), and

[0075] Q₁ to Q₃ and Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0076] In an exemplary embodiment of the present invention, R₁ to R₁₀ may each independently be selected from the group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and —Si(Q₁)(Q₂)(Q₃), and

[0077] Q₁ to Q₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0078] In an exemplary embodiment of the present invention, R₁ to R₁₀ may each independently be selected from the group represented by Formula 2, hydrogen, and deuterium, but exemplary embodiments of the present invention are not limited thereto.

[0079] In an exemplary embodiment of the present invention, two selected from R₁ to R₁₀ in Formula 1 may each be the group represented by Formula 2,

[0080] the other substituents (e.g., substituents that are selected from R₁ to R₁₀ and are not the group represented by Formula 2) may be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and —Si(Q₁)(Q₂)(Q₃), and


[0081] Q₁ to Q₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0082] In an exemplary embodiment of the present invention, two selected from R₁ to R₁₀ may each be the group represented by Formula 2, and the other substituents (e.g., substituents that are selected from R₁ to R₁₀ and are not the

group represented by Formula 2) may each be hydrogen, but exemplary embodiments of the present invention are not limited thereto.

[0083] In an exemplary embodiment of the present invention, the condensed cyclic compound may be represented by Formula 1A below:

<Formula 1A>



[0084] In Formula 1A,

[0085] according to an exemplary embodiment of the present invention, X₁, X₂, L₁, L₂, L₁₁, a₁, a₂, a₁₁, Ar₁, Ar₂, and R₃ to R₁₀ are the same as described above, and L₃, L₄, L₁₂, a₃, a₄, a₁₂, Ar₃, and Ar₄ are the same as L₁, L₂, L₁₁, a₁, a₂, a₁₁, Ar₁ and Ar₂, respectively.

[0086] In an exemplary embodiment of the present invention, in Formula 1A,

[0087] X₁ and X₂ may each be O;

[0088] a₁ to a₄, a₁₁, and a₁₂ may each independently be selected from an integer selected from 0 or 1;

[0089] R₃ to R₁₀ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and —Si(Q₁)(Q₂)(Q₃), and

[0090] Q₁ to Q₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0091] In an exemplary embodiment of the present invention, in Formula 1A,


[0092] X₁ and X₂ may each be O; and

[0093] a₁ to a₄, a₁₁, and a₁₂ may each independently be 0 or 1; and


[0094] R₃ to R₁₀ may each be hydrogen, but exemplary embodiments of the present invention are not limited thereto.

[0095] In an exemplary embodiment of the present invention, the condensed cyclic compound may be represented by Formula 1Aa or 1Ab:

<Formula 1Aa>



<Formula 1Ab>



[0096] In Formulae 1Aa and 1Ab,

[0097] according to an exemplary embodiment of the present invention, X_1 , X_2 , L_1 , L_2 , L_{11} , a_1 , a_2 , a_{11} , Ar_1 , Ar_2 , and R_3 to R_{10} are the same as described above,

[0098] L_3 , L_4 , L_{12} , a_3 , a_4 , a_{12} , Ar_3 , and Ar_4 are the same as L_1 , L_2 , L_{11} , a_1 , a_2 , a_{11} , Ar_1 , and Ar_2 , respectively,

[0099] Z_1 may be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro

group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, and $-Si(Q_{31})(Q_{32})(Q_{33})$.

[0100] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, and

[0101] d_4 may be an integer selected from 1 to 4.

[0102] In an exemplary embodiment of the present invention, in Formulae 1Aa and 1Ab,

[0103] X_1 and X_2 may each be O;

[0104] a_1 to a_4 and a_{12} may each independently be an integer selected from 0 or 1;

[0105] R_3 to R_{10} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, and $-Si(Q_1)(Q_2)(Q_3)$, and

[0106] Q_1 to Q_3 may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0107] In an exemplary embodiment of the present invention, in Formulae 1Aa and 1Ab,


[0108] X_1 and X_2 may each be O;

[0109] a_1 to a_4 and a_{12} may each independently be an integer selected from 0 or 1; and


[0110] R_3 to R_{10} may each be hydrogen, but exemplary embodiments of the present invention are not limited thereto.

[0111] The condensed cyclic compound may be one of Compounds 1 to 48, but exemplary embodiments of the present invention are not limited thereto:

1





2



-continued



3

4





5

6









7

8





-continued



-continued



15

16





17

18



19


20




-continued

27



28




29



30





31



32



33



-continued



34

35





36

37



38



39



-continued



40

41





42

43





44

45




46

47



-continued

48



[0112] The condensed cyclic compound represented by Formula 1 has excellent charge transport ability and thermal stability. Accordingly, an organic light-emitting device using the condensed cyclic compound represented by Formula 1 may have relatively low driving voltage, relatively high efficiency, relatively high luminance, and a relatively long lifespan.

[0113] An exemplary synthesis method for the condensed cyclic compound represented by Formula 1 is provided below.

[0114] At least one condensed cyclic compound of Formula 1 may be positioned between a pair of electrodes of an organic light-emitting device. For example, the condensed cyclic compound may be included in at least one of a hole transport region and an emission layer. In an exemplary embodiment of the present invention, the condensed cyclic compound of Formula 1 may be used as a material for a capping layer located outside a pair of electrodes of an organic light-emitting device.

[0115] Accordingly, an organic light-emitting device according to an exemplary embodiment of the present invention may include a first electrode, a second electrode facing the first electrode, and an organic layer between the first electrode and the second electrode. The organic layer may include an emission layer. The organic layer may include the condensed cyclic compound represented by Formula 1.

[0116] According to an exemplary embodiment of the present invention, the organic layer may include more than one condensed cyclic compound represented by Formula 1. For example, two or more identical compounds represented by Formula 1 may be included in the organic layer. Alternatively, two or more compounds represented by Formula 1 that are different from each other may be included in the organic layer.

[0117] For example, the organic layer may include, as the condensed cyclic compound, only Compound 1. Compound 1 may be in a hole transport layer of the organic light-emitting device. In an exemplary embodiment of the present invention, the organic layer may include, as the condensed cyclic compound, Compound 1 and Compound 2. Compound 1 and Compound 2 may exist in a same layer as each other (for example, Compound 1 and Compound 2 may all exist in a hole transport layer), or different layers (for

example, Compound 1 may exist in a hole transport layer and Compound 2 may exist in an emission layer).

[0118] The organic layer may include i) a hole transport region that is disposed between the first electrode (anode) and the emission layer and includes at least one of a hole injection layer, a hole transport layer, an emission auxiliary layer, and an electron blocking layer, and ii) an electron transport region that is disposed between the emission layer and the second electrode (cathode) and includes at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer. The hole transport region may include at least one condensed cyclic compound represented by Formula 1. For example, the hole transport region of the organic light-emitting device may include a hole transport layer, and the hole transport layer may include at least one condensed cyclic compound represented by Formula 1.

[0119] The organic light-emitting device may include at least one selected from a first capping layer located in a pathway along which light generated in an emission layer proceeds toward the outside through the first electrode and a second capping layer located in a pathway along which light generated in an emission layer proceeds toward the outside through the second electrode. The first capping layer and the second capping layer may each include at least one condensed cyclic compound represented by Formula 1.

[0120] For example, the organic light-emitting device may have a stacked structure including a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order, a stacked structure including a first capping layer, a first electrode, an organic layer, and a second electrode which are sequentially stacked in this stated order, or a stacked structure including a first capping layer, a first electrode, an organic layer, a second electrode, and a second capping layer which are sequentially stacked in this stated order. The first capping layer and the second capping layer may each include the condensed cyclic compound represented by Formula 1.

[0121] The organic layer may have a single layer structure or a multi-layered structure. For example, the organic layer may include a plurality of layers located between the first electrode and the second electrode of an organic light-emitting device. However, exemplary embodiments of the present invention are not limited thereto, and the organic layer may include materials other than an organic material.

[0122] FIG. 1 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention.

[0123] Referring to FIG. 1, an organic light-emitting device 10 may include a first electrode 110, an organic layer 150, and a second electrode 190.

[0124] The structure of the organic light-emitting device 10 according to an exemplary embodiment of the present invention and a method of manufacturing the organic light-emitting device 10 will be described in more detail below with reference to FIG. 1.

[0125] Referring to FIG. 1, a substrate may be located under the first electrode 110 or above the second electrode 190. The substrate may have relatively strong mechanical strength, relatively high thermal stability, relatively high transparency, relatively high surface smoothness, relatively high ease of handling, and relatively high water resistance. For example, the substrate may be a glass substrate or a plastic substrate.

[0126] The first electrode 110 may be formed by depositing or sputtering a material for forming the first electrode 110 on the substrate. When the first electrode 110 is an anode, the material for a first electrode may be selected from materials with a relatively high work function to facilitate hole injection.

[0127] The first electrode 110 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode 110 is a transmissive electrode, a material for forming a first electrode may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO₂), zinc oxide (ZnO), and any combinations thereof, but exemplary embodiments of the present invention are not limited thereto. When the first electrode 110 is a semi-transmissive electrode or a reflective electrode, as a material for forming the first electrode 110, magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), or any combination thereof may be used. However, the material for forming the first electrode 110 is not limited thereto.

[0128] The first electrode 110 may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode 110 may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode 110 is not limited thereto.

[0129] The organic layer 150 may be located on the first electrode 110. The organic layer 150 may include an emission layer.

[0130] The organic layer 150 may further include a hole transport region between the first electrode 110 and the emission layer, and an electron transport region between the emission layer and the second electrode 190.

[0131] The hole transport region may have a single-layered structure including a single layer including a single material, a single-layered structure including a single layer including a plurality of different materials, or a multi-layered


structure having a plurality of layers in which each of the plurality of layers include different materials from each other.

[0132] The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, or an electron blocking layer (EBL).


[0133] For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure. In the multi-layered structure, each of the layers may be sequentially stacked on the first electrode 110 in the stated order, but the structure of the hole transport region is not limited thereto.

[0134] The hole transport region may include at least one condensed cyclic compound represented by Formula 1.

[0135] In an exemplary embodiment of the present invention, the hole transport region may include at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), 13-NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4"-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:




-continued




TDATA


-continued




β -NPB




2-TNATA




TPD




Spiro-TPD



Spiro-NPB



NPB



methylated NPB

zylene group, a dibenzocarbazolyene group, a dibenzosilolyene group, and a pyridinylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃) and —N(Q₃₁)(Q₃₂), and

[0147] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group.

[0148] In an exemplary embodiment of the present invention, xa1 to xa4 may each independently be an integer selected from 0, 1, or 2.

[0149] In an exemplary embodiment of the present invention, xa5 may be an integer selected from 1, 2, 3, or 4.

[0150] In an exemplary embodiment of the present invention, R₂₀₁ to R₂₀₄ and Q₂₀₁ may each independently be selected from a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

[0151] a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl

group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), and —N(Q₃₁)(Q₃₂), and

[0152] Q₃₁ to Q₃₃ may be the same as described above.

[0153] In an exemplary embodiment of the present invention, at least one selected from R₂₀₁ to R₂₀₃ in Formula 201 may be selected from:

[0154] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a benzofuranyl group, and a dibenzothiophenyl group; and

[0155] a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a benzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; but exemplary embodiments of the present invention are not limited thereto.

[0156] In an exemplary embodiment of the present invention, in Formula 202, R₂₀₁ and R₂₀₂ may be chemically bonded to each other via a single bond, and/or R₂₀₃ and R₂₀₄ may be chemically bonded to each other via a single bond.


[0157] In an exemplary embodiment of the present invention, at least one selected from R₂₀₁ to R₂₀₄ in Formula 202 may be selected from:

[0158] a carbazolyl group; and


[0159] a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl

group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothio-phenyl group, but exemplary embodiments of the present invention are not limited thereto.


[0160] The compound represented by Formula 201 may be represented by Formula 201A:




[0161] In an exemplary embodiment of the present invention, the compound represented by Formula 201 may be represented by Formula 201A(1) below, but exemplary embodiments of the present invention are not limited thereto:




[0162] In an exemplary embodiment of the present invention, the compound represented by Formula 201 may be represented by Formula 201A-1 below, but exemplary embodiments of the present invention are not limited thereto:



[0163] In an exemplary embodiment of the present invention, the compound represented by Formula 202 may be represented by Formula 202A:



[0164] In an exemplary embodiment of the present invention, the compound represented by Formula 202 may be represented by Formula 202A-1:



[0165] In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1,


[0166] L₂₀₁ to L₂₀₃, xa1 to xa3, xa5, and R₂₀₂ to R₂₀₄ may be the same as described above,

[0167] R₂₁₁ and R₂₁₂ may be the same as in R₂₀₃, as described above,


[0168] R₂₁₃ to R₂₁₇ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C₁-C₁₀ alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiofenyl group, a dibenzofuranyl group, a dibenzothiofenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0169] The hole transport region may include at least one compound selected from Compounds HT1 to HT39, but exemplary embodiments of the present invention are not limited thereto:


HT1




HT2



HT3





HT4





-continued
HT9

HT10





HT11

HT12





-continued
HT13

HT14





HT15

HT16




HT17


HT18




-continued
HT19




HT20



HT21




HT22





-continued

HT23





HT24

HT25





HT26

HT27





-continued
HT28

HT29





HT30

HT31





HT32

HT33





HT34

HT35





-continued
HT36

HT37



HT38

HT39



[0170] A thickness of the hole transport region may be in a range of from about 100 Å to about 10,000 Å, for example, from about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer or a hole transport layer, a thickness of the hole injection layer may be in a range of from about 100 Å to about 9,000 Å, for example, from about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of from about 50 Å to about 2,000 Å, for example from about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, desired hole transporting characteristics may be obtained without a substantial increase in driving voltage.

[0171] The emission auxiliary layer may increase light-emission efficiency by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer, and the electron blocking layer may substantially block the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may include the materials as described above.

[0172] The hole transport region may include a charge-generation material for increasing conductive properties of the hole transport region. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

[0173] The charge-generation material may be, for example, a p-dopant.

[0174] In an exemplary embodiment of the present invention, a lowest unoccupied molecular orbital (LUMO) energy level of the p-dopant may be -3.5 eV or less.

[0175] The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but exemplary embodiments of the present invention are not limited thereto.


[0176] For example, the p-dopant may include at least one selected from:

[0177] a quinone derivative, such as tetracyanoquinodimethane (TCNQ) or 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);


[0178] a metal oxide, such as tungsten oxide or molybdenum oxide;

[0179] 1,4,5,8,9,12-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

[0180] a compound represented by Formula 221 below, but exemplary embodiments of the present invention are not limited thereto:



<Formula 221>



[0181] In Formula 221,

[0182] R_{221} to R_{223} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and at least one selected from R_{221} to R_{223} may include at least one substituent selected from a cyano group, $-F$, $-Cl$, $-Br$, $-I$, a C_1 - C_{20} alkyl group substituted with $-F$, a C_1 - C_{20} alkyl group substituted with $-Cl$, a C_1 - C_{20} alkyl group substituted with $-Br$, or a C_1 - C_{20} alkyl group substituted with $-I$.

[0183] When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In an exemplary embodiment of the present invention, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, or a blue emission layer, in which the two or more

layers are in direct contact with each other or are separated from each other. In an exemplary embodiment of the present invention, the emission layer may include two or more materials selected from a red light-emitting material, a green light-emitting material, and a blue light-emitting material, in which the two or more materials are mixed with each other in a single layer to emit white light.

[0184] The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

[0185] An amount of the dopant in the emission layer may be, in general, in a range of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host, but exemplary embodiments of the present invention are not limited thereto.

[0186] A thickness of the emission layer may be in a range of from about 100 Å to about 1,000 Å, for example, from about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.

[0187] In an exemplary embodiment of the present invention, the host may include a compound represented by Formula 301 below.



[0188] In Formula 301,

[0189] Ar_{301} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group, or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0190] $xb11$ may be an integer selected from 1, 2, or 3;

[0191] L_{301} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

[0192] $xb1$ may be an integer selected from 0 to 5;

[0193] R_{301} may be selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{60} alkyl group, a substituted or unsubstituted C_2 - C_{60} alkenyl group, a substituted or unsubstituted C_2 - C_{60} alkynyl group, a substituted or unsubstituted C_1 - C_{60} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{301})(Q_{302})(Q_{303})$, $-N(Q_{301})(Q_{302})$, $-B(Q_{301})(Q_{302})$, $-C(=O)(Q_{301})$, $-S(=O)_2(Q_{301})$, and $-P(=O)(Q_{301})(Q_{302})$;

[0194] $xb21$ may be an integer selected from 1 to 5; and

[0195] Q_{301} to Q_{303} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but exemplary embodiments of the present invention are not limited thereto.

[0196] In an exemplary embodiment of the present invention, Ar_{301} in Formula 301 may be selected from:

[0197] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group; and

[0198] a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothiophene group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$; and

[0199] Q_{31} to Q_{33} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but exemplary embodiments of the present invention are not limited thereto.

[0200] When $xb11$ in Formula 301 is two or more, two or more $Ar_{301}(s)$ may be chemically bonded to each other via a single bond.

[0201] In an exemplary embodiment of the present invention, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:

[0202] In Formulae 301-1 and 301-2,

[0203] A_{301} to A_{304} may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group,

[0204] X_{301} may be O, S, or $N-[(L_{304})_{xb4}-R_{304}]$,

[0205] R_{311} to R_{314} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, $-Si(Q_{31})(Q_{32})(Q_{33})$, $-N(Q_{31})(Q_{32})$, $-B(Q_{31})(Q_{32})$, $-C(=O)(Q_{31})$, $-S(=O)_2(Q_{31})$, and $-P(=O)(Q_{31})(Q_{32})$,

[0206] $xb22$ and $xb23$ may each independently be an integer selected from 0, 1, or 2,

[0207] L_{301} , $xb1$, R_{301} , and Q_{31} to Q_{33} may be the same as described above,

[0208] L_{302} to L_{304} may each independently be the same as L_{301} ,


[0209] $xb2$ to $xb4$ may each independently be the same as $xb1$, and

[0210] R_{302} to R_{304} may each independently be the same as R_{301} .


[0211] For example, L_{301} to L_{304} in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0212] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a

<Formula 301-1>



<Formula 301-2>



thiophenylene group, a furanylene group, a carbazolyene group, an indolyene group, an isoindolyene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolyene group, a dibenzocarbazolyene group, a dibenzosilolyene group, a pyridinylene group, an imidazolyene group, a pyrazolyene group, a thiazolyene group, an isothiazolyene group, an oxazolyene group, an isoxazolyene group, a thiadiazolyene group, an oxadiazolyene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolyene group, an benzoisothiazolyene group, a benzoxazolyene group, an benzoisoxazolyene group, a triazolyene group, a tetrazolyene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolyene group; and

[0213] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenyene group, a fluoranthenyene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenyene group, a hexacenyene group, a pentacenyene group, a thiophenylene group, a furanylene group, a carbazolyene group, an indolyene group, an isoindolyene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolyene group, a dibenzocarbazolyene group, a dibenzosilolyene group, a pyridinylene group, an imidazolyene group, a pyrazolyene group, a thiazolyene group, an isothiazolyene group, an oxazolyene group, an isoxazolyene group, a thiadiazolyene group, an oxadiazolyene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolyene group, an benzoisothiazolyene group, a benzoxazolyene group, an benzoisoxazolyene group, a triazolyene group, a tetrazolyene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolyene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyyl group, a pentacenyyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a

imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0214] Q₃₁ to Q₃₃ may be the same as described above.

[0215] In an exemplary embodiment of the present invention, R₃₀₁ to R₃₀₄ in Formulae 301, 301-1, and 301-2 may each independently be selected from:

[0216] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyyl group, a pentacenyyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0217] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyyl group, a pentacenyyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolyl group, an isoquinolyl group, a benzoquinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalyl group, a quinazolyl group, a cinnolyl group, a phenanthridinyl group, an acridinyl group, a


phenanthrolyl group, a phenaziny group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolyl group, a phenaziny group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂), and

[0218] Q₃₁ to Q₃₃ may be the same as described above.


[0219] In an exemplary embodiment of the present invention, the host may include an alkaline earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55), a Mg complex, and a Zn complex.

[0220] The host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and Compounds H1 to H55, but exemplary embodiments of the present invention are not limited thereto:


-continued




H2




H3




H4




H5



H6




H1




H7


-continued



-continued




-continued




H19

-continued




H23


H20



H24





H21




H25


H22




-continued




H26




H27




H28




-continued




H29



H30




H31




H32

-continued


H33




H34



H35




H36




-continued


H37




H38




H39




H40




-continued



H41




H42




H43


-continued




H44




H45



H46




H47




H48


-continued




H49




H50



H51




H52




H53

-continued




H54




H55

[0221] The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:



<Formula 401>



<Formula 402>

[0222] In Formulae 401 and 402,
 [0223] M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),
 [0224] L_{401} may be selected from ligands represented by Formula 402, and $xc1$ may be an integer selected from 1, 2, or 3. When $xc1$ is two or more, two or more $L_{401}(s)$ may be identical to or different from each other,
 [0225] L_{402} may be an organic ligand, and $xc2$ may be an integer selected from 0 to 4. When $xc2$ is two or more, two or more $L_{402}(s)$ may be identical to or different from each other,

[0226] X_{401} to X_{404} may each independently be nitrogen or carbon,

[0227] X_{401} and X_{403} may be chemically bonded to each other via a single bond or a double bond, and X_{402} and X_{404} may be chemically bonded to each other via a single bond or a double bond,

[0228] A_{401} and A_{402} may each independently be a C_5 - C_{60} carbocyclic group or a C_1 - C_{60} heterocyclic group,

[0229] X_{405} may be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{411})-*$, $*-C(Q_{411})(Q_{412})-*$, $*-C(Q_{411})=C(Q_{412})-*$, $*-C(Q_{411})=*$, or $*=C-*$, Q_{411} and Q_{412} may be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

[0230] X_{406} may be a single bond, O, or S,

[0231] R_{401} and R_{402} may each independently be selected from hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C_1 - C_{20} alkyl group, a substituted or unsubstituted C_1 - C_{20} alkoxy group, a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$, and Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a C_6 - C_{20} aryl group, and a C_1 - C_{20} heteroaryl group,

[0232] $xc11$ and $xc12$ may each independently be an integer selected from 0 to 10, and

[0233] $*$ and $*$ in Formula 402 each indicate a binding site to M in Formula 401.

[0234] In an exemplary embodiment of the present invention, A_{401} and A_{402} in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothiophene group, an benzisothiophene group, a benzoxazole group, an benzisoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

[0235] In an exemplary embodiment of the present invention, in Formula 402, X_{401} is nitrogen and X_{402} is carbon, or each of X_{401} and X_{402} may be nitrogen.

[0236] In an exemplary embodiment of the present invention, R_{401} and R_{402} in Formula 402 may each independently be selected from:

[0237] hydrogen, deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group;

[0238] a C_1 - C_{20} alkyl group, and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

[0239] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

[0240] a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, $-F$, $-Cl$, $-Br$, $-I$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and


[0241] $-Si(Q_{401})(Q_{402})(Q_{403})$, $-N(Q_{401})(Q_{402})$, $-B(Q_{401})(Q_{402})$, $-C(=O)(Q_{401})$, $-S(=O)_2(Q_{401})$, and $-P(=O)(Q_{401})(Q_{402})$, and

[0242] Q_{401} to Q_{403} may each independently be selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, and a naphthyl group, but exemplary embodiments of the present invention are not limited thereto.


[0243] In an exemplary embodiment of the present invention, when $xc1$ in Formula 401 is two or more, two $A_{401}(s)$ in two or more $L_{401}(s)$ may be chemically bonded to each other via X_{407} , which may be a linking group, or two $A_{402}(s)$ in two or more $L_{401}(s)$ may be chemically bonded to each other via X_{408} , which may be a linking group (see, e.g., Compounds PD1 to PD4 and PD7). X_{407} and X_{408} may each independently be a single bond, $*-O-*$, $*-S-*$, $*-C(=O)-*$, $*-N(Q_{413})-*$, $*-C(Q_{413})(Q_{414})-*$, or $*-C(Q_{413})=C(Q_{414})-*$ (e.g., Q_{413} and Q_{414} may each independently be hydrogen, deuterium, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), but exemplary embodiments of the present invention are not limited thereto.

[0244] L_{402} in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example, L_{402} may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), $-C(=O)$, isonitrile, $-CN$, and phosphorus (for example, phosphine, or phosphite), but exemplary embodiments of the present invention are not limited thereto.

[0245] In an exemplary embodiment of the present invention, the phosphorescent dopant may be selected from, for example, Compounds PD1 to PD25, but exemplary embodiments of the present invention are not limited thereto:




PD1




PD5

PD6




PD2

PD7




PD3

PD8





PD4


PD9








PD10



-continued




-continued




-continued

PD11




PD16

PD12




PD17

PD13




PD18

PD14




PD19

PD15




PD20

-continued



PD21

-continued





PD25

[0246] The fluorescent dopant may include an arylamine compound or a styrylamine compound.

[0247] The fluorescent dopant may include a compound represented by Formula 501 below.

PD22

<Formula 501>



[0248] In Formula 501,

[0249] Ar_{501} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0250] L_{501} to L_{503} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0251] x_{d1} to x_{d3} may each independently be an integer selected from 0 to 3;


[0252] R_{501} and R_{502} may each independently be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

[0253] x_{d4} may be an integer selected from 1 to 6.

[0254] In an exemplary embodiment of the present invention, Ar_{501} in Formula 501 may be selected from:

[0255] a heptalene group, a fluorene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthalene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group; and

PD24



[0256] a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indenophenanthrene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0257] In an exemplary embodiment of the present invention, L₅₀₁ to L₅₀₃ in Formula 501 may each independently be selected from:

[0258] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group; and

[0259] a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a benzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

[0260] In an exemplary embodiment of the present invention, R₅₀₁ and R₅₀₂ in Formula 501 may each independently be selected from:

[0261] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl

group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and


[0262] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, and —Si(Q₃₁)(Q₃₂)(Q₃₃); and

[0263] Q₃₁ to Q₃₃ may each be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.




[0264] In an exemplary embodiment of the present invention, xd4 in Formula 501 may be 2, but exemplary embodiments of the present invention are not limited thereto.

[0265] For example, the fluorescent dopant may be selected from Compounds FD1 to FD22:

FD1




-continued



-continued


FD2

FD5




FD3

FD6




FD4

FD7




-continued

FD8




-continued


FD11




FD12




FD9




FD13




FD14







FD10



FD15




-continued



-continued


FD16



FD20


FD17

FD18




FD21

FD19




FD22


[0266] In an exemplary embodiment of the present invention, the fluorescent dopant may be selected from the following compounds, but exemplary embodiments of the present invention are not limited thereto.




DPVBi




DPAVBi




TBPe




DCM



DCJTb



Coumarin 6



[0267] The electron transport region may have a single-layered structure including a single layer including a single material, a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a plurality of layers in which each of the plurality of layers include different materials from each other.

[0268] The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, but exemplary embodiments of the present invention are not limited thereto.

[0269] For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure. In each of these structures the layers may be sequentially stacked on an emission layer. However, exemplary embodiments of the present invention are not limited thereto.

[0270] The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound including at least one π electron-depleted nitrogen-containing ring.

[0271] The “ π electron-depleted nitrogen-containing ring” indicates a C_1 - C_{60} heterocyclic group having at least one $*-N=*$ moiety as a ring-forming moiety.

[0272] For example, the “ π electron-depleted nitrogen-containing ring” may be a 5-membered to 7-membered heteromonocyclic group having at least one $*-N=*$ moiety, a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one $*-N=*$ moiety are condensed with each other, or a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one $*-N=*$ moiety, is condensed with at least one C_5 - C_{60} carbocyclic group.

[0273] Examples of the π electron-depleted nitrogen-containing ring include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an benzoisothiazole, a benzoxazole, an benzoisoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, thiadiazol, an imidazopyridine, an imidazopyrimi-

dine, and an azacarbazole, but exemplary embodiments of the present invention are not limited thereto.

[0274] For example, the electron transport region may include a compound represented by Formula 601:



[0275] In Formula 601,

[0276] Ar_{601} may be a substituted or unsubstituted C_5 - C_{60} carbocyclic group or a substituted or unsubstituted C_1 - C_{60} heterocyclic group,

[0277] $xe11$ may be an integer selected from 1, 2, or 3,

[0278] L_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkylene group, a substituted or unsubstituted C_3 - C_{10} cycloalkenylene group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenylene group, a substituted or unsubstituted C_6 - C_{60} arylene group, a substituted or unsubstituted C_1 - C_{60} heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

[0279] $xe1$ may be an integer selected from 0 to 5,

[0280] R_{601} may be selected from a substituted or unsubstituted C_3 - C_{10} cycloalkyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkyl group, a substituted or unsubstituted C_3 - C_{10} cycloalkenyl group, a substituted or unsubstituted C_1 - C_{10} heterocycloalkenyl group, a substituted or unsubstituted C_6 - C_{60} aryl group, a substituted or unsubstituted C_6 - C_{60} aryloxy group, a substituted or unsubstituted C_6 - C_{60} arylthio group, a substituted or unsubstituted C_1 - C_{60} heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, $-Si(Q_{601})(Q_{602})(Q_{603})$, $-C(=O)(Q_{601})$, $-S(=O)_2(Q_{601})$, and $-P(=O)(Q_{601})(Q_{602})$,

[0281] Q_{601} to Q_{603} may each independently be a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and

[0282] $xe21$ may be an integer selected from 1 to 5.

[0283] In an exemplary embodiment of the present invention, at least one of $Ar_{601}(s)$ in the number of $xe11$ and $R_{601}(s)$ in the number of $xe21$ may include the π electron-depleted nitrogen-containing ring.

[0284] In an exemplary embodiment of the present invention, ring Ar_{601} in Formula 601 may be selected from:

[0285] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naph-

thacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

[0286] a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an benzoisoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q₃₁)(Q₃₂)(Q₃₃), —S(=O)₂(Q₃₁), and —P(=O)(Q₃₁)(Q₃₂); and


[0287] Q₃₁ to Q₃₃ may each independently be selected from a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0288] When xe11 in Formula 601 is two or more, two or more Ar601(s) may be chemically bonded to each other via a single bond.

[0289] In an exemplary embodiment of the present invention, Ar601 in Formula 601 may be an anthracene group.

[0290] In an exemplary embodiment of the present invention, a compound represented by Formula 601 may be represented by Formula 601-1:

<Formula 601-1>



[0291] In Formula 601-1,

[0292] X₆₁₄ may be N or C(R₆₁₄), X₆₁₅ may be N or C(R₆₁₅), X₆₁₆ may be N or C(R₆₁₆), at least one selected from X₆₁₄ to X₆₁₆ may be N,

[0293] L₆₁₁ to L₆₁₃ are the same as L₆₀₁,

[0294] xe611 to xe613 are the same as xe1,

[0295] R₆₁₁ to R₆₁₃ are the same as R₆₀₁,

[0296] R₆₁₄ to R₆₁₆ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

[0297] In an exemplary embodiment of the present invention, L₆₀₁ and L₆₁₁ to L₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0298] a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylene group, a phenazinylylene group, a benzimidazolylylene group, a benzoisothiazolylylene group, a benzoxazolylylene group, an benzoisoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group; and

[0299] a phenylene group, a naphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a diben-

zofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an benzoisothiazolylene group, a benzoxazolylene group, an benzoisoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylnyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; but exemplary embodiments of the present invention are not limited thereto.

[0300] In an exemplary embodiment of the present invention, x_{e1} and x_{e611} to x_{e613} in Formulae 601 and 601-1 may each independently be an integer selected from 0, 1, or 2.

[0301] In an exemplary embodiment of the present invention, R₆₀₁ and R₆₁₁ to R₆₁₃ in Formulae 601 and 601-1 may each independently be selected from:

[0302] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl

group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylnyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;






[0303] a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylnyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinylnyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl

group, a phenanthrolyl group, a phenaziny group, a benzimidazolyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

[0304] $-S(=O)_2(Q_{601})$ and $-P(=O)(Q_{601})(Q_{602})$, and


[0305] Q_{601} and Q_{602} are the same as described above.

[0306] The electron transport region may include at least one compound selected from Compounds ET1 to ET36, but exemplary embodiments of the present invention are not limited thereto:




-continued

ET6




-continued


ET9




ET7



ET10




ET8




-continued

ET11





-continued

ET14





ET15

ET12




ET13

ET16




-continued




ET17

-continued




ET20


ET18




ET21



ET19




ET22




-continued

ET23




-continued


ET26




ET24




ET27



ET25




ET28




-continued

ET29





-continued

ET32




ET33


ET30





ET31




ET34



-continued





[0307] In an exemplary embodiment of the present invention, the electron transport region may include at least one compound selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (bphen), Alq₃, BALq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ.

Alq₃


-continued

ET35



TAZ

ET36



NTAZ

[0308] A thickness of the buffer layer, the hole blocking layer, or the electron control layer may be in a range of from about 20 Å to about 1,000 Å, for example, from about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.


[0309] A thickness of the electron transport layer may be in a range of from about 100 Å to about 1,000 Å, for example, from about 150 Å to about 500 Å. When the thickness of the electron transport layer is within the ranges described (TAZ), above, the electron transport layer may have desired electron transport characteristics without a substantial increase in driving voltage.

[0310] The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a material including metal.


[0311] The material including metal may include at least one selected from alkali metal complex or alkaline earth-metal complex. The alkali metal complex may include a metal ion selected from a Li ion, a Na ion, a K ion, a Rb ion, and a Cs ion, and the alkaline earth-metal complex may include a metal ion selected from a Be ion, a Mg ion, a Ca ion, a Sr ion, and a Ba ion. A ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy

phenyloxazole, a hydroxy phenylthiazole, a hydroxy diphenyloxadiazole, a hydroxy diphenylthiadiazol, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but exemplary embodiments of the present invention are not limited thereto.

[0312] For example, the material including metal may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2.



ET-D1



ET-D2

[0313] The electron transport region may include an electron injection layer that facilitates the injection of electrons from the second electrode **190**. The electron injection layer may be in direct contact with the second electrode **190**.

[0314] The electron injection layer may have a single-layered structure including a single layer including a single material, a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a plurality of layers in which each of the plurality of layers include different materials from each other.

[0315] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof.

[0316] The alkali metal may be selected from Li, Na, K, Rb, or Cs. In an exemplary embodiment of the present invention, the alkali metal may be Li, Na, or Cs. In an exemplary embodiment of the present invention, the alkali metal may be Li or Cs, but exemplary embodiments of the present invention are not limited thereto.

[0317] The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

[0318] The rare earth metal may be selected from Sc, Y, Ce, Tb, Yb, and Gd.

[0319] The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare earth metal.

[0320] The alkali metal compound may be selected from alkali metal oxides, such as Li_2O , Cs_2O , or K_2O , and alkali metal halides, such as LiF, NaF, CsF, KF, LiI, NaI, CsI, or KI. In an exemplary embodiment of the present invention, the alkali metal compound may be selected from LiF, Li_2O , NaF, LiI, NaI, CsI, and KI, but exemplary embodiments of the present invention are not limited thereto.

[0321] The alkaline earth-metal compound may be selected from alkaline earth-metal oxides, such as BaO, SrO, CaO, $\text{Ba}_x\text{Sr}_{1-x}\text{O}$ ($0 < x < 1$), or $\text{Ba}_x\text{Ca}_{1-x}\text{O}$ ($0 < x < 1$). In an exemplary embodiment of the present invention, the alkaline earth-metal compound may be selected from BaO, SrO, and CaO, but exemplary embodiments of the present invention are not limited thereto.

[0322] The rare earth metal compound may be selected from YbF_3 , ScF_3 , ScO_3 , Y_2O_3 , Ce_2O_3 , GdF_3 , and TbF_3 . In an exemplary embodiment of the present invention, the rare earth metal compound may be selected from YbF_3 , ScF_3 , TbF_3 , YbI_3 , ScI_3 , and TbI_3 , but exemplary embodiments of the present invention are not limited thereto.

[0323] The alkali metal complex, the alkaline earth-metal complex, and the rare earth metal complex may include an ion of alkali metal, alkaline earth-metal, and rare earth metal as described above. A ligand coordinated with a metal ion of the alkali metal complex, the alkaline earth-metal complex, or the rare earth metal complex may be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenyloxazole, hydroxy phenylthiazole, hydroxy diphenyloxadiazole, hydroxy diphenylthiadiazol, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, but exemplary embodiments of the present invention are not limited thereto.

[0324] The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof, as described above. In an exemplary embodiment of the present invention, the electron injection layer may further include an organic material. An organic material, an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combinations thereof included in the electron injection layer may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

[0325] A thickness of the electron injection layer may be in a range of from about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have desired electron injection characteristics without a substantial increase in driving voltage.

[0326] The second electrode **190** may be located on the organic layer **150**. The second electrode **190** may be a cathode which is an electron injection electrode, and in this regard, a material included in the second electrode **190** may be selected from a metal, an alloy, an electrically conductive compound, and a mixture thereof, which may have a relatively low work function.

[0327] The second electrode **190** may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, but exemplary embodiments of the present invention are not limited thereto. The second electrode **190** may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

[0328] The second electrode **190** may have a single-layered structure, or a multi-layered structure including two or more layers.

[0329] FIG. 2 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention. FIG. 3 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention. FIG. 4 is a schematic view of an organic light-emitting device according to an exemplary embodiment of the present invention.

[0330] Referring to FIGS. 2 to 4, an organic light-emitting device **20** may include a first capping layer **210**, a first electrode **110**, an organic layer **150**, and a second electrode **190**, sequentially stacked in this stated order. An organic light-emitting device **30** may include a first electrode **110**, an organic layer **150**, a second electrode **190**, and a second capping layer **220**, sequentially stacked in this stated order. An organic light-emitting device **40** may include a first capping layer **210**, a first electrode **110**, an organic layer **150**, a second electrode **190**, and a second capping layer **220**, sequentially stacked in this stated order.

[0331] The first electrode **110**, the organic layer **150**, and the second electrode **190** described below with reference to FIGS. 2 to 4 may be substantially the same as the first electrode **110**, the organic layer **150**, and the second electrode **190** described above with reference to FIG. 1, and thus duplicative descriptions may be omitted below.

[0332] In the organic layer **150** of each of the organic light-emitting devices **20** and **40**, light generated in an emission layer may pass through the first electrode **110** and the first capping layer **210** toward the outside. As an example, the first electrode **110** may be a semi-transmissive electrode or a transmissive electrode. In the organic layer **150** of each of the organic light-emitting devices **30** and **40**, light generated in an emission layer may pass through the second electrode **190** and the second capping layer **220** toward the outside. As an example, the second electrode **190** may be a semi-transmissive electrode or a transmissive electrode.

[0333] The first capping layer **210** and the second capping layer **220** may increase external luminescent efficiency according to the principle of constructive interference.


[0334] The first capping layer **210** and the second capping layer **220** may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

[0335] At least one selected from the first capping layer **210** and the second capping layer **220** may each independently include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrine derivatives, phthalocyanine derivatives, a naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based


compound may be substituted with a substituent including at least one element selected from O, N, S, Se, Si, F, Cl, Br, and I. In an exemplary embodiment of the present invention, at least one selected from the first capping layer **210** and the second capping layer **220** may each independently include an amine-based compound.

[0336] In an exemplary embodiment of the present invention, at least one selected from the first capping layer **210** and the second capping layer **220** may each independently include the compound represented by Formula 201 or the compound represented by Formula 202.

[0337] In an exemplary embodiment of the present invention, at least one selected from the first capping layer **210** and the second capping layer **220** may each independently include a compound selected from Compounds HT28 to HT33 and Compounds CP1 to CP5, but exemplary embodiments of the present invention are not limited thereto.



-continued



[0338] In an exemplary embodiment of the present invention, at least one selected from the first capping layer 210 and the second capping layer 220 (see, e.g., FIG. 4) may include the condensed cyclic compound represented by Formula 1.

[0339] An organic light-emitting device according to an exemplary embodiment of the present invention is described above with reference to FIGS. 1-4. However, exemplary embodiments of the present invention are not limited thereto.

[0340] The hole transport region, the emission layer, or the electron transport region may be formed in a predetermined region by, for example, vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

[0341] When the hole transport region, the emission layer, or the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of from about 100° C. to about 500° C., at a vacuum degree of from about 10–8 torr to about 10–3 torr, and at a deposition rate of from about 0.01 Å/sec to about 100 Å/sec by taking into account a compound to be included in a layer to be formed, and the structure of a layer to be formed.

[0342] When the hole transport region, the emission layer, and the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of from about 2000 rpm to about 5000 rpm and at a heat treatment temperature of from about 80° C. to about 200° C. by taking into account a compound to be included in a layer to be formed, and the structure of a layer to be formed.

[0343] The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and

examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. A C₁-C₆₀ alkylene group used herein refers to a divalent group having the same structure as that of the C₁-C₆₀ alkyl group.

[0344] The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group having the same structure as that of the C₂-C₆₀ alkyl group, except for at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. A C₂-C₆₀ alkenylene group used herein refers to a divalent group having the same structure as that of the C₂-C₆₀ alkenyl group.

[0345] The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group having at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group, and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as that of the C₂-C₆₀ alkynyl group.

[0346] The term “C₁-C₆₀ alkoxy group” as used herein refers to a monovalent group represented by —OA₁₀₁ (e.g., A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

[0347] The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. A C₃-C₁₀ cycloalkylene group used herein refers to a divalent group having the same structure as that of the C₃-C₁₀ cycloalkyl group.

[0348] The term “C₁-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuranlyl group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

[0349] The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

[0350] The term “C₁-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Examples of the C₁-C₁₀ heterocycloalkenyl group are a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuranlyl group, and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkenyl group.

[0351] The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and a C₆-C₆₀ arylene group used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C₆-C₆₀ aryl group are a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be chemically bonded to each other.

[0352] The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C₁-C₆₀ heteroarylene group” as used herein refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Examples of the C₆-C₆₀ heteroaryl group are a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinoxalinyl group, and an isoquinolinyl group. When the C₆-C₆₀ heteroaryl group and the C₆-C₆₀ heteroarylene group each include two or more rings, the rings may be chemically bonded to each other.

[0353] The term “C₆-C₆₀ aryloxy group” as used herein indicates —OA₁₀₂ (e.g., A₁₀₂ is the C₆-C₆₀ aryl group), and the term “C₆-C₆₀ arylthio group” as used herein indicates —SA₁₀₃ (e.g., A₁₀₃ is the C₆-C₆₀ aryl group).

[0354] The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) having two or more rings condensed with each other, only carbon atoms as ring-forming atoms, and no aromaticity in its entire molecular structure. An example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

[0355] The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) having two or more rings chemically bonded to each other, at least one heteroatom selected from N, O, Si, P, and S, other than carbon atoms, as a ring-forming atom, and no aromaticity in its entire molecular structure. An example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

[0356] The term “C₅-C₆₀ carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms in which a ring-forming atom is a carbon atom only. The C₅-C₆₀ carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C₅-C₆₀ carbocyclic group may be a ring, such as benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In an exemplary embodiment of the present invention, depending on the number of substituents connected to the C₅-C₆₀ carbocyclic

group, the C₅-C₆₀ carbocyclic group may be a trivalent group or a quadrivalent group.

[0357] The term “C₁-C₆₀ heterocyclic group” as used herein refers to a group having the same structure as the C₁-C₆₀ carbocyclic group, except that as a ring-forming atom, at least one heteroatom selected from N, O, Si, P, and S is used in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

[0358] At least one substituent of the substituted C₅-C₆₀ carbocyclic group, substituted C₁-C₆₀ heterocyclic group, substituted C₃-C₁₀ cycloalkylene group, substituted C₁-C₁₀ heterocycloalkylene group, substituted C₃-C₁₀ cycloalkenylene group, substituted C₁-C₁₀ heterocycloalkenylene group, substituted C₆-C₆₀ arylene group, substituted C₁-C₆₀ heteroarylene group, substituted divalent non-aromatic condensed polycyclic group, substituted divalent non-aromatic condensed heteropolycyclic group, substituted C₁-C₆₀ alkyl group, substituted C₂-C₆₀ alkenyl group, substituted C₂-C₆₀ alkynyl group, substituted C₁-C₆₀ alkoxy group, substituted C₃-C₁₀ cycloalkyl group, substituted C₁-C₁₀ heterocycloalkyl group, substituted C₃-C₁₀ cycloalkenyl group, substituted C₁-C₁₀ heterocycloalkenyl group, substituted C₆-C₆₀ aryl group, substituted C₆-C₆₀ aryloxy group, substituted C₆-C₆₀ arylthio group, substituted C₁-C₆₀ heteroaryl group, substituted monovalent non-aromatic condensed polycyclic group and substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

[0359] deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group;

[0360] a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), and —P(=O)(Q₁₁)(Q₁₂);

[0361] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

[0362] a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy

group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), S(=O)₂(Q₂₁), and —P(=O)(Q₂₁)(Q₂₂);

[0363] —Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), or —P(=O)(Q₃₁)(Q₃₂); and

[0364] Q₁₁ to Q₁₃, Q₂₁ to Q₂₃ and Q₃₁ to Q₃₃ may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

[0365] The term “Ph” as used herein represents a phenyl group, the term “Me” as used herein represents a methyl group, the term “Et” as used herein represents an ethyl group, the term “ter-Bu” or “But,” as used herein, represents a tert-butyl group, and the term “OMe” as used herein represents a methoxy group.

[0366] The term “biphenyl group” used herein refers to a “phenyl group substituted with a phenyl group. The term “biphenyl” is a “substituted phenyl group” having a “C₆-C₆₀ aryl group” as a substituent.

[0367] A “terphenyl group” used herein refers to a “phenyl group substituted with a biphenyl group. The “terphenyl group” is a “phenyl group” having, as a substituent, a “C₆-C₆₀ aryl group substituted with a C₆-C₆₀ aryl group.”



[0368] * and * used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.

[0369] A compound according to an exemplary embodiment of the present invention and an organic light-emitting device according to an exemplary embodiment of the present invention will be described in more detail below with reference to Synthesis Examples and Examples. The wording “B was used instead of A” used in describing Synthesis Examples refers to that an identical molar equivalent of B was used in place of A.

EXAMPLES

Synthesis Example 1: Synthesis of Intermediate A

[0370]



Exemplary Synthesis of Intermediate A-1


[0371] 2.281 g (20 mmol) of 1,2-difluorobenzene, 3.211 g (20 mmol) of 2,3-dihydroxynaphthalene, 0.190 g (1 mmol) of CuI, 0.264 g (1.0 mmol) of 18-Crown-6, and 4.124 g (30 mmol) of potassium carbonate may be dissolved in 100 mL of DMF, and then, the mixture may be stirred at a temperature of 140° C. for 24 hours. The reaction solution may be cooled to room temperature, and then, the result may be subjected to an extraction process three times by using 60 mL of water and 60 mL of diethylether, and an organic layer separated therefrom may be dried by using magnesium sulfate and the residual obtained by evaporating a solvent therefrom was separation-purified by silica gel column chromatography to obtain 3.512 g (e.g., yield of 70%) of Intermediate A-1.

Exemplary Synthesis of Intermediate A


[0372] 3.512 g (15 mmol) of Intermediate A-1 may be dissolved in 50 mL of dichloromethane, and then, the mixture may be stirred at a temperature of 0° C. for 30 minutes. Then 2.385 g (15 mmol) of Br₂ may be slowly added dropwise thereto, and the reaction solution may be stirred at room temperature for 6 hours. 60 mL of water may be added thereto and the mixture may be extracted three times with 50 mL of dichloromethane. The obtained organic layer may be dried by using magnesium sulfate and the solvent may be evaporated. The obtained residue may be separation-purified by silica gel column chromatography to obtain 5.881 g (e.g., yield 100%) of Intermediate A.

Synthesis Example 2: Synthesis of Compound 1


[0373]



-continued




-continued



[0374] 3.921 g (10 mmol) of Intermediate A, 13.348 g (20 mmol) of amine 1, 0.452 g (0.5 mmol) of tris(dibenzylideneacetone)dipalladium ($\text{Pd}_2(\text{dba})_3$), 0.101 g (0.5 mmol) of PtBu_3 , and 0.961 g (10 mmol) of NaO^tBu may be dissolved in 150 ml of toluene, and then, the mixture may be stirred at a temperature of 90° C. for 4 hours. The reaction solution may be cooled to room temperature, and then, subjected to an extraction process three times by using 10 ml of water and 50 ml of diethyl ether. The obtained organic layer may be dried by using magnesium sulfate and the residual obtained by evaporating a solvent therefrom may be separation-purified by silica gel column chromatography to obtain 3.976 g (7 mmol, the yield of, e.g., 70%) of Compound 1.

Synthesis Example 3: Synthesis of Compound 12


[0375]



[0376] 6.307 g (7 mmol, the yield of 70%) of Compound 12 may be obtained in the same manner as used to prepare Compound 1 except that amine 2 may be used instead of amine 1.

Synthesis Example 4: Synthesis of Intermediate B

[0377]






[0378] In a nitrogen atmosphere, 3.921 g (10 mmol) of Intermediate A, 1.563 g (10 mmol) of 4-chlorophenyl boronic acid, 0.578 g (0.5 mmol) of $\text{Pd}(\text{PPh}_3)_4$, and 2.762 g (20 mmol) of K_2CO_3 may be dissolved in 100 mL of a mixed solution containing THF and H_2O (at a volumetric ratio of 2:1), and then, the mixture may be stirred at a temperature of 80° C. for 12 hours. The reaction solution may be cooled to room temperature, and then, 50 ml of water may be added thereto, and the resultant solution may be extracted three times by using 150 ml of ethylether. The obtained organic

solvent layer may be dried by using magnesium sulfate and the residual obtained by evaporating a solvent therefrom may be separation-purified by silica gel column chromatography to obtain 2.965 g (7 mmol, the yield of, e.g., 70%) of Intermediate B.

Synthesis Example 5: Synthesis of Compound 18


[0379]



[0380] 5.572 g (7 mmol, the yield of 70%) of Compound 18 may be obtained in the same manner as used to synthesize Compound 1 except that Intermediate B may be used instead of Intermediate A and amine 3 may be used instead of amine 1.

Synthesis Example 6: Synthesis of Compound 29


[0381]




[0382] 5.465 g (7 mmol, the yield of 70%) of Compound 29 may be obtained in the same manner as used to synthesize Compound 1 except that Intermediate B may be used instead of Intermediate A and amine 4 may be used instead of amine 1.

Synthesis Example 7: Synthesis of Compound 40

[0383]







[0384] 6.013 g (7 mmol, the yield of, e.g., 70%) of Compound 40 may be obtained in the same manner as used to synthesize Compound 1 except that Intermediate B may be used instead of Intermediate A and amine 5 may be used instead of amine 1.

Synthesis Example 8: Synthesis of Intermediate C


[0385]




[0386] 3.376 g (7 mmol, the yield of, e.g., 70%) of Intermediate C may be obtained in the same method as used to prepare Intermediate B except that Intermediate B may be used instead of Intermediate A and 4-chloro-2,5-dimethylphenyl boronic acid may be used instead of 4-chlorophenyl boronic acid.

Synthesis Example 9: Synthesis of Compound 48

[0387]





[0388] 6.728 g (7 mmol, the yield of, e.g., 70%) of Compound 48 may be obtained in the same manner as used to synthesize Compound 1 except that Intermediate C may be used instead of Intermediate A and amine 6 may be used instead of amine 1.

TABLE 1

Com- pound	¹ H NMR (CDCl ₃ , 300 MHz)	LC/MS	
		found	calc.
1	8.2(d, 2H), 7.8-7.5(m, 10H), 7.4-7.0 (m, 14H), 6.9 (d, 2H)	569.20	568.22
12	8.2(d, 2H), 8.0-7.6(m, 18H), 7.5-7.2 (m, 14H), 7.0 (m, 4H), 6.9 (d, 2H)	901.05	900.30
18	8.3(d, 1H), 8.2(d, 1H), 8.0-7.6(m, 14H), 7.5-7.2 (m, 14H), 7.0 (m, 8H), 6.9 (d, 2H)	796.98	796.31
29	8.3(d, 1H), 8.2-8.0(m, 5H), 7.8-7.4 (m, 14H), 7.3-7.0 (m, 10H), 6.9 (d, 2H)	781.05	780.26
40	8.3(d, 1H), 8.2-7.8(m, 7H), 7.7-7.2 (m, 18H), 7.1-6.9 (m, 6H), 6.8 (d, 2H)	859.05	858.21
48	8.3(d, 1H), 8.2(d, 1H), 8.0-7.6(m, 12H), 7.5-7.2 (m, 14H), 7.0 (m, 8H), 6.9 (d, 2H), 2.4(s, 3H), 2, 2(s, 3H)	961.21	960.28




Example 1

[0389] As a substrate and an anode, a glass substrate with Corning 15 Ωcm² (e.g., about 1,200 Å) ITO thereon may be cut to a size of 50 mm×50 mm×0.7 mm, sonicating the glass substrate by using isopropyl alcohol and pure water for 5 minutes each, and then irradiating UV light for 30 minutes thereto and exposing to ozone to clean. Then, the resultant glass substrate may be loaded onto a vacuum deposition apparatus.


[0390] 2-TNATA may be vacuum-deposited on the ITO anode to form a hole injection layer having a thickness of about 600 Å, and then, Compound 1 may be vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of about 300 Å.

[0391] 9,10-di-naphthalene-2-yl-anthracene (ADN) acting as a host and N,N,N',N'-tetraphenyl-pyrene-1,6-diamine (TPD) acting as a dopant may be co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of about 300 Å.

[0392] Thereafter, Alq₃ may be deposited on the emission layer to form an electron transport layer having a thickness of about 300 Å, and LiF may be deposited on the electron transport layer to form an electron injection layer having a thickness of about 10 Å, and Al may be vacuum-deposited thereon to form a cathode having a thickness of about 3,000 Å, thus forming an organic light-emitting device.



-continued



Examples 2 to 6 and Comparative Examples 1 and 2

[0393] Organic light-emitting devices may be manufactured in the same manner as in Example 1, except that in forming a hole transport layer, the compounds shown in Table 2 may be used instead of Compound 1.

Evaluation Example 1

[0394] The driving voltage, current density, efficiency, emission color, and half-lifespan of the organic light-emitting devices manufactured according to Examples 1 to 6, and Comparative Examples 1 and 2 may be measured by using Kethley MU 236 and a brightness photometer PR650, and exemplary results thereof are shown in Table 2.

TABLE 2

	Hole transport layer	Driving voltage (V)	Current density (mA/cm ²)	Brightness (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
Example 1	Compound 1	5.66	50	3356	6.53	Blue	320
Example 2	Compound 12	5.49	50	3591	6.45	Blue	340
Example 3	Compound 18	5.90	50	3493	6.27	Blue	317
Example 4	Compound 29	5.60	50	3075	6.15	Blue	300
Example 5	Compound 40	5.52	50	3175	6.35	Blue	325
Example 6	Compound 48	5.84	50	3020	6.04	Blue	315
Comparative Example 1	NPB	6.99	50	2745	5.27	Blue	266
Comparative Example 2	Compound A	9.48	50	1305	2.61	Blue	163





TABLE 2-continued





Hole transport layer	Driving voltage (V)	Current density (mA/cm ²)	Brightness (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
						
18						
						
29						
						
40						
						
48						

TABLE 2-continued

Hole transport layer	Driving voltage (V)	Current density (mA/cm ²)	Brightness (cd/m ²)	Efficiency (cd/A)	Emission color	Half lifespan (hr @ 100 mA/cm ²)
<p style="text-align: center;">NPB</p>						
<p style="text-align: center;">A</p>						


[0395] Referring to Table 2, the organic light-emitting devices prepared according to Examples 1 to 6 may have lower driving voltage, higher intensity of brightness, higher efficiency, and longer lifespan than the organic light-emitting devices prepared according to Comparative Examples 1 and 2.

[0396] An organic light-emitting device including the condensed cyclic compound according to an exemplary embodiment of the present invention may have a relatively low driving voltage, relatively high brightness, relatively high efficiency, and a relatively long lifespan.


[0397] While the present invention has been shown and described with reference to the exemplary embodiments thereof, it will be apparent to those of ordinary skill in the art that various changes in form and detail may be made thereto without departing from the spirit and scope of the present invention.

What is claimed is:

1. A condensed cyclic compound represented by Formula 1:



-continued



wherein, in Formulae 1 and 2,

X₁ and X₂ are each independently O or S,

L₁, L₂, and L₁₁ are each independently selected from a substituted or unsubstituted C₃-C₁₀ cycloalkylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkylene group, a substituted or unsubstituted C₃-C₁₀ cycloalkenylene group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenylene group, a substituted or unsubstituted C₆-C₆₀ arylene group, a substituted or unsubstituted C₁-C₆₀ heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

a₁, a₂, and a₁₁ are each independently an integer selected from 0, 1, 2, 3, 4, or 5,

Ar₁ and Ar₂ are each independently selected from a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl

group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, or a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

R₁ to R₁₀ are each independently selected from a group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or each independently unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁)(Q₂)(Q₃), —B(Q₁)(Q₂), —C(=O)(Q₁), —S(=O)₂(Q₁), or —P(=O)(Q₁)(Q₂), and at least two selected from R₁ to R₁₀ are each the group represented by Formula 2, and

at least one selected from the substituted C₃-C₁₀ cycloalkylene group, the substituted C₁-C₁₀ heterocycloalkylene group, the substituted C₃-C₁₀ cycloalkenylene group, the substituted C₁-C₁₀ heterocycloalkenylene group, the substituted C₆-C₆₀ arylene group, the substituted C₁-C₆₀ heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₁-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, or the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, and a C₁-C₆₀ alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀

heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₁₁)(Q₁₂)(Q₁₃), —N(Q₁₁)(Q₁₂), —B(Q₁₁)(Q₁₂), —C(=O)(Q₁₁), —S(=O)₂(Q₁₁), or —P(=O)(Q₁₁)(Q₁₂);

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q₂₁)(Q₂₂)(Q₂₃), —N(Q₂₁)(Q₂₂), —B(Q₂₁)(Q₂₂), —C(=O)(Q₂₁), —S(=O)₂(Q₂₁), or —P(=O)(Q₂₁)(Q₂₂); or

—Si(Q₃₁)(Q₃₂)(Q₃₃), —N(Q₃₁)(Q₃₂), —B(Q₃₁)(Q₃₂), —C(=O)(Q₃₁), —S(=O)₂(Q₃₁), or —P(=O)(Q₃₁)(Q₃₂),

wherein Q₁ to Q₃, Q₁₁ to Q₁₃, Q₂₁ to Q₂₃, and Q₃₁ to Q₃₃ are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, or a terphenyl group.

2. The condensed cyclic compound of claim 1, wherein X₁ and X₂ are each O.

3. The condensed cyclic compound of claim 1, wherein L₁, L₂, and L₁₁ are each independently selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a


pyrenylene group, a chrysenylene group, a naphthacenylylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a pyrrolylylene group, a thiophenylene group, a furanylylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylylene group, a benzothiofenylene group, an benzoisothiazolylene group, a benzoxazolylene group, an benzoisoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzosilolylene group, a dibenzocarbazolylene group, a thiadiazolylene group, an imidazopyridinylene group, or an imidazopyrimidinylene group; or

a phenylene group, a pentalenylylene group, an indenylene group, a naphthylene group, an azulenylylene group, a heptalenylene group, an indacenylylene group, an acenaphthylene group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylylene group, a picenylene group, a perylenylene group, a pentaphenylene group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenylylene group, an ovalenylylene group, a pyrrolylylene group, a thiophenylene group, a furanylylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a pyridinylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, an isoindolylene group, an indolylene group, an indazolylene group, a purinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a carbazolylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, a benzofuranylylene group, a benzothiofenylene group, an benzoisothiazolylene group, a benzoxazolylene group, an benzoisoxazolylene group, a triazolylene group, a tetrazolylene group, an oxadiazolylene group, a triazinylene group, a dibenzofuranylylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a thiadiazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, or $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

diazolylene group, an imidazopyridinylene group, and an imidazopyrimidinylene group, each substituted with at least one selected from deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenylylene group, a heptalenyl group, an indacenylylene group, an acenaphthyl group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenylylene group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylylene group, a pentacenylylene group, a rubicenylylene group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, or $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,












wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenylylene group, a spiro-bifluorenylylene group, a benzofluorenylylene group, a dibenzofluorenylylene group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, or a carbazolyl group.

4. The condensed cyclic compound of claim 1, wherein L_1 , L_2 , and L_{11} are each independently selected from groups represented by Formulae 3-1 to 3-28:




Formula 3-1

-continued




-continued


Formula 3-2




Formula 3-3




Formula 3-4



Formula 3-5




Formula 3-6




Formula 3-7

Formula 3-8




Formula 3-9


Formula 3-10



Formula 3-11



Formula 3-12



Formula 3-13

Formula 3-14

Formula 3-15

Formula 3-16

Formula 3-17


Formula 3-18

Formula 3-19


Formula 3-20

Formula 3-21


-continued




Formula 3-22




Formula 3-23




Formula 3-24




Formula 3-25



Formula 3-26



Formula 3-27



Formula 3-28


wherein, in Formulae 3-1 to 3-28,

Y_1 is O, S, C(Z_3)(Z_4), N(Z_5), or Si(Z_6)(Z_7),


Z_1 to Z_7 are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, or —Si(Q_{31})(Q_{32})(Q_{33}),

Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, d_2 is an integer selected from 1 or 2, d_3 is an integer selected from 1 to 3, d_4 is an integer selected from 1 to 4, d_5 is an integer selected from 1 to 5, d_6 is an integer selected from 1 to 6, d_8 is an integer selected from 1 to 8, and * and *' each indicate a binding site to a neighboring atom.


5. The condensed cyclic compound of claim 1, wherein L_1 , L_2 , and L_{11} are each independently selected from groups represented by Formulae 4-1 to 4-15:




Formular 4-1




Formular 4-2




Formular 4-3




Formular 4-4




Formular 4-5



Formular 4-6




Formular 4-7




Formular 4-8


-continued




Formular 4-9




Formular 4-10




Formular 4-11




Formular 4-12



Formular 4-13



Formular 4-14



Formular 4-15

wherein * and *' in Formulae 4-1 to 4-15 each indicate a binding site to a neighboring atom.

6. The condensed cyclic compound of claim 1, wherein a1, a2, and a11 are each independently an integer selected from 0, 1, or 2.

7. The condensed cyclic compound of claim 1, wherein Ar₁ and Ar₂ are each independently selected from:

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a


dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzoisothiazolyl group, a benzoxazolyl group, a benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group; or

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, a benzoisothiazolyl group, a benzoxazolyl group, a benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a thiadiazolyl group, an imidazopyridinyl group, and an imidazopyrimidinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group,


a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacacenyl group, a pentacacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an benzoisothiazolyl group, a benzoxazolyl group, an benzoisoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzosilolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a thiadiazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$, $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{C}(=\text{O})(\text{Q}_{31})$, $-\text{S}(=\text{O})_2(\text{Q}_{31})$, or $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$,

wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, or a carbazolyl group.


8. The condensed cyclic compound of claim 1, wherein Ar_1 and Ar_2 are each independently selected from groups represented by Formulae 5-1 to 5-24:




Formular 5-1




Formular 5-2




Formular 5-3




Formular 5-4




Formular 5-5




Formular 5-6




Formular 5-7




Formular 5-8




Formular 5-9




Formular 5-10



Formular 5-11




Formular 5-12




Formular 5-13

-continued


-continued




Fomular 5-14




Fomular 5-15




Fomular 5-16




Fomular 5-17




Fomular 5-18




Fomular 5-19




Fomular 5-20




Fomular 5-21



Fomular 5-22



Fomular 5-23



Fomular 5-24

wherein, in Formulae 5-1 to 5-24,

Y_{31} is O, S, $C(Z_{33})(Z_{34})$, $N(Z_{35})$, or $Si(Z_{36})(Z_{37})$,

Z_{31} to Z_{37} are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, or —Si(Q_{31})(Q_{32})(Q_{33}),

Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

e_2 is an integer selected from 1 or 2,

e_3 is an integer selected from 1 to 3,

e_4 is an integer selected from 1 to 4,


e_5 is an integer selected from 1 to 5,

e_6 is an integer selected from 1 to 6,


e_7 is an integer selected from 1 to 7, and

* indicates a binding site to a neighboring atom.


9. The condensed cyclic compound of claim 1, wherein Ar_1 and Ar_2 are each independently selected from groups represented by Formulae 6-1 to 6-25:




Fomular 6-1




Fomular 6-2




Fomular 6-3



Fomular 6-4












Fomular 6-5




Fomular 6-6

-continued




-continued


Formular 6-7




Formular 6-8




Formular 6-9




Formular 6-10




Formular 6-11



Formular 6-12




Formular 6-13



Formular 6-14

Formular 6-15



Formular 6-16

Formular 6-17

Formular 6-18

Formular 6-19

Formular 6-20


Formular 6-21

Formular 6-22


Formular 6-23

-continued

Formular 6-24



Formular 6-25



wherein * in Formulae 6-1 to 6-25 indicates a binding site to a neighboring atom.

10. The condensed cyclic compound of claim 1, wherein R_1 to R_{10} are each independently selected from:

the group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, or a C_1 - C_{20} alkoxy group;

a C_1 - C_{20} alkyl group and a C_1 - C_{20} alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, or —Si(Q_{31})(Q_{32})(Q_{33});

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, or a dibenzocarbazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, and a dibenzocarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a C_1 - C_{20} alkyl group, or a C_1 - C_{20} alkoxy group, and —Si(Q_{31})(Q_{32})(Q_{33}); or —Si(Q_1)(Q_2)(Q_3),

wherein Q_1 to Q_3 and Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group.

11. The condensed cyclic compound of claim 1, wherein R_1 to R_{10} are each independently selected from the group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl

group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q_1)(Q_2)(Q_3),

wherein Q_1 to Q_3 are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group.

12. The condensed cyclic compound of claim 1, wherein two selected from R_1 to R_{10} are each the group represented by Formula 2,

other substituents that are not selected from R_1 to R_{10} are selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, or —Si(Q_1)(Q_2)(Q_3),


wherein Q_1 to Q_3 are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group.

13. The condensed cyclic compound of claim 1, wherein two selected from R_1 to R_{10} are each the group represented by Formula 2, and

other substituents that are not selected from R_1 to R_{10} are each hydrogen.

14. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by Formula 1A:

<Formula 1A>




wherein, in Formula 1A,

X_1 , X_2 , L_1 , L_2 , L_{11} , a_1 , a_2 , a_{11} , Ar_1 , Ar_2 , and R_3 to R_{10} are the same as in claim 1, and


L_3 , L_4 , L_{12} , a_3 , a_4 , a_{12} , Ar_3 , and Ar_4 are the same as L_1 , L_2 , L_{11} , a_1 , a_2 , a_{11} , Ar_1 , and Ar_2 , respectively.

15. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is represented by Formula 1Aa or Formula 1Ab:

<Formula 1Aa>



<Formula 1Ab>



wherein, in Formulae 1Aa and 1Ab,

X_1 , X_2 , L_1 , L_2 , L_{11} , a_1 , a_2 , a_{11} , Ar_1 , Ar_2 , and R_3 to R_{10} are the same as in claim 1, and

L_3 , L_4 , L_{12} , a_3 , a_4 , a_{12} , Ar_3 , and Ar_4 are the same as L_1 , L_2 , L_{11} , a_1 , a_2 , a_{11} , Ar_1 , and Ar_2 , respectively, and


Z_1 is selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a carbazolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, or —Si(Q_{31})(Q_{32})(Q_{33}),

wherein Q_{31} to Q_{33} are each independently selected from a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and


d_4 is an integer selected from 1 to 4.

16. The condensed cyclic compound of claim 1, wherein the condensed cyclic compound is one of Compounds 1 to 48:


1



2




3




-continued

4




-continued


7




5




8




6



9




-continued





10

-continued




13

11





14

15




16

12




-continued


17




18



19




20




-continued


21




22



23




24




-continued

25




-continued


29




26




30




27




31




28



32




-continued




33


-continued




37




34




38



36




35



39


-continued

40




-continued


44




45




41




46




42



43




47



-continued

48



17. An organic light-emitting device, comprising:
 a first electrode;
 a second electrode facing the first electrode; and
 an organic layer between the first electrode and the second electrode
 wherein the organic layer comprises an emission layer and at least one of the condensed cyclic compounds of claim 1.

18. The organic light-emitting device of claim 17, wherein the first electrode is an anode, the second electrode is a cathode, and the organic layer further comprises a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode, wherein the hole transport region comprises a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof, and the electron transport region comprises a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, an electron injection layer, or any combination thereof.

19. The organic light-emitting device of claim 18, wherein the hole transport region comprises the at least one of the condensed cyclic compounds of claim 1.

20. The organic light-emitting device of claim 18, wherein the hole transport region comprises the hole transport layer, and the hole transport layer comprises the at least one of the condensed cyclic compounds of claim 1.

* * * * *

专利名称(译)	缩合环状化合物和包括其的有机发光器件		
公开(公告)号	US20190140184A1	公开(公告)日	2019-05-09
申请号	US16/182827	申请日	2018-11-07
[标]申请(专利权)人(译)	三星显示有限公司		
申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
当前申请(专利权)人(译)	三星DISPLAY CO. , LTD.		
[标]发明人	HAN SANGHYUN KIM SOOYON KIM YOUNGKOOK KIM JONGWOO JUNG HYEJIN HWANG SEOKHWAN		
发明人	HAN, SANGHYUN KIM, SOOYON KIM, YOUNGKOOK KIM, JONGWOO JUNG, HYEJIN HWANG, SEOKHWAN		
IPC分类号	H01L51/00 C07D319/14 C07D407/14 C07D409/14		
CPC分类号	H01L51/0061 C07D319/14 C07D407/14 C07D409/14 H01L51/5012 H01L51/5056 H01L51/5072 H01L51/5096 H01L51/5088 H01L51/5092 H01L51/0073 H01L51/0067 H01L51/0074		
优先权	1020170147603 2017-11-07 KR		
外部链接	Espacenet USPTO		

摘要(译)

缩合环状化合物和包含其的有机发光装置。由式1表示的耦合环状化合物：

10

190
150
110