(19)





(11) **EP 2 438 848 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent: 19.03.2014 Bulletin 2014/12 (51) Int Cl.: A61B 5/00 ^(2006.01) G06F 19/00 ^(2011.01)

A61B 5/0402^(2006.01)

- (21) Application number: 11184156.5
- (22) Date of filing: 06.10.2011

(54) Computer-implemented method for remote evaluation of ambulatory electrocardiographic monitoring

Computerumgesetztes Verfahren zur Telediagnostik der ambulanten Elektrokardiografie-Überwachung

Procédé informatique pour évaluer à distance la surveillance électrocardiographique en ambulatoire

- (84) Designated Contracting States: AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
- (30) Priority: 08.10.2010 US 901461
- (43) Date of publication of application: 11.04.2012 Bulletin 2012/15
- (73) Proprietor: Cardiac Science Corporation Bothell, WA 98021 (US)
- (72) Inventors:
 - Bardy, Gust H.
 - Carnation, Washington 98014 (US) • Bishay, Jon Mikalson
 - Seattle, Washington 98119 (US)

- (74) Representative: Bethenod, Marc et al Novagraaf Technologies SA
 122, Rue Edouard Vaillant
 92593 Levallois-Perret Cedex (FR)
- (56) References cited:
 - WO-A1-2010/066507US-A1- 2003 083 559US-A1- 2004 034 284US-A1- 2007 255 153US-A1- 2007 276 270
 - LIEBERMAN JONATHON: "How telemedicine is aiding prompt ECG diagnosis in primary care", BRITISH JOURNAL OF COMMUNITY NURSING, vol. 13, no. 3, 1 March 2008 (2008-03-01), pages 123-126, XP009155082, ISSN: 1462-4753

2 438 848 B1 Ч

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

Field

[0001] This application relates in general to ambulatory electrocardiographic monitoring and, in particular, to a computer-implemented method for evaluating ambulatory electrocardiographic monitoring of cardiac rhythm disorders.

Background

[0002] Cardiologists are experts in the diagnosis and treatment of heart rhythm disorders, such as tachycardia (excessively fast and abnormal heart rates) and bradycardia (excessively slow and abnormal heart rates). Disorders of the heart's electrical system are the primary cause of heart rhythm disorders, which, in turn, cause ineffective pumping of the blood, thereby placing a patient at risk of loss of consciousness and sudden death. Under managed care, access to medical specialists, such as cardiologists and cardiac electrophysiologists (cardiologists who focus purely on the heart's electrical system), requires a patient's primary care provider to first make a referral. Primary care physicians typically undertake their own diagnostic testing and screening before referring a patient. Evaluating the patient thoroughly often involves the ordering of tests. Each step preceding cardiac specialist referral, though, adds on an additional 30 percent of administrative overhead costs on average, while a failure by a primary care provider to act may force patients to seek assistance through the emergency room, thereby increasing emergency services' already overwhelming burden.

[0003] The diagnosis of cardiac rhythm disorders is challenging to address in the brief period of in-clinic exposure to the primary care provider due to the intermittent and variable nature of most heart rhythm disorders. Moreover, a patient's symptoms, such as syncope, presyncope, palpitations, angina, dyspnea, and fatigue, are consistent with many health problems, other than heart rhythm disorders, including benign causes. The ability of a primary care provider to make the appropriate diagnosis can be extremely challenging and the tools to aid in this diagnosis difficult to access. Moreover, an overall decline in the skills required to diagnose and treat heart rhythm disease in the general medical community has made primary care providers ill-equipped to make the kinds of classic arrhythmia diagnoses required to prevent or treat a serious illness. Furthermore, primary care providers are artificially limited in their ability to make patient referrals to cardiac specialists due to managed care restrictions, third-party provider rules, and practical obstacles in accessing needed diagnostic tools that are usually not in the same physical location as the primary care provider.

[0004] Current forms of diagnostic testing available inclinic to primary care providers, typically only the stand-

ard 12-lead electrocardiogram (ECG), are inadequate to identify cardiac rhythm disorders in the context of today's limitations in time, finances, and obstacles in the patient referral network. For instance, the conventional 12-lead ECG test is merely a "snapshot" of a patient's heart rhythm, lasting approximately 10 seconds. While 12-lead ECG testing can identify many forms of persistently-presented heart disease, 12-lead ECG testing is of nominal assistance in identifying episodic heart rhythm disorders 10 that rarely are present during clinic visits.

[0005] Other forms of ECG testing are more effective at identifying cardiac rhythm disorders but are seldom ordered by primary care providers because of the administrative and cost hurdles incurred in attempting the use

15 of such tools. For instance, U.S. Patent No. 3,215,136 issued November 2, 1965 to Holter et al. discloses an electrocardiographic recording and playback means, more commonly known as a "Holter monitor" named after its inventor, Norman Holter in 1949. Holter monitor is now

20 used synonymously with long-term heart rhythm recording, usually lasting at least 24 hours. With use of a Holter monitor, episodes of ventricular tachycardia, asystolic intervals, and ectopic heart beats are sensed and recorded by electrodes placed on the patient's skin. ECG rhythm

25 disturbances recorded by the Holter monitor can then be identified. In less constrained times, such abnormalities would be seen by cardiologists, who have been contracted to overread the Holter recordings and provide diagnostic information to the referring primary care provider.

30 These cardiologists would, in turn, care for the patient should a serious rhythm disorder be identified. Times have changed, however, and this older model of health care is no longer easily used or, if used, taken to the next step of referral and care by a cardiac specialist because 35 of logistical and cost constraints.

[0006] The value of ambulatory ECG or Holter monitoring and the appropriate use of this technology, especially by primary care providers, suffers from several drawbacks. First, under managed care, ambulatory ECG 40 monitoring is usually physically inaccessible to primary care providers in their offices. Such monitoring requires a referral to a testing laboratory, plus an additional referral to a cardiologist to interpret any findings, presuming the results of the Holter recording actually make their way

45 back to the referring primary care provider, which is not necessarily a guaranteed event. Further obstacles arise by virtue of the need for the patient to be compliant with accessing, properly using, and returning the monitor to the pickup location, as well as communication burdens 50 that arise between the referring physician, the overseeing cardiologist, and feedback to the patient following the outcome of the test. This communication complexity often leads to patients either not receiving the diagnostic equipment, not having the test results communicated to 55 them, or not having an appropriate follow up with a knowledgeable physician. Too commonly, no one calls the patient post-monitoring. In essence, the patient is left out-

side the health care channel and must wait until, and if,

he is informed of test results and follow up care. In addition, the reading of recorded ECG data usually requires an analysis by a heart disease specialist, which incurs additional cost and loss of time in making the patient diagnosis. Such back-end burdens can dissuade or even prevent a primary healthcare physician from ordering ambulatory ECG monitoring in the first place.

[0007] U.S. Patent application, Publication No. 2007/0255153, filed November 1, 2007, to Kumar et al. ; U.S. Patent application, Publication No. 2007/0225611, filed February 6, 2007, to Kumar et al.; and U.S. Patent application, Publication No. 2007/0249946, filed February 6, 2007, to Kumar et al. disclose a non-invasive cardiac monitor and methods of using continuously recorded cardiac data. A heart monitor suitable for use in primary care includes a self-contained and sealed housing. Continuously recorded cardiac monitoring is provided through a sequence of simple detect-store-offload operations that are performed by a state machine. The housing is adapted to remain affixed to a mammal from at least seven days up through four weeks. The heart monitor can include an activation or event notation button, the actuation of which increases the fidelity of the ECG information stored in the memory. The monitor is specifically intended to provide monitoring continuously and without interruption over an extended period, after which the stored information can be retrieved and analyzed offline to identify ECG events, including determining the presence of an arrhythmia.

[0008] Finally, U.S. Patent application, Publication No. 2008/0284599, filed April 28, 2006, to Zdeblick et al. and U.S. Patent application, Publication No. 2008/0306359, filed December 11, 2008, to Zdeblick et al., disclose a pharma-informatics system for detecting the actual physical delivery of a pharmaceutical agent into a body. An integrated circuit is surrounded by pharmacologically active or inert materials to form a pill, which dissolve in the stomach through a combination of mechanical action and stomach fluids. As the pill dissolves, areas of the integrated circuit become exposed and power is supplied to the circuit, which begins to operate and transmit a signal that may indicate the type, A signal detection receiver can be positioned as an external device worn outside the body with one or more electrodes attached to the skin at different locations. The receiver can include the capability to provide both pharmaceutical ingestion reporting and psychological sensing in a form that can be transmitted to a remote location, such as a clinician or central monitoring agency.

[0009] Therefore, a need remains for a way to provide primary care providers with direct on-hand access to ambulatory ECG monitoring without adding significant procedural obstacles, cost or time burdens, and which work within the confines of managed care, while simultaneously providing important diagnostic information and appropriate care directly to the patient through a closed health care loop.

[0010] US 2007/0255153 A1 discloses a method of ob-

taining and evaluating cardiac data, wherein a patient is provided at a health care provider with a cardiac monitoring patch having a unique patch identification. At the conclusion of the monitoring period the patient may mail the patch, using an envelope, to a processing center or health care provider for evaluation. As a result of the evaluation, the primary care physician may refer the patient to a cardiologist.

10 Summary

[0011] The invention is defined in the appended claims. [0012] A computer-implemented method for inexpensive and readily accessible ambulatory ECG monitoring 15 is provided. The ambulatory ECG monitoring is particularly adapted for use in managed healthcare by primary care physicians and their patients. Each ambulatory ECG monitor includes leadless sensing electrodes and low cost recording circuitry for short term or extended wear 20 monitoring. The patient wears the monitor for a set observation period and preferably maintains an electronic, vocal, or paper diary contemporaneous to monitoring. A unique identifier is assigned to the monitor that is used throughout the remainder of the diagnosis and referral 25 process. The patient sends the monitor (or its ECG data

recordings) and diary, if created, to a monitoring, consultation, and specialist referral center ("referral center") that retrieves and evaluates the recorded ECG data. In one embodiment, the referral center retains a cardiac specialist to interpret any ectopic findings and provide a for-

mal medical diagnosis. As a key and unique feature of this system, when appropriate, the patient is directly referred by the referral center to a cardiac specialist and an appointment is set up for definitive patient evaluation

and care, if the monitoring results indicate, rather than having the patient report back to his primary care physician. Additionally, the patient can proactively track the status of and make inquiries concerning his test results through the unique identifier, rather than relying on his
primary care provider's office staff or the physician himself. The primary care physician is also informed, but is not required to be involved in any manner. This process results in a rapid diagnosis, and importantly, rapid access

to a specialist, who can initiate definitive patient care, 45 when needed while enabling the patient to play an active role throughout each stage of the monitoring process. [0013] One embodiment provides a computer-implemented method for evaluating ambulatory electrocardiographic (ECG) monitoring of cardiac rhythm disorders. 50 A patient is registered online with a referral center and medical records for the patient are assembled in a centralized database maintained by the referral center. A tracking number is assigned to the monitor in relation to the ambulatory electrocardiographic (ECG) monitoring 55 and the tracking number is matched to the patient in the database. An ambulatory ECG monitor that includes leadless integrated sensing electrodes independently suspended from a flexible housing that encloses ECG

10

15

20

25

30

35

recording circuitry connected to the electrodes, is registered to the patient. An electrocardiogram is retrieved from the recording circuitry. The electrocardiogram and the medical records for the patient are evaluated against diagnostic criteria, which includes statistical correlations of cardiac disease states and cardiac rhythm patterns. Upon making a finding of a cardiac rhythm abnormality when the diagnostic criteria is met, the patient is referred to a cardiac rhythm specialist online, which includes sending the cardiac rhythm abnormality finding.

[0014] Primary care physicians are empowered with a type of ambulatory ECG monitoring that, in conjunction with a referral center, ensures proper data interpretation and medical follow up. A primary care physician need only apply an ambulatory ECG monitor in-clinic or provide a monitor to a patient through a prescription called into a pharmacy or other dispensary point-of-sale. Subspecialty expertise in arrhythmia diagnosis need not be resident in the provider's clinic, nor must the patient be referred to a separate ambulatory ECG testing laboratory. The low cost of each monitor encourages use when patient symptoms urge access to ambulatory ECG monitoring data. The backup system of support for the general physician helps minimize the risk of misdiagnosis and the need to even establish a referral, which is often not a simple decision or a simple process to ensure. Additionally, the combination of low cost and convenience of access to expertise encourages testing when appropriate to evaluating new medications or other changes important for the conduct of high-quality medical care.

[0015] Another key feature is that patients are empowered with the ability to self-screen a potential arrhythmic condition through ambulatory ECG monitoring. Access to cardiac rhythm expertise is difficult for a variety of reasons. Patients save both the costs and inconvenience of undertaking intermediate diagnostic testing, as typically required when undergoing conventional Holter-type ambulatory ECG monitoring, as well as avoid the risk of non-reimbursement that arises when they seek help outside their managed care plan. Patients are able to stay informed of their test results and follow on care without having to passively wait for follow up to occur. Moreover, wasted time is avoided by all interested parties.

[0016] Finally, as part of the system employed by primary care providers, cardiac specialists are empowered with receiving complete patient referrals and critical ECG data that enable them to effectively diagnose and treat arrhythmic conditions without the usual repetitive phone calls and requests to access medical information between doctors offices. Medical information and patientgenerated diary entries are communicated to the referral center as part of the ambulatory ECG monitoring process, which is provided to cardiac specialists as part of a complete referral.

[0017] Still other embodiments will become readily apparent to those skilled in the art from the following detailed description, wherein are described embodiments by way of illustrating the best mode contemplated. Accordingly,

the drawings and detailed description are to be regarded as illustrative in nature and not as restrictive.

Brief Description of the Drawings

[0018]

FIGURE 1 is a diagram showing, by way of example, a method for evaluating ambulatory electrocardiographic monitoring of cardiac rhythm disorders in accordance with one embodiment.

FIGURE 2 is a flow diagram showing the tracking of actions by a referral center as used in conjunction with the method of FIGURE 1.

FIGURE 3 is a screen shot showing, by way of example, a Web page for the tracking of actions by a referral center as used in conjunction with the method of FIGURE 1.

FIGURE 4 is a diagram showing, by way of example, in-clinic application of the ambulatory ECG monitor of FIGURE 1.

FIGURE 5 is a diagram showing, by way of example, an ambulatory ECG monitor in accordance with one embodiment.

FIGURE 6 is a diagram showing, by way of example, delivery of the ambulatory ECG monitor of FIGURE 1.

FIGURE 7 is a diagram showing, by way of example, evaluation of patient ECG data retrieved following the ambulatory monitoring of FIGURE 1.

FIGURE 8 is a block diagram showing, by way of example, levels of triage use by the referral center of FIGURE 6.

FIGURE 9 is a diagram showing a reimbursement scheme for health care provisioning participants.

Detailed Description

[0019] To the average internal medicine or family prac-40 tice physician, conventional forms of ambulatory ECG monitoring are only indirectly accessible and are rarely used because of the cost and inconvenience of access, diagnosis and follow-up referral to patient and physician alike. This situation hampers the evaluation of patients 45 with sporadic symptoms like syncope, palpitations, and, dyspnea, that are often not present at the time of visiting the doctor. Equipping these physicians with proper diagnostic tools is critical to proper patient management. Such tools must be highly-accessible, yet low-cost to en-50 sure use when needed. At the same time, patients need a sense of transparency in healthcare provisioning. Undertaking diagnostic testing is an accepted part of the healthcare process, yet once testing has been completed, patients can feel that they are at the mercy of the 55 healthcare system in terms of receiving follow up information and guidance. Such feelings of helplessness are only exacerbated as the waiting time for information and referral after test completion continues for days or weeks.

[0020] Both of these primary care physician and patient needs can be met through a system that forms a closedloop using ambulatory ECG monitoring. FIGURE 1 is a diagram showing, by way of example, a method for evaluating ambulatory electrocardiographic monitoring of cardiac rhythm disorders in accordance with one embodiment. Briefly, an ambulatory ECG monitor is packaged as an adhesive patch that can be applied in-clinic by a primary care provider, or at home by the patient. ECG data is recorded and subsequently read by a monitoring, consultation, and specialist referral center ("referral center") with which the patient can automatically be enrolled using tracking information specific to the patient and embedded into the monitor using, for instance, a radio frequency identification tag. Based on the recorded ECG data, diagnosis and referral to cardiac or other medical specialists, as appropriate, are made. The referrals are made by the referral center, not the physician who applied or prescribed the ECG monitor. Likewise, the referrals do not require further in-clinic primary care services, which is a key feature of that is particularly helpful to the patient. Moreover, the patient can track and make inquiries regarding the status of his test results on demand throughout each stage of diagnosis and referral by tracking the information and the results of the information contained in his ECG monitor. Thus, by assuring access to expert medical specialist referral by automatically scheduling an appointment for the patient with the most pertinent physician based on the patient's ECG findings and by enabling the patient to be a part of the post-monitoring process, the key omissions in conventional patient management are resolved.

[0021] The scenario for use will take the following course. A patient 12, possibly suffering from cardiac arrhythmia, experiences classic symptoms, such as syncope, presyncope, palpitations, angina, dyspnea, and fatigue (step A) will become concerned and sees his primary care physician 14 for an in-clinic appointment. During the appointment, the physician 14 or an assistant, such as a nurse, applies an ambulatory ECG monitor 15 to the patient's chest when presented with indications for ambulatory ECG monitoring (step B), as further described below with reference to FIGURE 4. (The size of the monitor 15 in FIGURE 1 et seq. is exaggerated for clarity.) Alternatively, the physician 14 may give the patient 12 the monitor 15 to be applied later on by the patient 12 himself or by an assistant in another place, such as at home. In a further embodiment, the patient 12 could self-screen by obtaining an ambulatory ECG monitoring kit over-the-counter or, if required, under a prescription called into a pharmacy by his primary care physician, such as described in U.S. Patent application US 2012/089417 A1, entitled "Computer-Implemented System and Method for Mediating Patient-Initiated Physiological Monitoring". The self-screening monitoring kit would include instructions on how to either physically return the monitor 15 or to electronically transfer the recorded physiological data from the monitor 15 to the referral center following monitoring.

[0022] Either prior to seeing the physician 14 or following application of the monitor 15, the patient 12 is registered with the referral center, if he was not already registered before. The patient 12 also receives instructions, such as from his physician's nurse 16, regarding physically sending the monitor 15 (or its ECG data recordings) to the referral center following monitoring and on how to track the status of the test results and follow on care

¹⁰ online (step C). A tracking number ("TN") 19 is assigned to the monitor 15 that follows the monitor 15 throughout monitoring of the heart beat, reading its output, follow up, diagnosis and referral, which is also provided to the patient 12. The nurse 16 electronically matches the tracking

¹⁵ number 19 to the patient 12 in a database centrally maintained by the referral center online. The nurse 16 also uploads the patient's personal medical information, thereby creating a complete medical record that can be used by a cardiac specialist in the event of referral.

20 [0023] The patient 12 is given a sealable envelope 18 or similar container in which to physically enclose and send off the used monitor 15 to the referral center. The envelope 18 includes an RFID tag 20 and a detachable tracking ticket 22. The nurse 16 uses an RFID transceiv-

er-equipped computer, including a desktop, notebook, or tablet computer, or mobile computing device, such as a smart telephone, to read from and record into the RFID tag 22. The computer includes those components conventionally found in general purpose programmable de vices, such as a central processing unit, volatile memory.

vices, such as a central processing unit, volatile memory, input and output ports, user display, keyboard or other input device, network interface, and non-volatile mass storage. Other components are possible. The nurse 16 records the tracking number 19 into the envelope's RFID

tag 20 and onto the tracking ticket 22. Alternatively, the tracking number 19 could originate as a pre-generated code already printed on the tracking ticket 22 and be recorded into both the envelope's RFID tag 20 and an RFID tag provided with the monitor 15, as further described *infra*. The patient 12 is instructed to detach the tracking ticket 22 on which the tracking number 19 has been recorded prior to sending off the monitor 15. In a further embodiment, the patient 12 electronically trans-

mits the recorded ECG data to the referral center following completion of monitoring.

[0024] At the same time that post-monitoring instructions are received, the patient 12 is given a diary 17 to help chronicle physical symptoms, subjective feelings, and activities at the time of symptomatic onset during
⁵⁰ monitoring. A related challenge facing physicians is a lack of contemporaneous subjective data from the patient himself concerning his complaints at the time of occurrence. To address this concern, the patient 12 is expected to maintain the diary 17 throughout the time that the monitor 15 is worn. In one embodiment, the patient 12 is either provided with a form of electronic or traditional paper diary 17 or instructed on accessing a dictation service that makes diary entries for the patient. In a further embodi-

5

ment, where the patient obtains a monitor 15 over-thecounter or through a prescription called in by the primary care provider 14, the monitoring kit includes a diary 17. [0025] The diary 17 can be implemented in the form of software, technology-assisted dictation, or conventional writing that is later electronically transcribed. A softwareimplemented diary can be provided in several forms. For instance, the diary software could be distributed on machine-readable media, or downloaded online, which, when installed on a computing device, locally executes an electronic diary application. The computing device can be a computer workstation, personal computer, personal digital assistant, programmable or "smart" mobile telephone, such as an iPhone or Blackberry, respectively licensed by Apple Inc., Cupertino, CA and Research In Motion, Waterloo, Ontario, Canada, intelligent digital media player, mobile tablet computer, or other programmable stationary or portable electronic device. The computer includes those components conventionally found in general purpose programmable devices, such as a central processing unit, volatile memory, input and output ports, user display, keyboard or other input device, network interface, and non-volatile mass storage. Other components are possible. Alternatively, the diary software could be a virtual application loaded from a remote server, such as a Web-based application that executes in a Web browser on the patient's computing device. As well, the diary software could be part of an existing software application, like a personal information manager and communications program, such as Outlook, licensed by Microsoft Corporation, Redmond, WA, or a social networking and microblogging service, such as Twitter, licensed by Twitter, Inc., San Bruno, CA. Preferably once a day or as appropriate, the patient 12 types or dictates entries into the diary 17. A technology-assisted dictated diary uses an existing electronics infrastructure to accept spoken diary entries from the patient 12, who is asked to telephone a call monitoring center to dictate his diary entries daily. This form of diary is particularly apropos for those patients who lack access to or knowledge of using a computing device. When dictated, voice recognition software executing at the call center converts patient's spoken words into text, or the speech can be manually transcribed into text offline by a third party transcription service. A conventional written diary is typically a bound notebook or other form of printed material into which the patient 12 manually chronicles his daily activities and physical complaints. Diary entries are later converted into electronic form by the referral center or a third party transcription service. Other forms of diary are also possible. [0026] During monitoring (step D), the patient 12 engages in activities of daily living, while the monitor 15 unobtrusively monitors and records ECG data. Continuous and uninterrupted wear of the monitor 15 over the entire course of monitoring may be impracticable for every patient. Skin sensitivities, allergies, irritation, and similar factors have an effect on a patient's ability to tolerate the wearing of the monitor 15 for an extended period.

Similarly, oil on the skin's surface, perspiration, and overall physical hygiene can affect monitor adhesion. Thus, each monitor 15 includes a flexible housing and standoffseparated skin adhesion assembly. The housing can be separated from the skin adhesion assembly to allow the patient 12 to reposition or replace the skin adhesion assembly. Either the same housing or a new housing can be used during successive periods of monitoring. When

the same housing is reused, the recording circuitry compensates for disconnection and reconnection of the sensing electrodes by stopping recording of ECG data during the gap in monitoring, as sensed by disconnection from the set of sensing electrodes. The recording circuitry thereafter resumes recording upon being reconnected.

¹⁵ If necessary, the patient 12 may choose to take a break and allow her skin to "breathe" between applications of the skin adhesion layer.

[0027] Throughout the monitoring period, the patient ideally makes regular entries in his diary 17, as described *infra,* which provides context that can help relate patient symptoms to the recorded ECG data, even where there are gaps in the monitoring due to replenishment of the skin adhesive layer or other factors. In a further embodiment, the monitor 15 includes an actimetry sensor, which

measures gross motor activity undertaken by the patient, such as through walking, running, changing posture or sleep position, and other body motions, that can be temporally matched with diary entries during post-monitoring evaluation. Monitoring ends when the patient 12 removes
the monitor 15 from his chest. Thereafter, the patient 12 sends the monitor 15 (or its ECG data recordings) to the referral center, along with his completed diary 17 (step E), as further described below with reference to FIGURE

6. Upon receipt, a laboratory technician 23 or other personnel retrieves the recorded ECG data 25 and, if provided, the actimetry, from the monitor 15 (step F) for analysis, post-monitoring diagnosis, follow up, and referral, as further described below with reference to FIGURE 7.
[0028] The tracking number 19 allows both the referral
center and the patient 15 to follow the use of the monitor

center and the patient 15 to follow the use of the monitor 15 from application through reading, diagnosis, and referral. In-clinic, the primary care physician 14 or his assistant use a computer, including a desktop, notebook, or tablet computer, or mobile computing device, such as

45 a smart telephone, to access the RFID tags of the monitor 15 and envelope 18. The ability ascertain the whereabouts and status of the monitor 15 and its recorded data at any point in the process is empowering. Thus, in response to an inquiry sent to the referral center by the 50 patient 12 and keyed to the tracking number 19, the patient 12 can determine the status of ambulatory ECG monitoring, data retrieval, ECG data evaluation, any clinical findings, including findings of a cardiac rhythm abnormality, or a referral to an appropriate medical special-55 ist. Primary care physicians can establish patient compliance. Cardiac specialists are able to reconstruct the steps leading up to a referral. The referral center maintains positive control. And patients are granted the ability

10

15

20

to close the loop on their health care provisioning by knowing where they are in the diagnostic process, and, equally important, where they are headed in the process of their follow-up, subsequent care and therapy. In sum, patients are provided closure for their problem, which is increasingly infrequent in today's healthcare system. **[0029]** Each monitor 15 and its recorded data are tracked using a combination of the tracking number 19 and a series of scheduled actions. The actions are tracked and coordinated by the referral center through an online database. FIGURE 2 is a flow diagram showing the tracking of actions 30 by a referral center as used in conjunction with the method 10 of FIGURE 1. The actions include:

(1) Each monitor 15 is expected to be sent off by the patient 12 and received at the referral center within a fixed amount of time following completion of monitoring (block 31). For instance, a period of 48 hours may be allotted for a patient 12 to provide the monitor 15 or its recorded data to the referral center, where the data is transmitted electronically in lieu of physically sending the monitor 15. Upon the expiry of 48 hours or other allotted time (block 32), the referral center will initiate follow up (block 33), which can include contacting the carrier, such as Federal Express, or the patient 12. Follow up may be electronic, for instance, by accessing an online package tracking database or sending text messages, or via phone or mail communication.

(2) Analysis and diagnosis is performed by the referral center within a fixed amount of time upon receipt of each monitor 15 or its recorded data (block 34). For instance, a period of 24 hours may be allotted for completing data analysis and diagnosis. Failure to complete analysis and diagnosis within the allotted time (block 35) is flagged for internal follow up by the referral center (block 36). A physician, either primary care or referred, or the patient 12 is permitted to inquire with the referral center into the nature of any delay.

(3) Follow up with the patient 12 is by the referral center is guaranteed within a fixed amount of time following completion of analysis and diagnosis (block 37). For instance, a period of 24 hours may be allotted for patient follow up, which can include contacting the patient 12 with status information, generating a referral to a cardiac specialist, and no-tifying the patient 12 of their cardiac specialist appointment. Failure to complete patient feedback within the allotted time (block 38) is flagged for internal follow up by the referral center (block 39). A physician, either primary care or specialist, or the patient 12 is permitted to inquire with the referral center into the nature of any delay.

(4) Where a referral to a cardiac specialist is generated, the test results and patient medical records must be sent to the referred physician within a fixed amount of time following patient follow up (block 40). For instance, a period of 24 hours may be allotted for forwarding the test results. The test results can be transmitted electronically to the referred physician. Patients 12 and authorized physicians can access the test results through a secure Web page. Failure to forward test results within the allotted time (block 41) is flagged for internal follow up by the referral center (block 42). A physician, either primary care or specialist, or the patient 12 is permitted to inquire with the referral center into the nature of any delay.

(5) The patient 12 is expected to confirm his appointment with the referred physician within a fixed amount of time following patient follow up (block 43). For instance, a period of 24 hours may be allotted for the patient 12 to confirm. The patient 12 can confirm electronically, for instance, by email or text message, or manually. Upon the expiry of 24 hours or other allotted time (block 44), the referral center will initiate follow up with the patient 12 (block 45). Follow up may similarly be electronic, voice or mail.

[0030] In one embodiment, the referral center provides 25 Web-based applications for physicians, their staff, and patients that enable them to access the database with proper authorizations and privacy protections, as further described below with reference to FIGURE 3. Additionally, the referral center can send electronic status up-30 dates via email, text message, or automated telephone calls to the patients, as well as others, upon the completion of each of the foregoing actions, as well as other actions. In a further embodiment, the physicians, their staff, and the patients can also contact the referral center 35 manually, such as via a toll free telephone number. For instance, a patient 12 may wish to speak to an on-call nurse at the referral center concerning the test results or their scheduled referral. Other ways to interface with the referral center are possible.

40 [0031] In one embodiment, status can be checked by accessing the referral center's Web site and entering the tracking number. FIGURE 3 is a screen shot showing, by way of example, a Web page 47 for the tracking of actions 30 by a referral center as used in conjunction with

⁴⁵ the method 10 of FIGURE 1. The Web page is viewable using a Web browser or similar application with online access as executed using a Web-capable device, including computer workstation, personal computer, personal digital assistant, programmable or "smart" mobile telephone, intelligent digital media player, mobile tablet com-

puter, or other programmable stationary or portable electronic device.

[0032] The Web page 47 displays the status of the actions 30 undertaken following completion of ambulatory
 ⁵⁵ ECG monitoring. The patient 12 retrieves the Web page 47 by entering the tracking number 48 or other identifying information. The status of the testing 49 is then presented in a secure manner than ensures patient privacy, for in-

stance, by using secure socket layer (SSL) communications security. The status information can include, for instance, a summary of the patient information on file and a history of the actions undertaken by the referral center. Other kinds of status information, forms of processing status inquiries, and of presenting status information are possible.

[0033] The monitor provides frontline primary care physicians with a simple and low-cost form of ambulatory ECG monitoring without the obstacles inherent in acquiring the device, the complexities and inefficiencies of calling and contacting referral offices, and the unacceptable costs in terms of time and repetitive travel that are attendant to conventional ambulatory ECG monitors. FIG-URE 4 is a diagram showing, by way of example, in-clinic application 50 of the ambulatory ECG monitor of FIGURE 1. To encourage proactive diagnosis of arrhythmias or other conditions of medical concern, the ambulatory ECG monitors are provided as single use monitoring packages that are low cost and readily dispensable, such as described in commonly-assigned U.S. Patent application US 2012/089001 A1, entitled "Ambulatory Electrocardiographic Monitor and Method of Use", and U.S. Patent application US 2012/089000 A1, entitled "Ambulatory Electrocardiographic Monitor for Providing Ease of Use in Women and Method of Use". Other types of ambulatory ECG monitors could also be used.

[0034] The low cost and convenience of access of ambulatory ECG monitors encourages rapid patient diagnostic testing, which is currently underutilized in primary care medicine because of the inconvenience and difficulty generally encountered in accessing conventional ambulatory ECG monitors and the difficulty of diagnosis and receiving actionable information. Here, a primary care physician 14 can keep a set of the pre-packaged ambulatory ECG monitors 52 readily at-hand (step A), for instance, in his laboratory coat pocket 53 or on an examination room table. Ambulatory ECG monitoring can be undertaken in a wider range of circumstances than practicable with conventional forms of ambulatory monitoring, which entail significant care giver, patient, and financial burden. For instance, a primary care provider 14 can use the ambulatory ECG monitors 52 for evaluating the efficacy of new medications or other changes to patient care. Similarly, a series of ambulatory ECG monitors 52 can be used over the course of several months to establish a trending analysis for a patient 12 with particularly sporadic symptoms, such as episodic paroxysmal atrial fibrillation. The threshold of use is low enough and the back up reading support and expert referral, when necessary, are integral parts of the system. Thus, a primary care provider 14 can freely apply the ambulatory ECG monitors 52 without distraction from managed health care limits or concerns of being outside his range of expertise, both in the case of interpreting ambulatory ECG monitoring data and, importantly, in managing its results. In turn, the patient 12 receives in a timely manner the full spectrum of appropriate responses

to his symptoms.

[0035] During examination, as appropriate, the primary care provider 14 takes a patient history 55 and performs a physical examination 39 (step **B**). A complete patient history 55 indicates how frequent the patients presumed arrhythmia arises, its duration, the severity of symptoms, heart rate, drug and dietary history, systematic illnesses, and family history of rhythm disturbances. The patient history 55 elicits major cardiovascular symptoms and

¹⁰ how they may have varied over time. Classic cardiovascular symptoms can include chest discomfort, dyspnea, fatigue, edema, palpitations, syncope, coughing, hemoptysis, and cyanosis. The physical examination 39 includes findings from physical inspection, palpitation, per-

¹⁵ cussion, and auscultation, which may help provide clues that link disparate aspects of the patient's presentation. Findings from the patient medical history 55 and physical examination 39 are used during subsequent online diagnosis by a server operated as part of the referral center,

²⁰ as further described below with reference to FIGURE 7. [0036] Through use of the monitor 15, both physicians and patients enjoy an ease-of-use not found with conventional ambulatory ECG monitors. Ambulatory ECG monitoring can be initiated when indicated by patient con-

dition or when otherwise considered appropriate by the physician 14. Referring briefly ahead to FIGURE 5, each monitor includes a pair of leadless sensing electrodes 63a, 63b exposed on the underside of an adhesive layer 62 and connected to low cost ECG recording circuitry
integrated in a compact single-use housing 61. Each

monitor also includes an RFID tag 66 containing a unique identifier for the monitor that is either integrated with the ECG recording circuitry, or is embedded into the monitor's housing, such as within a foam-constructed cover.

The RFID tag 66 is used during monitoring to associate a monitor 15 to a patient-specific tracking number 19 that can be used by the patient 12, referral center, and physician or staff to track the physical whereabouts of the monitor 15 and to determine the post-monitoring status
 of diagnosis and follow up care. The RFID tag 66 is accessed using standard RFID transmitter and receiver

units. Referring back to FIGURE 4, the monitor 15 is [0037] placed on the patient's chest at midline, covering the 45 center third of the sternum between the manubrium and the xiphoid process that is s found at the inferior limit of the sternum (step C). The monitor 15 can be applied by the primary care physician 14, or an assistant, such as a nurse. This unique location for ECG monitor application 50 and the monitor's small size allow for a uniformity of applicability by minimally trained physicians or even lay individuals. Application of the monitor 15 is no more difficult than applying a bandage over a wound. Monitor application takes a few seconds and is significantly more time 55 and cost efficient than both making a cardiac specialist referral, which often requires writing letters and making telephone calls, or sending a patient to a hospital or major cardiology practice to access an ambulatory ECG monitor. Unlike application of the twelve leads used by standard Holter systems, non-expert individuals have the ability to successfully apply the monitor 15 without needing medical training or separate instruction. In addition, the midline sternum-centered location is ideal for demonstrating key ECG features of atrial activity, such as Pwaves, as well as ventricular activity, such as R-waves. [0038] In a further embodiment, ambulatory ECG monitoring can be initiated through self-care. Where available without a prescription, the patient 15 can simply obtain and apply an ambulatory ECG monitoring kit over-thecounter. Alternatively, the primary care provider 14 can call in a prescription to a pharmacy or other dispensary point-of-sale for the patient 12 to obtain a single use ambulatory ECG monitoring kit. The patient 12 may have already performed his own research as to his condition or spoken to a physician, nurse practitioner, or other health care professional about his perceived health concerns through telephonic "on-call" patient services. The patient 12 can request his primary care provider 14 to call in a prescription for an ambulatory ECG monitor. The patient thus avoids the need to see the primary care provider 14 in-person through an in-clinic appointment and can apply the monitor 15 himself.

[0039] Physical application of the monitor 15 is the only step required to initiate ambulatory ECG monitoring. Minimal to no primary care provider involvement need be undertaken prior to starting ambulatory ECG monitoring. Conventional follow up appointments with an ECG monitoring laboratory and a primary care physician-referred cardiac specialist are unnecessary, at least at this point. Instead, the patient 12 takes care of returning the monitor 15, or, in a further embodiment, just the ECG data recorded by the monitor 15, for reading and follow up by the referral center, along with the diary 17 as maintained by the patient 12 throughout monitoring.

[0040] Finally, contemporaneous to application of the monitor 15, the primary care provider 14, or an assistant, electronically matches the monitor 15 to the patient 12 using the monitor's unique identifier that is stored in the RFID tag 66 (step **D**). Each monitor 15 is registered for use by the patient and is generally for one-time use. An RFID transceiver-equipped computer 58, including a desktop, notebook, or tablet computer, or mobile computing device, such as a smart telephone, scans the RFID tag 66 and records the monitor's unique identifier into a database centrally maintained by the referral center on-line.

[0041] A small, anatomically adaptive, and single-use ambulatory ECG monitor is applied in-clinic by a primary care provider 14, at home by the patient 12, or by other healthcare or lay individuals to record ECG data over an extended time period, while the patient 12 engages in activities of daily living. FIGURE 5 is a diagram showing, by way of example, an ambulatory ECG monitor 60 in accordance with one embodiment. The monitor 60 is placed on the midline of the patient's chest and centered over the sternum. Conforming fit and secure adhesion to the inherently uneven surface of a midline sternum-centered location are provided through two interconnected structures: a flexible housing for the electronic circuitry and standoff-separated skin adhesion assembly specif-

⁵ ically designed to adapt to a variable contour sternal surface. A waterproof housing 61 encloses ECG recording circuitry. The housing 61 is independently suspended through a set of standoffs that forms a gap 64 of about 2.5mm (0.lin) between an adhesive layer 62 that con-

formably adheres to the patient's skin and the housing 61. A set of at least two electrodes 63a, 63b are received in holes or "gel wells" formed through the standoffs and are exposed at the bottom surface of the adhesive layer 62.

¹⁵ [0042] Physically, when viewed from above, the monitor 60 has an elongated triangular shape with rounded vertices with dimensions of approximately 3.8cm (1.5in) wide and 7.6cm (3.0in) long. The monitor 60 weighs about 14.2g (0.5oz) when assembled with electrodes and

²⁰ a waterproof housing for the ECG recording circuitry, although a weight of up to 28g (1.0oz) would be acceptable. The electrodes 63a, 63b are spaced less than 6cm apart, and, when adhered to the patient, are aligned and placed midline sternum-centered. In one embodiment, two elec-

trodes are used, although three or more electrodes could also be used. When adhered onto a patient's sternum, the narrowest part of the monitor 41 faces downwards towards the patient's feet. On a female patient, the narrow part of the monitor 60 fits partway into the upper intermammary cleft. In a further embodiment, the monitor 60 includes a patient-operable button 65 that allows the patient 12 to set an electronic marker in the recorded ECG, such as when she feels the onset of a symptom or episode. During post-monitoring evaluation, each electronic marker is matched to the ECG and diary entries.

[0043] Ambulatory ECG monitoring begins with clinic application of a monitor and ends upon removal of the monitor by the patient, who must then provide the recorded ECG data to the referral center. FIGURE 6 is a diagram
showing, by way of example, delivery 70 of the ambulatory ECG monitor 15 of FIGURE 1. The referral center 71 is an organizational entity typically independent from the primary care provider's clinic. The referral center 71 provides ambulatory ECG data reading, analysis, diag-

⁴⁵ nosis, and patient referral and follow up, such as described in commonly-assigned U.S. Patent application US 2012/089412 A1, entitled "Computer-Implemented System and Method for Facilitating Patient Advocacy through Online Healthcare Provisioning".

50 [0044] To encourage appropriate use by primary care physicians and patients alike, the referral center 71 offers several channels for receiving used monitors 15. First, the monitor 15 can be physically sent to the referral center 71 via a package carrier, delivery service, or postal service. Delivery can be as simple as placing the monitor 15 into a pre-addressed sealable envelope 18 received from the primary care physician's office and dropping the envelope 18 into delivery box 71, post office, shipping cent-

er, and the like. Second, the sealable envelope 18 could be sent using a courier or delivery service that is contracted to specifically deliver used monitors 15 to the referral center 71. The patient 12 places the monitor 15 into the sealable envelope 18, which is then dropped into a kiosk 72 or other pickup location designated by the courier or package delivery service. Finally, the patient 12 could just return the monitor 15 in the sealable envelope 18 to administrative personnel 73 at the clinic, a "help" desk at a hospital, or other locations manned by health care staff.

[0045] In a further embodiment, not forming part of the present invention, instead of physically returning the used monitor 15 to the referral center 71, the recorded ECG data can be delivered electronically. For example, the monitor 15 can include a USB or similar standardized data exchange interface 74 for plugging into a computer 75, including a desktop, notebook, or tablet computer, or mobile computing device, such as a smart telephone. Alternatively, the monitor 15 could include a removable standardized memory module, such as a Secure Data memory card. The personal computer reads the recorded ECG data and patient information and electronically transmits the data read to the referral center, either by dedicated private communications circuit or publiclyavailable data communications network, such as the Internet. As well, the monitor 15 could incorporate a wired data connection, such as an IEEE 802.3 Ethernet interface, or a wireless transmitter using, for instance, Bluetooth or Wireless Fidelity ("WiFi") technology to wirelessly relay the recorded ECG data to the referral center 71. Other forms of physical or electronic delivery of recorded ECG data are possible.

[0046] The low cost ambulatory ECG monitor facilitates flexible implementation with consistent and clinically-sound medical follow up that ensures resolution of the patient's arrhythmic heart disorder concerns, in other words, a closed loop system from the patient's perspective. FIGURE 7 is a diagram showing, by way of example, evaluation 80 of patient ECG data retrieved following the ambulatory monitoring of FIGURE 1. The referral center performs data interpretation and follow up with the patient directly.

[0047] The referral center first processes receipt of the monitor 15. If the monitor 15 is physically sent in a sealable envelope 18, a laboratory specialist 81 first ensures that the tracking number or other confirmatory information assigned to the monitor 15 correctly matches the sealable envelope 18 (step **A**). If mismatched, the laboratory specialist 81 notifies the primary care provider 14 that a problem exists with either the wrong sealable envelope having been provided to the patient 12, or incorrect encoding of the monitor 15. If the monitor 15 and sealable envelope 18 pairing is confirmed, the laboratory specialist 81 electronically retrieves the recorded ECG data 82 from the monitor 15 into a computer workstation or other intermediate electronic data repository (step **B**). The laboratory specialist 81 also processes any further patient medical information provided with the sealable envelope 18, such as the patient's diary 17. The first two steps are skipped if the recorded ECG data is electronically sent directly to the referral center 71.

⁵ **[0048]** The referral center 71 maintains a database 83 within which recorded ECG data 82, medical records and other quantitative information, and diary entries and other qualitative information are stored for each patient 12, as identified using the tracking number 19. A server 84, or

¹⁰ similar centralized computational device, evaluates the recorded ECG data 82 to identify cardiac rhythm abnormalities. The server includes those components conventionally found in general purpose programmable devices, such as a central processing unit, volatile memory, input

¹⁵ and output ports, user display, keyboard or other input device, network interface, and non-volatile mass storage. Other components are possible.

[0049] The electrocardiogram recorded by the monitor 15, like all electrocardiograms, is amplified, filtered, and digitized over the range of 0.05 Hertz (Hz) to 150 Hz bandwidth for adults, and up to a 250 Hz bandwidth for children with a sampling rate of between one Kilohertz (KHz) and two KHz. ECG tracings are compared by the server 84 to medical diagnostic criteria to identify specific

²⁵ cardiac abnormalities. The criteria may be the sole basis for initial diagnosis. However, the initial diagnosis may also be correlated against any clinical, physiological, or pharmacological findings available in the patient's electronic medical records. The most common diagnostic cri-

³⁰ teria are based on statistical correlations between well known cardiac and other potential disease states with particular cardiac rhythm patterns appearing in ECGs and are well represented in many textbooks of cardiology, such as P. Libby et al., "Braunwald's Heart Disease

 A Textbook of Cardiovascular Medicine," Chs. 11 and 12 (8th ed. 2008). Innumerable diagnostic algorithms have been devised to identify various disease states for these disorders. Consequently, for purposes of both clarity and brevity, the discussion herein focuses on the se quence of events that unfold should a particular rhythm

disorder become manifest on an ECG. [0050] In diagnosing cardiac arrhythmia, some arrhythmias are highly symptomatic, yet are not associated with any adverse outcomes, such as premature ventricu-

⁴⁵ lar complexes. Other arrhythmias may present with no symptoms, yet may still present a significant risk of stroke, such as atrial fibrillation. Accordingly, in addition to the diagnostic criteria of the purely ECG-based data, the patient's medical history and the daily entries in the patient's diary 17 can be persuasive in reaching a diagnosis, or guiding further diagnostic testing or physician referral. Inquiry into the patient's medical history examines:

 mode of onset of arrhythmic episodes: For example, palpitations that occur while exercising, or when frightened or angry, signify different cardiac proclivities than palpitations that occur at rest or which

10

35

awaken the patient from sleep. Similarly, lightheadedness or syncope when frightened or having blood drawn is significantly different than syncope without warning, with the latter carrying a more serious implication and is more often associated with serious rhythm abnormalities.

- mode of termination of ectopic episodes: Palpitations reliably terminated by breath holding or the performance of the Valsalva or other vagal maneuvers can be a reliable indicator of a congenital rhythm abnormality, like atrioventricular nodal re-entry.

Other inquiries are possible, such as those regarding cardiac or lung disease status, or overall health. The diary entries can be temporally matched to cardiac rhythm patterns in the recorded electrocardiogram, which can assist with pairing physiological symptoms identified in the ambulatory ECG data to the patient's activities of daily living and contemporaneous symptomatic complaints. Other uses of the diary entries and patient medical history are possible.

[0051] Based on established diagnostic criteria, a set of diagnostic findings 85 is generated by the server 84 (step **D**). The findings 85, along with the patient's diary entries, can then be provided to a clinical specialist 86 (step E), such as a cardiologist, who has pre-contracted with or been retained by the referral center 71 to interpret diagnostic findings 85 and to make a formal medical diagnosis 87. The diary entries help temporally link objective symptomatic data recorded by the monitor to subjective observations made by the patient concerning his activities and physical complaints during the monitoring period. Diary entries are particularly helpful in placing patient symptomatic complaints in context. Unfortunately, a common outcome in modern managed care medical practices, where patients rarely see the same physician, key circumstances attendant to a complaint may be overlooked or forgotten by the original attending physician who ordered the ambulatory ECG recording in the first place. By storing the patient's medical history, electronic medical records, ambulatory ECG recording, and diary entries in one centralized database 83 maintained by the referral center 71 and by pre-contracting with expert assistance in the geographic area in which the patient lives, a patient can be given immediate and, importantly, pertinent referrals and thereby pertinent medical care.

[0052] Based on the diagnosis 87, the referral center 71, and not the initial source of ECG monitor application, that is, the primary care physician 14, performs medical follow up directly with the patient 12, which can include sending the diagnosis 87 to one or more clinical specialists 88a-c (step \mathbf{F}), such as cardiologists or cardiac electrophysiologists when the findings indicate a cardiac rhythm disorder, and notifying the patient 12 of the follow up referrals that were arranged. This step not only jump starts the patient management process in the event of a serious diagnosis, but also ensures that patients receive appropriate care and are not lost in the confusing shuffle

of modern medicine. The patient 12 remains an active part of the care provisioning and can inquire at any stage as to the whereabouts and status of his test results and follow up and the health care loop is thereby closed, without the patient being left in the open, that is, as a passive actor awaiting notification that may, or may not, happen. The primary care provider 14 is also notified of the type of follow up taken, but is not a part of the continuing health care provisioning process. Other forms of patient follow up are possible.

[0053] The referral center 71 triages medical follow up using, for instance, a three-tiered referral scheme. FIG-URE 8 is a block diagram showing, by way of example, levels of triage 90 use by the referral center 71 of FIGURE

¹⁵ 6. At the highest level "red" 91, a potentially life-threatening cardiac rhythm disorder is diagnosed, such as ventricular tachycardia, asystole or heart block. An appointment is automatically booked with a clinical specialist 88a-c, and the patient 12 is notified of the appointment

²⁰ immediately. His primary care provider 14 is also provided the diagnosis 87 and notice of the patient's cardiac specialist appointment and the urgency behind the referral. In a further embodiment, several clinical specialists 88a-c can receive the diagnosis 87 and the clinical spe-

cialist 88a-c that is ultimately assigned the referral is selected based on a selection criteria, such as first-to-respond or round-robin. Other selection criteria are possible, for instance, closest geographic location, earliest appointment opening, field of interest or expertise, such as
 a cardiac electrophysiologist specializing in heart rhythm

disorders, and the like.

[0054] At the second level "yellow" 92, a non-cardiac rhythm diagnosis 87 is made and the patient 12 is referred to a non-cardiac rhythm specialist (not shown). For ex-

ample, an ST segment depression warrants referral to a cardiologist specializing in coronary disease management, and not to a cardiac electrophysiologist. As well, widening of the QRS complex may best indicate the need for a referral to a heart failure expert, while QT interval

40 variation indicates appropriate follow up with an internal medicine physician to consider, for instance, drug-induced QT prolongation and the need for alternative drug therapy, for example, a different antibiotic. A loss of consciousness, despite normal laboratory ECG findings, im-

⁴⁵ plies non-heart rhythm problems and may indicate a need for referral to a neurologist or vascular surgeon. Other types of healthcare disorders can be diagnosed and referred to the appropriate physician in a timely and efficient manner without wasting time for either the patient or the ⁵⁰ primary care provider.

[0055] At the lowest level "green" 93, the patient 12 is considered normal or healthy, or at least with no discernible illness. No referral need be made and the patient 12 can be reassured, such as through a message from the referral center 71, that their heart rhythm is normal or that the symptoms may arise from a non-rhythm problem. Other levels of referrals and triage are possible.

[0056] Reimbursement for medical services provided

is a reality of managed care. The ambulatory ECG monitor 15 can be used within a medical insurance reimbursement structure that compensates the major service providers commensurate with the work performed. FIGURE 9 is a diagram showing a reimbursement scheme 100 5 for health care provisioning participants 14, 71, 86. Each of the primary care provider 14, the referral center 71, and the retained clinical specialist 86 provide services respectively in the initiation of ambulatory ECG monitoring, data collection and initial analysis, and interpretative 10 over read of a patient's ambulatory ECG data 82, activities that can entitle each party to some form of compensable reimbursement. For instance, the primary care provider 14 can receive reimbursement comparable to inclinic laboratory testing for application of a monitor 15 to 15 the patient 12. The referral center 71 bears the onus of the costs for consumable goods, infrastructure, and liaison between the patient 12, the primary care provider 14, and, when appropriate, cardiac specialists 97a-c. 20 Thus, the referral center 71 can receive reimbursement for providing the consumable goods, that is, the monitors 15, sealable envelopes 18, and diaries 17 to the primary care providers 14, or through over-the-counter or pharmaceutical channels, as appropriate.. The referral center 25 71 also can receive reimbursement for consolidating patient medical histories and electronic medical records and for the reading, processing, and storage of recorded ECG data 82. Finally, the retained clinical specialist 86 can receive reimbursement comparable to a professional reading of medical findings. Other reimbursement 30 schemes and tiers are possible.

[0057] While the invention has been particularly shown and described as referenced to the embodiments thereof, those skilled in the art will understand that changes in form and detail may be made therein without departing from the scope of the invention as defined in the appended claims.

Claims

1. A computer-implemented method (10) for evaluating ambulatory electrocardiographic (ECG) monitoring of cardiac rhythm disorders, comprising:

registering a patient (12) online with a referral center (71) and assembling medical records for the patient (12) in a centralized database (83) maintained by the referral center (71); providing an ambulatory ECG monitor (15) that comprises leadless integrated sensing electrodes (63a, 63b) independently suspended from a flexible housing (61) that encloses ECG recording circuitry connected to the electrodes (63a, 63b), to the patient (12);

assigning a tracking number (19) to the monitor (15) in relation to the ambulatory ECG monitoring and matching the tracking number (19) to the patient (12) in the database (83); recording the tracking number (19) into a RFID tag (20) included in an envelope (18) with which to send the monitor (15) to the referral center (71) for the evaluation following the completion of the ambulatory ECG monitoring; providing the envelope (18) comprising the RFID tag (20) to the patient (12) providing the tracking number (19) to the patient; tracking the physical whereabouts of the monitor (15) through the tracking number (19) of the RFID tag (20) prior to receipt of the monitor (15) for the evaluation;

retrieving an electrocardiogram (25) from the recording circuitry at the referral center (71); evaluating the electrocardiogram (25) and the medical records for the patient (12) against diagnostic criteria comprising statistical correlations of cardiac disease states and cardiac rhythm patterns at the referral center (71); upon making a finding (85) of a cardiac rhythm abnormality when the diagnostic criteria is met, referring the patient (12) to a cardiac rhythm specialist (86) online, including sending the cardiac rhythm abnormality finding (85); and in response to an online inquiry by the patient (12), determining a status of the evaluation and referral keyed to the tracking number (19).

- 2. A method (10) according to Claim 1, wherein the status concerns one or more of the ambulatory ECG monitoring, the retrieval of the electrocardiogram (25), the evaluation of the electrocardiogram (25), the finding (85) of a cardiac rhythm abnormality, and the referral to a cardiac rhythm specialist (86).
- **3.** A method (10) according to Claim 2, further comprising:
- setting a time period for one or more of the completion of the ambulatory ECG monitoring, the retrieval of the electrocardiogram (25), the evaluation of the electrocardiogram (25), the finding (85) of a cardiac rhythm abnormality, and the referral to a cardiac rhythm specialist (86); and undertaking a follow up upon expiry of one or more of the time periods.
 - **4.** A method (10) according to Claim 1, wherein the monitor (15) is placed on the patient's chest at midline, covering the center third of the sternum and centered between the manubrium and the xiphoid process on the inferior border of the sternum.
- ⁵⁵ **5.** A method according to claim 1, further comprising; providing a status upon demand regarding the analysis and referral to the patient (12) keyed to the tracking number (19); and

50

15

upon making a finding (85) of a cardiac rhythm abnormality when the diagnostic criteria is met, undertaking direct referral, comprising:

setting an appointment for the patient (12) with a cardiac rhythm specialist (86) online, including sending the electrocardiogram (25), medical records, and cardiac rhythm abnormality finding (85); and

notifying the patient (12) of the appointment with confirmation.

6. A method (10) according to Claim 5, the diagnostic criteria further comprising at least one of:

identifying physiological symptoms comprising one or more of arrhythmic episodes and ectopic episodes, as recorded during the monitoring period;

determining a mode of onset of each arrhythmic ²⁰ episode;

determining a mode of termination of each ectopic episode; and

comparing the physiological symptoms, the mode of onset of each arrhythmic episode, and ²⁵ the mode of termination of each ectopic episode to statistical correlations of cardiac rhythm patterns indicating possible cardiac disease states.

7. A method (10) according to Claim 5, further compris- ³⁰ ing one or more of:

timing receipt of the monitor (15) from the patient (12) following the completion of the ambulatory ECG monitoring;

timing performance of the analysis and referral of the patient (12);

timing notifying the patient (12);

timing sending of the electrocardiogram (25), medical records, and cardiac rhythm abnormality finding (85); and

timing confirmation of the patient's appointment,

wherein a time period for follow up is triggered if any one of the timings exceed a set time period. 45

- **8.** A method (10) according to Claim 7, wherein the follow up is undertaken by the referral center (71).
- **9.** A method (10) according to Claim 5, further compris- ⁵⁰ ing:

registering an electronically-stored diary (17) to the patient (12) with the referral center (71) under the tracking number (19);

recording entries into the diary (17) as provided by the patient (12) contemporaneous to the ambulatory ECG monitoring; temporally matching the diary entries to cardiac rhythm patterns found in the electrocardiogram (25); and

pairing the physiological symptoms with the diary entries.

- **10.** A method (10) according to Claim 9, further comprising one or more of:
- directly recording the diary entries into the diary in electronic form;
 electronically transcribing the diary entries from dictated verbal form; and
 - electronically transcribing the diary entries from written form.
 - **11.** A method (10) according to Claim 5, further comprising:

sensing disconnection of the recording circuitry from the electrodes (63a, 63b) during the monitoring; and

sending reconnection of the recording circuitry to electrodes (63a, 63b) and resuming the monitoring.

12. A method (10) according to Claim 5, wherein the medical records for the patient (12) comprise one or more of monitored patient identification data, prior electrocardiograms (25), laboratory results, physical examination results, exercise testing results, and medical history.

35 Patentansprüche

 Computerumgesetztes Verfahren (10) zur Auswertung der ambulanten Elektrokardiografie- (EKG) Überwachung von Herzrhythmusstörungen, umfassend:

> Online-Registrierung eines Patienten (12) bei einem Überweisungszentrum (71) und Zusammenstellung der Krankenakten des Patienten (12) in einer zentralisierten Datenbank (83), die im Überweisungszentrum (71) unterhalten wird; Bereitstellung eines ambulanten EKG-Überwachungsgeräts (15), das kabellose integrierte Messelektroden (63a, 63b) umfasst, die unabhängig von einem flexiblen Gehäuse (61), das einen mit den Elektroden (63a, 63b) verbundenen EKG-Aufzeichnungsschaltkreis umgibt, herabhängen, an den Patienten (12);

> Zuweisung einer Verfolgungsnummer (19) für das Überwachungsgerät (15) in Bezug auf die ambulante EKG-Überwachung und Abstimmung der Verfolgungsnummer (19) mit dem Patienten (12) in der Datenbank (83);

55

Aufzeichnung der Verfolgungsnummer (19) in einem RFID-Tag (20), das in einem Umschlag (18) steckt, mit dem das Überwachungsgerät (15) nach Abschluss der ambulanten EKG-Überwachung zur Auswertung an das Überweisungszentrum (71) geschickt werden soll;

Bereitstellung des Umschlags (18), der das RFID-Tag (20) umfasst, an den Patienten (12); Bereitstellung der Verfolgungsnummer (19) an den Patienten;

Verfolgung des Verbleibs des Überwachungsgeräts (15) durch die Verfolgungsnummer (19) des RFID-Tags (20) vor Erhalt des Überwachungsgeräts (15) zur Auswertung;

Abrufen eines Elektrokardiogramms (25) aus dem Aufzeichnungsschaltkreis im Überweisungszentrum (71);

Auswertung des Elektrokardiogramms (25) und der Krankenakten des Patienten (12) gegen Diagnosekriterien mit statistischen Korrelationen von Herzkrankheitszuständen und Herzrhythmusmustern im Überweisungszentrum (71);

bei Nachweis eines Befundes (85) einer Herzrhythmusstörung, wenn die Diagnosekriterien ²⁵ erfüllt sind, Online-Überweisung des Patienten (12) an einen Spezialisten für Herzrhythmusstörungen (86),

einschließlich Übersendung des Befundes der Herzrhythmusstörung (85); und

als Reaktion auf eine Online-Anfrage des Patienten (12) Bestimmung eines Status der Auswertung und Überweisung,

der mit der Verfolgungsnummer (19) verbunden ist.

- Verfahren (10) nach Anspruch 1, worin der Status eines oder mehrere der Aufgaben von ambulanter EKG-Überwachung, Abrufen des Elektrokardiogramms (25), Auswertung des Elektrokardiogramms (25), Befund (85) einer Herzrhythmusstörung und Überweisung an einen Spezialisten (86) für Herzrhythmusstörungen betrifft.
- 3. Verfahren (10) nach Anspruch 2, ferner umfassend: ⁴⁵

Festsetzung einer Zeitspanne für eine oder mehrere der Aufgaben von Beendigung der ambulanten EKG-Überwachung,

Abrufen des Elektrokardiogramms (25), Auswertung des Elektrokardiogramms (25), Befund (85) einer Herzrhythmusstörung und Überweisung an einen Spezialisten (86) für Herzrhythmusstörungen; und

Durchführung einer Kontrolle nach Ablauf von ⁵⁵ einer oder

mehreren der Zeitspannen.

- 4. Verfahren (10) nach Anspruch 1, worin das Überwachungsgerät (15) auf der Mittellinie der Brust des Patienten angebracht wird, so dass es das mittlere Drittel des Brustbeins überdeckt und zwischen dem Manubrium und dem Schwertfortsatz an der unteren Brustbeingrenze zentriert ist.
- 5. Verfahren nach Anspruch 1 ferner umfassend:
- 10

15

30

35

40

50

5

auf Verlangen Bereitstellung eines Status an den Patienten (12) in Bezug auf die Analyse und Überweisung, der mit der Verfolgungsnummer (19) verbunden ist; und

bei Nachweis eines Befundes (85) einer Herzrhythmusstörung, wenn die Diagnosekriterien erfüllt sind, direkte Überweisung, umfassend:

Online-Vereinbarung eines Termins für den Patienten (12) bei einem Spezialisten (86) für Herzrhythmusstörungen, einschließlich Übersendung des Elektrokardiogramms (25), der Krankenakten und des Befundes (85) der Herzrhythmusstörung; und Benachrichtigung des Patienten (12) über den Termin mit Bestätigung.

- **6.** Verfahren (10) nach Anspruch 5, wobei die Diagnosekriterien ferner zumindest eines der Folgenden umfassen:
 - Identifizierung von physiologischen Symptomen, umfassend eine oder mehrere von arrhythmischen Episoden und ektopischen Episoden, wie während des Überwachungszeitraums aufgezeichnet;

Bestimmung eines Modus des Einsetzens jeder arrhythmischen Episode;

Bestimmung eines Modus des Endes jede ektopischen Episode; und

Vergleich der physiologischen Symptome, des Modus des Einsetzens jeder arrhythmischen Episode und des Modus des Endes jeder ektopischen Episode mit statistischen Korrelationen von Herzrhythmusmustern als Hinweis auf mögliche Herzkrankheitszustände.

7. Verfahren (10) nach Anspruch 5, ferner umfassend eines oder mehrere der Folgenden:

Terminierung des Erhalts des Überwachungsgeräts (15) von dem Patienten (12) nach Abschluss der ambulanten EKG-Überwachung; Terminierung der Durchführung der Analyse und Überweisung des Patienten (12);

Terminierung der Benachrichtigung des Patienten (12);

Terminierung der Übersendung des Elektrokardiogramms (25), der Krankenakten und des Be-

10

20

25

35

fundes (85) der Herzrhythmusstörung; und Terminierung der Bestätigung des Termins des Patienten, worin eine Zeitspanne für die Kontrolle eingeleitet wird, wenn einer der Termine eine festgesetzte Zeitspanne überschreitet.

- Verfahren (10) nach Anspruch 7, worin die Kontrolle von dem Überweisungszentrum (71) durchgeführt wird.
- **9.** Verfahren (10) nach Anspruch 5, ferner umfassend:

Registrierung eines elektronisch gespeicherten Tagebuchs (17) für den Patienten (12) bei dem Überweisungszentrum (71) unter der Verfol-¹⁵ gungsnummer (19);

Erfassung von Einträgen in das Tagebuch (17) die von dem Patienten (12) bereitgestellt werden, zeitgleich mit der ambulanten EKG-Überwachung;

zeitlicher Vergleich der Tagebucheinträge mit den in dem Elektrokardiogramm (25) gefunden Herzrhythmusmustern; und

Paarung der physiologischen Symptome mit den Tagebucheinträgen.

10. Verfahren (10) nach Anspruch 9, ferner umfassend eines oder mehrere der Folgenden:

direkte Erfassung der Tagebucheinträge in dem ³⁰ Tagebuch in elektronischer Form;

elektronische Transkription der Tagebucheinträge aus einer diktierten mündlichen Form; und elektronische Transkription der Tagebucheinträge aus der schriftlichen Form.

11. Verfahren (10) nach Anspruch 5, ferner umfassend:

Nachweis eines Ablösens des Aufzeichnungsschaltkreises von den Elektroden (63a, 63b) 40 während der Überwachung; und Sendung einer Wiederverbindung des Aufzeichnungsschaltkreises an die Elektroden (63a, 63b) und Wiederaufnahme der Überwachung. 45

 Verfahren (10) nach Anspruch 5, worin die Krankenakten des Patienten (12) eine oder mehrere von überwachten Patientenidentifikationsdaten, früheren Elektrokardiogrammen (25), Laborergebnissen, ⁵⁰ Ergebnissen körperlicher Untersuchungen, Ergebnisse von Belastungstests und Krankengeschichte umfassen.

Revendications

1. Procédé mis en oeuvre par ordinateur (10) pour

l'évaluation du monitorage électrocardiographique (ECG) ambulatoire des troubles du rythme cardiaque, comprenant :

enregistrement d'un patient (12) en ligne auprès d'un centre d'orientation (71) et regroupement des dossiers médicaux du patient (12) dans une base de données (83) centralisée maintenue par le centre d'orientation (71) ;

fourniture au patient (12) d'un moniteur d'ECG ambulatoire (15) qui comprend des électrodes (63a, 63b) de détection intégrées sans fil suspendues indépendamment d' un boîtier souple (61) qui renferme un circuit d'enregistrement d'ECG raccordé aux électrodes (63a, 63b) ;

affectation d'un numéro de suivi (19) au moniteur (15) en relation avec le monitorage d'ECG ambulatoire et mise en correspondance du numéro de suivi (19) avec le patient (12) dans la base de données (83) ;

enregistrement du numéro de suivi (19) dans une étiquette RFID (20) incluse dans une enveloppe (18) avec laquelle il faudra envoyer le moniteur (15) au centre d'orientation (71) pour l'évaluation après avoir terminé le monitorage d'ECG ambulatoire ;

fourniture de l'enveloppe (18) comprenant l'étiquette RFID (20) au patient (12) ;

fourniture du numéro de suivi (19) au patient ; suivi des lieux où se trouvent le moniteur (15) par l'intermédiaire du numéro de suivi (19) de l'étiquette RFID (20) avant la réception du moniteur (15) pour l'évaluation ;

récupération d'un électrocardiogramme (25) auprès du circuit d'enregistrement au centre d'orientation (71) ;

évaluation de l'électrocardiogramme (25) et des dossiers médicaux du patient (12) par rapport à des critères de diagnostic comprenant les corrélations statistiques des états de maladies cardiaques et les modèles de rythme cardiaque au centre d'orientation (71) ;

sur résultat constaté (85) d'une anomalie du rythme cardiaque lorsque les critères de diagnostic sont remplis, orientation du patient (12) en ligne vers un spécialiste du rythme cardiaque (86), y compris envoi du résultat constaté (85) d'anomalie du rythme cardiaque ; et

en réponse à une demande en ligne par le patient (12), détermination d'un état de l'évaluation et de l'orientation codé avec le numéro de suivi (19).

 Procédé (10) selon la revendication 1, selon lequel l'état concerne une ou plusieurs des opérations suivantes : monitorage d'ECG ambulatoire, récupération de l'électrocardiogramme (25), évaluation de l'électrocardiogramme (25), résultat constaté (85)

10

15

30

45

d'une anomalie du rythme cardiaque et orientation vers un spécialiste du rythme cardiaque (86).

3. Procédé (10) selon la revendication 2, comprenant en outre :

définition d'une période pour l'achèvement d'une ou plusieurs des opérations suivantes : monitorage d'ECG ambulatoire, récupération de l'électrocardiogramme (25), évaluation de l'électrocardiogramme (25), résultat constaté (85) d'une anomalie du rythme cardiaque et orientation vers un spécialiste du rythme cardiaque (86) ; et réalisation d'un suivi à l'expiration d'une ou plu-

sieurs des périodes.

- Procédé (10) selon la revendication 1, selon lequel le moniteur (15) est placé sur la poitrine du patient sur la ligne médiane, couvrant le tiers central du sternum et centré entre le manubrium et l'appendice xiphoïde sur la limite inférieure du sternum.
- 5. Procédé (10) selon la revendication 1, comprenant en outre :

fourniture à la demande au patient (12) d'un état concernant l'analyse et l'orientation codé avec le numéro de suivi (19) ; et

sur résultat constaté (85) d'une anomalie du rythme cardiaque lorsque les critères de diagnostic sont remplis, réalisation d'une orientation directe comprenant :

prise en ligne d'un rendez-vous pour le patient (12) avec un spécialiste du rythme cardiaque (86), y compris envoi de l'électrocardiogramme (25), des dossiers médicaux et du résultat constaté (85) d'anomalie du rythme cardiaque ; et 40

notification au patient (12) du rendez-vous avec confirmation.

 Procédé (10) selon la revendication 5, les critères de diagnostic comprenant en outre au moins l'une des étapes suivantes :

> identification de symptômes physiologiques comprenant un ou plusieurs des épisodes d'arythmie et des épisodes ectopiques, tels ⁵⁰ qu'ils ont été enregistrés pendant la période de monitorage ;

détermination d'un mode d'amorçage de chaque épisode d'arythmie ;

détermination d'un mode de terminaison de chaque épisode ectopique ; et

comparaison des symptômes physiologiques, du mode d'amorçage de chaque épisode d'arythmie et du mode de terminaison de chaque épisode ectopique avec des corrélations statistiques des modèles de rythme cardiaque indiquant des états possibles de maladie cardiaque.

- **7.** Procédé (10) selon la revendication 5, comprenant en outre une ou plusieurs des étapes suivantes :
 - minutage de la réception du moniteur (15) de la part du patient (12) après avoir terminé le monitorage d'ECG ambulatoire ; minutage de la réalisation de l'analyse et de

l'orientation du patient (12) ;

- minutage de la notification du patient (12) ; minutage de l'envoi de l'électrocardiogramme (25), des dossiers médicaux et du résultat constaté (85) d'une anomalie du rythme cardiaque ; et
- minutage de la confirmation du rendez-vous du patient, une période de suivi étant déclenchée si l'un quelconque des minutages dépasse une période fixée.
- ²⁵ 8. Procédé (10) selon la revendication 7, selon lequel le suivi est entrepris par le centre d'orientation (71).
 - **9.** Procédé (10) selon la revendication 5, comprenant en outre :
 - enregistrement d'un journal (17) stocké électroniquement au patient (12) auprès du centre d'orientation (71) sous le numéro de suivi (19) ; enregistrement des entrées dans le journal (17) telles qu'elles sont fournies par le patient (12) au moment du monitorage d'ECG ambulatoire ; mise en correspondance temporelle des entrées du journal avec les modèles de rythme cardiaque trouvés dans l'électrocardiogramme (25) ; et

appariement des symptômes physiologiques avec les entrées du journal.

10. Procédé (10) selon la revendication 9, comprenant en outre une ou plusieurs des étapes suivantes :

enregistrement direct des entrées du journal dans le journal sous forme électronique ; transcription électronique des entrées du journal depuis la forme verbale dictée ; et transcription électronique des entrées du journal depuis la forme écrite.

11. Procédé (10) selon la revendication 5, comprenant en outre :

détection de la déconnexion du circuit d'enregistrement des électrodes (63a, 63b) pendant

le monitorage ; et détection de la reconnexion du circuit d'enregistrement aux électrodes (63a, 63b) et poursuite du monitorage.

12. Procédé (10) selon la revendication 5, selon lequel les dossiers médicaux du patient (12) comprennent un ou plusieurs des éléments suivantes :

données d'identification du patient soumis au ¹⁰ monitorage, électrocardiogrammes (25) précédents, résultats de laboratoire, résultats d'examens physiques, résultats de test d'effort et antécédents médicaux.

15

20

25

30

35

40

45

50









Fig. 3.



Fig. 5.

<u>60</u>







18 Post 71











Fig. 9.



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 3215136 A, Holter [0005]
- US 20070255153 A, Kumar [0007]
- US 20070225611 A, Kumar [0007]
- US 20070249946 A, Kumar [0007]
- US 20080284599 A, Zdeblick [0008]
- US 20080306359 A, Zdeblick [0008]

Non-patent literature cited in the description

• P. LIBBY et al. Braunwald's Heart Disease - A Textbook of Cardiovascular Medicine. 2008 [0049]

- US 20070255153 A1 [0010]
- US 2012089417 A1 [0021]
- US 2012089001 A1 [0033]
- US 2012089000 A1 [0033]
- US 2012089412 A1 [0043]

patsnap

专利名称(译)	计算机实现的动态心电图监测远程评估方法		
公开(公告)号	EP2438848B1	公开(公告)日	2014-03-19
申请号	EP2011184156	申请日	2011-10-06
[标]申请(专利权)人(译)	心脏科学公司		
申请(专利权)人(译)	心脏科技股份有限公司		
当前申请(专利权)人(译)	心脏科技股份有限公司		
[标]发明人	BARDY GUST H BISHAY JON MIKALSON		
发明人	BARDY, GUST H. BISHAY, JON MIKALSON		
IPC分类号	A61B5/00 A61B5/0402 G06F19/00		
CPC分类号	A61B5/0006 A61B5/0402 A61B5/0432 A61B2562/08 G16H10/60 G16H40/20 G16H80/00		
优先权	12/901461 2010-10-08 US		
其他公开文献	EP2438848A3 EP2438848A2 EP2438848B8		
外部链接	Espacenet		

摘要(译)

提供了一种用于评估心律紊乱的动态心电图(ECG)监测的计算机实现 的方法。患者(12)在线注册并且组装患者(12)的医疗记录。包括独 立悬挂在柔性外壳上的无引线集成感测电极的移动ECG监视器(15)被 注册到患者(12)。监视器被发送到转诊中心,例如使用预先设定的可 密封信封(18)。从记录电路检索心电图(25)。根据诊断标准评估心 电图(25)和医疗记录。在满足诊断标准时进行发现时,患者(12)在 线被称为心律专家,其包括发送心律异常发现。因此,医生和患者都享 有传统动态心电监护仪所不具备的易用性(15)。通过绕过确定是否需 要转诊并单独建立转诊,患者(12)可以比通过常规患者转诊更完整和 更快速地治疗,特别是对于严重疾病,患者成本更低且延迟更少。

