



(11) **EP 2 049 007 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
19.09.2018 Bulletin 2018/38

(51) Int Cl.:
A61B 5/00 ^(2006.01) **A61K 31/519** ^(2006.01)
A61K 31/7076 ^(2006.01) **A61P 9/08** ^(2006.01)

(21) Application number: **07765052.1**

(86) International application number:
PCT/EP2007/005923

(22) Date of filing: **04.07.2007**

(87) International publication number:
WO 2008/003479 (10.01.2008 Gazette 2008/02)

(54) **METHODS, COMPOSITIONS, UNIT DOSAGE FORMS, AND KITS FOR PHARMACOLOGIC STRESS TESTING WITH REDUCED SIDE EFFECTS**

VERFAHREN, ZUSAMMENSETZUNGEN, EINHEITSDOSISFORMEN UND KITS FÜR
PHARMAKOLOGISCHE BELASTUNGSTESTS MIT VERRINGERTEN NEBENWIRKUNGEN

PROCÉDÉS, COMPOSITIONS, FORMES PHARMACEUTIQUES UNITAIRES ET KITS POUR
ÉPREUVES DE STRESS PHARMACOLOGIQUE ASSOCIÉS À DES EFFETS SECONDAIRES
RÉDUITS

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE
SI SK TR**

(30) Priority: **05.07.2006 US 818928 P**
02.07.2007 US 772684

(43) Date of publication of application:
22.04.2009 Bulletin 2009/17

(73) Proprietor: **Adenobio N.V.**
1118 BH Amsterdam Schiphol (NL)

(72) Inventor: **GORNY, Philippe**
75116 Paris (FR)

(74) Representative: **J A Kemp**
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(56) References cited:
EP-A- 0 984 280 WO-A-00/78774
WO-A-02/39124 US-A- 5 919 614

- **CONRADSON ET AL: "Cardiovascular effects of infused adenosine in man: potentiation by dipyridamole" ACTA PHYSIOLOGICA SCANDINAVICA, OXFORD, GB, vol. 129, no. 3, 1987, pages 387-391, XP009090152**
- **NOTT M W: "THE POSSIBLE ROLE OF ADENOSINE IN THE CORONARY DILATOR ACTION OF SOME PYRIMIDO PYRIMIDINES AND PTERIDINES" BRITISH JOURNAL OF PHARMACOLOGY, vol. 39, no. 2, 1970, pages 287-295, XP009090150 Sweden**
- **MAHMARIAN J J ET AL: "Myocardial perfusion imaging during pharmacologic stress testing" CARDIOLOGY CLINICS, W.B. SAUNDERS COMPANY, PHILADELPHIA, US, vol. 12, no. 2, 1994, pages 223-245, XP009090182 ISSN: 0733-8651**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

1. BACKGROUND

[0001] Functional assessment of myocardium, in particular the evaluation of the myocardium's oxygen status, is important in guiding therapeutic decisions in the care of patients with cardiac ischemia. In current clinical practice, myocardial ischemia status is most often assessed using non-invasive nuclear perfusion imaging methodologies, such as planar scintigraphy or single photon emission computed tomography (SPECT), with thallium and technetium as the most frequently used isotopes. More recently, positron emission tomography (PET) with rubidium-82 has been gaining recognition as providing improved images with less radiation. Semi-invasive transesophageal doppler echography is also useful to study the motion of ventricular walls, and non-invasive transthoracic doppler echocardiography is an easy and non-invasive methodology for measurement of coronary flow reserve.

[0002] These functional tests typically require that the patient's heart be "stressed", either through controlled exercise or by pharmacologic means, and are thus generically and colloquially known as "stress tests". Pharmacological stressors for functional assessment of myocardium act through coronary vasodilation: by dilating normal vessels to a greater extent than diseased vessels, these agents establish a shunt, or "myocardial steal", that produces differential increases in blood flow in healthy vs. diseased arteries in patients with coronary artery disease, optimizing the discriminatory imaging of cardiac muscle areas in need of oxygen supply.

[0003] Adenosine and dipyridamole are coronary vasodilators, each of which is approved for individual use as a pharmacologic stressor for stress testing. Adenosine acts directly by stimulating adenosine purinergic PI receptors on the arterial wall. Dipyridamole is believed to work indirectly by blocking reuptake of adenosine at the cellular level, leading to an increase in endogenous adenosine concentration in the blood. Dipyridamole produces similar near-maximal coronary hyperemia to that produced by exogenous adenosine, but less quickly.

[0004] To ensure near-maximal coronary vasodilation, and to provide sufficient time for the acquisition of cardiac images, adenosine is infused for 6 minutes at a dosage rate of 140 $\mu\text{g/kg}$ patient body weight/min; dipyridamole is infused for 4 minutes at 140 $\mu\text{g/kg}$ patient body weight/min. Thus, the total recommended dose of adenosine is 0.84 mg/kg, and the total recommended dose for dipyridamole is 0.56 mg/kg at the minimum and 0.80 mg/kg on average in a 4 minute infusion. If vasodilation is insufficient, the total dose of dipyridamole can be increased up to 0.95 mg/kg, administered over a 6 minute infusion.

[0005] Although infused for only a few minutes, compounds that stimulate adenosine receptors are accompanied by numerous uncomfortable adverse effects. With adenosine, the most frequently reported are flushing (44%), chest pain or chest discomfort (40%), dyspnea (28%), headache (18%), throat or neck or jaw discomfort (15%), and gastrointestinal discomfort (13%); other side effects are less frequent.

[0006] The adverse effects of adenosine are dose-dependent. Symptoms such as heat sensation, flushed face, dyspnea and chest pain increase as adenosine dosage is increased from 60 to 140 $\mu\text{g/kg/min}$, in a six minute infusion. Chest pain typically appears at doses of 90 $\mu\text{g/kg/min}$, and becomes frequent at 120 $\mu\text{g/kg/min}$. At a dosage of 70 $\mu\text{g/kg/min}$ or less, it has been noted that adenosine adverse reactions are very few and of mild intensity. However, when administered by intravenous perfusion at 70 $\mu\text{g/kg/min}$ or less, or even at 90 - 120 $\mu\text{g/kg/min}$, adenosine shows reduced efficacy, and is not recommended for stress testing at such reduced dosages.

[0007] The side effect profile of dipyridamole is similar, but with adverse events occurring less often. However, dipyridamole side effects last longer, are more difficult to manage, and thus more frequently require the administration of intravenous aminophylline as an antidote.

[0008] Because dipyridamole is understood to act by increasing endogenous adenosine, use of both adenosine and dipyridamole at full intravenous dosage is contraindicated. Analogously, oral intake of dipyridamole prior to an adenosine pharmacologic stress testing is generally avoided.

[0009] In an effort to reduce side effects at maximally effective agonist doses, adenosinergic agents are being developed that are selective for the A_{2a} receptor subtype. See, e.g., U.S. Pat. Nos. 6,531,457; 6,448,235; 6,322,771; and 5,877,180. Specific compounds in development include regadenoson, binodenoson and apadenoson (BMS068645). However, despite their receptor selectivity, only modest reductions in side effects have been observed with these compounds. In addition, the compounds have a longer duration of action than adenosine; accordingly, the side effects, e.g., flushing, headache, and dyspnea, are longer lasting. Thus, although more specific than adenosine, these agents may be more likely to trigger prolonged side effects, and to require administration of pharmacologic antidotes, than is adenosine itself, whose side effects rapidly dissipate once administration is stopped. Moreover, none of these selective agents has yet been approved for clinical use.

[0010] There thus exists a continuing need in the art for injectable agents that can be used for pharmacologic stress testing that have the rapid onset and short half-life of adenosine, and thus can be managed clinically in the same manner as adenosine, and that provide maximal efficacy with reduced side effects.

2. SUMMARY

[0011] Although dipyridamole is believed to act indirectly by increasing adenosine concentration, and in clinical practice has a side effect profile similar to that of adenosine, I have now discovered that extremely low parenteral doses of dipyridamole - on the order of 5% of the dose now used clinically in cardiac imaging studies - can potentiate the vasodilation effects of adjunctively administered adenosine without commensurate potentiation of adenosine's side effects. This permits adenosine to be used at reduced dosage to effect coronary vasodilation, *e.g.*, for functional assessment of myocardial function, with equal or superior efficacy as compared to current protocols, yet with reduced side effects. Moreover, the clinical and hemodynamic effects advantageously stop less than one minute after cessation of adenosine administration.

[0012] Accordingly, described herein are methods, compositions, unit dosage forms and kits that exploit this newly discovered phenomenon.

[0013] The present invention provides a pharmaceutical composition comprising adenosine and dipyridamole in an adenosine:dipyridamole weight ratio of 2:1 to 10:1, wherein the composition is formulated for parenteral administration to an individual subject and wherein adenosine and dipyridamole are present in amounts that permit adenosine to be administered at a dosage rate of 35 to 100 $\mu\text{g/kg/min}$ and dipyridamole to be administered at a dosage rate of 3.5 to 50 $\mu\text{g/kg/min}$, wherein the administration is continuous for a period of less than 6 minutes.

[0014] The present invention also provides a unit dosage form containing 2 to 50 ml of the composition according to any one of claims 1 to 5, wherein the composition is a sterile fluid.

[0015] The present invention further provides a unit dosage form containing 5 to 60 mg of adenosine and 0.5 to 30 mg of dipyridamole, wherein the composition is a solid capable of sterile reconstitution in a physiologically acceptable solvent or solution for parenteral administration to an individual subject, and wherein the adenosine:dipyridamole weight ratio is from 2:1 to 10:1.

[0016] The present invention further provides adenosine and dipyridamole for concurrent administration, for use in a method of effecting coronary vasodilation for cardiac diagnosis, wherein the adenosine and dipyridamole are concurrently administered parenterally to an individual subject at an adenosine:dipyridamole weight ratio of 2:1 to 10:1, and wherein adenosine is administered at a dosage rate of 35 to 100 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 3.5 to 50 $\mu\text{g/kg/min}$, wherein the administration is continuous for a period of less than 6 minutes.

[0017] In a first aspect of the disclosure, methods of effecting coronary vasodilation for cardiac diagnosis are provided. The method comprises concurrently administering adenosine and dipyridamole, wherein adenosine and dipyridamole are administered parenterally at an adenosine:dipyridamole weight ratio of 2: 1 to 10:1.

[0018] In the compositions disclosed herein the adenosine:dipyridamole ratio may be 2: 1 to 4: 1, such as 4: 1 ; in various other embodiments, the adenosine dipyridamole ratio is 7:1 or 8: 1.

[0019] In a method disclosed herein, adenosine is administered at a dosage rate of 35 to 100 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 3.5-50 $\mu\text{g/kg/min}$. In some embodiments, adenosine is administered at a dosage rate of 70 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 10 $\mu\text{g/kg/min}$; adenosine is administered at a dosage rate of 70 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 8.75 $\mu\text{g/kg/min}$; adenosine is administered at a dosage rate of 50 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 12.5-25 $\mu\text{g/kg/min}$.

[0020] Adenosine and dipyridamole may be parenterally administered continuously for a period of at least 2 minutes, typically less than 6 minutes. In certain embodiments, adenosine and dipyridamole are parenterally administered continuously for a period of 4 minutes.

[0021] In some embodiments, adenosine and dipyridamole are administered as a single composition. In other embodiments, adenosine and dipyridamole are administered from separate compositions.

[0022] Disclosed herein, at least one adenosine and dipyridamole may be administered by intravenous infusion. Disclosed herein, at least one adenosine and dipyridamole may be administered by intra-atrial or intra-arterial administration. The dosage of that agent can be adjusted downward using a dosage multiplier of 1/200 - 1/400, as compared to the intravenous dosage. When adenosine and/or dipyridamole is administered by intra-coronary administration, the dosage of that agent can be adjusted downward using a dosage multiplier of 1/200 - 1/400, as compared to the intravenous dosage.

[0023] Certain embodiments further comprises the step of assessing cardiac function. Assessing cardiac function may include use of one or more techniques selected from the group consisting of: electrocardiography, M mode echography, two dimensional echography, three dimensional echography, echo-doppler, cardiac imaging, planar (conventional) scintigraphy, single photon emission computed tomography (SPECT), dynamic single photon emission computed tomography, positron emission tomography (PET), first pass radionuclide angiography, equilibrium radionuclide angiography, nuclear magnetic resonance (NMR) imaging, perfusion contrast echocardiography, digital subtraction angiography (DSA), and ultrafast x-ray computed tomography (CINE CT). In certain embodiments, functional assessment is performed by SPECT; in other embodiments, the assessment is performed by PET.

[0024] Disclosed herein are methods that further comprise assessment of cardiac function, assessing cardiac function

includes parenteral administration of an isotope. The isotope is typically administered no less than 2.5 minutes after the concurrent parenteral administration of adenosine and dipyridamole has begun. The isotope can be administered no less than 2.55 minutes, no less than 2.6 minutes, no less than 2.65 minutes, no less than 2.7 minutes, no less than 2.75 minutes after the concurrent administration of adenosine and dipyridamole has begun. The isotope can be administered about 2.5 - 2.75 minutes after the concurrent parenteral administration of adenosine and dipyridamole has begun.

[0025] In a second aspect of the disclosure, methods are provided for effecting coronary vasodilation for cardiac diagnosis. These methods comprising: (i) parenterally administering dipyridamole; and (ii) concurrently or sequentially thereafter parenterally administering an adenosine receptor agonist. Each of dipyridamole and the adenosine receptor agonist is administered at a dosage lower than that required for maximal coronary vasodilation when administered as a single agent by identical parenteral route.

[0026] Disclosed herein, the adenosine receptor agonist may be selected from the group consisting of: adenosine, adenosine triphosphate (ATP), adenosine diphosphate (ADP), adenosine monophosphate (AMP), and pro-drugs and pharmaceutically acceptable salts of adenosine or AMP, ADP, ATP.

[0027] Each route of parenteral administration may be independently selected from the group consisting of: intra-arterial, intravenous, and atrial administration. Dipyridamole may be administered by intravenous or intra-arterial bolus injection. Dipyridamole may be administered as an intravenous or intra-arterial bolus at a dosage of no more than 140 $\mu\text{g/kg}$, no more than 50 $\mu\text{g/kg}$, even no more than 40 $\mu\text{g/kg}$, and typically at a dosage of at least 14 $\mu\text{g/kg}$. For example, dipyridamole may be administered as an intravenous or intra-arterial bolus at a dosage of 23 to 60 $\mu\text{g/kg}$, such as 35 $\mu\text{g/kg}$ or 40 $\mu\text{g/kg}$.

[0028] In various embodiments, dipyridamole is administered by intravenous infusion over 1 or 2 minutes.

[0029] Disclosed herein, the adenosine receptor agonist administration may be begun after completion of dipyridamole administration, such as between 30 seconds and 2 minutes after dipyridamole injection or infusion.

[0030] Disclosed herein, dipyridamole may be administered in admixture with the adenosine receptor agonist by intravenous infusion over 2 to 6 minutes, such as for 4 minutes.

[0031] The adenosine receptor agonist of the invention is adenosine, administered by intravenous infusion at a dosage rate of 35 $\mu\text{g/kg/min}$ - 100 $\mu\text{g/kg/min}$. In the invention, adenosine is administered at a dosage rate no more than 100 $\mu\text{g/kg/min}$. In some embodiments, adenosine is administered at a dosage rate of no more than about 70 $\mu\text{g/kg/min}$, even no more than about 50 $\mu\text{g/kg/min}$. The adenosine receptor agonist of the invention is adenosine, administered by intravenous infusion at a dosage rate of at least 35 $\mu\text{g/kg/min}$, even at least 50 $\mu\text{g/kg/min}$. For example, in some embodiments, adenosine is administered by intravenous infusion at a rate of about 50 $\mu\text{g/kg/min}$ to about 70 $\mu\text{g/kg/min}$, such as about 70 $\mu\text{g/kg/min}$.

[0032] In some embodiments, dipyridamole is administered intravenously and adenosine is administered intravenously.

[0033] In certain exemplary embodiments, the adenosine receptor agonist is adenosine, the total dose of dipyridamole is 23 to 40 $\mu\text{g/kg}$, and the dosage rate for adenosine is 50 to 70 $\mu\text{g/kg/min}$. For example, in some embodiments, the total dose of dipyridamole is 40 $\mu\text{g/kg}$ and the dosage rate for adenosine is 70 $\mu\text{g/kg/min}$.

[0034] The method disclosed herein may further comprise the step of: assessing cardiac function. Assessing cardiac function may include parenteral administration of an isotope, and the isotope is administered after 2 minutes, when dipyridamole and the adenosine receptor agonist are administered sequentially, and after 2.5 minutes, after 2.55 minutes, after 2.6 minutes, after 2.65 minutes, after 2.75 minutes, after 2.8 minutes, after 2.85 minutes, after 2.9 minutes or after 2.95 minutes but before 3 minutes, when dipyridamole and the adenosine receptor agonist are administered concurrently. In some embodiments, isotope is administered at about 2.5 - 2.75 minutes after sequential administration of dipyridamole and the adenosine receptor agonist.

[0035] In another aspect, pharmaceutical compositions comprising adenosine and dipyridamole are presented. The compositions comprise adenosine and dipyridamole in adenosine:dipyridamole weight ratios of 2: 1 to 10: 1, such as 2: 1 to 4:1. In some embodiments, the ratio is about 7:1 or 8: 1.

[0036] Adenosine and dipyridamole of the invention are present in amounts that permit adenosine to be administered at a dosage rate of 35 to 100 $\mu\text{g/kg/min}$ and dipyridamole to be administered at a dosage rate of 3.5 to 50 $\mu\text{g/kg/min}$.

[0037] The composition may be a sterile fluid, such as a sterile fluid suitable for parenteral administration, such as intravenous administration. In some embodiments, adenosine and dipyridamole are present at concentrations that permit direct intravenous administration, without dilution.

[0038] In various embodiments, adenosine and dipyridamole are present at concentrations that permit administration of adenosine at a dosage rate of 70 $\mu\text{g/kg/min}$ and dipyridamole at a dosage rate of 8.75 to 10 $\mu\text{g/kg/min}$. In some embodiments, adenosine and dipyridamole are present at concentrations that permit administration of adenosine at a dosage rate of 50 $\mu\text{g/kg/min}$ and dipyridamole at a dosage rate of 12.5 to 25 $\mu\text{g/kg/min}$.

[0039] In a range of embodiments of the pharmaceutical compositions here provided, the concentration of adenosine is 1 to 10 mg/ml. Usefully, the concentration of adenosine is 3 mg/ml or 4 mg/ml, even 5 mg/ml, or 7 mg/ml.

[0040] In certain embodiments, the concentration of dipyridamole is about 0.1 to 5 mg/ml, such as: 0.375 to 0.428 mg/ml; 0.5 to 0.571 mg/ml; 0.625 to 0.714 mg/ml; 0.75 to 0.857 mg/ml; and 0.875 to 1 mg/ml. The concentration may,

for example, be 1 mg/ml.

[0041] In another aspect, unit dosage forms are provided that contain pharmaceutical compositions as above-described, comprising adenosine and dipyridamole.

[0042] In some embodiments, the unit dose contains 2 to 50 ml of the pharmaceutical composition formulated as a sterile fluid, typically a sterile, nonpyrogenic, solution suitable for parenteral administration. In some embodiments, the unit dose contains 2 ml, 3 ml, 4 ml, 7 ml, 8 ml or 14 ml.

[0043] In some embodiments, the unit dose contains 5 to 60 mg of adenosine and 0.5 to 30 mg of dipyridamole; the composition is a solid capable of sterile reconstitution in a physiologically acceptable solvent or solution.

[0044] In exemplary embodiments, for example, the unit dosage contains 14 mg of adenosine and 2 mg of dipyridamole, 21 mg of adenosine and 3 mg of dipyridamole; 28 mg of adenosine and 4 mg of dipyridamole; 35 mg of adenosine and 5 mg of dipyridamole; 42 mg of adenosine and 6 mg of dipyridamole; 56 mg of adenosine and 8 mg of dipyridamole; 20 mg of adenosine and 5 to 10 mg of dipyridamole; 30 mg of adenosine and 7.5 to 15 mg of dipyridamole; 40 mg of adenosine and 10 to 20 mg of dipyridamole.

[0045] In a further aspect of the disclosure, unit doses of dipyridamole are provided. In various embodiments, dipyridamole is provided in solution at a concentration of about 0.1 to 5 mg/ml.

[0046] The dipyridamole concentration may be usefully about 0.5 mg/ml. Among these embodiments of the disclosure are unit dosage forms that contain 3 mg dipyridamole in 6 ml; 4 mg dipyridamole in 8 ml; 5 mg dipyridamole in 10 ml; 6 mg dipyridamole in 12 ml; 8 mg dipyridamole in 16 ml. In some embodiments, the unit dose contains dipyridamole at a concentration between about 3 mg/ml and about 5 mg/ml, such as 3 mg/ml or 4 mg/ml, usefully in volumes of 1 ml or of 2 ml, providing unit dosage forms containing 6 mg dipyridamole in 2 ml and 8 mg dipyridamole in 2 ml.

[0047] In a further aspect of the disclosure, unit doses of adenosine are provided. The unit doses are formulated in sterile fluid composition, and the dose packaging permits sterile introduction of a second fluid in a volume at least 15% that of the adenosine composition. The second fluid usefully comprises dipyridamole.

[0048] The unit dose may contain 21 mg adenosine in 6 ml; 28 mg adenosine in 6 ml; 42 mg adenosine in 12 ml; or 56 mg adenosine in 12 ml.

[0049] The unit dose may comprise adenosine at a concentration of about 4 mg/ml. The unit dose may contain 28 mg adenosine in 7 ml; 56 mg adenosine in 14 ml.

[0050] Also disclosed herein are kits. The kits comprise at least one unit dose of dipyridamole and at least one unit dose of adenosine. The at least one unit dose of dipyridamole may be a unit dose as above-described, and the unit dose of adenosine is a unit dose as above-described.

3. DETAILED DESCRIPTION

3.1 Overview

[0051] Although dipyridamole is believed to act indirectly by increasing adenosine concentration, and in clinical practice has a side effect profile similar to that of adenosine, I have now discovered that extremely low parenteral-subclinical doses of dipyridamole - on the order of 5% of the dose now used clinically in cardiac imaging studies - can potentiate the vasodilation effects of adjunctively administered adenosine without commensurate potentiation of adenosine's side effects. This permits adenosine to be used at reduced dosage to effect coronary vasodilation, e.g., for functional assessment of myocardial function, with equal or superior efficacy as compared to current protocols, yet with reduced side effects, of short duration.

[0052] In the first of two clinical studies reported in detail below (Example 1), the hemodynamic effects of administering dipyridamole and adenosine intravenously as a combined (albeit, sequentially administered) pharmacological stressor were compared to the effects of administering adenosine alone in 40 consecutive patients suffering from ischemic heart disease. Each patient served as his own control. Dipyridamole was administered as an intravenous bolus. Adenosine was administered immediately thereafter by continuous intravenous infusion for three minutes. Each of the two agents was administered at a dosage lower than its clinically preferred dosage when used as a single agent for myocardial perfusion imaging: dipyridamole at 4 - 6% of its single-agent total dose, adenosine at one half its single-agent dosage rate.

[0053] Effects were measured using noninvasive transthoracic doppler echocardiography (TTDE). The measured blood flow velocities (known as reflecting coronary blood flow values), whether peak or mean, were 1.5 to 4% lower in absolute values than those measured upon administration of adenosine alone at its standard dosage rate. However, these differences were not statistically significant ($p > 0.05$): there was no statistical difference between the current standard treatment - infusion of adenosine alone at 140 $\mu\text{g/kg/min}$ - and sequential bolus administration of dipyridamole at 4 - 6% of its typical single-agent total dose followed by adenosine infusion at 70 $\mu\text{g/kg/min}$.

[0054] In addition, among the first series of 30 patients, three (3) patients received the adenosine infusion two minutes after the dipyridamole bolus, rather than immediately thereafter, and two (2) patients were injected with the two agents concurrently in the same infusion line using a "Y" connector. No differences were seen as compared to the sequential

administration protocol.

[0055] A significant reduction was seen in the incidence of chest pain among the 40 patients in this study, as compared to the number reporting chest pain upon administration of adenosine alone at 140 $\mu\text{g/kg/min}$. In addition, the severity of the three main adverse side effects - chest pain, dyspnea, and flushing - cumulated across all dipyridamole doses, was reduced by 31.6% with the sequential combination as compared to standard adenosine treatment. This decrease was statistically significant ($p = 0.001$).

[0056] As reported in detail in Example 2, below, 27 patients have now been assessed in a subsequent Phase II study comparing dipyridamole-adenosine combination administration to adenosine alone (Adenoscan®, Astellas) as the pharmacologic stressor in coronary patients undergoing single photon emission computed tomography (SPECT) imaging studies.

[0057] Data from initial patients who participated in a preliminary dose-finding study demonstrated that either bolus intravenous administration of dipyridamole at 35 $\mu\text{g/kg}$ over 20 - 30 seconds, followed by intravenous infusion of adenosine at 70 $\mu\text{g/kg/min}$ (tested in 3 patients) or the concurrent administration of the two drugs at the same dosage (tested in 5 patients), provided images comparable to Adenoscan (adenosine at 140 $\mu\text{g/kg/min}$) in 7 consecutive patients, while under-scoring Adenoscan in one patient, but within the acceptable limits defined by the protocol. The combination stressor of 40 $\mu\text{g/kg}$ dipyridamole followed by 70 $\mu\text{g/kg/min}$ adenosine tested in 10 patients or their concurrent administration at the same dosage (adenosine 70 $\mu\text{g/kg/min}$ with dipyridamole 10 $\mu\text{g/kg/min}$) tested in 9 patients showed equivalent, and sometimes better results, than Adenoscan in terms of imaging efficacy in 10 and 9 consecutive patients respectively.

[0058] Significant reduction in both the occurrence and the severity of chest pain with the dipyridamole-adenosine combination, as compared to adenosine alone, was observed, as was reduction in ST changes on EKG.

[0059] The data from these two studies demonstrate that the sequential bolus administration of dipyridamole at 28 to 40 $\mu\text{g/kg}$ - well below the total dose infused when dipyridamole is used as a single agent stressor - followed by infusion of adenosine at 70 $\mu\text{g/kg/min}$ (50% less than its usual dosage), is equally efficacious in providing coronary vasodilation for imaging studies, while causing fewer side effects. The data also demonstrate that dipyridamole and adenosine may be combined in a single infusion, over 4 minutes, to similar effect. Among the side effects reduced by the combination of the present disclosure are chest pain, and the risk of significant heart blockage.

[0060] Accordingly, described herein are methods, pharmaceutical compositions, unit dosage forms, and kits that exploit this discovery, combining adenosine with dipyridamole at dosages at which some of the most frequent side effects of both adenosine and dipyridamole, notably cardiac side effects, are significantly reduced, while maintaining optimal coronary vasodilation for the diagnosis of myocardial ischemia.

3.2 Methods of effecting coronary vasodilation

[0061] Methods of effecting coronary vasodilation for cardiac diagnosis are disclosed herein.

[0062] The methods may comprise parenterally administering dipyridamole and concurrently or sequentially thereafter parenterally administering an adenosine receptor agonist. Each of dipyridamole and the adenosine receptor agonist is administered at a dosage lower than that required for maximal coronary vasodilation when the respective agent is administered individually by identical parenteral route. Dipyridamole and the adenosine receptor agonist are administered in amounts, at weight ratios, and for a time, sufficient to achieve the desired therapeutic or diagnostic effect.

[0063] Programmable syringe pumps or micropumps, as are typical in clinical practice, are usefully employed to facilitate parenteral administration in precise dosage.

[0064] The route of parenteral administration is chosen based upon the desired clinical effect, as further described below. Disclosed herein at least one of dipyridamole and the adenosine receptor agonist may be administered by intravenous infusion. In other embodiments of the disclosure, at least one of dipyridamole and the adenosine receptor agonist is administered by intra-arterial infusion, such as intra-coronary infusion, or by intra-atrial infusion. In these latter embodiments of the disclosure, the active is administered at a lower rate, and at a lower dosage, than for intravenous infusion, as further described below. In yet other embodiments of the disclosure, at least one of the actives is administered as a perfusate.

[0065] At least one of dipyridamole and the adenosine receptor agonist may be infused over a period of time of at least 1 minute, typically at least 2 minutes, 3 minutes, 4 minutes, 5 minutes, even at least 6 minutes. As used herein, "continuous infusion" intends infusion over a period of at least 2 minutes.

[0066] Dipyridamole may be administered by intravenous infusion at an infusion rate from 3.5 $\mu\text{g/kg/min}$ to 50 $\mu\text{g/kg/min}$. All dosage ranges described herein include the upper and lower recited limits, and nonintegral intermediary values. Thus, in some embodiments of the disclosure, dipyridamole is infused at a rate of at least 3.5 $\mu\text{g/kg/min}$, at least 4 $\mu\text{g/kg/min}$, at least 5 $\mu\text{g/kg/min}$, at least 6 $\mu\text{g/kg/min}$, at least 7 $\mu\text{g/kg/min}$, at least 7.5 $\mu\text{g/kg/min}$, at least 8 $\mu\text{g/kg/min}$, at least 8.75 $\mu\text{g/kg/min}$, at least 9 $\mu\text{g/kg/min}$, at least 10 $\mu\text{g/kg/min}$, at about 11 $\mu\text{g/kg/min}$, at least 11.25 $\mu\text{g/kg/min}$, at least 12 $\mu\text{g/kg/min}$, at least 12.5 $\mu\text{g/kg/min}$, at least 13 $\mu\text{g/kg/min}$, at least 13.75 $\mu\text{g/kg/min}$, at least 14 $\mu\text{g/kg/min}$, at

least 15 $\mu\text{g/kg/min}$, at least 16 $\mu\text{g/kg/min}$, at least 16.25 $\mu\text{g/kg/min}$, at least 17 $\mu\text{g/kg/min}$, and at least 17.5 $\mu\text{g/kg/min}$, at least 18 $\mu\text{g/kg/min}$, at least 19 $\mu\text{g/kg/min}$, at least 20 $\mu\text{g/kg/min}$, at least 21 $\mu\text{g/kg/min}$, at least 22 $\mu\text{g/kg/min}$, at least 23 $\mu\text{g/kg/min}$, at least 24 $\mu\text{g/kg/min}$, at least 25 $\mu\text{g/kg/min}$, at least 26 $\mu\text{g/kg/min}$, at least 27 $\mu\text{g/kg/min}$, at least 28 $\mu\text{g/kg/min}$, at least 29 $\mu\text{g/kg/min}$, at least 30 $\mu\text{g/kg/min}$, at least 31 $\mu\text{g/kg/min}$, at least 32 $\mu\text{g/kg/min}$, at least 33 $\mu\text{g/kg/min}$, at least 34 $\mu\text{g/kg/min}$, at least 35 $\mu\text{g/kg/min}$, at least 36 $\mu\text{g/kg/min}$, at least 37 $\mu\text{g/kg/min}$, at least 38 $\mu\text{g/kg/min}$, at least 39 $\mu\text{g/kg/min}$, at least 40 $\mu\text{g/kg/min}$, at least 41 $\mu\text{g/kg/min}$, at least 42 $\mu\text{g/kg/min}$, at least 43 $\mu\text{g/kg/min}$, at least 44 $\mu\text{g/kg/min}$, at least 45 $\mu\text{g/kg/min}$, at least 46 $\mu\text{g/kg/min}$, at least 47 $\mu\text{g/kg/min}$, at least 48 $\mu\text{g/kg/min}$, at least 49 $\mu\text{g/kg/min}$, at least 50 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0067] In some embodiments of the disclosure, dipyridamole is infused intravenously at a rate of no more than 50 $\mu\text{g/kg/min}$, no more than 49 $\mu\text{g/kg/min}$, no more than 48 $\mu\text{g/kg/min}$, no more than 47 $\mu\text{g/kg/min}$, no more than 46 $\mu\text{g/kg/min}$, no more than 45 $\mu\text{g/kg/min}$, no more than 44 $\mu\text{g/kg/min}$, no more than 43 $\mu\text{g/kg/min}$, no more than 42 $\mu\text{g/kg/min}$, no more than 41 $\mu\text{g/kg/min}$, no more than 40 $\mu\text{g/kg/min}$, no more than 39 $\mu\text{g/kg/min}$, no more than 38 $\mu\text{g/kg/min}$, no more than 37 $\mu\text{g/kg/min}$, no more than 36 $\mu\text{g/kg/min}$, no more than 35 $\mu\text{g/kg/min}$, no more than 34 $\mu\text{g/kg/min}$, no more than 33 $\mu\text{g/kg/min}$, no more than 32 $\mu\text{g/kg/min}$, no more than 31 $\mu\text{g/kg/min}$, no more than 30 $\mu\text{g/kg/min}$, of no more than 29 $\mu\text{g/kg/min}$, no more than 28 $\mu\text{g/kg/min}$, no more than 27 $\mu\text{g/kg/min}$, no more than 26 $\mu\text{g/kg/min}$, no more than 25 $\mu\text{g/kg/min}$, no more than 24 $\mu\text{g/kg/min}$, no more than 23 $\mu\text{g/kg/min}$, no more than 22 $\mu\text{g/kg/min}$, no more than 21 $\mu\text{g/kg/min}$, no more than 20 $\mu\text{g/kg/min}$, no more than 19 $\mu\text{g/kg/min}$, no more than 18 $\mu\text{g/kg/min}$, no more than 17.5 $\mu\text{g/kg/min}$, no more than 17 $\mu\text{g/kg/min}$, no more than 16.25 $\mu\text{g/kg/min}$, no more than 16 $\mu\text{g/kg/min}$, no more than 15 $\mu\text{g/kg/min}$, no more than 14 $\mu\text{g/kg/min}$, no more than 13.75 $\mu\text{g/kg/min}$, no more than 13 $\mu\text{g/kg/min}$, no more than 12.5 $\mu\text{g/kg/min}$, no more than 12 $\mu\text{g/kg/min}$, no more than 11.25 $\mu\text{g/kg/min}$, no more than 11 $\mu\text{g/kg/min}$, no more than 10 $\mu\text{g/kg/min}$, no more than 9 $\mu\text{g/kg/min}$, no more than 8.75 $\mu\text{g/kg/min}$, no more than 8 $\mu\text{g/kg/min}$, no more than 7.5 $\mu\text{g/kg/min}$, no more than 7 $\mu\text{g/kg/min}$, no more than 6 $\mu\text{g/kg/min}$, no more than 5 $\mu\text{g/kg/min}$, no more than 4 $\mu\text{g/kg/min}$, no more than 3.5 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0068] In some embodiments of the disclosure, dipyridamole is administered as a bolus, typically over a period of about 20 - 30 seconds.

[0069] In some of these embodiments of the disclosure, dipyridamole is administered as an intravenous bolus. In such embodiments of the disclosure, dipyridamole is administered at a dosage between 14 $\mu\text{g/kg}$ to 140 $\mu\text{g/kg}$. In various embodiments of the disclosure, dipyridamole is administered intravenously as a bolus at a dosage between 28 $\mu\text{g/kg}$ and 40 $\mu\text{g/kg}$.

[0070] Thus, in certain embodiments of the disclosure, dipyridamole is administered as an intravenous bolus at a dose of at least 14 $\mu\text{g/kg}$, at least 20 $\mu\text{g/kg}$, at least 25 $\mu\text{g/kg}$, at least 28 $\mu\text{g/kg}$, at least 29 $\mu\text{g/kg}$, at least 30 $\mu\text{g/kg}$, at least 31 $\mu\text{g/kg}$, at least 32 $\mu\text{g/kg}$, at least 33 $\mu\text{g/kg}$, at least 34 $\mu\text{g/kg}$, at least 35 $\mu\text{g/kg}$, at least 36 $\mu\text{g/kg}$, at least 37 $\mu\text{g/kg}$, at least 38 $\mu\text{g/kg}$, at least 39 $\mu\text{g/kg}$, at least 40 $\mu\text{g/kg}$, at least 45 $\mu\text{g/kg}$, at least 50 $\mu\text{g/kg}$, at least 55 $\mu\text{g/kg}$, at least 60 $\mu\text{g/kg}$, at least 65 $\mu\text{g/kg}$, even at least 70, 80, 90, 100, 110, 120, 130, even 140 $\mu\text{g/kg}$, with intermediate doses permissible.

[0071] In some embodiments of the disclosure, dipyridamole is administered intravenously as a bolus at a dosage of no more than 140 $\mu\text{g/kg}$, 130 $\mu\text{g/kg}$, 120 $\mu\text{g/kg}$, 110 $\mu\text{g/kg}$, 100 $\mu\text{g/kg}$, 90 $\mu\text{g/kg}$, 80 $\mu\text{g/kg}$, 70 $\mu\text{g/kg}$, even no more than 60 $\mu\text{g/kg}$, even no more than 55 $\mu\text{g/kg}$, no more than 50 $\mu\text{g/kg}$, no more than 45 $\mu\text{g/kg}$, no more than 40 $\mu\text{g/kg}$, no more than 39 $\mu\text{g/kg}$, no more than 38 $\mu\text{g/kg}$, no more than 37 $\mu\text{g/kg}$, no more than 36 $\mu\text{g/kg}$, no more than 35 $\mu\text{g/kg}$, no more than 34 $\mu\text{g/kg}$, no more than 33 $\mu\text{g/kg}$, no more than 32 $\mu\text{g/kg}$, no more than 31 $\mu\text{g/kg}$, no more than 30 $\mu\text{g/kg}$, no more than 29 $\mu\text{g/kg}$, no more than 28 $\mu\text{g/kg}$, no more than 25 $\mu\text{g/kg}$, no more than 20 $\mu\text{g/kg}$, no more than 14 $\mu\text{g/kg}$, with intermediate values permissible.

[0072] When administered to a human being, the dosages of dipyridamole useful in the methods of the present disclosure can be expressed in μg by multiplying the dosage, expressed as $\mu\text{g/kg}$, by the weight of the individual. For example, for a human weighing 50 kg, the dosage of dipyridamole useful in the present methods can be expressed as ranging between 700 to 7,000 μg ; for a human being weighing 60 kg, the dosage of dipyridamole can be expressed as ranging between 840 to 8,400 μg ; for a human being weighing 75 kg, the dosage of dipyridamole can be expressed as ranging between 1,050 to 10,500 μg ; and for a human being weighing 100 kg, the dosage can be expressed as ranging between 1,400 to 14,000 μg .

[0073] In various embodiments of the disclosure, dipyridamole is infused intra-arterially at an infusion rate of no more than 0.07 $\mu\text{g/kg/min}$, no more than 0.06 $\mu\text{g/kg/min}$, no more than 0.05 $\mu\text{g/kg/min}$, no more than 0.04 $\mu\text{g/kg/min}$, no more than 0.03 $\mu\text{g/kg/min}$, no more than 0.02 $\mu\text{g/kg/min}$, or no more than 0.01 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0074] In various embodiments of the disclosure, the adenosine receptor agonist is selected from the group consisting of adenosine, and adenosine donors (that is, compounds that can be metabolized to adenosine), including natural donors such as adenosine triphosphate (ATP), adenosine diphosphate (ADP), and adenosine monophosphate (AMP), each at approximately the same dosages as adenosine, and any synthetic molecule that is capable of being metabolized to adenosine, and pharmaceutically acceptable salts thereof.

[0075] Typically, adenosine is used. For convenience, its particular use will hereafter be described, without intending thereby to limit the described methods to use of adenosine as the adenosine receptor agonist.

[0076] In some embodiments of the disclosure, adenosine is administered by intravenous infusion at an infusion rate between 35 $\mu\text{g/kg/min}$ to 100 $\mu\text{g/kg/min}$. Thus, in some embodiments of the disclosure, adenosine is infused at a rate of at least 35 $\mu\text{g/kg/min}$, at least 40 $\mu\text{g/kg/min}$, at least 45 $\mu\text{g/kg/min}$, at least 50 $\mu\text{g/kg/min}$, at least 55 $\mu\text{g/kg/min}$, at least 60 $\mu\text{g/kg/min}$, at least 65 $\mu\text{g/kg/min}$, at least 70 $\mu\text{g/kg/min}$, at least 75 $\mu\text{g/kg/min}$, at least 80 $\mu\text{g/kg/min}$, at least 85 $\mu\text{g/kg/min}$, at least 90 $\mu\text{g/kg/min}$, at least 95 $\mu\text{g/kg/min}$, and at least 100 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0077] In various embodiments of the disclosure, adenosine is infused intravenously at a rate of no more than 100 $\mu\text{g/kg/min}$, no more than 95 $\mu\text{g/kg/min}$, no more than 90 $\mu\text{g/kg/min}$, no more than 85 $\mu\text{g/kg/min}$, no more than 80 $\mu\text{g/kg/min}$, no more than 75 $\mu\text{g/kg/min}$, no more than 70 $\mu\text{g/kg/min}$, no more than 65 $\mu\text{g/kg/min}$, no more than 60 $\mu\text{g/kg/min}$, no more than 55 $\mu\text{g/kg/min}$, no more than 50 $\mu\text{g/kg/min}$, no more than 45 $\mu\text{g/kg/min}$, no more than 40 $\mu\text{g/kg/min}$, no more than 35 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0078] When administered to a human being, the dosage rate of adenosine can be expressed in $\mu\text{g/min}$ by multiplying the dosage rate expressed in $\mu\text{g/kg/min}$ by the weight of the individual. For example, for a human being weighing 50 kg, the dosage of adenosine useful in the practice of the present methods can be expressed as ranging between 1,750 to 5,000 $\mu\text{g/min}$; for a human being weighing 60 kg, the dosage rate of adenosine can be expressed as ranging between 2,100 to 6,000 $\mu\text{g/min}$; for a human being weighing 75 kg, the dosage of adenosine can be expressed as ranging between 2,625 to 7,500 $\mu\text{g/min}$; and for a human being weighing 100 kg, the dosage of adenosine can be expressed as ranging between 3,500 to 10,000 $\mu\text{g/min}$.

[0079] In some embodiments of the disclosure, adenosine is administered by intra-arterial infusion, such as intracoronary infusion, at an infusion rate about 200- to 400-fold lower than intravenous infusion. Thus, in some embodiments of the disclosure, adenosine is infused at a rate of at least 0.50 $\mu\text{g/kg/min}$, at least 0.45 $\mu\text{g/kg/min}$, at least 0.40 $\mu\text{g/kg/min}$, 0.35 $\mu\text{g/kg/min}$, at least 0.30 $\mu\text{g/kg/min}$, at least 0.25 $\mu\text{g/kg/min}$ at least 0.20 $\mu\text{g/kg/min}$, at least 0.15 $\mu\text{g/kg/min}$, at least 0.10 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0080] In various embodiments of the disclosure, adenosine is infused intra-arterially and in particular intracoronarily at an infusion rate of no more than 0.10 $\mu\text{g/kg/min}$, no more than 0.15 $\mu\text{g/kg/min}$, no more than 0.20 $\mu\text{g/kg/min}$, no more than 0.25 $\mu\text{g/kg/min}$, no more than 0.30 $\mu\text{g/kg/min}$, no more than 0.35 $\mu\text{g/kg/min}$, no more than 0.40 $\mu\text{g/kg/min}$, no more than 0.45 $\mu\text{g/kg/min}$, even no more than 0.50 $\mu\text{g/kg/min}$, with intermediate values permissible.

[0081] In various embodiments of the disclosure, the methods presented herein comprise parenterally administering dipyridamole; and concurrently or sequentially thereafter parenterally administering an adenosine receptor agonist, such as adenosine, at an adenosine:dipyridamole (A:D) weight ratio of about 2:1 to about 10:1. In various embodiments of the disclosure, the methods comprise concurrently administering adenosine and dipyridamole at an adenosine: dipyridamole ratio of about 2: 1 to about 10: 1.

[0082] In some embodiments of the disclosure, the ratio is 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, even 10:1, with nonintegral ratios between 2:1 and 10:1 permissible. In certain embodiments of the disclosure, the methods comprise concurrent infusion of adenosine and dipyridamole at an A:D ratio of 6:1 to 8:1, preferably 7:1. For certain methods further described below, embodiments usefully comprise concurrent parenteral infusion of adenosine and dipyridamole at an A:D weight ratio of 2:1 to 4:1.

[0083] In some embodiments of the disclosure, dipyridamole is administered by intra-coronary infusion: in such embodiments, dipyridamole is typically administered at an infusion rate which is 1/7 that of adenosine, and between 0.01 to 0.07 $\mu\text{g/kg/min}$. Thus, in some embodiments of the disclosure, dipyridamole is infused intra-arterially at a rate of at least 0.01 $\mu\text{g/kg/min}$, at least 0.02 $\mu\text{g/kg/min}$ at least 0.03 $\mu\text{g/kg/min}$ at least 0.04 $\mu\text{g/kg/min}$ at least 0.05 $\mu\text{g/kg/min}$ at least 0.06 $\mu\text{g/kg/min}$ at least 0.07 $\mu\text{g/kg/min}$.

[0084] In various embodiments of the disclosure, dipyridamole is administered as an intravenous bolus, and adenosine is administered thereafter as an intravenous infusion.

[0085] In certain such embodiments of the disclosure, dipyridamole is administered over 20 - 30 seconds, and adenosine is thereafter infused for 2 to 6 minutes.

[0086] In these embodiments of the disclosure, the total dose of dipyridamole given intravenously as a bolus is typically between 1/16 to 1/24, e.g., 1/20 (5%), that of the total recommended standard dose when dipyridamole is used as a single agent (standard single agent dose: 0.56 mg - 0.80 mg/kg). In these embodiments of the disclosure, the total dose of intravenously infused adenosine is typically 25% to 50% that of the total recommended standard dose when adenosine is used as a single agent (standard single agent dose: 0.84 mg/kg).

[0087] In typical embodiments of the disclosure, dipyridamole is administered as an IV bolus over 20 - 30 seconds at a dosage of 14 to 60 $\mu\text{g/kg}$, followed immediately (that is, as soon as clinically practicable, typically within about 5 to 30 seconds) by the infusion of adenosine at a dosage of 35 to 100 $\mu\text{g/kg/min}$ for a period of 3 to 6 minutes. The duration of adenosine administration is determined by the chosen imaging methodology, as is well known in the art.

[0088] In some embodiments of the disclosure, an intravenous dipyridamole bolus of 28 - 40 $\mu\text{g/kg}$ is followed immediately - that is, as soon as clinically practicable, typically within about 5 to 30 seconds - by intravenous infusion of adenosine at 50 - 70 $\mu\text{g/kg/min}$ for 2 to 6 minutes. In certain embodiments of the disclosure, an IV dipyridamole bolus

of 40 $\mu\text{g/kg}$ is followed immediately by intravenous administration of adenosine at 70 $\mu\text{g/kg/min}$ for 4 minutes.

[0089] In sequential administration embodiments of the disclosure, adenosine infusion may be delayed as long as 2 - 10 minutes after dipyridamole bolus, typically no more than 5 minutes after dipyridamole bolus.

[0090] In sequential administration embodiments of the disclosure, dipyridamole may be injected manually as a bolus via a syringe, although programmable administration (e.g., by micropump) is also possible. When administered by micropump, dipyridamole may be injected over 1 or 2 minutes prior to adenosine infusion. Adenosine infusion is typically accomplished using a programmable device so as to ensure its measured delivery.

[0091] In some embodiments of the disclosure, dipyridamole and adenosine are administered concurrently.

[0092] In certain concurrent administration embodiments of the disclosure, dipyridamole and adenosine are in separate unit dosage forms, and are mixed prior to administration and infused together in a single composition.

[0093] For example, in some embodiments of the disclosure, a volume of dipyridamole corresponding to a dosage of 14 to 60 $\mu\text{g/kg}$ is sampled and a volume of adenosine corresponding to a dosage of 35 to 100 $\mu\text{g/kg/min}$ is similarly sampled and the two mixed in the same syringe. The mixture is then infused over 3 to 6 minutes. The duration of intravenous administration is determined by the chosen imaging methodology, as is well known in the art.

[0094] In some embodiments, a volume of dipyridamole corresponding to 28 - 40 $\mu\text{g/kg}$ is sampled and a volume of adenosine corresponding to a dosage of 50 to 70 $\mu\text{g/kg/min}$ is similarly sampled, and the two mixed in the same syringe. The mixture is then infused over 3 to 6 minutes. In certain embodiments, a volume of dipyridamole corresponding to 40 $\mu\text{g/kg}$ is sampled and a volume of adenosine corresponding to a dosage of 70 $\mu\text{g/kg/min}$ is similarly sampled and the two mixed in the same syringe. The mixture is then infused over 4 minutes. Thus, as is described further below, in another aspect, specific unit dosage forms of adenosine are provided, usefully copackaged with specific unit dosage forms of dipyridamole, so as to facilitate the sequential sampling and mixture of both actives in the same syringe.

[0095] In other embodiments of the disclosure, the total volume of the dipyridamole unit dosage form is injected into the adenosine vial and the two are mixed. Thus, as described below, in another aspect the disclosure provides unit dosage forms of adenosine packaged so as to permit the sterile introduction of an appropriate volume of dipyridamole.

[0096] In embodiments in which the two actives are in admixture in a single composition prior to administration, the volume to administer is usefully calculated using adenosine dose tables as reference.

[0097] In other embodiments, dipyridamole and adenosine are concurrently administered from separate compositions. Usefully, the two agents may be introduced into the same infusion line using a Y connector (at the same dosages as set forth above).

3.3 Methods of pharmacological stress testing

[0098] Vasodilation that is achieved according to the above-described methods will often be used as a pharmacological stressor in cardiac stress tests. Accordingly, the methods disclosed herein further comprise the step of assessing cardiac function.

[0099] Any method suitable for assessing cardiac function in cardiac stress testing may be used.

[0100] In various embodiments, for example, assessing cardiac function includes use of one or more techniques selected from the group consisting of: electrocardiography, echography (M mode, two-dimensional, and three dimensional), echo-doppler, cardiac imaging, including planar (conventional) scintigraphy, single photon emission computed tomography (SPECT), dynamic single photon emission computed tomography (D-SPECT™ Cardiac Scan), positron emission tomography (PET), radionuclide angiography (first pass and equilibrium studies utilizing, e.g., technetium-99m-labeled red blood cells), nuclear magnetic resonance (NMR) imaging, perfusion contrast echocardiography, digital subtraction angiography (DSA), and ultrafast x-ray computed tomography (CINE CT).

SPECT and PET present certain advantages.

[0101] SPECT studies can be performed using any of the isotopes known to be suitable for such studies, such as thallium-201, technetium sestamibi, tetrofosmine. PET studies can be performed using any of the isotopes known to be suitable for such studies, such as for example rubidium-82, nitrogen-13, fluorine-18, carbon-11, boron-11, and oxygen-15.

[0102] Typically, isotope is injected during the infusion of adenosine, and imaging begins after the end of the infusion. In some embodiments, the isotope is administered no less than about 2.5 minutes after adenosine infusion has begun.

3.4 Pharmaceutical compositions

[0103] In another aspect, pharmaceutical compositions that are useful in the above-described methods are provided.

[0104] In typical embodiments, the pharmaceutical composition comprises adenosine and dipyridamole in an adenosine:dipyridamole (A:D) weight ratio of about 2:1 to about 10:1, with intermediate (including nonintegral) values permissible. In certain embodiments in which adenosine is intended to be administered at 70 $\mu\text{g/kg/min}$, the ratio is usefully

about 7:1, 8:1, 9:1 and 10:1, with intermediate and nonintegral ratios permissible. In other embodiments, in which adenosine is to be administered at 50 µg/kg/min or less, A:D ratios are usefully about 2:1, 3:1, and 4:1, with intermediate and nonintegral ratios between 2:1 and 4:1 permissible. For certain clinical methods, the composition usefully comprises adenosine and inosine at an A:D weight ratio of about 7:1.

[0105] In certain embodiments, the pharmaceutical composition is suitable for intravenous, intra-atrial, or intra-arterial infusion.

[0106] The composition may, for example, be in the form of a sterile, nonpyrogenic, fluid composition.

[0107] In typical fluid embodiments, the concentration of adenosine is at least about 1 mg/ml, 2 mg/ml, 3 mg/ml, 4 mg/ml and possibly 5 mg/ml, with intermediate, nonintegral, values permissible. These embodiments typically have a pH of about 3.5 to about 8. In other typical fluid embodiments with readily a lower pH (e.g., pH 2 -3.5), adenosine concentration can be higher, even at least about 5 mg/ml, 6 mg/ml, 7 mg/ml, 8 mg/ml, 9 mg/ml and even 10 mg/ml, with intermediate, nonintegral, values permissible. In typical pharmaceutical composition embodiments, adenosine is present at a concentration of about 3 mg/ml, 4 mg/ml, 5 mg/ml, or 7 mg/ml.

[0108] In various fluid embodiments, the concentration of dipyridamole is at least about 0.1 mg/ml, and may usefully be as high as 2.5 mg/ml, and even 5 mg/ml. The concentration may, in certain embodiments, be at least about 0.1 mg/ml, 0.2 mg/ml, 0.3 mg/ml, 0.4 mg/ml, 0.5 mg/ml, 0.6 mg/ml, 0.7 mg/ml, 0.8 mg/ml, 0.9 mg/ml, 1 mg/ml, or more, including, e.g., 1.1 mg/ml, 1.2 mg/ml, 1.3 mg/ml, 1.4 mg/ml, 1.5 mg/ml, 1.6 mg/ml, 1.7 mg/ml, 1.8 mg/ml, 1.9 mg/ml, 2 mg/ml, 2.1 mg/ml, 2.2 mg/ml, 2.3 mg/ml, or 2.4 mg/ml, 2.5 mg/ml, 3 mg/ml, 3.5 mg/ml, 4 mg/ml, 4.5 mg/ml, 5 mg/ml, with intermediate and nonintegral values permissible (e.g., 0.43, 0.57, 0.71 or 0.86 mg/ml).

[0109] In certain embodiments, the composition comprises adenosine at a concentration of about 3 mg/ml, and dipyridamole at a concentration of about 0.375 - 0.428 mg/ml (A:D ratios of 8:1 and 7:1), which may be rounded to 0.38 - 0.43 mg/ml. In one embodiment, for example, the composition comprises adenosine at a concentration of about 3 mg/ml and dipyridamole at a concentration of about 0.43 mg/ml (ratio 7:1). In another embodiment, the composition comprises adenosine at a concentration of about 4 mg/ml and dipyridamole at a concentration of about 0.5 - 0.57 mg/ml (ratios of about 8:1 to 7:1). In another embodiment, the composition comprises adenosine at a concentration of about 5 mg/ml and dipyridamole at a concentration of about 0.62 - 0.71 mg/ml (ratios of 8:1 and 7:1). In another embodiment the composition comprises adenosine at a concentration of about 6 mg/ml and dipyridamole at a concentration of about 0.86 mg/ml (ratio 7:1). In another embodiment, the composition comprises adenosine at a concentration of about 7 mg/ml and dipyridamole at a concentration of about 1 mg/ml (ratio 7:1), and so on, up to adenosine concentrations as high as 10 mg/ml.

[0110] In other embodiments, the composition is dry, and suitable for reconstitution prior to infusion by addition of a sterile fluid into which both dipyridamole and adenosine are readily solubilized. Usefully, the composition comprises adenosine and dipyridamole in amounts suitable to permit reconstitution in the enclosing vessel to the adenosine and dipyridamole concentrations above-described.

[0111] Whether fluid or dry, the pharmaceutical composition may further comprise carriers and excipients suitable for intravenous, intra-atrial, or intra-arterial administration, as are well known in the art. Among such excipients are those used in currently approved dipyridamole and adenosine compositions, such as tartaric acid, hydrochloric acid and polyethylene glycol (macrogol 600). Others are permissible, such as, for example, mannitol. See, http://www.adenosin.com/en/en_SPC_05.pdf (Item Development AB, 2005). See also, Remington: The Science and Practice of Pharmacy, 21st ed. (2005), Lippincott Williams & Wilkins (ISBN: 0781746736).

[0112] The compositions may further comprise additional actives, and in some embodiments, may further comprise contrast agents, including ultrasound and MRI contrast agents.

[0113] In embodiments intended for continuous intravenous infusion in the methods above-described, adenosine is typically present in the pharmaceutical composition at a concentration, or in a weight amount, that permits adenosine to be infused at a rate between about 35 µg/kg/min to about 100 µg/kg/min.

[0114] In some of these embodiments, adenosine is present in an amount that permits infusion at a rate of at least 35 µg/kg/min, at least 40 µg/kg/min, at least 45 µg/kg/min, at least 50 µg/kg/min, at least 55 µg/kg/min, at least 60 µg/kg/min, at least 65 µg/kg/min, at least 70 µg/kg/min, at least 75 µg/kg/min, at least 80 µg/kg/min, at least 85 µg/kg/min, at least 90 µg/kg/min, at least 95 µg/kg/min, and 100 µg/kg/min, with intermediate and nonintegral values permissible.

[0115] In some embodiments, adenosine is present in the composition in an amount that permits infusion at a rate of no more than 100 µg/kg/min, no more than 95 µg/kg/min, no more than 90 µg/kg/min, no more than 85 µg/kg/min, no more than 80 µg/kg/min, no more than 75 µg/kg/min, no more than 70 µg/kg/min, no more than 65 µg/kg/min, no more than 60 µg/kg/min, no more than 55 µg/kg/min, no more than 50 µg/kg/min, no more than 45 µg/kg/min, no more than 40 µg/kg/min, no more than 35 µg/kg/min, with intermediate and nonintegral values permissible.

[0116] In embodiments intended for continuous intravenous infusion, dipyridamole is typically present in the pharmaceutical composition at a concentration, or in a weight amount, that permits dipyridamole to be infused at a rate between 3.5 µg/kg/min to 50 µg/kg/min.

[0117] In some of these embodiments, dipyridamole is present in an amount that permits infusion at a rate of at least

3.5 µg/kg/min, at least 4 µg/kg/min, at least 5 µg/kg/min, at least 6 µg/kg/min, at least 7 µg/kg/min, at least 7.5 µg/kg/min, at least 8 µg/kg/min, at least 8.75 µg/kg/min, at least 9 µg/kg/min, at least 9.25 µg/kg/min, at least 9.50 µg/kg/min at least 10 µg/kg/min, at least 11 µg/kg/min, at least 11.25 µg/kg/min, at least 12 µg/kg/min, at least 12.5 µg/kg/min, at least 13 µg/kg/min, at least 13.75 µg/kg/min, at least 14 µg/kg/min, at least 15 µg/kg/min, at least 16 µg/kg/min, at least 16.25 µg/kg/min, at least 17 µg/kg/min, and at least 17.5 µg/kg/min, at least 18 µg/kg/min, at least 19 µg/kg/min, at least 20 µg/kg/min, at least 21 µg/kg/min, at least 22 µg/kg/min, at least 23 µg/kg/min, at least 24 µg/kg/min, at least 25 µg/kg/min, at least 26 µg/kg/min, at least 27 µg/kg/min, at least 28 µg/kg/min, at least 29 µg/kg/min, at least 30 µg/kg/min, at least 31 µg/kg/min, at least 32 µg/kg/min, at least 33 µg/kg/min, at least 34 µg/kg/min, at least 35 µg/kg/min, at least 36 µg/kg/min, at least 37 µg/kg/min, at least 38 µg/kg/min, at least 39 µg/kg/min, at least 40 µg/kg/min, at least 41 µg/kg/min, at least 42 µg/kg/min, at least 43 µg/kg/min, at least 44 µg/kg/min, at least 45 µg/kg/min, at least 46 µg/kg/min, at least 47 µg/kg/min, at least 48 µg/kg/min, at least 49 µg/kg/min, 50 µg/kg/min, with intermediate and nonintegral values permissible.

[0118] In some embodiments, dipyridamole is present in the composition in an amount that permits intravenous infusion at a rate of no more than 50 µg/kg/min, no more than 49 µg/kg/min, no more than 48 µg/kg/min, no more than 47 µg/kg/min, no more than 45 µg/kg/min, no more than 44 µg/kg/min, no more than 43 µg/kg/min, no more than 42 µg/kg/min, no more than 41 µg/kg/min, no more than 40 µg/kg/min, no more than 39 µg/kg/min, no more than 38 µg/kg/min, no more than 37 µg/kg/min, no more than 36 µg/kg/min, no more than about 35 µg/kg/min, no more than about 34 µg/kg/min, no more than about 33 µg/kg/min, no more than about 32 µg/kg/min, no more than about 31 µg/kg/min, no more than about 30 µg/kg/min, of no more than about 29 µg/kg/min, no more than about 28 µg/kg/min, no more than about 27 µg/kg/min, no more than about 26 µg/kg/min, no more than about 25 µg/kg/min, no more than about 24 µg/kg/min, no more than about 23 µg/kg/min, no more than about 22 µg/kg/min, no more than about 21 µg/kg/min, no more than about 20 µg/kg/min, no more than about 19 µg/kg/min, no more than about 18 µg/kg/min, no more than about 17.5 µg/kg/min, no more than about 17 µg/kg/min, no more than about 16.25 µg/kg/min, no more than about 16 µg/kg/min, no more than about 15 µg/kg/min, no more than about 14 µg/kg/min, no more than about 13.75 µg/kg/min, no more than about 13 µg/kg/min, no more than about 12.5 µg/kg/min, no more than about 12 µg/kg/min, no more than about 11.25 µg/kg/min, no more than about 11 µg/kg/min, no more than about 10 µg/kg/min, no more than about 9.25 µg/kg/min, no more than about 9.50 µg/kg/min no more than about 9 µg/kg/min, no more than about 8.75 µg/kg/min, no more than about 8 µg/kg/min, no more than about 7.5 µg/kg/min, no more than about 7 µg/kg/min, no more than about 6 µg/kg/min, no more than about 5 µg/kg/min, no more than about 4 µg/kg/min, no more than about 3.5 µg/kg/min, with intermediate and nonintegral values permissible.

3.5 Unit dosage forms

3.5.1 Dipyridamole:adenosine combined compositions

[0119] The pharmaceutical compositions described herein are usefully packaged in a unit dosage form adapted for use in the methods above-described.

[0120] In embodiments in which the pharmaceutical composition is in the form of a liquid suitable for parenteral infusion, the composition may, for example, be packaged in volumes of 2 - 50 ml. Convenient unit dosage forms contain 2 to 14 ml, typically 2, 3, 4, 5, 6, 7, 8, or 14 ml. Unit dosage forms containing volumes as low as 1 ml, and unit dosage forms containing higher volumes, such as 15 or 20 ml, are also possible. Intermediate and nonintegral volumes are permissible.

[0121] Table 1 below lists certain useful unit dosage form embodiments of the adenosine:dipyridamole pharmaceutical compositions herein described.

Table 1				
Adenosine concentration (mg/ml)	Volume (ml)	Total amount of adenosine per dosage unit (mg)	Maximal patient weight (kgs)	Total amount of dipyridamole (A:D ratio 7:1) (mg)
3	7	21	75	3
	14	42	150	6
4	7	28	100	4
	14	56	200	8
5	7	35	125	5
6	7	42	150	6

(continued)

Table 1				
Adenosine concentration (mg/ml)	Volume (ml)	Total amount of adenosine per dosage unit (mg)	Maximal patient weight (kgs)	Total amount of dipyridamole (A:D ratio 7:1) (mg)
7	2	14	50	2
	3	21	75	3
	4	28	100	4
	5	35	125	5
	6	42	150	6
	8	56	200	8

[0122] Thus, in some embodiments, the unit dosage form usefully contains 14 mg of adenosine and 2 mg of dipyridamole in 2 ml; 21 mg of adenosine and 3 mg of dipyridamole in 3 or 7 ml; 28 mg of adenosine and 4 mg of dipyridamole in 4 or 7 ml; 35 mg of adenosine and 5 mg of dipyridamole in 5 or 7 ml; 42 mg of adenosine and 6 mg of dipyridamole in 6 or 7 ml; 56 mg of adenosine and 8 mg of dipyridamole in 8 or 14 ml.

[0123] Various embodiments usefully package 9 mg or 12 mg adenosine respectively in 3 ml or 4 ml for light weight patients (e.g., children). Other embodiments usefully package 20 mg adenosine in 5 ml total volume (20mg/5ml), 24 mg in 6 ml total volume (24 mg/6ml), and 25 mg in 5 ml total volume (25 mg/5ml). Various unit dose embodiments of the pharmaceutical compositions described herein contain 30 mg adenosine in a total volume of 6 ml (30 mg/6 ml), 30 mg/10 ml, or 32 mg/8 ml. Other unit dose embodiments usefully contain adenosine at 36 mg/9 ml, 40 mg/10 ml, 40 mg/8 ml, 44 mg/11 ml, 45 mg/15 ml and 50 mg/10 ml.

[0124] In certain embodiments, the unit dosage form contains, in a total of 10 ml, 30 mg of adenosine and 3 to 5 mg of dipyridamole, with intermediate and nonintegral amounts of dipyridamole permissible (e.g., 3.75 mg or 4.28 mg, for A:D ratios of 8:1 and 7:1). In other embodiments, a 6 ml vial or ampule contains 24 mg of adenosine with 3 mg of dipyridamole (ratio 8:1). In other embodiments, an 8 ml vial or ampule will usefully contain 32 mg of adenosine with 3.2 to 5.3 mg of dipyridamole, with intermediate and nonintegral amounts of dipyridamole permissible (e.g., 4 mg to 4.57 mg, for ratios of 8:1 and 7:1).

[0125] The container for unit dosage embodiments is typically adapted for use with standard intravenous infusion sets.

[0126] In other embodiments, the unit dosage form contains adenosine and dipyridamole as solids suitable for reconstitution.

[0127] Whether liquid or dry, the unit dosage form is typically sterile and nonpyrogenic.

3.5.2 Dipyridamole unit dosage forms

[0128] In typical embodiments of the methods herein described, dipyridamole is administered at about 5% of the dose at which it is currently administered as a single agent. Accordingly, in another aspect, the disclosure provides novel unit dosage forms of dipyridamole.

[0129] Convenient unit dosage forms of dipyridamole (as single active) are vials, ampules, or prefilled syringes, usefully with 0.1 ml graduations, containing dipyridamole at a concentration of 0.5 mg/ml. At a dipyridamole dose of 35 - 40 µg/kg, a 5 mg/10 ml vial, ampule, or prefilled syringe is sufficient for almost all clinical needs, and is convenient for immediate and accurate dose adjustment. Also useful are unit dose forms of dipyridamole containing 4 mg/8 ml; 3.5 mg/7 ml; 3 mg/6 ml; 5 mg/10 ml; 6 mg/12 ml and 8 mg/16 ml.

[0130] Unit dosage forms containing dipyridamole at a concentration of 1 mg/ml -- for example, unit dosage forms containing 6 mg dipyridamole in 6 ml, 5 mg/5 ml, 4 mg/4 ml, 3 mg/3 ml ~ also find use, but fine tuning the dose may be more difficult at this higher concentration. Adjustment may be optimally obtained after dilution of dipyridamole, e.g., in saline solution.

[0131] Preparations with dipyridamole concentrations over 1 mg/ml, such as 2 mg/ml, 2.5 mg/ml, 3 mg/ml, 3.5 mg/ml, 4 mg/ml, 4.5 mg/ml, even 5 mg/ml are feasible, and are particularly convenient if it is intended to mix the whole of the dipyridamole unit dose into the adenosine unit dose, prior to administration. In this case, the 2 mg/ml, 3 mg/ml, 4 mg/ml, 5 mg/ml, 6 mg/2 ml or 8 mg/2 ml unit dosage forms (or prefilled syringes) are particularly useful.

[0132] In other embodiments disclosed herein, dipyridamole unit dosage forms contain dipyridamole at a lower concentration, such as 0.1 mg/ml. Usefully, such unit dosage forms contain 6 mg dipyridamole in 60 ml (6 mg/60 ml), 5

mg/50 ml, 4 mg/40 ml, and 3 mg/30 ml. Dipyridamole preparations at a concentration of 0.2, 0.3, and 0.4 mg/ml are also permissible.

3.5.3 Adenosine unit dosage forms

[0133] As described above, dipyridamole and adenosine may be usefully provided in separate pharmaceutical compositions, and then combined prior to administration. A dipyridamole composition may be usefully introduced into a unit dose of adenosine, and the combined composition then administered.

[0134] Thus, unit dosage forms of adenosine are disclosed, in which adenosine is formulated in sterile fluid composition, and in which the dose packaging permits sterile introduction of a second fluid in a volume at least 15% that of the adenosine composition.

[0135] Adenosine may be present at any of the concentrations at which it is present in the pharmaceutical compositions above-described - *e.g.*, usefully from 1 mg/ml to 5 mg/ml either as directly packaged, or as thereafter will be achieved upon introduction of an appropriate amount of dipyridamole composition.

[0136] For example, an adenosine unit dosage form may contain 21 mg adenosine in 6 ml (21 mg/6 ml). This will reconstitute to a desired 21 mg adenosine/7 ml (3 mg/ml adenosine) composition upon introduction of 1 ml of 3 mg/ml dipyridamole (*e.g.*, the entirety of a unit dose of dipyridamole containing 1 ml dipyridamole at 3 mg/ml). In another embodiment of the disclosure, an adenosine unit dosage form contains 42 mg/12 ml. This will reconstitute to a desired 42 mg/14 ml (3 mg/ml) adenosine upon introduction of a 6 mg/2 ml dipyridamole unit dose. In another embodiment of the disclosure, the adenosine unit dosage form contains 28 mg adenosine/6 ml, which will reconstitute to 28 mg/7 ml (4 mg/ml adenosine) upon introduction of a 4 mg/ml dipyridamole unit dose. In another embodiment of the disclosure, the adenosine unit dosage form contains 56 mg adenosine/12 ml, which will reconstitute to 56 mg/14 ml (4 mg/ml adenosine) upon introduction of an 8 mg/2 ml dipyridamole unit dose. In another embodiment of the disclosure, the adenosine unit dosage form contains 28 mg/3 ml or 35 mg/4 ml which will reconstitute to 28 mg/4 ml and 35 mg/5 ml upon introduction of a 4 or 5 mg/ml dipyridamole unit dose respectively (7 mg/ml adenosine). In another embodiment of the disclosure, the adenosine unit dosage form contains 42 mg/4 ml or 56 mg/6 ml which will reconstitute to 42 mg/6 ml and 56 mg/8 ml upon introduction of a 6 mg/2 ml or 8 mg/2 ml dipyridamole unit dose respectively (7 mg/ml adenosine). The following table summarizes the exemplary unit dosage forms.

Table 2

Initial adenosine vial volume (ml)	Volume (ml) of dipyridamole solution added to each adenosine vial	Final adenosine vial volume after addition of dipyridamole	Final adenosine concentration (mg/ml)	Total amount of adenosine per unit dosage form (mg)	Total amount of dipyridamole (A:D ratio 7:1) (mg)	Maximal patient's weight covered by the composition (kgs)
6	1	7	3	21	3	75
12	2	14	3	42	6	150
6	1	7	4	28	4	100
12	2	14	4	56	8	200
3	1	4	7	28	4	100
4	1	5	7	35	5	125
4	2	6	7	42	6	150
6	2	8	7	56	8	200

[0137] In certain embodiments of the disclosure, dipyridamole and adenosine are sequentially sampled from separate unit dosage forms, and mixed in the same syringe. In these embodiments, convenient adenosine unit dosage forms are 28 mg of adenosine in 7 ml and 56 mg of adenosine in 14 ml (4 mg/ml adenosine).

[0138] Typically, dosing and sampling are thereafter determined according to adenosine tables, and not dipyridamole tables.

3.6 Kits

[0139] In another aspect, kits are disclosed in which one or more unit doses of dipyridamole, such as those above-described, are packaged with one or more unit doses of adenosine, such as those above-described. Typically, the kit will comprise an equal number of dipyridamole and adenosine doses.

[0140] The unit dose of dipyridamole disclosed herein may be packaged in a pre-packed syringe, and the adenosine unit dose may be packaged as a vial with an injection port, such as a septum, permitting sterile introduction of dipyridamole into the adenosine dose.

[0141] The kit disclosed herein may further include one or more of an adenosine dosage table, one or more needles, diluent, and infusion sets.

4. EXAMPLES

4.1 Example 1

[0142] The effects of administering dipyridamole and adenosine intravenously as a combined (albeit, sequentially administered) pharmacological stressor were compared to the effects of administering adenosine alone in 40 consecutive patients suffering from ischemic heart disease. In the combined administration, each of the two agents was administered at a dosage lower than its clinically preferred dosage when used as a single agent for myocardial perfusion imaging. Effects were measured using noninvasive transthoracic doppler echocardiography (TTDE).

[0143] Primary efficacy end-points were peak and mean diastolic flow velocities (measured as reflecting coronary blood flow values). The secondary end-point was patient tolerance to the procedure. The protocol was designed as follows.

[0144] Forty (40) consecutive patients suffering from ischemic heart disease were enrolled. Each patient served as his own control.

[0145] Adenosine was administered by IV infusion for three minutes at the standard single-agent infusion rate of 140 $\mu\text{g/kg/min}$.

[0146] After a stabilization period of five minutes, patients then received an IV injection of dipyridamole at a total dose of either 23 $\mu\text{g/kg}$, 28 $\mu\text{g/kg}$, or 35 $\mu\text{g/kg}$, administered as a bolus over about 20 - 30 seconds. These total doses are between about 4 - 6% of the lowest single-agent total dose of dipyridamole (i.e., 0.56 mg/kg, infused over a total of 4 minutes).

[0147] The bolus injection of dipyridamole was followed immediately by an IV infusion of adenosine at 70 $\mu\text{g/kg/min}$ for 3 minutes. This dose is half the standard single-agent dosage rate of 140 $\mu\text{g/kg/min}$.

[0148] Blood flow velocity was measured in the left anterior descending coronary artery (LAD) at four time points: (i) before initial adenosine infusion (spontaneous flow at rest), (ii) during the initial 140 $\mu\text{g/kg/min}$ adenosine infusion, (iii) before the sequential administration of dipyridamole and adenosine (during the stabilization period), and (iv) during the 70 $\mu\text{g/kg/min}$ adenosine infusion, subsequent to dipyridamole bolus injection.

[0149] Results are given in Tables 3 - 6. Abbreviations used in the table are defined below:

ADE:	adenosine alone at 140 $\mu\text{g/kg/min}$
SC:	sequential combination of dipyridamole followed by adenosine at 70 $\mu\text{g/kg/min}$
PV:	peak velocity (cm/sec)
MV:	mean velocity (cm/sec)
max:	velocity under stress conditions
min:	velocity at rest (under basal conditions)
():	standard deviation
D%:	velocity differential (peak or mean), as percentage of maximum peak or mean velocity

[0150] Table 3 presents results comparing dipyridamole 28 $\mu\text{g/kg}$ IV bolus (over 20 - 30 seconds) followed by adenosine infusion at 70 $\mu\text{g/kg/min}$ ("DIP5" sequential combination) as compared to adenosine infusion alone at the standard single-agent dose of 140 $\mu\text{g/kg/min}$, in 30 patients.

Table 3								
	Peak velocity ADE		Peak velocity SC		Mean velocity ADE		Mean velocity SC	
	Max	Min	Max	Min	Max	Min	Max	Min
Mean (s.d.)	82.6 (20.7)	30.6 (8.5)	81 (21)	30.6 (9.6)	61.4 (15.1)	23.1 (6.1)	60.2 (15.3)	22.9 (7.3)
Max velocity differential	PV : 82.6 - 81 = 1.6				MV : 61.4 - 60.2 = 1.2			
Max velocity differential (D%)	1.9%				1.9 %			
P value	0.217				0.201			

[0151] Table 4 presents results comparing dipyridamole (35 µg/kg) bolus (administered over 20 - 30 seconds), followed by adenosine, administered by infusion at a rate of 70 µg/kg/min ("DIP4" sequential combination) as compared to adenosine infusion alone at the standard single-agent dose of 140 µg/kg/min, in 5 patients.

Table 4								
	Peak velocity ADE		Peak velocity SC		Mean velocity ADE		Mean velocity SC	
	Max	Min	Max	Min	Max	Min	Max	Min
Mean (s.d.)	80 (22.3)	28 (7.4)	78.8 (15.3)	28 (7.7)	60.4 (15.5)	22.2 (5)	57.8 (14.4)	23 (5.7)
Max velocity differential	PV: 80-78.8 = 1.2				MV: 60.4 - 57.8 = 2.6			
Max velocity differential (D%)	1.5%				4.3%			
P value	0.863				0.448			

[0152] Table 5 presents results comparing dipyridamole (23 µg/kg), administered as a bolus over 20 - 30 seconds, followed by adenosine, administered by infusion at a rate of 70 µg/kg/min ("DIP6" sequential combination) as compared to adenosine infusion alone at the standard single-agent dose of 140 µg/kg/min, in 5 patients.

Table 5								
	Peak velocity ADE		Peak velocity SC		Mean velocity ADE		Mean velocity SC	
	Max	Min	Max	Min	Max	Min	Max	Min
Mean (s.d.)	107 (36.5)	37.6 (12.1)	105.4 (34.3)	38.6 (16.5)	80.2 (25.4)	28.2 (11.1)	79.2 (24.2)	30 (14.7)
Max velocity differential	PV: 107 - 105.4 = 1.6				MV: 80.2 - 79.2 = 1			
Max velocity differential (D%)	1.5%				1.2%			
P value	0.842				0.771			

[0153] Table 6 presents results cumulated from all 40 patients:

Table 6								
	Peak velocity ADE		Peak velocity SC		Mean velocity ADE		Mean velocity SC	
	Max	Min	Max	Min	Max	Min	Max	Min
Mean (s.d.)	85.3 (24)	31.2 (9)	83.8 (23.3)	31.3 (10.5)	63.6 (17.3)	23.7 (6.8)	62.3 (17.3)	23.7 (8.4)
Max velocity differential	PV: 85.3 - 83.8 = 1.5				MV: 63.6 - 62.3 = 1.3			
Max velocity differential (D%)	1.75%				2%			
P value	0.314				0.109			

[0154] The measured blood flow velocities, whether peak or mean, were 3 to 4% lower, in absolute values, than those of adenosine alone (see Tables 3 to 6). However, these differences were not statistically significant (all P values > 0.05) : there was no statistical difference between the standard treatment - infusion of adenosine alone at 140 µg/kg/min - and sequential bolus administration of dipyridamole (at 4 - 6% its typical single-agent total dose) followed by adenosine at 70 µg/kg/min, whether assessed separately for each of the three tested dipyridamole doses (Tables 3 - 5), or cumulated across all dipyridamole doses (Table 6).

[0155] Table 7 shows the number and frequency of occurrence among all 40 patients of the three adverse events most commonly observed in clinical practice upon administration of adenosine alone at 140 µg/kg/min: chest pain, dyspnea, and flushing.

Table 7			
Adverse event	ADE # patients reporting events (frequency)	All SC (DIP 4/5/6) # patients reporting events (frequency)	% reduction
Chest pain	9 (22.5%)	5 (12.5%)	-44%
Dyspnea	20 (50%)	18 (45%)	slight
Flushing	21 (52.5%)	17 (42.5%)	slight

[0156] Table 8 presents descriptive statistics and analysis of the mean global visual analogue scale (VAS) score for the three main adverse events (each one being scored from 0 to 10 with a total score of 30). VAS scores provide a measure of patient self-assessment of pain intensity/discomfort.

Table 8				
		n	Mean	SD
Group DIP4	ADE	5	5.4	1.949
	SC	5	3	3.742
Group DIP5	ADE	30	5.95	3.705
	SC	30	4.383	3.38
Group DIP6	ADE	5	8.2	1.095
	SC	5	4.4	2.408

[0157] Although not tabulated in Table 8, the severity of the three main adverse symptoms, cumulated across all dipyridamole doses, was reduced by 31.6% with the sequential combination as compared to adenosine alone. This decrease was statistically significant (p=0.001). No difference was observed as among the different doses of dipyridamole: all sequential combinations reduced mean severity on the VAS of each of the three main adverse symptoms, as compared to adenosine alone at 140 µg/kg/min.

[0158] As shown in Table 7 the number (and frequency) of adverse events related to the stimulation of A1 receptors, mainly chest pain, was reduced by 44% and its severity (not shown) decreased by 60% with the sequential combination treatment as compared to adenosine alone. The number (and frequency) of adverse events related to the stimulation of A2a receptors, mainly dyspnea and flushing, did not decrease. However their severity (not shown) decreased by 24 and 38% respectively, with the sequential combination compared to adenosine alone.

[0159] The mean coronary flow reserve (ratio of maximal-stimulated coronary blood flow "CBF" to baseline-resting CBF equivalent to peak and mean blood flow velocities ratios) of the 40 patients enrolled in the study was above 2, which indicates that the observed reduction in side effects with sequential treatment was drug dependent, and not flawed by the ischemic status of the studied population.

[0160] Although not shown in the tabular data, EKG was not significantly different with the sequential combination treatment as compared to standard single-agent adenosine, and remained unchanged in all the patients. Vital signs (heart rate, systolic and diastolic blood pressures) changed similarly with the two methods. However, the heart rate increase and blood pressures decreases were less pronounced with the sequential combination than with adenosine alone.

[0161] It should be noted that in the first series of 30 patients (DIP5 group), three (3) patients received the adenosine infusion 2 minutes after the dipyridamole bolus, and two (2) patients were injected the two agents (dipyridamole and adenosine) concurrently in the same infusion line using a "Y" connector. These two modalities appeared equally effective and as effective as the immediate sequential administration protocol.

[0162] In a first control experiment, 5 patients were treated according to a modification of the experimental protocol, in which the initial adenosine infusion at 140 µg/kg/min was followed by a five (5) minute stabilization period, and thereafter by a second, single-agent, adenosine infusion at 140 µg/kg/min, without the use of dipyridamole. Results are shown in Table 9. Abbreviations used in the table are defined below:

PV: peak velocity (cm/sec)

MV: mean velocity (cm/sec)

max: velocity under stress conditions

min: velocity at rest (under basal conditions)

ADE1: first adenosine infusion

ADE2: second adenosine infusion

Table 9								
	PV-ADE1		PV-ADE2		MV-ADE1		MV-ADE2	
Patient ID	Max	Min	Max	Min	Max	Min	Max	Min
MART	68	34	60	31	49	25	45	23
LIEN	65	23	66	22	48	17	48	16
GRAND	102	30	103	29	75	22	75	22
NGHI	73	30	63	26	53	23	47	21
CORD	70	28	62	25	50	20	48	20
Mean	75.6	29	70.8	26.6	55	21.4	52.6	20.4
Max velocity differential	PV: 75.6 (ADE1) - 70.8 (ADE2) = 4.8				MV: 55 (ADE1) - 52.6 (ADE2) = 2.4			
P value	0.12				0.11			

[0163] No statistically significant differences ($P > 0.05$) were noted in velocity measurements as between first and second adenosine infusions. No significant differences were noted in the occurrence of subjective symptoms (data not shown). These data confirm previous literature reports that adenosine, administered acutely, does not induce tachyphylaxis. The data serve to validate the protocol design.

[0164] In a second set of control experiments, dipyridamole was administered alone by bolus injection to 5 patients at the dosage of 28, 35 or 40 $\mu\text{g/kg}$ after a three minute adenosine infusion at 140 $\mu\text{g/kg/min}$, and again after a 3 minute stabilization period. Data are shown in Table 10.

Table 10											
Patient ID	DIP dose ($\mu\text{g/kg}$)	PV-ADE		PV-DIP		MV-ADE		MV-DIP		Symptoms under ADE	Symptoms under DIP alone
		Max	Min	Max	Min	Max	Min	Max	Min		
GROS	28	84	27	26	23	64	20	21	19	Flushing 8/10	None
COCH	35	84	32	40	30	61	23	28	22	Flushing 6/10	None
WURI	35	88	25	23	21	71	21	20	18	None	None
STUR	40	92	33	35	33	67	23	25	23	Dyspnea 5/10	None
TALL	40	112	33	40	34	86	25	29	24	None	None

[0165] Dipyridamole did not modify peak and mean diastolic velocities. No symptoms were recorded during a follow-up of 10 minutes. The data demonstrate that intravenous bolus administration of dipyridamole as a single agent in the dosage range used in the experimental protocol, has no detectable effects; doses of dipyridamole at 28 to 40 $\mu\text{g/kg}$ alone do not induce significant hemodynamic and clinical effects.

4.2 Example 2

[0166] A Phase II study was initiated to compare dipyridamole-adenosine combination administration (also termed herein, at all doses, Adenosoft™) to adenosine alone (Adenoscan®, Astellas) as a pharmacologic stressor in coronary patients undergoing single photon emission computed tomography (SPECT) imaging studies. The study, which is on-going, is a mono-center, single-blind, 2-arm, cross-over trial.

[0167] All patients underwent a first SPECT imaging study using adenosine as single agent pharmacologic stressor at 140 $\mu\text{g/kg/min}$, according to standard clinical protocol. Only those patients in whom an ischemic zone was detected were declared eligible for the second test, and were enrolled in the study if other inclusion criteria were satisfied.

[0168] In the second test, eligible patients were stressed pharmacologically by either bolus administration of dipyridamole over 20 - 30 seconds, followed by adenosine infusion at 70 $\mu\text{g/kg/min}$ or their concurrent administration. SPECT images were acquired as per the standard approach performed the preceding week.

[0169] Randomization of SPECT images, and their analysis by two blinded readers, took place every 10 patients. Anonymous and randomized images were assessed using the standard 17-segment model and the semi-quantitative visual score method on a 5-point scale (from 0 to 4). Safety of the procedure was analyzed as in the preceding hemodynamic study (Example 1) using a visual scale, focusing on the three most common symptoms seen with adenosine, as well as on EKG changes and other usual cardiac parameters.

[0170] The study is on-going. It will include a total of about 60 patients. Preliminary results are as follows.

[0171] About 10 patients whose participation permitted the study protocol details to be finalized have been excluded from study statistics. However they provided the following information, summarized below.

[0172] The 70 $\mu\text{g/kg/min}$ adenosine + 28 $\mu\text{g/kg}$ dipyridamole combination provided images comparable to those provided by Adenoscan in 3 patients, but scored less well in two patients, at a level deemed unacceptable according to the study protocol. The 70 $\mu\text{g/kg/min}$ adenosine + 35 $\mu\text{g/kg}$ dipyridamole combination provided images comparable to Adenoscan in 7 consecutive patients, while under-scoring Adenoscan in one patient, but within the acceptable limits defined by the protocol.

[0173] In five patients of this initial series, adenosine and dipyridamole were mixed together before administration and infused over 4 minutes. This modality was successful in all the patients.

[0174] The combination stressor of 40 $\mu\text{g/kg}$ dipyridamole and 70 $\mu\text{g/kg/min}$ adenosine was tested and showed equivalent, and sometimes better results, than Adenoscan in terms of imaging efficacy in 19 patients (10 sequential and 9 concurrent administrations). Table 11 shows the stress scores (correlated to ischemia defects) for the first 10 patients of this 40 $\mu\text{g/kg}$ series (denominated, "Series I -sequential administration").

Table 11			
Series I	Stress score A (Adenoscan)	Stress score B (SC)	Delta score
Patient 1	5	4	1
Patient 2	13	11	2
Patient 3	5	5	0
Patient 4	4	4	0
Patient 5	7	7.5	-0.5
Patient 6	5	5	0
Patient 7	4	3	1
Patient 8	14	14	0
Patient 9	5	12.5	-7.5
Patient 10	5	6	-1
Total			-5

[0175] The "stress score" is the image of ischemia in one or more of the 17 cardiac segments, as scored from 0 to 4 by the blinded reader using the following scale: normal perfusion = 0, mild reduction in counts = 1, moderate reduction in counts = 2, severe reduction = 3, absence of uptake = 4). Normally, a zone of ischemia is certain when a score ≥ 4 is noted. A stress score sums the scores of all abnormal segments.

[0176] "Stress score A" is the image score difference between stress conditions between adenosine alone and rest conditions. "Stress score B" is the image score difference between stress conditions with the combination stressor - dipyridamole bolus at 40 $\mu\text{g/kg}$ over 20 - 30 seconds, followed by adenosine infusion at 70 $\mu\text{g/kg/min}$ - and resting conditions. "Delta score" is the difference between "Stress score A" and "Stress score B." It must not be >2 .

[0177] Since thallium is currently deemed the best isotope to test myocardial viability at rest, and sestamibi the best isotope to detect myocardial defects under stress conditions, a dual isotope myocardial scintigraphy technique was used for this study.

[0178] As seen from the data in Table 11, no patient had a delta score greater than 2: *i.e.*, in no patient did the combination stressor perform significantly worse as compared to Adenoscan. In three (3) patients, the delta score was negative, indicating that the ischemic defects were better visualized with the combination stressor than with Adenoscan alone. Total delta score for the ten patients, which is the primary study end-point, is below 2 and even negative

[0179] The second 40 $\mu\text{g/kg}$ series corresponding to the concurrent administration of adenosine 70 $\mu\text{g/kg/mn}$ with dipyridamole 10 $\mu\text{g/kg/mn}$ is expected to provide similar results (a preliminary analysis made by a technician who does not participate to the trial indicates that images are comparable in all the patients). However the randomized analysis of this series of images by the two blinded readers with corresponding delta scores has not yet been performed (one patient missing to get to 10)..

[0180] With respect to patient tolerance of the procedure, initial results are shown in Table 12 ($n = 27$ patients). Data from these initial results indicate that there is not much difference in symptomatology between (i) 35 $\mu\text{g/kg}$ dipyridamole combined with 70 $\mu\text{g/kg/min}$ adenosine, and (ii) 40 $\mu\text{g/kg}$ of dipyridamole combined with 70 $\mu\text{g/kg/min}$ adenosine. Results are therefore cumulated, and include data from 8 patients treated with 35 $\mu\text{g/kg}$ dipyridamole plus adenosine at 70 $\mu\text{g/kg/min}$, and from 19 patients treated with 40 $\mu\text{g/kg}$ dipyridamole, plus adenosine at 70 $\mu\text{g/kg/min}$ (SC). Given the similar adverse effect profile, the 40 $\mu\text{g/kg}$ dipyridamole dose is currently preferred, since it tends to show greater efficacy. As noted above, all 27 patients were first imaged using Adenoscan alone (ADE).

Table 12			
Adverse events	ADE (# patients reporting event)	SC (# patients reporting event)	% reduction
Chest pain	14	8	- 43 %
Dyspnea	11	9	None

(continued)

Table 12

Adverse events	ADE (# patients reporting event)	SC (# patients reporting event)	% reduction
Flushing	14	14	None
Note: Reduction in the number of chest pain events among the patients (n=19) who received dipyridamole 40µg/kg with adenosine 70µg/kg/mn was -45% with no difference regarding the number of other adverse events. # = number of			

[0181] Table 13 gives adverse event severity cumulated scores, evaluated on a visual scale (going from 0 to 10), for the first 27 patients.

Table 13

	Chest pain	Dyspnea	Flushing
Adenosine	60	40	43.5
Combination	18.5	30	44.5
% Change	-69%	-25%	+ 2.3 %
Note: % of changes among the patients (n=19) who received dipyridamole 40µg/kg with adenosine 70µg/kg/mn were very similar to those of the table above			

[0182] At this point in time, no difference in the occurrence of A₂ receptor-related side effects, or severity of symptoms, has become clearly apparent as between the combination stressor and Adenoscan. In contrast, the reduction in the occurrence (- 43 %) and severity (-69 %) of chest pain (an A₁ receptor-related adverse effect) has already become manifest in the first 27 patients. So too has reduction in ST changes on EKG, which are fewer and less severe during testing with the combination as compared to Adenoscan: in 27 patients tested to date, only 6 patients have ST variations. The difference in ST-variation for each patient alone is not significant, but the total difference has achieved significance, with a clear trend in favor of the combination. In the table below (Table 14) ST-changes/baseline are expressed in millimeters.

Table 14

Patient ID	Adenosine 140µg/kg/mn	Combination
N° 5	0.5	0
N°12	1	0.1
N°14	0.8	0.3
N°16	1	0.7
N°17	1.2	0.9
N°24	1	0
Total	5.5	2

4.3 Example 3

[0183] A study was undertaken to assess pharmacological properties of a pharmaceutical composition comprising adenosine and dipyridamole at an A:D weight ratio of 7:1. In particular, the study was conducted to assess the properties of a composition comprising adenosine at a concentration of 7 mg/ml and dipyridamole at a concentration of 1mg/ml. This particular formulation is very convenient, since it permits the use of plain figures for calculating concentrations, maximal volumes, and weights -- which is useful to reduce dosing errors in the clinical setting -- while also covering a wide range of needs, as shown in the table below:

Table 15				
Adenosine concentration (mg/ml)	Volume (ml)	Total amount of adenosine per unit (mg)	Maximal patient weight covered (kgs)	Total amount of dipyridamole per unit (mg)
7	2	14	50	2
	3	21	75	3
	4	28	100	4
	5	35	125	5
	6	42	150	6
	8	56	200	8

Dipyridamole is poorly soluble in saline and is unstable in the long term in solvents at pH > 4. Adenosine compositions currently used in clinical practice have pH > 4, and are thus poorly suited to addition of dipyridamole. Therefore, a more acidic pH was chosen. The lower pH also increases adenosine solubility above 4 mg/ml, which is the upper acceptable adenosine concentration limit in saline.

[0184] The following composition was prepared:

Table 16			
Component	Final concentration	Supplier	Reference
Adenosine	7mg/ml	Sigma Aldrich	Eur.Ph. 01/2005:1486
Dipyridamole	1mg/ml	Sigma Aldrich	Eur.Ph. 01/2005: 1199
Polyethyleneglycol 600	50mg/ml	SASOL	Eur.Ph.. 01/2005:1444
Tartaric acid	2mg/ml	Sigma Aldrich	Eur.Ph.. 01/2005:0460
Water for injection	-- (1 ml)	Aguettant	AMM: 319 508.5

After sonication for 2 minutes and magnetic stirring for 10 minutes, the solution became completely clear with a pH of 3.6 and an osmolality of 151 mosmol/kg.

[0185] In summary, adenosine (Adenoscan, Astellas) is the standard pharmacological stressor used in cardiac imaging to induce near-maximal coronary vasodilation. At its recommended dosage rate of 140 $\mu\text{g/kg/min}$, its use is attended by numerous uncomfortable side effects. The data from these studies demonstrate that the sequential bolus administration of dipyridamole at 28 - 40 $\mu\text{g/kg}$ - well below the total dose infused when dipyridamole is used as a single agent stressor - followed by infusion of adenosine at 70 $\mu\text{g/kg/min}$, is equally efficacious in providing coronary vasodilation for imaging studies, while causing fewer side effects. The data also demonstrate that dipyridamole and adenosine may be combined in a single infusion, over 4 minutes, to similar effect. Among the side effects reduced by the combination disclosed herein are chest pain, and the risk of significant heart blockage.

Claims

1. A pharmaceutical composition comprising adenosine and dipyridamole in an adenosine:dipyridamole weight ratio of 2:1 to 10:1, wherein the composition is formulated for parenteral administration to an individual subject and wherein adenosine and dipyridamole are present in amounts that permit adenosine to be administered at a dosage rate of 35 to 100 $\mu\text{g/kg/min}$ and dipyridamole to be administered at a dosage rate of 3.5 to 50 $\mu\text{g/kg/min}$, wherein the administration is continuous for a period of less than 6 minutes.
2. The pharmaceutical composition according to claim 1, in which the ratio is 7:1.
3. The pharmaceutical composition according to claim 1, wherein adenosine and dipyridamole are present in amounts that permit adenosine to be administered at a dosage rate of 70 $\mu\text{g/kg/min}$ and dipyridamole to be administered at

a dosage rate of 8.75 to 10 $\mu\text{g/kg/min}$.

4. The pharmaceutical composition according to claim 1 or 2, wherein the concentration of adenosine is 1 to 10 mg/ml, or wherein the concentration of adenosine is 3 mg/ml, 4 mg/ml, 5 mg/ml, or 7 mg/ml.
5. The pharmaceutical composition according to any one of claims 1, 2 and 4, wherein the concentration of dipyridamole is 0.1 to 5 mg/ml, or wherein the concentration of dipyridamole is 1 mg/ml.
6. A unit dosage form containing 2 to 50 ml of the composition according to any one of claims 1 to 5, wherein the composition is a sterile fluid.
7. A unit dosage form containing 5 to 60 mg of adenosine and 0.5 to 30 mg of dipyridamole, wherein the composition is a solid capable of sterile reconstitution in a physiologically acceptable solvent or solution for parenteral administration to an individual subject, and wherein the adenosine:dipyridamole weight ratio is from 2:1 to 10:1.
8. Adenosine and dipyridamole for concurrent administration, for use in a method of effecting coronary vasodilation for cardiac diagnosis, wherein the adenosine and dipyridamole are concurrently administered parenterally to an individual subject at an adenosine:dipyridamole weight ratio of 2:1 to 10:1, and wherein adenosine is administered at a dosage rate of 35 to 100 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 3.5 to 50 $\mu\text{g/kg/min}$, wherein the administration is continuous for a period of less than 6 minutes.
9. Adenosine and dipyridamole for use according to claim 8 wherein adenosine is administered at a dosage rate of 50 to 70 $\mu\text{g/kg/min}$.
10. Adenosine and dipyridamole for use according to claim 8 wherein adenosine is administered at a dosage rate of 70 $\mu\text{g/kg/min}$ and dipyridamole is administered at a dosage rate of 10 $\mu\text{g/kg/min}$.
11. Adenosine and dipyridamole for use according to claim 8
 - (a) wherein adenosine and dipyridamole are parenterally administered continuously for a period of at least 2 minutes;
 - (b) wherein adenosine and dipyridamole are parenterally administered continuously for less than 6 minutes; or
 - (c) wherein adenosine and dipyridamole are parenterally administered continuously for a period of 4 minutes.
12. Adenosine and dipyridamole for use according to claim 8 wherein adenosine and dipyridamole are administered as a single composition; or wherein adenosine and dipyridamole are administered from separate compositions.
13. Adenosine and dipyridamole for use according to claim 8 wherein the adenosine and dipyridamole compositions are administered by intravenous infusion.
14. Adenosine and dipyridamole for use according to claim 8 wherein the adenosine:dipyridamole ratio is 7:1.
15. Adenosine and dipyridamole for use according to claim 8 further comprising the step of: assessing cardiac function, optionally wherein assessing cardiac function includes use of one or more techniques selected from the group consisting of: electrocardiography, M mode echography, two dimensional echography, three dimensional echography, echo-doppler, cardiac imaging, planar (conventional) scintigraphy, single photon emission computed tomography (SPECT), dynamic single photon emission computed tomography, positron emission tomography (PET), first pass radionuclide angiography, equilibrium radionuclide angiography, nuclear magnetic resonance (NMR) imaging, perfusion contrast echocardiography, digital subtraction angiography (DSA), and ultrafast x-ray computed tomography (CINE CT).

Patentansprüche

1. Pharmazeutische Zusammensetzung, umfassend Adenosin und Dipyridamol in einem Adenosin:Dipyridamol-Gewichtsverhältnis von 2:1 bis 10:1, wobei die Zusammensetzung für parentale Verabreichung an ein individuelles Subjekt formuliert ist und wobei Adenosin und Dipyridamol in Mengen vorhanden sind, die zulassen, dass Adenosin mit einer Dosierungsrate von 35 bis 100 $\mu\text{g/kg/min}$ verabreicht wird und Dipyridamol mit einer Dosierungsrate von

3,5 bis 50 $\mu\text{g/kg/min}$ verabreicht wird, wobei die Verabreichung für einen Zeitraum von weniger als 6 Minuten kontinuierlich ist.

2. Pharmazeutische Zusammensetzung nach Anspruch 1, in der das Verhältnis 7:1 ist.
3. Pharmazeutische Zusammensetzung nach Anspruch 1, wobei Adenosin und Dipyridamol in Mengen vorhanden sind, die zulassen, dass Adenosin mit einer Dosierungsrate von 70 $\mu\text{g/kg/min}$ verabreicht wird und Dipyridamol mit einer Dosierungsrate von 8,75 bis 10 $\mu\text{g/kg/min}$ verabreicht wird.
4. Pharmazeutische Zusammensetzung nach Anspruch 1 oder 2, wobei die Konzentration von Adenosin 1 bis 10 mg/ml ist oder wobei die Konzentration von Adenosin 3 mg/ml, 4 mg/ml, 5 mg/ml oder 7 mg/ml ist.
5. Pharmazeutische Zusammensetzung nach einem der Ansprüche 1, 2 und 4, wobei die Konzentration von Dipyridamol 0,1 bis 5 mg/ml ist oder wobei die Konzentration von Dipyridamol 1 mg/ml ist.
6. Einheitsdosierungsform, die 2 bis 50 ml der Zusammensetzung nach einem der Ansprüche 1 bis 5 enthält, wobei die Zusammensetzung ein steriles Fluid ist.
7. Einheitsdosierungsform, die 5 bis 60 mg Adenosin und 0,5 bis 30 mg Dipyridamol enthält, wobei die Zusammensetzung ein Feststoff ist, der zu steriler Rekonstitution in einem/einer physiologisch annehmbaren Lösungsmittel oder Lösung für parentale Verabreichung an ein individuelles Subjekt fähig ist, und wobei das Adenosin:Dipyridamol-Gewichtsverhältnis von 2:1 bis 10:1 ist.
8. Adenosin und Dipyridamol für gleichzeitige Verabreichung zur Verwendung in einem Verfahren des Bewirkens von koronarer Vasodilatation für eine Herzdiagnose, wobei das Adenosin und Dipyridamol einem individuellen Subjekt bei einem Adenosin:Dipyridamol-Gewichtsverhältnis von 2:1 bis 10:1 gleichzeitig parental verabreicht werden, und wobei Adenosin mit einer Dosierungsrate von 35 bis 100 $\mu\text{g/kg/min}$ verabreicht wird und Dipyridamol mit einer Dosierungsrate von 3,5 bis 50 $\mu\text{g/kg/min}$ verabreicht wird, wobei die Verabreichung für einen Zeitraum von weniger als 6 Minuten kontinuierlich ist.
9. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8, wobei Adenosin mit einer Dosierungsrate von 50 bis 70 $\mu\text{g/kg/min}$ verabreicht wird.
10. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8, wobei Adenosin mit einer Dosierungsrate von 70 $\mu\text{g/kg/min}$ verabreicht wird und Dipyridamol mit einer Dosierungsrate von 10 $\mu\text{g/kg/min}$ verabreicht wird.
11. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8,
 - (a) wobei Adenosin und Dipyridamol für einen Zeitraum von mindestens 2 Minuten kontinuierlich parental verabreicht werden;
 - (b) wobei Adenosin und Dipyridamol weniger als 6 Minuten lang kontinuierlich parental verabreicht werden; oder
 - (c) wobei Adenosin und Dipyridamol für einen Zeitraum von 4 Minuten kontinuierlich parental verabreicht werden.
12. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8, wobei Adenosin und Dipyridamol als eine einzige Zusammensetzung verabreicht werden; oder wobei Adenosin und Dipyridamol aus separaten Zusammensetzungen verabreicht werden.
13. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8, wobei die Adenosin- und Dipyridamol-Zusammensetzungen durch intravenöse Infusion verabreicht werden.
14. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8, wobei das Adenosin:Dipyridamol-Verhältnis 7:1 ist.
15. Adenosin und Dipyridamol zur Verwendung nach Anspruch 8, ferner umfassend den folgenden Schritt: Bewerten von Herzfunktion, wobei optional das Bewerten von Herzfunktion das Verwenden von einer oder mehreren Techniken umfasst, ausgewählt aus der Gruppe bestehend aus: Elektrokardiographie, M-Modus-Echographie, zweidimensionaler Echographie, dreidimensionaler Echographie, Echo-Doppler, Herzbildgebung, planarer (herkömmlicher) Szintigraphie, Einzelphotonen-Emissions-Computertomographie (SPECT), dynamischer Einzelphotonen-Emissions-Computertomographie, Positronenemissionstomographie (PET), First-Pass-Radionuklidangiographie, Äquilib-

rium-Radionuklidangiographie, Kernspinresonanztomographie (NMR), Durchblutungs-Kontrastechokardiographie, digitaler Subtraktionsangiographie (DSA) und ultraschneller Röntgen-Computertomographie (CINE CT).

5 Revendications

1. Composition pharmaceutique comprenant de l'adénosine et du dipyridamole en un rapport en poids adénosine:dipyridamole de 2:1 à 10:1, la composition étant formulée pour l'administration parentérale à un sujet individuel et l'adénosine et le dipyridamole étant présents en quantités qui permettent à l'adénosine d'être administrée à un taux de dosage de 35 à 100 $\mu\text{g/kg/min}$ et au dipyridamole d'être administré à un taux de dosage de 3,5 à 50 $\mu\text{g/kg/min}$, l'administration étant continue pendant une période inférieure à 6 minutes.
2. Composition pharmaceutique selon la revendication 1, dans laquelle le rapport est de 7:1.
3. Composition pharmaceutique selon la revendication 1, dans laquelle l'adénosine et le dipyridamole sont présents en quantités qui permettent à l'adénosine d'être administrée à un taux de dosage de 70 $\mu\text{g/kg/min}$ et au dipyridamole d'être administré à un taux de dosage de 8,75 à 10 $\mu\text{g/kg/min}$.
4. Composition pharmaceutique selon la revendication 1 ou 2, dans laquelle la concentration d'adénosine est de 1 à 10 mg/ml ou dans laquelle la concentration d'adénosine est de 3 mg/ml, 4 mg/ml, 5 mg/ml ou 7 mg/ml.
5. Composition pharmaceutique selon l'une quelconque des revendications 1, 2 et 4, dans laquelle la concentration de dipyridamole est de 0,1 à 5 mg/ml, ou dans laquelle la concentration de dipyridamole est de 1 mg/ml.
6. Forme posologique unitaire contenant 2 à 50 ml de la composition selon l'une quelconque des revendications 1 à 5, dans laquelle la composition est un fluide stérile.
7. Forme posologique unitaire contenant 5 à 60 mg d'adénosine et 0,5 à 30 mg de dipyridamole, dans laquelle la composition est un solide capable d'une reconstitution stérile dans un solvant ou une solution physiologiquement acceptable pour l'administration parentérale à un sujet individuel, et dans laquelle le rapport en poids adénosine:dipyridamole est de 2:1 à 10:1.
8. Adénosine et dipyridamole pour l'administration concurrente, destinés à une utilisation dans un procédé permettant d'effectuer une vasodilatation coronarienne pour un diagnostic cardiaque, l'adénosine et le dipyridamole étant administrés de manière concurrente par voie parentérale à un sujet individuel à un rapport en poids adénosine:dipyridamole est de 2:1 à 10:1, et l'adénosine étant administrée à un taux de dosage de 35 à 100 $\mu\text{g/kg/min}$ et le dipyridamole étant administré à un taux de dosage de 3,5 à 50 $\mu\text{g/kg/min}$, l'administration étant continue pendant une période inférieure à 6 minutes.
9. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8, l'adénosine étant administrée à un taux de dosage de 50 à 70 $\mu\text{g/kg/min}$.
10. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8, l'adénosine étant administrée à un taux de dosage de 70 $\mu\text{g/kg/min}$. et le dipyridamole étant administré à un taux de dosage de 10 $\mu\text{g/kg/min}$.
11. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8,
 - (a) l'adénosine et le dipyridamole étant administrés par voie parentérale en continu pendant une période d'au moins 2 minutes ;
 - (b) l'adénosine et le dipyridamole étant administrés par voie parentérale en continu pendant moins de 6 minutes ;
 - ou
 - (c) l'adénosine et le dipyridamole étant administrés par voie parentérale en continu pendant une période de 4 minutes.
12. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8, l'adénosine et le dipyridamole étant administrés sous la forme d'une composition unique ; ou l'adénosine et le dipyridamole étant administrés sous la forme de compositions séparées.

13. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8, les compositions d'adénosine et de dipyridamole étant administrées par perfusion intraveineuse.

5

14. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8, le rapport adénosine:dipyridamole étant de 7:1.

10

15. Adénosine et dipyridamole destinés à une utilisation selon la revendication 8, comprenant en outre l'étape d'évaluation de la fonction cardiaque, l'évaluation de la fonction cardiaque comprenant éventuellement l'utilisation d'une ou de plusieurs techniques choisies dans le groupe constitué par: l'électrocardiographie, l'échographie en mode M, l'échographie bidimensionnelle, l'échographie tridimensionnelle, l'écho-doppler, l'imagerie cardiaque, la scintigraphie planaire (classique), la tomographie d'émission monophotonique (TEMP), la tomographie d'émission monophotonique dynamique, la tomographie par émission de positrons (TEP), l'angiographie isotopique de premier passage, l'angiographie isotopique à l'équilibre, l'imagerie par résonance magnétique nucléaire (RMN), l'échocardiographie avec perfusion de produit de contraste, l'angiographie par soustraction numérique (ASN) et la tomographie à rayons X ultra-rapide (CINE CT).

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6531457 B [0009]
- US 6448235 B [0009]
- US 6322771 B [0009]
- US 5877180 A [0009]

Non-patent literature cited in the description

- Remington: The Science and Practice of Pharmacy.
Lippincott Williams & Wilkins, 2005 [0111]

专利名称(译)	用于药理学压力测试的方法，组合物，单位剂型和试剂盒，具有降低的副作用		
公开(公告)号	EP2049007A1	公开(公告)日	2009-04-22
申请号	EP2007765052	申请日	2007-07-04
[标]申请(专利权)人(译)	ADENOBIO		
申请(专利权)人(译)	ADENOBIO N.V.		
当前申请(专利权)人(译)	ADENOBIO N.V.		
[标]发明人	GORNYPHILIPPE		
发明人	GORNYPHILIPPE		
IPC分类号	A61B5/00 A61K31/519 A61K31/7076 A61P9/08		
CPC分类号	A61K31/519 A61K31/7076 A61K49/0004		
优先权	60/818928 2006-07-05 US 11/772684 2007-07-02 US		
其他公开文献	EP2049007B1		
外部链接	Espacenet		

摘要(译)

提出了包括给予包含腺苷和双嘧达莫的药物组合物的方法，以及包括作为推注给予双嘧达莫与作为输注给予的腺苷的组合给药的方法，两者均以低于其各自的单一药剂剂量的剂量，用于检测存在和/或评估药理学压力测试期间心肌缺血的严重程度。该方法可用于利用腺苷的血管舒张能力，在该剂量下，与腺苷相关的副作用显著降低，同时实现最佳冠状动脉灌注。还提供了可用于实施所述方法的组合物，单位剂型和试剂盒。