



(11) **EP 3 242 588 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
16.10.2019 Bulletin 2019/42

(51) Int Cl.:
A61B 5/00 (2006.01) G01N 33/48 (2006.01)
G01N 33/53 (2006.01) G01N 33/68 (2006.01)

(21) Application number: **16735507.2**

(86) International application number:
PCT/US2016/012725

(22) Date of filing: **08.01.2016**

(87) International publication number:
WO 2016/112337 (14.07.2016 Gazette 2016/28)

(54) **BLOOD BASED BIOMARKERS FOR DIAGNOSING ATHEROSCLEROTIC CORONARY ARTERY DISEASE**

BLUTBASIERTE BIOMARKER ZUR DIAGNOSTIZIERUNG ATHEROSKLEROTISCHER
KORONARARTERIENERKRANKUNG

BIOMARQUEURS SANGUINS POUR LE DIAGNOSTIC DE CORONAROPATHIE
ATHÉROSCLÉREUSE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(74) Representative: **Conroy, John
Fish & Richardson P.C.
Highlight Business Towers
Mies-van-der-Rohe-Straße 8
80807 München (DE)**

(30) Priority: **09.01.2015 US 201562101445 P**

(43) Date of publication of application:
15.11.2017 Bulletin 2017/46

(56) References cited:
**WO-A1-2011/063470 WO-A1-2014/135696
WO-A2-02/089657 US-A1- 2013 338 031
US-A1- 2014 303 236**

(73) Proprietor: **Global Genomics Group, LLC
Atlanta, GA 30355 (US)**

(72) Inventors:
• **VOROS, Szilard
Atlanta, GA 30355 (US)**
• **BROWN, Bradley O.
Atlanta, GA 30355 (US)**
• **MARVASTY, Idean B.
Atlanta, GA 30355 (US)**

- **LABORDE CARLOS M ET AL: "Plasma metabolomics reveals a potential panel of biomarkers for early diagnosis in acute coronary syndrome", METABOLOMICS, SPRINGER NEW YORK LLC, US, vol. 10, no. 3, 26 October 2013 (2013-10-26), pages 414-424, XP036158507, ISSN: 1573-3882, DOI: 10.1007/S11306-013-0595-9 [retrieved on 2013-10-26]**
- **ALBRECHT ET AL.: 'Metabolite profiling reveals new insights into the regulation of serum urate in humans' METABOLOMICS vol. 10, no. 1, 2014, pages 141 - 151, XP036158475 DOI: 10.1007/S11306-013-0565-2**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 242 588 B1

Description**TECHNICAL FIELD**

[0001] This invention relates to non-invasive methods for diagnosing atherosclerotic coronary artery disease ("AS-CAD") and/or detecting a coronary atherosclerotic plaque in a patient, and more particularly to methods of using metabolomic blood-based biomarkers and their combinations to identify patients having atherosclerotic CAD and/or as having a coronary atherosclerotic plaque.

BACKGROUND

[0002] Cardiovascular disease (CVD) remains the leading cause of morbidity and mortality both in the United States and worldwide. Based on the 2014 Heart Disease and Stroke Update by the American Heart Association (AHA), 83.6 million American adults have at least one type of CVD (>1 in 3 prevalence). Of these adults, 15.4 million suffer from coronary heart disease (CHD), with the following breakdown: myocardial infarction (MI): 7.6 million and angina pectoris (AP): 7.8 million. It is estimated that by 2030, 43.9% of the US population will have some form of CVD.

[0003] The underlying cause of CVD is atherosclerotic coronary artery disease (ASCAD), which begins with the development of an atherosclerotic plaque in the coronary arterial vasculature. Atherosclerosis is of unquestionable importance, in terms of human health and societal cost. It is responsible for coronary artery disease (CAD) and cerebrovascular disease, both of which are leading causes of morbidity and mortality worldwide. Atherosclerosis is also responsible for peripheral arterial disease, a leading medical cause of limb-loss.

[0004] The development and progression of atherosclerotic CAD follows two distinct paths that may lead from one to another, and lead to two distinct clinical syndromes. One of the two broad clinical syndromes is "exertional angina", also called "stable angina" or "stable CAD." In this context, progressively growing atherosclerotic plaque leads to progressively worsening coronary arterial luminal stenosis, or narrowing, that starts to impede blood flow to the myocardium (heart muscle). When luminal stenosis, or narrowing, reaches a critical limit, approximately 70% diameter stenosis, there is significant pressure drop over the stenosis, and myocardial perfusion (blood flow to the heart muscle) is compromised, especially during periods of increased myocardial oxygen demand, such as during exertion or emotional states. This myocardial oxygen "supply-demand mismatch" leads to myocardial ischemia, when blood flow to the heart muscle can no longer keep up with demand. It is myocardial ischemia itself that triggers pain receptors in the heart, leading to the clinical symptoms of "angina pectoris", or chest pain. While this process typically causes chest pain during exertion or emotional states, this typically does not lead to acute coronary syndromes, such as sudden cardiac death (SCD), myocardial infarction (MI; heart attack) or unstable angina (UA). Typical treatment options include medical interventions to slow the heart rate with beta blockers or calcium channel blockers, and to improve blood flow with drugs like nitrates. Typical interventional approaches are aimed at relieving the stenosis, by either percutaneously placing a stent inside the stenosis (percutaneous coronary intervention; PCI) to open the narrowing, or to bypass the stenotic (narrowed) segment using arterial or venous grafts during coronary artery bypass grafting (CABG) surgery. As pointed out above, the root cause of this pathophysiology and clinical course of "exertional angina" is coronary arterial atherosclerosis, or atherosclerotic CAD.

[0005] The second broad path in pathophysiology with a corresponding clinical picture is referred to as "acute coronary syndromes" (ACS), which encompasses sudden cardiac death (SCD), myocardial infarction (MI) and unstable angina (UA). In this context, the atherosclerotic plaque (the root cause) in the coronary artery can get disrupted either through rupture, or erosion. When a coronary atherosclerotic plaque gets disrupted, this triggers the formation of a thrombus (clot) over the disrupted region, which may partially or completely occlude the entire coronary artery, leading to abrupt cutoff of blood supply to the myocardium (acute myocardial ischemia). This pathophysiology may manifest in sudden cardiac death (SCD), myocardial infarction (MI), or resting angina. Typical treatment options are commonly interventional options, where the acutely disrupted plaque is covered up with a stent during an invasive procedure. Just as in the case of "exertional angina", the root cause of acute coronary syndromes is atherosclerotic plaque in the coronary arterial vasculature.

[0006] Accordingly, the initiation and progression of atherosclerotic plaques are the root cause of the clinical manifestations of atherosclerotic CAD and CHD. Thus, there remains a need for improved diagnostic methods for detecting an atherosclerotic plaque and for diagnosing atherosclerotic CAD.

[0007] Laborde et al., *Metabolomics* (2014) 10:414-424, WO 2011/063470 A1 and WO 2014/135696 A1 disclose metabolite biomarkers for diagnosis of coronary heart disease.

SUMMARY

[0008] The present invention is defined by the independent claims. The dependent claims depict additional embodi-

ments of the invention.

[0009] In one aspect, the disclosure provides non-invasive methods for evaluating a human subject for having atherosclerotic coronary artery disease (ASCAD) or as having a coronary atherosclerotic plaque. ASCAD includes, for example, atherosclerosis. The methods include measuring the levels of each analyte within a panel of analyte biomarkers in a biological sample obtained from a subject, wherein the panel of analyte biomarkers is selected from the group consisting of:

(i) a panel of biomarkers comprising valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4) and mannose;

, wherein the levels, including relative levels, or the occurrences of the plurality of analyte biomarkers are indicative of having ASCAD and/or for the presence of a coronary atherosclerotic plaque. Any of the panels above may further include the analysis of age and gender.

[0010] In another aspect, the disclosure provides a method comprising measuring by mass spectrometry the levels of each of a plurality of metabolites in a blood sample obtained from a human subject, wherein the plurality of metabolites is selected from the group consisting of (i) valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4) and mannose.

[0011] Also provided in aspects of the disclosure are panels of biomarkers useful for evaluating a human subject for having ASCAD or as having a coronary atherosclerotic plaque. By a "panel of biomarkers" it is meant a collection, or combination, of two or more molecular entities, e.g. two, three, four, five, or more than five entities, whose representation in a sample is associated with ASCAD or a coronary atherosclerotic plaque. The panel of biomarkers described herein may be used to diagnose ASCAD or identify the presences of a coronary atherosclerotic plaque., to provide a prognosis to a patient having ASCAD or a coronary atherosclerotic plaque, to provide a prediction of the responsiveness of a patient with ASCAD or a coronary atherosclerotic plaque to a medical therapy, to monitor a patient having ASCAD or a coronary atherosclerotic plaque, to treat a patient having ASCAD or a coronary atherosclerotic plaque, etc.

[0012] In some embodiments, the methods further comprise of performing a comparison between the measured levels of the analytes in the biological sample with one or more reference samples, said references being representative of matched (e.g., age, gender, etc.) human subjects.

[0013] In some embodiments, the methods further include identifying a subject as having ASCAD or as having a coronary atherosclerotic plaque if the measured levels of the analytes in the biological sample are increased or decreased relative to the amounts of the analytes in the reference samples.

[0014] In some embodiments, the methods include instructing a healthcare professional (e.g., a physician, physician assistant, nurse practitioner, nurse and case manager) to complete a non-invasive cardiovascular evaluation on a subject identified as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque. The further non-invasive cardiovascular evaluation can confirm whether the subject has or does not have atherosclerotic coronary artery disease or a coronary atherosclerotic plaque.

[0015] In some embodiments, the methods further comprise of performing a non-invasive cardiovascular evaluation on a subject identified as having ASCAD or as having a coronary atherosclerotic plaque to confirm the subject does or does not have atherosclerotic coronary artery disease or a coronary atherosclerotic plaque. The non-invasive cardiovascular evaluation can be a procedure selected from the group consisting of cardiovascular computed tomography (CT) imaging, an exercise stress test, a pharmacologic stress test, myocardial perfusion imaging, stress echocardiography, and cardiovascular magnetic resonance imaging.

[0016] In yet another embodiment, compositions for use in treating atherosclerotic coronary artery disease further comprise selectively administering a composition comprising an effective amount of a therapeutic agent to a subject identified as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque to treat the subject, wherein the therapeutic agent is selected from the group consisting of a statin, cholesterol absorption inhibitors, niacin-derivatives, omega-3-fatty acid compounds, bile acid sequestrants, PCSK9 inhibitor, anti-platelet agents, aldosterone blockers, angiotensin-converting enzyme (ACE) inhibitors, angiotensin-receptor blockers (ARBs), beta blockers, diuretics, digitalis, hydralazine and nitrates, warfarin and aspirin.

[0017] For subjects identified as having ASCAD or as having a coronary atherosclerotic plaque, the methods may further comprise selecting a treatment plan for a subject. For example, the compositions for use in treating atherosclerotic coronary artery disease may further comprise selecting a treatment plan for a subject, which comprises selectively administering a composition comprising an effective amount of a therapeutic agent, and/or performing a non-invasive cardiovascular evaluation, which comprises, for example, performing a procedure selected from the group consisting of cardiovascular computed tomography (CT) imaging, an exercise stress test, a pharmacologic stress test, myocardial perfusion imaging, stress echocardiography, and cardiovascular magnetic resonance imaging. The treatment plan can comprise prescribing to the subject therapeutic lifestyle changes to improve cardiovascular health.

[0018] In some embodiments, the level of each analyte is measured using mass spectrometry (MS) analysis. The MS analysis method can be liquid chromatography-tandem mass spectrometry (UPLC-MS/MS), chromatography-mass

spectrometry (GC-MS), or nuclear magnetic resonance (NMR).

[0019] In some aspects, the methods disclosed herein further comprise selecting a subject identified as having ASCAD or as having a coronary atherosclerotic plaque for treatment. In some embodiments, the treatment comprises selectively administering a composition comprising an effective amount of a therapeutic agent selected from the group consisting of a statin, cholesterol absorption inhibitors, niacin-derivatives, omega-3-fatty acid compounds, bile acid sequestrants, PCSK9 inhibitors, anti-platelet agents, aldosterone blockers, angiotensin-converting enzyme (ACE) inhibitors, angiotensin-receptor blockers (ARBs), beta blockers, diuretics, digitalis, hydralazine and nitrates, warfarin and aspirin to a subject identified as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque to treat the subject. In some embodiments, the treatment comprises performing a non-invasive cardiovascular evaluation on a subject identified as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque to confirm whether the subject does or does not have atherosclerotic coronary artery disease or a coronary atherosclerotic plaque. In some embodiments, the treatment plan comprises prescribing therapeutic lifestyle changes to improve cardiovascular health.

[0020] In another aspect, the invention provides a kit for evaluating a human subject for having ASCAD or as having a coronary atherosclerotic plaque. The kits include reagents suitable for determining levels of a plurality of analytes in a test sample (e.g., reagents suitable for determining levels of the metabolomic biomarkers disclosed herein); optionally one or more control samples comprising predetermined levels of the same analytes, wherein comparison of the levels of the analytes in a test sample with levels in the control samples identifies a subject as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque; and instructions for use of the kit in the method described herein.

[0021] As used herein, the term "biological sample" or "sample" refers to a sample obtained or derived from a subject. By way of example, the sample may be selected from the group consisting of body fluids, blood, whole blood, plasma, serum, mucus secretions, urine or saliva. In some embodiments the sample is, or comprises a blood sample. The preferred biological source for detection of the biomarkers is a blood sample, a serum sample or a plasma sample.

[0022] The term "subject" as used herein refers to a mammal. A subject therefore refers to, for example, dogs, cats, horses, cows, pigs, guinea pigs, humans and the like. When the subject is a human, the subject may be referred to herein as a patient. The subject can be symptomatic (e.g., the subject presents symptoms associated with atherosclerotic CAD, such as, for example, chest pain, angina, angina equivalent, dyspnea, or dyspnea on exertion; and/or presents risk factors associated with coronary artery disease, such as, for example, male gender, hypertension, dyslipidemia, diabetes, post-menopausal state in females, smoking, or a family history of coronary artery disease), or the subject can be asymptomatic (e.g., the subject does not present symptoms associated with atherosclerotic CAD).

[0023] As used herein, "obtain" or "obtaining" can be any means whereby one comes into possession of the sample by "direct" or "indirect" means. Directly obtaining a sample means performing a process (e.g., performing a physical method such as extraction) to obtain the sample. Indirectly obtaining a sample refers to receiving the sample from another party or source (e.g., a third party laboratory that directly acquired the sample). Directly obtaining a sample includes performing a process that includes a physical change in a physical substance, e.g., a starting material, such as a blood, e.g., blood that was previously isolated from a patient. Thus, obtain is used to mean collection and/or removal of the sample from the subject. Furthermore, "obtain" is also used to mean where one receives the sample from another who was in possession of the sample previously.

[0024] In some embodiments, the reference sample is obtained from at least one individual not suffering from a cardiovascular disease. In some other embodiments, the reference sample is obtained from at least one individual previously diagnosed as having a cardiovascular disease (e.g., atherosclerotic coronary artery disease (ASCAD) or a coronary atherosclerotic plaque). In some embodiments, the reference sample comprises a predetermined, statistically significant reference analyte levels.

[0025] In some embodiments, the determining step is performed using a gradient boosting algorithm. In some embodiments, the determining step is performed using generalized linear modeling.

[0026] In yet another example, the methods further comprise modifying the subject's clinical record to identify the subject as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque. Preferably, the clinical record is stored in a computer readable medium.

[0027] A summary of the relevant analysis models described herein are provided below:

a. Atherosclerosis in Native CAD - A^{nCAD}

- i. Significant in Fasting & Non-Fasting Combined - A_{FNF}^{nCAD}
- ii. Independently Significant in Fasting and Non-Fasting - A_{IFNF}^{nCAD}
- iii. Significant in Fasting - A_F^{nCAD}

b. Atherosclerosis in All CAD (including revascularization) - A^{aCAD}

i. Significant in Fasting & Non-Fasting Combined - A_{FNF}^{aCAD}

ii. Independently Significant in Fasting and Non-Fasting - A_{IFNF}^{aCAD}

iii. Significant in Fasting - A_F^{aCAD}

c. 50% stenosis in Native CAD

i. Significant in Fasting & Non-Fasting Combined - S_{FNF}^{nCAD}

ii. Independently Significant in Fasting and Non-Fasting - S_{IFNF}^{nCAD}

iii. Significant in Fasting - S_F^{nCAD}

d. 50% stenosis in ALL CAD (including revascularization)

i. Significant in Fasting & Non-Fasting Combined - S_{FNF}^{aCAD}

ii. Independently Significant in Fasting and Non-Fasting - S_{IFNF}^{aCAD}

iii. Analytes Significant in Fasting - S_F^{aCAD} .

[0028] The section headings used herein are for organizational purposes only and are not to be construed as limiting the described subject matter in any way. When definitions of terms in references appear to differ from the definitions provided in the present teachings, the definition provided in the present teachings shall control. It will be appreciated that there is an implied "about" prior to metrics such as temperatures, concentrations, and times discussed in the present teachings, such that slight and insubstantial deviations are within the scope of the present teachings herein. In this application, the use of the singular includes the plural unless specifically stated otherwise. Also, the use of "comprise," "comprises," "comprising," "contain," "contains," "containing," "include," "includes," and "including" are not intended to be limiting. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention. The articles "a" and "an" are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0029] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. In case of conflict between references mentioned herein and the present specification, the present specification, including definitions, will control.

[0030] Other features and advantages of the invention will be apparent from the following detailed description and figures, and from the claims.

DESCRIPTION OF DRAWINGS

[0031]

Figure 1 is a graph demonstrating a box plot of Diamond Forrester Score by clinical group.

Figure 2 is a graph demonstrating a ROC curve for Diamond Forrester Score and Native CAD (AUC = 0.5; Threshold = inf; Sensitivity = 0; Specificity = 1; Accuracy = 0.57).

Figure 3 is a graph demonstrating a box plot of Morise Score by clinical group.

Figure 4 is a graph demonstrating a ROC curve for Morise Score of Native CAD (AUC = 0.58; Threshold = 20; Sensitivity = 0.04; Specificity = 0.98; Accuracy = 0.58).

Figure 5 is a graph demonstrating a ROC curve for the Clinical Model of Native CAD (AUC = 0.79; Threshold = 0.5; Sensitivity = 0.49; Specificity = 0.9; Accuracy = 0.73).

Figure 6 is a series of graphs that provide Partial Dependence Plots for the eleven clinical variables for Clinical Model of Native CAD.

Figure 7 is a graph demonstrating a ROC curve for the Metabolomic Model of Native CAD (AUC = 0.78; Threshold = 0.39; Sensitivity = 0.82; Specificity = 0.64; Accuracy = 0.72).

Figure 8 is a series of graphs that provide Partial Dependence Plots for the six clinical variables for Metabolomic Model of Native CAD.

Figure 9 is a series of graphs that provide Partial Dependence Plots for the ten clinical variables for Clinical-Metab-

olomic Model of Native CAD.

Figure 10 is a graph demonstrating a ROC curve for the Clinical-Metabolomic Model of Native CAD (AUC = 0.82; Threshold = 0.5; Sensitivity = 0.56; Specificity = 0.88; Accuracy = 0.74).

Figure 11 is a series of graphs that provide Partial Dependence Plots for the seven clinical variables for Lipidomic Model of Native CAD.

Figure 12 is a graph demonstrating a ROC curve for Lipidomic Model of Native CAD (AUC = 0.74; Threshold = 0.46; Sensitivity = 0.6; Specificity = 0.79; Accuracy = 0.71).

Figure 13 is a series of graphs that provide Partial Dependence Plots for the seven clinical variables for Clinical-Lipidomic Model of Native CAD.

Figure 14 is a graph demonstrating a ROC curve for the Clinical-Lipidomic Model of Native CAD (AUC = 0.76; Threshold = 0.47; Sensitivity = 0.52; Specificity = 0.84; Accuracy = 0.7).

Figure 15 is a series of graphs that provide Partial Dependence Plots for the six clinical variables for Clinical-Metabolomic-Lipidomic Model of Native CAD.

Figure 16 is a graph demonstrating a ROC curve for the Clinical-Metabolomics-Lipidomic Model of Native CAD (AUC = 0.78; Threshold = 0.47; Sensitivity = 0.52; Specificity = 0.89; Accuracy = 0.73).

Figure 17 is a graph demonstrating a ROC curve for Diamond Forrester Score and All CAD (AUC = 0.53; Threshold = 0.51; Sensitivity = 0.39; Specificity = 0.72; Accuracy = 0.55).

Figure 18 is a graph demonstrating a ROC curve for Morise Score of All CAD (AUC = 0.64; Threshold = 12; Sensitivity = 0.73; Specificity = 0.48; Accuracy = 0.61).

Figure 19 is a series of graphs that provide Partial Dependence Plots for the eleven clinical variables for Clinical Model of All CAD.

Figure 20 is a graph demonstrating a ROC curve for the Clinical Model of All CAD (AUC = 0.82; Threshold = 0.49; Sensitivity = 0.76; Specificity = 0.74; Accuracy = 0.75).

Figure 21 is a series of graphs that provide Partial Dependence Plots for the sixteen clinical variables for Metabolomics Model of All CAD.

Figure 22 is a graph demonstrating a ROC curve for Metabolomics Model of All CAD (AUC = 0.85; Threshold = 0.48; Sensitivity = 0.9; Specificity = 0.65; Accuracy = 0.78).

Figure 23 is a series of graphs that provide Partial Dependence Plots for the twelve clinical variables for Clinical-Metabolomics Model of All CAD.

Figure 24 is a graph demonstrating a ROC curve for the Clinical-Metabolomic Model of All CAD (AUC = 0.83; Threshold = 0.56; Sensitivity = 0.69; Specificity = 0.81; Accuracy = 0.75).

Figure 25 is a series of graphs that provide Partial Dependence Plots for the thirteen clinical variables for Lipidomics Model of All CAD.

Figure 26 is a graph demonstrating a ROC curve for Lipidomic Model of All CAD (AUC = 0.79; Threshold = 0.48; Sensitivity = 0.82; Specificity = 0.64; Accuracy = 0.73).

Figure 27 is a series of graphs that provide Partial Dependence Plots for the eleven clinical variables for Clinical Lipidomics Model of All CAD.

Figure 28 is a graph demonstrating a ROC curve for the Clinical-Lipidomic Model of All CAD (AUC = 0.8; Threshold = 0.54; Sensitivity = 0.64; Specificity = 0.84; Accuracy = 0.74).

Figure 29 is a series of graphs that provide Partial Dependence Plots for the twelve clinical variables for Clinical-Metabolomic-Lipidomic Model of All CAD.

Figure 30 is a graph demonstrating a ROC curve for the Clinical-Metabolomics-Lipidomic Model of All CAD (AUC = 0.82; Threshold = 0.46; Sensitivity = 0.84; Specificity = 0.63; Accuracy = 0.74).

Figure 31 is a schematic diagram describing a diagnostic algorithm in patient suspected of having atherosclerotic CAD.

Figure 32 is a schematic diagram describing a diagnostic algorithm evaluating a subject for having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque.

Figure 33 is a graph demonstrating a box plot of Diamond Forrester Score by clinical group .

Figure 34 is a graph demonstrating a ROC curve for Diamond Forrester Score and A^{nCAD} (AUC = 0.45; Threshold = 0.00; Sensitivity = 1.00; Specificity = 0.00; Accuracy = 0.62).

Figure 35 is a graph demonstrating a ROC curve for the Metabolomic Model of A_{FNF}^{nCAD} (AUC = 0.82; Threshold = 0.45; Sensitivity = 0.85; Specificity = 0.61; Accuracy = 0.76).

Figure 36 is a graph demonstrating a ROC curve for the Metabolomic Model of A_{IFNF}^{nCAD} (AUC = 0.80; Threshold = 0.45; Sensitivity = 0.85; Specificity = 0.64; Accuracy = 0.77).

Figure 37 is a graph demonstrating a ROC curve for the Metabolomic Model of A_F^{nCAD} (AUC = 0.81; Threshold = 0.48; Sensitivity = 0.87; Specificity = 0.58; Accuracy = 0.76).

Figure 38 is a graph demonstrating a ROC curve for Diamond Forrester Score and A^{aCAD} (AUC = 0.45; Threshold

=0.00; Sensitivity =1.00; Specificity =0.00; Accuracy =0.64).

Figure 39 is a graph demonstrating a ROC curve for the Metabolomic Model of A_{FNF}^{aCAD} (AUC =0.83; Threshold =0.40; Sensitivity =0.93; Specificity =0.50; Accuracy =0.78).

Figure 40 is a graph demonstrating a ROC curve for the Metabolomic Model of A_{IFNF}^{aCAD} (AUC =0.80.; Threshold =0.47; Sensitivity =0.85; Specificity =0.63; Accuracy =0.77).

Figure 41 is a graph demonstrating a ROC curve for the Metabolomic Model of A_F^{aCAD} (AUC =0.82; Threshold =0.56; Sensitivity =0.83; Specificity =0.64; Accuracy =0.76).

Figure 42 is a graph demonstrating a ROC curve for Diamond Forrester Score and S^{nCAD} (AUC =0.45; Threshold =0.94; Sensitivity =0.03; Specificity =1.00; Accuracy =0.78).

Figure 43 is a graph demonstrating a ROC curve for the Metabolomic Model of S_{FNF}^{nCAD} (AUC =0.73; Threshold =0.45; Sensitivity =0.21; Specificity =0.96; Accuracy =0.79).

Figure 44 is a graph demonstrating a ROC curve for the Metabolomic Model of S_{IFNF}^{nCAD} (AUC =0.76; Threshold =0.42; Sensitivity =0.30; Specificity =0.96; Accuracy =0.81).

Figure 45 is a graph demonstrating a ROC curve for the Metabolomic Model of S_F^{nCAD} (AUC =0.67 ; Threshold =0.90; Sensitivity =0.04; Specificity =1.0; Accuracy =0.78).

Figure 46 is a graph demonstrating a ROC curve for Diamond Forrester Score and S^{aCAD} (AUC =0.45; Threshold =0.94; Sensitivity =0.03; Specificity =1.00; Accuracy =0.75).

Figure 47 is a graph demonstrating a ROC curve for the Metabolomic Model of S_{FNF}^{aCAD} (AUC =0.78; Threshold =0.51; Sensitivity =0.34; Specificity =0.94; Accuracy =0.78).

Figure 48 is a graph demonstrating a ROC curve for the Metabolomic Model of S_{IFNF}^{aCAD} (AUC =0.78; Threshold =0.54; Sensitivity =0.23; Specificity =0.97; Accuracy =0.77).

Figure 49 is a graph demonstrating a ROC curve for the Metabolomic Model of S_F^{aCAD} (AUC =0.74; Threshold =0.74; Sensitivity =0.22; Specificity =0.96; Accuracy =0.77).

DETAILED DESCRIPTION

[0032] The initiation and progression of atherosclerotic plaques is the root cause of the clinical manifestations of atherosclerotic CAD and CHD. The first step and the "sine qua non" of atherosclerosis, is the retention of atherogenic lipoprotein particles, which is followed by a compensatory, secondary inflammatory response. The infiltration of the coronary arterial wall by monocytes and macrophages creates a feedback loop that further triggers the progression of atherosclerosis. The macrophages in the coronary arterial wall start the phagocytosis of atherogenic lipoprotein particles and the uncontrolled uptake of atherogenic lipids and lipoproteins leads to apoptosis and necrosis of the macrophages, creating a lipid-rich necrotic core (LRNC). It is this state of the LRNC that is most susceptible to rupture and erosions. Later stages of the atherosclerotic process involve healing through calcification and fibrosis.

[0033] It follows that blood-based biomarkers provide insight into the atherosclerotic process at two levels. First, blood-based biomarkers can assess causal factors that lead to the development of atherosclerosis; and second, blood-based biomarkers can assess the consequences and responses to the initial steps in the atherosclerotic process.

[0034] In contrast to the biological paradigm outlined above, the clinical diagnosis of atherosclerotic CAD is predicated on population-based probability-driven clinical risk stratification (Figure 31).

[0035] The clinical diagnostic algorithm depends on clinical risk- and symptom-based risk stratification. Asymptomatic patients are evaluated through global risk scores, such as the Framingham Risk Score, or the more recently developed and introduced Pooled Risk Estimator. These global risk scores attempt to estimate the probability of a cardiovascular event occurring over a period of time, but are not designed to make the diagnosis of atherosclerotic CAD.

[0036] The clinical evaluation of symptomatic patients, those who are suspected of having atherosclerotic CAD based on their symptoms, is driven by risk scores such as the Diamond-Forrester algorithm. The Diamond-Forrester system classifies patients into one of three categories to provide a pre-test likelihood of obstructive CAD (low [$<15\%$], intermediate [$15-85\%$] or high [$>85\%$]), based on age, gender and clinical attributes of chest pain. In patients with low pre-test likelihood of CAD, no further evaluation is recommended. The largest group of patients is those with intermediate pre-test likelihood of CAD. In this group, the recommendation is to perform a myocardial perfusion imaging (MPI) study to assess the hemodynamic consequence of any potential luminal stenosis. While the published sensitivity and specificity of is approximately 80% and 80% in the research studies, real-life accuracy of MPI is vastly lower than this, with specificity often under 30% (Thomas GS, et al. A Blood-Based Gene Expression Test for Obstructive Coronary Artery Disease Tested in Symptomatic Nondiabetic Patients Referred for Myocardial Perfusion Imaging The COMPASS Study. Circulation: Cardiovascular Genetics 2013;6(2): 154-162.). Patients with high pre-test likelihood of CAD are referred for invasive coronary angiography to define the degree and extent of luminal stenosis, and to further refer for percutaneous or surgical

revascularization, as appropriate.

[0037] Importantly, in this paradigm, none of the diagnostic approaches have been designed to diagnose the root cause of atherosclerotic CAD, the atherosclerotic plaque itself. However, in the context of unbiased, hypothesis-free discovery studies using specific phenotyping tools for the detection of atherosclerotic plaque itself, it is now possible to

[0038] The blood-based biomarkers disclosed herein allow for detection of an atherosclerotic plaque per se and allow for early diagnosis of disease and provide an opportunity to significantly change the downstream diagnostic evaluation and treatment process.

[0039] The blood-based biomarkers identified herein are designed to detect atherosclerotic CAD by directly detecting atherosclerotic plaque and are calibrated to have a high negative predictive value. Asymptomatic subjects at risk for atherosclerotic CAD, or symptomatic patients suspected of having atherosclerotic CAD are candidates for the biomarker testing. If the test is negative, this excludes the presence of atherosclerotic CAD, and no further testing is warranted. If the test is positive, this indicates the presence of an atherosclerotic plaque and the presence of atherosclerotic CAD, and the patient can be referred for cardiovascular CT, or other non-invasive cardiovascular diagnostic testing to confirm the results. If the cardiovascular CT or other non-invasive cardiovascular diagnostic testing detects non-obstructive plaque and non-obstructive CAD, the patient is prescribed medical therapy to treat the factors that lead to the development of the plaque, in order to halt or reverse the further progression of atherosclerotic CAD. If, on the other hand, the cardiovascular CT examination or other non-invasive cardiovascular diagnostic testing shows the presence of obstructive atherosclerotic plaque and obstructive CAD, the patient is referred for invasive coronary angiography, with or without functional testing, to determine if the plaque causes hemodynamically significant stenosis, and coronary revascularization may be performed, as appropriate. Therefore, in this paradigm, the blood-based biomarker plays a central role in determining patient management. Of note, it is well-known that approximately 39% of all invasive coronary angiograms in the United States show no significant coronary artery disease, and another ~20% have non-obstructive CAD (Patel M, et al., Low Diagnostic Yield of Elective Coronary Angiography. N Engl J Med. 2010 Mar 11; 362(10): 886-95). The proportion of negative or non-obstructive CT angiograms are even higher; therefore, the blood-based biomarker that can exclude the presence of atherosclerotic CAD will avoid a very large number of unnecessary invasive and CT coronary angiograms, and other non-invasive cardiovascular diagnostic tests.

[0040] Previously, metabolomics approaches have been attempted in the context of CVD, but not in the context of atherosclerotic CAD. Cheng et al. have identified branched-chain amino acids, other hydrophobic amino acids, tryptophan breakdown products, and nucleotide metabolites associated with cardiovascular disease. (Cheng S, et al., Metabolite Profiling Identifies Pathways Associated With Metabolic Risk in Humans / Clinical Perspective. Circulation 2012;125(18):2222-2231). Rather than evaluating atherosclerosis, Bodi et al. have used myocardial ischemia as an outcomes variable to identify metabolic signatures. (Bodi V, Sanchis J, Morales JM, Marrachelli VG, Nunez J, Forteza MJ et al. Metabolomic Profile of Human Myocardial Ischemia by Nuclear Magnetic Resonance Spectroscopy of Peripheral Blood Serum: A Translational Study Based on Transient Coronary Occlusion Models. J Am Coll Cardiol 2012;59(18): 1629-1641)

[0041] The terms "decrease", "decreased", "reduced", "reduction" or "downregulated" are all used herein generally to mean a decrease by a statistically significant amount. However, for avoidance of doubt, "reduced", "reduction", "decreased" or "decrease" means a decrease by at least 10% as compared to a reference level, for example a decrease by at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% decrease (i.e. absent level as compared to a reference sample), or any decrease between 10-100% as compared to a reference level, or at least about a 0.5-fold, or at least about a 1.0-fold, or at least about a 1.2-fold, or at least about a 1.5-fold, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold decrease, or any decrease between 1.0-fold and 10-fold or greater as compared to a reference level.

[0042] The terms "increased", "increase" or "up-regulated" are all used herein to generally mean an increase by a statistically significant amount; for the avoidance of any doubt, the terms "increased" or "increase" means an increase of at least 10% as compared to a reference level, for example an increase of at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% increase or any increase between 10-100% as compared to a reference level, or at least about a 0.5-fold, or at least about a 1.0-fold, or at least about a 1.2-fold, or at least about a 1.5-fold, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold increase, or any increase between 1.0-fold and 10-fold or greater as compared to a reference level.

[0043] The present invention relates to methods useful for the characterization of (e.g., clinical evaluation, diagnosis, classification, prediction, profiling) of atherosclerotic CAD and or coronary atherosclerotic plaque based on the levels or occurrence of certain analytes referred to herein as biomarkers, or analyte biomarkers. As used herein, levels refer to the amount or concentration of an analyte in a sample (e.g., a plasma or serum sample) or subject. Whereas, occurrence refers to the presence or absence of a detectable analyte in a sample. Thus, level is a continuous indicator of amount,

whereas occurrence is a binary indicator of an analyte. In some cases, an occurrence may be determined using a threshold level above which a biomarker is present and below which a biomarker is absent.

[0044] The analyte biomarkers described herein are particularly useful for characterizing (e.g., assessing or evaluating) a subject for having atherosclerotic CAD or as having a coronary atherosclerotic plaque in a non-invasive manner.

[0045] The invention relates to the discovery of a plurality of biomarkers that are useful for characterizing an atherosclerosis-related metabolic disorder. Thus, in some aspects, the invention provides method comprising measuring the level of each analyte within a selected panel of analyte biomarkers, wherein a panel comprises a plurality of analyte biomarkers, in a biological sample obtained from a subject. The number of biomarkers, or metabolites, in the plurality (at least two) may be 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 11 or more, 12 or more, 13 or more, 14 or more, 15 or more, 16 or more, 17 or more, 18 or more, 19 or more, or 20 or more, e.g., 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, or more.

[0046] The methods described herein are useful for identifying whether a subject has ASCAD or has a coronary atherosclerotic plaque. Stated another way, the methods described herein are useful for determining the probability a subject has atherosclerotic CAD or has a coronary atherosclerotic plaque, the methods relying on differences in the relative amounts of the analyte biomarker panels obtained from the subject, wherein the probability is determined using a gradient boosting algorithm or generalized linear model, as described herein.

[0047] Moreover, the methods described herein are useful for diagnosing whether a subject has atherosclerotic CAD or has a coronary atherosclerotic plaque. As used herein, diagnosing includes both diagnosing and aiding in diagnosing. Thus, other diagnostic criteria may be evaluated in conjunction with the results of the methods in order to make a diagnosis.

[0048] According to some embodiments, the method comprises determining the amount (i.e., determining the level, measuring the amount, or measuring the level) of each (i.e., all) analyte within a panel of analyte biomarkers. In some embodiments, any of the panels below may further include the analysis of the age and/or gender of the subject.

[0049] In some embodiments, the analytes are selected from the group consisting of valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4), mannose. Thus, in some embodiments, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4), mannose.

[0050] In some examples, the analytes are selected from the group consisting of acisoga, o.cresol.sulfate, threonate, cysteine.glutathione.disulfide. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: acisoga, o.cresol.sulfate, threonate, cysteine.glutathione.disulfide.

[0051] In some examples, the analytes are selected from the group consisting of N.acetylphenylalanine, N.acetylleucine, valylleucine, xylitol, 2.hydroxyhippurate (salicylurate), glycylphenylalanine, serylleucine, fucose. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: N.acetylphenylalanine, N.acetylleucine, valylleucine, xylitol, 2.hydroxyhippurate (salicylurate), glycylphenylalanine, serylleucine, fucose.

[0052] In some examples, the analytes are selected from the group consisting of glutamate, acisoga, valylleucine, mannose, glucuronate, urate, valerylcarnitine (C5), 12.HETE. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: glutamate, acisoga, valylleucine, mannose, glucuronate, urate, valerylcarnitine (C5), 12.HETE.

[0053] In some examples, the analytes are selected from the group consisting of o.cresol.sulfate, acisoga, cysteine.glutathione.disulfide, glucose, 1.nonadecanoylglycerophosphocholine (19:0), threonate. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: o.cresol.sulfate, acisoga, cysteine.glutathione.disulfide, glucose, 1.nonadecanoylglycerophosphocholine (19:0), threonate.

[0054] In some examples, the analytes are selected from the group consisting of N.acetylphenylalanine, N.acetylleucine, glycylphenylalanine, valylleucine, xylitol, 12.HETE, 2.hydroxyhippurate (salicylurate). Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: N.acetylphenylalanine, N.acetylleucine, glycylphenylalanine, valylleucine, xylitol, 12.HETE, 2.hydroxyhippurate (salicylurate).

[0055] In some examples, the analytes are selected from the group consisting of N.acetylthreonine, tigloylglycine, glycerate, valerylcarnitine (C5), threonate, 2.hydroxyhippurate (salicylurate), salicylate, tartronate (hydroxymalonated). Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: N.acetylthreonine, tigloylglycine, glycerate, valerylcarnitine (C5), threonate, 2.hydroxyhippurate (salicylurate), salicylate, tartronate (hydroxymalonated).

[0056] In some examples, the analytes are selected from the group consisting of N.acetyl glycine, 3.ethylphenylsulfate. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: N.acetyl glycine, 3.ethylphenylsulfate.

[0057] In some examples, the analytes are selected from the group consisting of serotonin (5HT), N.acetylputrescine, leucylleucine, propionylglycine (C3), cholate, glycocholenate.sulfate, asparagine, 3.ethylphenylsulfate. Accordingly, in

some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: serotonin (5HT), N.acetylputrescine, leucylleucine, propionylglycine (C3), cholate, glycocholate.sulfate, asparagine, 3 .ethylphenylsulfate.

[0058] In some examples, the analytes are selected from the group consisting of acisoga, mannose, valeryl carnitine (C5), 1.linoleoyl.GPE (18.2), glycoirsodeoxycholate, threonate, 2.hydroxyhippurate (salicylurate), salicylate. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: acisoga, mannose, valeryl carnitine (C5), 1.linoleoyl.GPE (18.2), glycoirsodeoxycholate, threonate, 2.hydroxyhippurate (salicylurate), salicylate.

[0059] In some examples, the analytes are selected from the group consisting of N.acetyl glycine, threonate, 2.hydroxyhippurate (salicylurate), 3.ethylphenylsulfate, salicylate. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: N.acetyl glycine, threonate, 2.hydroxyhippurate (salicylurate), 3.ethylphenylsulfate, salicylate.

[0060] In some examples, the analytes are selected from the group consisting of asparagine, taurine, acisoga, 1.oleoylglycerol (18:1), cholesterol, 2.hydroxyhippurate (salicylurate), 6.oxopiperidine.2.carboxylic.acid, propionylglycine (C3). Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: asparagine, taurine, acisoga, 1.oleoylglycerol (18:1), cholesterol, 2.hydroxyhippurate (salicylurate), 6.oxopiperidine.2.carboxylic.acid, propionylglycine (C3).

[0061] In some examples, the analytes are selected from the group consisting of valylleucine, isoleucylvaline, glutamine-leucine, X-12212, glucuronate, and glutamate. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: valylleucine, isoleucylvaline, glutamine-leucine, X-12212, glucuronate, and glutamate.

[0062] In some examples, the analytes are selected from the group consisting of glucose, gamma-glutamyl transferase, uric acid, low density lipoprotein cholesterol, triglycerides, lipoprotein (a) mass, alkaline phosphatase and apolipoprotein B. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: glucose, gamma-glutamyl transferase, uric acid, low density lipoprotein cholesterol, triglycerides, lipoprotein (a) mass, alkaline phosphatase and apolipoprotein B.

[0063] In some examples, the analytes are selected from the group consisting of glutamate, X-16132, isoleucylalanine, isoleucylphenylalanine, lipoprotein (a) mass, small dense low density lipoprotein cholesterol, X-12212, ADSGEGDFX-AEGGGVR (SEQ ID NO: 1), glutamine. leucine, and threonylleucine. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: glutamate, X-16132, isoleucylalanine, isoleucylphenylalanine, lipoprotein (a) mass, small dense low density lipoprotein cholesterol, X-12212, ADSGEGDFX-AEGGGVR (SEQ ID NO: 1), glutamine-leucine, and threonylleucine.

[0064] In some examples, the analytes are selected from the group consisting of CER22:1, HexCER18:1, TG15:0, SM16:0, CER20:0, DHC20:1, and CER18:0. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: CER22:1, HexCER18:1, TG15:0, SM16:0, CER20:0, DHC20:1, and CER18:0.

[0065] In some examples, the analytes are selected from the group consisting of CER20:0, apolipoprotein B, SM16:0, lipoprotein (a) mass, DHC20L1, alkaline phosphatase, and CER18:0. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: CER20:0, apolipoprotein B, SM16:0, lipoprotein (a) mass, DHC20L1, alkaline phosphatase, and CER18:0

[0066] In some examples, the analytes are selected from the group consisting of glutamate, X-16132, isoleucylalanine, Lipoprotein (a) mass, X-12212, and glutamine-leucine. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: glutamate, X-16132, isoleucylalanine, Lipoprotein (a) mass, X-12212, and glutamine-leucine.

[0067] In some examples, the analytes are selected from the group consisting of oleic2, N-terminal pro-brain natriuretic peptide, lipoprotein (a) mass, insulin, glycated hemoglobin A1C, cismontaotl, small dense low density lipoprotein cholesterol, behenic2, adiponectin, and gamma-glutamyl transferase. Thus, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: oleic2, N-terminal pro-brain natriuretic peptide, lipoprotein (a) mass, insulin, glycated hemoglobin A1C, cismontaotl, small dense low density lipoprotein cholesterol, behenic2, adiponectin, and gamma-glutamyl transferase.

[0068] In some examples, the analytes are selected from the group consisting of isoleucylalanine, glucuronate, glycine, isoleucylphenylalanine, mannose, X-21452, 1-oleoylglycerophosphoglycerol, X-21335, 7-methylxanthine, X-12729, acisoga, alpha-hydroxyisovaleroyl carnitine, 2-aminoadipate, X-18914, tigloylglycine, and pyridoxal. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: isoleucylalanine, glucuronate, glycine, isoleucylphenylalanine, mannose, X-21452, 1-oleoylglycerophosphoglycerol, X-21335, 7-methylxanthine, X-12729, acisoga, alpha-hydroxyisovaleroyl carnitine, 2-aminoadipate, X-18914, tigloylglycine, and pyridoxal.

[0069] In some examples, the analytes are selected from the group consisting of isoleucylalanine, glycine, isoleucyl-

phenylalanine, oleic2, X-21335, methylxanthine, aminoadipate, mannose, acisoga, arachidonoyl.GPE.20.4, and triglycerides. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: isoleucylalanine, glycine, isoleucylphenylalanine, oleic2, X-21335, methylxanthine, aminoadipate, mannose, acisoga, arachidonoyl.GPE.20.4, and triglycerides

[0070] In some examples, the analytes are selected from the group consisting of DHC18:0, DG18:1n9, total diacylglycerol, TG20:3n9, CE20:4n6, CER18:1, DHC20:1, CE18:2n6, PL18:2n6, SM18:0, FA14:1n5, PL15:0, and CE16:0. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: DHC18:0, DG18:1n9, total diacylglycerol, TG20:3n9, CE20:4n6, CER18:1, DHC20:1, CE18:2n6, PL18:2n6, SM18:0, FA14:1n5, PL15:0, and CE16:0.

[0071] In some examples, the analytes are selected from the group consisting of glycated hemoglobin A1C, total diacylglycerol, cismontotl, lipoprotein (a) mass, TG20.3n9, oleic2, DHC18.1, N-terminal pro-brain natriuretic peptide, gamma-glutamyl transferase, and CER18. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: glycated hemoglobin A1C, total diacylglycerol, cismontotl, lipoprotein (a) mass, TG20.3n9, oleic2, DHC18.1, N-terminal pro-brain natriuretic peptide, gamma-glutamyl transferase, and CER18.

[0072] In some examples, the analytes are selected from the group consisting of isoleucylalanine, glycine, cismontotl, lipoprotein (a) mass, X-21452, isoleucylphenylalanine, glycated hemoglobin A1C, X-21335, X7.methylxanthine, acisoga and X-12729. Accordingly, in some examples, the methods disclosed herein comprise measuring the levels of the following analytes in the sample: isoleucylalanine, glycine, cismontotl, lipoprotein (a) mass, X-21452, isoleucylphenylalanine, glycated hemoglobin A1C, X-21335, X7.methylxanthine, acisoga and X-12729.

[0073] Thus, in a representative embodiment, the disclosure provides a method for evaluating a subject for having ASCAD or as having a coronary atherosclerotic plaque in a subject, the method comprising (a) obtaining a plasma or serum sample from a human subject; (b) determining the amount of each analyte from a panel of analyte biomarkers; (c) comparing the relative amount of the analytes in the biological sample with the relative amount of the analytes in a control sample; and (d) determining the probability the subject has atherosclerotic coronary artery disease or a coronary atherosclerotic plaque if the relative amount of the analytes in the biological sample are increased or decreased relative to the amount of the analytes in a control sample.

[0074] In one aspect, the disclosure provides a method for evaluating a subject for having atherosclerotic coronary artery disease (ASCAD) or as having a coronary atherosclerotic plaque in a subject, the method comprising measuring the level of each analyte within a selected panel of analyte biomarkers in a biological sample obtained from a subject. In some embodiments, the method measuring the level of one or more analytes from a panel of analyte biomarkers, wherein the panel of analyte biomarkers is selected from the group consisting of:

(i) valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4), mannose;

[0075] In some embodiments, analysis of any of the panels take into account the age and/or gender of the subject.

[0076] In one aspect, the disclosure provides a method for evaluating a subject for having atherosclerotic coronary artery disease (ASCAD) or as having a coronary atherosclerotic plaque in a subject, the method comprising measuring the level of each analyte within a selected panel of analyte biomarkers in a biological sample obtained from a subject. In some examples, the method measuring the level of one or more analytes from a panel of analyte biomarkers, wherein the panel of analyte biomarkers is selected from the group consisting of

(xiii) linoleic2, Lipoprotein (a) cholesterol, Apolipoprotein B, Alkaline phosphatase, Lipoprotein (a) mass, MHDM2, B.Sitosterol, Campesterol,

Lipoprotein (a) mass, Gamma-glutamyl transferasepalmleic2, Glucose, ProInsulin, cholesterol, Fish Oil, High Sensitive C Reactive Protein, Small dense Low Density Lipoprotein Cholesterol, Uric acid, Low Density Lipoprotein Cholesterol, Insulin, Triglycerides, and Vitamin D;

(xiv) 1-myristoylglycerol (14:0), glycerol 3-phosphate (G3P), serylleucine, 1-nonadecanoylglycerophosphocholine(19:0), glycine, theobromine, 1-oleoylglycerol (18:1), glycyltryptophan, threonate, 1-oleoylglycerophosphoglycerol, guanidinosuccinate, threonylleucine, 1-oleoyl-GPC (18:1), histidylphenylalanine, tigloylglycine, 1-stearoylglycerophosphoglycerol, hydroxybutyrylcarnitine, tryptophylphenylalanine, 2-aminooctanoate, imidazole lactate, urate, 2-arachidonoyl-GPE (20:4), imidazole propionate, valylglycine, 2-docosahexaenoylglycerophosphoethanolamine, indolepropionate, valylisoleucine, 2-hydroxybutyrate (AHB), isobutyrylglycine (C4), valylleucine, 2prime-deoxyuridine, isoleucylalanine, X - 12212, 3 7-dimethylurate, isoleucylglycine, X - 12472, 3-ethylphenylsulfate, isoleucylisoleucine, X - 12524, 3-hydroxy-2-ethylpropionate, isoleucylleucine, X - 12544, 3-methyl-2-oxobutyrate, isoleucylphenylalanine, X - 12824, 3-methylglutaryl carnitine-1, isoleucylvaline, X - 14056, 3-methylxanthine, leucylglycine, X - 14291, 4-hydroxyphenylacetate, leucylserine, X - 15245, 7-methylurate, mannose, X - 16129, 7-methylxanthine, methyl glucopyranoside (alpha + beta), X - 16132, acisoga, methyl indole-3-acetate, X - 17178,

ADSGEGDFXAEGGGVR (SEQ ID NO: 1), N2 N2-dimethylguanosine, X - 21289, alpha-glutamyltyrosine, N4-acetylcytidine, X - 21335, alpha-hydroxyisovaleroyl carnitine, N-acetylalanine, X - 21365, alpha-ketobutyrate, N-acetyl-beta-alanine, X - 21452, alpha-ketoglutarate, N-acetylputrescine, X - 21626, asparagine, N-acetylthreonine, X - 21662, carnitine, N-acetylvaline, xanthine, 1-myristoylglycerol (14:0), glycerol 3-phosphate (G3P), serylleucine, 1-nonadecanoylglycerophosphocholine(19:0), glycine, theobromine, 1-oleoylglycerol (18:1), glycytryptophan, threonate, 1-oleoylglycerophosphoglycerol, guanidinosuccinate, threonylleucine, 1-oleoyl-GPC (18:1), histidylphenylalanine, tigloylglycine, 1-stearoylglycerophosphoglycerol, hydroxybutyrylcarnitine, tryptophylphenylalanine, 2-aminooctanoate, imidazole, lactate, and urate;

(xv) CE16:1n7, DG16:0, DHC20:1, HexCER16:0, SM16:0, CE18:2n6, DG18:0, DHC24:0, HexCER18:1, SM20:1, CE20:4n6, DG18:1n9, DHC24:1, LacCER16:0, TG15:0, CER18:0, DG20:3n9, FA18:3n6, PL15:0, TG16:0, CER20:0, DHC18:0, FA20:3n6, PLdm16:0, total Triacylglycerol. CER22:1, DHC18:1, HexCER14:0, and PLdm18:1n9;

(xvi) Glycated hemoglobin A1C, Lipoprotein (a) mass, nervonic2, B.Sitosterol, Adiponectin, Triglycerides, Omega 6 total, behenic2, Campesterol, N-terminal pro-brain natriuretic peptide, oleic2, Desmosterol, palmleic2, cismontol, Gamma-glutamyl transferase, ProInsulin, Glucose, Small dense Low Density Lipoprotein Cholesterol, transpalm2, High Density Lipoprotein Cholesterol, Uric acid, High Density Lipoprotein Fraction 3, Insulin, Vitamin D, linoleic2, Lipoprotein (a) cholesterol, and MHDm2;

(xvii) 1-3-dimethylurate, glycerol 3-phosphate (G3P), pyroglutamylglutamine, 1 7-dimethylurate, glycine, pyroglutamylglycine, 12-HETE, glycyphenylalanine, pyruvate, 1-linoleoyl-GPE (18:2), glycytryptophan, S-adenosylhomocysteine (SAH), 1-methylurate, guanidinosuccinate, salicylate, 1-nonadecanoylglycerophosphocholine(19:0), hexanoylcarnitine (C6), serylleucine, 1-oleoylglycerol (18:1), histidylphenylalanine, succinylcarnitine (C4), 1-oleoylglycerophosphoglycerol, homostachydrine, threonate, 2-aminoadipate, hydroxybutyrylcarnitine, threonylleucine, 2-aminobutyrate, imidazole propionate, thymol sulfate, 2-aminooctanoate, indolepropionate, tigloylglycine, 2-arachidonoyl-GPE (20:4), isobutyrylglycine (C4), tryptophylglycine, 2-docosahexaenoylglycerophosphoethanolamine, isoleucylalanine, tryptophylphenylalanine, 2-hydroxybutyrate (AHB), isoleucylglycine, tyrosylglutamine, 2-hydroxyhippurate (salicylurate), isoleucylisoleucine, urate, 2-linoleoyl-GPE (18:2), isoleucylleucine, valerylalanine (C5), 2prime-deoxyuridine, isoleucylphenylalanine, valylglycine, 3-ethylphenylsulfate, isovalerylglycine, valylisoleucine, 3-hydroxyisobutyrate, kynurenine, valylleucine, 3-methyl-2-oxobutyrate, leucylaspartate, valylvaline, 3-methylglutarylalanine-1, leucylglycine, X - 11429, 3-methylglutarylalanine-2, leucylserine, X - 11444, 3-methylxanthine, lysylleucine, X - 11787, 4-hydroxyphenylacetate, mannose, X - 11945, 5alpha-androstan-3beta 17beta-diol monosulfate 2, methyl glucopyranoside (alpha + beta), X - 12212, 7-methylurate, methyl indole-3-acetate, X - 12472, 7-methylxanthine, N2 N2-dimethylguanosine, X - 12729, acetylcarnitine (C2), N4-acetylcytidine, X - 12824, acisoga, N6-carbamoylthreonyladenosine, X - 14056, ADSGEGDFXAEGGGVR, N-acetylalanine, X - 15245, alpha-glutamyltyrosine, N-acetylglutamine, X - 15492, alpha-hydroxyisovaleroyl carnitine, N-acetylisoleucine, X - 16129, alpha-ketobutyrate, N-acetylneuraminic acid, X - 16132, alpha-ketoglutarate, N-acetylthreonine, X - 17178, beta-tocopherol, N-acetylvaline, X - 17690, butyrylcarnitine (C4), N-methyl proline, X - 18914, carnitine, octanoylcarnitine (C8), X - 18922, cysteine-glutathione disulfide, oleic ethanolamide, X - 19438, cytidine, O-sulfo-L-tyrosine, X - 21289, fucose, phenylalanyl aspartate, X - 21335, gamma-glutamylisoleucine, prolylphenylalanine, X - 21365, gamma-glutamylvaline, propionylcarnitine (C3), X - 21367, glucose, propionylglycine (C3), X - 21452, glucuronate, pseudouridine, X - 21471, glutamate, pyridoxal, X - 21626, glutamine-leucine, pyridoxate, and xanthine; and

(xviii) CE16:0, DG18:0, DHC18:1, DHC26:1, PL24:0, CE16:1n7, DG18:1n9, DHC20:0, FA14:1n5, SM18:0, CE18:1n9, DG20:0, DHC20:1, PL15:0, TG15:0, CE18:2n6, DG20:2n6, DHC22:1, PL18:2n6, TG20:3n9, CE20:4n6, DG20:3n9, DHC24:1, PL20:3n9, Total Diacylglycerol, CER18:0, DHC18:0, DHC26:0, PL20:4n6, Total Triacylglycerol, and CER18:1.

[0077] The term "determining the amount of each analyte" as used herein refers to determining at least one characteristic feature of at least one metabolite comprised by the sample referred to herein. Characteristic features in accordance with the present invention are features which characterize the physical and/or chemical properties including biochemical properties of a metabolite. Such properties include, e.g., molecular weight, viscosity, density, electrical charge, spin, optical activity, elementary composition, chemical structure, capability to react with other compounds, capability to elicit a response in a biological read out system (e.g., induction of a reporter gene) and the like. Values for said properties may serve as characteristic features and can be determined by techniques well known in the art. Moreover, the characteristic feature may be any feature which is derived from the values of the physical and/or chemical properties of a metabolite by standard operations, e.g., mathematical calculations such as multiplication, division, gradient boosting, generalized linear modeling, or logarithmic calculus. Most preferably, the at least one characteristic feature allows the determination and/or chemical identification of the said at least one metabolite.

[0078] The analytes comprised by a biological sample may be determined in accordance with the present invention quantitatively or qualitatively. For qualitative determination, the presence or absence of the metabolite will be determined by a suitable technique. Moreover, qualitative determination may, preferably, include determination of the chemical

structure or composition of the metabolite. For quantitative determination, either the precise amount of the analyte(s) present in the biological sample will be determined or the relative amount of the analyte(s) will be determined, preferably, based on the value determined for the characteristic feature(s) referred to herein above. The relative amount may be determined in a case where the precise amount of a metabolite can or shall not be determined. In said case, it can be determined whether the amount in which the analyte(s) is present is increased or decreased with respect to a second sample comprising said analyte(s) in a second amount. Quantitatively analyzing an analyte(s), thus, also includes what is sometimes referred to as semi-quantitative analysis of a metabolite.

[0079] Typically, the analyte level is determined by measuring the level of the metabolite in a body fluid (clinical sample), e.g., blood, serum, plasma, or urine. The level can be determined by, for example, mass spectrometry (MS), ELISA, immunoassays, enzymatic assays, spectrophotometry, colorimetry, fluorometry, bacterial assays, compound separation techniques, or other known techniques for determining the presence and/or quantity of an analyte.

[0080] Compound separation techniques yield a time resolved separation of the analytes comprised by the sample. Suitable techniques for separation to be used include, for example, all chromatographic separation techniques such as liquid chromatography (LC), high performance liquid chromatography (HPLC), gas chromatography (GC), thin layer chromatography, size exclusion or affinity chromatography. These techniques are well known in the art and can be applied by the person skilled in the art. In some embodiments, the methods utilize LC and/or GC chromatographic techniques including, for example, gas chromatography mass spectrometry (GC-MS), liquid chromatography mass spectrometry (LC-MS), liquid chromatography-tandem mass spectrometry (UPLC-MS/MS), direct infusion mass spectrometry or Fourier transform ion-cyclotron-resonance mass spectrometry (FT-ICR-MS), capillary electrophoresis mass spectrometry (CE-MS), high-performance liquid chromatography coupled mass spectrometry (HPLC-MS), quadrupole mass spectrometry, any sequentially coupled mass spectrometry, such as MS-MS or MS-MS-MS, inductively coupled plasma mass spectrometry (ICP-MS), pyrolysis mass spectrometry (Py-MS), ion mobility mass spectrometry or time of flight mass spectrometry (TOF). In some embodiments, LC-MS and/or GC-MS. As an alternative or in addition to mass spectrometry techniques, the following techniques may be used for compound determination: nuclear magnetic resonance (NMR), magnetic resonance imaging (MRI), Fourier transform infrared analysis (FT-IR), ultra violet (UV) spectroscopy, refraction index (RI), fluorescent detection, radiochemical detection, electrochemical detection, light scattering (LS), dispersive Raman spectroscopy or flame ionization detection (FID). These techniques are well known to the person skilled in the art and can be applied without further ado. In some embodiments, the methods disclosed herein shall be, optionally, assisted by automation. For example sample processing or pre-treatment can be automated by robotics. Data processing and comparison can be assisted by suitable computer programs and databases. Automation as described herein allows using the method of the present invention in high-throughput approaches.

[0081] "Determining" methods include, for example, sending a clinical sample(s) to a commercial laboratory for measurement or the use of commercially available assay kits. Commercially available assay kits are known in the art. For example, Quest Diagnostics, Sigma Aldrich, CATAChem Inc., Eton Bioscience Inc., and Bio Vision Research Products are exemplary suppliers of such assays. Exemplary kits and suppliers will be apparent to the skilled artisan.

[0082] In some cases, the methods disclosed herein involve comparing levels or occurrences to a reference. The reference can take on a variety of forms. In some cases, the reference comprises predetermined values for the plurality of analytes (e.g., each of the plurality of analytes). The predetermined value can take a variety of forms. It can be a level or occurrence of an analyte obtained from a subject known to have atherosclerotic CAD or to have a coronary atherosclerotic plaque (e.g., a symptomatic subject), or obtained from a subject known not to suffer from atherosclerotic CAD or known to not have a coronary atherosclerotic plaque (e.g., an asymptomatic subject). It can be a level or occurrence of an analyte obtained from a subject having no previous history of coronary artery disease. It can be a level or occurrence in the same subject, e.g., at a different time point. A predetermined value that represents a level(s) of an analyte is referred to herein as a predetermined level. A predetermined level can be single cut-off value, such as a median or mean. It can be a range of cut-off (or threshold) values, such as a confidence interval. It can be established based upon comparative groups, such as where the risk in one defined group is a fold higher, or lower, (e.g., approximately 2-fold, 4-fold, 8-fold, 16-fold or more) than the risk in another defined group. It can be a range, for example, where a population of subjects (e.g., control subjects) is divided equally (or unequally) into groups, such as a low-risk group, a medium-risk group and a high-risk group, or into quartiles, the lowest quartile being subjects with the lowest risk and the highest quartile being subjects with the highest risk, or into n-quartiles (i.e., n regularly spaced intervals) the lowest of the n-quartiles being subjects with the lowest risk and the highest of the n-quartiles being subjects with the highest risk. Moreover, the reference could be a calculated reference, most preferably the average or median, for the relative or absolute amount of an analyte of a population of individuals comprising the subject to be investigated. The absolute or relative amounts of the analytes of said individuals of the population can be determined as specified elsewhere herein. How to calculate a suitable reference value, preferably, the average or median, is well known in the art. The population of subjects referred to before shall comprise a plurality of subjects, preferably, at least 5, 10, 50, 100, 1,000 subjects. It is to be understood that the subject to be diagnosed by the method of the present invention and the subjects of the said plurality of subjects are of the same species.

[0083] Subjects associated with predetermined values are typically referred to as control subjects (or controls). A control subject does not have atherosclerosis or atherosclerotic CAD. In some cases it may be desirable that control subject is a symptomatic subject, and in other cases it may be desirable that a control subject is an asymptomatic subject. Thus, in some cases the level of an analyte in a subject being greater than or equal to the level of the analyte in a control subject is indicative of a clinical status (e.g., indicative of CAD diagnosis). In other cases the level of an analyte in a subject being less than or equal to the level of the analyte in a control subject is indicative of a clinical status. More generally, the combination of raised and lowered levels of a predefined group of analytes is indicative of clinical status. The amount of the greater than and the amount of the less than is usually of a sufficient magnitude to, for example, facilitate distinguishing a subject from a control subject using the disclosed methods. Typically, the greater than, or the less than, that is sufficient to distinguish a subject from a control subject is a statistically significant greater than, or a statistically significant less than. In cases where the level of an analyte in a subject being equal to the level of the metabolite in a control subject is indicative of a clinical status, the "being equal" refers to being approximately equal (e.g., not statistically different).

[0084] The predetermined value can depend upon a particular population of subjects (e.g., human subjects) selected. For example, an apparently healthy population will have a different 'normal' range of metabolites than will a population of subjects which have, or are likely to have, atherosclerotic CAD or a coronary atherosclerotic plaque. Accordingly, the predetermined values selected may take into account the category (e.g., healthy, at risk, diseased, age, gender, etc.) in which a subject (e.g., human subject) falls. Appropriate ranges and categories can be selected with no more than routine experimentation by those of ordinary skill in the art. In some cases a predetermined value of a biomarker is a value that is the average for a population of healthy subjects (human subjects) (e.g., human subjects who have no apparent signs and symptoms of CAD). The predetermined value will depend, of course, on the particular analyte (biomarker) selected and even upon the characteristics of the population in which the subject lies. In characterizing likelihood, or risk, numerous predetermined values can be established.

[0085] A level, in some embodiments, may itself be a relative level that reflects a comparison of levels between two states. Relative levels that reflect a comparison (e.g., ratio, difference, logarithmic difference, percentage change, etc.) between two states (e.g., healthy and diseased) may be referred to as delta values. The use of relative levels is beneficial in some cases because, to an extent, they exclude measurement related variations (e.g., laboratory personnel, laboratories, measurements devices, reagent lots/preparations, assay kits, etc.). However, the invention is not so limited.

[0086] Analyte levels and/or reference levels may be stored in a suitable data storage medium (e.g., a database) and are, thus, also available for future diagnoses. This also allows efficiently diagnosing prevalence for a disease because suitable reference results can be identified in the database once it has been confirmed (in the future) that the subject from which the corresponding reference sample was obtained did have CAD or a coronary atherosclerotic plaque. As used herein a "database" comprises data collected (e.g., analyte and/or reference level information and /or patient information) on a suitable storage medium. Moreover, the database, may further comprise a database management system. The database management system is, preferably, a network-based, hierarchical or object-oriented database management system. More preferably, the database will be implemented as a distributed (federal) system, e.g. as a Client-Server-System. More preferably, the database is structured as to allow a search algorithm to compare a test data set with the data sets comprised by the data collection. Specifically, by using such an algorithm, the database can be searched for similar or identical data sets being indicative of atherosclerotic CAD or a coronary atherosclerotic plaque (e.g. a query search). Thus, if an identical or similar data set can be identified in the data collection, the test data set will be associated with atherosclerotic CAD or coronary atherosclerotic plaques. Consequently, the information obtained from the data collection can be used to diagnose CAD or based on a test data set obtained from a subject. More preferably, the data collection comprises characteristic values of all analytes comprised by any one of the groups recited above.

[0087] In some embodiments, the methods disclosed herein further comprise modifying the subject's clinical record to identify the subject as having or as not having atherosclerotic coronary artery disease, or as having or as not having a coronary atherosclerotic plaque. The clinical record may be stored in any suitable data storage medium (e.g., a computer readable medium).

[0088] The invention also may provide kits for evaluating analyte biomarkers in a subject. The kits of the invention can take on a variety of forms. Typically, the kits will include reagents suitable for determining levels of a plurality of analytes biomarkers (e.g., those disclosed herein) in a sample. Optionally, the kits may contain one or more control samples. Typically, a comparison between the levels of the biomarkers in the subject and levels of the biomarkers in the control samples is indicative of a clinical status (e.g., diagnosis or likelihood of having atherosclerotic CAD or as having a coronary atherosclerotic plaque). Also, the kits, in some cases, will include written information (indicia) providing a reference (e.g., predetermined values), wherein a comparison between the levels of the biomarkers in the subject and the reference (predetermined values) is indicative of a clinical status. In some cases, the kits comprise software useful for comparing biomarker levels or occurrences with a reference (e.g., a prediction model). Usually the software will be provided in a computer readable format such as a compact disc, but it also may be available for downloading via the internet. However, the kits are not so limited and other variations with will be apparent to one of ordinary skill in the art.

[0089] The present methods can also be used for selecting a treatment and/or determining a treatment plan for a subject, based on the occurrence or levels of certain analytes relevant to CAD. In some examples, using the method disclosed herein, a health care provider (e.g., a physician) identifies a subject as having or at risk of having for having atherosclerotic CAD or as having a coronary atherosclerotic plaque and, based on this identification the health care provider determines an adequate management plan for the subject. In some examples, using the method disclosed herein, a health care provider (e.g., a physician) diagnoses a subject as having a atherosclerotic CAD or as having a coronary atherosclerotic plaque based on the occurrence or levels of certain analytes in a clinical sample obtained from the subject, and/or based on a classification of a clinical sample obtained from the subject. By way of this diagnosis the health care provider determines an adequate treatment or treatment plan for the subject. In some embodiments, compositions for use in methods of treating atherosclerotic coronary artery disease further include administering the treatment to the subject.

[0090] In some embodiments, the invention relates to identifying subjects who are likely to have successful treatment with a particular drug dose, formulation and/or administration modality. Other embodiments include evaluating the efficacy of a drug using the metabolomic profiling methods of the present invention. In some embodiments, the metabolomic profiling methods are useful for identifying subjects who are likely to have successful treatment with a particular drug or therapeutic regiment. For example, during a study (e.g., a clinical study) of a drug or treatment, subjects who have a CAD or coronary atherosclerotic plaque may respond well to the drug or treatment, and others may not. Disparity in treatment efficacy is associated with numerous variables, for example genetic variations among the subjects. In some embodiments, subjects in a population are stratified based on the metabolomic profiling methods disclosed herein. In some embodiments, resulting strata are further evaluated based on various epidemiological, and or clinical factors (e.g., response to a specific treatment). In some embodiments, stratum, identified based on a metabolic profile, reflect a subpopulation of subjects that response predictably (e.g., have a predetermined response) to certain treatments. In further embodiments, samples are obtained from subjects who have been subjected to the drug being tested and who have a predetermined response to the treatment. In some cases, a reference can be established from all or a portion of the analytes from these samples, for example, to provide a reference metabolic profile. A sample to be tested can then be evaluated (e.g., using a prediction model) against the reference and classified on the basis of whether treatment would be successful or unsuccessful. A company and/or person testing a treatment (e.g., compound, drug, and life-style change) could discern more accurate information regarding the types or subtypes of CAD for which a treatment is most useful. This information also aids a healthcare provider in determining the best treatment plan for a subject.

[0091] In some examples, treatment for the atherosclerotic CAD or coronary atherosclerotic plaques is to administer to the subject a composition comprising an effective amount of at therapeutic agent and/or to instruct the subject to adopt at least one therapeutic lifestyle change (e.g., change in diet or exercise). Therapeutic compounds suitable for treating CAD or coronary atherosclerotic plaques are well known in the art and some are disclosed herein. Non-limiting examples include statins, cholesterol absorption inhibitors, niacin-derivatives, omega-3-fatty acid compounds, bile acid sequestrants, PCSK9 antagonists, anti-platelet agents and aspirin. Appropriate lifestyle changes to improve cardiovascular health are also well known in the art. Non-limiting examples include increased physical activity, caloric intake restriction, nutritional meal planning, and weight reduction. However, the invention is not so limited and other appropriate treatments will be apparent to one of ordinary skill in the art.

[0092] When a therapeutic agent or other treatment is administered, it is administered in an amount effective to treat CAD or reduce the likelihood (or risk) of future CAD or CAD events. An effective amount is a dosage of the therapeutic agent sufficient to provide a medically desirable result. The effective amount will vary with the particular condition being treated, the age and physical condition of the subject being treated, the severity of the condition, the duration of the treatment, the nature of the concurrent therapy (if any), the specific route of administration and the like factors within the knowledge and expertise of the health care practitioner. For example, an effective amount can depend upon the degree to which a subject has abnormal levels of certain analytes (e.g., analytes as described herein) that are indicative of atherosclerotic CAD or a coronary atherosclerotic plaque. It should be understood that the therapeutic agents of the invention are used to treat and/or prevent atherosclerotic CAD or a coronary atherosclerotic plaque. Thus, in some cases, they may be used prophylactically in human subjects at risk of developing an atherosclerotic CAD or a coronary atherosclerotic plaque. Thus, in some cases, an effective amount is that amount which can lower the risk of, slow or perhaps prevent altogether the development of atherosclerotic CAD or a coronary atherosclerotic plaque. It will be recognized when the therapeutic agent is used in acute circumstances, it is used to prevent one or more medically undesirable results that typically flow from such adverse events. Methods for selecting a suitable treatment and an appropriate dose thereof will be apparent to one of ordinary skill in the art.

[0093] The invention further provides for the communication of assay results or diagnoses or both to technicians, physicians or patients, for example. In certain embodiments, computers will be used to communicate assay results or diagnoses or both to interested parties, e.g., physicians and their patients. In some instances, the methods further comprise providing communicating the patient status (i.e. as having or not having ASCAD or the presence or absence of a coronary atherosclerotic plaque) as a report. Thus, in some instances, the subject methods may further include a

step of generating or outputting a report providing the results of the subject methods, which report can be provided in the form of an electronic medium (e.g., an electronic display on a computer monitor), or in the form of a tangible medium (e.g., a report printed on paper or other tangible medium). Any form of report may be provided, e.g. as known in the art. In some embodiments of the invention, a diagnosis based on the presence or absence in a test subject of any the biomarkers of Tables 30, 32, 34, 36, 38, 40, 42, 44, 46, 48 or 50 is communicated to the subject as soon as possible after the diagnosis is obtained. The diagnosis may be communicated to the subject by the subject's treating physician. Alternatively, the diagnosis may be sent to a test subject by email or communicated to the subject by phone. The diagnosis may be sent to a test subject by in the form of a report. A computer may be used to communicate the diagnosis by email or phone. In certain embodiments, the message containing results of a diagnostic test may be generated and delivered automatically to the subject using a combination of computer hardware and software which will be familiar to artisans skilled in telecommunications.

[0094] A "report," as described herein, is an electronic or tangible document which includes report elements that provide information of interest relating to a diagnosis assessment, a prognosis assessment, a treatment assessment, a monitoring assessment, etc. and its results. A subject report can be completely or partially electronically generated. A subject report includes at least a subject assessment, e.g., a diagnosis as to whether a subject has a high likelihood of having ASCAD or as having a coronary atherosclerotic plaque; or a prognosis assessment, e.g. a prediction of the responsiveness of a patient to therapy; and/or a suggested course of treatment to be followed. A subject report can further include one or more of: 1) information regarding the testing facility; 2) service provider information; 3) subject data; 4) sample data; 5) an assessment report, which can include various information including: a) test data, where test data can include i) the biomarker levels of one or more biomarkers; and/or ii) the biomarker signatures for one or more panel of biomarkers described herein.

[0095] The report may include information about the testing facility, which information is relevant to the hospital, clinic, or laboratory in which sample gathering and/or data generation was conducted. This information can include one or more details relating to, for example, the name and location of the testing facility, the identity of the lab technician who conducted the assay and/or who entered the input data, the date and time the assay was conducted and/or analyzed, the location where the sample and/or result data is stored, the lot number of the reagents (e.g., kit, etc.) used in the assay, and the like. Report fields with this information can generally be populated using information provided by the user.

[0096] The report may include information about the service provider, which may be located outside the healthcare facility at which the user is located, or within the healthcare facility. Examples of such information can include the name and location of the service provider, the name of the reviewer, and where necessary or desired the name of the individual who conducted sample gathering and/or data generation. Report fields with this information can generally be populated using data entered by the user, which can be selected from among pre-scripted selections (e.g., using a drop-down menu). Other service provider information in the report can include contact information for technical information about the result and/or about the interpretive report.

[0097] The report may include a subject data section, including subject medical history as well as administrative subject data (that is, data that are not essential to the diagnosis, prognosis, or treatment assessment) such as information to identify the subject (e.g., name, subject date of birth (DOB), gender, mailing and/or residence address, medical record number (MRN), room and/or bed number in a healthcare facility), insurance information, and the like), the name of the subject's physician or other health professional who ordered the susceptibility prediction and, if different from the ordering physician, the name of a staff physician who is responsible for the subject's care (e.g., primary care physician).

[0098] The report may include a sample data section, which may provide information about the biological sample analyzed, such as the source of biological sample obtained from the subject (e.g. blood, e.g., whole blood, fractionated blood, plasma, serum, etc.), how the sample was handled (e.g. storage temperature, preparatory protocols) and the date and time collected. Report fields with this information can generally be populated using data entered by the user, some of which may be provided as pre-scripted selections (e.g., using a drop-down menu).

[0099] It will also be readily appreciated that the reports can include additional elements or modified elements. For example, where electronic, the report can contain hyperlinks which point to internal or external databases which provide more detailed information about selected elements of the report. For example, the patient data element of the report can include a hyperlink to an electronic patient record, or a site for accessing such a patient record, which patient record is maintained in a confidential database. This latter embodiment may be of interest in an in-hospital system or in-clinic setting. When in electronic format, the report is recorded on a suitable physical medium, such as a computer readable medium, e.g., in a computer memory, zip drive, CD, DVD, flash drive, etc.

[0100] It will be readily appreciated that the report can include all or some of the elements above, with the proviso that the report generally includes at least the elements sufficient to provide the analysis requested by the user (e.g., a diagnosis, a prognosis, or a prediction of responsiveness to a therapy).

[0101] The methods described herein may be used on samples collected from patients in a clinical trial and the results of the test used in conjunction with patient outcomes in order to determine whether subgroups of patients are more or less likely to show a response to a new drug than the whole group or other subgroups. Further, such methods can be

used to identify from clinical data the subsets of patients who can benefit from therapy. Additionally, a patient is more likely to be included in a clinical trial if the results of the test indicate a higher likelihood that the patient will be responsive to medical treatment, and a patient is less likely to be included in a clinical trial if the results of the test indicate a lower likelihood that the patient will be responsive to medical treatment.

[0102] The methods described herein can be used alone or in combination with other clinical methods for patient stratification known in the art to provide a diagnosis, a prognosis, or a prediction of responsiveness to therapy. For example, clinical parameters that are known in the art for diagnosing ASCAD may be incorporated into the ordinarily skilled artisan's analysis to arrive at an ovarian cancer assessment with the subject methods.

EXAMPLES

[0103] The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

[0104] The Genetic Loci and the Burden of Atherosclerotic Lesions study (GLOBAL; NCT01738828) is an international multi-center, prospective study, which enrolled 7526 patients (approximately half of whom are atherosclerotic CAD cases; the other half are without atherosclerotic CAD) who were referred for coronary CT angiography, for assessment of suspected coronary artery disease (CAD) at a total of 48 clinical sites. Previous genetic and genomic studies of CAD have used patient history of CAD or MI (myocardial infarction), or invasive coronary angiography in order to determine case/ control status. In the GLOBAL study, cardiovascular CT is utilized, including non-contrast-enhanced coronary computerized tomography (CT), also known as coronary artery calcium (CAC) scoring, and secondly contrast-enhanced CT angiography (CTA). The increased sensitivity of this imaging technique for the detection of coronary plaque has been shown to allow the re-classification of controls as cases (sponsor's own data).

[0105] The following Examples describe statistical analyses of the pilot GLOBAL data, consisting of metabolomic data for 1096 patients from a total of 26 clinical sites. A subset of 748 patients made up the Discovery Set; a further 348 participants made up the Validation Set. The aims of the GLOBAL Pilot Discovery cohort were to assess preliminary detection of strong biomarker-associations for atherosclerosis, in patients phenotyped using advanced cardiac CT-based imaging.

[0106] Patients presenting for a CTA, for an assessment of possible CAD, were identified as potential study candidates and were screened for eligibility according to the study inclusion and exclusion criteria. Subjects were assessed for cardiovascular disease risk factors including gender, age, hypertension, hyperlipidemia, diabetes, and smoking, as well as a number of other conditions, and signed the appropriate Institutional Review Board-approved informed consent form. The general inclusion criteria were as follows (1) Ages 18-90; (2) White and of Non-Hispanic or Non-Latino origin; and (3) Referred for coronary CT angiography to evaluate presence of CAD. The general exclusion criteria were as follows: (1) Use of immunosuppressive or immunomodulatory therapy including any dose of systemic corticosteroids in the preceding 30 days (except if steroids were administered as pre-medication prior to contrast administration for CT scan within 24 hours); (2) Chemotherapy in the preceding year; (3) Major surgery in the preceding 2 months; (4) Blood or blood product transfusion in the preceding 2 months; (5) Subjects for whom coronary CT angiography is contra-indicated per institutional standard of care; (6) Subjects with previous coronary arterial revascularization (percutaneous coronary intervention (PCI) or coronary artery bypass graft (CABG)); (7) Subjects with atrial fibrillation/flutter or frequent irregular or rapid heart rhythms, which occurred within the past 3 months; (8) Subjects with a pacemaker or implantable cardioverter-defibrillator implant; (9) Active congestive heart failure or the presence of known non-ischemic cardiomyopathy; and (10) Known genetic disorders of atherosclerosis, lipid or lipoprotein metabolism. The subjects must have met all of the general inclusion criteria to be enrolled in the study. If any of the general exclusion criteria were met, the subject was excluded from the study.

[0107] Each patient was characterized in terms of biomarkers by the application of the following methods: whole genome sequencing (WGS), whole genome methylation (WGM), whole blood based transcriptome sequencing (WTS), unbiased proteomics, unbiased metabolomics, unbiased lipidomics and lipoprotein proteomics. In addition, more conventional biomarkers were measured.

Sample Preparation: Metabolomics:

[0108] Samples were stored at -70°C until processed. Sample preparation was carried out as described previously. (Evans AM, et. al., Integrated, nontargeted ultrahigh performance liquid chromatography/electrospray ionization tandem mass spectrometry platform for the identification and relative quantification of the small-molecule complement of biological systems. *Anal Chem* 2009;81(16):6656-6667) Briefly, recovery standards were added prior to the first step in the extraction process for quality control purposes. To remove protein, dissociate small molecules bound to protein or trapped in the precipitated protein matrix, and to recover chemically diverse metabolites, proteins were precipitated with methanol under vigorous shaking for 2 min (Glen Mills Genogrinder 2000) followed by centrifugation. The resulting extract was

divided into four fractions: one for analysis by ultra high performance liquid chromatography-tandem mass spectrometry (UPLC-MS/MS; positive mode), one for analysis by UPLC-MS/MS (negative mode), one for analysis by gas chromatography-mass spectrometry (GC-MS), and one sample is reserved for backup. Three types of controls were analyzed in concert with the experimental samples: samples generated from a pool of human plasma (extensively characterized by Metabolon, Inc.) served as technical replicate throughout the data set; extracted water samples served as process blanks; and a cocktail of standards spiked into every analyzed sample allowed instrument performance monitoring. Instrument variability was determined by calculating the median relative standard deviation (RSD) for the standards that are added to each sample prior to injection into the mass spectrometers (median RSD = 5%; n = 30 standards). Overall process variability was determined by calculating the median RSD for all endogenous metabolites (i.e., non-instrument standards) present in 100% of the pooled human plasma samples (median RSD = 11%; n = 610 metabolites). Experimental samples and controls were randomized across the platform run.

[0109] Mass Spectrometry Analysis. Non-targeted MS analysis was performed at Metabolon, Inc. Extracts were subjected to either GC-MS or UPLC-MS/MS⁷. The chromatography was standardized and, once the method was validated no further changes were made. As part of Metabolon's general practice, all columns are purchased from a single manufacturer's lot at the outset of experiments. All solvents were similarly purchased in bulk from a single manufacturer's lot in sufficient quantity to complete all related experiments. For each sample, vacuum-dried samples were dissolved in injection solvent containing eight or more injection standards at fixed concentrations, depending on the platform. The internal standards were used both to assure injection and chromatographic consistency. Instruments were tuned and calibrated for mass resolution and mass accuracy daily. The UPLC-MS/MS platform utilized a Waters Acquity UPLC with Waters UPLC BEH C18-2.1×100 mm, 1.7 μm columns and a Thermo Scientific Q-Exactive high resolution/accurate mass spectrometer interfaced with a heated electrospray ionization (HESI-II) source and Orbitrap mass analyzer operated at 35,000 mass resolution. The sample extract was dried then reconstituted in acidic or basic LC-compatible solvents, each of which contained eight or more injection standards at fixed concentrations to ensure injection and chromatographic consistency. One aliquot was analyzed using acidic, positive ion-optimized conditions and the other using basic, negative ion-optimized conditions in two independent injections using separate dedicated columns. Extracts reconstituted in acidic conditions were gradient eluted using water and methanol containing 0.1% formic acid, while the basic extracts, which also used water/methanol, contained 6.5mM ammonium bicarbonate. The MS analysis alternated between MS and data-dependent MS2 scans using dynamic exclusion, and the scan range was from 80-1000 m/z.

[0110] The samples destined for analysis by GC-MS were dried under vacuum desiccation for a minimum of 18 hours prior to being derivatized under dried nitrogen using bistrimethyl-silyltrifluoroacetamide. Derivatized samples were separated on a 5% phenyldimethyl silicone column with helium as carrier gas and a temperature ramp from 60° to 340°C within a 17-min period. All samples were analyzed on a Thermo-Finnigan Trace DSQ MS operated at unit mass resolving power with electron impact ionization and a 50-750 atomic mass unit scan range.

[0111] Compound Identification, Quantification, and Data Curation. Metabolites were identified by automated comparison of the ion features in the experimental samples to a reference library of chemical standard entries that included retention time, molecular weight (m/z), preferred adducts, and in-source fragments as well as associated MS spectra and curated by visual inspection for quality control using software developed at Metabolon. (DeHaven CD, et. al., Organization of GC/MS and LC/MS metabolomics data into chemical libraries. J Cheminform 2010;2(1):9) Identification of known chemical entities was based on comparison to metabolomic library entries of purified standards. Over 2,500 commercially available purified standard compounds have been acquired and registered into LIMS for distribution to both the LC/MS and GC/MS platforms for determination of their detectable characteristics. An additional 250 mass spectral entries have been created for structurally unnamed biochemicals, which have been identified by virtue of their recurrent nature (both chromatographic and mass spectral). These compounds have the potential to be identified by future acquisition of a matching purified standard or by classical structural analysis. Peaks were quantified using area-under-the-curve. Raw area counts for each metabolite in each sample were normalized to correct for variation resulting from instrument inter-day tuning differences by the median value for each run-day, therefore, setting the medians to 1.0 for each run. This preserved variation between samples but allowed metabolites of widely different raw peak areas to be compared on a similar graphical scale. Missing values are imputed with the observed minimum after normalization.

Sample Preparation: Lipidomics

[0112] TrueMass® Lipomic Panel. Lipids were extracted in the presence of authentic internal standards by the method of Folch et al. using chloroform:methanol (2:1 v/v). For the separation of neutral lipid classes [FFA, TAG, DAG, CE], a solvent system consisting of petroleum ether/diethyl ether/acetic acid (80:20:1) was employed. Individual phospholipid classes within each extract [PC, PE] were separated using the Agilent Technologies 1100 Series LC. Each lipid class was transesterified in 1% sulfuric acid in methanol in a sealed vial under a nitrogen atmosphere at 100°C for 45 minutes. The resulting fatty acid methyl esters were extracted from the mixture with hexane containing 0.05% butylated hydroxytoluene and prepared for GC by sealing the hexane extracts under nitrogen. Fatty acid methyl esters were separated

and quantified by capillary GC (Agilent Technologies 6890 Series GC) equipped with a 30 m DB 88 capillary column (Agilent Technologies) and a flame ionization detector.

[0113] *TrueMass® Ceramides Panel.* Deuterium-labelled internal standards were added to the samples and the mixture was solubilized in methanol followed by a crash extraction. A bilayer was formed with the addition of KCl in water, and the organic layer is removed and concentrated under nitrogen. The extract was spun, filtered, and split into 2 injections - one for ceramides and one for sphingosines. The extract was injected onto an Agilent C8 column connected to an Agilent 1290 Infinity LC and ABI 4000 QTRAP. The analytes were ionized via positive electrospray and the mass spectrometer was operated in the tandem MS mode. The absolute concentration of each sphingolipid was determined by comparing the peak to that of the relevant internal standard.

[0114] *Mass Spectrometry Analysis.* Non-targeted MS analysis was performed at Metabolon, Inc. Extracts were subjected to either GC-MS or UPLC-MS/MS⁷. The chromatography was standardized and, once the method was validated no further changes were made. As part of Metabolon's general practice, all columns are purchased from a single manufacturer's lot at the outset of experiments. All solvents were similarly purchased in bulk from a single manufacturer's lot in sufficient quantity to complete all related experiments. For each sample, vacuum-dried samples were dissolved in injection solvent containing eight or more injection standards at fixed concentrations, depending on the platform. The internal standards were used both to assure injection and chromatographic consistency. Instruments were tuned and calibrated for mass resolution and mass accuracy daily. The UPLC-MS/MS platform utilized a Waters Acquity UPLC with Waters UPLC BEH C18-2.1×100 mm, 1.7 µm columns and a Thermo Scientific Q-Exactive high resolution/accurate mass spectrometer interfaced with a heated electrospray ionization (HESI-II) source and Orbitrap mass analyzer operated at 35,000 mass resolution. The sample extract was dried then reconstituted in acidic or basic LC-compatible solvents, each of which contained eight or more injection standards at fixed concentrations to ensure injection and chromatographic consistency. One aliquot was analyzed using acidic, positive ion-optimized conditions and the other using basic, negative ion-optimized conditions in two independent injections using separate dedicated columns. Extracts reconstituted in acidic conditions were gradient eluted using water and methanol containing 0.1% formic acid, while the basic extracts, which also used water/methanol, contained 6.5mM ammonium bicarbonate. The MS analysis alternated between MS and data-dependent MS2 scans using dynamic exclusion, and the scan range was from 80-1000 m/z.

[0115] The samples destined for analysis by GC-MS were dried under vacuum desiccation for a minimum of 18 hours prior to being derivatized under dried nitrogen using bistrimethyl-silyltrifluoroacetamide. Derivatized samples were separated on a 5% phenyldimethyl silicone column with helium as carrier gas and a temperature ramp from 60° to 340°C within a 17-min period. All samples were analyzed on a Thermo-Finnigan Trace DSQ MS operated at unit mass resolving power with electron impact ionization and a 50-750 atomic mass unit scan range.

[0116] *Compound Identification, Quantification, and Data Curation.* Metabolites were identified by automated comparison of the ion features in the experimental samples to a reference library of chemical standard entries that included retention time, molecular weight (m/z), preferred adducts, and in-source fragments as well as associated MS spectra and curated by visual inspection for quality control using software developed at Metabolon. (DeHaven et al.) Identification of known chemical entities was based on comparison to metabolomic library entries of purified standards. Over 2,500 commercially available purified standard compounds have been acquired and registered into LIMS for distribution to both the LC/MS and GC/MS platforms for determination of their detectable characteristics. An additional 250 mass spectral entries have been created for structurally unnamed biochemicals, which have been identified by virtue of their recurrent nature (both chromatographic and mass spectral). These compounds have the potential to be identified by future acquisition of a matching purified standard or by classical structural analysis. Peaks were quantified using area-under-the-curve. Raw area counts for each metabolite in each sample were normalized to correct for variation resulting from instrument inter-day tuning differences by the median value for each run-day, therefore, setting the medians to 1.0 for each run. This preserved variation between samples but allowed metabolites of widely different raw peak areas to be compared on a similar graphical scale. Missing values are imputed with the observed minimum after normalization.

Example 1. Discovery

Analysis Populations

[0117] The "Full Analysis Set" ("FAS") consisted of pilot study patients with clinical data and a CT-based designation of either Revascularization CAD case, Native CAD case, or Control (N = 336 for the FAS group).

[0118] The "Native CAD Set" was the subset of the FAS with Native CAD as verified by CT, who had analyte (metabolomic and lipidomic) data (N = 120 for the Native CAD Set). These were subjects without previous revascularization procedures, such as percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG).

[0119] The "Revasc CAD Set" was the subset of the FAS who had undergone previous revascularization, such as percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG), and who had analyte data.

[0120] The "All CAD Set" was the union of the Native CAD Set and the Revasc CAD Set.

[0121] The "Control Set" was the subset of FAS who had a calcium score of zero and were designated a Control after inspection of CT data, and who had analyte data. (N = 279 for the Control Set plus Native CAD Set). These subjects had no discernable atherosclerotic CAD based on the combined inspection of the non-contrast CT and the contrast-enhanced CT data.

[0122] It is noted that by design, the only racial group represented in the study was White. Therefore, race-based sub-populations were not defined. The data was not split into training and validation sets.

I: GLOBAL Pilot Study

A. Study Endpoints

[0123] For the GLOBAL Pilot Discovery Cohort, there were two primary endpoints in the analysis: (1) Native CAD; and (2) All CAD (Native or Revascularization). All analyses were applied to both primary endpoints.

B. Statistical Hypothesis

[0124] The null hypothesis of no association, between the metabolite or lipid and the endpoint, was tested against the two-sided alternative that association exists.

C. Multiple Comparisons and Multiplicity

[0125] False discovery rate (FDR) q-values were calculated (Benjamini and Hochberg, 1995). Associations with FDR $q < 0.05$ were considered preliminary associations. In some circumstances, test results with raw $p < 0.05$ were reported as well.

D. Missing Data

[0126] Endpoint data was not imputed. Potential covariates with more than 5% missing data were excluded. Potential covariates with less than 5% missing data were imputed to the mean.

[0127] Metabolites with more than 10% missing data were excluded from the main analyses. Missing values for metabolites and lipids with less than 10% missing were imputed to the observed minimum after normalization.

E. Covariates

[0128] Twelve potential covariates were referred to as "clinical variables": Age; Gender; Body Mass Index (BMI); Type II Diabetes; Current smoker; Fasting; Statin use; Niacin use; Fibrate use; Ezetimibe use; Fish oil use; and Bile acid sequestrant use.

[0129] A further set of seventy-seven potential covariates, listed in Table 1, were referred to as "conventional blood-based biomarker variables".

Table 1:

Serum Test Name	Full name
Adiponectin	Adiponectin
ALB	Albumin
ALP	Alkaline phosphatase
ALT/GPT	Alanine transaminase
APO A1	Apolipoprotein A-I
APO B	Apolipoprotein B
AST/GOT	Aspartate transaminase
Calcium	Calcium
C-PEP	C-Peptide
CREAT	Creatinine

EP 3 242 588 B1

(continued)

Serum Test Name	Full name
DBILI	Direct Bilirubin
GGT	Gamma-glutamyl transferase
Glucose	Glucose
HCY	Homocysteine
HDL3	High Density Lipoprotein Fraction 3
HDL-C	High Density Lipoprotein Cholesterol
hsCRP	High Sensitive C Reactive Protein
Insulin	Insulin
LDH	Lactate dehydrogenase
LDL-C	Low Density Lipoprotein Cholesterol
LP(a) Mass	Lipoprotein (a) mass
LP(a)-C	Lipoprotein (a) cholesterol
LP(a)-P	Lipoprotein (a) particles number
Lp-PLA2	Lipoprotein-associated phospholipase A2
NEFA	Fatty acids
NTPROBNP	N-terminal pro-brain natriuretic peptide
OCN	Osteocalcin
PHOS	Phosphorous
PTH	Parathyroid hormone
sdLDL	Small dense Low Density Lipoprotein Cholesterol
TBILI	Total Bilirubin
Total Cholesterol	Total Cholesterol
TP	Total Protein
TRIG	Triglycerides
Uric Acid	Uric Acid
VIT D	Vitamin D
Plasma Test name	Full name
B-Sitosterol	B-Sitosterol
Campesterol	Campesterol
Cholestanol	Cholestanol
Desmosterol	Desmosterol
Leptin	Leptin
MPO	Myeloperoxidase
ProInsulin	ProInsulin
Whole Blood Test name	Full name
AA2	arachidonic acid
AAEPA	arachidonic divided by EPA
alphalin2	alphalinoleic acid

(continued)

Whole Blood Test name	Full name
arachidic2	arachidic acid
behenic2	behenic acid
cismontotl	palmitoleic + oleic + eicosenoic + nervonic
dcopentn32	docosapentaenoic-n3 acid
dcopentn62	docosapentaenoic-n6 acid
DHA2	docosahexaenoic acid
dihomolin2	dihomolinoleic acid
docosat2	docosatetraenoic acid
eicosad2	eicosadienoic acid
eicosen2	eicosenoic acid
EPA2	eicosapentaenoic acid
gammalin2	gamma linoleic acid
HbA1C	Glycated hemoglobin A1C
ligno2	lignoceric acid
linoleic2	linoleic acid
myristic2	myristic acid
nervonic2	nervonic acid
O3Index	Omega 3 index
O3total	Omega 3 total
O6O3	O6 total divided by O3 total
O6total	Omega 6 total
oleic2	oleic acid
palmitic2	palmitic acid
palmleic2	palmitoleic acid
satttotal	myristic + palmitic + behenic + lignoceric + arachadic + stearic
stearic2	stearic acid
tomegaprct	Total fatty acids
translin2	translinoleic acid
transol2	transoleic acid
transpalm2	transpalmitoleic acid
transtotal	trans palmitoleic + trans oleic + trans linoleic

[0130] As described below, a selected subset of these variables were used as covariates in the covariate-adjusted association analysis and/or as inputs for the prediction modeling analyses.

F. Analysis of Subgroups

[0131] The first primary endpoint was addressed using a subset of the FAS. Specifically the *Native CAD Set* was compared to the *Control Set*. The majority of the secondary analyses involved the *Native CAD Set* only. Additional subgroups were considered during the analysis of secondary endpoints arising from CT.

II. Demographic and Baseline Characteristics

[0132] The baseline and demographic characteristics of patients in the pilot study were tabulated. Continuous variables were summarized by the mean and standard error; binary variables were summarized by the count and percentage.

[0133] Table 2 shows general patient characteristics by clinical group (Revasc CAD vs. Native CAD vs. Control). A Kruskal-Wallis test was performed to investigate homogeneity of continuous measures; a Pearson's chi-squared test was conducted for binary measures; unadjusted p-values are reported.

[0134] Table 3 shows general patient characteristics by primary endpoint specifically, Native CAD vs. Control, and All CAD vs. Control. A Mann-Whitney test was performed to investigate homogeneity of continuous measures; a Pearson's chi-squared test was conducted for binary measures; unadjusted p-values are reported.

Table 2:

	All Controls	Native CAD	Revasc CAD	P-value
N	160	120	58	
Age				
mean (SE)	54.79 (0.79)	55.12 (0.87)	58.4 (1.22)	0.5768
SBP				
mean (SE)	131.26 (1.36)	134.86 (1.59)	137.88 (2.81)	0.2157
DBP				
mean (SE)	80.19 (0.77)	81.92 (1.07)	80.19 (1.34)	0.2659
Male				
N (%)	71 (44.38)	51 (42.50)	36 (62.07)	0.0351
Hypertension				
N (%)	85 (53.12)	82 (68.33)	51 (89.47)	3.06e-06
Dyslipidemia				
N (%)	80 (50.63)	85 (71.43)	55 (96.49)	8.7e-10
Diabetes (Any)				
N (%)	9 (5.62)	19 (15.83)	19 (32.76)	1.55e-06
Type I Diabetes				
N (%)	0 (0.00)	1 (0.83)	0 (0.00)	0.4021
Type II Diabetes				
N (%)	9 (5.62)	18 (15.00)	19 (32.76)	1.4e-06
Current Smoker				
N (%)	17 (10.62)	25 (20.83)	8 (13.79)	0.0571
Former Smoker				
N (%)	36 (22.50)	42 (35.00)	33 (56.90)	9.04e-06
Chest Pain				
N (%)	109 (68.12)	78 (65.00)	40 (68.97)	0.8158
Angina Equivalent				
N (%)	67 (41.88)	48 (40.00)	31 (53.45)	0.2125
Shortness of Breath				
N (%)	42 (26.25)	28 (23.33)	17 (29.31)	0.6797

(continued)

Family History of CAD				
N (%)	80 (50.00)	72 (60.00)	37 (63.79)	0.1031
Fasting				
N (%)	40 (25.00)	36 (30.00)	22 (37.93)	0.1696
Statin				
N (%)	49 (30.63)	59 (49.17)	51 (87.93)	5.52e-13
Niacin				
N (%)	5 (3.12)	1 (0.83)	2 (3.45)	0.3843
Fibrate				
N (%)	4 (2.50)	7 (5.83)	5 (8.62)	0.1330
Ezetimibe				
N (%)	4 (2.50)	3 (2.50)	5 (8.62)	0.0722
Fish Oil				
N (%)	18 (11.25)	23 (19.17)	6 (10.34)	0.1147
Bile Acid Sequestrant				
N (%)	3 (1.88)	0 (0.00)	0 (0.00)	0.1857
Aspirin				
N (%)	59 (36.88)	52 (43.33)	47 (81.03)	3.72e-08
Clopidogrel				
N (%)	3 (1.88)	4 (3.33)	28 (48.28)	2.6e-24
Vitamin K Antagonist				
	All Controls	Native CAD	Revasc CAD	P-value
N (%)	4 (2.50)	7 (5.83)	6 (10.34)	0.0568
Nitrate				
N (%)	4 (2.50)	4 (3.33)	13 (22.41)	1.36e-07
Beta Blocker				
N (%)	51 (31.87)	46 (38.33)	49 (84.48)	1.53e-11
ACE Inhibitor				
N (%)	32 (20.00)	35 (29.17)	32 (55.17)	3.01e-06

Table 3:

	All CAD	All Controls	P-value	Native CAD	All Controls	P-value
N	178	160		120	160	
Age						
mean (SE)	56.19 (0.72)	54.79 (0.79)	0.1968	55.12 (0.87)	54.79 (0.79)	0.7808
SBP						
mean (SE)	135.84 (1.41)	131.26 (1.36)	0.0273	134.86 (1.59)	131.26 (1.36)	0.0707

EP 3 242 588 B1

(continued)

5	DBP						
	mean (SE)	81.36 (0.84)	80.19 (0.77)	0.4625	81.92 (1.07)	80.19 (0.77)	0.2828
10	Male						
	N (%)	87 (48.88)	71 (44.38)	0.4722	51 (42.50)	71 (44.38)	0.8482
15	Hypertension						
	N (%)	133 (75.14)	85 (53.12)	3.98e-05	82 (68.33)	85 (53.12)	0.0145
20	Dyslipidemia						
	N (%)	140 (79.55)	80 (50.63)	5.08e-08	85 (71.43)	80 (50.63)	8.00e-04
25	Diabetes (Any)						
	N (%)	38 (21.35)	9 (5.62)	5.97e-05	19 (15.83)	9 (5.62)	0.0089
30	Type I Diabetes						
	N (%)	1 (0.56)	0 (0.00)	1.0000	1 (0.83)	0 (0.00)	0.8850
35	Type II Diabetes						
	N (%)	37 (20.79)	9 (5.62)	9.62e-05	18 (15.00)	9 (5.62)	0.0153
40	Current Smoker						
	N (%)	33 (18.54)	17 (10.62)	0.0584	25 (20.83)	17 (10.62)	0.0279
45	Former Smoker						
	N (%)	75 (42.13)	36 (22.50)	2.00e-04	42 (35.00)	36 (22.50)	0.0297
50	Chest Pain						
	N (%)	118 (66.29)	109 (68.12)	0.8086	78 (65.00)	109 (68.12)	0.6736
55	Angina Equivalent						
	N (%)	79 (44.38)	67 (41.88)	0.7229	48 (40.00)	67 (41.88)	0.8471
	Shortness of Breath						
	N (%)	45 (25.28)	42 (26.25)	0.9371	28 (23.33)	42 (26.25)	0.6757
	Family History of CAD						
	N (%)	109 (61.24)	80 (50.00)	0.0491	72 (60.00)	80 (50.00)	0.1233
	Fasting						
	N (%)	58 (32.58)	40 (25.00)	0.1573	36 (30.00)	40 (25.00)	0.4265
	Statin						
	N (%)	110 (61.80)	49 (30.63)	1.87e-08	59 (49.17)	49 (30.63)	0.0024
	Niacin						
	N (%)	3 (1.69)	5 (3.12)	0.6094	1 (0.83)	5 (3.12)	0.3716
	Fibrate						
	N (%)	12 (6.74)	4 (2.50)	0.1148	7 (5.83)	4 (2.50)	0.2670
	Ezetimibe						
	N (%)	8 (4.49)	4 (2.50)	0.4871	3 (2.50)	4 (2.50)	1.0000
	Fish Oil						
	N (%)	29 (16.29)	18 (11.25)	0.2379	23 (19.17)	18 (11.25)	0.0923

(continued)

Bile Acid Sequestrant						
N (%)	0 (0.00)	3 (1.88)	0.2097	0 (0.00)	3 (1.88)	0.3567
Aspirin						
N (%)	99 (55.62)	59 (36.88)	8.00e-04	52 (43.33)	59 (36.88)	0.3321
Clopidogrel						
N (%)	32 (17.98)	3 (1.88)	2.97e-06	4 (3.33)	3 (1.88)	0.6989
Vitamin K Antagonist						
N (%)	13 (7.30)	4 (2.50)	0.0770	7 (5.83)	4 (2.50)	0.2670
Nitrate						
N (%)	17 (9.55)	4 (2.50)	0.0141	4 (3.33)	4 (2.50)	0.9587
Beta Blocker						
N (%)	95 (53.37)	51 (31.87)	1.00e-04	46 (38.33)	51 (31.87)	0.3187
ACE Inhibitor						
N (%)	67 (37.64)	32 (20.00)	6.00e-04	35 (29.17)	32 (20.00)	0.1015

III. Exploratory Data Analyses for Metabolites and Complex Lipids

[0135] Sample preparation and mass spectrometry analyses were conducted by Metabolon, Inc. The raw data contained a total of 1088 analytes, measured for 336 pilot study participants. Two pilot study participants had no data.

[0136] Of the 1088 analytes, 705 metabolites and 183 complex lipids (888 total) had less than 10% missing data. All 336 patients had less than 10% missing data. Statistical analyses were therefore applied to 888 analytes and 336 patients. There were 57 patients in the "Revasc CAD Set", 120 patients in the "Native CAD Set" and 159 patients in the "Control Set".

[0137] The data was normalized in advance of receipt. A logarithm (base 2) transformation was applied and histograms were created to show the distribution of expression by analyte (data not shown).

[0138] The metabolomics and lipidomics data were generated in a single batch, so no batch-correction was applied, however, a principal components analysis (PCA) was performed, to look for evidence of any site effects. Variables were centered and scaled to unit variance. The inventors evaluated the presence of absence of site effects and concluded that there were no systematic differences in the data obtained from different sites.

IV. Univariate Analysis of Metabolites and Complex Lipids

A. Methods

[0139] Univariate association analysis was performed for all primary and secondary endpoints using a Mann-Whitney test, and an FDR correction was applied. Listings were produced showing p-values and q-values for all analytes.

[0140] Any analyte with $q < 0.05$ was considered a preliminary association. For each, a box-plot was generated to show the distribution of expression by clinical group. A heat-map was generated to show the preliminary metabolite associations for each endpoint: on the y-axis, patients were grouped by endpoint-value, on the x-axis a dendrogram showed clustering of metabolites with $q < 0.05$. Similarly, a heat-map was generated to show the preliminary complex lipid associations for each endpoint.

B. Results

[0141] Table 4 shows details of metabolites with $q < 0.05$ for the primary endpoint of Native CAD versus Controls; Table 5 shows details of complex lipids with $q < 0.05$.

[0142] Table 6 shows details of metabolites with $q < 0.05$ for the primary endpoint of All CAD versus Controls; Table 7 shows details of complex lipids with $q < 0.05$.

Table 4:

Analyte	Biochemical	Sub-pathway	p-value	q-value
168	Isoleucylphenylalanine	Dipeptide	7.87E-05	0.0207
163	Isoleucylalanine	Dipeptide	4.48E-05	0.0207
190	Pyroglutamylglutamine	Dipeptide	0.0001	0.0207
15	Glutamate	Glutamate Metabolism	0.0002	0.0219
1	Glycine	Glycine Serine and Threonine Metabolism	0.0004	0.0412

Table 5:

Analyte	Biochemical	Sub-pathway	p-value	q-value
594	CER18:0	Sphingolipid Metabolism	9.57E-05	0.0207
581	SM16:0	Sphingomyelin	0.0004	0.0437

Table 6:

Analyte	Biochemical	Sub-pathway	p-value	q-value
168	Isoleucylphenylalanine	Dipeptide	1.94E-06	0.0010
163	Isoleucylalanine	Dipeptide	5.10E-06	0.0010
1	Glycine	Glycine Serine and Threonine Metabolism	4.36E-06	0.0010
15	Glutamate	Glutamate Metabolism	9.42E-06	0.0014
426	1-oleoylglycerol (18:1)	Monoacylglycerol	5.84E-05	0.0059
236	Mannose	Fructose Mannose and Galactose Metabolism	9.30E-05	0.0080
238	Glucuronate	Aminosugar Metabolism	0.0001	0.0093
151	Alpha-glutamyltyrosine	Dipeptide	0.0002	0.0124
176	Leucylserine	Dipeptide	0.0003	0.0129
35	3-methylglutarylcamitine-1	Lysine Metabolism	0.0004	0.0143
219	Pyruvate	Glycolysis Gluconeogenesis and Pyruvate Metabolism	0.0004	0.0144
207	Valylleucine	Dipeptide	0.0004	0.0152
315	Propionylglycine (C3)	Fatty Acid Metabolism (also BCAA Metabolism)	0.0005	0.0153
147	Gamma-glutamylvaline	Gamma-glutamyl Amino Acid	0.0005	0.0154
190	Pyroglutamylglutamine	Dipeptide	0.0005	0.0155
134	Cysteine-glutathione disulfide	Glutathione Metabolism	0.0006	0.0159
320	Hydroxybutyrylcarnitine	Fatty Acid Metabolism(Acyl Carnitine)	0.0006	0.0161
167	Isoleucylleucine	Dipeptide	0.0007	0.0168
218	Glucose	Glycolysis Gluconeogenesis and Pyruvate Metabolism	0.0008	0.0177
159	Glycyltryptophan	Dipeptide	0.0008	0.0180

EP 3 242 588 B1

(continued)

Analyte	Biochemical	Sub-pathway	p-value	q-value
1068	X - 21452		0.0009	0.0184
215	ADSGEGDFXAEAGGVR	Fibrinogen Cleavage Peptide	0.0009	0.0186
1053	X - 21365		0.0010	0.0188
206	Valylisoleucine	Dipeptide	0.0010	0.0189
242	Alpha-ketoglutarate	TCA Cycle	0.0011	0.0196
85	Alpha-hydroxyisovaleroyl carnitine	Leucine Isoleucine and Valine Metabolism	0.0013	0.0207
700	2prime-deoxyuridine	Pyrimidine Metabolism Uracil containing	0.0014	0.0214
165	Isoleucylglycine	Dipeptide	0.0015	0.0217
128	Acisoga	Polyamine Metabolism	0.0015	0.0218
680	Urate	Purine Metabolism (Hypo)Xanthine/ Inosine containing	0.0015	0.0219
1077	X - 21626		0.0015	0.0220
793	Thymol sulfate	Food Component/Plant	0.0015	0.0221
92	Tigloylglycine	Leucine Isoleucine and Valine Metabolism	0.0018	0.0243
354	12-HETE	Eicosanoid	0.0018	0.0245
975	X - 16129		0.0019	0.0256
97	3-methyl-2-oxobutyrate	Leucine Isoleucine and Valine Metabolism	0.0020	0.0265
205	Valylglycine	Dipeptide	0.0023	0.0292
26	Imidazole propionate	Histidine Metabolism	0.0026	0.0314
333	Carnitine	Carnitine Metabolism	0.0029	0.0343
108	2-hydroxybutyrate (AHB)	Methionine Cysteine SAM and Taurine Metabolism	0.0033	0.0370
322	Hexanoylcarnitine (C6)	Fatty Acid Metabolism (Acyl Carnitine)	0.0034	0.0381
945	X - 12824		0.0034	0.0383
1028	X - 21289		0.0035	0.0384
694	N2 N2-dimethylguanosine	Purine Metabolism Guanine containing	0.0037	0.0402
166	Isoleucylisoleucine	Dipeptide	0.0039	0.0412
32	2-amino adipate	Lysine Metabolism	0.0039	0.0416
422	Glycerol 3-phosphate (G3P)	Glycerolipid Metabolism	0.0040	0.0422
737	2-hydroxyhippurate (salicylurate)	Benzoate Metabolism	0.0045	0.0454
795	Methyl glucopyranoside (alpha + beta)	Food Component/Plant	0.0051	0.0486
196	Threonylleucine	Dipeptide	0.0052	0.0491

Table 7:

Analyte	Biochemical	Sub-pathway	p-value	q-value
662	CE18:2n6	Cholesterol Ester	3.01E-05	0.0037
561	PL18:2n6	Phospholipids	0.0002	0.0110
666	CE20:4n6	Cholesterol Ester	0.0005	0.0154
485	Total Diacylglycerol	Diacylglycerol	0.0005	0.0157
619	DHC24:1	Sphingolipid Metabolism	0.0006	0.0160
517	TG15:0	Triacylglycerol	0.0007	0.0171
496	DG18:1n9	Diacylglycerol	0.0007	0.0173
615	DHC20:1	Sphingolipid Metabolism	0.0008	0.0178
582	SM18:0	Sphingomyelin	0.0010	0.0190
594	CER18:0	Sphingolipid Metabolism	0.0013	0.0208
515	Total Triacylglycerol	Triacylglycerol	0.0019	0.0250
547	PL15:0	Phospholipids	0.0049	0.0473
489	DG18:0	Diacylglycerol	0.0050	0.0480

V. Prediction Modeling for Primary Endpoints

A. Methods

[0143] Gradient boosting is an approach to determine a regression function that minimizes the expectation of a loss function. (Friedman JH., Greedy function approximation: a gradient boosting machine. *Ann Statist* 29(5): 1189-1232 (2001); Friedman JH., Stochastic gradient boosting. *Comput Stat Data Anal* 38(4): 367-378(2002)). It is an iterative method, in which the negative gradient of the loss function is calculated, a regression model is fitted, the gradient descent step size is selected, and the regression function is updated. The gradient is approximated by means of a regression tree, which makes use of covariate information, and at each iteration the gradient determines the direction in which the function needs to move, in order to improve the fit to the data.

[0144] Two rounds of gradient boosting were applied for discovery purposes, in order to select the most highly predictive metabolites and lipids respectively: (1) *Metabolomics Model*: Metabolomics data only; and (2) *Lipidomics Model*: Lipidomics data only.

[0145] A further four rounds of gradient boosting were applied in order to quantify the relative benefit of different levels of model complexity: (3) *Clinical Model*: Clinical variables, plus conventional blood-based biomarker variables; (4) *Clinical-Metabolomic Model*: Clinical variables, conventional blood-based biomarker variables, plus metabolomics data; (5) *Clinical-Lipidomic Model*: Clinical variables, conventional blood-based biomarker variables, plus lipidomics data; and (6) *Clinical-Metabolomic-Lipidomic Model*: Clinical variables, conventional blood-based biomarker variables, plus metabolomics and lipidomics.

[0146] To simplify the search space, variables were filtered to include only those exhibiting nominal univariate association (raw $p < 0.05$) for the endpoint of interest. The loss function was assumed Bernoulli, due to the binary nature of the primary endpoints. A learning rate (λ) was introduced to dampen proposed moves and to protect against over-fitting. The optimal number of iterations, given by T , was determined by 5-fold cross-validation. The minimum number of observations in each terminal node was 10. Two-way interactions were allowed. Random sub-sampling, without replacement, of half of the observations was applied to achieve variance reduction in gradient estimation.

[0147] The estimated performance of the derived models was summarized by the cross-validated (CV) estimate of loss (error), based upon the optimal number of iterations/ trees. The variables selected, and their estimated relative influences were tabulated. Partial dependence plots were generated for variables exhibiting greatest influence.

[0148] In this way, six prediction models were derived for each primary endpoint, and preliminary estimates of model performance were obtained using the same data.

[0149] The six models were used to generate probability predictions for each patient. For each model, the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) were calculated for the range of predicted probability thresholds. A Receiver Operating Characteristic (ROC) curve was generated to plot sensitivity as a function of (1-specificity). The optimal classification threshold was determined on the basis of accuracy, defined as the proportion

of correct predictions. In addition, the Area Under the Curve (AUC) was estimated.

[0150] Calibration performance was assessed graphically: plots were generated to show rates of CAD versus predicted risk. A Hosmer-Lemeshow C-test was performed.

[0151] The performance of model-based predictions were compared to the performance of probability predictions obtained by Diamond-Forrester scoring and Morise scoring. (Diamond GA, Forrester JS. Analysis of probability as an aid in the clinical diagnosis of coronary-artery disease. N Engl J Med. 1979 Jun 14; 300(24): 1350-8; Morise AP, Jalis F. Evaluation of pretest and exercise test scores to assess all-cause mortality in unselected patients presenting for exercise testing with symptoms of suspected coronary artery disease. J Am Coll Cardiol. 2003 Sep 3;42(5):842-50).

Detailed Results for Native CAD

[0152] The results show that the Diamond-Forrester score provides poor prediction of the GLOBAL phenotypes (**Figure 1**). The estimates of AUC and accuracy for prediction of Native CAD indicate that performance is no better than assigning all patients an equal probability of 0.57 of disease. (**Figure 2**) Some modest improvement in performance was obtained for All CAD, but performance remained weak. (**Figure 17**)

[0153] The Morise Score also performs poorly for prediction of Native CAD, but better for the prediction of All CAD. (**Figures 3 and 4**) For the latter, accuracy of 0.61 was obtained, which compares favorably to the minimum accuracy of 0.53, obtained by assigning all patients an equal probability of disease. (**Figure 18**)

Clinical Model

[0154] Of the 888 analytes measured (705 metabolites and 183 complex lipids), 28 clinical variables exhibited a nominal univariate association (raw $p < 0.05$) for Native CAD (Table 8). Table 9 provides the filtered list of the 28 clinical variables for All CAD.

Table 8:

Age	linoleic2	Log.Lp.a..C..mg.dL.	Male
Apo.B	Log.ALP	Log.Lp.a..mass	MHDM2
B.Sitosterol	Log.Campesterol	Log.Lp.a..P..nmol.L.	SMKCURR
BMI	Log.GGT	Log.palmleic2	Statin
fast	Log.Glucose	Log.ProInsulin	Total.cholesterol
FishOil	Log.hsCRP	Log.sdLDL.C	Uric.acid
LDL.C	Log.Insulin	Log.Trig	Vit.D..ng.mL.

[0155] Of the 28 clinical variables exhibiting a nominal univariate association (raw $p < 0.05$) for Native CAD, a panel of ten clinical variables were predictive for probability of CAD. Table 9 provides the relative influence of the eleven clinical variables for Clinical Model of Native CAD. **Figure 5** provides a ROC curve for the Clinical Model of Native CAD. **Figure 6** provides Partial Dependence Plot for the eleven clinical variables for Clinical Model of Native CAD.

Table 9:

Variable	Relative Influence	Direction of Change
Log.Lp.a..P..nmol.L.	19.09	Elevated
Apo.B	15.25	Elevated
Statin	13.52	Elevated
Log.ALP	10.50	Elevated
Log.Lp.a..mass	7.93	Elevated
Log.Trig	6.71	Elevated
LDL.C	6.55	Elevated
Uric.acid	6.19	Elevated

(continued)

Variable	Relative Influence	Direction of Change
Log.GGT	5.50	Elevated
Log.Glucose	4.63	Elevated
SMKCURR	4.13	Elevated

Metabolomics Model

[0156] Of the 888 analytes measured, 105 clinical variables exhibited a nominal univariate association (raw $p < 0.05$) for Native CAD. Table 10 provides a filtered list of the 105 metabolomic variables for Native CAD.

Table 10:

1-myristoylglycerol (14:0)	glycerol 3-phosphate (G3P)	serylleucine
1-nonadecanoylglycerophosphocholine(19:0)	glycine	theobromine
1-oleoylglycerol (18:1)	glycyltryptophan	threonate
1-oleoylglycerophosphoglycerol	guanidosuccinate	threonylleucine
1-oleoyl-GPC (18:1)	histidylphenylalanine	tigloylglycine
1-stearoylglycerophosphoglycerol	hydroxybutyrylcarnitine	tryptophylphenylalanine
2-aminooctanoate	imidazole lactate	urate
2-arachidonoyl-GPE (20:4)	imidazole propionate	valylglycine
2-docosahexaenoylglycerophosphoethanolamine	indolepropionate	valylisoleucine
2-hydroxybutyrate (AHB)	isobutyrylglycine (C4)	valylleucine
2prime-deoxyuridine	isoleucylalanine	X-12212
37-dimethylurate	isoleucylglycine	X-12472
3-ethylphenylsulfate	isoleucylisoleucine	X-12524
3-hydroxy-2-ethylpropionate	isoleucylleucine	X-12544
3-methyl-2-oxobutyrate	isoleucylphenylalanine	X-12824
3-methylglutaryl carnitine-1	isoleucylvaline	X-14056
3-methylxanthine	leucylglycine	X - 14291
4-hydroxyphenylacetate	leucylserine	X-15245
7-methylurate	mannose	X-16129
7-methylxanthine	methyl glucopyranoside (alpha + beta)	X-16132
acisoga	methyl indole-3-acetate	X-17178
ADSGEGDFXAEGGGVR	N2 N2-dimethylguanosine	X-21289
alpha-glutamyltyrosine	N4-acetylcytidine	X-21335
alpha-hydroxyisovaleryl carnitine	N-acetylalanine	X-21365
alpha-ketobutyrate	N-acetyl-beta-alanine	X-21452
alpha-ketoglutarate	N-acetylputrescine	X-21626
asparagine	N-acetylthreonine	X - 21662
carnitine	N-acetylvaline	xanthine
1-myristoylglycerol (14:0)	glycerol 3-phosphate (G3P)	serylleucine

(continued)

1-nonadecanoylglycerophosphocholine(19:0)	glycine	theobromine
1-oleoylglycerol (18:1)	glycyltryptophan	threonate
1-oleoylglycerophosphoglycerol	guanidinosuccinate	threonylleucine
1-oleoyl-GPC (18:1)	histidylphenylalanine	tigloylglycine
1-stearoylglycerophosphoglycerol	hydroxybutyrylcarnitine	tryptophylphenylalanine
2-aminooctanoate	imidazole lactate	urate

[0157] Of the 105 clinical variables exhibiting a nominal univariate association for Native CAD, a panel of six metabolomic variables were predictive for probability of CAD. Table 11 provides the relative influence of the six clinical variables for the Metabolomics Model of Native CAD. **Figure 7** provides a ROC curve for the Metabolomics Model of Native CAD. **Figure 8** provides Partial Dependence Plot for the six clinical variables for Metabolomics Model of Native CAD.

Table 11:

Variable	Relative Influence	Direction of Change
glutamate	23.34	Elevated
glucuronate	19.02	Elevated
'X-12212'	15.63	Decreased
'glutamine-leucine'	15.12	Decreased
isoleucylvaline	14.46	Decreased
valylleucine	12.43	Decreased

Clinical-Metabolomic Model

[0158] For the Clinical-Metabolomic Model, a panel of ten clinical variables were predictive for probability of CAD. Table 12 provides the relative influence of the ten clinical variables for the Clinical-Metabolomic Model of Native CAD. Figure 9 provides Partial Dependence Plot for ten clinical variables for Clinical-Metabolomic Model of Native CAD. **Figure 10** provides a ROC curve for the Clinical-Metabolomic Model of Native CAD.

Table 12:

Variable	Relative Influence	Direction of Change
glutamate	13.13	Elevated
X-16132	11.78	Elevated
isoleucylalanine	11.43	Decreased
isoleucylphenylalanine	11.11	Decreased
Log.Lp.a..mass	9.38	Elevated
Log.sdLDL.C	9.04	Elevated
X-12212	8.84	Decreased
ADSGEGDFXAEGGGVR.	8.82	Elevated
glutamine.leucine	8.66	Decreased
threonylleucine	7.82	Decreased

Lipidomics Model

[0159] Of the 138 lipids evaluated, 29 exhibited a nominal univariate association (raw $p < 0.05$) for Native CAD. Table

13 provides a filtered list of the 29 lipidomic variables for Native CAD.

Table 13:

CE16:1n7	DG16:0	DHC20:1	HexCER16:0	SM16:0
CE18:2n6	DG18:0	DHC24:0	HexCER18:1	SM20:1
CE20:4n6	DG18:1n9	DHC24:1	LacCER16:0	TG15:0
CER18:0	DG20:3n9	FA18:3n6	PL15:0	TG16:0
CER20:0	DHC18:0	FA20:3n6	PLdm16:0	Total Triacylglycerol
CER22:1	DHC18:1	HexCER14:0	PLdm18:1n9	

[0160] For the Lipidomics Model, a panel of seven variables were predictive for probability of CAD. Table 14 provides the relative influence of seven lipidomic variables for the Lipidomics Model of Native CAD. **Figure 11** provides Partial Dependence Plot for seven clinical variables for Lipidomics Model of Native CAD. **Figure 12** provides a ROC curve for the Lipidomics Model of Native CAD.

Table 14:

Variable	Relative Influence	Direction of Change
CER18:0	29.10	Elevated
DHC20:1	17.10	Elevated
CER20:0	12.54	Elevated
SM16:0	12.25	Decreased
TG15:0	10.54	Decreased
HexCER18:1	10.31	Elevated
CER22:1	8.15	Elevated

Clinical-Lipidomic Model

[0161] A panel of seven clinical variables were predictive for probability of CAD in the Clinical-Lipidomic Model. Table 15 provides the relative influence of those seven clinical variables for the Clinical-Lipidomics Model of Native CAD. **Figure 13** provides Partial Dependence Plot for seven clinical variables for Clinical-Lipidomics Model of Native CAD. **Figure 14** provides a ROC curve for the Clinical-Lipidomic Model of Native CAD.

Table 15:

Variable	Relative Influence	Direction of Change
CER18.0	25.81	Elevated
Log.ALP	14.97	Elevated
DHC20.1	14.88	Elevated
Log.Lp.a..P..nmol.L.	12.85	Elevated
SM16.0	10.61	Decreased
Apo.B	10.49	Elevated
CER20.0	10.40	Elevated

Clinical-Metabolomic-Lipidomic Model

[0162] A panel of six clinical variables were predictive for probability of CAD in the Clinical-Metabolomic-Lipidomic Model. Table 16 provides the relative influence of those six variables for the Clinical-Metabolomic-Lipidomic Model of

Native CAD. Figure 15 provides Partial Dependence Plot for six clinical variables for Clinical-Metabolomic-Lipidomics Model of Native CAD. Figure 16 provides a ROC curve for the Clinical-Metabolomic-Lipidomic Model of Native CAD.

Table 16:

Variable	Relative Influence	Direction of Change
glutamate	20.76	Elevated
X-16132	18.63	Elevated
isoleucylalanine	18.08	Decreased
Log.Lp.a..mass	14.83	Elevated
X-12212	13.98	Decreased
glutamine.leucine	13.71	Decreased

Detailed Results for All CAD

Clinical Model

[0163] 35 clinical variables exhibited a nominal univariate association (raw $p < 0.05$) for Native CAD (Table 17). Table 17 provides a filtered list of the 35 clinical variables for All CAD.

Table 17:

Age	Log.A1C	Log.Lp.a..mass	nervonic2
B.Sitosterol	Log.Adiponectin	Log.Lp.a..P..nmol.L.	06total
behenic2	Log.Campesterol	Log.NT.ProBNP	oleic2
BMI	Log.Desmosterol	Log.palmleic2	SMKCURR
cismontotl	Log.GGT	Log.ProInsulin	Statin
Fast	Log.Glucose	Log.sdLDL.C	transpalm2
FishOil	Log.HDL.C	Log.Trig	Uric.acid
HDL3.C	Log.Insulin	Male	Vit.D..ng.mL.
linoleic2	Log.Lp.a..C..mg.dL.	MHDM2	

[0164] Of the 35 clinical variables exhibiting a nominal univariate association (raw $p < 0.05$) for ALL CAD, a panel of eleven clinical variables were predictive for probability of CAD. Table 18 provides the relative influence of the eleven clinical variables for Clinical Model of All CAD. **Figure 19** provides Partial Dependence Plot for eleven clinical variables for Clinical Model of All Native CAD. **Figure 20** provides a ROC curve for the Clinical Model of All CAD.

Table 18:

Variable	Relative Influence	Direction of Change
oleic2	12.78	Elevated
Log.NT.ProBNP	12.00	Elevated
Log.Lp.a..mass	11.62	Elevated
Log.Insulin	10.00	Elevated
Log.A1C	9.97	Elevated
cismontotl	8.93	Elevated
Statin	8.54	Elevated
Log.sdLDL.C	8.21	Elevated

(continued)

Variable	Relative Influence	Direction of Change
behenic2	6.23	Elevated
Log.Adiponectin	6.11	Decreased
Log.GGT	5.61	Elevated

Metabolomics Model

[0165] As shown in Table 19, a filtered list of 138 metabolomic variables a nominal univariate association (raw p<0.05) for All CAD.

Table 19:

13-dimethylurate	glycerol 3-phosphate (G3P)	pyroglutamylglutamine
17-dimethylurate	glycine	pyroglutamylglycine
12-HETE	glycylphenylalanine	pyruvate
1-linoleoyl-GPE (18:2)	glycyltryptophan	S-adenosylhomocysteine (SAH)
1-methylurate	guanidinosuccinate	salicylate
1-nonadecanoylglycerophosphocholine(19:0)	hexanoylcarnitine (C6)	serylleucine
1-oleoylglycerol (18:1)	histidylphenylalanine	succinylcarnitine (C4)
1-oleoylglycerophosphoglycerol	homostachydrine	threonate
2-aminoadipate	hydroxybutyrylcarnitine	threonylleucine
2-aminobutyrate	imidazole propionate	thymol sulfate
2-aminooctanoate	indolepropionate	tigloylglycine
2-arachidonoyl-GPE (20:4)	isobutyrylglycine (C4)	tryptophylglycine
2-docosahexaenoylglycerophosphoethanolamine	isoleucylalanine	tryptophylphenylalanine
2-hydroxybutyrate (AHB)	isoleucylglycine	tyrosylglutamine
2-hydroxyhippurate (salicylurate)	isoleucylisoleucine	urate
2-linoleoyl-GPE (18:2)	isoleucylleucine	valeryl carnitine (C5)
2prime-deoxyuridine	isoleucylphenylalanine	valylglycine
3-ethylphenylsulfate	isovaleryl glycine	valylisoleucine
3-hydroxyisobutyrate	kynurenine	valylleucine
3-methyl-2-oxobutyrate	leucylaspartate	valylvaline
3-methylglutaryl carnitine-1	leucylglycine	X - 11429
3-methylglutaryl carnitine-2	leucylserine	X - 11444
3-methylxanthine	lysylleucine	X - 11787
4-hydroxyphenylacetate	mannose	X - 11945
5alpha-androstan-3beta 17beta-diol monosulfate 2	methyl glucopyranoside (alpha + beta)	X - 12212
7-methylurate	methyl indole-3-acetate	X - 12472
7-methylxanthine	N2 N2-dimethylguanosine	X - 12729

(continued)

acetylcarnitine (C2)	N4-acetylcytidine	X - 12824
acisoga	N6-carbamoylthreonyladenosine	X - 14056
ADSGEGDFXAEGGGVR	N-acetylalanine	X - 15245
alpha-glutamyltyrosine	N-acetyl glycine	X - 15492
alpha-hydroxyisovaleroyl carnitine	N-acetyl isoleucine	X - 16129
alpha-ketobutyrate	N-acetylneuraminate	X - 16132
alpha-ketoglutarate	N-acetylthreonine	X - 17178
beta-tocopherol	N-acetylvaline	X - 17690
butyrylcarnitine (C4)	N-methyl proline	X - 18914
carnitine	octanoylcarnitine (C8)	X - 18922
cysteine-glutathione disulfide	oleic ethanolamide	X - 19438
cytidine	O-sulfo-L-tyrosine	X - 21289
fucose	phenylalanyl aspartate	X - 21335
gamma-glutamylisoleucine	prolyl phenylalanine	X - 21365
gamma-glutamylvaline	propionyl carnitine (C3)	X - 21367
glucose	propionyl glycine (C3)	X - 21452
glucuronate	pseudouridine	X - 21471
glutamate	pyridoxal	X - 21626
glutamine-leucine	pyridoxate	xanthine

[0166] Of the 138 clinical variables exhibiting a nominal univariate association for All CAD, a panel of sixteen metabolomic variables were predictive for probability of CAD. Table 20 provides the relative influence of the sixteen clinical variables for the Metabolomics Model of Native CAD. **Figure 21** provides Partial Dependence Plot for sixteen clinical variables for Metabolomics Model of All CAD. **Figure 22** provides a ROC curve for the Metabolomics Model of All CAD.

Table 20:

Variable	Relative Influence	Direction of Change
isoleucylalanine	9.75	Decreased
glucuronate	8.31	Elevated
glycine	8.11	Decreased
isoleucylphenylalanine	7.69	Decreased
mannose	7.65	Elevated
'X-21452'	6.93	Decreased
'1-oleoylglycerophosphoglycerol'	6.50	Elevated
'X-21335'	5.94	Elevated
'7-methylxanthine'	5.77	Elevated
'X-12729'	5.46	Decreased
acisoga	5.46	Elevated
'alpha-hydroxyisovaleroyl carnitine'	5.11	Elevated
'2-aminoadipate'	4.80	Elevated
'X-18914'	4.78	Decreased

(continued)

Variable	Relative Influence	Direction of Change
tigloylglycine	4.16	Decreased
pyridoxal	3.56	Decreased

Clinical-Metabolomic Model

[0167] A panel of twelve clinical variables were predictive for probability of CAD in the Clinical-Metabolomic Model for All CAD. Table 21 provides the relative influence of the twelve clinical variables for the Clinical-Metabolomic Model of All CAD and **Figure 23** provides Partial Dependence Plots for the twelve clinical variables for Clinical-Metabolomic Model of All CAD. **Figure 24** provides a ROC curve for the Clinical-Metabolomic Model of All CAD.

Table 21:

Variable	Relative Influence	Direction of Change
isoleucylalanine	12.92	Decreased
glycine	10.75	Decreased
isoleucylphenylalanine	10.15	Decreased
oleic2	9.55	Elevated
X-21335	8.22	Elevated
X3.methylxanthine	7.97	Elevated
X2.aminoadipate	7.76	Elevated
mannose	7.28	Elevated
acisoga	6.73	Elevated
X2.arachidonoyl.GPE...20.4..	6.51	Elevated
Log.Lp.a..P..nmol.L.	6.16	Elevated
Log.Trig	6.01	Elevated

Lipidomics Model

[0168] 31 lipidomic variables exhibited a nominal univariate association (raw $p < 0.05$) for All CAD. Table 22 provides a filtered list of the 29 lipidomic variables for All CAD.

Table 22:

CE16:0	DG18:0	DHC18:1	DHC26:1	PL24:0
CE16:1n7	DG18:1n9	DHC20:0	FA14:1n5	SM18:0
CE18:1n9	DG20:0	DHC20:1	PL15:0	TG15:0
CE18:2n6	DG20:2n6	DHC22:1	PL18:2n6	TG20:3n9
CE20:4n6	DG20:3n9	DHC24:1	PL20:3n9	Total Diacylglycerol
CER18:0	DHC18:0	DHC26:0	PL20:4n6	Total Triacylglycerol
CER18:1				

[0169] A panel of thirteen variables were predictive for probability of CAD. Table 23 provides the relative influence of the thirteen lipidomic variables for the Lipidomics Model of All CAD. **Figure 25** provides Partial Dependence Plots for thirteen clinical variables for Lipidomics Model of All CAD. **Figure 26** provides a ROC curve for the Lipidomics Model of All CAD.

Table 23:

Variable	Relative Influence	Direction of Change
'DHC18:0'	14.54	Elevated
'DG18:1n9'	12.23	Elevated
'Total Diacylglycerol'	11.50	Elevated
'TG20:3n9'	8.14	Elevated
'CE20:4n6'	7.92	Elevated
'CER18:1'	7.45	Elevated
'DHC20:1'	7.28	Elevated
'CE18:2n6'	6.24	Decreased
'PL18:2n6'	5.97	Decreased
'SM18:0'	5.34	Elevated
'FA14:1n5'	5.31	Decreased
'PL15:0'	4.66	Decreased
'CE16:0'	3.41	Elevated

Clinical-Lipidomic Model

[0170] A panel of eleven clinical variables were predictive for probability of CAD in the Clinical-Lipidomic Model for All CAD. Table 24 provides the relative influence of those eleven clinical variables for the Clinical-Lipidomics Model of All CAD. **Figure 27** provides Partial Dependence Plot for the eleven clinical variables for Clinical-Lipidomics Model of All CAD. **Figure 28** provides a ROC curve for the Clinical-Lipidomic Model of All CAD.

Table 24: Relative influence for Clinical-Lipidomic Model of All CAD

Variable	Relative Influence	Direction of Change
Log.A1C	15.78	Decreased below a cutpoint and Elevated over another cutpoint represents risk
Total.Diacylglycerol	11.49	Elevated
cismontotl	11.08	Elevated
Log.Lp.a..mass	9.27	Elevated
Statin	8.99	Elevated
TG20.3n9	8.13	Elevated
oleic2	7.67	Elevated
DHC18.1	7.47	Elevated
Log.NT.ProBNP	7.12	Elevated
Log.GGT	6.68	Elevated
CER18.0	6.32	Elevated

Clinical-Metabolomic-Lipidomic Model

[0171] A panel of twelve clinical variables were predictive for probability of CAD in the Clinical-Metabolomic-Lipidomic Model of All CAD. Table 25 provides the relative influence of the twelve variables for the Clinical-Metabolomic-Lipidomic Model of All CAD. **Figure 29** provides Partial Dependence Plots for the twelve clinical variables for Clinical-Metabolomic-Lipidomic Model of All CAD. **Figure 30** provides a ROC curve for the Clinical-Metabolomic-Lipidomic Model of Native CAD.

Table 25:

Variable	Relative Influence	Direction of Change
isoleucylalanine	12.49	Decreased
Glycine	10.39	Decreased
Cismontotl	9.55	Elevated
Log.Lp.a..mass	9.01	Elevated
X-21452	8.82	Decreased
isoleucylphenylalanine	7.81	Decreased
Log.A1C	7.59	Elevated
X-21335	7.49	Elevated
X7.methylxanthine	7.12	Elevated
BMI	7.10	Elevated
acisoga	6.83	Elevated
X-12729	5.79	Decreased

[0172] Throughout the examples described herein, a number of analytes are identified as "X-" compounds. The several X-compound are described in further detail in Table 26.

Table 26:

name	lib_id	comp_id	quant_mass	rt	tentative details of unnamed metabolites, if available	spectra
X- 12212	LC/MS Neg	46358	229.01793	3.47	sulfated	56.99574:5.1 59.98531:1 60.99317:0.6 68.99663:0.8 79.95761:1.3 80.96536:2 93.03481:0.3 108.02196:0.8 121.06626:0.2 122.03769:1.9 122.96783:0.7 130.04274:0.3 131.05064:0.5 134.03756:0.2 140.9786:0.5 147.04556:2.2 148.05347:3.8 149.06116:100 150.06449:11.1 162.9604:0.2 166.95785:0.4 184.96803:0.4 188.9398:2.5 206.95033:1.5 229.01806:0.7
X- 12472	LC/MS Neg	46628	241.11935	2.38		58.0299:7.8 59.014:0.8 67.0304:3.4 72.00935:3.9 72.99325:1.2 74.0249:14.5 82.03117:6 84.0456:29.5 86.02505:3.2 95.05055:1 97.0409:3.2 98.02495:4

EP 3 242 588 B1

5

10

15

20

25

30

35

40

45

50

55

						99.05655:4 101.0721:17.3 107.0253:3.5 109.0408:33.5 112.0771:1.1 125.0357:13.6 127.0514:95.7 128.03545:38.2 128.05505:5.6 145.062:100 146.06585:4.4
X- 12524	LC/M S Neg	46621	205.1599	5.54		189.12873:8.1 190.13217:1.1 205.16003:100 206.16342:18.1
X- 12544	LC/M S Pos	34112	211.0962	4.25	C11H14O4	53.0391:1 55.0183:0.9 55.05465:0.1 57.03405:0.2 65.0389:1.5 66.0445:0.2 67.05455:1 77.03865:1.3 78.0465:1.1 79.0544:11.9 80.05987:0.5 81.03375:0.1 81.06995:0.2 90.04655:0.2 91.05435:15.9 92.05913:0.9 93.07015:1.4 95.0493:7.3 95.08605:0.3 96.05265:0.4 103.0544:10.1 104.05735:1.1 105.0337:0.5 105.0448:1.2 105.07005:27.5 106.0414:0.6 106.07335:3.1 107.0494:0.8 107.0793:0.2 107.0859:0.2 108.05715:2.7 109.06545:1.5 115.0544:0.5 117.07:0.5 118.04135:0.8 119.04965:0.8 120.05715:4.8 121.02855:0.4 121.0651:11.8 122.0319:0.5 122.06715:0.9 123.0808:1.3 133.065:1.5 133.0841:2.2 134.0904:49.5 135.04445:0.8 135.081:8.4 135.0966:24.5 136.05205:1.2 136.08425:0.9 137.0598:0.1 139.0754:0.8 149.0235:1.4 149.0601:0.8 149.0793:0.1 150.027:3.5 150.0678:7.8 150.0853:1 151.0756:100 152.0789:10.6 152.09625:0.4 165.0912:44.2 166.09455:5.4 193.0861:2.2 194.0897:0.3
X- 12824	LC/M S Neg	46471	243.1351 3	2.72		56.99603:4 58.02977:8.1 59.0142:1 61.98863:2.7 67.0303:1.4 74.02507:5.4 74.99407:1.4 79.9578:1 82.03013:3.6 83.05065:0.8 83.0618:4.6 84.0457:12.3 92.9294:1.9 97.066:3.4 101.07217:12.2 109.0408:14.8 110.0447:0.8 113.0973:0.8 114.09257:100 115.0766:1.1 115.09597:7.3 125.03593:2.4 127.0515:39.4 128.03547:11.2 128.05517:1.4 128.1083:1.4 136.9402:1.6 145.06197:23.5

EP 3 242 588 B1

						147.03285:2.3 162.05645:6.2 162.8935:1.3 164.89587:9.3 181.13493:14.6 182.1383:1.9 188.93987:2 206.9505:1.3 210.9053:1.1 242.8729:8.5
X- 14056	LC/M S Pos	46632	192.0325 4	0.9		55.01832:1 55.93465:1.6 56.04993:2.9 57.98756:2.5 58.99541:96.7 59.99693:1.3 67.04195:1 71.01293:0.4 72.93729:1.1 73.01092:2.7 73.02854:0.5 76.02167:2.7 77.00568:1.1 85.01076:2.6 85.02846:0.6 86.99001:38.5 87.99308:1.2 88.03939:35.8 89.04262:1.2 90.94774:0.7 100.02161:100 101.00566:5.5 101.02507:4.3 105.00054:26.1 106.00399:0.6 110.98994:1.4 113.00562:1.3 118.03217:0.9 128.01649:3.4 129.00051:4.7 146.02709:40.1 147.01132:1 147.03064:2.4 175.0062:0.7
X- 14291	LC/M S Pos	46414	195.1127 4	2.26		55.9349:30 70.06541:100 71.06908:4 98.06023:67.3 99.06351:3.7 195.11306:26.4 196.11652:3.3
X- 15245	LC/M S Neg	46661	233.0303 3	0.55		57.03447:14.6 59.014:4.4 67.01913:3.7 71.01412:4.3 73.02963:9.6 81.03482:4 83.01418:1.7 83.05046:4.8 87.0089:10.8 94.99334:4.2 99.00891:17.3 101.02453:100 102.02798:5.2 111.00896:11.4 127.04026:4.4 127.99036:1.3 128.98645:2.7 145.00866:32.6 188.99888:6.3 233.15516:13.1 234.1583:1.9
X- 16129	LC/M S Pos	46642	287.6506 2	1.39		55.05471:6.7 56.04994:9.1 57.05657:0.5 60.05593:9.2 68.04974:1.3 70.06546:33.3 71.06867:1.2 72.08095:72 73.08427:3.4 84.04451:100 85.02856:1 85.04719:3.9 87.05555:14 88.07577:1.7 96.04446:7.1 97.07616:0.7 102.05503:7.8 106.08634:1.4 112.08699:4.9 113.07096:3.6 114.1027:1.3 115.05029:7.7 115.08666:3.5 116.07066:11.7 127.05026:1.6 129.1023:18.8 130.09762:3.7 130.10606:1 137.59745:0.8 141.06593:7.5 142.09755:3.1 157.0973:1.1 158.09249:8.3 169.06086:6.2 175.11909:17.5 274.18752:3 331.20942:1.3

EP 3 242 588 B1

5

10

15

20

25

30

35

40

45

50

55

X- 17178	LC/M S Neg	46406	661.2552 4	2.31		58.02997:2 60.99327:1.4 67.03042:2.4 70.03061:14.1 71.02544:6.5 72.00932:5.2 73.04091:8.8 74.02488:21.8 82.02999:23.3 83.0253:4.4 84.00973:9.5 85.04103:3.2 86.02505:3.3 94.02995:3.5 95.02522:12.2 96.00928:47.6 96.04563:4.8 97.04086:4.3 98.02489:8.3 99.0202:6.6 104.03548:12.9 109.04084:24.6 110.02489:5.7 110.03631:4.1 111.02022:4.4 111.04392:4.4 112.04059:4.4 113.03584:24.4 114.01979:10 115.08781:4.5 116.05071:7 118.06646:2.9 123.05663:11.8 124.02836:1.5 124.04055:31.7 125.03579:4 125.07224:6 127.05149:100 128.03551:44.4 128.05521:4.8 129.03898:2.5 137.03613:2.2 137.07225:83.7 138.05649:5.8 138.07584:6.3 139.01529:3.1 139.05149:12.5 141.06716:14.4 145.06521:2.2 146.06133:12.9 147.04553:2 151.08791:9 153.0671:6.1 154.09889:39.1 155.08291:2.8 155.10235:3.3 160.07707:6.5 164.07199:3 165.05452:6.2 166.06246:67.9 167.04649:32.4 167.06622:4.5 168.05006:2.4 172.07704:15.7 172.10929:2.8 181.0623:4 184.07311:34.2 185.07305:8.5 189.06719:1.9 190.08768:2.5 194.09386:3.7 222.07612:9.2 265.1308:56.4 266.13418:7.8 283.14148:8.1
X- 21289 (renamed as X- 14465)	LC/M S Pos	46310	211.1435 3	3.17		53.00271:0.2 53.03908:0.3 55.01833:0.5 55.05472:0.5 55.93482:2.4 57.07036:0.5 61.03997:0.8 67.05449:0.8 68.04967:0.7 69.0701:2.3 70.06538:100 71.0624:0.3 71.06913:3.7 71.08566:0.3 72.08094:0.3 72.93724:0.6 74.95295:0.2 77.0387:0.2 79.05434:1 80.04958:0.6 81.0336:0.9 81.06996:0.7 82.06524:0.6 84.08092:0.2 86.09651:23.6 87.09881:1 90.94778:1.4 91.05433:1.2 93.06995:0.6 95.04921:2.3 95.08559:0.4 96.08085:0.3 97.06483:0.3 98.04746:0.3

EP 3 242 588 B1

5

10

15

20

25

30

35

40

45

50

55

						98.06009:0.8 99.09154:0.2 104.95094:0.3 105.03354:4.9 105.04462:0.3 105.06992:0.9 105.93487:0.4 106.03689:0.3 109.06487:0.5 110.06012:1 112.02687:1 113.07095:0.5 114.05501:0.3 114.09139:10.1 115.09475:0.7 119.08549:0.4 121.084:0.3 125.07085:0.3 126.07878:0.6 126.12785:0.2 127.08668:1.1 135.04423:0.8 137.07117:0.7 138.05506:0.2 138.12781:11.2 139.13129:1.1 149.05985:0.9 150.08528:0.4 153.06583:1.7 154.07373:14.1 155.07773:1 155.15428:3.2 166.12275:1.4 183.14922:4.1 184.15265:0.4 194.11745:1.5 211.14404:5.7 212.14739:0.7
X- 21335 (renamed as X- 16132)	LC/M S Neg	46380	313.4437 4	0.95		56.99582:14.9 58.02984:4.1 59.01394:1.1 59.98543:2.2 67.03042:1.4 70.03066:16.5 71.01414:2 74.02489:39.5 82.03012:33.4 84.00923:7.4 84.04569:5.6 85.04109:10.7 87.05662:1.8 91.02232:2 93.00156:1.9 94.03002:2.2 95.02524:17.5 96.00927:100 96.04572:4.5 97.01255:4.5 97.04092:1.6 98.02491:9.8 98.06135:1.5 109.04084:17.5 110.02493:4.6 110.03621:2.7 110.06129:1.7 111.0202:2 111.04401:4.3 111.05659:3.9 112.02798:3.9 112.04051:6.7 113.03583:54.1 113.0722:5 114.01982:8 114.03913:2 115.08784:5.7 123.05656:6.9 124.04057:23.7 125.03584:2.9 125.07221:6.3 127.05151:60.1 128.03553:84.9 128.05516:3.3 129.03903:5 136.04062:1.6 137.07226:70.9 138.05631:4.5 138.07575:5.3 139.0153:1.3 139.05146:9.4 141.06721:51 142.07062:3 149.03607:1.3 151.08797:4.4 153.06716:3.8 154.09886:38 155.08295:2.2 155.10235:2 162.96043:1.9 165.05463:4.7 166.06244:6.8 167.04641:13.1 172.1095:1.9 181.06215:2 183.0414:3 184.0732:3.1 185.05707:17.2 188.93964:23.8 206.95024:14.2 222.0761:6.7 223.02876:1.8 265.13086:38.6 266.13405:5.8 283.14168:4.3

EP 3 242 588 B1

5

10

15

20

25

30

35

40

45

50

55

X- 21365	LC/M S Pos	46410	160.1332 6	0.91		53.03916:2.2 55.05479:100 56.05004:1.4 56.05833:4.8 57.03401:1.2 57.07041:0.7 58.06576:6.6 59.04963:26.7 59.07262:1.1 60.05298:0.7 60.08128:80.5 61.0848:2.3 67.05459:0.8 69.03384:0.6 69.07022:0.5 70.06544:0.7 71.04943:0.6 72.04467:2.2 72.93739:0.5 73.06505:1.2 83.04937:37.1 84.05268:2.1 84.08102:1.9 84.95992:1.4 87.04431:3.6 91.05443:0.6 95.08577:0.8 101.05997:86.6 101.10755:2.5 102.06325:5.1 105.0701:0.8 160.1335:35.9 161.13707:3.5
X- 21452 (renamed as X- 02269)	LC/M S Neg	46497	255.0876 1	1.49	hydroxy metabolite of CMPF	55.0189:100 57.03468:17.2 59.01397:19.3 71.05051:5.3 73.02978:5.6 84.02189:7.9 85.02967:47.7 86.037:28.8 93.03473:9.8 107.05039:37 109.06599:68.2 139.07662:16.6 147.08171:24 149.09737:16.6 153.05596:11.8 167.10795:23.2 211.09786:31.9 237.8701:8.4 254.91917:11.4 255.23316:50.7 256.23649:11.2
X- 21626	LC/M S Neg	46694	696.7766 1	2.4		58.02983:1.8 66.03515:0.8 67.03025:2.6 68.05086:0.6 70.03052:13 71.01414:1.9 71.02533:6 72.00938:0.6 72.04573:0.6 73.04118:8.9 74.02484:2.5 80.05081:1 82.01751:1.3 82.03001:22 83.02528:4.1 83.03341:0.7 84.00975:10.7 84.04565:2 85.04097:2.3 86.02505:1.2 87.0566:1.2 88.04059:60.8 89.0437:1.8 91.05553:1.5 94.03005:2.8 94.06643:0.5 95.02526:12.4 96.0093:43.2 96.04566:4 97.01273:1.6 97.0409:2.8 98.0124:1.6 98.02495:8.3 98.06133:0.6 99.0202:6.9 99.05659:7.8 100.04064:0.9 101.07217:5.3 106.03:0.5 107.02526:1.6 108.03323:0.6 109.02861:1.4 109.04085:24.3 110.01249:0.9 110.02493:5.3 110.03632:4.1 110.04395:1.2 110.0612:1.9 111.02026:6.2 111.04409:4.8 111.05643:2.2 112.02816:1.1

EP 3 242 588 B1

5

10

15

20

25

30

35

40

45

50

55

						112.04056:4.5 113.03584:22.1
						113.07218:2.2 114.01981:9.4
						114.03904:0.6 115.08788:4.2
						116.05076:5.2 118.06643:3.2
						121.04083:1.1 122.02497:0.6
						123.05659:9.8 124.02795:1.3
						124.0406:29.6 125.03585:3.8
						125.04412:1.7 125.07225:6.1
						127.05153:100
						128.03555:38.3 128.05529:5.3
						129.03913:2 130.06567:0.7
						131.04677:1.1 135.05658:1.9
						136.04068:1 137.03619:2
						137.07226:78.8 138.05655:5.3
						138.07601:5.7 139.01528:2.4
						139.05147:10.5 139.08788:0.7
						140.07209:0.9 140.08279:0.7
						141.03084:0.9 141.0672:13.4
						142.06758:1 145.06211:10.8
						146.06132:11.6 147.04558:0.9
						147.06506:1 149.03591:1.7
						151.05172:1 151.08795:8.9
						152.07198:0.9 152.08264:1
						153.03092:1.3 153.06711:4.8
						154.09888:37.3 155.08325:3.3
						155.10263:2.7 156.0783:1.8
						157.0626:1.1 158.06144:2.7
						158.09379:0.7 160.07705:3.7
						163.05154:1 163.08804:2
						164.07197:1.2 165.05471:6.6
						166.06249:50.9
						167.04654:26.1 167.06708:3.4
						167.08272:1.3 168.05029:1.8
						171.09326:1 172.07742:0.7
						172.10944:2.6 173.07229:1.5
						175.08778:0.8 177.0674:1.8
						180.07808:0.7 181.06225:3.1
						184.0732:27.4 185.05684:1.4
						185.07237:15.4 186.05647:9.5
						186.07532:1.7 187.06032:0.9
						188.07211:1 189.06713:2.3
						193.02589:0.9 194.09383:3.8
						195.07808:0.9 195.11422:1.1
						201.06731:1 202.09936:2.3
						203.08317:3.2 210.05257:0.7
						212.1046:1 213.10414:0.6
						215.08308:0.8 220.10956:1.4
						222.07619:11.7 222.12498:0.9
						223.0805:1.2 229.09872:1.1
						231.11421:5.9 232.11622:1
						238.08375:1.9 238.12027:1.2
						243.11435:1.3 248.10805:1.3
						256.11026:1.9 265.13091:53.2
						266.13433:7.5 274.11996:1.7
						276.10381:1 283.14159:7.7
						284.14486:1 292.1308:0.9

						313.13127:7.1 314.13535:0.9 331.14176:2.3
5	X- 21662	LC/M S Pos	46731	510.3558 3	5.62	15- methylpalmit oyl-GPC
10						53.0027:0.7 55.0547:2.7 56.0499:4.1 56.96523:0.4 57.03436:1.2 57.0703:7.3 58.06545:2.8 58.07365:0.2 59.04953:3.2 59.0734:0.6 60.08115:36.5 61.08445:1.2 65.0389:0.2 67.05448:3.6 69.0701:2.1 70.06533:0.7 71.0731:22.1 71.0857:3.2 72.07738:0.7 73.0285:1.4 79.05432:1.4 81.07:3.3 83.08558:1.2 85.10112:1.2 86.09653:100 87.04413:0.5 87.09423:0.3 87.0998:4.8 89.05973:1.8 91.05425:1.3 93.06997:1.3 95.0492:0.2 95.0856:3.1 96.0889:0.2 97.10122:0.6 98.98423:17.9 104.10703:73.8 105.06988:1.1 105.11032:3.7 107.08547:0.5 109.10108:1 119.0854:1 124.9999:65.1 126.0032:1.3 131.08543:0.4 133.10083:0.3 143.01033:0.3 163.01535:0.3 184.07342:44 185.07693:2.2
15						
20						
25						

30 **[0173]** For Table 26, the data is in the following format: **name; lib_id; comp_id; quant_mass; rt; tentative details of unnamed metabolites, if available; and spectra**. The spectra data format (x:y) indicates the mass of the ion ($m/z = x$) and relative peak intensity (y). Each ion in the spectra is separated by a space.

Methods: Univariate Analysis of Secondary Endpoints

35

Example 2. Discovery and Validation

Analysis Populations

40 **[0174]** The "Discovery-Full Analysis Set" ("Discovery FAS") consisted of pilot study patients with clinical data and a CT-based designation of either Revascularization CAD case, Native CAD case, or Control (N = 748 for the Discovery-FAS group).

45 **[0175]** The "Discovery-Native CAD Set" was the subset of the Discovery-FAS with Native CAD as verified by CT, who had analyte (metabolomic) data (N = 366 for the Discovery-Native CAD Set). These were subjects without previous revascularization procedures, such as percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG).

[0176] The "Discovery-Revasc CAD Set" was the subset of the Discovery-FAS who had undergone previous revascularization, such as percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG), and who had analyte data (N=44).

50 **[0177]** The "Discovery-All CAD Set" was the union of the Discovery-Native CAD Set and the Discovery-Revasc CAD Set (N=410).

[0178] The "Discovery-Control Set" was the subset of Discovery-FAS who had a calcium score of zero and were designated a Control after inspection of CT data, and who had analyte data. (N=338 for the Discovery-Control Set.)

55 **[0179]** The "Validation-Full Analysis Set" ("Validation-FAS") consisted of pilot study patients with clinical data and a CT-based designation of either Revascularization CAD case, Native CAD case, or Control (N = 348 for the Validation-FAS group).

[0180] The "Validation-Native CAD Set" was the subset of the Validation-FAS with Native CAD as verified by CT, who had analyte (metabolomic) data (N = 207 for the Validation-Native CAD Set). These were subjects without previous

revascularization procedures, such as percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG).

[0181] The "Validation-Revasc CAD Set" was the subset of the Validation-FAS who had undergone previous revascularization, such as percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG), and who had analyte data (N=15).

[0182] The "Validation-All CAD Set" was the union of the Validation-Native CAD Set and the Validation-Revasc CAD Set (N=222).

[0183] The "Validation-Control Set" was the subset of Validation-FAS who had a calcium score of zero and were designated a Control after inspection of CT data, and who had analyte data. (N=126 for the Validation-Control Set)

[0184] It is noted that by design, the only racial group represented in the study was White. Therefore, race-based subpopulations were not defined.

A. Study Endpoints

[0185] For the GLOBAL Pilot Discovery Cohort, there were four primary endpoints in the analysis: (1) Native CAD; (2) All CAD (Native or Revascularization); (3) 50% Stenosis without Revascularization; (4) 50% Stenosis or Revascularization. All analyses were applied to all primary endpoints.

B. Statistical Hypothesis

[0186] The null hypothesis of no association, between the metabolite or lipid and the endpoint, was tested against the two-sided alternative that association exists.

C. Multiple Comparisons and Multiplicity

[0187] False discovery rate (FDR) q-values were calculated (Benjamini and Hochberg, 1995). Associations with FDR $q < 0.05$ were considered preliminary associations. In some circumstances, test results with raw $p < 0.05$ were reported as well.


D. Missing Data

[0188] Endpoint data was not imputed. Potential covariates with more than 5% missing data were excluded. Potential covariates with less than 5% missing data were imputed to the mean.

[0189] Metabolites with more than 10% missing data were excluded from the main analyses. Missing values for metabolites and lipids with less than 10% missing were imputed to the observed minimum after normalization.

E. Analysis of Subgroups

[0190] The first and third primary endpoints were addressed using a subset of the FAS. Specifically the *Native CAD Set* and the *Control Set* were considered to the exclusion of the Revasc. CAD Set. For the purposes of discovery, further subsets were created on the basis of participants' fasting status, where patients were categorized as Fasting if they had not eaten for eight or more hours. The remainder, either known not to be fasted, or with unknown fasting status were categorized as 'Non-Fasting'.



I. Demographic and Baseline Characteristics

[0191] The baseline and demographic characteristics of patients in the pilot study were tabulated. Continuous variables were summarized by the mean and standard error; binary variables were summarized by the count and percentage.

[0192] Table 27 shows general patient characteristics for the Discovery Set by clinical group (Revasc CAD vs. Native CAD vs. Control). A Kruskal-Wallis test was performed to investigate homogeneity of continuous measures; a Pearson's chisquared test was conducted for binary measures; unadjusted p-values are reported.

[0193] Table 28 shows general patient characteristics for the Validation Set by clinical group (Revasc CAD vs. Native CAD vs. Control). A Kruskal-Wallis test was performed to investigate homogeneity of continuous measures; a Pearson's chisquared test was conducted for binary measures; unadjusted p-values are reported.

Table 27:

	All Controls	Native CAD	Revasc CAD	P-value
N	338	366	44	
Age				
mean (SE)	53.8 (0.57)	58.02 (0.54)	59.55 (1.40)	3.93E-07
SBP				
mean (SE)	129.62 (0.91)	132.6 (0.92)	128.09 (2.32)	0.0550
DBP				
mean (SE)	79.03 (0.56)	79.73 (0.58)	75.12 (1.57)	0.0402
Male				
N (%)	151 (44.67)	195 (53.28)	30 (68.18)	0.0037
Hypertension				
N (%)	172(51.04)	244 (66.85)	40 (90.91)	1.61E-08
Dyslipidemia				
N (%)	184 (55.42)	259 (71.15)	43 (100.00)	4.95E-10
Diabetes (Any)				
N (%)	25 (7.40)	54 (14.75)	10 (22.73)	8.00E-04
Type I Diabetes				
N (%)	1 (0.30)	1 (0.27)	0 (0.00)	0.9377
Type II Diabetes				
N (%)	24 (7.10)	53 (14.48)	10 (22.73)	6.00E-04

EP 3 242 588 B1

(continued)

	All Controls	Native CAD	Revasc CAD	P-value
Current Smoker				
N (%)	39 (11.54)	67 (18.31)	5 (11.36)	0.0331
Former Smoker				
N (%)	84 (24.85)	130 (35.52)	21 (47.73)	5.00E-04
Chest Pain				
N (%)	221 (65.38)	212 (57.92)	30 (68.18)	0.0850
Angina Equivalent				
N (%)	126 (37.28)	122 (33.33)	19 (43.18)	0.3115
Shortness of Breath				
N (%)	74 (21.89)	71 (19.40)	8 (18.18)	0.6635
Family History of CAD				
N (%)	179 (52.96)	223 (60.93)	28 (63.64)	0.0710
Fasting				
N (%)	120 (35.50)	207 (56.56)	44 (100.00)	8.28E-18
Statin				
N (%)	111 (32.84)	184 (50.27)	38 (86.36)	1.28E-12
Niacin				
N (%)	5 (1.48)	4 (1.09)	3 (6.82)	0.0164
Fibrate				
N (%)	12 (3.55)	21 (5.74)	3 (6.82)	0.3254
Ezetimibe				
N (%)	6 (1.78)	11 (3.01)	8 (18.18)	7.97E-08
Fish Oil				
N (%)	26 (7.69)	50 (13.66)	5 (11.36)	0.0388
Bile Acid Sequestrant				
N (%)	3 (0.89)	4 (1.09)	0 (0.00)	0.7705
Aspirin				
N (%)	98 (28.99)	157 (42.90)	34 (77.27)	3.15E-10
Clopidogrel				
N (%)	7 (2.07)	11 (3.01)	15 (34.09)	5.21E-22
Vitamin K Antagonist				
N (%)	7 (2.07)	17 (4.64)	2 (4.55)	0.1629
Nitrate				
N (%)	7 (2.07)	16 (4.37)	11 (25.00)	5.57E-11
Beta Blocker				
N (%)	114 (33.73)	140 (38.25)	34 (77.27)	1.68E-07
ACE Inhibitor				
N (%)	63 (18.64)	106 (28.96)	24 (54.55)	3.13E-07

Table 28:

	All Controls	Native CAD	Revasc CAD	P-value
N	126	207	15	
Age				
mean (SE)	49.48 (0.93)	60.83 (0.59)	68.6 (2.05)	3.35E-22
SBP				
mean (SE)	127.25 (1.55)	131.46 (1.10)	131.8 (4.16)	0.0276
DBP				
mean (SE)	77.33 (0.98)	78.99 (0.77)	78.87 (3.16)	0.2091
Male				
N (%)	46 (36.51)	138 (66.67)	15 (100.00)	1.36E-09
Hypertension				
N (%)	51 (40.48)	132 (63.77)	13 (86.67)	9.44E-06
Dyslipidemia				
N (%)	65 (53.72)	152 (74.88)	14 (100.00)	1.33E-05
Diabetes (Any)				
N (%)	12 (9.52)	44 (21.36)	4 (26.67)	0.0134
Type I Diabetes				
N (%)	1 (0.79)	3 (1.45)	1 (6.67)	0.1954
Type II Diabetes				
N (%)	11 (8.73)	41 (19.81)	3 (20.00)	0.0244
Current Smoker				
N (%)	20 (15.87)	28 (13.53)	0 (0.00)	0.238
Former Smoker				
N (%)	25 (19.84)	78 (37.68)	8 (53.33)	6.00E-04
Chest Pain				
N (%)	96 (76.19)	110 (53.14)	7 (46.67)	7.78E-05
Angina Equivalent				
N (%)	46 (36.51)	53 (25.60)	5 (33.33)	0.1037
Shortness of Breath				
N (%)	28 (22.22)	30 (14.49)	7 (46.67)	0.0038
Family History of CAD				
N (%)	53 (42.06)	129 (62.32)	6 (40.00)	8.00E-04
Fasting				
N (%)	126 (100.00)	207 (100.00)	15 (100.00)	NA
Statin				
N (%)	39 (30.95)	125 (60.39)	13 (86.67)	2.28E-08
Niacin				
N (%)	1 (0.79)	3 (1.45)	0 (0.00)	0.7872

(continued)

	All Controls	Native CAD	Revasc CAD	P-value
Fibrate				
N (%)	4 (3.17)	7 (3.38)	1 (6.67)	0.7797
Ezetimibe				
N (%)	1 (0.79)	5 (2.42)	0 (0.00)	0.4745
Fish Oil				
N (%)	8 (6.35)	23 (11.11)	1 (6.67)	0.3251
Bile Acid Sequestrant				
N (%)	0 (0.00)	0 (0.00)	0 (0.00)	NA
Aspirin				
N (%)	28 (22.22)	99 (47.83)	9 (60.00)	4.91E-06
Clopidogrel				
N (%)	1 (0.79)	2 (0.97)	4 (26.67)	3.16E-11
Vitamin K Antagonist				
N (%)	6 (4.76)	3 (1.45)	1 (6.67)	0.1432
Nitrate				
N (%)	3 (2.38)	10 (4.83)	3 (20.00)	0.0084
Beta Blocker				
N (%)	36 (28.57)	77 (37.20)	12 (80.00)	4.00E-04
ACE Inhibitor				
N (%)	26 (20.63)	59 (28.50)	9 (60.00)	0.0039

II. Exploratory Data Analyses for Metabolites

[0194] Sample preparation and mass spectrometry analyses were conducted by Metabolon, Inc. The raw data contained a total of 1088 analytes, measured for 1096 pilot study participants.

[0195] Of the 1088 analytes (including unnamed metabolites and complex lipids), 481 named metabolites had less than 10% missing data. All 1096 patients had less than 10% missing data for these metabolites. Statistical analyses were therefore applied to 481 analytes and 1096 patients. The data was normalized in advance of receipt. A logarithm (base 2) transformation was applied and histograms were created to show the distribution of expression by analyte (data not shown).

[0196] The metabolomics data were generated in multiple batches; however, a principal components analysis (PCA) showed no evidence of any systematic site effects.

III. Prediction Modeling for Primary Endpoints

[0197] Methods. Patients in the Discovery-FAS Set were categorized according to whether they had fasted for at least eight hours. By this criteria, a total of 377 participants were Fasted and 371 were Non-Fasted. Association testing, with adjustment for age and gender was conducted for the four primary endpoints, and nominal associations were defined in three ways as follows:

1 Significant in Fasting and Non-Fasting combined

2 Significant in Fasting and Non-Fasting independently

3 Significant in Fasting alone

[0198] It is emphasized that, at this stage, 'significant' pertains to any association with raw, unadjusted $p < 0.05$.

[0199] In this way, twelve scenarios were considered as follows:

a) Atherosclerosis in Native CAD - A^{nCAD}

c. Significant in Fasting & Non-Fasting Combined - A_{FNF}^{nCAD}

c. Independently Significant in Fasting and Non-Fasting - A_{IFNF}^{nCAD}

c. Significant in Fasting - A_F^{nCAD}

b) Atherosclerosis in All CAD (including revascularization) - A^{aCAD}

c. Significant in Fasting & Non-Fasting Combined - A_{FNF}^{aCAD}

c. Independently Significant in Fasting and Non-Fasting - A_{IFNF}^{aCAD}

c. Significant in Fasting - A_F^{aCAD}

c) 50% stenosis in Native CAD - S^{nCAD}

c. Significant in Fasting & Non-Fasting Combined - S_{FNF}^{nCAD}

c. Independently Significant in Fasting and Non-Fasting - S_{IFNF}^{nCAD}

c. Significant in Fasting - S_F^{nCAD}

d) 50% stenosis in ALL CAD (including revascularization) - S^{aCAD}

c. Significant in Fasting & Non-Fasting Combined - S_{FNF}^{aCAD}

c. Independently Significant in Fasting and Non-Fasting - S_{IFNF}^{aCAD}

c. Analytes Significant in Fasting - S_F^{aCAD}

[0200] When more than 9 variables had $p < 0.05$, Age and Gender were added to the variables, and gradient boosting (see below) was applied to select 9 predictors.

[0201] Twelve prediction models were obtained by generalized linear (logistic) regression as follows. When fewer than nine variables had $p < 0.05$, Age and Gender were added to the variables, and the full model was fitted. Otherwise, the nine variables selected by gradient boosting variables were combined with Age and Gender in a generalized linear (logistic) model.

[0202] Gradient boosting is an approach to determine a regression function that minimizes the expectation of a loss function. (Freidman JH (2001) and Friedman JH (2002)) It is an iterative method, in which the negative gradient of the loss function is calculated, a regression model is fitted, the gradient descent step size is selected, and the regression function is updated. The gradient is approximated by means of a regression tree, which makes use of covariate information, and at each iteration the gradient determines the direction in which the function needs to move, in order to improve the fit to the data.

[0203] The loss function was assumed Bernoulli, due to the binary nature of the primary endpoints. A learning rate (λ) was introduced to dampen proposed moves and to protect against over-fitting. The optimal number of iterations, given by T , was determined by 5-fold cross-validation. The minimum number of observations in each terminal node was 10. Two-way interactions were allowed. Random sub-sampling, without replacement, of half of the observations was applied to achieve variance reduction in gradient estimation.

[0204] For current purposes, 50 rounds of gradient boosting were run for each scenario, and the nine variables most often showing highest estimated relative influence were taken forwards to generalized linear modeling.

[0205] The twelve models were used to generate probability predictions for each patient in the Validation-FAS. For each model, the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) were calculated for the range of predicted probability thresholds. A Receiver Operating Characteristic (ROC) curve was generated to plot sensitivity as a function of (1-specificity). The optimal classification threshold was determined on the basis of accuracy, defined as the proportion of correct predictions. In addition, the Area Under the Curve (AUC) and accuracy was estimated (Tables 27, 28, 29, 30 for the four primary endpoints, respectively).

[0206] The performance of model-based predictions were compared to the performance of probability predictions obtained by Diamond-Forrester scoring. (Diamond and Forrester (1979)).

Detailed Results for Native CAD

[0207] The results show that the Diamond-Forrester score provides poor prediction of the GLOBAL phenotypes (Figures 34, 38, 42, 46). The estimates of AUC and accuracy for prediction of Native CAD indicate that performance is no better than assigning all patients as 'at risk' of disease, by which 62% of predictions in the Validation Set for Native CAD (*Validation-Native CAD plus Validation-Control*) are correct, and 64% of predictions in the Validation Set for All CAD (*Validation-Native CAD plus Validation-Revasc. CAD plus Validation-Control*), are correct.

Metabolomics Model**[0208]**

I. Atherosclerosis in Native CAD - A^{nCAD}

a. Significant in Fasting & Non-Fasting Combined - A_{FNF}^{nCAD}

i. Of the 481 analytes measured, 83 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for

A_{FNF}^{nCAD} . Table 29 provides a list of the 83 metabolomic variables for A_{FNF}^{nCAD} .

Table 29:

glutamate	serylleucine	1-linoleoyl-GPC (18:2)
acisoga	3-methoxytyrosine	1-methylguanosine
threonate	prolylhydroxyproline	12 13-DiHOME
urate	valerylcarnitine (C5)	O-sulfo-L-tyrosine
mannose	caproate (6:0)	erucamide
oleic ethanolamide	tigloylglycine	inositol 1-phosphate (I1P)
cysteine-glutathione disulfide	guanidinosuccinate	isoleucylvaline
pyroglutamylglutamine	isobutyrylglycine (C4)	gamma-tocopherol
valylleucine	glycocholate sulfate*	1-eicosenoylglycerophosphocholine (20:1n9)*
butyrylcarnitine (C4)	o-cresol sulfate	tyrosylglutamine
cytidine	N-acetylthreonine	indolepropionate
palmitoyl ethanolamide	leucylglycine	gamma-glutamylvaline
phenylalanylvaline	2-hydroxybutyrate (AHB)	2-aminoadipate
hydroxybutyrylcarnitine*	leucylaspartate	aspartate
1-nonadecanoylglycerophosphocholine (19:0)	1-arachidoyl-GPC (20:0)	N6-carbamoylthreonyladenine
glycine	N6-methyladenosine	methyl glucopyranoside (alpha + beta)
propionylglycine (C3)	hexanoylcarnitine (C6)	myo-inositol
pseudouridine	valylisoleucine	alpha-ketobutyrate
ADSGEGDFAEGGGVR*	beta-alanine	S-adenosylhomocysteine (SAH)
2-hydroxyhippurate (salicylurate)	1-linoleoyl-GPE (18:2)*	1-oleoylglycerol (18:1)
alpha-glutamyltyrosine	gamma-glutamylglutamate	tartronate (hydroxymalonate)
fucose	3-hydroxy-2-ethylpropionate	3-methylglutaryl carnitine-2
glucuronate	adenine	1-methylurate
3-methylglutaryl carnitine-1	xylitol	N-acetyl-beta-alanine
xanthine	N2 N2-dimethylguanosine	histidyltryptophan

(continued)

12-HETE	methyl indole-3-acetate	1-oleoyl-GPC (18:1)*
glucose	homostachydrine*	3-hydroxydecanoate
salicylate	phenylacetylglutamine	--

Of the 83 metabolomic variables exhibiting a nominal univariate association for A_{FNF}^{nCAD} , a panel of eight metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 30 provides the relative influence of the eight metabolomic variables, in combination with age and gender, for the

Metabolomics Model of A_{FNF}^{nCAD} . **Figure 35** provides a ROC curve for the Metabolomics Model of A_{FNF}^{nCAD} . Table 53 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 30:

Variable	Relative Influence	Direction of Change
valylleucine	28.88	Decreased
glutamate	14.47	Elevated
acisoga	14.25	Elevated
urate	10.39	Elevated
glucuronate	9.26	Elevated
age	6.68	Elevated
fucose	6.18	Elevated
Butyrylcarnitine (C4)	4.72	Elevated
mannose	4.46	Elevated
Male gender	0.70	Present§
§ The term "present" conveys that male gender was taken into account in the prediction model, with 'relative influence' denoting the association of male gender with the outcome (i.e., ASCAD or the presents of a coronary atherosclerotic plaque).		

b. Independently Significant in Fasting and Non-Fasting - A_{IFNF}^{nCAD}

i. Of the 481 analytes measured, 4 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for A_{IFNF}^{nCAD} . Table 31 provides a list of the 4 metabolomic variables for A_{IFNF}^{nCAD} .

Table 31:

acisoga	o.cresol.sulfate	Cysteine.glutathione.disulfide
threonate	--	--

Of the 4 metabolomic variables exhibiting a nominal univariate association for A_{IFNF}^{nCAD} , a panel of all four metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 32 provides the relative influence of the four metabolomic variables, in combination with age and gender, for the

Metabolomics Model of A_{IFNF}^{nCAD} . **Figure 36** provides a ROC curve for the Metabolomics Model of A_{IFNF}^{nCAD} . Table 53 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 32:

Variable	Relative Influence	Direction of Change
acisoga	40.74	Elevated
age	20.77	Elevated
cysteine.glutathione.disulfide	18.87	Decreased
threonate	12.67	Decreased
o.cresol.sulfate	4.49	Elevated
Male gender	2.45	Present

c. Significant in Fasting - A_F^{nCAD}

i. Of the 481 analytes measured, 34 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for A_F^{nCAD} . Table 33 provides a list of the 34 metabolomic variables for A_F^{nCAD} .

Table 33:

xylitol	valylleucine	cysteine-glutathione disulfide
3-carboxy-4-methyl-5-propyl-2-furanpropanoate (CMPF)	N-acetylleucine	threonate
serylleucine	alpha-glutamyltyrosine	fucose
phenylalanylvaline	4-androsten-3alpha 17alradiol monosulfate 2	adenosine
12-HETE	inositol 1-phosphate (I1P)	valylisoleucine
glycocholenate sulfate*	1-docosaheptaenoyl-GPC* (22:6)*	phenylalanylserine
oleic ethanolamide	2-hydroxyhippurate (salicylurate)	gamma-tocopherol
acisoga	salicylate	palmitoyl ethanolamide
leucylglycine	phenylalanylglycine	hydroquinone sulfate
N-acetylphenylalanine	glycylphenylalanine	propionylglycine (C3)
o-cresol sulfate	3-methoxytyrosine	--
histidyltryptophan	adenine	--

Of the 34 metabolomic variables exhibiting a nominal univariate association for A_F^{nCAD} , a panel of eight metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 34 provides the relative influence of the eight metabolomic variables, in combination with age and gender, for the

Metabolomics Model of A_F^{nCAD} . **Figure 37** provides a ROC curve for the Metabolomics Model of A_F^{nCAD} . Table 53 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 34:

Variable	Relative Influence	Direction of Change
N.acetylphenylalanine	22.27	Elevated
age	18.18	Elevated
valylleucine	17.61	Decreased
xylitol	8.07	Elevated

(continued)

Variable	Relative Influence	Direction of Change
2.hydroxyhippurate (salicylurate)	6.97	Elevated
N.acetylleucine	6.15	Elevated
serylleucine	6.15	Decreased
fucose	6.06	Elevated
glycylphenylalanine	4.97	Decreased
Male gender	3.56	Present

II. Atherosclerosis in All CAD (inc revasc) - A^{aCAD} a. Significant in Fasting & Non-Fasting Combined - A_{FNF}^{aCAD} i. Of the 481 analytes measured, 92 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for A_{FNF}^{aCAD} . Table 35 provides a filtered list of the 92 metabolomic variables for A_{FNF}^{aCAD} .

Table 35:

acisoga	tyrosylglutamine	N6-carbamoylthreonyladen osine
Glutamate	xanthine	2-linoleoyl-GPC* (18:2)*
Threonate	beta-alanine	3-methyl-2-oxobutyrate
Mannose	isobutyrylglycine (C4)	methyl glucopyranoside (alpha + beta)
Urate	3-methylglutaryl carnitine-1	serylleucine
cysteine-glutathione disulfide	valerylcarnitine (C5)	caproate (6:0)
oleic ethanolamide	1-linoleoyl-GPC (18:2)	N-methyl proline
pyroglutamylglutamine	hexanoylcarnitine (C6)	laurylcarnitine (C12)*
butyrylcarnitine (C4)	2-hydroxybutyrate (AHB)	o-cresol sulfate
Cytidine	1-arachidoyl-GPC (20:0)	gamma-glutamylglutamate
hydroxybutyrylcarnitine*	guanidinosuccinate	N-acetyl-beta-alanine
alpha-glutamyltyrosine	fucose	1-eicosenoylglycerophosphocholine (20:1n9)*
2-hydroxyhippurate (salicylurate)	phenylacetylglutamine	N-acetyl glycine
Valylleucine	3-methylglutaryl carnitine-2	seryltyrosine
propionylglycine (C3)	glycohyocholate	4-guanidinobutanoate
1-nonadecanoylglycerophosphocholine (19:0)	N6-methyladenosine	S-methylcysteine
Glycine	N2 N2-dimethylguanosine	isoleucylvaline
12-HETE	gamma-glutamylvaline	adenine
pseudouridine	leucylaspartate	1-methylurate
Salicylate	2-hydroxyoctanoate	xylitol
Glucose	alpha-ketobutyrate	phenylalanylalanine
ADSGEGDFXAEGGGVR*	glycocholate sulfate*	O-sulfo-L-tyrosine
1-linoleoyl-GPE (18:2)*	valylisoleucine	erucamide
Phenylalanylvaline	homostachydrine*	pregnanediol-3-glucuronide

(continued)

Tigloylglycine	methyl indole-3-acetate	3-hydroxy-2-ethylpropionate
Glucuronate	leucylglycine	pyridoxal
palmitoyl ethanolamide	N-acetylthreonine	1-oleoyl-GPC (18:1)*
1-oleoylglycerol (18:1)	2-hydroxydecanoate	2prime-deoxyuridine
1213-DiHOME	1-methylguanosine	threonylphenylalanine
3-methoxytyrosine	prolylhydroxyproline	2-aminoadipate
2-linoleoyl-GPE* (18:2)*	prolylglycine	--

Of the 92 metabolomic variables exhibiting a nominal univariate association for A_{FNF}^{aCAD} , a panel of eight metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 36 provides the relative influence of the eight metabolomic variables combined with age and gender for the

Metabolomics Model of A_{FNF}^{aCAD} . **Figure 39** provides a ROC curve for the Metabolomics Model of A_{FNF}^{aCAD} . Table 54 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 36:

Variable	Relative Influence	Direction of Change
valylleucine	26.79	Decreased
acisoga	16.87	Elevated
glutamate	13.93	Elevated
urate	9.74	Elevated
glucuronate	8.74	Elevated
mannose	7.13	Elevated
age	6.24	Elevated
12.HETE	5.03	Decreased
Valerylcarnitine (C5)	4.81	Elevated
Male gender	0.72	Present

b. Independently Significant in Fasting and Non-Fasting - A_{IFNF}^{aCAD}

i. Of the 481 analytes measured, 6 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for

A_{IFNF}^{aCAD} . Table 37 provides a list of the 6 metabolomic variables for A_{IFNF}^{aCAD} .

Table 37:

threonate	threonate	cysteine-glutathione disulfide
o-cresol sulfate	1-nonadecanoylglycerophosphocholine (19:0)	glucose

Of the 6 metabolomic variables exhibiting a nominal univariate association for A_{IFNF}^{aCAD} , a panel of all six metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 38 provides the relative influence of the six metabolomic variables, in combination with age and gender, for the

Metabolomics Model of A_{IFNF}^{aCAD} . **Figure 40** provides a ROC curve for the Metabolomics Model of A_{IFNF}^{aCAD} . Table 54 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of

predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 38:

Variable	Relative Influence	Direction of Change
acisoga	39.32	Elevated
age	17.31	Elevated
1.nonadecanoylglycerophosphocholine (19:0)	12.00	Decreased
cysteine.glutathione.disulfide	10.91	Decreased
threonate	10.71	Decreased
glucose	6.64	Elevated
Male gender	2.06	Present
o.cresol.sulfate	1.05	Elevated

c. Significant in Fasting - A_F^{aCAD}

i. Of the 481 analytes measured, 48 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for

A_F^{aCAD} . Table 39 provides a list of the 48 metabolomic variables for A_F^{aCAD} .

Table 39:

12-HETE	N-acetylphenylalanine	1-arachidonylglycerol
alpha-glutamyltyrosine	N-acetylleucine	Pyroglutamylvaline
salicylate	4-androsten-3alpha 17alphiadiol monosulfate 2	phenylalanyltryptophan
2-hydroxyhippurate (salicylurate)	o-cresol sulfate	methyl indole-3-acetate
acisoga	phenylalanylvaline	Histidyltryptophan
3-carboxy-4-methyl-5-propyl-2-furanpropanoate (CMPF)	leucylglycine	4-ethylphenyl sulfate
threonate	phenylalanylglycine	1-myristoylglycerol (14:0)
glycocholenate sulfate*	propionylglycine (C3)	inositol 1-phosphate (I1P)
xylitol	mannitol	1-nonadecanoylglycerophosphocholine(19:0)
1-docosaheptaenoyl-GPC* (22:6)*	serylleucine	Glucose
phenylalanyls erine	hydroquinone sulfate	N-stearoyltaurine
3-methoxytyrosine	adenosine	Valylisoleucine
oleic ethanolamide	2-hydroxydecanoate	beta-alanine
cysteine-glutathione disulfide	tyrosylglutamine	N-acetylglycine
glycylphenylalanine	N-octanoylglycine	Allantoin
valylleucine	adenine	Phenylalanylphenylalanine

Of the 48 metabolomic variables exhibiting a nominal univariate association for A_F^{aCAD} , a panel of seven metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 40 provides the relative influence of the seven metabolomic variables, in combination with age and gender, for

the Metabolomics Model of A_F^{aCAD} . **Figure 41** provides a ROC curve for the Metabolomics Model of A_F^{aCAD} . Table

54 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 40:

Variable	Relative Influence	Direction of Change
age	21.21	Elevated
valylleucine	20.76	Decreased
N.acetylphenylalanine	18.59	Elevated
2.hydroxyhippurate (salicylurate)	12.20	Elevated
N.acetylleucine	6.10	Elevated
12.HETE	5.96	Decreased
xylitol	5.40	Elevated
glycylphenylalanine	5.39	Decreased
Male gender	4.40	Present

III. 50% stenosis in Native CAD - S^{nCAD}

a. Significant in Fasting & Non-Fasting Combined - S_{FNF}^{nCAD}

i. Of the 481 analytes measured, 49 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for

S_{FNF}^{nCAD} . Table 41 provides a list of the 49 metabolomic variables for S_{FNF}^{nCAD} .

Table 41:

threonate	serotonin (5HT)	5alpha-androstan-3alpha 17beta-diol disulfate
N-acetylglycine	xanthine	1-stearoyl-GPC (18:0)
glycerate	2-oleoyl-GPE* (18:1)*	serine
isobutyrylglycine (C4)	4-guanidinobutanoate	acisoga
valerylcarnitine (C5)	leucylleucine	mannose
fumarate	cholate	valylleucine
1-nonadecanoylglycerophosphocholine(19:0)	propionylglycine (C3)	gamma-tocopherol
tartronate (hydroxymalonate)	glycocholate	3-ethylphenylsulfate
2-hydroxyhippurate (salicylurate)	N-octanoylglycine	glutamate
1-arachidoyl-GPC (20:0)	glycoursodeoxycholate	sphingosine 1-phosphate
threitol	isovaleryllycine	carnitine
N-(2-furoyl)glycine	pregnanediol-3-glucuronide	arabonate
tigloylglycine	5alpha-androstan-3beta 17beta-diol monosulfate 2	cyclo(leu-pro)
salicylate	arabinose	indoleacetylglutamine
N-acetylthreonine	1-linoleoyl-GPE (18:2)*	prolylglycine
xylonate	5-HETE	--
xylose	hydroquinone sulfate	--

Of the 49 metabolomic variables exhibiting a nominal univariate association for S_{FNF}^{nCAD} , a panel of eight metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 42 provides the relative influence of the eight metabolomic variables, in combination with age and gender, for the

Metabolomics Model of S_{FNF}^{nCAD} . **Figure 43** provides a ROC curve for the Metabolomics Model of S_{FNF}^{nCAD} . Table 55 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 42:

Variable	Relative Influence	Direction of Change
Age	37.04	Elevated
valerylcarnitine (C5)	14.32	Elevated
N.acetylthreonine	10.39	Elevated
tigloylglycine	8.72	Decreased
2.hydroxyhippurate (salicylurate)	7.06	Elevated
glycerate	6.42	Decreased
salicylate	5.67	Decreased
threonate	5.58	Decreased
tartronate (hydroxymalonated);	4.25	Elevated
Male gender	0.55	Present

b. Independently Significant in Fasting and Non-Fasting - S_{IFNF}^{nCAD}

i. Of the 481 analytes measured, 2 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for S_{IFNF}^{nCAD} . Table 43 provides a list of the 2 metabolomic variables for S_{IFNF}^{nCAD} .

Table 43:

N-acetylglycine	3-ethylphenylsulfate	--
-----------------	----------------------	----

Of the 2 metabolomic variables exhibiting a nominal univariate association for S_{IFNF}^{nCAD} , a panel of both variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 44 provides the relative influence of the two metabolomic variables in combination with age and gender for the Metabolomics Model

of S_{IFNF}^{nCAD} . **Figure 44** provides a ROC curve for the Metabolomics Model of S_{IFNF}^{nCAD} . Table 55 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 44:

Variable	Relative Influence	Direction of Change
age	67.33	Elevated
N.acetylglycine	14.67	Decreased
3.ethylphenylsulfate	12.88	Elevated
Male gender	5.12	Elevated

c. Significant in Fasting - S_F^{nCAD}

i. Of the 481 analytes measured, 28 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for

S_F^{nCAD} . Table 45 provides a filtered list of the 28 metabolomic variables for S_F^{nCAD} .

Table 45:

leucylleucine	valylisoleucine	7-methylguanine
asparagine	glycocholate sulfate*	cyclo(leu-pro)
glycerate	arabitol	Methionine
threitol	N-acetylglycine	propionylglycine (C3)
cholate	serotonin (5HT)	Serine
N-octanoylglycine	xylose	2-oleoyl-GPE* (18:1)*
xylolate	N-acetylputrescine	Tigloylglycine
isobutyrylglycine (C4)	arabonate	3-ethylphenylsulfate
isovaleryl-glycine	lysine	--
fumarate	N-(2-furoyl)glycine	--

Of the 28 metabolomic variables exhibiting a nominal univariate association for S_F^{nCAD} , a panel of eight metabolomic variables were selected as best predictors; they were combined with age and gender in a prediction model for CAD. Table 46 provides the relative influence of the eight metabolomic variables, in combination with age and gender, for the

Metabolomics Model of S_F^{nCAD} . Figure 45 provides a ROC curve for the Metabolomics Model of S_F^{nCAD} . Table 55 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 46:

Variable	Relative Influence	Direction of Change
age	24.44	Elevated
leucylleucine	21.83	Decreased
serotonin (5HT)	11.37	Elevated
N.acetylputrescine	9.68	Decreased
glycocholate.sulfate	8.56	Decreased
propionylglycine (C3)	6.95	Decreased
cholate	6.14	Decreased
asparagine	5.73	Elevated
3.ethylphenylsulfate	4.79	Elevated
Male gender	0.50	Present

IV. 50% stenosis in ALL CAD (inc revasc) - S^{aCAD}

a. Significant in Fasting & Non-Fasting Combined - S_{FNF}^{aCAD}

i. Of the 481 analytes measured, 72 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for S_{FNF}^{aCAD} . Table 47 provides a list of the 72 metabolomic variables for S_{FNF}^{aCAD} .

Table 47:

threonate	prolylglycine	2-hydroxydecanoate
1-linoleoyl-GPE (18:2)*	N-octanoylglycine	glutamate

(continued)

	N-acetylglycine	threitol	N-acetylthreonine
5	glycoursodeoxycholate	fumarate	taurine
	2-hydroxyhippurate (salicylurate)	pregnanediol-3-glucuronide	1-oleoylplasmenylethanol amine*
	salicylate	1-oleoyl-GPI (18:1)*	1-palmitoyl-GPE (16:0)
	2-linoleoyl-GPE* (18:2)*	serotonin (5HT)	N-acetylglutamate
10	mannose	xylionate	13-HODE + 9-HODE
	tigloylglycine	cyclo(leu-pro)	1-palmitoyl-GPI* (16:0)*
	2-linolenoylglycerophosphocholine(18: 3n3)*	glycerate	hydroquinone sulfate
15	1-nonadecanoylglycerophosphocholine (19: 0)	tartrate (hydroxymalonnate)	caprylate (8:0)
	2-oleoyl-GPE* (18:1)*	xylose	1-stearoyl-GPC (18:0)
	isovaleryl glycine	glycohyocholate	glycochenodeoxycholate
20	isobutyrylglycine (C4)	glucose	p-cresol sulfate
	N-(2-furoyl)glycine	xanthine	12-HETE
	glycocholate	cyclo(L-phe-L-pro)	5-hydroxyindoleacetate
	acisoga	beta-alanine	arabonate
25	4-guanidinobutanoate	pyridoxate	2-hydroxyoctanoate
	1-arachidoyl-GPC (20:0)	tartrate	urate
	propionyl glycine (C3)	1-linoleoyl-GPC (18:2)	valylleucine
30	valeryl carnitine (C5)	pyridoxal	carnitine
	1-oleoylglycerol (18:1)	cholate	1-linoleoyl-GPI* (18:2)*
	1-oleoyl-GPE (18:1)	serine	N-acetylputrescine
35	arabinose	homostachydrine*	succinate

Of the 72 metabolomic variables exhibiting a nominal univariate association for S_{FNF}^{aCAD} , a panel of eight metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 48 provides the relative influence of the eight metabolomic variables in combination with age and gender for the Metabolomics Model of S_{FNF}^{aCAD} . **Figure 47** provides a ROC curve for the Metabolomics Model of S_{FNF}^{aCAD} . Table 56 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 48:

Variable	Relative Influence	Direction of Change
Age	18.38	Elevated
glycoursodeoxycholate	16.52	Decreased
acisoga	12.81	Elevated
2.hydroxyhippurate (salicylurate)	10.33	Elevated
1.linoleoyl.GPE (18:2)	10.26	Decreased
valeryl carnitine (C5),	8.91	Elevated
threonate	7.13	Decreased

(continued)

Variable	Relative Influence	Direction of Change
mannose	7.12	Elevated
salicylate	7.02	Elevated
Male gender	1.52	Present

b. Independently Significant in Fasting and Non-Fasting - S_{IFNF}^{aCAD}
 i. Of the 481 analytes measured, 5 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for S_{IFNF}^{aCAD} . Table 49 provides a filtered list of the 5 metabolomic variables for S_{IFNF}^{aCAD} .

Table 49:

N-acetylglycine	threonate	Salicylate
2-hydroxyhippurate (salicylurate)	3-ethylphenylsulfate	--

Of the 5 metabolomic variables exhibiting a nominal univariate association for S_{IFNF}^{aCAD} , a panel of all five metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 50 provides the relative influence of the five metabolomic variables in combination with age and gender for the

Metabolomics Model of S_{IFNF}^{aCAD} . **Figure 48** provides a ROC curve for the Metabolomics Model of S_{IFNF}^{aCAD} . Table 56 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 50:

Variable	Relative Influence	Direction of Change
Age	40.82	Elevated
2-hydroxyhippurate (salicylurate)	19.56	Elevated
threonate	14.84	Decreased
salicylate	12.23	Elevated
Male gender	7.00	Present
N.acetylglycine	3.13	Decreased
3-ethylphenylsulfate	2.42	Elevated

c. Analytes Significant in Fasting - S_F^{aCAD}

i. Of the 481 analytes measured, 40 metabolomic variables exhibited a nominal univariate association (raw $p < 0.05$) for S_F^{aCAD} . Table 51 provides a filtered list of the 40 metabolomic variables for S_F^{aCAD} .

Table 51:

N-octanoylglycine	salicylate	Dimethylglycine
1-oleoylglycerol (18:1)	7-methylguanine	xylonite
isovalerylglycine	lysine	Phenylalanylphenylalanine
N-acetylglycine	glycoursodeoxycholate	Valylisoleucine
2-linolenoylglycerophosphocholine(18: 3n3)*	3-indoxyl sulfate	Glycerate
asparagine	6-oxopiperidine-2-carboxylic acid	1-arachidonylglycerol

(continued)

isobutyrylglycine (C4)	1-arachidonoylglycerophosphate	Fumarate
cyclo(leu-pro)	2-hydroxyhippurate (salicylurate)	3-ethylphenylsulfate
cholate	threitol	7-HOCA
serotonin (5HT)	methionine	Taurine
threonate	acisoga	Cholesterol
N-acetylputrescine	tigloylglycine	Arabitol
propionylglycine (C3)	1-linoleoylglycerol (18:2)	--
2-oleoyl-GPE* (18:1)*	1-oleoyl-GPI (18:1)*	--

Of the 40 metabolomic variables exhibiting a nominal univariate association for S_F^{aCAD} , a panel of eight metabolomic variables were selected as best predictors; these were combined with age and gender in a prediction model for CAD. Table 52 provides the relative influence of the eight metabolomic variables in combination with age and gender for the

Metabolomics Model of S_F^{aCAD} . **Figure 49** provides a ROC curve for the Metabolomics Model of S_F^{aCAD} . Table 56 provides the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) for the range of predicted probability thresholds; Area Under the Curve (AUC) and accuracy was estimated.

Table 52:

Variable	Relative Influence	Direction of Change
age	15.37	Elevated
cholesterol	15.19	Decreased
1.oleoylglycerol (18:1)	15.12	Elevated
acisoga	14.01	Elevated
2.hydroxyhippurate (salicylurate)	9.47	Elevated
asparagine	8.18	Elevated
taurine	7.93	Decreased
6.oxopiperidine.2.carboxylic.acid	7.50	Elevated
propionylglycine (C3)	6.66	Decreased
Male gender	0.56	Present

For each model below, the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) were calculated for the range of predicted probability thresholds (Tables 53, 54, 55, 56). A Receiver Operating Characteristic (ROC) curve was generated to plot sensitivity as a function of (1-specificity). The optimal classification threshold was determined on the basis of accuracy, defined as the proportion of correct predictions. In addition, the Area Under the Curve (AUC) and accuracy was estimated (Tables 53, 54, 55, 56 for Native CAD, All CAD, 50% stenosis in Native CAD, and 50% stenosis in All CAD, respectively). The first row for each model indicates the performance of the maximum accuracy threshold, the optimal balance between sensitivity and specificity. Those models with a second row were optimized for a high negative predictive value (NPV).

Table 53:

Model	AUC	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Accuracy
DF	0.45	1.00	0.00	0.62	N/A	0.62
A_{FNF}^{nCAD}	0.82	0.85	0.61	0.78	0.71	0.76
		0.99	0.09	0.64	0.92	0.65

EP 3 242 588 B1

(continued)

Model	AUC	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Accuracy
A_{IFNF}^{nCAD}	0.80	0.85	0.64	0.80	0.72	0.77
		1.00	0.10	0.65	0.93	0.66
A_F^{aCAD}	0.81	0.87	0.58	0.77	0.74	0.76
		0.99	0.26	0.69	0.94	0.72
DF = Diamond-Forrester						

Table 54:

Model	AUC	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Accuracy
DF	0.45	1.00	0.00	0.64	N/A	0.64
A_{FNF}^{aCAD}	0.83	0.93	0.50	0.77	0.81	0.78
		1.00	0.16	0.68	0.95	0.69
A_{IFNF}^{aCAD}	0.81	0.85	0.66	0.81	0.71	0.78
		1.00	0.07	0.65	0.90	0.66
A_F^{aCAD}	0.82	0.83	0.64	0.80	0.68	0.76
		1.00	0.16	0.68	0.95	0.69
DF = Diamond-Forrester						

Table 55:

Model	AUC	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Accuracy
DF	0.45	0.03	1.00	1.00	0.78	0.78
S_{FNF}^{nCAD}	0.73	0.21	0.96	0.64	0.80	0.79
		0.95	0.30	0.29	0.95	0.45
S_{IFNF}^{nCAD}	0.76	0.30	0.96	0.72	0.82	0.81
		0.93	0.37	0.31	0.95	0.50
S_F^{nCAD}	0.67	0.04	1.00	1.00	0.78	0.78
		0.96	0.20	0.26	0.95	0.38
DF = Diamond-Forrester						

Table 56:

Model	AUC	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Accuracy
DF	0.45	0.03	1.00	1.00	0.74	0.75
S_{FNF}^{aCAD}	0.78	0.34	0.94	0.66	0.80	0.78

(continued)

Model	AUC	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Accuracy
		0.96	0.30	0.33	0.95	0.47
S_{IFNF}^{aCAD}	0.78	0.23	0.97	0.72	0.78	0.77
		0.96	0.30	0.33	0.95	0.47
S_F^{aCAD}	0.74	0.22	0.96	0.69	0.78	0.77
		0.97	0.22	0.31	0.95	0.42
DF = Diamond-Forrester						

Claims

1. A method for evaluating a human subject for having atherosclerotic coronary artery disease (ASCAD) or as having a coronary atherosclerotic plaque, the method comprising:

measuring the levels of each analyte within a panel of analyte biomarkers in a biological sample obtained from a subject, wherein the panel of analyte biomarkers comprises valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4) and mannose;

comparing the measured levels of the analytes in the biological sample with one or more reference samples, wherein the references are representative of matched human subjects; and

identifying the subject as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque if measured levels of the analytes in the biological sample are increased or decreased relative to the amounts of the analytes in the reference samples.
2. The method of claim 1, wherein the levels are measured using mass spectrometry (MS) analysis, optionally wherein the MS analysis comprises liquid chromatography-tandem mass spectrometry (UPLC-MS/MS) or gas chromatography-mass spectrometry (GC-MS).
3. The method of any one of claims 1 to 2, wherein the CAD comprises atherosclerosis, optionally wherein the subject presents with symptoms of atherosclerotic CAD comprising chest pain, angina, angina equivalent, dyspnea, or dyspnea on exertion, and optionally wherein the subject has no previous history of coronary artery disease.
4. The method of any one of claims 1 to 3, wherein the biological sample is a blood sample, a serum sample or a plasma sample.
5. The method of any one of claims 1 to 4, wherein the subject presents with risk factors associated with coronary artery disease selected from the group consisting of male gender, hypertension, dyslipidemia, diabetes, and a family history of coronary artery disease.
6. The method of any one of claims 1 to 5, wherein the reference sample is obtained from at least one individual not suffering from a cardiovascular disease, or wherein the reference sample comprises predetermined, statistically significant reference analyte levels.
7. The method of any one of claims 1 to 6, further comprising modifying the subject's clinical record to identify the subject as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque, optionally wherein the clinical record is stored in a computer readable medium.
8. The method of claim 1, wherein the identifying step is performed using a gradient boosting algorithm, a decision tree model, or a linear regression analyses.
9. A kit for evaluating a human subject for having atherosclerotic coronary artery disease (ASCAD) or as having a coronary atherosclerotic plaque, the kit comprising:

reagents suitable for determining levels of a plurality of analytes in a test sample, wherein the plurality of analytes comprise a panel of biomarkers comprising valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4) and mannose; and instructions for use of the kit in the method of any of claims 1 to 8.

5
10. The kit of claim 9 further comprising:

one or more control samples comprising predetermined levels of the same analytes, wherein comparison of the levels of the analytes in a test sample with levels in the control samples identifies a subject as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque.

10
11. A composition for use in a method of treating a subject having atherosclerotic coronary artery disease (ASCAD) or having a coronary atherosclerotic plaque, the method comprising:

15 measuring the levels of each analyte within a panel of analyte biomarkers in a biological sample obtained from a subject, wherein the panel of analyte biomarkers comprises valylleucine, glutamate, acisoga, urate, glucuronate, fucose, butyrylcarnitine (C4) and mannose;
comparing the measured levels of the analytes in the biological sample with one or more reference samples, wherein the references are representative of matched human subjects;
20 identifying the subject as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque if measured levels of the analytes in the biological sample are increased or decreased relative to the amounts of the analytes in the reference samples; and
selectively administering a composition comprising an effective amount of a therapeutic agent selected from the group consisting of a statin, cholesterol absorption inhibitors, niacin-derivatives, omega-3-fatty acid compounds, bile acid sequestrants, anti-platelet agents, aldosterone blockers, angiotensin-converting enzyme
25 (ACE) inhibitors, angiotensin-receptor blockers (ARBs), beta blockers, diuretics, digitalis, hydralazine and nitrates, warfarin and aspirin to the subject identified as having atherosclerotic coronary artery disease or as having a coronary atherosclerotic plaque to treat the subject.

30
12. The composition for use of claim 11, wherein the therapeutic agent is a statin.

Patentansprüche

35
1. Verfahren zur Evaluierung eines menschlichen Individuums auf eine atherosklerotische Koronararterienerkrankung (ASCAD) oder auf atherosklerotischen Koronar-Plaque, wobei das Verfahren das Folgende umfasst:

Messen der Spiegel jedes Analyten innerhalb einer Gruppe von Analyten-Biomarkern in einer biologischen Probe, die von einem Individuum erhalten wurde, wobei die Gruppe von Analyten-Biomarkern Valylleucin, Glutamat, Acisoga, Urat, Glucuronat, Fucose, Butyrylcarnitin (C4) und Mannose umfasst;
40 Vergleichen der gemessenen Spiegel der Analyten in der biologischen Probe mit einer oder mehreren Bezugsproben, wobei die Bezugsproben für passende menschliche Individuen repräsentativ sind; und
Identifizieren des Individuums als Individuum mit einer atherosklerotischen Koronararterienerkrankung oder mit atherosklerotischen Koronar-Plaque, wenn gemessene Spiegel der Analyten in der biologischen Probe bezüglich der Mengen der Analyten in den Bezugsproben erhöht oder verringert sind.

45
2. Verfahren nach Anspruch 1, wobei die Spiegel mittels Massenspektrometrie (MS) -Analyse gemessen werden, wobei fakultativ die MS-Analyse Flüssigkeitschromatographie-Tandem-Massenspektrometrie (UPLC-MS/MS) oder Gaschromatographie-Massenspektrometrie (GC-MS) umfasst.

50
3. Verfahren nach einem der Ansprüche 1 bis 2, wobei die Koronararterienerkrankung Atherosklerose umfasst, wobei fakultativ das Individuum Symptome einer atherosklerotischen Koronararterienerkrankung zeigt, einschließlich Brustschmerzen, Angina, Angina-Äquivalent, Dyspnoe oder Dyspnoe bei Anstrengung, und wobei das Individuum fakultativ keine Koronararterienerkrankung in der Anamnese hat.

55
4. Verfahren nach einem der Ansprüche 1 bis 3, wobei die biologische Probe eine Blutprobe, eine Serumprobe oder eine Plasmaprobe ist.

5. Verfahren nach einem der Ansprüche 1 bis 4, wobei das Individuum Risikofaktoren im Zusammenhang mit Koro-

nararterienerkrankung aufweist, die aus der aus männliches Geschlecht, Hypertonie, Dyslipidämie, Diabetes und Koronararterienerkrankung in der Familienanamnese bestehenden Gruppe ausgewählt sind.

6. Verfahren nach einem der Ansprüche 1 bis 5, wobei die Bezugsprobe von mindestens einem Individuum erhalten wurde, das nicht an einer kardiovaskulären Erkrankung leidet, oder wobei die Bezugsprobe vorgegebene statistisch signifikante Bezugsanalytenspiegel umfasst.

7. Verfahren nach einem der Ansprüche 1 bis 6, ferner umfassend das Modifizieren der Krankenakte des Individuums zur Identifizierung des Individuums als Individuum mit einer atherosklerotischen Koronararterienerkrankung oder mit atherosklerotischem Koronar-Plaque, wobei fakultativ die Krankenakte in einem computerlesbaren Medium gespeichert wird.

8. Verfahren nach Anspruch 1, wobei der Identifizierungsschritt unter Verwendung eines Gradient-Boosting-Algorithmus, eines Entscheidungsbaum-Modells oder einer linearen Regressionsanalyse durchgeführt wird.

9. Kit zur Evaluierung eines menschlichen Individuums auf eine atherosklerotische Koronararterienerkrankung (ASCAD) oder auf atherosklerotischen Koronar-Plaque, wobei das Kit das Folgende umfasst:

Reagenzien, die zur Bestimmung von Spiegeln einer Vielzahl von Analyten in einer Testprobe geeignet sind, wobei die Vielzahl von Analyten eine Gruppe von Biomarkern umfasst, umfassend Valylleucin, Glutamat, Acisoga, Urat, Glucuronat, Fucose, Butyrylcarnitin (C4) und Mannose; und Anweisungen zur Verwendung des Kits im Verfahren nach einem der Ansprüche 1 bis 8.

10. Kit nach Anspruch 9, ferner umfassend:

eine oder mehrere Kontrollproben, die vorgegebene Spiegel der gleichen Analyten umfassen, wobei ein Vergleich der Spiegel der Analyten in einer Testprobe mit Spiegeln in den Kontrollproben ein Individuum als Individuum mit einer atherosklerotischen Koronararterienerkrankung oder mit atherosklerotischem Koronar-Plaque identifiziert.

11. Zusammensetzung zur Verwendung in einem Verfahren zur Behandlung eines Individuums mit atherosklerotischer Koronararterienerkrankung (ASCAD) oder mit atherosklerotischem Koronar-Plaque, wobei das Verfahren das Folgende umfasst:

Messen der Spiegel jedes Analyten innerhalb einer Gruppe von Analyten-Biomarkern in einer biologischen Probe, die von einem Individuum erhalten wurde, wobei die Gruppe von Analyten-Biomarkern Valylleucin, Glutamat, Acisoga, Urat, Glucuronat, Fucose, Butyrylcarnitin (C4) und Mannose umfasst; Vergleichen der gemessenen Spiegel der Analyten in der biologischen Probe mit einer oder mehreren Bezugsproben, wobei die Bezugsproben für passende menschliche Individuen repräsentativ sind; Identifizieren des Individuums als Individuum mit einer atherosklerotischen Koronararterienerkrankung oder mit atherosklerotischem Koronar-Plaque, wenn gemessene Spiegel der Analyten in der biologischen Probe bezüglich der Mengen der Analyten in den Bezugsproben erhöht oder verringert sind; und selektives Verabreichen einer Zusammensetzung, die eine wirksame Menge eines therapeutischen Mittels umfasst, das aus der aus einem Statin, Cholesterinabsorptionshemmern, Niacin-Derivaten, Omega-3-Fettsäureverbindungen, Gallensäure-Sequestriermitteln, Thrombozytenaggregationshemmern, Aldosteronblockern, Angiotensin-Converting-Enzyme (ACE) Hemmern, Angiotensin-Rezeptorenblockern (ARBs), Betablockern, Diuretika, Digitalis, Hydralazin und Nitraten, Warfarin und Acetylsalicylsäure bestehenden Gruppe ausgewählt ist, an das Individuum, das als Individuum mit einer atherosklerotischen Koronararterienerkrankung oder mit atherosklerotischem Koronar-Plaque identifiziert wurde, zur Behandlung des Individuums.

12. Zusammensetzung zur Verwendung nach Anspruch 11, wobei das therapeutische Mittel ein Statin ist.

Revendications

1. Procédé pour évaluer si un sujet humain est atteint d'une coronaropathie athéroscléreuse (ASCAD) ou atteint d'une plaque athéroscléreuse coronaire, le procédé consistant :

à mesurer les niveaux de chaque analyte dans un panel de biomarqueurs d'analyte dans un échantillon biologique obtenu d'un sujet, dans lequel le panel de biomarqueurs d'analyte comprend la valylleucine, le glutamate,

l'acisoga, l'urate, le glucuronate, le fucose, la butyrylcarnitine (C4) et la mannose ;

à comparer les niveaux mesurés des analytes dans l'échantillon biologique avec un ou plusieurs échantillons de référence, dans lequel les références sont représentatives de sujets humains appariés ; et

à identifier chez le sujet une coronaropathie athéroscléreuse ou une plaque athéroscléreuse coronaire si des niveaux mesurés des analytes dans l'échantillon biologique sont augmentés ou abaissés par rapport aux quantités d'analytes dans les échantillons de référence.

2. Procédé selon la revendication 1, dans lequel les niveaux sont mesurés à l'aide d'une analyse par spectrométrie de masse (MS), facultativement dans lequel l'analyse par spectrométrie de masse comprend une chromatographie en phase liquide couplée à une spectrométrie de masse tandem (UPLC-MS/MS) ou une chromatographie en phase gazeuse couplée à une spectrométrie de masse (GC-MS).

3. Procédé selon l'une quelconque des revendications 1 à 2, dans lequel la coronaropathie (CAD) comprend l'athérosclérose, facultativement dans lequel le sujet présente des symptômes d'une coronaropathie athéroscléreuse comprenant la douleur thoracique, l'angine, un équivalent angineux, la dyspnée ou la dyspnée lors d'un exercice, et facultativement dans lequel le sujet ne présente pas d'historique de coronaropathie.

4. Procédé selon l'une quelconque des revendications 1 à 3, dans lequel l'échantillon biologique est un échantillon de sang, un échantillon de sérum ou un échantillon de plasma.

5. Procédé selon l'une quelconque des revendications 1 à 4, dans lequel le sujet présente des facteurs de risque associés à la coronaropathie sélectionnés dans le groupe constitué par le sexe mâle, l'hypertension, la dyslipidémie, les diabètes et les antécédents familiaux de coronaropathie.

6. Procédé selon l'une quelconque des revendications 1 à 5, dans lequel l'échantillon de référence est obtenu à partir d'au moins un individu qui ne souffre pas d'une maladie cardiovasculaire ou dans lequel l'échantillon de référence comprend des niveaux d'analyte de référence prédéterminés statistiquement significatifs.

7. Procédé selon l'une quelconque des revendications 1 à 6, consistant en outre à modifier le dossier clinique du sujet pour identifier chez le sujet une coronaropathie ou une plaque athéroscléreuse, facultativement dans lequel le dossier clinique est stocké dans un support lisible par ordinateur.

8. Procédé selon la revendication 1, dans lequel l'étape d'identification est réalisée à l'aide d'un algorithme d'amplification de gradient, d'un modèle d'arbre de décision ou d'analyses de régression linéaire.

9. Kit pour évaluer si un sujet humain est atteint d'une coronaropathie athéroscléreuse (ASCAD) ou atteint d'une plaque athéroscléreuse coronaire, le kit comprenant :

des réactifs appropriés pour déterminer des niveaux d'une pluralité d'analytes dans un échantillon pour essai, dans lequel la pluralité d'analytes comprennent un panel de biomarqueurs comprenant la valylleucine, le glutamate, l'acisoga, l'urate, le glucuronate, le fucose, la butyrylcarnitine (C4) et la mannose ; et des instructions pour l'utilisation du kit dans le procédé selon l'une quelconque des revendications 1 à 8.

10. Kit selon la revendication 9, comprenant en outre :

un ou plusieurs échantillons témoins comprenant des niveaux prédéterminés des mêmes analytes, dans lequel une comparaison des niveaux des analytes dans un échantillon pour essai avec des niveaux dans les échantillons témoins identifie un sujet atteint d'une coronaropathie athéroscléreuse ou atteint d'une plaque athéroscléreuse coronaire.

11. Composition destinée à être utilisée dans un procédé de traitement d'un sujet atteint d'une coronaropathie athéroscléreuse (ASCAD) ou atteint d'une plaque athéroscléreuse coronaire, le procédé consistant :

à mesurer les niveaux de chaque analyte dans un panel de biomarqueurs d'analyte dans un échantillon biologique obtenu d'un sujet, dans laquelle le panel de biomarqueurs d'analyte comprend la valylleucine, le glutamate, l'acisoga, l'urate, le glucuronate, le fucose, la butyrylcarnitine (C4) et la mannose ; à comparer les niveaux mesurés des analytes dans l'échantillon biologique avec un ou plusieurs échantillons de référence, dans laquelle les références sont représentatives de sujets humains appariés ; et à identifier chez le sujet une coronaropathie athéroscléreuse ou une plaque athéroscléreuse coronaire si des

niveaux mesurés des analytes dans l'échantillon biologique sont augmentés ou abaissés par rapport aux quantités d'analytes dans les échantillons de référence ; et

à administrer de manière sélective une composition comprenant une quantité efficace d'un agent thérapeutique sélectionné dans le groupe constitué par une statine, des inhibiteurs de l'absorption du cholestérol, des dérivés de niacine, des composés d'acide gras oméga 3, des séquestrants d'acides biliaires, des antiagrégants plaquettaires, des bloquants d'aldostérone, des inhibiteurs de l'enzyme de conversion de l'angiotensine (ACE), des bloquants des récepteurs de l'angiotensine (ARB), des bêta-bloquants, des diurétiques, une digitale, l'hydralazine et des nitrates, la warfarine et l'aspirine au sujet identifié comme étant atteint d'une coronaropathie athéroscléreuse (ASCAD) ou atteint d'une plaque athéroscléreuse coronaire pour traiter le sujet.

12. Composition destinée à être utilisée selon la revendication 11, dans laquelle l'agent thérapeutique est une statine.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10










FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16





FIG. 17

**FIG. 18**




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34

Native CAD PV2
PD1-PD2 Joint Logistic




FIG. 35

Native CAD / PV2
 $p < 0.05$ in PD1 and PD2




FIG. 36




FIG. 37




FIG. 38

All CAD PV2
PD1PD2 Joint Logistic




FIG. 39

All CAD / PV2
 $p < 0.05$ in PD1 and PD2




FIG. 40




FIG. 41




FIG. 42

50% Stenosis (Exc. Revasc) PV2
PD1-PD2 Joint Logistics




FIG. 43

50% Stenosis (No Revasc) PV2
 $p < 0.05$ in PD1 and PD2




FIG. 44

50% Stenosis (Exc. Revasc)
PV2 PD2 Logistic




FIG. 45

Diamond Forrester Score in 50% Stenosis or Revasc.
N=348




FIG. 46

50% Stenosis or Revasc PV2
PD1PD2 Joint Logistic




FIG. 47

50% Stenosis (Inc.Revasc) / PV2
 $p < 0.05$ in PD1 and PD2




FIG. 48




FIG. 49

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2011063470 A1 [0007]
- WO 2014135696 A1 [0007]

Non-patent literature cited in the description

- **LABORDE et al.** *Metabolomics*, 2014, vol. 10, 414-424 [0007]
- **THOMAS GS et al.** A Blood-Based Gene Expression Test for Obstructive Coronary Artery Disease Tested in Symptomatic Nondiabetic Patients Referred for Myocardial Perfusion Imaging The COMPASS Study. *Circulation: Cardiovascular Genetics*, 2013, vol. 6 (2), 154-162 [0036]
- **PATEL M et al.** Low Diagnostic Yield of Elective Coronary Angiography. *N Engl J Med.*, 11 March 2010, vol. 362 (10), 886-95 [0039]
- **CHENG S et al.** Metabolite Profiling Identifies Pathways Associated With Metabolic Risk in Humans / Clinical Perspective. *Circulation*, 2012, vol. 125 (18), 2222-2231 [0040]
- **BODI V ; SANCHIS J ; MORALES JM ; MARRACHELLI V ; NUNEZ J ; FORTEZA MJ et al.** Metabolomic Profile of Human Myocardial Ischemia by Nuclear Magnetic Resonance Spectroscopy of Peripheral Blood Serum: A Translational Study Based on Transient Coronary Occlusion Models. *J Am Coll Cardiol*, 2012, vol. 59 (18), 1629-1641 [0040]
- **EVANS AM.** Integrated, nontargeted ultrahigh performance liquid chromatography/electrospray ionization tandem mass spectrometry platform for the identification and relative quantification of the small-molecule complement of biological systems. *Anal Chem*, 2009, (16), 6656-6667 [0108]
- **DEHAVEN CD.** Organization of GC/MS and LC/MS metabolomics data into chemical libraries. *J Cheminform*, 2010, vol. 2 (1), 9 [0111]
- **FRIEDMAN JH.** Greedy function approximation: a gradient boosting machine. *Ann Statist*, 2001, vol. 29 (5), 1189-1232 [0143]
- **FRIEDMAN JH.** Stochastic gradient boosting. *Comput Stat Data Anal*, 2002, vol. 38 (4), 367-378 [0143]
- **DIAMOND GA ; FORRESTER JS.** Analysis of probability as an aid in the clinical diagnosis of coronary-artery disease. *N Engl J Med.*, 14 June 1979, vol. 300 (24), 1350-8 [0151]
- **MORISE AP ; JALISIS F.** Evaluation of pretest and exercise test scores to assess all-cause mortality in unselected patients presenting for exercise testing with symptoms of suspected coronary artery disease. *J Am Coll Cardiol.*, 03 September 2003, vol. 42 (5), 842-50 [0151]

专利名称(译)	基于血液的生物标志物用于诊断动脉粥样硬化性冠状动脉疾病		
公开(公告)号	EP3242588A4	公开(公告)日	2018-10-17
申请号	EP2016735507	申请日	2016-01-08
[标]发明人	VOROS SZILARD BROWN BRADLEY O MARVASTY IDEAN B		
发明人	VOROS, SZILARD BROWN, BRADLEY O. MARVASTY, IDEAN B.		
IPC分类号	A61B5/00 G01N33/48 G01N33/53		
CPC分类号	A61K31/00 G01N33/492 G01N33/6812 G01N33/6893 G01N2800/324 G01N2560/00 G01N2570/00 G01N2800/60 A61K31/37 A61K31/616		
代理机构(译)	康罗伊, JOHN		
优先权	62/101445 2015-01-09 US		
其他公开文献	EP3242588A1 EP3242588B1		
外部链接	Espacenet		

摘要(译)

在一些方面，本发明涉及用于评估患有动脉粥样硬化性冠状动脉疾病（ASCAD）或具有冠状动脉粥样硬化斑块的人类受试者的方法。在一些方面，本发明涉及用于诊断，分类，分析和治疗动脉粥样硬化CAD和/或冠状动脉粥样硬化斑块的方法和试剂盒。