





US 20080050709A1


(19) **United States**(12) **Patent Application Publication**
Aruffo et al.(10) **Pub. No.: US 2008/0050709 A1**(43) **Pub. Date: Feb. 28, 2008**(54) **INTEGRATED BLOOD SUGAR CONTROL,
BLOOD PRESSURE CONTROL AND HEART
FAILURE SELF-CARE SYSTEM AND
METHOD**(75) Inventors: **Sylvia Aruffo**, Highland Park, IL (US);
Shirley Grey, Northbrook, IL (US);
Carol Outland, Carol Stream, IL (US);
Judith Franks Farah, Northbrook, IL
(US)Correspondence Address:
BELL, BOYD & LLOYD, LLP
P.O. Box 1135
CHICAGO, IL 60690 (US)**Related U.S. Application Data**(60) Provisional application No. 60/806,201, filed on Jun.
29, 2006. Provisional application No. 60/806,200,
filed on Jun. 29, 2006. Provisional application No.
60/806,203, filed on Jun. 29, 2006.**Publication Classification**(51) **Int. Cl.**
G09B 23/28 (2006.01)
A47B 81/00 (2006.01)
A61B 5/00 (2006.01)
(52) **U.S. Cl.** **434/262; 312/234; 600/485**(57) **ABSTRACT**

Methods and apparatus for blood sugar control, blood pressure control and heart failure care are disclosed. The methods and apparatus enable a user to provide blood sugar control, blood pressure control, and heart failure care to himself/herself. For example, after being treated by a doctor for a heart problem, a person may use the disclosed system to prevent additional heart related problems.

(73) Assignee: **CAREGUIDE SYSTEMS, INC.**, Buf-
falo Grove, IL (US)(21) Appl. No.: **11/771,793**(22) Filed: **Jun. 29, 2007**




Handle on Health - Heart Failure, Blood Pressure, & Sugar Map




Pill Organizer
Organizador de
medicinas




Magnetic Board
Affiche
de noticias



Record Book
Libro de registro



Educational Materials
Libros de instrucciones



Blood Pressure Monitor
Monitor de
tensión arterial

Scale
Balanza

Fig. 1B



Step One ²¹⁰


Know the process

The process to control your health is just like controlling *production at work* or *grades at school* or *success at sports*:

Test ²¹² First, you do a TEST to see where you are today.

Plan ²¹⁴ Then you make a PLAN to practice and get a better score the next time.


Test ²¹⁶ You TEST again and see if the plan is working.



If the score is better, keep at it. If it isn't, change the plan.

Now you're going to TEST - PLAN - TEST for your health.


²²⁰ Three monitors for TESTS



Three areas for PLANS




²³⁰ To make a plan, you need goals. Find the **Short Story** in the box. The **Short Story** explains what's happening in your body, so Goals and Plans will make sense.



When you finish the **Short Story**, come back to this paper and open it up for Step Two.

Fig. 2A

Fig. 2B




Step Two
Ready the Tools 240

1

Find the Record Book 242

This is the *one place* where you write:
Daily Tracking and Doctor Visits.



Write on Pages 2 and 4 all your **Alert** and **Alarm** numbers.

You will need: **2** Weight **Alarm** numbers,
2 Blood Pressure **Alarm** numbers, and
2 Blood Sugar **Alert** numbers.
2 Blood Sugar **Alarm** numbers.

You need these numbers when you do your tests.

If you don't have all these numbers, call the clinic and ask.

2

Get out the Magnet Boards 244

Copy the **ALARM** numbers and how often to test
your sugar when you're sick.

Stick the Magnet Boards on the refrigerator.

3

Find the Wallet Card 246

Fill in the spaces. Put the Card
in your purse or wallet.



You will Need these numbers to know
when you have an Emergency and what
to do.



Step Three

Take the Tests

1 Weight Test

- In the morning, before breakfast, use the toilet.
- Take off any clothes.
- Put the scale on a hard floor, not carpet.
- Step on the scale.
- Write your first *Weight* score on Page 20 in the Record Book.

2 Blood Pressure Test

- If you don't have a Blood Pressure Monitor, most drug stores have a free station to check it.
- Write your first *Pressure* score on Page 20 in the Record Book.

How I Tested		Start
		SUNDAY
Number of pillows last night		
WEIGHT	167	
BLOOD	128/84	
PULSE	72	
BLOOD SUGAR		
Finger		
Stick spot		

3 Blood Sugar Test

Find a large bottle with a screw cap. Write on it, **SHARPS do not recycle**. Use it to throw away used lancets and needles so no one gets stuck when they handle the trash.

- Choose a clean place near a sink.
Bring the Record Book, pen, clean towel, soap, a lancet and a test strip.
Get the Monitor and directions that came with it.



If you can't find the monitor directions, look on the device for a phone number. Call and they will send you new directions.

- Wash your hands with warm water and soap. Dry them on the clean towel. **Don't use alcohol.** It makes skin tough and harder to stick.
- Stick the **side** of your finger. Follow the Monitor directions to see the blood sugar score on the screen. The score will stay on the screen while you open the Record Book.
- Turn to Page 18 in the Record Book. Follow the steps there to record your first test.

How I Tested	
HGT	5'10"
WT	167
BLOOD	128/84
PULSE	72
BLOOD SUGAR	100
Finger	
Stick spot	

Fig. 2C

Fig. 2D



Step Four

Make a Plan

280

The Plan lets you reach the **Six Goals** in the Short Story:

- | | | |
|-----|---|--|
| 282 | 1 | Pump up the heart. <i>Exercise.</i> |
| 284 | 2 | Lighten the load. <i>Eat less salt. Take water pills.</i> |
| 286 | 3 | Shorten the trip. <i>Watch what you eat. Move more.</i> |
| 288 | 4 | Widen the road. <i>Stop smoking, relax. Take heart pills.</i> |
| 290 | 5 | Clear the strip. <i>Eat more fiber and less fat. Take cholesterol pills.</i> |
| | 6 | Steady as she goes! <i>Regular amounts of everything on time.</i> |

The Plan has three Parts: *watch the meals*, *make the moves*, *take the meds*



292 In the Folder you'll find a section for each of the 3 areas of The Plan. Follow the steps to make your own Plan.

When you've made The Plan, the *Record Book* keeps track: **How I Tested** and **What I Did**

Daily Tracking

You already have in the Record Book:

- The doctor's limit numbers
- Your first scores

Now day by day you put in:

- What you do to follow The Plan
- Test scores to see if The Plan is working.

Doctor Visits

Take the Record Book to every clinic visit so the doctor can see if The Plan is working.

Don't leave the clinic until you know:

- Doctor's last test scores
- Any changes to daily limit numbers
- Any changes to meals, moves or meds.

Fig. 3A

112

The Short Story on Pressure, Arteries and Sugar



1

"Pressure" is one thing pushing on another. Blood pushing on the inside walls of arteries is *blood pressure*.

3/0

Blood pushes on the arteries when the heart pushes on the blood. The push is like a pump: *push and rest, push and rest.*

3/2


Pressure is higher on *push* and lower on *rest*. That's why blood pressure has two numbers: a higher one for *push* and a lower one for *rest*.

For example:

Healthy blood pressure could be 120 for when the heart pushes, and 80 for when the heart rests.

3/4

These are good numbers: enough pressure to move blood through clear, open arteries.



The heart pushes blood through the arteries to feed all parts of the body.

Blood leaves the heart with its cargo like THRU TRAFFIC on the highway.

The coronary arteries make a short loop back to feed the heart itself, like getting off the highway at the first exit.

2

Sometimes the heart pushes harder. The top number goes up, *more than 120*.

3/20

When the heart pushes harder, it doesn't rest well between pushes. So the bottom number goes up, *more than 80*.

It's normal for pressure to go up when you need extra energy, like at work. These short times of high pressure are not a problem.

High blood pressure all the time is a problem.





Fig. 3B

What Makes Blood Pressure Stay Up



Blood carries food and oxygen to all parts of your body.
But if the blood can't flow through, you have a problem.

Fig. 3C

When not enough food and oxygen get to your feet or hands,
they complain by "falling asleep."

You shake more blood down into them and they're OK.

When not enough food and oxygen get to the heart,
it complains, too—a lot louder.

You get chest pain, or even a "heart attack."

But you *can't* just shake more blood into the heart.

Here's what you do instead:



What Makes Blood Pressure Go Down





Fig. 3D


The Scoop on Blood Sugar



1

You know you have sugar in your blood. How did it get there? From what you ate.

You can guess sweets like candy and cake put sugar in your blood. But a lot of other food has *carbs* and *carbs turn into sugar* after you eat.





2

Your body uses sugar when you move during the day, like a car uses gas. When the gas tank gets low, you feel hungry. You fill up your stomach like a gas tank. The carbs in the food change into sugar and enter your blood.



A car doesn't burn all the gas at once. The car uses gas slowly, a little at a time. A car has a system to control the gas supply.

This system lets just the right amount of gas into the engine. *If the system isn't working, you flood the engine with gas.*



3

Your body doesn't burn all the sugar at once. The body uses sugar slowly, a little at a time. A body has a system to control the sugar supply.

Insulin lets just the right amount of sugar into the blood. *If the insulin isn't working, you flood the blood with sugar.*




When your test shows **High Blood Sugar** that means your body's insulin isn't working right, or you ate and didn't move enough. So the insulin let too much of that sugar into the blood.


You've flooded your engine.

Fig. 3E

4 Now suppose you drive a long way without stopping for gas. The tank goes to empty. If your stomach is empty, you get **Low Blood Sugar**. You went too long without eating or moved more than usual. The body does have a spare tank, the liver. Most people tap into that spare tank when they run low.



But it's insulin that opens the spare tank.
If insulin doesn't work, you can't tap the spare tank.
 Even if insulin *works*, the liver won't open if it's busy.
 The liver gets busy when you drink alcohol.
So never drink alcohol on an empty stomach!



When your test shows **Low Blood Sugar** that means you didn't eat enough, moved more than usual, or insulin couldn't tap the liver for extra sugar.

You've run out of gas.

- 5** Controlling blood sugar means making everything regular:
- Eat regular size meals so you don't flood the engine.
 - Eat at regular times so you don't run out of gas.
 - Exercise the same amount every day so you use all the gas in the tank.
 - Take your pills and shots on time so the system lets just the right amount of gas into the tank.

Now Put it All Together ✓ 700

- 1** Lighten the load. *Eat less salt. Take water pills.*
- 2** Shorten the trip. *Watch what you eat. Move more.*
- 3** Widen the road. *Stop smoking, relax. Take heart pills.*
- 4** Clear the strip. *Eat more fiber and less fat. Take cholesterol pills.*
- 5** Steady as she goes! *Regular amounts of everything, on time.*




Fig. 3F

Why are Pressure, Arteries and Sugar a "Triple Threat?"




We all know people who should watch what they eat, get more exercise and take medicine, *—but they don't.* It seems like they get away with it.

But when you have all three: clogged arteries, high pressure, and runaway sugar, it's not likely you can get away with much.


There's a reason, something you have known since you were a small child—*sugar makes things sticky!*



① Start with the fact you have more cholesterol than other people do...



② ...and you often have more sugar in your blood than other people do.



③ Extra sugar makes more cholesterol stick to your arteries



④ More stuck cholesterol means even tighter space to squeeze blood through



Your blood pressure goes even higher



More cholesterol, more sugar, more pressure: a triple threat. Life is just not fair—but you *can* get control. This kit will guide you step by step, to beat the threat and get on with living your life.

It's **TEST-PLAN-TEST** from now on.


Fig. 3G

800

Tests: Find Out If Your Plan is Working

You saw on the first page of this kit, **Step One**, that you do **three tests at home**: Sugar, Pressure, and Weight. There are two more tests that the clinic does: a different Sugar test, and a Fat test.

Many drugstores have a poster that says "**A B C**". Your clinic may have it, too. The poster is about **TESTS**.




A is for **A** 1 C. That's the name of the *clinic's* sugar test.

B is for **B**lood Pressure. You take your Pressure test at home, and the clinic takes it, too.

B is also for **B**lood Sugar. The poster here means the sugar test you do at home.

C is for **C**holesterol. The poster only says *cholesterol*, but in fact the clinic test reports both cholesterol and triglycerides. It's actually a test for *fat*.



The clinic's fat test reports:

- **Total** cholesterol
- **Two kinds** of cholesterol and
- Triglycerides


The two kinds of cholesterol are **LDL** and **HDL**.

LD stands for **Low Density**. *Density* means how tightly packed something is.

Low Density is loose pack. *Loose* is the kind that sticks. Think of styrofoam packing. Loose bits stick to your hand. LDL sticks to your arteries. **Low** is the *bad* cholesterol.

High Density is hard pack. *Hard* pack doesn't stick. Those same styrofoam bits, hard packed, don't stick. HDL doesn't stick to arteries. **High** is the *good* cholesterol.

Think of something you don't like that starts with the letter **L**. Then you'll never forget which cholesterol is bad.



W is for **W**eight. The poster doesn't mention this test, but when you read **Step One**, you know to weigh in every day. The clinic will test your weight, too.

How to Read Clinic Test Results

The clinic should give you a copy of your test results.
If they don't, ask for them!



The Sugar Test

Health Care people call red blood Hemoglobin.
Sugar sticks to hemoglobin. The sugar builds up over three months.
Then it flushes out and starts over again.

The clinic sugar test tells how much sugar has built up on the hemoglobin.
If you know the Hemoglobin A1C score from the clinic, you can tell what your average daily score for the last three months was on the sugar test you do at home.

If your A1c score is	then your average daily score was
6	135
6.5	153
7	170
7.5	188
8	205
8.5	223

Here is a real test result for a man 56 years old.

The clinic sugar test is called **Hemoglobin A1C**

The man's doctor said his goal is 6.5 Is his Plan working?

Lab Results:

Patient Name:

Chemistry Profile: Glucose: 110 (goal < 99)
Electrolytes:
Liver Tests:
Kidney Tests:
Proteins: } all normal

Hemoglobin A1C (3 month diabetic blood test): 6.3

The Fat Test

Health Care people call all kinds of fat Lipids.

Here is a real result for a woman 63 years old.

1 Less than 150 is good for Triglycerides.

Is she OK?

2 Less than 200 is good for Total Cholesterol.

How is she doing?

3 For Bad Cholesterol, less than 70 is the goal.

Should she change her Plan?

4 For Good Cholesterol, most women need more than 50. Men need more than 40.


Is this woman OK?

LIPID CHEMISTRY		
	ABN/LOW	NORMAL HIGH
COLLECTED 09/11/2003 09:00		
Triglycerides	86	
Cholesterol	199	
LDL		128 H
HDL Cholesterol	54	
<40 mg/dL:	low HDL-cholesterol (major risk factor for CHD)	
>=60 mg/dL:	high HDL-cholesterol	

The Volume Test

Health Care people call the volume test Ejection Fraction.

The doctor sees the results as part of an Echocardiogram. The goal is 55% or more.
What was this person's first score?



The Volume Test: 18%, but the next year he was OK.
The Fat Test: No
Answers: The Sugar Test: No

Fig 3H

106



Fig. 4A

Fig. 4B

TEST

Ask these questions every clinic visit

Answers you need to do your tests

Ask the Doctor		VISIT 1	2	3	4	5	6	7	8
1104	When should I test my Blood Sugar?								
	Before Meals								
	1 hour After Meals								
	Bedtime								
1112	Where should my Blood Sugar number be?								
	HIGH LIMIT:								
	LOW LIMIT:								
1114	I should call you if my Blood Sugar number is:								
	ALARM								
1114	Top Alarm								
	Bottom Alarm								
When I am sick, I should test my sugar every _____ hours.									
Do I need a prescription for ketone test strips?									

PAGE 1

[illegible]

1120

1122

Reminders to the Clinic to do their tests

1124

When to do the Test	Test Name	My goal is	How it really came out LAB RESULTS							
			VISIT 1	2	3	4	5	6	7	8
Every visit	Weight									
Every visit	Blood Pressure	Less than 120/80								
Every visit	Foot Check									
At least two times a year	Hemoglobin A1c	Less than 7								
	Total Cholesterol	Less than 200								
	HDL (good)	Women: more than 50 Men: more than 40								
	LDL (bad)	Less than 100								
	Triglycerides	Less than 150								
Once a year	Microalbumin	Less than 30								

1126

ABOUT MY NEXT APPOINTMENT

What day and time?

PAGE 2

Fig 4C

1200
1202

PLAN

The last pages were about tests.
This page starts the Plan: MOVES, MEDS AND MEALS.

ABOUT FOOD VISIT 1 2 3 4 5 6 7 8

What should my meal plan be? Write the answers on the top of Page 7.

How much cholesterol in one day? Less than 300 is good.								
Is a salt substitute with potassium OK?								
How much salt in one day? Less than 1500 is good.								

ABOUT MOVES

What kind of exercise should I do?								
How long each time?								
How many times a week?								

Do I need these shots? Who should I get to give them to me?

A flu shot?	Date I got one		A pneumonia shot?	Date I got one	
-------------	----------------	----------------------	-------------------	----------------	----------------------

Should I cut my own toenails? If yes, show me how.

If not, tell me a foot doctor to see:

NAME	
ADDRESS	TELEPHONE

Show the foot doctor the foot test you do on the back page of this Record Book.

How often should I see the eye doctor?

NAME	
ADDRESS	TELEPHONE

How often should I see the dentist?

NAME	
ADDRESS	TELEPHONE

Do you have any special instructions for me?

Fig. 4D

1204
1206
1208
Pills for my Heart 1214 1212

Name of the Pill	Strength	How Many to take at One Time	When to Take It			
			Breakfast	Lunch	Dinner	Bedtime
		1210				

If the Doctor changes your pills, cross out the old; write in the new. Tell any allergies you have _____

Pills for Blood Sugar 1216

Name of the Pill	Strength	How Many to take at One Time	When to Take It			
			Breakfast	Lunch	Dinner	Bedtime

If the Doctor changes your pills, cross out the old and write in the new. Tell any allergies you have _____

Fig. 4E

Other Prescription Meds ¹²¹⁸

Name of the Med	Strength	How Many to take at One Time	When to Take It			
			Breakfast	Lunch	Dinner	Bedtime

Things I Take on my Own ¹²²² ¹²²⁴ ¹²²⁶ ¹²²⁰ ¹²²⁸

Kind of thing	NAME	HOW MUCH I TAKE	WHEN I TAKE IT
VITAMINS			
HERBS			
OTHER			

Fig. 4E (cont.)

Fig. 4E (cont.)

Ask the doctor: Where on my body should I give myself the shots?

Every Day Insulin

Start Date	What Kind of Insulin	How Much to take at One Time	When to Take It
		1236 units	
		units	
		units	
		units	
		units	
		units	
		units	
		units	
		units	
		units	
		units	
		units	

... the doctor may want you to take different insulin when your Blood Sugar test score is too high.

"It Depends" Insulin

	If your Blood Sugar test is more than	Take this kind of Insulin	How much to take at one time
Date the doctor gave me these numbers:			units
			units
			units
			units
			units
Date the doctor gave me new numbers:			units
			units
			units
			units
			units

1200

MEAL PLAN		HOW MANY SERVINGS or CARBS Which do you use?								
		VISIT	1	2	3	4	5	6	7	8
Breakfast	Protein									
	Fat									
	Fruit									
	Vegetables									
	Grain & Starch									
	Milk									
	Total									
Lunch	Protein									
	Fat									
	Fruit									
	Vegetables									
	Grain & Starch									
	Milk									
	Total									
Dinner	Protein									
	Fat									
	Fruit									
	Vegetables									
	Grain & Starch									
	Milk									
	Total									
Snack										

Fig. 4F


Fig. 4G

1300

Write the First BLOOD SUGAR Test Score

Turn back to Page 1. Find the LOW and HIGH numbers your doctor told you.

Write the numbers in the boxes on the Chart on this page.



2 What was your test score?

1311 If the test was more than the **HIGH** limit, write the time and the score in the gray space on top.

TOO HIGH

1314a

1314b

1316a

If the test was in between, write the time and the score in the white space in the middle.

1312b

1312a

1316b

1316c

TOO LOW

If the test was less than the **LOW** limit, write the time and score in the gray space on the bottom.

PAGE 7

1302

Take ONE WEEK to see the difference food makes!

On the next 2 pages, you'll see a chart for the whole week. Start tomorrow to track your MEDS and your MEALS.

During the week, read Step Four, "Make a Plan."

When you have a whole week of tests, you'll see at a glance how you're doing most of the time:

IN CONTROL
or ON ALERT.

When you see how what you eat changes your Blood Sugar, you'll want to make a plan.

1 Write the date and the HIGH and LOW numbers. Remember the ALARM number.

Weekly Record Start Date: 9/

TUESDAY WEDNESDAY

2 Write the time you take each Blood Sugar med.

MEALS

Time 1:00

Time 2:00

Time 3:00

3 Write everything you eat and drink.

3 eggs

2 bananas

orange juice

apple

bread

chicken

meat

black

4 Test your Blood Sugar at least once a day.

Write the time and the score: **IN CONTROL** or **ON ALERT.**

How I Tested

TOO HIGH

TOO LOW

Time 1:00

Time 2:00

Time 3:00

If you test at or beyond your ALARM, follow the emergency instructions on the Magnet Board.

1304

What I Ate Weekly Record
Start Date:

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
MEDS							
BREAKFAST							
LUNCH							

PAGE 8

1306

1308

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
DINNER							
SNACK							

How I Tested

TOO HIGH							
TOO LOW							

1310

1314

1312

Fig. 4B (cont.)

Fig. 4H

Now the week is over. Look at the days when your Blood Sugar was IN CONTROL. What did you eat on those days? When it went high or low, what did you eat? You should see a pattern.

How to Keep the Record Book


From now on, you don't need to write everything, but just enough to track:



14110

 Every week, prepare for the TESTS:

fill in the date, and the numbers for the
HIGH and **LOW** blood sugar limits.



PAGE 9


14120

 Every Day Record the Tests

Weight - weigh yourself and write your score on the next page.

BLOOD PRESSURE - Take your pressure and record your reading.

Blood Sugar - Test. Write the times and the scores.



 Every Day Track the Plan

MEALS

14140


Level I: Put a check if you followed the Poster. Write what you ate from the red side so you can think of ways to resist.

14142

Level: II Put a check if you controlled the serving size. Write what foods went over the limit so you can think of how to control them.

14144

Level III: Put a check if you followed your meal plan. If you had anything not on the plan, write what and how much. Also write if you skipped a meal.



MOVES: Write what you did to get moving and for how long.
1450



MEDS: If you haven't done it yet, read the blue pages starting with "The Right Meds at the Right Time."
1460 You've written the names of your meds and the time you take them on Pages 4, 5, and 6.

If you take anything else, or take a med at a different time, write what and at what time on the chart. Write if you skipped a med. Write where you stuck yourself to test your blood sugar.

If you take shots, write the shot spot on the Chart.

Turn the page to start.

When this Book is full, use RECORD BOOK 2.



PAGE 10

Fig. 4 H (cont.)

How to Test your Feet

*It's best to have someone do this test for you.
If you do it alone, you may need a mirror.*

1 Get a pencil. Use the Red Filament below.
Hold the Filament by the paper handle.


2 Look at the circles on the pictures of feet below.
You'll touch each of those places on your feet.

*If you have a sore, callous or scar in a circle,
touch the skin beside it, not the spot itself.*

3 Touch the Filament to the first circle place.
Push hard enough to make the Filament bend.
Hold it there while you count to two.

4 Can you feel the Filament?
If you can't, put an X in the circle.
Repeat for each circle on both feet.

IF YOU HAVE ANY Xs, CALL THE DOCTOR



Check off each month after you do the test:

January ☐	April ☐	July ☐	October ☐
February ☐	May ☐	August ☐	November ☐
March ☐	June ☐	September ☐	December ☐

The LEAP Foot Screening Filament in this Kit is endorsed by the American Diabetes Association, the American Pharmaceutical Association and the American Podiatric Medical Association.

Fig. 4J




Fig. 4K

Fig. 5

a balancing act Day By Day



Every Day

Shopping List

Once A Week

Fill Pill Organizer

Check meds and other supplies

Make a Meal Plan for the week

Make a list and go shopping

Once A Month

Do the LEAP test on feet

Cut toenails (if doctor said it's OK)

Pharmacy Number: _____

Emergency Action Plan

If you feel sick or strange,
dizzy or jittery,
test your blood sugar.

If it's Too High

If it's Too Low

If the Alarm number is more than 300
Call for help

If the Alarm number is less than 70
Call for help

If it's not that high yet: Flush the sugar out:
Every hour, take a big drink with
no sugar and no caffeine

If it's not that low yet: Get some sugar in:
• Chew 3 glucose tabs (in the Kit!) or
• Drink 1/2 cup of fruit juice or
• Take a spoonful of sugar, honey or syrup or
• Suck 8 small or 2 large hard candies
not chocolate--it's too slow

If the doctor said "test for ketones," do it now.
If the test shows ketones, call the Doctor

Test again every _____ hours

Sick Days

If you have a fever, sneezing and coughing,
you've caught a bug!
If you have vomiting or diarrhea, call the
Doctor.
Always let someone know you're sick so they
can check on you.

• Keep taking the blood sugar meds.
You need them more than ever.
• Every hour, take a big drink with no sugar
and no caffeine
• Test your blood sugar every _____ hours.

Blood Sugar Doctor: _____

Heart Doctor: _____

Call the Heart Doctor if:

- I've gained _____ pounds in a day.
- I've gained _____ pounds in a week.
- My blood pressure is above _____
- My blood pressure is below _____
- I feel tired and weak and my blood sugar is OK.

Call 911 if:

- ▶ A heavy weight is crushing my chest
- ▶ Pain spreads to my shoulders, across my back, neck, arm or jaw
- ▶ Chest pain keeps coming back
- ▶ It's really hard to breathe
- ▶ I'm sweaty, cold and clammy
- ▶ I have indigestion or vomiting

Things to remember to ask the doctor or nurse:

Next Appointment: _____

2150 ⚡ **ON SICK DAYS**

- 1 Tell someone who can check on you
- 2 Keep taking your meds!
- 3 Take water, diet drinks every hour
- 4 Test your blood sugar every 4 hours

CALL YOUR DOCTOR IF

- The blood sugar test is over _____
- You can't stop vomiting or diarrhea

WALLET CARD

2160 ⚡ **EMERGENCY MEDICAL INFORMATION**

NAME _____

Day Phone _____

Blood Sugar Doctor _____

EMERG PHONE _____

Heart Doctor _____

EMERG PHONE _____

Fig. 6A




Fig. 6B

Fig. 7A

124
124

How to "Watch What You Eat"

People often say "I watch what I eat" —when they really don't.

Level One: The Poster 3100

Put up the Poster in the kitchen. Eat more from the green side and eat less from the red side.




Eating Less Salt 3120

Less salt in cooking

New flavors in the pot 3122

Check out other spices and sauces. Try lemon and other juices.



Less salt from a shaker

New flavors on the table 3124

There may be new things that work for the whole family.




Less in the cupboards & fridge

Salty food off the shelf 3126

- Bouillon
- Catsup
- Cheese
- Chili Sauce
- Cold Cuts
- Frozen dinners

- Mustard
- Olives
- Pickles
- Salad Dressing
- Sausages
- Soy Sauce



Eat More "Soaker" Carbs 3130

Some carbs pull cholesterol out of your arteries! *Call them Soakers.*

Soaker Grains 3132

- Barley
- Corn Grits
- Dark Rye Bread
- Oat Bran, Rolled Oats
- Nuts and Seeds
- Rice Bran, Brown Rice
- Popcorn

Soaker Fruits 3134

- Apples
- Apricots
- Bananas
- Blackberries
- Blueberries

- Oranges
- Pears
- Plums
- Strawberries
- Tangerines

Forget Grapefruit. It stops cholesterol meds from working!

Soaker Vegetables 3136

- Asparagus
- Broccoli
- Brussels Sprouts
- Cabbage
- Carrots
- Lentils

- Peas
- Pinto Beans
- Potatoes
- String Beans
- Summer Squash
- Sweet Potatoes

- Winter Squash
- Zucchini

Some soakers are "light" and some are "heavy." Check the Poster!



Did the doctor say use fake salt with **potassium**? Look for it with spices at the store. Use it at the table. You may never even notice it's not salt.




Fig. 7B

Level Two: The Plate

3200

3202

Restaurant meals can be "super-sized" or "all you can eat." Helpings that big may seem normal.

But when a health care person says "one serving," it means a very specific size, not just any helping you put on your plate.

Helping = a scoop of food, any size.

Serving = a scoop of food a certain size.

"Portion" means the same as **Serving**.

"Exchange" means the same as **Serving**

1 Find the Portion Plate. On the Plate are playing pieces from games and sports. These are always the same, standard size.

2 Put the Portion Plate on top of your real plate. Get used to the sizes. Picture these sizes whenever you put food on a plate.

3 To find the serving size for other foods, use the Loose Food Guide.





Fig. 7C



How Many Servings a Day?

A **serving** is just enough for the right balance of nutrition in a food. A serving is Nature's Pill: a "dose" of nutrition.

Protein  Most people need 3-4 servings of meat, fish or eggs a day. <i>That's one each meal.</i>	Vegetable Fat  Most people need 3-6 servings a day of oil or margarine. <i>That's one or two each meal.</i>	Vegetables  Most people need 2-3 servings of vegetables a day. <i>That's one each meal.</i>
--	---	---

~3270

Some people need just one serving of a food and others need more, just like some people need one pill and others need more. *The doctor decides. The clinic will tell you.*

Milk & Yogurt  Most people need 2-3 servings of milk or yogurt a day. <i>That's one each meal.</i>	Grains & Starch  Most people need 5-6 servings of grain and starch a day. <i>That's two each meal.</i>	Fruits  Most people need 2-3 servings of fruit a day. <i>That's one each meal.</i>
--	--	--

3272

3274

3276

3278

3280

3282

Fig. 7E

Counting Tools 3310

How much salt, cholesterol or carbs are in what you want to eat? *With just three tools, you can spot how much is in any food.*

3320

1 If the food comes in a package, check the

Nutrition Facts Label


Find the words **Cholesterol**, **Sodium** and **Total Carbohydrate** or **Total Carb**.



The number right after those words is how much is in ONE serving.

Nutrition Facts	
Serving Size 1 cup [240 ml]	
Servings per container 8	
Amount Per Serving	
Calories 100	Calories from Fat 25
% Daily Value*	
Total Fat 2.5g	4%
Saturated Fat 1.5g	8%
Cholesterol 10mg	3%
Sodium 125mg	5%
Total Carbohydrate 12g	4%

Count Total Carb



NOT Net Carb

Number here	Words here
Nutrition Facts Serving size: 1 bar (32 g), Amount Per Serving: Calories 260, Fat Cal 130, Total Fat 14 g (22% DV), Sat. Fat 2 g (10% DV), Cholesterol 0 mg (0% DV), Sodium 140 mg (6% DV), Total Carb. 28 g (9% DV), Fiber 2 g (5% DV), Sugars 22 g, Protein 6 g, Vitamin A 0%, Vitamin C 0%, Calcium 2%, Iron 2%. Percent Daily Values (DV) are based on a 2,000 calorie diet.	

3330

2 If the food doesn't have a package, look it up in the

Loose Food Guide

When you buy fresh fruit and vegetables or go out to a restaurant, it doesn't come in a package.

Use the *Loose Food Guide* to find how much salt, cholesterol and how many carbs are in a serving.
15 carbs = 1 serving.



3340

3 If you're going out to grab a bite, take the

Fast Food Guide

The book is a little tricky: First, see how many *servings* are really in your order. Skip the first number, the one the book calls a "serving." Go to the last number. The book calls a real serving an exchange.

In the photo, 4c means 4 servings of carbs.
Run your finger back to see how many carbs that is.
Sodium and cholesterol are on the line, too.

Servings here	Carbs here	Cholesterol here	Sodium here
1	15	30	100
2	30	60	200
3	45	90	300
4	60	120	400
5	75	150	500
6	90	180	600
7	105	210	700
8	120	240	800
9	135	270	900
10	150	300	1000

Fig. 7F


Read this page to get the basic idea of counting. Turn the page for practice counting. Then you'll make a meal plan with all the right numbers.

3350

Count Salt

For most people, the salt limit is 1500 mg a day. That's 500mg a meal. Usually 5 servings make a meal. So 100mg each serving is usually the limit.

Check the counting tools on foods you like. One brand may have too much, another may be just right.



3360

Count Cholesterol

For most people, the cholesterol limit is 300mg. That's 100mg a meal. If you have 2 servings a meal, 50mg a serving is the limit.

An egg has 225mg. So if you eat an egg for breakfast, you've only left yourself 1 serving of meat or fat for the rest of the day!

You may decide it's easier to eat just egg whites—no cholesterol.



Nutrition Facts		Amount Per Serving	% Daily Value
Serving Size 1/2 cup (125ml) condensed soup		50g	4%
Servings about 2.5			
Calories 60			
Total Fat	1.5g	2%	
Sat. Fat	0.5g	1%	
Cholest.	15mg	3%	

OK!



Too much!


3370

Count Carbs

For most people, the carb limit is 150g a day. That's 50g a meal. If you have 3 servings a meal, 16g of carbs a serving is the limit.

Remember the "free" vegetables! You don't have to count their carbs

That means you can always get more servings without going over the carb limit.



Now Practice Counting all Three: Salt, Cholesterol and Carbs $\checkmark$ 3390

Here's Someone's Dinner. How much salt, cholesterol and carbs are in this meal?



	Pinto Beans	Formal Slice	Fried Chicken	White Rice	Cola Drink
salt					
cholesterol					
carbs					

Find the numbers for salt, cholesterol and carbs.
If the food has a package, use the Nutrition Facts label.
If the food doesn't have a package, use the Loose Food Guide or the Fast Food Guide.


How does this meal look...

- if the limit for one meal is 500 mg of salt?
- if the limit for cholesterol is 100mg?
- if the limit for carbs is 50g?

The "Real Meal" Deal

Before you turn the page and start planning your own meals, find the Deck of Cards in the Kit. Get a friend or family member to play it with you—or try a game of solitaire.

*Play the game first.
Then making meal plans
will be much easier!*




Primo Beans	(Tomato Slice)	(Fried Chicken)	White Rice	Cold Drink	
260	0	375	9	50	664
0	0	77	0	0	77
19	0	7	45	39	110

Add 'em up:

ANSWERS

Fig. 7G

Practice Counting Carbs



How many carbs?

How many carbs? _____

How many carbs? _____

3384

0
0
0
0
0
0
0
0
0

How many carbs?

carbs? _____ How

low many carbs? _____

2

How many carbs? _____

low many carbs? _____

How many carbs?

How many carbs? _____

How many carbs? _____

—

H
?

How many carbs? _____

ANSWERS Top Row
Brown Bread: 1 slice = 12 carbs
Tortillas: 3 tortillas = 32 carbs
Sugar-Free Jell-O: 0 carbs

Note!
One tortilla is like 1 slice bread
Sugar-free gelatin has no carbs!

ANSWERS Middle Row
 1 Orange = 15 carbs
 1 ear yellow corn = 14 carbs
 1 medium potato = 15 carbs
 1 bowl of lettuce = Not enough
 to count. *Lettuce is a free food*

ANSWERS Bottom Row
1 slice pizza = 32 carbs
1 medium DQ = 63 carbs
1 Big Mac = 47 carbs

8-8333

Fig. 7H

Plan Meals for a Week at a Time

3600

Enough practice!! *Time for a real plan.*

- 1 Check Record Book on Page 14.
Find your own limits for salt, cholesterol and servings or carbs.

The "rules" for planning real meals are just like the card game, *except now you use real numbers from the clinic.*

- 2 Find the Meal Planner in the kit. Copy the limit numbers from the Record Book into the spaces on the Meal Planner.

- 3 Take the Loose Food Guide to the kitchen.
Use the Guide and the packages of food there to plan meals with the right numbers.

If you don't have packages of foods you want to eat at home, take the Meal Planner to the store. Lay the Meal Planner in the grocery cart. Write a plan as you check package labels.

ABOUT FOOD		Visit	1	2	3	4	5	6	7	8
What should my meal plan be?		Write the answers below								
How much salt in one day?										
Less than 1500mg is good										
Do I need to eat foods with a high or low potassium content?										
Is a salt substitute in my potassium diet?										
How much cholesterol in one day?										
Less than 300mg is good										


MEAL PLANNER		HOW MANY SERVINGS OF CARBS (Which do you like?)							
Breakfast	Visit	1	2	3	4	5	6	7	8
Protein									
Fat									
Fruit									
Vegetables									
Grain & Starch									
NAH									
Total									

PAGE 14

LUNCH		HOW MANY SERVINGS OF CARBS (Which do you like?)							
Protein	Visit	1	2	3	4	5	6	7	8
Fat									
Fruit									
Vegetables									
Grain & Starch									
NAH									
Total									

DINNER		HOW MANY SERVINGS OF CARBS (Which do you like?)							
Protein	Visit	1	2	3	4	5	6	7	8
Fat									
Fruit									
Vegetables									
Grain & Starch									
NAH									
Total									

PAGE 15



How to Enjoy a Night Out

3790

The goal is to wake up feeling good about yourself the next day.

- 1 Before you order, ask yourself how the choice will make you feel in the morning.
- 2 Wait staff are ready to make changes —everybody does it today! So ask for changes, like dressing on the side or lemon slices instead of dressing.
- 3 Ask for a take home bag when you place your order. Cut portions in half and put half away before you eat the meal.
- 4 Watch out for drinks:
 - Sugary soft drinks send blood sugar too high.
 - Alcohol—believe it or not!—sends blood sugar too low.

Fool them all: order sparkling water with a lime. If you take alcohol, sip just 1 glass a long time.



126

Fig. 8

How to "Watch What You Eat"



3114 Watch Out for Salt **Eat less** **Eat more** 3112

Salty Snacks



Food in Cans



Salt Shakers



Snacks with no Salt



Cans with no Salt or Frozen Vegetables



Spices with no salt




3116 Watch Out for Cholesterol

Red Meat, Egg yolks



Fat that comes from Animals



Fish, Chicken, Pork, Egg whites



Fat that comes from Plants



3118 Watch Out for Carbs

Sugar



Sugar-Free




"Whites"



"Browns"



Heavy Vegetables
They're healthy, but have a lot of carbs. A small helping is good.




Light Vegetables



Heavy Fruits


All dried fruit

Try to eat just half



Light Fruits








Fig. 10A

Fig. 10B

3340

If it's not in a package and not fast food,
look here.

Calculated from U.S. Dept of Agriculture Tables of Food Values

Grain & Starches

3341

	Serving, Portion or Exchange	Calories	Sodium Content(mg)	Cholesterol (mg)	Number of Carbs
Bagels, plain	1/4 of a 4" bagel	80	130	0	17
Biscuits, plain or buttermilk, made from recipe	1-1/4" biscuit	106	174	2	13
Bread, combread, made from recipe, with lowfat (2%) milk	1/2 piece	86	214	26	14
Bread, Indian (Navajo) fry	5" piece	297	313	6	21
Bread, pita, white	4" pita	77	150	0	16
Cornmeal, whole-grain, yellow	1/8 cup	55	6	0	12
Corn, sweet, white, cooked, boiled, drained no salt	1 ear	83	13	0	19
Corn, sweet, yellow, frozen, kernels on cob, cooked, boiled, drained, no salt	1 ear	59	3	0	14
Couscous, cooked	1/2 cup	88	4	0	18

Potato, baked, with skin, no salt	1/2 potato	57	3	0	21
Potato, baked, no skin, no salt	1/2 cup	74	4	0	16
Potatoes, boiled, cooked in skin, no salt	1/2 potato	59	2	0	14
Potatoes, peeled then boiled, no salt	1/2 potato	58	3	0	13
Rice, brown, long-grain, cooked	1/3 cup	72	1	0	15
Rice, white, long-grain, cooked	1/3 cup	68	3	0	15
Sweet potato, cooked, baked, in skin, no salt	1/2 potato	65	8	0	15
Sweet potato, cooked, boiled, no skin, no salt	1/2 potato	60	10	0	13
Wheat flour, white, all-purpose, enriched, bleached	1/4 cup	114	1	0	21

Proteins

Not enough carbs to count, but watch calories, sodium and cholesterol


3342

Cheese, blue	1 oz	100	395	21	1
Cheese, camembert	1 1/4 oz	114	320	27	0
Cheese, cheddar	1 oz	114	176	30	0
Cheese, cottage, creamed, large or small curd	1 cup	216	851	32	6
Cheese, cream	1 tbsp	51	43	16	0
Cheese, feta	1 oz	75	316	25	1
Cheese, low fat, cheddar or Colby	1 oz	49	174	6	1

130 ↘

*Make educated choices within
your meal plan...*

Tips for Healthier Eating



Indispensable to
human health

**BD Getting Started™
Fast Food Guide**

Fig 11A

www.krispykreme.com
Brands are trademarks of their respective holders.

Krispy Kreme Doughnuts®

Food Item

Serving	Weight (g)	Calories	Carbs (g)	Fiber (g)
Glazed Mini Cruller	3 each	230	32	1
Chocolate Enrobed Doughnut Holes	54	270	37	2
Honey Bun	1 each	96	410	44
Coconut Crème Pie	1 each	128	450	61
Peach Pie	1 each	126	370	51
Cherry Pie	1 each	128	410	56
Apple Pie	1 each	126	400	54

www.mcdonalds.com

McDonald's®

Food Item

Serving	Weight (g)	Calories	Carbs (g)	Fiber (g)
Breakfast Items				
Cheddar Bacon Sausage McMuffin®	1 each	NA	560	27
Egg McMuffin	1 each	NA	290	27
Sausage McMuffin	1 each	NA	360	26
Sausage McMuffin with Egg	1 each	NA	440	27
English Muffin	1 each	NA	140	25
Sausage Biscuit	1 each	NA	410	30
Sausage Biscuit with Egg	1 each	NA	490	31
Bacon, Egg & Cheese Biscuit	1 each	NA	480	31
Biscuit	1 each	NA	240	30
Ham and Egg Cheese Bagel	1 each	NA	550	58
Spanish Omelet Bagel	1 each	NA	690	60
Steak and Egg Cheese Bagel	1 each	NA	700	57
Sausage	1 each	NA	170	0
Scrambled Eggs (2)	1 serving	NA	160	1
Hash Browns	1 each	NA	130	14
Hotcakes (Plain)	1 serving	NA	340	58
Hotcakes (Margarine, 2 pats, & Syrup)	1 serving	NA	600	104
Breakfast Burrito	1 each	NA	290	24
Lowfat Apple Bran Muffin	1 each	NA	300	61
Apple Danish	1 each	NA	340	47
Cheese Danish	1 each	NA	400	45
Cinnamon Roll	1 each	NA	390	50
Sandwiches				
Chicken Breast Parmesan	1 each	NA	490	47
Hamburger	1 each	NA	280	35
Cheeseburger	1 each	NA	330	36
Quarter Pounder®	1 each	NA	430	37
Quarter Pounder® with Cheese	1 each	NA	530	38

62

Exchanges: c=carb; m=medium fat meat, lm=lean meat, vlm=very lean meat; f=fat

Protein (g)	Fat (g)	% Cals from Fat	Saturated Fat (g)	Cholesterol (mg)	Sodium (mg)	Exchanges carb-meat-fat
2	10	39	3	10	190	2c-2f
3	17	57	8	15	240	2c-2f
5	24	53	6	10	170	3c-4f
3	21	42	7	0	220	4c-3f
3	17	41	5	0	220	3c-3f
4	19	42	6	0	200	4c-2f
4	19	43	5	0	220	4c-2f

Protein (g)	Fat (g)	% Cals from Fat	Saturated Fat (g)	Cholesterol (mg)	Sodium (mg)	Exchanges carb-meat-fat
26	39	63	15	285	1430	2c-3m-4f
17	12	37	4.5	235	790	2c-2m
13	23	58	8	45	740	2c-1.5m-3f
19	28	57	10	255	890	2c-2.5m-3f
4	2	13	0	0	240	2c
10	28	61	8	35	930	2c-1m-4f
16	33	61	10	245	1010	2c-2m-4f
20	31	58	10	250	1410	2c-2m-4f
4	11	41	2.5	0	640	2c-2f
26	23	38	8	255	1390	4c-3m-1f
27	38	50	14	275	1570	4c-3m-4f
38	35	45	13	290	1290	4c-4m-1f
6	16	85	5	35	290	1m-2f
13	11	62	3.5	425	170	2m
1	8	55	1.5	0	330	1c-1f
9	8	21	1.5	20	630	4c-1f
9	17	26	3	20	770	7c-3f
13	16	50	6	170	680	1.5c-2m-1f
6	3	9	0.5	0	380	4c
5	15	40	3	20	340	3c-2.5f
7	21	47	5	40	400	3c-4f
6	18	42	5	65	310	3.5c-3f
29	21	39	7	60	1300	3c-3m-1f
12	10	32	4	30	590	2c-1.5m
15	14	38	6	45	830	2c-1.5m-1f
23	21	44	8	70	840	2.5c-3m-1f
28	30	51	13	95	1310	2.5c-3.5m-2f

63

Fig. 11B

Introduction to Fast Food Guide ~ 3347 Dining Out

BD provides this Guide for informational purposes only. It is not intended to be a substitute for professional medical advice, diagnosis or treatment. Do not disregard professional advice or delay in seeking it because of something you read in this guide.

Eating at fast food restaurants can be a challenge for someone with diabetes. Meals can be high in fat, sodium, calories, and carbohydrate. Finding fast food that fits into your meal plan takes planning. If you know the nutritional content of the menu items you want to eat, you can make educated choices within your meal plan. This booklet provides you with the nutritional information for the following fast food restaurants: Arby's®, Baskin Robbins®, Burger King®, Carl's Jr.®, Dairy Queen®, Domino's Pizza®, Dunkin' Donuts®, Einstein Bros Bagels®, Jack-In-The-Box®, Jamba Juice®, Kentucky Fried Chicken®, Krispy Kreme Doughnuts®, McDonald's®, Panda Express®, Pizza Hut®, Starbucks®, Subway®, Taco Bell®, and Wendy's®.

The nutrition information was obtained from the individual restaurants. Menu items were analyzed by computer and not by chemical analysis of the product. Soft drinks are not listed in each of the fast food restaurants. Refer to Einstein's cold beverage section for a sampling of these items. Specialty drinks like milkshakes, smoothies, and coffees have been included. For more detailed information you may visit the restaurant's web site found at the top of each page.

The nutritional information includes serving sizes (weight), calories, carbohydrates, fiber, protein, fat, percent calories from fat, saturated fat, cholesterol, and sodium. Meal exchanges are also included. The legend for the Exchanges are: c=carb (starch, fruit, milk and vegetables); m=medium fat meat, lm=lean meat, vlm=very lean meat; f=fat.

A Registered Dietitian or Certified Diabetes Educator can teach you how to interpret this guide so that you can make food choices that best fit into your meal plan.

Tips for Healthier Eating

- Know your meal plan

Meal plans should be based on the eating habits of the person with diabetes. For most adults, 2-4 carbohydrate servings (30-60g) per meal and 1-2 carbohydrate servings (15-30g) for snacks are reasonable amounts. Children, teenagers and athletes need more calories and may need to eat more carbohydrates. Check with your diabetes educator for the proper amount of carbohydrates for meals and snacks.

Healthy ranges of fat are 30-60g for children, 40-80g for teenagers, 30-60g for women, 40-80g for men, and 80-120g for athletes or those with very high activity levels. People over the age of two generally should get no more than 30 percent of their calories from fat. Limiting saturated fat to less than 10% (no more than 20g for adults) is also important for a healthy heart. The following chart will help you determine the maximum fat intake for your calorie level.

Calories	Fat	% Cals from Fat
1200	30g	23%
1500	40g	24%
1800	50g	25%
2000	60g	27%
2200	70g	28%
2500	80g	29%
2800	90g	29%
3000	100g	30%

~ 3348

- Sodium (Salt) should be limited to 2400mg/day. People with high blood pressure should generally eat 2000mg or less/day. Discuss your sodium limits with your diabetes educator.

- Fill up on fiber - choose salads, vegetables, and whole grain breads. Bring fruit or raw vegetables from home to add to your meal.

- To reduce fat, calories, and sodium, limit fried foods, cheese, bacon, pickles, olives, mayonnaise, tartar sauces, sour cream, and "special sauces". Order sauces, gravy, and dressings on the side and use a small amount. Choose a small order of fries instead of a large. A side salad or a serving of vegetables is a better choice than fries. Eat fresh vegetables, light mayonnaise, and reduced calorie dressings whenever possible.

- Choose grilled or baked chicken or fish instead of beef. Grilled chicken sandwiches or salads are usually the best fast food choices. When ordering beef, choose the smallest burger without special sauces and cheese. Ask for extra lettuce, tomatoes, and onions.

- Control portions - consider splitting a meal with a friend or take half home for another meal. Avoid super-sizing!

- Speak up - don't be afraid to ask questions to make sure that you are getting what you really want. Ask how a sandwich is prepared or if the restaurant offers a low fat salad dressing selection.

- If possible, exercise more to help burn additional calories and control blood sugar levels.

Fig. 11C

- Test blood sugar levels 2-3 hours after your meal to evaluate its effect.
- A high fat meal may cause your blood sugar to rise 6 to 8 hours after eating.

You can work fast food into your diet once in a while and still maintain good health. The basis of a healthy diet is one that is made up of a variety of foods such as whole grains, fruits and vegetables, lean meats, low fat dairy products, and healthy fats. Choosing foods that are not processed foods and are high in fiber will improve the quality of your diet and keep sodium low.

Sample Menu Items

Breakfast:
 McDonald's - 400 calories, 60g carbohydrate, 12g fat, 800mg sodium
 Egg McMuffin
 6 oz Orange Juice

Dunkin Donuts - 400 calories, 70g carbohydrate, 4.5g fat, 270mg sodium
 Bagel with 2T cream cheese

Einstein's - 300 calories, 55g carbohydrate, 4.5g fat, 270mg sodium
 Challah Roll

Krispy Kreme Donuts - 230 calories, 33g carbohydrate, 9g fat, 85mg sodium
 Cinnamon Twist

Lunch and Dinner:
 Subway - 300 calories, 40g carbohydrate, 6g fat, 900 mg sodium
 Roasted Chicken Breast on Country Wheat (6" sub)
 Water

Wendy's - 400 calories, 40g carbohydrate, 10g fat, 900mg sodium
 Grilled Chicken Sandwich
 Side Salad
 Water

Pizza Hut - 400 calories, 60g carbohydrate, 16g fat, 1200mg sodium
 2 Slices Cheese Pizza w/Veggies
 (order pizza with 1/2 the amount of cheese)

Taco Bell - 370 calories, 54g carbohydrate, 12g fat, 1100mg sodium
 Bean Burrito
 Raw Veggies (i.e. baby carrots) from home

Carl's Jr - 400 calories, 45g carbohydrate, 12g fat, 450mg sodium
 Charbroiled Chicken Salad-To-Go
 1/2 Plain Baked Potato w/Chives and 2T Sour Cream

KFC - 350 calories, 35g carbohydrate, 8g fat, 1300mg sodium
 Tender Roast Sandwich without sauce
 Mean Greens

Arby's - 400 calories, 45g carbohydrate, 17g fat, 1000mg sodium
 Regular Roast Beef Sandwich with Garden Salad

Arby's - 450 calories, 45g carbohydrate, 15g fat, 1200mg sodium
 Grilled Chicken Salad
 Child Size Homestyle Fries

Dairy Queen - 400 calories, 35g carbohydrate, 14g fat, 1400mg sodium
 Grilled Chicken Salad with Fat Free Italian Dressing (to reduce sodium, use less salad dressing)
 1/2C. DQ Chocolate Soft Serve Ice Cream

Jack-In-The-Box - 400 calories, 65g carbohydrate, 11g fat, 900mg sodium
 Chicken Fajita Pita
 Fruit (from home)

Snacks:
 Starbucks - 160 calories, 23g carbohydrate, 1g fat, 200mg sodium
 Grande Nonfat Latte

Baskin Robbins - 180 calories, 31g carbohydrate, 2.5g fat, 100mg sodium
 Low fat Espresso 'n Cream Regular Scoop Ice Cream

Taco Bell - 190 calories, 19g carbohydrate, 7g fat, 480mg sodium
 Soft Chicken Taco

Fig. 11D

Fig. 12A




How to play THE REAL MEAL DEAL

Playing the game helps you to figure out and remember salt, cholesterol and carb contents of foods for planning meals.

In this card game, the Suits are Food Groups.

- Protein
- Milk & Yogurt
- Sauces & Spices
- Fruit
- Vegetables
- Heavy Vegetables
- Sweets
- Fat
- Grains & Starch


Each card is a food in a Food Group. The top right corner of the card tells you the Nutrition Facts: how much salt, cholesterol and carbs are in one serving of the food.



Goal of the Game: From the cards in your hand, plan meals (and snacks, if you want) for a day.

Nutrition Goals:

- For each meal, there must be at least: 40 carbs
- For each day, there must be at least: 2 servings from each Food Group
- and no more than: 1500 mg of salt
300 mg of cholesterol
150 g of carbs



THE WINNER is the first one who lays down the most cards to plan 3 meals without going over the limits.

Tip: Use the Tally Sheet to keep track of the Nutrition Goals as you go!

TO PLAY:

- 1** The Dealer shuffles and deals 10 cards to each player, then places the rest in a DRAW PILE, face down.
- 2** The First Player lays down cards for a day's meals and snacks, matching the Nutrition Goals.
- 3** During a turn, a Player may:
 - Move cards from meal to meal but not pick them up again
 - Trade cards with other Players, if they agree
- 4** To end the turn, a Player
 - Places any unwanted cards in the DISCARD PILE next to the DRAW PILE
 - Draws enough cards from the DRAW PILE to have 10 in hand again

When the DRAW PILE is gone, Players turn over the DISCARD PILE and keep going.




Fig. 13

136 → **Meal Planner** - THIS IS A SAMPLE. Write your own plan on the back.

3612 → FOOD GROUP 3614 → SERVING SIZE 3616 → How many 3618 → How much 3620 → How much 3622 →

3610 →

	FOOD GROUP	SERVING SIZE	How many	SERVINGS/CARBS?	How much	How much	
					Salt?	Chol?	
BREAKFAST	protein egg beaters	1 3616 1	0	0	2		
	fat butter	1 TBsp 1	0	51	0		
	fruit orange	1 1	13	1	0		
	vegetables peppers, onions	FREE! FREE!	FREE!	FREE!	FREE!		
	grain & starch whole wheat toast	1 slice 1	13	148	0		
	milk skim milk	1 cup 1	12	128	6		
	spices garlic powder	1 tsp 1	2	0	0		
	sweets marmalade, low sugar	1 TBsp 1	0	0	6		
3630 → BREAKFAST TOTALS: 7				40	328	14	
LUNCH	protein tuna fish, water pkg.	2 ounces 1	0	202	2		
	fat mayonnaise	1 Tablespoon 1	2	125	5		
	fruit blueberries	1/2 cup 1/2	11	5	0		
	vegetables lettuce,	1/4 head 1	4	0	0		
	grain & starch whole wheat, low carb	2 slices 2	18	230	01		
	milk sugar-free yogurt	6 ounces 1	12	119	11		
	spices						
	sweets Olive oil & vinegar	2 TBsp 1	0	0	0		
3640 → LUNCH TOTALS: 8				47	681	18	
DINNER	protein chicken breast	3 ounces 1	0	128	146		
	fat						
	fruit strawberries	1/2 cup 1	10	1	0		
	vegetables asparagus	4 spears 1	2	0	0		
	grain & starch brown rice	2/3 cup 2	2	2	2		
	milk skim milk	4 ounces 1	6	64	3		
	spices						
	sweets sugar free ice cream	1/2 cup 1	20	50	0		
3650 → DINNER TOTALS: 7				50	245	146	
SNACK?	milk 1%	1/2 cup 1	6	62	6	3660	
Whole Day's Totals				23	143	1316	18
							3670

The Shopping List

Make copies before you use this.

Circle the ones to buy. Write other things below.

Cut the Salt, buy:

Cut the Fat, buy:

Basil Leaves	Ginger Root	Chicken
Cayenne	Horseradish	Fish
Pepper	root	Turkey
Celery Seed	Lemon Grass	Egg Whites
Chives	Lemons	Fat free milk and
Cumin	Limes	sour cream
Curry Powder	Mint	Fat free yogurt
Dill	Onions	Margarine in tubs
Fennel	Parsley	Canola and other
Garlic	Hot Peppers	vegetable oils
Green Onions	Tarragon	Baking fat substitute

Buy Free Vegetables

Fruits to buy:

Asparagus	Leeks	Apples
Broccoli	Lettuce	Apricots
Brussel Sprouts	Mushrooms	Blueberries
Cabbage	Onions	Oranges
Cauliflower	Parsley	Plums
Celery	Peppers	Raspberries
Cilantro	Spinach	Strawberries
Cucumber	Sprouts	Tangerines
Eggplant	Tomatoes	Buy sweeteners
Green beans	Watercress	
Greens	Yellow squash	
	Zucchini	

The Shopping List

Make copies before you use this.

Circle the ones to buy. Write other things below.

Cut the Salt, buy:

Cut the Fat, buy:

Basil Leaves	Ginger Root	Chicken
Cayenne	Horseradish	Fish
Pepper	root	Turkey
Celery Seed	Lemon Grass	Egg Whites
Chives	Lemons	Fat free milk and
Cumin	Limes	sour cream
Curry Powder	Mint	Fat free yogurt
Dill	Onions	Margarine in tubs
Fennel	Parsley	Canola and other
Garlic	Hot Peppers	vegetable oils
Green Onions	Tarragon	Baking fat substitute

Buy Free Vegetables

Fruits to buy:

Asparagus	Leeks	Apples
Broccoli	Lettuce	Apricots
Brussel Sprouts	Mushrooms	Blueberries
Cabbage	Onions	Oranges
Cauliflower	Parsley	Plums
Celery	Peppers	Raspberries
Cilantro	Spinach	Strawberries
Cucumber	Sprouts	Tangerines
Eggplant	Tomatoes	Buy sweeteners
Green beans	Watercress	
Greens	Yellow squash	
	Zucchini	

Fig 14A

Fig. 14B

Meal Planner

FOOD GROUP

3760

BREAKFAST	protein
	fat
	fruit
	veggies
	grain & starch
	milk
LUNCH	protein
	fat
	fruit
	veggies
	grain & starch
	milk
DINNER	protein
	fat
	fruit
	veggies
	veggies
	grain & starch
milk	
SNACK ?	

Meal Planner

FOOD GROUP

BREAKFAST	protein
	fat
	fruit
	veggies
	grain & starch
	milk
LUNCH	protein
	fat
	fruit
	veggies
	grain & starch
	milk
DINNER	protein
	fat
	fruit
	veggies
	veggies
	grain & starch
milk	
SNACK ?	

Get Out and Walk



Check the Record Book, Page 3. Did the doctor say walking is right for you? The page also should say how many minutes a day the doctor wants you to walk. If you haven't asked yet, don't start until you ask.

Why Walk? 23802

Most people can do it!

Walking gets the blood flowing in your legs and feet.

It helps bring the blood sugar down if it's too high.

You'll notice the payoff in 3 months or less:

- You'll have more energy
- You'll sleep better

Walk Every Day 23810

You need six things to be a regular walker:

1 Shoes: Comfortable shoes, the right size, made of canvas or soft leather. Have a professional help you fit them. Wear socks without seams. Put them on the Shopping List!

2 Time: Pick a regular time of day to walk. Stick to it. If you try for every day, you'll probably make it 5 times a week. Five times a week is good!

3 Watch or Pedometer: Time yourself out and back or count how many steps you take to see your progress.

4 Water: Take a water bottle - Drink often; you need extra while exercising.

5 Emergency Tools:

- Make a "Just in Case" bag: put in the things you eat for low blood sugar, like crackers or Glucose Tabs
- Slip in the Emergency Card

Keep these things in the same place so you can find them fast when you're ready to go.

6 Guts: Go! Find someone to walk with you--you'll be more likely to get up and out. But even if you have to go alone, just go--try a shopping mall.

If you ever feel faint, dizzy or can't talk easily, stop. Rest until you feel OK again.



Fig 15A

Make Walking a Habit

It's good to have a way to keep going day after day. The best way is to keep track of the program. Here's how:

1 On your next walk, time yourself or get a pedometer to count your steps.

2 The charts are walking plans.
Pick your level: • Beginner
• Mover
• Expert

3 Walk slowly at first. Then speed up. At the half way point, turn back while you keep walking fast. Then slow down. You're back where you started.



Beginner	SLOW	FAST	FAST	SLOW	TOTAL
every day Week 1	1 minute	1 minute	1 minute	1 minute	4 minutes
every day Week 2	1 minute	2 minute	2 minute	1 minute	6 minutes
every day Week 3	2 minutes	2 minutes	2 minutes	2 minutes	8 minutes
every day Week 4	2 minutes	3 minutes	3 minutes	2 minutes	10 minutes
every day Week 5	3 minutes	3 minutes	3 minutes	3 minutes	12 minutes
every day Week 6	3 minutes	4 minutes	4 minutes	3 minutes	14 minutes

Remember, if you ever feel faint, dizzy or can't talk easily, stop. Rest until you feel OK again.



4 When you get back home, go straight to the Record Book. Write how many minutes or steps you walked.

5 Do the same walk every day for a week. The next week, add one minute to each fast part. You will add 2 minutes to each week by adding a minute to each fast part.

Fig. 15B

Walking Plans cont'd

3826-5 MOVER

	SLOW	FASTER	FAST	SLOWER	TOTAL
every day Week 1	5 minutes	3 minutes	TURN BACK 3 minutes	5 minutes	16 minutes
every day Week 2	5 minutes	4 minutes	TURN BACK 4 minutes	5 minutes	18 minutes
every day Week 3	5 minutes	5 minutes	TURN BACK 5 minutes	5 minutes	20 minutes
every day Week 4	5 minutes	6 minutes	TURN BACK 6 minutes	5 minutes	22 minutes
every day Week 5	5 minutes	7 minutes	TURN BACK 7 minutes	5 minutes	24 minutes
every day Week 6	5 minutes	8 minutes	TURN BACK 8 minutes	5 minutes	26 minutes

Don't over do it! If you can talk or sing easily while walking, you are exercising at the right pace. Breathing deeply is OK, but if you are puffing, panting, feel faint or dizzy, stop. Rest until you feel OK again.

3828-5 EXPERT

	SLOW	FAST	FAST	SLOW	TOTAL
every day Week 1	5 minutes	10 minutes	TURN BACK 10 minutes	5 minutes	30 minutes
every day Week 2	5 minutes	11 minutes	TURN BACK 11 minutes	5 minutes	32 minutes
every day Week 3	5 minutes	12 minutes	TURN BACK 12 minutes	5 minutes	34 minutes
every day Week 4	5 minutes	13 minutes	TURN BACK 13 minutes	5 minutes	36 minutes
every day Week 5	5 minutes	14 minutes	TURN BACK 14 minutes	5 minutes	38 minutes
every day Week 6	5 minutes	16 minutes	TURN BACK 16 minutes	5 minutes	40 minutes

6 Remember, after you walk, write in the Record Book how many minutes or steps you walked so you can keep track of your progress.




Fig. 15 B (cont.)

When You Travel


1 Ask your doctor for extra med prescriptions. Keep them in your wallet in case your meds are lost or stolen.

2 Call the home or hotel where you're going. Get a hospital number you can call if you have an emergency.

3 Check that you have enough supplies to pack. Take double what you need—just in case. If you're short, buy more.

4 Pack in a small bag:

- All the supplies and meds you'll need
- Glucose Tabs or snacks, like breakfast bars
- Your ID and Emergency Medical Card
- Emergency phone number of the place you're going.



5 Wear a Medical ID Alert Bracelet when traveling.



If you change the time you eat, make the same change in the time you take your medicines.

If you're going by air, all your supplies are welcome on the plane! You just need to follow these rules:

If you take:

You must also take:

Syringes

Insulin

Insulin

The box the insulin came in with the prescription label

Insulin Pens


The box the pen came in with the prescription label

Insulin Pump

The box the insulin came in with the prescription label

Lancets

Caps on the Lancets and hard Sharps disposal container



If you've lost the box or label, call the drug store or mail order supplier to get a copy.

Call the airline to see if any rules have changed—or are different for international travel:

American Diabetic Association: 1-703-549-1500

Transportation Security Administration: 1-866-289-9673

At the Airport

Tell any one who asks that your extra carry-on is for diabetes.

If you wear a pump, ask them to check just by looking. You don't have to take it off. If you have a problem, ask for the supervisor.

If there is still a problem, ask for the FAA Grounds Security Commissioner. You have a right to private screening.



Fig. 15C

Fig. 16A

Protect Your Skin and Feet




Step by Step

Get into the Daily Habit 2 3850

There are five things to do for skin and feet every day:

3852 **1 WASH:** Run water and test the temperature. It should be warm, not too hot or too cold. Use a sudsy washcloth on your feet. For a bath or shower, get out in ten minutes or less.



3854 **2 DRY:** Pat yourself dry--don't rub. Pat under your breasts and arms, between your legs, and between your fingers and toes.



3856 **3 SMOOTH:** Rub Lotion all over, but NOT between your toes. *There's natural oil there already.* When you buy lotion, choose one that *alcohol* is not in the first five ingredients



3858 **4 CHECK:** Look for changes:
 • Blisters or sore places
 • Thick, hard places
 • Cracks or cuts that don't heal
 • Red, white, black or purple spots



Have someone else help you check your head and the bottoms of your feet.

If you're by yourself, use the Mirror to look carefully at your head and the bottoms of your feet. If you see any skin changes, call the clinic.



3860 **5 DRESS** Always wear fresh, clean hose or socks. Shake out shoes before you put them on.

Put "extra slippers" on your Shopping List. Put a pair of slippers any place where you might take off your shoes: by the TV, in the bathroom, near the bed. NEVER GO BAREFOOT.

Fig. 16 B

3870
Other times for Skin and Feet

3872 **TWICE A WEEK**

Wash your hair--or more often.

3874 **ONCE A WEEK**

After a bath or shower, *if the doctor said it's OK to cut your own toenails*, get your toenail clippers. *If you don't remember if you can cut your own toenails, check the Record Book, page 3.*

Check if your clippers are the right kind, with the cutters straight across, like the picture here.

If they are not the right kind, stop. Put "Straight-edge Toenail Clippers" on your Shopping List.

In the Record Book on Page 4, it says to have someone at the clinic show you how to cut your toenails. *Just as a reminder: Use straight-edge clippers. Put the cutting edge across the end of your toenail, even with the end of your toe. Cut the nail straight across.*

3880 **ONCE A MONTH**

Go to the back page of the Record Book. Do the LEAP test.

It's better to have someone else do it--the idea is to see if you can feel the stick. If you do it yourself, you know exactly when it touches you!

3882 **SUMMERTIME**

When you go outside:

- Use sunscreen with SPF 15 or higher. Check the label.
- Wear long sleeves and a hat.
- Use bug spray.

3884 **WINTERTIME**

- Use a humidifier or put bowls of water near the heat registers. Dry skin can crack--water in the air helps skin stay soft.
- Bundle up when you go outside!



The Right Meds at the Right Time




Organize Them ~ 4010

Where do you keep medicines? Have you made a list?
Bottles and papers may be all over the house.


- 4020 **1** Bring everything you take for health—even vitamins or special oils—to a table, with the Record Book and a pen or pencil.



- 4030 **2** Sort the meds into groups, like this:
- 4032 Vitamins, herbs, teas, anything you take the doctor might not know about
 - 4034 Blood Sugar meds
 - ☐ Shots
 - ☐ Pills
 - 4036 Heart meds
 - 4038 Other prescription meds
 - 4040 Mystery meds—anything you don't know what it's for!




- 4050 **3** Open the Record Book Page 4, *Meds I Take*.
In the first section, write the names of the vitamins and other things you take without a prescription.



Now you're ready to write the prescription meds.
If you already have a list, just tape it to the page.
But if you don't always remember what each med is for, or if you want a fresh start, follow the steps below.

Know What They're For ~ 4100

- 4** You saw in the Short Story that you do five things to keep your body rolling along:
1. Charge up the heart,
 2. Lighten the load
 3. Open the road
 4. Don't flood the engine
 5. Don't run out of gas.



There's a Med for each one. Turn the page and get clear on which med does what.

Fig. 17A

Which Med Does What *a matching game..*

Start with a bottle from the heart meds. Match the name on the bottle to a name on the list. *All meds have 2 names:*


The **brand** name in red.

The **generic** name is next to it in black.

If a name on a bottle matches a name on the page, write it in the Record Book.

Turn the bottle to see how much to take and when.

Write that in the Record Book, too.



Heart Meds


1 To Charge Up the Heart

Check your Mystery Meds.
Are any of them here?

These pills make the heart pump strong, like a body builder.

Digoxin	digitalis
Lanoxin	

If you don't have one of these meds, ask the doctor if you need digitalis.



These pills make the heart pump steady, like a clock ticks.

Betapace	sotalol
Codarone	amiodarone
Pacerone	
Tikosyn	dofetilide
KayCiel	
K-Lor	potassium
K-Dur	
Mag-Ox	magnesium oxide
Uro-Mag	

If you don't have any of these meds, ask the doctor if you need an antidysrhythmic.



These make each pump count for more, like using a lower gear on a bicycle going uphill.

Coreg	carvedilol
Lopressor	metoprolol
Toprol-XL	
Tenormin	atenolol

If you don't have any of these meds, ask the doctor if you need a beta blocker.



Fig. 17B

Fig. 17C

2 To Lighten the Load

Check your Mystery Meds.
Are any of them here?

4130

These pills flush out extra water, like an open hydrant.

4132

Aldactone	spironolactone
Bumex	bumetanide
Demadex	torsemide
Lasix	furosemide
HydroDIURIL	hydrochlorothiazide
Inspira	epplerenone
Maxizide	triamt/HCTZ
Zaroxolyn	metolazone



If you don't have any of these meds,
ask the doctor if you need a water pill.

These pills make blood thinner, like mixing turpentine in paint.

4134

Aspirin	acetylsalicylic acid, aspirin
Ecotrin	
Coumadin	warfarin
Plavix	clopidogrel
Ticlid	ticlopidine

If you don't have any of these meds,
ask the doctor if you need an anticoagulant.



Fig. 17D

3 To Open the RoadCheck your Mystery Meds.
Are any of them here?

These pills open arteries and veins, like an empty well paved freeway.

Accupril	quinapril
Altace	ramipril
Capoten	captopril
Lotensin	benazepril
Mavik	trandolapril
Monopril	fosinopril
Prinivil	lisinopril
Univasc	moexipril
Vasotec	enalapril
Zestril	lisinopril

If you don't have any of these meds,
ask the doctor if you need an ace inhibitor.

These pills keep arteries and veins from getting tight, like stretching.

Atacand	candesartan
Avapro	irbesartan
Benicar	olmesartan
Cozaar	losartan
Diovan	valsartan

If you don't have any of these meds, ask the doctor
if you need an angio II receptor blocker.

Norvasc	amlodipine	Cardizem	diltiazem
Calan		Dilacor-XR	
Isoptin	verapamil	Plendil	felodipine
Verelan			

If you don't have any of these meds, ask the doctor if you need a calcium channel blocker.

These pills relax the arteries, like blocking out stress.

Nitro Bid	
Nitrostat	nitroglycerin
Nitrogl	

If you don't have any of these meds, ask the doctor
if you need nitroglycerin for chest pain.

Apresoline	hydrALAZINE
Iso-bid	
Isordil	isosorbide dinitrate

If you don't have any of these meds, ask the doctor if you need a vasodilator.

These pills stop fat build up.

These pills attack animal fat, like a matador.

Lescol	fluvastatin
Lipitor	atorvastatin
Mevacor	lovastatin
Pravachol	pravastatin
Vytorin	ezetimibe and simvastatin
Zocor	simvastatin

If you don't have any of these meds, ask the doctor if you need a statin.



These pills attack vegetable fat, like cutting weeds.

Atromid-S	clofibrate
Lopid	gemfibrozil
Tricor	fenofibrate

If you don't have any of these meds, ask the doctor if you need a fibrate.



This pill stops cholesterol before it enters the blood.

Zetia | ezetimibe

If you are not on this pill, ask the doctor if this would help.

Fig. 17E

Blood Sugar Meds

4 To Keep from Flooding the Engine

4220 These pills put a lid on the sugar. They stop it from going up too far, like a ceiling stops balloons.

4222

Actos	pioglitazone
Avandia	rosiglitazone
Glucophage	
Glucophage XR	metformin
Riomet	
Glyset	miglitol
Precose	acarbose



These pills put a leash on the sugar. They pull it back down, like pulling down on balloon strings.

4224

Amaryl	glimepiride
Diabinese	chlorpropamide
DiaBeta	
Glynase	glyburide
Micronase	
Glucotrol	
Glucotrol XL	glipizide
Prandin	repaglinide
Starlix	nateglinide



4226 This shot can be a boost for people just taking pills for blood sugar control.

Byetta	exenatide
--------	-----------

4228 This shot can be a boost for people taking pills or insulin for blood sugar control.

Apidra	insulin glulisine
Symlin	pramlintide acetate

4229 This inhaled insulin can be a boost for people taking pills or insulin for blood sugar control.


EXUBERA	insulin human [rDNA origin]
---------	-----------------------------

Insulin--it's the only thing that can open the spare tank in addition to keeping the engine from flooding. Turn the page to learn how the different kinds of insulin work.

4230

✓ 4350

✓ 4300



4312




4314

4316

4318

Five Ways to Keep Your Medicine Plan on Track: ~ 44100

- 44110 **1 Be ready for emergencies**
- When you first open the nitroglycerin pills, mark the date on the Calendar. Count 5 months ahead. Make a note to order new pills that day.
- Fill the NITRO NECKLACE with 6 nitro pills.
 - Wear the Necklace.
 - Put new pills in every month.
 - Keep the bottle in a dark, dry place.



- 4420 **2 Be ready for the week**
- Find the **Pill Organizer**. Check the times you wrote in the **Record Book** for taking each pill. Fill the boxes with the right pills at the right time for each day, for the week.

When you go out for the day, carry the day's strip of boxes with you. Put your finger in the hole on the back of the tray and pop out the strip.

If you use insulin and are going out, pack a carry bag with meds, alcohol pads, a syringe, insulin and the Record Book.

- 4430 **3 Be ready to fight temptation**
- When you feel better, a voice from somewhere says, "You're OK now. Stop taking those stupid meds."
- Don't listen!** The voice is wrong. The reason you feel better is that you are taking the meds.
- If you stop, you'll slide back into feeling sick and then slide back into the hospital.*



- 4440 **4 Be ready to make a record**
- Keep the Record Book in your lunchbox, purse, briefcase, or wherever you can reach it when you need to write in it during the day.




Fig. 17G

How to Give Insulin

CAREGUIDE

Step by Step

Prepare for Care:




1 Collect the things you need: the Record Book, a pen, insulin, syringe, an alcohol pad, a clean, fresh towel and "SHARPS" container.

2 Take everything to a clean, flat place to work near a sink. Wash your hands with soap and water. Dry them on the clean towel.

3 Look at the picture to learn the parts of each item. If you can't see the numbers on the syringe, get someone to help you.

4 Turn to Page 3 of the Record Book. Check which insulin you need and how much.

4538 If the kind of insulin you have is not the same, call your doctor or clinic now.



5 Look at the date on the insulin to be sure the date is still in the future.

If the date is past, call the pharmacy and get new insulin.





Fig. 18 A

Pick a Spot to Start ⁴⁵⁵⁰

The colored areas on the drawings of the body show places where you can give insulin. Use a different spot on the body every time so the skin doesn't get hard or sore.

1 Look how the insulin areas are divided into separate, small squares, or "shot spots." Imagine each shot spot has a number going across, row by row.



2 ⁴⁵⁵⁸ Check the Record Book, to see which areas you were told are best for you to use. Pick an area of your body to start.

3 ⁴⁵⁶⁰ Wash the shot spot you are going to use this time with soap and water.

In the future when you give insulin, follow the pretend row, using a different shot spot each time. When you get to the end of a row, imagine the next row and start again. Use all the shot spots in one area and then move to another area.

Fig. 18B

Fill the Syringe



- 4572
- 1 Roll the insulin between your hands to mix. Never shake it.

If you have a new bottle, take off the colored cap. Leave on the rubber stopper and the metal band under the cap.


- 4574
- 2 Use an alcohol pad to wipe the rubber stopper on the bottle.

- 4576
- 3 Take the caps off the plunger and the needle. Lay them on the table.



- 4578
- 4 Check the Record Book again to be sure of the kind of insulin and number of units you need. Find that same number on the barrel of the syringe.

- 4580
- 5 Pull back the plunger until the end, inside the barrel, is at the number.



- 4582
- 6 Hold the insulin bottle firm on the table. Push the needle all the way through the rubber stopper. Then push the plunger all the way into the syringe.

You are filling the bottle with air.

Fig. 18C

- 4584
- 7** Turn the bottle and syringe upside down. Pull the plunger back to the number of units of insulin you need.

If you see air bubbles, push the plunger in to put the insulin back into the bottle. Slowly pull it out again to your number. Repeat until there are no air bubbles. Check again that the plunger is at the right number.

- 4586
- 8** Put the bottle down. Hold the barrel to pull the syringe out of the bottle. Lay the syringe on the table. Don't let the needle touch anything.



Give the Injection 4590

- 4591 **1** Find your shot spot again. Gently pinch up a fold of clean, dry skin.
- 4592 **2** Hold the syringe so the needle will go straight in. Push the needle in all the way. Push the plunger in all the way. Let go of the skin.
- 4593 **3** Pull the needle out. Press tissue over the shot spot.




- 4** Put the syringe in the "SHARPS" container.

4594

- 4595 **5** Put the "SHARPS" container where children can't reach it

Before the container is full, call City Hall to find out the rules for throwing away medical sharps.



- 4596 **6** Write the time and the shot spot in your Record Book in the top section for Meds.

If you did anything different, write the kind of insulin you used and the number of units you gave yourself.

Fig. 18D

Automatic Blood Pressure Monitor



Step by Step


Page 1

Read all the steps before you begin. When this guide first tells you to use an item, its name appears in red. Keep everything in the Kit until the Careguide tells you to use it.

Prepare for Care ~ 5010

5012 **1** Get the machine ready. The first time you take your blood pressure, or any time the "REPLACE BATTERIES" light comes on, you'll need to put in batteries.

- Lay the Monitor face down on a table. Press down on the word "OPEN." Slide the cover up and off.
- Put in 4 batteries. Match the + and - marks on the batteries to the + and - marks inside the case.
- Lay the cover flat on the back of the machine so that all the batteries show. Slide the cover down over the batteries until it snaps in.




5014 **2** Get yourself ready. Bring a pen and the Journal to the table. If you just had a smoke, ate something, exercised, showered or if you feel stress, relax for at least 30 minutes.



Put the Cuff on ~ 5020

5022 **3** Use your left arm every time unless you know a good reason why not. If you have bulky clothes on your arm, take them off. Don't just push the sleeve up! Sit next to the table. Put both feet flat on the floor.

5024 **4** Pull the sticky tab open on the Cuff, so the metal bar can slide back and forth. Notice the square of hooks on the end of the tab.



5026 **5** Hold the cuff with the white stripe and the tube on the bottom, pointing down your arm. Slip your arm through the cuff.



Fig. 19A

Automatic Blood Pressure Monitor




Step by Step

Page 2


5028  Push the cuff up until the bottom edge is 1 inch above the bend inside your elbow.

Move the cuff around until the tube is right above the bend. Run the tube straight down to your hand, palm up.



5030  Pull the loose flap of the cuff until the cuff is snug around your arm.

While you pull, move the cuff around to keep it 1 inch above the bend in your elbow and to keep the tube running straight down the center of your arm.



5032  Press the flap against the cuff to hold it tight.


The cuff is tight enough if you can only squeeze two fingers under the edge at bend.

The **whole** square of hooks should fold over and stick to the Cuff. If only **part** of the square of hooks folds over the metal bar, the Cuff is too small for you.

If the cuff is too small for you, call 1-877-CAREKIT (toll free) and order a large cuff.

Use the Monitor

5042  Push the tube into the hole on the left side of the Monitor.



5044  Put the Monitor on the table so you can see the screen.

Put your elbow on the table with the palm of your hand up. The cuff should be level with your heart.



If you sit up straight, it should be level!

Fig. 19B

Automatic Blood Pressure Monitor




Step by Step

Page 3




5046 Press the red **POWER** button with your right hand.

A lot of numbers come up, then change to a zero. Three beeps tell you the machine is ready.




5048 Press the blue **START** button with your right hand.

The cuff will get tight, but not too tight. Stay still while it works. Don't talk, eat or smoke. If this is the first time you are using the machine, it may start and stop a few times, getting used to you. That's normal.



If you think anything is really wrong, like the cuff gets so tight it hurts, press **POWER. The machine will stop.**

When the Monitor is finished, it shows three numbers on the screen. The number in the middle is a smaller size than the numbers on the ends.



5050 Open the Journal.

Find the space called **Pulse**. Write the number you see in the middle of the screen in the space for pulse.

Find the space called **Blood Pressure**. Write the first number, a slash and the last number in the space.

WEIGHT	Week of
	SUN
BLOOD PRESSURE	143 / 79
PULSE	57
EXERCISE	



5052 Press the **POWER** button to turn off the Monitor.

If you want to take another blood pressure reading, wait 15 minutes for your arm to go back to normal.



If this is the first time you have used the Monitor, you have one more job. Turn the page to see how to Set the Pumping Target on the Monitor.

Fig. 19C

Automatic Blood Pressure Monitor




Step by Step



Set the Pumping Target ~ 5060

You only need to set the Pumping Target once.


- 5062 **1** Press the **POWER** button. When you hear the beeps and see a zero on the screen, press the white **MEMORY** button on the left side of the machine.



Your last blood pressure reading will come up.

- 5064 **2** What is the big number on the left of the screen? Add 30 to it to get your "PUMPING TARGET."

For example: if your number is 143 like the photo, add 30 = 173. If your number is 151, add 30 = 181. Remember that number for the next step.




- 5066 **3** Press the white **SET** button.

The screen will show a number on the right side. Is that number lower than your Pumping Target? Each time you press the SET button, the number goes up. Keep pressing the SET button to the *first* number that is the *same* or *higher* than your Pumping Target. Then just stop pressing. The Monitor is now set.

If you miss the number, don't worry. Keep pressing the SET button! The screen goes up to 280 and starts over.

- 5068 **4** Read the small book in the Monitor box now. Look for the section on "error messages," so you'll know what to do if you see one.



SPECIAL INSTRUCTIONS FROM MY DOCTOR OR NURSE

Fig. 19A

How To Quit Smoking



Step by Step

KNOW THE FACTS *they're surprising! ~5110*

FACT

Most people who try to quit succeed. But it takes more than one try.

FACT

Just as many people succeed without signing up for a program as succeed with a program.

FACT

Heavy smokers succeed in quitting just as often as light smokers.

FACT

Most people who succeed do it when they have some other big change in their life.



THAT MEANS

1. If you keep trying, chances are 2 out of 3 you'll succeed.
2. You can sign up for a program OR do it on your own.
3. It doesn't matter how much you smoke right now.
4. You can use "my health" as the big change in your life.

64% of all people who try to quit succeed after multiple tries.
Stanley Schachter, Columbia University: American Psychologist, April, 1982.

SO GET READY ~5120

- 1 If you don't already have toothpicks, sugarless gum and diet juice, put them on the Shopping List.

If you don't think you can stop absolutely this week, put a different brand with more nicotine on the Shopping List, to change the taste and make you feel sick.

- 2 Set the date to stop. Mark it down so you can't kid yourself about when you said you were going to stop. Tell family and friends you're going to quit for two days.

- 3 Tell yourself you're only going to quit for two days --it's no big deal.






Fig. 20A

DO IT: QUIT ✓ 5130

1 Get rid of all cigarettes around you. If you can't bring yourself to throw them away yet, at least store the lighter or matches far away from the cigarettes.



2 When you have the urge to smoke - ask yourself why:

If you smoke to be social, INSTEAD:

- Visit a non-smoker friend
- Sign up with a community volunteer group

If you smoke to relieve stress, INSTEAD:

- Squeeze a "stress ball"
- Do some exercises in your chair
- Ride a bike
- Play some ball



If you smoke to keep from eating, INSTEAD:

- Bite a toothpick
- Chew some sugarless gum
- Drink some water
- Drink diet juice

3 At the end of Day Two, ask yourself how you did. Don't kid yourself. If you smoked fewer than usual, GREAT--but you still haven't quit. The goal is to stop completely. Change something in the plan to make it work better.



Repeat the "Do It" Steps 1-2-3 every two days.

BRACE YOURSELF ✓ 5140

If you feel sick at first, don't worry. It won't last.

If you feel like having one, don't panic. The urge will pass.

If you have a bad day, don't beat on yourself. Just start again.

Watch for the good changes that will come in few weeks:

- More energy
- Easier to breathe
- Everything smells and tastes better.



Fig. 20B

5240

3 ACT


- Negative neighbor? Walk away.
- Task too big? Break it down. Get help.
- Stakes too high? Get feedback early.
- Lonely? Volunteer.
- Sad? Get a pet.
- Nervous? Cut out coffee and sugar.
- Made a mistake? Admit it and move on.
- No time? Say NO--suggest someone else.
- Frowning? See a funny video on Google.
- Stuck? Walk around--outside if you can.
- Overwhelmed? Make a "TO DO" list.

Whatever your stress,
someone else has it, too.
Find them.

5250


4 RATE

Where does
today's stress
rate on
this meter?



Four steps to calm: **B-S-A-R**
Breathe, **S**tretch, **A**ct, **R**ate.
For stress, **BSAR**!
To do anything else would be bizarre.

Fig. 21A



You know bringing down stress does good things for your health.

Stress is getting to you when...

- Headaches come often
- Worry grabs you and won't let go
- You get easily distracted
- Others say you're "touchy"
- Sleep won't come at night
- You care so much you just *don't care*
- Muscles are too tight or twitching
- Breathing is hard
- An upset stomach makes diarrhea
- It all seems so hopeless

Now follow the Four Steps to calm

1 Breathe


- A. Close your eyes.
- B. Breathe in while you count to 5 and raise your shoulders up and back.
- C. Breathe out while you count to 10 and slowly lower your shoulders.

REPEAT 5 TIMES



2 STRETCH

- A. Lace your fingers, palms in.
- B. Reach to the ceiling, palms down.
- C. Turn your palms up. *Reach higher.*
- D. Lean to the left.
- E. Lean to the right.



- F. Lift legs.
- G. Turn feet in circles.



- H. Roll head to the left.
- I. Roll head to the right.

Fig. 21B

**INTEGRATED BLOOD SUGAR CONTROL,
BLOOD PRESSURE CONTROL AND HEART
FAILURE SELF-CARE SYSTEM AND METHOD**

PRIORITY CLAIM

[0001] This application is a non-provisional of, claims priority to and the benefit of U.S. Provisional Patent Application Ser. No. 60/806,201 filed Jun. 29, 2006; U.S. Provisional Patent Application Ser. No. 60/806,200 filed Jun. 29, 2006; and U.S. Provisional Patent Application Ser. No. 60/806,203 filed Jun. 29, 2006, the entire contents of which are incorporated herein by reference.

COPYRIGHT NOTICE

[0002] A portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner has no objection to the photocopy reproduction by anyone of the patent document or the patent disclosure in exactly the form it appears in the Patent and Trademark Office patent file or records, but otherwise reserves all copyright rights whatsoever.

TECHNICAL FIELD

[0003] The present application relates in general to methods, systems and apparatus for integrated blood sugar control, blood pressure control and heart failure self-care by a person.

BACKGROUND

[0004] The health care system in the United States has dramatically changed in the last several decades. Lengthy hospital stays after surgical and other medical procedures have decreased or have been eliminated, at least in part, to reduce overall health care cost. When patients leave hospitals or do not stay in hospitals, the patients often require more professional home care and/or more self-care at home. Self-care at home has necessitated the need for better, more advanced, more user friendly, and more easier-to-use self-care systems, apparatus and methods. For example, the assignee of the present application has developed and distributed many individual self-care systems. Such self-care systems have typically been designed for self-care after an individual surgical or other medical procedure (such as the Post Surgery Drain System described in U.S. Pat. No. 6,926,708).

[0005] Self-care systems, apparatus and methods have also been distributed for individual medical conditions (such as for asthma control, blood pressure control, blood sugar control, and healthy foot care). The assignee of the present application has developed and marketed several individual self-care systems, each focused solely on self-care for an individual medical condition.

[0006] However, many people have multiple, different, simultaneously existing medical conditions. Multiple, different, simultaneously existing medical conditions can be related to each other, or may be unrelated to each other. For instance, a person may have heart disease and high blood pressure. These conditions may be considered to be related, in a sense, because they both relate to the heart. On the other hand, a person may have asthma and high blood pressure. These conditions may be, considered to be in a sense, unrelated.

[0007] Patients with multiple, different, simultaneously existing medical conditions can use the known individual self-care systems to individually care for each of their conditions. For instance, a person with heart disease and high blood pressure may use an individual heart disease self-care system and may also use an individual high blood pressure self-care system. While these self-care systems may have some overlapping instructions, they may also have contradictory or inconsistent instructions. The concurrent use of these two individual different self-care systems may also require the person to duplicate certain efforts. Using two or more different self-care systems may also cause the person to become confused, discouraged, or overwhelmed. Moreover, many people tend to be able to handle only a limited amount of information and instructions. Using such multiple individual systems greatly lessens the likelihood that the patient will strictly follow either of the self-care systems. Moreover, even if a person is compliant with both individual self-care systems, the systems may be contradictory to a certain extent or may not be effective as possible. The problems get even more complicated when the person has more than two conditions.

[0008] Unfortunately, prior to the present invention, there were no known self-care systems which effectively provide single integrated systems for self-care by people with the multiple different medical conditions which the present disclosure addresses.

[0009] Prior to the present invention, these problems were not being addressed in the healthcare industry. Rather, the medical literature has been first discussing substantial problems with known clinical practice guidelines or treatment protocols. Clinical practice guidelines and treatment protocols are substantially different than self-care systems. Clinical practice guidelines and treatment protocols are standardized guidelines for health care professionals (such as doctors and nurses) to follow when they are treating patients. These clinical practice guidelines and treatment protocols are not intended to be and are not written for patients to follow for self-care and generally are not readily understandable or usable by people other than medically trained professionals.

[0010] Milliman Inc., and McKesson Corp. are two nationally recognized companies that are currently distributing clinical practice guidelines and treatment protocols for healthcare professionals (such as doctors and nurses). Every hospital in the United States must adopt one of these two sets of protocols to receive accreditation. These protocols are only for use in hospitals and only for use by healthcare professionals.

[0011] The medical literature has been discussing substantial problems with such types of known clinical practice guidelines or treatment protocols. For instance, the Aug. 10, 2005, article entitled "Clinical Practice Guidelines and Quality of Care for Older Patients with Multiple Co-morbid Diseases: Implications for Pay-for-Performances" by Darer J. Boyd CM, et al. published by JAMA provides that:

[0012] CONTEXT: Clinical practice guidelines (CPGs) [for doctors to follow] have been developed to improve the quality of health care for many chronic conditions. Pay-for-performance initiatives assess physician adherence to interventions that may reflect CPG recommendations.

- [0013] **OBJECTIVE:** To evaluate the applicability of CPGs to the care of older individuals with several comorbid diseases.
- [0014] **DATA SOURCES:** The National Health Interview Survey and a nationally representative sample of Medicare beneficiaries (to identify the most prevalent chronic diseases in this population); the National Guideline Clearinghouse (for locating evidence-based CPGs for each chronic disease).
- [0015] **STUDY SELECTION:** Of the 15 most common chronic diseases, we have selected hypertension, chronic heart failure, stable angina, atrial fibrillation, hypercholesterolemia, diabetes mellitus, osteoarthritis, chronic obstructive pulmonary disease, and osteoporosis, which are usually managed in primary care, choosing CPGs promulgated by national and international medical organizations for each.
- [0016] **DATA EXTRACTION:** Two investigators independently assessed whether each CPG addressed older patients with multiple comorbid diseases, goals of treatment, interactions between recommendations, burden to patients and caregivers, patient preferences, life expectancy, and quality of life. Differences were resolved by consensus. For a hypothetical 79-year-old woman with chronic obstructive pulmonary disease, type 2 diabetes, osteoporosis, hypertension, and osteoarthritis, we aggregated the recommendations from the relevant CPGs.
- [0017] **DATA SYNTHESIS:** Most CPG's did not modify or discuss the applicability of their recommendations for older patients with multiple comorbidities. Most also did not comment on burden, short- and long-term goals, and the quality of the underlying scientific evidence, nor give guidance for incorporating patient preferences into treatment plans. If the relevant CPGs were followed, the hypothetical patient would be prescribed 12 medications (costing her 406 dollars per month) and a complicated nonpharmacological regimen. Adverse interactions between drugs and diseases could result.
- [0018] **CONCLUSIONS:** This review suggests that adhering to current CPGs in caring for an older person with several comorbidities may have undesirable effects. Basing standards for quality of care and pay for performance on existing CPGs could lead to inappropriate judgment of the care provided to older individuals with complex comorbidities and could create perverse incentives that emphasize the wrong aspects of care for this population and diminish the quality of their care. Developing measures of the quality of the care needed by older patients with complex comorbidities is critical to improving their care.
- [0019] The Aug. 4, 2005 article, entitled "Following Clinical Practice Guidelines for Older Adults With Several Illnesses" which discusses the JAMA study further explains that:
- [0020] Current clinical practice guidelines [for doctors to follow] are not written with older adults with multiple illnesses in mind, according to a study in the August 10 issue of *JAMA*.
- [0021] The aging of the population and the increasing prevalence of chronic diseases pose challenges to the development and application of clinical practice guidelines (CPGs), according to background information in the article. In 1999, 48 percent of Medicare beneficiaries aged 65 years or older had at least 3 chronic medical conditions and 21 percent had 5 or more.
- [0022] Clinical practice guidelines are based on clinical evidence and expert consensus to help decision making about treating specific diseases. Most CPGs address single diseases in accordance with modern medicine's focus on disease and pathophysiology. However, physicians who care for older adults with multiple diseases must strike a balance between following CPGs and adjusting recommendations for individual patients' circumstances. Difficulties escalate with the number of diseases the patient has. The limitations of current single-disease CPGs may be highlighted by the growth of pay-for-performance initiatives, which reward practitioners for providing specific elements of care. Because the specific element of care may be based on single-disease CPGs, pay-for-performance may create incentives for ignoring the complexity of multiple comorbid (co-existing illnesses) chronic diseases and dissuade clinicians from providing optimal care for individuals with multiple comorbid diseases.
- [0023] Cynthia M. Boyd, M.D., M.P.H., from the Center of Aging and Health, Johns Hopkins University, Baltimore, and colleagues examined how CPGs address comorbidity in older patients and explored what happens when multiple single-disease CPGs are applied to a hypothetical 79-year-old woman with 5 common chronic diseases. Selection of these diseases were based on data from the National Health Interview Survey and a nationally representative sample of Medicare beneficiaries (to identify the most prevalent chronic diseases in this population). The National Guideline Clearinghouse was used to locate evidence based CPGs for each chronic disease. Of the 15 most common chronic diseases, the researchers focused on CPGs for hypertension, chronic heart failure, stable angina, atrial fibrillation, hypercholesterolemia, diabetes mellitus, osteoarthritis, chronic obstructive pulmonary disease, and osteoporosis.
- [0024] Two investigators independently assessed whether each CPG addressed older patients with comorbidities, goals of treatment, interactions between recommendations, burden to patients and caregivers, patient preferences, life expectancy, and quality of life. For a hypothetical 79-year-old woman with chronic obstructive pulmonary disease, type 2 diabetes, osteoporosis, hypertension, and osteoarthritis, the authors aggregated the recommendations from the relevant CPGs.
- [0025] The researchers found that most CPGs did not modify or discuss the applicability of their recommendations for older patients with multiple comorbidities. Most also did not comment on burden, short- and long-term goals, and the quality of the underlying scientific evidence, nor give guidance for incorporating patients preferences into treatment plans. If the relevant CPGs were followed, the hypothetical patient would be

prescribed 12 medications (costing her \$406 per month) and a complicated nonpharmacological regimen. Adverse interactions between drugs and diseases could result.

[0026] “For the present, widely used CPGs offer little guidance to clinicians caring for older patients with several chronic diseases. The use of CPGs as the basis for pay-for-performance initiatives that focus on specific treatments for single diseases may be particularly unsuited to the care of older individuals with multiple chronic diseases. Quality improvement and pay-for-performance initiatives within the Medicare system should be designed to improve the quality of care for older patients with multiple chronic diseases; a critical first step is research to define measures of the quality of care needed by this population, including care coordination, education, empowerment for self-management and shared decision making based on the individual circumstances of older patients,” the authors conclude. (JAMA. 2005; 294:716-724. <http://www.jamamedia.org>.)

[0027] In an accompanying editorial, Patrick J. O’Connor, M.D., M.P.H., of the HealthPartners Research Foundation, Minneapolis, commented on the JAMA study by Boyd et al. as follows:

[0028] Despite their limitations, evidence-based CPGs remain an important and necessary tool in the effort to improve health care quality. Strategies to address the limitations of current CPGs need to be developed and implemented, including providing recommendations based on level of evidence for particular patient groups and considering the potential economic and personal burden on the patient and caregiver as well as potential interactions with comorbid conditions. Future CPGs could be improved by including explicit information such as the number needed to treat to obtain a specified benefit, and should also be crafted more systematically to consider the influence of patient-specific factors such as age, life expectancy, and comorbidity on anticipated benefits of interventions. In addition, CPGs could include information on cost of various potential therapies, which may influence patient preferences and patient adherence to therapeutic regimens. Such modifications will increase the value of CPGs to clinicians and patients at the point of care, especially when physicians have too much to do [in a given office visit].

[0029] Encouraging customization of care in complex clinical scenarios respects the individuality of patients and the professional judgment of highly skilled physicians and minimizes the problem of overtreating patients most susceptible to drug interactions, drug adverse effects, and medical error. Boyd and colleagues have presented these important ‘in the trenches’ issues in a clear and compelling way. Physicians and designers of CPGs owe it to themselves and their patients to consider these issues carefully and to craft CPGs and pay-for-performance accountability measures that will reinforce excellent clinical care while being mindful of resource use and being respectful of patient preferences and priorities.

[0030] The Aug. 17, 2005, article entitled “Guidelines May Fail to Meet Needs of Elderly Patients With Comor-

bidity” by Karia Gale from Reuters Health also addresses the JAMA study in the following manner:

[0031] Current clinical practice guidelines [for doctors to follow] are designed to manage single diseases, offering little guidance to clinicians caring for older patients who have several chronic illnesses, authors of a new study suggest.

[0032] “Following clinical practice guidelines for single diseases in patients with multiple chronic conditions is very complex and costly and may lead to adverse consequences, including polypharmacy with its associated risks of adverse effects and drug interactions and even hospitalizations.” Lead investigator Dr. Cynthia M. Boyd told Reuters Health.

[0033] This is especially pertinent, she added, because pay-for-performance incentives may be based on quality of care standards created for the management of single diseases, whereas half of patients over age 65 have three or more chronic conditions. The care of these patients accounts for almost 90% of Medicare’s annual budget.

[0034] “Rewarding physicians based on what is good care for younger patients with single diseases is unrealistic,” the researcher added. “Performance incentives based on this model may penalize physicians caring thoughtfully for older patients and may impact the quality of care those patients receive.”

[0035] For their study, Dr. Boyd from Johns Hopkins University in Baltimore and her associates identified the most recently released evidence-based guidelines for hypertension, chronic heart failure, stable angina, atrial fibrillation, hypercholesterolemia, diabetes, osteoarthritis, chronic obstructive pulmonary disease (COPD), and osteoporosis.

[0036] They found that only guidelines for diabetes, chronic heart failure, angina, and hypercholesterolemia gave general guidance for patients with several comorbid conditions. None discussed the burden of comprehensive treatment on patients or caregivers, and only the guidelines for chronic heart failure explicitly discussed end-of-life treatment.

[0037] Dr. Boyd’s group used guidelines to develop a treatment plan for a hypothetical 79-year-old woman with osteoporosis, osteoarthritis, type 2 diabetes, hypertension, and COPD.

[0038] If all the recommendations were followed, the patient would require 12 separate medications taken as 19 doses at five times during a typical day. Without any insurance coverage for prescription drugs, that would amount to approximately \$400 per month. If she were a typical Medicare patient, her costs with the new Medicare drug benefit would still add up to more than \$3700 per year.

[0039] “We need to think less about individual disease and more about individual people who are living longer with multiple chronic conditions,” Dr. Boyd said. More research is needed, she added, to form “reasonable estimates of risks, benefits and burdens that are specific to them and their individual circumstances and preferences.”

[0040] Dr. Patrick J. O'Connor, from HealthPartners Research Foundation in Minneapolis, Minn., agrees with this assessment, according to his accompanying editorial.

[0041] "Ideally," he writes, "clinical practice guidelines would help physicians select from among multiple evidence-based recommendations those with the greatest potential benefit to a given patient."

[0042] This medical literature and commentary clearly explains the lack of and need for integrated treatment protocols and clinical practice guidelines for healthcare professionals (such as doctors and nurses) to follow in treating patients with multiple different diseases or different medical conditions. Thus, there is clearly a need for integrated treatment protocols and clinical practice guidelines for healthcare professionals (such as doctors and nurses).

[0043] Similarly, the existing literature does not appear to suggest that the healthcare industry prior to the present invention has considered integrated protocols for patient self-care. In February, 2005, Milliman and Robertson introduced individual treatment protocols for use by healthcare professionals outside of hospitals and for individual chronic conditions. Health plans have been buying these treatment protocols to start individual disease management programs. However, these treatment protocols for healthcare professionals to follow are not the same as self-care programs that patients (rather than healthcare professionals) must follow at home and usually alone to treat themselves. These treatment protocols would be completely unusable by a person attempting provide self-care for multiple simultaneously existing medical conditions.

[0044] Accordingly, there is a substantial need for self-care systems for treating multiple simultaneously existing different medical conditions.

SUMMARY

[0045] The present disclosure provides systems, methods and apparatus that enable a user to provide integrated self-care for blood sugar control, blood pressure control and heart failure care.

[0046] In one embodiment, the present disclosure provides an integrated blood sugar control, blood pressure control and heart failure care self care system. In one embodiment, the self-care system includes (a) a personal health folder (b) a process guide, (c) an education guide, and (d) various programs specifically designed to address these multiple health conditions of the user. In one embodiment, the process guide includes at least one record book, one or more magnet boards and a wallet card. In one embodiment, the programs included in the system include (i) a test program, (ii) a meal program, (iii) a move or exercise program, (iv) a medication program, and (v) a general health program. In one embodiment, scales, blood pressure monitor guide and blood pressure monitor are included in the system to be used in the test program. In one embodiment, the meal program includes a food guide, a food poster, a loose food guide, a fast food guide, a portion plate, a meal planner game, a meal planner and a shopping list. In one embodiment, the move or exercise program includes a skin and foot care guide and a walking guide. In one embodiment, the move program further includes skin lotion, mirror and a foot filament guide.

In one embodiment, the medication program includes a medication guide, an insulin administration guide, a pill organizer and glucose tablets. In one embodiment, the general health program includes a tobacco addiction guide and a stress management guide.

[0047] It should be appreciated that one significant benefit of the present disclosure is the increased likelihood that the patient will actually read and use this single set of materials which is substantially less volume than multiple sets of materials for each condition. When a chronic care program delivers multiple sets of materials, the sheer amount often discourages the person from trying at all. The more conditions a person has, the more likely the person will be overwhelmed. In the combination of any disease with Heart Failure, the mere name of the disease can cause a person to give up trying to manage their health. Even if they do read through everything, they will not learn the most important issue of having such multiple diseases; that is, a person who adds Hypertension to their Heart Failure is more likely to have more episodes of the heart pumping ineffectively than a person who has heart failure alone. A person who adds both Hypertension and Diabetes to Heart Failure is even more likely to suffer frequent episodes than a person who has heart failure and just one of the other diseases. None of the three sets of independent materials will alert the person to their heightened risk.

[0048] The present disclosure avoids another danger that the person will read one set and put the others away for later, often never coming back to them. For instance, if the person reads Hypertension first, they will seek to avoid salt, but the material is not likely to alert them to problems with potassium. The heart functions by alternating activities of sodium and potassium, but potassium is not a concern until a person's health has seriously deteriorated. Hypertension is such a common disease that individual instructions do not assume that the person has advanced, serious co-existing conditions. The person with hypertension and heart failure must ask their doctor whether they need a diet high or low in potassium, whether to seek or avoid potassium-containing foods, and whether to take a potassium supplement.

[0049] It is most likely, if any material is skipped or "put away for later," it will be the Diabetes materials. A significant percentage of people with diabetes are in denial about their disease and will deliberately avoid materials that are labeled "Diabetes." In this way, they miss crucial information about controlling carbohydrates in their diet.

[0050] By integrating the three sets of instructions about diet, the present disclosure enables (and to certain extent forces) the person to see the entire picture at once.

[0051] Of the three most important lifestyle changes for such chronic diseases and the most difficult to manage is diet. With three chronic diseases, three sets of instructions, the challenge can be daunting, even with sincere effort. For example, all other educational program's individual sets of instructions attempting to teach how to read Nutritional Facts labels, teach all of the lines of information. The present system teaches just the essential lines for the particular disease. If a person only has attend to a single set of essential data from the entire label, the person is more likely to sustain attention. However, if all three sets teach the entire Nutritional Facts label, it will be much more difficult to extract what is essential (i.e., a person may miss that there is

something different to learn about each one). They may see a similar section begin in a new set of materials, assume they already know what is there, and skip it. The present system teaches what is essential for all three diseases combined using a single system.

[0052] For one embodiment of the present triple-condition system or kit, the diet section can be integrated and radically simplified with a card game for learning meal planning. In one such embodiment, the suits are the food groups, each card represents a serving of a particular food, and the essential nutritional values of salt, cholesterol and carbs are the number values on the card. The goal is to plan a day's meals without going over the limits for salt, cholesterol and carbs. By playing the game, a person learns what combinations of foods will work together for a successful day's diet plan. The delight and familiarity of a card game increases the likelihood that the person will both attempt the challenge of meal planning and the likelihood that they will understand and follow the meal planning instructions.

[0053] A further benefit of the integrated approach disclosed herein is that the basic understanding of the multiple diseases is simplified and coherent; that is, both Heart Failure and Hypertension mean the heart is not pumping blood adequately through the body, but for different reasons. Education about Heart Failure typically emphasizes the excess amount of fluid. People with Hypertension can be assumed to have cholesterol sticking to arteries and clogging the pathways. Education about Diabetes alone will fail to explain the impact of Diabetes on the heart disease. Extra sugar in the blood makes everything more sticky and thick, compounding the difficulty of pumping blood throughout the body. That simple, integrated statement can be a breakthrough understanding for a patient and will not be found in three separate sets of material. Thus, the present disclosure not only provides one integrated set of materials, but also an integrated understanding of how the three conditions relate to one another. The disclosure may reconcile what may be contradictions and highlight what others may miss.

[0054] Additional features and advantages of the present system are described in, and will be apparent from, the following Detailed Description and the figures.

BRIEF DESCRIPTION OF THE DRAWINGS

[0055] FIG. 1A is a diagram illustrating the relationship between the components of an embodiment of the disclosed integrated system.

[0056] FIG. 1B is a front view of components included in an embodiment of the disclosed integrated system.

[0057] FIGS. 2A, 2B, 2C and 2D are front views of a process guide included in an embodiment of the disclosed integrated system.

[0058] FIGS. 3A, 3B, 3C, 3D, 3E, 3F, 3G and 3H are front views of an education guide included in an embodiment of the disclosed integrated system.

[0059] FIGS. 4A, 4B, 4C, 4D, 4E, 4F, 4G, 4H, 4I, 4J and 4K are front views of a record book included in an embodiment of the disclosed integrated system.

[0060] FIG. 5 is a front view of a magnetic board of one embodiment of the present disclosure.

[0061] FIGS. 6A and 6B is a front view of a wallet card of one embodiment of the present disclosure.

[0062] FIGS. 7A, 7B, 7C, 7D, 7E, 7F, 7G and 7H is a front view of a food guide included in an embodiment of the disclosed integrated system.

[0063] FIG. 8 is a front view of a food poster included in an embodiment of the disclosed integrated system.

[0064] FIG. 9 is a front view of a portion plate included in an embodiment of the disclosed integrated system.

[0065] FIGS. 10A and 10B are front views of a loose food guide included in an embodiment of the disclosed integrated system.

[0066] FIGS. 11A, 11B, 11C and 11D are front views of a fast food guide included in an embodiment of the disclosed integrated system.

[0067] FIGS. 12A, 12B and 12C are front views of a food selection game included in an embodiment of the disclosed integrated system.

[0068] FIG. 13 is a front view of a sample meal planner included in an embodiment of the disclosed integrated system.

[0069] FIGS. 14A and 14B is a front view of a shopping list included in an embodiment of the disclosed integrated system.

[0070] FIGS. 15A and 15B are front views of a walking guide included in an embodiment of the disclosed integrated system.

[0071] FIGS. 16A and 16B are front views of a skin and foot care guide included in an embodiment of the disclosed integrated system.

[0072] FIGS. 17A, 17B, 17C, 17D, 17E, 17F and 17G are front views of a medication guide included in an embodiment of the disclosed integrated system.

[0073] FIGS. 18A, 18B, 18C and 18D are front views of an insulin administration guide included in an embodiment of the disclosed integrated system.

[0074] FIGS. 19A, 19B, 19C and 19D are front views of an automatic blood pressure monitor guide included in an embodiment of the disclosed integrated system.

[0075] FIGS. 20A and 20B are front views of a tobacco addiction guide included in an embodiment of the disclosed integrated system.

[0076] FIGS. 21A and 21B are front views of a stress management guide included in an embodiment of the disclosed integrated system.

DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

[0077] The system and method described herein provide for integrated blood sugar control, blood pressure control and heart failure self-care of a single person by that person. It should also be appreciated that the system and method of the present disclosure can be employed by another person such as a care-giver, an in-home care provider, a health-care provider for the person being cared for. The disclosed method and system includes a collection of devices and

instructions a user may use to simultaneously provide blood sugar control, blood pressure control and heart failure care in an integrated manner. Many of the instructions are in the form of step-by-step guides with illustrations to guide the user through use of the system and method.

[0078] It should be appreciated that the different embodiments of the present invention may include: (a) less than all of the components described below, (b) more than the components described below, or (c) one or more substitute or alternative components for one or more of the components described below.

[0079] One embodiment of the system of the present disclosure is a system 100 of integrated components illustrated in FIGS. 1A and 1B. The illustrated integrated components system 100 includes a personal health folder 102, a primary step-by-step process or preparation guide 104 including a medical record keeping book 106, magnet boards 108 and wallet card 110, an education guide 112 discussing blood pressure, blood sugar and the heart, testing tools including scales 114, blood pressure monitor 116 and a blood pressure monitor guide 118 and secondary step-by-step program guides 120. The secondary step-by-step program guides 120 include a meal program including a food guide 124, a food poster 126, a loose food guide 128, a fast food guide 130, a portion plate 132, a meal planner game 134, meal planners 136 and shopping lists 138, a move program including a walking guide 140 and skin and foot guide 142, skin lotion 144, mirror 146 and foot filament guide 148, a medication program including a medication guide 150, an insulin administration guide 152, a pill organizer 154 and glucose tablets 156, and a general health program including a tobacco addiction guide 158 and stress management guide 160. Preferably, all of these items are packaged together in a suitable container such as a cardboard box. FIG. 1B illustrates examples of some of the components of the integrated system 100.

[0080] The self-care system 100 includes a personal health folder 102 to hold the record books, program guides and other materials components of the system 100; and a four-step preparation guide 104 illustrated in FIGS. 2A to 2D. Step one 210 of the preparation guide 104 is to know the process to control your health, step two 240 is to get the tools ready, step three 260 is to take the tests and step four 280 is to make a plan.

[0081] In step one 210, the preparation guide 104 presents the process of performing an assessment test 212 to determine the current condition of the user, making a plan 214 to improve the test score, and performing a follow-up test 216 to reassess whether the plan is working or needs to be changed. The preparation guide illustrates three monitors 220 used to test the health of the user. In the illustrated embodiment, a glucometer or blood sugar monitor, a blood pressure monitor and scales are illustrated. The monitors may or may not be included with the system 100.

[0082] Step one 210 goes on to instruct the user to read the education guide 112 included in the system. More specifically, for the purpose of educating the user about the heart, blood pressure and blood sugar, the system 100 includes an education information guide 112 on the heart, blood pressure and blood sugar illustrated in FIGS. 3A to 3H.

[0083] The education guide 112 is entitled "The Short Story on the Heart, Pressure and Sugar." As illustrated in

FIGS. 3A to 3C, the education guide 112 includes a description of the function of the heart and what constitutes heart failure 300. The education guide also includes a definition of blood pressure 310, an explanation of why blood pressure is measured with two numbers 312 and what the two numbers represent 314, an explanation of high blood pressure 320, an explanation of what makes blood pressure go up 400, and an explanation of what makes blood pressure go down 500. In FIG. 3D, the education guide 112 further provides information about the presence and function of sugar in the blood 600, and the role of insulin and other factors that affect the amount of sugar in the blood 650. In addition, in FIGS. 3E and 3F, the education guide provides an overview of the relationship between blood sugar, blood pressure and heart failure 700.

[0084] As shown in FIG. 3G, the education guide 112 provides an explanation for the different tests required in the test-plan-test process 800 described in the preparation guide 104 including the blood sugar test, blood pressure test, cholesterol, weight, and volume. The education guide 112 explains that certain tests can be performed by the user and by health care professionals. Tests such as blood sugar, blood pressure, and weight can be performed by the user and by health care professionals.

[0085] As shown in FIG. 3H, the education guide 112 further explains that other tests are typically performed by health care professionals but provides instructions on how to read those test results 900. Certain tests, such as a sugar test 910 such as hemoglobin A1C test, a fat or lipids test 920 and a volume test 930 are typically performed only by health care professionals. In addition, the education guide 112 illustrates example lab reports 914, 924 and 934 and asks the user one or more questions 912, 922 and 932 related to the reading of the lab reports 912, 922 and 932 providing answers 950 to those questions 914, 924 and 934 in the education guide 112. In particular, FIG. 3H also illustrates an example of a lab report 912 providing a hemoglobin A1C score 918. As described in the education guide 112, the hemoglobin A1C score is a measure of how much sugar has built up on hemoglobin indicating the average daily blood sugar score. To this end, a table 916 is provided setting forth the relationship between a hemoglobin A1C score and an average daily blood sugar score.

[0086] FIG. 3H also illustrates an example of a lab report 922 providing a lipid chemistry profile. The fat test 920 measures cholesterol and triglyceride levels in the blood. Referring back to FIG. 3G, in the explanation of what cholesterol tests mean 804, the education guide 114 explains that there are two types of cholesterol. A first type of cholesterol is low density cholesterol (i.e., LDL). According to the education guide 112, low density cholesterol sticks to the arteries. This type of cholesterol is unhealthy and may need to be reduced. A second type of cholesterol is high density cholesterol (i.e., HDL). The education guide 112 explains that high density cholesterol is tough bits of hard fat that does not stick to the arteries and that this type of cholesterol is healthy and may need to be increased.

[0087] As shown in FIG. 3H further illustrates an example of a lab report 932 providing an ejection fraction graph. Ejection fraction is another test performed by health care professionals to determine the volume of blood that is pumped through the heart on each beat.

[0088] The education guide 112 attempts to make the explanation of complicated physiological concepts simple enough for one without an intimate knowledge of medical science to understand the concepts. For example, in the definition of pressure 310, mentioned above, the educational guide 112 defines “pressure” as one thing pushing on another thing and “blood pressure” as blood pushing on the inside walls of the arteries. The educational guide goes on to explain in simple terms that pressure is higher during a push state and lower during a rest state. Blood pressure is typically measured during both states. Therefore, two different numbers are measured. For example, a blood pressure measurement of 120 during the push state and 80 during the rest state is typically considered a healthy blood pressure. These and other simple explanations in the system 100 enable users to understand their multiple conditions enough to motivate the user to follow the steps to help themselves. The instructions provided in the system also explain to users what they need to do without individually addressing each separate medical condition.

[0089] Turning to FIG. 3B, in the example explanation of what makes blood pressure go up 400, the education guide 112 explains that chronic high blood pressure may be caused by extra water in the bloodstream 410 (i.e., too much to pump), excess body fat 420 (i.e., too far to pump), constricted arteries 430 (i.e., too narrow to get through), and/or block arteries 440 (i.e., too clogged to get through). The education guide 112 goes on to explain that any carbohydrates not burned up by exercise are made into a form of fat known as triglycerides. The education guide also explains that animal fat is made into cholesterol by the body which can stick to the walls of arteries to clog the channel. Pictures are included to illustrate each of the concepts taught in the educational guide 114.

[0090] Turning to FIG. 3C, in the example explanation of what makes blood pressure go down 500, the education guide explains that each of the four things described that cause blood pressure to increase can be addressed in a different way to cause blood pressure to go back down. To decrease extra water in the bloodstream 510 (i.e., lighten the load—less to pump), a person may reduce salt intake and/or take medications that reduce water in the body. To decrease excess body fat 520 (i.e., shorten the trip—less distance to pump), the person may need to lose weight by watching what they eat and moving more. To open arteries 530 (i.e., widen the road—more room to get through), the person may need to stop smoking, learn to relax to reduce stress, and/or take heart medications to widen the arteries. To unblock arteries 540 (i.e., clear the strip—less clogged to get through), the person may need to eat more fiber, eat less fat, exercise, and/or take cholesterol medication.

[0091] FIG. 3D illustrates five concepts related to blood sugar. The first concept includes how sugar gets into the blood. The second concept includes how the body uses sugar. The third concept includes how insulin affects how the body uses sugar and what is meant by high blood sugar. The fourth concept includes what is meant by low blood sugar and what causes low blood sugar. The fifth concept includes what is necessary to control blood sugar.

[0092] By educating the user about basic concepts associated with heart failure, blood pressure, and blood sugar, it is believed that the user will be more likely to understand

why each of the programs (discussed below) are extremely important to follow. As a result of this understanding, the user is more likely to follow each of the steps in the single integrated programs.

[0093] Returning to step two 240 of the preparation guide 104, to further prepare to initiate the system 100, the preparation guide 104 instructs the user to refer to the record book 242, magnet boards 244 and wallet card 246, each included with the system 100.

[0094] The record book 106 is illustrated in FIGS. 4A to 4K. The record book 106 includes a cover, and sections directed to the assessment tests 212, the plan 214 and the follow-up tests 216 according to the process described in step one 210 of the preparation guide 104 illustrated in FIG. 2A. In particular, as illustrated in FIGS. 4B and 4C, the test section 1100 of the record book 106 includes a place to record answers to recommended questions 1102 to be asked of the doctor on each clinic visit and a place to record blood test results 1120. The plan section 1200 of the record book 106 includes a place to record information about meals 1202, moves or exercise 1203, and medications 1204. The follow-up test section 1300 includes instructions on how to keep the medical records 1400, and a plurality of weekly record pages 1402.

[0095] The assessment tests section 1100 includes questions to ask the doctor or health care professional on each clinic visit and tests the user should expect the clinic to perform on each clinic visit. FIG. 4B illustrates a place to record answers to questions for the doctor 1102. The record book 106 includes questions about blood sugar 1104, weight 1106 and blood pressure 1108 and corresponding blanks 1110 to record the answer from the health care professional from each clinic visit. For example, the questions about blood sugar 1104 the user should ask of the health care provider include when blood sugar should be tested, what the blood sugar level should be including high and low limits of the blood sugar level 1112, what the high and low blood sugar alarm limits 1114 should be beyond which the user should call the doctor, how often the blood sugar should be checked when sick, and whether a prescription for ketone strips is needed. Questions about weight 1106 include what is a healthy weight for the user and how much weight can be gained in a week before the user would be required to notify the doctor. Questions about blood pressure 1108 include how often to take a blood pressure reading, what the blood pressure top and bottom numbers should be, at what top and bottom numbers blood pressure alarm limits should the user call the doctor, and what a pulse reading should be.

[0096] FIG. 4C illustrates a table of tests 1120. The system 100 recommends that the user remind the health care provider to perform the tests in the table 1120 on a periodic basis. Tests that are recommended to be performed on each clinic visit are weight, blood pressure, and skin and feet check. Tests recommended to be performed two times per year include a hemoglobin A1C level, and a lipid profile including total cholesterol, good cholesterol (i.e., HDL), bad cholesterol (i.e., LDL), and triglycerides. A test recommended to be performed once a year includes a microalbumin level. As illustrated in FIG. 4C, a place to record blood test results 2104 is also provided. For each of the blood test types, the record keeping book 106 includes a target score (e.g., less than 200 for total cholesterol). Questions about the

next appointment **1126** may also be included such as the date and/or time of the next appointment.

[0097] The plan section **1200** of the record book **106** is illustrated in FIGS. **4D** to **4F**. The plan section **1200** includes a portion directed to questions to ask the doctor or health care provider **1202** and **1203** illustrated in FIG. **4D**, a meds portion **1204** illustrated in FIG. **4E** and a meal portion **1206** illustrated in FIG. **4F**.

[0098] The first part of the plan section **1202** include questions to be asked of the health care provider about food, exercise, vaccinations, foot care, as well as referrals and frequency of visits to foot doctors, eye doctors and dentists. Questions about food may include a question on how much cholesterol and salt should be consumed each day and whether potassium is an acceptable salt substitute. Questions about exercise may include what type of exercises the user should be doing, how long the user should be exercising each time, and/or how often the user should be exercising.

[0099] FIG. **4E** illustrates the med portion **1204** of the plan section **1200** of the record book **106** where a user can record information about medications. For each of the medications, the record keeping book **106** includes a blank for a name of the medication **1206**, a dosage or strength of the medication **1208**, the amount of medication or number of pills to take **1210**, and when to take the medication **1212** (e.g., breakfast, lunch, dinner, bedtime). Preferably, the medications are grouped by the function they perform. In this example, the groups include pills for the heart **1214**, pills for blood sugar **1216**, other prescription meds **1218**. The meds portion also includes a table to record non-prescription supplements **1220**, such as vitamins and herbs, taken by the user. The table provides space for the user to record the kind of supplement **1222**, name of each supplement **1224**, how much of each supplement is taken **1226** and when each supplement is taken **1228**.

[0100] In addition, the meds portion of the record book includes a table to record information about insulin administration. A table is provided for "every day" insulin **1230** and for "it depends" insulin **1240**. The table for "every day" insulin **1230** includes a column, each for entries as to start dates **1232**, what kinds of insulin is being taken **1234**, how much (in units) to take at one time **1236** and when to take the insulin **1238**. A separate table is provided for the "it depends" insulin **1240**. "It depends" insulin may be a different insulin the doctor may want the user to take if the blood sugar test score is too high. The table **1240** provides a place to enter a blood sugar test score threshold **1242** and the date that threshold was provided by the health care provider **1244a** and **1244b**. The table **1240** also provides a place to enter the kind of insulin **1246** and how much (in units) of that kind of insulin should be taken by the user **1248** if the blood sugar test score of the user is more than that threshold score. Additionally, the record book provides a section **1250** to record where the doctor has instructed the user to administer the insulin.

[0101] The plan section **1200** of the record book **106** further includes a meal plan section **1206** illustrated in FIG. **4F**. The meal plan section **1206** provides the user a table to record the recommended number of servings or number of carbohydrates for each food group and total for breakfast, lunch, dinner and snack(s) recommended by the health care provider. For example, the food groups may include protein, fat, fruit, vegetables, grain and starch, and milk.

[0102] The follow-up test section **1300** of the record book **106** illustrated in FIGS. **4G** to **4I** includes a section **1302** illustrated in FIG. **4G** that enables a user to focus on a one week test period to determine whether the blood sugar is "in control" or "on alert." In particular, the record book provides a table **1304** with spaces enabling a user to track medications **1306** and meals **1308** and their effect on blood sugar test scores **1310** on a daily basis through a period of one week. The user is instructed to record in a space provided in the record book what medications the user took and what food the user ate for breakfast, lunch, dinner and any snacks for each day of the first test week.

[0103] In addition, the user is instructed **1311** to record blood sugar test scores **1310** corresponding to each day of the test week. The blood sugar test scores are recorded in a spatial relationship to the blood sugar limits **1112** provided by the user's health care provider in the assessment tests section **1110** of the record book **106** illustrated in FIG. **4B**. For example, the low blood sugar limit **1312a** is indicated by line **1312b**, and the high blood sugar limit **1314a** is indicated by line **1314b**. A blood sugar score above the high limit **1314a** is recorded above line **1314b** in space **1316a**. A blood sugar score between the high limit **1314a** and the low limit **1312a** is recorded between lines **1314b** and **1312b** in space **1316b**. A blood sugar score below the low limit **1312a** is recorded below line **1312b** in space **1316c**.

[0104] FIG. **4H** illustrates instructions on how to keep the medical records **1400** along with pictures of the section corresponding to the instructions. FIG. **4I** illustrates one example of a plurality of weekly record pages **1402**. The table includes columns for each day of a week and is divided into two sections: the "How I tested" section **1404** and the "What I did" section **1406**.

[0105] As illustrated in FIG. **4H**, the first step **1410** discussed in the record book **106** is to prepare for the tests each week by recording in the "How I tested" section **1404** the date **1408** and the numbers for the high **1312** and low **1314** blood sugar limits **1112** provided to the user by the health care provider. The second step **1420** is to record the results of the tests in the "How I tested" section **1404**.

[0106] The weekly record illustrated in FIG. **4I** includes a row of spaces for each test where the user may enter the test results for each day of the week. For example, the test result on Sunday for weight can be entered in space **1410**; the test result on Sunday for blood pressure can be entered in spaces **1412a** and **1412b**; the location where the user obtained the specimen can be entered in space **1413**; and the test result on Sunday for blood sugar can be entered in space **1414**. Again, the user is instructed to record blood sugar test scores corresponding to each day and in spatial relation along a vertical axis to the high and low blood sugar limits provided by the health care provider as described above. In addition, the test result on Sunday for a pulse reading can be entered in space **1416**.

[0107] As illustrated in FIG. **4H**, the third step **1430** is to track the plan in the "What I did" section **1406** of the medical record tables illustrated in FIG. **4I**, which includes spaces to enter information about the user's meals **1440**, moves or exercise **1450**, and medications **1460**. In FIG. **4H**, for meals, the user is instructed to record information at three levels. The first level **1442** asks the user to indicate if the food poster described below included in the system was

followed and what the user ate from the list of foods recommended to be avoided. The second level **1444** asks the user to indicate if the user controlled serving size according to the recommendations of the system and what the user ate that went over the serving size limit. The third level **1446** asks the user to indicate if the user followed the meal plan and to record what and how much the user ate that was not on the meal plan.

[**0108**] In the moves or exercise section **1450**, the user is instructed to record what kind of exercise **1452**, how long each exercise was performed **1454** and how many steps until the user is short of breath **1456**. The type of exercise, the amount of exercise and the number of steps until being short of breath may be recorded each day on the weekly medical record **1400**. On a Sunday, for example, this information can be recorded in blanks **1462**, **1464**, and **1466**, respectively. By recording the type and amount of exercise performed on various days of the week, the user is able to track progress from week to week.

[**0109**] In the med section **1460** of the “What I Did Section” **1406**, the user is instructed to record in the weekly medical record **1402** illustrated in FIG. 4I, any variation in taking the medications the user has recorded in the meds section **1204** of the record book **106** illustrated in FIG. 4E and described above. Variations can include taking a medication that is not listed in the meds section **1204**, taking a medication at a different time or not at all, or any other deviation from the plan.

[**0110**] As illustrated in FIG. 4I, the user is prompted to record any occurrence of an annoying cough each day on the weekly medical record **1400**, and, on a Sunday for example, in blank **1472**. The user is also prompted to record the number of cigarettes he/she smokes each day (if any) on the weekly medical record **1400**, and, on a Sunday for example, in blank **1474**. By recording the number of cigarettes smoked each day, the user is able to track progress from week to week. The user is further prompted to record whether their shoes or belt feel tighter than usual **1376**. This may be recorded each day on the weekly medical record **1400**, and, on a Sunday for example, in blank **1476**.

[**0111**] Referring to FIG. 4J, the record book **106** includes a description of how a user can test his or her feet **1490**. The instructions refer to a filament guide **146** which may or may not be provided with the system **100** to contact different areas of the bottom of the feet to determine if the user is able to detect the contact.

[**0112**] An example of a filament guide is illustrated in FIG. 14K. The filament guide may include a filament **1492** having a stiffness that enables the filament to stand erect at a length of one to two inches. The system **100** can include any suitable filament or probe with any suitable stiffness. The filament may be attached to a handle **1494** which provides a diagram illustrating areas on the feet to be tested with the filament **3872**. The record book **106** also provides diagrams **1496** illustrating areas **1498** on the feet to be tested.

[**0113**] Referring back to FIG. 4J, the instructions instruct the user to touch the filament to an area of the foot indicated by a circle pushing hard enough for the filament to bend. The filament is held in place against the foot for approximately two seconds. If the user cannot feel the filament, the instruc-

tions instruct the user to place an “x” in the circle on the diagram **1498** in the record book **106**. The instructions **1490** further instruct the user to repeat this method for each circle on both feet and to call the doctor if an “x” is indicated in any of the circles. In addition, the instructions include a list of months for the user to indicate when a foot test has been performed for that month.

[**0114**] Returning to the second step **240** of the preparation guide **104** of the system **100**, in addition to the record book **106**, the second step **240** provides instructions **244** and **246** for a magnetic board **108** and a wallet card **110** illustrated in FIGS. 5 and 6A and 6B, respectively. The magnetic board **108** may be hung from a refrigerator as a convenient reference for information. In the illustrated embodiment, the information included on the magnet boards is divided into three sections. The first section **2010** includes a list of, and space to add, what needs to be done on a daily, weekly and monthly basis. The second section **2020** of the magnetic board **108** includes an emergency action plan tailored to respond to symptoms associated with abnormal blood sugar levels. The emergency action plan includes alarm numbers **1114** for high **2030** and low **2032** blood sugar levels provided by the health care provider and recorded in the assessment tests section **1100** of the record book **106** illustrated in FIG. 4B, along with instructions **2030** describing what to do if the symptoms associated with abnormal blood sugar levels occur. The emergency action plan also includes prescription refill telephone numbers **2040** along with an emergency telephone number **2050** for the blood sugar doctor.

[**0115**] A third section of the magnet board **108** includes a place to record the telephone number of the heart doctor **2110** and a list of events **2120** related to the health of the user. If one or more of the events **2120** occur, the user is instructed to call the heart doctor or call emergency medical services. The events may include a change for the worse, weight gain in excess of certain threshold limits in a day or a week, use of more pillows to sleep, worsening cough, swelling legs, difficulty breathing, blood pressure above or below certain threshold limits, feeling faint or dizzy with normal blood sugar, experiencing chest pains. The magnet board **108** also includes an area to note questions to ask the healthcare provider **2130** and a place to record information concerning the next appointment **2140**.

[**0116**] Referring again to FIG. 2B, another tool described in the second step **240** of the preparation guide **104** of the system **100** includes a wallet card **110** illustrated in FIGS. 6A and 6B. The wallet card **110** may include a front and back side with specific instructions **2150** and emergency medical information **2160**. The instructions may include what to do when the user is not feeling well, what to do **2170** if the user's blood sugar level exceeds certain specified limits including how to manage high or low blood sugar and when to call a doctor. The emergency medical information **2160** may include the identification and phone number of the user and names and phone numbers of doctors such as the heart doctor and blood sugar doctor.

[**0117**] As illustrated in FIG. 2C, step three **260** of the preparation guide **104** of the system **100** is to “Take the Tests.” In the illustrated embodiment, three tests are listed and described: the weight test **262**, the blood pressure test **264** and the blood sugar test **266**.

[0118] The weight test instructions 262 instructs the user to weigh himself/herself each morning before breakfast, and after using the toilet. The instructions 262 indicate that the user should place a scale on a hard floor (not carpeting) and to remove any clothing before weighing. A suitable set of scales 114 may be provided with the system 100. The weight is then recorded in a blank 1410 corresponding to the current day on the weekly record page 1402 as described above. The instructions 262 also tell the user to watch this recorded weight to make sure that the weight does not change (e.g., go up over time).

[0119] The blood pressure instructions 264 instruct the user to take his/her blood pressure every day and to record the blood pressure in the record book 106. A suitable blood pressure monitor 116 may be provided with the system 100. The blood pressure may be recorded each day in two blanks 1412a and 1412b on the weekly record page 1402 as described above and illustrated in FIG. 4I. The first blank 1412a is for recording blood pressure during the heart's push state, and the second blank 1412b is for recording blood pressure during the heart's rest state. An additional space 1416 is provided for the user to enter a pulse reading each day. A digital blood pressure monitor 116 is included in the system 100 for measuring blood pressure, and instructions 118 for using the blood pressure monitor 116 are described below.

[0120] The blood sugar instructions 266 instruct the user to obtain a blood sugar monitor which may or may not be included in the system 100. In addition, the user is instructed to obtain a lancet, a test strip, a clean towel, soap, a writing instrument such as a pen and the record book. The preparation guide 104 further instructs the user to wash and dry his or her hands with warm water and soap, and not alcohol, at a clean place near a sink. The user is then instructed to stick the side of a finger and to follow the monitor instructions to obtain a blood sugar measurement. The user is then instructed to record in the space provided 1413 and illustrated in FIG. 4I where the user obtained a sample and the blood sugar reading. The blood sugar is recorded in spatial relation 1414 to the high and low blood sugar limits provided by the health care provider in the weekly record 1402 in the record book 106 illustrated in FIG. 4B.

[0121] FIG. 2D illustrates the fourth step 280 of the preparation guide 104. The fourth step includes making a plan. As indicated in step one 210 of the system 100, and in the record book 106, the plan includes three parts: "watch the meals," "make the moves," and "take the meds." The Personal Health Folder includes a section for each of these parts where the user may formulate a plan. According to the preparation guide 104, the plan will enable the user to achieve the five goals discussed in the short story described above. The five goals include: to lighten the load 282 by eating less salt and taking water pills, to shorten the trip 284 by watching what you eat and exercising more, to widen the road 286 by stopping smoking, relaxing and taking heart pills, to clear the strip 288 by eating more fiber and less fat and taking cholesterol pills, and to keep it steady 290 by doing regular amounts of everything on time.

[0122] Once the user has created a plan, step four 280 of the preparation guide 104 describes how the record book 106 enables the user to keep track of what the user does to follow the plan in the "What I did" section 1406 of the

record book 106 and how the plan is working in the "How I tested" 1404 section of the record book 106 illustrated in FIG. 4I. The user is also instructed in the preparation guide 104 to take the record book 106 to every clinic visit to enable the doctor to review the information with the patient to determine if the plan is working and make revisions if necessary. The preparation guide 104 further reminds the user to obtain from the health care provider the updated test scores and any changes to limit numbers, diet, exercise or medication.

[0123] FIGS. 7A to 14B illustrate a meal program included in the system 100. The meal program includes a food guide 124, a food poster 126, a loose food guide 128, a fast food guide 130, a portion plate 132, a meal planner game 134, a meal planner 136, and a shopping list 138.

[0124] The meal program of the system 100 includes a food guide 124 illustrated at FIGS. 7A to 7G. The food guide 124 instructs the user of the system 100 how to watch what the user eats and, specifically, how to reduce salt, cholesterol and carbs in the diet. The food guide includes three levels: the poster 3100, the plate 3200, and the meal plan 3300. In the first level 3100 of the meal program, the user is taught how to eat less of salt, cholesterol and carbohydrates using a poster 126 that illustrates different foods categorized by their content. The second level 3200 of the meal program instructs or teaches the user how to control serving size using a portion plate 132 to illustrate the proper serving size for typical foods. The third level 3300 of the food guide teaches the user how to plan a menu for the day and to count how much salt, fat and carbohydrates are in foods eaten by the user.

[0125] As described in level one 3100 of the food guide 124, a food poster 126 is included in the system 100 and is illustrated in FIG. 8. Referring to FIG. 8, the food poster 126 includes illustrations of food items. The food items are categorized by whether to eat more or less of the food. The food items are also categorized by what they contain or do not contain. The illustrated food poster is divided into a red section of foods 3110 and a green section of foods 3112 as an alternative to the foods in the red section 3110. The user is further instructed to eat more foods from the green side 3112 of the poster and eat less foods from the red side 3110 of the poster. The red or "eat less" 3110 section and the green or "eat more" 3112 section of the poster 126 are divided into three sections. The three sections include "watch out for salt" 3114, watch out for cholesterol" 3116 and "watch out for carbs" 3118.

[0126] In the "watch out for salt" section 3114, examples of food items containing salt in the red section 3110 and alternatives to those foods in the green section 3112 are illustrated. Examples of food products such as salty snacks, food in cans and salt shakers are illustrated as foods to eat less. Examples of food products such as snacks with no salt, canned food with no salt or frozen vegetables, and spices with no salt are illustrated as foods to eat more.

[0127] In the "watch out for cholesterol" section 3116, examples of food items containing cholesterol in the red section 3110 and alternatives to those foods in the green section 3112 are illustrated. Examples of red meat and egg yolks, and foods with fat from animals are illustrated as foods to eat less. Examples of fish chicken, pork and egg whites, and foods with fats that come from plants are illustrated as foods to eat more.

[0128] In the “watch out for carbs” section 3118, examples of food items containing carbs in the red section 3110 and alternatives to those foods in the green section 3112 are illustrated. Examples of sugar, white flour, heavy vegetables and heavy fruits are illustrated as foods to eat less. Examples of sugar-free foods, brown flour, light vegetables and light fruits are illustrated as foods to eat more. The user of the meal program is instructed by the food guide 124 to put up the poster 126 in the kitchen.

[0129] Referring again to the food guide 124 and FIG. 7A, the poster is explained or supplemented by instructions in the food guide to eat less salt 3120. The user is instructed to use less salt in cooking 3122, less salt from a shaker 3124, and less salt in the cupboards and refrigerator 3126. To use less salt in cooking, the food guide recommends using new flavors and other spices and sauces, such as lemon and other juices. In addition, the instructions recommend that the user remove salty food from the cupboards and refrigerator, such as bouillon, ketchup, cheese, chili sauce, cold cuts, frozen dinners, mustard, olives, pickles, salad dressing, sausages and soy sauce. The instructions also teach the user to get new flavors such as garlic, lemon and limes, basil, cilantro and onion into the home. As described above, the user may refer to the “About Food” section 1202 of the plan section 1200 of the record book 106 illustrated in FIG. 4D to determine how much salt a user is limited to in a single day and whether a potassium salt substitute is acceptable with a doctor’s approval.

[0130] As illustrated in FIG. 7A, the food guide 124 discusses eating foods with potassium 3130. The food guide 124 provides a list of foods with high potassium 3132 for the user if the user has been advised by the doctor to eat foods with more potassium. Examples of food with high potassium include: apricots, avocado, banana, cantaloupe, honeydew, kiwi, lima beans, meat, poultry, fish, milk, oranges, juice, potatoes, prunes, spinach, tomatoes and winter squash. The instructions also provide a list of foods with low potassium 3134 for the user if the user has been advised by the doctor to eat foods with less potassium. Examples of food with low potassium include: apples, bell peppers, blueberries, cabbage, cranberries, cucumbers, fruit cocktail, grapes, green beans, iceberg lettuce, mushrooms, canned peaches, fresh pineapple and plums. In some cases, the user has been advised by the health provider to use fake salt with potassium. The food guide 124 illustrates where a user may look on a product to determine if the product contains potassium salt 3136.

[0131] As illustrated in FIG. 7B, the second level 3200 of the food guide 124 includes the plate 132. This section 3200 of the food guide 124 describes proper serving sizes for different types of food and a proper number of servings per day. The instructions 3202 discuss the difference between a “helping” and a “serving.” According to the instructions 3202, a helping includes a scoop of food of any size. In contrast, a serving or portion includes a scoop of food a certain size.

[0132] The food guide 124 employs a portion plate 132 illustrated at FIG. 9 to teach proper serving sizes for different types of food. Referring to FIG. 9, on the portion plate 132 are pictured playing pieces from games and sports providing the user a reference to different amounts included in a serving size. For example, a hockey puck 3210 represents

about ½ cup; a combo lock 3220 represents about ⅓ cup; a baseball 3230 represents about 1 cup; a golf ball 3240 represents about a ¼ cup; a deck of cards 3250 represents about 3 ounces; and four stacked checkers 3260 represent about 1 tablespoon. The portion plate 132 also indicates typical foods that should be consumed by the user in the amount specified on the portion plate 132. For example, the instructions indicate that a proper serving size for cooked oatmeal, applesauce, peas, corn, chili or sweet potatoes is ½ cup, or the size of a hockey puck 3210 illustrated on the portion plate 132. The proper serving size for baked beans, cooked rice, cooked noodles, bran cereal or turkey stuffing is ⅓ a cup, or the size of the combo lock 3220 illustrated on the portion plate 132. The proper serving size for milk, melon, plain yogurt or blueberries can be 1 cup, or the size of the baseball 3230 illustrated on the portion plate 132. The proper serving size for granola or nuts and seeds can be ¼ cup, or the size of the golf ball illustrated on the portion plate 132. The proper serving size for meat, fish, chicken or turkey can be 3 ounces, or the size of a deck of cards 3250 illustrated on the portion plate 132. A proper serving size for margarine, syrup, honey, jam, oils or fat free dressing can be 1 tablespoon, or the size of four checkers 3260 illustrated on the portion plate 132.

[0133] Referring to FIG. 7C, after discussing serving size, the food guide discusses how many servings a user should have per day 3270. The instructions 3270 indicate that the number of servings may vary between different people. The instructions 3270 further illustrate different variations in serving sizes based on the type of food. The instructions list typical numbers of servings for protein 3272, vegetable fat 3274, vegetables 3276, milk and yogurt 3278, grains and starch 3280, and fruits 3282. For protein 3272, the instructions indicate that most people need three to four servings of meat, fish, or eggs a day, or one serving each meal. For vegetable fat 3274, the guide indicates that most people need three to six servings a day of oil or margarine, or one or two servings per meal. For vegetables 3276, the guide indicates that most people need two to three servings of vegetables a day, or one serving for each meal. For milk and yogurt 3278, the guide indicates that most people need two to three servings of milk or yogurt a day, or one serving for each meal. For grains and starch 3280, the guide indicates that most people need five to six serving of grains and starch a day, or two servings per meal. For fruits 3282, the guide indicates that most people need two to three servings of fruit a day, or one serving per meal.

[0134] As illustrated in FIG. 7D, the third level 3300 of the meal program described in the food guide 124 includes the plan. The food guide 124 teaches the user how to track the amount of salt, fat and carbohydrates the user consumes. In particular, the instructions provide a description of various counting tools 3310 the user of the program can use to count how much salt, cholesterol or carbohydrates are in certain foods.

[0135] The counting tool instructions 3310 illustrated in FIG. 7E include instructions for using the nutrition facts labels on a food package 3320, instructions 3330 for using the loose food guide 128 and the instructions for using the fast food guide 130. The nutrition facts label included on most food packages lists the amount of cholesterol, sodium or salt and carbohydrates or total carbohydrates.

[0136] For those foods that may not have nutritional facts labeling, the system 100 includes a loose food guide 128 and a fast food guide 130 illustrated in FIGS. 10 and 11, respectively. As illustrated in FIGS. 10A and 10B, the loose food guide 128 enables the user to determine nutritional facts for foods typically purchased without packaging. The loose food guide 128 lists the calories, sodium, cholesterol, and carbohydrate content per serving, portion or exchange of grains and starches, proteins, fats and nuts, fruit, milk and yogurt, light and heavy vegetables, sweets and any other food that is not contained in a package. FIG. 10B illustrates an example of a table 3340 listing these amounts for grains and starches 3341, and proteins 3342.

[0137] Similarly, the fast food guide 130 included in the system 100 enables the user to determine the amount of calories, carbohydrates, fiber, protein, fat, percent of calories from fat, saturated fat, cholesterol, and sodium in a typical serving size, a real serving or exchange size of food items available at restaurants or fast food establishments. For example, FIG. 11B illustrates a table 3344 listing the nutritional amounts for food items at Krispy Kreme® 3345 and McDonalds® 3346. As illustrated in FIG. 11C, the fast food guide 130 also includes instructions on maintaining a healthy diet 3347 and a conversion table 3348 that enables the user to determine recommended fat intake for different calorie levels. In addition, the fast food guide 130 includes a listing 3349 of sample menu items from various fast food establishments that fall within certain nutritional parameters, as illustrated in FIG. 11D.

[0138] Referring back to the food guide 124, as illustrated in FIG. 7F, the food guide also includes examples of how to count salt 3350, cholesterol 3360 and carbohydrates 3370. The food guide instructions also include typical limits per day, per meal, and per serving for salt, cholesterol and carbohydrates. For example, according to the food guide the salt limit for most people is 1500 mg per day or 500 mg per meal. The counting salt instructions 3350 point out that, since there are approximately five servings per meal, each serving has a limit of 100 mg of salt. The instructions 3350 also point out that different brands of the same food product can often have different salt amounts. According to the food guide, the cholesterol limit for most people is 300 mg per day or 100 mg per meal. If a person has two servings of food with cholesterol in a meal, the limit for each serving is approximately 50 mg of cholesterol. According to the food guide, the carbohydrate limit for most people is approximately 150 grams per day or 50 grams per meal. If the user has three servings of food that include carbohydrates per meal, the average number of carbohydrates per serving is 16. The instructions 3370 do not ask the user to count carbohydrates for “free” fruits and vegetables. The food guide also provides and illustrates examples of the number of carbohydrates associated with proper serving sizes of different foods.

[0139] As illustrated in FIG. 7G, the food guide 124 of the meal program provides a user an opportunity to practice counting carbohydrates in various food products. The food guide illustrates packaged 3382, loose food 3384 and fast food 3386 products and provides the number of carbohydrates in each of these products 3388. In addition, the meal program provides the user an opportunity to determine the amount of salt cholesterol and carbohydrates in an example meal 3390.

[0140] In one embodiment, the meal program includes a game, referred to in the food guide 124, to teach a user how to plan meals within the limits for the amount of salt, cholesterol and carbohydrates in each of those meals. In particular, FIGS. 12A to 12C illustrate a food selection or meal planner game 134 included in the meal program. The food selection game 134 is entitled “The Real Meal Deal” and enables a player to learn about salt, cholesterol and carbohydrate content in foods for planning meals. FIG. 12A illustrates instructions for the game. The game includes a plurality of cards. Examples of the cards are illustrated in FIG. 12B. Each card 3410 indicates a particular food or food product 3420 in a food group 3430 such as protein, fruit, sweets, milk & yogurt, vegetables fat, sauces and spices, heavy vegetables and grains and starch. In addition, the card includes nutrition facts 3440 of the food product 3420, and the serving size 3450 of the food product 3420. The nutrition facts are customized for the conditions being addressed by the system and include how much salt, cholesterol and carbohydrates are contained in one serving of the food product. For example, card 3430, illustrated in FIG. 12B indicates strawberries in the fruit food group. The nutrition facts 3440 include a salt content 3442 of 1, a cholesterol content 3444 of 0 and a carbohydrate content 3446 of 10 included in a serving size 3450 of one-half cup. FIG. 12B illustrates other examples of cards in the food selection game.

[0141] To begin the game, a dealer shuffles and deals to each player the same number of cards, such as ten cards. The remaining non-dealt cards are placed in a “Draw Pile” face down. Each player attempts to select from the ten cards indicating food products to be included in a meal or a snack. The goal of the game is to plan three meals without exceeding certain nutrition limits. In the illustrated embodiment, each meal must include 40 carbohydrates. The meals for each day may include at least two servings from at least one food group and contain no more than 1500 mg of salt, no more than 300 mg of cholesterol, and no more than 150 g of carbohydrates corresponding to the recommended daily limits described in the food guide and illustrated in FIG. 7F.

[0142] During play of the game, players take turns presenting cards for a day’s meals and snacks which match the nutrition goals. The players are permitted to move cards from one meal to another meal but may not pick them up again. Players may also trade cards with other players if each agrees to the trade. As illustrated in FIG. 12B, a wild card 3460 may also be included in the game. The wild card 3460 indicates that the player may “copy any card in your hand” and “count the points a second time.” Therefore, the wild card enables a player to choose any card in his hand and count the numbers associated with salt, cholesterol and carbohydrate content a second time to meet the desired limits to be achieved in the game. Tally sheets 3470 illustrated in FIG. 12C may be provided to enable the players to keep track of the nutritional goals throughout the game.

[0143] A turn ends when the player places any unwanted cards in a “Discard Pile” and draws enough cards from the “Draw Pile” to replace the unwanted cards and to return the number of cards in the players hand to ten cards. When there are no more cards in the “Draw Pile,” the cards from the “Discard Pile” are then used. The first player who lays down the most cards to plan three meals without exceeding the nutritional limits is the winner. Once a winner has been

declared, the game ends. Alternatively, play may continue to enable other players to attempt to plan three meals within the nutritional limits.

[0144] Once the user has had an opportunity to use the counting tools **3310** described in the food guide **124** and practiced planning meal with foods having less than the recommended limits for salt, cholesterol and carbohydrates, the user may use the meal planner **136** included in the system **100** to prepare actual meal plans. Referring to FIG. 7H, the food guide provides instruction **3600** in preparing meal plans that conform to the limits of serving size, number of servings and the amount of salt, cholesterol and carbohydrates in each of those meals. The food guide refers the user to the record book to assist the user in determining the limits recommended by the health care provider that have been recorded in the record book.

[0145] To assist the user in planning meals within these nutritional limits, the system includes a sample meal planner **136** illustrated in FIG. 13. The sample meal planner **136** lists for each meal the food groups, serving sizes for each of the food groups, how many servings in each of the food groups, how many carbohydrates in each of the food groups, how much salt in each of the food groups and how much cholesterol in each of the food groups. The food groups include: protein, fat, fruit, vegetables, grain and starch, milk, spices and sweets. In a meal planner without the example food information entered provided with the system **100**, the meal planner **136** includes space to record numbers of servings **3616**, total amounts of carbohydrates **3618** and amounts of salt **3620** and cholesterol **3622** from each of the foods and food groups represented in each meal. The meals include breakfast, lunch, dinner and snacks. Each of the breakfast totals **3630**, lunch totals **3640**, dinner totals **3650** and snack totals **3660** are added to determine the total for each of these parameters for the day **3670**. Before the user completes the meal planner, the food guide instructs the user to copy the limit number for cholesterol and salt that have been entered into the plan section **1202** of the record book illustrated in FIG. 4D and described above. The limit numbers are recorded in the appropriate spaces on the meal planner. The user is instructed to plan meals within the proper limits based on the nutritional information on food product packaging and the information provided in the loose food guide **128**.

[0146] The meal program also includes an example shopping list **138** and meal plans illustrated in FIGS. 14A and 14B. The shopping list **138** provides a list of alternative foods to those with salt **3710** and fat **3720** for easy reference when shopping in a grocery store. The shopping list also includes a listing of free vegetables **3730** and fruits **3740**. The shopping list further includes an abbreviated meal planner **3760** with space to list foods in each food group for each meal including breakfast, lunch, dinner and snacks.

[0147] Referring back to FIG. 7H, the food guide **124** illustrates instructions on how to survive a night eating out **3790**. For example, instead of regular salad dressing on a salad, the instructions **3790** suggest low fat salad dressing on the side or lemon juice squeezed from a lemon slice. In addition, the instructions suggest that the user cut the served portion in half and put half of the meal away before eating the meal. The instructions **3790** further advise the user that soft-drink consumption may cause blood sugar to increase

above a normal level and that alcohol consumption may decrease blood sugar below a normal level.

[0148] The system **100** includes a move or exercise program. The exercise program includes a walking guide **140**, a skin and foot care guide **142**, skin lotion **144**, mirror **146** and foot filament guide **148**.

[0149] FIGS. 15A, 15B and 15C illustrate a secondary step-by-step walking guide **140**. The secondary step-by-step walking guide **140** includes instructions on why people should walk **3802**, items people need to have to be a regular walker **3810**, how to make walking a habit **3820** and instructions for travel **3850**.

[0150] FIG. 15A illustrates instructions on why people should walk **3802**. The instructions **3802** inform the user that walking is good exercise that almost anyone can do. The instructions **3802** indicate that walking is good exercise because walking gets the blood flowing in the feet and legs. Walking also helps reduce high blood pressure. After several months of regular walking, the user can expect to have more energy and sleep better.

[0151] FIG. 15A also illustrates instructions on things people need to have to be a regular walker **3810**. The instructions **3810** indicate that the user should have walking shoes, a regular walking time, a watch, water, and guts. The walking shoes should be the right size, comfortable, and made of canvas or soft leather. The user should walk at a regular walking time seven days per week and be content with sticking to the walking routine at least five times per week. The instructions also advise the user to use a watch and/or pedometer to measure time and distance or number of steps of the walk. The instructions further suggest taking a water bottle and to drink often. In addition the user is instructed to prepare a bag of emergency tools including the emergency wallet card **110** and what is necessary to treat any medical condition such as glucose tablets **56** or snacks to treat low blood sugar. The instructions **3810** also indicate that the user may want to find a friend to walk with. In this manner, the user may be more likely to stick to the walking commitment. Other suggestions included in the sample instructions **3810** include walking at a mall, not over-doing the walking routine, and walking at a pace where talking is still comfortable.

[0152] As illustrated in FIG. 15B, the "Make Walking a Habit **3820** section of the walking guide **140** presents instructions **3821** for using walking plans **3822** for different levels of walkers. In particular, the section includes three six-week programs for a "beginner" level **3824**, a "mover" level **3826** and an "expert" level **3828**. For each level and for each week, the plans **3822**, include increments of relative speeds **3830**, such as slow, slower, fast or faster, and the duration at each speed **3832**. The plans **3822** also include at least two increments to walk in one direction and two increments to turn back and walk the opposite direction. For example, at the beginner level **3824** every day of the first week, the user is instructed to walk slow for one minute and fast for one minute. The user is then instructed to turn back and walk fast one minute and slow one minute for a total time of four minutes. At the expert level **3828** every day of sixth week, the user is instructed to walk slow for five minutes and fast for sixteen minutes. The user is then instructed to turn back and walk fast for sixteen minutes and slow for five minutes for a total time of forty minutes. The

instructions **3821** throughout this section of the walking guide **140** advise the user what to do if warning signs such as feeling faint, dizzy occur. The user is also instructed to record in section **1203** of the plan section **1200** of the record book **104**.

[**0153**] FIG. **15C** illustrates advice on traveling **3840**. The instructions **3820** include four steps. The first step is to obtain a hospital number to call in case of an emergency at the destination. The second step is to take double the amount of supplies that are anticipated. The third step is to pack the following in a small bag: (1) all needed supplies and medications; (2) the glucose tablets **56**, included with the system **100** or snacks; (3) food with protein for a meal; (4) phone number for the user's health care provider; and (5) an emergency phone number at the destination. The fourth step is to obtain and wear an alert bracelet. The instructions **3820** also provide information on carrying insulin and blood sugar testing equipment on an airplane, including syringes, insulin pen, lancets and an insulin pump.

[**0154**] The exercise program of the system **100** also includes a step-by-step skin and foot care guide **142** illustrated in FIGS. **16A** and **16B** which provides instructions on how the user can protect skin and feet. The instructions are organized by how often different tasks should be performed. The instructions include daily instructions **3850**, and instruction for other frequencies such as instructions for twice a week **3872**, instructions for once a week **3874**, instructions for once a month **3880**, instructions for summer **3882** and instructions for winter **3884**.

[**0155**] The "Protect Your Skin and Feet" instruction guide recommends five tasks that a user perform for skin and feet each day illustrated in FIG. **16A** with explanatory photographs. The five tasks include: wash **3852**, dry **3854**, smooth **3856**, check **3858** and dress **3860**. The wash instruction **3852** instructs the user to run water and test the temperature. The water should be warm and not too hot or too cold. The instructions further provide that a user use a sudsy wash cloth on the feet and to limit the time in a bath or shower to ten minutes or less. The dry instructions **3854** instruct the user to pat dry the skin including hard to reach areas and not to rub the skin. The smooth instructions **3856** instruct the user to rub an alcohol-free skin lotion **144** provided in the system **100** over the entire body but not between the toes. In the illustrated embodiment, the skin lotion **144** is provided. The check instruction **3858** instructs the user to look for changes including blisters or sore places, thick hard places, cracks or cuts that don't heal, and red, white, black or purple spots. The instructions **3858** also suggest that the user have someone else check areas that cannot be visualized by the user. In the illustrated embodiment, the system **100** includes a mirror **146** for the user to use to look carefully at areas, such as the head, bottoms of feet and any other areas of the body difficult to view without the mirror **146**. The mirror **146** included with the system **100** can be any suitable mirror. The user is instructed to contact the health care provider if the user notices any skin changes. The dress instruction **3860** instructs the user to always wear fresh, clean hose or socks and to shake out shoes before putting them on. The instructions **3860** also recommend purchasing extra slippers and placing the slippers any place wear the user might take off his or her shoes such as by a TV, in the bathroom, near a bed, to avoid going barefoot.

[**0156**] Additional instructions **3870** included in the skin and feet guide include washing hair at least twice a week **3872**. If the user's health care provider permits, as documented in the record book **106** illustrated at FIG. **4D**, the instructions **3870** advise users to cut their toenails once a week **3874** after a bath or shower. The instructions **3870** also provide a description and an illustration of the type of clippers the user should use. The instructions **3870** recommend straight-edge toenail clippers and provide a detailed description on how to cut the toenails. The system **100** may or may not include a pair of clippers.

[**0157**] The instructions further recommend that the user perform a leap test **2880** once a month and to indicate in the record book when the leap test is performed in accordance with the description of FIG. **4J**. The leap test includes sticking the bottom of the foot to determine if the stick is felt by the user and when the stick is felt by the user.

[**0158**] The summer instructions **3882** recommend that the user use sunscreen which may or may not be included in the system **100** with SPF 15 or higher, to wear long sleeves and a hat, and to use bug spray when the user goes outside. The winter instructions **3884** recommend that the user use a humidifier or put bowls of water near the heat registers to prevent dry skin from cracking. The instructions further advise the user to bundle up when the user goes outside.

[**0159**] The system **100** includes a medication program. The medication program includes a medication guide **150**, an insulin administration guide **152**, a pill organizer **154** and glucose tablets **156**.

[**0160**] FIGS. **17A** to **17G** illustrate a secondary step-by-step medication guide **150** entitled, "The Right Meds at the Right Time." The medication guide includes four sections. The first section **4010**, illustrated in FIG. **17A**, describes how the user can organize his or her medications. The second section **4100**, illustrated in FIGS. **17A**, **17B**, **17C**, **17D**, **17E**, and **17F** provides an explanation to the user of what the medications are for. The third section **4400**, illustrated in FIG. **17G**, provides instructions to keep the medicine plan on track.

[**0161**] As illustrated in FIG. **17A**, the medication guide instructs the user to organize the medications that the user is taking into five groups. The five groups include: medications that the doctor may not know the user has **4032** such as over-the-counter medications, vitamins, herbs and supplements; blood sugar medications **4034**; heart medications **4036**; other prescription medications **4038**; and any "mystery" medications whose function is unknown to the user **4040**. The medication guide **150** instructs the user to record each of the medications in the respective sections in the record book **106** as described above and as illustrated in FIG. **4E**.

[**0162**] Once the user has organized the medications, the medication guide **150** instructs the user to understand what each medication is for **4100**. The medication guide **150** categorizes each medication by whether it is a heart medication **4110** or blood sugar medication **4210** as illustrated in FIGS. **17B** to **17F** and provides a list of each of the drugs for each category.

[**0163**] The three types of heart medications include medications that charge up the heart **4120** illustrated in FIG. **17B**, medications that lighten the load **4130** illustrated in FIG.

17C and medications that open the road 4140 illustrated in FIG. 17D. These three types of heart medications correspond to the concepts related to the heart presented in the education guide 112 described above and illustrated in FIGS. 3B and 3C. The medications that charge up the heart 4120 include pills that make the heart pump strong 4122 (i.e., digitalis), pills that make the heart pump steady 4124 (i.e., anti-disrhythmics) and pills that make each pump more efficient 4126 (i.e., beta blockers). Beta-blockers reduce a person's heart rate. The instructions liken this to using a lower gear on a bicycle to pedal uphill.

[0164] The system tries to get the user to understand that: (i) the heart medications that lighten the load 4130 include pills that flush out extra water 4132 (i.e., diuretics) and pills that make blood thinner 4134 (i.e., anti-coagulants), (2) diuretics or water pills flush out extra water from the body; and, (3) with less fluid to pump, the heart does not have to work as hard. The system and method disclosed herein provides this in an easy to understand and remember form to better enable the user to understand why they need to do certain things and show things are related.

[0165] The heart medications that open the road 4140 include pills that open arteries and veins 4142 (i.e., ace inhibitors), pills that keep arteries and veins from getting tight 4144 (i.e., angio II receptor blockers and calcium channel blockers), pills that relax the arteries 4146 (i.e., nitroglycerin and vasodilators), and pills that stop fat build-up 4148 (i.e., statins and fibrates). Ace inhibitors facilitate the opening of blood vessels. Calcium channel blockers relax a person's veins and arteries, which makes the veins and arteries wider and easier for blood to pass through. Nitroglycerin works quickly to relax a person's veins and arteries. Nitroglycerin is especially helpful to reduce chest pain by getting blood to the heart quickly. Statins reduce the amount of cholesterol the body produces. Fibrates reduce the amount of cholesterol that gets into the blood stream.

[0166] The medications that keep the user from "flooding the engine" 4220 are the blood sugar medications 4210. The list of blood sugar medications 4210 is illustrated in FIG. 17E and includes pills that prevent blood sugar from increasing too much 4222, pills that decrease blood sugar 4224, shots for people taking pills for blood sugar 4226, shots for people taking pills or insulin for blood sugar 4228, and insulin 4230.

[0167] The medication guide 150 includes a discussion 4300 on how insulin works 4310 and which insulin is right for the user 4350. This discussion is illustrated in FIG. 17F. The medication guide 150 graphically compares the onset and duration of the action and effects of each type of insulin. There are four different types of insulin. There is long-lasting 4318, medium 4314, short 4316, and quick 4318 types of insulin. To simplify the concepts for the user, the medication guide 150 also provides an analogy to different types of vehicles to illustrate the onset and duration of action of insulin. The medication guide 150 also discusses the relationship between the number of shots and adherence to the meal and exercise program 4320. For example, if a user wants to take fewer insulin shots, the guide recommends the user to plan to eat meals when the insulin is working the most, and plan exercise at the same time everyday. The user may only be given one kind of insulin, such as a medium or combination of a medium and short type of insulin. Alter-

natively, if the user wants a more flexible schedule, the doctor may give the user two kinds of insulin, long-lasting and quick insulin. The long-lasting insulin will be administered once a day and the quick insulin administered before or after a meal.

[0168] Referring to FIG. 17G and the "Four Ways to Keep Your Medicine Plan on Track" section 4400, the secondary step-by-step medication guide 150 includes instructions on being ready for the week 4410, being ready for "feeling traps" 4420, being ready to make a record 4430, and being ready for clinic visits 4440.

[0169] FIG. 17G also illustrates instructions on being ready for each week 4420. The instructions 4420 inform the user to pick one day a week (e.g., every Monday) to fill the pill organizer 154. The record book 106 may be used to ensure the correct pills are put into the pill organizer 154 in a way that corresponds to the right time of day for each pill to be taken. A portion of the pill organizer 154 corresponding to four time periods for each day (e.g., morning, noon, evening, bedtime) may be removed from the pill organizer 154 at the beginning of each day and replaced each night.

[0170] FIG. 17G also illustrates instructions on being ready for the "Feeling Traps" to fight temptation 4430 and to make a record 4440. The instructions 4430 inform the user that feeling better is not a reason to stop taking medications. On the contrary, the reason the person is feeling better is because he/she is taking the medications. A cycle that some people enter is to stop taking medications when they feel better only to wind up back in the hospital. The instructions 4440 also instruct the user to keep the record book 106 available to enter information.

[0171] FIG. 17G also illustrates instructions on being ready for doctor visits 4450. The instructions 4450 inform the user to record medication consumption in the record book 106 and to bring the record book 106 and the user's medication containers to each doctor visit. In this manner, the doctor can review the record book 106 and medication containers to determine if the right medications are being taken in the right quantities and frequencies.

[0172] The system 100 includes an insulin administration guide 152 on how to give insulin illustrated in FIGS. 18A to 18D. The instruction guide includes four sections: the "Prepare for Care" section 4510, the "Pick A Spot To Start" section 4550, the "Fill The Syringe" section 4570 and the "Give the Injection" section 4580.

[0173] In the "Prepare for Care" section illustrated in FIG. 18A, five steps are listed in the instructions to prepare to administer insulin. The first step 4512 instructs the user to collect the things the user needs to administer insulin including the record book, a pen, insulin, syringe, an alcohol pad, a clean fresh towel and a "sharps" container. The second step 4514 includes instructing the user to take the items to a clean flat place near a sink to work and to wash hands with soap and water and dry them on a clean towel. The third step 4516 instructs the user to look at the picture to learn the parts of each item. To this end, the instruction guide includes illustrations of components used in administering insulin including a vial of insulin and a syringe, the cap 4518, rubber stopper 4520 and metal band 4522 of a vial 4524 of insulin are identified. Also included is an illustration of a syringe 4526 and the needle 4528, barrel 4530, and

plunger 4532 of the syringe, along with the needle cap 4534 and the plunger cap 4536. The fourth step 4538 of the instructions instructs the user to reference the page in the record book 106 illustrated in FIG. 4E, as discussed above, where the user recorded the type of insulin and the amount of insulin to be administered. If the type of insulin the user has is not the same as that recorded in the record book 106, the insulin administration guide instructs the user to call the doctor or clinic immediately. The fifth step 4540 of the instructions advises the user to check the expiration date of the insulin.

[0174] The next section of the insulin administration guide includes the "Pick A Spot To Start" section 4550 illustrated in FIG. 18B. The instructions provide three steps in picking a spot to start. The first step 4552 instructs the user to refer to drawings indicating areas of the body where the user can give insulin, noting that the insulin administration areas 4554 illustrated are divided into separate small squares or "shot spots" 4556. Each shot spot 4556 is given a number going across row by row.

[0175] The second step 4558 instructs the user to check the record book 106 to see which doctor recommends areas are best to inject their insulin. The user is then instructed to pick an area 4554 of the body to begin a series of injections. The third step 4560 is to wash with soap and water the selected shot spot 4556 to be used. The user is instructed that a different spot on the body should be used each time an injection is given following a pretend row 4562. When the user comes to the end of a row 4562, the user is instructed to start on the next row 4564 using all the shot spots 4556 in one area 4554 before moving to another area.

[0176] The next section of instructions of the insulin administration guide includes a "Fill The Syringe" section 4570 illustrated in FIG. 18C. The "Fill The Syringe" section 4570 includes eight steps. The first step 4522 instructs the user to roll the insulin between the user's hands to mix and not to shake the insulin. The user is then instructed to take off the colored cap 4518 exposing the rubber stopper 4520 and metal band 4522 under the cap 4518 on the vial 4524. In the second step 4574, the user is instructed to use an alcohol pad to wipe the rubber stopper 4524 on the vial 4524. In the third step 4576, the user is instructed to take the caps 4536, 4534 off the plunger and the needle and lay them on the table. In the fourth step 4578, the user is instructed to check the record book 106 again to be sure the kind of insulin and number of units to be given is correct. The user is further instructed to find the same number on the barrel 4530 of the syringe 4526. The fifth step 4580 instructs the user to pull back the plunger 4532 until the end inside the barrel 4530 is at the number. The sixth step 4582 instructs the user to hold the insulin vial 4524 firm on the table, push the needle 4528 all the way through the rubber stopper 4520 and then push the plunger 4532 all the way into the syringe 4526 to fill the vial 4524 with air. The seventh step 4584 instructs the user to turn the vial 4524 and syringe 4526 upside down, and pull the plunger 4532 back to the number of units of insulin needed. If the user notices air bubbles, the user is instructed to push the plunger 4532 in to put the insulin back into the vial 4524. The instructions further instruct the user to slowly pull the plunger 4532 out again to the number of units needed and repeat until there are no air bubbles. The eighth step 4586 instructs the user to put the vial 4524 down and to hold the barrel 4530 to pull the needle

4528 out of the vial 4524. The instructions further provide that the user lay the syringe 4526 on the table preventing the needle 4528 from touching anything.

[0177] The fourth section of the insulin administration guide 152 includes the "Give The Injection" section 4590 includes six steps illustrated in FIG. 18D. The first step 4591 instructs the user to find the shot spot identified in the second section and to gently pinch up a fold of clean, dry skin. The second step 4592 instructs the user to hold the syringe so the needle will go straight in, push the needle in all the way, push the plunger in all the way injecting the insulin, and let go of the skin. The third step 4593 instructs the user to pull the needle out and press the tissue over the shot spot. The fourth step 4594 instructs the user to put the syringe in the sharps container. The fifth step 4595 instructs the user to place the sharps container where children cannot reach it. Before the container is full, the user is instructed to contact the local government to find out the rules for throwing away medical sharps. The sixth step 4596 instructs the user to document the time and shot spot 4556 in the record book 106 in the section for meds illustrated in FIG. 4E. If anything is done different than what is recorded in the record book 106, the user is instructed to record the kind of insulin used and the number of units given in the appropriate section of the record book 106 illustrated in FIG. 4I.

[0178] In the illustrated embodiment, the system 100 includes a pill organizer 154. The pill organizer 154 includes compartments for twenty-eight doses or four compartments for each of seven days of a week. The pill organizer 154 further includes labels indicative of each of the seven days and each of the four time periods of each of the seven days. It should be appreciated that any suitable pill organizer may be included in the system 100.

[0179] In addition, the illustrated embodiment of the system 100 includes glucose tablets 156. The glucose tablets may be in any suitable form and include any suitable dose.

[0180] In the illustrated embodiment, the system 100 includes a blood pressure monitor 116 and instructions for the blood pressure monitor 118. Instructions 118 for the blood pressure monitor 116 are illustrated in FIGS. 19A to 19D.

[0181] The blood pressure monitor 116 may be any type of suitable blood pressure monitor. For example, the blood pressure monitor 116 may be a manual blood pressure monitor 116 or an automatic blood pressure monitor. A manual blood pressure monitor must be manually pumped (e.g., by hand). The instructions 118 are for an automatic blood pressure monitor, which pumps automatically. The instructions 118 indicate that the first time the user is instructed to use a particular item, the name of that item appears in red.

[0182] A first section 5010 of the blood pressure monitor instructions 118 tells the user how to prepare for a blood pressure reading. A first step 5012 tells the user to prepare the blood pressure monitor 116 for use by inserting fresh batteries. A second step 5014 tells the user to have a writing instrument and the medical record book 106 handy. In addition, the second step 5014 tells the user to relax for thirty minutes if he/she just smoked a cigarette, ate something, exercised, showered, or feels stressed.

[0183] A second section 5020 of the blood pressure monitor instructions 118 tells the user how to put the cuff on

his/her arm. A third step **5022** tells the user to use the left arm (unless there is a good reason not to), remove bulky clothing, and to sit next to a table with both feet flat on the floor. A fourth step **5024** tells the user to pull open the sticky tab (e.g., velcro tab) on the cuff, so a metal bar can slide back and forth. A fifth step **5026** tells the user to slip his/her arm through the cuff while holding the cuff with the white strip and the tube on the bottom, pointing down the user's arm.

[**0184**] Turning to FIG. 19B, a sixth step **5028** tells the user to push the cuff up until the bottom edge of the cuff is about one inch above the bend inside the user's elbow. A seventh step **5030** tells the user to pull the loose flap against the cuff until the cuff is snug around the user's arm. An eighth step **5032** tells the user to press the flap against the cuff to hold the cuff tight.

[**0185**] A third section **5040** of the blood pressure monitor instructions **118** tells the user how to use the blood pressure monitor **116**. A ninth step **5042** tells the user to push the tube into the hole on the left side of the monitor **116**. A tenth step **5044** tells the user to place the blood pressure monitor **116** on the table where the display can be seen and to put the user's elbow on the table with the palm up and the cuff at the level of the user's heart.

[**0186**] Turning to FIG. 19C, an eleventh step **5046** tells the user to press the red power button on the blood pressure monitor **116** with the user's right hand. A twelfth step **5048** tells the user to press the blue start button on the blood pressure monitor **116** with the user's right hand. The twelfth step **5048** also explains that the cuff will tighten and explains how to read the numbers from the blood pressure monitor **116**. A thirteenth step **5050** tells the user how to record the pulse and blood pressure readings in the medical record book **106**. A fourteenth step **5052** tells the user to press the red power button again to turn the blood pressure monitor **106** off.

[**0187**] Turning to FIG. 19D, a fourth section **5060** of the blood pressure monitor instructions **118** tells the user how to set a pumping target on the blood pressure monitor **116**. Typically, setting the pumping target only needs to be performed after the first use of the blood pressure monitor **116**. A first step **5062** tells the user to turn the blood pressure monitor **116** on and press the white memory button. Pressing the memory button brings up the user's last blood pressure reading. A second step **5064** tells the user to calculate the user's actual pumping target by adding thirty to the last blood pressure reading. For example, if the last blood pressure reading was one hundred forty-three, adding thirty results in an actual pumping target of **173**. A third step **5066** tells the user to press the set button and check the set target number that appears on the right hand side of the screen. If the set target number is lower than the actual pumping target, then the user is instructed to keep pressing the set button until the set target number reaches or exceeds (for the first time) the actual pumping target. A fourth step **5068** tells the user to consult a list of error messages to further familiarize the user with the blood pressure monitor **116**. The blood pressure monitor instructions **118** also include a blank portion **5070** where the user may record any special instructions from his/her doctor about the blood pressure monitor **116** and/or taking the blood pressure readings.

[**0188**] FIGS. 20A and 20B illustrate a secondary step-by-step tobacco addiction guide **158** included in the system **100**

illustrated embodiment of the guide to assist the user in stop smoking. The tobacco addiction guide **158** includes facts about quitting smoking **5110**, instructions on preparing to quit smoking **5120**, instructions on quitting smoking **5130**, and instructions on preparing for the effects of quitting smoking **5140**.

[**0189**] FIG. 20A illustrates facts about quitting smoking **5110**. The facts **5110** inform the user that most people who attempt to quit smoking succeed. However, quitting may take more than one attempt. In fact, about two thirds of people who try to quit succeed after multiple attempts. The facts **5110** also inform the user that just as many people succeed in quitting smoking without signing up for a program as those who succeed with a program. The facts **5110** also inform the user that heavy smokers succeed in quitting just as often as light smokers. So, it does not matter how much the person currently smokes. The facts **5110** also inform the user that most people who successfully quit smoking quit when they have some other big change in their life. For example, beginning to use the system **100** may be the big change.

[**0190**] FIG. 20A also illustrates instructions on preparing to quit smoking **5120**. The instructions **5120** instruct the user to add toothpicks, sugarless gum, and diet juice to the person's grocery shopping list. In addition, the instructions **5120** suggest that if the person does not feel that he/she can stop smoking right away, to switch to a brand of cigarettes that contain more nicotine to change the taste associated with smoking and make the person feel sick. The instructions **5120** also tell the user to pick a date to stop smoking completely and to mark that date on their calendar. The instructions **5120** also instruct the user to tell friends and family (and himself/herself) that he/she only intends to quit for two days (i.e., that quitting is not a big deal).

[**0191**] FIG. 20B illustrates instructions on quitting smoking **5130**. The instructions **5130** instruct the user to dispose of all cigarettes. Alternatively, the user may store any existing cigarettes in a separate place, away from lighters and matches. When the urge to smoke occurs, the instructions **5130** have the user ask himself/herself why they smoke. If the user smokes to be social, the instructions **5130** suggest visiting a non-smoking friend and/or joining a community volunteer group. If the user smokes to relieve stress, the instructions **5130** suggest squeezing a stress ball, doing some exercises (e.g., in the user's chair at work and/or at home), riding a bike, and/or playing a sport. If the user smokes to keep from eating, the instructions **5130** suggest biting on a toothpick, chewing sugarless gum, drinking some water, and/or drinking diet juice. After two days, the user performs a self-assessment and makes any adjustments that are needed to quit smoking.

[**0192**] FIG. 20B also illustrates instructions on preparing for the effects of quitting smoking **5140**. The instructions **5140** inform the user that if he/she feels sick, not to worry because the feeling will pass. The instructions **5130** also inform the user that if the user feels the urge to have a cigarette, not to panic because the urge will pass. The instructions **5140** also inform the user that if the user has a bad day, not to worry and to just start the program over again. The instructions **5140** encourage the user by informing the user that by quitting smoking, the user should notice

several positive benefits such as having more energy, having an easier time breathing, and that things will smell and taste better.

[0193] FIG. 21 illustrates a secondary step-by-step guide 160. The stress management guide 160 includes an explanation of how to recognize stress 5210 and ways to manage stress such as breathing exercises 5220, stretching exercises 5230, and actions to avoid stress 5240, and how to rate stress 5250.

[0194] The explanation of how to recognize stress 5210 informs the user that everyone experiences some stress. Stress is the body's natural reaction to tension, pressure, and/or change. Small amounts of stress make life more interesting and less boring. However, excess stress, especially prolonged and unrelieved stress, can be mentally and physically unhealthy. The stress management guide includes a listing of signs of stress 5210 such as headaches, upset stomach, hopeless feelings, etc. Many of these symptoms may also be caused by medical conditions, but the guide 160 points out that effective management of the stress may improve health. In addition, the stress management guide 160 may include a scale on which to rate the stress of the user at a particular time 5250.

[0195] The guide 160 includes anti-stress exercises such as breathing exercises 5220 and stretching exercises 5230. For example, the breathing exercises may include closing eyes, and breathing in or out for periods of time while moving certain parts of the body such as the shoulders. The stretching exercises may include a finger fan exercise, an upper back stretch, an ear-to-shoulder exercise, an overhead reach exercise, a knee-pull exercise, and a waist-bend exercise. The finger fan exercise includes extending the arms and spreading the fingers. The upper back stretch includes sitting up straight with the fingers interlocked behind the head and bring the elbows back. The ear-to-shoulder exercise includes lowering the ear to the shoulder. The overhead reach exercise includes raising the arms over the head with interlaced fingers. The knee-pull exercise includes pulling one knee at a time up to the chest in a seated position. The waist-bend exercise includes bending from side-to-side at the waist with the arms extended over the head.

[0196] The instructions on ways to manage stress 5240 include actions such as thinking positively and being around other positive people, avoiding being overly demanding on oneself and getting help when needed, writing and following a reasonable to do list each day, dividing big tasks into smaller more manageable tasks, eating a healthier diet, minimizing and/or eliminating consumption of coffee and sugar, getting plenty of sleep, getting enough exercise, making time to relax, avoiding tobacco, alcohol, and drugs, using mistakes and setbacks as opportunities to learn, avoiding high stress tasks, talking to a friend about disappointments and frustration before they build up and sharing your successes, admitting when you are wrong, eliminating everyday sources of stress such as loud music and clutter, having fun and laugh, knowing it is okay to cry, practicing deep breathing, being active during everyday tasks such as taking the stairs instead of the elevator and/or not sitting when talking on the telephone, and doing anti-stress stretches every day.

[0197] The stress management guide 160 may include other examples of how to act in response to stressful

circumstances 5240 such as those listed in FIG. 21: if you have a negative neighbor, walk away, if a task is too big, break it down and get help; if the stakes are too high, get feedback early; if lonely, volunteer; if sad, get a pet; if nervous, cut out coffee and sugar; if a mistake is made, admit it and move on; if no time, say no and suggest someone else; if frowning, enjoy some comedy; if feeling stuck, walk around, outside if possible; if overwhelmed, make a "to do" list. These and other such examples may be included in the stress management guide 160.

[0198] It should be appreciated that any of the materials included in the system 100 may be included in any suitable form or format. Elements of the system may be included as a software product, stored on any suitable paper form, in a computer readable form storage device such as a CD-ROM, DVD. The instructions may be provided in additional video form.

[0199] It should be appreciated that the disclosed integrated system or any elements thereof may be provided in any suitable sensory form and on any suitable medium and combinations thereof. For example, any element of the disclosed integrated system may be provided in audio, visual, or tactile form such as Braille. Any element of the disclosed integrated system may be provided on paper, on any suitable machine or computer readable form such as CD-ROM, DVD, or any other suitable physical or electronic medium.

[0200] It should be appreciated that the system, apparatus and method disclosed herein provides: (a) an easy to understand and remember method for better enabling the user to understand why they need to do certain things and how things are related, (b) instructions which integrate the care for multiple medical conditions, and (c) directions on meal planning and other activities for multiple medical conditions, all without overlapping for contrary instructions. The system and apparatus are thus configured to enable a user to simultaneously provide integrated self care for multiple different medical conditions including diabetes, high blood pressure and heart failure.

[0201] In summary, methods and apparatus for blood sugar control, blood pressure control and heart failure care have been provided. The foregoing description has been presented for the purposes of illustration and description. It is not intended to be exhaustive or to limit the invention to the exemplary embodiments disclosed. Many modifications and variations are possible in light of the above teachings. It is intended that the scope of the invention be limited not by this detailed description, but rather by the claims appended hereto.

The invention is claimed as follows:

1. An integrated blood sugar control, blood pressure control and heart failure self-care system, the system comprising:

- a magnet board, the magnet board including a portion to write a telephone number and a label indicating the telephone number is a doctor's telephone number;
- a blood pressure monitor;
- a pill organizer;
- a stress management guide;

a primary step-by-step guide, the primary step-by-step guide including a plurality of primary steps associated with heart, blood pressure and blood sugar including at least three of:

- (1) a dietary program;
- (2) a medicine compliance program;
- (3) a tobacco addiction program;
- (4) a stress management program;
- (5) an exercise program;
- (6) a skin and foot care program; and
- (6) a medical record keeping program; and

at least one secondary step-by-step guide, the secondary step-by-step guide including a plurality of secondary steps associated with at least one of the primary steps; and

a container for storing the above components.

2. The system of claim 1, wherein the blood pressure monitor includes a digital blood pressure monitor.

3. The system of claim 1, including a set of blood pressure monitor instructions, the blood pressure monitor instructions including at least three of:

- (a) at least one instruction for getting the blood pressure monitor ready for use;
- (b) at least one instruction for getting a user prepared;
- (c) at least one instruction for attaching a cuff of the blood pressure monitor to the user; and
- (d) at least one instruction for operating the blood pressure monitor.

4. The system of claim 1, wherein the pill organizer includes a twenty-eight dose pill organizer.

5. The system of claim 1, wherein the twenty-eight dose pill organizer includes: four compartments for each of seven days of a week, and labels indicative of each of the seven days and four time periods during each of the seven days.

6. The system of claim 1, wherein the stress management guide includes a pocket-sized stress management guide.

7. The system of claim 1, wherein the primary steps of the primary step-by-step guide includes steps associated with:

- (1) the dietary program;
- (2) the medicine compliance program;
- (3) the tobacco addiction program;
- (4) the stress management program;
- (5) the exercise program; and
- (6) the medical record keeping program.

8. The system of claim 1, wherein the plurality of secondary steps includes dietary program steps, the dietary program steps including at least three of:

- (a) instructions on how to eat less salt;
- (b) instructions on how to eat less cholesterol;
- (c) instructions on how to eat carbohydrates;
- (d) a food poster;
- (e) a loose food guide;

(f) a fast food guide;

(g) a portion plate;

(h) a food selection game;

(i) a meal planner; and

(e) a shopping list.

9. The system of claim 1, wherein the plurality of secondary steps includes medicine compliance program steps, the medicine compliance program steps including at least two of:

- (a) instructions on how to organize pills;
- (b) information on the purpose of different types of pills; and

(c) instructions on how to administer insulin.

10. The system of claim 1, wherein the plurality of secondary steps includes tobacco addiction program steps, the tobacco addiction program steps including at least three of:

- (a) facts about people who try to quit smoking;
- (b) a shopping list;
- (c) instructions on marking a calendar; and
- (d) instructions on alternatives to smoking.

11. The system of claim 1, wherein the plurality of secondary steps includes stress management program steps, the stress management program steps including at least three of:

- (a) an explanation of stress;
- (b) a list of stress signs;
- (c) a list of ways to manage stress; and
- (d) a plurality of anti-stress exercises.

12. The system of claim 1, wherein the plurality of secondary steps includes exercise program steps, the exercise program steps including at least two of:

- (a) information about the benefits of walking;
- (b) a list of things needed for walking.

13. The system of claim 1, wherein the plurality of secondary steps includes medical record keeping program steps, the medical record keeping program steps including at least three of:

- (a) medication compliance recording;
- (b) weight recording;
- (c) blood pressure recording;
- (d) blood sugar recording;
- (e) exercise recording;
- (f) tobacco consumption recording; and
- (g) pain recording.

14. The system of claim 1, wherein the plurality of secondary steps includes a medical record keeping step and the system includes a medical record keeping form, the medical record keeping form including at least three of:

- (a) a place to record answers to questions posed to a doctor;

- (b) a place to record blood test results;
- (c) a place to record prescription medication information;
- (d) a place to record blood pressure readings;
- (e) a place to record blood sugar readings;
- (e) a place to record body weights;
- (f) a place to record tobacco consumption; and
- (g) a place to record pain information.

15. A method for integrated blood sugar control, blood pressure control and heart failure care, the method comprising:

providing a blood pressure monitor;

providing a pill organizer;

providing a stress management guide;

providing a primary step-by-step guide, the primary step-by-step guide including a plurality of primary steps associated with heart, blood pressure and blood sugar including at least three of:

- (1) a dietary program;
- (2) a medicine compliance program;
- (3) a tobacco addiction program;
- (4) a stress management program;
- (5) a skin and foot care program;
- (6) an exercise program; and
- (7) a medical record keeping program; and

providing at least one secondary step-by-step guide, the secondary step-by-step guide including a plurality of secondary steps associated with at least one of the primary steps.

16. A system for integrated blood sugar control, blood pressure control and heart failure self-care by a user, the system comprising:

a blood pressure monitor;

a primary step-by-step guide associated with the blood pressure monitor, the primary step-by-step guide including a plurality of primary steps associated with blood pressure, blood sugar and heart failure including at least five of:

- (1) a dietary program;
- (2) a medicine compliance program;
- (3) a tobacco addiction program;
- (4) a stress management program;
- (5) a skin and foot care program
- (5) an exercise program; and
- (6) a medical record keeping program;

at least one secondary step-by-step guide, the secondary step-by-step guide including a plurality of secondary steps associated with at least one of the primary steps; and a container for storing the above components.

* * * * *

专利名称(译)	综合血糖控制，血压控制和心力衰竭自我护理系统和方法		
公开(公告)号	US20080050709A1	公开(公告)日	2008-02-28
申请号	US11/771793	申请日	2007-06-29
[标]申请(专利权)人(译)	CAREGUIDE SYST		
申请(专利权)人(译)	CAREGUIDE系统公司.		
当前申请(专利权)人(译)	CAREGUIDE系统公司.		
[标]发明人	ARUFFO SYLVIA GREY SHIRLEY OUTLAND CAROL FARAH JUDITH FRANKS		
发明人	ARUFFO, SYLVIA GREY, SHIRLEY OUTLAND, CAROL FARAH, JUDITH FRANKS		
IPC分类号	G09B23/28 A47B81/00 A61B5/00		
CPC分类号	A61B5/021 G09B19/00 A61B5/14532		
优先权	60/806201 2006-06-29 US 60/806200 2006-06-29 US 60/806203 2006-06-29 US		
外部链接	Espacenet USPTO		

摘要(译)

公开了用于血糖控制，血压控制和心力衰竭护理的方法和装置。该方法
和装置使用户能够向他/她自己提供血糖控制，血压控制和心力衰竭护
理。例如，在由医生治疗心脏问题之后，人们可以使用所公开的系统来
防止额外的心脏相关问题。

