



US 20030225162A1

(19) **United States**

(12) **Patent Application Publication**  
**Schreiber et al.**

(10) **Pub. No.: US 2003/0225162 A1**

(43) **Pub. Date: Dec. 4, 2003**

(54) **METHODS FOR THE DIAGNOSIS OF HEMODIALYSIS PATIENTS AND UTILIZATION OF SUCH METHODS TO IMPROVE THE ADMINISTRATION OF INTRAVENOUS LEVOCARNITINE TREATMENTS TO HEMODIALYSIS PATIENTS**

(75) Inventors: **Brian Schreiber**, Neenah, WI (US);  
**Yvonne T. Lewis**, New Orleans, LA (US)

Correspondence Address:

**HOGAN & HARTSON LLP  
IP GROUP, COLUMBIA SQUARE  
555 THIRTEENTH STREET, N.W.  
WASHINGTON, DC 20004 (US)**

(73) Assignee: **Sigma-Tau Pharmaceuticals, Inc.**,  
Gaithersburg, MD

(21) Appl. No.: **10/355,263**

(22) Filed: **Jan. 31, 2003**

**Related U.S. Application Data**

(60) Provisional application No. 60/352,505, filed on Jan. 31, 2002.

**Publication Classification**

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 31/198**; A61B 5/00;  
G06F 19/00; G01N 33/48;  
G01N 33/50


(52) **U.S. Cl.** ..... **514/565**; 600/300; 702/19

(57) **ABSTRACT**

The disclosed invention pertains to methods for diagnosing clinical conditions that are common in hemodialysis patients and that may be related to abnormal carnitine metabolism resulting from hemodialysis. Further, the present invention pertains to methods for monitoring and improving the administration of therapeutic intravenous levocarnitine to such patients. Clinical algorithms have been developed for the clinical symptoms seen in end-stage renal disease (ESRD) patients that may be related to carnitine deficiency. Monitoring tools to assist healthcare professionals in implementing the clinical algorithms are also provided.

***Patient Diagnosis Tool***

***Summary***



**Summary**

**Patient Diagnosis Tool**




Figure 1

# Cardiomyopathy

# Patient Diagnosis




Figure 2a




Figure 2b

# Cardiomyopathy

# Charting Tool

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

### TREATMENT CONSIDERATIONS

	<u>YES</u>	<u>NO</u>	<u>NRT*</u>
• CONTROL VOLUME	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• CONTROL BLOOD PRESSURE	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• TREAT ARRHYTHMIA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• ADJUST HCT/HGB	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• REDUCE AFTER-LOAD	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• EVALUATE PRE-DIALYSIS PLASMA CARNITINE LEVEL	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• BELOW NORMAL PRE-DIALYSIS PLASMA CARNITINE LEVEL (<40-50 μMOL/L)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

*Re-evaluate by clinical exam to be certain that therapies given are adequate.*

Signature \_\_\_\_\_ Date \_\_\_\_\_

Figure 3

\*NRT = No response to therapy

# Cardiac Arrhythmias Patient Diagnosis




Figure 4

# Cardiac Arrhythmias

# Charting Tool

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

Underlying Medical Condition:

- |   | <u>YES</u>               | <u>NO</u>                | <u>NRT*</u>              |
|---|--------------------------|--------------------------|--------------------------|
| • Ischemic heart disease                              | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Thyroid disease                                     | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Electrolyte abnormal                                | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Chronic lung disease                                | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Drugs (digoxin, theophylline, antiarrhythmics, etc) | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Dialysis Related Effect:

- |  | <u>YES</u>               | <u>NO</u>                | <u>NRT*</u>              |
|--|--------------------------|--------------------------|--------------------------|
| • Optimize volume status   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Adjust dialysate (K+)  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Inadequate dialysis  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Oxygen on dialysis   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Evaluate pre-dialysis plasma carnitine level                     | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Below normal pre-dialysis plasma carnitine level (<40-50 μmol/L) | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Re-evaluate by clinical exam to be certain that therapies given are adequate.

Signature \_\_\_\_\_ Date \_\_\_\_\_

Figure 5

\*NRT = No response to therapy

# Muscle Weakness

# Patient Diagnosis




Figure 6

# Muscle Weakness

# Charting Tool

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

- WEAKNESS IN ARTHRITIC OR INJURED JOINT
- SENSORY ABNORMALITY PRESENT
- STRETCH REFLEXES
- IS MUSCLE WEAKNESS RELATING TO MYOPATHY?

<u>YES</u>	<u>NO</u>	<u>NRT*</u>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

*If yes, then proceed to Diagnosis Tool on Myopathy.*

*If no, then proceed to Diagnosis Tool on Malaise/Fatigue.*

*Re-evaluate by clinical exam to be certain that therapies given are adequate.*

Signature \_\_\_\_\_ Date \_\_\_\_\_

\*NRT = No response to therapy

Figure 7

# Patient Diagnosis

# Muscle Myopathy




Figure 8

**Muscle Myopathy** **Charting Tool**

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

*After ruling out muscle weakness as outlined in the algorithm, proceed as follows:*

	<u>Yes</u>	<u>No</u>	<u>NRT*</u>
• Infectious/inflammatory/malignant	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Thyroid disorder	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Cushing's Disease	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Alcohol/Drugs	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Uncontrolled hyperparathyroidism	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Osteomalacia	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Aluminum toxicity	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Evaluate pre-dialysis plasma carnitine level	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Below normal pre-dialysis plasma carnitine level (<40-50 μmol/L)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

*Re-evaluate by clinical exam to be certain that therapies given are adequate.*


Signature \_\_\_\_\_ Date \_\_\_\_\_

**Figure 9**

\*NRT = No response to therapy

**Malaise/Fatigue**

**Patient Diagnosis**



**Figure 10**

**Malaise/Fatigue**

**Charting Tool**

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

After ruling out muscle weakness as outlined in the algorithm, proceed as follows:

Rule out superimposed non-renal illness/condition:

- |                            | <u>Yes</u>               | <u>No</u>                | <u>NRT*</u>              |
|----------------------------|--------------------------|--------------------------|--------------------------|
| • Depression               | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Cancer                   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Congestive heart failure | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Thyroid disease          | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • COPD                     | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Uncontrolled diabetes    | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Infection                | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Sedative Drugs           | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Evaluate Dialysis Specific Factors:

- |  | <u>Yes</u>               | <u>No</u>                | <u>NRT*</u>              |
|--|--------------------------|--------------------------|--------------------------|
| • Inadequate dialysis  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| - Inappropriate dialysis weight                                    | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| - Under-dialyzed   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Incompletely compensated anemia (EPO dose adjusted)              | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Evaluate pre-dialysis plasma carnitine level                     | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Below normal pre-dialysis plasma carnitine level (<40-50 μmol/L) | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

\*\* If no adequate response see Patient Diagnosis Algorithm for Delayed/Diminished Response to EPO

**Re-evaluate by clinical exam to be certain that therapies given are adequate.**

Signature \_\_\_\_\_ Date \_\_\_\_\_

Figure 11

\*NRT = No response to therapy




Figure 12

**Delayed/Diminished EPO Response** **Charting Tool**

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

*If the patient fails to respond or to maintain a response to EPO doses within the recommended dosing range, evaluate/consider the following etiologies:*

	<u>Yes</u>	<u>No</u>	<u>NRT*</u>
• Iron deficiency	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Underlying infectious, inflammatory or malignant process	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Occult blood loss	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Underlying hematologic disease (i.e., thalassemia, refractory anemia or other myelodysplastic disorders)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Hemolysis	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Osteitis fibrosa cystica	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Aluminum intoxication	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Vitamin deficiencies (folic acid or vitamin B12)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Evaluate pre-dialysis plasma carnitine level	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
• Below normal pre-dialysis plasma carnitine level (< 40-50 μmol/L)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**Re-evaluate by clinical exam to be certain that therapies given are adequate.**

Signature \_\_\_\_\_ Date \_\_\_\_\_

Figure 13

\*NRT = No response to therapy

**PATIENT DIAGNOSIS**

1400

**HYPOTENSION**





Figure 14a



Continued from Fig. 14a

Figure 14b

**Dialysis Related Hypotension** **Charting Tool**

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_

Acute Episode:

- |   | <u>YES</u>               | <u>NO</u>                | <u>NRT*</u>              |
|---|--------------------------|--------------------------|--------------------------|
| • Trendelenberg position                  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • IV fluid                                | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Decrease or Zero UFR                    | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Consider Oxygen                         | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Consider causes as in chronic sustained | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Chronic Sustained

- |                  | <u>YES</u>               | <u>NO</u>                | <u>NRT*</u>              |
|------------------|--------------------------|--------------------------|--------------------------|
| • Cardiac (echo) | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Pericarditis   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • CHF            | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Valve          | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Ischemia       | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Infections     | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Hemotologic    | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| • Sepsis         | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

*Re-evaluate by clinical exam to be certain that therapies given are adequate.*

Signature \_\_\_\_\_ Date \_\_\_\_\_

**Figure 15**

\*NRT = No response to therapy

**Clinical Monitoring Tool**

Patient Name \_\_\_\_\_ I.D. # \_\_\_\_\_  
 Diagnosis: \_\_\_\_\_ Dialysis Start Date: \_\_\_\_\_

Parameter	Baseline	One Month	Three Months	Six Months
Clinical Laboratory	Hemoglobin			
	Hematocrit			
	Trans Saturation			
	RBC			
	Albumin			
	Glucose			
	Phosphorus			
	BUN			
	KTV or URR			
	Pre-dialysis Plasma Carnitine (µmol/L)			
	EPO Dosage			
	Iron Dosage			
	Carnitor® (levocarnitine) Injection Dosage			
	Intradialytic Cramping (frequency)			
Patient Assessment	Hypotensive Episodes (frequency)			
	SGA Score			
	Ejection Fraction (%)			
	Appetite Changes			
	Fatigue			
	Physical Appearance			
	Hospitalizations (changes in frequency)			
Meals				

Continue therapy: Yes  No   
 Signature \_\_\_\_\_ Date \_\_\_\_\_

Figure 16

**METHODS FOR THE DIAGNOSIS OF  
HEMODIALYSIS PATIENTS AND UTILIZATION  
OF SUCH METHODS TO IMPROVE THE  
ADMINISTRATION OF INTRAVENOUS  
LEVOCARNITINE TREATMENTS TO  
HEMODIALYSIS PATIENTS**

**CROSS REFERENCE TO RELATED  
APPLICATIONS**

[0001] The present application claims priority from U.S. Provisional Patent Application Serial No. 60/352,505, **METHODS FOR THE DIAGNOSIS OF HEMODIALYSIS PATIENTS AND UTILIZATION OF SUCH METHODS TO IMPROVE THE ADMINISTRATION OF INTRAVENOUS LEVOCARNITINE TREATMENTS TO HEMODIALYSIS PATIENTS**, filed Jan. 31, 2002, which is hereby incorporated by reference. Furthermore, the present application is related in subject matter to U.S. Pat. No. 6,335,369, **TREATING CHRONIC UREMIC PATIENTS UNDERGOING PERIODICAL DIALYSIS**, issued Jan. 1, 2002, the specification of which is hereby incorporated by reference in its entirety.

**FIELD OF THE INVENTION**

[0002] The present invention relates to methods for diagnosing and treating various conditions prevalent in hemodialysis patients with end stage renal disease. In particular, the present invention pertains to methods for diagnosing clinical conditions that are common in hemodialysis patients and that may be related to abnormal carnitine metabolism resulting from hemodialysis; thus, the present invention provides methods for monitoring and improving the administration of therapeutic intravenous levocarnitine to such patients.

**BACKGROUND OF THE INVENTION**

[0003] Studies indicate that more than 70% of the carnitine present in the plasma of a hemodialysis patient can be removed during a dialysis session. Carnitine is a naturally occurring substance in the human body required for energy metabolism at the cellular level because it transports fatty acid-derivatives into the inner aspect of the mitochondria to produce energy and removes various acyl moieties from the mitochondria and cells. Dialytic loss of carnitine by patients undergoing hemodialysis is thought to be attributable to the compound's relatively small molecular weight, high water solubility, and poor protein binding. Carnitine levels are further diminished in end stage renal disease patients by reduced renal synthesis and reduced intake of meat and dairy foods.

[0004] Primary and secondary carnitine deficiency syndromes are well-described entities whose clinical sequelae could result in serious illness or death. Carnitine deficiency syndromes are characterized by such diverse symptoms as cardiomyopathy, muscle weakness, lipid storage myopathy, hepatic dysfunction, encephalopathy, failure to thrive, and recurrent infections.

[0005] As described in U.S. Pat. No. 6,335,369, the administration of levocarnitine may be beneficially used to treat carnitine deficiency in patients with end stage renal disease ("ESRD") who are undergoing regular hemodialysis. Chronic uremic patients undergoing periodical hemodi-

alysis are treated with carnitine or one of its salts to prevent or treat carnitine deficiency in patients with end stage renal disease. An effective dose of levocarnitine (the recommended starting intravenous dosage is typically 10-20 mg of drug per kg of patient dry body weight) is administered intravenously via a venous return line after each dialysis session. Initiation of such levocarnitine injection therapy may be prompted by pre-dialysis plasma free carnitine concentrations that are below normal (normal concentrations being approximately 40-50 micromoles/liter ( $\mu\text{mol/L}$ )). Such intravenous administration of levocarnitine to end stage renal disease patients on hemodialysis results in increased plasma carnitine concentrations and thereby makes it possible to correct for the loss of plasma carnitine which otherwise takes place during hemodialysis therapy. Importantly, immediately after giving intravenous carnitine to a patient at the end of a dialysis session, carnitine levels in the patient's plasma rise to high levels and then return to a pre-dialysis baseline levels after approximately 10-24 hours (presumably because the carnitine has entered the tissues). In this manner, it is possible to avoid tissue carnitine depletion, which is a long-term consequence of repeated losses of carnitine from plasma that the patient undergoes during successive dialytic sessions over a prolonged period of time.

[0006] Prior studies have listed specific subsets of patients in whom intravenously administered carnitine had been associated with improvement in clinical parameters. These subsets of patients have included those with cardiomyopathy, skeletal muscle weakness/myopathy, anemia of uremia unresponsive to or requiring large doses of erythropoietin (EPO), lack of energy (which has a negative effect upon quality of life), severe and persistent muscle cramps, and/or intradialytic hypotensive episodes. Despite these studies, the use of levocarnitine in dialysis patients is sometimes limited.

[0007] Cardiomyopathy, muscle weakness, and fatigue are often due to factors other than carnitine deficiency. The ability to methodically analyze these common conditions is critical to the appropriate application of this therapy. Non-critical use of carnitine is both medically incorrect and economically wasteful. Thus, there remains a need in the art for methods to differentiate other etiologies for clinical conditions that have been associated with dialysis related carnitine deficiency and for which intravenous levocarnitine therapy is being considered in the treatment of dialysis patients.

**SUMMARY OF THE INVENTION**

[0008] In light of the above-described and other deficiencies, the present invention as disclosed herein is intended to help health care professionals in the diagnosis of ESRD patients undergoing dialysis with diagnosing clinical conditions and symptoms that are common in hemodialysis patients and to help document the medical necessity of selected treatments. Clinical algorithms have been developed for the clinical symptoms seen in ESRD patients that may be related to carnitine deficiency. These symptoms include secondary cardiomyopathy, dialysis related hypotension, cardiac arrhythmia, muscle wasting or weakness, muscle cramping, protein catabolism, lack of energy, and delayed or diminished response to erythropoietin ("EPO"). Monitoring tools to assist health care professionals in implementing the clinical algorithms are also provided.

[0009] ESRD patients with the above conditions, as well as those with poor erythropoietin response, who have demonstrated an inadequate response to appropriate therapy are prescribed a course of intravenous levocarnitine in a recommended range of drug dose and duration of therapy. Once patients have been on this therapy in recommended dose and duration a clinical outcome monitoring tool is used to evaluate a response.

[0010] Embodiments of the present invention provide methods for monitoring and improving the administration of therapeutic levocarnitine to ESRD patients by providing clinical algorithms and accompanying tools that are adapted to assist health care professionals in assessing patient status from such symptoms and identifying appropriate therapeutic actions to take. These methods can also be incorporated as a computer-based software application that records the data for individual patients and systematically guides the health care practitioner through the required steps.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0011] FIG. 1 is a schematic diagram depicting the basic patient symptoms and associated diagnostic modules of the present invention.

[0012] FIG. 2 is a schematic diagram depicting the steps for a cardiomyopathy diagnostic module.

[0013] FIG. 2b is a schematic diagram depicting alternate steps for a cardiomyopathy diagnostic module according to one embodiment of the invention.

[0014] FIG. 3 is an example of a charting tool for implementing a cardiomyopathy clinical algorithm according to the embodiments of the present invention.

[0015] FIG. 4 is a schematic diagram depicting the steps for a cardiac arrhythmias diagnostic module.

[0016] FIG. 5 is an example of a charting tool for implementing a cardiac arrhythmias clinical algorithm according to the embodiments of the present invention.

[0017] FIG. 6 is a schematic diagram depicting the steps for a muscle weakness diagnostic module.

[0018] FIG. 7 is an example of a charting tool for implementing a muscle weakness clinical algorithm according to the embodiments of the present invention.

[0019] FIG. 8 is a schematic diagram depicting the steps for a muscle myopathy diagnostic module.

[0020] FIG. 9 is an example of a charting tool for implementing a muscle myopathy clinical algorithm according to the embodiments of the present invention.

[0021] FIG. 10 is a schematic diagram depicting the steps for a malaise/fatigue diagnostic module.

[0022] FIG. 11 is an example of a charting tool for implementing a malaise/fatigue clinical algorithm according to the embodiments of the present invention.

[0023] FIG. 12 is a schematic diagram depicting the steps for a delayed or diminished erythropoietin ("EPO") response diagnostic module.

[0024] FIG. 13 is an example of a charting tool for implementing a delayed or diminished EPO response clinical algorithm according to the embodiments of the present invention.

[0025] FIG. 14 is part one of a schematic diagram depicting the steps for a dialysis related hypotension diagnostic module.

[0026] FIG. 15 is part two of the schematic diagram depicting the steps for a dialysis related hypotension diagnostic module.

[0027] FIG. 16 is an example of a charting tool for implementing a dialysis related hypotension clinical algorithm according to the embodiments of the present invention.

[0028] FIG. 17 is an example of a clinical outcome monitoring tool to track a patient's response to levocarnitine therapy according to the embodiments of the present invention.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0029] As shown in FIGS. 1-16, embodiments of the present invention are methods that help health care professionals to associate common symptoms identified in end stage renal disease ("ESRD") patients with one or more appropriate diagnosing algorithms and their associated tools for monitoring and improving the potential administration of therapeutic levocarnitine to such patients.

[0030] As shown in FIG. 1, symptoms can be associated with one of six diagnosing algorithm and tool sets, namely, the pairs for cardiomyopathy (FIGS. 2-3), cardiac arrhythmia (FIGS. 4-5), muscle myopathy (FIGS. 6-9), malaise/fatigue (FIGS. 10-11), delayed/diminished response to EPO (FIGS. 12-13), and dialysis related hypotension (FIGS. 14-16). In use, a health care professional applies the appropriate diagnosing algorithm (for example, the cardiomyopathy algorithm of FIG. 2) depending upon the symptom(s) (e.g., secondary cardiomyopathy) presented by the ESRD patient to determine the patient's status and potential suitability for levocarnitine administration. Additionally, the health care professional concurrently with the application of the diagnosing algorithm uses the appropriate monitoring and treatment tool (in the case of the above example, the cardiomyopathy tool illustrated by FIG. 3) to record the patient's response to the diagnostics (and, potentially, treatments) performed according to the related diagnosing algorithm (i.e., FIG. 2 for cardiomyopathy). FIGS. 2-16 describe the application of the diagnosing algorithms and tools according to the invention in detail.

[0031] FIG. 2 depicts the appropriate diagnosing algorithm for cardiomyopathy 200 to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of secondary cardiomyopathy as shown in FIG. 1. According to the algorithm in FIG. 2, a patient with these symptoms is evaluated by clinical exam 210 and considered for a series of treatments 220 known to those skilled in the art that include controlling fluid volume during dialysis, controlling blood pressure, treating arrhythmia, adjusting hematocrit (HCT)/hemoglobin (HGB), and reducing after-load. If these treatments provide adequate improvement upon evaluation 230, then they are continued 235, and the patient will continue to be monitored in subsequent clinical exams 210. If the treatments 220 do not provide adequate improvement upon evaluation 240, then the health care professional will evaluate the patient's pre-dialysis

plasma carnitine concentration **241**, determine if the pre-dialysis carnitine level is below normal **242**, and, if so, initiate levocarnitine injection therapy **243**. The steps of algorithm **200** are charted in **FIG. 3**, such that a health care professional can effectively monitor the diagnostic process.

[**0032**] In one embodiment of the appropriate diagnosing algorithm for cardiomyopathy **200**, step **220** is made of additional steps. One step is considering the possibility of global Hypokinesia **221**, and if present, implementing appropriate treatments such as strict volume control, angiotensin-converting enzyme (ACE) and angiotensin receptor blocker (ARB) treatments, beta blocker treatment, arrhythmia treatment, maximum HCT, (not to be excessive), afterload reduction, or strict blood pressure control. Another step is determining if there may be a diastolic dysfunction **224**, and if so, treating appropriately **225**. An additional step is determining if regional abnormalities are present with/without suggestive symptoms through a cardiac catheterization or non-invasive screen **223**. If a remedial lesion **227** is found, then surgery or angioplasty **228** is required. If no remedial lesion is found **226**, then the patient is re-evaluated by clinical exam **229**.

[**0033**] **FIG. 4** depicts the appropriate diagnosing algorithm **400** to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of cardiac arrhythmia as shown in **FIG. 1**. According to the algorithm in **FIG. 4**, a patient with these symptoms is evaluated by clinical exam **410**, and the health care professional determines whether the symptoms are related to an underlying medical condition **420** or are related to effects from dialysis **430**. If the arrhythmia symptoms are related to an underlying medical condition, then appropriate treatment **425** relating to the underlying condition must first be provided, and the patient will continue to be monitored in subsequent clinical exams **410**. If the arrhythmia symptoms are determined to be related to dialysis treatment **430**, a series of adjustments to the dialysis process **440** known to those skilled in the art are considered that include optimizing the volume status, adjusting the dialysate if possible, prolonging dialysis to moderate fluid shifts, and administering oxygen during dialysis. If these changes to the dialysis process control the arrhythmia symptoms upon evaluation **450**, then they are continued **455**, and the patient will continue to be monitored in subsequent clinical exams **410**. If changes to the dialysis process do not control the arrhythmia symptoms **450**, then the health care professional will evaluate the patient's pre-dialysis plasma carnitine concentration **461**, determine if the pre-dialysis carnitine level is below normal **462**, and, if so, initiate levocarnitine injection therapy **463**. The steps of algorithm **400** are charted in **FIG. 5**, such that a health care professional can effectively monitor the diagnostic process.

[**0034**] **FIG. 6** depicts the appropriate diagnosing algorithm **600** to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of muscle weakness as shown in **FIG. 1**. Different clinical algorithms are available for symptoms of muscle weakness, as compared to symptoms of general fatigue. According to the algorithm in **FIG. 6**, a health care professional examines the muscle groups of a patient with these symptoms **610** to determine if the patient exhibits demonstrable muscle weakness. If no demonstrable muscle weakness is identified **620**, then the patient is evaluated for malaise/fatigue **625** in accordance with the steps of **FIG. 10**.

If demonstrable muscle weakness is identified **630**, then the health care professional must rule out the possible effects of an arthritic or injured joint **640** or a sensory abnormality **645** as the cause of the apparent muscle weakness before proceeding to further evaluation. If there are no sensory abnormalities or joint weakness, then the stretch reflexes of the patient are evaluated **650**. If the evaluation shows decreased proximal stretch reflexes or predominant proximal weakness, then the patient is evaluated for muscle myopathy **670** in accordance with the steps of **FIG. 8**. The steps of algorithm **600** are charted in **FIG. 7**, such that a health care professional can effectively monitor the diagnostic process.

[**0035**] **FIG. 8** depicts the appropriate diagnosing algorithm **800** to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of muscle weakness/wasting or muscle cramping as shown in **FIG. 1**. According to the algorithm in **FIG. 8**, a patient with these symptoms is evaluated by clinical exam **810**, to determine the cause or origin of the muscle cramping of the demonstrated muscle weakness beyond the evaluation of **FIG. 6**. The health care professional must rule out the possible symptoms of muscle myopathy unrelated to dialysis before proceeding to further evaluation of possible carnitine deficiency.

[**0036**] Reviewing the steps in **FIG. 8**, possible acquired infection, inflammation, or malignancy **820** must be considered and, if present, treated using skills known in the art. Also, thyroid disorders **830** must be considered and, if present, treated accordingly. Furthermore, Cushing's disease **840** must be considered and, if present, treated with appropriate drug therapy or surgery as determined by the health care professional. Additionally, alcohol or drug use **850** may be the cause of the muscle myopathy symptoms, and, if so, the use of the offending agent must be eliminated. Uncontrolled hyperparathyroidism **860** is yet another possible cause that must be considered, and if present, treated appropriately with suppressive therapy or surgery. Also, osteomalacia **870** must be considered and, if present, treated with appropriate treatment such as vitamin D therapy. Moreover, aluminum toxicity **880** must be considered as a possible cause of muscle myopathy, and, if present, treated with appropriate means such as chelation therapy. Finally, if none of the previous etiologies are present, a carnitine deficiency **890** is a likely cause of muscle myopathy symptoms. As shown in **FIG. 8**, the health care professional will evaluate the patient's pre-dialysis plasma carnitine concentration **891**, determine if the pre-dialysis carnitine level is below normal **892**, and, if so, initiate levocarnitine injection therapy **893**. The steps of algorithm **800** are charted in **FIG. 9**, such that a health care professional can effectively monitor the diagnostic process.

[**0037**] **FIG. 10** depicts the appropriate diagnosing algorithm **1000** to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of protein catabolism or lack of energy as shown in **FIG. 1**. According to the algorithm in **FIG. 10**, a patient with these symptoms is evaluated by clinical exam **1010** to rule out a non-renal illness or condition as the cause or origin of the symptoms before evaluating factors specific to dialysis **1020**. The health care professional must consider inadequate dialysis **1030** and incompletely compensated anemia **1040** as possible causes of a patient's symptoms of malaise or fatigue. If both inadequate dialysis **1030** and incompletely

compensated anemia **1040** are excluded as possible causes **1060**, then the health care professional will evaluate the patient's pre-dialysis plasma carnitine concentration **1061**, determine if the pre-dialysis carnitine level is below normal **1062**, and, if so, initiate levocarnitine injection therapy **1063**.

[0038] Still referring to **FIG. 10**, if inadequate dialysis **1030** is a possible cause of the malaise or fatigue symptoms, the dialysis must be adjusted **1050**. If the adjusted dialysis **1050** relieves the symptoms of malaise or fatigue **1051**, then the successful measures should be continued **1052** and the patient re-evaluated as necessary **1080**. If the adjusted dialysis **1050** does not relieve the symptoms of malaise or fatigue **1055**, then the health care professional will evaluate the patient's pre-dialysis plasma carnitine concentration **1061**, determine if the pre-dialysis carnitine level is below normal **1062**, and, if so, initiate levocarnitine injection therapy **1063**. If incompletely compensated anemia **1040** is a possible cause of the malaise or fatigue symptoms, the EPO dose must be adjusted **1070**. If the adjusted EPO dose **1070** produces an adequate response (e.g., relieves the symptoms of malaise or fatigue) **1071**, then the successful measures should be continued **1072** and the patient reevaluated as necessary **1080**. If the adjusted EPO dose **1070** produces an inadequate response (e.g., does not relieve the symptoms of malaise or fatigue) **1073**, then the patient is evaluated for delayed/diminished response to EPO **1074** in accordance with the steps of **FIG. 12**. The steps of algorithm **1000** are charted in **FIG. 11**, such that a health care professional can effectively monitor the diagnostic process.

[0039] **FIG. 12** depicts the appropriate diagnosing algorithm **1200** to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of delayed or diminished response to EPO as shown in **FIG. 1**. According to the algorithm in **FIG. 12**, a patient is evaluated by clinical exam **1210** to rule out symptoms of delayed or diminished response to EPO that are unrelated to reduced carnitine levels before proceeding to further evaluation of possible carnitine deficiency.

[0040] Reviewing the steps in **FIG. 12**, iron deficiency **1220** must be considered and, if present, treated accordingly with, for example, supplemental iron. Also, possible acquired infection, inflammation, or malignancy **1230** must be considered and, if present, treated using skills known in the art. Furthermore, occult blood loss **1240** must be considered and, if present, the source of bleeding identified and treated with appropriate measures. Additionally, an underlying hematologic disease **1250**, such as thalassemia, refractory anemia, or other myelodysplastic disorders, may be the cause of the delayed or diminished response to EPO symptoms, and, if so, these diseases must be treated accordingly. Hemolysis **1260** is yet another possibly cause that must be considered, and if present, treated appropriately. Also, osteitis fibrosa cystica **1270** must be considered and, if present, treated with standard medical treatment. Moreover, aluminum intoxication **1280** must be considered as a possible cause of delayed or diminished response to EPO, and, if present, treated with appropriate means such as chelation therapy. Likewise, vitamin deficiencies **1285**, such as a lack of folic acid or vitamin B12, must be considered and, if present, treated accordingly with, for example, supplemental vitamins. Finally, if none of the previous indications are present, a carnitine deficiency **1290** is a likely cause of the

delayed or diminished response to EPO symptoms. As shown in **FIG. 12**, the health care professional will evaluate the patient's pre-dialysis plasma carnitine concentration **1291**, determine if the pre-dialysis carnitine level is below normal **1292**, and, if so, initiate levocarnitine injection therapy **1293**. The steps of algorithm **1200** are charted in **FIG. 13**, such that a health care professional can effectively monitor the diagnostic process.

[0041] **FIG. 14a** depicts the appropriate diagnosing algorithm **1400** to determine a patient's status and potential suitability for levocarnitine administration given the symptoms of dialysis related hypotension as shown in **FIG. 1**. According to the algorithm in **FIG. 14a**, a patient with these symptoms is evaluated by clinical exam **1410** for hypotension (e.g., sustained pre- or post-dialysis blood pressure less than 110 systolic or a decline in systolic blood pressure greater than 20 mm Hg which is either (a) associated with symptoms referable to hypotension, (b) requires intervention, or (c) compromises the intended dialysis prescription). From the examination **1410**, the health care professional determines whether the symptoms of hypotension are an acute episode **1420** or a chronic sustained condition **1450**. If the symptoms indicate an acute episode **1420**, then the health care professional initiates immediate corrective measures **1430**, such as repositioning said the patient into a Trendelenberg position, using IV fluid, decreasing or zeroing ultrafiltration rate (UFR) for the dialysis, considering use of oxygen, or considering the causes as in chronic sustained step **1450**. If the acute episode is not indicative of repetitive intradialytic hypotension, then intermittent acute treatment is continued **1445**. If the acute episode **1420** is indicative of repetitive intradialytic hypotension, then the health care professional must apply initial measures **1490** known to those skilled in the art as appropriate to the patient's situation, such as using a machine with ultrafiltration control, counseling patient to limit salt, aiming for intradialytic weight gain greater than 1 kg/d, re-evaluating dry weight and avoiding ultrafiltration below that value, using bicarbonate containing dialysate, and reviewing anemia treatment to see if improvements are necessary.

[0042] Continuing with **FIG. 14a**, if the symptoms are a chronic sustained condition **1440**, then the health care professional conducts a series of diagnostics **1450**, including electrocardiograms, evaluating for pericarditis, testing for congestive heart failure (CHF), considering possible valve problems, checking for ischemia, testing for infections, considering hematologic disorders, and evaluating for sepsis. If the diagnostics **1450** indicate persistent hypotension despite treatment of the underlying condition, then the health care professional must apply initial measures **1490** known to those skilled in the art as appropriate to the patient's situation. If the diagnostics **1450** indicate hypotension is related to the underlying condition, then the treatment of the underlying condition is continued **1475**, as necessary.

[0043] Referring to **FIG. 14b**, if the initial measures of step **1490** show inadequate improvement **1500**, the health care professional must apply additional measures **1510** according to his skill in the art and patient tolerance, such as withholding pre hemodialysis blood pressure medications, lowering dialysate temperature, omitting specific foods or glucose containing solutions, adjusting dialysate calcium, conducting high sodium or sodium gradient dialysis, using a blood volume monitor, adding additional treatment(s) per

week, and conducting sequential treatments. If these measures provide adequate improvement **1560** for the patient's conditions, then the successful measures should be continued **1570** and the patient reevaluated as necessary **1580**. If the additional measures of step **1510** show inadequate improvement **1520**, then health care professional must consider pharmacological therapy measures **1530**, such as Midodrine, Caffeine, Florinef, and Pseudoephedrine. If these pharmacological therapy measures provide adequate improvement **1560** for the patient's conditions, then the successful measures should be continued **1570** and the patient re-evaluated as necessary **1580**. If the measures of step **1530** show inadequate improvement **1540**, then the health care professional will evaluate the patient's pre-dialysis plasma carnitine concentration **1541**, determine if the pre-dialysis carnitine level is below normal **1542**, and, if so, initiate levocarnitine injection therapy **1543**. If the levocarnitine injection therapy **1543** provides adequate improvement **1560** for the patient's conditions, then the successful measures should be continued **1570** and the patient re-evaluated as necessary **1580**. If the measures of step **1543** show inadequate improvement **1545**, then the health care professional will reevaluate the underlying pathology, consider Parkinsons Disease, and evaluate transplant urgency. The steps of algorithm **1400** are charted in **FIG. 15**, such that a health care professional can effectively monitor the diagnostic process.

[**0044**] ESRD patients with the above conditions, as well as those with erythropoietin resistance, who are unresponsive to standard therapy are prescribed levocarnitine, generally 20 mg/kg/dialysis session. Once patients are on the therapy, the clinical outcome monitoring tool, as shown in **FIG. 16**, is used to evaluate a patient's response. The monitoring tool provides a map of key indicators from laboratory test results, patient intake of medication and supplements, and patient assessments. These key indicators are measured preferably before initiating levocarnitine injection therapy, after one month, after three months and after six months from the start of therapy.

[**0045**] As will be readily understood by one of ordinary skill in the art, the present invention can preferably be automated with software as is known in the art such that a computing device can be adapted to interact with health care professionals via a software interface and assist such professionals in appropriately utilizing and applying the algorithms and tools disclosed herein. In this manner, the health care professional can be automatically prompted to consider appropriate follow-up questions or considerations, or prompted to take appropriate diagnostic actions based upon the application of the algorithms and tools herein disclosed to the feedback data provided by the user. Once the health care professional has provided sufficient feedback data to the software automating the present invention, he or she can then be provided with a well-defined clinical picture that helps him or her identify a recommended course of action with respect to therapeutic carnitine administration.

[**0046**] Various modifications of the embodiments herein disclosed will be readily apparent to one skilled in the art after reading the above. For example, algorithms and tools automated by software according to alternative embodiments of the present invention may be adapted to store patient data and thereby automatically create patient logs and charts based upon the feedback data entered by the users

of tools. Any and all such modifications are intended to be covered by the application as claimed.

What is claimed is:

1. A method for diagnosing clinical conditions common in a hemodialysis patient comprising the steps of:

(a) associating symptoms of said patient with a symptom category selected from the group of secondary cardiomyopathy, dialysis related hypotension, cardiac arrhythmia, muscle wasting or weakness, muscle cramping, protein catabolism, lack of energy, and delayed or diminished response to erythropoietin; and

(b) applying an appropriate clinical diagnosis algorithm for said symptom category.

2. The method of claim 1 wherein said step of applying an appropriate clinical diagnosis algorithm includes use of monitoring tools to assist healthcare professionals in implementing the clinical algorithms.

3. The method of claim 2 wherein said monitoring tools are incorporated into a computer program and accessed via computer that records the data for individual patents.

4. The method of claim 1 further comprising the step of evaluating said patient's pre-dialysis plasma carnitine concentration, determining if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy.

5. The method of claim 4 further comprising the step of monitoring said levocarnitine injection therapy with a monitoring tool that provides a map of key indicators from laboratory test results, patient intake of medication and supplements, and patient assessments.

6. The method of claim 1 wherein said appropriate clinical algorithm for said symptom categories of secondary cardiomyopathy and hypotension during dialysis comprises the steps of:

(a) evaluating said patient by a clinical exam, wherein a types of treatments are considered that include controlling fluid volume during dialysis, controlling blood pressure, treating arrhythmia, adjusting hematocrit (HCT)/hemoglobin (HGB), and reducing after-load;

(b) if any of said types of treatments provide adequate improvement upon evaluation, then continuing the improving treatment and monitoring the patient in subsequent clinical exams; and

(c) if none of said types of treatments provide adequate improvement upon evaluation, then evaluating said patient's pre-dialysis plasma carnitine concentration, determining if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy.

7. The method of claim 6 wherein the results of each completed step are recorded in a charting tool that contains the steps of the appropriate clinical algorithm for said symptom categories of secondary cardiomyopathy and hypotension during dialysis.

8. The method of claim 1 wherein said appropriate clinical algorithm for said symptom category cardiac arrhythmia comprises the steps of:

- (a) evaluating said patient by a clinical exam;
- (b) determining whether the symptoms are related to an underlying medical condition or to effects from dialysis;
- (c) if the arrhythmia symptoms are related to an underlying medical condition, then providing appropriate treatment relating to the underlying condition, and monitoring the patient in subsequent clinical exams;
- (d) if the arrhythmia symptoms are related to dialysis treatment, adjusting the dialysis process using techniques from the group comprised of optimizing the volume status, adjusting the dialysate if possible, prolonging dialysis to moderate fluid shifts, and administering oxygen during dialysis;
- (e) if any of said adjustments to the dialysis process control the arrhythmia symptoms upon evaluation, then continuing the improving adjustments and monitoring the patient in subsequent clinical exams; and
- (f) if any of said adjustments to the dialysis process do not control the arrhythmia symptoms, then evaluating the patient's pre-dialysis plasma carnitine concentration, determining if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy.

9. The method of claim 8 wherein the results of each completed step are recorded in a charting tool that contains the steps of the appropriate clinical algorithm for said symptom category cardiac arrhythmia.

10. The method of claim 1 wherein said appropriate clinical algorithm for said symptom categories of muscle wasting or weakness and muscle cramping comprises the steps of:

- (a) examining the muscle groups of a patient with said muscle wasting or weakness and muscle cramping symptoms to determine if the patient exhibits demonstrable muscle weakness;
- (b) if no demonstrable muscle weakness is identified, then evaluating said patient for malaise/fatigue;
- (c) if demonstrable muscle weakness is identified, then examining for possible effects of an arthritic joint, an injured joint, or a sensory abnormality as the cause of the apparent muscle weakness;
- (d) if arthritic joints, injured joints, or sensory abnormalities are found, then providing appropriate treatment relating to the underlying condition;
- (e) if no sensory abnormalities, arthritic joints, or injured joints are found, then evaluating the stretch reflexes of said patient; and
- (f) if the evaluation shows decreased proximal stretch reflexes or predominant proximal weakness, then evaluating said patient for muscle myopathy.

11. The method of claim 10 wherein the results of each completed step are recorded in a charting tool that contains the steps of the appropriate clinical algorithm for said symptom categories of muscle wasting or weakness and muscle cramping.

12. The method of claim 10 wherein said step of evaluating said patient for malaise/fatigue comprises the steps of:

- (a) evaluating a patient by clinical exam to rule out a non-renal illness or condition as the cause or origin of the malaise/fatigue symptoms;
  - (b) evaluating factors specific to dialysis, including inadequate dialysis and incompletely compensated anemia;
  - (c) if inadequate dialysis is a possible cause of the malaise or fatigue symptoms, adjusting the dialysis accordingly;
  - (d) if the adjusted dialysis relieves the symptoms of malaise or fatigue, then continuing the successful measures and re-evaluating the patient's condition as necessary;
  - (e) if the adjusted dialysis does not relieve the malaise/fatigue symptoms or if both inadequate dialysis and incompletely compensated anemia are excluded as possible causes of the malaise/fatigue symptoms, then evaluating the patient's pre-dialysis plasma carnitine concentration, determining if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy;
  - (f) if incompletely compensated anemia is a possible cause of the malaise or fatigue symptoms, then adjusting the erythropoietin dose accordingly;
  - (g) if the adjusted erythropoietin dose relieves the malaise/fatigue symptoms, then continuing the successful erythropoietin dosage and reevaluating the patient as necessary;
  - (h) if the adjusted erythropoietin dose does not relieve the malaise/fatigue symptoms, then evaluating the patient for delayed/diminished response to erythropoietin.
13. The method of claim 12 wherein the results of each completed step are recorded in a charting tool that contains the steps of the appropriate clinical algorithm evaluating said patient for malaise/fatigue.
14. The method of claim 12 wherein said appropriate clinical algorithm for said symptom category delayed or diminished response to erythropoietin comprises the steps of:
- (a) evaluating said patient for iron deficiency and, if present, treating accordingly;
  - (b) evaluating said patient for possible acquired infection, inflammation, or malignancy and, if present, treating said infection, inflammation, or malignancy using skills known in the art;
  - (c) evaluating said patient for occult blood loss and, if present, identifying the source of bleeding and treating with appropriate measures;
  - (d) evaluating said patient for an underlying hematologic disease, such as thalassemia, refractory anemia, or other myelodysplastic disorders and, if present, treating with appropriate measures;
  - (e) evaluating said patient for hemolysis and, if present, treating the hemolysis appropriately;
  - (f) evaluating said patient for osteitis fibrosa cystica and, if present, treating with standard medical treatment;
  - (g) evaluating said patient for aluminum intoxication and, if present, treating with appropriate means, such as chelation therapy;

- (h) evaluating said patient for vitamin deficiencies and, if present, treating accordingly; and
- (i) if none of the previous indications are present, evaluating the patient's pre-dialysis plasma carnitine concentration, determining if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy.

15. The method of claim 14 wherein the results of each completed step are recorded in a charting tool that contains the steps of said appropriate clinical algorithm for said symptom category delayed or diminished response to erythropoietin.

16. The method of claim 10 wherein said step of evaluating said patient for muscle myopathy comprises the steps of:

- (a) evaluating for possible acquired infection, inflammation, or malignancy and, if present, treating said acquired infection, inflammation, or malignancy;
- (b) evaluating for possible thyroid disorders and, if present, treating said thyroid disorder;
- (c) evaluating for Cushing's disease and, if present, treating said Cushing disease with drug therapy or surgery;
- (d) evaluating for alcohol or drug use, and, if present, eliminating the alcohol or drug use;
- (e) evaluating for uncontrolled hyperparathyroidism, and, if present, treating said hyperparathyroidism with suppressive therapy or surgery;
- (f) evaluating for osteomalacia and, if present, treating with appropriate treatment;
- (g) evaluating for aluminum toxicity, and, if present, treating said aluminum toxicity with appropriate means; and
- (h) if none of the previous etiologies are present, evaluating the patient's pre-dialysis plasma carnitine concentration, determine if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy.

17. The method of claim 16 wherein the results of each completed step are recorded in a charting tool that contains the steps of the appropriate clinical algorithm evaluating said patient for muscle myopathy.

18. The method of claim 1, wherein said appropriate clinical algorithm for said symptom category dialysis related hypotension comprises the steps of:

- (a) evaluating said patient by a clinical exam;
- (b) determining whether symptoms of hypotension are an acute episode **1420** or a chronic sustained condition;
- (c) if the symptoms indicate an acute episode, then initiating immediate corrective measures;
- (d) if the acute episode is not indicative of repetitive intradialytic hypotension, then continuing intermittent acute treatment;

- (e) if the acute episode is indicative of repetitive intradialytic hypotension, then the applying one or more initial measures selected from the group of: using a machine with ultrafiltration control, counseling patient to limit salt, aiming for intradialytic weight gain greater than 1 kg/d, re-evaluating dry weight and avoiding ultrafiltration below the dry weight value, using bicarbonate containing dialysate, and reviewing anemia treatment to see if improvements are necessary;

- (f) if the symptoms are a chronic sustained condition, then conducting one or more diagnostic selected from the group of: electrocardiograms, evaluating for pericarditis, testing for congestive heart failure (CHF), considering possible valve problems, checking for ischemia, testing for infections, considering hematologic disorders, and evaluating for sepsis;

- (g) if any of said diagnostics indicates persistent hypotension despite treatment of the underlying condition, then applying said initial measures;

- (h) if any of said diagnostics indicate hypotension is related to an underlying condition, then continuing treatment of the underlying condition, as necessary;

- (i) if said initial measure provide inadequate improvement, applying one or more additional measures selected from the group of: withholding pre hemodialysis blood pressure medications, lowering dialysate temperature, omitting specific foods or glucose containing solutions, adjusting dialysate calcium, conducting high sodium or sodium gradient dialysis, using a blood volume monitor, adding additional treatment(s) per week, and conducting sequential treatments;

- (j) if said additional measures provide adequate improvement for the patient's conditions, then continuing the successful measures and reevaluating the patient as necessary;

- (k) if said additional measures of step provide inadequate improvement, then considering one or more pharmacological therapy measures selected from the group of: Midodrine, Caffeine, Florinef, and Pseudoephedrine.

- (l) if said pharmacological therapy measures provide adequate improvement for said patient's conditions, then continuing the successful measures and reevaluating the patient as necessary; and

- (m) if said pharmacological therapy measures provide inadequate improvement, then evaluating the patient's pre-dialysis plasma carnitine concentration, determine if the pre-dialysis carnitine level is below normal, and, if so, initiating levocarnitine injection therapy.

19. The method of claim 18 wherein the results of each completed step are recorded in a charting tool that contains the steps of the appropriate clinical algorithm evaluating said patient for dialysis related hypotension.

专利名称(译)	血液透析患者的诊断方法和利用这些方法改善静脉注射左卡尼汀治疗血液透析患者的方法		
公开(公告)号	<a href="#">US20030225162A1</a>	公开(公告)日	2003-12-04
申请号	US10/355263	申请日	2003-01-31
[标]申请(专利权)人(译)	SIGMA TAU PHARMLS		
申请(专利权)人(译)	SIGMA-TAU制药公司.		
[标]发明人	SCHREIBER BRIAN LEWIS VYONNE T		
发明人	SCHREIBER, BRIAN LEWIS, VYONNE T.		
IPC分类号	A61B5/00 A61K31/198 G01N33/48 G01N33/50 G06F19/00		
CPC分类号	A61K31/198 Y10S707/99943 G06F19/3456		
优先权	60/352505 2002-01-31 US		
其他公开文献	US7333894		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

所公开的发明涉及用于诊断血液透析患者中常见的临床病症的方法，该方法可能与血液透析引起的异常肉毒碱代谢有关。此外，本发明涉及用于监测和改善治疗性静脉内左卡尼汀对这些患者的给药的方法。已经针对可能与肉毒碱缺乏有关的终末期肾病 ( ESRD ) 患者中出现的临床症状开发了临床算法。还提供了用于帮助医疗保健专业人员实施临床算法的监视工具。

**Patient Diagnosis Tool**

**Summary**

