



US007321862B2

(12) **United States Patent**  
**Rosenfeld et al.**

(10) **Patent No.:** **US 7,321,862 B2**  
(45) **Date of Patent:** **\*Jan. 22, 2008**

(54) **SYSTEM AND METHOD FOR  
PATIENT-WORN MONITORING OF  
PATIENTS IN GEOGRAPHICALLY  
DISPERSED HEALTH CARE LOCATIONS**

(75) Inventors: **Brian A. Rosenfeld**, Baltimore, MD  
(US); **Michael Breslow**, Lutherville,  
MD (US)

(73) Assignee: **Visicu, Inc.**, Baltimore, MD (US)

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 282 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **11/268,706**

(22) Filed: **Nov. 7, 2005**

(65) **Prior Publication Data**

US 2006/0085227 A1 Apr. 20, 2006

**Related U.S. Application Data**

(63) Continuation-in-part of application No. 10/946,548,  
filed on Sep. 21, 2004, now Pat. No. 7,256,708, and  
a continuation-in-part of application No. 10/654,668,  
filed on Sep. 4, 2003, which is a continuation-in-part  
of application No. 09/443,072, filed on Nov. 18, 1999,  
now Pat. No. 6,804,656.

(60) Provisional application No. 60/141,520, filed on Jun.  
23, 1999.

(51) **Int. Cl.**  
**G06F 19/00** (2006.01)

(52) **U.S. Cl.** ..... **705/3; 705/2; 600/300**


(58) **Field of Classification Search** ..... **705/2-3;**  
**600/300**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,646,606	A	*	2/1972	Buxton et al.	600/483
4,365,199	A		12/1982	McNair	
4,489,387	A		12/1984	Lamb et al.	
4,731,725	A		3/1988	Suto et al.	
4,838,275	A		6/1989	Lee	
4,852,570	A		8/1989	Levine	
4,878,175	A		10/1989	Norden-Paul et al.	
5,255,187	A		10/1993	Sorensen	
5,321,800	A		6/1994	Lesser	
5,348,008	A		9/1994	Bornn et al.	
5,544,649	A		8/1996	David et al.	
5,544,661	A		8/1996	Davis et al.	
5,574,828	A		11/1996	Hayward et al.	
5,576,952	A		11/1996	Stutman et al.	
5,619,991	A		4/1997	Sloane	
5,678,562	A		10/1997	Sellers	
5,701,894	A		12/1997	Cherry et al.	
5,715,449	A		2/1998	Peters, Jr. et al.	
5,724,580	A		3/1998	Levin et al.	
5,729,204	A		3/1998	Fackler et al.	
5,772,585	A		6/1998	Lavin	
5,812,983	A		9/1998	Kumagai	
5,822,544	A		10/1998	Chaco et al.	
5,823,948	A		10/1998	Ross, Jr. et al.	
5,832,450	A		11/1998	Myers et al.	
5,839,438	A		11/1998	Graettinger et al.	
5,842,978	A		12/1998	Levy	
5,867,821	A		2/1999	Ballantyne et al.	
5,868,669	A		2/1999	Iliff	
5,899,855	A		5/1999	Brown	
5,942,986	A		8/1999	Shabot et al.	
5,987,519	A		11/1999	Peifer et al.	
6,024,699	A		2/2000	Surwit et al.	
6,093,146	A		7/2000	Filangeri	
6,102,856	A		8/2000	Groff et al.	
6,154,668	A		11/2000	Pedersen et al.	
6,168,563	B1		1/2001	Brown	
6,215,403	B1		4/2001	Chan et al.	
6,225,901	B1		5/2001	Kail, IV	
6,230,142	B1		5/2001	Benigno et al.	
6,234,964	B1		5/2001	Iliff	
6,238,338	B1		5/2001	DeLuca et al.	
6,245,013	B1		6/2001	Minoz et al.	
6,254,536	B1		7/2001	Devito	
6,278,999	B1		8/2001	Knapp	



6,292,698 B1 9/2001 Duffin et al.  
 6,304,788 B1 10/2001 Eady et al.  
 6,315,719 B1 11/2001 Rode et al.  
 6,364,834 B1 4/2002 Reuss et al.  
 6,385,589 B1 5/2002 Trusheim et al.  
 6,398,728 B1 6/2002 Bardy  
 6,442,433 B1 8/2002 Linberg  
 6,533,724 B2 3/2003 McNair  
 6,741,264 B1 5/2004 Lesser  
 6,835,176 B2 12/2004 McNair  
 6,893,396 B2 5/2005 Schulze et al.  
 2002/0002473 A1 1/2002 Schrier et al.  
 2002/0177758 A1 11/2002 Schoenberg et al.  
 2002/0177759 A1 11/2002 Schoenberg et al.  
 2002/0187483 A1 12/2002 Hoffman et al.  
 2002/0193667 A1 12/2002 McNair  
 2003/0036687 A1 2/2003 Schoenberg et al.  
 2004/0030578 A1 2/2004 Cross et al.  
 2004/0063031 A1 4/2004 Gallucci et al.  
 2004/0078366 A1 4/2004 Crooks et al.  
 2004/0193451 A1 9/2004 McNair  
 2004/0197813 A1 10/2004 Hoffman et al.  
 2004/0199333 A1 10/2004 Hoffman et al.  
 2004/0225201 A1 11/2004 McNair  
 2004/0236604 A1 11/2004 McNair  
 2005/0027563 A1 2/2005 Fackler et al.  
 2005/0049891 A1 3/2005 Wilson  
 2005/0060191 A1 3/2005 Parkins et al.  
 2005/0075794 A1 4/2005 Hoffman et al.  
 2005/0075904 A1 4/2005 Wagner et al.  
 2005/0076060 A1 4/2005 Finn et al.  
 2005/0125098 A1 6/2005 Wang et al.  
 2005/0125256 A1 6/2005 Schoenberg et al.  
 2005/0206518 A1 9/2005 Welch et al.  
 2005/0228241 A1 10/2005 McNair  
 2005/0251418 A1 11/2005 Fox, Jr. et al.  
 2005/0267351 A1 12/2005 Humphrey et al.  
 2005/0283062 A1 12/2005 Hoffman et al.  
 2006/0031018 A1 2/2006 Bush et al.  
 2006/0036542 A1 2/2006 McNair

FOREIGN PATENT DOCUMENTS

WO WO 98/29790 7/1998  
 WO WO 99/13766 3/1999  
 WO 00/79466 A2 12/2000

OTHER PUBLICATIONS

Terry Ann Capuano et al. Remote Telemetry, Jul. 1995, Nursing Management, vol. 26, No. 7, p. 26.\*  
 Terry Ann Capuano, et al., Remote Telemetry, Nursing Management, Vol. 26, No. 7, Jul. 1995, p. 26.  
 Valeriy Nenov and John Klopp, Remote Access to Neurosurgical ICU Physiological Data using the World Wide web, health Care in the Information Age, 1996, pp. 242-249.  
 Betty L. Grundy, et al., Telemedicine in Critical Care: An Experiment in Health Care Delivery, JACEP, vol. 6, Oct. 1977, pp. 439-444.  
 Susan L. Mabry, et al., Integrated Medical Analysis System, Proceedings of the 1997 Winter Simulation Conference., 1997, pp. 1167-1168.  
 Simon M. Kaplan and Geraldine Fitzpatrick, Designing Support for Remote Intensive-Care Telehealth Using the Locales Framework, ACM, 1997, pp. 173-184.  
 Douglas A. Perednia, Telemedicine Technology and Clinical Applications, JAMA, vol. 6, Feb. 8, 1995, p.  
 Silvia Miksch, Artificial Intelligence for Decision Support: Needs, Possibilities, and Limitations in ICU, 10th Postgraduate Course in Critical Care Medicine A.P.I.C.E. '95, Springer 1995 pp. 1-11.  
 Ho Sung Lee, et al., Remote Patient Monitoring Service through World-Wide Web, Proceedings—19th Int.  
 Silvia Miksch, Artificial Intelligence for Decision Support: Needs, Possibilities, and Limitations in ICU, 10th Postgraduate Course in Critical Care Medicine APICE '95, Springer, 1995.

Doctors use 'remote control' to monitor ICU patients, CNN.com. technology=computing, Aug. 21, 2000, <http://www.cnn.com/2000/TECH/computing/08/21/icu.t.t/>.  
 Finding Value in Intensive Care, From Afar, The New York Times on the Web, Jul. 27, 1999, [www.visicu.com/companynews/0799\\_nytimes.htm](http://www.visicu.com/companynews/0799_nytimes.htm).  
 Remote Monitoring of ICU Patients Lowers Mortality Rates, Complications, Johns Hopkins Newsrelease, Mar. 20, 2001, <http://www.newswise.com/articles/2001/3/ICU.JHM.html>.  
 Brian A. Rosenfeld, M.D., FCCM, FCCP, et al., Intensive care unit telemedicine: Alternate paradigm for providing continuous intensivist care, Critical Care Medicine, vol. 28, No. 12, 2000 p. 3925.  
 Benjamin Berg, Dale Vincent, and Donald Hudson, Remote Clinical Care Consultation: Telehealth Projection of Clinical Specialty Expertise, Tripler Army Medical Center, Honolulu.  
 Xin Li, et al., A World Wide Web Telemedicine System, SPIE vol. 2711 p. 427-439.  
 Guidelines for Intensive Care Unit Design, Critical Care Medicine, Mar. 1995; 23(3):582-588.  
 Michael Breslow, et al., Effect of a Multiple-Site Intensive Care Unit Telemedicine Program on clinical and Economic Outcomes: An Alternative Paradigm for Intensivist Staffing, Critical Care Medicine 2004 vol. 32, No. 1.  
 Richard Brill, et al., Critical care Delivery in the Intensive Care Unit: Defining Clinical Roles and the Best Practice Model, Critical Care Medicine 2001 vol. 29, No. 10.  
 M. Michael Shabot, et al., Decision Support Systems in Critical Care, 1994, Springer-Verlag Publishing, New York.  
 Rosenfeld, et al. Intensive care unit telemedicine: alternate paradigm for providing continuous intensivist care, Dec. 28, 2000, [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov).  
 Definitions of Intensive Care Unit (ICU) on the Web, Apr. 2004, [www.google.com](http://www.google.com) and other websites.  
 Grundy, Betty Lou; Jones, Paul; Lovitt, Ann; "Telemedicine in critical care: Problems in design, implementation and assessment" Jul. 1982. Critical Care Medicine vol. 10, No. 7.  
 Tsien, C.L. and Fackler, J.C., "Poor prognosis for existing monitors in the intensive care unit," Critical Care Medical Journal, vol. 25, No. 4, (1997) (p. 614-619).  
 Tsien, C.L. "Reducing False Alarms in the Intensive Care Unit: A Systematic Comparison of Four Algorithms," Proceedings Annual AMIA Fall Symposium (1997).  
 Kohane, I. S. and Halmowitz, I.J., "Hypothesis-Driven Data Abstraction with Trend Templates," Proceedings Annual AMIA Symposium on Computer Applications in Medical Care (1994), (p. 444-448).  
 Hetherington, Laurel Traynowicz; "High tech meets high touch: telemedicine's contribution to patient wellness"; Spring, 1998; Nursing Administration Quarterly, vol. 22, No. 3.  
 Tsien, Christine L., "TrendFinder: Automated Detection of Alarmable Trends", Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Massachusetts; Jun. 2000.  
 Hosseinzadeh, Abolfazl, "A Rule-Based System for Vital Sign Monitoring in Intensive Care", Department of Electrical Engineering, McGill University, Montreal; Nov. 1993.  
 Aukburg, S.J. et al., "Automation of Physiological Data Presentation and Alarms in the Post Anesthesia Care Unit." In Symposium on Computer Applications in Medical Care, Nov. 5-8, 1989, Washington, DC; pp. 580-582.  
 Benis, A. M. et al., "Improved Detection of Adverse Cardiovascular Trends with the Use of a Two-Variable Computer Alarm," *Critical Care Medicine*, vol. 8, No. 2, Jun. 1980: 341-344.  
 Bierman, M. I. et al., "Pulse Oximetry in the Postoperative Care of Cardiac Surgical Patients; A Randomized Controlled Trial," *Chest*, vol. 102, No. 5, Nov. 1992: 1367-1370.  
 Bradshaw, K. E., "Computerized Alerting System Warns of Life-Threatening Events." In Symposium on Computer Application in Medical Care, Oct. 25-26, 1986, Washington, DC; pp. 403-409.  
 Chizeck, H. J., "Modeling, Simulation and Control in a Data Rich Environment." In Symposium on Computer Applications in Medical Care. Oct. 25-26, 1986, Washington, DC; pp. 65-69.  
 Coiera, E., "Intelligent Monitoring and Control of Dynamic Physiological Systems." *Artificial Intelligence in Medicine*, vol. 5, 1993: pp. 1-8.

- Colvin, J. R. et al., "Microcomputer-Controlled Administration of Vasodilators Following Cardiac Surgery: Technical Considerations." *J. Cardiothoracic Anesthesia*, vol. 3, No. 1, Feb. 1989: pp. 10-15.
- Coplin, W. M. et al., "Accuracy of Continuous Jugular Bulb Oximetry in the Intensive Care Unit." *Neurosurgery*, vol. 42, No. 3, Mar. 1998: 533-540.
- Crew, A. D. et al., "Preliminary Clinical Trials of a Computer-Based Cardiac Arrest Alarm." *Intensive Care Med*, vol. 17, 1991: 359-364.
- Garfinkel, D. et al., "PONI: An Intelligent Alarm System for Respiratory and Circulation Management in the Operating Rooms." In Symposium on Computer Applications in Medical Care, Nov. 6-9, 1988, Washington, DC; pp. 13-17.
- Garfinkel D. et al., "Patient Monitoring in the Operating Room: Validation of Instrument Reading by Artificial Intelligence Methods." In Symposium on Computer Applications in Medical Care, Nov. 5-8, 1989, Washington, Dc; pp. 575-579.
- Guedes de Oliveira, P. et al., "The Role of Computer Based Techniques in Patient Monitoring: Technical Note." *Acta Neurochir*, vol. 55, 1992 (Suppl.): 18-20.
- Hahnel, J. et al., "Can a Clinician Predict the Technical Equipement a Patient will Need During Intensive Care Unit Treatment? An Approach to Standardize and Redesign the Intensive Care Unit Workstation." *J Clinical Monitoring*, vol. 8, No. 1, Jan. 1992: 1-6.
- Hall, G. L. & P.B. Colditz, "Continuous Physiological Monitoring: An Integrated System for Use in Neonatal Intensive Care." *Australian Physical & Engineering Sciences in Medicine*, vol. 18, No. 3, 1995; 139-142.
- Hayes-Roth, B. et al., "Guardian: An Experimental System for Intelligent ICU Monitoring." In Symposium on Computer Applications in Medical Care, Nov. 5-9, 1994, Washington, DC; p. 1004.
- Irazuzta, Jose, "Monitoring in Pediatric Intensive Care." *Indian J. Pediatrics*, vol. 60, 1993: 55-65.
- Jans, R. et al., "A Low Cost ECG Central Station for Intensive Care." *Australian Physical & Engineering Sciences in Medicine*, vol. 13, No. 1, 1990: 31-35.
- Jastremski, M. et al., "A Model for Technology Assessment as Applied to Closed Loop Infusion Systems" *Critical Care Medicine*, vol. 23, No. 10, Oct. 1995: 1745-1755.
- Klass, M. A. & E. Y. Cheng, "Early Response to Pulse Oximetry Alarms with Telemetry." *J. Clinical Monitoring*, vol. 10, No. 3, May 1994: 178-180.
- Koski, E. M. J. et al., "A Knowledge-Based Alarm System for Monitoring Cardiac Operated Patients—Assessment of Clinical Performance." *International J Clinical Monitoring and Computing*, vol. 11, 1994: 79-83.
- Koski, E. M. J. et al., "Development of an Expert System for Haemodynamic Monitoring: Computerized Symbolism of On-Line Monitoring Data." *International J. Clinical Monitoring and Computing*, vol. 8, 1992: 289-293.
- Laffel, G. et al., "Using Control Charts to Analyze Serial Patient-Related Data." *Quality Management in Health Care*, vol. 3, No. 1, Fall 1994: 70-77.
- L'Esrange, P. R. et al., "A Microcomputer System for Physiological Data Collection and Analysis." *Australian Dental Journal*, vol. 38, No. 5, Oct. 1993: 400-405.
- M. de Beer, N. A. et al., "Clinical Evaluation of a Method for Automatic Detection and Removal of Artifacts in Auditory Evoked Potential Monitoring." *J Clinical Monitorin*, vol. 11, No. 6, Nov. 1995: 381-391.
- Makivirta, A. et al., "The Median Filter as a Preprocessor for a Patient Monitor Limit Alarm System in Intensive Care." *Computer Methods and Programs in Biomedicine*, vol. 34, No. 2/3, Feb./Mar. 1991: 139-144.
- Makivirta, A. & E. M. J. Koski, "Alarm-Inducing Variability in Cardiac Postoperative Data and the Effects of Prealarm Delay." *Critical Care Medicine*, vol. 8, No. 6, May 1994: 153-162.
- Martin, J. F., "Closed-Loop Control of Arterial Pressure During Cardiac Surgery." *J. Clinical Monitoring*, vol. 8, No. 3, Jul. 1992: 252-253.
- Meyer, C. "Visions of Tomorrow's ICU." *American J. Nursing*, Apr. 1993: 27-31.
- Nenov, V. I. et al., "Computer Applications in the Intensive Care Unit." *Neurosurgery Clinics of North America*, vol. 5, No. 4, Oct. 1994: 811-827.
- Nobel, J. J., "Physiologic Monitoring Systems, Acute Care." *Pediatric Emergency Care*, vol. 8, No. 4, Aug. 1992: 235-237.
- Orr, J. A. & Westenskow, D. R., "A Breathing Circuit Alarm System Based on Neural Networks." *J. Clinical Monitoring*, vol. 10, No. 2, Mar. 1994: 101-109.
- Pappert, D. et al., "Preliminary Evaluation of a New Continuous Intra-Arterial Blood Gas Monitoring Device." *Acta Anaesthesiologica Scandinavica*, Suppl. 107, vol. 39, 1995: 67-70.
- Rampil, I. J., "Intelligent Detection of Artifact." *The Automated-Anesthesia Record and Alarm Systems*, Chapter 17, 1987: 175-190.
- Runciman, W. B. et al., "The Pulse Oximeter: Applications and Limitations—An Analysis of 2000 Incident Reports." *Anaesthesia and Intensive Care*, vol. 21, No. 5, Oct. 1993: 543-550.
- Sailors, R. M., "A Model-Based Simulator for Testing Rule-Based Decision Support Systems for Mechanical Ventilation of ARDS Patients." In Symposium on Computer Applications in Medical Care, Nov. 5-9, 1994, Washington, DC; p. 1007.
- Sanklecha, M., "The Pulse Oximeter." *Indian J. Pediatrics*, vol. 60, No. 3, 1993: 469-470.
- Schnapp, L. M. & N. J. Cohen, "Pulse Oximetry; Uses and Abuses." *Chest*, vol. 98, No. 5, Nov. 1990: 1244-1250.
- Simpson, R. L., "Automating the ICU: Facing the Realities." *Nursing Management*, vol. 23, No. 3, Mar. 1992: 24-26.
- Sittig, D. F. & M. Factor, "Physiological Trend Detection and Artifact Rejection: A Parallel Implementation of a Multi-State Kalman Filtering Algorithm." In Symposium on Computer Applications in Medical Care, Nov. 5-8, 1989, Washington, DC; pp. 569-574.
- Stoodley, K. D. C. et al., "Problems in the Development of a Computerized Ward Monitoring System for a Pediatric Intensive Care Unit." *International J Clinical Monitoring and Computing*, vol. 8, 1992: 281-287.
- Sukavaara, T. et al., "A Knowledge-based Alarm System for Monitoring Cardiac Operated Patients—Technical Construction and Evaluation." *International J. Clinical Monitoring and Computing*, vol. 10, 1993: 117-126.
- Szaflarski, N. L., "Emerging Technology in Critical Care: Continuous Intra-Arterial Blood Gas Monitoring." *American J. Critical Care*, vol. 5, No. 1, Jan. 1996: 55-65.
- Uckun, S., "Intelligent Systems in Patient Monitoring and Therapy Management." *International J. Clinical Monitoring and Computing*, vol. 11, 1994: 241-253.
- Webb, R. K., "Medical Decision Making and Decision Analysis." *Anesthesia and Intensive Care*, vol. 16, No. 1, Feb. 1988: 107-109.
- Yien, H. et al., "Spectral Analysis of Systemic Arterial Pressure and Heart Rate Signals as a Prognostic Tool for the Prediction of Patient Outcome in the Intensive Care Unit." *Critical Care Medicine*, vol. 25, No. 2, 1997: 258-266.
- Tsien, Christine L. and James Fackler, "Poor Prognosis for Existing Monitors in the Intensive Care Unit" *Critical Care Medicine*, vol. 25, No. 4, 1997: 614-619.
- Tsien, Christine L., "Reducing False Alarm in the Intensive Care Unit: A Systematic Comparison of Algorithms" Proceedings *AMIA Symposium*, 1997. pp. 9-14 (unnumbered).
- Tsien, Christine L., "Reducing False Alarms in the Intensive Care Unit: A Systematic Comparison of Four Algorithms" Proceedings Annual *AMIA Fall Symposium* (1997), p. 894.
- Tsien, Christine L. and James C. Fackler "An Annotated Data Collection System to Support Intelligent Analysis of Intensive Care Unit Data." Proceedings of the Second International Symposium on Advances in Intelligent Data Analysis, Reasoning about Data; Aug. 4-6, 1997; X. Liu, P. R. Cohen, and M. R. Berthold, Eds.; Springer-Verlag, London, UK; pp. 111-121.
- Zhao, Ruilin, "A Model-Based Expert System for Interpretation of Hemodynamic Data from ICU Patients." Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology; May 18, 1997 (pp. 1-121).
- Angus, D.C., et al. "Caring for the critically ill patient. Current and projected workforce requirements for care of the critically ill and

- patients with pulmonary disease: can we meet the requirements of an aging population?" (abstract) *JAMA*. Dec. 6, 2000; 284(21):2762-70.
- Celi, Leo Anthony, et al. "The eICU: It's Not Just Telemedicine." *Crit. Care Med.* 2001, vol. 29, No. 8 (Suppl.).
- Hanson, C.W. 3rd, et al. "Effects of an organized critical care service on outcomes and resource utilization: a cohort study." (abstract) *Crit. Care Med.* Feb. 1999; 27(2):270-4.
- Manthous, C.A., et al. "Effects of a medical intensivist on patient care in a community teaching hospital." (abstract) *Mayo Clin. Proc.* May 1997; 72(5):391-9.
- Pronovost, P.J., et al. "Organizational characteristics of intensive care units related to outcomes of abdominal aortic surgery." (abstract) *JAMA*. Apr. 14, 1999; 281(14):1310-7.
- Pronovost, Peter J., et al. "Physician Staffing Patterns and Clinical Outcomes in Critically Ill Patients: A Systematic Review." *JAMA*. Nov. 6, 2002; 288(17):2151-62.
- "Remote Control." *Modern Healthcare*, Feb. 25, 2002 (4 pages).
- Breslow, Michael J., et al. "Technology Strategies to Improve ICU Practice." *Seminars in Anesthesia* 24: 59-70, 2005.
- Breslow, Michael J. "Remote ICU Care Programs; Current Status." *J. Crit. Care*: 22, 66-76, 2007.
- Cantrell, Mark. "Nurses Keep Watch from miles away or at the bedside." *Nursing Spectrum*, Jan. 2005 (4 pages).
- Editors: M. Michael Shabot and Reed M. Gardner, *Computers and Medicine: Decision Support Systems in Critical Care*, 1994, Springer-Verlag New York, Inc. New York.
- Gilad J. Kuperman, M.D. and Reed M. Gardner, Ph.D., *The Help System: A Snapshot in Time*, 1988, Dept. of Biophysics, LDS Hospital, Salt Lake City, Utah.
- Project Leaders: Benoit Dawant, Ph.D. and John A. Morris, Jr. M.D., *Vanderbilt University Simon Project Website*, 2004, Vanderbilt University, Nashville, Tennessee.
- Greg Borzo, *Web Technology: Coming Soon to a Hospital Near You*, *American Medical News*, Nov. 18, 1996, American Medical Association [www.amednews.com](http://www.amednews.com).
- Abstract: J.E. Gray, C. Safran, R.B. Davis, G. Pomilio-Weitzner, J.E. Stewart, L. Zaccagnini and D. Pursley, *Baby Care Link: Using the Internet and Telemedicine to Improve Care for High-risk Infants*, Dec. 2000, *Pediatrics*, vol. 106, No. 6, pp. 1318-1324.
- Abstract: Ray Duncan and Jeffrey J. Pomerance, *Computer Assistance in Delivery of Patient Care in a Neonatal Intensive Care Unit, The Use of Computers in Perinatal Medicine*, Chapter 19, pp. 337-351, 1982, Praeger Publishers, New York, NY.
- Abstract: Ray Duncan, MD, *Computer Assisted Care in the Neonatal Intensive Care Unit*, *Proceedings of the 17th Annual Symposium on Computer Applications in Medical Care*, Nov. 1993, p. 929, American Medical Informatics Association.
- Abstract: Metnitz PG, Laback P, Popow C, Laback O, Lenz K, Hiesmayr M, *Computer assisted data analysis in intensive care: the ICDEV project—development of a scientific database system for intensive care (Intensive Care Data Evaluation Project)*, *International Journal of Clinical Monitoring and Computing*, 1995, vol. 12, No. 3, pp. 147-159.
- Abstract: Paul H. Peristein, MD, Neil K. Edwards, MS, Harry D. Atherton, MS, James M. Sutherland, MD, *Computer Assisted Newborn Intensive Care*, *Pediatrics*, Apr. 1976, vol. 57, No. 4, pp. 494-501.
- Abstract: Edward H. Shortliffe, MD, PHD, *Computer Programs to Support Clinical Decision Making*, *JAMA*, Jul. 3, 1987, vol. 258, No. 1, pp. 61-66.
- Abstract: Merz U, Peschgens T, Budde R, Kretschmann F, Homchen H V, *Computer-assisted monitoring in the neonatal intensive care unit [German]*, *Klin Padiatr*, Nov./Dec. 1995, vol. 207, No. 6, pp. 331-333.
- Abstract: Charles Safran, MD, Francois Herrman, MD, David Rind, MD, Hollis B. Kowaloff, BA, Howard L. Bleich, MD, and Warner V. Slack, MD, *Computer-Based Support for Clinical Decision Making*, *M.D. Computing*, 1990, vol. 7, No. 5, pp. 319-322.
- Abstract: Reed M. Gardner, PHD, *Computerized Management of Intensive Care Patients*, *M.D. Computing*, 1986, vol. 3, No. 1, pp. 36-51.
- Abstract: F. John Lewis; Steven Deller; Michael Quinn; Benjamin Lee; Raymond Will; and John Raines, *Continuous Patient Monitoring with a Small Digital Computer*, *Computers and Biomedical Research*, 1972, vol. 5, pp. 411-428.
- Abstract: N. Fumai, C. Collet, M. Petroni, K. Roger, A. Lam, E. Saab, A. 'S. Malowany, F. A. Carenvale, R. D. Gottesman, *Database Design of an Intensive Care Unit Patient Data Management System*, *Proceedings of the Fourth Annual IEEE Symposium on Computer-Based Medical Systems*, May 12, 1991, pp. 78-85, IEEE Computer Society Press, Los Alamitos, CA.
- Abstract: George Hripcsak; Paul D. Clayton; Robert A. Jenders; James J. Cimino; and Stephen B. Johnson, *Design of a Clinical Event Monitor*, *Computers and Biomedical Research*, Jun. 1996, vol. 29, No. 3, pp. 194-221.
- Abstract: David M. Rind, MD; Roger Davis, SCD; and Charles Safran, MD, *Designing Studies of Computer-Based Alerts and Reminders*, *MD Computing*, 1995, vol. 12, No. 2, pp. 122-126.
- Abstract: Dwayne R. Westenkow, PHD, *Automating Patient Care with Closed-Loop Control*, *M.D. Computing*, 1986, vol. 3, No. 2, pp. 14-20.
- Abstract: Snowden S. Brownlee KG, Dear P R, *An expert system to assist neonatal intensive care*, *I Med Eng Technol Mar.-Apr. 1997*;21(2):67-73, vol. 21, No. 3, pp. 67-73.
- Abstract: A. Aifredo Morales, Engr., Ms , James Gray, MD, MS , Charles Safran, MD, *An Application Server Approach for Integration of Clinical Systems*, *Proceedings of the AMIA 1999 Annual Symposium*, 1999, AMIA.
- Abstract: Kang Wang, PHD; Isaac Kohane, MD, PHD; Karen L. Bradshaw, BS; James Facider, MD, *A Real Time Patient Monitoring System on the World Wide Web*, *Proceedings of the 1996 AMIA Annual Fall Symposium*, Nov. 1996, pp. 729-732, Hanley and Belfus, Inc.
- Abstract: Michael Factor, David H. Gelernter, Craig E. Kolb, Perry L. Miller and Dean F. Sittig, *Real-Time Data Fusion in the Intensive Care Unit*, *IEEE Computer*, Nov. 1991, pp. 45-53.
- Editor: Judy G. Ozbolt, Ph.D., *A Conference of the American Medical Informatics Association, Symposium on Computer Applications in Medical Care*, Nov. 5-9, 1994, Hanley & Belfus, Inc. Medical Publishers, Philadelphia, PA.
- W. Hsueh-Fen Young, Reed M. Gardner, Thomas D. East and Kristi Turner, *Computerized Ventilator Data Selection: Artifact Rejection and Data Reduction*, *Int'l Journal of Clinical Monitoring and Computing* 1997, 14: 165-176, Kluwer Academic Publishers, Netherlands.
- Randolph A. Miller, M.D. and Reed M. Gardner, Ph.D., *Summary Recommendations for Responsible Monitoring and Regulation of Clinical Software Systems*, *Annals of Internal Medicine*, Nov. 1997, vol. 127, No. 9.
- Reed M. Gardner, T. Allan Pryor and Homer R. Warner, *The HELP Hospital Information System: Update 1998*, *Intl Journal of Medical Informatics* 1999, vol. 54, pp. 169-182, Elsevier Science Ireland Ltd., Ireland.
- Martin Spikoff, *Systems Aid Rural Health Delivery*, *QIPhysician.com*, Sep. 2003.
- Abstract: Jerome P. Kassirer, MD, *The Next Transformation in the Delivery of Health Care (Editorial)*, *NEJM*, Jan. 5, 1995, vol. 332, No. 1, pp. 52-54.
- Abstract: Lorene S. Avila; M. Michael Shabot, *Keys to the successful implementation of an ICU patient data management system*, *International Journal of Clinical Monitoring and Computing*, 1988, vol. 5, pp. 15-25.
- Abstract: Reed M. Gardner, MD; M. Michael Shabot, MD, *Computerized ICU Data Management Pitfalls and Promises*, *International Journal of Clinical Monitoring and Computing*, 1990, vol. 7, pp. 99-105.
- Karl W. Thomas, M.D., Charles S. Dayton, B.S., R.Ph., and Michael W. Peterson, M.D., *Evolution of Internet-Based Clinical Decision Support Systems*, *Journal of Medical Internet Research* 1999, vol. 1, University of Iowa, Iowa City, Iowa.
- Abstract: C. J. McDonald, *Protocol-Based Computer Reminders, The Quality of Care and The Non-Perfectability of Man*, *The New England Journal of Medicine*, Dec. 9, 1976, vol. 295, No. 24, 1351-1355.

- Abstract: T.D. East, A.H. Morris, C.J. Wallace, T.P. Clemmer, J.F. Orme, Jr., L.K. Weaver, S. Henderson and D.F. Sittig, A Strategy for Development of Computerized Critical Care Decision Support Systems, *Intl Journal of Clinical Monitoring and Computing*, 1991-92, vol. 8, No. 4, 263-269.
- Dr. Ramana Reddy and Dr. V. "Juggy" Jagannathan, Secure Collaboration Technology for Rural Clinical Telemedicine, National Library of Medicine, Oct. 8, 1996 Press Release, West Virginia University, West Virginia.
- West Virginia University Research Corporation, Secure Collaboration Technology for Rural Clinical Telemedicine: Final Report, National Library of Medicine.
- Martin J. Tobin, M.D., Principles and Practice of Intensive Care Monitoring, 1998, McGraw-Hill Inc.,
- Peter J. Haug, Reed M. Gardner, and R. Scott Evans, "Hospital-Based Decision Support" in *Clinical Decision Support Systems: Theory and Practice*, ETA S. Berner [ed.], Springer-Verlag New York, Inc., New York, NY, pp. 77-103.
- Clement J. McDonald, M.D. and William M. Tierney, M.D., Computer-Stored Medical Records: Their Future Role in Medical Practice, *JAMA*, Jun. 17, 1988, pp. 3433-3440, vol. 259, No. 23.
- Gilad J. Kuperman, Reed M. Gardner, and T. Allan Pryor, HELP: A Dynamic Hospital Information System, 1991, Springer-Verlag New York, Inc., New York, NY.
- M. Michael Shabot, M.D., Mark Lobue, B.S., and Jeannie Chen, Pharm.D., Wireless Clinical Alerts for Physiologic, Laboratory and Medication Data, Department of Enterprise Information Services, Surgery and Pharmacy Cedars-Sinai Health System, Los Angeles, CA.
- Chaoxin Sima, Ravi Raman, Ramana Reddy, William Hunt and Sumitra Reddy, Vital Signs Services for Secure Telemedicine Applications, Concurrent Engineering Research Center, West Virginia University, Morgantown, WV.
- Dickey Seidlitz Johnson, Jane Ranzenberger, Ruth D. Herbert, Reed M. Gardner, and Terry P. Clemmer, A Computerized Alert Program for Acutely Ill Patients, *Journal of Nursing Administration*, Jun. 1980, pp. 26-35.
- Reed M. Gardner, Ph.D., Blair J. West, M.S., T. Allan Pryor, Ph.D., Keith G. Larsen, R.Ph., Homer R. Warner, M.D., Terry P. Clemmer, M.D., James F. Orme, Jr. M.D., Computer-Based ICU Data Acquisition as an Aid to Clinical Decision-Making, *Critical Care Medicine*, 1982, pp. 823-830, vol. 10, No. 12, The Williams & Wilkins Co.
- Reed M. Gardner and Terry P. Clemmer, Computerized Protocols Applied to Acute Patient Care, 1977, Mediad Inc., Tarrytown, NY.
- Karen E. Bradshaw, Reed M. Gardner, and T. Allan Pryor, Development of a Computerized Laboratory Alerting System, *Computers and Biomedical Research* 22, 575-587, 1989, Academic Press, Inc.
- Terry P. Clemmer and Reed M. Gardner, Medical Informatics in the Intensive Care Unit: State of the Art 1991, *International Journal of Clinical Monitoring and Computing* 8: 237-250, 1992, Kluwer Academic Publishers, Netherlands.
- Reed M. Gardner, Ph.D., David V. Ostler, and O. Hank Duffy, M.D., Computers in the Emergency Room, *Internal Medicine for the Specialist*, vol. 8, No. 3, Mar. 1987.
- Dean F. Sittig, Nathan L. Pace, Reed M. Gardner, Eduardo Beck, and Alan H. Morris, Implementation of a Computerized Patient Advice System Using the HELP Clinical Information System, *Computers and Biomedical Research* 22, 474-487, 1989, Academic Press Inc.
- P. D. Clayton, R. Scott Evans, T. Pryor, R. M. Gardner, P. J. Haug, O. B. Wigertz, and H. R. Warner, Bringing HELP to the Clinical Laboratory—Use of an Expert System to Provide Automatic Interpretation of Laboratory Data, *Ann Clin Biochem* 1987; 24: Supplement.
- D. F. Sittig, Ph.D., R. M. Gardner, Ph.D., N. L. Pace, M.D., M. Bombino, M. D., and A. H. Morris, M.D., Compas: A Computerized Patient Advice System to Direct Ventilatory Care, *Medical Informatics* 88: Computers in Clinical Medicine, Sep. 13-15, 1988, British Medical Informatics Society, London.
- Karen E. Bradshaw, Ph.D., Dean F. Sittig, Ph.D., Reed M. Gardner, Ph.D., T. Allan Pryor, Ph.D., and Marge Budd, M.S., Improving Efficiency and Quality in a Computerized ICU, 1988 SCAMC, Inc.
- Dean F. Sittig, Ph.D., C. Gregory Elliott, M.D., C. Jane Wallace, R.N., B.S.N., Polly Bailey, R.N., Reed M. Gardner, Ph.D., Computerized Screening for Identification of Adult Respiration Distress Syndrome (ARDS) Patients, 1988 SCAMC, Inc.
- R. Scott Evans, Ph.D., Reed M. Gardner, Ph.D., John P. Burke, M.D., Stanley L. Pestotnik, R.P.H., Robert A. Larsen, M.D., David C. Classen, M.D., and Paul D. Clayton, Ph.D., A Computerized Approach to Monitor Prophylactic Antibiotics, 1987, SCAMC, Inc.
- Susan Henderson, B.A., Thomas D. East, Ph.D., Alan H. Morris, M.D., Reed M. Gardner, Ph.D., Performance evaluation of computerized clinical protocols for management of arterial hypoxemia in ARDS patients, LDS Hospital, and University of Utah, Salt Lake City, UT.
- Thomas D. East, Ph.D., Susan Henderson, B.A., Alan H. Morris, M.D., Reed M. Gardner, Ph.D., Implementation Issues and Challenges for Computerized Clinical Protocols for Management of Mechanical Ventilation in ARDS Patients, LDS Hospital, Salt Lake City, UT.
- C. Gregory Elliott, M.D., Deon Simmons, R.R.T., C. Duwayne Schmidt, M.D., Kip Enger, B.S., C.R.T.T., Loren Greenway, B.S., R.R.T., and Reed M. Gardner, Ph.D., Computer-Assisted Medical Direction of Respiratory Care, *Respiratory Management*, vol. 19, No. 2.
- H. Keller and CH. Trendelenburg, Data Presentation Interpretation, *Clinical Biochemistry Principles, Methods, Applications*, Walter-deGruyter & Co., 1989.
- Reed M. Gardner, Ph.D., Karen W. Hollingsworth, R.N., M.S., C.C.R.N., ECG and Pressure Monitoring: How to Obtain Optimal Results, 295-305.
- Reed M. Gardner, Ph.D., Dean F. Sittig, M.S., Marge C. Budd, R.N., M.S., Computers in the Intensive Care Unit: Match or Mismatch?, 248-259.
- Emmanuel Furst, Ph.D., Cardiovascular Technology, *The Journal of Cardiovascular Nursing*, Nov. 1989, 68-78.
- Dean F. Sittig, Reed M. Gardner, Nathan L. Pace, Alan H. Morris, and Eduardo Beck, Computerized Management of Patient Care in a Complex, Controlled Clinical Trial in the Intensive Care Unit, *Computer Methods and Programs in Biomedicine* 30, 1989, 77-84.
- Karen E. Bradshaw, Ph.D., Dean F. Sittig, Ph.D., Reed M. Gardner, Ph.D., T. Allan Pryor, Ph.D., and Marge Budd, R.N., M.S., Computer-Based Data Entry for Nurses in the ICU, *Clinical Computing*, Nov. 1988.
- Robert A. Larsen, M.D., R. Scott Evans, Ph.D., John P. Burke, M.D., Stanley L. Pestotnik, R.Ph., Reed M. Gardner, Ph.D., David C. Classen, M.D., Improved Perioperative Antibiotic Use and Reduced Surgical Wound Infections Through Use of Computer Decision Analysis, *Computer Applications for Surgical Prophylaxis/Larsen et al.*
- R. M. Gardner, Computers in the ICU and Surgery—Keeping Real-Time Patient Records for Decision Making.
- Thomas D. East, Ph.D., Alan H. Morris, M.D., Terry Clemmer, M.D., James F. Orme, M.D., C. Jane Wallace, B.S.N., Susan Henderson, B.A., Dean F. Sittig, Ph.D., Reed M. Gardner, Ph.D., Development of Computerized Critical Care Protocols—A Strategy That Really Works!, 1990 LDS Hospital, Salt Lake City, UT.
- R. Scott Evans, Ph.D., John P. Burke, M.D., Stanley L. Pestotnik, R.Ph., David C. Classen, M.D., Ronald L. Menlove, Ph.D., and Reed M. Gardner, Ph.D., Prediction of Hospital Infections and Selection of Antibiotics Using an Automated Hospital Database, 1990, SCAMC, Inc. 663-667.
- Susan E. Henderson, B.A., Robert O. Crapo, M.D., Thomas D. East, Ph.D., Alan H. Morris, M.D., C. Jane Wallace, R.N., Reed M. Gardner, Ph.D., Computerized Clinical Protocols in an Intensive Care Unit: How Well are They Followed?, 1990, SCAMC, Inc., LDS Hospital, Salt Lake City, UT.
- Reed M. Gardner, PHD, Russell K. Hulse, RPH, MBA, Keith G. Larsen, RPH, Assessing The Effectiveness Of A Computerized Pharmacy System, 1990, SCAMC, Inc., 668-672.
- Reed M. Gardner, "Patient-Monitoring Systems", *Medical Informatics: Computer Applications in Health Care*, E.H. Shortliffe and L.E. Perrealt (eds.), G. Wiederhold and L.M. Fagan (assoc. eds.) (Reading, MA: Addison-Wesley, 1990).

- Reed M. Gardner, Olaf K. Golubjatnikov, R. Myron Laub, Julie T. Jacobson, and R. Scott Evans, Computer-Critiqued Blood Ordering Using the HELP System, *Computers and Biomaterial Research* 23, 514-528, 1990, Academic Press, Inc.
- Karen E. Tate, PH.D., Reed M. Gardner, PH.D., and Lindell K. Weaver, M.D., A Computerized Laboratory Alerting System, *Clinical Computing*, 1990, vol. 7, No. 5, 296-301.
- Dean F. Sittig, Reed M. Gardner, Alan H. Morris, and C. Jane Wallace, Clinical Evaluation of Computer-Based Respiratory Care Algorithms, *International Journal of Clinical Monitoring and Computing* 7, 1990, 177-185, Kluwer Academic Publishers, Netherlands.
- R. Scott Evans, Stanley L. Pestotnik, John P. Burke, Reed M. Gardner, Robert A. Larsen, and David C. Classen, Reducing Title Duration Of Prophylactic Antibiotic Use Through Computer Monitoring Of Surgical Patients, DICP, *The Annals of Pharmacotherapy*, Apr. 1990, vol. 24, 351-354, Harvey Whitney Books Company, Cincinnati, OH.
- Reed M. Gardner, and M. Michael Shabot, Computerized ICU Data Management: Pitfalls and Promises, *International Journal of Clinical Monitoring and Computing* 7: 99-105, 1990, Kluwer Academic Publishers, Netherlands.
- Stanley L. Pestotnik, R.Ph., R. Scott Evans, PH.D., John P. Burke, M.D., Reed M. Gardner, PH.D., David C. Classen, M.D., Therapeutic Antibiotic Monitoring: Surveillance Using a Computerized Expert System, *The American Journal of Medicine*, Jan. 1990, vol. 88, 43-48.
- Gil Kuperman, MD, Brent James, MD, MSTAT, Julie Jacobsen, MT (ASCP), Reed M. Gardner, PH.D., Continuous Quality Improvement Applies To Medical Care: Experiences At LDS Hospital, *Medical Decision Making*, Oct. - Dec. 1991, 60-65, vol. 11, No. 4.
- Susan Henderson, Robert O. Crapo, C. Jane Wallace, Thomas D. East, Alan H. Morris & Reed M. Gardner, Performance Of Computerized Protocols For The Management Of Arterial Oxygenation In An Intensive Care Unit, *International Journal of Clinical Monitoring and Computing* 8, 1992, 271-180, Kluwer Academic Publishers, Netherlands.
- Reed M. Gardner, Ph.D., William L. Hawley, Thomas D. East, Ph.D., Thomas A. Oniki, B.S., Hsueh-Fen W. Young, B.S., Real Time Data Acquisition: Experience With The Medical Information Bus (MIB), LDS Hospital, University of Utah, Salt Lake City, UT.
- Eric F. Lepage, MD, Reed M. Gardner, PH.D., R. Myron Laub, MD, Julie T. Jacobson, MT(ASCP), Assessing The Effectiveness Of A Computerized Blood Order Consultation System, *LDS Hospital*, 1992, 33-37, AMIA, Inc.
- R. Scott Evans, Ph.D., Stanley L. Pestotnik, R.Ph., David C. Classen, M.D., Sheron B. Bass, B.S.N. Ronald L. Menlove, Ph.D., Reed M. Gardner, Ph.D., and John P. Burke, M.D., Development Of A Computerized Adverse Drug Event Monitor, LDS Hospital and University of Utah, Salt Lake City, UT.
- E. Lepage, R. Traineau, Ph. Marchetti, M. Benbunan, R. M. Gardner, Development Of A Computerized Knowledge Based System Integrated To A Medical Workstation: Application To Blood Transfusion, *MEDINFO*, 1992, 585-590, Elsevier Science Publishers B. V.
- Reed M. Gardner, Ph.D. and R. Scott Evans, Ph.D., Computer-Assisted Quality Assurance, *Group Practice Journal*, May/Jun. 1992, 41(3), 8-11.
- Thomas D. East, Ph.D., W. Hsueh-Fen Young, M.S., and Reed M. Gardner, Ph.D., Digital Electronic Communication between ICU Ventilators and Computers and Printers, *Respiratory Care*, Sep. 1992, vol. 37 No. 9, 1113-1123.
- Reed M. Gardner, *Computers in Critical Care*, Wellcome Trends in Hospital Pharmacy, Jul. 1992.
- T. Allan Pryor, Reed M. Gardner and W. Clinton Day, Computer System for Research and Clinical Application to Medicine, AFIPS—Conference Proceedings, vol. 33, 1968, 809-816.
- Homer R. Warner, M.D., Reed M. Gardner and Alan F. Toronto, M.D., Computer-Based Monitoring of Cardiovascular Functions in Postoperative Patients, Supplement II to *Circulation*, Apr. 1968, vols. 37 & 38, 68-74.
- Russell M. Nelson, Homer R. Warner, Reed E. Gardner and J. D. Mortensen, Computer Based Monitoring of Patients Following Cardiac Surgery, *Computers in Cardiology*, Jul.-Aug. 1969, vol. 5, No. 4, 926-930.
- Reed M. Gardner, Computerized Patient Monitoring at LDS Hospital—An Evaluation, *Proceedings of the San Diego Biomedical Symposium* 1971, vol. 10, 151-159.
- Reed M. Gardner, Monitoring of Physiological Data in a Clinical Environment, *Annual Review of Biophysics and Bioengineering*, 1972, vol. 1, 211-224.
- Reed M. Gardner, Computerized Intensive Care Monitoring at LDS Hospital—Progress and Development, 97-105.
- Reed M. Gardner, Donald R. Bennet, and Richard B. Vorce, Eight-Channel Data Set for Clinical EEG Transmission Over Dial-Up Telephone Network, *IEEE Transactions on Biomedical Engineering*, May 1974, vol. BME-21, No. 3, 246-249.
- Reed M. Gardner, George H. Cannon, Alan H. Morris, Kenneth R. Olsen, W. Gary Price, Computerized Blood Gas Interpretation and Reporting System, *Computer Magazine*, Jan. 1975, 39-45.
- Russell K. Hulse, Stephen J. Clark, J. Craig Jackson, Homer R. Warner and Reed M. Gardner, Computerized Medication Monitoring System, *American Journal of Hospital Pharmacy* 33, Oct. 1976, 1061-1064.
- Reed M. Gardner, Ph.D., *Computers in the ICU*, *Medical Electronics*, Jun. 1984, 129-135.
- Robert D. Andrews, M.S., M.T., Reed M. Gardner, Ph.D., Sandy M. Metcalf, R.R.T., and Deon Simmons, R.R.T., Computer Charting: An Evaluation of a Respiratory Care Computer System, *Respiratory Care*, Aug. 1985, vol. 30, No. 8, 695-707.
- Reed M. Gardner, Ph.D., Computerized Data Management and Decision Making in Critical Care, *Symposium on Critical Care*, Aug. 1985, vol. 65, No. 4, 1041-1051.
- Reed M. Gardner, David P. Scoville, Blair J. West, Beth Bateman, Robert M. Cundick, Jr., Terry P. Clemmer, Integrated Computer Systems for Monitoring of the Critically Ill, 1977, 301-307.
- T. Allan Pryor, Reed M. Gardner, Paul D. Clayton, Homer R. Warner, A Distributed Processing System for Patient Management, *Computers in Cardiology*, Sep. 1978, 325-328.
- Reed M. Gardner, Ph.D., Terry P. Clemmer, M.D., Keith G. Larsen, R.Ph., and Dickey S. Johnson, R.N., Computerized Alert System Use in Clinical Medicine, *IEEE Session* 6, 1979, 136-140.
- T. Allan Pryor, Homer R. Warner, Reed M. Gardner, Help—A Total Hospital Information System.
- T. P. Clemmer, R. M. Gardner, J. F. Orme, Jr., Computer Support in Critical Care Medicine, 1980.
- Scott R. Cannon, and Reed M. Gardner, Experience with a Computerized Interactive Protocol System Using HELP, *Computers and Biomedical Research* 13, 1980, 399-409, Academic Press, Inc.
- T. Allan Pryor, Paul D. Clayton, Reed M. Gardner, Randy Waki, and Homer R. Warner, HELP—A Hospital-Wide System for Computer-Based Support of Decision Making, Jan. 1981.
- T. A. Pryor, R. M. Gardner, P. D. Clayton and H. R. Warner, The HELP System Proceedings of the Sixth Annual Symposium on Computer Applications in Medical Care, Oct.-Nov. 1982, 19-27, IEEE.
- Reed M. Gardner, Information Management—Hemodynamic Monitoring, *Seminars in Anesthesia*, Dec. 1983, vol. 2, No. 4, 287-299.
- T. A. Pryor, R. M. Gardner, P. D. Clayton, H. R. Warner, The HELP System, *Journal of Medical Systems*, 1983, vol. 7, No. 2, 87-102.
- Reed M. Gardner, Blair J. West, T. Allan Pryor, Distributed Data Base and Network for ICU Monitoring, *IEEE Computers in Cardiology*, Sep. 18-24, 1984, 305-307.
- Reed M. Gardner, T. Allan Pryor, Paul D. Clayton, and R. Scott Evans, Integrated Computer Network for Acute Patient Care, *Symposium on Computer Application in Medical Care*, Nov. 4-7, 1984.
- Reed M. Gardner, Tomorrow's Electronic Hospital is Here Today, *IEEE Spectrum*, Jun. 1984, 101-103.
- Karen E. Bradshaw, Reed M. Gardner, Terry P. Clemmer, Jams F. Orme, Frank Thomas, and Blair J. West, Physician Decision Making—Evaluation of Data Used in a Computerized ICU, *International Journal of Clinical Monitoring and Computing* 1, 1984, 81-91.

- Terry P. Clemmer, M.D., and Reed M. Gardner, Ph.D., Data Gathering, Analysis, and Display in Critical Care Medicine, *Respiratory Care*, Jul. 1985, vol. 30, No. 7, 586-601.
- Reed M. Gardner, Ph.D., and William L. Hawley, Standardizing Communications and Networks in the ICU, *Patient Monitoring and Data Management*, 1985, 59-63.
- R. Scott Evans, Reed M. Gardner, Allan R. Bush, John P. Burke, Jay A. Jacobson, Robert A. Larsen, Fred A. Meier, and Homer R. Warner, Development of a Computerized Infectious Disease Monitor (CIDM), *Computers and Biomedical Research* 18, 185, 103-113.
- Reed M. Gardner, Ph.D., Susan M. Monis, Paul Oehler, Monitoring Direct Blood Pressure: Algorithm Enhancements, 607-610.
- R. Scott Evans, PHD, Robert A. Larsen, MD, John P. Burke, MD, Reed M. Gardner, PHD, Frederick A. Meier, MD, Jay A. Jacobson, MD, Marlyn T. Conti, BSN, Julie T. Jacobsen, MT, Russell K. Hulse, RPH, Computer Surveillance of Hospital-Acquired Infections and Antibiotic Use, *Journal of the American Medical Association*, Aug. 22-29, 1986, vol. 256, No. 8, 1007-1011.
- Reed M. Gardner, Computerized Management of Intensive Care Patients, Images, Signals, and Devices, 1986, vol. 3, No. 1.
- R. Whiting, L. Hayes, The Practice of Telemedicine—The TARDIS Perspective, *Informatics in Healthcare—Australia*, Jul./Aug. 1997, vol. 6, No. 3, 103-106.
- Monique Frize, Robin Walker, Clinical Decision- Support Systems for Intensive Care Units Using Care-Based Reasoning.
- Ho Sung Lee, Seung Hun Park, and Eung Je Woo, Remote Patient Monitoring Service Through World-Wide Web, *Proceedings—19<sup>th</sup> International Conference—IEEE/EMBS*, Oct. 30 - Nov. 2, 1997, 928-931.
- Betty L. Grundy, M.D., Pauline Crawford, R.N., Paul K. Jones, Ph.D., May Lou Kiley, Ph.D., Arnold Reisman, Ph.D., Yoh-Han Pao, Ph.D., Edward L. Wilkerson, M.D., J. S. Gravenstein, M.D., Telemedicine in Critical Care: An Experiment in Health Care Delivery, Oct. 1977, 6:10.
- Betty Lou Grundy, M.D., Paul K. Jones, Ph.D., and Ann Lovitt, M.D., Telemedicine in Critical Care: Problems in Design, Implementation, and Assessment, *Critical Care Medicine*, Jul. 1982, vol. 10, No. 7, 471-475.
- Geraldine Fitzpatrick, TARDIS Evaluation: Report on Final Usage Evaluation of the TARDIS Telehealth System, Jul. 24, 1998, Issue No. 1.0.
- Abstract Marie Delima, R.D., M. Michael Shabot, M.D., FACS, FCCM, FACMI, Karen Morris, R.N., Janet Mould, R.N., Eden Torre-Javier, R.N., Mark Lobue, B.A. and Jeannie Chen, Pharm.D., Successful Implementation of a Multiple-ICU Clinical Information System in a Tertiary Care Medical Center.
- Xin Li, Daniel J. Valentino, George J. So, Robert Lufkin, Rickey K. Taira, A World Wide Web Telemedicine System, *SPIE* vol. 2711, 427-439.
- Stephen M. Ayres, M.S., F.C.C.M., Ake Grenvik, M.D., Ph.D., F.C.C.M., Peter R. Holbrook, M.D., F.C.C.M., William C. Shoemaker, M.D., F.C.C.M., *Textbook of Critical Care*, 3<sup>rd</sup> Edition, 1995, Harcourt Brace & Company.
- Karen B. Tate, Ph.D., Reed M. Gardner, Ph.D., Kurt Scherting, Nurses, Pagers, and Patient-Specific Criteria; Three Keys to Improved Critical Value Reporting, 1995, 164-168, AMIA, Inc.
- Karen E. Tate, Ph.D., Reed M. Gardner, Ph.D., Computers, Quality, and the Clinical Laboratory: A Look at Critical Value Reporting, 17<sup>th</sup> Annual Symposium on Computer Applications in Medical Care, Oct. 30 - Nov. 3, 1993, 193-197.
- Peter J. Haug, Reed M. Gardner, Karen E. Tate, R. Scott Evans, Thomas D. East, Gilad Kuperman, T. Allan Pryor, Stanley M. Huff, and Homer R. Warner, Decision Support in Medicine: Examples from the HELP System, *Computers and Biomedical Research* 27, 1994, 396-418.
- Thomas D. East, Ph.D., C. Jane Wallace, R.N., M.S., Allan H. Morris, M.D., Reed M. Gardner, Ph.D., and Dwayne R. Westenskow, Ph.D., *Computers in Critical Care, New Technologies in Critical Care*, Jun. 1995, vol. 7, No. 2, 203-216.
- Reed M. Gardner, Ph.D., Bette B. Maack, R.R.A., R. Scott Evans, Ph.D., and Stanley M. Huff, M.D., Computerized Medical Care: The HELP System at LDS Hospital, *Journal of AHIMA*, Jun. 1992, 63(6):68-78.
- Reed M. Gardner, Ph.D., Integrated Computerized Records Provide Improved Quality of Care with Little Loss of Privacy, *Journal of the AMIA*, Jul./Aug. 1994, vol. 1, No. 4, 320-322.
- S Reddy, M. Niewiadomska-Bugaj, Y V Reddy, H C Galfalvy, V Jagannathan, R Raman, K. Srinivas, R. Shank, T. Davis, S. Friedman, MD, B. Merkin, MD, M. Kilkenny, MD, Experience with Aretmis—An Internet-Based Telemedicine System, *AMIA*, 1997, 759-763.
- Patrick R. Norris, M.S., Benoit M. Dawant, Ph.D., Antoine Geissbuhler, M.D., Web-Based Data Integration and Annotation in the Intensive Care Unit, 1997.
- H. C. Galfalvy, M.S., S. M. Reddy, Ph.D., M. Niewiadomska-Bugaj, Ph.D., S. Friedman, M.D., Evaluation of Community Care Network (CNN) System in a Rural Health Care Setting, 1995, *AMIA Inc.*, 698-702.
- K. Major, M. Shabot, S. Cunneen, Wireless Critical Alerts and Patient Outcomes in the Surgical Intensive Care Unit; *The American Surgeon*, 2000; p. 1057-1060.
- M. Shabot, M. Lobue, Cedars-Sinai Medical Center Critical Alerting System, Feb. 2004; p. 1-16.
- Shabot MM, LoBue M, Leyerly BJ, Dubin SB. Inferencing strategies for automated ALERTS on critically abnormal laboratory and blood gas data, *SCAMC* 1989; 13:54—57.
- APACHE® III Equation Update (Version III-J) 2002, pp. 1-22.
- APACHE® III Equation Update (Version III-I) 2003, pp. 1-13.
- O. Kostopoulou, M. Wildman, Sources of Variability in Uncertain Medical Decisions In the ICU: A Process Tracing Study, *Qual. Saf. Health Care* 2004, 13:272-280.
- A. Seiver, *Critical Care Computing: Past, Present, and Future; Critical Care Clinics*, vol. 16, No. 4, Oct. 2000, pp. 1-17.
- J. Fisher, S. Harbarth, A. Agathe, A. Benn, S. Ringer, D. Goldmann, and S. Fancani, Quantifying Uncertainty: Physicians' Estimates of Infection in Critically Ill Neonates and Children; *Clinical Infectious Diseases* 2004;38, pp. 1383-1390.
- N. Halpern, S. Pastores, R. Greenstein, *Critical Care Medicine in the United States 1985-2000: An Analysis of Bed Numbers, Use, and Cost; Critical Care Medicine* 2004, vol. 32, No. 6, pp. 1254-1259.
- J.Mrus, Getting Beyond Diagnostic Accuracy: Moving Toward Approaches That Can Be Used in Practice; *Clinical Infectious Diseases* 2004;38, pp. 1391-1393.
- B. Leyerly, M. Shabot, Integrated Computerized Databases for Medical Data Management Beyond the Bedside, *International Journal of Clinical Monitoring and Clinical Computing* 1990;7, pp 83-89.
- M.Shabot, M. Lobue, B. Leyerly, S. Dubin, Decision Support Alerts For Clinical Laboratory and Blood Gas Data, *Int. J. Clinical Monitoring and Computing* 1990;7, pp. 27-31.
- M. Shabot, M. Lobue, Real-Time Wireless Decision Support Alerts on a Palmtop PDA; *Proc. Ann. Symp. Compt Appl. Med Care* 1995, pp. 174-179.
- G. Kuperman, D. Sittig, M. Shabot, J. Teich, Clinical Decision Support for Hospital and Critical Care, pp. 174-179.
- W. Bates, M. Cohen, L. Leape, J. Overhage, M. Shabot, T. Sheridan, Reducing the Frequency of Errors In Medicine, *J. American Medical Informatics Assn.* 2001;8 pp. 299-308.
- M. Shabot, B. Leyerly, M. Lobue, Automatic Extraction of Intensity Intervention Scores From A Computerized Surgical ICU Flowsheet, *Am. J. Surg* 1987;154:1, pp. 72-76.

\* cited by examiner

Primary Examiner—Robert W Morgan

(74) Attorney, Agent, or Firm—Roberts, Mardula & Wertheim, LLC

(57)

#### ABSTRACT

A patient-worn monitoring system and method for geographically dispersed health care locations. A patient-worn monitoring system comprises a network, a body-worn monitoring station, a monitoring station server, a remote command center, and a rules engine. The network comprises a

first sub-network and a second sub-network. The body-worn monitoring station comprises monitoring equipment. The body worn monitoring station monitors data elements from a patient assigned to a health care location and sends monitored data elements to a monitoring station server via the first sub-network. The monitoring station server receives the monitored data elements from the body worn monitoring station and relays the monitored data elements to the remote command center via the second sub-network. The remote command center receives the monitored data elements from the monitoring station server, associates the monitored data elements with the patient assigned to the health care location, accesses patient data elements indicative of a medical

condition associated with the patient, and establishes a patient-specific rule associated with the patient. The rules engine selects data elements from the monitored data elements and the patient data elements associated with the patient and applies the patient-specific rule to the selected data elements continuously and simultaneously. A determination is made whether the patient-specific rule for the patient has been contravened. An alert is issued from the remote command center in the event the patient-specific rule for the patient has been contravened.

**46 Claims, 67 Drawing Sheets**




FIG. 1A





FIG. 2A



F

FIG. 2B




FIG. 3





FIG. 4A







FIG. 4C







FIG. 6A






FIG. 6B

FIG. 7





FIG. 8A







Fig.9






Fig.10





Fig. 11

- ▶ Hot Phone Access
- ▶ Scanner Control
- ▶ Audio/Video Switcher and Camera Control
- ▶ Remote Surveillance Display

FIG. 12






FIG. 13A




FIG. 13B

FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19






FIG. 20








FIG. 21B

FIG. 22




FIG. 23




FIG. 24A





FIG. 24B






FIG. 25A




FIG. 25B




FIG. 26A





FIG. 26B






FIG. 26D

FIG. 26E





FIG. 27A







FIG. 27B

FIG. 28A






FIG. 29A




FIG. 29B





FIG. 30A






FIG.30B

FIG. 31A





FIG. 31B







FIG. 32






FIG. 33B

FIG. 33C





FIG. 34A







FIG. 34B

FIG. 34C






FIG. 35A




FIG. 35B




FIG. 36




FIG. 37




FIG. 38




FIG. 39




FIG. 40




FIG. 41A




FIG. 41B




FIG. 42




FIG. 43

**SYSTEM AND METHOD FOR  
PATIENT-WORN MONITORING OF  
PATIENTS IN GEOGRAPHICALLY  
DISPERSED HEALTH CARE LOCATIONS**

RELATIONSHIP TO OTHER APPLICATIONS

This application is a continuation in part of application Ser. No. 10/654,668 filed Sep. 4, 2003 and a continuation in part of application Ser. No. 10/946,548 filed Sep. 21, 2004 now U.S. Pat. No. 7,256,708, both of which are continuations in part of application Ser. No. 09/443,072 filed Nov. 18, 1999, now U.S. Pat. No. 6,804,656 issued Oct. 12, 2004, which claims the benefit of U.S. Provisional Application No. 60/141,520, filed Jun. 23, 1999. The Ser. Nos. 10/654,668, 10/946,548, 09/443,072, and the 60/141,520 applications are hereby incorporated by reference in their entirety for all purposes.

FIELD OF THE INVENTION

This invention relates generally to the care of patients in health care locations, as for example and without limitation, Intensive Care Units (ICUs). More particularly this invention is a system and method for care of the critically ill that combines a real-time, multi-node telemedicine network and an integrated, patient care management system to enable specially-trained Intensivists to provide 24-hour/7-day-per-week patient monitoring and management to multiple, geographically dispersed ICUs from both on-site and remote locations.

BACKGROUND OF THE INVENTION

While the severity of illness of ICU patients over the past 15 years has increased dramatically, the level of and type of physician coverage in most ICUs has remained constant. Most ICU patients receive brief minutes of attention during morning rounds from physicians with limited critical care experience. During the remainder of the day and night, nurses are the primary caregivers, with specialists called only after patient conditions have started to deteriorate. The result of this mismatch between severity of illness and physician coverage is an unacceptably high ICU mortality rate (10% nationwide), and a high prevalence of avoidable errors that result in clinical complications. In 1998, an Institute of Medicine Roundtable determined that avoidable patient complications were the single largest problem in medical care delivery. In another prominent 1998 study of 1000 patients, 46% experienced an avoidable adverse event in care, with 40% of these errors resulting in serious disability or death.

The physicians who can remedy this situation are in critically short supply. Numerous studies have shown that Intensivists (physicians who have trained and board certified in Critical Care Medicine) can markedly improve patient outcomes. However, only one-third of all ICU patients ever has an Intensivist involved in their care, and the number of Intensivists would need to increase tenfold (nationally) to provide 24-hour coverage to all ICU patients. With the rapid aging of the population, this shortfall of expertise is going to increase dramatically.

Even where Intensivists are present (and especially where they are not), patients suffer from unnecessary variation in practice. There is little incentive for physicians to develop and conform to evidence-based best practices (it takes significant work and a change in behavior to develop and

implement them). This variation contributes to sub-optimal outcomes, in both the quality and cost of care delivered to ICU patients.

What is needed is a redesigning of the critical care regimen offered to patients in an ICU. Rather than the consultative model where a periodic visit takes place and the doctor then goes away, a more active 24-hour intensivist managed care is required. Further, technology that leverages the intensivists' expertise and standardizes the care afforded to patients in an ICU is required. Further, continuous feedback to improve the practice of intensivists in an ICU is necessary to provide the intervention required to minimize adverse events. This invention seeks to provide new methods for managing and delivering care to the critically ill.

Attempts to automate various aspects of patient care have been the subject of various inventions. For example, U.S. Pat. No. 5,868,669 to Iliff was issued for "Medical Diagnostic and Treatment Advice System." The disclosed invention is for a system and method for providing knowledge based medical diagnostic and treatment advice to the general public over a telephone network.

U.S. Pat. No. 5,823,948 to Ross, Jr. et al was issued for "Medical Records Documentation, Tracking and Order Entry System". The disclosed invention is for a system and method that computerizes medical records, documentation, tracking and order entries. A teleconferencing system is employed to allow patient and medical personnel to communicate with each other. A video system can be employed to videotape a patient's consent.

U.S. Pat. No. 4,878,175 to Norden-Paul et al. was issued for "Method for Generating Patient-Specific Flowsheets By Adding/Deleting Parameters." The disclosed invention is for an automated clinical records system for automated entry of bedside equipment results, such as an EKG monitor, respirator, etc. The system allows for information to be entered at the bedside using a terminal having input means and a video display.

U.S. Pat. No. 5,544,649 to David et al. was issued for "Ambulatory Patient Health Monitoring Techniques Utilizing Interactive Visual Communications." The disclosed invention is for an interactive visual system, which allows monitoring of patients at remote sites, such as the patient's home. Electronic equipment and sensors are used at the remote site to obtain data from the patient, which is sent to the monitoring site. The monitoring site can display and save the video, audio and patient's data.

U.S. Pat. No. 5,867,821 to Ballantyne et al. was issued for "Method and Apparatus for Electronically Accessing and Distributing Personal Health Care Information and Services in Hospitals and Homes." The disclosed invention is for an automated system and method for distribution and administration of medical services, entertainment services, and electronic health records for health care facilities.

U.S. Pat. No. 5,832,450 to Myers et al. issued for "Electronic Medical Record Using Text Database." The disclosed invention is for an electronic medical record system, which stores data about patient encounters arising from a content generator in freeform text.

U.S. Pat. No. 5,812,983 to Kumagai was issued for "Computer Medical File and Chart System." The disclosed invention is for a system and method which integrates and displays medical data in which a computer program links a flow sheet of a medical record to medical charts.

U.S. Pat. No. 4,489,387 to Lamb et al. was issued for "Method and Apparatus for Coordinating Medical Procedures." The disclosed invention is for a method and appa-

ratus that coordinates two or more medical teams to evaluate and treat a patient at the same time without repeating the same steps.

U.S. Pat. No. 4,731,725 to Suto et al. issued for "Data Processing System which Suggests a Pattern of Medical Tests to Reduce the Number of Tests Necessary to Confirm or Deny a Diagnosis." The disclosed invention is for a data processing system that uses decision trees for diagnosing a patient's symptoms to confirm or deny the patient's ailment.

U.S. Pat. No. 5,255,187 to Sorensen issued for "Computer Aided Medical Diagnostic Method and Apparatus." The disclosed invention is for an interactive diagnostic system which relies on color codes which signify the presence or absence of the possibility of a disease based on the symptoms a physician provides the system.

U.S. Pat. No. 5,553,609 to Chen et al. issued for "Intelligent Remote Visual Monitoring System for Home Health Care Service." The disclosed invention is for a computer-based remote visual monitoring system, which provides in-home patient health care from a remote location via ordinary telephone lines.

U.S. Pat. No. 5,842,978 to Levy was issued for "Supplemental Audio Visual Emergency Reviewing Apparatus and Method." The disclosed invention is for a system which videotapes a patient and superimposes the patient's vital statistics onto the videotape.

While these inventions provide useful records management and diagnostic tools, none of them provides a comprehensive method for monitoring and providing real time critical care at disparate ICUs. In short, they are NOT designed for critical care. Further, none of these inventions provide for the care of a full time intensivist backed by appropriate database and decision support assistance in the intensive care environment. What would be useful is a system and method for providing care for the critically ill that maximizes the presence of an intensivist trained in the care of the critically ill. Further such a system would standardize the care in ICUs at a high level and reduce the mortality rate of patients being cared for in ICUs.

#### SUMMARY OF THE INVENTION

The present invention provides a core business of Continuous Expert Care Network (CXCN) solution for hospital intensive care units (ICUs). This e-solution uses network, database, and decision support technologies to provide 24-hour connectivity between Intensivists and ICUs. The improved access to clinical information and continuous expert oversight leads to reduced clinical complications, fewer medical errors, reduced mortality, reduced length of stay, and reduced overall cost per case.

The technology of the present invention as explained below can be implemented all at once or in stages. Thus the technology, as more fully explained below is available in separate components to allow for the fact that hospitals may not be able to implement all of the technology at once. Thus modular pieces (e.g. videoconferencing, vital sign monitoring with smart alarms, hand-held physician productivity tools, etc.) can be implemented, all of which can add value in a stand-alone capacity. First amongst these offerings will be an Intensivist Decision Support System, a stand-alone software application that codifies evidence-based, best practice medicine for 150 common ICU clinical scenarios. These support algorithms are explained more fully below.

The "Command Center" model, again as more fully set forth below, will ultimately give way to a more distributed remote management model where Intensivists and other

physicians can access ICU patients and clinicians (voice, video, data) from their office or home. In this scenario, the present invention will be available in hospital applications that centralize ICU information, and offer physicians web-based applications that provide them with real-time connectivity to this information and to the ICUs. This access and connectivity will enable physicians to monitor and care for their patients remotely. These products will be natural extensions and adaptations of the present invention and the existing applications disclosed herein that those skilled in the art will appreciate and which do not depart from the scope of the invention as disclosed herein.

The present invention addresses these issues and shortcomings of the existing situation in intensive care, and its shortfalls via two major thrusts. First, an integrated video/voice/data network application enables continuous real-time management of ICU patients from a remote setting. Second, a client-server database application—integrated to the remote care network—provides the data analysis, data presentation, productivity tools and expert knowledge base that enable a single Intensivist to manage the care of up to 40 patients simultaneously. The combination of these two thrusts—care management from a remote location and new, technology-enhanced efficiency of Intensivist efforts—allows health care systems to economically raise the standard of care in their ICUs to one of 24.times.7 continuous Intensivist oversight.

It is therefore an object of the present invention to reduce avoidable complications in an ICU.

It is a further object of the present invention to reduce unexplained variations in resource utilization in an ICU.

It is a further objective of the present invention to mitigate the serious shortage of intensivists.

It is yet another objective of the present invention to reduce the occurrence of adverse events in an ICU.

It is a further objective of the present invention to standardize the care at a high level among ICUs.

It is yet another objective of the present invention to reduce the cost of ICU care.

It is yet another objective of the present invention to dramatically decrease the mortality in an ICU.

It is yet another objective of the present invention to bring information from the ICU to the intensivist, rather than bring the intensivist to the ICU.

It is a further objective of the present invention to combine tele-medical systems comprising two-way audio/video communication with a continuous real time feed of clinical information to enable the intensivist to oversee care within the ICU.

It is a further objective of the present invention to allow intensivists to monitor ICUs from a site remote from each individual ICU.

It is a further objective of the present invention to bring organized detailed clinical information to the intensivist, thereby providing standardized care in the ICU.

It is yet another objective of the present invention to utilize knowledge-based software to use rules, logic, and expertise to provide preliminary analysis and warnings for the intensivists.

The present invention comprises a command center/remote location, which is electronically linked to ICUs remote from the command center/remote location. The command center/remote location is manned by intensivists 24 hours a day, seven days per week. Each ICU comprises a nurse's station, to which data flows from individual beds in the ICU. Each patient in the ICU is monitored by a video camera, as well as by clinical monitors typical for the intensive care

unit. These monitors provide constant real time patient information to the nurse's station, which in turn provides that information over a dedicated T-1 (high bandwidth) line to the ICU command center/remote location. As noted earlier, the command center/remote location is remote from the ICU, thereby allowing the command center/remote location to simultaneously monitor a number of patients in different ICUs remote from the command center/remote location.

At each command center/remote location, video monitors exist so that the intensivist can visually monitor patients within the ICU. Further, the intensivist can steer and zoom the video camera near each patient so that specific views of the patient may be obtained, both up close and generally. Audio links allow intensivists to talk to patients and staff at an ICU bed location and allow those individuals to converse with the intensivist.

Clinical data is constantly monitored and presented to the command center/remote location in real time so that the intensivist can not only monitor the video of the patient but also see the vital signs as transmitted from the bedside. The signals from the clinical data and video data are submitted to a relational database, which comprises 1) standardized guidelines for the care of the critically ill, 2) various algorithms to support the intensive care regimen, 3) order writing software so that knowledge-based recommendations and prescriptions for medication can be made based upon the clinical data, and 4) knowledge-based vital-sign/hemodynamic algorithms that key the intensivist to engage in early intervention to minimize adverse events.

The advantage of the present invention is that intensivists see all patients at a plurality of ICU's at all times. Further, there is a continuous proactive intensivist care of all patients within the ICU, thereby minimizing adverse events. Intervention is triggered by evidence-based data-driven feedback to the intensivist so that standardized care can be provided across a plurality of ICUs.

The economic benefits of the present invention are manifold. For the first time, 24-hour a day, seven day a week intensivist care for patients in an ICU can be obtained. Further, more timely interventions in the care of the patients can be created by the knowledge-based guidelines of the present invention, thereby minimizing complications and adverse events. This in turn will lead to a reduced mortality within the ICU, and hence, a reduced liability cost due to the dramatic reduction in avoidable errors in health care.

By providing timely interventions, the length of stay within the ICU can be greatly reduced, thereby allowing more critically ill patients to be cared for in the ICU.

In addition, by reviewing and standardizing the care afforded to patients in an ICU, a more standardized practice across a variety of ICUs can be achieved. This will lead to more cost-effective care within the ICU, and reduced ancillary cost for the care of the critically ill.

The overall architecture of the present invention comprises a "pod." The pod comprises a tele-medicine command center/remote location connected to a plurality multiple ICUs at various locations. The connection between the command center/remote location and the ICUs is via a dedicated wide-area network linking the ICUs to the command center/remote location and a team of intensivists who integrate their services to provide 24-hour, seven day a week care to all of the pod ICUs.

The pod is connected via a wide-area network using dedicated T-1 lines, for example, with redundant backup. This network provides reliable, high speed secure transmission of clinical data and video/audio signals between each

patient room and the command center/remote location. The use of a T-1 line is not meant as a limitation. It is expected that more and higher bandwidth networks will become available. Such high bandwidth networks would come within the scope of the invention as well.

Each patient room is equipped with a pan/tilt/zoom video camera with audio and speaker to enable full videoconferencing capability. In addition, computer workstations are dedicated for exclusive physician use in each ICU, preferably at the nurse's station. Intensivists use the workstations to view patient information, consult decision support information, record their notes, and generate patient orders.

The patient management software used by intensivists is provided across the pod. Updates and changes made to the record are available at both the ICU and the command center/remote location for any given patient.

Each command center/remote location contains at least three workstations: one for the intensivist, one for the critical care registered nurse, and one for a clerk/administrative person.

The intensivist workstation comprises separate monitors for displaying ICU video images of patients and/or ICU personnel, output from bedside monitoring equipment, patient clinical data comprising history, notes, lab reports, etc., and decision support information. The staff at the command center/remote location are able to activate and control the cameras in each patient's room so that appropriate visual views of the patient can be generated.

Intensivists are able to switch between rooms and patients and can monitor at least two rooms simultaneously via the video screens. Patient data such as X-ray and ECG images are scanned and transmitted to the command center/remote location upon request of the intensivist.

Remote patient management is utilized in the present invention's critical care program to supplement traditional onsite care. The rationale underlying the remote patient management of the present invention is that critically ill patients are inherently unstable and require continuous expert care that is not now offered in existing ICU monitoring regimens. Further, remote monitoring allows a single intensivist to care for patients in multiple ICU locations, thereby creating an efficiency that makes continuous care feasible.

Remote intensivist care of the present invention is proactive. Intensivists will order needed therapies and check results of tests and monitor modalities in a more timely fashion than is currently offered. Patients can be observed visually when needed using the ceiling-mounted cameras in each room.

Command center/remote location personnel communicate with ICU staff through videoconferencing and through "hot phones," which are dedicated telephones directly linked between the command center/remote location and the ICU. These communications links are used to discuss patient care issues and to communicate when a new order has been generated.

Intensivists document important events occurring during their shift in progress notes generated on the command center/remote location computer terminal.

Intensivists detect impending problems by intermittently screening patient data, including both real time and continuously stored vital sign data. Patient severity of illness determines the frequency with which each patient's data is reviewed by the intensivists.

Embodiments of the present invention provide a system for providing continuous, expert network health care services from a remote location. The system comprises a

plurality of health care locations, at least one remote command center for managing health care at the plurality of health care locations, and at least one network. The plurality of health care locations is electronically connected to said-at least one remote command center by the network. By way of illustration and not as limitation, a health care location may be a hospital, a nursing home, a mobile health care facility, a space-based health care facility, a residence, an emergency room, an intensive care unit, a field health care facility, an operating room, and a step-down unit. Additionally, the mobile health care facility is selected from the group consisting of a ship, a helicopter, an ambulance, a MASH unit, and a tent hospital. The at least one remote command center provides intensivists monitoring of the plurality of health care locations 24 hours per days seven days per week. In an embodiment, a patient monitoring system for geographically dispersed health care locations comprises a network, a body-worn monitoring station, a monitoring station server, a remote command center, and a rules engine. The network comprises a first sub-network and a second sub-network, either of which may be wired or wireless. By way of illustration and not as a limitation, a wireless sub-network may be a Bluetooth-compatible network, an 802.11-compatible network, and a short-range, slotted protocol network. The body-worn monitoring station comprises monitoring equipment and is adapted for monitoring data elements from a patient assigned to a health care location and sending monitored data elements to a monitoring station server via the first sub-network. By way of illustration and not as a limitation, the monitoring equipment may be an ECG sensor, a pulse oximeter, a thermometer, a respiration sensor, a heart rate sensor, an audio sensor, and a tachodynamometer. The monitoring station server is adapted for receiving the monitored data elements from the body worn monitoring station and relaying the monitored data elements to the remote command center via the second sub-network. In an embodiment, the monitored data elements are associated with a patient identifier. The remote command center is adapted for receiving the monitored data elements from the monitoring station server, associating the monitored data elements with the patient assigned to the health care location, accessing patient data elements indicative of a medical condition associated with the patient, and establishing a patient-specific rule associated with the patient. The rules engine is adapted for selecting data elements from the monitored data elements and the patient data elements associated with the patient, applying the patient-specific rule to the selected data elements continuously and simultaneously, determining whether the patient-specific rule for the patient has been contravened, and issuing an alert from the remote command center in the event the patient-specific rule for the patient has been contravened. In another embodiment, the monitoring station server is wearable by the patient. For example and not as a limitation, the patient may be a fetus carried by an expectant mother and the monitoring equipment may comprise a tachodynamometer.

The remote command center further comprises a patient care management system for monitoring and treating individual patients at any of said plurality of healthcare locations. The patient care management system further comprises a data server/data warehouse for storing and analyzing data from the at least one remote command center.

Each of the plurality of health care locations further comprises patient monitoring equipment electronically connected to the at least one remote command center over the network. In another embodiment of the present invention

each health care location further comprises a nurses' station electronically connected to said monitoring equipment and to the at least one remote command center over the network. In still another embodiment of the present invention, the healthcare locations comprise intensive care units (ICU's).

Optionally, the patient care management system further comprises a relational database for storing a plurality of decision support algorithms and for prompting intensivists to provide care to patients based upon any of the decision support algorithms. The algorithms are selected from the group consisting of algorithms for treating Acalculous Cholecystitis, Acute Pancreatitis Algorithms, Acute Renal Failure-Diagnosis, Acute Renal Failure-Management & Treatment, Adrenal Insufficiency, Agitation and Anxiety, Depression & Withdrawal, Aminoglycoside Dosing and Therapeutic Monitoring, an Amphotericin-B Treatment Guidelines, Analgesia, Antibiotic Classification & Costs, Antibiograms Algorithm, Antibiotic associated Colitis Algorithm, ARDS: Hemodynamic Management, ARDS: Steroid Use, ARDS: Ventilator Strategies, Asthma, Bleeding Patient, Bloodstream Infections, Blunt Cardiac Injury, Bradyarrhythmias, Brain Death, Bronchodilator Use in Ventilator Patients, Bronchoscopy & Thoracostomy Guidelines, Candiduria, Cardiogenic Shock, CardioPulmonary Resuscitation Guideline, Catheter Related Septicemia, a Catheter Replacement Strategies, Cervical Cord Injury, Congestive Heart Failure, COPD Exacerbation & Treatment, CXR (Indications), Dealing with Difficult patients and families, Diabetic Ketoacidosis, Dialysis, Diuretic Use, Drug Changes with Renal Dysfunction, Emergency Cardiac Pacing, Endocarditis Diagnosis and Treatment, Endocarditis Prophylaxis, End of Life Decisions, Endotracheal Tubes & Tracheotomy, Ethical Guidelines, Febrile Neutropenia, FUO, Fluid Resuscitation, Guillain-Barre Syndrome, Heparin, Heparin-Induced Thrombocytopenia, Hepatic Encephalopathy, Hepatic Failure, HIV+ Patient Infections, Hypercalcemia Diagnosis and Treatment, Hyperglycemia Insulin Treatment, Hyperkalemia: Etiology & Treatment, Hypermagnesemia: Etiology & Treatment, Hypertensive Crisis, Hypokalemia: Etiology & Treatment, Hyponatremia: Etiology & Treatment, Hypothermia, Identification of Cervical Cord Injury, Implantable Cardio-defibrillator, Intra-Aortic Balloon Device, Intracerebral Hemorrhage, Latex Allergy, Magnesium Administration, Management of Hypotension, Inotropes, Management of Patients with Ascites, Empiric Meningitis, Meningitis, a Myasthenia Gravis, Myocardial Infarction, Myocardial Infarction with left bundle branch block, Necrotizing Soft Tissue Infections, Neuromuscular Blockers, Neuromuscular Complications of Critical Illness, Non-Infectious Causes of Fever, Non-Traumatic Coma, Noninvasive Modes of Ventilation, Nutritional Management, Obstetrical Complication, Oliguria, Open Fractures, Ophthalmic Infections, Organ Procurement Guidelines, PA Catheter Guideline and Troubleshooting, Pancreatitis, Penetrating Abdominal Injury, Penetrating Chest Injury, Penicillin Allergy, Permanent Pacemaker and Indications, Pneumonia Community Acquired, Pneumonia Hospital Acquired, Post-Op Bleeding, Post-Op Hypertension, Post-Op Management of Abdominal Post-Op Management of Carotid, Post-Op Management of Open Heart, Post-Op Management of Thoracotomy, Post-Op Myocardial Ischemia (Non-Cardiac Arrhythmias after Cardiac Surgery), Post-Op Power Weaning, Pressure Ulcers, Pulmonary Embolism Diagnosis, Pulmonary Embolism Treatment, Respiratory Isolation, Sedation, Seizure, Status Epilepticus, Stroke, Sub-Arachnoid Hemorrhage, Supra-Ventricular Tachyarrhythmia, Supra-Ventricular Tachycardia, Wide Complex QRS Tachycardia,

Therapeutic Drug Monitoring, Thrombocytopenia, Thrombolytic Therapy, Transfusion Guidelines, Traumatic Brain Injury, Assessment of Sedation, Sedation, Septic Shock, Bolus Sliding, Scale Midazolam, Short Term Sedation Process, Sinusitis, SIRS, Spinal Cord Injury, Steroid Replacement Strategy, Thyroid Disease, Transplant Infection Prophylaxis, Transplant Related Infections, Treatment of Airway Obstruction, Unknown Poisoning, Unstable Angina, Upper GI Bleeding Stress Prophylaxis, Vancomycin, Upper GI Bleeding Non-Variceal, Upper GI Bleeding Variceal, Use of Hematopoietic Growth Factors, Ventilator Weaning, Ventilator Weaning Protocol, Venous Thrombosis Diagnosis and Treatment, Venous Thromboembolism Prophylaxis, Ventricular Arrhythmia, Warfarin, Warfarin Dosing, and Wound Healing Strategies.

In yet another embodiment of the present invention, the patient care management system further comprises order writing software for providing knowledge-based recommendations and prescriptions for medication based upon the clinical data. In another embodiment of the present invention, the patient care management system further comprises knowledge-based vital sign/hemodynamic algorithms that prompt said intensivist to engage in early intervention.

Embodiments of the present invention provide methods for continuous expert critical care. Patients are monitored in a plurality of ICU's. Information from the patient monitoring is communicated to at least one command center over a first network. The information from the patient monitoring is received and analyzed at the command center over the first network; and guidance is provided from the command center to the plurality of ICU's to take actions regarding patient care. In another embodiment of the present invention, providing guidance from the command center further comprises an intensivist reviewing decision support algorithms that provide guidance for treating a plurality of critical care conditions. The algorithms are taken from the group consisting of algorithms for treating Acalculous Cholecystitis, Acute Pancreatitis Algorithm, Acute Renal Failure-Diagnosis, Acute Renal Failure-Management & Treatment, Adrenal Insufficiency, Agitation and Anxiety, Depression & Withdrawal, Aminoglycoside Dosing and Therapeutic Monitoring, an Amphotericin-B Treatment Guidelines, Analgesia, Antibiotic Classification & Costs, Antibigrams Algorithm, Antibiotic associated Colitis Algorithm, ARDS: Hemodynamic Management, ARDS: Steroid Use, ARDS: Ventilator Strategies, Asthma, Bleeding Patient, Bloodstream Infections, Blunt Cardiac Injury, Bradyarrhythmias, Brain Death, Bronchodilator Use in Ventilator Patients, Bronchoscopy & Thoracentesis Guidelines, Candiduria, Cardiogenic Shock, CardioPulmonary Resuscitation Guideline, Catheter Related Septicemia, a Catheter Replacement Strategies, Cervical Cord Injury, Congestive Heart Failure, COPD Exacerbation & Treatment, CXR (Indications), Dealing with Difficult patients and families, Diabetic Ketoacidosis, Dialysis, Diuretic Use, Drug Changes with Renal Dysfunction, Emergency Cardiac Pacing, Endocarditis Diagnosis and Treatment, Endocarditis Prophylaxis, End of Life Decisions, Endotracheal Tubes & Tracheotomy, Ethical Guidelines, Febrile Neutropenia, FUO, Fluid Resuscitation, Guillain-Barre Syndrome, Heparin, Heparin-Induced Thrombocytopenia, Hepatic Encephalopathy, Hepatic Failure, HIV+ Patient Infections, Hypercalcemia Diagnosis and Treatment, Hyperglycemia Insulin Treatment, Hyperkalemia: Etiology & Treatment, Hyponatremia: Etiology & Treatment, Hypertensive Crisis, Hypokalemia: Etiology & Treatment, Hyponatremia: Etiology & Treatment, Hypothermia, Identification of Cervical Cord Injury, Implantable Cardio-

defibrillator, Intra-Aortic Balloon Device, Intracerebral Hemorrhage, Latex Allergy, Magnesium Administration, Management of Hypotension, Inotropes, Management of Patients with Ascites, Empiric Meningitis, Meningitis, a Myasthenia Gravis, Myocardial Infarction, Myocardial Infarction with left bundle branch block, Necrotizing Soft Tissue Infections, Neuromuscular Blockers, Neuromuscular Complications of Critical Illness, Non-Infectious Causes of Fever, Non-Traumatic Coma, Noninvasive Modes of Ventilation, Nutritional Management, Obstetrical Complications, Oliguria, Open Fractures, Ophthalmic Infections, Organ Procurement Guidelines, PA Catheter Guideline and Troubleshooting, Pancreatitis, Penetrating Abdominal Injury, Penetrating Chest Injury, Penicillin Allergy, Permanent Pacemaker and Indications, Pneumonia Community Acquired, Pneumonia Hospital Acquired, Post-Op Bleeding, Post-Op Hypertension, Post-Op Management of Abdominal, Post-Op Management of Carotid, Post-Op Management of Open Heart, Post-Op Management of Thoracotomy, Post-Op Myocardial Ischemia, (Non-Cardiac Arrhythmias after Cardiac Surgery), Post-Op Power Weaning, Pressure Ulcers, Pulmonary Embolism Diagnosis, Pulmonary Embolism Treatment, Respiratory Isolation, Sedation, Seizure, Status Epilepticus, Stroke, Sub-Arachnoid Hemorrhage, Supraventricular Tachyarrhythmia, Supra-Ventricular Tachycardia, Wide Complex QRS Tachycardia, Therapeutic Drug Monitoring, Thrombocytopenia, Thrombolytic Therapy, Transfusion Guidelines, Traumatic. Brain Injury, Assessment of Sedation, Sedation, Septic Shock, Bolus Sliding Scale Midazolam, Short Term Sedation Process, Sinusitis, SIRS, Spinal Cord Injury, Steroid Replacement Strategy, Thyroid Disease, Transplant Infection Prophylaxis, Transplant Related Infections, Treatment of Airway Obstruction, Unknown Poisoning, Unstable Angina, Upper GI Bleeding Stress Prophylaxis, Vancomycin, Upper GI Bleeding Non-Variceal, Upper GI Bleeding Variceal, Use of Hematopoietic Growth Factors, Ventilator Weaning, Ventilator Weaning Protocol, Venous Thrombosis Diagnosis and Treatment, Venous Thromboembolism Prophylaxis, Ventricular Arrhythmia, Warfarin, Warfarin Dosing, and Wound Healing Strategies.

In another embodiment, a method further comprises a data server/data warehouse storing and analyzing patient data from the at least one command center and providing analysis in results over a second network to the at least one command center.

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1A illustrates the logical data structure for billing, insurance and demographic information.

FIG. 1B illustrates the logical data structure for billing, insurance and demographic information (cont).

FIG. 2A illustrates the command center logical data structure.

FIG. 2B illustrates the command center logical data structure (cont).

FIG. 3 illustrates the logical data structure for creating a medical history.

FIG. 4A illustrates the logical data structure for creating notes relating to patient treatment and diagnosis.

FIG. 4B illustrates the logical data structure for creating notes relating to patient treatment and diagnosis (cont).

FIG. 4C illustrates the logical data structure for creating notes relating to patient treatment and diagnosis (cont).

FIG. 5 illustrates the logical data structure for entry of medical orders.

FIG. 6A illustrates the logical data structure for patient care, laboratory testing and diagnostic imaging.

FIG. 6B illustrates the logical data structure for patient care, laboratory testing and diagnostic imaging (cont).

FIG. 7 illustrates the logical data structure for categories of information that are permitted to be presented to intensivists and other care givers by the system.

FIG. 8A illustrates the logical data structure for documenting patient vital signs.

FIG. 8B illustrates the logical data structure for documenting patient vital signs (cont).

FIG. 9 illustrates the distributed architecture of the present invention.

FIG. 10 illustrates the system architecture of the present invention.

FIG. 11 illustrates the decision support algorithm for diagnosis and treatment of pancreatitis.

FIG. 12 illustrates the vital signs data flow.

FIG. 13A illustrates capture and display of diagnostic imaging.

FIG. 13B illustrates establishing videoconferencing in the present invention.

FIG. 14 illustrates the physician resources order writing data interface of the present invention.

FIG. 15 illustrates the physician resources database data interface of the present invention.

FIG. 16 illustrates the automated coding and billing system integrated with the workflow and dataflow of the present invention.

FIG. 17 illustrates the order writing data flow of the present invention.

FIG. 18 illustrates the event log flow of the present invention.

FIG. 19 illustrates the smart alarms implementation of the present invention.

FIG. 20 illustrates the procedure note creation and line log for the present invention.

FIGS. 21A-B illustrate the acalculous cholecystitis decision support algorithm.

FIG. 22 illustrates the adrenal insufficiency decision support algorithm.

FIG. 23 illustrates the blunt cardiac injury decision support algorithm.

FIGS. 24A-B illustrate the candiduria decision support algorithm.

FIGS. 25A-B illustrate the cervical spine injury decision support algorithm.

FIGS. 26A-B illustrate the oliguria decision support algorithm.

FIGS. 26C-D illustrate the oliguria decision support algorithm (cont).

FIG. 26E illustrates the oliguria decision support algorithm (cont).

FIGS. 27A-B illustrate the open fractures decision support algorithm.

FIGS. 28A-B illustrate the pancreatitis decision support algorithm.

FIGS. 29A-B illustrate the penicillin allergy decision support algorithm.

FIGS. 30A-B illustrate the post-op hypertension decision support algorithm.

FIG. 31A illustrates the pulmonary embolism decision support algorithm.

FIG. 31B illustrates the pulmonary embolism decision support algorithm (cont).

FIG. 32 illustrates the seizure decision support algorithm.

FIGS. 33A-B illustrate the SVT determination decision support algorithm.

FIG. 33C illustrates the SVT unstable decision support algorithm.

FIGS. 34A-B illustrate the wide complex QRS Tachycardia decision support algorithm.

FIG. 34C illustrates the wide complex QRS Tachycardia decision support algorithm (cont).

FIG. 35A illustrates the assessment of sedation decision support algorithm.

FIG. 35B illustrates the assessment of sedation decision support algorithm (cont).

FIG. 36 illustrates the bolus sliding scale midazolam decision support algorithm.

FIG. 37 illustrates the sedation assessment algorithm decision support algorithm.

FIG. 38 illustrates the short term sedation process, decision support algorithm.

FIG. 39 illustrates the respiratory isolation decision support algorithm.

FIG. 40 illustrates the empiric meningitis treatment decision support algorithm.

FIG. 41A illustrates the ventilator weaning decision support algorithm.

FIG. 41B illustrates the ventilator weaning decision support algorithm (cont).

FIG. 42 illustrates the warfarin dosing decision support algorithm.

FIG. 43 illustrates the HIT-2 diagnostic decision support algorithm.

DEFINITIONS OF TERMS AND DATA

In the following Detailed Description of the Invention, a number of modules and procedures are described. For purposes of definitions, the following module definitions apply and are more fully amplified in the descriptions of the figures that follow.

Term Definitions

Following are a series of definitions for certain terms used in this specification:

Insurance carrier: This is a table of all the valid insurance carriers listed in the system of the present invention.

Patient guarantor: Provides the insurance guarantor information for a given patient.

Patient information: Provides demographic information for each patient.

Medical event date history: This contains the various disorders of the patient and the dates associated with major medical events relating to those disorders.

Medical history: Contains non-major system medical history of a patient.

Drug: Contains what medication and allergies have been identified for a patient at admission.

Address: Contains the address or addresses for a given patient.

Patient visit: There may be multiple records for any given patient, since the patient may visit the ICU on more than one occasion. This file contains a record of each visit to an ICU by a patient.

Physician-patient task: Contains the task that had been defined for each patient.

Present illness: This contains a textual description of the patient illness for the specific ICU visit.

Physical exam: This contains the information gathered as a result of a physical examination of the patient during the admission to the ICU.

Surgical fluids: This provides all the information related to the fluids provided during surgery.

Surgery: This contains all information pertaining to any surgical procedure performed on a patient while the patient is at the ICU.

Patient admit: This provides general information that needs to be gathered when a patient is admitted into the ICU.

Medical orders: This provides the general information for all types of medical orders associated with a given patient.

Daily treatment: This contains the treatment provided for a given patient on a given day.

Daily diagnosis: This contains the daily diagnosis for a given patient, which includes neurological, cardiological, pulmonary, renal, endocrinological, and any other diagnosis that may be associated with a patient.

Vital sign information is also critical to the administration of care in the ICU. A number of different modules collect information relating to patient vital signs. For example:

Patient admit: This provides the general information that needs to be gathered when a patient is admitted to the ICU

Patient visit: This contains a record of each visit to an ICU by a patient.

Patient: Provides demographic information for each patient.

Vital sign header: This contains general information related to the vital sign data for the particular patient.

Vital sign: Contains the vital sign data taken at specific intervals for a given patient.

Hospital: This contains identifying information for a particular hospital where the care is given.

ICU bed: Contains the association for identifying which beds are in a given ICU

Command center/remote location definitions and modules have also been created for the present invention to allow for the orderly storage and retrieval and entering of data. For example:

Physician-physician (such as nurses and LPN and the like):

Contains the names of all of the physicians and physician extenders for the command center/remote location as well as for ICUs associated with the command center/remote location.

Communication: Contains all of the various types of communication vehicles used to contact an individual physician or physician extender.

Physician role: Contains the role a physician is playing for a given patient, (i.e., primary care, consultant, etc.)

Patient: Provides demographic information for each patient.

Command center/remote location: Provides identifying information for a particular command center/remote location.

Hospital: Contains identifying information for a particular hospital wherein an ICU is located.

ICU: Contains identifying information for an ICU at a hospital.

ICU bed: Contains the association for identifying which beds are in a given hospital.

ICU patient location: Provides the association between an ICU and a patient and identifies where a patient is located within an ICU in a particular hospital.

The order entry functionality of the present invention provides a critical service for obtaining information on the patient during admission, medical orders, and procedures provided to the patient during the ICU stay. For example:

Radiology: Contains all radiology performed on a particular patient.

Radiology results: Contains the results of each radiology test performed on the particular patient.

Drugs: Contains all relevant information for all the drugs that a patient has been administered.

5 Laboratory: Contains all laboratory tests ordered for a patient.

Microbiology result: Contains the results of microbiology organisms taken on a patient.

10 Laboratory result: Contains the results for a laboratory test ordered for a particular patient.

#### DETAILED DESCRIPTION OF THE INVENTION

15 The present invention is a system and method for remote monitoring of ICU's from a distant command center/remote location. By monitoring a plurality of ICU's remotely, intensivists can better spread their expertise over more ICU beds that heretofore achievable. The presence of 24-hour a day/7 day-per-week intensivist care dramatically decreases the mortality rates associated with ICU care.

Referring to FIGS. 1A and 1B, the Billing and Demographic data structure of the present invention is illustrated. Patient demographic information **9010** is collected on the particular patient. This information comprises all the typical kinds of information one would normally gather on a patient such as first name, last name, telephone number, marital status, and other types of information. Patient insurance information **9012** is collected and associated with the patient demographic information **9010**. Patient insurance information **9012** relates to information on the type of accident and related information such as employment, employer name, place of service, and other information that would relate to the accident that actually occurred (if at all) and which would have to be reported to an insurance agency. This information is associated with the patient demographic information which assigns the unique patient ID to the particular patient.

Insurance plan information **9008** is also created and stored and comprises insurance carrier ID's, the plan name, policy number, and group number. This information on the insurance plan **9008** is also associated with the patient ID and demographic information **9010**.

Physician information **9002** is also created and stored for each physician associated with the system of the present invention. Information such as first and last name, credentials, and other information concerning the physician is saved. In addition, the physician's role is identified **9004** and information concerning the physician and the physician's role is associated with the particular patient via the patient ID stored in the demographic information **9010**.

Patients are entered into the hospital by a hospital representative **9006** who has a representative ID which also is ultimately associated with the patient ID. In addition, communications data **9000** is stored concerning how a representative can be reached (cell phone, home phone etc.).

Referring now to FIG. 1B, the Overall Billing and Insurance data structure is illustrated. An insurance provider number **9014** is also stored in the system. Each physician is given a provider number and provider ID by each insurance company. Thus data must be stored regarding the ID that is given to a particular physician by each insurance provider. This information is also stored and can be associated ultimately with treatment of the patient.

Each patient admitted to the hospital and to the ICU has a patient visit ID associated with the patient **9017**. This visit ID has patient ID information, ICU information, admission

date, and other information relevant to the specific visit. This information is illustrated in FIG. 1B. The visit ID **9017** is associated with the patient ID **9010** so that each visit can be tracked by patient.

Insurance carrier information **9018** is stored by the system and is associated with the insurance plan information **9008** as appropriate. Thus the particular insurance carrier with its name, address, and other identifying information **9018** is associated with the type of plan **9008** carried by the patient. The insurance carrier information **9018** together with the insurance plan information **9008** is associated with the patient via the patient ID information **9010**.

Patient address information **9020** and **9022** are collected for each individual patient and associated with the patient demographic information **9010**. If there is a patient guarantor, this information is obtained and stored with information on the guarantor **9026**. Such information as the guarantor's first and last name, date of birth, and other information is stored and is illustrated in FIG. 1B. Further, the guarantor's address **9024** is also collected and ultimately associated with the patient demographic information **9010**.

Referring to FIGS. 2A and 2B, the Command Center logical data structure is illustrated. The various information associated with demographic and insurance information is again used to manage the care and operations of the command center. Therefore, communications information **9000** is combined with physician and physician extender (i.e. nurse, LPN and the like) information **9002** and physician role **9004** to be associated with the demographic information **9010**. The patient visit information **9017** together with this information is associated with the patient's location which has a unique identifier **9030**. Each location ID has patient ID information and visit ID information associated with it.

Referring now to FIG. 2B, the Command Center logical data structure illustration continues. Each ICU bed has an associated location ID which comprises hospital ICU information, room number, and bed number **9038**. In addition, and as described earlier, instrumentation such as cameras are also associated with the particular patient. Therefore the camera setting **9040** will have a location ID relating to the ICU bed as well as have camera value settings and associated camera identifier information.

Each ICU bed **9038** is associated with an ICU **9032**. Each ICU has information associated with it that uniquely identifies the ICU as being associated with the particular hospital, and having particular phone numbers, fax numbers, work space addresses, and other information, that help to identify the ICU.

As noted above, each ICU is associated with a hospital **9034**. Each hospital has a unique identifier, as well as its own name, address, and other identifying information. Further, since each hospital ICU is to be coordinated through a remote command center, information on the remote command center **9036** is associated with the hospital information. Each command center has a unique ID and has associated address information stored as well.

Thus in the Command Center logical data structure, patient ID information **9010** is linked to a patient location **9030** which in turn is associated with an ICU bed **9038** each of which beds are uniquely associated an ICU **9032** which is associated with a hospital **9034** which in turn has the ICU managed by a command center **9036**.

An integral part of the system of the present invention is the recording of medical history. Referring to FIG. 3, the logical relationship among data elements for medical history is illustrated. Patient visit information **9017** combined with the physician-physician extender information **9002** is com-

bined with specific note-taking information **9042**. The note information comprises the date and time the notes are taken as well as the note type. The note ID is fed information from the medical history item **9044**, which has its own unique medical ID associated with it. This information comprises medical text, category of information, and other information relevant to the medical history. As noted, this information for medical history **9044** is associated with a note ID **9042**, which in turn is associated with the patient visit and physician information **9017** and **9002**.

Referring to FIG. 4A, 4B, and 4C, the note-keeping logical data structure of the present invention is illustrated. As noted earlier, the note ID **9042** combines information from visit ID, treating physician, and other information relating to the time the note was entered. Other information is associated with the note ID. Referring first to FIG. 4A, the patient visit information **9017**, is associated with the note ID **9042**. Various procedural information **9046** is kept by the system of the present invention and is associated with the visit ID **9017**. Physicians are able to create free text patient illness notations **9048** and associate them with the note **9042**. Similarly, free text information regarding functioning of the system **9050** is permitted and also associated with notes regarding the particular patient and procedure **9042**.

Specific notes regarding, for example, surgical procedures are also kept. Surgery notes **9054** are associated with a particular note ID and have such information as anesthesia, surgical diagnosis, elective information, and other related surgical information. Surgical fluids **9052** administered during the course of surgery are associated with the surgery information **9054**. Additionally, any surgical complications **9056** are noted and also associated with the surgery which in turn has an associated note ID.

Referring now to FIG. 4B, the logical data structure for notes and its description is continued. An assessment plan **9058** is created and associated with the same note ID for the particular patient. The plan has a free text field that allows a physician to create the appropriate assessment plan and associate it with a note ID **9042**.

Various daily notes are also kept and associated with the individual note ID **9042**. For example, the daily mental state **9060** is recorded to document the mental state of the patient. The daily treatment **9062** administered to the patient is associated with the unique note ID. The daily diagnosis **9068** is also created and associated with unique note ID **9042**.

Any unstable conditions are also noted **9070** and records kept of those conditions. Similarly mortality performance measures (MPM) information **9072** is kept and associated with the unique note ID. To the extent that any physical exam **9074** is administered, that physical exam and any free text created by the physician is associated with the unique ID and records kept. Allergy information **9076** for the particular patient is also created and stored along with the allergy type, and allergy name. This information is uniquely associated with the note ID. Referring now to FIG. 4C, the Logical Data Structure for the Notes Creation and Storage description is continued. A specific note item record **9078** is also kept and associated with unique note ID. This note item comprises the principal diagnosis, the chief complaint, the past history of the patient, the reason for the note, and various other identifications and flags of information which help in documenting the patient's condition.

Any drugs that are administered to the patient, including dosage, type, and number **9086** is kept and associated with the unique note ID **9042**.

Procedural note items are also documented **9082**. Procedural notes involve the procedural type, the principal diag-

nosis, the procedural location, procedural indications, and other information of a procedural nature. Procedural description information **9088** is kept as input to the procedural note item. This information is also associated with a procedural evaluation **9084** which comprises text describing the procedural evaluation that occurred. These three items, the procedural description **9088**, procedural evaluation **9084**, and procedural note items **9082**, are all uniquely associated with the note ID **9042**.

Referring now to FIG. 5, the Logical Data Structure of the Medical Order Functionality of the Present Invention is illustrated. Each medical order **9092** has a unique order ID associated with it. This information derives its uniqueness from the visit ID, the representative ID, and various information about the date in which the order was created and other such relevant information. Any non-drug orders **9090** are associated with a unique non-drug order ID. The order is classified, identified, and free text can be created by the physician to describe the order. This information in the non-drug order **9090** is associated with the unique medical order for that particular patient **9092**.

Again physician and physician extender identification information **9002** is also uniquely associated with the medical order to identify the physician involved in creating the particular order in question.

Drug orders **9094** are created each with its own unique drug order ID. Various information is collected as part of the drug order including the type of drug, the dosage, start date, frequency, stop date, to name but a few elements typical of a drug order. The drug order information **9094** is associated with the unique medical order ID **9092** assigned to that particular patient. All of the medical order information is associated with patient visit information **9017** which allows that information to be uniquely identified with a particular patient for a particular visit.

Referring again to FIG. 4C, the system is also capable of annotating and storing various log items **9080**. For example, an event log item is given a number, a patient profile item has its own number, as do neurological, cardiographic, pulmonary, renal, and other events can have log items associated with them and may be used as input to any of the note taking of the present invention.

Referring to FIGS. 6A and 6B, the logical data structure of the patient care functionality of the present invention is illustrated. Each patient visit with its unique ID **9017** has a number of other pieces of information associated with it. For example, physician-patient tasks are tracked **9098** and have a unique task ID associated with them. The patient code status **9096** is documented and associated with the physician-patient task **9098** task ID. This information is uniquely associated with the patient visit via the the patient visit ID **9017**.

Laboratory information **9100** has a unique lab ID associated with it. That information is keyed to the visit ID and records the specimen taken, the date it was taken, and various other information germane to the laboratory procedure involved. Other lab procedures **9102** are also documented with another unique ID. "Other" lab ID is associated with the laboratory ID **9100** which again is uniquely associated with the particular patient.

Microbiological studies **9104** are documented together with the date and the date taken and the type of study involved. Any study of microorganisms **9106** is documented with a unique microorganism ID. Micro sensitivities **9108** which record the sensitivity to microorganisms and certain antibiotics is recorded and associated with the microorganism ID **9106**. This information in turn is associated with a

microbiological study **9104**, all of which is associated with the unique patient visit ID **9107**.

Respiratory studies **9101** are also recorded with unique identification numbers and a description. This information is again associated with the patient visit ID **9017**.

Referring now to FIG. 6B, the logical data structure of the patient care functionality of the Present Invention is further illustrated. Other organism studies **9118** are also conducted to determine any other conditions associated with microorganisms that might exist with the particular patient. This other organism information **9118** is associated with the microorganism studies **9106** which in turn is associated with the microbiology category of information of the present invention **9104**.

Various diagnostic imaging also takes place and is recorded. This image information **9114** has unique image ID associated with each image and comprises associated information such as the image type, the date performed, and other information relevant to the diagnostic imagery. The result of the image taken **9116** is also uniquely identified with the image ID and a unique image result ID. This information is associated with the image information **9114** which again is uniquely associated with the patient visit ID.

Various intake and output for the patient's biological functioning is recorded **9110**. Intake and output total **9112** is recorded and uniquely associated with the intake/output identification note **9110**. Intake/output totals **9112** also comprised the weight the total taken in, the total out, and five-day cumulative totals for biological functioning of the particular patient.

Referring to FIG. 7, The Logical Data Structure Concern with Reference Information for the present invention is illustrated. This data structure allows only certain ranges of data to be input by care givers into the system. This is accomplished by having categories of information **9120** each category capable of having only certain values. Similarly, each type of data **9126** associated with each category is only permitted to have certain values. This combination of Category and Type results in a Combined ID **9122** which can be used in combination with certain values **9128** to create a value and combination **9124** that can be presented to a care giver viewing and entering data. This effectively limits errors in data entry by only allowing certain values to be entered for given types of data. For example, if only milligrams of a medication are supposed to be administered, this data structure prevents a care giver from administering kilograms of material since it is not a permitted range of data entry. The "nextkey" function **9027** is the function that keeps track of the ID's that are given during the administration of the present invention. This function insures that only unique ID's are given and that no identical ID's are given to two different patients for example.

Referring to FIG. 8A, the Logical Data Structure of the Vital Signs Functionality of the Present Invention is illustrated. Vital sign header information **9120** is created and uniquely associated with the visit ID for the particular patient. This header information comprises a date-time stamp combined with hospital information, medical reference numbers, and identification of the patient. Vital sign details **9122** are also created and uniquely date-time stamped and associated with the particular visit ID for the patient. This information comprises all manner of vital sign information relating to blood pressure, respiration, and other factors. Vital sign information is associated with the patient visit **9017** and the demographic information concerning the patient **9016**. Such associations of information can be the basis for later studies.

Referring to FIG. 8B, Additional Vital Sign Logical Data Structures are illustrated. For example, a vital sign log header **9120** is created using the unique hospital ID and medical record numbers. Other information, such as a patient name and date-time stamp, are also stored. Vital sign log details **9124** are created and associated with the vital sign log header **9120**. For example, blood pressure measurements, respiration, and other factors are all detailed for a particular hospital ID. It should be noted that all vital sign data is logged in and kept by the systems of the present invention. Where vital sign information is received but cannot be associated with a particular patient, such communications are noted as errors.

Vital sign error details **9126** are also recorded and associated with a particular hospital. Information and the vital sign error detail also comprises heart rate, blood pressure, and other information. This information is associated with a vital sign error header **9130** which is associated with the hospital identifier and the patient first and last name and other information. Various vital sign error codes **9128** exist with the present invention and are used in association with the vital sign error detail **9126**. This information however relates to communications of vital sign data that are deemed "errors" as noted above.

Care Net patient location **9132** is recorded and associated with a particular hospital ID and location ID for the particular patient. Carenet is a proprietary product designation of Hewlett-Packard and is kept by the system of the present invention since it identifies the equipment from which measurements come. The ICU bed information **9038** is associated with the Care Net patient location **9132**.

Referring to FIG. 9, the distributed architecture of the present invention is shown. In concept, the distributed architecture comprises a headquarters component **200**, a command center/remote location **202**, and a hospital ICU **204**, which, while represented as a single hospital in this illustration, in the preferred embodiment comprises several hospital ICUs at different locations. The headquarters unit **200** comprises a database server and data warehouse functionality, together with a patient information front end. The patient information front end **206** provides patient specific information to the command center/remote location. The database server/warehouse function **208** comprises the amassed information of a wide variety of patients, in their various conditions, treatments, outcomes, and other information of a statistical nature that will assist clinicians and intensivists in treating patients in the ICU. The headquarters' function also serves to allow centralized creation of decision support algorithms and a wide variety of other treatment information that can be centrally managed and thereby standardized across a variety of command center/remote locations. Further, the database server/data warehousing functionality **208** serves to store information coming from command center/remote locations replicating that data so that, in the event of a catastrophic loss of information at the command center/remote location, the information can be duplicated at the command center/remote location once all systems are up and running.

At the hospital ICU **204**, each patient room **232**, **234** has a series of bedside monitors and both video and audio monitoring of each patient in the patient room. Each ICU further has a nurse's station with a video camera and monitor **230** so that videoconferencing can go on between the nurses and doctors at the nursing station and those intensivists at the command center/remote location. The monitoring equipment at the ICU is served by a monitor server **236**, which receives and coordinates the transmission of all bedside

monitoring and nurses station communication with the command center/remote location. Finally, each ICU has a patient information front end **228**, which receives and transmits to the command center/remote location information concerning the identity and other characteristics of the patient.

Command center/remote location **202** comprises its own video capture and monitoring capability **212** in order to allow the intensivists to view the patients and information from the bedside monitoring as well as to have videoconferencing with the nursing station and with patients as the need arises. Information from the monitor server **236** at the hospital ICU is served to an HL7 (the language for transmitting hospital/patient/diagnostic data) gateway **214** to a database server **222**. In this fashion, information from the bedside monitors can be stored for current and historical analysis. Monitor front ends **216** and **218** allow technicians and command center/remote location personnel to monitor the incoming data from the patient rooms in the ICU. Information from the patient information front end **228** is provided to an application server **224**, having its own patient information front end **226** for aggregating and assembling information in the database **222** that is associated with individual patients in the ICU.

It is expected that there will be a great deal of concurrent hospital data that is necessary to the implementation of the present invention. It is therefore expected that there will be a legacy database system **210** having a front end **220** from which intensivists and command center/remote location personnel can retrieve legacy database information.

Referring to FIG. 10, a system architecture of one embodiment of the present invention is illustrated. Headquarters **200** comprises an application server **238**, an NT file server **240**, and Sun SPARC Enterprise 250242 and Enterprise network management system **244**, a Cisco 3600 router **246**, a Cisco 2924 switch **248**, and a hot phone **250**. The application server **238** is designed to monitor and update those applications used at the command center/remote location. The NT file server serves to monitor, store, and replicate information coming from the command center/remote locations. The SPARC Enterprise **250** server **242** is a disc storage server, for storing and serving information, such as practice guidelines, algorithms, patient information, and all matter of other information records that must be stored in order to support the present invention. As explained below, the SPARC Enterprise **250** server and other components are such as routers and switches are commonly used in the ICU, the command center/remote location, and the headquarters. For example:

The Cisco 3600 router is a multi-function device that combines dial access, routing, and local area network (LAN) to LAN services, as well as the multi-service integration of voice, video, and data in the same device. This is necessary, since the various command center/remote locations, headquarters, and intensive care units all must integrate and transmit video, audio, and data among the various entities.

The Cisco 7204 is a router which provides high speed LAN interconnect, virtual private networks, and Internet access, all of which is required for providing the communication in the network of the present invention; and

The Cisco 2924 switch is an autosensing fast ethernet switch, allowing networked multimedia and virtual LAN support. Multi-level security is also offered in the switch to prevent unauthorized users from gaining access and altering switch configuration. These components are also identified in the figures (below).

The particular commercial systems named here are given as but some examples of equipment available today. The function of these equipment is the important factor. Other similar or improved equipment can also be utilized.

The network management system **244** allows the entire traffic and condition of the network to be monitored and to allow maintenance to take place. The router **246** and switch **248** is used for communication with the various command center/remote locations that are served by the Headquarters component. The Headquarters component interacts via frame relay with the command center/remote location **202**.

Command center/remote location **202** comprises an applications server **262** for the purpose of running various applications for the intensivists and command center/remote location staff. The NT file server **264** at the command center/remote location allows patient files, historical files, algorithms, practice standards, and guidelines, to be served to the clinicians and intensivists to assist in monitoring the patients. The Sun SPARC Enterprise 250266 is used to for storage purposes as noted above. The Enterprise network management system **268** monitors the overall health of the network of command center/remote locations and intensive care units as well as the functionality of the individual pieces of equipment within the command center/remote location. A Cisco 2924 switch **256** and Cisco 7204 router **258**, combined with the Cisco 3600 router **260** allows for point to point communication over a T1 line, with a plurality of intensive care units located remotely from the command center/remote location. Hot phones **252** and **254** allow communication with the headquarters and the intensive care unit.

Intensive care unit **204** comprises a Cisco 2924 switch **272** for the purpose of interfacing with the various audio-video feeds **274**, **276** from the various patient rooms and the nursing station. A local work station **280** is connected to a scanner **282** which allows data to be input, scanned, and communicated via the point to point T1 communications to the command center/remote location. Further, the workstation **280** provides for textual advice and patient orders to be delivered to the intensive care unit for execution. The intensive care unit also comprises a laser printer **284** for the printing of patient orders and other information relevant to the care of intensive care patients. Referring to FIG. **11**, the videoconferencing/surveillance/imaging components of the present invention are illustrated. The hospital ICU **204** comprises a series of video cameras **290**, which are located in patient rooms and at the nurse's station. Control for the cameras is provided through an RS424 to RS232 converter **288**, with instructions for imaging emanating from the workstation at the command center/remote location **252** through the ICU workstation **280** through a multi-port serial controller **286**. Video feed from the video cameras **290** is provided to an audio-video switcher **292**, which in turn provides its output to the multi-port serial controller **286** for subsequent viewing at the nurse's station and at the command center/remote location. Of equal importance is a microphone feed from the patient and from the nurses. That microphone **296** provides its signal to an audio line amplifier **294**, which in turn provides an audio feed to the audio-video switcher **292**. In this way, a patient can provide information, as can nurses who are visiting the patient during the course of patient care. It is also important that information of an audio nature be fed to the intensive care unit, both to the patient rooms and to the nurse's station. To do this, the multi-port serial controller **286** provides an audio signal to a reverse audio switcher **298**, which in turn provides information to speakers **300** that are located at the nurse's station as well as at the bedside of the patients. Information to the

reverse audio switcher is provided an audio amplifier **302** from information from a video code **304**, which in turn is connected to the workstation at the ICU. As noted earlier, a scanner **282** is provided, so that information can be scanned and provided to the command center/remote location **202** and a hot telephone **278** communicates with a telephone **252** at the command center/remote location.

Referring to FIG. **12** the vital signs data flow is illustrated. The monitoring system at each ICU bedside comprises a monitoring system for monitoring the vital signs for the patient. The vital sign monitoring system **450** captures vital sign data **452** and transmits that vital sign data **454** using the HL7 language (the standard processing language for hospital data and information). The processor at the ICU processes the vital sign data for transmission and storage purposes and transmits that information to the remote location. Vital sign data is then loaded into the data base **458**. The data base for each individual patient is then reviewed and process rules are applied **460** to the vital sign data. These process rules relate to certain alarming conditions which, if a certain threshold is reached, provides an alarm to the intensivist on duty. The vital sign alarm **462** is then displaced to the intensivist who can then take appropriate action. A typical type of rule processing of the vital sign data might be if blood pressure remains at a certain low level for an extended period of time, or if heart rate remains high for an extended period of time. In addition a wide range of other rules are provided which will provide an audible alarm to the intensivist before a critical situation is reached.

In addition to the information being provided to the alarming system for the intensivist, the vital sign data **464** is also transmitted **466** into a database warehouse **468** comprising vital sign data **470** from not only the individual patient but from all of the patients being cared for in the ICU. This database warehouse provides the ability to do data mining for trends that can give rise to additional process rules and vital sign thresholding. In addition to the transmission of vital sign data **454** to the remote site, the vital sign data is displayed in real time at the ICU **472**.

Referring to FIG. **13A** the diagnostic imaging interaction is illustrated. X-rays for example, are created and transmitted to the command center **472**. Additionally, the information could be ACT scan, MRI, or any other method of medical diagnostic imaging. The x-ray image is captured at the command center **474** where it is stored and in addition displayed on the image monitor **476** for the intensivist to review.

Referring to FIG. **13B** the interactive video session is illustrated. A video conferencing session is established **478** regarding a particular patient in an ICU bed. Using the video cameras in each room and/or at the nurse's station at the ICU, the patient and/or the nurse can be viewed **480**. On the other end of the video conferencing session is the intensivist who can then both visually and orally communicate with the patient and/or nurse **482**.

Referring to FIG. **14** the physician resources and order writing data interface is illustrated. The user interface **484** allows the physicians to access physician resources **486**. These resources provide guidelines for the treatment of the critically ill. In this example the intensivist is requested to enter the antibiotic associated with colitis **488**. The system then generates a request for a fecal leukocyte test **490**. This request is translated into an order writing module **496** which results in the actual order for the test **502**. Since the order needs to be transmitted to the appropriate organization for execution, an appropriate order is generated to the microbiology laboratory **500** in this instance. The order results are

then achieved **506** and the completion of the order is reported to the order writing assignment manager **496**. In addition, the order writing module **502** also results in a task list **504** of orders for various other individuals in laboratories. In addition, user interface **484** allows the physician to re-enter the physician resources module at any particular location with results of the tests. These tests are then fed into the system to continue with the diagnostic algorithm processing of the patient test results **494**. The user interface also allows interaction with the resident database **498**.

Referring to FIG. **15** the physician resources database data interface is illustrated. User interface **508** allows the intensivist to interact with the physician resources data base **510**. In this example, resident data base **524** which comprises the identification and background of the resident admitting the patient causes an admission diagnosis **526** to be created. In this example a diagnosis of pancreatitis is illustrated. This diagnosis of pancreatitis **522** alerts the physician resources module **510** which causes an entry for the topic pancreatitis **512**. The diagnosis algorithm for pancreatitis **514** is then retrieved and a request for an Apache II score **516** is requested. The system also requests information for operative data **528** describing what if any operations have taken place with respect to this patient, vital sign data **530**, request for laboratory information **532**, past medical history for the patient **534** and patient demographics **536**. All this information is provided to the Apache II score assignment manager **538** which assigns an Apache II score based upon weighted composite up to twenty five different variables. This Apache II score is provided to the Apache II score request module **516**. If the severity based Apache II score is greater than or equal to eight the diagnostic of the system continue **520**. If the Apache II score is less than eight, the patient is triaged to a non-ICU bed **518** since the patient will not necessarily require intensive care thereby saving relatively scarce resources of the ICU for those who are truly critically ill.

Referring to FIG. **16** the automated coding/billing work flow and data flow is illustrated. Clearly ICUs must be paid for the care that they give. At the outset of the visit **540** the user interface **542** allows for the input of International Classification of Diseases, Ninth Revision (ICD 9) diagnosis code information concerning complexity of the case, whether the patient is stable, whether the physician involved is the attending physician or consulting physician and all other manner of information required for billing purposes. In addition, resident data **544** is input such as patient demographics, insurance information, physician, guarantor, the date that the service is provided. All this information is provided to the data manager **546** which assembles the required data element for subsequent processing. The data manager sends the demographic, physician, guarantor, insurance and related information to a bill generator **548** which begins to assemble the information to subsequently generate a bill. Clinical information is provided to the current procedural terminology (CPT) code assignment manager which assigns codes based upon the scores and user input for bill generation purposes. A history of present illness (HPI) score **560** is generated along with a review of systems (ROS) score **562**. A past, family, and/or social history (PFSH) score **564** is generated along with a score relating to the physical exam **566**. A mortality prediction model (MPM) score **568** which is a score relating to the severity of the illness is also generated. All of these various scores are provided to the CPT assignment manager **558**. Periodically information is downloaded for management reports **556**. Once all of the information for the CPT code assignment is generated that

information is provided to the bill generator **548** which assembles all the data elements needed to generate a Health Care Financing Administration (HCFA) 1500 claim form. The input for the bill generator is then verified **550** where the physician can disagree with code assignments return progress notes and generally review the bill. This smart processing of the HCFA 1500 claim form allows for fewer mistakes to be made. If there is any error or additional information that is required, the verification process fails the proposed claim form and information regarding that failure is provided back to the resident data for completion of any missing items. Once an invoice has been verified as having the appropriate information to be submitted the HCFA 1500 claim form is generated **554**. Additional information is written to a billing data file **552** for importation to the patient accounting system of the present invention.

Referring to FIG. **17** the order writing data flow is illustrated. Order entry user interface **600** allows the intensivist to order procedures and medication to assist the patients in the ICU. For example, the intensivist can order an ECG **604**. Thereafter the order is reviewed and a digital signature relating to the intensivist is supplied **606**. Once reviewed and signed off, the order is approved **607** and sent to the data output system **610**. Thereafter the data output system prints the order to the printer in the ICU **616**. For record keeping purposes the order is exported in the HL7 language to the hospital data system **618**. In addition the data output system adds an item to the data base that will subsequently cause an intensivist to check the ECG results. This notification to the task list is provided to the database **614**. In addition, as part of the database an orders file relating to the specific patient is also kept. The fact that an ECG has been ordered is entered in the orders file for that patient.

In a similar fashion using the order entry user interface **600** the intensivist can order medications **602** for a patient. The medication order then is provided to an order checking system **608**. The order checking system retrieves information from the database **614** relating to allergies of the patient and medication list which includes medications which are already being administered to the patient. This allows for the order checking system to check for drug interactions. Further laboratory data is extracted from the database **614** and the order checking system checks to insure that there will be no adverse impact of the recommended dosage upon the renal function of the patient. Once the order checking system **608** is completed, the order is okayed and provided to the order review and signature module **606**. In this module the digital signature of the intensivist is affixed to the order electronically and the order is approved **607**. Thereafter it is provided to the data output system **610** where again the orders are printed for ICU and **616** and for the hospital data system. In this case, any medications that are ordered are then provided to the medications list file in the database **614** so that the complete list of all medications that are being administered to the ICU patient is current.

Referring to FIG. **18** the event log is illustrated. The database **620** contains all manner of notes and data relating to the particular patient that is admitted to the ICU. For example, admission notes **622** are taken upon admission of the patient and stored in the file that is specific to that patient. Progress notes **624** are created during the patients stay within the ICU to note the progress the patient is making giving the various treatments. Procedural notes **626** are also created by the intensivist to note what procedures have taken place and what if any events have occurred associated with those procedures. Laboratory data such as positive blood

cultures are also stored in the file 628 in the database 620. Further x-ray data 630 and abnormal CT Scan results are stored in the database.

The result of these individual files are then provided to an event log manager 632. For example, admission notes might contain operations performed. Progress notes 624 might relate to the operations performed. This information is provided to the event log manager 632. Admission information is also input to the event log manager as are a listing of the procedures administered to the patient. To the extent there are positive blood cultures in the laboratory data 628 those are provided to the event log manager 632 as are abnormal CT scan results. All of this information is made available through the user interface 634. Thus the event log presents in a single location key clinical information from throughout a patient's stay in the ICU. The event log user interface provides caregivers with a snapshot view of all salient events since admission. All relevant data on procedures and laboratory tests, etc. are presented chronologically.

Referring to FIG. 19 the smart alarms of the present invention are illustrated. The smart alarm system constantly monitors physiologic data (collected once per minute from the bedside monitors) and all other clinical information stored in the database (labs, medications, etc). The periodicity of the collection of data is stated for illustrative purposes only. It is well within the scope of the present invention to collect physiological data at more frequent time intervals. Thus, monitor 636 provides information in HL7 form to the interface engine 638. The physiological data is then formatted by the interface engine for storage in the database 640 where all patient information is maintained. The rules engine 642 searches for patterns of data indicative of clinical deterioration.

One family of alarms looks for changes in vital signs over time, using pre-configured thresholds. These thresholds are patient-specific and setting/disease-specific. For example, patients with coronary artery disease can develop myocardial ischemia with relatively minor increases in heart rate. Heart rate thresholds for patients with active ischemia (e.g. those with unstable angina in a coronary care unit) are set to detect an absolute heart rate of 75 beats per minute. In contrast, patients with known coronary artery disease in a surgical ICU have alarms set to detect either an absolute heart rate of 95 beats per minute or a 20% increase in heart rate over the baseline. For this alarm, current heart rate, calculated each minute based on the median value over the preceding 5 minutes, is compared each minute to the baseline value (the median value over the preceding 4 hours). Physiologic alarms can be based on multiple variables. For example, one alarm looks for a simultaneous increase in heart rate of 25% and a decrease in blood pressure of 20%, occurring over a time interval of 2 hours. For this alarm, thresholds were initially selected based on the known association between changes in these two variables and adverse clinical events. Actual patient data were then evaluated to determine the magnitude of change in each variable that yielded the best balance between sensitivity and specificity. This process was used to set the final thresholds for the rules engine.

Alarms also track additional clinical data in the patient database. One alarm tracks central venous pressure and urine output, because simultaneous decreases in these two variables can indicate that a patient is developing hypovolemia. Other rules follow laboratory data (e.g. looking for need to exclude active bleeding and possibly to administer blood).

The purpose of the rules engine is to facilitate detection of impending problems and to automate problem detection thereby allowing for intervention before a condition reaches a crisis state.

Referring to FIG. 20 the procedural note-line log is illustrated. This log allows clinicians to evaluate the likelihood that a given procedure might result in further complications. In this example presented in this FIG. 20 a catheter removal is illustrated. When a new catheter is inserted in a patient 648 a procedural note is created on the procedure note creation user interface 646. The note is reviewed and a digital signature is attached to the note to associate the note with a particular intensivist 654. The procedure is then approved and is provided to the data output system 656. The procedural note is then printed on the printer in the ICU 658 and is exported in HL7 language to the hospital data system 660. In addition, this also triggers a billing event and the data output system provides appropriate output to the billing module 662 to generate an invoice line item. In addition, the note is stored in the emergency medical record associated with the patient in the database 664. In addition, the line log is updated in the database 664 to show what procedure was administered to a patient at what time. If there is an existing catheter, that is displayed to the intensivist at the procedure note creation user interface 646. This would show an existing catheter changed over a wire 650. That information is provided to the line id module 652 which extracts information from the line log in the database 664. This information results in a note being created and provided to the note review and signature module 664. Thus the line log contains, for each patient, relevant information about all in-dwelling catheters, including type and location of the catheter, insertion date, the most recent date that the catheter was changed over a wire, and the date the catheter was removed. This information helps clinicians evaluate the likelihood that a given catheter is infected and guides its subsequent management of that procedure.

Evidence-based Guidelines, Algorithms, and Practice Standards Decision Support Algorithms

In order to standardize treatment across ICUs at the highest possible level, decision support algorithms are used in the present invention. These include textual material describing the topic, scientific treatments and possible complications. This information is available in real time to assist in all types of clinical decisions from diagnosis to treatment to triage.

All connections among components of the present invention are presently with a high bandwidth T-1 line although this is not meant as a limitation. It is anticipated that other existing and future high bandwidth communication capabilities, both wired and wireless, as well as satellite communications will be suitable for the communications anticipated for the present invention.

As noted earlier, a key objective of the present invention is to standardize care and treatment across ICUs. This is effective in the present invention by providing decision support to intensivists as well as information concerning the latest care and practice standards for any given condition. As noted in Table I below, a wide variety of conditions is noted. Each of the conditions has an associated guideline of practice standard that can be presented to the intensivist who might be faced with that particular condition in a patient. These guidelines of practice standards can be accessed at the command center/remote location or at the ICU to assist in the treatment of the patient. Thus, the general categories of cardiovascular, endocrinology, general, gastrointestinal,

hematology, infectious diseases, neurology, pharmacology, pulmonary, renal, surgery, toxicology, trauma all have guidelines and practice standards associated with them.

TABLE 1

EVIDENCE-BASED GUIDELINES ALGORITHMS & PRACTICE STANDARDS DECISION SUPPORT
<u>CARDIOVASCULAR</u>
BRADYARRHYTHMIAS
CARDIOGENIC SHOCK
CARDIO-PULMONARY RESUSCITATION GUIDELINES
CONGESTIVE HEART FAILURE
EMERGENCY CARDIAC PACING
FLUID RESUSCITATION
HYPERTENSIVE CRISIS
IMPLANTABLE CARDIO-DEFIBRILLATOR INDICATIONS
INTRA-AORTIC BALLOON DEVICE INDICATIONS
MAGNESIUM ADMINISTRATION IN PATIENTS
MANAGEMENT OF HYPOTENSION, INOTROPES
MYOCARDIAL INFARCTION
MI WITH LEFT BUNDLE BRANCH BLOCK
PA CATHETER GUIDELINES & TROUBLE-SHOOTING
PERMANENT PACEMAKER INDICATIONS
PULMONARY EMBOLISM DIAGNOSIS
PULMONARY EMBOLISM TREATMENT
SUPRA-VENTRICULAR TACHYARRHYTHMIAS
UNSTABLE ANGINA
VENOUS THROMBOEMBOLISM PROPHYLAXIS
VENOUS THROMBOSIS: DIAGNOSIS & TREATMENT
VENTRICULAR ARRHYTHMIAS
<u>ENDOCRINOLOGY</u>
ADRENAL INSUFFICIENCY
DIABETIC KETOACIDOSIS
HYPERCALCEMIA: DIAGNOSIS & TREATMENT
HYPERGLYCEMIA: DIAGNOSIS & TREATMENT
STEROID REPLACEMENT STRATEGIES
THYROID DISEASE
<u>GENERAL</u>
DEALING WITH DIFFICULT PATIENTS AND FAMILIES
END OF LIFE DECISIONS
ETHICAL GUIDELINES
PRESSURE ULCERS
ORGAN PROCUREMENT INDICATIONS AND SALVAGE
<u>GASTROINTESTINAL</u>
ANTIBIOTIC ASSOCIATED COLITIS
HEPATIC ENCEPHALOPATHY
HEPATIC FAILURE
MANAGEMENT OF PATIENTS WITH ASCITES
NUTRITIONAL MANAGEMENT
ACUTE PANCREATITIS
UPPER GI BLEEDING: STRESS PROPHYLAXIS
UPPER GI BLEEDING: NON-VARICEAL
UPPER GI BLEEDING: VARICEAL
<u>HEMATOLOGY</u>
HEPARIN
HEPARIN-INDUCED THROMBOCYTOPENIA
THE BLEEDING PATIENT
THROMBOCYTOPENIA
THROMBOLYTIC THERAPY
TRANSFUSION GUIDELINES
USE OF HEMATOPOETIC GROWTH FACTORS
WARFARIN
<u>INFECTIOUS DISEASES</u>
ACALCULUS CHOLECYSTITIS
ANTIBIOGRAMS
BLOODSTREAM INFECTIONS
CANDIDURIA
CATHETER RELATED SEPTICEMIA
CATHETER REPLACEMENT STRATEGIES
ENDOCARDITIS PROPHYLAXIS
ENDOCARDITIS DIAGNOSIS AND TREATMENT
FEBRILE NEUTROPENIA
FUO

TABLE 1-continued

EVIDENCE-BASED GUIDELINES ALGORITHMS & PRACTICE STANDARDS DECISION SUPPORT	
5	HIV+ PATIENT INFECTIONS
	MENINGITIS
	NECROTIZING SOFT TISSUE INFECTIONS
	NON-INFECTIOUS CAUSES OF FEVER
	OPHTHALMIC INFECTIONS
10	PNEUMONIA, COMMUNITY ACQUIRED
	PNEUMONIA, HOSPITAL ACQUIRED
	SEPTIC SHOCK
	SINUSITIS
	SIRS
	TRANSPLANT INFECTION PROPHYLAXIS
15	TRANSPLANT-RELATED INFECTIONS
	<u>NEUROLOGY</u>
	AGITATION, ANXIETY, DEPRESSION & WITHDRAWAL
	DIAGNOSIS AND TREATMENT
	BRAIN DEATH
	GUILLAIN-BARRE SYNDROME
20	INTRACEREBRAL HEMORRHAGE
	MYASTHENIA GRAVIS
	NEUROMUSCULAR COMPLICATIONS OF CRITICAL ILLNESS
	NON-TRAUMATIC COMA
	SEDATION
	STATUS EPILEPTICUS
25	STROKE
	SUB-ARACHNOID HEMORRHAGE
	<u>PHARMACOLOGY</u>
	AMINOGLYCOSIDE DOSING AND THERAPEUTIC MONITORING
	AMPHOTERICIN-B TREATMENT GUIDELINES
30	ANALGESIA
	ANTIBIOTIC CLASSIFICATION & COSTS
	DRUG CHANGES WITH RENAL DYSFUNCTION IDENTIFICATION AND MANAGEMENT
	PENICILLIN ALLERGY
	NEUROMUSCULAR BLOCKERS
35	VANCOMYCIN
	THERAPEUTIC DRUG MONITORING
	<u>PULMONARY</u>
	ARDS: HEMODYNAMIC MANAGEMENT
	ARDS: STEROID USE
40	ARDS: VENTILATOR STRATEGIES
	ASTHMA
	BRONCHODILATOR USE IN VENTILATOR PATIENTS
	BRONCHOSCOPY & THORACENTESIS GUIDELINES
	CHRONIC OBSTRUCTIVE PULMONARY DIAGNOSIS AND TREATMENT
	CXR (INDICATIONS)
45	NONINVASIVE MODES OF VENTILATION INDICATIONS AND TREATMENT
	ENDOTRACHEAL TUBES & TRACHEOTOMY
	TREATMENT OF AIRWAY OBSTRUCTION
	VENTILATOR WEANING TREATMENT
	<u>RENAL</u>
50	ACUTE RENAL FAILURE: DIAGNOSIS
	ACUTE RENAL FAILURE: MANAGEMENT & TREATMENT
	DIALYSIS
	DIURETIC USE
	HYPERKALEMIA: ETIOLOGY & TREATMENT
55	HYPERNATREMIA: ETIOLOGY & TREATMENT
	HYPOKALEMIA: ETIOLOGY & TREATMENT
	HYPONATREMIA: ETIOLOGY & TREATMENT
	OLIGURIA
	<u>SURGERY</u>
60	OBSTETRICAL COMPLICATIONS
	DISSECTING AORTIC ANEURYSM
	POST-OPERATIVE HYPERTENSION
	POST-OPERATIVE MYOCARDIAL ISCHEMIA (NON-CARDIAC
	ARRHYTHMIAS AFTER CARDIAC SURGERY
	POST-OPERATIVE BLEEDING
	POST-OPERATIVE MANAGEMENT OF ABDOMINAL SURGERY
65	POST-OPERATIVE MANAGEMENT OF OPEN HEART SURGERY
	POST-OPERATIVE MANAGEMENT OF THORACOTOMY SURGERY

TABLE 1-continued

EVIDENCE-BASED GUIDELINES ALGORITHMS & PRACTICE STANDARDS DECISION SUPPORT
POST-OPERATIVE POWER WEANING
POST-OPERATIVE MANAGEMENT OF CAROTID SURGERY
WOUND HEALING STRATEGIES
<u>TOXICOLOGY</u>
ACETAMINOPHEN OVERDOSE
ANAPHYLAXIS
COCAINE TOXICITY
ALCOHOL WITHDRAWAL
HYPERTHERMIA
LATEX ALLERGY
UNKNOWN POISONING
<u>TRAUMA</u>
ABDOMINAL COMPARTMENT SYNDROME
BLUNT ABDOMINAL INJURY
BLUNT AORTIC INJURY
BLUNT CARDIAC INJURY
DVT PROPHYLAXIS
EXTREMITY COMPARTMENT SYNDROME
HEAD INJURY
HYPOTHERMIA
IDENTIFICATION OF CERVICAL CORD INJURY
SPINAL CORD INJURY
OPEN FRACTURES
PENETRATING ABDOMINAL INJURY
PENETRATING CHEST INJURY

Referring to FIGS. 21A-B, the acalculous cholecystitis decision support algorithm of the present invention is illustrated. If an intensivist suspects that acalculous cholecystitis may be present, the intensivist may not be certain of all of the aspects that would be indicative of this particular condition. Therefore, the intensivist is lead through a decision support algorithm, which first causes the intensivist to determine if the patient is clinically infected, either febrile or leukocytosis **800**. If this criterion is not met, the intensivist is prompted that it is unlikely that the patient has acalculous cholecystitis **802**.

If the patient is clinically infected **800**, the intensivist is prompted to determine whether the patient has had a previous cholecystectomy **804**. If patient has had a previous cholecystectomy, the intensivist is prompted that it is very unlikely that the patient has acalculous cholecystitis **806**. Alternatively, if a patient has not had a previous cholecystectomy, the intensivist is prompted to determine whether the patient has any of seven (7) risk factors, specifically: 1) Prolonged intensive care unit (ICU) stay (defined as greater than six (6) days); 2) recent surgery (particularly aortic cross clamp procedures); 3) hypotension; 4) positive end-expiratory pressure (PEEP) greater than ten (10) centimeters (cm); 5) transfusion greater than six (6) units of blood; 6) inability to use the gastrointestinal (GI) tract for nutrition; or 7) immunosuppression (AIDS, transplantation, or leukemia) **808**. If the patient has none of these seven risk factors, the intensivist is prompted that the patient probably does not have acalculous cholecystitis **810**.

If the patient has any of the seven risk factors **808**, the intensivist is prompted to determine whether the patient has any of the following symptoms: right upper quadrant (RUQ) tenderness; elevated alkalinephosphatase; elevated bilirubin; or elevated liver transaminases **812**. If the patient has none of these four (4) symptoms **812**, the intensivist is prompted to consider other more likely sources of infection (see fever of unknown origin or FUO) **814**. If the infection remains undiagnosed following an alternative work-up, the intensivist is prompted to re-enter the algorithm **814**.

If the patient has any of these four (4) symptoms **812**, the intensivist is prompted to determine whether alternative intra-abdominal infectious sources are more likely **816**. If alternative intra-abdominal infectious sources are not more likely, the intensivist is prompted to determine whether the patient is sufficiently stable to go for a test **826**. If the patient is sufficiently stable to go for a test, the intensivist is prompted to perform an mso4 Cholescintigraphy **836**. The normal AC is excluded **838**. If the test indicates an abnormality, the intensivist is prompted to consider a cholecystectomy or precutaneous drainage **840**. If the patient is not sufficiently stable to go for a test, the intensivist is prompted to perform a bedside ultrasound **828**. If no other infectious etiologies are identified and no abnormalities of the gall-bladder are noted but: a) the patient remains ill **830**, the intensivist is prompted to consider empiric cholecystostomy **832**. If no other infectious etiologies are identified and no abnormalities of the gall bladder are noted but: b) the patient is improving **830**, the intensivist is prompted to continue to observe the patient **834**.

If alternative intra-abdominal infectious sources are more likely **816**, the intensivist is prompted to determine whether the patient is sufficiently stable to go for a test **818**. If the patient is sufficiently stable to go for a test **818**, the intensivist is prompted to perform an abdominal CT scan **820**. If no other infectious etiologies are apparent and the test: a) demonstrates abnormalities of the gall-bladder but not diagnostic; or b) no gall-bladder abnormalities are noted **822**, the intensivist is prompted to maintain continued observation of the patient **824**. Alternatively, if neither of these criteria is met **822**, the intensivist is prompted to perform an mso4 cholescintigraphy **836**. Normal AC is excluded **838**. If the test is abnormal, the intensivist is prompted to consider cholecystectomy or precutaneous drainage **840**. If the patient is not sufficiently stable to go for a test, the intensivist is prompted to perform a bedside ultrasound **828**. If no other infectious etiologies are identified and no abnormalities of the gall-bladder are noted but: a) the patient remains ill **830**, the intensivist is prompted to consider empiric cholecystostomy **832**. If no other infectious etiologies are identified and no abnormalities of the gall bladder are noted but: b) the patient is improving **830**, the intensivist is prompted to continue to observe the patient **834**.

Referring to FIG. 22, the adrenal insufficiency decision support algorithm of the present invention is illustrated. When an intensivist suspects an adrenal problem may be presented in a patient, the intensivist may initiate the adrenal insufficiency decision support algorithm which prompts questions concerning all aspects of the condition. First the intensivist is prompted to determine whether the patient is either hypotensive and/or has been administered pressors for forty-eight hours or longer **900**. If neither condition is met, the system advises the intensivist that it is unlikely that an adrenal problem is present **902**.

If one or both conditions are met, the intensivist is asked whether an obvious cause for hypotensive blood pressure or treatment with pressors are manifested, such as hypovolemia or low blood volume, myocardial dysfunction, or spinal injury **904**. If at least one of these obvious causes is present, the intensivist is alerted by the system that the underlying cause must first be treated **906**. If treatment of a suspected underlying cause is reversed, yet the hypotension or pressor need persists, the intensivist is further directed to determine whether other adrenal problems have occurred in the patient's history **908**, **910**, **912**.

In order to examine prior treatment issues, the intensivist is first prompted by the system to determine if the patient has been treated with steroids in the previous six months for at least a two week period **908**. Next, the intensivist is

prompted to determine whether the patient has hyponatremia or hyperkalemia **910**. The intensivist is also prompted to determine whether the patient has experienced anticoagulation or become coagulopathic prior to the hypotension or pressor treatment **912**. According to the responses provided by the intensivist to the system queries or blocks **908**, **910**, and **912**, the system calculates a treatment action **914** as follows: The array of possible responses to diagnosis questions **908**, **910**, and **912** are given a Decision Code as shown in Table 1A: Adrenal Insufficiency Considerations, below.

TABLE 1A

Adrenal Insufficiency Considerations			
Question 1 908	Question 2 910	Question 3 912	Decision Code
N	N	N	A
N	N	Y	A
N	Y	N	B
N	Y	Y	C
Y	Y	Y	C
Y	N	N	D
Y	Y	N	B
Y	N	Y	D
Y	Y	Y	C

The possible decision codes of Table 1A are as follows:

Decision Code	Treatment Action
A	Do cosyntropin stim test
B	Consider possible Adrenal Insufficiency. Give decadron 5 mg IV, so cosyntropin stim test and empirically treat with hydrocortisone 50 mg IV every 8 hours until stim test results return.
C	Consider possible Adrenal Insufficiency, secondary to adrenal hemorrhage. Give decadron 5 mg IV, so cosyntropin stim test and empirically treat with hydrocortisone 50 mg IV every 8 hours until stim test results return.
D	Do cosyntropin stim test, may empirically treat with hydrocortisone 25-50 mg IV every 8 hours until stim test results return

Besides specialized treatment actions listed in the decision codes above, the intensivist is directed to administer a cosyntropin stimulation test **914** in order to see how much cortisone the adrenal gland is producing.

After performing the cosyntropin stimulation test, the intensivist is prompted to enter the patient's level of cortisol before administering cosyntropin and thirty minutes afterwards **916**. The software analyzes the test results as follows: The results in Table 2, shown below, are shown as having certain decision codes A through F.

TABLE 2

Cosyntropin Stimulation Test Results		
basal (A)	basal (B)	basal (C)
<15	15-20	>25
stim (D)	stim (E)	stim (F)
<5	5-10	>10

Depending upon the outcome of the analysis of Table 2, one of the treatment actions, shown below in Table 3, will be displayed **918**.

TABLE 3

Cosyntropin Test Result Treatment Actions	
Decision Code	Treatment Action
A + D	Adrenal insufficiency diagnosed - treat with hydrocortisone 50 mg IV every 8 hours and consider endocrine consult
A + E	Probable Adrenal insufficiency- treat with hydrocortisone
B + D	25-50 mg IV every 8 hours and taper as intercurrent illness improves
A + F	Possible Adrenal insufficiency- consider treatment with
B + E	hydrocortisone 25 mg IV every 8 hours and taper as intercurrent illness improves
A + F	Adrenal insufficiency unlikely- would not treat
B + F	
C + E	
C + F	

Referring to FIG. 23, the blunt cardiac injury decision support algorithm of the present invention is illustrated. If an intensivist suspects that blunt cardiac injury may be present, the intensivist may not be certain of all aspects that would be critical to or indicative of this particular condition. Therefore, the intensivist is lead through a decision support algorithm, which first causes the intensivist to determine whether any of seven (7) risk factors are present: 1) was thoracic impact greater than fifteen (15) mph; 2) was the steering wheel deformed; 3) was there precordial ecchymosis, contusions, or abrasions; 4) was marked precordial tenderness present; 5) was there a fractured sternum; 6) were bilateral rib/costal cartilage fractures present; 7) were thoracic spine fractures present **1000**. If none of the 7 risk factors are present, the intensivist is prompted that no further evaluation is necessary **1002**. If any of the 7 risk factors are present, the intensivist is prompted to obtain an electrocardiogram (ECG) and chest X-ray (CXR) **1004**.

Once the results of the ECG and CXR are obtained, the intensivist is prompted to determine: whether the ECG results are abnormal, with abnormal being defined as anything other than sinus rhythm, including ectopy and unexplained sinus tachycardia (greater than 100 beats/minute); and whether the CXR results are abnormal, with abnormal being defined as any skeletal or pulmonary injury, especially cardiac enlargement **1006**. If either the ECG or CXR is not abnormal, the intensivist is prompted that a monitored bed is unnecessary for the patient **1008**. If either the ECG or CXR is abnormal, the intensivist is prompted to determine whether there is any hemodynamic instability (hemodynamic instability being defined as the absence of hypovolemia, spinal cord injury, or sepsis) that cannot be explained by hypovolemia, spinal cord injury, or sepsis **1010**.

If this criterion is not met, the intensivist is prompted: that the patient should be in a monitored bed; that the ECG should be repeated at 24 hours; that, at any time, if unexplained hemodynamic instability is present, the intensivist should request a stat echo; and that, if blunt thoracic aortic injury is also suspected, a transesophageal echocardiogram (TEE) is favored over a transthoracic echocardiogram (TTE) **1012**. Once the results of these tests are obtained, the intensivist is prompted further to determine whether ectopy, arrhythmia, or abnormality is present on the ECG **1014**. If none of these criteria are met, the intensivist is prompted that cardiac injury is excluded **1016**. If any of these criteria are met, the intensivist is prompted that he should consider monitoring the patient for an additional 24 hours **1018**.

If the internist determines that there is any hemodynamic instability that cannot be explained by hypovolemia, spinal cord injury, or sepsis **1010**, he is prompted: to perform a stat echo; and, if blunt thoracic aortic injury is also suspected, that a transesophageal echocardiogram (TEE) is favored over a transthoracic echocardiogram (TTE) **1020**. Once the results of the stat echo are obtained, the intensivist is prompted to determine whether the echo is abnormal with possible causes for the abnormality being: pericardial effusion (tamponade; hypokineses or akinesis (wall motion); dilatation or reduced systolic function; acute valvular dysfunction; and/or chamber rupture **1022**. If the stat echo is abnormal, the intensivist is prompted to treat as indicated for the particular cause of the abnormality **1026**. If the stat echo is not abnormal, the intensivist is prompted to continue to monitor the patient and repeat the ECG at 24 hours **1024**.

Once the results of the ECG are obtained, the intensivist is prompted to determine whether ectopy, arrhythmia, or abnormality are present on the ECG **1014**. If any of these criteria are not met, the intensivist is prompted that cardiac injury is excluded **1016**. If any of these criteria are met, the intensivist is prompted that he should consider monitoring the patient for an additional 24 hours **1018**.

Referring to FIGS. 24A-B, the candiduria decision support algorithm, which is yet another decision support algorithm of the present invention is illustrated. In the candiduria decision support algorithm, the intensivist is presented with the criteria for diagnosing candiduria, or severe fungal infection. First, the intensivist determines whether the patient has any medical conditions that render the patient prone to fungal infections, such as diabetes, GU anatomic abnormality, renal transplant, or pyuria **1100**. If there are no such conditions, the intensivist is next prompted by the system to look for dissemination or spreading of the fungal infection **1102**. If the infection does not seem to have spread, the intensivist is prompted to change the patient's catheter and test for pyuria after twenty four hours have passed **1104**.

The intensivist is prompted by the system to determine whether the patient can have P.O. **1106**. If the patient can take P.O., the system next prompts the intensivist to determine whether azoles, an organic compound for inhibiting fungal growth, have been administered in the past three days to fight the infection **1108**. If azoles have been previously administered, the systemic infection diagnosis is confirmed and the intensivist is referred to the systemic amphotericin dosing algorithm **1110**. If azoles have not been previously administered, directions for the proper treatment dosage of fluconazole (a type of azole) is provided to the intensivist along with adjustments for the species of fungus found **1112**. Where the patient cannot take P.O., the intensivist is again referred to the systemic amphotericin dosing algorithm **1114**.

When the patient does have some condition prone to fungal infection, the intensivist is prompted to determine what other signs of dissemination are exhibited in the patient **1116**. The intensivist is prompted to see if the patient can take P.O. If the patient cannot take P.O., the intensivist is referred to the systemic amphotericin dosing algorithm **1120**. If the patient can take P.O., the intensivist is prompted to check whether azoles have been administered in the previous three days **1122**. If azoles have been administered, the systemic infection is confirmed and the intensivist is referred to the systemic amphotericin dosing algorithm **1124**. If no azoles have been administered previously, the intensivist is given instructions for administering fluconazole to treat the fungal infection **1126**.

If there is no evidence of dissemination, the intensivist is still prompted to determine whether the patient can take P.O. **1128**. Where the patient cannot take P.O., directions are provided to administer amphotericin bladder washing procedures **1130**. If the patient cannot take P.O., the intensivist is prompted to determine whether azoles have been administered in the previous three days **1132**. If azoles have been administered, the systemic infection is confirmed and the intensivist is referred to the systemic amphotericin dosing algorithm **1134**. If no azoles have been administered previously, the intensivist is given instructions for administering fluconazole to treat the fungal infection **1136**.

Referring to FIGS. 25A-B, the Cervical Spine Injury decision support algorithm of the present invention is illustrated. If an intensivist suspects that a cervical spine injury may be present, the intensivist may not be certain of all of the factors that would be indicative of this particular condition. Therefore, the intensivist is lead through a decision support algorithm, which first prompts the intensivist to determine if the patient is awake, alert, not intoxicated, and has no mental status changes **1200**. If these criteria are met, the intensivist is prompted to determine whether the patient has any neck pain **1202**. If the patient does not have any neck pain, the intensivist is prompted to determine whether the patient has any other pain which would distract from his or her neck pain **1204**. If this criterion is not met, the intensivist is prompted to determine whether the patient has any neurologic deficits **1206**. If this criterion is not met, the intensivist is prompted that a stable C-spine is present if the patient can flex, extend, move neck left/right without pain and without neck tenderness to palpitation **1208**. The intensivist is prompted further that he can remove the collar **1208**.

Alternatively, if the patient does have neck pain **1202**, the intensivist is prompted to order 3 x rays **1210** consisting of: 1) lateral view revealing the base of the occiput to the upper border of the first thoracic vertebra; 2) anteroposterior view revealing spinous processes of the second cervical through the first thoracic vertebra; and 3) an open mouth odontoid view revealing the lateral masses of the first cervical vertebra and entire odontoid process **1210**. If the x rays are normal the intensivist is prompted to consider extension then flexion lateral x rays; if normal he is prompted that he can remove the collar; if abnormal, he is prompted to obtain a surgical consult **1212**. If the x rays are abnormal, the intensivist is prompted to obtain a surgical consult and order a CT scan **1214**. If the x rays are indeterminate, the intensivist is prompted to order a CT scan **1216**.

Alternatively, if the patient has no other pain which would distract from their neck pain **1204**, the intensivist is prompted to order 3 x rays (the same types of x rays described in **1210** above with the same prompting based on normal, abnormal, or indeterminate x rays) **1218**.

If the patient does have neurologic deficits **1206**, the intensivist is prompted to determine whether the neurologic deficit is referable to the cervical spine **1226**. If this criterion is not met, the intensivist is prompted to order 3 x rays (the same types of x rays described in **1210** above with the same prompting based on normal, abnormal, or indeterminate x rays) **1218**. If the neurologic deficit is referable to the cervical spine **1226**, the intensivist is prompted that the patient should obtain immediate spine trauma surgery consult and CT or MRI (if available) **1228**.

Alternatively, if the intensivist determines that the patient does not pass the criteria of being awake, alert, not intoxicated and having no mental status changes **1200**, the intensivist is prompted to determine whether the patient has severe head trauma **1232**. If this criterion is met, the inten-

sivist is prompted to order CT of the neck with head CT **1236**. If this criterion is not met, the intensivist is prompted to determine whether the patient has any neurologic deficit referable to the cervical spine **1234**. If the intensivist determines that the patient does have a neurologic deficit referable to the cervical spine, the intensivist is prompted that the patient should obtain immediate spine trauma surgery consult and CT or MRI (if available) **1228**. If the intensivist determines that the patient does not have a neurologic deficit referable to the cervical spine **1234**, he is prompted to order 3 x rays (the same types of x rays described in **1210** above with the same prompting based on normal, abnormal, or indeterminate x rays) **1218**.

Referring to FIG. **26A-B**, the Oliguria decision support algorithm of the present invention is illustrated. If an intensivist suspects that Oliguria may be present, the intensivist may not be certain of all of the aspects that would be indicative of this particular condition. Therefore, the intensivist is lead through a decision support algorithm, which first causes the intensivist to determine if the patient is oliguric, with the criterion being passage of less than 25 cc of urine in a period of 2 hours **1300**. If this criterion is met the intensivist is prompted to determine whether the patient is anuric (the criterion for which is passage of less than 10 cc of urine in a 2 hour period) in spite of fluid administration **1302**.

If this criterion is met, the intensivist is prompted to determine whether the urinary catheter is working by flushing the catheter **1304**. The intensivist is then prompted to determine whether the catheter is functioning **1306**. If the catheter is not functioning, the intensivist is prompted to replace or reposition the catheter **1308**. If the catheter is functioning, the intensivist is prompted to determine whether the patient has a history of: 1) renal stone disease; 2) abdominal, pelvic, or retroperitoneal cancer; or 3) recent pelvic or retroperitoneal surgery **1310**. If any of these criteria are met, the intensivist is prompted to perform the following actions: 1) do renal ultrasound emergently to rule out obstruction; 2) while waiting for ultrasound, administer fluid at the rate of 7-15 ml/kg of bodyweight; and 3) send urine for specific gravity determination **1312**. Based on the renal ultrasound test results, the intensivist is prompted to determine whether an obstruction is present **1314**. If an obstruction is determined to be present, the intensivist is prompted to consult a urologist immediately **1316**.

Alternatively, if the intensivist determines that the patient does not have a history of: 1) renal stone disease; 2) abdominal, pelvic, or retroperitoneal cancer; or 3) recent pelvic or retroperitoneal surgery **1310**, the intensivist is prompted to determine whether: 1) the patient has a history of heart failure or known ejection fraction of less than 30 percent; or 2) there are rales on the physical exam **1318**.

Alternatively, if following the renal ultrasound test, the intensivist determines that there is no obstruction the intensivist is prompted to determine whether: 1) the patient has a history of heart failure or known ejection fraction of less than 30 percent; or 2) there are rales on the physical exam **1318**.

If the intensivist determines that the patient is not anuric **1302**, then the intensivist is prompted to determine whether: 1) the patient has a history of heart failure or known ejection fraction of less than 30 percent; or 2) whether there are rales on the physical examination **1318**. If neither of these criteria is met, the intensivist is prompted to administer fluids to the patient at the rate of 10-20 ml/kg of bodyweight **1320** and send the patient's urine sample for a specific gravity test **1322** as more fully described in FIGS. **26B-C**.

Alternatively, if the patient does: 1) have a history of heart failure or known ejection fraction less than 30 percent; or 2) there are rales on the physical exam **1318**, the intensivist is prompted to determine whether there has been a chest x-ray (CXR) in the last 6 hours **1324**. If this criterion is not met, the intensivist is prompted to determine whether there has been a change in respiratory status **1326**. If there has been no change in the respiratory status, the intensivist is prompted to administer 7-15 ml of fluids per kg of bodyweight **1328** and to send the patient's urine sample for a specific gravity test.

Alternatively, if the intensivist determines that there has been a change in respiratory status **1326**, the intensivist is prompted to: 1) do a chest x-ray; and 2) determine whether there is evidence of edema or congestion **1334**. If there is evidence of edema or congestion **1334**, the intensivist is prompted to: 1) insert a PA catheter to measure wedge pressure and liver function to direct fluid replacement; and 2) send urine creatinine and sodium **1332**.

If the intensivist determines that there has been a CXR in the last 6 hours **1324**, the intensivist is prompted to determine whether there is evidence of edema or congestion **1330**. If there is no evidence of edema or congestion, the intensivist is prompted to administer 7-15 ml of fluids per kg of bodyweight **1328** and send the patient's urine for a specific gravity test **1322**.

Alternatively, if the intensivist determines there is evidence of edema or congestion **1330**, the intensivist is prompted to: 1) insert a PA catheter to measure wedge pressure and liver function to direct fluid replacement; and 2) send urine creatinine and sodium **1332**.

Referring now to FIG. **26C-D**, the oliguria algorithm description continues. Following the specific gravity test of the patient's urine, the intensivist is prompted to determine whether the results indicate the specific gravity is less than 1.018. If this criterion is met, the intensivist is prompted to: 1) send blood and urine immediately to test for blood urea nitrogen (BUN), creatinine, electrolytes, and Hgb, and spot urine for creatinine, sodium, and sediment; and 2) administer 5-10 ml of fluid per kg of bodyweight **1356**. Once the results of these tests are obtained, the intensivist is prompted to determine what is the Hgb **1338**.

If the Hgb has increased by more than 1.5 gm/dl compared to the previous Hgb **1340**, the intensivist is prompted to: 1) administer fluids 5-10 ml/kg of bodyweight and follow the urine output closely **1342**. Following this, the intensivist is prompted to determine whether the labs confirm renal failure by use of the formula  $FE.sub.Na=Urine\ Na.times.Serum\ Creatinine/Urine\ Creatinine.times.Serum\ Na.times.100$  **1344**.

If the Hgb is within 1.5 gm/dl from the previous Hgb or no comparison **1352**, the intensivist is prompted to determine what is the mean blood pressure **1354**. If the mean blood pressure is determined to be within 20 percent or higher than the baseline blood pressure **1356**, the intensivist is prompted to determine whether the labs confirm renal failure **1344**. If the mean blood pressure is determined to be greater than 20 percent below the baseline pressure **1358**, the intensivist is prompted to give additional fluids and consider invasive hemodynamic monitoring **1360**. Following this, the intensivist is prompted to determine whether the labs confirm renal failure by use of the formula  $FE.sub.Na=Urine\ Na.times.Serum\ Creatinine/Urine\ Creatinine.times.Serum\ Na.times.100$  **1344**.

Alternatively if the Hgb has decreased by 1.5 gm/dl compared to the previous Hgb **1362**, the intensivist is prompted to: 1) transfuse PRBCs as needed; 2) look for

source of bleeding and check PT, aPTT, & platelet count **1364**. Following this, the intensivist is prompted to determine what is the mean blood pressure **1354**. If the mean blood pressure is determined to be greater than 20 percent below the baseline pressure **1358**, the intensivist is prompted to give additional fluids and consider invasive hemodynamic monitoring **1360**. Following this, the intensivist is prompted to determine whether the labs confirm renal failure by use of the formula  $FE.sub.Na=Urine\ Na.times.Serum\ Creatinine/Urine\ Creatinine.times.Serum\ Na.times.\ 100$  **1344**.

If the labs do not confirm renal failure, as indicated by  $FE.sub.Na.ltoreq.1$  percent **1346**, the intensivist is prompted to: 1) continue to administer fluids and follow urine output; and 2) recheck creatinine in 6-12 hours **1348**.

Alternatively, if the labs do confirm renal failure, as indicated by  $FE.sub.Na.>1$  percent **1350**, the intensivist is prompted to: 1) place central venous pressure (CVP); 2) Assure adequate intravascular volume; 3) give trial of diuretics: 40 mg lasix IV, if no response in 1 hour, give hydrodiuril 500 mg IV, wait 20-30 minutes then give 100 mg lasix, if persistent oliguria, restrict: 1) fluids; 2) potassium & phosphate; if diuresis ensues, restrict only potassium & phosphate; in both situations, adjust all renally excreted medications; and 4) see acute renal failure **1350**.

Referring now to FIG. 26E, the oliguria algorithm description continues. Alternatively, following the specific gravity test of the patient's urine, the intensivist is prompted to determine whether the results indicate the specific gravity is greater than or equal to 1.018 **1336**. If this criterion is not met **1364**, the intensivist is prompted to determine whether the urine is dark or tea colored **1366**. If this criterion is met, the intensivist is prompted to: 1) check creatinine phosphokinase; and 2) force fluids to induce diuresis **1368**.

If the intensivist determines that the urine is not dark or tea colored, the intensivist is prompted to: 1) administer 10-20 ml of fluids per kg of bodyweight; and 2) check Hgb **1370**. The intensivist is then prompted to determine what is the Hgb **1372**.

If the Hgb is determined to be greater than 1.5 gm/dl higher than the previous Hgb **1374**, the intensivist is directed to: 1) force fluids; and 2) continue to follow the urine output **1376**.

Alternatively, if the Hgb is determined to be within 1.5 gm/dl of the last Hgb or there is no Hgb for comparison **1378**, the intensivist is prompted to determine what is the mean blood pressure **1380**. If the mean blood pressure is determined to be 20 percent or higher than the baseline pressure **1382**, the intensivist is prompted to: 1) continue to administer fluids; 2) follow urine output; and 3) check creatinine in 6-12 hours **1384**. If the mean blood pressure is determined to be greater than 20 percent below the baseline pressure **1386**, the intensivist is prompted to: 1) continue to push fluids; 2) consider invasive hemodynamic monitoring; and 3) if post-op abdominal trauma, consider abdominal compartment syndrome **1388**.

If the Hgb is determined to be greater than 1.5 gm/dl below the previous Hgb **1390**, the intensivist is prompted to: 1) transfuse blood as needed; 2) look for bleeding source; 3) check PT, aPPT & platelet count; 4) continue to push fluids; and 5) recheck Hgb in 1-2 hours **1392**.

Referring to FIG. 27A-B, the open fractures decision support algorithm of the present invention is illustrated. Open fractures are where bone, cartilage, or a tooth break and push through the skin surface. The intensivist is first prompted by the system to determine whether the patient has an open fracture **1500**. If one has occurred, the intensivist must then determine whether the wound is contaminated

with soil, or was inflicted in a barnyard **1502** in order to address higher risk of infection. If the wound is contaminated with soil, or was inflicted in a barnyard, the intensivist is prompted to administer a high dose of penicillin to the antibiotics prescribed **1504**. The intensivist is also prompted to take several treatment steps **1506**. These treatment steps include administering tetanus prophylaxis, such as antitoxin injection, monitoring staphylococcus aureus until twenty-four hours after surgery, caring for the wound within six hours, and where the injury is found to be more severe during surgery, the intensivist is prompted to administer aminoglycosides for seventy two hours.

If the wound is not contaminated with soil, or was inflicted in a barnyard, the intensivist is next prompted to determine the severity of the wound **1508**. To do so, the intensivist must determine the length of the wound and corresponding soft tissue damage. If the wound is either less than one centimeter and clean or greater than a centimeter long without extensive soft tissue damage, the intensivist is prompted to take several treatment steps **1506** as previously described. Where the soft tissue damage is extensive or amputation has occurred, the intensivist is prompted by the system to make further determinations **1510**, **1512**, **1514** about the wound caused by the fracture. The intensivist is prompted to determine if enough soft tissue coverage is remaining for the wound to close and heal **1510**, if any arterial repair is needed **1512**, and if extensive soft tissue damage with periosteal injury, and bone exposure **1514**. If there is adequate soft tissue coverage, the intensivist is advised that risk of infection is low and directed to take treatment actions **1516**. If arterial damage requiring repair is present, the intensivist is advised by the system that risk of infection is moderate to high and given treatment instructions **1518**. Where there is soft tissue injury with periosteal stripping and bone exposure, the intensivist is alerted by the system that risk of infection is high and given treatment instructions **1520**. The treatment instructions in each case **1516**, **1518**, **1520** include administering tetanus prophylaxis, such as antitoxin injection, caring for the wound within six hours, and performing: monitoring for staphylococcus aureus, and administering aminoglycosides and high doses of penicillin, all for seventy two hours before and after any operative procedures.

If the intensivist has determined that no exposed fracture has occurred, the system next prompts the intensivist to determine whether there is any evidence of neuro-vascular damage **1522**. If there is evidence of neuro-vascular damage, the intensivist is prompted to consult with a neurosurgeon or vascular surgeon immediately **1524**. If the intensivist determines there is no evidence of neuro-vascular damage to the patient, the system next prompts the intensivist to determine whether the patient has compartment syndrome **1526**. If there is evidence of compartment syndrome seen in the patient, the intensivist is prompted to consult orthopedics right away **1528**. If there is no evidence of compartment syndrome seen in the patient, the intensivist is still prompted to consult orthopedics, but without any prompt for time sensitivity **1530**.

Referring to FIGS. 28A-B, the Pancreatitis diagnostic algorithm of the present invention is illustrated. To evaluate whether a patient has pancreatitis, the intensivist is first prompted to examine whether severe epigastric abdominal pains and amylase levels three times greater than normal are present in the patient **1600**. If neither or one of the conditions is present, the intensivist is prompted to consider other

causes of the abdominal pain, such as mesenteric ischemia, a perforated ulcer, intestinal obstruction, biliary colic, or an ectopic pregnancy **1602**.

If severe epigastric abdominal pains and amylase levels three times greater than normal are present, the intensivist is next prompted to provide the Ranson Criteria which is a criteria associated with the severity of pancreatitis and the potential outcome or prognosis at that particular level of severity, or Apache II score which is also a score associated with the severity of the disease and the potential prognosis at a particular level of the patient **1604**. If the patient has a Ranson Criteria less than three or an Apache II score of less than eight, the intensivist is prompted by the system to consider removing the patient from the Intensive Care Unit **1606**. However, if the patient has a Ranson Criteria greater than three or an Apache II score of greater than eight, the intensivist is instructed to perform an abdominal ultrasound test within twenty-four hours **1607**. If the results of the ultrasound test show a biliary obstruction, the intensivist is instructed to consider performing an ERCP to find and remove any gallstones **1608**.

If the abdominal ultrasound results do not show any biliary obstruction, intensivist is next prompted to perform more diagnostic tests **1610**. The intensivist is directed to perform a Dynamic IV contrast and an abdominal Tomography (CT) scan. If the intensivist does not suspect a surgical condition exists, such as a perforated ulcer, mesenteric infarction or pancreatic infection, the tests may be performed after three days have passed. If the intensivist does suspect a surgical condition exists, the tests should be performed within three days. In either case, if the patient has creatinine levels greater than or equal to 2 milligrams per dl, the intensivist should not perform the Dynamic IV contrast test.

Once the CT scan is performed, the intensivist is prompted to determine whether necrotizing pancreatitis is present **1612**. The intensivist is next required to determine whether the patient has improved since admission **1614**. If no improvement has been seen, the intensivist is directed to perform percutaneous fluid aspiration and do a gram stain culture the collected fluid **1616**. If the culture shows infection **1618**, the intensivist is directed to perform surgical debridement of the pancreas **1620**. If the results of the culture are sterile **1622**, the intensivist is directed to closely follow up on the patient's condition **1624** and watch for clinical deterioration **1626**. If the patient does further deteriorate, the intensivist is then instructed to perform a surgical debridement of the pancreas **1628**. If the patient does not deteriorate, the intensivist is still prompted to closely follow the patient's condition **1630**.

Where the CT scan does not show signs of necrotizing pancreatitis **1612**, the intensivist is prompted by the system to closely observe the patient **1632**. The intensivist is also prompted to check whether clinical deterioration is occurring **1634**. If no deterioration is observed, the intensivist continues to observe the patient's condition **1636**. If clinical deterioration is occurring **1634**, the intensivist is directed to perform percutaneous fluid aspiration and do a gram stain culture the collected fluid **1616**. If the culture shows infection **1618**, the intensivist is directed to order surgical debridement of the pancreas **1620**. If the results of the culture are sterile **1622**, the intensivist is directed to closely follow up on the patient's condition **1624** and watch for clinical deterioration **1626**. If the patient does further deteriorate, the intensivist is then prompted to order a surgical debridement of the pancreas **1628**. If the patient does not deteriorate, the intensivist is still directed by the system to closely follow the patient's condition **1630**.

Referring to FIGS. 29A-B, the penicillin allergy diagnosis algorithm of the present invention is illustrated. In order to

diagnose a penicillin allergy, the intensivist is first prompted to determine whether the patient has a history suggestive of previous penicillin or cephalosporin anaphylaxis **1700**. Various known reactions, including angioedema, flushing, pruritis, airway obstruction, syncope, and hypertension, are displayed for the intensivist's review. If the patient has previously had any of these reactions, the intensivist is prompted to determine whether the patient has ever taken synthetic or partially synthetic antibiotics, such as ampicillin, amoxicillin, duricef or kefzol, without any anaphylaxis symptoms **1702**. If the patient has taken synthetics without reaction, the intensivist is advised by the system that penicillin or cephalosporin may be administered **1716**. If the patient has reacted to synthetic or partially synthetic antibiotics, the intensivist is next prompted to determine whether the patient needs penicillin or cephalosporin specifically **1704**.

If the patient is not required to have penicillin or cephalosporin, the intensivist is prompted to administer the synthetic antibiotics **1706**. If the patient does need penicillin or cephalosporin, the intensivist is directed by the system to consider consulting with an allergist or immunologist and perform skin tests for reactions **1708**. Next, the intensivist is prompted to enter whether the skin test was positive **1710**. If the results are negative, the intensivist is further directed by the system to administer penicillin or cephalosporin with caution, to consider pretreatment with benadryl or prednisone to counter any reaction, and to closely monitor the patient **1712**. If the results of the skin test are positive, the intensivist is prompted by the system to perform desensitization procedures **1714**.

If the patient does not have a history suggestive of previous penicillin or cephalosporin anaphylaxis **1700**, the intensivist is prompted to determine whether the patient has previously experienced skin-level reactions, such as exfoliative dermatitis, Stevens Johnson Syndrome, or Toxic Epidermal Necrolysis, when given penicillin or cephalosporin **1718**. If the patient has previously experienced one of these reactions, the intensivist is directed by the system to administer an alternative antibiotic **1720**. If the patient has not experienced one of these reactions, the intensivist is prompted to determine whether there is a history of any rash when given penicillin or cephalosporin **1722**. If the patient has not previously had a rash when given penicillin or cephalosporin, the intensivist is advised that the patient will most likely be able to take penicillin or cephalosporin **1724**.

If the patient has previously experienced a rash when given penicillin or cephalosporin, the intensivist is prompted to determine whether the rash presented when the patient was given ampicillin or amoxycillin **1726**. If the rash resulted from ampicillin or amoxycillin, the intensivist is next prompted to determine whether the rash was urticarial **1728**. If the rash was not urticarial, the intensivist is advised by the system that the patient probably can take penicillin or cephalosporin, but should be closely monitored **1730**. If the rash was urticarial, the intensivist is prompted to determine whether or not the patient needs penicillin or cephalosporin **1704**.

If the patient is not required to have penicillin or cephalosporin, the intensivist is directed by the system to administer the synthetic antibiotics **1706**. If the patient does need penicillin or cephalosporin, the intensivist is directed to consider consulting with an allergist or immunologist and perform skin tests for reactions **1708**. Next, the intensivist is prompted to enter whether the skin test was positive **1710**. If the results are negative, the intensivist is further directed to administer penicillin or cephalosporin with caution, to consider pretreatment with benadryl or prednisone to counter any reaction, and to closely monitor the patient

1712. If the results of the skin test are positive, the intensivist is directed to perform desensitization procedures 1714.

Referring to FIG. 30A-B, the Post-Op Hypertension decision support algorithm of the present invention is illustrated. If an intensivist determines that there may be a possibility of post-op hypertension, the intensivist may not be certain of all aspects that would be involved in this particular condition. Therefore, the intensivist is lead through a decision support algorithm which prompts the intensivist to determine the appropriate care to be given.

Initially, the intensivist is prompted to determine whether the patient is hypertensive (BP greater than 20 percent above mean baseline) 1800. If this criterion is met, the intensivist is prompted to determine whether the patient has any of the causes of reversible hypertension: 1) hypercapnia; 2) bladder distension; 3) pain; 4) increased ICP; 5) drugs (pressors, cocaine, ketamine and chronic MAO use with indirect acting vasopressors); 6) automatic hyperreflexia; or 7) volume overload 1802. If any of these criteria are met, the intensivist is prompted to first treat those specific etiologies and, if pressure remains high, re-enter algorithm 1804.

Alternatively, if none of these criteria are met 1802, the intensivist is prompted to determine whether the patient is at risk of injury from post-op hypertension (i.e., vascular surgery, coronary artery disease, neurosurgery, ocular surgery, etc.) 1806. If this criterion is not met 1806, the intensivist is prompted to determine whether the BP is greater than 40 percent above mean baseline 1808. If this criterion is not met, the intensivist is prompted that the patient may not need BP treatment 1810.

If the BP is greater than 40 percent above the mean baseline 1808, the intensivist is prompted to determine whether the patient is in pain 1812. If this criterion is met 1812, the intensivist is prompted to treat pain and continue 1814. Following this prompt 1814, the intensivist is prompted next to determine whether the patient is actively bleeding or at significant risk for post-op bleeding (i.e., "moist closure" or high drain output) 1816. If either of these criteria is met 1816, the intensivist is prompted to use only short acting agents including emolol and nitroprusside as needed until bleeding has abated 1818.

Alternatively, if neither of these criteria is met 1816, the intensivist is prompted to determine whether the patient is tachycardic (absolute greater than 90 bpm or (relative greater than 15 percent over baseline)) 1820. If either of these criteria is met 1820, the intensivist is prompted to go to Decision Table C, which is programmed for the condition of a high heart rate. If neither of these criteria is met 1820, the intensivist is prompted to eliminate (NOT C) Table C and proceed to the next decision point 1820.

TABLE C

		HR ↑					
	CAD	Y	Y	Y	N	N	N
	RAD	N	Y	Y	N	Y	N
	↓EF	N	N	Y	N	Y	Y
Treatment	1 <sup>ST</sup>	L	E	L	L	A	E
	2 <sup>ND</sup>	E	L	A	N	N	A

The intensivist is prompted next to determine whether the patient is bradycardic (absolute less than 60 bpm) 1822. If this criterion is met, the intensivist is prompted to go to Decision Table B, which is programmed for the condition of a low heart rate.

TABLE B

		HR ↓					
	CAD	Y	Y	Y	N	N	N
	RAD	N	Y	Y	N	Y	N
	↓EF	N	N	Y	N	Y	Y
Treatment	1 <sup>ST</sup>	N	N	A	N	A	A
	2 <sup>ND</sup>	S	S	S	H	H	H

If this criterion is not met, the intensivist is prompted to eliminate (NOT B) Table B and proceed to the next decision point 1822. [Note: If NOT C and NOT B, the intensivist is prompted to go to Table A by default, i.e., If NOT C and NOT B Then A].

TABLE A

		HR (nl)					
	CAD	Y	Y	Y	N	N	N
	RAD	N	Y	Y	N	Y	N
	↓EF	N	N	Y	N	Y	Y
Treatment	1 <sup>ST</sup>	L	E	A	N	A	A
	2 <sup>ND</sup>	N	N	E	A	N	N

The intensivist is prompted next to determine, sequentially, table input values for CAD, RAD, and EF.

In these decision tables, the letter references have the following meanings: L=labelalol, E=esmolol, A=enalapril, N=nicardipine, H=hyrdalazine, S=nitroprusside. The reference to 1.sup.st and 2.sup.nd means that treatment should begin with the 1.sup.st drug and add or substitute the 2.sup.nd drug as needed.

Using the above decision tables, the intensivist is prompted to determine whether the patient has known coronary artery disease (CAD) or 3 or more risk factors for CAD 1824. If either of these criteria is met 1824, the intensivist is prompted to enter a "Y" or "YES" for CAD into the table selected above in 1820 and 1822. If neither of these criteria is met, the intensivist is prompted to enter a "N" or "NO" for CAD into the table selected above in 1820 and 1822.

Next, the intensivist is prompted to determine whether the patient has known reactive airway disease (RAD) 1826. If this criterion is met 1826, the intensivist is prompted to enter a "Y" or "YES" for RAD into the table selected above in 1820 and 1822. If this criterion is not met, the intensivist is prompted to enter a "N" or "NO" for RAD into the table selected above in 1820 and 1822.

Next, the intensivist is prompted to determine whether the patient has known EF less than 30 percent or a history of systolic heart failure 1828. If either of these criteria is met 1828, the intensivist is prompted to enter a "Y" or "YES" for EF into the table selected above in 1820 and 1822. If neither of these criteria is met 1828, the intensivist is prompted to enter a "N" or "NO" for EF into the table selected above in 1820 and 1822.

Based on the table selected in 1820 and 1822 above, and the table inputs determined from 1824, 1826, and 1828, the intensivist is prompted with the proper medication to administer for the 1.sup.st and 2.sup.nd treatment.

If the patient is not in pain 1812, the intensivist is prompted to employ the procedures described above in 1816.

If the patient is at risk of injury from post-op hypertension 1806, the intensivist is prompted to determine whether the blood pressure is greater than 40 percent above baseline

1830. If this criterion is met 1830, the intensivist is prompted to employ the procedures described above in 1812.

Alternatively, if this criterion is not met 1830, the intensivist is prompted to determine whether the patient is in pain 1836. If this criterion is met 1836, the intensivist is prompted to treat pain and reevaluate following analgesia and, if still hypertensive, to continue algorithm 1838. Following this action 1838, the intensivist is prompted to employ the procedures described above in 1816. If the patient is not in pain 1836, the intensivist is prompted to employ the procedures described above in 1816.

If the patient is determined not to be hypertensive 1800, the intensivist is prompted to determine whether the patient requires their BP controlled near baseline (i.e., neurosurgery, carotid surgery, thoracic aorta surgery) 1832. If this criterion is not met 1832, the intensivist is prompted that the patient probably does not need treatment 1834.

Alternatively, if this criterion is met 1832, the intensivist is prompted to employ the procedures described above in 1836.

Referring to FIG. 31A, the pulmonary embolism diagnosis algorithm is illustrated. If a pulmonary embolism is suspected, the intensivist is first prompted to determine whether the patient is hemodynamically unstable 2900. If the patient is hemodynamically unstable, the intensivist is directed by the system to consider performing an immediate transthoracic echocardiogram, pulmonary angiogram and treatment consistent with massive pulmonary embolism 2902. If the patient is not hemodynamically unstable, the intensivist is prompted to perform a VQ scan and perform further assessment of the patient 2904.

In order to further assess the patient, the intensivist is prompted to respond to a series of questions 2906, 2908, 2910, 2912. The intensivist is prompted to determine whether any of the following patient conditions are present: Dyspnea, Worsening chronic dyspnea, Pleuritic chest pain, Chest pain that is non-retro sternal & non-pleuritic, O.sub.2 saturation<92% on room air that corrects with 40% O.sub.2 supplementation, Hemoptysis, or Pleural rub 2906. The intensivist is also prompted to determine whether any risk factors are in the patient's history, such as: Surgery within 12 weeks, Immobilization (complete bed rest) for >3 days within 4 weeks, Previous DVT or objectively diagnosed PE, Lower extremity fracture & immobilization within 12 weeks, Strong family history of DVT or PE(.gtoreq.2 family members with objective proven events or 1.sup.st degree relative with hereditary thrombophilia), Cancer (treatment within the last 6 months or palliative stages), Postpartum, or Lower extremity paralysis 2908. Further, the intensivist must determine whether the patient has any of the following symptoms: Heart rate>90 beats/min, Temp.gtoreq.38.0, CXR free of abnormalities (edema, pneumonia, pneumothorax), or Leg symptoms c/w DVT, syncope, blood pressure less than 90 mm Hg with heart rate greater than 100 beats/min, receiving mechanical ventilation and/or oxygen supplementation greater than 40%, and new onset or right heart failure (-JVP, new S1, Q3, T3, or RBBB) 2910. The intensivist is also queried by the system to consider alternative diagnosis that may be more likely than pulmonary embolism. To do so, the intensivist is prompted to consider conditions that simulate major pulmonary embolism, such as myocardial infarction, acute infection with COPD, septic Shock, dissecting aortic aneurysm, or occult hemorrhage. The intensivist is additionally prompted to consider conditions that simulate minor pulmonary embolism, such as acute bronchitis, pericarditis, viral pleurisy, pneumonia, and esophageal spasm 2912.

Referring to FIG. 31B, the pulmonary embolism algorithm description continues. The intensivist enters the answers to the assessment queries posed 2906, 2908, 2910, 2912 into the system. If two or more responses to the patient condition query 2906 were answered yes and one or more questions were answered yes from: Heart rate>90 beats/min, Temp.gtoreq.38.0, CXR free of abnormalities, or Leg symptoms c/w DVT of the symptoms query 2910, the intensivist is informed that a typical pulmonary embolism is present 2914. Next, the system compares this response to the answer to the alternative diagnosis query 2912. If an alternative diagnosis is at least as likely as pulmonary embolism 2916, the intensivist is also given a low probability 2918 to moderate probability 2920 risk factor. If an alternative diagnosis is less likely than pulmonary embolism 2922, the intensivist is given a moderate 2924 to high 2926 probability risk factor.

If less than two yes answers resulted from the patient conditions 2906, the intensivist is advised by the system that an atypical pulmonary embolism may be present 2928. Next, the system compares this response to the answer to the alternative diagnosis query 2912. If an alternative diagnosis is at least as likely as pulmonary embolism 2930, the intensivist is told there is no risk and low probability 2932 or some risk with a low probability 2934 risk factor. If an alternative diagnosis is less likely than pulmonary embolism 2934, the intensivist is given a no risk and low probability 2938 to risk but moderate probability 2940.

If at least one answer to the symptoms of syncope, blood pressure less than 90 mm Hg with heart rate greater than 100 beats/min, receiving mechanical ventilation and/or oxygen supplementation greater than 40%, and new onset or right heart failure 2910 is yes, the intensivist is prompted with a message that severe pulmonary embolism is occurring 2942. Next, the system compares this response to the answer to the alternative diagnosis query 2912. If an alternative diagnosis is at least as likely as pulmonary embolism 2944, the intensivist is told there is a moderate probability of pulmonary embolism 2946. If an alternative diagnosis is less likely than pulmonary embolism 2948, the intensivist is notified that a high probability of pulmonary embolism is present 2950.

Once the risk factors and probabilities are determined the system compares this information to the VQ scan results. This comparison is performed according to the following Table 4 below.

TABLE 4

Input	Clinical Probability		
	High	Moderate	Low
V/Q Scan			
High	A	A	B
Intermediate	B	C	C
Low	B	C	E
Normal	E	E	E

Where the VQ scan column and the risk column intersect, a letter code is assigned to various treatment instructions. The treatment instructions are as follows.

- A=Pulmonary embolus diagnosed. Begin treatment
- E=Pulmonary embolus excluded

B=Proceed with the following work-up:

- 1) Perform spira CT (If patient has renal insufficiency [creatinine > 2.0], consider going directly to pulmonary angiogram to reduce the potential dye load). If positive begin treatment,
- 2) If negative, assess for DVT using compression ultrasound or venography. If positive begin treatment,
- 3) If negative, perform pulmonary angiogram. If positive begin treatment, if negative diagnosis excluded.

C=Proceed with the following work-up:

- 1) Perform spiral CT. If positive begin treatment,
- 2) If negative, assess for DVT using compression ultrasound or venography. If positive begin treatment,
- 3) If negative perform D-dimer assay (elisa only). If negative diagnosis excluded, If positive, perform serial ultrasound of the lower extremities.

Once the correlation is made, the instructions associated with the letter code are displayed by the system to prompt the intensivist with diagnosis and treatment instructions.

Referring to FIG. 32, the seizure decision support algorithm of the present invention is illustrated. If an intensivist encounters seizure in a patient, he may not be certain of all of the aspects and the timelines that are critical to treating this particular condition. Therefore, the intensivist is lead through a decision support algorithm, which divides the treatment sequence into three segments: 0-30 minutes; 30-60 minutes; and beyond 60 minutes.

At the onset of a seizure, in the 0-30 minute segment of the algorithm, the intensivist is prompted to give the patient lorazepam (0.1 mg/kg of bodyweight) in 2 mg boluses up to 8 mg **2000**. Subsequently, the intensivist is prompted to give the patient phenytoin (18-20 mg/kg of bodyweight) at 50 mg/min of fosphenytoin (18-20 mg/kg of bodyweight) at 150 mg/min followed by 5 mg/kg of bodyweight/day through separate IV line **2002**.

During the 30-60 minute segment of the algorithm, the intensivist is prompted to: reload additional phenytoin or fosphenytoin (10 mg/kg of bodyweight) maintaining previous infusion; and give additional lorazepam (0.05 mg/kg of bodyweight) **2004**. Subsequently, the intensivist is prompted to begin continuous EEG monitoring **2006**.

The intensivist is then prompted to determine whether the patient is hemodynamically stable **2008**. If hemodynamically stable, the intensivist is prompted to administer propofol 1-2 mg/kg of bodyweight bolus followed by 2-10 mg/kg/hr **2010**.

At the 60 minute segment of the algorithm, the intensivist is prompted that if seizure activity stops, he should taper either midazolam or propofol over the next 12-24 hours while maintaining phenytoin but if seizures persist, he is prompted to move to the pentobarbital coma block **2012**.

Under pentobarbital coma, the intensivist is prompted to administer 10-15 mg/kg/hr and to maintain until seizure control is achieved on EEG **2014**. The intensivist is prompted further that the patient usually requires PA catheter and pressors to maintain hemodynamic control **2014**.

Alternatively, if the patient is determined to be hemodynamically unstable **2016**, the intensivist is prompted to utilize fluids and pressors as needed (phynylephrine or dopamine) midazolam 0.2 mg/kg bolus followed by 0.1-2.0 mg/kg/hr **2018**.

At the 60 minute segment of the algorithm, the intensivist is prompted that if seizure activity stops, he should taper either midazolam or propofol over the next 12-24 hours while maintain phenytoin but if seizures persist, he is prompted to move to the pentobarbital coma block **2012**.

Under pentobarbital coma, the intensivist is prompted to administer 10-15 mg/kg/hr and to maintain until seizure control is achieved on EEG **2014**. The intensivist is prompted further that the patient usually requires PA catheter and pressors to maintain hemodynamic control **2014**.

Referring to FIGS. 33A-B, the supra ventricular tachycardia (SVT) decision support algorithm of the present invention is illustrated. If an intensivist determines that SVT is present, the intensivist may not be certain of all aspects that would be involved in treating this particular condition. Therefore, the intensivist is lead through a decision support algorithm which prompts the intensivist to determine the appropriate care to be given.

Initially, the intensivist is prompted to determine whether SVT is stable or unstable **2100**. If SVT is stable **2102**, the intensivist is prompted to determine whether the patient has a regular or irregular rhythm **2102**. If the patient has a regular rhythm **2104**, the intensivist is prompted to determine whether there is a wide complex or a narrow complex **2104**. If the intensivist determines that there is a wide complex **2106**, the intensivist is prompted to administer adenosine 6 mg/12 mg (if needed) **2108**. Following the administering of adenosine **2108**, the intensivist is prompted to consider that if the patient converts to sinus rhythm (SR) to consider re-entrant junctional or WPW re-entrant. If the wide complex recurs, treat the patient with esmolol or Ca+2 blockers.

Alternatively; if no effect, the intensivist is prompted to consider V-tach **2112**. Next, the intensivist is prompted to: 1) load procainamide 150 mg over 10 min, then 1 mg/min infusion; and 2) synchronized cardiovert **2114**.

Alternatively, if the wide complex slows, the intensivist is prompted to consider SVT w/aberrancy and continue to slow with esmolol or Ca+2 blockers **2116**.

The intensivist is prompted next to administer esmolol/calcium blockers and link to ventricular rate control **2118**. The intensivist is prompted next to determine whether there has been a conversion to SR **2120**. If there is no conversion to SR in 24 hours, the intensivist is prompted to add antiarrhythmic agent and consider anticoagulation **2122**. The intensivist is prompted next to determine whether there has been conversion to SR. If conversion to SR, the intensivist is prompted to continue maintenance antiarrhythmic agent during hospitalization **2124**. If no conversion to SR, the intensivist is prompted to cardiovert while on antiarrhythmic & following heparinization **2126**.

If the patient has a regular rhythm **2104**, the intensivist is prompted to determine whether there is a wide complex or a narrow complex **2104**. If the intensivist determines that there is a narrow complex **2128**, the intensivist is prompted to administer adenosine 6 mg/12 mg (if needed) **2130**. If administering the adenosine **2130** slows the ventricular rate only and the atrial rate persists, the intensivist is prompted to consider atrial flutter and continue to slow with esmolol or Ca+2 blockers **2132**. The intensivist is prompted next to employ the procedures described above in **2118**.

If administering the adenosine **2130** converts the patient to SR, the intensivist is prompted to consider re-entrant sinus or junctional and if recurs, treat with esmolol or Ca+2 blockers **2134**.

If administering the adenosine **2130** slows both atrial and ventricular rates the intensivist is prompted that there is a probable sinus tachycardia **2136**. The intensivist is prompted next to continue to slow with esmolol **2138**. The intensivist is prompted next to employ the procedures described above in **2118**.

If SVT is stable **2102**, the intensivist is also prompted to determine whether the patient has a regular or irregular rhythm **2102**. If the patient has an irregular rhythm **2140**, the intensivist is prompted that if no p waves, there is probable Atrial fibrillation **2142**. The intensivist is prompted next to

slow ventricular response with esmolol or Ca+2 blockers **2144**. The intensivist is prompted next to employ the procedures described above in **2118**.

If the patient has an irregular rhythm **2140**, the intensivist is prompted to determine whether there are more than 3 p wave types MAT—and to treat underlying lung dz. and avoid theophylline compounds **2146**. The intensivist is prompted next to slow rate with Ca+2 blockers only **2148**. The intensivist is prompted next to employ the procedures described above in **2118**.

Referring now to FIG. **33C**, the description of the SVT decision algorithm continues. If SVT is unstable **2101**, the intensivist is prompted to determine whether the patient has SBP less than 80, ischemia, mental status changes **2150**. The intensivist is prompted next to perform synchronous cardioversion (100 J, 200 J, 300 J) **2152**. The intensivist is prompted next that if sinus rhythm: 1) correct reversible etiologies; 2) consider starting IV antiarrhythmic for maintenance of sinus rhythm **2154**. Alternatively, following **2152**, the intensivist is prompted next that if continued SVT: 1) correct reversible etiologies; 2) load IV antiarrhythmic (see dosing guidelines) and repeat DC cardioversion **2156**.

For example, and without limitations, wide complex QRS Tachycardia is also addressed in the decision support algorithm of the present invention. Referring to FIGS. **34A-B**, the wide complex QRS tachycardia decision support algorithm is illustrated. If an intensivist determines that there may be a possibility of wide complex QRS tachycardia, the intensivist may not be certain of all aspects that would be involved in this particular condition. Therefore, the intensivist is lead through a decision support algorithm which prompts the intensivist to determine the appropriate care to be given.

Initially, the intensivist is prompted to determine whether the patient is hemodynamically stable (no angina, heart failure, or hypotension (systolic less than 80 mm)) **2200**. If this criterion is not met, the intensivist is prompted to go to the cardio-pulmonary guidelines algorithm which is generally known to those skilled in the art.

Alternatively, if this criterion is met, the intensivist is prompted to determine whether the patient is within 7 days of a myocardial infarction or at risk for myocardial ischemia **2202**. If the patient is not within 7 days of a myocardial infarction or at risk for myocardial ischemia **2202**, the intensivist is prompted to determine whether the wide complex QRS rhythm is sustained (greater than 30 seconds) **2234**. If this criterion is not met, the intensivist is prompted to determined whether the QRS is monomorphic **2236**. If the QRS is monomorphic **2236**, the to intensivist is prompted to determine whether the patient has structural heart disease **2242**. If the patient has structural heart disease **2242**, the intensivist is prompted to: 1) monitor closely; 2) look for reversible etiologies; and 3) consider antiarrhythmic therapy **2244**. If the patient does not have structural heart disease **2242**, the intensivist is prompted to: 1) monitor closely; 2) look for reversible etiologies; and 3) if recurs and symptomatic may require further testing (prolonged holter or EP study) **2246**.

If the QRS is not monomorphic **2236**, the intensivist is prompted to determine whether the QT is prolonged **2238**. If this criterion is met, the intensivist is prompted to: 1) check K; 2) give Mg; and 3) consider overdrive pacing **2240**. If the

intensivist determines that the QT is not prolonged, **2238**, the intensivist is prompted to employ the procedures described above in **2242**.

If the wide complex QRS rhythm is sustained **2234**, the intensivist is prompted to determine whether the rhythm is polymorphic or irregular **2208**. If the rhythm is polymorphic or irregular, the intensivist is prompted to consider atrial fibrillation with accessory pathway conduction and load with procainamide and get a cardiology consultation **2210**. If the rhythm is not polymorphic or irregular, the intensivist is prompted with the question of whether he wishes to: 1) perform ECG diagnosis; or 2) administer adenosine diagnostically **2220**. If the intensivist makes the determination to perform an ECG diagnosis **2220**, he is prompted to go to the ECG diagnosis algorithm **2300**.

If the intensivist makes the determination to administer adenosine diagnostically **2220**, he is prompted to go to the administer adenosine branch of the algorithm **2222**. If there is no effect, the intensivist is prompted that there is probable VT and to determine whether the VT is monomorphic **2224**. If the VT is monomorphic **2224**, the intensivist is prompted to load with procainamide and perform synchronous cardioversion **2226**.

Alternatively, if the VT is not monomorphic **2224**, the intensivist is prompted to load with lidocaine and perform immediate cardioversion **2228**.

If the ventricular response is slowed after administering adenosine **2222**, the intensivist is prompted to consider SVT with aberrancy and treat with esmolol or Ca blockers **2230**.

If the ventricular response converts to sinus rhythm after administering adenosine **2222**, the intensivist is prompted: to consider re-entrant mechanism with BBB or WPW; and, 1) if WPW consult cardiology for possible ablation **2232**.

If the patient is within 7 days of a myocardial infarction or at risk for myocardial ischemia **2202**, the intensivist is prompted to determine whether the wide complex is sustained (30 seconds) **2204**. If the wide complex is not sustained **2204**, the intensivist is prompted to determine whether the patient: 1) symptomatic; 2) tachycardia runs are frequent; or 3) the tachycardia rates are rapid (greater than 180) **2212**. If none of these criteria is met, the intensivist is prompted to observe **2216**. Alternatively, if any of these criteria is met **2212**, the intensivist is prompted to: 1) administer lidocaine 100-200 mg & 1-4 mg/min infusion; and 2) amiodarone **2214**.

If the wide complex is sustained **2204**, the intensivist is prompted to determine whether the rate is greater than 140/min **2206**. If this criterion is not met **2206**, the intensivist is prompted: to consider accelerated idioventricular, and that in some patients this can lead to hemodynamic compromise; and that 1) he can perform overdrive pacing if needed **2218**.

Alternatively, if this criterion is met, the intensivist is prompted to follow the procedures in **2208**.

If the intensivist makes the determination to perform ECG Diagnosis **2220**, he is prompted to go to the ECG Diagnosis branch of the algorithm **2220**. Referring now to FIG. **34C**, in the ECG Diagnosis branch, the intensivist is prompted to determine whether the patient has known pre-excitation syndrome **2300**. If this criterion is met, the intensivist is prompted to determine whether the QRS complexes are predominantly negative in leads V4-V6 **2302**. If the QRS complexes are predominantly negative in leads V4-V6, the intensivist is prompted that there is probable VT **2304**.

If the QRS complexes are not predominantly negative in leads V4-V6 **2302**, the intensivist is prompted to determine whether there is a QR complex in one or more of leads

V2-V6**2306**. If this criterion is met, the intensivist is prompted that there is probable VT **2308**.

Alternatively, if this criterion is not met **2306**, the intensivist is prompted to determine whether there are more QRS complexes than P waves **2310**. If there are more QRS complexes than P waves **2310**, the intensivist is prompted that there is probable VT **2312**. If there are not more QRS complexes than P waves **2310**, the intensivist is prompted: to consider pre-excited SVT; and that he may wish to perform EP study **2314**.

If the intensivist determines that the patient does not have known pre-excitation syndrome **2300**, the intensivist is prompted to determine whether there is an RS complex present in any precordial lead **2316**. If this criterion is not met **2316**, the intensivist is prompted that there is probable VT **2318**.

Alternatively, if this criterion is met **2316**, the intensivist is prompted to determine whether the R to S interval is greater than 100 MS in any one precordial lead **2320**. If this criterion is met, the intensivist is prompted that there is probable VT **2322**.

If the R to S interval is not greater than 100 MS in any one precordial lead **2320**, the intensivist is prompted to determine whether there is evidence of atrioventricular dissociation **2324**. If this criterion is met, the intensivist is prompted that there is probable VT **2326**.

Alternatively, if there is no evidence of atrioventricular dissociation **2324**, the intensivist is prompted to determine whether V-1 is negative and V-6 positive and QRS greater than 0.14 mSEC **2328**. If these criteria are met, the intensivist is prompted that there is probable VT **2330**.

If none of these criteria is met **2328**, the intensivist is prompted that the situation may represent SVT with aberrancy or underlying bundle branch block **2332**.

Referring to FIG. 35A, the assessment of sedation algorithm of the present invention is illustrated. If an intensivist encounters a need for sedation, he may not be certain of all of the aspects and the timelines that are critical to this particular process. Therefore, the intensivist is lead through a decision support algorithm, which prompts the intensivist to address a number of factors in the process **3100**.

The intensivist is prompted initially to go to the Scoring section of the algorithm **3100**. The intensivist is prompted to proceed through a number of scorings **3102** and to first score the patient's alertness with points being allocated in the following manner: asleep/unresponsive=0; responsive to voice=1; and hyperresponsive=2 **3104**.

The intensivist is prompted next to score the patient's movement with points being allocated in the following manner: no spontaneous movement=0; spontaneous movement=1; and pulls at lines, tubes, dressings=2 **3106**.

The intensivist is prompted next to score the patient's respiration based on whether the patient is mechanically ventilated or spontaneously breathing with points being allocated as subsequently discussed. If the patient is mechanically ventilated, the intensivist is prompted to allocate points in the following manner: no spontaneous ventilation=0; spontaneous ventilations and synchronous with ventilator=1; or spontaneous ventilations with cough or dysynchrony>10 percent of breaths=2 **3108**. Alternatively, if the patient is spontaneously breathing, the intensivist is prompted to allocate points in the following manner: respiration rate (RR)<10=0; RR=10-30=1; or RR>30=2 **3108**.

The intensivist is prompted next to score the patient's heart rate with points being allocated in the following

manner: >20 percent below mean for last 4 hr=0; within 20 percent mean for last 4 hr=1; or >20 percent above mean for last 4 hr=2 **3110**.

The intensivist is prompted next to score the patient's blood pressure with points being allocated in the following manner: MAP>20 percent for last 4 hr=0; MAP within 20 percent mean for last 4 hr=1; or MAP>20 percent above mean for last 4 hr=2 **3112**.

The intensivist is prompted next to determine the sedation score by the following formula: SEDATION SCORE=alertness+movement+respirations+heart rate+ blood pressure **3114**. In one embodiment, respiratory rate, heart rate, and BP can be computer linked to monitor data thereby simplifying the sedation scoring assessment. The nursing observations are deemed intuitive and the nursing burden in sedation scoring can be minimal by using this point scoring.

Referring now to FIG. 35B, the sedation assessment algorithm description continues. The intensivist is prompted then to continue the sedation assessment by moving to the Pain Assessment section of the algorithm **3116**.

In the Pain Assessment section, the intensivist is prompted to determine whether the patient is conscious, communicative, and acknowledging pain **3118**. If any of these criteria is not met, the intensivist is prompted to determine: whether the sedation score is greater than 2 and the patient: is known to be in pain before becoming uncommunicative; or S/p recent surgery; or having tissue ischemia or infarct; or has wounds; or has large tumor possibly impinging on nerves. If the answer to either of these two questions is YES, the intensivist is prompted to treat for pain **3118**. The intensivist is prompted then to continue the assessment by moving to the Delirium Assessment section of the algorithm **3118**.

In the Delirium Assessment section, the intensivist is prompted to determine whether the sedation score is greater than 2 AND the patient has: day/night reversal with increased agitation at night OR eyes open and "awake" but disoriented; or eyes open and "awake" but pulling at lines, tubes, or dressings OR difficult to sedate prior to ventilator weaning OR paradoxical response to benzodiazepines. If these criteria is met, the intensivist is prompted to consider butyropheneone **3120**.

Referring to FIG. 36, the Bolus sliding scale algorithm is illustrated. If an intensivist encounters a need for sedation, the algorithm for which may contain a reference to the bolus sliding scale for midazolam, he may not be certain of all of the aspects which are critical to this scale. Therefore, the intensivist is lead through a decision support algorithm, which prompts the intensivist through the use of the scale **3200**.

If lorazepam is less than 0-2 mg IV q 6 hr, then the intensivist is prompted to give midazolam 1-2 mg q 5 min until adequately sedated **3202**.

Alternatively, if lorazepam equals 2-4 mg IV q 4 hr, then the intensivist is prompted to give midazolam 2 mg q 5 min until adequately sedated **3202**.

Alternatively, if lorazepam is greater than 10 mg IV q 4 hr, then the intensivist is prompted to give midazolam 5 mg q 5 min until adequately AND consider fentanyl and/or droperidol or Haldol for synergy despite delirium and pain assessment **3202**.

Yet another decision support routine is the sedation algorithm. Referring to FIG. 37, the sedation process decision support algorithm is illustrated. If an intensivist determines that a patient will require sedation, the intensivist may not be certain of all aspects that would be involved in this particular

process. Therefore, the intensivist is lead through a decision support algorithm, which prompts the intensivist to conduct a sedation assessment based on: 1) scoring; 2) pain; and 3) delirium (see Assessment of Sedation algorithm) **3300**.

Following completion of the sedation assessment process **3300**, the intensivist is prompted to determine whether the patient is in pain **3302**. If this criterion is met, the intensivist is prompted to administer bolus morphine, fentanyl, other narcotic, start patient controlled analgesic (PCA) or epidural analgesia as indicated **3324**. If the patient is not in pain **3302** or after administering bolus morphine, fentanyl, other narcotic, start patient controlled analgesic (PCA) or epidural analgesia as indicated **3324**, the intensivist is prompted to determine whether the patient is delirious **3304**.

If the intensivist determines that the patient is delirious **3304**, he is prompted to administer droperidol 2.5-5 mg q 30 min pm and that he may consider IV Haldol not to exceed 30 mg/24 hr **3326**. If the patient is not delirious or after following the procedures in **3326**, the intensivist is prompted to determine whether the patient will need sedation for more than the next 24 hours **3306**. If the patient will not need sedation for more than the next 24 hours **3306**, the process continues as described in FIG. **38**.

Alternatively, if the patient will need sedation for more than the next 24 hours **3306**, the intensivist is prompted to determine whether the sedation score is 8-10 **3308**. If this criterion is met, the intensivist is prompted to employ the Bolus sliding scale midazolam and increase lorazepam by 20 percent **3328** (see Bolus sliding scale midazolam algorithm—FIG. **36**). Subsequently, the intensivist is prompted to reassess sedation in 4 hr **3330**.

Alternatively, if the patient will need sedation for more than the next 24 hours **3306**, the intensivist is prompted to determine whether the sedation score is 8-10 **3308**. If this criterion is met, the intensivist is prompted to employ the Bolus sliding scale midazolam and increase lorazepam by 20 percent **3328** (see Bolus sliding scale midazolam algorithm—FIG. **42**). Subsequently, the intensivist is prompted to reassess sedation in 4 hr **3330**.

If the sedation score is not 8-10, the intensivist is prompted to determine whether the sedation score is greater than or equal to the last Sed Scr after sedative bolus or increase **3310**. If this criterion is met, the intensivist is prompted to employ the procedures described above in **3328** and **3330**.

If the sedation score is not greater than or equal to the last Sed Scr after sedative bolus or increase **3310**, the intensivist is prompted to determine whether four (4) or more midaz boluses have been given since last q 4 hr assessment **3312**. If this criterion is met, the intensivist is prompted to employ the procedures described above in **3328** and **3330**.

Alternatively, if less than four (4) midaz boluses have been given since last q 4 hr assessment **3312**, the intensivist is prompted to determine whether the patient is adequately sedated **3314**. If this criterion is not met, the intensivist is prompted to employ the procedure described in **3328** and **3330**.

If the intensivist determines that the patient is adequately sedated **3314**, the intensivist is prompted to determine whether the sedation score is 0-2 **3316**. If this criterion is met, the intensivist is prompted to decrease lorazepam by 20 percent **3332** and reassess sedation in 4 hr **3334**.

Alternatively, if the sedation score is not 0-2 **3316**, the intensivist is prompted to determine whether the sedation score is less than or equal to the last Sed Scr after sedative

decrease **3318**. If this criterion is met, the intensivist is prompted to employ the procedure described in **3332** and **3334**.

If the sedation score is not less than or equal to the last Sed Scr after sedative increase **3318**, the intensivist is prompted to determine whether the patient is clinically oversedated **3320**. If the patient is clinically oversedated **3320**, the intensivist is prompted to employ the procedure described in **3332** and **3334**. If the patient is not clinically oversedated **3320**, the intensivist is prompted to reassess sedation in 4 hr **3322**.

Referring to FIG. **38**, the short term sedation process decision support algorithm of the present invention is illustrated. If an intensivist determines that a patient will not require sedation past the next 24 hour period, the intensivist may not be certain of all aspects that would be involved in this particular process. Therefore, the intensivist is lead through a decision support algorithm, which prompts the intensivist to conduct a sedation assessment based on: 1) scoring; 2) pain; and 3) delirium (see Assessment of Sedation algorithm) **3100**.

Following completion of the sedation assessment process **3100**, the intensivist is prompted to decrease lorazepam by 20 percent from baseline per day **3102**. The intensivist is prompted next to determine whether the patient is in pain **3104**. If this criterion is met, the intensivist is prompted to administer bolus morphine or fentanyl **3122**. If the patient is not in pain or after administering bolus morphine or fentanyl **3122**, the intensivist is prompted to determine whether the patient is delirious **3106**.

If the intensivist determines that the patient is delirious, he is prompted to administer droperidol 2.5-5 mg q30 min pm **3124**. If the patient is not delirious or after administering droperidol **3124**, the intensivist is prompted to determine whether the sedation score is 8-10 **3108**.

If this criterion is met, the intensivist is prompted to employ the Bolus sliding scale midazolam (see Bolus sliding scale midazolam algorithm) and begin midazolam infusion or begin propofol 1-2 mg/kg bolus and 5-50 mcg/kg/min infusion **3126**. Subsequently, the intensivist is prompted to reassess sedation in 1 hr **3128**.

If the sedation score is not 8-10, the intensivist is prompted to determine whether the sedation score is greater than or equal to the last Sed Scr after sedative bolus or increase **3110**. If this criterion is met, the intensivist is prompted to employ the procedures described above in **3126** and **3128**.

If the intensivist determines that the sedation score is not greater than the last sedation score after sedative bolus or increase **3110**, the intensivist is prompted to determine whether the patient is adequately sedated **3112**. If this criterion is not met, the intensivist is prompted to employ the procedures described above in **3126** and **3128**.

If the intensivist determines that the patient is adequately sedated **3112**, he is prompted to determine whether the sedation score is 0-2 **3114**. If this criterion is met, the intensivist is prompted to determine if the patient has been sedated for more than 72 hours **3130**. If the patient has not been sedated for more than 72 hours **3130**, the intensivist is prompted to hold midazolam or propofol and hold or decrease lorazepam by 50 percent **3132**. The intensivist is prompted subsequently to reassess sedation in 1 hour **3134**.

Alternatively, if the intensivist determines that the patient has been sedated for more than 72 hours **3130**, the intensivist is prompted to hold midazolam or propofol and decrease lorazepam by 20 percent per day **3136**. The intensivist is prompted subsequently to reassess sedation in 1 hour **3134**.

Alternatively, if the intensivist determines that the sedation score is not 0-2 **3114**, the intensivist is prompted to determine whether the sedation score is less than or equal to the last sedation screening after sedative decrease **3116**. If this criterion is met, the intensivist is prompted to determine whether the patient has been sedated for more than 72 hours and to follow the procedures described above in **3130**.

If the intensivist determines that the sedation score is not less than or equal to the last Sed Scr after sedative decrease **3116**, the intensivist is prompted to determine whether the patient is clinically oversedated **3118**. If this criterion is met, the intensivist is prompted to determine whether the patient has been sedated for more than 72 hours and to follow the procedures described above in **3130**. If this criterion is not met, the intensivist is prompted to reassess sedation in 1 hr **3120**.

Referring to FIG. **39**, the respiratory isolation decision support algorithm is illustrated. If an intensivist determines that there may be a need for respiratory isolation, the intensivist may not be certain of all aspects that would be involved in this process. Therefore, the intensivist is lead through a decision support algorithm which prompts the intensivist to determine the need for respiratory isolation based upon: a) clinical assessment; and/or b) smear/culture findings **3500**.

Pursuing the clinical assessment branch of the decision support algorithm, the intensivist is prompted to determine whether the patient has known mTB (mycobacterium tuberculosis) **3502**. If this criterion is met, the intensivist is prompted to determine whether the patient has been compliant with their medications for over 2 weeks and is clinically responding **3512**. If the patient has not been compliant with their medications for over 2 weeks and is not clinically responding **3512**, the intensivist is prompted that isolation is required **3514**. If the patient has been compliant with their medications and is clinically responding **3512**, the intensivist is prompted that no isolation is required **3516**.

Alternatively, if the patient does not have known mTB **3502**, the intensivist is prompted to determine whether the patient has known mycobacterial disease other than TB **3504**. If this criterion is met, the intensivist is prompted to determine whether the patient has new CXR (chest x ray) findings and symptoms (cough 2 weeks, fever, weight loss) **3518**. If the patient does not have new CXR findings and symptoms **3518**, the intensivist is prompted that no isolation is required **3520**. If the patient does have new CXR findings and symptoms **3518**, the intensivist is prompted that isolation is required **3522**.

If the patient does not have known mycobacterial disease other than TB **3504**, the intensivist is prompted to determine whether there is a new cavitory lesion on CXR **3506**. If this criterion is met, the intensivist is prompted that isolation is required **3524**.

Alternatively, if there is no new cavitory lesion on CXR **3506**, the intensivist is prompted to determine whether there are pulmonary infiltrates or whether the patient is HIV (human immunodeficiency virus) positive **3508**. If neither of these criteria is met, the intensivist is prompted that no isolation is required **3510**. If either of these criteria is met, the intensivist is prompted to determine whether the patient has new CXR findings and symptoms (cough 2 weeks, fever, weight loss) and at high risk: 1) known mTB exposure; 2) homeless; 3) prisoner; 4) travel to area with multi-drug resistant TB **3526**. If these criteria are met, the intensivist is prompted that isolation is required **3528**. Alternatively, if these criteria are not met, the intensivist is prompted that no isolation is required **3530**.

Pursuing the smear/culture branch of the decision support algorithm **3500**, the intensivist is prompted to determine whether the AFB (acid-fast bacilli) smear is positive **3532**. If the AFB smear is not positive, the intensivist is prompted that: no isolation is required; await culture results; if culture negative, no isolation required; if culture positive and patient has mycobacterial disease other than TB (MOTT no isolation is required; if the culture is positive and the patient does not have MOTT consult ID **3534**.

Alternatively, if the AFB smear is positive, the intensivist is prompted to determine whether the patient has known mycobacterial disease other than TB **3536**. If this criterion is not met, the intensivist is prompted that isolation is required **3538**. If this criterion is met, the intensivist is prompted: to isolate until results of NAP test are in; if mTB positive isolate the patient; if no mTB, no isolation is required **3540**.

Referring to FIG. **40**, the empiric meningitis treatment decision support algorithm of the present invention is illustrated. If the intensivist is treating a patient for meningitis, the intensivist is prompted to answer a series of queries by the system to properly address medication and dosage. First, the intensivist is prompted to determine whether the patient has suffered a head trauma or undergone neurosurgery **3700**. The answer to this question is input 1 to table x below. The intensivist is next prompted to determine whether the patient is allergic to penicillin or is from an area where penicillin resistant staphylococcus pneumoniae is prevalent **3702**. The answer to this question becomes input 2 to table x below. The intensivist must also determine whether the patient is immunocompromised **3704**, and the answer becomes input 3 to table x below. The intensivist determines if the patient is over fifty years of age **3706**, with the answer being input 4 in table x below. Lastly, the intensivist is prompted to determine whether the patient has altered mental status **3708**, and the answer becomes input 5 in table x below. The inputs to each of these prompts **3702**, **3704**, **3706**, **3708** is compared to a dosage database according to the Table 5 below.

TABLE 5

Meningitis Input-Output Table

Input	Combinations	Output
1	1 = yes 2 = no	A) vancomycin 1.5-2 gm IV q 12 h + ceftazidime 2 gm IV q 8 hr or cefapime 2 gm IV q 8 hr
2	1 = yes 2 = no	B) vancomycin 1.5-2 gm IV q 12 h + aztreonam 0.5-2 gm IV q 6-8 hr
3	1 = no 2 = no 3 = no 4 = yes	ampicillin 2 gm IV q 4 h + ceftriaxone 2 gm IV q12 cefotaxime 2 gm IV q 6 h
4	1 = no 2 = no 3 = no 4 = no	ceftriaxone 2 gm IV q 12 hr or cefotaxime 2 gm IV q 6 hr
5	1 = no 2 = no 3 = yes	ampicillin 2 gm IV q 4 hr + ceftazidime 2 gm IV q 8 hr or cefipime 2 gm IV q 8 hr
6	1 = no 2 = yes 3 = no 4 = yes	vancomycin 1.5-2 gm IV q 12 hr + chloramphenicol 1 gm IV q 6 hr
7	1 = no 2 = yes 3 = no 4 = no	

TABLE 5-continued

Meningitis Input-Output Table		
Input	Combinations	Output
8	1 = no 2 = yes 3 = yes	
9	5 = yes to inputs 3-8	add to output consider acyclovir 10 mg/kg IV q 8 h

In the Meningitis Input-Output Table, possible combinations of the five inputs are listed. For the conditions manifested in the patient, different drugs and dosages will be required. The proper treatment for each combination is listed in the output column of Table 5. After the algorithm runs the comparison, the output is displayed on the computer screen, prompting the intensivist with the proper treatment **3712**.

Referring to FIG. 41A, the ventilator weaning decision support algorithm of the present invention is illustrate The ventilator weaning decision support algorithm is used to determine whether an intensive care unit patient can return to breathing unassisted, and discontinue use of a ventilator. Such a determination weures evaluation of the patient by the intensivist over the course of several days.

To begin the decision process of whether to wean a patient from ventilator use, the intensivist is prompted to conduct daily screening, preferably during the hours of 06:00 a.m. to 10:00 a.m. **3800**. The daily screen prompts the intensivist to determine whether: the patients P/F ratio is greater than 200, the patient's positive end-expiratory pressure (PEEP) is less than or equal to 5, whether cough suctioning has been adequate and/or spontaneous, infusions with vasopressors have been necessary, and continuous infusions of sedatives or neuromuscular blocking agents have been necessary **3800**. If all conditions **3802** are answered no, the intensivist is directed by the system to repeat the daily screen **3805** the following morning. If all the conditions of the daily screen are met **3802**, the intensivist is prompted to perform additional tests.

If the patient has satisfied the daily screen, the intensivist is next directed to conduct a rapid shallow breathing test **3804**. To perform the test, the intensivist is directed to change the ventilator setting to continuous positive airway pressure (CPAP) less than or equal to 5. In other words, there is no intermittent mandatory ventilation or pressure support provided for the patient. The patient is given one minute to reach a steady state of breathing. Then the intensivist measures the ratio of breaths per minute to tidal volume (f/V.sub.T). The intensivist next is prompted to determine whether the patient's f/V.sub.T is less than or equal to 105 breathes per minute **3806**. If the patient's f/V.sub.T is greater than 105 breathes per minute, the intensivist is prompted to return to performing daily screening the following morning **3808**.

If the patient's f/V.sub.T is less than or equal to 105 breathes per minute, the intensivist is next directed to perform a trial of spontaneous breathing. Here, the intensivist can either insert a T-Piece in the patient's airway or reduce the patient's CPAP to less than or equal to 5 over the course of two hours. The intensivist is prompted to observe the patient periodically in order to evaluate if the patient is breathing without assistance **3810**. The intensivist is prompted to perform a periodic assessment by determining whether: the patient's breathing characteristics are greater

than 35 breaths per minute for 5 minutes, or SpO.sub.2 is less than 90%, or the patient's Heart Rate (HR) is grater than 140, or HR deviates from the baseline breathing rate by more than 20%, or the patient's SBP is outside the range of 90 to 180. If any of the conditions are met, the intensivist is directed by the system to terminate ventilator weaning **3812**. If the conditions are not met, the patient is further assessed.

In further assessment, the intensivist is prompted to determine whether the patient has been able to breathe spontaneously for two hours, keep a clear airway, and does not have any procedures scheduled within twenty-four hours that would require the patient to be intubated **3814**. If the patient meets all of these criteria **3814**, the intensivist is notified by the system that the patient may be extubated **3816**. If the patient does not meet one or more of the criteria **3814**, the intensivist is prompted to perform steps for progressive weaning **3818**.

Referring to FIG. 41B, the ventilator weaning decision support algorithm of the present invention is further illustrated. The intensivist, at his or her discretion may choose either T-piece progressive weaning or pressure support progressive weaning. In order to perform T-piece progressive weaning, the intensivist is directed to repeat the trial of spontaneous breathing (as previously described **3810**). The intensivist can either insert a T-piece in the patient's airway or reduce the patient's CPAP to less than or equal to 5 over the course of two hours. The intensivist is prompted to perform periodic assessment of the patient by either a two hour or 30 minute trial **3820**.

In order to perform pressure support progressive weaning, the intensivist is first prompted to observe whether the patient's pressure support (PS) rating is equal to eighteen plus or minus the positive end-expiratory pressure (PEEP). Next, the intensivist is directed by the system to regulate the pressure values in order to keep the patient's respiratory rate (RR) between twenty and thirty. Next, the intensivist is directed by the system to decrease the patient's pressure support by 2-4 centimeters of water two times per day. Once the patient maintains pressure support for at least two hours, the intensivist is prompted to further pursue extubating the patient **3822**.

After either T-Piece progressive weaning **3820** or pressure support progressive weaning **3822**, the intensivist is next prompted to perform a periodic assessment of the patient. Here, the intensivist must determine whether whether: the patient's breathing characteristics are greater than 35 breaths per minute for 5 minutes, or SpO.sub.2 is less than 90%, or the patient's HR is grater than 140, or HR deviates from the baseline breathing rate by more than 20%; or the patient's SBP is outside the range of 90 to 180. Where the patient meets any of these criteria, the intensivist is prompted to terminate weaning. If the patient meets none of these criteria; the intensivist is prompted to further assess the patient's ability to breath spontaneously **3824**.

In further assessment, the intensivist is prompted to determine whether the patient has been able to breathe spontaneously for two hours, keep a clear airway, and does not have any procedures scheduled within twenty-four hours that would require the patient to be intubated **3826**. If the patient meets all of these criteria **3814**, the intensivist is notified by the system that the patient may be extubated **3828**. If the patient does not meet one or more of the criteria **3826**, the intensivist is directed by the system to allow the patient to rest for at least twelve hours at A/C, the last level of pressure support the patient achieved **3830**. The intensivist is prompted to resume progressive weaning the following day **3832**.

Referring to FIG. 42, the Warfarin Dosing Algorithm of the present invention is illustrated. The intensivist is first prompted to give the initial dose and determine subsequent dosage each day 3900. When the intensivist determines subsequent dosage, he is first prompted to determine the patient's target INR 3902. If the patient's target INR ranges from 2.0 to 3.0, the intensivist is prompted by the system to make further determinations relevant to dosage. The intensivist is directed by the system to determine whether the patient is taking drugs that effect prothrombin time 3904, the baseline INR value 3906, and whether rapid anticoagulation is required 3908. Each answer is assigned a point value, and the total points are tabulated. If the point value is greater than one, the system refers to the 10 milligram load target database for dosing. If the point value is less than one, the system refers to the 5 milligram load target database for dosing 3910.

At the initial INR determination 3902, if the patient's INR was initially between 1.5 and 2.0, the system refers to the 5 milligram load target database for dosing. If the patient's INR was initially between 3.0 and 4.0, the system refers to the 10 milligram load target database for dosing 3910. Next the intensivist is prompted to enter the day of treatment 3912 and the patient's INR 3914. Depending on whether the system has been directed to the 5 milligram load target or the 10 milligram load target, a comparison is run 3916 according to the following tables.

TABLE 6

5 mg Load Target INR 1.5-2.0				
Day	<1.5	1.5-2	2-2.5	>2.5
2	5	1.25-2.5	0	0
3	5-7.5	1.25-2.5	0-1.25	0
4	10-(Check to see whether pt has received vit K)	1.25-2.5	0-1.25	0
5	10 (Check to see whether pt Has received vit K)	2.5-5	0-2.5	0-1.25
6	15 Obtain hematology consultation.	2.5-5	1.25-2.5	0-1.25

TABLE 7

10 mg Load Target INR 3.0-4.0					
Day	<1.5	1.5-2	2-2.5	2.5-3	>3
2	10	7.5-10	5-7.5	2.5-5.0	0-2.5
3	10-15	7.5-10	5-7.5	2.5-5	2.5-5
4	10-15 (Check to see whether pt has received vit K)	7.5-12.5	5-10	5-7.5	2.5-5
5	15 (Check to see whether pt has received vit K)	10-12.5	7.5-10	5-7.5	2.5-5
6	15-20 obtain hematology consultation.	10-15	7.5-12.5	5-10	5-7.5

The appropriate dosage and instructions is displayed on the computer screen to the intensivist 3918.

Referring to FIG. 43, the heparin-induced thrombocytopenia (HIT) decision support algorithm of the present invention is illustrated. The intensivist is prompted to observe whether the patient's platelet count has dropped

50% or more over seventy-two hours while being treated with heparin, and whether any other obvious causes of platelet reduction might be present 4100. If such a drop has not occurred, the intensivist is notified by the system that the patient most likely does not have HIT, but monitoring of the platelet count should continue 4102. If the patient's platelet count has drastically dropped, the intensivist is prompted to determine whether the patient has been treated with heparin for more than three days 4104. Regardless of the answer, the intensivist is next prompted to determine if the patient has been treated with heparin in the preceding three months 4106. If the patient has not received heparin in the preceding three months, the intensivist is notified by the system that HIT is not likely to be the cause of the platelet drop. The intensivist is also prompted to monitor platelet count for infection or other thrombocytopenia-causing drugs, and to consider stopping heparin therapy if the platelet count drops below 50,000 per cubic millimeter 4108.

If the patient has received heparin in the last three days 4104, the intensivist is further prompted to look for signs of thrombosis, or blood clotting 4110. If the patient shows signs of thrombosis, the intensivist is notified by the system that the patient is likely to have HIT. Accordingly, the intensivist is prompted to stop administering heparin and flush any drug administration equipment that would contain heparin traces. The intensivist is also provided instructions by the system to treat a patient still requiring anticoagulation treatment with alternate drugs and methods 4112.

Where the patient does not show signs of thrombosis 4110, the intensivist is prompted to check for heparin resistance 4114. Signs of heparin resistance include inability to hold aPTT though heparin doses have been increase. If the patient shows signs of heparin resistance, the intensivist is prompted to consider stopping heparin treatment and to consider treating a patient still requiring anticoagulation treatment with alternate drugs and methods 4116. If the patient does not show signs of heparin resistance, the intensivist is notified by the system that the patient possibly has HIT. The intensivist is accordingly prompted to continue monitoring for thrombosis, consider infection or other drugs that cause thrombocytopenia, and to consider stopping heparin therapy if the platelet count drops below 50,000 per cubic millimeter 4118

Results

The structure of the present invention and its efficacy have yielded striking results in practice. In a research setting, deployment of certain rudimentary aspects of the present the invention designed to experimentally test the approach described and developed in detail above, yielded unprecedented improvements in clinical and economic outcomes: 50% improvement in severity adjusted mortality, 40% improvement in clinical complication rates, 30% improvement in ICU length of stay, and 30% improvement in overall ICU cost of care.

A system and method for accounting and billing patients in a hospital environment has been shown. It will be apparent to those skilled in the art that other variations of the present invention are possible without departing from the scope of the invention as disclosed. For example, one can envision different ratios of command center/remote location to ICU's, other decision support algorithms that would be used by intensivists, other types of remote monitoring of not only hospitalized patients but other types of hospital functions as well as industrial functions where critical expertise is in limited supply but where that expertise must be applied

to ongoing processes. In such cases a system such as that described can be employed to monitor processes and to provide standardized interventions across a number of geographically dispersed locations and operations and to provide billing for services provided. Further, any reference to claim elements in the singular, for example, using the articles "a," "an," or "the" is not to be construed as limiting the element to the singular.

What is claimed is:

1. A patient monitoring system for geographically dispersed health care locations comprising a network, a body-worn monitoring station, a monitoring station server, a remote command center, and a rules engine, wherein,

the network comprises a first sub-network and a second sub-network;

the body-worn monitoring station comprises monitoring equipment, and wherein the body worn monitoring station comprises instructions for:

monitoring data elements from a patient assigned to a health care location; and sending monitored data elements to a monitoring station server via the first sub-network;

the monitoring station server comprises instructions for: receiving the monitored data elements from the body worn monitoring station; and

relaying the monitored data elements to the remote command center via the second sub-network;

the remote command center comprises instructions for: receiving the monitored data elements from the monitoring station server;

associating the monitored data elements with the patient assigned to the health care location;

accessing patient data elements indicative of a medical condition associated with the patient;

establishing a patient-specific rule associated with the patient; and

the rules engine comprises instructions for:

selecting data elements from the monitored data elements and the patient data elements associated with the patient;

applying the patient-specific rule to the selected data elements continuously and simultaneously;

determining in an automated fashion 24 hours per day 7 days per week from the remote command center whether the patient-specific rule for the patient has been contravened; and

issuing an alert from the remote command center in the event the patient-specific rule for the patient has been contravened.

2. The patient monitoring system of claim 1, wherein the patient specific rule for the patient comprises an algorithm.

3. The patient monitoring system of claim 1, wherein the selected data elements comprise a physiological data element of the patient and a clinical data element of the patient.

4. The patient monitoring system of claim 1, wherein the selected data elements comprise a physiological data element of the patient and a medication data element of the patient.

5. The patient monitoring system of claim 1, wherein the selected data elements comprise a physiological data element of the patient and a laboratory data element of the patient.

6. The patient monitoring system of claim 1, wherein the selected data elements comprise a clinical data element of the patient and a laboratory data element of the patient.

7. The patient monitoring system of claim 1, wherein the selected data elements comprise a physiological data element of the patient and another physiological data element of the patient.

8. The patient monitoring system of claim 1, wherein the selected data elements comprise at least two data elements of the patient selected from the group consisting of a physiological data element, a clinical data element of the patient, a medication data element of the patient, and a laboratory data element of the patient.

9. The patient monitoring system of claim 1, wherein the first and second sub-networks are selected from the group consisting of a wired network and a wireless network.

10. The patient monitoring system of claim 9, wherein the wireless network is selected from the group consisting of a Bluetooth-compatible network, an 802.11-compatible network, and a short-range, slotted protocol network.

11. The patient monitoring system of claim 1, wherein the monitoring station server is wearable by the patient.

12. The patient monitoring system of claim 11, wherein the first sub-network is a wired network and the second sub-network is a wireless network.

13. The patient monitoring system of claim 11, wherein the first sub-network is a wireless network and the second sub-network is a wireless network.

14. The patient monitoring system of claim 1, wherein the patient is a fetus carried by an expectant mother and wherein the monitoring equipment comprises a tachodynamometer.

15. The patient monitoring system of claim 1, wherein the monitoring equipment is at least one sensor selected from the group consisting of an ECG sensor, a pulse oximeter, a thermometer, a respiration sensor, a heart rate sensor, and an audio sensor.

16. The patient monitoring system of claim 1, wherein the monitored data elements are associated with a patient identifier.

17. The patient monitoring system of claim 1, wherein the health care location is selected from the group consisting of a hospital, a nursing home, a mobile health care facility, a space-based health care facility, a residence, an emergency room, an intensive care unit, a field health care facility, an operating room, and a step-down unit.

18. The patient monitoring system of claim 17, wherein the mobile health care facility is selected from the group consisting of a ship, a helicopter, an ambulance, a MASH unit, and a tent hospital.

19. The patient monitoring system of claim 1, wherein monitoring equipment comprises physiological sensors and wherein monitored data elements comprise physiological data elements.

20. The patient monitoring system of claim 1, wherein the remote command center further comprises:

an external network interface, wherein the external network interface comprises instructions for connecting to an external network; and

instructions for providing a health care provider access to the remote command center via the external network.

21. The patient monitoring system of claim 20, wherein the external network is selected from the group consisting of a wired network, a wireless network, a cable network, a fiber optic network, and the Internet.

22. The patient monitoring system of claim 20, wherein the health care provider is selected from the group consisting of a physician, a nurse, a clinician, a diagnostician, and a intensivist.

23. The patient monitoring system of claim 20, wherein the remote command center further comprises instructions

for sending the health care provider the alert if the patient-specific rule for the hospitalized patient has been contravened.

24. A method for monitoring patients assigned to geographically dispersed health care locations comprising:

associating a body-worn monitoring station with a patient assigned to a health care location, wherein the body-worn monitoring station comprises monitoring equipment;

monitoring data elements from a patient assigned to a health care location;

sending monitored data elements to a monitoring station server via a first sub-network;

relaying the monitored data elements from the monitoring station server to a remote command center via a second sub-network;

receiving the monitored data elements at the remote command center from the monitoring station server;

associating the monitored data elements with the patient assigned to the health care location;

accessing patient data elements indicative of a medical condition associated with the patient;

establishing a patient-specific rule associated with the patient;

selecting data elements from the monitored data elements and the patient data elements associated with the patient;

applying the patient-specific rule to the selected data elements continuously and simultaneously;

determining in an automated fashion 24 hours per day 7 days per week at the remote command center whether the patient-specific rule for the patient has been contravened;

and

issuing an alert from the remote command center in the event the patient-specific rule for the patient has been contravened.

25. The method of claim 24, wherein the patient specific rule for the patient comprises an algorithm.

26. The method of claim 24, wherein the selected data elements comprise a physiological data element of the patient and a clinical data element of the patient.

27. The method of claim 24, wherein the selected data elements comprise a physiological data element of the patient and a medication data element of the patient.

28. The method of claim 24, wherein the selected data elements comprise a physiological data element of the patient and a laboratory data element of the patient.

29. The method of claim 24, wherein the selected data elements comprise a clinical data element of the patient and a laboratory data element of the patient.

30. The method of claim 24, wherein the selected data elements comprise a physiological data element of the patient and another physiological data element of the patient.

31. The method of claim 24, wherein the selected data elements comprise at least two data elements of the patient selected from the group consisting of a physiological data

element, a clinical data element of the patient, a medication data element of the patient, and a laboratory data element of the patient.

32. The method of claim 24, wherein the first and second sub-networks are selected from the group consisting of a wired network and a wireless network.

33. The method of claim 24, wherein the monitoring station server is wearable by the patient.

34. The method of claim 32, wherein wireless network is selected from the group consisting of a Bluetooth-compatible network, an 802.11-compatible network, and a short-range, slotted protocol network.

35. The method of claim 33, wherein the first sub-network is a wired network and the second sub-network is a wireless network.

36. The method of claim 33, wherein the first sub-network is a wireless network and the second sub-network is a wireless network.

37. The method of claim 24, wherein the patient is a fetus carried by an expectant mother and wherein the monitoring equipment comprises a tachodynamometer.

38. The method of claim 24, wherein the monitoring equipment is at least one sensor selected from the group consisting of an ECG sensor, a pulse oximeter, a thermometer, a respiration sensor, a heart rate sensor, and an audio sensor.

39. The method of claim 24, wherein the monitored data elements are associated with a patient identifier.

40. The method of claim 24, wherein the health care location is selected from the group consisting of a hospital, a nursing home, a mobile health care facility, a space-based health care facility, a residence, an emergency room, an intensive care unit, a field health care facility, an operating room, and a step-down unit.

41. The method of claim 40, wherein the mobile health care facility is selected from the group consisting of a ship, a helicopter, an ambulance, a MASH unit, and a tent hospital.

42. The method of claim 24, wherein monitoring equipment comprises physiological sensors and wherein monitored data elements comprise physiological data elements.

43. The method of claim 24 further comprising:

interfacing with an external network; and

providing a health care provider access to the remote command center via the external network.

44. The method of claim 43, wherein the external network is selected from the group consisting of a wired network, a wireless network, a cable network, a fiber optic network, and the Internet.

45. The method of claim 43, wherein the health care provider is selected from the group consisting of a physician, a nurse, a clinician, a diagnostician, and a intensivist.

46. The method of claim 43 further comprising sending the health care provider the alert if the patient-specific rule for the hospitalized patient has been contravened.

\* \* \* \* \*

专利名称(译)	用于在地理上分散的医疗保健场所中患者佩戴式监测患者的系统和方法		
公开(公告)号	<a href="#">US7321862</a>	公开(公告)日	2008-01-22
申请号	US11/268706	申请日	2005-11-07
[标]申请(专利权)人(译)	VISICU		
申请(专利权)人(译)	VISICU INC.		
当前申请(专利权)人(译)	VISICU INC.		
[标]发明人	ROSENFELD BRIAN A BRESLOW MICHAEL		
发明人	ROSENFELD, BRIAN A. BRESLOW, MICHAEL		
IPC分类号	G06F19/00 A61B5/00 A61B5/0205 G08B21/04		
CPC分类号	A61B5/412 G06F19/322 G06F19/325 G06F19/327 G06F19/3418 G06F19/3425 G06F19/345 G06F19/3481 G06Q50/22 G06Q50/24 G08B21/0476 A61B5/0205 G06F19/328 A61B5/002 A61B5/0022 A61B5/0006 A61B5/4094 A61B5/411 G06Q40/08 G08B21/182 G16H10/60 G16H20/30 G16H40/20 G16H40/67 G16H50/20 G16H70/20 G16H80/00		
审查员(译)	MORGAN, ROBERTW		
优先权	09/443072 2004-10-12 US 60/141520 1999-06-23 US		
其他公开文献	US20060085227A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

用于地理上分散的医疗保健场所的患者佩戴的监测系统和方法。患者佩戴的监测系统包括网络，体佩监测站，监测站服务器，远程命令中心和规则引擎。该网络包括第一子网和第二子网。体佩监测站包括监测设备。身体佩戴的监测站监测来自分配给医疗保健位置的患者的数据元素，并经由第一子网将监测的数据元素发送到监测站服务器。监测站服务器从身体佩戴的监测站接收监测的数据元素，并通过第二子网将监测的数据元素中继到远程命令中心。该远程命令中心从监测站服务器接收监控的数据元素，将监控的数据元素与分配给医疗保健位置的患者相关联，访问指示与患者相关的医疗状况的患者数据元素，并建立患者特定的规则与病人有关。规则引擎从监视的数据元素和与患者相关联的患者数据元素中选择数据元素，并且连续且同时地将患者特定的规则应用于所选择的数据元素。确定是否违反了患者的患者特异性规则。如果患者的患者特定规则已经发生，则从远程命令中心发出警报违反。

