



US010179028B2

(12) **United States Patent**
Levin et al.

(10) **Patent No.:** **US 10,179,028 B2**
(45) **Date of Patent:** ***Jan. 15, 2019**

(54) **METHODS FOR TREATING PATIENTS VIA
RENAL NEUROMODULATION**

(71) Applicant: **Medtronic Ardian Luxembourg
S.a.r.l.**, Luxembourg (LU)

(72) Inventors: **Howard R. Levin**, Teaneck, NJ (US);
Mark Gelfand, New York, NY (US)

(73) Assignee: **Medtronic Ardian Luxembourg
S.a.r.l.**, Luxembourg (LU)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **15/878,746**

(22) Filed: **Jan. 24, 2018**

(65) **Prior Publication Data**

US 2018/0228535 A1 Aug. 16, 2018

Related U.S. Application Data

(63) Continuation of application No. 15/476,867, filed on
Mar. 31, 2017, now Pat. No. 9,907,611, which is a
(Continued)

(51) **Int. Cl.**
A61B 5/00 (2006.01)
A61B 5/044 (2006.01)
(Continued)

(52) **U.S. Cl.**
CPC **A61B 18/1492** (2013.01); **A61B 18/04**
(2013.01); **A61M 5/142** (2013.01); **A61M**
5/14276 (2013.01); **A61M 5/1723** (2013.01);
A61N 1/05 (2013.01); **A61N 1/0551**
(2013.01); **A61N 1/326** (2013.01);
(Continued)

(58) **Field of Classification Search**

CPC **A61B 18/1492**; **A61B 18/04**; **A61B**
2018/00404; **A61B 2018/00511**; **A61B**
2018/00434; **A61B 2018/00577**; **A61N**
1/403; **A61N 1/40**; **A61N 1/36135**; **A61N**
1/36125; **A61N 5/00**; **A61N 1/05**; **A61N**
1/0551; **A61N 1/36114**; **A61N 1/36117**;
A61N 1/326; **A61N 1/3627**;
(Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,130,758 A 9/1938 Rose
2,276,995 A 3/1942 Milinowski
(Continued)

FOREIGN PATENT DOCUMENTS

DE 3151180 8/1982
EP 0811395 12/1997
(Continued)

OTHER PUBLICATIONS


U.S. Appl. No. 95/002,110, filed Aug. 29, 2012, Demarais et al.
(Continued)

Primary Examiner — Mark W Bockelman

(57) **ABSTRACT**

A method and apparatus for treatment of heart failure, hypertension and renal failure by stimulating the renal nerve. The goal of therapy is to reduce sympathetic activity of the renal nerve. Therapy is accomplished by at least partially blocking the nerve with drug infusion or electro-stimulation. Apparatus can be permanently implanted or catheter based.

20 Claims, 9 Drawing Sheets



Related U.S. Application Data

continuation of application No. 15/095,220, filed on Apr. 11, 2016, which is a continuation of application No. 14/846,480, filed on Sep. 4, 2015, which is a continuation of application No. 13/617,994, filed on Sep. 14, 2012, which is a continuation of application No. 13/361,019, filed on Jan. 30, 2012, which is a continuation of application No. 11/688,178, filed on Mar. 19, 2007, now Pat. No. 8,131,372, which is a continuation of application No. 11/144,173, filed on Jun. 3, 2005, now Pat. No. 7,647,115, which is a continuation of application No. 10/408,665, filed on Apr. 8, 2003, now Pat. No. 7,162,303.

- (60) Provisional application No. 60/442,970, filed on Jan. 29, 2003, provisional application No. 60/415,575, filed on Oct. 3, 2002, provisional application No. 60/370,190, filed on Apr. 8, 2002.

(51) Int. Cl.

A61B 5/0452 (2006.01)
A61B 5/0402 (2006.01)
A61B 18/14 (2006.01)
A61M 5/142 (2006.01)
A61M 5/172 (2006.01)
A61N 1/05 (2006.01)
A61N 1/32 (2006.01)
A61N 5/00 (2006.01)
A61B 18/00 (2006.01)
A61N 1/40 (2006.01)
A61N 1/36 (2006.01)
A61N 1/362 (2006.01)
A61M 1/36 (2006.01)
A61B 18/00 (2006.01)

(52) U.S. Cl.

CPC *A61N 1/36114* (2013.01); *A61N 1/36117* (2013.01); *A61N 1/36125* (2013.01); *A61N 1/36135* (2013.01); *A61N 1/40* (2013.01); *A61N 1/403* (2013.01); *A61N 5/00* (2013.01); *A61B 2018/00404* (2013.01); *A61B 2018/00434* (2013.01); *A61B 2018/00511* (2013.01); *A61B 2018/00577* (2013.01); *A61M 1/3627* (2013.01); *A61M 2210/1082* (2013.01); *A61N 1/36007* (2013.01); *A61N 1/3627* (2013.01)

(58) Field of Classification Search

CPC *A61N 1/36007*; *A61M 5/142*; *A61M 5/14276*; *A61M 5/1723*; *A61M 1/3627*; *A61M 2210/1082*

See application file for complete search history.

(56) References Cited

U.S. PATENT DOCUMENTS

2,276,996 A 3/1942 Milinowski
 3,043,310 A 7/1962 Milinowski
 3,127,895 A 4/1964 Kendall et al.
 3,181,535 A 5/1965 Milinowski
 3,270,746 A 9/1966 Kendall et al.
 3,329,149 A 7/1967 Kendall et al.
 3,522,811 A 8/1970 Schwartz et al.
 3,563,246 A 2/1971 Puharich et al.
 3,650,277 A 3/1972 Sjostrand et al.
 3,670,737 A 6/1972 Pearo
 3,760,812 A 9/1973 Timm et al.
 3,774,620 A 11/1973 Hansjurgens et al.
 3,794,022 A 2/1974 Nawracaj et al.

3,800,802 A 4/1974 Berry et al.
 3,803,463 A 4/1974 Cover
 3,894,532 A 7/1975 Morey
 3,895,639 A 7/1975 Rodler et al.
 3,897,789 A 8/1975 Blanchard
 3,911,930 A 10/1975 Hagfors et al.
 3,952,751 A 4/1976 Yarger
 3,987,790 A 10/1976 Eckenhoff et al.
 4,011,861 A 3/1977 Enger
 4,026,300 A 5/1977 DeLuca et al.
 4,055,190 A 10/1977 Tany et al.
 4,071,033 A 1/1978 Nawracaj et al.
 4,105,017 A 8/1978 Ryaby et al.
 4,141,365 A 2/1979 Fischell et al.
 4,266,532 A 5/1981 Ryaby et al.
 4,266,533 A 5/1981 Ryaby et al.
 4,305,115 A 12/1981 Armitage et al.
 4,315,503 A 2/1982 Ryaby et al.
 4,360,019 A 11/1982 Fortner et al.
 4,379,462 A 4/1983 Borkan et al.
 4,405,305 A 9/1983 Stephen et al.
 4,454,883 A 6/1984 Fellus et al.
 4,467,808 A 8/1984 Brighton et al.
 4,487,603 A 12/1984 Harris
 4,530,840 A 7/1985 Tice et al.
 4,587,975 A 5/1986 Salo et al.
 4,602,624 A 7/1986 Naples et al.
 4,608,985 A 9/1986 Crish et al.
 4,649,936 A 3/1987 Ungar et al.
 4,671,286 A 6/1987 Renault et al.
 4,674,482 A 6/1987 Waltonen et al.
 4,692,147 A 9/1987 Duggan
 4,709,698 A 12/1987 Johnston et al.
 4,715,852 A 12/1987 Reinicke et al.
 4,774,967 A 10/1988 Zanakos et al.
 4,791,931 A 12/1988 Slate
 4,816,016 A 3/1989 Schulte et al.
 4,852,573 A 8/1989 Kennedy
 4,865,845 A 9/1989 Eckenhoff et al.
 4,890,623 A 1/1990 Cook et al.
 4,976,711 A 12/1990 Parins et al.
 4,979,511 A 12/1990 Terry, Jr.
 4,981,146 A 1/1991 Bertolucci
 4,998,532 A 3/1991 Griffith
 5,006,119 A 4/1991 Acker et al.
 5,014,699 A 5/1991 Pollack et al.
 5,019,034 A 5/1991 Weaver et al.
 5,057,318 A 10/1991 Magruder et al.
 5,058,584 A 10/1991 Bourgeois et al.
 5,059,423 A 10/1991 Magruder et al.
 5,061,492 A 10/1991 Okada et al.
 5,094,242 A 3/1992 Gleason et al.
 5,111,815 A 5/1992 Mower
 5,112,614 A 5/1992 Magruder et al.
 5,125,928 A 6/1992 Parins et al.
 5,131,409 A 7/1992 Lobarev et al.
 5,137,727 A 8/1992 Eckenhoff
 5,188,837 A 2/1993 Domb
 5,193,048 A 3/1993 Kaufman et al.
 5,193,539 A 3/1993 Schulman et al.
 5,193,540 A 3/1993 Schulman et al.
 5,199,428 A 4/1993 Obel et al.
 5,203,326 A 4/1993 Collins et al.
 5,213,098 A 5/1993 Bennett et al.
 5,215,086 A 6/1993 Terry, Jr. et al.
 5,231,988 A 8/1993 Wernicke et al.
 5,234,692 A 8/1993 Magruder et al.
 5,234,693 A 8/1993 Magruder et al.
 5,251,634 A 10/1993 Weinberg
 5,251,643 A 10/1993 Osypka et al.
 5,263,480 A 11/1993 Wernicke et al.
 5,269,303 A 12/1993 Wernicke et al.
 5,282,468 A 2/1994 Klepinski
 5,282,785 A 2/1994 Shapland et al.
 5,286,254 A 2/1994 Shapland et al.
 5,299,569 A 4/1994 Wernicke et al.
 5,300,068 A 4/1994 Rosar et al.
 5,304,120 A 4/1994 Crandell et al.
 5,304,206 A 4/1994 Baker, Jr. et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,317,155 A	5/1994	King	5,814,079 A	9/1998	Kieval
5,324,255 A	6/1994	Passafaro et al.	5,824,087 A	10/1998	Aspden et al.
5,324,316 A	6/1994	Schulman et al.	5,836,935 A	11/1998	Ashton et al.
5,334,193 A	8/1994	Nardella	RE35,987 E	12/1998	Harris et al.
5,335,657 A	8/1994	Terry, Jr. et al.	5,843,069 A	12/1998	Butler et al.
5,338,662 A	8/1994	Sadri	5,860,974 A	1/1999	Abele
5,351,394 A	10/1994	Weinberg	5,861,021 A	1/1999	Thome et al.
5,358,514 A	10/1994	Schulman et al.	5,865,787 A	2/1999	Shapland et al.
5,368,591 A	11/1994	Lennox et al.	5,871,449 A	2/1999	Brown
5,370,680 A	12/1994	Proctor	5,891,181 A	4/1999	Zhu et al.
5,389,069 A	2/1995	Weaver	5,893,885 A	4/1999	Webster et al.
5,397,308 A	3/1995	Ellis et al.	5,906,636 A	5/1999	Casscells, III et al.
5,397,338 A	3/1995	Grey et al.	5,906,817 A	5/1999	Moullier et al.
5,400,784 A	3/1995	Durand et al.	5,913,876 A	6/1999	Taylor et al.
5,405,367 A	4/1995	Schulman et al.	5,916,154 A	6/1999	Hobbs et al.
5,423,744 A	6/1995	Gencheff et al.	5,916,239 A	6/1999	Geddes et al.
5,429,634 A	7/1995	Narciso, Jr.	5,919,187 A	7/1999	Guglielmi et al.
5,433,739 A	7/1995	Sluijter et al.	5,924,997 A	7/1999	Campbell
5,439,440 A	8/1995	Hofmann	5,928,272 A	7/1999	Adkins et al.
5,454,782 A	10/1995	Perkins	5,935,075 A	8/1999	Casscells et al.
5,454,809 A	10/1995	Janssen	5,944,710 A	8/1999	Dev et al.
5,458,568 A	10/1995	Racchini et al.	5,954,719 A	9/1999	Chen et al.
5,458,626 A	10/1995	Krause	5,983,131 A	11/1999	Weaver et al.
5,458,631 A	10/1995	Xavier	5,983,141 A	11/1999	Sluijter et al.
5,470,352 A	11/1995	Rappaport	6,004,269 A	12/1999	Crowley et al.
5,472,406 A	12/1995	de la Torre et al.	6,006,134 A	12/1999	Hill et al.
5,478,303 A	12/1995	Foley-Nolan et al.	6,009,877 A	1/2000	Edwards
5,484,400 A	1/1996	Edwards et al.	6,010,613 A	1/2000	Walters et al.
5,494,822 A	2/1996	Sadri	6,024,740 A	2/2000	Lesh
5,498,238 A	3/1996	Shapland et al.	6,026,326 A	2/2000	Bardy
5,499,971 A	3/1996	Shapland et al.	6,036,687 A	3/2000	Laufer et al.
5,505,700 A	4/1996	Leone et al.	6,041,252 A	3/2000	Walker et al.
5,507,724 A	4/1996	Hofmann et al.	6,051,017 A	4/2000	Loeb et al.
5,507,791 A	4/1996	Sit'ko et al.	6,058,328 A	5/2000	Levine et al.
5,531,778 A	7/1996	Maschino et al.	6,058,331 A	5/2000	King
5,540,730 A	7/1996	Terry, Jr. et al.	6,066,134 A	5/2000	Eggers et al.
5,540,734 A	7/1996	Zabara	6,073,048 A	6/2000	Kieval et al.
5,553,611 A	9/1996	Budd et al.	6,077,227 A	6/2000	Miesel et al.
5,560,360 A	10/1996	Filler et al.	6,086,527 A	7/2000	Talpade
5,569,198 A	10/1996	Racchini	6,091,995 A	7/2000	Ingle et al.
5,571,147 A	11/1996	Sluijter et al.	6,117,101 A	9/2000	Diederich et al.
5,571,150 A	11/1996	Wernicke et al.	6,122,548 A	9/2000	Starkebaum et al.
5,573,552 A	11/1996	Hansjurgens et al.	6,123,718 A	9/2000	Tu et al.
5,584,863 A	12/1996	Rauch et al.	6,135,999 A	10/2000	Fanton et al.
5,588,964 A	12/1996	Imran et al.	6,146,380 A	11/2000	Racz et al.
5,589,192 A	12/1996	Okabe et al.	6,149,620 A	11/2000	Baker et al.
5,599,345 A	2/1997	Edwards et al.	6,161,048 A	12/2000	Sluijter et al.
5,618,563 A	4/1997	Berde et al.	6,171,306 B1	1/2001	Swanson et al.
5,626,576 A	5/1997	Janssen	6,178,349 B1	1/2001	Kieval
5,626,862 A	5/1997	Brem et al.	6,190,353 B1	2/2001	Makower et al.
5,628,730 A	5/1997	Shapland et al.	6,192,889 B1	2/2001	Morrish
5,634,462 A	6/1997	Tyler et al.	6,205,361 B1	3/2001	Kuzma et al.
5,634,899 A	6/1997	Shapland et al.	6,208,894 B1	3/2001	Schulman et al.
5,672,174 A	9/1997	Gough et al.	6,214,032 B1	4/2001	Loeb et al.
5,688,266 A	11/1997	Edwards et al.	6,219,577 B1	4/2001	Brown, III et al.
5,689,877 A	11/1997	Grill, Jr. et al.	6,224,592 B1	5/2001	Eggers et al.
5,690,691 A	11/1997	Chen et al.	6,238,702 B1	5/2001	Berde et al.
5,700,282 A	12/1997	Zabara	6,245,026 B1	6/2001	Campbell et al.
5,700,485 A	12/1997	Berde et al.	6,246,912 B1	6/2001	Sluijter et al.
5,704,908 A	1/1998	Hofmann et al.	6,251,130 B1	6/2001	Dobak, III et al.
5,707,400 A	1/1998	Terry, Jr. et al.	6,254,598 B1	7/2001	Edwards et al.
5,711,326 A	1/1998	Thies et al.	6,258,087 B1	7/2001	Edwards et al.
5,713,847 A	2/1998	Howard, III et al.	6,259,952 B1	7/2001	Sluijter et al.
5,722,401 A	3/1998	Pietroski et al.	6,269,269 B1	7/2001	Ottenhoff et al.
5,723,001 A	3/1998	Pilla et al.	6,272,377 B1	8/2001	Sweeney et al.
5,725,563 A	3/1998	Klotz et al.	6,272,383 B1	8/2001	Grey et al.
5,728,396 A	3/1998	Peery et al.	6,273,886 B1	8/2001	Edwards et al.
5,747,060 A	5/1998	Sackler et al.	6,280,377 B1	8/2001	Talpade
5,755,750 A	5/1998	Petruska et al.	6,287,304 B1	9/2001	Eggers et al.
5,756,115 A	5/1998	Moo-Young et al.	6,287,608 B1	9/2001	Levin et al.
5,772,590 A	6/1998	Webster, Jr.	6,292,695 B1	9/2001	Webster, Jr. et al.
5,792,187 A	8/1998	Adams	6,304,777 B1	10/2001	Ben-Haim et al.
5,800,464 A	9/1998	Kieval	6,304,787 B1	10/2001	Kuzma et al.
5,807,306 A	9/1998	Shapland et al.	6,306,423 B1	10/2001	Donovan et al.
5,810,802 A	9/1998	Panescu et al.	6,314,325 B1	11/2001	Fitz
			6,322,558 B1	11/2001	Taylor et al.
			6,322,559 B1	11/2001	Daulton et al.
			6,326,020 B1	12/2001	Kohane et al.
			6,326,177 B1	12/2001	Schoenbach et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

6,328,699	B1	12/2001	Eigler et al.	6,916,656	B2	7/2005	Walters et al.
6,334,069	B1	12/2001	George et al.	6,927,049	B2	8/2005	Rubinsky et al.
6,347,247	B1	2/2002	Dev et al.	6,936,047	B2	8/2005	Nasab et al.
6,353,763	B1	3/2002	George et al.	6,939,345	B2	9/2005	KenKnight et al.
6,356,786	B1	3/2002	Rezai et al.	6,939,346	B2	9/2005	Kannenberget al.
6,356,787	B1	3/2002	Rezai et al.	6,958,060	B2	10/2005	Mathiesen et al.
6,366,808	B1	4/2002	Schroeppelet al.	6,969,388	B2	11/2005	Goldman et al.
6,366,815	B1	4/2002	Haugland et al.	6,972,013	B1	12/2005	Zhang et al.
6,393,324	B2	5/2002	Gruzdowich et al.	6,978,174	B2	12/2005	Gelfand et al.
6,400,982	B2	6/2002	Sweeney et al.	6,985,774	B2	1/2006	Kieval et al.
6,405,079	B1	6/2002	Ansarinia	6,994,700	B2	2/2006	Elkins et al.
6,405,732	B1	6/2002	Edwards et al.	6,994,706	B2	2/2006	Chornenky et al.
6,413,255	B1	7/2002	Stern	7,004,911	B1	2/2006	Tu et al.
6,415,183	B1	7/2002	Scheiner et al.	7,054,685	B2	5/2006	Dimmer et al.
6,415,187	B1	7/2002	Kuzma et al.	7,063,679	B2	6/2006	Maguire et al.
6,438,423	B1	8/2002	Rezai et al.	7,081,114	B2	7/2006	Rashidi
6,442,424	B1	8/2002	Ben-Haim et al.	7,081,115	B2	7/2006	Taimisto
6,449,507	B1	9/2002	Hill et al.	7,083,614	B2	8/2006	Fjield et al.
6,450,942	B1	9/2002	Lapanashvili et al.	7,122,019	B1	10/2006	Kesten et al.
6,461,314	B1	10/2002	Pant et al.	7,155,284	B1	12/2006	Whitehurst et al.
6,464,687	B1	10/2002	Lshikawa et al.	7,162,303	B2	1/2007	Levin et al.
6,473,644	B1	10/2002	Terry, Jr. et al.	7,191,015	B2	3/2007	Lamson et al.
6,482,619	B1	11/2002	Rubinsky et al.	7,373,204	B2	5/2008	Gelfand et al.
6,488,679	B1	12/2002	Swanson et al.	7,444,183	B2	10/2008	Knudson et al.
6,506,189	B1	1/2003	Rittman, III et al.	7,617,005	B2	11/2009	Demarais et al.
6,508,774	B1	1/2003	Acker et al.	7,620,451	B2	11/2009	Demarais et al.
6,514,226	B1	2/2003	Levin et al.	7,647,115	B2	1/2010	Levin et al.
6,516,211	B1	2/2003	Acker et al.	7,653,438	B2	1/2010	Deem et al.
6,517,811	B2	2/2003	John et al.	7,717,948	B2	5/2010	Demarais et al.
6,522,926	B1	2/2003	Kieval et al.	7,756,583	B2	7/2010	Demarais et al.
6,522,932	B1	2/2003	Kuzma et al.	8,131,371	B2	3/2012	Demarais et al.
6,524,607	B1	2/2003	Goldenheim et al.	8,131,372	B2	3/2012	Levin et al.
6,534,081	B2	3/2003	Goldenheim et al.	8,145,317	B2	3/2012	Demarais et al.
6,536,949	B1	3/2003	Heuser	8,150,518	B2	4/2012	Levin et al.
6,542,781	B1	4/2003	Koblish et al.	8,150,519	B2	4/2012	Demarais et al.
6,564,096	B2	5/2003	Mest	8,150,520	B2	4/2012	Demarais et al.
6,571,127	B1	5/2003	Ben-Haim et al.	8,175,711	B2	5/2012	Demarais et al.
6,592,567	B1	7/2003	Levin et al.	8,350,846	B2	1/2013	Mejdreich et al.
6,599,256	B1	7/2003	Acker et al.	9,907,611	B2	3/2018	Levin et al.
6,600,954	B2	7/2003	Cohen et al.	2001/0044596	A1	11/2001	Jaafar
6,600,956	B2	7/2003	Maschino et al.	2002/0002329	A1	1/2002	Avitall
6,601,459	B1	8/2003	Jenni	2002/0026222	A1	2/2002	Schauerte et al.
6,605,084	B2	8/2003	Acker et al.	2002/0026228	A1	2/2002	Schauerte
6,613,045	B1	9/2003	Laufer et al.	2002/0032468	A1	3/2002	Hill et al.
6,615,071	B1	9/2003	Casscells, III et al.	2002/0038137	A1	3/2002	Stein
6,616,624	B1	9/2003	Kieval	2002/0040204	A1	4/2002	Dev et al.
6,620,151	B2	9/2003	Blischak et al.	2002/0045853	A1	4/2002	Dev et al.
6,622,041	B2	9/2003	Terry, Jr. et al.	2002/0065541	A1	5/2002	Fredricks et al.
6,622,731	B2	9/2003	Daniel et al.	2002/0072782	A1	6/2002	Osorio et al.
6,635,054	B2	10/2003	Fjield et al.	2002/0087208	A1	7/2002	Koblish et al.
6,654,636	B1	11/2003	Dev et al.	2002/0107515	A1	8/2002	Edwards et al.
6,666,845	B2	12/2003	Hooper et al.	2002/0107553	A1	8/2002	Hill et al.
6,669,655	B1	12/2003	Acker et al.	2002/0116030	A1	8/2002	Rezai
6,671,556	B2	12/2003	Osorio et al.	2002/0120304	A1	8/2002	Mest
6,672,312	B2	1/2004	Acker	2002/0139379	A1	10/2002	Edwards et al.
6,676,657	B2	1/2004	Wood	2002/0165532	A1	11/2002	Hill et al.
6,681,136	B2	1/2004	Schuler et al.	2002/0165586	A1	11/2002	Hill et al.
6,684,105	B2	1/2004	Cohen et al.	2002/0169413	A1	11/2002	Keren et al.
6,690,971	B2	2/2004	Schauerte et al.	2002/0177846	A1	11/2002	Mulier et al.
6,692,738	B2	2/2004	MacLaughlin et al.	2002/0183682	A1	12/2002	Darvish et al.
6,697,670	B2	2/2004	Chomenky et al.	2002/0183684	A1	12/2002	Dev et al.
6,711,444	B2	3/2004	Koblish	2002/0188325	A1	12/2002	Hill et al.
6,718,208	B2	4/2004	Hill et al.	2002/0198512	A1	12/2002	Seward
6,735,471	B2	5/2004	Hill et al.	2003/0004549	A1	1/2003	Hill et al.
6,738,663	B2	5/2004	Schroeppelet al.	2003/0009145	A1	1/2003	Struijker-Boudier et al.
6,749,598	B1	6/2004	Keren et al.	2003/0018367	A1	1/2003	DiLorenzo
6,752,805	B2	6/2004	Maguire et al.	2003/0040774	A1	2/2003	Terry et al.
6,786,904	B2	9/2004	Doscher et al.	2003/0045909	A1	3/2003	Gross et al.
6,795,728	B2	9/2004	Chornenky et al.	2003/0050635	A1	3/2003	Truckai et al.
6,845,267	B2	1/2005	Harrison et al.	2003/0050681	A1	3/2003	Pianca et al.
6,850,801	B2	2/2005	Kieval et al.	2003/0060848	A1	3/2003	Kieval et al.
6,862,479	B1	3/2005	Whitehurst et al.	2003/0060857	A1	3/2003	Perrson et al.
6,865,416	B2	3/2005	Dev et al.	2003/0060858	A1	3/2003	Kieval et al.
6,869,431	B2	3/2005	Maguire et al.	2003/0074039	A1	4/2003	Puskas
6,885,888	B2	4/2005	Rezai	2003/0100924	A1	5/2003	Foreman et al.
				2003/0120270	A1	6/2003	Acker
				2003/0125790	A1	7/2003	Fastovsky et al.
				2003/0150464	A1	8/2003	Casscells
				2003/0158584	A1	8/2003	Cates et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2003/0181897	A1	9/2003	Thomas et al.	2006/0036218	A1	2/2006	Goodson et al.
2003/0181963	A1	9/2003	Pellegrino et al.	2006/0041277	A1	2/2006	Deem et al.
2003/0195507	A1	10/2003	Stewart et al.	2006/0041283	A1	2/2006	Gelfand et al.
2003/0199747	A1	10/2003	Michlitsch et al.	2006/0067972	A1	3/2006	Kesten et al.
2003/0199767	A1	10/2003	Cespedes et al.	2006/0069323	A1	3/2006	Elkins et al.
2003/0199768	A1	10/2003	Cespedes et al.	2006/0074453	A1	4/2006	Kieval et al.
2003/0199806	A1	10/2003	Kieval	2006/0079859	A1	4/2006	Elkins et al.
2003/0199863	A1	10/2003	Swanson et al.	2006/0085046	A1	4/2006	Rezai et al.
2003/0204161	A1	10/2003	Ferek-Petric	2006/0085054	A1	4/2006	Zikorus et al.
2003/0216792	A1	11/2003	Levin et al.	2006/0089674	A1	4/2006	Walters et al.
2003/0220521	A1	11/2003	Reitz et al.	2006/0095029	A1	5/2006	Young et al.
2003/0233099	A1	12/2003	Danaek et al.	2006/0100667	A1	5/2006	Machado et al.
2003/0236443	A1	12/2003	Cespedes et al.	2006/0106429	A1	5/2006	Libbus et al.
2004/0010289	A1	1/2004	Biggs et al.	2006/0111754	A1	5/2006	Rezai et al.
2004/0010303	A1	1/2004	Bolea et al.	2006/0116720	A1	6/2006	Knoblich
2004/0019364	A1	1/2004	Kieval et al.	2006/0121016	A1	6/2006	Lee
2004/0019371	A1	1/2004	Jaafar et al.	2006/0121610	A1	6/2006	Rubinsky et al.
2004/0064090	A1	4/2004	Keren et al.	2006/0135998	A1	6/2006	Libbus et al.
2004/0064091	A1	4/2004	Keren et al.	2006/0136004	A1	6/2006	Cowan et al.
2004/0065615	A1	4/2004	Hooper et al.	2006/0149350	A1	7/2006	Patel et al.
2004/0073238	A1	4/2004	Makower	2006/0155344	A1	7/2006	Rezai et al.
2004/0082978	A1	4/2004	Harrison et al.	2006/0167437	A1	7/2006	Valencia
2004/0101523	A1	5/2004	Reitz et al.	2006/0167498	A1	7/2006	DiLorenzo
2004/0106953	A1	6/2004	Yomtov et al.	2006/0167499	A1	7/2006	Palti
2004/0111080	A1	6/2004	Harper et al.	2006/0189941	A1	8/2006	Seward et al.
2004/0127942	A1	7/2004	Yomtov et al.	2006/0189960	A1	8/2006	Kesten et al.
2004/0162590	A1	8/2004	Whitehurst et al.	2006/0190044	A1	8/2006	Libbus et al.
2004/0163655	A1	8/2004	Gelfand et al.	2006/0206149	A1	9/2006	Yun
2004/0167415	A1	8/2004	Gelfand et al.	2006/0206150	A1	9/2006	Demarais et al.
2004/0167509	A1	8/2004	Taimisto	2006/0212076	A1	9/2006	Demarais et al.
2004/0176699	A1	9/2004	Walker et al.	2006/0212078	A1	9/2006	Demarais et al.
2004/0176757	A1	9/2004	Sinelnikov et al.	2006/0229677	A1	10/2006	Moffitt et al.
2004/0193228	A1	9/2004	Gerber	2006/0235474	A1	10/2006	Demarais
2004/0215186	A1	10/2004	Cornelius et al.	2006/0265014	A1	11/2006	Demarais et al.
2004/0220511	A1	11/2004	Scott et al.	2006/0265015	A1	11/2006	Demarais et al.
2004/0243102	A1	12/2004	Berg et al.	2006/0271111	A1	11/2006	Demarais et al.
2004/0243206	A1	12/2004	Tadlock	2006/0276852	A1	12/2006	Demarais et al.
2004/0249416	A1	12/2004	Yun et al.	2007/0066957	A1	3/2007	Demarais et al.
2004/0254616	A1	12/2004	Rossing et al.	2007/0066972	A1	3/2007	Ormsby et al.
2005/0010263	A1	1/2005	Schauerte	2007/0083239	A1	4/2007	Demarais et al.
2005/0021092	A1	1/2005	Yun et al.	2007/0129760	A1	6/2007	Demarais et al.
2005/0038409	A1	2/2005	Segal et al.	2007/0129761	A1	6/2007	Demarais et al.
2005/0049542	A1	3/2005	Sigg et al.	2007/0135875	A1	6/2007	Demarais et al.
2005/0065562	A1	3/2005	Rezai	2007/0142864	A1	6/2007	Libbus et al.
2005/0065573	A1	3/2005	Rezai	2007/0156200	A1	7/2007	Kornet et al.
2005/0065574	A1	3/2005	Rezai	2007/0173899	A1	7/2007	Levin et al.
2005/0075681	A1	4/2005	Rezai et al.	2007/0208382	A1	9/2007	Yun
2005/0080409	A1	4/2005	Young et al.	2007/0265687	A1	11/2007	Deem et al.
2005/0080459	A1	4/2005	Jacobson et al.	2007/0282376	A1	12/2007	Shuros et al.
2005/0096710	A1	5/2005	Kieval	2007/0288070	A1	12/2007	Libbus et al.
2005/0153885	A1	7/2005	Yun et al.	2008/0004673	A1	1/2008	Rossing et al.
2005/0154418	A1	7/2005	Kieval et al.	2008/0015659	A1	1/2008	Zhang et al.
2005/0171523	A1	8/2005	Rubinsky et al.	2008/0039904	A1	2/2008	Bulkes et al.
2005/0171574	A1	8/2005	Rubinsky et al.	2008/0091255	A1	4/2008	Caparso et al.
2005/0171575	A1	8/2005	Dev et al.	2008/0140150	A1	6/2008	Zhou et al.
2005/0187579	A1	8/2005	Danek et al.	2008/0213331	A1	9/2008	Gelfand et al.
2005/0197624	A1	9/2005	Goodson et al.	2008/0255642	A1	10/2008	Zarins et al.
2005/0209548	A1	9/2005	Dev et al.	2008/0319513	A1	12/2008	Pu et al.
2005/0209642	A1	9/2005	Palti	2009/0024195	A1	1/2009	Rezai et al.
2005/0228460	A1	10/2005	Levin et al.	2009/0036948	A1	2/2009	Levin et al.
2005/0234523	A1	10/2005	Levin et al.	2009/0062873	A1	3/2009	Wu et al.
2005/0240126	A1	10/2005	Foley et al.	2009/0076409	A1	3/2009	Wu et al.
2005/0240173	A1	10/2005	Palti	2010/0010567	A1	1/2010	Deem et al.
2005/0240228	A1	10/2005	Palti	2010/0057150	A1	3/2010	Demarais et al.
2005/0240241	A1	10/2005	Yun et al.	2010/0137860	A1	6/2010	Demarais et al.
2005/0245882	A1	11/2005	Elkins et al.	2010/0137952	A1	6/2010	Demarais et al.
2005/0245892	A1	11/2005	Elkins et al.	2010/0168731	A1	7/2010	Wu et al.
2005/0251212	A1	11/2005	Kieval et al.	2010/0168739	A1	7/2010	Wu et al.
2005/0261672	A1	11/2005	Deem et al.	2010/0174282	A1	7/2010	Demarais et al.
2005/0267010	A1	12/2005	Goodson et al.	2010/0191112	A1	7/2010	Demarais et al.
2005/0282284	A1	12/2005	Rubinsky et al.	2010/0222851	A1	9/2010	Deem et al.
2006/0004417	A1	1/2006	Rossing et al.	2010/0222854	A1	9/2010	Demarais et al.
2006/0004430	A1	1/2006	Rossing et al.	2010/0241183	A1	9/2010	DiLorenzo
2006/0025821	A1	2/2006	Gelfand et al.	2010/0249773	A1	9/2010	Clark et al.
2006/0030814	A1	2/2006	Valencia et al.	2010/0268307	A1	10/2010	Demarais et al.
				2011/0060324	A1	3/2011	Wu et al.
				2011/0112400	A1	5/2011	Emery et al.
				2011/0137298	A1	6/2011	Nguyen et al.
				2011/0178570	A1	7/2011	Demarais

(56) References Cited

U.S. PATENT DOCUMENTS

2011/0200171	A1	8/2011	Beetel et al.
2011/0202098	A1	8/2011	Demarais et al.
2011/0257564	A1	10/2011	Demarais et al.
2011/0264011	A1	10/2011	Wu et al.
2011/0264075	A1	10/2011	Leung et al.
2012/0130345	A1	5/2012	Levin et al.
2013/0013024	A1	1/2013	Levin et al.
2013/0218029	A1	8/2013	Cholette et al.
2013/0253603	A1	9/2013	Levin et al.

FOREIGN PATENT DOCUMENTS

EP	2092957	8/2009
WO	WO-85/01213	3/1985
WO	WO-91/04725	4/1991
WO	WO-9220291	11/1992
WO	WO-93/02740	2/1993
WO	WO-93/07803	4/1993
WO	WO-94/00188	1/1994
WO	WO-9407446	4/1994
WO	WO-94/11057	5/1994
WO	WO-95/25472	9/1995
WO	WO-9531142	11/1995
WO	WO-95/33514	12/1995
WO	WO-96/00039	1/1996
WO	WO-96/04957	2/1996
WO	WO-96/11723	4/1996
WO	WO-97/13463	4/1997
WO	WO-97/13550	4/1997
WO	WO-1997036548	10/1997
WO	WO-97/49453	12/1997
WO	WO-98/37926	9/1998
WO	WO-98/42403	10/1998
WO	WO-98/43700	10/1998
WO	WO-98/43701	10/1998
WO	WO1998042403	10/1998
WO	WO-98/48888	11/1998
WO	WO-99/33407	7/1999
WO	WO-99/51286	10/1999
WO	WO-99/52424	10/1999
WO	WO-01/26729	4/2001
WO	WO-2001022897	4/2001
WO	WO-0170114	9/2001
WO	WO-02/09808	2/2002
WO	WO-02/26314	4/2002
WO	WO-02/053207	7/2002
WO	WO-02/070039	9/2002
WO	WO-02/070047	9/2002
WO	WO-02/085448	10/2002
WO	WO-2002085192	10/2002
WO	WO-03/018108	3/2003
WO	WO-2003022167	3/2003
WO	WO-03/028802	4/2003
WO	WO-03/063692	8/2003
WO	WO-03/071140	8/2003
WO	WO-03/076008	9/2003
WO	WO-03/082403	10/2003
WO	WO-2003/082080	10/2003
WO	WO-2004/026370	4/2004
WO	WO-2004/026371	4/2004
WO	WO-2004/026374	4/2004
WO	WO-2004/030718	4/2004
WO	WO-2004/032791	4/2004
WO	WO-2004/107965	12/2004
WO	WO-2005/014100	2/2005
WO	WO-2005/016165	2/2005
WO	WO-05/032646	4/2005
WO	WO-2005/032646	4/2005
WO	WO-2005030072	4/2005
WO	WO-2005041748	5/2005
WO	WO-2005/065284	7/2005
WO	WO-2005/084389	9/2005
WO	WO-2005/097256	10/2005
WO	WO-2005/110528	11/2005
WO	WO-2005/123183	12/2005

WO	WO-2006/007048	1/2006
WO	WO-2006/018528	2/2006
WO	WO-2006/022790	3/2006
WO	WO-2006/031899	3/2006
WO	WO-2006041847	4/2006
WO	WO-2006041881	4/2006
WO	WO-2006105121	10/2006
WO	WO-2007008954	1/2007
WO	WO-2007035537	3/2007
WO	WO-2007078997	7/2007
WO	WO-2007086965	8/2007
WO	WO-2007103879	9/2007
WO	WO-2007103881	9/2007
WO	WO-2007121309	10/2007
WO	WO-2007146834	12/2007
WO	WO-2008003058	1/2008
WO	WO-2008049084	4/2008
WO	WO-2008061150	5/2008
WO	WO-2008061152	5/2008
WO	WO-2008070413	6/2008
WO	WO-2010078175	7/2010

OTHER PUBLICATIONS

U.S. Appl. No. 95/002,209, filed Sep. 13, 2012, Levin et al.

U.S. Appl. No. 95/002,233, filed Sep. 13, 2012, Levin et al.

U.S. Appl. No. 95/002,243, filed Sep. 13, 2012, Levin et al.

U.S. Appl. No. 95/002,253, filed Sep. 13, 2012, Demarais et al.

U.S. Appl. No. 95/002,255, filed Sep. 13, 2012, Demarais et al.

U.S. Appl. No. 95/002,292, filed Sep. 14, 2012, Demarais et al.

U.S. Appl. No. 95/002,327, filed Sep. 14, 2012, Demarais et al.

U.S. Appl. No. 95/002,335, filed Sep. 14, 2012, Demarais et al.

U.S. Appl. No. 95/002,336, filed Sep. 14, 2012, Levin et al.

U.S. Appl. No. 95/002,356, filed Sep. 14, 2012, Demarais et al.

"2011 Edison Award Winners." Edison Awards: Honoring Innovations & Innovators, 2011, 6 pages, <http://www.edisonawards.com/BestNewProduct_2011.php>.

"2012 top 10 advances in heart disease and stroke research: American Heart Association/American Stroke Association Top 10 Research Report." American Heart Association, Dec. 17, 2012, 5 pages, <<http://newsroom.heart.org/news/2012-top-10-advances-in-heart-241901>>.

"Ardian(R) Receives 2010 EuroPCR Innovation Award and Demonstrates Further Durability of Renal Denervation Treatment for Hypertension." PR Newswire, Jun. 3, 2010, 2 pages, <<http://www.prnewswire.com/news-releases/ardianr-receives-2010-europcr-innovation-award-and-demonstrates-further-durability-of-renal-denervation-treatment-for-hypertension-95545014.html>>.

"Boston Scientific to Acquire Vessix Vascular, Inc.: Company to Strengthen Hypertension Program with Acquisition of Renal Denervation Technology." Boston Scientific: Advancing science for life—Investor Relations, Nov. 8, 2012, 2 pages, <<http://phx.corporate-ir.net/phoenix.zhtml?c=62272&p=irol-newsArticle&id=1756108>>.

"Cleveland Clinic Unveils Top 10 Medical Innovations for 2012: Experts Predict Ten Emerging Technologies that will Shape Health Care Next Year." Cleveland Clinic, Oct. 6, 2011, 2 pages, <http://my.clevelandclinic.org/media_relations/library/2011/2011-10-6-cleveland-clinic-unveils-top-10-medical-innovations-for-2012.aspx>.

"Does renal denervation represent a new treatment option for resistant hypertension?" Interventional News, Aug. 3, 2010, 2 pages, <<http://www.cxvascular.com/in-latest-news/interventional-news--latest-news/does-renal-denervation-represent-a-new-treatment-option-for-resistant-hypertension>>.

"Iberis—Renal Sympathetic Denervation System: Turning innovation into quality care." [Brochure], Terumo Europe N.V., 2013, Europe, 3 pages.

"Neurotech Reports Announces Winners of Gold Electrode Awards." Neurotech business report, 2009, 1 page, <<http://www.neurotechreports.com/pages/goldelectrodes09.html>>.

"Quick. Consistent. Controlled. OneShot renal Denervation System" [Brochure], Covidien: positive results for life, 2013, (n. 1.), 4 pages.

(56)

References Cited

OTHER PUBLICATIONS

- "Renal Denervation Technology of Vessix Vascular, Inc. been acquired by Boston Scientific Corporation (BSX) to pay up to \$425 Million." Vessix Vascular Pharmaceutical Intelligence: A blog specializing in Pharmaceutical Intelligence and Analytics, Nov. 8, 2012, 21 pages, <<http://pharmaceuticalintelligence.com/tag/vessix-vascular/>>.
- "The Edison Awards™" Edison Awards: Honoring Innovations & Innovators, 2013, 2 pages, <<http://www.edisonawards.com/Awards.php>>.
- "The Future of Renal denervation for the Treatment of Resistant Hypertension." St. Jude Medical, Inc., 2012, 12 pages.
- "Vessix Renal Denervation System: So Advanced It's Simple." [Brochure], Boston Scientific: Advancing science for life, 2013, 6 pages.
- Asbell, Penny, "Conductive Keratoplasty for the Correction of Hyperopia." *Tr Am Ophth Soc*, 2001, vol. 99, 10 pages.
- Badoer, Emilio, "Cardiac afferents play the dominant role in renal nerve inhibition elicited by volume expansion in the rabbit." *Am J Physiol Regul Integr Comp Physiol*, vol. 274, 1998, 7 pages.
- Bengel, Frank, "Serial Assessment of Sympathetic Reinnervation After Orthotopic Heart Transplantation: A longitudinal Study Using PET and C-11 Hydroxyephedrine." *Circulation*, vol. 99, 1999, 7 pages.
- Benito, F., et al. "Radiofrequency catheter ablation of accessory pathways in infants." *Heart*, 78:160-162 (1997).
- Bettmann, Michael, Carotid Stenting and Angioplasty: A Statement for Healthcare Professionals From the Councils on Cardiovascular Radiology, Stroke, Cardio-Thoracic and Vascular Surgery, Epidemiology and Prevention, and Clinical Cardiology, American Heart Association, *Circulation*, vol. 97, 1998, 4 pages.
- Bohm, Michael et al., "Rationale and design of a large registry on renal denervation: the Global Symplicity registry." *EuroIntervention*, vol. 9, 2013, 9 pages.
- Brosky, John, "EuroPCR 2013: CE-approved devices line up for renal denervation approval." *Medical Device Daily*, May 28, 2013, 3 pages, <http://www.medicaldevicedaily.com/servlet/com.accumedia.web.Dispatcher?next=bioWorldHeadlines_article&forceid=83002>.
- Davis, Mark et al., "Effectiveness of Renal Denervation Therapy for Resistant Hypertension." *Journal of the American College of Cardiology*, vol. 62, No. 3, 2013, 11 pages.
- Dubuc, M., et al., "Feasibility of cardiac cryoablation using a transvenous steerable electrode catheter." *J Interv Cardiac Electrophysiol*, 2:285-292 (1998).
- Final Office Action; U.S. Appl. No. 12/827,700; dated Feb. 5, 2013, 61 pages.
- Geisler, Benjamin et al., "Cost-Effectiveness and Clinical Effectiveness of Catheter-Based Renal Denervation for Resistant Hypertension." *Journal of the American College of Cardiology*, col. 60, No. 14, 2012, 7 pages.
- Gelfand, M., et al., "Treatment of renal failure and hypertension." U.S. Appl. No. 60/442,970, filed Jan. 29, 2003, 23 pages.
- Gertner, Jon, "Meet the Tech Duo That's Revitalizing the Medical Device Industry." *Fast Company*, Apr. 15, 2013, 6:00 AM, 17 pages, <<http://www.fastcompany.com/3007845/meet-tech-duo-thats-revitalizing-medical-device-industry>>.
- Golwyn, D. H., Jr., et al. "Percutaneous Transcatheter Renal Ablation with Absolute Ethanol for Uncontrolled Hypertension or Nephrotic Syndrome: Results in 11 Patients with End-Stage Renal Disease." *JVIR*, 8: 527-533 (1997).
- Hall, W. H., et al. "Combined embolization and percutaneous radiofrequency ablation of a solid renal tumor." *Am. J. Roentgenol*, 174: 1592-1594 (2000).
- Han, Y.-M, et al., "Renal artery embolization with diluted hot contrast medium: An experimental study." *J Vasc Interv Radiol*, 12: 862-868 (2001).
- Hansen, J. M., et al. "The transplanted human kidney does not achieve functional reinnervation." *Clin. Sci*, 87: 13-19 (1994).
- Hendee, W. R. et al. "Use of Animals in Biomedical Research: The Challenge and Response." *American Medical Association White Paper* (1988) 39 pages.
- Hering, Dagmara et al., "Chronic kidney disease: role of sympathetic nervous system activation and potential benefits of renal denervation." *EuroIntervention*, vol. 9, 2013, 9 pages.
- Imimdtanz, "Medtronic awarded industry's highest honor for renal denervation system." The official blog of Medtronic Australasia, Nov. 12, 2012, 2 pages, <<http://97waterlooad.wordpress.com/2012/11/12/medtronic-awarded-industrys-highest-honour-for-renal-denervation-system/>>.
- Kaiser, Chris, AHA Lists Year's Big Advances in CV Research, *medpage Today*, Dec. 18, 2012, 4 pages, <<http://www.medpagetoday.com/Cardiology/PCI/36509>>.
- Kompanowska, E., et al., "Early Effects of renal denervation in the anaesthetised rat: Natriuresis and increased cortical blood flow." *J Physiol*, 531. 2:527-534 (2001).
- Lee, S. J., et al. "Ultrasonic energy in endoscopic surgery." *Yonsei Med J*, 40:545-549 (1999).
- Lin, Dominik et al., "Renal denervation suppresses ventricular arrhythmias during acute ventricular ischemia in pigs." *Heart Rhythm*, vol. 0, No. 0, 2013, 6 pages.
- Lustgarten, D. L., et al., "Cryothermal ablation: Mechanism of tissue injury and current experience in the treatment of tachyarrhythmias." *Progr Cardiovasc Dis*, 41:481-498 (1999).
- Mabin, Tom et al., "First experience with endovascular ultrasound renal denervation for the treatment of resistant hypertension." *EuroIntervention*, vol. 8, 2012, 5 pages.
- Mahfoud, Felix et al., "Ambulatory Blood Pressure Changes after Renal Sympathetic Denervation in Patients with Resistant Hypertension." *Circulation*, 2013, 25 pages.
- Mahfoud, Felix et al., "Expert consensus document from the European Society of Cardiology on catheter-based renal denervation." *European Heart Journal*, 2013, 9 pages.
- Mahfoud, Felix et al., "Renal Hemodynamics and Renal Function After Catheter-Based Renal Sympathetic Denervation in Patients With Resistant Hypertension." *Hypertension*, 2012, 6 pages.
- Medical-Dictionary.com, Definition of "Animal Model," <http://medical-dictionary.com> (search "Animal Model"), 2005, 1 page.
- Medtronic, Inc., Annual Report (Form 10-K) (Jun. 28, 2011) 44 pages.
- Millard, F. C., et al, "Renal Embolization for ablation of function in renal failure and hypertension." *Postgraduate Medical Journal*, 65, 729-734, (1989).
- Oliveira, V., et al., "Renal denervation normalizes pressure and baroreceptor reflex in high renin hypertension in conscious rats." *Hypertension*, 19:II-17-II-21 (1992).
- Ong, K. L., et al. "Prevalence, Awareness, Treatment, and Control of Hypertension Among United States Adults 1999-2004." *Hypertension*, 49: 69-75 (2007) (originally published online Dec. 11, 2006).
- Orniston, John et al., "First-in-human use of the OneShot™ renal denervation system from Covidien." *EuroIntervention*, vol. 8, 2013, 4 pages.
- Orniston, John et al., "Renal denervation for resistant hypertension using an irrigated radiofrequency balloon: 12-month results from the Renal Hypertension Ablation System (RHAS) trial." *EuroIntervention*, vol. 9, 2013, 5 pages.
- Pedersen, Amanda, "TCT 2012: Renal denervation device makers play show and tell." *Medical Device Daily*, Oct. 26, 2012, 2 pages, <http://www.medicaldevicedaily.com/servlet/com.accumedia.web.Dispatcher?next=bioWorldHeadlines_article&forceid=80880>.
- Peet, M., "Hypertension and its Surgical Treatment by bilateral supradiaphragmatic splanchnicectomy" *Am J Surgery* (1948) pp. 48-68.
- Renal Denervation (RDN), Symplicity RDN System Common Q&A (2011), 4 pages, <http://www.medtronic.com/rdn/mediakit/RDN%20FAQ.pdf>.
- Schlaich, Markus et al., "Renal Denervation in Human Hypertension: Mechanisms, Current Findings, and Future Prospects." *Curr Hypertens Rep*, vol. 14, 2012, 7 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Schmid, Axel et al., "Does Renal Artery Supply Indicate Treatment Success of Renal Denervation." *Cardiovasc Intervent Radiol*, vol. 36, 2013, 5 pages.
- Schmieder, Roland E. et al., "Updated ESH position paper on interventional therapy of resistant hypertension." *EuroIntervention*, vol. 9, 2013, 9 pages.
- Sievert, Horst, "Novelty Award EuroPCR 2010." *Euro PCR*, 2010, 15 pages.
- Stella, A., et al., "Effects of reversible renal denervation on haemodynamic and excretory functions on the ipsilateral and contralateral kidney in the cat." *Hypertension*, 4:181-188 (1986).
- Stouffer, G. A. et al., "Catheter-based renal denervation in the treatment of resistant hypertension." *Journal of Molecular and Cellular Cardiology*, vol. 62, 2013, 6 pages.
- Swartz, J. F., et al., "Radiofrequency endocardial catheter ablation of accessory atrioventricular pathway atrial insertion sites." *Circulation*, 87: 487-499 (1993).
- Uchida, F., et al., "Effect of radiofrequency catheter ablation on parasympathetic denervation: A comparison of three different ablation sites." *PACE*, 21:2517-2521 (1998).
- Verloop, W. L. et al., "Renal denervation: a new treatment option in resistant arterial hypertension." *Neth Heart J*, Nov. 30, 2012, 6 pages, <<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3547427/>>.
- Weinstock, M., et al., "Renal denervation prevents sodium retention and hypertension in salt sensitive rabbits with genetic baroreflex impairment." *Clinical Science*, 90:287-293 (1996).
- Wilcox, Josiah N., Scientific Basis Behind Renal Denervation for the Control of Hypertension. ICI 2012, Dec. 5-6, 2012, 38 pages.
- Worthley, Stephen et al., "Safety and efficacy of a multi-electrode renal sympathetic denervation system in resistant hypertension: the EnligHTN I trial." *European Heart Journal*, vol. 34, 2013, 9 pages.
- Worthley, Stephen, "The St. Jude Renal Denervation System Technology and Clinical Review." The University of Adelaide Australia, 2012, 24 pages.
- Zuern, Christine S., "Impaired Cardiac Baroreflex Sensitivity Predicts Response to Renal Sympathetic Denervation in Patients with Resistant Hypertension." *Journal of the American College of Cardiology*, 2013, doi: 10.1016/j.jacc.2013.07.046, 24 pages.
- Miller, Reed, "Finding a Future for Renal Denervation With Better Controlled Trials." *Pharma & Medtech Business Intelligence*, Article # 01141006003, Oct. 6, 2014, 4 pages.
- Papademetriou, Vasilios, "Renal Denervation and Symplicity HTN-3: "Dubium Sapientiae Initium" (Doubt Is the Beginning of Wisdom)," *Circulation Research*, 2014; 115: 211-214.
- Papademetriou, Vasilios et al., "Renal Nerve Ablation for Resistant Hypertension: How Did We Get Here, Present Status, and Future Directions." *Circulation*. 2014; 129: 1440-1450.
- Papademetriou, Vasilios et al., "Catheter-Based Renal Denervation for Resistant Hypertension: 12-Month Results of the EnligHTN I First-in-Human Study Using a Multielectrode Ablation System." *Hypertension*. 2014; 64: 565-572.
- Doumas, Michael et al., "Renal Nerve Ablation for Resistant Hypertension: The Dust Has Not Yet Settled." *The Journal of Clinical Hypertension*. 2014; vol. 16, No. 6, 2 pages.
- Messerli, Franz H. et al., "Renal Denervation for Resistant Hypertension: Dead or Alive?" *Healio: Cardiology today's Intervention*, May/June. 2014, 2 pages.
- 2003 European Society of Hypertension—European Society of Cardiology guidelines for the management of arterial hypertension, Guidelines Committee, *Journal of Hypertension* 2003, vol. 21, No. 6, pp. 1011-1053.
- Aars, H. and S. Akre, Reflex Changes in Sympathetic Activity and Arterial Blood Pressure Evoked by Afferent Stimulation of the Renal Nerve, Feb. 26, 1999, *Acta physiol. Scand.*, vol. 78, 1970, pp. 184-188.
- Abramov, G.S. et al., Alteration in sensory nerve function following electrical shock, *Burns* vol. 22, No. 8, 1996 Elsevier Science Ltd., pp. 602-606.
- Achar, Suraj, M.D., and Suriti Kundu, M.D., Principles of Office Anesthesia: Part I. Infiltrative Anesthesia, Office Procedures, *American Family Physician*, Jul. 1, 2002, vol. 66, No. 1, pp. 91-94.
- Advanced Neuromodulation Systems' Comparison Chart, Dec. 16, 2008, pp. 1.
- Advances in the role of the sympathetic nervous system in cardiovascular medicine, 2001 SNS Report, No. 3, Springer, Published with an educational grant from Servier, pp. 1-8.
- Aggarwal, A. et al., Regional sympathetic effects of low-dose clonidine in heart failure. *Hypertension*. 2003;41:553-7.
- Agnew, William F. et al., Evolution and Resolution of Stimulation-Induced Axonal Injury in Peripheral Nerve, May 21, 1999, *Muscle & Nerve*, vol. 22, Oct. 1999, John Wiley & Sons, Inc. 1999, pp. 1393-1402.
- Ahadian, Farshad M., M.D., Pulsed Radiofrequency Neurotomy: Advances in Pain Medicine, Current Pain and Headache Reports 2004, vol. 8, 2004 Current Science Inc., pp. 34-40.
- Alexander, B.T. et al., Renal denervation abolishes hypertension in low-birth-weight offspring from pregnant rats with reduced uterine perfusion, *Hypertension*, 2005; 45 (part 2): pp. 754-758.
- Alford, J. Winslow, M.D. and Paul D. Fadale, M.D., Evaluation of Postoperative Bupivacaine Infusion for Pain Management After Anterior Cruciate Ligament Reconstruction, *The Journal of Arthroscopic and Related Surgery*, vol. 19, No. 8, Oct. 2003 Arthroscopy Association of North America, pp. 855-861.
- Allen, E.V., Sympathectomy for essential hypertension, *Circulation*, 1952, 6:131-140.
- Amersham Health. Hypaque-Cysto, 2003, 6 pages.
- Andrews, B.T. et al., The use of surgical sympathectomy in the treatment of chronic renal pain. *Br J Urol*. 1997; 80: 6-10.
- Antman, Elliott M. and Eugene Braunwald, Chapter 37—Acute Myocardial Infarction, Heart Disease—A Textbook of Cardiovascular Medicine, 5th Edition, vol. 2, 1997, Edited by Eugene Braunwald, pp. 1184-1288.
- Archer, Steffen et al., Cell Reactions to Dielectrophoretic Manipulation, Mar. 1, 1999, Biochemical and Biophysical Research Communications, 1999 Academic Press, pp. 687-698.
- Arentz, T. et al., Incidence of pulmonary vein stenosis 2 years after radiofrequency catheter ablation of refractory atrial fibrillation. *European Heart Journal*. 2003. 24; pp. 963-969.
- Arias, M.D., Manuel J., Percutaneous Radio-Frequency Thermocoagulation with Low Temperature in the Treatment of Essential Glossopharyngeal Neuralgia, *Surg. Neurol.* 1986, vol. 25, 1986 Elsevier Science Publishing Co., Inc., pp. 94-96.
- Aronofsky, David H., D.D.S., Reduction of dental postsurgical symptoms using nonthermal pulsed high-peak-power electromagnetic energy, *Oral Surg.*, Nov. 1971, vol. 32, No. 5, pp. 688-696.
- Aspelin, Peter, M.D., Ph.D. et al., Nephrotoxic Effects in High-Risk Patients Undergoing Angiography, Feb. 6, 2003, *New England Journal of Medicine* 2003, vol. 348, No. 6, 2003 Massachusetts Medical Society, pp. 491-499.
- Atrial Fibrillation Heart and Vascular Health on Yahoo! Health. 2 pgs. <URL: http://health.yahoo.com/topic/heart/overview/article/healthwise/hw160872;_ylt=AiBT43EY74HQ7ft3jAb4C.sPu7cF> Feb. 21, 2006.
- Augustyniak, Robert A. et al., Sympathetic Overactivity as a Cause of Hypertension in Chronic Renal Failure, Aug. 14, 2001, *Journal of Hypertension* 2002, vol. 20, 2002 Lippincott Williams & Wilkins, pp. 3-9.
- Awvrad, Ziad M., FRCS and Bashir A. Atiyat, GBA, JBA, Pain relief using continuous bupivacaine infusion in the paravertebral space after loin incision, May 15, 2004, *Saudi Med J* 2004, vol. 25 (10), pp. 1369-1373.
- Badyal, D. K., H. Lata and A.P. Dadhich, Animal Models of Hypertension and Effect of Drugs, Aug. 19, 2003, *Indian Journal of Pharmacology* 2003, vol. 35, pp. 349-362.
- Baker, Carol E. et al., Effect of pH of Bupivacaine on Duration of Repeated Sciatic Nerve Blocks in the Albino Rat, *Anesth Analg*, 1991, vol. 72, The International Anesthesia Research Society 1991, pp. 773-778.
- Balazs, Tibor, Development of Tissue Resistance to Toxic Effects of Chemicals, Jan. 26, 1974, *Toxicology*, 2 (1974), Elsevier/North-Holland, Amsterdam, pp. 247-255.

(56)

References Cited

OTHER PUBLICATIONS

- Barajas, L. Innervation of the renal cortex. *Fex Proc.* 1978;37:1192-201.
- Barrett, Carolyn J. et al., Long-term control of renal blood flow: what is the role of the renal nerves?, Jan. 4, 2001, *Am J Physiol Regulatory Integrative Comp Physiol* 280, 2001, The American Physiological Society 2001, pp. R1534-R1545.
- Barrett, Carolyn J. et al., What Sets the Long-Term Level of Renal Sympathetic Nerve Activity, May 12, 2003, *Integrative Physiology, Circ Res.* 2003, vol. 92, 2003 American Heart Association, pp. 1330-1336.
- Bassett, C. Andrew L. et al., Augmentation of Bone Repair by Inductively Coupled Electromagnetic Fields, May 3, 1974, *Science*, vol. 184, pp. 575-577.
- Bassett, C. Andrew L., Fundamental and Practical Aspects of Therapeutic Uses of Pulsed Electromagnetic Fields (PEMFs), *Critical Reviews in Biomedical Engineering*, vol. 17, Issue 5, 1989, pp. 451-514.
- Beebe, Stephen J. et al., Nanosecond pulsed electric fields modulate cell function through intracellular signal transduction mechanisms, Apr. 8, 2004, *Physiol. Meas.* 25, 2004, IOP Publishing Ltd. 2004, pp. 1077-1093.
- Beebe, Stephen J., et al., Nanosecond Pulsed Electric Field (nsPEF) Effects on Cells and Tissues: Apoptosis Induction and Tumor Growth Inhibition, Oct. 11, 2001, *IEEE Transactions on Plasma Science*, vol. 30, No. 1, Feb. 2002, IEEE 2002, pp. 286-292.
- Bello-Reuss, E. et al., Acute unilateral renal denervation in rats with extracellular volume expansion, *Departments of Medicine and Physiology, University of North Carolina School of Medicine.* F26-F32 Jul. 1975.
- Bello-Reuss, E. et al., Effect of renal sympathetic nerve stimulation on proximal water and sodium reabsorption, *J Clin Invest*, 1976;57:1104-1107.
- Bello-Reuss, E. et al., Effects of Acute Unilateral Renal Denervation in the Rat, *J Clin Invest*, 1975;56:208-217.
- Berde, C. et al., *Local Anesthetics, Anesthesia*, Chapter 13, 5th addition, Churchill-Livingston, Philadelphia 2000, pp. 491-521.
- Bhadra, Niloy and Kevin L. Kilgore, Direct Current Electrical Conduction Block of Peripheral Nerve, Feb. 25, 2004, *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, vol. 12, No. 3, Sep. 2004, pp. 313-324.
- Bhandari, A. and Elias, M., Loin pain hematuria syndrome: Pain control with RFA to the Splanchnic plexus, *The Pain Clinic*, 2000, vol. 12, No. 4, pp. 323-327.
- Bhatt, Deepak L. et al., Rhabdomyolysis Due to Pulsed Electric Fields, May 11, 1989, *Plastic and Reconstructive Surgery* Jul. 1990, pp. 1-11.
- Bichet, D., et al., Renal intracortical blood flow and renin secretion after denervation by 6-hydroxydopamine. *Can J Physiol Pharmacol.* 1982;60:184-92.
- Bigler, D. et al., Tachyphylaxis during postoperative epidural analgesia—new insights, Apr. 15, 1987, Letter to the Editor, *Acta Anaesthesiol Scand.* 1987, vol. 31, pp. 664-665.
- Binder, Allan et al., Pulsed Electromagnetic Field Therapy of Persistent Rotator Cuff Tendinitis, *The Lancet*, Saturday Mar. 31, 1984, The Lancet Ltd., pp. 695-698.
- Black, M.D., Henry R., Resistant Hypertension 2004, presentation at Rush University Medical Center, Jul. 15, 2004, 40 pages.
- Blad, B., et al., An Electrical Impedance index to Assess Electroporation in Tissue, *Tissue and Organ (Therapy)*, 2001, Oslo, www.bl.uk <http://www.bl.uk> British Library, pp. 31-34.
- Blair, M. L. et al, Sympathetic activation cannot fully account for increased plasma renin levels during water deprivation, Sep. 23, 1996, *Am. J. Physiol.*, vol. 272, 1997, The American Physiological Society 1997, pp. R1197-R1203.
- Blomberg, S.G., M.D., PhD, Long-Term Home Self-Treatment with High Thoracic Epidural Anesthesia in Patients with Severe Coronary Artery Disease, Mar. 29, 1994, *Anesth Analg* 1994, vol. 79, 1994 International Anesthesia Research Society, pp. 413-421.
- Boehmer, J.P., Resynchronization Therapy for Chronic CHF: Indications, Devices and Outcomes. Penn State College of Medicine: Penn State Heart and Vascular Institute. Transcatheter Cardiovascular Therapeutics 2005, 31 slides.
- Bourge, R.C., Heart Failure Monitoring Devices: Rationale and Status 28 pages, Feb. 2001.
- Braunwald, E., Heart Disease, A Textbook of Cardiovascular Medicine, 5th Ed., vol. 2, 1997, pp. 480-481, 824-825, 1184-1288 and 1923-1925, W.B. Saunders Company.
- Bravo, E.L., et al., Renal denervation for resistant hypertension, *American Journal of Kidney Diseases*, 2009, 3 pgs.
- Bunch, Jared T. et al. Mechanisms of Phrenic Nerve Injury During Radiofrequency Ablation at the Pulmonary Vein Orifice. *Journal of Cardiovascular Electrophysiology*. vol. 16, No. 12. pp. 1318-1325. Dec. 2005.
- Burkoff, D., Interventional Device-Based Therapy for CHF Will Redefine Current Treatment Paradigms. Columbia University. 2004. 32 slides.
- Burns, J. et al., Relationship between central sympathetic drive and magnetic resonance imaging-determined left ventricular mass in essential hypertension. *Circulation*. 2007;115:1999-2005.
- Cahana, A. et al., Acute Differential Modulation of Synaptic Transmission and Cell Survival During Exposure to Pulsed and Continuous Radiofrequency Energy, May 2003, *The Journal of Pain*, vol. 4, No. 4, © 2003 by the American Pain Society, pp. 197-202.
- Cahana, Alex, M.D., Pulsed Radiofrequency: A Neurobiologic and Clinical Reality, May 17, 2005, *Anesthesiology* 2005, vol. 103, No. 6, Dec. 2005, 2005 American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc., p. 1311.
- Calaresu, F.R. et al., Haemodynamic Responses and Renin Release During Stimulation of Afferent Renal Nerves in the Cat, Aug. 12, 1975, *J. Physiol.* 1976, vol. 255, pp. 687-700.
- Cameron, Tracy. Micromodular Implants to Provide Electrical Stimulation of Paralyzed Muscles and Limbs. *IEEE Transactions on Biomedical Engineering*, vol. 44, No. 9, Sep. 1997, pp. 781-790.
- Campese, V.M. et al., Renal afferent denervation prevents hypertension in rats with chronic renal failure. *Hypertension*. 1995;25:878-82.
- Campese, V.M. et al., Renal Afferent Denervation Prevents the Progression of Renal Disease in the Renal Ablation Model of Chronic Renal Failure in the Rat, *Am J Kidney Dis.* 1995;26:861-5.
- Campese, V.M., a new model of neurogenic hypertension caused by renal injury: pathophysiology and therapeutic implications, *Clin Exp Nephrol* (2003) 7: 167-171, Japanese Society of Nephrology 2003.
- Campese, V.M., Neurogenic factors and hypertension in chronic renal failure, *Journal of Nephrology*, vol. 10, No. 4, 1997, Societa Italiana di Nefrologia, pp. 184-187.
- Campese, V.M., Neurogenic factors and hypertension in renal disease. *Kidney Int.* 2000;57 Suppl 75:S2-3.
- Canbaz, S. et al., Electrophysiological evaluation of phrenic nerve injury during cardiac surgery—a prospective, controlled clinical study. *BioMed Central.* 5 pgs. 2004.
- Cardiac Glycosides, Heart Disease—A Textbook of Cardiovascular Medicine vol. 2, Edited by Eugene Braunwald, 5th Edition, 1997 WB Saunders Company, pp. 480-481.
- Carls, G. et al., Electrical and magnetic stimulation of the intercostal nerves: a comparative study, *Electromyogr. clin. Neurophysiol.* 1997, vol. 37, pp. 509-512.
- Carlson, Scott H. and J. Michael Wyss, e-Hypertension—Opening New Vistas, Introductory Commentary, *Hypertension* 2000, vol. 35, American Heart Association, Inc. 2000, p. 538.
- Carson, P., Device-based Treatment for Chronic Heart Failure: Electrical Modulation of Myocardial Contractility. *Transcatheter Cardiovascular Therapeutics* 2005, 21 slides.
- Chang, Donald C., Cell poration and cell fusion using an oscillating electric field, *Biophysical Journal*, vol. 56, Oct. 1989, Biophysical Society, pp. 641-652.
- Chen, S.A. et al., Initiation of atrial fibrillation by ectopic beats originating from the pulmonary veins: electrophysiological characteristics, pharmacological responses, and effects of radiofrequency ablation, *Circulation*, 1999, 100:1879-1886.

(56)

References Cited

OTHER PUBLICATIONS

- Chin, J.L. et al., Renal autotransplantation for the loin pain-hematuria syndrome: long term follow up of 26 cases, *J Urol*, 1998, vol. 160, pp. 1232-1236.
- Chiou, C.W. et al., Efferent Vagal Innervation of the Canine Atria and Sinus and Atrioventricular Nodes. *Circulation*. Jun. 1997. 95(11):2573-2584. Abstract only. 2 pgs.
- Chobanian, Aram V. et al., Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure, Nov. 6, 2003, *Hypertension* 2003, vol. 42, 2003 American Heart Association, Inc., pp. 1206-1252.
- Clinical Trials in Hypertension and Renal Diseases, Slide Source, www.hypertensiononline.org, 33 pages Aug. 13, 2001.
- Conradi, E. and Ines Helen Pages, Effects of Continuous and Pulsed Microwave Irradiation on Distribution of Heat in the Gluteal Region of Minipigs, *Scand J Rehab Med*, vol. 21, 1989, pp. 59-62.
- Converse, R.L., Jr. et al., Sympathetic Overactivity in Patients with Chronic Renal Failure, *N Engl J Med*. Dec. 31, 1992, vol. 327 (27), pp. 1912-1918.
- Cosman, E.R., Jr. et al., Electric and Thermal Field Effects in Tissue Around Radiofrequency Electrodes, *Pain Medicine*, vol. 6, No. 6, 2005, American Academy of Pain Medicine, pp. 405-424.
- Cosman, E.R., Ph.D. Comment on the History of the Pulsed Radiofrequency Technique for Pain Therapy, *Anesthesiology* Dec. 2005, vol. 103, No. 6, 2005 American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc., p. 1312.
- Crawford, William H. et al., Pulsed Radio Frequency Therapy of Experimentally Induced Arthritis in Ponies, Dec. 18, 1989, *Can. J. Vet. Res.* 1991, vol. 55, pp. 76-85.
- Curtis, J.J. et al., Surgical therapy for persistent hypertension after renal transplantation, *Transplantation*, 1981, 31(2):125-128.
- Dahm, Peter et al., Efficacy and Technical Complications of Long-Term Continuous Intraspinal Infusions of Opioid and/or Bupivacaine in Refractory Nonmalignant Pain . . . , Oct. 6, 1997, *The Clinical Journal of Pain*, vol. 14, No. 1, 1998, Lippincott-Raven Publishers 1998, pp. 4-16.
- Dahm, Peter O. et al., Long-Term Intrathecal Infusion of Opioid and/or Bupivacaine in the Prophylaxis and Treatment of Phantom Limb Pain, *Neuromodulation*, vol. 1, No. 3, 1998, International Neuromodulation Society 1998, pp. 111-128.
- Dang, Nicholas C. et al., A Novel Approach to Increase Total Urine Output in Heart Failure: Renal Nerve Blockade, ACC 2005 poster; 1 page.
- Daniel, Alan and Honig, Carl R. Does Histamine Influence Vaso-dilation Caused by Prolonged Arterial Occlusion or Heavy Exercise? *The Journal of Pharmacology and Experimental Therapeutics*. vol. 215 No. 2. Aug. 21, 1980. pp. 533-538.
- Davalos, R. et al., Electrical Impedance Tomography for Imaging Tissue Electroporation, Jul. 25, 2003, *IEEE Transactions on Bio-medical Engineering*, vol. 51, No. 5, May 2004, IEEE 2004, pp. 761-767.
- Davalos, R.V. et al., Tissue Ablation with Irreversible Electroporation, Sep. 7, 2004, *Annals of Biomedical Engineering*, Feb. 2005, vol. 33, No. 2, 2005 Biomedical Engineering Society, pp. 223-231.
- De Leeuw, Peter W. et al., Renal Vascular Tachyphylaxis to Angiotensin II: Specificity of the Response for Angiotensin, Dec. 28, 1981, *Life Sciences*, vol. 30, 1982 Pergamon Press Ltd., pp. 813-819.
- Deng, Jingdong et al., The Effects of Intense Submicrosecond Electrical Pulses on Cells, Nov. 26, 2002, *Biophysical Journal*, vol. 84, Apr. 2003, Biophysical Society 2003, pp. 2709-2714.
- Denton, Kate M. et al., Differential Neural Control of Glomerular Ultrafiltration, Jan. 30, 2004, *Proceedings of the Australian Physiological and Pharmacological Society Symposium: Hormonal, Metabolic and Neural Control of the Kidney, Clinical and Experimental Pharmacology and Physiology* (2004) 31, pp. 380-386.
- Dev, Nagendu B., Ph.D. et al., Intravascular Electroporation Markedly Attenuates Neointima Formation After Balloon Injury of the Carotid Artery in the Rat, *Journal of Interventional Cardiology*, vol. 13, No. 5, 2000, pp. 331-338.
- Dev, Nagendu B., Ph.D. et al., Sustained Local Delivery of Heparin to the Rabbit Arterial Wall with an Electroporation Catheter, May 5, 1998, *Catheterization and Cardiovascular Diagnosis*, vol. 45, 1998, Wiley-Liss, Inc. 1998, pp. 337-345.
- Devereaux, R.B. et al., Regression of Hypertensive Left Ventricular Hypertrophy by Losartan Compared With Atenolol: The Losartan Intervention for Endpoint Reduction in Hypertension (LIFE) Trial, *Circulation*, 2004, vol. 110, pp. 1456-1462.
- Dibona, Gerald F. and Linda L. Sawin, Role of renal nerves in sodium retention of cirrhosis and congestive heart failure, Sep. 27, 1990, *Am. J. Physiol.* 1991, vol. 260, 1991 the American Physiological Society, pp. R298-R305.
- Dibona, Gerald F. and Susan Y. Jones, Dynamic Analysis of Renal Nerve Activity Responses to Baroreceptor Denervation in Hypertensive Rats, Sep. 19, 2000, *Hypertension* Apr. 2001, American Heart Association, Inc. 2001, pp. 1153-1163.
- Dibona, Gerald F. and Ulla C. Kopp, Neural Control of Renal Function, *Physiological Reviews*, vol. 77, No. 1, Jan. 1997, the American Physiological Society 1997, pp. 75-197.
- Dibona, Gerald F. and Ulla C. Kopp, Role of the Renal Sympathetic Nerves in Pathophysiological States, *Neural Control of Renal Function*, vol. 77, pp. 142-197 Jan. 1997.
- Dibona, Gerald F., Functionally Specific Renal Sympathetic Nerve Fibers: Role in Cardiovascular Regulation, Mar. 6, 2001, *American Journal of Hypertension*, 2001, vol. 14, 2001 American Journal of Hypertension, Ltd. Published by Elsevier Science Inc., pp. 163S-170S.
- Dibona, Gerald F., L.L. Sawin, Effect of renal nerve stimulation on NaCl and H₂O transport in Henle's loop of the rat, 1982, *American Physiological Society*, F576-F580, 5 pgs.
- Dibona, Gerald F., Nervous Kidney—Interaction Between Renal Sympathetic Nerves and the Renin-Angiotensin System in the Control of Renal Function, Jun. 21, 2000, *Hypertension* 2000, vol. 36, 2000 American Heart Association, Inc., pp. 1083-1088.
- Dibona, Gerald F., Neural Control of the Kidney—Past, Present and Future, Nov. 4, 2002, *Novartis Lecture, Hypertension* 2003, 41 part 2, 2002 American Heart Association, Inc., pp. 621-624.
- DiBona, Gerald F., Neural Control of the Kidney: Functionally Specific Renal Sympathetic Nerve Fibers, *Starling Lecture, Am J Physiol Regulatory Integrative Comp Physiol*, 2000, 279, 2000 The American Physiological Society, pp. R1517-R1524.
- Dibona, Gerald F., Peripheral and Central Interactions between the Renin-Angiotensin System and the Renal Sympathetic Nerves in Control of Renal Function, *Annals New York Academy of Sciences*, pp. 395-406 Jan. 25, 2006.
- Dibona, Gerald F., Renal Innervation and Denervation: Lessons from Renal Transplantation Reconsidered, *Artificial Organs*, vol. 11, No. 6, Raven Press, Ltd., 1987 International Society for Artificial Organs, pp. 457-462.
- Dibona, Gerald F., Sympathetic Nervous System and the Kidney in Hypertension, *Current Opinion in Nephrology and Hypertension* 2002, vol. 11, 2002 Lippincott Williams & Wilkins, pp. 197-200.
- Dibona, Gerald F., The Sympathetic Nervous System and Hypertension, Dec. 4, 2003, *Hypertension Highlights, Hypertension* Feb. 2004, vol. 43, 2004 American Heart Association, Inc., pp. 147-150.
- Dibona, Gerald, L.L. Sawin, Effect of renal denervation on dynamic autoregulation of renal blood flow, Feb. 12, 2004, *AmJ Physiol Renal Physiol* 286, pp. F1209-1218.
- Dong, Jun et al. Incidence and Predictors of Pulmonary Vein Stenosis Following Catheter Ablation of Atrial Fibrillation Using the Anatomic Pulmonary Vein Ablation Approach: Results from Paired Magnetic Resonance Imaging. *Journal of Cardiovascular Electrophysiology*. vol. 16, No. 8, Aug. 2005. pp. 845-852.
- Dorros, Gerald, M.D., Renal Artery Stenting State of the Art, presentation, TCT, Washington D.C., Sep. 2003, 27 pages.
- Dueck, Ron, M.D., Noninvasive Cardiac Output Monitoring, *The Cardiopulmonary and Critical Care Journal*, Chest, vol. 120, sec. 2, Aug. 2001, American College of Chest Physicians 2005, pp. 339-341, 5 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Dunn, Matthew D. et al., Laparoscopic Nephrectomy in Patients With End-Stage Renal Disease and Autosomal Dominant Polycystic Kidney Disease, Oct. 25, 1999, American Journal of Kidney Diseases, vol. 35, No. 4 Apr. 2000, National Kidney Foundation, Inc. 2000, pp. 720-725.
- Durand, D.M., Electric Field Effects in Hyperexcitable Neural Tissue: A Review, Radiation Protection Dosimetry, vol. 106, No. 4, 2003 Nuclear Technology Publishing, pp. 325-331.
- Effects of Renal Failure on the Cardiovascular System, 5th Edition Heart Disease, A Textbook of Cardiovascular Medicine, vol. 2, Edited by Eugene Braunwald, 1997, W.B. Saunders Company, pp. 1923-1925.
- Electrical Stimulation for the Treatment of Chronic Wounds, Radiation Protection Standard, Maximum Exposure Levels to Radiofrequency Fields—3 KHz to 300 GHz, Radiation Protection Series No. 3, Australian Radiation Protection and Nuclear Safety Agency, Apr. 1996, 322 pgs.
- Electroporation (Electroporation), Cyto Pulse Sciences, Inc., <http://www.cytopulse.com/electroporation.html> (last accessed Mar. 3, 2005), 3 pgs.
- Electroporation based Technologies and Treatments, ESPE Newsletter No. 6, QLK 02002-2003, Jan. 2005, www.cliniporator.com, 4 pgs.
- End-stage renal disease payment policies in traditional Medicare, Chapter 8, Report to the Congress: Medicare Payment Policy, Mar. 2001, Medpac, pp. 123-138.
- Epidemiology of Renal Disease in Hypertension, slide presentation by hypertensiononline.org, 21 pages Mar. 30, 2001.
- Erdine, Serap and Alev Arat-Ozkan, Resistant Hypertension, European Society of Hypertension Scientific Newsletter: Update on Hypertension Management 2003, vol. 4, No. 15, 2 pages.
- Esler, M. et al., Mechanism of elevated plasma noradrenaline in the course of essential hypertension. J Cardiovasc Pharmacol. 1986;8:S39-43.
- Esler, M. et al., Noradrenaline release and the pathophysiology of primary human hypertension. Am J Hypertens. 1989; 2:140S-146S.
- Esler, M. et al., Sympathetic nerve biology in essential hypertension, Clin and Exp Pharmacology and Physiology, 2001, 28:986-989.
- European Examination Report; European Patent Application No. 07799148.7; Applicant: Ardian, Inc.; dated Jan. 19, 2010, 4 pgs.
- European Examination Report; European Patent Application No. 09156661.2; Applicant: Ardian, Inc.; dated Jan. 19, 2010, 6 pgs.
- European Search Report; European Patent Application No. 05806045.0; Applicant: Ardian, Inc.; dated Sep. 22, 2009, 8 pgs.
- European Search Report; European Patent Application No. 05811851.4; Applicant: Ardian, Inc.; dated Oct. 1, 2009, 7 pgs.
- European Search Report; European Patent Application No. 06847926.0; Applicant: Ardian, Inc.; dated Feb. 10, 2010, 6 pgs.
- European Search Report; European Patent Application No. 07757925.8; Applicant: Ardian, Inc.; dated Apr. 29, 2010, 9 pgs.
- European Search Report; European Patent Application No. 07798341.9; Applicant: Ardian, Inc.; dated Aug. 4, 2011, 6 pgs.
- European Search Report; European Patent Application No. 07799148.7; Applicant: Ardian, Inc.; dated Jul. 23, 2009, 6 pgs.
- European Search Report; European Patent Application No. 07868755.5; Applicant: Ardian, Inc.; dated Jul. 28, 2010, 7 pgs.
- European Search Report; European Patent Application No. 09156661.2; Applicant: Ardian, Inc.; dated Jul. 23, 2009, 6 pgs.
- European Search Report; European Patent Application No. 09167937.3; Applicant: Ardian, Inc.; dated Nov. 11, 2009, 6 pgs.
- European Search Report; European Patent Application No. 09168202.1; Applicant: Ardian, Inc.; dated Nov. 11, 2009, 5 pgs.
- European Search Report; European Patent Application No. 09168204.7; Applicant: Ardian, Inc.; dated Nov. 19, 2009, 6 pgs.
- Evelyn, K.A. et al., Effect of thoracolumbar sympathectomy on the clinical course of primary (essential) hypertension, Am J Med, 1960;28:188-221.
- Ex parte Quayle Office Action; U.S. Appl. No. 11/144,173; dated May 28, 2009, 4 pgs.
- Fact Book Fiscal Year 2003, National Institutes of Health National Heart, Lung, and Blood Institute, Feb. 2004, 197 pgs.
- Fajardo, J. et al., Effect of chemical sympathectomy on renal hydroelectrolytic handling in dogs with chronic caval constriction. Clin Physiol Biochem. 1986;4:252-6.
- Fareed, Jawed, Ph.D. et al., Some Objective Considerations for the Use of Heparins and Recombinant Hirudin in Percutaneous Transluminal Coronary Angioplasty, Seminars in Thrombosis and Hemostasis 1991, vol. 17, No. 4, 1991 by Thieme Medical Publishers, Inc., pp. 455-470.
- Ferguson, D.R. et al., Responses of the pig isolated renal artery to transmural electrical stimulation and drugs, Dec. 7, 1984, Br. J. Pharmac. 1985, vol. 84, The Macmillan Press Ltd. 1985, pp. 879-882.
- Fernandez-Ortiz, Antonio, et al., A New Approach for Local Intravascular Drug Delivery—Iontophoretic Balloon, Intravascular Iontophoretic Local Delivery, Circulation, vol. 89, No. 4, Apr. 1994, pp. 1518-1522.
- Fields, Larry E. et al., The Burden of Adult Hypertension in the United States 1999 to 2000—A Rising Tide, May 18, 2004, American Heart Association 2004, Hypertension Oct. 2004, pp. 1-7.
- Final Office Action; U.S. Appl. No. 11/233,814; dated Jan. 29, 2009, 11 pgs.
- Final Office Action; U.S. Appl. No. 11/266,993; dated Jan. 8, 2010, 7 pgs.
- Final Office Action; U.S. Appl. No. 11/363,867; dated May 1, 2009, 8 pgs.
- Final Office Action; U.S. Appl. No. 11/451,728; dated Jan. 13, 2009, 7 pgs.
- Final Office Action; U.S. Appl. No. 11/599,649; dated Jan. 15, 2009, 10 pgs.
- Final Office Action; U.S. Appl. No. 11/599,723; dated Apr. 5, 2010, 17 pgs.
- Final Office Action; U.S. Appl. No. 11/599,890; dated Apr. 29, 2009, 9 pgs.
- Fischell, Tim A. et al., Ultrasonic Energy: Effects on Vascular Function and Integrity, Circulation: Journal of the American Heart Association. 1991. 84;pp. 1783-1795.
- Freeman, Scott A. et al., Theory of Electroporation of Planar Bilayer Membranes: Predictions of the Aqueous Area, Change in Capacitance, and Pore-Pore Separation, Feb. 23, 1994, Biophysical Journal, Jul. 1994, vol. 67, 1994 by the Biophysical Society, pp. 42-56.
- Fukuoka, Yuko et al., Imaging of neural conduction block by neuromagnetic recording, Oct. 16, 2002, Clinical Neurophysiology, vol. 113, 2002, Elsevier Science Ireland Ltd. 2002, pp. 1985-1992.
- Fuster, Valentin et al. ACC/AHA/ESC Practice Guidelines: ACA/AHA/ESC 2006 Guidelines for the Management of Patients with Atrial Fibrillation. JACC vol. 48, No. 4, Aug. 15, 2006.
- Gami, Apoor S., M.D. and Vesna D. Garovic, M.D., Contrast Nephropathy After Coronary Angiography, Mayo Clin Proc. 2004, vol. 79, 2004 Mayo Foundation for Medical Education and Research, pp. 211-219.
- Gattone II, Vincent H. et al., Contribution of Renal Innervation to Hypertension in Polycystic Kidney Disease in the Rat, University of Chicago Section of Urology, 16 pages, Mar. 17, 2008.
- Gaylor, D.C. et al., Significance of Cell Size and Tissue Structure in Electrical Trauma, Jan. 26, 1988, J. theor. Biol. 1988, vol. 133, 1988 Academic Press Limited, pp. 223-237.
- Gazdar, A.F. and G.J. Dammin, Neural degeneration and regeneration in human renal transplants, NEJM, Jul. 30, 1970, 283:222-244.
- Gehl, Julie et al., In Vivo Electroporation of Skeletal Muscle: Threshold, Efficacy and Relation to Electric Field Distribution, Biochimica et Biophysica Acta, 1428, 1999, Elsevier Science B.V. 1999, pp. 233-240, www.elsevier.com/locate/bba <<http://www.elsevier.com/locate/bba>>.
- Getts, R.T. et al., Regression of left ventricular hypertrophy after bilateral nephrectomy, Nephrol Dial Transplant, 2006, vol. 21, pp. 1089-1091.

(56)

References Cited

OTHER PUBLICATIONS

- Ghonaime, El-sayed A. et al., Percutaneous electrical nerve stimulation: an alternative to TENS in the management of sciatica, Apr. 26, 1999, *Pain* 1999, vol. 83, 1999 International Association for the Study of Pain / Published by Elsevier Science B.V., pp. 193-199.
- Gimple, M.D., Lawrence et al., Effect of Chronic Subcutaneous or Intramural Administration of Heparin on Femoral Artery Restenosis After Balloon Angioplasty in Hypercholesterolemic Rabbits, Laboratory Investigation, *Circulation*, vol. 86, No. 5, Nov. 1992, pp. 1536-1546.
- Goldberger, Jeffrey J. et al., New technique for vagal nerve stimulation, Jun. 2, 1999, *Journal of Neuroscience Methods* 91, 1999, Elsevier Science B.V. 1999, pp. 109-114.
- Gorunov, F.E. et al., The Use of Pulsed and Continuous Short Wave Diathermy (Electric Field) in Medical Rehabilitation of the Patients with Guillan-Barre Syndrome and Other Peripheral Myelinopathies, May 6, 1994, 5 pages (most of article in Russian language).
- Gottschalk, C.W., Renal nerves and sodium excretion, *Ann. Rev. Physiol.*, 1979, 41:229-240.
- Greenwell, T.J. et al., The outcome of renal denervation for managing loin pain haematuria syndrome. *BJU International*, 2004; 4 pgs.
- Gruberg, Luis, M.D. et al., The Prognostic Implications of Further Renal Function Deterioration Within 48 h of Interventional Coronary Procedures in Patients with Pre-existent Chronic Renal Insufficiency, Jun. 19, 2000, *Journal of the American College of Cardiology* 2000, vol. 36, No. 5, 2000 by the American College of Cardiology, pp. 1542-1548.
- Guimaraes, Sarfim. Vascular Adrenoceptors: An Update. pp. 319-356, Jun. 1, 2001.
- Haissaguerre, M. et al., Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins, *New England Journal of Medicine*, 1998, 339: 659-666.
- Hajjar, Ihab, M.D., M.S. and Theodore A. Kotchen, M.D., Trends in Prevalence, Awareness, Treatment, and Control of Hypertension in the United States, 1988-2000, *JAMA*, Jul. 9, 2003, vol. 290, No. 2, pp. 199-206.
- Hammer, Leah W. Differential Inhibition of Functional Dilation of Small Arterioles by Indomethacin and Glibenclamide. Hypertension. Feb. 2001 Part II. pp. 599-603.
- Hampers, C.L. et al., A hemodynamic evaluation of bilateral nephrectomy and hemodialysis in hypertensive man, *Circulation*. 1967;35:272-288.
- Hamza, M.D., Mohamed A. et al., Effect of the Duration of Electrical Stimulation on the Analgesic Response in Patients with Low Back Pain, *Anesthesiology*, vol. 91, No. 6, Dec. 1999, American Society of Anesthesiologists, Inc. 1999, pp. 1622-1627.
- Han, Hyo-Kyung and Gordon L. Amidon, Targeted Prodrug Design to Optimize Drug Delivery, Mar. 21, 2000, *AAPS PharmSci* 2000, 2 (1) article 6, pp. 1-11.
- Hansen, J.M. et al., The transplanted human kidney does not achieve functional reinnervation, *Clin Science*, 1994, vol. 87, pp. 13-20.
- Hasking, G.J. et al., Norepinephrine spillover to plasma in patients with congestive heart failure: evidence of increased overall and cardiorenal sympathetic nervous activity. *Circulation*. 1986;73:615-21.
- Hausberg, M. et al., Sympathetic nerve activity in end-stage renal disease, *Circulation*, 2002, 106: 1974-1979.
- Heart Arrhythmia Heart and Vascular Health on Yahoo! Health. 13 pgs. <URL: http://health.yahoo.com/topic/heart/overview/article/mayoclinic/21BBE2B0-128D-4AA2-A5CE215065586678;_ylt=Aqd9M5rNyHD0sbPomHXFhLcPu7cF> Feb. 16, 2005.
- Heart Disease and Stroke Statistics—2004 Update, American Heart Association, American Stroke Association, Dallas, Texas, 2003 American Heart Association, 52 pgs.
- Heida, Tjitske, et al., Investigating Membrane Breakdown of Neuronal Cells Exposed to Nonuniform Electric Fields by Finite-Element Modeling and Experiments, May 9, 2002, *IEEE Transactions on Biomedical Engineering*, vol. 49, No. 10, Oct. 2002, IEEE 2002, pp. 1195-1203.
- Heuer, G.J., The surgical treatment of essential hypertension, *Annals of Surgery*, 1936; 104 (4): 771-786.
- Higuchi, Yoshinori, M.D., Ph.D. et al, Exposure of the Dorsal Root Ganglion in Rats to Pulsed Radiofrequency Currents Activates Dorsal Horn Lamina I and II Neurons, Dec. 4, 2001, *Experimental Studies, Neurosurgery*, vol. 50, No. 4, Apr. 2002, pp. 850-856.
- Hildebrand, Keith R., D.V.M., Ph.D. et al., Stability, Compatibility, and Safety of Intrathecal Bupivacaine Administered Chronically via an Implantable Delivery System, May 18, 2001, *The Clinical Journal of Pain*, vol. 17, No. 3, 2001 Lippincott Williams & Wilkins, Inc., pp. 239-244.
- Hing, Esther, M.P.H. and Kimberly Middleton, B.S.N., M.P.H., National Hospital Ambulatory Medical Care Survey: 2001 Outpatient Department Summary, Aug. 5, 2003, Advance Data from Vital and Health Statistics, No. 338, CDC, 32 pages.
- Hodgkin, Douglas D. et al., Electrophysiologic Characteristics of a Pulsed Iontophoretic Drug-Delivery System in Coronary Arteries, *Journal of Cardiovascular Pharmacology*. 29(1):pp. 39-44, Jan. 1997, Abstract, 2 pgs.
- Hopp, F.A. et al., Respiratory Responses to Selective Blockade of Carotid Sinus Baroreceptors in the Dog, Jun. 22, 2005, *Am J Physiol Regul Integr Comp Physiol* 1998, vol. 275, 2005 American Physiological Society, pp. R10-R18.
- Hortobagyi, Gabriel N., Randomized Trial of High-Dose Chemotherapy and Blood Cell Autographs for High-Risk Primary Breast Carcinoma, *Journal of the National Cancer Institute*, vol. 92, No. 3, Feb. 2, 2000, pp. 225-233.
- Horwich, Tamara, M.D., New Advances in the Diagnosis and Management of Acute Decompensated Heart Failure, the heart.org satellite program, Rapid Review, CME Symposium presented on Nov. 8, 2004 at the Sheraton New Orleans Hotel, 4 pages.
- Huang, Wann-Chu et al. Renal Denervation Prevents and Reverses Hyperinsulinemia-Induced Hypertension in Rats, Mar. 25, 1998, *Hypertension* 1998, vol. 32, 1998 American Heart Association, pp. 249-254.
- Huang, Yifei et al., Remodeling of the chronic severely failing ischemic sheep heart after coronary microembolization: functional, energetic, structural and cellular responses, Jan. 8, 2004, *Am J Physiol. Heart Circ. Physiol.* 2004, vol. 286, 2004 the American Physiological Society, pp. H2141-H2150.
- Hughes, Gordon B., M.D. et al., A Comparative Study of Neuropathologic Changes Following Pulsed and Direct Current Stimulation of the Mouse Sciatic Nerve, Jun. 27, 1980, *American Journal of Otolaryngology*, Nov. 1980, vol. 1, No. 5, pp. 378-384.
- Hypertension and Renal Disease: Mechanisms, Slide Show by www.hypertensiononline.org, 22 pages Mar. 30, 2001.
- Hypertension Incidence and Prevalence, Age-Specific Rates, by Gender, B.C., 2001/2002, Graph, Chronic Disease Management, May 2003, British Columbia Ministry of Health Services, 1 page.
- Implantable Neurostimulation Systems, Medtronic Neurological, Jan. 18, 1999, 6 pages. http://medtronic.com/neuro/paintherapies/pain_treatment_ladder/pdf/implantable_brochure.pdf.
- Implantable Pump—The Medtronic MiniMed 2007 Implantable Insulin Pump System, Medtronic MiniMed 2004, 4 pgs.
- International Search Report and Written Opinion for PCT/US2009/069334; Applicant: Ardian, Inc.; dated Mar. 1, 2010, 10 pgs.
- International Search Report and Written Opinion, PCT/US05/35693, dated Mar. 8, 2006, Applicant: Ardian, Inc., 29 pgs.
- International Search Report and Written Opinion, PCT/US05/35757, dated Dec. 27, 2006, Applicant: Ardian, Inc., 8 pgs.
- International Search Report and Written Opinion, PCT/US06/36120, dated Jun. 25, 2008, Applicant: Ardian, Inc., 10 pgs.
- International Search Report and Written Opinion, PCT/US06/41889, dated Oct. 20, 2008, Applicant: Ardian, Inc., 7 pgs.
- International Search Report and Written Opinion, PCT/US06/48822, dated Aug. 15, 2008, Applicant: Ardian, Inc., 12 pgs.
- International Search Report and Written Opinion, PCT/US07/633222, dated Mar. 3, 2008, Applicant: Ardian, Inc., 10 pgs.

(56)

References Cited

OTHER PUBLICATIONS

- International Search Report and Written Opinion, PCT/US07/63324, dated Oct. 10, 2008, Applicant: Ardian, Inc., 10 pgs.
- International Search Report and Written Opinion, PCT/US07/66539, dated Jan. 28, 2008, Applicant: Ardian, Inc., 6 pgs.
- International Search Report and Written Opinion, PCT/US07/70799, dated Jul. 2, 2008, Applicant: Ardian, Inc., 7 pgs.
- International Search Report and Written Opinion, PCT/US07/72396, dated Aug. 27, 2008, Applicant: Ardian, Inc., 9 pgs.
- International Search Report and Written Opinion, PCT/US07/84701, dated Aug. 21, 2008, Applicant: Ardian, Inc., 11 pgs.
- International Search Report and Written Opinion, PCT/US07/84705, dated Jul. 28, 2008, Applicant: Ardian, Inc., 12 pgs.
- International Search Report and Written Opinion, PCT/US07/84708, dated Aug. 11, 2008, Applicant: Ardian, Inc., 9 pgs.
- International Search Report, PCT/US02/0039, dated Sep. 11, 2002, Applicant: Advanced Neuromodulation Systems, Inc.
- International Search Report, PCT/US02/25712, dated Apr. 23, 2003, Applicant: Cyberonics, Inc.
- International Search Report, PCT/US03/08014, dated Sep. 23, 2003, Applicant: The General Hospital Corporation.
- International Search Report, PCT/US03/09764, dated Oct. 28, 2003, Applicant: CVRX, Inc.
- International Search Report, PCT/US04/38498, dated Feb. 18, 2005, Applicant: G & L Consulting, LLC, 4 pgs.
- Introduction to Autonomic Pharmacology, Chapter 3, Part 2 Autonomic Pharmacology, pp. 18-26, May 24, 2002.
- Isovue: Data Sheet. Regional Health Limited. 8 pgs. Mar. 11, 2003.
- Israili, Z.H., Clinical pharmacokinetics of angiotensin II (AT) receptor blockers in hypertension, *Journal of Human Hypertension*, 2000, Macmillan Publishers Ltd., vol. 14, pp. S73-S86.
- Janda, J., Impact of the electrical stimulation apparatus rebox on the course of ischemic renal damage in rats, *British Library—The world's knowledge* pp. 252-254 (translated and untranslated versions) 1996.
- Janssen, Ben J.A. et al., Effects of complete renal denervation and selective afferent renal denervation on the hypertension induced by intrarenal norepinephrine infusion in conscious rats, Jan. 4, 1989, *Journal of Hypertension* 1989, vol. 7, No. 6, Current Science Ltd, pp. 447-455.
- Jia, Jianping et al., Cold injury to nerves is not due to ischaemia alone, *Brain*, 121, pp. 989-1001. 1998.
- Jia, Jianping et al., The pathogenesis of non-freezing cold nerve injury: Observations in the rat, *Brain*, 120; pp. 631-646. 1997.
- Jin, Yuanzhe et al., Pulmonary Vein Stenosis and Remodeling After Electrical Isolation for Treatment of Atrial Fibrillation: Short- and Medium-Term Follow-Up, *PACE*, vol. 27., Oct. 2004, pp. 1362-1370.
- Johansson, Bjorn, Electrical Membrane Breakdown, A Possible Mediator of the Actions of Electroconvulsive Therapy, *Medical Hypotheses* 1987, vol. 24, Longman Group UK Ltd 1987, pp. 313-324.
- Joles, J.A. et al., Causes and Consequences of Increased Sympathetic Activity in Renal Disease, *Hypertension*, 2004;43:699-706.
- Jorgensen, William A. et al., Electrochemical Therapy of Pelvic Pain: Effects of Pulsed Electromagnetic Fields (PEMF) on Tissue Trauma, *Eur J Surg* 1994, Suppl 574, vol. 160, 1994 Scandinavian University Press, pp. 83-86.
- Joshi, R. P. and K. H. Schoenbach, Mechanism for membrane electroporation irreversibility under high-intensity, ultrashort electrical pulse conditions, Nov. 11, 2002, *Physical Review E* 66, 2002, The American Physical Society 2002, pp. 052901-1-052901-4.
- Joshi, R. P. et al., Improved energy model for membrane electroporation in biological cells subjected to electrical pulses, Apr. 9, 2002, *Physical Review E*, vol. 65, 041920-1, 2002 The American Physical Society, 8 pages.
- Joshi, R. P. et al., Self-consistent simulations of electroporation dynamics in biological cells subjected to ultrashort electrical pulses, Jun. 21, 2001, *Physical Review E*, vol. 64, 011913, 2001 The American Physical Society, pp. 1-10.
- Joye, James D. et al., In Vivo Study of Endovascular Cryotherapy for the Prevention of Restenosis, 4 pages, 2003.
- Kanduser, Masa et al., Effect of surfactant polyoxyethylene glycol (C12E8) on electroporation of cell line DC3F, Aug. 20, 2002, *Colloids and Surfaces A: Physicochem. Eng. Aspects* 214, 2003, Elsevier Science B.V. 2002, pp. 205-217.
- Kassab, S. et al., Renal denervation attenuates the sodium retention and hypertension associated with obesity, *Hypertension*, 1995, 25:893-897.
- Katholi, R.E. et al., Importance of the renal nerves in established two-kidney, one clip Goldblatt hypertension, *Hypertension*, 1982, 4 (suppl II):II-166-II-174.
- Katholi, R.E. et al., Role of the renal nerves in the pathogenesis of one-kidney renal hypertension in the rat, *Hypertension*, 1981, 3(4) 404-409.
- Katholi, R.E., Renal nerves and hypertension: an update, *Fed Proc.*, 1985, 44:2846-2850.
- Katholi, Richard E., Renal nerves in the pathogenesis of hypertension in experimental animals and humans, *Am. J. Physiol.* vol. 245, 1983, the American Physiological Society 1983, pp. F1-F14.
- Kaye, D.M. et al., Functional and neurochemical evidence for partial cardiac sympathetic reinnervation after cardiac transplantation in humans, *Circulation*, 1993, vol. 88, pp. 1101-1109.
- Kelleher, Catherine L. et al., Characteristics of Hypertension in Young Adults with Autosomal Dominant Polycystic Kidney Disease Compared with the General U.S. Population, Jun. 9, 2004, *American Journal of Hypertension* 2004, pp. 1029-1034.
- King, Ronald W. P., Nerves in a Human Body Exposed to Low-Frequency Electromagnetic Fields, Jun. 7, 1999, *IEEE Transactions on Biomedical Engineering*, vol. 46, No. 12, Dec. 1999, IEEE 1999, pp. 1426-1431.
- Kinney, Brian M., M.D., High-Tech Healing—The evolution of therapeutic electromagnetic fields in plastic surgery, *Plastic Surgery Products*, Jun. 2004, pp. 32-36, 3 pages.
- Kirchheim, H. et al., Sympathetic modulation of renal hemodynamics, renin release and sodium excretion, *Klin Wochenschr*, 1989, 67:858-864.
- Klein, K. et al., Impaired autocrine regulation of hypothalamic norepinephrine release in experimental uremia, *J Am Soc Nephrol*, 2005;16:2081-7.
- Knot, H. J. et al., Regulation of arterial diameter and wall [Ca²⁺] in cerebral arteries of rat by membrane potential and intravascular pressure, *The Journal of Physiology*, 1998, 508; pp. 199-209.
- Kok, Lai Chow et al., Effect of Heating on Pulmonary Veins: How to Avoid Pulmonary Vein Stenosis, *Journal of Cardiovascular Electrophysiology*, vol. 14, No. 3, Mar. 2003, pp. 250-254.
- Kok, R. J. et al., Specific Delivery of Captopril to the Kidney with the Prodrug Captopril-Lysozyme, Aug. 16, 1998, *Journal of Pharmacology and Experimental Therapeutics*, vol. 288, No. 1, 1999 by The American Society for Pharmacology and Experimental Therapeutics, pp. 281-285.
- Kon, V. Neural Control of Renal Circulation, *Miner Electrolyte Metab*, 1989;15:33-43.
- Koomans, H.A., et al., Sympathetic hyperactivity in chronic renal failure: a wake-up call, *J Am Soc Nephrol*, 2004;15:524-37.
- Kopp, U. et al., Dietary sodium loading increases arterial pressure in afferent renal-denervated rats, *Hypertension*, 2003, 42:968-973.
- Kopp, U.C. et al., Renal sympathetic nerve activity modulates afferent renal nerve activity by PGE₂-dependent activation of alpha₁- and alpha₂-adrenoceptors on renal sensory nerve fibers, *Am J Physiol Regul Integr Comp Physiol*, 2007;293:R1561-72.
- Koyama, Shozo et al., Relative Contribution of Renal Nerve and Adrenal Gland to Renal Vascular Tone During Prolonged Canine Hemorrhagic Hypotension, Sep. 24, 1992, *Circulatory Shock* 1993, vol. 39, Wiley-Liss, Inc. 1993, pp. 269-274.
- Kozak, Lola Jean, Ph.D et al., National Hospital Discharge Survey: 2001 Annual Summary with Detailed Diagnosis and Procedure Data, Vital and Health Statistics, Series 13 No. 156, Jun. 2004, CDC, 206 pages.
- Kumagai, K. et al. New Approach to Pulmonary Vein Isolation for Atrial Fibrillation Using a Multielectrode Basket Catheter, *Circulation Journal*, 2006;70:88-93.

(56)

References Cited

OTHER PUBLICATIONS

- Lafayette, Richard A., M.D., How Does Knocking Out Angiotensin II Activity Reduce Renal Injury in Mice?, Jun. 14, 1999, Journal Club, American Journal of Kidney Diseases, vol. 35, No. 1, Jan. 2000, National Kidney Foundation, Inc. 2000, pp. 166-172.
- Lavie, Peretz, Ph.D. and Victor Hoffstein, M.D., Sleep Apnea Syndrome: A Possible Contributing Factor to Resistant Hypertension, Jun. 2001, Sleep 2001, vol. 24, No. 6, pp. 721-725.
- Le Noble, J.L. et al., Pharmacological evidence for rapid destruction of efferent renal nerves in rats by intrarenal infusion of 6-hydroxydopamine. *J Hypertens Suppl.* 1985;3:S137-40.
- Lee, Michael A. (editor). SPORTSMed. Connecticut State Medical Society Committee on the Medical Aspects of Sports. Fall/Winter 2005. 10 pgs.
- Lee, Raphael C. et al., Biophysical Injury Mechanisms in Electronic Shock Trauma, *Annu. Rev. Biomed. Eng.*, 2000, vol. 2, Copyright © 2000 by Annual Reviews, pp. 477-509.
- Lee, Raphael C. et al., Clinical Sequelae Manifested in Electrical Shock Survivors, Presentation by the Electrical Trauma Research Program, The University of Chicago, 37 pages Dec. 24, 2004.
- Lee, Raphael C. et al., Membrane Biology and Biophysics, Chapter 25, *Surgical Research*, 2001 Academic Press, pp. 297-305.
- Lee, Raphael C., M.D., Sc.D. and Michael S. Kolodney, S.B., Electrical Injury Mechanisms: Electrical Breakdown of Cell Membranes, Oct. 1, 1986, Plastic and Reconstructive Surgery, Nov. 1987, vol. 80, No. 5, pp. 672-679.
- Lenoble, L.M. et al., Selective efferent chemical sympathectomy of rat kidneys. *Am J Physiol.* 1985;249:R496-501.
- Ligtenberg, Gerry M.D. et al., Reduction of Sympathetic Hyperactivity by Enalapril in Patients with Chronic Renal Failure, Apr. 29, 1999, New England Journal of Medicine 1999, vol. 340, No. 17, 1999 Massachusetts Medical Society, pp. 1321-1328.
- Lin, Vernon W. H. et al., High intensity magnetic stimulation over the lumbosacral spine evokes antinociception in rats, Apr. 16, 2002, *Clinical Neurophysiology*, vol. 113, 2002 Elsevier Science Ireland Ltd., pp. 1006-1012.
- Lipfert, Peter, M.D. et al., Tachyphylaxis to Local Anesthetics Does Not Result from Reduced Drug Effectiveness at the Nerve Itself, Aug. 3, 1988, *Anesthesiology* 1989, vol. 70, pp. 71-75.
- Lohmeier, Thomas E. and Drew A. Hildebrandt, Renal Nerves Promote Sodium Excretion in Angiotensin-Induced Hypertension, Oct. 20, 1997, *Hypertension* 1998, vol. 31, part 2, 1998 American Heart Association, Inc., pp. 429-434.
- Lohmeier, Thomas E. et al., Prolonged Activation of the Baroreflex Produces Sustained Hypotension, Harry Goldblatt Award, Nov. 26, 2003, *Hypertension* 2004, vol. 43, Part 2, 2004 American Heart Association, Inc., pp. 306-311.
- Lohmeier, Thomas E. et al., Renal Nerves Promote Sodium Excretion During Long-Term Increases in Salt Intake, Oct. 23, 1998, *Hypertension* 1999, vol. 33, part II, 1999 American Heart Association, Inc., pp. 487-492.
- Lohmeier, Thomas E. et al., Sustained influence of the renal nerves to attenuate sodium retention in angiotensin hypertension, Apr. 13, 2001, *Am J Physiol Regulatory Integrative Comp Physiol*, vol. 281, 2001 the American Physiological Society, pp. R434-R443.
- Lohmeier, Thomas E., et al., Baroreflexes prevent neurally induced sodium retention in angiotensin hypertension, *American Journal Physiol Regulatory Integrative Comp Physiol*, vol. 279, 2000 the American Physiological Society, pp. R1437-R1448.
- Lohmeier, Thomas E., Interactions Between Angiotensin II and Baroreflexes in Long-Term Regulation of Renal Sympathetic Nerve Activity, *Circulation Research*, Jun. 27, 2003, American Heart Association, Inc. 2003, pp. 1282-1284.
- Luff, S.E. et al., Two types of sympathetic axon innervating the juxtaglomerular arterioles of the rabbit and rat kidney differ structurally from those supplying other arteries, May 1, 1991, *Journal of Neurocytology* 1991, vol. 20, 1991 Chapman and Hall Ltd., pp. 781-795.
- Luippold, G. et al., Chronic renal denervation prevents glomerular hyperfiltration in diabetic rats, *Nephrol Dial Transplant* (2004) 19:342-347.
- Lundborg, C. et al., Clinical experience using intrathecal (IT) bupivacaine infusion in three patients with complex regional pain syndrome type I (CRPS-I), *Acta Anaesthesiol Scand* 1999, vol. 43, pp. 667-678.
- Maeder, Micha, M.D. et al., Contrast Nephropathy: Review Focusing on Prevention, Jun. 22, 2004, *Journal of the American College of Cardiology* Nov. 2, 2004, vol. 44, No. 9, 2004 by the American College of Cardiology Foundation, pp. 1763-1771.
- Malpas, Simon C., What sets the long-term level of sympathetic nerve activity: is there a role for arterial baroreceptors?, *Invited Review, Am J Physiol Regul Integr Comp Physiol* 2004, vol. 286, 2004 the American Physiological Society, pp. R1-R12.
- Mancia, G., Grassi, G., Giannattasio, C., Seravalle, G., Sympathetic activation of pathogenesis of hypertension and progression of organ damage, *Hypertension* 1999, 34 (4 Pt 2): 724-728.
- Marenzi, Giancarlo, M.D. et al., The Prevention of Radiocontrast-Agent-Induced Nephropathy by Hemofiltration, *New England Journal of Medicine*, Oct. 2, 2003, vol. 349 (14), 2003 Massachusetts Medical Society, pp. 1333-1340.
- Market for infusion pumps grows with an aging population, *NWL 97-01, The BBI Newsletter*, vol. 20, No. 2, Feb. 1, 1997, American Health Consultants, Inc., pp. 6.
- Martin, Jason B. et al., Gene Transfer to Intact Mesenteric Arteries by Electroporation, Mar. 27, 2000, *Journal of Vascular Research* 2000, vol. 37, 2000 S. Karger AG, Basel, pp. 372-380.
- McCreery, Douglas B. et al., Charge Density and Charge Per Phase as Cofactors in Neural Injury Induced by Electrical Stimulation, *IEEE Transactions on Biomedical Engineering*, vol. 17, No. 10, Oct. 1990, pp. 996-1000.
- McCullough, Peter A., M.D., MPH et al., Acute Renal Failure after Coronary Intervention: Incidence, Risk Factors and Relationship to Mortality, Apr. 14, 1997, *Am J Med.* 1997, vol. 103, 1997 Excerpta Medica, Inc., pp. 368-375.
- McMurray, John J.V., M.D. and Eileen O'Meara, M.D., Treatment of Heart Failure with Spironolactone—Trial and Tribulations, Aug. 5, 2004, *New England Journal of Medicine*, vol. 351, No. 6, 2004 Massachusetts Medical Society, pp. 526-528.
- McRobbie, D. and M.A. Foster, Thresholds for biological effects of time-varying magnetic fields, Dec. 16, 1983, *Clin. Phys. Physiol. Meas.* 1984, vol. 5, No. 2, 1984 The Institute of Physics, pp. 67-78.
- Medtronic Neurostimulation Systems, Expanding the Array of Pain Control Solutions, informational pamphlet, 1999 Medtronic, Inc., 6 pages.
- Medtronic, Spinal Cord Stimulation, Patient Management Guidelines for Clinicians, Medtronic, Inc. 1999, 115 pages.
- Medtronic, SynchroMed Infusion System—Clinical Reference Guide for Pain Therapy, Medtronic, Inc. 1998, 198 pages.
- Mehran, Roxana, Renal insufficiency and contrast nephropathy: The most common, least understood risk factor, *Cardiovascular Research Foundation*, Columbia University Medical Center, 2005, 86 slides.
- Mess, Sarah A., M.D. et al., Implantable Baclofen Pump as an Adjuvant in Treatment of Pressure Sores, Mar. 1, 2003, *Annals of Plastic Surgery*, vol. 51, No. 5, Nov. 2003, Lippincott Williams & Wilkins 2003, pp. 465-467.
- Micro ETS Hyperhidrosis USA Hyperhidrosis USA. 2 pgs. <URL: <http://www.hyperhidrosis-usa.com/Index.html>>. Nov. 6, 2006.
- Mihran, Richard T. et al., Temporally-Specific Modification of Myelinated Axon Excitability in Vitro Following a Single Ultrasound Pulse, Sep. 25, 1989, *Ultrasound in Med. & Biol.* 1990, vol. 16, No. 3, pp. 297-309.
- Miklavčič, D. et al, A Validated Model of in Vivo Electric Field Distribution in Tissues for Electrochemotherapy and for DNA Electrotansfer for Gene Therapy, *Biochimica et Biophysica Acta*, 1523, 2000, pp. 73-83, <<http://www.elsevier.com/locate/bba>>.
- Mitchell, G. A. G., The Nerve Supply of the Kidneys, Aug. 20, 1949, *Acta Anatomica*, vol. X, Fasc. ½, 1950, pp. 1-37.
- Morrissey, D.M. et al., Sympathectomy in the treatment of hypertension: Review of 122 cases, *Lancet.* 1953;1:403-408.

(56)

References Cited

OTHER PUBLICATIONS

- Moss, Nicholas G., Renal function and renal afferent and efferent nerve activity, *Am. J. Physiol.* 1982, vol. 243, 1982 the American Physiological Society, pp. F425-F433.
- Munglani, Rajesh, The longer term effect of pulsed radiofrequency for neuropathic pain, Jun. 8, 1998, *Pain* 80, 1999, International Association for the Study of Pain 1999, Published by Elsevier Science B.V., pp. 437-439.
- Naropin (ropivacaine HCl) Injection, RX only Description, AstraZeneca 2001, 3 pages.
- National High Blood Pressure Education Program, 1995 Update of the Working Group Reports on Chronic Renal Failure and Renovascular Hypertension, presentation, 13 pages.
- National Kidney Foundation, Are You At Increased Risk for Chronic Kidney Disease?, 2002 National Kidney Foundation, Inc., 14 pages.
- Nelson, L. et al., Neurogenic Control of Renal Function in Response to Graded Nonhypotensive Hemorrhage in Conscious Dogs, Sep. 13, 1992, *Am J. Physiol.* 264, 1993, American Physiological Society 1993, pp. R661-R667.
- Nikolsky, Eugenia, M.D. et al., Radiocontrast Nephropathy: Identifying the High-Risk Patient and the Implications of Exacerbating Renal Function, *Rev Cardiovasc Med.* 2003, vol. 4, Supp. 1, 2003 MedReviews, LLC, pp. S7-S14.
- Non-Final Office Action; U.S. Appl. No. 10/408,665; dated Mar. 21, 2006, 14 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/129,765; dated May 18, 2007, 10 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/129,765; dated Sep. 10, 2007, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/129,765; dated Oct. 6, 2006, 30 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/133,925; dated Oct. 8, 2008, 41 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/144,173; dated Apr. 5, 2007, 33 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/144,173; dated Sep. 10, 2007, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/144,298; dated Oct. 29, 2009, 8 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/144,298; dated Apr. 5, 2007, 33 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/144,298; dated Sep. 10, 2007, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/144,298; dated Dec. 29, 2008, 7 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/145,122; dated Apr. 11, 2007, 33 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/145,122; dated Sep. 10, 2007, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/189,563; dated May 28, 2009, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/233,814; dated Jun. 17, 2008, 12 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/252,462; dated Feb. 22, 2010, 6 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/266,993; dated Jul. 8, 2009, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/266,993; dated Dec. 30, 2008, 7 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/363,867; dated Sep. 25, 2008, 10 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/368,553; dated May 18, 2010, 4 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/368,553; dated Oct. 7, 2009, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/368,809; dated Dec. 3, 2009, 4 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/368,949; dated Jun. 11, 2010, 6 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/368,971; dated Aug. 24, 2010, 9 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/451,728; dated Jun. 12, 2008, 41 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/451,728; dated Jul. 2, 2009, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/451,728; dated Dec. 28, 2009, 7 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/504,117; dated Mar. 31, 2009, 10 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/599,649; dated Mar. 30, 2009, 10 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/599,649; dated Jun. 23, 2008, 9 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/599,723; dated Jun. 26, 2009, 17 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/599,723; dated Oct. 15, 2010, 16 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/599,882; dated Jul. 6, 2009, 13 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/688,178; dated Jun. 28, 2010, 5 pgs.
- Non-Final Office Action; U.S. Appl. No. 11/840,142; dated Apr. 3, 2009, 13 pgs.
- Non-Final Office Action; U.S. Appl. No. 12/567,521; dated Sep. 3, 2010, 9 pgs.
- Non-Final Office Action; U.S. Appl. No. 12/616,708; dated Sep. 16, 2010, 10 pgs.
- Non-Final Office Action; U.S. Appl. No. 12/725,375; dated Oct. 12, 2010, 14 pgs.
- Nozawa, T. et al., Effects of Long Term Renal Sympathetic Denervation on Heart Failure After Myocardial Infarction in Rats, Sep. 22, 2001, *Heart Vessels*, 2002, 16, Springer-Verlag 2002, pp. 51-56.
- O'Hagan, K.P. et al., Renal denervation decreases blood pressure in DOCA-treated miniature swine with established hypertension, *Am J Hypertens.*, 1990, 3:62-64.
- Onesti, G. et al., Blood pressure regulation in end-stage renal disease and anephric man, *Circ Res Suppl.*, 1975, 36 & 37: 145-152.
- Osborn, et al., Effect of renal nerve stimulation on renal blood flow autoregulation and antinatriuresis during reductions in renal perfusion pressure, in *Proceedings of the Society for Experimental Biology and Medicine*, vol. 168, 77-81, 1981. (Abstract).
- Packer, Douglas L. et al., Clinical Presentation, Investigation, and Management of Pulmonary Vein Stenosis Complication Ablation for Atrial Fibrillation, *Circulation: Journal of the American Heart Association*. Feb. 8, 2005. pp. 546-554.
- Page, I.H. et al., The Effect of Renal Denervation on the Level of Arterial Blood Pressure and Renal Function in Essential Hypertension. *J Clin Invest.* 1935;14:27-30.
- Page, I.H., et al., The Effect of Renal Efficiency of Lowering Arterial Blood Pressure in Cases of Essential Nephritis, *Hospital of the Rockefeller Institute*, Jul. 12, 1934, 7 pgs.
- Palmer, Biff, F., M.D., Managing Hyperkalemia Caused by Inhibitors of the Renin-Angiotensin-Aldosterone System, Aug. 5, 2004, *The New England Journal of Medicine* 2004, vol. 351;6, 2004 Massachusetts Medical Society, pp. 585-592.
- Pappone, Carlo et al., [2005][P2-70] Safety Report of Circumferential Pulmonary Vein Ablation. A 9-Year Single-Center Experience on 6,442 Patients with Atrial Fibrillation, Abstract only. 1 page, May 2005.
- Pappone, Carlo et al., [2004][759] Pulmonary Vein Denervation Benefits Paroxysmal Atrial Fibrillation Patients after Circumferential Ablation, Abstract only. 1 page, Jan. 5, 2004.
- Pappone, Carol and Santinelli, Vincenzo. Multielectrode basket catheter: A new tool for curing atrial fibrillation? *Heart Rhythm*, vol. 3, Issue 4, pp. 385-386. Apr. 2006.
- Peacock, J.M. and R. Orchardson, Action potential conduction block of nerves in vitro by potassium citrate, potassium tartrate and potassium oxalate, May 6, 1998, *Journal of Clinical Periodontology*, Munksgaard 1999, vol. 26, pp. 33-37.
- Petersson, M. et al., Long-term outcome in relation to renal sympathetic activity in patients with chronic heart failure. *Eur Heart J.* 2005;26:906-13.

(56)

References Cited

OTHER PUBLICATIONS

- Pettersson, A. et al., Renal interaction between sympathetic activity and ANP in rats with chronic ischaemic heart failure, Nov. 25, 1988, *Acta Physiol Scand* 135, pp. 487-492.
- PHCL 762 Pharmacology of the Autonomic Nervous System, Chapter 2 and 6.8 in Mosby, <http://www.kumc.edu/research/medicine/pharmacology/CAI/phc1762.html>, last accessed Aug. 24, 2004, 14 pgs.
- Pitt, B. et al., Effects of Eplerenone, Enalapril, and Eplerenone/Enalapril in Patients With Essential Hypertension and Left Ventricular Hypertrophy: The 4E-Left Ventricular Hypertrophy Study, *Circulation*, 2003, vol. 108, pp. 1831-1838.
- Pliquett, U., Joule heating during solid tissue electroporation, Oct. 22, 2002, *Med. Biol. Eng. Comput.*, 2003, vol. 41, pp. 215-219.
- Podhajsky R.J. et al, The Histologic Effects of Pulsed and Continuous Radiofrequency Lesions at 42 C to Rat Dorsal Root Ganglion and Sciatic Nerve, *Spine*, vol. 30, No. 9, 2005, Lippincott Williams & Wilkins Inc., pp. 1008-1013.
- Pope, Jill. Fixing a Hole: Treating Injury by Repairing Cells. The New York Academy of Sciences. Jul. 6, 2006. 6 pgs.
- Popovic, Jennifer R. and Margaret J. Hall, 1999 National Hospital Discharge Survey, Apr. 24, 2001, *Advance Data*, No. 319, CDC, pp. 1-17 & 20.
- Practice Guidelines Writing Committee and ESH/ESC Hypertension Guidelines Committee, Practice Guidelines for Primary Care Physicians: 2003 ESH/ESC Hypertension Guidelines, Published in *Journal of Hypertension* 2003, vol. 21, No. 10: 1011-1053, European Society of Hypertension 2003, pp. 1779-1786.
- Programmable Infusion System, Pumps and Pump Selection, Medtronic Pain Therapies, Medtronic, Inc. Sep. 5, 2001, 2 pgs.
- Pucihar, Gorazd et al., The influence of medium conductivity on electroporation and survival of cells in vitro, May 31, 2001, *Bioelectrochemistry*, vol. 54, 2001, Elsevier Science B.V. 2001, pp. 107-115.
- Pulmonary Concepts in Critical Care Breath Sounds, <http://mbob.tripod.com/breath.htm>, last accessed Aug. 23, 2004, 5 pages.
- Pulmonary Function Testing, <http://jan.ucc.nau.edu/~daa/lecture/pft.htm>, last accessed Aug. 23, 2004, 8 pages.
- Purerfellner, Helmut and Martinek, Martin. Pulmonary vein stenosis following catheter ablation of atrial fibrillation. *Current Opinion in Cardiology*. 20; pp. 484-490. 2005.
- Purerfellner, Helmut et al., Pulmonary Vein Stenosis by Ostial Irrigated-Tip Ablation: Incidence, Time Course, and Prediction, *Journal of Cardiovascular Electrophysiology*. vol. 14, No. 2, Feb. 2003. pp. 158-164.
- Raji, A. R. M. and R. E. M. Bowden, Effects of High-Peak Pulsed Electromagnetic Field on the Degeneration and Regeneration of the Common Peroneal Nerve in Rats, *The Journal of Bone and Joint Surgery* Aug. 1983, vol. 65-B, No. 4, 1983 British Editorial Society of Bone and Joint Surgery, pp. 478-492.
- Ram, C. Venkata S., M.D., Understanding refractory hypertension, May 15, 2004, *Patient Care* May 2004, vol. 38, pp. 12-16, 7 pages from <http://www.patientcareonline.com/patcare/content/printContentPopup.jsp?id=108324>.
- Ravalia, A. et al., Tachyphylaxis and epidural anaesthesia, *Edgware General Hospital, Correspondence*, p. 529, Jun. 1989.
- Renal Parenchymal Disease, Ch. 26, 5th Edition *Heart Disease, A Textbook of Cardiovascular Medicine* vol. 2, Edited by Eugene Braunwald, WB Saunders Company, pp. 824-825 1997.
- Ribstein, Jean and Michael H. Humphreys, Renal nerves and cation excretion after acute reduction in functioning renal mass in the rat, Sep. 22, 1983, *Am. J. Physiol.*, vol. 246, 1984 the American Physiological Society, pp. F260-F265.
- Richebe, Philippe, M.D. et al., Immediate Early Genes after Pulsed Radiofrequency Treatment: Neurobiology in Need of Clinical Trials, Oct. 13, 2004, *Anesthesiology* Jan. 2005, vol. 102, No. 1, 2004 American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc., pp. 1-3.
- Rihal, Charanjit S. et al., Incidence and Prognostic Importance of Acute Renal Failure After Percutaneous Coronary Intervention, Mar. 6, 2002, *Circulation* May 14, 2002, vol. 10, 2002 American Heart Association, Inc., pp. 2259-2264.
- Rosen, S.M. et al., Relationship of Vascular Reactivity to Plasma Renin Concentration in Patients with Terminal Renal Failure, *Proc. Dialysis Transplant Forum* 1974, pp. 45-47.
- Roth, Bradley J. and Peter J. Basser, A Model of the Stimulation of a Nerve Fiber by Electromagnetic Induction, *IEEE Transactions on Biomedical Engineering*, vol. 37, No. 6, Jun. 1990, pp. 588-597.
- Rudin, Asa, M.D. et al., Postoperative Epidural or Intravenous Analgesia after Major Abdominal or Thoraco-Abdominal Surgery, *The Journal of the American Society of Anesthesiologists, Inc.*, *Anesthesiology* 2001, vol. 95, A-970, 1 page.
- Rudnick, Michael R. et al., Contrast-induced nephropathy: How it develops, how to prevent it, *Cleveland Clinic Journal of Medicine* Jan. 2006, vol. 73, No. 1, pp. 75-87.
- Rump, L.C., The Role of Sympathetic Nervous Activity in Chronic Renal Failure, *J Clinical Basic Cardiology* 2001, vol. 4, pp. 179-182.
- Ruohonen, Jarmo et al., Modeling Peripheral Nerve Stimulation Using Magnetic Fields, *Journal of the Peripheral Nervous System*, vol. 2, No. 1, 1997, Woodland Publications 1997, pp. 17-29.
- Saad, Eduardo B. et al., Pulmonary Vein Stenosis After Radiofrequency Ablation of Atrial Fibrillation: Functional Characterization, Evolution, and Influence of the Ablation Strategy, *Circulation*. 108; pp. 3102-3107. 2003.
- Sabbah, Hani N., Animal Models for Heart Failure and Device Development, Henry Ford Health System. 24 slides, Oct. 17, 2005.
- Schauerte, P et al., Focal atrial fibrillation: experimental evidence for a pathophysiologic role of the autonomic nervous system, *Journal of Cardiovascular Electrophysiology*. 12(5). May 2001. Abstract only. 2 pgs.
- Schauerte, P et al., Catheter ablation of cardiac autonomic nerves for prevention of vagal atrial fibrillation, *Circulation*. 102(22). Nov. 28, 2000. Abstract only. 2 pgs.
- Schauerte, P et al., Transvenous parasympathetic nerve stimulation in the inferior vena cava and atrioventricular conduction, *Journal of Cardiovascular Electrophysiology*. 11(1). Jan. 2000. Abstract only. 2 pgs.
- Scheiner, Avram, Ph.D., The design, development and implementation of electrodes used for functional electrical stimulation, Thesis paper, Case Western Reserve University, May 1992, 220 pages.
- Scherlag, BJ and Po, S., The intrinsic cardiac nervous system and atrial fibrillation, *Current Opinion in Cardiology*. 21(1):51-54, Jan. 2006. Abstract only. 2 pgs.
- Schlaich, M.P. et al., Relation between cardiac sympathetic activity and hypertensive left ventricular hypertrophy. *Circulation*. 2003;108:560-5.
- Schlaich, M.P. et al., Sympathetic augmentation in hypertension: role of nerve firing, norepinephrine reuptake, and angiotensin neuromodulation, *Hypertension*, 2004, 43:169-175.
- Schmitt, Joseph et al., Intravascular Optical Coherence Tomography—Opening a Window into Coronary Artery Disease, *LightLab Imaging, Inc. Business Briefing: European Cardiology* 2005.
- Schoenbach, Karl H. et al, Intracellular Effect of Ultrashort Electrical Pulses, Dec. 26, 2000, *Bioelectromagnetics*, vol. 22, 2001, Wiley-Liss, Inc. 2001, pp. 440-448.
- Schrier, Robert et al., Cardiac and Renal Effects of Standard Versus Rigorous Blood Pressure Control in Autosomal-Dominant Polycystic Kidney Disease, Mar. 23, 2002, *Journal of the American Society of Nephrology, American Society of Nephrology* 2002, pp. 1733-1739.
- Scremin, Oscar U., M.D., Ph.D. and Daniel P. Holschneider, M.D., 31 & 32.. An Implantable Bolus Infusion Pump for the Neurosciences, *FRP*, Apr. 2005, 3 pages.
- Sensorcaine—MPF Spinal Injection, informational document, AstraZeneca 2001, 2 pgs.
- Shah, D.C., Haissaguerre, M., Jais, P., Catheter ablation of pulmonary vein foci for atrial fibrillation: pulmonary vein foci ablation for atrial fibrillation, *Thorac Cardiovasc Surg*, 1999, 47 (suppl. 3): 352-356.
- Shannon, J.L. et al., Studies on the innervation of human renal allografts, *J Pathol*. 1998, vol. 186, pp. 109-115.

(56)

References Cited

OTHER PUBLICATIONS

- Shlipak, M.G. et al., The clinical challenge of cardiorenal syndrome. *Circulation*. 2004;110:1514-7.
- Shupak, Naomi M., Therapeutic Uses of Pulsed Magnetic-Field Exposure: A Review, *Radio Science Bulletin* Dec. 2003, No. 307, pp. 9-32.
- Shu-Qing, Liu et al., Old spinal cord injury treated by pulsed electric stimulation, General Hospital of Beijing Command, Beijing, Dec. 6, 1990, 5 pages (full article in Chinese; abstract on last page).
- Siegel, RJ et al., Clinical demonstration that catheter-delivered ultrasound energy reverses arterial vasoconstriction, *Journal of the American College of Cardiology*. 1992. 20; 732-735. Summary only. 2 pgs.
- Simpson, B. et al., Implantable spinal infusion devices for chronic pain and spasticity: an accelerated systematic review, ASERNIP-S Report No. 42, Adelaide, South Australia, ASERNIP-S, May 2003, 56 pages.
- Sisken, B.F. et al., 229.17 Influence of Non-Thermal Pulsed Radiofrequency Fields (PRF) on Neurite Outgrowth, *Society for Neuroscience*, vol. 21, 1995, 2 pages.
- Skeie, B. et al., Effect of chronic bupivacaine infusion on seizure threshold to bupivacaine, Dec. 28, 1986, *Acta Anaesthesiol Scand* 1987, vol. 31, pp. 423-425.
- Skopec, M., A Primer on Medical Device Interactions with Magnetic Resonance Imaging Systems, Feb. 4, 1997, CDRH Magnetic Resonance Working Group, U.S. Department of Health and Human Services, Food and Drug Administration, Center for Devices and Radiological Health, Updated May 23, 1997, 17 pages, <http://www.fda.gov/cdrh/ode/primerf6.html>, (last accessed Jan. 23, 2006).
- Slappendel, Robert et al., The efficacy of radiofrequency lesioning of the cervical spinal dorsal root ganglion in a double blinded randomized study, Jun. 26, 1997, *Pain* 73, 1997 International Association for the Study of Pain, Elsevier Science B.V., pp. 159-163.
- Sluijter, M.D., Ph.D., Pulsed Radiofrequency, May 17, 2005, *Anesthesiology* Dec. 2005, vol. 103, No. 6, 2005 American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc., pp. 1313-1314.
- Sluijter, M.D., Ph.D., Radiofrequency Part 1: The Lumbosacral Region, Chapter 1 Mechanisms of Chronic Pain and part of Chapter 2 Spinal Pain, 2001 FlivoPress SA, Meggen (LU), Switzerland, pp. 1-26.
- Sluijter, M.D., Ph.D., Radiofrequency Part 2: Thoracic and Cervical Region, Headache and Facial Pain, various pages from, FlivoPress SA, Meggen (LU), Switzerland, 13 pages 2002.
- Sluijter, M.D., Ph.D., The Role of Radiofrequency in Failed Back Surgery Patients, *Current Review of Pain* 2000, vol. 4, 2000 by Current Science Inc., pp. 49-53.
- Smithwick, R.H. et al., Hypertension and associated cardiovascular disease: comparison of male and female mortality rates and their influence on selection of therapy, *JAMA*, 1956, 160:1023-1033.
- Smithwick, R.H. et al., Splanchicectomy for essential hypertension, *Journal Am Med Assn*, 1953;152:1501-1504.
- Smithwick, R.H., Surgical treatment of hypertension, *Am J Med* 1948, 4:744-759.
- Sobotka, Paul A., Treatment Strategies for Fluid Overload, CHF Patients, CHF Solutions. Transcatheter Cardiovascular Therapeutics 2005. 20 slides.
- Solis-Herruzo, J.A. et al., Effects of lumbar sympathetic block on kidney function in cirrhotic patients with hepatorenal syndrome, *Journal of Hepatology*, 1987; 5: 167-173.
- Souza, D.R.B. et al., Chronic experimental myocardial infarction produces antinatriuresis by a renal nerve-dependent mechanism, Oct. 14, 2003, *Brazilian Journal of Medical and Biological Research* 2004, vol. 37, pp. 285-293.
- Standl, Thomas, M.D., et al., Patient-controlled epidural analgesia reduces analgesic requirements compared to continuous epidural infusion after major abdominal surgery, Aug. 29, 2002, *Canada Journal of Anesthesia* 2003, vol. 50 (3), pp. 258-264.
- Steffen, W. et al., Catheter-delivered high intensity, low frequency ultrasound induces vasodilation in vivo, *European Heart Journal*. 1994. 15; pp. 369-376.
- Steg, PG et al., Pulsed ultraviolet laser irradiation produces endothelium-independent relaxation of vascular smooth muscle, *Circulation: Journal of the American Heart Association*. 1989. pp. 189-197.
- Stone, Gregg W., M.D. et al., Fenoldopam Mesylate for the Prevention of Contrast-Induced Nephropathy, *JAMA* Nov. 5, 2003, vol. 290, No. 17, 2003 American Medical Association, pp. 2284-2291.
- Strojek, K. et al., Lowering of microalbuminuria in diabetic patients by a sympatheticoplegic agent: novel approach to prevent progression of diabetic nephropathy? *J Am Soc Nephrol*. 2001;12:602-5.
- Summary, Critical Reviews in Biomedical Engineering, vol. 17, Issue 5, 1989, pp. 515-529.
- Sung, Duk Hyun, M.D. et al., Phenol Block of Peripheral Nerve Conduction: Titrating for Optimum Effect, Jun. 27, 2000, *Arch. Phys. Med. Rehabil.* vol. 82, May 2001, pp. 671-676.
- Taka, Tomomi et al., Impaired Flow-Mediated Vasodilation in vivo and Reduced Shear-Induced Platelet Reactivity in vitro in Response to Nitric Oxide in Prothrombotic, Stroke-Prone Spontaneously Hypertensive Rats, *Pathophysiology of Haemostasis and Thrombosis*. Dec. 23, 2002. pp. 184-189.
- Taler, Sandra J. et al., Resistant Hypertension, Comparing Hemodynamic Management to Specialist Care, Mar. 12, 2002, *Hypertension* 2002, vol. 39, 2002 American Heart Association, Inc., pp. 982-988.
- Tamborero, David et al., Incidence of Pulmonary Vein Stenosis in Patients Submitted to Atrial Fibrillation Ablation: A Comparison of the Selective Segmental Ostial Ablation vs. the Circumferential Pulmonary Veins Ablation, *Journal of Interventional Cardiac Electrophysiology*. 14; pp. 41-25. 2005.
- Tay, Victoria KM, et al., Computed tomography fluoroscopy-guided chemical lumbar sympathectomy: Simple, safe and effective, Oct. 31, 2001, *Diagnostic Radiology, Australasian Radiology* 2002, vol. 46, pp. 163-166.
- Terashima, Mitsuyasu et al. Feasibility and Safety of a Novel CryoPlasty™ System. Poster. 1 page, Mar. 15, 2002.
- Thatipelli et al., CT Angiography of Renal Artery Anatomy for Evaluating Embolic Protection Devices, *Journal of Vascular and Interventional Radiology*, Jul. 2007, pp. 842-846.
- The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial, ALLHAT Research Group, *JAMA*, 2002, vol. 288, pp. 2981-2997.
- Thomas, John R. and Oakley, E. Howard N. Chapter 15: Nonfreezing Cold Injury Medical Aspects of Harsh Environments, vol. 1. pp. 467-490, 2001.
- Thompson, Gregory W., et al., Bradycardia Induced by Intravascular Versus Direct Stimulation of the Vagus Nerve, Aug. 24, 1997, *The Society of Thoracic Surgeons* 1998, pp. 637-642.
- Thrasher, Terry N., Unloading arterial baroreceptors causes neurogenic hypertension, Dec. 4, 2001, *Am J. Physiol Regulatory Integrative Comp Physiol*, vol. 282, 2002 the American Physiological Society, pp. R1044-R1053.
- Tokuno, Hajime A. et al., Local anesthetic effects of cocaethylene and isopropylcocaine on rat peripheral nerves, Oct. 7, 2003, *Brain Research* 996, 2004, Elsevier B.V. 2003, pp. 159-167.
- Trapani, Angelo J. et al., Neurohumoral interactions in conscious dehydrated rabbit, *Am. J. Physiol*. 254, 1988, the American Physiological Society 1988, pp. R338-R347.
- Trock, David H. et al., The Effect of Pulsed Electromagnetic Fields in the Treatment of Osteoarthritis of the Knee and Cervical Spine. Report of Randomized, Double Blind, Placebo Controlled Trials, Mar. 22, 1994, *The Journal of Rheumatology* 1994, vol. 21, pp. 1903-1911.
- Troiano, Gregory C. et al., The Reduction in Electroporation Voltages by the Addition of a Surfactant to Planar Lipid Bilayers, May 12, 1998, *Biophysical Journal*, vol. 75, Aug. 1998, the Biophysical Society 1998, pp. 880-888.
- Trumble, Dennis R. and James A. McGovern, Comparison of Dog and Pig Models for Testing Substernal Cardiac Compression Devices, Nov. 2003, *ASAIO Journal* 2004, pp. 188-192.
- Tsai, E., Intrathecal drug delivery for pain indications, technique, results, *Pain Lecture presentation*, Jun. 8, 2001, 31 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Uematsu, Toshihiko, M.D., Ph.D., F.I.C.A. et al., Extrinsic Innervation of the Canine Superior Vena Cava, Pulmonary, Portal and Renal Veins, *Angiology—Journal of Vascular Diseases*, Aug. 1984, pp. 486-493.
- United States Renal Data System, USRDS 2003 Annual Data Report: Atlas of End-Stage Renal Disease in the United States, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 2003, 593 pages.
- Upadhyay, Pramod, Electroporation of the skin to deliver antigen by using a piezo ceramic gas igniter, Jan. 27, 2001, *International Journal of Pharmaceutics*, vol. 217, 2001 Elsevier Science B.V., pp. 249-253.
- Valente, John F. et al., Laparoscopic renal denervation for intractable ADPKD-related pain, Aug. 24, 2000, *Nephrol Dial Transplant* 2001, vol. 16, European Renal Association—European Dialysis and Transplant Association, p. 160.
- Van Antwerp, Bill and Poonam Gulati, Protein Delivery from Mechanical Devices Challenges and Opportunities, Medtronic presentation, 19 pages, Jul. 2003.
- Velazquez, Eric J., An international perspective on heart failure and left ventricular systolic dysfunction complicating myocardial infarction: the Valiant registry, Aug. 5, 2004, *European Heart Journal* vol. 25, 2004 Elsevier, pp. 1911-1919.
- Velez-Roa, Sonia, M.D. et al., Peripheral Sympathetic Control During Dobutamine Infusion: Effects of Aging and Heart Failure, Jul. 7, 2003, *Journal of the American College of Cardiology*, vol. 42, No. 9, 2003, American College of Cardiology Foundation 2003, pp. 1605-1610.
- Villarreal, Daniel et al., Effects of renal denervation on postprandial sodium excretion in experimental heart failure, Oct. 29, 1993, *Am J Physiol* 266, 1994, pp. R1599-R1604.
- Villarreal, Daniel et al., Neurohumoral modulators and sodium balance in experimental heart failure, Nov. 6, 1992, *Am. J. Physiol*, vol. 264, 1993, pp. H1187-H1193.
- Vonend, O. et al., Moxonidine treatment of hypertensive patients with advanced renal failure. *J Hypertens*. 2003;21:1709-17.
- Wagner, C.D. et al., Very low frequency oscillations in arterial blood pressure after autonomic blockade in conscious dogs, Feb. 5, 1997, *Am J Physiol Regul Integr Comp Physiol* 1997, vol. 272, 1997 the American Physiological Society, pp. 2034-2039.
- Wald, Jan D., Ph.D. et al., *Cardiology Update*: 2003, Sep. 11, 2003, AG Edwards 2003, 120 pages.
- Wang, Xi et al., Alterations of adenylyl cyclase and G proteins in aortocaval shunt-induced heart failure, Jul. 2004, *Am J Physiol Heart Circ Physiol* vol. 287, 2004 the American Physiological Society, pp. H118-H125.
- Weaver, James C., Chapter 1 Electroporation Theory, Concepts and Mechanisms, *Methods in Molecular Biology*, vol. 55, Plant Cell Electroporation and Electrofusion Protocols, Edited by J.A. Nickoloff, Humana Press Inc., pp. 3-28, 1995.
- Weaver, James C., Electroporation: A General Phenomenon for Manipulating Cells and Tissues, Oct. 22, 1992, *Journal of Cellular Biochemistry*, vol. 51, 1993 Wiley-Liss, Inc., pp. 426-435.
- Weiner, Richard L., M.D., Peripheral nerve neurostimulation, *Neurosurg. Clin. N. Am.* vol. 14, 2003, Elsevier, Inc. 2003, pp. 401-408.
- Weisbord, Steven D., M.D. and Paul M. Palevsky, M.D., Radiocontrast-Induced Acute Renal Failure, Jul. 10, 2004, *Journal of Intensive Care Medicine* 2005, vol. 20 (2), 2005 Sage Publications, pp. 63-75.
- Whitelaw, G.P., Kinsey, D., Smithwick, R.H., Factors influencing the choice of treatment in essential hypertension: surgical, medical, or a combination of both, *Am J Surg*, 1964, 107:220-231.
- Wilson, D.H. et al., The Effects of Pulsed Electromagnetic Energy on Peripheral Nerve Regeneration, *Annals New York Academy of Sciences*, Oct. 1974, pp. 575-585.
- Wolinsky, Harvey, M.D. PhD and Swan N. Thung, M.D., Use of a Perforated Balloon Catheter to Deliver Concentrated Heparin Into the Wall of the Normal Canine Artery, Aug. 30, 1989, *JACC* 1990, vol. 15, 1990 by the American College of Cardiology, pp. 475-481.
- Wyss, J. Michael et al., Neuronal control of the kidney: Contribution to hypertension, Apr. 8, 1991, *Can. J. Physiol. Pharmacol.* 1992;70: 759-770.
- Yamaguchi, Jun-ichi, M.D. et al., Prognostic Significance of Serum Creatinine Concentration for In-Hospital Mortality in Patients with Acute Myocardial Infarction Who Underwent Successful Primary Percutaneous Coronary Intervention (from the Heart Institute of Japan Acute Myocardial Infarction [HIJAMI] Registry), Feb. 24, 2004, *The American Journal of Cardiology* vol. 93, Jun. 15, 2004, 2004 by Excerpta Medica, Inc., pp. 1526-1528.
- Ye, Richard D., M.D., Ph.D., *Pharmacology of the Peripheral Nervous System*, E-425 MSB, 6 pages, Jan. 2000.
- Zucker, Irving H. et al., The origin of sympathetic outflow in heart failure: the roles of angiotensin II and nitric oxide, *Progress in Biophysics & Molecular Biology*, vol. 84, 2004, Elsevier Ltd. 2003, pp. 217-232.
- Zundert, Jan Van, M.D. FIPP and Alex Cahana, M.D. DAAPM, Pulsed Radiofrequency in Chronic Pain Management: Looking for the Best Use of Electrical Current, *Pain Practice* 2005, vol. 5, Issue 2, 2005 World Institute of Pain, pp. 74-76.
- Ye, S. et al., A limited renal injury may cause a permanent form of neurogenic hypertension. *Am J Hypertens*. 1998;11:723-8.
- Ye, Shaohua et al., Renal Injury Caused by Intrarenal Injection of Pheno Increases Afferent and Efferent Renal Sympathetic Nerve Activity, Mar. 12, 2002, *American Journal of Hypertension*, Aug. 2002, vol. 15, No. 8, 2002 the American Journal of Hypertension, Ltd. Published by Elsevier Science Inc., pp. 717-724.
- Yong-Quan, Dong et al., The therapeutic effect of pulsed electric field on experimental spinal cord injury, *Beijing Army General Hospital of People's Liberation Army*, Beijing, 5 pages (full article in Chinese; abstract on last page) Mar. 30, 1992.
- Young, James B., M.D., FACC, Management of Chronic Heart Failure: What Do Recent Clinical Trials Teach Us?, *Reviews in Cardiovascular Medicine*, vol. 5, Suppl. 1, 2004, MedReviews, LLC 2004, pp. S3-S9.
- Yu, Wen-Chung et al., Acquired Pulmonary Vein Stenosis after Radiofrequency Catheter Ablation of Paroxysmal Atrial Fibrillation. *Journal of Cardiovascular Electrophysiology*. vol. 12, No. 8. Aug. 2001. pp. 887-892.
- Zanchetti, A. et al., Neural Control of the Kidney—Are There Reno-Renal Reflexes?, *Clin. and Exper. Hyper. Theory and Practice*, A6 (1&2), 1984, Marcel Dekker, Inc. 1984, pp. 275-286.
- Zanchetti, A. et al., Practice Guidelines for Primary Care Physicians: 2003 ESH/ESC Hypertension Guidelines, *Journal of Hypertension*, vol. 21, No. 10, 2003, pp. 1779-1786.
- Zanchetti, A.S., Neural regulation of renin release: Experimental evidence and clinical implications in arterial hypertension, *Circulation*, 1977, 56(5) 691-698.
- Zimmermann, Ulrich, *Electrical Breakdown, Electroporation and Electrofusion*, Rev. Physiol. Biochem. Pharmacol., vol. 105, Springer-Verlag 1986, pp. 175-256.
- Zoccali, C. et al., Plasma norepinephrine predicts survival and incident cardiovascular events in patients with end-stage renal disease. *Circulation*. 2002;105:1354-9.
- Ahmed, Humera et al., Renal Sympathetic Denervation Using an Irrigated Radiofrequency Ablation Catheter for the Management of Drug-Resistant Hypertension, *JACC Cardiovascular Interventions*, vol. 5, No. 7, 2012, pp. 758-765.
- Avital et al., "The creation of linear contiguous lesions in the atria with an expandable loop catheter," *Journal of the American College of Cardiology*, 1999; 33; pp. 972-984.
- Beale et al., "Minimally Invasive Treatment for Varicose Veins: A Review of Endovenous Laser Treatment and Radiofrequency Ablation". *Lower Extremity Wounds* 3(4), 2004, 10 pages.
- Blessing, Erwin et al., Cardiac Ablation and Renal Denervation Systems Have Distinct Purposes and Different Technical Requirements, *JACC Cardiovascular Interventions*, vol. 6, No. 3, 2013, 1 page.
- ClinicalTrials.gov, Renal Denervation in Patients with uncontrolled Hypertension in Chinese (2011), 6pages. www.clinicaltrials.gov/ct2/show/NCT01390831.

(56)

References Cited

OTHER PUBLICATIONS

Excerpt of Operator's Manual of Boston Scientific's EPT-1000 XP Cardiac Ablation Controller & Accessories, Version of Apr. 2003, (6 pages).

Excerpt of Operator's Manual of Boston Scientific's Maestro 30000 Cardiac Ablation System, Version of Oct. 17, 2005, (4 pages).

Schneider, Peter A., "Endovascular Skills—Guidewire and Catheter Skills for Endovascular Surgery," Second Edition Revised and Expanded, 10 pages, (2003).

Kandarpa, Krishna et al., "Handbook of Interventional Radiologic Procedures", Third Edition, pp. 194-210 (2002).

ThermoCool Irrigated Catheter and Integrated Ablation System, Biosense Webster (2006), 6 pages.

Mount Sinai School of Medicine clinical trial for Impact of Renal Sympathetic Denervation of Chronic Hypertension, Mar. 2013, 11 pages. <http://clinicaltrials.gov/ct2/show/NCT01628198>.

Opposition to European Patent No. EP2092957, Granted Jan. 5, 2011, Date of Opposition Oct. 5, 2011, 26 pages.

Opposition to European Patent No. EP1802370, Granted Jan. 5, 2011, Date of Opposition Oct. 5, 2011, 20 pages.

Opposition to European Patent No. EP2037840, Granted Dec. 7, 2011, Date of Opposition Sep. 7, 2012, 25 pages.

Oz, Mehmet, Pressure Relief, TIME, Jan. 9, 2012, 2 pages. <www.time.com/time/printout/0,8816,2103278,00.html>.

Prochnau, Dirk et al., Catheter-based renal denervation for drug-resistant hypertension by using a standard electrophysiology catheter; Euro Intervention 2012, vol. 7, pp. 1077-1080.

Papademetriou, Vasilios, Renal Sympathetic Denervation for the Treatment of Difficult-to-Control or Resistant Hypertension, Int. Journal of Hypertension, 2011, 8 pages.

Holmes et al., Pulmonary Vein Stenosis Complicating Ablation for Atrial Fibrillation: Clinical Spectrum and Interventional Considerations, JACC: Cardiovascular Interventions, 2: 4, 2009, 10 pages.

Purerfellner, Helmut et al., Incidence, Management, and Outcome in Significant Pulmonary Vein Stenosis Complicating Ablation for Atrial Fibrillation, Am. J. Cardiol, 93, Jun. 1, 2004, 4 pages.

Tsao, Hsuan-Ming, Evaluation of Pulmonary Vein Stenosis after Catheter Ablation of Atrial Fibrillation, Cardiac Electrophysiology Review, 6, 2002, 4 pages.

Wittkamp et al., "Control of radiofrequency lesion size by power regulation," Journal of the American Heart Association, 1989, 80: pp. 962-968.

Zheng et al., "Comparison of the temperature profile and pathological effect at unipolar, bipolar and phased radiofrequency current configurations," Journal of Interventional Cardiac Electrophysiology, 2001, pp. 401-410.

U.S. Appl. No. 60/852,787, filed Oct. 18, 2006, 105 pages.

Pieper et al., "Design and Implementation of a New Computerized System for Intraoperative Cardiac Mapping," Journal of Applied Physiology, 1991, vol. 71, No. 4, pp. 1529-1539.

Remo, Benjamin F. et al., "Safety and Efficacy of Renal Denervation as a Novel Treatment of Ventricular Tachycardia Storm in Patients with Cardiomyopathy," Heart Rhythm, 2014, 11(4), 541-6.


Dodge, et al., "Lumen Diameter of Normal Human Coronary Arteries Influence of Age, Sex, Anatomic Variation, and Left Ventricular Hypertrophy or Dilation", Circulation, 1992, vol. 86 (1), pp. 232-246.


Opposition to European Patent No. 2465470, Granted Oct. 28, 2015, Date of Opposition Jul. 27, 2016, 34 pp.

U.S. Appl. No. 11/363,867, filed Feb. 27, 2006, 70 pp.

U.S. Appl. No. 60/813,589, filed Dec. 29, 2005, 62 pp.

Ureter, <https://en.wikipedia.org/wiki/Ureter>, Jun. 2016, 6 pp.

*Fig. 1*

*Fig. 2*




Fig. 4







Fig. 3

*Fig. 5**Fig. 6**Fig. 7*




Fig. 8




Fig. 9




Fig. 10






Fig. 11

*Fig. 12**Fig. 13*




Fig. 14




Fig. 15




Fig. 16




Fig. 17

METHODS FOR TREATING PATIENTS VIA RENAL NEUROMODULATION

RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 15/476,867, filed Mar. 31, 2017, now U.S. Pat. No. 9,907,611, which is a continuation of U.S. application Ser. No. 15/095,220 filed Apr. 11, 2016, now abandoned, which is a continuation of U.S. application Ser. No. 14/846,480, filed Sep. 4, 2015, now abandoned, which is a continuation of U.S. application Ser. No. 13/617,994 filed Sep. 14, 2012, now abandoned, which is a continuation of U.S. application Ser. No. 13/361,019 filed Jan. 30, 2012, now abandoned, which is a continuation of U.S. application Ser. No. 11/688,178 filed Mar. 19, 2007, now U.S. Pat. No. 8,131,372, which is a continuation of U.S. application Ser. No. 11/144,173, filed Jun. 3, 2005, now U.S. Pat. No. 7,647,115, which is a continuation of U.S. application Ser. No. 10/408,665, filed Apr. 8, 2003, now U.S. Pat. No. 7,162,303, which claims benefit of U.S. Provisional Appl. No. 60/370,190 filed Apr. 8, 2002, and claims benefit of U.S. Provisional Appl. No. 60/415,575, filed Oct. 3, 2002, and claims benefit of U.S. Provisional Appl. No. 60/442,970 filed Jan. 29, 2003.

FIELD OF THE INVENTION

This invention relates to methods and apparatus for treatment of congestive heart failure, chronic renal failure and hypertension by nerve stimulation. In particular, the invention relates to the improvement of these conditions of patients by blocking signals to the renal (kidney) nerve.

BACKGROUND OF THE INVENTION

The Heart Failure Problem:

Congestive Heart Failure (CHF) is a form of heart disease still increasing in frequency. According to the American Heart Association, CHF is the "Disease of the Next Millennium". The number of patients with CHF is expected to grow even more significantly as an increasing number of the "Baby Boomers" reach 50 years of age. CHF is a condition that occurs when the heart becomes damaged and reduces blood flow to the organs of the body. If blood flow decreases sufficiently, kidney function becomes impaired and results in fluid retention, abnormal hormone secretions and increased constriction of blood vessels. These results increase the workload of the heart and further decrease the capacity of the heart to pump blood through the kidney and circulatory system. This reduced capacity further reduces blood flow to the kidney, which in turn further reduces the capacity of the blood. It is believed that the progressively-decreasing perfusion of the kidney is the principal non-cardiac cause perpetuating the downward spiral of the "Vicious Cycle of CHF". Moreover, the fluid overload and associated clinical symptoms resulting from these physiologic changes are predominant causes for excessive hospital admissions, terrible quality of life and overwhelming costs to the health care system due to CHF.

While many different diseases may initially damage the heart, once present, CHF is split into two types: Chronic CHF and Acute (or Decompensated-Chronic) CHF. Chronic Congestive Heart Failure is a longer term, slowly progressive, degenerative disease. Over years, chronic congestive heart failure leads to cardiac insufficiency. Chronic CHF is clinically categorized by the patient's ability to exercise or perform normal activities of daily living (such as defined by

the New York Heart Association Functional Class). Chronic CHF patients are usually managed on an outpatient basis, typically with drugs.

Chronic CHF patients may experience an abrupt, severe deterioration in heart function, termed Acute Congestive Heart Failure, resulting in the inability of the heart to maintain sufficient blood flow and pressure to keep vital organs of the body alive. These acute CHF deteriorations can occur when extra stress (such as an infection or excessive fluid overload) significantly increases the workload on the heart in a stable chronic CHF patient. In contrast to the stepwise downward progression of chronic CHF, a patient suffering acute CHF may deteriorate from even the earliest stages of CHF to severe hemodynamic collapse. In addition, Acute CHF can occur within hours or days following an Acute Myocardial Infarction (AMI), which is a sudden, irreversible injury to the heart muscle, commonly referred to as a heart attack.

Normal Kidney Function:

The kidneys are a pair of organs that lie in the back of the abdomen on each side of the vertebral column. Kidneys play an important regulatory role in maintaining the homeostatic balance of the body. The kidneys function like a complex chemical plant. The kidneys eliminate foreign chemicals from the body, regulate inorganic substances and the extracellular fluid, and function as endocrine glands, secreting hormonal substances like renin and erythropoietin.

The main functions of the kidney are to maintain the water balance of the body and control metabolic homeostasis. Healthy kidneys regulate the amount of fluid in the body by making the urine more or less concentrated, thus either reabsorbing or excreting more fluid, respectively. In case of renal disease, some normal and important physiological functions become detrimental to the patient's health. This process is called overcompensation. In the case of Chronic Renal Failure (CRF) patients overcompensation often manifests in hypertension (pathologically high blood pressure) that is damaging to heart and blood vessels and can result in a stroke or death.

The functions of the kidney can be summarized under three broad categories: a) filtering blood and excreting waste products generated by the body's metabolism; b) regulating salt, water, electrolyte and acid-base balance; and c) secreting hormones to maintain vital organ blood flow. Without properly functioning kidneys, a patient will suffer water retention, reduced urine flow and an accumulation of wastes toxins in the blood and body.

The primary functional unit of the kidneys that is involved in urine formation is called the "nephron". Each kidney consists of about one million nephrons. The nephron is made up of a glomerulus and its tubules, which can be separated into a number of sections: the proximal tubule, the medullary loop (loop of Henle), and the distal tubule. Each nephron is surrounded by different types of cells that have the ability to secrete several substances and hormones (such as renin and erythropoietin). Urine is formed as a result of a complex process starting with the filtration of plasma water from blood into the glomerulus. The walls of the glomerulus are freely permeable to water and small molecules but almost impermeable to proteins and large molecules. Thus, in a healthy kidney, the filtrate is virtually free of protein and has no cellular elements. The filtered fluid that eventually becomes urine flows through the tubules. The final chemical composition of the urine is determined by the secretion into and reabsorption of substances from the urine required to maintain homeostasis.

Receiving about 20% of cardiac output, the two kidneys filter about 125 ml of plasma water per minute. This is called the Glomerular Filtration Rate (GFR) and is the gold standard measurement of the kidney function. Since measurement of GFR is very cumbersome and expensive, clinically, the serum creatinine level or creatinine clearance are used as surrogates to measure kidney function. Filtration occurs because of a pressure gradient across the glomerular membrane. The pressure in the arteries of the kidney pushes plasma water into the glomerulus causing filtration. To keep the GFR relatively constant, pressure in the glomerulus is held constant by the constriction or dilatation of the afferent and efferent arterioles, the muscular walled vessels leading to and from each glomerulus.

Abnormal Kidney Function in CHF:

The kidneys maintain the water balance of the body and control metabolic homeostasis. The kidneys regulate the amount of fluid in the body by making the urine more or less concentrated, thus either reabsorbing or excreting more fluid, respectively. Without properly functioning kidneys, a patient will suffer water retention, reduced urine flow and an accumulation of wastes toxins in the blood and body. These conditions resulting from reduced renal function or renal failure (kidney failure) are believed to increase the workload of the heart. In a CHF patient, renal failure will cause the heart to further deteriorate as the water build-up and blood toxins accumulate due to the poorly functioning kidneys and in turn, cause the heart further harm.

In a CHF patient, for any of the known cause of heart dysfunction, the heart will progressively fail and blood flow and pressure will drop in the patients circulatory system. In the acute heart failure, the short-term compensations serve to maintain perfusion to critical organs, notably the brain and the heart that cannot survive prolonged reduction in blood flow. In chronic heart failure, these same responses that initially aided survival in acute heart failure can become deleterious.

A combination of complex mechanisms contribute to the deleterious fluid overload in CHF. As the heart fails and blood pressure drops, the kidneys cannot function owing to insufficient blood pressure for perfusion and become impaired. This impairment in renal function ultimately leads to a decrease in urine output. Without sufficient urine output, the body retains fluids and the resulting fluid overload causes peripheral edema (swelling of the legs), shortness of breath (from fluid in the lungs), and fluid in the abdomen, among other undesirable conditions in the patient.

In addition, the decrease in cardiac output leads to reduced renal blood flow, increased neurohormonal stimulus, and release of the hormone renin from the juxtaglomerular apparatus of the kidney. This results in avid retention of sodium and thus volume expansion. Increased rennin results in the formation of angiotensin, a potent vasoconstrictor.

Heart failure and the resulting reduction in blood pressure reduces the blood flow and perfusion pressure through organs in the body, other than the kidneys. As they suffer reduced blood pressure, these organs may become hypoxic causing the development of a metabolic acidosis which reduces the effectiveness of pharmacological therapy as well as increases the risk of sudden death.

This spiral of deterioration that physicians observe in heart failure patients is believed to be mediated, in large part, by activation of a subtle interaction between heart function and kidney function, known as the renin-angiotensin system. Disturbances in the heart's pumping function results in decreased cardiac output and diminished blood flow. The

kidneys respond to the diminished blood flow as though the total blood volume was decreased, when in fact the measured volume is normal or even increased. This leads to fluid retention by the kidneys and formation of edema causing fluid overload and increased stress on the heart.

Systemically, CHF is associated with an abnormally elevated peripheral vascular resistance and is dominated by alterations of the circulation resulting from an intense disturbance of sympathetic nervous system function. Increased activity of the sympathetic nervous system promotes a downward vicious cycle of increased arterial vasoconstriction (increased resistance of vessels to blood flow) followed by a further reduction of cardiac output, causing even more diminished blood flow to the vital organs.

In CHF via the previously explained mechanism of vasoconstriction, the heart and circulatory system dramatically reduces blood flow to kidneys. During CHF, the kidneys receive a command from higher neural centers via neural pathways and hormonal messengers to retain fluid and sodium in the body. In response, to stress on the heart, the neural centers command the kidneys to reduce their filtering functions. While in the short term, these commands can be beneficial, if these commands continue over hours and days they can jeopardize the person's life or make the person dependent on artificial kidney for life by causing the kidneys to cease functioning.

When the kidneys do not fully filter the blood, a huge amount of fluid is retained in the body resulting in bloating (fluid in tissues), and increases the workload of the heart. Fluid can penetrate into the lungs and the patient becomes short of breath. This odd and self-destructive phenomenon is most likely explained by the effects of normal compensatory mechanisms of the body that improperly perceive the chronically low blood pressure of CHF as a sign of temporary disturbance such as bleeding.

In an acute situation, the organism tries to protect its most vital organs, the brain and the heart, from the hazards of oxygen deprivation. Commands are issued via neural and hormonal pathways and messengers. These commands are directed toward the goal of maintaining blood pressure to the brain and heart, which are treated by the body as the most vital organs. The brain and heart cannot sustain low perfusion for any substantial period of time. A stroke or a cardiac arrest will result if the blood pressure to these organs is reduced to unacceptable levels. Other organs, such as kidneys, can withstand somewhat longer periods of ischemia without suffering long-term damage. Accordingly, the body sacrifices blood supply to these other organs in favor of the brain and the heart.

The hemodynamic impairment resulting from CHF activates several neurohormonal systems, such as the renin-angiotensin and aldosterone system, sympatho-adrenal system and vasopressin release. As the kidneys suffer from increased renal vasoconstriction, the filtering rate (GFR) of the blood drops and the sodium load in the circulatory system increases. Simultaneously, more renin is liberated from the juxtaglomerular of the kidney. The combined effects of reduced kidney functioning include reduced glomerular sodium load, an aldosterone-mediated increase in tubular reabsorption of sodium, and retention in the body of sodium and water. These effects lead to several signs and symptoms of the CHF condition, including an enlarged heart, increased systolic wall stress, an increased myocardial oxygen demand, and the formation of edema on the basis of fluid and sodium retention in the kidney. Accordingly,

sustained reduction in renal blood flow and vasoconstriction is directly responsible for causing the fluid retention associated with CHF.

In view of the physiologic mechanisms described above it is positively established that the abnormal activity of the kidney is a principal non-cardiac cause of a progressive condition in a patient suffering from CHF.

Growing population of late stage CHF patients is an increasing concern for the society. The disease is progressive, and as of now, not curable. The limitations of drug therapy and its inability to reverse or even arrest the deterioration of CHF patients are clear. Surgical therapies are effective in some cases, but limited to the end-stage patient population because of the associated risk and cost. There is clearly a need for a new treatment that will overcome limitations of drug therapy but will be less invasive and costly than heart transplantation.

Similar condition existed several decades ago in the area of cardiac arrhythmias. Limitations of anti-arrhythmic drugs were overcome by the invention of heart pacemakers. Widespread use of implantable electric pacemakers resulted in prolonged productive life for millions of cardiac patients. So far, all medical devices proposed for the treatment of CHF are cardio-centric i.e., focus on the improvement of the heart function. The dramatic role played by kidneys in the deterioration of CHF patients has been overlooked by the medical device industry.

Neural Control of Kidneys:

The autonomic nervous system is recognized as an important pathway for control signals that are responsible for the regulation of body functions critical for maintaining vascular fluid balance and blood pressure. The autonomic nervous system conducts information in the form of signals from the body's biologic sensors such as baroreceptors (responding to pressure and volume of blood) and chemoreceptors (responding to chemical composition of blood) to the central nervous system via its sensory fibers. It also conducts command signals from the central nervous system that control the various innervated components of the vascular system via its motor fibers.

Experience with human kidney transplantation provided early evidence of the role of the nervous system in the kidney function. It was noted that after the transplant, when all the kidney nerves are totally severed, the kidney increased the excretion of water and sodium. This phenomenon was also observed in animals when the renal nerves were cut or chemically destroyed. The phenomenon was called "denervation diuresis" since the denervation acted on a kidney similar to a diuretic medication. Later the "denervation diuresis" was found to be associated with the vasodilatation the renal arterial system that led to the increase of the blood flow through the kidney. This observation was confirmed by the observation in animals that reducing blood pressure supplying the kidney could reverse the "denervation diuresis".

It was also observed that after several months passed after the transplant surgery in successful cases, the "denervation diuresis" in transplant recipients stopped and the kidney function returned to normal. Originally it was believed that the "renal diuresis" is a transient phenomenon and that the nerves conducting signals from the central nervous system to the kidney are not essential for the kidney function. Later, new discoveries led to the different explanation. It is believed now that the renal nerves have a profound ability to regenerate and the reversal of the "denervation diuresis" shall be attributed to the growth of the new nerve fibers supplying kidneys with the necessary stimuli.

Another body of research that is of particular importance for this application was conducted in the period of 1964-1969 and focused on the role of the neural control of secretion of the hormone renin by the kidney. As was discussed previously, renin is a hormone responsible for the "vicious cycle" of vasoconstriction and water and sodium retention in heart failure patients. It was demonstrated that increase (renal nerve stimulation) or decrease (renal nerve denervation) in renal sympathetic nerve activity produced parallel increases and decreases in the renin secretion rate by the kidney, respectively.

In summary, it is known from clinical experience and the large body of animal research that the stimulation of the renal nerve leads to the vasoconstriction of blood vessels supplying the kidney, decreased renal blood flow, decreased removal of water and sodium from the body and increased renin secretion. These observations closely resemble the physiologic landscape of the deleterious effects of the chronic congestive heart failure. It is also known that the reduction of the sympathetic renal nerve activity, achieved by denervation, can reverse these processes.

It was established in animal models that the heart failure condition results in the abnormally high sympathetic stimulation of the kidney. This phenomenon was traced back to the sensory nerves conducting signals from baroreceptors to the central nervous system. Baroreceptors are the biologic sensors sensitive to blood pressure. They are present in the different locations of the vascular system. Powerful relationship exists between the baroreceptors in the carotid arteries (supplying brain with arterial blood) and the sympathetic nervous stimulus to the kidneys. When the arterial blood pressure was suddenly reduced in experimental animals with heart failure, the sympathetic tone increased. Nevertheless the normal baroreflex alone, cannot be responsible for the elevated renal nerve activity in chronic CHF patients. If exposed to the reduced level of arterial pressure for a prolonged time baroreceptors normally "reset" i.e. return to the baseline level of activity until a new disturbance is introduced. Therefore, in chronic CHF patients the components of the autonomic nervous system responsible for the control of blood pressure and the neural control of the kidney function become abnormal. The exact mechanisms that cause this abnormality are not fully understood but, its effects on the overall condition of the CHF patients are profoundly negative.

End Stage Renal Disease Problem:

There is a dramatic increase in patients with end-stage renal disease (ESRD) due to diabetic nephropathy, chronic glomerulonephritis and uncontrolled hypertension. In the US alone, 372,000 patients required dialysis in the year 2000. There were 90,000 new cases of ESRD in 1999 with the number of patients on dialysis is expected to rise to 650,000 by the year 2010. The trends in Europe and Japan are forecasted to follow a similar path. Mortality in patients with ESRD remains 10-20 times higher than that in the general population. Annual Medicare patient costs \$52,868 for dialysis and \$18,496 for transplantation. The total cost for Medicare patients with ESRD in 1998 was \$12.04 billion.

The primary cause of these problems is the slow relentless progression of Chronic Renal Failure (CRF) to ESRD. CRF represents a critical period in the evolution of ESRD. The signs and symptoms of CRF are initially minor, but over the course of 2-5 years, become progressive and irreversible. Until the 1980's, there were no therapies that could significantly slow the progression of CRF to ESRD. While some progress has been made in combating the progression to and

complications of ESRD in last two decades, the clinical benefits of existing interventions remain limited with no new drug or device therapies on the horizon.

Progression of Chronic Renal Failure:

It has been known for several decades that renal diseases of diverse etiology (hypotension, infection, trauma, autoimmune disease, etc.) can lead to the syndrome of CRF characterized by systemic hypertension, proteinuria (excess protein filtered from the blood into the urine) and a progressive decline in GFR ultimately resulting in ESRD. These observations suggested that CRF progresses via a common pathway of mechanisms, and that therapeutic interventions inhibiting this common pathway may be successful in slowing the rate of progression of CRF irrespective of the initiating cause.

To start the vicious cycle of CRF, an initial insult to the kidney causes loss of some nephrons. To maintain normal GFR, there is an activation of compensatory renal and systemic mechanisms resulting in a state of hyperfiltration in the remaining nephrons. Eventually, however, the increasing numbers of nephrons "overworked" and damaged by hyperfiltration are lost. At some point, a sufficient number of nephrons are lost so that normal GFR can no longer be maintained. These pathologic changes of CRF produce worsening systemic hypertension, thus high glomerular pressure and increased hyperfiltration. Increased glomerular hyperfiltration and permeability in CRF pushes an increased amount of protein from the blood, across the glomerulus and into the renal tubules. This protein is directly toxic to the tubules and leads to further loss of nephrons, increasing the rate of progression of CRF. This vicious cycle of CRF continues as the GFR drops, with loss of additional nephrons leading to further hyperfiltration and eventually to ESRD requiring dialysis. Clinically, hypertension and excess protein filtration have been shown to be two major determining factors in the rate of progression of CRF to ESRD.

Though previously clinically known, it was not until the 1980s that the physiologic link between hypertension, proteinuria, nephron loss and CRF was identified. In 1990s the role of sympathetic nervous system activity was elucidated. Afferent signals arising from the damaged kidneys due to the activation of mechanoreceptors and chemoreceptors stimulate areas of the brain responsible for blood pressure control. In response brain increases sympathetic stimulation on the systemic level resulting in the increased blood pressure primarily through vasoconstriction of blood vessels.

When elevated sympathetic stimulation reaches the kidney via the efferent sympathetic nerve fibers, it produces major deleterious effects in two forms:

A. Kidney is damaged by direct renal toxicity from the release of sympathetic neurotransmitters (such as norepinephrine) in the kidney independent of the hypertension.

B. Secretion of renin that activates Angiotensin II is increased leading to the increased systemic vasoconstriction and exacerbated hypertension.

Over time damage to the kidney leads to further increase of afferent sympathetic signals from the kidney to the brain. Elevated Angiotensin II further facilitates internal renal release of neurotransmitters. The feedback loop is therefore closed accelerating the deterioration of the kidney.

BRIEF DESCRIPTION OF THE INVENTION

A treatment of heart failure, renal failure and hypertension has been developed to arrest or slow down the progression of the disease. This treatment is expected to delay the morbid conditions and death often suffered by CHF patients and to

delay the need for dialysis in renal failure. This treatment is expected to control hypertension in patients that do not respond to drugs or require multiple drugs.

The treatment includes a device and method that reduces the abnormally elevated sympathetic nerve signals that contribute to the progression of heart and renal disease. The desired treatment should be implemented while preserving a patient's mobility and quality of life without the risk of major surgery.

The treatment breaks with tradition and proposes a counterintuitive novel method and apparatus of treating heart failure, renal failure and hypertension by electrically or chemically modulating the nerves of the kidney. Elevated nerve signals to and from the kidney are a common pathway of the progression of these chronic conditions.

Chronic heart and renal failure is treated by reducing the sympathetic efferent or afferent nerve activity of the kidney. Efferent nerves (as opposed to afferent) are the nerves leading from the central nervous system to the organ, in this case to the kidney. Sympathetic nervous system (as opposed to parasympathetic) is the part of the autonomic nervous system that is concerned especially with preparing the body to react to situations of stress or emergency that tends to depress secretion, decrease the tone and contractility of smooth muscle, and increase heart rate. In the case of renal sympathetic activity, it is manifested in the inhibition of the production of urine and excretion of sodium. It also elevates the secretion of renin that triggers vasoconstriction. This mechanism is best illustrated by the response of the body to severe bleeding. When in experimental animals, the blood pressure is artificially reduced by bleeding, and the sympathetic inhibition of the kidney is increased to maintain blood pressure with an ultimate goal of preserving the brain from hypotension. The resulting vasoconstriction and fluid retention work in synchrony to help the body to maintain homeostasis.

Efferent renal nerve activity is considered postganglionic, autonomic and exclusively sympathetic. In general, efferent sympathetic nerves can cause a variety of responses in the innervated organs. Studies of sympathetic renal nerves show that they have a strong tendency to behave as a uniform population that acts as vasoconstrictors. The renal postganglionic neurons are modulated by preganglionic (ganglion is a "knot" or agglomeration of nerve cells) nerves that originate from the brain and thoracic and upper lumbar regions of the spinal cord.

The preganglionic nerves have diverse function and are likely to have high degree of redundancy. Although different pathways exist to achieve reduced efferent renal nerve activity, the simplest way is to denervate the postganglionic nerves with an electric stimulus or a chemical agent. The same desired affect could be achieved by total surgical, electric or chemical destruction (ablation) of the nerve. For two reasons this is not a preferred pathway. As was described before, renal nerves regenerate and can grow back as soon as several months after surgery. Secondly, total irreversible denervation of the kidney can result in danger to the patient. Overdiuresis or removal of excess water from blood can result in the reduction of blood volume beyond the amount that can be rapidly replaced by fluid intake. This can result in hypovolemia and hypotension. Hypotension is especially dangerous in heart failure patients with the reduced capacity of the heart to pump blood and maintain blood pressure. In addition, the vasodilation of the renal artery resulting from the renal denervation will cause a significant increase in renal blood flow. In a healthy person, renal blood flow can amount to as much as 20% of the total

cardiac output. In heart failure patients cardiac output is reduced and the renal denervation can “steal” even larger fraction of it from circulation. This, in turn, can lead to hypotension. Also, in a heart failure patient the heart has limited ability to keep up with the demand for oxygenated blood that can be caused by even modest physical effort. Therefore a heart failure patient that can sustain the increased blood flow to the kidneys while at rest can face serious complications resulting from acute hypotension, if the demand for blood flow is increased by temperature change or exercise.

In view of the factors described above it is desired to have means to reduce the efferent sympathetic stimulation of the kidney in CHF patients in a reversible, controlled fashion preferably based on a physiologic feedback signal that is indicative of the oxygen demand by the body, blood pressure, cardiac output of the patient or a combination of these and other physiologic parameters.

The treatment also breaks with tradition and proposes a counterintuitive novel method and apparatus of treating chronic renal failure (CRF) with the goal of slowing down the progression of CRF to the ESRD by electrically or chemically altering the sympathetic neural stimulation entering and exiting the kidney. The described method and apparatus can be also used to treat hypertension in patients with renal disease or abnormal renal function.

To control the afferent nerve signals from the kidney to the brain and block efferent nerve stimuli from entering the kidney (without systemic side effects of drug therapy), a renal nerve stimulator is implanted and attached to an electrode lead placed around or close to the renal artery. Stimulation effectively blocks or significantly reduces both efferent and afferent signals traveling between the kidney, the autonomic nervous system and the central nervous system.

The benefits that may be possible by controlling renal nerve signals to reduce efferent overstimulation are:

a. The secretion of renin by kidney should be reduced by 40-50% translating into the proportionate reduction of systemic angiotensin II, resulting in the reduction of blood pressure in all hypertensive patients including patients refractory to drugs.

b. Similar to renoprotective mechanisms of ACE-I, the reduction of angiotensin II should result in slowed progression of intrarenal changes in glomerular structure and function independent of blood pressure control.

c. Similar to the effects of moxonidine, reduced efferent overstimulation should reduce damage by direct renal toxicity from the release of sympathetic neurotransmitters.

Following the reduction of the afferent sympathetic renal feedback to the brain, there is expected to be a marked reduction in the systemic efferent overstimulation. This will translate into the systemic vasodilation and reduction of hypertension independent of the renin-angiotensin II mechanism.

Renal nerve stimulation in hypertensive CRF patients is unlikely to cause clinically relevant episodes of hypotension. Systemic blood pressure is tightly controlled by feedbacks from baroreceptors in aorta and carotid sinuses. These mechanisms are likely to take over if the blood pressure becomes too low. In polycystic kidney disease (PKD) patients who underwent surgery for total denervation of kidneys, denervation resolved hypertension without postoperative episodes of hypotension.

Technique for Nerve Modulation

Nerve activity can be reversibly modulated in several different ways. Nerves can be stimulated with electric cur-

rent or chemicals that enhance or inhibit neurotransmission. In the case of electrical stimulation, a stimulator containing a power source is typically connected to the nerve by wires or leads. Leads can terminate in electrodes, cuffs that enclose the nerve or in conductive anchors (screws or hooks) that are embedded in tissue. In the later case, the lead is designed to generate sufficient electric field to alter or induce current in the nerve without physically contacting it. The electrodes or leads can be bipolar or unipolar. There are permanent leads that are implanted for months and years to treat a chronic condition and temporary leads used to support the patient during an acute stage of the disease. The engineering aspects of design and manufacturing of nerve stimulators, pacemakers, leads, anchors and nerve cuffs are well known.

Proposed clinical applications of nerve stimulation include: Depression, Anxiety, Alzheimer’s Disease, Obesity, and others. In all existing clinical applications except pain control, the targeted nerves are stimulated to increase the intensity of the transmitted signal. To achieve relief of hypertension and CRF signal traffic traveling to and from the kidney via renal nerves needs to be reduced. This can be achieved by known methods previously used in physiologic studies on animals. A nerve can be paced with electric pulses at high rate or at voltage that substantially exceed normal traffic. As a result, a nerve will be “overpaced”, run out of neurotransmitter substance and transmit less stimulus to the kidney. Alternatively relatively high voltage potential can be applied to the nerve to create a blockade. This method is known as “voltage clamping” of a nerve. Infusion of a small dose of a local anesthetic in the vicinity of the nerve will produce the same effect.

Ablation of conductive tissue pathways is another commonly used technique to control arterial or ventricular tachycardia of the heart. Ablation can be performed by introduction of a catheter into the venous system in close proximity of the sympathetic renal nerve subsequent ablation of the tissue. Catheter based ablation devices were previously used to stop electric stimulation of nerves by heating nerve tissue with RF energy that can be delivered by a system of electrodes. RF energy thus delivered stops the nerve conduction. U.S. Pat. No. 6,292,695 describes in detail a method and apparatus for transvascular treatment of tachycardia and fibrillation with nerve stimulation and ablation. Similar catheter based apparatus can be used to ablate the renal nerve with an intent to treat CRF. The method described in this invention is applicable to irreversible ablation of the renal nerve by electric energy, cold, or chemical agents such as phenol or alcohol.

Thermal means may be used to cool the renal nerve and adjacent tissue to reduce the sympathetic nerve stimulation of the kidney. Specifically, the renal nerve signals may be dampened by either directly cooling the renal nerve or the kidney, to reduce their sensitivity, metabolic activity and function, or by cooling the surrounding tissue. An example of this approach is to use the cooling effect of the Peltier device. Specifically, the thermal transfer junction may be positioned adjacent the vascular wall or a renal artery to provide a cooling effect. The cooling effect may be used to dampen signals generated by the kidney. Another example of this approach is to use the fluid delivery device to deliver a cool or cold fluid (e.g. saline).

BRIEF DESCRIPTION OF THE DRAWINGS

A preferred embodiment and best mode of the invention is illustrated in the attached drawings that are described as follows:

FIG. 1 illustrates the role of sympathetic renal nerve stimulation in congestive heart failure (CHF).

FIG. 2 illustrates the preferred implanted electrostimulation embodiment of the present invention.

FIG. 3 illustrates stimulation of renal nerves across the wall of the renal vein.

FIG. 4 illustrates the drug infusion blocking embodiment with an implanted drug pump.

FIG. 5 illustrates the arterial pressure based control algorithm for renal nerve modulation.

FIG. 6 illustrates electrostimulation of the renal nerve with an anodal block.

FIG. 7 illustrates different nerve fibers in a nerve bundle trunk.

FIG. 8 illustrates renal nerve modulation by blocking electric signals at one point and stimulating the nerve at a different point.

FIG. 9 illustrates transvenous stimulation of the renal nerve with electric field.

FIG. 10 illustrates an embodiment where the stimulation lead is placed using laparoscopic surgery.

FIG. 11 illustrates a patient controlled stimulation embodiment.

FIG. 12 illustrates the progression of CRF to ESRD.

FIG. 13 illustrates the physiologic mechanisms of CRF.

FIG. 14 illustrates stimulation of renal nerves in a patient with an implanted stimulator with a renal artery cuff electrode.

FIG. 15 illustrates the placement of a stimulation cuff on a renal artery end nerve plexus.

FIG. 16 illustrates the design of the cuff electrode that wraps around an artery.

FIG. 17 illustrates the interface between cuff electrodes and the renal artery surface.

DETAILED DESCRIPTION OF THE INVENTION

A method and apparatus has been developed to regulate sympathetic nerve activity to the kidney to improve a patient's renal function and overall condition, and ultimately to arrest or reverse the vicious cycle of CHF disease.

FIG. 1 illustrates the role of sympathetic renal nerves in heart failure. Neural pathways are indicated by solid lines, hormones by interrupted lines. Baroreceptors **101** respond to low blood pressure resulting from the reduced ability of the failing heart **103** to pump blood. Unloading of baroreceptors **101** in the left ventricle of the heart **103**, carotid sinus, and aortic arch (not shown) generates afferent neural signals **113** that stimulate cardio-regulatory centers in the brain **102**. This stimulation results in activation of efferent pathways in the sympathetic nervous system **118**. Sympathetic signals are transmitted to the spinal cord **106**, sympathetic ganglia **107** and via the sympathetic efferent renal nerve **109** to the kidney **111**. The increased activity of sympathetic nerves **108** also causes vasoconstriction **110** (increased resistance) of peripheral blood vessels.

In the kidney **111** efferent sympathetic nerve stimulation **109** causes retention of water (reduction of the amount of urine) and retention of sodium **112** an osmotic agent that is responsible for the expansion of blood volume. The sympathetic stimulation of the kidney stimulates the release of hormones renin **105** and angiotensin II. These hormones activate the complex renin-angiotensin-aldosterone system **117** leading to more deleterious hormones causing vasoconstriction **104** and heart damage **116**. The sympathetic stimulation of the hypothalamus of the brain **102** results in the

release of the powerful hormone vasopressin **114** that causes further vasoconstriction of blood vessels. Angiotensin **11** constricts blood vessels and stimulates the release of aldosterone from adrenal gland (not shown). It also increases tubular sodium reabsorption (sodium retention) in the kidney **111** and causes remodeling of cardiac myocytes therefore contributing to the further deterioration of the heart **103** and the kidney **111**.

It can be inferred from the FIG. 1 that the renal efferent sympathetic stimulation in heart failure is caused by low blood pressure and is a primary factor responsible for the most debilitating symptom of heart failure i.e. fluid overload. It also contributes to the progression of the disease. Acting through the volume overload and peripheral vasoconstriction (together increasing load on the heart) it accelerates the enlargement of the left ventricle that in turn results in the deteriorating ability of the heart to pump blood. Drugs used to treat heart failure address these issues separately. Diuretics are used to reduce fluid overload by reducing the reabsorption of sodium and increasing the excretion of water **112**. Vasodilators are used to reduce peripheral vasoconstriction **110** by reducing levels of angiotensin **117**. Inotropic agents are used to increase blood pressure and de-activate the signals from baroreceptors **101**. These drugs have limited affect and ultimately fail to control the progression and debilitating symptoms heart failure. The proposed invention corrects the neurohormonal misbalance in heart failure by directly controlling the sympathetic neural stimulation **109** of the kidney **111**.

FIG. 2 shows a patient **201** suffering from chronic congestive heart failure treated in accordance with the invention. An implantable device **202** is implanted in the patient's body. An implantable device can be an electric device similar to a pacemaker or nerve stimulator or a chemical substance infusion device. Such devices are well known in the field of medicine. Internal mechanism of the implantable device typically includes a battery **203**, an electronic circuit and (in the case of a drug delivery device) a reservoir with medication.

An example of an implantable drug infusion device is the implantable insulin pump system for treatment of diabetes sold as the MiniMed 2007 or the SynchroMed Infusion System used to control chronic pain, both manufactured by Medtronic Inc. The drug used in this embodiment can be a common local anesthetic such as Novocain or Lidocaine or a more long lasting equivalent anesthetic. Alternatively, a nerve toxin such as the botox can be used to block the nerve. An example of an implantable nerve stimulator is the Vagus Nerve Stimulation (VNSTM) with the Cyberonics NeuroCybernetic Prosthesis (NCP[®]) System used for treatment of epilepsy. It is manufactured by Cyberonics Inc. The internal mechanism of the implantable device typically includes a battery, an electronic circuit and (in the case of a drug delivery device), a reservoir with medication. Neurostimulation systems from different manufacturers are virtually identical across application areas, usually varying only in the patterns of stimulating voltage pulses, style or number of electrodes used, and the programmed parameters. The basic implantable system consists of a pacemaker-like titanium case enclosing the power source and microcircuitry that are used to create and regulate the electrical impulses. An extension lead attached to this generator carries the electrical pulses to the electrode lead that is implanted or attached to the nerves or tissues to be stimulated.

The implantable device **202** is equipped with the lead **204** connecting it to the renal nerve **205**. The lead can contain an electric wire system or a catheter for delivery of medication

or both. Renal nerve conducts efferent sympathetic stimulation from the sympathetic trunk **206** to the kidney **208**. Sympathetic trunk is connected to the patient's spinal cord inside the spine **207**. The connection can be located between the kidney **208** and the posterior renal or other renal ganglia (not shown) in the region of the 10.sup.th, 11.sup.th and 12.sup.th thoracic and 1.sup.th lumbar segments of the spine **207**.

The implantable device **202** is also equipped with the sensor lead **209** terminated with the sensor **210**. The sensor can be a pressure sensor or an oxygen saturation sensor. The sensor **210** can be located in the left ventricle of the heart **211**, right atrium of the heart or other cavity of the heart. It can also be located outside of the heart in the aorta **213**, the aortic arch **212** or a carotid artery **214**. If the sensor is a pressure sensor, it is used to supply the device **202** with the information necessary to safely regulate the sympathetic nerve signals to the kidney **208**. A venous blood oxygen saturation signal can be used in a similar way to control the sympathetic nerve traffic based on oxygen demand. The sensor will be placed in the right atrium of the heart or in the vena cava. More than one sensor can be used in combination to supply information to the device. Sensors can be inside the vascular system (blood vessels) or outside of it. For example, a motion sensor can be used to detect activity of the person. Such sensor does not require placement outside the implanted device case and can be integrated inside the sealed case of the device **202** as a part of the internal mechanism.

FIG. 3 shows external renal nerve stimulator apparatus **306** connected to the electrode tip **308** by the catheter **301**. A catheter is inserted via an insertion site **303** into the femoral vein **305** into the vena cava **302** and further into the renal vein **304**. The tip **308** is then brought into the electric contact with the wall of the vein **304**. Hooks or screws, similar to ones used to secure pacemaker leads, can be used to anchor the tip and improve the electric contact. The tip **308** can have one, two or more electrodes integrated in its design. The purpose of the electrodes is to generate the electric field sufficiently strong to influence traffic along the renal nerve **205** stimulating the kidney **208**.

Two potential uses for the embodiment shown on FIG. 3 are the acute short-term stimulation of the renal nerve and the implanted embodiment. For short term treatment, a catheter equipped with electrodes on the tip is positioned in the renal vein. The proximal end of the catheter is left outside of the body and connected to the electro stimulation apparatus. For the implanted application, the catheter is used to position a stimulation lead, which is anchored in the vessel and left in place after the catheter is withdrawn. The lead is then connected to the implantable stimulator that is left in the body and the surgical site is closed. Patients have the benefit of mobility and lower risk of infection with the implanted stimulator-lead system.

Similar to the venous embodiment, an arterial system can be used. Catheter will be introduced via the femoral artery and aorta (not shown) into the renal artery **307**. Arterial catheterization is more dangerous than venous but may achieve superior result by placing stimulation electrode (or electrodes) in close proximity to the renal nerve without surgery.

FIG. 4 shows the use of a drug infusion pump **401** to block or partially block stimulation of the kidney **208** by infiltrating tissue proximal to the renal nerve **205** with a nerve-blocking drug. Pump **401** can be an implanted drug pump. The pump is equipped with a reservoir **403** and an access port (not shown) to refill the reservoir with the drug by

puncturing the skin of the patient and the port septum with an infusion needle. The pump is connected to the infusion catheter **402** that is surgically implanted in the proximity of the renal nerve **205**. The drug used in this embodiment can be a common local anesthetic such as Novocain. If it is desired to block the nerve for a long time after a single bolus drug infusion, a nerve toxin such as botox (botulism toxin) can be used as a nerve-blocking drug. Other suitable nerve desensitizing agents may comprise, for example, tetrodotoxin or other inhibitor of excitable tissues.

FIG. 5 illustrates the use of arterial blood pressure monitoring to modulate the treatment of CHF with renal nerve blocking. The blood pressure is monitored by the computer controlled implanted device **202** (FIG. 2) using the implanted sensor **210**. Alternatively the controlling device can be incorporated in the external nerve stimulator **306** (FIG. 3) and connected to a standard blood pressure measurement device (not shown). The objective of control is to avoid hypotension that can be caused by excessive vasodilation of renal arteries caused by suppression of renal sympathetic stimulus. This may cause the increase of renal blood flow dangerous for the heart failure patient with the limited heart pumping ability. The control algorithm increases or decreases the level of therapy with the goal of maintaining the blood pressure within the safe range. Similarly the oxygen content of venous or arterial blood can be measured and used to control therapy. Reduction of blood oxygen is an indicator of insufficient cardiac output in heart failure patients.

FIG. 6 illustrates the principles of modulating renal nerve signal with an anodal block. Renal nerve **601** conducts efferent sympathetic electric signals in the direction towards the kidney **602**. Renal nerve **601** trunk is enveloped with two conductive cuff type electrodes: the anode **603** is a positive pole and the cathode **604** is a negative pole electrode. It is significant that the anode **603** is downstream of the cathode and closer to the kidney while the cathode is upstream of the anode and closer to the spine where the sympathetic nerve traffic is coming from. The electric current flowing between the electrodes opposes the normal propagation of nerve signals and creates a nerve block. Anode **603** and cathode **604** electrodes are connected to the signal generator (stimulator) **306** with wires **606**. This embodiment has a practical application even if the device for renal nerve signal modulation is implanted surgically. During surgery the renal nerve is exposed and cuffs are placed that overlap the nerve. The wires and the stimulator can be fully implanted at the time of surgery. Alternatively wires or leads can cross the skin and connect to the signal generator outside of the body. An implantable stimulator can be implanted later during a separate surgery or the use of an external stimulator can be continued.

Clinically used spiral cuffs for connecting to a nerve are manufactured by Cyberonics Inc. (Houston, Tex.) that also manufactures a fully implantable nerve stimulator operating on batteries. See also, e.g., U.S. Pat. No. 5,251,643. Various external signal generators suitable for nerve stimulation are available from Grass-Telefactor Astro-Med Product Group (West Warwick, R.I.). Nerve cuff electrodes are well known. See, e.g., U.S. Pat. No. 6,366,815. The principle of the anodal block is based on the observation that close to an anodal electrode contact the propagation of a nerve action potential can be blocked due to hyperpolarization of the fiber membrane. See e.g., U.S. Pat. Nos. 5,814,079 and 5,800,464. If the membrane is sufficiently hyperpolarized, action potentials cannot pass the hyperpolarized zone and are annihilated.

As large diameter fibers need a smaller stimulus for their blocking than do small diameter fibers, a selective blockade of the large fibers is possible. See e.g., U.S. Pat. No. 5,755,750. The activity in different fibers of a nerve in an animal can be selectively blocked by applying direct electric current between an anode and a cathode attached to the nerve.

Antidromic pulse generating wave form for collision blocking is an alternative means of inducing a temporary electric blockade of signals traveling along nerve fibers. See e.g., U.S. Pat. No. 4,608,985. In general, nerve traffic manipulation techniques such as anodal blocking, cathodal blocking and collision blocking are sufficiently well described in scientific literature and are available to an expert in neurology. Most of blocking methods allow sufficient selectivity and reversibility so that the nerve will not be damaged in the process of blocking and that selective and gradual modulation or suppression of traffic in different functional fibers can be achieved.

A nerve is composed of the axons of a large number of individual nerve fibers. A large nerve, such as a renal nerve, may contain thousands of individual nerve fibers, both myelinated and non-myelinated. Practical implementation of physiological blockade of selective nerve fibers in a living organism is illustrated by the paper "Respiratory responses to selective blockade of carotid sinus baroreceptors in the dog" by Francis Hopp. Both anodal block and local anesthesia by injection of bupivacaine (a common long-acting local anaesthetic, used for surgical anaesthesia and acute pain management) were applied to the surgically isolated and exposed but intact nerve leading from baroreceptors (physiologic pressure sensors) in the carotid sinus of the heart to the brain of an animal. Anodal block was induced using simple wire electrodes. Experiments showed that by increasing anodal blocking current from 50 to 350 microamperes signal conduction in C type fibers was gradually reduced from 100% to 0% (complete block) in linear proportion to the strength of the electric current. Similarly increasing concentration of injected bupivacaine (5, 10, 20 and 100 mg/ml) resulted in gradual blocking of the carotid sinus nerve activity in a dog. These experiments confirmed that it is possible to reduce intensity of nerve stimulation (nerve traffic) in an isolated nerve in controllable, reversible and gradual was by the application of electric current or chemical blockade. In the same paper it was described that smaller C type fibers were blocked by lower electric current and higher concentration of bupivacaine than larger C type fibers.

Gerald DiBona in "Neural control of the kidney: functionally specific renal sympathetic nerve fibers" described the structure and role of individual nerve fibers controlling the kidney function. Approximately 96% of sympathetic renal fibers in the renal nerve are slow conducting unmyelinated C type fibers 0.4 to 2.5 micrometers in diameter. Different fibers within this range carry different signals and respond to different levels of stimulation and inhibition. It is known that lower stimulation voltage of the renal nerve created untidiuretic effect (reduced urine output) while higher level of stimulation created vasoconstriction effect. Stimulation threshold is inversely proportional to the fiber diameter; therefore it is likely that elevated signal levels in larger diameter renal nerve C fibers are responsible for the retention of fluid in heart failure. Relatively smaller diameter C fibers are responsible for vasoconstriction resulting in the reduction of renal blood flow in heart failure.

FIG. 7 illustrates a simplified cross-section of the renal nerve trunk **601**. Trunk **601** consists of a number of indi-

vidual fibers. The stimulation electrode cuff **603** envelops the nerve trunk. Larger C type fiber **705** exemplifies fibers responsible for diuresis. There are also other fibers **702** that can be for example afferent fibers. Traffic along these fibers can be blocked by the application of lower blocking voltage or lower dose of anesthetic drug. The resulting effect will be diuresis of the CHF patient (secretion of sodium and water by the kidney) and the relief of fluid overload. Smaller C fiber **704** is responsible for the regulation of renal blood flow.

In clinical practice, it may be desired to modulate or block selectively or preferably the larger fibers **705**. This can be achieved with lower levels of stimulation. The patient can be relieved of excess fluid without significantly increasing renal blood flow since traffic in smaller C fibers will not be altered. Renal blood flow can amount to as much as 20% of cardiac output. In a CHF patient with a weakened heart significant increase of renal blood flow can lead to a dangerous decrease of arterial pressure if the diseased heart fails to pump harder to keep up with an increased demand for oxygenated blood. The nerve stimulator or signal generator **306** therefore is capable of at least two levels of stimulation: first (lower) level to block or partially block signals propagating in larger C fibers that control diuresis, and second (higher) level to block signals propagating in smaller C fibers that control renal vascular resistance and blood flow to the kidney. The later method of nerve traffic modulation with higher electric current levels is useful in preventing damage to kidneys in acute clinical situations where the vasoconstriction can lead to the ischemia of a kidney, acute tubular necrosis (ATN), acute renal failure and sometimes permanent kidney damage. This type of clinical scenario is often associated with the acute heart failure when hypotension (low blood pressure) results from a severe decompensation of a chronic heart failure patient. Acute renal failure caused by low blood flow to the kidneys is the most costly complication in patients with heart failure.

Similar differentiated response to modulation could be elicited by applying different frequency of electric pulses (overpacing) to the renal nerve and keeping the applied voltage constant. DiBona noted that renal fibers responsible for rennin secretion responded to the lowest frequency of pulses (0.5 to 1 Hz), fibers responsible for sodium retention responded to middle range of frequencies (1 to 2 Hz) and fibers responsible for blood flow responded to the highest frequency of stimulation (2 to 5 Hz). This approach can be used when the renal nerve block is achieved by overpacing the renal nerve by applying rapid series of electric pulses to the electrodes with the intent to fatigue the nerve to the point when it stops conducting stimulation pulses.

One embodiment of the method of treating heart failure comprises the following steps:

A. Introducing one or more electrodes in the close proximity with the renal nerve,

B. Connecting the electrodes to an electric stimulator or generator with conductive leads or wires,

C. Initiating flow of electric current to the electrodes sufficient to block or reduce signal traffic in the sympathetic efferent renal nerve fibers with the intention of increasing diuresis, reducing renal secretion of renin and vasodilation of the blood vessels in the kidney to increase renal blood supply.

FIG. 8 shows an alternative embodiment of the invention. In this embodiment the natural efferent signal traffic **804** entering the renal nerve trunk **601** is completely blocked by the anodal block device stimulator **306** using a pair of electrodes **604** and **603**. The third electrode (or pair of

electrodes) **803** is situated downstream of the block. The electrode is used to stimulate or pace the kidney. Stimulation signal is transmitted from the generator **306** via the additional lead wire **805** to the electrode **803**. The induced signal becomes the nerve input to the kidney. This way full control of nerve input is accomplished while the natural sympathetic tone is totally abolished.

FIG. **9** shows the transvenous embodiment of the invention using anodal blockade to modulate renal nerve traffic. Renal nerve **601** is located between the renal artery **901** and the renal vein **902**. It follows the same direction towards the kidney. Renal artery can branch before entering the kidney but in the majority of humans there is only one renal artery. Stimulation catheter or lead **903** is introduced into the renal vein **902** and anchored to the wall of the vein using a securing device **904**. The securing device can be a barb or a screw if the permanent placement of the lead **903** is desired. Electric field **904** is induced by the electric current applied by the positively charged anode **905** and cathode **906** catheter electrodes. Electrodes are connected to the stimulator (not shown) by wires **907** and **908** that can be incorporated into the trunk of the lead **903**. Electric field **904** is induced in the tissue surrounding the renal vein **902** and created the desired local polarization of the segment of the renal nerve trunk **601** situated in the close proximity of the catheter electrodes **905** and **907**. Similarly catheters or leads can be designed that induce a cathodal block, a collision block or fatigue the nerve by rapidly pacing it using an induced field rather than by contacting the nerve directly.

FIG. **10** shows an embodiment where the stimulation lead is placed using laparoscopic surgery. This technology is common in modern surgery and uses a small video-camera and a few customized instruments to perform surgery with minimal tissue injury. The camera and instruments are inserted into the abdomen through small skin cuts allowing the surgeon to explore the whole cavity without the need of making large standard openings dividing skin and muscle.

After the cut is made in the umbilical area a special needle is inserted to start insufflation. A pressure regulated CO₂ insufflator is connected to the needle. After satisfactory insufflation the needle is removed and a trocar is inserted through the previous small wound. This method reduces the recovery time due to its minimal tissue damage permitting the patient to return to normal activity in a shorter period of time. Although this type of procedure is known since the beginning of the 19th. century, it was not until the advent of high resolution video camera that laparoscopic surgery became very popular among surgeons. Kidney surgery including removal of donor kidneys is routinely done using laparoscopic methodology. It should be easy for a skilled surgeon to place the lead **903** through a tunnel in tissue layers **1001** surrounding the renal nerve **601**. This way lead electrodes **905** and **906** are placed in close proximity to the nerve and can be used to induce a block without major surgery.

FIG. **11** shows an implanted embodiment of the invention controlled by the patient from outside of the body. The implanted stimulation device **203** is an electric stimulation device to modulate the renal nerve signal but can be an implantable infusion pump capable of infusing a dose of an anesthetic drug on command. The implantable device **203** incorporates a magnetically activated switch such as a reed relay. The reed switch can be a single-pole, single-throw (SPST) type having normally open contacts and containing two reeds that can be magnetically actuated by an electromagnet, permanent magnet or combination of both. Such switch of extremely small size and low power requirements

suitable for an implanted device is available from Coto Technology of Providence, R.I. in several configurations. Switch is normally open preventing electric or chemical blockade of the renal nerve **209**. When the patient brings a magnet **1101** in close proximity to the body site where the device **202** is implanted the magnetic field **1103** acts on the magnetic switch **1102**. Switch is closed and blocking of the renal nerve is activated. The resulting reduction of the sympathetic tone commands the kidney **208** to increase the production of urine. Patient can use the device when they feel the symptoms of fluid overload to remove excess fluid from the body. The device **202** can be equipped with a timing circuit that is set by the external magnet. After the activation by the magnet the device can stay active (block renal nerve activity) for a predetermined duration of time to allow the kidney to make a desired amount of urine such as for an hour or several hours. Then the device will time out to avoid excessive fluid removal or adaptation of the renal nerve to the new condition.

FIG. **12** illustrates the progression of CRF to ESRD. Following the original injury to the kidney **1201** some nephrons **1202** are lost. Loss of nephrons lead to hyperfiltration **1203** and triggers compensatory mechanisms **1204** that are initially beneficial but over time make injury worse until the ESRD **1208** occurs. Compensatory mechanisms lead to elevated afferent and efferent sympathetic nerve signal level (increased signal traffic) **1207** to and from the kidney. It is the objective of this invention to block, reduce, modulate or otherwise decrease this level of stimulation.

The effect of the invented therapeutic intervention will be the reduction of central (coming from the brain) sympathetic stimulation **1206** to all organs and particularly blood vessels that causes vasoconstriction and elevation of blood pressure. Following that hypertension **1205** will be reduced therefore reducing continuous additional insult to the kidney and other organs.

FIG. **13** illustrates the physiologic mechanisms of CRF and hypertension. Injured kidney **1302** sends elevated afferent nerve **1306** signals to the brain **1301**. Brain in response increases sympathetic efferent signals to the kidney **1307** and to blood vessels **1311** that increase vascular resistance **1303** by vasoconstriction. Vasoconstriction **1303** causes hypertension **1304**. Kidney **1302** secretes renin **1310** that stimulates production of the vasoconstrictor hormone Angiotensin II **1305** that increases vasoconstriction of blood vessels **1303** and further increases hypertension **1304**. Hypertension causes further mechanical damage **1312** to the kidney **1302** while sympathetically activated neurohormones **1307** and angiotensin II causes more subtle injury via the hormonal pathway **1310**.

Invented therapy reduces or eliminates critical pathways of the progressive disease by blocking afferent **1306** and efferent **1307** signals to and from the kidney **1302**. Both neurological **1311** and hormonal **1309** stimulus of vasoconstriction are therefore reduced resulting in the relief of hypertension **1304**. As a result, over time the progression of renal disease is slowed down, kidney function is improved and the possibility of stroke from high blood pressure is reduced.

FIG. **14** shows a patient **201** suffering from CRF or renal hypertension treated in accordance with the invention. An implantable device **202** is implanted in the patient's body. An implantable device can be an electric nerve stimulator or a chemical substance (drug) infusion device. The implantable device **202** described above is equipped with the lead **204** connecting it to the renal nerve artery cuff **1401**. Cuff **1401** envelopes the renal artery **203** that anatomically serves

as a support structure for the renal nerve plexus. It is understood that there exist many varieties of electrode configurations such as wires, rings, needles, anchors, screws, cuffs and hooks that could all potentially be used to stimulate renal nerves. The cuff configuration **1401** illustrated by FIGS. **14**, **15**, **16** and **17** was selected for the preferred embodiment base on the information available to the inventors at the time of invention.

The lead conduit can be alternatively an electric wire or a catheter for delivery of medication or a combination of both. Renal nerve conducts efferent sympathetic stimulation from the sympathetic trunk **206** to the kidney **208**. Sympathetic trunk is connected to the patient's spinal cord inside the spine **207**. The lead to nerve connection can be located anywhere between the kidney **208** and the posterior renal or other renal ganglia (not shown) in the region of the 10.sup.th, 11.sup.th and 12.sup.th thoracic and 1.sup.st lumbar segments of the spine **207**. The stimulation lead **204** and the arterial nerve cuff **1401**, as selected for the preferred embodiment of the invention, can be placed using laparoscopic surgery.

FIG. **15** illustrates one possible embodiment of the renal nerve stimulation cuff electrode cuff. When the treated disease is CRF or hypertension it is the additional objective of this embodiment of the invention to selectively modulate nerve traffic in both afferent and efferent nerve fibers innervating the human kidney. Using existing selective modulation techniques it is possible to stimulate only afferent or efferent fibers. Different types of fibers have different structure and respond to different levels and frequency of stimulation. Anatomically renal nerve is difficult to locate in humans even during surgery. The autonomic nervous system forms a plexus on the external surface renal artery. Fibers contributing to the plexus arise from the celiac ganglion, the lowest splanchnic nerve, the aorticorenal ganglion and aortic plexus. The plexus is distributed with branches of the renal artery to vessels of the kidney, the glomeruli and tubules. The nerves from these sources, fifteen or twenty in number, have a few ganglia developed upon them. They accompany the branches of the renal artery into the kidney; some filaments are distributed to the spermatic plexus and, on the right side, to the inferior vena cava. This makes isolating a renal nerve difficult.

To overcome this anatomic limitation the preferred embodiment of the neurostimulation shown on FIG. **15** has an innovative stimulation cuff. The cuff **1401** envelopes the renal artery **203** and overlaps nerve fibers **1501** that form the renal plexus and look like a spider web. Cuff has at least two isolated electrodes **1402** and **1403** needed for nerve blocking. More electrodes can be used for selective patterns of stimulation and blocking. Electrodes are connected to the lead **204**. Renal artery **203** connects aorta **213** to the kidney **208**. It is subject to pulsations of pressure and therefore cyclically swells and contracts.

FIG. **16** further illustrates the design of the cuff **1401**. Cuff envelopes the renal artery **203**. Cuff is almost circumferential but has an opening **406**. When the artery cyclically swells with blood pressure pulses, the cuff opens up without damaging the nerve or pinching the artery. Opening **406** also allows placement of the cuff around the artery. Similar designs of nerve cuffs known as "helical" cuffs are well known, see e.g., U.S. Pat. Nos. 5,251,634; 4,649,936 and 5,634,462.

FIG. **17** shows the crosssection of the cuff **1401**. Cuff **1401** is made out of dielectric material. Two electrodes **1402** and **1403** form rings to maximize the contact area with the wall of the artery **203**.

Common to all the embodiments, is that an invasive device is used to decrease the level of renal nerve signals that are received by the kidney or generated by the kidney and received by the brain. The invention has been described in connection with the best mode now known to the applicant inventors. The invention is not to be limited to the disclosed embodiment. Rather, the invention covers all of various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

Heart failure, also called congestive heart failure (CHF) and chronic heart failure is a progressive heart disease characterized by low cardiac output, deterioration of heart muscle and fluid retention. Renal failure, also called chronic renal failure (CRF) is a progressive degenerative renal disease that is characterized by gradual loss of renal function that leads to the end stage renal disease (ESRD). ESRD requires dialysis for life. Hypertension is the chronic disease associated with high probability of stroke, renal failure and heart failure that is characterized by the abnormally high blood pressure.

A nerve in the context of this application means a separate nerve or a nerve bundle, nerve fiber, nerve plexus or nerve ganglion. Renal nerve is a part of the autonomic nervous system that forms a plexus on the external surface renal artery. Fibers contributing to the plexus arise from the celiac ganglion, the lowest splanchnic nerve, the aorticorenal ganglion and aortic plexus. The plexus is distributed with branches of the renal artery to blood vessels of the kidney, the glomeruli and tubules. The nerves from these sources, have a few ganglia developed upon them. They accompany the branches of the renal artery into the kidney; some filaments are distributed to the spermatic plexus and, on the right side, to the inferior vena cava.

Nerve stimulation, neurostimulation, nerve modulation and neuromodulation are equivalent and mean altering (reducing or increasing) naturally occurring level of electric signals propagating through the nerve. The electric signal in the nerve is also called nerve traffic, nerve tone or nerve stimulus.

Nerve block, blocking or blockade is a form of neuro-modulation and means the reduction or total termination of the propagation or conduction of the electric signal along the selected nerve. Nerve block can be pharmacological (induced by a drug or other chemical substance) or an electric block by electrostimulation. Electric nerve block can be a hyperpolarization block, cathodal, anodal or collision block. Overpacing a nerve can also induce a block. Overpacing means stimulating the nerve with rapid electric pulses at a rate that exceeds the natural cycling rate of the nerve polarization and depolarization. As a result of overpacing the nerve gets fatigued, reserves of the immediately available neurotransmitter substance in the nerve become exhausted, and the nerve becomes temporarily unable to conduct signals. Nerve block by the means listed above can result in the reduction of the nerve signal, in particular the renal sympathetic efferent or afferent tone that determines the electric stimulus received or generated by the kidney. The technique of the controlled reduction of the nerve signal or traffic, which results in less organ stimulation, is called nerve signal modulation. Nerve modulation means that the individual nerve fibers fire with a reduced frequency or that fewer of the nerve fibers comprising the renal nerve are actively conducting or firing. The increase of nerve traffic or nerve activity usually involves recruitment of larger number of fibers in the nerve; alternatively less stimulation is

21

associated with less active fibers. Denervation means blocking of the renal nerve conduction or the destruction of the renal nerve.

Lead is a medical device used to access the nerve designated for stimulation or blocking. It is usually a tubular device that is electrically insulated and includes multiple conductors or wires. Wires conduct stimulation or blocking signals from the stimulator to the designated nerve. Wires are terminated in electrodes. Electrodes are conductive terminals and can contact the nerve directly or contact the conductive tissue in the vicinity of the nerve. Electrodes can have different geometric configurations and can be made of different materials. The lead can include lumens or tubes for drug delivery to the nerve. A stimulator or an electrostimulator is an electric device used to generate electric signals that are conducted by the lead to the nerve. The stimulator can be implanted in the body or external. Electric signals can be a DC current, voltage, series of pulses or AC current or voltage. Electrodes can induce an electric field that affects the nerve and results in nerve blocking. Nerve cuff is a support structure that at least partially envelops the targeted nerve.

While the invention has been described in connection with what is presently considered to be the most practical and preferred embodiment, it is to be understood that the invention is not to be limited to the disclosed embodiment, but on the contrary, is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

We claim:

1. A method for treatment of a hypertensive human patient via renal denervation, the method comprising:
percutaneously introducing a catheter into a renal vein and proximate to a renal nerve of the patient; and thermally affecting the renal nerve of the patient via an energy transfer device carried by the catheter, wherein thermally affecting the renal nerve results in a therapeutically beneficial reduction in blood pressure of the patient.
2. The method of claim 1 wherein thermally affecting the renal nerve via the energy transfer device comprises reducing neural communication along the renal nerve.
3. The method of claim 1 wherein thermally affecting the renal nerve of the patient via the energy transfer device comprises ablating the renal nerve.
4. The method of claim 1 wherein thermally affecting the renal nerve of the patient via the energy transfer device comprises partially ablating the renal nerve.
5. The method of claim 1 wherein thermally affecting the renal nerve via the energy transfer device comprises reducing afferent neural signals along the renal nerve.
6. The method of claim 1 wherein thermally affecting the renal nerve via the energy transfer device comprises reducing efferent neural signals along the renal nerve.

22

7. The method of claim 1, further comprising monitoring a parameter of the catheter and/or a physiological parameter of the patient before and during thermally affecting the renal nerve.

8. The method of claim 1, further comprising removing the catheter from the patient after therapy to conclude the procedure.

9. The method of claim 1, further comprising anchoring at least one of the electrodes to an inner wall of the renal vein before delivery of the electric field.

10. The method of claim 9 wherein anchoring at least one electrode comprises securing the therapeutic element to the inner wall of the renal vein via a barb.

11. The method of claim 9 wherein anchoring the therapeutic element comprises securing the therapeutic element to the inner wall of the renal vein via a screw.

12. A method of treating a human patient, the method comprising:

delivering a catheter comprising one or more electrodes within a renal vein of the patient and proximate to neural fibers innervating a kidney of the patient; and at least partially ablating the neural fibers via an electric field delivered by the one or more electrodes, wherein at least partially ablating the neural fibers results in a therapeutically beneficial reduction in clinical symptoms of hypertension in the patient.

13. The method of claim 12 wherein delivering a catheter comprising one or more electrodes within a renal vein of the patient comprises positioning the one or more electrodes in contact with an inner wall of the renal vein.

14. The method of claim 12 wherein at least partially ablating the neural fibers via an electric field delivered by the one or more electrodes comprises delivering radiofrequency (RF) energy via the one or more electrodes to the neural fibers.

15. The method of claim 12 wherein at least partially ablating the neural fibers via an electric field delivered by the one or more electrodes comprises at least partially ablating the neural fibers via a monopolar electric field.

16. The method of claim 12 wherein at least partially ablating the neural fibers via an electric field delivered by the one or more electrodes comprises at least partially ablating the neural fibers via a bipolar electric field.

17. The method of claim 12 wherein at least partially ablating the neural fibers comprises blocking neural traffic to and/or from the kidney of the patient.

18. The method of claim 12 wherein at least partially ablating the neural fibers comprises denervating the kidney of the patient.

19. The method of claim 12, further comprising monitoring at least one condition of the patient via a sensor.

20. The method of claim 19, further comprising altering delivery of the electric field based on a monitored condition of the patient.

* * * * *

专利名称(译)	通过肾神经调节治疗患者的方法		
公开(公告)号	US10179028	公开(公告)日	2019-01-15
申请号	US15/878746	申请日	2018-01-24
申请(专利权)人(译)	美敦力公司ARDIAN卢森堡S.A.R.L.		
当前申请(专利权)人(译)	美敦力公司ARDIAN卢森堡S.A.R.L.		
[标]发明人	LEVIN HOWARD R GELFAND MARK		
发明人	LEVIN, HOWARD R. GELFAND, MARK		
IPC分类号	A61B5/00 A61B5/044 A61B5/0452 A61N1/40 A61B18/04 A61N5/00 A61B5/0402 A61B18/14 A61M5/142 A61M5/172 A61N1/05 A61N1/32 A61M1/36 A61N1/362 A61N1/36 A61B18/00		
CPC分类号	A61B18/1492 A61M5/142 A61M5/14276 A61M5/1723 A61N1/05 A61N1/0551 A61N1/326 A61N1/36114 A61N1/36117 A61N1/36125 A61N1/36135 A61N1/40 A61N1/403 A61N5/00 A61B18/04 A61N1/3627 A61B2018/00404 A61B2018/00434 A61B2018/00511 A61B2018/00577 A61M1/3627 A61M2210/1082 A61N1/36007		
优先权	60/442970 2003-01-29 US 60/415575 2002-10-03 US 60/370190 2002-04-08 US		
其他公开文献	US20180228535A1		
外部链接	Espacenet		

摘要(译)

一种通过刺激肾神经来治疗心力衰竭，高血压和肾衰竭的方法和设备。治疗的目的是降低肾神经的交感神经活动。通过药物输注或电刺激至少部分地阻断神经来完成治疗。装置可以永久植入或基于导管。

