

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 September 2008 (18.09.2008)

PCT

(10) International Publication Number
WO 2008/112942 A2

(51) International Patent Classification:

A61B 19/00 (2006.01) *A61B 17/08* (2006.01)

(21) International Application Number:

PCT/US2008/056921

(22) International Filing Date: 13 March 2008 (13.03.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/894,626 13 March 2007 (13.03.2007) US

(71) Applicants and

(72) Inventors: **HARRIS, Peter S.** [US/US]; 604 97th Place S.E., Bellevue, Washington 98004 (US). **RABIN, Barry Hal** [US/US]; 3660 W. 81st South, Idaho Falls, Idaho 83402 (US).

(74) Agents: **SPECKMAN, Ann W.** et al.; Speckman Law Group PLLC, 1201 Third Avenue, Suite 330, Seattle, Washington 98101 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(54) Title: METHODS AND DEVICES FOR REDUCING GASTRIC VOLUME

708




FIG. 7C

(57) Abstract: The present invention involves new interventional methods and devices for reducing gastric volume, and thereby treating obesity. The procedures are generally performed laparoscopically and may generally be described as laparoscopic plication gastroplasty (LPG) in which, after obtaining abdominal access, spaced apart sites on a gastric wall are engaged and approximated to create one or more tissue folds that are then secured to produce one or more plications projecting into the gastrointestinal space. The serosal tissue may optionally be treated during the procedure to promote the formation of a strong serosa-to-serosa bond that ensures the long-term stability of the tissue plication. These procedures are preferably carried out entirely extragastrically (i.e. without penetrating through the gastrointestinal wall), thereby minimizing the risks of serious complications. Methods for reversing the procedure are also disclosed.

WO 2008/112942 A2

5

METHODS AND DEVICES FOR REDUCING GASTRIC VOLUME

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority under 35 U.S.C. §119(e) to U.S. Provisional Patent Application No. 60/894,626 filed March 13, 2007.

10

Field of the Invention

15

The present invention relates generally to methods and devices for reducing the volume of a hollow body organ, such as gastric volume. One application of methods and devices of the present invention is treating obesity in a patient by effectively reducing the functional volume of the stomach.

Background and Description of the Prior Art

20

Obesity is rapidly reaching epidemic proportions in developed societies worldwide. There are currently over 1 billion overweight people globally, with 300 million of these people considered clinically obese. In the United States alone there are more than 50 million obese adults, and the numbers are expected to increase by more than 50% in the next decade. Morbid obesity (i.e. obesity in which there are secondary complications such as hypertension, diabetes, coronary artery disease, stroke, congestive heart failure, orthopedic problems and pulmonary insufficiency) not only affects quality of life, but also shortens life expectancy and costs the health care industry billions of dollars annually.

25

Interventional procedures and associated medical devices for treating morbid obesity in patients are well known in the art. In general, these interventional procedures promote weight loss by either (a) gastric restriction or volume reduction, (b) malabsorption, or (c) a combination of the foregoing. Gastric restriction or volume reduction methods promote weight loss by limiting the amount of food intake (i.e. the patient eats less), either due to physical space limitation or by inducing a feeling of early satiety in the patient. Malabsorption methods promote weight loss by limiting the uptake of nutrients (i.e. the patient digests less of what is eaten), usually by removing or bypassing a portion of the gastrointestinal (GI) tract.

5 Among the earliest interventional procedures directed at promoting weight loss were variations of the jejuno-ileal bypass developed in the 1950s. This surgery effectively bypasses the small intestine and is therefore a strictly malabsorption procedure, which poses serious risks. The bilopancreatic diversion procedure, which combines bypass of most of the
10 small intestine with a partial gastrectomy, is a combined volume reduction and malabsorption procedure that was developed in effort to reduce these risks, but it too had complications and its success was limited.

Roux-en-Y gastric bypass surgery is a commonly performed bariatric procedure, especially in the US. It was originally performed as an open interventional procedure, but it is now routinely performed laparoscopically. This procedure utilizes interventional stapling
15 and cutting devices to form a small stomach pouch, bypassing the lower part of the stomach, and creates a Roux-en-Y limb to attach the jejunum to the pouch. The Roux-en-Y procedure is predominantly a volume reduction method (the stomach pouch is typically ~25 cc in volume), although there is a significant malabsorption component.

Despite the proven efficacy of the Roux-en-Y procedure in terms of achieving weight
20 loss, and the recent laparoscopic improvements that have reduced the associated interventional risks, it remains a highly invasive procedure with substantial rates of morbidity. The rate of interventional mortality may be as high as 1%, and known complications include frequent pulmonary morbidity and anastomotic leaks that can be life threatening. Furthermore, the malabsorption component of the Roux-en-Y procedure can
25 negatively affect health because of reduced vitamin uptake, and the long-term consequences of malabsorption are not yet fully understood.

A variety of other interventional procedures have also been developed involving the use of interventional stapling to bring together and fasten opposing walls of the stomach in order to reduce its volume. Most involve malabsorption to a greater or lesser extent,
30 depending on the procedure. Examples of such procedures include the horizontal gastroplasty (HG) and vertical banded gastroplasty (VBG), as well as more recent variations such as the Magenstrasse and Mill (M&M) and laparoscopic sleeve gastrectomy (LSG) procedures that involve not only stapling, but cutting away and removal of the unused stomach portion, leaving behind a reduced volume tube or sleeve running more or less parallel to the lesser curvature between the esophagus and the pylorus. Surgically inserted

5 artificial sleeves that longitudinally traverse the stomach may achieve similar effective volume reductions while significantly increasing malabsorption. In any case, weight loss results achieved with these procedures may sometimes approach those of the Roux-en-Y, however these procedures are not easily performed, are difficult if not impossible to reverse, and still suffer from risks of serious complications, most frequently related to failure or
10 leakage of the staples, which can lead to dangerous infections and even death.

An alternative minimally invasive procedure recently growing in popularity involves the laparoscopic placement of an adjustable silicone ring around the upper portion of the stomach, thereby creating a small (e.g. 50-120 cc) pouch. The LAP-BAND® is one such commercially available restrictive device that, after placement, induces a feeling of early
15 satiety in the patient. Although considerably less invasive than the Roux-en-Y procedure, and potentially reversible, significantly less weight loss has been observed with laparoscopic banding. This procedure also suffers from a variety of limitations and shortcomings. For example, because the laparoscopic band does not actually reduce the volume of the stomach, some patients report a feeling of nearly constant hunger. Additionally, long-term
20 complications of the laparoscopic banding procedure may include tissue erosion, slippage of the band, infection, or lack of effectiveness, frequently requiring removal of the band after a period of time.

Another less invasive alternative to the above-mentioned procedures is the intragastric balloon. The intragastric balloon is an inflatable device that is deployed within
25 the stomach, thereby displacing a known internal volume. The advantages of this method are that it is minimally invasive, involves no malabsorption component, and requires no stapling, permanent reconfiguration or removal of tissue. While the correlation between apparent stomach volume reduction and weight loss is well established by the intragastric balloon method, the weight loss achieved is typically considerably less than with Roux-en-Y.
30 Furthermore, unless it is surgically fastened to the stomach wall, the balloon is free floating and frequent complications such as obstruction, mucosal erosion, nausea, vomiting and pain have been documented, with the result that intragastric balloons are usually removed within 6 months after initial placement.

In effort to develop even less invasive devices and procedures, more recently there
35 has been considerable interest in various transoral (or transesophageal) endoscopic

5 approaches for reducing stomach volume entirely from within the gastrointestinal lumen, without the need for abdominal incisions. In general, these approaches involve advancing an endoscope down the patient's esophagus and into the stomach, whereby various tools are then used to manipulate and reconfigure the stomach tissue in order to create one or more divisions or internal folds (also known as plications) within the stomach wall. To securely
10 hold the divisions or plications so formed, some form of sutures, staples, anchors, or other similar securing means are placed transesophageally through the stomach walls, and sophisticated endoscopic tools have been developed for such purposes. Tissue approximation and fixation devices for use in endoscopic procedures are described, for example, in U.S. Patent Publications 2004/0215216, 2007/0112364, 2005/0080438. Many
15 other types of endoscopic tissue approximation and fixation devices and fasteners are also known in the art.

While quite promising, endoscopic approaches for reducing stomach have various limitations and shortcomings. For example, they must be performed by highly skilled endoscopic surgeons and involve the use of large, complicated endoscopic devices that
20 require specialized training to deal with the restricted access and small working space. In order to access the stomach internally, devices must be passed down the patient's esophagus, accruing a substantial risk of perforating the esophagus and injuring adjacent organs. In addition, capturing and manipulating the tissue layers and accurately applying the securing means during a transesophageal procedure is not only difficult but also hazardous, due to the
25 significant risk of accidental injury to other organs, bleeding, etc., when piercing (intentionally or accidentally) the stomach wall. Because there is no extragastric visualization in these procedures, there is no advance warning of a developing life threatening situation that may require a rescue operation.

The stomach wall is comprised of four main tissue layers. The mucosal layer is the
30 innermost tissue layer, adjacent a submucosal connective tissue layer. The submucosal connective tissue layer interfaces with the muscularis layer, and the serosal layer covers the exterior (extragastric) surface. Prior art gastric reduction procedures involving tissue reconfiguration from inside the stomach require the placement of sutures, staples, or anchors during surgery to hold the reconfigured tissue in place strongly enough to sustain the tensile
35 loads imposed by normal movement of the stomach wall during ingestion and processing of

5 food. Because the mucosal and submucosal connective tissue layers are relatively weak and prone to elastic stretching during digestion, the securing means generally penetrate the stomach wall to engage at least the muscularis layer. For this reason, the prior art securing means are generally transgastric, passing one or more times completely through the stomach wall.

10 Proper use and placement of fasteners that penetrate the gastric wall is challenging and concentrates significant forces over a small surface area of mucosal tissue, thereby potentially causing the suture, staple or anchor to leak or tear through the tissue, with potentially disastrous consequences. It is well known that the fasteners used in these procedures frequently migrate, dislodge or even completely disappear over time, resulting in
15 partial or complete failure to maintain the gastrointestinal volume reduction, as well as possible complications. These are significant limitations and shortcomings of prior art bariatric procedures involving tissue reconfiguration.

Previously known interventional procedures for treating obesity through gastrointestinal volume reduction or malabsorption thus involve numerous risks, including
20 life-threatening post-operative complications (e.g. internal bleeding, infection), and long-term problems such as diarrhea, vitamin deficiency, electrolytic imbalance, unpredictable or insufficient weight loss, and gastrointestinal reflux disease (GERD). Given the above noted shortcomings, limitations and risks of prior art procedures, it is apparent there remains a need for safe, easy-to-perform and effective interventional procedures for reducing gastric volume,
25 as well as for devices enabling such procedures.

SUMMARY OF THE INVENTION

The methods and devices of the present invention represent a new approach for reducing gastric volume, and thereby treating obesity and other disorders of the
30 gastrointestinal tract, that is safe, effective, and overcomes many shortcomings and limitations of prior art procedures. In general, methods of the present invention involve reconfiguring a portion of the gastrointestinal tract (e.g., stomach wall) from the abdominal space, by contacting external tissue surfaces and drawing them toward one another to form one or more tissue invaginations, then approximating the shoulders of the extragastric tissue
35 forming the invagination to form a tissue fold or plication, and then securing the shoulders of

5 the extragastric tissue forming the plication to maintain a permanent plication. In preferred embodiments, the extragastric tissue is approximated such that external tissue surfaces abut one another to form the tissue plication, which extends into the internal gastric space. One or more plications may be formed to effectively reduce the circumference, and thereby cross-sectional area and volume, of the gastrointestinal lumen. One of the advantages of this
10 procedure is that the gastric volume is reduced without reducing the mucosal surface area involved in digestive absorption. In a preferred embodiment of the present invention, the portion of the gastric tissue that is reconfigured, according to the procedure described above, is the anterior surface or anterior wall of the stomach, which is readily accessible from the intra-abdominal space. In another preferred embodiment of the present invention, which may
15 allow for even greater gastric volume reduction, the portion of the gastric tissue that is reconfigured includes both the anterior surface and posterior surface of the stomach.

The methods of the present invention may be carried out using open interventional procedures, which are useful, for example, to penetrate the abdominal space and obtain access to difficult or remote regions of the abdomen and gastrointestinal tract, such as the
20 stomach. Alternatively, however, abdominal access to the gastrointestinal tract (e.g., stomach) is provided using conventional laparoscopic procedures that involve relatively minimal penetration of the abdominal space. Minimally invasive non-laparoscopic methods may also be used (i.e. wherein access to the abdominal cavity is achieved without establishing a pneumoperitoneum via insufflation) to access the external surface(s) of the
25 gastrointestinal tract. Numerous methods for accessing the internal abdominal space, and for monitoring intra-abdominal interventions (e.g., imaging and visualizing the intra-abdominal space and intervention) are known and may be used in conjunction with methods of the present invention.

According to one embodiment of the present invention, a method for reducing gastric
30 volume comprises obtaining access to an external surface of the gastrointestinal tract (e.g. stomach); invaginating and approximating the wall of the gastrointestinal tract from its external surface to create at least one plication therein; and fastening surfaces of the approximated gastrointestinal wall to one another to secure the plication(s). According to another embodiment, a method for reducing gastric volume comprises obtaining access to an
35 external surface of the gastrointestinal tract (e.g., stomach); invaginating and approximating

- 5 the wall of the gastrointestinal tract from its external surface by drawing external surfaces of the gastrointestinal tract toward one another to form a plication extending into the interior space of the gastrointestinal tract; and fastening the approximated surfaces of the gastrointestinal wall to one another to secure the plication(s). This methodology provides a significant reduction in the internal volume of the gastrointestinal tract (e.g., stomach)
10 without reducing the interior wall surface available for digestion and nutrient absorption.

The exterior serosal layer and adjacent muscularis layers of the gastrointestinal tract have relatively more strength than the submucosal and mucosal layers. In certain embodiments of methods of the present invention wherein external surfaces of the gastrointestinal wall are approximated to form a plication projecting into the internal space of
15 the gastrointestinal tract, fastening of the approximated portions of the gastrointestinal wall is accomplished by penetrating fewer than all of the layers of the gastric wall. In preferred embodiments, fastening of the approximated portions of the gastric wall is accomplished by penetrating at least the thin, tough serosal layer covering the exterior of the gastrointestinal lumen and, optionally, the serosal and muscularis layers, without penetrating the submucosal
20 and mucosal layers of the gastric wall. In these embodiments, the intragastric space is not breached during the procedure, and the mucosal layer of the gastrointestinal tract remains intact. This is advantageous not only because it simplifies the procedure, but also because it avoids a variety of known complications arising from prior art procedures that may result when transgastric methods are employed that puncture, damage or otherwise compromise the
25 mucosa during the intervention. Thus, according to another embodiment, a method for reducing gastric volume comprises obtaining access to an external surface of the gastrointestinal tract (e.g. stomach); invaginating and approximating the wall of the gastrointestinal tract from its external surface to form a plication extending into the interior space of the gastrointestinal tract; and fastening approximated surfaces of the gastrointestinal
30 wall to one another without penetrating all layers of the gastric wall to secure the plication(s). In one embodiment, the surfaces of the gastrointestinal wall are fastened to one another using fasteners that penetrate at least the serosal layer and preferably the serosal and muscularis layers of portions of the gastrointestinal wall forming the plication.

Additional embodiments of methods of the present invention, disclosed in detail
35 below, incorporate additional features for the purpose of improving the safety and

5 effectiveness and/or reducing the complexity and cost of the procedure. For example, in one embodiment of methods of the present invention, immediately prior to, or contemporaneously with the above mentioned invaginating and approximating steps, serosal tissue on surfaces of the gastrointestinal wall that adjoin to form the plication is treated to promote bonding or adhesion of adjoining tissue layers within the plication. In one
10 embodiment, bonding of adjoining tissue layers within the plication is accomplished by disrupting the serosal tissue and promoting a healing response therein. In one preferred embodiment, a serosal tissue treatment that involves serosal tissue disruption and/or promotion of the formation of a serosal-to-serosal bond is provided over substantially the gastrointestinal surface area involved in forming the one or more tissue folds.

15 It is known that serosal tissue is capable forming strong adhesions to itself, or adjacent tissues, following inadvertent disruption of or damage to the serosal tissue that occurs during surgery. Typically, such adhesions are considered an undesirable and sometimes dangerous complication of abdominal surgery, and avoiding inadvertent damage to the serosa to minimize the formation of adhesions is an important goal during abdominal
20 interventions. In contrast, in methods of the present invention, serosal tissue disruption and formation of the consequent adhesions may be optionally and intentionally promoted on targeted surface areas of the gastrointestinal lumen. When combined with the invaginating and approximating methods of the present invention, it has unexpectedly been discovered that serosal adhesions can be used beneficially for the purpose of providing a supplementary
25 or even primary securing means for the gastrointestinal reconfiguration. According to the present invention therefore, serosal tissue on surfaces of the gastrointestinal wall that form the plication may be treated to disrupt the serosal tissue and promote a healing response for the purpose of selectively promoting the formation of a serosa-to-serosa bond across the approximated tissue boundary within the gastrointestinal plication.

30 A strong serosa-to-serosa bond is typically formed after a relatively brief period of time (e.g. approximately 7 days after surgery). Once formed, this serosa-to-serosa bond is sufficiently strong to substantially resist the separation forces generated by the stomach during ingestion and digestion, and ensures the long-term integrity of the plication. The formation of a strong serosa-to-serosa bond in the gastric plication of the present invention
35 significantly improves the durability and lifespan of the plication, and consequently of the

5 gastric reduction, and offers a significant improvement compared to the (solely) mechanical fastening methods used in tissue approximation and plication in the prior art. Thus, in the present invention, the fasteners used during the intervention to initially secure the tissue fold serve as the sole structural support for securing the plication only during the brief healing phase following surgery. Following its formation, the serosa-to-serosa bond may provide the
10 primary structural support for securing the plication, and the fasteners initially placed to secure the plication may be removed, absorbed or, more typically, left in place within the patient to provide additional support for the plication.

In contrast to Roux-en-Y or other gastrectomy procedures involving stapling, it should be pointed out that the method of the present invention does not require cutting,
15 transection, anastomosis, or removal of any gastrointestinal tissues from the body. It is therefore possible that the gastric reduction accomplished during this procedure is interventionally reversible. For example, if at a later date the surgeon/patient elects to reverse the gastric reduction, it is possible to substantially restore the original gastrointestinal configuration using a simple and safe procedure wherein the plication is substantially
20 eliminated by removal of any remaining implanted securing means, followed by dissection of the serosa-to-serosa bond along the original line of tissue approximation, and subsequent localized treatment to prevent further formation of adhesions during post-operative healing.

A variety of novel devices, tools and systems are provided herein that enable a medical professional to engage and approximate soft body tissues during an interventional procedure, more safely and conveniently than possible using the prior art instruments. These inventive devices, tools and systems are useful for, among a variety of other possible interventional purposes, performing gastric reduction procedures by invaginating and approximating the wall of the gastrointestinal tract from its external surface to create at least one plication therein; and fastening surfaces of the approximated gastrointestinal wall to one
30 another to secure the plication(s).

Gastric reduction methods of the present invention are performed in the abdominal cavity and involve contacting and manipulating the gastrointestinal tract from its external surface. The methods are typically accomplished using minimally invasive laparoscopic techniques, and the devices and systems of the present invention are therefore generally
35 intended to be used in connection with laparoscopic techniques. However, any technique

5 that provides access to the intra-abdominal space and, particularly, the exterior surface of the gastrointestinal tract may be used, including natural orifice transluminal endoscopic surgery (NOTES) techniques and other minimally invasive non-laparoscopic techniques.

In one embodiment, a specialized device is provided for carrying out the tissue invagination and approximation steps; another device may optionally be provided for
10 disrupting and/or promoting the bonding of serosal tissue, and yet another device may be provided for securing the tissue plication(s). A device for invaginating and approximating gastric tissue of the present invention preferably comprises a tool having an actuation mechanism (generally on or in proximity to a handle) manipulable by an operator, at least one extendible member, and at least two tissue engagement mechanisms. Tissue engagement
15 mechanisms are generally provided at or in proximity to the distal end(s) of the device or extendible member(s), but may be provided at other locations. In one embodiment, the approximation device comprises at least one tissue engagement mechanism provided in association with a device shaft that is inserted at the site of the intervention, and another tissue engagement mechanism provided in association with an extendible member. In this
20 embodiment, tissue is approximated by engaging tissue at two spaced apart locations using the tissue engagement mechanisms and then moving the extendible member and the device shaft relative to one another to approximate the engaged tissue.

According to another embodiment, the approximation device of the present invention comprises at least one tissue engagement mechanism provided in association with each of at
25 least two extendible members. The extendible members are adjustable by the operator between an insertion (collapsed, pre-deployed) condition, in which they may be inserted into the abdominal space, and an expanded (extended, deployed) condition, in which the associated tissue engagement mechanisms are separated and positioned to engage two portions of tissue spaced apart from one another. The extendible member(s) are also
30 adjustable by the operator, by means of an actuation mechanism, following engagement of the two portions of tissue to draw together, or approximate, the two portions of tissue engaged by the tissue engagement mechanisms. The tissue engagement mechanisms are furthermore manipulable to release engaged tissue, and the extendible members are manipulable to reposition the members in a low profile, collapsed condition for withdrawal
35 of the device from the abdominal space. Thus, in operation, the distal portion of the tissue

5 invagination and approximation device is positioned in the abdominal space; a control feature
is actuated by the operator to adjust the extendible members from a low-profile, collapsed
condition to a desired extended condition; and the tissue engagement mechanisms are
positioned to engage the exterior surface of spaced-apart portions of the gastrointestinal tract
(e.g., stomach); a control feature is actuated by the operator to draw the tissue engagement
10 mechanisms together and approximate the two engaged portions of tissue; the engagement
mechanisms are disengaged from the tissue; and after repeating the above steps any desired
number of times, the extendible members are collapsed and the device is withdrawn from the
abdominal cavity.

In one embodiment, the device for invaginating and approximating gastrointestinal
15 tissue has a selection feature that allows the medical professional to select the degree of
separation of the extendible members in the expanded condition, and thereby select and
control placement of the tissue engagement mechanisms and the overall size of the one or
more tissue folds to provide a desired degree of gastric reduction. In another embodiment, a
20 variety of interchangeable tools may be provided, allowing the operator to select
approximation tools providing the desired placement of tissue engagement mechanisms and,
consequently, the overall size of the tissue fold(s).

Another tissue invagination and approximation device of the present invention
comprises a tool having at least two extendible members adjustable between a collapsed
insertion condition and an extended operating condition, and additionally comprising at least
25 one tissue invagination structure arranged and adjustable along an axis to contact and
invaginate tissue located generally at a midline between the tissue portions engaged by the
tissue engagement mechanisms. The tissue invagination structure is preferably axially
adjustable between a withdrawn insertion condition in which it does not extend substantially
beyond the terminal ends of the extendible members and an invaginating, projected
30 condition, in which the tissue invagination structure projects toward the midline of the tissue
surface engaged by the tissue engagement mechanisms. In one embodiment, the axial
movement of the tissue invagination structure may be coordinated with the extension of the
tissue engagement mechanisms such that, following engagement of two spaced apart portions
of tissue, the tissue invagination structure is extended to contact and invaginate tissue as the
35 approximation members are drawn together to approximate the two spaced apart tissue

5 portions. A selection feature may allow the medical professional to select the degree of extension of the invagination structure, thereby controlling the overall size of the tissue invagination and plication, and providing a desired degree of gastric reduction.

In yet another embodiment, a serosal treatment device may be provided and used separately from or in coordination with the tissue approximation and invagination device. A 10 serosal tissue treatment device, in one embodiment, is adapted to disrupt serosal tissue lying between spaced apart tissue surfaces engaged by the approximating members to promote healing and formation of a serosal-to-serosal bond between serosal tissue surfaces contacting one another in the plication formed during the tissue approximation. The serosal treatment device may utilize one or more mechanical structures, such as a discontinuous or a non-15 smooth surface structure, to disrupt serosal tissue and thereby promote serosal tissue adhesion. Additionally or alternatively, the serosal treatment device may be operated to facilitate application or administration of an agent that promotes serosal tissue disruption and/or healing in serosal-to-serosal bonds, or to administer a tissue bonding agent that promotes serosal-to-serosal tissue bonds. The serosal treatment device may incorporate an 20 alternative modality for serosal tissue treatment, e.g., by application of heat, RF radiation, ultrasound, electromagnetic radiation, or other types of radiating energy. In one embodiment, the serosal tissue treatment device may be integrated with the approximating members and/or the tissue invagination structure, as described more fully below.

A separate tissue securing or fastening device may be provided for fastening the two 25 adjacent portions of approximated tissue to one another to secure the plication. Suitable devices, such as suturing, stapling and other types of mechanical tissue fastening devices are well known in the art. The tissue fastening device, in one embodiment, is a multi-fire device that is capable of administering multiple fasteners, in multiple positions along a line of approximated tissue, without requiring removal from the abdominal space. Various types of 30 fasteners and fastening devices may be used, as described more fully below.

In another embodiment, an integrated device may be provided for carrying out the tissue invagination and approximation steps, and for optionally treating serosal tissue in the invaginated tissue, while a separate device may be provided for securing the tissue plication. This beneficially eliminates the need for at least one laparoscopic incision and trocar during 35 the procedure. In yet another embodiment, a single multi-functional device is provided that

5 comprises tools capable of invaginating and approximating tissue, optionally treating the serosal tissue to promote a healing response, and for securing the tissue fold to produce the plication. In this embodiment, a single minimally invasive laparoscopic device is provided, thereby minimizing the number of trocars needed to complete the procedure. For example, assuming one access port is needed for the video camera and one is needed for a grasper,
10 liver/organ manipulator, dissector, or other tissue manipulation device, the procedure may be completed using only 3 trocars. In another embodiment, the single integrated minimally invasive laparoscopic device may be optionally configured having one or more extra service channels through which the camera and other tissue manipulation devices may be inserted, thereby allowing the entire gastric reduction intervention to be completed using only a single
15 access port. In comparison, 5 or more laparoscopic incisions are commonly needed for the Roux-en-Y procedure. Using a multifunctional tool of the present invention, the gastric reduction procedure is less invasive, requires less time to complete and therefore reduces the risks attendant any intervention, speeds patient recovery, and reduces the overall cost of treatment.

20 Other embodiments of medical devices of the present invention further incorporate novel tool configurations, detailed below, that enable and simplify the steps of securing the one or more tissue folds created in order to produce the one or more plications in the wall of the gastrointestinal tract. In one embodiment, means are provided for delivering individual tissue anchors comprising a securing assembly. In yet another embodiment, individual tissue
25 anchors are reconfigured from a first state (e.g. a configuration used for delivery) to a second state (e.g. a deployed configuration). In yet another embodiment, the deployed securing assembly is configured to penetrate only the serosal and muscularis tissue layers, without penetrating completely through the wall of the gastrointestinal tract.

According to the brief summary provided above, it is apparent that methods and
30 devices of the present invention offer several advantages over the prior art. For example, because the one or more gastric tissue plications produced may achieve substantial therapeutic gastric reductions, it is possible to obtain weight loss results comparable to prior art procedures using an interventional alternative that may be performed using minimally invasive laparoscopic or non-laparoscopic abdominal access procedures, while at the same time avoiding a variety of complications associated with malabsorption, the long-term
35

5 presence of restrictive devices within the body, leakage or failure at transgastric anastomosis or anchoring sites, permanent restructuring of the gastrointestinal tract, and the like. Gastric reduction procedures of the present invention are therefore simpler, easier to perform, and safer than prior art interventional methods. In addition, the methods of the present invention, which may optionally be performed substantially or entirely extragastrically, may be carried
10 out by conventionally skilled laparoscopic surgeons, requiring minimal specialized training to achieve substantial gastric volume reduction and effective weight loss results, while significantly reducing the risk of injury or damage to neighboring organs and other complications. This is a significant advantage compared to prior art transesophageal endoluminal interventional methods.

15 While the present invention will be described more fully hereinafter with reference to the accompanying drawings, in which particular embodiments are shown and explained, it is to be understood that persons skilled in the art may modify the embodiments herein described while achieving the same methods, functions and results. Accordingly, the descriptions that follow are to be understood as illustrative and exemplary of specific structures, aspects and
20 features within the broad scope of the present invention and not as limiting of such broad scope.

BRIEF DESCRIPTION OF THE FIGURES

25 **Figure 1** schematically illustrates an interventional method according to one embodiment of the present invention, pre-procedure (Figures 1A-1A3), and post procedure (Figures 1B-1B2).

Figures 2A-2E2 schematically illustrate an exemplary interventional gastric reduction method according to one embodiment of the present invention.

30 **Figures 3A and 3B** show an organ having a plication and a cross sectional view of a plication, illustrating securing means applied according to one embodiment of the present invention.

Figures 4A and 4B show an organ having two plications and a cross sectional view of the multiple plications according to one embodiment of the present invention.

35 **Figures 5A-5F** illustrate operation of a medical device according to one embodiment of the present invention, wherein Figure 5A shows an overview; Figure 5B shows a close-up, distal end of the device in a collapsed state; Figure 5C shows a close-up, distal end of the

5 device in an extended state; Figure 5D shows the device in an extended state following tissue engagement; Figure 5E illustrates partial retraction of the extendible members and tissue engagement mechanisms and actuation of a projecting serosal tissue treatment member during invagination and approximation; and Figure 5F illustrates complete retraction of the extendible members and full extension of the projecting serosal tissue treatment member to
10 form the plication.

Figures 6A-6D illustrate a medical device system according to one embodiment of the present invention, wherein Figure 6A shows separate tools positioning; Figure 6B shows the tissue fold created; Figure 6C shows the fasteners applied; and Figure 6D shows a plurality of fasteners.

15 **Figures 7A-7H** illustrate a medical device according to one embodiment of the present invention, wherein Figure 7A shows an overview; Figure 7B shows the distal end in collapsed state; Figure 7C shows the distal end in expanded state; Figure 7D shows the tissue engagement; Figure 7E shows the tissue invagination and approximation; Figure 7F shows the tissue fold created; Figure 7G shows the securing means applied, with the distal
20 end retracted to collapsed state; and Figure 7H shows a plurality of securing means.

25 **Figures 8A-8E** illustrate a medical device according to another embodiment of the present invention, wherein Figure 8A shows the distal end in collapsed state with a helical fastener; Figure 8B shows a tissue fold created; Figure 8C shows the fasteners applied and the distal end retracted to collapsed state; and Figure 8D shows a plurality of fasteners applied.

Figures 9A and 9B illustrate an embodiment of the present invention, wherein Figure 9A shows a first tissue fold created and first fastener applied to produce first plication; and Figure 9B shows a second tissue fold created and a second fastener applied producing second plication.

30 **Figure 10** illustrates one embodiment of the present invention in which a plurality of helical fasteners have been applied to secure a tissue fold and thereby produce a plication.

Figure 11 shows another embodiment of the present invention involving articulation of the distal multi-functional tool assembly.

5 Methods of the present invention provide effective reduction of the functional volume
of the gastrointestinal tract (e.g., stomach) using an extragastric gastroplasty procedure. In
this procedure, a portion of the gastrointestinal tract is reconfigured by invaginating and
approximating tissue to form one or more tissue folds, and then securing the one or more
tissue folds in order to produce one or more plications. While the following detailed
10 descriptions refer in general to reducing the functional volume of the gastrointestinal tract,
the stomach in particular, it should be recognized that the invaginaton, approximation and
securing methods of the present invention may be used on other body tissues and for other
interventional purposes, within the scope of the present invention.

15 Gastric reduction procedures of the present invention generally access the
gastrointestinal tract via the abdominal cavity. This is most typically accomplished using
conventional laparoscopic techniques wherein the patient is anesthestetized, one or more
small incisions are made through the abdominal wall, and a pneumoperitoneum is established
by insufflation, thereby allowing the insertion of imaging devices and one or more
interventional instruments through laparoscopic ports, also known as trocars. Alternatively,
20 methods of the present invention may also be carried out when access to the abdominal
cavity and gastrointestinal tract is obtained using even less invasive, non-laparoscopic
techniques. A variety of such non-laparoscopic techniques may be utilized within the scope
of the present invention, typically involving grasping and lifting, or otherwise retracting the
abdominal wall to create sufficient working space within the abdominal cavity, without the
25 need for insufflation. Alternatively, the methods and devices of the present invention may
also be adapted for flexible endoscopic use, allowing access to the abdominal cavity and
external surface of the gastrointestinal tract to be obtained by first entering the body through
a natural orifice (e.g esophagus, anus or vagina), then penetrating through the wall of an
anatomical lumen into the abdominal cavity.

30 Once abdominal access has been obtained, the medical professional employs one or
more cameras or other imaging devices, along with a variety of tools known in the art, to
manipulate the internal organs and/or tissues to expose the region of the gastrointestinal tract
of interest. In preferred embodiments of the present invention, at least the anterior portion of
the stomach is exposed sufficiently to allow for its reconfiguration. This may require
35 dissection and/or removal of at least a portion of the omentum, and it may require lifting

5 and/or partial retraction of the liver, both of which are relatively simple interventional steps that are well known in the art. The subsequent reconfiguration and gastric reduction may then be performed, preferably using the devices and systems of the present invention, which are described in detail below.

Figure 1 schematically illustrates the relevant portion of the gastrointestinal tract 10 (anterior view), both pre-procedure (Figure 1A) and post-procedure (Figure 1B). To aid in the following discussion, it is helpful to first distinguish the various anatomical structures in Figure 1A. The stomach itself lies between the esophagus 105 and pylorus 110. The anterior wall 115 of the stomach is shown, along with the fundus 120, the greater curvature 125, and lesser curvature 130. Two cross-sectional views of the stomach are shown in 15 Figure 1A1 at X-X and in Figure 1A2 at Y-Y. It is helpful to point out the major tissue layers of the stomach wall, as illustrated in Figure 1A3. Starting intragastrically and moving outward, the innermost tissue layer is the mucosal tissue layer 150, then there is a submucosal connective tissue layer 152, the muscularis tissue layer 155, and the exterior serosal tissue layer 160 that covers the extragastric surface of the stomach.

20 Figure 1B illustrates a stomach following gastric reduction according to methods of the present invention. As shown in Figures 1B1 and 1B2, the stomach now exhibits a significantly reduced cross sectional area (e.g. at X-X and Y-Y) and the functional volume of the stomach has been decreased approximately 50% as a result of single fold 180 being placed in the anterior wall 115 of the stomach. As shown, fold 180 is located approximately 25 midway between the greater curvature 125 and lesser curvature 130, and extends approximately longitudinally from near fundus 120 to near pylorus 110. As can be seen in sections X-X and Y-Y of Figures 1B1 and 1B2, fold 180 was created by invaginating and approximating the tissue of the anterior wall 115 of the stomach so as to bring the serosal tissue layer 160 into contact with itself. Fasteners are then applied to the tissue brought 30 together to produce the plication in the wall of the stomach.

In a preferred embodiment of the present invention, a single fold and plication is produced in the above described manner and location, as illustrated in Figure 1B; however, in other embodiments, two or more such plications may be produced. Although the plication is illustrated as being formed approximately midway between the greater and lesser 35 curvatures of the stomach, it will be appreciated that other areas of the stomach or

5 gastrointestinal wall may be used, as may be necessary based on individual anatomy and the surgeon's desire to achieve the targeted functional gastric reduction, while minimizing the overall invasiveness of the procedure. According to the present invention the functional volume of the stomach is preferably decreased at least 20%, is more preferably decreased at least 30%, and is most preferably decreased at least 40%. In morbidly obese patients, a
10 functional volume reduction of 50% or more may be achieved in order to promote the desired excessive weight loss.

In **Figure 1B**, securing means comprising a row of individual staples **185** are placed substantially along the length of fold **180**. As shown in **Figure 1B2** at section **Y-Y**, staples **185** grasp tissue shoulders **195** that are formed where the opposing tissue layers of the tissue fold intersect the circumference of the stomach. As can also be seen in section **Y-Y**, according to a preferred embodiment of the present invention, staples **185** engage tissue shoulders **195** by penetrating only through serosal tissue layer **160** and underlying muscularis tissue layer **155**, without penetrating completely through the stomach wall to breach or otherwise compromise mucosal tissue layer **150**. As can also be seen in section **Y-Y**, according to another preferred embodiment of the present invention, the approximated tissue surfaces within the tissue fold are configured such that there is substantially intimate serosal-to-serosa contact within the plication **190**.

Figure 2 illustrates in greater detail the intermediate steps of the procedure, according to one embodiment of the present invention. **Figure 2A** and **Figure 2E** are identical to **Figure 1A** and **Figure 1B**, respectively, and are repeated for completeness. **Figure 2B**, **Figure 2C** and **Figure 2D** are helpful to explain other aspects of the intermediate steps. In **Figure 2B**, for example, prior to commencing with the reconfiguration portion of the procedure, the region of interest on anterior wall **115** may be visually identified, marked or mapped out to aid subsequent steps of the procedure. For example, it may be desirable to identify and/or indicate the target position and length of the fold centerline **202**, as well as the bounding lines **204** and **206** where the tissue will be contacted, engaged and/or secured. The location of bounding lines **204** and **206** define the depth of the tissue fold to be created, as well as the surface area of tissue that will be approximated during creation of the tissue fold. Identification, marking and/or mapping of the tissue structures and/or locations can be carried out according to methods well known in the art, for example, inks, dyes, adhesives,

5 implantable tags, clips, fasteners, radio-opaque markers, fluorescent markers, cauterizing marks, and the like, may be used.

Figure 2C schematically illustrates the early steps in the procedure, starting at one end of the target area (e.g. near the pylorus) and working progressively in one direction (e.g. toward the fundus). It should be recognized, however, that this progression is optional, and
10 that it is just as feasible to start near the fundus and work toward the pylorus, to start anywhere along the length of the intended fold and work in both directions, or any combination of the foregoing. To form a tissue fold, the tissue is contacted and/or engaged at two or more locations, and various combinations of relative motions are then used to ensure the tissue is invaginated as the opposing tissue surfaces are approximated. Examples of such
15 combinations of relative motions include one or more motions selected from the group consisting of pushing motions, pulling motions, twisting motions, and shearing motions.

In Figure 2C, for example, tissue is contacted and engaged at locations 208 and 210 on opposite sides of a fold centerline location 212. Relative motion between central location 212 and the tissue contact and engagement locations 208 and 210, is represented in Figure
20 2C1 by pushing force vector 214 and pulling force vectors 216 and 218, respectively. These motions invaginate the tissue and approximate the opposing tissue surfaces, while bringing tissue shoulders 195 toward each other for subsequent securing. The relative motion illustrated may be achieved, for example, by holding central location 212 substantially stationary and pulling the tissue engagement points 208 and 210, or by holding the tissue engagement points 208 and 210 substantially stationary and pushing on the central location 212, or alternatively, any combination of pushing and pulling may be used to achieve the same effect.

After the tissue has been approximated to create the tissue fold 180 as described above, and tissue shoulders 195 have been brought together into proximity of one another, a
30 tissue fastener 185 is then applied at that location to secure the plication 190, as shown in Figure 2D. In Figure 2D, exemplary tissue fastener 185 is schematically shown as a box-type of interventional staple, similar in form and function to a box-type staple known in the art of interventional skin stapling for use in wound closure applications. However, it should be obvious to those skilled in the art that, within the scope of the present invention, a wide
35 variety of mechanical elements may be used as tissue fasteners 185 for the purpose of

5 anchoring, fastening, holding, attaching, or otherwise securing tissue surfaces **180** to produce plication **190**. Examples of suitable tissue fasteners that may be used include but are not limited to sutures, staples, screws, tacks (e.g. U-shaped, circular and helical fasteners), clips, hooks, clamps, t-tags, and the like. In a preferred embodiment of the present invention, tissue fasteners **185** are preferably applied at least directly across tissue shoulders **195** at
10 more than one location along the length of tissue fold **180**, more preferably at several relatively closely spaced locations to secure the plication.

The tissue engagement, approximation and fastening steps are repeated any number of times as is necessary to completely form and secure the one or more tissue plications. In the example provided herein, the final result is shown schematically in **Figure 2E**.

15 For convenience, the procedure may progress sequentially in one direction along the length of the intended fold, as illustrated in **Figure 2D**, effectively producing the plication in a manner similar to closing a zipper. However, sequential advancement is not required, and the surgeon may use discretion in deciding where to begin and how to advance the procedure.
20 At each of one or more locations along the length of the intended fold, the tissue is invaginated, approximated and secured with one or more tissue fasteners before moving to the next location. In one embodiment, a device may be provided that allows simultaneous or sequential placement of multiple tissue fasteners while the invaginating and approximating tool is placed and held at one location. Alternatively, in another embodiment, a device may be provided that allows placement of a single tissue fastener along a substantial length, or
25 even along the complete length, of the tissue fold, while the invaginating and approximating tool is held at one location.

According to one embodiment of the present invention, prior to securing the approximated tissue to produce the one or more plications, at least a portion of the surface area of the serosal tissue enfolded by the one or more plications is selectively treated to
30 promote serosal-to-serosal tissue bonding. There is a considerable body of clinical knowledge regarding the mechanisms of abdominal adhesion formation, and a variety of methods known to those skilled in the art may be used to selectively treat the serosal tissue surfaces to promote tissue adhesion of the serosal tissue layers adjoining one another inside the tissue fold forming the plication. Examples of such tissue treatments include but are not
35 limited to mechanical disruption methods (e.g. abrasion), energy deposition methods (e.g.

5 RF, ultrasonic, electromagnetic, and the like), methods involving treatment using liquids (e.g. chemicals, pharmaceuticals, adhesives, etc.) and methods involving treatment using solids (e.g. powders, films, etc.). Regardless of the tissue treatment method used, an important aspect of this embodiment is that serosal tissue bonding or adhesion is promoted over a sufficiently large interfacial surface area across the approximated tissue boundary
10 within the plication to achieve a strong and durable serosa-to-serosa bond post-operatively.

In yet another embodiment of the present invention, additional tissue fasteners may also be optionally applied while the tissues are being approximated to aid in forming, stabilizing and/or providing additional strength to the resulting tissue plication, as well as to further promote the formation of a strong serosa-to-serosa bond inside the plication. For
15 example, as illustrated in the enlarged cross sectional view X-X shown in **Figure 3B**, in addition to outer tissue fastener **305** (similar to the tissue fastener **185** described previously), one or more additional internal tissue fastener **310** may be applied across the contact area of the approximated tissue surfaces within the fold while it is being formed, such that after the plication is completed, the one or more additional internal tissue fasteners **310** are located
20 inside the plication for the purpose of better securing the tissue across the approximated tissue surfaces. Additional internal tissue fastener **310** may be identical to outer tissue fastener **305**, being placed by the same device, or in an alternative embodiment, additional internal tissue fastener **310** may have a different design and/or be placed using additional devices. Note that additional internal tissue fastener **310** also preferably penetrates only the
25 serosal and muscularis tissue layers. Although **Figure 3** illustrates the use of a box-type staple, as in the case of tissue fastener **185** described previously, this embodiment is merely illustrative and a wide variety of alternative fasteners exist that may be used for the outer tissue fastener **305** and additional internal tissue fastener **310**, within the scope of the present invention.

30 In yet another embodiment of the present invention, more than one tissue plication may be produced according to the previously described methods. For a variety of reasons, it may be advantageous in some cases to produce two or more plications. These advantages may include, for example, allowing a greater range of effective volume reductions in the stomach to be achieved, allowing smaller laparoscopic devices to be used, allowing the
35 surgeon more flexibility in positioning of the plications relative to the stomach or

5 surrounding organs, for reducing the maximum forces generated on the individual securing means, and so on. **Figures 4A and 4B** schematically show an example according to one embodiment of the present invention in which tissue two adjacent tissue folds **402** and **404** have been placed in the anterior wall of the stomach, running more or less parallel to one another. As can be seen in **Figure 4B** in the enlarged view of cross section X-X, tissue fold
10 **402** has been secured with tissue fastener **405** to produce a first plication **410**, whereas tissue fold **404** has been secured with tissue fastener **415** to produce a second plication **420**. It should be obvious to those skilled in the art that within the scope of the present invention, it is possible to produce any number of individual and separate plications in the manner described previously, each of which plication may be characterized individually in terms of
15 length, depth, position, number and type of fasteners placed, and so on, to achieve the intended interventional result.

Interventional Devices and Systems

Interventional devices for performing methods of the present invention are described herein that, taken together, comprise systems of the present invention. The devices and
20 systems of the present invention provide the ability to carry out the above described volume reduction procedures in a safe, efficient and minimally invasive manner, which is difficult or impossible to accomplish using prior art devices. It will be appreciated that while the devices and systems of the present invention are described below with respect to their use in gastric reduction methods of the present invention, they have utility and may be used for general
25 approximation and fastening of other types of soft body tissues and in other types of interventional procedures as well.

In general, at least one handheld interventional instrument is provided having one or more integrated tool assembly(ies) adapted for placement at an interventional site, such as within the abdominal cavity, in combination with one or more actuator(s) positioned
30 remotely from the tool assembly and providing operator control of the tool assembly(ies) during an intervention. The tool assembly is preferably capable of engaging tissue at two or more separate locations, and then invaginating and approximating tissue to effectively create a tissue fold between the tissue engagement locations. In one embodiment, the tool assembly comprises at least two tissue engagement mechanisms (e.g. clamps, grippers, forceps, jaws,
35 hooks, barbs, vacuum ports or the like, or combinations of these mechanisms) positioned at

5 or in proximity to the distal end of an elongate shaft of a laparoscopic device. The tissue engagement mechanisms may be positionable by means of a remote actuator, or they may be mounted on supporting members that may be positionable to engage desired tissue sites. Using this device, the laparoscopic shaft is positioned within the abdominal cavity, and the distal end of the shaft is positioned at a first desired tissue engagement site, where a tissue
10 engagement mechanism is engaged with the tissue. The operator then repositions the shaft by moving it to a second location, dragging the first engaged tissue location toward the second, and thereby approximating the first and second tissue locations. The approximated tissues may then be fastened to one another to secure the plication using fasteners applied with an independent device or an integrated assembly of the tissue approximation device.

15 In another embodiment, a first tissue engagement mechanism may be positioned at the distal end of the elongate shaft of a laparoscopic device, while a second tissue engagement mechanism may be positioned at the distal end of an extendible member that can be manipulated by an operator to move away from the axis of the device shaft to position the second tissue engagement mechanism at a second location, remote from the distal end of the
20 device. The extendible member may be substantially rigid, or it may be flexible, or it may have both substantially rigid and flexible portions, and it may either be deployable from inside the elongate shaft of the laparoscopic device, or attached near the distal end of the shaft by mechanical means. In one embodiment, a proximal end of an extendible member is attached near the distal end of the elongate shaft using a pivot connection, a hinge
25 connection, a flexible connection, or the like, that allows the extendible member to be operatively and selectively actuated to move its distal, operating end (comprising a tissue engagement member) away from the axis of the laparoscopic device to engage tissue. In operation, the distal end of the shaft of the laparoscopic device is first positioned at a desired tissue surface and the tissue is engaged at a first site. The extendible member and its
30 associated tissue engagement mechanism is then deployed, extending away from the axis of the shaft to independently engage tissue at a second location. The extendible arm and its associated tissue engagement mechanism is then retracted, under control of the operator, and the second engaged tissue location is drawn in toward the axis of the shaft and thereby approximated adjacent the first engaged tissue site. An invaginated tissue fold projecting

5 away from the distal end of the device and into the gastrointestinal space is created as the two tissue sites are drawn together and approximated.

In other embodiments, described in detail below, two or more such extendible members are provided on an interventional device, each extendible member having at least one tissue engagement mechanism, generally (but not necessarily) positioned at its distal end, 10 such that the engagement of tissue at multiple separate locations can be accomplished without requiring the shaft of the laparoscopic device itself to contact the tissue surface. The extendible members may be actuated and positioned separately and independently of one another, or they may be actuated and positioned simultaneously and in coordination with one another. Operation of this type of device involves deploying each of the extendible members 15 and their associated tissue engagement mechanisms, independently or in coordination, to contact the tissue engagement mechanisms at two locations on the tissue, then approximating the engaged tissue to form an invaginated tissue fold by moving at least one of the extendible members toward the other and, in some embodiments, by moving multiple extendible members toward a central location, thereby approximating the engaged tissue substantially 20 near the distal end of the device (or along a longitudinal axis extending therefrom).

Another embodiment that provides an alternative to using two or more extendible members to engage tissue involves the use of tethers. In this case, the distal end of the shaft of a laparoscopic instrument may be positioned to sequentially engage tissue at each of two or more locations using releasable tissue engagement mechanisms mounted on retrievable 25 tethers, wherein each tissue engagement mechanism, after being engaged in tissue, is released from the end of the shaft of the laparoscopic instrument, yet remains connected to the instrument by a tether (e.g. a suture, wire, or the like). This allows the instrument to be moved freely between each desired tissue engagement location to deploy two or more tissue engagement mechanisms at different tissue sites. Subsequently, the tethers may be 30 selectively retrieved, or retracted back toward the shaft of the device to draw the engaged tissue sites toward one another, thereby approximating the tissue sites. Alternatively a cinching member through which the flexible tethers pass may be slid distally down the length of tethers, causing the engaged tissue locations to move toward each other, thereby approximating tissue. Retrieval of the tether(s) and/or operation of the cinching member(s) is 35 under the control of an operator using associated actuation mechanisms.

5 It will be appreciated that methods and systems of the present invention may be used in connection with other diagnostic and therapeutic methods and devices. Methods of the present invention may thus be used, for example, in connection with conventional diagnostic and therapeutic methods and may involve the administration of diagnostic or therapeutic agents, agents for visualizing the interventional site, and the like. Similarly, device
10 components of the present invention may be used in connection with various procedures and agents that are known in the art. Certain device components that are intended for introduction to the interventional site, such as tissue engagement mechanisms, probes, extendible members, fasteners, and the like may be administered in association with various types of diagnostic or therapeutic agents, or may be coated or impregnated with such
15 materials. Suitable agents may include clotting agents, healing agents, hydrophobic and/or hydrophilic materials, agents promoting lubricity, and the like.

Figures 5A-5F illustrate an exemplary tissue approximation device according to one embodiment of the present invention. An overview of device 500 is shown in Figure 5A in the pre-deployed configuration, and Figure 5B shows a distal end of device 500 in the deployed configuration. Device 500 comprises an elongate tubular member 502 having at least one internal working channel 504, handle assembly 506 positioned at the proximal end, and approximating tool assembly 508 positioned at the distal end, wherein approximating tool assembly 508 is shown in the collapsed (i.e. pre-deployment or fully retracted) state, substantially confined within working channel 504. In the case of minimally invasive laparoscopic surgery, this low profile collapsed state configuration is useful for delivery of the instrument to and removal of the instrument from an internal site in the patient, such as the abdominal cavity, through a standard trocar. It is therefore generally desirable that the outer diameter of elongate tubular member 502 be as small as possible, preferably 15 mm or less, more preferably 12 mm or less and, in some embodiments, 5 mm or less. Also shown in Figure 5A, actuating mechanisms such as a trigger 510, slider 512, and plunger 514 are provided in connection with handle assembly 506. Also shown is rotating collar 516 that allows the orientation of handle assembly 506 to be independently adjusted by the operator relative to the orientation of approximating tool assembly 508.

Figure 5B shows an enlarged cross section view of the distal end of device 500, with approximating tool assembly 508 being shown in the collapsed state. In this configuration,

5 located along longitudinal axis **518** of working channel **504** are two (or more) extendible members **520**, and pushing member **522**, each being operatively connected to an actuating mechanism operated at the handle assembly **506**, as described below. Each of said extendible members **520** is configured at its distal end with a tissue engagement mechanism **524** comprising one or more mechanisms for controllably and selectively grasping, grabbing, 10 gripping, piercing, holding or otherwise engaging tissue. In the example shown, tissue engagement mechanism **524** incorporates a tissue hook **526**. Hook **526** has a generally pointed distal end for penetration of tissue and has a relatively short curved segment, thus limiting the degree of tissue penetration. Tissue engagement mechanisms having a generally pointed and sharp tissue penetration structure for penetrating tissue, such as the relatively 15 tough serosal layer forming the exterior gastric wall, are preferred in many embodiments.

Figure 5C shows an enlarged view of the distal end of tissue approximation device **500**, with approximating tool assembly **508** being shown in the extended state, i.e. after being deployed by the operator. In this embodiment, extendible members **520** open, or extend, along a predefined path as they're released from the distal end of the shaft. An actuating 20 mechanism such as plunger **514** is operatively connected to extendible members **520**, such that when plunger **514** is axially displaced into handle assembly **506**, extendible members **520** move distally along longitudinal axis **518** and thereby extend outward from working channel **504** beyond the end of elongate tubular member **502**. After deployment to an expanded state, each of extendible members **520** is positioned with its distal ends **524** spaced 25 apart from and positioned on opposite sides of longitudinal axis **518** from an opposing extendible member.

The degree of extension of the extendible members, and the spacing **521** between distal ends **524** of extendible members **520** may be governed by the degree of deployment out of shaft **502**. In some embodiments, both the degree of extension of distal ends **524** from 30 the shaft **504**, indicated as longitudinal spacing **519**, and the distance between extended distal ends **524** are selectively controllable by the operator to facilitate tissue engagement at desired locations, and to facilitate the creation of a tissue plication of the desired dimensions, thereby producing the desired gastric volume reduction.

Tissue approximating device **500** illustrated in **Figures 5A-5F** additionally comprises 35 a pushing member **522** operatively connected to an actuator, such as slider **512**, such that

5 when slider **512** is translated away from its proximal (fully retracted) position, the distal end
of pushing member **522** moves along longitudinal axis **518**, thereby extending out of working
channel **504** a distance **505** beyond the end of elongate tubular member **502**. The extension
of pushing member **522** facilitates invagination of a tissue fold and formation of a tissue
plication as two or more tissue sites are approximated. Pushing member **522** may be
10 operated independently of, or in coordination with, extendible members **520**. In one
embodiment, pushing member **522** is extended out of working channel **504** as the extendible
members **520** are extended and the tissue engagement mechanisms are positioned to engage
tissue.

15 Illustrative operation of a tissue approximation device **500** illustrated in **Figures 5A-**
5F is described below. Following insertion of the shaft into the intra-abdominal space and
positioning of the distal end of the shaft near a desired tissue approximation site, extendible
members **520** are deployed from a collapsed state to an expanded state to prepare the device
for subsequent tissue engagement steps. In one embodiment, extendible members **520** are
20 expanded by an actuator that pushes the members out of, or releases them from the shaft, as
follows. In this case, extendible members **520** are produced from a highly flexible and
elastically deformable material (e.g. flexible polymers, flexible metals, shape change
materials and combinations thereof may be used) and are made in a shape when in the
expanded state having an outward (i.e. away from longitudinal axis **518**) curvature. As the
25 extendible members **520** are released from the working channel **504**, they assume their
expanded state, and the distal tissue engagement mechanisms are brought into contact with
the tissue surface. Due to their flexible nature and outwardly curved shape, extendible
members **520** flex elastically and continue to assume a progressively more extended
condition as the operator continues releasing them from the shaft, causing distal arm portions
30 **524** to slide outward along the tissue surface, becoming spaced apart, until the distal tissue
engagement mechanisms are located in the desired positions for tissue engagement, as
described below.

35 In another embodiment, extendible members **520** are designed to be released from the
collapsed state to the expanded state in a self-actuating manner, automatically achieving the
desired tissue engagement configuration when extended out of working channel **504** beyond
the end of elongate tubular member **502**. Such self-actuating motions can be achieved by

5 various methods known in the art. For example, in one preferred embodiment of the present invention, extendible members **520** are produced from a highly elastic material (e.g. spring steel, hardened stainless steel, a shape change material such as a superelastic NiTi alloy, superelastic polymer, or the like) and are formed during manufacturing into the desired final deployed shape by mechanical and/or thermomechanical processing means known in the art.

10 Extendible members **520** are then biased (i.e. mechanical potential energy is stored, similar to a pre-loaded spring) by elastically deforming and loading them into working channel **504** to thereby provide the device in its collapsed state. As extendible members **520** are then pushed out of working channel **504** during deployment, the stored energy is released and extendible members **520** automatically return to the pre-determined shape desired for subsequent tissue engagement when brought into contact with the tissue surface. It will be appreciated that different assemblies of extendible members having different dimensions, different curvatures, different elastic properties, and the like may be provided for use in a tissue approximating device of the present invention and an operator may select an appropriate extendible member assembly having the desired dimensions and extension properties and

15 install the desired assembly in the working channel prior to an intervention.

20

In yet other embodiments, deployment of extendible members **520** from the collapsed state to the expanded state may be accomplished, by means of an actuating mechanism, by any combination of manual pushing to cause expansion and self-actuating expansion mechanisms. Factors that may be adjusted to optimize the above described reconfiguration and deployment motions include, for example, the cross sectional shape, curvatures, mechanical properties, length, etc. of extendible members **520**. It should also be obvious to those skilled in the art that, within the scope of the present invention, other mechanical actuation mechanisms of providing the desired reconfiguration and deployment to adjust the extendible members from the collapsed state to the expanded state may also be used. Such actuating mechanisms may comprise, for example, springs, levers, cams, gears, linkages, and the like may be used.

30 Distal ends **524** of extendible members **520** each incorporate one or more tissue engagement means configured to allow targeted tissue surface **535** to be selectively and controllably engaged by the device when actuated by the operator. Various tissue engagement mechanisms are known in the art may be employed to provide secure and robust

5 tissue engagement having sufficient strength, for example, to allow the tissue to be subsequently pulled or otherwise manipulated without disengaging, slipping or tearing. Tissue engagement mechanisms that may be used include, for example, hooks, barbs, grippers, teeth, clamps, jaws, clips, t-tags, and the like. According to one embodiment of the present invention, as shown in **Figure 5C**, tissue hooks **526** are located at the distal ends **524**,
10 and further comprise sharpened points **528** to promote tissue penetration. While extendible members **520** are in the expanded state, distal ends **524** and tissue hooks **526** are positioned such that sharpened points **528** curve slightly downward (distally) and inward (toward longitudinal axis **518**). As a result, when pushed slightly downward onto the surface of the tissue, elastic deformation of extendible members **520** causes distal ends **524** to first move
15 slightly outward. Then, when extendible members **520** are either lifted slightly (e.g. by the surgeon lifting device **500**) or alternatively, when retraction of the extendible members is initiated by the operator (as described below), tissue hooks **526** move slightly downward and inward, thereby causing sharpened points **528** to pierce, penetrate and securely engage the tissue at tissue engagement locations **530**, as shown in **Figure 5D**. Preferably, distal ends
20 **524**, tissue hooks **526** and sharpened points **528** are designed such that secure tissue engagement is achieved by penetrating only the serosal tissue surface **535** (i.e. the serosal tissue layer), or a combination of the serosal and muscularis tissue layers, without penetrating the mucosal tissue surface **540**.

While more complicated mechanical tissue engagement means may be employed in accordance with the present invention (e.g. hinged jaws, mechanical clamps, forceps, grippers, vacuum actuated mechanisms, and the like) there are several advantages to the embodiment described above, and similarly designed self-actuating embodiments. One advantage, for example, is that it is a simple, single component design having low production cost . Additionally, successful operation of this device is not particularly dependent upon operator technique (i.e. no sophisticated hand motions or unusual device manipulations are required), successful operation instead being more dependent upon device design factors that control, for example, the directions and magnitudes of the forces generated by extendible members **520** during the pushing and pulling motions involved in deployment and/or retraction of the device. Examples of design factors that may be optimized in the self-actuating design embodiments of the present invention include the shape, physical
30
35

5 dimensions, geometrical angles, surface finish, and the like, of extendible members **520**, distal ends **524**, tissue hooks **526**, and sharpened points **528**, as well as their materials of manufacture and mechanical properties.

In one embodiment, extendible members **520** have a non-circular, generally flattened cross section to effectively increase the lateral (i.e. out of plane) stiffness when extendible 10 members **520** are extended. Examples of suitable non-circular cross sectional shapes include square cross sections, rectangular cross sections, triangular cross sections, arcuate cross sections, hemispherical cross sections, oblong or flattened cross-sections, and combinations of the foregoing. The cross sectional shape, physical dimensions, mechanical properties, and so on, of extendible members **520** may be designed having variations along their length to 15 provide improved deployment, tissue engagement or retraction characteristics.

In another embodiment, extendible members **520** have a pre-determined shape when in the expanded state that includes at least two bends having radii of curvature in substantially opposing directions. Such a shape, as illustrated in **Figures 5C-5E** and explained above, may be utilized to initially give rise to a slight downward motion of distal 20 ends **524**, in addition to the inward motion that occurs during the retraction of extendible members **520** back into working channel **504**, wherein the combined initial downward and inward motions of distal ends **524** effectively promotes tissue penetration and secure tissue engagement of sharpened points **528** on tissue hooks **526** upon actuated retraction of extendible members **520**. The combined initial downward and inward motions of distal ends 25 **524** that promote tissue penetration and secure tissue engagement may also be achieved using other designs obvious to those skilled in the art;. This embodiment simplifies the operation, improves consistency, reduces procedural times and risk of complications, by minimizing reliance on individual operator technique and instead taking advantage of highly controlled and repeatable device motions.

30 After tissue has been securely engaged by approximating tool assembly **508**, as described above, the operator actuates device **500** to initiate the tissue invagination and approximation step, wherein the desired tissue fold is formed by bringing serosal tissue surfaces between the engaged tissue sites in contact with each other, so that the mucosal tissue surface **540** forms a plication extending into the gastrointestinal lumen. **Figure 5E** 35 illustrates this process. In the example provided, the operator selectively activates device

5 **500** remotely using trigger **510** provided within handle assembly **506**, which is operatively connected to extendible members **520** in a manner such that, as trigger **510** is squeezed, extendible members **520** are thereby controllably retracted and pulled back into working channel **504**, as indicated by retraction forces **531**. The mechanisms used to operatively connect trigger **510** to extendible members **520** may include various mechanical elements
10 known to those skilled in the art, such as gears, transmissions, levers, pivots, linkages, and the like, whether manual or automated, in order to provide the retraction forces at the working (distal) end of the device, while keeping the actuating mechanisms operated by the operator at a convenient level.

The retraction of extendible members **520** causes tissue engagement locations **530** to
15 be gradually pulled inward toward longitudinal axis **518**. In one device embodiment that incorporates a pushing member, the operator may selectively and independently actuate pushing member **522** from within handle assembly **506** (i.e. using slider **512**) as the tissue engagement locations are drawn toward one another. The pushing member is extended distally along longitudinal axis **518** to contact and push against the tissue, e.g. with pushing
20 force **532**, at a location between tissue engagement points **530**. This promotes tissue invagination in the desired manner while the engaged tissue is approximated, as shown in
Figure 5E. Once extendible members **520** have been fully retracted by complete actuation of trigger **510**, the tissue engagement locations **530** have been brought into approximation near the distal end of elongate tubular member **502** to create tissue fold **540** as shown in
25 **Figure 5F**. In this illustration, pushing member **522** is shown remaining in the fully extended position.

The combination of extendible members and a pushing member in devices of the present invention, enabling the combined action of pulling tissue engagement points **530** toward one another via retraction of extendible members **520** while simultaneously having
30 the user selectable option to push against the tissue between tissue engagement points **530** with pushing member **522** promotes creation of a uniform and consistent tissue fold, as shown in **Figure 5F**. In preferred embodiments of the present invention therefore, operation of the device in the described manner effectively approximates opposing serosal tissue surfaces **535** inside the tissue fold, providing substantially intimate serosa-to-serosa contact,

- 5 without forming wrinkles, bunches, gaps, or the like, and without penetrating the mucosal tissue surface **540**.

In other embodiments of the present invention, additional user selectable controls may be optionally provided within handle assembly **506**. For example, controls may be optionally provided to allow the surgeon to adjust the span **521** of extendible members **520** 10 when in the expanded state, and the distal extension distance **505** and pushing force **532** of pushing member **522**. Independent, operator controlled actuation mechanisms may be provided for each of the more than one extendible member **520**, and the actuation mechanisms may control the speed and force that may be used to retract extendible members **520**, as well as other operating parameters. It should also be recognized that the actuation 15 means described above are exemplary, and that other actuation and control mechanisms that are known to those skilled in the art may be used and are considered within the scope of the present invention. For example, actuation may be accomplished manually by one or more various means known in the art (e.g. triggers, levers, buttons, knobs, or the like) or by one or more various powered means known in the art (e.g. AC or DC electric motors, compressed 20 gas, vacuum, or the like), or by any combination of the foregoing.

As described previously, according to one embodiment of the present invention, it is desirable to selectively and therapeutically treat the serosal tissue layer to promote bonding or adhesion of the serosal layers that abut one another within the plication. This may be accomplished using device **500** in various ways. For example, in one embodiment illustrated 25 in **Figures 5C-5F**, the distal tip and/or lateral surfaces of pushing member **522** may be used to mechanically disturb and disrupt the thin layer of mesothelial cells that form the outermost covering of the serosa. Since the layer of mesothelial cells covering the serosa is quite thin and fragile, it is easily disrupted, and pushing member **522** may be scraped, dragged or otherwise frictionally moved across the surface of the tissue to produce the desired 30 disruption. To further aid in disrupting the serosal tissue surface and promote tissue adhesion, pushing member **522** may be modified, for example, by incorporating roughening features **523**, illustrated as protuberances in **Figures 5C-5F**. As will be obvious to those skilled in the art, a wide variety of such roughening features and arrangements may be used 35 to accomplish the desired serosal treatment, for example, ridges, bumps, bristles, teeth, scales, serrations, and the like may be used.

5 The optional serosal treatment described above may be carried out before the tissue fold is formed, after the tissue fold is formed but prior to the securing means is applied, after the tissue fold is formed and the securing means is applied, or any combination of the foregoing. For example, prior to actuating extendible members **520** to engage tissue, the distal end of pushing member **522** may be moved across substantially the identified area of
10 serosal tissue to be included within the tissue fold in a sweeping or painting type of motion. Alternatively, the lateral surfaces of pushing member **522** contact and slide across the opposing serosal tissue surfaces of the tissue fold when pushing member **522** is retracted from within the tissue fold (as is evident in **Figure 5F**), thereby disrupting at least a substantial portion of the serosal tissue surface during normal device operation. In this case,
15 roughening features **523** present on the lateral surfaces of pushing member **522** may ensure more uniform and consistent serosal treatment, leading to a more effective and stronger serosa-to-serosa tissue bond.

20 In another serosal treatment embodiment, ports may be provided near the distal tip of shaft **502** and/or along pushing member **522** such that, when the shaft and/or pushing member lumen is connected to a supply of source material (e.g., a liquid reservoir located within or attached to the proximal handle assembly **506**), the device provides controlled dispensing of a chemical or therapeutic agent (e.g. liquid, gas, solid powder, solid film, or combinations thereof) onto the tissue surface that promotes tissue bonding and adhesion.
25 Alternatively, the distal tip of shaft **502** and/or pushing member **522** may optionally incorporate an energy deposition mechanism capable of delivering energy to the target tissue. Exemplary energy deposition mechanisms include, for example, components capable of RF cauterizing, electro-cauterizing, ultrasonic vibration, and the like.

30 According to the present invention, once the tissue has been approximated and the desired tissue fold has been created as described above, fasteners are then applied to secure the plication. This is most conveniently accomplished while approximating tool assembly **508** is held in place by the operator to maintain the tissue in a stable, folded configuration. In one embodiment, a separate interventional instrument may be introduced through a separate trocar, and its distal tip may be positioned immediately adjacent approximating tool assembly **508**. This instrument is then actuated to apply a fastener directly into and across the shoulders of the approximated tissue forming the tissue fold, thereby securing the plication.
35

5 In this embodiment illustrated in **Figure 6A**, a system **600** of the present invention comprises two separate handheld devices, each device capable of being actuated using controls located at their respective proximal handle assemblies. A first device **620** incorporates an approximating tool assembly **625** which may be substantially similar to approximating tool assembly **508**, described above, at its distal end, and a second device **640** incorporates a
10 fastening tool assembly **645** at its distal end, capable of applying a fastener to the tissue fold to secure the plication. A wide variety of a suitable fasteners are known to those skilled in the art and may be suitably be used as fasteners within the broad scope of the present invention. Exemplary fasteners comprise, for example, sutures, box-type staples, U-shaped or hemispherical fasteners, helical fasteners, clips, tacks, wall anchors, t-tags, and the like. A
15 commercially available laparoscopic stapler, suturing device or tack applicator may be used to secure the tissue fold.

Accordingly, the laparoscopic interventional stapler shown in **Figure 6A** comprises an elongate tubular shaft **650** having at its proximal end a handle assembly **655** containing user controls, actuation mechanisms, and so on, and having at its distal end a fastening tool assembly **645**, which incorporates mechanisms known in the art for feeding, deploying, forming and applying to the target tissue a plurality of fasteners. These fasteners are most commonly made from stainless steel, titanium or NiTi, although other materials may also be used (e.g. other biocompatible alloys, polymers, bioabsorbable materials, and the like). Typically, a plurality of such staples would be provided within a disposable (i.e. single patient use) cartridge that is loaded at the distal end of the device, allowing multiple staples to be placed consecutively by the operator without removing the device from the patient.
25

Figure 6B shows a close up view of the distal ends of device **620** and device **640**, indicating the preferred relative positioning of approximating tool assembly **625** and fastening tool assembly **645**, respectively, according to one embodiment of the present invention. In this view, approximating tool assembly **625** has previously been deployed, the tissue has been engaged, and the extendible members have been retracted (these steps being carried out e.g. as described in **Figure 5**), in order to create tissue fold **660**. Shoulders **665** of tissue fold **660** are approximated near the distal tip of approximating tool assembly **625**, and are held in position, ready for the tissue fastener to be applied by fastening tool assembly
30 **645**. The cross sectional view of **Figure 6C** shows a close up of the distal tip of fastening
35

5 tool assembly 645. In this example, a box-type staple in the pre-deployed state 670 is shown loaded within the within fastening tool assembly 645. Prior to applying the staple, fastening tool assembly 645 is positioned such that staple legs 671 of box-type staple in pre-deployed state 670 are positioned substantially perpendicular to, and in contact with, shoulders 665 of the tissue fold. When the surgeon fires the stapler using actuation means provided within the
10 proximal handle assembly, extendible pistons 642 extend distally, deforming staple legs 671 around stationary anvil 644 and thereby reconfiguring the box-type staple into deployed state 675 as it is ejected from the device. As the staple is deployed, it penetrates the tissue and simultaneously pulls opposing tissue shoulders 665 toward one another, as shown. Note in this example that the box-type staple in deployed state 675 engages only the outermost layers
15 of gastric tissue, i.e. serosal layer 535 and/or the muscularis tissue layers (not shown), and that there is no penetration through the gastric wall, which preserves the mucosal tissue layer 540 intact. **Figure 6D** schematically illustrates a plication being secured using several consecutively repeated applications of the above described procedure. Approximating tool assembly 625 and fastening tool assembly 645 are shown, along with a multiplicity of
20 individual box-type staples in the deployed state 675 that have been applied and which are arranged in a substantially continuous row extending along the length of tissue shoulders 665 to secure plication 690 projecting into the gastrointestinal space. The depth 680 below the surface and spacing 685 between the individual staple placements may be selectively controlled by the operator.

25 In another embodiment of the present invention, the tissue approximating and fastening functions described above requiring the use of two separately operable handheld interventional instruments are combined into a single multi-functional device having one or more integrated tools capable of invaginating and approximating tissue to create a tissue fold, as well as one or more integrated tools for applying fasteners to secure the plication. By
30 combining these functions conveniently in a single handheld device, the overall procedure is simplified, and it can be performed without requiring extensive operator training. Furthermore, the need for one laparoscopic access port is eliminated, which provides a significant advantage.

35 **Figures 7A-7H** illustrates such an integrated device and its operation, according to one embodiment of the present invention. Device 700 comprises an elongate tubular member

5 702 having internal working channel 704 and handle assembly 706 positioned at the proximal end. At the distal end of device 700 is multi-functional tool assembly 708, shown in the collapsed (i.e. pre-deployment or fully retracted) state in **Figure 7A**. It is generally desirable that the outer diameter of elongate tubular member 702 be as small as possible, preferably 20 mm or less, more preferably 15 mm or less and, in some embodiments, 12 mm
10 or less. The embodiment illustrated in **Figure 7A**, illustrates actuating mechanisms used to operate the device, namely first trigger 710, second trigger 711, slider 712, and plunger 714 provided in connection with handle assembly 706. Also shown is rotating collar 716 that allows the orientation of handle assembly 706 to be independently adjusted by the user relative to the orientation of approximating tool assembly 708.

15 A close up cross sectional view of the distal end of device 700 is shown in **Figure 7B**, illustrating details of multi-functional tool assembly 708 in the collapsed state. Multi-functional tool assembly 708 combines substantially similar structural and functional elements as previously illustrated in and described with reference to **Figures 5 and 6**. Accordingly, in this configuration, located along longitudinal axis 718 of working channel
20 704 are two (or more) extendible members 720, and (optional) pushing member 722, each being operatively connected to actuating mechanisms accessible to an operator at handle assembly 706. Each of the extendible members 720 is configured at its distal end with a distal tip 724, and each distal tip 724 incorporates one or more tissue engagement mechanisms whose working function is to controllably and selectively grasp, grab, grip, pierce, hold or otherwise engage tissue. In the example shown, distal tips 724 incorporate tissue hooks 726. Box-type staples in pre-deployed state 730 are loaded into working channel 704 and are configured (using, for example, guide channels and a spring loading mechanism) to slidably move toward the distal end of multi-functional tool assembly 708 and into the pre-fire position 731 as staples are sequentially ejected from the device. Pistons 732
25 are positioned at the distal end of shaft 733, and, along with stationary anvil 734, are used to deform staple legs 735 and thereby reconfigure and eject the staples when the device is actuated by the user, as described below.
30

35 **Figure 7C** illustrates a close up view of multi-functional tool assembly 708 having extendible members and tissue engagement mechanisms in the extended state, i.e. after being deployed by the operator. In the embodiment illustrated, plunger 714 is operatively

5 connected to extendible members **720**, such that when plunger **714** is pushed into handle assembly **706**, extendible members **720** move distally along longitudinal axis **718** and thereby extend outwardly from working channel **704** and beyond the end of elongate tubular member **702**. During deployment to the extended state, each of extendible members **720** is positioned such that distal tips **724** are spaced apart from one another and positioned on
10 opposite sides of longitudinal axis **718**. In the example shown, extendible members **720** have a flattened cross sectional configuration to increase lateral stiffness and prevent undesirable out-of-plane bending during deployment. Distal tips **724** of the extendible members **720** may comprise multiple tissue hooks **726**, which facilitate secure tissue engagement and help to prevent undesired out-of-plane bending of extendible members **720** during deployment. Both
15 the longitudinal positioning **719** and spacing **721** of arm tips **724** may be selectively controlled by the user to facilitate the desired positioning of tissue engagement members **726** and the subsequent size and position of the tissue plication formed by approximating the tissue.

Device **708** additionally incorporates pushing member **722**, which is operatively
20 connected to slider **712**, such that when slider **712** is pushed from its proximal (fully retracted) position, the distal end of pushing member **720** moves along longitudinal axis **718**, thereby extending out of working channel **704** a user selectable distance **705** beyond the end of elongate tubular member **702**. The pushing member facilitates invagination and folding of the tissue between the engaged portions and may, additionally, function to disrupt the serosal
25 tissue surface, or facilitate application of a tissue bonding promoter, as described above. Operation of the pushing member may be independent of, or coordinated with, extension and retraction of the extendible members and tissue engagement mechanisms..

The steps of deploying device **700**, engaging tissue, and invaginating and approximating tissue to create a tissue fold are substantially similar to what was previously
30 described with reference to **Figures 5D-5F**. For the sake of clarity, these sequential steps are again illustrated in **Figures 7D-7F** with reference to operation of multi-functional tool assembly **708**. After the tissue has been approximated and the fold has been created, device **700** is positioned in a suitable location for the subsequent step of applying one or more fasteners to secure the plication. Accordingly, similar to corresponding **Figure 6C**, **Figure**
35 **7G** illustrates the distal portion of device **700** after the device has been actuated from within

5 handle assembly 706 using a second trigger 711, which is operatively connected to extendible shaft 733. The actuation, as described previously, forms and ejects a box-type staple, reconfiguring it by deformation from the pre-deployed state 730 to the deployed state 736, and securely implanting the staple within the tissue as described previously. This results
10 in penetration and pulling together of the opposing tissue shoulders 765, which thereby secures the created tissue plication 790 projecting into the gastrointestinal space. Tissue hooks 726 may then be operatively disengaged from the tissue using a slight forward actuation of plunger 714 located within handle assembly 706, after which extendible members 720 may be completely retracted back into the shaft of the device by full reverse actuation of plunger 714. Pushing member 722 may also be completely retracted back into
15 the device, using reverse actuation of slider 712. The serosal tissue layer may be treated to promote bonding during manipulation of the pushing member, as discussed previously. The next in line pre-loaded staple in the pre-deployed state 730 automatically (for example, via spring pressure) moves into the pre-fire position 731, and the device is therefore fully prepared and ready for repeating the entire sequence at the next tissue location selected by
20 the operator, as shown in **Figure 7G**.

As illustrated in **Figure 7H** (substantially similar to **Figure 6D**), after repeating the procedural steps described above using multi-functional tool assembly 708, a plurality of staples in the deployed state 736 are implanted into and across tissue shoulders 765, securing plication 790 projecting into the gastrointestinal space. One or more such plications may be
25 produced in this manner, each having the desired length, depth, etc., and each having a selectable number of implanted fasteners, fastener depth, fastener-to-fastener spacing, and so on, as previously described. Using the devices of the present invention in this manner, the operator is therefore able to achieve the desired gastric reduction laparoscopically and without ever needing to fully penetrate the gastric wall or otherwise compromise the internal
30 mucosal tissue layer.

Figure 8A illustrates a close up view of the distal end of a tissue approximation device according to another embodiment of the present invention. In this case, the device is a handheld instrument that is designed and operates similarly to device 700, and incorporates multi-functional tool assembly 808 at a distal end of the shaft. Multi-functional tool assembly 808 is similar to multi-functional tool assembly 708 described above, with the

5 notable exception that the fasteners used in this embodiment are helical fasteners, shown as helical fastener **810** in **Figure 8A**, as an alternative to the box-type staple described previously. Helical fastener **810** may be formed from wire having desirable characteristics (e.g. strength, stiffness, surface finish, anti-friction coatings, drug eluting coatings, and so on) and includes body **812**, sharpened leading tip **814** and proximal end **816**. Fastener body **812**
10 may have one or more screw- or coil-type turns, and is additionally characterized by length **818** and diameter **820**, which may be optimized according to the desired depth and width of tissue penetration desired for various interventional procedures. Length **818** is preferably between 1 mm and 50 mm, more preferably between 2 mm and 40 mm and, in many embodiments, between 3 mm and 30 mm. Diameter **820** is preferably between 1 mm and 20
15 mm, more preferably between 2 mm and 15 mm and, in many embodiments, between 3 mm and 12 mm. Sharpened tip **814** is configured to aid in tissue penetration during deployment. Proximal end **816** is typically configured to allow operative engagement directly or indirectly to a rotating shaft located within the working channel of the elongate tubular member of device **800**, such that when rotatably actuated from within the handle assembly, the helical
20 fastener rotates as it exits the distal end of the device, thereby penetrating the tissue. Helical fastener **810** may be fabricated from any suitable biocompatible material known in the art, for example stainless steel, Ti, NiTi, or the like may be used, as well as other materials such as polymers, ceramics, and combinations of the foregoing.

In using the device illustrated in **Figures 8A-8D**, the steps of deploying the device,
25 engaging tissue and approximating tissue to create a tissue fold are substantially identical to what was described above regarding device **700**, and illustrated in **Figures 7A-7H**. After the tissue fold has been created, multi-functional tool assembly **808** is in position and ready to apply the securing means, as illustrated on **Figure 8B**. **Figure 8C** shows multi-functional tool assembly **808** immediately after helical fastener **810** has been applied to the tissue fold
30 to produce plication **830**, illustrating the preferred placement location and orientation of helical fastener **810** between tissue shoulders **840**. It is important that diameter **818** of helical fastener **810** be sized appropriately relative to the thickness of the tissue, and that proper orientation of the device is maintained (i.e. substantially perpendicular to the tissue surface and co-planar with the opposing tissue surfaces within the tissue fold), such that tissue on
35 both sides of the tissue fold are repeatedly and consistently engaged as the helical fastener is

5 deployed into the tissue during actuated rotation of device **800**. Preferably, diameter **818** is approximately comparable to tissue thickness **850**, more preferably it is between 0.5x and 1.5x tissue thickness **850**, but in any case it is most preferably maintained at less than twice the tissue thickness **850** to avoid penetration completely through the stomach wall. Similar to **Figure 7H**, **Figure 8D** shows plication **830** projecting into the gastrointestinal space that
10 was produced as a result of the repeated placement of device **800** and actuation of multi-functional tool assembly **808**, wherein a plurality of helical fasteners **810** have been applied, as described previously.

There are advantages to using helical fasteners as securing means in methods and devices of the present invention. The mechanisms incorporated into devices for loading, feeding and deploying helical fasteners into the target tissue are simple to construct (e.g. few moving parts), compact, reliable, and easy to use. In general, helical fasteners require only rotation for deployment, and they don't necessarily involve reconfiguration from a pre-deployed state to a deployed state, as in the case of spring-type or deforming-type fasteners. Also, helical fasteners may be deployed such that the fastener repeatedly engages tissue at
20 multiple points of contact over a relatively large surface area on the opposing tissue surfaces. This leads to effective load distribution and tends to reduce the maximum forces generated on both the tissue and fastener, resulting in less likelihood that either the tissue or the fasteners will fail. The use of helical fasteners may thus increase the mechanical robustness of the plication produced and improve the long-term prognosis for a successful interventional
25 outcome.

In certain situations, it may be desirable and advantageous to (optionally) provide additional reinforcement to the opposing tissue surfaces within the tissue fold and resulting plication. Such additional reinforcement not only results in stronger securement of the plication and greater load distribution, but it may also provide stabilization against
30 undesirable or excessive tissue motions, more intimate serosa-to-serosa contact and bonding, and increased rigidity to the gastrointestinal lumen (which may reduce the amount of stretching that occurs during digestion). Additional reinforcement may be accomplished using the methods and devices of the present invention by applying additional fasteners at a location within the plication as it is being produced, as illustrated in **Figures 9A** and **9B**.
35 **Figure 9A** illustrates that multi-functional tool assembly **808** has been used to place first

5 helical fastener **910**, creating first plication **920** having depth **925**, using the procedures described previously. Next, rather than move to the next tissue location to repeat the procedure (e.g. as shown in **Figure 8D**), multi-functional tool assembly **808** is instead maintained at substantially the same tissue location and tissue approximation is repeated a second time, creating a second tissue fold directly over top of the initial plication. As shown
10 in **Figure 9B**, a second helical fastener **930** is then applied, thereby producing extended plication **940** having depth **945** (greater than depth **925**), and having first helical fastener **910** completely inside the plication, acting as an additional securing means interior to the plication. This procedure may be repeated as many times as desired by the operator, resulting in the successive placement of interior fasteners and extension of the depth of the
15 plication. Beyond the stated benefits of the additional interior fasteners, a significant advantage of building up the plication depth in this manner is that the maximum designed working span of the device (e.g. spacing **721** of arm tips **724** in **Figure 7C**) may be reduced, resulting in a more compact and reliably operating device.

As will be obvious to those skilled in the art, the concept of providing additional
20 reinforcement to a plication through placement of interior securing means can be extended according to other embodiments of the present invention. For example, **Figure 10** illustrates a cross sectional view of a laparoscopically produced plication **1010** projecting into the gastrointestinal space that was created entirely extragastrically using multi-functional tool assembly **808**. A plurality of helical fasteners **810** have been placed at various locations
25 along the length and depth of the plication, thereby ensuring substantially intimate serosa-to-serosa contact over substantially the entire tissue contact area inside the plication. In addition, using the devices of the present invention, the surgeon has complete flexibility while performing the procedure to accommodate natural patient-to-patient anatomical variations in organ shape, tissue thickness, texture, presence of defects, and the like.

30 In another embodiment of the present invention illustrated in **Figure 11**, device **1100** is substantially similar in many functional aspects to the previously described devices. Device **1100** has elongate tubular member **1102** having handle assembly **1106** at its proximal end and multi-functional tool assembly **1108** at its distal end. Handle assembly **1106** further comprises the various actuating means that are operatively connected to and useful for
35 controlling the extendible elements of multi-functional tool assembly **1108**, namely first

5 trigger **1110** (used for actuating retraction of extendible members), second trigger **1112** (used for actuating deployment of fasteners), slider **1114** (used for actuating the pushing member), and plunger **1116** (used for actuating deployment of the extendible members). Rotating collar **1118** permits handle assembly **1106** to pivot around the longitudinal axis **1120** of elongate tubular member **1102** in a user selectable fashion.

10 In device **1100**, at least one multi-functional tool assembly **1108** is operatively connected to the distal end of elongate tubular member **1102** at articulating joint **1122**. Articulating joint **1122** incorporates a flexible coupling along elongate tubular member **1102**, as well as flexible internal components that operatively connect the actuating mechanisms of handle assembly **1106** to multi-functional tool assembly **1108**. This feature allows multi-functional tool assembly **1108** to be adjustably positioned by the user at tip angle **1124** relative to longitudinal axis **1120**, as shown. Preferably, tip angle **1124** is adjustable between 0 and ± 90 degrees and, in some embodiments, tip angle **1124** is adjustable between 0 and ± 60 degrees, while in yet other embodiments, tip angle **1124** is adjustable between 0 and ± 45 degrees. Any type of articulating joint design known to those skilled in the art may be used, 15 e.g. hinge joints, ball joints, universal joints, bellows joints, and the like, may be used. In the example shown, articulating joint **1122** allows multi-functional tool assembly **1108** to pivot around a single axis perpendicular to longitudinal axis **1120**, meaning that tip angle **1124** can be adjusted only within a fixed plane. For convenience, in **Figure 11** this is shown as the plane of handle assembly **1106**; however, since handle assembly **1106** can rotate around 20 longitudinal axis **1120** (by adjusting rotating collar **1118**), the operator has complete relational control between handle position and distal tip orientation, which is extremely useful for rapid, safe and efficient device operation. While a single articulating joint and multifunctional tool assembly is illustrated in **Figure 11**, it will be appreciated that multiple multifunctional tool assemblies and multiple articulating joints may be provided in 25 30 interventional tools of the present invention.

It will be appreciated that while methods and devices of the present invention have been described specifically with reference to reducing gastric volume by invaginating and approximating a wall of the gastrointestinal tract to create at least one plication therein, there are many other applications for both methods and devices of the present invention. More generally, methods and devices of the present invention may be used to approximate and,

- 5 optionally, fasten two tissue locations, and may be used in connection with a wide variety of tissue sites, and all of these applications are encompassed by the methods and devices of the present invention.

WE CLAIM:

1. A method for reducing gastric volume comprising: accessing an external surface of a wall of the gastrointestinal tract; invaginating and approximating the wall of the gastrointestinal tract from a location in proximity to its external surface to create at least one plication therein; and fastening external surfaces of the gastrointestinal wall to one another to secure the plication(s).
2. A method of claim 1, comprising invaginating and approximating the wall of the gastrointestinal tract from a location in proximity to its external surface by drawing at least one portion of an external surface of the gastrointestinal tract toward another portion of the external surface to form a plication extending into the interior space of the gastrointestinal tract.
3. A method of claim 1, comprising invaginating and approximating the wall of the gastrointestinal tract from a location in proximity to its external surface by drawing at least two portions of external surfaces of the gastrointestinal tract toward one another to form a plication extending into the interior space of the gastrointestinal tract.
4. A method of claim 1, additionally comprising penetrating fewer than all of the layers of the gastro-intestinal wall to fasten the at least two portions of the external surfaces of the gastrointestinal wall to one another.
5. A method of claim 4, comprising fastening the at least two portions of the external surfaces of the gastrointestinal wall to one another by penetrating at least the serosal layer of the gastrointestinal wall.
6. A method of claim 4, comprising fastening the at least two portions of the external surfaces of the gastrointestinal wall to one another by penetrating at least the serosal and muscularis layers of the gastrointestinal wall.
7. A method of claim 1, additionally comprising treating an external surface of the gastrointestinal tract between the at least two portions forming the plication to promote bonding.
8. A method of claim 7, additionally comprising administering an agent that promotes bonding of an external surface of the gastrointestinal tract between the at least two portions forming the plication.

9. A method of claim 1, additionally comprising administering an agent that promotes bonding of an external surface of the gastrointestinal tract between the at least two portions forming the plication.

10. A method of claim 1, additionally comprising applying energy that promotes bonding to an external surface of the gastrointestinal tract between the at least two portions forming the plication.

11. A method of claim 1, wherein at least a first tissue engagement mechanism and a second tissue engagement mechanism are positioned at a first tissue site and at a second tissue site, respectively, on the wall of the gastrointestinal tract and the wall is invaginated and approximated by moving the two tissue engagement members with respect to one another.

12. A method of claim 11, wherein the first and second tissue engagement mechanisms are positioned at the first and second tissue sites by moving them from a collapsed gastrointestinal tract access condition to an extended tissue engagement condition.

13. A method of claim 11, wherein at least one of the first and second tissue engagement mechanisms are associated with at least one extendible member, and the wall is invaginated and approximated by retracting the at least one extendible member.

14. A method of claim 1, wherein the fastening involves application of at least one fastener selected from the group consisting of: sutures, staples, screws, tacks, clips, hooks, clamps, t-tags, and helical fasteners.

15. A method of claim 1, wherein the fastening involves application of at least one agent that promotes tissue bonding.

16. A method for reducing gastric volume comprising: accessing an external surface of a wall of the gastrointestinal tract; engaging the external wall of the gastrointestinal tract at a first tissue site and at a second tissue site spaced apart from the first tissue site; manipulating tissue at the first and/or second tissue sites to approximate the first and second tissue sites and form a fold in the gastrointestinal tract; and fastening tissue in proximity to the approximated first and second tissue sites to secure the fold.

17. A method according to claim 16, wherein manipulating tissue at the first and/or second tissue sites involves moving tissue engaged at the first or second tissue sites toward tissue engaged at the other tissue site.

18. A method according to claim 16, wherein manipulating tissue at the first and/or second tissue sites involves moving tissue at both the first and second tissue sites toward tissue engaged at the other tissue site.

19. A method for reducing gastric volume comprising: accessing an external surface of a wall of the gastrointestinal tract; manipulating tissue at a first tissue site and/or at a second tissue site spaced from the first tissue site to approximate the first and second tissue sites and form a fold in the gastrointestinal tract; fastening tissue in proximity to the approximated first and second tissue sites to secure the fold; manipulating tissue at a third and/or a fourth tissue site spaced apart from one another and spaced on opposite tissue surfaces with respect to the fold to approximate the third and fourth tissue sites and form an extension of the fold; and fastening tissue in proximity to the approximated third and fourth tissue sites to secure the extension of the fold.

20. A method according to claim 19, wherein the fastening involves application of at least one fastener selected from the group consisting of: sutures, staples, screws, tacks, clips, hooks, clamps, t-tags, and helical fasteners.

21. A method for engaging, approximating and fastening tissue, comprising: accessing a tissue surface; engaging tissue at a first tissue site using a first tissue engagement mechanism associated with a tissue approximation tool; repositioning the first tissue engagement mechanism and the first tissue site to a second tissue location to approximate tissue at the first and second tissue sites; engaging tissue at the second tissue location using a second tissue engagement mechanism associated with the tissue approximation tool; and fastening tissue at the first and second tissue sites to secure the first and second tissue sites to one another.

22. A method for treating obesity in a patient, comprising: accessing an external surface of a wall of the patient's gastrointestinal tract; invaginating and approximating the wall of the patient's gastrointestinal tract from a location in proximity to its external surface to create at least one plication therein; and fastening external surfaces of the gastrointestinal wall to one another to secure the plication(s) and thereby reduce the volume of the gastrointestinal tract.

23. A device for engaging and approximating tissue comprising: an approximating tool assembly adapted for insertion into a body cavity and having at least two tissue engagement mechanisms; at least one extendible member adjustable between a collapsed condition and an extended condition and having at least one associated tissue engagement mechanism; and an actuating mechanism operatively connected to the approximating tool assembly and adapted to position at least one of the two tissue engagement mechanisms at a tissue site remote from another tissue engagement mechanism.

24. A device of claim 23, comprising at least two extendible members adjustable between a collapsed condition and an extended condition, each of the extendible members having at least one associated tissue engagement mechanism.

25. A device of claim 23, additionally comprising an actuating mechanism operatively connected to the approximating tool assembly and adapted to move the at least two tissue engagement mechanisms relative to one another to approximate tissue engaged by the tissue engagement mechanisms.

26. A device of claim 25, wherein a common actuating mechanism is used to position at least one of the two tissue engagement mechanisms at a tissue site remote from another tissue engagement mechanism and to move the two tissue engagement mechanisms relative to one another to approximate tissue engaged by the tissue engagement mechanisms.

27. A device of claim 23, wherein the tissue engagement mechanisms are selected from the group consisting of clamps, grippers, forceps, jaws, hooks, barbs, vacuum ports, and combinations thereof.

28. A device of claim 23 additionally comprising a longitudinal shaft, wherein the at least one extendible member is substantially retained within the longitudinal shaft in a collapsed condition.

29. A device of claim 28, wherein the at least one extendible member is adjustable to an extended condition by moving it in relation to a distal opening in the longitudinal shaft.

30. A device of claim 28, wherein the at least one extendible member moves away from the axis of the longitudinal shaft as it moves in relation to a distal opening in the longitudinal shaft.

31. A device of claim 23, wherein the at least one extendible member is formed from a highly elastic material and assumes a predetermined configuration upon adjustment to an extended condition.

32. A device of claim 23, wherein the at least one extendible member has a non-circular cross-sectional profile.

33. A device of claim 23, wherein each of the at least two tissue engagement mechanisms comprises at least one sharpened tip for penetrating tissue.

34. A device of claim 33, wherein the at least one sharpened tip on each of the tissue engagement mechanisms is configured to penetrate fewer than all of the layers of the gastric wall.

35. A device of claim 23, wherein the tissue engagement mechanisms are configured to penetrate fewer than all of the layers of the gastric wall.

36. A device of claim 23, wherein the at least one extendible member comprises a tether.

37. A device of claim 23, additionally comprising a longitudinal shaft and at least two extendible members, wherein the at least two extendible members are substantially retained within the longitudinal shaft in a collapsed condition, and additionally comprising a pusher member operatively connected to the approximating tool assembly and adapted to move along an axis of the longitudinal shaft.

38. A device of claim 37, wherein axial movement of the pusher member is coordinated with movement of the extendible members.

39. A device of claim 23, additionally comprising a tissue treatment mechanism positionable to administer a treatment to tissue lying between at least the tissue sites engaged by the tissue engagement mechanisms.

40. A device of claim 39, wherein the tissue treatment mechanism disrupts the tissue surface.

41. A device of claim 39, additionally comprising a longitudinal shaft and a pusher member adapted to move along an axis of the longitudinal shaft, wherein the tissue treatment mechanism is provided on the pusher mechanism.

42. A device of claim 39, wherein the tissue treatment mechanism comprises at least one port for administration of a tissue treatment agent.

43. A device of claim 23, additionally comprising at least one open longitudinal channel for passage of accessory devices or tools.

44. A device for engaging, approximating and fastening tissue comprising: an approximating tool assembly adapted for insertion into a body cavity and having at least two extendible members adjustable between an insertion condition and an extended condition, each of the extendible members comprising at least one tissue engagement mechanism; an actuating mechanism adapted for positioning at least one of the two tissue engagement mechanisms at a tissue site remote from another tissue engagement mechanism; an actuating mechanism adapted to move the two tissue engagement mechanisms relative to one another to approximate tissue engaged by the tissue engagement mechanisms; and a fastening tool for fastening approximated tissue.

45. A device according to claim 44, wherein the fastening tool comprises at least one port for administration of an agent that promotes bonding of tissue.

46. A device according to claim 44, wherein the fastening tool comprises an actuator for deploying fasteners to the approximated tissue.

47. A device according to claim 44, wherein the fastening tool stores and deploys a single fastener.

48. A device according to claim 44, wherein the fastening tool stores and deploys multiple fasteners without reloading.

49. A device according to claim 44, which is adapted to receive a cartridge loaded with a plurality of fasteners and the fastening tool has a mechanism for feeding and deploying multiple fasteners sequentially.

50. A device according to claim 44, wherein the fasteners are selected from the group consisting of: sutures, staples, screws, tacks, clips, hooks, clamps, t-tags, and helical fasteners.

51. A device according to claim 44, additionally comprising at least one open longitudinal channel for passage of accessory devices or tools.

52. A device for engaging and approximating tissue comprising: an approximating tool assembly adapted for insertion into a body cavity and having at least two tissue engagement mechanisms positioned in proximity to a distal, insertion end of the tool; and a handle assembly provided remotely from the approximating tool assembly and comprising an actuator operatively connected to the approximating tool assembly.

53. A device of claim 52, additionally comprising an elongate tubular member provided between the handle assembly and the approximating tool assembly, wherein the approximating tool assembly is operatively connected to the elongate tubular member at an articulating joint.

54. A device of claim 53, wherein the approximating tool assembly is positionable at an angle of between 0 and ± 90 degrees relative to the axis of the elongate tubular member.

55. A device of claim 52, additionally comprising an elongate tubular member provided between the handle assembly and the approximating tool assembly, wherein the handle assembly is movable around a longitudinal axis of the elongate tubular member in a user selectable fashion.

56. A device of claim 52, additionally comprising a fastening tool for fastening approximated tissue.

1/24




FIG. 1A




FIG. 1B

FIG. 1A1




FIG. 1A2




FIG. 1B1















FIG. 1B2

FIG. 1A3

2/24



3/24





FIG. 3A



X-X (2x)

FIG. 3B




FIG. 4A




FIG. 4B

4/24




FIG. 5A

5/24




FIG. 5B

6/24




FIG. 5C

7/24




FIG. 5D

8/24




FIG. 5E

9/24




FIG. 5F

10/24




FIG. 6A

11/24





FIG. 6B

12/24**FIG. 6C**

13/24




FIG. 6D

14/24






FIG. 7A

15/24**FIG. 7B**

16/24708**FIG. 7C**

17/24

FIG. 7D




FIG. 7E







FIG. 7F



18/24**FIG. 7G**

19/24**FIG. 7H**

20/24









FIG. 8A

21/24**FIG. 8B****FIG. 8C****FIG. 8D**

22/24**FIG. 9A****FIG. 9B**

23/24





FIG. 10

24/24**FIG. 11**

专利名称(译)	用于减少胃容积的方法和装置		
公开(公告)号	EP2129301A4	公开(公告)日	2015-03-25
申请号	EP2008743869	申请日	2008-03-13
[标]申请(专利权)人(译)	哈里斯PETER小号 拉宾BARRY HAL		
申请(专利权)人(译)	哈里斯 , PETER S. 拉宾BARRY HAL		
当前申请(专利权)人(译)	哈里斯 , PETER S. 拉宾BARRY HAL		
[标]发明人	HARRIS PETER S RABIN BARRY HAL		
发明人	HARRIS, PETER S. RABIN, BARRY HAL		
IPC分类号	A61B17/08 A61B17/00 A61B17/064 A61B17/068 A61B17/10 A61F5/00		
CPC分类号	A61F5/0086 A61B17/00234 A61B17/0644 A61B17/0684 A61B17/083 A61B17/10 A61B2017/00349 A61B2017/00818 A61B2017/0649 A61F5/0089		
优先权	60/894626 2007-03-13 US		
其他公开文献	EP2129301A2		
外部链接	Espacenet		

摘要(译)

本发明涉及用于减少胃容量从而治疗肥胖症的新的介入方法。该手术通常通过腹腔镜进行，并且通常可描述为腹腔镜折叠胃成形术 (LPG)，其中在获得腹部通路后，胃壁上的间隔开的部位被接合并且近似以产生一个或多个组织褶皱，然后将其固定以产生一个或多个褶皱伸入胃肠道空间。可任选地在手术期间处理浆膜组织以促进形成强的浆膜 - 浆膜结合，从而确保组织折叠的长期稳定性。这些方法优选完全经胃外进行（即不穿透胃肠壁），从而使严重并发症的风险最小化。还公开了用于逆转该过程的方法。