

(19) 日本国特許庁(JP)

(12) 公表特許公報(A)

(11) 特許出願公表番号

特表2005-504512

(P2005-504512A)

(43) 公表日 平成17年2月17日(2005.2.17)

(51) Int. Cl. ⁷	F I	テーマコード (参考)
C 1 2 N 15/02	C 1 2 N 15/00	2 G O 4 5
C O 7 K 16/18	C O 7 K 16/18	4 B O 2 4
C 1 2 P 21/08	C 1 2 P 21/08	4 B O 6 4
G O 1 N 33/53	G O 1 N 33/53	4 H O 4 5
G O 1 N 33/577	G O 1 N 33/577	
	審査請求 未請求 予備審査請求 有	(全 100 頁) 最終頁に続く

(21) 出願番号	特願2002-585482 (P2002-585482)	(71) 出願人	503096568 ダコサイトメーション・デンマーク・アク ティーゼルスカプ D a k o C y t o m a t i o n D e n m a r k A / S
(86) (22) 出願日	平成14年4月25日 (2002. 4. 25)	(74) 代理人	100095832 弁理士 細田 芳徳
(85) 翻訳文提出日	平成15年10月24日 (2003. 10. 24)	(72) 発明者	ジェンセン, スジー, スティーン デンマーク国 グロストラップ ディーケ イー-2600 ラーゲヴェイ 20
(86) 国際出願番号	PCT/DK2002/000267	(72) 発明者	ビー, カート デンマーク国 バレラップ ディーケー イー-2750 ソエガーズヴェイ 2
(87) 国際公開番号	W02002/088185		最終頁に続く
(87) 国際公開日	平成14年11月7日 (2002. 11. 7)		
(31) 優先権主張番号	PA 2001 00656		
(32) 優先日	平成13年4月26日 (2001. 4. 26)		
(33) 優先権主張国	デンマーク (DK)		

(54) 【発明の名称】 p a n 特異的モノクローナル抗体

(57) 【要約】

本発明は、ヒトの血液サンプル中におけるグリケーションされたヘモグロビンの全ての構造バリエーションの量を測定するためのpan 特異的モノクローナル抗体を記載する。ある一人の個体の血液中におけるヘモグロビンのグリケーションの割合を測定することによって、糖尿病患者におけるグルコースレベルコントロールの有用な指標が提供される。

【特許請求の範囲】

【請求項 1】

式：

グリコシル-(NH)-Val-His-またはグリコシル-(NH)-Val-His-AA、

式中、グリコシル-(NH)-Val は非酵素的にグリケーションされたバリン残基を表し、His はヘモグロビンの天然型 - 鎖の 2 番目のアミノ酸を表し、AAは 1 ~ 3 個のアミノ酸残基を意味する

のグリケーションされたペプチド残基と特異的に結合する抗体結合部位を含有してなる、pan 特異的である、すなわちグリコシルヘモグロビンの全ての構造バリエーションと反応することができる、モノクローナル抗体またはその断片。

10

【請求項 2】

AAが、ヒトヘモグロビンの全ての構造バリエーションのN-末端の - 鎖の一部に対応する 1 ~ 3 個のアミノ酸配列である請求項 1 記載のモノクローナル抗体またはその断片。

【請求項 3】

AAが、-Leu-Aa、-Leu-Thr-Aa、-Leu-Thr-Pro-Aa であり、Aaが、AA基中の残りの数のアミノ酸を意味する請求項 2 記載のモノクローナル抗体またはその断片。

【請求項 4】

前記抗体が、Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) にアクセス番号DSM ACC2495 として寄託されているハイブリドーマDAK Hb1c-1により得ることができる前記請求項いずれか 1 つに記載のモノクローナル抗体またはその断片。

20

【請求項 5】

前記抗体が、pan 特異的、すなわちHbA1c、HbC1c およびHbS1c を含むヒトヘモグロビンの全ての構造バリエーションと反応することができる、前記請求項いずれか 1 つに記載のモノクローナル抗体またはその断片。

【請求項 6】

前記抗体が請求項 7 ~ 11 いずれか 1 つに記載の方法により得ることができる、前記請求項いずれか 1 つに記載のモノクローナル抗体またはその断片。

【請求項 7】

抗体が免疫原性キャリア物質に化学的に連結されたグリケーションされたペプチドにより得ることができる抗原を含有する免疫原に対して惹起され、該グリケーションされたペプチドが、ヒトヘモグロビンのN-末端の - 鎖に対応する 2 ~ 5 個のアミノ酸単位を有する、モノクローナル抗体またはその断片の産生方法。

30

【請求項 8】

免疫原が、式：

[フルクトシル-(NH)-Val-His-AA-R] n-キャリア

式中、フルクトシル-(NH)-Valは非酵素的にグリケーションされたバリン残基を表し、His はヘモグロビンの天然型 - 鎖の 2 番目のアミノ酸を表し、AAは 1 つ以上のさらなるアミノ酸残基を意味し、R は結合基であり、キャリアは免疫原性キャリア物質であり、n は 1 ~ 該キャリア上で利用可能なカップリング部位の数の整数である、

である請求項 7 記載のモノクローナル抗体またはその断片の産生方法。

40

【請求項 9】

AAが、ヒトヘモグロビンの全ての構造バリエーションのN-末端の - 鎖の一部に対応する 1 ~ 3 個のアミノ酸配列である請求項 8 記載のモノクローナル抗体またはその断片の産生方法。

【請求項 10】

前記キャリアが、ヒトヘモグロビン以外の免疫原性タンパク質またはペプチドである請求項 8 記載のモノクローナル抗体またはその断片の産生方法。

【請求項 11】

抗体が、ヒトヘモグロビンA1c、C1c およびS1c の - 鎖のグリケーションされたN-末端

50

ペプチド配列への立体構造的なアクセスを提供するのに十分露出された後のヒトヘモグロビンA1c、C1cおよびS1cの - 鎖のグリケーションされたN-末端ペプチド配列に特異的に結合する前記請求項いずれか1つに記載のモノクローナル抗体またはその断片の使用法。

【請求項12】

前記グリケーションされたペプチド配列が、物理的または化学的な変性または消化により抗体結合部位に対して露出されている請求項11記載のモノクローナル抗体またはその断片の使用法。

【請求項13】

前記グリケーションされたペプチド配列が、カオトロピック剤を用いた変性により抗体結合部位に対して露出されている請求項11または12記載のモノクローナル抗体またはその断片の使用法。

【請求項14】

カオトロピック剤が、グアニジン、尿素、チオシアン酸カリウムまたは界面活性剤からなる群より選ばれる請求項13記載のモノクローナル抗体またはその断片の使用法。

【請求項15】

前記抗体が、固相に吸着されているヒトヘモグロビンの全ての構造バリエーションの - 鎖のグリケーションされたN-末端のペプチド配列に特異的に結合する前記請求項いずれか1つに記載のモノクローナル抗体またはその断片の使用法。

【請求項16】

固相が、好ましくはマイクロタイプレート、微粒子、膜、チューブ、ピン、チップまたはディスクからなる群より選ばれる物質から構成される請求項15記載のモノクローナル抗体またはその断片の使用法。

【請求項17】

ヒト血液中のグリケーションされたヘモグロビンのレベルの決定のための前記請求項いずれか1つに記載のモノクローナル抗体またはその断片の使用。

【請求項18】

ヒト血液中のグリケーションされたHbS および/またはHbC のレベルの決定のための前記請求項いずれか1つに記載のモノクローナル抗体またはその断片の使用。

【請求項19】

ヒト血液中のグリケーションされたヘテロ接合型HbA、HbS および/またはHbC のレベルの決定のための前記請求項いずれか1つに記載のモノクローナル抗体またはその断片の使用。

【請求項20】

二粒子比濁アッセイにおいて測定されるグリケーションされたヘモグロビンのレベルの決定のための前記請求項いずれか1つに記載のモノクローナル抗体またはその断片の使用。

【発明の詳細な説明】

【0001】

本発明は、ヒトの血液サンプル中におけるグリケーションされたヘモグロビンの全ての構造バリエーションの量の測定方法に関する。ある一人の個体の血液におけるヘモグロビンのグリケーションの割合を測定することによって、糖尿病患者におけるグルコースレベルコントロールの有用な指標が提供される。

【0002】

特に、本発明はモノクローナル抗体またはその断片の調製方法に関するものであり、これはヒトのヘモグロビンなどのグリケーションされたN-末端ペプチド残基を特異的に認識する。

【0003】

糖尿病患者は、グルコースを通常の方法で代謝することができず、その結果、患者の血液中および尿中にグルコースが蓄積する。従来より、このような体液中のグルコースレベルは糖尿病の状況の状態の指標として取り扱われており、言い換えるとこのレベルは、服用

10

20

30

40

50

するインスリンまたは他の薬剤の量の指針として、または患者の食生活を変更する必要性の指針として用いられている。

【0004】

直近の食事の時間および内容、直近のインスリン注射などに依存して、グルコースレベルが広い範囲で変動するという点を除いては、これはそれなりに十分役に立っている。このように、示される値はその瞬間の状況を反映することになり、このことは、糖尿病の状況の長期間の状態と正確には合致しないかもしれない。

【0005】

糖尿病の状況の別の結果として、糖尿病患者の血液中のグリケーションされたヘモグロビンの量が増加するということが知られている。ヘモグロビン(Hb)は、アミノ酸の四本の鎖(サブユニット)からなる四量体タンパク質であり、鎖のそれぞれは約143単位であり、合計で約64000g/molの分子量を有する。ヘモグロビン中の α -サブユニットのN-末端のバリンは、グルコースと反応することが可能である。グルコースとバリンの α -アミノ基とに関する非酵素的な反応によって、ヘモグロビンのグリケーションが生じる。反応物の間に Schiff 塩基が形成された後、グルコースはアマドリ転位を受けて1-デオキシフルクト-バリンを形成する。最終的な複合体での1-デオキシフルクトースとバリンとの間の結合は共有結合であり、不可逆性である。グリケーション反応は、反応物、たとえばヘモグロビンとグルコースレベルに支配される。糖尿病ではない個体についての、糖尿病のコントロールと合併症のトライアル(Diabetes Control and Complication Trial)(DCCT)の基準範囲は、約4-6%のグリケーションされたヘモグロビンである。 α -鎖のN-末端上に1-デオキシフルクト-バリンを伴うヘモグロビン四量体は、グリケーションされたヘモグロビン;たとえばHb1cまたはHbA1c、HbC1cまたはHbS1cと認められる。

10

20

【0006】

HbCの特徴は、サハラ以南の地域において30%が患者であり、アフリカ系アメリカ人の2.3%が患者であることである。合衆国におけるHbの最も一般的なバリエーションであるHbSの特徴は、アフリカ系アメリカ人の7.8%が患者であることである。ひとまとめにすると、アメリカ黒人の少なくとも10%が、HbCの特徴またはHbSの特徴のいずれかを持っている。世界人口では混血がどんどん増加しつつある。したがって、グリケーションされたヘモグロビンの全ての構造バリエーションと同様に反応する抗体を持つことがますます重要となる。もし、利用可能な抗体がグリケーションされたヘモグロビンの全ての構造バリエーションとは反応しない場合、HbSおよび/またはHbCの特徴を備える患者からのHb1cの割合の測定が不正確になるだろう。

30

【0007】

糖尿病患者におけるグルコースレベルは、血液中のグルコースレベルに直接依存するグリケーションの割合を高めるのに十分に高く、これは糖尿病の状況の重篤度を反映する。それゆえに、ヘモグロビンA1cのレベルは約6%から12%に上昇する。ヘモグロビンAが循環する寿命は約120日間であることから、グリケーションされたヘモグロビンの測定には、その期間の平均的なグルコースレベルを反映するという意味があるであろう。特に、グルコースが多い食事によって、グリケーションされたヘモグロビンまたは血清アルブミンのレベルが即座に高く反映されるわけではない。したがって、グリケーションされたヘモグロビンの含有量の測定によって、循環しているグルコースの平均的なレベルのより本当に近い臨床像が与えられ、したがって患者の長期間の状況のより本当に近い臨床像が与えられる。

40

【0008】

グリケーションされたヘモグロビンレベルを測定するために用いられる一般的な方法には、溶血サンプルをボロン酸塩アフィニティカラムに通す工程が含まれる。この方法によって、HbA_{1c}と、リジン残基ならびにヘモグロビンの α -鎖および β -鎖の両者のN-末端のバリン残基上で形成されるケトアミン構造体とを含む、グリケーションされたヘモグロビンの総量を測定する。このカラムを洗浄し、そして分光光度法でグリケーションされたヘモグロビンを測定する。

50

【0009】

機器によって自動的に分析するために、微粒子強化免疫比濁法 (Microparticle enhanced turbidimetric immunoassay technique) を利用することができる。これらの機器は、ある特定の診断用アッセイのために用意したのではなく、そして極めて長いサンプルのサイクル時間に備えている。現在のところ、グリケーションされたヘモグロビンの測定のためのこの方法の実用性は、ありきたりの臨床分析として高まっている。これらの免疫アッセイのいくつかは、微粒子強化競合免疫比濁阻害法 (microparticle enhanced competitive turbidimetric inhibition immunoassay technique) の原理に基づいている。

【0010】

米国特許第4,247,533号には、HbA1cの注射によって、ヒツジの中でHbA1cに対する抗体を報告されるように惹起させたことと、グリケーションされていないヘモグロビンを吸着させることでポリクローナル抗体を供給し、これによってHbA1cとグリケーションされていないヘモグロビンとを区別するといった分析方法が開示されている。次いで、このような抗体は、サンプル中のグリケーションされたヘモグロビンの割合を測定するための試験の基礎を形成する。

10

【0011】

従来技術では、ヘモグロビンA_{1c}抗体と、その抗体がどのようにしてグリケーションされたヘモグロビンのバリエーションAと反応することができるかについてだけが記載されている。グリケーションされたヘモグロビンの他のバリエーションの影響力については言及されず、そしてグリケーションされたヘモグロビンの全ての構造バリエーションに対するモノクローナル抗体についても考慮されていない。この抗体はHbS1cおよびHbC1cを認識しないという事実が原因で、HbS1cおよびHbC1cを有する被験体において、グリケーションされたヘモグロビンのパーセンテージの免疫アッセイ測定をこの抗体を用いて行うことによって、事実上低いグリケーションされたヘモグロビンのパーセンテージが生じてしまう。

20

【0012】

Hb1cの全ての構造バリエーションについて特異的であるHb1c抗体は、pan特異的とみなされる。

【0013】

したがって、上記で認められた欠点を現さない上記の種類抗体についての必要性が残っている。

30

【0014】

pan特異的である抗体、すなわちグリケーションされたヘモグロビンの全ての構造バリエーションについて特異的な抗体を提供することが、したがって本発明の目的である。

【0015】

HbA1c、HbC1cおよびHbS1cの免疫アッセイ測定法に基づいて、患者の長期間の血液の糖レベルを測定するための抗体を提供することも目的である。

【0016】

患者の全血サンプル中のグリケーションされたヘモグロビンであるHbA1c、HbC1cおよびHbS1cの含有量を測定するための、このような抗体を提供することは、本発明の別の目的である。

40

【0017】

これらの目的および他の目的は、特許請求の範囲に規定される本発明によって達成される。

【0018】

本発明は、pan特異的、すなわちグリケーションされたヘモグロビンの全ての構造バリエーションと反応することが可能であるモノクローナル抗体またはその断片に関する。このグリケーションされたヘモグロビンの構造バリエーションとしては、HbA1c、HbC1cおよびHbS1cが好ましい。

【0019】

このモノクローナル抗体が、HbSおよびHbCを持つ患者におけるグリケーションされたヘ

50

モグロビンレベルの測定に関する現行の免疫アッセイの改良をもたらすことが好ましい。

【0020】

本発明のモノクローナル抗体は、その抗体がヒトのヘモグロビンA、CおよびSのβ鎖のグリケーションされたN-末端のペプチド配列と特異的に結合する抗体結合部位を含有するという特徴を有する。

【0021】

本発明のモノクローナル抗体試薬は、グリケーションされたヘモグロビン中の線状エピトープについてのその特異的な結合親和性に特徴を有する。したがって、本明細書で用いられるような「抗体試薬」または「抗体結合部位」という用語は、このようなペプチドのエピトープについて特異的なモノクローナル抗体結合部位を含有するあらゆる物質を示すことになり、そしてこの物質はどのようにして得られたものであってもよい。したがって、このような用語には、全ての抗体および適切な断片または多官能性形態が包含される。全ての抗体の形態である場合、たとえばIgG、IgMなどの公知の免疫グロブリンのいずれかのクラスおよびサブクラスに属することが可能である。ペプチドのエピトープについての特異的な結合親和性を有する、いずれかのそのような免疫グロブリンのあらゆる断片を用いることが可能であり、たとえば、IgGの断片は従来よりFab、Fab'およびF(ab')₂として知られている。さらに、免疫グロブリンまたはその断片の凝集体、ポリマー、誘導体、複合体(conjugates)およびハイブリッドを、適切である場合には用いることが可能である。さらに、単鎖抗体または組み換え手段によって得られるあらゆる他の断片を、適切である場合には用いることが可能である。

10

20

【0022】

グリケーションされた残基は、HbA1c、HbS1cおよびHbC1cの顕著な構造上の特徴である。本発明の抗体には、グルコースと末端のアミンとの間の反応生成物のアマドリ転位で形成される1-デオキシフルクトシル糖質単位を最低限含有するエピトープまたは決定部位と、天然型HbA1c、HbS1cおよびHbC1cの配列に対応する位置にあるHbA1c、HbS1cおよびHbC1cのN-末端の配列の少なくとも1つのアミノ酸単位を含有する、そこから伸びているペプチド配列とが必要である。エピトープを特徴付けるペプチド配列中の他のアミノ酸単位は、天然型HbA1c、HbS1cおよびHbC1c配列中に見られるアミノ酸単位と同一であってもよく、異なってもよい。このように、このエピトープには、抗体に対する少なくとも二つの接触部位または結合部位という特徴があり、このような部位はグリケーションされたN-末端のHbA1c、HbS1cおよびHbC1c配列に対して特有である。

30

【0023】

この抗体は、次の式：

グリコシル-(NH)-Val-His-またはグリコシル-(NH)-Val-His-AA、

式中、グリコシル-(NH)-Valは非酵素的にグリケーションされたバリン残基を表し、Hisはヘモグロビンの天然型β鎖の2番目のアミノ酸を表し、AAは1つ以上のさらなるアミノ酸残基である、

のグリケーションされたペプチド残基と特異的に結合することが好ましい。

【0024】

1つの好ましい態様において、AAとは、ヒトのヘモグロビンの全ての構造バリエーションのN-末端のβ鎖の一部に対応する1~3個のアミノ酸の配列である。AAとしては、-Leu-Aa、-Leu-Thr-Aa、-Leu-Thr-Pro-Aaであることが好ましく、そしてAaとは、AA基中のアミノ酸の残りの数を意味する。

40

【0025】

本発明のモノクローナル抗体またはその断片としては、ハイブリドーマDAK Hb1c-1を用いて得ることができるものが好ましく、このハイブリドーマは、Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ)にアクセッション番号DSM ACC2495として寄託されている。

【0026】

さらに本発明は、モノクローナル抗体またはその断片を産生する方法に関し、ここで、こ

50

の抗体はpan 特異的、すなわちグリケーションされたヘモグロビンの全ての構造バリエーションと反応することが可能である。

【0027】

抗体またはその断片は、特許請求の範囲に規定された方法によって得られ得るものであることが好ましい。

【0028】

本発明のモノクローナル抗体またはその断片を産生する方法は、免疫原性キャリア物質に化学的に連結させたグリケーションされたペプチドを含有する免疫原に対する抗体を惹起させる、ここで該グリケーションされたペプチドがヒトのヘモグロビンの - 鎖のN-末端に対応する2 ~ 5 個のアミノ酸単位を有する、工程を含む。

10

この免疫原は次の式：

[フルクトシル-(NH)-Val-His-AA-R]_n - キャリア

式中、フルクトシル-(NH)-Valは非酵素的にグリケーションされたバリン残基を表し、Hisはヘモグロビンの天然型 - 鎖の2番目のアミノ酸を表し、AAは1つ以上のさらなるアミノ酸残基を意味し、Rは結合基であり、キャリアは免疫原性キャリア物質であり、nは1 ~ 当該キャリア上で利用可能なカップリング部位の数までの整数である、であることが好ましい。

【0029】

1つの好ましい態様において、AAとは1つ以上のさらなるアミノ酸残基である。別の好ましい態様において、AAとは、ヒトのヘモグロビンの全ての構造バリエーションの - 鎖のN-末端部分に対応する1 ~ 3 個のアミノ酸の配列である。

20

【0030】

結合基であるRは、本質的には、都合が良くかつ安定したあらゆる構造であり得る。このような結合基であるRは、通常は、水素を除いて1 ~ 約20の間の原子を含有し、そして窒素、酸素および硫黄などのヘテロ原子を含む脂肪族鎖の形態である。様々な基を介してこのグリケーションされた残基を結合させて、メチレン、エーテル、チオエーテル、イミノなどを含む結合鎖のRを形成させることが可能である。当業者であれば、そこから免疫原を調製するために選択するところの幅広い種類の結合基を分かっているだろう。

【0031】

好ましい態様において、キャリアはヒトのヘモグロビン以外の免疫原性タンパク質またはペプチドである。

30

【0032】

最も一般的な意味において適切な免疫グロブリンの産生を刺激するために用いられる免疫原は、免疫原性キャリア物質に化学的に連結させられた1つ以上のグリケーションされたペプチド残基を含有する。グリケーションされたペプチド残基にカップリングすることが可能な官能基を有することが通常知られているあらゆるものから、この免疫原性キャリア物質を選択することが可能である。十分なサイズと免疫原性を持つ、たとえば炭水化物、多糖類、リポ多糖、核酸などの他の物質を同様に用いることも可能ではあるが、多くの場合、キャリアはタンパク質またはポリペプチドである。主として、免疫原性タンパク質および免疫原性ポリペプチドの分子量は4000g/mol ~ 10000000g/molの間であり、15000g/molを超える量であることが好ましく、そして50000g/molを超える量であることがより普通である。別の種の血流内に導入する場合、一般的に、1つの動物種から得られるタンパク質が免疫原性となる。特に有用なタンパク質は、アルブミン、グロブリン、酵素、ヘモシアニン、グルテリン、明らかに非タンパク質性の組成を有するタンパク質などである。

40

【0033】

好ましい態様において、測定されるサンプル中でタンパク質を適切に変性または消化することによって、本発明のモノクローナル抗体またはその断片に接触可能な天然型HbA1c、HbS1cおよびHbC1c分子のグリケーションされたN-末端のペプチド残基を作製する。

【0034】

別の好ましい態様において、天然由来のグリケーションされたヘモグロビン、たとえばHb

50

A1c、HbS1c および HbC1c を化学的にまたは酵素的に消化することによって、グリケーションされた断片を産生することが可能である。伝統的なカップリング手段、たとえばグルタルアルデヒドまたはカルボジイミドを用いて、キャリアにこの断片をカップリングさせることが可能であり、そしてこの複合体を免疫原として用いる。

【0035】

本発明の抗体またはその断片を、次の工程によって産生することも好ましい：

【0036】

A. ペプチド免疫原の調製

HbA、S および C の -鎖上にある N-末端のバリン残基のグリケーション部位に対する抗体を得るために、免疫原として短いペプチド（五量体）を用いることが好ましい。この抗原としては、Hb1c の四量体にシステインを付加したものの、すなわちフルクトシル-V-H-L-T-C が好ましい場合がある。

10

【0037】

B. ヒト Hb1c ペプチドによるマウスの免疫化

モノクローナル抗体を分泌するハイブリドーマを生育するために用いる一般的な方法は、当業者に十分知られている。

【0038】

実際に利用した方法の実例は、ジャーナル・オブ・イムノロジカル・メソッズ (Journal of Immunological Methods)、1980、39:285 に記載されている。

【0039】

CF1 マウスと BALB/c マウスとの交配によるメスの F1 ハイブリッドを用いることが好ましいが、他の系統のマウスを用いても構わない。この免疫化計画およびペプチド-キャリア複合体のレベルを、十分な量の抗原的に刺激されたリンパ球が形成されるように選択することができる。好ましい態様においては、Hb1c ペプチドとの複合体の状態の 50 μ g の精製タンパク質誘導体 (PPD、結核菌 (*M. tuberculosis*) 由来の変性タンパク質の、一部が未知の混合物) によって、二週間の間隔を空けて四回、腹腔経路でマウスを免疫化する。最終免疫の 3 ~ 5 日後、融合のために、動物から脾臓細胞を抽出する。

20

【0040】

他の種からのハイブリドーマによって産生されるモノクローナル抗体も好ましく用いることができる。

30

【0041】

C. 細胞融合

免疫化マウスから脾臓を無菌的に取り出し、そして脾臓細胞の懸濁液を調製する。適切な融合プロモーターの存在下で、この脾臓細胞を適切な細胞株由来のマウスミエローマ細胞と融合させる。好ましい態様において、ミエローマ細胞に対する脾臓細胞の好ましい割合は、約 20:1 ~ 約 2:1 であり、そして約 10^8 個の脾臓細胞あたり 0.5 ~ 1.5 ml の量で、適切に融合培地を用いる。

【0042】

細胞融合のために用いられるマウスミエローマ細胞は十分に知られている。好ましいマウスミエローマ細胞として、P3-X63-Ag 8.653 を用いてもよい。

40

【0043】

融合プロモーターとしては、平均分子量が 1000 ~ 4000 g/mol のポリエチレングリコールが好ましい。技術的に公知の他の融合プロモーターを用いることも可能である。好ましい態様において、平均分子量が 1500 のポリエチレングリコールを用いることができる。

【0044】

D. 融合した細胞のスクリーニング

個々に分かれている容器（たとえばマイクロタイタープレート）中で、融合していない脾臓細胞、融合していないマウスミエローマ細胞および融合化ハイブリドーマ細胞から構成される混合物を、融合していないマウスミエローマ細胞を生育させない選択培地で好ましくは希釈し、そして融合していないマウスミエローマ細胞を死滅させるのに十分な期間（

50

約1週間)培養する。薬物耐性(たとえば、8-アザグアニンに対する耐性)を有し、かつ融合していないマウスミエローマ細胞を生育させない培養培地、たとえばHAT培地を用いる。選択培地中で、融合していないミエローマ細胞は死滅する。融合していない脾臓細胞は形質転換されなかった細胞であることから、ある特定の期間(1週間)が経過すると、これらは徐々に死滅する。他方、融合した細胞は、親のミエローマ細胞の腫瘍としての性質と、親の脾臓細胞の性質との両方を有することから、この選択培地中でこれらが生存することができる。

【0045】

E.ELISAによるグリケーションされたヘモグロビンの測定

特異的な免疫性認識の原理、およびモノクローナル抗体と、抗体がそれに対して特有かつ特異的に結合する抗原性エピトープとの間の反応に基づくこの測定は好ましいものである。たとえば、ELISA試験によって認識および結合を検出してもよく、この場合、固相支持体、たとえばプラスチック製のウェルの底部に血液サンプルまたはキャリブレーターを固定する。ヒトのヘモグロビンを固定化することによって、抗体の結合のために抗原性エピトープを露出させてもよい。抗体、酵素標識化二次抗体および酵素の基質を、固定化抗原(ヒトのヘモグロビン)と相互作用させてもよく、それによって、抗体-抗原複合体を形成させることが好ましい。何回かの希釈、インキュベーション工程および洗浄工程、その後結合した試薬と遊離している試薬とを分離させる。抗体の抗原との結合と、その結果として生じる酵素の基質との相互作用反応の結果として、発色反応が生じる。色素の形成によって血液サンプル中にHb1cが存在することが示され、そして発色の強度によって、そのサンプル中のグリケーションされたエピトープの定量測定を提供する。技術的に知られている他の免疫アッセイ形式によってグリケーションされたヘモグロビンを測定するために、本発明の抗体を用いてもよい。

10

20

【0046】

F.所望の抗体を産生するハイブリドーマ細胞のクローニングとその抗体の産生

限界希釈法などの適切な方法によって、所望の抗体を産生することができるハイブリドーマ細胞をクローニングし、そして所望の抗体を次の方法によって産生することが可能である。適切な培地中で、ある特定の時間ハイブリドーマ細胞を培養し、そしてその培養物の上清からこのハイブリドーマ細胞によって産生されるモノクローナル抗体を単離することが可能である。

30

【0047】

本発明はさらに、モノクローナル抗体またはその断片を用いる方法にも関する。

【0048】

未知の量のグリケーションされたヘモグロビンを含有する血液サンプルと反応させるという通常の方法で、本発明の抗体またはその断片を好ましく用いることが可能であり、そしてグリケーションの程度を測定するために較正された標準と、反応の程度を比較することが可能である。蛍光、免疫アッセイなどによって、HbA1c、HbS1cおよびHbC1cにおけるグリケーションされたエピトープについてのそれらの結合力を失うことなく公知の方法にてモノクローナル抗体に読み取り可能な基を適切に結合させることによって、データを出力することが可能である。

40

【0049】

本発明のモノクローナル抗体は、グリケーションされたN-末端のペプチド残基の全ての構造バリエーションに、好ましくはHbA、HbSおよびHbCで見られるものに特異的に結合する。エピトープが都合良く露出している場合、この抗体は天然型HbA1c、HbS1cおよびHbC1c分子中のエピトープに結合することが可能である。エピトープに立体構造的にアクセスすることは、任意の効果的な方法で達成することが可能である。無処理タンパク質におけるエピトープの露出は、少なくともエピトープの領域中で物理的または化学的変性または消化を行うことによって達成されると理解されている。このような変性または消化をエピトープの領域に限定することが可能であり、またはこのような変性または消化には、より全体的なすなわちさらに実質的に完全な三次構造の変性、およびタンパク質の二次構造の変

50

性、あるいはタンパク質の部分消化または完全消化を包含することができる。

【0050】

好ましい態様において、この抗体は、ヒトヘモグロビンA1c、C1c およびS1c の - 鎖のグリケーションされたN-末端ペプチド配列への立体構造的なアクセスを提供するに十分な程度に露出させた後、ヒトヘモグロビンA1c、C1c およびS1c の - 鎖のグリケーションされたN-末端ペプチド配列に特異的に結合する。

【0051】

加熱、超音波処理、高pHまたは低pHへの暴露などの物理的手段によるタンパク質の通常の方法を含む様々な方法によって、変性を成し遂げることが可能であり、溶液中での薬剤またはカオトロープを用いての相互作用による化学的変性が好ましい。

10

【0052】

たとえば、グアニジンと加熱、グアニジンとSDS、またはグアニジンとジチオスレイトールのように、化学的手段および/または化学的かつ物理的手段を組み合わせることであれば、タンパク質の変性を最も効率的に成し遂げることが可能である。グアニジンのような特に強力なカオトロープが好ましい。当然のことながら、抗体の結合のための溶液に接触しやすい露出されるエピトープが含まれているわずかな量のタンパク質の実質的な不溶化、凝集または沈殿生成という結果を招く変性条件は、避けることになる。有用な免疫結合をもたらすために、溶液中または懸濁液中に十分な量の変性タンパク質が残っていなければならない。必要な可溶化の程度は、意図されるまたは所望される結合の状況に依存する。

20

【0053】

好ましい態様においては、物理的または化学的変性または消化によって、抗体結合部位に対してグリケーションされたペプチド配列を露出させる。

【0054】

別の好ましい態様においては、カオトロピック剤を用いての変性によって、抗体結合部位に対してグリケーションされたペプチド配列を露出させる。このカオトロピック剤としては、限定されないが、グアニジン、尿素および種々の界面活性剤（例えば、ドデシル硫酸ナトリウム（SDS）など）が挙げられ、他には、限定されないが、デオキシコール酸塩および特定の胆汁酸塩、3-（3-コラミドプロピル(cholamidopropyl)）-ジメチル-アンモニオ-1-プロパンスルホン酸塩、メタノール、プロパノール、アセトニトリルなどの有機溶媒、ならびにチオシアン酸カリウムなどの特定の塩が挙げられる。たとえばTriton X（登録商標）、Tween（登録商標）、nonidet NP-40（登録商標）およびオクチルグルコシドなどの非イオン性界面活性剤もタンパク質の変性剤として機能することが可能である。

30

【0055】

得られる水性混合物中であらゆるHbA1c、HbS1c およびHbC1c を変性させるのに十分なレベルのカオトロープが存在する水溶液を用いて、特定の血液サンプルの有意な量のHbA1c、HbS1c およびHbC1c を変性させて、サンプル（たとえば、全血または赤血球の溶血物）との混合による抗体結合のためにグリケーションされたエピトープを露出させる。全血がサンプルである場合には、カオトロープは赤血球を溶血させることにも役立ち、その結果、ヘモグロビンが放出され、そしてプロテアーゼが不活化される。

40

【0056】

混合物を短時間加熱することによって、変性プロセスを顕著に促進させてもよい。37 以下の温度では、カオトロープによる変性には1時間～数時間かかる可能性があり、それに対して50 よりも高い温度では、1分間かそれよりも短時間で十分な変性に到達することができることが分かった。抗体および他のタンパク質性試薬の著しい変性を防止して、その後混合物へ添加するために、別々の工程として、または、カオトロープがこのような試薬の変性を実質的に無効にし、さらに有意な程度のHbA1c、HbS1c およびHbC1c が再び天然型になるのを防ぐことによってエピトープの露出を防止するであろうレベルに試薬溶液を添加することによって、サンプル-カオトロープ混合物を希釈することが普通である。

50

【0057】

好ましい態様においては、抗体は、固相に吸着されているグリケーションされたヒトヘモグロビンの - 鎖中のグリケーションされたN-末端のペプチド配列と特異的に結合する。ここで、この固相としては、ポリスチレン、ジビニルベンゼン、ブタジエン、ポリカーボネート、ポリアクリルアミド、ポリアクリル酸、ポリアクリルアミド、ポリエチレン、ポリプロピレン、フッ素化ポリマー、ポリアミドまたはそのブロックコポリマー、金、炭素、セルロース、誘導体化セルロースまたはガラスからなる群より選択されるものが好ましい。

【0058】

好ましい態様において、ヒトの血液中のグリケーションされたヘモグロビンレベルを測定するために、モノクローナル抗体またはその断片を用いてもよい。さらに、ヒトの血液中のグリケーションされたHbS および/またはHbC レベルを測定するために、モノクローナル抗体またはその断片を用いることもまた好ましい。

10

【0059】

さらに好ましい態様において、ヒトの血液中のグリケーションされたヘテロ接合型HbA、HbS および/またはHbC レベルを測定するために、モノクローナル抗体またはその断片を用いてもよい。

【0060】

さらに好ましい態様においては、グリケーションされたヘモグロビンレベルを測定するために、二つのマイクロスフェアによる比濁法 (turbidimetric two-microsphere method) が提供される。この方法は、サンプルと、モノクローナル抗体またはその断片が結合しているマイクロスフェアとを混合する、ここで後者は免疫粒子と称される、工程、または、サンプルと、マイクロスフェアとを混合し、次いで免疫マイクロスフェアと混合する工程を含む。得られる免疫凝集物の濁度を測定する。

20

【0061】

マイクロスフェアは、好ましくは、ポリスチレン、ジビニルベンゼン、ブタジエン、ポリカーボネート、ポリアクリルアミド、ポリアクリル酸、ポリアクリルアミド、ポリエチレン、ポリプロピレン、フッ素化ポリマー、ポリアミドまたはそのブロックコポリマー (co-block polymer)、金またはシリカからなる群より選択される。

【0062】

この例の方法において用いられるマイクロスフェアは0.01~5 μm の粒子サイズを有し、これは、通常用いられる緩衝液中で分散させることによって用いられる。

30

【0063】

次の非限定的な実施例は、本発明の特定の特徴および態様を明らかにすることを目的とするものである。

【0064】

実施例 1 :

国際特許出願W086/03494号に記載されているとおりの、シェパードおよびアサトン (Shepard and Atherton) の固相法を用いて、1-デオキシフルクトシル-Val-His-AA ペプチドを合成した。フルオレニルメトキシカルボニル (Fmoc) アミノ酸を用いて、カップリング試薬としてPyBOP (ベンゾトリアゾール-1-イル-オキシ-トリス-ピロリジノ-ホスホリウムヘキサフルオロホスフェート) およびHOBt (N-ヒドロキシ-ベンゾトリアゾール) を使用して、このペプチドを合成した。

40

【0065】

トリチル基 (trt) で側鎖を保護したシステインおよびヒスチジン、ならびに第三ブチル基 (tBu) で側鎖を保護したスレオニンを用いた。ロイシンは側鎖を保護せずに用いた。

【0066】

公知の方法を修正することによって、フルクトースアミノ酸複合体であるN-(デオキシ-D-フルクトース-1-イル)-L-バリンを合成した。要約すれば、ジ-イソプロピリデンで保護されたグルコースをトリフレートとして活性化し、そしてバリンのベンジルエステル

50

と結合させた。得られた2,3:4,5-ジ-0- イソプロピリデン-N- (デオキシ-D- フルクトース-1- イル)-L- バリン- ベンジルエステルを、エタノール中のPd/C上で水素化し、凍結乾燥し、そして最後に、Fmoc保護を伴わないペプチド合成に用いた。TLC 分析および¹Hまたは¹³C のNMR スペクトルによって、生成物の純度を確認した。

【0067】

自動化固相合成 (カルバイオケム- ノババイオケム社 (Calbiochem-Novabiochem Ltd) の結晶化装置) によってこのペプチドを合成し、TFA (トリフルオロ酢酸) -TES (トリエチルシラン) - グリセロール-H₂O (90%:4%:3%:3%) によって樹脂から切断し、沈殿させ、そしてメチル- 第三ブチル- エーテルで洗浄し、遠心分離に付し、そして凍結乾燥した。

【0068】

HPLC (C₁₈ 逆相カラム、0.1%のTFA 中の典型的な5-25% のMeCNの段階的勾配を使用) によって精製を行った。MALDI-TOF 質量分析法によってこのペプチドの実態を確認し、そして分析用HPLCによる分析によって、その純度が90% よりも高いことを最終的に保証した。

【0069】

実施例 2 :

ギバ財団 (Giba Foundation) のシンポジウム1986, 119:25に記載されたとおり、最初の免疫化の四週間前に、カルメット- グラン杆菌 (BCG) ワクチンを注射して、マウスにおける免疫応答の準備を行った。

【0070】

ペプチド免疫原を作製するために、キャリア混合物のツベルクリンPPD に複合化させた。グルタルジアルデヒド、すなわちCHO-(CH₂)₃-CHO を用いて、ペプチドとキャリア (PPD) とを複合化させた。グルタルジアルデヒドは、ペプチドおよびキャリア上のアミノ基と反応する。

【0071】

したがって、これらの免疫原は次の式を有する :

フルクトシル-Val-His-Leu-Thr-Cys-CH-(CH₂)₃-CH-PPD

【0072】

実施例 3 :

CF1 マウスとBALB/cマウスとの交配によるメスのF1ハイブリッド (12週齢) をBCG-ワクチンで準備し、そして上記のペプチド-PPD複合体によって免疫化した。リン酸塩緩衝液 (PBS) 中に溶解したペプチド-PPD複合体と水酸化アルミニウム (1mg/用量) との50 μg の混合物によって、マウスの腹腔内に四回免疫化した。グリケーションされたヘモグロビンに対する高力価の抗体を有するマウスを、融合のために選択した。最終免疫化の三日後に、この免疫化マウス由来の脾臓細胞を、細胞融合のために用いた。

【0073】

選択した免疫化マウス由来の脾臓細胞およびBALB/c起源のミエローマ細胞 (P3-X63-Ag 8.653) を約3:1 の割合で混合し、そして50% のポリエチレングリコール1500 (ロッシュ・ダイアグノスティクス (Roche Diagnostics GmbH) 株式会社の製品) の存在下で融合させた (ジェイ・ダブリュ・ゴッディング (J. W. Goding) : モノクローナル抗体 : 原理と実践、アカデミック・プレス社 (Academic Press)、サンディエゴ、1996)。この融合した細胞を、10% のFCS、ヒポキサンチン、アミノプテリンおよびチミジンを含有するRPMI-1640 培地 (HAT 培地) 中に懸濁して、1 × 10⁶ 細胞/mL のレベルとした。この懸濁液を、96- 穴のマイクロプレート (コスター (Costar) 社) に、ウェル当たり200 μL の量で分配した。

【0074】

この細胞を、CO₂ インキュベーター (5%のCO₂、37 °C) 中でインキュベーションを行った。14日後、ハイブリドーマ細胞のスクリーニングを行った。

【0075】

抗原として4.7%のHbA1c および16.7% のHbA1c を含有するヘモグロビンキャリアプレートでコーティングしたマイクロタイタープレートを用いるELISA 法によって、ハイブリド-

10

20

30

40

50

マ細胞の培養物から産生された抗体を含有する上清を検出した。120 µg/mLのキャリブレーターを、炭酸塩/重炭酸塩緩衝液 (pH9.6) を用いて、37 °C で15分間かけて、プラスチック製のマイクロタイターのウェル上に固定した。結合しなかった抗原を取り除く洗浄工程の後に、0.1%のTween20 を含有する50mMのトリス-HCl緩衝液に希釈したハイブドーマの培養物の上清をそれぞれのウェルに添加し、そして室温で一時間反応させた。結合しなかった抗体を取り除く別の洗浄工程の後に、ヤギの抗マウス抗体と酵素である西洋ワサビペルオキシダーゼとの複合体を添加した。インキュベーション工程および三度目の洗浄工程の後に、基質である3,3',5,5'-テトラメチルベンジジン (TMB) との反応によって、結合している複合体を検出した。硫酸を用いてこの反応を停止させ、そして450nmにおける吸光度を測定した。グリケーションされたヘモグロビンキャリブレーターである4.7%のHbA1c および16.7%のHbA1c と反応する、強度が高くなっている陽性のウェルからの細胞だけを選択した。

10

【0076】

これらの陽性細胞を、限界希釈法によって3回クローニングした。得られたクローンであるDAK Hb1c-1を90%のFCS、10%のDMSO中に懸濁し、そして-150 °C で貯蔵した。

【0077】

実施例4:

線状抗体エピトープマッピングとは、ポリペプチド配列またはタンパク質配列に由来する、可能性のある全てのペプチドの体系的なスクリーニングである。エピトープマッピングによって、所定の抗体に対する相互作用部位を形成するアミノ酸および誘導体の直線状の広がりに関する情報が得られる。

20

【0078】

ヘモグロビンの α -鎖のグリケーションされたエピトープについてのDAK Hb1c-1の特異性は確定されている。

【0079】

1~5アミノ酸長のペプチドを含有するペプチドライブラリーを、アミノ酸のN-末端の保護基としてFmocを用いて、固相 (アビメド (Abimed) 社の膜) 上でC-末端からN-末端まで手動で合成した。ペプチド合成の詳細な説明については、実施例1に記載する。次の四つのペプチドを合成した:

【0080】

フルクトシル-Hb の α -鎖

N-末端1-デオキシフルクトシル-ヘモグロビンの α -鎖の1~5アミノ酸長のペプチドアナログ。

30

【0081】

Hbの α -鎖

N-末端ヘモグロビンの α -鎖の1~5アミノ酸長のペプチドアナログ (1-デオキシフルクトシル α -鎖を引いたもの; 基幹コントロール)。

【0082】

Hbの β -鎖

N-末端ヘモグロビンの β -鎖の5アミノ酸長のペプチドアナログ (陰性の配列コントロール)。

40

【0083】

フルクトシル-Hb の β -鎖

N末端のパリン残基における1-デオキシフルクトシルを含むヘモグロビンの β -鎖の5アミノ酸長のペプチドアナログ (陰性の1-デオキシフルクトシル配列コントロール)。

【0084】

ペプチドのスポットを伴った膜を、DAK Hb1c-1および無関係な抗体 (陰性の抗体コントロール、抗トロポニンI、ダコ (DAKO) 社、製品番号09528) とともにインキュベートした。その後、膜をHRP複合化ヤギの抗マウス抗体 (ダコ社、製品番号P0447) とともにインキュベートして、3-アミノ-9-エチルカルバゾールを基質として用いて、目に見えるよう

50

にした。

【 0 0 8 5 】

【 表 1 】

アミノ酸の数	5	4	3	2	1
ペプチドの名称					
フルクトシル-Hb β-鎖	++	++	++	+	-
Hb β-鎖	-	-	-	-	-
Hb α-鎖	-	N.D.	N.D.	N.D.	N.D.
フルクトシル-Hb α-鎖	-	N.D.	N.D.	N.D.	N.D.

10

【 0 0 8 6 】

表 1 : 抗Hb_{1c}抗体のクローンDAK Hb1c-1のエピトープマッピングのためのスポットペプチドライブラリー。抗Hb_{1c}抗体のクローンDAK Hb1c-1とのインキュベーションと、それに続く視覚化の後の、ペプチドのスポットの強度を示す。高い強度(++)、中程度の強度(+)、陰性(-)、測定せず(N.D.)。

【 0 0 8 7 】

DAK Hb1c-1は、1-デオキシフルクトシル・ヘモグロビン β-鎖の二量体とは穏やかに反応し、そして1-デオキシフルクトシル・ヘモグロビン β-鎖三量体~五量体とは強く反応する(表1を参照すること)。DAK Hb1c-1は、1-デオキシフルクトシル-バリンそのものとは反応しない。DAK Hb1c-1は、ヘモグロビン β-鎖のフルクトシル化されていないペプチド、1-デオキシフルクトシル・ヘモグロビン β-鎖五量体およびヘモグロビン β-鎖五量体とは反応しない。DAK Hb1c-1とは対照的に、陰性の抗体コントロールは、いずれのペプチドとも反応しなかった。

20

【 0 0 8 8 】

要約すれば、ヘモグロビン β-鎖の1-デオキシフルクトシル化N-末端部分についての、抗Hb1c抗体のクローンDAK Hb1c-1の特異性が実証された。

【 0 0 8 9 】

実施例 5 :

全血サンプル、赤血球の溶血物またはキャリブレーター由来のペプチド、タンパク質を、96-穴のマイクロタイタープレートの表面に結合させた。結合しなかったタンパク質を取り除く洗浄工程に続いて、モノクローナル抗体であるDAK Hb1c-1を添加した。インキュベーションの後に、結合しなかった抗体を洗い落とし、そしてヤギの抗マウス抗体と酵素である西洋ワサビペルオキシダーゼとの複合体を添加した。別の洗浄工程の後に、基質である3,3',5,5'-テトラメチルベンジジン(TMB)との反応によって、結合している複合体を検出した。酸を添加してこの反応を停止させ、分光測光的に読み取られる比色定量の終点を得た。

30

【 0 0 9 0 】

実施例 6 :

1-デオキシフルクトシルをヘモグロビン β-鎖のアミノ末端に結合させなかった点を除いて、HbA₀はHbA_{1c}と同じである。

40

【 0 0 9 1 】

DAK Hb1c-1の、HbA₀ならびにHbC₀の17マーのペプチドおよびHbS₀の17マーのペプチドとの反応を、上記のELISAによって分析した。

【 0 0 9 2 】

HbA₀またはHbC₀の17マーのペプチドもしくはHbS₀の17マーのペプチドのいずれかでコーティングしたマイクロタイタープレートにDAK Hb1c-1を添加した場合、測定される値はバックグラウンドの値よりも高くはなかった。

【 0 0 9 3 】

50

結論としては、DAK Hb1c-1は、HbA₀、HbC₀またはHbS₀との反応性を示さなかった。

【0094】

実施例7：

DAK Hb1c-1と天然型HbA_{1c}との溶液中での反応を、競合的ELISA法で分析した。溶液中で、DAK Hb1c-1抗体と一緒にHbA_{1c}キャリアレーターのプレインキュベーションを行い、何らかの抗原-抗体複合体が形成されるかどうかを試験した。次いで、抗原-抗体の溶液をマイクロタイターのウェルに移した。ここには同一の抗原を固相上に吸着させた(固定化抗原)。溶液中の抗原は過剰に存在した。溶液中で抗原と複合体を形成していない抗体は、この固定化抗原と結合することが可能であった。上記の二次抗体によって、結合しているDAK Hb1c-1抗体を検出した。コントロールとして、溶液中でHbA_{1c}キャリアレーターとのプレインキュベーションを行っていない同一量の抗体を、マイクロタイタープレートの1つのレーンに添加し、続いて上記のプロトコールに付した。

10

【0095】

溶液中でHbA_{1c}と一緒にDAK Hb1c-1抗体のプレインキュベーションを行っても、何らの効果も見られなかったことから、DAK Hb1c-1は、溶液中で天然型HbA_{1c}とは結合しないとの結論を出した。

【0096】

実施例8：

DAK Hb1c-1およびHEM13の、ヘモグロビンの構造バリエーションを含むヒトの血液との反応性を、等電点電気泳動(IEF)とそれに続くイムノブロットによって分析した。

20

【0097】

ヘモグロビンの構造バリエーションを分離する目的で、IEFポリアクリルアミドゲル(ファストゲル(PhastGel)IEF-5-8、ファルマシア社)を用いるファルマシア社のファストシステム(Pharmacia's PhastSystem)によって、HbCA、HbSSおよびHbAAを含む赤血球の溶血物を電気泳動的に分離した。次いで、このゲルを押し付けて、PVDF膜上へのタンパク質のブロッティングを行い、そして、ブロッキング剤を用いて非特異的染色をブロッキングした。次いで、DAK Hb1c-1(図1Aを参照すること)およびHEM13(図1Bを参照すること)のそれぞれと一緒に、この膜のインキュベーションを行った。HRP-複合化ヤギの抗マウス免疫グロブリン(ダコA/S社、製品番号P0447)を二次抗体として用いた。最後に、ダコ社の液体DAB+基質-色素原システム(Liquid DAB+ Substrate-Chromogen System)(ダコA/S社、製品番号K3468)を用いて、HRP活性を目に見えるようにした。

30

【0098】

ホモ接合型HbSSを持つ、糖尿病ではない患者からの赤血球の溶血物を、モノクローナル抗体であるクローンDAK Hb1c-1を用いて分析して、強度が高く明確なHbS_{1c}のバンドを同定した(図1Aを参照すること)。対照的に、ホモ接合型HbSSを持つ患者からの赤血球の溶血物をHEM13を用いて分析すると、HbS_{1c}がわずかに染色されることが分かった(図1Bを参照すること)。

【0099】

ヘテロ接合型ヘモグロビンCAを持つ、糖尿病ではない個体からの赤血球の溶血物を、DAK Hb1cおよびHEM13を用いて分析すると、HbC_{1c}のそれぞれ高強度のバンドおよび中程度の強度のバンドが目に見えるように現れた。DAK Hb1cおよびHEM13の両者については、ほぼ同じ強度の強いHbA_{1c}のバンドが観察された。

40

【0100】

コントロールとして、ヘモグロビンAを持つ糖尿病患者からの赤血球の溶血物のプールについても試験を行った。CおよびSとは関係のない、ヘモグロビンの他のバリエーションがこの患者のプールの一部であってもよい。DAK Hb1c-1およびHEM13の両者について、ほぼ同一の強さのHbA_{1c}のバンドが現れるまで、基質と一緒にそれぞれの膜のインキュベーションを行った。

【0101】

最終的に、我々は、我々のpan特異的モノクローナルグリケーションされたヘモグロビン

50

抗体であるクローン DAK Hb1c-1 が、ヒトの血液サンプル由来の HbA_{1c}、HbS_{1c} および HbC₁ と反応することを実証した。

【0102】

実施例 9：

成熟ヘモグロビンは四量体タンパク質であり、二つの α -サブユニットおよび二つの β -サブユニット、 $\alpha_2\beta_2$ からなる。Hb の構造バリエーションは、ペプチドの一次構造中に、遺伝的に決められた変異を有する。A、S および C は、 β -鎖の N-末端から 6 番目のアミノ酸残基だけが異なっている：

【0103】

【化 1】

HbA _{1c}	fructosyl-V-H-L-T-P-E-E-K-T-A-V-N-A-L-W-G-K.....
HbC _{1c}	fructosyl-V-H-L-T-P-K-E-K-T-A-V-N-A-L-W-G-K.....
HbS _{1c}	fructosyl-V-H-L-T-P-V-E-K-T-A-V-N-A-L-W-G-K.....

10

【0104】

抗体が pan 特異的であるか、すなわち抗体がグリケーションされたヘモグロビンの全ての構造バリエーションと反応するかどうかを試験するために、我々は、DAK Hb1c-1 抗体の、グリケーションされたヘモグロビン A、C および S の β -サブユニットの N-末端に対応する三種類の 17 マーのグリケーションされたペプチドとの反応について、ELISA 法によって試験した (図 2)。これらのペプチド配列を上記に示す。コントロールとして、ヘモグロビン A を持つ糖尿病患者からの赤血球の溶血物のプールについても、DAK Hb1c-1 の ELISA 法により試験をした。

20

【0105】

精製したヘモグロビン A_{1c} を用いてマウスを免疫化することによって、モノクローナル Hb A_{1c} 抗体のクローン HEM13 を惹起した。我々は、DAK Hb1c-1 および HbA_{1c} 抗体クローン HEM13 の反応と、グリケーションされたヘモグロビンペプチドおよび赤血球抽出物であるヘモグロビン A とを比較した。

【0106】

DAK Hb1c-1 のグリケーションされたヘモグロビンペプチドに対する親和性は、滴定曲線を比較することによって判断されるように、同じペプチドに対する HEM13 の親和性よりも明らかに強い (図 2 および図 3 を参照すること)。

30

【0107】

DAK Hb1c-1 は三種類のペプチドの全てについて同じように反応する (図 1) のに対して、HEM13 は A_{1c}、S_{1c} および C_{1c} ペプチドについて異なる反応性を示す (図 3)。これらの結果は、DAK Hb1c-1 抗体が pan 特異的であることを実証する。

【0108】

この実施例では、グリケーションされたヘモグロビン A、C および S の構造バリエーションの β -鎖の全ての最初の 4 個のアミノ酸に 1 つの無関係なアミノ酸 (ここではシステイン) を加えたものに対応するところの 5 マーのグリケーションされたペプチドを注射することによって、グリケーションされたヘモグロビンの全ての構造バリエーションについて同じように反応する抗体を惹起することが可能であることを実証する。

40

【0109】

実施例 10：

「ERL 教育プログラム (ERL Educational Programme)」からの血液サンプルを試験した。陽イオン交換クロマトグラフィーによって、それぞれのサンプルにおける HbA_{1c} のパーセンテージを測定した (表 2 中の IFCC/DCCT-値；IFCC：国際臨床化学連合 (International Federation of Clinical Chemistry)；DCCT：糖尿病のコントロールと合併症のトライアル)。ヘテロ接合型のサンプルは二つの異なるヘモグロビン α -サブユニット、たとえば AS または AC から構成されるのに対して、ホモ接合型の血液サンプルは二つの同一の α -サブユニット、たとえば AA または SS から構成される。

50

【 0 1 1 0 】

【 表 2 】

	IFCC	DCCT	DAK-Hb1c-1
HbAA	5.0	6.0	5.4
HbAA	5.7	6.5	6.2
HbAA	6.3	7.0	6.5
HbAA	7.0	7.6	7.9
HbAA	7.5	8.0	8.5
HbAA	8.1	8.4	8.7
HbAA	8.8	8.9	9.4
HbAA	9.4	9.4	9.7
HbAA	10.0	9.8	10.0
HbAS	4.3	5.5	4.7
HbAC	4.3	5.5	4.0

表2：血液サンプル中のHb1cのパーセンテージ。血液サンプルACおよびASは、ヘテロ接合型ヘモグロビンを含む。

10

【 0 1 1 1 】

上記のELISA法の形式にて、DAK Hb1c-1を用いて、全血サンプル中のHbA1cのパーセンテージを4-10%の範囲内で測定した。IFCCおよびDCCTによって推奨されている標準的な方法によって測定された値と同じレベルでのHbA1cのパーセンテージを得た。

20

【 0 1 1 2 】

要約すれば、当該抗体は、HbA1c、HbC1cおよびHbS1cに対して正確な特異性を有することが実証される。

【 0 1 1 3 】

実施例 1 1 :

二つのマイクロスフェアによる比濁法によって、グリケーションされたヘモグロビンレベルを測定した。この準備において、グリケーションされたヘモグロビンを非共有結合的にマイクロスフェアに吸着させ、次いで、マイクロスフェアに共有結合したグリケーションされたヘモグロビン抗体を伴う抗原を検出した。免疫マイクロスフェアと複合化した抗原マイクロスフェアを構成する、得られるネットワークの濁度を測定した。

30

【 0 1 1 4 】

出願人又は代理人 の書類記号	P062 W0 01	国際出願番号	RO/DK 25 APR 2002
-------------------	------------	--------	-------------------

寄託された微生物または他の生物学的材料に関する表示

(PCT Rule 136a)

A. 下記になされる表示は、明細書の以下で言及された寄託された微生物または他の生物学的材料に関する 頁 <u>7</u> , 行 <u>19</u>	
B. 寄託の同定 さらなる寄託が別用紙に同定されている <input type="checkbox"/>	
寄託機関の名称 DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH	
寄託機関のあて名（郵便番号及び国名も記載） Mascheroder Weg 1b D-38124 Braunschweig	
寄託の日付 6 March 2001	アクセッション番号 DSM ACC2495
C. さらなる表示（該当しない場合空白のままにする） この情報は別用紙に記載されている <input type="checkbox"/>	
D. 表示がなされる指定国（表示が全ての指定国に対するものではない場合）	
E. 表示の別の提示（該当しない場合空白のままにする）	
下記の表示は後日国際事務局に提出される（表示の一般的種類例えば「寄託のアクセッション番号」を明記）	

10

20

30

受理官庁記入欄	国際事務局記入欄
<input type="checkbox"/> この用紙は国際出願として受理された	<input type="checkbox"/> この用紙を国際事務局で受理した日 16 MAY 2002 (16.05.02)
権限のある職員	権限のある職員 Jean-Claude MARTIN

Form PCT/RO/134 (July1998)

【 図面の簡単な説明 】

【 0 1 1 5 】

【 図 1 】

【 図 2 】

【 図 3 】

40

【 図 1 】




Figure 1

【 図 2 】




Figure 2 : 日立試薬により測定される、DAK Hb1c-1と、ヘモグロビン A1 (HbA1) を持つ患者からの赤血球の溶血物ならびにヘモグロビン A1c、C1c および S1c に対応する 17- α -糖化ヘマチドとの反応性

【 図 3 】




Figure 3 : 日立試薬により測定される、HEM13と、ヘモグロビン A1 (HbA1) を持つ患者からの赤血球の溶血物ならびにヘモグロビン A1c、C1c および S1c に対応する 17- α -糖化ヘマチドとの反応性

【国際公開パンフレット】

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number
WO 02/088185 A2(51) International Patent Classification: **C07K 16/00**

(21) International Application Number: PCT/DK02/00267

(22) International Filing Date: 25 April 2002 (25.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2001 00656 26 April 2001 (26.04.2001) DK(71) Applicant (for all designated States except US): **DAKO-CYTOMATION DENMARK A/S** [DK/DK]; Produktionsvej 42, DK-2600 Glostrup (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JENSEN, Sussie, Steen** [DK/DK]; Røngevej 20, DK-2600 Glostrup (DK).
PIL, Kurt [DK/DK]; Soegaardsvej 2, DK-2750 Ballerup (DK).
SIMONSEN, Anna, Carina, Wiborg [DK/DK]; Sverigesvej 13A, DK-2800 Kgs. Lyngby (DK).(74) Agent: **HERMANSEN, Erik**; Dako A/S, Produktionsvej 42, DK-2600 Glostrup (DK).

(81) Designated States (national): AF, AG, AI, AM, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CI, CN, CO, CR, CU, CZ (utility model), DE (utility model), DL, DK (utility model), DM, DZ, EC, EE

(utility model), EB, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO utility model (GH), ARIPO patent (GH), ARIPO utility model (GM), ARIPO patent (GM), ARIPO utility model (KE), ARIPO patent (KE), ARIPO utility model (LS), ARIPO patent (LS), ARIPO utility model (MW), ARIPO patent (MW), ARIPO utility model (MZ), ARIPO patent (MZ), ARIPO utility model (SD), ARIPO patent (SD), ARIPO utility model (SL), ARIPO patent (SL), ARIPO utility model (SZ), ARIPO patent (SZ), ARIPO utility model (TZ), ARIPO patent (TZ), ARIPO utility model (UG), ARIPO patent (UG), ARIPO utility model (ZM), ARIPO patent (ZM), ARIPO utility model (ZW), ARIPO patent (ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IL, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BI, CI, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
*without international search report and to be republished upon receipt of that report**For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/088185 A2

(54) Title: PAN-SPECIFIC MONOCLONAL ANTIBODY

(57) Abstract: The invention describes pan-specific monoclonal antibodies for the determination of the amount of all structural variants of glycosylated hemoglobin in human blood samples. The determination of the rate of glycation of hemoglobin in an individual's blood provides a useful index of glucose level control in diabetics.

WO 02/088185

PCT/DK02/00267

Pan-specific monoclonal antibody

This invention relates to the determination of the amount of all structural variants of glycated hemoglobin in human blood samples. The determination of the rate of glycation of hemoglobin in an individual's blood provides a useful index of glucose level control in diabetics.

In particular, the present invention concerns the preparation of monoclonal antibodies or a fragment thereof, which recognize specifically the glycated N-terminal peptide residue in such human hemoglobin.

Patients afflicted with diabetes are incapable of metabolizing glucose in a conventional manner resulting in a build-up of glucose in their blood and urine. Conventionally, the glucose level in such body fluids is taken as a measure of the state of the diabetic condition, which in turn is used as a guide for the amount of insulin or other agent to be taken or of the need to change the patient's diet.

This works moderately well except that the glucose level may fluctuate widely in dependence on time and content of the last meal, the last insulin injection, and the like. Thus, the reading will reflect an instantaneous condition, which may not truly identify the longer-term state of the diabetic condition.

It is known that another effect of the diabetic condition is an increase in the amount of glycated hemoglobin in the blood of the diabetic. Hemoglobin (Hb) is a protein tetramer consisting of four chains (subunits) of amino acids, each of about 143 units and having a total

WO 02/088185

PCT/DK02/00267

2

molecular weight of approximately 64,000 g/mol. The N-terminal valine of the β -subunit in hemoglobin can react with glucose. The glycation of hemoglobin occurs by a non-enzymatic reaction involving glucose and the alpha-amino group of valine. Following a Schiff base formation between the reactants, the glucose undergoes an Amadori rearrangement forming 1-deoxyfructo-valine. The final complex binding between 1-deoxyfructose and valine is covalent and irreversible. The glycation reaction is governed by the concentration of the reactants, e.g., hemoglobin and glucose. The Diabetes Control and Complication Trial (DCCT) reference interval for non-diabetic individuals is approximately 4-6% glycated hemoglobin. Hemoglobin tetramers with a 1-deoxyfructo-valine on the N-terminus of a β -chain are identified as being glycated hemoglobin; e.g. Hb1c or HbA1c, HbC1c or HbS1c.

HbC trait has a prevalence of 30% in parts of sub-Saharan and 2.3% among African Americans. HbS trait, the most commonly variant of Hb in US, has a prevalence of 7.8% of African Americans. All together, at least 10% of black Americans have either HbC or HbS trait. The world population is getting more and more mixed with respect to race. Therefore, it is increasingly important to have an antibody that reacts with all structural variants of glycated hemoglobin in the same manner. If the available antibody do not react with all structural variants of glycated hemoglobin, Hb1c percent measurements from patients with traits of HbS and/or HbC will be incorrect.

Glucose levels in diabetics are sufficiently high to increase the rate of glycation depending directly on the glucose level in the blood, which reflects the severity

WO 02/088185

PCT/DK02/00267

3

of the diabetic condition. Therefore, the hemoglobin A1c level is raised to about 6 to 12%. Since the circulating life span of hemoglobin A is about 120 days, a glycated hemoglobin measurement will give a value, which reflects
5 an average glucose level for that period. Notably a meal high in glucose will not be reflected in instantly high levels of glycated hemoglobin or serum albumin. Thus, measurement of the glycated-hemoglobin content gives a truer picture of the average circulating glucose levels
10 and thus a truer picture of the long-term condition of the patient.

A common method used to determine the level of glycated hemoglobin involves passing a lysed blood sample through
15 a boronate affinity column. The method determines the total glycated hemoglobin, including HbA_{1c} and ketoamine structures formed on lysine and N-terminal valine residues of both the α - and β -chains of hemoglobin. The column is washed and the glycated hemoglobin determined
20 spectrophotometrically.

Microparticle enhanced turbidimetric immunoassay techniques are available for automatic analyses by
instruments. These instruments are not dedicated to a
25 certain diagnostic assay and have a very high sample cycle time. Currently, the utility of this technique for determination of glycated hemoglobin is growing as routine clinical analysis. Some of these immunoassays are based on the principle of microparticle enhanced competitive
30 turbidimetric inhibition immunoassay techniques.

US Patent no. 4,247,533 discloses an analytical technique wherein antibodies to HbA_{1c} were reportedly raised in a sheep by injection of HbA_{1c} and absorbed with nonglycated

WO 02/088185

PCT/DK02/00267

4

hemoglobin to provide polyclonal antibodies, which distinguished between HbA_{1c} and nonglycated hemoglobin. Such antibodies then form the basis for a test to determine the proportion of glycated hemoglobin in a sample.

In the prior art only the hemoglobin A_{1c} antibody and how the antibody is able to react with the A variant of glycated hemoglobin is described. The influence of the other variants of glycated hemoglobin is not mentioned and a monoclonal antibody against all structural variants of glycated-hemoglobin has neither been considered. Due to the fact that the antibody does not recognise HbS_{1c} and HbC_{1c}, immunoassay measurement of glycated-hemoglobin percentages in subjects with HbS_{1c} and HbC_{1c} using that antibody will lead to truly decreased percentages of glycated hemoglobin.

An Hb_{1c} antibody, which is specific for all structural variants of Hb_{1c}, is considered as pan specific.

Accordingly, there remains a need for an antibody of the above-mentioned kind, which does not exhibit the above-identified drawbacks.

It is accordingly an objective of the present invention to provide an antibody, which is pan-specific, i.e. specific for all structural variants of glycated hemoglobin.

It is also the objective to provide an antibody for determining the long-term blood sugar level of a patient based on an immunoassay determination of HbA_{1c}, HbC_{1c} and HbS_{1c}.

WO 02/088185

PCT/DK02/00267

5

It is another objective of the invention to provide such an antibody for determining the content of glycosylated hemoglobin, HbA_{1c}, HbC_{1c} and HbS_{1c}, in a patient's whole
5 blood sample.

These and the other objects are achieved by the invention as defined in the claims.

10 The invention relates to a monoclonal antibody or a fragment thereof, which is pan-specific, i.e. able to react with all structural variants of glycosylated-hemoglobin. The structural variants of glycosylated-hemoglobin are preferably HbA_{1c}, HbC_{1c} and HbS_{1c}.

15 The monoclonal antibody preferably leads to improvement of existing immunoassay with respect to measurements of the level of glycosylated hemoglobin in patients with HbS and HbC.

20 The monoclonal antibody of the present invention is characterized in that the antibody comprises an antibody combining site, which binds specifically to the glycosylated N-terminal peptide sequence of human hemoglobin A, C and
25 S β -chains.

The monoclonal antibody reagent of the present invention is characterized by its specific binding affinity for the linear epitope in glycosylated hemoglobin. Therefore, as used
30 herein the term "antibody reagent" or "antibody combining site" will refer to any material however obtained which comprises a monoclonal antibody combining site specific for such peptide epitope. Such term therefore includes whole antibodies as well as appropriate fragments or

WO 02/088185

PCT/DK02/00267

6

polyfunctionalized forms thereof. When in the form of whole antibody, it can belong to any of the classes and subclasses of known immunoglobulins, e.g. IgG, IgM and so forth. Any fragment of any such immunoglobulins which
5 retains specific binding affinity for the peptide epitope can also be employed, for instance, the fragments of IgG conventionally known as Fab, Fab' and F(ab')₂. In addition, aggregates, polymers, derivatives, conjugates, and hybrids of immunoglobulins, or their fragments can be
10 used where appropriate. Furthermore, single chain antibodies or any other fragment obtained by recombinant means can be used where appropriate.

The glycosylated residue is the distinguishing structural
15 feature of HbA_{1c}, HbS_{1c} and HbC_{1c}. An antibody of the present invention requires an epitope or determinant site comprising minimally the 1-deoxyfructosyl carbohydrate unit, formed upon Amadori rearrangement of the reaction product between glucose and the terminal amine, and a
20 peptide sequence extending therefrom comprising at least one of the amino acid units of the HbA_{1c}, HbS_{1c} and HbC_{1c} N-terminal sequence in the position corresponding to the native HbA_{1c}, HbS_{1c} and HbC_{1c} sequence. The other amino acid units in the peptide sequence characterizing the
25 epitope may be the same or different as those appearing in the native HbA_{1c}, HbS_{1c} and HbC_{1c} sequence. In this way, the epitope is characterized by at least two contact or binding sites with the antibody, which sites are unique to the glycosylated N-terminal HbA_{1c}, HbS_{1c} and HbC_{1c}
30 sequence.

The antibody will preferably specifically bind a glycosylated peptide residue of the formula:

WO 02/088185

PCT/DK02/00267

7

Glycosyl-(NH)-Val-His- or Glycosyl-(NH)-Val-His-AA,

wherein Glycosyl-(NH)-Val represents a non-enzymatically glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin and AA means one or more additional amino acid residues.

In a preferred embodiment AA is a sequence of from 1 to 3 amino acids corresponding to a part of the N-terminal β -chain of all structural variants of human hemoglobin. AA is preferably -Leu-Aa; -Leu-Thr-Aa; -Leu-Thr-Pro-Aa, and Aa designates the remaining number of amino acids in the AA group.

A preferred monoclonal antibody or a fragment thereof according to the invention is obtainable using the hybridoma DAK Hb1c-1, which is deposited at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) as accession no. DSM ACC2495.

The invention also relates to a method of producing a monoclonal antibody or a fragment thereof, wherein the antibody is pan-specific, i.e. able to react with all structural variants of glycosylated hemoglobin.

The antibody or a fragment thereof may preferably be obtainable by the method as defined in claims.

The method of producing a monoclonal antibody or a fragment thereof according to the invention comprising the step of raising an antibody against an immunogen comprising a glycosylated peptide chemically linked to an immunogenic carrier material, the glycosylated peptide having

WO 02/088185

PCT/DK02/00267

8

from 2 to 5 amino acid units corresponding to the N-terminal in the β -chain of human hemoglobin.
The immunogen is preferably of the formula:

5 [Fructosyl-(NH)-Val-His-AA-R]_n-Carrier

wherein Fructosyl-(NH)-Val represents a nonenzymatic glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin, AA means one or more additional amino acid residues, R is a linking group, Carrier is an immunogenic carrier material, and n is an integer from 1 to the number of available coupling sites on said carrier.

15 In a preferred embodiment AA is one or more additional amino acid residues. In another preferred embodiment AA is a sequence of from 1 to 3 amino acids corresponding to the N-terminal part of the β -chain of all structural variants of human hemoglobin.

20 Linking group R can be essentially any convenient and stable structure. Such linking group R will usually be in the form of an aliphatic chain comprising between 1 and approximately 20 atoms, excluding hydrogen, and including heteroatoms such as nitrogen, oxygen, and sulfur. The glycosylated residue can be joined through a variety of groups to form linking chain R, including methylene, ether, thioether, imino, and the like. One skilled in the art will have a wide variety of linking groups from which
25 to choose to prepare the immunogen.
30

In a preferred embodiment the carrier is an immunogenic protein or peptide other than human hemoglobin.

WO 02/088185

PCT/DK02/00267

9

The immunogen used to stimulate production of appropriate immunoglobulins in the most general sense will comprise one or more of the glycosylated peptide residues chemically linked to an immunogenic carrier material. The immunogenic carrier material can be selected from any of those conventionally known having functional groups available for coupling to the glycosylated peptide residue. In most cases, the carrier is a protein or polypeptide, although other materials such as carbohydrates, polysaccharides, lipopolysaccharides, nucleic acids, and the like of sufficient size and immunogenicity can likewise be used. For the most part, immunogenic proteins and polypeptides will have molecular weights between 4,000 g/mol and 10,000,000 g/mol, preferably greater than 15,000 g/mol, and more usually greater than 50,000 g/mol. Generally, proteins taken from one animal species will be immunogenic when introduced into the blood stream of another species. Particularly useful proteins are albumins, globulins, enzymes, hemocyanins, glutelins, proteins having significant nonproteinaceous constituents, and the like.

In a preferred embodiment the glycosylated N-terminal peptide residue of the native HbA_{1c}, HbS_{1c} and HbC_{1c} molecule is made accessible to the monoclonal antibody or a fragment thereof of the present invention by appropriate denaturation or digestion of the protein in the sample to be assayed.

In another preferred embodiment the glycosylated fragment can be produced by chemical or enzymatic digestion of naturally occurring glycosylated hemoglobin, e.g. HbA_{1c}, HbS_{1c} and HbC_{1c}. This fragment can be coupled to a carrier using classical coupling procedures, e.g.,

WO 02/088185

PCT/DK02/00267

10

glutaraldehyde or carbodiimide, and the conjugate used as an immunogen.

- 5 The antibody or a fragment thereof according to the invention may preferably be produced by the following step:

A. Preparation of peptide immunogen

10

To obtain antibodies against the glycation site of N-terminal valine residue on the β -chain of Hb A, S and C a short peptide (pentamer) is preferably used as an immunogen. The antigen may preferably be Hb1c 4-mer plus
15 cysteine, i.e. fructosyl-V-H-L-T-C.

B. Immunization of mice with human Hb1c peptides

20

The general method used to develop hybridomas secreting monoclonal antibodies is well known to a person skilled in the art.

25

An illustration of the techniques utilized in the practice is described in Journal of Immunological Methods, 1980, 39:285.

30

Female F1 hybrids of CF1 x BALB/c mice may preferably be used, but mice of other strains may also be used. The immunization plan and the concentration of peptide-carrier complex may be selected to form a sufficient amount of antigenically stimulated lymphocytes. In a preferred embodiment mice are immunized intraperitoneally with 50 micrograms of purified protein derivatives (PPD, a

WO 02/088185

PCT/DK02/00267

11

partly unknown mixture of denatured proteins from *M. tuberculosis*) in complex with the Hb1c peptide four times at 2-week intervals. 3-5 days after the final immunization, the spleen cells are extracted from the animal for fusion.

Monoclonal antibodies produced by hybridomas from other species may also preferably be used.

10

C. Cell fusion

The spleen is aseptically taken out from the immunized mouse, and a suspension of the spleen cells is prepared. The spleen cells are fused with mouse myeloma cells from a suitable cell line in the presence of a suitable fusion promoter. In a preferred embodiment the preferred ratio of the spleen cells to the myeloma cells is from about 20:1 to about 2:1, and a fusion medium is suitably used in an amount of 0.5 to 1.5 ml per about 10^8 spleen cells.

The mouse myeloma cells used for cell fusion are well known. P3-X63-Ag 8.653 may preferably be used as a mouse myeloma cell.

25

The fusion promoter is preferably polyethylene glycol having an average molecular weight of 1000 to 4000 g/mol. Other fusion promoters known in the art can also be used. In a preferred embodiment polyethylene glycol having an average molecular weight of 1500 may be used.

30

WO 02/088185

PCT/DK02/00267

12

D. Screening of the fused cells

In a separate container (such as a microtiter plate), a mixture composed of the unfused spleen cells, unfused mouse myeloma cells and the fused hybridoma cells is preferably diluted with a selective medium not supporting the unfused mouse myeloma cells and cultured for a period sufficient to kill the unfused mouse myeloma cells (about 1 week). A culture medium having drug resistance (for example, resistance to 8-azaguanine) and not supporting the unfused mouse myeloma cells, such as HAT medium, is used. In the selective medium, the unfused myeloma cells die. Since the unfused spleen cells are untransformed cells, they die away after a certain period of time (1 week). The fused cells, on the other hand, can survive in the selective medium since they have both the tumoral nature of the parent myeloma cells and the nature of the parent spleen cells.

20

E. Determination of glycosylated hemoglobin by ELISA

The determination is preferably based on the principle of specific immunologic recognition and reaction between a monoclonal antibody and the antigenic epitope to which the antibody uniquely and specifically binds. The recognition and binding may be detected, for example, by an ELISA test, wherein blood samples or calibrators are immobilized on a solid phase support, such as the bottom of a plastic well. By immobilizing human hemoglobin the antigenic epitope may be exposed for antibody binding. The antibody, secondary enzyme-labeled antibody and enzyme substrate may interact with the immobilized antigen (human hemoglobin) thereby antibody-antigen

WO 02/088185

PCT/DK02/00267

13

complexes are preferably formed. A number of dilution, incubation and washing steps, then allow separation of bound and free reagents. A colour forming reaction takes place as a result of binding of the antibody to antigen
5 and the consequent reaction of the enzyme upon its substrate interaction. The formation of colour indicates the presence of Hb1c in the blood sample, and the intensity of the colour provides a quantitative measure of the amount of glycated epitope in the sample. The
10 antibody of the present invention may be used to measure glycated hemoglobin in other immunological assay formats known in the art.

15 F. Cloning of the hybridoma cells producing the desired antibody and the production of the antibody

Hybridoma cells capable of producing the desired antibody are cloned by a suitable method such as limiting
20 dilution method, and the desired antibody can be produced by the following method. Hybridoma cells are cultured in a suitable medium for a certain period of time, and the monoclonal antibody produced by the hybridoma cells can be isolated from the supernatant of the culture.

25

The invention also relates to a method of using a monoclonal antibody or a fragment thereof.

30 The antibodies or a fragment thereof according to the invention can preferably be used in conventional manner to react with blood samples containing unknown quantities of glycated hemoglobin and the extent of reaction can be compared with calibrated standards to determine the

WO 02/088185

PCT/DE02/00267

14

extent of glycation. The read-out can be by fluorescence, by immunoassay, or the like, by joining suitably readable groups to the monoclonal antibodies in known manner without loss of their binding power for the glycated epitope in HbA1c, HbS1c and HbC1c.

The monoclonal antibodies of the present invention are specific for binding to all structural variants of glycated N-terminal peptide residue preferable found in HbA, HbS and HbC. The antibodies are able to bind to the epitope in the native HbA1c, HbS1c and HbC1c molecule when the epitope is appropriately exposed. Steric access to the epitope can be obtained in any effective manner. Exposure of the epitope in the intact protein is understood to be accomplished by a physical or chemical denaturation or digestion at least in the region of the epitope. Such denaturation or digestion can be localized to the region of the epitope or can involve a more general, or even substantially complete denaturation of the tertiary, and additionally the secondary structure of the protein, or partial or complete digestion of the protein.

In a preferred embodiment the antibody binds specifically to the glycated N-terminal peptide sequence in the β -chain of human hemoglobin A1c, C1c and S1c after having been exposed sufficiently to provide steric access thereto.

Denaturation can be accomplished in a variety of ways including conventional treatment of the protein by physical means such as heat, sonication, high or low pH and, as is preferable, chemical denaturation by interaction with an agent or chaotrope in solution.

WO 02/088185

PCT/DK02/00267

15

Protein denaturation can be most effectively accomplished if combinations of chemical and/or chemical and physical means are used, e.g. guanidine and heat, guanidine and SDS, or guanidine and dithiothreitol. Particularly strong chaotropes such as guanidine are preferred. Of course, denaturing conditions which result in substantial insolubilization, aggregation, or precipitation of the protein implying an insignificant amount of the exposed epitope is accessible to the solution for antibody binding will be avoided. A sufficient amount of the denatured protein must remain in solution or suspension in order to obtain useful immunobinding. The extent of solubilization necessary will depend upon the circumstances of the intended or desired binding.

15 In a preferred embodiment the glycated peptide sequence is exposed to the antibody binding site by physical or chemical denaturation or digestion.

20 In another preferred embodiment the glycated peptide sequence is exposed to the antibody binding site by denaturation with a chaotropic agent. The chaotropic agents include, without limitation, guanidine, urea, and various detergents such as sodium dodecylsulfate (SDS) and others, without limitation, including deoxycholate and certain bile salts, 3-(3-cholamidopropyl)-dimethyl-ammonio-1-propanesulfonate, organic solvents such as methanol, propanol, acetonitrile and certain salts such as potassium thiocyanate. Non-ionic detergents such as Triton X[®], Tween[®], nonidet NP-40[®], and octyl-glucosides can also function as protein denaturants.

A significant amount of HbA1c, HbS1c and HbC1c in a particular blood sample can be denatured to expose the

WO 02/088185

PCT/DK02/00267

16

glycated epitope for antibody binding by combining the sample, e.g., whole blood or red cell hemolysate, with an aqueous solution of the chaotrope present at sufficient concentration to denature any HbA1c, HbS1c and HbC1c in the resulting aqueous mixture. Where whole blood is the sample, the chaotrope also serves to lyse red blood cells to release hemoglobin and to inactivate proteases.

The denaturation process may be significantly accelerated by heating the mixture for a short period of time. It has been found that at temperatures below 37°C, denaturation by the chaotrope can take from one to several hours, whereas at temperatures above 50°C sufficient denaturation can be attained in a minute or less. In order to prevent significant denaturation of the antibody and other proteinaceous reagents to be subsequently added to the mixture, the sample-chaotrope mixture will normally be diluted as a separate step or by addition of reagent solutions to a level that the chaotrope is substantially ineffective to denature such reagents, yet will preserve the exposure of the epitope by preventing significant renaturation of HbA1c, HbS1c and HbC1c.

In a preferred embodiment the antibody binds specifically to a glycated N-terminal peptide sequence in the β -chain of glycated human hemoglobin, which has been adsorbed to a solid phase, wherein the solid phase preferably is selected from the group consisting of polystyrene, divinylbenzene, butadiene, polycarbonate, polyacrylamide, polyacrylic acid, polyacryl amide, polyethylene, polypropylene, fluorinated polymers, poly amides, or co-block polymers hereof, gold, carbon, cellulose, derivatized cellulose, or glass.

WO 02/088185

PCT/DK02/00267

17

In a preferred embodiment the monoclonal antibody or a fragment thereof may be used for the determination of the level of glycated hemoglobin in human blood. The monoclonal antibody or a fragment thereof may also preferably be used for the determination of the level of glycated HbS and/or HbC in human blood.

In a further preferred embodiment the monoclonal antibody or a fragment thereof may be used for the determination of the level of glycated heterozygous HbA, HbS and/or HbC in human blood.

In a further preferred embodiment there is provided a turbidimetric two-microsphere method for determination of the level of glycated hemoglobin. This method comprises mixing a sample with microspheres and microspheres on which the monoclonal antibody or a fragment thereof is bound; the latter is referred to as immunoparticles, or with microspheres and subsequently with immunomicrospheres. The resulting immunoagglutinates is measured turbidimetrically.

The microspheres are preferably selected from the group consisting of polystyrene, divinylbenzene, butadiene, polycarbonate, polyacrylamide, polyacrylic acid, polyacryl amide, polyethylene, polypropylene, fluorinated polymers, poly amides, or co-block polymers hereof, gold, or silica.

Microspheres to be used in the method of the present example have a particle size of from 0.01 to 5 micrometers, which are used by dispersing in a usually used buffer solution.

WO 02/068185

PCT/DK02/00267

18

The following experimental non-limiting examples are intended to illustrate certain features and embodiments of the invention.

5 **Example 1:**

The 1-deoxyfructosyl-Val-His-AA peptide was synthesized using the solid-phase method of Sheppard and Atherton as described in international patent application no. WO 86/03494. The peptide was synthesized with fluorenylmethoxycarbonyl (Fmoc) amino acids, using PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate) and HOBt (N-hydroxy-benzotriazole) as coupling reagents.

15 Cysteine and histidin were used with trityl (trt), and threonin with tertiary butyl (tBu) side-chain protection. Leucin was used without side-chain protection.

20 The fructose amino acid conjugate, N-(deoxy-D-fructos-1-yl)-L-valine, was synthesized by modifications of known methods. In short, Di-isopropyliden protected glucose was activated as the trifluate and conjugated to the benzyl ester of valine. The resulting 2,3:4,5-Di-O-isopropyliden-N-(deoxy-D-fructos-1-yl)-L-valine-benzyl ester was hydrogenated over Pd/C in ethanol, lyophilized, and finally used in the peptide synthesis without Fmoc protection. The purity of the product was verified by TLC analysis and ^1H or ^{13}C NMR spectra.

30 The peptide was synthesized by automated solid phase synthesis (Crystal instrument from Calbiochem-Novabiochem Ltd) cleaved from the resin with TFA (trifluoroacetic acid) - TBS (triethylsilan) - Glycerol - H_2O (90% : 4% :

WO 02/088185

PCT/DK02/00267

19

3% : 3%), precipitated and washed with methyl-tertiary-butyl-ether, centrifuged and lyophilised.

Purification was performed by HPLC (C₁₈ reversed-phase
5 column, using a step-wise gradient of typically 5-25%
MeCN in 0.1% TFA). The identity of the peptide was
verified by MALDI-TOF mass spectroscopy and the purity
was finally assured by an analytical HPLC analysis to be
better than 90%.

10

Example 2:

Four weeks before first immunization Bacille Calmette
Guérin (BCG) vaccine was injected to prime the immune
15 response in the mice as described in Giba Foundation
Symposium, 1986, 119: 25.

To make the peptide immunogen it was conjugated to a
carrier mixture, Tuberculin PPD. Peptide and carrier
20 (PPD) were conjugated with glutardialdehyde, i.e. CHO-
(CH₂)₃-CHO. Glutardialdehyde reacts with amino-groups on
the peptide and the carrier.

The immunogens thus have the following formula:

25

Fructosyl-Val-His-Leu-Thr-Cys-CH-(CH₂)₃-CH-PPD

Example 3:

30

Female F1 hybrids of CF1 x BALB/c mice (12 weeks old)
primed with BCG-vaccine were immunized with the peptide-
PPD complex described above. Mice were immunized four
times intraperitoneally with a mixture of 50 micrograms

WO 02/088185

PCT/DE02/00267

20

of peptide-PPD complex dissolved in phosphate buffered saline (PBS) and aluminium hydroxide (1 mg/dose). A mouse with high titer of antibodies against glycosylated hemoglobin was selected for fusion. Three days after the final immunization, spleen cells from the immunized mouse was used for cell fusion.

Spleen cells from the selected immunized mouse and myeloma cells of BALB/c origin (P3-X63-Ag 8.653) were mixed in a ratio of about 3:1 and fused in the presence of 50% polyethylene glycol 1500 (a product of Roche Diagnostics GmbH) (J. W. Goding: Monoclonal Antibodies: Principle and Practice, Academic Press, San Diego, 1996). The fused cells were suspended in RPMI-1640 medium, containing 10% FCS, hypoxanthine, aminopterin and thymidine (HAT media) to give a concentration of 1×10^6 cells/mL. The suspension was distributed to 96-well microplates (Costar) in an amount of 200 μ L per well.

The cells were incubated in a CO₂ incubator (5% CO₂, 37°C). After 14 days hybridoma cells were screened.

Supernatants containing antibodies produced by the hybridoma cell cultures were detected by an ELISA technique using microtiter plates coated with hemoglobin calibrator containing 4.7% HbA_{1c} and 16.7% HbA_{1c} as antigens. 120 μ g/mL of the calibrators were immobilized onto plastic microtiter wells using a carbonate/bicarbonate buffer (pH 9.6) for 15 minutes at 37°C. After a washing step that removes unbound antigen the hybridoma culture supernatants diluted in 50 mM Tris-HCl buffer containing 0.1% Tween 20 were added to each well and allowed to react for one hour at room temperature. After another washing step to remove unbound

WO 02/068185

PCT/DK02/00267

21

antibody a conjugate of a goat anti-mouse antibody and the enzyme horseradish peroxidase was added. After incubation and a third washing step the bound conjugate was detected by reaction with the substrate 3,3',5,5'-tetramethylbenzidine (TMB). The reaction was stopped with sulphuric acid and the absorbance at 450 nm was measured. Only cells from positive wells reacting with glycosylated hemoglobin calibrators, 4.7% HbA1c and 16.7% HbA1c, with increasing intensity were selected.

10 These positive cells were cloned three times by limiting dilution method. The resulting clone, DAK Hb1c-1, was suspended in 90% FCS; 10% DMSO and stored at -150°C.

15 **Example 4:**

A linear antibody epitope mapping is the systematic screening of all possible peptides derived from a polypeptide or protein sequence. Epitope mapping yields information on the linear stretch of amino acids and derivatives that form an interaction site to a given antibody.

25 The specificity of DAK Hb1c-1 for the glycosylated epitope of hemoglobin β -chain has been determined.

A peptide library containing 1-5 amino acid long peptides manually synthesized from the C- to the N-terminal end on a solid phase (Abimed membrane) using Fmoc as protecting group of the N-terminal end of amino acids. Detailed description of peptide synthesis is described in Example 1. The following four peptides were synthesized:

Fructosyl-Hb β -chain

WO 02/088185

PCT/DK02/00267

22

1-5 amino acid long peptide analogs of N-terminal 1-deoxyfructosyl-hemoglobin β -chain.

Hb β -chain

- 5 1-5 amino acid long peptide analog of N-terminal hemoglobin β -chain (minus 1-deoxyfructosyl β -chain; backbone control).

Hb α -chain

- 10 5 amino acid long peptide analog of N-terminal hemoglobin α -chain (negative sequence control).

Fructosyl-Hb α -chain

- 15 5 amino acid long peptide analogs of hemoglobin α -chain incl. 1-deoxyfructosyl at the N-terminal valine residue (negative 1-deoxyfructosyl sequence control).

The membranes with peptide spots were incubated with DAK Hbc-1 and an irrelevant antibody (a negative antibody control, anti-troponin I, DAKO, product no. O 9528). Thereafter, the membranes were incubated with HRP conjugated goat-anti-mouse antibodies (DAKO, product no. P 0447) and visualized using 3-amino-9-ethylcarbazole as substrate.

25

<i>Number of Amino Acids</i>	5	4	3	2	1
<i>Peptide name</i>					
Fructosyl-Hb β -chain	++	++	++	+	-
Hb β -chain	-	-	-	-	-
Hb α -chain	-	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>
Fructosyl-Hb α -chain	-	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>	<i>N.D.</i>

WO 02/088185

PCT/DK02/00267

23

Table 1: Spot peptide library for epitope mapping of anti-Hb1c antibody clone DAK Hb1c-1. Intensity of peptide spots after incubation with anti-Hb1c antibody clone DAK Hb1c-1 followed by visualization are shown; strong intensity (++) , medium intensity (+) , negative (-) , not determined (N.D.).

10 DAK Hb1c-1 reacts moderately with 1-deoxyfructosyl 2-mer of hemoglobin β -chain, strongly with 1-deoxyfructosyl 3-mer to 5-mer hemoglobin β -chain (see Table 1). DAK Hb1c-1 does not react with 1-deoxyfructosyl-valine alone. DAK Hb1c-1 does not react with non-fructosylated peptides of hemoglobin β -chain, 1-deoxyfructosyl 5-mer hemoglobin α -chain and 5-mer hemoglobin α -chain. In contrast to DAK Hb1c-1, the negative antibody control did not react with any of the peptides.

20 In summary, the specificity of anti-Hb1c antibody clone DAK Hb1c-1 for the 1-deoxyfructosylated N-terminal part of hemoglobin β -chain has been demonstrated.

Example 5:

25 Peptides, proteins from a whole blood sample, red cellhemolysate or calibrators were allowed to bind to the surface of a 96-well microtiter plate. Following a washing step to remove unbound proteins, the monoclonal antibody, DAK Hb1c-1, was added. After incubation, the unbound antibody was washed away, and a conjugate of a 30 goat-anti-mouse antibody and the enzyme horseradish peroxidase was added. After another washing step the bound conjugate was detected by reaction with 3,3',5,5'-tetramethylbenzidine (TMB) substrate. The reaction was

WO 02/088185

PCT/DK02/00267

24

stopped by adding acid to give a colorimetric end point that was read spectrophotometrically.

Example 6:

5 HbA₀ is identical to HbA_{1c} except that there is no 1-deoxyfructosyl linked to the amino-terminus of the hemoglobin β -chain.

10 The reaction of DAK Hb1c-1 with HbA₀ and 17-mer peptides of HbC₀ and HbS₀ was analysed by the ELISA described above.

15 When DAK Hb1c-1 was added to microtiter plates coated with either HbA₀ or 17-mers peptides of HbC₀ or HbS₀ the values measured were no higher than the background values.

20 In conclusion, no reactivity of DAK Hb1c-1 with HbA₀, HbC₀ or HbS₀ was demonstrated.

Example 7:

25 The reaction of DAK Hb1c-1 with native HbA_{1c} in solution was analyzed in a competitive ELISA. HbA_{1c} calibrator in solution was pre-incubated with DAK Hb1c-1 antibody to test if any antigen-antibody complexes were formed. The solution of antigen-antibody was then transferred to microtiter-wells, where the same antigen was adsorbed on
30 the solid phase (immobilized antigen). The antigen in solution was in excess. The antibody not in complex with the antigen in solution was able to bind to the immobilized antigen. A secondary antibody as described above detected the bound DAK Hb1c-1 antibody. As a

WO 02/088185

PCT/DK02/00267

25

control the same amount of antibody not pre-incubated with HbA_{1c} calibrator in solution was added to one lane of the microtiter plate and followingly subjected to the protocol described above.

5

No effect of pre-incubation of DAK Hb1c-1 antibody with HbA_{1c} in solution was observed, concluding that DAK Hb1c-1 does not bind to native HbA_{1c} in solution.

10 Example 8:

The reactivity of DAK Hb1c-1 and HEM13 with human blood containing structural variants of hemoglobin have been analysed by isoelectric focusing (IEF) followed by immunoblotting.

15

Red cell hemolysate containing HbCA, HbSS and HbAA were electrophoretically separated by Pharmacia's PhastSystem using IEF polyacrylamide gel (PhastGel IEF - 5-8, Pharmacia) in order to separate structural variants of hemoglobin. Following, the gels were subjected to press blotting of proteins onto a PVDF membrane and blocked for non-specific staining with a blocking agent. The membranes were then incubated with DAK Hb1c-1 (see figure 1 A) and HEM13 (see figure 1 B), respectively. HRP-conjugated goat anti-mouse immunoglobulin (DAKO A/S, product no. P 0447) was used as secondary antibody. Finally, HRP activity were visualized using DAKO liquid DAB+ Substrate-Chromogen System (DAKO A/S, product no. K 3468).

25
30

A distinct HbS_{1c} band with strong intensity was identified analysing red cell hemolysate from a non-diabetic patient with homozygous HbSS with the monoclonal

WO 02/088185

PCT/DK02/00267

26

antibody, clone DAK Hb1c-1 (see Figure 1, A). In contrast, weak HbS_{1c} staining was revealed with HEM13 analyzing red cell hemolysate from a patient with homozygous HbSS (see Figure 1 B).

5

Analyzing red cell hemolysate from a non-diabetic individual with heterozygous hemoglobin CA, HbC_{1c} bands with strong and moderate intensity were visualized with DAK Hb1c and HEM 13, respectively. A strong HbA_{1c} band with approximately the same intensity was observed for both DAK Hb1c and HEM13.

As a control a pool of red cell hemolysate from diabetes patients with hemoglobin A was also tested. Other variants of hemoglobin not related to C and S may be a part of this patient pool. Each membrane was incubated with substrate until HbA_{1c} bands with approximately the same intensity were revealed on both DAK Hb1c-1 and HEM 13.

20

Conclusively, we have demonstrated that our pan-specific monoclonal glycosylated hemoglobin antibody, clone DAK Hb1c-1, reacts with HbA_{1c}, HbS_{1c} and HbC_{1c} from human blood samples.

25

Example 9:

Adult hemoglobin is a tetrameric protein, consisting of two α -subunits and two β -subunits, $\alpha_2\beta_2$. Structural variants of Hb contain genetically determined changes in the primary structure of the peptides. A, S and C differ only in the sixth amino acid residue from the N-terminal of the β -chain:

30

WO 02/088185

PCT/DK02/00267

27

HbA_{1c} fructosyl-V-H-L-T-P-E-E-K-T-A-V-N-A-L-W-G-K.....
HbC_{1c} fructosyl-V-H-L-T-P-K-E-K-T-A-V-N-A-L-W-G-K.....
HbS_{1c} fructosyl-V-H-L-T-P-V-E-K-T-A-V-N-A-L-W-G-K.....

5 To test if the antibody is pan-specific, i.e. reacting
with all structural variants of glycosylated hemoglobin, we
have tested the reaction of DAK Hb1c-1 antibody with
three 17-mer glycosylated peptides corresponding to the N-
terminal end of glycosylated hemoglobin A, C and S β -subunits
10 by ELISA (Figure 2). The peptide sequences are shown
above. As a control a pool of red cell hemolysate from
diabetes patients with hemoglobin A was also tested by
DAK Hb1c-1 ELISA.

15 Monoclonal HbA_{1c} antibody clone HEM13 has been raised by
immunization of mice with purified hemoglobin A_{1c}. We
have compared the reaction of DAK Hb1c-1 and HbA_{1c}
antibody clone HEM13 with the glycosylated hemoglobin
peptides and erythrocyte extract of hemoglobin A.

20 The affinity of DAK Hb1c-1 for the glycosylated hemoglobin
peptides is significantly stronger than the affinity of
HEM13 for the same peptides as judged by comparing the
titration curves (see Figure 2 and Figure 3).

25 HEM13 shows different reactions with the A_{1c}, S_{1c} and C_{1c}
peptides (Figure 3) whereas DAK Hb1c-1 reacts in the same
manner with all three peptides (Figure 1). These results
demonstrates that the DAK Hb1c-1 antibody is pan-
30 specific.

This example verifies that an antibody that reacts in the
same manner with all structural variants of glycosylated
hemoglobin can be raised by injection of a glycosylated 5-mer

WO 02/088185

PCT/DK02/00267

28

peptide corresponding to the first 4 amino acids of all structural variants of glycated hemoglobin A, C and S β -chain plus one irrelevant amino acid (here cysteine).

5 **Example 10:**

Blood samples from the "ERL Educational Programme" have been tested. The HbA_{1c}-percentage of each sample has been determined by cation-exchange chromatography (IFCC/DCCT-values in table 2; IFCC: International Federation of
10 Clinical Chemistry; DCCT: Diabetes Control and Complication Trial). A homozygotic blood sample consists of two identical β -subunits, e.g. AA or SS, whereas heterozygotic samples consists of two different
15 hemoglobin β -subunits, e.g. AS or AC.

	IFCC	DCCT	DAK-HbA _{1c} -1
HbAA	5.0	6.0	5.4
HbAA	5.7	6.5	6.2
HbAA	6.3	7.0	6.5
HbAA	7.0	7.6	7.9
HbAA	7.5	8.0	8.5
HbAA	8.1	8.4	8.7
HbAA	8.8	8.9	9.4
HbAA	9.4	9.4	9.7
HbAA	10.0	9.8	10.0
HbAS	4.3	5.5	4.7
HbAC	4.3	5.5	4.0

Table 2: Percentage of Hb_{1c} in blood samples. The blood sample AC and AS contain heterozygotic hemoglobin.

20 At a range from 4-10% the HbA_{1c} percentage was measured in whole blood samples using DAK Hb_{1c}-1 in the above-mentioned ELISA format. HbA_{1c} percentages at the same

WO 02/088185

PCT/DK02/00267

29

levels as the values determined by reference methods recommended by IFCC and DCCT were obtained.

In summary, it is demonstrated that the antibody in question has the correct specificity against HbA1c, HbC1c and HbS1c.

Example 11:

- 10 The level of glycated hemoglobin was measured in a two-microsphere turbidimetry assay. In this set-up glycated hemoglobin was non-covalently adsorbed to microspheres followed by detection of the antigen with glycated hemoglobin antibodies covalently bound to microspheres.
- 15 The resulting network constituting antigen microspheres complexed with immunomicrospheres were measured turbidimetrically.

WO 02/088185

30

PCT/DK02/00267

Applicant's or agent's file reference P062 W0 01	International application No. RO/DK 2 5 APR 2002
---	--

**INDICATIONS RELATING TO DEPOSITED MICROORGANISMS
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>7</u> , line <u>19</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH	
Address of depository institution (including postal code and country) Mascheroder Weg 1b D-38124 Braunschweig	
Date of deposit 6 March 2001	Accession Number DSM ACC2495
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: 16 MAY 2002 (16.05.02) Authorized officer <i>Jean-Luc MARTIN</i>
--	--

Claims:

1. A monoclonal antibody or a fragment thereof, wherein said antibody is pan-specific, i.e. able to react with
5 all structural variants of glycosyl-hemoglobin, said antibody or a fragment thereof comprises an antibody combining site, which binds specifically to a glycosylated peptide residue of the formula:

10 Glycosyl-(NH)-Val-His- or Glycosyl-(NH)-Val-His-AA,

wherein Glycosyl-(NH)-Val represents a non-enzymatically glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin and AA means 1
15 to 3 amino acid residues.

2. A monoclonal antibody or a fragment thereof according to claim 1, wherein AA is a sequence of from 1 to 3 amino acids corresponding to a part of the N-terminal β -chain
20 of all structural variants of human hemoglobin.

3. A monoclonal antibody or a fragment thereof according to claim 2, wherein AA is -Leu-Aa; -Leu-Thr-Aa; -Leu-Thr-Pro-Aa, and Aa designates the remaining number of amino
25 acids in the AA group.

4. A monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody is obtainable by hybridoma DAK Hblc-1 deposited at
30 Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) as accession no. DSM ACC2495.

5. A monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody

WO 02/088185

32

PCT/DK02/00267

is pan-specific, i.e. able to react with all structural variants of human hemoglobin including HbA1c, HbC1c and HbS1c.

5 6. A monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody being obtainable by a method as defined in any one of the claims 7-11.

10 7. A method of producing a monoclonal antibody or a fragment thereof, wherein said antibody has been raised against an immunogen comprising an antigen obtainable by a glycosylated peptide chemically linked to an immunogenic carrier material, the glycosylated peptide having from 2 to 5
15 amino acid units corresponding to the N-terminal β -chain of human hemoglobin.

8. A method of producing a monoclonal antibody or a fragment thereof according to claim 7, wherein the
20 immunogen is of the formula:

[Fructosyl-(NH)-Val-His-AA-R] $_n$ -Carrier

wherein Fructosyl-(NH)-Val represents a non-enzymatic
25 glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin, AA means one or more additional amino acid residues, R is a linking group, Carrier is an immunogenic carrier material, and n is an integer from 1 to the number of available coupling
30 sites on said carrier.

9. A method of producing a monoclonal antibody or a fragment thereof according to claim 8, wherein AA is a sequence of from 1 to 3 amino acids corresponding to a

part of the N-terminal β -chain of all structural variants of human hemoglobin.

10. A method of producing a monoclonal antibody or a fragment thereof according to claim 8, wherein said Carrier is immunogenic proteins or peptides other than human hemoglobin.

11. A method of using a monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody binds specifically to said glycosylated N-terminal peptide sequence of human hemoglobin A1c, C1c and S1c β -chains after having been exposed sufficiently to provide steric access thereto.

12. A method of using a monoclonal antibody or a fragment thereof according to claim 11, wherein said glycosylated peptide sequence is exposed to the antibody combining site by physical or chemical denaturation or digestion.

13. A method of using a monoclonal antibody or a fragment thereof according to claims 11-12, wherein said glycosylated peptide sequence is exposed to the antibody combining site by denaturation with a chaotropic agent.

14. A method of using a monoclonal antibody or a fragment thereof according to claim 13, wherein the chaotropic agent is selected from the group consisting of guanidine, urea, potassium-thiocyanate, or detergent.

15. A method of using a monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody binds specifically to a glycosylated N-terminal peptide sequence of the β -chain of all

structural variants of human hemoglobin, which has been adsorbed to a solid phase.

16. A method of using a monoclonal antibody or a fragment thereof according to claim 15, wherein the solid phase preferably is constituted by a material selected from the group consisting of microtiter plates, micro particles, membranes, tubes, pins, chips or discs.
- 10 17. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycosylated hemoglobin in human blood.
- 15 18. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycosylated HbS and/or HbC in human blood.
- 20 19. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycosylated heterozygous HbA, HbS and/or HbC in human blood.
- 25 20. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycosylated hemoglobin measured in a two-particle turbidimetric assay.

WO 02/088185

PCT/DK02/00267

1/3




Figure 1




Figure 2: Reactivity of DAK Hb1c-1 with red cell hemolysate from patients with hemoglobin AA (HbAA) and 17-mer glycosylated peptides corresponding to hemoglobin A1c, C1c and S1c as determined by an ELISA test.

WO 02/088185

PCT/DK02/00267

3/3




Figure 3. Reactivity of HEM13 with red cell hemolysate from patients with hemoglobin AA (HbAA) and 17-mer glycosylated peptides corresponding to hemoglobin A1c, C1c and S1c as determined by an ELISA test.

【国際公開パンフレット(コレクション)】

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number
WO 2002/088185 A3(51) International Patent Classification: C07K 16/28,
G01N 33/577, 33/68, 33/72(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD,
SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.(21) International Application Number:
PCT/DK2002/000267

(22) International Filing Date: 25 April 2002 (25.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2001 00656 26 April 2001 (26.04.2001) DK(71) Applicant (for all designated States except US): DAKO-
CYTOMATION DENMARK A/S [DK/DK]; Produk-
tionsvej 42, DK-2600 Glostrup (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JENSEN, Susste,
Steen [DK/DK]; Raagevej 20, DK-2600 Glostrup (DK).
PII, Kurt [DK/DK]; Soegaardsvej 2, DK-2750 Ballerup
(DK). SIMONSEN, Anna, Carlna, Wiborg [DK/DK];
Sverigesvej 13A, DK-2800 Kgs. Lyngby (DK).(84) Designated States (regional): ARIPO utility model
(GH), ARIPO patent (GH), ARIPO utility model (GM),
ARIPO patent (GM), ARIPO utility model (KE), ARIPO
patent (KE), ARIPO utility model (LS), ARIPO patent
(LS), ARIPO utility model (MW), ARIPO patent (MW),
ARIPO utility model (MZ), ARIPO patent (MZ), ARIPO
utility model (SD), ARIPO patent (SD), ARIPO utility
model (SL), ARIPO patent (SL), ARIPO utility model
(SZ), ARIPO patent (SZ), ARIPO utility model (TZ),
ARIPO patent (TZ), ARIPO utility model (UG), ARIPO
patent (UG), ARIPO utility model (ZM), ARIPO patent
(ZM), ARIPO utility model (ZW), ARIPO patent (ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

(74) Agent: HERMANSEN, Erik; Dako A/S, Produktionsvej
42, DK-2600 Glostrup (DK).(88) Date of publication of the international search report:
19 February 2004(81) Designated States (national): AE, AG, AL, AM, AT (uti-
lity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (uti-
lity model), DE, DK (utility model), DK, DM, DZ, EC, EEFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 2002/088185 A3

(54) Title: PAN-SPECIFIC MONOCLONAL ANTIBODY

(57) Abstract: The invention describes pan-specific monoclonal antibodies for the determination of the amount of all structural variants of glycosylated hemoglobin in human blood samples. The determination of the rate of glycosylation of hemoglobin in an individual's blood provides a useful index of glucose level control in diabetics.

【国際公開パンフレット(コレクトバージョン)】

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number
WO 02/088185 A2

(43) International Publication Date
7 November 2002 (07.11.2002)

PCT

(51) International Patent Classification: C07K 16/00

(utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(21) International Application Number: PCT/DK02/00267

(22) International Filing Date: 25 April 2002 (25.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
FA 2001 00656 26 April 2001 (26.04.2001) DK

(71) Applicant (for all designated States except US): DAKO-CYTOMATION DENMARK A/S [DK/DK]; Produktionsvej 42, DK-2600 Glostrup (DK).

(72) Inventors; and
(75) Inventors/Applicants (for US only): JENSEN, Susste, Steen [DK/DK]; Raagevej 20, DK-2600 Glostrup (DK).
PH, Kurt [DK/DK]; Soegaardsvej 2, DK-2750 Ballerup (DK).
SIMONSEN, Anna, Carina, Wilborg [DK/DK]; Sverigesvej 13A, DK-2800 Kgs. Lyngby (DK).

(84) Designated States (regional): ARIPO utility model (GH), ARIPO patent (GH), ARIPO utility model (GM), ARIPO patent (GM), ARIPO utility model (KE), ARIPO patent (KE), ARIPO utility model (LS), ARIPO patent (LS), ARIPO utility model (MW), ARIPO patent (MW), ARIPO utility model (MZ), ARIPO patent (MZ), ARIPO utility model (SD), ARIPO patent (SD), ARIPO utility model (SL), ARIPO patent (SL), ARIPO utility model (SZ), ARIPO patent (SZ), ARIPO utility model (TZ), ARIPO patent (TZ), ARIPO utility model (UG), ARIPO patent (UG), ARIPO utility model (ZM), ARIPO patent (ZM), ARIPO utility model (ZW), ARIPO patent (ZW).
Eurasian patent (AM, AZ, BY, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(74) Agent: HERMANSEN, Erik; Dako A/S, Produktionsvej 42, DK-2600 Glostrup (DK).

Published: without international search report and to be republished upon receipt of that report

(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

60450120020



WO 02/088185 A2



(54) Title: PAN-SPECIFIC MONOCLONAL ANTIBODY

(57) Abstract: The invention describes pan-specific monoclonal antibodies for the determination of the amount of all structural variants of glycosylated hemoglobin in human blood samples. The determination of the rate of glycation of hemoglobin in an individual's blood provides a useful index of glucose level control in diabetics.

WO 02/088185

PCT/DK02/00267

2

Pan-specific monoclonal antibody

This invention relates to the determination of the amount of all structural variants of glycated hemoglobin in human blood samples. The determination of the rate of glycation of hemoglobin in an individual's blood provides a useful index of glucose level control in diabetics.

In particular, the present invention concerns the preparation of monoclonal antibodies or a fragment thereof, which recognize specifically the glycated N-terminal peptide residue in such human hemoglobin.

Patients afflicted with diabetes are incapable of metabolizing glucose in a conventional manner resulting in a build-up of glucose in their blood and urine. Conventionally, the glucose level in such body fluids is taken as a measure of the state of the diabetic condition, which in turn is used as a guide for the amount of insulin or other agent to be taken or of the need to change the patient's diet.

This works moderately well except that the glucose level may fluctuate widely in dependence on time and content of the last meal, the last insulin injection, and the like. Thus, the reading will reflect an instantaneous condition, which may not truly identify the longer-term state of the diabetic condition.

It is known that another effect of the diabetic condition is an increase in the amount of glycated hemoglobin in the blood of the diabetic. Hemoglobin (Hb) is a protein tetramer consisting of four chains (subunits) of amino acids, each of about 143 units and having a total

WO 02/088185

PCT/DK02/00267

2

molecular weight of approximately 64,000 g/mol. The N-terminal valine of the β -subunit in hemoglobin can react with glucose. The glycation of hemoglobin occurs by a non-enzymatic reaction involving glucose and the alpha-amino group of valine. Following a Schiff base formation between the reactants, the glucose undergoes an Amadori rearrangement forming 1-deoxyfructo-valine. The final complex binding between 1-deoxyfructose and valine is covalent and irreversible. The glycation reaction is governed by the concentration of the reactants, e.g., hemoglobin and glucose. The Diabetes Control and Complication Trial (DCCT) reference interval for non-diabetic individuals is approximately 4-6% glycated hemoglobin. Hemoglobin tetramers with a 1-deoxyfructo-valine on the N-terminus of a β -chain are identified as being glycated hemoglobin; e.g. Hb1c or HbA1c, HbC1c or HbS1c.

HbC trait has a prevalence of 30% in parts of sub-Saharan and 2.3% among African Americans. HbS trait, the most commonly variant of Hb in US, has a prevalence of 7.8% of African Americans. All together, at least 10% of black Americans have either HbC or HbS trait. The world population is getting more and more mixed with respect to race. Therefore, it is increasingly important to have an antibody that reacts with all structural variants of glycated hemoglobin in the same manner. If the available antibody do not react with all structural variants of glycated hemoglobin, Hb1c percent measurements from patients with traits of HbS and/or HbC will be incorrect.

Glucose levels in diabetics are sufficiently high to increase the rate of glycation depending directly on the glucose level in the blood, which reflects the severity

WO 02/088185

PCT/DK02/00267

4

3

of the diabetic condition. Therefore, the hemoglobin A1c level is raised to about 6 to 12%. Since the circulating life span of hemoglobin A is about 120 days, a glycated hemoglobin measurement will give a value, which reflects an average glucose level for that period. Notably a meal high in glucose will not be reflected in instantly high levels of glycated hemoglobin or serum albumin. Thus, measurement of the glycated-hemoglobin content gives a truer picture of the average circulating glucose levels and thus a truer picture of the long-term condition of the patient.

A common method used to determine the level of glycated hemoglobin involves passing a lysed blood sample through a boronate affinity column. The method determines the total glycated hemoglobin, including HbA_{1c} and ketoamine structures formed on lysine and N-terminal valine residues of both the α - and β -chains of hemoglobin. The column is washed and the glycated hemoglobin determined spectrophotometrically.

Microparticle enhanced turbidimetric immunoassay techniques are available for automatic analyses by instruments. These instruments are not dedicated to a certain diagnostic assay and have a very high sample cycle time. Currently, the utility of this technique for determination of glycated hemoglobin is growing as routine clinical analysis. Some of these immunoassays are based on the principle of microparticle enhanced competitive turbidimetric inhibition immunoassay techniques.

US Patent no. 4,247,533 discloses an analytical technique wherein antibodies to HbA_{1c} were reportedly raised in a sheep by injection of HbA_{1c} and absorbed with nonglycated

WO 02/088185

PCT/DK02/00267

4

hemoglobin to provide polyclonal antibodies, which distinguished between HbA1c and nonglycated hemoglobin. Such antibodies then form the basis for a test to determine the proportion of glycated hemoglobin in a sample.

In the prior art only the hemoglobin A_{1c} antibody and how the antibody is able to react with the A variant of glycated hemoglobin is described. The influence of the other variants of glycated hemoglobin is not mentioned and a monoclonal antibody against all structural variants of glycated-hemoglobin has neither been considered. Due to the fact that the antibody does not recognise HbS1c and HbC1c, immunoassay measurement of glycated-hemoglobin percentages in subjects with HbS1c and HbC1c using that antibody will lead to truly decreased percentages of glycated hemoglobin.

An Hb1c antibody, which is specific for all structural variants of Hb1c, is considered as pan specific.

Accordingly, there remains a need for an antibody of the above-mentioned kind, which does not exhibit the above-identified drawbacks.

It is accordingly an objective of the present invention to provide an antibody, which is pan-specific, i.e. specific for all structural variants of glycated hemoglobin.

It is also the objective to provide an antibody for determining the long-term blood sugar level of a patient based on an immunoassay determination of HbA1c, HbC1c and HbS1c.

WO 02/088185

PCT/DK02/00267

6

5

It is another objective of the invention to provide such an antibody for determining the content of glycosylated hemoglobin, HbA1c, HbC1c and HbS1c, in a patient's whole blood sample.

These and the other objects are achieved by the invention as defined in the claims.

10 The invention relates to a monoclonal antibody or a fragment thereof, which is pan-specific, i.e. able to react with all structural variants of glycosylated-hemoglobin. The structural variants of glycosylated-hemoglobin are preferably HbA1c, HbC1c and HbS1c.

15 The monoclonal antibody preferably leads to improvement of existing immunoassay with respect to measurements of the level of glycosylated hemoglobin in patients with HbS and HbC.

20 The monoclonal antibody of the present invention is characterized in that the antibody comprises an antibody combining site, which binds specifically to the glycosylated N-terminal peptide sequence of human hemoglobin A, C and S β -chains.

25 The monoclonal antibody reagent of the present invention is characterized by its specific binding affinity for the linear epitope in glycosylated hemoglobin. Therefore, as used herein the term "antibody reagent" or "antibody combining site" will refer to any material however obtained which comprises a monoclonal antibody combining site specific for such peptide epitope. Such term therefore includes whole antibodies as well as appropriate fragments or

WO 02/088185

PCT/DK02/00267

7

6

polyfunctionalized forms thereof. When in the form of whole antibody, it can belong to any of the classes and subclasses of known immunoglobulins, e.g. IgG, IgM and so forth. Any fragment of any such immunoglobulins which retains specific binding affinity for the peptide epitope can also be employed, for instance, the fragments of IgG conventionally known as Fab, Fab' and F(ab')₂. In addition, aggregates, polymers, derivates, conjugates, and hybrids of immunoglobulins, or their fragments can be used where appropriate. Furthermore, single chain antibodies or any other fragment obtained by recombinant means can be used where appropriate.

The glycosylated residue is the distinguishing structural feature of HbA_{1c}, HbS_{1c} and HbC_{1c}. An antibody of the present invention requires an epitope or determinant site comprising minimally the 1-deoxyfructosyl carbohydrate unit, formed upon Amadori rearrangement of the reaction product between glucose and the terminal amine, and a peptide sequence extending therefrom comprising at least one of the amino acid units of the HbA_{1c}, HbS_{1c} and HbC_{1c} N-terminal sequence in the position corresponding to the native HbA_{1c}, HbS_{1c} and HbC_{1c} sequence. The other amino acid units in the peptide sequence characterizing the epitope may be the same or different as those appearing in the native HbA_{1c}, HbS_{1c} and HbC_{1c} sequence. In this way, the epitope is characterized by at least two contact or binding sites with the antibody, which sites are unique to the glycosylated N-terminal HbA_{1c}, HbS_{1c} and HbC_{1c} sequence.

The antibody will preferably specifically bind a glycosylated peptide residue of the formula:

WO 02/088185

PCT/DK02/00267

7

Glycosyl-(NH)-Val-His- or Glycosyl-(NH)-Val-His-AA,

wherein Glycosyl-(NH)-Val represents a non-enzymatically glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin and AA means one or more additional amino acid residues.

In a preferred embodiment AA is a sequence of from 1 to 3 amino acids corresponding to a part of the N-terminal β -chain of all structural variants of human hemoglobin. AA is preferably -Leu-Aa; -Leu-Thr-Aa; -Leu-Thr-Pro-Aa, and Aa designates the remaining number of amino acids in the AA group.

A preferred monoclonal antibody or a fragment thereof according to the invention is obtainable using the hybridoma DAK Hblc-1, which is deposited at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) as accession no. DSM ACC2495.

The invention also relates to a method of producing a monoclonal antibody or a fragment thereof, wherein the antibody is pan-specific, i.e. able to react with all structural variants of glycosylated hemoglobin.

The antibody or a fragment thereof may preferably be obtainable by the method as defined in claims.

The method of producing a monoclonal antibody or a fragment thereof according to the invention comprising the step of raising an antibody against an immunogen comprising a glycosylated peptide chemically linked to an immunogenic carrier material, the glycosylated peptide having

WO 02/088185

PCT/DK02/00267

9

8

from 2 to 5 amino acid units corresponding to the N-terminal in the β -chain of human hemoglobin.
The immunogen is preferably of the formula:

5 [Fructosyl-(NH)-Val-His-AA-R]_n-Carrier

wherein Fructosyl-(NH)-Val represents a nonenzymatic glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin, AA means one or
10 more additional amino acid residues, R is a linking group, Carrier is an immunogenic carrier material, and n is an integer from 1 to the number of available coupling sites on said carrier.

15 In a preferred embodiment AA is one or more additional amino acid residues. In another preferred embodiment AA is a sequence of from 1 to 3 amino acids corresponding to the N-terminal part of the β -chain of all structural variants of human hemoglobin.

20 Linking group R can be essentially any convenient and stable structure. Such linking group R will usually be in the form of an aliphatic chain comprising between 1 and approximately 20 atoms, excluding hydrogen, and including
25 heteroatoms such as nitrogen, oxygen, and sulfur. The glycosylated residue can be joined through a variety of groups to form linking chain R, including methylene, ether, thioether, imino, and the like. One skilled in the art will have a wide variety of linking groups from which
30 to choose to prepare the immunogen.

In a preferred embodiment the carrier is an immunogenic protein or peptide other than human hemoglobin.

WO 02/088185

PCT/DK02/00267

/o

9

The immunogen used to stimulate production of appropriate immunoglobulins in the most general sense will comprise one or more of the glycosylated peptide residues chemically linked to an immunogenic carrier material. The immunogenic carrier material can be selected from any of those conventionally known having functional groups available for coupling to the glycosylated peptide residue. In most cases, the carrier is a protein or polypeptide, although other materials such as carbohydrates, polysaccharides, lipopolysaccharides, nucleic acids, and the like of sufficient size and immunogenicity can likewise be used. For the most part, immunogenic proteins and polypeptides will have molecular weights between 4,000 g/mol and 10,000,000 g/mol, preferably greater than 15,000 g/mol, and more usually greater than 50,000 g/mol. Generally, proteins taken from one animal species will be immunogenic when introduced into the blood stream of another species. Particularly useful proteins are albumins, globulins, enzymes, hemocyanins, glutelins, proteins having significant nonproteinaceous constituents, and the like.

In a preferred embodiment the glycosylated N-terminal peptide residue of the native HbA_{1c}, HbS_{1c} and HbC_{1c} molecule is made accessible to the monoclonal antibody or a fragment thereof of the present invention by appropriate denaturation or digestion of the protein in the sample to be assayed.

In another preferred embodiment the glycosylated fragment can be produced by chemical or enzymatic digestion of naturally occurring glycosylated hemoglobin, e.g. HbA_{1c}, HbS_{1c} and HbC_{1c}. This fragment can be coupled to a carrier using classical coupling procedures, e.g.,

WO 02/088185

PCT/DK02/00267

//

10

glutaraldehyde or carbodiimide, and the conjugate used as an immunogen.

- 5 The antibody or a fragment thereof according to the invention may preferably be produced by the following step:

A. Preparation of peptide immunogen

10

To obtain antibodies against the glycation site of N-terminal valine residue on the β -chain of Hb A, S and C a short peptide (pentamer) is preferably used as an immunogen. The antigen may preferably be Hb1c 4-mer plus
15 cysteine, i.e. fructosyl-V-H-L-T-C.

B. Immunization of mice with human Hb1c peptides

- 20 The general method used to develop hybridomas secreting monoclonal antibodies is well known to a person skilled in the art.

An illustration of the techniques utilized in the practice is described in Journal of Immunological
25 Methods, 1980, 39:285.

Female F1 hybrids of CF1 x BALB/c mice may preferably be used, but mice of other strains may also be used. The
30 immunization plan and the concentration of peptide-carrier complex may be selected to form a sufficient amount of antigenically stimulated lymphocytes. In a preferred embodiment mice are immunized intraperitoneally with 50 micrograms of purified protein derivatives (PPD, a

WO 02/088185

PCT/DK02/00267

/2

11

partly unknown mixture of denatured proteins from *M. tuberculosis*) in complex with the Hb1c peptide four times at 2-week intervals. 3-5 days after the final immunization, the spleen cells are extracted from the animal for fusion.

Monoclonal antibodies produced by hybridomas from other species may also preferably be used.

10

C. Cell fusion

The spleen is aseptically taken out from the immunized mouse, and a suspension of the spleen cells is prepared.

15 The spleen cells are fused with mouse myeloma cells from a suitable cell line in the presence of a suitable fusion promoter. In a preferred embodiment the preferred ratio of the spleen cells to the myeloma cells is from about 20:1 to about 2:1, and a fusion medium is suitably used

20 in an amount of 0.5 to 1.5 ml per about 10^8 spleen cells.

The mouse myeloma cells used for cell fusion are well known. P3-X63-Ag 8.653 may preferably be used as a mouse myeloma cell.

25

The fusion promoter is preferably polyethylene glycol having an average molecular weight of 1000 to 4000 g/mol. Other fusion promoters known in the art can also be used.

30 In a preferred embodiment polyethylene glycol having an average molecular weight of 1500 may be used.

WO 02/088185

PCT/DK02/00267

/3

12

D. Screening of the fused cells

In a separate container (such as a microtiter plate), a mixture composed of the unfused spleen cells, unfused mouse myeloma cells and the fused hybridoma cells is preferably diluted with a selective medium not supporting the unfused mouse myeloma cells and cultured for a period sufficient to kill the unfused mouse myeloma cells (about 1 week). A culture medium having drug resistance (for example, resistance to 8-azaguanine) and not supporting the unfused mouse myeloma cells, such as HAT medium, is used. In the selective medium, the unfused myeloma cells die. Since the unfused spleen cells are untransformed cells, they die away after a certain period of time (1 week). The fused cells, on the other hand, can survive in the selective medium since they have both the tumoral nature of the parent myeloma cells and the nature of the parent spleen cells.

20

E. Determination of glycated hemoglobin by ELISA

The determination is preferably based on the principle of specific immunologic recognition and reaction between a monoclonal antibody and the antigenic epitope to which the antibody uniquely and specifically binds. The recognition and binding may be detected, for example, by an ELISA test, wherein blood samples or calibrators are immobilized on a solid phase support, such as the bottom of a plastic well. By immobilizing human hemoglobin the antigenic epitope may be exposed for antibody binding. The antibody, secondary enzyme-labeled antibody and enzyme substrate may interact with the immobilized antigen (human hemoglobin) thereby antibody-antigen

WO 02/088185

PCT/DK02/00267

14

13

complexes are preferably formed. A number of dilution, incubation and washing steps, then allow separation of bound and free reagents. A colour forming reaction takes place as a result of binding of the antibody to antigen and the consequent reaction of the enzyme upon its substrate interaction. The formation of colour indicates the presence of Hb1c in the blood sample, and the intensity of the colour provides a quantitative measure of the amount of glycated epitope in the sample. The antibody of the present invention may be used to measure glycated hemoglobin in other immunological assay formats known in the art.

15 F. Cloning of the hybridoma cells producing the desired antibody and the production of the antibody

Hybridoma cells capable of producing the desired antibody are cloned by a suitable method such as limiting dilution method, and the desired antibody can be produced by the following method. Hybridoma cells are cultured in a suitable medium for a certain period of time, and the monoclonal antibody produced by the hybridoma cells can be isolated from the supernatant of the culture.

25

The invention also relates to a method of using a monoclonal antibody or a fragment thereof.

30 The antibodies or a fragment thereof according to the invention can preferably be used in conventional manner to react with blood samples containing unknown quantities of glycated hemoglobin and the extent of reaction can be compared with calibrated standards to determine the

WO 02/088185

PCT/DK02/00267

15

14

extent of glycation. The read-out can be by fluorescence, by immunoassay, or the like, by joining suitably readable groups to the monoclonal antibodies in known manner without loss of their binding power for the glycated epitope in HbA1c, HbS1c and HbC1c.

The monoclonal antibodies of the present invention are specific for binding to all structural variants of glycated N-terminal peptide residue preferable found in HbA, HbS and HbC. The antibodies are able to bind to the epitope in the native HbA1c, HbS1c and HbC1c molecule when the epitope is appropriately exposed. Steric access to the epitope can be obtained in any effective manner. Exposure of the epitope in the intact protein is understood to be accomplished by a physical or chemical denaturation or digestion at least in the region of the epitope. Such denaturation or digestion can be localized to the region of the epitope or can involve a more general, or even substantially complete denaturation of the tertiary, and additionally the secondary structure of the protein, or partial or complete digestion of the protein.

In a preferred embodiment the antibody binds specifically to the glycated N-terminal peptide sequence in the β -chain of human hemoglobin A1c, C1c and S1c after having been exposed sufficiently to provide steric access thereto.

Denaturation can be accomplished in a variety of ways including conventional treatment of the protein by physical means such as heat, sonication, high or low pH and, as is preferable, chemical denaturation by interaction with an agent or chaotrope in solution.

Protein denaturation can be most effectively accomplished if combinations of chemical and/or chemical and physical means are used, e.g. guanidine and heat, guanidine and SDS, or guanidine and dithiothreitol. Particularly strong chaotropes such as guanidine are preferred. Of course, denaturing conditions which result in substantial insolubilization, aggregation, or precipitation of the protein implying an insignificant amount of the exposed epitope is accessible to the solution for antibody binding will be avoided. A sufficient amount of the denatured protein must remain in solution or suspension in order to obtain useful immunobinding. The extent of solubilization necessary will depend upon the circumstances of the intended or desired binding.

15 In a preferred embodiment the glycated peptide sequence is exposed to the antibodybinding site by physical or chemical denaturation or digestion.

20 In another preferred embodiment the glycated peptide sequence is exposed to the antibody binding site by denaturation with a chaotropic agent. The chaotropic agents include, without limitation, guanidine, urea, and various detergents such as sodium dodecylsulfate (SDS) and others, without limitation, including deoxycholate and certain bile salts, 3-(3-cholamidopropyl)-dimethyl-ammonio-1-propanesulfonate, organic solvents such as methanol, propanol, acetonitrile and certain salts such as potassium thiocyanate. Non-ionic detergents such as Triton X[®], Tween[®], nonidet NP-40[®], and octyl-glucosides can also function as protein denaturants.

A significant amount of HbA1c, HbS1c and HbC1c in a particular blood sample can be denatured to expose the

WO 02/088185

PCT/DK02/00267

17

16

glycated epitope for antibody binding by combining the sample, e.g., whole blood or red cell hemolysate, with an aqueous solution of the chaotrope present at sufficient concentration to denature any HbA1c, HbS1c and HbC1c in the resulting aqueous mixture. Where whole blood is the sample, the chaotrope also serves to lyse red blood cells to release hemoglobin and to inactivate proteases.

The denaturation process may be significantly accelerated by heating the mixture for a short period of time. It has been found that at temperatures below 37°C, denaturation by the chaotrope can take from one to several hours, whereas at temperatures above 50°C sufficient denaturation can be attained in a minute or less. In order to prevent significant denaturation of the antibody and other proteinaceous reagents to be subsequently added to the mixture, the sample-chaotrope mixture will normally be diluted as a separate step or by addition of reagent solutions to a level that the chaotrope is substantially ineffective to denature such reagents, yet will preserve the exposure of the epitope by preventing significant renaturation of HbA1c, HbS1c and HbC1c.

In a preferred embodiment the antibody binds specifically to a glycated N-terminal peptide sequence in the β -chain of glycated human hemoglobin, which has been adsorbed to a solid phase, wherein the solid phase preferably is selected from the group consisting of polystyrene, divinylbenzene, butadiene, polycarbonate, polyacrylamide, polyacrylic acid, polyacryl amide, polyethylene, polypropylene, fluorinated polymers, poly amides, or co-block polymers hereof, gold, carbon, cellulose, derivatized cellulose, or glass.

WO 02/088185

PCT/DK02/00267

18

17

In a preferred embodiment the monoclonal antibody or a fragment thereof may be used for the determination of the level of glycosylated hemoglobin in human blood. The monoclonal antibody or a fragment thereof may also preferably be used for the determination of the level of glycosylated HbS and/or HbC in human blood.

In a further preferred embodiment the monoclonal antibody or a fragment thereof may be used for the determination of the level of glycosylated heterozygous HbA, HbS and/or HbC in human blood.

In a further preferred embodiment there is provided a turbidimetric two-microsphere method for determination of the level of glycosylated hemoglobin. This method comprises mixing a sample with microspheres and microspheres on which the monoclonal antibody or a fragment thereof is bound; the latter is referred to as immunoparticles, or with microspheres and subsequently with immunomicrospheres. The resulting immunoagglutinates is measured turbidimetrically.

The microspheres are preferably selected from the group consisting of polystyrene, divinylbenzene, butadiene, polycarbonate, polyacrylamide, polyacrylic acid, polyacryl amide, polyethylene, polypropylene, fluorinated polymers, poly amides, or co-block polymers hereof, gold, or silica.

Microspheres to be used in the method of the present example have a particle size of from 0.01 to 5 micrometers, which are used by dispersing in a usually used buffer solution.

WO 02/088185

PCT/DK02/00267

19

18

The following experimental non-limiting examples are intended to illustrate certain features and embodiments of the invention.

5 **Example 1:**

The 1-deoxyfructosyl-Val-His-AA peptide was synthesized using the solid-phase method of Sheppard and Atherton as described in international patent application no. WO 86/03494. The peptide was synthesized with fluorenylmethoxycarbonyl (Fmoc) amino acids, using PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate) and HOBT (N-hydroxy-benzotriazole) as coupling reagents.

15 Cysteine and histidin were used with trityl (trt), and threonin with tertiary butyl (tBu) side-chain protection. Leucin was used without side-chain protection.

20 The fructose amino acid conjugate, N-(deoxy-D-fructos-1-yl)-L-valine, was synthesized by modifications of known methods. In short, Di-isopropyliden protected glucose was activated as the trifluate and conjugated to the benzyl ester of valine. The resulting 2,3:4,5-Di-O-isopropyliden-N-(deoxy-D-fructos-1-yl)-L-valine-benzyl ester was hydrogenated over Pd/C in ethanol, lyophilized, and finally used in the peptide synthesis without Fmoc protection. The purity of the product was verified by TLC analysis and ¹H or ¹³C NMR spectra.

30 The peptide was synthesized by automated solid phase synthesis (Crystal instrument from Calbiochem-Novabiochem Ltd) cleaved from the resin with TFA (trifluoroacetic acid) - TES (triethylsilan) - Glycerol - H₂O (90% : 4% :

WO 02/088185

PCT/DK02/00267

20

19

3% : 3%), precipitated and washed with methyl-tertiary-butyl-ether, centrifuged and lyophilised.

5 Purification was performed by HPLC (C_{18} reversed-phase column, using a step-wise gradient of typically 5-25% MeCN in 0.1% TFA). The identity of the peptide was verified by MALDI-TOF mass spectroscopy and the purity was finally assured by an analytical HPLC analysis to be better than 90%.

10

Example 2:

15 Four weeks before first immunization Bacille Calmette Guérin (BCG) vaccine was injected to prime the immune response in the mice as described in Giba Foundation Symposium, 1986, 119: 25.

20 To make the peptide immunogen it was conjugated to a carrier mixture, Tuberculin PPD. Peptide and carrier (PPD) were conjugated with glutardialdehyde, i.e. $CHO-(CH_2)_3-CHO$. Glutardialdehyde reacts with amino-groups on the peptide and the carrier.

The immunogens thus have the following formula:

25

Fructosyl-Val-His-Leu-Thr-Cys-CH- $(CH_2)_3$ -CH-PPD

Example 3:

30

Female F1 hybrids of CF1 x BALB/c mice (12 weeks old) primed with BCG-vaccine were immunized with the peptide-PPD complex described above. Mice were immunized four times intraperitoneally with a mixture of 50 micrograms

WO 02/088185

PCT/DK02/00267

21

20

of peptide-PPD complex dissolved in phosphate buffered saline (PBS) and aluminium hydroxide (1 mg/dose). A mouse with high titer of antibodies against glycosylated hemoglobin was selected for fusion. Three days after the final immunization, spleen cells from the immunized mouse was used for cell fusion.

Spleen cells from the selected immunized mouse and myeloma cells of BALB/c origin (P3-X63-Ag 8.653) were mixed in a ratio of about 3:1 and fused in the presence of 50% polyethylene glycol 1500 (a product of Roche Diagnostics GmbH) (J. W. Goding: Monoclonal Antibodies: Principle and Practice, Academic Press, San Diego, 1996). The fused cells were suspended in RPMI-1640 medium, containing 10% FCS, hypoxanthine, aminopterin and thymidine (HAT media) to give a concentration of 1×10^6 cells/mL. The suspension was distributed to 96-well microplates (Costar) in an amount of 200 μ L per well.

The cells were incubated in a CO₂ incubator (5% CO₂, 37°C). After 14 days hybridoma cells were screened.

Supernatants containing antibodies produced by the hybridoma cell cultures were detected by an ELISA technique using microtiter plates coated with hemoglobin calibrator containing 4.7% HbA1c and 16.7% HbA1c as antigens. 120 μ g/mL of the calibrators were immobilized onto plastic microtiter wells using a carbonate/bicarbonate buffer (pH 9.6) for 15 minutes at 37°C. After a washing step that removes unbound antigen the hybridoma culture supernatants diluted in 50 mM Tris-HCl buffer containing 0.1% Tween 20 were added to each well and allowed to react for one hour at room temperature. After another washing step to remove unbound

WO 02/088185

PCT/DK02/00267

22

21

antibody a conjugate of a goat anti-mouse antibody and the enzyme horseradish peroxidase was added. After incubation and a third washing step the bound conjugate was detected by reaction with the substrate 3,3',5,5'-tetramethylbenzidine (TMB). The reaction was stopped with sulphuric acid and the absorbance at 450 nm was measured. Only cells from positive wells reacting with glycosylated hemoglobin calibrators, 4.7% HbA1c and 16.7% HbA1c, with increasing intensity were selected.

These positive cells were cloned three times by limiting dilution method. The resulting clone, DAK Hb1c-1, was suspended in 90% FCS; 10% DMSO and stored at -150°C.

Example 4:

A linear antibody epitope mapping is the systematic screening of all possible peptides derived from a polypeptide or protein sequence. Epitope mapping yields information on the linear stretch of amino acids and derivatives that form an interaction site to a given antibody.

The specificity of DAK Hb1c-1 for the glycosylated epitope of hemoglobin β -chain has been determined.

A peptide library containing 1-5 amino acid long peptides manually synthesized from the C- to the N-terminal end on a solid phase (Abimed membrane) using Fmoc as protecting group of the N-terminal end of amino acids. Detailed description of peptide synthesis is described in Example 1. The following four peptides were synthesized:

Fructosyl-Hb β -chain

WO 02/088185

PCT/DK02/00267

23

22

1-5 amino acid long peptide analogs of N-terminal 1-deoxyfructosyl-hemoglobin β -chain.

Hb β -chain

- 5 1-5 amino acid long peptide analog of N-terminal hemoglobin β -chain (minus 1-deoxyfructosyl β -chain; backbone control).

Hb α -chain

- 10 5 amino acid long peptide analog of N-terminal hemoglobin α -chain (negative sequence control).

Fructosyl-Hb α -chain

- 15 5 amino acid long peptide analogs of hemoglobin α -chain incl. 1-deoxyfructosyl at the N-terminal valine residue (negative 1-deoxyfructosyl sequence control).

20 The membranes with peptide spots were incubated with DAK Hblc-1 and an irrelevant antibody (a negative antibody control, anti-troponin I, DAKO, product no. O 9528). Thereafter, the membranes were incubated with HRP conjugated goat-anti-mouse antibodies (DAKO, product no. P 0447) and visualized using 3-amino-9-ethylcarbazole as substrate.

25

Number of Amino Acids	5	4	3	2	1
Peptide name					
Fructosyl-Hb β -chain	++	++	++	+	-
Hb β -chain	-	-	-	-	-
Hb α -chain	-	N.D.	N.D.	N.D.	N.D.
Fructosyl-Hb α -chain	-	N.D.	N.D.	N.D.	N.D.

WO 02/088185

PCT/DK02/00267

24

23

Table 1: Spot peptide library for epitope mapping of anti-Hb1c antibody clone DAK Hb1c-1. Intensity of peptide spots after incubation with anti-Hb1c antibody clone DAK Hb1c-1 followed by visualization are shown; strong intensity (++), medium intensity (+), negative (-), not determined (N.D.).

DAK Hb1c-1 reacts moderately with 1-deoxyfructosyl 2-mer of hemoglobin β -chain, strongly with 1-deoxyfructosyl 3-mer to 5-mer hemoglobin β -chain (see Table 1). DAK Hb1c-1 does not react with 1-deoxyfructosyl-valine alone. DAK Hb1c-1 does not react with non-fructosylated peptides of hemoglobin β -chain, 1-deoxyfructosyl 5-mer hemoglobin α -chain and 5-mer hemoglobin α -chain. In contrast to DAK Hb1c-1, the negative antibody control did not react with any of the peptides.

In summary, the specificity of anti-Hb1c antibody clone DAK Hb1c-1 for the 1-deoxyfructosylated N-terminal part of hemoglobin β -chain has been demonstrated.

Example 5:

Peptides, proteins from a whole blood sample, red cellhemolysate or calibrators were allowed to bind to the surface of a 96-well microtiter plate. Following a washing step to remove unbound proteins, the monoclonal antibody, DAK Hb1c-1, was added. After incubation, the unbound antibody was washed away, and a conjugate of a goat-anti-mouse antibody and the enzyme horseradish peroxidase was added. After another washing step the bound conjugate was detected by reaction with 3,3',5,5'-tetramethylbenzidine (TMB) substrate. The reaction was

WO 02/088185

PCT/DK02/00267

25

24

stopped by adding acid to give a colorimetric end point that was read spectrophotometrically.

Example 6:

5

HbA₀ is identical to HbA_{1c} except that there is no 1-deoxyfructosyl linked to the amino-terminus of the hemoglobin β -chain.

10 The reaction of DAK Hb1c-1 with HbA₀ and 17-mer peptides of HbC₀ and HbS₀ was analysed by the ELISA described above.

15 When DAK Hb1c-1 was added to microtiter plates coated with either HbA₀ or 17-mers peptides of HbC₀ or HbS₀ the values measured were no higher than the background values.

20 In conclusion, no reactivity of DAK Hb1c-1 with HbA₀, HbC₀ or HbS₀ was demonstrated.

Example 7:

25 The reaction of DAK Hb1c-1 with native HbA_{1c} in solution was analyzed in a competitive ELISA. HbA_{1c} calibrator in solution was pre-incubated with DAK Hb1c-1 antibody to test if any antigen-antibody complexes were formed. The solution of antigen-antibody was then transferred to microtiter-wells, where the same antigen was adsorbed on
30 the solid phase (immobilized antigen). The antigen in solution was in excess. The antibody not in complex with the antigen in solution was able to bind to the immobilized antigen. A secondary antibody as described above detected the bound DAK Hb1c-1 antibody. As a

WO 02/088185

PCT/DK02/00267

26

25

control the same amount of antibody not pre-incubated with HbA_{1c} calibrator in solution was added to one lane of the microtiter plate and followingly subjected to the protocol described above.

5

No effect of pre-incubation of DAK Hb1c-1 antibody with HbA_{1c} in solution was observed, concluding that DAK Hb1c-1 does not bind to native HbA_{1c} in solution.

10 **Example 8:**

The reactivity of DAK Hb1c-1 and HEM13 with human blood containing structural variants of hemoglobin have been analysed by isoelectric focusing (IEF) followed by
15 immunoblotting.

Red cell hemolysate containing HbCA, HbSS and HbAA were electrophoretically separated by Pharmacia's PhastSystem using IEF polyacrylamide gel (PhastGel IEF - 5-8,
20 Pharmacia) in order to separate structural variants of hemoglobin. Following, the gels were subjected to press blotting of proteins onto a PVDF membrane and blocked for non-specific staining with a blocking agent. The membranes were then incubated with DAK Hb1c-1 (see figure
25 1 A) and HEM13 (see figure 1 B), respectively. HRP-conjugated goat anti-mouse immunoglobulin (DAKO A/S, product no. P 0447) was used as secondary antibody. Finally, HRP activity were visualized using DAKO liquid DAB+ Substrate-Chromogen System (DAKO A/S, product no. K
30 3468).

A distinct HbS_{1c} band with strong intensity was identified analysing red cell hemolysate from a non-diabetic patient with homozygous HbSS with the monoclonal

WO 02/088185

PCT/DK02/00267

27

26

antibody, clone DAK Hb1c-1 (see Figure 1, A). In contrast, weak HbS_{1c} staining was revealed with HEM13 analyzing red cell hemolysate from a patient with homozygous HbSS (see Figure 1 B).

5

Analyzing red cell hemolysate from a non-diabetic individual with heterozygous hemoglobin CA, HbC_{1c} bands with strong and moderate intensity were visualized with DAK Hb1c and HEM 13, respectively. A strong HbA_{1c} band with approximately the same intensity was observed for both DAK Hb1c and HEM13.

10

As a control a pool of red cell hemolysate from diabetes patients with hemoglobin A was also tested. Other variants of hemoglobin not related to C and S may be a part of this patient pool. Each membrane was incubated with substrate until HbA_{1c} bands with approximately the same intensity were revealed on both DAK Hb1c-1 and HEM 13.

15

20

Conclusively, we have demonstrated that our pan-specific monoclonal glycosylated hemoglobin antibody, clone DAK Hb1c-1, reacts with HbA_{1c}, HbS_{1c} and HbC_{1c} from human blood samples.

25

Example 9:

Adult hemoglobin is a tetrameric protein, consisting of two α -subunits and two β -subunits, $\alpha_2\beta_2$. Structural variants of Hb contain genetically determined changes in the primary structure of the peptides. A, S and C differ only in the sixth amino acid residue from the N-terminal of the β -chain:

30

WO 02/088185

PCT/DK02/00267

2f

27

HbA_{1c} fructosyl-V-H-L-T-P-E-E-K-T-A-V-N-A-L-W-G-K.....
HbC_{1c} fructosyl-V-H-L-T-P-K-E-K-T-A-V-N-A-L-W-G-K.....
HbS_{1c} fructosyl-V-H-L-T-P-V-E-K-T-A-V-N-A-L-W-G-K.....

5 To test if the antibody is pan-specific, i.e. reacting with all structural variants of glycosylated hemoglobin, we have tested the reaction of DAK Hb1c-1 antibody with three 17-mer glycosylated peptides corresponding to the N-terminal end of glycosylated hemoglobin A, C and S β -subunits
10 by ELISA (Figure 2). The peptide sequences are shown above. As a control a pool of red cell hemolysate from diabetes patients with hemoglobin A was also tested by DAK Hb1c-1 ELISA.

15 Monoclonal HbA_{1c} antibody clone HEM13 has been raised by immunization of mice with purified hemoglobin A_{1c}. We have compared the reaction of DAK Hb1c-1 and HbA_{1c} antibody clone HEM13 with the glycosylated hemoglobin peptides and erythrocyte extract of hemoglobin A.

20 The affinity of DAK Hb1c-1 for the glycosylated hemoglobin peptides is significantly stronger than the affinity of HEM13 for the same peptides as judged by comparing the titration curves (see Figure 2 and Figure 3).

25 HEM13 shows different reactions with the A_{1c}, S_{1c} and C_{1c} peptides (Figure 3) whereas DAK Hb1c-1 reacts in the same manner with all three peptides (Figure 1). These results demonstrates that the DAK Hb1c-1 antibody is pan-
30 specific.

This example verifies that an antibody that reacts in the same manner with all structural variants of glycosylated hemoglobin can be raised by injection of a glycosylated 5-mer

WO 02/088185

PCT/DK02/00267

29

28

peptide corresponding to the first 4 amino acids of all structural variants of glycated hemoglobin A, C and S β -chain plus one irrelevant amino acid (here cysteine).

5 Example 10:

Blood samples from the "ERL Educational Programme" have been tested. The HbA_{1c}-percentage of each sample has been determined by cation-exchange chromatography (IFCC/DCCT-
 10 values in table 2; IFCC: International Federation of Clinical Chemistry; DCCT: Diabetes Control and
 15 hemoglobin β -subunits, e.g. AS or AC.

	IFCC	DCCT	DAK-Hb1c-1
HbAA	5.0	6.0	5.4
HbAA	5.7	6.5	6.2
HbAA	6.3	7.0	6.5
HbAA	7.0	7.6	7.9
HbAA	7.5	8.0	8.5
HbAA	8.1	8.4	8.7
HbAA	8.8	8.9	9.4
HbAA	9.4	9.4	9.7
HbAA	10.0	9.8	10.0
HbAS	4.3	5.5	4.7
HbAC	4.3	5.5	4.0

Table 2: Percentage of Hb_{1c} in blood samples. The blood sample AC and AS contain heterozygotic hemoglobin.

20 At a range from 4-10% the HbA_{1c} percentage was measured in whole blood samples using DAK Hb_{1c}-1 in the above-mentioned ELISA format. HbA_{1c} percentages at the same

WO 02/088185

PCT/DK02/00267

30

29

levels as the values determined by reference methods recommended by IFCC and DCCT were obtained.

In summary, it is demonstrated that the antibody in question has the correct specificity against HbA1c, HbC1c and HbS1c.

Example 11:

- 10 The level of glycated hemoglobin was measured in a two-microsphere turbidimetry assay. In this set-up glycated hemoglobin was non-covalently adsorbed to microspheres followed by detection of the antigen with glycated hemoglobin antibodies covalently bound to microspheres.
- 15 The resulting network constituting antigen microspheres complexed with immunomicrospheres were measured turbidimetrically.

WO 02/088185

30

PCT/DK02/00267

31

Applicant's or agent's file reference P062 WD 01	International application No. RO/DK 2 5 APR 2002
---	---

**INDICATIONS RELATING TO DEPOSITED MICROORGANISMS
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>7</u> , line <u>19</u> .	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH	
Address of depository institution (including postal code and country) Mascheroder Weg 1b D-38124 Braunschweig	
Date of deposit 6 March 2001	Accession Number DSM ACC2495
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general names of the indications e.g. "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on: 16 MAY 2002 (16.05.02)
Authorized officer	Authorized officer <i>Jean-Luc MARTIN</i>

WO 02/088185

31

PCT/DK02/00267

32

Claims:

1. A monoclonal antibody or a fragment thereof, wherein said antibody is pan-specific, i.e. able to react with all structural variants of glycosyl-hemoglobin, said antibody or a fragment thereof comprises an antibody combining site, which binds specifically to a glycosylated peptide residue of the formula:
- 10 Glycosyl-(NH)-Val-His- or Glycosyl-(NH)-Val-His-AA,
- wherein Glycosyl-(NH)-Val represents a non-enzymatically glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin and AA means 1 to 3 amino acid residues.
2. A monoclonal antibody or a fragment thereof according to claim 1, wherein AA is a sequence of from 1 to 3 amino acids corresponding to a part of the N-terminal β -chain of all structural variants of human hemoglobin.
3. A monoclonal antibody or a fragment thereof according to claim 2, wherein AA is -Leu-Aa; -Leu-Thr-Aa; -Leu-Thr-Pro-Aa, and Aa designates the remaining number of amino acids in the AA group.
4. A monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody is obtainable by hybridoma DAK Hb1c-1 deposited at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) as accession no. DSM ACC2495.
5. A monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody

WO 02/088185

32

PCT/DK02/00267

33

is pan-specific, i.e. able to react with all structural variants of human hemoglobin including HbA1c, HbC1c and HbS1c.

5 6. A monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody being obtainable by a method as defined in any one of the claims 7-11.

10 7. A method of producing a monoclonal antibody or a fragment thereof, wherein said antibody has been raised against an immunogen comprising an antigen obtainable by a glycosylated peptide chemically linked to an immunogenic carrier material, the glycosylated peptide having from 2 to 5
15 amino acid units corresponding to the N-terminal β -chain of human hemoglobin.

8. A method of producing a monoclonal antibody or a fragment thereof according to claim 7, wherein the
20 immunogen is of the formula:

[Fructosyl-(NH)-Val-His-AA-R] $_n$ -Carrier

wherein Fructosyl-(NH)-Val represents a non-enzymatic
25 glycosylated valine residue, His represents the second amino acid in the native β -chain of hemoglobin, AA means one or more additional amino acid residues, R is a linking group, Carrier is an immunogenic carrier material, and n is an integer from 1 to the number of available coupling
30 sites on said carrier.

9. A method of producing a monoclonal antibody or a fragment thereof according to claim 8, wherein AA is a sequence of from 1 to 3 amino acids corresponding to a

part of the N-terminal β -chain of all structural variants of human hemoglobin.

10. A method of producing a monoclonal antibody or a fragment thereof according to claim 8, wherein said carrier is immunogenic proteins or peptides other than human hemoglobin.

11. A method of using a monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody binds specifically to said glycosylated N-terminal peptide sequence of human hemoglobin A1c, C1c and S1c β -chains after having been exposed sufficiently to provide steric access thereto.

12. A method of using a monoclonal antibody or a fragment thereof according to claim 11, wherein said glycosylated peptide sequence is exposed to the antibody combining site by physical or chemical denaturation or digestion.

13. A method of using a monoclonal antibody or a fragment thereof according to claims 11-12, wherein said glycosylated peptide sequence is exposed to the antibody combining site by denaturation with a chaotropic agent.

14. A method of using a monoclonal antibody or a fragment thereof according to claim 13, wherein the chaotropic agent is selected from the group consisting of guanidine, urea, potassium-thiocyanate, or detergent.

15. A method of using a monoclonal antibody or a fragment thereof according to any one of the preceding claims, wherein said antibody binds specifically to a glycosylated N-terminal peptide sequence of the β -chain of all

structural variants of human hemoglobin, which has been adsorbed to a solid phase.

16. A method of using a monoclonal antibody or a fragment thereof according to claim 15, wherein the solid phase preferably is constituted by a material selected from the group consisting of microtiter plates, micro particles, membranes, tubes, pins, chips or discs.
17. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycated hemoglobin in human blood.
18. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycated HbS and/or HbC in human blood.
19. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycated heterozygous HbA, HbS and/or HbC in human blood.
20. Use of a monoclonal antibody or a fragment thereof according to anyone of the preceding claims for the determination of the level of glycated hemoglobin measured in a two-particle turbidimetric assay.

WO 02/088185

PCT/DK02/00267 36

1/3




Figure 1




Figure 2: Reactivity of DAK Hb1c-1 with red cell hemolysate from patients with hemoglobin AA (HbAA) and 17-mer glycosylated peptides corresponding to hemoglobin A1c, C1c and S1c as determined by an ELISA test.




Figure 3. Reactivity of HEM13 with red cell hemolysate from patients with hemoglobin AA (HbAA) and 17-mer glycated peptides corresponding to hemoglobin A1c, C1c and S1c as determined by an ELISA test.

【 国際調査報告 】

INTERNATIONAL SEARCH REPORT		International Classification No PCT/DK 02/00267
A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07K16/28 G01N33/577 G01N33/68 G01N33/72		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07K G01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, WPI Data, EPO-Internal, PAJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 34 39 610 A (BOEHRINGER MANNHEIM GMBH) 30 April 1986 (1986-04-30) page 5, column 26-29 page 6, line 10-21 page 7, column 14-20 examples 4,6	1-20
X	US 4 727 036 A (KNOWLES WILLIAM J ET AL) 23 February 1988 (1988-02-23) column 1, line 47 -column 2, line 15 column 3, line 60 -column 4, line 10 column 10, line 13 -column 11, line 27 claims	1-20
A	EP 0 315 864 A (MILES INC) 17 May 1989 (1989-05-17) page 2, line 10-20 page 7, line 55 -page 11, line 39	20
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents:		
A document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (see specifics) *C* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed		*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *S* document member of the same patent family
Date of the actual completion of the international search 12 May 2003		Date of mailing of the international search report 19/05/2003
Name and mailing address of the ISA European Patent Office, P.B. 5518 Patentlaan 2 NL - 2280 HV Rijswijk Tel: (+31-70) 340-2040, Tx: 31 651 epo nl, Fax: (+31-70) 340-3018		Authorized officer COVONE-VAN HEES, M

INTERNATIONAL SEARCH REPORT		International application No. PCT/DK 02/00267
Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)		
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:		
1.	<input checked="" type="checkbox"/> Claims Nos.:	because they relate to subject matter not required to be searched by this Authority, namely:
		Although claims 7-19 are directed to a method of treatment and/or diagnosis of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	<input type="checkbox"/> Claims Nos.:	because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.	<input type="checkbox"/> Claims Nos.:	because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)		
This International Searching Authority found multiple inventions in this International application, as follows:		
1.	<input type="checkbox"/> As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.	
2.	<input type="checkbox"/> As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.	
3.	<input type="checkbox"/> As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:	
4.	<input type="checkbox"/> No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:	
Remark on Protest		<input type="checkbox"/> The additional search fees were accompanied by the applicant's protest.
		<input type="checkbox"/> No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.
PCT/DK 02/00267

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
DE 3439610	A	30-04-1986	DE 3439610 A1	30-04-1986
US 4727036	A	23-02-1988	US 4647654 A	03-03-1987
			AT 80951 T	15-10-1992
			AT 98776 T	15-01-1994
			AU 3375589 A	17-08-1989
			AU 594651 B2	15-03-1990
			AU 4926085 A	08-05-1986
			CA 1339952 A1	14-07-1998
			DE 3586679 D1	29-10-1992
			DE 3586679 T3	15-10-1998
			DE 3587687 D1	27-01-1994
			DE 3587687 T2	07-04-1994
			DK 130791 A , B,	04-07-1991
			DK 494085 A	30-04-1986
			EP 0185870 A2	02-07-1986
			EP 0316306 A2	17-05-1989
			ES 8705633 A1	16-07-1987
			ES 8800435 A1	01-01-1988
			FI 854187 A , B,	30-04-1986
			FI 104376 B1	14-01-2000
			IE 63731 B1	14-06-1995
			IE 63768 B1	14-06-1995
			IL 76827 A	29-11-1990
			IL 91034 A	29-11-1990
			JP 2858534 B2	17-02-1999
			JP 7051087 A	28-02-1995
			JP 7023891 B	15-03-1995
			JP 61172064 A	02-08-1986
			NZ 213959 A	27-03-1990
			NZ 228194 A	27-03-1990
			US 4658022 A	14-04-1987
EP 0315864	A	17-05-1989	US 4970171 A	13-11-1990
			AT 83561 T	15-01-1993
			AU 2465588 A	11-05-1989
			CA 1340220 A1	15-12-1998
			DE 3876761 D1	28-01-1993
			DE 3876761 T2	22-04-1993
			DK 622788 A	10-05-1989
			EP 0315864 A1	17-05-1989
			ES 2043762 T3	01-01-1994
			IE 62086 B1	14-12-1994
			IL 87682 A	18-08-1992
			JP 1155268 A	19-06-1989
			JP 2683572 B2	03-12-1997

フロントページの続き

(51)Int.Cl.⁷ F I テーマコード(参考)
 G 0 1 N 33/72 G 0 1 N 33/72 A

(81)指定国 AP(GH,GM,KE,LS,MW,MZ,SD,SL,SZ,TZ,UG,ZM,ZW),EA(AM,AZ,BY,KG,KZ,MD,RU,TJ,TM),EP(AT, BE,CH,CY,DE,DK,ES,FI,FR,GB,GR,IE,IT,LU,MC,NL,PT,SE,TR),OA(BF,BJ,CF,CG,CI,CM,GA,GN,GQ,GW,ML,MR,NE,SN, TD,TG),AE,AG,AL,AM,AT,AU,AZ,BA,BB,BG,BR,BY,BZ,CA,CH,CN,CO,CR,CU,CZ,DE,DK,DM,DZ,EC,EE,ES,FI,GB,GD,GE, GH,GM,HR,HU,ID,IL,IN,IS,JP,KE,KG,KP,KR,KZ,LC,LK,LR,LS,LT,LU,LV,MA,MD,MG,MK,MN,MW,MX,MZ,NO,NZ,OM,PH,P L,PT,RO,RU,SD,SE,SG,SI,SK,SL,SZ,TJ,TM,TN,TR,TT,TZ,UA,UG,US,UZ,VN,YU,ZA,ZM,ZW

(72)発明者 シモンセン, アンナ, カリーナ, ウィボーグ
 デンマーク国 ケイジーエス・リングビー ディーケイ - 2 8 0 0 スヴェリゲスヴェイ 1 3 エ
 イ

Fターム(参考) 2G045 AA13 DA45 DA48 FB03
 4B024 AA11 BA43 GA05 HA15
 4B064 AG27 CA10 CA20 CC24 CD20 DA13
 4H045 AA11 BA10 CA40 DA76 EA50 FA72 FA74

专利名称(译)	泛特异性单克隆抗体		
公开(公告)号	JP2005504512A	公开(公告)日	2005-02-17
申请号	JP2002585482	申请日	2002-04-25
[标]申请(专利权)人(译)	DAKOCYTOMATION丹麦		
申请(专利权)人(译)	DakoCytomation , 丹麦 , ACTY洛杉矶萝卜		
[标]发明人	ジェンセン,スージー,ステイーン ピー,カート シモンセン,アンナ,カリーナ,ウィボーグ		
发明人	ジェンセン,スージー,ステイーン ピー,カート シモンセン,アンナ,カリーナ,ウィボーグ		
IPC分类号	G01N33/53 C07K16/00 C07K16/18 C07K16/28 C12N15/02 C12P21/08 G01N33/577 G01N33/72		
CPC分类号	C07K16/00 C07K16/28 C07K2317/34		
FI分类号	C12N15/00.C C07K16/18.ZNA C12P21/08 G01N33/53.V G01N33/577.B G01N33/72.A		
F-TERM分类号	2G045/AA13 2G045/DA45 2G045/DA48 2G045/FB03 4B024/AA11 4B024/BA43 4B024/GA05 4B024/HA15 4B064/AG27 4B064/CA10 4B064/CA20 4B064/CC24 4B064/CD20 4B064/DA13 4H045/AA11 4H045/BA10 4H045/CA40 4H045/DA76 4H045/EA50 4H045/FA72 4H045/FA74		
优先权	200100656 2001-04-26 DK		
其他公开文献	JP2005504512A5 JP4318458B2		
外部链接	Espacenet		

摘要(译)

本发明描述了泛特异性单克隆抗体，用于测定人血液样品中糖化血红蛋白的所有结构变体的量。确定个体血液中血红蛋白的糖化速率提供了糖尿病患者中葡萄糖水平控制的有用指标。