



(11)

EP 2 949 673 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
02.12.2015 Bulletin 2015/49

(21) Application number: **15172601.5**

(22) Date of filing: **27.04.2010**

(51) Int Cl.:
C07K 16/28 ^(2006.01) **A61K 39/395** ^(2006.01)
A61P 35/02 ^(2006.01) **C07K 16/46** ^(2006.01)
G01N 33/53 ^(2006.01) **G01N 33/574** ^(2006.01)
G01N 33/577 ^(2006.01) **C12N 15/02** ^(2006.01)
C12N 15/09 ^(2006.01) **C12P 21/08** ^(2006.01)
C12R 1/91 ^(2006.01)

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK SM TR**

(30) Priority: **27.04.2009 US 172923 P**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
10769764.1 / 2 426 148

(71) Applicant: **Kyowa Hakko Kirin Co., Ltd.**
Tokyo 100-8185 (JP)

(72) Inventors:
• **TAWARA, Tomonori**
Tokyo 194-8533 (JP)

• **TAKAYANAGI, Shinichiro**
Tokyo 194-8533 (JP)
• **INAGAKI, Yoshimasa**
Tokyo 194-8533 (JP)

(74) Representative: **Tuxworth, Pamela M.**
J A Kemp
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

Remarks:
This application was filed on 17-06-2015 as a
divisional application to the application mentioned
under INID code 62.

(54) **ANTI-IL-3RA ANTIBODY FOR USE IN TREATMENT OF BLOOD TUMOR**

(57) The present invention provides an antibody to human IL-3R α chain, which does not inhibit IL-3 signaling and binds to B domain of the human IL-3R α chain but does not bind to C domain of the human IL-3R α chain; a composition for preventing or treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood of a subject, which comprises the

antibody to human IL-3R α as an active ingredient; and a method for treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood, which comprises administering, to a subject, a composition comprising the IL-3R α antibody as an active ingredient.

EP 2 949 673 A1

Description

TECHNICAL FIELD

5 **[0001]** This invention relates to an antibody to human IL-3R α protein (another name: human CD 123). The invention also relates to an invention of a therapeutic agent and diagnostic agent for myelocytic malignant tumors, particularly acute myeloid leukemia (AML), which comprises a human IL-3R α antibody as an active ingredient.

BACKGROUND OF THE INVENTION

10

Regarding malignant tumor:

[0002] A malignant tumor (cancer) is the first leading cause of death in Japan and the number of patients is increasing every year, and the development of a drug and a therapeutic method having high efficacy and safety is strongly desired. As the cause of forming a malignant tumor, there is a mutation of DNA by radiation, ultraviolet rays and various carcinogenic substances. Studies on malignant tumors have been focused on molecular biological identification of these genetic changes. As a result, it is considered that tumorigenic transformation is induced by accumulation of a large number of mutations and the like. It has been shown by a cell line model and the like that some decisive mutations directly connected with the tumorigenic transformation. Regarding leukemia as one of the objective diseases of the invention, many chromosomal abnormalities have been identified and classified. In many of the case, translocation of chromosome is found and the some genes associated with chromosomal translocation have already been identified in principle chromosomal translocations. By analyses of functions of the translocation related genes, a case has been found that these genes are concerned in the onset of leukemia.

25 Regarding cancer stem cell:

[0003] On the other hand, a so-called cancer stem cell hypothesis has been proposed for a long time from the viewpoint of cell biology, stating that stem cell is the origin of a malignant tumor similar to the normal tissue. The stem cell is defined as a cell having autonomous replication ability and pluripotency and generally divided roughly into totipotency stem cell and tissue stem cell. Tissue stem cells are originated from specific tissues and organs such as of blood system, liver, nerve system and the like and present at an extremely low frequency. Among them, hematopoietic stem cell has been studied most frequently. It has been reported that a hematopoietic system can be reconstructed over a long period of time by transplanting one hematopoietic stem cell into a mouse in which the hematopoietic system was destructed by a lethal dose of irradiation (Non-patent Document 1). Different from the normal stem cell, studies on cancer stem cells have been delayed for a prolonged period of time since their true nature could not be found. However, a cancer stem cell has been identified for the first time in acute myeloid leukemia, in 1997 by Dick *et al.* Thereafter, the presence of cancer stem cells has been reported in various malignant tumors. In summing up, cancer stem cells are present at a frequency of several % or less of the whole tumor and rare as well as normal stem cells. It is considered that the remaining cells which form the tumor are tumor precursor cells in which proliferation ability is limited or tumor cells.

40 **[0004]** By these reports, it was shown that hierarchy is present even in tumor similar to the normal tissue, and the cancer stem cell residing at this peak (origin) has strong tumor forming ability.

Characteristics and therapeutic problems of cancer stem cells:

45 **[0005]** In summing up many reports, it is considered that cancer stem cells are maintaining various characteristics possessed by the normal stem cells. Examples of similarities include the rarity of the cells, a microenvironment (niche) in which the cell exists, expression of a multiple drug resistance gene, cell cycle arrest, and the like.

[0006] Particularly, the characteristics that they express a group of multiple drug resistance genes and are at the interphase of cell cycle similar to the normal stem cells could become a therapeutically great problem. A multiple drug resistance gene BCRP is a pump which impairs the drug efficacy by eliminating various antitumor agents into outside of cells, and a method for collecting stem cells making use of the activity has been reported (Non-patent Document 2). In addition, their presence at the interphase of cell cycle under a state of "arresting" (Non-patent Document 3) is causing reduction of sensitivity for many antitumor agents and radiations which targets the quick cell growth of cancer (Non-patent Documents 4 and 5).

55 **[0007]** Based on the above characteristics, it is considered that the cancer stem cell which exhibiting resistance to the therapy is a cause of tumor regeneration.

Regarding molecular target drug

[0008] Three main courses of the treatment of a malignant tumor include of antitumor agent therapy, radiation therapy and excision. The blood tumor is limited to the antitumor agent therapy and radiation therapy, and as described in the above, the cancer stem cell can have a resistance to these treatments. Another problem is that side effects are large since these two treatments affect the entire body. It is a molecular target drug that is expected as a resolving means for this problem. It has a possibility to reduce side effects by exhibiting its drug efficacy only in the cell expressing the target molecule.

[0009] Examples of typical drugs of the molecular target drug in the field of blood diseases include imatinib and rituximab. Imatinib targets at a leukemia-causing factor called Bcr-Abl produced by a chromosomal abnormality (Philadelphia chromosome) which is observed in 95% of CML patients. This is a low molecular weight drug which induces suicide of leukemia cell by inhibiting function of Bcr-Abl. Rituximab is a therapeutic antibody which recognizes CD20 as a surface molecule on a B cell and has an antitumor effect on a malignant tumor of B cell (non-Hodgkin lymphoma and the like). On the other hand, molecular target drugs for AML are few, and there is only an agent gemtuzumab-ozogamicin (Mylotarg) in which an antibiotic calicheamicin is linked to a monoclonal antibody for CD33 known as an AML cell surface antigen. However, it is the present situation that the use of Mylotarg is limited because of its strong toxicity which is considered to be derived from calicheamicin in addition to the problem that therapeutic range is narrow. Based on the above, it can be said that discovery of a new target gene and development of a therapeutic agent for this are important inventions which directly lead to the possibility of therapy and expansion of the choices.

[0010] As the embodiment of molecular target drugs, various substances have been studied and developed such as a therapeutic antibody and a low molecular weight drug, as well as a peptide drug, a biological protein preparation such as cytokine, an siRNA, aptamer and the like. When an antibody is used as a therapeutic agent, due to its specificity, it is useful in treating pathological conditions in which the disordered cell expresses a specific antigen. The antibody binds to a protein expressing on the cell surface as its antigen and effectively acts upon the bound cell. The antibody has a characteristic of long blood half life and high specificity for its antigen and is also markedly useful as an antitumor agent. For example, when an antibody targets at a tumor-specific antigen, it can be expected that the administered antibody accumulates into the tumor and thereby attacks the tumor cell via complement-dependent cytotoxicity (CDC) and antibody-dependent cellular cytotoxicity (ADCC). In addition, by binding a radioactive substance, a cytotoxic substance and the like to an antibody, it becomes possible to transfer an agent efficiently to the tumor part and thereby to allow to act thereon. At the same time, it can decrease the amount of the reached agent to non-specific other tissues and reduction of side effects can also be expected. Inhibition of tumor growth or regression of tumor can be expected by administering an antibody having agonistic activity when a tumor-specific antigen has an activity to induce cell death, or by administering an antibody having neutralization activity when a tumor-specific antigen relates to in the growth and survival of cells. Due to the above characteristics, it is considered that antibodies are suited in applying as antitumor agents.

Regarding therapeutic antibodies:

[0011] In the original antibody preparation, a mouse was used as the animal to be immunized. However, use of mouse antibodies as drugs is limited due to a large number of reasons. A mouse antibody which can be recognized as a foreign substances in the human body can induce so-called "human anti-mouse antibody" namely "HAMA" response (Non-patent Document 6). Further, the Fc region of mouse antibody is not effective for the attack on disease cells via human complement or cellular toxicity.

[0012] As one of the approaches for avoiding such problems, a chimeric antibody has been developed (Patent Documents 1 and 2). The chimeric antibody contains parts of antibodies derived from two or more species (mouse antibody variable region, human antibody constant region and the like). An advantageous point of such a chimeric antibody is that it keeps the characteristics of mouse antibody but can stimulate human complement or cellular cytotoxicity since it has human Fc. However, it is known that such a chimeric antibody still induces "human anti-chimeric antibody" namely "HACA" response (Non-patent Document 7).

[0013] Further, it has been developed a recombinant antibody in which only a part of a substituted antibody is a complementarity determining region ("CDR") (Patent Documents 3 and 4). By the use of a CDR grafting technique, an antibody comprising mouse CDR and human variable region framework and human constant region, so-called "humanized antibody" (Non-patent Document 8). Further, by the use of a human antibody producing mouse or by a screening using a human antibody library, broadly utilized techniques have been provided also regarding preparation of complete human antibodies (Non-patent Documents 9 and 10).

Regarding IL-3R α :

[0014] IL-3R α is the α chain of IL-3 receptor, belongs to a cytokine receptor family and shows weak affinity for IL-3

as its ligand. By forming a hetero receptor with its β chain (CD131, hereinafter also referred to as IL-3R β), an IL-3 receptor has a strong binding and transfers a signal such as growth, differentiation and the like into a cell through intracellular region of the β chain. IL-5 receptor α chain and GM-CSF receptor α chain share the β chain in common.

[0015] IL-3R α is a type I membrane protein of single-pass transmembrane, and it is known based on the sequence that an IL-3 binding site and a fibronectin type III site are present in the extramembrane region. It is known that there is no structure which can transfer a signal in the intramembrane region. Though three-dimensional structure of IL-3R α has not been analyzed yet, it can be assumed that structures of cytokine receptors are similar between families since position of cysteine residue which forms the structurally important S-S bond is preserved in most cases. Among the same cytokine receptors, crystalline structures of IL-13 receptor α chain, IL-4 receptor α chain and GM-CSF receptor α chain have been analyzed. Based on the information of these cytokine receptor families, it can be assumed that the extramembrane region of IL-3R α is roughly divided into 3 domains (A-B-C domains). It is known that an antibody 7G3 which recognizes A domain of human IL-3R α blocks IL-3 signaling (Non-patent Document 11). In addition, expression of an A domain-deficient IL-3R α molecule has been reported (Non-patent Document 12), and as a matter of course, an antibody which recognizes A domain cannot recognize A domain-deficient IL-3R α . In addition, it is considered that C domain is the root of IL-3R α molecule and has a high possibility to three-dimensionally inhibit association of IL-3R β with IL-3R α .

[0016] IL-3 is the only a ligand which is known as a ligand of IL-3R α . IL-3 is a hematopoietic factor which is known to accelerate colony formation of the following: erythrocyte, megakaryocyte, neutrophil, eosinophil, basophil, mast cell and a monocyte system cell. It is known that IL-3 also stimulates a precursor cell having pluripotency, but IL-3 is rather said to accelerate a differentiation of not an immature stem cell having autonomous replication ability but a precursor cell committed to differentiation.

[0017] It is known that IL-3R α relates to the growth and differentiation of myeloid cells by forming a heterodimer with β chain and thereby transferring the IL-3 signaling into the cell via the Serine/Threonine phosphorylation pathway. It is known that IL-3R α is expressed in Granulocyte-Macrophage Progenitor (GMP) or Common Myeloid Progenitor (CMP) among hematopoietic precursor cells and induces growth and differentiation into neutrophil and macrophage systems via the IL-3 signaling. On the other hand, it has been reported that the Megakaryocyte Erythroid Progenitor (MEP) presenting in the downstream of CMP does not express IL-3R α different from the GMP which is also present in the downstream.

[0018] Regarding the AML stem cell, Bonnet and Dick have reported that the AML stem cell is present in the CD34 positive CD38 negative fraction (Non-patent reference 13). Further, by comparing with the same fraction (CD34 positive CD38 negative) of normal stem cell, Jordan *et al.* have found that IL-3R α is highly expressed in the AML stem cell (Non-patent reference 14). A high potential of IL-3R α as a marker of not only AML stem cell but also leukemia stem cell has also been reported in the plural of reports thereafter (Non-patent references 15 and 16). In the treatment of cancers including leukemia, it is important that only the cancer cells are removed without injuring normal cells as many as possible, and it is considered that this difference in the expression of IL-3R α between normal stem cell and leukemia stem cell is useful in the treatment targeting at the leukemia stem cell.

[0019] Regarding IL-3R β which forms a heterodimer with IL-3R α , there is no report that IL-3R β is highly expressed leukemia stem cell, and also in the case of a microarray in which expression of mRNA in leukemia stem cell and normal stem cell is compared in fact, IL-3R β is not identified as a molecule in which its expression is increased in leukemia stem cell (Non-patent reference 17).

[0020] Regarding IL-3R β which forms a heterodimer with IL-3R α , there is no report that IL-3R β is highly expressed leukemia stem cell, and also in the case of a microarray in which expression of mRNA in leukemia stem cell and normal stem cell is compared in fact, IL-3R β is not identified as a molecule of which expression is increased in leukemia stem cell (Non-patent reference 18).

[0021] The presence of a leukemia cell which depends on IL-3 has been known for a long time, and the old studies are studies focused on a blast cell which occupies most of the leukemia cells. According to the recent studies on leukemia stem cell, it is said that the leukemia stem cell acquires antitumor agent resistance by exhaustively suppressing its growth. In addition, it is considered that an IL-3 reactive blast cell has high proliferation ability so that it is assumed that such a cell is effective in the general treatment using an antitumor agent.

[0022] As a candidate of the agent targeting at an IL-3R receptor, the IL-3 itself was administered for a long time to patients of hematopoietic insufficiency but it did not become a drug as a result. A clinical trial for a fusion protein in which diphtheria toxin is added to IL-3 is in progress aiming leukemia as a target of the disease. Regarding the IL-3 and diphtheria toxin-IL-3 fusion, these are not suitable as the agents which are targeting at cells in which expression of IL-3R α is specifically increased, since IL-3 binds strongly not a protein of IL-3R α alone but a hetero protein of IL-3R α and β due to properties of IL-3. On the other hand, as a candidate of an agent targeting at IL-3R α , a first phase result of an IL-3R α human mouse chimeric antibody 7G3 has been reported (Non-patent Document 19). Since the 7G3 chimeric antibody uses for the purpose of blocking of IL-3 signaling as the mechanism of AML therapy, this is not an agent aimed at removing IL-3R α positive cells. Also, although some other IL-3R α antibodies are known (9F5 (Becton Dickinson), 6H6 (SANTA CRUZ BIOTECHNOLOGY) and AC 145 (Miltenyi-Biotech)), these do not have the ability to remove the

cells highly expressing IL-3R α .

CITATION LIST

5 Patent Document

[0023]

10 Patent Document 1 : EP Published Patent Application 120694
 Patent Document 2 : EP Published Patent Application No. 125023
 Patent Document 3 : GB Patent application No. GB2188638A
 Patent Document 4 : US Patent No. 5,585,089

15 Non-Patent Document

[0024]

20 Non-patent Document 1 : Osawa M et al., Science. 273:2 42-5 (1996)
 Non-patent Document 2 : Goodell MA et al., J Exp Med. 183: 1797-806 (1996)
 Non-patent Document 3 : Yamazaki S et al., EMBO J. 25: 3515-23 (2006)
 Non-patent Document 4 : Ishikawa F et al., Nat Biotechnol. 25:1315-21. (2007)
 Non-patent Document 5 : Bao S et al., Nature. 444: 756-60 (2006)
 Non-patent Document 6 : Schiff et al., Canc. Res., 45, 879-885 (1985)
 Non-patent Document 7 : Bruggemann et al., J. Exp. Med., 170:2153-2157 (1989)
 25 Non-patent Document 8 : Riechmann et al., Nature, 332:323-327 (1988)
 Non-patent Document 9 : Ishida I et al., Cloning Stem Cells. 4:91-102 (2002)
 Non-patent Document 10 : Wu et al., J Mol Biol. 19:151-62 (1999)
 Non-patent Document 11 : Sun et al., Blood, 87:83 (1996)
 Non-patent Document 12 : Chen et al., J Biol Chem, 284: 5763(2009)
 30 Non-patent Document 13 : Bonnet et al., Nat Med, 1997; 3: 730
 Non-patent Document 14 : Jordan et al., Leukemia, 2000; 14: 1777
 Non-patent Document 15 : Haematologica, 2001; 86:1261
 Non-patent Document 16 : LeukLymphoma, 2006; 47:207
 Non-patent Document 17 : Majeti et al., Proc Natl Acad Sci USA. 2009; 106:3396
 35 Non-patent Document 18 : Majeti et al., Proc Natl Acad Sci USA. 106:3396(2009)
 Non-patent Document 19 : Blood, 2008 112 (11): Abstract 2956

SUMMARY OF THE INVENTION

40 TECHNICAL PROBLEMS

[0025] An object of the invention is to provide a therapeutic agent which can remove leukemia stem cells alone and also can hardly exhibit adverse effects upon normal cells (shows fewer side effects). Specifically, the present invention provides an antibody to human IL-3R α chain, which does not inhibit IL-3 signaling and binds to B domain of human IL-3R α chain but does not bind to C domain; a composition comprising the antibody; and a therapeutic method or detection method comprising the antibody.

SOLUTION TO PROBLEMS

50 [0026] The invention relates to the following (1) to (9).

(1) An antibody to a human IL-3R α chain, which does not inhibit IL-3 signaling and binds to B domain of human IL-3R α chain but does not bind to C domain.
 (2) The antibody described in the above-mentioned (1), further having high antibody-dependent cellular cytotoxicity (ADCC).
 55 (3) The antibody described in the above-mentioned (1) or (2), wherein the high antibody-dependent cellular cytotoxicity (ADCC) shows a specific lysis rate of 10% at an antibody concentration of 0.01 μ g/ml, by a Colon-26/hCD123 ADCC assay method which uses PBMC cultured with IL-2.

(4) The antibody described in any one of the above-mentioned (1) to (3), which comprises amino acid sequences of CDRs of heavy chain and CDRs of light chain selected from the group consisting of the following (a) to (e);

- (a) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:113 to 115, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:131 to 133, respectively,
- (b) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:116 to 118, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs: 134 to 136, respectively,
- (c) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs: 119 to 121, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:137 to 139, respectively,
- (d) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs: 122 to 124, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:140 to 142, respectively, and
- (e) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs: 125 to 127, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs: 143 to 145, respectively.

(5) The antibody described in any one of the above-mentioned (1) to (4), which comprises the heavy chain variable region and light chain variable region selected from the group consisting of the following (a) to (f);

- (a) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:53 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:55;
- (b) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence represented by SEQ ID NO:57 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:59;
- (c) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence represented by SEQ ID NO:61 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:63;
- (d) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:65 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:67;
- (e) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 138 in the amino acid sequence represented by SEQ ID NO:69 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:71 and;
- (f) a heavy chain variable region and/or light chain variable region, which comprise amino acid sequences in which 1 to 3 amino acid residues are deleted, substituted, added or inserted in the heavy chain variable region and/or light chain variable region shown by the above (a) to (e).

(6) A composition for preventing or treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood of a subject, which comprises the IL-3R α antibody described in any one of (1) to (5) as an active ingredient.

(7) A method for treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood, which comprises administering, to a subject, a composition comprising the IL-3R α antibody described in any one of (1) to (5) as an active ingredient.

(8) A composition for detecting a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood of a biological sample from a subject, which comprises the IL-3R α antibody described in any one of (1) to (5).

(9) The composition or method described in any one of (1) to (5), wherein the aforementioned blood tumor is acute myeloid leukemia (AML).

ADVANTAGEOUS EFFECT OF THE INVENTION

[0027] The invention can provide an antibody to human IL-3R α chain, which does not inhibit IL-3 signaling and binds to B domain of human IL-3R α chain but does not bind to C domain; a composition which comprises said antibody and a therapeutic method or detection method using said antibody.

BRIEF DESCRIPTION OF THE DRAWINGS

[0028]

[Fig. 1] and [Fig. 2] Figs. 1 and 2 are results of a flow cytometry analysis of a cell expressing an IL-3R α /GM-CSFR α chimeric protein using a labeled anti-IL-3R α antibody.

[Fig. 3] Fig. 3 is a result of a flow cytometry analysis of a cell expressing an IL-3R α /GM-CSFR α chimeric protein using a labeled anti-IL-3R α antibody.

[Fig. 4] Fig. 4 is a graph in which, among the nucleotide and amino acid sequences of A and B domains of human IL-3R α molecule, parts of regions in which the regions 1 to 7 arranged on the outside of the molecule are substituted by the GM-CSFR α sequence are shown by dotted lines.

[Fig. 5] Fig. 5 is a result of a cell growth test for examining blocking activity of IL-3 signaling. The ordinate represents the cell growth inhibition ratio (%) and the abscissa represents various IL-3R α antibody names.

[Fig. 6] Fig. 6 is a result of a colony assay test for examining blocking activity of IL-3 signaling. GM, E, and GEMM show results using Granulocyte/Macrophage system, Erythroid system colony and mixed colonies, respectively.

[Fig. 7] Fig. 7 is a result of examining anti-tumor effects of various human antibodies in a tumor bearing model. The ordinate represents the number of MOLM13 cells, and the abscissa represents the various IL-3R α antibody names.

[Fig. 8] and [Fig. 9] Figs. 8 and 9 are results of ADCC test for IL-3R α expressing cell lines using anti-IL-3R α antibody. PBMC not cultured with IL-2 was used in Fig. 8, and PBMC cultured with IL-2 was used in Fig. 9.

[Fig. 10] Fig. 10 is a result of flow cytometry analysis of cells expressing a *Macaca fascicularis* IL-3R α by anti-IL-3R α antibody and a PE-labeled anti-human IgG secondary antibody. The upper column shows the cells expressing *Macaca fascicularis* IL-3R α , and the lower column shows the cells expressing human IL-3R α .

MODE FOR CARRYING OUT THE INVENTION

(Detailed description of specified desirable embodiments)

[0029] Headings of the sections to be used in this specification are only for the purpose of organization and should not be interpreted as limitation to the main subject to be described. All of the cited references cited in this application are clearly incorporated by references into this specification for optional purposes.

(Outline)

[0030] This invention relates to an antibody to human IL-3R α chain, which does not inhibit IL-3 signaling and binds to B domain of human IL-3R α chain (hereinafter referred to as IL-3R α) but does not bind to C domain.

[0031] IL-3 receptors (hereinafter referred to as IL-3R), particularly IL-3R α , are expressed on the cell surface of a leukemia stem cell. In general, IL-3 receptor β chain (hereinafter referred to as IL-3R β) transfers IL-3 signaling into the cell and therefore induces growth and differentiation.

[0032] Accordingly, there is a possibility that inhibition of IL-3 signaling cause side effects such as inhibition of normal hematopoietic action by a normal stem cell. Thus, as a new therapeutic method which targets at leukemia stem cell, it is preferable that the method targets at IL-3R α and also does not inhibit IL-3 signaling.

(IL-3R α)

[0033] IL-3R α gene is a type I transmembrane protein which belongs to a cytokine receptor family. In normal cells, the IL-3R α molecule is expressed on a part of hematopoietic precursor cells, basophil, a part of dendritic cells and the like. In the case of tumors, it is known to be expressed in a hematopoietic system cancer and leukemia. As examples of tumors which express IL-3R α , it is known that IL-3R α is expressed on the blast cell of AML and CML in blastic crisis phase, and in the case of a differentiation marker-negative CD34 positive CD38 negative fraction considered to be a leukemia stem cell, in AML, CML, MDS, ALL and SM. In blood, IL-3 which is a known ligand of IL-3R α is expressed on an activated T cell, a natural killer cell, a mast cell and a part of cells of megakaryocyte system. In addition, the IL-3R α is also called CD123. The IL-3R α includes a mammal (e.g., the primates and human) type IL-3R α . The IL-3R α sequence includes polymorphic variants. Specific examples of the full length human IL-3R α include the following amino acid sequences.

MVLLWLTL⁵LLIALPCLLQTKEDPNPPITNLRMKAKAQQLTWDLNRNVTDIECVK
 DADYSMPAVNNSYCQFGAISLCEVTNYTVRVANPPFSTWILFPENSGKPWAGAE
 NLTCWIHDVDFLSCSWAVGPGAPADVQYDLYLNVANRRQQYECLHYKTDAQG
 TRIGCRFDDISRLSSGSQSSHILVRGRSAAFGIPCTDKFVVFSQIEILTPPNMTAKC
 NKTHSFMHWKMRSHFNRKFRYELQIQKRMQPVITEQVRDRTSFQLLNPGTYTV
 QIRARERVYEFLSAWSTPQRFEC¹⁰DQEEGANTRAWRTSLLIALGTLLALVCVFVIC
 RRYLVMQRLFPRI¹⁵PHMKDPIGDSFQNDKLVVWEAGKAGLEECLVTEVQVVQKT
 (SEQ ID NO:1)

[0034] Specific examples of amino acid sequence of the extracellular region of human IL-3R α include the following amino acid sequence.

MVLLWLTL²⁰LLIALPCLLQTKEDPNPPITNLRMKAKAQQLTWDLNRNVTDIECVK
 DADYSMPAVNNSYCQFGAISLCEVTNYTVRVANPPFSTWILFPENSGKPWAGAE
 NLTCWIHDVDFLSCSWAVGPGAPADVQYDLYLNVANRRQQYECLHYKTDAQG
 TRIGCRFDDISRLSSGSQSSHILVRGRSAAFGIPCTDKFVVFSQIEILTPPNMTAKC
 NKTHSFMHWKMRSHFNRKFRYELQIQKRMQPVITEQVRDRTSFQLLNPGTYTV
 QIRARERVYEFLSAWSTPQRFEC²⁵DQEEGANTRAWRTSL (SEQ ID NO:2)

[0035] In addition, the extracellular region of IL-3R α is divided into three domains of AtoC.

[0036] A domain comprises a region from glutamine (Q) at position 18 to serine (S) at position 100 in the amino acids of SEQ ID NO:2, and B domain comprises from glycine (G) at position 101 to serine (S) at position 203 in the amino acids of SEQ ID NO:2 and C domain that from glutamine (Q) at position 204 to leucine (L) at position 308 in the amino acids of SEQ ID NO:2.

[0037] Further, in A domain and B domain, the following 7 regions are arranged on the outside of the molecule.

[0038] The region 1 is from aspartic acid (D) at position 55 to proline (P) at position 61 in the amino acids of SEQ ID NO:2, the region 2 is from valine (V) at position 63 to phenylalanine (F) at position 70 in the amino acids of SEQ ID NO:2, the region 3 is from serine (S) at position 91 to glutamic acid (E) at position 98 in the amino acids of SEQ ID NO:2, the region 4 is from proline (P) at position 97 to tryptophan (W) at position 104 in the amino acids of SEQ ID NO:2, the region 5 is from cysteine (C) at position 122 to proline (P) at position 128 in the amino acids of SEQ ID NO:2, the region 6 is from isoleucine (I) at position 182 to serine (S) at position 188 in the amino acids of SEQ ID NO:2 and the region 7 is from glycine (G) at position 192 to lysine (K) at position 198 in the amino acids of SEQ ID NO:2.

[0039] Accordingly, examples of the antibody of the invention include an antibody which binds to an amino acid sequence of positions 101 to 203 in the amino acids of SEQ ID NO:2 which is the extracellular region of IL-3R α , but does not bind to an amino acid sequence of positions 204 to 308, and an antibody which further binds to amino acid sequences of positions 182 to 188 and positions 192 to 198 in the amino acid sequence of SEQ ID NO:2.

[0040] The antibody of the invention binds to the above-mentioned specific regions of the extracellular region of IL-3R α and does not inhibit IL-3 signaling.

[0041] The term "does not inhibit IL-3 signaling" as used in the invention means that it does not inhibit the intracellular signal through IL-3R by IL-3, and it includes a case in which the association of IL-3 with IL-3R is not inhibited and the binding of IL-3R α chain and β chain is not inhibited. Specifically, it means that the cell growth inhibition ratio shown by Fig. 5 according to the analysis in Example 8 is 40% or more, preferably 60% or more, further preferably 80% or more, when the antibody concentration is set to 10 μ g/ml. According to this specification, the terms "blocking of IL-3 signaling" and "inhibition of IL-3 signaling" are used as the same meaning and not discriminated, and the blocking activity of IL-3 signaling means the ability to inhibit IL-3 signaling.

[0042] Also, the antibody of the invention has high antibody-dependent cellular cytotoxicity (ADCC) in addition to the above-mentioned properties.

[0043] The IL-3R α antibody having ADCC activity means an antibody which binds to a cell expressing IL-3R α to kill

the IL-3R α -expressing cell via an effector cell having cytotoxicity such as NK cell and the like.

[0044] The high ADCC activity means that the specific lysis rate is 10% or more at an antibody concentration of 0.01 μ g/ml or less when measured by a Colon-26/hCD123 ADCC assay method which uses PBMC cultured with IL-2.

[0045] The specific lysis rate means a value obtained by measuring the lysis rate of a target cell by an antibody and specifically it can be calculated in accordance with the following Example 11.

[0046] Examples of the cell expressing IL-3R α , include blood cancer cells (acute myeloid leukemia (AML) cells, chronic myeloid leukemia (CML) cells, myeloid splastic syndromes (MDS) cells, acute lymphoid leukemia (ALL) cells, chronic lymphoid leukemia (CLL) cells, multiple myeloma (multiple myeloma: MM) cells, systemic mastocytoma (SM) cells etc.), regulatory T cells (such as CD4-positive CD25-positive cell), antigen presenting cells (such as dendritic cells, monocytes, macrophages and similar cells thereto (hepatic stellate cells, osteoclasts, microglial cells, the major epidermal phagocytic cells, dust cells (alveolar macrophages), etc.)), basophils and the like.

[0047] In addition, AML cell, CML cell, ALL cell, CLL cell, MDS cell, SM cell, MM cell, various lymphoma cells include their cancer stem cells.

[0048] The cancer stem cell is one of the cell groups constituting tumor. For example, in acute myeloid leukemia (AML) it is represented by Lineage(-)CD34(+)CD38(-) myeloid cell. Accordingly, since the antibody of the invention has high ADCC activity, it induces reduction or elimination of cells expressing IL-3R α .

[0049] Also, the IL-3R α antibody of the invention includes an IL-3R α antibody which has CDRs of heavy chain and CDRs of light chain selected from the group consisting of the following (a) to (e);

- (a) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs:113 to 115, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:131 to 133, respectively,
- (b) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs:116 to 118, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:134 to 136, respectively,
- (c) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs:119 to 121, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:137 to 139, respectively,
- (d) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs:122 to 124, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:140 to 142, respectively, and
- (e) CDR 1 to 3 of heavy chain are the amino acid sequences represented by SEQ ID NOs:125 to 127, respectively, and CDR 1 to 3 of light chain are the amino acid sequences represented by SEQ ID NOs:143 to 145, respectively.

[0050] In addition, the antibody of the invention includes an IL-3R α antibody which comprises the heavy chain variable region and the light chain variable region selected from the group consisting of the following (a) to (f) (shown in parentheses are names of the antibodies which are described in the following Examples from which each of variable regions are derived);

- (a) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence represented by SEQ ID NO:53 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence represented by SEQ ID NO:55 (name of antibody: Old4)
- (b) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence represented by SEQ ID NO:57 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence represented by SEQ ID NO:59 (name of antibody: Old5)
- (c) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence represented by SEQ ID NO:61 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence represented by SEQ ID NO:63 (name of antibody: Old17)
- (d) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence represented by SEQ ID NO:65 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence represented by SEQ ID NO:67 (name of antibody: Old19)
- (e) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 138 in the amino acid sequence represented by SEQ ID NO:69 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence represented by SEQ ID NO:71 (name of antibody: New102) and
- (f) a heavy chain variable region and/or light chain variable region, which comprise amino acid sequences in which 1 to 3 amino acid residues are deleted, substituted, added or inserted in the heavy chain variable region and/or light

chain variable region shown by (a) to (e).

(Antibody)

[0051] The antibody is used in a most broad sense and includes a monoclonal antibody, a polyclonal antibody, a multivalent antibody, a multispecific antibody (e.g., bispecific antibody) and also antibody fragments as long as these exhibit the desired biological activity.

[0052] The antibody contains a mature heavy chain or light chain variable region sequence. In addition, the antibody also includes a modified form and variant form such as substitutions within or outside of a constant region, a complementary determining region (CDR) or a framework (FR) region antibody of a mature heavy or light chain variable region sequence of the antibody, and the like. In a specific embodiment, the substitution includes a conservative amino acid substitution is included in the substitution.

[0053] In addition, the antibody also includes a subsequence of the mature heavy chain or light chain variable region sequence. In a specific embodiment, the subsequence is selected from Fab, Fab', F(ab')₂, Fv, Fd, single chain Fv (scFv), disulfide bond Fv (sdFv) and VL or VH.

[0054] In addition, the antibody also includes a heterogeneous domain. In a specific embodiment, the heterogeneous domain includes a tag, a detectable label or a cytotoxic agent.

[0055] Examples of the antibody include a monoclonal antibody and a polyclonal antibody and any isotype or subclass thereof. In a specific embodiment, the aforementioned antibody is an isotype of IgG (e.g., IgG1, IgG2, IgG3 or IgG4), IgA, IgM, IgE or IgD. The "monoclonal" antibody means an antibody that is based upon, obtained from a single clone including a eukaryote clone, a prokaryote clone or a phage clone or derived from a single clone including a eukaryote clone, a prokaryote clone or a phage clone, based on a single clone including a eukaryote clone, a prokaryote clone or a phage clone. Accordingly, the "monoclonal" antibody is a structurally defined substance and not a method by which it is produced.

[0056] The IL-3R α antibody, anti-IL-3R α and anti-IL-3R α antibody mean an antibody which specifically binds to IL-3R α . The specific binding means that it is selective for the epitope presenting in IL-3R α . The specific binding can be distinguished from non-specific binding using a known assay in the technical field (e.g., immunoprecipitation, ELISA, flow cytometry, Western blotting).

[0057] When all or a part of antigen epitopes to which an IL-3R α antibody specifically binds are present in different proteins, there is a possibility that this antibody can bind to the different proteins. Therefore, there is a possibility that the IL-3R α antibody specifically binds to other protein having high sequence or structural homology to IL-3R α epitope depending on the sequence or structural homology to IL-3R α epitope. Accordingly, there is a possibility that IL-3R α antibody binds to a different protein when an epitope having sufficient sequence or structural homology is present in the different protein.

[0058] The IL-3R α antibody includes isolated and purified antibodies. The antibody of the invention including an isolated or purified IL-3R α antibody includes human.

[0059] The term "(be) isolated" to be used as a modifier of a composition means that the composition is prepared by the hand of man or separated from one or more other components in *in vivo* environment presenting in nature generally by one or more manipulative steps or processes. In general, a composition separated in this manner does not substantially contain one or more materials with which they normally associate in nature, such as one or more proteins, nucleic acids, lipids, carbohydrates and cell membranes. Because of this, the isolated composition is separated from other biological components in the cells of the organism in which the composition naturally occurs, or from the artificial medium in which it is produced (e.g., by synthesis or cell culture). For example, an isolated IL-3R α antibody can be obtained from an animal in which the antibody is produced (e.g., non-transgenic mammals or transgenic mammals (rodents (mouse) or the ungulates (cattle)) and is separated from other polypeptides and nucleic acids. Accordingly, it is considered that the serum containing an antibody obtained from such an animal is isolated. The term "(be) isolated" does not exclude alternative physical forms, and for example, an isolated antibody could include antibody subsequences and chimeras, multimers or derivatized forms.

[0060] The term "(be) purified" to be used as a modifier of a composition refers to a composition which is free of most of or substantially all of the materials with which it typically associates in nature. In general, a purified antibody is obtained from the components generally presenting in the antibody environment. Because of this, it is considered that an antibody supernatant which is separated from a cell culture mixture of an antibody producing hybridoma is purified. Accordingly, the "(be) purified" does not require absolute purity and is context specific. Furthermore, the "(be) purified" composition can be combined with one or more other molecules. Because of this, the term "(be) purified" does not exclude combination of composition. The purity can be determined by an optional appropriate method such as UV spectrometry, chromatography (e.g., HPLC, gas phase), gel electrophoresis (e.g., silver or Coomassie staining), sequence analysis (peptide and nucleic acid) and the like.

[0061] The "(be) purified" protein and nucleic acid include a protein and a nucleic acid which are obtained by a standard

purification method. Also, a protein and a nucleic acid obtained by recombination expression in a host cell and chemical synthesis are also included in this term. In addition, the "(be) purified" can also refer to a composition in which the level of contaminants is lower than the level which is acceptable to a regulatory agency for administration to human or non-human animals, such as the Food and Drug Administration (FDA).

[0062] The IL-3R α antibody includes an antibody which binds to IL-3R α and modulates function or activity of IL-3R α *in vivo* or *in vitro* (e.g., in a subject). In the specification, the "to modulate" and the grammatical variations thereof when used in relation to the activity or function of IL-3R α mean that the IL-3R α activity or function is detectably affected, modified or altered but does not include inhibition of IL-3 signaling. Accordingly, the IL-3R α antibody which modulates the activity or function of IL-3R α is an antibody that provides influence, modification or alteration such that one or more of the IL-3R α activity or function can be detected without inhibiting IL-3 signaling, and such an activity or function of IL-3R α can include, for example, binding of IL-3R α with an IL-3R α ligand (e.g., IL-3), an IL-3R α -mediated signal transfer or an IL-3R α -mediated cell response or a cell response that can be modulated by IL-3R α , or the activity or function of other IL-3R α described in the specification or, otherwise, is commonly known or can be known.

[0063] Examples of various non-limited IL-3R α activities and functions which can be modulated include IL-3R α mediated signal transduction or IL-3R α mediated cellular response, cellular response which can be modulated via IL-3R α , cell proliferation or cell expansion (e.g., AML cell, CML cell, ALL cell, CLL cell, MDS cell, MM cell, SM cell, various lymphoma cells, monocytes, macrophages, mast cells, basophils, helper T cells, regulatory T cells, natural killer cells, myeloid progenitor cells and lymphoid progenitor cells), cell survival or apoptosis (e.g., AML cell, CML cell, ALL cell, CLL cell, MDS cell, MM cell, SM cell, various lymphoma cells, monocytes, macrophages, mast cells, basophils, helper T cells, regulatory T cells, natural killer cells, myeloid progenitor cells and lymphoid progenitor cells), cytokines (e.g., Th1, Th2 and non-Th1/Th2 cytokines) and interferon expression or production, expression or production of anti-apoptosis protein or proapoptosis protein, treatment, suppression or improvement of disorder, disease, physiological condition, pathological condition and symptom. Specific cytokines to be modulated are not limited and examples include IL-1, IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-14, IL-16, IL-17, IL-23, IL-26, TNF- α , and interferon γ (*in vitro* or *in vivo*). Specific anti-apoptosis proteins and proapoptosis proteins are not limited and examples include Bcl-xL, Bcl-2, Bad, Bim, and Mcl-1.

[0064] Therefore, examples of anti-IL-3R α antibody described in the present specification include an antibody which modulates IL-3R α mediated signal transduction or IL-3R α mediated cellular response, cellular response which can be modulated via IL-3R α , cell proliferation or cell growth (e.g., AML cell, CML cell, ALL cell, CLL cell, MDS cell, MM cell, SM cell, various lymphoma cells, monocytes, macrophages, mast cells, basophils, helper T cells, regulatory T cells, natural killer cells, myeloid progenitor cells and lymphoid progenitor cells), cell survival or apoptosis (e.g., AML cell, CML cell, ALL cell, CLL cell, MDS cell, MM cell, SM cell, various lymphoma cells, monocytes, macrophages, mast cells, basophils, helper T cells, regulatory T cells, natural killer cells, myeloid progenitor cells and lymphoid progenitor cells), cytokines (e.g., Th1, Th2 and non-Th1/Th2 cytokines) and interferon expression or production, expression or production of anti-apoptosis protein or proapoptosis protein, treatment, suppression or improvement of disorder, disease, physiological condition, pathological condition and symptom. In the specific embodiments, anti-IL-3R α antibody of the present invention can modulate expansion or survival of AML cell, number of other blood cancer cell (e.g., CML cell, ALL cell, MDS cell, MM cell, SM cell or various lymphoma cell), growth or survival of non-cancer blood cell such as monocytes, macrophages, mast cells, basophils, helper T cells, regulatory T cells, natural killer cells, myeloid progenitor cells and lymphoid progenitor cells, and reduces, disappears or depletes AML cell, CML cell, ALL cell, CLL cell, MDS cell, MM cell, SM cell, or various lymphoma cells.

[0065] The IL-3R α antibody includes a modified form such as a substitution product (e.g., an amino acid substitution product) which is also called as "variant", an addition product, deletion product (e.g., a subsequence or fragment) and the like. Such modified antibody forms and variants retain at least partial function or activity of the IL-3R α antibody shown by the invention, such as binding with IL-3R α , or modulation of activity or function (e.g., IL-3R α signal transfer) of IL-3R α . Accordingly, the modified IL-3R α antibody can retain the ability to modulate, for example, at least partial of IL-3R α binding or one or more of the IL-3R α functions or activities (e.g., signal transfer, cell response and the like).

[0066] According to this specification, the term "to alter" ("to modify") and the grammatical variations thereof means that the composition derivates a reference composition. The modified proteins, nucleic acids and other compositions can have higher or lower activities than a reference unmodified protein, nucleic acid or other composition or can have a different function from a reference unmodified protein, nucleic acid or other composition.

[0067] Such an antibody containing an amino acid substitution can be encoded by nucleic acid. Accordingly, the present invention also provides a nucleotide sequence encoding an antibody containing an amino acid substitution.

[0068] The term "identity" or "identical" means that two or more referenced substances are the same. Accordingly, when two protein sequences (e.g., IL-3R α antibodies) are identical, they have the same amino acid sequences at least within the referenced regions or portion. The term "identical region" means an identical region of two or more referenced substances. Thus, when two protein sequences are identical over one or more sequence regions, they have identity within the regions. "Substantial identity" means that a molecule is structurally or functionally conserved such that the molecule has or is predicted to have at least partial function or activity of one or more of reference molecule functions

or activities or relevant/corresponding region or a portion of the reference molecule to which it shares identity. Thus, polypeptides having substantial identity (e.g., IL-3R α antibodies) have or are predicted to have at least a part of the activity or function as a referenced polypeptide (e.g., IL-3R α antibody). For example, in a specific embodiment, it is considered that an IL-3R α antibody having one or more modifications (e.g., deletion, substitution, addition or insertion of 1 to 3 amino acid residues) which retain at least partial activity or function of the unmodified IL-3R α antibody has substantial identity to the reference IL-3R α antibody.

[0069] Due to variations between structurally related protein and functionally related protein, the amount of sequence identity required to retain functions or activity on the protein, region and function or activity of the region. In the case of protein, an activity or function can be retained by the presence of merely 30% of amino acid sequence identity, but in general, higher identity of 50%, 60%, 75%, 85%, 90%, 95%, 96%, 97% or 98%, to the reference sequence is present. The extent of identity between two sequences can be verified using a computer program or mathematic algorithm conventionally known in the technical field. In such an algorithm which calculates ratio of sequence identity (homology), in general, sequence gaps and mismatches over the comparison region are accounted. For example, BLAST (e.g., BLAST 2.0) retrieval algorithm (e.g., see Altschul et al., J. Mol. Biol., 215: 403 (1990), publicly available through NCBI) has the following illustrative retrieval parameters: mismatch-2; gap start 5; gap elongation 2. In the polypeptide sequence comparison, the BLASTP algorithm is typically used in combination with a scoring matrix such as PAM 100, PAM 250, BLOSUM 62, BLOSUM50.FASTA (e.g., FASTA 2 and FASTA 3) and the like, and SSEARCH sequence comparison program is also used for determining the extent of identity (Pearson et al., Proc. Natl. Acad. Sci. USA, 85: 2444 (1988); Pearson, Methods Mol. Bio., 132: 185 (2000); and Smith et al., J. Mol. Biol., 147: 195 (1981)). A program has also been developed for determining protein structural similarity using topological mapping based on Delaunay (Bostick et al., Biochem. Biophys. Res. Commun., 304: 320 (2003)).

[0070] A "conservative substitution" is a substitution of one amino acid by a biologically, chemically or structurally similar residue. Biological similarity means that a biological activity such as IL-3R α binding activity is not destroyed by the substitution. Structural similarity means that amino acids have side chain with similar length (e.g., alanine, glycine and serine) or have similar size. Chemical similarity means that the residues have the same charge or are hydrophilic or hydrophobic. Specific examples include substitution of one hydrophobic residue such as isoleucine, valine, leucine, and methionine with other residue, or the substitution of one polar residue with other residue such as the substitution of arginine with lysine, the substitution of glutamic acid with aspartic acid, or the substitution of glutamine with asparagine, and the substitution of serine with threonine.

[0071] In addition, examples of the modified antibody include peptide mimetics having one or more D-amino acids substituted with L-amino acids (and a mixture thereof), structural and functional analogs such as synthesized or non-natural amino acids or amino acid analogs, and derivatized form thereof. Examples of modification include a cyclic structure such as an end-to-end amide bond between the amino and carboxy-terminus of the molecule or intra- or inter-molecular disulfide bond or intramolecular or intermolecular disulfide bond.

[0072] Additional non-limiting specific examples of the amino acid modifications include partial sequence (subsequence) and fragment of IL-3R α . Exemplary subsequence and fragment of IL-3R α include a part of the IL-3R α sequence to which the exemplary IL-3R α antibody of the invention binds. Also, the exemplary subsequence and fragment of IL-3R α include an immunogenicity region such as a part of the IL-3R α to which the exemplary IL-3R α antibody of the invention binds.

[0073] According to the invention, there is provided a nucleic acid encoding an IL-3R α antibody subsequence of fragment which retains at least a part of the function or activity of the IL-3R α antibody and an unmodified or reference IL-3R α antibody. In this specification, the term "subsequence" or "fragment" means a portion of a full length molecule. The subsequence of the IL-3R α antibody encoding IL-3R α antibody has amino acids of smaller than those of the full length IL-3R α antibody by at least one (e.g., deletion of one or more inner or terminal amino acids from the amino terminus or carboxy terminus). The subsequence of IL-3R α antibody has amino acids of smaller than those of the full length IL-3R α antibody by at least one. The nucleic acid subsequence has nucleotides of smaller than those of the full length comparative nucleic acid sequence by at least one. Accordingly, the subsequence can be an optional length within the full length of native IL-3R α .

[0074] The IL-3R α antibody subsequence and fragment can have a binding affinity as the full length antibody, a binding specificity as the full length antibody or one or more activities or functions as the full length antibody, such as the function or activity of an IL-3R α antagonist or agonist antibody. The terms "functional subsequence" and "functional fragment" in the case of referring to the antibody mean an antibody portion which retains one or more functions or activities as the full length reference antibody, such as at least a part of the function or activity of IL-3R α antibody. For example, an antibody subsequence which binds to IL-3R α or a fragment of IL-3R α is considered a functional subsequence.

[0075] The antibody subsequence and fragment can be combined. For example, a VL or VH subsequence can be connected by a linker sequence and thereby can form a VL-VH chimeric body. A combination of single chain Fv(scFv) subsequences can be connected by a linker sequence and thereby can form a scFv-scFv-chimeric body. The IL-3R α antibody subsequence and fragment include a single chain antibody or variable region alone or in combination with all

or a portion of other IL-3R α antibody subsequence.

[0076] The antibody subsequence and fragment can be prepared by hydrolysis of the antibody by its proteolysis for example by a pepsin or papain digestion of the whole antibody. The antibody subsequence and fragment obtained by enzymatic cleavage with pepsin provide a 5S fragment represented by F(ab')₂. This fragment can be further cleaved using a thiol reducing agent to form a 3.5S Fab' monovalent fragment. Alternatively, an enzymatic cleavage using pepsin directly produces two monovalent Fab' fragments and Fc fragment (see e.g., US Patent No. 4,036,945 and US Patent No. 4,331,647; and Edelman et al., *Methods Enzymol.*, 1: 422 (1967)). Other methods of cleaving an antibody, such as separation of heavy chain for forming a monovalent light chain-heavy chain fragment, further cleavage of the fragment or other enzymatic or chemical method may be used.

[0077] A protein and an antibody, as well as subsequence thereof and fragment can be prepared using a genetic engineering. The technology includes the full or partial gene encoding a protein or an antibody is expressed in a host cell such as a COS cell and *E. Coli*. A recombinant host cell synthesizes the full or subsequence such as scFv (such as Whitlow et al, In: *Methods: A Companion to Methods in Enzymology* 2:97 (1991), Bird et al, *Science* 242:423 (1988); and US Patent No.4,946,778). A single chain Fv and an antibody can be prepared in accordance with the procedure as described in US Patent No. 4,946,778 and US Patent No. 5,258,498; Huston et al, *Methods Enzymol* 203:46 (1991); Shu et al, *Proc. Natl. Acad. Sci. USA* 90:7995 (1993); and Skerra et al, *Science* 240:1038 (1988).

[0078] The modified form includes a derivatized sequence such as amino acids in which the free amino groups form amine hydrochloride, p-toluenesulfonyl group and carbobenzoquinone group; the free carboxy groups which form a salt or methyl and ethyl ester; and the free hydroxyl groups form an O-acyl or O-alkyl derivative, and naturally existing amino acid derivatives such as 4-hydroxyproline (derivative of proline), 5-hydroxylysine (derivative of lysine), homoserine (derivative of serine), ornithine (derivative of lysine) and the like. The modification can be carried out using a method conventionally known in the technical field (e.g., site-specific deletion or insertion mutagenesis based on PCR, chemical modification and mutagenesis, crosslinking and the like).

[0079] Addition products and insertion products are included in the modified forms of protein (e.g., antibody), nucleic acid and other compositions. For example, the addition can be a covalent or non-covalent bond with any type of molecules of protein (e.g., antibody), nucleic acid or other compositions. In general, addition and insertion confer different function or activity.

[0080] Fusion (chimeric) polypeptides or nucleic acid sequences are included in the addition product and insertion product, and these are sequences having one or more molecules which are generally not present in the reference native (wild type) sequence covalently attached to the aforementioned sequence. A specific example is an amino acid sequence of other protein (e.g., an antibody) for producing a multifunctional protein (e.g., a multispecific antibody).

[0081] Also, the antibody of the invention include a chimeric or fusion product in which one or more additional domains are covalently linked thereto in order to confer a different or complementary function or activity. Examples of the antibody include a chimeric or fusion product which does not naturally present in natural and in which two or more amino acid sequences are mutually bonded.

[0082] According to the invention, there are provided an IL-3R α antibody which contains a heterologous domain and a nucleic acid that encodes the IL-3R α antibody. The heterologous domain can be an amino acid addition product or insertion product, but does not limited to an amino acid residue. Accordingly, the heterologous domain can be composed of any one of various different types of small or large functional parts. Such a part includes a nucleic acid, a peptide, a carbohydrate, a lipid or small organic compound such as a drug, a metal (gold, silver) and the like.

[0083] Non-limiting specific examples of the heterologous domain include a tag, a detectable label and a cytotoxic agent. Specific examples of the tag and detectable label include T7-, His-, myc-, HA- and FLAG-tags; enzymes (horse-radish peroxidase, urease, catalase, alkaline phosphatase, β -galactosidase, chloramphenicol transferase); enzyme substrates; ligands (e.g., biotin); receptors (avidin); radionuclide (e.g., C14, S35, P32, P33, H3, I125 and I131); electron density reagents; energy transfer molecules; paramagnetic labels; fluorophore (fluorescein, Rhodamine, Phycoerythrin); chromophore; chemiluminescence agents (imidazole, luciferase) and bioluminescence agents. Specific examples of the cytotoxic agent include diphtheria toxin (diphtheria, toxin), cholera toxin and lysine.

[0084] A linker sequence may be inserted between the protein (e.g., an antibody), nucleic acid or other composition and the addition product or insertion product (e.g., a heterologous domain) so that the two substances maintain at least a part of different function or activity. The linker sequence may have one or more properties which can accelerate either of the domains or can carry out mutual reaction with either of the domains, and such characteristics include impossibility to form a flexible structure and an ordered secondary structure or hydrophobic property or charging property. Examples of the amino acids which are generally found in the flexible protein regions include glycine, asparagine and serine. Other amino acids close to neutral such as threonine and alanine may also be used in the linker sequence. The length of the linker sequence can be varied (e.g., see US Patent No. 6,087,329). The linker further include chemical crosslinking agents and binding agents (conjugating agents) such as a sulfo-succinimidyl derivative (sulfo-SMCC, sulfo-SMPB), disuccinimidyl suberate (DSS), disuccinimidyl glutarate (DSG) and disuccinimidyl tartarate (DST).

[0085] Further examples of the addition include any one of glycosylation, fatty acid, lipid, acetylation, phosphorylation,

amidation, formylation, ubiquitination and derivatization by a protecting or blocking group and a large number of chemical modifications. Other substitutions and possibilities can be easily understood by those skilled in the art and are considered to be within the scope of the invention.

[0086] Such a modified sequence can be prepared using recombinant DNA techniques which mediate cell expression *or in vitro* translation. Polypeptides and nucleic acid sequences can also be prepared by a conventionally known method in the technical field such as chemical synthesis using an automatic peptide synthesizer (see e.g., Applied Biosystems, Foster City CA).

[0087] Modified and variant antibodies such as substitution products, subsequences addition products and the like can maintain detectable activity of IL-3R α antibody. In an embodiment, the modified antibody has the activity to bind to IL-3R α molecule and induces reduction or elimination of IL-3R α expression cells by an immune system mainly centering on an effector cell. The modified antibody relates to the functional control of IL-3R α expression cells and induces survival, growth, resting, cell death and the like of the cells. The cell death includes apoptosis, necrosis, autophagy and the like.

(Screening method of IL-3R α)

[0088] According to the invention, there are further provided a cell-free method and a cell-based method (e.g., *in vivo* or *in vitro*) which screen, detect and identify IL-3R α (e.g., in a solution or by a solid phase). These methods can be carried out in a solution *in vitro* using a biomaterial or sample, and *in vivo* for example using a sample of an animal-derived cell (e.g., lymphocyte). In an embodiment, the method comprises a step of contacting a biomaterial or sample with an antibody bound to IL-3R α under a condition of allowing binding of the antibody with IL-3R α and a step of assaying for the antibody bound to IL-3R α . The presence of IL-3R α is detected by binding of the antibody to bind to IL-3R α . In an embodiment, IL-3R α is present in a cell or tissue. In another embodiment, the aforementioned biomaterial or sample is obtained from a mammal analyte.

[0089] The term "contacting" when it is used in relation to the composition such a protein (e.g., IL-3R α antibody), a material, a sample or treatment means a direct or indirect interaction between the composition (e.g., IL-3R α antibody) and other referenced substance. Specific examples of the direct interaction include bonding. Specific examples of the indirect interaction include a case in which the composition acts upon an intermediate molecule and this intermediate molecule then acts upon the referenced substance. Accordingly, for example, contacting a cell (e.g., lymphocyte) to IL-3R α antibody includes to allow the antibody to bind to the cell (e.g., through binding to IL-3R α) or to allow the antibody to act on an intermediate substance, followed by the action of this intermediate substance upon the cell.

[0090] The terms "assaying" and "measuring" and grammatical variations thereof are synonymously used in the specification and mean either of qualitative measurement and quantitative measurement or both of qualitative measurement and quantitative measurement. When these terms are used in relation to binding, they include any means of evaluating relative amount, affinity or specificity of binding including various methods which are described in the specification and conventionally known in the technical field. For example, binding of the IL-3R α antibody with IL-3R α can be assayed or measured by a flow cytometry assay.

(Production of antibody)

[0091] The invention also provides a method for producing a human IL-3R α antibody having cytotoxicity for IL-3R α positive cells. In an embodiment, the method comprises administering a human IL-3R α extracellular region conjugated with a human IL-3R α recombinant protein or an IL-3R α gene introduced cell into animals capable of expressing human immunoglobulin (e.g., transgenic mice or transgenic cattle); screening the animal for expression of a human IL-3R α antibody; selecting the animal producing the human IL-3R α antibody; and isolating the antibody from the selected animal.

[0092] The IL-3R α protein suitable for the antibody production can be produced by any one of various standard protein purification and recombinant expression techniques. For example, the IL-3R α sequence can be prepared by standard peptide synthesis techniques such as a solid phase synthesis. In order to facilitate purification of the expressed or synthesized protein, a portion of the protein may contain an amino acid sequence such as a FLAG tag, a T7 tag, a polyhistidine sequence or the like. The protein is expressed inside the cells and can be purified. The protein can be expressed by a recombination method as a part of a further large protein (e.g., a fusion or chimeric product). The embodiment of the IL-3R α suitable for generating immune response includes IL-3R α subsequences such as an immunogenicity fragment. Further embodiment of IL-3R α includes an IL-3R α expressing cell, an IL-3R α containing preparation or cell extract or fraction and a partially purified IL-3R α .

[0093] The method for preparing polyclonal antibody and monoclonal antibody is conventionally known in the technical field. For example, IL-3R α or its immunogenicity fragment used for immunizing an animal by optionally conjugating with a carrier such as keyhole limpet hemocyanin (KLH) or ovalbumin (e.g., BSA) or mixing with an adjuvant such as complete Freund's adjuvant or incomplete Freund's adjuvant. By isolating a spleen cell derived from an immunized animal which responds to IL-3R α , it can be fused with myeloma cell using hybridoma techniques. The monoclonal antibodies produced

by hybridomas can be screened for reactivity with IL-3R α or immunogenicity fragment thereof.

[0094] The animal which can be immunized includes the primates, mouse, rat, rabbit, goat, sheep, cattle and guinea pig. The initial and any optionally subsequent immunization may be by intravenous route, intraperitoneal route, intramuscular route or subcutaneous route. Further, in order to increase immune response, the antigen can be conjugated with other protein such as keyhole limpet hemocyanin (KLH), thyroglobulin and tetanus toxoid, or can be mixed with an adjuvant such as complete Freund's adjuvant, incomplete Freund's adjuvant and the like. The initial and any optionally subsequent immunization may be through intraperitoneal route, intramuscular route, intraocular route or subcutaneous route. The immunization may be at the same concentration or different concentration of an IL-3R α preparation or at regular or irregular intervals.

[0095] The animal includes those which are genetically modified to include human gene loci, and a human antibody can be prepared using the same. Examples of the transgenic animals with one or more human immunoglobulin genes, are described for example in US Patent No. 5,939,598, WO02/43478 and WO02/092812. Using conventional hybridoma technique, an spleen cells which are isolated from immunized mouse having high responders to the antigen and are fused with myeloma cell. A monoclonal antibody which binds to IL-3R α can be obtained.

[0096] The method for producing a human polyclonal antibody and a human monoclonal antibody is further described (see, such as Kuroiwa et al, Nat. Biotechnol. 20:889 (2002); WO98/24893; WO92/01047; WO96/34096; WO96/33735; US Patent No. 5,413,923; US Patent No. 5,625,126; US Patent No. 5,633,425; US Patent No. 5,569,825; US Patent No. 5,661,016; US Patent No. 5,545,806; US Patent No. 5,814,318; US Patent No. 5,885,793; US Patent No. 5,916,771; and US Patent No.5,939,598).

[0097] The term "human" when it is used in reference to an antibody means that amino acid sequence of the antibody is completely the human amino acid sequence, namely is human heavy chain and human light chain variable regions and human constant region. Accordingly, all of the amino acids are human amino acids or present in the human antibody. An antibody which is a non-human antibody can be made into a complete human antibody by substituting the non-human amino acid residues with the amino acid residues which are present in the human antibody. The amino acid residues which are present in the human antibody, CDR region map and human antibody consensus residues are well known in the technical field (see e.g., Kabat, Sequences of Proteins of Immunological Interest, 4th edition, US Department of Health and Human Services, Public Health Service (1987); Chothia and Lesk (1987)). A consensus sequence of human VH subgroup III based on the investigation carried out using 22 known human VH III sequences as the object and a consensus sequence of human VL κ chain subgroup I based on the investigation carried out using 30 known human κ chain I sequences as the object are described in Padlan, Mol. Immunol., 31: 169 (1994) and Padlan, Mol. Immunol., 28: 489 (1991). Accordingly, the human antibody includes an antibody in which one or more amino acid residues have been substituted with one or more amino acids existing in an optional other human antibody.

[0098] Examples of the anti-IL-3R α antibody include antibodies prepared using a known method in the technical field, such as CDR-grafting (EP 239,400; WO91/09967; US Patent No. 5,225,539; US Patent No.5,530,101; and US Patent No.5,585,089), veneering, resurfacing (EP592,106; EP519,596; Padlan, Molecular Immunol. 28: 489 (1991); Studnicka et al., Protein Engineering 7: 805 (1994); Roguska et al., Proc. Nat'l Acad. Sci. USA 91: 969 (1994)) and chain shuffling (US Patent No.5,565,332). In order to produce a humanized antibody, human consensus sequence (Padlan, Mol. Immunol. 31:169 (1994); and Padlan, Mol. Immunol. 28: 489 (1991)) has been used (Carter et al., Proc. Natl. Acad. Sci. USA 89: 4285 (1992); and Presta et al, J. Immunol. 151:2623(1993)).

[0099] The term "humanized" when it is used in relation to an antibody means that amino acid sequence of the antibody has one or more non-human amino acid residues (e.g., mouse, rat, goat, rabbit and the like) of complement determining region (CDR) which specifically binds to a desired antigen in an acceptor human immunoglobulin molecule and one or more human amino acid residues (amino acid residues which are flanked with CDR) in Fv framework region (FR). The antibody called "primatized" is within the scope of meaning of "humanized", except that amino acid residues of the acceptor human immunoglobulin molecule and framework region can be any primate amino acid residues (e.g., monkey, gibbon, gorilla, chimpanzee, orangutan, macaque monkey) in addition to any human residues. Human FR residues of immunoglobulin can be substituted with corresponding non-human residues. Accordingly, for example, in order to alter, generally to improve, antigen affinity or specificity, residues in the CDR or human framework region can be substituted with corresponding residues from the non-human CDR or framework region donor antibody. The humanized antibody can contain residues which cannot be found in the human antibody and donor CDR or framework sequence. For example, it can be predicted that FR substitution at a particular position which cannot be found in human antibody or donor non-human antibody can improve binding affinity or specific human antibody at this position. Antibody framework and CDR substitutions based on the molecular modeling are conventionally known in the technical field, for example by the modeling of interaction of CDR and framework residues to identify framework residues important for antigen binding and the sequence comparison for identifying unusual framework residues at the specific position (see e.g., US Patent No. 5,585,089; and Riechmann et al., Nature, 332:323 (1988)).

[0100] Chimeric antibodies are included in the IL-3R α antibody. According to this specification, the term "chimeric" and the grammatical variations thereof when it is used in relation to antibodies mean that amino acid sequence of the

antibody contains one or more portion which is derived from two or more different species, is obtained or isolated from two or more different species or is based on two or more different species. For example, a portion of the antibody can be human (e.g., constant region) and other portion of the antibody can be non-human (e.g., a mouse heavy chain or a mouse light variable region). Accordingly, an example of the chimeric antibody includes an antibody in which the different portion of the antibody is derived from a different species. Different from the humanized or primatized antibody, the chimeric antibody can have a sequence of different species in an arbitrary region of the antibody.

[0101] The method for producing a chimeric antibody is known in the technical field (such as Morrison, Science 229: 1202 (1985); Oi et al., BioTechniques 4: 214 (1986); Gillies et al., J. Immunol. Methods 125: 191 (1989); US Patent No. 5,807,715; US Patent No. 4,816,567; and US Patent No. 4,816,397). For example, in Munro, Nature 312: 597 (1984); Neuberger et al., Nature 312: 604 (1984); Sharon et al., Nature 309: 364 (1984); Morrison et al., Proc. Nat'l. Acad. Sci. USA 81: 6851 (1984); Boulianne et al., Nature 312: 643 (1984); Capon et al., Nature 337: 525 (1989); and Traunecker et al., Nature 339: 68 (1989), a chimeric antibody in which a variable region of antibody derived from one species is replaced by a variable region of antibody derived from another species.

[0102] In addition, the anti-IL-3R α antibody can be prepared by hybridoma technique, recombinant technique, and phage display technique, and a combination thereof (see US Patent No. 4,902,614, US Patent No. 4,543,439, and US Patent No. 4,411,993; and also see Monoclonal Antibodies. Hybridomas: A New Dimension in Biological Analyses, Plenum Press, Kennett, McKearn, and Bechtol et al., 1980, and Harlow et al., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, the second edition, 1988).

[0103] The human anti-human IL-3R α antibody of the invention was produced using chromosome-transferred mice (KM mice (trademark)) immunized with various forms of soluble form of recombinant human IL-3R α proteins or cell lines expressing IL-3R α (WO02/43478, WO02/092812, and Ishida et al., IBC's 11th Antibody Engineering Meeting, Abstract (2000)). Since the human anti-human IL-3R α antibody detectably stains not a non-transformed parent cell line but a human IL-3R α stable transfectant cell line, such as Jurkat-IL-3R α cell and L929-IL-3R α cell, the antibody specifically is shown to bind to human IL-3R α .

[0104] The antibody of the invention can have κ light chain sequence or λ light chain sequence, full length of either one of them as present in naturally existing antibody, a mixture thereof (namely a fusion product of κ chain sequence and λ chain sequence) and subsequences/fragments thereof. The naturally presenting antibody molecules contain two κ light chains or two λ light chains.

[0105] The invention provides a method for preparing an antibody which specifically binds to IL-3R α . In a specific embodiment, the method for preparing IL-3R α antibody comprises administering human IL-3R α , a subsequence thereof or a fragment thereof (e.g., IL-3R α extracellular region), conjugated with a human Fc recombinant protein if necessary, to animals which can express human immunoglobulin (e.g., transgenic mice or transgenic cattle), screening the animals for their expression of human IL-3R α antibody, selecting an animal which produces human IL-3R α antibody and isolating the antibody from the selected animal. In an embodiment, whether or not the human IL-3R α antibody has an IL-3R α antagonist or agonist activity is judged by this method.

[0106] The effector activity means an antibody-dependent activity induced via Fc region of antibody, and such as antibody-dependent cellular cytotoxicity (ADCC activity), complement-dependent cytotoxicity (CDC activity), antibody-dependent phagocytosis (ADP activity) by phagocytes such as macrophage and dendritic cell, and the like, are known.

[0107] As a method for controlling effector activity of the anti-IL-3R α monoclonal antibody of the invention, examples include a method which controls the amount of the fucose (also called core fucose) which is bound to N-acetylglucosamine (GlcNAc) through α -1,6 bond in a reducing end of a complex-type N-linked sugar chain which is bound to asparagine (Asn) at position 297 of an Fc region of an antibody (WO2005/035586, WO2002/31140, and WO00/61739), a method in which is controlled by modifying amino acid residues of Fc region of the antibody, and the like. The effector activity can be controlled by applying any one of these methods to the anti-IL-3R α monoclonal antibody of the invention.

[0108] By controlling the content of the core fucose of complex-type N-linked sugar chain of Fc of the antibody, effector activity of the antibody can be increased or decreased. As a method for reducing the content of the fucose which binds to the complex-type N-linked sugar chain which is bound to Fc of the antibody, defucosylation (defucosylated or non-fucosylated) can be mentioned. The defucosylation is to express an antibody using CHO cell from which α 1,6-fucosyltransferase gene is deleted, and an antibody to which fucose is not bound can be obtained. The antibody to which fucose is not bound has high ADCC activity. On the other hand, as a method for increasing the content of the fucose which binds to the complex-type N-linked sugar chain to which Fc of the antibody is bound, the antibody to which fucose is bound can be obtained by expressing the antibody using a host cell in which α 1,6-fucosyltransferase gene is introduced. The antibody to which fucose is bound has the ADCC activity lower than that of the antibody to which fucose is not bound.

[0109] In addition, ADCC activity and CDC activity can be increased or decreased by modifying amino acid residues of the Fc region of the antibody. For example, CDC activity of the antibody can be increased by using the amino acid sequence of the Fc region described in US 2007/0148165. Also, ADCC activity or CDC activity can be increased or decreased by carrying out the amino acid modification described in US 6,737,056, US 7,297,775 and US 7,317,091. Further, an antibody in which effector activity of the antibody is controlled can be obtained by using the above-mentioned

methods in combination in one antibody.

[0110] According to the invention, the nucleotide sequence of the invention such as of a vector and the like is further provided. In an embodiment, the vector comprises a nucleic acid sequence encoding an IL-3R α antibody or a subsequence or fragment thereof.

[0111] The nucleic acid can have various lengths. The length of the nucleic acid encoding the IL-3R α antibody of the present invention or the subsequence thereof is generally about 100 to 600 nucleotides, or any numerical value or range within encompassing such lengths the above described range; 100 to 150, 150 to 200, 200 to 250, 250 to 300, 300 to 350, 350 to 400, 400 to 450, 450 to 500, 500 to 550 or 550 to 600 nucleotide length, or any numerical value or range or value within or encompassing such length the above described range. Examples of the length of nucleic acid encoding the IL-3R α antibody of the present invention or the subsequence thereof include generally 10 to 20, 20 to 30, 30 to 50, 50 to 100, 100 to 150, 150 to 200, 200 to 250, 250 to 300, 300 to 400, 400 to 500, 500 to 600 nucleotides and any numerical value or range within or encompassing such length.

[0112] The terms "nucleic acid" and "polynucleotide" means at least two or more ribo- or deoxy-ribo nucleic acid base pairs (nucleotide) linked which are through a phosphoester bond or equivalent. The nucleic acid includes polynucleotide and polynucleoside. The nucleic acid includes a single molecule, a double molecule, a triple molecule, a circular molecule or a linear molecule. Examples of the nucleic acid include RNA, DNA, cDNA, a genomic nucleic acid, a naturally existing nucleic acid and a non-natural nucleic acid such as a synthetic nucleic acid, but are not limited. Short nucleic acids and polynucleotides (e.g., 10 to 20, 20 to 30, 30 to 50, 50 to 100 nucleotides) are commonly called "oligonucleotides" or "probes" of single-stranded or double-stranded DNA.

[0113] Nucleic acid can be prepared using various standard cloning techniques and chemical synthesis techniques. Examples of the techniques include but are not limited to, nucleic acid amplification such as polymerase chain reaction (PCR), with genomic DNA or cDNA targets using primers (e.g., a degenerate primer mixture) which can be annealed with an antibody encoding sequence. In addition, nucleic acid can also be prepared by chemical synthesis (e.g., solid phase phosphoramidite synthesis) or transcription from a gene. Thereafter, the prepared sequence can be expressed by a cell (e.g., a host cell such as yeast, bacteria or eukaryote (an animal or mammal cell or in a plant)) after the sequence cloned into a plasmid and then amplified, or the sequence is translated *in vitro*.

[0114] A vector is a vehicle which can be manipulated by insertion or incorporation of nucleic acid. Examples of the vector include a plasmid vector, a virus vector, a prokaryote (bacterium) vector and a eukaryote (plant, fungi, mammals) vector. The vector can be used for *in vitro* or *in vivo* expression of nucleic acid. Such a vector is called "expression vector" and is useful for the transfer of nucleic acid including a nucleic acid which encodes an IL-3R α antibody or its subsequence or fragment and the expression of an encoded protein by *in vitro* (e.g., in a solution or on solid phase), by a cell or by *in vivo* in a subject.

[0115] In addition, the vector can also be used for manipulation of nucleic acids. For genetic manipulation, an inserted nucleic acid can be transcribed or translated using a "cloning vector" *in vitro* (e.g., in a solution or on solid phase), in a cell or *in vivo* in a subject.

[0116] In general, the vector contains an origin of replication for amplification in a cell *in vitro* or *in vivo*. Control elements such as an expression control element present in the vector can be included in order to facilitate transcription and translation, if necessary.

[0117] A vector can include a selection marker. The "selection marker" is a gene which allows for the selection of a cell containing the gene. "Positive selection" means a process for selecting a cell containing the selection marker due to a positive selection. Drug resistance is an example of the positive selection marker, and a cell containing the marker will survive in culture medium containing the drug and a cell which does not contain the marker will die. Examples of the selection marker include drug resistance genes such as *neo* which provides resistance to G418; *hygr* which provides resistance to hygromycin; *puro* which provides resistance to puromycin, and the like. Other positive selection marker includes genes which enable identification or screening of a cell containing the marker. Examples of these genes include a fluorescent protein (GFP and GFP-like chromophore, luciferase) gene, *lacZ* gene, alkaline phosphatase gene, and a surface marker such as CD8. "Negative selection" means a process for killing cells which contain negative selection markers by exposing to an appropriate negative selection agent. For example, a cell containing a herpes simplex virus thymidine kinase (HSV-tk) gene (Wigler et al., Cell, 11: 223 (1977)) is sensitive to a drug ganciclovir (GANC). Similarly, *gpt* gene makes a cell sensitive to 6-thioxanthine.

[0118] The virus vector includes those which are based on retroviral (a lentivirus for infecting not only dividing cells but also non-dividing cells), foamy virus (US Patent No. 5,624,820, US Patent No. 5,693,508, US Patent No. 5,665,577, US Patent No. 6,013,516 and US Patent No. 5,674,703; WO 92/05266 and WO 92/14829), adenovirus (US Patent No. 5,700,470, US Patent No. 5,731,172 and US Patent No. 5,928,944), adeno-associated virus (AAV) (US Patent No. 5,604,090), a herpes simplex virus vector (US Patent No. 5,501,979), a cytomegalovirus (CMV) system vector (US Patent No. 5,561,063), reovirus, rotavirus genome, simian virus 40 (SV40) or papilloma virus (Cone et al., Proc. Natl. Acad. Sci. USA, 81:6349 (1984); Eukaryotic Viral Vectors, Cold Spring Harbor Laboratory, edited by Gluzman, 1982; Sarver et al., Mol. Cell. Biol., 1: 486 (1981); US Patent No. 5,719,054). Adenovirus efficiently infects a slowly replicating

and/or terminally differentiated cell, and can be used to target the slowly replicating cell and/or terminally differentiated cell. Additional examples of virus vectors useful for expression include parbovirus, Norwalk virus, corona virus, paramyxovirus and rhabdo virus, toga virus (e.g., Sindbis virus and Semliki forest virus) and vesicular stomatitis virus (VSV).

[0119] A vector comprising a nucleotide acid can be expressed when the nucleic acid is connected to expression elements so as to function. The term "connected so as to function" (operably linked) means that a physical or functional relation between the elements referred to that permit them to operate in their intended fashion. Accordingly, the nucleic acid "operably linked" to an expression control element means that the control element modulates nucleic acid transcription and, as appropriate, translation of the transcription product.

[0120] The "expression control element" or "expression control sequence" is a polynucleotide which influences upon expression of an operably linked nucleic acid. Promoters and enhancers are non-limiting specific examples of expression controlling elements and sequences. The "promoter" is a cis-acting DNA regulatory region which can initiate transcription of downstream (3' direction) nucleic acid sequence. A nucleotide which accelerates transcription initiation is included in the promoter sequence. The enhancer also regulates nucleic acid expression but acts at a distance from the transcription initiation site of the nucleic acid to which it is operably linked. When the enhancer is present in either the 5' or 3' end of the nucleic acid as well as within the nucleic acid (e.g., intron or coding sequence), the enhancer further functions. Additional examples of the expression control element include a leader sequence and a fusion partner sequence, an internal ribosome entry site (IRES) element for preparing multigene, or polycistronic message, splicing signal of intron, maintenance of correct reading frame of gene to enable inframe translation of mRNA, polyadenylation signal which produces proper polyadenylation of the transcription product of interest, and stop codons.

[0121] Examples the expression control element include a "constitutional" element in which transcription of an operably linked nucleic acid occurs without the presence of signals or stimulus. The expression control element which confers expression in response to the signal or stimulus and increase or decrease expression of the operably linked nucleic acid is "adjustable". The adjustable element which increases expression of the operably linked nucleic acid in response to a signal or stimulus is called "inducible element". The adjustable element which decreases expression of the operably linked nucleic acid in response to a signal or stimulus is called "repressor element" (namely, the signal decreases the expression; and the expression increases when the signal is removed or not present).

[0122] Examples of the constitutional promoter for bacterial expression include an inducing promoter, such as T7 and pL, plac, ptrp and ptac (ptrp-lac hybrid promoter) of bacteriophage λ and the like. For insect cell system, a constitutional or an inducible promoter (e.g., ecdysone) can be used. The constitutional promoter for yeast include an inducing promoter such as ADH, LEU2, GAL and the like (e.g., see Ausubel et al., In: Current Protocols in Molecular Biology, Vol. 2, Chapter 13, Greene Publish. Assoc. & Wiley Interscience edition, 1988; Grant et al., In: Methods in Enzymology, 153: 516-544 (1987) Wu & Grossman, 1987, Acad. Press, N.Y.; Glover, DNA Cloning, Vol. 11, Chapter 3, IRL Press, Wash., D.C., 1986; Bitter, In: Methods in Enzymology, 152: 673-684 (1987), edited by Berger & Kimmel, Acad. Press, N.Y.; and Strathern et al., The Molecular Biology of the Yeast *Saccharomyces*, edited by Cold Spring Harbor Press, Vol. 1 and Vol. 11 (1982)).

[0123] For the expression in mammals, a constitutional promoter derived from a virus or other origin can be used. For example, inducible promoters derived from CMV, SV40, or a viral long terminal repeated sequence (LTR), or mammal cell genome (e.g., metallothionein IIA promoter; heat shock promoter, steroid/thyroid hormone/retinoic acid responding element) or mammal virus (e.g., adenovirus late promoter; mouse breast cancer virus LTR) can be used.

[0124] Examples of the expression control element include an element which is active in a specific tissue or cell types, and such an element is called "tissue specific expression control element". In general, the tissue specific expression control element is more active in specific cells or tissue types, and this is because this tissue specific expression control element is recognized by a transcription activating protein which is active in the specific cell or tissue types or by other transcription factor, as compared to other cells or tissue types. Non-limiting specific examples of such an expression control element are hexokinase II, COX-2, α -fetoprotein, carcinoembryonic antigen, DE3/MUC1, prostate specific antigen, C-erbB2/neu, glucose-dependent insulin secretion stimulatory polypeptide (GIP), telomerase reverse transcriptase and a promoter such as hypoxia-responsive promoter.

[0125] According to the invention, a host cell transformed or transfected with IL-3R α nucleic acid or vector of the invention is provided. Examples of the host cells, but are not limited to, include prokaryotic cell and eukaryotic cell, such as, bacteria, fungi (yeast), and cells of plants, insects and animals (e.g., mammals such as primates, human and the like). Non-limiting examples of transformed cell include a bacteria transformed with a recombinant bacteriophage nucleic acid, a plasmid nucleic acid or cosmid nucleic acid expression vector; a yeast transformed with a recombinant yeast expression vector; a plant cell infected with a recombinant virus expression vector (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with a recombinant plasmid expression vector (e.g., Ti plasmid); an insect cell infected with a recombinant virus expression vector (e.g., baculovirus); and an animal cell infected with a recombinant virus expression vector (e.g., retrovirus, adenovirus, vaccinia virus) or a transformed animal cell manipulated for stable expression. CHO cell is a non-limiting example of a mammal host cell which expresses an IL-3R α antibody and its subsequence thereof and fragment. The host cell may be a plurality or population of cells from a primary cell-separated

line, an isolated secondary cell or subcultured cell, or an established cell line or immortalized cell culture.

[0126] The term "be transformed" or be transfected" when it is used in reference to a cell (e.g., host cell) or an organism means a change of gene in a cell after incorporation of an exogenous molecule, such as a protein or a nucleic acid (e.g., transgene), into the cell. Accordingly, the "transfected" or "transformed" cell is a cell into which the exogenous molecule is introduced by the hand of man by, for example, by recombinant DNA techniques or a progeny thereof.

[0127] The nucleic acid or protein can be transfected or transformed (expressed) in the cell or a progeny thereof stably or temporarily. The introduced protein can be expressed by growing the cell, or transcribing the nucleic acid. Since there is a possibility that a mutation occurs during replication, there is a case that a progeny of the transfected or transformed cell is not identical to the parent cell.

[0128] In general, a vector is used in the cell transfection or transformation. The vector can be included in a viral particle or vesicle and can be optionally directed demands to a specific cell types by including a protein on the particle or vesicle surface which binds to a target cell ligand or receptor. Accordingly, a cell can be used as a target by preparing the viral particle or vesicle itself or the viral surface protein, for the purpose of an *in vitro*, *ex vivo* or *in vivo* transfection or transformation. Accordingly, the vector includes *in vitro*, *in vivo* and *ex vivo* delivering techniques of viral and non-viral vectors into a cell, tissue or organ.

[0129] In addition, introduction of a nucleic acid into a target cell (e.g., a host cell) can also be carried out by a method conventionally known in the technical field, such as osmotic shock (e.g., calcium phosphate), electroporation, microinjection, cell fusion and the like. The introduction of nucleic acid and polypeptide *in vitro*, *ex vivo* and *in vivo* can also be carried out using other techniques. For example, a polymer substance such as polyester, polyamide, polyacetal, hydrogel, polyvinyl pyrrolidone, ethylene-vinyl acetate, methyl cellulose, carboxymethylcellulose, protamine sulfate, or lactide/glycolide copolymer, polylactide/glycolide copolymer, or ethylene vinyl acetate copolymer and the like. The nucleic acid can be enclosed in a hydroxymethyl cellulose or gelatin-microcapsule, or a microcapsule prepared using poly(methyl methacrylate) microcapsule, or in a colloid system, respectively, by a coacervation technique or by interfacial polymerization. The colloid dispersion system includes a system based on a polymer complex, nanocapsule, microsphere, beads and lipid (oil-in-water type emulsion, micelle, mixed micelle, liposome and the like).

[0130] The liposome for introducing various compositions into cells is conventionally known in the technical field, and for example, phosphatidylcholine, phosphatidylserine, lipofectin and DOTAP are included therein (e.g., US Patent No. 4,844,904, US Patent No. 5,000,959, US Patent No. 4,863,740 and US Patent No. 4,975,282; and GIBCO-BRL, Gaithersburg, Md.). Piperazine based amphiphilic cationic lipids which is useful in gene therapy (see e.g., US Patent No. 5,861,397) are also known. A cationic lipid system is also known (see e.g., US Patent No. 5,459,127). In this specification, the polymer substance, microcapsule and colloid dispersion system (liposome and the like) are collectively called as "vesicle".

[0131] In addition, examples of the suitable techniques which can be used in the method for producing an antibody are affinity purification, non-modified gel purification, HPLC or RP-HPLC, size exclusion, purification by protein A column and an optional combination of these techniques. An IL-3R α antibody isotype can be determined using ELISA assay, and for example, human Ig can be identified using mouse Ig absorbed anti-human Ig.

[0132] Binding affinity can be determined by association (K_a) and dissociation (K_d) rates. The equilibrium affinity constant K_D is the ratio of K_a/K_d . The association (K_a) and dissociation (K_d) rates can be measured using surface plasmon resonance (SPR) (Rich and Myszk, Curr. Opin. Biotechnol., 11: 54 (2000); Englebienne, Analyst., 123: 1599 (1998)). Instrumentation and methods for real time detection and monitoring of association rate are conventionally known and commercially available (BiaCore 2000, Biacore AB, Uppsala, Sweden; and Malmqvist, Biochem. Soc. Trans., 27:335 (1999)). The K_D value can be defined as the IL-3R α antibody concentration required to saturate one half of the binding site (50%) on IL-3R α .

(Crossing property in primates)

[0133] Currently, although as many as 500 therapeutic antibodies are being developed in the world, it is said that human antibodies have a high possibility to be able to avoid problems of immunogenicity. However, on the other hand, there are many cases in which drug efficacy of human antibodies are not exhibited at all in rodents. In that case, there are many cases in that primates have to be used in toxicity tests, and furthermore the reactivity is found only in chimpanzee is not rare in many cases. When the pharmacological reaction can be found only in chimpanzee, the toxicity test is further significantly constrained. In the first place, facilities where chimpanzee experiments can be carried out are considerably limited, individuals are infected with HIV in many cases and there are also problems of labor hygiene of workers involved in the experiments. In addition, regarding chimpanzee, there are large limitations that anatomy test after final drug administration cannot be carried out and of reproductive toxicity test is also impossible to carry out and the like. Accordingly, the ability to verify drug efficacy in monkey (*Macaca fascicularis* and/or *Macaca mulatta*) is useful from the viewpoint of advancing toxicity tests which are essential for developing pharmaceuticals.

[0134] Regarding the method for confirming monkey crossreactivity with monkey, it can be confirmed by a conven-

tionally known method such as immunochemical tissue staining method, solid phase enzyme immunoassay (hereinafter, "ELISA"), flow cytometry (FCM) and the like.

(Pharmaceutical composition)

[0135] Antibodies can be included in a pharmaceutical composition. In an embodiment, an antibody comprises a pharmaceutically acceptable carrier, a stabilizer or a filler and is prepared in the form of aqueous solution or as a freeze-dried preparation. Typically, an appropriate amount of a pharmaceutically acceptable salt is used for isotonicity of the pharmaceutical preparation. Examples of the acceptable carrier, stabilizer or filler include a buffer solution such as phosphate, citrate and other organic acid and the like; a low molecular weight (less than 10 in the number of residues) polypeptide; a protein such as serum albumin, gelatin, immunoglobulin and the like; a hydrophilic polymer such as polyvinyl pyrrolidone; an amino acid such as glycine, glutamine, asparagine, histidine, arginine, lysine and the like; a monosaccharide, disaccharides and other carbohydrates such as glucose, mannose, dextrin and the like; a chelating agent such as EDTA and the like; saccharides such as sucrose, mannitol, trehalose, sorbitol and the like; a salt forming counter ion such as sodium and the like; a metal complex (e.g., Zn-protein complex); an antiseptic (octadecyl dimethylbenzylammonium chloride; hexamethonium chloride; banzalconium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); and/or a nonionic surfactant such as TWEEN™, PLURONICS™, polyethylene glycol (PEG) and the like.

(Therapeutic use of antitumor substance which targets IL-3R α , expression cells)

[0136] Examples of the diseases for which the therapeutic use is examined, but are not limited thereto, include the diseases which can be considered to treat by binding or targeting IL-3R α -expressing blood tumor cells (AML cell, CML cell, MDS cell, ALL cell, CLL cell, multiple myeloma cell and the like), mastocyte, basophile, helper T cell (e.g., Th1 cell, Th17 cell), regulatory T cell (e.g., CD4 positive CD25 positive cell), antigen presenting cell (e.g., dendritic cell, monocyte-macrophage and related cells (hepatic stellate cell, osteoclast, microglia, intraepidermal macrophage, dust cell (alveolar phagocyte) and the like)).

[0137] Examples of the disease for which therapeutic use is examined include a blood disease in which expression of IL-3R α is found in bone marrow or peripheral blood. Specific example may include acute myeloid leukemia (AML). Based on the FAB classification (French-American-British criteria) which can determine which stage of the cell among the cells in the course of differentiating into various blood cells from the hematopoietic stem cell caused tumorigenic transformation, the acute myeloid leukemia is classified into disease types of M0 (micro-differentiation type myeloblastic leukemia), M1 (undifferentiated myeloblastic leukemia), M2 (differentiated myeloblastic leukemia), M3 (acute promyelocytic leukemia), M4 (myelomonocytic leukemia), M5 (monocytic leukemia), M6 (erythroleukemia), M7 (megakaryocytic leukemia) and subtypes thereof. In addition, further examples of diseases include acute lymphocytic leukemia, atypical leukemia, chronic lymphocytic leukemia, adult T cell leukemia, NK/T cell lymphoma, granular lymphocytosis (LGL leukemia), polycythemia vera, essential thrombocythemia, hypereosinophilic syndrome, Hodgkin lymphoma, non-Hodgkin lymphoma, follicular lymphoma, MALT lymphoma, mantle cell lymphoma, diffuse large B-cell lymphoma, Burkitt lymphoma, lymphoblastic lymphoma and Catsleman disease.

[0138] The method of the invention which comprises administration or delivery of an IL-3R α antibody and an antitumor substance which targets an IL-3R α expression cell can be carried out by any acceptable method. In a specified embodiment, these are administered to a subject, locally, regionally or systemically.

[0139] In addition, regarding the IL-3R α antibody, the antitumor substance which targets IL-3R α expression cell for treating the above-mentioned diseases can also be considered to combine with other therapeutic agent suitable for the same disease (typically a chemotherapeutic agent) or be administered in combination with radiotherapy. Examples of the suitable other therapeutic agent include a chemotherapeutic agent such as cytarabine (Ara-C), an anthracycline system antitumor agent (typically, daunorubicin (DNR), idarubicin (IDA)) and the like, a differentiation induction therapeutic agent such as all-trans retinoic acid (ATRA), arsenious acid, Am80 (tamibarotene), gemtuzumab-ozogamicin (ozogamicin conjugate anti-CD33 antibody), topotecan, fludarabine, cyclosporine, mitoxantrone (MIT), interferon and imatinib, but are not limited thereto, and also include a combination with a therapeutic method considered to be clinically effective.

[0140] Mammals (e.g., human) are included in the subject which can be treated by the invention. In a specified embodiment, it is a subject who is a candidate of blood tumor or a subject who received treatment of the blood tumor, a subject having a possibility causing IL-3R α -mediated cellular response or a subject who received treatment of the IL-3R α -mediated cellular response, a subject who is a candidate of a myelocytic malignant tumor or a subject who received treatment of the myelocytic malignant tumor or a subject who is a candidate of acute myeloid leukemia or a subject who received treatment of the acute myeloid leukemia.

[0141] According to this specification, the terms "treat", "treating", "treatment" and the grammatical variations thereof

mean a protocol, a planning, a process or an improving method which is carried out on each subject who is desirable to obtain physiological effect or good outcome on the patient. Accordingly, the method of the invention includes a treatment and a treating method which produce measurable improvement or beneficial effect, particularly on a disorder, a disease, pathology, a condition of a disease or a symptom of a given subject. The measurable improvement or profitable effect is objective or subjective, immoderate, transient or long-term improvement of any one of disorders, diseases, pathology, conditions of a disease or symptoms, or a reduction in onset, severity, duration or frequency of adverse symptom related to or caused by disorders, diseases, physiological conditions, pathology or state. According to the method of the invention, there is a possibility that its effect is not always exhibited immediately, but eventual improvement or beneficial effect is found a little later with the lapse of time, so that stabilization or amelioration in a give subject will occur.

[0142] Unless otherwise noted, all of the technical terms and scientific terms used in this specification have the same meanings of those which are generally evident for persons in the technical field to which the invention is related. Methods and materials similar or equivalent to those described in this specification can be used in the operations or examinations of the invention, but those which are described in this specification are suitable methods and materials.

EXAMPLES

Example 1 Preparation of human, *Macaca fascicularis* or *Macaca mulatta* IL-3R α , expression cell

(Molecular cloning of IL-3R α cDNA and preparation of expression vector)

[0143] Human IL-3R α cDNA was amplified from a blood cell-derived DNA (CLONTECH Human MTC Panel) by PCR using ExTaq (TAKARA BIO INC.). As a PCR device, GeneAmp PCR System 9700 (Applied Biosystems, hereinafter, the PCR device is the same in this specification) was used. Regarding the PCR, after a denaturation step at 94°C for 5 minutes, a three step reaction at 94°C 30 seconds-55°C 30 seconds-72°C 2 minutes was carried out 40 cycles and then a reaction at 99°C for 30 seconds was carried out. The PCR primers used are as follows.

IL-3R α _Fw: 5'-CGGCAATTGCCACCATGGTCCTCCTTTGGCTCAC-3' (SEQ ID NO:3)

IL-3R α _Re: 5'-ATTGCGGCCGCTCAAGTTTTCTGCACGACCT-3' (SEQ ID NO:4)

[0144] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A band at around 1.2 kb was cut out and extracted using JetSob (Genomed). The extracted DNA was digested with *MfeI* and *NotI*, mixed with pEGFP-N1 vector (Clontech) or pEF6/Myc-His vector which had been digested with *EcoRI* and *NotI* and ligated using TaKaRa Ligation Kit. Regarding the transformation, the ligation sample and a DH10B competent cell were mixed and spread on LB plate (containing kanamycin). Insert check of the pEGFP-N1 vector was carried out by colony direct PCR using LA Taq (Takara Shuzo Co., Ltd.). Regarding the PCR, after a denaturation step at 94°C for 5 minutes, a three step reaction at 94°C 30 seconds-55°C 30 seconds-72°C 2 minutes was carried out 40 cycles and then a reaction at 99°C for 30 seconds was carried out. Regarding the primers used, IL-3R α -Fw and IL-3R α -Re were used.

[0145] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. Using a colony from which amplification at around 1.2 kb was obtained, nucleotide sequence was determined by a direct sequencing method. In the reaction of sequence samples, BigDye(R) Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems) and GeneAmp PCR System 9700 (Applied Biosystems) were used (these were used in the all DNA sequence analyses in this specification). Regarding the primers, IL-3R α -Fw, IL-3R α -Re and the following primer were used. IL-3R α _seqF1: 5'-GTCTTCACTACAAAACGGAT-3' (SEQ ID NO:5)

[0146] ABI 3700XL DNA analyzer (Applied Biosystems) was used as the sequence analyzing device. A clone having the same sequence of the coding region of GenBank association number NP-002174.1 was selected and a plasmid DNA was extracted by a Miniprep method. The vector names were pEGFR-N1/hCD123 and pEF6/Myc-His/hCD123, respectively.

[0147] The sequence of the insert (*MfeI* to *NotI*) was as follows.

```
CAATTGCCACCATGGTCCTCCTTTGGCTCACGCTGCTCCTGATCGCCCTGCCC
TGTCTCCTGCAAACGAAGGAAGATCCAAACCCACCAATCACGAACCTAAGG
ATGAAAGCAAAGGCTCAGCAGTTGACCTGGGACCTTAACAGAAATGTGACC
GATATCGAGTGTGTTAAAGACGCCGACTATTCTATGCCGGCAGTGAACAATAG
CTATTGCCAGTTTGGAGCAATTTCTTATGTGAAGTGACCAACTACACCGTCC
GAGTGGCCAACCCACCATTCTCCACGTGGATCCTCTTCCCTGAGAACAGTGG
```

GAAGCCTTGGGCAGGTGCGGAGAATCTGACCTGCTGGATTTCATGACGTGGAT
 TTCTTGAGCTGCAGCTGGGCGGTAGGCCCGGGGGCCCCCGCGGACGTCCAG
 5 TACGACCTGTACTTGAACGTTGCCAACAGGCGTCAACAGTACGAGTGTCTTC
 ACTACAAAACGGATGCTCAGGGAACACGTATCGGGTGTCGTTTTCGATGACAT
 CTCTCGACTCTCCAGCGGTTCTCAAAGTTCCCACATCCTGGTGCGGGGCGAGG
 10 AGCGCAGCCTTCGGTATCCCCTGCACAGATAAGTTTGTCTCTTTTCACAGAT
 TGAGATATTAACCTCCACCCAACATGACTGCAAAGTGTAATAAGACACATTCTT
 TTATGCACTGGAAAATGAGAAGTCATTTCAATCGCAAATTTTCGCTATGAGCTT
 15 CAGATACAAAAGAGAATGCAGCCTGTAATCACAGAACAGGTCAGAGACAGA
 ACCTCCTTCCAGCTACTCAATCCTGGAACGTACACAGTACAAATAAGAGCCC
 GGGAAAGAGTGTATGAATTCTTGAGCGCCTGGAGCACCCCCCAGCGCTTCGA
 GTGCGACCAGGAGGAGGGCGCAAACACACGTGCCTGGCGGACGTCGCTGCT
 20 GATCGCGCTGGGGACGCTGCTGGCCCTGGTCTGTGTCTTCGTGATCTGCAGA
 AGGTATCTGGTGATGCAGAGACTCTTTCCCCGCATCCCTCACATGAAAGACC
 CCATCGGTGACAGCTTCCAAAACGACAAGCTGGTGGTCTGGGAGGCGGGCA
 25 AAGCCGGCCTGGAGGAGTGTCTGGTGACTGAAGTACAGGTCGTGCAGAAAA
 CTTGAGCGGCCGC (SEQ ID NO:6)

[0148] The *Macaca fascicularis* and *Macaca mulatta* cDNA samples were amplified from a *Macaca fascicularis* bone marrow-derived cDNA or *Macaca mulatta* bone marrow-derived cDNA by a PCR method using LA Taq (TAKARA BIO INC). GeneAmp PCR System 9700 (Applied Biosystems) was used as the PCR device. Regarding the PCR, after a denaturation step at 95°C for 1 minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 70 seconds was carried out 40 cycles and then a reaction at 72°C for 2 minutes was carried out. Subsequences were obtained through BLAST retrieval for the public data base of *Macaca mulatta* genome (<http://www.hgsc.bnm.tmc.edu/blast.hgsc>), based on the hIL-3Rα cDNA sequence to design primers. The used primer sequences were as follows.

Rhe123Fw1:CGGCAATTGCCACCATGACCCTCCTTTGGCTGACGCTG (SEQ ID NO:7)

Rhe123Rv1:TATATTGCGCCGCTCAAGTTTCTCCACCACCTGCAC (SEQ ID NO:8)

[0149] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A band at around 1.2 kb was cut out and the DNA was extracted using Gel Extraction Kit (QIAGEN). The thus extracted DNA was mixed with pGEM-T Easy vector (Promega) and ligated using TaKaRa Ligation Kit. Regarding the transformation, the ligation sample and a DH10B competent cell were mixed and spread on LB plate (containing ampicillin). Insert check of the pGEM-T Easy vector was carried out by colony direct PCR using LA Taq (Takara Shuzo Co., Ltd.). Regarding the PCR, after a denaturation step at 95°C for 1 minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 1 minute was carried out 35 cycles and then a reaction at 72°C for 2 minutes was carried out. The following were used as the primers.

T7: TAATACGACTCACTATAGGG (SEQ ID NO:9)

SP6: GATTTAGGTGACACTATAG (SEQ ID NO: 10)

[0150] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. Using a colony from which amplification at around 1.2 kb was obtained, nucleotide sequence was determined by a direct sequencing method. As the PCR primers, T7 and SP6 were used. A clone showing no mutation by PCR was selected and its plasmid DNA was extracted by the Miniprep method. The thus obtained DNA was digested with *MfeI* and *NotI*, mixed with pEGFP-N1 vector (Clontech) which had been cleaved with *EcoRI* and *NotI* and ligated using TaKaRa Ligation Kit. Regarding the transformation, the ligation sample and a DH10B competent cell were mixed and spread on LB plate (containing kanamycin).

[0151] Insert check of the pEGFP-N1 vector was carried out by a colony direct PCR using La Taq (Takara Shuzo Co., Ltd.). Regarding the PCR, after a denaturation step at 94°C for 5 minutes, a three step reaction at 94°C 30 seconds-55°C 30 seconds-72°C 2 minutes was carried out 40 cycles and then a reaction at 99°C for 30 seconds was carried out. Regarding the used PCR primers, Rhe123Fw1 and Rhe123Rv1 were used.

[0152] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. Using a colony from which amplification at around 1.2 kb was obtained, nucleotide sequence was determined by a direct sequencing method. BigDye(R) Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems) and GeneAmp PCR System 9700 (Applied Biosystems) were used in the reaction of sequence sample (these were used in all of the DNA sequence analyses in this specification). As the primers, Rhe123Fw1 and Rhe123Rv1 were used. The vectors were named pEGFR-N1/cyCD123 and pEGFR-N1/rhCD123, respectively.

[0153] The sequence of the insert (*MefI* to *NotI*) of *Macaca fascicularis* IL-3R α was as follows.

```

10  CAATTGCCACCATGACCCTCCTTTGGCTGACGCTGCTCCTGGTCGCCACGCC
    CTGTCTCCTGCAAACGAAGGAGGATCCAAATGCACCAATCAGGAATCTAAGG
    ATGAAAGAAAAGGCTCAGCAGTTGATGTGGGACCTGAACAGAAACGTGACC
15  GACGTGGAGTGTATCAAAGGCACCGACTATTCTATGCCGGCAATGAACAACA
    GCTATTGCCAGTTCGGAGCCATTTCCTTATGTGAAGTGACCAACTACACCGTC
    CGAGTGGCCAGTCCCCCGTTCTCCACGTGGATCCTCTTCCCTGAGAACAGTG
20  GGACGCCTCAGGCAGGCGCGGAGAATCTGACCTGCTGGGTTCATGACGTGG
    ATTTCTTGAGCTGCAGCTGGGTGGCAGGCCCGGCGGCCCGCTGACGTCCA
    GTACGACCTGTACTTGAACAATCCCAACAGCCACGAACAGTACAGGTGCCTT
25  CACTACAAAACGGATGCTCGGGGAACACAGATCGGGTGTCTGGTTCGATGACA
    TCGCTCGACTCTCCCGCGGTTCTCAAAGTTCCCACATCCTGGTGAGGGGCAG
    GAGCGCAGCCGTCAGTATCCCCTGCACAGATAAGTTTGTCTTCTTTTCACAGA
    TTGAGAGATTAACCTCCACCCAACATGACTGGAGAGTGTAATGAGACACATTC
30  CTTTCATGCACTGGAAAATGAAAAGTCATTTCATCGCAAATTCCGCTATGAGC
    TTCGGATCCAAAAGAGAATGCAGCCTGTAAGGACAGAACAGGTCAGAGACA
    CAACCTCCTTCCAGCTACCCAATCCTGGAACGTACACAGTGCAAATAAGAGC
35  CCGGGAAACAGTGTATGAATTCTTGAGTGCCTGGAGCACCCCCCAGCGCTTC
    GAGTGCAGACCAGGAGGAGGGCGCGAGCTCGCGTGCCTGGCGGACGTCGCTG
    CTGATCGCGCTGGGGACGCTGCTGGCCTTGCTCTGTGTGTTTCCTCATCTGCAG
40  AAGGTATCTGGTGATGCAGAGGCTGTTTCCCCGCATCCCACACATGAAAGAC
    CCCATCGGTGACACCTTCCAACAGGACAAGCTGGTGGTCTGGGAGGCGGGC
    AAAGCCGGCCTGGAGGAGTGTCTGGTGTCTGAAGTGCAGGTGGTGGAGAAA
45  ACTTGAGCGGCCGC (SEQ ID NO:11)

```

[0154] The sequence of the insert (*MefI* to *NotI*) of *Macaca mulatta* IL-3R α was as follows.

CAATTGCCACCATGACCCTCCTTTGGCTGACGCTGCTCCTGGTCGCCACGCC
 CTGTCTCCTGCAAACCAAGGAGGATCCAAATGCACCAATCAGGAATCTAAGG
 5 ATGAAAGAAAAGGCTCAGCAGTTGATGTGGGACCTGAACAGAAACGTGACC
 GACGTGGAGTGTATCAAAGGCACCGACTATTCTATGCCGGCAATGAACGACA
 GCTATTGCCAGTTCGGAGCCATTTCTTATGTGAAGTGACCAACTACACCGTC
 10 CGAGTGGCCAGTCCTCCGTTCTCCACGTGGATCCTCTTCCCTGAGAACAGTG
 GGACGCCTCGGGCAGGCGCGGAGAATTTGACCTGCTGGGTTCATGACGTGG
 ATTTCTTGAGCTGCAGCTGGGTGGTAGGCCCGGCGGCCCCCGCTGACGTCCA
 GTACGACCTGTACTTGAACAATCCCAACAGCCACGAACAGTACAGGTGCCTT
 15 CGCTACAAAACGGATGCTCGGGGAACACAGATCGGGTGTCTCGGTTCGATGACA
 TCGCTCGACTCTCCCGCGGTTCTCAAAGTTCCACATCCTGGTGAGGGGCGAG
 GAGCGCAGCCGTCAGTATCCCCTGCACAGATAAGTTTGTCTTCTTTTCACAGA
 20 TTGAGAGATTAACCTCCACCCAACATGACTGGAGAGTGTAATGAGACACATTC
 CTTTCATGCACTGGAAAATGAAAAGTCATTTCAATCGCAAATTCCTACTATGAGC
 TTCGGATCCAAAAGAGAATGCAGCCTGTAAGGACAGAACAGGTCAGAGACA
 25 CAACCTCCTTCCAGCTACCCAATCCTGGAACGTACACAGTGCAAATAAGAGC
 CCGGGAAACAGTGTATGAATTCTTGAGTGCCTGGAGCACCCCCCAGCGCTTC
 GAGTGCAGACCAGGAGGAGGGCGCGAGCTCGCGTGCCTGGCGGACGTCTGCTG
 30 CTGATCGCGCTGGGGACGCTGCTGGCCTTGCTCTGTGTGTTCTCATCTGCAG
 AAGGTATCTGGTGATGCAGAGGCTGTTTCCCCGCATCCACACATGAAAGAC
 CCCATCGGTGACACCTTCCAACAGGACAAGCTGGTGGTCTGGGAGGCGGGC
 35 AAAGCCGGCCTGGAGGAGTGTCTGGTGTCTGAAGTGACAGGTGGTGGAGAAA
 ACTTGAGCGGCCGC (SEQ ID NO:12)

(Preparation of IL-3R α forced expression cell line)

[0155] L929 cell (manufactured by ATCC) and Colon-26 cell (manufactured by ATCC) were infected with pEGFP-N1 vector/hCD123 or pEF6/Myc-His vector/hCD123 using electroporation (BTX). Specifically, 10 to 20 μ g of DNA was mixed with one hundred thousand cells and allowed to react at 300 V and 950 μ F. Regarding the cells, drug resistant cells were selected using neomycin (Calbiochem) for pEGFP-N1/hCD123 or blasticidin (Invitrogen) for pEF6/Myc-His/hCD123. Regarding the thus selected cells, a GFP-positive cell or a cell highly expressing IL-3R α (CD123) was further selected by sorting using flow cytometry (FAC SVantage, FACSria and the like, BD Biosciences) and named as L929/hCD123 and Colon-26/hCD123, respectively.

[0156] Regarding the preparation of *Macaca fascicularis* IL-3R α and *Macaca mulatta* IL-3R α forced expression cells, these were also prepared using L929 and Colon-26 in the same manner as the case of human IL-3R α forced expression cell and named L929/cyCD123, Colon-26/cyCD123, L929/rhCD123 and Colon-26/rhCD123. Example 2 Preparation of soluble form of IL-3R α extracellular region

(Preparation of soluble form of human IL-3R α extracellular region expression vector)

[0157] A cDNA encoding the extracellular region of human IL-3R α was amplified by a PCR method, and FLAG tag was connected to its downstream. Specifically, the cDNA encoding the extracellular region of human IL-3R α was amplified by PCR using pEF6/Myc-His/hCD123 plasmid DNA as the template and using Platinum Pfu polymerase (Invitrogen). Regarding the PCR, after a denaturation step at 96°C for 2 minutes, a three step reaction at 96°C 20 seconds-55°C 30

seconds-68°C 65 seconds was carried out 30 cycles. The primers used were IL-3R α -Fw and the following primer.

hIL-3R α sol-FLAG-NotI: 5'- ATTGCGGCCGCTCACTTATCGTCGTCATCCTTGTAGTCCCGCCAGGCACGTGT
GTTTG-3' (SEQ ID NO: 13)

[0158] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. The DNA was extracted using JetSorb (Genomed). Thus purified DNA was digested with *MfeI* and *NotI* and again subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). A band of around 1.0 kb was cut out and the DNA was extracted using JetSorb (Genomed). The obtained DNA was mixed with a pTracer-CMV/Bsd vector, which had been cleaved using the same enzymes of the purified DNA, and ligated using TaKaRa Ligation Kit. Regarding the transformation, the ligation sample and a DH10B competent cell were mixed and spread on LB plate (containing ampicillin). Insert check was carried out by colony direct PCR using LA Taq (Takara Shuzo Co., Ltd.). Regarding the PCR, after a denaturation step at 95°C for 1 minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 40 seconds was carried out 35 cycles and then an elongation reaction at 72°C for 2 minutes was carried out. The PCR primers used were IL-3R α -Fw and IL-3R α sol-FLAG-NotI.

[0159] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A plasmid DNA was extracted by the Miniprep method from a colony in which amplification of around 1.0 kb was obtained. It was found by a DNA sequence analysis that the purified plasmid DNA has the sequence identical to the corresponding region of GenBank accession number NP-002174.1.

[0160] The sequence of the insert (*MfeI* to *NotI*) was as follows.

```

CAATTGCCACCATGGTCCTCCTTTGGCTCACGCTGCTCCTGATCGCCCTGCCC
TGTCTCCTGCAAACGAAGGAAGATCCAAACCCACCAATCACGAACCTAAGG
ATGAAAGCAAAGGCTCAGCAGTTGACCTGGGACCTTAACAGAAATGTGACC
GATATCGAGTGTGTTAAAGACGCCGACTATTCTATGCCGGCAGTGAACAATAG
CTATTGCCAGTTTGGAGCAATTTCTTATGTGAAGTGACCAACTACACCGTCC
GAGTGGCCAACCCACCATTCTCCACGTGGATCCTCTTCCCTGAGAACAGTGG
GAAGCCTTGGGCAGGTGCGGAGAATCTGACCTGCTGGATTTCATGACGTGGAT

TTCTTGAGCTGCAGCTGGGCGGTAGGCCCGGGGGCCCCCGCGGACGTCCAG
TACGACCTGTACTTGAACGTTGCCAACAGGCGTCAACAGTACGAGTGTCTTC
ACTACAAAACGGATGCTCAGGGAACACGTATCGGGTGTCGTTTCGATGACAT
CTCTCGACTCTCCAGCGGTTCTCAAAGTTCCACATCCTGGTGCGGGGCAGG
AGCGCAGCCTTCGGTATCCCCTGCACAGATAAGTTTGTCGTCTTTTCACAGAT
TGAGATATTAACCTCCACCCAACATGACTGCAAAGTGTAATAAGACACATTCCT
TTATGCACTGGAAAATGAGAAGTCATTTCAATCGCAAATTTTCGCTATGAGCTT
CAGATACAAAAGAGAATGCAGCCTGTAATCACAGAACAGGTCAGAGACAGA
ACCTCCTTCCAGCTACTCAATCCTGGAACGTACACAGTACAAATAAGAGCCC
GGGAAAGAGTGTATGAATTCTTGAGCGCCTGGAGCACCCCCCAGCGCTTCGA
GTGCGACCAGGAGGAGGGCGCAAACACACGTGCCTGGCGGGACTACAAGG
ATGACGACGATAAGTGAGCGGCCGC (SEQ ID NO:14)

```

(Preparation of soluble form of human IL-3R α protein)

[0161] Plasmid DNA of pTracer CMV expression vector containing soluble form of IL-3R α sequence was purified using QIAGEN Plasmid Maxi Kit. A CHORas1 cell was used as a host cell for expression. The CHORas1 cell was cultured with shaking using SFM II medium (Invitrogen) (37°C, 5% CO₂).

[0162] A PEI method was used in the gene introduction. Polyethylenimine, Linear, MW 25,000 (Polysciences) was weighed and dissolved in PBS while adjusting to around pH 7.0 with HCl (1 g/l). The obtained solution was stirred for 1

hour and then sterilized by filtering through a membrane filter having a pore size of 0.22 μm , MILLEX-GV (Millipore). Then, 1 mg of the purified plasmid DNA was mixed with 20 ml of Opti-Pro SFM (Invitrogen) to obtain Solution A. Solution B was prepared by mixing 2.5 ml of PEI solution (1 g/l) with 20 ml of Opti-Pro SFM (Invitrogen). After solution A and Solution B were mixed, and then allowed to stand still for 10 minutes, the obtained solution was added to CHORas1 cells (1,000,000 cells per 1 ml). After six days, the cell supernatant was recovered and used for the protein purification.

[0163] Purification of the soluble form of human IL-3R α protein was carried out by the following method. A culture supernatant containing soluble form of IL-3R α protein was recovered by centrifugation 6 days after the gene introduction and passed through a filter. The obtained solution was diluted 5 times with Tris buffered saline (TBS), an Anti-FLAG column was prepared using anti-FLAG M2 Agarose Affinity Gel (Sigma) and the solution was applied thereto using HiLoad Pump P-50 (Pharmacia Biotech). Elution was carried out using FLAG peptide (Sigma) and in accordance with the manual. The eluate was fractioned into several fractions, each fraction was subjected to SDS-PAGE (MultiGel II Mini 10/20% gradient gel; Cosmo Bio Co., Ltd.) under a reducing condition, and then silver staining and Western blotting were carried out. A silver staining reagent "Daiichi" (Daiichi Pure Chemicals Co., Ltd.) was used for the silver staining. Anti-FLAG M2 antibody (Sigma) and an alkaline phosphatase-labeled rabbit anti-mouse immunoglobulin antibody were used for the Western blotting. A fraction in which the protein of interest was found was concentrated using Amicon Ultra-4 10K (Millipore), and gel filtration chromatography was carried out using Superdex 200 gp (GE Healthcare). After fractionation, each fraction was subjected to SDS-PAGE (MultiGel II Mini 10/20% gradient gel; Cosmo Bio Co., Ltd.) under a reducing condition, and then silver staining and Western blotting were carried out. A silver staining reagent "Daiichi" (Daiichi Pure Chemicals Co., Ltd.) was used in the silver staining. Anti-FLAG M2 antibody (Sigma) and an alkaline phosphatase-labeled rabbit anti-mouse immunoglobulin antibody were used in the Western blotting. A fraction in which the protein of interest was found was concentrated using Amicon Ultra-4 10K (Millipore) and washed with PBS. By carrying out sterilization by filtration using a membrane filter MILLEX-GV (Millipore) having a pore size of 0.22 μm , a soluble form of human IL-3R α protein was obtained. As a result of Limulus test using Limulus ES-II Kit Wako (Wako Pure Chemical Industries, Ltd.), endotoxin was not detected. Regarding concentration of the soluble form of human IL-3R α protein, absorbance at 280 nm was measured and 1 mg/ml was calculated as 1.4 OD.

Example 3 Preparation of anti-human IL-3R α human antibody using human antibody producing mouse

(Human antibody producing mouse)

[0164] The mouse used in the immunization has a genetic background of homozygote on both of endogenous Ig heavy chain and κ light chain disruptions and also simultaneously keeps the 14th chromosomal fragment containing human Ig heavy chain locus (SC20) and human Ig κ chain transgene (KCo5). This mouse was prepared by the crossing of a line A mouse having the human Ig heavy chain locus onto a line B mouse having the human Ig κ chain transgene. The line A is a homozygote on both of endogenous Ig heavy chain and κ light chain disruptions, is a mouse line which maintain the 14th chromosomal fragment (SC20) which can be transmitted to progeny and is described for example in a report by Tomizuka *et al.* [Tomizuka *et al.*, Proc. Natl. Acad. Sci. USA, 2000, Vol. 97: 722]. Also, the line B is a homozygote on both of endogenous Ig heavy chain and κ light chain disruptions, is a mouse line (transgenic mouse) which maintains a human Ig κ chain transgene (KCo5) and is described in a report by such as Fishwild *et al.* [Nat. Biotechnol (1996), 114: 845].

[0165] An individual in which human Ig heavy chain and κ light chain were simultaneously detected in serum, obtained by the crossing of a line A male mouse onto a line B female mouse or the crossing of a line A female mouse onto a line B male mouse, [Ishida & Lonberg, IBC's 11th Antibody Engineering, Abstract 2000] was used in the following immune test. In this connection, the aforementioned human antibody producing mouse (referred to as KM mouse) can be obtained from Kyowa Hakko Kirin Co., Ltd. by establishing a contract.

(Preparation of human monoclonal antibody to human IL-3R α)

[0166] Regarding the preparation of monoclonal antibody in this Example, it was prepared in accordance with a general method described in A Guide to Monoclonal Antibody Experimental Operations (written in Japanese) (edited by Tamie Ando *et al.* published by Kodansha, 1991) and the like. For the IL-3R α as an immunogen, an IL-3R α expressing L929 cell (CCL-1, ATCC), an IL-3R α expressing Colon-26 cell (Cell Resource Center for Biomedical Research Institute of Development, Aging and Cancer Tohoku University) or a soluble form of human IL-3R α Fc fusion protein was used. As an animal to be immunized, the above-mentioned KM mouse was used.

[0167] For the purpose of preparing human monoclonal antibody to human IL-3R α , the KM mouse was immunized with the IL-3R α expression L929 cell or IL-3R α expression Colon-26 cell prepared in Example 1, intraperitoneally at a dose of 1×10^7 cells/animal every 1 week to 2 weeks in a total of 4 times. Three days before the extraction of spleen which is described below, 20 μg /mouse individual of the soluble form of human IL-3R α protein was administered through

the caudal vein.

[0168] After the spleen was surgically obtained from the immunized mouse, the spleen was put into PBS and minced on a mesh (cell strainer, FALCON) using a syringe piston. After the cell suspension was passed through the mesh and was centrifuged, the obtained precipitated cells were re-suspended in Red Blood Cell Lysing Buffer (Sigma). After 5 minutes of incubation at room temperature, serum-free DMEM medium (Invitrogen) containing 350 mg/ml sodium bi-carbonate, 50 units/ml penicillin and 50 µg/ml streptomycin (hereinafter referred to as "serum-free DMEM medium") was added thereto to precipitate the cells. By suspending again in the serum-free DMEM medium, the number of cells was measured.

[0169] On the other hand, a myeloma cell SP2/0 (ATCC No. CRL-1581) was cultured at 37°C in the presence of 5% carbon dioxide using DMEM medium (Invitrogen) containing 10% FCS (Invitrogen), 50 units/ml penicillin and 50 µg/ml streptomycin (hereinafter referred to as "serum-containing DMEM medium"). The SP2/0 cells were washed with serum-free DMEM medium. In the same manner, the cells were suspended in serum-free DMEM medium to measure the number of cells. After the suspension of the recovered spleen-derived cells and a suspension of the mouse myeloma were mixed at a cell number ratio of 5:1, the mixed suspension was centrifuged, and then the supernatant was completely removed. As a fusion agent, 50% (w/v) polyethylene glycol 1500 (Boehringer-Mannheim) was slowly added to the obtained pellet while stirring the pellet with the tip of a pipette, and then serum-free DMEM medium heated to 37°C in advance was slowly added thereto. Furthermore, an appropriate amount of serum-free DMEM medium was slowly added thereto. Thereafter, the obtained solution was allowed to stand still at 37°C for 5 minutes in the presence of 5% carbon dioxide. After centrifugation, the supernatant was removed and thus obtained fused cells were suspended in DMEM medium (Invitrogen) containing 10% FCS (Invitrogen), penicillin-streptomycin-glutamine (Sigma), IL-6 (5 ng/ml) and 2-mercaptoethanol (Invitrogen) (hereinafter, referred to as "IL-6-containing DMEM medium") and cultured at 37°C in the presence of 5% carbon dioxide. On the next day, the cells were recovered by pipetting, and precipitated by centrifugation. The obtained cell pellet was re-suspended in the IL-6-containing DMEM medium. The suspended cells were subjected to limiting dilution on a 96-well plate and cultured for about 7 days to 14 days. The culture supernatant was used in the hybridoma screening described in the following example.

(Screening of hybridoma producing a human monoclonal antibody which binds to human IL-3Rα)

[0170] Screening of hybridoma was carried out using the cell supernatant prepared in the above example. The method was, in short, carried out by a flow cytometry in which a human IL-3Rα stable expression cell line was used.

[0171] Specifically, a combination of human IL-3Rα expression L929 cell and parent cell line L929 cell or a combination of human IL-3Rα expression Colon-26 cell and parent cell line Colon-26 cell, was mixed with the supernatant of hybridoma and allowed to stand still at 4°C for 30 minutes. After washing the obtained cells twice with a staining medium (Dulbecco's PBS containing 2% fetal calf serum, 2 mM EDTA, 0.05% NaN₃), Goat F(ab')₂ Anti-Human IgG-PE (Southern Biotech) as a secondary antibody was added thereto and allowed to stand still at 4°C for 30 minutes. After washing twice with the staining medium, the obtained cells were analyzed by FACSCalibur (BD Biosciences). A hybridoma which reacted with only human IL-3Rα expression L929 cell was collected.

[0172] The selected hybridoma was subjected to limiting dilution, and screening was carried out using its culture supernatant. Specifically, each of human IL-3Rα expression L929 cell and parent cell line L929 cell was mixed with the supernatant of hybridoma and allowed to stand still at 4°C for 30 minutes. After washing twice with the staining medium, Goat F(ab')₂ Anti-Human kappa-PE (Dako) as a secondary antibody was added thereto and allowed to stand still at 4°C for 30 minutes. After washing twice with the staining medium, the cells were analyzed by FACSCalibur (BD Biosciences). A hybridoma which reacted with only human IL-3Rα expression L929 cell was collected.

Example 4 Preparation of recombinant anti-human IL-3Rα human antibody (Obtaining of anti-human IL-3Rα human antibody gene from hybridoma and preparation of expression vector)

[0173] From the hybridoma obtained in Example 3, clone names Old 4, Old5, Old17, Old19, New102 and Old6 were cultured using eRDF medium (Kyokuto Pharmaceutical Industrial Co., Ltd.) containing 10 ng/ml IL-6 (R & D Systems) and 10% Fetal Bovine Serum (SIGMA) and then cells were collected by centrifugation. To the obtained cells, and then TRIZOL (GIBCO) was added and total RNA was extracted in accordance with the instructions. Cloning of variable region of the antibody cDNA was carried out using SMART RACE cDNA Amplification Kit (Clontech) in accordance with the instructions attached thereto.

[0174] In addition, variable region was cloned from a hybridoma ATCC, HB12009, which produces an anti-human IL-3Rα mouse antibody 7G3 and was used as a control. By connecting the thus obtained cDNA with a DNA encoding human IgG1 constant region, a chimeric antibody expression vector was prepared. Specifically, the cells were collected by centrifugation from the cryopreserved hybridoma, and then TRIZOL (GIBCO) was added thereto to extract total RNA in accordance with the instructions. Cloning of variable region of the antibody cDNA was carried out using mouse IgG

antibody-specific primes in addition to the SMART RACE cDNA Amplification Kit (Clontech), in accordance with the instructions attached thereto.

[0175] Using 5 µg of the total RNA as a template, the 1st strand cDNA was prepared.

5 1) Synthesis of the 1st strand cDNA

[0176] Total RNA 5µgm/3µL

5'CDS 1µL

SMART oligo 1µL

10 **[0177]** After the reaction mixture comprising the above compositions was incubated at 72°C for 2 minutes,

5×Buffer 2µL

DTT 1µL

DNTP mix 1µL and

SuperscriptII 1µL

15 were added to the reaction mixture and incubated at 42°C for 1.5 hours.

[0178] To the obtained mixture, 100 µl of Tricine Buffer was further added and incubated at 72°C for 7 minutes.

2) Amplification of heavy chain gene and light chain gene by PCR and construction of expression vector of recombinant antibody

20

[0179] For amplification of cDNA, Z-Taq manufactured by Takara was used.

cDNA 2µL

10×Z-Taq Buffer 5µL

dNTPmix 4µL

25 Z-Taq1µL

Primer 1

Primer 2

[0180] A reaction solution comprising the above-mentioned composition was adjusted to a final volume of 50 µl with re-distilled water to be subjected to PCR.

30 **[0181]** For amplification of the heavy chain, UPM (SMART RACE cDNA Amplification Kit; manufactured by Clontech) and hh-6 primer (5'-GGTCCGGGAGATCATGAGGGTGTCTT-3') (SEQ ID NO:15) were used and a cycle of reactions at 98°C 1 second and 68°C 30 seconds was repeated 30 times. Further, using 1 µl of the obtained reaction solution as a template and using NUP (SMART RACE cDNA Amplification Kit; manufactured by Clontech) and hh-3 primer (5'-GTGCACGCCGCT GGT CAGGGCGCCTG-3') (SEQ ID NO:16) as primers, a cycle of reactions at 98°C 1 second and 68°C 30 seconds was repeated 20 times. Thereafter, the amplified PCR products were purified using PCR purification kit (QIAGEN), and nucleotide sequences was determined using hh-4 (5'-GGT GCC AGG GGG AAG ACC GAT GG-3') (SEQ ID NO: 17) as a primer. The following specific primers were synthesized based on sequence information, and the sequences were also determined from the opposite direction using the following primers.

Old4 heavy chain specific primer Fw

40 (5'-AGAGAGAGAGGTTCGACCACCATGGACTGGACCTGGAGGTTCTCTTTG T -3') (SEQ ID NO:18)

Old4 heavy chain specific primer Rv

(5'-AGAGAGAGAGGCTAGCTGAAGAGACGGTGACCATTGTCCC -3') (SEQ ID NO:19)

Old5 heavy chain specific primer Fw

(5'-AGAGAGAGAGGTTCGACCACCATGGACTGGACCTGGAGGTTCTCT TTG T -3') (SEQ ID NO:20)

45 Old5 heavy chain specific primer Rv

(5'-AGAGAGAGAGGCTAGCTGAAGAGACGGTGACCATTGTCCC-3') (SEQ ID NO:21)

Old17 heavy chain specific primer Fw

(5'-AGAGAGAGAGGTTCGACCACCATGGACTGGACCTGGAGGTTCTCT TTG T -3') (SEQ ID NO:22)

Old 17 heavy chain specific primer Rv

50 (5'-AGAGAGAGAGGCTAGCTGAGGAGACGGTGACAAGGGTTCCC-3') (SEQ ID NO:23)

Old19 heavy chain specific primer Fw

(5'-AGAGAGAGAGGTTCGACCACCATGGACTGGACCTGGAGGTTCTCT TTG T -3') (SEQ ID NO:24)

Old19 heavy chain specific primer Rv

(5'-AGAGAGAGAGGCTAGCTGAGGAGACGGTGACCAGGGTTC -3') (SEQ ID NO:25)

55 New102 heavy chain specific primer Fw

(5'-AGAGAGAGAGGTTCGACCACCATGGACTGGACCTGGAGGTTCTCTTTG T -3') (SEQ ID NO:26)

New102 heavy chain specific primer Rv

(5'-AGAGAGAGAGGCTAGCTGAGGAGACGGTGACCAGGGTT -3') (SEQ ID NO:27)

Old6 heavy chain specific primer Fw

(5'-AGAGAGAGAGGTTCGACCCACCATGGAAGTGGGGCTCCGCTG-3') (SEQ ID NO:28)

Old6 heavy chain specific primer Rv

(5'-AGAGAGAGAGGCTAGCTGAGGAGACGGTGACCAGGGTTC-3') (SEQ ID NO:29)

5 **[0182]** For the amplification of the heavy chain of mouse antibody 7G3, UPM (SMART RACE cDNA amplification Kit; manufactured by Clontech) and mH-Rv1 primer (5'-ATTTTG TCG ACC KYG GTS YTG CTG GCY GGGTG-3') (SEQ ID NO:30) were used and a cycle of reactions at 98°C 1 second and 68°C 30 seconds was repeated 30 times. Further, using 1 µl of this reaction solution as a template and using NUP (SMART RACE cDNA Amplification Kit; manufactured by Clontech) and mH-Rv2 primer (5'-GCACACYRCTGGACAGGGATCCAGAGTTCC-3') (SEQ ID NO:31), a cycle of reactions at 98°C 1 second and 68°C 30 seconds was repeated 20 times. Thereafter, the amplified PCR products were purified using PCR purification kit (QIAGEN), and the nucleotide sequence of heavy chain variable region was determined by using mH-Rv2 primer (SEQ ID NO:31) as a primer. The following specific primers were synthesized based on sequence information, and the sequences were also determined from the opposite direction using the following primers.

7G3 heavy chain specific primer Fw

15 (5'-AGAGAGAGAGGTTCGACCCACCATGGGATGGAGCTGGATCTTTCTC-3') (SEQ ID NO:32)

7G3 heavy chain specific primer Rv

(5'-AGAGAGAGAGGCTAGCTGCAGAGACAGTGACCAGAGTCCC-3') (SEQ ID NO:33)

20 **[0183]** PCR was carried out using the above-mentioned specific primers (98°C 1 second, 60°C 30 seconds, 72°C 30 seconds), and heavy chain amplification cDNA fragment was digested with *Sall* and *NheI* and inserted into a N5KG1-Val Lark vector [a modified vector of NSKG1 (US6,001,358, Idec Pharmaceuticals)] which had been cleaved with the same enzymes. By determining the sequence using the vector as a template, it was found that the inserted sequence is identical to the one determined by direct sequence.

25 **[0184]** The light chain was amplified using UPM (SMART RACE cDNA Amplification Kit; manufactured by Clontech) and hk-2 primer (5'-GTT GAAGCT CTT TGT GAC GGG CGA GC-3') (SEQ ID NO:34) and repeating a cycle of reactions at 98°C 1 second and 68°C 30 seconds 30 times. Further, using 1 µl of this reaction solution as a template and using NUP (SMART RACE cDNA Amplification Kit; manufactured by Clontech) and hk-6 (5'-TGGCGGGAAGATGAAGACAGATGGT G-3') (SEQ ID NO:35), a cycle of reactions at 98°C 1 second and 68°C 30 seconds was repeated 20 times. Thereafter, the amplified PCR products were purified using PCR purification kit (QIAGEN), and the nucleotide sequence was determined using hk-6 primer. The following specific primers were synthesized based on sequence information, and the sequences were determined also from the opposite direction.

Old4 light chain specific primer Fw

(5'-AGAGAGAGAGATCTCTCACCATGGACATGAGGGTCC CCG CTC AGC -3') (SEQ ID NO:36)

Old4 light chain specific primer Rv

(5'-AGAGAGAGAGCGTACGTTTGATCTCCAGCTTGGTCC CCT G -3') (SEQ ID NO:37)

35 Old5 light chain specific primer Fw

(5'-AGA GAGAGAGATCTCTCACCATGGACATGAGGGTCCCCG CTC AGC -3') (SEQ ID NO:38)

Old5 light chain specific primer Rv

(5'-AGAGAGAGAGCGTACGTTTGATCTCCAGCTTGGTCC CCT G -3') (SEQ ID NO:39)

Old17 light chain specific primer Fw

40 (5'-AGAGAGAGAGATCTCTCACCATGGACATGAGGGTCC TCG CTC AG -3') (SEQ ID NO:40)

Old 17 light chain specific primer Rv

(5'-AGAGAGAGAGCGTACGTTTGATCTCCAGCTTGGTCC CCT G -3') (SEQ ID NO:41)

Old19 light chain specific primer Fw

(5'-AGAGAGAGAGATCTCTCACCATGGACATGAGGGTCCCTCGCTCAG-3') (SEQ ID NO:42)

45 Old 19 light chain specific primer Rv

(5'-AGAGAGAGAGCGTACGTTTGATTTCACCTTGGTCC CTT GGC -3') (SEQ ID NO:43)

New102 light chain specific primer Fw

(5'-AGAGAGAGAGATCTCTCACCATGGACATGAGGGTCC TCG CTC AG -3') (SEQ ID NO:44)

New102 light chain specific primer Rv

50 (5'-AGAGAGAGAGCGTACGTTTGATCTCCAGCTTGG TCC CCT G -3') (SEQ ID NO:45)

Old6 light chain specific primer Fw

(5'-AGAGAGAGAGATCTCTCACCATGGACATGAGGGTCCCCGCTCAGC-3') (SEQ ID NO:46)

Old6 light chain specific primer Rv

(5'-AGAGAGAGAGCGTACGTTTGATATCCACTTTGGTCCCAGGGC-3') (SEQ ID NO:47)

55 **[0185]** Light chain of the mouse antibody 7G3 was amplified using UPM (SMART RACE cDNA amplification Kit; manufactured by Clontech) and mK-Rv1 primer mK_Rv1 (5'-TT GAA GCT CTT GAC AAT GGG TGAAGT TGAT-3') (SEQ ID NO:48) and repeating a cycle of reactions at 98°C 1 second and 68°C 30 seconds 30 times. Further, using 1 µl of this reaction solution as a template and using NUP (SMART RACE cDNA Amplification Kit; manufactured by

Clontech) and mK-Rv2 (5'-GTAGGTGCTGTCTTTGCTGTCCTGATCAGT-3') (SEQ ID NO:49), a cycle of reactions at 98°C 1 second and 68°C 30 seconds was repeated 20 times. Thereafter, the amplified PCR products were purified using PCR purification kit (QIAGEN), and the nucleotide sequence was determined using mK-Rv2 primer. The following specific primers were synthesized based on sequence information, and the sequences were also determined from the opposite direction.

7G3 light chain specific primer Fw

(5'-AGAGAGAGAGAGATCTCACCATGGAATCACAGACTCAGGTCCTC-3') (SEQ ID NO:50)

7G3 light chain specific primer Rv

(5'-AGAGAGAGAGCGTACGTTTTATTTCCAGCTTGGTCCCCC-3') (SEQ ID NO:51)

[0186] PCR was carried out using the above-mentioned specific primers (98°C 1 second, 60°C 30 seconds, 72°C 30 seconds), and a light chain amplification cDNA fragment was digested with *Bgl*II and *Bs*WI and inserted into a N5KG1-Val Lark vector which had been cleaved with the same enzymes. By determining the sequence using the vector as a template, it was found that the inserted sequence is identical to the one determined by direct sequence.

[0187] Each of DNA molecules encoding the heavy chain variable region and light chain variable region of Old4 and amino acid sequences of the heavy chain variable region and light chain variable region are shown in the following.

<Old4 heavy chain variable region>

[0188]

GACCCGTCGACCACCAATGGACTGGACCTGGAGGTTCTCTTTGTGGTGGCAG
CAGCTACAGGTGTCCAGTCCCAGGTCCAGCTGCTACAGTCTGGGGCTGAGGT
GAAGAAGCCTGGGTCCTCGGTGAAGGTCTCATGCAAGGCTTCTGGAGGCAC
CTTCAGCACCTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGGCTT
GAGTGGATGGGAGGGATCATCCCTATCTTTGGTATAGTAACTACGCACAGAA
GTTCCAGGGCAGAGTCACGATTACCGCGGACGAATCCACGAGTACAGCCTAC
ATGGAAGTGAAGCAGCCTGAGATCTGAGGACACGGCCGTGTATTATTGTGCGA
GAGGGGGGGGCTCGGGCCCAGATGTTCTTGATATCTGGGGCCAAGGGACAAT
GGTCACCGTCTCTTCAGCTAGCACCAA (SEQ ID NO:52)

<Old4 heavy chain variable region>

[0189]

MDWTWRFLFVVAATGVQSQVQLLQSGAEVKKPGSSVKVSKASGGTFSTYAI
SWVRQAPGQGLEWMGGIPIFGIVNYAQKFQGRVTITADESTSTAYMELSSLRSE
DTAVYYCARGGGSGPDVLDIWGQGMVTVSSASTX (SEQ ID NO:53)

[0190] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 16 from the 5'-terminal of SEQ ID NO:52, and a boundary between the antibody variable region and the constant region is located between adenine (A) at position 432 and guanine (G) at position 433 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to serine (S) residue at position 139 from the N-terminal of SEQ ID NO:53, and the constant region is on and after alanine (A) at position 140. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of heavy chain was estimated to be up to serine (S) at position 19 from the N-terminal of SEQ ID NO:53. The N-terminal of the mature form was considered to be glutamine (Q) at position 20 of SEQ ID NO:53.

<Old4 light chain variable region>

[0191]

CACAGATCTCTCACCATGGACATGAGGGTCCCCGCTCAGCTCCTGGGGCTCC
 TGCTGCTCTGGCTCCCAGGTGCCAGATGTGTCATCTGGATGACCCAGTCTCCA
 5 TCCTTACTCTCTGCATCTACAGGAGACAGAGTCACCATCAGTTGTCTGGATGA
 GTCAGGGCATTAGGAGTTATTTAGCCTGGTATCAGCAAAAACCAGGGAAAGC
 CCCTGAGCTCCTGATCTATGCTGCATCCACTTTGCAAAGTGGGGTCCCATCAA
 10 GGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCT
 GCAGTCTGAAGATTTTGCAACTTATTACTGTCAACAGTATTATAGTTTCCCGTA
 CACTTTTGGCCAGGGGACCAAGCTGGAGATCAAACGTACGGTGG (SEQ ID
 NO:54)

<Old4 light chain variable region>

[0192]

MDMRVPAQLLGLLLLWLPGARCVIWMQSPSLLSASTGDRVTISCRMSQGIRSY
 LAWYQQKPGKAPELLIYAASLTQSGVPSRFSGSGSGTDFTLTISLQSEDFATYYC
 20 QQYYSFPTYTFGQGTKLEIKRTVX (SEQ ID NO:55)

[0193] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position 16 from the 5'-terminal of SEQ ID NO:54, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 402 and cytosine (C) at position 403 from the 5'-terminal. In the light chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 129 from the N-terminal of SEQ ID NO:55, and the constant region is on and after arginine (R) at position 130. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of light chain was estimated to be up to cysteine (C) at position 22 from the N-terminal of SEQ ID NO:55. The N-terminal of the mature form was considered to be valine (V) at position 23 of SEQ ID NO:55.

[0194] Each of DNA molecules encoding the heavy chain variable region and the light chain variable region of Old5 and amino acid sequences of the heavy chain variable region and light chain variable region was shown in the following.

<Old5 heavy chain variable region>

[0195]

GTCGACCACCATGGACTGGACCTGGAGGTTCTCTTTGTGGTGGCAGCAGCT
 ACAGGTGTCCAGTCCCAGGTCCAGCTGGTGCAGTCTGGGGCTGAGGTGAAG
 45 AAGCCTGGGTCTCGGTGAAGGTCTCATGCAAGGCTTCTGGAGGCACCTTCA
 GCACCTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGGCTTGAGTG
 GATGGGAGGGCTCATCCCTATCTTTGATATAGAAAACCTACGCACAGAAGTTCC
 AGGGCAGAGTCACGATTACCGCGGACGAATCCACGAGCACAGTCTATATGGA
 50 ACTGAGCAGCCTGAGATCTGAGGACACGGCCATGTATTACTGTGCGAGAGGG
 GGGGGTTTCGGGCCCTGATGTTCTTGATATCTGGGGCCAAGGGACAATGGTCA
 CCGTCTCTTCAAGCTAGC (SEQ ID NO:56)

<Old5 heavy chain variable region>

[0196]

MDWTWRFLFVVAATGVQSQVQLVQSGAEVKKPGSSVKVSCKASGGTFSTYAI
 SWVRQAPGQGLEWMGGLIPFDIENYAQKFQGRVTITADESTSTVYMELSSLRSE
 5 DTAMYYCARGGGSGPDVLDIWGQGTMTVTVSSAS (SEQ ID NO:57)

[0197] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 16 from the 5'-terminal of SEQ ID NO:56, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 427 and guanine (G) at position 428 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to serine (S) residue at position 139 from the N-terminal of SEQ ID NO:57, and the constant region is on and after alanine (A) at position 140. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of heavy chain was estimated to be up to serine (S) at position 19 from the N-terminal of SEQ ID NO:57. The N-terminal of the mature form was considered to be glutamine (Q) at position 20 of SEQ ID NO:57.

<Old5 light chain variable region>

[0198]

CACAGATCTCTCACCATGGACATGAGGGTCCCCGCTCAGCTCCTGGGGCTCC
 TGCTGCTCTGGCTCCCAGGTGCCAGATGTGTCATCTGGATGACCCAGTCTCCA
 TCCTTACTCTCTGCATCTACAGGAGACAGAGTCACCATCAGTTGTCTGGATGA
 25 GTCAGGGCATTAGGAGTTATTTAGCCTGGTATCAGCAAAAACCAGGGAAAGC
 CCCTGAGCTCCTGATCTATGCTGCATCCACTTTGCAAAGTGGGGTCCCATCAA
 GGTTCAAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCT
 30 GCAGTCTGAAGATTTTGCAACTTATTACTGTCAACAGTATTATAGTTTCCCGTA
 CACTTTTGGCCAGGGGACCAAGCTGGAGATCAAACGTACGGTGG (SEQ ID
 NO:58)

<Old5 light chain variable region>

[0199]

MDMRVPAQLLGLLLLWLPGARCVIWMQTQSPSLLSASTGDRVTISCRMSQGIRSY
 40 LAWYQQKPGKAPELLIYAASLTQSGVPSRFSGSGSGTDFTLTISLQSEDFATYYC
 QQYYSFPTYTFGQGTKLEIKRTVX (SEQ ID NO:59)

[0200] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position 16 from the 5'-terminal of SEQ ID NO:58, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 402 and cytosine (C) at position 403 from the 5'-terminal. In the light chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 129 from the N-terminal of SEQ ID NO:59, and the constant region is on and after arginine (R) at position 130. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of light chain was estimated to be up to cysteine (C) at position 22 from the N-terminal of SEQ ID NO:59. The N-terminal of the mature form was considered to be valine (V) at position 23 of SEQ ID NO:59.

[0201] Each of DNA molecules encoding the heavy chain variable region and the light chain variable region of Old17 and the amino acid sequences of the heavy chain variable region and the light chain variable region are shown in the following.

<Old17 heavy chain variable region>

[0202]

5 GACCCGTCGACCACCAATGGACTGGACCTGGAGGTTCTCTTTGTGGTGGCAG
 CAGCTACAGGTGTCCAGTCCCAGGTCCAGCTGGTGCAGTCTGGGGCTGAGG
 10 TGAAGAAGCCTGGGTCCTCGGTGAAGGTCTCCTGCAAGACTTCTGGAGGCA
 CCTTCAGCAACTTTGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGGCT
 TGAGTGGATGGGAGGGATCATCCCTATCTTTGGTTCAACAACTACGCACAG
 15 AAGTTCCAGGGCAGAGTCACGATTAACGCGGACGAATCCACGAGCACAGCC
 TACATGGAGCTGAGCAGTCTGAGATCTGAGGACACGGCCGTGTATTACTGTG
 CGGGTGGAGACAAATATGGTCCTTACTACTTTCACTACTGGGGCCAGGGAAC
 CTTGTACACCGTCTCCTCAGCTAGC (SEQ ID NO:60)

<Old17 heavy chain variable region>

[0203]

25 MDWTWRFLFVVAATGVQSQVQLVQSGAEVKKPGSSVKVSKTSGGTFSNFAI
 SWVRQAPGQGLEWMGGIPIFGSTNYAQKFQGRVTINADESTSTAYMELSSLRSE
 30 DTAVYYCAGGDKYGPYYFHYWGQGLVTVSSAS (SEQ ID NO:61)

[0204] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 16 from the 5'-terminal of SEQ ID NO:60, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 432 and guanine (G) at position 433 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to serine (S) residue at position 139 from the N-terminal of SEQ ID NO:61, and the constant region is on and after alanine (A) at position 140. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of heavy chain was estimated to be up to serine (S) at position 19 from the N-terminal of SEQ ID NO:61. The N-terminal of the mature form was considered to be glutamine (Q) at position 20 of SEQ ID NO:61.

<Old17 light chain variable region>

[0205]

45 AGATCTCTCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGC
 TGCTCTGTTTCCCAGGTGCCAGATGTGACATCCAGATGACCCAGTCTCCATCC
 TCACTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGTCGGGCGAGTC
 50 AGGGTATTAGCAGCTGGTTAGCCTGGTATCAGCAGAAACCAGAGAAAGCCCC
 TAAGTCCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGT
 TCAGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCA
 GCCTGAAGATTTTGCAACTTATTACTGCCAACAGTATAATAGTTACCCGTACA
 55 CTTTTGGCCAGGGGACCAAGCTGGAGATCAAACGTACGGT (SEQ ID NO:62)

<Old17 light chain variable region>

[0206]

MDMRVLAQLLGLLLLCFPGARCDIQMTQSPSSLSASVGDRVITITCRASQGISSW
LAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYC
QQYNSYPYTFGQGGTKLEIKRTX (SEQ ID NO:63)

[0207] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position 19 from the 5'-terminal of SEQ ID NO:62, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 399 and cytosine (C) at position 400 from the 5'-terminal. In the light chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 129 from the N-terminal of SEQ ID NO:63, and the constant region is on and after arginine (R) at position 130. By a gene sequence estimation software (Signal P ver. 2), it was estimated that the signal sequence of light chain is up to cysteine (C) at position 22 from the N-terminal of SEQ ID NO:63. It is considered that the N-terminal of the mature form is aspartic acid (D) at position 23 of SEQ ID NO:63.

[0208] Each of DNA molecules encoding the heavy chain variable region and the light chain variable region of Old19 and the amino acid sequences of the heavy chain variable region and light chain variable region was shown in the following.

<Old19 heavy chain variable region>

[0209]

TCGACCCCATGGACTGGACCTGGAGGTTCTCTTTGTGGTGGCAGCAGCTAC
AGGTGTCCAGTCCCAGGTCCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAA
GCCTGGGTCCTCGGTGAAGGTCTCCTGCAAGGCTTCTGGAGGCACCTTCAGC
AGCTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGGCTTGAGTGGG
TGGGAGGGATCATCCCTATCTTTGGTACAGCAAACCTACGCACAGAAGTTCCA
GGGCAGAGTCACGATTACCGCGGACGAATCCACGAGCACAGCCTACATGGA
GCTGAGCAGCCTGAGATCTGAGGACACGGCCGTGTATTACTGTGCGAGAGG
ACACAAATATGGCCCCTACTACTTTGACTACTGGGGCCAGGGAACCCTGGTC
ACCGTCTCCTCAGCTAGCACCAAG (SEQ ID NO:64)

<Old19 heavy chain variable region>

[0210]

MDWTWRFLFVVAATGVQSQVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAI
SWVRQAPGQGLEWVGGIPIFGTANYAQKFQGRVTITADESTSTAYMELSSLRSE
DTAVYYCARGHKYGPYYFDYWGQGLTVTVSSASTK (SEQ ID NO:65)

[0211] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 9 from the 5'-terminal of SEQ ID NO:64, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 425 and guanine (G) at position 426 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to serine (S) residue at position 139 from the N-terminal of SEQ ID NO:65, and the constant region is on and after alanine (A) at position 140. By a gene sequence estimation software (Signal P ver. 2), it was estimated that the signal sequence of heavy chain is up to serine (S) at position 19 from the N-terminal of SEQ ID NO:65. It is considered that the N-terminal of the mature form is the glutamine (Q) at position 20 of SEQ ID NO:65.

<Old19 light chain variable region>

[0212]

5 AGATCTCTCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGC
 TGCTCTGTTTCCCAGGTGCCAGATGTGACATCCAGATGACCCAGTCTCCATCC
 TCACTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGTTCGGGCGAGTC
 10 AGGGTATTAGCAGCTGGTTAGCCTGGTATCAGCAGAAACCAGAGAAAGCCCC
 TAAGTCCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGT
 TCAGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCA
 15 GCCTGAAGATTTTGCAACTTATTACTGCCAACAGTATAATAGTTACCCTCGGA
 CGTTCGGCCAAGGGACCAAGGTGGAAATCAAACGTACGGTGGCT (SEQ ID
 NO:66)

20 <Old19 light chain variable region>

[0213]

25 MDMRVLAQLLGLLLLCFPGARCDIQMTQSPSSLSASVGDRVITICRASQGISSW
 LAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYC
 QQYNSYPRTFGQGTKVEIKRTVA (SEQ ID NO:67)

30 [0214] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position
 13 from the 5'-terminal of SEQ ID NO:66, and a boundary between the variable region and the constant region of the
 antibody is located between adenine (A) at position 399 and cytosine (C) at position 400 from the 5'-terminal. In the light
 chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 129 from the N-terminal
 of SEQ ID NO:67, and the constant region is on and after arginine (R) at position 130. By a gene sequence estimation
 35 software (Signal P ver. 2), it was estimated that the signal sequence of light chain is up to cysteine (C) at position 22
 from the N-terminal of SEQ ID NO:67. It is considered that the N-terminal of the mature form is aspartic acid (D) at
 position 23 of SEQ ID NO:67.

[0215] Each of DNA molecules encoding the heavy chain variable region and the light chain variable region of New102
 and the amino acid sequences of the heavy chain variable region and light chain variable region was shown in the following.

40

<New102 heavy chain variable region>

[0216]

45 TCGACCACCAATGGACTGGACCTGGAGGTTCTCTTTGTGGTGGCAGCAGCTA
 CAGGTGTCCAGTCCCAGGTCCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGA

50

55

AGCCTGGATCCTCGGTGAAGGTCTCCTGCATGGCTTCAGGAGGCACCGTCAG
 CAGCTACGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGGCTTGAGTG
 5 GATGGGAGAGATCATCCCTATCTTTGGTATAGTAAACTACGCACAGAAGTTCC
 AGGGCAGAGTCACGATTACCGCGGACGAATCCACGAACACAGCCTACATGG
 AGCTGAGCAGCCTGAGATCTGAGGACACGGCCATATATTACTGTGCGAGAGA
 10 GACAGCAGTGGCTGGTATTCTTGGTTACTGGGGCCAGGGAACCCTGGTCACC
 GTCTCCTCAGCTAGCACCAAG (SEQ ID NO:68)

<New102 heavy chain variable region>

[0217]

MDWTWRFLFVVAATGVQSQVQLVQSGAEVKKPGSSVKVSCMASGGTVSSYA
 ISWVRQAPGQGLEWMGEIPIFGIVNYAQKFQGRVTITADESTNTAYMELSSLRS
 20 EDTAIYYCARETAVAGILGYWGQGLTVTVSSASTK (SEQ ID NO:69)

[0218] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 9 from the 5'-terminal of SEQ ID NO:68, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 423 and guanine (G) at position 424 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to serine (S) residue at position 138 from the N-terminal of SEQ ID NO:69, and the constant region is on and after alanine (A) at position 139. By a gene sequence estimation software (Signal P ver. 2), it was estimated that the signal sequence of heavy chain is up to serine (S) at position 19 from the N-terminal of SEQ ID NO:69. It is considered that the N-terminal of the mature form is glutamine (Q) at position 20 of SEQ ID NO:69.

<New102 light chain variable region>

[0219]

AGATCTCTCACCATGGACATGAGGGTCCTCGCTCAGCTCCTGGGGCTCCTGC
 TGCTCTGTTTCCCAGGTGCCAGATGTGACATCCAGATGACCCAGTCTCCATCC
 40 TCACTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGTCGGGCGAGTC
 AGGGTATTAGCAGCTGGTTAGCCTGGTATCAGCAGAAACCAGAGAAAGCCCC
 TAAGTCCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCAAGGT
 TCAGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCA
 45 GCCTGAAGATTTTGCAACTTATTACTGCCAACAGTATAATAGTTACCCGTACA
 CTTTGGCCAGGGGACCAAGCTGGAGATCAAACGTACGGTGGCTGCA (SEQ
 ID NO:70)

<New102 light chain variable region>

[0220]

MDMRVLAQLLGLLLLCFPGARCDIQMTQSPSSLSASVGDRVITICRASQGISSW
 LAWYQQKPEKAPKSLIYAASSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYC
 QQYNSYPYTFGQGTKLEIKRTVAA (SEQ ID NO:71)

[0221] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position 13 from the 5'-terminal of SEQ ID NO:70, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 399 and cytosine (C) at position 400 from the 5'-terminal. In the light chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 129 from the N-terminal of SEQ ID NO:71, and the constant region is on and after arginine (R) at position 130. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of light chain was estimated to be up to cysteine (C) at position 22 from the N-terminal of SEQ ID NO:71. The N-terminal of the mature form was considered to be aspartic acid (D) at position 23 of SEQ ID NO:71.

[0222] Each of DNA molecules encoding the heavy chain variable region and the light chain variable region of Old6 and amino acid sequences of the heavy chain variable region and the light chain variable region was shown in the following.

<Old6 heavy chain variable region>

[0223]

CGACCCACCATGGAAGTGGGGCTCCGCTGGGTTTTCTTGTTGCTATTTTAGA
 AGGTGTCCAGTGTGAGGTGCAGTTGGTGGAGTCTGGGGGAGGCCTGGTCAA
 GCCTGGGGGGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTCAGT
 AGCCATAACATGAACTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGG
 GTCTCATCCATTAGTAGTAGTAGTTACATATATTATGCAGACTCAGTGAAG
 GGCCGATTCACCATCTCCAGAGACAACGCCAAGAACTCACTGTATCTGCAAA
 TGAACAGCCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGG
 ACTGGGGCTACTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTC
 AGCTAGC (SEQ ID NO:72)

<Old6 heavy chain variable region>

[0224]

MELGLRWVFLVAILEGVQCEVQLVESGGGLVKPGGSLRLSCAASGFTFSSHNMN
 WVRQAPGKGLEWVSSISSSSSYIYYADSVKGRFTISRDNKNSLYLQMNSLRAE
 DTAVYYCAREDWGYFDYWGQGLVTVSSASTK (SEQ ID NO:73)

[0225] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 10 from the 5'-terminal of SEQ ID NO:72, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 417 and guanine (G) at position 418 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to serine (S) residue at position 136 from the N-terminal of SEQ ID NO:73, and the constant region is on and after alanine (A) at position 137. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of heavy chain was estimated to be up to cysteine (C) at position 19 from the N-terminal of SEQ ID NO:73. The N-terminal of the mature form was considered to be glutamic acid (E) at position 20 of SEQ ID NO:73.

<Old6 light chain variable region>

[0226]

AGATCTCTCACCATGGACATGAGGGTCCCCGCTCAGCTCCTGGGGCTTCTGC
 TGCTCTGGCTCCCAGGTGCCAGATGTGCCATCCAGTTGACCCAGTCTCCATCC
 5 TCCCTGTCTGCATCTGTAGGAGACAGAGTCACCATCACTTGCCGGGCAAGTC
 AGGGCATTAGCAGTGATTTAGCCTGGTATCAGCAGAAACCAGGGAAAGCTCC
 TAAGCTCCTGATCTATGATGCCTCCAGTTTGGAAAGTGGGGTCCCATCAAGGT
 10 TCAGCGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCA
 GCCTGAAGATTTTGCAACTTATTACTGTCAACAGTTTAATAGTTACCCATTAC
 TTTCGGCCCTGGGACCAAAGTGGATATCAAACGTACGGT (SEQ ID NO:74)

<Old6 light chain variable region>

[0227]

MDMRVPAQLLGLLLLWLPGARCAIQLTQSPSSLSASVGDRVTTITCRASQGISSDL
 AWYQQKPGKAPKLLIYDASSLESGVPSRFSGSGSGTDFTLTISLQPEDFATYYC
 QQFNSYPFTFGPGTKVDIKRTVAA (SEQ ID NO:75)

[0228] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position 13 from the 5'-terminal of SEQ ID NO:74, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 399 and cytosine (C) at position 400 from the 5'-terminal. In the light chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 129 from the N-terminal of SEQ ID NO:75, and the constant region is on and after arginine (R) at position 130. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of light chain was estimated to be up to cysteine (C) at position 23 from the N-terminal of SEQ ID NO:75. The N-terminal of the mature form was considered to be alanine (A) at position 24 of SEQ ID NO:75.

[0229] Each of DNA molecules encoding the heavy chain variable region and the light chain variable region of 7G3 and the amino acid sequences of the heavy chain variable region and light chain variable region was shown in the following.

<7G3 heavy chain variable region>

[0230]

GTCGACCACCATGGGATGGAGCTGGATCTTTCTCTTTCTCGTGTCAGGAACT
 GGAGGTGTCCTCTCTGAGGTCCAGCTGCAACAGTCTGGACCTGAGCTGGTG
 AAGCCTGGGGCTTCAGTAAAGATGTCCTGCAAGGCTTCTGGATACACCTTCA
 45 CTGACTACTACATGAAGTGGGTGAAACAGAGCCATGGAAAGAGCCTTGAGT
 GGATTGGAGATATTATTCCTAGCAATGGTGCCACTTTCTACAACCAGAAGTTC
 50 AAGGGCAAGGCCACTTTGACTGTGGACAGATCCTCCAGCACAGCCTACATGC
 ACCTCAACAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTACAAGATCG
 CATTTACTGCGGGCCTCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCA
 55 CTGTCTCTGCAGCTAGC (SEQ ID NO:76)

<7G3 heavy chain variable region>

[0231]

5 MGWSWIFLFLVSGTGGVLSEVQLQQSGPELVKPGASVKMSCKASGYTFTDYY
 MKWVKQSHGKSLEWIGDIIPSNQATFYNQKFKGKATLTVDRSSSTAYMHLNSLT
 10 SEDSAVYYCTRSHLLRASWFAYWGQGLTVTSAAS (SEQ ID NO:77)

[0232] The translation initiation site of the heavy chain DNA is an ATG codon which starts from adenine (A) at position 16 from the 5'-terminal of SEQ ID NO:76, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 427 and guanine (G) at position 428 from the 5'-terminal. In the heavy chain amino acid sequence, the heavy chain variable region is up to alanine (A) residue at position 139 from the N-terminal of SEQ ID NO:77, and the constant region is on and after alanine (A) at position 140. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of heavy chain was estimated to be up to serine (S) at position 19 from the N-terminal of SEQ ID NO:77. The N-terminal of the mature form was considered to be glutamic acid (E) at position 20 of SEQ ID NO:77.

<7G3 light chain variable region>

[0233]

25 AGATCTCACCATGGAATCACAGACTCAGGTCCTCATGTCCCTGCTGTTCTGGG
 TATCTGGTACCTGTGGGGACTTTGTGATGACACAGTCTCCATCCTCCCTGACT
 GTGACAGCAGGAGAGAAGGTCAGTATGAGCTGCAAGTCTAGTCAGAGTCTG
 30 TTAAACAGTGGAAATCAAAAGAACTACTTGACCTGGTATCTGCAGAAACCAG
 GGCAGCCTCCTAAATTGTTGATCTATTGGGCATCCACTAGGGAATCTGGGGTC
 CCTGATCGCTTCACAGGCAGTGGATCTGGAACAGATTTCACTCTCACCATCA
 GCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGTCAGAATGATTATAGT
 35 TATCCGTACACGTTTCGGAGGGGGGACCAAGCTGGAAATAAAACGT (SEQ ID
 NO:78)

<7G3 light chain variable region>

[0234]

45 MESQTQVLMSLLFWVSGTCGDFVMTQSPSSLTVTAGEKVTMSCKSSQSLLNSG
 NQKNYLWYLYKPGQPPKLLIYWASTRESGVDPDRFTGSGSGTDFTLTISVQAE
 DLAVYYCQNDYSYPYTFGGGTKLEIKR (SEQ ID NO:79)

[0235] The translation initiation site of the light chain DNA is an ATG codon which starts from adenine (A) at position 11 from the 5'-terminal of SEQ ID NO:78, and a boundary between the variable region and the constant region of the antibody is located between adenine (A) at position 409 and cytosine (C) at position 410 from the 5'-terminal. In the light chain amino acid sequence, the light chain variable region is up to lysine (K) residue at position 133 from the N-terminal of SEQ ID NO:79, and the constant region is on and after arginine (R) at position 134. By a gene sequence estimation software (Signal P ver. 2), the signal sequence of light chain was estimated to be up to the glycine (G) at position 22 from the N-terminal of SEQ ID NO:79. The N-terminal of the mature form was considered to be aspartic acid (D) at position 22 of SEQ ID NO:79.

(Preparation of recombinant type antibodies)

[0236] Cells expressing a recombinant antibody were prepared by introducing each of the constructed six recombinant antibody expression vectors into a host cell. HEK293F (Invitrogen) was used as a host cell for expression.

[0237] Each expression vector was introduced into HEK293F using 293Fectin (Invitrogen). The HEK293F was cultured under conditions of 5% CO₂ and 37°C using a shaker, and the culture supernatant was recovered about 5 days after culturing. The recovered culture supernatant was subjected to affinity purification using rmp Protein A (Amersham-Pharmacia Biotech). PBS as an adsorption buffer and 0.02 M of glycine buffer (pH 3) as an elution buffer, using 0.8×40 cm column (Bio-Rad Laboratories) and the like depending on the amount for the purification. The elution fraction was adjusted to about pH 7.2 by adding 1 M of Tris (pH 9.0). The thus prepared antibody solution was substituted to PBS using a dialysis membrane (10,000 cut, Spectrum Laboratories) and subjected to sterilization by filtration using a membrane filter MILLEX-GV having a pore size of 0.22 μm (Millipore), thereby obtaining a purified human anti-IL-3R α monoclonal antibody. The concentration of the purified antibody was calculated by measuring absorbance at 280 nm and regarding 1 mg/ml as 1.4 OD.

[0238] A summary of amino acid sequences and SEQ ID NOs of each human antibody CDR (complementarity-determining region) was shown in Table 1.

[Table 1]

Heavy chain variable region	SEQ ID NO CDR 1 CDR 2 CDR 3	CDR 1	CDR 2	CDR3
Old4	113 114 115	TYAIS	GIPIFGIVNYAQKFQG	GGGSGPDVLDI
Old5	116 117 118	TYAIS	GLIPIFDIENYAQKFQG	GGGSGPDVLDI
Old17	119 120 121	NFAIS	GIPIFGSTNYAQKFQG	GDKYGPYYFHY
Old19	122 123 124	SYAIS	GIPIFGTANYAQKFQG	GHKYGPYYFDY
New102	125 126 127	SYAIS	EIPIFGIVNYAQKFQG	ETAVAGILGY
Old6	128 129 130	SHNMN	SISSSSYIYYADSVKG	EDWGYFD
Light chain variable region	SEQ ID NO CDR 1, CDR 2, CDR 3	CDR 1	CDR 2	CDR 3
Old4	131, 132, 133	RMSQGIRSYLA	AASLQS	QQYYSFPYT
Old5	134, 135, 136	RMSQGIRSYLA	AASLQS	QQYYSFPYT
Old17	137, 138, 139	RASQGISSWLA	AASLQS	QQYNSYPYT
Old19	140, 141, 142	RASQGISSWLA	AASLQS	QQYNSYPRT
New102	143, 144, 145	RASQGISSWLA	AASLQS	QQYNSYPYT
Old6	146, 147, 148	RASQGISSDLA	DASSLES	QQFNSYPFT

Example 5 Purification of anti-IL-3R α human antibody from hybridoma culture supernatant

[0239] A hybridoma was cultured after adapting from the IL-6-containing DMEM medium used in Example 3 to E-RDF medium (Kyokuto Pharmaceutical Industrial Co., Ltd.). An antibody was purified from the culture supernatant. The purification of the antibody was carried out in accordance with Example 4.

[0240] Firstly, a hybridoma producing a human anti-IL-3R α monoclonal antibody was adapted to eRDF medium (Kyokuto Pharmaceutical Industrial Co., Ltd.) containing 10 ng/ml IL-6 and 10% fetal calf serum (FCS: SIGMA). Next, the obtained hybridoma was adapted to eRDF medium (Kyokuto Pharmaceutical Industrial Co., Ltd.) containing bovine insulin (5 μ g/ml, GIBCO BRL), human transferrin (5 μ g/ml, GIBCO BRL), ethanolamine (0.01 mM, SIGMA), sodium selenite (2.5×10^{-5} mM, SIGMA) and 1% low IgG FCS (HyClone). This adapted hybridoma was cultured in a flask, and the culture supernatant was recovered. The recovered supernatant was subjected to 10 μ m and 0.2 μ m filters (Gelman Sciences Inc.) to remove useless articles such as a hybridoma and the like. The antibody was purified from the thus recovered supernatant by the same method as Example 4.

Example 6 Calculation of association and dissociation constants using the purified anti-IL-3R α human antibody

[0241] The association and dissociation constants of the purified anti-IL-3R α antibody were analyzed using an analyzer which is based on a surface plasmon resonance principal (Biacore, GE Healthcare, hereinafter GE). Briefly, an anti-human antibody or anti-mouse antibody was immobilized on a CM5 sensor tip, then an anti-IL-3R α human or mouse antibody was applied thereto to allow to bind, then the soluble form of IL-3R α protein prepared in Example 2 was applied thereto, and the association and dissociation were observed using Biacore 2000. Through the whole test steps, the test method of GE Healthcare for the calculation of association and dissociation constants was basically employed.

[0242] Specifically, CM5 (research grade) was used for a sensor tip (each GE). Firstly, the CM5 tip was activated by applying an equivalent mixture of 400 mM EDC (N-ethyl-N'-(3-dimethylaminopropyl)carbodiimidehydrochloride) and 100 mM NHS (N-hydroxysuccinimide) to the CM5 tip. Next, an antibody to the human antibody attached to the Human Antibody Capture Kit (GE) (hereinafter, referred to as anti-human antibody antibody) was diluted with the solution attached to the kit and applied thereto, thereby immobilizing the required amount of the anti-human antibody antibody to the CM5 tip. Regarding the mouse antibody to be used as a control, an antibody to the mouse antibody attached to the Mouse Antibody Capture Kit (GE) (hereinafter, referred to as anti-mouse antibody antibody) was diluted with the solution attached to the kit and applied thereto, thereby immobilizing a necessary amount thereof to the CM5 tip. Next, the surface of the activated tip was blocked and inactivated by applying 1 M of ethanolamine dihydrochloride thereto. By the above steps until this, preparation of a CM5 sensor tip which can measure the dissociation constant K_D was completed.

[0243] Next, each of anti-IL-3R α antibodies, , was diluted to give a concentration of 5 μ g/ml with HBS-FP buffer (GE) one kind per one flow cell and applied, thereby allowing it to bind to the immobilized anti-human antibody antibody or anti-mouse antibody antibody. Next, the soluble form of IL-3R α protein was applied thereto. In order to dissociate the bound anti-IL-3R α antibody and soluble form of IL-3R α protein, 3 M MgCl₂ attached to Human Antibody Capture Kit or pH 1.7 of Glycine-HCl attached to Mouse Antibody Capture Kit was applied in the amount attached to the kit. The steps until this were regarded as one step. By repeating the same steps using two or more concentrations of the soluble form of IL-3R α protein, data for calculating association and dissociation constants (sensorgram) were obtained.

[0244] The concentration of the soluble form of human IL-3R α protein applied to a subject was calculated as described in Example 2 by measuring the absorbance at 280 nm and regarding 1 mg/ml as 1.4 OD. The molecular weight of the soluble form of human IL-3R α protein was calculated as follows. Regarding molecular weight of human IL-3R α protein, it has been reported that it comprises 360 amino acid residues, has six N-type sugar chain binding sites and the molecular weight is 70 kDa (The Cytokine Facts Book second edition, Academic Press). Accordingly, the molecular weight of the soluble form of human IL-3R α protein was calculated as about 63 kDa by subtracting molecular weights of the amino acids of the transmembrane region and the intracellular region from 70 kDa known as a reference information and adding, to the resulting value, the molecular weight of the amino acids of the Flag sequence.

[0245] In the analysis, Biaevaluation software (GE) was used and Biaevaluation Software Handbook was referred. Specifically, by carrying out simultaneous analysis of kinetics analysis, employing basically the 1:1 Langmuir binding reaction model and fitting, association rate constant (K_a) and dissociation rate constant (K_d) were calculated, and the value of dissociation constant K_D was calculated by the calculation of K_d/K_a .

[0246] The results are shown in the following table 2.

[Table 2]

Antibody name	K _a	K _d	K _D
Human antibodies			
Old4	3.88×10^5	5.15×10^{-4}	1.33×10^{-9}
Old5	7.17×10^5	4.72×10^{-4}	6.58×10^{-10}
Old7	2.08×10^5	2.98×10^{-4}	1.43×10^{-9}
Old19	1.54×10^5	4.99×10^{-4}	3.24×10^{-9}
New102	6.02×10^5	4.80×10^{-4}	7.98×10^{-10}
Old6	1.71×10^6	2.15×10^{-5}	1.26×10^{-9}
Chimeric antibody			
7G3	2.48×10^5	4.66×10^{-4}	1.88×10^{-9}
Mouse antibodies			
7G3	1.68×10^5	9.52×10^{-5}	5.66×10^{-10}
9F5	7.13×10^4	6.5×10^{-5}	9.11×10^{-10}
107D2.08	4.16×10^5	2.03×10^{-5}	4.88×10^{-8}
AC145	7.66×10^4	4.26×10^{-5}	5.57×10^{-8}
L-16	8.13×10^5	4.16×10^{-5}	5.12×10^{-9}

Example 7 Epitope analysis of anti-human IL-3R α human antibody(Preparation of IL-3R α /GM-CSFR α chimeric protein expression cell)

[0247] In order to carry out epitope analysis of IL-3R α antibody, a chimeric protein in which a portion of the extra-membrane region of IL-3R α was replaced by GM-CSFR α was expressed in a cell, and binding activity of each anti-IL-3R α antibody to the cell was analyzed. In brief, firstly, the IL-3R α molecule and GM-CSFR α molecule were divided into three regions (A, B and C domains from the above-mentioned N-terminal), secondly vectors which express molecules in which each of the A, B and C domains of the IL-3R α molecule was replaced by the corresponding domain of GM-CSFR α molecule were respectively constructed, thirdly, these were forcedly expressed in HEK293F cell, and fourthly, whether or not each anti-IL-3R α antibody labeled with a fluorescence dye binds thereto was observed by flow cytometry.

(Preparation of GM-CSFR/pEF6/Myc-HisC plasmid DNA)

[0248] A cDNA of human GM-CSFR receptor α chain (GM-CSFR α , CD116) was amplified from a spleen-derived cDNA (CLONTECH Human MTC Panel) by a PCR method using KOD-Plus-Ver. 2 (Toyobo Co., Ltd.). As a PCR device, GeneAmp PCR System 9700 (Applied Biosystems) was used. Regarding the PCR, after a denaturation step at 94°C for 2 minutes, a three step reaction at 98°C 10 seconds-55°C 30 seconds-68°C 75 seconds was carried out 35 cycles. The PCR primers used are as follows. hCD116Fw-MfeI:

5'-CGGCAATTGCCACCATGCTTCTCCTGGTGACAAGCCT-3' (SEQ ID NO:80) hCD116Rv-NotI:

5'-ATTGCGGCCGCTCAGGTAATTCCTTACGG-3' (SEQ ID NO:81)

[0249] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (TAE buffer). DNA was visualized by ethidium bromide staining. A band of at around 1.2 kb was cut out, and the DNA was extracted using JETsorb Kit (Genomed, Bad Oeynhausen, Germany) and then digested with *NotI* and *MfeI*. A pEF6/Myc-HisC plasmid DNA (Invitrogen) was digested with *EcoRI* and *NotI*. Each DNA was subjected to 0.8% agarose gel electrophoresis and bands of at around 1.2 kb and around 6 kb were cut out, and the DNA molecules were extracted using JETsorb Kit (Genomed, Bad Oeynhausen, Germany). Then, 0.5 μ l of a pEF6/Myc-HisC plasmid DNA-derived DNA solution and 4 μ l of a PCR product-derived DNA solution were mixed and subjected to ligation using TaKaRa Ligation Kit (TAKARA BIO INC.). Regarding the transformation, a ligation sample and DH5 alpha competent cells were mixed and spread on an LB plate. Insertion check was carried out by colony direct PCR using LA Taq (TAKARA BIO INC.). Regarding the PCR, after a denaturation step at 94°C for 5 minutes, a three step reaction at 94°C 30 seconds-55°C 30 seconds-72°C

2 minutes was carried out 40 cycles and then a treatment at 99°C for 30 minutes was carried out.

[0250] The PCR primers used were as follows.

hCD116Fw-MfeI: 5'-CGGCAATTGCCACCATGCTTCTCCTGGTGACAAGCCT-3' (SEQ ID NO:82)

hCD116Rv-NotI: 5'-ATTGCGGCCGCTCAGGTAATTCCTTCACGG-3' (SEQ ID NO:83)

[0251] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. Using a colony from which amplification of around 1.2 kb was obtained, nucleotide sequence was determined by a direct sequencing method. In the reaction of sequence samples, BigDye(R) Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems) and GeneAmp PCR System 9700 (Applied Biosystems) were used (these were used in the all DNA sequence analyses in this specification). The PCR primers used

are as follows.

hCD116Rv-NotI: 5'-ATTGCGGCCGCTCAGGTAATTCCTTCACGG-3' (SEQ ID NO:85)

hCD116SeqFw1: 5'-TGAAGTGTACCTGGGCGAGG-3' (SEQ ID NO:86)

hCD116SeqFw2: 5'-CTGGCACGGAAACCTACTG-3' (SEQ ID NO:87)

hCD116SeqRv1: 5'-CCTGAATTTGGATAAAGCAG-3' (SEQ ID NO:88)

[0252] ABI 3700XL DNA analyzer (Applied Biosystems) was used as a sequence analyzing device (this was used in the all DNA sequence analyses in this specification). By selecting a clone in which mutation in the amino acid sequence by PCR was not found, plasmid DNA was extracted by Largeprep method (QIAGEN).

(Preparation of IL-3RA-FLAG/pEGFP-N1)

[0253] The full length cDNA of human IL-3R α (CD123) was amplified by PCR and FLAG tag was linked to its downstream (IL-3RA-FLAG/pEGFP-N1).

[0254] Human IL-3RA cDNA was amplified by a PCR method using hCD123/pEGFP-N1 plasmid DNA as a template and LA Taq (TAKARA BIO INC.). Regarding the PCR, after a denaturation step at 95°C for 30 seconds, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 60 seconds was carried out 10 cycles and then 2 minutes of an elongation reaction was carried out. The PCR primers used are as follows.

T7: 5'-TAATACGACTCACTATAGGG -3' (SEQ ID NO:89) hCD123-C-FLAG-R1:

5'-TCGTCATCGTCCTTGTAGTCAGTTTCTGCACGACCTGTA-3' (SEQ ID NO:90)

[0255] Using 2 μ l of the PCR product as a template, amplification was carried out by a PCR method using LA Taq (TAKARA BIO INC.). Regarding the PCR, after a denaturation step at 95°C for 1 minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 60 seconds was carried out 15 cycles and then an elongation reaction at 72°C for 2 minutes was carried out. The PCR primers used are as follows. IL-3R α _Fw:

5'-CGGCAATTGCCACCATGGTCCTCCTTTGGCTCAC-3' (SEQ ID NO:91) C-FLAG-NotR2:

5'-AAAAGCGGCCGCTCACTTGTGTCATCGTCCTTGTAGTC-3' (SEQ ID NO:92)

[0256] The thus obtained PCR products were subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A band of at around 1 kb was cut out, and the DNA was extracted using Wizard SV Gel and PCR Clean-Up System. The whole amount of the extracted DNA was digested with *MfeI* and *NotI* and subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A band of at around 1 kb was cut out, and the DNA was extracted using Wizard SV Gel and PCR Clean-Up System. Then, 5 μ l of the thus extracted IL-3RA-FLAG cDNA and 1 μ l of the pEGFP-N1 plasmid DNA which had been cleaved with *EcoRI* and *NotI* were mixed and ligated using TaKaRa Ligation Kit (TAKARA BIO INC.). Regarding the transformation, a ligation sample and DH10B competent cells were mixed and spread on an LB plate (containing kanamycin). Insertion check was carried out by colony direct PCR using LA Taq (TAKARA BIO INC.). Regarding the PCR, after a denaturation step at 95°C for 1 minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 60 seconds was carried out 35 cycles and then an elongation reaction at 72°C for 2 minutes was carried out. The PCR primers used are as follows.

pEGFP-N1-Fw: 5'-CGTGTACGGTGGGAGGTCTA-3' (SEQ ID NO:93)

pEGFP-N1-Re: 5'-TTTATGTTTCAGGTTTCAGG-3' (SEQ ID NO:94)

[0257] A plasmid DNA was extracted by a Miniprep method from a colony in which amplification of around 0.8 kb was obtained.

[0258] It was found by a DNA sequence analysis that the purified IL-3RA-FLAG/pEGFP-N1 plasmid DNA does not have a mutation caused by the PCR and that the FLAG tag is present therein. The primers used in the DNA sequence analysis are as follows.

pEGFP-N1-Fw: 5'-CGTGTACGGTGGGAGGTCTA-3' (SEQ ID NO:95)

pEGFP-N1-Re: 5'-TTTATGTTTCAGGTTTCAGG-3' (SEQ ID NO:96)

(Domain mapping of IL-3R α)

[0259] As a result of BLASTP search (database: Protein Data Bank proteins (pdb)), IL-13 receptor alpha chain (IL-13R α) was hit with the highest score (PDB: 3BPNC; Chain C, Crystal Structure of the IL4-IL4r-IL13ra Ternary Complex). Using the PDB file down-loaded from Protein Data Bank and a graphic software RasMol, three-dimensional structure of the IL-13R α , protein was visualized and three domains constituting the extracellular region (the above-mentioned A, B and C domains) were divided. Using a Multiple Alignment software MUSCLE, IL-3R α amino acid sequence and IL-13R α amino acid sequence were compared and IL-3R α extracellular region was also divided into three domains. Further, GM-CSFR α and IL-3R α were compared in the same manner and GM-CSFR α extracellular region was also divided into three domains.

[0260] In order to assign epitopes of anti-human IL-3R α human antibodies, proteins in which each of the three domains of IL-3R α was replaced one by one by said domains of GM-CSFR α were prepared and expressed on the cell membrane and the presence or absence of antibody binding was confirmed.

[0261] Using the IL-3RA-FLAG/pEGFP-N1 plasmid DNA as a template, amplification was carried out by a PCR method which uses PrimeSTAR(R) HS DNA Polymerase (TAKARA BIO INC.). Regarding the PCR, a two step reaction at 98°C 10 seconds-68°C 6 minutes was carried out 25 cycles. The PCR primers used are as follows.

A domain deficiency;

CD123R11pEGFPN1: AAAGGTACCGAATTCGAAGCTTGAGCTC (SEQ ID NO:97)

CD123F11: AAAGGTACCGGGAAGCCTTGCGCAGGT (SEQ ID NO:98)

B domain deficiency;

CD123R12-2: AAAGGTACCACTGTTCTCAGGGAAGAGGAT (SEQ ID NO:99)

CD123F12-2: AAAGGTACCCAGATTGAGATATTAAGTCC (SEQ ID NO:100)

C domain deficiency;

CD123R13: AAAGGTACCTGAAAAGACGACAACTT (SEQ ID NO:101)

CD123F13: AAAGGTACCTCGCTGCTGATCGCGCTG (SEQ ID NO:102)

[0262] The thus obtained PCR product was subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. After confirming the amplification, this was purified using Wizard SV Gel and PCR Clean-Up System. The thus obtained DNA was digested with *KpnI* and *DpnI*, purified using Wizard SV Gel and PCR Clean-Up System and ligated using TaKaRa Ligation Kit. Regarding the transformation, a ligation sample and DH10B competent cells were mixed and spread on an LB plate (containing kanamycin). Insert check was carried out by colony direct PCR using LA Taq (TAKARA BIO INC.). Regarding the PCR, after a denaturation step at 95°C for 1 minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 40 seconds was carried out 38 cycles and then an elongation reaction at 72°C for 2 minutes was carried out. The PCR primers used are as follows.

pEGFP-N1-Fw: 5'-CGTGTACGGTGGGAGGTCTA-3' (SEQ ID NO:103)

pEGFP-N1-Re: 5'-TTTATGTTTCAGGTTTCAGG -3' (SEQ ID NO:104)

[0263] The thus obtained PCR product was subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A plasmid DNA was extracted by the Miniprep method from a colony in which amplification at around 1 kb was obtained.

[0264] Using the GM-CSFR/pEF6/Myc-HisC plasmid DNA as a template, amplification was carried out by a PCR method which uses PrimeSTAR(R) HS DNA Polymerase (TAKARA BIO INC.). Regarding the PCR, a two step reaction at 98°C 10 seconds-68°C 30 seconds was carried out 25 cycles. The PCR primers used are as follows.

A domain insertion;

GM-CSFRF11: AAAGGTACCGCCACCATGCTTCTCCTGGTGACA (SEQ ID NO:105)

GM-CSFR11: AAAGGTACCTGAATTTGGATAAAGCAG (SEQ ID NO:106)

B domain insertion;

GM-CSFRF12: AAAGGTACCGGAAGGGAGGGTACCGCT (SEQ ID NO:107)

GM-CSFR12: AAAGGTACCCTTTGTGTCCAAAAGTGA (SEQ ID NO:108)

C domain insertion;

GM-CSFRF13: AAAGGTACCAAAATAGAACGATTCAAC (SEQ ID NO:109)

GM-CSFR13: AAAGGTACCAATGTACACAGAGCCGAG (SEQ ID NO:110)

[0265] The thus obtained PCR product was subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. After confirming the amplification, this was purified using Wizard SV Gel and PCR Clean-Up System.

[0266] The thus obtained DNA was digested with *KpnI* and then purified using QIAquick Gel Extraction Kit (QIAGEN), mixed with IL-3RA-FLAG/pEGFP-N1 plasmid DNA in which the corresponding domain was deleted (already cleaved with *KpnI* and purified) and ligated using TaKaRa Ligation Kit. Regarding the transformation, a ligation sample and DH10B competent cells were mixed and spread on an LB plate (containing kanamycin). Insert check was carried out by colony direct PCR using LA Taq (TAKARA BIO INC.). Regarding the PCR, after a denaturation step at 95°C for 1

minute, a three step reaction at 95°C 15 seconds-56°C 15 seconds-72°C 40 seconds was carried out 38 cycles and then an elongation reaction at 72°C for 2 minutes was carried out. The PCR primers used are as follows.

pEGFP-N1-Fw: 5'-CGTGACGGTGGGAGGTCTA-3' (SEQ ID NO:111)

pEGFP-N1-Re: 5'-TTTATGTTTCAGGTTTCAGG-3' (SEQ ID NO:112)

[0267] The thus obtained PCR product was subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. A plasmid DNA was extracted by the Miniprep method from a colony in which amplification at around 1 kb was obtained.

(Labeling of anti-IL-3R α human antibody with fluorescence dye)

[0268] In order to determine binding activity of anti-human IL-3R α human antibodies, each human antibody was labeled with a fluorescence dye AlexaFluor488 (Molecular Probe, Invitrogen). Regarding the labeling method, it was carried out in accordance with the manual provided by Invitrogen, and regarding the detection, fluorescence was detected by FL1 of a flow cytometry (FACS Calibur, BD Biosciences).

[0269] Specifically, 1/10 volume of 1 M Na₂CO₃ was added to an antibody solution dissolved in PBS. Next, a predetermined amount of the antibody solution described in the manual was added to a container containing powder of AlexaFluor488 to which tetrafluorophenyl (TFP) had been added, and allowed to undergo the reaction in the dark at room temperature for 1 hour while stirring. Next, after a gel filtration column (NAP-10 and the like, GE Healthcare) was sufficiently substituted with PBS, the solution of antibody reacted with AlexaFluor488 was added thereto while the buffer of the antibody solution was substituted with PBS. An antibody fraction which showed yellow green was obtained. Regarding the AlexaFluor488-labeled anti-human IL-3R α antibody obtained in the above manner, the absorbances at wavelengths of 280 nm and 494 nm (A280 and A494, respectively) were measured using a spectrometer, and the antibody concentration was calculated by the following calculation formula.

$$\text{Antibody concentration (mg/ml)} = (A_{280} - A_{494} \times 0.11) / 1.4$$

(Flow cytometry analysis of IL-3R α /GM-CSFR α chimeric protein expression cell using labeled anti-IL-3R α antibody)

[0270] HEK293T cell (ATCC CRL 1268) was used for the preparation of the IL-3R α /GM-CSFR α chimeric protein expression cell. Using 293Fectin (Invitrogen), the plasmid DNA obtained in the above was introduced as an expression vector into the HEK293T. The HEK293T to which the expression vector was introduced was cultured using a shaker under conditions of 5% CO₂ and 37°C after 2 days of the introduction, the obtained protein was used for in the flow cytometry analysis.

[0271] From 100,000 to 1,000,000 cells of the chimeric protein expression cell were allowed to react for 30 minutes on ice with at a concentration of 1 μ g/ml with an AlexaFluor488-labeled human antibody or a commercially available FITC-labeled anti-IL-3R α mouse antibody (7G3 or 9F5: both from BD Biosciences, 6H6: Acris Antibodies, AC 145: Milteny Biotech, 107D2.08: Dendritics). A staining medium (Dulbecco's PBS supplemented with 2% fetal bovine serum, 2 mM EDTA and 0.05% NaN₃) was used for the dilution of antibodies and cells. Next, the cells reacted with antibodies were washed three times with the staining medium and whether the labeled antibody bound to the cell was confirmed by flow cytometry.

[0272] The results are shown in Figs. 1 and 2. Reactions of 7G3, 9F5, 6H6 and AC 145 antibodies disappeared only in the cells expressing a protein in which A domain was replaced by GM-CSFR α . On the other hand, reactions of Old4, Old5, Old19 and New102 antibodies disappeared in the cells expressing a protein in which B domain was replaced by GM-CSFR α . Regarding Old19, its reaction with the cells expressing a protein in which A domain was replaced by GM-CSFR α also disappeared. Regarding the Old6 and 107D2.08, their reaction to B domain- and C domain-substituted protein expression cells disappeared.

[0273] Based on the above, it was shown a possibility that 7G3, 9F5, 6H6 and AC 145 recognized A domain, and Old4, Old5 and New102 recognized B domain, Old19 recognized A domain and B domain, and Old6 and 107D2.08 recognized B domain and C domain. Accordingly, the reactivity of various anti-IL-3R α antibodies to A to C domains of IL-3R α was as the following Table 3.

[Table 3]

	7G3	9F5	6H6	AC 145	107D2.08	Old19	New102	Old4	Old5	Old6	Old28
No substitution	++	++	++	++	++	++	++	++	++	++	++
A domain substitution	-	-	-	-	++	-	++	++	++	++	+
B domain substitution	++	++	++	+	-	-	-	-	-	-	-

(continued)

	7G3	9F5	6H6	AC 145	107D2.08	Old19	New102	Old4	Old5	Old6	Old28
C domain substitution	++	++	++	++	-	++	++	++	++	-	++

Example 8 Analysis of IL-3 signaling blocking activity of anti-IL-3R α antibodies

[0274] In order to examine whether the thus obtained IL-3R α antibodies inhibit IL-3 signaling, a cell line TF-1 (DMSZ no. ACC344) which grows IL-3- or GM-CSF-dependently was used. Specifically, the TF-1 cell was diluted with RPMI 1640 medium containing 1 ng/ml of IL-3 and 10% fetal calf serum (TF-1 medium) and dispensed on a 96-well plate. Further, various IL-3R α antibodies and a human serum-derived IgG as a negative control antibody were diluted with the TF-1 medium, transferred to the 96-well plate and added in such a manner that final concentration of each antibody gave a final concentration of 10 or 100 μ g/ml. As a control, a well of cell-free medium alone and a well to which the TF-1 cell was added were prepared. After 3 days of culturing under an environment of 37°C and 5% CO₂, CelltiterGlo (Promega) was added thereto in an amount equivalent to the medium. After 30 minutes of still standing, the emission was determined using a plate reader (ARBO, Perkin Elmer).

[0275] For the growth inhibition ratio, the following calculation was carried out.

$$\frac{(\text{emission of sample} - \text{well with no cells})}{(\text{well to which TF-1 cell alone was added} - \text{well with no cells})} \times 100 (\%)$$

[0276] Regarding the commercially available antibodies 9F5, 6H6 and 107D2.08, an NAP-5 column was used for the purpose of substituting the buffer with PBS. Specifically, 0.5 ml of an antibody solution was added to the NAP-5 column sufficiently substituted with PBS. Next, by adding 1.0 ml of PBS, the solution discharged from the column was recovered. By carrying out sterilization by filtration using a membrane filter MILLEX-GV (Millipore) having a pore size of 0.22 μ m, an antibody dissolved in PBS as a solvent was obtained.

[0277] The results are shown in Fig. 5. It was found that the antibody Old4, antibody Old5, antibody Old17, antibody Old19, antibody New102, antibody 9F5 and antibody 6H6 did not inhibit the IL-3 signaling, and on the other hand, it was found that the antibody 7G3, antibody Old6 and antibody 107D2.08 inhibited IL-3 signaling. Example 9 Examination of influence upon colony forming ability using anti-IL-3R α human antibody

[0278] A colony assay was carried out to find whether various IL-3R α antibodies have effects upon the colony forming ability by hematopoietic precursor cells.

[0279] In brief, 400 cells/ml of cord blood-derived CD34 positive cell (AllCells) was added to a Methocult medium (Stem Cell Technologie) supplemented with erythropoietin, IL-3, G-CSF and Stem Cell Factor, and the number of colonies was measured 14 days to 16 days thereafter. The colonies were counted by dividing them into a granulocyte/macrophage system colony (CFU-GM), an erythroid system colony (BFU-E) and mixed colonies (CFU-Mix or CFU-GEMM). Regarding the classification method of colony types, a manual provided by Stem Cell Technologie or various textbooks on hematology was used as references.

[0280] As the antibodies, each of the chimeric 7G3 antibody as an antibody in which IL-3 signaling blocking activity was found in Example 8 and the New102 antibody as an antibody in which the blocking activity was not found was used.

[0281] The results are shown in Fig. 6. In the colony assay in which erythropoietin, IL-3, G-CSF and Stem Cell Factor were added, decrease in the number of colonies and decrease in the colony size were found by the addition of the 7G3 antibody which had the blocking activity of IL-3 signaling. On the other hand, change in the number of colonies by the addition of the New102 antibody was not found. Based on this result, it is considered that the influence upon the normal hematopoietic function is small and side effects are few when the IL-3 signaling is not inhibited or blocked.

Example 10 Antitumor effect in a mouse tumor bearing model using anti-IL-3R α , human antibody

[0282] The thus obtained anti-IL-3R α antibody was administered to mouse tumor bearing model and its antitumor effect was examined. In brief, a leukemia cell was transferred into a mouse through its caudal vein, the antibody was administered thereto on the next day, and about 3 weeks thereafter, the number of leukemia cells in bone marrow cells collected from a bone of the mouse was counted.

[0283] Specifically, 0.01 ml equivalent of anti-asialoGM1 antiserum (Wako Pure Chemical Industries, Ltd.) was diluted with physiological saline and administered to SCID mouse (CLEA Japan Inc.) (Day -1). On the next day, 500,000 cells of a cell line of acute myeloid leukemia, MOLM13 (ATCC), were transplanted through caudal vein (Day 0). Further on

the next day (Day 1), 10 μ g of the anti-IL-3R α antibody was intraperitoneally administered. The mouse was sacrificed on Day 21, bone marrow was collected from thighbones and shinbones and the bone marrow cells were stained with FITC-labeled human CD45 antibody and PE-labeled anti-IL-3R α (both from BD Biosciences). Specifically, the antibody was added to about 1,000,000 cells of the bone marrow cell, to give a final concentration of 1 μ g/ml for each and allowed to stand still on ice for 30 minutes under shade. Thereafter, the cells stained with the antibody was washed 3 times using a staining medium (a solution prepared by adding 2% fetal bovine serum, 0.05% sodium azide and 2 mM EDTA to PBS (GIBCO)), and a human CD45 positive and human IL-3R α positive cell was detected by a flow cytometry (FACSCalibur, BD Biosciences). Also, at the time of collecting mouse bone marrow, the number of bone marrow cells was counted using TURK solution. Further, the absolute number of the MOLM13 cells contained in one thighbone was counted by simultaneously adding quantified fluorescent beads (Flow-Count, Beckman Coulter) at the time of the above-mentioned antibody staining.

[0284] The results are shown in Fig. 7. It was found that the number of MOLM13 cells in the thighbone bone marrow in each antibody administered group is markedly reduced in comparison with the Vehicle group to which the antibody was not administered. This result shows that the anti-IL-3R α antibody has a possibility as a therapeutic agent for leukemia.

Example 11 Toxicity test of IL-3R α expression cell by anti-IL-3R α antibody

[0285] In order to measure antibody-mediated cytotoxicity (antibody-dependent cellular cytotoxicity, hereinafter referred to as ADCC), this was carried out in the presence of antibody using a human peripheral blood mononuclear cells (peripheral blood mononuclear cells, hereinafter PBMC) as an effector.

[0286] Peripheral blood was collected from a healthy volunteer and an anticoagulant was added thereto. The blood was put statically on Ficoll-Plaque Plus (GE Healthcare) such that the interface was not disturbed and centrifuged at 2,000 rpm for 20 minutes using a large centrifuge (CF9RX, Hitachi, Ltd.) The intermediate layer containing the cells was collected and washed with PBS, platelets were removed by centrifugation at 900 rpm for 20 minutes, and the peripheral blood-derived mononuclear cells (PBMC) were used as an effector.

[0287] Further, PBMC cultured overnight under conditions of 37°C and 5% CO₂ using RPMI 1640 medium containing 10% fetal bovine serum to which human IL-2 (Peprtech) was added to a final concentration of 4 ng/ml (40 IU/ml or more) was also used as an effector of ADCC assay.

[0288] In the method, in simple, a target cell is cultured in the presence of an antibody and PBMC and the lysis rate of specific target cell by the antibody is measured.

[0289] The following "Colon-26/hCD123 ADCC assay method" was used for the measurement of the lysis rate. Specifically, a target cell was labeled with ⁵¹Cr by culturing the IL-3R α forced expression Colon-26 cell as a target cell at 37°C for 1 hour in the presence of 5% CO₂, together with sodium chromate labeled with a radioisotope ⁵¹Cr (Na₂⁵¹CrO₄, Perkin Elmer, NEZ030S). The labeled target cell was washed 3 times to remove excess ⁵¹Cr and then suspended in the medium and transferred to a 96-well plate to which antibodies had been added in advance at various concentrations. PBMC was suspended in the medium and transferred to the plate to which the target cell and antibodies had been added (effector/target ratio = 100). As the antibodies, the anti-IL-3R α antibody purified in Example 4 was used, and human serum-derived IgG (SIGMA) as a negative control. As various controls, a well of the medium and target cell alone, a well of PBMC and target alone and a well supplemented with Triton-X were prepared. The 96-well plate filled with mixed solutions was cultured at 37°C for 4 hours in the presence of 5% CO₂.

[0290] After centrifugation of the plate, 50 μ l of each supernatant was transferred to a scintillator-containing 96-well plate (Lumaplate-TM, Perkin Elmer) and dried at 56°C for 2 hours. The plate was sealed (TopSeal-A, Packard) and measured using a microplate reader (TopCount, Perkin Elmer).

[0291] Regarding the lysis rate of the target cell, amount of ⁵¹Cr in the sodium chromate released into the medium due to the lysis of cells was measured. That is, the "specific lysis rate" was calculated by dividing a value obtained by subtracting the value of a well to which the antibody was not added from the value of each well, by a value obtained by subtracting the value of a well to which the antibody was not added from the value of a well to which Triton-X was added (specific lysis rate is set to 100%).

[0292] The results are shown in Fig. 8 and Fig. 9. In each IL-3R α antibodies, the ADCC activity for the target cell was found depending on the concentration. Also, these antibodies exhibited higher ADCC activity than the chimeric 7G3 antibody as a control. This shows that the IL-3R α antibody exhibit high ADCC activity for IL-3R α expression cells and has a possibility of a treatment in which drug efficacy is a removal of the IL-3R α positive cell.

Example 12 Affinity test of anti-IL-3R α antibody to monkey IL-3R α protein

[0293] Regarding the presence or absence of binding of the thus obtained anti-human IL-3R α antibody to monkey IL-3R α , whether or not the anti-human IL-3R α antibody prepared in Example 7 binds to the *Macaca fascicularis* IL-3R α forced expression cell prepared in Example 1 was analyzed using flow cytometry.

[0294] Specifically, 2×10^5 cells of the monkey IL-3R α forced expression L929 cell were allowed to react with 100 μ l so that a final concentration of 10 μ g/ml, of the anti-human IL-3R α antibody at 4°C for 30 minutes. The antibodies used are anti-dinitrophenol (DNP) human IgG1 antibody (manufactured by this firm) as a negative control and Old4, Old5, Old17, Old19, New102 and chimeric 7G3 antibodies. Thereafter, this was washed 3 times using a staining medium (Dulbecco's PBS supplemented with 2% fetal bovine serum, 2 mM EDTA and 0.05% NaN₃). Next, a PE-labeled anti-human antibody λ chain specific antibody (Southern Bio) was allowed to undergo the reaction in the staining medium at a final concentration of 1 μ g/ml and washed 3 times with the staining medium in the same manner. Finally, the cells were mixed with the staining medium and whether the presence or absence of PE positive was analyzed by flow cytometry.

[0295] The results are shown in Fig. 10. It was found that the anti-human IL-3R α human antibodies, Old4, Old5, Old17, Old19, New102 and chimeric 7G3 antibodies, react with the *Macaca fascicularis* IL-3R α .

Example 13 Detailed epitope analysis of anti-human IL-3R α human antibodies

(Preparation of IL-3R α /GM-CSFR α chimeric protein expression cell)

[0296] In order to carry out further detailed epitope analysis of IL-3R α antibodies, a chimeric protein in which a region smaller than the domain of IL-3R α extra-membrane region was replaced by GM-CSFR α was expressed in a cell and affinity of each anti-IL-3R α antibody to the cell was analyzed. In brief, firstly, a region considered to be positioned at outside based on a three dimensional structure prediction of IL-3R α molecule was determined, secondly, vectors which express IL-3R α molecules in which the small region was replaced by GM-CSFR α were respectively constructed, thirdly, these were forcedly expressed in HEK293F cell and fourthly, whether or not each anti-IL-3R α antibody labeled with fluorescence dye binds thereto was observed by flow cytometry.

(Domain mapping of IL-3R α)

[0297] Among the 3 domains divided according to Example 7, A and B domains which were recognized by the obtained antibodies Old19 and New102 were selected and analyzed in detail. Based on the three dimensional structure of IL-4 receptor alpha chain (IL-4R α , CD124) (PDB: 3BPN; Chain C, Crystal Structure of the IL4-IL4r-IL3ra Ternary Complex), three dimensional structure of IL-3R α was subjected to homology modeling using SWISS-MODEL (<http://swissmodel.expasy.org/> SWISS-MODEL.html). The predicted IL-3R α protein structure was visualized using a graphic software RasMol (<http://rasmol.org/>) and 7 regions considered to be positioned at extracellular amino acid region of IL-3R α molecule were determined (Fig. 4).

[0298] In order to specify epitope of anti-human IL-3R α human antibody, a protein in which corresponding regions of GM-CSFR α were replaced by the 6 regions of IL-3R α divided as described in the above was prepared and expressed on the cell membrane, and the presence or absence of binding of antibodies was determined.

[0299] Using the IL-3R α -Flag/pEGFP-N1 plasmid DNA as a template, amplification was carried out by a PCR method which uses PrimeSTAR(R) HS DNA polymerase (TAKARA BIO INC.). Regarding the PCR, a two step reaction at 98°C 10 seconds-68°C 5 minutes was carried out 25 cycles. The PCR primers used are as follows.

Region 1 Deficiency;

CD123-Fw21: CGTGGAACCCGAGTGAACAATAGCTATT (SEQ ID NO:149)

CD123-Re21: ACTCTGTTCTTTTAACACACTCGATATCG (SEQ ID NO:150)

Region 2 Deficiency;

CD123-Fw22: CTTTATCCAAATAACAGTGGGAAGCCTTG (SEQ ID NO:151)

CD123-Re22: CAGTTTCTGTTGGAATGGTGGGTTGGCCACT (SEQ ID NO:152)

Region 3 Deficiency;

CD123-Fw23: AGGGAGGGTACCGGTGCGGAGAATCTGACCTGCT (SEQ ID NO: 153)

CD123-Re23: TCCTGAATTTGGATAGAAGAGGATCCACGTGG (SEQ ID NO:154)

Region 4 Deficiency;

CD123-Fw24: GGTCCGACGGCCCCCGCGGACGTCCAGTA (SEQ ID NO:155)

CD123-Re24 : CCTCGCCCAGGTACAGCTCAAGAAATCCACGT (SEQ ID NO:156)

Region 5 Deficiency;

CD123-Fw25: ACGGAACCAGCGCAGCCTTCGGTATCCCCT (SEQ ID NO:157)

CD123-Re25: TAACCAGAAAGTGGGAACCTTGAGAACC (SEQ ID NO:158)

Region 6 Deficiency;

CD123-Fw26: TCTTTGATTCATTTGTCGTCTTTTCACA (SEQ ID NO:159)

CD123-Re26: ATTGGATGCCGAAGGCTGCGCTCCTGCCC (SEQ ID NO:160)

[0300] The thus obtained PCR product was subjected to 0.8% agarose gel electrophoresis (135 V, 15 minutes, TAE buffer). DNA was visualized by ethidium bromide staining. After confirming the amplification, purification was carried out

using Wizard SV Gel and PCR Clean-Up System. The thus obtained DNA was subjected to phosphorylation using polynucleotide kinase (New England Biolabs) and to ethanol precipitation and then a part thereof was allowed to undergo the reaction using TaKaRa Ligation Kit. Regarding the transformation, a ligation sample and DH10B competent cell were mixed and spread on an LB plate (containing kanamycin). A plasmid DNA was extracted from the thus obtained colony by the Miniprep method and digested with *XhoI* and *NotI* and the insert was verified.

(Flow cytometry analysis of IL-3R α /GM-CSFR α chimeric protein expression cell using labeled anti-IL-3R α antibody)

[0301] HEK293T cell was used in the preparation of IL-3R α /GM-CSFR α chimeric protein expression cell. The plasmid DNA obtained in the above was introduced as an expression vector into the HEK293T. HEK293T introduced with the expression vector was cultured under an environment of 5% CO₂ and 37°C and used in the flow cytometry analysis 2 days after the introduction.

[0302] Each of Alexa Flour488-labeled human antibodies or commercially available FITC-labeled anti-IL-3R α mouse antibodies (7G3 and 9F5: both available from BD Biosciences, 6H6: from Acris Antibodies) at a concentration of 1 μ g/ml was allowed to react for 30 minutes on ice with 100,000 to 1,000,000 cells of the chimeric protein expression cell. A staining medium (Dulbecco's PBS containing 2% fetal calf serum, 2 mM EDTA and 0.05% NaN₃) was used for diluting the antibodies and cells. Next, the cells reacted with each antibody were washed 3 times with the staining medium, and whether or not the labeled antibody bound to the cells was confirmed by flow cytometry.

[0303] The results are shown in Fig. 3. The reaction of the antibody c7G3 disappeared only in the case of the protein expression cell in which the region 1 was replaced by GM-CSFR α . Reaction of the Old19 disappeared in the case of the protein expression cell in which the region 2 and region 3 in A domain and region 5 and region 6 in B domain were replaced by GM-CSFR α . Reaction of the New102 disappeared in the case of the protein expression cell in which the region 5 and region 6 of B domain was replaced by GM-CSFR α .

[0304] Based on the above, it was shown a possibility that the antibody Old19 recognized the regions 2 and 3 of A domain and regions 5 and 6 of B domain, and the antibody New102 recognized the regions 5 and 6 of B domain. The above results are summarized as Table 4.

[Table 4]

Region (domain)	Replacing sequence	7G3	9F5	6H6	Old19	New102
Region 1 (A)	55-DADYSMP-61	-	++	++	++	++
Region 2 (A)	91-STWLFPE-98	++	++	++	-	++
Region 3 (A-B)	97-PENSGKPWA-104	++	++	++	-	++
Region 4 (B)	122-CSWAVGPG-128	++	++	++	++	++
Region 5 (B)	182-ILVRGRS-188	++	++	++	-	-
Region 6 (B)	192-GIPCTDK-198	++	++	++	-	-

INDUSTRIAL APPLICABILITY

[0305] According to the invention, there is provided an antibody to human IL-3R α protein (another name: human CD 123) and a therapeutic agent and a diagnostic agent for myelocytic malignant tumors, particularly acute myeloid leukemia (AML), which comprises a human IL-3R α antibody as an active ingredient.

FREE TEXT OF SEQUENCE LISTING

[0306]

SEQ ID NO : 3 : IL-3R α _Fw primer
 SEQ ID NO : 4 : IL-3R α _Re primer
 SEQ ID NO : 5 : IL-3R α _seqF1 primer
 SEQ ID NO : 6 : Insert (MfeI from NotI)
 SEQ ID NO : 7 : Rhe123Fw1 primer
 SEQ ID NO : 8 : Rhe123Rv1 primer
 SEQ ID NO : 9 : T7 primer
 SEQ ID NO : 10 : SP6 primer

EP 2 949 673 A1

SEQ ID NO : 11 : Insert (*MefI* to *NotI*) of *Macaca fascicularis* IL-3R α
 SEQ ID NO : 12 : Insert (*MefI* to *NotI*) of *Macaca mulatta* IL-3R α
 SEQ ID NO : 13 : hIL-3R α so1-FLAG-*NotI* primer
 SEQ ID NO : 14 : Insert (*MejI* to *NotI*)
 5 SEQ ID NO : 15 : hh-6 primer
 SEQ ID NO : 16 : hh-3 primer
 SEQ ID NO : 17 : hh-4 primer
 SEQ ID NO : 18 : Old4 heavy chain specific primer Fw
 SEQ ID NO : 19 : Old4 heavy chain specific primer Rv
 10 SEQ ID NO : 20 : Old5 heavy chain specific primer Fw
 SEQ ID NO : 21 : Old5 heavy chain specific primer Rv
 SEQ ID NO : 22 : Old17 heavy chain specific primer Fw
 SEQ ID NO : 23 : Old17 heavy chain specific primer Rv
 SEQ ID NO : 24 : Old19 heavy chain specific primer Fw
 15 SEQ ID NO : 25 : Old19 heavy chain specific primer Rv
 SEQ ID NO : 26 : New102 heavy chain specific primer Fw
 SEQ ID NO : 27 : New102 heavy chain specific primer Rv
 SEQ ID NO : 28 : Old6 heavy chain specific primer Fw
 SEQ ID NO : 29 : Old6 heavy chain specific primer Rv
 20 SEQ ID NO : 30 : mH_Rv1 primer
 SEQ ID NO : 31 : mH_Rv2 primer
 SEQ ID NO : 32 : 7G3 heavy chain specific primer Fw
 SEQ ID NO : 33 : 7G3 heavy chain specific primer Rv
 SEQ ID NO : 34 : hk-2 primer
 25 SEQ ID NO : 35 : hk-6 primer
 SEQ ID NO : 36 : Old4 light chain specific primer Fw
 SEQ ID NO : 37 : Old4 light chain specific primer Rv
 SEQ ID NO : 3 : Old5 light chain specific primer Fw
 SEQ ID NO : 39 : Old5 light chain specific primer Rv
 30 SEQ ID NO : 40 : Old17 light chain specific primer Fw
 SEQ ID NO : 41 : Old17 light chain specific primer Rv
 SEQ ID NO : 42 : Old19 light chain specific primer Fw
 SEQ ID NO : 43 : Old 19 light chain specific primer Rv
 SEQ ID NO : 44 : New102 light chain specific primer Fw
 35 SEQ ID NO : 45 : New102 light chain specific primer Rv
 SEQ ID NO : 46 : Old6 light chain specific primer Fw
 SEQ ID NO : 47 : Old6 light chain specific primer Rv
 SEQ ID NO : 48 : mK_Rv1 primer
 SEQ ID NO : 49 : mK_Rv2 primer
 40 SEQ ID NO : 50 : 7G3 light chain specific primer Fw
 SEQ ID NO : 51 : 7G3 light chain specific primer Rv
 SEQ ID NO : 80 : hCD116Fw-MfeI primer
 SEQ ID NO : 81 : hCD116Rv-*NotI* primer
 SEQ ID NO : 82 : hCD116Fw-MfeI primer
 45 SEQ ID NO : 83 : hCD116Rv-*NotI* primer
 SEQ ID NO : 84 : hCD116Fw-MfeI primer
 SEQ ID NO : 85 : hCD116Rv-*NotI* primer
 SEQ ID NO : 86 : hCD116SeqFw1 primer
 SEQ ID NO : 87 : hCD116SeqFw2 primer
 50 SEQ ID NO : 88 : hCD116SeqRv1 primer
 SEQ ID NO : 89 : T7 primer
 SEQ ID NO : 90 : hCD123-C-FLAG-R1 primer
 SEQ ID NO : 91 : IL-3R α _Fw primer
 SEQ ID NO : 92 : C-FLAG-NotR2 primer
 55 SEQ ID NO : 93 : pEGFP-N1-Fw primer
 SEQ ID NO : 94 : pEGFP-N1-Re primer
 SEQ ID NO : 95 : pEGFP-N1-Fw primer
 SEQ ID NO : 96 : pEGFP-N1-Re primer

SEQ ID NO : 97 : CD123R11pEGFPN1 primer
 SEQ ID NO : 98 : CD123F11 primer
 SEQ ID NO : 99 : CD123R12-2 primer
 SEQ ID NO : 100 : CD123F12-2 primer
 SEQ ID NO : 101 : CD123R13 primer
 SEQ ID NO : 102 : CD123F13 primer
 SEQ ID NO : 103 : pEGFP-N1-Fw primer
 SEQ ID NO : 104 : pEGFP-N1-Re primer
 SEQ ID NO : 105 : GM-CSFRF11 primer
 SEQ ID NO : 106 : GM-CSFR11 primer
 SEQ ID NO : 107 : GM-CSFRF12 primer
 SEQ ID NO : 108 : GM-CSFR12 primer
 SEQ ID NO : 109 : GM-CSFRF13 primer
 SEQ ID NO : 110 : GM-CSFR13 primer
 SEQ ID NO : 111 : pEGFP-N1-Fw primer
 SEQ ID NO : 112 : pEGFP-N1-Re primer
 SEQ ID NO : 149 : CD123-Fw21 primer
 SEQ ID NO : 150 : CD123-Re21 primer
 SEQ ID NO : 151 : CD 123-Fw22 primer
 SEQ ID NO : 152 : CD123-Re22 primer
 SEQ ID NO : 153 : CD123-Fw23 primer
 SEQ ID NO : 154 : CD123-Re23 primer
 SEQ ID NO : 155 : CD123-Fw24 primer
 SEQ ID NO : 156 : CD123-Re24 primer
 SEQ ID NO : 157 : CD123-Fw25 primer
 SEQ ID NO : 158 : CD123-Re25 primer
 SEQ ID NO : 159 : CD123-Fw26 primer
 SEQ ID NO : 160 : CD123-Re26 primer

ASPECTS OF THE INVENTION

[0307]

1. An antibody to a human IL-3R α chain, which does not inhibit IL-3 signaling and binds to B domain of the human IL-3R α chain but does not bind to C domain of the human IL-3R α chain.

2. The antibody according to aspect 1, further having high antibody-dependent cellular cytotoxicity (ADCC).

3. The antibody according to aspect 1 or 2, wherein the high antibody-dependent cellular cytotoxicity (ADCC) shows a specific lysis rate of 10% at an antibody concentration of 0.01 μ g/ml, by a Colon-26/hCD123 ADCC assay method using PBMC cultured with IL-2.

4. The antibody according to any one of aspects 1 to 3, which comprises amino acid sequences of CDRs of heavy chain and CDRs of light chain selected from the group consisting of the following (a) to (e);

- (a) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs: 113 to 115, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs: 131 to 133, respectively,
- (b) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs: 116 to 118, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs: 134 to 136, respectively,
- (c) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs: 119 to 121, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs: 137 to 139, respectively,
- (d) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs: 122 to 124, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs: 140 to 142, respectively, and
- (e) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs: 125 to 127, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs: 143 to 145, respectively.

5. The antibody according to any one of aspects 1 to 4, which comprises a heavy chain variable region and a light chain variable region selected from the group consisting of the following (a) to (f);

(a) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:53 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:55;

(b) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:57 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:59;

(c) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:61 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:63;

(d) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:65 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:67;

(e) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 138 in the amino acid sequence of SEQ ID NO:69 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:71 and;

(f) a heavy chain variable region and/or light chain variable region, which comprise amino acid sequences in which 1 to 3 amino acid residues are deleted, substituted, added or inserted in the heavy chain variable region and/or light chain variable region shown by the above (a) to (e).

6. A composition for preventing or treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood of a subject, which comprises the IL-3R α antibody according to any one of aspects 1 to 5 as an active ingredient.

7. A method for treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood, which comprises administering, to a subject, a composition comprising the IL-3R α antibody according to any one of aspects 1 to 5 as an active ingredient.

8. A composition for detecting a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood of a biological sample from a subject, which comprises the IL-3R α antibody according to any one of aspects 1 to 5.

9. The composition or method according to any one of aspects 1 to 5, wherein the blood tumor is acute myeloid leukemia (AML).

EP 2 949 673 A1

SEQUENCE LISTING

<110> Kyowa Hakko Kirin Co., Ltd.

<120> Anti-IL-3RA Antibody for use in treatment of blood tumor

<130> N115530A-EP

<150> US 61/172923

<151> 2009-04-27

<160> 160

<170> PatentIn version 3.3

<210> 1

<211> 378

<212> PRT

<213> Homo sapiens

<400> 1

Met Val Leu Leu Trp Leu Thr Leu Leu Leu Ile Ala Leu Pro Cys Leu
1 5 10 15

Leu Gln Thr Lys Glu Asp Pro Asn Pro Pro Ile Thr Asn Leu Arg Met
20 25 30

Lys Ala Lys Ala Gln Gln Leu Thr Trp Asp Leu Asn Arg Asn Val Thr
35 40 45

Asp Ile Glu Cys Val Lys Asp Ala Asp Tyr Ser Met Pro Ala Val Asn
50 55 60

Asn Ser Tyr Cys Gln Phe Gly Ala Ile Ser Leu Cys Glu Val Thr Asn
65 70 75 80

Tyr Thr Val Arg Val Ala Asn Pro Pro Phe Ser Thr Trp Ile Leu Phe
85 90 95

Pro Glu Asn Ser Gly Lys Pro Trp Ala Gly Ala Glu Asn Leu Thr Cys
100 105 110

Trp Ile His Asp Val Asp Phe Leu Ser Cys Ser Trp Ala Val Gly Pro
115 120 125

Gly Ala Pro Ala Asp Val Gln Tyr Asp Leu Tyr Leu Asn Val Ala Asn
130 135 140

Arg Arg Gln Gln Tyr Glu Cys Leu His Tyr Lys Thr Asp Ala Gln Gly
145 150 155 160

Thr Arg Ile Gly Cys Arg Phe Asp Asp Ile Ser Arg Leu Ser Ser Gly

EP 2 949 673 A1

	165	170	175
5	Ser Gln Ser Ser His Ile Leu Val Arg Gly Arg Ser Ala Ala Phe Gly 180 185 190		
10	Ile Pro Cys Thr Asp Lys Phe Val Val Phe Ser Gln Ile Glu Ile Leu 195 200 205		
15	Thr Pro Pro Asn Met Thr Ala Lys Cys Asn Lys Thr His Ser Phe Met 210 215 220		
20	His Trp Lys Met Arg Ser His Phe Asn Arg Lys Phe Arg Tyr Glu Leu 225 230 235 240		
25	Gln Ile Gln Lys Arg Met Gln Pro Val Ile Thr Glu Gln Val Arg Asp 245 250 255		
30	Arg Thr Ser Phe Gln Leu Leu Asn Pro Gly Thr Tyr Thr Val Gln Ile 260 265 270		
35	Arg Ala Arg Glu Arg Val Tyr Glu Phe Leu Ser Ala Trp Ser Thr Pro 275 280 285		
40	Gln Arg Phe Glu Cys Asp Gln Glu Glu Gly Ala Asn Thr Arg Ala Trp 290 295 300		
45	Arg Thr Ser Leu Leu Ile Ala Leu Gly Thr Leu Leu Ala Leu Val Cys 305 310 315 320		
50	Val Phe Val Ile Cys Arg Arg Tyr Leu Val Met Gln Arg Leu Phe Pro 325 330 335		
55	Arg Ile Pro His Met Lys Asp Pro Ile Gly Asp Ser Phe Gln Asn Asp 340 345 350		
	Lys Leu Val Val Trp Glu Ala Gly Lys Ala Gly Leu Glu Glu Cys Leu 355 360 365		
	Val Thr Glu Val Gln Val Val Gln Lys Thr 370 375		
	<210> 2 <211> 308 <212> PRT <213> Homo sapiens		
	<400> 2		
	Met Val Leu Leu Trp Leu Thr Leu Leu Leu Ile Ala Leu Pro Cys Leu		

EP 2 949 673 A1

	1			5					10				15			
5	Leu	Gln	Thr	Lys	Glu	Asp	Pro	Asn	Pro	Pro	Ile	Thr	Asn	Leu	Arg	Met
				20					25					30		
	Lys	Ala	Lys	Ala	Gln	Gln	Leu	Thr	Trp	Asp	Leu	Asn	Arg	Asn	Val	Thr
			35					40					45			
10	Asp	Ile	Glu	Cys	Val	Lys	Asp	Ala	Asp	Tyr	Ser	Met	Pro	Ala	Val	Asn
	50						55					60				
15	Asn	Ser	Tyr	Cys	Gln	Phe	Gly	Ala	Ile	Ser	Leu	Cys	Glu	Val	Thr	Asn
	65					70					75					80
	Tyr	Thr	Val	Arg	Val	Ala	Asn	Pro	Pro	Phe	Ser	Thr	Trp	Ile	Leu	Phe
20					85					90					95	
	Pro	Glu	Asn	Ser	Gly	Lys	Pro	Trp	Ala	Gly	Ala	Glu	Asn	Leu	Thr	Cys
				100					105					110		
25	Trp	Ile	His	Asp	Val	Asp	Phe	Leu	Ser	Cys	Ser	Trp	Ala	Val	Gly	Pro
			115					120					125			
	Gly	Ala	Pro	Ala	Asp	Val	Gln	Tyr	Asp	Leu	Tyr	Leu	Asn	Val	Ala	Asn
30							135						140			
	Arg	Arg	Gln	Gln	Tyr	Glu	Cys	Leu	His	Tyr	Lys	Thr	Asp	Ala	Gln	Gly
35	145					150					155					160
	Thr	Arg	Ile	Gly	Cys	Arg	Phe	Asp	Asp	Ile	Ser	Arg	Leu	Ser	Ser	Gly
					165					170					175	
40	Ser	Gln	Ser	Ser	His	Ile	Leu	Val	Arg	Gly	Arg	Ser	Ala	Ala	Phe	Gly
				180					185					190		
	Ile	Pro	Cys	Thr	Asp	Lys	Phe	Val	Val	Phe	Ser	Gln	Ile	Glu	Ile	Leu
45			195					200					205			
	Thr	Pro	Pro	Asn	Met	Thr	Ala	Lys	Cys	Asn	Lys	Thr	His	Ser	Phe	Met
		210					215					220				
50	His	Trp	Lys	Met	Arg	Ser	His	Phe	Asn	Arg	Lys	Phe	Arg	Tyr	Glu	Leu
	225					230					235					240
	Gln	Ile	Gln	Lys	Arg	Met	Gln	Pro	Val	Ile	Thr	Glu	Gln	Val	Arg	Asp
55					245					250					255	

EP 2 949 673 A1

Arg Thr Ser Phe Gln Leu Leu Asn Pro Gly Thr Tyr Thr Val Gln Ile
260 265 270

5 Arg Ala Arg Glu Arg Val Tyr Glu Phe Leu Ser Ala Trp Ser Thr Pro
275 280 285

10 Gln Arg Phe Glu Cys Asp Gln Glu Glu Gly Ala Asn Thr Arg Ala Trp
290 295 300

Arg Thr Ser Leu
305

15 <210> 3
<211> 34
<212> DNA
<213> Artificial

20 <220>
<223> primer IL-3RA_Fw

<400> 3
cggcaattgc caccatgggc ctcctttggc tcac 34

25 <210> 4
<211> 31
<212> DNA
<213> Artificial

30 <220>
<223> IL-3RA_Re

<400> 4
attgcggccg ctcaagtttt ctgcacgacc t 31

<210> 5
<211> 20
<212> DNA
<213> Artificial

<220>
<223> primer IL-3RA_seqF1

45 <400> 5
gtcttcacta caaaacggat 20

<210> 6
<211> 1156
<212> DNA
<213> Artificial

<220>
<223> insert sequence

55 <400> 6
caattgccac catgttcctc ctttggctca cgctgtcct gatcgccctg cctgtctcc 60

EP 2 949 673 A1

	tgcaaacgaa ggaagatcca aaccaccaa tcacgaacct aaggatgaaa gcaaaggctc	120
	agcagttgac ctgggacctt aacagaaatg tgaccgatat cgagtgtgtt aaagacgccg	180
5	actattctat gccggcagtg aacaatagct attgccagtt tggagcaatt tccttatgtg	240
	aagtgaccaa ctacaccgtc cgagtggcca acccaccatt ctccacgtgg atcctcttcc	300
	ctgagaacag tgggaagcct tgggcaggtg cggagaatct gacctgctgg attcatgacg	360
10	tggatttctt gagctgcagc tgggcggtag gcccgggggc ccccgaggac gtccagtacg	420
	acctgtactt gaacgttgcc aacaggcgtc aacagtacga gtgtcttcac tacaaaacgg	480
15	atgctcaggg aacacgtatc ggggtgtcgtt tcgatgacat ctctcgactc tccagcgggtt	540
	ctcaaagttc ccacatcctg gtgcggggca ggagcgcagc cttcggtatc ccctgcacag	600
	ataagtttgt cgtcttttca cagattgaga tattaactcc acccaacatg actgcaaagt	660
20	gtaataagac acattccttt atgcactgga aaatgagaag tcatttcaat cgcaaatttc	720
	gctatgagct tcagatacaa aagagaatgc agcctgtaat cacagaacag gtcagagaca	780
	gaacctcctt ccagctactc aatcctggaa cgtacacagt acaaataaga gcccgggaaa	840
25	gagtgtatga attcttgagc gcctggagca cccccagcg cttcgagtgc gaccaggagg	900
	agggcgcaaa cacacgtgcc tggcggacgt cgctgctgat cgcgctgggg acgctgctgg	960
	ccctgggtctg tgtcttcgtg atctgcagaa ggtatctggt gatgcagaga ctctttcccc	1020
30	gcatccctca catgaaagac cccatcggtg acagcttcca aaacgacaag ctggtggtct	1080
	gggaggcggg caaagccggc ctggaggagt gtctggtgac tgaagtacag gtcgtgcaga	1140
35	aaacttgagc ggccgc	1156
	<210> 7 <211> 38 <212> DNA <213> Artificial	
40	<220> <223> primer Rhe123Fw1	
	<400> 7 cggcaattgc caccatgacc ctcttttggc tgacgctg	
45		38
	<210> 8 <211> 38 <212> DNA <213> Artificial	
50	<220> <223> primer Rhe123Rv1	
55	<400> 8 tatattgcgg ccgctcaagt tttctccacc acctgcac	
		38

EP 2 949 673 A1

<210> 9
 <211> 20
 <212> DNA
 <213> Artificial
 5
 <220>
 <223> primer T7
 <400> 9
 10 taatacgcact cactataggg 20
 <210> 10
 <211> 19
 <212> DNA
 15 <213> Artificial
 <220>
 <223> primer SP6
 <400> 10
 20 gatttaggtg acactatag 19
 <210> 11
 <211> 1156
 25 <212> DNA
 <213> Artificial
 <220>
 <223> insert sequence
 30 <400> 11
 caattgccac catgaccctc ctttggctga cgctgctcct ggtcgccacg ccctgtctcc 60
 tgcaaacgaa ggaggatcca aatgcaccaa tcaggaatct aaggatgaaa gaaaaggctc 120
 35 agcagttgat gtgggacctg aacagaaaacg tgaccgacgt ggagtgtatc aaaggcaccg 180
 actattctat gccggcaatg aacaacagct attgccagtt cggagccatt tccttatgtg 240
 aagtgaacaa ctacaccgtc cgagtggcca gtcccccggt ctccacgtgg atcctcttcc 300
 40 ctgagaacag tgggacgcct caggcaggcg cggagaatct gacctgctgg gttcatgacg 360
 tggatttctt gagctgcagc tgggtggcag gcccgggcgg ccccgctgac gtccagtacg 420
 acctgtactt gaacaatccc aacagccacg aacagtacag gtgccttcac tacaaaacgg 480
 45 atgctcgggg aacacagatc ggggtgtcggg togatgacat cgctcgactc tcccgcgggt 540
 ctcaaagttc ccacatcctg gtgaggggca ggagcgcagc cgtcagtatc ccctgcacag 600
 ataagtttgt cttcttttca cagattgaga gattaactcc acccaacatg actggagagt 660
 50 gtaatgagac acattccttc atgcactgga aaatgaaaag tcatttcaat cgcaaattcc 720
 gctatgagct tcggatccaa aagagaatgc agcctgtaag gacagaacag gtcagagaca 780
 55 caacctcctt ccagctaccc aatcctggaa cgtacacagt gcaaataaga gcccgggaaa 840
 cagtgtatga attcttgagt gcctggagca cccccagcg cttcgagtgc gaccaggagg 900

EP 2 949 673 A1

	agggcgcgag ctcgcggtgcc tggcggacgt cgctgctgat cgcgctgggg acgctgctgg	960
	ccttgctctg tgtgttcctc atctgcagaa ggtatctggt gatgcagagg ctgtttcccc	1020
5	gcatcccaca catgaaagac cccatcggtg acaccttcca acaggacaag ctggtggtct	1080
	gggagggcggg caaagccggc ctggaggagt gtctggtgtc tgaagtgcag gtggtggaga	1140
	aaacttgagc ggccgc	1156
10	<210> 12 <211> 1156 <212> DNA <213> Artificial	
15	<220> <223> insert sequence	
	<400> 12 caattgccac catgaccctc ctttggtga cgctgctcct ggtcgccacg ccctgtctcc	
20	tgcaaaccaa ggaggatcca aatgcaccaa tcaggaatct aaggatgaaa gaaaaggctc	60
	agcagttgat gtgggacctg aacagaaacg tgaccgacgt ggagtgtatc aaaggcaccg	120
	actattctat gccggcaatg aacgacagct attgccagtt cggagccatt tccttatgtg	180
25	aagtgaccaa ctacaccgtc cgagtggcca gtcctccgtt ctccacgtgg atcctcttcc	240
	ctgagaacag tgggacgcct cgggcaggcg cggagaattt gacctgctgg gttcatgacg	300
30	tggatttctt gagctgcagc tgggtggtag gcccgggcggc ccccgctgac gtccagtacg	360
	acctgtactt gaacaatccc aacagccacg aacagtacag gtgccttcgc taaaaacgg	420
	atgctcgggg aacacagatc ggggtgctggt tcgatgacat cgctcgactc tcccgcggtt	480
35	ctcaaagttc ccacatcctg gtgaggggca ggagcgcagc cgtcagtatc ccctgcacag	540
	ataagtttgt cttcttttca cagattgaga gattaactcc acccaacatg actggagagt	600
	gtaatgagac acattccttc atgcaactgga aaatgaaaag tcatttcaat cgcaaattcc	660
40	actatgagct tcggatccaa aagagaatgc agcctgtaag gacagaacag gtcagagaca	720
	caacctcctt ccagctaccc aatcctggaa cgtacacagt gcaaataaga gcccgggaaa	780
	cagtgtatga attccttgagt gcctggagca cccccagcg cttcgagtgc gaccaggagg	840
45	agggcgcgag ctcgcggtgcc tggcggacgt cgctgctgat cgcgctgggg acgctgctgg	900
	ccttgctctg tgtgttcctc atctgcagaa ggtatctggt gatgcagagg ctgtttcccc	960
50	gcatcccaca catgaaagac cccatcggtg acaccttcca acaggacaag ctggtggtct	1020
	gggagggcggg caaagccggc ctggaggagt gtctggtgtc tgaagtgcag gtggtggaga	1080
	aaacttgagc ggccgc	1156
55	<210> 13 <211> 58	

EP 2 949 673 A1

<212> DNA
 <213> Artificial

 <220>
 5 <223> primer hIL-3RAsol-FLAG-NotI

 <400> 13
 attgcggccg ctcaattatc gtcgtcatcc ttgtagtccc gccaggcacg tgtgtttg 58

 10 <210> 14
 <211> 961
 <212> DNA
 <213> Artificial

 15 <220>
 <223> insert sequence

 <400> 14
 caattgccac catggtcctc ctttggtcctc cgctgctcct gatcgccctg ccctgtctcc 60
 20 tgcaaacgaa ggaagatcca aaccacacaa tcacgaacct aaggatgaaa gcaaaggctc 120
 agcagttgac ctgggacctt aacagaaatg tgaccgatat cgagtgtgtt aaagacgccg 180
 actattctat gccggcagtg aacaatagct attgccagtt tggagcaatt tccttatgtg 240
 25 aagtgaccaa ctacaccgtc cgagtggcca acccaccatt ctccacgtgg atcctcttcc 300
 ctgagaacag tgggaagcct tgggcaggtg cggagaatct gacctgctgg attcatgacg 360
 tggatttctt gagctgcagc tgggcggtag gcccgggggc ccccgcgac gtccagtacg 420
 30 acctgtactt gaacgttgcc aacaggcgtc aacagtacga gtgtcttcac tacaaaacgg 480
 atgctcaggg aacacgtatc ggggtgctgtt tcgatgacat ctctcgactc tccagcggtt 540
 ctcaaagttc ccacatcctg gtgcggggca ggagcgcagc cttcggtatc ccctgcacag 600
 35 ataagtttgt cgtcttttca cagattgaga tattaactcc acccaacatg actgcaaagt 660
 gtaataagac acattccttt atgcaactgga aaatgagaag tcatttcaat cgcaaatttc 720
 40 gctatgagct tcagatacaa aagagaatgc agcctgtaat cacagaacag gtcagagaca 780
 gaacctcctt ccagctactc aatcctggaa cgtacacagt acaaataaga gcccgggaaa 840
 gagtgtatga attcttgagc gcctggagca cccccagcg cttcgagtgc gaccaggagg 900
 45 agggcgcaaa cacacgtgcc tggcgggact acaaggatga cgacgataag tgagcggccg 960
 c 961

 50 <210> 15
 <211> 27
 <212> DNA
 <213> Artificial

 <220>
 55 <223> primer hh-6

 <400> 15

ggtccgggag atcatgaggg tgcctt 27

 5 <210> 16
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 10 <223> primer hh-3

 <400> 16
 gtgcacgccg ctggtcaggg cgcctg 26

 15 <210> 17
 <211> 23
 <212> DNA
 <213> Artificial

 20 <220>
 <223> primer hh-4

 <400> 17
 ggtgccaggg ggaagaccga tgg 23

 25 <210> 18
 <211> 49
 <212> DNA
 <213> Artificial

 30 <220>
 <223> primer Fw spcific for Old4 Heavy chain

 <400> 18
 35 agagagagag gtcgaccacc atggactgga cctggagggtt cctctttgt 49

 <210> 19
 <211> 40
 <212> DNA
 40 <213> Artificial

 <220>
 <223> primer Rv specific for Old4 heavy chain

 <400> 19
 45 agagagagag gctagctgaa gagacggtga ccattgtccc 40

 <210> 20
 <211> 49
 50 <212> DNA
 <213> Artificial

 <220>
 <223> primer Fw specific for Old5 heavy chain

 55 <400> 20
 agagagagag gtcgaccacc atggactgga cctggagggtt cctctttgt 49

EP 2 949 673 A1

<210> 21
 <211> 40
 <212> DNA
 <213> Artificial
 5
 <220>
 <223> primer Rv specific for Old5 heavy chain
 <400> 21
 10 agagagagag gctagctgaa gagacggtga ccattgtccc 40
 <210> 22
 <211> 49
 <212> DNA
 15 <213> Artificial
 <220>
 <223> primer Fw specific for Old17 Heavy chain
 <400> 22
 20 agagagagag gtcgaccacc atggactgga cctggaggtt cctctttgt 49
 <210> 23
 <211> 41
 <212> DNA
 25 <213> Artificial
 <220>
 <223> primer Rv specific for Old 17 heavy chain
 <400> 23
 30 agagagagag gctagctgag gagacggtga caagggttcc c 41
 <210> 24
 <211> 49
 <212> DNA
 35 <213> Artificial
 <220>
 <223> primer Fw specific for Old19 heavy chain
 <400> 24
 40 agagagagag gtcgaccacc atggactgga cctggaggtt cctctttgt 49
 <210> 25
 <211> 39
 <212> DNA
 45 <213> Artificial
 <220>
 <223> primer Rv specific for Old19 heavy chain
 <400> 25
 50 agagagagag gctagctgag gagacggtga ccagggttc 39
 <210> 26
 <211> 49
 <212> DNA
 55

EP 2 949 673 A1

<213> Artificial
 <220>
 <223> primer Fw specific for New102 Heavy chain
 5
 <400> 26
 agagagagag gtcgaccacc atggactgga cctggagggtt cctctttgt 49
 <210> 27
 10 <211> 38
 <212> DNA
 <213> Artificial
 <220>
 15 <223> primer Rv specific for New102 heavy chain
 <400> 27
 agagagagag gctagctgag gagacggtga ccagggtt 38
 <210> 28
 20 <211> 41
 <212> DNA
 <213> Artificial
 <220>
 25 <223> primer Fw specific for Old6 heavy chain
 <400> 28
 agagagagag gtcgaccac catggaactg gggctccgct g 41
 30 <210> 29
 <211> 39
 <212> DNA
 <213> Artificial
 35 <220>
 <223> primer Rv specific for Old6 heavy chain
 <400> 29
 agagagagag gctagctgag gagacggtga ccagggttc 39
 40 <210> 30
 <211> 32
 <212> DNA
 <213> Artificial
 45 <220>
 <223> primer mH_Rv1
 <400> 30
 50 attttgtcga ckyggtst gctggcyggg tg 32
 <210> 31
 <211> 30
 <212> DNA
 55 <213> Artificial
 <220>

EP 2 949 673 A1

<223> primer mH_Rv2
 <400> 31
 gcacacyrct ggacagggat ccagagttcc 30
 5
 <210> 32
 <211> 44
 <212> DNA
 <213> Artificial
 10
 <220>
 <223> primer Fw specific for 7G3 heavy chain
 <400> 32
 agagagagag gtcgaccacc atgggatgga gctggatctt tctc 44
 15
 <210> 33
 <211> 40
 <212> DNA
 <213> Artificial
 20
 <220>
 <223> primer Rv specific for 7G3 heavy chain
 <400> 33
 agagagagag gctagctgca gagacagtga ccagagtccc 40
 25
 <210> 34
 <211> 26
 <212> DNA
 <213> Artificial
 30
 <220>
 <223> primer hk-2
 <400> 34
 gttgaagctc tttgtgacgg gcgagc 26
 35
 <210> 35
 <211> 26
 <212> DNA
 <213> Artificial
 40
 <220>
 <223> primer hk-6
 <400> 35
 tggcgggaag atgaagacag atggtg 26
 45
 <210> 36
 <211> 45
 <212> DNA
 <213> Artificial
 50
 <220>
 <223> primer Fw specific for Old4 light chain
 <400> 36
 55

EP 2 949 673 A1

agagagagag atctctcacc atggacatga ggggtccccgc tcagc 45

5 <210> 37
<211> 40
<212> DNA
<213> Artificial

<220>
10 <223> primer Rv specific for Old4 light chain

<400> 37
agagagagag cgtacgtttg atctccagct tgggtcccctg 40

15 <210> 38
<211> 45
<212> DNA
<213> Artificial

<220>
20 <223> primer Fw specific for Old5 light chain

<400> 38
agagagagag atctctcacc atggacatga ggggtccccgc tcagc 45

25 <210> 39
<211> 40
<212> DNA
<213> Artificial

30 <220>
<223> primer Rv specific for Old5 light chain

<400> 39
35 agagagagag cgtacgtttg atctccagct tgggtcccctg 40

<210> 40
<211> 44
<212> DNA
40 <213> Artificial

<220>
<223> primer Fw specific for Old17 light chain

<400> 40
45 agagagagag atctctcacc atggacatga ggggtcctcgc tcag 44

<210> 41
<211> 40
50 <212> DNA
<213> Artificial

<220>
<223> primer Rv specific for light chain

55 <400> 41
agagagagag cgtacgtttg atctccagct tgggtcccctg 40

EP 2 949 673 A1

<210> 42
 <211> 44
 <212> DNA
 <213> Artificial
 5
 <220>
 <223> primer Fw specific for Old19 light chain
 <400> 42
 10 agagagagag atctctcacc atggacatga gggtcctcgc tcag 44
 <210> 43
 <211> 42
 <212> DNA
 15 <213> Artificial
 <220>
 <223> primer Rv specific for Old19 light chain
 <400> 43
 20 agagagagag cgtacgtttg atttccacct tggtccttg gc 42
 <210> 44
 <211> 44
 <212> DNA
 25 <213> Artificial
 <220>
 <223> primer Fw specific for New102 light chain
 <400> 44
 30 agagagagag atctctcacc atggacatga gggtcctcgc tcag 44
 <210> 45
 <211> 40
 <212> DNA
 35 <213> Artificial
 <220>
 <223> primer Rv specific for New102 light chain
 <400> 45
 40 agagagagag cgtacgtttg atctccagct tggtccttg 40
 <210> 46
 <211> 44
 <212> DNA
 45 <213> Artificial
 <220>
 <223> primer Fw specific for Old6 light chain
 <400> 46
 50 gagagagaga tctctcacca tggacatgag ggtccccgct cagc 44
 <210> 47
 <211> 42
 <212> DNA
 55

EP 2 949 673 A1

<213> Artificial
 <220>
 <223> primer Rv specific for Old6 light chain
 5
 <400> 47
 agagagagag cgtacgtttg atatccactt tgggtcccagg gc 42
 <210> 48
 10 <211> 30
 <212> DNA
 <213> Artificial
 <220>
 15 <223> primer mK_Rv1
 <400> 48
 ttgaagctct tgacaatggg tgaagttgat 30
 <210> 49
 20 <211> 30
 <212> DNA
 <213> Artificial
 <220>
 25 <223> primer mK_Rv2
 <400> 49
 gtaggtgctg tctttgctgt cctgatcagt 30
 <210> 50
 30 <211> 44
 <212> DNA
 <213> Artificial
 <220>
 35 <223> primer Fw specific for 7G3 light chain
 <400> 50
 agagagagag agatctcacc atggaatcac agactcaggt cctc 44
 <210> 51
 40 <211> 40
 <212> DNA
 <213> Artificial
 <220>
 45 <223> primer Rv specific for 7G3 light chain
 <400> 51
 50 agagagagag cgtacgtttt atttccagct tgggtccccc 40
 <210> 52
 <211> 443
 <212> DNA
 55 <213> Homo sapiens
 <400> 52

EP 2 949 673 A1

```

gacccgtcga ccaccatgga ctggacctgg aggttcctct ttgtgggtggc agcagctaca      60
ggtgtccagt cccaggtcca gctgctacag tctggggctg aggtgaagaa gcctgggtcc      120
5  tcggtgaagg tctcatgcaa ggcttctgga ggcaccttca gcacctatgc tatcagctgg      180
gtgcgacagg cccctggaca agggcttgag tggatgggag ggatcatccc tatctttggt      240
atagtaaact acgcacagaa gttccagggc agagtcacga ttaccgcgga cgaatccacg      300
10 agtacagcct acatggaact gagcagcctg agatctgagg acacggccgt gtattattgt      360
gcgagagggg ggggctcggg cccagatgtt cttgatatct ggggccaagg gacaatggtc      420
accgtctctt cagctagcac caa                                             443

```

15

```

<210> 53
<211> 143
<212> PRT
<213> Homo sapiens

```

20

```

<220>
<221> misc_feature
<222> (143)..(143)
<223> Xaa can be any naturally occurring amino acid

```

25

<400> 53

```

Met Asp Trp Thr Trp Arg Phe Leu Phe Val Val Ala Ala Ala Thr Gly
1          5          10          15

```

30

```

Val Gln Ser Gln Val Gln Leu Leu Gln Ser Gly Ala Glu Val Lys Lys
          20          25          30

```

35

```

Pro Gly Ser Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe
          35          40          45

```

40

```

Ser Thr Tyr Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
          50          55          60

```

45

```

Glu Trp Met Gly Gly Ile Ile Pro Ile Phe Gly Ile Val Asn Tyr Ala
65          70          75          80

```

```

Gln Lys Phe Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser
          85          90          95

```

50

```

Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
          100          105          110

```

```

Tyr Tyr Cys Ala Arg Gly Gly Gly Ser Gly Pro Asp Val Leu Asp Ile
          115          120          125

```

55

```

Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser Ala Ser Thr Xaa

```

EP 2 949 673 A1

130 135 140

5
 <210> 54
 <211> 412
 <212> DNA
 <213> Homo sapiens

10
 <400> 54
 cacagatctc tcacatgga catgagggtc cccgctcagc tcctggggct cctgctgctc 60
 tggtcccag gtgccagatg tgtcatctgg atgacccagt ctccatcctt actctctgca 120
 tctacaggag acagagtcac catcagttgt cggatgagtc agggcattag gagttattta 180
 gcctggtatc agcaaaaacc agggaaagcc cctgagctcc tgatctatgc tgcattccact 240
 ttgcaaagtg ggggcccatc aaggttcagt ggcagtggat ctgggacaga tttcactctc 300
 accatcagca gcctgcagtc tgaagatttt gcaacttatt actgtcaaca gtattatagt 360
 20 tccccgtaca cttttggcca ggggaccaag ctggagatca aacgtacggt gg 412

25
 <210> 55
 <211> 133
 <212> PRT
 <213> Homo sapiens

30
 <220>
 <221> misc_feature
 <222> (133)..(133)
 <223> Xaa can be any naturally occurring amino acid

35
 <400> 55
 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1 5 10 15

40
 Leu Pro Gly Ala Arg Cys Val Ile Trp Met Thr Gln Ser Pro Ser Leu
 20 25 30

45
 Leu Ser Ala Ser Thr Gly Asp Arg Val Thr Ile Ser Cys Arg Met Ser
 35 40 45

50
 Gln Gly Ile Arg Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60

55
 Ala Pro Glu Leu Leu Ile Tyr Ala Ala Ser Thr Leu Gln Ser Gly Val
 65 70 75 80

60
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 85 90 95

65
 Ile Ser Ser Leu Gln Ser Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 100 105 110

EP 2 949 673 A1

Tyr Tyr Ser Phe Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
115 120 125

5 Lys Arg Thr Val Xaa
130

10 <210> 56
<211> 433
<212> DNA
<213> Homo sapiens

15 <400> 56
gtcgaccacc atggactgga cctggagggt cctctttgtg gtggcagcag ctacagggtgt 60
ccagtcccag gtccagctgg tgcagttctgg ggctgagggtg aagaagcctg ggtcctcgggt 120
gaaggtctca tgcaaggctt ctggaggcac cttcagcacc tatgctatca gctgggtgctg 180
20 acaggcccct ggacaagggc ttgagtggat gggaggggtc atccctatct ttgatataga 240
aaactacgca cagaagttcc agggcagagt cacgattacc gcggacgaat ccacgagcac 300
agtctatatg gaactgagca gcctgagatc tgaggacacg gccatgtatt actgtgctgag 360
25 aggggggggt tcgggccctg atgttcttga tatctggggc caagggacaa tggtcaccgt 420
ctcttcagct agc 433

30 <210> 57
<211> 141
<212> PRT
<213> Homo sapiens

35 <400> 57

Met Asp Trp Thr Trp Arg Phe Leu Phe Val Val Ala Ala Ala Thr Gly
1 5 10 15

40 Val Gln Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20 25 30

45 Pro Gly Ser Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe
35 40 45

Ser Thr Tyr Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
50 55 60

50 Glu Trp Met Gly Gly Leu Ile Pro Ile Phe Asp Ile Glu Asn Tyr Ala
65 70 75 80

55 Gln Lys Phe Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser
85 90 95

EP 2 949 673 A1

Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met
100 105 110

5 Tyr Tyr Cys Ala Arg Gly Gly Gly Ser Gly Pro Asp Val Leu Asp Ile
115 120 125

10 Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser Ala Ser
130 135 140

<210> 58
<211> 412
<212> DNA
15 <213> Homo sapiens

<400> 58
cacagatctc tcaccatgga catgagggtc cccgctcagc tcctggggct cctgctgctc 60
20 tggctcccag gtgccagatg tgatcatctgg atgacccagt ctccatcctt actctctgca 120
tctacaggag acagagtcac catcagttgt cggatgagtc agggcattag gagttattta 180
gcctgggtatc agcaaaaacc agggaaagcc cctgagctcc tgatctatgc tgcattccact 240
25 ttgcaaagtg ggggtcccatc aagggttcagt ggcagtggat ctgggacaga tttcactctc 300
accatcagca gcctgcagtc tgaagatttt gcaacttatt actgtcaaca gtattatagt 360
30 ttcccgtaca cttttggcca ggggaccaag ctggagatca aacgtacggt gg 412

<210> 59
<211> 133
<212> PRT
35 <213> Homo sapiens

<220>
<221> misc_feature
40 <222> (133)..(133)
<223> Xaa can be any naturally occurring amino acid

<400> 59

45 Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
1 5 10 15

Leu Pro Gly Ala Arg Cys Val Ile Trp Met Thr Gln Ser Pro Ser Leu
20 25 30

50 Leu Ser Ala Ser Thr Gly Asp Arg Val Thr Ile Ser Cys Arg Met Ser
35 40 45

55 Gln Gly Ile Arg Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
50 55 60

EP 2 949 673 A1

Ala Pro Glu Leu Leu Ile Tyr Ala Ala Ser Thr Leu Gln Ser Gly Val
65 70 75 80

5 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

10 Ile Ser Ser Leu Gln Ser Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

Tyr Tyr Ser Phe Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
115 120 125

15 Lys Arg Thr Val Xaa
130

20 <210> 60
<211> 438
<212> DNA
<213> Homo sapiens

25 <400> 60
gacccgtcga ccaccatgga ctggacctgg aggttcctct ttgtggtggc agcagctaca 60
ggtgtccagt ccaggtcca gctggtgcag tctggggctg aggtgaagaa gcctgggtcc 120
tcggtgaagg tctcctgcaa gacttctgga ggcaccttca gcaactttgc tatcagctgg 180
30 gtgcgacagg cccctggaca agggcttgag tggatgggag ggatcatccc tatctttggt 240
tcaacaaact acgcacagaa gttccagggc agagtcacga ttaacgcgga cgaatccacg 300
agcacagcct acatggagct gagcagtctg agatctgagg acacggccgt gtattactgt 360
35 gcgggtggag acaaatatgg tccttactac tttcactact ggggccaggg aacccttgtc 420
accgtctcct cagctagc 438

40 <210> 61
<211> 141
<212> PRT
<213> Homo sapiens

45 <400> 61
Met Asp Trp Thr Trp Arg Phe Leu Phe Val Val Ala Ala Ala Thr Gly
1 5 10 15

50 Val Gln Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20 25 30

Pro Gly Ser Ser Val Lys Val Ser Cys Lys Thr Ser Gly Gly Thr Phe
35 40 45

55 Ser Asn Phe Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu

EP 2 949 673 A1

50

55

60

5

Glu Trp Met Gly Gly Ile Ile Pro Ile Phe Gly Ser Thr Asn Tyr Ala
65 70 75 80

10

Gln Lys Phe Gln Gly Arg Val Thr Ile Asn Ala Asp Glu Ser Thr Ser
85 90 95

15

Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
100 105 110

Tyr Tyr Cys Ala Gly Gly Asp Lys Tyr Gly Pro Tyr Tyr Phe His Tyr
115 120 125

20

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser
130 135 140

25

<210> 62
<211> 407
<212> DNA
<213> Homo sapiens

30

<400> 62
agatctctca ccatggacat gagggtcctc gctcagctcc tggggctcct gctgctctgt 60
ttcccagggtg ccagatgtga catccagatg acccagtctc catcctcact gtctgcatct 120
gtaggagaca gagtcacat cacttgctcg gcgagtcagg gtattagcag ctggttagcc 180
tggtatcagc agaaaccaga gaaagcccct aagtcctga tctatgctgc atccagtttg 240
caaagtgggg tcccatcaag gttcagcggc agtggatctg ggacagattt cactctcacc 300
atcagcagcc tgcagcctga agattttgca acttattact gccaacagta taatagttac 360
ccgtacactt ttggccaggg gaccaagctg gagatcaaac gtacggt 407

40

<210> 63
<211> 132
<212> PRT
<213> Homo sapiens

45

<220>
<221> misc_feature
<222> (132)..(132)
<223> Xaa can be any naturally occurring amino acid

50

<400> 63

Met Asp Met Arg Val Leu Ala Gln Leu Leu Gly Leu Leu Leu Leu Cys
1 5 10 15

55

Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
20 25 30

EP 2 949 673 A1

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45

5 Gln Gly Ile Ser Ser Trp Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys
 50 55 60

10 Ala Pro Lys Ser Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
 65 70 75 80

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 85 90 95

15 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 100 105 110

20 Tyr Asn Ser Tyr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
 115 120 125

25 Lys Arg Thr Xaa
 130

30 <210> 64
 <211> 437
 <212> DNA
 <213> Homo sapiens

<400> 64
 tcgaccccat ggactggacc tggaggttcc tctttgtggt ggcagcagct acaggtgtcc 60
 35 agtcccaggt ccagctggtg cagtctgggg ctgaggtgaa gaagcctggg tcctcgggtga 120
 aggtctcctg caaggcttct ggaggcacct tcagcagcta tgctatcagc tgggtgcgac 180
 agggccctgg acaagggtt gagtgggtgg gagggatcat ccctatcttt ggtacagcaa 240
 40 actacgcaca gaagttccag ggcagagtca cgattaccgc ggacgaatcc acgagcacag 300
 cctacatgga gctgagcagc ctgagatctg aggacacggc cgtgtattac tgtgcgagag 360
 45 gacacaaata tggcccctac tactttgact actggggcca gggaaccctg gtcaccgtct 420
 cctcagctag caccaag 437

50 <210> 65
 <211> 143
 <212> PRT
 <213> Homo sapiens

<400> 65

55 Met Asp Trp Thr Trp Arg Phe Leu Phe Val Val Ala Ala Ala Thr Gly
 1 5 10 15

EP 2 949 673 A1

Val Gln Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20 25 30

5 Pro Gly Ser Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe
35 40 45

10 Ser Ser Tyr Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
50 55 60

Glu Trp Val Gly Gly Ile Ile Pro Ile Phe Gly Thr Ala Asn Tyr Ala
65 70 75 80

15 Gln Lys Phe Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser
85 90 95

20 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
100 105 110

Tyr Tyr Cys Ala Arg Gly His Lys Tyr Gly Pro Tyr Tyr Phe Asp Tyr
115 120 125

25 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
130 135 140

30 <210> 66
<211> 411
<212> DNA
<213> Homo sapiens

35 <400> 66
agatctctca ccatggacat gagggtcctc gctcagctcc tggggctcct gctgctctgt 60
ttcccaggtg ccagatgtga catccagatg acccagtctc catcctcact gtctgcatct 120
gtaggagaca gagtcacat cacttgctcg gcgagtcagg gtattagcag ctggttagcc 180
tggtatcagc agaaaccaga gaaagcccct aagtcctga tctatgctgc atccagtttg 240
caaagtgggg tcccatcaag gttcagcggc agtggatctg ggacagattt cactctcacc 300
45 atcagcagcc tgcagcctga agattttgca acttattact gccaacagta taatagttac 360
cctcggacgt tcggccaagg gaccaaggtg gaaatcaaac gtacggtggc t 411

50 <210> 67
<211> 133
<212> PRT
<213> Homo sapiens

55 <400> 67
Met Asp Met Arg Val Leu Ala Gln Leu Leu Gly Leu Leu Leu Leu Cys
1 5 10 15

EP 2 949 673 A1

Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
20 25 30

5 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
35 40 45

10 Gln Gly Ile Ser Ser Trp Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys
50 55 60

Ala Pro Lys Ser Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
65 70 75 80

15 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

20 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

25 Tyr Asn Ser Tyr Pro Arg Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
115 120 125

Lys Arg Thr Val Ala
130

30 <210> 68
<211> 435
<212> DNA
<213> Homo sapiens

35 <400> 68
tcgaccacca tggactggac ctggagggttc ctctttgtgg tggcagcagc tacaggtgtc 60
cagtcccagg tccagctggg gcagtctggg gctgaggtga agaagcctgg atcctcggtg 120
40 aaggtctcct gcatggcttc aggaggcacc gtcagcagct acgctatcag ctgggtgcga 180
caggccccctg gacaagggct tgagtggatg ggagagatca tccctatctt tggatatagta 240
aactacgcac agaagttcca gggcagagtc acgattaccg cggacgaatc cacgaacaca 300
45 gcctacatgg agctgagcag cctgagatct gaggacacgg ccatatatta ctgtgcgaga 360
gagacagcag tggctggtat tcttggttac tggggccagg gaaccctggt caccgtctcc 420
tcagctagca ccaag 435

50 <210> 69
<211> 142
<212> PRT
55 <213> Homo sapiens
<400> 69

EP 2 949 673 A1

Met Asp Trp Thr Trp Arg Phe Leu Phe Val Val Ala Ala Ala Thr Gly
1 5 10 15

5 Val Gln Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20 25 30

10 Pro Gly Ser Ser Val Lys Val Ser Cys Met Ala Ser Gly Gly Thr Val
35 40 45

15 Ser Ser Tyr Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
50 55 60

Glu Trp Met Gly Glu Ile Ile Pro Ile Phe Gly Ile Val Asn Tyr Ala
65 70 75 80

20 Gln Lys Phe Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Asn
85 90 95

25 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Ile
100 105 110

Tyr Tyr Cys Ala Arg Glu Thr Ala Val Ala Gly Ile Leu Gly Tyr Trp
115 120 125

30 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
130 135 140

35 <210> 70
<211> 414
<212> DNA
<213> Homo sapiens

40 <400> 70
agatctctca ccatggacat gagggctcctc gctcagctcc tggggctcct gctgctctgt 60
ttcccagggtg ccagatgtga catccagatg acccagtctc catcctcact gtctgcatct 120
gtaggagaca gagtcaccat cacttgctcgg gcgagtcagg gtattagcag ctgggttagcc 180
45 tggtatcagc agaaaccaga gaaagccctt aagtccttga tctatgctgc atccagtttg 240
caaagtgggg tcccatcaag gttcagcggc agtggatctg ggacagattt cactctcacc 300
atcagcagcc tgcagcctga agattttgca acttattact gccaacagta taatagttac 360
50 ccgtacactt ttggccaggg gaccaagctg gagatcaaac gtacggtggc tgca 414

55 <210> 71
<211> 134
<212> PRT
<213> Homo sapiens

EP 2 949 673 A1

<400> 71

Met Asp Met Arg Val Leu Ala Gln Leu Leu Gly Leu Leu Leu Leu Cys
1 5 10 15

Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
20 25 30

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
35 40 45

Gln Gly Ile Ser Ser Trp Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys
50 55 60

Ala Pro Lys Ser Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
65 70 75 80

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

Tyr Asn Ser Tyr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
115 120 125

Lys Arg Thr Val Ala Ala
130

<210> 72

<211> 423

<212> DNA

<213> Homo sapiens

<400> 72

cgaccaccca tggaactggg gctccgctgg gttttccttg ttgctatattt agaaggtgtc 60

cagtgtgagg tgcagttggt ggagtctggg ggaggcctgg tcaagcctgg ggggtccctg 120

agactctcct gtgcagcctc tggattcacc ttcagtagcc ataacatgaa ctgggtccgc 180

caggctccag ggaaggggct ggagtgggtc tcatccatta gtagtagtag tagttacata 240

tattatgcag actcagtga gggccgattc accatctcca gagacaacgc caagaactca 300

ctgtatctgc aaatgaacag cctgagagcc gaggacacgg ctgtgtatta ctgtgcgaga 360

gaggactggg gctactttga ctactggggc caggggaaccc tggtcaccgt ctcctcagct 420

agc 423

<210> 73

<211> 140

EP 2 949 673 A1

<212> PRT
<213> Homo sapiens

<400> 73

5 Met Glu Leu Gly Leu Arg Trp Val Phe Leu Val Ala Ile Leu Glu Gly
1 5 10 15

10 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
20 25 30

15 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35 40 45

Ser Ser His Asn Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
50 55 60

20 Glu Trp Val Ser Ser Ile Ser Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala
65 70 75 80

25 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
85 90 95

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100 105 110

30 Tyr Tyr Cys Ala Arg Glu Asp Trp Gly Tyr Phe Asp Tyr Trp Gly Gln
115 120 125

35 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
130 135 140

<210> 74
<211> 407
<212> DNA
<213> Homo sapiens

<400> 74

agatctctca ccatggacat gaggggtcccc gctcagctcc tggggcttct gctgctctgg 60

45 ctcccaggtg ccagatgtgc catccagttg acccagctct catcctccct gtctgcatct 120

gtaggagaca gagtcacat cacttgccgg gcaagtcagg gcattagcag tgatttagcc 180

tggtatcagc agaaaccagg gaaagctcct aagctcctga tctatgatgc ctccagtttg 240

50 gaaagtgggg tcccatcaag gttcagcggc agtggatctg ggacagattt cactctcacc 300

atcagcagcc tgcagcctga agattttgca acttattact gtcaacagtt taatagttac 360

ccattcactt tcggccctgg gaccaaagtg gatatcaaac gtacggt 407

<210> 75

EP 2 949 673 A1

<211> 134
<212> PRT
<213> Homo sapiens

5

<400> 75

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
1 5 10 15

10

Leu Pro Gly Ala Arg Cys Ala Ile Gln Leu Thr Gln Ser Pro Ser Ser
20 25 30

15

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
35 40 45

Gln Gly Ile Ser Ser Asp Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
50 55 60

20

Ala Pro Lys Leu Leu Ile Tyr Asp Ala Ser Ser Leu Glu Ser Gly Val
65 70 75 80

25

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

30

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

Phe Asn Ser Tyr Pro Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile
115 120 125

35

Lys Arg Thr Val Ala Ala
130

40

<210> 76
<211> 433
<212> DNA
<213> Homo sapiens

<400> 76

gtcgaccacc atgggatgga gctggatctt tctctttctc gtgtcaggaa ctggaggtgt 60

45

cctctctgag gtccagctgc aacagtctgg acctgagctg gtgaagcctg gggcttcagt 120

aaagatgtcc tgcaaggctt ctggatacac cttcactgac tactacatga agtgggtgaa 180

50

acagagccat ggaaagagcc ttgagtggat tggagatatt attcctagca atgggtgccac 240

tttctacaac cagaagttca agggcaaggc cactttgact gtggacagat cctccagcac 300

agcctacatg cacctcaaca gcctgacatc tgaggactct gcagtctatt actgtacaag 360

55

atcgcattta ctgcgggcct cctggtttgc ttactggggc caaggggactc tggtcactgt 420

ctctgcagct agc 433

EP 2 949 673 A1

<210> 77
 <211> 141
 <212> PRT
 <213> Homo sapiens

<400> 77

Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Val Ser Gly Thr Gly Gly
 1 5 10 15

Val Leu Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys
 20 25 30

Pro Gly Ala Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35 40 45

Thr Asp Tyr Tyr Met Lys Trp Val Lys Gln Ser His Gly Lys Ser Leu
 50 55 60

Glu Trp Ile Gly Asp Ile Ile Pro Ser Asn Gly Ala Thr Phe Tyr Asn
 65 70 75 80

Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Arg Ser Ser Ser
 85 90 95

Thr Ala Tyr Met His Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val
 100 105 110

Tyr Tyr Cys Thr Arg Ser His Leu Leu Arg Ala Ser Trp Phe Ala Tyr
 115 120 125

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Ala Ser
 130 135 140

<210> 78
 <211> 412
 <212> DNA
 <213> Homo sapiens

<400> 78

agatctcacc atggaatcac agactcaggt cctcatgtcc ctgctgttct gggatatctgg 60
 tacctgtggg gactttgtga tgacacagtc tccatcctcc ctgactgtga cagcaggaga 120
 gaaggtcact atgagctgca agtctagtca gagtctgtta aacagtggaa atcaaaagaa 180
 ctacttgacc tggatatctgc agaaaccagg gcagcctcct aaattgttga tctattgggc 240
 atccactagg gaatctgggg tccctgatcg cttcacaggc agtggatctg gaacagattt 300
 cactctcacc atcagcagtg tgcaggctga agacctggca gtttattact gtcagaatga 360

ttatagttat ccgtacacgt tcggagggggg gaccaagctg gaaataaaac gt

412

<210> 79

<211> 134

<212> PRT

<213> Homo sapiens

<400> 79

Met Glu Ser Gln Thr Gln Val Leu Met Ser Leu Leu Phe Trp Val Ser
1 5 10 15

Gly Thr Cys Gly Asp Phe Val Met Thr Gln Ser Pro Ser Ser Leu Thr
20 25 30

Val Thr Ala Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser
35 40 45

Leu Leu Asn Ser Gly Asn Gln Lys Asn Tyr Leu Thr Trp Tyr Leu Gln
50 55 60

Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg
65 70 75 80

Glu Ser Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp
85 90 95

Phe Thr Leu Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr
100 105 110

Tyr Cys Gln Asn Asp Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr
115 120 125

Lys Leu Glu Ile Lys Arg
130

<210> 80

<211> 37

<212> DNA

<213> Artificial

<220>

<223> primer hCD116Fw-MfeI

<400> 80

cggcaattgc caccatgctt ctcctgggtga caagcct

37

<210> 81

<211> 31

<212> DNA

<213> Artificial

<220>
 <223> primer hCD116Rv-NotI

 <400> 81
 5 attgcggccg ctcaggtaat ttccttcacg g 31

 <210> 82
 <211> 37
 10 <212> DNA
 <213> Artificial

 <220>
 <223> primer hCD116Fw-MfeI

 15 <400> 82
 cggcaattgc caccatgctt ctcctggtga caagcct 37

 <210> 83
 <211> 31
 20 <212> DNA
 <213> Artificial

 <220>
 <223> primer hCD116Rv-NotI

 25 <400> 83
 attgcggccg ctcaggtaat ttccttcacg g 31

 <210> 84
 <211> 37
 30 <212> DNA
 <213> Artificial

 <220>
 <223> primer hCD116Fw-MfeI

 35 <400> 84
 cggcaattgc caccatgctt ctcctggtga caagcct 37

 <210> 85
 <211> 31
 40 <212> DNA
 <213> Artificial

 <220>
 <223> primer hCD116Rv-NotI

 45 <400> 85
 attgcggccg ctcaggtaat ttccttcacg g 31

 50 <210> 86
 <211> 20
 <212> DNA
 <213> Artificial

 55 <220>
 <223> primer hCD116SeqFw1

	<400> 86	
	tgaactgtac ctgggcgagg	20
5	<210> 87	
	<211> 20	
	<212> DNA	
	<213> Artificial	
10	<220>	
	<223> primer hCD116SeqFw2	
	<400> 87	
	ctggcacgga aaacctactg	20
15	<210> 88	
	<211> 20	
	<212> DNA	
	<213> Artificial	
20	<220>	
	<223> primer hCD116SeqRv1	
	<400> 88	
	cctgaatttg gataaagcag	20
25	<210> 89	
	<211> 20	
	<212> DNA	
	<213> Artificial	
30	<220>	
	<223> primer T7	
	<400> 89	
35	taatacgact cactataggg	20
	<210> 90	
	<211> 40	
	<212> DNA	
40	<213> Artificial	
	<220>	
	<223> primer hCD123-C-FLAG-R1	
45	<400> 90	
	tcgtcatcgt ccttgtagtc agttttctgc acgacctgta	40
	<210> 91	
	<211> 34	
50	<212> DNA	
	<213> Artificial	
	<220>	
	<223> primer IL-3RA_Fw	
55	<400> 91	
	cggcaattgc caccatgggc ctcctttggc tcac	34

5
 <210> 92
 <211> 39
 <212> DNA
 <213> Artificial
 <220>
 <223> primer C-FLAG-NotR2
 <400> 92
 10 aaaagcggcc gctcacttgt cgtcatcgtc cttgtagtc 39
 <210> 93
 <211> 20
 <212> DNA
 15 <213> Artificial
 <220>
 <223> primer pEGFP-N1-Fw
 <400> 93
 20 cgtgtacggg gggaggtcta 20
 <210> 94
 <211> 19
 25 <212> DNA
 <213> Artificial
 <220>
 <223> primer pEGFP-N1-Re
 30 <400> 94
 tttatgtttc aggttcagg 19
 <210> 95
 35 <211> 20
 <212> DNA
 <213> Artificial
 <220>
 <223> primer pEGFP-N1-Fw
 40 <400> 95
 cgtgtacggg gggaggtcta 20
 <210> 96
 45 <211> 19
 <212> DNA
 <213> Artificial
 <220>
 50 <223> primer pEGFP-N1-Re
 <400> 96
 tttatgtttc aggttcagg 19
 55 <210> 97
 <211> 28

<212> DNA
 <213> Artificial

5 <220>
 <223> primer CD123R11pEGFPN1

<400> 97
 aaaggtaccg aattcgaagc ttgagctc 28

10 <210> 98
 <211> 27
 <212> DNA
 <213> Artificial

15 <220>
 <223> primer CD123F11

<400> 98
 aaaggtaccg ggaagccttg ggcaggt 27

20 <210> 99
 <211> 30
 <212> DNA
 <213> Artificial

25 <220>
 <223> primer CD123R12-2

30 <400> 99
 aaaggtacca ctgttctcag ggaagaggat 30

<210> 100
 <211> 29
 <212> DNA
 <213> Artificial

35 <220>
 <223> primer CD123F12-2

40 <400> 100
 aaaggtaccc agattgagat attaactcc 29

45 <210> 101
 <211> 27
 <212> DNA
 <213> Artificial

<220>
 <223> primer CD123R13

50 <400> 101
 aaaggtacct gaaaagacga caaactt 27

55 <210> 102
 <211> 27
 <212> DNA
 <213> Artificial

<220>
 <223> primer CD123F13

 <400> 102
 5 aaaggtacct cgctgctgat cgcgctg 27

 <210> 103
 <211> 20
 10 <212> DNA
 <213> Artificial

 <220>
 <223> primer pEGFP-N1-Fw

 15 <400> 103
 cgtgtacggg gggaggtcta 20

 <210> 104
 <211> 19
 20 <212> DNA
 <213> Artificial

 <220>
 <223> primer pEGFP-N1-Re

 25 <400> 104
 tttatgtttc aggttcagg 19

 <210> 105
 <211> 33
 30 <212> DNA
 <213> Artificial

 <220>
 <223> primer GM-CSFRF11

 35 <400> 105
 aaaggtaccg ccaccatgct tctcctggtg aca 33

 <210> 106
 <211> 27
 40 <212> DNA
 <213> Artificial

 <220>
 <223> primer GM-CSFR11

 45 <400> 106
 aaaggtacct gaatttggat aaagcag 27

 50 <210> 107
 <211> 27
 <212> DNA
 <213> Artificial

 55 <220>
 <223> primer GM-CSFRF12

<400> 107
 aaaggtaccg gaaggaggagg taccgct 27

5 <210> 108
 <211> 27
 <212> DNA
 <213> Artificial

10 <220>
 <223> primer GM-CSFR12
 <400> 108
 aaaggtaccc tttgtgtcca aaagtga 27

15 <210> 109
 <211> 27
 <212> DNA
 <213> Artificial

20 <220>
 <223> primer GM-CSFR13
 <400> 109
 aaaggtacca aaatagaacg attcaac 27

25 <210> 110
 <211> 27
 <212> DNA
 <213> Artificial

30 <220>
 <223> primer GM-CSFR13
 <400> 110
 aaaggtacca atgtacacag agccgag 27

35 <210> 111
 <211> 20
 <212> DNA
 <213> Artificial

40 <220>
 <223> primer pEGFP-N1-Fw
 <400> 111
 cgtgtacggt gggaggtcta 20

45 <210> 112
 <211> 19
 <212> DNA
 <213> Artificial

50 <220>
 <223> primer pEGFP-N1-Re
 <400> 112
 tttatgtttc aggttcagg 19

55

5
 <210> 113
 <211> 5
 <212> PRT
 <213> Homo sapiens
 <400> 113
 Thr Tyr Ala Ile Ser
 1 5

10
 <210> 114
 <211> 17
 <212> PRT
 <213> Homo sapiens
 <400> 114
 Gly Ile Ile Pro Ile Phe Gly Ile Val Asn Tyr Ala Gln Lys Phe Gln
 1 5 10 15

20
 Gly
 25
 <210> 115
 <211> 11
 <212> PRT
 <213> Homo sapiens
 <400> 115
 Gly Gly Gly Ser Gly Pro Asp Val Leu Asp Ile
 1 5 10

35
 <210> 116
 <211> 5
 <212> PRT
 <213> Homo sapiens
 <400> 116
 Thr Tyr Ala Ile Ser
 1 5

45
 <210> 117
 <211> 17
 <212> PRT
 <213> Homo sapiens
 <400> 117
 Gly Leu Ile Pro Ile Phe Asp Ile Glu Asn Tyr Ala Gln Lys Phe Gln
 1 5 10 15

55
 Gly

EP 2 949 673 A1

<210> 118
 <211> 11
 <212> PRT
 <213> Homo sapiens

5

<400> 118

Gly Gly Gly Ser Gly Pro Asp Val Leu Asp Ile
 1 5 10

10

<210> 119
 <211> 5
 <212> PRT
 <213> Homo sapiens

15

<400> 119

Asn Phe Ala Ile Ser
 1 5

20

<210> 120
 <211> 17
 <212> PRT
 <213> Homo sapiens

25

<400> 120

Gly Ile Ile Pro Ile Phe Gly Ser Thr Asn Tyr Ala Gln Lys Phe Gln
 1 5 10 15

30

Gly

35

<210> 121
 <211> 11
 <212> PRT
 <213> Homo sapiens

40

<400> 121

Gly Asp Lys Tyr Gly Pro Tyr Tyr Phe His Tyr
 1 5 10

45

<210> 122
 <211> 5
 <212> PRT
 <213> Homo sapiens

50

<400> 122

Ser Tyr Ala Ile Ser
 1 5

55

<210> 123
 <211> 17
 <212> PRT

EP 2 949 673 A1

<213> Homo sapiens

<400> 123

5 Gly Ile Ile Pro Ile Phe Gly Thr Ala Asn Tyr Ala Gln Lys Phe Gln
1 5 10 15

Gly

10

<210> 124

<211> 11

<212> PRT

15 <213> Homo sapiens

<400> 124

20 Gly His Lys Tyr Gly Pro Tyr Tyr Phe Asp Tyr
1 5 10

<210> 125

<211> 5

<212> PRT

25 <213> Homo sapiens

<400> 125

30 Ser Tyr Ala Ile Ser
1 5

<210> 126

<211> 17

<212> PRT

35 <213> Homo sapiens

<400> 126

40 Glu Ile Ile Pro Ile Phe Gly Ile Val Asn Tyr Ala Gln Lys Phe Gln
1 5 10 15

Gly

45 <210> 127

<211> 10

<212> PRT

<213> Homo sapiens

<400> 127

50 Glu Thr Ala Val Ala Gly Ile Leu Gly Tyr
1 5 10

55 <210> 128

<211> 5

<212> PRT

EP 2 949 673 A1

<213> Homo sapiens
 <400> 128
 5 Ser His Asn Met Asn
 1 5
 <210> 129
 10 <211> 17
 <212> PRT
 <213> Homo sapiens
 <400> 129
 15 Ser Ile Ser Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15
 Gly
 20
 <210> 130
 <211> 7
 <212> PRT
 25 <213> Homo sapiens
 <400> 130
 Glu Asp Trp Gly Tyr Phe Asp
 1 5
 30
 <210> 131
 <211> 11
 <212> PRT
 35 <213> Homo sapiens
 <400> 131
 Arg Met Ser Gln Gly Ile Arg Ser Tyr Leu Ala
 1 5 10
 40
 <210> 132
 <211> 7
 <212> PRT
 45 <213> Homo sapiens
 <400> 132
 Ala Ala Ser Thr Leu Gln Ser
 1 5
 50
 <210> 133
 <211> 9
 <212> PRT
 55 <213> Homo sapiens
 <400> 133

EP 2 949 673 A1

Gln Gln Tyr Tyr Ser Phe Pro Tyr Thr
1 5

5
<210> 134
<211> 11
<212> PRT
<213> Homo sapiens

10
<400> 134

Arg Met Ser Gln Gly Ile Arg Ser Tyr Leu Ala
1 5 10

15
<210> 135
<211> 7
<212> PRT
<213> Homo sapiens

20
<400> 135

Ala Ala Ser Thr Leu Gln Ser
1 5

25
<210> 136
<211> 9
<212> PRT
<213> Homo sapiens

30
<400> 136

Gln Gln Tyr Tyr Ser Phe Pro Tyr Thr
1 5

35
<210> 137
<211> 11
<212> PRT
<213> Homo sapiens

40
<400> 137

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
1 5 10

45
<210> 138
<211> 7
<212> PRT
<213> Homo sapiens

50
<400> 138

Ala Ala Ser Ser Leu Gln Ser
1 5

55
<210> 139
<211> 9
<212> PRT
<213> Homo sapiens

EP 2 949 673 A1

<400> 139

Gln Gln Tyr Asn Ser Tyr Pro Tyr Thr
1 5

<210> 140

<211> 11

<212> PRT

<213> Homo sapiens

<400> 140

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
1 5 10

<210> 141

<211> 7

<212> PRT

<213> Homo sapiens

<400> 141

Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 142

<211> 9

<212> PRT

<213> Homo sapiens

<400> 142

Gln Gln Tyr Asn Ser Tyr Pro Arg Thr
1 5

<210> 143

<211> 11

<212> PRT

<213> Homo sapiens

<400> 143

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
1 5 10

<210> 144

<211> 7

<212> PRT

<213> Homo sapiens

<400> 144

Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 145

<211> 9
 <212> PRT
 <213> Homo sapiens
 5 <400> 145
 Gln Gln Tyr Asn Ser Tyr Pro Tyr Thr
 1 5
 10 <210> 146
 <211> 11
 <212> PRT
 <213> Homo sapiens
 15 <400> 146
 Arg Ala Ser Gln Gly Ile Ser Ser Asp Leu Ala
 1 5 10
 20 <210> 147
 <211> 7
 <212> PRT
 <213> Homo sapiens
 25 <400> 147
 Asp Ala Ser Ser Leu Glu Ser
 1 5
 30 <210> 148
 <211> 9
 <212> PRT
 <213> Homo sapiens
 35 <400> 148
 Gln Gln Phe Asn Ser Tyr Pro Phe Thr
 1 5
 40 <210> 149
 <211> 29
 <212> DNA
 <213> Artificial
 45 <220>
 <223> primer CD123-Fw21
 <400> 149
 cgtggaaccc gcagtgaaca atagctatt 29
 50 <210> 150
 <211> 30
 <212> DNA
 <213> Artificial
 55 <220>
 <223> primer CD123-Re21

	<400> 150 actctgttct ttttaacaca ctcgatatcg	30
5	<210> 151 <211> 29 <212> DNA <213> Artificial	
10	<220> <223> primer CD123-Fw22	
	<400> 151 ctttatccaa ataacagtgg gaagccttg	29
15	<210> 152 <211> 31 <212> DNA <213> Artificial	
20	<220> <223> primer CD123-Re22	
	<400> 152 cagtttctgt tggaatggtg ggttggccac t	31
25	<210> 153 <211> 34 <212> DNA <213> Artificial	
30	<220> <223> primer CD123-Fw23	
35	<400> 153 agggagggtgta ccggtgcgga gaatctgacc tgct	34
40	<210> 154 <211> 32 <212> DNA <213> Artificial	
	<220> <223> primer CD123-Re23	
45	<400> 154 tcctgaattt ggatagaaga ggatccacgt gg	32
50	<210> 155 <211> 29 <212> DNA <213> Artificial	
	<220> <223> primer CD123-Fw24	
55	<400> 155 ggtccgacgg cccccgcgga cgtccagta	29

5	<210> 156 <211> 32 <212> DNA <213> Artificial	
	<220> <223> primer CD123-Re24	
10	<400> 156 cctcgcccag gtacagctca agaatccac gt	32
15	<210> 157 <211> 30 <212> DNA <213> Artificial	
	<220> <223> primer CD123-Fw25	
20	<400> 157 acggaaccag cgcagccttc ggtatcccct	30
25	<210> 158 <211> 28 <212> DNA <213> Artificial	
	<220> <223> primer CD123-Re25	
30	<400> 158 taaccagaaa gtgggaactt tgagaacc	28
35	<210> 159 <211> 28 <212> DNA <213> Artificial	
	<220> <223> primer CD123-Fw26	
40	<400> 159 tctttgattc atttgtcgtc ttttcaca	28
45	<210> 160 <211> 29 <212> DNA <213> Artificial	
50	<220> <223> primer CD123-Re26	
55	<400> 160 attggatgcc gaaggctgcg ctccctgccc	29

Claims

1. A method for producing an antibody to a human IL-3R α chain which does not inhibit IL-3 signaling and binds to B domain of the human IL-3R α chain but does not bind to C domain of the human IL-3R α chain.
2. The method according to claim 1, wherein the human IL-3R α antibody further has high antibody-dependent cellular cytotoxicity (ADCC).
3. The method according to claim 1 or 2, wherein the high antibody-dependent cellular cytotoxicity (ADCC) shows a specific lysis rate of 10% at an antibody concentration of 0.01 μ g/ml, by a Colon-26/hCD123 ADCC assay method using PBMC cultured with IL-2.
4. The method according to any one of claims 1 to 3, wherein the human IL-3R α antibody comprises amino acid sequences of CDRs of heavy chain and CDRs of light chain selected from the group consisting of the following (a) to (e);
 - (a) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:113 to 115, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs:131 to 133, respectively,
 - (b) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:116 to 118, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs: 134 to 136, respectively,
 - (c) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:119 to 121, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs:137 to 139, respectively,
 - (d) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:122 to 124, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs:140 to 142, respectively, and
 - (e) CDR 1 to 3 of heavy chain are the amino acid sequences of SEQ ID NOs:125 to 127, respectively, and CDR 1 to 3 of light chain are the amino acid sequences of SEQ ID NOs:143 to 145, respectively.
5. The method according to any one of claims 1 to 4, wherein the human IL-3R α antibody comprises a heavy chain variable region and a light chain variable region selected from the group consisting of the following (a) to (e);
 - (a) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:53 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:55;
 - (b) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:57 and a light chain variable region comprising an amino acid sequence from valine (V) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:59;
 - (c) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:61 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:63;
 - (d) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 139 in the amino acid sequence of SEQ ID NO:65 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:67; and
 - (e) a heavy chain variable region comprising an amino acid sequence from glutamine (Q) at position 20 to serine (S) at position 138 in the amino acid sequence of SEQ ID NO:69 and a light chain variable region comprising an amino acid sequence from aspartic acid (D) at position 23 to lysine (K) at position 129 in the amino acid sequence of SEQ ID NO:71.
6. The method of any one of claims 1 to 5, wherein the method comprises:
 - administering human IL-3R α , a subsequence thereof or a fragment thereof to animals which can express human immunoglobulin;
 - screening the animals for their expression of human IL-3R α antibody;
 - selecting an animal which produces human IL-3R α antibody; and
 - isolating the human IL-3R α antibody from the selected animal.

7. The method of claim 6, wherein the human IL-3R α , subsequence thereof or fragment thereof is conjugated with a human Fc recombinant protein.

8. The method of claim 6 or 7, wherein the fragment of human IL-3R α is the IL-3R α extracellular region.

9. The method according to any one of claims 1 to 5, wherein the human IL-3R α antibody is produced by a recombinant technique.

10. The method according to any one of claims 1 to 5, wherein the method comprises:

- transforming a host cell with vector comprising a nucleic acid sequence encoding the antibody;
- expressing the antibody by growing the cell; and
- purifying the antibody.

11. Use of the IL-3R α antibody as defined in any one of claims 1 to 5 for the manufacture of a pharmaceutical composition for use in a method for preventing or treating a blood tumor in which a cell expressing IL-3R α is found in bone marrow or peripheral blood of a subject.

12. The use according to claim 11, wherein the blood tumor is acute myeloid leukemia (AML).

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9





Fig. 10





EUROPEAN SEARCH REPORT

Application Number
EP 15 17 2601

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	SUN Q ET AL: "Monoclonal Antibody 7G3 Recognizes the N-Terminal Domain of the Human Interleukin-3 (IL-3) Receptor alpha-Chain and Functions as a Specific IL-3 Receptor Antagonist", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 87, no. 1, 1 January 1996 (1996-01-01), pages 83-92, XP008154678, ISSN: 0006-4971 * the whole document *	1-12	INV. C07K16/28 A61K39/395 A61P35/02 C07K16/46 G01N33/53 G01N33/574 G01N33/577 C12N15/02 C12N15/09 C12P21/08 C12R1/91
A	WO 97/24373 A1 (MEDVET SCIENCE PTY LIMITED [AU]; LOPEZ ANGEL F [AU]) 10 July 1997 (1997-07-10) * page 3, lines 22-23; example 15 *	1-12	
A	ROBERTS ANDREW W ET AL: "A Phase 1 and Correlative Biological Study of CSL360 (anti-CD123 mAb) in AML", BLOOD, vol. 112, no. 11, November 2008 (2008-11), pages 1015-1016, XP009161393, & 50TH ANNUAL MEETING OF THE AMERICAN-SOCIETY-OF-HEMATOLOGY; SAN FRANCISCO, CA, USA; DECEMBER 06 -09, 2008 ISSN: 0006-4971 * the whole document *	1-12	TECHNICAL FIELDS SEARCHED (IPC) C07K A61K
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 14 September 2015	Examiner Domingues, Helena
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



EUROPEAN SEARCH REPORT

 Application Number
 EP 15 17 2601

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	ISHIDA I ET AL: "Production of human monoclonal and polyclonal antibodies in TransChromo animals", CLONING AND STEM CELLS, MARY ANN LIEBERT, LARCHMONT, US, vol. 4, no. 1, 1 January 2002 (2002-01-01), pages 91-102, XP002970309, ISSN: 1536-2302, DOI: 10.1089/153623002753632084 * the whole document *	1-12	
A	WO 02/43478 A2 (MEDAREX INC [US]; KIRIN BREWERY [JP]; TOMIZUKA KAZUMA [JP]; ISHIDA ISA) 6 June 2002 (2002-06-06) * examples 1-12 *	1-12	
A	WO 2007/005874 A2 (MEDAREX INC [US]; KORMAN ALAN J [US]; SELBY MARK J [US]; WANG CHANGYU) 11 January 2007 (2007-01-11) * example 1 *	1-12	
A	LONBERG N: "Human monoclonal antibodies from transgenic mice", HANDBOOK OF EXPERIMENTAL PHARMACOLOGY, SPRINGER VERLAG, BERLIN, DE, vol. 181, 1 January 2008 (2008-01-01), pages 69-97, XP009099873, ISSN: 0171-2004, DOI: 10.1007/978-3-540-73259-4_4 * the whole document *	1-12	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (IPC)
Place of search Munich		Date of completion of the search 14 September 2015	Examiner Domingues, Helena
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 15 17 2601

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-09-2015

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9724373 A1	10-07-1997	US 6177078 B1	23-01-2001
		WO 9724373 A1	10-07-1997
WO 0243478 A2	06-06-2002	AT 378403 T	15-11-2007
		AU 3942202 A	11-06-2002
		AU 2002239422 B2	07-12-2006
		CA 2430013 A1	06-06-2002
		CN 1487996 A	07-04-2004
		CN 101940189 A	12-01-2011
		DE 60131456 T2	10-07-2008
		DK 1354034 T3	25-03-2008
		EP 1354034 A2	22-10-2003
		EP 1916303 A1	30-04-2008
		ES 2295228 T3	16-04-2008
		ES 2405944 T3	04-06-2013
		HK 1064708 A1	15-10-2010
		IL 155977 A	15-06-2009
		JP 3523245 B1	26-04-2004
		JP 2004515230 A	27-05-2004
		KR 20030074634 A	19-09-2003
		MX PA03004793 A	03-12-2004
		PT 1354034 E	28-02-2008
		TW I313299 B	11-08-2009
		US 2002199213 A1	26-12-2002
		US 2006026698 A1	02-02-2006
		US 2006037093 A1	16-02-2006
		US 2011003319 A1	06-01-2011
		US 2014380515 A1	25-12-2014
		WO 0243478 A2	06-06-2002
WO 2007005874 A2	11-01-2007	AU 2006265108 A1	11-01-2007
		BR PI0613361 A2	04-01-2011
		CA 2612241 A1	11-01-2007
		CN 101248089 A	20-08-2008
		CN 104356236 A	18-02-2015
		EA 200800229 A1	30-06-2008
		EP 1907424 A2	09-04-2008
		HR P20080053 A2	31-08-2009
		JP 5252635 B2	31-07-2013
		JP 2008544755 A	11-12-2008
		JP 2013150606 A	08-08-2013
		KR 20080045674 A	23-05-2008
		KR 20140002041 A	07-01-2014
		KR 20150082674 A	15-07-2015
		NZ 564592 A	25-11-2011
		SG 163554 A1	30-08-2010

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 15 17 2601

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-09-2015

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		UA 99701 C2	25-09-2012
		US 2009055944 A1	26-02-2009
		US 2013122014 A1	16-05-2013
		WO 2007005874 A2	11-01-2007
		ZA 200710919 A	29-10-2008

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 120694 A [0023]
- EP 125023 A [0023]
- GB 2188638 A [0023]
- US 5585089 A [0023] [0098] [0099]
- US 4036945 A [0076]
- US 4331647 A [0076]
- US 4946778 A [0077]
- US 5258498 A [0077]
- US 6087329 A [0084]
- US 5939598 A [0095] [0096]
- WO 0243478 A [0095] [0103]
- WO 02092812 A [0095] [0103]
- WO 9824893 A [0096]
- WO 9201047 A [0096]
- WO 9634096 A [0096]
- WO 9633735 A [0096]
- US 5413923 A [0096]
- US 5625126 A [0096]
- US 5633425 A [0096]
- US 5569825 A [0096]
- US 5661016 A [0096]
- US 5545806 A [0096]
- US 5814318 A [0096]
- US 5885793 A [0096]
- US 5916771 A [0096]
- EP 239400 A [0098]
- WO 9109967 A [0098]
- US 5225539 A [0098]
- US 5530101 A [0098]
- EP 592106 A [0098]
- EP 519596 A [0098]
- US 5565332 A [0098]
- US 5807715 A [0101]
- US 4816567 A [0101]
- US 4816397 A [0101]
- US 4902614 A [0102]
- US 4543439 A [0102]
- US 4411993 A [0102]
- WO 2005035586 A [0107]
- WO 200231140 A [0107]
- WO 0061739 A [0107]
- US 20070148165 A [0109]
- US 6737056 B [0109]
- US 7297775 B [0109]
- US 7317091 B [0109]
- US 5624820 A [0118]
- US 5693508 A [0118]
- US 5665577 A [0118]
- US 6013516 A [0118]
- US 5674703 A [0118]
- WO 9205266 A [0118]
- WO 9214829 A [0118]
- US 5700470 A [0118]
- US 5731172 A [0118]
- US 5928944 A [0118]
- US 5604090 A [0118]
- US 5501979 A [0118]
- US 5561063 A [0118]
- US 5719054 A [0118]
- US 4844904 A [0130]
- US 5000959 A [0130]
- US 4863740 A [0130]
- US 4975282 A [0130]
- US 5861397 A [0130]
- US 5459127 A [0130]
- US 6001358 A [0183]

Non-patent literature cited in the description

- **OSAWA M et al.** *Science*, 1996, vol. 273:2, 42-5 [0024]
- **GOODELL MA et al.** *J Exp Med.*, 1996, vol. 183, 1797-806 [0024]
- **YAMAZAKI S et al.** *EMBO J.*, 2006, vol. 25, 3515-23 [0024]
- **ISHIKAWA F et al.** *Nat Biotechnol.*, 2007, vol. 25, 1315-21 [0024]
- **BAO S et al.** *Nature*, 2006, vol. 444, 756-60 [0024]
- **SCHIFF et al.** *Canc. Res.*, 1985, vol. 45, 879-885 [0024]
- **BRUGGEMANN et al.** *J. Exp. Med.*, 1989, vol. 170, 2153-2157 [0024]
- **RIECHMANN et al.** *Nature*, 1988, vol. 332, 323-327 [0024]
- **ISHIDA I et al.** *Cloning Stem Cells*, 2002, vol. 4, 91-102 [0024]
- **WU et al.** *J Mol Biol.*, 1999, vol. 19, 151-62 [0024]
- **SUN et al.** *Blood*, 1996, vol. 87, 83 [0024]
- **CHEN et al.** *JBiol Chem*, 2009, vol. 284, 5763 [0024]
- **BONNET et al.** *Nat Med*, 1997, vol. 3, 730 [0024]
- **JORDAN et al.** *Leukemia*, 2000, vol. 14, 1777 [0024]
- *Haematologica*, 2001, vol. 86, 1261 [0024]

- *LeukLymphoma*, 2006, vol. 47, 207 [0024]
- **MAJETI et al.** *Proc Natl Acad Sci USA.*, 2009, vol. 106, 3396 [0024]
- *Blood*, 2008, vol. 112 (11 [0024]
- **ALTSCHUL et al.** *J. Mol. Biol.*, 1990, vol. 215, 403 [0069]
- **PEARSON et al.** *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 2444 [0069]
- **PEARSON.** *Methods Mol. Bio.*, 2000, vol. 132, 185 [0069]
- **SMITH et al.** *J. Mol. Biol.*, 1981, vol. 147, 195 [0069]
- **BOSTICK et al.** *Biochem. Biophys. Res. Commun.*, 2003, vol. 304, 320 [0069]
- **EDELMAN et al.** *Methods Enzymol.*, 1967, vol. 1, 422 [0076]
- **WHITLOW et al.** *Methods: A Companion to Methods in Enzymology*, 1991, vol. 2, 97 [0077]
- **BIRD et al.** *Science*, 1988, vol. 242, 423 [0077]
- **HUSTON et al.** *Methods Enzymol*, 1991, vol. 203, 46 [0077]
- **SHU et al.** *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 7995 [0077]
- **SKERRA et al.** *Science*, 1988, vol. 240, 1038 [0077]
- **KUROIWA et al.** *Nat. Biotechnol.*, 2002, vol. 20, 889 [0096]
- **KABAT.** Sequences of Proteins of Immunological Interest. US Department of Health and Human Services, 1987 [0097]
- **PADLAN.** *Mol. Immunol.*, 1994, vol. 31, 169 [0097]
- **PADLAN.** *Mol. Immunol.*, 1991, vol. 28, 489 [0097]
- **PADLAN.** *Molecular Immunol.*, vol. 28, 489 [0098]
- **STUDNICKA et al.** *Protein Engineering* 7, 1994, 805 [0098]
- **ROGUSKA et al.** *Proc. Nat'l Acad. Sci. USA*, 1994, vol. 91, 969 [0098]
- **CARTER et al.** *Proc. Natl. Acad. Sci. USA*, 1992, vol. 89, 4285 [0098]
- **PRESTA et al.** *J. Immunol.*, 1993, vol. 151, 2623 [0098]
- **RIECHMANN et al.** *Nature*, 1988, vol. 332, 323 [0099]
- **MORRISON.** *Science*, 1985, vol. 229, 1202 [0101]
- **OI et al.** *BioTechniques*, 1986, vol. 4, 214 [0101]
- **GILLIES et al.** *J. Immunol. Methods*, 1989, vol. 125, 191 [0101]
- **MUNRO.** *Nature*, 1984, vol. 312, 597 [0101]
- **NEUBERGER et al.** *Nature*, 1984, vol. 312, 604 [0101]
- **SHARON et al.** *Nature*, 1984, vol. 309, 364 [0101]
- **MORRISON et al.** *Proc. Nat'l. Acad. Sci. USA*, 1984, vol. 81, 6851 [0101]
- **BOULIANNE et al.** *Nature*, 1984, vol. 312, 643 [0101]
- **CAPON et al.** *Nature*, 1989, vol. 337, 525 [0101]
- **TRAUNECKER et al.** *Nature*, 1989, vol. 339, 68 [0101]
- **KENNETT ; MCKEARN ; BECHTOL et al.** Monoclonal Antibodies. Hybridomas: A New Dimension in Biological Analyses. Plenum Press, 1980 [0102]
- **HARLOW et al.** Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1988 [0102]
- **ISHIDA et al.** *IBC's 11th Antibody Engineering Meeting*, 2000 [0103]
- **WIGLER et al.** *Cell*, 1977, vol. 11, 223 [0117]
- **CONE et al.** *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 6349 [0118]
- Eukaryotic Viral Vectors. Cold Spring Harbor Laboratory, 1982 [0118]
- **SARVER et al.** *Mol. Cell. Biol.*, 1981, vol. 1, 486 [0118]
- **AUSUBEL et al.** Current Protocols in Molecular Biology. Greene Publish. Assoc. & Wiley Interscience, 1988, vol. 2 [0122]
- **GRANT et al.** *Methods in Enzymology*. Acad. Press, 1987, vol. 153, 516-544 [0122]
- **GLOVER.** DNA Cloning. IRL Press, 1986, vol. 11 [0122]
- **BITTER.** *Methods in Enzymology*. Acad. Press, 1987, vol. 152, 673-684 [0122]
- **STRATHERN et al.** The Molecular Biology of the Yeast *Saccharomyces*. Cold Spring Harbor Press, 1982, 1, , 2 [0122]
- **RICH ; MYSZKA.** *Curr. Opin. Biotechnol.*, 2000, vol. 11, 54 [0132]
- **ENGLEBIENNE.** *Analyst*, 1998, vol. 123, 1599 [0132]
- **MALMQVIST.** *Biochem. Soc. Trans.*, 1999, vol. 27, 335 [0132]
- **TOMIZUKA et al.** *Proc. Natl. Acad. Sci. USA*, 2000, vol. 97, 722 [0164]
- **FISHWILD et al.** *Nat. Biotechnol*, 1996, vol. 14, 845 [0164]
- **ISHIDA ; LONBERG.** *IBC's 11th Antibody Engineering*, 2000 [0165]
- A Guide to Monoclonal Antibody Experimental Operations. Kodansha, 1991 [0166]
- The Cytokine Facts Book. Academic Press [0244]

专利名称(译)	抗il-3ra抗体用于治疗血肿瘤		
公开(公告)号	EP2949673A1	公开(公告)日	2015-12-02
申请号	EP2015172601	申请日	2010-04-27
[标]申请(专利权)人(译)	协和酸酵麒麟有限公司		
申请(专利权)人(译)	协和发酵工业株式会社CO. , LTD.		
当前申请(专利权)人(译)	协和发酵工业株式会社CO. , LTD.		
[标]发明人	TAWARA TOMONORI TAKAYANAGI SHINICHIRO INAGAKI YOSHIMASA		
发明人	TAWARA, TOMONORI TAKAYANAGI, SHINICHIRO INAGAKI, YOSHIMASA		
IPC分类号	C07K16/28 A61K39/395 A61P35/02 C07K16/46 G01N33/53 G01N33/574 G01N33/577 C12N15/02 C12N15/09 C12P21/08 C12R1/91		
CPC分类号	C07K16/2866 C07K2317/732 C07K2317/75 G01N33/57407 G01N33/57426 G01N33/57492 C07K16 /3061 C07K2317/21 C07K2317/33 C07K2317/51 C07K2317/515 C07K2317/56 C07K2317/565 C07K2317/92 G01N2333/7155		
审查员(译)	DOMINGUES , HELENA		
优先权	61/172923 2009-04-27 US		
外部链接	Espacenet		

摘要(译)

本发明提供了针对人IL-3Rα链的抗体，其不抑制IL-3信号传导并结合人IL-3Rα链的B结构域，但不结合人IL-3Rα链的C结构域。 1.一种用于预防或治疗血肿瘤的组合物，其中在受试者的骨髓或外周血中发现表达IL-3Rα的细胞，其包含人IL-3Rα抗体作为活性成分;以及治疗血肿瘤的方法，其中在骨髓或外周血中发现表达IL-3Rα的细胞，该方法包括给受试者施用包含IL-3Rα抗体作为活性成分的组合物。

MVLLWLTLLIALPCLLQTKEDPNPPITNLRMKAKAQQLTWDLNRNVTDIECVK
DADYSMPAVNNSYCQFGAISLCEVTNYTVRVANPPFSTWLFPENSGKPPWAGAE
NLTCWIHDVDFLSCSWAVGPGAPADVQYDLYLVANRRQQYECLHYKTDAGG
TRIGCRFDDISRLSSGSQSSHILVRGRSAAFGHPCTDKFVVFQIEILTPPNMTAKC
NKTHSFMIHWKMRSHFNKFRYELQIQKMQPVITEQVRDRTSFOLLNPGTYTV