

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 July 2008 (10.07.2008)

PCT

(10) International Publication Number
WO 2008/082974 A2

- (51) International Patent Classification:
C12Q 1/00 (2006.01)
- (21) International Application Number:
PCT/US2007/088070
- (22) International Filing Date:
19 December 2007 (19.12.2007)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
11/618,495 29 December 2006 (29.12.2006) US
- (71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; Dept 0377, Bldg Ap6a-1a, 100 Abbott Park Rd, Abbott Park, IL 60064 (US).
- (71) Applicants and
(72) Inventors: **GRENIER, Frank C** [US/US]; 218 S Second Avenue, Libertyville, Illinois 60048 (US). **WORKMAN, Ryan F** [US/US]; 1296 S Pleasant Hill Court, Libertyville, IL 60048 (US). **SYED, Hina N** [IN/US]; 1164 Vista Drive, Gurnee, Illinois 60031 (US). **ALI, Salman** [US/US]; 5 Kingman Lane, Hoffman Estates, Illinois 60169 (US).
- (74) Agent: **BARTNICKI, Audrey L.**; Dept. 377 Bldg AP6A-1, 100 Abbott Park Road, Abbott Park, Illinois 60064 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report



WO 2008/082974 A2

(54) Title: NON-DENATURING LYSIS REAGENT

(57) Abstract: The invention provides a lysis reagent and method for preparing a test sample for use in an assay, wherein the method yields a homogeneous lysis mixture suitable for use in automated pipetting systems without the need for a centrifugation step. The lysis reagent includes a glycol and an alcohol. Other aspects of the invention include related immunoassays and test kits.

NON-DENATURING LYSIS REAGENT

TECHNICAL FIELD

This invention relates to a non-denaturing lysis reagent useful, for example, in diagnostic immunoassays to determine the concentration levels of an immunosuppressant drug in a test sample.

BACKGROUND

Many analytes of clinical interest are taken up by cells or become complexed with one or more other components of the test sample. Accordingly, to obtain an accurate measurement of the amount of analyte present in the sample, it is preferable to treat the sample, and/or conduct the assay under conditions, such that the analyte is released from the cells or other component(s) for detection in the assay.

For example, immunosuppressant drugs such as tacrolimus, everolimus, temsolorimus and cyclosporine are effective for the treatment of organ or tissue rejection following transplant surgery, of graft versus host disease and of autoimmune diseases in humans. During immunosuppressant drug therapy, monitoring the blood concentration levels of the immunosuppressant is an important aspect of clinical care because insufficient drug levels lead to graft (organ or tissue) rejection and excessive levels lead to undesired side effects and toxicities. Blood levels of immunosuppressant are therefore measured so drug dosages can be adjusted to maintain the drug level at the appropriate concentration. Diagnostic assays for determination of immunosuppressant blood levels have thus found wide clinical use.

Initially, the immunosuppressant must be extracted and separated from the other components of the patient sample. The bulk of the immunosuppressant drug in the patient sample is present in a complex with various "carrier" molecules, such as binding proteins. Sirolimus, tacrolimus and cyclosporine are found predominately in the red blood cells of patient specimens and are associated with specific binding proteins, FKBP for sirolimus and tacrolimus, and cyclophilin for cyclosporine. To ensure an accurate measurement of the total drug concentration in the specimen, the

drug bound to the binding proteins is preferably liberated prior to quantitation. This has been addressed by using detergents to lyse cells and/or organic solvents to denature the sample proteins.

Following its extraction from the binding proteins, the drug can be measured in a number of different ways, including by immunoassay or chromatography with absorbance or mass spectrophotometric detection. Immunoassays for immunosuppressant drugs are available in a variety of formats, but all use the binding of an antibody or binding protein (*e.g.* FKBP) to the immunosuppressant drug. A commonly used immunoassay is an assay which involves the binding of a first antibody to the immunosuppressant and the binding of labeled immunosuppressant (*e.g.* acridinium-sirolimus) to the remaining free antibody binding sites, followed by quantitation by detection of the label.

SUMMARY

The invention provides a method for preparing a test sample for use in an assay, the method comprising contacting the test sample with a lysis reagent to form a lysis mixture. The lysis reagent includes a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof, and at least one alcohol having five or fewer carbons is included in the lysis reagent or added to the lysis mixture. The alcohol can, for example, be methanol, ethanol, and/or propanol. In particular embodiments, the test sample includes a human blood sample.

Advantages of the sample preparation method of the invention include that the sample can be prepared for analysis without a centrifugation step and/or without the use of a detergent.

In preferred embodiments, the alcohol is included in the lysis reagent. In variations of such embodiments, the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4 or, preferably, about 4:1 to about 1:2. The test sample can be added to any such lysis reagent at a ratio in the range of about 2:1 to about 1:2.

Where the method is carried out prior to an assay for an analyte that is bound to one or more binding proteins in the test sample, the method can additionally include contacting the test sample or the lysis mixture with an agent that releases the analyte

from the binding protein(s). The releasing agent can, for example, be an agent that competes with the analyte for binding to the binding protein(s). In an exemplary embodiment, the analyte includes an immunosuppressant drug, and the agent includes a different, but structurally similar, immunosuppressant drug. Where the analyte
5 includes a non-protein molecule, the agent can, for example, include a protease that degrades the binding protein(s).

Another aspect of the invention is a lysis reagent mixture, which includes a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof and at least one alcohol having five or fewer carbons. In exemplary
10 embodiments, the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4 or, preferably, about 4:1 to about 1:2. This glycol-alcohol mixture is typically added to a test sample, in exemplary embodiments, at a ratio in the range of about 2:1 to about 1:2. Accordingly, the term "lysis reagent mixture" also includes the mixture that results from the addition of the glycol-alcohol mixture to the test sample. The alcohol can, for
15 example, be methanol, ethanol, and/or propanol. In particular embodiments, the test sample includes a human blood sample.

The invention also provides a method for assessing the presence or concentration of an analyte in a test sample. This method entails contacting the test sample with a lysis reagent to form a lysis mixture and assaying the lysis mixture for
20 the analyte. The lysis reagent includes a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof, and at least one alcohol having five or fewer carbons is included in the lysis reagent or added to the lysis mixture. The alcohol can, for example, be methanol, ethanol, and/or propanol. In particular embodiments, the test sample includes a human blood sample.

25 In exemplary embodiments, the alcohol is included in the lysis reagent, and the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4 or, preferably, about 4:1 to about 1:2. This glycol-alcohol mixture is typically added to a test sample, in exemplary embodiments, at a ratio in the range of about 2:1 to about 1:2.

Advantages of the sample preparation method of the invention include that the
30 lysis mixture is a homogeneous mixture, suitable for automated pipetting, without the

need for a centrifugation step and/or without the use of a detergent. However, the test sample or the lysis mixture can be contacted with a detergent, if desired.

The lysis mixture can, for example, be analyzed by immunoassay. In exemplary embodiments, the analyte detected includes an immunosuppressant drug, such as, for example, sirolimus, tacrolimus, everolimus, temsorolimus, zotarolimus, cyclosporine, or analogs of any of these compounds.

In particular embodiments, the assay detects an analyte that is bound to one or more binding proteins in the test sample. In such embodiments, the method can additionally include contacting the test sample or the lysis mixture with an agent that releases the analyte from the binding protein(s). The releasing agent can, for example, be an agent that competes with the analyte for binding to the binding protein(s). In an exemplary embodiment, the analyte includes an immunosuppressant drug, and the agent includes a different, but structurally similar, immunosuppressant drug. Where the analyte includes a non-protein molecule, the agent can, for example, include a protease that degrades the binding protein(s).

Another aspect of the invention is a test kit including: (a) at least one antibody or protein capable of binding specifically to at least one analyte; (b) a lysis reagent including a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof; and (c) at least one alcohol having five or fewer carbons. The alcohol can, for example, be methanol, ethanol, and/or propanol. In preferred embodiments, the lysis reagent and alcohol(s) are combined and packaged in a single container. In variations of such embodiments, the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4, preferably about 4:1 to about 1:2. The test kit can, optionally, contain a control composition including the at least one analyte of (a) and/or a detergent.

In exemplary embodiments, the analyte detected includes an immunosuppressant drug, such as, for example, sirolimus, tacrolimus, everolimus, temsorolimus, zotarolimus, cyclosporine, or analogs of any of these compounds.

In particular embodiments, the test kit additionally includes an agent that releases the analyte from one or more binding proteins in the test sample. The releasing agent can, for example, be an agent that competes with the analyte for binding to the

binding protein(s). In an exemplary embodiment, the analyte includes an immunosuppressant drug, and the agent includes a different, but structurally similar, immunosuppressant drug. Where the analyte includes a non-protein molecule, the agent can, for example, include a protease that degrades the binding protein(s).

- 5 An exemplary, preferred test kit according to the invention includes: (a) at least one antibody or protein capable of binding specifically to at least one immunosuppressant drug selected from the group consisting of sirolimus, tacrolimus, everolimus, temsoralimus, zotarolimus and cyclosporine; (b) a lysis reagent including propylene glycol and ethanol at a ratio in the range of about 4:1 to about 1:2; and (c)
10 control composition including the at least one immunosuppressant drug of (a).

DETAILED DESCRIPTION

The invention relates to a non-denaturing lysis reagent that can be mixed with a test sample to produce a homogeneous lysis mixture. This approach is superior to previous extraction methods that rely on the use of detergents or denaturants.

- 15 The use of detergents can be problematic in particular formats because the quantity of detergent needed to quickly lyse and fragment cells may cause foaming, which is unacceptable for samples that must be pipetted by most automated pipetting systems, and may interfere with immunochemistry in samples to be analyzed immunoassay. The lysis reagent of the invention produces a sample that is not
20 susceptible to foaming and eliminates the need for detergents, thus avoiding detergent driven interferences in assay immunochemistry.

- The use of denaturants requires subsequent centrifugation steps to remove precipitated blood constituents, which reduces the efficiency of this approach. Additionally, the use of organic solvents at the concentration required can lead to
25 sample evaporation that is significant enough to affect analyte concentration. The lysis reagent of the invention produces a homogeneous mixture that is suitable for use in automated pipetting systems without the need for a centrifugation step and eliminates the use of substantial concentrations of volatile organic solvents.


Definitions

Terms used in the claims and specification are defined as set forth below unless otherwise specified.


An “immunosuppressant drug” or “immunosuppressant”, as used herein, refers to a therapeutic compound, either small molecule or antibody based, that has the same or similar chemical structure to either rapamycin (sirolimus) or cyclosporine, also known as cyclosporine A. Any known or hereafter developed analog of either rapamycin or cyclosporine is considered an immunosuppressant herein. Preferred immunosuppressants include sirolimus, tacrolimus, everolimus, temsolorimus, zotarolimus and cyclosporine. Tacrolimus and cyclosporine are calcineurin inhibitors that suppress early activation of the immune system’s T lymphocytes through inhibition of cytokines such as interleukin 2. In contrast, the primary target of sirolimus, everolimus and zotarolimus is mammalian target of rapamycin (mTOR), a specific cell-cycle regulatory protein. The inhibition of mTOR leads to suppression of cytokine-driven T-lymphocyte proliferation.

The chemical formula of cyclosporine is in Formula A. The chemical formula of sirolimus (rapamycin) is in Formula B. The chemical formula of the structural difference of everolimus (RAD) from sirolimus is in Formula C.


A



B



C



Numerous derivatives or analogs of cyclosporine have been prepared. The invention comprises lysis reagents, lysis methods, assays and assay kits for cyclosporine or any of its analogs.

Numerous derivatives or analogs of rapamycin have been prepared. For example, these include the preparation of ester mono- and di-ester derivatives of rapamycin (PCT International Application WO 92/05179), 27-oximes of rapamycin (European Patent EP 0 467606); 42-oxo analog of rapamycin (U.S. Patent No.


5,023,262); bicyclic rapamycins (U.S. Patent No. 5,120,725); rapamycin dimers (U.S. Patent No. 5,120,727); silyl ethers of rapamycin (U.S. Patent No. 5,120,842); and arylsulfonates and sulfamates (U.S. Patent No. 5,177,203). Rapamycin was recently synthesized in its naturally occurring enantiomeric form (K. C. Nicolaou et al., *J. Am. Chem. Soc.*, 1993, *115*, 4419-4420; S. L. Schreiber, *J. Am. Chem. Soc.*, 1993, *115*, 7906-7907; S. J. Danishefsky, *J. Am. Chem. Soc.*, 1993, *115*, 9345-9346. The invention comprises lysis reagents, lysis methods, assays and assay kits for rapamycin or any of its analogs.

Another immunosuppressant analog of rapamycin is FK-506, also known as tacrolimus, which was isolated from a strain of *S. tsukubaensis*. FK-506's chemical formula is published in European Patent EP 0 293 892 B1. Analogs of FK-506 include the related natural products FR-900520 and FR-900523, which differ from FK-506 in their alkyl substituent at C-21, and were isolated from *S. hygrosopicus yakushimnaensis*. Another analog, FR-900525, produced by *S. tsukubaensis*, differs from FK-506 in the replacement of a pipercolic acid moiety with a proline group. The invention comprises lysis reagents, lysis methods, assays and assay kits for FK-506 or any of its analogs. Temsorolimus is another ester derivative of sirolimus which can be monitored with the invention.

ABT-578 [40-epi-(1-tetrazolyl)-rapamycin], known better today as zotarolimus, is a semi-synthetic macrolide triene antibiotic derived from rapamycin. Zotarolimus' structure is shown in Formula D.

Formula D.

The isomers of zotarolimus



Pyran 1 (N-1 isomer)

Oxepane 2

(Ia)

(Ib)

As used herein with reference to an immunosuppressant drugs, the term “structurally similar” indicates that the drugs have sufficiently similar structures that the drugs bind competitively to at least one common binding partner (*e.g.*, a binding protein).

5 The term “test sample” refers to a component, tissue or fluid of an animal’s body that is the source of the immunosuppressant drug analyte. These components, tissues and fluids include human and animal body fluids such as whole blood, serum, plasma, synovial fluid, cerebrospinal fluid, urine, lymph fluids, and various external secretions of the respiratory, intestinal and genitourinary tracts, tears, saliva, milk,
10 white blood cells, myelomas and the like; biological fluids such as cell culture supernatants; fixed tissue specimens; and fixed cell specimens. Preferably, the test sample is a human peripheral blood sample.

As used herein, an “antibody” refers to a protein consisting of one or more polypeptides substantially encoded by immunoglobulin genes or fragments of
15 immunoglobulin genes. This term encompasses polyclonal antibodies, monoclonal antibodies, and fragments thereof, as well as molecules engineered from immunoglobulin gene sequences. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as myriad immunoglobulin variable region genes. Light chains are classified as either
20 kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

A typical immunoglobulin (antibody) structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each
25 pair having one “light” (about 25 kD) and one “heavy” chain (about 50 - 70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms “variable light chain (VL)” and “variable heavy chain (VH)” refer to these light and heavy chains respectively.

30 Antibodies exist as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for

example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce F(ab')₂, a dimer of Fab which itself is a light chain joined to VH-CH1 by a disulfide bond. The F(ab')₂ may be reduced under mild conditions to break the disulfide linkage in the hinge region thereby converting the (Fab')₂ dimer into a Fab' monomer. The Fab' monomer is essentially a Fab with part of the hinge region (see, 5 Fundamental Immunology, W.E. Paul, ed., Raven Press, N.Y. (1993), for a more detailed description of other antibody fragments). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such Fab' fragments may be synthesized de novo either chemically or by utilizing 10 recombinant DNA methodology.

Thus, the term "antibody," as used herein, also includes antibody fragments either produced by the modification of whole antibodies or synthesized de novo using recombinant DNA methodologies. The term "antibody" also encompasses single chain antibodies (antibodies that exist as a single polypeptide chain), more preferably single 15 chain Fv antibodies (sFv or scFv), in which a variable heavy and a variable light chain are joined together (directly or through a peptide linker) to form a continuous polypeptide. The single chain Fv antibody is a covalently linked VH-VL heterodimer which may be expressed from a nucleic acid including VH- and VL-encoding sequences either joined directly or joined by a peptide-encoding linker (Huston, et al. 20 (1988) Proc. Nat. Acad. Sci. USA, 85: 5879-5883). While the VH and VL are connected to each as a single polypeptide chain, the VH and VL domains associate non-covalently. The scFv antibodies and a number of other structures convert the naturally aggregated, but chemically separated, light and heavy polypeptide chains from an antibody V region into a molecule that folds into a three dimensional structure 25 substantially similar to the structure of an antigen-binding site are known to those of skill in the art (see *e.g.*, U.S. Patent Nos. 5,091,513, 5,132,405, and 4,956,778).

"Analyte," as used herein, refers to the substance to be detected, which may be suspected of being present in the test sample. The analyte can be any substance for which there exists a naturally occurring specific binding partner or for which a specific 30 binding partner can be prepared. Thus, an analyte is a substance that can bind to one or more specific binding partners in an assay.

A “binding partner,” as used herein, is a member of a binding pair, i.e., a pair of molecules wherein one of the molecules binds to the second molecule. Binding partners that bind specifically are termed “specific binding partners.” In addition to the antigen and antibody binding partners commonly used in immunoassays, other specific
5 binding partners can include biotin and avidin, carbohydrates and lectins, complementary nucleotide sequences, effector and receptor molecules, cofactors and enzymes, enzyme inhibitors and enzymes, and the like. Immunoreactive specific binding partners include antigens, antigen fragments, antibodies and antibody
10 fragments, both monoclonal and polyclonal, and complexes thereof, including those formed by recombinant DNA methods.

The term “specific binding” is defined herein as the preferential binding of binding partners to another (*e.g.*, a polypeptide and a ligand (analyte), two polypeptides, a polypeptide and nucleic acid molecule, or two nucleic acid molecules) at specific sites. The term “specifically binds” indicates that the binding preference
15 (*e.g.*, affinity) for the target molecule/sequence is at least 2-fold, more preferably at least 5-fold, and most preferably at least 10- or 20-fold over a non-specific target molecule (*e.g.* a randomly generated molecule lacking the specifically recognized site(s)).

An antibody that specifically binds an immunosuppressant drug is said to be
20 “specific for” that immunosuppressant drug.

The term “capture agent” is used herein to refer to a binding partner that binds to analyte, preferably specifically. Capture agents can be attached to a solid phase. As used herein, the binding of a solid phase-affixed capture agent to analyte forms a “solid phase-affixed complex.”

25 The term “labeled detection agent” is used herein to refer to a binding partner that binds to analyte, preferably specifically, and is labeled with a detectable label or becomes labeled with a detectable label during use in an assay.

A “detectable label” includes a moiety that is detectable or that can be rendered detectable.

As used with reference to a labeled detection agent, a “direct label” is a detectable label that is attached, by any means, to the detection agent.

As used with reference to a labeled detection agent, an “indirect label” is a detectable label that specifically binds the detection agent. Thus, an indirect label
5 includes a moiety that is the specific binding partner of a moiety of the detection agent. Biotin and avidin are examples of such moieties that are employed, for example, by contacting a biotinylated antibody with labeled avidin to produce an indirectly labeled antibody.

As used herein, the term “indicator reagent” refers to any agent that is contacted
10 with a label to produce a detectable signal. Thus, for example, in conventional enzyme labeling, an antibody labeled with an enzyme can be contacted with a substrate (the indicator reagent) to produce a detectable signal, such as a colored reaction product.

As used herein, a “glycol analog” is any glycol having from two to six carbon atoms.

15 A lysis mixture is said to be “homogeneous” when it is sufficiently free of large particulates to allow accurate and reliable pipetting (either manually or using an automated system).

I. Sample Collection and Processing

The methods of the invention are generally carried out on test samples derived
20 from an animal, preferably a mammal, and more preferably a human.

The methods of the invention can be carried out using any sample that may contain the analyte of interest (*e.g.*, an immunosuppressant drug), such as a blood sample.

The sample is collected by any standard technique and then contacted with a
25 lysis reagent to form a lysis mixture. The lysis reagent includes a glycol having from two to six carbon atoms. At least one alcohol having five or fewer carbons is included in the lysis reagent or added to the lysis mixture. In preferred embodiments, the lysis reagent includes the alcohol(s). Glycols suitable for use in the lysis reagent include, for example, ethylene glycol, propylene glycol, and analogs thereof thereof, as well as

mixtures of such glycols. Alcohols suitable for use in the invention include, for example, methanol, ethanol, propanol, and mixtures thereof. In particular
embodiments, the ratio of glycol to alcohol is in the range of about 5:1 to about 1:5, 4:1
to about 1:4, 2:1 to about 1:2, or about 1:1 (volume:volume). In more particular
5 embodiments, the ratio of glycol to alcohol is in the range of about 4:1 to about 1:2.

The lysis mixture can be formed by any mixing technique at any desirable
temperature to contact any chosen amount of the sample with the lysis reagent. The
sample is contacted with a sufficient volume of lysis reagent to lyse the cells in the
sample and produce a homogeneous mixture. For a lysis reagent wherein the ratio of
10 glycol to alcohol is in the range of about 4:1 to about 1:4, as described above, sample
can be added to the lysis reagent at a ratio in the range of about 2:1 to about 1:2, about
1:3, about 1:4, about 1:5, or about 1:10; *e.g.*, about 1:1 (volume:volume), or any other
range including these values as endpoints, depending on the lysis reagent composition.
For example, about 100 μL to about 600 μL of blood sample can be mixed with about
15 50 μL to about 1200 μL of the lysis reagent for up to about five minutes. In certain
embodiments, the lysis mixture is formed by mixing 150 μL of blood sample with 300
 μL of lysis reagent and vortexing vigorously for from about 5 to about 10 seconds. In
preferred embodiments, lysis is complete in less than a minute at room temperature.
The lysis mixture is then assayed for the analyte using a suitable assay. In preferred
20 embodiments, the lysis mixture is produced, ready for analysis, without the need to
centrifuge the sample.

The lysis reagent of the invention can be employed without any added
detergent. However, in certain embodiments, one or more detergents can be added, if
desired. Detergents typically do not foam in the presence of the lysis reagent, and thus
25 lysis mixtures prepared according to the invention are amenable to automated pipetting,
regardless of whether a detergent is included. If included in a lysis mixture intended
for immunoassay, the detergent is preferably present at a concentration that does not
interfere with the immunochemistry. Preferably, the detergent is a non-ionic detergent,
such as saponin, and is employed at a concentration in the range of about 0.01% to
30 0.1%, more preferably about 0.1%. U.S. Patent No. 5,650,288 (issued July 22, 1997 to
MacFarlane and Jensen; hereby incorporated by reference in its entirety for its

teachings regarding detergent use) describes the use of detergents and particularly in immunoassays for immunosuppressants.

In particular embodiments, where the analyte is bound to one or more binding proteins in the test sample, the method can additionally entail contacting the test sample with an agent that releases the analyte from the binding protein(s). This agent can be included in the lysis reagent, if desired. The agent can, for example, be one that competes with the analyte for binding to the binding protein(s). The agent is generally selected so that it will not affect the results of the assay to be carried out. Thus, for instance, if the assay is an immunoassay, the agent is typically one that the relevant antibody does not cross-react with. Where the analyte is an immunosuppressant drug, the agent can be a different, but structurally similar, immunosuppressant drug. For example, sirolimus and tacrolimus both bind FKBP, and, for this reason, sirolimus can be used to release tacrolimus from FKBP and vice versa. Any subsequent immunoassay will generally employ an antibody that distinguishes between sirolimus and tacrolimus. U.S. Patent No. 6,187,547 (issued Feb. 13, 2001 to Legay and Wenger; incorporated herein by reference in its entirety for its teachings regarding immunosuppressant drug competition) describes "binding competitors" useful for releasing immunosuppressant drugs from binding proteins. Examples include: [Thr², Leu⁵, D-Hiv⁸, Leu¹⁰]-Ciclosporin, which can release cyclosporine.

Where the analyte is a non-protein molecule, a protease can be employed to release the analyte from binding protein(s). The protease used in the method should be one that can degrade the binding protein, thereby releasing the analyte for assay, and can be inactivated without adversely affecting the sensitivity and the precision of the assay to be carried out. Care should be taken in obtaining enzymes free from other contaminating enzymes that might not be inactivated by the method of inactivation used. Otherwise, any residual proteolytic activity could degrade an antibody used in a subsequent immunoassay. Exemplary proteases include proteinase K, subtilisin, dispase, thermolysin, trypsin, ficin, bromelain, and combinations thereof.

Proteinase K (Sigma Chemical Co., St. Louis, Mo.) is a nonspecific, Ca-dependent protease that can be inactivated by heat (65°C or higher) and by specific protease inhibitors, such as phenyl methyl sulfonyl fluoride (PMSF, Boehringer

Mannheim, Indianapolis, Ind.) or diisopropylfluorophosphate (DFP, Calbiochem, La Jolla, Calif.). Subtilisin (Sigma) is also a nonspecific, Ca-dependent protease that can be inactivated by heat (55°C or higher), although it can be inhibited by acidic pH or a specific protease inhibitor, such as PMSF, DFP or aprotinin.

5 Dispase (Boehringer Mannheim or Sigma or Calbiochem) and thermolysin (Sigma or Boehringer Mannheim) are Ca-dependent metallo-proteases, which can be inactivated by EDTA, at a concentration of about 5 mM, for example. When dispase and thermolysin combined are used as the protease, proteolysis is preferably inactivated by addition of a divalent cation chelator, such as EDTA, at a concentration of about 5
10 mM, for example, in the presence of a zinc salt, *e.g.*, ZnSO₄, at a concentration of about 40 mM, for example.

 Trypsin (Worthington Biochemical Corp., Freehold, N.J.) cleaves proteins specifically at the carboxyl side of lysine or arginine residues and can be inhibited by heat (90°C or higher) or specifically inhibited by many agents, including aprotinin
15 (Aprotinin injection formerly marketed as Trasylol® by Bayer, West Haven, CT; inhibitor still available from Calbiochem, La Jolla, CA, and other vendors), leupeptin (Sigma-Aldrich, St. Louis, MO or Boehringer Mannheim), PMSF, or specific trypsin inhibitors derived from soybean, lima bean or egg white (Worthington Biochemical Corp., Freehold, N.J. or Sigma-Aldrich, St. Louis, MO). Ficin is a thiol protease that
20 can be inactivated by HgCl₂, at a concentration of about 2 mM, for example. Bromelain is also a thiol protease and can be inactivated by bromelain inhibitor (Sigma-Aldrich, St. Louis, MO).

 In particular embodiments, the concentration of protease is high enough to degrade to the binding proteins within about 30 min, preferably within about 20 min,
25 yet low enough to allow efficient inactivation of the enzyme. Accordingly, the concentration of protease is preferably be in the range of about 0.5 to 2.0 units/mL, more preferably about 1 unit/mL.

 After lysis and release from binding proteins, if applicable, the analyte can be measured using any standard technique for detecting that analyte, *e.g.*, immunoassay or
30 chromatography with absorbance or mass spectrophotometric detection. For detection of immunosuppressant drugs, immunoassays are conveniently employed.

II. Immunoassays

A. In General

Immunoassays according to the invention can be used for the qualitative identification and/or the quantification of analyte in a test sample. These methods are applicable, for example, to immunoassays of immunosuppressant drugs, such as rapamycin (sirolimus), tacrolimus, everolimus, temsorolimus, zotarolimus, cyclosporine, and analogs of any of these compounds. Such immunoassays can be carried out by combining a lysis reagent with the test sample to form a lysis mixture, as described above. The lysis mixture can be contacted with at least one antibody specific for the analyte under conditions suitable for binding of the antibody to the analyte, if present, to form an assay mixture, and binding of the antibody to the analyte is then detected.

In certain embodiments, enhanced assay sensitivity can be achieved by contacting the lysis mixture with the antibody in the presence of a salt concentration of greater than about 0.4 M (*e.g.*, from about 0.5 M to about 5.0 M). In particular embodiments, the salt concentration is less than or equal to about 4.0 M (*e.g.*, from about 0.5 M to about 4.0 M). In exemplary embodiments, the salt concentration is about 2.0 M (*e.g.*, from about 1.5 M to about 2.5 M, particularly about 1.8 M, about 1.9 M, about 2.0 M, about 2.1 M, or about 2.2M). Suitable salts can include, for example, any of the following anions: fluoride, chloride, bromide, iodide, thiocyanate, acetate, citrate, and bisulfate. In particular embodiments, the salt includes a monovalent anion, such as, for example: fluoride, chloride, bromide, iodide, thiocyanate, and acetate. In preferred embodiments, the salt includes chloride, *e.g.*, a chloride salt of an alkali metal (*e.g.*, lithium, sodium, potassium, rubidium, cesium). Generally, the salt employed is soluble under the assay conditions. Sodium chloride is highly soluble under most conditions, and can thus be conveniently used to enhance assay sensitivity in a wide variety of immunoassays according to the invention.

The salt can be provided to the assay mixture in any convenient manner and can be present before, or added after, contact between the lysis mixture and the antibody. In particular embodiments, the salt is provided in an assay diluent, which can also optionally include one or more other components, in addition to water (such as, for

example, a buffer). The salt concentration in the assay diluent will vary, depending on the desired final salt concentration and on the amount of diluent added to the assay mixture. For example, an assay diluent having a salt concentration of about 4.0 M could be added to an equal volume of assay mixture to provide a final salt
5 concentration of about 2.0 M.

B. Antibodies

In immunoassays for the qualitative or quantitative detection of an analyte in a test sample, at least one antibody that binds to the analyte is contacted with a lysis mixture suspected of containing the analyte to form an antibody-analyte immune
10 complex. To detect immunosuppressant drugs, any suitable antibodies that bind to the particular drug can be used in immunoassay according to the invention. Antibodies to each of rapamycin (sirolimus), tacrolimus, zotarolimus, cyclosporine and everolimus are known in the art and/or are commercially available, and any of these can be used. It is preferred to use the monoclonal antibody that is a component of Abbott Laboratories'
15 commercially available IMx® Sirolimus assay (Abbott Laboratories, Abbott Park, IL) for measuring sirolimus, or any other Sirolimus assay kit marketed by Abbott Laboratories (*e.g.*, for use on a different commercial automated platform).

An exemplary protocol for producing an antibody specific for an immunosuppressant drug is as follows. Female RBf/Dnj mice are administered 3
20 monthly boosts of a drug-27-CMO-tetanus toxoid immunogen followed by an immunization with drug-42-HS-tetanus toxoid preparation on the 4th month. Seven months later, an intrasplenic pre-fusion boost is administered to the animal using the drug-27-CMO-tetanus toxoid immunogen 3 days prior to the fusion. Splenic B-cells are then isolated and used in a standard polyethylene (PEG) fusion with the SP2/0
25 myeloma. Confluent cultures are screened for anti-drug activity 10-14 days later in a microtiter EIA and positive cultures are then cloned using limiting dilution cloning technique. The resulting clones are isolated and scaled up in IMDM w/FBS (Invitrogen Corp., Carlsbad, CA) tissue culture medium and the secreted antibody is affinity purified using Protein A. An exemplary, preferred antibody generated using sirolimus
30 as the drug can be used in immunoassays for sirolimus, everolimus and zotarolimus.

An exemplary, preferred antibody for use in immunoassays for tacrolimus is described in M. Kobayashi et al., "A Highly Sensitive Method to Assay FK-506 Levels in Plasma", at pp 23 –29 of "FK-506 A Potential Breakthrough in Immunosuppression", *A Transplantation Proceedings Reprint*, Supplement 6, Vol. XIX, October, 1987, Editors T. Starzl, L. Makowka and S. Todo, published by Grune & Stratton, Inc., Philadelphia, PA.

An exemplary, preferred antibody for use in immunoassays for cyclosporine is the monoclonal antibody that is a component of Abbott Laboratories' commercially available AxSYM® cyclosporine assay for measuring cyclosporine.

10 **C. Detection**

The antibody-analyte immune complexes can then detected using any suitable technique. For example, an antibody can be labeled with a detectable label to detect the presence of the antibody-analyte complex. The selection of a particular label is not critical, but the chosen label must be capable of producing a detectable signal either by itself or in conjunction with one or more additional substances.

Useful detectable labels, their attachment to antibodies and detection techniques therefore are known in the art. Any detectable label known in the art can be used. For example, the detectable label can be a radioactive label, such as, ^3H , ^{125}I , ^{35}S , ^{14}C , ^{32}P , ^{33}P ; an enzymatic label, such as horseradish peroxidase, alkaline peroxidase, glucose 6-phosphate dehydrogenase, etc.; a chemiluminescent label, such as, acridinium derivatives, luminol, isoluminol, thioesters, sulfonamides, phenanthridinium esters, etc.; a fluorescent label, such as, fluorescein (5-fluorescein, 6- carboxyfluorescein, 3'6-carboxyfluorescein, 5(6)-carboxyfluorescein, 6-hexachloro-fluorescein, 6-tetrachlorofluorescein, fluorescein isothiocyanate, etc.), rhodamine, phycobiliproteins, R-phycoerythrin, quantum dots (zinc sulfide-capped cadmium selenide), a thermometric label or an immuno-polymerase chain reaction label. An introduction to labels, labeling procedures and detection of labels is found in Polak and Van Noorden, *Introduction to Immunocytochemistry*, 2nd ed., Springer Verlag, N.Y.(1997) and in Haugland, *Handbook of Fluorescent Probes and Research Chemi* (1996), which is a combined handbook and catalogue published by Molecular Probes, Inc., Eugene, Oregon, each of which is incorporated herein by reference. Preferred labels for use

with the invention are chemiluminescent labels such as acridinium-9-carboxamide. Additional detail can be found in Mattingly, P. G., and Adamczyk, M. (2002) Chemiluminescent *N*-sulfonylacridinium-9-carboxamides and their application in clinical assays, in *Luminescence Biotechnology: Instruments and Applications* (Dyke, K. V., Ed.) pp 77–105, CRC Press, Boca Raton.

The detectable label can be bound to the analyte, analyte analog, or antibody either directly or through a coupling agent. An example of a coupling agent that can be used is EDAC (1-ethyl-3-(3-dimethylaminopropyl) carbodiimide, hydrochloride), which is commercially available from Sigma-Aldrich (St. Louis, MO). Other coupling agents that can be used are known in the art. Methods for binding a detectable label to an antibody are known in the art. Additionally, many detectable labels can be purchased or synthesized that already contain end groups that facilitate the coupling of the detectable label to the antibody, such as, N10-(3-sulfopropyl)-N-(3-carboxypropyl)-acridinium-9-carboxamide, otherwise known as CPSP-Acridinium Ester or N10-(3-sulfopropyl)-N-(3-sulfopropyl)-acridinium-9-carboxamide, otherwise known as SPSP-Acridinium Ester.

Alternatively, a second antibody that binds to analyte and that contains a detectable label can be added to the lysis mixture and used to detect the presence of the antibody-analyte complex. Any suitable detectable label can be used in this embodiment.

D. Exemplary Formats

The immunoassays of the invention can be conducted using any format known in the art, such as, but not limited to, a sandwich format, a competitive inhibition format (including both forward or reverse competitive inhibition assays) or a fluorescence polarization format. The exemplary formats described below are described in terms of assaying an immunosuppressant drug. However, as those of skill in the art appreciate, the described formats are applicable to any analyte.

In immunoassays for the quantitative detection of an immunosuppressant, such as a preferred sandwich type format, at least two antibodies are employed to separate and quantify the drug in the lysis mixture. More specifically, the at least two antibodies

bind to different parts of the drug, forming an immune complex which is referred to as a “sandwich”. Generally, one or more antibodies can be used to capture the immunosuppressant in the test sample (these antibodies are frequently referred to as a “capture” antibody or “capture” antibodies) and one or more antibodies is used to bind
5 a detectable (namely, quantifiable) label to the sandwich (these antibodies are frequently referred to as the “detection” antibody or “detection” antibodies). In a sandwich assay, it is preferred that both antibodies binding to the drug are not diminished by the binding of any other antibody in the assay to its respective binding site. In other words, antibodies should be selected so that the one or more first
10 antibodies brought into contact with a lysis mixture suspected of containing an immunosuppressant do not bind to all or part of the binding site recognized by the second or subsequent antibodies, thereby interfering with the ability of the one or more second or subsequent antibodies to bind to the drug. In a sandwich assay, the antibodies, and preferably, the at least one capture antibody, are used in molar excess
15 amounts relative to the maximum amount of drug expected in the lysis mixture. For example, from about 5 $\mu\text{g/mL}$ to about 1 mg/mL of antibody per mL of solid phase containing solution can be used.

In one embodiment, the at least one first capture antibody can be bound to a solid support which facilitates the separation of the first antibody-drug complex from
20 the test sample. The solid support or “solid phase” used in the inventive immunoassay is not critical and can be selected by one skilled in the art. A solid phase or solid support, as used herein, refers to any material that is insoluble, or can be made insoluble by a subsequent reaction. Useful solid phases or solid supports are known to those in the art and include the walls of wells of a reaction tray, test tubes, polystyrene
25 beads, magnetic beads, nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, and Duracytes® (a registered trademark of Abbott Laboratories, Abbott Park, Ill.), which are red blood cells “fixed” by pyruvic aldehyde and formaldehyde, and others. Suitable methods for immobilizing peptides on solid phases include ionic, hydrophobic, covalent interactions and the like.
30 The solid phase can be chosen for its intrinsic ability to attract and immobilize the capture agent. Alternatively, the solid phase can comprise an additional receptor which has the ability to attract and immobilize the capture agent. The additional receptor can

include a charged substance that is oppositely charged with respect to the capture agent itself or to a charged substance conjugated to the capture agent. As yet another alternative, the receptor can be any specific binding partner which is immobilized upon (attached to) the solid phase and which has the ability to immobilize the capture agent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture agent to a solid phase material before the performance of the assay or during the performance of the assay.

Any solid support known in the art can be used, including but not limited to, solid supports made out of polymeric materials in the forms of wells, tubes or beads. The antibody (or antibodies) can be bound to the solid support by adsorption, by covalent bonding using a chemical coupling agent or by other means known in the art, provided that such binding does not interfere with the ability of the antibody to bind the drug. Moreover, if necessary, the solid support can be derivatized to allow reactivity with various functional groups on the antibody. Such derivatization requires the use of certain coupling agents such as, but not limited to, maleic anhydride, N-hydroxysuccinimide and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide.

It is within the scope of the present invention that the solid phase also can comprise any suitable porous material with sufficient porosity to allow access by detection antibodies and a suitable surface affinity to bind antigens. Microporous structure generally are preferred, but materials with gel structure in the hydrated state may be used as well. Such useful solid supports include but are not limited to nitrocellulose and nylon. It is contemplated that such porous solid supports described herein preferably are in the form of sheets of thickness from about 0.01 to 0.5 mm, preferably about 0.1 mm. The pore size may vary within wide limits, and preferably is from about 0.025 to 15 microns, especially from about 0.15 to 15 microns. The surface of such supports may be activated by chemical processes which cause covalent linkage of the antigen or antibody to the support. The irreversible binding of the antigen or antibody is obtained, however, in general, by adsorption on the porous material by hydrophobic forces.

After the lysis mixture suspected of containing or containing the immunosuppressant is brought into contact with the at least one first capture antibody,

the resulting assay mixture is incubated to allow for the formation of a first capture antibody (or multiple antibody)-drug complex. The incubation can be carried out at any suitable pH, including a pH of from about 4.5 to about 10.0, at any suitable temperature, including from about 2°C to about 45°C, and for a suitable time period
5 from at least about one (1) minute to about eighteen (18) hours, preferably from about 4-20 minutes, most preferably from about 17-19 minutes.

After the addition of a detection agent and the formation of a labeled complex, the amount of label in the complex is quantified using techniques known in the art. For example, if an enzymatic label is used, the labeled complex is reacted with a substrate
10 for the label that gives a quantifiable reaction such as the development of color. If the label is a radioactive label, the label is quantified using a scintillation counter. If the label is a fluorescent label, the label is quantified by stimulating the label with a light of one color (which is known as the “excitation wavelength”) and detecting another color (which is known as the “emission wavelength”) that is emitted by the label in response
15 to the stimulation. If the label is a chemiluminescent label, the label is quantified detecting the light emitted either visually or by using luminometers, x-ray film, high speed photographic film, a CCD camera, etc. Once the amount of the label in the complex has been quantified, the concentration of drug in the test sample can be determined by use of a standard curve that has been generated, for example, using serial
20 dilutions of immunosuppressant drug of known concentration. Other than using serial dilutions of the drug, the standard curve can be generated gravimetrically, by mass spectroscopy and by other techniques known in the art.

In a preferred forward competitive format, an aliquot of labeled drug, or analogue thereof, of a known concentration is used to compete with the drug present in
25 a test sample for binding to the antibody. In a forward competition assay, an immobilized antibody can either be sequentially or simultaneously contacted with the test sample and a labeled drug or drug analogue thereof. The drug or drug analogue can be labeled with any suitable detectable label, including those detectable labels discussed above. In this assay, the capture antibody can be immobilized on to a solid
30 support using the techniques discussed previously herein. Alternatively, the capture antibody can be coupled to an antibody, such as an antispecies antibody, that has been immobilized on to a solid support, such as a microparticle.

The labeled drug or drug analogue, the lysis mixture and the antibody are incubated under conditions similar to those described above in connection with the sandwich assay format. Two different types of antibody-drug complexes are then generated. Specifically, one of the antibody-drug complexes generated contains a detectable label while the other antibody-drug complex does not contain a detectable label. The antibody-drug complex can be, but does not have to be, separated from the remainder of the assay mixture prior to quantification of the detectable label. Regardless of whether the antibody-drug complex is separated from the remainder of the assay mixture, the amount of detectable label in the antibody-drug complex is then quantified. The concentration of drug in the test sample can then be determined by comparing the quantity of detectable label in the antibody-drug complex to a standard curve. The standard curve can be generated using serial dilutions of the drug of known concentration, by mass spectroscopy, gravimetrically and by other techniques known in the art.

The antibody-drug complex can be separated from the assay mixture by binding the antibody to a solid support, such as the solid supports discussed above in connection with the sandwich assay format, and then removing the remainder of the assay mixture from contact with the solid support.

In a reverse competition assay, an immobilized immunosuppressant drug or analogue thereof can either be sequentially or simultaneously contacted with a lysis mixture and at least one labeled antibody. The antibody can be labeled with any suitable detectable label, including those detectable labels discussed above. The drug or drug analogue can be bound to a solid support, such as the solid supports discussed above in connection with the sandwich assay format.

The immobilized drug or drug analogue, lysis mixture, and at least one labeled antibody are incubated under conditions similar to those described above in connection with the sandwich assay format. Two different types of antibody-drug complexes are then generated. Specifically, one of the antibody-drug complexes generated is immobilized and contains a detectable label while the other antibody-drug complex is not immobilized and contains a detectable label. The non-immobilized antibody-drug complex and the remainder of the assay mixture are removed from the presence of the

immobilized antibody-drug complex through techniques known in the art, such as washing. Once the non-immobilized antibody-drug complex is removed, the amount of detectable label in the immobilized antibody-drug complex is then quantified. The concentration of drug in the test sample can then be determined by comparing the quantity of detectable label in the antibody-drug complex to a standard curve. The standard curve can be generated using serial dilutions of the drug of known concentration, by mass spectroscopy, gravimetrically and by other techniques known in the art.

In a fluorescence polarization assay, in one embodiment, an antibody or functionally active fragment thereof is first contacted with an unlabeled lysis mixture containing the immunosuppressant drug to form an unlabeled antibody-drug complex. The unlabeled antibody-drug complex is then contacted with a fluorescently labeled drug or analogue thereof. The labeled drug or drug analogue competes with any unlabeled drug in the assay mixture for binding to the antibody or functionally active fragment thereof. The amount of labeled antibody-drug complex formed is determined and the amount of drug in the test sample determined via use of a standard curve.

The use of scanning probe microscopy (SPM) for immunoassays also is a technology to which the immunoassay methods of the present invention are easily adaptable. In SPM, in particular in atomic force microscopy, a capture agent is affixed to a solid phase having a surface suitable for scanning. The capture agent can, for example, be adsorbed to a plastic or metal surface. Alternatively, the capture agent can be covalently attached to, *e.g.*, derivatized plastic, metal, silicon, or glass according to methods known to those of ordinary skill in the art. Following attachment of the capture agent, the lysis mixture is contacted with the solid phase, and a scanning probe microscope is used to detect and quantify solid phase-affixed complexes. The use of SPM eliminates the need for labels which are typically employed in immunoassay systems. Such a system is described in U.S. App. No. 662,147, which is incorporated herein by reference.

Immunoassays according to the invention can also be carried out using a MicroElectroMechanical System (MEMS). MEMS are microscopic structures integrated onto silicon that combine mechanical, optical, and fluidic elements with

electronics, allowing convenient detection of an analyte of interest. An exemplary MEMS device suitable for use in the invention is the Protiveris' multicantilever array. This array is based on chemo-mechanical actuation of specially designed silicon microcantilevers and subsequent optical detection of the microcantilever deflections.

5 When coated on one side with a binding partner, a microcantilever will bend when it is exposed to a solution containing the complementary molecule. This bending is caused by the change in the surface energy due to the binding event. Optical detection of the degree of bending (deflection) allows measurement of the amount of complementary molecule bound to the microcantilever.

10 In other embodiments, immunoassays according to the invention are carried out using electrochemical detection. A basic procedure for electrochemical detection has been described by Heineman and coworkers. This entailed immobilization of a primary antibody (Ab, rat-anti mouse IgG), followed by exposure to a sequence of solutions containing the antigen (Ag, mouse IgG), the secondary antibody conjugated to an
15 enzyme label (AP-Ab, rat anti mouse IgG and alkaline phosphatase), and p-aminophenyl phosphate (PAPP). The AP converts PAPP to p-aminophenol (PAP_R , the "R" is intended to distinguish the reduced form from the oxidized form, PAP_O , the quinoneimine), which is electrochemically reversible at potentials that do not interfere with reduction of oxygen and water at pH 9.0, where AP exhibits optimum activity.
20 PAP_R does not cause electrode fouling, unlike phenol whose precursor, phenylphosphate, is often used as the enzyme substrate. Although PAP_R undergoes air and light oxidation, these are easily prevented on small scales and short time frames. Picomole detection limits for PAP_R and femtogram detection limits for IgG achieved in microelectrochemical immunoassays using PAPP volumes ranging from 20 μ L to 360
25 μ L have been reported previously. In capillary immunoassays with electrochemical detection, the lowest detection limit reported thus far is 3000 molecules of mouse IgG using a volume of 70 μ L and a 30 minute or 25 minute assay time.

Various electrochemical detection systems are described in U.S. Patent Nos. 7,045,364 (issued May 16, 2006; incorporated herein by reference), 7,045,310
30 (issued May 16, 2006; incorporated herein by reference), 6,887,714 (issued May 3, 2005; incorporated herein by reference), 6,682,648 (issued January 27, 2004;

incorporated herein by reference); 6,670,115 (issued December 30, 2003; incorporated herein by reference).

In particular embodiments, useful, for example, for simultaneously assaying multiple analytes in one test sample, the solid phase can include a plurality of different capture agents. Thus, for example, the solid phase can have affixed thereon a plurality of antibodies, wherein each is intended to test for the presence of different analytes in the sample. In an exemplary embodiment, the solid phase can consist of a plurality of different regions on a surface, wherein each region has a particular antibody affixed therein.

Multiplex formats can, but need not, employ a plurality of labels, wherein each label is used for the detection of a particular analyte. For example, multiple, different analytes can be detected without using a plurality of labels where a plurality of capture agents, such as antibodies, are affixed to the solid phase at different known locations, based on specificity. Because the specificity of the capture agent at each location is known, the detection of a signal at a particular location can be associated with the presence of analyte bound at that location. Examples of this format include microfluidic devices and capillary arrays, containing different capture agents at different locations along a channel or capillary, respectively, and microarrays, which typically contain different capture agents arranged in a matrix of spots (“target elements”) on a surface of a solid support. In particular embodiments, each different capture agent can be affixed to a different electrode, which can, for example, be formed on a surface of a solid support, in a channel of a microfluidic device, or in a capillary.

III. Test Kits

The invention also provides test kits for assaying test samples for an analyte. Test kits according to the invention include one or more reagents useful for practicing one or more immunoassays according to the invention. A test kit generally includes a package with one or more containers holding the reagents, as one or more separate compositions or, optionally, as admixture where the compatibility of the reagents will allow. The test kit can also include other material(s), which may be desirable from a user standpoint, such as a buffer(s), a diluent(s), a standard(s), and/or any other material useful in sample processing, washing, or conducting any other step of the assay.

In particular embodiments, test kits of the invention can include: (a) at least one antibody or protein capable of binding specifically to at least one analyte; and (b) a lysis reagent comprising: a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof; and at least one alcohol having five or fewer carbons. In exemplary embodiments, useful for carrying out immunoassays for immunosuppressant drugs, the antibody can be specific for rapamycin (sirolimus), tacrolimus, everolimus, temsorolimus, zotarolimus, cyclosporine, or analogs of any of these compounds.

In certain embodiments, the lysis reagent includes methanol, ethanol, propanol, or a mixture of any of these alcohols. In exemplary embodiments, the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4, more particularly, in the range of about 4:1 to about 1:2. In one embodiment, the lysis reagent uses a glycol level that is between about 10% and about 40% of the lysis reagent. In another embodiment, alcohol is used alone in the lysis reagent, without glycol added.

If desired, the test kit can additionally include a control composition that includes the analyte being assayed.

In particular embodiments, test kits according to the invention can include one or more detergents and/or agents that release the analyte from one or more binding proteins in the test sample. Suitable detergents or detergent combinations include non-ionic detergents, such as saponin, as described above. Suitable releasing agents include agents that compete with the analyte for binding to one or more binding proteins, as described above, and proteases, which can be used to degrade binding proteins and liberate non-protein analytes. Exemplary proteases include proteinase K, subtilisin, dispase, thermolysin, trypsin, ficin, bromelain, and combinations thereof. Any detergents or proteases provided in kits of the invention should be provided in a manner that facilitates the production of a lysis mixture containing the components in suitable concentration, as described above.

Kits according to the invention can include a solid phase and a capture agent that is affixed to the solid phase or that becomes solid phase-affixed during the assay. In exemplary embodiments, the solid phase includes one or more microparticles or electrodes. Where such kits are to be employed for conducting sandwich

immunoassays, the kits can additionally include a labeled detection agent. In certain embodiments, the test kit includes at least one direct label, such as acridinium-9-carboxamide. Test kits according to the invention can also include at least one indirect label. If the label employed generally requires an indicator reagent to produce a detectable signal, the test kit preferably includes one or more suitable indicator reagents.

Test kits according to the invention preferably include instructions for carrying out one or more of the immunoassays of the invention. Instructions included in kits of the invention can be affixed to packaging material or can be included as a package insert. While the instructions are typically written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to, electronic storage media (*e.g.*, magnetic discs, tapes, cartridges, chips), optical media (*e.g.*, CD ROM), and the like. As used herein, the term “instructions” can include the address of an internet site that provides the instructions.

Of course, it goes without saying that any of the exemplary formats herein, and any assay or kit according to the invention can be adapted or optimized for use in automated and semi-automated systems (including those in which there is a solid phase comprising a microparticle), as described, *e.g.*, in U.S. Patent Nos. 5,089,424 and 5,006,309, and as, *e.g.*, commercially marketed by Abbott Laboratories (Abbott Park, IL) including but not limited to Abbott Laboratories’ ARCHITECT®, AxSYM®, IMX®, ABBOTT PRISM®, and Quantum II platforms, as well as other platforms.

Additionally, the assays and kits of the present invention optionally can be adapted or optimized for point of care assay systems, including Abbott Laboratories’ Point of Care (i-STAT®) electrochemical immunoassay system. Immunosensors and methods of manufacturing and operating them in single-use test devices are described, for example in U.S. Patent 5,063,081 and published U.S. Patent Applications 20030170881, 20040018577, 20050054078, and 20060160164 (incorporated by reference herein for their teachings regarding same).

EXAMPLES

The following examples are offered to illustrate, but not to limit, the claimed invention.

Example

5 **Lysis and Immunoassay of a Blood Sample for Sirolimus**

This example illustrates the use of a non-denaturing lysis reagent to lyse a blood sample and carry out an automated immunoassay for sirolimus.

A lysis reagent is prepared by mixing propylene glycol and ethanol at a volume:volume ratio of 4:1, then diluted with water until the propylene glycol is
10 between 10% and 40% (vol). Lysis of a blood sample is accomplished by mixing 100
μL of sample with 200 μL of lysis reagent by vortexing vigorously for 5 – 10 seconds.
Lysis is complete in less than 1 min at room temperature. The resulting lysis mixture is
assayed on an automated ARCHITECT® i2000® analyzer (Abbott Laboratories,
Abbott Park, Illinois) by:

- 15 (1) Mixing 10 – 40 μL of the lysis mixture with 50 μL of microparticles coated with goat anti-mouse antibody (from Sigma-Aldrich, St. Louis, Missouri) and mouse anti-sirolimus antibody (prepared as described below).
- (2) Incubating the assay mixture for approximately 18 minutes at 33 – 38 degrees C. The sirolimus in the sample binds the anti-sirolimus antibody on the
20 microparticles.
- (3) Adding 20 μL of acridinium-sirolimus conjugate to the reaction mixture.
- (4) Incubating the reaction mixture for approximately 4 minutes at 33 – 38 degrees C. The acridinium-sirolimus conjugate binds free anti-sirolimus binding sites.
- (5) Washing the microparticles with a phosphate buffer.
- 25 (6) Adding Pre-trigger (acid solution) and Trigger (basic solution) to cause the captured acridinium-sirolimus label to emit light, which is measured by the instrument.

The sirolimus-binding antibody was produced as follows: Female RBf/Dnj mice were administered 3 monthly boosts of a sirolimus-27-CMO-tetanus toxoid
30 immunogen, followed by an immunization with sirolimus-42-HS-tetanus toxoid

preparation on the 4th month. Seven months later, an intrasplenic pre-fusion boost was administered to the animals using the sirolimus-27-CMO-tetanus toxoid immunogen 3 days prior to the fusion. Splenic B-cells were isolated and used in a standard PEG fusion with the SP2/0 myeloma. Confluent cultures were screened for anti-sirolimus activity 10-14 days later in a microtiter EIA, and positive cultures were cloned using limiting dilution cloning technique. Isolated clones were scaled up in IMDM w/FBS (Invitrogen Corp., Carlsbad, CA) tissue culture medium, and secreted antibody was affinity purified using Protein A.

10 It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

In addition, the commonly owned, co-pending application U.S. Provisional Application Serial Number 60/882,732 filed on December 29, 2006 is explicitly incorporated by reference in its entirety for its teachings regarding a diagnostic test for the detection of a molecule or drug in whole blood.

The commonly owned, co-pending application U.S. Provisional Application Serial Number 60/878,017 filed on December 29, 2006 is explicitly incorporated by reference in its entirety for its teachings regarding a non-denaturing lysis reagent for use with capture-in-solution immunoassay.

The commonly owned, co-pending application U.S. Provisional Application Serial Number 60/882,863 filed on December 29, 2006 is explicitly incorporated by reference in its entirety for its teachings regarding an improved assay for immunosuppressant drugs.

The commonly owned, co-pending application U.S. Nonprovisional Application Serial Number 11/490,624 filed on July 21, 2006 is explicitly incorporated by reference in its entirety for its teachings regarding an extractive reagent composition.

In addition, all other publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

What is claimed is:

1. A method for preparing a test sample for use in an assay, the method comprising contacting the test sample with a lysis reagent to form a lysis mixture, the lysis reagent comprising a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof;

wherein at least one alcohol having five or fewer carbons is included in the lysis reagent or added to the lysis mixture.

2. The method of claim 1, wherein the alcohol is included in the lysis reagent.

3. The method of claim 1, wherein the test sample comprises a human blood sample.

4. The method of claim 1, wherein the alcohol is selected from the group consisting of methanol, ethanol, and propanol.

5. The method of claim 2, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4.

6. The method of claim 2, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:2.

7. The method of claim 2, wherein the test sample is added to the lysis reagent at a ratio in the range of about 2:1 to about 1:2.

8. The method of any of claims 1 to 7, wherein the method does not comprise centrifuging the lysis mixture.

9. The method of any of claims 1 to 7, wherein the method does not comprise contacting the test sample or the lysis mixture with a detergent.

10. The method of any of claims 1 to 7, wherein the method comprises contacting the test sample or the lysis mixture with a detergent.

11. The method of any of claims 1 to 7, wherein the assay detects an analyte that is bound to one or more binding proteins in the test sample, the method additionally comprising contacting the test sample or the lysis mixture with an agent that releases the analyte from said one or more binding proteins.

12. The method of claim 11, wherein the agent competes with the analyte for binding to said one or more binding proteins.

13. The method of claim 12, wherein the analyte comprises an immunosuppressant drug, and the agent comprises a different, but structurally similar, immunosuppressant drug.

14. The method of claim 11, wherein the analyte comprises a non-protein molecule, and the agent comprises a protease that degrades said one or more binding proteins.

15. A lysis reagent mixture comprising:
a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof; and
at least one alcohol having five or fewer carbons.

16. The lysis reagent mixture of claim 15, wherein the mixture additionally comprises a test sample.

17. The lysis reagent mixture of claim 16, wherein the test sample comprises a human blood sample.

18. The lysis reagent mixture of claim 15, wherein the alcohol is selected from the group consisting of methanol, ethanol, and propanol.

19. The lysis reagent mixture of any of claims 15 to 18, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4.

20. The lysis reagent mixture of any of claims 15 to 18, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:2.

21. The lysis reagent mixture of claim 16 or 17, wherein the ratio of the test sample to the lysis reagent is in the range of about 2:1 to about 1:2.
22. A method for assessing the presence or concentration of an analyte in a test sample, the method comprising:
 - (a) contacting the test sample with a lysis reagent to form a lysis mixture, the lysis reagent comprising a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof, wherein at least one alcohol having five or fewer carbons is included in the lysis reagent or added to the lysis mixture; and
 - (b) assaying the lysis mixture for the analyte.
23. The method of claim 22, wherein the alcohol is included in the lysis reagent.
24. The method of claim 22, wherein the lysis mixture is a homogeneous mixture.
25. The method of claim 22, wherein the assay comprises an immunoassay.
26. The method of claim 22, wherein the analyte comprises an immunosuppressant drug.
27. The method of claim 26, wherein the immunosuppressant drug is selected from the group consisting of sirolimus, tacrolimus, everolimus, temsorolimus, zotarolimus, cyclosporine, and analogs of any of these compounds.
28. The method of claim 22, wherein the test sample comprises a human blood sample.
29. The method of claim 22, wherein the alcohol is selected from the group consisting of methanol, ethanol, and propanol.
30. The method of claim 23, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4.

31. The method of claim 30, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:2.
32. The method of claim 23, wherein the test sample is added to the lysis reagent at a ratio in the range of about 2:1 to about 1:2.
33. The method of any of claims 22 to 32, wherein the method does not comprise centrifuging the lysis mixture.
34. The method of any of claims 22 to 32, wherein the method does not comprise contacting the test sample or the lysis mixture with a detergent.
35. The method of any of claims 22 to 32, wherein the method comprises contacting the test sample or the lysis mixture with a detergent.
36. The method of any of claims 22 to 32, wherein the assay detects an analyte that is bound to one or more binding proteins in the test sample, the method additionally comprising contacting the test sample or the lysis mixture with an agent that releases the analyte from said one or more binding proteins.
37. The method of claim 36, wherein the agent competes with the analyte for binding to said one or more binding proteins.
38. The method of claim 37, wherein the analyte is an immunosuppressant drug and the agent comprises a different, but structurally similar, immunosuppressant drug.
39. The method of claim 36, wherein the analyte comprises a non-protein molecule, and the agent comprises a protease that degrades said one or more binding proteins.
40. A test kit comprising:
- (a) at least one antibody or protein capable of binding specifically to at least one analyte;
 - (b) a lysis reagent comprising a glycol selected from the group consisting of ethylene glycol, propylene glycol, and an analog thereof; and

(c) at least one alcohol having five or fewer carbons.

41. The test kit of claim 40, wherein the lysis reagent and alcohol(s) are combined and packaged in a single container.

42. The test kit of claim 40, wherein the analyte comprises an immunosuppressant drug.

43. The test kit of claim 42, wherein the immunosuppressant drug is selected from the group consisting of sirolimus, tacrolimus, everolimus, temsorolimus, zotarolimus, cyclosporine, and analogs of any of these compounds.

44. The test kit of claim 40, additionally comprising a control composition comprising the at least one analyte of (a).

45. The test kit of claim 40, wherein the alcohol is selected from the group consisting of methanol, ethanol, and propanol.

46. The test kit of claim 41, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:4.

47. The test kit of claim 40, wherein the ratio of glycol to alcohol is in the range of about 4:1 to about 1:2.

48. The test kit of claim 40, additionally comprising a detergent.

49. The test kit of claim 40, wherein the test kit additionally comprises an agent that releases the analyte from one or more binding proteins in the test sample.

50. The test kit claim 49, wherein the agent competes with the analyte for binding to said one or more binding proteins.

51. The test kit of any of claims 40 to 50, wherein the agent analyte is an immunosuppressant drug and the agent comprises a different, but structurally similar, immunosuppressant drug.

52. The test kit claim 49, wherein the agent comprises a protease that degrades said one or more binding proteins.

53. A test kit comprising:

(a) at least one antibody or protein capable of binding specifically to at least one immunosuppressant drug selected from the group consisting of sirolimus, tacrolimus, everolimus, temsorolimus, zotarolimus and cyclosporine;

(b) a lysis reagent comprising propylene glycol and ethanol at a ratio in the range of about 4:1 to about 1:2; and

(c) a control composition comprising the at least one immunosuppressant drug of (a).

54. The test kit of claim 53, wherein the at least one immunosuppressant drug comprises sirolimus.

专利名称(译)	非变性裂解试剂		
公开(公告)号	EP2118655A4	公开(公告)日	2010-05-05
申请号	EP2007869487	申请日	2007-12-19
[标]申请(专利权)人(译)	雅培公司 GRENIER FRÇ WORKMAN RYAN°F 赛义德·HINAñ ALI SALMAN		
申请(专利权)人(译)	亚培 GRENIER , FRANK C. WORKMAN , RYAN F. SYED , HINA N. ALI , 萨勒曼		
当前申请(专利权)人(译)	亚培 GRENIER , FRANK C. WORKMAN , RYAN F. SYED , HINA N. ALI , 萨勒曼		
[标]发明人	GRENIER FRANK C WORKMAN RYAN F SYED HINA N ALI SALMAN		
发明人	GRENIER, FRANK C WORKMAN, RYAN F SYED, HINA N ALI, SALMAN		
IPC分类号	G01N33/53 G01N33/50 G01N33/94		
CPC分类号	G01N33/5306 G01N33/9493		
优先权	11/618495 2006-12-29 US		
其他公开文献	EP2118655B1 EP2118655A2		
外部链接	Espacenet		

摘要(译)

本发明提供了裂解试剂和制备用于测定的测试样品的方法，其中该方法产生适用于自动移液系统的均匀裂解混合物，而不需要离心步骤。裂解试剂包括二醇和醇。本发明的其他方面包括相关的免疫测定和测试试剂盒。

