



(11) **EP 1 629 281 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
29.10.2008 Bulletin 2008/44

(21) Application number: **04753487.0**

(22) Date of filing: **27.05.2004**

(51) Int Cl.:
G01N 33/53 (2006.01)

(86) International application number:
PCT/US2004/016667

(87) International publication number:
WO 2004/110250 (23.12.2004 Gazette 2004/52)

(54) **DIAGNOSTIC METHOD FOR ATHEROSCLEROSIS**
DIAGNOSEVERFAHREN FÜR ATHEROSKLEROSE
PROCEDE DE DIAGNOSTIC DE L'ATHEROSCLEROSE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PL PT RO SE SI SK TR

(30) Priority: **30.05.2003 US 474731 P**

(43) Date of publication of application:
01.03.2006 Bulletin 2006/09

(73) Proprietor: **Purdue Research Foundation West Lafayette, IN 47906 (US)**

(72) Inventor: **LOW, Philip, Stewart West Lafayette, IN 47906 (US)**

(74) Representative: **Elsy, David et al Withers & Rogers LLP Goldings House 2 Hays Lane London SE1 2HW (GB)**

(56) References cited:
WO-A-02/087424 US-A1- 2002 127 181

- **TURK M J ET AL: "Folate-targeted imaging of activated macrophages in rats with adjuvant-induced arthritis" ARTHRITIS AND RHEUMATISM, LIPPINCOTT, PHILADELPHIA, US, vol. 46, no. 7, July 2002 (2002-07), pages 1947-1955, XP002284074 ISSN: 0004-3591**
- **LIN-WU YANI ET AL: "Identification and analysis of macrophage-derived foam cells from human atherosclerotic lesions by using a "Mock" FL3 channel in flow cytometry" 1997, CYTOMETRY, VOL. 29, NR. 2, PAGE(S) 155-161 , XP002385778 ISSN: 0196-4763 * table 1 ***
- **ANTOHE FELICIA ET AL: "Increased uptake of folate conjugates by activated macrophages in experimental hyperlipemia" May 2005 (2005-05), CELL & TISSUE RESEARCH, VOL. 320, NR. 2, PAGE(S) 277-285 , XP002385779 ISSN: 0302-766X**
- **KENNEDY MICHAEL D ET AL: "Evaluation of folate conjugate uptake and transport by the choroid plexus of mice." May 2003 (2003-05), PHARMACEUTICAL RESEARCH (DORDRECHT), VOL. 20, NR. 5, PAGE(S) 714-719 , XP002385780 ISSN: 0724-8741 * figure 1 ***

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 629 281 B1

Description

FIELD OF THE INVENTION

5 **[0001]** This invention relates to a composition for use in a method for identifying/monitoring active atherosclerotic plaques. More particularly, ligands that bind to activated macrophages are conjugated to a chromophore or to a chemical moiety capable of emitting radiation for administration to a diseased host for identifying/monitoring active atherosclerotic plaques using a catheterization-based device.

10 BACKGROUND AND SUMMARY OF THE INVENTION

[0002] Activated macrophages can participate in the immune response by nonspecifically engulfing and killing foreign pathogens within the macrophage, by displaying degraded peptides from foreign proteins on the macrophage cell surface where they can be recognized by other immune cells, and by secreting cytokines and other factors that modulate the function of T and B lymphocytes, resulting in further stimulation of immune responses. Activated macrophages can also contribute to the pathophysiology of disease in some instances. For example, activated macrophages can contribute to atherosclerosis, rheumatoid arthritis, autoimmune disease states, and graft versus host disease.

15 **[0003]** Atherosclerosis is initiated when a fatty streak forms within a blood vessel wall. Formation of fatty streaks is believed to result from accumulation of lipoprotein particles in the intima layer of the blood vessel wall, the layer of the vessel wall underlying the luminal endothelial cell layer. Lipoprotein particles can associate with extracellular matrix components in the intima layer and can become inaccessible to plasma antioxidants, resulting in oxidative modification of the lipoprotein particles. Such oxidative modification may trigger a local inflammatory response resulting in adhesion of activated macrophages and T lymphocytes to the luminal endothelium followed by migration into the intima layer. The oxidized lipoprotein particles themselves can act as chemoattractants for cells of the immune system, such as macrophages and T cells, or can induce cells in the vascular wall to produce chemoattractants. The atherosclerotic lesion then forms a fibrous cap with a lipid-rich core filled with activated macrophages. Atherosclerotic lesions that are unstable are characterized by local inflammation, and lesions that have ruptured and have caused fatal myocardial infarction are characterized by an infiltration of activated macrophages and T lymphocytes.

25 **[0004]** The present invention relates to the use of a composition comprising a conjugate of the general formula

30

L-X

wherein the group L comprises a folate receptor binding ligand and the group X comprises a chromophore capable of emitting light or a chemical moiety capable of emitting radiation in the manufacture of a composition for identifying/monitoring active atherosclerotic plaques associated with blood vessel walls wherein the plaques comprise activated macrophages having accessible binding sites for the vitamin.

35 **[0005]** In accordance with the invention a ligand, that binds to a receptor which is preferentially expressed/presented on the surface of activated macrophages relative to resting macrophages, is conjugated to a chromophore or a chemical moiety capable of emitting radiation and the ligand conjugates are administered to a patient being evaluated for atherosclerosis. The ligand conjugates bind to activated macrophages associated with active atherosclerotic plaques and emit light (i.e., ligand-chromophore conjugates) or radiation (i.e., ligand-chemical moiety conjugates) and are detected using a catheter-based device. Accordingly, the ligand conjugates can be used to distinguish active atherosclerotic plaques containing activated macrophages from inactive plaques.

40 **[0006]** Methods are not presently available for distinguishing active and inactive atherosclerotic plaques. Because many unstable (i.e., active) atherosclerotic plaques, capable of rupturing and causing acute atherosclerotic syndromes do not produce luminal narrowing of blood vessels, particularly in the coronary circulation, the method of the present invention represents a significant advance in diagnosing the risk of myocardial infarction, and in evaluating the need for clinical intervention, in patients suffering from atherosclerosis.

45 **[0007]** A method is provided of identifying/monitoring active atherosclerotic plaques associated with blood vessel walls wherein the plaques comprise activated macrophages having accessible binding sites for a ligand. The method comprises the steps of administering to a patient being evaluated for atherosclerosis an effective amount of a composition comprising a conjugate of the general formula

50

L-X

55

wherein the group L comprises the ligand and the group X comprises a chromophore capable of emitting light under predetermined conditions, allowing sufficient time for the ligand conjugate to bind to activated macrophages associated with the active plaques, subjecting the blood vessel walls to the predetermined conditions using a catheter-based device,

and identifying active plaques by detecting light emitted by the chromophore using a catheter-based device.

[0008] A method is provided of identifying/monitoring active atherosclerotic plaques associated with blood vessel walls where the plaques comprise activated macrophages having accessible binding sites for a ligand. The method comprises the steps of administering to a patient suffering from atherosclerosis an effective amount of a composition comprising a conjugate of the general formula

L-X

wherein the group L comprises the ligand and the group X comprises a chemical moiety capable of emitting radiation, allowing sufficient time for the ligand conjugate to bind to the activated macrophages associated with the active plaques, and identifying active plaques by detecting radiation emitted by the chemical moiety using a catheter-based device.

[0009] The ligand can be any ligand that binds to a receptor which is preferentially expressed/presented on the surface of activated macrophages relative to resting macrophages. Such ligands include vitamins selected from the group consisting of folate, biotin, vitamin B₁₂, riboflavin, thiamine, and other vitamin receptor binding ligands.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010]

Fig. 1 shows the uptake of a folate-targeted ^{99m}Tc chelating chemical moiety (EC20) in the organs of atherosclerotic Watanabe rabbits.

Fig. 2 shows folate-FITC retention by macrophages harvested from hyperlipemic hamsters (H) compared to animals maintained on either normal (high folate) diet (N) or folic acid (FA) deficient diet (D).

Fig. 3 shows folate-FITC retention by macrophages harvested from hyperlipemic mice (H) compared to animals maintained on either normal (high folate) diet (N) or folic acid (FA) deficient diet (D).

Fig. 4 shows the serum levels of folate-FITC in hyperlipemic hamsters (H) compared to animals maintained on either normal (high folate) diet (N) or folic acid (FA) deficient diet (D).

Fig. 5 shows total serum cholesterol and triglyceride levels in hyperlipemic hamsters (H) compared to animals maintained on either normal (high folate) diet (N) or folic acid (FA) deficient diet (D).

Figs. 6a and 6b show fluorescence analysis of tissue fragments from atherosclerotic lesion-prone areas of Golden Syrian hamsters fed the atherosclerotic diet.

Figs. 7a and 7b show fluorescence analysis of macrophages harvested from the peritoneum of folic acid-Texas Red treated hypercholesterolemic hamsters.

Fig. 8 shows folate-FITC retention by U937 cells grown in either normal medium (a) or hyperlipidemic medium (b) or normal medium supplemented with LPS (c).

DETAILED DESCRIPTION OF THE INVENTION

[0011] The present invention relates to the use of a composition for identifying/monitoring active atherosclerotic plaques in blood vessel walls. In accordance with the invention a ligand that binds to a receptor which is preferentially expressed/presented on the surface of activated macrophages relative to resting macrophages, is conjugated to a chromophore capable of emitting light or a chemical moiety capable of emitting radiation and the ligand conjugates are administered to a patient being evaluated for atherosclerosis. The ligand conjugates bind to activated macrophages associated with active atherosclerotic plaques. The light or radiation emitted by the ligand-chromophore conjugate or the chemical moiety, respectively, is detected using a catheter-based device. Accordingly, the ligand conjugates can be used to distinguish active atherosclerotic plaques, containing activated macrophages, from inactive plaques wherein the plaques are present in the arteries or veins of a patient being evaluated for atherosclerosis.

[0012] In accordance with the invention, the word "catheter" means any catheter, guidewire, or other device capable of transluminal delivery (i.e., delivery into the lumen of blood vessels) of optical energy or of radiation, and/or any catheter, guidewire, or other device capable of detecting, in the lumen of blood vessels, light or radioactivity emitted from the ligand conjugates used in accordance with the method of the present invention, and/or any catheter, guidewire, or other device capable of delivering a therapeutic drug to the lumen of blood vessels.

[0013] In accordance with the present invention, the ligand conjugates can be formed from a wide variety of ligands, including any ligand that binds to a receptor expressed or presented on the surface of activated macrophages that is not expressed/presented or is not present in significant amounts on the surface of resting macrophages. Such ligands include N-formyl peptides (e.g., f-Met-Leu-Phe), high mobility group factor I protein (HMGB1), hyaluronan fragments, HSP-70, toll-like receptor ligands, scavenger receptor ligands, co-receptors for antigen presentation, ligands that bind to the CD68, BER-MAC3, RFD7, CD4, CD14, and HLA-D markers on activated macrophages, ligands that bind to

urokinase plasminogen activator receptors (e.g., the WX-360 peptide), antibodies, or fragments thereof, that bind preferentially to activated macrophages, and vitamins or receptor-binding vitamin analogs/derivatives. The ligand conjugates are capable of preferentially binding to activated macrophages compared to resting macrophages due to preferential expression of the receptor for the ligand on activated macrophages.

5 **[0014]** Acceptable vitamin moieties that can be used as ligands in accordance with the invention include niacin, pantothenic acid, folic acid, riboflavin, thiamine, biotin, vitamin B₁₂, and the lipid soluble vitamins A, D, E and K. These vitamins, and their receptor-binding analogs and derivatives, constitute the targeting entity that can be coupled with a chromophore or a chemical moiety, capable of emitting radiation, to form the ligand conjugates for use in accordance with the invention. Preferred vitamin moieties include folic acid, biotin, riboflavin, thiamine, vitamin B₁₂, and receptor-binding analogs and derivatives of these vitamin molecules, and other related vitamin receptor-binding molecules (see U.S. Patent No. 5,688,488).

[0015] Exemplary of a vitamin analog is a folate analog containing a glutamic acid residue in the D configuration (folic acid normally contains one glutamic acid in the L configuration linked to pteric acid).

10 **[0016]** In the ligand conjugates of the general formula L-X in accordance with the present invention, the group L is a ligand capable of binding to activated macrophages as compared to resting macrophages as described above. In one embodiment the activated macrophage binding ligand is folic acid, a folic acid analog/derivative or other folate receptor binding molecules. In another embodiment the activated macrophage binding ligand is a specific monoclonal or polyclonal antibody or Fab or scFv (i.e., a single chain variable region) fragments of an antibody capable of preferential binding to activated macrophages as compared to resting macrophages.

20 **[0017]** Activated macrophages express a 38 kD GPI-anchored folate receptor that binds folate and folate-derivatized compounds with subnanomolar affinity (i.e., < 1 nM). Importantly, covalent conjugation of small molecules, proteins, and even liposomes to folic acid does not alter the vitamin's ability to bind the folate receptor. Because most cells use an unrelated reduced folate carrier to acquire the necessary folic acid, expression of the folate receptor is restricted to a few cell types. With the exception of kidney, choroid plexus, and placenta, normal tissues express low or nondetectable levels of the folate receptor. However, many malignant tissues, including ovarian, breast, bronchial, and brain cancers express significantly elevated levels of the receptor. Also, it has recently been reported that the folate receptor β , the nonepithelial isoform of the folate receptor, is expressed in active form on activated, but not resting synovial macrophages.

25 **[0018]** The binding site for the ligand can include receptors for any ligand molecule, or a derivative or analog thereof, capable of preferentially binding to a receptor uniquely expressed or preferentially expressed/presented on the surface of activated macrophages. A surface-presented protein uniquely expressed or preferentially expressed by activated macrophages is a receptor that is either not present or is present at insignificant concentrations on resting macrophages providing a means for preferential detection of activated macrophages. Accordingly, any receptor that is upregulated on activated macrophages compared to resting macrophages, or which is not expressed/presented on the surface of resting macrophages, or any receptor that is not expressed/presented on the surface of resting macrophages in significant amounts could be used for targeting. In one embodiment the site that binds the ligand conjugates used in accordance with the present invention is a vitamin receptor, for example, the folate receptor, which binds folate or an analog or derivative thereof.

30 **[0019]** In accordance with the invention the ligand conjugates can bind with high affinity to receptors on activated macrophages. The high affinity binding can be inherent to the ligand or the binding affinity can be enhanced by the use of a chemically modified ligand (i.e., an analog or a derivative) or by the particular chemical linkage, in the ligand conjugate, between the ligand and the chromophore or between the ligand and the chemical moiety capable of emitting radiation.

35 **[0020]** The chemical linkage in the ligand conjugate between the ligand and the chromophore or between the ligand and the chemical moiety can be a direct linkage or can be through an intermediary linker. If present, an intermediary linker can be any biocompatible linker known in the art. Typically, the linker comprises about 1 to about 30 carbon atoms, more typically about 2 to about 20 carbon atoms. Lower molecular weight linkers (i.e., those having an approximate molecular weight of about 30 to about 300) are typically employed.

40 **[0021]** Generally, any manner of forming a complex between the ligand and the chromophore, between the ligand and the chemical moiety capable of emitting radiation, between a linker and the ligand, or between a linker and the chromophore or chemical moiety capable of emitting radiation can be utilized in accordance with the present invention.

45 With or without a linker, the complex can be formed by conjugation of the components of the conjugate, for example, through hydrogen, ionic, or covalent bonds. Covalent bonding of the components of the conjugate can occur, for example, through the formation of amide, ester, disulfide, or imino bonds between acid, aldehyde, hydroxy, amino, sulfhydryl, or hydrazo groups. Also, in accordance with this invention a linker can comprise an indirect means for associating the ligand with the chromophore/chemical moiety, such as by connection through spacer arms or bridging molecules. Both direct and indirect means for association should not prevent the binding of the ligand to the receptor on the activated macrophages for operation of the method of the present invention. Alternatively, the ligand conjugate can be one comprising a liposome wherein the chemical moiety capable of emitting radiation, for example, is contained within a liposome which is itself covalently linked to the activated macrophage-binding ligand.

[0022] In the embodiment where the ligand is folic acid, an analog/derivative of folic acid, or any other folate receptor binding molecule, the folate ligand can be conjugated to the chromophore/chemical moiety by an art-recognized procedure that utilizes trifluoroacetic anhydride to prepare γ -esters of folic acid via a pteroyl azide intermediate. This procedure results in the synthesis of a folate ligand, conjugated to the chromophore/chemical moiety only through the γ -carboxy group of the glutamic acid groups of folate. Alternatively, folic acid analogs can be coupled by art-recognized procedures through the α -carboxy moiety of the glutamic acid group or both the α and γ carboxylic acid entities.

[0023] The amount of the conjugate effective for use in accordance with the method of the invention depends on many parameters, including the molecular weight of the conjugate, its route of administration, and its tissue distribution. In accordance with the invention an "effective amount" of the ligand conjugate is an amount sufficient to bind to activated macrophages and to be useful in the identification/monitoring of active atherosclerotic plaques. The effective amount of the ligand conjugate to be administered to a patient being evaluated for atherosclerosis can range from about 1 ng/kg to about 10 mg/kg, or from about 10 μ g/kg to about 1 mg/kg, or from about 100 μ g/kg to about 500 μ g/kg.

[0024] The ligand conjugate can be administered in one or more doses (e.g., about 1 to about 3 doses) prior to the catheterization procedure. The number of doses depends on the molecular weight of the conjugate, its route of administration, and its tissue distribution, among other factors. When used for identification/monitoring of active atherosclerotic plaques, the catheterization procedure is typically performed about 1 to about 6 hours post-administration of the ligand conjugate targeted to activated macrophages, but the catheterization procedure can be performed at any time post-administration of the ligand conjugate as long as binding of the ligand conjugate to activated macrophages is detectable.

[0025] The ligand conjugates administered in accordance with the method of this invention are preferably administered parenterally to the patient being evaluated for atherosclerosis, for example, intravenously, intradermally, subcutaneously, intramuscularly, or intraperitoneally, in combination with a pharmaceutically acceptable carrier. Alternatively, the conjugates can be administered to the patient being evaluated for atherosclerosis by other medically useful procedures such as in an orally available formulation. In accordance with the invention, a "patient being evaluated for atherosclerosis" means any patient suspected of having atherosclerosis, whether symptomatic or not, who would benefit from an evaluation using the method of the present invention.

[0026] The conjugates used in accordance with this invention of the formula L-X are used in one aspect of this invention to formulate diagnostic compositions comprising effective amounts of the conjugate and an acceptable carrier therefor. Examples of parenteral dosage forms include aqueous solutions of the conjugate, for example, a solution in isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carriers such as alcohols, glycols, esters and amides. The parenteral compositions for use in accordance with this invention can be in the form of a reconstitutable lyophilizate comprising the one or more doses of the ligand conjugate. Any orally available dosage forms known in the art can also be used.

[0027] The activated macrophage-targeted conjugates used for identifying/monitoring disease states mediated by activated macrophages in accordance with this invention are formed to target and, thus, to concentrate the ligand conjugate at the site of activated macrophage populations (i.e., activated macrophages adhering to the luminal endothelial layer of the plaque or activated macrophages present in the lipid-rich core of the plaque) in the patient being evaluated for atherosclerosis.

[0028] In one embodiment of the invention active atherosclerotic plaques comprising activated macrophages are identified/monitored in a patient being evaluated for atherosclerosis by administering a conjugate of the formula L-X wherein L comprises a ligand capable of preferentially binding to activated macrophages, compared to resting macrophages, and X comprises a chromophore or a chemical moiety capable of emitting radiation. The inner lining of a patient's blood vessels is thereafter examined with a catheter-based device capable of detecting a localized concentration of the chromophore/chemical moiety conjugated to the ligand bound to activated macrophages.

[0029] The ligand conjugates are typically administered as a diagnostic composition comprising a ligand conjugate and a pharmaceutically acceptable carrier. The composition is typically formulated for parenteral administration and is administered to the patient in an amount effective to enable detection of the locale of activated macrophages. The nature of the chromophore/chemical moiety component of the ligand conjugate is dictated by the methodology used for catheter-based detection of the active atherosclerotic plaques. Thus, for example, the chromophore can comprise a fluorophore, such as fluorescein, (see PCT publication number WO 01/074382, for a description of a ligand-fluorophore conjugate) or another chromophore such as an hematoporphyrin, or a derivative thereof, or a Raman enhancing dye or agent, or a long wavelength fluorescent dye with optical properties that allow detection through many layers of tissue. The component of the ligand conjugate used for detection can also be a chemical moiety, such as a chelating moiety and a metal cation, for example, a radionuclide. It should be noted that the method of the present invention can be used for detecting light or radioactivity emitted from ligand conjugates bound both at the surface of atherosclerotic plaques and below the surface.

[0030] Such conjugates wherein the group L is folic acid, a folic acid analog/derivative, or another folic acid receptor binding ligand are described in detail in U.S. Patent No. 5,688,488. That patent, as well as related U.S. Patents Nos. 5,416,016 and 5,108,921, describe methods and examples for preparing conjugates useful in accordance with the

present invention. The present macrophage-targeted ligand conjugates can be prepared and used following general protocols described in those earlier patents.

5 [0031] In the embodiment where the ligand conjugate comprises a chromophore for use in identifying/monitoring active atherosclerotic plaques, the blood vessel walls can be subjected to predetermined conditions to detect locations on the inner linings of blood vessels where the ligand-chromophore conjugates are concentrated (i.e., active atherosclerotic plaques). Such predetermined conditions include any conditions known in the art to be useful for the detection of a chromophore, such as a fluorophore, using a catheter-based device. For example, the blood vessel walls can be subjected to radiation, in the ultraviolet, visible, or infrared region of the spectrum, from a laser. Catheter-based techniques employing optical fibers for the pulsed or steady state illumination of atherosclerotic plaques with laser radiation of a given wavelength can be used. A signal generated by the fluorescent light emitted by the ligand conjugates is then conveyed by one or more of the optical fibers to the end of the catheter where it can be analyzed to yield information about the atherosclerotic plaque being evaluated. The light emitted can be analyzed using art-recognized techniques as described below to identify/monitor the atherosclerotic plaque being evaluated.

10 [0032] In view of the increase in folate receptor levels during macrophage activation, a ligand conjugate comprising a ^{99m}Tc chelating chemical moiety targeted to activated macrophages using a vitamin, such as folate, can be used to detect active plaques *in vivo*. Such a ligand conjugate is described in U.S. Patent Application No. 60/378,571. Typically the activated macrophage-targeted ligand conjugate is administered to a patient, and following a period of time sufficient (e.g., from about 1 to about 24 hours) for the ligand conjugate to bind to activated macrophages associated with the active plaques, the patient is subjected to the catheterization procedure and identification/monitoring of active plaques is enabled by the targeted ligands conjugate.

15 [0033] Active atherosclerotic plaques can be identified/monitored in accordance with the method of the invention by, for example, spectral analysis of fluorescence emitted by the chromophore where the fluorescence emission is stimulated by radiation from, for example, a laser (e.g., laser-induced fluorescence spectroscopy), or by analysis of radioactivity emitted by the chemical moiety. Exemplary analytical techniques are described in U.S. Pats. Nos. 4,718,417 and 4,785,806, and in U.S. Patent Application Publication No. US 2003-0162234 A1, but any technique useful for analyzing light or radioactivity emitted from an atherosclerotic plaque to identify/monitor the atherosclerotic plaque in accordance with the invention can be used. In one embodiment, the fluorescence or radioactivity analysis is used to control an ablation laser, and accordingly, the ablation laser is activated, automatically or manually, after the diagnostic laser.

20 [0034] A variety of lasers known in the art can be used in the method of the invention. Exemplary lasers include holmium-doped yttrium aluminum garnet (YAG), holmium-doped yttrium lithium fluoride (YLF), and thulium-doped YAG and thulium-doped YLF. Further details regarding these and other suitable lasers are disclosed in U.S. Pats. Nos. 4,917,084 and 4,950,266,

25 [0035] The methods described in U.S. Pats. Nos. 5,217,456, 5,275,594, 5,562,100, 6,167,297, 6,217,847, 6,246,901, 6,387,350, 6,507,747, can also be used to stimulate emission of light from ligand-chromophore conjugates in accordance with the present invention and to detect/analyze light or radioactivity emitted from the ligand conjugates.

30 [0036] The method of the present invention can be used alone or in combination with any other method(s) known in the art for the detection/analysis/ablation of atherosclerotic plaques. For example, the invention can be used in combination with methods to ablate atherosclerotic plaques in cases where active plaques cause narrowing of blood vessels. In such cases, the ligand conjugates of the present invention can be used not only to identify active atherosclerotic plaques as compared to inactive plaques, but also to distinguish between atherosclerotic and normal tissue to help in ablation procedures. Thus, the present invention can be used to analyze both the physiological and the morphological state of atherosclerotic plaques. For example, angioplasty involves the nonsurgical widening of a vessel narrowed by plaque deposition, and laser energy, for example, directed through optical fibers in a catheter-based device, can be used to ablate or partially remove the plaque deposits. Catheter-based devices for ablating plaques using laser energy are described in U.S. Patents Nos. 4,817,601, 4,850,351, and 4,950,266.

35 [0037] When laser energy is used to ablate an atherosclerotic plaque, thermal damage to normal tissue is a serious risk because the energy level of radiation emitted from lasers used for ablation of plaque can damage or destroy normal tissue with the possibility of inadvertent perforation of an artery. Accordingly, the ligand conjugates of the present invention can be used to not only identify active atherosclerotic plaques, but to distinguish between atherosclerotic plaques and normal tissue to avert damage to normal tissue during plaque ablation. Pulsed laser emission can also be used whenever continuous laser exposure might damage the tissue.

40 [0038] The method of the present invention can also be used in combination with other techniques for differentiating between atherosclerotic plaques (e.g., fibrous plaque, calcified plaque, and lipid plaque) and normal tissue during plaque ablation. Such techniques include techniques based on analysis of laser-induced calcium photoemission from calcified plaque and laser-induced fluorescence from noncalcified plaque. Other such techniques include the analysis of fluorescence (e.g., laser-induced fluorescence), at selected wavelengths from tissue in an artery, with or without the use of a dye to enhance the contrast between the fluorescence emitted from atherosclerotic plaques and the fluorescence emitted from normal tissue (see U.S. Patents Nos. 4,641,650, 4,718,417, and 4,785,806.

[0039] Other laser-based techniques that can be used in combination with the method of the present invention to differentiate between atherosclerotic plaques and normal tissue include techniques utilizing laser-induced Raman light scattering and laser-induced plasma photoemission. Any other type of technique employing diagnostic and/or ablation lasers known in the art can also be used in combination with the method of the present invention (see U.S. Patents Nos. 4,817,601 and 4,850,351).

[0040] The method of the present invention can also be used in combination with any other method(s) known in the art for the detection/analysis/ ablation of atherosclerotic plaques, including the methods described in U.S. Patents Nos. 5,217,456, 5,275,594, 5,562,100, 6,167,297, 6,217,847, 6,246,901, 6,387,350, 6,507,747. Furthermore, the invention can be used to guide the positioning of therapeutic drugs and nucleic acid constructs positioned in the same catheter assembly or a different catheter assembly (see U.S. Patent Application Publication No. US 2002-0192157 A1).

EXAMPLE 1

Analysis of EC20 Uptake in the Organs of Atherosclerotic Rabbits

[0041] The Watanabe Heritable Hyperlipidemic rabbit model (i.e., a model for atherosclerosis) was used to determine whether binding of a folate-targeted ^{99m}Tc chelating chemical moiety (EC20; see U.S. Patent Application No. 60/378,571, could be detected in the arteries of atherosclerotic rabbits. Watanabe rabbits are LDL receptor deficient and, thus, provide a model for hypercholesterolaemia and atherosclerosis.

[0042] Watanabe rabbits (2 rabbits each 6 months old) were purchased from HRP-Covance (PA, USA). The rabbits were kept on a folate deficient diet for 3 weeks at which time the biodistribution of EC20 was evaluated. For the EC20 biodistribution studies, each rabbit was administered with 3.4 mCi of ^{99m}Tc and 7.4 x 10⁻⁹ moles of EC20 (EC20 in Fig. 1). One rabbit (i.e., control) also received 1000-fold excess of free folic acid (EC20 + FF in Fig. 1). The total injection volume for each rabbit was 400 μl via the marginal ear vein. The rabbits were sacrificed 4 hours after injection, and organs were extracted for EC20 biodistribution analysis (i.e., the radioactivity per gram of tissue was measured; see Fig. 1).

[0043] The results presented in Fig. 1 show that specific binding of ^{99m}Tc-EC20, competed by a 1000-fold excess of free folate, was detected in the aortic arch, the inferior vena cava, the superior vena cava, the pulmonary artery, the thoracic aorta, and the abdominal aorta of the Watanabe atherosclerotic rabbits.

EXAMPLE 2

Materials

[0044] Fmoc protected amino acid derivatives, Fmoc-glycine loaded Wang resin, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and N-hydroxybenzotriazole were purchased from Novabiochem (San Diego, CA). N¹⁰-trifluoroacetylptericoic acid was synthesized from folic acid (Sigma Chemical Company, St. Louis, MO) according to a previously published report (Godwin, 1972). The synthesis of gamma-carboxy linked folate-fluorescein (folate-FITC) has been described in a previous publication (Kennedy et al., 2003). RPMI Medium 1640 with L-glutamine, but without folic acid was from Gibco-BRL (Grand Island, N.Y., U.S.A.); folic acid and triglyceride reagent GPO-PAP were from DIALAB (Vienna, Austria); INFINITY CHOLESTEROL reagent and other chemicals were obtained from Sigma Chemical Co. (St. Louis, MO).

EXAMPLE 3

Synthesis of folate-Texas red conjugate

[0045] Standard Fmoc peptide chemistry was used to synthesize a folate-derivatized peptide linked to Texas Red via the gamma carboxyl of folic acid. The sequence Gly-Lys-(γ)Glu-ptericoic acid was constructed by Fmoc chemistry using HBTU and N-hydroxybenzotriazole as the activating agents along with diisopropylethylamine as the base and 20% piperidine in DMF for deprotection of the Fmoc groups. Fmoc-protected lysine containing a 4-methyltrityl protecting group on the ε-amine was linked to Fmoc-protected glycine attached to a Wang resin. An α-t-Boc protected N-α-Fmoc glutamic acid was then linked to the peptide to provide a γ-linked conjugate on folate after attaching N¹⁰-trifluoroacetylptericoic acid to the peptide. The methoxytrityl protecting group on the ε-amine of lysine was removed with 1% trifluoroacetic acid in dichloromethane to allow attachment of Texas Red. Texas Red N-hydroxysuccinimide (Molecular Probes, Eugene, OR) in DMF was reacted overnight with the peptide and then washed thoroughly from the peptide resin beads. The folic acid-Texas Red peptide was then cleaved from the resin with 95% trifluoroacetic acid:2.5% water:2.5% triisopropylsilane solution. Diethyl ether was used to precipitate the product and the precipitant was collected by centrifugation. The product was then washed twice with diethyl ether and dried under vacuum overnight. The product was then analyzed and

EP 1 629 281 B1

confirmed by mass spectroscopic analysis ($[M]$ calculated: 1423, found: 1422). In order to remove the N - trifluoroacetyl protecting group, the product was dissolved in 5 mls of water containing 0.5 mls of 10% ammonium hydroxide with the pH adjusted to 9.5-10.0 and stirred for 30 minutes at room temperature. The product was then precipitated using isopropanol/ether and the precipitant was collected by centrifugation. The product was then added to a G-10 Sephadex gel filtration column (1.5 x 15 cm) using water as the eluent. The product peaks were collected and lyophilized.

EXAMPLE 4

Animal model

[0046] Male healthy RAP mice and Golden Syrian Hamsters were selected for the experiments. Animals were divided in three experimental groups: animals fed with (i) normal diet (N) containing 0.1 g folic acid/ 100 kg; (ii) folic acid deficient diet (D), and (iii) folic acid deficient diet administered together with hypercholesterolemic food containing 3% cholesterol and 15% butter (H) for 6 months. Because commercial animal diets are supplemented with supraphysiological concentrations of folic acid, serum folate levels in these "normal" animals will often exceed natural levels by forty-fold (Wang et al.). Such high serum folate contents results in suppression of FR expression.

EXAMPLE 5

Folate Receptors on Macrophages in Hyperlipidemic Animals

[0047] Folic acid conjugated to either FITC (FA-FITC) or to Texas Red (FA-TR) were injected i.p. (10 μ g/ 100g body weight), and after 4 hours animals were bled (under light ether anesthesia) and then killed by carotid section. A macrophage-enriched suspension (also containing some mast cells, PMN, and monocytes) was then obtained by peritoneal lavage (Aviram, 1989). Briefly, the peritoneal cavity was injected with 10 ml cold phosphate buffered saline (PBS), pH 7.4, and gently massaged for 2 min. The suspension of peritoneal cells was filtered through nylon cloth and washed two times with PBS by centrifugation at 250 x g, 10 minutes at 4°C. The pellet was then solubilized in cold lysis buffer (1% NP-40, 50mM TRIS-HCl, 2mM EDTA, 1mM DTT, PMSF, protease inhibitors), cleared by centrifugation, and the fluorescence was quantified with a RF-5001 PC spectrofluorimeter using a standard curve. Protein concentration was determined by Amido Black and the results were expressed as a ratio of ng FA/mg protein for each group of animals. The same determinations were performed for the serum sample collected from each animal (ng FA-FITC/serum protein). In control experiments animals were injected with the FA-FITC conjugate in the presence of a 100 X excess of free folic acid. The total cholesterol and serum triglyceride concentration after six months of diet were measured using specific reagent kits.

[0048] In order to determine whether peritoneal macrophages from Golden Syrian hamsters express functional folate receptors, folate-FITC was injected into the hamsters intraperitoneally and uptake of the folate conjugate by macrophages extracted 4 hours later was examined by quantitative fluorimetry. The results in Fig. 2 show a significant increase in the folate conjugate's retention by macrophages harvested from hyperlipemic hamsters (H) compared to animals maintained on either normal (high folate) diet (N) or FA deficient diet (D). Importantly, the same differences were not detected in macrophages isolated from mice maintained on a similar diet (Fig. 3), a result that may be explained by the fact that mice are resistant to atherosclerosis, in contradistinction to hamsters which develop lesions similar to those of humans.

[0049] In all experimental groups, the serum level of FA-FITC 4 hours post injection was measured. The data indicate a higher concentration of the folate conjugate in serum of hyperlipemic hamsters, suggesting either an acquired deficiency in a clearance mechanism or an increase in folate conjugate binding by serum proteins has occurred in response to the high fat diet (see Fig. 4). The hyperlipidemic condition of the hamsters maintained on the hypercholesterolemic diet was also confirmed by the high level of both total serum cholesterol and triglyceride (Fig. 5). Taken together, the results in Figs. 2-5 show that the hypercholesterolemic diet induces abnormalities in folate homeostasis, including at least an increase in folate uptake by macrophages and a decrease in clearance of the conjugates from serum.

EXAMPLE 6

Preparation of tissue sections

[0050] Tissue fragments from atherosclerotic lesion-prone areas (aorta and valves) were collected from each experimental group and processed for fluorescence microscopy. The aorta was exposed, the branching arteries were cut off, and loose adventitial tissue was removed *in situ*. The vessel was then cut open and thoroughly washed with cold sterile PBS. Aorta and cardiac valves were carefully excised and all segments were fixed in 4% p-formaldehyde in PBS for 90 min at room temperature. The tissues were then immersed in OTC medium, flash-frozen in liquid nitrogen, and crysec-

tioned. The frozen sections were stained with oil red O and counterstained with hematoxylin (Mancini, 1995). Similar semithin cryosections were examined both in phase contrast and fluorescent microscopy using appropriate filters.

5 [0051] To determine whether common sites of atherosclerosis might be affected by altered folate homeostasis, tissue fragments from atherosclerotic lesion-prone areas were collected from Golden Syrian hamsters fed on the atherosclerotic diet and processed for fluorescence microscopy. Although animals maintained on either the normal or low folate diet did not develop atherosclerotic lesions and displayed normal serum concentrations of total cholesterol and triglycerides, within 6 months of transfer to the hyperlipidemic diet hamsters developed extensive lesions on both the aorta and cardiac valves. The lesions displayed a heterogeneous structure and are heavily laden with lipid, as revealed by Oil Red staining of the semithin cryosections (Fig. 6a). Importantly, adjacent unstained sections examined by fluorescence microscopy reveal significant uptake of the folate-Texas Red conjugate, as indicated by the prominent red fluorescence of the fatty lesions (Fig. 6b). Thus, intense FA-TR uptake is seen in distinct foci throughout the atherosclerotic valve, while low level FA-TR retention is observed rather uniformly across the fatty lesions. It is possible that the intense fluorescent spots constitute active microdomains where younger activated macrophages are accumulating, whereas the less intensely fluorescent regions may correspond to areas enriched in lipid laden cells (foam cells) that have gradually become more quiescent.

EXAMPLE 7

Specific uptake of FA- Texas Red in cultured peritoneal macrophages

20 [0052] Hamster peritoneal macrophages ($10\text{-}20 \times 10^6$ per hamster) were harvested from peritoneal fluid and washed by centrifugation 3 X in PBS at $1000 \times g$ for 10 minutes. Cells were suspended in 10 ml of RPMI Medium 1640 containing L-glutamine but lacking folic acid and supplemented with 10% fetal bovine serum (heat inactivated at 56°C for 30 minutes); 100 U/ml penicillin, and 100 $\mu\text{g/ml}$ streptomycin. Folic acid was left out of the culture medium to prevent down-
25 regulation of cell surface FR, which usually occurs when cells are cultured in the high folate concentrations present in unmodified RPMI. The peritoneal macrophages were then plated into 50 mm Petri dishes and cultured in a humidified incubator (5% CO_2 , 95% air). After 2 hours of incubation, the cells were washed to remove non-adherent cells, and further incubated under similar conditions for an additional 18 hours. The cells were briefly washed, mounted with slow fade and examined under a Texas red filter in the fluorescent microscope (Nikon).

30 [0053] Macrophages harvested from the peritoneum of FA-TR treated hypercholesterolemic hamsters were also found to be significantly more fluorescent (Fig. 7a) than macrophages similarly isolated from animals fed a normal lipid diet, demonstrating that the elevated uptake previously observed for folate-FITC (Fig. 2) can also be replicated using an independent folate conjugate, i.e., FA-TR. Since activated, but not resting macrophages have been reported to express a folate receptor (Nakashima-Matsushita, 1999), the data support the hypothesis that atherosclerosis is an inflammatory
35 disease in which activated macrophages are abundantly present (Haynes, 2002).

EXAMPLE 8

Specific uptake of FA-FITC in the U937 cell line

40 [0054] To extend the study of live animal models, cell culture experiments were also conducted using U937 cells (Harris, 1985), a cell line that originated from the pleural fluid of a patient with diffuse histiocytic lymphoma and which exhibits many characteristics of monocytes. To facilitate their expression of folate receptors, U937 cells were grown in folate-deficient RPMI supplemented with either 5% FCS or with 5% serum from patients with high cholesterol and glucose levels (Chl 295 mg/dl, Glc 315 mg/dl). In parallel experiments aimed at evaluating FR expression following a more classical method of U937 cell activation, U937 cells were incubated in RPMI without folic acid but supplemented with 5% FCS in the presence or absence of 0.5 g/ml bacterial lipopolysaccharide (LPS). After 24 hours of activation either with patient's serum or LPS, cells were incubated with 2.5 g/ml FA-FITC. The cells were then washed with PBS and solubilized in lysis buffer (50 mM Tris-HCl, 1% NP-40, 2 mM EDTA, and 1mM DTT and protease inhibitors), and FA-FITC uptake was assayed by spectrophotofluorimetry. Protein concentration was determined by BCA methods and the results were expressed as ratio of ng FA-FITC/mg protein for each experimental condition.

50 [0055] U937 cells (Sundstrom, 1976), are cells derived from committed progenitors of the monocyte lineage. In preliminary experiments, it was verified that uptake of FA-FITC by U937 cells is specifically folate receptor mediated by demonstrating that addition of 100-fold excess free folic acid would block uptake of the FA-FITC conjugates. FA-FITC retention by U937 cells grown in either normal medium (RPMI supplemented with 5% fetal calf serum) or hyperlipidemic medium (RPMI supplemented with 5% serum from a hypercholesterolemic patient) was then evaluated. As shown in Fig. 8b, by the end of the 4 hour incubation, higher FA-FITC uptake (1.6 X) was observed in U937 cells grown in medium supplemented with the hyperlipidemic serum than in cells maintained in normal medium (Fig. 8a). Importantly, similar

5 results were obtained when the cells were exposed to bacterial lipopolysaccharide (Fig. 8c), a ligand known to induce secretion of chemokines responsible for differentiation of U937 cells into cells with characteristics of an activated macrophage (Wang, 1998). In this case, however, uptake of FA-FITC was 1.8 X higher than in the absence of stimulation (Fig. 8a, c). Taken together, these experiments suggest that hypercholesterolemic conditions induce over-expression of folic acid receptors on macrophages, in a manner similar to direct activation of the macrophages by treatment with lipopolysaccharide.

10 **Claims**

1. The use of a composition comprising a conjugate of the general formula



15 wherein the group L comprises a folate receptor binding ligand and the group X comprises a chromophore capable of emitting light in the manufacture of a composition for identifying/monitoring active atherosclerotic plaques associated with blood vessel walls wherein the plaques comprise activated macrophages having accessible binding sites for the vitamin.

20 2. The use of claim 1 wherein the emitted light is detectable using a catheter-based device.

3. The use of claim 1 or 2 wherein the chromophore is a fluorophore.

25 4. The use of claim 3 wherein the fluorophore is fluorescein.

5. The use of claims 1 or 2 wherein L is an antibody or an antibody fragment capable of binding to the accessible vitamin-binding site of the activated macrophage.

30 6. The use of a composition comprising a conjugate of the general formula



35 wherein the group L comprises a folate receptor binding ligand and the group X comprises a chemical moiety capable of emitting radiation in the manufacture of a composition for identifying/monitoring active atherosclerotic plaques associated with blood vessel walls wherein the plaques comprise activated macrophages having accessible binding sites for the vitamin.

7. The use of claim 6 wherein the radiation is detectable using a catheter-based device.

40 8. The use of claim 6 or 7 wherein the chemical moiety comprises a metal chelating moiety.

9. The use of claim 8 wherein the chemical moiety further comprises a metal cation.

45 10. The use of claim 9 wherein the metal cation is a radionuclide.

11. The use of claims 6 to 7 wherein L is an antibody or an antibody fragment capable of binding to the accessible vitamin-binding site of the activated macrophage.

50 **Patentansprüche**

1. Verwendung einer Zusammensetzung, umfassend ein Konjugat der allgemeinen Formel



55 bei der die Gruppe L einen Folatrezeptor-Bindungsliganden und die Gruppe X ein Chromophor umfasst, welches zum Aussenden von Licht befähigt ist, in der Herstellung einer Zusammensetzung zur Identifizierung/Beobachtung von arteriosklerotischen aktiven Plaques, die mit Blutgefäßwänden verbunden sind, wobei die Plaques aktivierte

EP 1 629 281 B1

Makrophagen umfassen, die für das Vitamin zugängliche Bindungsstellen aufweisen.

2. Verwendung nach Anspruch 1, wobei das emittierte Licht unter Verwendung einer auf einem Katheter basierenden Vorrichtung nachweisbar ist.
3. Verwendung nach Anspruch 1 oder 2, wobei das Chromophor ein Fluorophor ist.
4. Verwendung nach Anspruch 3, wobei das Fluorophor Fluorescein ist.
5. Verwendung der Ansprüche 1 oder 2, wobei L ein Antikörper oder ein Antikörperfragment ist, der zur Bindung an die für das Vitamin zugängliche Bindungsstelle des aktivierten Makrophagen befähigt ist.
6. Verwendung einer Zusammensetzung, umfassend ein Konjugat der allgemeinen Formel

L-X

wobei die Gruppe L einen Folatrezeptor-Bindungsliganden und die Gruppe X einen chemischen Rest aufweist, der zum Aussenden von Strahlung befähigt ist, in der Herstellung einer Zusammensetzung zur Identifizierung/Beobachtung arteriosklerotischer aktiver Plaques, die mit Blutgefäßwandungen assoziiert sind, wobei die Plaques aktivierte Makrophagen aufweisen, die für das Vitamin zugängliche Bindungsstellen besitzen.

7. Verwendung nach Anspruch 6, wobei die Strahlung durch Verwendung einer Vorrichtung, die auf einem Katheter basiert, nachweisbar ist.
8. Verwendung nach Anspruch 6 oder 7, wobei der chemische Rest einen Metall-Chelat bildenden Rest aufweist.
9. Verwendung nach Anspruch 8, wobei der chemische Rest weiterhin ein Metallkation aufweist.
10. Verwendung nach Anspruch 9, wobei das Metallkation ein Radionuklid ist.
11. Verwendung der Ansprüche 6 bis 7, wobei L ein Antikörper oder ein Antikörperfragment ist, der zur Bindung an die für das Vitamin zugängliche Bindungsstelle des aktivierten Makrophagen befähigt ist.

Revendications

1. Utilisation d'une composition comprenant un conjugué de la formule générale :

L-X

dans laquelle le groupe L comprend un ligand de liaison de récepteur d'acide folique et le groupe X comprend un chromophore capable d'émettre de la lumière, dans la fabrication d'une composition destinée à identifier / surveiller les plaques athéroscléreuses actives associées aux parois des vaisseaux sanguins dans laquelle les plaques comprennent des macrophages activés ayant des sites de liaison accessibles pour la vitamine.

2. Utilisation selon la revendication 1, dans laquelle la lumière émise est détectable en utilisant un dispositif à base de cathéter.
3. Utilisation selon la revendication 1 ou 2, dans laquelle le chromophore est un fluorophore.
4. Utilisation selon la revendication 3, dans laquelle le fluorophore est la fluorescéine.
5. Utilisation selon la revendication 1 ou 2, dans laquelle L est un anticorps ou un fragment d'anticorps capable de se lier au site de liaison de vitamine accessible du macrophage activé.
6. Utilisation d'une composition comprenant un conjugué de la formule générale :

L-X

EP 1 629 281 B1

dans laquelle le groupe L comprend un ligand de liaison de récepteur d'acide folique et le groupe X comprend un fragment chimique capable d'émettre un rayonnement, dans la fabrication d'une composition destinée à identifier / surveiller les plaques athéroscléreuses actives associées aux parois des vaisseaux sanguins dans laquelle les plaques comprennent des macrophages activés ayant des sites de liaison accessibles pour la vitamine.

7. Utilisation selon la revendication 6, dans laquelle le rayonnement est détectable en utilisant un dispositif à base de cathéter.
8. Utilisation selon la revendication 6 ou 7, dans laquelle le fragment chimique comprend un fragment de chélation de métal.
9. Utilisation selon la revendication 8, dans laquelle le fragment chimique comprend en outre un cation métallique.
10. Utilisation selon la revendication 9, dans laquelle le cation métallique est un radionucléide.
11. Utilisation selon la revendication 6 ou 7, dans laquelle L est un anticorps ou un fragment d'anticorps capable de se lier au site de liaison de vitamine accessible du macrophage activé.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6a

FIG. 6b




FIG. 7a

FIG. 7b




FIG. 8

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5688488 A [0014] [0030]
- WO 01074382 A [0029]
- US 5416016 A [0030]
- US 5108921 A [0030]
- US 378571 P [0032] [0041]
- US 4718417 A [0033] [0038]
- US 4785806 A [0033] [0038]
- US 20030162234 A1 [0033]
- US 4917084 A [0034]
- US 4950266 A [0034] [0036]
- US 5217456 A [0035] [0040]
- US 5275594 A [0035] [0040]
- US 5562100 A [0035] [0040]
- US 6167297 A [0035] [0040]
- US 6217847 B [0035] [0040]
- US 6246901 B [0035] [0040]
- US 6387350 B [0035] [0040]
- US 6507747 B [0035] [0040]
- US 4817601 A [0036] [0039]
- US 4850351 A [0036] [0039]
- US 4641650 A [0038]
- US 20020192157 A1 [0040]

专利名称(译)	动脉粥样硬化的诊断方法		
公开(公告)号	EP1629281A4	公开(公告)日	2006-08-30
申请号	EP2004753487	申请日	2004-05-27
[标]申请(专利权)人(译)	普渡研究基金会		
申请(专利权)人(译)	普渡研究基金会		
当前申请(专利权)人(译)	普渡研究基金会		
[标]发明人	LOW PHILIP STEWART		
发明人	LOW, PHILIP, STEWART		
IPC分类号	G01N33/53 A61B A61K39/395 A61K49/00 G01N33/50 G01N33/92		
CPC分类号	A61K49/0052 A61K49/0041 A61K49/0043 A61K49/006 A61K51/0474 A61K51/0497 G01N33/5061 G01N33/92 G01N2800/323		
代理机构(译)	ELSY, DAVID		
优先权	60/474731 2003-05-30 US		
其他公开文献	EP1629281A2 EP1629281B1		
外部链接	Espacenet		

摘要(译)

本发明涉及鉴定/监测与血管壁相关的活动性动脉粥样硬化斑块的方法，其中所述斑块包含具有配体的可接近结合位点的活化巨噬细胞。该方法包括给予正在评估动脉粥样硬化的患者施用有效量的组合物的步骤，所述组合物包含配体和能够在预定条件下发光的发色团的缀合物，允许配体缀合物有足够的时间结合活化的巨噬细胞。使用基于导管的装置使血管经受预定条件，并通过使用基于导管的装置检测由发色团发射的光来识别活性斑块。本发明还涉及类似的方法，其中能够发射辐射的化学部分与配体缀合。

