



(12) **EUROPEAN PATENT APPLICATION**
 published in accordance with Art. 153(4) EPC

(43) Date of publication:
06.04.2016 Bulletin 2016/14

(51) Int Cl.:
G01N 33/53 (2006.01) G01N 33/543 (2006.01)
C07K 16/00 (2006.01) C12Q 1/02 (2006.01)

(21) Application number: **14797879.5**

(86) International application number:
PCT/JP2014/057227

(22) Date of filing: **18.03.2014**

(87) International publication number:
WO 2014/185145 (20.11.2014 Gazette 2014/47)

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
 Designated Extension States:
BA ME

(72) Inventors:
 • **OHNISHI, Arata**
Chiba-shi
Chiba 263-8555 (JP)
 • **MINAMIMOTO, Takafumi**
Chiba-shi
Chiba 263-8555 (JP)
 • **SUHARA, Tetsuya**
Chiba-shi
Chiba 263-8555 (JP)

(30) Priority: **16.05.2013 JP 2013104124**

(74) Representative: **Maikowski & Ninnemann**
Postfach 15 09 20
10671 Berlin (DE)


(71) Applicant: **Resvo Inc.**
Tokyo 143-0016 (JP)

(54) **BIOMARKER FOR PSYCHIATRIC AND NEUROLOGICAL DISORDERS**

(57) The purpose of the present invention is to provide: a biomarker for psychiatric and neurological disorders, in particular, a biomarker for diagnosing psychiatric and neurological disorders; and a test kit and test method for psychiatric and neurological disorders. The inventors discovered that there is a significant increase in the concentration of free κ immunoglobulin chains and free λ immunoglobulin chains in the blood samples from patients with psychiatric and neurological disorders. Consequently, the present invention provides a biomarker for psychiatric and neurological disorders that includes

at least one selected from the group consisting of free κ immunoglobulin chains, free λ immunoglobulin chains, and fragments thereof. Using this biomarker for psychiatric and neurological disorders enables the psychiatric and neurological disorders to be easily tested and diagnosed with the blood samples from the subjects. The biomarker enables more effective tests and diagnoses to be conducted combining inflammatory cytokines in the blood of patients with psychiatric and neurological disorders.

[Fig.5]



Description

Technical Field

5 **[0001]** The present invention relates to a biomarker for a psychiatric/neurological disorder, particularly a biomarker for diagnosing a psychiatric/neurological disorder. The present invention also relates to a test kit for a psychiatric/neurological disorder for detecting the biomarker, and a test method for a psychiatric/neurological disorder using the biomarker.

10 Background Art

[0002] Diagnosis criteria, such as DMSIV-TR, ICD-10, and PANSS, are currently used as indications in diagnosing psychiatric/neurological disorders, such as schizophrenia, autism, and Alzheimer's disease. However, diagnoses using these criteria are performed by, for example, a medical examination by interview with patients and thus have had the problem of producing variations depending on the evaluator. Thus, there has been a need for a biomarker as an objective biological indication.

15 **[0003]** Attempts have also been previously made to search for biomarkers for diagnosing psychiatric/neurological disorders. For example, Patent Literature 1 discloses a schizophrenia marker. Patent Literature 2 discloses a biomarker for diagnosing and monitoring mental disorder. However, these markers are not put to practical use in view of accuracy, easiness to practice, cost, and the like.

Citation List

25 Patent Literature

[0004]

Patent Literature 1: Japanese Patent Laid-Open No. 2012-013415

Patent Literature 2: Japanese Translation of PCT Application No. 2011-506995

30 Non-patent Literature

[0005]

35 Non-patent Literature 1: Serum reference intervals and diagnostic ranges for free kappa and free lambda immunoglobulin light chains: relative sensitivity for detection of monoclonal light chains. Katzmann JA, Clark RJ, Abraham RS, Bryant S, Lymp JF, Bradwell AR, Kyle RA. Clin Chem. 2002 Sep; 48(9): 1437-44.

40 Non-patent Literature 2: Assessment of proliferating cell nuclear antigen and its relationship with proinflammatory cytokines and parameters of disease activity in multiple myeloma patients. Tsirakis G, Pappa CA, Kaparou M, Katsomitrou V, Hatzivasili A, Alegakis T, Xekalou A, Stathopoulos EN, Alexandrakis MG. Eur J Histochem. 2011; 55(3): e21.

45 Non-patent Literature 3: Serum proinflammatory mediators at different periods of therapy in patients with multiple myeloma. Kuku I, Bayraktar MR, Kaya E, Erkurt MA, Bayraktar N, Cikim K, Aydogdu I. Mediators Inflamm. 2005 Aug 14; 2005(3): 171-4.

Summary of Invention

Technical Problem

50 **[0006]** An object of the present invention is to provide a biomarker for a psychiatric/neurological disorder, particularly a biomarker for diagnosing a psychiatric/neurological disorder, a test kit for a psychiatric/neurological disorder for detecting the biomarker, and a test method using the biomarker.

Solution to Problem

55 **[0007]** In view of the above-described problems, the present inventors conducted studies for identifying a biomarker for a psychiatric/neurological disorder. As a result, the present inventors have discovered that the blood concentration of the free κ immunoglobulin chain (containing the constant region) and the free λ immunoglobulin chain (containing the

constant region) is significantly increased in patients with psychiatric/neurological disorders compared to that in healthy individuals. Thus, the present invention provides a biomarker for a psychiatric/neurological disorder, comprising at least one selected from the group consisting of the free κ immunoglobulin chain, the free λ immunoglobulin chain, and fragments thereof. The biomarker can be used to readily conduct the test, diagnosis, and the like of a psychiatric/neurological disorder with a blood sample of interest.

[0008] In one embodiment, the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 1 (constant region) and the free λ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 6 (constant region). In one embodiment, the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 2 or 3 and the free λ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 6.

[0009] It is known that the free κ immunoglobulin chain increases also in the serum of patients having multiple myeloma (Non-patent Literature 1). It is known that in this case, an inflammatory cytokine increases simultaneously (Non-patent Literatures 2 and 3). The present inventors have surprisingly discovered that in patients with psychiatric/neurological disorders, the inflammatory cytokine does not increase despite an increase in the free κ immunoglobulin chain. Thus, the present invention provides a biomarker for a psychiatric/neurological disorder, comprising an inflammatory cytokine in addition to the above biomarker. In one embodiment, the inflammatory cytokine is at least one selected from the group consisting of IL-6, IL-1 beta, and TNF-alpha.

[0010] The present invention also provides a test kit for a psychiatric/neurological disorder, comprising a substance capable of detecting the biomarker for a psychiatric/neurological disorder. In addition, the present invention provides a test method for a psychiatric/neurological disorder, using the biomarker as an indication. In one embodiment, the test method for a psychiatric/neurological disorder according to the present invention comprises the steps of (a) measuring the amount of the biomarker for a psychiatric/neurological disorder according to the present invention in a blood sample from a subject and (b) comparing the amount with that from a normal subject or with the median value of that from a normal subject.

Advantageous Effects of Invention

[0011] The biomarker for a psychiatric/neurological disorder according to the present invention can be easily collected and quantified from a blood sample from a subject, and thus the use of the biomarker for a psychiatric/neurological disorder according to the present invention enables the rapid and simple diagnosis and the like of a psychiatric and neurological disorder.

Brief Description of Drawings

[0012]

[Figure 1] Figure 1 is a diagram explaining a κ immunoglobulin chain. Figure 1 (a) shows the construction and approximate molecular weight of the κ immunoglobulin chain. Figure 1 (b) shows the expression place and location of the κ immunoglobulin chain. The κ immunoglobulin chain (Ig κ) consists of a constant region of about 12 kDa called a κ immunoglobulin chain C region (Ig κ -C) and a variable region of about 12 kDa (Figure 1 (a)), and one in free form and one expressed in IgG (also including IgA and IgM) or on B cells are identified (Figure 1 (b)).

[Figure 2] Figure 2 shows the results of separating serum proteins obtained from a group of maternal immune activation model rats prepared by administering Poly IC to their mother bodies (schizophrenia and autism model animals) (MIA rats) (denoted by "Poly IC" in the figure) and a group of control rats prepared by administering saline to their mother bodies (denoted by "Saline" in the figure), by a 2-dimensional electrophoresis method. A difference was observed in a spot of about 25 kDa (Ref. Spot 1115).

[Figure 3] Figure 3 shows the results of the LC MS/MS measurement of the spot (Ref. Spot 1115, molecular weight: about 25 kDa) in which a difference was observed in the 2-dimensional electrophoresis gel image shown in Figure 2. These results showed that a concordant difference existed in the level of Ig κ -C expression in sera between 3 individuals in the MIA rat group and 3 individuals in the control rat group.

[Figure 4] Figure 4 shows the results of denaturing sera from the MIA rat group (denoted by "Poly IC" in the figure) and the control rat group (denoted by "Saline" in the figure) with SDS alone (100°C, 5 minutes), followed by Western blotting with an anti-Ig κ -C antibody. A significant difference was observed in a band of about 25 kDa ((2) in the figure) between the MIA rats and the control rats. The band was probably a free κ immunoglobulin chain since it was detected without the cleavage of S-S bonds or hydrogen bonds.

[Figure 5] Figure 5 shows the results of comparing the concentrations (mg/l) of a free κ immunoglobulin chain (A in the figure) and a free λ immunoglobulin chain (B in the figure) in sera from a group of schizophrenia patients (SZ) and a normal group. The concentrations of the free κ immunoglobulin chain (containing a constant region) and the free λ immunoglobulin chain (containing a constant region) significantly increased in schizophrenia patients (SZ).

As a result of comparing the concentration of the free κ immunoglobulin chain + the free λ immunoglobulin chain (C in the figure), a further significant difference was observed ($P = 0.00063$).

Description of Embodiments

(1. Definition)

[0013] As used herein, the term "psychiatric/neurological disorder" has the meaning generally used in the art, and examples thereof include schizophrenia, autism, Alzheimer's disease, cognitive abnormality, bipolar disorder (manic-depressive psychosis), disorders of cranial nerve development, cognitive impairment ascribed to neuropathy due to infection during pregnancy, mental disorder ascribed to impaired immunity, epilepsy, idiopathic mental disorder, toxic mental disorder, intellectual disability (mental retardation), psychopathy, and neurosis. In addition, the "psychiatric/neurological disorder" also includes syphilitic mental disorder, senile mental disorder, cerebrovascular mental disorder, mental disorder due to head injury, atypical endogenous psychosis, endocrine mental disorder and exogenous reaction type, and involuntional mental disorder. In the present invention, the psychiatric/neurological disorder is preferably schizophrenia, autism, or Alzheimer's disease, most preferably schizophrenia.

[0014] As used herein, the term "biomarker for a psychiatric/neurological disorder" refers to a living substance capable of being an indication for determining whether the psychiatric/neurological disorder have been developed or can be potentially developed, or determining prognosis after developing the disorders. For example, the biomarker for a psychiatric/neurological disorder can be used to diagnose the psychiatric/neurological disorder by measuring its amount. The biomarker for a psychiatric/neurological disorder may comprise a plurality of biomarkers. For example, when 2 or more living substances change in quantity in a subject with a psychiatric/neurological disorder, a combination of the 2 or more living substances can be a biomarker for the psychiatric/neurological disorder. For example, when certain 2 living substances simultaneously increase in a subject with another disorder or condition other than a psychiatric/neurological disorder but one of them increases while the other not increasing in a subject with the psychiatric/neurological disorder, a combination of the 2 living substances can also be a biomarker for the psychiatric/neurological disorder. The biomarker for a psychiatric/neurological disorder can be used, by measuring its amount, for the diagnosis and test of the psychiatric/neurological disorder, the screening of subjects having the psychiatric/neurological disorder, the establishment of an efficient therapeutic method, the screening of a candidate agent useful for the psychiatric/neurological disorder, and the development of model animals for the psychiatric/neurological disorder.

[0015] The term " κ immunoglobulin chain" and the term " λ immunoglobulin chain" mean the 2 types (κ type and λ type) of an immunoglobulin L chain, respectively. They are herein also referred to as "Ig κ " and "Ig λ ", respectively. The κ immunoglobulin chain (Ig κ) is a polypeptide consisting of a constant region of about 12 kDa called a κ immunoglobulin chain C region (Ig κ -C) and a variable region of about 12 kDa (see (a) in Figure 1), and one in free form and one expressed in IgG or on B cells are known (see (b) in Figure 1). The λ immunoglobulin chain (Ig λ) is an isoform of the κ immunoglobulin chain (Ig κ), and consists of a constant region and a variable region as is the case with the κ immunoglobulin chain (Ig κ).

[0016] The term " κ immunoglobulin chain C region" is a polypeptide forming the constant region of a κ immunoglobulin chain or a free κ immunoglobulin chain. The polypeptide is herein also referred to as "Ig κ chain C" or "Ig κ -C". As used herein, the " λ immunoglobulin chain C region" is a polypeptide forming the constant region of a λ immunoglobulin chain or a free λ immunoglobulin chain. The polypeptide is also referred to as "Ig λ chain C" or "Ig λ -C".

[0017] As used herein, the term "free κ immunoglobulin chain" and the term "free λ immunoglobulin chain" mean a κ immunoglobulin chain in free form and a λ immunoglobulin chain in free form, respectively (see (b) in Figure 1). The term "free" means being present alone without expression in IgG or on B cells or the like. The free κ immunoglobulin chain contains the κ immunoglobulin chain C region. The free κ immunoglobulin chain is herein also referred to as "free Ig κ ". The free λ immunoglobulin chain is also referred to as "free Ig λ ". The free λ immunoglobulin chain contains the λ immunoglobulin chain C region.

[0018] In the context of the biomarker for a psychiatric/neurological disorder according to the present invention herein, "fragment" refers to a polypeptide containing a partial amino acid sequence of a full-length protein (or a polypeptide). Thus, the "fragments" is a polypeptide which has a length within the full length of the biomarker for a psychiatric/neurological disorder according to the present invention and is at least one amino acid residue shorter, for example, 1 to 100, 1 to 50, 1 to 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid residue shorter, than the full length. Preferably, the fragment of the free κ immunoglobulin chain is the κ immunoglobulin chain C region or a site capable of being an antigen in detection with an antibody. Preferably, the fragment of the free λ immunoglobulin chain is the λ immunoglobulin chain C region or a site capable of being an antigen in detection with an antibody.

[0019] As used herein, the term "measurement" includes detection and quantification. The term "quantification" also includes semi-quantification. As used herein, the term "blood sample" refers to blood collected from a subject, and includes serum, plasma, and whole blood.

[0020] As used herein, the term "subject" means any mammal, for example, humans, rats, mice, guinea pigs, rabbits,

monkeys (for example, marmosets, particularly common marmosets, or rhesus monkeys), dogs, cats, or miniature pigs. The subject is preferably a human. As used herein, the term "normal subject" means a subject not having developed or potentially not developing a psychiatric/neurological disorder, for example, a human, a rat, a mouse, a guinea pig, a rabbit, a monkey (for example, marmosets, particularly common marmosets, or rhesus monkeys), a dog, a cat, or a miniature pig not having developed or potentially not developing a psychiatric/neurological disorder. The normal subject is preferably a human. The "normal subject" and "normal control" can be herein synonymously used. When the normal subject is a human, the normal subject is herein also referred to as a healthy individual.

(2. Biomarker for Psychiatric/Neurological Disorder According to Present Invention)

[0021] The biomarker for a psychiatric/neurological disorder according to the present invention comprises at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof. Preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof. More preferably, the biomarker for a psychiatric/neurological disorder according to the present invention comprises a free κ immunoglobulin chain or a fragment thereof. Alternatively, more preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a free κ immunoglobulin chain or a fragment thereof. More preferably, the biomarker for a psychiatric/neurological disorder according to the present invention comprises a free λ immunoglobulin chain or a fragment thereof. Alternatively, more preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a free λ immunoglobulin chain or a fragment thereof. More preferably, the biomarker for a psychiatric/neurological disorder according to the present invention comprises a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof. Alternatively, more preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof.

[0022] In another embodiment, the biomarker for a psychiatric/neurological disorder according to the present invention comprises at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof, and an inflammatory cytokine. Preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof, and an inflammatory cytokine. More preferably, the biomarker for a psychiatric/neurological disorder according to the present invention comprises a free κ immunoglobulin chain or a fragment thereof, and an inflammatory cytokine. Alternatively, more preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a free κ immunoglobulin chain or a fragment thereof, and an inflammatory cytokine. More preferably, the biomarker for a psychiatric/neurological disorder according to the present invention comprises a free λ immunoglobulin chain or a fragment thereof, and an inflammatory cytokine. Alternatively, more preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a free λ immunoglobulin chain or a fragment thereof, and an inflammatory cytokine. More preferably, the biomarker for a psychiatric/neurological disorder according to the present invention comprises a free κ immunoglobulin chain or a fragment thereof, a free λ immunoglobulin chain or a fragment thereof, and an inflammatory cytokine. Alternatively, more preferably, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a free κ immunoglobulin chain or a fragment thereof, a free λ immunoglobulin chain or a fragment thereof, and an inflammatory cytokine.

[0023] In one embodiment, the inflammatory cytokine is at least one selected from the group consisting of IL-6, IL-1 beta, and TNF-alpha. Preferably, the inflammatory cytokine is at least two selected from the group consisting of IL-6, IL-1 beta, and TNF-alpha. More preferably, the inflammatory cytokine is IL-6, IL-1 beta, and TNF-alpha. In one embodiment, the biomarker for a psychiatric/neurological disorder according to the present invention is present in a blood sample collected from a subject.

[0024] The amino acid sequence of a free κ immunoglobulin chain is well-known to those of ordinary skill in the art. The human-derived free κ immunoglobulin chain comprises a constant region (κ immunoglobulin chain C region). For example, the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 1 (C region). Preferably, the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 2 or 3. Those of ordinary skill in the art can detect and quantify the free κ immunoglobulin chain using a substance capable of recognizing the constant region (κ immunoglobulin chain C region) or an antigen site thereof. Examples of the substance include Anti-Ig κ light chain antibody (MAR-C8) (Abcam Co., Ltd.) and FREELITE κ chain (MBL Co., Ltd., code number: BS-LK016BD).

[0025] The constant region (κ immunoglobulin chain C region) preferably comprises the amino acid sequence of SEQ ID NO: 1. Alternatively, the constant region (κ immunoglobulin chain C region) preferably consists of the amino acid sequence of SEQ ID NO: 1. More preferably, the constant region (κ immunoglobulin chain C region) comprises the amino acid sequence of SEQ ID NO: 2 or 3. The antigen site of the constant region (κ immunoglobulin chain C region)

preferably comprises the amino acid sequence of SEQ ID NO: 2 or 3. Alternatively, the antigen site of the constant region (κ immunoglobulin chain C region) preferably consists of the amino acid sequence of SEQ ID NO: 2 or 3. The fragment of the free κ immunoglobulin chain preferably comprises the constant region (κ immunoglobulin chain C region) or an antigen site thereof. Alternatively, the fragment of the free κ immunoglobulin chain preferably consists of the constant region (κ immunoglobulin chain C region) or an antigen site thereof.

[0026] A free κ immunoglobulin chain comprising the amino acid sequence of SEQ ID NO: 4 (rat-derived) or SEQ ID NO: 5 (rat-derived) can be used to test, diagnose, and/or determine whether psychiatric/neurological disorder model animals (particularly, rats) have developed or can potentially develop the psychiatric/neurological disorder. It is useful, particularly for the preparation, quality control, or the like of psychiatric/neurological disorder model animals. The test method can be similarly performed by the method shown in the section "Test Method for Psychiatric/Neurological Disorder" below. Thus, in another embodiment, the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 4 (C region). Preferably, the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 5. The constant region (κ immunoglobulin chain C region) preferably comprises the amino acid sequence of SEQ ID NO: 4. Alternatively, the constant region (κ immunoglobulin chain C region) preferably consists of the amino acid sequence of SEQ ID NO: 4. The antigen site of the constant region (κ immunoglobulin chain C region) preferably comprises the amino acid sequence of SEQ ID NO: 5. Alternatively, the antigen site of the constant region (κ immunoglobulin chain C region) preferably consists of the amino acid sequence of SEQ ID NO: 5. The fragment of the free κ immunoglobulin chain preferably comprises the constant region (κ immunoglobulin chain C region) or an antigen site thereof. Alternatively, the fragment of the free κ immunoglobulin chain preferably consists of the constant region (κ immunoglobulin chain C region) or an antigen site thereof.

[0027] The amino acid sequence of a free λ immunoglobulin chain is also well-known to those of ordinary skill in the art. The free λ immunoglobulin chain comprises a constant region (λ immunoglobulin chain C region). For example, the free λ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 6. Preferably, the free λ immunoglobulin chain comprises the antigen site of the constant region (λ immunoglobulin chain C region). Those of ordinary skill in the art can readily detect and quantify the free λ immunoglobulin chain using a substance capable of recognizing the constant region (λ immunoglobulin chain C region) or an antigen site thereof. Examples of the substance include Anti-Ig λ light chain (human) pAb-FITC (MBL Co., Ltd.) and EREELITE λ chain (MBL Co., Ltd., code number: BS-LK018BD).

[0028] The constant region (λ immunoglobulin chain C region) preferably comprises the amino acid sequence of SEQ ID NO: 6. Alternatively, the constant region (λ immunoglobulin chain C region) preferably consists of the amino acid sequence of SEQ ID NO: 6. The fragment of the free λ immunoglobulin chain preferably comprises the λ immunoglobulin chain C region or an antigen site thereof. Alternatively, the fragment of the free λ immunoglobulin chain preferably consists of the λ immunoglobulin chain C region or an antigen site thereof.

[0029] The relation between SEQ ID NO and the κ immunoglobulin chain C region or the λ immunoglobulin chain C region is as follows.

- SEQ ID NO: 1 - κ immunoglobulin chain C region (human-derived)
- SEQ ID NO: 2 - antigen site (human-derived)
- SEQ ID NO: 3 - antigen site (human-derived)
- SEQ ID NO: 4 - κ immunoglobulin chain C region (rat-derived)
- SEQ ID NO: 5 - antigen site (rat-derived)
- SEQ NO: 6 - λ immunoglobulin chain C region (human-derived)

[0030] The free κ immunoglobulin chain is a polypeptide consisting of a constant region (λ immunoglobulin chain C region) of about 12 kDa and a variable region of about 12 kDa (see Figure 1 (a)). Thus, the measurement of the polypeptide amount of the κ immunoglobulin chain C region will mean to measure the amount of the free κ immunoglobulin chain. Similarly, the measurement of the polypeptide amount of the λ immunoglobulin chain C region will mean to measure the amount of the free λ immunoglobulin chain.

[0031] In another embodiment, the biomarker for a psychiatric/neurological disorder According to the present invention comprises a polypeptide containing the antigen site of a free κ immunoglobulin chain, or a fragment thereof. Alternatively, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a polypeptide containing the antigen site of a free κ immunoglobulin chain, or a fragment thereof. In still another embodiment, the biomarker for a psychiatric/neurological disorder According to the present invention comprises a polypeptide containing a free κ immunoglobulin chain C region, or a fragment thereof. Alternatively, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a polypeptide containing a free κ immunoglobulin chain C region, or a fragment thereof. In another embodiment, the biomarker for a psychiatric/neurological disorder According to the present invention comprises a polypeptide containing the antigen site of a free λ immunoglobulin chain, or a fragment thereof. Alternatively, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a polypeptide containing the antigen site of a free λ immunoglobulin chain, or a fragment thereof. In still another embodiment,

the biomarker for a psychiatric/neurological disorder According to the present invention comprises a polypeptide containing a free λ immunoglobulin chain C region, or a fragment thereof. Alternatively, the biomarker for a psychiatric/neurological disorder according to the present invention consists of a polypeptide containing a free λ immunoglobulin chain C region, or a fragment thereof.

5 **[0032]** Those of ordinary skill in the art will understand that deletion, addition, variation, or substitution in the amino acid sequence of each of the polypeptides having the above amino acid sequences can be permitted provided that the uniqueness of the polypeptide, i.e., a specific amino acid sequence characterizing the polypeptide (e.g., a motif, a domain, a box sequence, or the like), or the like, is retained. The number of such deletions, additions, variations, or substitutions is not limited (for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) provided that the uniqueness of the biomarker
10 for a psychiatric/neurological disorder According to the present invention is retained. In general, variation or substitution between amino acids having the same properties has no or little effect on the function of the polypeptide; thus, a polypeptide containing the variation or substitution can be equated with the original polypeptide (for example, a polypeptide specified by any of the above SEQ NOS). The properties of such amino acids can be categorized, for example, as follows: acidic amino acids (aspartic acid and glutamic acid), basic amino acids (lysine, histidine, and arginine), hydrophobic (aliphatic) amino acids (glycine, alanine, valine, leucine, and isoleucine), sulfur-containing amino acids (methionine and cysteine), aromatic amino acids (phenylalanine, tryptophan, and tyrosine), acidic amino acid amides (asparagine and glutamine), or neutral amino acids (glycine, alanine, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan, proline, serine, threonine, cysteine, methionine, asparagine, and glutamine).

20 **[0033]** For example, an amino acid sequence in which Glu at the 105th position and Cys at the 106th position are deleted in the amino acid sequence of SEQ NO: 1 is also included within the scope of the present invention.

[0034] Those of ordinary skill in the art will understand that the biomarker for a psychiatric/neurological disorder according to the present invention can also include one modified, for example, with phosphorylation, sugar chain addition, amidation, ubiquitination, or acetylation.

25 **[0035]** The biomarker for a psychiatric/neurological disorder According to the present invention can be present in a blood, mine, or spinal fluid sample from a subject. The sample is preferably a blood sample (serum, plasma, or whole blood). The blood sample is more preferably serum. The collection of a blood sample from a subject can be performed by a common method well-known to those of ordinary skill in the art. The blood sample can be collected, for example, using a syringe.

30 **[0036]** All of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof in the blood sample of the subject increase significantly. Thus, their change can be measured to determine the subject as having developed or capable of potentially developing the psychiatric/neurological disorder.

35 **[0037]** The free κ immunoglobulin chain may also increase, for example, in patients with multiple myeloma. However, it is known that an inflammatory cytokine increases simultaneously in this case (Non-patent Literatures 1 to 3). On the other hand, the present inventors have found that in a subject with psychiatric/neurological disorder, the free κ immunoglobulin chain increases while an increase in an inflammatory cytokine is not observed. Thus, a psychiatric/neurological disorder can be more clearly determined by the combination of the free κ immunoglobulin chain and the inflammatory cytokine, i.e., by using the amount of the free κ immunoglobulin chain or a fragment thereof and the amount of the inflammatory cytokine as indications. The amount of the free λ immunoglobulin chain or a fragment thereof and the amount of the inflammatory cytokine may also be used as indications. Most preferably, the three of the free κ immunoglobulin chain or a fragment thereof, the free λ immunoglobulin chain or a fragment thereof, and the inflammatory cytokine are used as indications.

40 **[0038]** The biomarker for a psychiatric/neurological disorder According to the present invention is preferably a biomarker for schizophrenia, autism, disorders of cranial nerve development, cognitive impairment ascribed to neuropathy due to infection during pregnancy, or mental disorder ascribed to impaired immunity. Most preferably it is a biomarker for schizophrenia.

45 **[0039]** The details of the determination of the development or potential development of a psychiatric/neurological disorder will be described in "Test Method for Psychiatric/Neurological Disorder" below.

(3. Test Kit for Psychiatric/Neurological Disorder)

50 **[0040]** The present invention provides a test kit for a psychiatric/neurological disorder. The test kit can be used to measure the amount of the biomarker for a psychiatric/neurological disorder According to the present invention in a blood sample from a subject to determine whether the subject has developed or is at risk of developing the psychiatric/neurological disorder.

55 **[0041]** The test kit of the present invention contains a substance capable of detecting the biomarker for a psychiatric/neurological disorder according to the present invention. When the biomarker for a psychiatric/neurological disorder comprises a plurality of biomarkers, the kit can contain a plurality of substances capable of detecting the plurality of biomarkers respectively. The substance is not particularly limited; however, it can be any molecule having affinity to the

biomarker for a psychiatric/neurological disorder according to the present invention. The substance can be a substance capable of detecting a free κ immunoglobulin chain C region or an antigen site thereof. The substance can be a substance capable of detecting a free λ immunoglobulin chain C region or an antigen site thereof. The substance includes, for example, an antibody, a low molecular compound, a high molecular compound, a protein, a peptide, and a nucleic acid.

The substance is preferably labeled with any marker, such as a dye or fluorescent dye, a radioactive isotope, and an enzyme so that the biomarker for a psychiatric/neurological disorder According to the present invention can be detected. The substance is more preferably an antibody. Preferably, the antibody is an anti-Ig κ antibody, an anti-Ig κ -C antibody, an anti-Ig λ antibody, or an anti-Ig λ -C antibody. The antibody may be a monoclonal antibody or a polyclonal antibody.

[0042] Examples of the substance capable of detecting the biomarker for a psychiatric/neurological disorder according to the present invention include Anti-Ig κ light chain antibody (MAR-C8) (Abcam Co., Ltd.), FREELITE κ chain (MBL Co., Ltd., code number: BS-LK016BD), Anti-Ig λ light chain (human) pAb-FITC (MBL Co., Ltd.), and FREELITE λ chain (MBL Co., Ltd., code number: BS-LK018BD).

[0043] The antibody may be prepared by a method well-known to those of ordinary skill in the art. For example, the biomarker for a psychiatric/neurological disorder According to the present invention can be administered to a mammal, such as a rabbit, for immunization to prepare an antibody from its serum. The anti-Ig κ antibody can be prepared by administering the polypeptide of the κ immunoglobulin chain C region or an antigen site thereof to a mammal, such as a rabbit. For example, the anti-Ig κ antibody is prepared by administering a polypeptide comprising the amino acid sequence of any one of SEQ ID NOS: 1 to 5 or a polypeptide consisting of the amino acid sequence of any one of SEQ ID NOS: 1 to 5. The anti-Ig λ antibody can be prepared by administering the polypeptide of the λ immunoglobulin chain C region or an antigen site thereof to a mammal, such as a rabbit. For example, the anti-Ig λ antibody is prepared by administering a polypeptide comprising the amino acid sequence of SEQ ID NO: 6 or a polypeptide consisting of the amino acid sequence of SEQ ID NO: 6. The antibody may be labeled with any marker, such as a dye (e.g., a fluorescent dye), a radioactive isotope, and an enzyme.

[0044] The test kit for a psychiatric/neurological disorder according to the present invention may further contain a labeled secondary antibody. The labeling can be performed using any marker, such as a dye (e.g., a fluorescent dye), a radioactive isotope, and an enzyme.

[0045] In a particular embodiment, the substance capable of detecting the biomarker for a psychiatric/neurological disorder according to the present invention, for example, an antibody, is immobilized on a solid-phase support. The solid-phase support may be made of any appropriate material, such as a plastic (e.g., polystyrene resin, polycarbonate resin, polyethylene resin, or polypropylene resin) or glass. The solid-phase support may be a substrate of any shape, such as a dish, a well, a strip, or a chip. The substrate surface may be optionally coated and modified.

[0046] The test kit for a psychiatric/neurological disorder according to the present invention may contain instructions showing a method for measuring the amount of the biomarker for a psychiatric/neurological disorder according to the present invention.

[0047] In addition, the test kit for a psychiatric/neurological disorder according to the present invention may contain, for example, reagents (for example, a buffer solution, a blocking reagent, an enzyme substrate, and a fluorescent reagent) and/or devices or instruments (for example, a container, a reaction device, a fluorescence measuring instrument) used in the test method for a psychiatric/neurological disorder.

[0048] The test kit for a psychiatric/neurological disorder according to the present invention is preferably a test kit for schizophrenia, autism, disorders of cranial nerve development, cognitive impairment ascribed to neuropathy due to infection during pregnancy, or mental disorder ascribed to impaired immunity. Most preferably, it is a test kit for schizophrenia.

[0049] The test kit for a psychiatric/neurological disorder according to the present invention can be used for determining the development or potential development of a psychiatric/neurological disorder and prognosis after developing the disorder. The details of the determination will be described in the item of "Test Method for Psychiatric/Neurological Disorder" below. The test kit for a psychiatric/neurological disorder according to the present invention can be used for diagnosing a psychiatric/neurological disorder. The test kit for a psychiatric/neurological disorder according to the present invention can be used in a test method for a psychiatric/neurological disorder shown below.

(4. Test Method for Psychiatric/Neurological Disorder)

[0050] The present invention provides a test method for a psychiatric/neurological disorder. The test method uses the amount of the biomarker for a psychiatric/neurological disorder according to the present invention in a blood sample from a subject as an indication. The test method of the present invention is useful for determining the development or potential development of a psychiatric/neurological disorder and prognosis after developing the disorder. The test method of the present invention can also be used for diagnosing a psychiatric/neurological disorder. The test method of the present invention can also provide data necessary for the treatment, prevention, diagnosis, differentiation, or prognosis estimation of a psychiatric/neurological disorder. The test method of the present invention can also provide information

useful for determining the development or potential development of a psychiatric/neurological disorder and prognosis after developing the disorder. In addition, a treatment method and a treatment regimen for a subject can be decided from the results obtained using the test method of the present invention. Further, a primary screening of a psychiatric/neurological disorder in a medical checkup and the like can be performed using the test method of the present invention. A screening of a psychiatric/neurological disorder can also be carried out in the first contact of patients in psychiatry.

[0051] The test method for a psychiatric/neurological disorder according to the present invention is preferably a test method for schizophrenia, autism, disorders of cranial nerve development, cognitive impairment ascribed to neuropathy due to infection during pregnancy, or mental disorder ascribed to impaired immunity. Most preferably, it is a test kit for schizophrenia.

[0052] The test method of the present invention can comprise a step of measuring the amount of the biomarker for a psychiatric/neurological disorder according to the present invention in a subject and a step of comparing the measured amount of the biomarker for a psychiatric/neurological disorder according to the present invention with that in a normal subject. The test method of the present invention can comprise the steps of (a) measuring the amount of at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof in a blood sample from a subject and (b) comparing the amount with that from a normal subject. Preferably, the test method of the present invention can comprise the steps of (a) measuring the amount of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject and (b) comparing the amount with that from a normal subject. The comparing step comprises a step of comparison with the median value in the normal control group. The amount is preferably an amount by combining, i.e., the total amounts of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject.

[0053] In another embodiment, the test method of the present invention can comprise the steps of (a) measuring the amount of at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof and the amount of an inflammatory cytokine in a blood sample from a subject and (b) comparing the amount with that from a normal subject. More preferably, the test method of the present invention comprises the steps of (a) measuring the amount of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof and the amount of an inflammatory cytokine in a blood sample from a subject and (b) comparing these amounts with those from a normal subject. The comparing step comprises a step of comparison with the median value in the normal control group. The amount is preferably an amount by combining, i.e., the total amounts of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject.

[0054] The details of a psychiatric/neurological disorder, a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof, and an inflammatory cytokine have been described in the section "Biomarker for Psychiatric/Neurological Disorder According to Present Invention" and will not be again described.

[0055] In the step (a) of the test method of the present invention, the measurement includes detection and quantification. The level of the quantification can be a level that enables the determination of the development or potential development of a psychiatric/neurological disorder or the determination of prognosis after developing the disorders. The quantification may be semi-quantification. The measurement of a free κ immunoglobulin chain may be the detection or quantification of a free κ immunoglobulin chain C region and/or an antigen site thereof. The measurement of a free λ immunoglobulin chain may be the detection of a free λ immunoglobulin chain C region and/or an antigen site thereof. The measurement may be the measurement of the amount of mRNA encoding the biomarker for a psychiatric/neurological disorder according to the present invention.

[0056] In the step (a) of the test method of the present invention, the measurement of the amount of the biomarker for a psychiatric/neurological disorder according to the present invention can be carried out by a common method well-known to those of ordinary skill in the art. Examples thereof include a one- or two-dimensional electrophoresis method, a Western blotting method, an enzyme linked immunosorbent assay (ELISA), a sandwich ELISA method, an immunoelectrophoresis method, single radial immunodiffusion (SRID), a radioimmunoassay (RIA), an enzyme immunoassay (EIA), a latex immunoassay (LIA), and a fluorescence immunoassay (FIA). The amount of the biomarker for a psychiatric/neurological disorder according to the present invention can also be measured using the detection kit for a psychiatric/neurological disorder according to the present invention. The amount of the biomarker for a psychiatric/neurological disorder according to the present invention can also be measured using, for example, a device (e.g., a radiation counter, an optical measurement device, or a fluorescence intensity measuring device) and an image analysis software which are well-known to those of ordinary skill in the art. The amount of the biomarker for a psychiatric/neurological disorder according to the present invention may be measured once or a plurality of times.

[0057] In the step (a) of the test method of the present invention, the amount of the biomarker for a psychiatric/neurological disorder according to the present invention may be a concentration (e.g., mg/l, μ g/l, or pg/ml), a radiation intensity, or a fluorescence intensity. The unit of the amount of the biomarker for a psychiatric/neurological disorder

according to the present invention will depend on the measurement method. In a certain embodiment, the amount of the biomarker for a psychiatric/neurological disorder according to the present invention may be a relative ratio to that in a normal subject.

5 [0058] The blood sample may be serum, plasma, or whole blood. The blood sample is preferably serum. The method for collecting the blood sample may be a common method, such as collection using a syringe or the like from a subject, and is not particularly limited. The blood sample collected from a subject may be denatured and/or stored by a method well-known to those of ordinary skill in the art.

10 [0059] Before measuring the blood sample, the blood sample collected from a subject can be prepared using a reagent and a means which are well-known to those of ordinary skill in the art. Examples of the reagent include buffers, such as acetic acid-acetate, phosphoric acid-phosphate, citric acid-citrate, or tris(hydroxymethyl)aminomethane (tris); denaturing agents, such as urea and/or thiourea; reducing agents, such as dithiothreitol (DTT); and surfactants, such as sodium lauryl sulfate (SDS). Examples of the means include filtration, ultrafiltration, centrifugation, homogenization, incubation, shaking, and lyophilization.

15 [0060] In the test method of the present invention, when the free κ immunoglobulin chain, the free λ immunoglobulin chain, and fragments thereof in a blood sample from a subject are measured, denaturation (cleavage of S-S bonds) by DTT may not be conducted. Urea and/or thiourea may also not be used. For example, the blood sample may be denatured only with SDS. The reason therefor is to eliminate the κ immunoglobulin chain and the λ immunoglobulin chain expressed in IgG, IgA, IgM and the like and efficiently measure only the free κ immunoglobulin chain, the free λ immunoglobulin chain, and fragments thereof.

20 [0061] In the step (a) of the test method of the present invention, the subject is not particularly limited. The subject is preferably a human. The subject is, for example, a subject for whom it is necessary to be determined whether or not the subject has developed a psychiatric/neurological disorder. The subject may be a subject determined as having developed a psychiatric/neurological disorder using conventional diagnosis criteria, such as DMSIV-TR, ICD-10, and PANSS. The subject may be a subject not determined as having developed a psychiatric/neurological disorder using conventional diagnosis criteria, such as DMSIV-TR, ICD-10, and PANSS. In addition, the subject may be a subject for whom no information about a psychiatric/neurological disorder is obtained. It may be confirmed in advance whether the subject has a cancer and/or an inflammatory disorder.

25 [0062] In preparing a blood sample, the blood sample is preferably incubated at about 60°C, about 70°C, about 80°C, about 90°C, about 100°C, about 110°C, or about 120°C for about 1 minute, about 2 minutes, about 3 minutes, about 4 minutes, about 5 minutes, about 6 minutes, about 7 minutes, about 8 minutes, about 9 minutes, or about 10 minutes. Most preferably, it may be incubated at about 100°C for about 5 minutes.

30 [0063] In the step (b), the normal subject is a subject not having developed or potentially not developing a psychiatric/neurological disorder. For example, the subject is a subject determined as not having developed a psychiatric/neurological disorder using conventional diagnosis criteria, such as DMSIV-TR, ICD-10, and PANSS. The method for the comparison in the step (b) may be a method well-known to those of ordinary skill in the art and is not particularly limited. For example, the measured value obtained in the step (a) can be compared with the measured value of the biomarker for a psychiatric/neurological disorder according to the present invention in a normal subject obtained in advance. Preferably, the results obtained by a two-dimensional electrophoresis method or an enzyme linked immunosorbent assay (ELISA) are compared. When the biomarker for a psychiatric/neurological disorder according to the present invention comprises a plurality of biomarkers, the values measured therefor may be added together for comparison. The measurement of the biomarker for a psychiatric/neurological disorder according to the present invention in a normal subject may be the same as the measurement method described in the step (a).

35 [0064] In most normal subjects, an inflammatory cytokine is below the limit of detection. Thus, the comparison of the amount of the inflammatory cytokine with that from a normal subject may be simply whether or not the inflammatory cytokine is detected. No detection of the inflammatory cytokine can be determined as indicating no change in the cytokine amount as compared with that from a normal subject.

40 [0065] Using the test method of the present invention, whether or not a subject has developed or can potentially develop a psychiatric/neurological disorder can be determined as follows. When the amount of a free κ immunoglobulin chain or a fragment thereof in a blood sample from a subject increases compared to that from a normal subject or is larger than the median value of that from a normal control group, the subject can be determined as having developed or potentially developing a psychiatric/neurological disorder. Alternatively, when the amount of a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject increases compared to that from a normal subject, the subject can be determined as having developed or potentially developing a psychiatric/neurological disorder. Alternatively, when the amount of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject increases compared to that from a normal subject or is larger than the median value of that from a normal control group, the subject can be determined as having developed or potentially developing a psychiatric/neurological disorder.

45 [0066] Using the test method of the present invention, whether or not a subject has developed or can potentially

develop a psychiatric/neurological disorder can also be determined as follows. When the amount of a free κ immunoglobulin chain or a fragment thereof in a blood sample from a subject increases compared to that from a normal subject or is larger than the median value of that from a normal control group, and the amount of an inflammatory cytokine does not change compared to that from a normal subject or is not detected, the subject can be determined as having developed or potentially developing a psychiatric/neurological disorder. Alternatively, when the amount of a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject increases compared to that from a normal subject or is larger than the median value of that from a normal control group, and the amount of an inflammatory cytokine does not change or is not detected compared to that from a normal subject, the subject can be determined as having developed or potentially developing a psychiatric/neurological disorder. Alternatively, when the amount of a free κ immunoglobulin chain or a fragment thereof and a free λ immunoglobulin chain or a fragment thereof in a blood sample from a subject increases compared to that from a normal subject and the amount of an inflammatory cytokine does not change or is not detected compared to that from a normal subject, the subject can be determined as having developed or potentially developing a psychiatric/neurological disorder.

[0067] "Increase" can be increasing about 1.5-fold, about 2-fold, about 3-fold, about 4-fold, about 5-fold, about 10-fold, about 15-fold, about 20-fold, about 30-fold, about 50-fold, about 100-fold, or more. "Not change" refers to that the amount of the biomarker for a psychiatric/neurological disorder according to the present invention is the same or is not significantly different between the amount from a subject in the step (a) and that from a normal subject in the step (b). "Not significantly different" refers to, for example, that the p value is less than 0.05, or the difference is about 0.1%, about 0.2%, about 0.3%, about 0.4%, about 0.5%, about 1%, about 2%, about 3%, about 4%, about 5%, about 10%, about 20%, or about 30%.

[0068] "Larger than the median value" means, for example, larger than the median value in the normal control group. The median value in the normal control group may be a known median value as described, for example, in Non-patent Literature 1, or a value calculated from the actual measurement in the normal control group.

Examples

(5. Examples)

[0069] Examples of the present invention will be described below. The following Examples are described for a better understanding of the scope of claims of the present invention, and not intended to limit the scope of claims of the present invention.

Example 1

[0070] Poly IC (polyriboinosinic-polyribocytidilic acid) (Sigma Co., Ltd.) dissolved in saline was intraperitoneally administered in an amount of 4 mg/kg to a Wistar rat at day 15 of gestation before normal delivery to prepare maternal immune activation model rats (MIA rats) which were schizophrenic and autistic model animals. The MIA rat is one of psychiatric/neurological disorder model animals (see Limin Shi et al, J. Neurosci. 2003 Jan 1; 23(1): 297-302 and Arata Oh-Nishi et al, Brain Research, Volume 1363, 6 December 2010, Pages 170-179). Only an equal volume of saline was intraperitoneally administered to another Wistar rat at day 15 of gestation before normal delivery to prepare control rats (hereinafter, referred to as "control" rats) as a control group. The MIA rats had significantly attenuated PPI compared to the control rats and were confirmed to be capable of being used as psychiatric/neurological disorder model animals.

[0071] The sera of the MIA rats (12-week old) and the control rats (12-week old) were collected and denatured with urea/thiourea, DTT, and SDS, followed by separating protein by a 2-dimensional electrophoresis method. Electrophoresis gel images are shown in Figure 2. As a result of analyzing the images using an image analysis software (Progenesis PG220; from Shimadzu Corporation), spot 1115 (about 25 kDa) was found to be significantly increased in the MIA rats (described as "Poly IC" in the figure) compared to that in the control rats (described as "Saline" in the figure). The spot was cut out, and the protein was identified by a LC MS/MS measurement method; as a result, it was found to be a κ immunoglobulin chain C region (Ig κ -C) (Figure 3).

[0072] The free κ immunoglobulin chain consists of a constant region (κ immunoglobulin chain C region) of about 12 kDa and a variable region of about 12 kDa. The variable region frequently changes in the amino acid sequence and thus is not detected by a mass spectrometer (MS). The spot in which the significant difference was seen was observed around about 25 kDa and thus it may be actually a free κ immunoglobulin chain.

Example 2

[0073] A rat-derived κ immunoglobulin chain C region (SEQ ID NO: 4) was used as an antigen and administered to a rabbit for immunization to prepare an antibody (anti-Ig κ -C antibody). Using the same method as that in Example 1,

MIA rats and control rats were prepared. The sera of the MIA rats (12-week old, N=7) and the control rats (12-week old, N=7) were denatured only with SDS (100°C, 5 minutes) and subjected to electrophoresis with SDS-polyacrylamide gel to separate protein. Thereafter, Western blotting was carried out (gel: 10 to 20%, primary antibody (1:1,000), secondary antibody (1:5,000)) (Figure 4). The anti-Ig κ-C antibody was used as a primary antibody, and an anti-rabbit IgG-HRP (from Chemi-Con Co., Ltd.) was used as a secondary antibody.

[0074] The intensity of a band of about 25 kDa significantly increased in the MIA rats ("Poly IC" in Figure 4) compared to that in the control rats ("Saline" in Figure 4) (Figure 4(2)). The band of about 25 kDa is the free κ immunoglobulin chain since it is detected without the cleavage of S-S bonds or hydrogen bonds. These results clearly show that the free κ immunoglobulin chain serves as a biomarker for a psychiatric/neurological disorder.

Example 3

[0075] The amounts of inflammatory cytokines in the serum were measured in the MIA rats ("Poly IC" in the table, 12-week old, N=7) and the control rats ("Saline" in the table, 12-week old, N=7) of Example 2. The measurement of the amount of inflammatory cytokines used Quantikine ELASA Rat IL-1β, IL-6, and TNF-α (from R&D Co., Ltd.).

[Table 1]

Table 1. Concentration of Inflammatory Cytokine in Serum

	IL-6 (μg/ml)	IL-1 beta (pg/ml)	TNF-alpha (pg/ml)
Saline	N.D	N.D	N.D
Poly IC	N.D	N.D	N.D
	(N.D<250)	(N.D<96)	(N.D<25)

[0076] No increase of the inflammatory cytokines in sera was observed in the MIA rats ("Poly IC" in the table) and the control rats ("Saline" in the table).

Example 4

[0077] The usefulness of the biomarker for a psychiatric/neurological disorder according to the present invention was tested in a group of schizophrenia patients (SZ) and a group of healthy individuals (Normal).

[0078] Serum samples from the SZ and Normal groups, collected according to an IRB-approved protocol, which was given sufficient ethical consideration, were purchased from PrecisionMed Inc. (USA) (acting import agent: Wako Pure Chemical Industries, Ltd.).

[0079] The amounts of the free κ immunoglobulin chain and the free λ immunoglobulin chain in the sera were measured using FREELITE κ Chain (MBL Co., Ltd., code number: BS-LK016BD) and FREELITE λ chain (MBL Co., Ltd., code number: BS-LK018BD) according to the instructions of the provider. The inflammatory cytokines were measured using AlphaLASA Human IL-1 beta, IL-6, and TNF-alpha Kits (from PerkinElmer Inc., code numbers: AL223 C/F, AL220 C/F, and AL208 C/F).

[0080] Basic data and the data summarizing average values in the group of schizophrenia patients (SZ) and the normal group are shown in Tables 2 and 3 below (psychopathological score: PANSS score).

[Table 2]

Table 2. Basic Data in Schizophrenia Patient(SZ) and Normal Groups

ID(#)	Race	Age	Age of Onset	Positive Symptom	Negative Symptom	General Psychopathology	IL-6(pg/ml)	IL-1 beta (pg/ml)	TNF-alpha (pg/ml)
10226006	Caucasian	36	21	19	16	34	ND	ND	ND
10226007	Caucasian	35	16	18	16	36	ND	ND	ND
10226029	Caucasian	36	16	17	22	45	ND	ND	ND
10226046	Caucasian	51	20	20	19	41	ND	ND	ND
10226066	Caucasian	61	26	8	15	25	ND	ND	ND
10226067	Caucasian	41	27	24	20	44	ND	ND	ND
10226016	Caucasian	45	45	26	24	53	ND	ND	ND
10226047	Caucasian	58	23	17	14	33	ND	ND	ND
10226063	Caucasian	41	20	23	12	44	ND	ND	ND
10226084	Caucasian	36	17	15	12	27	ND	ND	ND
10226089	Caucasian	38	18	18	19	39	ND	ND	ND
Normal									
7516	Caucasian	25	-	-	-	-	ND	ND	ND
7520	Caucasian	55	-	-	-	-	ND	ND	ND
7538	Caucasian	44	-	-	-	-	ND	ND	ND
7543	Caucasian	62	-	-	-	-	ND	ND	ND
7551	Caucasian	49	-	-	-	-	ND	ND	ND
7522	Hispanic	39	-	-	-	-	ND (ND<10)	ND (ND<3)	ND
7552	Hispanic	40	-	-	-	-	ND (ND<10)	ND (ND<3)	ND (ND<10)

[Table 3]

Table 3. Summary of Basic Data

	Normal (N=7)	SZ (N=11)
Age	44.8±4.8	43.45±2.79
Age of Onset		22.6±2.7
Positive Symptom		18.6±1.4
Negative Symptom		17.1±1.1
General Psychopathological Score		38.2±2.5
Average ±S.E.M		

[0081] There was no statistically significant difference in age, and the inflammatory cytokines (IL-6, IL-1 beta, and TNF-alpha) were not detected in both of the SZ and normal groups.

[0082] Figure 5 and Table 4 show the results of comparing the concentrations of the free κ immunoglobulin chain (A in Figure 5) and the free λ immunoglobulin chain (B in Figure 5) in the sera of the group of schizophrenia patients (SZ) and the normal group. In the schizophrenia patients (SZ), the concentrations of the free κ immunoglobulin chain and the free λ immunoglobulin chain significantly increased ($P = 0.0012$ and $P = 0.04$, respectively). These results clearly show that each the free κ immunoglobulin chain and the free λ immunoglobulin chain individually serves as a biomarker for a psychiatric/neurological disorder. As a result of comparing the concentration of the free κ immunoglobulin chain + the free λ immunoglobulin chain in the group of schizophrenia patients (SZ) and the normal group (C in Figure 5), a further significant difference was observed ($P = 0.00063$). These results show that it is particularly useful for determining a psychiatric/neurological disorder to combine the free κ immunoglobulin chain and the free λ immunoglobulin chain.

[Table 4]

Table 4. Concentration of Free Ig κ and λ in Serum of Schizophrenia Patient

	Normal	SZ
Free Ig κ (mg/l)	7.89±0.46 (N=6)	14.09±1.41** (N=11)
Free Ig λ (mg/l)	10.89±0.94 (N=7)	14.04±0.99* (N=11)
Free Ig κ + Free Ig λ (mg/l)	17.79±1.16 (N=6)	28.14±2.07*** (N=11)
(Average ± S.E.M)		

[0083] In addition, the concentration of free Ig κ in the sera from the 11 patients of the SZ group had higher values (minimum: 9.11 mg/l, maximum: 23.8 mg/l) compared with the median value (7.5 mg/l) of the concentration of free Igh in the sera from healthy individuals aged 40 to 49 years described in Non-patent Literature 1. On the other hand, the median value of the concentration of free Ig λ in the sera from the healthy individuals aged 40 to 49 years is 12.8 mg/l, and 8 of 11 patients in the SZ group had a higher value than the median value (minimum: 9.05 mg/l, maximum: 20.8 mg/l).

Industrial Applicability

[0084] The biomarker of the present invention in a blood sample can be measured to perform the primary screening of a psychiatric/neurological disorder in a medical checkup and the like. The screening of a psychiatric/neurological disorder can also be carried out in the first contact of patients in psychiatry.

EP 3 002 588 A1

SEQUENCE LISTING

<110> National Institute of Radiological Sciences
 5 <120> Biomarker for Psycho-neurologic Disease
 <130> 0749005W01
 <160> 6
 10 <170> PatentIn version 3.5
 <210> 1
 <211> 106
 <212> PRT
 15 <213> Homo sapiens
 <400> 1
 Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 1 5 10 15
 20 Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 20 25 30
 25 Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 35 40 45
 30 Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 50 55 60
 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 65 70 75 80
 35 His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 85 90 95
 40 Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 100 105
 <210> 2
 <211> 20
 45 <212> PRT
 <213> Homo sapiens
 <400> 2
 50 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
 1 5 10 15
 Thr Leu Thr Leu
 20
 55 <210> 3

EP 3 002 588 A1

<211> 21
 <212> PRT
 <213> Homo sapiens

5 <400> 3

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
 1 5 10 15

10 Asn Arg Gly Glu Cys
 20

15 <210> 4
 <211> 106
 <212> PRT
 <213> Rattus norvegicus

<400> 4

20 Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser Met Glu Gln
 1 5 10 15

25 Leu Thr Ser Gly Gly Ala Thr Val Val Cys Phe Val Asn Asn Phe Tyr
 20 25 30

Pro Arg Asp Ile Ser Val Lys Trp Lys Ile Asp Gly Ser Glu Gln Arg
 35 40 45

30 Asp Gly Val Leu Asp Ser Val Thr Asp Gln Asp Ser Lys Asp Ser Thr
 50 55 60

35 Tyr Ser Met Ser Ser Thr Leu Ser Leu Thr Lys Val Glu Tyr Glu Arg
 65 70 75 80

His Asn Leu Tyr Thr Cys Glu Val Val His Lys Thr Ser Ser Ser Pro
 85 90 95

40

Val Val Lys Ser Phe Asn Arg Asn Glu Cys
 100 105

45 <210> 5
 <211> 20
 <212> PRT
 <213> Rattus norvegicus

50 <400> 5

Ser Val Thr Asp Gln Asp Ser Lys Asp Ser Thr Tyr Ser Met Ser Ser
 1 5 10 15

55 Thr Leu Ser Leu
 20

<210> 6
 <211> 94
 <212> PRT
 <213> Homo sapiens

<400> 6

Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln Ala
 1 5 10 15

Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly Ala
 20 25 30

Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val Lys Ala Gly Val
 35 40 45

Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala Ser
 50 55 60

Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Lys Ser Tyr
 65 70 75 80

Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr
 85 90

Claims

1. A biomarker for a psychiatric/neurological disorder, comprising at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof.
2. A biomarker for a psychiatric/neurological disorder, comprising at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof, wherein the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 2 or 3 and the free λ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 6.
3. A biomarker for a psychiatric/neurological disorder, comprising at least one selected from the group consisting of a free κ immunoglobulin chain, a free λ immunoglobulin chain, and fragments thereof, wherein the free κ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 1 and the free λ immunoglobulin chain comprises the amino acid sequence of SEQ ID NO: 6.
4. The biomarker for a psychiatric/neurological disorder according to any one of claims 1 to 3, further comprising an inflammatory cytokine.
5. The biomarker for a psychiatric/neurological disorder according to claim 4, wherein the inflammatory cytokine is at least one selected from the group consisting of IL-6, IL-1 beta, and TNF-alpha.
6. The biomarker for a psychiatric/neurological disorder according to any one of claims 1 to 5, wherein the psychiatric/neurological disorder is schizophrenia, autism, manic-depressive psychosis, cognitive impairment ascribed to neuropathy due to infection during pregnancy, or Alzheimer's disease.
7. The biomarker for a psychiatric/neurological disorder according to any one of claims 1 to 5, wherein the psychiatric/neurological disorder is schizophrenia.

EP 3 002 588 A1

8. A test kit for a psychiatric/neurological disorder, comprising a substance capable of detecting the biomarker for a psychiatric/neurological disorder according to any one of claims 1 to 7.

5

9. The test kit according to claim 8, wherein the substance is an antibody.

10. The test kit according to claim 8 or 9, wherein the substance is immobilized on a solid-phase support.

10

11. A test method for a psychiatric/neurological disorder, using the amount of the biomarker for a psychiatric/neurological disorder according to any one of claims 1 to 7 in a blood sample from a subject as an indication.

12. A test method for a psychiatric/neurological disorder, comprising the steps of:

(a) measuring the amount of the biomarker for a psychiatric/neurological disorder according to any one of claims 1 to 7 in a blood sample from a subject; and

15

(b) comparing the amount with the amount of that from a normal subject.

13. The test method according to any one of claim 11 or 12, wherein the blood sample is serum.

20

25

30

35


40

45


50

55


[Fig. 1]




[Fig. 2]




[Fig. 3]



[Fig. 4]



[Fig.5]



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2014/057227

A. CLASSIFICATION OF SUBJECT MATTER

G01N33/53(2006.01)i, G01N33/543(2006.01)i, C07K16/00(2006.01)n, C12Q1/02(2006.01)n

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

G01N33/53, G01N33/543, C07K16/00, C12Q1/02

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Jitsuyo Shinan Koho 1922-1996 Jitsuyo Shinan Toroku Koho 1996-2014
Kokai Jitsuyo Shinan Koho 1971-2014 Toroku Jitsuyo Shinan Koho 1994-2014

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAplus/MEDLINE/EMBASE/BIOSIS (STN), JSTPlus/JMEDPlus/JST7580 (JDreamIII)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	O.C. Fagnart et al, Free kappa and lambda light	1-3, 8-13
Y	chain levels in the cerebrospinal fluid of	4-5
A	patients with multiple sclerosis and other	6-7
	neurological diseases, Journal of	
	Neuroimmunology, 1988.08, Volume 19, Issues	
	1-2, 119-132	
Y	WO 2012/078623 A2 (RIDGE DIAGNOSTICS, INC.),	4-5
	14 June 2012 (14.06.2012),	
	page 11, line 27 to page 12, line 14	
	& US 2012/0178118 A1 & CA 2820616 A	
A	I.V.L. Narasimha Rao et al, SERUM	1-13
	IMMUNOGLOBULINS AND SCHIZOPHRENIA, Indian	
	J Psychiatry, 1985, 27(4), 325-328	

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
11 June, 2014 (11.06.14)Date of mailing of the international search report
24 June, 2014 (24.06.14)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2014/057227

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Popovic M et al, Light chain deposition disease restricted to the brain: The first case report, Hum Pathol, 2007.01, 38(1), 179-184	1-13
A	WO 2011/144934 A1 (CAMBRIDGE ENTERPRISE LTD.), 24 November 2011 (24.11.2011), claims & US 2013/0178385 A1 & GB 201008340 D & EP 2572195 A & CA 2799663 A	1-13
A	US 2012/0238936 A1 (Roderick A. Hyde), 20 September 2012 (20.09.2012), paragraph [0077] & US 2012/0238837 A1 & US 2012/0238836 A1 & US 2012/0238839 A1 & US 2012/0238835 A1 & US 2012/0238838 A1 & WO 2012/125728 A2	1-13

Form PCT/ISA/210 (continuation of second sheet) (July 2009)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 2012013415 A [0004]
- JP 2011506995 W [0004]

Non-patent literature cited in the description

- **KATZMANN JA ; CLARK RJ ; ABRAHAM RS ; BRYANT S ; LYMP JF ; BRADWELL AR ; KYLE RA.** *Clin Chem.*, September 2002, vol. 48 (9), 1437-44 [0005]
- **TSIRAKIS G ; PAPPA CA ; KAPAROU M ; KATSOMITROU V ; HATZIVASILI A ; ALEGAKIS T ; XEKALOU A ; STATHOPOULOS EN ; ALEXANDRAKIS MG.** *Eur J Histochem.*, 2011, vol. 55 (3), e21 [0005]
- **KUKU I ; BAYRAKTAR MR ; KAYA E ; ERKURT MA ; BAYRAKTAR N ; CIKIM K ; AYDOGDU I.** *Mediators Inflamm.*, 14 August 2005, vol. 2005 (3), 171-4 [0005]
- **LIMIN SHI et al.** *J. Neurosci.*, 01 January 2003, vol. 23 (1), 297-302 [0070]
- **ARATA OH-NISHI et al.** *Brain Research*, 06 December 2010, vol. 1363, 170-179 [0070]

专利名称(译)	用于精神和神经疾病的生物标志物		
公开(公告)号	EP3002588A1	公开(公告)日	2016-04-06
申请号	EP2014797879	申请日	2014-03-18
[标]申请(专利权)人(译)	RESVO INC		
申请(专利权)人(译)	resvo Inc.		
当前申请(专利权)人(译)	石川彻		
[标]发明人	OHNISHI ARATA MINAMIMOTO TAKAFUMI SUHARA TETSUYA		
发明人	OHNISHI, ARATA MINAMIMOTO, TAKAFUMI SUHARA, TETSUYA		
IPC分类号	G01N33/53 G01N33/543 C07K16/00 C12Q1/02 C07K16/42 G01N33/68		
CPC分类号	G01N33/6857 C07K16/42 G01N33/6863 G01N33/6869 G01N33/6893 G01N33/6896 G01N2333/525 G01N2333/5412 G01N2333/545 G01N2800/2821 G01N2800/301 G01N2800/302 G01N2800/304		
代理机构(译)	MAIKOWSKI & Ninnemann		
优先权	2013104124 2013-05-16 JP		
其他公开文献	EP3002588A4		
外部链接	Espacenet		

摘要(译)

本发明的目的是提供：用于精神病和神经病症的生物标志物，特别是用于诊断精神病和神经病症的生物标志物；以及用于精神和神经疾病的测试试剂盒和测试方法。发明人发现，来自患有精神病和神经病症的患者的血液样品中游离κ免疫球蛋白链和游离λ免疫球蛋白链的浓度显著增加。因此，本发明提供用于精神病和神经病症的生物标志物，其包括选自游离κ免疫球蛋白链，游离λ免疫球蛋白链及其片段中的至少一种。使用该生物标志物用于精神病学和神经学病症使得能够容易地测试和诊断来自受试者的血液样品的精神病学和神经学病症。生物标志物使得能够在患有精神病和神经病症的患者的血液中结合炎症细胞因子进行更有效的测试和诊断。

[Fig.5]

