



(11) **EP 2 021 799 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
20.07.2011 Bulletin 2011/29

(51) Int Cl.:
G01N 33/53 (2006.01) C07K 14/00 (2006.01)
C07K 16/00 (2006.01) G01N 33/68 (2006.01)

(21) Application number: **07811859.3**

(86) International application number:
PCT/US2007/067626

(22) Date of filing: **27.04.2007**

(87) International publication number:
WO 2007/143295 (13.12.2007 Gazette 2007/50)

(54) **INTERLEUKIN-33 (IL-33) FOR THE DIAGNOSIS AND PROGNOSIS OF CARDIOVASCULAR DISEASE**

INTERLEUKIN-33 (IL-33) ZUR DIAGNOSE UND VORHERSAGE VON HERZ-GEFÄSS-ERKRANKUNGEN

INTERLEUKINE-33 (IL-33) UTILISÉE POUR LE DIAGNOSTIC ET LE PRONOSTIC DES MALADIES CARDIOVASCULAIRES

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR

(56) References cited:
WO-A-2005/079844 WO-A-2007/127749
WO-A-2007/130627 WO-A-2007/131031
US-A1- 2008 003 199

(30) Priority: **27.04.2006 US 795473 P**

(43) Date of publication of application:
11.02.2009 Bulletin 2009/07

(73) Proprietor: **Critical Care Diagnostics, Inc.**
New York, New York 10019 (US)

(72) Inventors:
• **SNIDER, James**
Pleasanton, CA 94588 (US)
• **JACOBSON, Sven**
New York, New York 10001 (US)

(74) Representative: **Fleck, Barbara**
Marks & Clerk LLP
62-68 Hills Road
Cambridge
CB2 1LA (GB)

- **SHIMPO MASAHISA ET AL: "Serum levels of the interleukin-1 receptor family member ST2 predict mortality and clinical outcome in acute myocardial infarction" CIRCULATION, vol. 109, no. 18, 11 May 2004 (2004-05-11), pages 2186-2190, XP002523097**
- **SCHMITZ J ET AL: "IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines" IMMUNITY, CELL PRESS, US, vol. 23, no. 5, 1 November 2005 (2005-11-01), pages 479-490, XP002432892 ISSN: 1074-7613**
- **KAKKAR R ET AL: "The IL-33/ST2 pathway: Therapeutic target and novel biomarker" NATURE REVIEWS DRUG DISCOVERY 2008 GB, vol. 7, no. 10, 2008, pages 827-840, XP002530680 ISSN: 1474-1776 1474-1784**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 021 799 B1

Description**TECHNICAL FIELD**

5 [0001] This invention pertains to *in vitro* methods and compositions for the diagnosis and prognosis of cardiovascular conditions.

BACKGROUND

10 [0002] Despite significant advances in therapy, cardiovascular disease remains the single most common cause of morbidity and mortality in the developed world. Thus, prevention and therapy of cardiovascular conditions such as heart failure, myocardial infarction and stroke is an area of major public health importance. Currently, several risk factors for future cardiovascular disorders have been described and are in wide clinical use in the detection of subjects at high risk. Such screening tests include evaluations of total and HDL cholesterol levels. However, a large number of cardiovascular disorders occur in subjects with apparently low to moderate risk profiles, and ability to identify such patients is limited. Moreover, accumulating data suggests that the beneficial effects of certain preventative and therapeutic treatments for patients at risk for or known to have cardiovascular disorders differs in magnitude among different patient groups.

15 [0003] Shimpo et al ("Serum levels of interleukin-1 receptor family member ST2 predict mortality in clinical outcome in acute myocardial infarction", *Circulation*, Vol. 109, No. 18, 11 May 2004, pages 2186-2190) relates to a study designed to test the hypothesis that ST2 serum levels are associated with risk of death and heart failure in patients presenting with ST elevation myocardial infarction, but is not concerned with diagnosis. Schmitz et al ("IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines", *Immunity*, Vol. 23, No. 5, 1 November 2005, pages 479-490) reports the induction of IL-33 expression in primary lung or dermal fibroblasts and keratinocytes upon TNF- α and IL-1- β activation. Induction in such cell types correlates with higher endogenous levels of IL-33 in lung and skin.

SUMMARY

20 [0004] The present invention includes *in vitro* methods for the use of interleukin-33 (IL-33) in the diagnosis and prognosis of cardiovascular conditions including acute coronary syndrome (ACS), coronary artery disease (CAD), myocardial infarction, heart failure, angina, cardiac hypertrophy, arteriosclerosis, myocarditis, pericarditis, endocarditis, stroke and/or pulmonary embolism.

25 [0005] In one aspect, the invention provides *in vitro* methods for diagnosing a cardiac disease in a subject. The methods include determining a level of IL-33 in a sample from a subject, wherein the level of IL-33 in the sample is indicative of whether the subject has a cardiovascular disease (i.e., correlates to the presence or absence of cardiovascular disease). The methods include determining a level of IL-33 in the sample; and comparing the level of IL-33 in the sample to a reference level of IL-33. The level of IL-33 in the sample as compared to the reference indicates whether the subject has a cardiovascular disease.

30 [0006] In another aspect, the invention provides *in vitro* methods for determining the severity of a cardiovascular disease in a subject. The methods include determining a level of IL-33 in a sample from a subject, wherein the level of IL-33 in the sample is indicative of (correlates to) the severity of the cardiovascular disease in the subject. The methods include determining a level of IL-33 in the sample; and comparing the level of IL-33 in the sample to a reference level of IL-33. The level of IL-33 in the sample as compared to the reference indicates the severity of the cardiovascular disease in the subject.

35 [0007] In some embodiments of the methods described herein, the sample comprises blood, serum or plasma. In some embodiments, the sample comprises serum.

[0008] The reference level represents a level in a subject who does not have cardiovascular disease. In some methods, the reference level represents a level in a subject with a preselected severity of cardiovascular disease.

40 [0009] In some embodiments of the methods described herein, determining a level of IL-33 in the sample includes determining a level of one, two, or all three of mature IL-33, pre-IL-33, and pro-IL-33. Determining a level of IL-33 in the sample can include contacting a binding composition to the sample, wherein the binding composition specifically binds to IL-33, and measuring or determining the specific binding of the binding composition to the sample. Suitable binding compositions include antibodies that bind specifically to IL-33 polypeptide, and oligonucleotide probes that bind specifically to IL-33 polynucleotide.

45 [0010] In some embodiments, the cardiac disease diagnosed by a method described herein is acute coronary syndrome (ACS), myocardial infarction, heart failure, angina, cardiac hypertrophy, arteriosclerosis, myocarditis, pericarditis, endocarditis, stroke, and/or pulmonary embolism.

50 [0011] In some embodiments, the methods described herein also include determining a level in the sample of one or

more other biomarkers, e.g., biomarkers selected from the group consisting of ST2, NT-proBNP, BNP, NT-proANP, and ANP, troponin, IMA, IL-6, CRP, creatinine, D-dimers, BUN, liver function enzymes, albumin, and bacterial endotoxin.

[0012] We also describe herein kits for diagnosing cardiovascular disease. The kits include an antibody that specifically binds to IL-33, and/or an oligonucleotide probe that specifically binds to a nucleic acid encoding IL-33, and instructions for use in a method described herein.

[0013] "Upregulated," as used herein, refers to increased expression of a gene and/or its encoded polypeptide. "Increased expression" refers to increasing (i.e., to a detectable extent) replication, transcription, and/or translation of IL-33, since upregulation of any of these processes results in an increase in concentration/amount of the polypeptide encoded by the gene. Conversely, "downregulation," or "decreased expression" as used herein, refers to reduced replication, transcription, and/or translation of the IL-33 gene and/or its encoded polypeptide. The upregulation or downregulation of gene expression can be directly determined by detecting an increase or decrease, respectively, in the level of mRNA for the gene, or the level of protein expression of the gene-encoded polypeptide, using any suitable means known to the art, such as nucleic acid hybridization or antibody detection methods, respectively, and in comparison to controls. "Expression," as used herein, refers to nucleic acid and/or polypeptide expression.

[0014] A "cardiac cell," as used herein, refers to a cardiomyocyte.

[0015] As used herein, a "subject" is a mammal or a non-human mammal. In general, human nucleic acids, polypeptides, and human subjects are preferred for use in diagnosing patients.

[0016] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0017] Other features and advantages of the invention will be apparent from the following detailed description and figures, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018]

FIG. 1 is a representation of the nucleic acid sequence of human IL-33 (SEQ ID NO:1).

FIG. 2 is a representation of the amino acid sequence of human IL-33 (SEQ ID NO:2).

DETAILED DESCRIPTION

[0019] It has previously been described that levels of the interleukin-1 receptor-like 1 (IL1RL1) protein can be used to diagnose cardiovascular disease and determine the prognosis for a patient with cardiovascular disease. The ligand for IL1RL1 has been described, and named IL-33 (see, e.g., Schmitz et al., *Immunity* 23(5):479-90 (2005); U.S. Pat. Pub. No. 2005/0203046). The present methods include the measurement of levels of IL-33 for the diagnosis and prognosis of cardiovascular disease.

General Methodology

[0020] In general, the methods described herein include evaluating levels of IL-33 in a biological sample (e.g., a blood, serum, plasma, urine, or body tissue sample) from a subject, e.g., a mammal, e.g., a human. These levels provide diagnostic information indicating whether the subject has a cardiac disease, as described herein. In some embodiments, the level of IL-33 is determined once, e.g., at presentation. In some embodiment, the level of IL-33 is determined at any one or more of 1, 2, 3, 4, 5, 6, 7, 8, 12, 18, and/or 24 hours, and/or at 1-7 days after the onset of symptoms.

[0021] In embodiments where the level of IL-33 is determined more than once, the higher level can be used, or the change in levels can be determined and used. Levels of IL-33 can be determined multiple times to evaluate a subject's response to a treatment. For example, a level of IL-33 taken after administration of a treatment, e.g., one or more doses or rounds of a treatment, can be compared to levels of IL-33 before the treatment was initiated, e.g., a baseline level. The change in IL-33 levels would indicate whether the treatment was effective; e.g., a reduction in IL-33 levels would indicate that the treatment was effective.

[0022] In some embodiments, the level of IL-33 can be measured in patients previously diagnosed with a cardiovascular condition who are undergoing treatment on an outpatient basis, to monitor the condition of the patient and determine whether their condition is improving, stable or deteriorating. Levels of IL-33 can be measured at one or more times and compared to previous measurements, where an increase in IL-33 relative to an earlier measurement would indicate that the patient's condition is deteriorating, suggesting that the current treatment should be adjusted or possibility that the patient should be admitted to a hospital for inpatient treatment.

[0023] Evaluating circulating levels of IL-33 in a subject typically includes obtaining a biological sample, e.g., serum or blood, from the subject. Levels of IL-33 in the sample can be determined by measuring levels of polypeptide, either the pre-IL33, proIL-33 or IL-33 peptide, in the sample, using methods known in the art and/or described herein, e.g., immunoassays such as enzyme-linked immunosorbent assays (ELISA). Alternatively, levels of IL-33 mRNA can be measured, again using methods known in the art and/or described herein, e.g., by quantitative PCR or Northern blotting analysis.

[0024] For example, a method as described herein, e.g., for diagnosis or prognosis of cardiac disease, can include contacting a sample from a subject, e.g., a sample including blood, serum, plasma, urine, or body tissue from the subject, with a binding composition (e.g., an antibody or oligonucleotide probe) that specifically binds to a polypeptide or nucleic acid of IL-33. The methods can also include contacting a sample from a control subject, normal subject, or normal tissue or fluid from the test subject, with the binding composition, e.g., to provide a reference or control. Moreover the method can additionally include comparing the specific binding of the composition to the test subject with the specific binding of the composition to the normal subject, control subject, or normal tissue or fluid from the test subject. Expression or activity of IL-33 in a test sample or test subject can also be compared with that in a control sample or control subject. A control sample can include, e.g., a sample from a non-affected subject, or it subject who has a cardiac disease of known severity. Expression or activity from a control subject or control sample can be provided as a predetermined value, e.g., acquired from a statistically appropriate group of control subjects.

[0025] An antibody that "binds specifically to" an antigen, binds preferentially to the antigen in a sample containing other proteins. The term "antibody" as used herein refers to an immunoglobulin molecule or immunologically active portion thereof, i.e., an antigen-binding portion. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab')₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The antibody can be polyclonal, monoclonal, recombinant, e.g., a chimeric or humanized, fully human, non-human, e.g., murine, monospecific, or single chain antibody. In some embodiments it has effector function and can fix complement.

[0026] An "oligonucleotide probe" (also referred to simply as a "probe") is a nucleic acid that is at least 10, and less than 200 (typically less than about 100 or 50) base pairs in length. A probe that "binds specifically to" a target nucleic acid hybridizes to the target under high stringency conditions. As used herein, the term "hybridizes under high stringency conditions" describes conditions for hybridization and washing. As used herein, high stringency conditions are 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C. Methods for performing nucleic acid hybridization assays are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6.

[0027] Detection can be facilitated by coupling (e.g., physically linking) the antibody or probe to a detectable substance (e.g., antibody labeling). Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride, quantum dots, or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ¹²⁵I, ¹³¹I, ³⁵S or ³H.

[0028] Diagnostic assays can be used with biological matrices such as live cells, cell extracts, cell lysates, fixed cells, cell cultures, bodily fluids, or forensic samples. Conjugated antibodies useful for diagnostic or kit purposes, include antibodies coupled to dyes, isotopes, enzymes, and metals, see, e.g., Le Doussal et al., *New Engl. J. Med.* 146:169-175 (1991); Gibellini et al., *J. Immunol.* 160:3891-3898 (1998); Hsing and Bishop, *New Engl. J. Med.* 162:2804-2811 (1999); and Everts et al., *New Engl. J. Med.* 168:883-889 (2002). Various assay formats exist, such as radioimmunoassays (RIA), ELISA, and lab on a chip (U.S. Pat. Nos. 6,176,962 and 6,517,234).

[0029] Known techniques in biochemistry and molecular biology can be used in the methods described herein (see, e.g., Maniatis et al., *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1982); Sambrook and Russell, *Molecular Cloning*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001); Wu, *Recombinant DNA*, Vol. 217, Academic Press, San Diego, Calif (1993); and Ausbel et al., *Current Protocols in Molecular Biology*, Vols. 1-4, John Wiley and Sons, Inc. New York, N.Y. (2001)).

[0030] Once a level of IL-33 has been determined, the level is compared to a reference level. In some embodiments, e.g., where the level of IL-33 is determined using an ELISA, the reference level will represent a threshold level, above which the subject can be diagnosed with cardiac disease. The reference level chosen may depend on the methodology used to measure the levels of IL-33.

[0031] In some embodiments, the level of IL-33 is used to determine the level of severity of cardiac disease in a subject. The level of IL-33 typically increases with the severity of the cardiac disease, therefore, higher levels indicate more severe disease. Thus, the reference levels can represent levels of IL-33 in subjects with cardiac disease of known severity.

[0032] In some embodiments, the reference level is a range of levels.

[0033] In some embodiments, both levels of Interleukin 1 Receptor-Like 1 (IL1RL1) and IL-33 are determined, and the information from the comparison of both biomarkers with their respective reference levels provides cumulative information regarding the presence of cardiac disease in the subject, and/or the presence of a severe disease in the subject. In some embodiments, the ratio of IL1RL1 to IL-33 may be determined, and the ratio compared to a reference ratio that represents a threshold ratio above which the subject has cardiac disease.

[0034] Also described herein are kits that include a reagent for the detection of one or more of the IL-33 polypeptide (s) or nucleic acid, e.g., an anti-IL-33 antibody (i.e., an antibody that binds specifically to IL-33), or a nucleic acid probe complementary to all or part of the IL-33 nucleic acid, and instructions for use.

Interleukin-33 (IL-33)

[0035] IL-33 was recently identified as the ligand for IL1RL1, and the presence of increased levels of IL-33 in various inflammatory disorders has been described (see Schmitz et al., *Immunity* 23(5):479-90 (2005); U.S. Pat. Pub. No. 2005/0203046).

[0036] IL-33 protein is expressed as an inactive molecule, pre-IL-33, that is activated after cleavage by Caspase I resulting in the active IL-33 peptide as well as the cleavage peptide product, pro-IL-33. Therefore, the methods described herein can include measuring, one, two, or all three of mature IL-33, pro-IL-13, and/or pro-IL-33, all of which are included in the term "IL-33" as used herein.

[0037] The nucleic acid sequence of human IL-33 can be found at GenBank Acc. No. NM_033439.2 (Figure 1, SEQ ID NO:1), and the polypeptide sequence is at GenBank Acc. No. NP_254274 (Figure 2, SEQ ID NO:2). Additional information is available in the public databases at GeneID: 90865, MIM ID # *608678, and UniGene No. Hs.348390. IL-33 is also known as Chromosome 9 Open Reading Frame 26 (C9ORP26); Nuclear Factor from High Endothelial Venules (NFHEV); and Interleukin 33. See also Baekkevold et al., *Am. J. Path.* 163: 69-79 (2003).

[0038] Methods for measuring levels of IL-33 polypeptide and nucleic acid are known in the art, see, e.g., Schmitz et al., *Immunity* 23(5):479-90 (2005); U.S. Pat. Pub. No. 2005/0203046,

Interleukin 1 Receptor-like 1 (IL1RL1)

[0039] In the methods described herein, IL1 RL1 can be measured in addition to IL-33. The ratio of IL1 RL1 to IL-33 can also be determined. The IL1 RL1 gene is a member of the interleukin-1 receptor family, whose protein product exists both as a trans-membrane form, as well as a soluble receptor that is detectable in serum (Kieser et al., *FEBS Lett.* 372 (2-3): 189-93 (1995); Kumar et al., *J. Biol. Chem.* 270(46):27905-13 (1993); Yanagisawa et al., *FEBS Lett.* 302(1):51-3 (1992); Kuroiwa et al., *Hybridoma* 19(2):151-9 (2000)). ST2 was recently described to be markedly up-regulated in an experimental model of heart failure (Weinberg et al., *Circulation* 106(23):2961-6 (2002)), and preliminary results suggest that ST2 concentration may be elevated in those with chronic severe HF (Weinberg et al., *Circulation* 107(5):721-6 (2003)) as well as in those with acute myocardial infarction (MI) (Shimpo et al., *Circulation* 109(18):2186-90 (2004)).

[0040] The transmembrane form of IL1RL1 is thought to play a role in modulating responses of T helper type 2 cells (Lohning et al., *Proc. Natl. Acad. Sci. U. S. A.* 95(12):6930-5 (1998); Schmitz et al., *Immunity* 23(5):479-90 (2005)), and may play a role in development of tolerance in states of severe or chronic inflammation (Brint et al., *Nat. Immunol.* 5(4): 373-9 (2004)), while the soluble form of IL1RL1 is up-regulated in growth stimulated fibroblasts (Yanagisawa et al., 1992, supra). Experimental data suggest that the IL1RL1 gene is markedly up-regulated in states of myocyte stretch (Weinberg et al., 2002, supra) in a manner analogous to the induction of the BNP gene (Bruneau et al., *Cardiovasc. Res.* 28(10): 1519-25 (1994)).

[0041] Tominaga, *FEBS Lett.* 258:301-304 (1989), isolated murine genes that were specifically expressed by growth stimulation in BALB/c-3T3 cells; they termed one of these genes St2 (for Growth Stimulation-Expressed Gene 2). The St2 gene encodes two protein products: ST2, which is a soluble secreted form; and ST2L, a transmembrane receptor form that is very similar to the interleukin- 1 receptors. The HUGO Nomenclature Committee designated the human homolog, the cloning of which was described in Tominaga et al., *Biochim. Biophys. Acta* 1171:215-218 (1992), as Interleukin 1 Receptor-Like 1 (IL1RL1). The two terms are used interchangeably herein.

[0042] The mRNA sequence of the shorter, soluble isoform of human ST2 can be found at GenBank Acc. No. NM_003856.2, and the polypeptide sequence is at GenBank Acc. No. NP_003847.2; the mRNA sequence for the longer form of human ST2 is at GenBank Acc. No. NM_016232.4; the polypeptide sequence is at GenBank Acc. No. NP_057316.3. Additional information is available in the public databases at GeneID: 9173, MIM ID # 601203, and UniGene No. Hs.66. In general, in the methods described herein, the soluble form of ST2 polypeptide is measured.

[0043] Methods for detecting and measuring ST2 are known in the art, e.g., as described in U.S. Pat. Pub. Nos. 2003/0124624, 2004/0048286 and 2005/0130136. Kits for measuring ST2 polypeptide are also commercially available, e.g., the ST2 ELISA Kit manufactured by Medical & Biological Laboratories Co., Ltd. (MBL International Corp., Woburn,

MA), no. 7638. In addition, devices for measuring ST2 and other biomarkers are described in U.S. Pat. Pub. No. 2005/0250156.

[0044] In some embodiment, the methods include determining the identity of the nucleotide sequence at RefSNP ID: rs1041973.

5

Other Biomarkers

[0045] The methods described herein can also include measuring levels of other biomarkers in addition to IL-33. Suitable biomarkers include NT-proBNP, BNP, NT-proANP, and ANP troponin, GK-MB, Myo, IMA, IL-6, CRP, creatinine; D-dimers, and/or BUN,. Methods for measuring these biomarkers are known in the art, see, e.g., U.S. Pat. Pub. Nos. 2004/0048286 and 2005/0130136 to Lee et al.; Dhalla et al., Mol. Cell Biochem. 87:85-92 (1989); Moe et al., Am: Heart J. 139:587-95 (2000).

10

[0046] In these embodiments, levels of IL-33 and one or more additional biomarkers are determined, and the information from the comparison of the biomarkers with their respective reference levels provides additional information regarding the presence of cardiovascular disease in the subject, and/or the level of severity of the disease in the subject.

15

Cardiovascular Disease

[0047] The methods described herein are useful in the diagnosis and prognosis of subjects with cardiovascular disease, e.g., MI, ACS, CAD, HF and/or stroke. In subjects with cardiovascular disease, a diagnosis is often made, and the extent of any cardiac tissue damage determined, using one or more of the following methods: electrocardiogram (ECG) - single or repeated over several hours; echocardiography; coronary angiography; nuclear ventriculography (e.g., radionuclide ventriculography (RNV) or multiple gate acquisition scan (MUGA)).

20

[0048] In addition, some biomarkers have proven to be by-products of heart damage or in conjunction with coronary artery disease (CAD), and therefore-useful for diagnosis of cardiac disease. These biomarkers include Troponin I (TnI) and troponin T (TnT); creatine phosphokinase (CPK) and CPK-MB; ischemia modified albumin (TMA) and serum myoglobin. The present invention provides additional methods of diagnosing cardiovascular disease, and prognostic methods for determining the severity of disease in a subject. Thus, the methods described herein can include determining IL-33 levels as part of a diagnostic effort, e.g., to determine whether a subject has cardiovascular disease and/or can include determining IL-33 levels in a subject for whom a diagnosis of cardiovascular disease has already been made, e.g., to determine the severity of the disease.

25

30

[0049] Risk factors for cardiovascular disease include smoking, hypertension, high fat diet, poor blood cholesterol levels, especially high LDL ("bad") cholesterol and low HDL ("good") cholesterol, diabetes, male gender, age, heredity, and being overweight/obese. Biomarkers for increased risk include elevated homocysteine, C-reactive protein, and fibrinogen levels.

35

Myocardial Infarction (MI)

[0050] A myocardial infarction (MI) occurs when an area of heart muscle becomes necrotic (dies) or is permanently damaged because of an inadequate supply of oxygen to that area (ischemia). Most heart attacks are caused by a clot that blocks one of the coronary arteries. Clots usually form in a coronary artery that has been previously narrowed from changes related to atherosclerosis; atherosclerotic plaque (buildup) inside the arterial wall sometimes cracks, and this triggers the formation of a clot, also called a thrombus.

40

[0051] A clot in the coronary artery interrupts the flow of blood and oxygen to the heart muscle, resulting in ischemia of the tissue leading to the death of heart cells in that area (necrosis). The damaged heart muscle loses its ability to contract, and the remaining heart muscle needs to compensate for that weakened area

45

Acute Coronary Syndrome (ACS)

[0052] ACS is a term that is used to cover any group of clinical symptoms associated with acute myocardial ischemia. Patients with ACS include those whose clinical presentations the following range of diagnoses: unstable angina, non-ST-elevation myocardial infarction (NSTEMI), and ST-elevation myocardial infarction (STEMI). Myocardial ischemia is most often due to atherosclerotic plaques, as described above for MI.

50

Heart Failure (HF)

55

[0053] Heart failure is a pathophysiologic state in which the heart, via an abnormality of cardiac function (detectable or not), fails to pump blood at a rate commensurate with the requirements of the metabolizing tissues, and/or pumps

only from an elevated diastolic filling pressure.

[0054] Heart failure may be causal by myocardial failure but may also occur in the preserve of near-normal cardiac function under conditions of high demand, e.g., stress, Inadequate adaptation of the cardiac myocytes to increased wall stress to maintain adequate cardiac output following myocardial injury, is generally the meting event in CHF. HF may have an acute onset or may occur over several months to years, due to a primary disturbance in myocardial contractility or an excesssive hemodynamic burden placed on the ventricle, or both.

Special Populations

[0055] Certain populations of may benefit particularly from the methods described herein. These subjects include people for whom BNP or NT-proBNP is less useful, such as in those with impaired renal function (Anwaruddin et al., J. Am. Coll. Cardiol. 47(1):91-7 (2006); McCullough et al., Am. J. Kidney Dis. 41 (3):571-9 (2003)), or in those who are overweight (Body Mass Index (BMI) of 25-29) or obese (BMI \geq 30) (Krauser et al., Am. Heart J. 149(4):744-50 (2005); McCord et at, Arch. Intern. Med. 164(20):2247-52 (2004)). It is known and accepted in the field that patients with a high BMI usually have levels of natriuretic peptide that are lower than expected relative to a normal body mass patient for the same level of disease; the exact mechanism for this phenomerion is not known. IL-33 levels may not be affected by renal failure or high BMI. The methods described herein can include determining a subject's BMI, and if the subject is overweight or obese, selecting the patient for determination of IL-33 levels, as described herein.

EXAMPLES

[0056] The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

Example 1: Detection and Measurement of IL-33 in the Serum

[0057] A blood sample is collected from a subject, and serum is prepared from the sample using standard methods. A labeled monoclonal antibody to IL-33 (e.g., as described U.S. Pat. App. Pub. No. 2005/0203046, incorporated herein by reference in its entirety) in is added to the sample and incubated for a sufficient amount of time for binding to occur. The antibody/IL-33 complexes are then detected using standard methods, and the amount of IL-33 present is quantified.

OTHER EMBODIMENTS

[0058] It is to be understood that while the invention has been described in with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

SEQUENCE LISTING

[0059]

<110> Critical Care Dagnostics, Inc.

<120> INTERLEUKIN-33 (IL- 33) FOR THE DIAGNOSIS AND PROGNOSIS OF CARDIOVASCULAR DISEASE

<130> 20060- 004WO1

<140> PCT/ US2007/ 067626<141>2007- 04- 27

<150> US 60/795,473<151> 2006-04-27

<160> 2

<170> Fast SEQ for Windows Version 4.0

<210> 1

<211> 2644

<212> DNA

EP 2 021 799 B1

<213> Homo sapiens

<400> 1

5 caacagaat a ct gaaaaat g aagcct aaaa t gaagt at t c aaccaacaaa at t t ccacag 60
 caaagt ggaa gaacacagca agcaaagcct t gt gt t t caa gct gggaaaa t cccaacaga 120
 aggccaaaga agt t t gcccc at gt act t t a t gaagct ccg ct ct ggcct t at gat aaaaa 180
 aggaggcct g t t act t t agg agagaaacca ccaaaaggcc t t cact gaaa acaggt agaa 240
 agcacaaaaag acat ct ggt a ct cgct gcct gt caacagca gt ct act gt g gagt gct t t g 300
 cct t t ggt at at caggggt c cagaaaat at a ct agagcact t cat gat t ca agt at cacag 360
 10 gaat t t cacc t at t acagag t at ct t gct t ct ct aagcac at acaat gat caat ccat t a 420
 ct t t t gct t t ggaggat gaa agt t at gaga t at at gt t ga agact t gaaa aaagat gaaa 480
 agaaagat aa ggt gt t act g agt t act at g agt ct caaca cccct caaat gaat caggt g 540
 acgggt gt t ga t ggt aagat g t t aat ggt aa ccct gagt cc t acaaaagac t t ct ggt t gc 600
 at gccaaaca caaggaacac t ct gt ggagc t ccat aagt g t gaaaaacca ct gccagacc 660
 aggcct t ct t t gt cct t cat aat at gcact ccaact gt gt t t cat t t gaa t gcaagact g 720
 15 at cct ggagt gt t t at aggt gt aaagat a at cat ct t gc t ct gat t aaa gt agact ct t 780
 ct gagaat t t gt gt act gaa aat at ct t gt t t aagct ct c t gaaact t ag t t gat ggaaa 840
 cct gt gagt c t t ggggt t gag t acccaaat g ct accact gg agaaggaat g agagat aaag 900
 aaagagacag gt gacat ct a agggaaat ga agagt gct t a gcat gt gt gg aat gt t t t cc 960
 at at t at gt a t aaaaaat at t t t t t ct aat c ct ccagt t at t ct t t t at t t ccct ct gt at 1020
 aact gcat ct t caat acaag t at cagt at a t t aat aggg t at t ggt aaa gaaacggt ca 1080
 20 acat t ct aaa gagat acagt ct gacct t t a ct t t t ct ct a gt t t cagt cc agaaagaact 1140
 t cat at t t ag agct aaggcc act gaggaaa gagccat agc t t aagt ct ct at gt agacag 1200
 ggat ccat t t t aaagagct a ct t agagaaa t aat t t t cca cagt t ccaaa cgat aggct c 1260
 aaacact aga gct gct agt a aaaagaagac cagat gct t c acagaat t at cat t t t t t ca 1320
 act ggaat aa aacaccaggt t t gt t t gt ag at gt ct t agg caacact cag agcagat ct c 1380
 cct t act gt c aggggat at g gaact t caaa ggcccacat g gcaagccagg t aacat aaat 1440
 25 gt gt gaaaaa gt aaagat aa ct aaaaaat t t agaaaaat a aat ccagt at t t gt aaagt g 1500
 aat aact t ca t t t ct aat t g t t t aat t t t t aaaat t ct ga t t t t t at at a t t gagt t t aa 1560
 gcaaggcat t ct t acacgag gaagt gaagt aaat t t t agt t cagacat aa aat t t cact t 1620
 at t aggaat a t gt aacat gc t aaaact t t t t t t t t t aa agagt act ga gt cacaacat 1680
 gt t t t agagc at ccaagt ac cat at aat cc aact at cat g gt aaggccag aaat ct t ct a 1740
 accct accaga gccct agat ga gacaccgaat t aacat t aaa at t t cagt aa ct gact gt cc 1800
 30 ct cat gt cca t ggcct acca t cct t ct ga cctt ggct t c cagggacct a t gt ct t t t aa 1860
 t act cact gt cacat t gggc aaagt t gct t ct aat cct t a t t t cccat gt gcacaagt ct 1920
 t t t t gt at t c cagct t cct g at aacact gc t t act gt gga at at t cat t t gacat ct gt c 1980
 t ct t t t cat t t ct t t t aact accat gccct t gat at at ct t t t gcacct g ct gaact t ca 2040
 t t t ct gt at c acct gacct c t ggt gccaa aacgt t t at t ct gct t t gt c t gt t gt agaa 2100
 35 t t t t agat aa agct at t aat ggcaat at t t t t t gct aaa cgt t t t t gt t t t t t act gt c 2160
 act agggcaa t aaaaat t t at act caaccat at aat aacat t t t t t aact a ct aaaggagt 2220
 agt t t t t at t t t aaagt ct t agcaat t t ct at t acaact t t ct t agact t aacact t at 2280
 gat aaat gac t aacat agt a acagaat ct t t at gaaat at gacct t t t ct gaaaaat acat 2340
 act t t t acat t t ct act t t a t t gagacct a t t agat gt aa gt gct agt ag aat at aagat 2400
 aaaagaggct gagaat t acc at acaagggt at t acaact g t aaaaact t t at ct t t gt t 2460
 40 t cat t gt t ct gt caat aat t gt t accaaag agat aaaaaat aaaagcagaa t gt at at cat 2520

45 cccat ct gaa aaacact aat t at t gacat g t gcat ct gt a caat aaact t aaaaat gat t a 2580
 t t aaat aat c aaat at at ct act acat t gt t t at at t at t gaat aaagt a t at t t t ccaa 2640
 at gt 2644

<210> 2

<211> 270

50 <212> PRT

<213> Homo sapiens

<400> 2

55

Met 1 Lys Pro Lys Met 5 Lys Tyr Ser Thr Asn 10 Lys Ile Ser Thr Ala Lys
 Trp Lys Asn Thr 20 Ala Ser Lys Ala Leu 25 Oys Phe Lys Leu Gly 30 Lys Ser
 5 G n G n Lys 35 Ala Lys G u Val Oys 40 Pro Met Tyr Phe Met 45 Lys Leu Arg
 Ser G y Leu Met 50 Ile Lys Lys 55 G u Ala Oys Tyr Phe 60 Arg Arg G u Thr
 Thr Lys Arg Pro Ser 65 Leu 70 Lys Thr Gly Arg Lys 75 His Lys Arg His Leu
 10 Val Leu Ala Ala Oys 85 G n G n G n Ser Thr 90 Val G u Oys Phe Ala Phe
 Gly Ile Ser Gly 100 Val G n Lys Tyr Thr 105 Arg Ala Leu His Asp Ser Ser
 Ile Thr G y 115 Ile Ser Pro Ile Thr 120 G u Tyr Leu Ala Ser 125 Leu Ser Thr
 15 Tyr Asn Asp G n Ser 130 Ile Thr Phe Ala Leu G u Asp 140 G u Ser Tyr G u
 Ile Tyr Val G u Asp 145 Leu 150 Lys Lys Asp G u Lys 155 Lys Asp Lys Val Leu
 Leu Ser Tyr Tyr G u 165 Ser G n His Pro Ser 170 Asn G u Ser G y Asp 175 Gly
 20 Val Asp G y Lys 180 Met Leu Met Val Thr 185 Leu Ser Pro Thr Lys Asp Phe
 Trp Leu His 195 Ala Asn Asn Lys G u 200 His Ser Val G u Leu 205 His Lys Cys
 G u Lys Pro Leu Pro Asp 215 G n Ala Phe Phe Val Leu His Asn Met His
 25 Ser Asn Oys Val Ser Phe 220 G u Cys Lys Thr Asp 225 Pro G y Val Phe Ile
 225 G y Val Lys Asp Asn 230 His Leu Ala Leu Ile 235 Lys Val Asp Ser Ser 240 G u
 Asn Leu Oys Thr 245 G u Asn Ile Leu Phe 250 Lys Leu Ser G u Thr 255
 30 260 265 270

Claims

- 35 1. An in vitro method of aiding in the diagnosis of a cardiovascular disease in a subject, the method comprising determining a level of IL-33 in a sample comprising blood, serum or plasma from a subject, wherein the level of IL-33 in the sample is indicative of whether the subject has a cardiovascular disease and further comprising comparing the level of IL-33 in the sample to a reference level, wherein the reference level represents a level in a subject who has or does not have cardiovascular disease.
- 40 2. An in vitro method of aiding in determining the severity of a cardiovascular disease in a subject, the method comprising determining a level of IL-33 in a sample comprising blood, serum or plasma from a subject, wherein the level of IL-33 in the sample is indicative of the severity of the cardiovascular disease in the subject and further comprising comparing the level of IL-33 in the sample to a reference level, wherein the reference level represents a level in a subject with a preselected severity of cardiovascular disease.
- 45 3. The method of claim 1 or 2, wherein determining a level of IL-33 in the sample comprises determining a level of one, two, or all three of mature IL-33, pre-IL-33, and pro-IL-33
- 50 4. The method of claim 1 or 2, wherein determining a level of IL-33 in the sample comprises contacting a binding composition to the sample, wherein the binding composition specifically binds to IL-33, and measuring or determining the specific binding of the binding composition to the sample.
- 55 5. The method of claim 4, wherein the binding composition comprises an antibody that binds specifically to IL-33 polypeptide.
6. The method of claim 5, wherein the binding composition comprises an oligonucleotide probe that binds specifically to IL-33 polynucleotide.

7. The method of claim 1 or 2, wherein the cardiovascular disease is acute coronary syndrome (ACS), myocardial infarction, heart failure, angina, cardiac hypertrophy, arteriosclerosis, myocarditis, pericarditis, endocarditis, stroke and/or pulmonary embolism.
- 5 8. The method of claim 1 or 2, further comprising determining a level in the sample of one or more other biomarkers.
9. The method of claim 8, wherein the other biomarkers are selected from the group consisting of ST2, NT-proBNP, BNP, NT-proANP, and ANP, troponin, IMA, IL-6, CRP, creatinine, D-dimers, BUN, liver function enzymes, albumin, and bacterial endotoxin.
- 10 10. A method of monitoring the condition of a subject previously diagnosed with a cardiovascular condition, the method comprising:
- 15 determining a first level of IL-33 in a sample comprising blood, serum or plasma from a subject;
determining at least one subsequent level of IL-33 in a sample comprising blood, serum or plasma from the subject; and
- 20 comparing the level of IL-33 in the subsequent sample to the level of IL-33 in the first sample, wherein a difference in the IL-33 level in the subsequent sample relative to the level in the first sample is correlated with a change in the patient's condition.
11. The method of claim 10, wherein the subject is undergoing treatment on an outpatient basis.
12. The method of claim 10, wherein an increase in IL-33 in the subsequent sample relative to the level in the first sample indicates that the patient's condition is deteriorating.
- 25 13. The method of claim 12, wherein an increase in IL-33 in the subsequent sample relative to the level in the first sample indicates that the current treatment should be adjusted.
- 30 14. The method of claim 12, wherein an increase in IL-33 in the subsequent sample relative to the level in the first sample indicates that the patient should be admitted to a hospital for inpatient treatment.

Patentansprüche

- 35 1. In vitro-Verfahren zum Unterstützen des Diagnostizierens einer kardiovaskulären Erkrankung bei einem Subjekt, wobei das Verfahren das Bestimmen eines IL-33-Spiegels in einer Probe, die Blut, Serum oder Plasma von einem Subjekt umfasst, wobei der IL-33-Spiegel in der Probe dafür indikativ ist, ob das Subjekt an einer kardiovaskulären Krankheit leidet, und des Weiteren das Vergleichen des IL-33-Spiegels in der Probe mit einem Bezugsspiegel umfasst, wobei der Bezugsspiegel einen Spiegel bei einem Subjekt darstellt, das an einer kardiovaskulären Krankheit leidet oder nicht.
- 40 2. In vitro-Verfahren zum Unterstützen des Bestimmens der Schwere einer kardiovaskulären Erkrankung bei einem Subjekt, wobei das Verfahren das Bestimmen eines IL-33-Spiegels in einer Probe, die Blut, Serum oder Plasma von einem Subjekt umfasst, wobei der IL-33-Spiegel in der Probe für die Schwere der kardiovaskulären Krankheit bei dem Subjekt indikativ ist und des Weiteren das Vergleichen des IL-33-Spiegels in der Probe mit einem Bezugsspiegel umfasst, wobei der Bezugsspiegel einen Spiegel bei einem Subjekt darstellt, das an einer vorher gewählten Schwere einer kardiovaskulären Krankheit leidet.
- 45 3. Verfahren nach Anspruch 1 oder 2, wobei das Bestimmen eines IL-33-Spiegels in der Probe das Bestimmen eines Spiegels von einem, zwei oder allen drei von reifem IL-33, Prä-IL-33 und Pro-IL-33 umfasst.
- 50 4. Verfahren nach Anspruch 1 oder 2, wobei das Bestimmen eines IL-33-Spiegels in der Probe das Kontaktieren einer Bindungszusammensetzung mit der Probe, wobei die Bindungszusammensetzung sich spezifisch an IL-33 bindet, und das Messen oder Bestimmen der spezifischen Bindung der Bindungszusammensetzung an die Probe umfasst.
- 55 5. Verfahren nach Anspruch 4, wobei die Bindungszusammensetzung einen Antikörper umfasst, der sich spezifisch an IL-33-Polypeptid bindet.

EP 2 021 799 B1

6. Verfahren nach Anspruch 5, wobei die Bindungszusammensetzung eine Oligonucleotidsonde umfasst, die sich spezifisch an IL-33-Polynucleotid bindet.
- 5 7. Verfahren nach Anspruch 1 oder 2, wobei die kardiovaskuläre Krankheit das akute Koronarsyndrom (ACS), Herzinfarkt, Herzversagen, Angina, Herzhypertrophie, Arteriosklerose, Myokarditis, Perikarditis, Endokarditis, Schlaganfall und/oder Lungenembolie ist.
- 10 8. Verfahren nach Anspruch 1 oder 2, des Weiteren das Bestimmen eines Spiegels in der Probe eines oder mehrerer anderer Biomarker umfassend.
- 15 9. Verfahren nach Anspruch 8, wobei die anderen Biomarker aus der Gruppe ausgewählt sind bestehend aus ST2, NT-proBNP, BNP, NT-proANP und ANP, Troponin, IMA, IL-6, CRP, Creatinin, D-Dimeren, BUN, Leberfunktionsenzymen, Albumin und Bakterienendotoxin.
- 20 10. Verfahren zum Überwachen des Zustands eines Subjekts, das zuvor als an einer kardiovaskulären Beschwerde leidend diagnostiziert worden ist, wobei das Verfahren folgendes umfasst:
- das Bestimmen eines ersten IL-33-Spiegels in einer Probe, die Blut, Serum oder Plasma von einem Subjekt umfasst;
- das Bestimmen mindestens eines darauffolgenden IL-33-Spiegels in einer Probe, die Blut, Serum oder Plasma von einem Subjekt umfasst; und
- das Vergleichen des IL-33-Spiegels in der darauffolgenden Probe mit dem IL-33-Spiegel in der ersten Probe, wobei ein Unterschied des IL-33-Spiegels in der darauffolgenden Probe mit Bezug auf den Spiegel in der ersten Probe mit einer Änderung des Zustands der Patienten korreliert.
- 25 11. Verfahren nach Anspruch 10, wobei das Subjekt eine Behandlung auf ambulanter Basis erfährt.
- 30 12. Verfahren nach Anspruch 10, wobei eine Erhöhung von IL-33 in der darauffolgenden Probe mit Bezug auf den Spiegel in der ersten Probe anzeigt, dass der Zustand des Patienten sich verschlechtert.
- 35 13. Verfahren nach Anspruch 12, wobei eine Erhöhung von IL-33 in der darauffolgenden Probe mit Bezug auf den Spiegel in der ersten Probe anzeigt, dass die gegenwärtige Behandlung angepasst werden sollte.
14. Verfahren nach Anspruch 12, wobei eine Erhöhung von IL-33 in der darauffolgenden Probe mit Bezug auf den Spiegel in der ersten Probe anzeigt, dass der Patient zur stationären Behandlung in ein Krankenhaus aufgenommen werden sollte.

Revendications

- 40 1. Procédé *in vitro* pour faciliter le diagnostic d'une maladie cardio-vasculaire chez un sujet, le procédé comprenant la détermination d'un taux d'IL-33 dans un échantillon comprenant du sang, du sérum ou du plasma d'un sujet, dans lequel le taux d'IL-33 dans l'échantillon indique si le sujet a une maladie cardio-vasculaire, et comprenant en outre la comparaison du taux d'IL-33 dans l'échantillon avec un taux de référence, dans lequel le taux de référence représente un taux chez un sujet qui a ou n'a pas une maladie cardio-vasculaire.
- 45 2. Procédé *in vitro* pour faciliter la détermination de la sévérité d'une maladie cardio-vasculaire chez un sujet, le procédé comprenant la détermination d'un taux d'IL-33 dans un échantillon comprenant du sang, du sérum ou du plasma d'un sujet, dans lequel le taux d'IL-33 dans l'échantillon est indicatif de la sévérité de la maladie cardio-vasculaire chez le sujet, et comprenant en outre la comparaison du taux d'IL-33 dans l'échantillon avec un taux de référence, dans lequel le taux de référence représente un taux chez un sujet ayant une sévérité présélectionnée de maladie cardio-vasculaire.
- 50 3. Procédé selon la revendication 1 ou 2, dans lequel la détermination d'un taux d'IL-33 dans l'échantillon comprend la détermination d'un taux d'une, de deux ou de toutes les trois parmi IL-33 mature, pré-IL-33 et pro-IL-33.
- 55 4. Procédé selon la revendication 1 ou 2, dans lequel la détermination d'un taux d'IL-33 dans l'échantillon comprend la mise en contact d'une composition de liaison avec l'échantillon, dans lequel la composition de liaison se lie

EP 2 021 799 B1

spécifiquement à IL-33, et la mesure ou la détermination de la liaison spécifique de la composition de liaison à l'échantillon.

- 5
5. Procédé selon la revendication 4, dans lequel la composition de liaison comprend un anticorps qui se lie spécifiquement au polypeptide IL-33.
6. Procédé selon la revendication 5, dans lequel la composition de liaison comprend une sonde oligonucléotidique qui se lie spécifiquement au polynucléotide IL-33.
- 10
7. Procédé selon la revendication 1 ou 2, dans lequel la maladie cardio-vasculaire est un syndrome coronaire aigu (SCA), un infarctus du myocarde, une défaillance cardiaque, une angine de poitrine, une hypertrophie cardiaque, une artériosclérose, une myocardite, une péricardite, une endocardite, une attaque et/ou une embolie pulmonaire.
- 15
8. Procédé selon la revendication 1 ou 2, comprenant en outre la détermination d'un taux d'un ou plusieurs autres biomarqueurs dans l'échantillon.
- 20
9. Procédé selon la revendication 8, dans lequel les autres biomarqueurs sont choisis dans le groupe constitué par ST2, NT-proBNP, BNP, NT-proANP et ANP, troponine, IMA, IL-6, CRP, créatinine, D-dimères, BUN, enzymes de la fonction hépatique, albumine et endotoxine bactérienne.
- 25
10. Procédé pour suivre l'affection d'un patient préalablement diagnostiqué avec une affection cardio-vasculaire, le procédé comprenant:
- 30
- la détermination d'un premier taux d'IL-33 dans un échantillon comprenant du sang, du sérum ou du plasma d'un sujet;
 - la détermination d'au moins un taux subséquent d'IL-33 dans un échantillon comprenant du sang, du sérum ou du plasma du sujet; et
 - la comparaison du taux d'IL-33 dans l'échantillon subséquent avec le taux d'IL-33 dans le premier échantillon, dans lequel une différence entre le taux d'IL-33 dans l'échantillon subséquent et le taux dans le premier échantillon est corrélée avec un changement de l'affection du patient.
- 35
11. Procédé selon la revendication 10, dans lequel le sujet subit un traitement de façon ambulatoire.
- 40
12. Procédé selon la revendication 10, dans lequel une augmentation d'IL-33 dans l'échantillon subséquent par rapport au taux dans le premier échantillon indique que l'affection du patient se détériore.
- 45
13. Procédé selon la revendication 12, dans lequel une augmentation d'IL-33 dans l'échantillon subséquent par rapport au taux dans le premier échantillon indique que le traitement en cours devrait être ajusté.
- 50
14. Procédé selon la revendication 12, dans lequel une augmentation d'IL-33 dans l'échantillon subséquent par rapport au taux dans le premier échantillon indique que le patient devrait être admis dans un hôpital pour un traitement avec hospitalisation.
- 55

FIG. 1

```

1 caacagaata cggaaaantg aagcctaaaa tgaagtattc aaccaaaaaa atttcacag
61 caaaggggaa gaacacagca agcaaagcct tgtgtttcaa gctgggaaaa toccaacaga
121 aggcdaaaga agtttgcccc atgtacttta tgaagctccg ctctggcctt atgataaaaa
181 agggggcctg ttacttttagg agagaaaacca ccaaaaggcc ttcactgaaa acaggtagaa
241 agcacaaaag acatctggta ctctctgccc gtcaacagca gctctctgtg gagggctttg
301 cctttgggtat aacagggggtc cagaaabata cttagagcact tcatgaltca agtatcacag
361 gaatttcacc tattacagag tatcttgcct ctctaagcac atacaatgat caatccatta
421 cttttgcttt ggaggatgaa agttatgaga tatatgttga agacttgaaa aaagatgaaa
481 agaaagataa ggtgttactg agttactatg agtctnaaca cccctcaaat gaatcaggtg
541 acggtggtga tggtaagatg ttaatggtaa ccoctgagtc taaaaaagac ttctggttgc
601 atgccaacaa caaggaacac ctctgtggagc tccataagtg taaaaacca ctgccagacc
661 aggccttctt tgtccctcat aatctgcact ccaactgtgt tucattttaa tgcagaactg
721 atcctggagt gtttataggt gtaaggata atcatcttgc tctgattaaa gtgactctt
781 ctgagaatit gtgtactgaa aatatcttgt ttaagctctc tgaacttag ttgatggaaa
841 cctggagctc ttgggtggag tacccaaatg ctaccactgg agaaggaatg agagataaag
901 aagagagcag gtgacatcta agggaaatga agagagctta gcatgtgpyg gcatttttcc
961 atattatgta taaaaatatt ttttcaatc ctccagttar tcttttattt cctctongtat
1021 aactgcctct tcaacacaaq tatcagtata ttaaataggg tattggtaaa gaaacggta
1081 acattctaaa gagatacagt cagaacttta cttttctcta gtttcagtc agaaagaact
1141 tcatctttag agctaaggcc actgaggaaa gagccatage ttaagtctct atgtagacag
1201 ggcctcaatt taagagagca cttagagaaa taattttcca cagttccaaa cgtatagctc
1261 aaacctaga gctgctagta aaaagaagac cagatgcttc acagaattat catcttttca
1321 actggaaata aacaccaggt ttgtttgtag atgtcttagg caacactcag agcagatctc
1381 ccttactgtc aggggatatg gaactcaaaa ggcccacatg gcaagccagg taacataaet
1441 gtgggaaaaa gtaagataa ctuaaaaaat tagaaaaata aatccagat ttgtaaagtg
1501 ataacctca tttctaatgg ttaattttt aaaaattctg tttttatata ttgagtttaa
1561 gcaagcctatt cttacacgag gaagtgaagt eaattttagt ccagacataa aatttcaact
1621 attaggaata tghtaacatg taanaacttt ttttttttaa agagtactga gtcaaacat
1681 gttttagagc aaccaagtae catataatcc aactatctag gtaagggcag aaactctca
1741 aacbaacaga gcctaganga gacaccyaat taacattaaa atttcagtaa ctgactgtcc
1801 ctcatgtcca tggcctacca tccctctgta cccggcttc cagggaccca tgtcttttaa
1861 tactcactgt cacattgggc aaagttgctt ctaatectta tttcccagt gcaaaagtct
1921 tttgtattc cagcttctg ataacactgc ttactgtgga atattcaatt gacatctgtc
1981 tctttctatt tcttttaact accatgcctc tgaatatact tttgacctg ctgacttca
2041 tttctgtate acctgacctc tggatgcccc aactgtttat cngctttgtc tctgagaa
2101 ttttagataa egctattaat ggcattatit tttngotaaa cgtttttgtt ttttactgtc
2161 actagggcaa caaaatttat acctaacct atataaacat tttttaacta ctaaaagggt
2221 agtcttatt ttaagctctt agcaatttct attacaactt ttcttagact taactctat
2281 gataaatgac taacatagta acagaactct tatgaaatat gaccttttct gaaaatacat
2341 acttttacat ttctactata ttgagaccta ttgatgtaa gtgctagtag aatataagat
2401 aaaagaggtc gagaattacc atacaagggg attacaactg taaaacaatt tatctttgtt
2461 tcatgttct gtcaataaatt gttaccaaaq agataaaaaa aaaagcagaa tgtatatcat
2521 cccatctgaa aaacactaat tattgacatg tgcactctga caataaacn aaatgatta
2581 ttaataatc aatatatct actacattgt ttatattatt gaataaagta tattttccaa
2641 atgt (SEQ ID NO:1)

```

FIG. 2

```

1 mkpkmkystn kistakwkn askalcflg ksqkakevc pmyfmklrsg imikkeacyt
61 rrehtkrpsl kgrkhrhl vlaacqqqt vecfagisg vqkytralhd siltgispit
121 aylaslstyn dqsitfaled esyeyvedl kkdexkdkvl lyyesqhrs nesgdgvdgk
181 mlmvtlapk dfwihanne hveihkcek plpdqaffvl hmhancvsv ecktdpyvfi
241 gvkdnhlali kvdsenict enilfklsr (SEQ ID NO:2)

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20050203046 A [0019] [0035] [0038] [0057]
- US 6176962 B [0028]
- US 6517234 B [0028]
- US 20030124624 A [0043]
- US 20040048286 A [0043] [0045]
- US 20050130136 A [0043] [0045]
- US 20050250156 A [0043]
- US 2007067626 W [0059]
- US 60795473 B [0059]

Non-patent literature cited in the description

- **Shimpo et al.** Serum levels of interleukin-1 receptor family member ST2 predict mortality in clinical outcome in acute myocardial infarction. *Circulation*, 11 May 2004, vol. 109 (18), 2186-2190 [0003]
- **Schmitz et al.** IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity*, 01 November 2005, vol. 23 (5), 479-490 [0003]
- **Schmitz et al.** *Immunity*, 2005, vol. 23 (5), 479-90 [0019] [0035] [0038] [0040]
- Current Protocols in Molecular Biology. John Wiley & Sons, 1989, 6.3.1-6.3.6 [0026]
- **Le Doussal et al.** *New Engl. J. Med.*, 1991, vol. 146, 169-175 [0028]
- **Gibellini et al.** *J. Immunol.*, 1998, vol. 160, 3891-3898 [0028]
- **Hsing ; Bishop.** *New Engl. J. Med.*, 1999, vol. 162, 2804-2811 [0028]
- **Everts et al.** *New Engl. J. Med.*, 2002, vol. 168, 883-889 [0028]
- **Maniatis et al.** Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1982 [0029]
- **Sambrook ; Russell.** Molecular Cloning. Cold Spring Harbor Laboratory Press, 2001 [0029]
- **Wu.** Recombinant DNA. Academic Press, 1993, vol. 217 [0029]
- **Ausbel et al.** Current Photocols in Molecular Biology. John Wiley and Sons, Inc, 2001, vol. 1-4 [0029]
- **Baekkevold et al.** *Am. J. Path.*, 2003, vol. 163, 69-79 [0037]
- **Kieser et al.** *FEBS Lett.*, 1995, vol. 372 (2-3), 189-93 [0039]
- **Kumar et al.** *J. Biol. Chem.*, 1993, vol. 270 (46), 27905-13 [0039]
- **Yanagisawa et al.** *FEBS Lett.*, 1992, vol. 302 (1), 51-3 [0039]
- **Kuroiwa et al.** *Hybridoma*, 2000, vol. 19 (2), 151-9 [0039]
- **Weinberg et al.** *Circulation*, 2002, vol. 106 (23), 2961-6 [0039]
- **Weinherg et al.** *Circulation*, 2003, vol. 107 (5), 721-6 [0039]
- **Shimpo et al.** *Circulation*, 2004, vol. 109 (18), 2186-90 [0039]
- **Lohning et al.** *Proc. Natl. Acad. Sei. U. S. A.*, 1998, vol. 95 (12), 6930-5 [0040]
- **Brint et al.** *Nat. Immunol.*, 2004, vol. 5 (4), 373-9 [0040]
- **Bruneau et al.** *Cardiovasc. Res.*, 1994, vol. 28 (10), 1519-25 [0040]
- **Tominaga.** *FEBS Lett.*, 1989, vol. 258, 301-304 [0041]
- **Torminaga et al.** *Biochim. Biophys. Acta*, 1992, vol. 1171, 215-218 [0041]
- **Dhalla et al.** *Mol. Cell Biochem.*, 1989, vol. 87, 85-92 [0045]
- **Moe et al.** *Am. Heart J.*, 2000, vol. 139, 587-95 [0045]
- **Anwaruddin et al.** *J. Am. Coll. Cardiol.*, 2006, vol. 47 (1), 91-7 [0055]
- **McCullough et al.** *Am. J. Kidney Dis.*, 2003, vol. 41 (3), 571-9 [0055]
- **Krauser et al.** *Am. Heart J.*, 2005, vol. 149 (4), 744-50 [0055]
- **McCord.** *Arch. Intern. Med.*, 2004, vol. 164 (20), 2247-52 [0055]

专利名称(译)	白细胞介素-33 (il-33) 用于心血管疾病的诊断和预后		
公开(公告)号	EP2021799B1	公开(公告)日	2011-07-20
申请号	EP2007811859	申请日	2007-04-27
[标]申请(专利权)人(译)	重症监护诊断股份有限公司		
申请(专利权)人(译)	急救诊断, INC.		
当前申请(专利权)人(译)	急救诊断, INC.		
[标]发明人	SNIDER JAMES JACOBSON SVEN		
发明人	SNIDER, JAMES JACOBSON, SVEN		
IPC分类号	G01N33/53 C07K14/00 C07K16/00 G01N33/68		
CPC分类号	C12Q1/6883 G01N33/6869 G01N2800/32 Y10T436/143333		
优先权	60/795473 2006-04-27 US		
其他公开文献	EP2021799A2 EP2021799A4		
外部链接	Espacenet		

摘要(译)

本发明包括使用白介素33 (IL-33) 诊断心血管疾病的方法, 所述疾病包括急性冠状动脉综合征 (ACS), 心肌梗塞和/或心力衰竭, 心绞痛, 心脏肥大, 动脉硬化, 心肌炎, 全心炎, 心内膜炎, 中风和/或肺栓塞, 并确定此类疾病的严重程度 (预后) 。

```

caacagaat a ct gaaaaat g aagcct aaaa tgaagt at c aaccaacaaa at t tccacag 60
caaa g gaa gaacacagca agcaagcct t g t l t caa gcl g g g a a a t c c c a c a g a 120
agccaaaga agt t t g c c c at g t a c l t t a t g a a g t c e g c t c t g g c t t at g a t a a a a 180
aggaggcct g t t a c t t a g g a g a a a c c a c a a a g g c c t t c a c t g a a a c a g g t a g a a 240
agcaaaaaag acat ct ggt a ct c g c t g c c t g t c a a c a g c a g t c t a c t g t g g a g t g c t t g 300
c c t t t g g t a t at c a g g g t c c a g a a a t a c t a g a g c a c t t c a t g a t t c a a g t a t c a c a g 360
g a a t t c a c c t a t t a c a g a g t a t c t t g c t t c t a g c a c at a c a a t g a t c a a t c c a t t a 420
c t t t g c t t t g g a g g a t g a a a g t t a t g a g a t a t a t g t g a a g a c t t g a a a a a a g a t g a a a 480
a g a a g a t a a g g t g t a c t g a g t a c t a t g a g t c t c a a c a c c c t c a a a t g a a t c a g g t g 540
a c g g t g t g a t g g t a a g a t g t t a a t g g t a a c c c t g a g t c c t a c a a a a g a c t t c t g g t t g c 600
at g c c a a c a a c a a g g a a c a c t c t g t g a g c t c c a t a a g t g t g a a a a a c c a c t g c c a g a c c 660
a g g c c t c t t t g t c c t c a t a a t a t g c a c t c c a a c t g t g t t c a t t t g a a t g c a a g a c t g 720
at c c t g a g t g t t a t a g g t g t a a g g a t a a t c a t c t t g c t c t g a t t a a a g t a g a c t c t t 780
c t g a g a a t t t g t a c t g a a a a t a t c t t g t t a a g c t c t t g a a a c t t a g t g a t g g a a a 840
c c t g t g a g t t t g g g t g a g t a c c a a a t g c t a c c a c t g g a g a a g a a t g a g a t a a a g 900
a a a g a g a c a g g t g a c a t c t a a g g g a a a t g a a g a g t g c t t a g c a t g t g t g g a a t g t t t c c 960
at a t t a t g t a t a a a a t a t t t t c t a a t c c t c a g t t a t t c t t t t a t t t c c t c t g t a t 1020
a a c t g c a t c t t c a a t a a a g t t a t c a g t a t a t t a a t a g g g t a t t g g t a a a g a a c g g t c a 1080
a c a t t c t a a a g a g a t a c a g t c t g a c c t t t a c t t t c t c t a g t t c a g t c c a g a a a g a a c t 1140
t c a t a t t a g a g t a a g g c c a c t g a g g a a a g a g c c a t a g c t t a a g t c t c t a t g t a g a c a g 1200
g g a t c c a t t t t a a g a g a t a c t t a g a g a a t a a t t t t c c a c a g t t c c a a a c g a t a g g c t c 1260
a a a c a c t a g a g c t g c t a g t a a a a g a a g a c c a g a t g c t t c a c a g a a t t a t c a t t t t t t c a 1320
a c t g g a a t a a a a c a c c a g g t t t g t t t g t a g a t g t c t t a g g c a a c a c t c a g a c a g a t c t c 1380
c c t t a c t g t c a g g g a t a t g g a a c t c a a a g g c c a a c t g g c a a g c a g g t a a a c a t a a a t 1440
g t g t g a a a a g t a a g a t a a c t a a a a a a t t t a g a a a a t a a t c a a g t a t t g t a a a g t g 1500
a a t a a c t t c a t t t c t a a t t g t t t a a t t t t a a a a t t c t g a t t t t a t a t a t t g a g t t t a a 1560
g c a a g g c a t t c t t a c a c a g a g a a g t g a a g t a a a t t t a g t t c a g a c a t a a a a t t c a c t t 1620
a t t a g g a a t a t g t a a c a t g c t a a a c t t t t t t t t t t t t a a a g a g t a c t g a t c a c a c a t 1680
g t t t a g a g c a t c c a a g t a c c a t a t a a t c c a a c t a t c a t g t a a g g c c a g a a a t c t t c a 1740
a c c t a c c a g a g c c t a g a t g a g a c a c c a a t t a a c a t t a a a a t t t c a g t a a c t g a c t g t c c 1800
c t c a t g t c c a t g g c c t a c c a t c c c t t c t g a c c c t g g c t t c c a g g a c a c t a t g t c t t t t a a 1860
t a c t c a c t g t c a c a t t g g g c a a g t t g c t t c a t a a c t t a t t t c c a t g t g c a a a g t c t 1920
t t t t g t a t t c c a g c t t c c t g a t a a c a c t g c t t a c t g t g g a a t a t t c a t t t g a c a t c t g c 1980
t c t t t c a t t t c t t t a a c t a c c a t g c c c t g a t a t a t c t t t t g c a c c t g c t g a a c t t c a 2040
t t t c t g a t c a c c t g a c c t t g g a t g c c a a a a c g t t a t t c t g t t t g t c t g t t g t a g a a 2100
t t t t a g a t a a a g c t a t t a a t g g c a a t a t t t t t g c t a a a c g t t t t g t t t t t a c t g t c 2160
a c t a g g g c a a t a a a a t t t a t a c t c a a c c a t a t a a t a a c a t t t t t t a a c t a c t a a a g g a g t 2220
a g t t t t a t t t a a a g t c t t a g c a a t t t c t a t a c a a c t t t c t t a g a c t t a a c a c t t a t 2280
g a t a a a t g a c t a a a c t a g t a c a g a a t c t t t a t g a a a t a t g a c c t t t c t g a a a a t a c a t 2340
a c t t t t a e a t t t c t a c t t t a t t g a g a c c t a t t a g a t g t a a g t c t a g t a g a a t a t a g a t 2400
a a a a g a g g c t g a g a a t t a c c a t a c a a g g g t a t t a c a a c t g t a a a a c a a t t t a c t t t g t t 2460
t c a t t g t t c t g t c a a t a a t t g t t a c c a a a g a g a t a a a a a t a a a a g c a g a a t g t a t a t c a t 2520

```