

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 November 2006 (16.11.2006)

PCT

(10) International Publication Number
WO 2006/119569 A1

(51) International Patent Classification:
G01N 33/53 (2006.01) C12Q 1/68 (2006.01)

(74) Agent: FB RICE & CO; Level 23, 200 Queen Street,
Melbourne, Victoria 3000 (AU).

(21) International Application Number:
PCT/AU2006/000617

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 11 May 2006 (11.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/679,745 11 May 2005 (11.05.2005) US
60/689,745 9 June 2005 (09.06.2005) US
60/725,365 11 October 2005 (11.10.2005) US

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): GE-
NETIC TECHNOLOGIES LINTIED [AU/AU]; 60-66
Hanover Street, Fitzroy, Victoria 3065 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BOEHMER,
Ralph, Michael [DE/AU]; 17/31 York Street, St Kilda
West, Victoria 3182 (AU). ALLMAN, Richard [GB/AU];
2/32 Linacre Road, Hampton, Victoria 3188 (AU).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHODS OF ENRICHING FETAL CELLS

(57) Abstract: The present invention relates to methods of enriching fetal cells from a pregnant female. The present invention relates to removing, from a sample, cells that comprise at least one MHC molecule. The present invention also relates to methods that rely on using telomerase, mRNA encoding components thereof, as well as telomere length, as markers for fetal cells. Enriched fetal cells can be used in a variety of procedures including, detection of a trait of interest such as a disease trait, or a genetic predisposition thereto, gender typing and parentage testing.

WO 2006/119569 A1

METHODS OF ENRICHING FETAL CELLS

FIELD OF THE INVENTION

The present invention relates to methods of enriching fetal cells from a pregnant female. Enriched fetal cells can be used in a variety of procedures including, detection of a trait of interest such as a disease trait, or a genetic predisposition thereto, gender
5 typing and parentage testing.

BACKGROUND OF THE INVENTION

Fetal testing for chromosomal abnormalities is often performed on cells obtained using amniocentesis, or alternatively, Chorionic Villus Sampling (CVS).
10 Amniocentesis is a procedure used to retrieve fetal cells from the fluid that surrounds the fetus. This relatively invasive procedure is performed after the 12th week of pregnancy. There is about 0.5% increased risk of miscarriage following amniocentesis. CVS is a prenatal test in which cells surrounding an embryo are removed in order to examine the chromosomes. CVS is relatively less invasive, and can be performed as
15 early as 10 weeks from conception. There is about 1% increased risk of miscarriage following CVS.

Fetal therapy is in its very early stages and the possibility of very early tests for a wide range of disorders would undoubtedly greatly increase the pace of research in this area. Current fetal surgical techniques have improved, making fetal surgery for
20 some genetic problems like spina bifida and cleft palate very feasible. In addition, relatively simple effective fetal treatment is currently available for other disorders such as 21-hydroxylase deficiency (treatment with dexamethasone) and holocarboxylase synthetase (treatment with biotin) deficiencies, as long as detection can take place early enough.

At least some fetal cell types such as platelets, trophoblasts, erythrocytes and leucocytes have been shown to cross the placenta and circulate in maternal blood (Douglas *et al.*, 1959; Schroder, 1975). Maternal blood represents a non-invasive source of fetal cell types, however the isolation of fetal cells from maternal blood is hampered by the scarcity of such fetal cells in the maternal circulation, as well as the
30 lack of a marker that identifies all fetal cells, rather than merely a sub-population. A variety of methods have been proposed for isolation or enrichment of fetal cells in maternal blood. These methods include centrifugation techniques, immunoaffinity techniques, and fluorescent *in situ* hybridization (FISH) methods. However, these methods suffer from a number of deficiencies.

A fetal specific antibody is yet to be identified which can be used to reliably and reproducibly enrich fetal cells. This problem can be overcome with the method described by Simons (US 5,153,117 and US 5,447,842), based on a negative selection approach that does not require knowledge about fetal cell types and fetal cell numbers.
5 However, the Simons method is operationally difficult and expensive to perform, due to the need to HLA type the mother, as well as due to the fact that high-quality specific HLA antibodies are not commercially available.

There is a need in the art for new methods for the enrichment and identification of fetal cells.

10

SUMMARY OF THE INVENTION

It is generally considered that Class I Major Histocompatibility Complex (MHC) molecules (human Class I MHC molecules are also known in the art as Class I Human Leukocyte Antigens (HLA)) are expressed on most, if not all, nucleated cell
15 types. Notably, at least the Class I MHC molecules HLA-G and HLA-C have been found to be expressed on some types of fetal trophoblasts (Shorter *et al.*, 1993; King *et al.*, 1996). However, it has surprisingly been found that depleting a sample using an agent which binds MHC molecules results in an enriched population of fetal cells. Furthermore, it has been determined that telomerase and telomeres can be considered as
20 a marker of fetal cells. This enables these molecules to be targeted in procedures for detecting and isolating fetal cells. When combined together, these procedures enhance the purity of enriched fetal cell populations.

Accordingly, in a first aspect the present invention provides a method of enriching fetal cells from a sample, the method comprising

- 25 i) depleting maternal cells by removing cells that express at least one MHC molecule on their surface, and
ii) selecting fetal cells by
a) selecting cells that express telomerase, and/or
b) selecting cells based on telomere length.

30 Steps i) and ii) can be performed in any order. Thus, one step may be performed on the sample obtained from the mother, and the other step on the remaining cell population. Alternatively, the steps may be performed simultaneously.

In another aspect, the present invention provides a method of enriching fetal cells from a sample, the method comprising removing from the sample cells that
35 express at least one MHC molecule on their surface.

Preferably, the MHC molecule is a Class I MHC molecule.

In a further preferred embodiment, all cells expressing at least one Class I MHC molecule are removed.

In a particularly preferred embodiment, the Class I MHC molecule is HLA-A. In another preferred embodiment, the Class I MHC molecule is HLA-B. In a further preferred embodiment, the Class I MHC molecule is HLA-A and HLA-B.

An advantage of the above aspects of the invention when compared to that of Simons (US 5,153,117) is that it is not necessary to determine the genotype of MHC alleles of the mother, father and/or fetus. Thus, in a particularly preferred embodiment, the genotype of an MHC allele is not determined for the mother, father and/or fetus. More preferably, the genotype of an MHC allele is not determined for the mother.

In another embodiment, the method comprises

i) contacting cells in the sample with an agent that binds at least one MHC molecule, and

ii) removing cells bound by the agent.

In a further preferred embodiment, the method comprises contacting the sample with i) an agent that binds at least one Class I MHC molecule, and ii) an agent that binds at least one Class II MHC molecule.

In another preferred embodiment, the agent binds:

i) a monomorphic determinant of HLA-A molecules,

ii) a monomorphic determinant of HLA-B molecules, or

iii) a monomorphic determinant of HLA-A and HLA-B molecules.

In one embodiment, the agent does not bind HLA-C.

In another embodiment, the agent binds a monomorphic determinant of HLA-A, HLA-B and HLA-C molecules. Preferably, the agent that binds a monomorphic determinant of HLA-A, HLA-B and HLA-C molecules is used at sub-saturating concentrations.

In a further embodiment, more than two agents are used which bind different isotypes of the same class or sub-class of MHC molecule. Preferably, collectively the agents bind all isotypes (alleles) of the same class or sub-class of MHC molecule.

In one embodiment, the two agents are an antibody that binds HLA-Bw4 and an antibody that binds HLA-Bw6.

Compounds have been shown to associate *in situ* with MHC molecules, and hence these compounds can be targeted using the methods of the invention. Accordingly, in another embodiment, the method comprises

i) contacting cells in the sample with an agent that binds a compound that associates with an MHC molecule, and

ii) removing cells bound by the agent.

For example, the compound could be a ligand, for example a protein ligand, that binds an MHC molecule.

The binding of the agent to a maternal cell can be detected directly or indirectly.

5 Direct detection relies on the agent being bound to a detectable label or isolatable label. Indirect detection relies on a further factor, for example a detectably labelled secondary antibody, which binds the agent/maternal cell complex. Preferably, the label is selected from, but not limited to, the group consisting of: a fluorescent label, a radioactive label, a paramagnetic particle (such as a magnetic bead), a chemiluminescent label, a label
10 that is detectable by virtue of a secondary enzymatic reaction, and a label that is detectable by virtue of binding to a molecule.

Labelled cells can be removed from the sample using any technique known in the art. In one embodiment, the step of removing cells comprises detecting the label and removing the labeled cells.

15 In a further embodiment, the detectable label or isolatable label is a fluorescent label, wherein the step of removing cells comprises performing fluorescence activated cell sorting.

In another embodiment, the detectable label or isolatable label is a paramagnetic particle such as a magnetic bead, wherein the step of removing cells comprises
20 exposing the labelled cells to a magnetic field.

The agent can be any compound which specifically binds MHC expressed on the surface of a maternal cell. Typically, the agent will be an antibody or antibody fragment.

In another embodiment, the maternal cells bound by an antibody which binds an
25 MHC molecule are removed by killing the cells using complement-dependent lysis.

In another aspect, the present invention provides a method of enriching fetal cells from a sample, the method comprising selecting cells from the sample that express telomerase.

Telomerase is a protein/RNA complex. In one embodiment, the method
30 comprises detecting a protein component of telomerase. Preferably, the protein component is telomere reverse transcriptase (TERT). Examples of other proteins which may form part of the telomerase protein/RNA complex are: TEP-1 (telomerase associated protein-1) and 14-3-3 protein.

A protein component of telomerase can be detected using any technique known
35 in the art. Preferably, the cell is exposed to a polypeptide (more preferably, an antibody) which binds telomerases, especially TERT. Using an antibody as an

example, the antibody bound to telomerase may be detected directly or indirectly. Direct detection relies on the antibody being detectably labelled. Indirect detection relies on a further factor, for example a detectably labelled secondary antibody, which binds the anti-telomerase antibody/telomerase complex.

5 In another embodiment, the method comprises detecting an RNA component of telomerase. In yet another embodiment, the method comprises detecting an mRNA encoding a protein component of telomerase.

RNA/mRNA can be detected using any technique known in the art. Typically, the cells are exposed to a labelled probe which hybridizes to the RNA/mRNA. The
10 probe can be of any length or structure as long as it is capable of hybridizing the target RNA or mRNA.

Telomeres prior to birth can be considered to be at maximum length. After birth, with each cell division, they get progressively shorter. Telomeres generally remain until death, however, they just get shorter with time. It has been determined
15 that telomeres are attractive targets to use in identifying fetal cells, (1) because they provide an age-discriminant for cell selection, namely young cells can be separated from older cells (fetal from maternal), and (2) because probes can be designed with a relatively low coefficient of variation and good signal:noise ratio.

Thus, in yet another aspect, the present invention provides a method of
20 enriching fetal cells from a sample, the method comprising selecting cells from the sample based on telomere length.

In one embodiment, the method comprises contacting cells with a detectably labelled probe that binds telomeres.

In another embodiment, about 1 to about 100 cells, more preferably about 1 to
25 about 20 cells and even more preferably about 1 to about 10 cells, are selected, wherein the selected cells have been bound by more probe than the other cells in the sample. In this embodiment, a probe is used that will bind in approximate proportion (by number) to the length of the telomere. Thus, the selected cells are the most intensely labelled cells.

30 The sample can be obtained from any source known in the art to potentially contain fetal cells. Examples include, but are not limited to, blood, cervical mucous or urine. Preferably, the sample is maternal blood.

When the sample is maternal blood it is preferred that the method further
35 comprises isolating from the maternal blood sample a cell fraction comprising nucleated cells.

In some cases, particularly when performing procedures which detect RNA or DNA, it is preferred that the cells are fixed and permeabilized.

Fetal cell enrichment using the methods of the invention may be further enhanced by negatively selecting for cells that express at least one other maternal cell marker. As outlined above, this marker may be an MHC molecule. In an embodiment, 5 the method further comprises removing from the sample red blood cells, lymphocytes, and/or cancer cells. In a particularly preferred embodiment, the method further comprises removing hemopoietic cells from the sample. Preferably, the method further comprises contacting cells in the sample with an agent that binds a hemopoietic cell.

10 Examples of hemopoietic cells that can be removed include, but are not limited to, T cells, a B cells, macrophages, neutrophils, dendritic cells and/or basophils.

Preferably, the agent binds a cell surface protein of the cell. Such cell surface proteins are known to those skilled in the art. Examples of cell surface proteins include, but are not limited to, CD3, CD4, CD8, CD10, CD14, CD15, CD45 and CD56.

15 In a particularly preferred embodiment, the method further comprises contacting cells in the sample with an agent that binds CD45, and removing cells bound by the agent that binds CD45. Such embodiments can be performed using similar techniques to those described herein for depletion using an agent which binds at least one MHC molecule.

20 The methods of the invention can also be used in combination with further methods of positively selecting for fetal cells by targeting molecules expressed by fetal cells but not by (or only a small proportion of) maternal cells. Thus, in a further embodiment, the method further comprises contacting the cells with an agent that binds fetal cells, and selecting cells bound by the agent that binds fetal cells. Examples of 25 such markers include, but are not limited to, trophoblast specific proteins, fetal or embryonal hemoglobin, and fetal nucleated red blood cell specific proteins.

The sample can be obtained during any stage of pregnancy. If the sample is to be screened to determine if the fetus has a genetic defect, the detection of which may lead to the pregnancy being terminated, it is preferred that the sample is obtained from 30 the mother in the first trimester of pregnancy, preferably between week 8 and week 12.

The labelled fetal cells can be selected using any method known in the art. In many instances the procedure for selection is linked to the nature of the label. For example, where the label used emits a fluorescent signal the cells can be selected by, but not limited to, fluorescence activated cell sorting, fluorescence microscopy, or laser 35 microdissection.

In another aspect, the present invention provides a method of detecting a fetal cell(s) in a sample, the method comprising analysing a candidate cell for the expression of telomerase.

In a further aspect, the present invention provides a method of detecting a fetal cell(s) in a sample, the method comprising analysing a candidate cell for the presence of telomeres and/or analysing the length of the telomeres in a candidate cell.

In a further aspect, the present invention provides an enriched population of fetal cells obtained by a method according to the invention.

In another aspect, the present invention provides a composition comprising fetal cells of the invention, and a carrier.

In yet another aspect, the present invention provides for the use of an agent that binds at least one MHC molecule, and/or an agent that binds a compound that associates with an MHC molecule, for enriching fetal cells from a sample.

In another aspect, the present invention provides for the use of an agent that binds telomerase for enriching fetal cells from a sample.

In yet a further aspect, the present invention provides for the use of an agent that binds telomeres for enriching fetal cells from a sample.

Fetal cells enriched/detected using a method of the invention can be used to analyse the genotype of the fetus. Thus, in another aspect, the present invention provides a method for analysing the genotype of a fetal cell at a locus of interest, the method comprising

i) obtaining enriched fetal cells using a method according to the invention and/or detecting a fetal cell using a method of the invention, and

ii) analysing the genotype of at least one fetal cell at a locus of interest.

The genotype of the fetus can be determined using any technique known in the art. Examples include, but are not limited to, karyotyping, hybridization based procedures, and/or amplification based procedures.

The genotype of a fetal cell can be analysed for any purpose. Typically, the genotype will be analysed to detect the likelihood that the offspring will possess a trait of interest. Preferably, the fetal cell is analysed for a genetic abnormality linked to a disease state, or predisposition thereto. In one embodiment, the genetic abnormality is in the structure and/or number of chromosomes. In another embodiment, the genetic abnormality encodes an abnormal protein. In another embodiment, the genetic abnormality results in decreased or increased expression levels of a gene.

In at least some instances, the enrichment methods of the invention will not result in a pure fetal cell population. In other words, some maternal cells may remain.

Thus, in a preferred embodiment the methods of diagnosis (determination, analysis etc) further comprises identifying a cell as a fetal cell. This analysis may positively identify maternal or fetal cells. In the case of positively identifying maternal cells, the non-labelled cells will be fetal cells. Alternatively, both maternal and fetal cells are positively identified using different selectable markers, or a marker that results in a different level of signal between maternal and fetal cells is used. These procedures can be performed using any technique known in the art. For example, for male fetal cells a Y-chromosome specific probe can be used. In another example, telomere length is analysed. In a further embodiment, maternal cells are identified using an agent, such as an antibody, that binds a Class I MHC molecule. Other methods suitable to perform this embodiment are described herein.

The enriched/detected fetal cells can be used to determine the sex of the fetus. As a result, in a further aspect the present invention provides a method of determining the sex of a fetus, the method comprising

- i) obtaining enriched fetal cells using a method according to the invention and/or detecting a fetal cell using a method of the invention, and
- ii) analysing of at least one fetal cell to determine the sex of the fetus.

The analysis of the fetal cells to determine the sex of the fetus can be performed using any technique known in the art. For example, Y-chromosome specific probes can be used, and/or the cells karyotyped.

The enriched fetal cells can also be used to identify the father of the fetus. Accordingly, in a further aspect, the present invention provides a method of determining the father of a fetus, the method comprising

- i) obtaining enriched fetal cells using a method according to the invention and/or detecting a fetal cell using a method of the invention,
- ii) determining the genotype of the candidate father at one or more loci,
- iii) determining the genotype of the fetus at one or more of said loci, and
- iv) comparing the genotypes of ii) and iii) to determine the probability that the candidate father is the biological father of the fetus.

Whilst in some cases it may not be essential that the genotype of the mother also be analysed, for accuracy it is preferred that the method further comprises determining the genotype of the mother at one or more of said loci.

Analysis of the genotype of the candidate father, fetus or mother can be performed using any technique known in the art. One preferred technique is performing DNA fingerprinting analysis using probes/primers which hybridize to

tandemly repeated regions of the genome. Another technique is to analyse the HLA/MHC region of the genome.

In a further aspect, the present invention provides a kit for enriching fetal cells from a sample, the kit comprising

- 5 i) an agent that binds at least one MHC molecule, and/or an agent that binds a compound that associates with an MHC molecule, and/or an agent that binds a hemopoietic cell, and
- ii) a molecule which binds to telomerase, and/or which hybridizes to a polynucleotide encoding a protein component of said telomerase, and/or which
- 10 hybridizes to telomeres.

In yet another aspect, the present invention provides a kit for enriching fetal cells from a sample, the kit comprising an agent that binds at least one MHC molecule, and/or an agent that binds a compound that associates with an MHC molecule, and/or an agent that binds a hemopoietic cell.

- 15 Preferably, the agent that binds at least one MHC molecule is an antibody.

In another embodiment, the kit comprises

- i) an agent that binds all HLA-A molecules,
- ii) an agent that binds all HLA-B molecules, and/or
- iii) an agent that binds all HLA-A and HLA-B molecules.
- 20 Preferably, at least one agent is linked to a magnetic bead.

In yet another aspect, the present invention provides a kit for detecting a fetal cell, the kit comprising a molecule which binds to telomerase, and/or which hybridizes to a polynucleotide encoding a protein component of said telomerase, and/or which hybridizes to telomeres.

- 25 Preferably, the molecule is selected from the group consisting of; an anti-telomerase antibody, a polynucleotide which hybridizes to mRNA encoding a protein component of telomerase, a polynucleotide which hybridizes to an RNA component of telomerase, or a polynucleotide which hybridizes to telomeric DNA on the chromosome.

- 30 Preferably, the molecule is detectably labelled.

In a further aspect, the present invention provides a kit for detecting a genetic abnormality in a fetal cell, the kit comprising

- i) a molecule for detecting a fetal cell, wherein the molecule binds to telomerase, which hybridizes to a polynucleotide encoding a protein component of said telomerase,
- 35 or which hybridizes to telomeres, and
- ii) at least one reagent for detecting said genetic abnormality.

As will be apparent, preferred features and characteristics of one aspect of the invention are applicable to many other aspects of the invention.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated
5 element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

The invention is hereinafter described by way of the following non-limiting Examples and with reference to the accompanying figures.

10 **BRIEF DESCRIPTION OF THE ACCOMPANYING DRAWINGS**

Figure 1 - Shows a statistics of total numbers of male fetal cells in 10 ml blood samples. Only samples containing male cells are plotted. Fetal cell numbers range from just about 1 cell to more than 100 cells.

Figure 2 - Shows for HLA depletion, the dependence of fetal cell numbers on
15 gestational age.

Figure 3 - Shows fetal cell numbers together with total cell numbers found in the non-retained fraction of the magnetic column.

Figure 4 - Enrichment of fetal cells using combinations of an anti-HLA antibody and an anti-CD45 antibody.

20 **Figure 5** - Data used to produce Figure 4.

Figure 6 - Effect of auxiliary depletion with CD45 paramagnetic beads.

Figure 7 - Total maternal blood cell contamination after depletion with anti-HLA antibodies +/- CD45 antibodies.

Figure 8 - Comparison between different anti-HLA Class I antibodies.

25 **Figure 9** - Detection of male fetal cells using a RED Y-FISH probe.

Figure 10 - Selection of fetal cells using an anti-telomerase antibody.

KEY TO THE SEQUENCE LISTING

SEQ ID NO: 1 - Human telomerase reverse transcriptase (Genbank Accession No.
30 AAC51724).

SEQ ID NO:2 - mRNA encoding human telomerase reverse transcriptase (Genbank Accession No. NM_003219).

SEQ ID NO: 3 - RNA component of human telomerase (nucleotides 799 to 1248 of (Genbank Accession No. AF047386).

DETAILED DESCRIPTION OF THE INVENTION

General Techniques

Unless specifically defined otherwise, all technical and scientific terms used herein shall be taken to have the same meaning as commonly understood by one of
5 ordinary skill in the art (e.g., in cell culture, fetal cell biology, molecular genetics, immunology, immunohistochemistry, protein chemistry, nucleic acid hybridization, flow cytometry, and biochemistry).

Unless otherwise indicated, the recombinant protein, cell culture, and immunological techniques utilized in the present invention are standard procedures,
10 well known to those skilled in the art. Such techniques are described and explained throughout the literature in sources such as, J. Perbal, *A Practical Guide to Molecular Cloning*, John Wiley and Sons (1984), J. Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbour Laboratory Press (1989), T.A. Brown (editor), *Essential Molecular Biology: A Practical Approach*, Volumes 1 and 2, IRL
15 Press (1991), D.M. Glover and B.D. Hames (editors), *DNA Cloning: A Practical Approach*, Volumes 1-4, IRL Press (1995 and 1996), and F.M. Ausubel *et al.* (editors), *Current Protocols in Molecular Biology*, Greene Pub. Associates and Wiley-Interscience (1988, including all updates until present), Ed Harlow and David Lane (editors) *Antibodies: A Laboratory Manual*, Cold Spring Harbour Laboratory, (1988),
20 and J.E. Coligan *et al.* (editors) *Current Protocols in Immunology*, John Wiley & Sons (including all updates until present), and are incorporated herein by reference.

Major Histocompatibility Complex

The major histocompatibility complex (MHC) includes at least three classes of
25 genes. Class I and II genes encode antigens expressed on cell surface, whilst class III genes encode several components of the complement system. Classes I and II antigens are glycoproteins that present peptides to T lymphocytes. Human MHC molecules are also known in the art as Human Leukocyte Antigens (HLA). Thus, the terms "HLA" and "MHC" are often used interchangeably herein.

30 Human and murine class I molecules are heterodimers, consisting of a heavy alpha chain (45kD) and a light chain, beta-2-globulin (12kD). Class I molecules are found on most, if not all, nucleated cells. The alpha chain can be divided into three extracellular domains, alpha1, alpha2 and alpha3, in addition to the transmembranous and cytoplasmic domains. The alpha3 domain is highly conserved, as is beta-2-
35 microglobulin. Both alpha3 domain and beta-2-microglobulin are homologous to the CH3 domain of human immunoglobulin.

Class II molecules are heterodimeric glycoproteins, alpha chain (34kD) and beta chain (29kD). Each chain has 2 extracellular domains, together with the transmembranous and cytoplasmic domains. The membrane-proximal alpha2 and beta2 domains are homologous to immunoglobulin CH domain. Class II molecules are
5 less commonly expressed when compared to Class I, typically being found in dendritic cells, B lymphocytes, macrophages, and a few other cell types.

There are 3 class I loci (B,C,A) in the short arm of human chromosome 6, and 4 loci (K, D(L), Qa, Tla) in murine chromosome 17. These loci are highly polymorphic. The variable residues are clustered in 7 subsequences, 3 in alpha1 domain and 4 in
10 alpha2 domain. There are 3 major human class II loci (HLA-DR, HLA-DO, HLA-DP) and 2 murine loci (H-2I-A, H-2I-E). All class II beta chains are polymorphic. Human HLA-DQ alpha chain is also polymorphic.

Preferably, at least some methods of the invention utilize an agent (preferably an antibody) which binds at least one MHC molecule. Preferably, the agent binds an
15 extracellular portion of the MHC molecule. This has at least two advantages, i) the method of the invention can be used to enrich live fetal cells, and ii) an additional step of ensuring that the agent passes through the cell membrane (for example having to fix and permeabilize the cell) is not required.

Preferably, the agent is capable of binding at least one Class I HLA molecule.
20 In one embodiment, the agent is capable of binding HLA-A, HLA-B and HLA-C molecules. In a preferred embodiment, the agent is capable of binding HLA-A and/or HLA-B molecules. In a further embodiment, at least two different agents can be used that bind the same or different Classes or sub-classes of MHC molecules.

As used herein, a "monomorphic determinant" refers to a region of a group
25 proteins that is highly conserved between at least 90%, more preferably at least 95%, more preferably at least 99%, and even more preferably 100% of the group which can be recognised by a suitable binding agent such as an antibody. The region can be a continuous stretch of amino acids, and/or a group of highly conserved amino acids that, upon protein folding, are closely associated. For example, a "monomorphic
30 determinant" of a Class I MHC molecule is a region of the proteins (isotypes) encoded by different alleles of Class I MHC genes that is highly conserved between the different proteins of the Class and that can be bound by the same antibody.

As used herein, a "sub-class" of a MHC molecule is a distinct type of MHC molecules of a particular Class. For example, HLA-A molecules and HLA-B
35 molecules are each considered herein as a sub-class of Class I MHC molecules.

Telomeres and Telomerases

Telomeres consist of DNA-protein complexes that are located at the ends of eukaryotic chromosomes and function to provide protection against genome instability promoting events such as degradation of the terminal regions of chromosomes, fusion
5 of a telomere with another telomere or broken DNA end, or inappropriate recombination. Telomeres prior to birth can be considered to be at maximum length. After birth, with each cell division, they get progressively shorter (Vaziri *et al.*, 1994). Telomeric DNA comprises tandem repeats of DNA, in humans the 6-base pair sequence TTAGGG, that form a molecular scaffold containing binding sites for
10 telomeric proteins, resulting in a dynamic DNA-protein complex at the telomere.

Telomerase is an enzyme concerned with the formation, maintenance, and renovation of telomeres at the ends of chromosomes. Telomerase acts as an RNA-dependent DNA polymerase that synthesizes telomeric DNA sequences and consists of
15 two essential components; the first being the functional RNA component (in humans also known as hTR - see SEQ ID NO:3) and the other being the catalytic protein (in humans also known as hTERT - see SEQ ID NO:1). Hence, telomerase is a ribonucleoprotein. Telomerase regulates the proliferative capacity of cells. Telomerase is now classed as a tumour-associated antigen. It may also play a role in the clonal expansion of lymphocytes in response to viral infection.

20 In biochemical terms, telomerase acts as a telomerase reverse transcriptase (TERT). It transcribes RNA into DNA and is the reverse-transcribing enzyme specific to the telomeric sequence. It has two unique features: it is able to recognize a single-stranded (G-rich) telomere primer and it is able to add multiple telomeric repeats to its end by using its RNA moiety as a template.

25 The correlation between telomerase activity, telomere lengths, and cellular replicative capacity has led to the theory that maintenance of telomere lengths by telomerase acts as a molecular clock to control replicative capacity and senescence.

The RNA components of human and other telomerases have been cloned and characterized (WO 96/01835). However, the characterization of all the protein
30 components of telomerase has been difficult. Despite this, a number of proteins that may interact with TERT have been identified and include TEP-1 (telomerase associated protein 1) (Harrington *et al.*, 1997) and 14-3-3 proteins (Seimiya *et al.*, 2000).

As used herein, the term "telomerase" refers to at the least the ribonucleoprotein comprising the functional RNA component and the reverse transcriptase. However, at
35 least in some instances this term may also encompass other proteins which may form part of the telomerase complex such as the TEP-1 and 14-3-3 proteins.

Agent

The present invention relies on the use of various agents which bind molecules expressed by maternal or fetal cells. These agents can be of any structure or composition as long as they are capable binding to a target molecule. In one embodiment, the agents useful for the present invention are proteins. Preferably, the protein is an antibody or fragment thereof.

In an embodiment, it is preferred that an agent is used that binds at least one MHC molecule, and that this agent is an anti-MHC antibody. Preferably, the antibody binds an extracellular portion of the MHC molecule. In another embodiment, the antibody binds specifically to a protein component of telomerase, preferably the reverse transcriptase.

Antibodies useful for the methods of the invention can be monoclonal or polyclonal antibodies. Antibodies useful for the methods of the invention can readily be produced using techniques known in the art. Alternatively, at least some anti-MHC antibodies can be obtained from commercial sources such as US Biological (Massachusetts, USA) and Chemicon International Inc. (California, USA). Furthermore, at least some anti-telomerase antibodies can be obtained from commercial sources such as Abcam Ltd (Cambridge, UK) and Calbiochem (California, USA).

The term "binds specifically" refers to the ability of the antibody to bind to a target ligand (such as telomerase or an MHC molecule) but not other proteins in the sample.

If polyclonal antibodies are desired, a selected mammal (e.g., mouse, rabbit, goat, horse, etc.) is immunised with a suitable immunogenic polypeptide (for example, the extracellular domain of HLA-A can be used when an anti-MHC antibody is desired, or a protein comprising the sequence provided in SEQ ID NO:1 when an anti-telomerase antibody is required). Serum from the immunised animal is collected and treated according to known procedures. If serum containing polyclonal antibodies contains antibodies to other antigens, the polyclonal antibodies can be purified by immunoaffinity chromatography. Techniques for producing and processing polyclonal antisera are known in the art.

Monoclonal antibodies can also be readily produced by one skilled in the art. The general methodology for making monoclonal antibodies by hybridomas is well known. Immortal antibody-producing cell lines can be created by cell fusion, and also by other techniques such as direct transformation of B lymphocytes with oncogenic

DNA, or transfection with Epstein-Barr virus. Panels of monoclonal antibodies produced can be screened for various properties; i.e., for isotype and epitope affinity.

An alternative technique involves screening phage display libraries where, for example the phage express single chain antibodies (scFv) fragments on the surface of
5 their coat with a large variety of complementarity determining regions (CDRs). This technique is well known in the art.

For the purposes of this invention, the term "antibody", unless specified to the contrary, includes fragments of whole antibodies which retain their binding activity for a target antigen. Such fragments include Fv, F(ab') and F(ab')₂ fragments, as well as
10 scFv. Furthermore, the antibodies and fragments thereof may be humanised antibodies, for example as described in EP-A-239400.

Preferably, agents used in the methods of the present invention are bound to a detectable label or isolatable label. Alternatively, the agent is not directly labelled but detected using indirect methods such as using a detectably labelled secondary antibody
15 which specifically binds the agent.

The terms "detectable" and "isolatable" label are generally used herein interchangeably. Some labels useful for the methods of the invention cannot readily be visualized (detectable) but nonetheless can be used to enrich (isolate) fetal cells (for example a paramagnetic particle).

Exemplary labels that allow for direct measurement of antibody binding include
20 radiolabels, fluorophores, dyes, magnetic beads, chemiluminescers, colloidal particles, and the like. Examples of labels which permit indirect measurement of binding include enzymes where the substrate may provide for a coloured or fluorescent product. Additional exemplary labels include covalently bound enzymes capable of providing a
25 detectable product signal after addition of suitable substrate. Examples of suitable enzymes for use in conjugates include horseradish peroxidase, alkaline phosphatase, malate dehydrogenase and the like. Where not commercially available, such antibody-enzyme conjugates are readily produced by techniques known to those skilled in the art. Further exemplary detectable labels include biotin, which binds with high affinity
30 to avidin or streptavidin; fluorochromes (e.g., phycobiliproteins, phycoerythrin and allophycocyanins; fluorescein and Texas red), which can be used with a fluorescence activated cell sorter; haptens; and the like.

Examples of fluorophores which can be used to label antibodies includes, but are not limited to, Fluorescein Isothiocyanate (FITC), Tetramethyl Rhodamine
35 Isothiocyanate (TRITC), R-Phycoerythrin (R-PE), AlexaTM, Dyes, Pacific BlueTM, Allophycocyanin (APC), and PerCPTM.

The label may also be a quantum dot. In the context of antibody labelling they are used in exactly the same way as fluorescent dyes. Quantum Dots are developed and marketed by several companies, including, Quantum Dot Corporation (USA) and Evident Technologies (USA). Examples of antibodies labelled with quantum dots are
5 described in Michalet *et al.* (2005) and Tokumasu and Dvorak (2003).

As noted above, in some embodiments the agent is not directly labelled. In this instance, cells are identified using another factor, typically a detectably labeled secondary antibody. The use of detectably labeled secondary antibodies in methods of detecting a marker of interest are well known in the art. For example, if an anti-MHC
10 antibody or anti-telomerase antibody was produced from a rabbit, the secondary antibody could be an anti-rabbit antibody produced from a mouse.

As used herein, the term "sub-saturating concentrations" of an agent such as an antibody means that the number of molecules of the agent is less, preferably significantly less, than the number of target molecules (for example MHC Class I
15 molecules) in a sample. Thus, in this situation only a small fraction of target antigens per cell get an agent bound to them. For example, in some embodiments the ratio of agent to target is less than 1:10, 1:100, 1:1000, or 1:10000. Sub-saturating concentrations of an agent can readily be determined by the skilled person using standard techniques.

20 Maternal cells bound by an antibody can be killed, and thus depleted from a sample, by complement-dependent lysis. For example, antibody labelled cells can be incubated with rabbit complement at 37°C for 2 hr. Commercial sources for suitable complement systems include Calbiochem, Equitech-Bio and Pel Freez Biologicals. Suitable anti-MHC antibodies for use in complement-dependent lysis are known in the
25 art, for example the W6/32 antibody mentioned in the Examples can be used for this procedure.

Labelling of Fetal Cells using a Probe which binds the RNA Component of Telomerase, the mRNA encoding a Protein Component of Telomerase, or Telomeres

30 A probe from use in a method of the invention will typically be DNA, RNA or a mixture thereof. However, the probe may comprise modifications which are usually designed to reduce the likelihood of degradation. Such modifications are typically the use of nucleotide analogs and/or altered linker groups. Nucleic acid analogs which can be used in probes of the invention include phosphoramidate, phosphorothioate,
35 phosphorodithioate, O-methylphosphoroamidite linkages, and peptide nucleic acid backbones and linkages. Other analog nucleic acids include those with positive

backbones, non-ionic backbones, and non-ribose backbones. Probes containing one or more carbocyclic sugars are also useful in the methods of the invention.

Preferably a probe used in the methods of the invention is at least 15 nucleotides in length, more preferably at least 20 nucleotides in length, more preferably at least 25 nucleotides in length, more preferably at least 50 nucleotides in length, and even more preferably at least 100 nucleotides in length.

In one embodiment, the probe is capable of hybridizing to a mRNA encoding human TERT (SEQ ID NO:2) or the RNA component of human telomerase (SEQ ID NO:3). The probes of these embodiment are of sufficient length and specificity that there is little, if any, background hybridization to non-target DNA or RNA in the cells of the sample being analysed. Such probes can readily be designed by the skilled person.

In another embodiment, the probe hybridizes to telomeres. As outlined above, human telomeres are repeats of TTAGGG. Thus, probes useful for this embodiment of the invention comprise multiple repeats of this sequence, or the reverse complement thereof. Typically, probes which hybridize telomeres are reasonably long, being at least 1kb, at least 5kb, at least 20kb, at least 50kb, at least 100kb, or at least 200kb in length. Whilst non-fetal cells will also comprise telomeres, fetal cells can still be detected by selecting cells which produce a greater signal upon hybridization with the telomere probe.

Particularly preferred are peptide nucleic acid (PNA) probes which includes peptide nucleic acid analogs. These backbones are substantially non-ionic under neutral conditions, in contrast to the highly charged phosphodiester backbone of naturally occurring nucleic acids. The PNA backbone exhibits improved hybridization kinetics. PNAs have larger changes in the melting temperature (T_m) for mismatched versus perfectly matched basepairs. DNA and RNA typically exhibit a 2-4°C drop in T_m for an internal mismatch. With the non-ionic PNA backbone, the drop is closer to 7-9. Similarly, due to their non-ionic nature, hybridization of the bases attached to these backbones is relatively insensitive to salt concentration. In addition, PNAs are not degraded by cellular enzymes, and thus can be more stable.

Probes can contain any detection moiety that facilitates the detection of the probe when hybridized to a target nucleic acid sequence (either genomic DNA, mRNA or the RNA component of telomerase). Effective detection moieties include both direct and indirect labels as described below.

Probes can be directly labeled with a detectable label. Examples of detectable labels include, but are not limited to, a fluorescent or chemiluminescent compound,

such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme (e.g., as commonly used in an ELISA), biotin, digoxigenin, and radioactive isotopes, e.g., ^{32}P , and ^3H . The detectable label may also be a quantum dot. Fluorophores can be directly labeled following covalent attachment to a nucleotide by incorporating the labeled
5 nucleotide into the probe with standard techniques such as nick translation, random priming, and PCR labeling. Alternatively, nucleotides within the probe can be transaminated with a linker. The fluorophore can then be covalently attached to the transaminated nucleotides. Useful probe labeling techniques are described in Molecular Cytogenetics: Protocols and Applications, Y.-S. Fan, Ed., Chap. 2,
10 "Labeling Fluorescence *In Situ* Hybridization Probes for Genomic Targets", L. Morrison et. al., p. 21-40, Humana Press, 2002, incorporated herein by reference.

Examples of fluorophores that can be used in the methods described herein include, but are not limited to, 7-amino-4-methylcoumarin-3-acetic acid (AMCA), Texas RedTM (Molecular Probes, Inc., Eugene, Oreg.); 5-(and-6)-carboxy-X-
15 rhodamine, lissamine rhodamine B, 5-(and-6)-carboxyfluorescein; fluorescein-5-isothiocyanate (FITC); 7-diethylaminocoumarin-3-carboxylic acid, tetramethylrhodamine-5-(and-6)-isothiocyanate; 5-(and-6)-
carboxytetramethylrhodamine; 7-hydroxycoumarin-3-carboxylic acid; 6-[fluorescein 5-(and-6)-carboxamido]hexanoic acid; N-(4,4-difluoro-5,7-dimethyl-4-bora-3a, 4a diaza-
20 3-indacenepropionic acid; cosin-5-isothiocyanate; erythrosine-5-isothiocyanate; 5-(and-6)-carboxyrhodamine 6G; and CascadeTM blue acetylazide (Molecular Probes, Inc., Eugene, Oreg.).

When multiple probes are used, fluorophores of different colours can be chosen such that each probe in a set can be distinctly visualized. For example, activated
25 maternal lymphocytes could be distinguished from fetal cells using such a multiple probe approach.

Probes labeled with a fluorescent moiety can be viewed with a fluorescence microscope and an appropriate filter for each fluorophore, or by using dual or triple band-pass filter sets to observe multiple fluorophores. Any suitable microscopic
30 imaging method can be used to visualize the hybridized probes, including automated digital imaging systems, such as those available from MetaSystems or Applied Imaging. Alternatively, techniques such as flow cytometry can also be used to examine the hybridization pattern of the probes.

Probes can also be labeled indirectly, e.g., with biotin or digoxigenin by means
35 well known in the art. However, secondary detection molecules or further processing are then required to visualize the labeled probes. For example, a probe labeled with

biotin can be detected by avidin conjugated to a detectable marker, e.g., a fluorophore. Additionally, avidin can be conjugated to an enzymatic marker such as alkaline phosphatase or horseradish peroxidase. Such enzymatic markers can be detected in standard calorimetric reactions using a substrate for the enzyme. Substrates for alkaline phosphatase include 5-bromo-4-chloro-3-indolylphosphate and nitro blue tetrazolium. Diaminobenzoate can be used as a substrate for horseradish peroxidase.

Digoxigenin PNA probes are available commercially for flow cytometric measurement of telomere length by DAKO Cytomation. Digoxigenin conjugated hybridisations may be detected using anti-digoxigenin fluorescently labelled antibodies. Digoxigenin containing nucleic acid probes can also be produced using a Dig-RNA labelling kit (Roche).

With regard to the detection of telomere length using, for example, a fluorescently labelled PNA probe, a preferred embodiment of the invention is selecting cells that are the most brightly labelled. For instance, in an embodiment fetal cells will typically have a about 1.3 to about 1.5 greater signal than maternal cells. Flow cytometry can be used to measure telomere length (for example, as described by Schmid *et al.*, 2002; Baerlocher *et al.*, 2002; Baerlocher *et al.*, 2003; Cabuy *et al.*, 2004), with analysis algorithms such as those described by De Pauw *et al.* (1998) and Narath *et al.* (2005) being suitable to distinguish the more highly labelled fetal cells from the less labelled maternal cells.

Labelled Fetal Cell Detection and Isolation

As used herein, the terms "enriching" and "enriched" are used in their broadest sense to encompass the isolation of the fetal cells such that the relative concentration of fetal cells to non-fetal cells in the treated sample is greater than a comparable untreated sample. Preferably, the enriched fetal cells are separated from at least 10%, more preferably at least 20%, more preferably at least 30%, more preferably at least 40%, more preferably at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 90%, more preferably at least 95%, and even more preferably at least 99% of the non-fetal cells in the sample obtained from the mother. Most preferably, the enriched cell population contains no maternal cells (namely, pure). The terms "enrich" and variations thereof are used interchangeably herein with the term "isolate" and variations thereof. Furthermore, a population of cells enriched using a method of the invention may only comprise a single fetal cell. In addition, the enrichment methods of the invention may be used to isolate a single fetal cell.

Maternal cells expressing at least one type of MHC molecule can be depleted from the sample, by a variety of techniques well known in the art, including cell sorting, especially fluorescence-activated cell sorting (FACS), by using an affinity reagent bound to a substrate (e.g., a plastic surface, as in panning), or by using an
5 affinity reagent bound to a solid phase particle which can be isolated on the basis of the properties of the beads (e.g., colored latex beads or magnetic particles). These same procedures can be used to enrich for cells using telomerase, and/or telomere length, as a marker. Naturally, the procedure used to remove the maternal cells will depend upon how the cells have been labelled.

10 For removal of maternal cells by cell sorting, the cells are labeled directly or indirectly with a substance which can be detected by a cell sorter, preferably a dye. Preferably, the dye is a fluorescent dye. A large number of different dyes are known in the art, including fluorescein, rhodamine, Texas red, phycoerythrin, and the like. Any detectable substance which has the appropriate characteristics for the cell sorter may be
15 used (e.g., in the case of a fluorescent dye, a dye which can be excited by the sorter's light source, and an emission spectra which can be detected by the cell sorter's detectors). Again, similar techniques can be used to enrich cells using telomerase, and/or telomere length, as a marker.

In flow cytometry, a beam of laser light is projected through a liquid stream that
20 contains cells, or other particles, which when struck by the focussed light give out signals which are picked up by detectors. These signals are then converted for computer storage and data analysis, and can provide information about various cellular properties. Cells labelled with a suitable dye are excited by the laser beam, and emit light at characteristic wavelengths. This emitted light is picked up by detectors, and
25 these analogue signals are converted to digital signals, allowing for their storage, analysis and display.

Many larger flow cytometers are also "cell sorters", such as fluorescence-activated cell sorters (FACS), and are instruments which have the ability to selectively deposit cells from particular populations into tubes, or other collection vessels. In a
30 particularly preferred embodiment, the cells are isolated using FACS. This procedure is well known in the art and described by, for example, Melamed, *et al.* (1990) *Flow Cytometry and Sorting* Wiley-Liss, Inc., New York, N.Y.; Shapiro (2003) *Practical Flow Cytometry*, 4 ed, Wiley-Liss, Hoboken, NJ.; and Robinson, *et al.* (1993) *Handbook of Flow Cytometry Methods* Wiley-Liss, New York, N.Y.

35 In order to sort cells, the instruments electronics interprets the signals collected for each cell as it is interrogated by the laser beam and compares the signal with sorting

criteria set on the computer. If the cell meets the required criteria, an electrical charge is applied to the liquid stream which is being accurately broken into droplets containing the cells. This charge is applied to the stream at the precise moment the cell of interest is about to break off from the stream, then removed when the charged droplet has
5 broken from the stream. As the droplets fall, they pass between two metal plates, which are strongly positively or negatively charged. Charged droplets get drawn towards the metal plate of the opposite polarity, and deposited in the collection vessel, or onto a microscope slide, for further examination.

The cells can automatically be deposited in collection vessels as single cells or
10 as a plurality of cells, e.g. using a laser, e.g. an argon laser (488 nm) and for example with a Flow Cytometer fitted with an Autoclone unit (Coulter EPICS Altra, Beckman-Coulter, Miami, Fla., USA). Other examples of suitable FACS machines useful for the methods of the invention include, but are not limited to, MoFlo™ High-speed cell sorter (Dako-Cytomation Ltd), FACS Aria™ (Becton Dickinson), ALTRA™ Hyper sort
15 (Beckman Coulter) and CyFlow™ sorting system (Partec GmbH).

For removal of maternal cells from a sample using solid-phase particles, any particle with the desired properties may be utilized. For example, large particles (e.g., greater than about 90-100 µm in diameter) may be used to facilitate sedimentation. Preferably, the particles are "magnetic particles" (i.e., particles which can be collected
20 using a magnetic field). Typically, maternal cells labelled with the magnetic probe are passed through a column, held within a magnetic field. Labelled cells are retained in the column (held by the magnetic field), whilst unlabelled cells pass straight through and are eluted at the other end. Magnetic particles are now commonly available from a variety of manufacturers including Dynal Biotech (Oslo, Norway) and Milteni Biotech
25 GmbH (Germany). An example of magnetic cell sorting (MACS) is provided by Al-Mufti *et al.* (1999). Yet again, similar techniques can be used to enrich cells using telomerase, and/or telomere length, as a marker.

Laser-capture microdissection can also be used to selectively remove labelled maternal cells on a slide using methods of the invention. Methods of using laser-
30 capture microdissection are known in the art (see, for example, U.S. 20030227611 and Bauer *et al.*, 2002).

As the skilled person will appreciate, maternal cells can be labelled with one type of label, and fetal cells with another type of label, and the respective cells types identified and/or depleted/selected on the basis of the different labelling. For example,
35 maternal cells can be labelled as described herein such that they produce a fluorescent

green signal, and maternal cells can be labelled as described herein such that they produce a fluorescent red signal.

Following enrichment, the cells can be cultured *in vitro* to expand fetal cells numbers using techniques known in the art. For example culturing in RPMI 1640
5 media (Gibco).

Sample and Preparation of Cells

As used herein, the term "sample" refers to material taken directly from the pregnant female (such as blood), as well as such material that has already been partially
10 purified. Examples of such partial purification include the removal of at least some non-cellular material, removal of maternal red blood cells, and/or removal of maternal lymphocytes. Thus, the term "sample" is used herein broadly to include a sample obtained after depletion of maternal cells using, for example, an anti-MHC antibody, but before selection based on the expression of telomerase or telomere length (or *vice*
15 *versa*). In some embodiments, the cells in the sample are cultured *in vitro* before a method of the invention is performed.

The methods of the invention can be performed on any pregnant female of any species, wherein the genome of the species comprises a major histocompatibility complex and/or fetal cells of the organism produce telomerase. Preferably, the female
20 is a mammal. Preferred mammals include, but are not limited to, humans, livestock animals such as sheep, cattle and horses, as well as companion animals such as cats and dogs.

In a preferred embodiment, the sample comprising fetal cells is obtained from a pregnant woman in her first trimester of pregnancy. In one embodiment the sample can
25 be a blood sample which is prevented from clotting such as a sample containing heparin or, preferably, ACD solution. The sample is preferably stored at 0 to 4°C until use to minimize the number of dead cells, cell debris and cell clumps. The number of fetal cells in the sample varies depending on factors including the age of the fetus. Typically, from 7 to 20 ml of maternal blood provides sufficient fetal cells upon
30 separation from maternal cells. Preferably, 30 ml or more blood is drawn to ensure sufficient cells without the need to draw an additional sample.

In another embodiment, the fetal cells are obtained from the cervical mucous of the mother as, for example, generally described in WO 03/020986, WO 2004/076653 or WO 2005/047532.

35 In a preferred embodiment, red blood cells are removed from a sample comprising, or derived from, maternal blood. Red blood cells can be removed using

any technique known in the art. Red blood cells (erythrocytes) may be depleted by, for example, density gradient centrifugation over Percoll, Ficoll, or other suitable gradients. Red blood cells may also be depleted by selective lysis using commercially available lysing solutions (eg, FACSlyse™, Becton Dickinson), Ammonium Chloride
5 based lysing solutions or other osmotic lysing agents.

Fetal nucleated red cells, if potentially present in the sample, can be protected from ammonium chloride lysis by acetazolamide (Orskoff lysis).

The purity of recovered fetal cells may be increased by depleting the sample of
10 maternal cells using auxiliary agents which bind maternal cell markers other than MHC molecules. The essential feature for choosing such markers for this purpose is that they are not expressed on at least the majority of fetal cells. This auxiliary depletion is performed before, during or after the steps of the invention. Those skilled in the art are aware that the types of nucleated maternal cells in maternal blood include B cells, T cells, monocytes, macrophages dendritic cells and stem cells, each characterised by a
15 specific set of surface markers that can be targeted for depletion. Preferably, the maternal cell population or maternal cells are further depleted by exposing a maternal sample or a nucleated cellular fraction thereof to an antibody that binds to a cellular marker on the maternal cell for a time and under conditions sufficient to form an antibody-maternal cell complex and isolating the antibody-maternal cell complex. As
20 with other embodiments described herein, the antibody-maternal cell complex is preferably isolated by contacting said complex with a readily detectable and/or a readily isolatable label. Examples of non-MHC molecules which can be targeted to possibly further deplete the sample of maternal cells include, but are not limited to, CD3, CD4, CD8, CD10, CD14, CD15, CD45, CD56 and proteins described by
25 Blaschitz *et al.* (2000). Such further maternal cell specific agents can readily be used in combination with an agent that binds at least one MHC molecule. For example, magnetic beads can be produced which have both anti-MHC and anti-CD45 antibodies attached thereto.

It has been shown that telomerase activity can be detected in cancerous cells
30 (see, for example, Satyanarayana *et al.*, 2004). Thus, when selecting cells for the presence of telomerase or telomere length it is preferred that the sample does not comprise cancerous cells. Such cells can be avoided by screening the individual for cancer before the method of the invention is performed. Such screening can be performed by any method known in the art including analysing the patient, or a sample
35 therefrom, for cancer markers. As the skilled person would be aware, such cancer markers could also be used in methods of removing cancer cells from the sample.

A cancer marker is a molecule which has been shown to be expressed, and/or overexpressed, by a cancer cell. Examples of cancer markers include, but are not limited to, CA 15-3 (marker for numerous cancers including breast cancer), CA 19-9 (marker for numerous cancers including pancreatic cancer and biliary tract tumours),
5 CA 125 (marker for various cancers including ovarian cancer), calcitonin (marker for various tumours including thyroid medullary carcinoma), catecholamines and metabolites (phaeochromoctoma), CEA (marker for various cancers including colorectal cancers and other gastrointestinal cancers), epithelial growth factor (EGF) and/or epithelial growth factor receptor (EGFR) (both associated with colon cancer),
10 A33 colonic epithelial antigen (colon cancer), hCG/beta hCG (marker for various cancers including germ-cell tumours and choriocarcinomas), 5HIAA in urine (carcinoid syndrome), PSA (prostate cancer), serotonin (carcinoid syndrome) NY-ESO-1 (marker of oesophageal cancer), thyroglobulin (thyroid carcinoma), and the CT antigens such as MAGE (associated with many liver cancers and melanomas), GAGE
15 (hepatocarcinoma), SSX2 (sarcoma) differentiation antigens (such as Melan A/MART1, GP100 and tyrosinase), mutational antigens (such as CDK4, β -catenin), amplification antigens (such as P53 and Her2), and splice variant antigens (such as ING1).

A number of researches have identified that telomerase activity in lymphocytes
20 can lead to false positives when investigating whether a patient has cancer (see, for example, Kavalier *et al.*, 1998; Matthews *et al.*, 2001; Seki *et al.*, 2001; Sidransky, 2002; Trulsson *et al.*, 2003). Accordingly, when selecting cells for the presence of telomerase or telomere length, at least in some circumstances it will be useful to avoid such cells in the sample, and/or take measures to differentially label lymphocytes.
25 Furthermore, when selecting cells for the presence of telomerase or telomere length, it may be useful to ensure the pregnant female does not have an infection which may lead to elevated levels of activated lymphocytes. Lymphocytes can be removed and/or labelled using any technique known in the art. For example, Seki *et al.* (2001) removed peripheral blood lymphocytes by Ficoll-Isopaque gradient centrifugation before
30 performing a telomerase assay to detect cancer cells. A similar procedure could be used in the present instance. In another example, the cells are pre-sorted by targeting cell surface markers on lymphocytes with a suitable antibody and separating the bound cell. Alternatively, the antibody specific for the lymphocytes could be labeled with a different label than that used to detect the fetal cells, allowing for the two cell types to
35 differentiated as maternal lymphocytes will be doubly labelled whereas the fetal cells will only be labelled with, for example, an antibody which binds telomerase. There are

a number of lymphocyte markers which could be used to avoid the false detection of maternal lymphocytes. Suitable T-cell markers include, but are not limited to, CD2, CD3, CD4, CD5, CD6, CD7, CD8, CD25, CD 56, CD94 and CD158a. Suitable B-cell markers include, but are not limited to, CD19 and CD20.

5 The methods of the invention may include the step of fixing and permeabilizing the cells in the sample. Such procedures are known to those skilled in the art. For example, fixation may involve initial paraformaldehyde fixation followed by treatment with detergents such as Saponin, TWEEN-based detergents, Triton X-100, Nonidet NP40, NP40 substitutes, or other membrane disrupting detergents. Permeabilization
10 may also involve treatment with alcohols (ethanol or methanol). Initial fixation may also be in ethanol. Combined fixation/permeabilization may also be performed using commercially available kits, including DAKO-Intrastain™, Caltag's Fix & Perm reagents, Ortho Diagnostic's Permeafix™.

 In other embodiments, such as when electroporation or quantum dots are used to
15 deliver a detectably labelled anti-telomerase antibody to the cells, it is not necessary to fix and permeabilize the cells. As a result, in some embodiments, the methods of the invention can detect and/or isolate live cells. Such isolated live cells could be cultured *in vitro* to expand fetal cells numbers using techniques known in the art. For example culturing in RPMI 1640 media (Gibco).

20 Methods for using electroporation to deliver a labelled antibody to a live cell are known in the art (see, for example, Berglund and Starkey, 1989).

Additional Procedures for the Positive Selection of Fetal Cells

 The methods of the invention can include the additional step of positively
25 selecting fetal cells beyond selection based on telomerases or telomere length. Such positive selection relies on targeting molecules produced by fetal cells but not by (or only a small proportion of) the remaining maternal cells. As the skilled person will appreciate, the procedures described above for removing maternal cells expressing at least one MHC molecule are readily adapted for the positive selection of fetal cells
30 expressing a particular cell marker.

 For example, fetal cells are selected using cytokeratin-7, a marker on virtually all trophoblast types. Another marker that covers many types of fetal trophoblasts is HLA-G. Further trophoblast-specific antibodies are commercially available, although none of them covers all types of trophoblasts.

35 In a further example, fetal/embryonic hemoglobin can be used as a marker for fetal nucleated red cells.

Depending on fetal cell types present, such markers can be combined.

Uses

Enriched fetal cells comprise the same genetic DNA make up of the somatic
5 cells of the fetus, and hence fetal cells isolated using the methods of the invention can
be analysed for traits of interest and/or abnormalities using techniques known in the art.
Such analysis can be performed on any cellular material that enables the trait, or
predisposition thereto, to be detected. Preferably, this material is nuclear DNA,
however, at least in some instances it may be informative to analyse RNA or protein
10 from the isolated fetal cells. Furthermore, the DNA may encode a gene, or may encode
a functional RNA which is not translated, or the DNA analysed may even be an
informative non-transcribed sequence or marker.

In one preferred embodiment, chromosomal abnormalities are detected. By
"chromosomal abnormality" we include any gross abnormality in a chromosome or the
15 number of chromosomes. For example, this includes detecting trisomy in chromosome
21 which is indicative of Down's syndrome, trisomy 18, trisomy 13, sex chromosomal
abnormalities such as Klinefelter syndrome (47, XXY), XYY or Turner's syndrome,
chromosome translocations and deletions, a small proportion of Down's syndrome
patients have translocation and chromosomal deletion syndromes include Pradar-Willi
20 syndrome and Angelman syndrome, both of which involve deletions of part of
chromosome 15, and the detection of mutations (such as deletions, insertions,
transitions, transversions and other mutations) in individual genes. Other types of
chromosomal problems also exist such as Fragile X syndrome, hemophilia, spinal
muscular dystrophy, myotonic dystrophy, Menkes disease and neurofibromatosis, which
25 can be detected by DNA analysis.

The phrase "genetic abnormality" also refers to a single nucleotide substitution,
deletion, insertion, micro-deletion, micro-insertion, short deletion, short insertion,
multinucleotide substitution, and abnormal DNA methylation and loss of imprint
(LOI). Such a genetic abnormality can be related to an inherited genetic disease such
30 as a single-gene disorder (e.g., cystic fibrosis, Canavan, Tay-Sachs disease, Gaucher
disease, Familial Dysautonomia, Niemann-Pick disease, Fanconi anemia, Ataxia
telaugiostasia, Bloom syndrome, Familial Mediterranean fever (FMF), X-linked
spondyloepiphyseal dysplasia tarda, factor XI), an imprinting disorder [e.g., Angelman
Syndrome, Prader-Willi Syndrome, Beckwith-Wiedemann syndrome, Myoclonus-
35 dystonia syndrome (MDS)], or to predisposition to various diseases (e.g., mutations in
the BRCA1 and BRCA2 genes). Other genetic disorders which can be detected by

DNA analysis are known such as thalassaemia, Duchenne muscular dystrophy, connexin 26, congenital adrenal hypoplasia, X-linked hydrocephalus, ornithine transcarbamylase deficiency, Huntington's disease, mitochondrial disorder, mucopolysaccharidosis I or IV, Norrie's disease, Rett syndrome, Smith-Lemli Optiz
5 syndrome, 21-hydroxylase deficiency or holocarboxylase synthetase deficiency, diastrophic displasia, galactosialidosis, gangliosidosis, hereditary sensory neuropathy, hypogammaglobulinaemia, hypophosphatasia, Leigh's syndrome, aspartylglucosaminuria, metachromatic leukodystrophy Wilson's disease, steroid sulfatase deficiency, X-linked adrenoleukodystrophy, phosphorylase kinase deficiency
10 (Type VI glycogen storage disease) and debranching enzyme deficiency (Type III glycogen storage disease). These and other genetic diseases are mentioned in *The Metabolic and Molecular Basis of Inherited Disease*, 8th Edition, Volumes I, II, III and IV, Scriver, C. R. *et al.* (eds), McGraw Hill, 2001. Clearly, any genetic disease where the gene has been cloned and mutations detected can be analysed.

15 The methods of the present invention can also be used to determine the sex of the fetus. For example, staining of the isolated fetal cells with a Y-chromosome specific marker will indicate that the fetus is male, whereas the lack of staining will indicate that the fetus is female.

In yet another use of the invention, the methods described herein can be used for
20 paternity testing. Where the paternity of a child is disputed, the procedures of the invention enable this issue to be resolved early on during pregnancy. Many procedures have been described for parentage testing which rely on the analysis of suitable polymorphic markers. As used herein, the phrase "polymorphic markers" refers to any nucleic acid change (e.g., substitution, deletion, insertion, inversion), variable number
25 of tandem repeats (VNTR), short tandem repeats (STR), minisatellite variant repeats (MVR) and the like. Typically, parentage testing involves DNA fingerprinting targeting informative repeat regions, or the analysis of highly polymorphic regions of the genome such as HLA loci.

30 Analysis of Fetal Cells

Fetal cells enriched/detected using the methods of the invention can be analysed by a variety of procedures, however, typically genetic assays will be performed. Genetic assay methods include the standard techniques of karyotyping, analysis of methylation patterns, restriction fragment length polymorphism assays, sequencing and
35 PCR-based assays, as well as other methods described below.

Chromosomal abnormalities, either in structure or number, can be detected by karyotyping which is well known in the art. Karyotyping analysis is generally performed on cells which have been arrested during mitosis by the addition of a mitotic spindle inhibitor such as colchicine. Preferably, a Giemsa-stained chromosome spread
5 is prepared, allowing analysis of chromosome number as well as detection of chromosomal translocations.

The genetic assays may involve any suitable method for identifying mutations or polymorphisms, such as: sequencing of the DNA at one or more of the relevant positions; differential hybridisation of an oligonucleotide probe designed to hybridise at
10 the relevant positions of either the wild-type or mutant sequence; denaturing gel electrophoresis following digestion with an appropriate restriction enzyme, preferably following amplification of the relevant DNA regions; S1 nuclease sequence analysis; non-denaturing gel electrophoresis, preferably following amplification of the relevant DNA regions; conventional RFLP (restriction fragment length polymorphism) assays;
15 selective DNA amplification using oligonucleotides which are matched for the wild-type sequence and unmatched for the mutant sequence or vice versa; or the selective introduction of a restriction site using a PCR (or similar) primer matched for the wild-type or mutant genotype, followed by a restriction digest. The assay may be indirect, ie capable of detecting a mutation at another position or gene which is known to be linked
20 to one or more of the mutant positions. The probes and primers may be fragments of DNA isolated from nature or may be synthetic.

A non-denaturing gel may be used to detect differing lengths of fragments resulting from digestion with an appropriate restriction enzyme. The DNA is usually amplified before digestion, for example using the polymerase chain reaction (PCR)
25 method and modifications thereof.

Amplification of DNA may be achieved by the established PCR methods or by developments thereof or alternatives such as the ligase chain reaction, QB replicase and nucleic acid sequence-based amplification.

An "appropriate restriction enzyme" is one which will recognise and cut the
30 wild-type sequence and not the mutated sequence or *vice versa*. The sequence which is recognised and cut by the restriction enzyme (or not, as the case may be) can be present as a consequence of the mutation or it can be introduced into the normal or mutant allele using mismatched oligonucleotides in the PCR reaction. It is convenient if the enzyme cuts DNA only infrequently, in other words if it recognises a sequence which
35 occurs only rarely.

In another method, a pair of PCR primers are used which hybridise to either the wild-type genotype or the mutant genotype but not both. Whether amplified DNA is produced will then indicate the wild-type or mutant genotype (and hence phenotype).

A preferable method employs similar PCR primers but, as well as hybridising to
5 only one of the wild-type or mutant sequences, they introduce a restriction site which is not otherwise there in either the wild-type or mutant sequences.

In order to facilitate subsequent cloning of amplified sequences, primers may have restriction enzyme sites appended to their 5' ends. Thus, all nucleotides of the primers are derived from the gene sequence of interest or sequences adjacent to that
10 gene except the few nucleotides necessary to form a restriction enzyme site. Such enzymes and sites are well known in the art. The primers themselves can be synthesized using techniques which are well known in the art. Generally, the primers can be made using synthesizing machines which are commercially available.

PCR techniques that utilize fluorescent dyes may also be used to detect genetic
15 defects in DNA from fetal cells isolated by the methods of the invention. These include, but are not limited to, the following five techniques.

i) Fluorescent dyes can be used to detect specific PCR amplified double stranded DNA product (e.g. ethidium bromide, or SYBR Green I).

ii) The 5' nuclease (TaqMan) assay can be used which utilizes a specially
20 constructed primer whose fluorescence is quenched until it is released by the nuclease activity of the Taq DNA polymerase during extension of the PCR product.

iii) Assays based on Molecular Beacon technology can be used which rely on a specially constructed oligonucleotide that when self-hybridized quenches fluorescence (fluorescent dye and quencher molecule are adjacent). Upon hybridization to a specific
25 amplified PCR product, fluorescence is increased due to separation of the quencher from the fluorescent molecule.

iv) Assays based on Amplifluor (Intergen) technology can be used which utilize specially prepared primers, where again fluorescence is quenched due to self-hybridization. In this case, fluorescence is released during PCR amplification by
30 extension through the primer sequence, which results in the separation of fluorescent and quencher molecules.

v) Assays that rely on an increase in fluorescence resonance energy transfer can be used which utilize two specially designed adjacent primers, which have different fluorochromes on their ends. When these primers anneal to a specific PCR amplified
35 product, the two fluorochromes are brought together. The excitation of one fluorochrome results in an increase in fluorescence of the other fluorochrome.

If required, methods for the extraction of DNA from fixed samples for genetic analysis are also known to those skilled in the art. For example, US patent application 20040126796 discloses a method for the extraction of DNA from tissues and other samples, such as formalin-fixed tissue. The isolation of DNA from fixed samples for use in PCR has also been described by Lehman and Kreipe (2001) and Fitzgerald *et al.* (1993).

Fetal cells, or an enriched cell population of fetal cells, obtained using a method of the invention can be placed into wells of a microtitre plate (one cell per well) and analysed independently. Preferably, each cell not only screened for a trait(s) of interest, but screened to confirm/detect that the cell in a particular well is a fetal cell. In this instance, multiplex analysis can be performed as generally described by Finlay *et al.* (1996, 1998 and 2001).

Kits

The present invention also provides a kits for enriching fetal cells from a sample. In one example, the kit comprises i) an agent that binds at least one MHC molecule, an agent that binds a compound that associates with an MHC molecule, and/or an agent that binds a hemopoietic cell, and ii) a molecule which binds to telomerase, and/or which hybridizes to a polynucleotide encoding a protein component of said telomerase, and/or which hybridizes to telomeres. Other examples are described herein.

In one embodiment, a kit of the present invention includes, a single agent in an amount sufficient for at least one enrichment and/or detection procedure. Kits containing multiple agents are also contemplated by the present invention. The multiple agents may bind different MHC molecules of the same Class, and/or bind unrelated molecules (such as one agent that binds a monomorphic determinant of HLA-A molecules and another agent that binds CD45). Such agents may be bound to detectable or isolatable labels. For ease of use, multiple agents are typically bound to the same detectable or isolatable label.

In one embodiment, the agent(s) are each linked to magnetic beads. Different agents may be linked to different beads such that a single type of bead comprises different types of agents, or beads may be produced that only comprises a single type of agent and these beads mixed with other beads that have linked thereto a single type, but different, agent.

The kit may further comprise components for analysing the genotype of a fetal cell, determining the father of a fetus, and/or determining the sex of the fetus.

Typically, the kits will also include instructions recorded in a tangible form (e.g., contained on paper or an electronic medium), for example, for using a packaged agent for enriching fetal cells from a sample. The instructions will typically indicate the reagents and/or concentrations of reagents and at least one enrichment method
5 parameter which might be, for example, the relative amounts of agents to use per amount of sample. In addition, such specifics as maintenance, time periods, temperature and buffer conditions may also be included.

EXAMPLES

10 **EXAMPLE 1 - Enrichment of fetal cells using mouse anti-human HLA Class 1 antigen antibodies**

Materials and Methods

Blood

Blood samples were obtained from a private abortion clinic and the Royal
15 Children's Hospital (RCH) (Melbourne, Australia). Sample collection was anonymous, with donors de-identified. Whilst samples from the abortion clinic were specified to be pre-abortive, it was later determined that some of the samples that yielded higher numbers of fetal cells had been obtained post-abortion.

Blood samples (8-16 ml) were drawn into vacuum collection tubes with EDTA
20 as anti-coagulant. The samples were processed either fresh or after overnight storage at 4°C.

Magnetic cell separation

Mononuclear cells were isolated by density gradient (Ficoll 1.077)
25 centrifugation, and the entire samples were magnetically labelled with either of the following three procedures:

1. Cells were exposed to saturating amounts of a biotinylated antibody against a HLA Class 1 epitope common to all HLA-A, B and C (US Biological; Cat # H6098-39F2; Mouse anti-Human HLA Class 1 Antigen ABC /Biotin; IgG2a;
30 Clone 3H2211). Cells were then washed and labelled with saturating amounts of streptavidin-coated paramagnetic particles (Molecular Probes / Invitrogen; Cat# C-21476 "Captive").
2. Cells were exposed to saturating amounts of paramagnetic particles coated with antibodies to CD45 (Miltenyi Cat# 130-090-872; concentrated "whole-blood"
35 anti-human CD45 beads).

3. Procedures 1 and 2 were combined: Cells were first labelled to the HLA antibody, followed with a simultaneous exposure to Captivate and CD45 magnetic beads.

Magnetically labelled cell samples were passed through a magnetised column 5 (Miltenyi, LS columns Cat# 130-042-401), retaining all labelled cells. The non-adhered as well as the adhered fractions were collected, pelleted and frozen at -80°C until further use.

Detection of fetal cells by quantitative PCR (Q-PCR)

10 DNA was prepared using a commercial kit (Qiagen Cat# 51204 "FlexiGene DNA kit), and Q-PCR was performed on a real-time PCR machine (StrataGene MX3000), targeting a Y-chromosome-specific multi-copy sequence. Parallel reactions targeting a gender-unspecific sequence were performed to quantitate the total amount of DNA in the sample. All PCR reagents were from Qiagen. By comparison with 15 standards derived from known amounts of pure male-derived DNA, the total numbers of male (=fetal) cells and all cells in the magnetically separated preparations were calculated.

Results and Discussion

20 Among 30 samples processed with HLA Class 1 cell depletion, 14 samples contained male cells. Since about half of all fetuses are female, a nearly 50% detection rate of male cells indicates that fetal cells are retrieved in nearly every maternal sample.

Among 9 samples processed with CD45 cell depletion, 5 samples contained 25 male cells. Again, this is a approximately 50% detection rate and indicates that fetal cells are always retrieved.

Provided in Figure 1 are statistics of total numbers of male fetal cells in 10 ml blood samples. Only samples containing male cells are plotted. Fetal cell numbers range from just about 1 cell to more than 100 cells.

30 Figure 2 shows, for HLA depletion, the dependence of fetal cell numbers on gestational age.

Figure 3 provides fetal cell numbers together with total cell numbers found in the non-retained fraction of the magnetic column. The numbers vary from less than 1000 to about 100,000. With an approximate 10 million cells in the starting population of mononuclear cells, this is a dramatic enrichment. Also shown are the controls in 35 which 1% of the retained cell fraction was examined for fetal cells. The presence of

some occasional fetal cells in 1% of the controls indicates that not all fetal cells are retrieved with these procedures, but that some are CD45 + and HLA Cl.1 +.

EXAMPLE 2 - Enrichment of fetal cells using mouse anti-human HLA Class 1 antigen ABC clone 39-F2 or clone W6/32, both in combination with an anti-CD45 antibody

Unless stated to the contrary the procedures used were the same as those described above for Example 1.

Blood at different gestational ages was subjected to gradient centrifugation, removing erythrocytes, then labelled with either of two clones (39-F2 and W6/32) of biotinylated monoclonal antibody, each directed against a different monomorphic determinant of HLA-A,B,C. [US Biological, USA]. Subsequently, the cells were labelled simultaneously with streptavidin-coated paramagnetic beads ("Captive", Molecular Probes, USA) and paramagnetic beads coated with an antibody against the CD45 antigen [Miltenyi, Germany].

The labelled cells were passed through a magnetic column [Miltenyi], and the non-attached cells were subjected to quantitative PCR, targeting a Y-chromosome-specific sequence.

Figures 4 and 5 show total fetal (male) cell numbers per 10 ml of blood, plotted as a function of gestational age (GA). Nearly half of all blood samples (with unknown fetal gender) yielded a Y-signal. Only the positive samples (with at least one male cell) are shown.

These results complement those provided in Example 1. They provide the following additional information:

- a. The fetal cell enrichment achieved by this method does not depend on a special HLA-ABC epitope targeted by one particular HLA-ABC antibody clone, but can be achieved with different antibodies targeting different epitopes on the Class 1 antigens.
- b. An enhanced enrichment of fetal cells is obtained using a combination of antibodies that bind MHC molecules and hemopoietic cells (in this case an anti-CD45 antibody) (Figure 5).
- c. Similar numbers of fetal cells are found at gestational ages 7 and 8 weeks, and even as early as week 6 there are fetal cells to be found.

EXAMPLE 3 - Further studies showing enrichment of fetal cells using mouse anti-human HLA Class 1 antigen ABC antibodies

Materials and Methods

Blood

5 Blood of pregnant women was obtained from a private abortion clinic and the Royal Children's Hospital (RCH) (Melbourne, Australia). Maternal blood samples were drawn in steady-state pregnancy, prior to any testing or abortive procedure that could release fetal cells into the maternal circulation. For use as a model system, other samples were drawn during or after termination of pregnancy (post-termination
10 samples) to provide blood samples with increased numbers of fetal cells due to fetal hemorrhage. Sample collection was anonymous, with donors de-identified.

Blood samples (8-16 ml) were drawn into vacuum collection tubes with EDTA as anti-coagulant. The samples were processed either fresh or after overnight storage at 4°C.

15

Magnetic cell separation

Mononuclear cells were isolated by density gradient (Ficoll 1.083) centrifugation, and the entire samples were magnetically labelled with either of the following three procedures:

20 1. Cells were exposed to saturating amounts of one of the following biotinylated antibodies against a HLA Class 1 epitope common to all HLA-A, B and C:

- a. US Biological; Cat # H6098-39F2; Mouse anti-Human HLA Class 1 Antigen ABC ; (Data code: F2)
- 25 b. USBiological; Cat# H6098-60B; Mouse anti-Human HLA Class 1 Antigen ABC (Data code: 60B)
- c. EBioscience; Cat# 13-9983-82; Mouse anti-Human HLA Class 1 Antigen ABC; clone W6/32 (Data code: W6) ,or

30 antibodies against epitopes on HLA-B locus: one Lambda; mouse anti-human Bw4 (cat # BIH0007; mouse anti-human IgG2a); mouse anti-human Bw6 (cat #BIH0038; mouse anti-human IgG3) (Data code Bw4/6).

35 Cells were then washed and labelled with saturating amounts of streptavidin-coated paramagnetic particles (Molecular Probes / Invitrogen; Cat# C-21476 "Captive").

2. Cells were exposed to saturating amounts of paramagnetic particles coated with antibodies to CD45 (Miltenyi Cat# 130-090-872; concentrated "whole-blood" anti-human CD45 beads).
3. Procedures 1 and 2 were combined: Cells were first labelled to the HLA antibody, followed with a simultaneous exposure to Captivate and CD45 magnetic beads.

Magnetically labelled cell samples were passed through a magnetised column (Miltenyi, LS columns Cat# 130-042-401), retaining all labelled cells. The non-retained as well as the retained fractions were collected, pelleted and frozen at -80°C until further use.

Detection of fetal cells by quantitative PCR (Q-PCR)

DNA was prepared using a commercial kit (Qiagen Cat# 51204 "FlexiGene DNA kit), and Q-PCR was performed on a real-time PCR machine (StrataGene MX3000), targeting a Y-chromosome-specific multi-copy sequence. Parallel reactions targeting a gender-unspecific sequence were performed to quantitate the total amount of DNA in the sample. All PCR reagents were from Qiagen. By comparison with standards derived from known amounts of pure male-derived DNA, the total numbers of male (=fetal) cells and all cells in the magnetically separated preparations were estimated. At low fetal cell numbers (<10), this method was found to under-estimate fetal cell numbers.

Results and Discussion

Table 1 shows the fetal cell detection rates in steady-state maternal blood samples collected between week 7 and 14 of pregnancy. 101 samples were processed with HLA Class 1 and CD45 cell depletion. As many as 43 samples produced a clear Y-chromosome-specific signal, indicating that they contained at least 1 fetal cell. Since about half of all fetuses are female, a nearly 50% detection rate of male cells indicates that fetal cells are retrieved in nearly every maternal sample. Considering that the PCR method was found to under-estimate fetal cell numbers in the range from 1-10, we suggest that the true fetal cell recovery is higher than the detection rate, probably 100%.

Table 1: Detection of fetal cells in steady-state, first trimester maternal blood.

GA	samples	Y signal	
7	17	7	
8	46	17	
9	9	6	
10	17	6	
12	7	5	
11	2	1	
13	2	0	
14	1	1	
total	101	43	42.60%

The data indicates that fetal cells can be found as early as 7 weeks GA, a result that appears to be a dramatic improvement over any other published results.

Similar to Example 2, Figure 6 shows the effect of the auxiliary use of CD45 depletion in addition to cell depletion with HLA Class I antibody, using post-termination blood samples, which serve as a model system with increased numbers of fetal cells. Nucleated blood cells were incubated with biotinylated antibody to HLA Class I antigen (Bw4+6), followed by incubation with streptavidin ferrofluid. Half of the sample was simultaneously incubated with paramagnetic beads binding to CD45 antigen, the other half served as control. Total numbers of remaining cells, as well as the numbers of male cells, were determined by Q-PCR. The ratios of cell numbers after HLA+CD45 depletion were divided by cell numbers after only HLA depletion and are shown as % of control. The plot of ALL vs. Y values for each sample (insert graph of Figure 6) shows the lack of correlation between the two values. The graph shows that the auxiliary depletion by CD45 beads reduced the total remaining cell numbers to 1 percent of controls (HLA depletion only), while the numbers of fetal cells are only reduced by about 50%.

In a further experiment, maternal blood (10-12 ml) was processed by density gradient (1.083), nucleated cells were labelled with anti-HLA Bw4+Bw6 / biotin, then depleted with streptavidin ferrofluid +/- anti-CD45 paramagnetic beads. Total remaining cell numbers (mostly maternal, of course) were determined by Q-PCR. The results are provided in Figure 7 with median values and approximate range being shown. The data show that depletion with HLA+CD45 results in total cell numbers of

about 100 cells on average, which implies that average fetal cell purity is > 1%, whenever any number of fetal cells are present (Figure 7). These relatively few contaminating maternal cells and resulting high fetal cell purity will enable positive fetal cell markers to positively identify fetal cells by microscopy or single-cell PCR techniques.

Figure 8 provides a comparison between different HLA-Class I antibodies with respect to fetal cell recovery and total cell depletion. Three of the antibodies (F2, 60B and W6/32) are directed to 3 different epitopes common to all HLA-A, B and C antigens. Bw4/6 is a mixture of specific antibodies to Bw4 and Bw6. A person is Bw4, Bw6 or both, so that the combination of both ensures antibody binding for each blood donor. The data show that there is little difference between the different antibodies, which implies a wide choice of commercially available antibodies for this method.

Example 4 - Detection of Human Telomerase Reverse Transcriptase protein (hTERT) by monoclonal or polyclonal antibodies

Two protocols are provided below for the detection of human telomerase reverse transcriptase protein (hTERT) by monoclonal or polyclonal antibodies. As the skilled person would be aware, many of individual procedures described below are interchangeable between the two protocols.

Protocol 1

Cells from blood of a pregnant female are separated from plasma by centrifugation. Red cells are depleted on Percoll density gradients. Cells are fixed and permeabilized using a commercial kit – DAKO-Intrastain. The cells are washed again in PBS, and then incubated with monoclonal anti-telomerase antibody (Abcam Ltd, Cambridge, UK) for 1 hour at room temperature.

The cells are then washed in PBS (150 mM NaCl, 10 mM phosphate buffer) containing 0.5% bovine serum albumin (BSA), and a Fluorescein Isothiocyanate (FITC) fluorescently labelled secondary antibody which binds the monoclonal antibody is added for 1 hour at room temperature. Cells are washed in PBS containing 0.5% BSA.

Cells are analysed and labelled cells separated using fluorescence activated cell sorting on a MoFlo High-speed cell sorter (Dako-Cytomation, Ltd).

Protocol 2

Cells from blood of a pregnant female are separated from plasma by centrifugation. Red cells are depleted by selective lysis using Becton Dickinson FACSLyse solution. Cells are fixed in paraformaldehyde (about 1.5%) for 24 hours at 5 4°C. Cells are washed in PBS and permeabilised using 0.05% Triton X-100 in PBS for 30 min at room temp.

The cells are washed again in PBS, and then incubated with a polyclonal antisera comprising anti-telomerase antibodies (Calbiochem, California, USA) which are labelled with magnetic beads (DynaL Biotech) for 1 hour at room temperature.

10 The cells are then washed in PBS containing 0.5% bovine serum albumin (BSA), and analysed and labelled cells separated using magnetic activated cell sorting.

Example 5 - Detection of hTERT mRNA by hybridisation

Cells from blood of a pregnant female are separated from plasma by 15 centrifugation. Red cells are depleted on 70% Percoll density gradients. Cells are fixed and permeabilized using a commercial kit – Caltag Fix & Perm.

The cell suspension is centrifuged (1000g, 5 min), and the cells resuspended in 500 µl ice-cold methanol and incubated for 10 min at 4°C. The cells are centrifuged at 1000g for 5 min, and resuspended in 500 µl 0.2% Triton X-100/TE buffer (TE = 20 Tris/EDTA buffer (10mM Tris / 1mM EDTA pH 7.2). The cells are centrifuged again at 1000g for 5 min, and the supernatant carefully removed. Cells are washed once in 500 µl TE and centrifuged at 1000g for 5 min. Cells are resuspended in 5 µl of TE (avoiding bubbles).

20 µl of riboprobe comprising fluorescein-UTP is added in hybridization buffer 25 (50% Formamide, 10 mM Tris (pH 7.0), 5 mM EDTA, 10% Dextran Sulphate, 1 µg/µl tRNA). The riboprobe has a sequence which is complementary to the mRNA encoding hTERT, and is produced using techniques known in the art (Sambrook *et al.*, *supra*). The hybridization proceeds for 12 hours at 45°C. Cells are washed with 2 x SSC buffer, and pelleted at 1000g for 5 min. As much supernatant as possible is removed, and the 30 cells resuspended in 200 µl 2x SSC/ 0.3% NP40.

The cells are incubated at 37°C for 30 min. The cells are then centrifuged at 1000g 5 min, and the supernatant carefully removed. The cells are then resuspended in 200 µl 2x SSC/ 0.3% NP40 and incubated at room temp for 30 min. The cells are then centrifuged at 1000g 5 min, and the supernatant carefully removed. The cells are then 35 resuspended in 2.5 µl of TE.

Cells are analysed and labelled cells separated using fluorescence activated cell sorting on a MoFlo High-speed cell sorter (Dako-Cytomation, Ltd).

Example 6 - Determination of Telomere length with a PNA hybridisation probe

5 Cells from blood are separated from plasma by centrifugation. Red cells are depleted on Percoll density gradients. Cells are fixed and permeabilized using a commercial kit – Caltag Fix & Perm. 100 µl of the resulting fixed cells are placed in an eppendorf tube and centrifuged (1000g, 5 min).

10 Cells are resuspended in 500 µl ice-cold methanol and incubated for 10 min at 4°C, and the centrifuged at 1000g for 5 min. The cells are resuspended in 500 µl 0.2% Triton X-100/TE buffer, centrifuge at 1000g for 5 min, and then the supernatant carefully removed.

15 Cells are washed once in 500 µl TE and centrifuged at 1000g for 5 min. Cells are resuspended in 5 µl of TE (avoiding bubbles). 20 µl of PNA (Dako Telomere PNA kit/FITC, Dako-Cytomation) in hybridization buffer is added and co-denatured at 80°C for 20 min in thermocycler.

20 Hybridization is allowed to proceed for 12 hours at 37°C. The cells are then washed with 2 x SSC buffer, and pelleted at 1000g for 5 min. As much supernatant as possible is removed, and the cells resuspended in 200 µl 2x SSC/ 0.3% NP40. The cells are incubated at 37°C for 30 min, centrifuged at 1000g for 5 min, and as much supernatant as possible removed. The cells are then resuspended in 200 µl 2x SSC/ 0.3% NP40, and incubated at room temp for 30 min. The cells are then centrifuged at 1000g for 5 min. As much supernatant as possible is removed and the cells resuspended in 2.5 µl TE.

25 Cells are analysed and labelled cells separated using fluorescence activated cell sorting on a MoFlo High-speed cell sorter (Dako-Cytomation, Ltd).

Example 7 - Labelling fetal cells using anti-telomerase antibody

30 10 ml samples of peripheral blood were obtained by venupuncture from female volunteers during the first trimester of pregnancy.

Red cells were depleted by density gradient centrifugation over a gradient of 70% Percoll. The collected cells were washed in PBS containing 5% BSA and then fixed overnight in 2% paraformaldehyde at 4°C.

35 Cells were washed in PBS then permeabilized using 0.05% Triton X-100 (in PBS) for 30 min at 4°C. Cells were washed once in PBS and resuspended in 500 µl PBS. Anti-Telomerase polyclonal antibody (Abcam Ltd, Cambridge, UK) (10 µg

contained in 100 µl H₂O) was added to the cells and incubated at 4°C for 4 hours. Cells were washed twice in PBS and resuspended in 500 µl PBS. 100 µl Goat-anti-rabbit IgG-FITC was added and the cells incubated for 1 hour at 4°C. Cells were washed twice in PBS and resuspended in 5 ml PBS.

5 Cells were then analysed and sorted using a Dako-Cytomation MoFlo high speed cell sorter. Sort gates were set on cells expressing the top 5% of fluorescence values for this initial experiment.

Male fetal cells are labelled with RED (Spectrum Orange™) Y-FISH probe (Vysis, USA) and Green (Spectrum Green™) X-FISH probe (Vysis, USA). Male fetal
10 cells are those which express 1 Red and 1 Green FISH signal.

As can be seen from Figure 9, male fetal cells were double stained for X and Y-chromosome markers showing that anti-telomerase antibodies can be used to isolate fetal cells from maternal blood.

In a further experiment, whole blood samples (10 ml) were depleted of
15 erythrocytes by density gradient centrifugation, labeled with anti-telomerase polyclonal antibody and analysed by FACS. Cells in the region encompassing the top 5% of fluorescence intensities were sorted (Figure 10) and assessed for fetal cell content by fluorescence *in situ* hybridization.

A study of 10 samples gave male fetal cells in 50% of cases with fetal cell
20 numbers ranging from 1 – 10. These figures are similar to those obtained using the HLA negative selection approach described in Example 3.

Example 8 - Combined depletion of maternal cells expressing MHC and selection of cells based on expression of telomerase and/or telomere length

25 Two protocols are provided below, however, as the skilled person would be aware many of individual procedures described below are interchangeable between the two protocols. Considering the present disclosure, other protocols can readily be devised.

30 Protocol 1

Cells from blood of a pregnant female are separated from plasma by centrifugation. Red cells are depleted on Percoll density gradients. Cells are fixed and permeabilized using a commercial kit – DAKO-Intrastain. The cells are washed again in PBS, and then incubated with monoclonal anti-telomerase antibody (Abcam Ltd,
35 Cambridge, UK) for 1 hour at room temperature.

The cells are then washed in PBS (150 mM NaCl, 10 mM phosphate buffer) containing 0.5% bovine serum albumin (BSA), and a Fluorescein Isothiocyanate (FITC) fluorescently labelled secondary antibody which binds the monoclonal antibody is added for 1 hour at room temperature. Cells are washed in PBS containing 0.5%
5 BSA.

Cells are analysed and labelled cells separated using fluorescence activated cell sorting on a MoFlo High-speed cell sorter (Dako-Cytomation, Ltd).

Following the above positive selection, the enriched fetal cell population is depleted for at least some of the remaining maternal cells expressing MHC molecules.
10 To achieve this end, cells are exposed to saturating amounts of the following biotinylated antibodies against a HLA Class 1 epitope common to all HLA-A, B and C:

- a. US Biological; Cat # H6098-39F2; Mouse anti-Human HLA Class 1 Antigen ABC ; (Data code: F2)
- b. USBiological; Cat# H6098-60B; Mouse anti-Human HLA Class 1 Antigen
15 ABC (Data code: 60B), and
- c. EBioscience; Cat# 13-9983-82; Mouse anti-Human HLA Class 1 Antigen ABC; clone W6/32 (Data code: W6).

Cells are then washed and labelled with saturating amounts of streptavidin-coated paramagnetic particles (Molecular Probes / Invitrogen; Cat# C-21476
20 "Captive"). Magnetically labelled cell samples are passed through a magnetised column (Miltenyi, LS columns Cat# 130-042-401), retaining all labelled cells. Cells passing through the column include the further enriched fetal cell population and are collected for further analysis.

Analysis to confirm the presence of fetal cells may be by Fluorescence in situ
25 hybridisation or by quantitative PCR

Protocol 2

Maternal blood samples (8-16 ml) are drawn into vacuum collection tubes with EDTA as anti-coagulant. The samples are processed either fresh or after overnight
30 storage at 4°C.

Mononuclear cells are isolated by density gradient (Ficoll 1.083) centrifugation, and the entire samples are magnetically labelled with antibodies against epitopes on HLA-B locus: one Lambda; mouse anti-human Bw4 (cat # BIH0007; mouse anti-human IgG2a); mouse anti-human Bw6 (cat #BIH0038; mouse anti-human IgG3) (Data
35 code Bw4/6).

Cells are then washed and labelled with saturating amounts of streptavidin-coated paramagnetic particles (Molecular Probes / Invitrogen; Cat# C-21476 "Captivate"). Magnetically labelled cell samples are passed through a magnetised column (Miltenyi, LS columns Cat# 130-042-401), retaining all labelled cells. Cells
5 passing through the column are collected for further processing for telomere length.

Following the above negative selection, fetal cells in the enriched fetal cell population are selected on the basis of telomere length. To achieve this end, cells are washed once in 500 μ l TE and centrifuged at 1000g for 5 min. Cells are resuspended in 5 μ l of TE (avoiding bubbles). 20 μ l of PNA (Dako Telomere PNA kit/FITC, Dako-
10 Cytomation) in hybridization buffer is added and co-denatured at 80°C for 20 min in thermocycler.

Hybridization is allowed to proceed for 12 hours at 37°C. The cells are then washed with 2 x SSC buffer, and pelleted at 1000g for 5 min. As much supernatant as possible is removed, and the cells resuspended in 200 μ l 2x SSC/ 0.3% NP40. The
15 cells are incubated at 37°C for 30 min, centrifuged at 1000g for 5 min, and as much supernatant as possible removed. The cells are then resuspended in 200 μ l 2x SSC/ 0.3% NP40, and incubated at room temp for 30 min. The cells are then centrifuged at 1000g for 5 min. As much supernatant as possible is removed and the cells resuspended in 2.5 μ l TE.

20 Cells are analysed and labelled cells separated using fluorescence activated cell sorting on a MoFlo High-speed cell sorter (Dako-Cytomation, Ltd).

Analysis to confirm the presence of fetal cells may be by Fluorescence in situ hybridisation or by quantitative PCR

25

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as
30 illustrative and not restrictive.

All publications discussed above are incorporated herein in their entirety.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of
35 these matters form part of the prior art base or were common general knowledge in the

field relevant to the present invention as it existed before the priority date of each claim of this application.

REFERENCES

- Al-Mufti, R. *et al.* (1999) *Am. J. Med. Genet.* 85:66-75.
- Baerlocher *et al.* (2002) *Cytometry* 47: 89-99.
- 5 Baerlocher *et al.* (2003) *Cytometry* 55A: 1-6.
- Bauer *et al.* (2002) *Int. J. Legal Med.* 116:39-42.
- Berglund and Starkey (1989) *J Immunol Methods* 125: 79-87.
- Blaschitz *et al.* (2000) *Placenta* 21:733-741.
- Cabuy *et al.* (2004) *Cytometry* 62A:150-161.
- 10 De Pauw *et al.* (1998) *Cytometry* 32:163-169.
- Douglas *et al.* (1959) *Am. J. Obstet. Gynec.* 78:960-973.
- Findlay *et al.* (1996) *Hum Reprod Update* 2: 137-152.
- Findlay *et al.* (1998) *J Clin Pathol Mol Pathol* 51: 164-167.
- Findlay *et al.* (2001) *Mol Cellul Endocrinol* 183: S5-S12.
- 15 Fitzgerald *et al.* (1993) *Biotechniques* 15:128-133.
- Harrington *et al.* (1997) *Science* 275: 973-977.
- Kavaler *et al.* (1998) *Cancer* 82: 708-714.
- King *et al.* (1996) *The Journal of Immunology* 156:2068-2076.
- Lehman and Kreipe (2001) *Methods* 25:409-418.
- 20 Matthews *et al.* (2001) *J Pathol* 194: 183-193.
- Michalet *et al.* (2005) *Science* 307:538-544.
- Narath *et al.* (2005) *Cytometry* 68A:113-120.
- Satyanarayana *et al.* (2004) *Cell Cycle* 3: 1138-1150.
- Schmid *et al.* (2002) *Cytometry* 49: 96-105.
- 25 Schroder (1975) *J. Med. Genet.* 12:230-242.
- Seimiya *et al.* (2000) *EMBO J* 19: 2652-2661.
- Seki *et al.* (2001) *Clinical Cancer Research* 7: 1976-1981.
- Shorter *et al.* (1993) *Placenta* 14:571-582.
- Sidransky (2002) *Nature Reviews Cancer* 2: 210-219.
- 30 Tokumasu and Dvorak (2003) *J. Microsc.* 211:256-261.
- Trulsson *et al.* (2003) *Am J Surg* 186: 83-88.
- Vaziri *et al.* (1994) *PNAS* 91:9857-9860.

CLAIMS

1. A method of enriching fetal cells from a sample, the method comprising
 - i) depleting maternal cells by removing cells that express at least one MHC
- 5 molecule on their surface, and
 - ii) selecting fetal cells by
 - a) selecting cells that express telomerase, and/or
 - b) selecting cells based on telomere length.
- 10 2. A method of enriching fetal cells from a sample, the method comprising removing from the sample cells that express at least one MHC molecule on their surface.
3. The method of claim 1 or claim 2, wherein the MHC molecule is a Class I MHC
- 15 molecule.
4. The method of claim 3, wherein the Class I MHC molecule is HLA-A and/or HLA-B.
- 20 5. The method of claim 4 which comprises
 - i) contacting cells in the sample with an agent that binds at least one MHC molecule, and
 - ii) removing cells bound by the agent.
- 25 6. The method of claim 5, wherein the MHC molecule is a Class I MHC molecule.
7. The method of claim 6, wherein the Class I molecule is HLA-A and/or HLA-B.
8. The method of claim 5, wherein the method comprises contacting the sample
- 30 with i) an agent that binds at least one Class I MHC molecule, and ii) an agent that binds at least one Class II MHC molecule.
9. The method of claim 5, wherein the agent binds:
 - i) a monomorphic determinant of HLA-A molecules,
 - ii) a monomorphic determinant of HLA-B molecules, or
 - 35 iii) a monomorphic determinant of HLA-A and HLA-B molecules.

10. The method according to any one of claims 7 to 9, wherein the agent does not bind HLA-C.
- 5 11. The method of claim 5, wherein the agent binds a monomorphic determinant of HLA-A, HLA-B and HLA-C molecules.
12. The method of claim 11, wherein the agent is used at sub-saturating concentrations.
- 10 13. The method of claim 5, wherein more than two agents are used which bind different alleles of the same class of MHC molecule.
14. The method of claim 13, wherein collectively the agents bind all alleles of the
15 same class of MHC molecule.
15. The method according to any one of claims 1 to 14, wherein the method comprises
- 20 i) contacting cells in the sample with an agent that binds a compound that associates with an MHC molecule, and
- ii) removing cells bound by the agent.
16. The method according to any one of claims 1 to 15, wherein the genotype of an
25 MHC allele is not determined for the mother, father and/or fetus.
17. The method according to any one of claims 5 to 16, wherein the agent is an antibody or antibody fragment.
18. The method according to any one of claims 5 to 17, wherein the agent is bound
30 to a detectable label or isolatable label.
19. The method according to any one of claims 5 to 18, wherein the method further comprises binding to the agent a detectable label or isolatable label.
- 35 20. The method of claim 18 or claim 19, wherein the label is selected from the group consisting of: a fluorescent label, a radioactive label, a paramagnetic particle, a

chemiluminescent label, a label that is detectable by virtue of a secondary enzymatic reaction, and a label that is detectable by virtue of binding to a molecule.

21. The method according to any one of claims 18 to 20, wherein the step of
5 removing cells comprises detecting the label and removing the labeled cells.

22. The method of claim 21, wherein the detectable label or isolatable label is a
fluorescent label and wherein the step of removing cells comprises performing
fluorescence activated cell sorting.

10

23. The method of claim 21, wherein the detectable label or isolatable label is a
paramagnetic particle and wherein the step of removing cells comprises exposing the
labelled cells to a magnetic field.

15 24. The method according to any one of claims 2 to 23, wherein the method further
comprises contacting the cells with an agent that binds fetal cells, and selecting cells
bound by the agent that binds fetal cells.

25. A method of enriching fetal cells from a sample, the method comprising
20 selecting cells from the sample that express telomerase.

26. The method of claim 1 or claim 25, wherein the method comprises detecting a
protein component of telomerase.

25 27. The method of claim 26, wherein the protein component of telomerase is
telomere reverse transcriptase (TERT), telomerase associated protein-1 (TEP-1), or 14-
3-3 protein.

28. The method of claim 26 or claim 27 which comprises exposing the cells to an
30 antibody which specifically binds a protein component of telomerase.

29. The method of claim 28, wherein the antibody is detectably labelled.

30. The method of claim 28, wherein the method comprises exposing the cells to a
35 detectably labelled secondary antibody which binds the antibody.

31. The method of claim 1 or claim 25, wherein the method comprises detecting an RNA component of telomerase.
32. The method of claim 1 or claim 25, wherein method comprises detecting an
5 mRNA encoding a protein component of telomerase.
33. The method of claim 31 or claim 32, wherein the method comprises exposing the cells to a labelled probe which hybridizes to the RNA or mRNA.
- 10 34. The method of claim 33, wherein the probe is a PNA probe.
35. A method of enriching fetal cells from a sample, the method comprising selecting cells based on telomere length.
- 15 36. The method of claim 1 or claim 35, wherein the method comprises contacting cells with a detectably labelled probe that binds telomeres.
37. The method of claim 36, wherein about 1 to about 100 cells are selected, and wherein the selected cells have been bound by more probe than the other cells in the
20 sample.
38. The method according to any one of claims 1 to 37, wherein the sample is maternal blood, cervical mucous or urine.
- 25 39. The method according to any one of claims 1 to 38, wherein the method further comprises removing from the sample red blood cells, lymphocytes, and/or cancer cells.
40. The method according to any one of claims 1 to 39, wherein the method further comprises removing hemopoietic cells from the sample.
30
41. The method of claim 40, wherein the method comprises contacting cells in the sample with an agent that binds a hemopoietic cell.
42. The method of claim 40 or claim 41, wherein the hemopoietic cell is selected
35 from the group consisting of: a T cell, a B cell, a macrophage, a neutrophil, a dendritic cell and a basophil.

43. The method of claim 42, the agent binds a cell surface protein of the cell selected from the group consisting of: CD3, CD4, CD8, CD10, CD14, CD15, CD45 and CD56.
- 5
44. The method according to any one of claims 1 to 43, wherein the sample was obtained from the mother in the first trimester of pregnancy.
45. A method of detecting a fetal cell(s) in a sample, the method comprising
10 analysing a candidate cell for the expression of telomerase.
46. A method of detecting a fetal cell(s) in a sample, the method comprising analysing a candidate cell for the presence of telomeres and/or analysing the length of the telomeres in a candidate cell.
- 15
47. An enriched population of fetal cells obtained by a method according to any one of claims 1 to 44.
48. A composition comprising fetal cells according to claim 47, and a carrier.
- 20
49. Use of an agent that binds at least one MHC molecule, and/or an agent that binds a compound that associates with an MHC molecule, for enriching fetal cells from a sample.
- 25
50. Use of an agent that binds telomerase for enriching fetal cells from a sample.
51. Use of an agent that binds telomeres for enriching fetal cells from a sample.
52. A method for analysing the genotype of a fetal cell at a locus of interest, the
30 method comprising
- i) obtaining enriched fetal cells using a method according to any one of claims 1 to 44, and/or detecting a fetal cell using a method of claim 45 or claim 46, and
 - ii) analysing the genotype of at least one fetal cell at a locus of interest.
- 35
53. The method of claim 52, wherein the method comprises karyotyping, hybridization based procedures, and/or amplification based procedures.

54. The method of claim 52 or claim 53, wherein the fetal cell is analysed for a genetic abnormality linked to a disease state, or predisposition thereto.
- 5 55. A method of determining the sex of a fetus, the method comprising
i) obtaining enriched fetal cells using a method according to any one of claims 1 to 44, and/or detecting a fetal cell using a method of claim 45 or claim 46, and
ii) analysing at least one fetal cell to determine the sex of the fetus.
- 10 56. A method of determining the father of a fetus, the method comprising
i) obtaining enriched fetal cells using a method according to any one of claims 1 to 44, and/or detecting a fetal cell using a method of claim 45 or claim 46, and
ii) determining the genotype of the candidate father at one or more loci,
iii) determining the genotype of the fetus at one or more of said loci, and
15 iv) comparing the genotypes of ii) and iii) to determine the probability that the candidate father is the biological father of the fetus.
57. The method according to any one of claims 52 to 56, wherein the method further comprises identifying a cell obtained using a method according to any one of claims 20 to 35 as a fetal cell.
58. A kit for enriching fetal cells from a sample, the kit comprising
i) an agent that binds at least one MHC molecule, and/or an agent that binds a compound that associates with an MHC molecule, and/or an agent that binds a
25 hemopoietic cell, and
ii) a molecule which binds to telomerase, and/or which hybridizes to a polynucleotide encoding a protein component of said telomerase, and/or which hybridizes to telomeres.
- 30 59. A kit for enriching fetal cells from a sample, the kit comprising an agent that binds at least one MHC molecule, and/or an agent that binds a compound that associates with an MHC molecule, and/or an agent that binds a hemopoietic cell.
60. The kit of claim 58 or claim 59, wherein the agent that binds at least one MHC
35 molecule is an antibody.

61. The kit of claim 60, wherein the kit comprises
i) an agent that binds all HLA-A molecules,
ii) an agent that binds all HLA-B molecules, and/or
iii) an agent that binds all HLA-A and HLA-B molecules.
- 5
62. The kit according to any one of claims 58 to 61, wherein at least one agent is linked to a magnetic bead.
63. A kit for detecting a fetal cell, the kit comprising a molecule which binds to
10 telomerase, and/or which hybridizes to a polynucleotide encoding a protein component of said telomerase, and/or which hybridizes to telomeres.
64. The kit of claim 58 or claim 64, wherein the molecule is selected from the group consisting of; an anti-telomerase antibody, a polynucleotide which hybridizes to mRNA
15 encoding a protein component of telomerase, a polynucleotide which hybridizes to an RNA component of telomerase, or a polynucleotide which hybridizes to telomeric DNA on the chromosome.
65. A kit for detecting a genetic abnormality in a fetal cell, the kit comprising
20 i) a molecule for detecting a fetal cell, wherein the molecule binds to telomerase, which hybridizes to a polynucleotide encoding a protein component of said telomerase, or which hybridizes to telomeres, and
ii) at least one reagent for detecting said genetic abnormality.




Figure 1




Figure 2

HLA depletion

GA depleted fraction			1% of retained	
	<i>Y</i>	<i>all</i>	<i>Y</i>	<i>all</i>
8	4.7	4308	nd	nd
8	99	9550	nd	nd
8	133	171794	13	312717
8	517	44004	12	384559
8	90	3057	0	144771
8	20	380000	0	178000
9	140	2480	nd	nd
9	24	40000	0	23000
10	1.4	4800	0	14000
10	2.5	566	0	61000
10	1.1	45000	2.2	116000
10	1	149	nd	nd
10	2	457	1.5	23000

CD45 depletion

GA depleted fraction			1% of retained	
	<i>Y</i>	<i>all</i>	<i>Y</i>	<i>all</i>
8	6.7	110000	0	60000
8	1	875	0	188000
9	650	113000	1	150000
9	7.2	1025	0	163000
11	7.45	3300	0	125000

Figure 3




Figure 4

Clone 39-F alone		Clone 39-F + CD45		Clone W6/32 + CD45	
GA	Y cells all cells	GA	Y cells all cells	GA	Y cells all cells
8	4.7 4308	7	34 937	8	36 4241
8	99 9550	7	2 1906	8	716 10956
8	133 171794	7	217 2624	8	1 12685
8	517 44004	7	2.2 1407	9	1 3498
8	90 3057	7.5	1.2 1334	9	4 564
8	20 380000	8	1 3586		
9	140 2480	8	35 415		
9	24 40000	8	47 1500		
10	1.4 4800	11	1.6 1939		
10	2.5 566	15	6.3 537		
10	1.1 45000				
10	1 149				
10	2 457				

Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10

SEQUENCE LISTING

<110> Genetic Technologies Ltd

<120> Methods of enriching fetal cells

<130> 504338

<150> 60/679745

<151> 2005-05-11

<150> 60/689745

<151> 2005-06-09

<150> 60/725365

<151> 2005-10-11

<160> 3

<170> PatentIn version 3.3

<210> 1

<211> 1132

<212> PRT

<213> Homo sapiens

<400> 1

Met Pro Arg Ala Pro Arg Cys Arg Ala Val Arg Ser Leu Leu Arg Ser
1 5 10 15

His Tyr Arg Glu Val Leu Pro Leu Ala Thr Phe Val Arg Arg Leu Gly
 20 25 30

Pro Gln Gly Trp Arg Leu Val Gln Arg Gly Asp Pro Ala Ala Phe Arg
 35 40 45

Ala Leu Val Ala Gln Cys Leu Val Cys Val Pro Trp Asp Ala Arg Pro
 50 55 60

Pro Pro Ala Ala Pro Ser Phe Arg Gln Val Ser Cys Leu Lys Glu Leu
 65 70 75 80

Val Ala Arg Val Leu Gln Arg Leu Cys Glu Arg Gly Ala Lys Asn Val
 85 90 95

Leu Ala Phe Gly Phe Ala Leu Leu Asp Gly Ala Arg Gly Gly Pro Pro
 100 105 110

Glu Ala Phe Thr Thr Ser Val Arg Ser Tyr Leu Pro Asn Thr Val Thr
 115 120 125

Asp Ala Leu Arg Gly Ser Gly Ala Trp Gly Leu Leu Leu Arg Arg Val
 130 135 140

Gly Asp Asp Val Leu Val His Leu Leu Ala Arg Cys Ala Leu Phe Val
 145 150 155 160

Leu Val Ala Pro Ser Cys Ala Tyr Gln Val Cys Gly Pro Pro Leu Tyr
 165 170 175

Gln Leu Gly Ala Ala Thr Gln Ala Arg Pro Pro Pro His Ala Ser Gly
 180 185 190

Pro Arg Arg Arg Leu Gly Cys Glu Arg Ala Trp Asn His Ser Val Arg
 195 200 205

Glu Ala Gly Val Pro Leu Gly Leu Pro Ala Pro Gly Ala Arg Arg Arg
 210 215 220

Gly Gly Ser Ala Ser Arg Ser Leu Pro Leu Pro Lys Arg Pro Arg Arg
 225 230 235 240

Gly Ala Ala Pro Glu Pro Glu Arg Thr Pro Val Gly Gln Gly Ser Trp
 245 250 255

Ala His Pro Gly Arg Thr Arg Gly Pro Ser Asp Arg Gly Phe Cys Val
 260 265 270

Val Ser Pro Ala Arg Pro Ala Glu Glu Ala Thr Ser Leu Glu Gly Ala
 275 280 285

Leu Ser Gly Thr Arg His Ser His Pro Ser Val Gly Arg Gln His His
 290 295 300

Ala Gly Pro Pro Ser Thr Ser Arg Pro Pro Arg Pro Trp Asp Thr Pro
 305 310 315 320

Cys Pro Pro Val Tyr Ala Glu Thr Lys His Phe Leu Tyr Ser Ser Gly
 325 330 335

Asp Lys Glu Gln Leu Arg Pro Ser Phe Leu Leu Ser Ser Leu Arg Pro
 340 345 350

Ser Leu Thr Gly Ala Arg Arg Leu Val Glu Thr Ile Phe Leu Gly Ser
 355 360 365

Arg Pro Trp Met Pro Gly Thr Pro Arg Arg Leu Pro Arg Leu Pro Gln
 370 375 380

Arg Tyr Trp Gln Met Arg Pro Leu Phe Leu Glu Leu Leu Gly Asn His
 385 390 395 400

Ala Gln Cys Pro Tyr Gly Val Leu Leu Lys Thr His Cys Pro Leu Arg
 405 410 415

Ala Ala Val Thr Pro Ala Ala Gly Val Cys Ala Arg Glu Lys Pro Gln

	805		810		815
Ala Val Arg Ile Arg Gly Lys Ser Tyr Val Gln Cys Gln Gly Ile Pro					
	820		825		830
Gln Gly Ser Ile Leu Ser Thr Leu Leu Cys Ser Leu Cys Tyr Gly Asp					
	835		840		845
Met Glu Asn Lys Leu Phe Ala Gly Ile Arg Arg Asp Gly Leu Leu Leu					
	850		855		860
Arg Leu Val Asp Asp Phe Leu Leu Val Thr Pro His Leu Thr His Ala					
	865		870		880
Lys Thr Phe Leu Arg Thr Leu Val Arg Gly Val Pro Glu Tyr Gly Cys					
	885		890		895
Val Val Asn Leu Arg Lys Thr Val Val Asn Phe Pro Val Glu Asp Glu					
	900		905		910
Ala Leu Gly Gly Thr Ala Phe Val Gln Met Pro Ala His Gly Leu Phe					
	915		920		925
Pro Trp Cys Gly Leu Leu Leu Asp Thr Arg Thr Leu Glu Val Gln Ser					
	930		935		940
Asp Tyr Ser Ser Tyr Ala Arg Thr Ser Ile Arg Ala Ser Leu Thr Phe					
	945		950		960
Asn Arg Gly Phe Lys Ala Gly Arg Asn Met Arg Arg Lys Leu Phe Gly					
	965		970		975
Val Leu Arg Leu Lys Cys His Ser Leu Phe Leu Asp Leu Gln Val Asn					
	980		985		990
Ser Leu Gln Thr Val Cys Thr Asn Ile Tyr Lys Ile Leu Leu Leu Gln					

995	1000	1005
Ala Tyr Arg Phe His Ala Cys Val Leu Gln Leu Pro Phe His Gln		
1010	1015	1020
Gln Val Trp Lys Asn Pro Thr Phe Phe Leu Arg Val Ile Ser Asp		
1025	1030	1035
Thr Ala Ser Leu Cys Tyr Ser Ile Leu Lys Ala Lys Asn Ala Gly		
1040	1045	1050
Met Ser Leu Gly Ala Lys Gly Ala Ala Gly Pro Leu Pro Ser Glu		
1055	1060	1065
Ala Val Gln Trp Leu Cys His Gln Ala Phe Leu Leu Lys Leu Thr		
1070	1075	1080
Arg His Arg Val Thr Tyr Val Pro Leu Leu Gly Ser Leu Arg Thr		
1085	1090	1095
Ala Gln Thr Gln Leu Ser Arg Lys Leu Pro Gly Thr Thr Leu Thr		
1100	1105	1110
Ala Leu Glu Ala Ala Ala Asn Pro Ala Leu Pro Ser Asp Phe Lys		
1115	1120	1125

Thr Ile Leu Asp
1130

<210> 2

<211> 4015

<212> RNA

<213> Homo sapiens

<400> 2

gcagcgcugc guccugcugc gcacguggga agcccuggcc ccggccaccc ccgcaugcc 60

gcgcgcccc cgugccgag ccgugcgcuc ccugcugcgc agccacuacc gcgaggugcu 120

gccgcuggcc acguucgugc ggcgccuggg gccccagggc uggcggcugg ugcagcgcgg 180

ggacccggcg gcuuuccgcg cguggguggc ccagugccug gugugcugc ccugggacgc 240

acggccgccc cccgccgccc ccuccuuccg ccaggugucc ugccugaagg agcugguggc 300

ccgagugcug cagaggcugu gcgagcgcgg cgcaagaac gugcuggccu ucggcuucgc 360

gcugcuggac ggggccccgcg ggggcccccc cgaggccuuc accaccagcg ugcgcagcua 420

ccugcccaac acggugaccg acgcacugcg ggggagcggg gcgugggggc ucugcugcgc 480

ccgcgugggc gacgacgugc ugguuaccu gcuggcacgc ugcgcgcucu uugugcuggu 540

ggcucccagc ugcgccuacc aggugugcgg gccgccgcug uaccagcucg gcgcugccac 600

ucaggcccgg cccccgccac acgcuagugg accccgaagg cgucugggau gcgaacgggc 660

cuggaaccu agcgucaggg aggccggggu cccccugggc cugccagccc cgggugcgag 720

gaggcgcggg ggcagugcca gccgaagucu gccguugccc aagaggcca ggcguggcgc 780

ugccccugag ccggagcggg cggccguugg gcaggggucc ugggcccacc cgggacggac 840

gcguggaccg agugaccgug guuucugugu ggugucaccu gccagaccg ccgaagaagc 900

caccucuug gagggugcgc ucucuggcac gcgccacucc caccuaccg ugggccgcca 960

gcaccacgcg ggcccccau ccacaucgcg gccaccacgu cccugggaca cgccuugucc 1020

cccgguguac gccgagacca agcacuuccu cuacuccuca ggcgacaagg agcagcugcgc 1080

gcccuccuuc cuacucagcu cucugaggcc cagccugacu ggcgcucgga ggcucgugga 1140

gaccaucuuu cuggguucca ggcccuggau gccagggacu ccccgcaggu ugccccgccu 1200

gccccagcgc uacuggcaaa ugcggccccc guuucuggag cugcuuggga accacgcgca 1260

gugcccuac ggggugcucc ucaagacgca cugcccgcug cgagcugcgg ucaccccagc 1320

agccgguguc ugugcccggg agaagcccca gggcucugug gcggcccccg aggaggagga 1380

cacagacccc cgucgccugg ugcagcugcu ccgccagcac agcagccccc ggaggugua 1440

cggcuucgug cgggccugcc ugcgccggcu ggugcccca gccucuggg gcuccaggca 1500

caacgaacgc cgcuuccuca ggaacaccaa gaagucauc ucccugggga agcaugccaa 1560

gcucucgcug caggagcuga cguggaagau gagcugcgg gacugcguu ggucgcgag 1620

gagcccaggg guuggcugug uuccggccgc agagcaccgu cugcgugagg agauccuggc 1680

caaguuccug cacuggcuga ugagugugua cgucgucgag cugcucaggu cuuucuuua 1740

ugucacggag accacguuuc aaaagaacag gcucuuuuuc uaccggaaga gugucuggag 1800

caaguugcaa agcauuggaa ucagacagca cuugaagagg gugcagcugc gggagcuguc 1860

ggaagcagag gucaggcagc aucgggaagc caggcccgcc cugcugacgu ccagacuccg 1920

cuucaucccc aagccugacg ggcugcggcc gauugugaac auggacuacg ucgugggagc 1980

cagaacguuc cgagagaaa agagggccga gcgucucacc ucgaggguga aggcacuguu 2040

cagcugcuc aacuacgagc gggcgcgggc ccccggccuc cugggcgccu cugucuggg 2100

ccuggacgau auccacaggg ccuggcgcac cuucgugcug cgugugcggg cccaggaccc 2160

gccgccugag cuguacuug ucaaggugga ugugacggg gcguacgaca ccaucccca 2220

ggacaggcuc acggagguca ucgccagcau caucaaacc cagaacacgu acugcgugcg 2280

ucgguaugcc gugguccaga aggccgcca ugggcacguc cgcaaggccu ucaagagcca 2340

cgucucuacc uugacagacc uccagccgua caugcgacag uucguggcuc accugcagga 2400

gaccagcccg cugagggaug ccgucgucau cgagcagagc uccucccuga augaggccag 2460

caguggccuc uucgacgucu uccuacgcuu caugugccac cacgccgugc gcaucagggg 2520

caaguccuac guccagugcc aggggauccc gcagggcucc auccucucca cgugcucug 2580

cagccugugc uacggcgaca uggagaaca gcuguuugcg gggauucggc gggacgggcu 2640

gcuccugcgu uugguggaug auuucuuguu ggugacaccu caccucacc acgcgaaaac 2700

cuuccucagg acccuggucc gaggugucc ugaguauggc ugcgugguga acuugcggaa 2760

gacaguggug aacuucccug uagaagacga ggcccugggu ggcacggcuu uuguucagau 2820

gccggcccac ggccuauucc ccuggugcgg ccugcugcug gauaccgga ccuggaggu 2880

gcagagcgac uacuccagcu augcccggac cuccaucaga gccagucuca ccuucaccg 2940

cggcuucaag gcugggagga acaugcgucg caaacucuuu ggggucuuugc ggcugaagug 3000

ucacagccug uuucuggauu ugcaggugaa cagccuccag acggugugca ccaacauca 3060

caagauccuc cugcugcagg cguacagguu ucacgcaugu gugcugcagc ucccuuuca 3120

ucagcaaguu uggaagaacc ccacauuuuu ccugcgugc aucucugaca cggccuccu 3180

cugcuacucc auccugaaag ccaagaacgc agggauugc cugggggcca agggcgccg 3240

cggcccucug cccuccgagg ccgugcagug gcugugccac caagcauucc ugcucaagcu 3300

gacucgacac cgugucaccu acgugccacu ccugggguca cucaggacag cccagacgca 3360
 gcugagucgg aagcucccg ggcagacgcu gacugcccug gaggccgag ccaacccggc 3420
 acugcccuca gacuucaaga ccauccugga cugauggcca cccgcccaca gccaggccga 3480
 gagcagacac cagcagcccu gucacgccg gcucuacguc ccagggaggg aggggcggcc 3540
 cacaccagg cccgcaccgc ugggagucug aggccugagu gaguguuugg ccgaggccug 3600
 cauguccggc ugaaggcuga guguccggcu gaggccugag cgagugucca gccaagggcu 3660
 gaguguccag cacaccugcc gucuucacuu cccacaggc uggcgcucgg cuccaccca 3720
 gggccagcuu uuccucacca ggagcccggc uccacucucc cacauaggaa uaguccaucc 3780
 ccagauucgc cauuguucac cccucgcccu gccuuccuuu gccuuccacc cccaccaucc 3840
 agguaggagac ccugagaagg acccugggag cucugggaau uuggagugac caaaggugug 3900
 cccuguacac aggcgaggac ccugcaccug gauggggguc ccuguggguc aaauuggggg 3960
 gaggugcugu gggaguaaaa uacugaauau augaguuuuu caguuuugaa aaaaa 4015

<210> 3

<211> 451

<212> RNA

<213> Homo sapiens

<400> 3

ggguugcgga gggugggccu gggaggggug guggccauuu uuugucuaac ccuaacugag 60

aagggcguag ggcggugcu uuugcucucc ggcgcuguu uuucucgug acuuucagcg 120

ggcggaaaag ccucggccug ccgccuucca ccguucauuc uagagcaaac aaaaaauguc 180

agcugcuggc ccguucgccc cucccgggga ccugcggcgg gucgccugcc cagccccga 240

accccgccug gaggccgagg ucggcccggg gcuucuccgg aggcacccac ugccaccgag 300

aagaguuggg cucugucagc cgcgggucuc ucgggggaga gggcgagguu caggccuuc 360

aggccgagg aagaggaacg gagcgagucc ccgcgcgagg cgcgauuccc ugagcugugg 420

gacgugcacc caggacucgg cucacacaug c 451

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU2006/000617

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl.		
G01N 33/53 (2006.01) C12Q 1/68 (2006.01)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) BIOTECHABS, BIOSIS, MEDLINE, WPIDS, CAPLUS (telomerase, hTERT, major histocompatibility complex (MHC), human leuckocyte antigen (HLA), maternal circulation, maternal blood)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHAETZLEIN, S. et al. 2004. Telomere length is reset during early mammalian embryogenesis. Proc. Natl. Acad. Sci. USA. 101(21) : 8034-8038. (see whole document)	25-48, 52-57, 63-65
X	SOLIER, C. et al. 2002. Secretion of pro-apoptotic intron 4-retaining soluble HLA-G1 by human villous trophoblast. European Journal of Immunology. 32 : 3576-3586. (see whole document, especially p3577, paragraph bridging columns)	2-24, 47-49, 52-62
X	KOUMANTAKI, Y. et al. 2001. Microsatellite analysis provides efficient confirmation of fetal trophoblast isolation from maternal circulation. Prenatal Diagnosis. 21 : 566-570. (see whole document, especially p567, column 2, paragraph 2)	2-24, 52-57, 59-62
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents:		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 22 June 2006	Date of mailing of the international search report 28 JUN 2006	
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaaustralia.gov.au Facsimile No. (02) 6285 3929	Authorized officer David Olde Telephone No : (02) 6283 2569	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU2006/000617

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WEETMAN, A. P. 1999. The immunology of pregnancy. Thyroid. 9(7): 643-646. (see whole document, especially p644, column 1, paragraph 3)	2-24, 47-49, 52-57, 59-62
X	WO 1997015687 A (GERON CORPORATION) 1 May 1997. (see whole document, especially p8, lines 5-10 and lines 22-25)	25-34, 45, 47, 48, 52-57, 63-65

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

The international application does not comply with the requirements of unity of invention because it does not relate to one invention or to a group of inventions so linked as to form a single general inventive concept.

Continued in supplemental box

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

Supplemental Box

(To be used when the space in any of Boxes I to VIII is not sufficient)

Continuation of Box No:

Note that Rule 13.2 states that where a group of inventions is claimed in one and the same international application, the requirement of unity of invention referred to in Rule 13.1 shall be fulfilled only where there is a technical relationship among those inventions involving one or more of the same corresponding special technical features. The expression "special technical features" shall mean those technical features that define a contribution which each of the claimed inventions, considered as a whole, makes over the prior art.

The ISA has identified 3 separate inventions as follows:

Invention 1

Claims 1 (partially), 2 (completely), 3-24, 47-49, 52-62 (partially) relate to methods of enriching fetal cells by depleting or removing cells that express at least one major histocompatibility complex (MHC) molecule on their surface.

Invention 2

Claims 1, 3-24 (partially), 45 (completely), 47-50, 58, 60, 62-65 (partially) relate to methods of positively selecting or enriching for fetal cells by presence of telomerase.

Invention 3

Claims 1, 3-24 (partially), 46 (completely), 47-51, 58, 60 and 62-65 (partially) relate to methods of positively selecting or enriching for fetal cells by telomere length.

The only feature common to all the claims is selecting or enriching for fetal cells. However this concept is not novel as it is disclosed in the prior art. See for example the following documents identified in the International Search Report:

D2: SOLIER, C. et al. 2002. European Journal of Immunology. **32**: 3576-3586.

D3: KOUMANTAKI, Y. et al. 2001. Prenatal Diagnosis. **21**: 566-570.

D4: WEETMAN, A. P. 1999. Thyroid. **9(7)**: 643-646.

Accordingly, this feature cannot constitute a special technical feature within the meaning of PCT Rule 13.2, since it makes no contribution over the prior art.

Because the common feature does not satisfy the requirement for being a special technical feature it follows that it cannot provide the necessary technical relationship between the identified inventions. Therefore the claims do not satisfy the requirement of unity of invention *a posteriori*.

As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/AU2006/000617

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member					
WO	9715687	AU	11781/95	AU	12090/95	AU	13307/95
		AU	36924/95	AU	43740/93	AU	60582/98
		AU	63808/96	AU	71836/98	AU	89495/98
		AU	89496/98	CA	2135648	CA	2173872
		CA	2202618	CA	2245461	CA	2245462
		EP	0642591	EP	0728207	EP	0789780
		HK	1011384	JP	11123100	JP	11127874
		JP	11243998	JP	2000116388	JP	2002101898
		JP	2005198659	US	5489508	US	5580726
		US	5629154	US	5639613	US	5645986
		US	5648215	US	5686245	US	5686306
		US	5693474	US	5695932	US	5707795
		US	5744300	US	5804380	US	5830644
		US	5837453	US	5840495	US	5863726
		US	5891639	US	5989807	US	6007989
		US	6194206	US	6368789	US	6391554
		US	6551774	US	2002127634	US	2003175766
		US	2003190638	US	2004198659	WO	9323572
		WO	9513381	WO	9513382	WO	9513383
		WO	9613610				

Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001.

END OF ANNEX

专利名称(译)	富集胎儿细胞的方法		
公开(公告)号	EP1886138A1	公开(公告)日	2008-02-13
申请号	EP2006721493	申请日	2006-05-11
[标]申请(专利权)人(译)	遗传技术有限公司		
申请(专利权)人(译)	遗传技术有限公司		
当前申请(专利权)人(译)	遗传技术有限公司		
[标]发明人	BOEHMER RALPH MICHAEL ALLMAN RICHARD		
发明人	BOEHMER, RALPH, MICHAEL ALLMAN, RICHARD		
IPC分类号	G01N33/53 C12Q1/68 G01N33/569 G01N33/68		
CPC分类号	C12Q1/6881 C12Q1/6879 C12Q1/6883 C12Q2600/158 G01N33/5002 G01N33/5044 G01N33/5091 G01N33/5094 G01N33/54326 G01N33/56966 G01N33/573		
优先权	60/679745 2005-05-11 US 60/689745 2005-06-09 US 60/725365 2005-10-11 US		
其他公开文献	EP1886138A4		
外部链接	Espacenet		

摘要(译)

本发明涉及从怀孕女性中富集胎儿细胞的方法。本发明涉及从样品中除去包含至少一种MHC分子的细胞。本发明还涉及依赖于使用端粒酶，编码其组分的mRNA以及端粒长度作为胎儿细胞标记物的方法。富集的胎儿细胞可用于多种程序，包括检测感兴趣的性状，例如疾病性状，或遗传易感性，性别分类和亲子鉴定。