

(19)



(11)

EP 1 410 012 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
13.02.2013 Bulletin 2013/07

(51) Int Cl.:
G01N 33/68 (2006.01) C12Q 1/68 (2006.01)

(21) Application number: **02756158.8**

(86) International application number:
PCT/US2002/018479

(22) Date of filing: **10.06.2002**

(87) International publication number:
WO 2002/103352 (27.12.2002 Gazette 2002/52)

(54) DETERMINATION OF PREGNANCY STATUS

BESTIMMUNG DES SCHWANGERSCHAFTSZUSTANDS

TEST D'ETAT DE GESTATION

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
 MC NL PT SE TR**

(30) Priority: **19.06.2001 US 299553 P**

(43) Date of publication of application:
21.04.2004 Bulletin 2004/17

(73) Proprietor: **Idaho Research Foundation
 Moscow,
 Idaho 83844-3003 (US)**

(72) Inventor: **OTT, Troy, L.
 Moscow, ID 83843 (US)**

(74) Representative: **Gowshall, Jonathan Vallance et al
 Forresters
 Skygarden
 Erika-Mann-Strasse 11
 80636 München (DE)**

(56) References cited:
US-A- 4 705 748

- **OTT T L ET AL: "Effects of the estrous cycle and early pregnancy on uterine expression of Mx protein in sheep (ovis aries)" BIOLOGY OF REPRODUCTION, SOCIETY FOR THE STUDY OF REPRODUCTION, CHAMPAIGN, IL, US, vol. 59, 1998, pages 784-794, XP002962277 ISSN: 0006-3363**
- **BAZER ET AL: "Maternal recognition of pregnancy: Comparative aspects" PLACENTA, W.B. SAUNDERS, vol. 19, 1998, pages 375-386, XP005419052 ISSN: 0143-4004**

- **SPENCER THOMAS E ET AL: "Expression of interferon regulatory factors one and two in the ovine endometrium: Effects of pregnancy and ovine interferon tau" BIOLOGY OF REPRODUCTION, vol. 58, no. 5, May 1998 (1998-05), pages 1154-1162, XP002408450 ISSN: 0006-3363**
- **DEISENHAMMER F ET AL: "Bioavailability of interferon beta 1b in MS patients with and without neutralizing antibodies" NEUROLOGY, vol. 52, no. 6, 12 April 1999 (1999-04-12), pages 1239-1243, XP002408451 ISSN: 0028-3878**
- **CHARD T: "Interferon in pregnancy." JOURNAL OF DEVELOPMENTAL PHYSIOLOGY. MAY 1989, vol. 11, no. 5, May 1989 (1989-05), pages 271-276, XP009075225 ISSN: 0141-9846**
- **YANKEY S J ET AL: "Expression of the antiviral protein Mx in peripheral blood mononuclear cells of pregnant and brad, non-pregnant ewes" JOURNAL OF ENDOCRINOLOGY, BRISTOL, GB, vol. 170, 2001, pages R7-R11, XP002962276 ISSN: 0022-0795**
- **HICKS BRADY A ET AL: "Expression of the uterine Mx protein in the cyclic and pregnant cow, gilt, and mare" BIOLOGY OF REPRODUCTION, vol. 64, no. Supplement 1, 2001, page 267, XP009075329 & THIRTY-FOURTH ANNUAL MEETING OF THE SOCIETY FOR THE STUDY OF REPRODUCTION; OTTAWA, ONTARIO, CANADA; JULY 28-AUGUST 01, 2001 ISSN: 0006-3363**
- **HICKS B A ET AL: "Expression of the uterine Mx protein in cyclic and pregnant cows, gilts, and mares." JOURNAL OF ANIMAL SCIENCE. JUN 2003, vol. 81, no. 6, June 2003 (2003-06), pages 1552-1561, XP002408478 ISSN: 0021-8812**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 410 012 B1

- HANSEN ET AL.: 'Transient ubiquitin cross-reactive protein gene expression in the bovine endometrium' ENDOCRINOLOGY vol. 138, no. 11, 1997, pages 5079 - 5082, XP002955567
- JOHNSON ET AL.: 'Pregnancy and interferon-tau induce conjugation of bovine ubiquitin cross-reactive protein to cytosolic uterine proteins' BIOLOGY OF REPRODUCTION vol. 58, 1998, pages 898 - 904, XP002955568
- OLIVEIRA JODO F ET AL: "Expression of interferon (IFN)-stimulated genes in extrauterine tissues during early pregnancy in sheep is the consequence of endocrine IFN-tau release from the uterine vein", ENDOCRINOLOGY, vol. 149, no. 3, March 2008 (2008-03), pages 1252-1259, ISSN: 0013-7227
- GIFFORD C A ET AL: "Regulation of interferon-stimulated genes in peripheral blood leukocytes in pregnant and bred, nonpregnant dairy cows", JOURNAL OF DAIRY SCIENCE, vol. 90, no. 1, January 2007 (2007-01), pages 274-280, ISSN: 0022-0302

Description

Field of the Invention

[0001] The invention relates to methods for determining the pregnancy status of ruminant and non-ruminant ungulates, and non-hoofed ruminant animals.

Background of the Invention

[0002] In the rearing of livestock, it is very important to accurately determine the pregnancy status of bred animals. In particular, it is the accurate and early identification of failed pregnancy of an animal that has been bred that is economically important. Presently, once an animal is bred, for example a cow, pregnancy status is determined by such methods as palpation, which does not provide an accurate determination of pregnancy status until after 30 days following breeding. Because cattle have an estrous cycle of about 21 days, this means that with presently available methods at least one opportunity for breeding an animal that fails to conceive, the estrus period immediately following the failed breeding, will be missed.

[0003] This has important economic consequences for the cattle breeding industry, especially for the dairy industry. Efficient milk production farming requires that cows be successfully bred to become pregnant 80-100 days after calving. Dairy cows, however, have a low fertility rate with artificial insemination, requiring, on average, 2.5 to 3 inseminations per conception. Therefore, a significant need exists for a method by which a dairy farmer may accurately determine that an animal is not pregnant without missing an opportunity to re-breed the animal at the next estrus period following an unsuccessful breeding.

[0004] Sasser, U.S. Patent No. 4,705,748, discloses a method for determining pregnancy by detecting a protein produced by a conceptus. By this method, cattle were determined to be pregnant as early as day 27 following breeding. Sasser does not disclose the diagnosis of pregnancy prior to the time when a subsequent estrus period will have commenced in non-pregnant cattle and does not disclose an early determination of non-pregnancy.

[0005] Maternal recognition of pregnancy in ungulates involves local and systemic gene regulation by the conceptus that results in reduced or altered production of the luteolytic signal, prostaglandin F₂ α ((PGF₂ α); Yankey et al., Expression of the antiviral protein Mx in peripheral blood mononuclear cells of pregnant and bred, non-pregnant ewes. *Journal of Endocrinology* 170, R7-R11 (2001); Bazer et al., Regulation of endometrial responsiveness to estrogen and progesterone by pregnancy recognition signals during the peri-implantation period. In *Molecular and Cellular Aspects of Peri-implantation Processes*, pp 27-47. Ed S.K. Dey. Springer-Verlag, New York, Inc. (1995)). This is in contrast to pregnancy recognition in primates, which involves a direct luteo-

trophic effect on the corpus luteum (CL) by conceptus-produced chorionic gonadotropin (Bazer et al. 1995). The signal for maternal recognition in ungulates is the secretion by the conceptus of interferon-tau (IFN τ) during the second and third week of pregnancy (Bazer et al., 1995; Godkin et al., *J. Reprod. Fert.* 65:141-150(1982)). IFN τ prevents increases in endometrial estrogen and oxytocin receptors, to abrogate oxytocin-induced luteolytic pulses of PGF₂ α , and maintains CL function (Spencer et al., *Endocrinology* 136:4932-4944 (1995)).

[0006] IFN τ is a member of the Type I IFN family, which also includes IFN α , β , and ω (Samuel, *Virology* 183:1-11 (1991)), and, more recently, interferon δ (Lefevre, F., et al., *Biochimie* 80:779-788 (1998)). IFN τ signaling through the Type I IFN receptor and Janus Kinase (JAK)-signal transducer and activator of transcription (STAT) signal transduction pathway (Stewart et al., *Endocrinology* 142:98-107 (2001)) induces a number of genes in the ovine uterus including 2',5' oligoadenylate synthetase (Johnson et al., *Biol. Reprod.* 64:1392-1399 (2001)), β 2-microglobulin (Vallet et al., *J. Endocrinol.* 130:R1-4 (1991)), IFN regulatory factor 1 (Spencer et al., 1998), ubiquitin cross-reactive protein (Johnson et al., *Biol. Reprod.* 62:622-627(2000)), and Mx protein (Charleston and Stewart, *Gene* 137:327-331(1993); Ott et al., *Biol. Reprod.* 59:784-794 (1998)). While the functions of many of these proteins in the antiviral response are well characterized, their roles during early pregnancy are not.

[0007] Mx proteins are monomeric GTPases, which, depending on the species of animal and type of virus, are potent inhibitors of viral replication (Samuel, *Virology* 183:1-11 (1991)). The sequences of Mx proteins from various species, including sheep, cattle, pigs, and horses, are publicly available through GenBank and have been assigned GenBank Accession numbers X66093, U88329, M65087, and U55216, respectively. Although the antiviral effects of Mx are generally directed against negative-stranded RNA viruses (e.g. orthomyxovirus), their expression is induced in all cells that possess Type I IFN receptors and has been used to distinguish between bacterial and viral infection (Haller et al., *Rev. Sci. Tech.* 17:220-230 (1998)). Recently Mx mRNA and protein were shown to be elevated from epithelium (by day 13) to myometrium (by day 15) within the uterine wall in pregnant ewes and levels remained elevated through day 25 (Ott et al., *Biol. Reprod.* 59:784-794 (1998)). In addition, Mx mRNA levels were elevated in the corpus luteum in response to injections of rolIFN τ into the uterine lumen (Spencer et al., *Biol Reprod* 61:464-470 (1999)).

[0008] These results indicated that IFN τ was either: 1) acting directly on all uterine cell types (i.e., epithelial, stromal and myometrial) and on the CL; or 2) inducing substances (cytokines) that have paracrine/endocrine effects on uterine cells and other organs including the ovaries; or 3) affecting components of the uterine mucosal and circulating immune systems which then affect the various uterine cells and CL.

[0009] It is impractical, however, to measure the level

of Mx protein in uterine tissue as a test for evaluating pregnancy status. Besides being an invasive and time and labor intensive process, the disruption of uterine tissues necessary to determine the uterine levels of Mx would tend to have a deleterious effect on a pregnancy.

[0010] A significant need exists for a reliable, reproducible, and non-invasive method for determining pregnancy or lack of pregnancy in domestic livestock.

Summary of the Invention

[0011] The invention is defined in the claims. It has been discovered that the expression of the genes encoding for several proteins, herein referred to as "pregnancy-induced proteins", including 2', 5' oligoadenylate synthetase, β 2-microglobulin, IFN regulatory factor 1, ubiquitin cross-reactive protein, and Mx protein, increases significantly in certain animals during the first month of pregnancy. It has further been discovered that the increase in the expression of the pregnancy-induced proteins does not occur in animals that are not pregnant.

[0012] In many animals, the increase in expression of the pregnancy induced protein is due to the secretion by the embryo of a hormone, Type I interferon, that is the signal from the embryo to the mother of its existence, referred to as the signal for maternal recognition of pregnancy. Different type I interferons, including Interferon alpha ($IFN\alpha$), Interferon beta ($IFN\beta$), Interferon omega ($IFN\omega$), Interferon delta ($IFN\delta$), and Interferon tau ($IFN\tau$) are secreted by the embryos of different species. For example, $IFN\tau$ is secreted as a pregnancy recognition hormone in ruminants and $IFN\delta$ is secreted in swine. In other species, such as horses and other equidae, although the pregnancy-induced protein Mx protein is detectable in the uterus during early pregnancy, to date the secretion by the equine conceptus of a Type I interferon has not been demonstrated. Rather, in equines and other species whose conceptuses do not produce a Type I IFN, it is possible that the uterus produces Type I interferon in response to the presence of the embryo.

[0013] In one embodiment, the invention is a method for determining the pregnancy status of a ruminant or non-ruminant ungulate animal or a non-hoofed ungulate animal. According to this embodiment of the invention, the level of expression of a pregnancy induced protein during early pregnancy is determined and compared to the level of the expression of that pregnancy induced protein during the same period in a non-pregnant female animal of the same species. The pregnancy induced protein that is determined and compared is a Type I interferon-induced protein as set out in the claims. Most preferably, the pregnancy induced protein is Mx protein or ubiquitin cross-reactive protein.

[0014] As used herein, the term "pregnancy-induced protein" refers to a protein that is expressed by a maternal gene and which expression is induced in response to the presence of a pregnancy. A pregnancy-induced protein is distinct from a protein that is produced by the concep-

tus, unless such protein is also expressed by a maternal gene and this maternal expression is induced in response to the presence of a pregnancy.

[0015] As used herein, the term "early pregnancy" refers to that time during or following the period of pregnancy recognition signaling in which the level of a pregnancy induced protein is elevated in a pregnant animal compared to a non-pregnant animal of the same species. Although animals that are bred unsuccessfully, that is do not become pregnant, may or may not undergo a period of pregnancy recognition signaling, the term "early pregnancy" is used herein with regards to non-pregnant animals to refer to the period of time in which there would be an early pregnancy if the breeding had been successful. Typically, the period of early pregnancy, as used in relation to the method of the invention, ends at about the end of the first month following conception.

[0016] As used herein the "period of pregnancy recognition signaling" refers to that time during which the embryo secretes a protein or hormone, the secretion of which causes recognition by the mother of the existence of the embryo. Although animals that are bred unsuccessfully, that is do not become pregnant, may or may not undergo a period of pregnancy recognition signaling, the term is used herein with regards to non-pregnant animals to refer to the period of time in which, had the breeding been successful, biochemical signaling would be occurring between the conceptus and the uterus.

[0017] In accordance with the invention, an animal that is pregnant will exhibit a markedly higher level of expression of one or more pregnancy-induced proteins, such as Mx protein or ubiquitin cross-reactive protein, during early pregnancy, such as during the period of pregnancy recognition signaling, than will a non-pregnant animal of the same species. A non-pregnant animal, whether or not the animal has been bred will exhibit about the baseline level of the pregnancy-induced protein expression, including Mx protein, during this period.

[0018] In another embodiment, the disclosure provides a kit for determining the reproductive status of an animal of a species in which the conceptus secretes a protein or hormone as a signal for maternal recognition of pregnancy. According to this embodiment of the disclosure, the kit includes a receptacle for holding a test sample, one or more reagents which when combined with the test sample enable an operator to visually determine the level of one or more pregnancy-induced protein, such as Mx protein or ubiquitin cross-reactive protein, in the test sample, and instructions for determining the level of the protein in the sample. Preferably the kit further contains instructions that enable the operator to determine the pregnancy status of the animal based on the determined level of protein in the sample.

[0019] The invention is further illustrated below with reference to Mx protein. One skilled in the art will understand that the disclosure below is applicable to other pregnancy-induced proteins, such as other Type I interferon-induced proteins such as $IFN\tau$ induced proteins,

including 2', 5' oligoadenylate synthetase, β 2-microglobulin, IFN regulatory factor 1, and ubiquitin cross-reactive protein, as well as to the illustrated Mx protein. Therefore, in the following disclosure, at the mention of the term "Mx protein", any other pregnancy-induced protein may be substituted.

Brief Description of the Figures

[0020]

Figure 1 is a Northern Blot analysis of Mx mRNA from PBMC (peripheral blood mononuclear cells) at day 26 following artificial insemination in ewes. Lanes 1-6 represent pregnant ewes and lanes 8-13 represent non-pregnant ewes. Mx mRNA migrated at - 2.5 kb.

Figure 2 is a graph showing the results of a slot blot analysis of total cellular RNA isolated from PBMC at day 26 following artificial insemination in ewes. Mx mRNA levels were about 4 fold greater in pregnant versus bred, non-pregnant ewes at D 26 ($P < 0.01$). PSPB (pregnancy-specific protein B) levels (\blacklozenge) confirmed pregnancy status and were correlated with number of lambs born. (CNTS in Fig. 2 refers to photon counts on a chemiluminescent Northern blot.)

Figure 3 is a bar graph showing the expression of Mx mRNA in PBMC from artificially inseminated pregnant and non-pregnant ewes from Day 0 to Day 30 following the artificial insemination. (CNTS in Fig. 3 refers to photon counts on a chemiluminescent slot blot.)

Figure 4 is a Western Blot analysis of Mx protein expression from PBMC in pregnant and non-pregnant ewes 15 and 18 days following artificial insemination.

Detailed Description of the Invention

[0021] The present invention is applicable to any non-ruminant or ruminant ungulate animal or non-hoofed ruminant animal species that secretes increased levels of one or more pregnancy induced proteins during early pregnancy. The animal species is one that secretes a Type I interferon as a signal for maternal recognition of pregnancy. Most preferably, the animal species is one that secretes the hormone IFN τ as a signal for maternal recognition of pregnancy. In these animals, the increased levels of Type I interferon, such as IFN τ , induces an increased expression of Mx protein during the period of pregnancy recognition signaling. According to the invention, the lack of an increase in expression of a pregnancy-induced protein, such as Mx protein, in a suitable animal during early pregnancy, preferably the period of time which would encompass the period of pregnancy recognition signaling following a breeding, is a positive indication that the animal is not pregnant. Conversely, a negative result, that is the presence of an increase in expres-

sion of the protein during this period is an indication that the animal is pregnant.

[0022] In this application, the lack of an increased expression of the pregnancy induced protein, such as Mx protein, is referred to as a positive result whereas the presence of an increased expression of the protein is referred to as a negative result. This terminology, which might at first appear to be contrary to the usual usage of the terms "positive" and "negative" result, is utilized herein because it is the finding of non-pregnancy, rather than of pregnancy, which is of most concern to a farmer or rancher or other person engaged in animal husbandry. If an animal is determined to be pregnant, no additional work is expended to ensure that she is indeed pregnant, outside of watching her to look for signs that the pregnancy has been terminated. In contrast, if an animal is determined to be not pregnant, then she must be further evaluated for the onset of her next estrus and will be bred again. Therefore, it is the finding of non-pregnancy that provides the impetus for additional labor to be expended upon the animal to ensure that she does indeed become pregnant.

[0023] As stated above, the invention is applicable to any female animal belonging to a species that produces increased levels of a pregnancy-induced protein during early pregnancy, such as an animal of a species in which a Type I interferon such as IFN τ is the sole signal or one of more than one signal for maternal recognition of pregnancy. Animals suitable for the method of the invention are ruminant or non-ruminant ungulates and non-hoofed ruminants. The ungulates may be ruminants, such as cattle, sheep, goats, yak, water buffalo, and bison. Included among the ungulate ruminants suitable for the invention are also non-domesticated ungulates such as antelopes, gazelles, elk, reindeer, moose, bighorn sheep, giraffes, and other members of the cattle, sheep, and goat families. Non-hoofed ruminant non-ungulates suitable for the method of the invention include bactrian and dromedary camels and other camellids, such as llamas, alpacas, and vicunas. Ungulate non-ruminants suitable for the invention include domesticated and non-domesticated swine and horses.

[0024] In accordance with the method of the invention, during an appropriate time period following breeding, an animal is tested for the presence, or more precisely for the lack of presence, of an increased expression of a pregnancy-induced protein, exemplified hereafter as Mx protein. The test may be performed utilizing any cell in which Mx protein is expressed or in any bodily fluid in which Mx protein is found.

[0025] When determining whether an animal has or does not have an increase in expression of Mx protein, a comparison is made to the level of expression in animals that are known to be not pregnant. The comparison may be made, for example, by running a side-by-side comparison of a test sample from an animal that has been bred and which the pregnancy status is uncertain. Alternatively and preferably, the comparison is made by

testing a sample from an animal that has been bred and which the pregnancy status is uncertain and comparing the level of Mx protein in the sample to the level of Mx protein known to be present in non-pregnant animals, that is using a historical control.

[0026] The invention is illustrated herein with reference to determining the increased expression of Mx protein in peripheral blood mononuclear cells (PBMC). However, any cell in which Mx protein is expressed, including other nucleated cells present in the bloodstream, may be utilized in place of PBMC. Likewise, it is conceived that increased Mx expression in accordance with the invention may be determined by analysis of fluids, such as milk, saliva, urine, or nasal, ocular, or vaginal secretions, or whole blood, plasma, or serum.

[0027] Although the period of maternal recognition in most species occurs about the same time, for example at days 12 to 14 following breeding in ewes and days 15 to 18 in cows, the period of embryonic signaling that results in maternal recognition of pregnancy varies somewhat amongst different species. Accordingly, the actual dates following breeding on which the method of the invention may be effectively practiced will vary. For example, the period of maternal recognition of pregnancy signaling in domesticated sheep typically begins about day 11 following breeding and continues to about day 21. In cattle, this period typically begins about day 13 and continues to about day 35.

[0028] When comparing the level of Mx protein expression in a test sample to the baseline, (non-pregnant) level of Mx protein expression, typically a doubling or higher in Mx protein expression over the baseline is a negative test, that is the animal is not determined to be not pregnant. Usually, at peak levels of Mx protein expression during the period of pregnancy recognition signaling, a pregnant animal will have levels of Mx protein expression that are up to four or five times, or higher, that of baseline.

[0029] In accordance with the invention, the method may be practiced at any time commencing with the onset of the period of signaling until the time that the level of the pregnancy induced protein is no longer elevated in pregnant animals compared to non-pregnant animals of the same species. It is during this time that pregnant animals have an increased expression of Mx protein compared to non-pregnant animals. Thus, in sheep and cattle, the time period for comparing the level of Mx protein expression to determine pregnancy status is between 12 and 30 days following breeding. A more preferred time period is between days 12 and 21 following breeding. A most preferred time period between days 15 and 21, and a most preferred time is on day 18 in cattle and day 15 in sheep.

[0030] The level of Mx protein expression may be determined by any method that permits this determination to be made. Suitable methods include detecting the Mx protein itself, such as by ELISA test, an assay based on Mx protein function, or a Western blot. Suitable methods also include detecting increased levels of Mx mRNA,

such as by Northern blot, slot blot, or PCR. In a preferred embodiment, the level of Mx protein expression is determined by detecting the level of Mx protein present in a sample by a colorimetric assay based, for example, on the binding of an antibody to the Mx protein, similarly to the methods that are used in human home pregnancy diagnostic kits.

[0031] The method of the invention is an accurate, reproducible test that predictably determines that an animal is not pregnant, and may likewise be used to determine that an animal is pregnant. With regards to Mx protein in particular, but not necessarily the other pregnancy induced proteins, the only source of false negative results that would erroneously indicate that the animal is pregnant, that is an increased level of Mx protein expression, is if the animal is suffering from a severe viral infection.

[0032] The kit of the disclosure is preferably based on an enzyme linked assay (ELISA), such as what is known as an "immunometric" or "sandwich" assay. Such an assay involves "sandwiching" a ligand (such as an antigen) with two or more receptor molecules (such as antibodies) which complex with the ligand in a non-interfering manner and at different epitopic sites. Examples of such assays are described in David et al., U.S. Pat. No. 4,486,530.

Alternatively, the kit may be based on chemiluminescence assays, enhanced luminescence assays, and radioimmunoassays. In a preferred embodiment, the kit includes a package, which package houses a test surface, such as a slide or multiple test wells, that is bound to an antibody that will bind to an epitope of the protein of interest, such as Mx protein, a container housing a second antibody that will bind to a second epitope of the protein, which second antibody is labeled, a container housing a standard sample having a baseline concentration of the protein, a reagent that when contacted to the labeled second antibody permits the relative amount of the protein present to be visualized, and instructions for use of the kit to determine whether a test sample contains an amount of Mx protein indicative of pregnancy or non-pregnancy status.

[0033] The kit of the disclosure for determining pregnancy status by determining the relative level of a pregnancy induced protein, such as Mx protein, in a test sample compared to a control may be formulated in many different ways, which ways will be apparent to those skilled in the art upon reading the description herein. It is intended that these various formulations of the kit of the disclosure are included in the invention.

[0034] The invention is further described in the following illustrative, non-limiting, examples. The examples describe the method of the invention with reference to Mx protein. However, the method of the invention is applicable to other pregnancy induced proteins.

55 Example 1 Animal Models

[0035] Sixty (60) mature, white-faced, ewes from the U.S. Sheep Experiment Station (USSES, Dubois ID)

were synchronized and bred either by transcervical or laparoscopic artificial insemination ("AI"). Laparoscopic AI was performed according to the procedure disclosed in Stellflug et al., *J. Anim. Sci.* 79:568-573 (2001). The day of artificial insemination was designated Day 0 (D0) At 26 days after AI (D26), blood (10 ml) was collected by jugular venipuncture into EDTA-containing vacutainer tubes (Sherwood Medical, St. Louis MO). PBMC were isolated as described below in Example 2. Pregnancy was determined by assaying serum for pregnancy-specific protein B (PSPB; Biotracking Inc, Moscow ID) and lambing dates and number of lambs born were recorded.

Example 2. PBMC isolation

[0036] Blood was kept on ice until processed. Samples were centrifuged at 300 x g for 20 min at 4 C. The buffy coat was removed and resuspended in 0.87% Tris-NH₄CL lysis buffer at a 1 to 5 ratio. Samples were incubated for 5 min at 37°C and centrifuged at 300 x g for 10 min. The supernatant was removed and pellets were washed with 10 ml 1X PBS and centrifuged for 10 min at 300 x g. After removal of supernatant, cell pellets were either frozen at -80°C for protein extraction, or lysed with 2 ml TRIZOL (Life Technologies, Grand Island NY) and stored at -80°C for RNA extraction.

Example 3. RNA extraction, Northern and slot-blot analysis.

[0037] Total cellular RNA was extracted using TRIZOL according to manufacture's instructions. RNA was quantified by A260:280 ratio. To establish size and number of Mx transcripts in PBMC, RNA (5 µg) was electrophoresed in a 1% agarose/0.615 M formaldehyde gel and transferred to a nylon membrane (Nytran, Schleicher & Schuell, Keene NH) by capillary blotting. For quantification of Mx mRNA levels in PBMC, RNA (5 µg) was transferred to a nylon membrane by vacuum filtration (Minifold II, Schleicher & Schuell, Keene NH). Blots were probed with a biotin-labeled ovine Mx anti-sense cRNA probe (Ott et al., *Biol. Reprod.* 59:784-794 (1998)) using the North2South Hybridization kit (Pierce, Rockford IL) and chemiluminescent signal was quantified using a Bio-Rad Fluor-S Multilmager system and Quantity One software (Bio-Rad, Hercules CA). Slot-blots were stripped and re-probed with an ovine 18s rRNA cRNA probe to correct for variations in RNA loading.

[0038] Northern blot analysis, as shown in Fig. 1, detected a single, approximately 2.5 kD, band in PBMC isolated from pregnant and bred, non-pregnant ewes, which agrees with the known size of the ovine uterine Mx cDNA (Charleston and Stewart, *Gene* 137:327-331 (1993); Ott et al., *Biol. Reprod.* 59:784-794 (1998)).

[0039] Slot blot analysis, as shown in Fig. 2, of total cellular RNA isolated from PBMC collected at D26 post-AI showed a fourfold increase in Mx mRNA levels in pregnant versus bred, non-pregnant (n=26) ewes (P<0.01).

In addition, ewes carrying multiples (triplets or quads; n=10) had higher Mx mRNA levels than those carrying singles (n=10) or twins (n=9; P<0.05). Results from the PSPB (pregnancy-specific protein B) assay confirmed pregnancy status and, as reported previously, levels of PSPB were correlated with number of lambs born (Willard et al., *J. Anim. Sci.* 73:960-966 (1995)).

Example 4. Temporal expression of Mx protein mRNA during early pregnancy in sheep

[0040] A second study examined the temporal expression of Mx mRNA during early pregnancy in sheep, as shown in Fig. 3. Thirty four (34) mature Suffolk ewes were synchronized and bred by laparoscopic AI. Blood (20 ml) was collected by jugular venipuncture at D0, and every three days from D9 to D30, and PBMC were isolated. Pregnancy was confirmed by real-time ultrasonography and PSPB assay at D30. Results shown in Fig. 3 are a representative subset of all ewes and depict results from four pregnant and four bred, non-pregnant ewes during the first 30 days following insemination. This allowed analyzing all replicates on a single blot to eliminate problems associated with signal intensity between blots. Results showed Mx mRNA levels increased in pregnant ewes beginning at D15 (P<0.01). Levels peaked at D21 and gradually declined thereafter. At D30, Mx levels in pregnant ewes remained elevated two-fold compared to bred, non-pregnant ewes (P<0.01).

Example 5. Protein isolation and Western blot analysis.

[0041] Total cellular protein was extracted using M-PER reagent (Pierce, Rockford IL), according to manufacturers instructions. Protein concentration of samples was quantified by BCA assay (Pierce, Rockford IL) with bovine serum albumen as the standard. Proteins (8 µg/sample) from PBMC isolated from pregnant and bred, non-pregnant ewes at D15 and D18 were separated by 12% SDS-PAGE and electrophoretically transferred to a nitrocellulose membrane (BA83, Schleicher & Schuell, Keene NH). Following blocking of non-specific binding sites in 5% non-fat dry milk in Tris-buffered saline and Tween 20 (TBST) for 2 hours at 25°C, membranes were incubated with a 1:1000 dilution of a polyclonal rabbit ovine Mx peptide antiserum (#90618-2; 0.7 µg/ml) at 4°C overnight. Goat anti-rabbit IgG (0.8 µg/ml) labeled with horseradish peroxidase was used at a 1:200,000 dilution as secondary antibody. Chemiluminescent signal was developed using the West Femto Maximum Sensitivity Substrate (Pierce, Rockford IL) and quantified using the Fluor-S Multilmager system and Quantity One software. **[0042]** As shown in Fig. 4, Mx protein (-75 kDa) was not detected in either D15 or D18 open (non-pregnant) ewes, but was strongly up-regulated in PBMC from pregnant ewes on both days. Two additional bands (-48 and 36 kDa) were detected in PBMC from pregnant ewes.

Example 6. Analysis.

[0043] Chemiluminescent signal was analyzed using GLM procedures of SAS (Version 8.1, SAS Inc, Gary NC). The model included, where appropriate, status (pregnant versus bred, non-pregnant), ewe nested within status, day (0, 9, 12, 15, 18, 21, 24, 27, and 30) and appropriate interactions. Error terms in the F test were according to the expectation of mean squares for error. Signal for 18s rRNA was run as a covariate in the model to correct for variations in loading. Results are reported as adjusted Least Squares Means (LSM) and pooled standard errors.

Example 7. Cattle

[0044] Thirty three dairy cows were bred by artificial insemination and their levels of Mx mRNA were determined by the method described above in Examples 2 and 3. On day 15, levels of Mx mRNA were found to be about the same in cows that were later determined to be pregnant and in cows that were later determined not to be pregnant. On day 18, levels of Mx mRNA were found to be have increased markedly in cows later determined to be pregnant to about three times the level found in cows later determined not to be pregnant.

[0045] The results demonstrate a rapid and sustained activation of Mx gene expression in response to pregnancy recognition signaling, and indicate that, in addition to local effects of IFN τ , there is rapid systemic response in sheep and cattle. In addition, Mx expression did not increase in PBMC when pregnancy was not established (bred, non-pregnant animals). These findings are significant because pregnancy recognition signaling by IFN τ was heretofore considered to result solely from local regulation of endometrial gene expression (Stewart et al., *Endocrinology* 142:98-107 (2001); Johnson et al., *Biol. Reprod.* 64:1392-1399 (2001); Vallet et al., *J. Endocrinol.* 130:R1-4 (1991); Spencer et al., *Biol. Reprod.* 58:1154-1162 (1998); Johnson et al., *Biol. Reprod.* 62:622-627 (2000); Charleston and Stewart, *Gene* 137:327-331(1993); Ott et al., *Biol. Reprod.* 59:784-794 (1998); Spencer et al., *Biol. Reprod.* 61:464-470 (1999)) and suppression of estrogen and oxytocin receptor expression to abrogate luteolytic pulses of PGF $_{2\alpha}$. The methods and kits of the invention therefore provide new and economically important methods of non-pregnancy, and pregnancy, determinations in livestock.

Claims

1. A method for determining pregnancy status in a ruminant or non-ruminant ungulate animal or non-hoofed ruminant animal comprising determining the level of extra-uterine expression of a pregnancy-induced protein selected from the group consisting of 2', 5' oligoadenylate synthetase, ubiquitin cross-re-

active protein, and Mx protein in a body fluid collected from the animal during the period of 12 to 30 days following breeding of the animal and comparing the level of expression of said protein in the animal to that of a non-pregnant animal of the same species.

5

2.

The method of claim 1 wherein the animal is a ruminant.

10

3.

The method of claim 2 wherein the ruminant is selected from the group consisting of bactrian camel, dromedary camel, llama, alpaca, and vicuna.

15

4.

The method of claim 2 wherein the animal is an ungulate.

5.

The method of claim 1 wherein the animal is selected from the group consisting of cattle, sheep, goats, yaks, water buffalos, bison, antelopes, and deer.

20

6.

The method of claim 5 wherein the animal is a ewe or a cow.

25

7.

The method of claim 1 wherein the animal is an ungulate.

8.

The method of claim 7 wherein the animal is a swine.

30

9.

The method of claim 1 wherein the protein is Mx protein.

35

10.

The method of claim 1 wherein the level of expression of the protein is determined by determining the level of mRNA coding for the protein.

11.

The method of claim 10 wherein the determination of mRNA is by Northern blot analysis, slot-blot analysis, or polymerase chain reaction.

40

12.

The method of claim 1 wherein the level of expression of the protein is determined by determining the level of the protein.

13.

The method of claim 12 wherein the determination of the level of production of the protein is by evaluating the binding of an antibody to the protein or by an assay based on a function of the protein.

45

14.

The method of claim 12 wherein the level of production of the protein is detected by a colorimetric assay.

50

15.

The method of claim 1 wherein the level of expression of the protein is determined by determining the expression of the protein in a cell of the animal.

55

16.

The method of claim 15 wherein the cell is a nucleated blood cell.

17. The method of claim 16 wherein the cell is a peripheral blood mononuclear cell.
18. The method of claim 1 wherein the bodily fluid is selected from the group consisting of milk, saliva, urine, or nasal, ocular, or vaginal secretions.
19. The method of claim 1 wherein the bodily fluid is blood, plasma, or serum.
20. The method of claim 1 wherein the animal is determined to be not pregnant if the level of expression of the protein in the animal is not elevated compared to the level of expression of the protein in a non-pregnant animal of the same species.
21. The method of claim 1 wherein the animal is determined to be pregnant if the level of expression of protein in the animal is two times or more the level of expression of the protein in a non-pregnant animal of the same species.
22. The method of claim 21 wherein the animal is determined to be pregnant if the level of expression of the protein in the animal is four times or more the level of expression of the protein in a non-pregnant animal of the same species.
23. The method of claim 1 wherein the comparison is by side-by-side comparison of the level of expression of the protein in the animal and the level of expression in an animal known to be pregnant or not pregnant.
24. The method of claim 1 wherein the comparing is by using an historical control.
25. The method of claim 1 which is performed during the period between 15 and 30 days after breeding of the animal.
26. The method of claim 25 which is performed during the period between 12 and 21 days after breeding of the animal.
27. The method of claim 26 which is performed during the period between 18 and 21 days after breeding of the animal.

Patentansprüche

1. Ein Verfahren zum Bestimmen eines Trächtigkeitszustands in einem wiederkäuenden oder nicht wiederkäuenden Huftier oder nicht behuften wiederkäuenden Tier, das Bestimmen des Niveaus der extraterinen Expression eines trächtigkeitsinduzierten Proteins, das aus der aus 2',5'-Oligoadenylat-

Synthetase, Ubiquitin Cross-Reactive Protein und Mx-Protein bestehenden Gruppe ausgewählt ist, in einer Körperflüssigkeit, die dem Tier während der Zeit von 12 bis 30 Tagen nach der Züchtung des Tiers entnommen wird, und Vergleichen des Niveaus der Expression des Proteins im Tier mit demjenigen eines nicht trächtigen Tiers derselben Spezies beinhaltet.

2. Verfahren nach Anspruch 1, wobei das Tier ein Wiederkäuer ist.
3. Verfahren nach Anspruch 2, wobei der Wiederkäuer aus der aus Baktrischer Kamelstute, Dromedarstute, Lamastute, Alpakastute und Vikunjastute bestehenden Gruppe ausgewählt ist.
4. Verfahren nach Anspruch 2, wobei das Tier ein Huftier ist.
5. Verfahren nach Anspruch 1, wobei das Tier aus der aus Rindkühen, Ziegenweibchen, Geißen, Yaks, Wasserbüffelkühen, Bisonkühen, Antilopenkühen und Hirschkühen bestehenden Gruppe ausgewählt ist.
6. Verfahren nach Anspruch 5, wobei das Tier ein Mutterschaf oder eine Kuh ist.
7. Verfahren nach Anspruch 1, wobei das Tier ein Huftier ist.
8. Verfahren nach Anspruch 7, wobei das Tier ein Schwein ist.
9. Verfahren nach Anspruch 1, wobei das Protein Mx-Protein ist.
10. Verfahren nach Anspruch 1, wobei das Niveau der Expression des Proteins durch Bestimmen des Niveaus von mRNA-Codierung für das Protein bestimmt wird.
11. Verfahren nach Anspruch 10, wobei die Bestimmung von mRNA durch eine Northern-Blot-Analyse, eine Slot-Blot-Analyse oder eine Polymerase-Kettenreaktion erfolgt.
12. Verfahren nach Anspruch 1, wobei das Niveau der Expression des Proteins durch Bestimmen des Niveaus des Proteins bestimmt wird.
13. Verfahren nach Anspruch 12, wobei die Bestimmung des Niveaus der Produktion des Proteins durch Beurteilen der Bindung eines Antikörpers an das Protein oder durch eine Prüfung basierend auf einer Funktion des Proteins erfolgt.

14. Verfahren nach Anspruch 12, wobei das Niveau der Produktion des Proteins durch eine kolorimetrische Prüfung erfolgt.
15. Verfahren nach Anspruch 1, wobei das Niveau der Expression des Proteins durch Bestimmen der Expression des Proteins in einer Zelle des Tiers bestimmt wird.
16. Verfahren nach Anspruch 15, wobei die Zelle eine kernhaltige Blutzelle ist.
17. Verfahren nach Anspruch 16, wobei die Zelle eine mononukleäre Zelle des peripheren Blutes ist.
18. Verfahren nach Anspruch 1, wobei die Körperflüssigkeit aus der aus Milch, Speichel, Urin oder Nasen-, Augen- oder Scheidensekreten bestehenden Gruppe ausgewählt ist.
19. Verfahren nach Anspruch 1, wobei die Körperflüssigkeit Blut, Plasma oder Serum ist.
20. Verfahren nach Anspruch 1, wobei bestimmt wird, dass das Tier nicht trächtig ist, falls das Niveau der Expression des Proteins im Tier verglichen mit dem Niveau der Expression des Proteins in einem nicht trächtigen Tier derselben Spezies nicht erhöht ist.
21. Verfahren nach Anspruch 1, wobei bestimmt wird, dass das Tier trächtig ist, falls das Niveau der Expression des Proteins im Tier mindestens zwei Mal so hoch wie das Niveau der Expression des Proteins in einem nicht trächtigen Tier derselben Spezies ist.
22. Verfahren nach Anspruch 21, wobei bestimmt wird, dass das Tier trächtig ist, falls das Niveau der Expression des Proteins im Tier mindestens vier Mal so hoch wie das Niveau der Expression des Proteins in einem nicht trächtigen Tier derselben Spezies ist.
23. Verfahren nach Anspruch 1, wobei der Vergleich durch einen Parallelvergleich des Niveaus der Expression des Proteins im Tier und des Niveaus der Expression in einem Tier, von dem bekannt ist, dass es trächtig oder nicht trächtig ist, erfolgt.
24. Verfahren nach Anspruch 1, wobei das Vergleichen durch Nutzung einer Verlaufskontrolle erfolgt.
25. Verfahren nach Anspruch 1, das während der Zeit zwischen 15 und 30 Tagen nach der Züchtung des Tiers durchgeführt wird.
26. Verfahren nach Anspruch 25, das während der Zeit zwischen 12 und 21 Tagen nach der Züchtung des Tiers durchgeführt wird.

27. Verfahren nach Anspruch 26, das während der Zeit zwischen 18 und 21 Tagen nach der Züchtung des Tiers durchgeführt wird.

Revendications

1. Procédé de détermination de l'état de gestation d'un animal ongulé ruminant ou non-ruminant ou d'un animal non-ongulé ruminant, comprenant les étapes consistant à déterminer le niveau d'expression extra-utérine d'une protéine induite par la gestation, sélectionnée parmi le groupe consistant en la 2',5'-oligoadénylate synthétase, l'ubiquitine protéine réaction croisée et la protéine Mx dans un fluide organique collecté sur l'animal au cours de la période des 12 à 30 jours suivant la mise à la reproduction de l'animal, et à comparer le niveau d'expression de ladite protéine chez l'animal avec celui d'un animal non gestant de la même espèce.
2. Procédé conforme à la revendication 1, où l'animal est un ruminant.
3. Procédé conforme à la revendication 2, où le ruminant est sélectionné parmi le groupe consistant en le chameau bactrien, le dromadaire, le lama, l'alpaca et la vigogne.
4. Procédé conforme à la revendication 2, où l'animal est un ongulé.
5. Procédé conforme à la revendication 1, où l'animal est sélectionné parmi le groupe consistant en des bovins, des moutons, des chèvres, des yacks, des buffles d'Inde, des bisons, des antilopes et des cerfs.
6. Procédé conforme à la revendication 5, où l'animal est une brebis ou une vache.
7. Procédé conforme à la revendication 1, où l'animal est un ongulé.
8. Procédé conforme à la revendication 7, où l'animal est un porc.
9. Procédé conforme à la revendication 1, où la protéine est la protéine Mx.
10. Procédé conforme à la revendication 1 où le niveau d'expression de la protéine est déterminé en déterminant le niveau de l'ARNm codant pour la protéine.
11. Procédé conforme à la revendication 10, où la détermination de l'ARNm s'effectue par analyse de transfert de Northern, analyse Slot blot ou réaction en chaîne de la polymérase.

12. Procédé conforme à la revendication 1, où le niveau d'expression de la protéine est déterminé en déterminant le niveau de la protéine.
13. Procédé conforme à la revendication 12, où la détermination du niveau de production de la protéine s'effectue par l'évaluation de la fixation d'un anticorps à la protéine ou par un essai basé sur une fonction de la protéine. 5
14. Procédé conforme à la revendication 12, où le niveau de production de la protéine est détecté par un essai colorimétrique. 10
15. Procédé conforme à la revendication 1, où le niveau d'expression de la protéine est déterminé en déterminant l'expression de la protéine dans une cellule de l'animal. 15
16. Procédé conforme à la revendication 15, où la cellule est une cellule sanguine nucléée. 20
17. Procédé conforme à la revendication 16, où la cellule est une cellule mononucléée de sang périphérique. 25
18. Procédé conforme à la revendication 1, où le fluide organique est sélectionné parmi le groupe consistant en le lait, la salive, l'urine, ou les sécrétions nasales, oculaire ou vaginales. 30
19. Procédé conforme à la revendication 1, où le fluide organique est le sang, le plasma ou le sérum.
20. Procédé conforme à la revendication 1, où l'animal est déterminé n'être pas gestant si le niveau d'expression de la protéine chez l'animal n'est pas élevé par comparaison avec le niveau d'expression de la protéine chez un animal non gestant de la même espèce. 35
21. Procédé conforme à la revendication 1, où l'animal est déterminé être gestant si le niveau d'expression de la protéine chez l'animal est égal à deux fois ou plus le niveau d'expression de la protéine chez un animal non gestant de la même espèce. 40
22. Procédé conforme à la revendication 21, où l'animal est déterminé être gestant si le niveau d'expression de la protéine chez l'animal est égal à quatre fois ou plus le niveau d'expression de la protéine chez un animal non gestant de la même espèce. 50
23. Procédé conforme à la revendication 1, où la comparaison est une comparaison côte à côte du niveau d'expression de la protéine chez l'animal et le niveau d'expression de la protéine chez un animal connu pour être gestant ou non gestant. 55
24. Procédé conforme à la revendication 1, où la comparaison s'effectue à l'aide d'un contrôle historique.
25. Procédé conforme à la revendication 1, qui est mis en oeuvre au cours de la période de 15 à 30 jours après la mise à la reproduction de l'animal.
26. Procédé conforme à la revendication 25, qui est mis en oeuvre au cours de la période de 12 à 21 jours après la mise à la reproduction de l'animal.
27. Procédé conforme à la revendication 26, qui est mis en oeuvre au cours de la période de 18 à 21 jours après la mise à la reproduction de l'animal.




Figure 1




Figure 2




Figure 3




Figure 4

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4705748 A [0004]
- US 4486530 A, David [0032]

Non-patent literature cited in the description

- **YANKEY et al.** Expression of the antiviral protein Mx in peripheral blood mononuclear cells of pregnant and bred, non-pregnant ewes. *Journal of Endocrinology*, 2001, vol. 170, R7-R11 [0005]
- Regulation of endometrial responsiveness to estrogen and progesterone by pregnancy recognition signals during the peri-implantation period. **BAZER et al.** Molecular and Cellular Aspects of Peri-implantation Processes. Springer-Verlag, New York, Inc, 1995, 27-47 [0005]
- **GODKIN et al.** *J. Reprod. Fert.*, 1982, vol. 65, 141-150 [0005]
- **SPENCER et al.** *Endocrinology*, 1995, vol. 136, 4932-4944 [0005]
- **SAMUEL.** *Virology*, 1991, vol. 183, 1-11 [0006] [0007]
- **LEFEVRE, F. et al.** *Biochimie*, 1998, vol. 80, 779-788 [0006]
- **STEWART et al.** *Endocrinology*, 2001, vol. 142, 98-107 [0006] [0045]
- **JOHNSON et al.** *Biol. Reprod.*, 2001, vol. 64, 1392-1399 [0006] [0045]
- **VALLET et al.** *J. Endocrinol.*, 1991, vol. 130, R1-4 [0006] [0045]
- **JOHNSON et al.** *Biol. Reprod.*, 2000, vol. 62, 622-627 [0006] [0045]
- **CHARLESTON ; STEWART.** *Gene*, 1993, vol. 137, 327-331 [0006] [0038] [0045]
- **OTT et al.** *Biol. Reprod.*, 1998, vol. 59, 784-794 [0006] [0007] [0037] [0038] [0045]
- **HALLER et al.** *Rev. Sci. Tech.*, 1998, vol. 17, 220-230 [0007]
- **SPENCER et al.** *Biol Reprod*, 1999, vol. 61, 464-470 [0007]
- **STELLFLUG et al.** *J. Anim. Sci.*, 2001, vol. 79, 568-573 [0035]
- **WILLARD et al.** *J. Anim. Sci.*, 1995, vol. 73, 960-966 [0039]
- **SPENCER et al.** *Biol. Reprod.*, 1998, vol. 58, 1154-1162 [0045]
- **SPENCER et al.** *Biol. Reprod.*, 1999, vol. 61, 464-470 [0045]

专利名称(译)	确定怀孕状况		
公开(公告)号	EP1410012B1	公开(公告)日	2013-02-13
申请号	EP2002756158	申请日	2002-06-10
申请(专利权)人(译)	爱达荷州研究基金会		
当前申请(专利权)人(译)	爱达荷州研究基金会		
[标]发明人	OTT TROY L		
发明人	OTT, TROY, L.		
IPC分类号	G01N33/68 C12Q1/68 G01N33/48 C12Q1/02 C12Q1/6876 G01N33/52 G01N33/53 G01N33/566		
CPC分类号	C12Q1/6876 C12Q2600/158 G01N33/689 G01N2333/4715 G01N2800/368		
优先权	60/299553 2001-06-19 US		
其他公开文献	EP1410012A4 EP1410012A1		
外部链接	Espacenet		

摘要(译)

用于确定动物是否未怀孕或在繁殖后怀孕的方法和试剂盒。在需要妊娠状态信息的动物中测定妊娠诱导蛋白的表达水平，并将该水平与未怀孕的动物中的水平进行比较。




Figure 1




Figure 2