

(19)



(11)

EP 2 423 310 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

17.12.2014 Bulletin 2014/51

(21) Application number: **10767122.4**

(22) Date of filing: **22.04.2010**

(51) Int Cl.:

C12N 15/09 ^(2006.01) **A61K 31/7088** ^(2006.01)
A61K 38/00 ^(2006.01) **A61K 48/00** ^(2006.01)
A61P 35/00 ^(2006.01) **C07K 14/82** ^(2006.01)
C12N 5/07 ^(2010.01) **C12Q 1/04** ^(2006.01)
C12Q 1/06 ^(2006.01) **G01N 33/68** ^(2006.01)
G01N 33/53 ^(2006.01)

(86) International application number:

PCT/JP2010/057149

(87) International publication number:

WO 2010/123065 (28.10.2010 Gazette 2010/43)

(54) **CANCER ANTIGEN HELPER PEPTIDE**

KREBSANTIGEN-HELPERPEPTID

PEPTIDE AUXILIAIRE ANTIGÉNIQUE DE CANCER

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK SM TR**

(30) Priority: **23.04.2009 JP 2009105286**

(43) Date of publication of application:

29.02.2012 Bulletin 2012/09

(73) Proprietor: **International Institute of Cancer
Immunology, Inc.**

Suita-shi, Osaka 564-0053 (JP)

(72) Inventor: **SUGIYAMA, Haruo**

**Minoo-shi
Osaka 562-0036 (JP)**

(74) Representative: **Von Kreisler Selting Werner -
Partnerschaft
von Patentanwälten und Rechtsanwälten mbB
Deichmannhaus am Dom
Bahnhofsvorplatz 1
50667 Köln (DE)**

(56) References cited:

**WO-A1-02/28414 WO-A1-2005/045027
WO-A1-2008/105462 WO-A2-01/25273
WO-A2-03/037060 JP-T- 2005 518 192
JP-T- 2009 511 637 US-A1- 2008 070 835**

- **KNIGHTS ASHLEY JOHN ET AL: "Prediction of an HLA-DR-binding peptide derived from Wilms' tumour 1 protein and demonstration of in vitro immunogenicity of WT1(124-138)-pulsed dendritic cells generated according to an optimised protocol", CANCER IMMUNOLOGY AND IMMUNOTHERAPY, SPRINGER-VERLAG, BERLIN, DE, vol. 51, no. 5, 1 July 2002 (2002-07-01), pages 271-281, XP002473128, ISSN: 0340-7004, DOI: 10.1007/S00262-002-0278-2**
- **FUJIKI F. ET AL.: 'Identification and characterization of a WT1 (Wilms Tumor Gene) protein-derived HLA-DRB1*0405-restricted 16-mer helper peptide that promotes the induction and activation of WT1-specific cytotoxic T lymphocytes' J.IMMUNOTHER. vol. 30, no. 3, 2007, pages 282 - 293, XP008167201**
- **KNIGHTS A.J. ET AL.: 'Prediction of an HLA-DR-binding peptide derived from Wilms' tumour 1 protein and demonstration of in vitro immunogenicity of WT1(124-138)-pulsed dendritic cells generated according to an optimised protocol' CANCER IMMUNOL.IMMUNOTHER. vol. 51, no. 5, 2002, pages 271 - 281, XP002473128**
- **FUMIHIRO FUJIKI ET AL.: 'WT1 Tokuiteki CD4+ Helper T Saibo o HLA-class II Kosokusei ni Yudo Dekiru WT1 Peptide no Dotei to sono Yuyosei no Kento' THE JAPANESE SOCIETY FOR IMMUNOLOGY GAKUJUTSU SHUKAI KIROKU vol. 35, 2005, page 187, XP008168936**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 423 310 B1

- GUO Y. ET AL.: 'Direct recognition and lysis of leukemia cells by WT1-specific CD4+ T lymphocytes in an HLA class II-restricted manner' BLOOD vol. 106, no. 4, 2005, pages 1415 - 8, XP055100496
- FUJIKI F. ET AL.: 'A WT1 protein-derived, naturally processed 16-mer peptide, WT1(332), is a promiscuous helper peptide for induction of WT1-specific Th1-type CD4(+) T cells' MICROBIOL.IMMUNOL. vol. 52, no. 12, 2008, pages 591 - 600, XP002590600
- LEHE C. ET AL.: 'The Wilms' tumor antigen is a novel target for human CD4+ regulatory T cells: implications for immunotherapy' CANCER RES. vol. 68, no. 15, 2008, pages 6350 - 6359, XP055082460

- MAY R.J. ET AL.: 'Peptide epitopes from the Wilms' tumor 1 oncoprotein stimulate CD4+ and CD8+ T cells that recognize and kill human malignant mesothelioma tumor cells' CLIN.CANCER RES. vol. 13, no. 15 PT, 2007, pages 4547 - 4555, XP002544528

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Description

Technical Field

[0001] The present invention relates to a WT1 helper peptide, a polynucleotide encoding the peptide, WT1-specific helper T cells induced by the peptide, a pharmaceutical composition for treating/preventing cancer comprising them and the like.

Background Art

[0002] The WT1 gene (Wilms' tumor 1 gene) is a gene identified as a causative gene of a Wilms' tumor which is a kidney cancer in childhood (Non-Patent Documents 1 and 2), and is a transcription factor having a zinc finger structure. At first, the WT1 gene was considered to be a cancer suppressor gene. However, subsequent investigation showed that the above gene rather serves as a cancer gene in hematopoietic organ tumors and solid cancers (Non-Patent Documents 3 to 6).

[0003] Since the WT1 gene is highly expressed in many malignant tumors, a WT1 gene product which is a self-protein having no mutation has been verified for existence or non-existence of immunogenicity in vivo. As a result, it has been shown that a protein derived from the WT1 gene highly expressed in tumor cells is fragmented by intracellular processing and the resulting peptide forms a complex with an MHC class I molecule which is displayed on the cell surface, and that cytotoxic T cells (hereinafter also referred to as CTLs) recognizing such a complex can be induced by WT1 peptide vaccination (Non-Patent Documents 7 to 9). It has also been shown that mice immunized with a WT1 peptide or a WT1 cDNA reject implanted WT1 gene-expressing tumor cells in a high rate (Non-Patent Documents 7 and 10) but normal tissues endogenously expressing the WT1 gene are not damaged by induced CTLs (Non-Patent Document 7). Heretofore, it has been strongly suggested that it is possible to induce WT1-specific CTLs in not only mice but also human, and that such CTLs have a cytotoxic activity against tumor cells highly expressing the WT1 gene, but have no cytotoxic activity against normal cells endogenously expressing the WT1 gene (Non-Patent Documents 7 and 10 to 14).

[0004] On the other hand, it is reported that the presence of helper T cells specific to a cancer antigen is important in order to induce the CTLs effectively (Non-Patent Document 15). The helper T cells (CD4-positive T cells) are induced, proliferated, and activated by recognizing a complex of an MHC class II molecule with an antigen peptide on antigen presenting cells. Activated helper T cells produce cytokines such as IL-2, IL-4, IL-5, IL-6, or an interferon (IFN), and promote proliferation, differentiation and maturation of B cells and other subsets of T cells. Thus, it is considered that an antigen peptide binding to an MHC class II molecule effectively activates CTLs and others through induction of helper T cells and enhances an immune function (Non-Patent Document 16). Heretofore, only an antigen peptide binding to HLA-DRB1*0401 and HLA-DRB1*0405 of an MHC class II molecule has been reported with respect to WT1 (Non-Patent Document 17 and Patent Document 1), and it was necessary to find antigen peptides to other subtypes.

Prior Art Documents

Non-patent Documents

[0005]

Patent Document 1: International Publication No. WO 2005/045027

Non-Patent Documents:

[0006]

Non-Patent Document 1: Daniel A. Haber et al., Cell. 1990 Jun 29; 61(7): 1257-69.
 Non-Patent Document 2: Call KM et al., Cell. 1990 Feb 9; 60(3) : 509-20.
 Non-Patent Document 3: Menke AL et al., Int Rev Cytol. 1998; 181: 151-212. Review.
 Non-Patent Document 4: Yamagami T et al., Blood. 1996 Apr 1; 87(7): 2878-84.
 Non-Patent Document 5: Inoue K et al., Blood. 1998 Apr 15; 91(8): 2969-76.
 Non-Patent Document 6: Tsuboi A et al., Leuk Res. 1999 May; 23(5): 499-505.
 Non-Patent Document 7: Oka Y et al., J Immunol. 2000 Feb 15; 164(4): 1873-80.
 Non-Patent Document 8: Melief CJ et al., Immunol Rev. 1995 Jun; 145: 167-77.
 Non-Patent Document 9: Ritz J, J Clin Oncol. 1994 Feb; 12(2): 237-8.
 Non-Patent Document 10: Tsuboi A et al., J Clin Immunol. 2000 May; 20(3): 195-202.

Non-Patent Document 11: Oka Y et al., Immunogenetics. 2000 Feb; 51(2): 99-107.
 Non-Patent Document 12: Ohminami H et al., Blood. 2000 Jan 1; 95(1): 286-93.
 Non-Patent Document 13: Gao L et al., Blood. 2000 Apr 1; 95(7): 2198-203.
 Non-Patent Document 14: Ohminami H et al., Blood. 2000 Jan 1; 95(1): 286-93.
 Non-Patent Document 15: Cancer. Res. 62: 6438, 2002
 Non-Patent Document 16: J. Immunol. Immunother., 24: 195, 2001
 Non-Patent Document 17: Cancer. Immunol. Immunother. 51: 271, 2002

[0007] WO2005/045027 discloses WT1 tumour derived peptides binding MHC class II with the sequence KRYFKLSH-LQMHSRKH. Cancer Immunol Immunother, vol, 51, 2002, 271-281 discloses WT1 tumour derived peptides binding MHC class II with the sequence PQQMGSDVRDLNALL.

Disclosure of the Invention

Problems to be Solved by the Invention

[0008] Accordingly, an object to be achieved by the present invention is to provide a peptide inducing WT1-specific helper T cells by binding to various MHC class II molecules, a polynucleotide encoding the peptide, WT1 helper T cells induced by the peptide, and a pharmaceutical composition for treating/preventing cancer comprising them.

Means for Solving the Problems

[0009] The present inventors have intensively studied to achieve the above object. As a result, they has found that a peptide having a portion of a sequence of contiguous amino acids encoding a WT1 protein functions as a cancer antigen helper peptide, in other words, the peptide is displayed on antigen presenting cells by binding to an MHC class II molecule and induces WT1-specific helper T cells, and showed that the peptide can be used in a pharmaceutical composition for treating/preventing cancer.

[0010] Thus, the present invention provides:

1. A peptide which consists of an amino acid sequence consisting of contiguous amino acids derived from a WT1 protein and induces WT1-specific helper T cells by binding to an MHC class II molecule, wherein the amino acid sequence is selected from:

- (a) the amino acid sequence depicted in SEQ ID NO:3;
- (b) the amino acid sequence depicted in SEQ ID NO:4;
- (c) the amino acid sequence depicted in SEQ ID NO:5; and
- (d) an amino acid sequence in which one amino acid only is substituted, deleted or added in the amino acid sequences depicted in (a) to (c).

2. The peptide according to 1, wherein the amino acid sequence is the amino acid sequence depicted in SEQ ID NO:3.

3. The peptide according to 1 or 2, wherein

- (i) the MHC class II molecule is selected from DRB1*0101, DRB1*0405, DRB1*0802, DRB1*0803, DRB1*0901, DRB1*1201, DRB1*1403, DRB1*1501, DRB1*1502, DPB1*0201, DPB1*0202, DPB1*0402, DPB1*0501, DPB1*0901, DQB1*0301, DQB1*0302, DQB1*0401, DQB1*0501, DQB1*0601, DQB1*0602, and DRB5*0102; or
- (ii) the MHC class II molecule is selected from DRB1*0101, DRB1*0405, DRB1*1502, DPB1*0201, DPB1*0202, and DQB1*0601.

4. A polynucleotide encoding the peptide according to any one of 1 to 3.

5. An expression vector comprising the polynucleotide according to 4.

6. An antibody against the peptide according to any one of 1 to 3.

7. A pharmaceutical composition for use in a method of treating or preventing cancer, comprising the peptide according to any one of 1 to 3, the polynucleotide according to 4, or the vector according to 5.

8. A peptide according to any one of 1 to 3, a polynucleotide according to 4, or a vector according to 5, for use in a method of treating or preventing cancer.

9. Antigen presenting cells which display the peptide according to any one of 1 to 3 through the MHC class II molecule according to 3.

10. A method for inducing antigen presenting cells, which comprises culturing immature antigen presenting cells in the presence of the peptide according to any one of 1 to 3, and inducing antigen presenting cells, which display the peptide through the MHC class II molecule according to 3, from the immature antigen presenting cells.

11. WT1-Specific helper T cells which are induced by the peptide according to any one of 1 to 3.

12. A method for inducing WT1-specific helper T cells, which comprises culturing peripheral blood mononuclear cells in the presence of the peptide according to any one of 1 to 3, and inducing WT1-specific helper T cells from the peripheral blood mononuclear cells.

13. A kit for inducing WT1-specific helper T cells, comprising, as an essential ingredient, the peptide according to any one of 1 to 3.

14. A kit for preventing or treating cancer, comprising, as an essential ingredient, the peptide according to any one of 1 to 3, the polynucleotide according to 4, or the vector according to 5.

15. A method for determining the presence or amount of WT1-specific helper T cells in a subject having the MHC class II molecule according to 3, said method comprising the steps of:

- (a) reacting the peptide according to any one of 1 to 3 with a sample derived from the subject; and then
- (b) determining the presence or amount of a cytokine contained in the sample.

Effects of the Invention

[0011] According to the present invention, it is possible to obtain WT1 helper peptides which bind to many types of MHC class II molecules such as DRB1*0101, DRB1*0405, DRB1*0802, DRB1*0803, DRB1*0901, DRB1*1201, DRB1*1403, DRB1*1501, DRB1*1502, DPB1*0201, DPB1*0202, DPB1*0402, DPB1*0501, DPB1*0901, DQB1*0301, DQB1*0302, DQB1*0401, DQB1*0501, DQB1*0601, DQB1*0602, and DRB5*0102. A pharmaceutical composition for treating/preventing cancer including them and the like may be provided. It becomes possible to induce WT1-specific helper T cells in vivo and in vitro in various subjects (in particular, most Japanese have the above molecules). Since WT1-specific helper T cells are induced by the present invention, it is also possible to activate T cells and B cells effectively in cancer highly expressing the WT1.

Brief Description of the Drawings

[0012]

Fig. 1 shows the results obtained by measuring cell proliferation after stimulating each peptide-specific T cell line, which was prepared by pulsing with each of three peptides (mWT1₃₅, mWT1₈₆, and mWT1₂₉₄), with each peptide. In the drawing, the symbol "-" shows no peptide stimulation.

Fig. 2 shows the results obtained by measuring cell proliferation after stimulating each peptide-specific T cell line, which was prepared by pulsing with three peptides (mWT1₃₅, mWT1₈₆, and mWT1₂₉₄), with each, corresponding peptide in the presence of an anti-MHC class I or II antibody. In the drawing, the symbol "-" shows no peptide stimulation. The symbol "cpm" in the ordinate shows counts per minute.

Fig. 3 shows the results obtained by measuring cell proliferation of each WT1 peptide-specific T cell line in response to C1498 cells, C1498 cells pulsed with three peptides (mWT1₃₅, mWT1₈₆, and mWT1₂₉₄), as well as C1498 cells having forced expression of a WT1 protein. The symbol "cpm" in the ordinate shows counts per minute.

Fig. 4 shows the results obtained by measuring an IFN- γ producing ability in each peptide-specific T cell line prepared by pulsing with three peptides (mWT1₃₅, mWT1₈₆, and mWT1₂₉₄).

Fig. 5 shows the results obtained by measuring a CTL cytotoxic activity of three peptides (mWT1₃₅, mWT1₈₆, and mWT1₂₉₄). ● shows the results of experiments carried out using RMA-S cells pulsed with a WT1₁₂₆ peptide (MHC class I-restricted peptide). ○ shows the results of experiments carried out using control RMA-S cells.

Fig. 6 shows a time-series schematic drawing when carrying out tumor implantation and immunization in a tumor implantation experiment. Immunization with an mWT1₃₅ helper peptide was carried out on the 7th, 14th and 21st days after subcutaneous implantation of WT1-expressing leukemia cells to mice, and dissection was carried out on the 29th day. Downward white arrows show time points at which a control (PBS) was intradermally administered (IFA/30 μ l). Downward black arrows show time points at which an mWT1₃₅ helper peptide was intradermally administered (50 μ M/IFA/30 μ l).

Fig. 7 shows tumor sizes in mice immunized with an mWT1₃₅ helper peptide and a proportion of disease-free mouse populations. In mice immunized with an mWT1₃₅ helper peptide, 4 of 10 mice were disease-free. On the other hand, in mice immunized with a control, there was no disease-free mouse in 9 mice.

Fig. 8 shows a disease-free survival rate in mice immunized with an mWT1₃₅ helper peptide.

Fig. 9 shows cytotoxic activity of CTLs in mice immunized with an mWT1₃₅ helper peptide. • shows the results of experiments carried out using RMAS cells pulsed with an mWT1₁₂₆ peptide (MHC I peptide). ○ shows the results of experiments carried out using control RMAS cells. The numerical in parenthesis represents a tumor size (mm).

Fig. 10 shows cytotoxic activity of mWT1-specific CTLs in control mice. ● shows the results of experiments carried out using RMAS cells pulsed with an mWT1₁₂₆ peptide (MHC I peptide). ○ shows the results of experiments carried out using control RMAS cells. The numeral in parenthesis represents a tumor size (mm).

Fig. 11 shows cytotoxic activity of mWT1₁₂₆ peptide-specific CTLs (left) and a proportion of WT1₁₂₆ tetramer-positive T cells (right) when an mWT1₃₅ peptide was administered.

Fig. 12 shows the results obtained by measuring cell proliferation by WT1₃₅ peptide stimulation in peripheral blood mononuclear cells of each healthy subject having MHC class II molecules.

Fig. 13 shows the results obtained by measuring cell proliferation when a Responder [PBMCs derived from a DRB1*0101/0405-, DPB1*0201/0402-, and DQB1*0401/0501-positive healthy subject (healthy subject A)] was treated with a Stimulator [PBMCs derived from a DRB1*0405/0901-, DPB1*0201/0501-, and DQB1*0303/0401-positive healthy subject (healthy subject B)]. The ordinate shows the amount of ³H-thymidine incorporated (cpm). The abscissa shows the types of various antibodies added (no antibody, anti-HLA-DR antibody, anti-HLA-DP antibody, and anti-HLA-DQ antibody).

Fig. 14 shows the results obtained by measuring cell proliferation when a Responder [PBMCs derived from a DRB1*0101/0405-, DPB1*0201/0402-, and DQB1*0401/0501-positive healthy subject (healthy subject A)] was treated with a Stimulator [PBMCs derived from a DRB1*0405/0803-, DPB1*0202/0501-, and DQB1*0401/0601-positive healthy subject (healthy subject G)]. The ordinate shows the amount of ³H-thymidine incorporated (cpm). The abscissa shows the types of various antibodies added (no antibody, anti-HLA-DR antibody, anti-HLA-DP antibody, and anti-HLA-DQ antibody).

Fig. 15 shows the results obtained by measuring cell proliferation when a Responder [PBMCs derived from a healthy subject having DRB1*0101/0405, DPB1*0201/0402, and DQB1*0401/0501 (healthy subject A)] was treated with a Stimulator [PBMCs derived from a DRB1*0101/0803-, DPB1*0501/0501-, and DQB1*0501/0601-positive healthy subject (healthy subject H)]. The ordinate shows the amount of ³H-thymidine incorporated (cpm). The abscissa shows the types of various antibodies added (no antibody, anti-HLA-DR antibody, anti-HLA-DP antibody, and anti-HLA-DQ antibody).

Fig. 16 shows the results obtained by measuring an IFN- γ producing ability when a Responder [PBMCs derived from a DRB1*0405/0803-, DPB1*0202/0501-, and DQB1*0401/0601-positive healthy subject (healthy subject G)] was treated with a Stimulator (L cells having a DQB1*0601 gene introduced). The ordinate shows a proportion of an amount of IFN- γ in T cells. The abscissa shows the presence or absence (+ or -) of a pulse with a WT1₃₅ peptide.

Fig. 17 shows the results obtained by measuring cell proliferation when a Responder [PBMCs derived from a DRB1*1502/1502-, DPB1*0201/0901-, and DQB1*0601/0601-positive healthy subject (healthy subject D)] was treated with a Stimulator (PBMCs derived from the same healthy subject as in the Responder). The ordinate shows the amount of ³H-thymidine incorporated (cpm). The abscissa shows the types of various antibodies added (no antibody, anti-HLA-DR antibody, anti-HLA-DP antibody, and anti-HLA-DQ antibody).

Fig. 18 shows the results obtained by measuring an IFN- γ producing ability when a Responder [PBMCs derived from a DRB1*0101/1501-, DPB1*0201/0402-, and DQB1*0501/0602-positive healthy subject (healthy subject I)] was treated with a Stimulator (PBMCs derived from the same healthy subject as in the Responder). The ordinate shows a proportion of an amount of IFN- γ in T cells. The abscissa shows the presence or absence (+ or -) of a pulse with a WT1₃₅ peptide.

[0013] The present disclosure relates to a peptide having an amino acid sequence consisting of amino acids derived from a mouse or human WT1 protein. The WT1 gene is highly expressed, for example, in hematopoietic organ tumors such as leukemia; myelodysplastic syndrome, multiple myeloma, and malignant lymphoma; solid cancers such as stomach cancer, bowel cancer, lung cancer, breast cancer, germ-cell cancer, liver cancer, skin cancer, bladder cancer, prostate cancer, uterus cancer, cervical cancer, and ovary cancer. Disclosed peptides may be present in cancer cells

expressing the WT1 gene in a large amount.

[0014] A peptide of the disclosure is a peptide which has an amino acid sequence consisting of contiguous amino acids derived from the human WT1 protein depicted in SEQ ID NO:2, retains an ability to bind to the MHC class II molecules as shown below, and has an ability to induce WT1-specific helper T cells. There is no particular limitation on the amino acid sequence and length of the peptide as long as the peptide has the above features. However, too long peptide is susceptible to a protease action, and too short peptide can not bind to a peptide accommodating groove well. The length of the peptide is preferably 10 to 25 amino acids, more preferably 15 to 21 amino acids, further preferably 16 to 20 amino acids, for example, of 17 amino acids, 18 amino acids, or 19 amino acids. Specific examples of the peptide of the present invention are those having the amino acid sequence depicted in SEQ ID NO:3; the amino acid sequence depicted in SEQ ID NO:4; and the amino acid sequence depicted in SEQ ID NO:5.

[0015] Also, the peptide of the present disclosure includes variants of the above peptides. The variants may contain, for example, a peptide selected from the group consisting of peptides having an amino acid sequence which has substitution, deletion or addition of several amino acids, for example, 1 to 9, preferably 1 to 5, 1 to 4, 1 to 3, more preferably 1 to 2 amino acids, further preferably one amino acid in one of the above amino acid sequences. Substitution of amino acids in peptides may be carried out at any positions and with any types of amino acids. Conservative amino acid substitution is preferred. For example, a Glu residue may be substituted with an Asp residue, a Phe residue with a Tyr residue, a Leu residue with an Ile residue, an Ala residue with a Ser residue, and a His residue with an Arg residue. Addition or deletion of amino acids may be carried out preferably at the N-terminus and the C-terminus in peptides, but may be carried out in an interior sequence. A preferred specific example of the peptide of the disclosure has the sequence of SEQ ID NO:3. In this regard, all the above peptides must retain an ability to bind to an MHC class II molecule and have an ability to induce WT1-specific helper T cells.

[0016] In this connection, the MHC class II molecule to which the peptide of the disclosure binds may belong to any subclass of HLA-DR, HLA-DQ, and HLA-DP. Preferably, the MHC class II molecule is one selected from the group consisting of DRB1*0101, DRB1*0405, DRB1*0802, DRB1*0803, DRB1*0901, DRB1*1201, DRB1*1403, DRB1*1501, DRB3*1502, DPB1*0201, DPB1*0202, DPB1*0402, DPB1*0501, DPB1*0901, DQB1*0301, DQB1*0302, DQB1*0401, DQB1*0501, DQB1*0601, DQB1*0602, and DRB5*0102. More preferably, the MHC class II molecule is DRB1*0101, DRB1*0405, DRB1*1403, DRB1*1502, DPB1*0201, DPB1*0202, DPB1*0901, DQB1*0301, DQB1*0601 or DRB5*0102, and most preferably, DRB1*0101, DRB1*0405, DRB1*1502, DPB1*0201, DPB1*0202, or DQB1*0601. In the present specification, a peptide which retains an ability to bind to an MHC class II molecule and has an ability to induce WT1-specific helper T cells is referred to as a WT1 helper peptide. Also, in the Examples described below, a peptide having the amino acid sequence depicted in SEQ ID NO:3 is referred to as a WT1₃₅ peptide, WT1₃₅ helper peptide or WT1₃₅ peptide.

[0017] Also, the peptide of the disclosure may be a peptide having an amino acid sequence consisting of contiguous amino acids derived from the mouse WT1 protein depicted in SEQ ID NO:1, and the above amino acid sequence may be a peptide (SEQ ID NO:6) in which an amino acid residue at position 9 in the amino acid sequence depicted in SEQ ID NO:4 is substituted with leucine; or a peptide (SEQ ID NO:7) in which an amino acid residue at position 11 in the amino acid sequence depicted in SEQ ID NO:5 is substituted with serine. Moreover, the peptide of the disclosure may contain a peptide selected from the group consisting of peptides having an amino acid sequence which has substitution, deletion or addition of several amino acids, for example, 1 to 9, preferably 1 to 5, 1 to 4, 1 to 3, more preferably 1 to 2 amino acids, further preferably one amino acid in the amino acid sequence depicted in SEQ ID NO:6 or SEQ ID NO:7. In the Examples described below, a peptide having the amino acid sequence depicted in SEQ ID NO:6 is also referred to as an mWT1₈₆ peptide or an mWT1₈₆ helper peptide, and a peptide having the amino acid sequence depicted in SEQ ID NO:7 as an mWT1₂₉₄ peptide or an mWT1₂₉₄ helper peptide.

[0018] The peptide of the disclosure may be derived from a WT1 protein, and may consist of the above sequence of contiguous amino acids or comprise the sequence. Thus, the peptide of the disclosure may be, for example, a peptide consisting of the above amino acid sequence itself, or a WT1 protein comprising the above amino acid sequence or a portion thereof. Also, the peptide of the disclosure may be that obtained by modification of the above amino acid sequence. Amino acid residues in the above amino acid sequence can be modified by a known method. Such modification may be, for example, esterification, alkylation, halogenation, phosphorylation, sulfonation, amidation and the like on a functional group in a side chain of an amino acid residue constituting a peptide. Also, it is possible to bind various substances to the N-terminus and/or C-terminus of a peptide containing the above amino acid sequence. For example, an amino acid, a peptide, an analog thereof and the like may be bound to the peptide. In case these substances are bound to the peptide of the disclosure they may be treated, for example, by an enzyme in vivo and the like or by a process such as intracellular processing so as to finally generate a peptide consisting of the above amino acid sequence, which is displayed on cell surface as a complex with an MHC class II molecule, thereby being able to obtain an induction effect of helper T cells. These substances may be those regulating solubility of the peptide of the present invention, those improving stability of the peptide such as protease resistance, those allowing specific delivery of the peptide of the disclosure, for example, to a given tissue or organ, or those having an enhancing action of an uptake efficiency of antigen

presenting cells or other action. Also, these substances may be those increasing an ability to induce CTLs, for example, helper peptides other than the peptide of the disclosure.

[0019] The modification of the peptide of the disclosure may be modification of an amino group on an N-terminal amino acid or of a carboxyl group on a C-terminal amino acid of the peptide. Modifying groups of an amino group on an N-terminal amino acid include, for example, one to three alkyl groups having 1 to 6 carbon atoms, phenyl groups, cycloalkyl groups, and acyl groups. Specific examples of the acyl group include an alkanoyl group having 1 to 6 carbon atoms, an alkanoyl group having 1 to 6 carbon atoms substituted with a phenyl group, a carbonyl group substituted with a cycloalkyl group having 5 to 7 carbon atoms, an alkylsulfonyl group having 1 to 6 carbon atoms, a phenylsulfonyl group, an alkoxy carbonyl group having 2 to 6 carbon atoms, an alkoxy carbonyl group substituted with a phenyl group, a carbonyl group substituted with a cycloalkoxy group having 5 to 7 carbon atoms, a phenoxy carbonyl group and the like. Peptides having modification of a carboxyl group on a C-terminal amino acid include, for example, esterified and amidated peptides. Specific examples of the ester include an alkyl ester having 1 to 6 carbon atoms, an alkyl ester having 0 to 6 carbon atoms substituted with a phenyl group, a cycloalkyl ester having 5 to 7 carbon atoms and the like, and specific examples of the amide include an amide, an amide substituted with one or two alkyl groups having 1 to 6 carbon atoms, an amide substituted with one or two alkyl groups having 0 to 6 carbon atoms substituted with a phenyl group, an amide forming a 5- to 7-membered azacycloalkane including a nitrogen atom of the amide group, and the like.

[0020] Also, the modification of the peptide of the disclosure may be carried out by binding amino acid residues to each other through a bond other than a peptide bond such as a carbon-carbon bond, a carbon-nitrogen bond, and a carbon-sulfur bond. Moreover, the peptide of the disclosure may contain one or more D-amino acids.

[0021] The above-mentioned peptides, variant peptides and modified peptides according to the disclosure are illustrative only, and those skilled in the art can easily assume, prepare, evaluate and use other variations of the above peptides.

[0022] The peptide of the disclosure can be synthesized using a method routinely used in the art or a modified method thereof. Such a synthesis method is disclosed, for example, in Peptide Synthesis, Interscience, New York, 1966; The Proteins, Vol. 2, Academic Press Inc., New York, 1976; Peptide Synthesis, Maruzen Co., Ltd., 1975; Basis and Experiments of Peptide Synthesis, Maruzen Co., Ltd., 1985; Development of Medicines (continuation), Vol. 14, Peptide Synthesis, Hirokawa Shoten Co., 1991 and the like. Also, the peptide of the disclosure can be prepared using a genetic engineering technique on the basis of information of a nucleotide sequence encoding the peptide of the disclosure. Such a genetic engineering technique is well known to those skilled in the art. Such a technique can be conducted according to a method described in literatures [Molecular Cloning, T. Maniatis et al., CSH Laboratory (1983); DNA Cloning, DM. Glover, IRL PRESS (1985)] as described above or a method described below, and other methods.

[0023] It is possible to determine whether the peptide of the disclosure or a candidate peptide thereof binds to the above MHC class II molecule and induces helper T cells, by a known method such as, for example, a method described in Cancer Immunol. Immunother. 51:271 (2002), or a method described in the Examples of the present specification, and other methods.

[0024] Since the peptide of the disclosure activates helper T cells (CD4-positive T cells), the peptide induces and maintains differentiation of CTLs and exerts an action of activating effector cells such as macrophages. Accordingly, it is possible to use the peptide of the disclosure for effective treatment or prevention of cancer.

[0025] In another aspect, the present disclosure relates to a polynucleotide encoding a WT1 helper peptide (hereinafter also referred to as a WT1 polynucleotide). The polynucleotide may be a DNA or an RNA. The base sequence of the polynucleotide can be determined on the basis of the amino acid sequence of the above WT1 helper peptide. The polynucleotide can be prepared, for example, by a method for DNA or RNA synthesis, a PCR method and the like.

[0026] The polynucleotide includes a polynucleotide which hybridizes with a complementary sequence of a polynucleotide encoding the peptide of the disclosure under a stringent condition and encodes a peptide having an activity comparable to that of the peptide of the disclosure. As to the term "hybridize under a stringent condition", hybridization used herein can be carried out according to a conventional method described, for example, in Molecular Cloning, 2nd edition, Sambrook J., Frisch E. F., Maniatis T., Cold Spring Harbor Laboratory press and the like. Also, the "stringent condition" includes, for example, a condition wherein a hybrid is formed in a solution containing $6 \times \text{SSC}$ ($10 \times \text{SSC}$ is a solution containing 1.5 M NaCl and 0.15 M trisodium citrate) and 50% formamide at 45°C and then washed with $2 \times \text{SSC}$ at 50°C (Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6) and the like.

[0027] In still another aspect, the disclosure relates to an expression vector comprising a polynucleotide (hereinafter also referred to as a WT1 expression vector). The type of expression vectors, other sequences contained in addition to the above polynucleotide sequence and the like can be appropriately selected depending on the type of hosts into which the expression vectors are introduced, the purpose of the introduction and the like. Examples of the expression vector include plasmids, phage vectors, virus vectors and the like. In case the host is Escherichia coli cells, examples of the vector include plasmid vectors such as pUC118, pUC119, pBR322, and pCR3, as well as phage vectors such as λ ZAPII, and λ gt11. In case the host is yeast cells, examples of the vector include pYES2, pYEUra3 and the like. In case the host is insect cells, examples of the vector pAcSGHisNT-A and the like. In case the host is animal cells, examples of the

vector include plasmid vectors such as pKCR, pCDM8, pGL2, pcDNA3.1, pRc/RSV, and pRc/CMV, virus vectors such as a retrovirus vector, an adenovirus vector, and an adeno-associated virus vector. The vector may optionally contain factors such as an expression-inducible promoter, a gene encoding a signal sequence, a marker gene for selection, and a terminator. Also, a sequence expressed as a fusion protein with thioredoxin, a His tag, GST (glutathione S-transferase) and the like may be added to the vector for easy isolation and purification. In this case, it is possible to use a GST-fused protein vector (pGEX4T, etc.) having a suitable promoter (lac, tac, trc, trp, CMV, SV40 early promoter, etc.) functional in host cells, a vector (pcDNA3.1/Myc-His, etc.) having a tag sequence such as Myc and His, and also a vector (pET32a) expressing a fusion protein with thioredoxin and a His tag and the like.

[0028] When the expression vector is administered to a subject to produce a WT1 helper peptide in vivo, WT1-specific helper T cells induced by the peptide produce various cytokines (for example, IL-2, IL-4, IL-5, IL-6, or an interferon (IFN), etc.), and promote proliferation, differentiation and maturation of B cells and other T cells. Accordingly, tumor cells which have an MHC class I molecule and highly express WT1 can be damaged specifically using the WT1 expression vector.

[0029] In another aspect, the disclosure relates to an antibody against a WT1 helper peptide (hereinafter also referred to as a WT1 antibody). The antibody may be either of a polyclonal antibody or a monoclonal antibody. A method for preparing such an antibody is already known, and the antibody of the present invention can be prepared according to such a conventional method as well (Current protocols in Molecular Biology, Ausubel et al. (ed.), 1987, John Wiley and Sons (pub.), Section 11.12-11.13, Antibodies; A Laboratory Manual, Lane, H. D. et al. (ed.), Cold Spring Harbor Laboratory Press (pub.), New York, 1989).

[0030] The present invention relates to a pharmaceutical composition for treating or preventing cancer, comprising a WT1 helper peptide, WT1 polynucleotide, or WT1 expression vector. The WT1 gene is highly expressed, for example, in hematopoietic organ tumors such as leukemia, myelodysplastic syndrome, multiple myeloma, and malignant lymphoma, as well as in solid cancers such as stomach cancer, bowel cancer, lung cancer, breast cancer, germ-cell cancer, liver cancer, skin cancer, bladder cancer, prostate cancer, uterus cancer, cervical cancer, and ovary cancer, and therefore, it is possible to use the pharmaceutical composition of the disclosure for treating or preventing cancer expressing the WT1 gene. When the pharmaceutical composition is administered to a subject having an MHC class II molecule, WT1-specific helper T cells induced by a WT1 helper peptide contained in the pharmaceutical composition produce various cytokines (for example, IL-2, IL-4, IL-5, IL-6, or an interferon (IFN), etc.), and promote proliferation, differentiation and maturation of B cells and other subsets of T cells. Accordingly, tumor cells which have an MHC class I molecule and highly express WT1 can be damaged specifically using the peptide.

[0031] A pharmaceutical composition may comprise, for example, a carrier, an excipient and the like, in addition to the above WT1 helper peptide, WT1 polynucleotide, or WT1 expression vector as an effective component. The WT1 helper peptide contained in the pharmaceutical composition induces WT1-specific helper T cells, and thus the pharmaceutical composition may comprise a suitable adjuvant or may be administered together with a suitable adjuvant in order to enhance the induction efficiency. Examples of preferred adjuvant include, but are not limited to, a Freund's complete or incomplete adjuvant, aluminium hydroxide and the like. Also, the pharmaceutical composition may also comprise a known cancer antigen peptide other than a WT1 helper peptide such as, for example, a WT1₁₂₆ peptide inducing WT1-specific CTLs, as an effective component (Oka et al, "Cancer immunotherapy targeting Wilms' tumor gene WT1 product", Journal of Immunology, 164:1873-1880, 2000; and Oka et al., "Human cytotoxic T-lymphocyte responses specific for peptides of the wild-type Wilms' tumor gene (WT1) product", Immunogenetics, 51: 99-107, 2000).

[0032] Moreover, a pharmaceutical composition may be administered in combination with a known cancer antigen peptide. For example, a known cancer antigen peptide, for example, a WT1₁₂₆ peptide can be administered before or after the administration of the pharmaceutical composition. The pharmaceutical composition has a feature that activates B cells or other T cells by inducing WT1-specific helper T cells, and therefore, it is possible to further enhance an activity of CTLs induced by administering a known cancer antigen peptide, and to remarkably increase therapeutic effects.

[0033] A method for administering the pharmaceutical composition can be appropriately selected depending on conditions such as the type of diseases, the state of subjects, and the targeted sites. Examples of the administration method includes, but are not limited to, intradermal administration, subcutaneous administration, intramuscular administration, intravenous administration, transnasal administration, oral administration and the like. Also, the administration method may be a lymphocyte therapy or a DC (dendritic cell) therapy. The amount of a peptide contained in the pharmaceutical composition, the form and administration frequency of the pharmaceutical composition and the like can be appropriately selected depending on conditions such as the type of diseases, the state of subjects, and the targeted sites. In general, the amount of a peptide administered per dose is 0.0001 mg to 1000 mg, and preferably 0.001 mg to 10,000 mg.

[0034] In another aspect, the disclosure relates to a method for treating or preventing cancer, which comprises administering an effective amount of a pharmaceutical composition to a subject having the above MHC class II molecule. Cancers to be treated or prevented may be any cancers as long as they express the WT1 gene and include, for example, hematopoietic organ tumors such as leukemia, myelodysplastic syndrome, multiple myeloma, and malignant lymphoma, as well as solid cancers such as stomach cancer, bowel cancer, lung cancer, breast cancer, germ-cell cancer, liver cancer, skin cancer, bladder cancer, prostate cancer, uterus cancer, cervical cancer, and ovary cancer.

[0035] In another aspect, the disclosure relates to use of the above WT1 helper peptide, WT1 polynucleotide, or WT1 expression vector for treating or preventing cancer.

[0036] In still another aspect, the disclosure relates to use of the WT1 helper peptide for preparing a pharmaceutical composition for treating or preventing cancer.

[0037] In still another aspect, the disclosure relates to use of the WT1 polynucleotide or WT1 expression vector for preparing a pharmaceutical composition containing the above WT1 polynucleotide or WT1 expression vector.

[0038] In another aspect, the disclosure relates to cells including the above WT1 helper peptide, WT1 polynucleotide, or WT1 expression vector. The cells can be prepared, for example, by transforming host cells such as *Escherichia coli* cells, yeast cells, insect cells, and animal cells using the above expression vector. Transformation of host cells with an expression vector can be carried out using various methods properly selected. The peptide can be prepared by culturing transformed cells, and recovering and purifying a WT1 helper peptide produced.

[0039] In still another aspect, the disclosure relates to antigen presenting cells (for example, dendritic cells, B-lymphocytes, macrophages, etc.), which display the above WT1 helper peptide through the above MHC class II molecule. The antigen presenting cells are induced by the above WT1 helper peptide. WT1-specific helper T cells are efficiently induced using the antigen presenting cells.

[0040] In still another aspect, the disclosure relates to a method for inducing antigen presenting cells which display a WT1 helper peptide through an MHC class II molecule, said method comprising culturing immature antigen presenting cells in the presence of a WT1 helper peptide, and inducing antigen presenting cells, which display the WT1 helper peptide through the above MHC class II molecule, from the immature antigen presenting cells. The immature antigen presenting cells refer to cells which can become antigen presenting cells such as, for example, dendritic cells, B-lymphocytes, and macrophages upon maturation. Subjects from which the immature antigen presenting cells derive may be any subjects as long as they have the above MHC class II molecule. Since the immature antigen presenting cells are contained, for example, in peripheral blood mononuclear cells and the like, such cells may be cultured in the presence of WT1 helper peptide.

[0041] In another aspect, the present disclosure relates to a method for treating or preventing cancer, which comprises administering antigen presenting cells, which display a WT1 helper peptide through the above MHC class II molecule, to a subject having the same molecule as the above MHC class II molecule. The administration method of the antigen presenting cells can be appropriately selected depending on conditions such as the type of diseases, the state of subjects, and the targeted sites. Examples of the method include, but are not limited to, intravenous administration, intradermal administration, subcutaneous administration, intramuscular administration, transnasal administration, oral administration and the like.

[0042] In still another aspect, the disclosure relates to a method for preventing or treating cancer by induction of antigen presenting cells which display a WT1 helper peptide through the above MHC class II molecule, said method comprising the steps of:

- (a) reacting a sample with a nucleotide sequence encoding an amino acid sequence (SEQ ID NO:2) of a WT1 protein or a nucleic acid having a partial sequence thereof or the above WT1 helper peptide;
- (b) obtaining antigen presenting cells which display a WT1 helper peptide contained in the sample through the above MHC class II molecule; and
- (c) administering the antigen presenting cells to a subject having the same molecule as the above MHC class II molecule.

[0043] Samples in the above method may be any samples as long as they have a possibility of containing lymphocytes or dendritic cells and include, for example, subject-derived samples such as blood, cell culture solutions and the like. The reaction in the above method may be carried out using a conventional technique, and preferably using electroporation. Obtainment of the antigen presenting cells can be carried out using a method known to those skilled in the art. Culturing conditions of cells in a sample in each step can be determined properly by those skilled in the art. The administration method of the antigen presenting cells may be as described above.

[0044] In further aspect, the disclosure relates to WT1-specific helper T cells induced by WT1 helper peptide. The helper T cells are induced, proliferated, and activated when recognizing a complex of a WT1 helper peptide with an MHC class II molecule. The activated WT1-specific helper T cells produce cytokines such as IL-2, IL-4, IL-5, IL-6, or an interferon (IFN), and promote proliferation, differentiation and maturation of B cells and other subsets of T cells. Accordingly, tumor cells which have an MHC class I molecule and highly express WT1 can be damaged specifically using the helper T cells.

[0045] In another aspect, the disclosure relates to a method for inducing WT1-specific helper T cells, which comprises culturing peripheral blood mononuclear cells in the presence of a WT1 helper peptide, and inducing the WT1-specific helper T cells from the peripheral blood mononuclear cells. Subjects from which the peripheral blood mononuclear cells derive may be any subjects as long as they have the above MHC class II molecule. By culturing the peripheral blood

mononuclear cells in the presence of a WT1 helper peptide, WT1-specific helper T cells are induced from precursor cells of helper T cells in the peripheral blood mononuclear cells. It is possible to treat or prevent hematopoietic organ tumors and solid cancers in a subject by administering the WT1-specific helper T cells to a subject having the above MHC class II molecule. In this connection, the peripheral blood mononuclear cells include immature antigen presenting cells which are precursor cells of antigen presenting cells (for example, precursor cells of dendritic cells, B-lymphocytes, macrophages, etc.) Since the immature antigen presenting cells are contained, for example, in peripheral blood mononuclear cells and the like, such cells may be cultured in the presence of WT1 helper peptide.

[0046] In still another aspect, the disclosure relates to a kit for inducing WT1-specific helper T cells, comprising WT1 helper peptide as an essential ingredient. Preferably, the kit is used in the above method for inducing WT1-specific helper T cells. The kit may comprise, for example, an obtaining means of peripheral blood mononuclear cells, an adjuvant, a reaction vessel and others, in addition to WT1 helper peptide. In general, the kit is accompanied with an instruction manual. It is possible to induce WT1-specific helper T cells efficiently using the kit.

[0047] In still another aspect, the disclosure relates to a method for treating or preventing cancer, which comprises administering WT1-specific helper T cells to a subject having MHC class II molecule. The administration method of the WT1-specific helper T cells can be appropriately selected depending on conditions such as the type of diseases, the state of subjects, and the targeted sites. Examples of the administration method includes, but are not limited to, intravenous administration, intradermal administration, subcutaneous administration, intramuscular administration, transnasal administration, oral administration and the like.

[0048] Furthermore, the disclosure relates to a kit for preventing or treating cancer, comprising WT1 helper peptide, WT1 polynucleotide, or WT1 expression vector as an essential ingredient. The kit is a kit characterized by induction of antigen presenting cells which display WT1 helper peptide through MHC class II molecule. Also, the kit may comprise, for example, an obtaining means of samples, a reaction vessel and others, in addition to the above essential ingredient. In general, the kit is accompanied with an instruction manual. Antigen presenting cells which display a WT1 helper peptide through MHC class II molecule can be obtained efficiently using the kit, and be used for treating or preventing cancer by their administration.

[0049] In another aspect, the disclosure relates to a method for determining the presence or amount of WT1-specific helper T cells in a subject having MHC class II molecule, said method comprising the steps of:

- (a) reacting a complex of WT1 helper peptide with MHC class II molecule with a sample derived from the subject; and then
- (b) determining the presence or amount of helper T cells recognizing the complex contained in the sample.

[0050] Samples derived from subjects may be any samples as long as they have a possibility of containing lymphocytes and include, for example, body fluids such as blood and lymph fluid, tissues and the like. The complex of WT1 helper T cells with an MHC class II molecule may be, for example, in the form of tetramer, pentamer and the like, for example, using a method known to those skilled in the art such as a biotin-streptavidin method. The presence or amount of helper T cells recognizing such a complex can be determined by a method known to those skilled in the art. In this aspect, the above complex may be labeled. As a label, known labels such as a fluorescent label and a radioactive label can be used. By labeling, the presence or amount of helper T cells can be determined simply and rapidly. Using a method of this aspect of the disclosure, it becomes possible to make a diagnosis, a prognosis and the like of cancer.

[0051] Accordingly, the disclosure also provides a composition comprising a complex of a WT1 helper peptide with MHC class II molecule for determining the presence or amount of WT1-specific helper T cells in a subject having the above MHC class II molecule.

[0052] Also, the disclosure provides a kit comprising a complex of a WT1 helper peptide with MHC class II molecule for determining the presence or amount of WT1-specific helper T cells in a subject having MHC class II molecule.

[0053] In still another aspect, the disclosure relates to a method for determining the presence or amount of WT1-specific helper T cells in a subject having MHC class II molecule, said method comprising the steps of:

- (a) reacting WT1 helper peptide with a sample derived from the subject; and then
- (b) determining the presence or amount of a cytokine contained in the sample.

[0054] Samples derived from subjects may be any samples as long as they have a possibility of containing lymphocytes and include, for example, peripheral blood mononuclear cells, blood, body fluids, tissues and others, and preferably peripheral blood mononuclear cells. The reaction in the above step (a) can be carried out by reacting WT1 helper peptide in the sample derived from a subject using a conventional technique. Culturing conditions of cells in a sample in each step can be determined properly by those skilled in the art. The presence or amount of a cytokine contained in a sample can be measured by a method known to those skilled in the art. The cytokine may be one capable of being induced by helper T cells such as interferon- γ and interleukin-10. The cytokine may be labeled. As a label, known labels such as a

fluorescent label and a radioactive label can be used. Using the presence or amount of the above cytokine as an indicator, it becomes possible to determine the presence or amount of WT1-specific helper T cells simply and rapidly.

[0055] In further aspect, the disclosure relates to a method for obtaining WT1-specific helper T cells using a complex of a WT1 helper peptide with MHC class II molecule, said method comprising the steps of:

- (a) reacting a sample with the complex; and
- (b) obtaining helper T cells which are contained in the sample and recognize the complex.

[0056] The complex of a WT1 helper peptide with MHC class II molecule is as described above. Samples may be any samples as long as they have a possibility of containing lymphocytes and include, for example, subject-derived samples such as blood, cell culture solutions and the like. Obtainment of helper T cells recognizing the complex can be carried out, for example, using a method known to those skilled in the art such as FACS and MACS. It is possible to culture the resulting WT1-specific helper T cells and to use them for treating or preventing various cancers.

[0057] Accordingly, the disclosure also relates to WT1-specific helper T cells, which can be obtained by a method for obtaining WT1-specific helper T cells using a complex of a WT1 helper peptide with MHC class II molecule.

[0058] Moreover, the disclosure relates to a kit for obtaining WT1-specific helper T cells, comprising a complex of a WT1 helper peptide with MHC class II molecule.

[0059] In still another aspect, the disclosure relates to a method for diagnosing cancer, which comprises using WT1-specific helper T cells, antigen presenting cells which display a WT1 helper peptide through MHC class II molecule, or WT1 antibody. Preferably, the WT1-specific helper T cells are used for the method for diagnosing cancer. For example, helper T cells, antigen presenting cells or antibody can be incubated with a sample derived from a subject having MHC class II molecule, or administered to a subject having MHC class II molecule, and then, for example, the location, site, amount and the like of the helper T cells, antigen presenting cells or antibody can be determined to diagnose cancer. The helper T cells, antigen presenting cells or antibody may be labeled. By labeling, it is possible to carry out the method for diagnosing cancer efficiently.

[0060] In still another aspect, the disclosure relates to a kit for diagnosing cancer, comprising WT1-specific helper T cells, antigen presenting cells which display a WT1 helper peptide through MHC class II molecule, or an antibody against a WT1 helper peptide or an antibody against a polynucleotide encoding the peptide, as an essential ingredient.

[0061] The invention will be described specifically and described in detail below by way examples, but they should not be construed as limiting the invention.

Example 1

Selection of candidate WT1 peptides binding to MHC class II molecules

[0062] In order to search peptide sequences which bind to MHC class II molecules, a method as shown by Rammensee et al. was used (Rammensee et al, Immunogenetics 41:178-228, 1995). Specifically, selection was carried out using the programs described in the right end column in the Tables together with the law of Rammensee et al. By the method, WT1₃₅ peptides were narrowed down to peptide sequences as shown in Tables 1 and 2, WT1₈₆ peptides to peptide sequences as shown in Tables 3 and 4, and WT1₂₉₄ peptides to peptide sequences as shown in Tables 5 and 6. The left end column in Tables 1 to 6 shows "suitability" as a candidate peptide sequence. The more the number of "○" is, the higher the suitability is in the law of Rammensee et al. No mark shows poor suitability. Also, the group of amino acids in parenthesis of the column of "candidate peptide sequences binding to MHC class II molecules" in Tables 1 to 6 shows that one amino acid can be selected from the group of amino acids listed in the parenthesis. For example, the description [FLM] means one amino acid selected from the group of amino acids F, L and M. Also, the description [VYI(AL)] means one amino acid selected from the group of amino acids V, Y and I, or one amino acid selected from the group of amino acids A and L. "x" shows that it may be any amino acid. The right end column shows "program name" of programs used for listing candidate peptide sequences.

[Table 1]

Suitability	Types of MHC class II molecules	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₋₃₅ peptides)	Program name
○○○	DPA1*0102/DPB1*0201	[FLMVWY]xxx[FLMY]xx[IAMV]	SYFPEITHI Marsh2000, Chiciz 1997
	DPA1*0103/DPB1*0201	[YLVFK]xx[DSQT]x[YFWV]xx[LV]	
○	DPA1*0103/DPB1*0201	[FLM]xxx[FL]xx[IA]	Marsh2000, Rotzsche 1994
○	DPA1*0201/DPB1*0401	[FLYM(IVA)]xxxxx[FLY(MVIA)]xx[VYI(AL)]	Marsh2000
○	DPA1*0201/DPB1*0401	[FLYMIVA]xxxxxx[FLY(MVIA)]xx[VYIAL]	SYFPEITHI
○	DPA1*0201/DPB1*0901	[RK]xxxx[AGL]xx[LV]	Marsh2000
	DPB1*0301	x[R]xxxxxxx	Marsh2000
○	DQA1:0101/DQB1*0501	[L]xxx[YFW]	Marsh2000
	DQA1:0102/DQB1*0602	xxxxx[LIV(APST)]xx[AGST(LIVP)]	Marsh2000
○	DQA1:0301/DQB1*0301	xx[AGST]x[AVLI]	Marsh2000
○	DQA1:0301/DQB1*0301	[DEW]xx[AGST]x[ACLM]	SYFPEITHI
	DQA1:0301/DQB1*0302	[RK]xxx[AG]xx[NED]	Marsh2000
○	DQA1:0301/DQB1*0302	[TSW]xxxxxxx[RE]	SYFPEITHI
○○○	DQA1:0501/DQB1*0201	[FWYILV]xx[DELVIH]x[PDE(H)][ED]x[FYWVILM]	Marsh2000
○○○	DQA1:0501/DQB1*0201	[FWYILV]xx[DELVIH]x[PDEHPA][DE]x[FYWVILM]	SYFPEITHI
○○○	DQA1:0501/DQB1*0301	[FYIMLV]xxx[VLIMY]x[YFMLV]	Marsh2000
○	DQA1:0501/DQB1*0301	[WYAVM]xx[A]x[AIVTS]xxx[QN]	SYFPEITHI
○○○	DQB1*0602	[AFCLMNQSTVWYDE]x[AFGLMNQSTWYCD]x[AFGLMNSTWY]x[LIVAPST]xx[ASTGLIVP]	SYFPEITHI
	DRB1*0101	[YFWLIMVA]xx[LMAIVN]x[AGSTCP]xx[LAINVFMW]	Marsh2000
○○○	DRB1*0101	[YVLFIAMW]xx[LAIVMNQ]x[AGSTCP]xx[LAINVY]	SYFPEITHI
○	DRB1*0102	[ILVM]xx[ALM]x[AGSTCP]xx[ILAMYW]	Marsh2000
	DRB1*0102	[ILVM]xx[ALM]x[AGSTP]xx[ILAMYW]	SYFPEITHI
○	DRB1*0301	[LIFMV]xx[D]x[KR(EQN)]x[L] [YLF]	Marsh2000, Malcherek 1993
	DRB1*0301	[LIFMV]xx[D]x[KREQN]xx[YLF]	SYFPEITHI Marsh2000, Chiciz 1992
○	DRB1*0301 or DRB3*0201	[FILVY]xx[DNQT]	
	DRB1*0401	[FLV]xxxxxxx[NQST]	Marsh2000
○○○	DRB1*0401 or DRB4	[FYWILVM]xx[FWILVADE]x[NSTQHR]xx[k]	Marsh2000, Friede 1996
○	DRB1*0401 or DRB4*0101	[FYW]xxxxxxx[ST]	Marsh2000, Verreck 1995
○○○	DRB1*0401 or DRB4*0101	[FYWILVM]xx[PWILVADE]x[NSTQHR][DEHKNQRSTYACI LMV]x[DEHKNQRSTYACILMV]	SYFPEITHI

55	Suitability	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₋₃₅ peptides)			Program name
50		Types of MHC class II molecules	Candidate peptide sequences binding to MHC class II molecules		
45		DRB1*0402 or DRB4	[VILM]xx[YFWILMRNH]x[INSTQHK]x[RKHNQPI]x[H]	Marsh2000	
		DRB1*0402 or DRB4	[VILM]xx[YFWILMRN]x[NQSTK]x[RKHNQPI]x[DEHLNQRS TYCILMVHA]	SYFPEITHI	
	○○	DRB1*0404 or DRB4	[VILM]xx[FYWILVMADE]x[NTSQR]xx[K]	Marsh2000	
	○○	DRB1*0404 or DRB4	[VILM]xx[FYWILVMADE]x[NTSQR]xx[K]	SYFPEITHI	
	○○○	DRB1*0405 or DRB4	[FYWVILM]xx[VILMDE]x[INSTQKD]xxx[DEQ]	Marsh2000	
	○○○	DRB1*0405 or DRB4	[FYWVILM]xx[VILMDE]x[INSTQKD]xxx[DEQ]	SYFPEITHI	
		DRB1*0405 or DRB4*0101	[Y]xxxx[VT]xxx[D]	Marsh2000	
	○○○	DRB1*0407 or DRB4	[FYW]xx[AVTK]x[NTDS]xxx[QN]	Marsh2000	
	○○○	DRB1*0407 or DRB4	[FYW]xx[AVTK]x[NTDS]xxx[QN]	SYFPEITHI	
5					

EP 2 423 310 B1

[Table 2]

Candidate peptide sequences binding to various MHC class II molecules (WT₁₃₅ peptides)

5	Suitability	Types of MHC class II molecules	Candidate peptide sequences binding to MHC class II molecules	Program name
		DRB1*0701	[FILVY] xxxx [NST]	Marsh2000
	○	DRB1*0701	[FYWILV]xx[DEHKNQRSTY]x[NST]x[VILYF]	SYFPEITHI
		DRB1*0801	[FILVY]xxx[HKR]	Marsh2000
10	○	DRB1*0901 or DRB4*0101	[YFWL]xx[AS]	Marsh2000
	○○○	DRB1*0901 or DRB4*0101	[WYFL]xx[AVS]	SYFPEITHI
	○○○	DRB1*1101	[YF]xx[LVMAFY]x[RKH]xx[AGSP]	Marsh2000
15	○○○	DRB1*1101	[WYF]xx[LVMAF]x[RKH]xx[AGSP]	SYFPEITHI
		DRB1*1101 or DRB3*0202	[YF]xxxx[RK]x[RK]	Marsh2000
	○○○	DRB1*1104	[ILV]xx[LVMAFY]x[RKH]xx[AGSP]	Marsh2000
	○○○	DRB1*1104	[ILV]xx[LVMAFY]x[RKH]xx[AGSP]	SYFPEITHI
20		DRB1*1201 or DRB3	[ILFY(V)]x[LNM(VA)]xx[VY(FIN)]xx[YFM(IV)]	Marsh2000
		DRB1*1201 or DRB3	[ILFYV]x[LMNVA]xx[VYFINA]xx[YFMIV]	SYFPEITHI
25	○	DRB1*1301	[IVF]xx[YWLVAM]x[RK]xx[YFAST]	Marsh2000
	○	DRB1*1301	[ILV]xx[LVMAWY]x[RK]xx[YFAST]	SYFPEITHI
		DRB1*1301 or DRB3*0101	[ILV]xxxx[RK]xx[Y]	Marsh2000
	○	DRB1*1302	[YFVAI]xx[YWLVAM]x[RK]xx[YFAST]	Marsh2000
30	○	DRB1*1302	[YFVAI]xx[LVMAWY]x[RK]xx[YFAST]	SYFPEITHI
		DRB1*1302 or DRB3*1301	[ILFY]xxxx[RK]xx[Y]	Marsh2000
	○	DRB1*1501	[LVI]xx[FYI]xx[ILVMF]	Marsh2000
	○	DRB1*1501	[LVI]xx[FYI]xx[ILVMF]	SYFPEITHI
35		DRB1*1501 or DRB5*0101	[ILV]xxxxxxxx[HKR]	Marsh2000
	○	DRB3*0202	[YFIL]xx[N]x[ASPDE]xx[LVISG]	Marsh2000
	○	DRB3*0202	[YFIL]xx[N]x[ASPDE]xx[LVISG]	SYFPEITHI
40	○	DRB3*0301	[ILV]xx[N]x[ASPDE]xx[ILV]	Marsh2000
	○	DRB3*0301	[ILV]xx[N]x[ASPDE]xx[ILV]	SYFPEITHI
	○○	DRB5*0101	[FYLM]xx[QVIM]xxxx[RK]	Marsh2000
	○○	DRB5*0101	[FYLM]xx[QVIM]xxxx[RK]	SYFPEITHI

45

50

55

[Table 3]

Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₋₁₈₆ peptides) Candidate peptide sequences binding to MHC class II molecules	Program name
	DPA1*0102/DPB1*0201	DPw2	unknown	Marsh2000
	DPA1*0102/DPB1*0201	DPw2	[FLMVWY]-x-x-x-[FLMY]-x-x-[IAMV]	SYFPEITHI
	DPA1*0103/DPB1*0201	DPw2	[FLM]-x-x-x-[FL]-x-x-[IA]	Marsh2000
○	DPA1*0103/DPB1*0201	DPw2	[YLVFK]-x-x-[DSQT]-x-[YFWV]-x-x-[LV]	Marsh2000
	DPA1*0103/DPB1*0201	DPw2	unknown	SYFPEITHI
	DPA1*0103/DPB1*0201	DPw2	unknown	SYFPEITHI
○○	DPA1*0201/DPB1*0401	DPw4	[FLYM(IVA)]-x-x-x-x-x-[FLY(MVIA)]-x-x-[VYI(AL)]	Marsh2000
○○	DPA1*0201/DPB1*0401	DPw4	[FLYM(IVA)]-x-x-x-x-x-[FLY(MVIA)]-x-x-x-x-x-[VYI(AL)]	SYFPEITHI
○○○	DPA1*0201/DPB1*0901	DPw4	[RK]-x-x-x-x-[AGL]-x-x-[LV]	Marsh2000
	DPB1*0301	DPw3	x-[R]-x-x-x-x-x-x-x-x	Marsh2000
	DPB1*0301	DPw3	unknown	SYFPEITHI
	DQA1*0101/DQB1*0501	DQ5(1)	[L]-x-x-x-[YFW]	Marsh2000a
	DQA1*0101/DQB1*0501	DQ5(1)	unknown	SYFPEITHI
○○○	DQA1*0102/DQB1*0602	DQ6(1)	x-x-x-x-x-[LIV(APST)]-x-x-[AGST(LIVP)]	Marsh2000
	DQA1*0301/DQB1*0301	DQ7(3)	x-x-[AGST]-x-[AVLI]	Marsh2000
○○○	DQA1*0301/DQB1*0301	DQ7(3)	[DEW]-x-x-[AGST]-x-[ACLM]	SYFPEITHI
○	DQA1*0301/DQB1*0302	DQ8(3)	[RK]-x-x-x-x-[AG]-x-x-[NED]	Marsh2000
	DQA1*0301/DQB1*0201	DQ8(3)	[TSW]-x-x-x-x-x-x-x-[RE]	SYFPEITHI
○○○	DQA1*0501/DQB1*0201	DQ2	[FWYILV]-x-x-[DELVIH]-x-[PDE(H)]-[ED]-x-[FYWVILM]	Marsh2000
○○○	DQA1*0501/DQB1*0201	DQ2	[FWYILV]-x-x-[DELVIH]-x-[PDEH-PA]-[DE]-x-[FWYILVM]	SYFPEITHI
○○○	DQA1*0501/DQB1*0301	DQ7(3)	[FYIMLV]-x-x-x-[VLIMY]-x-x-[YFM-LVI]	Marsh2000
○	DQA1*0501/DQB1*0301	DQ7(3)	[WYAVM]-x-x-[A]-x-[AIVTS]-x-x-x-[QN]	SYFPEITHI

(continued)

Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₈₆ peptides) Candidate peptide sequences binding to MHC class II molecules	Program name
○○○	DQB1*0602	DQ6(1)	[AFCILMNQSTVWYDE]-x-[AF- GILMNQSTVWYCDE]-[AFGILM- NQSTVWY]-x-[LIVAPST]-x- x-[ASTGLIVP] [YFWLIMVA]-x- x-[LMAIVN]-x-[AGSTCP]-x- x-[LAIVNFYMW] [YVLFIAMW]-x-x-[LAIVM- NQ]-x-[AGSTCP]-x-x-[LAIVNFY] [ILVM]-x-x-[ALM]-x-[AGSTCP]-x- x-[ILAMYW] [ILVM]-x-x-[ALM]-x-[AGSTP]-x- x-[ILAMYW] [LIFMV]-x- x-[D]-x-[KR(EQN)]-x-[L]-[YLF] [LIFMV]-x-x-[D]-x-[KREQN]-x- x-[YLF] [FILVY]-x-x-[DNQT] unknown [FLV]-x-x-x-x-x-x-[NQST] unknown [FYWILVM]-x-x-[FWIL- VADE]-x-[INSTQHR]-x-x-[K] [FYW]-x-x-x-x-x-x-[ST] [FYWILVM]-x-x-[PWIL- VADE]-x-[NSTQHR]-[DEHKN- QRSTYACILMV]-x-[DEHKN- QRSTYACILMV] [VILM]-x-x-[YFWILM- RNH]-x-[NSTQHK]-x-[RKHN- QP]-x-[H]	SYFPEITHI
○○○	DRB1*0101	DR1		Marsh2000
○○○	DRB1*0101	DR1		SYFPEITHI
○○○	DRB1*0102	DR1		Marsh2000
○○○	DRB1*0102	DR1		SYFPEITHI
○○	DRB1*0301	DR17(3)		Marsh2000
○○	DRB1*0301	DR17 (3)		SYFPEITHI
○○	DRB1*0301 or DRB3*0201	DR17 (3)		Marsh2000
	DRB1*0301 or DRB3*0201	DR17 (3)		SYFPEITHI
	DRB1*0401	DR4		Marsh2000
	DRB1*0401	DR4		SYFPEITHI
○○	DRB1*0401 or DRB4	DR4		Marsh2000
	DRB1*0401 or DRB4*0101	DR4		Marsh2000
○○○	DRB1*0401 or DRB4*0101	DR4		SYFPEITHI
○○	DRB1*0402 or DRB4	DR4		Marsh2000

Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to MHC class II molecules	Program name
○○○	DRB1*0402 or DRB4	DR4	<p>Candidate peptide sequences binding to MHC class II molecules (WT₁₋₁₈₆ peptides)</p> <p>[VILM]-x-x-[YFWILM-RN]-x-[NQSTK]-[RKHN-QP]-x-[DEHLNQRSTYCIILMVHA]</p>	SYFPEITHI

EP 2 423 310 B1

[Table 4]

Candidate peptide sequences binding to various MHC class II molecules (WT₁₈₆ peptides)

5	Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to MHC class II molecules	Program name
	○	DRB1*0404or DRB4	DR4	[VILM]-x-x[FYWILVMADE]-x-[NTSQR]-x-x-[K]	Marsh2000
10	○	DRB1*0404or DRB4	DR4	[VILM]-x-x[FYWILVMADE]-x-[NTSQR]-x-x-[K]	SYFPEITHI
	○	DRB1*0405or DRB4	DR4	[FYWVILM]-x-x-[VILMDE]-x-[NSTQKD]-x-x-x-[DEQ]	Marsh2000
15	○	DRB1*0405or DRB4	DR4	[FYWVILM]-x-x-[VILMDE]-x-[NSTQKD]-x-x-x-[DEQ]	SYFPEITHI
		DRB1*0405or DRB4*0101	DR4	[Y]-x-x-x-x-[VT]-x-x-x-[D]	Marsh2000
		DRB1*0405or DRB4*0101	DR4	unknown	SYFPEITHI
20		DRB1*0407or DRB4	DR4	[FYW]-x-x-[AVTK]-x-[NTDS]-x-x-x-[QN]	Marsh2000
		DRB1*0407or DRB4	DR4	[FYW]-x-x-[AVK]-x-[NTDS]-x-x-x-[QN]	SYFPEITHI
25	○	DRB1*0701	DR7	[FILVY]-x-x-x-x-[NST]	Marsh2000
		DRB1*0701	DR7	[FYWILV]-x-x-[DEHKNQRSTY]-x-[NST]-x-x-[VILYF]	SYFPEITHI
		DRB1*0801	DR8	[FILVY]-x-x-x-[HKR]	Marsh2000
		DRB1*0801	DR8	unknown	SYFPEITHI
30		DRB1*0901or DRB4*0101	DR9	[YFWL]-x-x-[AS]	Marsh2000
	○	DRB1*0901or DRB4*0101	DR9	[WYFL]-x-x-[AVS]	SYFPEITHI
	○○	DRB1*1101	DR11 (5)	[YF]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	Marsh2000
	○○	DRB1*1101	DR11 (5)	[WYF]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	SYFPEITHI
35		DRB1*1101or DRB3*0202	DR11 (5)	[YF]-x-x-x-x-[RK]-x-[RK]	Marsh2000
	○	DRB1*1104	DR11 (5)	[ILV]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	Marsh2000
	○	DRB1*1104	DR11 (5)	[ILV]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	SYFPEITHI
40		DRB1*1201or DRB3	DR12 (5)	[ILFY(V)]-x-[LNM(VA)]-x-x-[VY(FIN)]-x-x-[YFM(IV)]	Marsh2000
	○○	DRB1*1201or DRB3	DR12 (5)	[ILFYV]-x-[LMNVA]-x-x-[VYFINA]-x-x-[YFMIV]	SYFPEITHI
45	○	DRB1*1301	DR13 (6)	[IVF]-x-x-[YWLVAAM]-x-[RK]-x-x-[YFAST]	Marsh2000
	○	DRB1*1301	DR13 (6)	[IVF]-x-x-[LVMAWY]-x-[RK]-x-x-[YFAST]	SYFPEITHI
		DRB1*1301or DRB3*0101	DR13 (6)	[ILV]-x-x-x-x-[RK]-x-x-[Y]	Marsh2000
		DRB1*1301or DRB3*0101	DR13 (6)	unknown	SYFPEITHI
50	○	DRB1*1302	DR13(6)	[YFVAI]-x-x-[YWLVAAM]-x-[RK]-x-x-[YFAST]	Marsh2000
	○	DRB1*1302	DR13 (6)	[YFVAI]-x-x-[LVMAWY]-x-[RK]-x-x-[YFAST]	SYFPEITHI
		DRB1*1302or DRB3*0301	DR13 (6)	[ILFY]-x-x-x-x-[RK]-x-x-[Y]	Marsh2000
55		DRB1*1302or DRB3*0301	DR13(6)	unknown	SYFPEITHI
	○	DRB1*1501	DR15(2)	[LVI]-x-x-[FYI]-x-x-[ILVMF]	Marsh2000
	○	DRB1*1501	DR15(2)	[LVI]-x-x-[FYI]-x-x-[ILVMF]	SYFPEITHI

EP 2 423 310 B1

(continued)

Candidate peptide sequences binding to various MHC class II molecules (WT1₁₈₆ peptides)

5	Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to MHC class II molecules	Program name
	○	DRB1*1501 or DRB5*0101	DR15(2)	[ILV]-x-x-x-x-x-x-x-[HKR]	Marsh2000
	○○○	DRB3*0202	DR52	[YFIL]-x-x-[N]-x-[ASPDE]-x-x-[LVISG]	Marsh2000
10	○○○	DRB3*0202	DR52	[YFIL]-x-x-[N]-x-[ASPDE]-x-x-[LVISG]	SYFPEITHI
	○○○	DRB3*0301	DR52	[ILV]-x-x-[N]-x-[ASPDE]-x-x-[ILV]	Marsh2000
	○○○	DRB3*0301	DR52	[ILV]-x-x-[N]-x-[ASPDE]-x-x-[ILV]	SYFPEITHI
		DRB5*0101	DR51	[FYLM]-x-x-[QVIM]-x-x-x-x-[RK]	Marsh2000
15		DRB5*0101	DR51	[FYLM]-x-x-[QVIM]-x-x-x-x-[RK]	SYFPEITHI

20

25

30

35

40

45

50

55

[Table 5]

Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₂₉₄ peptides) Candidate peptide sequences binding to MHC class II molecules	Program name
○○	DPA1*0102/DPB1*0201	DPw2	unknown	Marsh2000
○○	DPA1*0102/DPB1*0201	DPw2	[FLMVWY]-x-x-x-[FLMY]-x-x-[AMV]	SYFPEITHI
○	DPA1*0103/DPB1*0201	DPw2	[FLM]-x-x-x-[FL]-x-x-[IA]	Marsh2000
○	DPA1*0103/DPB1*0201	DPw2	[YLVFK]-x-x-[DSQT]-x-[YFWV]-x-x-[LV]	Marsh2000
○○○	DPA1*0103/DPB1*0201	DPw2	unknown	SYFPEITHI
○○○	DPA1*0103/DPB1*0201	DPw2	unknown	SYFPEITHI
○○○	DPA1*0201/DPB1*0401	DPw4	[FLYM(IVA)]-x-x-x-x-x-[FLY(MVIA)]-x-x-[VYI(AL)]	Marsh2000
○○○	DPA1*0201/DPB1*0401	DPw4	[FLYMIVA]-x-x-x-x-x-[FLYM-VIA]-x-x-[VYIAL]	SYFPEITHI
○○○	DPA1*0201/DPB1*0901	DPw4	[RK]-x-x-x-x-[AGL]-x-x-[LV]	Marsh2000
○	DPA1*0201/DPB1*0901	DPw3	x-[R]-x-x-x-x-x-x-x	Marsh2000
	DPA1*0301	DPw3	unknown	SYFPEITHI
	DQA1*0101/DQB1*0501	DQ5 (1)	[L]-x-x-x-[YFW]	Marsh2000
	DQA1*0101/DQB1*0501	DQ5 (1)	unknown	SYFPEITHI
	DQA1*0102/DQB1*0602	DQ6 (1)	x-x-x-x-x-[LIV(APST)]-x-x-[AGST(LVP)]	Marsh2000
	DQA1*0301/DQB1*0301	DQ7 (3)	x-x-[AGST]-x-[AVLI]	Marsh2000
	DQA1*0301/DQB1*0301	DQ7 (3)	[DEW]-x-x-[AGST]-x-[ACLM]	SYFPEITHI
	DQA1*0301/DQB1*0302	DQ8 (3)	[RK]-x-x-x-x-[AG]-x-x-[NED]	Marsh2000
○	DQA1*0301/DQB1*0302	DQ8 (3)	[TSW]-x-x-x-x-x-x-x-[RE]	SYFPEITHI
○○○	DQA1*0501/DQB1*0201	DQ2	[FWYILV]-x-x-[DELVIH]-x-[PDE(H)]-[ED]-x-[FYWVILM]	Marsh2000
○○○	DQA1*0501/DQB1*0201	DQ2	[FWYILV]-x-x-[DELVIH]-x-[PDEH-PA]-[DE]-x-[FWYILVM]	SYFPEITHI
○○○	DQA1*0501/DQB1*0301	DQ7 (3)	[FYIMLV]-x-x-x-[VLIMY]-x-[YFM-LVI]	Marsh2000
○○○	DQA1*0501/DQB1*0301	DQ7 (3)	[WYAVM]-x-x-[A]-x-[AIVTS]-x-x-x-[QN]	SYFPEITHI

(continued)

Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₂₉₄ peptides) Candidate peptide sequences binding to MHC class II molecules	Program name
○○○	DQB1*0602	DQ6 (1)	[AFCILMNQSTVWYDE]-x-[AF- GILMNQSTVWYCDE]-[AFGILM- NQSTVWY]-x-[LIVAPST]-x- x-[ASTGLIVP] [YFWLIMVA]-x- x-[LMAIVN]-x-[AGSTCP]-x- x-[LAIVNFYMW] [YVLFIAMW]-x-x-[LAIVM- NQ]-x-[AGSTCP]-x-x-[LAIVNFY] [ILVM]-x-x-[ALM]-x-[AGSTCP]-x- x-[LAMYYW] [ILVM]-x-x-[ALM]-x-[AGSTP]-x- x-[LAMYYW] [LIFMV]-x- x-[D]-x-[KR(EQN)]-x-[L]-YLF] [LIFMV]-x-x-[D]-x-[KREQN]-x- x-[YLF] [FILVY]-x-x-[DNQT]	SYFPEITHI Marsh2000 SYFPEITHI Marsh2000 SYFPEITHI Marsh2000 SYFPEITHI Marsh2000 SYFPEITH Marsh2000 Marsh2000 SYFPEITHI Marsh2000
○○○	DRB1*0101	DR1		
○○○	DRB1*0101	DR1		
	DRB1*0102	DR1		
	DRB1*0102	DR1		
○○	DRB1*0301	DQ17 (3)		
○○	DRB1*0301	DQ17 (3)		
○○	DRB1*0301 or DRB3*0201	DQ17 (3)		
○○	DRB1*0301 or DRB3*0201	DQ17 (3)		
○○○	DRB1*0401	DR4		
○○○	DRB1*0401	DR4		
○○○	DRB1*0401 or DRB4	DR4		
○○	DRB1*0401 or DRB4*0101	DR4		
○○○	DRB1*0401 or DRB4*0101	DR4		
○○○	DRB1*0402 or DRB4	DR4		

	(continued)						
	Candidate peptide sequences binding to various MHC class II molecules (WT ₁₂₉₄ peptides)						
	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to MHC class II molecules	Program name			
○○○	DRB1*0402 or DRB4	DR4	[VILM]-x-x-[YFWILM]-	SYFPEITHI			
			RN]-x-[NQSTK]-[RKHN-				
			QP]-x-[DEHLNQRSTYCILMVHA]				

[Table 6]

Candidate peptide sequences binding to various MHC class II molecules (WT₁₂₉₄ peptides)

5	Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to MHC class II molecules	Program name
	○	DRB1*0404or DRB4	DR4	[VILM]-x-x-[FYWILVMADE]-x-[NTSQR]-x-x-[K]	Marsh2000
10	○	DRB1*0404or DRB4	DR4	[VILM]-x-x-[FYWILVMADE]-x-[NTSQR]-x-x-[K]	SYFPEITHI
	○○○	DRB1*0405or DRB4	DR4	[FYWVILM]-x-x-[VILMDE]-x-[NSTQKD]-x-x-x-[DEQ]	Marsh2000
15	○○○	DRB1*0405or DRB4	DR4	[FYWVILM]-x-x-[VILMDE]-x-[NSTQKD]-x-x-x-[DEQ]	SYFPEITHI
		DRB1*0405or DRB4*0101	DR4	[Y]-x-x-x-x-[VT]-x-x-x-[D]	Marsh2000
		DRB1*0405or DRB4*0101	DR4	unknown	SYFPEITHI
20	○○○	DRB1*0407or DRB4	DR4	[FYW]-x-x-[AVTK]-x-[NTDS]-x-x-x-[QN]	Marsh2000
	○○○	DRB1*0407or DRB4	DR4	[FYW]-x-x-[AVK]-x-[NTDS]-x-x-x-[QN]	SYFPEITHI
	○	DRB1*0701	DR7	[FILVY]-x-x-x-x-[NST]	Marsh2000
25	○	DRB1*0701	DR7	[FYWILV]-x-x-[DEHKNQRSTY]-x-[NST]-x-x-[VILYF]	SYFPEITHI
	○	DRB1*0801	DR8	[FILVY]-x-x-x-[HKR]	Marsh2000
		DRB1*0801	DR8	unknown	SYFPEITHI
	○	DRB1*0901or DRB4*0101	DR9	[YFWL]-x-x-[AS]	Marsh2000
30	○	DRB1*0901or DRB4*0101	DR9	[WYFL]-x-x-[AVS]	SYFPEITHI
	○○	DRB1*1101	DR11 (5)	[YF]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	Marsh2000
	○○	DRB1*1101	DR11 (5)	[WYF]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	SYFPEITHI
35	○○○	DRB1*1101or DRB3*0202	DR11 (5)	[YF]-x-x-x-x-[RK]-x-[RK]	Marsh2000
		DRB1*1104	DR11 (5)	[ILV]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	Marsh2000
		DRB1*1104	DR11(5)	[ILV]-x-x-[LVMAFY]-x-[RKH]-x-x-[AGSP]	SYFPEITHI
40	○○	DRB1*1201or DRB3	DR12(5)	[ILFY(V)]-x-[LNM(VA)]-x-x-[VY(FIN)]-x-x-[YFM(IV)]	Marsh2000
	○○	DRB1*1201or DRB3	DR12 (5)	[ILFYV]-x-[LMNVA]-x-x-[VYFINA]-x-x-[YFMIV]	SYFPEITHI
		DRB1*1301	DR13(6)	[IVF]-x-x-[YWLVAAM]-x-[RK]-x-x-[YFAST]	Marsh2000
45		DRB1*1301	DR13(6)	[ILV]-x-x-[LVMAWY]-x-[RK]-x-x-[YFAST]	SYFPEITHI
		DRB1*1301or DRB3*0101	DR13(6)	[ILV]-x-x-x-x-[RK]-x-x-[Y]	Marsh2000
		DRB1*1301or DRB3*0101	DR13(6)	unknown	SYFPEITHI
50	○	DRB1*1302	DR13(6)	[YFVAI]-x-x-[YWLVAAM]-x-[RK]-x-x-[YEAST]	Marsh2000
	○	DRB1*1302	DR13(6)	[YFVAI]-x-x-[LVMAWY]-x-[RK]-x-x-[YEAST]	SYFPEITHI
	○	DRB1*1302or DRB3*0301	DR13(6)	[ILFY]-x-x-x-x-[RK]-x-x-[Y]	Marsh2000
55		DRB1*1302or DRB3*0301	DR13(6)	unknown	SYFPEITHI
	○○	DRB1*1501	DR15(2)	[LVI]-x-x-[FYI]-x-x-[ILVMF]	Marsh2000
	○○	DRB1*1501	DR15(2)	[LVI]-x-x-[FYI]-x-x-[ILVMF]	SYFPEITHI

(continued)

Candidate peptide sequences binding to various MHC class II molecules (WT1₂₉₄ peptides)

5	Suitability	Types of MHC class II molecules	Serotype	Candidate peptide sequences binding to MHC class II molecules	Program name
		DRB1*1501 or DRB5*0101	DR15(2)	[ILV]-x-x-x-x-x-x-x-[HKR]	Marsh2000
	○○○	DRB3*0202	DR52	[YFIL]-x-x-[N]-x-[ASPDE]-x-x-[LVISG]	Marsh2000
10	○○○	DRB3*0202	DR52	[YFIL]-x-x-[N]-x-[ASPDE]-x-x-[LVISG]	SYFPEITHI
	○	DRB3*0301	DR52	[ILV]-x-x-[N]-x-[ASPDE]-x-x-[ILV]	Marsh2000
	○	DRB3*0301	DR52	[ILV]-x-x-[N]-x-[ASPDE]-x-x-[ILV]	SYFPEITHI
	○○○	DRB5*0101	DR51	[FYLM]-x-x-[QVIM]-x-x-x-x-[RK]	Marsh2000
15	○○○	DRB5*0101	DR51	[FYLM]-x-x-[QVIM]-x-x-x-x-[RK]	SYFPEITHI

[0063] Next, candidate WT1 peptides were visually selected from Tables 1 to 6, peptides as shown in the following Table 7 were identified as preferred candidate peptides for MHC class II molecules, and actual functions of these peptides were analyzed as described below.

[Table 7]

Identification of peptide candidates for mouse MHC class II molecules

	WT1 ₃₅	WAPVLDFAPPGASAYGSL (SEQ ID NO:3)	18 mer	MW 1819.01
	WT1 ₈₆	EQCLSAFTLHFSGQFTG (SEQ ID NO:6)	17 mer	MW 1944.01
25	WT1 ₂₉₄	FRGIQDVRVSGVAPTLVR (SEQ ID NO:7)	19 mer	MW 2126.48

Preparation of WT1 peptide-specific cell lines and measurement of cell proliferation ability

[0064] First, the above WT1 peptides were emulsified with a Freund's incomplete adjuvant (Montanide ISA 51), and mice were intradermally inoculated with each WT1 peptide in an amount corresponding to 100 μg/mouse. The immunization was carried out 3 times at intervals of one week, the spleen was removed after 1 week of the final immunization, and spleen cells were prepared. The spleen cells were stimulated 3 times at intervals of 10 days using spleen cells of non-immunized mice, which were pulsed with the same WT1 peptide as that used for immunization of each mouse and irradiated, as a stimulator. Then, the 4th stimulation was carried out using spleen cells of non-immunized mice, which were pulsed with each peptide (WT1₃₅, WT1₈₆ or WT1₂₉₄ peptide) as shown in Table 7 and irradiated, as a stimulator, and proliferation reaction in response to each stimulator was measured by a ³H incorporation experiment. An OVA (ovalbumin) peptide irrelevant to WT1 peptides was used as a control peptide. As a result, mouse spleen cells immunized with a WT1₃₅ peptide, a WT1₈₆ peptide or a WT1₂₉₄ peptide each responded to the stimulator pulsed with a WT1₃₅ peptide, a WT1₈₆ peptide or a WT1₂₉₄ peptide, and proliferated (Fig. 1A to 1C).

[0065] As described above, spleen cells were stimulated in vitro 3 times at intervals of 10 days using spleen cells of non-immunized mice, which were pulsed with each WT1 peptide and irradiated. When the 4th stimulation was then carried out using spleen cells of non-immunized mice, which were pulsed with each peptide described above and irradiated, as a stimulator, and proliferation reaction was measured, an MHC class I antibody (D^b antibody) or an MHC class II antibody (A^b antibody) was added to the culture solution and ³H incorporation was measured. As a result, the proliferation reaction in response to the stimulator pulsed with each of a WT1₃₅ peptide, a WT1₈₆ peptide and a WT1₂₉₄ peptide was suppressed by the addition of an MHC class II antibody (Fig. 2A to 2C).

[0066] As described above, spleen cells were stimulated in vitro 3 times at intervals of 10 days using spleen cells of non-immunized mice, which were pulsed with each WT1 peptide and irradiated. Then, the proliferation reaction was measured by ³H incorporation using irradiated C1498 cells not expressing any WT1 protein, C1498 cells pulsed with each of the above WT1 peptides, or C1498 cells expressing a WT1 protein by introduction of a WT1 gene, as a stimulator. As a result, the proliferation reaction was produced in response to C1498 cells pulsed with the same WT1 peptide as that used in immunization in vivo and C1498 cells expressing a WT1 protein by introduction of a WT1 gene (Fig. 3). This revealed that a WT1₃₅ peptide, a WT1₈₆ peptide and a WT1₂₉₄ peptide are produced by an intracellular process of an endogenous WT1 protein and displayed on an MHC class II molecule. From the above facts, it was shown that these three WT1 peptides are MHC class II-restricted WT1 peptides.

Measurement of IFN- γ producing ability

[0067] As described above, spleen cells were stimulated in vitro 3 times at intervals of 10 days using spleen cells of non-immunized mice, which were pulsed with each WT1 peptide and irradiated. Then, the concentration of IFN- γ and IL-4 in a culture supernatant was measured using an ELISA kit (BIOSOURCE Immunoassay Kit, Invitrogen). As a result, spleen cells of two separate mice responded to spleen cells of non-immunized mice which were pulsed with each WT1 peptide and irradiated, and produced interferon- γ but little interleukin-4 (Fig. 4). This revealed that these three types of WT1 peptides induce Th1 type of WT1-specific helper T cells.

Example 2

Measurement of WT1-specific cytotoxic T cells (CTLs)

[0068] Mice were immunized 3 times with a WT1₁₂₆ peptide (MHC class I) alone, a WT1₁₂₆ peptide (MHC class I) + a WT1₃₅ peptide (MHC class II), a WT1₁₂₆ peptide (MHC class I) + a WT1₈₆ peptide (MHC class II), or a WT1₁₂₆ peptide (MHC class I) + a WT1₂₉₄ peptide (MHC class II), and spleen cells of the mice were prepared. Then, the spleen cells were stimulated once in vitro using a WT1₁₂₆ peptide (MHC class I), and on 6th day, cytotoxic activity was measured using RMAS cells pulsed with a WT1₁₂₆ peptide (MHC class I) as a target cell. RMAS cells not pulsed with a WT1₁₂₆ peptide (MHC class I) were used as a control target cell. As a result, mouse spleen cells immunized with a WT1₁₂₆ peptide (MHC class I) + a WT1 helper peptide (MHC class II) induced WT1-specific cytotoxic T cells more strongly as compared with mouse spleen cells immunized with a WT1₁₂₆ peptide (MHC class I) alone (Fig. 5). This demonstrated that the three WT1 peptides (MHC class II) are a WT1-specific helper peptides.

Example 3

Tumor implantation experiment

[0069] WT1-expressing C1498 leukemia cells were subcutaneously implanted in mice in a proportion of 2.5×10^5 cells per mouse, and 50 μ g/mouse of a WT1₃₅ helper peptide was intradermally administered together with a Freund's incomplete adjuvant, once a week, 3 times in total, starting from one week after the implantation (Fig. 6). As a control, a physiological saline instead of the WT1₃₅ helper peptide was intradermally administered together with a Freund's incomplete adjuvant. The size of a subcutaneous tumor was measured over time, and the disease-free survival rate was calculated up to the 29th day after the subcutaneous implantation. As a result, the tumor expanded in all mice of the control group, while proliferation of the tumor was completely suppressed in 4 of 10 mice of the WT1₃₅ helper peptide (MHC class II)-immunized group (Fig. 7). Also, a significant difference ($p < 0.05$) was recognized between the WT1₃₅ helper peptide-immunized group and the control group (Fig. 8). This demonstrated that the WT1₃₅ helper peptide (MHC class II) is a WT1 peptide having an ability to induce tumor immunization in vivo.

[0070] Next, mice were dissected on the 29th day after starting the above experiment, the spleen was excised, and a WT1-specific immune response was analyzed using spleen cells. Briefly, the spleen was excised when mice of the WT1₃₅ helper peptide (MHC class II)-immunized group and the control group were dissected, and spleen cells were prepared. The spleen cells were stimulated once with a WT1₁₂₆ peptide (MHC class I), and on the 6th day after the stimulation, cytotoxic activity of the spleen cells was measured using RMAS cells pulsed with a WT1₁₂₆ peptide (MHC class I) as a target cell. As a control, the cytotoxic activity of the spleen cells was measured using RMAS cells as a target cell. As a result, WT1-specific cytotoxic T cells were induced in all 4 mice of the WT1₃₅ helper peptide (MHC class II)-immunized group (Fig. 9). On the other hand, the WT1-specific cytotoxic T cells were very weakly induced in 3 mice of the control group (Fig. 10). The WT1-specific cytotoxic T cells were not induced in one mouse. Also, it was clear that the induction of the WT1-specific cytotoxic T cells was lower as compared with the WT1₃₅ helper peptide (MHC class II)-immunized group (Figs. 9 and 10). This shows that WT1-specific helper T cells were induced by administration of a WT1₃₅ class II helper peptide, and by the action of the WT1-specific helper T cells, WT1-specific cytotoxic T cells induced by immune-responding to a WT1 protein expressed by implanted tumor cells were strongly amplified in vivo. Thus, the results demonstrated the usefulness of the WT1₃₅ helper peptide.

[0071] Next, specific cytolysis was analyzed in mice of the above WT1₃₅ helper peptide (MHC class II)-immunized group and control group. Briefly, the degree of cytolysis (%) obtained by subtracting the rate of cytolysis (%) when target cells were RMAS cells from the rate of cytolysis (%) when target cells were RMAS cells pulsed with a WT1₁₂₆ peptide (MHC class I) in the above experiments was used as the specific cytolysis (%) (Fig. 11, left). Also, the above-prepared spleen cells and a fluorescence-labeled WT1 tetramer (H-2Db WT1 Tetramer-RFMPNAPYL-PE) were incubated at 4°C for 20 minutes, washed, then stained with fluorescence-labeled CD3 and CD8 antibodies, again washed, and analyzed by FACS. CD3-positive, CD8-positive, and WT1 tetramer-positive cells were served as WT1-specific cytotoxic T cells

(Fig. 11, right). As a result, significantly high WT1-specific cytotoxic T cells ($p < 0.05$) were induced in spleen cells of mice of the WT1₃₅ helper peptide (MHC class II)-immunized group as compared with spleen cells of mice of the control group (Fig. 11).

Example 4

Measurement of proliferation ability of WT1-specific cytotoxic T cells (CTLs) in human

[0072] Peripheral blood mononuclear cells were prepared from 6 healthy subjects having DRB1, DPB1, DQB1 or DRB5 subclass molecules as shown in Fig. 12. To the peripheral blood mononuclear cells, a WT1₃₅ helper peptide was added, and the cells were cultured for one week. Then, the peripheral blood mononuclear cells were stimulated 4 times in total at intervals of one week using identical subject-derived peripheral blood mononuclear cells, which were pulsed with a WT1₃₅ helper peptide and irradiated, as a stimulator, and ³H incorporation was measured on the 6th day. In all 6 healthy subjects, peripheral blood mononuclear cells responded to a WT1₃₅ helper peptide and proliferated (Fig. 12). This showed that the WT1₃₅ helper peptide has a function to bind to the mentioned HLA class II molecules and cause proliferation reaction. In this connection, the mouse WT1₈₆ peptide and WT1₂₉₄ peptide differ from the human WT1₈₆ peptide (SEQ ID NO:4) and WT1₂₉₄ peptide (SEQ ID NO:5) in one amino acid at the positions enclosed in squares, as shown in Table 8.

[Table 8]

Differences in sequences between mouse and human WT1 ₃₅ , WT1 ₈₆ and WT1 ₂₉₄ peptides			
mWT1 ₃₅	Mouse	WAPVLDFAPPGASAYGSL (SEQ ID NO:3)	18-mer
hWT1 ₃₅	Human	WAPVLDFAPPGASAYGSL (SEQ ID NO:3)	
mWT1 ₈₆	Mouse	EQCLSAFTLHFSGQFTG (SEQ ID NO:6)	17-mer
hWT1 ₈₆	Human	EQCLSAFTVHFSGQFTG (SEQ ID NO:4)	
mWT1 ₂₉₄	Mouse	FRGIQDVRRVSGVAPTLVR (SEQ ID NO:7)	19-mer
hWT1 ₂₉₄	Human	FRGIQDVRRVPGVAPTLVR (SEQ ID NO:5)	

Example 5

HLA class II molecule-restrictedness of WT1₃₅ peptide

[0073] In order to determine HLA class II molecule-restrictedness of a WT1₃₅ peptide, a further experiment was carried out by a method well known to those skilled in the art as briefly described below. First, peripheral blood mononuclear cells (PBMCs) derived from a healthy subject [a DRB1*0101/0405-, DPB1*0201/0402-, and DQB1*0401/0501-positive healthy subject (hereinafter referred to as healthy subject A)] were stimulated 5 times with a WT1₃₅ peptide to prepare a Responder. Next, peripheral blood mononuclear cells (PBMCs) derived from another healthy subject different in an HLA class II type [a DRB1*0405/0901-, DPB1*0201/0501-, and DQB1*0303/0401-positive healthy subject (referred to as healthy subject B)] were pulsed with the WT1₃₅ peptide to prepare a Stimulator, and cell proliferation [the amount of ³H-thymidine incorporated (cpm)] was measured. The measurement was carried out under conditions of no addition of an antibody, addition of an anti-HLA-DR antibody (+a-DR), addition of an anti-HLA-DP antibody (+a-DP), or addition of an anti-HLA-DQ antibody (+a-DQ). A common HLA class II type, which is positive in both the Responder and Stimulator, shows restrictedness of the WT1₃₅ peptide. As a result of the experiments, it was shown that the WT1₃₅ peptide is DRB1*0405-restricted because the proliferation was suppressed under a condition having addition of an anti-DR antibody, and DRB1*0405 was common in healthy subjects A and B, as shown in Fig. 13.

[0074] Next, an experiment was carried out under the same conditions as those of the above experiment, except that PBMCs derived from a healthy subject different from healthy subject A [DRB1*0405/0803-, DPB1*0202/0501-, and DQB1*0401/0601-positive healthy subject (referred to as healthy subject G)] were used as a Stimulator. As a result, it was shown that the WT1₃₅ peptide is DRB1*0405-, DPB1*0201- and DPB1*0202-restricted because the proliferation was suppressed under a condition having addition of an anti-HLA-DR antibody or an anti-HLA-DP antibody, and DRB1*0405, DPB1*0201 and DPB1*0202 were common in healthy subject A and healthy subject G (DPB1*0201 and DPB1*0202 have a high analogy and are cross-reactive, and therefore, they are considered as a common molecule), as shown in Fig. 14.

[0075] Next, an experiment was carried out under the same conditions as those of the above experiment, except that PBMCs derived from a healthy subject different from healthy subject A [DRB1*0101/0803, DPB1*0501/0501-, DQB1*0501/0601-positive (referred to as healthy subject H)] were used as a Stimulator. As a result, it was shown that

the WT₁₃₅ peptide is DRB1*0101-restricted because the proliferation was suppressed under a condition having addition of an anti-HLA-DR antibody, and DRB1*0101 was common in healthy subject A and healthy subject H, as shown in Fig. 15.

[0076] Moreover, PBMCs derived from healthy subject G were used as a Responder and L cells having a DQB1*0601 gene introduced were used as a Stimulator, in order to determine restrictedness of a WT₁₃₅ peptide. The difference in an amount of IFN- γ produced in the presence or absence of a pulse with a WT₁₃₅ peptide of L cells was measured. A proportion of intracellular IFN- γ production was measured using FACS which is a technique well known to those skilled in the art. As a result, it was shown that the WT₁₃₅ peptide is DQB1*0601-restricted because the Responder was activated by the pulse with a WT₁₃₅ peptide on L cells, as shown in Fig. 16.

[0077] Next, an experiment was carried out as described above using PBMCs derived from the same healthy subject as a Responder and a Stimulator. The types of HLA class II molecules possessed by healthy subjects used in this experiment were summarized in Table 9 below.

[Table 9]

Types of HLA class II molecules possessed by healthy subjects used in this experiment

Healthy subject No.	DRB1	DPB1	DQB1
A	*0101/0405	*0201/0402	*0401/0501
B	*0405/0901	*0201/0501	*0303/0401
C	*0802/1201	*0201/0501	*0301/0302
D	*1502/1502	*0201/0901	*0601/0601
E	*0405/0901	*0202/0501	*0303/0401
F	*1403/1502	*0201/0901	*0301/0601
G	*0405/0803	*0202/0501	*0401/0601
H	*0101/0803	*0501/-	*0501/0601
I	*0101/1501	*0201/0402	*0501/0602

[0078] As a result, it was found that addition of an anti-DR antibody or an anti-DP antibody, when the experiment was carried out using PBMCs derived from healthy subjects A to E, resulted in reduction of the amount of ³H-thymidine incorporated (cpm), and therefore, in suppression of the proliferation. Also, addition of only an anti-DR antibody, when PBMCs derived from healthy subject F were used, resulted in suppression of the proliferation. Moreover, addition of only an anti-HLA-DP antibody, when PBMCs derived from healthy subject G were used, resulted in suppression of the proliferation. By an experiment using healthy subject A, it was shown that the WT₁₃₅ peptide is DRB1*0101- or 0405-restricted, and DPB1*0201- or 0402-restricted. By an experiment using healthy subject B, it was shown that the WT₁₃₅ peptide is DRB1*0405- or 0901-restricted, and DPB1*0201- or 0501-restricted. By an experiment using healthy subject C, it was shown that the WT₁₃₅ peptide is DRB1*0802- or 1201-restricted, and DPB1*0201- or 0501-restricted. By an experiment using healthy subject D, it was shown that the WT₁₃₅ peptide is DRB1*1502-restricted because the DRB1*1502 is a homozygote (Fig. 17). In addition, it was shown that the WT₁₃₅ peptide is DPB1*0201- or 0901-restricted. By an experiment using healthy subject E, it was shown that the WT₁₃₅ peptide is DRB1*0405- or 0901-restricted, and DPB1*0202- or 0501-restricted. By an experiment using healthy subject F, it was shown that the WT₁₃₅ peptide is DRB1*1403- or 1502-restricted. By an experiment using healthy subject G, it was shown that the WT₁₃₅ peptide is DPB1*0202- or 0501-restricted.

[0079] Also, the difference in an amount of IFN- γ produced in the presence or absence of a pulse with a WT₁₃₅ peptide was measured using PBMCs derived from healthy subject I as a Responder and a Stimulator. A proportion of intracellular IFN- γ production was measured using FACS which is a technique well known to those skilled in the art. As a result, a proportion of an amount of IFN- γ remarkably increased by the pulse with a WT₁₃₅ peptide (Fig. 18). This shows that the WT₁₃₅ peptide is restricted by any one of DRB1* 0101, DRB1*1501, DPB1* 0201, DPB1* 0402, DQB1*0501, and DQB1*0602.

Industrial Applicability

[0080] The present invention provides WT1 peptides which are restricted by many types of MHC class II molecules, polynucleotides encoding the peptides, pharmaceutical compositions containing them and the like. Thus, they can be utilized in the field of pharmaceuticals, for example, in the field of the development and production of prophylactic or therapeutic drugs for various hematopoietic organ tumors and solid tumors which highly express a WT1 gene.

[Sequence Listing Free Text]

SEQUENCE LISTING

5 **[0081]**

 <110> International Institute of Cancer Immunology, Inc.

 <120> Cancer antigen helper peptide

10 <130> 669667

 <150> JP 2009-105286

 <151> 2009-04-23

15 <160> 8

 <170> PatentIn version 3.2

20 <210> 1

 <211> 449

 <212> PRT

 <213> Mus musculus

25 <400> 1

30

35

40

45

50

55

EP 2 423 310 B1

Met Gly Ser Asp Val Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Ser
1 5 10 15

Ser Leu Gly Gly Gly Gly Gly Gly Cys Gly Leu Pro Val Ser Gly Ala
20 25 30

Arg Gln Trp Ala Pro Val Leu Asp Phe Ala Pro Pro Gly Ala Ser Ala
35 40 45

Tyr Gly Ser Leu Gly Gly Pro Ala Pro Pro Pro Ala Pro Pro Pro Pro
50 55 60

Pro Pro Pro Pro His Ser Phe Ile Lys Gln Glu Pro Ser Trp Gly Gly
65 70 75 80

Ala Glu Pro His Glu Glu Gln Cys Leu Ser Ala Phe Thr Leu His Phe
85 90 95

Ser Gly Gln Phe Thr Gly Thr Ala Gly Ala Cys Arg Tyr Gly Pro Phe
100 105 110

Gly Pro Pro Pro Pro Ser Gln Ala Ser Ser Gly Gln Ala Arg Met Phe
115 120 125

Pro Asn Ala Pro Tyr Leu Pro Ser Cys Leu Glu Ser Gln Pro Thr Ile
130 135 140

Arg Asn Gln Gly Tyr Ser Thr Val Thr Phe Asp Gly Ala Pro Ser Tyr
145 150 155 160

EP 2 423 310 B1

	Gly	His	Thr	Pro	Ser	His	His	Ala	Ala	Gln	Phe	Pro	Asn	His	Ser	Phe	
					165					170					175		
5	Lys	His	Glu	Asp	Pro	Met	Gly	Gln	Gln	Gly	Ser	Leu	Gly	Glu	Gln	Gln	
				180					185					190			
10	Tyr	Ser	Val	Pro	Pro	Pro	Val	Tyr	Gly	Cys	His	Thr	Pro	Thr	Asp	Ser	
			195					200					205				
15	Cys	Thr	Gly	Ser	Gln	Ala	Leu	Leu	Leu	Arg	Thr	Pro	Tyr	Ser	Ser	Asp	
		210					215					220					
20	Asn	Leu	Tyr	Gln	Met	Thr	Ser	Gln	Leu	Glu	Cys	Met	Thr	Trp	Asn	Gln	
	225					230					235					240	
25	Met	Asn	Leu	Gly	Ala	Thr	Leu	Lys	Gly	Met	Ala	Ala	Gly	Ser	Ser	Ser	
					245					250					255		
30	Ser	Val	Lys	Trp	Thr	Glu	Gly	Gln	Ser	Asn	His	Gly	Ile	Gly	Tyr	Glu	
				260					265					270			
35	Ser	Glu	Asn	His	Thr	Ala	Pro	Ile	Leu	Cys	Gly	Ala	Gln	Tyr	Arg	Ile	
			275					280					285				
40	His	Thr	His	Gly	Val	Phe	Arg	Gly	Ile	Gln	Asp	Val	Arg	Arg	Val	Ser	
		290					295					300					
45	Gly	Val	Ala	Pro	Thr	Leu	Val	Arg	Ser	Ala	Ser	Glu	Thr	Ser	Glu	Lys	
	305					310					315					320	
50	Arg	Pro	Phe	Met	Cys	Ala	Tyr	Pro	Gly	Cys	Asn	Lys	Arg	Tyr	Phe	Lys	
				325						330					335		
55	Leu	Ser	His	Leu	Gln	Met	His	Ser	Arg	Lys	His	Thr	Gly	Glu	Lys	Pro	
				340					345					350			
60	Tyr	Gln	Cys	Asp	Phe	Lys	Asp	Cys	Glu	Arg	Arg	Phe	Ser	Arg	Ser	Asp	
			355					360					365				
65	Gln	Leu	Lys	Arg	His	Gln	Arg	Arg	His	Thr	Gly	Val	Lys	Pro	Phe	Gln	
		370					375					380					
70	Cys	Lys	Thr	Cys	Gln	Arg	Lys	Phe	Ser	Arg	Ser	Asp	His	Leu	Lys	Thr	
	385					390					395					400	

EP 2 423 310 B1

His Thr Arg Thr His Thr Gly Lys Thr Ser Glu Lys Pro Phe Ser Cys
405 410 415

5 Arg Trp His Ser Cys Gln Lys Lys Phe Ala Arg Ser Asp Glu Leu Val
420 425 430

10 Arg His His Asn Met His Gln Arg Asn Met Thr Lys Leu His Val Ala
435 440 445

15 Leu

<210> 2
<211> 449
<212> PRT
<213> Homo sapiens

20

25 Met Gly Ser Asp Val Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Pro
1 5 10 15

Ser Leu Gly Gly Gly Gly Gly Cys Ala Leu Pro Val Ser Gly Ala Ala
20 25 30

30 Gln Trp Ala Pro Val Leu Asp Phe Ala Pro Pro Gly Ala Ser Ala Tyr
35 40 45

35 Gly Ser Leu Gly Gly Pro Ala Pro Pro Pro Ala Pro Pro Pro Pro
50 55 60

40 Pro Pro Pro Pro His Ser Phe Ile Lys Gln Glu Pro Ser Trp Gly Gly
65 70 75 80

45 Ala Glu Pro His Glu Glu Gln Cys Leu Ser Ala Phe Thr Val His Phe
85 90 95

Ser Gly Gln Phe Thr Gly Thr Ala Gly Ala Cys Arg Tyr Gly Pro Phe
100 105 110

50 Gly Pro Pro Pro Pro Ser Gln Ala Ser Ser Gly Gln Ala Arg Met Phe
115 120 125

55 Pro Asn Ala Pro Tyr Leu Pro Ser Cys Leu Glu Ser Gln Pro Ala Ile
130 135 140

EP 2 423 310 B1

	Arg	Asn	Gln	Gly	Tyr	Ser	Thr	Val	Thr	Phe	Asp	Gly	Thr	Pro	Ser	Tyr	145	150	155									160
5	Gly	His	Thr	Pro	Ser	His	His	Ala	Ala	Gln	Phe	Pro	Asn	His	Ser	Phe		165	170								175	
10	Lys	His	Glu	Asp	Pro	Met	Gly	Gln	Gln	Gly	Ser	Leu	Gly	Glu	Gln	Gln		180	185							190		
15	Tyr	Ser	Val	Pro	Pro	Pro	Val	Tyr	Gly	Cys	His	Thr	Pro	Thr	Asp	Ser		195	200						205			
20	Cys	Thr	Gly	Ser	Gln	Ala	Leu	Leu	Leu	Arg	Thr	Pro	Tyr	Ser	Ser	Asp		210	215					220				
25	Asn	Leu	Tyr	Gln	Met	Thr	Ser	Gln	Leu	Glu	Cys	Met	Thr	Trp	Asn	Gln		225	230					235			240	
30	Met	Asn	Leu	Gly	Ala	Thr	Leu	Lys	Gly	Val	Ala	Ala	Gly	Ser	Ser	Ser		245	250						255			
35	Ser	Val	Lys	Trp	Thr	Glu	Gly	Gln	Ser	Asn	His	Ser	Thr	Gly	Tyr	Glu		260	265						270			
40	Ser	Asp	Asn	His	Thr	Thr	Pro	Ile	Leu	Cys	Gly	Ala	Gln	Tyr	Arg	Ile		275	280						285			
45	His	Thr	His	Gly	Val	Phe	Arg	Gly	Ile	Gln	Asp	Val	Arg	Arg	Val	Pro		290	295					300				
50	Gly	Val	Ala	Pro	Thr	Leu	Val	Arg	Ser	Ala	Ser	Glu	Thr	Ser	Glu	Lys		305	310					315			320	
55	Arg	Pro	Phe	Met	Cys	Ala	Tyr	Pro	Gly	Cys	Asn	Lys	Arg	Tyr	Phe	Lys		325	330						335			
	Leu	Ser	His	Leu	Gln	Met	His	Ser	Arg	Lys	His	Thr	Gly	Glu	Lys	Pro		340	345						350			
	Tyr	Gln	Cys	Asp	Phe	Lys	Asp	Cys	Glu	Arg	Arg	Phe	Ser	Arg	Ser	Asp		355	360						365			
	Gln	Leu	Lys	Arg	His	Gln	Arg	Arg	His	Thr	Gly	Val	Lys	Pro	Phe	Gln		370	375						380			

EP 2 423 310 B1

Cys Lys Thr Cys Gln Arg Lys Phe Ser Arg Ser Asp His Leu Lys Thr
385 390 395 400

5 His Thr Arg Thr His Thr Gly Lys Thr Ser Glu Lys Pro Phe Ser Cys
405 410 415

10 Arg Trp Pro Ser Cys Gln Lys Lys Phe Ala Arg Ser Asp Glu Leu Val
420 425 430

15 Arg His His Asn Met His Gln Arg Asn Met Thr Lys Leu Gln Leu Ala
435 440 445

Leu

20 <210> 3
<211> 18
<212> PRT
<213> Homo sapiens

25 <400> 3

Trp Ala Pro Val Leu Asp Phe Ala Pro Pro Gly Ala Ser Ala Tyr Gly
1 5 10 15

30

Ser Leu

35 <210> 4
<211> 17
<212> PRT
<213> Homo sapiens

<400> 4

40

Glu Gln Cys Leu Ser Ala Phe Thr Val His Phe Ser Gly Gln Phe Thr
1 5 10 15

45

Gly

50 <210> 5
<211> 19
<212> PRT
<213> Homo sapiens

<400> 5

55

Phe Arg Gly Ile Gln Asp Val Arg Arg Val Pro Gly Val Ala Pro Thr
1 5 10 15

Leu Val Arg

<210> 6
 <211> 17
 <212> PRT
 <213> Mus musculus

<400> 6

Glu Gln Cys Leu Ser Ala Phe Thr Leu His Phe Ser Gly Gln Phe Thr
 1 5 10 15

Gly

<210> 7
 <211> 19
 <212> PRT
 <213> Mus musculus

<400> 7

Phe Arg Gly Ile Gln Asp Val Arg Arg Val Ser Gly Val Ala Pro Thr
 1 5 10 15

Leu Val Arg

<210> 8
 <211> 9
 <212> PRT
 <213> Mus musculus

<400> 8

Arg Met Phe Pro Asn Ala Pro Tyr Leu
 1 5

Claims

1. A peptide which consists of an amino acid sequence consisting of contiguous amino acids derived from a WT1 protein and induces WT1-specific helper T cells by binding to an MHC class II molecule, wherein the amino acid sequence is selected from:
 - (a) the amino acid sequence depicted in SEQ ID NO:3;
 - (b) the amino acid sequence depicted in SEQ ID NO:4;
 - (c) the amino acid sequence depicted in SEQ ID NO:5; and
 - (d) an amino acid sequence in which one amino acid only is substituted, deleted or added in the amino acid sequences depicted in (a) to (c).
2. The peptide according to claim 1, wherein the amino acid sequence is the amino acid sequence depicted in SEQ ID NO:3.

3. The peptide according to claim 1 or 2, wherein

- (i) the MHC class II molecule is selected from DRB1*0101, DRB1*0405, DRB1*0802, DRB1*0803, DRB1*0901, DRB1*1201, DRB1*1403, DRB1*1501, DRB1*1502, DPB1*0201, DPB1*0202, DPB1*0402, DPB1*0501, DPB1*0901, DQB1*0301, DQB1*0302, DQB1*0401, DQB1*0501, DQB1*0601, DQB1*0602, and DRB5*0102; or
(ii) the MHC class II molecule is selected from DRB1*0101, DRB1*0405, DRB1*1502, DPB1*0201, DPB1*0202, and DQB1*0601.

4. A polynucleotide encoding the peptide according to any one of claims 1 to 3.

5. An expression vector comprising the polynucleotide according to claim 4.

6. An antibody against the peptide according to any one of claims 1 to 3.

7. A pharmaceutical composition for use in a method of treating or preventing cancer, comprising the peptide according to any one of claims 1 to 3, the polynucleotide according to claim 4, or the vector according to claim 5.

8. A peptide according to any one of claims 1 to 3, a polynucleotide according to claim 4, or a vector according to claim 5, for use in a method of treating or preventing cancer.

9. Antigen presenting cells which display the peptide according to any one of claims 1 to 3 through the MHC class II molecule according to claim 3.

10. A method for inducing antigen presenting cells, which comprises culturing immature antigen presenting cells in the presence of the peptide according to any one of claims 1 to 3, and inducing antigen presenting cells, which display the peptide through the MHC class II molecule according to claim 3, from the immature antigen presenting cells.

11. WT1-Specific helper T cells which are induced by the peptide according to any one of claims 1 to 3.

12. A method for inducing WT1-specific helper T cells, which comprises culturing peripheral blood mononuclear cells in the presence of the peptide according to any one of claims 1 to 3, and inducing WT1-specific helper T cells from the peripheral blood mononuclear cells.

13. A kit for inducing WT1-specific helper T cells, comprising, as an essential ingredient, the peptide according to any one of claims 1 to 3.

14. A kit for preventing or treating cancer, comprising, as an essential ingredient, the peptide according to any one of claims 1 to 3, the polynucleotide according to claim 4, or the vector according to claim 5.

15. A method for determining the presence or amount of WT1-specific helper T cells in a subject having the MHC class II molecule according to claim 3, said method comprising the steps of:

- (a) reacting the peptide according to any one of claims 1 to 3 with a sample derived from the subject; and then
(b) determining the presence or amount of a cytokine contained in the sample.

Patentansprüche

1. Peptid, das aus einer Aminosäuresequenz, die aus aufeinanderfolgenden Aminosäuren besteht, welche von einem WT1-Protein abgeleitet sind, besteht und WT1-spezifische Helfer-T-Zellen durch Bindung an ein MHC-Klasse-II-Molekül induziert, wobei die Aminosäuresequenz ausgewählt ist aus:

- (a) der in SEQ ID Nr. 3 gezeigten Aminosäuresequenz;
(b) der in SEQ ID Nr. 4 gezeigten Aminosäuresequenz;
(c) der in SEQ ID Nr. 5 gezeigten Aminosäuresequenz; und
(d) einer Aminosäuresequenz, bei der gegenüber den in (a) bis (c) gezeigten Aminosäuresequenzen nur eine einzige Aminosäure substituiert, deletiert oder addiert ist.

2. Peptid gemäß Anspruch 1, wobei es sich bei der Aminosäuresequenz um die in SEQ ID Nr. 3 gezeigte Aminosäuresequenz handelt.

3. Peptid gemäß Anspruch 1 oder 2, wobei

(i) das MHC-Klasse-II-Molekül aus DRB1*0101, DRB1*0405, DRB1*0802, DRB1*0803, DRB1*0901, DRB1*1201, DRB1*1403, DRB1*1501, DRB1*1502, DPB1*0201, DPB1*0202, DPB1*0402, DPB1*0501, DPB1*0901, DQB1*0301, DQB1*0302, DQB1*0401, DQB1*0501, DQB1*0601, DQB1*0602 und DRB5*0102 ausgewählt ist; oder

(ii) das MHC-Klasse-II-Molekül aus DRB1*0101, DRB1*0405, DRB1*1502, DPB1*0201, DPB1*0202 und DQB1*0601 ausgewählt ist.

4. Polynucleotid, das das Peptid gemäß einem der Ansprüche 1 bis 3 codiert.

5. Expressionsvektor, der das Polynucleotid gemäß Anspruch 4 umfasst.

6. Antikörper gegen das Peptid gemäß einem der Ansprüche 1 bis 3.

7. Pharmazeutische Zusammensetzung zur Verwendung in einem Verfahren zur Behandlung oder Prävention von Krebs, umfassend das Peptid gemäß einem der Ansprüche 1 bis 3, das Polynucleotid gemäß Anspruch 4 oder den Vektor gemäß Anspruch 5.

8. Peptid gemäß einem der Ansprüche 1 bis 3, Polynucleotid gemäß Anspruch 4 oder Vektor gemäß Anspruch 5 zur Verwendung in einem Verfahren zur Behandlung oder Prävention von Krebs.

9. Antigen-präsentierende Zellen, die das Peptid gemäß einem der Ansprüche 1 bis 3 über das MHC-Klasse-II-Molekül gemäß Anspruch 3 präsentieren.

10. Verfahren zum Induzieren von Antigen-präsentierenden Zellen, umfassend das Kultivieren von unreifen Antigen-präsentierenden Zellen in Gegenwart des Peptids gemäß einem der Ansprüche 1 bis 3 und Induzieren von Antigen-präsentierenden Zellen, die das Peptid über das MHC-Klasse-II-Molekül gemäß Anspruch 3 präsentieren, ausgehend von den unreifen Antigen-präsentierenden Zellen.

11. WT1-spezifische Helfer-T-Zellen, die durch das Peptid gemäß einem der Ansprüche 1 bis 3 induziert sind.

12. Verfahren zum Induzieren von WT1-spezifischen Helfer-T-Zellen, umfassend das Kultivieren von mononukleären Zellen des peripheren Bluts in Gegenwart des Peptids gemäß einem der Ansprüche 1 bis 3 und Induzieren von WT1-spezifischen Helfer-T-Zellen ausgehend von den mononukleären Zellen des peripheren Bluts.

13. Kit zum Induzieren von WT1-spezifischen Helfer-T-Zellen, umfassend das Peptid gemäß einem der Ansprüche 1 bis 3 als wesentlichen Bestandteil.

14. Kit zur Prävention oder Behandlung von Krebs, umfassend das Peptid gemäß einem der Ansprüche 1 bis 3, das Polynucleotid gemäß Anspruch 4 oder den Vektor gemäß Anspruch 5 als wesentlichen Bestandteil.

15. Verfahren zum Bestimmen der Anwesenheit oder Menge von WT1-spezifischen Helfer-T-Zellen bei einem Patienten, der das MHC-Klasse-II-Molekül gemäß Anspruch 3 aufweist, wobei das Verfahren die Schritte umfasst:

(a) Umsetzen des Peptids gemäß einem der Ansprüche 1 bis 3 mit einer Probe, die von dem Patienten stammt; und dann

(b) Bestimmen der Anwesenheit oder Menge eines in der Probe enthaltenen Cytokins.

Revendications

1. Peptide qui consiste en une séquence d'acides aminés consistant en des acides aminés contigus dérivés d'une protéine WT1 et qui induit les cellules T auxiliaires spécifiques de WT1 en se liant à une molécule de classe II du CMH, dans lequel la séquence d'acides aminés est sélectionnée parmi :

- (a) la séquence d'acides aminés donnée dans l'ID SEQ N° 3 ;
- (b) la séquence d'acides aminés donnée dans l'ID SEQ N° 4 ;
- (c) la séquence d'acides aminés donnée dans l'ID SEQ N° 5 ; et
- (d) une séquence d'acides aminés dans laquelle un acide aminé uniquement est substitué, supprimé ou ajouté dans les séquences d'acides aminés données dans (a) à (c).

2. Peptide selon la revendication 1, dans lequel la séquence d'acides aminés est la séquence d'acides aminés donnée dans l'ID SEQ N° 3.

3. Peptide selon la revendication 1 ou 2, dans lequel

- (i) la molécule de classe II du CMH est sélectionnée parmi DRB1*0101, DRB1*0405, DRB1*0802, DRB1*0803, DRB1*0901, DRB1*1201, DRB1*1403, DRB1*1501, DRB1*1502, DPB1*0201, DPB1*0202, DPB1*0402, DPB1*0501, DPB1*0901, DQB1*0301, DQB1*0302, DQB1*0401, DQB1*0501, DQB1*0601, DQB1*0602 et DRB35*0102 ; ou
- (ii) la molécule de classe II du CMH est sélectionnée parmi DRB1*0101, DRB1*0405, DRB1*1502, DPB1*0201, DPB1*0202 et DQB1*0601.

4. Polynucléotide codant pour le peptide selon l'une quelconque des revendications 1 à 3.

5. Vecteur d'expression comprenant le polynucléotide selon la revendication 4.

6. Anticorps dirigé contre le peptide selon l'une quelconque des revendications 1 à 3.

7. Composition pharmaceutique à utiliser dans un procédé de traitement ou de prévention d'un cancer, comprenant le peptide selon l'une quelconque des revendications 1 à 3, le polynucléotide selon la revendication 4 ou le vecteur selon la revendication 5.

8. Peptide selon l'une quelconque des revendications 1 à 3, polynucléotide selon la revendication 4 ou vecteur selon la revendication 5, à utiliser dans un procédé de traitement ou de prévention d'un cancer.

9. Cellules présentatrices d'antigène qui affichent le peptide selon l'une quelconque des revendications 1 à 3 via la molécule de classe II du CMH selon la revendication 3.

10. Procédé d'induction de cellules présentatrices d'antigène, qui comprend la mise en culture de cellules immatures présentatrices d'antigène en présence du peptide selon l'une quelconque des revendications 1 à 3, et l'induction des cellules présentatrices d'antigène qui affichent le peptide via la molécule de classe II du CMH selon la revendication 3 à partir des cellules immatures présentatrices d'antigène.

11. Cellules T auxiliaires spécifiques de WT1 qui sont induites par le peptide selon l'une quelconque des revendications 1 à 3.

12. Procédé d'induction de cellules T auxiliaires spécifiques de WT1, qui comprend la mise en culture de cellules mononucléaires du sang périphérique en présence du peptide selon l'une quelconque des revendications 1 à 3, et l'induction des cellules T auxiliaires spécifiques de WT1 à partir des cellules mononucléaires du sang périphérique.


13. Kit d'induction de cellules T auxiliaires spécifiques de WT1, comprenant en tant qu'ingrédient essentiel le peptide selon l'une quelconque des revendications 1 à 3.

14. Kit de prévention ou de traitement d'un cancer, comprenant en tant qu'ingrédient essentiel le peptide selon l'une quelconque des revendications 1 à 3, le polynucléotide selon la revendication 4 ou le vecteur selon la revendication 5.


15. Procédé de détermination de la présence ou de la quantité de cellules T auxiliaires spécifiques de WT1 chez un sujet présentant une molécule de classe II du CMH selon la revendication 3, ledit procédé comprenant les étapes consistant à :

- (a) faire réagir le peptide selon l'une quelconque des revendications 1 à 3 avec un échantillon dérivé du sujet ; puis
- (b) déterminer la présence ou la quantité d'une cytokine contenue dans l'échantillon.


[Fig. 1]




[Fig. 2]




[Fig. 3]




[Fig. 4]




[Fig. 5]




[Fig. 6]




[Fig. 7]




[Fig. 8]




[Fig. 9]




[Fig. 10]




[Fig. 11]




[Fig. 12]




[Fig. 13]




[Fig. 14]




[Fig. 15]




[Fig. 16]



[Fig. 17]



[Fig. 18]



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2005045027 A [0005] [0007]
- JP 2009105286 A [0081]

Non-patent literature cited in the description

- **DANIEL A. HABER et al.** *Cell*, 29 June 1990, vol. 61 (7), 1257-69 [0006]
- **CALL KM et al.** *Cell*, 09 February 1990, vol. 60 (3), 509-20 [0006]
- **MENKE AL et al.** *Int Rev Cytol.*, 1998, vol. 181, 151-212 [0006]
- **YAMAGAMI T et al.** *Blood*, 01 April 1996, vol. 87 (7), 2878-84 [0006]
- **INOUE K et al.** *Blood*, 15 April 1998, vol. 91 (8), 2969-76 [0006]
- **TSUBOI A et al.** *Leuk Res.*, May 1999, vol. 23 (5), 499-505 [0006]
- **OKA Y et al.** *J Immunol.*, 15 February 2000, vol. 164 (4), 1873-80 [0006]
- **MELIEF CJ et al.** *Immunol Rev.*, June 1995, vol. 145, 167-77 [0006]
- **RITZ J.** *J Clin Oncol.*, February 1994, vol. 12 (2), 237-8 [0006]
- **TSUBOI A et al.** *J Clin Immunol.*, May 2000, vol. 20 (3), 195-202 [0006]
- **OKA Y et al.** *Immunogenetics*, February 2000, vol. 51 (2), 99-107 [0006]
- **OHMINAMI H et al.** *Blood*, 01 January 2000, vol. 95 (1), 286-93 [0006]
- **GAO L et al.** *Blood*, 01 April 2000, vol. 95 (7), 2198-203 [0006]
- *Cancer. Res.*, 2002, vol. 62, 6438 [0006]
- *J. Immunol. Immunother.*, 2001, vol. 24, 195 [0006]
- *Cancer. Immunol. Immunother.*, 2002, vol. 51, 271 [0006]
- *Cancer Immunol Immunother*, 2002, vol. 51, 271-281 [0007]
- Peptide Synthesis. Interscience, 1966 [0022]
- The Proteins. Academic Press Inc, 1976, vol. 2 [0022]
- Peptide Synthesis. Maruzen Co., Ltd, 1975 [0022]
- Basis and Experiments of Peptide Synthesis. Maruzen Co., Ltd, 1985 [0022]
- Development of Medicines (continuation). Peptide Synthesis. Hirokawa Shoten Co, 1991, vol. 14 [0022]
- **T. MANIATIS et al.** *Molecular Cloning*. CSH Laboratory, 1983 [0022]
- **DM. GLOVER.** *DNA Cloning*. IRL PRESS, 1985 [0022]
- *Cancer Immunol. Immunother.*, 2002, vol. 51, 271 [0023]
- *Molecular Cloning*. Cold Spring Harbor Laboratory press [0026]
- *Molecular Biology*. John Wiley & Sons, 1989, 6.3.1-6.3.6 [0026]
- *Current protocols in Molecular Biology*. John Wiley and Sons, 1987 [0029]
- *Antibodies; A Laboratory Manual*. Cold Spring Harber Laboratory Press, 1989 [0029]
- **OKA et al.** Cancer immunotherapy targeting Wilms' tumor gene WT1 product. *Journal of Immunology*, 2000, vol. 164, 1873-1880 [0031]
- **OKA et al.** Human cytotoxic T-lymphocyte responses specific for peptides of the wild-type Wilms' tumor gene (WT1) product. *Immunogenetics*, 2000, vol. 51, 99-107 [0031]
- **RAMMENSEE et al.** *Immunogenetics*, 1995, vol. 41, 178-228 [0062]

专利名称(译)	癌抗原辅助肽		
公开(公告)号	EP2423310B1	公开(公告)日	2014-12-17
申请号	EP2010767122	申请日	2010-04-22
[标]申请(专利权)人(译)	株式会社癌免疫研究所		
申请(专利权)人(译)	国际学院癌症免疫，Inc.的.		
当前申请(专利权)人(译)	国际学院癌症免疫，Inc.的.		
[标]发明人	SUGIYAMA HARUO		
发明人	SUGIYAMA, HARUO		
IPC分类号	C12N15/09 A61K31/7088 A61K38/00 A61K48/00 A61P35/00 C07K14/82 C12N5/07 C12Q1/04 C12Q1/06 G01N33/68 G01N33/53		
CPC分类号	A61K38/00 A61K48/00 C07K14/82 C07K16/32 C12N5/0635 C12N5/0636 C12N5/0639 C12N5/0645 G01N33/68 A61K39/00 A61K39/0011 A61K2039/55566 A61P1/04 A61P1/16 A61P11/00 A61P13/08 A61P13/10 A61P15/00 A61P17/00 A61P19/00 A61P35/00 A61P35/02 C07K14/4702 C07K14/4748 G01N33/57438 A61K38/17 A61K38/177 A61K38/1774 A61K39/001153 C07K16/18 C07K16/30 C12N5/0634 G01N33/56966		
优先权	2009105286 2009-04-23 JP		
其他公开文献	EP2423310A4 EP2423310A1		
外部链接	Espacenet		

摘要(译)	Met Gly Ser Asp Val Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Ser 1 5 10 15 Ser Leu Gly Gly Gly Gly Gly Cys Gly Leu Pro Val Ser Gly Ala 20 25 30 Arg Gln Trp Ala Pro Val Leu Asp Phe Ala Pro Pro Gly Ala Ser Ala 35 40 45 Tyr Gly Ser Leu Gly Gly Pro Ala Pro Pro Pro Ala Pro Pro Pro Pro 50 55 60 Pro Pro Pro Pro His Ser Phe Ile Lys Gln Glu Pro Ser Trp Gly Gly 65 70 75 80 Ala Glu Pro His Glu Glu Gln Cys Leu Ser Ala Phe Thr Leu His Phe 85 90 95 Ser Gly Gln Phe Thr Gly Thr Ala Gly Ala Cys Arg Tyr Gly Pro Phe 100 105 110 Gly Pro Pro Pro Ser Gln Ala Ser Ser Gly Gln Ala Arg Met Phe 115 120 125 Pro Asn Ala Pro Tyr Leu Pro Ser Cys Leu Glu Ser Gln Pro Thr Ile 130 135 140 Arg Asn Gln Gly Tyr Ser Thr Val Thr Phe Asp Gly Ala Pro Ser Tyr 145 150 155 160
-------	--

本发明涉及一种WT1肽，其具有由衍生自WT1蛋白的连续氨基酸组成的氨基酸序列，并通过与MHC II类分子结合诱导WT1特异性辅助T细胞，包含它们的药物组合物等。