



(11) **EP 2 663 639 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
13.09.2017 Bulletin 2017/37

(51) Int Cl.:
C12N 15/10 ^(2006.01) **C12Q 1/68** ^(2006.01)
C12Q 1/48 ^(2006.01) **G01N 33/53** ^(2006.01)
C07H 21/02 ^(2006.01)

(21) Application number: **12757380.6**

(86) International application number:
PCT/US2012/021264

(22) Date of filing: **13.01.2012**

(87) International publication number:
WO 2012/125220 (20.09.2012 Gazette 2012/38)

(54) **METHODS FOR ISOLATION, IDENTIFICATION, AND QUANTIFICATION OF MIRNAS**

VERFAHREN ZUR ISOLIERUNG, IDENTIFIKATION UND QUANTIFIZIERUNG VON MIRNAS

PROCÉDÉS D'ISOLEMENT, D'IDENTIFICATION ET DE QUANTIFICATION DES MIARN

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

US-A1- 2006 099 619 US-A1- 2008 076 674
US-A1- 2008 274 904 US-A1- 2010 015 604
US-A1- 2010 209 933

(30) Priority: **14.01.2011 US 201161432874 P**

(43) Date of publication of application:
20.11.2013 Bulletin 2013/47

(73) Proprietor: **Life Technologies Corporation**
Carlsbad, CA 92008 (US)

(72) Inventors:
• **XU, Tom**
Carlsbad, California 92008 (US)
• **TRINH, Christopher**
Carlsbad, California 92008 (US)
• **BORNARTH, Carole**
Carlsbad, California 92008 (US)
• **EVANS, Brian**
Carlsbad, California 92008 (US)
• **RATH, Mousumi**
Carlsbad, California 92008 (US)
• **TRAN, Kathy**
Carlsbad, California 92008 (US)

- **TONG A W ET AL: "MicroRNA profile analysis of human prostate cancers", CANCER GENE THERAPY, APPLETON & LANGE, GB, vol. 16, no. 3, 1 March 2009 (2009-03-01), pages 206-216, XP002573838, ISSN: 0929-1903, DOI: 10.1038/CGT.2008.77 [retrieved on 2008-10-24]**
- **CASTOLDI M ET AL: "Regulation of Gene Expression By Small RNAs; Chapter 12, How to Assay microRNA Expression A Technology Guide", 1 January 2009 (2009-01-01), REGULATION OF GENE EXPRESSION BY SMALL RNAS, BOCA RATON, US, PAGE(S) 215 - 240, XP008121457, ISBN: 978-0-8493-9169-9 * page 15, paragraph 12.7.1.4; figure 12.1e ***
- **JUN LU ET AL: "MicroRNA expression profiles classify human cancers", NATURE, vol. 435, no. 7043, 9 June 2005 (2005-06-09), pages 834-838, XP055027393, ISSN: 0028-0836, DOI: 10.1038/nature03702**
- **J. CHEN ET AL: "Highly sensitive and specific microRNA expression profiling using BeadArray technology", NUCLEIC ACIDS RESEARCH, vol. 36, no. 14, 1 January 2008 (2008-01-01), pages e87-e87, XP055004505, ISSN: 0305-1048, DOI: 10.1093/nar/gkn387**

(74) Representative: **Weber, Birgit**
Life Technologies GmbH
Frankfurter Strasse 129 B
64293 Darmstadt (DE)

(56) References cited:
WO-A1-2005/012523 US-A1- 2006 057 595

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 663 639 B1

- SHI R ET AL: "FACILE MEANS FOR QUANTIFYING MICRORNA EXPRESSION BY REAL-TIME PCR", BIOTECHNIQUES, INFORMA HEALTHCARE, US, vol. 39, no. 4, 1 October 2005 (2005-10-01), pages 519-525, XP001246434, ISSN: 0736-6205, DOI: 10.2144/000112010
- PO-HSUN CHEN: 'Identification of Dysregulated miRNA Targets in Human Prostate Cancer' PAPER FOR DOCTOR OF PHILOSOPHY DEGREE December 2009, DEPARTMENT OF BIOMEDICAL STUDIES, BAYLOR UNIVERSITY, pages 13 - 35, XP055121230
- XIAO-HUI HUANG ET AL.: 'Bead-based microarray analysis of microRNA expression in hepatocellular carcinoma: miR-338 is downregulated' HEPATOLOGY RESEARCH vol. 39, 2009, pages 786 - 794, XP055121231
- ALBERTO BISCONTIN ET AL.: 'New miRNA labeling method for bead-based quantification' BMC MOLECULAR BIOLOGY vol. 11, no. 44, 16 June 2010, XP055121235
- KATHY TRAN ET AL.: 'microRNA Expression Profiling Potential for Human Breast Cancer Diagnosis Using a Bead-based microRNA-specific Capturing Method' 2012 AMERICAN ASSOCIATION FOR CANCER RESEARCH MEETING 30 March 2012, XP055126903
- ALBRETSSEN C ET AL: "Applications of magnetic beads with covalently attached oligonucleotides in hybridization: Isolation and detection of specific measles virus mRNA from a crude cell lysate", ANALYTICAL BIOCHEMISTRY, ACADEMIC PRESS INC, NEW YORK, vol. 189, no. 1, 15 August 1990 (1990-08-15), pages 40-50, XP024826933, ISSN: 0003-2697, DOI: 10.1016/0003-2697(90)90041-7 [retrieved on 1990-08-15]
- WANG W X ET AL: "Focus on RNA isolation: Obtaining RNA for microRNA (miRNA) expression profiling analyses of neural tissue", BIOCHIMICA ET BIOPHYSICA ACTA. GENE REGULATORY MECHANISMS, ELSEVIER, AMSTERDAM, NL, vol. 1779, no. 11, 1 November 2008 (2008-11-01), pages 749-757, XP025678678, ISSN: 1874-9399, DOI: 10.1016/J.BBAGRM.2008.01.005 [retrieved on 2008-02-13]
- VICTORIA EL-KHOURY ET AL: "Assessing cellular and circulating miRNA recovery: the impact of the RNA isolation method and the quantity of input material", SCIENTIFIC REPORTS, vol. 6, 20 January 2016 (2016-01-20), page 19529, XP055353647, DOI: 10.1038/srep19529
- Agnieszka Podolska ET AL: "How the RNA isolation method can affect microRNA microarray results", Acta Biochimica Polonica, 6 December 2011 (2011-12-06), pages 1-6, XP055353649, Retrieved from the Internet: URL:<http://gene-quantification.eu/podolska-et-al-micro-rna-quality-2011.pdf> [retrieved on 2017-03-10]

Description

FIELD OF THE INVENTION

5 **[0001]** This application relates to a method for isolating an miRNA from a sample of interest, comprising: a) contacting a biological sample with a lysis buffer to form a sample lysate b) contacting the sample lysate with an anti-miRNA probe covalently attached to a bead; c) incubating the sample lysate - bead mixture under suitable conditions to form hybridized complexes between the miRNA in the lysate and the anti-miRNA probes on the beads; d) washing the beads under suitable conditions to remove the unbound sample lysate material; and e) eluting the miRNA bound to the anti-miRNA probes to form an isolated miRNA sample. In a further aspect, the invention relates to a method of diagnosing a disease or disease progression, comprising: a) identifying a set of miRNA markers that are differentially regulated during a causation or progression of a disease; b) isolating and measuring the levels of such miRNA markers from a suitable biological sample from a patient in need thereof by a method of any one of claims 1 to 12, at different time points if necessary; and c) diagnosing the disease or its progression over time in the patient.

BACKGROUND

20 **[0002]** MicroRNAs regulate virtually every aspect of biology, including developmental timing, differentiation, proliferation, antiviral defense and metabolism. MicroRNAs are ~22-nucleotide-long RNAs that are generated by sequential processing from longer transcripts that contain a stem-loop. One strand is loaded into the miRNA-induced silencing complex (miRISC), which contains the proteins argonaute (Ago) and Trnc6 (trinucleotide repeat-containing 6; GW182). The other strand is usually degraded. The mature miRNA guides the miRISC to partially complementary sequences, termed miRNA recognition elements (MREs), in target mRNAs to repress mRNA translation, promote transcript decay or both. MicroRNAs probably regulate the expression of most coding genes. miRNAs have been reported from all living organisms, including humans, bacteria, viruses, plants, worms, and others. About 3% of human genes encode for miRNAs and about 30% of genes are believed to be regulated by miRNAs. It has also been widely reported that an abnormal level of miRNA expression or presence is associated with diseases, such as cancer, heart attack, diabetes, etc.

25 **[0003]** Quantitative and qualitative isolation of miRNAs from various biological samples has been hampered for several reasons, including, but not limited to, labor-intensive and time consuming protocols; the nature of small size of miRNA leads to easy loss of the targets during extraction; miRNAs, like other RNAs, are not stable and can be degraded easily during processing and storage; the miRNA detection rate from real patient samples is usually low or undetectable, *i.e.*, not quantitative; and further the extracted miRNA targets normally are poorly correlated between related study objects (e.g. placenta vs. blood).

30 **[0004]** For example, most miRNA (or RNA), isolation reagents or kits that are commercially available (from QIAGEN - PAXGENE blood RNA kit, AMBION - LEUKOLOCK total RNA kit, and others), suffer from one or more of the following drawbacks: low sensitivity and detectability; low yield; not reproducible; large sample volume requirement; miRNA extraction does not scale well; time-consuming workflows, 4-5 hours workflow in general; low throughput and not automatable. A method for the analysis of miRNAs using probes coupled to beads is disclosed in Tong et al. (Cancer Gene Therapy (2009) 16, 206-216). Albretsen et al. (Analytical Biochemistry (1990) 189, 40-50) discloses the isolation of a virus mRNA from a crude cell lysate. Thus, there is a need in the field for better methods for isolating, identifying, and quantifying miRNAs.

BRIEF DESCRIPTION OF THE DRAWINGS

45 **[0005]**

Figure 1: Schematic for making anti-miRNA probe beads using chemical synthesis.

Figure 2: Schematic for enzymatic synthesis of anti-miRNA probe beads.

Figure 3A: Non-anti-probe beads hybridization with Cy3-labeled Let-7a.

50 Figure 3B: Anti-Let-7a-probe beads hybridization with Cy3-labeled Let-7a.

Figure 3C: FACS after de-hybridization of Cy3-labeled Let-7a bond from anti-Let-7a-probe beads.

Figure 4: Let-7 family miRNA profiling by C_t values.

Figure 5: miRNA detection after addition (spiking) to plasma.

Figure 6A: Quantitative real-time PCR C_t profiles of 96 miRNAs from blood.

55 Figure 6B: Quantitative real-time PCR C_t profiles of 95 miRNAs from blood.

Figure 7: Saliva miRNA profiling using anti-miRNA probe beads.

Figure 8: A384C6 Beads showing Sensitivity and Dynamic Range of anti-miRNA probe beads.

Figure 9: miRNA Detectability and reproducibility.

Figure 10A: Let7a - miRNA profiling from human bladder LCM samples; LCM area and Ct correlation.
 Figure 10B: Let7c - miRNA profiling from human bladder LCM samples; LCM area and Ct correlation.
 Figure 10C: miR-16 - miRNA profiling from human bladder LCM samples; LCM area and Ct correlation.
 Figure 11: Breast cancer FFPE sample miRNA profiling.
 Figure 12: Difference in expression of miRNA levels between breast cancer lines: MBA-MB-21 versus MCF-7 (delta Ct is shown).
 Figure 13: miRNA profiling from raw cow milk.
 Figure 14: miRNA profiling from urine from three donors.
 Figure 15: Megaplex A miRNA profiling from 25 microL serum.
 Figure 16: Megaplex B miRNA profiling from 25 microL serum.
 Figure 17: miRNA expression before and at 15-week of pregnancy.

SUMMARY OF THE INVENTION

[0006] As described above, the isolation, detection and quantification of miRNAs have been a daunting problem. Applicants have now developed a simple and elegant solution to the problem. In one embodiment, the invention relates to a method for isolating an miRNA from a sample of interest, comprising: a) contacting a biological sample with a lysis buffer to form a sample lysate b) contacting the sample lysate with an anti-miRNA probe covalently attached to a bead; c) incubating the sample lysate - bead mixture under suitable conditions to form hybridized complexes between the miRNA in the lysate and the anti-miRNA probes on the beads; d) washing the beads under suitable conditions to remove the unbound sample lysate material; and e) eluting the miRNA bound to the anti-miRNA probes to form an isolated miRNA sample.

[0007] In a further embodiment of the invention, the method further comprises identifying or quantifying the miRNA in the isolated miRNA sample.

[0008] In another embodiment of the invention, the identifying or quantifying the miRNA is comprises using reverse transcription followed by conventional or real-time or quantitative polymerase chain reaction (q-PCR) or other methods of amplification. In another embodiment, the identifying or quantifying a miRNA involves non-qPCR techniques.

[0009] In one embodiment of the invention, the beads are magnetic beads. In another embodiment of the invention, the beads are non-magnetic beads. In one embodiment of the invention, the beads are carboxylic acid functionalized beads. In another embodiment of the invention, the beads are DNA beads. In another embodiment of the invention, the beads are P1 DNA beads.

[0010] In one embodiment of the invention, the anti-miRNA probe is attached to the beads using chemical synthesis.

[0011] In another embodiment of the invention, the anti-miRNA probe is attached to the beads using enzymatic synthesis. In another embodiment of the invention, the enzymatic synthesis comprises: taking an oligonucleotide that has a sequence complementary to an miRNA of interest (anti-miRNA probe); adding a terminal didexoy nucleotide to the 3' end using a terminal transferase; adding a 5'-phosphate group to the oligonucleotide using a kinase; and ligating the oligonucleotide to DNA beads using a single-strand DNA ligase. In one embodiment, the kinase is a T4 polynucleotide kinase.

[0012] In one embodiment of the invention, a single species of anti-miRNA probe is attached to an individual bead. In another embodiment of the invention more than one species of anti-miRNA probes are attached to an individual bead.

[0013] In one embodiment of the invention, sample of interest is a biological sample. In another embodiment of the invention, the biological sample is selected from the group consisting of blood, serum, plasma, urine, saliva, cerebrospinal fluid, wound exudates, biopsies, autopsies, tissues, formalin-fixed, paraffin-embedded (FFPE) samples, or organs. In another embodiment of the invention, the biological sample is a biological fluid. In another embodiment of the invention, the biological fluid is blood. In another embodiment of the invention, the biological fluid is plasma. In another embodiment of the invention, the biological fluid is serum. In another embodiment of the invention, the biological fluid is saliva. In another embodiment, the sample could be a fossil or a fossilized rock or sediment.

[0014] In another embodiment, a method of diagnosing a disease or disease progression is disclosed, comprising: a) identifying a set of miRNA markers that are differentially regulated during a causation or progression of a disease; b) isolating and measuring the levels of such miRNA markers from a suitable biological sample from a patient in need thereof by a method of any one of claims 1 to 12, at different time points if necessary; and c) diagnosing the disease or its progression over time in the patient.

DETAILED DESCRIPTION OF THE INVENTION

[0015] It is to be understood that the particular methods, reaction mixtures, and / or systems described herein may vary. The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting. Further, unless defined otherwise, all technical and scientific terms used herein have the same meaning as

commonly understood by one of ordinary skill in the art to which this invention pertains. As used in this specification and the appended claims, the singular forms "a", "an", and "the" also include plural referents unless the context clearly provides otherwise. All numerical ranges are intended to encompass each individual value within the range as if each were separately listed (e.g., 10-20 may include one or more of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 and / or 20). In terms of concentration ranges, these encompass fractional ranges, e.g., all values between 10-20, represents as if each were individually written out (e.g., 10.8, 11.5). The term "approximately", when used to modify a group of numerical values, is meant to apply to each value individually unless otherwise indicated.

[0016] An "amplicon" typically refers to a molecule made by copying or transcribing another molecule, e.g., as occurs in transcription, cloning, and/or in a polymerase chain reaction ("PCR") (e.g., strand displacement PCR amplification (SDA), duplex PCR amplification, etc.) or other nucleic acid amplification technique. Typically, an amplicon is a copy of a selected nucleic acid or a portion thereof (e.g., a template or target nucleic acid) or is complementary thereto. The term "amplifying" or "amplification" in the context of nucleic acids typically refers to the production of multiple copies of a polynucleotide, or a portion of the polynucleotide, typically starting from a small amount of the polynucleotide (e.g., a single polynucleotide molecule), where the amplification products or amplicons are generally detectable. Any of several methods may be used to amplify the target nucleic acid from the sample. The term "amplifying" which typically refers to an "exponential" increase in target nucleic acid is being used herein to describe both linear and exponential increases in the numbers of a select target sequence of nucleic acid. The term "amplification reaction mixture" refers to an aqueous solution comprising the various reagents used to amplify a target nucleic acid. These include enzymes, aqueous buffers, salts, amplification primers, target nucleic acid, and nucleoside triphosphates. Depending upon the context, the mixture can be either a complete or incomplete amplification reaction mixture. The method used to amplify the target nucleic acid may be any available to one of skill in the art. Any *in vitro* means for multiplying the copies of a target sequence of nucleic acid may be utilized. These include linear, logarithmic, and / or any other amplification method. Exemplary methods include polymerase chain reaction (PCR; see, e.g., U.S. Patent Nos. 4,683,202; 4,683,195; 4,965,188; and / or 5,035,996), isothermal procedures (using one or more RNA polymerases (see, e.g., WO 2006/081222), strand displacement (see, e.g., U.S. Pat. No. RE39007E), partial destruction of primer molecules (see, e.g., WO2006087574)), ligase chain reaction (LCR) (see, e.g., Wu, et al., Genomics 4: 560-569 (1990) and / or Barany, et al. PNAS USA 88:189-193 (1991)), Q β RNA replicase systems (see, e.g., WO/1994/016108), RNA transcription-based systems (e.g., TAS, 3SR), rolling circle amplification (RCA) (see, e.g., U.S. Pat. No. 5,854,033; U.S. Pub. No. 2004/265897; Lizardi et al. Nat. Genet. 19: 225-232 (1998); and/or Banér et al. Nucleic Acid Res., 26: 5073-5078 (1998)), and strand displacement amplification (SDA) (Little, et al. Clin Chem 45:777-784 (1999)), among others. Many systems are suitable for use in amplifying target nucleic acids and are contemplated herein as would be understood by one of skill in the art.

[0017] Any of several methods may be used to detect target nucleic acids using various primers and / or probes. Many different reagents, systems, and / or detectable labels may be used in the methods described herein. These include, for example, TaqMan[®] systems, detectable label-quencher systems (e.g., FRET, salicylate /DTPA ligand systems (see, e.g., Oser et al. Angew. Chem. Int. Engl. 29(10):1167 (1990), displacement hybridization, homologous probes, assays described in EP 070685), molecular beacons (e.g., NASBA, Scorpion, locked nucleic acid (LNA) bases (Singh, et al. Chem Commun 4:455-456 (1998)), peptide nucleic acid (PNA) probes (Pellestor, et al. European J. Human Gen. 12:694-700 (2004)), Eclipse probes (Afonina, et al. Biotechniques 32:940-949 (2002)), light-up probes (Svanvik, et al. Anal Biochem 281:26-35 (2001)), molecular beacons (Tyagi, et al. Nat. Biotechnol. 14:303-308 (1996)), tripartite molecular beacons (Nutiu, et al. Nucleic Acids Res. 30:e94 (2002)), QuantiProbes (www.qiagen.com), HyBeacons (French, et al. Mol. Cell. Probes 15:363-374 (2001)), displacement probes (Li, et al. Nucleic Acids Res. 30:e5 (2002)), HybProbes (Cardullo, et al. PNAS 85:8790-8794 (1988)), MGB Alert (www.nanogen.com), Q-PNA (Fiandaca, et al. Genome Res. 11:609-611 (2001)), Plexor (www.Promega.com), LUX primers (Nazarenko, et al. Nucleic Acids Res. 30:e37 (2002)), Scorpion primers (Whitcombe, et al. Nat Biotechnol 17:804-807 (1999)), AmpliFluor (Sunrise) primers (Nazarenko, et al. Nucleic Acids Res. 25:2516-2521 (1997)), DzyNA primers (Todd, et al. Clin. Chem. 46:625-630 (2000)), and the like. In each of these assays, the generation of amplification products may be monitored while the reaction is in progress. An apparatus for detecting the signal generated by the detectable label may be used to detect, measure, and quantify the signal before, during, and / or after amplification. The particular type of signal may dictate the choice of detection method. For example, in some embodiments, fluorescent dyes are used to label probes and / or amplified products. The probes bind to single-stranded and / or double-stranded amplified products, and / or the dyes intercalate into the double-stranded amplified products, and consequently, the resulting fluorescence increases as the amount of amplified product increases. In some embodiments, the T_m is ascertained by observing a fluorescence decrease as the double-stranded amplified product dissociates and the intercalating dye is released therefrom. The amount of fluorescence may be quantitated using standard equipment such as a spectra-fluorometer, for example. The use of other methods and/or reagents is also contemplated herein.

[0018] One exemplary method for amplifying and detecting target nucleic acids is commercially available as TaqMan[®] (see, e.g., U.S. Pat. Nos. 4,889,818; 5,079,352; 5,210,015; 5,436,134; 5,487,972; 5,658,751; 5,210,015; 5,487,972; 5,538,848; 5,618,711; 5,677,152; 5,723,591; 5,773,258; 5,789,224; 5,801,155; 5,804,375; 5,876,930; 5,994,056;

6,030,787; 6,084,102; 6,127,155; 6,171,785; 6,214,979; 6,258,569; 6,814,934; 6,821,727; 7,141,377; and / or 7,445,900). TaqMan[®] assays are typically carried out by performing nucleic acid amplification on a target polynucleotide using a nucleic acid polymerase having 5'-3' nuclease activity, a primer capable of hybridizing to said target polynucleotide, and an oligonucleotide probe capable of hybridizing to said target polynucleotide 3' relative to said primer. The oligonucleotide probe typically includes a detectable label (e.g., a fluorescent reporter molecule) and a quencher molecule capable of quenching the fluorescence of said reporter molecule. Typically, the detectable label and quencher molecule are part of a single probe. As amplification proceeds, the polymerase digests the probe to separate the detectable label from the quencher molecule. The detectable label (e.g., fluorescence) is monitored during the reaction, where detection of the label corresponds to the occurrence of nucleic acid amplification (e.g., the higher the signal the greater the amount of amplification). Variations of TaqMan[®] assays (e.g., LNA[™] spiked TaqMan[®] assay) are known in the art and would be suitable for use in the methods described herein.

[0019] Another exemplary system utilizes double-stranded probes in displacement hybridization methods (see, e.g., Morrison et al. *Anal. Biochem.*, 18:231-244 (1989); and / or Li, et al. *Nucleic Acids Res.*, 30(2,e5) (2002)). In such methods, the probe typically includes two complementary oligonucleotides of different lengths where one includes a detectable label and the other includes a quencher molecule. When not bound to a target nucleic acid, the quencher suppresses the signal from the detectable label. The probe becomes detectable upon displacement hybridization with a target nucleic acid. Multiple probes may be used, each containing different detectable labels, such that multiple target nucleic acids may be queried in a single reaction.

[0020] Additional exemplary methods for amplifying and detecting target nucleic acids involve "molecular beacons", which are single-stranded hairpin shaped oligonucleotide probes. In the presence of the target sequence, the probe unfolds, binds and emits a signal (e.g., fluoresces). A molecular beacon typically includes at least four components: 1) the "loop", an 18-30 nucleotide region which is complementary to the target sequence; 2) two 5-7 nucleotide "stems" found on either end of the loop and being complementary to one another; 3) at the 5' end, a detectable label; and 4) at the 3' end, a quencher dye that prevents the detectable label from emitting a signal when the probe is in the closed loop shape (e.g., not bound to a target nucleic acid). Thus, in the presence of a complementary target, the "stem" portion of the beacon separates out resulting in the probe hybridizing to the target. Other types of molecular beacons are also known and may be suitable for use in the methods described herein. Molecular beacons may be used in a variety of assay systems. One such system is nucleic acid sequence-based amplification (NASBA[®]), a single step isothermal process for amplifying RNA to double stranded DNA without temperature cycling. A NASBA reaction typically requires avian myeloblastosis virus reverse transcriptase (AMV-RT or AMV), T7 RNA polymerase, RNase H, and two oligonucleotide primers. After amplification, the amplified target nucleic acid may be detected using a molecular beacon. Other uses for molecular beacons are known in the art and would be suitable for use in the methods described herein.

[0021] The Scorpion system is another exemplary assay format that may be used in the methods described herein. Scorpion primers are bi-functional molecules in which a primer is covalently linked to the probe, along with a detectable label (e.g., a fluorophore) and a quencher. In the presence of a target nucleic acid, the detectable label and the quencher separate which leads to an increase in signal emitted from the detectable label. Typically, a primer used in the amplification reaction includes a probe element at the 5' end along with a "PCR blocker" element (e.g., HEG monomer) at the start of the hairpin loop. The probe typically includes a self-complementary stem sequence with a detectable label at one end and a quencher at the other. In the initial amplification cycles (e.g., PCR), the primer hybridizes to the target and extension occurs due to the action of polymerase. The Scorpion system may be used to examine and identify point mutations using multiple probes that may be differently tagged to distinguish between the probes. Using PCR as an example, after one extension cycle is complete, the newly synthesized target region will be attached to the same strand as the probe. Following the second cycle of denaturation and annealing, the probe and the target hybridize. The hairpin sequence then hybridizes to a part of the newly produced PCR product. This results in the separation of the detectable label from the quencher and causes emission of the signal. Other uses for molecular beacons are known in the art and would be suitable for use in the methods described herein.

[0022] One or more detectable labels and / or quenching agents are typically attached to an oligonucleotide primer and / or probe. The detectable label may emit a signal when free or when bound to the target nucleic acid. The detectable label may also emit a signal when in proximity to another detectable label. Detectable labels may also be used with quencher molecules such that the signal is only detectable when not in sufficiently close proximity to the quencher molecule. For instance, in some embodiments, the assay system may cause the detectable label to be liberated from the quenching molecule. Any of several detectable labels may be used to label the primers and probes used in the methods described herein. As mentioned above, in some embodiments the detectable label may be attached to a probe, which may be incorporated into a primer, or may otherwise bind to amplified target nucleic acid (e.g., a detectable nucleic acid binding agent such as an intercalating or non-intercalating dye). When using more than one detectable label, each should differ in their spectral properties such that the labels may be distinguished from each other, or such that together the detectable labels emit a signal that is not emitted by either detectable label alone. Exemplary detectable labels include, for instance, a fluorescent dye or fluorophore (e.g., a chemical group that can be excited by light to emit

fluorescence or phosphorescence), "acceptor dyes" capable of quenching a fluorescent signal from a fluorescent donor dye, and the like. Suitable detectable labels may include, for example, fluorosceins (e.g., 5-carboxy-2,7-dichlorofluorescein; 5-Carboxyfluorescein (5-FAM); 5-HAT (Hydroxy Tryptamine); 5-Hydroxy Tryptamine (HAT); 6-JOE; 6-carboxyfluorescein (6-FAM); FITC); Alexa fluors (e.g., 350, 405, 430, 488, 500, 514, 532, 546, 555, 568, 594, 610, 633, 635, 647, 660, 680, 700, 750); BODIPY fluorophores (e.g., 492/515, 493/503, 500/510, 505/515, 530/550, 542/563, 558/568, 564/570, 576/589, 581/591, 630/650-X, 650/665-X, 665/676, FL, FL ATP, FI-Ceramide, R6G SE, TMR, TMR-X conjugate, TMR-X, SE, TR, TR ATP, TR-X SE), coumarins (e.g., 7-amino-4-methylcoumarin, AMC, AMCA, AMCA-S, AMCA-X, ABQ, CPM methylcoumarin, coumarin phalloidin, hydroxycoumarin, CMFDA, methoxycoumarin), calcein, calcein AM, calcein blue, calcium dyes (e.g., calcium crimson, calcium green, calcium orange, calcofluor white), Cascade Blue, Cascade Yellow; Cy[™] dyes (e.g., 3, 3.18, 3.5, 5, 5.18, 5.5, 7), cyan GFP, cyclic AMP Fluorosensor (FiCRhR), fluorescent proteins (e.g., green fluorescent protein (e.g., GFP, EGFP), blue fluorescent protein (e.g., BFP, EBFP, EBFP2, Azurite, mKAlama1), cyan fluorescent protein (e.g., ECFP, Cerulean, CyPet), yellow fluorescent protein (e.g., YFP, Citrine, Venus, YPet), FRET donor/acceptor pairs (e.g., fluorescein/tetramethylrhodamine, IAEDANS/fluorescein, EDANS/dabcyl, fluorescein/fluorescein, BODIPY FL/BODIPY FL, Fluorescein/QSY7 and QSY9), LysoTracker and LysoSensor (e.g., LysoTracker Blue DND-22, LysoTracker Blue-White DPX, LysoTracker Yellow HCK-123, LysoTracker Green DND-26, LysoTracker Red DND-99, LysoSensor Blue DND-167, LysoSensor Green DND-189, LysoSensor Green DND-153, LysoSensor Yellow/Blue DND-160, LysoSensor Yellow/Blue 10,000 MW dextran), Oregon Green (e.g., 488, 488-X, 500, 514); rhodamines (e.g., 110, 123, B, B 200, BB, BG, B extra, 5-carboxytetramethylrhodamine (5-TAMRA), 5 GLD, 6-Carboxyrhodamine 6G, Lissamine, Lissamine Rhodamine B, Phalloidine, Phalloidine, Red, Rhod-2, 5-ROX (carboxy-X-rhodamine), Sulphorhodamine B can C, Sulphorhodamine G Extra, Tetramethylrhodamine (TRITC), WT), Texas Red, Texas Red-X, VIC and other labels described in, e.g., US Pub. No. 2009/0197254), among others as would be known to those of skill in the art. Other detectable labels may also be used (see, e.g., US Pub. No. 2009/0197254), as would be known to those of skill in the art.

[0023] Nucleic acid binding agents may also be used to detect nucleic acids amplified using the methods described herein. Many suitable detectable nucleic acid binding agents are available to one of skill in the art and may be used alone or in combination with other agents and / or components of an assay system. Exemplary DNA binding agents may include, for example, acridines (e.g., acridine orange, acriflavine), actinomycin D (Jain, et al. J. Mol. Biol. 68:21 (1972)), anthramycin, BOBO[™]-1, BOBO[™]-3, BO-PRO[™]-1, chromomycin, DAPI (Kapuseinski, et al. Nuc. Acids Res. 6(112): 3519 (1979)), daunomycin, distamycin (e.g., distamycin D), dyes described in U.S. Pat. No. 7,387,887, ellipticine, ethidium salts (e.g., ethidium bromide), fluorcoumanin, fluorescent intercalators as described in U.S. Pat. No. 4,257,774, GelStar[®] (Cambrex Bio Science Rockland Inc., Rockland, Me.), Hoechst 33258 (Searle and Embrey, 1990, Nuc. Acids Res. 18:3753-3762), Hoechst 33342, homidium, JO-PRO[™]-1, LIZ dyes, LO-PRO[™]-1, mepacrine, mithramycin, NED dyes, netropsin, 4'6-diamidino- α -phenylindole, proflavine, POPO[™]-1, POPO[™]-3, PO-PRO[™]-1, propidium iodide, ruthenium polypyridyls, S5, SYBR[®] Gold, SYBR[®] Green I (U.S. Patent No. 5,436,134 and 5,658,751), SYBR[®] Green II, SYTOX blue, SYTOX green, SYTO[®] 43, SYTO[®] 44, SYTO[®] 45, SYTOX[®] Blue, TO-PRO[®]-1, SYTO[®] 11, SYTO[®] 13, SYTO[®] 15, SYTO[®] 16, SYTO[®] 20, SYTO[®] 23, thiazole orange (Aldrich Chemical Co., Milwaukee, Wis.), TOTO[™]-3, YO-PRO[®]-1, and YOYO[®]-3 (Molecular Probes, Inc., Eugene, OR), among others. SYBR[®] Green I (see, e.g., U.S. Pat. Nos. 5,436,134; 5,658,751; and / or 6,569,927), for example, has been used to monitor a PCR reaction by amplifying the target sequence in the presence of the dye, exciting the biological sample with light at a wavelength absorbed by the dye and detecting the emission therefrom; and, determining a melting profile of the amplified target sequence. The presence of amplified products and, therefore, the target sequence in the sample, may thereafter be determined by, for example, performing a melting curve analysis (e.g., non-linear least squares regression of the sum of multiple gaussians). It is to be understood that the use of the SYBR[®] Green dye is presented as an example and that many such dyes may be used in the methods described herein. Other nucleic acid binding agents may also be suitable as would be understood by one of skill in the art.

[0024] Nucleic acids "hybridize" or "anneal" in a base-pairing interaction of one polynucleotide with another polynucleotide (typically an antiparallel polynucleotide) that results in formation of a duplex or other higher-ordered structure, typically termed a hybridization complex. The primary interaction between the antiparallel polynucleotides is typically base specific, e.g., A/T and G/C, by Watson/Crick and/or Hoogsteen-type interactions. It is not a requirement that two polynucleotides have 100% complementarity over their full length to achieve hybridization. In some aspects, a hybridization complex can form from intermolecular interactions, or alternatively, can form from intramolecular interactions. Hybridization occurs due to a variety of well-characterized forces, including hydrogen bonding, solvent exclusion, and base stacking. An extensive guide to nucleic hybridization may be found in, Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes, part I, chapter 2, "Overview of principles of hybridization and the strategy of nucleic acid probe assays," Elsevier (1993).

[0025] A "mixture" refers to a combination of two or more different components. A "reaction mixture" refers a mixture that comprises molecules that can participate in and/or facilitate a given reaction or assay. To illustrate, an amplification reaction mixture generally includes a solution containing reagents necessary to carry out an amplification reaction, and typically contains primers, a biocatalyst (e.g., a nucleic acid polymerase, a ligase, etc.), dNTPs, and a divalent metal

cation in a suitable buffer. A reaction mixture is referred to as complete if it contains all reagents necessary to carry out the reaction, and incomplete if it contains only a subset of the necessary reagents. It will be understood by one of skill in the art that reaction components are routinely stored as separate solutions, each containing a subset of the total components, for reasons of convenience, storage stability, or to allow for application-dependent adjustment of the component concentrations, and that reaction components are combined prior to the reaction to create a complete reaction mixture. Furthermore, it will be understood by one of skill in the art that reaction components are packaged separately for commercialization and that useful commercial kits may contain any subset of the reaction or assay components, which includes the biomolecules of the invention.

[0026] A "moiety" or "group" refers to one of the portions into which something, such as a molecule, is or can be divided (e.g., a functional group, substituent group, or the like). For example, an oligonucleotide described herein includes at least one donor moiety and/or at least one acceptor moiety in certain embodiments.

[0027] The term "mutation" refers to a nucleic acid that has been altered in its nucleic acid sequence or an encoded protein product of a nucleic acid that has been altered in its amino acid sequence relative to an unaltered or native form of the nucleic acid or encoded protein product. Such alterations include, for example, point mutations or substitutions, deletions and insertions.

[0028] The term "nucleic acid" or "polynucleotide" refers to a polymer that can be corresponded to a ribose nucleic acid (RNA) or deoxyribose nucleic acid (DNA) polymer, or an analog thereof. This includes polymers of nucleotides such as RNA and DNA, as well as modified forms thereof, peptide nucleic acids (PNAs), locked nucleic acids (LNA™), and the like. In certain embodiments, a nucleic acid can be a polymer that includes multiple monomer types, e.g., both RNA and DNA subunits. A nucleic acid can be or can include, e.g., a chromosome or chromosomal segment, a vector (e.g., an expression vector), an expression cassette, a naked DNA or RNA polymer, the product of a polymerase chain reaction (PCR), an oligonucleotide, a probe, a primer, etc. A nucleic acid may be, e.g., single-stranded, double-stranded, triple-stranded (and the like) and is not limited to any particular length. Unless otherwise indicated, a particular nucleic acid sequence optionally comprises or encodes complementary sequences, in addition to any sequence explicitly indicated.

[0029] Nucleic acids are not limited to molecules having naturally occurring polynucleotide sequences or structures, naturally occurring backbones, and/or naturally occurring internucleotide linkages. For example, nucleic acids containing one or more carbocyclic sugars are also included within this definition (Jenkins et al. (1995) Chem. Soc. Rev. pp 169-176, which is incorporated by reference). To further illustrate, although a nucleic acid will generally contain phosphodiester bonds, in some cases nucleic acid analogs are included that have alternate backbones. These may include, without limitation, phosphoramidate (Beaucage et al. (1993) Tetrahedron 49(10): 1925 and the references therein; Letsinger (1970) J. Org. Chem. 35:3800; Sprinzl et al. (1977) Eur. J. Biochem. 81:579; Letsinger et al. (1986) Nucl. Acids Res. 14: 3487; Sawai et al. (1984) Chem. Lett. 805; Letsinger et al. (1988) J. Am. Chem. Soc. 110:4470; and Pauwels et al. (1986) *Chemica Scripta* 26:1419), phosphorothioate (Mag et al. (1991) Nucleic Acids Res. 19:1437 and U.S. Pat. No. 5,644,048), phosphorodithioate (Brill et al. (1989) J. Am. Chem. Soc. 111:2321), O-methylphosphoroamidite linkages (Eckstein, *Oligonucleotides and Analogues: A Practical Approach*, Oxford University Press (1992)), and peptide nucleic acid backbones and linkages (Egholm (1992) J. Am. Chem. Soc. 114:1895; Meier et al. (1992) Chem. Int. Ed. Engl. 31:1008; Nielsen (1993) Nature 365:566; and Carlsson et al. (1996) Nature 380:207). Other analog nucleic acids include those with positively charged backbones (Denpcy et al. (1995) Proc. Natl. Acad. Sci. USA 92:6097); non-ionic backbones (U.S. Pat. Nos. 5,386,023, 5,637,684, 5,602,240, 5,216,141 and 4,469,863; Angew (1991) Chem. Intl. Ed. English 30: 423; Letsinger et al. (1988) J. Am. Chem. Soc. 110:4470; Letsinger et al. (1994) Nucleoside & Nucleotide 13:1597; Chapters 2 and 3, ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research", Ed. Y. S. Sanghvi and P. Dan Cook; Mesmaeker et al. (1994) Bioorganic & Medicinal Chem. Lett. 4: 395; Jeffs et al. (1994) J. Biomolecular NMR 34:17; Tetrahedron Lett. 37:743 (1996)) and non-ribose backbones, including those described in U.S. Pat. Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, ASC Symposium Series 580, Carbohydrate Modifications in Antisense Research, Ed. Y. S. Sanghvi and P. Dan Cook, which are each incorporated by reference. Several nucleic acid analogs are also described in, e.g., Rawls, C & E News Jun. 2, 1997 page 35. Modifications of the ribose-phosphate backbone may be done to facilitate the addition of additional moieties, such as labeling moieties, or to alter the stability and half-life of such molecules in physiological environments.

[0030] In addition to naturally occurring heterocyclic bases that are typically found in nucleic acids (e.g., adenine, guanine, thymine, cytosine, and uracil), nucleic acid analogs also include those having non-naturally occurring heterocyclic or other modified bases. For instance, certain bases used in nucleotides that act as melting temperature (T_m) modifiers are optionally included. Exemplary of these are 7-deazapurines (e.g., 7-deazaguanine, 7-deazaadenine, and the like); pyrazolo[3,4-d]pyrimidines, propynyl-dN (e.g., propynyl-dU, propynyl-dC, and the like); hypoxanthine; inosine; xanthine; 8-aza derivatives of 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 7-deaza-8-aza derivatives of adenine, guanine, 2-aminopurine, 2,6-diaminopurine, 2-amino-6-chloropurine, hypoxanthine, inosine and xanthine; 6-azacytosine; 5-fluorocytosine; 5-chlorocytosine; 5-iodocytosine; 5-bromocytosine; 5-methylcytosine; 5-propynylcytosine; 5-bromovinyluracil; 5-fluorouracil; 5-chlorouracil; 5-iodouracil; 5-bromouracil; 5-trifluoromethyluracil; 5-methoxymethyluracil; 5-ethynyluracil; 5-propynyluracil; non-naturally occurring bases as de-

scribed by, for example, Seela et al. ((1991) *Helv. Chim. Acta* 74:1790; (1999) *Helv. Chim. Acta* 82:1640); Grein et al. ((1994) *Bioorg. Med. Chem. Lett.* 4:971-976), U.S. Pat. Nos. 5,484,908, 5,645,985, 5,990,303, 5,830,653, 6,639,059, 6,303,315, U.S. Pat. Appln. No. 2003/0092905, and the like.

[0031] "Nucleoside" typically refers to a nucleic acid component that comprises a base or basic group (e.g., comprising at least one homocyclic ring, at least one heterocyclic ring, at least one aryl group, and/or the like) covalently linked to a sugar moiety (e.g., a ribose sugar, etc.), a derivative of a sugar moiety, or a functional equivalent of a sugar moiety (e.g., an analog, such as carbocyclic ring). For example, when a nucleoside includes a sugar moiety, the base is typically linked to a 1'-position of that sugar moiety. As described above, a base can be naturally occurring (e.g., a purine base, such as adenine (A) or guanine (G), a pyrimidine base, such as thymine (T), cytosine (C), or uracil (U)), or non-naturally occurring (e.g., a 7-deazapurine base, a pyrazolo[3,4-d]pyrimidine base, a propynyl-dN base, etc.). Exemplary nucleosides include ribonucleosides, deoxyribonucleosides, dideoxyribonucleosides, carbocyclic nucleosides, and the like. A "nucleotide" refers to an ester of a nucleoside, e.g., a phosphate ester of a nucleoside. To illustrate, a nucleotide can include 1, 2, 3, or more phosphate groups covalently linked to a 5' position of a sugar moiety of the nucleoside. Nucleoside triphosphates and 2'-deoxynucleoside triphosphates or their chemically modified versions may be used as substrates for multiple-nucleotide extension by APP where, for example, one nucleotide is incorporated the extending strand can be further extended. 2',3'-dideoxynucleoside triphosphates, chemically modified versions thereof, or other suitable compounds (e.g., as in US 2005/0037398A1, acycloNMP) may be used as terminators for further extension may be used for single-nucleotide extension. 2',3'-dideoxynucleoside triphosphates may be labeled with radioactivity or fluorescence dye for differentiation from the 3' terminal dideoxynucleotide of oligonucleotide P*. Mixtures of nucleoside triphosphates or 2'-deoxynucleoside triphosphates and 2',3'-dideoxynucleoside triphosphates may also be used.

[0032] A "nucleotide incorporating biocatalyst" refers to a catalyst that catalyzes the incorporation of nucleotides into a nucleic acid. Nucleotide incorporating biocatalysts are typically enzymes. An "enzyme" is a protein- and/or nucleic acid-based catalyst that acts to reduce the activation energy of a chemical reaction involving other compounds or "substrates." A "nucleotide incorporating enzyme" refers to an enzyme that catalyzes the incorporation of nucleotides into a nucleic acid, e.g., during nucleic acid amplification or the like. Exemplary nucleotide incorporating enzymes include, e.g., polymerases, terminal transferases, reverse transcriptases (e.g., RQY polymerase described above), telomerases, polynucleotide phosphorylases, and the like. A "thermostable enzyme" refers to an enzyme that is stable to heat, is heat resistant, and retains sufficient catalytic activity when subjected to elevated temperatures for selected periods of time. For example, a thermostable polymerase retains sufficient activity to effect subsequent primer extension reactions when subjected to elevated temperatures for the time necessary to effect denaturation of double-stranded nucleic acids. Heating conditions necessary for nucleic acid denaturation are well known to persons skilled in the art and are exemplified in, for example, U.S. Pat. Nos. 4,683,202, 4,683,195, and 4,965,188. To further illustrate, a "thermostable polymerase" refers to an enzyme that is suitable for use in a temperature cycling reaction, such as a polymerase chain reaction ("PCR"). For a thermostable polymerase, enzymatic activity refers to the catalysis of the combination of the nucleotides in the proper manner to form primer extension products that are complementary to a template nucleic acid. Exemplary thermostable polymerases are described herein, and others may available to the skilled artisan may also be suitable.

[0033] TdT (terminal deoxynucleotidyl transferase) catalyzes the addition of nucleotides to the 3' terminus of a DNA molecule. Unlike most DNA polymerases it does not require a template. The preferred substrate of this enzyme is a 3'-overhang, but it can also add nucleotides to blunt or recessed 3' ends. Cobalt is a necessary cofactor, however the enzyme catalyzes reaction upon Mg and Mn administration *in vitro*.

[0034] CircLigase™ (Epicentre Biotechnologies), ssDNA Ligase is a thermostable ATP-dependent ligase that catalyzes intramolecular ligation (i.e. circularization) of ssDNA templates having a 5'-phosphate and a 3'-hydroxyl group. In contrast to T4 DNA Ligase and Ampligase™ DNA Ligase, which ligate DNA ends that are annealed adjacent to each other on a complementary DNA sequence, CircLigase ssDNA Ligase ligates ends of ssDNA in the absence of a complementary sequence. The enzyme is therefore useful for making circular ssDNA molecules from linear ssDNA. Circular ssDNA molecules can be used as substrates for rolling-circle replication or rolling-circle transcription. Linear ssDNA of >30 bases is circularized by CircLigase enzyme. Under standard reaction conditions, virtually no linear concatamers or circular concatamers are produced. In addition to its activity on ssDNA, CircLigase enzyme also has activity in ligating a single-stranded nucleic acid having a 3'-hydroxyl ribonucleotide and a 5'-phosphorylated ribonucleotide or deoxyribonucleotide.

[0035] An "oligonucleotide" refers to a nucleic acid that includes at least two nucleic acid monomer units (e.g., nucleotides), typically more than three monomer units, and more typically greater than ten monomer units. The exact size of an oligonucleotide generally depends on various factors, including the ultimate function or use of the oligonucleotide. Typically, the nucleoside monomers are linked by phosphodiester bonds or analogs thereof, including phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoranilidate, phosphoramidate, and the like, including associated counterions, e.g., H⁺, NH₄⁺, Na⁺, and the like, if such counterions are present. Typically, the 3' end linkage of P* is a phosphodiester bond. Oligonucleotides may be prepared by any suitable method, including, but not limited to, isolation of an existing or natural sequence, DNA replication or amplification, reverse tran-

scription, cloning and restriction digestion of appropriate sequences, or direct chemical synthesis by a method such as the phosphotriester method of Narang et al. ((1979) Meth. Enzymol. 68:90-99); the phosphodiester method of Brown et al. ((1979) Meth. Enzymol. 68:109-151); the diethylphosphoramidite method of Beaucage et al. ((1981) Tetrahedron Lett. 22:1859-1862); the triester method of Matteucci et al. ((1981) J. Am. Chem. Soc. 103:3185-3191); automated synthesis methods; or the solid support method described in U.S. Pat. No. 4,458,066, and / or other methods known to those skilled in the art.

[0036] A "primer nucleic acid" or "primer" is a nucleic acid that can hybridize to a target or template nucleic acid and permit chain extension or elongation using, e.g., a nucleotide incorporating biocatalyst, such as a polymerase under appropriate reaction conditions. A primer nucleic acid is typically a natural or synthetic oligonucleotide (e.g., a single-stranded oligodeoxyribonucleotide). Although other primer nucleic acid lengths are optionally utilized, they typically comprise hybridizing regions that range from about 8 to about 100 nucleotides in length. Shorter primer nucleic acids generally require lower temperatures to form sufficiently stable hybrid complexes with template nucleic acids. A primer nucleic acid that is at least partially complementary to a subsequence of a template nucleic acid is typically sufficient to hybridize with the template for extension to occur. A primer nucleic acid may be labeled, if desired, by incorporating a detectable label as described herein.

[0037] The term "probe nucleic acid" or "probe" refers to a labeled or unlabeled oligonucleotide capable of selectively hybridizing to a target or template nucleic acid under suitable conditions. Typically, a probe is sufficiently complementary to a specific target sequence contained in a nucleic acid sample to form a stable hybridization duplex with the target sequence under selected hybridization conditions. A hybridization assay carried out using a probe under sufficiently stringent hybridization conditions permits the selective detection of a specific target sequence. The term "hybridizing region" refers to that region of a nucleic acid that is exactly or substantially complementary to, and therefore capable of hybridizing to, the target sequence. For use in a hybridization assay for the discrimination of single nucleotide differences in sequence, the hybridizing region is typically from about 8 to about 100 nucleotides in length. Although the hybridizing region generally refers to the entire oligonucleotide, the probe may include additional nucleotide sequences that function, for example, as linker binding sites to provide a site for attaching the probe sequence to a solid support. A probe of the invention may be generally included in a nucleic acid that comprises one or more labels (e.g., donor moieties, acceptor moieties, and/or quencher moieties), such as a 5'-nuclease probe, a hybridization probe, a fluorescent resonance energy transfer (FRET) probe, a hairpin probe, or a molecular beacon, which can also be utilized to detect hybridization between the probe and target nucleic acids in a sample. In some embodiments, the hybridizing region of the probe is completely complementary to the target sequence. However, in general, complete complementarity is not necessary (e.g., nucleic acids can be partially complementary to one another); stable hybridization complexes may contain mismatched bases or unmatched bases. Modification of the stringent conditions may be necessary to permit a stable hybridization complex with one or more base pair mismatches or unmatched bases. Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001) provides guidance for suitable modification. Stability of the target/probe hybridization complex depends on a number of variables including length of the oligonucleotide, base composition and sequence of the oligonucleotide, temperature, and ionic conditions. One of skill in the art will recognize that, in general, the exact complement of a given probe is similarly useful as a probe. One of skill in the art will also recognize that, in certain embodiments, probe nucleic acids can also be used as primer nucleic acids.

[0038] The methods described herein may be useful for detecting and/or quantifying a variety of target nucleic acids from a test sample. A target nucleic acid is any nucleic acid for which an assay system is designed to identify or detect as present (or not), and/or quantify in a test sample. Such nucleic acids may include, for example, those of infectious agents (e.g., virus, bacteria, parasite, and the like), a disease process such as cancer, diabetes, or the like, or to measure an immune response. Exemplary "test samples" include various types of samples, such as biological samples. Exemplary biological samples include, for instance, a bodily fluid (e.g., blood, saliva, or spinal fluid), a tissue sample, a food (e.g., meat) or beverage (e.g., milk) product, or the like. Expressed nucleic acids may include, for example, genes for which expression (or lack thereof) is associated with medical conditions such as infectious disease (e.g., bacterial, viral, fungal, protozoal infections) or cancer. The methods described herein may also be used to detect contaminants (e.g., bacteria, virus, fungus, and / or protozoan) in pharmaceutical, food, or beverage products. The methods may be useful to, for example, detect minimal residual disease (e.g., rare remaining cancer cells during remission, especially mutations in the p53 gene or other tumor suppressor genes previously identified within the tumors), and / or measure mutation load (e.g., the frequency of specific somatic mutations present in normal tissues, such as blood or urine).

[0039] Kits for performing the methods described herein are also disclosed. The kit typically includes at least a pair of oligonucleotides (e.g., at least one of the pair being a P* oligonucleotide) for amplifying at least one target nucleic acid from a sample, one or more polyphosphorylating agents described herein, a biocatalyst (e.g., DNA polymerase) and / or corresponding one or more probes labeled with a detectable label. The kit may also include samples containing pre-defined target nucleic acids to be used in control reactions. The kit may also optionally include stock solutions, buffers, enzymes, detectable labels or reagents required for detection, tubes, membranes, and the like that may be used to complete the amplification reaction. In some embodiments, multiple primer sets are included. Other embodiments of

particular systems and kits are also contemplated.

[0040] Magnetic polymer particles are of general utility in various medical and biochemical fields, for example as transport vehicles for the delivery of pharmaceutical products, for diagnostic purposes, for separation and for synthetic purposes. Such particles rely upon their magnetic properties in order to perform these functions: in diagnostic assay applications, for example, application of a magnetic field to a sample containing an analyte bound to magnetic polymer particles allows the isolation of the analyte without the use of centrifugation or filtration; and in therapeutic applications, for example, application of a magnetic field to the patient may serve to target drug-carrying magnetic polymer particles to a desired body site.

[0041] By magnetic is meant herein that the polymer particles contain superparamagnetic crystals. Thus the magnetic polymer particles are magnetically displaceable but are not permanently magnetizable. Many processes for preparing magnetic polymer particles are known, a large number of which involve preparing maghemite- or magnetite-containing polymer particles from pre-formed magnetic iron oxides, e.g. magnetite. Some of processes involved are described in U.S. Pat. No. 4,654,267 (Ugelstad).

[0042] The magnetic polymer particles used in the process of the invention may be any magnetic polymer particle e.g. as described in U.S. Pat. No. 4,654,267. The particles are preferably porous to allow the presence of the superparamagnetic crystals in the pores thereof. The surface of the magnetic particles is normally functionalized to allow coupling of a ligand to the polymer particle, e.g. it may be functionalized to carry any known surface structure such as carboxyl groups, tosyl groups, amino groups, epoxy groups, maleamido groups, thiol groups etc. Hence, the surface may be amine functionalized. Alternatively, an amine functionalized surface can itself be further functionalized to carry other functional groups, e.g. COOH groups.

[0043] The polymer particle is preferably made from combinations of vinylic polymers (e.g. styrene), acrylates and/or methacrylates. The polymeric material may optionally be cross-linked, for example by incorporation of cross-linking agents, for example as comonomers, e.g. divinylbenzene (DVB) or ethyleneglycol dimethacrylate. Appropriate quantities of the cross-linking agents (e.g. comonomers) required will be well known to the skilled person. Preferably the polymer is a cross-linked styrenic polymer (e.g. a styrenedivinylbenzene polymer, surface functionalized by the use of a nitro-group containing comonomer, e.g. nitro-styrene, and subsequent reduction) or a cross-linked (meth)acrylic polymer surface functionalized by the use of an epoxy-group containing co-monomer (e.g. glycidylmethacrylate) and subsequent amination (e.g. by reaction with ethylene diamine).

[0044] The superparamagnetic crystals in the polymer particles used in the process of the invention may be of any material capable of being deposited in superparamagnetic crystalline form in the porous polymer particles. Magnetic iron oxides, e.g. magnetite or maghemite are preferred; however the crystals may be of mixed metal oxides or other magnetic material if desired. The total quantity of crystalline magnetic material present is generally more than 1%, preferably more than 3%, desirably more than or equal to 5% (by weight, e.g. up to 40% wt. The percentage is calculated on a Fe (or equivalent metal in the case of magnetic materials other than iron oxides) weight basis based upon the overall dry weight of the coated particles.

[0045] Polymer particles used in the methods of the present invention may generally have sizes (i.e. diameters) that are in the micrometer range, e.g. 0.3 to 100 μm , especially 0.5 to 50 μm , more especially 0.8 to 8 μm , e.g. 0.8 to 1.2 μm . 1 μm beads are referred.

[0046] Typically the porous particles used will have a surface area of at least 15 m^2/g (measured by the BET nitrogen absorption method), and more preferably at least 30 m^2/g , e.g. up to 700 m^2/g , when corrected to a mean particle diameter of 2.7 μm (i.e. multiply surface area by 2.7/MD, where MD is the mean diameter in micrometers). Similarly scaled, the particle pore volume is preferably at least 0.1 mL/g.

[0047] Typically, the polymer particles are spherical and substantially monodisperse before they are coated and especially preferably remain spherical and substantially monodisperse once they have been coated.

[0048] By substantially monodisperse, it is meant that for a majority of particles, the particles have a coefficient of variation (CV) of less than 20%, for example less than 15%, preferably less than 12%, more preferably less than 11%, still more preferably less than 10% and most preferably no more than about 8%, e.g. 2 to 5%. CV is determined in percentage as

$$\text{CV} = (100 \times \text{standard deviation}) / \text{mean}$$

where mean is the mean particle diameter and standard deviation is the standard deviation in particle size. CV is preferably calculated on the main mode, i.e. by fitting a monomodal distribution curve to the detected particle size distribution. Thus some particles below or above mode size may be discounted in the calculation which may for example be based on about 90% of total particle number (of detectable particles that is). Such a determination of CV may be performable on a Coulter or other particle size analyzer.

[0049] Functionalization of the polymeric material may take place after polymerization by, for example, nitration and subsequent reduction of the thus-formed nitro groups to pendant amine groups; or direct amination, for example by treatment with amino ethanol. As further alternatives, polymeric particles prepared by the well-known Ugelstad two-step swelling process and the improvements thereto disclosed in WO 00/61647 may be used. Porous polymer particles produced according to the processes described in this publication may have magnetic particles deposited in their pores by standard techniques.

[0050] As a further possibility, porous polymer particles may be prepared from nitro styrene and DVB, and magnetic material introduced as taught in U.S. Pat. No. 4,654,267.

[0051] The superparamagnetic polymer beads sold by Life Technologies (Invitrogen Dynal Biotech ASA), under the trade names DYNABEADS, especially DYNABEADS MYONE are especially preferred. DYNABEADS are particularly advantageous since they remain in suspension and do not exhibit magnetic particle sedimentation often associated with other magnetic beads. DYNABEADS also show excellent magnetic mobility compared to other magnetic particles in which high levels of iron are present. DYNABEADS exhibit beneficial kinetics allowing shorter reaction times and higher throughputs. Their non-specific binding is lower than other magnetic beads and their proper use results in a concentration of the desired material taking place resulting in easier and more efficient washing procedures. DYNABEADS MYONE beads are easy to automate and are monodisperse. DYNABEADS MYONE carboxylic acid beads are used generally; however, other functional groups may also be utilized.

[0052] Certain embodiments are further described in the following examples. These embodiments are provided as examples only and are not intended to limit the scope of the claims in any way.

EXAMPLES

Example 1

[0053] Covalent Bead - anti-miRNA Probe bead - Chemical Preparation.

[0054] Schematic is shown in Figure 1.

[0055] Reagents: EDC, miRNA oligo, and MYONE beads (Beads: 1 mL=10 mg=10 billion).

[0056] One ml of MYONE carboxylic acid beads (10 billion beads) are placed in 1.5ml microfuge tube. Beads are washed with 1000 μ L 10 mM NaOH 3 times. Beads are further washed with 10000 μ L deionized H₂O, 4 times. pH is adjusted to about 6.5-7. A reaction master mix is prepared in 1.5 mL tube, as follows:

add H₂O 80 μ L
add 20 μ L 5M NaCl
add 248 μ L DMSO

[0057] 1M EDC is prepared in 5mM aqueous Imidazole-HCl. One mg/5.2 μ L=18 mg/ 100 μ L 5mM aqueous Imidazole is prepared. 192 μ L (2X) of 0.5 mM aqueous anti-miRNA oligo is added to the master mix. Beads are suspended in reaction master mix and treated as follows:

remove the H₂O from the beads
add 100 μ L 1M EDC solution to the reaction master Mix
add the anti-miRNA oligo/Master Mix/EDC to the beads
immediately vortex and sonicate 3 times.
place the tube on a rotator.

[0058] The tube is rotated overnight at room temperature. The beads are further washed as follows: Wash with 1000 μ L DMSO. Wash 2 times with 1000 μ L deionized water. Wash 2 times with 1000 μ L 1X TEX (Tris, EDTA, Triton X-100). Wash 3 times with 1000 μ L 1X TEX and heat for 8 min at 80°C. Beads are stored in 1.00 mL 1X TEX at 4°C.

Example 2

Covalent anti-miRNA bead - Enzymatic Preparation

[0059] Schematic is shown in Figure 2.

3'-Modification and 5'-Phosphorylation of anti-miRNA Probes

[0060] Reagents: Terminal Transferase, ddTTP (TriLink),

EP 2 663 639 B1

miRNA-Probe	Sequence	Tm
Let-7aP	5'AACTATACAACCTACTACCTCA	46.41
Let-7bP	5'AACCACACAACCTACTACCTCA	54.8

3'-Modification Procedure

[0061]

Component	Let-7aP	Let-7bP	Final Concentration
10X Terminal Transferase Buffer	10	10	1X
10X Cobalt Chloride	10	10	1X
20U/ μ L Terminal Transferase	10	10	0.4U/ μ L
1mM ddTTP	10	10	100 μ M
5'P-Probe	30	30	~20 μ M
H ₂ O	30	30	
Total	100	100	
Input Yield Output	19.3OD x 30 μ L	15.5OD x 30 μ L	

[0062] The reaction is incubated at 37°C for 2hrs. The terminal transferase is inactivated by heating the reaction mix at 95°C for 5min.

Phosphorylation Procedure

Component	Let-7aPddT	Let-7bPddT
Carryover from above	100	100
10X Kinase Buffer	20	20
10mM ATP	20	20
10U/ μ L T4 Kinase	15	15
DI Water	45	45
Total (μ L)	200	200
Oligo Output (OD)	F3 1.5 x 50 μ L F4 25.9 x 50 μ L	F3 2.7 x 50 μ L F4 26.9 x 50 μ L

The reaction mix is incubated at 37°C for 1 hr. Purification is done using PD-10 column by desalting.

Bead Ligation with 5'P-Let-7a, b, c, d, e and f Probe-3'ddT

Reagents: SOLID™ P1 DNA Beads (ABI), 5'P-Let-7a, b, c, d, e and f Probes, Ligation Buffer (EpiCentre Biotechnologies), Ligase (EpiCentre Biotechnologies), ATP (Ambion), MnCl₂ (Teknova), PEG6000 (Emerald Biosystems).

Component	Stock Concentration	Volume for 2X (μ L)	Final Concentration (1X)
10X Ligation Buffer	10X	160	1X
50% PEG6000	50%	240	7.5%
25 mM MnCl ₂	25mM	160	2.5mM
250 μ M ATP	250 μ M	160	25 μ M
H ₂ O		80	
Total		800	

EP 2 663 639 B1

Pre-Treatment of beads

[0063] Transferred 300 μ L beads (10million/ μ L x 50 = 500million beads x 6) to a Lo-Bind tube. Placed the Lo-Bind tube in a magnetic rack for 1 min and removed the supernatant. Added 200 μ L 1X TEX (Tris-EDTA-Triton X-100) buffer. Sonicated the beads using Covaris Covalent Declump 1. Placed the Lo-Bind tube in a magnetic rack for 1 min and remove the supernatant. Added 150 μ L the Buffer mix and 150 μ L H₂O. Vortexed. Divided the beads into 6 tubes, 50 μ L each. Placed the Lo-Bind tubes in a magnetic rack for 1 min and removed the supernatant.

Added following reaction mix into the tube: Ligation Reaction

Component	7a	7b	7c	7d	7e	7f
Solid P1 Beads (500million beads)	500 million	500 million	500 million	500 million	500 million	500 million
2X Buffer Mix	100	100	100	100	100	100
5'P-Let-7P -3'-ddT A260	11.8 A260	10.5 A260	4.92 A260	4.44 A260	4.59 A260	4.55 A260
5'P-Let-7P -3'-ddT	40	40	80	80	80	80
H ₂ O	40	40	0	0	0	0
100U/ μ L CircLigase I	20	20	20	20	20	20
Total	200	200	200	200	200	200

Incubated reaction tubes at 60°C overnight on ThermoMixer at 1100rpm.

Bead Clean-up:

[0064] Denaturing Solution: Added 200 μ L Denaturant to 1800 μ L, 1X Denaturing buffer and mixed well. Placed the reaction tube in a magnetic rack for 1 min and remove the supernatant. Added 200 μ L 1X TEX buffer. Sonicated the beads using Covaris Covalent Declump 1. Placed the tube in a magnetic rack for 1 min and removed the supernatant. Added 200 μ L of the Denaturing Buffer, vortexed, and let stand for 1 min. Placed the tube in a magnetic rack for 1 min and removed the supernatant. Added 200 μ L 1X TEX buffer, vortexed and pulsed down. Placed the tube in a magnetic rack for 1 min and removed the supernatant. Repeat twice. Beads are stored (in 200 μ L TEX buffer, ~2.5 million beads/ μ L) at 4°C.

Example 3

Hybridization of anti- miRNA Beads (Hybridization with Cy3-labeled miRNA)

[0065] Fluorescent flow cytometry analysis is done to ensure that anti-miRNA probes are covalently attached to the beads and that Cy3-labeled miRNAs are effectively captured by the anti-miRNA beads.

Reagents: Anti-ath-miR159 Beads, Cy-3-ath-miR159a, Control: non-anti-probe beads, Cy3-ath-miR159a.

[0066] Hybridization is carried out using the following protocol:

[0067] Sonicate the beads with Covalent Declump 1 on Covaris. Aliquot 1 μ L of ath-miR159a beads and 1 μ L Solid P1beads into 1.5mL LoBind tubes separately. Add 100 μ L hybridization buffer. Add 5 μ L 100 μ M Cy3-ath-miR159a and vortex. Shake tubes at 1200rpm for 20min at room temperature. Magnet the beads for 1 mins and remove the supernatant. Add 200 μ L RNA Purification Wash Solution 1 (ABI). Vortex and incubate for 1min before placing tubes on magnetic rack. Magnet and after 1 min, remove the supernatant. Add 200 μ L RNA Purification Wash Solution 2 (ABI). Vortex and incubate for 1min before placing tubes on magnetic rack. Add 200 μ L 0.5X SSPE/T buffer (ABI). Vortex and incubate for 1min before placing tubes on magnetic rack. Briefly spin the tubes, magnet, and after 1 min, remove the supernatant. Add 100 μ L 0.5X SSPE/T Buffer, vortex, and sonicate with an ultrasonic bath briefly. ABI is Applied BioSystems, Inc. miRNAs may be eluted from the beads by de-hybridization using buffers that are known to one skilled in the art, e.g., 10 mM Tris:HCl or 10 mM Tris:HCl with 01 mM EDTA.

Fluorescent Flow Cytometry (FACS) (BD Bioscience) Analysis

[0068] Add about 250 μ L FACS read buffer. Add 10 μ L beads. Read. The beads are fluorescent and the center of the distribution peak of the beads is distinctively different from the control beads. Representative FACS profiles are shown in Figures 3A-3C.

EP 2 663 639 B1

Example 4

[0069]

5 Let-7 miRNA Profiling from Red Blood Cells and Plasma
 Reagents: P1-Let-7a-f Beads, Tempus Buffer (ABI), Finger Tip Fresh Blood
 Blood Sample Collection: 40µL of fresh finger blood sample was collected in a tube containing 0.4µL of 0.5M EDTA (pH 8.0) using a mini finger lancet.

10 [0070] Plasma and Red Blood Cells Separation and Lysis: The blood sample is centrifuged and the plasma is carefully pipetted out. 20µL plasma is mixed with 40µL Tempus buffer. The red blood cells are washed with 100µL 0.9% NaCl, centrifuged, and the supernatant is removed. Washing is repeated 2 more times. 10µL red blood cells are mixed with 20µL Tempus Buffer.

15 [0071] Hyb Sample Serial Dilution: 6µL of the lysate solution prepared above is diluted in 194µL Oligo dT hybridization buffer and vortexed. This is Hyb sample A (1µL blood in 100µL volume). The samples are serially diluted according to following table for the rest of Hyb samples.

Hyb Sample	Oligo dT Lys/Bind Buffer	Sample and Volume	Total Volume, µL	Blood, nL
A				1000
B	135	A: 15µL	150	100
C	135	B: 15µL	150	10
D	135	C: 15µL	150	1
E	135	D: 15µL	150	0.1

20 [0072] Hybridization: 5µL each of Let-7a thru Let-7f beads (~2.5million/µL, 12.5million total each) are combined in a LoBind tube, 200µL TEX buffer is added, and the tubes are vortexed. Tubes/beads are magnetized to remove the supernatant. 6µL of the mixed beads are added into each hybridization sample as prepared above. The beads are shaken at 1100rpm for 20min at room temperature. The beads are magnetized to remove the supernatants. The beads are washed according to the following table. Finally, 1X TE buffer is added and the beads are sonicated with Covalent Declump 1.

Test	Beads	Wash 1, 100µL	Wash 2, 100µL	Suspension, 100µL 1X TE
A	Let-7a-f	Buffer A	Buffer B	1X TE (0.125 million/µL)
B	Let-7a-f	Buffer A	Buffer B	1X TE (0.125 million/µL)
C	Let-7a-f	Buffer A	Buffer B	1X TE (0.125 million/µL)
D	Let-7a-f	Buffer A	Buffer B	1X TE (0.125 million/µL)
E	Let-7a-f	Buffer A	Buffer B	1X TE (0.125 million/µL)

35 [0073] Reverse Transcription (RT): The captured miRNAs (in suspension, above Table) are reverse transcribed to cDNA using TaqMan® MicroRNA Reverse Transcription Kit (ABI) following the recommended protocol. The reactions are incubated at 16°C/30 min, 42°C/30 min, and 85°C/5 min.

40 [0074] TaqMan PCR Reaction: The reverse-transcribed cDNA from the captured miRNAs are quantitatively analyzed using Applied Biosystems miRNA TaqMan assays and TaqMan Universal PCR Master Mix following the recommended protocols. TaqMan real-time PCR reactions are run on Applied Biosystems 7500 instrument using the default cycling parameters. The results are listed in Table 1 and Fig. 4.

Table 1 (C_t values for each Let-7 species are shown)

Blood Volume	Let-7a	Let-7b	Let-7c	Let-7d	Let-7e	Let-7f
NTC	40.00	40.00	40.00	40.00	40.00	39.00
1nL	33.21	32.89	34.48	36.17	39.03	34.34

EP 2 663 639 B1

(continued)

Blood Volume	Let-7a	Let-7b	Let-7c	Let-7d	Let-7e	Let-7f
10nL	31.01	30.99	36.54	32.01	36.08	34.33
100nL	27.74	27.68	34.12	28.24	37.28	30.27
1000nL	24.25	24.52	31.46	25.19	35.32	27.92
10000nL	21.78	21.03	27.50	22.46	32.57	24.19
Linear Data Points	4	4	3	5	2	4
Slope	3.12	3.30	3.31	3.42		3.28
R2						0.99

Example 5: miRNA spiking and recovery test from a plasma sample

[0075] A mixture of Let-7a-f miRNAs was spiked in plasma. The spiked plasma sample was serially diluted 10X with Hyb buffer. The miRNAs were captured and quantified by the beads and TaqMan PCR. Results are shown in Figure 5.

Example 6

[0076] Capturing and quantification of a panel of 96 miRNAs using anti miRNA beads

[0077] The following protocol was used. Dilute 5 μ L fresh blood in 490 μ L Hybridization Dilution buffer (ABI). Spike in 5 μ L of 1nM ath-miR159a RNA Artificial Template. Add anti miRNA beads prepared according to Example 2 with 96 anti miRNA oligonucleotides and shake at 1200rpm for 20min at room temperature to hybridize. Wash the beads with RNA Purification Wash Solution I and RNA Purification Wash Solution II as described previously. Add elution buffer and vortex to mix. Heat bead elution sample in 70°C for 2 minutes and then place immediately on ice. Perform reverse transcription and TaqMan real-time PCR as described in Example 4. The quantitative real-time PCR Ct profiles are shown in Figure 6A. A similar experiment is conducted on 50 microL of human plasma. The results are presented in Figure 6B. miRNA detectability and reproducibility was further confirmed as shown in Figure 9 and the table below:

	Run 1	Run 2
Total miRNA Tested	96	96
Detectable miRNA (Ct<35)	71	70
Undetectable miRNA	25	26
Detection Rate	74.0%	72.9%

Example 7

[0078] miRNA capturing and quantification from saliva. The following protocol was used.

[0079] Collect about 200 μ L saliva and centrifuge briefly. Take 100 μ L of the centrifuged saliva and add to 196 μ L Hybridization buffer. Vortex vigorously for 30 seconds. Spike in 4 μ L of 1nM ath-miR159a RNA Artificial Template. This is Sample E. Dilute Sample E serially according to following table using Hybridization buffer.

Hyb Sample	Hyb Dilution buffer μ L	Stock Sample and Volume	Total Volume μ L	Saliva Input nL/100 μ L
E		Prepared in step 1		33300
D	135	E: 15 μ L	150	3330
C	135	D: 15 μ L	150	333
B	135	C: 15 μ L	150	33.3
A	135	B: 15 μ L	150	3.33

[0080] Add anti miRNA beads to Hyb Samples A, B, C, D, and E prepared above. Shake the hybridization samples

EP 2 663 639 B1

in tubes at 1200rpm for 20min at room temperature. Wash the beads with RNA Purification Wash Solution I and RNA Purification Wash Solution II as described previously. Add elution buffer and vortex to mix. Heat bead elution sample in 70°C for 2 minutes and then place immediately on ice. Perform reverse transcription and TaqMan real-time PCR as described in Example 4. The quantitative real-time PCR Ct profiles are shown in table below and in Figure 7.

5

10

15

	Let-7a	Let-7b	Let-7c	Let-7d	Let-7e	Let-7f	ath-159a	miR320
NTC	40.00	36.58	37.30	40.00	40.00	40.00	37.13	40.00
3nL	34.65	33.11	35.49	38.60	40.00	37.15	29.54	34.09
30nL	31.84	32.13	34.83	38.60	40.00	35.43	27.25	34.24
300nL	29.44	29.77	31.80	34.09	40.00	34.13	23.75	34.04
3000nL	26.97	26.93	29.93	30.52	36.23	30.08	20.33	31.45
30000nL	24.03	24.07	26.96	27.67	33.80	27.50	17.04	28.66

Example 8 Protocols for miRNA Detection and Profiling Using Anti-Probe Beads

20

[0081] Two anti - miRNA bead Panels: A384C6: beads with 386 miRNA targets based on the ABI Megaplex A assay panel. B384C6: beads with 385 miRNA targets based on the ABI Megaplex B assay panel.

[0082] None-Human miRNA Controls: Each panel contains 6 non-human miRNAs: ath-miR159a, cel-miR-2, cel-miR-39, cel-miR-54, cel-miR-238, cel-lin-4. They can be used as external controls at user's choice.

25

[0083] Reagents in the kit: Lysis Buffer, ABC Buffer, Wash Buffer 1, Wash Buffer 2, Elution Buffer, LoBind Tubes (Eppendorf), Human Panel A Beads (1million/microl). Or Human Panel Panel B Beads.

[0084] Reagent Preparation: Add 6mL of Lysis Buffer to the ABC Buffer bottle to make the final ABC Buffer for use. Add 7mL of 100% Ethanol to the Wash Buffer 2 bottle to make the final Wash Buffer 2 for use.

[0085] Reagents required: TaqMan[®] MicroRNA Reverse Transcription Kit (ABI PN: 4366597), TaqMan[®] MicroRNA Assays (ABI PN: 4427975), TaqMan[®] Universal Master Mix II (ABI PN: 4440040)

30

miRNA Purification

35

[0086] Beads Preparation: Vortex the beads thoroughly to suspend the beads in solution. Sonicate the beads by an ultrasonic water bath for 1 minute. Vortex and then aliquot 80uL beads (80 million beads total) into a LoBind 1.5-mL microfuge tube. Magnet the beads for 1 minute and remove the supernatant.

[0087] Sample Preparation: Select (check mark) the sample type you are going to test. Based on your selection, prepare the sample for hybridization according to following procedures:

40

45

50

55

Sample Type	Sample Amount	Preparation
Whole Blood	10uL	Mix 10uL whole blood with 20uL Lysis Buffer and vortex for ~30 seconds. Add 120uL ABC Buffer and vortex for ~30 seconds. Total volume ~ 150 uL
Blood Lysate	30uL	Dilute 30uL blood lysate with 120uL ABC Buffer and vortex. Total volume ~ 150 uL
Plasma	50uL	Mix 50uL plasma with 100uL ABC Buffer and vortex for ~30 seconds. Total volume ~ 150 uL
Serum	50uL	Mix 50uL serum with 100uL ABC Buffer and vortex for ~30 seconds. Total volume ~ 150 uL
Cell Cultures	5 x 10 ⁴ cells in 50uL PBS	Mix the 50uL cell/PBS suspension with 150uL Lysis Buffer and vortex for ~30 seconds. Total volume ~200 uL
Solid Tissue	1-10mg	If needed, grind or homogenize the tissue. Add 30uL Lysis Buffer and vortex for ~30 seconds. Add 120uL ABC Buffer and vortex for ~30 seconds. Total volume ~ 150 uL

EP 2 663 639 B1

(continued)

Sample Type	Sample Amount	Preparation
FFPE	1 dissection (5-10um)	Wash the FFPE sample with xylene (2min x 2), 100%, 95% and 75% ethanol (1min each), H2O (1min). Add 30uL Lysis Buffer and vortex for ~30 seconds. Add 120uL ABC Buffer and vortex for ~30 seconds. Total volume ~ 150 uL
Raw Milk	100uL	Mix 100uL raw milk with 200uL Lysis Buffer and vortex for ~30 seconds. Total volume ~300 uL
Saliva	50uL	Centrifuge 15min at 2000rpm to remove debris. Mix 50uL saliva with 100uL Lysis Buffer and vortex for ~30 seconds. Total volume ~150 uL
Urine	50uL	Centrifuge 15min at 2000rpm to remove debris. Mix 50uL urine with 100uL Lysis Buffer and vortex for ~30 seconds. Total volume ~150 uL

[0088] (Optional) Spike in 2uL of 1nM external control miRNA (following six can be captured by the beads: ath-miR-159a, cel-miR-2, cel-miR-39, cel-miR-54, cel-miR-238, cel-Lin4) and vortex.

[0089] Hybridization: Transfer all the prepared sample into the bead tube prepared above. Using a ThermoMixer, shake the vial at 1200rpm at 30°C for 40 minutes.

[0090] Wash: Magnet the beads for 1 minute and remove the supernatant. Add 100uL Wash Buffer 1. Vortex to resuspend the beads into solution then pulse briefly. Magnet the beads for 1 minute and remove the supernatant. Add 100uL Wash Buffer 2. Vortex briefly to resuspend the beads into solution then pulse briefly. Magnet the beads for 1 minute and remove the supernatant. Repeat step c. Pulse the tube down to collect any residual liquid. Place the tube on magnetic rack for 20 sec, and remove any residual liquid using a fine pipette tip.

[0091] Elution: Add 100uL Elution Buffer and vortex to mix then pulse briefly. Using a ThermoMixer, shake the tube at 1200 rpm at 70°C for 3 minutes then place it immediately on the magnetic stand for 1 minute. Carefully transfer the supernatant (miRNA sample) into a clean LoBind tube. (Optional - this dilution step is only for a full 96-well plate run. The concentrated sample can be directly used). Add 500uL H2O to dilute the sample (total volume ~600uL). Place the sample vial on ice. Store the miRNA sample at -80°C if not used immediately.

Reverse Transcription Reaction

[0092] Following operations are for a panel of 96 miRNA assays in a 96-well plate. Aliquot each 5X RT primer into a 96-well PCR plate, 3uL per assay per well. Prepare miRNA RT mix:

Component	Stock	Volume: 1 Reaction	Volume: 100 Reactions
Nuclease-free dH ₂ O		4.16	416
25 mM each (100 mM total) dNTPs	25mM	0.15	15
50X MultiScribe Reverse Transcriptase	50X	1.0	100
10X RT Buffer	10X	1.5	150
20U/uL AB RNase Inhibitor	20U/uL	0.19	19
Total		7.0	700

[0093] Pipette 7uL miRNA RT mix to each well of the RT plate prepared above. Add 5uL miRNA sample to each well (final total RT volume 15uL/well). Cover the plate with Heavy-Duty sealing film, mix, and centrifuge. Incubate at 16°C/30 min, 42°C/30 min, 85°C/5 min, 4°C hold. Store at -20°C.

TAQMAN PCR Reaction

[0094] Following operations are for a panel of 96 miRNA assays in a 96-well plate. Aliquot 20X TaqMan miRNA assays into a 96-well PCR plate, 1uL per assay per well. Each miRNA assay location should match the corresponding RT plate location. Prepare TaqMan PCR mix.

EP 2 663 639 B1

Component	Stock	Volume (uL): 1 reaction	Volume (uL): 100 reaction
Nuclease-free dH ₂ O		7	700
2X Universal Master Mix II	2X	10	1000
Total		17	1700

[0095] Pipette the PCR mix to each well, 17uL/well. Transfer 2uL RT reaction solution obtained from the RT step into corresponding well location of the TaqMan plate. Total reaction volume 20uL. Run PCR: 95°C/10min, 95°C/15sec & 60°C/60sec for 40 cycles, Set FAM as reporter dye with ROX as Reference.

[0096] Representative results from various tissues, body fluids, cancer cell lines, breast cancer formalin-fixed paraffin-embedded (FFPE) samples, among others are shown in figures XX-XX and tables below. The miRNAs are isolated from various tissues using Laser Capture Microdissection (LCM) technique.

miRNA profiling from human bladder LCM samples

LCM Diameter	Area, um ²	Log(Area)	hsa-Let-7a	hsa-Let-7c	hsa-miR-16	hsa-miR-320	hsa-miR-92a
NTC	0		40.00	40.00	38.27	37.39	40.00
100um	7850	3.8949	35.34	37.56	34.20	35.61	40.00
200um	31400	4.4969	33.19	35.52	31.68	35.50	40.00
1000um	785000	5.8949	27.16	31.17	28.03	33.28	40.00

[0097] Example 9. microRNA profiling from urine

The profiling of microRNA in human urine samples during pregnancy at various stages of pregnancy showed that of the 384 miRNAs tested, 200 were detected in Urine samples using a Ct cutoff of 35. 18 pregnancy related miRNAs consistently showed increased expression with pregnancy progression to second trimester.

[0098] Expression of miR143 and miR145 in urine for patients with balloon injured artery and normal individuals was compared. miR143 and miR145 are significantly down regulated in balloon injured arteries in comparison to normal artery. The results are shown in the table below. A representative miRNA profiling is shown in Figure 17.

Assay ID	Workflow	Sample ID#	Avg.CT	StDev.Ct
miR-143	Indy	Normal	35.72	1.06
miR-143	Indy	Stent Artery	43.66	1.9
miR-143	TILDA	Normal	33.08	0.06
miR-143	TILDA	Stent Artery	39.61	0.55
miR-145	Indy	Normal	36.5	1.63
miR-145	Indy	Stent Artery	43.09	1.69
miR-145	TILDA	Normal	30.73	0.32
miR-145	TILDA	Stent Artery	36.84	0.53

Claims

1. A method for isolating an miRNA from a sample of interest, comprising:

- a) contacting a biological sample with a lysis buffer to form a sample lysate
- b) contacting the sample lysate with an anti-miRNA probe covalently attached to a bead;
- c) incubating the sample lysate- bead mixture under suitable conditions to form hybridized complexes between the miRNA in the lysate and the anti-miRNA probes on the beads;
- d) washing the beads under suitable conditions to remove the unbound sample lysate material; and

e) eluting the miRNA bound to the anti-miRNA probes to form an isolated miRNA sample.

2. The method according to claim 1, further comprising identifying or quantifying the miRNA in the isolated miRNA sample.

3. The method according to claim 2, wherein the identifying or quantifying the miRNA comprises using reverse transcription followed by real-time (RT) or quantitative polymerase chain reaction (q-PCR).

4. The method according to claim 1, wherein the beads are magnetic beads.

5. The method according to claim 4, wherein the beads are carboxylic acid beads.

6. The method according to claim 4, wherein the beads are DNA beads.

7. The method according to claim 1, wherein the anti-miRNA probe is attached to the beads using chemical synthesis or enzymatic synthesis.

8. The method according to claim 7, wherein the enzymatic synthesis comprises:

- a) obtaining an oligonucleotide that has a sequence complementary to an miRNA of interest (anti-miRNA probe);
- b) adding a terminal dideoxynucleotide to the 3' end of the oligonucleotide using a terminal transferase;
- c) adding a 5'-phosphate group to the oligonucleotide using a kinase; and
- d) ligating the oligonucleotide to DNA beads using a single-strand DNA ligase.

9. The method according to claim 1, wherein only one species of anti-miRNA probe is attached to an individual bead.

10. The method according to claim 1, wherein more than one species of anti-miRNA probes are attached to an individual bead.

11. The method according to claim 1, wherein the biological sample is selected from the group consisting of blood, serum, plasma, urine, saliva, cerebrospinal fluid, wound exudates, biopsies, autopsies, tissues, formalin-fixed, paraffin-embedded (PPFE) samples, and organs.

12. A method according to claim 11, wherein the biological sample is a biological fluid, and optionally wherein the biological fluid is blood or serum or saliva.

13. A method of diagnosing a disease or disease progression, comprising:

- a) identifying a set of miRNA markers that are differentially regulated during a causation or progression of a disease;
- b) isolating and measuring the levels of such miRNA markers from a suitable biological sample from a patient in need thereof by a method of any one of the previous claims, at different time points if necessary; and
- c) diagnosing the disease or its progression over time in the patient.

Patentansprüche

1. Verfahren zum Isolieren einer miRNA aus einer Probe von Interesse, umfassend:

- a) Inkontaktbringen einer biologischen Probe mit einem Lysepuffer zum Bilden eines Probenlysats
- b) Inkontaktbringen des Probenlysats mit einer Anti-miRNA-Sonde, die kovalent an ein Kügelchen gebunden ist;
- c) Inkubieren des Probenlysats-Kügelchen-Gemischs unter geeigneten Bedingungen zum Bilden hybridisierter Komplexe zwischen der miRNA in dem Lysat und den Anti-miRNA-Sonden auf den Kügelchen;
- d) Waschen der Kügelchen unter geeigneten Bedingungen zum Entfernen des ungebundenen Probenlysatmaterials; und
- e) Eluieren der an die Anti-miRNA-Sonden gebundenen miRNA zum Bilden einer isolierten miRNA-Probe.

2. Verfahren nach Anspruch 1, ferner umfassend das Identifizieren oder Quantifizieren der miRNA in der isolierten

miRNA-Probe.

- 5
3. Verfahren nach Anspruch 2, wobei das Identifizieren oder Quantifizieren der miRNA die Verwendung von reverser Transkription umfasst, gefolgt von Echtzeit (RT)- oder quantitativer Polymerasekettenreaktion (q-PCR).
4. Verfahren nach Anspruch 1, wobei die Kügelchen magnetische Kügelchen sind.
5. Verfahren nach Anspruch 4, wobei die Kügelchen Carbonsäurekügelchen sind.
- 10
6. Verfahren nach Anspruch 4, wobei die Kügelchen DNA-Kügelchen sind.
7. Verfahren nach Anspruch 1, wobei die Anti-miRNA-Sonde mittels chemischer Synthese oder enzymatischer Synthese an die Kügelchen gebunden ist.
- 15
8. Verfahren nach Anspruch 7, wobei die enzymatische Synthese umfasst:
- a) Beschaffen eines Oligonukleotids, das eine zu einer miRNA von Interesse komplementäre Sequenz aufweist (Anti-miRNA-Sonde);
 - b) Hinzufügen eines terminalen Didesoxynukleotids zum 3'-Ende des Oligonukleotids unter Verwendung einer terminalen Transferase;
 - c) Hinzufügen einer 5'-Phosphatgruppe zu dem Oligonukleotid unter Verwendung einer Kinase; und
 - d) Ligieren des Oligonukleotids an DNA-Kügelchen unter Verwendung einer Einzelstrang-DNA-Ligase.
- 20
9. Verfahren nach Anspruch 1, wobei nur eine Spezies von Anti-miRNA-Sonde an ein einzelnes Kügelchen gebunden ist.
- 25
10. Verfahren nach Anspruch 1, wobei mehr als eine Spezies von Anti-miRNA-Sonden an ein einzelnes Kügelchen gebunden ist.
- 30
11. Verfahren nach Anspruch 1, wobei die biologische Probe ausgewählt ist aus der Gruppe, bestehend aus Blut, Serum, Plasma, Urin, Speichel, Zerebrospinalflüssigkeit, Wundexsudaten, Biopsien, Autopsien, Geweben, Formalin-fixierten, Paraffineingebetteten (FFPE) Proben und Organen.
- 35
12. Verfahren nach Anspruch 11, wobei die biologische Probe eine biologische Flüssigkeit ist und wobei die biologische Flüssigkeit optional Blut oder Serum oder Speichel ist.
13. Verfahren zur Diagnose einer Erkrankung oder eines Krankheitsverlaufs, umfassend:
- a) Identifizieren eines Satzes von miRNA-Markern, die während einer Verursachung oder eines Fortschreitens einer Erkrankung differentiell reguliert sind;
 - b) Isolieren und Messen der Spiegel solcher miRNA-Marker aus einer geeigneten biologischen Probe von einem Patienten, der sie benötigt, durch ein Verfahren nach einem der vorhergehenden Ansprüche, bei Bedarf zu unterschiedlichen Zeitpunkten; und
 - c) Diagnostizieren der Krankheit oder ihres Fortschreitens im Laufe der Zeit in dem Patienten.
- 40
- 45

Revendications

- 50
1. Méthode pour isoler un micro-ARN d'un échantillon d'intérêt, comprenant :
- a) la mise en contact d'un échantillon biologique avec un tampon de lyse pour former un lysat de l'échantillon ;
 - b) la mise en contact du lysat de l'échantillon avec une sonde anti-micro-ARN attachée de manière covalente à une bille ;
 - c) l'incubation du mélange lysat de l'échantillon-bille dans des conditions appropriées pour former des complexes hybridés entre le micro-ARN dans le lysat et les sondes anti-micro-ARN sur les billes ;
 - d) le lavage des billes dans des conditions appropriées pour éliminer la substance de lysat de l'échantillon non liée ; et
 - e) l'éluion du micro-ARN lié aux sondes anti-micro-ARN pour former un échantillon de micro-ARN isolé.
- 55

EP 2 663 639 B1

2. Méthode selon la revendication 1, comprenant en outre l'identification ou la quantification du micro-ARN dans l'échantillon de micro-ARN isolé.
- 5 3. Méthode selon la revendication 2, dans laquelle l'identification ou la quantification du micro-ARN comprend l'utilisation d'une transcription inverse suivie d'une réaction en temps réel (RT) ou quantitative d'amplification en chaîne par polymérase (q-PCR).
4. Méthode selon la revendication 1, dans laquelle les billes sont des billes magnétiques.
- 10 5. Méthode selon la revendication 4, dans laquelle les billes sont des billes d'acide carboxylique.
6. Méthode selon la revendication 4, dans laquelle les billes sont des billes d'ADN.
- 15 7. Méthode selon la revendication 1, dans laquelle la sonde anti-micro-ARN est attachée aux billes à l'aide d'une synthèse chimique ou d'une synthèse enzymatique.
8. Méthode selon la revendication 7, dans laquelle la synthèse enzymatique comprend :
 - 20 a) l'obtention d'un oligonucléotide qui a une séquence complémentaire à un micro-ARN d'intérêt (sonde anti-micro-ARN) ;
 - b) l'addition d'un didésoxynucléotide terminal à l'extrémité 3' de l'oligonucléotide à l'aide d'une transférase terminale ;
 - c) l'addition d'un groupe 5'-phosphate à l'oligonucléotide à l'aide d'une kinase ; et
 - 25 d) la ligature de l'oligonucléotide à des billes d'ADN à l'aide d'une ADN ligase monocaténaire.
9. Méthode selon la revendication 1, dans laquelle seule une espèce de sonde anti-micro-ARN est attachée à une bille individuelle.
- 30 10. Méthode selon la revendication 1, dans laquelle plusieurs espèces de sondes anti-micro-ARN sont attachées à une bille individuelle.
11. Méthode selon la revendication 1, dans laquelle l'échantillon biologique est choisi dans le groupe constitué par du sang, du sérum, du plasma, de l'urine, de la salive, du liquide céphalorachidien, des exsudats de plaies, des biopsies, des autopsies, des tissus, des échantillons fixés au formol et enrobés de paraffine (PPFE) et des organes.
- 35 12. Méthode selon la revendication 11, dans laquelle l'échantillon biologique est un liquide biologique, et éventuellement dans laquelle le liquide biologique est du sang ou du sérum ou de la salive.
- 40 13. Méthode de diagnostic d'une maladie ou de l'évolution d'une maladie, comprenant :
 - 45 a) l'identification d'un ensemble de marqueurs de micro-ARN qui sont régulés différemment lors d'une causalité ou d'une évolution d'une maladie ;
 - b) l'isolement et la mesure des taux de tels marqueurs de micro-ARN d'un échantillon biologique approprié en provenance d'un patient ayant besoin d'une telle mesure par une méthode selon l'une quelconque des revendications précédentes, à différents moments si nécessaire ; et
 - 50 c) le diagnostic de la maladie ou de son évolution dans le temps chez le patient.
- 55




Figure 1: Schematic for making anti-miRNA probe beads using chemical synthesis




Figure 2: Schematic for enzymatic synthesis of anti-miRNA probe beads




Figure 3A: Non-anti-probe beads hybridization with Cy3-labeled Let-7a




Figure 3B: Anti-Let-7a-probe beads hybridization with Cy3-labeled Let-7a




Figure 3C: FACS after de-hybridization of Cy3-labeled Let-7a bond from anti-Let-7a-probe beads




Figure 4: Let-7 family miRNA profiling by Ct values




Figure 5: miRNA detection after addition (spiking) to plasma





Figure 6A: 96 miRNA profiling using anti miRNA beads




Figure 6B: Quantitative real-time PCR Ct profiles of 95 miRNAs from blood




Figure 7: Saliva miRNA profiling using anti-miRNA probe beads




Figure 8: A384C6 Beads Sensitivity and Dynamic Range




Figure 9: miRNA Detectability and reproducibility.




Figure 10A: Let7a - miRNA profiling from human bladder LCM samples; LCM area and Ct correlation




Figure 10B: Let7c - miRNA profiling from human bladder LCM samples; LCM area and Ct correlation




Figure 10C: miR-16 - miRNA profiling from human bladder LCM samples; LCM area and Ct correlation




Figure 11: Breast cancer FFPE sample miRNA profiling




Figure 12: Difference in expression of miRNA levels between breast cancer lines: MBA-MB-21 minus MCF-7




Figure 13: miRNA profiling from raw cow milk




Figure 14: miRNA profiling from urine




Figure 15: MegaPlex A miRNA Profiling from 25 microl Serum




Figure 16: MegaPlex B miRNA Profiling from 25 microl Serum




Figure 17: miRNA expression before and at 15-week of pregnancy.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4683202 A [0016] [0032]
- US 4683195 A [0016] [0032]
- US 4965188 A [0016] [0032]
- US 5035996 A [0016]
- WO 2006081222 A [0016]
- US RE39007 E [0016]
- WO 2006087574 A [0016]
- WO 1994016108 A [0016]
- US 5854033 A [0016]
- US 2004265897 A [0016]
- EP 070685 A [0017]
- US 4889818 A [0018]
- US 5079352 A [0018]
- US 5210015 A [0018]
- US 5436134 A [0018] [0023]
- US 5487972 A [0018]
- US 5658751 A [0018] [0023]
- US 5538848 A [0018]
- US 5618711 A [0018]
- US 5677152 A [0018]
- US 5723591 A [0018]
- US 5773258 A [0018]
- US 5789224 A [0018]
- US 5801155 A [0018]
- US 5804375 A [0018]
- US 5876930 A [0018]
- US 5994056 A [0018]
- US 6030787 A [0018]
- US 6084102 A [0018]
- US 6127155 A [0018]
- US 6171785 B [0018]
- US 6214979 B [0018]
- US 6258569 B [0018]
- US 6814934 B [0018]
- US 6821727 B [0018]
- US 7141377 B [0018]
- US 7445900 B [0018]
- US 20090197254 A [0022]
- US 7387887 B [0023]
- US 4257774 A [0023]
- US 6569927 B [0023]
- US 5644048 A [0029]
- US 5386023 A [0029]
- US 5637684 A [0029]
- US 5602240 A [0029]
- US 5216141 A [0029]
- US 4469863 A [0029]
- US 5235033 A [0029]
- US 5034506 A [0029]
- US 5484908 A [0030]
- US 5645985 A [0030]
- US 5990303 A [0030]
- US 5830653 A [0030]
- US 6639059 B [0030]
- US 6303315 B [0030]
- US 20030092905 A [0030]
- US 20050037398 A1 [0031]
- US 4458066 A [0035]
- US 4654267 A [0041] [0042] [0050]
- WO 0061647 A [0049]

Non-patent literature cited in the description

- **TONG et al.** *Cancer Gene Therapy*, 2009, vol. 16, 206-216 [0004]
- **ALBRETSSEN et al.** *Analytical Biochemistry*, 1990, vol. 189, 40-50 [0004]
- **WU et al.** *Genomics*, 1990, vol. 4, 560-569 [0016]
- **BARANY et al.** *PNAS USA*, 1991, vol. 88, 189-193 [0016]
- **LIZARDI et al.** *Nat.Genet.*, 1998, vol. 19, 225-232 [0016]
- **BANÉR et al.** *Nucleic Acid Res.*, 1998, vol. 26, 5073-5078 [0016]
- **LITTLE et al.** *Clin Chem*, 1999, vol. 45, 777-784 [0016]
- **OSER et al.** *Angew. Chem. Int. Engl.*, 1990, vol. 29 (10), 1167 [0017]
- **SINGH et al.** *Chem Commun*, 1998, vol. 4, 455-456 [0017]
- **PELLESTOR et al.** *European J. Human Gen.*, 2004, vol. 12, 694-700 [0017]
- **AFONINA et al.** *Biotechniques*, 2002, vol. 32, 940-949 [0017]
- **SVANVIK et al.** *Anal Biochem*, 2001, vol. 281, 26-35 [0017]
- **TYAGI et al.** *Nat. Biotechnol.*, 1996, vol. 14, 303-308 [0017]
- **NUTIU et al.** *Nucleic Acids Res.*, 2002, vol. 30, e94 [0017]
- **FRENCH et al.** *Mol. Cell. Probes*, 2001, vol. 15, 363-374 [0017]
- **LI et al.** *Nucleic Acids Res.*, 2002, vol. 30, e5 [0017]

- **CARDULLO et al.** *PNAS*, 1988, vol. 85, 8790-8794 [0017]
- **FIANDACA et al.** *Genome Res*, 2001, vol. 11, 609-611 [0017]
- **NAZARENKO et al.** *Nucleic Acids Res.*, 2002, vol. 30, e37 [0017]
- **WHITCOMBE et al.** *Nat Biotechnol*, 1999, vol. 17, 804-807 [0017]
- **NAZARENKO et al.** *Nucleic Acids Res.*, 1997, vol. 25, 2516-2521 [0017]
- **TODD et al.** *Clin. Chem.*, 2000, vol. 46, 625-630 [0017]
- **MORRISON et al.** *Anal. Biochem.*, 1989, vol. 18, 231-244 [0019]
- **LI et al.** *Nucleic Acids Res.*, 2002, vol. 30 (2), e5 [0019]
- **JAIN et al.** *J. Mol. Biol.*, 1972, vol. 68, 21 [0023]
- **KAPUSEINSKI et al.** *Nuc. Acids Res*, 1979, vol. 6 (112), 3519 [0023]
- **SEARLE ; EMBREY.** *Nuc. Acids Res*, 1990, vol. 18, 3753-3762 [0023]
- **JENKINS et al.** *Chem. Soc. Rev.*, 1995, 169-176 [0029]
- **BEAUCAGE et al.** *Tetrahedron*, 1993, vol. 49 (10), 1925 [0029]
- **LETSINGER.** *J. Org. Chem.*, 1970, vol. 35, 3800 [0029]
- **SPRINZL et al.** *Eur. J. Biochem.*, 1977, vol. 81, 579 [0029]
- **LETSINGER et al.** *Nucl. Acids Res.*, 1986, vol. 14, 3487 [0029]
- **SAWAI et al.** *Chem. Lett.*, 1984, vol. 805 [0029]
- **LETSINGER et al.** *J. Am. Chem. Soc.*, 1988, vol. 110, 4470 [0029]
- **PAUWELS et al.** *Chemica Scripta*, 1986, vol. 26, 1419 [0029]
- **MAG et al.** *Nucleic Acids Res.*, 1991, vol. 19, 1437 [0029]
- **BRILL et al.** *J. Am. Chem. Soc.*, 1989, vol. 111, 2321 [0029]
- **ECKSTEIN.** *Oligonucleotides and Analogues: A Practical Approach.* Oxford University Press, 1992 [0029]
- **EGHOLM.** *J. Am. Chem. Soc.*, 1992, vol. 114, 1895 [0029]
- **MEIER et al.** *Chem. Int. Ed. Engl*, 1992, vol. 31, 1008 [0029]
- **NIELSEN.** *Nature*, 1993, vol. 365, 566 [0029]
- **CARLSSON et al.** *Nature*, 1996, vol. 380, 207 [0029]
- **DENPCY et al.** *Proc. Natl. Acad. Sci. USA*, 1995, vol. 92, 6097 [0029]
- **ANGEW.** *Chem. Intl. Ed. English*, 1991, vol. 30, 423 [0029]
- **LETSINGER et al.** *Nucleoside & Nucleotide*, 1994, vol. 13, 1597 [0029]
- *Carbohydrate Modifications in Antisense Research.* ASC Symposium Series 580 [0029]
- **MESMAEKER et al.** *Bioorganic & Medicinal Chem. Lett.*, 1994, vol. 4, 395 [0029]
- **JEFFS et al.** *J. Biomolecular NMR*, 1994, vol. 34, 17 [0029]
- *Tetrahedron Lett.*, 1996, vol. 37, 743 [0029]
- **RAWLS.** *C & E News*, 02 June 1997, 35 [0029]
- **SEELA et al.** *Helv. Chim. Acta*, 1991, vol. 74, 1790 [0030]
- *Helv. Chim. Acta*, 1999, vol. 82, 1640 [0030]
- **GREIN et al.** *Bioorg. Med. Chem. Lett.*, 1994, vol. 4, 971-976 [0030]
- **NARANG et al.** *Meth. Enzymol.*, 1979, vol. 68, 90-99 [0035]
- **BROWN et al.** *Meth. Enzymol.*, 1979, vol. 68, 109-151 [0035]
- **BEAUCAGE et al.** *Tetrahedron Lett.*, 1981, vol. 22, 1859-1862 [0035]
- **MATTEUCCI et al.** *J. Am. Chem. Soc.*, 1981, vol. 103, 3185-3191 [0035]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual.* Cold Spring Harbor Laboratory Press, 2001 [0037]

专利名称(译)	用于分离, 鉴定和量化mirnas的方法		
公开(公告)号	EP2663639A4	公开(公告)日	2015-06-17
申请号	EP2012757380	申请日	2012-01-13
[标]申请(专利权)人(译)	生命技术公司		
申请(专利权)人(译)	Life Technologies公司		
当前申请(专利权)人(译)	Life Technologies公司		
[标]发明人	XU TOM TRINH CHRISTOPHER BORNARTH CAROLE EVANS BRIAN RATH MOUSUMI TRAN KATHY		
发明人	XU, TOM TRINH, CHRISTOPHER BORNARTH, CAROLE EVANS, BRIAN RATH, MOUSUMI TRAN, KATHY		
IPC分类号	C12N15/10 C12Q1/68 C12Q1/48 G01N33/53 C07H21/02		
CPC分类号	C07H21/02 C12N15/1006 C12Q1/6809 C12Q1/6834 C12Q1/6876 C12Q2600/178 C12Q2525/207 C12Q2563/149		
代理机构(译)	韦伯BIRGIT		
优先权	61/432874 2011-01-14 US		
其他公开文献	EP2663639B1 EP2663639A2		
外部链接	Espacenet		

摘要(译)

公开了用于分离, 鉴定和定量miRNA和其他小RNA的方法和组合物和试剂盒, 包括但不限于siRNA, mRNA和snRNA。还公开了诊断疾病或其进展的方法。