



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 206 493 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
15.11.2006 Bulletin 2006/46

(21) Application number: **00953792.9**

(22) Date of filing: **02.08.2000**

(51) Int Cl.:
C07K 16/00 (2006.01)

(86) International application number:
PCT/US2000/021036

(87) International publication number:
WO 2001/009190 (08.02.2001 Gazette 2001/06)

(54) **MONOCLONAL ANTIBODIES TO TACROLIMUS AND IMMUNOASSAY METHODS FOR TACROLIMUS**

MONOKLONALE ANTIKÖRPER GEGEN TACROLIMUS UND IMMUNOTESTVERFAHREN FÜR TACROLIMUS

ANTICORPS MONOCLONAUX DIRIGES CONTRE LE TACROLIMUS ET PROCEDES DE DOSAGE IMMUNOLOGIQUE DU TRACROLIMUS

(84) Designated Contracting States:
DE ES FR IT

(30) Priority: **03.08.1999 US 368010**

(43) Date of publication of application:
22.05.2002 Bulletin 2002/21

(73) Proprietor: **DADE BEHRING INC.**
Deerfield, Illinois 60015 (US)

(72) Inventors:
• **KASPER, Kenneth, C.**
Monte Sereno, CA 95030 (US)
• **JEONG, Henry**
Palo Alto, CA 94306 (US)
• **DAVALIAN, Dariush**
San Jose, CA 95124 (US)
• **LIU, Hshiou-Ting**
Milpitas, CA 95035 (US)
• **MILLER, Paul, L.**
South San Francisco, CA 94080 (US)
• **WILLIAMS, Denise, L.**
San Jose, CA 95123 (US)

(74) Representative: **Schweitzer, Klaus et al**
Zounek Plate Schweitzer
Patentanwaltskanzlei
Rheingastrasse 196
65203 Wiesbaden (DE)

(56) References cited:
EP-A- 0 293 892 **WO-A-94/25022**
US-A- 5 635 406

- **MURTHY JAYASIMHA N ET AL:** "Tacrolimus metabolite cross-reactivity in different tacrolimus assays." **CLINICAL BIOCHEMISTRY**, vol. 31, no. 8, November 1998 (1998-11), pages 613-617, XP000973332 ISSN: 0009-9120
- **ALAK ALA M:** "Measurement of tacrolimus (FK506) and its metabolites: A review of assay development and application in therapeutic drug monitoring and pharmacokinetic studies." **THERAPEUTIC DRUG MONITORING**, vol. 19, no. 3, June 1997 (1997-06), pages 338-351, XP000973285 ISSN: 0163-4356

EP 1 206 493 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionFIELD OF THE INVENTION

5 **[0001]** This invention is directed to monoclonal antibodies to tacrolimus, methods for producing such monoclonal antibodies, and derivatives of tacrolimus useful for producing such monoclonal antibodies.

BACKGROUND OF THE INVENTION

10 **[0002]** Tacrolimus is a macrolide isolated from *Streptomyces tsukubaensis*. Tacrolimus has the chemical name [3S-[3R*[E(1S*, 3S*, 4S*)], 4S*, 5R*, 8S*, 9E, 12R*, 14R*, 15S*, 16R*, 18S*, 19S*, 26aR*]]-5, 6, 8, 11, 12, 13, 14, 15, 16, 17, 18, 19, 24, 25, 26, 26a-hexadecahydro-5,19-dihydroxy-3-[2-(4-hydroxy-3-methoxycyclohexyl)-1-methylethenyl]-14, 16-dimethoxy-4, 10, 12, 18-tetramethyl-8-(2-propenyl)-15, 19-epoxy-3H-pyrido[2,1-c][1,4]-oxaazacyclotricosine-1,7,20,21(4H, 23H) tetrone. The structure of tacrolimus, giving the numbering, is shown below in Figure 16.

15 **[0003]** Tacrolimus is also known as FR-900506 or FK-506. Tacrolimus has immunosuppressive activity and antimicrobial activity.

[0004] The immunosuppressive activity of tacrolimus is particularly important and has led to the increasingly wide use of this drug. Immunosuppression is used clinically in a number of contexts, most importantly in preventing rejection in organ transplantation. Immunosuppressive drugs are also administered in prevention of Rh hemolytic disease of the newborn and in the treatment of autoimmune disorders. Tacrolimus inhibits T-cell activation by binding to a cytosolic protein known as FKBP (FK506 binding protein). The drug-binding protein complex stably associates with calcineurin. This inhibits the serine-threonine phosphatase activity of this Ca²⁺-dependent enzyme. This inhibits calcineurin-dependent activation of lymphokine expression, apoptosis, and degranulation (G. Wiederrecht et al., "The Mechanism of Action of FK-506 and Cyclosporin A," Ann. N.Y. Acad. Sci. 696:9-19 (1993)).

25 **[0005]** Tacrolimus can be administered intravenously in a short or continuous infusion or orally. Tacrolimus, like other immunosuppressant agents, has a spectrum of toxicity. The major toxicity associated with clinical use of the drug is nephrotoxicity. In addition, neurotoxicity can develop, associated with headache, tremor, insomnia, pain, or other symptoms. Additionally, gastrointestinal toxicity manifested by diarrhea or nausea can develop, as can cardiovascular toxicity manifested by hypertension.

30 **[0006]** Additionally, metabolic toxicity can develop as manifested by the development of such symptoms as hyperkalemia, hypomagnesemia, or hyperglycemia. In addition, long-term immunosuppression with tacrolimus can produce increased risk of all types of infections, not only the usual bacterial, viral, and fungal pathogens, but also various unusual opportunistic infections as well.

35 **[0007]** Additionally, there is an increased risk of lymphomas and related malignancies associated with the administration of tacrolimus (M.L. Cleary & J. Sklar, "Lymphoproliferative Disorders in Cardiac Transplant Recipients are Multiclonal Lymphomas," Lancet 2:49-493 (1984); L.J. Swinnen et al., "Increased Incidence of Lymphoproliferative Disorder After Immunosuppression with a Monoclonal Antibody OKT3 in Cardiac-Transplant Recipients," N. Engl. J. Med. 323: 1723-1728 (1990)). At least some of these malignancies are related to impaired immune responses to Epstein-Barr virus (B.Z. Katz et al., "Latent and Replicating Forms of Epstein-Barr virus DNA and Lymphomas in Lymphoproliferative Diseases," J. Infect. Dis., 160:589-598 (1989)).

40 **[0008]** The potency and the spectrum of toxicities of tacrolimus requires sensitive, reproducible, and reliable methods for monitoring the blood concentration of these compounds after administration to a patient, such a patient undergoing organ transplantation. It is important that such methods be sensitive enough to detect low concentrations of tacrolimus. It is also important that such methods be reliable and reproducible, and avoid interference from compounds such as metabolites of tacrolimus.

45 **[0009]** EP-A 293 892 and Clinical Biochemistry, Vol. 31, 1998, pg. 613-617 both disclose monoclonal and polyclonal antibodies to tacrolimus. However, these references do not disclose monoclonal antibodies to tacrolimus having limited cross-reactivity to all of the following tacrolimus metabolites; 15-demethyl tacrolimus; 31-demethyl tacrolimus; 13,31-didemethyl tacrolimus; 15,31-didemethyl tacrolimus; and 12-hydroxy tacrolimus.

50 **[0010]** Although antibodies and immunoassays to tacrolimus exist, and are described, for example, in U.S. Patent No. 5,532,137 to Niwa et al. there is still a need for the development of improved antibodies and immunoassays specific for tacrolimus. There is, particularly, a need for improved monoclonal antibodies to tacrolimus that can be used in developing a sensitive, reliable, and reproducible immunoassay for tacrolimus.

55 **[0011]** The development of a reliable immunoassay for tacrolimus is complicated by the fact that tacrolimus has a number of metabolites that are found in the blood of an individual being treated with tacrolimus. The conversion of tacrolimus to these metabolites involve demethylation, hydroxylation, and ring formation. It is important that antibodies to tacrolimus have as little cross-reactivity with these derivatives as possible.

[0012] There is therefore a need for the development of improved monoclonal antibodies to tacrolimus that are useful

for immunoassays for tacrolimus and that possess a minimal degree of cross-reactivity to tacrolimus metabolites. There is also a need for improved immunoassays using such monoclonal antibodies for the detection and determination of tacrolimus in the blood of patients being administered tacrolimus.

5 SUMMARY

[0013] We have discovered that tacrolimus, when derivatized in the non-binding domain, can be coupled to a high molecular weight carrier such as a protein for immunization to produce antibodies. The resulting antibody-producing cells can be used to generate monoclonal antibodies by cell fusion. These monoclonal antibodies have desirable properties, including reduced cross-reactivity with tacrolimus metabolites.

[0014] One embodiment of the present invention is a monoclonal antibody to tacrolimus that is a monoclonal antibody designated 1H6 and which has less than about 8% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus and 12-hydroxy tacrolimus.

[0015] Another embodiment of the present invention is a monoclonal antibody to tacrolimus that:

- (1) competes with the IgG₁λ monoclonal antibody designated 1H6 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody designated 1 H6 as measured by competition assays; and
- (2) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus, and 12-hydroxy tacrolimus. Preferably, the monoclonal antibody to tacrolimus competes at least about 90% as effectively on a molar basis as the monoclonal antibody designated 1 H6 and has less than about 8% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus and 12-hydroxy tacrolimus.

[0016] Another embodiment of the present invention is a hybridoma producing the IgG₁λ monoclonal antibody to tacrolimus designated 1H6 as described above.

[0017] Yet another embodiment of the present invention is a hybridoma producing a monoclonal antibody that:

- (1) competes with the IgG₁λ monoclonal antibody designated 1H6 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody designated 1 H6 as measured by competition assays; and
- (2) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus, and 12-hydroxy tacrolimus.

[0018] Another embodiment of the present invention is a monoclonal antibody that:

- (1) competes with the IgG₁λ monoclonal antibody designated 1H6 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody designated 1H6 as measured by competition assays; and
- (2) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus, and 12-hydroxy tacrolimus, wherein at least some of the constant regions of the antibody are replaced by human constant regions so that the monoclonal antibody is humanized.

[0019] Yet another embodiment of the present invention is a single-chain recombinant antibody (sFv) including therein the variable regions of an antibody to tacrolimus that:

- (1) competes with the IgG₁λ monoclonal antibody designated 1H6 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody designated 1 H6 as measured by competition assays; and
- (2) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus, and 12-hydroxy tacrolimus.

[0020] Still another embodiment of the present invention is a monoclonal antibody to tacrolimus produced by fusion of antibody-producing cells from an antibody-producing mammal immunized with tacrolimus derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus conjugated to a high molecular weight protein with a suitable fusion partner. Preferably, the carbon atom in the non-binding domain of tacrolimus is carbon 22. Preferably, the high molecular weight protein is keyhole limpet hemocyanin.

[0021] Yet another embodiment of the present invention is a monoclonal antibody to tacrolimus that is an IgG₁λ monoclonal antibody, that has a binding affinity for tacrolimus of about 3.7×10^9 liters/mole, that cross-reacts with 13-demethyl tacrolimus, and that has less than about 8% cross-reactivity to all of the following tacrolimus metabolites: 15-demethyl tacrolimus; 31-demethyl tacrolimus; 13, 31-didemethyl tacrolimus; 15, 31-didemethyl tacrolimus; and 12-hy-

droxy tacrolimus.

[0022] Another aspect of the present invention is antibodies, including polyclonal antibodies, produced by immunization of an antibody-producing mammal with tacrolimus derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus conjugated to a high molecular weight protein. Preferably, the carbon atom in the non-binding domain of tacrolimus is carbon 22. Preferably, the high molecular weight protein is keyhole limpet hemocyanin.

[0023] Another aspect of the present invention is a conjugate comprising an antibody according to the present invention directly or indirectly conjugated to a detectable label. The detectable label can be selected from the group consisting of an enzyme label, a radioactive label, a fluorescent label, a chemiluminescent label, a bioluminescent label, and a particulate label. In many applications, the label is preferably an enzyme label.

[0024] Still another aspect of the present invention is a method of detecting or determining tacrolimus comprising the steps of:

(1) providing a sample suspected of containing tacrolimus;

(2) reacting the sample with:

(a) an antibody according to the present invention; and

(b) optionally, a tacrolimus analogue; wherein one of the antibody or the tacrolimus analogue is labeled with a label producing a detectable signal; and

(3) observing or measuring one of:

(a) the signal associated with tacrolimus bound to antibody;

(b) the signal associated with tacrolimus unbound to antibody; or

(c) the total signal present;

in order to detect or determine the presence or concentration of tacrolimus in the sample.

[0025] Typically, in this method, the sample is reacted with a tacrolimus analogue labeled with an enzyme label and the total signal present is observed or measured to detect or determine the presence or concentration of tacrolimus in the sample.

[0026] Another aspect of the present invention is a test kit comprising, packaged in separate containers:

(1) an antibody according to the present invention; and

(2) a tacrolimus analogue labeled directly or indirectly with an enzyme label.

[0027] Yet another aspect of the present invention is a derivative of tacrolimus comprising tacrolimus that is derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus. Preferably, the carbon atom in the non-binding domain of tacrolimus is carbon 22.

[0028] Still another aspect of the present invention is a conjugate comprising a derivative of tacrolimus as described above conjugated to a high molecular weight protein. Preferably, the high molecular weight protein is keyhole limpet hemocyanin.

[0029] Yet another aspect of the present invention is a method of derivatizing tacrolimus comprising reacting tacrolimus with carboxymethoxylamine to produce a carboxymethyl oxime derivative of tacrolimus, the carboxymethyl oxime moiety being located at carbon atom 22.

[0030] Still another aspect of the present invention is a method of producing a conjugate of tacrolimus with a high molecular weight protein comprising:

(1) reacting tacrolimus with carboxymethoxylamine to produce a carboxymethyl oxime derivative of tacrolimus, the carboxymethyl oxime moiety being located at carbon atom 22;

(2) activating the carboxymethyl oxime to produce a reactive N-hydroxysuccinimide ester; and

(3) reacting the N-hydroxysuccinimide ester with the high molecular weight protein to produce the conjugate.

[0031] Another aspect of the present invention is a derivative of tacrolimus comprising tacrolimus substituted with a carboxymethyl oxime moiety at carbon atom 22 linked through a linker to a biotin moiety. In one preferred embodiment of this aspect of the present invention, the linker has the structure $\text{NH}_2\text{-CH}_2\text{-CH}_2\text{-NH-CO-(CH}_2\text{)}_5\text{-NH}_2$, and one amine group of the linker forms an amide bond with the carboxyl group of the carboxymethyl oxime and the other amine group of the linker forms an amide bond with the carboxyl group of the biotin.

[0032] Still another aspect of the present invention is a method of derivatizing tacrolimus comprising:

- (1) reacting tacrolimus with carboxymethoxylamine to produce a carboxymethyl oxime derivative of tacrolimus, the carboxymethyl oxime derivative being located at position 22;
- (2) activating the carboxymethyl oxime to produce a reactive N-hydroxysuccinimide ester; and
- (3) reacting the N-hydroxysuccinimide ester with the carboxyl group of biotin or a biotin derivative or analogue.

[0033] Yet another aspect of the present invention is a derivative of tacrolimus comprising tacrolimus that is derivatized with a bromoacetyl moiety at a carbon atom in the non-binding domain of tacrolimus. Preferably, the carbon atom in the non-binding domain is carbon atom 22. These derivatives can then be reacted with an enzyme or other protein to produce a conjugate containing the derivative conjugated to a protein, such as an enzyme. In one preferred embodiment, the protein is a cysteine-containing mite of glucose-6-phosphate dehydrogenase.

[0034] Accordingly, another aspect of the present invention is a method of derivatizing tacrolimus comprising:

- (1) reacting tacrolimus with carboxymethoxylamine to produce a carboxymethyl oxime derivative of tacrolimus, the carboxymethyl oxime derivative being located at position 22;
- (2) activating the carboxymethyl oxime to produce a reactive N-hydroxysuccinimide ester; and
- (3) reacting the N-hydroxysuccinimide ester with the trifluoroacetic acid salt of bromoacetyl ethylenediamine to produce a bromoacetyl derivative of tacrolimus.

BRIEF DESCRIPTION OF THE DRAWINGS

[0035] These and other features, aspects, and advantages of the present invention will become better understood with reference to the following description, appended claims, and accompanying drawings where:

Figure 1 is a mass spectrogram of the product resulting from the reaction of carboxymethoxylamine with the macrolide antibiotic FK-520, closely related in structure to tacrolimus;

Figure 2 is a portion of the mass spectrogram of Figure 1, shown in expanded resolution centered around the predominant peak;

Figure 3 is a mass spectrogram of the product resulting from the reaction of carboxymethoxylamine with tacrolimus;

Figure 4 is a portion of the mass spectrogram of Figure 3, shown in expanded resolution centered around the predominant peak;

Figure 5 is a ¹H NMR spectrum at 250 MHz in CDCl₃ of the product resulting from the reaction of carboxymethoxylamine with tacrolimus;

Figure 6 is a ¹³C NMR spectrum at 250 MHz in CDCl₃ of the product resulting from the reaction of carboxymethoxylamine with tacrolimus;

Figure 7 is a ¹³C NMR spectrum at 500 MHz in CDCl₃ of tacrolimus shown as a reference;

Figure 8 is a mass spectrogram of the product resulting from the reaction of LC-biotin with tacrolimus monooxime;

Figure 9 is a portion of the mass spectrogram of Figure 8, shown in expanded resolution centered around the predominant peak;

Figure 10 is a ¹H NMR spectrum at 250 MHz in CDCl₃ of the product resulting from the reaction of LC-biotin with tacrolimus monooxime;

Figure 11 is a calibration curve of a homogeneous enzyme immunoassay of tacrolimus using a monoclonal antibody produced by immunization of mice with a conjugate of tacrolimus derivatized at position 22 with a carboxymethyl oxime linked to keyhole limpet hemocyanin and cell fusion of the resulting antibody-producing cells with a fusion partner;

Figure 12 is a graph showing the correlation between the results for the assay of tacrolimus using a homogeneous enzyme immunoassay with the monoclonal antibody for which the calibration curve is shown in Figure 11 and the results for the assay of tacrolimus using a method employing gas chromatography and tandem mass spectroscopy (LC/MS/MS);

Figure 13 is a graph showing the correlation between the results for the assay of tacrolimus using the homogeneous enzyme immunoassay and using LC/MS/MS on a panel of 70 patients with liver damage to whom tacrolimus had been administered;

Figure 14 is a graph showing the correlation between the results for the assay of tacrolimus using the homogeneous enzyme immunoassay and using a commercially available immunoassay for tacrolimus for the panel of 70 patients of Figure 13;

Figure 15 is a graph showing a Bland-Altman difference analysis plot for the results of Figures 13-14; and

Figure 16 is a drawing of the structural formula for tacrolimus, showing the numbering of the molecule.

DESCRIPTIONDefinitions

- 5 **[0036]** As used herein, the terms defined below have the following meanings unless otherwise indicated:
- [0037]** "Antibody": As used herein the term "antibody" includes both intact antibody molecules of the appropriate specificity, and antibody fragments (including Fab, F(ab'), Fv and F(ab')₂), as well as chemically modified intact antibody molecules and antibody fragments, including hybrid antibodies assembled by *in vitro* association of subunits. Also included are single-chain antibody molecules generally denoted by the term sFv and humanized antibodies in which some or all of the originally non-human constant regions are replaced with constant regions originally derived from human antibody sequences. Both polyclonal and monoclonal antibodies are included unless otherwise specified; in a number of contexts, monoclonal antibodies are specifically specified. Additionally included are modified antibodies or antibodies conjugated to labels or other molecules that do not block or alter the binding capacity of the antibody.
- 10 **[0038]** "Nucleic Acid Sequence": The term "nucleic acid sequence" includes both DNA and RNA unless otherwise specified, and, unless otherwise specified, includes both double-stranded and single-stranded nucleic acids. Also included are hybrids such as DNA-RNA hybrids. In particular, reference to DNA includes RNA that has either the equivalent base sequence except for the substitution of uracil in RNA for thymine in DNA, or has a complementary base sequence except for the substitution of uracil for thymine, complementarity being determined according to the Watson-Crick base pairing rules. Additionally, a reference to a nucleic acid sequence includes its complement according to the Watson-Crick base pairing rules unless otherwise specified.
- 15 **[0039]** We have developed an improved monoclonal antibody to tacrolimus based on the use of tacrolimus derivatized at a carbon atom within the non-binding domain, preferably carbon-22. Polyclonal antibodies produced by immunization of antibody-producing animals with this immunogen are first made. The resulting antibody-producing cells are then used for cell fusion with a suitable fusion partner to produce hybridomas. The resulting monoclonal antibodies produced by
- 20 the hybridomas are particularly useful in immunoassays for the detection of tacrolimus.
- 25

I. DERIVATIVES OF TACROLIMUS

- 30 **[0040]** Accordingly, one aspect concerns derivatives of tacrolimus derivatized at a carbon atom within the non-binding domain of tacrolimus. Preferably, the derivative is derivatized at the carbon-22 position. One particularly preferred class of derivatives involves reacting the keto group at the 22 position with an amine to produce an oxime. A particularly preferred amine is carboxymethylamine. The reaction of tacrolimus with carboxymethyl amine produces a carboxymethyl oxime.
- 35 **[0041]** This reaction involves reacting tacrolimus with carboxymethyl amine in methanol in the presence of sodium acetate to give the oxime. Accordingly, another aspect of the present invention is a method of derivatizing tacrolimus comprising reacting tacrolimus with carboxymethyl amine to produce a carboxymethyl oxime derivative of tacrolimus, the oxime moiety being located at carbon-22. Further details of this reaction are given in Example 1.
- 40 **[0042]** Accordingly, another aspect concerns a conjugate comprising the derivative of tacrolimus which comprises tacrolimus derivatized with a carboxymethyl oxime moiety at position 22 conjugated to a high molecular weight protein through the oxime moiety. Typically, the high molecular weight protein is a protein that is a suitable carrier for haptens and can be, but is not necessarily limited to, proteins such as bovine serum albumin, thyroglobulin, ovalbumin, fibrinogen, or keyhole limpet hemocyanin. A particularly preferred carrier is keyhole limpet hemocyanin. Alternatively, the high molecular weight protein can be an enzyme, such as an enzyme producing a detectable signal, such as glucose-6-phosphate dehydrogenase or alkaline phosphatase. Such conjugates are particularly useful in immunoassays for tacrolimus, as discussed below.
- 45 **[0043]** The preparation of these conjugates is set out in more detail below in Example 2. In general however, a method of preparation of the conjugate of tacrolimus with the high molecular weight protein is as follows: (1) preparation of the carboxymethyl oxime of tacrolimus as described above; (2) activating the carboxymethyl oxime to produce a reactive N-hydroxysuccinimide ester; and (3) reacting the N-hydroxysuccinimide ester with the high molecular weight protein to produce the conjugate. This reaction is discussed in detail in Example 2. The activation of the carboxymethyl oxime to produce the N-hydroxysuccinimide ester typically is performed using a coupling agent such as a water-soluble carbodiimide. A preferred water-soluble carbodiimide is -3-(3-dimethylaminopropyl 1-ethyl-3-dimethylaminopropyl)-carbodiimide hydrochloride (EDAC). Other water-soluble carbodiimides are known in the art and can also be used.
- 50 **[0044]** In addition to forming conjugates with high molecular weight proteins such as keyhole limpet hemocyanin, the present invention also encompasses derivatives of tacrolimus substituted with a carboxymethyl oxime moiety at carbon 22 linked through a linker to a biotin moiety. The linker can take one of a number of forms and can have different lengths. The use of linkers between biotin and a hapten or antigen is well known in the art and need not be described further in detail here. In one particularly preferred derivative, the linker has the structure NH₂-CH₂-CH₂-NH-CO-(CH₂)₅-NH₂. One
- 55

of the amine groups of the linker forms an amide bond with the carboxyl group of the carboxymethyl oxime, and the other amine group forms an amide bond with the carboxyl group of the biotin.

[0045] The length of this spacer can be changed by inserting or deleting one or more CH₂ (methylene) groups at the two places in the spacer that have 2 or more methylene groups.

[0046] The derivatives can be formed by reacting a N-hydroxysuccinimide ester as described above with the carboxyl group of biotin or a biotin derivative or analogue. This reaction can be performed as described above.

[0047] In general, the method of forming such biotin derivatives comprises:

- (1) reacting tacrolimus with carboxymethoxylamine to produce a carboxymethyl oxime derivative of tacrolimus, the carboxymethyl oxime derivative being located at position 22;
- (2) activating the carboxymethyl oxime to produce a reactive N-hydroxysuccinimide ester; and
- (3) reacting the N-hydroxysuccinimide ester with the carboxyl group of biotin or a biotin derivative or analogue.

[0048] In one preferred alternative, as discussed below in Example 3, the oxime is reacted with EDAC and N-hydroxysuccinimide in dimethylformamide (DMF). The activated product is then reacted with the biotin containing the linker, such as LC-biotin.

[0049] Still another embodiment of conjugates of tacrolimus derivatized at a carbon atom within the non-binding domain of tacrolimus, preferably position 22, is a bromoacetyl derivative. The preparation of bromoacetyl derivatives of tacrolimus derivatized at position 22 is described in Example 4. In general, the preparation of such derivatives comprises:

- (1) reacting tacrolimus with carboxymethoxylamine to produce a carboxymethyl oxime derivative of tacrolimus, the carboxymethyl oxime derivative being located at position 22;
- (2) activating the carboxymethyl oxime to produce a reactive N-hydroxysuccinimide ester; and
- (3) reacting the N-hydroxysuccinimide ester with the trifluoroacetic acid salt of bromoacetyl ethylenediamine to produce a bromoacetyl derivative.

[0050] The carboxymethyl oxime derivative used in this method is prepared as described above.

[0051] Such bromoacetyl derivatives can be used to produce enzyme conjugates of tacrolimus by reacting the bromoacetyl moiety with a sulfhydryl group of an enzyme, such as a mutein of glucose 6-phosphate dehydrogenase that contains a cysteine residue. The bromoacetyl derivative can react with other cysteine groups in other enzymes or proteins to produce other enzyme or protein conjugates of tacrolimus.

II. MONOCLONAL AND POLYCLONAL ANTIBODIES AND HYBRIDOMAS

[0052] Another aspect of the present invention is monoclonal antibodies and hybridomas producing them, as well as polyclonal antibodies produced by immunization of antibody-producing animals with tacrolimus derivatives as described above.

[0053] One preferred embodiment of a monoclonal antibody to the present invention is a murine monoclonal antibody to tacrolimus that is a IgG₁λ monoclonal antibody designated 1H6. This monoclonal antibody reacts with 13-demethyl tacrolimus but has less than about 8% cross-reactivity to all of the following tacrolimus metabolites: 15-demethyl tacrolimus; 31-demethyl tacrolimus; 13, 31-didemethyl tacrolimus; 15, 31-didemethyl tacrolimus; and 12-hydroxy tacrolimus. This monoclonal antibody has an estimated affinity for binding to tacrolimus of 3.7 x 10⁹ liters/mole.

[0054] Another monoclonal antibody according to the present invention is a monoclonal antibody to tacrolimus that: (1) competes with the IgG₁λ monoclonal antibody designated 1H6 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody designated 1H6 as measured by competition assays; and (2) has less than about 10% cross reactivity with each of the following tacrolimus metabolites: 15-demethyl tacrolimus; 31-demethyl tacrolimus; 13, 31-didemethyl tacrolimus; 15, 31-didemethyl tacrolimus; and 12-hydroxy tacrolimus.

[0055] Preferably, the monoclonal antibody competes at least about 90% as effectively on a molar basis as the monoclonal antibody designated 1H6 and has less than about 8% cross reactivity with each of these tacrolimus metabolites.

[0056] Another aspect of the present invention is a hybridoma producing a monoclonal antibody as described above. This includes a hybridoma producing the IgG₁λ monoclonal antibody to tacrolimus designated as 1H6. This also includes a hybridoma producing a monoclonal antibody as described above which competes at least about 80% as effectively on a molar basis with that antibody and has a limited degree of cross-reactivity with tacrolimus metabolites as described above.

[0057] Another embodiment of the present invention is humanized monoclonal antibodies. In some applications it is preferred to have humanized monoclonal antibodies in which at least some of the constant regions of the antibody are replaced by human constant regions so that the monoclonal antibody is humanized. Typically, such procedures involve

the grafting of the murine complementarity-determining regions (CDRs) onto a human antibody. Additional alterations of individual amino acids within the framework may be necessary to recreate the antigen-binding site. Typically, this technique involves amplification of cDNA for the heavy and light variable chains of the murine hybridoma using the polymerase chain reaction (PCR) with a subset of synthetic oligonucleotide primers with a small level of degeneracy. Mutagenesis, if required, can be carried out by standard methods. Typically, humanized V genes carrying the variable regions are cloned into suitable expression vectors and expressed in cells such as COS cells or CHO-K1 cells. Other mutations can be made as needed. The preparation of such humanized monoclonal antibodies is generally known in the art and is described, for example in C.A.K. Borrebaeck, "Antibody Engineering" 2d. ed., Oxford University Press, New York, 1995), incorporated herein by this reference.

[0058] Therefore, another embodiment of the present invention is a single-chain recombinant antibody (sFv) including therein the variable regions of an antibody to tacrolimus that: (1) competes with the IgG₁λ monoclonal antibody designated 1H6 and described above at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody designated 1H6 as measured by competition assays; and (2) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus, and 12-hydroxy tacrolimus.

[0059] Strategies for producing sFv single-chain antibodies are well known in the art and are described, for example, in C.A.K. Borrebaeck, "Antibody Engineering," *supra*. In general, constructing an sFv involves manipulation of heavy- and light-chain variable regions, which must be connected at the gene level by an oligonucleotide sequence of codons for an appropriate peptide linker. The linker must connect the V_H and V_L domains of the chosen Fv without perturbing interdomain contacts or interfering with domain folding. A typically used linker is a 15-residue peptide that consists of 3 repeating units, each of which has four glycine residues followed by a serine residue (J.S. Huston et al., "Protein Engineering of Antibody Binding Sites: Recovery of Specific Activity in an Anti-Digoxin Single-Chain Fv Analog Produced in *Escherichia coli*," *Proc. Natl. Acad. Sci. USA* 85:5879-5883 (1988)). Typically, such single-chain antibodies are produced in inclusion bodies in bacteria. Activity typically requires refolding of the expressed polypeptides from inclusion bodies; conditions for performing this are generally well known.

[0060] Accordingly, such single-chain antibodies or sFv are also within the scope of the present invention.

[0061] Another embodiment of the present invention is a monoclonal antibody to tacrolimus produced by fusion of antibody-producing cells from an antibody-producing mammal immunized with tacrolimus derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus conjugated to a high molecular weight protein with a suitable fusion partner. Preferably, the carbon atom in the non-binding domain of tacrolimus is carbon 22. As indicated above, typically, the high molecular weight protein used for immunization is keyhole limpet hemocyanin.

[0062] The preparation of monoclonal antibodies is well known in the art and need not be described further here. For example, the production of monoclonal antibodies is described in J.W. Goding, "Monoclonal Antibodies: Principles and Practice" (2d. ed., Academic Press, London, 1986).

[0063] In general, the first step in the procedure is the production of polyclonal antibodies by standard techniques, such as immunization of an antibody-producing animal such as a mouse, a rat, a goat, a sheep, or a cow with the antigen. The antigen is typically coupled to a high molecular weight carrier such as a high molecular weight protein as discussed above. Immunization can be performed with or without an adjuvant such as complete Freund's adjuvant or other adjuvants such as monophosphoryl lipid A and synthetic trehalose dicorynomycolate adjuvant; it is generally preferred to immunize with an adjuvant. The next step is to isolate spleen cells from antibody-producing animals and fuse the antibody-producing spleen cells with an appropriate fusion partner, typically a myeloma cell, such as by the use of polyethylene glycol or other techniques. Typically, the myeloma cells used are those that grow normally in hypoxanthine-thymidine (HT) medium but cannot grow in hypoxanthine-aminopterin-thymidine (HAT) medium, used for selection of the fused cells. The next step is selection of the fused cells, typically by selection in HAT medium. The next step is to screen the cloned hybrids for appropriate antibody production using immunoassays such as enzyme-linked immunosorbent assay (ELISA) or other immunoassays appropriate for screening. Again, these steps are well known in the art and need not be described in further detail.

[0064] Another embodiment of the present invention is an antibody, which can be polyclonal antibody, to tacrolimus produced by immunization of an antibody-producing mammal with tacrolimus derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus conjugated to a high molecular weight protein as described above. Preferably, the carbon atom in the non-binding domain of tacrolimus is carbon 22.

[0065] Another aspect of the present invention is a conjugate comprising an antibody of the present invention as described above conjugated directly or indirectly to a detectable label. The antibody can be conjugated directly to the detectable label so that a covalent link exists between the label and the antibody. Such methods are well known in the art; such techniques are described, for example, in G.T. Hermanson, "Bioconjugate Techniques" (Academic Press, San Diego, 1996). In general, such techniques link reactive groups on the antibody and the label, typically through a linker or spacer. The length of the linker or spacer can be adjusted to preserve the activity and specificity of the antibody and to ensure that the label produces the detectable signal without interfering with the activity of the antibody.

[0066] Such labels are well known in the art and need not be described in detail. Such labels can include, but are not limited to, an enzyme label, a radioactive label, a fluorescent label, a chemiluminescent label, a bioluminescent label, or a particulate label.

[0067] Enzyme labels are well known in the art. Typically, the enzyme used is one that produces a visually detectable signal such as a colored product or an insoluble product. Among the enzymes that can be used are horseradish peroxidase, β -galactosidase, alkaline phosphatase, glucose oxidase, urease, catalase, lysozyme, malate dehydrogenase, glucose-6-phosphate dehydrogenase, and ribonuclease A. Other enzymes that can be used for immunoassays are well known in the art.

[0068] Among the particulate labels that can be used are latex labels and colloidal metal labels such as colloidal gold, silver, tin, and other metals.

[0069] A large variety of radioactive, fluorescent, chemiluminescent, and bioluminescent labels are known in the art. These labels need not be described further here.

[0070] As an alternative to a direct covalent linkage between the antibody and the label, the linkage can be indirect such as through a biotin-avidin linkage. The binding of biotin to avidin or its bacterial analog streptavidin is well understood in the art. This binding is highly specific and has an extremely high affinity. Typically, the antibody is conjugated to a biotin moiety and the label is bound to avidin or streptavidin. Other arrangements can also be used. The use of an avidin-biotin linkage is described, for example, in G.T. Hermanson, supra, pp. 570-592.

III. IMMUNOASSAYS

[0071] Another aspect of the present invention is immunoassays for tacrolimus using the antibodies described above. In general, such an immunoassay is a method of detecting or determining tacrolimus. The term "detecting" is used herein to refer to a qualitative assay that detects the presence or absence of tacrolimus in a sample, while the term "determining" refers to a quantitative or semiquantitative assay that determines the concentration of tacrolimus in the sample.

[0072] In general, such an immunoassay method comprises the steps of:

- (1) providing a sample suspected of containing tacrolimus;
- (2) reacting the sample with:

- (a) an antibody against tacrolimus as described above; and
- (b) optionally, with a tacrolimus analogue; wherein one of the antibody or the tacrolimus analog is labeled with a label producing a detectable signal; and

- (3) observing or measuring one of:

- (a) a signal associated with tacrolimus bound to antibody;
- (b) a signal associated with tacrolimus unbound to antibody; or
- (c) total signal present to detect or determine the presence or concentration of tacrolimus in the sample.

[0073] Such assays are described as either homogeneous or heterogeneous. In a heterogeneous assay, the antibody bound to antigen is separated from antibody unbound to antigen. This separation can be done by a number of steps well known in the art, such as differential solubility, reactivity with another antibody, or other properties. Such assays are well known in the art and need not be described further in detail here. If either the signal associated with tacrolimus bound to antibody or the signal associated with tacrolimus unbound to antibody is to be detected or determined, a heterogeneous assay is performed. By contrast, in a homogeneous assay, the total signal present is detected or determined. In a homogeneous assay, the existence of an antigen-antibody complex modulates the signal so that the signal level changes without a requirement of separating antigen bound to antibody from antigen unbound to antibody.

[0074] Although many heterogeneous immunoassay formats are well known in the art, in the context of the present invention, it is generally preferred to use a homogeneous immunoassay system. An example of a preferred homogeneous immunoassay system for the immunoassay of tacrolimus using antibodies according to the present invention is the homogeneous assay system known as EMIT, as described in D.D. Schottelius, "Homogeneous Immunoassay System (EMIT) for Quantitation of Antiepileptic Drugs in Biological Fluids" in Antiepileptic Drugs: Quantitative Analysis and Interpretation (C.E. Pippenger et al., Raven Press, New York, 1978, Ch. 10, pp:98-101, and further described in U.S. Patent No. 3,817,837 to Rubenstein et al..

[0075] In general, in this assay, an enzyme such as glucose-6-phosphate dehydrogenase is conjugated to the analyte (here, tacrolimus) to be assayed in such a fashion that it does not significantly alter the activity of the enzyme. However, when this analyte-enzyme conjugate is bound to the antibody for the analyte, the configuration is such that the active site of the enzyme is blocked and thus excludes the substrate. This results in the enzyme activity being reduced. When

the complex is not bound, the active site is available to interact with the substrate. If free analyte is present in the unknown sample, the free analyte then binds to the antibody. This prevents the binding of the analyte-enzyme conjugate and decreases the antibody-induced in activation of the enzyme activity in proportion to the concentration of analyte in the sample. Thus, in such a homogeneous assay, the greater the concentration analyte in the sample, the greater the signal produced by the enzyme label. This is a homogeneous assay as described above.

IV. TEST KITS

[0076] Another aspect of the present invention is a test kit, particularly for use in the EMIT homogeneous immunoassay described above.

[0077] Such a test kit comprises, packaged in separate containers:

- (1) an antibody as described above; and
- (2) a tacrolimus analogue labeled directly or indirectly with an enzyme label.

[0078] The tacrolimus analogue can be labeled either directly or indirectly; it is generally preferred that the tacrolimus analogue is labeled indirectly with a biotinylated tacrolimus molecule and a streptavidinylated enzyme. However, the tacrolimus can be coupled at carbon 22 with an enzyme suitable for use in the EMIT assay as described above.

[0079] Test kits can also include, in separately packaged containers, other ingredients such as substrates, buffers, stabilizers, coenzymes, and other ingredients required for the performance of the immunoassay.

[0080] The invention is illustrated by the following Examples. The Examples are for illustrative purposes only and are not intended to limit the invention.

EXAMPLES

Example 1

Preparation of Carbon-22-Substituted Derivatives of Tacrolimus

[0081] To determine the feasibility of derivatizing tacrolimus at carbon 22, as a model for this molecule, derivatization of the closely related molecule FK-520 was performed. FK-520 has the same structure as tacrolimus except that it has an ethyl instead of an allyl group at carbon 21.

[0082] For these experiments, reagents and solvents were commercial grades and were used as supplied without further purification. The reaction with FK-520 on a small scale (0.9 mg) was performed as follows: To a solution of FK-520 (0.9 mg, 1.1 μmol) in 0.23 mL of methanol was added sodium acetate (1.5 mg, 18.3 μmol , 16 equiv.) and carboxymethoxyl amine hydrochloride (3.0 mg, 13.7 μmol , 12 equiv.). The reaction was stirred at room temperature for 3 hours under argon. Solvent was then evaporated by purging with a stream of argon. Thin layer chromatography in 5:6:1.1:0.5% ($\text{CH}_2\text{Cl}_2/\text{EtOAc}/\text{MeOH}/\text{HOAc}$) on plates from Analtech (scored 10 x 20 cm, 250 microns) of the residue afforded 2 geometric isomers of FK-520 mono-oximes, enough to be analyzed by fast atom bombardment mass spectroscopy. LSIMS of $\text{C}_{45}\text{H}_{72}\text{N}_2\text{O}_{14}$ would be expected to give a $[\text{M}-\text{H}]^-$ of 863.3. The relevant portions of the mass spectroscopy results are shown in Figures 1 and 2. Figure 2 is an portion of the spectrum shown in Figure 1 expanded around the central peak.

[0083] Subsequently an analogous procedure was used to prepare the monooxime of tacrolimus. Tacrolimus was provided as a solid mixture in a capsule containing 1.02 mg of tacrolimus monohydrate and approximately a 6-fold excess of sodium dodecyl sulfate (SDS). The solid mixture from 30 capsules was combined and extracted 4 times with 12 ml each time of ethyl acetate. The combined organic extracts were concentrated under a vacuum to give 49 mg of crude tacrolimus which was used directly in the following reaction.

[0084] To the 49 mg of crude tacrolimus (theoretically 36.5 μmol) in 3.5 ml of methanol was added sodium acetate (16.7 mg, 204 μmol , 5.6 equiv.), and carboxymethoxyl amine hydrochloride (40 mg, 183 μmol , 5 equiv.). The reaction was stirred at room temperature for 12 hours under argon. Solvent and volatile materials were removed at reduced pressure to give crude solid residue. Preparative thin layer chromatography on pre-coated silica gel plates from Analtech (20 x 20 cm, 1000 microns) with 30:70:11:0.6 ($\text{CH}_2\text{Cl}_2/\text{EtOAc}/\text{MeOH}/\text{HOAc}$) of the crude product afforded geometric isomers of tacrolimus monooximes at white solid. LSIMS on $\text{C}_{46}\text{H}_{72}\text{N}_2\text{O}_{14}$ was expected to yield 899.4 as $[\text{M}+\text{Na}]^+$ and 915.4 as $[\text{N}+\text{K}]^+$. Results are shown in Figures 3 and 4. Figure 4 represents the data of Figure 3 at higher resolution centered around the relevant portion of the spectrum.

[0085] ^1H NMR at 250 MHz in CDCl_3 of the product of the reaction of carboxymethoxyl amine hydrochloride with tacrolimus is shown in Figure 5. ^{13}C NMR at 250 MHz in CDCl_3 of the product of the reaction of carboxymethoxyl amine hydrochloride with tacrolimus is shown in Figure 6. ^{13}C NMR at 500 MHz in CDCl_3 shown in Figure 7 was that of tacrolimus

as a reference (i.e. underivatized tacrolimus). All NMR spectra were recorded on Bruker instruments.

Example 2

5 Preparation of Tacrolimus-Keyhole Limpet Hemocyanin Conjugate

[0086] To a solution of tacrolimus monooxime (32.3 mg, 36.8 μmol) in 1.05 mL of anhydrous dimethylformamide was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDAC) (11 mg, 57.4 μmol , 1.5 equiv.) and N-hydroxysuccinimide (7.3 mg, 63.4 μmol , 1.7 equiv.). The reaction was stirred at room temperature for 1 hour under argon. The mixture was then added dropwise via a syringe to a solution of keyhole limpet hemocyanin (74 mg, 54% pure) in 5.0 mL of phosphate buffered saline (0.1 M, pH 8.0) and 0.25 mL of dimethylformamide. After stirring at room temperature for 2 hours, the resulting suspension was dialyzed (1 x 4 L, 4°C, 2 h) against PBS (10 mM, pH 7.0).

[0087] The resulting mixture was then extracted 3 X with CH_2Cl_2 to remove any trace amount of unreacted tacrolimus monooximes. Quantitative analysis of the mixture was conducted using bicinchoninic acid (BCA) protein assay solution to give 50 mg of immunogen in 8 ml of PBS (10 mM, pH 7.0).

[0088] Determination of the hapten number using the TNBS method (A.F.S.A. Habeeb, Anal. Biochem. 14:328 (1966)) gave a hapten number of 1300. The immunogen was immediately frozen using a dry ice-acetone bath and kept at -20°C for storage.

20 Example 3

Preparation of Tacrolimus-Biotin Conjugate

[0089] To a solution of tacrolimus monooxime (12 mg, 13.7 μmol) in 0.3 mL of anhydrous dimethylformamide was added EDAC (4 mg, 20.9 μmol , 1.5 equiv.) and N-hydroxysuccinimide (2.7 mg, 23.4 μmol , 1.7 equiv.). The reaction was stirred at room temperature for 1 h under argon. To this was added triethylamine (10 μL , 75 μmol) and a solution of LC-biotin (10 mg, 25 μmol , 2 equiv.) in 1.0 mL of DMF. Stirring was continued for another 3 h and the mixture was concentration under high vacuum to give a colorless residue. Reversed phase preparative thin-layer chromatography (PTLC) of the crude product on a C-18 plate from Whatman (PKLC-18F, 20 x 20 cm, 1000 microns) (13:7 MeOH/ H_2O) afforded 8 mg of the product as a white solid. The expected result from LSIMS of $\text{C}_{64}\text{H}_{103}\text{N}_7\text{O}_{16}\text{S}$ is $[\text{M}+\text{H}]^+$ of 1258.5. The mass spectroscopy results are shown in Figures 8 and 9. Figure 9 represents a section of the spectrum of Figure 8 at higher resolution centered around the region of interest. ^1H NMR at 350 MHz in CDCl_3 was performed; the results are shown in Figure 10.

[0090] This is an example of derivatization of tacrolimus at position 22.

35 Example 4

Preparation of Bromoacetyl Derivative of Tacrolimus

[0091] Another derivative of tacrolimus derivatized at position 22 is a bromoacetyl derivative. The bromoacetyl derivative can react a sulfhydryl residue of an enzyme or another protein to produce a tacrolimus-enzyme conjugate.

[0092] To form the bromoacetyl derivative of tacrolimus, first, to a solution of succinimidyl bromoacetate (740 mg, 3.14 mmol) in 6 ml of tetrahydrofuran (THF) at 4°C under argon was added, dropwise via a syringe, neat mono-N-BOC-ethylenediamine. After the addition was complete, the reaction was warmed to room temperature and stirred for 3 h. The reaction solution was then concentrated *in vacuo*, and the residue dissolved in ethyl acetate. The ethyl acetate layer was washed with H_2O once, saturated aqueous NaHCO_3 twice, and brine once, dried over MgSO_4 , and evaporated to dryness to give 440 mg of crude white solid. Purification on silica gel using Chromatotron (1:19 MeOH/ CH_2Cl_2) afforded 386 mg of pure bromoacetyl ethylenediamine-mono-t-BOC as a white solid.

[0093] To a solution of bromoacetyl ethylenediamine-mono-t-BOC (50 mg, 0.178 mmol) in 2 ml of CH_2Cl_2 at room temperature was added dropwise via a syringe neat trifluoroacetic acid (TFA) (0.29 ml, 3.76 mmol). The reaction was stirred for 3 h and then concentrated under vacuum. Trace amounts of TFA were azeotropically removed with toluene, and the resulting yellow oily product, the TFA salt of bromoacetyl ethylenediamine, was used directly in the next reaction.

[0094] To a solution of the carboxymethyloxime derivative of tacrolimus (derivatized at position 22) (100 mg, 0.14 mmol) and N-hydroxysuccinimide (NHS) (19 mg, 0.165 mmol) in 2 ml of THF under argon was added via a syringe neat diisopropyl carbodiimide (DIC) (23 μl , 0.147 mmol). The reaction was stirred at room temperature for 3 h and was then transferred via a syringe to a solution of the TFA salt of bromoacetyl ethylenediamine (0.178 mmol) in 3 mL of THF. To this reaction solution was then added neat diisopropylethylamine (DIEA) (40 μl), 0.23 mmol). Stirring was continued for 2.5 h, and the reaction was concentrated *in vacuo* to give a crude oil. Purification on preparative thin-layer chromatography

EP 1 206 493 B1

(TLC) (7:3:1 EtOAc/CH₂Cl₂/MeOH) afforded 44 mg (37%) of bromoacetyl tacrolimus as a colorless solid.

Example 5

5 Preparation of Monoclonal Antibody to Tacrolimus

[0095] Preparation of monoclonal antibody to tacrolimus was performed as follows. The immunogen was the tacrolimus conjugate with keyhole limpet hemocyanin of Example 2. This immunogen was used to immunize Balb/c mice. The first immunization was 25 µg in a volume of 200 µl with monophosphoryl lipid A and synthetic trehalose dicorynomycolate adjuvant (RIBI MPL + TDM Emulsion, RIBI ImmunoChem Research Inc.) intraperitoneally. Five weeks later a boost immunization was given with 25 µg of the immunogen in 200 µl of monophosphoryl lipid A and synthetic trehalose dicorynomycolate adjuvant intraperitoneally. Subsequently, after another 8 weeks, a pre-fusion boost was given of the 25 µg of the immunogen in 200 µl of Hanks' Balanced Salt Solution intravenously and intraperitoneally.

15 [0096] Three days later the fusion was performed by standard methods using a nonsecreting murine myeloma designated P3x63-AG8.653.

[0097] Cloning was done by standard methods.

20 [0098] The clones were screened by the following reverse ELISA immunoassay procedure according to the following protocol. Plates were coated with polyclonal goat anti-mouse IgG (IgG + IgA + IgM) (Zymed) at 5 µg/ml in phosphate buffered saline at 100 µl per well. Plate coating was performed for 2 hours or more at room temperature or overnight at about 4°C; the plates could be stored wrapped in film at about 4°C for several days. The plates were then flicked dry and blocked with 300 µl per well of blocking buffer diluent (0.5% bovine serum albumin, 0.05% Tween 20 in PBS). Plate blocking was performed by incubation for 15 minutes or more at room temperature with plate shaking. The plates were then flicked dry. The monoclonal antibody to be screened was then added to each well as follows: 50 µl per well of blocking buffer diluent was added along with 50 µl per well culture supernatant transferred from the corresponding well in the fusion growth plate. Incubation was for about 1 hour at room temperature with plates shaking. The plate was washed using a Titertek Plus plate washer with S20 stacker with the washing buffer being PBS with 0.05% Tween 20. An enzyme conjugate of tacrolimus covalently coupled to glucose-6-phosphate dehydrogenase diluted in blocking buffer diluent to 1:4000 was added at 100 µl per well. Incubation was performed for about 1 hour at room temperature with shaking. The plate was then washed and a chromogenic solution was added at a volume of 100 µl per well. The chromogenic solution contained 0.593 mM p-iodonitrotetrazolium violet, 0.02 M NAD, 0.033 M glucose-6-phosphate, 0.055 M Tris, 0.02% sodium azide, and a 1:4000 dilution of diaphorase (lipoyl dehydrogenase) (Sigma, St. Louis, MO). BSA was present at 1% (vol/vol) of a 5% w/vol BSA solution. BSA was used to help prevent rapid precipitation of reduced p-iodonitrotetrazolium violet.

25 [0099] From the screening a hybridoma producing a suitable monoclonal antibody was selected. This is designated as 1H6 and is a IgG₁λ antibody. This antibody has a binding affinity for tacrolimus of about 3.7 x 10⁹ liters/mole, that cross-reacts with 13-demethyl tacrolimus, and that has less than about 8% cross-reactivity to all of the following tacrolimus metabolites: 15-demethyl tacrolimus; 31-demethyl tacrolimus; 13, 31-didemethyl tacrolimus; 15, 31-didemethyl tacrolimus; and 12-hydroxy tacrolimus.

40 Example 6

Comparison of Cross-Reactivity with Tacrolimus Metabolites for Monoclonal Antibody of Example 4 and Other Antibodies

45 [0100] The antibody of Example 5, another antibody resulting from the cloning and designated 14H04, and other antibodies, were tested for cross-reactivity of tacrolimus metabolites (13-demethyl tacrolimus, 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13, 31-didemethyl tacrolimus, 15, 31-didemethyl tacrolimus, and 12-hydroxy tacrolimus). All of these metabolites except 12-hydroxy tacrolimus were tested at various levels in hemolysate containing 10 ng/ml of tacrolimus; 12-hydroxy tacrolimus was tested at 10 ng/ml with no tacrolimus in the sample. Methanolic extracts were used in this testing. The standard curve generated for this testing were verified using MORE immunosuppressive drug controls level 1-4, a standard control for these assays (More Diagnostics, Los Osos, California).

50 [0101] All antibodies showed cross-reactivity to 13-demethyl tacrolimus to varying degrees. Cross-reactivity was calculated by the following formula: (cross-reaction value ng/ml)-solvent control value (ng/ml) x 100/metabolite spiked target (ng/ml)=% cross-reactivity.

55 [0102] The results are shown in Table 1. With the exception of 13-demethyl tacrolimus, the 1 H6 antibody of Example 5 had less than 8% cross-reactivity with the other analytes tested.

TABLE 1

CROSS-REACTIVITY OF ANTIBODIES WITH TACROLIMUS METABOLITES						
Metabolite in Hemolysate	Tacrolimus Present	14H04	1H6	6D5-D11	6D5-D12	6D5-F12
13-demethyl tacrolimus 5.0 ng/mL	10 ng/mL	227%	78%	120%	87%	65%
15-demethyl tacrolimus 5.0 ng/mL	10 ng/mL	8%	5%	9%	4%	0
31-demethyl tacrolimus 5.0 ng/mL	10 ng/mL	31%	7%	13%	8%	0
13, 31 di demethyl tacrolimus 5.0 ng/mL	10 ng/mL	10%	0	42%	0	0
15, 31 di demethyl tacrolimus 5.0 ng/mL	10 ng/mL	0	0	29%	14%	0
12-OH tacrolimus 10 ng/mL	0 ng/mL	0	0	0	0	0

Example 7Homogeneous Immunoassay Using Monoclonal Antibody to Tacrolimus

[0103] A homogeneous immunoassay to tacrolimus based on the EMIT procedure was carried out according to the following protocol. A volume (200 μ l) of sample or calibration standard was pipetted into a microfuge tube. Into the same tube, 50 μ l of 300 mM CuSO_4 was pipetted along with 200 μ l of methanol. The mixture was capped and vortexed for 10 seconds and centrifuged for 5 minutes at 20800 x g. The supernatant was decanted into a sample cup of a Roche Cobas Mira analyzer according to standard parameters for that analyzer. The following reagents were placed into the reagent rack of the analyzer: Reagent A contained NaCl, Na_2EDTA , anti-tacrolimus monoclonal antibody (1H6), nicotinamide adenine dinucleotide (NAD), glucose-6-phosphate, detergent (Pluronic 25R2), sodium azide, and *n*-methylisothiazolone at pH 5.5. Reagent B was Tris, pH 8.2, with Na_2EDTA , detergent (Pluronic 25R2), sodium azide, and *n*-methylisothiazolone. Reagent C was a tacrolimus-glucose-6-phosphate dehydrogenase conjugate at pH 7.0, in Na_2HPO_4 , Na_2EDTA , bovine serum albumin, detergent (Pluronic 25R2), and sodium azide. The sample and reagent racks are loaded onto the analyzer and the run is started according to the parameters that have been preset. The enzyme rates are printed out during the run in terms of milli-od/min. The calibration curves are shown in Figure 11 for 1H6 and another monoclonal antibody designated as 14H04.

Example 8Correlation Between Assay for Tacrolimus Using Monoclonal Antibody of the Present Invention and EMIT Assay with LC-MS/MS Method

[0104] A series of samples from 70 patients to whom tacrolimus had been administered was assayed with the EMIT-homogeneous immunoassay using the 1 H6 monoclonal antibody of the present invention as compared with the LC/MS/MS method. The LC/MS/MS method is an assay method for tacrolimus that employs liquid chromatography followed by tandem mass spectroscopy (P.J. Taylor et al., "Sensitive, Specific Quantitative Analysis of Tacrolimus (FK506) in Blood by Liquid Chromatography-Electrospray Tandem Mass Spectrometry," *Clin. Chem.* 42: 279-285 (1996)). The results are shown in Figure 12. The results show a very high correlation with a correlation coefficient of 0.929.

Example 9Comparison of Results Using EMIT Homogeneous Immunoassay and LC/MS/MS Using Specimens from Patients with Liver Dysfunction

[0105] Although monoclonal antibody 1H6 has substantially minimized cross-reactivity to metabolites of tacrolimus other than 13-demethyl tacrolimus, some cross-reactivity remains with 13-demethyl tacrolimus. In order to determine whether or not the cross-reactivity with 13-demethyl tacrolimus would interfere with use of this antibody in an immu-

noassay, a panel of specimens from 70 patients who had severe liver dysfunction and were awaiting transplantation while on tacrolimus was assayed with the EMIT immunoassay using the 1 H6 monoclonal antibody, with the LC/MS/MS method, and with another commercially available immunoassay for tacrolimus, the IMx immunoassay produced by Abbott. Such samples typically contain higher levels of tacrolimus metabolites than do those from patients who have undergone organ transplantation but have normal liver function. Tacrolimus levels were measured on the panel of samples using these three assays.

[0106] The results are shown in Figure 13 for the comparison between the EMIT assay and the LC/MS/MS assay and in Figure 14 for the comparison between the EMIT assay and the IMx assay (Abbott). Deming's regression analysis was used to interpret the results; the results are shown in Table 2. The EMIT assay using the 1H6 monoclonal antibody had closer agreement with LC/MS/MS than the Abbott IMx assay based on the slope of the results (see Table 2) with the slope of 1.11 for EMIT and 1.45 for IMx. Average concentration of the 70 samples of 6.79 ng/ml by MC/MS/MS, 6.96 ng by EMIT using the 1 H6 monoclonal antibody, and 7.93 ng/ml by Abbott IMx. Two samples tested above range by IMx only. Bland-Altman analysis of the 70 results are shown in Figure 15 for EMIT using monoclonal antibody 1H6 and Abbott IMx versus LC/MS/MS. These plots illustrate that in general noticeable variability of results versus LC/MS/MS occurs with samples about 10 ng/ml. The pattern of difference for LC/MS/MS by immunoassay was similar for the EMIT assay using monoclonal antibody 1 H6 and the Abbott IMx assay.

Table 2

Correlation Between Results of Assays for Tacrolimus in Patients with Liver Damage		
Comparison	Slope/Intercept	Correlation
EMIT vs. LC/MS/MS	$y=1.11x-0.66$	0.910
IMx vs. LC/MS/MS	$y=1.47x-2.11$	0.925
EMIT vs. IMx	$y=0.77x+0.87$	0.953

[0107] Even in these patients containing high levels of tacrolimus metabolites, the assay using the monoclonal antibody 1 H6 in an EMIT homogeneous immunoassay showed a high degree of correlation relative to other methods. Therefore, liver dysfunction may not cause a severe impact on this assay using this antibody and the results imply that clinical management of patients by immunoassay can be accomplished similarly by using the EMIT immunoassay with the 1H6 monoclonal antibody or the previously-available Abbott IMx immunoassay for tacrolimus. This shows the clinical capability of the 1H6 monoclonal antibody.

ADVANTAGES OF THE PRESENT INVENTION

[0108] The present invention provides a monoclonal antibody to tacrolimus that minimizes cross-reactivity with tacrolimus metabolites. The monoclonal antibody of the present invention can be used in an improved immunoassay for tacrolimus that correlates well with other immunoassay methods and non-immunological assay methods for tacrolimus. An immunoassay employing a monoclonal antibody is suitable for clinical monitoring of patients receiving tacrolimus, including patients with impaired liver function who are expected to have high levels of tacrolimus metabolites in their serum.

Claims

1. A monoclonal antibody to tacrolimus that is an IgG₁λ monoclonal antibody that has a binding affinity for tacrolimus of about 3.7 x 10⁹ liters/mole, that cross-reacts with 13-demethyl tacrolimus, and that has less than about 8% cross-reactivity to all of the following tacrolimus metabolites; 15-demethyl tacrolimus; 31-demethyl tacrolimus; 13,31-didemethyl tacrolimus; 15,31-didemethyl tacrolimus; and 12-hydroxy tacrolimus.
2. A monoclonal antibody to tacrolimus that is produced by fusion of antibody-producing cells from an antibody-producing non-human mammal immunized with tacrolimus derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus conjugated to a high molecular weight protein with a suitable fusion partner and that monoclonal antibody:
 - (a) competes with the IgG₁λ monoclonal antibody according to claim 1 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody according to claim 1 as measured by competition assays; and

EP 1 206 493 B1

(b) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13,31-didemethyl tacrolimus, 15,31-didemethyl tacrolimus, and 12-hydroxy tacrolimus.

5 3. A single-chain recombinant antibody (sFv) including therein the variable regions of an antibody to tacrolimus that is produced by fusion of antibody-producing cells from an antibody-producing non-human mammal immunized with tacrolimus derivatized with a carboxymethyl oxime moiety at a carbon atom in the non-binding domain of tacrolimus conjugated to a high molecular weight protein with a suitable fusion partner and that antibody:

10 (a) competes with the IgG₁λ monoclonal antibody according to claim 1 at least about 80% as effectively on a molar basis as compared with the IgG₁λ monoclonal antibody according to claim 1 as measured by competition assays; and

(b) has less than about 10% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13,31-didemethyl tacrolimus, 15,31-didemethyl tacrolimus, and 12-hydroxy tacrolimus

15 4. The monoclonal antibody of claim 2, wherein at least some of the constant regions of the antibody are replaced by human constant regions so that the monoclonal antibody is humanized.

20 5. The monoclonal antibody of claim 2, 3 or 4, wherein the antibody competes at least about 90% as effectively on a molar basis as the monoclonal antibody according to claim 1 and has less than about 8% cross-reactivity with each of 15-demethyl tacrolimus, 31-demethyl tacrolimus, 13,31-didemethyl tacrolimus, 15,31-didemethyl tacrolimus and 12-hydroxy tacrolimus.

25 6. A hybridoma producing an IgG₁λ monoclonal antibody to tacrolimus that has a binding affinity for tacrolimus of about 3.7×10^9 liters/mole, that cross-reacts with 13-demethyl tacrolimus, and that has less than about 8% cross-reactivity to all of the following tacrolimus metabolites: 15-demethyl tacrolimus; 31-demethyl tacrolimus, 13,31-didemethyl tacrolimus; 15,31-didemethyl tacrolimus; and 12-hydroxy tacrolimus.

7. A hybridoma producing the monoclonal antibody of claim 2.

30 8. The antibody according to claim 2 or 3, wherein the tacrolimus is derivatized at carbon atom 22.

9. The antibody of claim 2 or 3 wherein the high molecular weight protein is keyhole limpet hemocyanin.

35 10. A conjugate comprising the antibody according to one of claims 1, 2, 3, 4 and 8 directly or indirectly conjugated to a detectable label.

11. The conjugate of claim 10 wherein the detectable label is selected from the group consisting of an enzyme label, a radioactive label, a fluorescent label, a chemiluminescent label, a bioluminescent label, and a particulate label.

40 12. The conjugate of claim 11 wherein the label is an enzyme label.

13. A method of detecting or determining tacrolimus comprising the steps of:

45 (a) providing a sample suspected of containing tacrolimus;

(b) reacting the sample with;

(i) the antibody according to one of claims 1, 2, 3, 4 and 8; and

(ii) optionally, a tacrolimus analogue;

50 wherein one of the antibody or the tacrolimus analogue is labeled with a label producing a detectable signal, and (c) observing or measuring one of:

(i) the signal associated with tacrolimus bound to antibody;

(ii) the signal associated with tacrolimus unbound to antibody; or

55 (iii) the total signal present;

in order to detect or determine the presence or concentration of tacrolimus in the sample.

14. The method of claim 13 wherein the sample is reacted with a tacrolimus analogue labeled with an enzyme label and the total signal present is observed or measured to detect or determine the presence or concentration of tacrolimus in the sample.

5 15. A test kit comprising, packaged in separate containers;

- (a) the antibody according to one of claims 1, 2, 3, 4 and 8, and
- (b) a tacrolimus analogue labeled directly or indirectly with an enzyme label.

10

Patentansprüche

15 1. Monoklonaler Antikörper gegen Tacrolimus, bei dem es sich um einen monoklonalen IgG₁λ-Antikörper mit einer Bindungsaffinität für Tacrolimus von etwa 3,7 x 10⁹ Liter/Mol handelt, der mit 13-Demethyltacrolimus kreuzreagiert und jeweils weniger als etwa 8 % Kreuzreaktivität gegenüber den folgenden Tacrolimusmetaboliten aufweist: 15-Demethyltacrolimus; 31-Demethyltacrolimus; 13,31-Didemethyltacrolimus; 15,31-Didemethyltacrolimus und 12-Hydroxytacrolimus.

20 2. Monoklonaler Antikörper gegen Tacrolimus, der durch Fusion antikörperproduzierender Zellen aus einem mit einer Carboxymethyloxim-Gruppierung an einem Kohlenstoffatom in der nichtbindenden Tacrolimusdomäne, die an ein Protein mit hohem Molekulargewicht konjugiert ist, derivatisiertem Tacrolimus immunisierten, antikörperproduzierenden nichtmenschlichen Säuger mit einem geeigneten Fusionspartner produziert wird, wobei der monoklonale Antikörper:

25 (a) mit dem monoklonalen IgG₁λ-Antikörper gemäß Anspruch 1 mit in bezug auf Molarität mindestens 80 % Effektivität im Vergleich zum monoklonalen IgG₁λ-Antikörper gemäß Anspruch 1, wie mit Wettbewerbstests gemessen, konkurriert und

(b) jeweils weniger als etwa 10 % Kreuzreaktivität mit 15-Demethyltacrolimus, 31-Demethyltacrolimus, 13,31-Didemethyltacrolimus, 15,31-Didemethyltacrolimus und 12-Hydroxytacrolimus aufweist.

30

3. Rekombinanter Einketten-Antikörper (sFv) mit den darin enthaltenen variablen Bereichen eines Antikörpers gegen Tacrolimus, der durch Fusion antikörperproduzierender Zellen aus einem mit einer Carboxymethyloxim-Gruppierung an einem Kohlenstoffatom in der nichtbindenden Tacrolimusdomäne, die an ein Protein mit hohem Molekulargewicht konjugiert ist, derivatisiertem Tacrolimus immunisierten, antikörperproduzierenden nichtmenschlichen Säuger mit einem geeigneten Fusionspartner produziert wird, wobei der monoklonale Antikörper:

35 (a) mit dem monoklonalen IgG₁λ-Antikörper gemäß Anspruch 1 mit in bezug auf Molarität mindestens 80 % Effektivität im Vergleich zum monoklonalen IgG₁λ-Antikörper gemäß Anspruch 1, wie mit Wettbewerbstests gemessen, konkurriert und

40 (b) jeweils weniger als etwa 10 % Kreuzreaktivität mit 15-Demethyltacrolimus, 31-Demethyltacrolimus, 13,31-Didemethyltacrolimus, 15,31-Didemethyltacrolimus und 12-Hydroxytacrolimus aufweist.

4. Monoklonaler Antikörper nach Anspruch 2, wobei wenigstens einige der konstanten Bereiche des Antikörpers durch menschliche konstante Bereiche ersetzt sind, so daß der monoklonale Antikörper humanisiert ist.

45

5. Monoklonaler Antikörper nach Anspruch 2, 3 oder 4, wobei der Antikörper in bezug auf Molarität mindestens etwa 90 % so effektiv wie der monoklonale Antikörper gemäß Anspruch 1 konkurriert und jeweils weniger als etwa 8 % Kreuzreaktivität mit 15-Demethyltacrolimus, 31-Demethyltacrolimus, 13,31-Didemethyltacrolimus, 15,31-Didemethyltacrolimus und 12-Hydroxytacrolimus aufweist.

50

6. Hybridomzelle, die einen monoklonalen IgG₁λ-Antikörper gegen Tacrolimus mit einer Bindungsaffinität für Tacrolimus von etwa 3,7 x 10⁹ Liter/Mol produziert, der mit 13-Demethyltacrolimus kreuzreagiert und jeweils weniger als etwa 8 % Kreuzreaktivität gegenüber den folgenden Tacrolimusmetaboliten aufweist: 15-Demethyltacrolimus; 31-Demethyltacrolimus; 13,31-Didemethyltacrolimus; 15,31-Didemethyltacrolimus und 12-Hydroxytacrolimus.

55

7. Hybridomzelle, die den monoklonalen Antikörper nach Anspruch 2 produziert.

8. Antikörper gemäß Anspruch 2 oder 3, wobei das Tacrolimus an Kohlenstoffatom 22 derivatisiert ist.

EP 1 206 493 B1

9. Antikörper nach Anspruch 2 oder 3, wobei es sich bei dem Protein mit hohem Molekulargewicht um Keyhole Limpet-Hämocyanin handelt.

5 10. Konjugat, umfassend den direkt oder indirekt an eine nachweisbare Markierung konjugierten Antikörper gemäß einem der Ansprüche 1, 2, 3, 4 und 8.

10 11. Konjugat nach Anspruch 10, wobei die nachweisbare Markierung aus der aus einer Enzymmarkierung, einer radioaktiven Markierung, einer Fluoreszenzmarkierung, einer Chemilumineszenzmarkierung, einer Biolumineszenzmarkierung und einer partikulären Markierung bestehenden Gruppe ausgewählt ist.

12. Konjugat nach Anspruch 11, wobei es sich bei der Markierung um eine Enzymmarkierung handelt.

13. Verfahren zum Nachweis oder zur Bestimmung von Tacrolimus, bei dem man in den folgenden Schritten:

15 (a) eine Probe bereitstellt, von der vermutet wird, daß sie Tacrolimus enthält,
(b) die Probe mit

20 i) dem Antikörper gemäß einem der Ansprüche 1, 2, 3, 4 und 8 und
ii) gegebenenfalls einem Tacrolimusanalogon

reagieren läßt, wobei entweder der Antikörper oder das Tacrolimusanalogon mit einer ein nachweisbares Signal produzierenden Markierung markiert ist, und

(c) entweder

25 i) das mit an Antikörper gebundenem Tacrolimus assoziierte Signal,
ii) das mit nicht an Antikörper gebundenem Tacrolimus assoziierte Signal oder
iii) das vorhandene Gesamtsignal beobachtet oder mißt,

um das Vorhandensein oder die Konzentration von Tacrolimus in der Probe nachzuweisen oder zu bestimmen.

30 14. Verfahren nach Anspruch 13, wobei man die Probe mit einem mit einer Enzymmarkierung markierten Tacrolimusanalogon reagieren läßt und das vorhandene Gesamtsignal zum Nachweis oder zur Bestimmung des Vorhandenseins oder der Konzentration von Tacrolimus in der Probe beobachtet oder mißt.

35 15. Testkit, umfassend in in getrennten Behältern abgepackter Form:

(a) den Antikörper gemäß einem der Ansprüche 1, 2, 3, 4 und 8 sowie
(b) ein direkt oder indirekt mit einer Enzymmarkierung markiertes Tacrolimusanalogon.

40 Revendications

45 1. Anticorps monoclonal dirigé contre le tacrolimus qui est un anticorps monoclonal IgG₁λ qui a une affinité de liaison pour le tacrolimus d'environ 3,7 x 10⁹ l/mole, qui réagit de manière croisée avec le 13-déméthyl tacrolimus, et qui a moins d'environ 8% de réactivité croisée avec tous les métabolites de tacrolimus suivants : le 15-déméthyl tacrolimus, le 31-déméthyl tacrolimus, le 13,31-didéméthyl tacrolimus, le 15,31-didéméthyl tacrolimus, et le 12-hydroxy tacrolimus.

50 2. Anticorps monoclonal dirigé contre le tacrolimus qui est produit par fusion de cellules productrices d'anticorps, à partir d'un mammifère non-humain producteur d'anticorps immunisé par un produit de dérivation du tacrolimus présentant un reste carboxyméthyl oxime au niveau d'un atome de carbone dans le domaine non-liant du tacrolimus conjugué à une protéine de haut poids moléculaire, avec un partenaire de fusion approprié et ledit anticorps monoclonal :

55 (a) agit par compétition avec l'anticorps monoclonal IgG₁λ selon la revendication 1 avec une efficacité d'au moins environ 80% sur une base molaire comparée à celle de l'anticorps monoclonal selon la revendication 1 telle que mesurée par des tests de compétition ; et

(b) a moins d'environ 10% de réactivité croisée avec l'un parmi le 15-déméthyl tacrolimus, le 31-déméthyl

EP 1 206 493 B1

tacrolimus, le 13,31-didéméthyl tacrolimus, le 15,31-didéméthyl tacrolimus, et le 12-hydroxy tacrolimus.

- 5
3. Anticorps recombinant à chaîne unique (sFv) incluant les régions variables d'un anticorps dirigé contre le tacrolimus qui est produit par fusion de cellules productrices d'anticorps à partir d'un mammifère non-humain producteur d'anticorps immunisé par un produit de dérivation du tacrolimus présentant un reste carboxyméthyl oxime au niveau d'un atome de carbone dans le domaine non-liant du tacrolimus conjugué à une protéine de haut poids moléculaire, avec un partenaire de fusion approprié et ledit anticorps monoclonal :
- 10
- (a) agit par compétition avec l'anticorps monoclonal IgG₁λ selon la revendication 1 avec une efficacité d'au moins environ 80% sur une base molaire comparée à celle de l'anticorps monoclonal selon la revendication 1 telle que mesurée par des tests de compétition ; et
- (b) a moins d'environ 10% de réactivité croisée avec l'un parmi le 15-déméthyl tacrolimus, le 31-déméthyl tacrolimus, le 13,31-didéméthyl tacrolimus, le 15,31-didéméthyl tacrolimus, et le 12-hydroxy tacrolimus.
- 15
4. Anticorps monoclonal selon la revendication 2, **caractérisé en ce qu'**au moins certaines des régions constantes de l'anticorps sont remplacées par des régions constantes humaines de sorte que l'anticorps monoclonal est humanisé.
- 20
5. Anticorps monoclonal selon la revendication 2, 3 ou 4, **caractérisé en ce que** l'anticorps agit par compétition avec une efficacité d'au moins environ 90% sur une base molaire comparée à celle de l'anticorps monoclonal selon la revendication 1 et a moins d'environ 8% de réactivité croisée avec l'un parmi le 15-déméthyl tacrolimus, le 31-déméthyl tacrolimus, le 13,31-didéméthyl tacrolimus, le 15,31-didéméthyl tacrolimus, et le 12-hydroxy tacrolimus.
- 25
6. Hybridome produisant un anticorps monoclonal IgG₁λ dirigé contre le tacrolimus qui a une affinité de liaison pour le tacrolimus d'environ 3,7 x 10⁹ l/mole, qui réagit de manière croisée avec le 13-déméthyl tacrolimus, et qui a moins d'environ 8% de réactivité croisée avec tous les métabolites du tacrolimus suivants : le 15-déméthyl tacrolimus, le 31-déméthyl tacrolimus, le 13,31-didéméthyl tacrolimus, le 15,31-didéméthyl tacrolimus, et le 12-hydroxy tacrolimus.
- 30
7. Hybridome produisant l'anticorps monoclonal selon la revendication 2.
- 35
8. Anticorps selon la revendication 2 ou 3, **caractérisé en ce que** la dérivation du tacrolimus est réalisée au niveau de l'atome de carbone 22.
9. Anticorps selon la revendication 2 ou 3, **caractérisé en ce que** la protéine de haut poids moléculaire est l'hémocyanine de patelle.
- 40
10. Conjugué comprenant l'anticorps selon l'une des revendications 1, 2, 3, 4 et 8, directement ou indirectement conjugué à un marqueur détectable.
- 45
11. Conjugué selon la revendication 10, **caractérisé en ce que** le marqueur détectable est choisi dans le groupe constitué par un marqueur enzymatique, un marqueur radioactif, un marqueur fluorescent, un marqueur chimioluminescent, un marqueur bioluminescent, et un marqueur particulaire.
12. Conjugué selon la revendication 11, **caractérisé en ce que** le marqueur est un marqueur enzymatique.
- 50
13. Méthode de détection ou de dosage du tacrolimus comprenant les étapes suivantes consistant à :
- (a) fournir un échantillon censé contenir le tacrolimus ;
- (b) faire réagir l'échantillon avec:
- 55
- (i) l'anticorps selon l'une des revendications 1, 2, 3, 4 et 8 ; et
- (ii) éventuellement, un analogue du tacrolimus ;
- caractérisée en ce que** l'un parmi l'anticorps ou l'analogue de tacrolimus est marqué par un marqueur produisant un signal détectable, et
- (c) observer ou mesurer l'un parmi :
- (i) le signal associé au tacrolimus lié à l'anticorps

EP 1 206 493 B1

- (ii) le signal associé au tacrolimus non lié à l'anticorps ; ou
- (iii) le signal total présent ;

afin de détecter la présence ou mesurer la concentration de tacrolimus dans l'échantillon.

- 5
- 14.** Méthode selon la revendication 13, **caractérisée en ce que** l'échantillon est mis à réagir avec un analogue du tacrolimus marqué par un marqueur enzymatique et **en ce que** le signal total présent est noté ou mesuré pour détecter la présence ou déterminer la concentration de tacrolimus dans l'échantillon.
- 10
- 15.** Kit d'analyse comprenant, conditionnés dans des conteneurs séparés :
- (a) l'anticorps selon l'une des revendications 1, 2, 3, 4 et 8, et
 - (b) un analogue du tacrolimus marqué directement ou indirectement par un marqueur enzymatique.

15

20

25

30

35

40

45

50

55

FIG. 1




FIG. 2




FIG. 3





FIG. 4






FIG. 6




FIG. 8





FIG. 9






FIG. 10

FIG. 11





FIG. 15






FIG. 12

FIG. 13





FIG. 14




FIG. 16

专利名称(译)	他克莫司的单克隆抗体和他克莫司的免疫测定方法		
公开(公告)号	EP1206493B1	公开(公告)日	2006-11-15
申请号	EP2000953792	申请日	2000-08-02
申请(专利权)人(译)	德灵INC.		
当前申请(专利权)人(译)	德灵INC.		
[标]发明人	KASPER KENNETH C JEONG HENRY DAVALIAN DARIUSH LIU HSHIOU TING MILLER PAUL L WILLIAMS DENISE L		
发明人	KASPER, KENNETH, C. JEONG, HENRY DAVALIAN, DARIUSH LIU, HSHIOU-TING MILLER, PAUL, L. WILLIAMS, DENISE, L.		
IPC分类号	C07K16/00 G01N33/53 A61K51/10 C07D498/18 C07K16/12 C07K16/44 C12N5/10 C12P21/08		
CPC分类号	A61K51/1093 C07K16/1292 Y10S436/815		
代理机构(译)	施韦泽KLAUS		
优先权	09/368010 1999-08-03 US		
其他公开文献	EP1206493A2		
外部链接	Espacenet		

摘要(译)

针对免疫抑制药物他克莫司的IgG1 λ 单克隆抗体具有改善的性质。特别是，这种命名为1H6的单克隆抗体与几种他克莫司代谢物的交叉反应性降低。该抗体适用于进行免疫测定，例如均相免疫测定，以检测或确定样品例如血液样品中他克莫司的存在或浓度。本发明进一步包括在分子的非结合部分衍生的他克莫司衍生物，其可用于免疫产生抗体的动物和产生这种单克隆抗体，以及在上述测定中用作他克莫司类似物的他克莫司的标记衍生物。本发明还包括用于检测他克莫司的免疫测定方法和用于进行这种免疫测定的测试试剂盒。

