



(11) **EP 1 348 029 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
07.03.2012 Bulletin 2012/10

(21) Application number: **02734847.3**

(22) Date of filing: **02.01.2002**

(51) Int Cl.:
C12N 15/11 (2006.01)

(86) International application number:
PCT/GB2002/000005

(87) International publication number:
WO 2002/059150 (01.08.2002 Gazette 2002/31)

(54) **MATERIALS AND METHODS RELATING TO PROTEIN AGGREGATION IN
NEURODEGENERATIVE DISEASE**

MATERIAL UND METHODEN, INVOLVIERT IN DIE PROTEIN-AGGREGATION
NEURODEGENERATIVER KRANKHEITEN

MOYENS ET PROCEDES ASSOCIES A L'AGREGATION DE PROTEINES DANS UNE MALADIE
NEURODEGENERATIVE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**

(30) Priority: **03.01.2001 GB 0100119**

(43) Date of publication of application:
01.10.2003 Bulletin 2003/40

(73) Proprietor: **WisTa Laboratories Ltd.**
Singapore 139948 (SG)

(72) Inventors:

- **WISCHIK, Claude Michel,**
University of Aberdeen
Foresterhill,
Aberdeen, Aberdeenshire
AB25 2ZD (GB)
- **RICKARD, Janet Elizabeth,**
University of Aberdeen
Foresterhill,
Aberdeen, Aberdeenshire
AB25 2ZD (GB)
- **HORSLEY, David,**
University of Aberdeen
Foresterhill,
Aberdeen, Aberdeenshire
AB25 2ZD (GB)
- **HARRINGTON, Charles Robert,**
University of Aberdeen
Foresterhill,
Aberdeen, Aberdeenshire
AB25 2ZD (GB)

- **THEURING, Franz,**
University Hospital Charite
Dorotheenstr. 94,
10117 Berlin (DE)
- **STAMER, Karsten,**
University Hospital Charite
Dorotheenstr. 94,
10117 Berlin (DE)
- **ZABKE, Claudia,**
WisTa Laboratories Ltd.
Forresterhill, Aberdeen AB25 2ZD
Scotland (GB)

(74) Representative: **Kremer, Simon Mark et al**
Mewburn Ellis LLP
33 Gutter Lane
London
EC2V 8AS (GB)

(56) References cited:
WO-A-01/53340 WO-A-01/95709
WO-A-96/30766 WO-A-99/62548

Remarks:

The file contains technical information submitted after
the application was filed and not included in this
specification

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

Technical field

[0001] The present invention concerns models, materials and methods relating to the aggregation of proteins associated with neurodegenerative disease.

Background art

[0002] Conditions of dementia such as Alzheimer's disease (AD) are frequently characterised by a progressive accumulation of intracellular and/or extracellular deposits of proteinaceous structures such as β -amyloid plaques and neurofibrillary tangles in the brains of affected patients. The appearance of these lesions largely correlates with pathological neurofibrillary degeneration and brain atrophy, as well as with cognitive impairment (Mukaetova-Ladinska, E.B. et al. (2000) Am. J. Pathol. Vol. 157, No. 2, 623-636).

[0003] Both neuritic plaques and neurofibrillary tangles contain paired helical filaments (PHFs), of which a major constituent is the microtubule-associated protein tau (Wischnik et al. (1988) PNAS USA 85, 4506). Plaques also contain extracellular β -amyloid fibrils derived from the abnormal processing of amyloid precursor protein (APP; Kang et al. (1987) Nature 325, 733). An article by Wischnik et al. (in "Neurobiology of Alzheimer's Disease", 2nd Edition (2000) Eds. Dawbarn, D. and Allen, S.J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford) discusses in detail the putative role of tau protein in the pathogenesis of neurodegenerative dementias.

[0004] Studies of Alzheimer's disease indicate that the loss of the normal form of tau (Mukaetova-Ladinska et al. (1993) Am. J. Pathol., 143, 565; Wischnik et al. (1995a) Neurobiol. Ageing, 16: 409; Lai et al. (1995b) Neurobiol. Ageing, 16: 433), accumulation of pathological PHFs (Mukaetova-Ladinska et al. (1993), *loc. cit.*; Harrington et al. (1994a) Dementia, 5, 215; Harrington et al. (1994b) Am. J. Pathol., 145, 1472; Wischnik et al., (1995a), *loc. cit.*) and loss of synapses in the mid-frontal cortex (Terry et al. (1991) Ann. Neurol., 30, 572) correlate with associated cognitive impairment. Furthermore, loss of synapses (Terry et al., *loc. cit.*) and loss of pyramidal cells (Bondareff et al. (1993) Arch. Gen. Psychiatry, 50: 350) both correlate with morphometric measures of tau-reactive neurofibrillary pathology, which parallels, at a molecular level, an almost total redistribution of the tau protein pool from a soluble to a polymerised form (PHFs) in Alzheimer's disease (Mukaetova-Ladinska et al. (1993), *loc. cit.*; Lai et al. (1995), *loc. cit.*).

[0005] Tau exists in alternatively-spliced isoforms, which contain three or four copies of a repeat sequence corresponding to the microtubule-binding domain (Goedert, M., et al. (1989) EMBO J. 8, 393-399; Goedert, M., et al. (1989) Neuron 3, 519-526). Tau in PHFs is proteolytically processed to a core domain (Wischnik, C.M., et

al. (1988) Proc. Natl. Acad. Sci. USA 85, 4884-4888; Wischnik et al. PNAS USA 1988, 85:4506-4510); Novak, M., et al. (1993) EMBO J. 12, 365-370) which is composed of a phase-shifted version of the repeat domain; only three repeats are involved in the stable tau-tau interaction (Jakes, R., et al. (1991) EMBO J. 10, 2725-2729). Once formed, PHF-like tau aggregates act as seeds for the further capture and provide a template for proteolytic processing of full-length tau protein (Wischnik et al. 1996 Proc Natl Acad Sci USA 93, 11213-11218).

[0006] In the course of their formation and accumulation, paired helical filaments (PHFs) first assemble to form amorphous aggregates within the cytoplasm, probably from early tau oligomers which become truncated prior to, or in the course of, PHF assembly (Mena, R., et al. (1995) Acta Neuropathol. 89, 50-56; Mena, R., et al. (1996) Acta Neuropathol. 91, 633-641). These filaments then go on to form classical intracellular neurofibrillary tangles. In this state, the PHFs consist of a core of truncated tau and a fuzzy outer coat containing full-length tau (Wischnik, C.M., et al. (1996b) in "Microtubule-associated proteins: modifications in disease", eds. Avila, J., Brandt, R. and Kosik, K. S. (Harwood Academic Publishers, Amsterdam) pp.185-241)). The assembly process is exponential, consuming the cellular pool of normal functional tau and inducing new tau synthesis to make up the deficit (Lai, R. Y. K., et al., (1995), Neurobiology of Ageing, Vol. 16, No. 3, 433-445). Eventually, functional impairment of the neurone progresses to the point of cell death, leaving behind an extracellular tangle. Cell death is highly correlated with the number of extracellular tangles (Wischnik et al. 2000, *loc. cit.*). As tangles are extruded into the extracellular space, there is progressive loss of the fuzzy outer coat of the neurone with corresponding loss of N-terminal tau immunoreactivity, but preservation of tau immunoreactivity associated with the PHF core (Figure 4a; also Bondareff, W. et al., (1994) J. Neuropath. Exper. Neurol., Vol. 53, No. 2, 158-164).

[0007] The phase shift which is observed in the repeat domain of tau incorporated into PHFs suggests that the repeat domain undergoes an induced conformational change during incorporation into the filament. During the onset of Alzheimer's disease, it is envisaged that this conformational change could be initiated by the binding of tau to a pathological substrate, such as damaged or mutated membrane proteins (see Figure 19a - also Wischnik, C.M., et al. (1997) in "Microtubule-associated proteins: modifications in disease", eds. Avila, J., Brandt, R. and Kosik, K. S. (Harwood Academic Publishers, Amsterdam) pp.185-241).

[0008] In the case of Alzheimer's disease, current pharmaceutical therapies are focused on symptomatic treatment of the loss of cholinergic transmission which results from neurodegeneration (Mayeux, R., et al. (1999) New Eng. J. Med. 341, 1670-1679). However, although the available treatments delay progression of the disease for up to six to eight months, they do not prevent it. The discovery of drugs that could prevent the aggre-

gation of tau which leads to neurodegeneration would provide a more effective strategy for prophylaxis or for inhibiting the progression of the disease, which would not require an immediate knowledge of the diverse upstream events that initiate the aggregation (Figure 19b).

Protein aggregation assays

[0009] Based on the putative model described above, WO 96/30766 describes an *in vitro* assay for tau aggregation in which a fragment of tau corresponding to the core repeat domain, which has been adsorbed to a solid phase substrate, is able to capture soluble full-length tau and bind tau with high affinity. This association confers stability against proteolytic digestion of the aggregated tau molecules. The process is self-propagating, and can be blocked selectively by prototype pharmaceutical agents (Wischik, C.M., *et al.* (1996), *loc. cit.*).

[0010] Although the *in vitro* assay described in WO 96/30766 enables the identification of inhibitors or modulators of tau-tau association, the present inventors have also recognized that cell-based models of Alzheimer's disease-like protein aggregation would be useful. Such cellular models could be used both in the primary screening of candidate modulators of tau-tau aggregation, and in the secondary screening of compounds already identified in the *in vitro* assay of WO 96/30766. Furthermore, the demonstration of tau aggregation in cells could also aid in the identification of normal cellular substrates which are involved in the initiation of pathological tau aggregation, which substrates could themselves be targets for pharmaceutical intervention.

[0011] However, numerous papers reporting the expression of various tau constructs in tissue culture models have failed to demonstrate aggregation (see e.g. Baum, L. *et al.*, (1995) *Mol. Brain Res.* 34:1-17).

[0012] These problems can be understood when it is considered that aggregation of soluble tau *in vitro* has so far only been achieved under non-physiological conditions and at high concentrations (reviewed in Wischik (2000), *loc. cit.*).

[0013] WO 96/30766 also describes two approaches for studying tau aggregation in a cellular environment. In the first approach, full-length tau or fragments of tau were stably expressed in cells. In the second approach, aggregated tau was transiently transfected into cells by use of lipofectin.

[0014] Although both of these approaches are useful for the study of tau-tau aggregation, they have some limitations. Transfection of aggregated tau into cells using lipofection is of variable efficiency, as is the production of aggregated tau itself. Moreover, the core tau fragment, which is the most efficient seed for tau aggregation, is found to be toxic when stably expressed in cells, leading to low expression levels. Thus, constitutive expression of the truncated tau fragment of the PHF core in eukaryotic cells is difficult to achieve. Transient expression systems permit the optimization of expression of tau, but the

inherent toxicity of the fragments renders even these systems unreliable. Longer fragments of tau are less toxic, but these do not reliably aggregate when expressed in cells.

[0015] Thus it would be desirable for an alternative model system to be developed, in which the interaction between e.g. tau molecules and the like could be investigated under physiological conditions, in a stable and controllable cell line, and which could be used to screen for potential diagnostic, prognostic or therapeutic agents of conditions such as Alzheimer's disease.

Animal models

[0016] Although *in vitro* and cell-based assays can be useful tools, animal models of Alzheimer's disease and related conditions can help to understand the relationship between the biochemical and pathological changes in the brain and impairment of memory and behavior. Furthermore, they enable the pathogenesis of the disease process to be examined *in vivo* and provide a model in which therapeutic strategies can be tested.

[0017] Specifically, animal models put cell-based systems in a pharmacokinetic context where, for example, potential therapeutics can be assessed not just in terms of particular biochemical activities against protein aggregation, but also in terms of toxicity, delivery and half-life at the site of action e.g. through the blood brain barrier.

[0018] Notwithstanding this, existing models fail to display the combination of tangles, plaques and cognitive impairment characteristic of AD. Aged dogs and non-human primates develop β -amyloidosis, but tau pathology is not a feature in these animals (Walker, 1997). Similarly, transgenic mice modeling amyloidosis have been created, but these fail to exhibit abnormal deposition of tau (Janus, Chishti, & Westaway, 2000). Conversely animal models of tau pathology fail to demonstrate amyloid pathology. The tangles that accumulate in the brains of rodents treated with aluminium differ in their ultrastructure from those found in AD (see above). Filamentous cytoskeletal changes associated with abnormally phosphorylated tau have been observed in aged baboons, bears, sheep and goats (Braak, Braak, & Strothjohann, 1994; Cork *et al.*, 1988; Nelson, Greenberg, & Saper, 1994; Nelson & Saper, 1995; Roertgen *et al.*, 1996; Schultz *et al.*, 2000).

[0019] Some experiments have implicated links between tau and amyloid in animal models. Hyperphosphorylated tau accumulates in the somatodendritic compartment of neurons in rat brains following chronic intraventricular infusion of okadaic acid, an inhibitor of protein phosphatase 2A (Arendt, Holzer, Fruth, Brijckner, & Gartner, 1995). Furthermore, the okadaic acid treatment also led to the formation of extracellular deposits of A β and memory impairment. Two other studies have implicated A β in the accumulation of tau in animal models. Microinjections of fibrillar A β in the cortex of aged rhesus monkeys caused the focal accumulations of intracellular

phosphorylated tau (Geula et al., 1998). This was dependent upon both age and species; the same result was not observed for rats or young rhesus monkeys. Secondly, focal deposits of tau have been observed in mice that were transgenic for APP carrying AD associated mutations (Sturchler-Pierrat et al., 1997).

[0020] The brains of tau-deficient mice appear immunohistochemically normal and axonal elongation was not affected in cultured neurons (Harada et al., 1994). However, microtubule stability was decreased and its organization altered in some small-calibre axons. Furthermore, an increase in microtubule-associated protein 1A (MAP 1A), which might compensate for a functional loss of tau in large-calibre axons, was found. Thus, tau seems to be crucial in the stabilization and organization of axonal microtubules in certain axons. Subsequent studies have demonstrated that tau-deficient mice exhibit signs similar to certain symptoms characteristic of frontotemporal dementia patients, i.e. personality changes (disinhibition/aggression) and deterioration of memory and executive function. The mice showed muscle weakness and impaired balance control, hyperactivity in a novel environment, and impairment in contextual fear conditioning (Ikegami, Harada, & Hirokawa, 2000). Spatial learning tasks, however, were unaffected in the mice as is the memory function in FTDP-17 patients.

[0021] Although neurofibrillary tangles and phenotypic alterations were not reported in animals transgenic for 3- or 4-repeat tau isoforms (Brion, Tremp, & Octave, 1999; Gotz et al., 1995), these findings may reflect low levels of protein expression.

[0022] Impaired motor function, in the presence or absence of tauopathy, has been observed in mice expressing higher levels of human tau protein (Ishihara et al., 1999; Spittaels et al., 1999). In these animals the filamentous inclusions did not exhibit the ultrastructural features of AD PHFs.

[0023] A recent transgenic mouse model expressing human tau with the P301L mutation develops neurofibrillary tangles, neuronal loss and motor dysfunction (Lewis et al., 2000). The tau inclusions in these mice show both straight and twisted ribbon filaments similar to those found in human patients. Attempts to combine this model with amyloidosis are underway by crossing these mice with those transgenic for APP. Other transgenic mouse models are discussed in US patents 5,912,410 and 5,898,094

[0024] Thus it can be seen that existing animal models fail to demonstrate any pathological evidence that the transgene leads to the accumulation of truncated tau protein encompassing the PHF domain. Although filaments are observed in tau carrying the P301L mutation, this corresponds to a form of tau that causes FTDP-17 and not AD. To date, no mutation in the tau gene has been identified which causes AD, and therefore approaches based purely on mutated protein for effect may be of limited relevance.

[0025] More particularly, there has been no clear dis-

closure of unmutated tau aggregating in these models, as evidenced by proteolytic processing to a truncated core fragment corresponding to that found in the actual AD PHF core. A system in which such proteolytic processing occurred in a pharmacokinetic context would thus provide a contribution to the art.

Disclosure of the invention

[0026] As discussed above, a conformational change of tau, rather than a chemical modification, is responsible for the progressive aggregation of tau molecules and their subsequent polymerization to form PHFs.

[0027] Surprisingly, the present inventors have now found that it is possible to reproduce this conformational change in a model system by causing the synthesis of a membrane-localised form of the protein of interest, e.g. tau, or fragments thereof containing the core aggregation domain.

[0028] It has been found, unexpectedly, that the association of e.g. tau or tau fragments with a cellular or organellar membrane, e.g. the plasma membrane of a cell or the membrane of the endoplasmic reticulum, can also seed the aggregation of further molecules of tau or tau fragments, thus propagating the formation of tau-tau aggregates in a manner somewhat analogous to that observed in the solid-phase assay system of WO 96/30766. This technique appears to have wide applicability to proteins involved in protein-aggregation type diseases.

[0029] Although not wishing to be limited by any particular mechanism, it is proposed by the inventors that the membrane-localised synthesis of the protein causes the high-affinity capture site of this protein to become exposed, by physically inducing the conformational change of tau that occurs upon its binding to a suitable substrate. The effect of this conformational change may be to create a sufficiently high local concentration of tau in its seeding conformation that further aggregation of the protein is promoted. These observations may have implications for all diseases in which protein aggregation plays a role.

[0030] In various aspects, the present invention provides, methods of modeling disease-like intracellular aggregation of an appropriate protein, or a core fragment thereof, the method being characterised by the step of synthesising a membrane-localised form of a protein which can initiate such aggregation. Such methods can be used, for instance, to identify modulators of the aggregation. The methods are performed in vitro, or in a cultured cell or cell line, or in non-human animals.

[0031] Thus, in aspects of the invention there are provided fusion polypeptides comprising: (i) an aggregating portion, which is derived from a protein associated with a disease in which the protein aggregates or initiates aggregation of protein through an induced conformational polymerisation interaction, (ii) a heterologous membrane-localising portion.

[0032] In this way, a protein such as tau, or a core

fragment of tau, or a corresponding protein as discussed below, can be synthesised in the form of a membrane-localized protein, where it can act as a seed for the intracellular aggregation and proteolytic processing of the fusion at an appropriate level for modeling aggregation of the actual protein from which it was derived. It may also seed aggregation with or between further molecules (e.g. full-length tau molecules), thus propagating the formation of protein aggregates.

[0033] Cells which express such membrane-associated protein aggregates can be used as models of disease-related protein aggregation, such as that observed in the cells of Alzheimer's-like diseases *in vivo*. This model system can be used in further studies, such as in screening assays for potential therapeutic, prognostic or diagnostic agents, as described in detail below.

[0034] Thus one aspect of the present invention there is disclosed a method of inducing, modeling or controlling aggregation of a protein associated with a disease in which the protein undergoes an induced conformational interaction, the method comprising the steps of:

(a) introducing to a membrane, a fusion polypeptide comprising: (i) an aggregation portion, which is derived from a first protein associated with a disease in which the protein undergoes an induced conformational polymerisation interaction, (ii) a heterologous membrane-localising portion, (b) causing or permitting an induced conformational polymerisation interaction between the fusion polypeptide and a further polypeptide, such as to cause aggregation of the said further polypeptide. Optionally the extent of aggregation may be monitored.

[0035] In this and other embodiments, the further polypeptide may, for example, be a further fusion protein, or may result from proteolytic degradation of the fusion polypeptide such as to yield a second protein capable of undergoing an induced conformational polymerisation interaction with said first protein.

[0036] Also disclosed, herein is a method of inducing, modeling or controlling aggregation of a (second) protein associated with a disease in which the protein undergoes an induced conformational interaction, the method comprising the steps of:

(a) introducing to a membrane, a fusion polypeptide comprising: (i) an aggregation portion, which is derived from a first protein associated with a disease in which the protein undergoes or initiates an induced conformational polymerisation interaction, (ii) a heterologous membrane-localising portion, (b) contacting the fusion polypeptide with a second protein capable of undergoing an induced conformational polymerisation interaction with said first protein, whereby interaction of the second protein with the membrane localized fusion protein causes aggregation of the second protein. Optionally the extent

of aggregation may be monitored.

[0037] Examples are disclosed below of first and second proteins, which may be derived from the same or different protein associated with a disease. In particular, the first protein will be one which (as a result of an induced conformational change) initiates and/or participates in the aggregation of further protein which is the same or different, and the second will be one which actually aggregates in the relevant disease.

[0038] In preferred embodiments there are disclosed cell-based method of inducing intracellular aggregation of a protein associated with a disease in which the protein undergoes an induced conformational interaction e.g. comprising the steps of: (a) introducing into a cell, such that it becomes localised at a membrane in the cell, a fusion polypeptide comprising: (i) an aggregation portion, which is derived from a first protein associated with a disease in which the protein undergoes an induced conformational polymerisation interaction, (ii) a heterologous membrane-localising portion, (b) introducing into the cell, a second protein capable of undergoing an induced conformational polymerisation interaction with said first protein, whereby interaction with of the second protein with the localized fusion protein causes intracellular aggregation and optionally proteolytic processing of either.

[0039] Again, optionally the extent of aggregation and/or proteolytic processing of the second protein may be monitored. This may be in the presence of one or more agents suspected of being capable of modulating (e.g. inhibiting or reversing) the aggregation.

[0040] Generally, speaking the method will be carried out by use of a nucleic acid encoding the fusion protein and/or the second protein.

[0041] Some preferred aspects and embodiments of the present invention will now be described in more detail.

Diseases and proteins

[0042] As stated above, the invention may employ any protein which is associated with a disease in which the protein undergoes an induced conformational polymerisation interaction i.e one in which a conformational change of the protein seeds the binding and aggregation of further protein molecules in a self-propagating manner. Once nucleation is initiated, an aggregation cascade may ensue which involves the induced conformational polymerisation of further protein molecules, which conformational change may render the aggregates more resistant to further proteolysis. The protein aggregates thus formed are thought to be a proximal cause of neurodegeneration, clinical dementia, and other pathological symptoms of this group of diseases. Purely for brevity, this group of conditions may be referred to herein as "Alzheimer's disease-like" conditions.

[0043] Preferred embodiments of the invention are based on tau protein. Where used herein, the term "tau protein" refers generally to any protein of the tau protein

family. Tau proteins are characterised as being one among a larger number of protein families which co-purify with microtubules during repeated cycles of assembly and disassembly (Shelanski et al. (1973) Proc. Natl. Acad. Sci. USA, 70., 765-768), and are known as microtubule-associated-proteins (MAPs). Members of the tau family share the common features of having a characteristic N-terminal segment, sequences of approximately 50 amino acids inserted in the N-terminal segment, which are developmentally regulated in the brain, a characteristic tandem repeat region consisting of 3 or 4 tandem repeats of 31-32 amino acids, and a C-terminal tail. The term is thus exemplified by, but not limited to, tau molecules which form PHFs (and, ultimately, neurofibrillary tangles) in Alzheimer's disease brains. Indeed, PHFs accumulate in the somatodendritic compartment, where the predominant microtubule-associated protein is MAP2 (Matus, A., in "Microtubules" [Hyams and Lloyd, eds.] pp 155-166, John Wiley and Sons, NY). MAP2 isoforms are almost identical to tau protein in the tandem repeat region, but differ substantially both in the sequence and extent of the N-terminal domain (Figures 7 and 8; Kindler and Garner (1994) Mol. Brain Res. 26, 218-224). Nevertheless, aggregation in the tandem-repeat region is not selective for the specific tau core amino acid sequence. Thus it will be appreciated that any discussion herein in relation to tau-tau aggregation should be taken as relating also to tau-MAP2 aggregation and MAP2-MAP2 aggregation.

[0044] Figure 20 shows a Table listing various other disease-associated aggregating proteins which may be used in the present invention. In each case the disease or diseases in which the initiation of aggregation and/or mutation of the protein(s) may play a role is also listed. The domain or mutation responsible for the disease activity is listed, and at least all or part of this minimal portion of the protein would preferably be encompassed by the protein used in the invention as the membrane-targeted 'capture' or 'seeding' protein.

[0045] As can be seen from the table, example diseases which are characterised by pathological protein aggregation include motor neurone disease and Lewy body disease. Furthermore, the pathogenesis of neurodegenerative disorders such as Pick's disease and Progressive Supranuclear Palsy appears to correlate with an accumulation of pathological truncated tau aggregates in the dentate gyrus and stellate pyramidal cells of the neocortex, respectively (Wischnik *et al.* 2000, *loc. cit.*).

[0046] Figure 21 describes proteins which may be involved in the initiation stages of diseases of the type discussed herein. For example, in Wischnik, C.M., *et al.* (1997) [supra] a variety of membrane proteins which may be involved in the initiation of tau capture and amyloid protein precursor in AD are discussed. These include presenilins and mitochondrial proteins such as core protein 2 of the *bc₁* enzyme complex (CP2), ATP-synthase subunit 9 (ATP S-9) and porin.

[0047] Thus it will be appreciated, in the light of the

above discussion, (and except where context requires otherwise) where the embodiments of the invention are described with respect to tau protein or tau-like proteins (e.g. MAP2) the description should be taken as applying equally to the other proteins discussed above (e.g. β -amyloid, synuclein, prion, appropriate mitochondrial protein etc.) or other proteins which may undergo a similar pathological aggregation by virtue of conformational change in a domain critical for propagation of the aggregation, or which imparts proteolytic stability to the aggregate this formed (article by Wischnik *et al.* (in "Neurobiology of Alzheimer's Disease", 2nd Edition (2000) Eds. Dawbarn, D. and Allen, S.J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford). All such proteins may be referred to herein as "aggregating disease proteins."

[0048] Likewise, where mention is made herein of "tau-tau aggregation", or the like, this may also be taken to be applicable to other "aggregating-protein aggregation", such as β -amyloid aggregation, prion aggregation and synuclein aggregation etc. Likewise "tau proteolytic degradation" and so on.

[0049] Proteins for use in the methods described above may, where appropriate (e.g. in the study of tau-tau binding) be isolated from brain tissues by methods known to those skilled in the art, for instance as described in the respective references. Soluble tau can be isolated either from human brain tissues, or from animal brain tissues, with a postmortem delay of less than 24 hours (see Wischnik *et al.* (1988); (1995a), *loc. cit.*). Microtubule proteins can be obtained by three cycles of temperature-dependent assembly-disassembly according to Shelanski *et al.* (1973, *loc. cit.*). Lipofection of the proteins may be carried out analogously to the methods described in W096/30766. More preferably the production of e.g. tau proteins and fragments thereof will be achieved by conventional recombinant DNA technology as described below.

40 Fragments, derivatives and controls

[0050] There is no requirement that the invention employ full-length protein; indeed this may be undesirable in certain contexts. Equally, the protein used may include an extended N- or C-terminus.

[0051] As shown in the Examples hereinafter, the nature of the aggregating portion of the fusion may be such that appropriate levels of aggregation (and possible proteolytic processing) are achieved even in the absence of further molecules.

[0052] Alternatively or additionally, further molecules (e.g. full length disease protein molecules) may be provided which participate in the aggregation.

[0053] In preferred embodiments of the present invention, the fusion polypeptide, as synthesized in either the cell-based method or the *in vitro* method described herein, comprises or consists essentially of a truncated fragment. Such a fragment will generally be a "core fragment"

which term refers to that part of the protein that is able to bind to further protein to initiate or propagate aggregation. In the case of disease proteins which aggregate, such core fragments are also likely to be those which contribute to the proteolytic stability of the aggregate.

[0054] Thus, for example, a "tau core fragment" is a tau fragment comprising a truncated tau protein sequence derived from the tandem repeat region ("core tandem repeat domain") and, which, in the appropriate conditions, is capable of binding to the tandem repeat region of a further tau protein or a MAP2 protein with high affinity.

[0055] Where the fusion polypeptide is derived from tau, the truncated fragment may preferably comprise a fragment from about 186-296 extending to about 390-441 of the full-length protein, more preferably any of the following portions: 186 to 390 or 391 or 441; 296 to 390 or 391 or 441 (see Figures 5 and 10). In the Examples hereinafter, the ligation of the tau fragment to the signal sequence destroyed amino acids 186-189, so that the starting amino acid was equivalent to 190.

[0056] In the case of APP (amyloid precursor protein), for instance, expression of a fragment of the APP that encompasses the A β domain of 1-40 or 1-42 amino acids as a fusion protein, may be preferred.

[0057] Other core fragments may be based e.g. on the domains discussed with reference to Figures 20 and 21.

[0058] The total length of the fusion polypeptide may be any which is appropriate to the assay and aggregation disease protein core fragment being used, but will generally be greater than 30, 40, 50, 60, 70, 80, 90, 100, 200, or 500 amino acids in length.

[0059] In all instances herein where a named protein (e.g. core fragment) or a recited nucleic acid sequence is discussed, a derivative or other variant of the corresponding reference protein (or nucleic acid) may be used as appropriate, provided that it retains appropriate characteristics of the reference sequence. Such derivatives will also share sequence identity with the reference sequence.

[0060] In principle, a derivative may be a chemical derivative which may be prepared from the functional groups occurring as side chains on the residues or the NB or C-terminal groups, by means known in the art. These derivatives may include aliphatic esters of the carboxyl groups, amides of the carboxyl groups by reaction with ammonia or with primary or secondary amines, N-acyl derivatives of free amino groups of the amino acid residues formed with acyl moieties (e.g. alkanoyl or carbocyclic aroyl groups) or O-acyl derivatives of free hydroxyl groups (for example that of seryl- or threonyl residues) formed with acyl moieties.

[0061] More commonly the derivative will be one by way of amino acid insertion, deletion, or addition of the reference sequence. For example, a tau protein, or tau core fragment, derivative will comprise at least a partial amino acid sequence resembling the tandem repeat region of the tau proteins, but in which one or more of the

amino acids of the natural tau or its fragments have been replaced or deleted, or into which other amino acids have been inserted.

[0062] Such changes may be made to enhance or ablate binding activity (the latter case being useful for control experiments). Controls may contain deletions of sequences or domains to see what effect on aggregation these may have.

[0063] Preferred derivatives may be those which incorporate mutations corresponding to those known or suspected to be associated with the disease state. These may include changes corresponding to P301S within the tau sequence (see Fig 5). Other mutations include G272V, G389R, P301L, N279K, S305N, V337M, G272V, K280 Δ , R406W (see also Wischik et al, 2000, supra).

[0064] Other preferred derivatives may include tandem repeats of the core-fragments discussed above, or binding domains within those fragments.

[0065] Yet further derivatives may be based on chimeric products based on multiple, related, disease proteins in which their sequences are mixed or combined. For example restriction enzyme fragments of tau could be ligated together with fragments of MAP2 or even of an unrelated gene to generate recombinant derivatives.

An alternative strategy for modifying the core fragments would employ PCR as described by Ho et al., 1989, Gene 77, 51-59 or DNA shuffling (Cramer et al., 1998 Nature 391).

30 *Membrane localizing sequence*

[0066] The heterologous membrane-localising or targeting portion of the fusion may be any appropriate signal sequence from any suitable protein e.g. a signal sequence deriving from rat albumin. Alternatively it may be a modified sequence derived from a native localizing portion. However, in each case it will be a sequence which does not occur 'naturally' with the aggregation portion. The presence of a signal sequence on a protein will direct the nascent polypeptide/ribosome complex to the ER membrane where it may be inserted into it. Translocation across the membrane of the ER, and into the ER lumen, may then occur depending on the hydrophobicity of the protein.

[0067] The signal sequence may, depending on the chosen host cell for expression of a membrane-localized form of the protein, be a prokaryotic signal sequence e.g. an alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leader. For yeast expression the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter being described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179), or the signal described in WO 90/13646. In mammalian cell expression, a mammalian signal sequence such as that from rat albumin may be used, or alternatively a viral leader sequence may be employed.

[0068] This membrane localising portion may also include a membrane anchor sequence and/or a stop-transfer sequence. Generally this may be preferred for proteins, the intrinsic (hydrophobic) properties of which, may otherwise lead them to be secreted or located extracellularly, when fused with a signal peptide. A membrane anchor domain can be incorporated to prevent the entire protein from being translocated into the lumen of the ER and ultimately secreted. In general the protein must be oriented on the cytosolic surface of any protein with which it is associated such that it can interact with exogenous protein in the cytoplasm.

Other binding partners

[0069] In one aspect of the invention, the membrane-localized expression of a protein such as tau or a core fragment of tau can be used to screen for specific binding partners of that protein or fragment, e.g. proteins other than tau or the core fragment of tau which can bind to the membrane-localised form of tau or a fragment thereof.

[0070] Thus, the present invention also relates to methods of screening for specific binding partners of the protein (e.g. tau) of interest, comprising causing or allowing the membrane-localized expression of the protein, contacting that protein with one or more potential binding partners, and determining whether or not the potential binding partner has bound to the membrane-localized protein.

[0071] The detection of binding between the protein of interest and a potential binding partner may be carried out using any suitable method, as described in detail above.

[0072] The method may include the further step of recovering the specific binding partner thus identified.

[0073] In this way, the present invention enables the identification of specific binding partners of proteins such as tau, which may represent further pathological binding partners of such proteins in Alzheimer's disease-like aggregates.

Use of nucleic acid constructs

[0074] In one preferred embodiment the (bound) fusion polypeptide (of step (a)) above is introduced into the cell by expression therein under the transcriptional control of a constitutive promoter, while the second ('free') protein (of step (b)) is introduced by expression therein under the transcriptional control of an inducible promoter, in the presence of the appropriate inducing agent. Alternatively, in the reverse, the (a) and (b) proteins may be expressed inducibly and constitutively respectively.

[0075] This system can be used to control the level or timing of aggregation in the cell.

[0076] Particularly preferred is the following configuration for a cell-based tau-tau binding assay: A tau fragment corresponding to the truncated tau unit of the core

PHF is constitutively expressed in a membrane-localised form in a host cell, so as to expose the high affinity tau-tau binding site within the tandem repeat region, and at a level which does not kill the cells - i.e. in a stable cell line. The tau fragment is preferably longer than the core repeat domain fragment (suitably amino acid residues 186-390 or 186-441 of full-length tau), which is able to achieve a balance between aggregation and toxicity.

[0077] Nucleic acids of, or for use in, the present invention may be provided isolated and/or purified from their natural environment, in substantially pure or homogeneous form, or free or substantially free of other nucleic acids of the species of origin. Where used herein, the term "isolated" encompasses all of these possibilities.

[0078] A nucleic acid encoding the fusion polypeptide will be at least partially synthetic in that it will comprise nucleic acid sequences which are not found together in nature (do not run contiguously) but which have been ligated or otherwise combined artificially.

[0079] Nucleic acid according to the present invention may be in the form of, or derived from, cDNA, RNA, genomic DNA and modified nucleic acids or nucleic acid analogs. Where a DNA sequence is specified, e.g. with reference to a figure, unless context requires otherwise the RNA equivalent, with U substituted for T where it occurs, is encompassed.

[0080] In particular, nucleic acids may comprise, consist or consist essentially of any of the sequences disclosed herein (which may be a gene, a genomic clone or other sequence, a cDNA, or an ORF or exon of any of these etc.) Where a nucleic acid (or nucleotide sequence) of the invention is referred to herein, the complement of that nucleic acid (or nucleotide sequence) will also be embraced by the invention. The 'complement' in each case is the same length as the reference, but is 100% complementary thereto whereby by each nucleotide is base paired to its counterpart i.e. G to C, and A to T or U.

[0081] Thus the invention also relates, in a further aspect, to a nucleic acid molecule which comprises a nucleotide sequence encoding a fusion protein discussed above e.g. a aggregating disease protein or its core tandem repeat domain (i.e. a protein or fragment thereof that is implicated in a neurodegenerative disorder such as Alzheimer's disease) linked to a signal sequence and, if necessary, a sequence encoding a membrane anchor domain or a stop-transfer sequence, whereby the encoded polypeptide will become inserted into the ER membrane.

[0082] The nucleic acid molecule encoding the protein of interest linked to the signal sequence may optionally also include an exogenous 3'UTR sequence, e.g. a sequence derived from rat globin. The nature of the 3'UTR appears to influence the efficiency of targeting of a protein to the ER in cells, and the 3'UTR sequence can thus be selected accordingly (Partridge, K.A., et al. (1999) *Cyto-technology* 30, 37-47).

[0083] As described above, the nucleic acids may encode derivatives or other variants sharing homology with

the reference sequences in question. Preferably, the nucleic acid and/or amino acid sequence in question would share about 50%, or 60%, or 70%, or 80% identity, most preferably at least about 90%, 95%, 96%, 97%, 98% or 99% of the sequence upon which the variant is based. Similarity or homology may be as defined and determined by the TBLASTN program, of Altschul et al. (1990) J. Mol. Biol. 215: 403-10, which is in standard use in the art, or, and this may be preferred, the standard program BestFit, which is part of the Wisconsin Package, Version 8, September 1994, (Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA, Wisconsin 53711) using the default parameters. One common formula for calculating the stringency conditions required to achieve hybridization between nucleic acid molecules of a specified sequence homology is: $T_m = 81.5^{\circ}\text{C} + 16.6\text{Log} [\text{Na}^+] + 0.41 (\% \text{G+C}) - 0.63 (\% \text{formamide}) - 600/\text{#bp in duplex}$.

[0084] Nucleic acid sequences which encode a polypeptide or peptide linked to a signal sequence in accordance with the present invention can be readily prepared by the skilled person using the information and references contained herein and techniques known in the art (for example, see Sambrook, Fritsch and Maniatis, "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory Press, 1989, and Ausubel et al., Short Protocols in Molecular Biology, John Wiley and Sons, 1992). These techniques include (i) the use of the polymerase chain reaction (PCR) to amplify samples of the relevant nucleic acid, e.g. from genomic sources, (ii) chemical synthesis, or (iii) preparation of cDNA sequences. The nucleic acid thus prepared can then be linked to a sequence encoding a suitable signal peptide, or can be inserted into a vector construct which includes such a signal peptide-encoding sequence. Other methods suitable for adaptation to the synthesis of the membrane-localized protein in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-695 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

[0085] DNA encoding e.g. tau core fragments may be generated and used in any suitable way known to those of skilled in the art, including by taking encoding DNA, identifying suitable restriction enzyme recognition sites either side of the portion to be expressed, and cutting out said portion from the DNA. Modifications to the protein (e.g. tau)-encoding sequences can be made, e.g. using site directed mutagenesis.

Constructs

[0086] In the cell-based assay embodiments of the present invention, the construction of a membrane-localized form of the protein of interest, e.g. tau or a fragment of tau, can be achieved by causing or allowing the expression in a cell of an expression construct or vector, which comprises nucleic acid encoding the given protein linked to a signal sequence and, where required, an anchor or stop-transfer sequence, and which optionally also

includes an exogenous 3'UTR sequence.

[0087] In general, nucleic acid encoding the signal sequence to which the nascent protein molecule is linked may be a component of the vector itself, or it may be a part of the exogenous protein-encoding DNA that is inserted into the vector.

[0088] The construct may include any other regulatory sequences or structural elements as would commonly be included in such a system, and as is described below. As well as the signal sequence, the vector components will usually include, but are not limited to, one or more of an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

[0089] Nucleic acid sequences which enable a vector to replicate in one or more selected host cells are well known for a variety of bacteria, yeast, and viruses. For Example, various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

[0090] Particularly preferred is an expression vector comprising a nucleic acid as described herein. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, phage, or any other suitable vector or construct which can be taken up by a cell and used to express a membrane-localised form of the aggregating disease protein, e.g. tau or tau core fragment.

[0091] Expression vectors usually contain a promoter which is operably linked to the protein-encoding nucleic acid sequence of interest, so as to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. "Operably linked" means joined as part of the same nucleic acid molecule, suitably positioned and oriented for transcription to be initiated from the promoter. DNA operably linked to a promoter is "under transcriptional control" of the promoter. Transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g. the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[0092] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA.

[0093] The promoter may be "inducible" - which is to say, as is well understood by those skilled in the art, expression is "switched on" or increased in response to an applied stimulus. The nature of the stimulus varies be-

tween promoters. Some inducible promoters cause little or undetectable levels of expression (or no expression) in the absence of the appropriate stimulus. Other inducible promoters cause detectable constitutive expression in the absence of the stimulus. Whatever the level of expression is in the absence of the stimulus, expression from any inducible promoter is increased in the presence of the correct stimulus. In experiments below, a Lac inducible promoter has been used.

[0094] Expression vectors of the invention may also contain one or more selection genes. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins e.g. ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*. An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the given membrane-targeted protein-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell, when wild-type DHFR is employed, is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA 77: 4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid Rp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast which lacks the ability to grow in tryptophan, for example, ATCC: No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Transformation

[0095] The method referred to above may therefore further include introducing the nucleic acid into a host cell. The introduction, which may (particularly for *in vitro* introduction) be generally referred to without limitation as "transformation", may employ any available technique. For eukaryotic cells, suitable techniques may include calcium phosphate transfection, DEAE-Dextran, electroporation, liposome-mediated transfection and transduction using retrovirus or other virus, e.g. vaccinia or, for insect cells, baculovirus.

[0096] The calcium treatment employing calcium chloride, as described in Sambrook *et al.*, *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989.

[0097] For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No.

4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g. polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature 336:348-352 (1988).

Host cells

[0098] Suitable host cells for use in the invention may include bacteria, eukaryotic cells such as mammalian and yeast cells, and baculovirus systems.

[0099] Mammalian cell lines available in the art for expression of a heterologous polypeptide include fibroblast 3T6 cells, HeLa cells, baby hamster kidney cells, COS cells, monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651), Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumour cells (MMT 060562, ATCC CCL51); and many others. Preferred may be any of the range of well known neuroblastomal cell lines e.g. N2A, N1E, SY5Y.

[0100] Suitable prokaryotic hosts include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, *Enterobacteriaceae* such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Eukaryotic microbes such as filamentous fungi or yeast are also suitable cloning or expression hosts for vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. The selection of the appropriate host cell is deemed to be within the skill in the art.

[0101] In a further aspect, the present invention provides a host cell containing heterologous nucleic acid of the invention as described above. The nucleic acid of the invention may be integrated into the genome (e.g. chromosome) of the host cell. Integration may be promoted by inclusion of sequences which promote recombination with the genome, in accordance with standard techniques. Alternatively, the nucleic acid may be on an extrachromosomal vector within the cell, or otherwise identifiably heterologous or foreign to the cell. Generally the cell line will be capable of expressing a membrane-localised form of a protein such as tau or derivative thereof, the cell line being transformed with nucleic acid encoding the protein or derivative thereof. Preferred cell-lines may be based on the fibroblast cell line, e.g. 3T6.

Expression products

[0102] Host cells transfected or transformed with expression or cloning vectors described herein for the production of the membrane-localized protein may be cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in "Mammalian Cell Biotechnology: a Practical Approach", M. Butler, ed. JRL Press, (1991) and Sambrook *et al*, *supra*.

[0103] Gene expression can be confirmed in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl Acad Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequence of the aggregating disease protein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

[0104] Gene expression, alternatively, may be measured by immunological methods such as immunohistochemical staining of cells or tissue sections, and assay of cell culture, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence of the aggregating disease polypeptide.

[0105] Thus one aspect of the present invention entails causing or allowing expression from the nucleic acids discussed herein, e.g. by culturing host cells (which may include cells actually transformed, although more likely the cells will be descendants of the transformed cells) under conditions for expression of the gene, so that the encoded polypeptide (or peptide) is produced.

[0106] The present invention also encompasses a method of making a membrane-localized polypeptide or peptide of the preceding aspect, the method including expression from nucleic acid as described above. As an alternative to growing in a culture a host cell which contains such a vector, under appropriate conditions which cause or allow expression of the polypeptide, the polypeptides and peptides may be expressed in an *in vitro* system, such as a reticulocyte lysate system, as described herein.

[0107] Generally, following production by expression, the encoded fusion polypeptide will be utilised in the environment of the cells in which it is expressed, i.e. without isolating it from the membranes in which it is localised. In this way, as discussed above, it has surprisingly been

found by the present inventors, the membrane-localized form of the protein can be used to seed the aggregation of further protein molecules in a manner which mimics Alzheimer's disease-like protein aggregation. Thus, cells displaying membrane-localised Alzheimer's disease-like protein aggregates in their membranes (e.g. ER membranes) can be used in further screening assays (for a discussion of which, see below).

[0108] Nevertheless, polypeptides of the invention may be isolated and/or purified from the host cell and/or culture medium, either in the form of a membrane fraction, or isolated from the membranes themselves, as the case may be. Proteins may be purified to essential homogeneity by known methods such as, for example, by centrifugation at different velocities, by precipitation with ammonium sulphate, by dialysis (at normal pressure or at reduced pressure), by preparative isoelectric focusing, by preparative gel electrophoresis or by various chromatographic methods such as gel filtration, high performance liquid chromatography (HPLC), ion exchange chromatography, reverse phase chromatography and affinity chromatography (e.g. on Sepharose(TM) Blue CL-6B or on carrier-bound monoclonal antibodies).

[0109] Thus the invention relates, in a further aspect, to a polypeptide encoded by a nucleic acid as provided in a preceding aspect e.g. a polypeptide or fragment thereof, (for instance based on tau or a core fragment of tau) linked to a signal peptide and (if necessary) an anchor and/or stop-transfer sequence. In a preferred embodiment, the polypeptide comprises aggregating disease protein e.g. tau protein, or a derivative thereof. More preferably, the protein comprises a tau core fragment, as described above.

In vitro model

[0110] The membrane-localised form of the protein of interest described above may be used in *in vitro* models.

[0111] For example it could be used analogously to the tau fragment used in the assay of WO 96/30766 which is incorporated herein by cross reference. As described above, in this assay for tau aggregation and inhibitors thereof a fragment of tau corresponding to the core repeat domain is adsorbed to a solid phase substrate. This is able to capture soluble full-length tau and bind tau with high affinity. The combination of these two strategies (membrane localisation, *in vitro* solid phase) may be advantageous in certain contexts.

[0112] As a further alternative to the cell-based methods described above, the modeling method can be performed *in vitro*, by causing the expression in an *in vitro* protein-expression system of a membrane-localised form of the protein of interest. For instance, an *in vitro* medium containing microsomes (which can be easily produced by fragmentation of the endoplasmic reticulum), ribosomes, tRNA molecules, and mRNA encoding the protein or protein fragment of interest linked to a signal sequence, as well as any other soluble factors required

for protein synthesis and translocation, could be used to synthesise a membrane-bound form of the protein. Here, in a similar way to the cell-based system described above, ribosomes would become attached to the microsomal membrane (which is equivalent to the ER membrane of the cell-based system) and cause the synthesis of the encoded polypeptide molecule.

[0113] In this latter embodiment, however, the construct which encodes the aggregating disease protein, the signal sequence and the anchor and/or stop transfer sequences (if required), are all preferably designed in such a way that the majority of the encoded protein molecule, when inserted into the microsomal membrane, projects from the cytosolic surface of the membrane rather than into the lumen. In this way, the protein molecule will be accessible both to further protein molecules (so allowing Alzheimer's disease-like aggregation to occur at the cytosolic surface of the microsome), and to test agents in a screening assay to screen for potential inhibitors or modulators of such aggregation.

[0114] The skilled person will be familiar with the design of suitable constructs in order to achieve desired protein topologies. Further discussion of this point will be given below.

[0115] Thus, the common feature linking the cell-based and *in vitro* model systems provided herein is the synthesis of a membrane-localised form of the aggregating disease protein or protein fragment. This common feature gives rise to a method of generating a seed molecule, which can propagate the aggregation of further protein molecules in a manner which mimics the pathological aggregation of proteins such as tau during the formation of e.g. neurofibrillary tangles.

Detection of induced polymerization interaction

[0116] In various embodiments, the progress aggregation (or modulation of aggregation) may be detected directly or indirectly by monitoring the concentration or level any one or more of the following species: an aggregate of the fusion and/or second protein (e.g. based on sedimentation co-efficients), or a proteolytic fragment of either of these (which will generally increase with aggregation); a species corresponding to the non-aggregated fusion polypeptide and/or second protein (which may decrease).

[0117] Thus, as exemplified with particular tau fusions and proteins (based on 40kDa 190-441 fragment and T40), aggregation can be monitored on the basis of increasing levels of a 25 kDa processed species, derived from both the fusion and longer proteins (see Figure 1).

[0118] It may be desirable that the second ('free') protein used in method is labeled, or otherwise distinguishable from the membrane-bound protein, e.g. immunologically distinct, to assist in detecting the degree of aggregation of the protein molecules.

[0119] For instance, the free form of the aggregating disease protein may be linked to a reporter molecule.

The term "reporter molecule", as used in the present specification, is intended to mean a molecule which, by its chemical nature, provides an analytically-detectable signal that allows the detection of antigen-bound antibody. Detection should preferably be quantifiable, to allow determination of the amount of antigen in the sample; this may be calculated in absolute terms, or may be performed in comparison with a standard (or series of standards) containing a known normal level of antigen. Examples are well known to those skilled in the art e.g. enzymes or fluorophores.

[0120] Thus the free form of the protein may be directly modified (e.g. marked with a radioactive or enzymatically-detectable label) or conjugated (e.g. to a fluorophore) in a domain of the molecule, for example the N-terminal segment, which is known not to be involved in the high affinity tau-tau binding site, so that it thereby functions both as the ligand in the tau-tau binding assay, and as the reporter molecule.

[0121] Where antibodies or fragments thereof are used in embodiments of the method of the present invention may be produced by conventional techniques. Polyclonal antibodies may be raised e.g. by injecting the corresponding tau antigen into an animal, preferably a rabbit, and recovering the antiserum by immunoaffinity purification, in which the polyclonal antibody is passed over a column to which the antigen is bound and is then eluted in a conventional manner. Preferably the invention will use monoclonal antibodies which are selective to tau epitopes may be prepared by the method of Kohler and Milstein. Suitable monoclonal antibodies to tau epitopes can be modified by known methods to provide Fab fragments or (Fab')₂ fragments, chimeric, humanised or single chain antibody embodiments. Antibodies according to the present invention may be modified in a number of ways. Indeed the term "antibody" should be construed as covering any binding substance having a binding domain with the required specificity. Thus the invention covers antibody fragments, derivatives, functional equivalents and homologues of antibodies, including synthetic molecules and molecules whose shape mimics that of an antibody enabling it to bind an antigen or epitope.

[0122] Generally speaking, where antibodies are employed for detection, the antibody may carry a reporter molecule. Alternatively, detection of binding may be performed by use of a second antibody capable of binding to a first unlabelled, tau-specific antibody. In this case, the second antibody is linked to a reporter molecule.

[0123] Antibodies may be used in any immunoassay system known in the art, including, but not limited to: radioimmunoassays, "sandwich" assays, enzyme-linked immunosorbent assays (ELISA); fluorescent immunoassays, protein A immunoassays, etc. Typically, an immunoblot method is used. Preferably the immunoassay is performed in the solid phase, as would be well known to the skilled person. For instance, an antibody may be adsorbed to e.g. an assay column, and the cellular sample may then be washed through the column

under conditions suitable for enabling binding to the solid-phase antibody of any aggregate of the protein of interest, e.g. a tau-tau aggregate, which is present in the membrane of the cell. Excess reagent is washed away, and the binding of aggregated protein to the column can then be detected by any suitable means, e.g. as exemplified above and below.

[0124] Preferred monoclonal antibodies are as follows:

- Those which recognise the N-terminal or C-terminal of the tau epitope permit measuring of binding between truncated and full-length tau species. Especially useful are antibodies recognising human-specific epitopes. One such monoclonal antibody (designated 27/499) recognises a human-specific epitope located in the region between Gly-16 and Gln-26 of tau, and thereby permits measurement of binding between full-length tau species, provided one is derived from a non-human source (Lai (1995); "The role of abnormal phosphorylation of tau protein in the development of neurofibrillary pathology in Alzheimer's disease", PhD Thesis, University of Cambridge).
- Those which recognise the core tau fragment truncated at Glu-391. An example is mAb 423 (Novak *et al.* (1993), *loc. cit.*). This antibody enables detection of the binding of a truncated core tau fragment terminating at Glu-391 to a similar fragment terminating at Ala-390, which is not recognised by mAb 423. This truncation occurs naturally in the course of PHF assembly in Alzheimer's disease (Mena *et al.* (1995), (1996), *loc. cit.*; Novak *et al.* (1993), *loc. cit.*; Mena *et al.* (1991), *loc. cit.*). Additionally, when tau is bound via the repeat domain *in vitro*, digestion with a protease (e.g. pronase) generates a fragment detectable by mAb 423 (see Wischik *et al.*, 1996, *loc. cit.*). Figure 17 shows various tau-derived species which may be generated in an assay according to the present invention, and the approximate truncation point in each. mAb 423 may thus be used to demonstrate that the molecular conformation of the tau-tau binding interaction generated in a membrane-localised model system as provided herein corresponds to that which occurs pathologically in the brain, and hence that such model systems according to the present invention represent viable models of Alzheimer's disease-like protein aggregation.
- Those which recognise a generic tau epitope in the repeat domain. A preferred embodiment utilises an antibody (e.g. mAb 7.51). Where tau-MAP2 or MAP2-MAP2 aggregation is to be detected, an antibody which detects a generic MAP2 epitope could be used. Antibody 7.51 recognises a generic tau epitope located in the antepenultimate repeat of tau (Novak *et al.* (1991) Proc. Natl. Acad. Sci. USA, 88, 5837-5841), which is occluded when tau is bound in

a PHF-like immunochemical configuration but can be exposed after formic acid treatment (Harrington *et al.* (1990), (1991), *loc. cit.*; Wischik *et al.* (1995a), *loc. cit.*). Normal soluble tau, or tau bound to microtubules, can be detected using mAb 7.51 without formic acid treatment (Harrington *et al.* (1991), *loc. cit.*; Wischik *et al.* (1995a), *loc. cit.*). Binding of full-length tau in the tau-tau binding assay is associated with partial occlusion of the mAb 7.51 epitope.

[0125] Antibody 27/342 recognises a non-species specific generic tau epitope located between Ser-208 and Ser238 which is partially occluded in the course of the tau-tau interaction (Lai, *loc. cit.*).

[0126] The binding sites of some monoclonal antibodies are shown in Figure 4.

Screening for Modulators and inhibitors

[0127] As described above, the invention is preferably concerned with use of a nucleic acid molecule, polypeptide or derivative thereof, a host cell, or an *in vitro* protein expression system, as provided herein, in a method of modeling Alzheimer's disease-like protein aggregation.

Methods of the invention may include the steps of causing the expression of the nucleic acid molecule in a cell-based or microsome-based expression system, so that the encoded polypeptide becomes localized in a membrane (e.g. the ER membrane) of the cell, or in the mitochondrial membrane, respectively.

[0128] In a preferred aspect, there is disclosed a method of identifying a therapeutic agent for treatment of a disease as discussed herein, which method comprises the step of providing the fusion polypeptide (and preferably, the second protein discussed above) and contacting the membrane-localized protein with test substances which are suspected of being inhibitors or modulators of the disease associated protein aggregation. In particular, the fusion polypeptides as produced by the modeling methods described above can be used to investigate and screen for potential inhibitors or modulators of Alzheimer's disease-like protein aggregation, such as pathological tau-tau association.

[0129] Thus the various methods described above may comprise the further steps of (c) contacting the fusion polypeptide and/or the second protein with one or more agents which it is desired to assess for ability to modulate or inhibit the aggregation; (d) monitoring the extent of aggregation, and optionally correlating the extent of aggregation with the modulatory activity of the agent(s).

[0130] Thus a method of identifying a modulator of (preferably intracellular) aggregation of a protein associated with a disease in which the protein undergoes an induced conformational interaction, may comprise performing a method for inducing aggregation as described above in the presence of one or more agents suspected of being capable of modulating (e.g. inhibiting or revers-

ing) the aggregation. The degree of aggregation (and optionally proteolytic processing) may be observed in the presence or absence of the agent, and the relative values correlated with its activity as a modulator.

[0131] Preferably, where the screening method employs cells which express a membrane-localised form of the protein of interest, such as tau, these may comprise, e.g., cells from a neuronal cell line, e.g. a neuroblastoma cell line, or cells from a fibroblast cell line. In a preferred embodiment, the cells are from a fibroblast 3T6 cell line.

[0132] In one embodiment, the present invention relates to a method of screening for agents which are capable of modulating or inhibiting Alzheimer's disease-like protein aggregation, e.g. pathological tau-tau association, the method comprising:

- (1) causing or allowing the synthesis of a membrane-localised form of the aggregating disease protein (e.g. tau or MAP2), or a core fragment thereof (e.g. the core tandem repeat domain of tau or MAP2);
- (2) contacting said fusion polypeptide or derivative with:

- (A) one or more agents suspected of being capable of modulating or inhibiting aggregation of the aggregating disease protein molecules, and
- (B) a further aggregating disease protein or core fragment thereof, which is capable of binding to the membrane-associated protein of step (1); and

- (3) detecting the binding, if any, of the protein of step (2)(b) with the fusion polypeptide of step (1).

[0133] The tau-like protein used in step (2B) above may, in certain embodiments, be a "free" form of the protein, which term is used herein to distinguish it from the membrane-bound aggregating disease protein of step (1). However, "free" is not intended to exclude the possibility that the aggregating disease protein is complexed with, or otherwise associated with, other molecules or moieties such as carbohydrate moieties or other peptides.

[0134] For example, a test substance may be added to a cellular system as described above, and the cells incubated for a period of time sufficient to allow binding and to demonstrate inhibition of binding. The bound tau complex can then be detected, e.g. using a suitably-labeled antibody such as MAb 7.51 in an immunoblot of total cell extract, or any other suitable detection method. If desired, the membrane-bound tau protein can be distinguished from aggregated free tau protein, for example by terminating the sequence of the membrane-bound tau protein at amino acid 390. The aggregated free tau will be proteolytically cleaved at amino acid Gly 391, which can be detected by an antibody specific to that epitope (e.g. MAb 423).

[0135] Where a screening method is employed for this

purpose, i.e. for the identification of modulatory/inhibitory compounds, a non-competitive or competitive assay may be used. For instance, in a competitive assay of the type well known in the art, the effect of a known inhibitor or modulator on Alzheimer's disease-like aggregation can be compared in the presence or absence of further test substances or agents, to determine the ability of the test substance to compete with the known inhibitor/modulator for binding to the protein of interest.

Choice of test compound

[0136] Compounds which are tested may be any which it is desired to assess for the relevant activity

[0137] The methods can serve either as primary screens, in order to identify new inhibitors/modulators, or as secondary screens in order to study known inhibitors/modulators in further detail.

[0138] Agents may be natural or synthetic chemical compounds. Antibodies which recognise an Alzheimer's disease-like protein aggregate and/or which modulate Alzheimer's disease-like protein aggregation form one class of putative inhibitory or modulatory compounds with respect to the aggregation process. More usually, relatively small chemical compounds, preferably which are capable of crossing the blood-brain barrier, will be tested.

[0139] The skilled person will appreciate that the amount of test substance or compound which is added in a screening assay according to this aspect of the invention will normally be determined by trial and error depending upon the type of compound used. It may be selected to be a level which could realistically be used in therapeutic context i.e. would be non-lethal to a patient.

Specificity of inhibition

[0140] Screening methods according to this aspect of the present invention may be used to screen for compounds which demonstrate the properties of selective competitive inhibition of pathological Alzheimer's disease-like protein aggregation (e.g. tau-tau or tau-MAP2 binding), without interference with the normal binding of e.g. tau or MAP2 to tubulin via the same region of the molecule (see WO 96/30766).

[0141] A suitable method for determining any possible interference of the binding of tau, MAP2 or a derivative thereof to tubulin by potential inhibitors/modulators of Alzheimer's disease-like protein aggregation, comprises contacting a preparation of depolymerised tubulin/MAP2 or taxol-stabilised microtubules with an agent suspected of being capable of modulating or inhibiting tau-tau or tau-MAP2 association, along with a protein or derivative thereof as mentioned above in step (2)(b), followed by detection of the tau-tubulin or MAP2-tubulin binding. Tau-tubulin binding could also, for example, be demonstrated by a normal cytoskeletal distribution, as described in e.g. WO 96/30766.

[0142] Methods for the preparation of tubulin proteins

or fragments thereof, possibly in combination with binding partners, are known in the art and are described e.g. by Slobada et al. (1976, in: Cell Mobility (R. Goldman, T. Pollard and J. Rosenbaum, eds.), Cold Spring Laboratory, Cold Spring Harbor, New York, pp 1171-1212).

Cell viability

[0143] Where the method utilizes a cell-based system, it may further include the step of testing the viability of the cells expressing the membrane-localised aggregating disease protein, e.g. by use of a lactate dehydrogenase assay kit (Sigma).

[0144] In the case where tau-tau, tau-MAP2 or MAP2-MAP2 aggregation is being investigated (see above, under 'specificity'), this step may also provide an indication of any interference by the test agent of the binding of tau or MAP2 to tubulin, since inhibition or interference of tau-tubulin or MAP2-tubulin binding will correlate to some extent with a decreased ability of the cells to divide, and thus with decreased cell viability.

Therapeutics and modes of administration

[0145] Performance of a screening assay method according to the various aspects above may be followed by isolation and/or manufacture and/or use of a compound, substance or molecule which tests positive for ability to interfere with or modulate disease related protein aggregation.

[0146] The compounds thus identified may be formulated into compositions for use in the diagnosis, prognosis or therapeutic treatment of Alzheimer's disease or the like (indeed any condition in which Alzheimer's disease-like protein aggregation occurs). Thus, the present invention provides the means to obtain pharmaceutical formulations comprising one or more inhibitory or modulatory compound as obtainable by a screening method as provided herein.

[0147] Following the identification of a substance or agent which modulates or affects such protein aggregation, the substance or agent may be investigated further. Other candidate inhibitory/modulatory compounds may be based on modelling the 3-dimensional structure of a polypeptide or peptide fragment and using rational drug design to provide potential inhibitor compounds with particular molecular shape, size and charge characteristics. Thus also disclosed herein is the use of an agent as identified using a screening method provided herein, or a derivative, active portion, analogue, variant or mimetic thereof which is able to inhibit or modulate Alzheimer's disease-like protein aggregation, in screening for, or designing, further inhibitors/modulators of Alzheimer's disease-like aggregation.

[0148] A compound which has been identified as described above, may be manufactured and/or may be used in the preparation, i.e. the manufacture or formulation, of a composition such as a medicament, pharma-

ceutical composition or drug. These may be administered to individuals.

[0149] Generally, such an inhibitor or modulator is provided in an isolated and/or purified form, i.e. substantially pure. This may include being in a composition where it represents at least about 90% active ingredient, more preferably at least about 95%, more preferably at least about 98%. Such a composition may, however, include inert carrier materials or other pharmaceutically and physiologically-acceptable excipients. As noted below, such a composition may include in addition to an inhibitory/modulatory compound as disclosed, one or more other molecules of therapeutic use.

[0150] In this way, a substance identified as described above, such as an inhibitor of tau-tau aggregation or other such Alzheimer's disease-like protein aggregation, may be provided for use in a method of treatment of the human or animal body of a condition in which Alzheimer's disease-like protein aggregation occurs.

[0151] Also described herein are a pharmaceutical composition, medicament, drug or other composition comprising a substance of the invention as described above, a method comprising administration of such a composition to a patient, e.g. for treatment or prophylaxis of Alzheimer's disease or an Alzheimer's disease-like condition, use of such a substance in the manufacture of a composition for administration, e.g. for treating Alzheimer's disease or similar treatment, and a method of making a pharmaceutical composition comprising admixing such a substance with a pharmaceutically acceptable excipient etc. as discussed below.

[0152] Whether it is a polypeptide, antibody, peptide, nucleic acid molecule, small molecule, mimetic or other pharmaceutically-useful compound as described herein that is to be given to an individual, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors.

[0153] A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

[0154] Pharmaceutical compositions as described herein, may include, in addition to active ingredient, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous or intravenous.

[0155] Pharmaceutical compositions for oral adminis-

tration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

[0156] For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

[0157] Examples of techniques and protocols mentioned above can be found in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed), 1980.

[0158] The agent may be administered in a localised manner to the brain or other desired site, or it may be delivered systemically in a manner such that it targets the brain or other cells. For instance, targeting therapies may be used to deliver the active agent more specifically to certain types of cell, by the use of targeting systems such as antibody or cell-specific ligands. Targeting may be desirable for a variety of reasons, for example if the agent is unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.

[0159] Instead of administering these agents directly, they may be administered in a precursor form, for conversion to the active form by an activating agent produced in, or targeted to, the cells to be treated.

[0160] An agent which is able to interfere with Alzheimer's disease-like protein aggregation may be provided in a kit, e.g. sealed in a suitable container which protects its contents from the external environment. Such a kit may include instructions for use.

Animals

[0161] Although much of the forgoing discussion has been concerned with cell-line or *in vitro*-based assays, the invention disclosed herein has particular utility in animal-based assays.

[0162] Thus host cells according to the present invention may be comprised in a transgenic animal, and the present invention further provides a transgenic animal, comprising cells which express a fusion polypeptide according to a preceding aspect, such as a membrane-localised form of the tau core fragment. Such animals may be prepared and/or used in analogous manner to those discussed in US 5,912,410 and 5,898,094.

[0163] The transgenic organisms of the invention all include within a plurality of their cells a cloned recom-

binant or synthetic DNA sequence which is believed to relate to the pathogenesis of a disease of protein aggregation. More specifically, the transgenic organisms contain specific sequences of exogenous genetic material, such as the sequences described above in detail which are comprised of a tissue specific promoter sequence and a sequence which encodes for production of a membrane targeted protein. Since it is possible to produce transgenic organisms of the invention utilizing one or more of the above-described sequences, a general description will be given of the production of transgenic organisms by referring generally to exogenous genetic material. This general description can be adapted by those skilled in the art in order to incorporate the above-described specific DNA sequences into organisms and obtain expression of those sequences utilizing the methods and materials described below. For more details regarding the production of transgenic organisms, and specifically transgenic mice, refer to U.S. Pat. No. 4,873,191, issued Oct. 10, 1989 and to the numerous scientific publications referred to and cited therein.

[0164] The exogenous genetic material may be placed in either the male or female pronucleus of the zygote. More preferably, it is placed in the male pronucleus as soon as possible after the sperm enters the egg. In other words, right after the formation of the male pronucleus when the pronuclei are clearly defined and are well separated, each being located near the zygote membrane. The male pronucleus of a fertilized mouse egg is the preferred site for addition of the exogenous genetic material of the present invention.

[0165] It is most preferred that the exogenous genetic material be added to the male DNA complement of the zygote prior to its being processed by the ovum nucleus or the zygote female pronucleus. It is thought that the ovum nucleus or female pronucleus release molecules which affect the male DNA complement, perhaps by replacing the protamines of the male DNA with histones, thereby facilitating the combination of the female and male DNA complements to form the diploid zygote.

[0166] Thus, it is preferred that the exogenous genetic material be added to the male complement of DNA or any other complement of DNA prior to its being affected by the female pronucleus. For example, the exogenous genetic material is added to the early male pronucleus, as soon as possible after the formation of the male pronucleus, which is when the male and female pronuclei are well separated and both are located close to the cell membrane. Alternatively, the exogenous genetic material could be added to the nucleus of the sperm after it has been induced to undergo decondensation. Sperm containing the exogenous genetic material could then be added to the ovum or the decondensed sperm could be added to the ovum with the exogenous genetic material being added as soon as possible thereafter.

[0167] For the purposes of this invention a zygote is essentially the formation of a diploid cell which is capable of developing into a complete organism. Generally, the

zygote will be comprised of an egg containing a nucleus formed, either naturally or artificially, by the fusion of two haploid nuclei from a gamete or gametes. Thus, the gamete nuclei must be ones which are naturally compatible, i.e., ones which result in a viable zygote capable of undergoing differentiation and developing into a functioning organism. Generally, a euploid zygote is preferred. If an aneuploid zygote is obtained, then the number of chromosomes should not vary by more than one with respect to the euploid number of the organism from which either gamete originated.

[0168] In addition to similar biological considerations, physical ones also govern the amount of exogenous genetic material which can be added to the nucleus of the zygote or to the genetic material which forms a part of the zygote nucleus. If no genetic material is removed, then the amount of exogenous genetic material which can be added is limited by the amount which will be absorbed without being physically disruptive. Generally, the volume of exogenous genetic material inserted will not exceed about 10 picoliters. The physical effects of addition must not be so great as to physically destroy the viability of the zygote. The biological limit of the number and variety of DNA sequences will vary depending upon the particular zygote and functions of the exogenous genetic material and will be readily apparent to one skilled in the art, because the genetic material, including the exogenous genetic material, of the resulting zygote must be biologically capable of initiating and maintaining the differentiation and development of the zygote into a functional organism.

[0169] The number of copies of the DNA sequences which are added to the zygote is dependent upon the total amount of exogenous genetic material added and will be the amount which enables the genetic transformation to occur. Theoretically only one copy is required; however, generally, numerous copies are utilized, for example, 1,000-20,000 copies of a gene, in order to insure that one copy is functional. As regards the present invention, there is generally an advantage to having more than one functioning copy of each of the inserted exogenous DNA sequences to enhance the phenotypic expression of the exogenous DNA sequences.

[0170] Any technique which allows for the addition of the exogenous genetic material into nucleic genetic material can be utilized so long as it is not destructive to the cell, nuclear membrane or other existing cellular or genetic structures. The exogenous genetic material is preferentially inserted into the nucleic genetic material by microinjection. Microinjection of cells and cellular structures is known and is used in the art.

[0171] Thus the present invention provides methods in which cloned recombinant DNA sequences encoding appropriate membrane targeting sequences may be injected into fertilized mammalian eggs (preferably mouse eggs). The injected eggs are implanted in pseudo pregnant females and are grown to term to provide transgenic mice whose cells express proteins related to the pathol-

ogy of the relevant disease. The injected sequences are constructed having promoter sequences connected so as to express the desired protein in specific tissues of the transgenic mammal (most notably in nerve tissue). Suitably, the vector is adapted for neurone-specific expression to produce Alzheimer's Disease-like aggregation such as is now known to occur in the human brain. Examples may include the prion specific promoter of Lewis et al (2000) or the neurospecific enolase promoter e.g. of Thunecke et al (2000).

[0172] Non-human animals of the invention may be homozygous or heterozygous for the fusion polypeptide. Mammalian animals include non-human primates, rodents, rabbits, sheep, cattle, goats, pigs. Rodents include mice, rats, and guinea pigs. Specifically provided are:

- Methods of preparing a transgenic animal model with enhanced or accelerated pathology for a disease of protein aggregation and the transgenic animal made by such methods.
- Methods of producing an F₁ generation by crossing a founder animal of either sex (F₀ generation) with an animal which is non-transgenic in respect of the proteins discussed herein, and is preferably wild-type). The offspring (F₁ generation) may then be screened and those which carry a transgene resulting in an enhanced pathology for the disease, are selected.
- Methods of producing an F₂ generation by crossing 2 F₁ animals of appropriate sex. The offspring (F₂ generation) may then be screened and those which carry a transgene in the appropriate dosage (i.e. hetero or homozygous) resulting in an enhanced pathology for the disease, are selected.
- A method of modulating the disease phenotype of a transgenic animal model, and the transgenic organism with the modulated phenotype, as described above.

[0173] Transgenic non-human mammals of the invention may thus be used for experimental purposes in studying Alzheimer's-like diseases, and in the development of therapies designed to alleviate the symptoms or progression of such conditions. By "experimental" it is meant permissible for use in animal experimentation or testing purposes under prevailing legislation applicable to the research facility where such experimentation occurs.

[0174] The animal may be used to screen for or test potential diagnostic, therapeutic or prognostic agents of Alzheimer's disease-like protein aggregation (see below for further discussion). This aspect of the invention thus provides an animal model which can aid in the investigation of Alzheimer's disease and Alzheimer's disease-like conditions. In the in vivo model system, the Alzheimer's disease-like condition is induced in the animal by causing or allowing a membrane-localised Alzheimer's disease-like protein aggregate to be synthesised in the cells. Using this model system, a compound suspected

of having a therapeutic effect in relation to Alzheimer's disease or an Alzheimer's disease-like condition can be administered to the animal, and any effects on the condition (e.g. improvements in symptoms, or any other suitable indicator) can be studied. In particular, the transgenic mammals of the present invention are useful in determining the effectiveness of pharmaceutical drugs with respect to their ability to decrease the amount of aggregates which form within the brain of the animal. These aggregates may be measured by biochemical and/or histological techniques used in measurements of AD in humans (see e.g. Mukaetova-Ladinska, E.B. *et al.* (2000) *loc cit.*).

[0175] Most preferably the aggregation (or inhibition thereof) is assessed by the production (or inhibition, or abolition, of production) of truncated fragments of tau resulting from proteolytic processing e.g. of endogenous, full length, tau. In preferred embodiments the aggregation of tau is evidenced by proteolytic processing to a truncated core fragment corresponding to that found in the actual AD PHF core. However the production of any one or more other fragments which are diagnostic products or bi-products of this process may also be used.

[0176] The mammals are thus useful in testing the efficacy of such drugs, in a pharmacokinetic context, in preventing the formation or reducing the amount of aggregation formed as well as eliminating or reducing aggregation already formed.

[0177] Generally speaking, a drug to be tested is administered to a control animal or group of animals which are not the transgenic animals of the invention and simultaneously to transgenic animals of the invention. The drug is preferably continuously administered over a period of time which is normally sufficient to effect the formation of aggregates in the brain of the animal. After administering the drug for a sufficient period of time the control animal(s) along with the transgenic animal(s) are sacrificed. Examination of the brain of the animals is made as described above.

[0178] In such a system, the effect of a potential inhibitor or modulator of protein aggregation can be determined or detected as described above e.g. by immunoassay, for instance to observe the inhibition of the formation of tau, or tau-like aggregates such as those detected in human brain tissue by the methods of e.g. Wischik *et al.* (1995; *loc. cit.*), Lai *et al.* (1995; *loc. cit.*), or Mukaetova (2000; *loc. cit.*) in which measurement of aggregated forms of tau protein is undertaken in brain tissues.

[0179] For instance, the production of aggregates within cells may be visualized histologically and by electron microscopy, and the prevention of these aggregates similarly assessed.

[0180] By comparing the amount of aggregate deposit within the control animal(s) to the amount of deposit within the transgenic mammal(s) of the invention a determination can be made with respect to the effectiveness of the drug. Comparative drug testing protocols known to

those skilled in the art can be used in connection with the transgenic mammals of the invention in order to test drugs. The final intracellular concentration of the drug may be selected to be appropriate to the precise disease protein and drug in question, but may be in a range which will ultimately be appropriate for clinical usage in terms of toxicity, uptake etc. (e.g. 1 μ M - 1mM more preferably 4 - 600 μ M).

[0181] The *in vivo* screening assay of the invention need not directly measure the degree of inhibition of, or modulation of, Alzheimer's disease-like protein aggregation caused by the compound being tested. Instead the downstream effect on the pathological condition itself, e.g. Alzheimer's disease, may be measured. For instance, it would be anticipated that there would be behavioral effects in the animals, such as defects of movement, power or tone, abnormal reflexes, evidence of defective memory or response to environmental cues.

[0182] In that the transgenic animals of the invention can be used to test the efficacy of drugs with respect to preventing disease-associated protein aggregation, the animals are valuable research tools with respect to allowing researchers to test the efficacy of such drugs in treating diseases such as Alzheimer's disease. If preferred, in all cases, the transgenic animal assay may be run in parallel with, or subsequent to, an *in vitro* or cell-based assay, e.g. according to preceding aspects of the present invention, in order to confirm that any effect on the condition itself is as a result of the inhibition of Alzheimer's disease-like protein aggregation and not merely a general toxic effect.

[0183] Thus, as is described in detail above, the present invention relates *inter alia* to the use of a nucleic acid molecule, polypeptide or derivative thereof, a host cell, a transgenic animal or an *in vitro* membrane-localized model system, all as provided herein, in methods of screening for test substances having an inhibitory or modulatory effect on Alzheimer's disease-like protein aggregation. In further aspects, the present invention extends to inhibitory or modulatory compounds obtainable using a screening method of a preceding aspect.

[0184] Various further aspects and embodiments of the present invention will be apparent to those skilled in the art in view of the present disclosure. Certain aspects and embodiments of the invention will now be illustrated by way of example only, and with reference to the following Figures and Examples.

Figures

[0185] Figure 1 shows a diagrammatic illustration of an assay according to the invention as described in Example 1 (cell based) and Example 3 (animal model).

(A) (1:) Assay based on constitutive expression of membrane-targeted sequence alone, in this case a Signal Sequence plus truncated tau (SStau190-441). The membrane associated mole-

cules combine leading to formation of a 25 kD fragment. As shown in Figure 17(a) the 25 kD fragment contains immunoreactivity associated with the repeat domain and the C-terminus, but not an epitope located between residues 208 - 238. The fragment length is ~ 175 residues and the likely N-terminal cleavage site is at residue ~266, the phase-shifted N-terminus of the first repeat found in tau protein isolated from the PHF-core.

(A) (2:) Assay based on second protein (T40 - full-length tau), the proteolytic cleavage of which is "seeded" by the presence of low background levels of SStau190-441 which is targeted to the endoplasmic reticulum by the signal sequence. T40 induced by IPTG is bound to the SStau190-441 species either at the membrane or after release of the SStau190-441-derived aggregate into the cytosol. This captured T40 is now further processed first to give rise to increased levels of a species with gel mobility and immunoreactivity identical to that of the SStau190-441-derived species. It is then further processed through the same pathway to generate increased levels of the 25 kD degradation product. Tau aggregation inhibitors block T40-derived production of the 25 kD fragment in the presence of IPTG.

(B) Assay based on second cleavage pattern of SStau190-441. In this case the membrane-associated molecules combine in a slightly different manner through the repeat region to give rise to three further truncation products (seen in Figure 17(a)) with gel mobilities 30 kD (~200 residues), 18 kD (~125 residues) and 12/14 kD (~95 residues).

(C) Assay based on constitutive expression and aggregation of SStau190-390. In this case, proteolytic processing gives rise to a single predominant species with gel mobility of 18 kD (~125 residues). This arises by aggregation of SStau190-390 molecules through the repeat domain, giving rise to a proteolytically protected fragment derived entirely from the repeat domain (ie ~266 - ~390). In a variant of the repeat domain binding pattern, illustrated in Figure 17(b)), a slightly different 18 kD species could arise by association of SStau190-390 with low background levels of T40, to give rise to a fragment equivalent to 4-repeats in length (ie ~125 residues), but extending by the equivalent of 1 repeat beyond 390 within the C-terminal tail of the molecule (ie ~295 - ~422). In a different scheme illustrated in Figure 16d, small quantities of a fragment consisting of only 3 repeats (ie ~95 residues) are able to form minority dimeric forms with atypical gel mobility corresponding to ~19/22 kD.

(D) Assay based on constitutive expression of

SStau296-390 in the presence of background T40. In this case, two predominant species are formed, one of 12/14 kD corresponding to the SStau296-390 fragment itself, and a further species with gel mobility of 18 kD. The latter cannot be a degradation product of SStau296-390. It is also unlikely to be a major form of the dimer (see Figure 16d) which has a predominant gel mobility corresponding to 24 kD. Therefore, the 18 kD is most likely to arise by truncation of T40 which has been captured by an SStau296-390 aggregate. This species again corresponds to the equivalent of 4 repeats (ie ~125 residues), and could arise through two possible binding interactions between T40 and the SStau296-390 aggregate as shown. In this model the rate limiting step for the formation of the 18 kD species from T40 may be the determined by the amount of aggregating, truncated, tau in the system.

(E) Assay based on constitutive expression of SStau296-390 within the transgenic mouse brain. As in the system described in (D), the predominant product is not the SStau296-390 itself, but a higher molecular weight species of 18 kD. For the reasons discussed above, this is unlikely to be a dimer or a degradation product of the SStau296-390 species introduced into the animal transgenically. Rather, it is most likely to arise by the capture and proteolytic processing of endogenous mouse tau. In this case, the mouse brain is known to express only the 3-repeat isoform. The 18 kD fragment generated in this case is most likely to be the form with the C-terminal extension giving rise to a species the equivalent of 4 repeats in length (~125 residues, or the fragment between residues ~266 and ~422 in the 3-repeat isoform). However, a slightly different binding pattern is also possible, in which the extension is into the N-terminus (ie the fragment between residues ~235 and ~390 in the 3-repeat isoform).

Figure 2. The inhibitory activities of thionine, chlorpromazine and tacrine were compared in the membrane-targeted cell model (production of 25 kD species). There is a clear distinction between active and inactive compounds (see Example 2).

Figure 3. The data from Figure 2 was analysed in greater detail mathematically using a standard inhibitory model, and a numerical procedure (Gauss-Newton method) to solve for KI. The data could be approximated very closely by assuming a Kd value for tau-tau binding within the cell of ~500 nM, and a corresponding KI value of ~33 nM. The significance of these values can be understood by comparing with what is known of the influence of phosphorylation on tau-tau binding in vitro (see Wischik, C.M., et al. (1997) *loc cit*).

Figure 4 (a) shows a schematic illustration of the structure of a paired helical filament (top) and the immunohistochemistry of neurofibrillary tangles during progression of Alzheimer's disease (bottom). Figure 4 (b) shows a schematic illustration of the binding sites of various monoclonal antibodies to different forms of N- and C-truncated tau.

Figure 5 (a) shows the nucleotide and predicted amino acid sequences of a human tau protein isoform. The sequence was deduced from cDNA clone httau40. Figure 5 (b) shows the sequences of some of the constructs of the present invention.

Figure 6 (a) shows constructs used for microinjection to prepare transgenic mice. These are (i) Thyl-sstau296-390, and (ii) Thyl-sstau186-441. Figure 6 (b) shows construct Thyl-SStau296-390 in more detail. This was cloned into the Thy-1 expression vector and was then microinjected into fertilized oocytes of NMRI mice using methods from: "Manipulating the mouse embryo", a laboratory manual, second edition, Cold Spring Harbour Press, CSH, N.Y. Hogan, B., Beddington, R., Costantini, F., Lacy, E., editors, 1994).

Figure 7 shows a schematic representation of tau, MAP2 (adult form), MAP2C (juvenile form) and high molecular weight tau (found in the PNS and neuroblastoma cell lines). These proteins share similar microtubule-binding domains, but differ substantially in sequence and extent of the N-terminal projection domain. The juvenile forms of tau and MAP2 have only 3 of the tandem repeats. A 4-repeat form of MAP2 also exists.

Figure 8 shows sequence differences in the tandem repeat region of human tau (upper line) and human MAP2 (lower line). Vertical arrows show the limits of the truncated PHF-core fragment terminating at Glu-391, and the tubulin-binding segments are shown underlined.

Figure 9 shows a construct according to one embodiment of the present invention. (a) The signal sequence of rat albumin was inserted in front of the start codon of the cDNA encoding rabbit globin to achieve membrane targeting of the globin mRNA. Figure 9(b) shows a fragment of the human tau cDNA encoding amino acids 190-441 was inserted to replace part of the globin sequence, to effect expression of a tau fragment with an N-terminal signal sequence. The globin 5' and 3' UTRs were retained in the construct.

Figure 10 shows a summary of tau fragments of the present invention which were expressed in host cells. Three fragments of tau, all containing the core

aggregation domain, were expressed with N-terminal signal sequences. For comparison, soluble forms of two of the fragments were also expressed, as well as full-length tau.

Figure 11 shows the localization of SStau190-390 after stable transfection of 3T6 cells. 3T6 cells isolated as stable clones after transfection with a cDNA encoding amino acids 190-390 of the longest isoform of human tau linked to an N-terminal signal sequence were incubated with 5 mM sodium butyrate for 20 h to increase protein expression, fixed with paraformaldehyde and labeled with mAb 7.51 directed against the repeat domain of tau. These cells show similar tau morphology to cells expressing SStau190-441, but cells isolated after stable transfection with SStau190-390 consistently showed higher levels of expression of the tau fragment, with a higher proportion showing large tau aggregates (see Example 1).

Figure 12 shows a Northern blot analysis of wild-type ("Wt") and transgenic mouse line 1 ("1") RNA with a specific probe showing a high expression level of transgenic RNA in transgenic mouse line 1. S26 control shows same loading amounts (see Example 3).

Figure 13 shows purification of the SStau290-391 by immunoprecipitation using mAb 7.51. 0.1g mouse brain was homogenized using 500µl lysis buffer (150mM NaCl, 0.01%SDS, 0.1% NP40, 50mM Tris, 100µg/ml PMSF, 1mM DTT, 1µg/ml Leupeptin). The suspension was then sonicated on ice for 1min and spun down with a benchtop (Eppendorf) centrifuge at 14,000rpm for 4min. This leads to the "S1" supernatant fraction. The first immunoprecipitation was undertaken using mAb 342 to deplete the preparation of endogenous mouse tau. The S1 fraction was incubated with 1/10 volume of antibody 342 (1:10), incubated for 2h at 4°C with rocking, then 100µl of ProteinA sepharose beads (10% v/v in lysis buffer) were added, incubated for 1h at 4°C, centrifuged with a benchtop centrifuge at 14,000rpm for 30sec. Supernatant was taken for further immunoprecipitation using mAb 7.51. 1/10 v/v of 7.51 was added, incubated for 2h at 4°C with rocking. 40µl of ProteinA sepharose beads were added, incubated for 1h at 4°C, centrifuged for 15sec at 14000rpm, washed 3 times with the above mentioned lysis buffer. Then, 30µl of laemmli sample buffer was added, heated to 85°C for 5min, centrifuged for 30sec. Supernatant was then loaded onto 15% SDS-Page gel. The IP751 lanes indicate a difference at -12-14 kD corresponding to presence of the SStau290-391 protein product in a transgenic animal from line 1. The dark background at higher molecular weight is due to presence of endogenous mouse immunoglobulins detected in the course of development of the immunoblot (see Example 3).

Figure 14 shows the S1 fraction prepared in lysis buffer as indicated above without immunoprecipitation. Immunoblotting using mAb 7.51 revealed in addition to the -12-14 kD band corresponding to the transgene a new more prominent species with gel mobility ~18 kD. This was seen in all line 1 individuals to varying degrees according to the age of the animal up to 15 months. Three are shown of ages 5 months, 4 months and 4 months respectively. Epitope mapping (not shown) demonstrated that the 18 kD species was recognised by mAb 7.51 (epitope in the repeat domain) and a polyclonal antiserum K9JA ("DAKO A/S", also recognising epitopes located in the repeat domain of tau). The 18 kD species was not recognised by the C-terminal polyclonal antiserum T46.1 or mAb 342, which recognises an epitope located between residues 208 - 238. The 18 kD species cannot arise as a degradation product of the transgene product, nor can it arise as a dimer of SStau296-390. The 18 kD species is therefore likely to be generated by aggregation-dependent proteolytic processing of endogenous mouse brain tau as shown in Figure 1.

Figure 15 shows two dimensional gel electrophoresis (2DE), silver stain and immunoblotting. 500mg of mouse brain was homogenized in 2ml of lysis buffer (see above), sonicated on ice, centrifuged at 14000rpm. The S1 fraction was taken and boiled for 5min at 95°C, centrifuged at 14000rpm for 5min, supernatant was taken and concentrated 10-fold using a centricon3000. 20% of this protein suspension was loaded onto a 2-DE gel. 1st dimension isoelectric focusing, 2nd dimension 15% SDS-PAGE. Gels were silver stained and in parallel gels were run under same conditions and blotted, then incubated with 7.51 antibody (see above). The black arrows show the 12-14 kD gene product and a doublet corresponding to the 18 kD species. In addition there is an as yet uncharacterised smear of lower gel mobility which also distinguished the transgenic animal from the wild-type. This smear is likely to contain several higher molecular weight tau products also derived from endogenous mouse tau. Other isolated higher molecular weight tau-immunoreactive spots are also variably observed, depending preparative protocol used (see Example 3).

Figure 16 shows the results obtained with the various cell lines of Example 1.

Figure 16 (a) shows an analysis of cells expressing tau fragments by immunoblotting. Cells stably expressing tau fragments, and also expressing full-length tau under the control of an IPTG-inducible promoter, were incubated with 5 mM IPTG or 5 mM sodium butyrate for 24 h before analysis on 15% SDS gels followed by immunoblotting with the mAb 7.51,

which binds to an epitope in the repeat domain of tau. The SStau190-441 protein has an apparent RMM of ~40K on gels, and is degraded to a fragment of ~25K. In contrast the soluble tau186-441 fragment and soluble T40 show different degradation profiles. Induction of full-length tau (I) in cells expressing SStau190-441 causes increased appearance the 40K, 25K proteins, and possibly a small amount of a 30K fragment. Increasing expression of the transfected protein by addition of sodium butyrate (B) also increases the amount of both fragments, and induces the appearance of the 30K fragment.

In Figure 16(b) The SStau190-390 protein has an apparent RMM of ~30K, with a small amount of a degradation product at ~18K. Induction of full-length tau causes a slight increase in the amount of the 30K protein. Butyrate treatment causes a larger increase in the amount of the 30K fragment, and also increases the amount of the 18 kD fragment. This pattern of degradation is not seen for the soluble 186-391 fragment or the full length T40.

Figure 16 (c) shows an analysis of cells expressing tau fragments by immunoblotting. Cells stably expressing the tau fragment SStau296-390 and also expressing full-length tau under the control of an IPTG-inducible promoter, were incubated with 5 mM IPTG ("I") or 5 mM sodium butyrate ("B") for 24 h before analysis on 15% SDS gels followed by immunoblotting with mAb 7.51, which binds an epitope in the repeat domain of tau. The unlabelled lane shows only a minor band with gel mobility corresponding to 18 kD. The lane designated "B" shows a prominent gel band with mobility corresponding to 18 kD and a minor band corresponding to 14 kD. The lane designated "I" shows a minor band with gel mobility corresponding to 18 kD. The derivation of the 18 kD band in this cell system is explained in Figure 1(d). As the tau fragment SStau296-390 has gel mobility of ~14 kD, the 18 kD species cannot arise as a degradation product of the introduced tau fragment, nor is there evidence of formation of the 24 kD dimer seen e.g. in Figure 16(d).

Figure 16 (d) shows how the tau dGAE fragment (residues 297 - 391) expressed bacterially in vitro aggregates spontaneously to form a series of higher order oligomers, of which the major form has gel mobility ~24 kD. There are minor atypical oligomers with gel mobility ~19/22 kD, which are most likely to be partially degraded forms of the dimer.

Figure 16(e) left hand panel replicates the effect shown in (a), wherein IPTG gives rise to increased levels of 25K, 30K and 40K proteins, for comparison. The same experiment conducted in the presence of thionine blocks production of the 25K protein, and

also reduces production of the 30K protein. Thionine does not, however, reduce detected levels of the 25K and 30K bands seen with butyrate. Therefore thionine is shown to block the capture and processing of full length tau that gives rise to the 25K band.

Figure 17 (a) shows the proteolytic processing of SStau190-441. Cells expressing SStau190-441 and induced to express full-length tau by treatment with IPTG (right-hand lane) were analysed with four antibodies recognising epitopes located along the length of tau. The gel shows analysis using mAb 7.51, which recognises the repeat domain of tau, reveal all of the tau fragments which contain the core aggregation domain. The remaining columns show a summary of the reactivity of the mAb 7.51 positive fragments with three other antibodies. The results are also shown in a schematic representation of the fragments produced by processing of full-length tau and SStau190-441. Only full-length tau is recognised by all four antibodies. A fragment slightly larger than the SStau190-441 is produced by N-terminal truncation of full-length tau. Two fragments smaller than SStau190-441 are produced by either N-terminal or C-terminal processing. The two smallest fragments only react with the antibody to the core domain, demonstrating proteolytic stability of this domain. Figure 17 (b) shows the proteolytic processing of SStau190-390. Cells expressing SStau190-390 and treated with sodium butyrate (right-hand lane) produce SStau190-390 and a single proteolytic product of ~18 kD. As shown on the table and schematically at the right, the 18 kD species is immunoreactive only with mAb 7.51. The probable boundaries of the fragment are shown on the right, with reasons indicated described in Figure 1.

Figure 18 shows the sedimentation of tau aggregates from cell extracts. Cells expressing (a) SStau190-441 or (b) SStau 190-390, and induced to express full-length tau, were homogenised and spun at 720g for 3 min to produce a post-nuclear supernatant (pns). The pns was spun at 12,000g for 2 min to sediment membranes and large aggregates, and the supernatant from this spin was spun at 200,000g for 22 min. The same analysis was performed on pns that had been treated with 0.1% Triton X-100 to solubilise membrane proteins. Comparison of the 12K supernatants without and with detergent (lanes 3, 7) shows that more protein is present in the supernatant after detergent extraction. In both cases, a significant amount of protein also appears in the pellet (lanes 4,8). Centrifugation of the 12K supernatants at 200Kg sediments the remaining protein (lanes 6, 10).

Figure 19 (a) shows a conceptual scheme wherein critical nucleating factors provide a 'seed' which in-

itiates tau capture, which then becomes autocatalytic. Figure 19 (b) shows a putative pathogenic model of Alzheimer's disease. Tau aggregation is a proximal process prior to failure of axonal transport and consequent neuronal death. The tau aggregation cascade can be triggered either by a seeding/nucleation event arising from upstream changes or from primary mutations in the tau gene.

Figure 20 shows a table listing proteins which play a role in diseases of protein aggregation. Also listed are the diseases themselves, the aggregating domain and/or mutation believed to be involved, and the putative (maximum) fibril subunit size. One or more literature references for each protein is given.

Figure 21 Sequence analysis of proteins in the PHF-core preparation after removal of acid-soluble PHF-tau. The residue was solubilised according to the scrapie-fibril protocol of Stahl et al (1990, Biochemistry 29, 8879-8884). Yields are given for a typical preparation from a single AD brain. The protein fragments and corresponding residues identified as copurifying with PHF-core tau protein in bands of apparent gel mobility are shown in kD. The non-tau proteins copurifying in the form of SDS- and electrophoresis-resistant complexes are: ubiquitin, porin and core-protein-2 of mitochondrial complex III ("core-2"). ATP-synthase subunit 9 ("ATP-9") was found in a form which Co-purified with the A β protein of plaques. Porin sequences were also identified from Asp-N * and CNBr** digests of the 34 kD species.

Examples

Example 1 - production of membrane bound protein

Introduction

[0186] An mRNA for a protein that is normally translated on free ribosomes can be targeted to membrane-bound ribosomes by inclusion of a signal sequence (SS) of a secreted protein; the nature of the 3'UTR also influences the efficiency of the targeting (Partridge, K.A., et al. (1999) Cytotechnology 30, 37-47). A protein that is normally soluble can therefore become inserted into the membrane of the endoplasmic reticulum. The present inventors have used this approach to construct a membrane-associated form of tau that creates the localised concentration of tau needed to promote aggregation.

[0187] Cell lines stably expressing these membrane-associated forms of tau show evidence of tau aggregation and proteolytic processing by the criteria of histology, sedimentation of insoluble forms of tau protein from cell extracts and analysis of proteolytic products by immunoblotting.

Overview of expression system

[0188] The expression system used for production of membrane associated tau fragments is summarised in Figure 9. The system is based on the observation that rabbit globin mRNA is translated on membrane-bound ribosomes if the signal sequence of rat albumin is inserted at the 5' end of the cDNA (Partridge, K.A., *et al.*, *loc. cit.*). The vector previously described for expression of globin (Figure 9a) was modified to incorporate fragments of the tau sequence downstream of the signal sequence, while at the same time maintaining the globin 3'UTR (Figure 9b).

[0189] Three truncation mutants of tau (amino acids 296-390, 190-441 and 190-390), referred to as SStau296-390, SStau190-441 and SStau190-390, respectively, were expressed using this system. The structures of these mutants, together with full-length tau and the equivalent soluble tau fragments used as controls, are summarized in Figures 5 and 10. The proteins were expressed constitutively in 3T6 cells that also express the full-length tau sequence under the control of an IPTG-inducible promoter.

Construction of SStau constructs.

[0190] pKSSGG as shown in Figure 9a and described by Partridge et al (1999) *loc. cit.*

[0191] PCR-based mutagenesis on pKSSGG to introduce an AgeI site at the globin start codon to make pKSSGGAgeI.

forward primer: 5'-gccttttcaccgggtgcatctgtcca-3'

reverse primer: 5'-tggacagatgcaccgggtgaaaaggc-3'

htau40 cDNA (see Goedert et al, 1989 *Neurone* 3, 519-526).

pKSSStau190-441

[0192] PCR-based mutagenesis on htau40 to introduce an AgeI site at nucleotide 1128 of htau40 (amino acid 186). Fragment cut with EcoRI (cuts the tau sequence beyond the stop codon), blunted, then with AgeI, ligated into pKSSGGAgeI cut with BamHI, blunted, then AgeI to make pKSSStau190-441. The ligation destroys the tau sequence between amino acids 186-189, so amino acid 190 is the start of the tau sequence.

forward primer: 5'-tctggtgaaccggtaaaatcgggg-3'

reverse primer: 5'-ccccgtattttaccggaacaccaga-3'

pKSSStau190-390

[0193] PCR-based mutagenesis on pKSSStau190-441 to introduce a BamHI site and stop codon at nucle-

otide 1740 (terminates the tau sequence at amino acid 390). Fragment cut with AgeI and BamHI and ligated into pKSSGGAgeI at the same sites to make pKSSStau190-390.

forward primer: 5'-ctgtactcgagctacgccccgtgg-3'

reverse primer: 5'-cggggcgtaggataagtagaagtcg-3'

10 pKSSStau296-390

[0194] PCR-based mutagenesis on pKSSStau190-390 to introduce an AgeI site at nucleotide 1450 (amino acid 293) of pKSSStau190-390. Fragment cut with AgeI and BamHI and ligated into pKSSGGAgeI at the same sites to make pKSSStau290-390.

forward primer: 5'-ggctcaccgggttaatatcaaac-3'

reverse primer: 5'-gtttgatattaaccgggtgagcc-3'

[0195] These constructs are in the expression vector pcDNA3.1, which contains the G418 resistance marker. All were subcloned in the pcDNA3.1zeo vector, which contains the zeocin resistance marker, for transfection into the T40 inducible cell line.

Construction of T40 inducible construct

[0196] This was based on the Stratagene system (Catalog No. 217450) in which a p3'SS vector (constitutively expressing the *lac I* gene, and selected for using hygromycin) is used in conjunction with pOPRSVICAT (which includes the strong RSV promoter with operator sequences from the *Lac* operon, and is selected for with neomycin). Briefly, the two vectors are incorporated into cells, such as 3T3 or 3T6 fibroblasts. The repression of the operon by *lac I* is relieved by the addition of IPTG. To prepare the T40 construct, the tau protein fragment was cloned into the pOPRSVICAT vector (which contains a NotI site) according to the manufacturer's instructions.

Results

[0197] The cells were analyzed by immunofluorescence labeling using an antibody directed against the core aggregation domain, mAb 7.51 (Novak, M., et al. (1991) *Proc. Natl. Acad. Sci. USA* 88, 5837-5841). By way of control, stably transfected 3T3 cells expressing the soluble tau fragment 186-441 were fixed with two different fixation conditions (a) paraformaldehyde or (b) glutaraldehyde and Triton X-100 and labeled with mAb 7.51 directed against the repeat domain of tau. After paraformaldehyde fixation, labeling for tau was diffuse throughout the cytoplasm (s previously reported by Schliwa et al., 1981, *PNAS USA* 78:1037-1041), whereas cells fixed in glutaraldehyde and Triton X-100 show labeling of the tau along microtubules. This labeling pattern was

typical for soluble fragments of tau; evidence of tau aggregation is, however, not observed.

[0198] 3T6 Cells stably transfected and expressing the membrane-targeted minimum aggregation domain, SStau296-390, were incubated with 5 mM sodium butyrate for 20 h to increase protein expression and fixed in paraformaldehyde. These became labeled by mAb 7.51 throughout the cytoplasm in a reticular pattern consistent with association with the endoplasmic reticulum (not shown). However, this protein was only expressed to detectable levels in a small proportion of the cells and aggregation of the protein was rarely seen.

[0199] In contrast, labeling of cells stably transfected with and expressing SStau190-441 or SStau190-390 reveals large aggregates, some of which have a linear appearance. These aggregates were often clustered in the perinuclear area, and sometimes have a filamentous form (Figure 11).

[0200] The proportion of cells expressing these aggregates is increased by addition of sodium butyrate, which increases the expression level of the protein (Gorman, C.M., et al. (1983) Nuc. Acids Res. 11, 7631-7648; Kruh, J. (1982) Mol. Cell. Biochem. 42, 65-82), but aggregates can also be found in cells expressing the protein constitutively.

[0201] Cells stably transfected with and expressing SStau190-390 were also fixed in glutaraldehyde in the presence of 0.2% Triton X-100 before labeling with 7.51. This fixation protocol allows visualization of microtubule-bound tau protein, but aggregates of tau are also seen, demonstrating that they are stable to detergent extraction i.e. their stability *per se* is not dependent on membrane association (not shown).

[0202] Proteins expressed in these cell lines have been analyzed by immunoblotting with mAb 7.51, and the effect of tau induction and butyrate treatment examined, compared to cell lines expressing full-length tau and soluble fragments of tau.

[0203] The SStau190-441 protein analyzed by immunoblotting with the mAb 7.51 has an apparent RMM of ~40K and shows a characteristic degradation product at ~25K, and a less prominent product at ~30K. The expression level of the protein can be increased by treatment with sodium butyrate, which increases the amount of all three products (Figure 16). Butyrate also occasionally induces the appearance of two smaller products at ~18 and ~14K (see Figure 17).

[0204] Induction of full-length tau protein in the presence of SStau190-491 increases the amount of products at 30K, and 25K, but there is no appearance of smaller products (Figure 16a).

[0205] Thus, considering Figure 16a, when the effects of butyrate and IPTG induction are compared, butyrate is seen to act non-specifically to increase production of constitutively expressed incorporated gene products (by a mechanism which is not well understood). This serves as a control for IPTG induction, which specifically induces only full-length tau. In the first panel, the effects on ex-

pression of the non-membrane targeted 186-441 fragment after treatment with butyrate or IPTG are compared. Butyrate ("B") produces more of the 186-441 fragment, and a closely related presumed truncation product, but IPTG ("I") does not.

[0206] In the third panel, the same conditions are examined in cells simply expressing inducible T40 under IPTG control as described previously. Here, only IPTG ("I") produces increased levels of T40, and approx. 45 kD spontaneous degradation products. Butyrate has no effect.

[0207] The middle panel shows the effect of both induction methods in cells which express both SS 190-441 constitutively and T 40 under an IPTG-inducible promoter. Without induction, there are low levels of T40 and the characteristic 40 kD and 25 kD bands derived from SS190-441. Butyrate ("B") induces a higher level of the 40, 30 kD and 25 kD bands. Induction of T40 by IPTG ("I"), which has no effect on constitutive expression of the 40 kD band, generates higher levels of both the 40 kD and 25 kD species through an aggregation dependent mechanism (see Figure 1). Therefore, induced T40 is being processed through the 40 kD pathway to give rise to the 25 kD truncation product, provided there are pre-existing low levels of SS 190-441.

[0208] Considering Figure 16b, the SStau190-390 protein analyzed by immunoblotting with the mAb 7.51 has an apparent RMM of 30K and shows a degradation product at ~18K, which approximately corresponds to the size of the minimum aggregation domain of PHFs. Expression level of the protein and its characteristic 18 kD degradation product is substantially increased by treatment with sodium butyrate. Once again, the equivalent soluble tau fragment is not degraded in the same way and does not cause production of the 18 kD degradation product i.e. induction of full-length tau protein in the presence of SStau190-390 slightly increases the amount of 30K protein, but there is no increased appearance of the smaller product (Figure 16b).

[0209] In both cases, the degradation pattern seen for the membrane-targeted tau fragments is different from that seen for the equivalent soluble protein. The membrane-targeted proteins can be degraded to a ~12 \ 14 K fragment equivalent to the core domain of PHFs (see e.g. Fig 17(a)). The aggregation of the tau fragments induced by membrane association is therefore conferring proteolytic stability to the core domain, as seen in PHFs.

[0210] Full-length tau protein which is induced in the absence of a membrane-localised tau fragment shows some degradation to a fragment of ~45K, but there is no production of smaller fragments. The enhanced production of smaller tau fragments seen when full-length tau is expressed with the SStau190-441 (Figure 16a) demonstrates processing of tau induced by interaction with the SStau190-441 aggregates.

[0211] The degradation products produced in cells expressing SStau190-441 have been analyzed using antibodies specific for epitopes which are both NB and C-

terminal to the aggregation domain. This analysis shows that the protein undergoes both N- and C-terminal truncations (Figure 17). The smallest fragment produced only reacts with the antibody against the core domain of PHFs and is of a similar size (~12-14K). The tau protein in these cells, therefore, appears to be undergoing PHF-like processing to the core fragment.

[0212] Both of the SStau proteins, together with their degradation products, can be sedimented from cell extracts even in the presence of the detergent Triton X-100, suggesting that the protein is aggregated rather than being associated with membranes (Figure 18). A fraction of the protein sediments at relatively low speed (12,000g), and the remainder behaves as particles with an S value of at least 50S, but does not sediment at 300S, consistent with aggregation to a complex of MW ~1-2x10⁶.

[0213] These results indicate the following pathway of events: aggregation, which is initially seeded by the membrane-localised protein, occurs at the cytosolic surface of the ER. The protein is cleaved at its N-terminus to remove the signal sequence, and this form is then free to appear as aggregates in the cytosol or as a microtubule-bound protein if aggregation has not occurred. Membrane-targeting of the tau-like protein, as taught herein, facilitates the capture of full-length tau which is, for instance, produced under the control of an inducible promoter. Further full-length tau is then channeled into the same proteolytic processing pathway, as indicated by production of the 25kD unit after induction of full-length tau (Figure 1).

Conclusion

[0214] Expressing a fragment of tau as a membrane-localised protein has overcome the barrier to aggregation in cells which is found using soluble proteins. The SS190-390 fragment, for example, is efficiently processed to shorter fragments including the 12-14K fragment, which is similar to that found in the core of PHFs.

[0215] The inventors have thus discovered that a signal sequence can be used to create a membrane-localised form of tau, that tau fragments thus expressed as membrane-localised proteins will form aggregates and that these tau aggregates have a proteolytically-stable core which is comparable to that in PHFs, both in relation to its size and to the presence of the 7.51 epitope. The inventors have also identified that the appearance of processed fragments is increased after induction of full-length tau protein, proving that there is capture of soluble tau into aggregates.

[0216] Thus, tau aggregation and processing have been achieved in cells, and the aggregates can be analyzed in assays to test for compounds which inhibit or modulate the aggregation.

Example 2 - screening inhibitors

[0217] Cells expressing the SS190-441 fragment as

well as inducible T40 under IPTG control were plated in 6 well dishes. One day after plating experimental drugs were added at indicated concentrations and 9 h later, IPTG was added at 5mM. 15h after IPTG addition, cells were washed with PBS then solubilised in gel buffer. Samples were separated by SDS-PAGE and immunoblotted using antibody 7.51. T40 induction in these cells leads to an increase in degradation products (as shown in Figure 1, and Figure 16c). The effect of the experimental drugs on this increase was analysed by quantification of the immunoblots.

[0218] Figure 2 shows how the inhibitory activities of thionine, chlorpromazine and tacrine were compared in the membrane-targeted cell model. There is a clear distinction between active and inactive compounds. The data was analysed in greater detail mathematically using a standard inhibitory model (Figure 3).

[0219] The values which the model yields are consistent with those which would be expected for an *in vivo* system in which the tau was phosphorylated, giving a resulting reduction in its binding affinity, and a concomitant increase in the apparent inhibitory potency of tau aggregation inhibitors.

[0220] This demonstrates the utility of the present invention in screening for inhibitors of the aggregation phenomena of proteins which undergo an induced conformation polymerisation in neuropathological disease. In particular the aggregation can be modified by aggregation inhibitors and their activity can be monitored by production of characteristic degradation products.

Example 3 - preparation and use of transgenic animal

[0221] Transgenic animals were prepared as follows.

Animals:

[0222] To obtain a great number of fertilized oocytes, 4 week old NMRI mice are treated with gonadotrophins to induce superovulation. Pregnant mare's serum (PMS) is used to mimic follicle-stimulating hormone (FSH) and human chorionic gonadotropin (hCG) is used to mimic luteinizing hormone (LH). PMS is administered intraperitoneally 2 days before mating and hCG is administered 46 hours later. After administration of hCG the female mice are mated with stud males. On the same day 10 week old NMRI mice are mated with vasectomized males, to obtain foster mothers for the reimplantation of the injected oocytes. Fertilized oocytes are collected from mice with copulation plug the following morning.

Collection of oocytes:

[0223] The mice are sacrificed by cervical dislocation and the oviduct is dissected. Newly ovulated oocytes, surrounded by cumulus cells, are found in the upper part of the oviduct (ampulla). One oviduct at a time is transferred into a petridish containing M2 medium and hyal-

oronidase. With a pair of watchmakers forceps the oocytes can be released from the oviduct. The hyaluronidase separates the eggs from the cumulus cells. The oocytes are then separated from any other cells and material in two successive cleaning steps through M2 medium. Until needed the oocytes are stored in M16 medium at 37°C in a 5% CO₂ incubator.

[0224] When collected the oocytes are in an early pronuclear stage wherein the male and female pronuclei are separated and distinguishable from the cytoplasm.

Preparation of the DNA construct for microinjection (see Figure 6):

[0225] The constructs contain an appropriate cDNA under the regulation of the Thyl-promoter which directs transgene expression to the brain. The plasmids were extracted from the host bacteria and purified by Qiagen Plasmid Purification Kit (sold by Qiagen, Hilden, Germany). The constructs were linearized with an appropriate endonuclease (NotI). The desired fragment was isolated from agarose gel by using Qiaquick™ Gel Extraction Kit (sold by Qiagen, Hilden, Germany). The constructs were diluted to 2 ng/μl in microinjection buffer.

Microinjection:

[0226] One micromanipulator is mounted on each side of an inverted microscope stage; one micromanipulator controls the holding pipette and the other the injection pipette. The eggs are placed in a depression slide injection chamber with M2 medium covered with paraffin oil. One egg is positioned on the holding pipette and the male pronucleus is injected with the injection pipette containing the DNA solution. After injection the oocytes are again cultured in M16 medium at 37°C in a 5% CO₂ incubator until retransfer.

Retransfer:

[0227] Pseudopregnant mice are anesthetized and the intact injected oocytes are retransferred into the oviduct of the mouse (10-15 oocytes into one oviduct). These foster mothers carried the implanted embryos to term.

Identification of transgenic mice:

[0228] Two different approaches to identify transgenic animals were used.

PCR:

[0229] Transgenic mice were detected with PCR using transgene specific primers.

Thy 1F2: gCA ggA ggT gCT CAg ggA CAg c
TautxR: CAC TgC CgC CTC CCg ggA CgT g

[0230] This primer combination yields a 415 bp PCR fragment only in mice carrying the Thyl-sstau296-390 transgene and a 711bp PCR fragment in the mice carrying the Thyl-sstau186-441 transgene.

[0231] Genomic DNA was prepared from 0.5 cm mouse tail using DNeasy™ Tissue Kit (Qiagen, Hilden, Germany)

Southern Blot

[0232] 10 μg genomic DNA of PCR positive tested animals were digested with EcoRI and electrophoresed on an 1% agarose gel. Southern blots were prepared and hybridized with 2 ³²P-labeled fragments from the promoter. These were obtained by digestion with XbaI and NcoI. The endogenous signal can be compared with the transgene signal to estimate the copy number.

[0233] In order to test the mode of integration the genomic DNA was digested with XbaI. Founder 1 and F1 animals showed single integration. Founder 86 and F1 animals showed double integration and could be split into two independent lines.

[0234] Blots were hybridized at 65°C for 20 hours and washed at 65°C 2x 1 hour.

RNA expression

[0235] To study RNA expression Northern blot analysis has been done. RNA was prepared of whole brain using Trizol™ (Gibco BRL, Paisley, Scotland). 20 μg RNA was electrophoresed and blotted. Hybridisation was performed against a 247bp PCR fragment. Primers were transgene specific: glob578-597 (AAG AAC AAT CAA ggg T CC CC), oligo20 (ATT Agg CAA CAT CCA TCA TAA ACC).

[0236] Blots were hybridized at 65°C for 20 hours and washed at 65°C 2x 1 hour.

[0237] Figures 12-15 show the results obtained with the mouse model and describe how an 18 kD species is generated by aggregation-dependent proteolytic processing of endogenous mouse brain tau through a pathway believed to correspond to that given in Figure 1 (E).

References

[0238]

Arendt et al. (1995) Neuroscience 69, 691-698.

Braak et al. (1994) Neuroscience letters 171, 1-4.

Brion et al. (1999) American Journal of Pathology 154, 255.270.

Cork et al. (1988) Journal of Neuropathology and Experimental Neurology 49, 629-641.

Duff et al. (2000) *Neurobiology of Disease* 7, 87-98.

Geula et al. (1998) *Nature Medicine* 4, 827-834.

Gotz et al. (1995) *EMBO Journal* 14, 1304-1313.

Harada et al. (1994) *Nature* 369, 488-491.

Ikegami et al. (2000) *Neuroscience letters* 279, 129-132.

Ishihara et al. (1999) *Neuron* 24, 751-762.

Janus et al. (2000) *Biochimica et Biophysica Acta* 1502, 63-75.

lewis et al. (2000) *Nature Genetics* 25, 402-405.

Nelson et al. (1995) *Neurobiology of Aging* 16, 315-323.

Roertgen et al. *Neurobiology of Aging* 17, 243-247.

Schultz et al. (2000) *Journal of Neuropathology and Experimental Neurology* 59, 39-52.

Spittaels et al. (1999) *American Journal of Pathology* 155, 2153-2165.

Sturchler-Pierrat et al. (1997) *Proceedings of the National Academy of Sciences, USA* 93, 13287-13292.

Thuncke et al. (2000) unpublished.

Additional References

[0239]

Abrahamson, M., Jonsdottir, S., Olafsson, I. & Grubb, A. (1992) Hereditary cystatin C amyloid angiopathy identification of the disease-causing mutation and specific diagnosis by polymerase chain reaction based analysis. *Human Genetics* 89, 377-380.

Booth, D.R., Sunde, M., Bellotti, V., Robinson, C.V., Hutchinson, W.L., Fraser, P.E., Hawkins, P.N., Dobson, C.M., Radford, S.E., Blake, C.C.F. & Pepys, M.B. (1997) Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis. *Nature* 385, 787-793.

Carrell, R.W. & Gooptu, B. (1998) Conformational changes and disease - serpins, prions and Alzheimer's. *Current Opinion in Structural Biology* 8, 799-809.

Chiti, F., Webster, P., Taddei, N., Clark, A., Stefani,

M., Ramponi, G. & Dobson, C. (1999) Designing conditions for in vitro formation of amyloid protofilaments and fibrils. *Proceedings of the National Academy of Sciences, USA* 96, 3590-3594.

Czech, C., Tremp, G. & Pradier, L. (2000) Presenilins and Alzheimer's disease: biological functions and pathogenic mechanisms. *Progress in Neurobiology* 60, 363-384.

Davis, R.L., Shrimpton, A.E., Holohan, P.D., Bradshaw, C., Feiglin, D., Collins, G.H., Sonderegger, P., Kinter, J., Becker, L.M., Lachawan, F., Krasnewich, D., Muenke, M., Lawrence, D.A., Yerby, M.S., Shaw, C.-M., Gooptu, B., Elliott, P.R., Finch, J.T., Carrell, R.W. & Lomas, D.A. (1999) Familial dementia caused by polymerization of mutant neuroserpin. *Nature* 401, 376-379.

DiFiglia, M., Sapp, E., Chase, K.O., Davies, S.W., Bates, G.P., Vonsattel, J.P. & Aronin, N. (1997) Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. *Science* 277, 1990-1993.

Dische, F.E., Wernstedt, C., Westermark, G.T., Westermark, P., Pepys, M.B., Rennie, J.A., Gilbey, S.G. & Watkins, P.J. (1988) Insulin as an amyloid-fibril protein at sites of repeated insulin injections in a diabetic patient. *Diabetologia* 31, 158-161.

Gasset, M., Bladwin, M.A., Lloyd, D.H., Abriel, J.-M., Holtzman, D.M., Cohen, F.E., Fletterick, R. & Prusiner, S.B. (1992) Predicted α -helical region of the prion protein when synthesized as peptides form amyloid. *Proceedings of the National Academy of Sciences, USA* 89, 10940-10944.

Glennner, G.G. & Wong, C.W. (1984) Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochemical and Biophysical Research Communications* 120, 885-890.

Goate, A., Chartier-Harlin, M.-C., Mullan, M., Brown, J., Crawford, F., Fidani, L., Giuffra, L., Haynes, A., Irving, N., James, L., Mant, R., Newton, P., Rooke, K., Roques, P., Talbot, C., Pericak-Vance, M., Roses, A., Williamson, R., Rossor, M., Owen, M. & Hardy, J. (1991) Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349, 704-706.

Gorevic, P.D., Casey, T.T., Stone, W.J., DiRaimondo, C.R., Prelli, F.C. & Frangione, B. (1985) b-2 Microglobulin is an amyloidogenic protein in man. *Journal of Clinical Investigation* 76, 2425-2429.

- Gustavsson, A., Engstrom, U. & Westermark, P. (1991) Normal transthyretin and synthetic transthyretin fragments form amyloid-like fibrils in vitro. *Biochemical and Biophysical Research Communications* 175, 1159-1164.
- Hutton, M., Lendon, C., Rizzu, P., Baker, M., Froelich, S., Houlden, H., Pickering-Brown, S., Chakraborty, S., Isaacs, A., Grover, A., Hackett, J., Adamson, J., Lincoln, S., Dickson, D., Davies, P., Petersen, R.C., Stevens, M., de Graaf, E., Wauters, E., van Baren, J., Hillebrand, M., Joosse, M., Kwon, J.M., Nowotny, P., Che, L.K., Norton, J., Morris, J.C., Reed, L.A., Trojanowski, J.Q., Basun, H., Lannfelt, L., Neystat, M., Fahn, S., Dark, F., Tannenberg, T., Dodd, P.R., Hayward, N., Kwok, J.B.J., Schofield, P.R., Andreadis, A., Snowden, J., Craufurd, D., Neary, D., Owen, F., Oostra, B.A., Hardy, J., Goate, A., van Swieten, J., Mann, D., Lynch, T. & Heutink, P. (1998) Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 393, 702-705.
- Johansson, B., Wernstedt, C. & Westermark, P. (1987) Atrial natriuretic peptide deposited as atrial amyloid fibrils. *Biochemical and Biophysical Research Communications* 148, 1087-1092.
- Lomas, D.A., Evans, D.L., Finch, J.T. & Carrell, R.W. (1992) The mechanism of α_1 -antitrypsin accumulation in the liver. *Nature* 357, 605-607.
- Maury, C.P. & Baumann, M. (1990) Isolation and characterization of cardiac amyloid in familial amyloid polyneuropathy type IV (Finnish): relation of the amyloid protein to variant gelsolin. *Biochimica et Biophysica Acta* 1096, 84-86.
- Paulson, H.L. (1999) Human genetics '99: trinucleotide repeats. *American Journal of Human Genetics* 64, 339-345.
- Pepys, M.B., Hawkins, P.N., Booth, D.R., Vigushin, D.M., Tennent, G.A., Soutar, A.K., Totty, N., Nguyen, O., Blake, C.C.F., Terry, C.J., Feast, T.G., Zalin, A.M. & Hsuan, J.J. (1993) Human lysozyme gene mutations cause hereditary systemic amyloidosis. *Nature* 362, 553-557.
- Polymeropoulos, M.H., Lavedan, C., Leroy, E., Ide, S.E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., Stenroos, E.S., Chandrasekharappa, S., Athanassiadou, A., Papaetropoulos, T., Johnson, W.G., Lazzarini, A.M., Duvoisin, R.C., Di Iorio, G., Golbe, L.I. & Nussbaum, R.L. (1997) Mutation in the α -synuclein gene identified in families with Parkinson's disease. *Science* 276, 2045-2047.
- Prusiner, S.B., Scott, M.R., DeArmond, S.J. & Cohen, F.E. (1998) Prion protein biology. *Cell* 93, 337-348.
- Shibata, N., Hirano, A., Kobayashi, M., Siddique, T., Deng, H.X., Hung, W.Y., Kato, T. & Asayama, K. (1996) Intense superoxide dismutase-1 immunoreactivity in intracytoplasmic hyaline inclusions of familial amyotrophic lateral sclerosis with posterior column involvement. *Journal of Neuropathology and Experimental Neurology* 55, 481-490.
- Sletten, K., Westermark, P. & Natvig, J.B. (1976) Characterization of amyloid fibril proteins from medullary carcinoma of the thyroid. *Journal of Experimental Medicine* 143, 993-998.
- Spillantini, M.G., Crowther, R.A., Jakes, R., Hasegawa, M. & Goedert, M. (1998) α -Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. *Proceedings of the National Academy of Sciences, USA* 95, 6469-6473.
- Uemichi, T., Liuepnicks, J.J. & Benson, M.D. (1994) Hereditary renal amyloidosis with a novel variant fibrinogen. *Journal of Clinical Investigation* 93, 731-736.
- Westermark, P., Engstrom, U., Johnson, K.H., Westermark, G.T. & Betsholtz, C. (1990) Islet amyloid polypeptide: pinpointing amino acid residues linked to amyloid fibril formation. *Proceedings of the National Academy of Sciences, USA* 87, 5036-5040.
- Westermark, P., Johnson, K.H., O'Brien, T.D. & Betsholtz, C. (1992) Islet amyloid polypeptide - a novel controversy in diabetes research. *Diabetologia* 35, 297-303.
- Westermark, P., Johnson, K.H. & Pitkanen, P. (1985) Systemic amyloidosis: A review with emphasis on pathogenesis. *Applied Physiology* 3, 55-68.
- Wischik, C.M., Novak, M., Thogersen, H.C., Edwards, P.C., Runswick, M.J., Jakes, R., Walker, J.E., Milstein, C., M., R. & Klug, A. (1988) Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer's disease. *Proceedings of the National Academy of Sciences, USA* 85, 4506-4510.

Claims

1. A method of inducing or modeling the pathological state of an aggregating disease protein (ADP) which is associated with a disease state in which the ADP

- aggregates pathologically through an induced conformational polymerisation interaction, the method being **characterised by** the step of providing a membrane-localisable fusion protein comprising (i) an aggregating portion, which is derived from the ADP
- (ii) a heterologous membrane-localising portion, wherein said method is performed *in vitro*, or is performed in a cultured cell or cell line, or is performed in a non-human animal.
2. A method as claimed in claim 1 wherein the aggregating portion of the fusion protein is derived from the ADP, and wherein membrane-localisation of the fusion protein causes the high-affinity capture site of the ADP protein to become exposed such that aggregation of further ADP is promoted, and pathological aggregation of the ADP in the disease state leads to proteolytic processing of the ADP to a core domain fragment.
 3. A method as claimed in claim 1 or claim 2 comprising the steps of:
 - (a) providing the fusion protein
 - (b) introducing it into a membrane.
 4. A method as claimed in any one of the preceding claims wherein the aggregation of the ADP in the disease state is associated with neurodegeneration and/or clinical dementia
 5. A method as claimed in claim 4 wherein the aggregating portion comprises, or consists essentially of, at least the core fragment of the ADP.
 6. A method as claimed in any one of the preceding claims wherein the membrane localising portion includes a membrane anchor sequence and/or a stop-transfer sequence.
 7. A method as claimed in any one of the preceding claims wherein the disease is selected from Pick's disease, Progressive Supranuclear Palsy, Alzheimer's disease.
 8. A method as claimed in any one of the preceding claims wherein the ADP is tau protein.
 9. A method as claimed in claim 8 wherein the aggregating portion of the fusion protein comprises a core fragment of tau, which fragment of tau extends from amino acids 186-296 to 390-441 of the full-length tau protein shown in Fig 5.
 10. A method as claimed in claim 9 wherein the aggregating portion of the fusion protein consists of amino acid residues 296-390, 190-441 or 190-390 of the full-length protein.
 11. A method as claimed in any one of claims 2 to 10 for screening for a specific binding partner of the ADP high-affinity capture site, the method comprising the steps of:
 - (a) providing the fusion protein in membrane localised form,
 - (b) contacting the fusion protein with one or more potential binding partners,
 - wherein said contacting step is intracellular and the binding partner is present within the cell,
 - (c) determining whether or not the potential binding partner binds to the fusion protein
 12. A method as claimed in any one of claims 1 to 10 for inducing aggregation of ADP molecules, the method comprising the steps of:
 - (a) providing the fusion protein in membrane localised form,
 - (b) causing or permitting an induced conformational polymerisation interaction between the fusion protein and a further ADP molecule which is ADP or a derivative thereof, and
 - (c) monitoring extent of aggregation of the further ADP molecule.
 13. A method as claimed in claim 12 wherein the membrane is a cell membrane and the aggregation occurs intracellularly
 14. A method as claimed in claim 13 wherein step (a) consists either of:
 - introducing the fusion protein into a cell, such that it becomes membrane localised at a membrane in the cell, or
 - introducing nucleic acid encoding the fusion protein into a cell or an ancestor thereof, such that the fusion protein is expressed in the cell and becomes membrane localised at a membrane in the cell.
 15. A method as claimed in claim 14 wherein the further ADP molecule is full length, soluble ADP which is native to the cell
 16. A method as claimed in claim 13 or claim 14 wherein the further ADP molecule is heterologous to the cell and the method comprises the step of introducing a nucleic acid encoding the further ADP molecule into the cell or an ancestor thereof, such that the further ADP molecule is expressed in the cell.
 17. A method as claimed in claim 16 wherein the fusion protein is expressed under the transcriptional control

of a constitutive promoter, and the further ADP molecule is expressed under the transcriptional control of an inducible promoter in the presence of the appropriate inducing agent.

18. A method as claimed in claim 16 or claim 17 wherein the further ADP molecule is an ADP derivative which is (i) a fragment the ADP, (ii) a mutant of the ADP, or (iii) full length, soluble ADP.
19. A method as claimed in claim 16 or claim 18 wherein the further ADP molecule incorporates mutations into the ADP corresponding to those known or suspected to be associated with the disease state.
20. A method as claimed in any one of claims 12 to 19 wherein the aggregation is monitored by monitoring the concentration or level of any one or more of the following species: (i) non-aggregated fusion protein and/or further ADP molecule, (ii) an aggregate of the fusion protein and/or further ADP molecule, (iii) a proteolytic fragment of the fusion protein and/or further ADP molecule resulting from aggregation thereof.
21. A method as claimed in claim 20 wherein the ADP is tau, and the aggregation is monitored on the basis of increasing levels of an approximately 18 or 25 kDa fragment of tau.
22. A method as claimed in claim 20 or claim 21 wherein the monitoring step comprises use of an antibody specific for any of (i) the fusion protein (ii) the further ADP molecule (iii) a proteolytic fragment of the fusion protein and/or further ADP molecule resulting from aggregation thereof.
23. A method as claimed in claim 22 wherein fusion protein is immunologically distinct from the further ADP protein, and the ADP is tau, and the antibody is selected from a monoclonal antibody which (i) is specific for a human-specific epitope located in the region between Gly-16 and Gln-26 of tau; (ii) is specific for the core tau fragment truncated at Glu-391; (iii) is specific for a generic tau epitope in the repeat domain; or (iv) is specific for a non-species specific generic tau epitope located between Ser-208 and Ser-238.
24. A method for identifying a modulator of aggregation of an aggregating disease protein (ADP) which is associated with a disease state in which the ADP aggregates pathologically through an induced conformational polymerisation interaction, which method comprises
 - (a) performing a method as claimed in any one of claims 12 to 23 in the presence of an agent

suspected of being capable of modulating the aggregation,
(b) monitoring the extent of aggregation, and
(c) correlating the extent of aggregation with the modulatory activity of the agent.

5

10

15

20

25

30

35

40

45

50

55

25. A method as claimed in claim 24 which is performed intracellularly
26. A method as claimed in claim 25 wherein the cell is from a neuronal cell line or a fibroblast cell line.
27. A method as claimed in claim 25 wherein the cell is comprised within a non-human transgenic animal.
28. An isolated nucleic acid molecule comprising a chimeric nucleotide sequence encoding a membrane-localisable fusion protein having (i) an aggregating portion, which is derived from an aggregating disease protein (ADP) which is associated with a disease state in which the ADP aggregates pathologically through an induced conformational polymerisation interaction (ii) a heterologous membrane-localising portion
29. A nucleic acid as claimed in claim 28 wherein the chimeric nucleotide sequence includes an exogenous 3'UTR sequence, and wherein the membrane-localising portion comprises a signal sequence and optionally sequence encoding a membrane anchor domain or a stop-transfer sequence
30. A nucleic acid as claimed in claim 28 or claim 29 wherein the aggregating portion nucleotide sequence encodes tau, or a core fragment of tau.
31. A nucleic acid as claimed in claim 30 wherein the aggregating portion nucleotide sequence encodes a truncated fragment of tau extending from between amino acids 186-296 to 390-441 of the full-length protein.
32. A nucleic acid as claimed in claim 31 wherein the aggregating portion nucleotide sequence encodes amino acids 296-390, 190-441 or 190-390.
33. A process for producing a nucleic acid as claimed in any one of claims 28 to 32 comprising the step of ligating a nucleic acid encoding the aggregating portion to a nucleic acid encoding the heterologous membrane-localising portion, and wherein the nucleic acid encoding the heterologous membrane-localising portion is part of a recombinant vector into which the nucleic acid encoding the aggregating portion is inserted.
34. A recombinant vector which comprises the nucleic

acid of any one of claims 28 to 32 wherein the nucleic acid is operably linked to a promoter for transcription in a host cell.

35. A vector as claimed in claim 34 wherein the promoter selected from: a tissue specific promoter; the prion specific promoter; or the neurospecific enolase promoter. 5
36. A method which comprises the step of introducing the vector of claim 34 or claim 35 into a host cell. 10
37. A host cell containing or transformed with a heterologous vector of claim 34 or claim 35. 15
38. A method of preparing a transgenic non-human animal model with enhanced or accelerated pathology for the disease state in which an aggregating disease protein (ADP) aggregates pathologically, which method comprises the step of introducing into a fertilized egg of the animal a vector as claimed any one of claim 34 or claim 35. 20
39. A transgenic non-human animal which includes within a plurality of its cells a promoter operably linked to a nucleotide sequence encoding a membrane-localisable fusion protein comprising (i) an aggregating portion, which is derived from an aggregating disease protein (ADP) which is associated with a disease state in which the ADP aggregates pathologically through an induced conformational polymerisation interaction (ii) a heterologous membrane-localising portion. 25 30
40. A cell or tissue sample of the animal of claim 39 comprising a promoter operably linked to a nucleotide sequence encoding the membrane-localisable fusion protein. 35
41. A method, animal or sample as claimed in any one of claims 38 to 40 wherein the animal is a rodent. 40
42. An isolated fusion protein which is encoded by the chimeric nucleotide sequence of any one of claims 28 to 32. 45
43. A method of making the protein of claim 42, which method comprises the step of causing or allowing expression from a nucleic acid of any one of claims 28 to 32 in a suitable host cell. 50
44. A method of inducing in the transgenic non-human animal of claim 39 a disease state in which the aggregating disease protein (ADP) aggregates pathologically, which method comprises causing or allowing expression of the nucleotide sequence encoding the membrane-localisable fusion protein. 55

45. A method for screening for an agent for use as a therapeutic, prognostic or diagnostic agent in a disease state in which an aggregating disease protein (ADP) aggregates pathologically, comprising carrying out a method as claimed in claim 44 in the presence of an agent suspected of being capable of modulating the aggregation.
46. A method as claimed in claim 27 or claim 45 wherein the agent is capable of crossing the blood-brain barrier.
47. A method as claimed in claims 27, 45 or 46 wherein any of the following is determined within the brain of the animal:
 - (i) the extent of proteolytic processing of the ADP,
 - (ii) the ability of the agent to decrease the amount of ADP aggregates which form,
 - (iii) the ability of the agent to eliminate or reduce the level of ADP aggregate already formed.
48. A method as claimed in any one of claims 44 to 47 wherein the ADP is tau and the disease state corresponds to Alzheimer's disease.
49. An animal as claimed in claim 39 wherein the ADP is tau and the disease state corresponds to Alzheimer's disease.

Patentansprüche

1. Verfahren zur Induktion oder Modellierung des pathologischen Zustands eines aggregierenden Krankheitsproteins (ADP), das mit einem Krankheitszustand assoziiert ist, in dem das ADP durch eine induzierte Konformationspolymerisationswechselwirkung pathologisch aggregiert, wobei das Verfahren durch den Schritt des Bereitstellens eines membranlokalisierbaren Fusionsproteins gekennzeichnet ist, das (i) einen aggregierenden Abschnitt, der von dem ADP stammt, und (ii) einen heterologen Membranlokalisierungsabschnitt umfasst, worin das Verfahren *in vitro* durchgeführt wird oder in einer kultivierten Zelle oder Zelllinie durchgeführt wird oder ein einem nichtmenschlichen Tier durchgeführt wird.
2. Verfahren nach Anspruch 1, worin der aggregierende Abschnitt des Fusionsproteins von dem ADP stammt und worin die Membranlokalisierung des Fusionsproteins dazu führt, dass die hochaffine Einfangstelle des ADP-Proteins freigelegt wird, so dass die Aggregation weiterer ADP gefördert wird, und die pathologische Aggregation des ADP im Krankheitszustand zu proteolytischer Prozessierung des

ADP zu einem Kerndomänenfragment führt.

3. Verfahren nach Anspruch 1 oder Anspruch 2, das die folgenden Schritte umfasst:

- (a) das Bereitstellen des Fusionsproteins,
(b) dessen Einführung in eine Membran.

4. Verfahren nach einem der vorangegangenen Ansprüche, worin die Aggregation des ADP im Krankheitszustand mit Neurodegeneration und/oder klinischer Demenz assoziiert ist.

5. Verfahren nach Anspruch 4, worin der aggregierende Abschnitt zumindest das Kernfragment des ADP umfasst oder im Wesentlichen daraus besteht.

6. Verfahren nach einem der vorangegangenen Ansprüche, worin der Membranlokalisierungsabschnitt eine Membranankersequenz und/oder eine Stop-Transfer-Sequenz umfasst.

7. Verfahren nach einem der vorangegangenen Ansprüche, worin die Krankheit aus der Pick-Krankheit, progressiver supranukleärer Blickparese und der Alzheimer-Krankheit ausgewählt ist.

8. Verfahren nach einem der vorangegangenen Ansprüche, woran das ADP Tau-Protein ist.

9. Verfahren nach Anspruch 8, worin der aggregierende Abschnitt des Fusionsproteins ein Kernfragment von Tau umfasst, wobei sich das Fragment von Tau von den Aminosäuren 186-296 bis 390-441 des in Fig. 5 gezeigten Tau-Proteins voller Länge erstreckt.

10. Verfahren nach Anspruch 9, worin der aggregierende Abschnitt des Fusionsproteins aus den Aminosäureresten 296-390, 190-441 oder 190-390 des Proteins voller Länge besteht.

11. Verfahren nach einem der Ansprüche 2 bis 10 zum Screenen auf einen spezifischen Bindungspartner der hochaffinen ADP-Einfangstelle, wobei das Verfahren die folgenden Schritte umfasst:

- (a) das Bereitstellen des Fusionsproteins in membranlokalisierter Form,
(b) das Kontaktieren des Fusionsproteins mit einem oder mehreren möglichen Bindungspartnern, worin der Schritt des Kontaktierens intrazellulär erfolgt und der Bindungspartner in der Zelle vorhanden ist,
(c) das Bestimmen, ob der mögliche Bindungspartner an das Fusionsprotein bindet oder nicht.

12. Verfahren nach einem der Ansprüche 1 bis 10 zur

Induktion der Aggregation von ADP-Molekülen, wobei das Verfahren die folgenden Schritte umfasst:

- (a) das Bereitstellen des Fusionsproteins in membranlokalisierter Form,
(b) das Verursachen oder Zulassen einer induzierten Konformationspolymerisationswechselwirkung zwischen dem Fusionsprotein und einem weiteren ADP-Molekül, bei dem es sich um ADP oder um ein Derivat davon handelt, und
(c) das Überwachen des Ausmaßes von Aggregation des weiteren ADP-Moleküls.

13. Verfahren nach Anspruch 12, worin die Membran eine Zellmembran ist und die Aggregation intrazellulär erfolgt.

14. Verfahren nach Anspruch 13, worin Schritt (a) aus einem der beiden folgenden Schritte besteht:

- dem Einführen des Fusionsproteins in eine Zelle, so dass es auf einer Membran in der Zelle membranlokalisiert wird, oder
dem Einführen von Nucleinsäure, die für das Fusionsprotein kodiert, in eine Zelle oder einen Vorläufer davon, so dass das Fusionsprotein in der Zelle exprimiert wird und auf einer Membran in der Zelle membranlokalisiert wird.

15. Verfahren nach Anspruch 14, worin das weitere ADP-Molekül lösliches ADP voller Länge ist, das in der Zelle nativ ist.

16. Verfahren nach Anspruch 13 oder Anspruch 14, worin das weitere ADP-Molekül heterolog für die Zelle ist und das Verfahren den Schritt des Einführens einer Nucleinsäure, die für das weitere ADP-Molekül kodiert, in die Zelle oder einen Vorläufer davon umfasst, so dass das weitere ADP-Molekül in der Zelle exprimiert wird.

17. Verfahren nach Anspruch 16, worin das Fusionsprotein unter der Transkriptionskontrolle eines konstitutiven Promotors exprimiert wird und das weitere ADP-Molekül unter der Transkriptionskontrolle eines induzierbaren Promotors in Gegenwart des entsprechenden Induktionsmittels exprimiert wird.

18. Verfahren nach Anspruch 16 oder Anspruch 17, worin das weitere ADP-Molekül ein ADP-Derivat ist, wobei es sich um (i) ein Fragment des ADP, (ii) eine Mutante des ADP oder (iii) lösliches ADP voller Länge handelt.

19. Verfahren nach Anspruch 16 oder Anspruch 18, worin das weitere ADP-Molekül Mutationen in das ADP einführt, die jenen entsprechen, die bekanntermaßen oder vermutlich mit dem Krankheitszustand as-

soziiert sind.

20. Verfahren nach einem der Ansprüche 12 bis 19, worin die Aggregation durch Überwachung der Konzentration oder der Menge einer oder mehrerer der folgenden Spezies überwacht wird: (i) des nichtaggregierten Fusionsproteins und/oder des weiteren ADP-Moleküls, (ii) eines Aggregats des Fusionsproteins und/oder des weiteren ADP-Moleküls, (iii) eines proteolytischen Fragments des Fusionsproteins und/oder des weiteren ADP-Moleküls, das aus dessen Aggregation hervorgeht. 5
21. Verfahren nach Anspruch 20, worin das ADP Tau ist und die Aggregation anhand zunehmender Mengen eines etwa 18 oder 25 kDa großen Fragments von Tau überwacht wird. 10
22. Verfahren nach Anspruch 20 oder Anspruch 21, worin der Schritt des Überwachens die Verwendung eines Antikörpers umfasst, der für Folgendes spezifisch ist: (i) das Fusionsprotein, (ii) das weitere ADP-Molekül, (iii) ein proteolytisches Fragment des Fusionsproteins und/oder des weiteren ADP-Moleküls, das aus dessen Aggregation hervorgeht. 15
23. Verfahren nach Anspruch 22, worin das Fusionsprotein sich immunologisch von dem weiteren ADP-Protein unterscheidet und das ADP Tau ist und der Antikörper aus (i) einem monoklonalen Antikörper, der für ein für Menschen spezifisches Epitop spezifisch ist, das in der Region zwischen Gly-16 und Gln-26 von Tau liegt, (ii) einem, der für das Kern-Tau-Fragment spezifisch ist, das an Glu-391 trunkiert ist, (iii) einem, der für ein allgemeines Tau-Epitop in der Wiederholungsdomäne spezifisch ist, und (iv) einem, der für ein nicht speziesspezifisches, allgemeines Tau-Epitop spezifisch ist, das zwischen Ser-208 und Ser-238 liegt, ausgewählt ist. 20
24. Verfahren zur Identifizierung eines Modulators der Aggregation eines aggregierenden Krankheitsproteins (ADP), das mit einem Krankheitszustand assoziiert ist, in dem das ADP durch eine induzierte Konformationspolymerisationswechselwirkung pathologisch aggregiert, wobei das Verfahren Folgendes umfasst: 25
- (a) das Durchführen eines Verfahrens nach einem der Ansprüche 12 bis 23 in Gegenwart eines Mittels, das vermutlich in der Lage ist, die Aggregation zu modulieren, 30
 - (b) das Überwachen des Ausmaßes der Aggregation und
 - (c) das Korrelieren des Ausmaßes der Aggregation mit der Modulierungsaktivität des Mittels. 35

25. Verfahren nach Anspruch 24, das intrazellulär

durchgeführt wird.

26. Verfahren nach Anspruch 25, worin die Zelle von einer neuronalen Zelllinie oder einer Fibroblastenzelllinie stammt. 40
27. Verfahren nach Anspruch 25, worin die Zelle in einem transgenen nichtmenschlichen Tier enthalten ist. 45
28. Isoliertes Nucleinsäuremolekül, umfassend eine chimäre Nucleotidsequenz, die für ein membranlokalisierbares Fusionsprotein mit (i) einem aggregierenden Abschnitt, der von einem aggregierenden Krankheitsprotein (ADP) stammt, das mit einem Krankheitszustand assoziiert ist, in dem das ADP durch eine induzierte Konformationspolymerisationswechselwirkung pathologisch aggregiert, und (ii) einem heterologen Membranlokalisierungsabschnitt kodiert. 50
29. Nucleinsäure nach Anspruch 28, worin die chimäre Nucleotidsequenz eine exogene 3'UTR-Sequenz umfasst und 55
- worin der Membranlokalisierungsabschnitt eine Signalsequenz und gegebenenfalls eine für eine Membranankerdomäne kodierende Sequenz oder eine Stop-Transfer-Sequenz umfasst.
30. Nucleinsäure nach Anspruch 28 oder Anspruch 29, worin die Nucleotidsequenz für das aggregierende Protein für Tau oder ein Kernfragment von Tau kodiert. 60
31. Nucleinsäure nach Anspruch 30, worin die Nucleotidsequenz für das aggregierende Protein für ein trunkiertes Fragment von Tau kodiert das sich von zwischen den Aminosäuren 186 und 296 bis 390 bis 441 des Proteins voller Länge erstreckt. 65
32. Nucleinsäure nach Anspruch 31, worin die Nucleotidsequenz für das aggregierende Protein für die Aminosäuren 296 bis 390, 190 bis 441 oder 190 bis 390 kodiert. 70
33. Verfahren zur Herstellung einer Nucleinsäure nach einem der Ansprüche 28 bis 32, umfassend den Schritt des Ligierens einer Nucleinsäure, die für den aggregierenden Abschnitt kodiert, an eine Nucleinsäure, die für den heterologen Membranlokalisierungsabschnitt kodiert, 75
- worin die Nucleinsäure, die für den heterologen Membranlokalisierungsabschnitt kodiert, Teil eines rekombinanten Vektors ist, in den die für den aggregierenden Abschnitt kodierende Nucleinsäure inseriert wird.
34. Rekombinanter Vektor, der eine Nucleinsäure nach

einem der Ansprüche 28 bis 32 umfasst, worin die Nucleinsäure mit einem Promotor zur Transkription in einer Wirtszelle operabel verbunden ist.

35. Vektor nach Anspruch 34, worin der Promotor aus einem gewebespezifischen Promotor, dem prionenspezifischen Promotor und dem neurospezifischen Enolasepromotor ausgewählt ist. 5
36. Verfahren, umfassend den Schritt des Einführens eines Vektors nach Anspruch 34 oder Anspruch 35 in eine Wirtszelle. 10
37. Wirtszelle, die einen heterologen Vektor nach Anspruch 34 oder Anspruch 35 enthält oder damit transformiert ist. 15
38. Verfahren zur Herstellung eines transgenen nicht-menschlichen Tiermodells mit verstärkter oder beschleunigter Pathologie für den Krankheitszustand, in dem ein aggregierendes Krankheitsprotein (ADP) pathologisch aggregiert, wobei das Verfahren den Schritt des Einführens eines Vektors nach einem der Ansprüche 34 und 35 in ein befruchtetes Ei des Tieres umfasst. 20 25
39. Transgenes nichtmenschliches Tier, das innerhalb der Vielzahl seiner Zellen einen Promotor umfasst, der an eine Nucleotidsequenz operabel gebunden ist, die für ein membranlokalisierbares Fusionsprotein kodiert, das (i) einen aggregierenden Abschnitt, der von einem aggregierenden Krankheitsprotein (ADP) stammt, das mit einem Krankheitszustand assoziiert ist, in dem das ADP durch eine induzierte Konformationspolymerisationswechselwirkung pathologisch aggregiert, und (ii) einen heterologen Membranlokalisierungsabschnitt umfasst. 30
40. Zell- oder Gewebeprobe des Tieres nach Anspruch 39, umfassend einen mit einer Nucleotidsequenz, die für das membranlokalisierbare Fusionsprotein kodiert, operabel verbundenen Promotor. 40
41. Verfahren, Tier oder Probe nach einem der Ansprüche 38 bis 40, worin das Tier ein Nagetier ist. 45
42. Isoliertes Fusionsprotein, das für eine chimäre Nucleotidsequenz nach einem der Ansprüche 28 bis 32 kodiert. 50
43. Verfahren zur Herstellung eines Proteins nach Anspruch 42, wobei das Verfahren den Schritt des Verursachens oder Zulassens von Expression einer Nucleinsäure nach einem der Ansprüche 28 bis 32 in einer geeigneten Wirtszelle umfasst. 55
44. Verfahren zur Induktion eines Krankheitszustands in einem transgenen nichtmenschlichen Tier nach

Anspruch 39, worin das aggregierende Krankheitsprotein (ADP) pathologisch aggregiert, wobei das Verfahren das Verursachen oder Zulassen von Expression der Nucleotidsequenz umfasst, die für das membranlokalisierbare Fusionsprotein kodiert.

45. Verfahren zum Screenen auf ein Mittel zur Verwendung als therapeutisches, prognostisches oder diagnostisches Mittel bei einem Krankheitszustand, in dem ein aggregierendes Krankheitsprotein (ADP) pathologisch aggregiert, umfassend das Durchführen eines Verfahrens nach Anspruch 44 in Gegenwart eines Mittels, das vermutlich in der Lage ist, die Aggregation zu modulieren.
46. Verfahren nach Anspruch 27 oder Anspruch 45, worin das Mittel in der Lage ist, die Blut-Hirn-Schranke zu passieren.
47. Verfahren nach Anspruch 27, 45 oder 46, worin im Gehirn des Tieres etwas von Folgendem ermittelt wird:
 - (i) das Ausmaß proteolytischer Prozessierung des ADP,
 - (ii) die Fähigkeit des Mittels, die Menge an sich bildenden ADP-Aggregaten zu verringern,
 - (iii) die Fähigkeit des Mittels, die Menge an bereits gebildetem ADP-Aggregat zu beseitigen oder zu reduzieren.
48. Verfahren nach einem der Ansprüche 44 bis 47, worin das ADP Tau ist und der Krankheitszustand der Alzheimer-Krankheit entspricht.
49. Tier nach Anspruch 39, worin das ADP Tau ist und der Krankheitszustand der Alzheimer-Krankheit entspricht.

Revendications

1. Procédé pour induire ou modéliser l'état pathologique d'une protéine pathologique d'agrégation (ADP) qui est associée à un état pathologique dans lequel l'ADP s'agrège de manière pathologique par une interaction de polymérisation conformationnelle induite, le procédé étant **caractérisé par** l'étape consistant à mettre à disposition une protéine de fusion localisable sur une membrane comprenant (i) une partie d'agrégation, qui est dérivée de l'ADP, (ii) une partie hétérologue de localisation sur une membrane, où ledit procédé est réalisé *in vitro*, ou est réalisé dans une cellule ou une lignée cellulaire en culture, ou est réalisé chez un animal non humain.
2. Procédé selon la revendication 1, dans lequel la par-

- tie d'agrégation de la protéine de fusion est dérivée de l'ADP, et dans lequel la localisation sur une membrane de la protéine de fusion entraîne l'exposition du site de capture de haute affinité de la protéine ADP de sorte que l'agrégation d'une autre ADP soit favorisée, et l'agrégation pathologique de l'ADP dans l'état pathologique entraîne la maturation protéolytique de l'ADP en un fragment de domaine central.
- 5
3. Procédé selon la revendication 1 ou la revendication 2, comprenant les étapes consistant à :
- 10
- (a) mettre à disposition la protéine de fusion
- (b) l'introduire dans une membrane.
- 15
4. Procédé selon l'une quelconque des revendications précédentes, dans lequel l'agrégation de l'ADP dans l'état pathologique est associée à une neurodégénérescence et/ou à une démence clinique.
- 20
5. Procédé selon la revendication 4, dans lequel la partie d'agrégation comprend, ou consiste essentiellement en, au moins le fragment central de l'ADP.
- 25
6. Procédé selon l'une quelconque des revendications précédentes, dans lequel la partie de localisation sur une membrane comprend une séquence d'ancrage à la membrane et/ou une séquence d'arrêt de transfert.
- 30
7. Procédé selon l'une quelconque des revendications précédentes, dans lequel la maladie est sélectionnée parmi la maladie de Pick, la paralysie supranucléaire progressive, la maladie d'Alzheimer.
- 35
8. Procédé selon l'une quelconque des revendications précédentes, dans lequel l'ADP est la protéine tau.
- 40
9. Procédé selon la revendication 8, dans lequel la partie d'agrégation de la protéine de fusion comprend un fragment central de tau, le fragment de tau s'étendant entre les acides aminés 186-296 à 390-441 de la protéine tau de pleine longueur montrée sur la Figure 5.
- 45
10. Procédé selon la revendication 9, dans lequel la partie d'agrégation de la protéine de fusion consiste en les résidus d'acides aminés 296-390, 190-441 ou 190-390 de la protéine de pleine longueur.
- 50
11. Procédé selon l'une quelconque des revendications 2 à 10, destiné à cribler un partenaire de liaison spécifique du site de capture de haute affinité de l'ADP, le procédé comprenant les étapes consistant à :
- 55
- (a) mettre à disposition la protéine de fusion sous une forme localisée sur une membrane,
- (b) mettre en contact la protéine de fusion avec un ou plusieurs partenaires de liaison potentiels, où ladite étape de mise en contact est intracellulaire et le partenaire de liaison est présent au sein de la cellule,
- (c) déterminer si le partenaire de liaison potentiel se lie ou non à la protéine de fusion.
12. Procédé selon l'une quelconque des revendications 1 à 10, destiné à induire l'agrégation de molécules ADP, le procédé comprenant les étapes consistant à :
- (a) mettre à disposition la protéine de fusion sous une forme localisée sur une membrane,
- (b) provoquer ou permettre une interaction de polymérisation conformationnelle induite entre la protéine de fusion et une molécule ADP supplémentaire qui est une ADP ou un dérivé de celle-ci, et
- (c) surveiller l'étendue de l'agrégation de la molécule ADP supplémentaire.
13. Procédé selon la revendication 12, dans lequel la membrane est une membrane cellulaire et l'agrégation a lieu dans le compartiment intracellulaire.
14. Procédé selon la revendication 13, dans lequel l'étape (a) consiste soit à :
- introduire la protéine de fusion dans une cellule de sorte qu'elle soit localisée sur une membrane dans la cellule, ou
- introduire un acide nucléique codant pour la protéine de fusion dans une cellule ou un ancêtre de celle-ci de sorte que la protéine de fusion soit exprimée dans la cellule et soit localisée sur une membrane au niveau d'une membrane dans la cellule.
15. Procédé selon la revendication 14, dans lequel la molécule ADP supplémentaire est une ADP soluble de pleine longueur qui est native pour la cellule.
16. Procédé selon la revendication 13 ou la revendication 14, dans lequel la molécule ADP supplémentaire est hétérologue pour la cellule et le procédé comprend l'étape qui consiste à introduire un acide nucléique codant pour la molécule ADP supplémentaire dans la cellule ou un ancêtre de celle-ci de sorte que la molécule ADP supplémentaire soit exprimée dans la cellule.
17. Procédé selon la revendication 16, dans lequel la protéine de fusion est exprimée sous le contrôle transcriptionnel d'un promoteur constitutif, et la molécule ADP supplémentaire est exprimée sous le contrôle transcriptionnel d'un promoteur inducible

en présence de l'agent d'induction approprié.

18. Procédé selon la revendication 16 ou la revendication 17, dans lequel la molécule ADP supplémentaire est un dérivé d'ADP qui est (i) un fragment de l'ADP, (ii) un mutant de l'ADP, ou (iii) une ADP soluble de pleine longueur. 5
19. Procédé selon la revendication 16 ou la revendication 18, dans lequel la molécule ADP supplémentaire incorpore des mutations dans l'ADP correspondant à celles qui sont connues ou suspectées d'être associées à l'état pathologique. 10
20. Procédé selon l'une quelconque des revendications 12 à 19, dans lequel l'agrégation est surveillée en surveillant la concentration ou le taux de l'une quelconque des espèces suivantes : (i) la protéine de fusion et/ou la molécule ADP supplémentaire non agrégée, (ii) un agrégat de la protéine de fusion et/ou de la molécule ADP supplémentaire, (iii) un fragment protéolytique de la protéine de fusion et/ou de la molécule ADP supplémentaire résultant de l'agrégation de celles-ci. 15
21. Procédé selon la revendication 20, dans lequel l'ADP est tau, et l'agrégation est surveillée en se basant sur des taux croissants d'un fragment de tau d'approximativement 18 ou 25 kDa. 20
22. Procédé selon la revendication 20 ou la revendication 21, dans lequel l'étape de surveillance consiste à utiliser un anticorps spécifique pour l'une quelconque de (i) la protéine de fusion, (ii) la molécule ADP supplémentaire, (iii) un fragment protéolytique de la protéine de fusion et/ou de la molécule ADP supplémentaire résultant de l'agrégation de celles-ci. 25
23. Procédé selon la revendication 22, dans lequel la protéine de fusion est immunologiquement distincte de la protéine ADP supplémentaire, et l'ADP est tau, et l'anticorps est sélectionné parmi un anticorps monoclonal qui (i) est spécifique pour un épitope spécifique à un humain localisé dans la région comprise entre Gly-16 et Gln-26 de tau ; (ii) est spécifique pour le fragment de tau central tronqué au niveau de Glu-391 ; (iii) est spécifique pour un épitope de tau générique dans le domaine de répétition ; ou (iv) est spécifique pour un épitope de tau générique non spécifique d'une espèce localisé entre Ser-208 et Ser-238. 30
24. Procédé pour identifier un modulateur de l'agrégation d'une protéine pathologique d'agrégation (ADP) qui est associée à un état pathologique dans lequel l'ADP s'agrège de manière pathologique par une interaction de polymérisation conformationnelle induite, le procédé consistant à 35
- (a) mettre en oeuvre un procédé selon l'une quelconque des revendications 12 à 23 en présence d'un agent suspecté d'être capable de moduler l'agrégation, 40
- (b) surveiller l'étendue de l'agrégation, et
- (c) corrélérer l'étendue de l'agrégation à l'activité modulatrice de l'agent. 45
25. Procédé selon la revendication 24, qui est réalisé dans le compartiment intracellulaire. 50
26. Procédé selon la revendication 25, dans lequel la cellule est issue d'une lignée cellulaire neuronale ou d'une lignée cellulaire de fibroblaste. 55
27. Procédé selon la revendication 25, dans lequel la cellule est comprise au sein d'un animal transgénique non humain.
28. Molécule d'acide nucléique isolée comprenant une séquence de nucléotides chimérique codant pour une protéine de fusion localisable sur une membrane ayant (i) une partie d'agrégation, qui est dérivée d'une protéine pathologique d'agrégation (ADP) qui est associée à un état pathologique dans lequel l'ADP s'agrège de manière pathologique par une interaction de polymérisation conformationnelle induite, (ii) une partie hétérologue de localisation sur une membrane.
29. Acide nucléique selon la revendication 28, dans lequel la séquence de nucléotides chimérique comprend une séquence UTR 3' exogène, et dans lequel la partie de localisation sur une membrane comprend une séquence signal et facultativement une séquence codant pour un domaine d'ancrage à une membrane ou une séquence d'arrêt de transfert.
30. Acide nucléique selon la revendication 28 ou la revendication 29, dans lequel la séquence de nucléotides de la partie d'agrégation code pour tau, ou pour un fragment central de tau.
31. Acide nucléique selon la revendication 30, dans lequel la séquence de nucléotides de la partie d'agrégation code pour un fragment tronqué de tau s'étendant entre les acides aminés 186-296 et 390-441 de la protéine de pleine longueur.
32. Acide nucléique selon la revendication 31, dans lequel la séquence de nucléotides de la partie d'agrégation code pour les acides aminés 296-390, 190-441 ou 190-390.
33. Procédé pour produire un acide nucléique selon l'une quelconque des revendications 28 à 32, comprenant l'étape qui consiste à ligaturer un acide nu-

- cléique codant pour la partie d'agrégation à un acide nucléique codant pour la partie hétérologue de localisation sur une membrane,
et dans lequel l'acide nucléique codant pour la partie hétérologue de localisation sur une membrane fait partie d'un vecteur recombinant dans lequel l'acide nucléique codant pour la partie d'agrégation est inséré.
- 34.** Vecteur recombinant qui comprend l'acide nucléique selon l'une quelconque des revendications 28 à 32, dans lequel l'acide nucléique est en liaison fonctionnelle avec un promoteur afin d'obtenir une transcription dans une cellule hôte.
- 35.** Vecteur selon la revendication 34, dans lequel le promoteur est sélectionné parmi : un promoteur spécifique à un tissu ; le promoteur spécifique aux prions ; ou le promoteur de l'énolase neurospécifique.
- 36.** Procédé qui comprend l'étape consistant à introduire le vecteur selon la revendication 34 ou la revendication 35 dans une cellule hôte.
- 37.** Cellule hôte contenant ou transformée avec un vecteur hétérologue selon la revendication 34 ou la revendication 35.
- 38.** Procédé pour préparer un modèle animal non humain transgénique présentant une pathologie amplifiée ou accélérée pour l'état pathologique dans lequel une protéine pathologique d'agrégation (ADP) s'agrège de manière pathologique, le procédé comprenant l'étape consistant à introduire dans un oeuf fertilisé de l'animal un vecteur selon l'une quelconque de la revendication 34 ou la revendication 35.
- 39.** Animal non humain transgénique qui comprend, au sein d'une pluralité de ses cellules, un promoteur en liaison fonctionnelle avec une séquence de nucléotides codant pour une protéine de fusion localisable sur une membrane comprenant (i) une partie d'agrégation, qui est dérivée d'une protéine pathologique d'agrégation (ADP) qui est associée à un état pathologique dans lequel l'ADP s'agrège de manière pathologique par une interaction de polymérisation conformationnelle induite, (ii) une partie hétérologue de localisation sur une membrane.
- 40.** Cellule ou échantillon de tissu de l'animal selon la revendication 39, comprenant un promoteur en liaison fonctionnelle avec une séquence de nucléotides codant pour la protéine de fusion localisable sur une membrane.
- 41.** Procédé, animal ou échantillon selon l'une quelconque des revendications 38 à 40, où l'animal est un rongeur.
- 42.** Protéine de fusion isolée qui est codée par la séquence de nucléotides chimérique selon l'une quelconque des revendications 28 à 32.
- 43.** Procédé pour fabriquer la protéine selon la revendication 42, le procédé comprenant l'étape qui consiste à provoquer ou à permettre une expression, à partir d'un acide nucléique selon l'une quelconque des revendications 28 à 32, dans une cellule hôte appropriée.
- 44.** Procédé pour induire, chez l'animal non humain transgénique selon la revendication 39, un état pathologique dans lequel la protéine pathologique d'agrégation (ADP) s'agrège de manière pathologique, le procédé consistant à provoquer ou à permettre l'expression de la séquence de nucléotides codant pour la protéine de fusion localisable sur une membrane.
- 45.** Procédé pour cribler un agent destiné à être utilisé en tant qu'agent thérapeutique, de pronostic ou de diagnostic dans un état pathologique dans lequel une protéine pathologique d'agrégation (ADP) s'agrège de manière pathologique, qui consiste à mettre en oeuvre un procédé selon la revendication 44 en présence d'un agent suspecté d'être capable de moduler l'agrégation.
- 46.** Procédé selon la revendication 27 ou la revendication 45, dans lequel l'agent est capable de traverser la barrière hémato-encéphalique.
- 47.** Procédé selon les revendications 27, 45 ou 46, dans lequel l'un quelconque des éléments suivants est déterminé dans le cerveau de l'animal :
- (i) l'étendue de la maturation protéolytique de l'ADP,
 - (ii) l'aptitude de l'agent à réduire la quantité d'agrégats d'ADP en formation,
 - (iii) l'aptitude de l'agent à éliminer ou à réduire le taux d'agrégats d'ADP déjà formés.
- 48.** Procédé selon l'une quelconque des revendications 44 à 47, dans lequel l'ADP est tau et l'état pathologique correspond à la maladie d'Alzheimer.
- 49.** Animal selon la revendication 39, dans lequel l'ADP est tau et l'état pathologique correspond à la maladie d'Alzheimer.





Figure 1a and 1b

(c) **Aggregation of SStau190-390 Produces an 18 kd Fragment**



(d) **Aggregation of SStau296-390: Interaction with T40 Produces Two Possible 18 kd Fragments, and a 12/14 kd Fragment**




Figure 1c and 1d




Figure 1e

Activities of compounds in the
 $\uparrow T40 + 40 \text{ kD} \Rightarrow \uparrow 25 \text{ kD}$ cell assay
 Results shown as relative production of 25 kD band





Figure 2

Thionine : memb-Tau model

↑T40 + 40 kD ⇒ ↑25 kD cell assay

Cellular activity predicted via standard inhibition model:

$$\text{activity} = [\text{tau}] / ([\text{tau}] + K_d * (1 + [\text{thionine}] / K_i))$$



Observed vs predicted activity

$r = 0.999$

Intracellular tau concentration

415 nM

Tau-tau binding affinity

500 nM

Thionine K_i

33 nM

Figure 3




Figure 4a

N- & C- terminal truncation of tau




Figure 4b

```

ATG GCT GAG CCC CGC CAG GAG TTC GAA GTG ATG GAA GAT CAC GCT GGG ACG TAC GGG TTG
Met ala glu pro arg gln glu phe glu val met glu asp his ala gly thr tyr gly leu
1      5      10      15      20
GGG GAC AGG AAA GAT CAG GGG GGC TAC ACC ATG CAC CAA GAC CAA GAG GGT GAC ACG GAC
gly asp arg lys asp gln gly gly tyr thr met his gln asp gln glu gly asp thr asp
25      30      35      40
GCT GGC CTG AAA GAA TCT CCC CTG CAG ACC CCC ACT GAG GAC GGA TCT GAG GAA CCG GGC
ala gly leu lys glu ser pro leu gln thr pro thr glu asp gly ser glu glu pro gly
45      50      55      60
TCT GAA ACC TCT GAT GCT AAG AGC ACT CCA ACA GCG GAA GAT GTG ACA GCA CCC TTA GTG
ser glu thr ser asp ala lys ser thr pro thr ala glu asp val thr ala pro leu val
65      70      75      80
GAT GAG GGA GCT CCC GGC AAG CAG GCT GCC GCG CAG CCC CAC ACG GAG ATC CCA GAA GGA
asp glu gly ala pro gly lys gln ala ala gln pro his thr glu ile pro glu gly
85      90      95      100
ACC ACA GCT GAA GAA GCA GGC ATT GGA GAC ACC CCC AGC CTG GAA GAC GAA GCT GCT GGT
thr thr ala glu ala gly ile gly asp thr pro ser leu glu asp glu ala ala gly
105      110      115      120
CAC GTG ACC CAA GCT CGC ATG GTC AGT AAA AGC AAA GAC GGG ACT GGA AGC GAT GAC AAA
his val thr gln ala arg met val ser lys ser lys asp gly thr gly ser asp asp lys
125      130      135      140
AAA GCC AAG GGG GCT GAT GGT AAA ACG AAG ATC GCC ACA CCG CGG GGA GCA GCC CCT CCA
lys ala lys gly ala asp gly lys thr lys ile ala thr pro arg gly ala ala pro pro
145      150      155      160
GGC CAG AAG GGC CAG GCC AAC GCC ACC AGG ATT CCA GCA AAA ACC CCG CCC GCT CCA AAG
gly gln lys gly gln ala asn ala thr arg ile pro ala lys thr pro pro ala pro lys
165      170      175      180
ACA CCA CCC AGC TCT GGT GAA CCT CCA AAA TCA GGG GAT CGC AGC GGC TAC AGC AGC CCC
thr pro pro ser gly glu pro pro lys ser gly asp arg ser gly tyr ser ser pro
185      190      195      200
GGC TCC CCA GGC ACT CCC GGC AGC CGC TCC CGC ACC CCG TCC CTT CCA ACC CCA CCC ACC
gly ser pro gly thr pro gly ser arg ser arg thr pro ser leu pro thr pro pro thr
205      210      215      220
CGG GAG CCC AAG AAG GTG GCA GTG GTC CGT ACT CCA CCC AAG TCG CCG TCT TCC GCC AAG
arg glu pro lys lys val ala val val arg thr pro pro lys ser pro ser ser ala lys
225      230      235      240
AGC CGC CTG CAG ACA GCC CCC GTG CCC ATG CCA GAC CTG AAG AAT GTC AAG TCC AAG ATC
ser arg leu gln thr ala pro val pro met pro asp leu lys asn val lys ser lys ile
245      250      255      260
GGC TCC ACT GAG AAC CTG AAG CAC CAG CCG GGA GGC GGG AAG GTG CAG ATA ATT AAT AAG
gly ser thr glu asn leu lys his gln pro gly gly gly lys val gln ile ile asn lys
265      270      275      280
AAG CTG GAT CTT AGC AAC GTC CAG TCC AAG TGT GGC TCA AAG GAT AAT ATC AAA CAC GTC
lys leu asp leu ser asn val gln ser lys cys gly ser lys asp asn ile lys his val
285      290      295      300
CCG GGA GGC GGC AGT GTG CAA ATA GTC TAC AAA CCA GTT GAC CTG AGC AAG GTG ACC TCC
pro gly gly gly ser val gln ile val tyr lys pro val asp leu ser lys val thr ser
305      310      315      320
AAG TGT GGC TCA TTA GGC AAC ATC CAT CAT AAA CCA GGA GGT GGC CAG GTG GAA GTA AAA
lys cys gly ser leu gly asn ile his his lys pro gly gly gly gln val glu val lys
325      330      335      340
TCT GAG AAG CTT GAC TTC AAG GAC AGA GTC CAG TCG AAG ATT GGG TCC CTG GAC AAT ATC
ser glu lys leu asp phe lys asp arg val gln ser lys ile gly ser leu asp asn ile
345      350      355      360
ACC CAC GTC CCT GGC GGA GGA AAT AAA AAG ATT GAA ACC CAC AAG CTG ACC TTC CGC GAG
thr his val pro gly gly gly asn lys lys ile glu thr his lys leu thr phe arg glu

```

Figure 5a

				365					370					375				380
AAC	GCC	AAA	GCC	AAG	ACA	GAC	CAC	GGG	GCG	GAG	ATC	GTG	TAC	AAG	TCG	CCA	GTG	TCT
asn	ala	lys	ala	lys	thr	asp	his	gly	ala	glu	ile	val	tyr	lys	ser	pro	val	ser
				385					390					395				400
GGG	GAC	ACG	TCT	CCA	CGG	CAT	CTC	AGC	AAT	GTC	TCC	TCC	ACC	GGC	AGC	ATC	GAC	GTA
gly	asp	thr	ser	pro	arg	his	leu	ser	asn	val	ser	ser	thr	gly	ser	ile	asp	met
				405					410					415				420
GAC	TCG	CCC	CAG	CTC	GCC	ACG	CTA	GCT	GAC	GAG	GTG	TCT	GCC	TCC	CTG	GCC	AAG	GGT
asp	ser	pro	gln	leu	ala	thr	leu	ala	asp	glu	val	ser	ala	ser	leu	ala	lys	gln
				425					430					435				440
TTG	TGA																	
leu	***																	

Figure 5a cont ...

square brackets indicates signal sequence
 bold indicates tubulin-binding segments

SStau186-441

[MKWVTFLLLLFISGSAFSPV]KSGDRSGYSSPGSPGTPGSRSRTPSLPTPP
 TREPKKVAVVRTPPKSPSSAKSRLQTAPVMPDLKNVKS**KIGSTENLKHQP**
GGGKVQIINKKLDSLNVQSKCGSKDNIKHVPGGGSVQIVYKPDLSKVTSK
CGSLGNIHHKPGGGQVEVKSEKLDKDRVQSKIGSLDNITHVPGGGNKKIE
 THKLTFRENAKAKTDHGAEIVYKSPVVS~~GD~~TSRHL~~SNV~~SSTGSIDMVDSP
 QLATLADEVASLAKQGL

SStau186-390

[MKWVTFLLLLFISGSAFSPV]KSGDRSGYSSPGSPGTPGSRSRTPSLPTPP
 TREPKKVAVVRTPPKSPSSAKSRLQTAPVMPDLKNVKS**KIGSTENLKHQP**
GGGKVQIINKKLDSLNVQSKCGSKDNIKHVPGGGSVQIVYKPDLSKVTSK
CGSLGNIHHKPGGGQVEVKSEKLDKDRVQSKIGSLDNITHVPGGGNKKIE
 THKLTFRENAKAKTDHGA

SStau296-390

[MKWVTFLLLLFISGSAFSPV]**NIKHVPGGGSVQIVYKPDLSKVTSKCGSLG**
NIHHKPGGGQVEVKSEKLDKDRVQSKIGSLDNITHVPGGGNKKIETHKLTFR
 ENAKAKTDHGA

Figure 5b




Figure 6a

Construct aa296-390 used for DNA microinjection into fertilized oocytes of mice




Figure 6b




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

**Northern blot analysis of construct aa296-390 (line 1)
shows high expression level**




Figure 12

**Pre-IP followed by IP7.51 shows
significant difference between wt and tg
at 12kD**




Figure 13

S1 fraction from three different mice of line 1 show 18kD




Figure 14

2-DE of boiled and concentrated S1 from mouse brain (6months) line 1 (aa296-390)




Figure 15




Figure 16a and 16b




Figure 16c

In vitro aggregation of
bacterially expressed dGAE




Figure 16d





Figure 16e

(a)



(b)




Figure 17




Figure 18




Figure 19a




Figure 19b

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
<i>Neurodegenerative disorders</i>				
Prion protein	Prion diseases (CJD, nvCJD, Fatal familial insomnia, Gerstmann-Strausler-Scheinker syndrome, Kuru)	<i>Inherited and sporadic forms</i> PrP ^{Sc} 27-30; many mutations Fibrillogenic domains: 113-120, 178-191, 202-218	27	Prusiner (1998) Gasset et al. (1992)
Tau protein	Alzheimer's disease, Down's syndrome, FTDP-17, CBD, post-encephalitic parkinsonism, Pick's disease, parkinsonism with dementia complex of Guam	<i>Inherited and sporadic forms</i> Truncated tau (tubulin-binding domain) 297-391 Mutations in tau in FTDP-17 Many mutations in presenilin proteins	10-12	Wischik et al. (1988) Hutton et al. (1998) Czech et al. (2000)
Amyloid β -protein	Alzheimer's disease, Down's syndrome	<i>Inherited and sporadic forms</i> Amyloid β -protein; 1-42(3); 11 mutations in APP in rare families	4	Glenner & Wong, (1984) Goate et al. (1991)
Huntingtin Ataxins (1, 2, 3, 7) Atrophin Androgen receptor	Huntington's disease Spinocerebellar ataxias (SCA1, 2, 3, 7) Dentatorubralpallidum atrophy (DRPLA) Spinal and bulbar muscular atrophy	N-termini of protein with expanded glutamine repeats Proteins with expanded glutamine repeats Proteins with expanded glutamine repeats Proteins with expanded glutamine repeats	40	DiFiglia et al. (1997) Paulson et al. (2000) Paulson et al. (2000) Paulson et al. (2000)
Neuroserpin α -Synuclein	Familial encephalopathy with neuronal inclusion bodies (FENIB) Parkinson's disease, dementia with Lewy bodies, multiple system atrophy	Neuroserpin; S49P, S52R <i>Inherited and sporadic forms</i> A53T, A30P in rare autosomal-dominant PD families	57 19	Davis et al. (1999) Spillantini et al. (1998) Polymeropoulos et al. (1997)
Cystatin C Superoxide dismutase 1	Hereditary cerebral angiopathy (Icelandic) Amyotrophic lateral sclerosis	Cystatin C less 10 residues; L68Q SOD1 mutations	12-13	Abrahamson et al. (1992) Shibata et al. (1996)
<i>Non-neurodegenerative disorders</i>				
Haemoglobin	Sickle cell anaemia Inclusion body haemolysis	Haemoglobin beta chain (S) Many mutations		Carrell & Gooptu (1998)
Serpins	α 1-Antitrypsin deficiency (emphysema, cirrhosis) Antithrombin deficiency (thromboembolic disease) C1-inhibitor deficiency (angioedema)	Mutations Mutations Mutations		Lomas et al. (1992) Carrell & Gooptu (1998) Carrell & Gooptu (1998)
Immunoglobulin light chain	Plasma cell dyscrasias (primary systemic AL amyloidosis)	light chain or fragments	0.5-25	Westermarck et al. (1985)
Serum amyloid A	Reactive, secondary systemic AA amyloidosis Chronic inflammatory disease	Variable N-terminal fragments of SAA	4.5-10	Westermarck et al. (1985)

(continued.....)

Figure 20

(.....continued)

Transthyretin	Familial amyloid polyneuropathy (systemic; FAP I)	Tetramer dissociated to conformational monomer variant Many mutations (some not associated with amyloid; several different types of disease)	10-14	Gustavsson et al. (1991)
	Senile cardiac amyloidosis	Normal transthyretin	10-14	Gustavsson et al. (1991)
Gelsolin	Familial amyloidosis - Finnish type (FAP IV)	D167Q leads to truncated 173-225/243 (critical residues 182-192)	9.5	Maury & Baumann (1990)
β 2-Microglobulin	Haemodialysis amyloidosis Prostatic amyloid	β 2-Microglobulin	12-25	Gorevic et al. (1985)
Apolipoprotein AI	Familial amyloid polyneuropathy (systemic; FAP II)	N-terminal 83-93 residues; G26R, W50R, L60R	9	Booth et al. (1995)
Lysozyme	Familial visceral amyloidosis	Lysozyme or fragments (with or without I56T, D67H)	14	Pepys et al. (1993)
Amylin (islet amyloid polypeptide)	Type II diabetes (NIDDM)	Fragment (critical core of 20-29); no mutations	3.9	Westermarck (1990)
Fibrinogen α -chain	Hereditary renal amyloidosis	Fibrinogen fragments	7-10	Uemichi et al. (1994)
Procalcitonin	Medullary carcinoma of thyroid	Calcitonin fragments	3.4	Sletten et al. (1976)
Atrial natriuretic factor	Cardiac amyloidosis	ANF, no mutants	3.5	Johansson et al. (1987)
Insulin	Injection localised amyloidosis	Insulin		Dische et al. (1988)
Other proteins forming amyloid	(<i>in vitro</i>)	Other proteins		Chiti et al. (1999)

Figure 20 cont ...

Gel Mobility (kD)	Yield (μg)	Proteins Identified	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
46/44	2.3	PHF-tau ubiquitin	Q ²⁶⁹	P	G	G	G	K	V	Q	I	V	Y		P ²⁸¹							
					F ⁴	V		T	L ⁸													
34/32	7.2	PHF-tau ubiquitin porin " " " "	Q ²⁶⁹	P	G	G	G	K	V	I	V	Y		P ²⁸¹								
					F ⁴	V		T	L ⁸													
			A ¹	V	P	T	Y	D	L	G ¹⁰												
			D ²⁹	L	K	T	K	E ³⁹														
			N ¹⁵⁵	F	E			N	L	A ²⁰⁸												
26/24	7.9	PHF-tau ubiquitin core-2	Q ²⁶⁹	P	G	G	G	K	V	Q	I	V	Y	K	P ²⁸¹					V ¹⁷		
					V ⁵	T	L	G	T											G ²²		
			G ¹⁶	A	P	P	Q	P	Q	D	L	E	F	T	K	L	P					
20/18	1.7	subunit 9	D ¹	I	D	A	K	I		A					A ¹⁴							
14/12	4.0	PHF-tau	H ²⁶⁹	Q/V	P	G	G	G	K/S	V	Q	I	V/I	Y	K	P	V	D	L	S	K	V ²⁸⁷
5		Aβ " " "	R ⁵	H	D	S	G	Y	E	V	H	H	Q	K	L	V	F	F	A		D	V ²⁴
			H ⁶	D	S	G	Y	E	V	H				L					A ²¹			
					Y ¹⁰	E	V	H	Q	L			F ¹⁹									
			Y ¹⁰	E	V	H	Q	L		F ¹⁹												
		subunit-9	I ²	T	A	A	F	I	G	A	G	A	A		G ¹⁷							
Pellet	2.6	PHF-tau ubiquitin subunit 9	P ²⁷⁰		K	V	Q	V	Y	K	P	V	D ²⁸³									
					F ⁴	V	L	T	G	T		T ¹⁴										
			I ²	D	T	A	A	K	F	I	G	A	G	A	A	T	V ¹⁶					
TOTAL	25.7																					

Figure 21

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9630766 A [0009] [0010] [0013] [0028] [0049] [0111] [0140] [0141]
- US 5912410 A [0023] [0162]
- US 5898094 A [0023] [0162]
- US 5010182 A [0067]
- EP 362179 A [0067]
- WO 9013646 A [0067]
- EP 117060 A [0084]
- EP 117058 A [0084]
- WO 8905859 A [0096]
- US 4399216 A [0097]
- US 4873191 A [0163]

Non-patent literature cited in the description

- Mukaetova-Ladinska, E.B. et al. *Am. J. Pathol.*, 2000, vol. 157 (2), 623-636 [0002]
- Wischik et al. *PNAS USA*, 1988, vol. 85, 4506 [0003]
- Kang et al. *Nature*, 1987, vol. 325, 733 [0003]
- The Molecular and Cellular Neurobiology Series. Wischik et al. *Neurobiology of Alzheimer's Disease*. Bios Scientific Publishers, 2000 [0003] [0047]
- Mukaetova-Ladinska et al. *Am. J. Pathol.*, 1993, vol. 143, 565 [0004]
- Wischik et al. *Neurobiol. Ageing*, 1995, vol. 16, 409 [0004]
- Lai et al. *Neurobiol. Ageing*, 1995, vol. 16, 433 [0004]
- Harrington et al. *Dementia*, 1994, vol. 5, 215 [0004]
- Harrington et al. *Am. J. Pathol.*, 1994, vol. 145, 1472 [0004]
- Terry et al. *Ann. Neurol.*, 1991, vol. 30, 572 [0004]
- Bondareff et al. *Arch. Gen. Psychiatry*, 1993, vol. 50, 350 [0004]
- Goedert, M. et al. *EMBO J.*, 1989, vol. 8, 393-399 [0005]
- Goedert, M. et al. *Neuron*, 1989, vol. 3, 519-526 [0005]
- Wischik, C.M. et al. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 4884-4888 [0005]
- Wischik et al. *PNAS USA*, 1988, vol. 85, 4506-4510 [0005]
- Novak, M. et al. *EMBO J.*, 1993, vol. 12, 365-370 [0005]
- Jakes, R. et al. *EMBO J.*, 1991, vol. 10, 2725-2729 [0005]
- Wischik et al. *Proc Natl Acad Sci USA*, 1996, vol. 93, 11213-11218 [0005]
- Mena, R. et al. *Acta Neuropathol.*, 1995, vol. 89, 50-56 [0006]
- Mena, R. et al. *Acta Neuropathol.*, 1996, vol. 91, 633-641 [0006]
- Wischik, C.M. et al. *Microtubule-associated proteins: modifications in disease*. Harwood Academic Publishers, 1996, 185-241 [0006]
- Lai, R. Y. K. et al. *Neurobiology of Ageing*, 1995, vol. 16 (3), 433-445 [0006]
- Bondareff, W. et al. *J. Neuropath. Exper. Neurol.*, 1994, vol. 53 (2), 158-164 [0006]
- Wischik, C.M. et al. *Microtubule-associated proteins: modifications in disease*. Harwood Academic Publishers, 1997, 185-241 [0007]
- Mayeux, R. et al. *New Eng. J. Med.*, 1999, vol. 341, 1670-1679 [0008]
- Baum, L. et al. *Mol. Brain Res.*, 1995, vol. 34, 1-17 [0011]
- Shelanski et al. *Proc. Natl. Acad. Sci. USA*, 1973, vol. 70, 765-768 [0043]
- Matus, A. *Microtubules*. John Wiley and Sons, 155-166 [0043]
- Kindler ; Garner. *Mol. Brain Res.*, 1994, vol. 26, 218-224 [0043]
- Ho et al. *Gene*, 1989, vol. 77, 51-59 [0065]
- Crameri et al. *Nature*, 1998, 391 [0065]
- Partridge, K.A. et al. *Cytotechnology*, 1999, vol. 30, 37-47 [0082] [0186]
- Altschul et al. *J. Mol. Biol.*, 1990, vol. 215, 403-10 [0083]
- Sambrook ; Fritsch ; Maniatis. *Molecular Cloning, A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1989 [0084]
- Ausubel et al. *Short Protocols in Molecular Biology*. John Wiley and Sons, 1992 [0084]
- Gething et al. *Nature*, 1981, vol. 293, 620-695 [0084]
- Mantei et al. *Nature*, 1979, vol. 281, 40-46 [0084]
- Urlaub et al. *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4216 [0094]
- Stinchcomb et al. *Nature*, 1979, vol. 282, 39 [0094]
- Kingsman et al. *Gene*, 1979, vol. 7, 141 [0094]
- Tschemper et al. *Gene*, 1980, vol. 10, 157 [0094]
- Jones. *Genetics*, 1977, vol. 85, 12 [0094]
- Shaw et al. *Gene*, 1983, vol. 23, 315 [0096]
- Graham ; van der Eb. *Virology*, 1978, vol. 52, 456-457 [0097]

- **Van Solingen et al.** *J. Bact.*, 1977, vol. 130, 946 [0097]
- **Hsiao et al.** *Proc. Natl. Acad. Sci. (USA)*, 1979, vol. 76, 3829 [0097]
- **Keown et al.** *Methods in Enzymology*, 1990, vol. 185, 527-537 [0097]
- **Mansour et al.** *Nature*, 1988, vol. 336, 348-352 [0097]
- **Urlaub ; Chasin.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4216 [0099]
- **Mather.** *Biol. Reprod.*, 1980, vol. 23, 243-251 [0099]
- **Mammalian Cell Biotechnology: a Practical Approach.** JRL Press, 1991 [0102]
- **Thomas.** *Proc. Natl Acad Sci. USA*, 1980, vol. 77, 5201-5205 [0103]
- **Novak et al.** *Proc. Natl. Acad. Sci. USA*, 1991, vol. 88, 5837-5841 [0124]
- **Slobada et al.** *Cell Mobility.* Cold Spring Harbor, 1976, 1171-1212 [0142]
- **Remington's Pharmaceutical Sciences.** 1980 [0157]
- **Manipulating the mouse embryo. a laboratory manual.** Cold Spring Harbour Press, 1994 [0185]
- **Stahl et al.** *Biochemistry*, 1990, vol. 29, 8879-8884 [0185]
- **Goedert et al.** *Neurone*, 1989, vol. 3, 519-526 [0191]
- **Novak, M. et al.** *Proc. Natl. Acad. Sci. USA*, 1991, vol. 88, 5837-5841 [0197]
- **Schliwa et al.** *PNAS USA*, 1981, vol. 78, 1037-1041 [0197]
- **Gorman, C.M. et al.** *Nuc. Acids Res.*, 1983, vol. 11, 7631-7648 [0200]
- **Kruh, J.** *Mol. Cell. Biochem*, 1982, vol. 42, 65-82 [0200]
- **Arendt et al.** *Neuroscience*, 1995, vol. 69, 691-698 [0238]
- **Braak et al.** *Neuroscience letters*, 1994, vol. 171, 1-4 [0238]
- **Brion et al.** *American Journal of Pathology*, 1999, vol. 154, 255-270 [0238]
- **Cork et al.** *Journal of Neuropathology and Experimental Neurology*, 1988, vol. 49, 629-641 [0238]
- **Duff et al.** *Neurobiology of Disease*, 2000, vol. 7, 87-98 [0238]
- **Geula et al.** *Nature Medicine*, 1998, vol. 4, 827-834 [0238]
- **Gotz et al.** *EMBO Journal*, 1995, vol. 14, 1304-1313 [0238]
- **Harada et al.** *Nature*, 1994, vol. 369, 488-491 [0238]
- **Ikegami et al.** *Neuroscience letters*, 2000, vol. 279, 129-132 [0238]
- **Ishihara et al.** *Neuron*, 1999, vol. 24, 751-762 [0238]
- **Janus et al.** *Biochimica et Biophysica Acta*, 2000, vol. 1502, 63-75 [0238]
- **lewis et al.** *Nature Genetics*, 2000, vol. 25, 402-405 [0238]
- **Nelson et al.** *Neurobiology of Aging*, 1995, vol. 16, 315-323 [0238]
- **Roertgen et al.** *Neurobiology of Aging*, vol. 17, 243-247 [0238]
- **Schultz et al.** *Journal of Neuropathology and Experimental Neurology*, 2000, vol. 59, 39-52 [0238]
- **Spittaels et al.** *American Journal of Pathology*, 1999, vol. 155, 2153-2165 [0238]
- **Sturchler-Pierrat et al.** *Proceedings of the National Academy of Sciences, USA*, 1997, vol. 93, 13287-13292 [0238]
- **Abrahamson, M. ; Jonsdottir, S. ; Olafsson, I. ; Grubb, A.** Hereditary cystatin C amyloid angiopathy identification of the disease-causing mutation and specific diagnosis by polymerase chain reaction based analysis. *Human Genetics*, 1992, vol. 89, 377-380 [0239]
- **Booth, D.R. ; Sunde, M. ; Bellotti, V. ; Robinson, C.V. ; Hutchinson, W.L. ; Fraser, P.E. ; Hawkins, P.N. ; Dobson, C.M. ; Radford, S.E. ; Blake, C.C.F.** Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis. *Nature*, 1997, vol. 385, 787-793 [0239]
- **Carrell, R.W. ; Gooptu, B.** Conformational changes and disease - serpins, prions and Alzheimer's. *Current Opinion in Structural Biology*, 1998, vol. 8, 799-809 [0239]
- **Chiti, F. ; Webster, P. ; Taddei, N. ; Clark, A. ; Stefani, M. ; Ramponi, G. ; Dobson, C.** Designing conditions for in vitro formation of amyloid protofilaments and fibrils. *Proceedings of the National Academy of Sciences, USA*, 1999, vol. 96, 3590-3594 [0239]
- **Czech, C. ; Tremp, G. ; Pradier, L.** Presenilins and Alzheimer's disease: biological functions and pathogenic mechanisms. *Progress in Neurobiology*, 2000, vol. 60, 363-384 [0239]
- **Davis, R.L. ; Shrimpton, A.E. ; Holohan, P.D. ; Bradshaw, C. ; Feiglin, D. ; Collins, G.H. ; Sonderegger, P. ; Kinter, J. ; Becker, L.M. ; Lacbawan, F.** Familial dementia caused by polymerization of mutant neuroserpin. *Nature*, 1999, vol. 401, 376-379 [0239]
- **DiFiglia, M. ; Sapp, E. ; Chase, K.O. ; Davies, S.W. ; Bates, G.P. ; Vonsattel, J.P. ; Aronin, N.** Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. *Science*, 1997, vol. 277, 1990-1993 [0239]
- **Dische, F.E. ; Wernstedt, C. ; Westermark, G.T. ; Westermark, P. ; Pepys, M.B. ; Rennie, J.A. ; Gilbert, S.G. ; Watkins, P.J.** Insulin as an amyloid-fibril protein at sites of repeated insulin injections in a diabetic patient. *Diabetologia*, 1988, vol. 31, 158-161 [0239]
- **Gasset, M. ; Bladwin, M.A. ; Lloyd, D.H. ; Abriel, J.-M. ; Holtzman, D.M. ; Cohen, F.E. ; Fletterick, R. ; Prusiner, S.B.** Predicted α -helical region of the prion protein when synthesized as peptides form amyloid. *Proceedings of the National Academy of Sciences*, 1992, vol. 89, 10940-10944 [0239]

- **Glennner, G.G. ; Wong, C.W.** Alzheimer's disease: initial report of the purification and characterisation of a novel cerebrovascular amyloid protein. *Biochemical and Biophysical Research Communications*, 1984, vol. 120, 885-890 [0239]
- **Goate, A. ; Chartier-Harlin, M.-C. ; Mullan, M. ; Brown, J. ; Crawford, F. ; Fidani, L. ; Giuffra, L. ; Haynes, A. ; Irving, N. ; James, L.** Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature*, 1991, vol. 349, 704-706 [0239]
- **Gorevic, P.D. ; Casey, T.T. ; Stone, W.J. ; DiRaimondo, C.R. ; Prelli, F.C. ; Frangione, B.** b-2 Microglobulin is an amyloidogenic protein in man. *Journal of Clinical Investigation*, 1985, vol. 76, 2425-2429 [0239]
- **Gustavsson, A. ; Engstrom, U. ; Westermark, P.** Normal transthyretin and synthetic transthyretin fragments form amyloid-like fibrils in vitro. *Biochemical and Biophysical Research Communications*, 1991, vol. 175, 1159-1164 [0239]
- **Hutton, M. ; Lendon, C. ; Rizzu, P. ; Baker, M. ; Froelich, S. ; Houlden, H. ; Pickering-Brown, S. ; Chakraverty, S. ; Isaacs, A. ; Grover, A.** Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature*, 1998, vol. 393, 702-705 [0239]
- **Johansson, B. ; Wernstedt, C. ; Westermark, P.** Atrial natriuretic peptide deposited as atrial amyloid fibrils. *Biochemical and Biophysical Research Communications*, 1987, vol. 148, 1087-1092 [0239]
- **Lomas, D.A. ; Evans, D.L. ; Finch, J.T. ; Carrell, R.W.** The mechanism of α_1 -antitrypsin accumulation in the liver. *Nature*, 1992, vol. 357, 605-607 [0239]
- **Maury, C.P. ; Baumann, M.** Isolation and characterization of cardiac amyloid in familial amyloid polyneuropathy type IV (Finnish): relation of the amyloid protein to variant gelsolin. *Biochimica et Biophysica Acta*, 1990, vol. 1096, 84-86 [0239]
- **Paulson, H.L.** Human genetics '99: trinucleotide repeats. *American Journal of Human Genetics*, 1999, vol. 64, 339-345 [0239]
- **Pepys, M.B. ; Hawkins, P.N. ; Booth, D.R. ; Vighushin, D.M. ; Tennent, G.A. ; Soutar, A.K. ; Totty, N. ; Nguyen, O. ; Blake, C.C.F. ; Terry, C.J.** Human lysozyme gene mutations cause hereditary systemic amyloidosis. *Nature*, 1993, vol. 362, 553-557 [0239]
- **Polymeropoulos, M.H. ; Lavedan, C. ; Leroy, E. ; Ide, S.E. ; Dehejia, A. ; Dutra, A. ; Pike, B. ; Root, H. ; Rubenstein, J. ; Boyer, R.** Mutation in the α -synuclein gene identified in families with Parkinson's disease. *Science*, 1997, vol. 276, 2045-2047 [0239]
- **Prusiner, S.B. ; Scott, M.R. ; DeArmond, S.J. ; Cohen, F.E.** Prion protein biology. *Cell*, 1998, vol. 93, 337-348 [0239]
- **Shibata, N. ; Hirano, A. ; Kobayashi, M. ; Sidique, T. ; Deng, H.X. ; Hung, W.Y. ; Kato, T. ; Asayama, K.** Intense superoxide dismutase-1 immunoreactivity in intracytoplasmic hyaline inclusions of familial amyotrophic lateral sclerosis with posterior column involvement. *Journal of Neuropathology and Experimental Neurology*, 1996, vol. 55, 481-490 [0239]
- **Sletten, K. ; Westermark, P. ; Natvig, J.B.** Characterization of amyloid fibril proteins from medullary carcinoma of the thyroid. *Journal of Experimental Medicine*, 1976, vol. 143, 993-998 [0239]
- **Spillantini, M.G. ; Crowther, R.A. ; Jakes, R. ; Hasegawa, M. ; Goedert, M.** α -Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. *Proceedings of the National Academy of Sciences, USA*, 1998, vol. 95, 6469-6473 [0239]
- **Uemichi, T. ; Liuepnicks, J.j. ; Benson, M.D.** Hereditary renal amyloidosis with a novel variant fibrinogen. *Journal of Clinical Investigation*, 1994, vol. 93, 731-736 [0239]
- **Westermark, P. ; Engstrom, U. ; Johnson, K.H. ; Westermark, G.T. ; Betsholtz, C.** Islet amyloid polypeptide: pinpointing amino acid residues linked to amyloid fibril formation. *Proceedings of the National Academy of Sciences, USA*, 1990, vol. 87, 5036-5040 [0239]
- **Westermark, P. ; Johnson, K.H. ; O'Brien, T.D. ; Betsholtz, C.** Islet amyloid polypeptide - a novel controversy in diabetes research. *Diabetologia*, 1992, vol. 35, 297-303 [0239]
- **Westermark, P. ; Johnson, K.H. ; Pitkanen, P.** Systemic amyloidosis: A review with emphasis on pathogenesis. *Applied Physiology*, 1985, vol. 3, 55-68 [0239]
- **Wischik, C.M. ; Novak, M. ; Thogersen, H.C. ; Edwards, P.C. ; Runswick, M.J. ; Jakes, R. ; Walker, J.E. ; Milstein, C. ; Klug, A.** Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer's disease. *Proceedings of the National Academy of Sciences, USA*, 1988, vol. 85, 4506-4510 [0239]

专利名称(译)	与神经退行性疾病中的蛋白质聚集有关的材料和方法		
公开(公告)号	EP1348029B1	公开(公告)日	2012-03-07
申请号	EP2002734847	申请日	2002-01-02
申请(专利权)人(译)	大学评议阿伯丁大学		
当前申请(专利权)人(译)	WISTA LABORATORIES LTD.		
[标]发明人	WISCHIK CLAUDE MICHEL UNIV OF ABERDEEN RICKARD JANET ELIZABETH UNIV OF ABERDEEN HORSLEY DAVID UNIV OF ABERDEEN HARRINGTON CHARLES ROBERT UNIV OF ABERDEEN THEURING FRANZ UNIV HOSPITAL CHARITE STAMER KARSTEN UNIV HOSPITAL CHARITE ZABKE CLAUDIA WISTA LAB LTD		
发明人	WISCHIK, CLAUDE MICHEL, UNIVERSITY OF ABERDEEN RICKARD, JANET ELIZABETH, UNIVERSITY OF ABERDEEN HORSLEY, DAVID, UNIVERSITY OF ABERDEEN HARRINGTON, CHARLES ROBERT, UNIVERSITY OF ABERDEEN THEURING, FRANZ, UNIVERSITY HOSPITAL CHARITE STAMER, KARSTEN, UNIVERSITY HOSPITAL CHARITE ZABKE, CLAUDIA, WISTA LABORATORIES LTD.		
IPC分类号	C12N15/11 A01K67/027 A61K45/00 A61P25/28 A61P43/00 C07K14/47 C07K19/00 C12N5/10 C12N15/09 C12N15/12 C12N15/85 C12P21/02 G01N33/15 G01N33/50 G01N33/53 G01N33/68		
CPC分类号	A01K67/0275 A01K2207/15 A01K2217/00 A01K2217/05 A01K2217/20 A01K2227/105 A01K2267/0312 A01K2267/0318 A61P25/28 C07K14/4711 C07K2319/00 C12N15/8509 G01N33/6896 G01N2500/00		
优先权	2001000119 2001-01-03 GB		
其他公开文献	EP1348029A2		
外部链接	Espacenet		

摘要(译)

公开了诱导或模拟聚集疾病蛋白 (ADP-例如tau蛋白) 的病理状态的方法, 其与ADP通过诱导的构象聚合相互作用在病理上聚集的疾病状态 (例如阿尔茨海默病) 相关, 该方法是本发明的特征在于提供膜可局部化的融合蛋白的步骤, 所述融合蛋白包含 (i) 衍生自ADP的聚集部分, 或来自启动ADP病理聚集的蛋白质, (ii) 异源膜定位部分。据信基于ADP的融合蛋白的膜定位导致ADP蛋白的高亲和力捕获位点暴露, 从而促进可能是系统天然或异源的其他ADP的聚集。该方法可以在体外或在细胞和动物模型中进行, 并且可以通过监测聚集例如用于筛选聚集过程的调节剂。通过监测聚集产生的ADP相关降解产物的产生。还提供了用于该方法或与该方法一起使用的材料, 方法。




Figure 1a and 1b