



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
**29.12.2004 Bulletin 2004/53**

(51) Int Cl.<sup>7</sup>: **C07K 16/18, G01N 33/573,  
G01N 33/577, G01N 33/68**

(21) Application number: **00305736.1**

(22) Date of filing: **07.07.2000**

(54) **Antibody against cleavage product of vimentin**

Antikörper gegen Spaltprodukte von Vimentin

Anticorps dirigé contre un produit de clivage de la vimentine

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

(30) Priority: **07.07.1999 JP 19323599**

(43) Date of publication of application:  
**10.01.2001 Bulletin 2001/02**

(73) Proprietor: **Riken**  
**Wako-shi, Saitama 351-0198 (JP)**

(72) Inventors:  
• **Morishima, Nobuhiro, c/o Riken  
Wako-shi, Saitama 351-0198 (JP)**  
• **Nakanishi, Keiko  
Omiya-shi, Saitama 331-0052 (JP)**  
• **Shibata, Takehiko, c/o Riken  
Wako-shi, Saitama 351-0198 (JP)**

(74) Representative: **Maschio, Antonio, Dr. et al  
D Young & Co  
120 Holborn  
London EC1N 2DY (GB)**

(56) References cited:  
• **MORISHIMA NOBUHIRO: "Changes in nuclear  
morphology during apoptosis correlate with  
vimentin cleavage by different caspases located  
either upstream or downstream of Bcl-2 action."  
GENES TO CELLS, vol. 4, no. 7, July 1999  
(1999-07), pages 401-414, XP000952783 ISSN:  
1356-9597**

- **PRASAD SARADA ET AL: "Intermediate filament  
proteins during carcinogenesis and apoptosis  
(Review)." INTERNATIONAL JOURNAL OF  
ONCOLOGY, vol. 14, no. 3, March 1999 (1999-03),  
pages 563-570, XP000952788 ISSN: 1019-6439**
- **PATENT ABSTRACTS OF JAPAN vol. 1996, no.  
02, 29 February 1996 (1996-02-29) & JP 07 258296  
A (SCIENCE & TECH AGENCY), 9 October 1995  
(1995-10-09)**
- **SCHMIDT M ET AL: "A monoclonal antibody  
directed against the head region of vimentin"  
BIOCHEMICAL AND BIOPHYSICAL RESEARCH  
COMMUNICATIONS, vol. 146, no. 3, 1987, pages  
1366-1374, XP002150823 ISSN: 0006-291X**
- **PRASAD SARADA C ET AL:  
"Apoptosis-associated proteolysis of vimentin  
in human prostate epithelial tumor cells."  
BIOCHEMICAL AND BIOPHYSICAL RESEARCH  
COMMUNICATIONS, vol. 249, no. 2, 19 August  
1998 (1998-08-19), pages 332-338, XP002150824  
ISSN: 0006-291X**
- **HASHIMOTO MAKOTO ET AL: "Rapid  
fragmentation of vimentin in human skin  
fibroblasts exposed to tamoxifen: A possible  
involvement of caspase-3." BIOCHEMICAL AND  
BIOPHYSICAL RESEARCH COMMUNICATIONS,  
vol. 247, no. 2, 18 June 1998 (1998-06-18), pages  
401-406, XP002150825 ISSN: 0006-291X**
- **NAKANISHI K. ET AL: 'Identification of a  
caspase-9 substrate and detection of its  
cleavage in programmed cell death during  
mouse development' THE JOURNAL OF  
BIOLOGICAL CHEMISTRY vol. 276, no. 44, 02  
November 2001, pages 41237 - 41244**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**Description****FIELD OF THE INVENTION**

[0001] The present invention relates to an antibody against the cleavage product of vimentin; a method for detecting apoptosis using this antibody; and a use for this antibody.

**BACKGROUND OF THE INVENTION**

[0002] Apoptosis is cell death in multicellular organisms. Surplus cells generated during developmental processes, cells no longer needed in an adult, cells damaged by radiation or chemical substances, or dangerous cells such as tumor cells are led to cell death by apoptosis, thus removed from a body.

[0003] Caspases are proteolytic enzymes (protease) that play a key role during apoptosis. Research on apoptosis has rapidly expanded since the 1990s. One of the key factors to have promoted the research is the identification of a caspase family of proteases that involves execution of apoptosis (Thornberry, N. & Lazebnik, Y, (1998) Science, 281, 1312-1316). At least 10 or more members of the caspase family are identified in mammals. Caspases are also shown to be present as an inactive precursor in normal cells. When apoptosis is initiated for a cell to die, an initiator caspase in the caspase family activates itself by limited proteolysis (processing). The activated initiator caspase activates another caspase by partially cleaving it, the cleaved caspase activates another caspase, and the process continues one after another. This amplification cascade mechanism is thought to achieve the whole activation. All caspases cleave the C-terminal side of a specific aspartic acid residue in protein, but the cleavage efficiency of each member of the caspase family varies depending on amino acid sequences near the cleavage site.

[0004] Apoptosis triggered by the stimulation of the anti-Fas antibody is the best analyzed apoptosis and thought to play a central role among adults (Nagata, S. (1997) Cell 88, 355-365). Caspase-8, involving execution of apoptosis is first activated among the caspase family in cells after stimulation with the anti-Fas antibody, and functions as an initiator (Boldin, M. P., Goncharov, T.M., Goltsev, Y.V. & Wallach, D. (1996) Cell 85, 803-815; Muzio, M., Chinnaiyan, A. M., Kishkel, F.C., O'Rourke, K., Shevchenko, A., Ni, J., Scaffidi, C., Bretz, J. D., Zhang, M., Gentz, R., Mann, M., Krammer, P. H., Peter, M. E. & Dixit, V. M. (1996) Cell, 85, 817-827).

[0005] Several methods for detecting activation of a caspase have been employed, such as 1) detecting processing of a caspase or activation using an antibody recognizing a caspase; or 2) measuring protease activity using a substrate analog. Any of these methods, however, have a drawback in that the ability to distinguish between members of caspase family is limited. In the method of 1), production of a specific antibody capable

of recognizing both an inactive precursor and an active type is often difficult.

[0006] MORISHIMA ('Changes in nuclear morphology during apoptosis correlate with vimentin cleavage by different caspases located either upstream or downstream of Bcl-2 action' GENES TO CELLS 1999, vol. 4, pages 401-414) describes the cleavage of vimentin by caspase-8 that occurs at multiple sites, e.g. between Asp259 and Val260 of the protein at an early stage of apoptosis. The detection of vimentin cleavage in apoptotic cells is carried out by Western blot analysis using a monoclonal anti-vimentin antibody that interacts with both the intact protein and with its cleaved fragments.

**SUMMARY OF THE INVENTION**

[0007] It is the objective of the present invention to provide an antibody against the cleavage product of vimentin that is a main component of an intracellular skeletal protein; a method for detecting apoptosis using said antibody; and the use of said antibody.

[0008] As a result of intensive and extensive research toward the above-mentioned objective, the inventors have finally found that vimentin in an apoptotic cell is specifically cleaved by caspase-8, and they have succeeded in producing an antibody capable of detecting the specific cleavage of vimentin.

[0009] That is, the present invention relates to an antibody or a fragment thereof that recognises an N-terminal fragment of vimentin wherein, the N-terminal fragment comprises at least 5 amino acid residues from Asp259 to the N-terminus of vimentin, and wherein, the C-terminal fragment comprises at least 5 amino acid residues from Val260 to the C-terminus of vimentin, characterised in that the antibody or the fragment thereof reacts with the cleavage product of vimentin obtained by the cleavage of intact vimentin between Asp259 and Val260 but does not react with intact vimentin.

[0010] The cleavage product of said vimentin is one cleaved by action of caspase-8.

[0011] The abovementioned antibody can be either a polyclonal antibody or a monoclonal antibody

[0012] Said vimentin comprises the sequences set forth as SEQ ID No. 1, SEQ ID No. 2, SEQ ID No. 3, SEQ ID No. 4.

[0013] Further, the present invention relates to a method for preparing a monoclonal antibody or a fragment thereof that reacts with a cleavage product of vimentin that is cleaved between Asp259 and Val260 but does not react with intact vimentin, comprising the steps of:

(a) eliciting an immune response in a non-human mammal by immunisation with an antigen polypeptide comprising N- or C-terminal fragments of vimentin wherein, the N-terminal fragment at least 5 amino acid residues from Asp259 to the N-terminus of vimentin, and wherein, the C-terminal fragment

comprises at least 5 amino acid residues from Val260 to the C-terminus of vimentin;

(b) preparing a hybridoma from lymphoid cells of the host;

(c) selecting a hybridoma which produces a monoclonal antibody reacting with a cleavage product of vimentin but not reacting with intact vimentin and

(d) optionally, the monoclonal antibody is processed into a fragment thereof and

(e) obtaining the antibody or the fragment thereof.

**[0014]** A monoclonal antibody or fragment thereof obtainable by the beforementioned method.

**[0015]** Further the present invention relates to a method for preparing a polyclonal antibody that reacts with a cleavage product of vimentin that is cleaved between Asp259 and Val260 but does not react with intact vimentin, comprising the steps of:

(a) eliciting an immune response in a non-human mammal by immunisation with an antigen polypeptide comprising N- or C-terminal fragments of vimentin wherein, the N-terminal fragment at least 5 amino acid residues from Asp259 to the N-terminus of vimentin, and wherein, the C-terminal fragment comprises at least 5 amino acid residues from Val260 to the C-terminus of vimentin;

(b) obtaining anti-serum from the blood of the non-human mammal host; and

(c) obtaining the antibody or a fragment thereof.

**[0016]** A polyclonal antibody or fragment thereof obtainable by the before mentioned method.

**[0017]** The cleavage product of vimentin is obtained by the action of caspase-8 and vimentin comprises the sequence set forth as SEQ ID No. 1, SEQ ID No. 2, SEQ ID No. 3, SEQ ID No. 4.

**[0018]** Further the present invention relates to a method for detecting caspase activity *in vitro* or a method for detecting apoptosis *in vitro*, comprising allowing the above mentioned antibody or a fragment thereof to react with cleavage product of vimentin and detecting the resulting reaction product.

**[0019]** Furthermore, the present invention relates to a kit for detecting apoptosis comprising the above mentioned antibody or a fragment thereof.

## BRIEF DESCRIPTION OF THE DRAWINGS

**[0020]**

Figure 1 shows a schematic diagram of the cleavage site by caspase in vimentin.

Figures 2 and 3 show photographs of Western blot analysis.

Figure 4 shows photographs of indirect immunofluorescence staining of apoptotic cells.

## DETAILED DESCRIPTION OF THE INVENTION

**[0021]** The present invention relates to an antibody specifically recognizing cleavage product of vimentin but not reacting with the whole vimentin (intact vimentin), a substrate of a caspase (e.g., caspase-8) activated at the initial stage of apoptosis. To detect the activity of caspases, the present inventors have produced an antibody reacting with a cleavage site of a substrate cleaved by a caspase by focusing on changes in the substrate (vimentin) not on the caspase itself. Therefore, activation of a caspase can now be detected with high sensitivity regardless of sample scale, such as tissue and cell. Among amino acid sequences of vimentin protein, the present inventors have identified an aspartic acid residue at a specific site which is cleaved by caspase-8. Oligopeptides having a sequence on N-terminal side or C-terminal side of the aspartic acid residue are chemically synthesized and used for immunizing a rabbit. After several times of immunization, anti-serum is obtained. Then an antibody that binds strongly to the oligopeptide used for immunization in anti-serum is purified. The purified antibody does not bind to an intact vimentin protein, but specifically bind to a product cleaved by caspase-8. By using this antibody in the Western blot analysis and immunostaining of cells and tissues, the detection of activation of caspase-8 becomes possible through cleavage of vimentin either *in vitro*, in tissue sample or in cell sample.

**[0022]** The term "antibody" in the present invention represents the whole antibody molecule or its fragments (e.g., Fab or F(ab')<sub>2</sub> fragment), which can bind to the cleavage product of vimentin as an antigen. This antibody of this invention may be a polyclonal or monoclonal antibody.

**[0023]** Antibodies of the present invention can be produced by various methods. Such production methods of antibodies are known in the art. (For example, see Harlow E. & Lane D., *Antibody*, Cold Spring Harbor Laboratory Press (1988))

1. Preparation of antibodies reacting with cleavage products of vimentin

(1) Preparation of antigens

**[0024]** Vimentin on which caspases act is an intermediate filament protein, characteristics in mesenchymal cells, such as fibroblasts and leukocytes.

**[0025]** Figure 1 is a scheme showing a site of an intermediate filament protein, vimentin, to be cleaved by a caspase when apoptosis is initiated by treatment with anti-Fas antibody. Caspases used in the present invention include caspase-8. Caspase-8 recognizes and cleaves C terminal side of Asp-259 of an amino acid sequence of vimentin in a human or mouse (SEQ ID NO: 3 for a human, SEQ ID NO:4 for a mouse). According to the present invention, the two types of antibody can

be produced against vimentin protein or peptide which is cleaved by caspase-8 between Asp-259 and Val-260: an antibody recognizing the resulting N-terminal fragment (referred to as V1 antibody) and an antibody recognizing the resulting C-terminal fragment (referred to as V2 antibody). See Figure 1.

**[0026]** When the V1 antibody is produced, a protein or peptide fragment that has the maximum of 259 amino acid residues, preferably 6 amino acid residues, or more preferably 5 amino acid residues, from Asp-259 to the 1<sup>st</sup> Met on the N-terminus among an amino acid sequence shown in SEQ ID NO:3 or 4 can be used as an antigen. Similarly, when the V2 antibody is produced, a protein or peptide fragment that has the maximum of 207 amino acid residues, preferably 6 amino acid residues, more preferably 5 amino acid residues, from Val-260 to Glu-466 on the C-terminus among an amino acid sequence shown in SEQ ID NO: 3 or 4 can be used as an antigen. Further to improve the antigenicity, each of the above protein or peptide fragment is preferably allowed to perform coupling reaction at the terminus other than the one resulted from the cleavage by a caspase. It is preferable for facilitating the coupling reaction that a Cys residue is bound to the protein or peptide terminal.

## (2) Production of monoclonal antibodies against cleavage product of vimentin

### (i) Recovery of antibody-producing cells

**[0027]** Proteins or peptides as produced in (1) are administered as antigens to mammals, such as a rat, mouse, or rabbit. A dosage of the antigen per animal is 0.1 to 100 mg when no adjuvant is used, and 1 to 100 µg when an adjuvant is used. The adjuvants include Freund's complete adjuvant (FCA), Freund's incomplete adjuvant (FIA), and aluminum hydroxide adjuvant. Immunization is principally carried out by intravenous, subcutaneous, or intraperitoneal injection. In addition intervals of immunization are not particularly limited. Immunization is carried out for 1 to 10 times, preferably 2 to 5 times, at intervals of from several days to several weeks, preferably 2 to 5 week-intervals. One to 60 days, preferably 1 to 14 days after the final immunization, antibody-producing cells are recovered. Examples of antibody-producing cells include spleen cells, lymph node cells, peripheral blood cells, preferably spleen cells or local lymph node cells.

### (ii) Cell fusion

**[0028]** The antibody-producing cells are allowed to fuse with myeloma cells so as to obtain hybridomas. Myeloma cells to be fused with the antibody-producing cells may include generally available established cell lines of animals, such as a mouse. Preferable established cell lines used herein have drug selectivity and cannot survive in HAT selection medium containing hypoxanthine,

aminopterin, and thymidine when unfused, but can survive therein only when fused with the antibody-producing cells. Specific examples of myeloma cells include mouse myeloma cell lines, such as X63Ag.8.653, NS1/1-Ag4-1, and NS0/1; and rat myeloma cell lines, such as YB 2/0.

**[0029]** Next, the myeloma cells are fused with the antibody-producing cells. Briefly,  $1 \times 10^6$  to  $1 \times 10^7$  cells/ml of antibody-producing cells are mixed with  $2 \times 10^5$  to  $2 \times 10^6$  cells/ml of myeloma cells in an animal cell culture medium such as serum-free DMEM or RPMI-1640. Preferable cell ratio of the antibody-producing cells to myeloma cells is 2 : 1 to 3 : 1. Then, fusion reaction is performed in the presence of a cell fusion promoter. The cell fusion promoter is, for example polyethylene glycol of an average molecular weight 1,000 to 6,000 daltons. Further, a commercially available cell fusion device using electric pulse stimulation, e.g., electroporation can be used to fuse antibody-producing cells with myeloma cells.

### (iii) Selection and cloning of the Hybridomas

**[0030]** Hybridomas of interest are selected from the fused cells. First, cell suspension is appropriately diluted with such as RPMI-1640 medium containing fetal calf serum, about  $3 \times 10^5$  cells/well are placed in microtiter plate wells, selection medium is added to each wells, and the cells are cultured while appropriately replacing selection medium. Cells growing around 14 days after the start of cultures in selection medium can be obtained as hybridomas.

**[0031]** Next, supernatants from hybridoma cell cultures are examined by screening for the presence of antibody reacting with cleavage product of vimentin. Screening of hybridomas may be performed by any of the conventional methods and not particularly limited. For example, a part of the culture supernatant contained in a well in which a hybridoma has grown may be collected and subjected to screening using an enzyme immunoassay, or radioimmunoassay.

**[0032]** Cloning of fused cells is performed by the limiting dilution method or the like. Finally, hybridomas which are cells producing monoclonal antibodies that react with cleavage product of vimentin but not with intact vimentin are established.

### (iv) Recovery of monoclonal antibodies

**[0033]** Monoclonal antibodies can be recovered from established hybridomas using conventional methods, such as a cell culture method, and ascites formation method.

**[0034]** In the cell culture method, hybridomas are grown in culture media for animal cells, such as 10 % fetal calf serum-containing RPMI-1640 or MEM medium, or serum-free medium under conventional culture conditions (for example, at 37 °C, under 5 % CO<sub>2</sub>) for 7

to 14 days. Then antibodies are recovered from the culture supernatants.

**[0035]** In the ascites formation method, about  $1 \times 10^7$  cells of hybridomas are administered intraperitoneally to animals belonging to the same species of the mammals from which myeloma cells are derived so that a large number of hybridomas are grown. After 1 to 2 weeks, ascitic fluid is collected.

**[0036]** When purification of antibodies is required in the above described recovery methods, antibodies can be purified by known methods such as an ammonium sulfate method, ion exchange chromatography, gel filtration, and affinity chromatography. These methods may be used independently or in combination.

### (3) Production of polyclonal antibodies against the cleavage product of vimentin

**[0037]** Antigens prepared as described above are administered to mammals, such as a rat, mouse, and rabbit. A dosage of antigens per animal is 0.1 to 100 mg when no adjuvant is used, and 10 to 1,000 pg when adjuvant is used. The adjuvants used in this invention include Freund's complete adjuvant (FCA), Freund's incomplete adjuvant (FIA), and aluminum hydroxide adjuvant. Immunization is principally carried out by intravenous, subcutaneous, or intraperitoneal injection. Intervals of immunization are not particularly limited. Immunization is carried out for 1 to 10 times, preferably 2 to 5 times, at intervals of from a few days to a few weeks, preferably 2 to 5 week-intervals. Six to 60 days after the final immunization, antibody titer is measured using ELISA (enzyme-linked immunosorbent assay), EIA (enzyme immunoassay) or RIA (radioimmuno assay). On a day that the maximum antibody titer is shown, blood is collected to obtain anti-serum.

**[0038]** Subsequently, reactivity of polyclonal antibodies in anti-serum against the cleavage product of vimentin is measured by ELISA. Polyclonal antibodies showing strong reactivity against the cleavage product of vimentin but showing no reactivity against intact vimentin are selected.

**[0039]** For example, polyclonal antibodies in anti-serum are applied to an affinity column to which the cleavage product of vimentin is fixed, thereby recovering antibodies (column adsorption fraction) reacting with the cleavage product of vimentin. Then the resulting antibodies are applied to an affinity column fixed with intact vimentin, thereby recovering antibodies that do not adsorb but flow out are recovered. Finally obtained antibodies are subjected to ELISA to confirm whether they react with the cleavage product of vimentin but do not react with intact vimentin.

### 2. Method for detecting apoptosis

**[0040]** The above antibody may be used to detect (e.g., quantify) the cleavage product of vimentin in the

present invention. For example, a sample containing the cleavage product of vimentin may be incubated with the monoclonal antibody or polyclonal antibody of the present invention, followed by an anti-mouse IgG antibody labeled with an enzyme such as horseradish peroxidase (HRP). The amount of the cleavage product of vimentin in the sample may be determined by measuring the intensity of color developed during the enzymatic reaction using a measuring device. The measured value can be used as an indicator for detecting the activity of caspase. A greater value measured represents an increased activity of caspase, indicating that apoptosis of the test cell has progressed.

**[0041]** The antibody of the present invention may also be used to detect apoptosis by reacting the antibody with cells in a biological sample. For example, the sample to be assayed may be incubated with the above antibody. The cleavage product of vimentin present in the sample may be detected and quantified by using an anti-mouse IgG antibody labeled with horseradish peroxidase (HRP) according to conventional methods.

**[0042]** A measured value greater than negative control indicates a higher activity of caspase. On the contrary, a measured value less than positive control indicates a lower activity of caspase. These measured values can be used as data for determining the progression of apoptosis. For example, they may be used as indicators of the progression of autoimmune diseases such as systemic lupus erythematosus (SLE), autoimmune hemolytic anemia and Basedow's disease or acquired immunodeficiency syndrome (AIDS).

### 3. Reagent containing the antibody of the present invention

**[0043]** The antibody against the cleavage product of vimentin may be used as various reagents in the present invention. For example, when the antibody is used as a reagent for detecting the cleavage product of vimentin, the detection may be carried out using the procedure shown in Section 2 above. The reagent according to the present invention may include, for example, an anti-mouse IgG antibody labeled with HRP, an anti-rabbit IgG antibody labeled with HRP, a substrate for HRP and a buffer, as well as the above antibody.

**[0044]** Alternatively, when the antibody of the present invention is used as a reagent for immunohistochemical staining, the detection may be carried out according to a conventional procedure for immunohistochemical staining. In this case, the reagent according to the present invention may include, for example, an anti-mouse IgG antibody labeled with HRP, a fluorescently labeled anti-mouse IgG antibody, a substrate for HRP and a buffer, as well as the above antibody. For example, various tissue microscopic sections obtained by biopsy of SLE patient may be prepared by conventional methods to react with the antibody of the present invention. These sections may be incubated with an anti-mouse

IgG antibody labeled with horseradish peroxidase (HRP) as a secondary antibody and then treated with a substrate 3,3'-diaminobenzidine to develop brown. The brown area found in the section indicates that caspase has become active in this area.

## EXAMPLES

**[0045]** The present invention is further described in the following examples. These examples are provided for illustrative purposes only, and are not intended to limit the scope of the invention.

### Example 1: Preparation of Antibodies

#### (a) Materials

**[0046]** Peptides used for immunization were obtained from Biologica Co. (Nagoya, Japan). Activated hemocyanin (Imject Maleimide-KLH) was obtained from Pierce (Rockford, IL, USA). FMP-activated cellulofine was obtained from Seikagaku Corp. (Tokyo, Japan). Other biochemical reagents were obtained from Sigma-Aldrich Japan (Tokyo, Japan).

#### (b) Coupling of Chemically Synthesized Peptides to Carrier Proteins

**[0047]** Synthesized peptides (2 mg) were dissolved in Dulbecco's phosphate buffer (hereinafter referred to as PBS; Harlow E. & Lane, D. (1988) "Antibody" Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA). These peptides have the following sequences, respectively: Cys-Gln-Ile-Asp-Val-Asp (SEQ ID NO: 1; referred to as sequence V1) and Val-Ser-Lys-Pro-Asp-Cys (SEQ ID NO: 2; referred to as sequence V2).

**[0048]** Sequences V 1 and V2 correspond to five amino acid residues that are immediately N- and C-terminal to the cleavage site by caspase-8 in vimentin protein, respectively and each having Cys residue for coupling reaction at the terminus other than the cleavage site by caspase-8 (Figure 1). Sequences V1 and V2 were dissolved in 400  $\mu$ l and 200  $\mu$ l PBS, respectively. Each solution was mixed with 200  $\mu$ l of activated hemocyanin (Imject Maleimide-KLH) to start coupling reaction. Each reaction mixture was rotated gently at room temperature for 2.5 hours. The reaction product was dialyzed against PBS at 4 °C to remove unreacted peptides. The resulting peptide-carrier conjugate was diluted with PBS to a final volume of 5 ml.

#### (c) Immunization

**[0049]** The above peptide-carrier conjugate was mixed well with Freund's adjuvant (Harlow E. & Lane, D. (1988) "Antibody" Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA) and subcutaneously injected into a New Zealand white rabbit for im-

munization. The peptide-carrier conjugate containing 400  $\mu$ g of the peptide was used per injection. The rabbit was immunized for 4 times at days 1, 9, 22 and 31. The circulating blood was collected after 8 days of the final immunization to prepare antiserum by conventional methods.

#### (d) Preparation of Affinity Column for Purifying Specific Antibodies

**[0050]** 0.3 g of an activated resin (FMP-activated cellulofine) were swelled in distilled water (50 ml) and filled into Econo-Column (Bio-Rad, Hercules, CA, USA). The resin in the column was further washed with 10 ml distilled water. One milligram of the synthesized peptide (sequence V 1 or V2) was dissolved in 10 ml coupling buffer (50 mM Na<sub>2</sub>CO<sub>3</sub>/NaHCO<sub>3</sub>, pH 8.5) and mixed with the resin in the column to start coupling reaction. The coupling reaction continued overnight at 4 °C with rotation. The resin was further incubated with 20 ml blocking buffer (50 mM Tris-HCl, pH 8.0, 0.1 M monoethanolamine) at room temperature for 4 hours with rotation in order to block unreacted resin. Subsequently, unreacted peptides were removed by washing sequentially with the following solutions (20 ml each): 1) distilled water; 2) 0.1M Gly-HCl (pH 2.5); 3) distilled water; 4) washing buffer (20 mM Tris-HCl, pH 7.5, 1 M NaCl, 1% Triton X-100) and 5) distilled water.

**[0051]** The affinity column prepared was washed with 10 ml elution buffer (0.1 M Gly-HCl, pH 2.5), 10 ml distilled water, followed by 20 ml TBS (20 mM Tris-HCl, pH 7.5, 0.15 M NaCl) immediately before used for affinity purification.

#### (e) Affinity Purification of Specific Antibodies from Antisera

**[0052]** The antiserum (3 ml for sequence V1, 6 ml for sequence V2) was loaded on the above column and mixed well with affinity resin therein. The column was allowed to stand at room temperature for 1 hour to cause binding the antibodies to the resin. To remove non-specifically bound proteins, the column was washed sequentially with the following solutions: 1) 10 ml TBS; 2) 30 ml washing buffer (supra); 3) 30 ml TBS and 4) 10 ml of 0.15 M NaCl. The specific antibodies were eluted with 4 ml elution buffer (supra). The protein fraction eluted from the column was mixed immediately with 1 M Tris (0.2 ml) on ice to recover its pH to a neutral range.

**[0053]** Finally, 4 ml of anti-V1 antibody (2.75 mg/ml) and anti-V2 antibody (0.15 mg/ml) were obtained, respectively. These purified antibodies were divided into aliquots and stored frozen at -20 °C.

### Example 2: Western Blot Analysis Using Specific Antibodies

**[0054]** Human T cell line Jurkat or SKW6.4 ( $4 \times 10^5$

cells/ml) was induced to undergo apoptosis by treatment with 200 ng/ml anti-Fas antibody (Medical & Biological Laboratories Co., Ltd., Nagoya, Japan). The treatment continued for 0 to 8 hours at the longest duration. The cells ( $2 \times 10^6$  each) were collected by centrifugation (1,000 rpm/min, 5 min) and then suspended in PBS. The cells were centrifuged again under the same condition as defined above to remove culture medium in the sample. Each cell sample was ultrasonically treated for 20 seconds in 40  $\mu$ l sample buffer (62.5 mM Tris-HCl, pH 6.8, 6 M Urea, 2% sodium dodecylsulfate (SDS), 10% glycerol, 0.003% bromophenol blue) for SDS-polyacrylamide gel electrophoresis. Proteins in each cell lysate sample were separated on 12% polyacrylamide gel electrophoresis (Laemmli, U.K. (1970) Nature, 227, 680-685) and transferred onto a nitrocellulose membrane using Semidry Blot transfer device (Nihon EIDO, Tokyo, Japan) to form Western blots.

**[0055]** The Western blots were immunostained by enzyme chemiluminescence labelling method (ECL) (Harlow E. & Lane, D. (1988) "Antibody" Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA). A primary antibody was a commercially available anti-vimentin antibody V9 (8.8  $\mu$ g/ml, Sigma-Aldrich Japan, Tokyo, Japan) or the anti-V1 or V2 antibody (0.2  $\mu$ g/ml). A secondary antibody was an anti-mouse or rabbit IgG antibody labeled with horseradish peroxidase (1000-fold dilution, Cappel, Durham, NC, USA). Following incubation with the secondary antibody, the Western blots were visualized using ECL Plus kit (Amersham Japan, Tokyo, Japan). Immunostaining was carried out according to the manufacturer's instruction. The results are shown in Figs. 2 and 3.

**[0056]** Figure 2 shows Western blots of Jurkat cell lysate treated with the anti-Fas antibody. Following electrophoresis, the blots were immunostained with the anti-vimentin antibody (V9). Upon treatment with the anti-Fas antibody for several hours, caspases cleaved vimentin (58 kDa) at the maximum of three sites into fragments of different size.

**[0057]** When the lysate of cells undergoing apoptosis was subjected to Western blot analysis using the anti-V1 or V2 antibody, these antibodies specifically stained only vimentin fragments generated by cleavage of the sequence Ile-Asp-Val-Asp (Figure 3). The anti-V1 antibody detected 30 kDa and 19.5 kDa fragments, while the anti-V2 antibody detected 27 kDa and 21 kDa fragments because other caspases caused further fragmentation almost simultaneously with or subsequently to the fragmentation by caspase-8. On the other hand, neither the anti-V1 nor V2 antibody reacted with intact vimentin (58 kDa) that did not undergo cleavage (see Figure 3, Lane of hour 0). Hence, Figure 3 shows that the antibody of the present invention recognizes only vimentin fragments found in cells undergoing apoptosis.

### Example 3: Indirect Immunofluorescence Staining of Apoptotic Cells with Specific Antibodies

**[0058]** Human T cell line Jurkat ( $4 \times 10^5$  cells/ml) was induced to undergo apoptosis by treatment with 200 ng/ml anti-Fas antibody (Medical & Biological Laboratories Co., Ltd., Nagoya, Japan). Following the treatment for 6 hours, the cells ( $1.5 \times 10^6$ ) were collected by centrifugation (1,000 rpm/min, 5 min) and then suspended in PBS. The cells were centrifuged again under the same condition as defined above to remove culture medium in the sample.

**[0059]** The collected cells were fixed in cold methanol at  $-20^\circ\text{C}$  for 2 min. The fixed cells were washed with PBS and subjected to indirect immunofluorescence staining (Harlow E. & Lane, D. (1988) "Antibody" Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA). First, the cells were incubated with the anti-V1 or V2 antibody (0.2  $\mu$ g/ml, in PBS containing 3% bovine serum albumin) at room temperature for 1 hour and washed with PBS to remove unbound antibodies. Next, a fluorescently labeled anti-rabbit IgG antibody (Alexa 448, Molecular Probes, Eugene, Oregon, USA) was used as a secondary antibody, which was diluted 500-fold and added to the above cells for incubation at room temperature for 30 min. The cells were washed with PBS to remove unbound secondary antibodies. PBS had contained a DNA-staining reagent DAPI (0.1  $\mu$ g/ml, Sigma-Aldrich) in order to visualize nuclear chromosomes. The cells were then examined by Fluorescence Microscope IX70 (Olympus Optical Co., Ltd., Tokyo, Japan).

**[0060]** The results are shown in Figure 4. Figure 4 shows a photograph of the cells are fluorescently stained with the anti-V2 antibody. The apoptotic cells could be identified by condensation and fragmentation of their nucleus when visualized using DAPI (see small arrows in Figure 4). The anti-V2 antibody was found to specifically stain only cells undergoing apoptosis. Cells that do not undergo apoptosis (see large arrows in Figure 4) are not stained with the anti-V2 antibody. As a control, Jurkat cells untreated with the anti-Fas antibody were similarly examined, indicating that none of these cells were stained with the anti-V2 antibody.

**[0061]** The antibody of the present invention was shown to react only with the cleavage product of vimentin, but not with intact vimentin.

**[0062]** The present invention can achieve the detection of the actual cleavage of substrates by caspase regardless of sample scale such as protein, cell or tissue. Hence, the present invention enables one 1) to observe the conversion of caspase-8 from its precursor into its active form; 2) to find the active form of caspase-8 to actually function in place within the cell; and 3) to selectively distinguish the cells in which caspase-8 has become active in cell populations.

**[0063]** The present invention provides the antibody that react with the cleavage product of vimentin, but not

with intact vimentin. By using the antibody of the present invention, the initiator caspase that is in active form and is cleaving proteins can be identified by detecting the cleavage product of vimentin.

[0064] Vimentin is highly expressed in various developing tissues. Also, increased expression of vimentin is often observed when the tissue becomes cancerous. Accordingly, the antibody of the present invention can be a useful reagent for examining caspase activity in developing tissues or cancer cells, for detecting apoptosis, or for deciding therapeutic regimens for apoptosis-related disease.

#### Free text of Sequence Listing

#### [0065]

SEQ ID NO.: 1: Synthetic peptide

SEQ ID NO.: 2: Synthetic peptide

#### Claims

1. An antibody or a fragment thereof that recognises an N- or C-terminal fragment of vimentin
  - wherein, the N-terminal fragment comprises at least 5 amino acid residues from Asp259 to the N-terminus of vimentin,
  - and wherein, the C-terminal fragment comprises at least 5 amino acid residues from Val260 to the C-terminus of vimentin,
  - characterised in that** the antibody or the fragment thereof reacts with the cleavage product of vimentin obtained by the cleavage of intact vimentin between Asp259 and Val260 but does not react with intact vimentin.
2. The antibody or a fragment thereof according to claims 1 wherein the cleavage product of vimentin is obtained by the action of caspase-8.
3. The antibody or a fragment thereof or the fragment thereof according to claim 1 or claim 2 wherein vimentin comprises the sequence set forth as SEQ ID No.1, SEQ ID No.2, SEQ ID No.3 or SEQ ID No.4.
4. The antibody or the fragment thereof according to any one of claims 1 to 3, wherein the antibody is a polyclonal antibody or a monoclonal antibody.
5. A method for preparing a monoclonal antibody or a fragment thereof that reacts with a cleavage product of vimentin that is cleaved between Asp259 and Val260 but does not react with intact vimentin, comprising the steps of:
  - (a) eliciting an immune response in a non-human mammal by immunisation with an antigen polypeptide comprising N- or C-terminal fragments of vimentin
    - wherein, the N-terminal fragment comprises at least 5 amino acid residues from Asp259 to the N-terminus of vimentin,
    - and wherein, the C-terminal fragment comprises at least 5 amino acid residues from Val260 to the C-terminus of vimentin;
  - (b) preparing a hybridoma from lymphoid cells of the host;
  - (c) selecting a hybridoma which produces a monoclonal antibody reacting with a cleavage product of vimentin but not reacting with intact vimentin; and
  - (d) optionally, the monoclonal antibody is processed into a fragment thereof; and
  - (e) obtaining the antibody or the fragment thereof.
6. A method for preparing a polyclonal antibody that reacts with a cleavage product of vimentin which is cleaved between Asp259 and Val260 but does not react with intact vimentin, comprising the steps of:
  - (a) eliciting an immune response in a non-human mammal by immunisation with an antigen polypeptide comprising N- or C-terminal fragments of vimentin cleaved between Asp259 and Val260
    - wherein, the N-terminal fragment comprises at least 5 amino acid residues from Asp259 to the N-terminus of vimentin,
    - and wherein, the C-terminal fragment comprises at least 5 amino acid residues from Val260 to the C-terminus of vimentin;
  - (b) obtaining anti-serum from the blood of the non-human mammal host; and
  - (c) obtaining the antibody or the fragment thereof.
7. A method according to claim 6 wherein the cleavage product of vimentin is obtained by the action of a caspase-8.
8. A method according to claim 6 or claim 7 wherein vimentin comprises the sequence set forth as SEQ ID No. 1, SEQ ID No. 2, SEQ ID No.3 or SEQ ID No.4.
9. A monoclonal antibody or a fragment thereof obtainable by the method according to any one of

claim 5.

10. A polyclonal antibody or a fragment thereof obtainable by the method according to any one of claims 6 to 8.

11. A method for detecting caspase activity in vitro, wherein the caspase recognises and cleaves vimentin between Asp259 and Val260, comprising the steps of:

(a) allowing the antibody or the fragment thereof of any one of claims 1 to 4 to react with a cleavage product of vimentin; and

(b) detecting the resulting reaction product.

12. A method for detecting apoptosis in vitro comprising the steps of:

(a) allowing the antibody or the fragment thereof of any one of claims 1 to 4 to react with a cleavage product of vimentin that is cleaved between Asp259 and Val260 by a caspase; and

(b) detecting the resulting reaction product.

13. A kit for detecting apoptosis comprising the antibody or the fragment of any one of claims 1 to 4.

14. A kit according to claim 13 comprising a first antibody, a second antibody labelled with an enzyme, substrate and buffer.

#### Patentansprüche

1. Antikörper oder ein Fragment davon, der ein N- oder C-terminales Fragment von Vimentin erkennt, wobei das N-terminale Fragment wenigstens 5 Aminosäurereste von Asp259 bis zu dem N-Terminus von Vimentin umfaßt

und wobei das C-terminale Fragment wenigstens 5 Aminosäurereste von Val260 bis zu dem C-Terminus von Vimentin umfaßt,

**dadurch gekennzeichnet, daß** der Antikörper oder das Fragment davon mit dem Spaltprodukt von Vimentin, das durch die Spaltung von intaktem Vimentin zwischen Asp259 und Val260 erhalten wird, reagiert, aber nicht mit intaktem Vimentin reagiert.

2. Antikörper oder ein Fragment davon nach Anspruch 1, wobei das Spaltprodukt von Vimentin durch die Wirkung von Caspase-8 erhalten wird.

3. Antikörper oder ein Fragment davon oder das Fragment davon nach Anspruch 1 oder Anspruch 2, wobei Vimentin die als SEQ ID No. 1, SEQ ID No. 2,

SEQ ID No. 3 oder SEQ ID No. 4 angegebene Sequenz umfaßt.

4. Antikörper oder Fragment davon nach einem der Ansprüche 1 bis 3, wobei der Antikörper ein polyclonaler Antikörper oder ein monoklonaler Antikörper ist.

5. Verfahren zur Herstellung eines monoklonalen Antikörpers oder eines Fragmentes davon, der mit einem Spaltprodukt von Vimentin, das zwischen Asp259 und Val260 gespalten ist, reagiert, aber nicht mit intaktem Vimentin reagiert, mit den Stufen, in denen man

(a) durch Immunisierung mit einem antigenen Polypeptid mit N- oder C-terminalen Fragmenten von Vimentin eine Immunantwort in einem nicht-menschlichen Säuger auslöst, wobei das N-terminale Fragment wenigstens 5 Aminosäurereste von Asp259 bis zu dem N-Terminus von Vimentin umfaßt und wobei das C-terminale Fragment wenigstens 5 Aminosäurereste von Val260 bis zu dem C-Terminus von Vimentin umfaßt,

(b) ein Hybridom aus Lymphoidzellen des Wirts herstellt,

(c) ein Hybridom auswählt, welches einen monoklonalen Antikörper produziert, der mit einem Spaltprodukt von Vimentin aber nicht mit intaktem Vimentin reagiert, und

(d) optional den monoklonalen Antikörper zu einem Fragment davon verarbeitet und

(e) den Antikörper oder das Fragment davon gewinnt.

6. Verfahren zur Herstellung eines polyclonalen Antikörpers, der mit einem Spaltprodukt von Vimentin, welches zwischen Asp259 und Val260 gespalten ist, aber nicht mit intaktem Vimentin reagiert, mit den Stufen, in denen man

(a) durch Immunisierung mit einem antigenen Polypeptid, das N- oder C-terminale Fragmente von Vimentin, welches zwischen Asp259 und Val260 gespalten ist, umfaßt, eine Immunantwort in einem nicht-menschlichen Säuger auslöst, wobei das N-terminale Fragment wenigstens 5 Aminosäurereste von Asp259 bis zu dem N-Terminus von Vimentin umfaßt, und wobei das C-terminale Fragment wenigstens 5 Aminosäurereste von Val260 bis zu dem C-Terminus von Vimentin umfaßt,

- (b) Antiserum aus dem Blut des nicht-menschlichen Säugers gewinnt und
- (c) den Antikörper oder das Fragment davon gewinnt.
7. Verfahren nach Anspruch 6, wobei das Spaltprodukt von Vimentin durch die Wirkung von Caspase-8 erhalten wird.
8. Verfahren nach Anspruch 6 oder Anspruch 7, wobei Vimentin die als SEQ ID No. 1, SEQ ID No. 2, SEQ ID No. 3 oder SEQ ID No. 4 angegebene Sequenz umfaßt.
9. Monoklonaler Antikörper oder Fragment davon, erhältlich nach dem Verfahren gemäß Anspruch 5.
10. Polyklonaler Antikörper oder Fragment davon, erhältlich nach dem Verfahren gemäß einem der Ansprüche 6 bis 8.
11. Verfahren zum Detektieren von Caspase-Aktivität in vitro, wobei die Caspase Vimentin erkennt und zwischen Asp259 und Val260 spaltet, mit den Stufen, in denen man
- (a) den Antikörper oder das Fragment davon nach einem der Ansprüche 1 bis 4 mit einem Spaltprodukt von Vimentin reagieren läßt und
- (b) das resultierende Reaktionsprodukt feststellt.
12. Verfahren zum Feststellen von Apoptose in vitro mit den Stufen, in denen man
- (a) den Antikörper oder das Fragment davon nach einem der Ansprüche 1 bis 4 mit einem Spaltprodukt von Vimentin, das von einer Caspase zwischen Asp259 und Val260 gespalten ist, reagieren läßt und
- (b) das resultierende Reaktionsprodukt feststellt.
13. Kit für die Detektion von Apoptose, welches den Antikörper oder das Fragment nach einem der Ansprüche 1 bis 4 umfaßt.
14. Kit nach Anspruch 13, welches einen ersten Antikörper, einen zweiten Antikörper, der mit einem Enzym markiert ist, Substrat und Puffer umfaßt.
- Revendications**
1. Anticorps ou un de ses fragments qui reconnaît un
- fragment N- ou C-terminal de la vimentine  
le fragment N-terminal comprenant au moins 5 résidus d'acides aminés de Asp259 jusqu'à l'extrémité N-terminale de la vimentine,  
et le fragment C-terminal comprenant 5 résidus d'acides aminés de Val260 jusqu'à l'extrémité C-terminale de la vimentine,  
ledit anticorps ou son fragment étant **caractérisé en ce qu'il** réagit avec le produit de clivage de la vimentine obtenu par clivage de la vimentine intacte entre Asp259 et Val260, mais ne réagit pas avec la vimentine intacte.
2. Anticorps ou un de ses fragments suivant la revendication 1, le produit de clivage de la vimentine étant obtenu par l'action de la caspase-8.
3. Anticorps ou un de ses fragments ou son fragment suivant la revendication 1 ou la revendication 2, la vimentine comprenant la séquence indiquée en tant que SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3 ou SEQ ID N° 4.
4. Anticorps ou un de ses fragments suivant l'une quelconque des revendications 1 à 3, ledit anticorps étant un anticorps polyclonal ou un anticorps monoclonal.
5. Procédé pour la préparation d'un anticorps monoclonal ou d'un de ses fragments qui réagit avec le produit de clivage de la vimentine qui est clivée entre Asp259 et Val260 mais qui ne réagit pas avec la vimentine intacte, comprenant les étapes consistant :
- (a) à engendrer une réponse immunitaire chez un mammifère non humain par immunisation avec un polypeptide antigénique comprenant le fragment N-terminal ou le fragment C-terminal de la vimentine  
le fragment N-terminal comprenant au moins 5 résidus d'acides aminés de Asp259 jusqu'à l'extrémité N-terminale de la vimentine,  
et le fragment C-terminal comprenant au moins 5 résidus d'acides aminés de Val260 jusqu'à l'extrémité C-terminale de la vimentine ;  
(b) à préparer un hybridome à partir de cellules lymphoïdes de l'hôte ;  
(c) à sélectionner un hybridome qui produit un anticorps monoclonal réagissant avec un produit de clivage de la vimentine mais ne réagissant pas avec la vimentine intacte ; et  
(d) facultativement, à transformer l'anticorps monoclonal en un de ses fragments ; et  
(e) à obtenir l'anticorps ou son fragment.
6. Procédé pour la préparation d'un anticorps polyclonal qui réagit avec un produit de clivage de la vi-

mentine qui est clivée entre Asp259 et Val260 mais qui ne réagit pas avec la vimentine intacte, comprenant les étapes consistant :

(a) à engendrer une réponse immunitaire chez un mammifère non humain par immunisation avec un polypeptide antigénique comprenant un fragment N-terminal ou un fragment C-terminal de la vimentine clivée entre Asp259 et Val260,

le fragment N-terminal comprenant au moins 5 résidus d'acides aminés de Asp259 jusqu'à l'extrémité N-terminale de la vimentine,

et le fragment C-terminal comprenant au moins 5 résidus d'acides aminés de Val260 jusqu'à l'extrémité C-terminale de la vimentine ;

(b) à obtenir un antisérum à partir du sang du mammifère hôte non humain ; et

(c) à obtenir l'anticorps ou son fragment.

7. Procédé suivant la revendication 6, dans lequel le produit de clivage de la vimentine est obtenu par l'action de la caspase-8.

8. Procédé suivant la revendication 6 ou la revendication 7, dans lequel la vimentine comprend la séquence indiquée en tant que SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3 ou SEQ ID N° 4.

9. Anticorps monoclonal ou un de ses fragments pouvant être obtenu par le procédé suivant l'une quelconque des revendications 5 à 8.

10. Anticorps polyclonal ou un de ses fragments pouvant être obtenu par le procédé suivant l'une quelconque des revendications 6 à 8.

11. Méthode pour la détection de l'activité de caspase in vitro, dans laquelle la caspase reconnaît et clive la vimentine entre Asp259 et Val260, comprenant les étapes consistant :

(a) à laisser l'anticorps ou son fragment suivant l'une quelconque des revendications 1 à 4 réagir avec un produit de clivage de la vimentine ; et

(b) à détecter le produit de réaction résultant.

12. Méthode pour détecter une apoptose in vitro, comprenant les étapes consistant :


(a) à laisser l'anticorps ou son fragment suivant l'une quelconque des revendications 1 à 4 réagir avec un produit de clivage de la vimentine qui est clivée entre Asp259 et Val260 par une caspase ; et


(b) à détecter le produit de réaction résultant.

13. Kit pour la détection d'une apoptose, comprenant l'anticorps ou son fragment suivant l'une quelconque des revendications 1 à 4.

14. Kit suivant la revendication 13, comprenant un premier anticorps, un second anticorps marqué avec une enzyme, un substrat et un tampon.

Figure 1





Cleavage of vimentin in Jurkat cells undergoing apoptosis

**FIG. 2**





FIG. 3



Small arrows indicate nuclei of apoptotic cells.

Large arrows indicate nuclei of normal cells.

**FIG. 4**

专利名称(译)	抗波形蛋白裂解产物的抗体		
公开(公告)号	<a href="#">EP1067142B1</a>	公开(公告)日	2004-12-29
申请号	EP2000305736	申请日	2000-07-07
[标]申请(专利权)人(译)	独立行政法人理化学研究所		
申请(专利权)人(译)	RIKEN		
当前申请(专利权)人(译)	RIKEN		
[标]发明人	MORISHIMA NOBUHIRO C O RIKEN NAKANISHI KEIKO SHIBATA TAKEHIKO C O RIKEN		
发明人	MORISHIMA, NOBUHIRO, C/O RIKEN NAKANISHI, KEIKO SHIBATA, TAKEHIKO, C/O RIKEN		
IPC分类号	C12N15/09 C07K16/18 C07K19/00 C12N15/02 C12P21/08 C12Q1/04 C12Q1/37 G01N33/53 G01N33/531 G01N33/577 G01N33/573 G01N33/68		
CPC分类号	C07K16/18		
优先权	1999193235 1999-07-07 JP		
其他公开文献	EP1067142A1		
外部链接	<a href="#">Espacenet</a>		

摘要(译)

本发明提供了与波形蛋白的裂解产物反应但不与完整波形蛋白反应的抗体。

Figure 1

