



**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**10.04.2013 Bulletin 2013/15**

(21) Application number: **03743713.4**

(22) Date of filing: **28.02.2003**

(51) Int Cl.:  
**G01N 33/576 (2006.01)**

(86) International application number:  
**PCT/US2003/006038**

(87) International publication number:  
**WO 2003/073822 (12.09.2003 Gazette 2003/37)**

(54) **METHODS OF DIAGNOSING LIVER FIBROSIS**

VERFAHREN ZUR DIAGNOSE VON LEBERFIBROSE

PROCEDES DE DIAGNOSTIC DE FIBROSE HEPATIQUE

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PT SE SI SK TR**

(30) Priority: **28.02.2002 US 87188**

(43) Date of publication of application:  
**22.12.2004 Bulletin 2004/52**

(73) Proprietor: **Nestec S.A.**  
**1800 Vevey (CH)**

(72) Inventors:  
• **ROSE, Steven, L.**  
**Escondido, CA 92029 (US)**  
• **OH, Esther, H.**  
**San Diego, CA 92029 (US)**  
• **WALSH, Michael, J.**  
**San Diego, CA 92130 (US)**

(74) Representative: **Rupp, Christian et al**  
**Mitscherlich & Partner**  
**Patent- und Rechtsanwälte**  
**Sonnenstraße 33**  
**80331 München (DE)**

(56) References cited:  
**WO-A-00/52479 US-B1- 6 631 330**

- **OH E H ET AL: "CLINICAL LABORATORY PERFORMANCE OF THE FIBROSPECT SERODIAGNOSTIC TEST FOR THE DETECTION OF LIVER FIBROSIS" HEPATOLOGY, WILLIAMS AND WILKINS, BALTIMORE, MD, US, vol. 36, no. 4, PART 02, 1 October 2002 (2002-10-01), page 566A, ABSTR.NO.1611, XP008053458 ISSN: 0270-9139**

- **PATEL K ET AL: "EVALUATION AND OPTIMIZATION OF A PANEL OF SERUM MARKERS FOR LIVER FIBROSIS IN CHRONIC HEPATITIS C PATIENTS" GASTROENTEROLOGY, ELSEVIER, PHILADELPHIA, PA, vol. 123, no. 1, SUPPL, 1 July 2002 (2002-07-01), page 48, XP008053514 ISSN: 0016-5085**
- **ZAMAN A ET AL: "ASSESSMENT OF PANEL OF SERUM MARKERS FOR HEPATIC FIBROSIS IN PATIENTS WITH CHRONIC HEPATITIS C" HEPATOLOGY, WILLIAMS AND WILKINS, BALTIMORE, MD, US, vol. 4, no. PART 02, 1 October 2002 (2002-10-01), page 558A, ABSTR.NO.1580, XP008053459 ISSN: 0270-9139**
- **AFDHAL N.H. ET AL: 'Urinary assays for desmosine and hydroxylsypylpyridinoline in the detection of cirrhosis' JOURNAL OF HEPATOLOGY vol. 27, no. 6, 1997, pages 993 - 1002, XP002978908**
- **IMBERT-BISMUT F. ET AL: 'Biochemical markers of liver fibrosis in patients with hepatitis C virus infection: a prospective study' THE LANCET vol. 357, 07 April 2001, pages 1069 - 1075, XP004234215**
- **MURAWAKI Y. ET AL: 'Diagnostic value of serum markers of connective tissue turnover for predicting histological staging and grading in patients with chronic hepatitis C' JOURNAL OF GASTROENTEROLOGY vol. 36, 2001, pages 399 - 406, XP002978909**
- **XUHUAI J. ET AL: 'Clinical significance of serum 7S collagen and type VI collagen levels for the diagnosis of hepatic fibrosis' CHINESE MEDICAL JOURNAL vol. 110, no. 3, 1997, pages 198 - 201, XP001019695**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- WALSH K.M. ET AL: 'Basement membrane peptides as markers of liver disease in chronic hepatitis C' JOURNAL OF HEPATOLOGY vol. 32, 2000, pages 325 - 330, XP002978910

**Description****BACKGROUND OF THE INVENTION****FIELD OF THE INVENTION**

**[0001]** The present invention relates generally to the fields of hepatology and fibrosis and, more specifically, to a panel of serological markers which together are diagnostic of liver fibrosis.

**BACKGROUND INFORMATION**

**[0002]** Progressive fibrosis of the liver, kidney, lungs and other organs frequently results in organ failure that leads to organ transplantation or death, affecting millions in the United States and worldwide. Hepatic fibrosis, for example, is the leading non-malignant gastrointestinal cause of death in the United States, and the progression of fibrosis is the single most important determinant of morbidity and mortality in patients with chronic liver disease. Furthermore, the process of fibrosis is common to liver diseases of many etiologies, including chronic viral hepatitis B and C, autoimmune liver disease such as autoimmune hepatitis, alcoholic liver disease, fatty liver disease; primary biliary cirrhosis; and drug-induced liver disease. The fibrosis seen in these disorders results from chronic insults to the liver such as viral infection, alcohol or drugs.

**[0003]** Hepatitis C, for example, is one of the leading causes of chronic liver disease in the United States, where an estimated 3.9 million people are chronically infected with hepatitis C virus (HCV) and approximately 30,000 new cases of acute HCV occur each year (Alter, Semin. Liver Dis. 15:5-14 (1995)). The prevalence of hepatitis C is estimated to be 1.8% in the United States, with up to 10,000 deaths per year likely resulting from chronic hepatitis C infection (Alter, *supra*, 1995).

**[0004]** While hepatic fibrosis is a reversible process resulting in the accumulation of extracellular matrix, liver cirrhosis is an irreversible process characterized by thick bands of matrix which completely encircle the parenchyma to form nodules. Untreated, fibrosis of the liver leads to cirrhosis and eventually end-stage liver disease or cancer. Cirrhosis of the liver is a common condition that frequently goes undetected. For example, in a large sample of the general Danish population, the prevalence of liver cirrhosis was 4.5%, of which one-third were undiagnosed at the time of death (Graudal, J. Intern. Med. 230:165-171 (1991)).

**[0005]** Timely and accurate diagnosis of liver fibrosis is important to effective medical treatment. As an example, patients with hepatitis C and cirrhosis are less likely to respond to treatment with  $\alpha$ -interferon compared to patients with less advanced disease (Davis, Hepatology 26(Suppl. 1):122-127S). Similarly, treatments for chronic HCV infection can be contra-indicated in patients with histologically advanced and decompensated disease (NIH Consensus Development Conference Panel Statement, Hepatology 26 (Suppl. 1):25-105S (1997)). The importance of early diagnosis is further emphasized by the serious early complications such as variceal rupture that are associated with cirrhosis; these complications can be prevented by early detection of cirrhosis (Calés and Pasqual, Gastroenterol. Clin. Biol. 12:245-254 (1988)).

**[0006]** Diagnosis of the presence or severity of fibrotic liver disease is difficult, with liver biopsy currently the most reliable method available. Unfortunately, liver biopsy has several limitations: pain in about 30% of patients; the risk of severe complications such as hemorrhage or infection; a death rate of 3 in 10,000; and the cost of hospitalization (Nord, Gastrointest. Endosc. 28:102-104 (1982); Cadranet et al., Hepatology 32:47-481 (2000); and Poynard et al., Can. J. Gastroenterol. 14:543-548 (2000)). Furthermore, slowly progressive diseases such as hepatitis C require repeated biopsies for continual assessment of disease progression, thus compounding the risks and costs of the procedure. Finally, biopsy can fail to detect disease because of the heterogeneous distribution of pathological changes in the liver; it is not surprising, then, that false negatives are seen in a significant percentage of cases biopsied (Nord, *supra*, 1982).

**[0007]** For years there has been a search for biochemical or serological markers which reflect fibrotic processes in liver disease and which can serve as a surrogate for liver biopsy. However, the performance of any single marker has not been good enough to substitute for the biopsy procedure in detecting or staging fibrosis. Thus, there is a need for a non-invasive method of diagnosing the presence or severity of liver fibrosis. The present invention satisfies this need by providing a convenient and reliable method for detection of liver fibrosis that is suitable for serial testing. Related advantages are provided as well.

**SUMMARY OF THE INVENTION**

**[0008]** The present invention provides a method of diagnosing the presence or severity of liver fibrosis in an individual by detecting  $\alpha$ 2-macroglobulin ( $\alpha$ 2-MG) in a sample from the individual; detecting hyaluronic acid (HA) in a sample from the individual; detecting tissue inhibitor of metalloproteinases-1 (TIMP-1) in a sample from the individual; and diagnosing

the presence or severity of liver fibrosis in the individual based on the presence or level of  $\alpha$ 2-MG, HA and TIMP-1. A method of the invention can be useful, for example, for differentiating no or mild (F0-F1) liver fibrosis from moderate to severe (F2-F4) liver fibrosis.

**[0009]** The methods of the invention for diagnosing the presence or severity of liver fibrosis can be useful in a variety of patient populations including, but not limited to, those with viral hepatitis, autoimmune liver disease such as autoimmune hepatitis, alcoholic liver disease, fatty liver disease and drug-induced liver disease. In one embodiment, a method of the invention is used to diagnose the presence or severity of liver fibrosis in an individual infected with hepatitis C virus.

**[0010]** A variety of means can be useful for detecting  $\alpha$ 2-MG, HA and TIMP-1 in a sample. In one embodiment, the invention is practiced by determining the level of  $\alpha$ 2-MG protein in a sample from the individual to be diagnosed using, for example, one or more  $\alpha$ 2-MG-specific binding agents such as anti- $\alpha$ 2-MG antibodies. In another embodiment, a method of the invention is practiced by determining the level of  $\alpha$ 2-MG activity in a sample from the individual.

**[0011]** A variety of means also can be used in a method of the invention to detect hyaluronic acid in a sample. In one embodiment, the invention is practiced by determining the level of HA in a sample, for example, using one or more HA-specific binding agents such as HA-binding proteins (HABPs) or anti-HA antibodies.

**[0012]** Similarly, a variety of means can be used in a method of the invention to detect TIMP-1 in a sample. In one embodiment, the invention is practiced by determining the level of TIMP-1 protein in a sample from the individual to be diagnosed. The level of TIMP-1 protein can be determined, for example, using one or more TIMP-1-specific binding agents such as anti-TIMP-1 antibodies. In another embodiment, the invention is practiced by assaying for TIMP-1 activity in a sample from the individual to be diagnosed.

**[0013]** The invention provides, for example, a method of diagnosing the presence or severity of liver fibrosis in an individual by determining the level of  $\alpha$ 2-MG protein in a sample from the individual; determining the level of HA in a sample from the individual; and determining the level of TIMP-1 protein in a sample from the individual; and diagnosing the presence or severity of liver fibrosis in the individual based on the levels of  $\alpha$ 2-MG protein, HA and TIMP-1 protein. If desired, the level of  $\alpha$ 2-MG protein, HA and TIMP-1 protein each can be determined using an enzyme-linked assay.

**[0014]** A variety of samples can be useful in practising the methods of the invention including, for example, blood, serum, plasma, urine, saliva and liver tissue. In one embodiment, a single sample is obtained from the individual to be diagnosed. Such a sample can be, for example, a serum sample. Such a sample also can be, for example, a tissue sample, for example, a liver biopsy sample.

**[0015]** The present invention further provides a method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual. The method includes the steps of contacting an appropriate dilution of a sample from the individual with anti- $\alpha$ 2-MG antibody under conditions suitable to form a first complex of  $\alpha$ 2-MG and anti- $\alpha$ 2-MG antibody; washing the first complex to remove unbound molecules; determining the amount of  $\alpha$ 2-MG-containing first complex; contacting an appropriate dilution of a sample from the individual with a HA-binding protein under conditions suitable to form a second complex of HA and HA-binding protein; washing the second complex to remove unbound molecules; determining the amount of HA-containing second complex; contacting an appropriate dilution of a sample from the individual with anti-TIMP-1 antibody under conditions suitable to form a third complex of TIMP-1 and anti-TIMP-1 antibody; washing the third complex to remove unbound molecules; determining the amount of TIMP-1-containing third complex; and differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in the individual based on the amounts of  $\alpha$ 2-MG, HA and TIMP-1-containing complexes.

**[0016]** The methods of the invention can be practiced by detecting the three markers  $\alpha$ 2-MG, HA and TIMP-1, without detecting additional serological markers, or can be combined with a detection method for one or more additional markers. Thus, in one embodiment, the invention is practiced by detecting  $\alpha$ 2-MG, HA and TIMP-1 and also detecting at least one of the following markers of fibrosis: N-terminal procollagen III propeptide (PIIINP), laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 or apoB. In a further embodiment, the presence or severity of liver fibrosis is diagnosed by detecting  $\alpha$ 2-MG, HA, TIMP-1 and YKL-40 in a sample from an individual.

**[0017]** The present invention also provides a method of monitoring the efficacy of anti-fibrotic therapy in a patient by detecting  $\alpha$ 2-macroglobulin in a sample from a patient administered an anti-fibrotic therapy; detecting hyaluronic acid (HA) in a sample from the patient; detecting tissue inhibitor of metalloproteinases-1 (TIMP-1) in a sample from the patient; and determining the presence or severity of liver fibrosis in the patient based on the presence or level of  $\alpha$ 2-MG, HA and TIMP-1, thereby monitoring the efficacy of anti-fibrotic therapy. Such a method can further include, if desired, comparing the presence or severity of liver fibrosis determined in step (d) to the presence or severity of liver fibrosis in the patient at an earlier time. The methods of the invention can be used to monitor, for example, the progression or regression of fibrosis over time in a patient treated with one or more anti-fibrotic therapies, or to compare, for example, the efficacies of two or more anti-fibrotic therapies.

**[0018]** In one embodiment, at most three markers of fibrosis are detected. In another embodiment, the method includes the step of detecting in a sample from the patient at least one marker selected from the group consisting of: PIIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand,

TGF- $\beta$ 1, IL-10, apoA1, apoA2, and apoB.

**[0019]** A variety of means can be useful for detecting  $\alpha$ 2-MG, HA and TIMP-1 in a method of the invention. Step (a) can be practiced, for example, by determining the level of  $\alpha$ 2-MG protein in the sample. In one embodiment, the level of  $\alpha$ 2-MG protein is determined using one or more anti- $\alpha$ 2-MG antibodies. Step (b) can be practiced, for example, by determining the level of HA in the sample. In one embodiment, the level of HA is determined using one or more HA-binding proteins. Step (c) can be practiced, for example, by determining the level of TIMP-1 protein in said sample. In one embodiment, the level of TIMP-1 protein is determined using one or more anti-TIMP-1 antibodies.

**[0020]** Further provided herein is a method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual by determining an  $\alpha$ 2-MG level in a sample from the individual; determining a HA level in a sample from the individual; determining a TIMP-1 level in a sample from the individual; and diagnosing the individual as having no or mild liver fibrosis when the  $\alpha$ 2-MG level is below an  $\alpha$ 2-MG cut-off value X1, the HA level is below a HA cut-off value Y1 or the TIMP-1 level is below a TIMP-1 cut-off value Z1; diagnosing the individual as having moderate to severe liver fibrosis when the  $\alpha$ 2-MG level is above an  $\alpha$ 2-MG cut-off value X2, the HA level is above a HA cut-off value Y2 and the TIMP-1 level is above a TIMP-1 cut-off value Z2; and diagnosing remaining individuals as having an indeterminate status.

**[0021]** The methods of the invention based on dual cut-off values for the levels of the  $\alpha$ 2-MG, HA and TIMP-1 markers can be useful in differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in a variety of patient populations. The methods of the invention can be useful, for example, in diagnosing an individual having a liver disease such as viral hepatitis, autoimmune liver disease such as autoimmune hepatitis, alcoholic liver disease, fatty liver disease or drug-induced liver disease. In one embodiment, the methods of the invention are used to differentiate no or mild liver fibrosis from moderate to severe liver fibrosis in an individual infected with hepatitis C virus. Samples useful in the methods of the invention include, but are not limited to, blood, serum, plasma, urine, saliva and liver tissue. In one embodiment, a method of the invention is practiced by determining the  $\alpha$ 2-MG level, HA level and TIMP-1 level in one or more serum samples from the individual to be diagnosed.

**[0022]** Thus, the present invention provides, for example, a method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual in which the differentiation is based on an X1 cut-off value between 1.8 and 2.2 mg/ml; a Y1 cut-off value between 31 and 39 ng/ml; a Z1 cut-off value between 900 and 1100 ng/ml; an X2 cut-off value between 1.8 and 2.2 mg/ml; a Y2 cut-off value between 54 and 66 ng/ml; and a Z2 cut-off value between 1415 and 1735 ng/ml. In a particular embodiment, the differentiation is based on an X1 cut-off value of 2.0 mg/ml; a Y1 cut-off value of 35 ng/ml; a Z1 cut-off value of 1000 ng/ml; an X2 cut-off value of 2.0 mg/ml; a Y2 cut-off value of 60 ng/ml; and a Z2 cut-off value of 1575 ng/ml. In another embodiment, the differentiation is based on an X1 cut-off value of 2.0 mg/ml; a Y1 cut-off value of 37 ng/ml; a Z1 cut-off value of 1100 ng/ml; an X2 cut-off value of 2.0 mg/ml; a Y2 cut-off value of 60 ng/ml; and a Z2 cut-off value of 1575 ng/ml. In a further embodiment, X1, Y1, Z1, X2, Y2 and Z2 are selected such that, in a population having up to 30% liver fibrosis prevalence, at least 65% of individuals in the population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with an accuracy of at least 90%. In another embodiment, X1, Y1, Z1, X2, Y2 and Z2 are selected such that, in a population having up to 30% liver fibrosis prevalence, at least 65% of individuals in said population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with a positive predictive value of at least 90% and a negative predictive value of at least 90%. In yet a further embodiment, X1, Y1, Z1, X2, Y2 and Z2 are selected such that, in a population having up to 10% fibrosis prevalence, at least 70% of individuals in the population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with an accuracy of at least 90%.

**[0023]** Also disclosed is a method of diagnosing the presence or severity of liver fibrosis in an individual by comparing a level of a first fibrotic marker X in the individual to a cut-off value X1 to determine whether the individual is positive for the first fibrotic marker X; comparing a level of a second fibrotic marker Y in the individual to a cut-off value Y1 to determine whether the individual is positive for the second fibrotic marker Y; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X and Y, where, in a population with up to 40% fibrosis prevalence, at least 65% of individuals in the population are diagnosed with an accuracy of at least 90%.

**[0024]** The method can include, if desired, comparing a level of a third fibrotic marker Z in the individual to a cut-off value Z1 to determine whether the individual is positive for the third fibrotic marker Z and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X, Y and Z. The first fibrotic marker is  $\alpha$ 2-MG, the second fibrotic marker is HA, and the third fibrotic marker is TIMP-1.

**[0025]** The levels of these at least three fibrotic markers are compared, and, in a further embodiment, the levels of exactly three fibrotic markers are compared. In additional embodiments, the levels of at least four or at least five fibrotic markers are compared. A method of the invention can be useful, for example, to differentiate no or mild liver fibrosis from moderate to severe liver fibrosis.

**[0026]** In a specific embodiment, a method of the invention serves to diagnose at least 65% of individuals in a population with up to 30% fibrosis prevalence with an accuracy of at least 93%. In a further embodiment, a method of the invention serves to diagnose at least 70% of individuals in a population with up to 20% fibrosis prevalence with an accuracy of at least 94%. In yet a further embodiment, a method of the invention serves to diagnose at least 70% of individuals in a

population with up to 10% fibrosis prevalence with an accuracy of at least 96%.

[0027] The present invention further provides a method of diagnosing the presence or severity of liver fibrosis in an individual by comparing a level of a first fibrotic marker X in the individual to a cut-off value X1 to determine whether the individual is positive for the first fibrotic marker X; comparing a level of a second fibrotic marker Y in the individual to a cut-off value Y1 to determine whether the individual is positive for the second fibrotic marker Y; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X and Y, where the cut-off values X1 and Y1 are optimized individually to give a desired performance characteristic, wherein the first fibrotic marker is  $\alpha$ 2-MG, the second fibrotic marker is HA, and the third fibrotic marker is TIMP-1. The method of the invention

[0028] includes the steps of comparing a level of a third fibrotic marker Z in the individual to a cut-off value Z1 to determine whether the individual is positive for the third fibrotic marker Z and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X, Y and Z, where the cut-off values X1, Y1 and Z1 are optimized individually to give a desired performance characteristic, wherein, levels of  $\alpha$ 2-MG, HA and TIMP-1 are compared. In another embodiment, the cut-off values are optimized using design of experiments (DOE) analysis. In further embodiments, the levels of exactly three, at least three, at least four, or at least five fibrotic markers are compared. A method of the invention can be useful, for example, in differentiating no or mild liver fibrosis from moderate to severe liver fibrosis.

[0029] Further provided by the invention is a method of diagnosing the presence or severity of liver fibrosis in an individual by comparing a level of the first fibrotic marker X in the individual to two cut-off values X1 and X2 to determine whether the individual is positive for the first fibrotic marker X; comparing a level of the second fibrotic marker Y in the individual to two cut-off values Y1 and Y2 to determine whether the individual is positive for the second fibrotic marker Y; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X and Y, where the cut-off values X1, Y1, X2 and Y2 are optimized individually to give a desired performance characteristic. A method of the invention further includes the steps of comparing a level of the third fibrotic marker Z in the individual to two cut-off values Z1 and Z2 to determine whether the individual is positive for the third fibrotic marker Z; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X, Y and Z, where the cut-off values X1, Y1, Z1, X2, Y2 and Z2 are optimized individually to give a desired performance characteristic. Cut-off values can be conveniently optimized, for example, using DOE analysis.

## **BRIEF DESCRIPTION OF THE DRAWINGS**

### **[0030]**

Figure 1 shows the nucleic acid sequence (SEQ ID NO: 1) and corresponding amino acid sequence (SEQ ID NO: 2) for mature human  $\alpha$ 2-macroglobulin available from Genbank accession M36501.

Figure 2 shows the nucleic acid sequence (SEQ ID NO: 3) and corresponding amino acid sequence (SEQ ID NO: 4) for human tissue inhibitor of metalloproteinases-1 (TIMP-1) available from Genbank accession NM\_003254.

## **DETAILED DESCRIPTION OF THE INVENTION**

[0031] As disclosed herein, the serum levels of a number of biochemical markers were analyzed in a patient population with confirmed hepatitis C and having a known Metavir stage (fibrosis score) of F0 to F4, where F0 represents very low or no fibrosis; F1, F2 and F3 represent intermediate fibrosis stages; and F4 represents severe fibrosis (Knodell et al., Hepatology 1:431-435 (1981)). See Tables 2 and 3. Using Design of Experiments (DOE) analysis for simultaneous variation of multiple cut-off values, a four-marker panel made up of hyaluronic acid (HA), PIIINP, collagen type IV and  $\alpha$ 2-macroglobulin ( $\alpha$ 2-MG) was identified which was capable of differentiating F0-F1 (no or mild) fibrosis from F2-F4 (moderate to severe) fibrosis with an accuracy of about 77% in a patient population with a fibrosis prevalence of 60%.

[0032] As further disclosed herein in Example I, two three-marker panels,  $\alpha$ 2-MG/HA/TIMP-1 and  $\alpha$ 2-MG/HA/YKL-40, also performed well in differentiating F0-F1 fibrosis from F2-F4 fibrosis when cut-offs were optimized using DOE analysis. In particular, the  $\alpha$ 2-MG/HA/TIMP-1 and  $\alpha$ 2-MG/HA/YKL-40 panels each performed better than the four-marker panel and were capable of differentiating F0-F1 from F2-F4 fibrosis with about 80% accuracy in the study population. As can be seen in Table 6, line 15, for example, the  $\alpha$ 2-MG/HA/TIMP-1 panel performed with a sensitivity of 83.48% and a specificity of 75.95% in the study population having 60% fibrosis prevalence. These results demonstrate that the  $\alpha$ 2-MG/HA/TIMP-1 three-marker panel can be useful for differentiating no or mild fibrosis from moderate to severe fibrosis.

[0033] Based on these findings, the present invention provides a method of diagnosing the presence or severity of liver fibrosis in an individual by detecting  $\alpha$ 2-MG in a sample from an individual; detecting HA in a sample from the individual; detecting TIMP-1 in a sample from the individual; and diagnosing the presence or severity of liver fibrosis in the individual based on the presence or levels of  $\alpha$ 2-MG, HA and TIMP-1. A method of the invention can be useful, for

example, for differentiating no or mild (F0-F1) liver fibrosis from moderate to severe (F2-F4) liver fibrosis.

#### Liver and other fibrotic disorders

**[0034]** The methods of the invention can be useful for diagnosing the presence or severity of liver fibrosis in a variety of individuals including those at risk for, or having one or more symptoms of, a liver disorder characterized by fibrosis. The methods of the invention can be used to diagnose liver fibrosis in an individual having, for example, viral hepatitis such as hepatitis A, B or C virus or a human immunodeficiency virus (HIV) such as HIV-1; chronic persistent hepatitis or chronic active hepatitis; autoimmune liver disease such as autoimmune hepatitis; alcoholic liver disease; fatty liver disease; non-alcoholic liver disease including non-alcoholic fatty liver disease and non-alcoholic steatohepatitis (NASH); primary biliary cirrhosis; primary sclerosing cholangitis, biliary atresia; liver disease resulting from medical treatment (drug-induced liver disease); or a congenital liver disease. The methods of the invention can be extremely useful, for example, in alleviating concerns of potential liver damage due to methotrexate treatment. Periodic monitoring of liver fibrosis in individuals treated with methotrexate or other drugs associated with risk of liver damage can be conveniently performed using the non-invasive methods of the invention, without the risks associated with liver biopsy.

**[0035]** In one embodiment, the methods of the invention are useful for differentiating individuals having a Metavir score of F0 or F1 from individuals having a Metavir score of F2, F3 or F4. Metavir scoring is a well accepted system for grading liver biopsy specimens and is described in Knodell, *supra*, 1981. F0 is equivalent to the absence of fibrosis; F1 signifies portal fibrosis without septa. F2 signifies portal fibrosis with a few septa. F3 signifies numerous septa without cirrhosis. F4 signifies cirrhosis.

**[0036]** It is understood that the diagnostic methods of the invention are applicable to a variety of individuals including individuals with chronic or active disease, individuals with one or more symptoms of fibrotic disease, asymptomatic or healthy individuals and individuals at risk for one or more fibrotic diseases. It further is clear to the skilled person that the methods of the invention can be useful, for example, to corroborate an initial diagnosis of disease or to gauge the progression of fibrosis in an individual with a previous definitive diagnosis of fibrotic disease. The methods of the invention can be used to monitor the status of fibrotic disease over a period of time and further can be used, if desired, to monitor the efficacy of therapeutic treatment. If desired, the results obtained from a sample from an individual undergoing therapy can be compared, for example, to the individual's baseline results prior to treatment, to results earlier during treatment, or to a historic or reference value.

**[0037]** The methods of the invention are useful for diagnosing the severity of liver in an individual. Thus, the methods of the invention can be useful for determining the "stage" or extent of liver. In one embodiment, a method of the invention is used to determine the Metavir score of an individual, for example, an individual with viral hepatitis C. As indicated above, Metavir scoring is a well established fibrosis scoring system using values of F0 (absence), F1 (portal fibrosis without septa), F2 (portal fibrosis with few septa), F3 (portal fibrosis with numerous septa in the absence of cirrosis) and F4 (cirrhosis). In other embodiments, a method of the invention is used to determine the Knodell score (histological activity index), Ishak score (modified histological activity index), or Scheuer classification of an individual, for example, an individual with viral hepatitis C. In a further embodiment, a method of the invention is used to determine the severity of liver fibrosis in an individual with NASH, for example, by determining the severity according to the Brunt proposal (Brunt et al., Am. J. Gastroenterol. 94:2467-2474. (1999)). It is understood that, where the severity of liver or other fibrosis is determined according to a method of the invention, any of the above or other art-accepted or clearly defined scoring systems can be useful in reporting results indicating the severity of fibrosis.

#### Samples

**[0038]** A variety of samples can be useful in practicing the methods of the invention including, for example, blood, serum, plasma, urine, saliva and liver tissue. In one embodiment, a single sample is obtained from the individual to be diagnosed. Such a sample can be, for example, a serum sample.

**[0039]** As used herein, the term "sample" means a biological specimen that contains one or more fibrotic markers such as  $\alpha$ 2-MG, HA or TIMP-1. A sample can be, for example, a fluid sample such as whole blood, plasma, saliva, urine, synovial fluid or other bodily fluid, or a tissue sample such as a lung, liver, kidney, prostate or breast tissue sample. One skilled in the art understands that fluid samples can be diluted, if desired, prior to analysis.

**[0040]** One skilled in the art understands that a single sample can be obtained from the individual to be diagnosed and can be subdivided prior to detecting  $\alpha$ 2-MG-, HA- and TIMP-1. One skilled in the art also understands that, if desired, two or more samples can be obtained from the individual to be diagnosed and that the samples can be of the same or a different type. In one embodiment,  $\alpha$ 2-MG-, HA- and TIMP-1 each are detected in serum samples. In another embodiment, a single serum sample is obtained from an individual and subdivided prior to detecting  $\alpha$ 2-MG-, HA- and TIMP-1.

$\alpha$ 2-macroglobulin

**[0041]** The methods of the invention rely, in part, on detecting  $\alpha$ 2-macroglobulin in a sample.  $\alpha$ 2-MG is a conserved, highly abundant component of plasma that functions as a broad spectrum protease-binding protein to clear active proteases from tissue fluids. Unlike active site protease inhibitors, members of the  $\alpha$ 2-macroglobulin family do not inactivate the catalytic activity of their protease substrates but act by physical entrapment of the target protease within the folds of the  $\alpha$ 2-MG family member.  $\alpha$ 2-MG is itself cleaved by target proteases; reorganization of the  $\alpha$ 2-MG molecule results in sequestering of the target protease within an internal pocket of the  $\alpha$ 2-MG molecule (Starkey et al., Biochem. J. 131:823-831 (1973)). While an  $\alpha$ 2-MG entrapped protease is sterically prevented from interacting with macromolecular substrates such as proteins, it remains active against low molecular mass substrates, such as amide and ester compounds, able to diffuse into the  $\alpha$ 2-MG cage to access the enzymatic site. Thus,  $\alpha$ 2-MG activity is characterized, in part, by the ability to inhibit proteolytic activity but not amidolytic activity of a protease substrate.  $\alpha$ 2-MG also is characterized by the ability to shield entrapped proteases from antibodies and high molecular mass active site inhibitors. For example; trypsin bound by  $\alpha$ 2-MG is protected from inhibition by soybean trypsin inhibitor (STI).

**[0042]** In contrast to the restricted specificity of active-site protease inhibitors,  $\alpha$ 2-MG acts on a broad spectrum of proteases with diverse substrate specificity and catalytic activity. Such target proteases include trypsin, subtilisin, chymotrypsin, plasmin, elastase, thermolysin and papain. Substrate diversity is determined, in part, by the  $\alpha$ 2-MG "bait" region, a highly flexible and solvent-exposed sequence of 30-40 residues that contains at least one site sensitive to cleavage by each of the major classes of proteolytic enzyme.

**[0043]** As used herein, the term " $\alpha$ 2-macroglobulin" is synonymous with " $\alpha$ 2-MG" and means a protein with significant structural homology to human  $\alpha$ 2-MG (SEQ ID NO: 2) and having broad spectrum protease inhibitory activity.  $\alpha$ 2-MG contains a unique thiol ester bond that is inactivated by small primary amines such as methylamine. Thus,  $\alpha$ 2-MG activity can be characterized, in part, by methylamine-sensitive protease inhibitory activity.  $\alpha$ 2-MG can be distinguished, if desired, from other members of the  $\alpha$ 2-macroglobulin family such as related protease-binding proteins and C3, C4 and C5 of the complement system (Sottrup-Jensen, " $\alpha$ 2-Macroglobulin and Related Thiol Ester Plasma Proteins," in Putnam (Ed.), The Plasma Proteins: Structure, Function and Generic Control Second edition, Orlando: Academic Press (1987), pages 191-291. It is understood that an assay for detecting  $\alpha$ 2-MG can be specific for  $\alpha$ 2-MG or can additionally detect one or more other members of the  $\alpha$ 2-macroglobulin family.

**[0044]** The methods of the invention rely, in part, on detecting  $\alpha$ 2-macroglobulin in a sample. As used herein, the phrase "detecting  $\alpha$ 2-MG" means any quantitative or qualitative assay for determining the presence of  $\alpha$ 2-MG. As used herein, the phrase "determining the level of  $\alpha$ 2-MG" means any direct or indirect quantitative assay for  $\alpha$ 2-MG.

**[0045]** Similarly, detecting any specified fibrotic marker in a sample means determining whether the marker is present in the sample, said fibrotic marker having a positive or negative correlation with liver fibrosis or with another fibrotic disorder such as are described herein above. It is understood that detection can refer to non-quantitative analysis, for example, the presence or absence of a particular trait, variable or biochemical or serological substance.

**[0046]** Diagnosis is based on analyzing the sample for the presence or level of the fibrotic marker or other characteristic and comparing it to a reference value, where the reference value serves to assist in differentiating those with a fibrotic disorder from other individuals. Where the fibrotic marker is a biochemical or serological marker, determining a "level" in a sample means quantifying the fibrotic marker by determining, for example, the relative or absolute amount of RNA, protein or activity of the fibrotic marker. Thus, determining a level in a sample encompasses, without limitation, analysis of relative and absolute RNA, protein and activity levels as well as other direct and indirect measurements of the fibrotic marker as discussed further below. It is understood that any assay useful for determining a "level" of a fibrotic marker also is useful for "detecting" the marker.

**[0047]** A variety of assays for detecting  $\alpha$ 2-MG are known in the art and include direct and indirect assays for  $\alpha$ 2-MG RNA,  $\alpha$ 2-MG protein and  $\alpha$ 2-MG activity.  $\alpha$ 2-MG can be detected, or an  $\alpha$ 2-MG level can be determined, for example, by analysis of  $\alpha$ 2-MG mRNA levels using routine techniques such as Northern analysis or RT-PCR, or other methods based on hybridization to a nucleic acid sequence that is complementary to a portion of the  $\alpha$ 2-MG coding sequence. For example, conditions and probes for Northern analysis and RNA slot blot hybridization of  $\alpha$ 2-MG RNA in human samples are described in Ortego et al., Exp. Eye Res. 65:289-299 (1997), and Simon et al., Cancer Res. 56:3112-3117 (1996), respectively.

**[0048]**  $\alpha$ 2-MG also can be detected, or an  $\alpha$ 2-MG level can be determined, by assaying for  $\alpha$ 2-MG protein by a variety of methods. Immunoassays, including radioimmunoassays, enzyme-linked immunoassays and two-antibody sandwich assays as described further below, are useful in the methods of the invention. For example, in nephelometry assays, complexes of  $\alpha$ 2-MG and anti- $\alpha$ 2-MG antibody result in increased light scatter that is converted to a peak rate signal, which is a function of the sample  $\alpha$ 2-MG concentration.  $\alpha$ 2-MG also can be detected, for example, by laser immunonephelometry using a Behring Nephelometer Analyzer (Fink et al., J. Clin. Chem. Clin. Biol. Chem. 27:261-276 (1989)) and rabbit anti-human  $\alpha$ 2-MG antiserum as described in Naveau et al., Dig. Diseases Sci. 39:2426-2432 (1994), or using the nephelometry assay commercially available from Beckman Coulter (Brea, CA; kit #449430). Furthermore,



monoclonal and polyclonal anti- $\alpha$ 2-MG antibodies useful in immunoassays can be readily obtained from a variety of sources. As examples, affinity purified goat anti-human  $\alpha$ 2-MG and peroxidase-labeled goat anti-human  $\alpha$ 2-MG antibodies suitable for immunoassays such as ELISA assays and western blotting are available from Cedarlane Laboratories Limited (Ontario, Canada; CL20010AP and CL20010APHP) and Affinity Biologicals Incorporated (Ontario, Canada; GAA2M-AP and GAA2M-APHRP). Levels of  $\alpha$ 2-MG protein also can be determined by quantifying the amount of purified  $\alpha$ 2-MG protein. Purification of  $\alpha$ 2-macroglobulin can be achieved, for example, by HPLC, alone or in combination with mass spectrophotometry, or as described, for example, in Hall and Roberts, *Biochem. J.* 171:27-38 (1978) or Imber and Pezzo, *J. Biol. Chem.* 256:8134-8139 (1981). Quantitation can be determined by well known methods including Bradford assays, Coomassie blue staining and assays for radiolabeled protein.

**[0049]** A variety of assays for  $\alpha$ 2-MG activity also can be useful for detecting  $\alpha$ 2-MG or determining a level of  $\alpha$ 2-MG in a sample according to a method of the invention.  $\alpha$ 2-MG can be detected or a level of  $\alpha$ 2-MG can be determined indirectly, for example, as a function of inhibition of target protease activity, without a corresponding inhibition of amidolytic activity. As discussed above,  $\alpha$ 2-MG-bound proteases retain the ability to hydrolyze amide and ester bonds of small substrates, even while high molecular mass substrates such as proteins cannot be hydrolyzed (see, for example, Armstrong et al., *Develop. Compar. Immunol.* 23:375-390 (1999)). As an example,  $\alpha$ 2-MG can be detected or the level of  $\alpha$ 2-MG can be determined by assaying for inhibition of trypsin, subtilisin, chymotrypsin, plasmin, elastase, thermolysin, or papain activity without inhibition of amidolytic activity. Convenient substrates to be analyzed include  $^{14}\text{C}$ -labeled casein and  $^{125}\text{I}$ -fibrin.

**[0050]** The characteristic of broad protease substrate specificity distinguishes  $\alpha$ 2-MG from inhibitors of protease active sites. Based on this characteristic,  $\alpha$ 2-MG can be detected or the level of  $\alpha$ 2-MG can be determined by assaying for inhibition of the activity of two or more proteases with different active site specificities.  $\alpha$ 2-MG can be detected or the level of  $\alpha$ 2-MG in a sample can be determined, for example, by analyzing the reduction in protease activity of two or more target proteases such as two or more of the following proteases: trypsin, subtilisin, chymotrypsin, plasmin, elastase, thermolysin and papain. Labeled protease substrates such as  $^{14}\text{C}$ -casein or  $^{125}\text{I}$ -fibrin can be useful in such methods (Armstrong et al., *supra*, 1999).

**[0051]**  $\alpha$ 2-MG also can be detected or the level of  $\alpha$ 2-MG determined based on the ability of  $\alpha$ 2-MG to shield a bound protease from an antibody or a high molecular weight inhibitor. A target protease such as trypsin, subtilisin, chymotrypsin, plasmin, elastase, thermolysin, or papain can be added to a plasma sample. Following removal of unbound protease, for example, by immunoprecipitation with anti-protease antibody, the amount of protease bound by  $\alpha$ 2-MG can be determined using a low molecular mass amide or ester substrate. The amount of hydrolyzed low molecular mass substrate is an indicator of the amount of protected,  $\alpha$ 2-MG-bound, protease and, therefore, of the concentration of  $\alpha$ 2-MG. Similarly, a sample can be reacted first with a protease such as trypsin and subsequently with excess protease inhibitor such as soybean trypsin inhibitor before assaying residual trypsin activity with a low molecular mass substrate, such as the amide BApNA ( $\text{N}^\alpha$ -benzoyl-DL-arginine p-nitroanilide (Ganrot, *Clin. Chem. Acta* 14:493-501 (1966); Armstrong et al., *J. Exp. Zool.* 236:1-9 (1985)). Trypsin not sequestered by  $\alpha$ 2-MG is inactivated by the trypsin inhibitor, with only  $\alpha$ 2-MG-protected trypsin remaining capable of substrate hydrolysis. Thus, a positive reaction in a soybean trypsin inhibitor assay detects  $\alpha$ 2-MG and is a quantitative measure of the amount of  $\alpha$ 2-MG (Armstrong et al., *supra*, 1999). One skilled in the art understands that the presence of low molecular mass protease inhibitors capable of inactivating  $\alpha$ 2-MG-bound enzyme can affect the results obtained with such an assay. It is further understood that these and other routine assays for  $\alpha$ 2-MG activity, as well as  $\alpha$ 2-MG RNA or protein levels, can be useful for detecting  $\alpha$ 2-MG or determining a level of  $\alpha$ 2-MG in a method of the invention.

#### Hyaluronic acid

**[0052]** The methods of the invention further rely, in part, on detecting hyaluronic acid or determining a level of hyaluronic acid in a sample. Hyaluronic acid, also known as hyaluronate or hyaluronan, is a high molecular weight polysaccharide with an unbranched backbone made up of alternating glucuronic acid and  $\beta(1,3)$ -N-acetylglucosamine moieties linked by  $\beta$ -1,4 linkages. Hyaluronic acid can have a length of a few to more than 1,000 dimeric units, with each dimeric unit having a molecular weight of about 450 D. Hyaluronic acid, which is produced principally by fibroblasts and other specialized connective tissue cells, plays a structural role in the connective tissue matrix. Furthermore, hyaluronic acid is widely distributed throughout the body and can be found as a free molecule in, for example, plasma, synovial fluid and urine. In plasma, hyaluronic acid has a relatively short half-life.

**[0053]** Serum HA levels can be elevated in liver diseases including cirrhosis (Bramley et al., *J. Hepatol.* 13:8-13 (1991); Ueno et al., *Gastroenterol.* 105:475-481 (1993); Oberti et al., *Gastroenterol.* 113:1609-1616 (1997); and McHutchison et al., *J. Gastroenterol. Hepatol.* 15:945-951 (2000)). Serum HA levels also can be elevated during synovial inflammation and cartilage destruction seen in rheumatoid arthritis; these levels have been found to correlate with disease activity and degree of synovial involvement (Kontinen et al., *Clin. Chimica Acta* 193:39-48 (1990); Poole et al., *Arthritis Rheum.* 37:1030-1038 (1994); Goldberg et al., *Arthritis Rheum.* 34: 799-807 (1991); and Emlen et al., *J. Rheum.* 23:974-978

(1996)). Elevated serum levels of HA also can be present, for example, in patients with osteoarthritis (OA), progressive systemic sclerosis (PSS) and systemic lupus erythematosus (SLE).

**[0054]** As used herein, the term "hyaluronic acid" is synonymous with "HA" and means a polymer of two or more dimeric units of alternating glucuronic acid and  $\beta(1,3)$ -N-acetylglucosamine moieties linked by  $\beta$ -1,4 linkages. As used herein, the phrase "detecting HA" means any quantitative or qualitative assay for determining the presence of HA, and the phrase "determining the level of HA" means any direct or indirect quantitative assay for HA. In view of the above, it is understood that the phrase "detecting HA" encompasses "determining the level of HA."

**[0055]** HA can be detected or a level of HA can be determined using one of a variety of well known assays based on HA-binding proteins or anti-HA antibodies, or by quantitation of purified HA. HA-binding proteins, for example, can be useful in detecting HA; a radiometric assay for HA based on  $^{125}\text{I}$ -labelled HA-binding protein is available from Pharmacia (Guehot et al., Clin. Chem. 42:558-563 (1996)). Other commercial assays based on HA-binding proteins are available, for example, from Corgenix (Westminster, CO; kit 029001). In addition, HA can be detected or a level of HA can be determined using hyaluronectin as described in Maingonnat and Delpech, Ann. Clin. Biochem. 28:305-306 (1991), or using the kit available from Nalgenunc International (Rochester, NY; Delpech and Bertrand, Anal. Biochem. 149:555-565 (1985)). Assays for detecting HA or determining a level of HA include a variety of competitive and non-competitive binding assays, for example, competitive binding assays using  $^{125}\text{I}$ -labeled HA binding protein; competitive binding assays based on alkaline phosphatase labeled-hyaluronectin (HN); and non-competitive binding assays based on peroxidase-labeled proteoglycan or peroxidase-labeled HA-binding protein, among others (Lindquist et al., Clin. Chem. 38: 127-132 (1992)). See, also, Delpech and Bertrand, supra, 1985; Engstrom-Laurent et al., Scand. J. Clin. Invest. 45:497-504 (1985); Brandt et al., Acta Otolaryn. 442 (Suppl.):31-35 (1987); Goldberg, Anal. Biochem. 174:448-458 (1988); Chichibu et al., Clin. Chim. Acta 181:317-324 (1989); Li et al., Conn. Tissue Res. 19:243-254 (1989); Poole et al., Arth. Rheum. 33:790-799 (1990); Poole et al., J. Biol. Chem. 260:6020-6025 (1985); and Laurent and Tengblad, Anal. Biochem. 109:386-394 (1980)). Assays for detecting HA or determining a level of HA in a sample can be performed using a variety of immunoassay formats, including radioimmunoassays and enzyme-linked immunoassays. Anti-HA antiserum useful in immunoassays can be, for example, affinity purified sheep anti-HA antiserum available from Biotrend (Cologne, Germany; #5029-9990).

**[0056]** A level of HA also can be determined by purifying HA from a sample, and quantifying the amount of purified polysaccharide. High performance liquid chromatography can be used alone or in conjunction with mass spectrophotometry. As an example, HPLC can be used to determine HA levels after digestion of samples containing an internal standard with hyaluronidase, separation by a reversed phase octadecylsilyl column and elution with 0.01 M tetrabutylammonium phosphate-acetonitrile (83:17, v/v) at pH 7.35 (Payan et al., J. Chromatogr. 566:9-18 (1991)).

**[0057]** HA levels have been shown to correlate with hyaluronidase levels (Bray et al., Am. Rev. Respir. Dis. 3:284-288 (1991)). Thus, HA can be detected or a level of HA can be determined indirectly by assaying for hyaluronidase activity. Assays for hyaluronidase activity are known in the art, as described in Bray et al., supra, 1991. One skilled in the art understands that these and other routine assays for determining hyaluronidase or HA levels are encompassed by the phrases "detecting HA" and "determining the level of HA" and can be useful in diagnosing the presence or severity of liver fibrosis according to a method of the invention.

#### TIMP-1

**[0058]** The methods of the invention also are based on detecting TIMP-1 in a sample and, in particular embodiments, on determining a level of TIMP-1 in a sample. Tissue inhibitors of metalloproteinases (TIMPs) regulate the activity of the matrix metalloproteinases (MMPs), which are an important group of ECM-degradative enzymes that include gelatinase A (MMP-2) and gelatinase B (MMP-9). In normal liver, matrix components such as collagens, fibronectin, laminin, tenascin, undulin and entactin are constantly remodeled by matrix degrading enzymes to control deposition of extracellular matrix. Elevation of TIMP levels results in inhibition of MMP activity and favors the accumulation of extracellular matrix. The TIMPs, which include TIMP-1, TIMP-2, TIMP-3 and TIMP-4, interact with the matrix metalloproteinases with a 1:1 stoichiometry and inhibit metalloprotease activity through reversible non-covalent binding. TIMP-1, TIMP-2 and TIMP-3 have similar MMP-inhibitory activities, inhibiting the proteolytic activity of collagenase, gelatinase, stromelysin, proteoglycanase and metalloelastases although their localization and regulation differ (Cawston et al., "Protein Inhibitors of Metalloproteinases" in Barrett and Salvesen (Eds), Proteinase Inhibitors Amsterdam Elsevier pages 589-610 (1986)).

**[0059]** Human TIMP-1 is a 184 amino acid sialoglycoprotein with a molecular weight of 28.5 kDa (Murphy et al., Biochem. J. 195:167-170 (1981); Dockerty et al., Nature 318:66-69 (1985); and Bodden et al., J. Biol. Chem. 269: 18943-18952 (1994)). TIMP-1 inhibits all active metalloproteinases, for example, interstitial collagenase MMP-1 as well as stromelysin and gelatinase B (MMP-9). The nucleic acid sequence (SEQ ID NO: 3) and corresponding amino acid sequence (SEQ ID NO: 4) of human TIMP-1 are shown in Figure 2.

**[0060]** As used herein, the term "tissue inhibitor of metalloproteinase-1" is synonymous with "TIMP-1" and means a protein with significant structural homology to human TIMP-1 (SEQ ID NO: 4) that inhibits the proteolytic activity of

metalloproteinases with a specificity similar to human TIMP-1. The presence of human TIMP-1 can be conveniently detected by the presence of epitopes, reactive with a known specific anti-TIMP-1 antibody such as 7-6Cl or 7-23G9.

**[0061]** As used herein, the phrase "detecting TIMP-1" means any quantitative or qualitative assay for determining the presence of TIMP-1, and the phrase "determining the level of TIMP-1" means any direct or indirect quantitative assay for TIMP-1. In view of the above, it is understood that the phrase "detecting TIMP-1" encompasses "determining the level of TIMP-1."

**[0062]** Assays for detecting TIMP-1 and for determining a level of TIMP-1 include well known assays for TIMP-1 RNA, protein and enzymatic activity. Methods of determining TIMP-1 RNA levels by Northern analysis or RT-PCR are well known in the art (Yoshiji et al., *Int. J. Cancer* 69:131-134 (1996); Janowska-Wieczorek et al., *Exp. Hematol.* 28:1274-1285 (2000); and Groft et al., *Br. J. Cancer* 85:55-63 (2001)) as described further below. TIMP-1 protein can be detected or the level of TIMP-1 protein can be conveniently determined, for example, by radioimmunoassay as described in Brophy et al., *Biochem. Biophys. Res. Comm.* 167:898-903 (1990) or by two-antibody sandwich assay as described in Murawaki et al., *Clinica Chimica Acta* 218:47-58 (1993). Plasma concentrations of TIMP-1 protein can be assayed by ELISA with a kit commercially available from Amersham Pharmacia (see, also Example III). Levels of TIMP-1 protein also can be determined by quantifying the amount of purified TIMP-1 protein. Purification of TIMP-1 can be achieved, for example, by HPLC, alone or in combination with mass spectrophotometry, or as described, for example, in Murphy et al., *Biochem. J.* 195:167-170 (1981), or Stricklin and Welgus, *J. Biol. Chem.* 258:12252-12258 (1983). TIMP-1 also can be detected or a level of TIMP-1 determined by assaying for inhibition of the activity of one or more metalloproteases, for example, using reverse gelatin zymography as described in Kossakowska et al., *Amer. J. Pathology* 153:1895-1902 (1998). Assays for TIMP-1 RNA, protein or activity are described further hereinbelow, and one skilled in the art understands that these and other routine assays for detecting TIMP-1 are encompassed by the methods of the invention.

#### Rule-in/Rule-out analysis

**[0063]** As disclosed herein, two sets of cut-off values can be used to increase the accuracy of an assay based on the  $\alpha$ 2-MG/HA/TIMP-1 three-marker panel. As set forth in Example II, a first set of cut-off values for  $\alpha$ 2-MG, HA and TIMP-1 were selected based on optimization for sensitivity in order to first rule out fibrosis, followed by analysis of the "positive" population using a second set of cut-off values optimized for specificity to determine the presence of significant fibrosis. Table 7 shows the results of the dual optimization strategy on the 194 HCV patient study population. The primary cut-offs were set at 2.0 mg/ml, 35 ng/ml and 1000 ng/ml for  $\alpha$ 2-MG, HA and TIMP-1, respectively, to achieve a high sensitivity in the primary analysis. Any samples with all three of  $\alpha$ 2-MG, HA and TIMP-1 levels above the primary cut-off values were tentatively indicated to be positive for F2-F4 fibrosis and were further evaluated using a second set of cut-off values of 2.0 mg/ml, 60 ng/ml and 1575 ng/ml for  $\alpha$ 2-MG, HA and TIMP-1, respectively, which were obtained by optimizing for specificity.

**[0064]** Using the second set of cut-off values optimized for high specificity, 54 of the 122 patients initially designated as positive for F2-F4 fibrosis were confirmed positive, only one of which was a false positive. In sum, of the 194 HCV patients in the study population, 72 were classified as negative (having F0-F1 fibrosis) and 54 were classified as positive (having F2-F4 fibrosis), while 68 samples had indeterminate results and were not classified. When the indeterminate samples were excluded, the  $\alpha$ 2-MG/HA/TIMP-1 panel performed with a positive predictive value of about 98% and a negative predictive value of about 79%. Furthermore, in a more typical patient population having 30% fibrosis prevalence, the same panel performs with positive and negative predictive values of close to 93%. These results indicate that the use of primary and secondary cut-off levels, whereby sensitivity is initially optimized followed by optimization for specificity, can increase the overall accuracy of a three-marker test, resulting in a panel test with about 93% accuracy for non-indeterminate samples, which make up about 70% of the samples tested.

**[0065]** Thus, the present invention provides a method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual by determining an  $\alpha$ 2-MG level in a sample from the individual; determining a HA level in a sample from the individual; determining a TIMP-1 level in a sample from the individual; and diagnosing the individual as having no or mild liver fibrosis when the  $\alpha$ 2-MG level is below an  $\alpha$ 2-MG cut-off value X1, the HA level is below a HA cut-off value Y1 or the TIMP-1 level is below a TIMP-1 cut-off value Z1; diagnosing the individual as having moderate to severe liver fibrosis when the  $\alpha$ 2-MG level is above an  $\alpha$ 2-MG cut-off value X2, the HA level is above a HA cut-off value Y2 and the TIMP-1 level is above a TIMP-1 cut-off value Z2; and diagnosing remaining individuals as having an indeterminate status.

**[0066]** The methods of the invention based on dual cut-off values for the levels of the  $\alpha$ 2-MG, HA and TIMP-1 markers can be useful in differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in a variety of patient populations. Such methods can be useful, for example, in diagnosing an individual having a liver disease such as viral hepatitis, autoimmune liver disease such as autoimmune hepatitis, alcoholic liver disease, fatty liver disease or drug-induced liver disease. In one embodiment, a method of the invention is used to differentiate no or mild liver fibrosis from moderate to severe liver fibrosis in an individual infected with hepatitis C virus. Samples useful in a method of the

invention based on dual cut-off values include, but are not limited to, blood, serum, plasma, urine, saliva and liver tissue. In one embodiment, a method of the invention is practiced by determining the  $\alpha$ 2-MG level, HA level and TIMP-1 level in one or more serum samples.

**[0067]** In a further embodiment, the present invention provides a method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual, where the differentiation is based on an X1 cut-off value between 1.8 and 2.2 mg/ml; a Y1 cut-off value between 31 and 39 ng/ml; a Z1 cut-off value between 900 and 1100 ng/ml; an X2 cut-off value between 1.8 and 2.2 mg/ml; a Y2 cut-off value between 54 and 66 ng/ml; and a Z2 cut-off value between 1415 and 1735 ng/ml. In another embodiment, the differentiation is based on an X1 cut-off value of 2.0 mg/ml; a Y1 cut-off value of 35 ng/ml; a Z1 cut-off value of 1000 ng/ml; an X2 cut-off value of 2.0 mg/ml; a Y2 cut-off value of 60 ng/ml; and a Z2 cut off value of 1575 ng/ml. In yet another embodiment, the differentiation is based on an X1 cut-off value of 2.0 mg/ml; a Y1 cut-off value of 37 ng/ml; a Z1 cut-off value of 1100 ng/ml; an X2 cut-off value of 2.0 mg/ml; a Y2 cut-off value of 60 ng/ml; and a Z2 cut-off value of 1575 ng/ml. In a further embodiment, X1, Y1, Z1, X2, Y2 and Z2 are selected such that, in a population having up to 30% liver fibrosis prevalence, at least 65% of individuals in the population are diagnosed as having no or mild fibrosis or moderate to severe fibrosis with an accuracy of at least 90%. In yet a further embodiment, X1, Y1, Z1, X2, Y2 and Z2 are selected such that, in a population having up to 10% liver fibrosis prevalence, at least 70% of individuals in the population are diagnosed as having no or mild fibrosis or moderate to severe fibrosis with an accuracy of at least 90%.

**[0068]** As set forth above, the methods of the invention are highly accurate for determining the presence or severity of fibrosis in a subgroup of the entire patient population assayed. For example, as shown in Table 7, the methods of the invention perform with more than 93% accuracy in determining the F0-F1 or F2-F4 fibrosis status in about 70% of a patient population having a liver fibrosis prevalence of 30%. The remaining 30% of the patient population are indicated to have an indeterminate status. As used herein, the term "indeterminate status" means that the individual cannot be confidently diagnosed with sufficient predictive value.

**[0069]** As used herein, the term "X1" or "X2" refers to an  $\alpha$ 2-MG cut-off value, against which an experimental  $\alpha$ 2-MG sample level is compared. Similarly, as used herein, the term "Y1" or "Y2" refers to an HA cut-off value, against which an experimental HA level is compared. The term "Z1" or "Z2," as used herein, refers to a TIMP-1 cut-off value against which an experimental TIMP-1 level is compared. X1, Y1 and Z1 cut-offs are combined to determine the presence or severity of fibrosis in a sample. Similarly, X2, Y2 and Z1 cut-off values are combined to determine the presence or severity of fibrosis in a sample. A sample having an  $\alpha$ 2-MG level less than X1, an HA level less than Y1, or a TIMP-1 level less than Z1 is classified as having F0-F1 fibrosis. A sample having an  $\alpha$ 2-MG level above X1, an HA level above Y1, and a TIMP-1 level above Z1 is possibly positive for F2-F4 fibrosis and warrants further analysis. Furthermore, a sample having an  $\alpha$ 2-MG level above X2, an HA level above Y2, and a TIMP-1 level above Z2 is classified as having F2-F4 fibrosis. A sample having an  $\alpha$ 2-MG level above X1, an HA level above Y1, and a TIMP-1 level above Z1 but one or more levels below X2, Y2 or Z2 is classified as having an "indeterminate status." It is understood that X2 generally is equal to or greater than X1; Y2 generally is equal to or greater than Y1; and Z2 generally is equal to or greater than Z1.

**[0070]** One skilled in the art can select  $\alpha$ 2-MG, HA and TIMP-1 cut-offs X1, Y1, Z1, X2, Y2 and Z2 to achieve one or more clinically useful parameters, such as a desired sensitivity or specificity, or a desired negative predictive value, positive predictive value or accuracy for a patient population having a particular fibrosis prevalence. Factorial Design Optimization, also known as Design of Experiments, methodology can be used, for example, to select the appropriate cut-off values. As disclosed herein in Example II, optimization software (DOE Keep It Simple Statistically from Air Academy Associates (Colorado Springs, CO) was used in a central composite design experiment to simultaneously vary the three cut-offs X1, Y1 and Z1, and then to simultaneously vary the three cut-offs X2, Y2 and Z2. In particular, the  $\alpha$ 2-MG cut-off was varied from 2.0 to 5.0 mg/ml; the HA cut-off was varied from 25-75 ng/ml; and the TIMP-1 cut-off was varied from 1000-1700 ng/ml. By comparing the test results determined for the 194 patients in the database (see Table 4) with the assigned X1, Y1 and Z1 cut-offs, each of the 194 samples were determined to be a true positive, true negative, false positive or false negative, and the clinical parameters of sensitivity, specificity, negative predictive value, positive predictive value and accuracy were determined for the study patient population. Although determination of the  $\alpha$ 2-MG, HA and TIMP-1 cut-off values is illustrated herein using the DOE KISS program, one skilled in the art understands that other computer programs for identifying cooperative interactions among multiple variables and for performing simultaneous equation calculations also can be used. For example, ECHIP optimization software, available from ECHIP, Incorporated (Hockessin, DE), or Statgraphics optimization software, available from STSC, Incorporated (Rockville, MD), also can be useful in determining  $\alpha$ 2-MG, HA and TIMP-1 cut-off values useful in the methods of the invention.

**[0071]** The clinical parameters of sensitivity, specificity, negative predictive value, positive predictive value and accuracy are calculated using true positives, false positives, true negatives and false negatives. A "true positive" sample is a sample positive for the indicated stage of fibrosis according to clinical biopsy, which is also diagnosed positive according to a method of the invention. A "false positive" sample is a sample negative for the indicated stage of fibrosis by biopsy, which is diagnosed positive according to a method of the invention. Similarly, a "false negative" is a sample positive for the indicated stage of fibrosis by biopsy, which is diagnosed negative according to a method of the invention. A "true

negative" is a sample negative for the indicated stage of fibrosis by biopsy, and also negative for fibrosis according to a method of the invention. See, for example, Motulsky (Ed.), *Intuitive Biostatistics* New York: Oxford University Press (1995).

**[0072]** As used herein, the term "sensitivity" means the probability that a diagnostic method of the invention gives a positive result when the sample is positive, for example, fibrotic with a Metavir score of F2-F4. Sensitivity is calculated as the number of true positive results divided by the sum of the true positives and false negatives. Sensitivity essentially is a measure of how well a method correctly identifies those with fibrotic disease. In a method of the invention, the X1, Y1, Z1, X2, Y2 and Z2 values can be selected such that the sensitivity of diagnosing an individual is at least about 70%, and can be, for example, at least 75%, 80%, 85%, 90% or 95% in at least 60% of the patient population assayed, or in at least 65%, 70%, 75% or 80% of the patient population assayed.

**[0073]** As used herein, the term "specificity" means the probability that a diagnostic method of the invention gives a negative result when the sample is not positive, for example, not of Metavir fibrosis stage F2-F4. Specificity is calculated as the number of true negative results divided by the sum of the true negatives and false positives. Specificity essentially is a measure of how well a method excludes those who do not have fibrosis. In a method of the invention, the cut-off values X1, Y1, Z1, X2, Y2 and Z2 can be selected such that, when the sensitivity is at least about 70%, the specificity of diagnosing an individual is in the range of 70-100%, for example, at least 75%, 80%, 85%, 90% or 95% in at least 60% of the patient population assayed, or in at least 65%, 70%, 75% or 80% of the patient population assayed. As illustrated in Example II, a specificity of greater than 98% and a sensitivity of about 77% were achieved in the non-indeterminate patient population, which was about 70% of the patient population having a fibrosis prevalence of 30%.

**[0074]** The term "negative predictive value," as used herein, is synonymous with "NPV" and means the probability that an individual diagnosed as not having fibrosis actually does not have the disease. Negative predictive value can be calculated as the number of true negatives divided by the sum of the true negatives and false negatives. Negative predictive value is determined by the characteristics of the diagnostic method as well as the prevalence of fibrosis in the population analyzed. In a method of the invention, the  $\alpha$ 2-MG, HA and TIMP-1 cut-off values can be selected such that the negative predictive value in a population having a liver fibrosis prevalence of up to 10% is in the range of 75-99% and can be, for example, at least 80%, at least 85%, at least 90%, or at least 95%, in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed. The  $\alpha$ 2-MG, HA and TIMP-1 cut-off values also can be selected such that the negative predictive value in a population having a liver fibrosis prevalence of up to 20% is in the range of 75-99% and can be, for example, at least 80%, at least 85%, at least 90%, or at least 95%, in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed. In addition,  $\alpha$ 2-MG, HA and TIMP-1 cut-off values can be selected such that the negative predictive value in a population having a liver fibrosis prevalence of up to 30% is in the range of 75-99% and can be, for example, at least 80%, at least 85%, at least 90%, or at least 95%, in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed.

**[0075]** The term "positive predictive value," as used herein, is synonymous with "PPV" and means the probability that an individual diagnosed as having fibrosis actually has the condition. Positive predictive value can be calculated as the number of true positives divided by the sum of the true positives and false positives. Positive predictive value is determined by the characteristics of the diagnostic method as well as the prevalence of fibrosis in the population analyzed. In a method of the invention, the  $\alpha$ 2-MG, HA and TIMP-1 cut-off values can be selected such that, in a patient population having up to 10% liver fibrosis prevalence, the positive predictive value of the method is at least about 75%, and can be at least 80%, at least 85%, at least 90% or at least 95% in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed. The  $\alpha$ 2-MG, HA and TIMP-1 cut-off values also can be selected such that, in a patient population having up to 20% liver fibrosis prevalence, the positive predictive value of the method is at least about 75%, and can be at least 80%, at least 85%; at least 90% or at least 95% in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed. Similarly, the  $\alpha$ 2-MG, HA and TIMP-1 cut-off values can be selected such that, in a patient population having up to 30% liver fibrosis prevalence, the positive predictive value of the method is at least about 75%, and can be at least 80%, at least 85%, at least 90% or at least 95% in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed.

**[0076]** Predictive values, including negative and positive predictive values, are influenced by the prevalence of the disease in the population analyzed. In the methods of the invention, the cut-off values X1, Y1, Z1, X2, Y2 and Z2 can be selected to produce a desired clinical parameter for a clinical population with a particular liver fibrosis prevalence. For example, cut-off values can be selected for a liver fibrosis prevalence of up to 10%, 12%, 15%, 18%, 20%, 25% or 30% which can be seen, for example, in a hepatologist's office. Cut-off values also can be selected for a liver fibrosis prevalence of up to 1%, 2%, 3%, 4%, 5%, 6%, 7% or 8%, which can be representative of the fibrosis prevalence seen in a general practitioner's office.

**[0077]** As used herein, the term "accuracy" means the overall agreement between the diagnostic method and the disease state. Accuracy is calculated as the sum of the true positives and true negatives divided by the total number of

sample results and is affected by the prevalence of fibrosis in the population analyzed. The  $\alpha$ 2-MG, HA and TIMP-1 cut-off values can be selected such that the accuracy of a method of the invention in a patient population having a liver fibrosis prevalence of up to 10% is at least about 80% and can be, for example, at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed. The  $\alpha$ 2-MG, HA and TIMP-1 cut-off values also can be selected such that the accuracy of a method of the invention in a patient population having a liver fibrosis prevalence of up to 20% is at least about 80% and can be, for example, at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed. Similarly, the  $\alpha$ 2-MG, HA and TIMP-1 cut-off values can be selected such that the accuracy of a method of the invention in a patient population having a liver fibrosis prevalence of up to 30% is at least about 80% and can be, for example, at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% in at least 60% of the patient population assayed, for example, in at least 65%, 70%, 75% or 80% of the patient population assayed.

#### Methods not limited to specific markers

**[0078]** Also disclosed is a method of diagnosing the presence or severity of liver fibrosis in an individual by comparing a level of a first fibrotic marker X in the individual to a cut-off value X1 to determine whether the individual is positive for the first fibrotic marker X; comparing a level of a second fibrotic marker Y in the individual to a cut-off value Y1 to determine whether the individual is positive for the second fibrotic marker Y; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X and Y, where, in a population with up to 40% fibrosis prevalence, at least 65% of individuals in the population are diagnosed with an accuracy of at least 90%.

**[0079]** The method can include, if desired, comparing a level of a third fibrotic marker Z in the individual to a cut-off value Z1 to determine whether the individual is positive for the third fibrotic marker Z and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X, Y and Z. In the method of the invention, the first fibrotic marker is  $\alpha$ 2-MG, the second fibrotic marker is HA, and the third fibrotic marker is TIMP-1.

**[0080]** In another embodiment, the levels of the at least three fibrotic markers are compared, and, in a further embodiment, the levels of exactly three fibrotic markers are compared to their respective cut-off values. In additional embodiments, the levels of at least four or at least five fibrotic markers are compared. A method of the invention can be useful, for example, to differentiate no or mild liver fibrosis from moderate to severe liver fibrosis.

**[0081]** In a specific embodiment, a method of the invention serves to diagnose at least 65% of individuals in a population with up to 30% fibrosis prevalence with an accuracy of at least 93%. In a further embodiment, a method of the invention serves to diagnose at least 70% of individuals in a population with up to 20% fibrosis prevalence with an accuracy of at least 94%. In yet a further embodiment, a method of the invention serves to diagnose at least 70% of individuals in a population with up to 10% fibrosis prevalence with an accuracy of at least 96%.

**[0082]** The methods of the invention provide unparalleled performance in diagnosing the presence or severity of liver fibrosis. While not all patients are provided with a diagnosis, the majority are diagnosed with extremely good accuracy. As an example, in a patient population with about 40% fibrosis prevalence, almost 70% of the population are diagnosed with more than 91% accuracy and with a positive predictive value of more than 96% and a negative predictive value of more than 89%. This excellent performance contrasts with alternative methods such as the method of Poynard et al., Lancet 357:1069 (2001). Using the method of Poynard et al. based on analysis of the six markers  $\alpha$ 2-MG,  $\alpha$ 2-globulin, total bilirubin,  $\gamma$ -globulin, apoA1 and GGT, only about 50% of a population having about 40% fibrosis prevalence are diagnosed, and only with an accuracy of about 89% (see Table 8). Thus, the methods of the invention provide an improvement, in that a significantly greater percentage of a patient population (about 70% as compared to about 50%) are diagnosed, and with an accuracy of more than 91% as compared to an accuracy of around 89% (see Table 8). Due to the novel performance characteristics of a method of the invention, biopsy is typically unnecessary in at least 65% of a patient population, and the patients diagnosed can have confidence in a diagnosis that is more than 90% accurate.

**[0083]** Like other methods of the invention, a method of the invention based on comparison of the at least three fibrotic markers can be used to diagnose the presence or severity of liver fibrosis in an individual having or suspected of having any liver disorder, including viral hepatitis, autoimmune liver disease such as autoimmune hepatitis, alcoholic liver disease, fatty liver disease or drug-induced liver disease, or any of the other liver diseases described herein above. Similarly a method of the invention based on comparison of the at least three fibrotic markers can be used to diagnose the presence or severity of fibrotic disorders including pulmonary fibrosis, kidney fibrosis, prostate fibrosis, breast fibrosis or a rheumatoid disease, or another fibrotic disorder described herein or known in the art.

**[0084]** A method of the invention relies on comparison of the level of a fibrotic marker to a predetermined cut-off value. For markers that positively correlate with fibrosis, positivity is indicated by a level that is greater than the predetermined cut-off value. For markers that negatively correlate with fibrosis, positivity is indicated by a level that is less than the predetermined cut-off value. Cut-off values useful in the methods of the invention can be determined as described herein, for example, using design of experiments (DOE) analysis.

**[0085]** As for the other diagnostic methods of the invention, these methods can be practiced using a variety of fibrotic markers known in the art or described herein. Such fibrotic markers include, without limitation,  $\alpha$ 2-MG, HA, TIMP-1, PIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 or ApoB. Additional serological, biochemical, clinical and echographic fibrotic markers are described herein above or are known in the art and can be included in any combination in a method of the invention. Furthermore, it is understood that comparison of the first and second fibrotic markers and any additional fibrotic markers can be performed simultaneously or in any order and using any combination of assay formats.

**[0086]** As described above, the "level" of a fibrotic marker can be a relative or absolute amount of, for example, RNA, protein or activity and can be a direct or indirect measurement of the fibrotic marker. In addition, the value of the level can be obtained from a secondary source, such as a physician or diagnostic laboratory or can be determined using any convenient sample and assay, including but not limited to those described herein above. Methods useful in determining the level of a fibrotic marker in order to perform the comparisons included in the methods of the invention encompass, for example, hybridization methods such as RT-PCR and RNA blot analysis, immunoassays including enzyme-linked immunosorbent assays (ELISAs) and radioimmunoassays (RIAs), sandwich immunoassays, quantitative western blotting and other standard assays for determining protein levels, and, where applicable, assays for the activity of the fibrotic marker. Such assays are routine in the art and described herein above.

**[0087]** The present invention further provides a method of diagnosing the presence or severity of liver fibrosis in an individual by comparing a level of a first fibrotic marker X in the individual to a cut-off value X1 to determine whether the individual is positive for the first fibrotic marker X; comparing a level of a second fibrotic marker Y in the individual to a cut-off value Y1 to determine whether the individual is positive for the second fibrotic marker Y; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X and Y, where the cut-off values X1 and Y1 are optimized individually to give a desired performance characteristic, wherein

**[0088]** The method of the invention includes the steps of comparing a level of a third fibrotic marker Z in the individual to a cut-off value Z1 to determine whether the individual is positive for the third fibrotic marker Z and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X, Y and Z, where the cut-off values X1, Y1 and Z1 are optimized individually to give a desired performance characteristic, wherein , levels of  $\alpha$ 2-MG, HA and TIMP-1 are compared. In other embodiments, the levels of exactly three, at least three, at least four, or at least five fibrotic markers are compared. A method of the invention can be useful, for example, in differentiating no or mild liver fibrosis from moderate to severe liver fibrosis. Cut-off values can be optimized as described herein, for example, using DOE analysis.

**[0089]** Further provided by the invention is a method of diagnosing the presence or severity of liver fibrosis in an individual by comparing a level of the first fibrotic marker X in the individual to two cut-off values X1 and X2 to determine whether the individual is positive for the first fibrotic marker X; comparing a level of the second fibrotic marker Y in the individual to two cut-off values Y1 and Y2 to determine whether the individual is positive for the second fibrotic marker Y; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X and Y, where the cut-off values X1, Y1, X2 and Y2 are optimized individually to give a desired performance characteristic. Such performance characteristics include particular sensitivities, specificities, PPVs, NPVs and accuracies, as described herein above.

**[0090]** The method of the invention further includes the steps of comparing a level of a third fibrotic marker Z in the individual to two cut-off values Z1 and Z1 to determine whether the individual is positive for the third fibrotic marker Z; and diagnosing the presence or severity of liver fibrosis in the individual based on positivity or negativity for X, Y and Z, where the cut-off values X1, Y1, Z1, X2, Y2 and Z2 are optimized individually to give a desired performance characteristic. In a method of the invention, cut-off values can be conveniently optimized, for example, using DOE analysis.

#### Methodology

**[0091]** A variety of means can be useful for detecting  $\alpha$ 2-MG, HA and TIMP-1 and for determining a level of  $\alpha$ 2-MG, HA and TIMP in a sample. In one embodiment, the invention is practiced by determining the level of  $\alpha$ 2-MG protein in a sample from the individual to be diagnosed using, for example, one or more  $\alpha$ 2-MG-specific binding agents such as anti- $\alpha$ 2-MG antibodies. In another embodiment, a method of the invention is practiced by assaying for  $\alpha$ 2-MG activity in a sample from the individual.

**[0092]** A variety of means also can be used in a method of the invention to detect HA or determine a level of HA in a sample. In one embodiment, the invention is practiced by determining the level of HA in a sample using one or more HA-specific binding agents such as HA-binding proteins or anti-HA antibodies.

**[0093]** Similarly, a variety of means can be used in a method of the invention to detect TIMP-1 or determine a level of TIMP-1 in a sample. In one embodiment, the invention is practiced by determining the level of TIMP-1 protein in a sample from the individual to be diagnosed. The level of TIMP-1 protein can be determined, for example, using one or more TIMP-1-specific binding agents such as anti-TIMP-1 antibodies. In another embodiment, the invention is practiced

by assaying for TIMP-1 activity in a sample from the individual to be diagnosed.

**[0094]** In a particular embodiment, the invention provides a method of diagnosing the presence or severity of liver fibrosis in an individual by determining the level of  $\alpha$ 2-MG protein in a sample from the individual; determining the level of HA in a sample from the individual; and determining the level of TIMP-1 protein in a sample from the individual; and diagnosing the presence or severity of liver fibrosis in the individual based on the levels of  $\alpha$ 2-MG protein, HA and TIMP-1 protein. If desired, the level of  $\alpha$ 2-MG protein, HA and TIMP-1 protein each can be determined using an enzyme-linked assay.

**[0095]** In a further embodiment, the present invention provides a method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual by contacting an appropriate dilution of a sample from the individual with anti- $\alpha$ 2-MG antibody under conditions suitable to form a first complex of  $\alpha$ 2-MG and anti- $\alpha$ 2-MG antibody; washing the first complex to remove unbound molecules; determining the amount of  $\alpha$ 2-MG-containing first complex; contacting an appropriate dilution of a sample from the individual with a HA-binding protein under conditions suitable to form a second complex of HA and HA-binding protein; washing the second complex to remove unbound molecules; determining the amount of HA-containing second complex; contacting an appropriate dilution of a sample from the individual with anti-TIMP-1 antibody under conditions suitable to form a third complex of TIMP-1 and anti-TIMP-1 antibody; washing the third complex to remove unbound molecules; determining the amount of TIMP-1-containing third complex; and differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in the individual based on the amounts of  $\alpha$ 2-MG, HA and TIMP-1-containing complexes.

**[0096]** It is understood that detecting  $\alpha$ 2-MG, HA and TIMP-1, or detecting  $\alpha$ 2-MG, HA and YKL-40, as discussed further below, can be accomplished by assaying for the amount of protein or polysaccharide directly, or, in the case of  $\alpha$ 2-MG and TIMP-1, can be determined by assaying for RNA levels or enzyme activity of a protease regulated by  $\alpha$ 2-MG or TIMP-1. Similarly, where one or more additional fibrotic markers is detected in a method of the invention, the marker can be assayed directly, or a precursor such as RNA, or a breakdown or proteolytic product, or an activity correlated with levels of the marker can be assayed. It is understood that determining a level of  $\alpha$ 2-MG, HA, TIMP-1 and YKL-40, or a level of any additional marker of fibrosis, can be performed using absolute values, for example, for RNA or protein levels or enzyme activity, or can be determined as relative values in comparison to one or more reference values.

**[0097]** It further is understood that each of the three fibrotic marker assays ( $\alpha$ 2-MG/HA/TIMP-1 or  $\alpha$ 2-MG/HA/YKL-40), as well as any additional assays, is performed independently of the others, in any order, and that any combination of assay formats is encompassed by the invention. As an example, a level of  $\alpha$ 2-MG and HA can be determined by assaying for the concentration of  $\alpha$ 2-MG and HA while a level of TIMP-1 is determined by assaying for TIMP-1 enzyme activity. As another example, a level of  $\alpha$ 2-MG can be determined using a radioimmunoassay, while levels of HA and TIMP-1 are determined using enzyme-linked assays. One skilled in the art understands that detection of the three fibrotic markers ( $\alpha$ 2-MG/HA/TIMP-1 or  $\alpha$ 2-MG/HA/YKL-40) and detection of any additional markers can be performed simultaneously or in any order. Furthermore, a single sample such as a serum sample can be obtained from an individual and subdivided into three portions for detecting  $\alpha$ 2-MG, HA and TIMP-1 or  $\alpha$ 2-MG, HA and TIMP-1, or the markers can be detected using different samples, which can be of the same or a different type and can be undiluted or diluted to the same or different extents. Where two or more samples are used, the samples are usually obtained from the individual within a relatively short time frame, for example, several days to several weeks.

#### RNA methods

**[0098]** Hybridization methods can be used to detect  $\alpha$ 2-MG or TIMP-1 mRNA or determine the level of  $\alpha$ 2-MG or TIMP-1 mRNA or the mRNA of another fibrotic marker useful in the invention such as YKL-40. Numerous methods are well known in the art for determining mRNA levels by specific or selective hybridization with a complementary nucleic acid probe. Such methods include solution hybridization procedures as well as solid-phase hybridization procedures in which the probe or sample is immobilized on a solid support. Specific examples of useful methods include amplification methods such as target and signal amplification methods and include PCR (polymerase chain reaction) and reverse-transcriptase-PCR (RT-PCR); transcription mediated amplification (Gen-Probe Incorporated; San Diego, CA); branched chain DNA (bDNA) amplification (Bayer Diagnostics; Emeryville, CA); strand displacement amplification (SDA; Becton Dickinson; Franklin Lakes, NJ); and ligase chain reaction (LCR) amplification (Abbott Laboratories; Abbott Park, IL). Additional methods useful in the invention include RNase protection; Northern analysis or other RNA blot, dot blot or membrane-based technology; dip stick; pin; and two-dimensional array immobilized onto a chip. Conditions are well known in the art for quantitative determination of mRNA levels using both solution and solid phase hybridization procedures as described, for example, in Ausubel et al., Current Protocols in Molecular Biology (Supplement 47), John Wiley & Sons, New York (1999).

**[0099]** The polymerase chain reaction (PCR) RT-PCR can be useful in the methods of the invention. PCR or RT-PCR can be performed with isolated RNA or crude or partially fractionated samples, for example, cells pelleted from a whole



blood sample. PCR methods are well known in the art as described, for example, in Dieffenbach and Dveksler, PCR Primer: A Laboratory Manual, Cold Spring Harbor Press, Plainview, New York (1995). Multisample formats such as two-dimensional arrays offer the advantage of analyzing numerous different samples in a single assay. Solid-phase dip stick-based methods also can be useful in the invention and offer the advantage of being able to rapidly analyze a fluid sample and obtain an immediate result.

**[0100]** Probes for detecting  $\alpha$ 2-MG and TIMP-1 mRNA or for determining  $\alpha$ 2-MG and TIMP-1 mRNA levels are well known in the art. One skilled in the art can use, for example, a probe corresponding to some or all of the human  $\alpha$ 2-MG nucleic acid sequence shown in Figure 1 (SEQ ID NO: 1) or some or all of the human TIMP-1 nucleic acid sequence shown in Figure 3, respectively. Appropriate conditions for various assay formats for detecting  $\alpha$ 2-MG and TIMP-1 mRNA or for determining  $\alpha$ 2-MG and TIMP-1 mRNA levels are well known in the art or can be established using routine methods. As an example, conditions and probes for Northern analysis of  $\alpha$ 2-MG RNA in human samples are described, for example, in Ortego et al., *supra*, 1997. As another example, conditions and probes for RNA slot blot hybridization to determine  $\alpha$ 2-MG RNA expression in human samples are described in Simon et al., *supra*, 1996. Similarly, Northern analysis of TIMP-1 RNA in human samples can be performed as described, for example, in Yoshiji et al., *supra*, 1996; RT-PCR assays for TIMP-1 in human samples also are well known in the art as described, for example, in Janowska-Wieczorek et al., *supra*, 2000, and Groft et al., *supra*, 2001. The skilled person understands that these and other assays can be useful for detecting  $\alpha$ 2-MG, TIMP-1 or YKL-40 RNA or for determining  $\alpha$ 2-MG, TIMP-1 or YKL-40 RNA levels or the levels of other fibrotic markers useful in the methods of the invention.

#### Immunoassays

**[0101]** A variety of immunoassay formats, including competitive and non-competitive immunoassay formats, antigen capture assays and two-antibody sandwich assays also are useful the methods of the invention (Self and Cook, Curr. Opin. Biotechnol. 7:60-65 (1996)). In one embodiment, a method of the invention relies on one or more antigen capture assays. In an antigen capture assay, antibody is bound to a solid phase, and sample is added such that  $\alpha$ 2-MG, HA, TIMP-1, YKL-40 or another fibrotic marker antigen is bound by the antibody. After unbound proteins are removed by washing, the amount of bound antigen can be quantitated, if desired, using, for example, a radioassay (Harlow and Lane, Antibodies A Laboratory Manual Cold Spring Harbor Laboratory: New York, 1988)). One skilled in the art understands that immunoassays useful in the invention are performed under conditions of antibody excess, or as antigen competitions, to quantitate the amount of antigen and, thus, determine a level of  $\alpha$ 2-MG, HA, TIMP-1 or YKL-40.

**[0102]** Enzyme-linked immunosorbent assays (ELISAs) can be useful in the methods of the invention. An enzyme such as horseradish peroxidase (HRP), alkaline phosphatase (AP),  $\beta$ -galactosidase or urease can be linked, for example, to an anti- $\alpha$ 2-MG, anti-HA, anti-TIMP-1 or anti-YKL-40 antibody or to a secondary antibody for use in a method of the invention. A horseradish-peroxidase detection system can be used, for example, with the chromogenic substrate tetramethylbenzidine (TMB), which yields a soluble product in the presence of hydrogen peroxide that is detectable at 450 nm. Other convenient enzyme-linked systems include, for example, the alkaline phosphatase detection system, which can be used with the chromogenic substrate *p*-nitrophenyl phosphate to yield a soluble product readily detectable at 405 nm. Similarly, a  $\beta$ -galactosidase detection system can be used with the chromogenic substrate *o*-nitrophenyl- $\beta$ -D-galactopyranoside (ONPG) to yield a soluble product detectable at 410 nm, or a urease detection system can be used with a substrate such as urea-bromocresol purple (Sigma Immunochemicals, St. Louis, MO). Useful enzyme-linked primary and secondary antibodies can be obtained from a number of commercial sources such as Jackson Immuno-Research (West Grove, PA) as described further below.

**[0103]** Chemiluminescent detection also can be useful for detecting  $\alpha$ 2-MG, HA, TIMP-1 or YKL-40 or for determining a level of  $\alpha$ 2-MG, HA, TIMP-1 or YKL-40 or another fibrotic marker according to a method of the invention. Chemiluminescent secondary antibodies can be obtained commercially from various sources such as Amersham.

**[0104]** Fluorescent detection also can be useful for detecting  $\alpha$ 2-MG, HA, TIMP-1 or YKL-40 or for determining a level of  $\alpha$ 2-MG, HA, TIMP-1 or YKL-40 or another fibrotic marker in a method of the invention. Useful fluorochromes include, without limitation, DAPI, fluorescein, Hoechst 33258, R-phycoerythrin, B-phycoerythrin, R-phycoerythrin, rhodamine, Texas red and lissamine. Fluorescein or rhodamine labeled  $\alpha$ 2-MG-, HA-, TIMP-1- or YKL-40-specific binding agents such as anti- $\alpha$ 2-MG, anti-HA, anti-TIMP-1, or anti-YKL-40 antibodies, or fluorescein- or rhodamine-labeled secondary antibodies can be useful in the invention. Useful fluorescent antibodies can be obtained commercially, for example, from Tago Immunologicals (Burlingame, CA) as described further below.

**[0105]** Radioimmunoassays (RIAs) also can be useful in the methods of the invention. Such assays are well known in the art. For example, Brophy et al., Biochem. Biophys. Res. Comm. 167:898-903 (1990)), describes a radioimmunoassay for detection of TIMP-1, and Pharmacia makes a radiometric assay for quantitation of HA using an  $^{125}$ I-labelled HA-binding protein (Guechot et al., Clin. Chem. 42:558-563 (1996). Radioimmunoassays can be performed, for example, with  $^{125}$ I-labeled primary or secondary antibody (Harlow and Lance, *supra*, 1988).

**[0106]** A signal from a detectable reagent can be analyzed, for example, using a spectrophotometer to detect color

from a chromogenic substrate; a radiation counter to detect radiation, such as a gamma counter for detection of  $^{125}\text{I}$ ; or a fluorometer to detect fluorescence in the presence of light of a certain wavelength. Where an enzyme-linked assay is used, quantitative analysis of the amount of  $\alpha 2$ -MG, HA, TIMP-1 or YKL-40 or another fibrotic marker can be performed using a spectrophotometer such as an EMAX Microplate Reader (Molecular Devices; Menlo Park, CA) in accordance with the manufacturer's instructions. It is understood that the assays of the invention can be automated or performed robotically, if desired, and that the signal from multiple samples can be detected simultaneously.

**[0107]** The methods of the invention also encompass the use of capillary electrophoresis based immunoassays (CEIA), which can be automated, if desired. Immunoassays also can be used in conjunction with laser-induced fluorescence as described, for example, in Schmalzing and Nashabeh, *Electrophoresis* 18:2184-93 (1997), and Bao, J. *Chromatogr. B. Biomed. Sci.* 699:963-80 (1997). Liposome immunoassays, such as flow-injection liposome immunoassays and liposome immunosensors, also can be used to detect  $\alpha 2$ -MG, HA, TIMP-1 or YKL-40 or to determine a level of  $\alpha 2$ -MG, HA, TIMP-1 or YKL-40 or another fibrotic marker according to a method of the invention (Rongen et al., *J. Immunol. Methods* 204: 105-133 (1997)).

**[0108]** Sandwich enzyme immunoassays also can be useful in the methods of the invention. In a two-antibody sandwich assay, a first antibody is bound to a solid support, and the antigen is allowed to bind to the first antibody. The amount of  $\alpha 2$ -MG, HA, TIMP-1, YKL-40 or another fibrotic marker antigen is quantitated by measuring the amount of a second antibody that binds the fibrotic marker.

**[0109]** As an example, a two-antibody sandwich immunoassay can be useful to determine a level of TIMP-1 as described in Murawaki et al., *supra*, 1993. Briefly, serum (25  $\mu\text{l}$ ) is diluted 41-fold with 10 mM sodium phosphate buffer, pH 7.0 (1.0 ml). The diluted sample (20  $\mu\text{l}$ ) is mixed with 0.3 ml of 10 mM sodium phosphate buffer, pH 7.0, containing 50 ng/ml monoclonal antibody (Fab of clone 7-6C1) labeled with horseradish peroxidase, 1% bovine serum albumin, 0.1% Tween 20, 0.1 M NaCl and 0.005% thimerosal. A 0.1 ml aliquot of the mixed solution is transferred to each microplate well previously coated with a second monoclonal antibody (clone 7-23G9) having a different epitope specificity, and the plate incubated for 30 minutes at room temperature without shaking. The plate is washed three times with 0.3 ml 10 mM sodium phosphate buffer, pH 7.0, containing 0.1% Tween 20 and 0.1 M NaCl. Peroxidase activity bound to the plate is assayed by a 15 minute incubation at room temperature with 0.1 ml 0.15 M citric acid sodium phosphate buffer, pH 4.9, containing 0.5 mg/ml *o*-phenylenediamine and 0.02%  $\text{H}_2\text{O}_2$ . After stopping the reaction by addition of 0.1 ml 2 N  $\text{H}_2\text{SO}_4$ , the absorbance at 492 nm is measured in a microplate reader using a standard of human serum TIMP-1. Linearity between the amount of TIMP-1 and absorbance at 492 nm is demonstrated by graphing with logarithmic scales and yields an assay range of about 1.5 to 300  $\mu\text{g}/\text{well}$ .

**[0110]** Quantitative western blotting also can be used to detect  $\alpha 2$ -MG, HA, TIMP-1 or YKL-40 or to determine a level of  $\alpha 2$ -MG, HA, TIMP-1 or YKL-40 or a level of another fibrotic marker antigen in a method of the invention. Western blots can be quantitated by well known methods such as scanning densitometry. As an example, protein samples are electrophoresed on 10% SDS-PAGE Laemmli gels. Primary murine monoclonal antibodies, for example, against human  $\alpha 2$ -MG, HA, TIMP-1 or YKL-40 are reacted with the blot, and antibody binding confirmed to be linear using a preliminary slot blot experiment. Goat anti-mouse horseradish peroxidase-coupled antibodies (BioRad) are used as the secondary antibody, and signal detection performed using chemiluminescence, for example, with the Renaissance chemiluminescence kit (New England Nuclear; Boston, MA) according to the manufacturer's instructions. Autoradiographs of the blots are analyzed using a scanning densitometer (Molecular Dynamics; Sunnyvale, CA) and normalized to a positive control. Values are reported, for example, as a ratio between the actual value to the positive control (densitometric index). Such methods are well known in the art as described, for example, in Parra et al., *J. Vasc. Surg.* 28:669-675 (1998).

#### Sources for antibodies

**[0111]** As described herein above, immunoassays including but not limited to enzyme-linked immunosorbent assays, radioimmunoassays and quantitative western analysis, can be useful in the diagnostic methods of the invention. Such assays rely on one or more antibodies, for example, anti- $\alpha 2$ -MG, anti-HA, anti-TIMP-1 or anti-YKL-40 antibodies. As used herein, the term "antibody" is used in its broadest sense to include polyclonal and monoclonal antibodies, as well as polypeptide fragments of antibodies that retain binding activity for  $\alpha 2$ -MG, HA, TIMP-1, YKL-40 or the relevant fibrotic marker antigen of at least about  $1 \times 10^5 \text{ M}^{-1}$ . One skilled in the art understands that antibody fragments such as anti- $\alpha 2$ -MG, anti-HA, anti-TIMP-1 and anti-YKL-40 antibody fragments and including Fab,  $\text{F(ab')}_2$  and Fv fragments can retain binding activity for the relevant fibrotic marker antigen and, thus, are included within the definition of the term antibody as used herein. Methods of preparing monoclonal and polyclonal antibodies are routine in the art, as described, for example, in Harlow and Lane, *supra*, 1988.

**[0112]** The term antibody, as used herein, also encompasses non-naturally occurring antibodies and fragments containing, at a minimum, one  $\text{V}_\text{H}$  and one  $\text{V}_\text{L}$  domain, such as chimeric antibodies, humanized antibodies and single chain  $\text{F}_\text{V}$  fragments (scFv) that specifically bind  $\alpha 2$ -MG, HA, TIMP-1, YKL-40 or the relevant fibrotic marker antigen. Such non-naturally occurring antibodies can be constructed using solid phase peptide synthesis, produced recombinantly or ob-

tained, for example, by screening combinatorial libraries consisting of variable heavy chains and variable light chains as described by Borrebaeck (Ed.), *Antibody Engineering* (Second edition) New York: Oxford University Press (1995).

[0113] A variety of useful anti- $\alpha$ 2-MG, anti-HA, anti-TIMP-1 and anti-YKL-40 monoclonal and polyclonal antibodies are well known in the art and, in many cases, are commercially available. For example, a nephelometry assay for  $\alpha$ 2-macroglobulin is available from Beckman Coulter (kit #449430), and affinity purified goat anti-human  $\alpha$ 2-MG and peroxidase-labeled goat anti-human  $\alpha$ 2-MG antibodies suitable for ELISA and western blotting are available, for example, from Cedarlane Laboratories Limited (CL20010AP and CL20010APHP) and Affinity Biologicals Incorporated (GAA2M-AP and GAA2M-APHRP). Similarly, affinity purified sheep anti-HA antiserum can be obtained from Biotrend (#5029-9990).

[0114] Anti-human TIMP-1 antibodies also are readily available from a variety of commercial sources. For example, the anti-human TIMP-1 monoclonal antibody 147-6D11 is suitable for ELISA or western blotting analysis and can be obtained from Medicorp, Inc. (Montreal, Canada), and the anti-human TIMP-1 monoclonal antibody MAB970 is available from R&D Systems, Inc., for use, for example, in western blotting or sandwich ELISA assays. MAB970 can be combined, for example, with biotinylated anti-human TIMP-1 antibody (BAF970) from R&D Systems, Inc., for detection of TIMP-1 by sandwich ELISA. In addition, rabbit anti-human TIMP-1 polyclonal antiserum and mouse anti-human monoclonal antibodies suitable, for example, for western blotting with enhanced chemiluminescence detection can be obtained from Research Diagnostics Inc. (RDI-TIMP1abr and RDI-TIMP1-C1).

#### Assays for activity

[0115] As discussed above, assays based on the activity of a fibrotic marker also can be useful for detecting  $\alpha$ 2-MG, HA or TIMP-1 or for determining a level of  $\alpha$ 2-MG, HA or TIMP-1 or another fibrotic marker and, therefore, are useful in the methods of the invention. As an example, a variety of assays for  $\alpha$ 2-MG activity can be useful for detecting  $\alpha$ 2-MG or determining a level of  $\alpha$ 2-MG in a sample in a method of the invention. Because  $\alpha$ 2-MG-bound proteases display inhibited proteolytic activity but retain the ability to hydrolyze amide and ester bonds of small substrates,  $\alpha$ 2-MG can be detected, or a level determined, by assaying for inhibition of trypsin, subtilisin, chymotrypsin, plasmin, elastase, thermolysin, or papain activity or the activity of another target protease without inhibition of amidolytic activity. Substrates such as labeled casein or labeled fibrin can be useful for assaying for inhibition of target protease activity. Furthermore, based on its broad protease substrate specificity, a level of  $\alpha$ 2-MG can be determined by assaying for inhibition of the activity of two or more target proteases using, for example,  $^{14}$ C-casein and  $^{125}$ I-fibrin (Armstrong et al., *supra*, 1999).  $\alpha$ 2-MG also can be detected or a level of  $\alpha$ 2-MG determined based on the ability of  $\alpha$ 2-MG to shield a bound protease from an antibody or a high molecular weight inhibitor. Following reaction of a sample with, for example, trypsin and then trypsin inhibitor, residual trypsin activity is assayed with a low molecular mass substrate such as the amide BApNA (Ganrot, *supra*, 1966; Armstrong et al., *supra*, 1985). Trypsin activity following treatment with trypsin inhibitor is indicative of  $\alpha$ 2-MG. These and other well known assays for  $\alpha$ 2-MG activity can be useful in the methods of the invention.

[0116] Similarly, assays for TIMP-1 activity are well known in the art. In particular, one assays for the ability to inhibit protease activity of a matrix metalloproteinase, for example, using reverse gelatin zymography. Reverse gelatin zymography is performed by including a gelatinase such as gelatinase A in a gel mix with the gelatin substrate. Conditioned media, such as conditioned media from baby hamster kidney cells can be used as a convenient source of gelatinase. Plasma samples are electrophoresed, and the resulting pattern analyzed, for example, with scanning digitization using a Hewlett Packard scanner. TIMP-1 activity is observed as a reduction of gelatin degradation. See, for example, Kosakowska et al., *supra*, 1998. The skilled person recognizes that these and other routine assays for TIMP-1 activity can be useful in the methods of the invention.

#### Additional markers

[0117] It is clear that the methods of the invention can be practiced, if desired, by detecting the three markers  $\alpha$ 2-MG, HA and TIMP-1 without assaying for any additional markers or evaluating any other clinical or echographic characteristics. In addition, these three assays can be used as a panel in combination with one or more additional fibrotic marker assays or evaluation of one or more clinical or echographic variables. In specific embodiments, the invention provides a method of diagnosing the presence or severity of liver fibrosis in an individual by detecting  $\alpha$ 2-MG, HA and TIMP-1 in a sample and also detecting at least one of the following markers: PIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 or ApoB. In one embodiment, a method of the invention for diagnosing the presence or severity of liver fibrosis includes the steps of detecting  $\alpha$ 2-MG, HA, TIMP-1 and YKL-40 in a sample. In a further embodiment, a method of the invention is limited to detecting  $\alpha$ 2-MG, HA, TIMP-1 and YKL-40, and no additional fibrotic markers are detected.

[0118] In view of the above, it is clear that assays for one or more additional biochemical or serological markers of fibrosis or evaluation of one or more clinical or echographic variables associated with fibrosis can be combined with

detection of  $\alpha$ 2-MG, HA, and TIMP-1 to diagnose the presence or severity of liver fibrosis. Examples of additional biochemical and serological markers include, yet are not limited to, PIIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 and apoB. Additional biochemical and serological markers useful in the invention include, without limitation, collagens such as collagen type I; fibronectin; vitronectin; endothelin; undulin; adhesion molecules such as selectins, vascular cell adhesion molecules (VCAMs) and intercellular adhesion molecules (ICAMs); pro-inflammatory cytokines such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ); pseudocholinesterase; manganese superoxide dismutase; N-acetyl- $\beta$ -glucosaminidase ( $\beta$ -NAG); glutathione peroxidase; connective tissue growth factor (CTGF); platelet derived growth factor (PDGF); PDGF receptor; monocyte chemotactic protein-1 (MCP-1); inducible nitric oxide synthetase; nitrotyrosine; bilirubin; ferritin and  $\alpha$ -feto-protein;  $\gamma$ -glutamyl transpeptidase (GGT); aspartate aminotransferase (AST); alanine aminotransferase (ALT); AST/ALT ratio; albumin;  $\gamma$ -globulins;  $\beta$  $\gamma$ -block; prothrombin index; Child-Pugh score; PGA index (prothrombin time, GGT concentration and apoA1 concentration); PGAA index (PGA score with  $\alpha$ 2-macroglobulin level); hemoglobin; mean corpuscular volume; lymphocyte count; cholesterol; urea; creatinine; sodium and platelet count.

**[0119]** A clinical or echographic variable also can be a fibrotic "marker" useful in the methods of the invention. Thus, analysis of one or more clinical or echographic variables can be combined with detection of  $\alpha$ 2-MG, HA and TIMP-1 to diagnose the presence or severity of liver fibrosis, or another fibrotic disorder as described hereinabove. As examples, such a clinical variable can be patient age or gender or the presence of palmar erythema, Dupuytren's contracture, finger clubbing, spider nevi, firm liver, splenomegaly or collateral circulation. Echographic variables useful in a method of the invention include, for example, liver length (right kidney), irregular liver surface, liver heterogeneity, spleen length, ascites or collateral circulation. See, for example, Oberti et al., *Gastroenterol.* 113:1609-1616 (1997). It is understood that the analysis of these and other well known clinical or echographic variables can be useful in a method of the invention. Furthermore, a method of the invention encompasses determination of the clinical or echographic variable, for example, liver palpation, or can rely on one or more historic, or previously determined clinical or echographic variables.

**[0120]** Assays for detection of biochemical or serological markers useful in the invention are well known in the art and in many cases commercially available. Such assays include, but are not limited to, amplification based methods such as RT-PCR and other methods for quantitative analysis of RNA levels; immunoassays such as radioimmunoassays, enzyme-linked assays, two-antibody sandwich assays and quantitative western analysis; and assays for biological activity such as enzyme activity. Assays for PIIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 and apoB are commercially available from various sources as summarized in. Table 1.

**[0121]** Assays for additional biochemical or serological markers that can be combined with detection of  $\alpha$ 2-MG, HA and TIMP-1 in a method of the invention also are well known in the art. Fibronectin, for example, can be conveniently assayed by turbidimetric assay available from Roche Diagnostics (Mannheim, Germany). Pseudocholinesterase (PCHE) can be assayed using standard methodology available from Boehringer. Levels of N-acetyl-  $\beta$ -glucosaminidase ( $\beta$ -NAG) can be determined by assaying for enzymatic activity using a kit available from Cortecs diagnostics. Manganese superoxide dismutase (Mn-SOD) levels can be conveniently determined by ELISA using a kit available, for example, from Bender MedSystem. Glutathione peroxidase levels can be determined by assaying for enzymatic activity using, for example, a kit available from Randox Laboratories Ltd (Oceanside, CA).

TABLE 1  
COMMERCIAL SOURCES FOR FIBROTIC MARKER ASSAYS

Marker	Company	Assay	Catalog number
PIIINP	Orion Diagnostica (Espoo, Finland)	RIA	05903
laminin	Chemicon Intl. (Temecula, CA)	ELISA	ECM310
tenascin	Chemicon Intl. (Temecula, CA)	ELISA	ECM320
collagen IV	Iatron Laboratories (Tokyo, Japan)	RIA	KCAD1
YKL-40	Metra Biosystems (Mountain View, CA)	ELISA	8020
MMP-3	Amersham Pharmacia (Piscataway, NJ)	ELISA	RPN 2613
MMP-2	Amersham Pharmacia (Piscataway, NJ)	ELISA	RPN 2617
MMP-9/TIMP-1 complex	SBA Sciences (Turku, Finland)	ELISA	MP2215
sFas ligand	Bender MedSystems Diagnostics (Vienna, Austria)	ELISA	BMS260/2
TGF- $\beta$ 1	R&D Systems (Minneapolis, MN)	ELISA	DB100

(continued)

TABLE 1  
COMMERCIAL SOURCES FOR FIBROTIC MARKER ASSAYS

Marker	Company	Assay	Catalog number
IL-10	R&D Systems (Minneapolis, MN)	ELISA	HS100B
apoA1	AlerChek, Inc. (Portland, ME)	ELISA	A70101
apoA2	AlerChek, Inc. (Portland, ME)	ELISA	A70102
apoB	Sigma Diagnostics (St. Louis, MO)	IT*	357-A
* designates immunoturbidimetric			

**[0122]** Total or direct bilirubin, GGT, AST and ALT levels can be determined using an autoanalyser such as Hitachi 917 Automate (Mannheim, Germany) with Roche Diagnostics reagents. Albumin levels can be determined, for example, by the bromocresol green method as described in Dumas et al., Clin. Chim. Acta 31:87-96 (1971); and ferritin and  $\alpha$ -fetoprotein levels can be conveniently determined using, for example, an immunoassay available from Boehringer. In addition, levels of  $\alpha_1$  globulin,  $\alpha_2$  globulin,  $\beta$  globulin and  $\gamma$ -globulin can be determined, for example, by serum protein electrophoresis in an automatic system (Hydrasys and Hyrys, Sebia; Issy-Les-Moulineaux, France). Methods of determining prothrombin activity also are well known in the art and include the clotting method available from Organon Technika (West Orange, NJ). PGA index can be determined as described in Poynard et al., Gastroenterol. 100:1397-1402 (1991), and PGAA index also can be determined by well known methods as described in Naveau et al., Dig. Dis. Sci. 39: 2426-2432 (1994)).

**[0123]** Platelet counts, lymphocyte counts, mean corpuscular volume and related variables can be determined by a variety of methodologies using, for example, a Bayer-Technicon H2 analyser (Bayer-Technicon Instruments; Tarrytown, NY). Cholesterol levels can be determined by standard methodologies available, for example, from Boehringer. Thus, it is clear to the skilled person that a variety of methodologies, including but not limited to the above, are well known in the art and can be useful in the diagnostic methods of the invention.

#### $\alpha$ 2-MG/HA/YKL-40 panel

**[0124]** Also disclosed is a method of diagnosing the presence or severity of liver fibrosis in an individual by detecting  $\alpha$ 2-MG in a sample; detecting HA in a sample; detecting YKL-40 in a sample; and diagnosing the presence or severity of liver fibrosis in the individual based on the presence or level of  $\alpha$ 2-MG, HA and YKL-40. The method can be useful, for example, or differentiating no or mild (F0-F1) liver fibrosis from moderate to severe (F2-F4) liver fibrosis.

**[0125]** Disclosed is a method of diagnosing the presence or severity of liver fibrosis in an individual by determining the level of  $\alpha$ 2-MG protein in a sample from the individual; determining the level of HA in a sample from the individual; determining the level of YKL-40 protein in a sample from the individual; and diagnosing the presence or severity of liver fibrosis in the individual based on the levels of  $\alpha$ 2-MG protein, HA and YKL-40 protein. If desired, the levels of  $\alpha$ 2-MG protein, HA and YKL-40 protein each can be determined using an enzyme-linked assay.

**[0126]** Thus, diagnostic methods are disclosed which rely, in part, on determining a level of the fibrotic marker YKL-40 in a sample. YKL-40, also known as human cartilage glycoprotein 39 (HC gp-39), is named for a molecular weight of 40 kDa and the amino-terminal sequence of the protein, tyrosine-lysine-leucine (YKL). This glycoprotein, a mammalian member of the chitinase family (18-glycosylhydrolases), is a lectin that binds heparin and chitin and is produced by chondrocytes, synovial cells, activated macrophages, neutrophils and MG-63 osteosarcoma cells (Hakala et al., J. Biol. Chem. 268:25803-15810 (1993); Nyirkos and Golds, Biochem. J. 268:265-268 (1990); Renkema et al., Eur. J. Biochem. 251:504-509 (1998); Volck et al., Proc. Assoc. Am. Physicians 110:351-360 (1998); and Johansen et al., J. Bone Miner. Res. 7:501-511 (1992)). The pattern of YKL-40 expression in normal and diseased tissue indicates that this glycoprotein can function in extracellular matrix remodelling or tissue inflammation (Nyirkos and Golds, *supra*, 1990; Renkema et al., *supra*, 1998; and Verheijden et al., Arthritis Rheum. 40:1115-1125 (1997)). Furthermore, YKL-40 mRNA is expressed in liver, and initial studies have shown that YKL-40 expression is elevated in patients with chronic liver disease and that increased serum YKL-40 can be associated with fibrosis and fibrogenesis (Johansen et al., Scand. J. Gastroenterol. 32: 582-590 (1997); and Johansen, J. Hepatol. 32:911-920 (2000)).

**[0127]** Methods of determining a level of YKL-40 in samples such as serum and synovial fluid are well known in the art. For example, a radioimmunoassay for YKL-40 based on a rabbit antibody raised against YKL-40 is described in Johansen et al., Br. J. Rheumatology 32:949-955 (1993). In addition, a sandwich enzyme immunoassay in a microliter stripwell format is commercially available from Metra Biosystems. In the Metra Biosystems assay, the Fab fragment of

a biotin-conjugated monoclonal anti-YKL-40 antibody binds to streptavidin on the strip and captures YKL-40 in a sample. Alkaline phosphatase-conjugated polyclonal anti-YKL-40 antiserum binds the captured YKL-40 antigen, and alkaline phosphatase activity is detected with p-nitrophenyl phosphate substrate as an indication of YKL-40 concentration. It is understood that the methods of the invention can be practiced with these or other routine assays for detecting or determining a level of YKL-40 RNA or protein.

[0128] The following examples are intended to illustrate but not limit the present invention.

## EXAMPLE I

### MARKER PANELS FOR NON-INVASIVE DIAGNOSIS OF LIVER FIBROSIS

[0129] This example demonstrates that several serological markers can be combined together as a panel which is useful in differentiating fibrosis stages F2, F3 and F4 from stages F0 and F1 in HCV-infected patients.

[0130] Serum samples from 194 HCV patients positive for hepatitis C virus by RNA and immuno-analysis and having elevated alanine aminotransferase (ALT) levels were chosen at random from an existing serum library. Each of the patients had had a liver biopsy as part of their care. Patient samples were chosen to allow comparison of other routine blood markers and physical examination results attendant to routine medical care, including HCV viral load.

[0131] The inclusion criteria for the study were that the patient 1) have a confirmed hepatitis C infection at the time of liver biopsy and serum draw; 2) have undergone a liver biopsy as part of their medical care independent of the study; and 3) have previously given informed consent. Patients who did not give informed consent or who were incarcerated were excluded from the study.

[0132] Fibrosis scores (Metavir stage) for the 194 patients were established by histopathologic examination of a needle biopsy specimen prior to therapy according to the criteria set forth in The French Metavir Cooperative Study Group, Hepatol. 20:15-20 (1994). All Metavir fibrosis scores were established by the same pathologist. For all analyses, Metavir scores of F0 and F1 were grouped together as "no/mild" fibrosis, while scores of F2, F3 and F4 were grouped together as "moderate/severe" fibrosis. The fibrosis prevalence in the 194 patient group was determined to be 60% was based on the proportion of F2-F4 scores in the group as shown in Table 2.

TABLE 2 COMPOSITION OF THE 194 HCV PATIENT STUDY POPULATION BY FIBROSIS STAGE		
Fibrosis stage	Number	Total F0-F1 or F2-F4
F0	38	F0-F1 = 78
F1	40	
F2	40	F2-F4 = 116
F3	39	
F4	37	
Total	194	Prevalence = 59.8%

[0133] As shown in the table above, the panel of HCV patient samples included 37 samples with very high fibrosis stage (F4); 39 samples from patients with very low or zero fibrosis stage (F0): and 158 samples from patients with fibrosis stage F1, F2 or F3.

[0134] Serum samples were assayed for the presence of several putative fibrosis markers, including laminin, YKL-40, HA, TIMP-1, PIIINP, type IV collagen and  $\alpha$ 2-MG. Assays were performed using commercial kits according to manufacturers' instructions (see Table 3). The results obtained for the 194 samples analyzed for laminin, YKL-40, HA, TIMP-1, PIIINP, collagen type IV and  $\alpha$ 2-MG are shown in Table 4.

TABLE 3 COMMERCIALLY AVAILABLE KITS FOR DETECTION OF FIBROSIS MARKERS			
Marker	Manufacturer	Assay type	Catalogue number
Laminin	Chemicon Intl. (Temecula, CA)	ELISA	ECM310
YKL-40	Metra Biosystems (Mountain View, CA)	ELISA	8020

(continued)

TABLE 3  
COMMERCIALY AVAILABLE KITS FOR DETECTION OF FIBROSIS MARKERS

Marker	Manufacturer	Assay type	Catalogue number
HA	Corgenix (Westminster, CO)	ELISA	029001
TIMP-1	Amersham Pharmacia (Piscataway, NJ)	ELISA	RPN 2611
PIIINP	Orion Diagnostica (Espoo, Finland)	RIA	05903
collagen IV	Iatron Laboratories (Tokyo, Japan)	RIA	KCAD1
$\alpha$ 2-MG	Beckman Coulter	Nephelometry	449430

TABLE 4

RAW DATA FROM 194 HCV PATIENTS ANALYZED FOR LEVELS OF LAMININ, YKL-40, HA, TIMP-1, PIIINP, COLLAGEN TYPE IV AND  $\alpha$ 2-MG

Sample ID	Patient	Laminin (ng/ml)	YKL-40 (ng/ml)	HA (ng/ml)	TIMP-1 (ng/ml)	PIIINP (ng/ml)	Coll IV (ng/ml)	$\alpha$ 2-MG (mg/ml)
100010	B-A	175.244	81.608	15.730	1308.802	2.288	1.737	3.03
100038	P-B	151.888	67.220	9.288	917.104	2.049	2.617	2.01
100044	C-B	187.811	60.757	44.127	1610.690	3.883	3.408	5.03
100059	T-B	232.082	51.002	22.583	1077.343	2.297	1.901	2.29
100069	N-C	285.269	131.726	73.851	2381.222	8.034	2.954	4.05
100077	H-C	268.685	47.709	18.066	1122.818	2.260	3.159	1.75
100090	B-C	263.426	26.370	47.339	1380.182	3.526	2.561	3.30
100127	D-D	279.580	166.113	105.505	1180.879	3.343	2.804	3.54
100167	G-F	274.533	482.708	341.132	2523.637	9.745	5.110	3.98
100175	B-G	266.903	95.808	27.721	1178.105	4.345	3.911	2.67
100178	M-G	211.613	159.040	25.669	1176.718	2.357	3.795	1.93
100182	T-G	246.686	55.391	8.889	1308.815	2.924	2.468	2.86
100198	A-G	226.372	48.441	13.901	1126.962	2.595	0.819	3.03
100209	J-G	288.524	83.925	5.051	1081.470	5.173	0.801	2.73
100229	S-H	253.561	110.020	46.568	1391.433	4.905	4.410	3.12
100238	T-J	229.781	38.076	29.516	1190.567	2.626	3.141	2.68
100245	D-K	279.768	270.250	171.481	2310.561	5.876	4.713	4.01
100247	C-K	244.559	131.482	10.219	1405.454	2.297	1.089	2.50
100250	J-K	262.136	101.729	54.821	1155.963	4.192	2.655	4.04
100252	J-K	260.998	61.366	57.275	1560.856	3.498	7.040	3.24
100253	E-K	292.189	173.917	168.768	1652.033	9.252	7.336	3.59
100254	W-K	288.551	477.560	102.775	1580.756	5.300	4.188	4.01
100271	M-L	278.201	89.900	69.651	1140.761	4.092	4.066	2.94
100276	V-L	257.309	176.112	12.196	1088.369	1.985	4.147	3.57
100284	M-L	224.339	130.263	31.822	1104.885	3.653	2.542	3.14
100290	D-L	199.542	541.552	50.429	1550.943	4.399	6.368	3.40
100294	TRL	281.501	217.328	200.436	2340.630	11.006	7.016	4.20
100301	P-M	285.543	430.475	29.772	1884.061	4.453	3.524	4.26
100313	J-M	301.751	188.062	45.539	1852.125	3.610	4.773	4.22
100323	M-M	223.002	626.140	144.334	2382.232	8.873	8.042	4.09
100334	K-R	173.320	63.317	38.516	1290.231	4.327	4.753	2.46
100339	S-M	184.085	36.125	12.049	1268.153	1.865	2.211	2.39
100340	K-M	206.582	57.496	22.015	974.475	2.315	0.819	2.80

EP 1 487 395 B1

(continued)

RAW DATA FROM 194 HCV PATIENTS ANALYZED FOR LEVELS OF LAMININ, YKL-40, HA, TIMP-1, PIIINP, COLLAGEN TYPE IV AND  $\alpha$ 2-MG

	Sample ID	Patient	Laminin (ng/ml)	YKL-40 (ng/ml)	HA (ng/ml)	TIMP-1 (ng/ml)	PIIINP (ng/ml)	Coll IV (ng/ml)	$\alpha$ 2-MG (mg/ml)
5	100341	T-M	257.580	76.834	91.748	1492.882	3.976	4.793	3.00
	100343	D-M	334.202	140.629	49.322	1098.199	4.092	2.412	3.12
	100357	P-M	419.291	27.368	14.971	784.932	4.225	0.801	2.01
10	100374	K-N	300.366	28.231	36.608	1697.678	5.478	6.690	3.16
	100379	T-O	233.496	75.711	26.906	1437.939	2.086	2.437	3.70
	100382	C-P	206.796	44.461	6.034	989.007	2.214	4.409	2.14
	100397	R-R	223.006	66.474	13.912	981.736	4.091	3.814	3.53
15	100410	R-S	224.775	36.605	37.499	1152.258	4.592	6.466	3.43
	100438	D-Q	228.008	149.349	75.452	1682.636	7.734	7.039	3.89
	100451	H-R	248.528	526.840	226.386	1961.029	10.505	7.113	3.36
	100453	A-R	225.862	65.956	33.169	1421.632	2.786	3.465	3.79
	100454	O-R	220.481	56.892	32.531	1125.960	3.003	7.014	3.68
20	100456	S-S	241.591	46.274	30.745	1337.355	5.182	2.411	2.81
	100466	S-S	210.562	48.605	34.443	1482.475	4.286	4.754	3.27
	100470	D-S	229.912	162.039	55.053	1684.159	6.806	3.640	3.10
	100485	C-T	229.811	113.523	38.389	1247.547	3.901	3.291	4.11
25	100486	M-T	265.326	281.257	706.557	2716.589	15.362	11.975	2.82
	100505	L-W	229.363	33.843	23.305	1149.367	3.397	2.889	3.29
	100519	R-W	204.646	68.632	10.431	845.571	3.852	6.815	2.52
	100547	S-G	223.959	75.711	8.257	1000.623	4.286	3.090	2.92
	100638	J-P	265.819	264.250	68.361	2095.698	15.945	7.418	4.67
30	100640	M-V	170.293	93.770	17.728	1200.584	4.755	5.561	3.50
	100006	L-A	135.628	75.349	79.430	1354.782	6.612	3.477	3.14
	100009	R-A	157.239	72.429	21.947	932.635	2.080	3.050	1.96
	100011	A-B	136.197	251.237	149.932	2004.294	7.600	4.853	3.57
35	100016	E-A	161.133	272.434	186.536	1900.341	9.341	9.071	3.08
	100021	E-AV	184.000	537.630	102.420	2456.883	4.863	6.157	3.97
	100023	C-B	126.346	194.523	47.976	1540.914	7.000	4.488	3.21
	100027	D-B	133.660	75.820	33.912	1519.528	2.966	3.286	3.51
	100030	R-B	140.584	50.007	153.135	1219.549	3.237	5.200	1.89
40	100035	K-B	124.645	37.383	60.934	1214.060	3.582	3.620	2.41
	100036	G-B	152.864	87.596	369.681	1305.790	3.163	4.391	2.71
	100041	M-B	168.422	42.376	143.372	1502.562	6.667	3.692	3.10
	100042	M-B	138.754	211.387	266.568	2899.870	8.233	8.559	3.98
45	100043	C-B	111.743	30.883	17.447	1168.327	5.488	2.343	2.76
	100045	V-B	164.940	241.063	221.249	2010.088	9.097	4.512	3.76
	100051	K-B	154.743	222.409	131.122	1600.554	4.863	4.075	3.43
	100055	D-B	146.817	110.018	84.447	1827.668	3.188	4.439	5.72
	100065	G-B	134.349	72.429	112.148	1455.905	3.353	3.002	4.02
50	100071	R-C	135.011	74.407	30.352	1485.573	2.820	3.120	3.29
	100073	G-C	146.785	63.761	43.312	1530.873	3.027	2.040	< 0.75
	100074	L-C	151.514	80.248	49.917	1647.700	5.036	3.286	3.11
	100078	R-C	163.144	213.365	45.839	1399.880	3.393	2.297	2.61
55	100081	A-C	147.915	45.862	56.686	1346.315	4.056	4.707	2.37
	100084	G-C	144.665	43.130	116.238	1736.670	5.337	6.702	< 0.75
	100091	P-C	171.782	215.249	33.321	1999.807	5.096	3.835	3.75
	100093	M-C	133.786	35.499	105.726	1499.707	5.983	5.876	2.85



EP 1 487 395 B1

(continued)

RAW DATA FROM 194 HCV PATIENTS ANALYZED FOR LEVELS OF LAMININ, YKL-40, HA, TIMP-1, PIIINP, COLLAGEN TYPE IV AND  $\alpha$ 2-MG

	Sample ID	Patient	Laminin (ng/ml)	YKL-40 (ng/ml)	HA (ng/ml)	TIMP-1 (ng/ml)	PIIINP (ng/ml)	Coll IV (ng/ml)	$\alpha$ 2-MG (mg/ml)
5	100099	S-C	174.239	49.159	28.163	1392.574	4.381	5.876	3.18
	100100	D-C	181.284	68.095	82.324	1613.489	5.552	5.547	3.71
	100103	J-C	151.396	74.849	55.720	1666.282	4.771	3.955	3.57
10	100104	C-C	128.182	38.890	13.719	1100.784	3.798	5.324	2.21
	100106	S-C	170.685	41.811	37.449	1280.697	5.096	4.464	2.52
	100107	J-C	103.835	27.397	32.334	1416.481	2.910	3.740	2.57
	100108	J-C	148.294	145.629	52.822	1884.389	4.484	3.597	4.62
15	100115	S-DLT	134.784	108.134	96.415	1597.696	8.860	5.225	3.48
	100121	R-D	149.335	74.878	34.109	1759.752	5.912	3.716	4.36
	100124	R-D	134.766	130.367	35.486	1569.219	2.489	4.877	1.21
	100125	B-D	170.790	67.078	97.770	2245.776	7.261	5.876	3.63
	100126	D-D	134.313	117.116	65.560	1970.476	2.775	3.788	3.04
20	100129	E-D	159.707	60.388	28.962	1651.995	5.195	4.902	3.41
	100131	J-D	155.166	119.774	31.852	1579.186	3.015	3.405	2.73
	100133	M-D	146.280	24.371	75.729	2098.560	3.225	3.788	1.74
	100135	H-E	167.472	41.600	66.767	1369.735	3.200	3.429	4.20
25	100137	D-E	158.406	25.104	68.740	1346.906	3.828	3.405	2.69
	100139	S-E	139.877	38.484	42.708	1388.605	4.215	2.814	4.05
	100140	L-E	158.942	30.695	181.056	1585.482	7.476	4.977	3.32
	100141	W-E	136.761	185.300	179.774	2045.873	9.097	9.872	2.89
	100142	R-E	119.383	62.037	16.170	888.744	3.286	2.673	2.07
30	100143	D-E	131.717	33.779	29.179	1072.170	2.978	3.144	1.91
	100147	D-E	132.426	77.159	35.912	1285.138	3.515	2.696	2.12
	100150	D-E	120.207	19.056	155.043	1488.729	2.298	4.196	3.66
	100151	D-F	125.885	35.735	51.625	1243.711	3.447	3.525	3.05
35	100155	C-H	146.728	29.137	54.330	1242.340	4.733	3.573	3.15
	100159	M-H	136.303	31.336	106.675	1567.716	4.151	3.525	3.51
	100161	JF-F	155.052	1710.890	572.598	1966.460	6.226	4.634	4.27
	100163	M-F	153.221	785.420	211.173	2167.501	8.269	5.698	3.57
	100175a	B-G	148.403	55.347	130.093	1282.502	5.296	4.537	2.79
40	100181	M-G	137.986	69.735	31.119	1384.651	2.813	3.097	1.89
	100183	D-G	168.842	181.909	58.358	1499.596	3.101	3.405	3.86
	100186	M-G	184.148	2258.120	347.854	5271.196	11.670	7.756	3.82
	100200	R-G	148.660	158.906	143.510	1939.499	5.530	6.080	4.02
45	100208	R-G	156.210	94.021	36.624	1379.174	5.339	4.366	< 0.75
	100221	ND-H	117.196	38.393	88.913	1375.112	2.610	5.274	3.76
	100222	J-H	106.131	34.544	31.603	1054.973	2.580	2.955	3.40
	100229a	S-H	125.123	53.139	73.989	1567.731	3.828	3.788	2.78
	100237	J-J	140.718	397.625	578.952	1824.407	11.836	6.675	2.46
50	100268	C-L	155.864	76.151	86.977	2060.140	10.963	6.310	3.66
	100270	T-L	176.060	24.738	38.749	1579.990	2.549	2.625	3.35
	100278	S-L	153.789	48.840	52.524	1367.051	3.039	3.167	1.91
	100279	L-LG	163.352	139.202	39.492	1223.652	2.921	4.099	2.37
55	100287	R-L	164.414	636.110	232.253	3285.078	14.450	9.448	4.19
	100291	MS-L	152.863	197.500	42.925	2144.445	2.180	1.993	3.78
	100293	D-L	147.479	104.509	27.209	1559.538	3.151	2.696	1.83
	100302	A-M	201.715	1021.070	159.330	3317.515	12.498	8.383	4.83

**EP 1 487 395 B1**

(continued)

**RAW DATA FROM 194 HCV PATIENTS ANALYZED FOR LEVELS OF LAMININ, YKL-40, HA, TIMP-1, PIIINP, COLLAGEN TYPE IV AND  $\alpha$ 2-MG**

	<b>Sample ID</b>	<b>Patient</b>	<b>Laminin (ng/ml)</b>	<b>YKL-40 (ng/ml)</b>	<b>HA (ng/ml)</b>	<b>TIMP-1 (ng/ml)</b>	<b>PIIINP (ng/ml)</b>	<b>Coll IV (ng/ml)</b>	<b><math>\alpha</math>2-MG (mg/ml)</b>
5	100306	JT-M	125.203	115.289	83.960	1722.987	3.842	2.413	4.16
	100307	D-M	128.095	23.612	10.882	1378.118	3.339	3.382	1.42
	100313a	J-McA	164.201	192.417	76.599	1966.287	3.828	5.523	3.89
10	100315	K-MF	153.427	113.449	112.430	2118.580	3.515	4.561	4.46
	100317	M-McM	165.245	94.693	144.632	1611.495	4.588	3.215	1.84
	100320	D-M	159.724	782.150	106.161	1581.345	13.215	5.597	2.33
	100322	R-M	120.098	39.914	51.444	1443.789	2.809	3.644	3.21
15	100327	K-R	168.143	194.521	101.231	1738.827	4.295	3.835	4.35
	100336	ES -M	165.374	135.711	36.329	1556.071	2.932	2.508	3.49
	100347	E-M	173.070	69.889	16.945	1710.951	4.808	3.859	3.29
	100348	A-M	207.186	75.06	301.583	1334.475	3.299	4.585	3.36
	100350	J-M	154.867	4.418	22.250	1388.371	4.087	3.238	1.60
20	100358	A-M	140.022	15.549	88.786	1247.147	4.502	4.682	1.57
	100365	A-M	96.324	26.329	43.344	1170.887	5.381	2.040	2.12
	100367	B-M	161.274	30.273	46.174	1469.088	3.089	2.790	3.59
	100388	A-P	230.782	275.681	938.015	4245.175	20.496	9.669	5.98
25	100395	D-R	125.908	24.226	15.309	1299.599	2.478	4.415	3.52
	100397a	R-R	179.186	56.479	100.853	1455.947	9.769	4.172	3.62
	100398	C-R	151.391	29.397	11.833	1100.156	2.652	2.932	2.28
	100403	L-P	179.196	321.607	350.713	2061.218	8.938	4.977	2.28
	100404	ML-P	179.163	1060.240	141.902	2248.495	5.959	4.123	4.07
30	100414	S-S	184.451	70.941	40.126	1048.761	2.549	3.026	3.18
	100424	A-P	158.538	62.439	167.519	1320.841	5.509	5.448	3.88
	100443	J-R	112.348	40.703	21.470	1045.054	2.663	5.647	4.34
	100450	M-R	186.892	200.744	203.399	1287.107	2.586	4.040	3.77
35	100472	T-S	127.877	119.759	21.867	797.753	2.787	3.962	1.97
	100474	J-S	118.319	55.427	19.699	939.914	2.287	3.524	3.15
	100482	J-T	125.011	33.428	39.749	1099.832	5.159	4.386	2.56
	100483	J-T	136.978	18.006	33.467	1003.831	2.586	4.485	2.06
	100488	M-T	178.106	51.908	123.723	1345.743	4.464	6.053	3.84
40	100495	J-V	180.283	219.322	93.680	1734.925	8.387	4.064	3.82
	100502	D-V	114.380	153.296	22.649	1136.728	2.774	4.757	4.47
	100503	J-W	149.928	453.095	92.918	1422.332	5.581	4.114	3.15
	100513	M-W	117.649	53.027	100.853	1335.730	5.092	4.757	3.74
45	100528	D-W	189.040	37.248	26.158	1103.495	4.642	3.215	2.90
	100530	M-W	106.100	40.234	20.490	924.291	3.297	3.988	2.29
	100534	D-A	135.702	37.357	91.410	1421.072	3.374	3.135	3.26
	100539	M-DB	167.910	474.960	104.470	215.8.238	4.957	5.566	3.99
	100540	A-B	135.980	54.815	183.589	1881.935	3.068	4.485	3.34
50	100546	D-F	113.363	75.942	47.682	1207.833	2.633	3.055	1.57
	100557	T-L	121.746	58.286	27.704	1297.060	3.842	2.060	1.83
	100560	C-N	194.265	142.696	91.041	2338.303	6.332	3.267	4.25
	100564	D-R	169.241	250.713	65.865	2407.901	5.502	4.534	3.52
55	100569	J-DC	160.145	64.634	43.744	1349.485	5.962	3.421	3.78
	100572	K-K	162.517	260.632	209.581	1729.746	9.292	6.191	2.70
	100585	K-Z	171.277	162.336	126.433	2030.404	8.907	4.881	4.71
	100594	R-M	119.193	216.295	42.261	1678.540	3.545	3.602	3.26

(continued)

**RAW DATA FROM 194 HCV PATIENTS ANALYZED FOR LEVELS OF LAMININ, YKL-40, HA, TIMP-1, PIIINP, COLLAGEN TYPE IV AND  $\alpha$ 2-MG**

Sample ID	Patient	Laminin (ng/ml)	YKL-40 (ng/ml)	HA (ng/ml)	TIMP-1 (ng/ml)	PIIINP (ng/ml)	Coll IV (ng/ml)	$\alpha$ 2-MG (mg/ml)
100603	M-S	119.071	61.460	34.332	1693.071	4.464	4.188	2.92
100611	P-F	178.856	269.956	92.721	896.077	3.960	3.524	3.96
100614	J-McA	204.794	245.159	322.970	3470.966	11.393	6.814	5.32
100617	C-W	159.292	51.343	38.850	1504.544	5.859	4.974	3.60
100630	E-AV	140.072	34.778	59.454	1091.420	1.969	3.161	3.34
100637	R-B	179.987	59.477	137.723	2077.095	4.726	3.002	4.24
101013	T-H	177.189	507.415	237.499	1556.336	9.381	8.910	3.36
101118	G-S	163.553	282.057	150.713	2348.845	6.231	7.161	3.93
101137	S-S	175.291	2049.970	715.601	3318.137	11.450	8.458	3.71
101257	M-F	155.324	40.730	49.441	1082.686	3.240	3.421	1.71
101275	J-C	121.598	143.292	45.227	1454.509	1.915	2.788	4.69
101284	R-F	123.312	16.825	42.048	1072.891	4.386	3.679	1.64
101321	H-P	180.159	180.091	367.681	2235.931	11.450	10.045	3.37
101322	A-P	133.640	269.262	244.520	2015.508	9.119	8.458	4.01
101335	L-S	164.947	64.238	208.719	2545.531	11.018	6.930	3.90
101336	L-S	156.847	741.300	72.664	1387.281	6.097	4.683	3.73
101351	P-F	128.701	12.461	99.050	1557.519	5.486	4.161	4.01
101441	R-H	104.993	176.909	354.512	1338.637	5.271	6.445	3.50
101478	G-S	142.642	63.543	149.266	1296.972	3.859	4.411	2.64
101565	D-A	132.117	74.355	187.250	1206.734	5.962	4.782	2.92

**[0135]** Clinical performance parameters were analyzed for the combinations of markers best able to differentiate the presence of significant fibrosis (F2-F4) from no/mild fibrosis (F0-F1) using various statistical algorithms. The statistical algorithms analyzed included univariate analysis, receiver operating characteristic curves (ROC), logistic regression, discriminant function analysis, and factorial design optimization.

**[0136]** The results of ROC analysis are shown in Table 5. The area under the curve (AUC) values represent relative diagnostic value of a single marker at the indicated cut-off. As can be seen by the decreasing AUC values, HA was shown to have the best diagnostic value when used alone at the indicated cut-off, followed by PIIINP, TIMP-1,  $\alpha$ 2-MG and collagen type IV;

<b>TABLE 5 ROC ANALYSIS</b>				
	AUC	Sensitivity	Specificity	Cutoff
HA	0.821	90.0%	62.0%	35.5 ng/ml
PIIINP	0.777	90.8%	39.2%	3.0 ng/ml
TIMP-1	0.773	90.8%	43.0%	1190.6 ng/ml
$\alpha$ 2-macroglobulin	0.722	90.5%	34.6%	2.4 mg/ml
Collagen Type IV-7S	0.726	90.8%	24.1%	2.79 ng/ml
YKL-40	0.696	90.8%	19.0%	34.5 ng/ml
Laminin	0.524	90.7%	16.5%	125.2 ng/ml

**[0137]** Clinical performance parameters for various combinations of fibrosis markers are shown in Table 6. The best subsets, including single markers as well as combinations of two to four markers and algorithms for discriminating F0-F1 from F2-F4, were generated by logistic regression. The markers included PIIINP,  $\alpha$ 2-MG, laminin and type IV collagen. As shown in Table 5, the diagnostic performance parameters (sensitivity, specificity, PPV and NPV) were similar for the

two, three and four-marker combinations identified by logistic regression in the study population, which had a fibrosis prevalence of about 60% (see lines 2-4 and 6-9).

**[0138]** As shown in Table 6, line 5, stepwise discriminant function analysis (SAS) resulted in identification of the 3-marker subset (PIIINP,  $\alpha$ 2-MG and laminin). The clinical performance of this combination was similar to the marker combinations identified using logistic regression.

**[0139]** Design of experiments software (DOE KISS, Build 8, Air Academy Associates) was used to simultaneously optimize the cutoffs of multiple variables to obtain the best performance of the panel of tests in predicting fibrosis. Using DOE KISS, a computer-aided central composite design for a combination of markers was generated; this design matrix consisted of a series of combinations of cutoffs for each of the markers in the combination. The results from these experiments (sensitivity, specificity and accuracy) for differentiating F0-F1 from F2-F4 fibrosis were recorded in the design sheet in DOE. Regression analysis was performed for each of the parameters (sensitivity, specificity and accuracy) to give cutoff values for each of the variables in the combination to achieve maximum performance for that parameter.

**[0140]** The five markers with best diagnostic performance in an ROC analysis (highest AUC) were HA, PIIINP, TIMP-1,  $\alpha$ 2-MG and type IV collagen (see Table 5). Cutoffs for each of the markers in this 5-marker panel were optimized for maximum accuracy. The results shown in Table 6, line 10, indicate that, at the optimum accuracy (69.6%), the specificity was too low to be useful (32.9%) while the sensitivity was high (94.8%). Similar results were obtained when the markers were optimized for sensitivity or specificity. Regression analysis showed that TIMP-1 did not have a significant effect on the accuracy, sensitivity or specificity of this 5-marker panel.

**[0141]** A similar 4-marker panel was analyzed by DOE as shown in Table 6, line 11. With TIMP-1 excluded, the four-marker panel was optimized for accuracy (77.8%) to give a sensitivity and specificity of 79.1% and 79.5%, respectively. These results demonstrate that the four-marker panel of HA, PIIINP,  $\alpha$ 2-MG and collagen IV has more value in differentiating F0-F1 fibrosis from F2-F4 fibrosis than a five-marker panel made up of HA, PIIINP,  $\alpha$ 2-MG, collagen IV and TIMP-1.

**[0142]** Several three-marker subsets of the four-marker panel also were analyzed by DOE. Line 12 shows the results obtained for the combination of HA, collagen and  $\alpha$ 2-MG with the results optimized for accuracy. This three-marker panel gave a very low specificity of less than 30%. In contrast, when a three-marker panel made up of HA, PIIINP and  $\alpha$ 2-MG was optimized for accuracy, performance was similar to the four-marker panel (compare lines 13 and 11 of Table 6).

**[0143]** A similar analysis of the two-marker panel of HA and  $\alpha$ 2-MG gave the results shown in line 14 of Table 6. This combination gave an improvement in specificity over the three-marker panel of HA, PIIINP and  $\alpha$ 2-MG (84.4% compared to 78.3%).

**TABLE 6**  
PERFORMANCE OF VARIOUS MARKER PANELS

	Markers	Method/ Model	Sens.	Spec.	Prevalence 59.3%				Prevalence 20%			
					PPV	NPV	Acc.		PPV	NPV	Acc.	
1	HA	N/A cut-off 60ug/ml	64.96%	82.05%	84.44%	60.95%			47.50%	90.35%	78.63%	
2	PIIINP	Logistic Best subset of 1	74.36%	58.97%	73.11%	60.53%			31.18%	90.20%	62.05%	
3	PIIINP, AMG	Logistic Best subset of 2	80.53%	63.63%	76.86%	68.54%			31.18%	90.20%	62.05%	
4	PIIINP, AMG, Laminin	Logistic Best subset of 3	78.76%	67.53%	78.44%	67.95%			37.75%	92.71%	69.78%	
5	PIIINP, AMG, Laminin	Discriminant Stepwise selection	78.76%	67.53%	78.44%	67.95%			37.75%	92.71%	69.78%	
6	PIIINP, AMG, Laminin, Coll IV	Logistic Best subset of 4	78.76%	64.94%	77.11%	67.09%			35.96%	92.44%	67.70%	
7	PIIINP, AMG, Iminin, YKL-40	Logistic Second best subset of 4	77.87%	67.53%	78.25%	67.05%			37.48%	92.43%	69.60%	
8	PIIINP, AMG, Coll IV, YKL-40	Logistic Third best subset of 4	78.76%	70.13%	79.82%	68.76%			39.73%	92.96%	71.86%	
9	PIIINP, AMG, TIMP-1	Logistic "Forced" selection	78.76%	64.94%	77.11%	67.09%			35.96%	92.44%	67.70%	
10	HA, PIIINP, AMG Coll IV, TIMP-1	DOE (for acc.) N/A	94.78%	32.91%	67.28%	81.25%	69.59%		26.1%	96.2%	45.3%	
11	HA, PIIINP, Coll IV, AMG	DOE (for acc.) N/A	79.13%	75.95%	82.73%	71.43%	77.84%		45.13%	93.57%	76.59%	
12	HA, CollIV, AMG	DOE (for acc.) N/A	95.65%	29.11%	66.27%	82.14%	68.56%		25.2%	96.4%	42.4%	
13	HA, PIIINP, AMG	DOE (for acc.) N/A	78.26%	75.95%	82.57%	70.59%	77.32%		44.86%	93.32%	76.41%	
14	HA, AMG (B)	DOE (for acc.) N/A.	84.35%	73.42%	82.20%	76.32%	79.90%		44.24%	94.94%	75.60%	
15	TIMP-1, HA, AMG	DOE (for acc.)	83.48%	75.95%	83.48%	75.95%	80.41%		46.46%	94.84%	77.46%	
16	YKL-40, HA, AMG	DOE (for acc.)	82.61%	75.95%	83.33%	75.00%	79.90%		46.20%	94.59%	77.28%	
17	HA, reflex with Coll IV and AMG	Logistic Positives reflexed	86.32%	70.52%	81.45%	77.46%	80.00%		42.26%	95.38%	73.68%	

**[0144]** TIMP-1, which was observed to be a good discriminator of fibrosis in the univariate analysis, was added to the two-marker panel. As shown in line 15, the performance of the HA,  $\alpha$ 2-MG and TIMP-1 three-marker panel was similar to that obtained with the two-marker panel, and the sensitivity was improved as compared to the three-marker HA/PIINP/ $\alpha$ 2-MG panel (83.5% sensitivity compared to 78.3%). Furthermore, in preliminary regression analysis, TIMP-1 contributed significantly to discrimination of fibrosis in a study population with a high prevalence of severe fibrosis.

**[0145]** Another three-marker panel, made up of HA,  $\alpha$ 2-MG and YKL-40, also was optimized for accuracy in differentiating F0-F1 from F2-F4 fibrosis. As shown in Table 6, line 16, this three marker panel had a performance similar to the  $\alpha$ 2-MG/HA/TIMP-1 panel.

**[0146]** In sum, these results indicate that a  $\alpha$ 2-MG/HA/TIMP-1 or  $\alpha$ 2-MG/HA/YKL-40 panel can be useful in differentiating F0-F1 from F2-F4 fibrosis.

## EXAMPLE II

### DUAL OPTIMIZATION STRATEGY FOR ANALYSIS OF THE $\alpha$ 2-MG/HA/TIMP THREE-MARKER PANEL

**[0147]** This example describes the use of multiple cut-offs for  $\alpha$ 2-MG-, HA- and TIMP-1 to achieve a relatively high degree of accuracy in a subset of a total patient population assayed.

**[0148]** Using the three-marker panel  $\alpha$ 2-MG/HA/TIMP-1 with cutoffs for  $\alpha$ 2-MG, HA- and TIMP-1 set at 35 ng/ml, 2 mg/ml, and 1000 ng/ml, respectively, samples were determined to be positive when all three of their variables were above the cut-off values, and were therefore negative when one or more of the  $\alpha$ 2-MG, HA or TIMP-1 levels were below the assigned cut-off value. As shown in Table 7, in the 194 patient population, there were a total of 72 negative results, 15 of which were false negatives, giving a negative predictive value (NPV) of 79% at the study prevalence of about 60% fibrosis%. At a prevalence of 30%, which is typical of the prevalence in a hepatology clinic, the negative predictive value is over 92%, which is useful in ruling out the presence of F2-F4 fibrosis (likelihood ratio 0.22).

**[0149]** Furthermore, of the 122 test positives using the 35 ng/ml, 2 mg/ml, and 1000 ng/ml cut-offs, 21 of the test positives were false, giving a positive predictive value (PPV) of 82.8%. However, at a more typical prevalence of 30% fibrosis, the positive predictive value falls to about 58% (see Table 7). Thus, in a population with a typical prevalence, a positive result would not have sufficient predictive value to be useful as a diagnostic.

**[0150]** In order to increase the positive predictive value for at least a subset of the total patient population, samples positive by the primary analysis were further evaluated for positivity for the three markers using a second set of cut-off values which were higher than the first set. By evaluating those samples positive after a primary analysis at higher cutoffs, the severe fibrosis samples within this group can be determined to be positive with a relatively high predictive value. Those samples that test negative by the secondary evaluation are considered "indeterminate" in that their fibrosis status cannot be determined with good predictive value.

TABLE 7									
PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH DUAL OPTIMIZATION STRATEGY IN POPULATIONS WITH VARIOUS DISEASE PREVALENCES									
Optimization for sensitivity to rule out fibrosis X1 = 2.0 mg/ml for $\alpha$ 2-MG Y1 = 35.00 ng/ml for HA Z1 = 1000.00 ng/ml for TIMP-1									
	Prevalence 0.598		Prevalence 0.300		Prevalence 0.200		Prevalence 0.100		
	Fib +	Fib -	Fib +	Fib -	Fib +	Fib -	Fib +	Fib -	
Test +	101	21	261	188	174	215	87	242	329
Test -	15	57	39	512	26	585	13	658	671
	116	78	300	700	200	800	100	900	1000
Sens.	87.07%	LR +	87.07%	LR +	87.07%	LR +	87.07%	LR +	3.23
Spec.	73.08%	LR -	73.08%	LR -	73.08%	LR -	73.08%	LR -	0.18
PPV	82.79%		58.09%		44.71%		26.43%		
NPV	79.17%		92.95%		95.76%		98.07%		
Accuracy	81.44%		77.27%		75.88%		74.48%		
TABLE 7									
PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH DUAL OPTIMIZATION STRATEGY IN POPULATIONS WITH VARIOUS DISEASE PREVALENCES									
Optimization for specificity to rule in fibrosis X1 = 2.0 mg/ml for $\alpha$ 2-MG Y1 = 60.00 ng/ml for HA Z1 = 1575.00 for TIMP-1 ng/ml									
	Prevalence -0.598		Prevalence 0.300		Prevalence 0.200		Prevalence 0.100		
	Fib +	Fib -	Fib +	Fib -	Fib +	Fib -	Fib +	Fib -	
Test +	53	1	137	9	91	10	46	12	57
Test Equiv	48	20	124	179	83	205	41	231	272
	101	21	261	188	174	215	87	242	329
Sens.	52.48%	LR +	52.48%	LR +	52.48%	LR +	52.48%	LR +	11.02
Spec.	95.24%	LR -	95.24%	LR -	95.24%	LR -	95.24%	LR -	0.50
PPV	98.15%		93.86%		89.91%		79.84%		
NPV	29.41%		59.11%		71.25%		84.80%		
Accuracy	59.84%		70.40%		76.12%		83.93%		

(continued)

TABLE 7															
PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH DUAL OPTIMIZATION STRATEGY IN POPULATIONS WITH VARIOUS DISEASE PREVALENCES															
Final performance after dual optimization															
Prevalence Fib +      Fib -		0.598		Prevalence Fib +      Fib -		0.300		Prevalence Fib +      Fib -		0.200		Prevalence Fib +      Fib -		0.100	
Sens. Spec. PPV NPV Accuracy	53	1	54	9	137	9	146	91	10	102	46	12	57	57	
	63	77	140	691	163	691	854	109	790	898	54	888	943	943	
	116	78	194	700	300	700	1000	200	800	1000	100	900	1000	1000	
	45.69%	LR +	35.64	LR +	45.69%	LR +	35.64	45.69%	LR +	35.64	45.69%	LR +	35.64	35.64	
	98.72%	LR -	0.55	LR -	98.72%	LR -	0.55	98.72%	LR -	0.55	98.72%	LR -	0.55	0.55	
	98.15%			93.86%				89.91%			79.84%				
	55.00%			80.92%				87.91%			94.24%				
	67.01%			82.81%				88.11%			93.42%				
TABLE 7															
PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH DUAL OPTIMIZATION STRATEGY IN POPULATIONS WITH VARIOUS DISEASE PREVALENCES															
Population Prevalence	194 0.598	Population Prevalence	1000 0.300	Population Prevalence	1000 0.200	Population Prevalence	1000 0.100								
False. Negative	15	False Negative	39	False Negative	26	False Negative	13								
False Positive	1	False Positive	9	False Positive	10	False Positive	12								
Total Incorrect	16	Total Incorrect	48	Total Incorrect	36	Total Incorrect	24								
Total Correct	110	Total Correct	649	Total Correct	676	Total Correct	703								
2nd Round															
Fib + Equivocal	48	Fib + Equivocal	124	Fib + Equivocal	83	Fib + Equivocal	41								
Fib - Equivocal	20	Fib - Equivocal	179	Fib - Equivocal	205	Fib - Equivocal	231								
Total Equivocal	68	Total Equivocal	304	Total Equivocal	288	Total Equivocal	272								
TABLE 7															
PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH DUAL OPTIMIZATION STRATEGY IN POPULATIONS WITH VARIOUS DISEASE PREVALENCES															
Final performance after dual optimization without equivocal															
Population Prevalence	194 0.598	Population Prevalence	1000 0.300	Population Prevalence	1000 0.200	Population Prevalence	1000 0.100								
False. Negative	15	False Negative	39	False Negative	26	False Negative	13								
False Positive	1	False Positive	9	False Positive	10	False Positive	12								
Total Incorrect	16	Total Incorrect	48	Total Incorrect	36	Total Incorrect	24								
Total Correct	110	Total Correct	649	Total Correct	676	Total Correct	703								



(continued)

Final performance after dual optimization without equivocal												
Test +	53	1	54	137	9	146	91	10	102	46 .	12	57
Test -	15	57	72	39	512	550	26	585	610	13	658	671
Sens.	68	58	<b>126</b>	176	521	<b>696</b>	117	595	<b>712</b>	59	669	<b>728</b>
Spec.	77.94%	LR +	45.21	77.94%	LR +	45.21	77.94%	LR +	45.21	77.94%	LR +	45.21
PPV	98.28%	LR -	0.22	98.28%	LR -	0.22	98.28%	LR -	0.22	98.28%	LR -	0.22
NPV	98.15%			93.86%			89.91%			79.84%		
	79.17%			92.95%			95.76%			98.07%		
Accuracy	87.30%			93.14%			94.93%			96.64%		
% of pop.												
Test +	27.8%			Test +	14.6%		Test +	10.2%		Test +	5.7%	
Test -	37.1%			Test -	55.0%		Test -	61.0%		Test -	67.1%	
Equiv	35.1%			Equiv	30.4%		Equiv	28.8%		Equiv	27.2%	

**[0151]** Table 7 shows performance of the  $\alpha$ 2-MG/HA/TIMP-1 panel assay with the dual optimization strategy. The primary cut-offs were set at 2.0 mg/ml, 35 ng/ml and 1000 ng/ml to achieve a relatively high sensitivity in the primary analysis. Any samples having all three of their  $\alpha$ 2-MG, HA and TIMP-1 levels above the assigned cut-off values were indicated to be positive. The 122 test positives obtained by the primary analysis were re-evaluated using 2.0 mg/ml, 60 ng/ml and 1575 ng/ml as the  $\alpha$ 2-MG, HA and TIMP-1 cut-offs and the criteria that the samples must have  $\alpha$ 2-MG, HA and TIMP-1 values above the assigned cut-off values to be positive.

**[0152]** Using the second set of cut-off values, 54 of the 122 patents were determined to be positive, only 1 of which was a false positive. The positive predictive value was 98.2% at 59.8% fibrosis prevalence, and was 93.9% at the more typical 30% fibrosis prevalence. In sum, of the 194 patients, 72 were classified as negative and 54 were classified as positive, while 68 samples had indeterminate results and could not be definitively classified. Furthermore, when the indeterminate samples are excluded, the three-marker assay has a positive predictive value of more than 93% and a negative predictive value of close to 93% in a typical population having a 30% fibrosis prevalence.

**[0153]** Table 8 shows a comparison of the performance of the  $\alpha$ 2-MG/HA/TIMP-1 three-marker panel with the six marker panel described in Poynard et al., Lancet 357:1069 (2001).

TABLE 8							
COMPARISON OF PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH 6 MARKER PANEL OF POYNARD ET AL.							
Prometheus				Poynard et al.			
	Biopsy				Biopsy		
	Fib +	Fib -			Fib +	Fib -	
Test +	53	1	54	Test + (>.08)	45	5	50
Test -	15	57	72	Test - (<0.20)	13	106	119
Equivocal	48	20	68	Equivocal	80	90	170
Total Pop.	116	78	194	Total Pop.	138	201	339
Prevalence	.05979			Prevalence	0.4071		
Sensitivity	0.7794			Sensitivity	0.7759		
Specificity	0.9828			Specificity	0.9550		
PPV	0.9815			PPV	0.900		
NPV	0.7917			NPV	0.8908		
Accuracy	0.8730			Accuracy	0.8935		
% Equivoc	.03505	68/194		% Equivoc	.0515	170/339	
False Pos	1	of 54 test +	1.85%	False Pos	5	of 50 test +	10.00%
False Neg	15	of 72 test -	20.83%	False Neg	13	of 119 test -	10.92%
Prometheus				Poynard et al.			
	Biopsy				Biopsy		
	Fib +	Fib -			Fib +	Fib -	
Test +	186	8	194	Test + (>.08)	133	15	147
Test -	53	433	486	Test - (<0.20)	38	313	351
Equivocal	168	152	320	Equivocal	236	265	501
	407	593	1000	Total Pop.	407	593	1000
Prevalence	0.4071			Prevalence	0.4071		
Sensitivity	0.7794			Sensitivity	0.7759		

(continued)

TABLE 8							
COMPARISON OF PERFORMANCE OF $\alpha$ 2-MG/HA/TIMP-1 PANEL WITH 6 MARKER PANEL OF POYNARD ET AL.							
Prometheus				Poynard et al.			
	Biopsy				Biopsy		
Specificity	0.9828			Specificity	0.9550		
PPV	.9607			PPV	0.9000		
NPV	.8917			NPV	0.8907		
Accuracy	.09113			Accuracy	0.8935		
% test pos	19.4%			% test pos	14.7%		
% test neg	48.6%			% test neg	35.1%		
% Equivoc	32.0			% Equivoc	50.1%		
False Pos	8	of 194 test +	3.93%	False Pos	15	of 147 test +	10.00%
False Neg	53	of 486 test -	10.83%	False Neg	38	of 35 test -	10.93%
* alpha2-macroglobulin, alpha2-globulin, total bilirubin, gama-globulin, apo A1 and GGT							

[0154] These results indicate that the  $\alpha$ 2-MG/HA/TIMP-1 three-marker panel can be useful in differentiating F0-F1 fibrosis from F2-F4 fibrosis with very good accuracy. These results further indicate that a combination fibrosis marker assay can be useful in determining the fibrosis status of a portion of the patients tested with very good accuracy, while the remaining patients are candidates for biopsy.

### EXAMPLE III

#### ASSAYS FOR $\alpha$ 2-MACROGLOBULIN, HYALURONIC ACID AND TISSUE INHIBITOR OF METALLOPROTEINASES-1

##### A. Quantitation of human $\alpha$ 2-macroglobulin ( $\alpha$ 2-MG)

[0155] Serum levels of human  $\alpha$ 2-macroglobulin were quantitated using the Beckman Array® 360 System as follows to determine  $\alpha$ 2-MG levels in the range of 0.75-270 mg/ml.

[0156] The Beckman Array® 360 system was used for determination of  $\alpha$ 2-MG concentrations. This system utilizes a nephelometer which measures the rate of light-scatter formation resulting from an immunoprecipitation reaction between  $\alpha$ 2-MG antigen in a sample with antibody to human  $\alpha$ 2-MG. After passing a beam of light through the solution in a flow cell, the intensity of light scattered by the formed macromolecular particles of insoluble complexes suspended in solution is detected and measured by the nephelometer. The increase in light scatter resulting from the antigen-antibody reaction is converted to a peak rate signal proportional to the  $\alpha$ 2-MG concentration in the sample. The resulting formation of complexes and the consequent change in the intensity of scattered light occurs at a rate that increases gradually at first, then rapidly, and finally proceeds through a peak rate of change for the component being analyzed.

[0157] Serum samples were drawn from fasting individuals and generally physically separated from cells within 2 hours from the time of collection as set forth in NCCLS publication H 18-A. Samples not assayed within 72 hours were stored frozen at -15°C to -20°C. Frozen samples were at most thawed one time. Grossly hemolyzed, highly lipemic or turbid specimens were rejected for further analysis.

[0158] Reagents were removed from storage at 4°C and used immediately. Buffers and Diluents were mixed thoroughly by inversion prior to being added to the instrument. Set-up, priming and calibration were performed according to the manufacturer's instructions with samples diluted 1:36. Relatively concentrated samples such as undiluted samples or 1:6 dilutions were generally avoided. Grossly lipemic sample were diluted 1:2 with diluent before assaying. Dust particles or other particulate matter, which can result in extraneous light-scattering signals, in the reaction solution were avoided. Prior to assaying samples, any air bubbles or foam in the sample cups and reagent bottle were removed by using a disposable transfer pipette or pipette tip to aspirate the bubbles. DTT was avoided in the work area.

[0159] Samples were analyzed for  $\alpha$ 2-MG concentration as follows. The Reagent Wheel (left wheel) on the instrument was loaded with AMG antiserum in space #2-. Dilution segments were loaded with 150  $\mu$ L of control or sample in the

wells on the larger side of the fan shaped segments. Segments and initial dilution control/sample cups were marked for identification. Bubbles were avoided while controls and serum samples were loaded.

**[0160]** Vigil™ Protein Control Levels 1 and 3 (3 drops) was placed in cups 1 and 3, respectively. Biorad Liquichek™ Immunology Control Level 2 (150 µL) was placed in cup 2. Patient samples (150 µL) were added to sequential cups. Segments were placed on right wheel beginning at position #1. Evaporation covers were placed over Reagent and Sample Wheels.

**[0161]** On the Master Screen menu, the RESULTS RECALL (F3) was selected before (F4) CLR CUR RUN. After returning to the MASTER SCREEN, the SAMPLE PROGRAM (F1) was selected. ENTER was selected when Reagent wheel #1 appeared and at each cup number. The control ID or sample Acc.# was entered. Test "2" was selected, and SAVE CUP (F1) was selected for each cup. START was selected to begin the analysis. At the end of the run, (Y) was selected in response to CLEAR CURRENT RUN & START NEXT RUN.

**[0162]** Results were reported by the Beckman Array® 360 in mg/dl using whole numbers in the Pros System. Samples were diluted routinely by the instrument 1:36. Samples greater than 750 mg/dl were assayed at a 1:216 dilution by the instrument. Samples having a concentration less than 75 mg/dl at a 1:36 dilution are reported as <75 mg/dl. At initial dilutions the Beckman analytical range was 75-750 mg/dl, while the extended range was 75-27,000 mg/dl. The range for normal individuals as verified at Prometheus Laboratories was 103-274 mg/dl.

**[0163]** Quality control was performed as follows. Three levels of controls were used: low, medium and high. Controls were within 2 standard deviations, except that runs were accepted with two controls within 2 standard deviations and the third control between 2 and 3 standard deviations. The controls used were Beckman Vigil I and III and Biorad Level II. Controls were assayed with each sample run.

**[0164]** The assay is calibrated every 14 days, and also when changes in reagent lots occur or when a major change has occurred in the instrument. Linearity is confirmed every 6 months with appropriate linearity material. This is done to ensure consistent performance over time and to comply with State and National standards.

**[0165]** Assay calibration verification is performed every 6 months to ensure consistency over time. A minimum of five verification samples including minimum, mid-point, and maximum concentrations are evaluated every 6 months. The coefficient of variation (%CV) of the verification sample results must be less than 15% in order to report out patient sample results.

#### B. Quantitation of Hyaluronic Acid (HA)

**[0166]** Serum levels of HA were determined using the Hyaluronic Acid (HA) Quantitative test kit (Catalog #029001) from Corgenix essentially as follows.

**[0167]** Serum samples were stored at -70°C. Multiple freeze/thaw cycles were avoided, with a maximum of 4 freeze/thaw cycles per sample. The kits were stored at 2-8°C.

**[0168]** Prior to use, the kit and patient samples were equilibrated to room temperature (18-28°C). The pouch of coated strips also was equilibrated to room temperature before opening. Wash solution (0.01 M PBS, pH 7.35 +/-0.1) was prepared by diluting the 33X PBS wash concentrate with distilled water and adjusting the pH of the final solution to pH 7.35 +/- 0.1.

**[0169]** All blanks, standards, controls and samples were assayed in duplicate. A water blank for calibration of the spectrophotometer was included with each plate and remained empty until addition of 200 µl water immediately prior to reading. Reaction buffer without serum sample was used for the reagent blank, which represented the 0 ng/ml HA reference solution, and was treated the same as patient samples and reference solutions in subsequent assay steps. Three known patient samples (low, middle and high) were run with each assay. In addition, 50 ng/ml HA, 100 ng/ml HA, 200 ng/ml HA, 500 ng/ml HA and 800 ng/ml HA reference solutions supplied with each kit were assayed as described further below.

**[0170]** HA reference solutions and patient samples were diluted 1:11 by addition of 25 µl reference solution or sample to 250 µl of reaction buffer and mixed by gentle vortexing. The diluted reference, samples and controls were added (100 µl) to each well. The water blank remained empty. The plate was covered and incubated for 60 minutes at room temperature. After the incubation was complete, the contents of the wells were removed by aspiration. Plates were washed four times with 1X wash solution while avoiding the plates drying out between washes. The plate was blotted vigorously on paper towels to remove residual buffer after the last wash.

**[0171]** HRP-conjugated HA binding protein solution (100 µl) was added to all wells except the water blank before covering the plate and incubating for 30 minutes at room temperature. After the incubation was complete, the plate was washed four times as described above. Substrate solution (100 µl 3,3',5,5'-tetramethylbenzidine and hydrogen peroxide, stabilized) was then added to each well except for the water blank. The covered plate was then incubated for 30 minutes at room temperature. The plate was kept in the dark.

**[0172]** The OD<sub>650</sub> of the 800 ng/ml HA standard was determined. For an OD less than 0.500, the substrate incubation was continued and the OD monitored to determine if the OD had reached 0.500. For an OD greater than 0.500 or after

one hour of substrate incubation even if the OD had not reached 0.500, the reactions were terminated by addition of 100  $\mu$ l of Stopping Solution (0.36 N sulfuric acid) to each well except the water blank. The stop solution was added in the same order and at approximately the same rate as addition of the substrate solution. Before reading the optical densities, 200  $\mu$ l distilled water was added to the water blank. The OD of each well was read at 450 nm (650 nm reference) within one hour after "zeroing" the plate reader against the water blank.

**[0173]** The following criteria were used to determine if the assay was reliable. The mean OD value of the reagent blank (zero standard) was less than 0.10. Readings greater than 0.10 were considered indicative of possible substrate or reagent contamination, and results were not reported under these conditions. The mean OD value of the 500 ng/ml HA reference was 0.800 or greater. Controls for the three known patient samples were within the following ranges: Low control: 78.6 to 117.2 ng/ml. Mid control: 148.5 to 214.1 ng/ml. High control: 297.8 to 460.7 ng/ml. Samples with HA concentrations greater than 800 ng/ml were further diluted and assayed a second time to obtain a more accurate result.

**[0174]** The known patient controls and samples were determined from a standard 4-parameter curve generated using Softmax and reported in ng/ml. The patient values were not reported if the concentration exceeded the concentration of the highest standard. For patient values greater than the concentration of the highest standard at a 1:11 dilution, samples were assayed at a 1:55 dilution and, if necessary, at higher dilution.

**[0175]** The HA ELISA assay is evaluated every six months to ensure consistent performance over time. A minimum of five samples with previously known HA values are evaluated in a blinded fashion to the operator. For the assay performance to be acceptable, results for negative samples must be negative, and results for positive samples must be positive and yield results within 15% of the previously obtained values. If greater than 20% of the validation samples fail the performance criteria, troubleshooting is implemented, and the assay is not used to report patient data until acceptable assay performance are reestablished.

#### C. Quantitation of Tissue Inhibitor of Metalloproteinases-1 (TIMP-1)

**[0176]** Serum levels of TIMP-1 were determined using the Biotrak™ test kit (Catalog# RPN2611) from Amersham Pharmacia Biotech (Piscataway, NJ) essentially as follows.

**[0177]** Kit contents were thawed and equilibrated to 20-25°C. Serum samples were stored frozen at -70°C. Repeated freeze-thaw cycles of the samples were minimized, with a maximum of six freeze-thaw cycles.

**[0178]** Assay reagents were prepared as follows and stored at 2-8°C for at most 7 days. Assay buffer 1 (0.1 M phosphate buffer, pH 7.5, with 0.9% (w/v) sodium chloride, 0.1% (w/v) BSA and 0.1% Tween-20) was prepared by adding distilled water to the assay buffer concentrate and adjusting the final volume to 100 ml.

**[0179]** Anti-TIMP-1 horseradish peroxidase conjugate was prepared in assay buffer 1 essentially as follows. To the stock bottle containing lyophilized conjugate, 11 ml diluted assay buffer 1 was added; the contents were mixed gently until completely dissolved while avoiding vigorous agitation and foaming. Wash buffer (0.1 M phosphate buffer, pH 7.5, containing 0.05% Tween-20) was prepared by adding distilled water to the wash buffer concentrate and bringing the final volume to 500 ml, followed by thorough mixing.

**[0180]** The 100 ng/ml TIMP-1 stock solution was prepared as follows and stored at 2-8°C. The lyophilized TIMP-1 standard was reconstituted in 0.1 M phosphate buffer, pH 7.5, containing 0.9% (w/v) sodium chloride, 0.1% (w/v) bovine serum albumin and 0.1% Tween-20 to make a standard TIMP-1 stock solution of 100 ng/ml. The contents were mixed gently until completely dissolved without vigorous agitation or foaming. Additional standards (1.565, 3.13, 6.25, 12.5, 25 and 50 ng/ml) for a standard curve were prepared fresh before each assay by two-fold serial dilution of the 100 ng/ml stock solution into assay buffer 1 in 1.2 ml dilution tubes. A zero standard (blank) was also prepared.

**[0181]** The pouch containing the microtiter plate was opened after equilibration to room temperature. All samples and standards were assayed in duplicate, and standards for a standard curve were present on each plate. On each plate, seven standards, two controls and a maximum of different 39 samples were present in duplicate.

**[0182]** Samples were diluted 1:120 in tubes by mixing 595  $\mu$ l assay buffer 1 with 5  $\mu$ l serum. The dilutions were mixed by vortexing. Using a multichannel pipettor, 100  $\mu$ l of blank, standards and diluted samples were added to individual wells on a microtiter plate. The plate was covered with the lid provided and incubated at room temperature for exactly two hours. Following the two hour incubation, the contents of the wells were aspirated, and each well was washed four times with wash buffer, with complete filling and aspiration of the wells after each wash. After the final wash, the plates were blotted on paper towels to remove residual wash buffer.

**[0183]** Peroxidase conjugate (100  $\mu$ l) was added to each well using a multichannel pipettor, and the covered plate incubated at room temperature for exactly two hours. After the incubation, the wells were aspirated and washed as before. Immediately upon conclusion of the incubation, 100  $\mu$ l of room temperature equilibrated TMB substrate (3,3', 5,5'-tetramethylbenzidine/ hydrogen peroxide in 20% (v/v) dimethylformamide) was added to each well. The plates were covered and incubated for exactly 30 minutes at room temperature. In some cases, the reactions were monitored at 630 nm. The reactions were stopped by addition of 100  $\mu$ l 1 M sulfuric acid to all wells. Absorbance was determined at 450 nm within 30 minutes.

**[0184]** Control and patient samples values were determined using a standard curve (4-parameter curve fit) generated using Softmax. Concentration values from the standard curve were multiplied by the dilution factor (120) to obtain actual concentrations, reported in ng/ml. Quality of the assay was confirmed using known serum samples. The low control was in the range of 668.1 to 979.9 ng/ml. The high control was in the range of 2677.9 to 3300.2 ng/ml. Patient values generally did not exceed the concentration in ng/ml of the highest standard. Where the patient value was greater than the concentration of the highest standard at a 1:120 dilution, the result was reported as greater than 120 times the concentration of the highest standard.

**[0185]** The TIMP-1 ELISA assay is validated every six months to ensure consistent performance over time. A minimum of five samples with previously known values are evaluated in a blinded fashion to the operator. Results for negative samples must be negative. Results for positive samples must be positive and must yield results within 15% of the previously obtained values. Where greater than 20% of the validation samples fail the performance criteria, troubleshooting is implemented. Further patient data are not reported until acceptable assay performance is reestablished.

#### SEQUENCE LISTING

##### **[0186]**

<110> Prometheus Laboratories, Inc.  
 Rose, Steven L.  
 Oh, Esther H.  
 Walsh, Michael J.

<120> Methods of Diagnosing Liver Fibrosis

<130> FP-PM 5505

<150> US 10/087,188

<151> 2002-02-28

<160> 4

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 2041

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)...(1932)

<400> 1

EP 1 487 395 B1

	ccc gcc ttc cta gct gtc cca gtg gag aag gaa caa gcg cct cac tgc	48
	Pro Ala Phe Leu Ala Val Pro Val Glu Lys Glu Gln Ala Pro His Cys	
	1 5 10 15	
5	atc tgt gca aac ggg cgg caa act gtg tcc tgg gca gta acc cca aag	96
	Ile Cys Ala Asn Gly Arg Gln Thr Val Ser Trp Ala Val Thr Pro Lys	
	20 25 30	
10	tca tta gga aat gtg aat ttc act gtg agc gca gag gca cta gag tct	144
	Ser Leu Gly Asn Val Asn Phe Thr Val Ser Ala Glu Ala Leu Glu Ser	
	35 40 45	
15	caa gag ctg tgt ggg act gag gtg cct tca gtt cct gaa cac gga agg	192
	Gln Glu Leu Cys Gly Thr Glu Val Pro Ser Val Pro Glu His Gly Arg	
	50 55 60	
20	aaa gac aca gtc atc aag cct ctg ttg gtt gaa cct gaa gga cta gag	240
	Lys Asp Thr Val Ile Lys Pro Leu Leu Val Glu Pro Glu Gly Leu Glu	
	65 70 75 80	
25	aag gaa aca aca ttc aac tcc cta ctt tgt cca tca ggt ggt gag gtt	288
	Lys Glu Thr Thr Phe Asn Ser Leu Leu Cys Pro Ser Gly Gly Glu Val	
	85 90 95	
30	tct gaa gaa tta tcc ctg aaa ctg cca cca aat gtg gta gaa gaa tct	336
35		
40		
45		
50		
55		

## EP 1 487 395 B1

	Ser	Glu	Glu	Leu	Ser	Leu	Lys	Leu	Pro	Pro	Asn	Val	Val	Glu	Glu	Ser	
				100					105								
5	gcc	cga	gct	tct	gtc	tca	gtt	ttg	gga	gac	ata	tta	ggc	tct	gcc	atg	384
	Ala	Arg	Ala	Ser	Val	Ser	Val	Leu	Gly	Asp	Ile	Leu	Gly	Ser	Ala	Met	
				115				120					125				
10	caa	aac	aca	caa	aat	ctt	ctc	cag	atg	ccc	tat	ggc	tgt	gga	gag	cag	432
	Gln	Asn	Thr	Gln	Asn	Leu	Leu	Gln	Met	Pro	Tyr	Gly	Cys	Gly	Glu	Gln	
				130				135				140					
15	aat	atg	gtc	ctc	ttt	gct	cct	aac	atc	tat	gta	ctg	gat	tat	cta	aat	480
	Asn	Met	Val	Leu	Phe	Ala	Pro	Asn	Ile	Tyr	Val	Leu	Asp	Tyr	Leu	Asn	
	145					150				155					160		
20	gaa	aca	cag	cag	ctt	act	cca	gag	atc	aag	tcc	aag	gcc	att	ggc	tat	528
	Glu	Thr	Gln	Gln	Leu	Thr	Pro	Glu	Ile	Lys	Ser	Lys	Ala	Ile	Gly	Tyr	
					165					170					175		
25	ctc	aac	act	ggt	tac	cag	aga	cag	ttg	aac	tac	aaa	cac	tat	gat	ggc	576
	Leu	Asn	Thr	Gly	Tyr	Gln	Arg	Gln	Leu	Asn	Tyr	Lys	His	Tyr	Asp	Gly	
				180					185					190			
30	tcc	tac	agc	acc	ttt	ggg	gag	cga	tat	ggc	agg	aac	cag	ggc	aac	acc	624
	Ser	Tyr	Ser	Thr	Phe	Gly	Glu	Arg	Tyr	Gly	Arg	Asn	Gln	Gly	Asn	Thr	
				195				200					205				
35	tgg	ctc	aca	gcc	ttt	gtt	ctg	aag	act	ttt	gcc	caa	gct	cga	gcc	tac	672
	Trp	Leu	Thr	Ala	Phe	Val	Leu	Lys	Thr	Phe	Ala	Gln	Ala	Arg	Ala	Tyr	
		210					215					220					
40	atc	ttc	atc	gat	gaa	gca	cac	att	acc	caa	gcc	ctc	ata	tgg	ctc	tcc	720
	Ile	Phe	Ile	Asp	Glu	Ala	His	Ile	Thr	Gln	Ala	Leu	Ile	Trp	Leu	Ser	
	225					230					235					240	
45	cag	agg	cag	aag	gac	aat	ggc	tgt	ttc	agg	agc	tct	ggg	tca	ctg	ctc	768
	Gln	Arg	Gln	Lys	Asp	Asn	Gly	Cys	Phe	Arg	Ser	Ser	Gly	Ser	Leu	Leu	
					245				250						255		
50	aac	aat	gcc	ata	aag	gga	gga	gta	gaa	gat	gaa	gtg	acc	ctc	tcc	gcc	816
	Asn	Asn	Ala	Ile	Lys	Gly	Gly	Val	Glu	Asp	Glu	Val	Thr	Leu	Ser	Ala	
				260				265						270			
55	tat	atc	acc	atc	gcc	ctt	ctg	gag	att	cct	ctc	aca	gtc	act	cac	cct	864
	Tyr	Ile	Thr	Ile	Ala	Leu	Leu	Glu	Ile	Pro	Leu	Thr	Val	Thr	His	Pro	
				275				280					285				
60	gtt	gtc	cgc	aat	gcc	ctg	ttt	tgc	ctg	gag	tca	gcc	tgg	aag	aca	gca	912
	Val	Val	Arg	Asn	Ala	Leu	Phe	Cys	Leu	Glu	Ser	Ala	Trp	Lys	Thr	Ala	
				290			295					300					
65	caa	gaa	ggg	gac	cat	ggc	agc	cat	gta	tat	acc	aaa	gac	ctg	ctg	gcc	960
	Gln	Glu	Gly	Asp	His	Gly	Ser	His	Val	Tyr	Thr	Lys	Asp	Leu	Leu	Ala	
	305					310				315						320	



EP 1 487 395 B1

	tat gct ttt gcc ctg gca ggt aac cag gac aag agg aag gaa gta ctc	1008
	Tyr Ala Phe Ala Leu Ala Gly Asn Gln Asp Lys Arg Lys Glu Val Leu	
	325 330 335	
5	aag tca ctt aat gag gaa gct gtg aag aaa gac aac tct gtc cat tgg	1056
	Lys Ser Leu Asn Glu Glu Ala Val Lys Lys Asp Asn Ser Val His Trp	
	340 345 350	
10	gag cgc cct cag aaa ccc aag gca cca gtg ggg gat ttt tac gaa ccc	1104
	Glu Arg Pro Gln Lys Pro Lys Ala Pro Val Gly Asp Phe Tyr Glu Pro	
	355 360 365	
15	cag gct ccc tct gct gag gtg gag atg aca tcc tat gtg ctc ctc gct	1152
	Gln Ala Pro Ser Ala Glu Val Glu Met Thr Ser Tyr Val Leu Leu Ala	
	370 375 380	
20	tat ctc acg gcc cag cca gcc cca acc tcg gag gac ctg acc tct gca	1200
	Tyr Leu Thr Ala Gln Pro Ala Pro Thr Ser Glu Asp Leu Thr Ser Ala	
	385 390 395 400	
25	acc aac atc gtg aag tgg atc acg aag cag cag aat gcc cag ggc ggt	1248
	Thr Asn Ile Val Lys Trp Ile Thr Lys Gln Gln Asn Ala Gln Gly Gly	
	405 410 415	
30	ttc tcc tcc acc cag gac aca gtg gtg gct ctc cat gct ctg tcc aaa	1296
	Phe Ser Ser Thr Gln Asp Thr Val Val Ala Leu His Ala Leu Ser Lys	
	420 425 430	
35	tat gga gca gcc aca ttt acc agg act ggg aag gct gca cag gtg act	1344
	Tyr Gly Ala Ala Thr Phe Thr Arg Thr Gly Lys Ala Ala Gln Val Thr	
	435 440 445	
40	atc cag tct tca ggg aca ttt tcc agc aaa ttc caa gtg gac aac aac	1392
	Ile Gln Ser Ser Gly Thr Phe Ser Ser Lys Phe Gln Val Asp Asn Asn	
	450 455 460	
45	aac cgc ctg tta ctg cag cag gtc tca ttg cca gag ctg cct ggg gaa	1440
	Asn Arg Leu Leu Leu Gln Gln Val Ser Leu Pro Glu Leu Pro Gly Glu	
	465 470 475 480	
50	tac agc atg aaa gtg aca gga gaa gga tgt gtc tac ctc cag aca tcc	1488
	Tyr Ser Met Lys Val Thr Gly Glu Gly Cys Val Tyr Leu Gln Thr Ser	
	485 490 495	
55	ttg aaa tac aat att ctc cca gaa aag gaa gag ttc ccc ttt gct tta	1536
	Leu Lys Tyr Asn Ile Leu Pro Glu Lys Glu Glu Phe Pro Phe Ala Leu	
	500 505 510	
60	gga gtg cag act ctg cct caa act tgt gat gaa ccc aaa gcc cac acc	1584
	Gly Val Gln Thr Leu Pro Gln Thr Cys Asp Glu Pro Lys Ala His Thr	
	515 520 525	
65	agc ttc caa atc tcc cta agt gtc agt tac aca ggg agc cgc tct gcc	1632
	Ser Phe Gln Ile Ser Leu Ser Val Ser Tyr Thr Gly Ser Arg Ser Ala	
	530 535 540	

EP 1 487 395 B1

tcc aac atg gcg atc gtt gat gtg aag atg gtc tct ggc ttc att ccc 1680  
 Ser Asn Met Ala Ile Val Asp Val Lys Met Val Ser Gly Phe Ile Pro  
 545 550 555 560  
 5  
 ctg aag cca aca gtg aaa atg ctt gaa aga tct aac cat gtg agc cgg 1728  
 Leu Lys Pro Thr Val Lys Met Leu Glu Arg Ser Asn His Val Ser Arg  
 565 570 575  
 10  
 aca gaa gtc agc agc aac cat gtc ttg att tac ctt gat aag gtg tca 1776  
 Thr Glu Val Ser Ser Asn His Val Leu Ile Tyr Leu Asp Lys Val Ser  
 580 585 590  
 15  
 aat cag aca ctg agc ttg ttc ttc acg gtt ctg caa gat gtc cca gta 1824  
 Asn Gln Thr Leu Ser Leu Phe Phe Thr Val Leu Gln Asp Val Pro Val  
 595 600 605  
 20  
 aga gat ctg aaa cca gcc ata gtg aaa gtc tat gat tac tac gag acg 1872  
 Arg Asp Leu Lys Pro Ala Ile Val Lys Val Tyr Asp Tyr Tyr Glu Thr  
 610 615 620  
 gat gag ttt gca att gct gag tac aat gct cct tgc agc aaa gat ctt 1920  
 Asp Glu Phe Ala Ile Ala Glu Tyr Asn Ala Pro Cys Ser Lys Asp Leu  
 625 630 635 640  
 25  
 gga aat gct tga agaccacaag gctgaaaagt gctttgctgg agtctctgttc 1972  
 Gly Asn Ala \*  
 30  
 tcagagctcc acagaagaca cgtgttttttg tatcttttaaa gacttgatga ataaacactt 2032  
 tttctggtc 2041  
 <210> 2  
 <211> 643  
 35  
 <212> PRT  
 <213> Homo sapiens  
 <400> 2  
 40  
 Pro Ala Phe Leu Ala Val Pro Val Glu Lys Glu Gln Ala Pro His Cys  
 1 5 10 15  
 Ile Cys Ala Asn Gly Arg Gln Thr Val Ser Trp Ala Val Thr Pro Lys  
 20 25 30  
 45  
 Ser Leu Gly Asn Val Asn Phe Thr Val Ser Ala Glu Ala Leu Glu Ser  
 35 40 45  
 Gln Glu Leu Cys Gly Thr Glu Val Pro Ser Val Pro Glu His Gly Arg  
 50 55 60  
 Lys Asp Thr Val Ile Lys Pro Leu Leu Val Glu Pro Glu Gly Leu Glu  
 65 70 75 80  
 50  
 Lys Glu Thr Thr Phe Asn Ser Leu Leu Cys Pro Ser Gly Gly Glu Val  
 85 90 95  
 Ser Glu Glu Leu Ser Leu Lys Leu Pro Pro Asn Val Val Glu Glu Ser  
 100 105 110  
 55  
 Ala Arg Ala Ser Val Ser Val Leu Gly Asp Ile Leu Gly Ser Ala Met  
 115 120 125  
 Gln Asn Thr Gln Asn Leu Leu Gln Met Pro Tyr Gly Cys Gly Glu Gln

## EP 1 487 395 B1

		130				135				140									
		Asn	Met	Val	Leu	Phe	Ala	Pro	Asn	Ile	Tyr	Val	Leu	Asp	Tyr	Leu	Asn		
		145					150					155					160		
5		Glu	Thr	Gln	Gln	Leu	Thr	Pro	Glu	Ile	Lys	Ser	Lys	Ala	Ile	Gly	Tyr		
					165						170					175			
		Leu	Asn	Thr	Gly	Tyr	Gln	Arg	Gln	Leu	Asn	Tyr	Lys	His	Tyr	Asp	Gly		
					180					185					190				
		Ser	Tyr	Ser	Thr	Phe	Gly	Glu	Arg	Tyr	Gly	Arg	Asn	Gln	Gly	Asn	Thr		
10					195				200					205					
		Trp	Leu	Thr	Ala	Phe	Val	Leu	Lys	Thr	Phe	Ala	Gln	Ala	Arg	Ala	Tyr		
		210						215					220						
		Ile	Phe	Ile	Asp	Glu	Ala	His	Ile	Thr	Gln	Ala	Leu	Ile	Trp	Leu	Ser		
		225				230						235					240		
15		Gln	Arg	Gln	Lys	Asp	Asn	Gly	Cys	Phe	Arg	Ser	Ser	Gly	Ser	Leu	Leu		
					245						250					255			
		Asn	Asn	Ala	Ile	Lys	Gly	Gly	Val	Glu	Asp	Glu	Val	Thr	Leu	Ser	Ala		
					260					265					270				
		Tyr	Ile	Thr	Ile	Ala	Leu	Leu	Glu	Ile	Pro	Leu	Thr	Val	Thr	His	Pro		
20					275				280					285					
		Val	Val	Arg	Asn	Ala	Leu	Phe	Cys	Leu	Glu	Ser	Ala	Trp	Lys	Thr	Ala		
					290				295				300						
		Gln	Glu	Gly	Asp	His	Gly	Ser	His	Val	Tyr	Thr	Lys	Asp	Leu	Leu	Ala		
		305				310						315					320		
25		Tyr	Ala	Phe	Ala	Leu	Ala	Gly	Asn	Gln	Asp	Lys	Arg	Lys	Glu	Val	Leu		
					325						330					335			
		Lys	Ser	Leu	Asn	Glu	Glu	Ala	Val	Lys	Lys	Asp	Asn	Ser	Val	His	Trp		
					340					345					350				
		Glu	Arg	Pro	Gln	Lys	Pro	Lys	Ala	Pro	Val	Gly	Asp	Phe	Tyr	Glu	Pro		
					355				360					365					
30		Gln	Ala	Pro	Ser	Ala	Glu	Val	Glu	Met	Thr	Ser	Tyr	Val	Leu	Leu	Ala		
					370				375				380						
		Tyr	Leu	Thr	Ala	Gln	Pro	Ala	Pro	Thr	Ser	Glu	Asp	Leu	Thr	Ser	Ala		
		385				390						395					400		
35		Thr	Asn	Ile	Val	Lys	Trp	Ile	Thr	Lys	Gln	Gln	Asn	Ala	Gln	Gly	Gly		
					405						410					415			
		Phe	Ser	Ser	Thr	Gln	Asp	Thr	Val	Val	Ala	Leu	His	Ala	Leu	Ser	Lys		
					420					425					430				
		Tyr	Gly	Ala	Ala	Thr	Phe	Thr	Arg	Thr	Gly	Lys	Ala	Ala	Gln	Val	Thr		
					435				440					445					
40		Ile	Gln	Ser	Ser	Gly	Thr	Phe	Ser	Ser	Lys	Phe	Gln	Val	Asp	Asn	Asn		
					450				455				460						
		Asn	Arg	Leu	Leu	Leu	Gln	Gln	Val	Ser	Leu	Pro	Glu	Leu	Pro	Gly	Glu		
		465				470						475					480		
		Tyr	Ser	Met	Lys	Val	Thr	Gly	Glu	Gly	Cys	Val	Tyr	Leu	Gln	Thr	Ser		
45					485						490					495			
		Leu	Lys	Tyr	Asn	Ile	Leu	Pro	Glu	Lys	Glu	Glu	Phe	Pro	Phe	Ala	Leu		
					500					505					510				
		Gly	Val	Gln	Thr	Leu	Pro	Gln	Thr	Cys	Asp	Glu	Pro	Lys	Ala	His	Thr		
					515				520					525					
50		Ser	Phe	Gln	Ile	Ser	Leu	Ser	Val	Ser	Tyr	Thr	Gly	Ser	Arg	Ser	Ala		
					530				535				540						
		Ser	Asn	Met	Ala	Ile	Val	Asp	Val	Lys	Met	Val	Ser	Gly	Phe	Ile	Pro		
		545				550						555					560		
		Leu	Lys	Pro	Thr	Val	Lys	Met	Leu	Glu	Arg	Ser	Asn	His	Val	Ser	Arg		
55					565						570					575			

Thr Glu Val Ser Ser Asn His Val Leu Ile Tyr Leu Asp Lys Val Ser  
 580 585 590  
 Asn Gln Thr Leu Ser Leu Phe Phe Thr Val Leu Gln Asp Val Pro Val  
 595 600 605  
 Arg Asp Leu Lys Pro Ala Ile Val Lys Val Tyr Asp Tyr Tyr Glu Thr  
 610 615 620  
 Asp Glu Phe Ala Ile Ala Glu Tyr Asn Ala Pro Cys Ser Lys Asp Leu  
 625 630 635 640  
 Gly Asn Ala

<210> 3

<211> 782

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (63)... (683)

<400> 3

aggggcctta gcgtgccgca tcgccgagat ccagcgccca gagagacacc agagaaccca 60  
 cc atg gcc ccc ttt gag ccc ctg gct tct ggc atc ctg ttg ttg ctg 107  
 Met Ala Pro Phe Glu Pro Leu Ala Ser Gly Ile Leu Leu Leu Leu  
 1 5 10 15  
 tgg ctg ata gcc ccc agc agg gcc tgc acc tgt gtc cca ccc cac cca 155  
 Trp Leu Ile Ala Pro Ser Arg Ala Cys Thr Cys Val Pro Pro His Pro  
 20 25 30  
 cag acg gcc ttc tgc aat tcc gac ctc gtc atc agg gcc aag ttc gtg 203  
 Gln Thr Ala Phe Cys Asn Ser Asp Leu Val Ile Arg Ala Lys Phe Val  
 35 40 45  
 ggg aca cca gaa gtc aac cag acc acc tta tac cag cgt tat gag atc 251  
 Gly Thr Pro Glu Val Asn Gln Thr Thr Leu Tyr Gln Arg Tyr Glu Ile  
 50 55 60  
 aag atg acc aag atg tat aaa ggg ttc caa gcc tta ggg gat gcc gct 299  
 Lys Met Thr Lys Met Tyr Lys Gly Phe Gln Ala Leu Gly Asp Ala Ala  
 65 70 75  
 gac atc cgg ttc gtc tac acc ccc gcc atg gag agt gtc tgc gga tac 347  
 Asp Ile Arg Phe Val Tyr Thr Pro Ala Met Glu Ser Val Cys Gly Tyr  
 80 85 90 95  
 ttc cac agg tcc cac aac cgc agc gag gag ttt ctc att gct gga aaa 395  
 Phe His Arg Ser His Asn Arg Ser Glu Glu Phe Leu Ile Ala Gly Lys  
 100 105 110  
 ctg cag gat gga ctc ttg cac atc act acc tgc agt ttc gtg gct ccc 443  
 Leu Gln Asp Gly Leu Leu His Ile Thr Thr Cys Ser Phe Val Ala Pro  
 115 120 125

tgg aac agc ctg agc tta gct cag cgc cgg ggc ttc acc aag acc tac 491  
 Trp Asn Ser Leu Ser Leu Ala Gln Arg Arg Gly Phe Thr Lys Thr Tyr  
 130 135 140

5 act gtt ggc tgt gag gaa tgc aca gtg ttt ccc tgt tta tcc atc ccc 539  
 Thr Val Gly Cys Glu Glu Cys Thr Val Phe Pro Cys Leu Ser Ile Pro  
 145 150 155

10 tgc aaa ctg cag agt ggc act cat tgc ttg tgg acg gac cag ctc ctc 587  
 Cys Lys Leu Gln Ser Gly Thr His Cys Leu Trp Thr Asp Gln Leu Leu  
 160 165 170 175

15 caa ggc tct gaa aag ggc ttc cag tcc cgt cac ctt gcc tgc ctg cct 635  
 Gln Gly Ser Glu Lys Gly Phe Gln Ser Arg His Leu Ala Cys Leu Pro  
 180 185 190

20 cgg gag cca ggg ctg tgc acc tgg cag tcc ctg cgg tcc cag ata gcc 683  
 Arg Glu Pro Gly Leu Cys Thr Trp Gln Ser Leu Arg Ser Gln Ile Ala  
 195 200 205

tgaatcctgc ccggagtgga actgaagcct gcacagtgtc caccctgttc ccactcccat 743  
 ctttcttcg gacaatgaaa taaagagtta ccaccagc 782

25 <210> 4  
 <211> 207  
 <212> PRT  
 <213> Homo sapiens

30 <400> 4

Met Ala Pro Phe Glu Pro Leu Ala Ser Gly Ile Leu Leu Leu Leu Trp  
 1 5 10 15  
 Leu Ile Ala Pro Ser Arg Ala Cys Thr Cys Val Pro Pro His Pro Gln  
 20 25 30  
 Thr Ala Phe Cys Asn Ser Asp Leu Val Ile Arg Ala Lys Phe Val Gly  
 35 40 45  
 Thr Pro Glu Val Asn Gln Thr Thr Leu Tyr Gln Arg Tyr Glu Ile Lys  
 50 55 60  
 40 Met Thr Lys Met Tyr Lys Gly Phe Gln Ala Leu Gly Asp Ala Ala Asp  
 65 70 75 80  
 Ile Arg Phe Val Tyr Thr Pro Ala Met Glu Ser Val Cys Gly Tyr Phe  
 85 90 95  
 His Arg Ser His Asn Arg Ser Glu Glu Phe Leu Ile Ala Gly Lys Leu  
 100 105 110  
 45 Gln Asp Gly Leu Leu His Ile Thr Thr Cys Ser Phe Val Ala Pro Trp  
 115 120 125  
 Asn Ser Leu Ser Leu Ala Gln Arg Arg Gly Phe Thr Lys Thr Tyr Thr  
 130 135 140  
 50 Val Gly Cys Glu Glu Cys Thr Val Phe Pro Cys Leu Ser Ile Pro Cys  
 145 150 155 160  
 Lys Leu Gln Ser Gly Thr His Cys Leu Trp Thr Asp Gln Leu Leu Gln  
 165 170 175  
 Gly Ser Glu Lys Gly Phe Gln Ser Arg His Leu Ala Cys Leu Pro Arg  
 180 185 190  
 55 Glu Pro Gly Leu Cys Thr Trp Gln Ser Leu Arg Ser Gln Ile Ala  
 195 200 205

## Claims

1. A method of diagnosing the presence or severity of liver fibrosis in an individual, comprising the steps of:

- (a) detecting  $\alpha$ 2-macroglobulin in a sample from said individual;
- (b) detecting hyaluronic acid (HA) in a sample from said individual;
- (c) detecting tissue inhibitor of metalloproteinases-1 (TIMP-1) in a sample from said individual; and
- (d) diagnosing the presence or severity of liver fibrosis in said individual based on the presence or level of  $\alpha$ 2-MG, HA and TIMP-1.

2. The method of claim 1, comprising detecting at most three markers of fibrosis.

3. The method of claim 1, further comprising detecting in a sample from said individual at least one marker selected from the group consisting of: PIIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2, and apoB.

4. The method of claim 3, wherein said marker is YKL-40.

5. The method of claim 1, further comprising detecting in a sample from said individual two or more markers selected from the group consisting of PIIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 and apoB.

6. The method of claim 1, wherein said individual has viral hepatitis.

7. The method of claim 6, wherein said individual is infected with hepatitis C virus.

8. The method of claim 6, wherein said individual is infected with hepatitis B virus.

9. The method of claim 1, wherein said individual has autoimmune liver disease.

10. The method of claim 1, wherein said individual has alcoholic liver disease.

11. The method of claim 1, wherein said individual has a fatty liver disease.

12. The method of claim 1, wherein said individual has drug-induced liver disease.

13. The method of claim 1, wherein step (a) comprises determining the level of  $\alpha$ 2-MG protein in said sample.

14. The method of claim 13, wherein the level of  $\alpha$ 2-MG protein is determined using one or more  $\alpha$ 2-MG-specific binding agents.

15. The method of claim 14, wherein the level of  $\alpha$ 2-MG protein is determined using one or more anti- $\alpha$ 2-MG antibodies.

16. The method of claim 1, wherein step (a) comprises determining a level of  $\alpha$ 2-MG activity.

17. The method of claim 1, wherein step (b) comprises determining the level of HA in said sample.

18. The method of claim 17, wherein the level of HA is determined using one or more HA-specific binding agents.

19. The method of claim 18, wherein the level of HA is determined using one or more HA-binding proteins.

20. The method of claim 18, wherein the level of HA is determined using one or more anti-HA antibodies.

21. The method of claim 1, wherein step (c) comprises determining the level of TIMP-1 protein in said sample.

22. The method of claim 21, wherein the level of TIMP-1 protein is determined using one or more TIMP-1-specific binding agents.

23. The method of claim 22, wherein the level of TIMP-1 protein is determined using one or more anti-TIMP-1 antibodies.

24. The method of claim 1, wherein step (c) comprises determining a level of TIMP-1 activity.

25. The method of claim 1,  
wherein step (a) comprises determining the level of  $\alpha$ 2-MG protein,  
wherein step (b) comprises determining the level of HA, and  
wherein step (c) comprises determining the level of TIMP-1 protein.

26. The method of claim 25, wherein the level of  $\alpha$ 2-MG protein, HA and TIMP-1 protein each is determined using an enzyme-linked assay.

27. The method of claim 1, wherein a single sample is obtained from said individual.

28. The method of claim 27, wherein said sample is selected from the group consisting of blood, serum, plasma, urine, saliva and liver tissue.

29. The method of claim 28, wherein said sample is a serum sample.

30. The method of claim 1, comprising differentiating no or mild liver fibrosis from moderate to severe liver fibrosis.

31. A method of differentiating no or mild liver fibrosis from moderate to severe liver fibrosis in an individual, comprising the steps of:

- (a) contacting an appropriate dilution of a sample from said individual with anti- $\alpha$ 2-MG antibody under conditions suitable to form a first complex of  $\alpha$ 2-MG and anti- $\alpha$ 2-MG antibody;
- (b) washing said first complex to remove unbound molecules;
- (c) determining the amount of  $\alpha$ 2-MG-containing first complex;
- (d) contacting an appropriate dilution of a sample from said individual with a HA-binding protein (HABP) under conditions suitable to form a second complex of HA and HABP;
- (e) washing said second complex to remove unbound molecules;
- (f) determining the amount of HA-containing second complex;
- (g) contacting an appropriate dilution of a sample from said individual with anti-TIMP-1 antibody under conditions suitable to form a third complex of TIMP-1 and anti-TIMP-1 antibody;
- (h) washing said third complex to remove unbound molecules;
- (i) determining the amount of TIMP-1-containing third complex; and
- (j) differentiating no/mild liver fibrosis from moderate/severe liver fibrosis in said individual based on the amounts of  $\alpha$ 2-MG, HA and TIMP-1-containing complexes.

32. A method of monitoring the efficacy of anti-fibrotic therapy in a patient, comprising the steps of:

- (a) detecting  $\alpha$ 2-macroglobulin in a sample from a patient administered an anti-fibrotic therapy;
- (b) detecting hyaluronic acid (HA) in a sample from said patient;
- (c) detecting tissue inhibitor of metalloproteinases-1 (TIMP-1) in a sample from said patient; and
- (d) determining the presence or severity of liver fibrosis in said patient based on the presence or level of  $\alpha$ 2-MG, HA and TIMP-1, thereby monitoring the efficacy of anti-fibrotic therapy.

33. The method of claim 32, further comprising comparing the presence or severity of liver fibrosis determined in step (d) to the presence or severity of liver fibrosis in said patient at an earlier time.

34. The method of claim 32, comprising detecting at most three markers of fibrosis.

35. The method of claim 32, further comprising detecting in a sample from said patient at least one marker selected from the group consisting of: PIINP, laminin, tenascin, collagen type IV, collagen type VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 complex, sFas ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2, and apoB.

36. The method of claim 32, wherein step (a) comprises determining the level of  $\alpha$ 2-MG protein in said sample.

37. The method of claim 36, wherein the level of  $\alpha$ 2-MG protein is determined using one or more anti- $\alpha$ 2-MG antibodies.
38. The method of claim 32, wherein step (b) comprises determining the level of HA in said sample..
- 5 39. The method of claim 38, wherein the level of HA is determined using one or more HA-binding proteins.
40. The method of claim 32, wherein step (c) comprises determining the level of TIMP-1 protein in said sample.
- 10 41. The method of claim 40, wherein the level of TIMP-1 protein is determined using one or more anti-TIMP-1 antibodies.
42. A method of differentiating no/mild liver fibrosis from moderate/severe liver fibrosis in an individual, comprising the steps of:
  - (a) determining an  $\alpha$ 2-MG level in a sample from said individual;
  - 15 (b) determining a HA level in a sample from said individual;
  - (c) determining a TIMP-1 level in a sample from said individual; and
  - (d) diagnosing said individual as having no/mild liver fibrosis when said  $\alpha$ 2-MG level is below an  $\alpha$ 2-MG cut-off value X1, said HA level is below a HA cut-off value Y1 or said TIMP-1 level is below a TIMP-1 cut-off value Z1,
  - 20 diagnosing said individual as having moderate/severe liver fibrosis when said  $\alpha$ 2-MG level is above an  $\alpha$ 2-MG cut-off value X2, said HA level is above a HA cut-off value Y2 and said TIMP-1 level is above a TIMP-1 cut-off value Z2, and diagnosing remaining individuals as having an indeterminate status.
43. The method of claim 42, wherein said individual has a disorder selected from the group consisting of viral hepatitis, autoimmune liver disease, alcoholic liver disease, fatty liver disease and drug-induced liver disease.
44. The method of claim 43, wherein said individual is infected with hepatitis C virus.
45. The method of claim 42, wherein said samples are independently selected from the group consisting of blood, serum, plasma, urine, saliva and liver tissue.
- 30 46. The method of claim 45, wherein said  $\alpha$ 2-MG, level, HA level and TIMP-1 level each is determined in a serum sample.
47. The method of claim 46,
  - 35 wherein X1 is a value between 1.8 and 2.2 mg/ml;
  - wherein Y1 is a value between 31 and 39 ng/ml;
  - wherein Z1 is a value between 900 and 1100 ng/ml;
  - wherein X2 is a value between 1.8 and 2.2 mg/ml;
  - wherein Y2 is a value between 54 and 66 ng/ml; and
  - 40 wherein Z2 is a value between 1415 and 1735 ng/ml.
48. The method of claim 47,
  - wherein X1 = 2.0 mg/ml;
  - wherein Y1 = 35 ng/ml;
  - 45 wherein Z1 = 1000 ng/ml;
  - wherein X2 = 2.0 mg/ml;
  - wherein Y2 = 60 ng/ml; and
  - wherein Z2 = 1575 ng/ml.
- 50 49. The method of claim 47,
  - wherein X1 = 2.0 mg/ml;
  - wherein Y1 = 37 ng/ml;
  - wherein Z1 = 1100 ng/ml;
  - wherein X2 = 2.0 mg/ml;
  - 55 wherein Y2 = 60 ng/ml; and
  - wherein Z2 = 1575 ng/ml.
50. The method of claim 42, wherein, in a population having up to 30% liver fibrosis prevalence, at least 65% of individuals



in said population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with an accuracy of at least 80%.

51. The method of claim 42, wherein, in a population having up to 30% liver fibrosis prevalence, at least 65% of individuals in said population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with an accuracy of at least 90%.

52. The method of claim 42, wherein, in a population having up to 30% liver fibrosis prevalence, at least 65% of individuals in said population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with a positive predictive value of at least 90% and a negative predictive value of at least 90%.

53. The method of claim 42, wherein, in a population having up to 10% liver fibrosis prevalence, at least 70% of individuals in said population are diagnosed as having no/mild fibrosis or moderate/severe fibrosis with an accuracy of at least 90%.

## Patentansprüche

1. Verfahren, um das Vorhandensein oder den Schweregrad von Leberfibrose in einem Subjekt zu diagnostizieren, Schritte umfassend, bei denen man:

- (a)  $\alpha$ 2-Makroglobulin in einer Probe von dem Subjekt nachweist;
- (b) Hyaluronsäure (HA) in einer Probe von dem Subjekt nachweist;
- (c) Gewebeinhibitor von Metalloproteinasen-1 (TIMP-1) in einer Probe des Subjekts nachweist; und
- (d) das Vorhandensein oder den Schweregrad von Leberfibrose in dem Subjekt basierend auf dem Vorhandensein oder der Menge von  $\alpha$ 2-MG, HA und TIMP-1 diagnostiziert.

2. Verfahren nach Anspruch 1, einen Schritt umfassend, bei dem man höchstens drei Fibrosemarker nachweist.

3. Verfahren nach Anspruch 1, einen weiteren Schritt umfassend, bei dem man in einer Probe von dem Subjekt mindestens einen Marker nachweist, der ausgewählt ist aus der Gruppe bestehend aus: PIINP, Laminin, Tenascin, Kollagen Typ IV, Kollagen Typ VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 Komplex, sFas Ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 und apoB.

4. Verfahren nach Anspruch 3, wobei der Marker YKL-40 ist.

5. Verfahren nach Anspruch 1, einen weiteren Schritt umfassend, bei dem man in einer Probe von dem Subjekt zwei oder mehr Marker nachweist, die ausgewählt sind aus der Gruppe bestehend aus: PIINP, Laminin, Tenascin, Kollagen Typ IV, Kollagen Typ VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 Komplex, sFas Ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 und apoB.

6. Verfahren nach Anspruch 1, wobei das Subjekt Virushepatitis hat.

7. Verfahren nach Anspruch 6, wobei das Subjekt mit Hepatitis C Virus infiziert ist.

8. Verfahren nach Anspruch 6, wobei das Subjekt mit Hepatitis B Virus infiziert ist.

9. Verfahren nach Anspruch 1, wobei das Subjekt autoimmune Lebererkrankung hat.

10. Verfahren nach Anspruch 1, wobei das Subjekt alkoholbedingte Lebererkrankung hat.

11. Verfahren nach Anspruch 1, wobei das Subjekt eine Fettlebererkrankung hat.

12. Verfahren nach Anspruch 1, wobei das Subjekt Medikamenteninduzierte Lebererkrankung hat.

13. Verfahren nach Anspruch 1, wobei Schritt (a) die Bestimmung der Menge des  $\alpha$ 2-MG Proteins in der Probe umfasst.

14. Verfahren nach Anspruch 13, wobei die Menge des  $\alpha$ 2-MG Proteins unter Verwendung eines oder mehrerer  $\alpha$ 2-

MG-spezifischer, bindender Agenzien bestimmt wird.

15. Verfahren nach Anspruch 14, wobei die Menge des  $\alpha$ 2-MG Proteins unter Verwendung eines oder mehrerer anti- $\alpha$ 2-MG-spezifischer Antikörper bestimmt wird.

16. Verfahren nach Anspruch 1, wobei Schritt (a) die Bestimmung eines Niveaus von  $\alpha$ 2-MG Aktivität umfasst.

17. Verfahren nach Anspruch 1, wobei Schritt (b) die Bestimmung der Menge von HA in der Probe umfasst.

18. Verfahren nach Anspruch 17, wobei die Menge von HA unter Verwendung eines oder mehrerer HA-spezifischer, bindender Agenzien bestimmt wird.

19. Verfahren nach Anspruch 18, wobei die Menge von HA unter Verwendung eines oder mehrerer HA-bindender Proteine bestimmt wird.

20. Verfahren nach Anspruch 18, wobei die Menge von HA unter Verwendung eines oder mehrerer anti-HA Antikörper bestimmt wird.

21. Verfahren nach Anspruch 1, wobei Schritt (c) die Bestimmung der Menge des TIMP-1 Proteins in der Probe umfasst.

22. Verfahren nach Anspruch 21, wobei die Menge des TIMP-1 Proteins unter Verwendung eines oder mehrerer TIMP-1-spezifischer, bindender Agenzien bestimmt wird.

23. Verfahren nach Anspruch 22, wobei die Menge des TIMP-1 Proteins unter Verwendung eines oder mehrerer anti-TIMP-1 Antikörper bestimmt wird.

24. Verfahren nach Anspruch 1, wobei Schritt (c) die Bestimmung eines Niveaus von TIMP-1 Aktivität umfasst.

25. Verfahren nach Anspruch 1, wobei Schritt (a) die Bestimmung der Menge von  $\alpha$ 2-MG Protein umfasst, wobei Schritt (b) die Bestimmung der Menge von HA umfasst, und wobei Schritt (c) die Bestimmung der Menge von TIMP-1 Protein umfasst.

26. Verfahren nach Anspruch 25, wobei die Menge von  $\alpha$ 2-MG Protein, HA und TIMP-1 Protein jeweils unter Verwendung eines enzymgekoppelten Assays bestimmt wird.

27. Verfahren nach Anspruch 1, wobei eine einzelne Probe von dem Subjekt entnommen wird.

28. Verfahren nach Anspruch 27, wobei die Probe ausgewählt ist aus der Gruppe bestehend aus Blut, Serum, Plasma, Urin, Speichel und Lebergewebe.

29. Verfahren nach Anspruch 28, wobei die Probe eine Serumprobe ist.

30. Verfahren nach Anspruch 1, einen Schritt umfassend, bei dem zwischen keiner oder leichter Leberfibrose und moderater bis schwerer Leberfibrose unterschieden wird.

31. Verfahren zur Differenzierung zwischen keiner oder leichter Leberfibrose und moderater bis schwerer Leberfibrose in einem Subjekt, Schritte umfassend, bei denen man:

- (a) eine geeignete Verdünnung einer Probe von dem Subjekt mit anti- $\alpha$ 2-MG Antikörper unter Bedingungen kontaktiert, die geeignet sind, um einen ersten Komplex aus  $\alpha$ 2-MG und anti- $\alpha$ 2-MG Antikörper zu bilden;
- (b) den ersten Komplex wäscht, um ungebundene Moleküle zu entfernen;
- (c) die Menge an  $\alpha$ 2-MG-beinhaltenden, ersten Komplex bestimmt;
- (d) eine geeignete Verdünnung einer Probe von dem Subjekt mit einem HA-bindenden Protein (HABP) unter Bedingungen kontaktiert, die geeignet sind, um einen zweiten Komplex aus HA und HABP zu bilden;
- (e) den zweiten Komplex wäscht, um ungebundene Moleküle zu entfernen;
- (f) die Menge an HA-beinhaltenden, zweiten Komplex bestimmt;
- (g) eine geeignete Verdünnung einer Probe von dem Subjekt mit anti-TIMP-1 Antikörper unter Bedingungen

kontaktiert, die geeignet sind, um einen dritten Komplex aus TIMP-1 und anti-TIMP-1 Antikörpern zu bilden;  
 (h) den dritten Komplex wäscht, um ungebundene Moleküle zu entfernen;  
 (i) die Menge an TIMP-1-beinhaltenden, dritten Komplex bestimmt; und  
 (j) basierend auf den Mengen von  $\alpha$ 2-MG, HA und TIMP-1-beinhaltenden Komplexen zwischen keiner/leichter  
 5 Leberfibrose und moderater/schwerer Leberfibrose in dem Subjekt unterscheidet.

32. Verfahren zur Überwachung der Wirksamkeit von antifibrotischer Therapie in einem Patienten, Schritte umfassend, bei denen man:

10 (a)  $\alpha$ 2-Makroglobulin in einer Probe eines Patienten nachweist, der eine antifibrotische Therapie erhielt;  
 (b) Hyaluronsäure (HA) in einer Probe von dem Patienten nachweist;  
 (c) Gewebeinhibitor von Metalloproteinasen-1 (TIMP-1) in einer Probe des Patienten nachweist; und  
 (d) basierend auf dem Vorhandensein oder der Menge von  $\alpha$ 2-MG, HA und TIMP-1 das Vorhandensein oder  
 15 den Schweregrad von Leberfibrose in dem Patienten bestimmt und dadurch die Wirksamkeit von antifibrotischer Therapie überwacht.

33. Verfahren nach Anspruch 32, einen weiteren Schritt umfassend, bei dem das Vorhandensein oder der Schweregrad von Leberfibrose in Schritt (d) mit dem Vorhandensein oder dem Schweregrad von Leberfibrose in dem Patienten zu einem früheren Zeitpunkt verglichen wird.

34. Verfahren nach Anspruch 32, einen Schritt umfassend, bei dem höchstens drei Fibrosemarker nachgewiesen werden.

35. Verfahren nach Anspruch 32, einen weiteren Schritt umfassend, bei dem in einer Probe von dem Patienten mindestens ein Marker nachgewiesen wird, der ausgewählt ist aus der Gruppe bestehend aus: PIINP, Laminin, Tenascin, Kollagen Typ IV, Kollagen TYP VI, YKL-40, MMP-3, MMP-2, MMP-9/TIMP-1 Komplex, sFas Ligand, TGF- $\beta$ 1, IL-10, apoA1, apoA2 und apoB.

36. Verfahren nach Anspruch 32, wobei Schritt (a) die Bestimmung der Menge von  $\alpha$ 2-MG Protein in der Probe umfasst.

37. Verfahren nach Anspruch 36, wobei die Menge von  $\alpha$ 2-MG Proteins unter Verwendung eines oder mehrerer anti- $\alpha$ 2-MG Antikörper bestimmt wird.

38. Verfahren nach Anspruch 32, wobei Schritt (b) die Bestimmung der Menge von HA in der Probe umfasst.

39. Verfahren nach Anspruch 38, wobei die Menge von HA unter Verwendung eines oder mehrerer HA-bindender Proteine bestimmt wird.

40. Verfahren nach Anspruch 32, wobei Schritt (c) die Bestimmung der Menge von TIMP-1 Protein in der Probe umfasst.

41. Verfahren nach Anspruch 40, wobei die Menge von TIMP-1 Protein unter Verwendung eines oder mehrerer anti-TIMP-1 Antikörper bestimmt wird.

42. Verfahren zur Differenzierung zwischen keiner/leichter und moderater/schwerer Leberfibrose in einem Subjekt, Schritte umfassend, bei denen man:

(a) eine Menge von  $\alpha$ 2-MG in einer Probe von dem Subjekt bestimmt;  
 (b) eine Menge von HA in einer Probe von dem Subjekt bestimmt;  
 (c) die Menge von TIMP-1 in einer Probe von dem Subjekt bestimmt; und  
 50 (d) das Subjekt als keine/leichte Leberfibrose aufweisend diagnostiziert, wenn die Menge von  $\alpha$ 2-MG unterhalb eines  $\alpha$ 2-MG Grenzwertes X1, die HA Menge unterhalb eines HA Grenzwertes Y1 oder die TIMP-1 Menge unterhalb eines TIMP-1 Grenzwertes Z1 ist,  
 das Subjekt als moderate/schwere Leberfibrose aufweisend diagnostiziert, wenn die Menge von  $\alpha$ 2-MG oberhalb eines  $\alpha$ 2-MG Grenzwertes X2, die HA Menge oberhalb eines HA Grenzwertes Y2 und die TIMP-1 Menge oberhalb eines TIMP-1 Grenzwertes Z2 ist,  
 55 und verbliebene Subjekte als einen unbestimmten Status aufweisend diagnostiziert.

43. Verfahren nach Anspruch 42, wobei das Subjekt eine Krankheit hat, die ausgewählt ist aus der Gruppe bestehend

aus Virushepatitis, autoimmune Lebererkrankung, alkoholbedingte Lebererkrankung, Fettlebererkrankung und Medikamenteninduzierte Lebererkrankung.

44. Verfahren nach Anspruch 43, wobei das Subjekt mit Hepatitis C Virus infiziert ist.

45. Verfahren nach Anspruch 42, wobei die Proben unabhängig ausgewählt sind aus der Gruppe bestehend aus Blut, Serum, Plasma, Urin, Speichel und Lebergewebe.

46. Verfahren nach Anspruch 45, wobei die Menge von  $\alpha$ 2-MG, die Menge von HA und die Menge von TIMP-1 jeweils in einer Serumprobe bestimmt wird.

47. Verfahren nach Anspruch 46,  
wobei X1 ein Wert zwischen 1,8 und 2,2 mg/ml ist;  
wobei Y1 ein Wert zwischen 31 und 39 ng/ml ist;  
wobei Z1 ein Wert zwischen 900 und 1100 ng/ml ist;  
wobei X2 ein Wert zwischen 1,8 und 2,2 mg/ml ist;  
wobei Y2 ein Wert zwischen 54 und 66 ng/ml ist; und  
wobei Z2 ein Wert zwischen 1415 und 1735 ng/ml ist.

48. Verfahren nach Anspruch 47,  
wobei X1 = 2,0 mg/ml;  
wobei Y1 = 35 ng/ml;  
wobei Z1 = 1000 ng/ml;  
wobei X2 = 2,0 mg/ml;  
wobei Y2 = 60 ng/ml; und  
wobei Z2 = 1575 ng/ml.

49. Verfahren nach Anspruch 47,  
wobei X1 = 2,0 mg/ml;  
wobei Y1 = 37 ng/ml;  
wobei Z1 = 1100 ng/ml;  
wobei X2 = 2,0 mg/ml;  
wobei Y2 = 60 ng/ml; und  
wobei Z2 = 1575 ng/ml.

50. Verfahren nach Anspruch 42, wobei in einer Population, die eine Leberfibrose-Prävalenz von bis zu 30% aufweist, mindestens 65% der Subjekte dieser Population mit einer Genauigkeit von mindestens 80% als keine/leichte Fibrose oder moderate/schwere Fibrose aufweisend diagnostiziert werden.

51. Verfahren nach Anspruch 42, wobei in einer Population, die eine Leberfibrose-Prävalenz von bis zu 30% aufweist, mindestens 65% der Subjekte dieser Population mit einer Genauigkeit von mindestens 90% als keine/leichte Fibrose oder moderate/schwere Fibrose aufweisend diagnostiziert werden.

52. Verfahren nach Anspruch 42, wobei in einer Population, die eine Leberfibrose-Prävalenz von bis zu 30% aufweist, mindestens 65% der Subjekte dieser Population mit einem positiven Vorhersagewert von mindestens 90% und einem negativen Vorhersagewert von mindestens 90% als keine/leichte Fibrose oder moderate/schwere Fibrose aufweisend diagnostiziert werden.

53. Verfahren nach Anspruch 42, wobei in einer Population, die eine Leberfibrose-Prävalenz von bis zu 10% aufweist, mindestens 70% der Subjekte dieser Population mit einer Genauigkeit von mindestens 90% als keine/leichte Fibrose oder moderate/schwere Fibrose aufweisend diagnostiziert werden.

## Revendications

1. Procédé de diagnostic de la présence ou de la gravité d'une fibrose hépatique chez un individu, comprenant les étapes de :

- (a) détection de l' $\alpha$ 2-macroglobuline dans un échantillon provenant dudit individu ;
- (b) détection de l'acide hyaluronique (HA) dans un échantillon provenant dudit individu ;
- (c) détection de l'inhibiteur tissulaire de métalloprotéinases 1 (TIMP-1) dans un échantillon provenant dudit individu ; et
- (d) diagnostic de la présence ou de la gravité d'une fibrose hépatique chez ledit individu, basé sur la présence ou le taux de  $\alpha$ 2-MG, HA et TIMP-1.

2. Procédé selon la revendication 1, comprenant la détection d'au maximum trois marqueurs de fibrose.

3. Procédé selon la revendication 1, en outre comprenant la détection dans un échantillon provenant dudit individu d'au moins un marqueur choisi dans le groupe constitué par : PIINP, laminine, ténascine, collagène de type IV, collagène de type VI, YKL-40, MMP-3, MMP-2, complexe MMP-9/TIMP-1, ligand sFas, TGF- $\beta$ 1, IL-10, apoA1 apoA2 et apoB.

4. Procédé selon la revendication 3, **caractérisé en ce que** ledit marqueur est YKL-40.

5. Procédé selon la revendication 1, en outre comprenant la détection dans un échantillon provenant dudit individu de deux ou plus de deux marqueurs choisis dans le groupe constitué par PIINP, laminine, ténascine, collagène de type IV, collagène de type VI, YKL-40, MMP-3, MMP-2, complexe MMP-9/TIMP-1, ligand sFas, TGF- $\beta$ 1, IL-10, apoA1 apoA2 et apoB.

6. Procédé selon la revendication 1, dans lequel ledit individu souffre d'hépatite virale.

7. Procédé selon la revendication 6, dans lequel ledit individu est infecté par le virus de l'hépatite C.

8. Procédé selon la revendication 6, dans lequel ledit individu est infecté par le virus de l'hépatite B.

9. Procédé selon la revendication 1, dans lequel ledit individu a une hépatopathie auto-immune.

10. Procédé selon la revendication 1, dans lequel ledit individu a une hépatopathie alcoolique.

11. Procédé selon la revendication 1, dans lequel ledit individu a une stéatose hépatique.

12. Procédé selon la revendication 1, dans lequel ledit individu a une hépatopathie d'origine médicamenteuse.

13. Procédé selon la revendication 1, dans lequel l'étape (a) comprend la détermination du taux de la protéine  $\alpha$ 2-MG dans ledit échantillon.

14. Procédé selon la revendication 13, dans lequel on détermine le taux de la protéine  $\alpha$ 2-MG en utilisant un ou plusieurs agents de liaison spécifiques de  $\alpha$ 2-MG.

15. Procédé selon la revendication 14, dans lequel on détermine le taux de la protéine  $\alpha$ 2-MG en utilisant un ou plusieurs anticorps anti- $\alpha$ 2-MG.

16. Procédé selon la revendication 1, dans lequel l'étape (a) comprend la détermination d'un taux d'activité  $\alpha$ 2-MG.

17. Procédé selon la revendication 1, dans lequel l'étape (b) comprend la détermination du taux de HA dans ledit échantillon.

18. Procédé selon la revendication 17, dans lequel on détermine le taux de HA en utilisant un ou plusieurs agents de liaison spécifiques de HA.

19. Procédé selon la revendication 18, dans lequel on détermine le taux de HA en utilisant une ou plusieurs protéines se liant à HA.

20. Procédé selon la revendication 18, dans lequel on détermine le taux de HA en utilisant un ou plusieurs anticorps anti-HA.

21. Procédé selon la revendication 1, dans lequel l'étape (c) comprend la détermination du taux de la protéine TIMP-1 dans ledit échantillon.
- 5 22. Procédé selon la revendication 21, dans lequel on détermine le taux de la protéine TIMP-1 en utilisant un ou plusieurs agents de liaison spécifiques de TIMP-1.
23. Procédé selon la revendication 22, dans lequel on détermine le taux de la protéine TIMP-1 en utilisant un ou plusieurs anticorps anti-TIMP-1.
- 10 24. Procédé selon la revendication 1, dans lequel l'étape (c) comprend la détermination d'un taux d'activité TIMP-1.
25. Procédé selon la revendication 1, dans lequel l'étape (a) comprend la détermination du taux de la protéine  $\alpha$ 2-MG, dans lequel l'étape (b) comprend la détermination du taux de HA, et  
15 dans lequel l'étape (c) comprend la détermination du taux de la protéine TIMP-1.
26. Procédé selon la revendication 25, on détermine chacun des taux de la protéine  $\alpha$ 2-MG, de HA et de la protéine TIMP-1 en utilisant un dosage à enzyme liée.
- 20 27. Procédé selon la revendication 1, dans lequel on obtient un seul échantillon à partir dudit individu.
28. Procédé selon la revendication 27, dans lequel ledit échantillon est choisi dans le groupe constitué par le sang, le sérum, le plasma, l'urine, la salive et des tissus hépatiques.
- 25 29. Procédé selon la revendication 28, dans lequel ledit échantillon est un échantillon de sérum.
30. Procédé selon la revendication 1, comprenant la distinction d'une fibrose hépatique légère ou absente d'avec une fibrose hépatique modérée à grave.
- 30 31. Procédé de distinction d'une fibrose hépatique légère ou absente d'avec une fibrose hépatique modérée à grave chez un individu, comprenant les étapes de :
  - (a1) mise en contact d'une dilution convenable d'un échantillon provenant dudit individu avec un anticorps anti- $\alpha$ 2-MG dans des conditions appropriées à la formation d'un premier complexe d' $\alpha$ 2-MG et d'anticorps anti- $\alpha$ 2-MG;  
35 (b) lavage dudit premier complexe pour l'élimination des molécules non fixées ;  
(c) détermination de la quantité du premier complexe contenant  $\alpha$ 2-MG;  
(d) mise en contact d'une dilution convenable d'un échantillon provenant dudit individu avec une protéine se liant à HA (HAPB) dans des conditions appropriées à la formation d'un deuxième complexe de HA et HAPB;  
40 (e) lavage dudit deuxième complexe pour l'élimination des molécules non fixées ;  
(f) détermination de la quantité du deuxième complexe contenant HA;  
(g) mise en contact d'une dilution convenable d'un échantillon provenant dudit individu avec un anticorps anti-TIMP-1 dans des conditions appropriées à la formation d'un troisième complexe de TIMP-1 et d'anticorps anti-TIMP-1 ;  
45 (h) lavage dudit troisième complexe pour l'élimination des molécules non fixées ;  
(i) détermination de la quantité du troisième complexe contenant TIMP-1; et  
(j) distinction d'une fibrose hépatique légère/absente d'avec une fibrose hépatique modérée/grave chez ledit individu sur la base des quantités des complexes contenant  $\alpha$ 2-MG, HA et TIMP-1.
- 50 32. Procédé de contrôle de l'efficacité d'une thérapie anti-fibrose chez un patient, comprenant les étapes de :
  - (a) détection de l' $\alpha$ 2-macroglobuline dans un échantillon provenant d'un patient auquel est administrée une thérapie anti-fibrose ;
  - (b) détection de l'acide hyaluronique (HA) dans un échantillon provenant dudit patient ;
  - 55 (c) détection de l'inhibiteur tissulaire de métalloprotéinases 1 (TIMP-1) dans un échantillon provenant dudit patient ; et  
(d) détermination de la présence ou de la gravité d'une fibrose hépatique chez ledit patient, basée sur la présence ou le taux de  $\alpha$ 2-MG, HA et TIMP-1, permettant ainsi de contrôler l'efficacité de la thérapie anti-fibrose.

33. Procédé selon la revendication 32, en outre comprenant la comparaison de la présence ou de la gravité d'une fibrose hépatique, déterminée dans l'étape (d), avec la présence ou la gravité d'une fibrose hépatique chez ledit patient à un moment antérieur.
- 5 34. Procédé selon la revendication 32, comprenant la détection d'au maximum trois marqueurs de fibrose.
35. Procédé selon la revendication 32, en outre comprenant la détection dans un échantillon provenant dudit patient d'au moins un marqueur choisi dans le groupe constitué par : PIINP, laminine, ténascine, collagène de type IV, collagène de type VI, YKL-40, MMP-3, MMP-2, complexe MMP-9/TIMP-1, ligand sFas, TGF- $\beta$ 1, IL-10, apoA1 apoA2 et apoB.
- 10 36. Procédé selon la revendication 32, dans lequel l'étape (a) comprend la détermination du taux de la protéine  $\alpha$ 2-MG dans ledit échantillon.
- 15 37. Procédé selon la revendication 36, dans lequel on détermine le taux de la protéine  $\alpha$ 2-MG en utilisant un ou plusieurs anticorps anti- $\alpha$ 2-MG.
38. Procédé selon la revendication 32, dans lequel l'étape (b) comprend la détermination du taux de HA dans ledit échantillon.
- 20 39. Procédé selon la revendication 38, dans lequel on détermine le taux de HA en utilisant une ou plusieurs protéines se liant à HA.
40. Procédé selon la revendication 32, dans lequel l'étape (c) comprend la détermination du taux de la protéine TIMP-1 dans ledit échantillon.
- 25 41. Procédé selon la revendication 40, dans lequel on détermine le taux de la protéine TIMP-1 en utilisant un ou plusieurs anticorps anti-TIMP-1.
- 30 42. Procédé de distinction d'une fibrose hépatique absente/légère d'avec une fibrose hépatique modérée/grave chez un individu, comprenant les étapes de :
- (a) détection d'un taux d' $\alpha$ 2-MG dans un échantillon provenant dudit individu ;
- (b) détection d'un taux de HA dans un échantillon provenant dudit individu ;
- 35 (c) détection d'un taux de TIMP-1 dans un échantillon provenant dudit individu ; et
- (d) diagnostic dudit individu comme n'ayant pas de fibrose hépatique/ayant une légère fibrose hépatique lorsque ledit taux d' $\alpha$ 2-MG est inférieur à une valeur seuil X1 d' $\alpha$ 2-MG, ledit taux de HA est inférieur à une valeur seuil Y1 de HA ou ledit taux de TIMP-1 est inférieur à une valeur seuil Z1 de TIMP-1,
- 40 diagnostic dudit individu comme ayant une fibrose hépatique modérée à grave lorsque ledit taux d' $\alpha$ 2-MG est supérieur à une valeur seuil X2 d' $\alpha$ 2-MG, ledit taux de HA est supérieur à une valeur seuil Y2 de HA et ledit taux de TIMP-1 est supérieur à une valeur seuil Z2 de TIMP-1,
- et diagnostic des individus restants comme ayant un état indéterminé.
- 45 43. Procédé selon la revendication 42, dans lequel ledit individu a une affection choisie dans le groupe constitué par l'hépatite virale, l'hépatopathie auto-immune, l'hépatopathie alcoolique, la stéatose hépatique et une hépatopathie d'origine médicamenteuse.
- 50 44. Procédé selon la revendication 43, dans lequel ledit individu est infecté par le virus de l'hépatite C.
45. Procédé selon la revendication 42, dans lequel lesdits échantillons sont choisis indépendamment dans le groupe constitué par le sang, le sérum, le plasma, l'urine, la salive et un tissu hépatique.
- 55 46. Procédé selon la revendication 45, dans lequel on détermine chacun des taux de  $\alpha$ 2-MG, HA et TIMP-1 dans un échantillon de sérum.
47. Procédé selon la revendication 46, dans lequel X1 est une valeur comprise entre 1,8 et 2,2 mg/ml ;

dans lequel Y1 est une valeur comprise entre 31 et 39 ng/ml ;  
 dans lequel Z1 est une valeur comprise entre 900 et 1 100 ng/ml ;  
 dans lequel X2 est une valeur comprise entre 1,8 et 2,2 mg/ml ;  
 dans lequel Y2 est une valeur comprise entre 54 et 66 ng/ml ; et  
 dans lequel Z2 est une valeur comprise entre 1 415 et 1 735 ng/ml.

48. Procédé selon la revendication 47,

dans lequel X1 = 2,0 mg/ml ;  
 dans lequel Y1 = 35 ng/ml ;  
 dans lequel Z1 = 1 000 ng/ml ;  
 dans lequel X2 = 2,0 mg/ml ;  
 dans lequel Y2 = 60 ng/ml ; et  
 dans lequel Z2 = 1 575 ng/ml.

49. Procédé selon la revendication 47,

dans lequel X1 = 2,0 mg/ml ;  
 dans lequel Y1 = 37 ng/ml ;  
 dans lequel Z1 = 1 100 ng/ml ;  
 dans lequel X2 = 2,0 mg/ml ;  
 dans lequel Y2 = 60 ng/ml ; et  
 dans lequel Z2 = 1 575 ng/ml.

50. Procédé selon la revendication 42, dans lequel, dans une population ayant jusqu'à 30 % de prévalence de fibrose hépatique, au moins 65 % des individus dans ladite population sont diagnostiqués come n'ayant pas de fibrose/ ayant une fibrose légère ou une fibrose modérée/grave avec une exactitude d'au moins 80 %.

51. Procédé selon la revendication 42, dans lequel, dans une population ayant jusqu'à 30 % de prévalence de fibrose hépatique, au moins 65 % des individus dans ladite population sont diagnostiqués come n'ayant pas de fibrose/ ayant une fibrose légère ou une fibrose modérée/grave avec une exactitude d'au moins 90 %.

52. Procédé selon la revendication 42, dans lequel, dans une population ayant jusqu'à 30 % de prévalence de fibrose hépatique, au moins 65 % des individus dans ladite population sont diagnostiqués come n'ayant pas de fibrose/ ayant une fibrose légère ou une fibrose modérée/grave avec une valeur prédictive positive d'au moins 90 % et une valeur prédictive négative d'au moins 90 %.

53. Procédé selon la revendication 42, dans lequel, dans une population ayant jusqu'à 10 % de prévalence de fibrose hépatique, au moins 70 % des individus dans ladite population sont diagnostiqués come n'ayant pas de fibrose/ ayant une fibrose légère ou une fibrose modérée/grave avec une exactitude d'au moins 90 %.



```

1 cccgccttcc tagctgtccc agtggagaag gaacaagcgc ctcactgcat ctgtgcaaac
61 gggcggaaca ctgtgtcctg ggcagtaacc ccaaagtcac taggaaatgt gaatttcact
121 gtgagcgagc aggcactaga gtctcaagag ctgtgtggga ctgaggtgcc ttcagttcct
181 gaacacggaa ggaaagacac agtcatcaag cctctgttgg ttgaacctga aggactagag
241 aaggaaacaa cattcaactc cctactttgt ccatcaggtg gtgaggtttc tgaagaatta
301 tccctgaaac tgccaccaa tgtggtagaa gaatctgccc gagcttctgt ctgagttttg
361 ggagacatat taggctctgc catgcaaaac acacaaaatc ttctccagat gccctatggc
421 tgtggagagc agaatatggc cctctttgct cctaaccatc atgtactgga ttatctaaat
481 gaaacacagc agcttactcc agagatcaag tccaaggcca ttggctatct caacactggc
541 taccagagac agttgaacta caaacactat gatggctcct acagcacctt tggggagcga
601 tatggcagga accagggcaa cacctggctc acagcctttg ttctgaagac ttttgcccaa
661 gctcgagcct acatcttcat cgatgaagca cacattaccc aagcctcat atggctctcc
721 cagaggcaga aggacaatgg ctgtttcagg agctctgggt cactgctcaa caatgccata
781 aaggaggagg tagaagatga agtgacctc tccgcctata tcaccatcgc ccttctggag
841 attcctctca cagtactca cctgtgtgtc cgcaatgccc tgttttgctt ggagtcagcc
901 tggaagacag cacaagaagg ggaccatggc agccatgtat ataccaaaga cctgctggcc
961 tatgcttttg ccttggcagg taaccaggac aagaggaagg aagtactcaa gtcacttaat
1021 gaggaagctg tgaagaaaga caactctgtc cattgggagc gccctcagaa acccaaggca
1081 ccagtggggg atttttacga acccaggct cctctgctg aggtggagat gacatcctat
1141 gtgctcctcg cttatctcac ggcccagcca gcccacacct cggaggacct gacctctgca
1201 accaacatcg tgaagtggat cacgaagcag cagaatgccc agggcggttt ctcctccacc
1261 caggacacag tgggtggctc ccatgctctg tccaaatatg gagcagccac atttaccagg
1321 actgggaagg ctgcacaggt gactatccag tcttcaggga cattttccag caaattccaa
1381 gtggacaaca acaaccgcct gttactgcag caggtctcat tgccagagct gcctggggaa
1441 tacagcatga aagtgcagag agaaggatgt gtctacctcc agacatcctt gaaatacaat
1501 attctcccag aaaaggaaga gttccccctt gctttaggag tgcagactct gcctcaaact
1561 tgtgatgaac ccaaagccca caccagcttc caaatctccc taagtgtcag ttacacaggg
1621 agccgctctg cctccaacat ggcgatcggt gatgtgaaga tggctctcgg cttcattccc
1681 ctgaagccaa cagtgaatat gcttgaaaga tctaaccatg tgagccggac agaagtcagc
1741 agcaaccatg tcttgattta ccttgataag gtgtcaaata agacactgag cttgttcttc
1801 acggttctgc aagatgtccc agtaagagat ctgaaaccag ccatagtga agtctatgat
1861 tactacgaga cggatgagtt tgcaattgct gagtacaatg ctccttgag caaagatctt
1921 ggaaatgctt gaagaccaca aggtgaaaa gtgctttgct ggagtcctgt tctcagagct
1981 ccacagaaga cacgtgtttt tgtatcttta aagacttgat gaataaacac tttttctggt
2041 c

```

A

PAFLAVPVEKEQAPHICANGRQTVSWAVTPKSLGNVNFTVSAEALESQELCGTEVPSVPEHGRKDTVIKPL  
LVEPEGLEKETTFNSLLCPSSGGEVSEELSLKLPPNVVEESARASVSVLGDILGSAMQNTQNLLQMPYGCGEQ  
NMVLFAPNIYVLDYLNQQLTPEIKSKAIGYLNQYQRLNYKHYDGSYSTFGERYGRNQGNLWLTAFVLK  
TFAQARAYIFIDEAHITQALIWLSQLQKDNQCFRSSGSLNNNAIKGGVEDEVTL SAYITIALLEIPLTVTHP  
VVRNALFCLESARKTAQEGDHGSHVYTKDLLAYAFALAGNQDKRKEVLKSLNEEA VKKDNSVHWERPQKPKA  
PVGDFYEPQAPSAEEMTSYVLLAYLTAQPAPTSDELTSATNIVKWITKQONAQGGFSSTQDTVVALHALSK  
YGAATFTRTGKAAQVTIQSSGTFSSKFQVDNNNRLLLQVSLPELPGEYS MKVTGEGCVYLQTS LKYNILPE  
KEEFPFALGVQTLPTCDEPKAHTSFQISLSVSYTGSRASNM AIVDKMVS GFIP LKPTVKMLERSNHVSR  
TEVSSNHVLIYLDKVSNTLSLFFTVLQDVPVRDLKPAIVKVYDYETDEF AIAEYNAPCSKDLGNA

B

FIGURE 1

```

1 aggggcctta gcgtgcegca tcgcgagat ccagcgccca gagagacacc agagaaccca
61 ccatggcccc ctttgagccc ctggcttctg gcatcctggt gttgctgtgg ctgatatgcc
121 ccagcagggc ctgcacctgt gtcccacccc acccacagac ggccttctgc aattccgacc
181 tcgtcatcag ggccaagttc gtggggacac cagaagtcaa ccagaccacc ttataccagc
241 gttatgagat caagatgacc aagatgtata aagggttcca agccttaggg gatgccgtg
301 acatccggtt cgtctacacc ccgccatgg agagtgtctg cggatacttc cacagggtccc
361 acaaccgcag cgaggagtgt ctcattgctg gaaaactgca ggatggactc ttgcacatca
421 ctacctgcag tttcgtggct ccctggaaca gcctgagctt agctcagcgc cggggcttca
481 ccaagacctt cactgttggt tgtgaggaat gcacagtgtt tccctgttta tccatccct
541 gcaaaactgca gagtggcact cattgcttgt ggacggacca gtcctccaa ggctctgaaa
601 agggcttcca gtcccgtcac cttgcctgcc tgcccgga gccagggctg tgcacctggc
661 agtccctgcg gtcccagata gcctgaatcc tgcccgaggt ggaactgaag cctgcacagt
721 gtccacctg ttccactec catcttctt ccggacaatg aaataaagag ttaccacca
781 gc

```

A

```

MAPFEPLASGILLLLWLIAPSRACCTVPPHPQTAFCSNDLVIRAKFVGTPEVNQTTLYQRYEIKMTKMY
KGFQALGDAADIRFVYTPAMESVCGYFHRSHNRSEEFLLIAGKLQDGLLHITTCFVAPWNSLSLAQRRG
FTKTYTVGCEECTVFPCLSI PCKLQSGTHCLWTDQLQSEKGFQSRHLACLPREPGLCTWQSLRSQIA

```

B

FIGURE 2

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 10087188 B [0186]

## Non-patent literature cited in the description

- **ALTER.** *Semin. Liver Dis.*, 1995, vol. 15, 5-14 [0003]
- **GRAUDAL.** *J. Intern. Med.*, 1991, vol. 230, 165-171 [0004]
- **DAVIS.** *Hepatology*, vol. 26 (1), 122-127S [0005]
- *Hepatology*, 1997, vol. 26 (1), 25-105S [0005]
- **CALÉS ; PASQUAL.** *Gastroenterol. Clin. Biol.*, 1988, vol. 12, 245-254 [0005]
- **NORD.** *Gastrointest. Endosc.*, 1982, vol. 28, 102-104 [0006]
- **CADRANEL et al.** *Hepatology*, 2000, vol. 32, 47-481 [0006]
- **POYNARD et al.** *Can. J. Gastroenterol.*, 2000, vol. 14, 543-548 [0006]
- **KNODELL et al.** *Hepatology*, 1981, vol. 1, 431-435 [0031]
- **BRUNT et al.** *Am. J. Gastroenterol.*, 1999, vol. 94, 2467-2474 [0037]
- **STARKEY et al.** *Biochem. J.*, 1973, vol. 131, 823-831 [0041]
- $\alpha$ 2-Macroglobulin and Related Thiol Ester Plasma Proteins. **SOTTRUP-JENSEN.** *The Plasma Proteins: Structure, Function and Genetic Control.* Academic Press, 1987, 191-291 [0043]
- **ORTEGO et al.** *Exp. Eye Res.*, 1997, vol. 65, 289-299 [0047]
- **SIMON et al.** *Cancer Res.*, 1996, vol. 56, 3112-3117 [0047]
- **FINK et al.** *J. Clin. Chem. Clin. Biol. Chem.*, 1989, vol. 27, 261-276 [0048]
- **NAVEAU et al.** *Dig. Diseases Sci.*, 1994, vol. 39, 2426-2432 [0048]
- **HALL ; ROBERTS.** *Biochem. J.*, 1978, vol. 171, 27-38 [0048]
- **IMBER ; PEZZO.** *J. Biol. Chem.*, 1981, vol. 256, 8134-8139 [0048]
- **ARMSTRONG et al.** *Develop. Compar. Immunol.*, 1999, vol. 23, 375-390 [0049]
- **GANROT.** *Clin. Chim. Acta*, 1966, vol. 14, 493-501 [0051]
- **ARMSTRONG et al.** *J. Exp. Zool.*, 1985, vol. 236, 1-9 [0051]
- **BRAMLEY et al.** *J. Hepatol.*, 1991, vol. 13, 8-13 [0053]
- **UENO et al.** *Gastroenterol.*, 1993, vol. 105, 475-481 [0053]
- **OBERTI et al.** *Gastroenterol.*, 1997, vol. 113, 1609-1616 [0053] [0119]
- **MCHUTCHISON et al.** *J. Gastroenterol. Hepatol.*, 2000, vol. 15, 945-951 [0053]
- **KONTTINEN et al.** *Clin. Chimica Acta*, 1990, vol. 193, 39-48 [0053]
- **POOLE et al.** *Arthritis Rheum.*, 1994, vol. 37, 1030-1038 [0053]
- **GOLDBERG et al.** *Arthritis Rheum.*, 1991, vol. 34, 799-807 [0053]
- **EMLEM et al.** *J. Rheum.*, 1996, vol. 23, 974-978 [0053]
- **GUECHOT et al.** *Clin. Chem.*, 1996, vol. 42, 558-563 [0055] [0105]
- **MAINGONNAT ; DELPECH.** *Ann. Clin. Biochem.*, 1991, vol. 28, 305-306 [0055]
- **DELPECH ; BERTRAND.** *Anal. Biochem.*, 1985, vol. 149, 555-565 [0055]
- **LINDQUIST et al.** *Clin. Chem.*, 1992, vol. 38, 127-132 [0055]
- **ENGSTROM-LAURENT et al.** *Scand. J. Clin. Lab. Invest.*, 1985, vol. 45, 497-504 [0055]
- **BRANDT et al.** *Acta Otolaryn.*, 1987, vol. 442, 31-35 [0055]
- **GOLDBERG.** *Anal. Biochem.*, 1988, vol. 174, 448-458 [0055]
- **CHICHIBU et al.** *Clin. Chim. Acta*, 1989, vol. 181, 317-324 [0055]
- **LI et al.** *Conn. Tissue Res.*, 1989, vol. 19, 243-254 [0055]
- **POOLE et al.** *Arth. Rheum.*, 1990, vol. 33, 790-799 [0055]
- **POOLE et al.** *J. Biol. Chem.*, 1985, vol. 260, 6020-6025 [0055]
- **LAURENT ; TENGBLAD.** *Anal. Biochem.*, 1980, vol. 109, 386-394 [0055]
- **PAYAN et al.** *J. Chromatogr.*, 1991, vol. 566, 9-18 [0056]
- **BRAY et al.** *Am. Rev. Respir. Dis.*, 1991, vol. 3, 284-288 [0057]

- Protein Inhibitors of Metalloproteinases. **CAWSTON et al.** Proteinase Inhibitors. Elsevier, 1986, 589-610 [0058]
- **MURPHY et al.** *Biochem. J.*, 1981, vol. 195, 167-170 [0059] [0062]
- **DOCKERTY et al.** *Nature*, 1985, vol. 318, 66-69 [0059]
- **BODDEN et al.** *J. Biol. Chem.*, 1994, vol. 269, 18943-18952 [0059]
- **YOSHIJI et al.** *Int. J. Cancer*, 1996, vol. 69, 131-134 [0062]
- **JANOWSKA-WIECZOREK et al.** *Exp. Hematol.*, 2000, vol. 28, 1274-1285 [0062]
- **GROFT et al.** *Br. J. Cancer*, 2001, vol. 85, 55-63 [0062]
- **BROPHY et al.** *Biochem. Biophys. Res. Comm.*, 1990, vol. 167, 898-903 [0062] [0105]
- **MURAWAKI et al.** *Clinica Chimica Acta*, 1993, vol. 218, 47-58 [0062]
- **STRICKLIN ; WELGUS.** *J. Biol. Chem.*, 1983, vol. 258, 12252-12258 [0062]
- **KOSSAKOWSKA et al.** *Amer. J. Pathology*, 1998, vol. 153, 1895-1902 [0062]
- Intuitive Biostatistics. Oxford University Press, 1995 [0071]
- **POYNARD et al.** *Lancet*, 2001, vol. 357, 1069 [0082] [0153]
- **AUSUBEL et al.** Current Protocols in Molecular Biology. John Wiley & Sons, 1999, vol. 47 [0098]
- **DIEFFENBACH ; DVEKSLER.** PCR Primer: A Laboratory Manual. Cold Spring Harbor Press, 1995 [0099]
- **SELF ; COOK.** *Curr. Opin. Biotechnol.*, 1996, vol. 7, 60-65 [0101]
- **HARLOW ; LANE.** Antibodies A Laboratory Manual. Cold Spring Harbor Laboratory, 1988 [0101]
- **SCHMALZING ; NASHABEH.** *Electrophoresis*, 1997, vol. 18, 2184-93 [0107]
- **BAO.** *J. Chromatogr. B. Biomed. Sci.*, 1997, vol. 699, 963-80 [0107]
- **RONGEN et al.** *J. Immunol. Methods*, 1997, vol. 204, 105-133 [0107]
- **PARRA et al.** *J. Vasc. Surg.*, 1998, vol. 28, 669-675 [0110]
- Antibody Engineering. Oxford University Press, 1995 [0112]
- **DOUMAS et al.** *Clin. Chim Acta*, 1971, vol. 31, 87-96 [0122]
- **POYNARD et al.** *Gastroenterol.*, 1991, vol. 100, 1397-1402 [0122]
- **NAVEAU et al.** *Dig. Dis. Sci.*, 1994, vol. 39, 2426-2432 [0122]
- **HAKALA et al.** *J. Biol. Chem.*, 1993, vol. 268, 25803-15810 [0126]
- **NYIRKOS ; GOLDS.** *Biochem. J.*, 1990, vol. 268, 265-268 [0126]
- **RENKEMA et al.** *Eur. J. Biochem.*, 1998, vol. 251, 504-509 [0126]
- **VOLCK et al.** *Proc. Assoc. Am. Physicians*, 1998, vol. 110, 351-360 [0126]
- **JOHANSEN et al.** *J. Bone Miner. Res.*, 1992, vol. 7, 501-511 [0126]
- **VERHEIJDEN et al.** *Arthritis Rheum.*, 1997, vol. 40, 1115-1125 [0126]
- **JOHANSEN et al.** *Scand. J. Gastroenterol.*, 1997, vol. 32, 582-590 [0126]
- **JOHANSEN.** *J. Hepatol.*, 2000, vol. 32, 911-920 [0126]
- **JOHANSEN et al.** *Br. J. Rheumatology*, 1993, vol. 32, 949-955 [0127]
- Hepatol. The French Metavir Cooperative Study Group, 1994, vol. 20, 15-20 [0132]

专利名称(译)	诊断肝纤维化的方法		
公开(公告)号	<a href="#">EP1487395B1</a>	公开(公告)日	2013-04-10
申请号	EP2003743713	申请日	2003-02-28
[标]申请(专利权)人(译)	普罗米修斯实验室		
申请(专利权)人(译)	普罗米修斯实验室，INC.		
当前申请(专利权)人(译)	NESTEC S.A.		
[标]发明人	ROSE STEVEN L OH ESTHER H WALSH MICHAEL J		
发明人	ROSE, STEVEN, L. OH, ESTHER, H. WALSH, MICHAEL, J.		
IPC分类号	G01N33/576 G01N33/53 G01N33/543 G01N33/569 G01N33/573 G01N33/68 G01N33/96		
CPC分类号	G01N33/5767 G01N33/6893 G01N2333/4713 G01N2333/8107 G01N2333/8146 G01N2800/085 G01N2800/52 Y02A90/24 Y02A90/26 Y10S436/811 Y10S436/82 Y10S706/924 Y10T436/143333		
代理机构(译)	RUPP，CHRISTIAN		
优先权	10/087188 2002-02-28 US		
其他公开文献	EP1487395A4 EP1487395A2		
外部链接	<a href="#">Espacenet</a>		

## 摘要(译)

本发明提供了通过检测来自个体的样品中的 $\alpha$ 2-巨球蛋白 (  $\alpha$ 2-MG ) 来诊断个体肝纤维化的存在或严重性的方法。检测来自个体的样品中的透明质酸 ( HA ) ;检测个体样品中金属蛋白酶-1 ( TIMP-1 ) 的组织抑制剂;并基于 $\alpha$ 2-MG，HA和TIMP-1的存在或水平诊断个体中肝纤维化的存在或严重性。

```

ccc gcc ttc cta gct gtc cca gtg gag aag gaa caa gcg cct cac tgc 48
Pro Ala Phe Leu Ala Val Pro Val Glu Lys Glu Gln Ala Pro His Cys
1 5 10 15
atc tgt gca aac ggg cgg caa act gtg tcc tgg gca gta acc cca aag 96
Ile Cys Ala Asn Gly Arg Gln Thr Val Ser Trp Ala Val Thr Pro Lys
20 25 30
tca tta gga aat gtg aat ttc act gtg agc gca gag gca cta gag tct 144
Ser Leu Gly Asn Val Asn Phe Thr Val Ser Ala Glu Ala Leu Glu Ser
35 40 45
caa gag ctg tgt ggg act gag gtg cct tca gtt cct gaa cac gga agg 192
Gln Glu Leu Cys Gly Thr Glu Val Pro Ser Val Pro Glu His Gly Arg
50 55 60
aaa gac aca gtc atc aag cct ctg ttg gtt gaa cct gaa gga cta gag 240
Lys Asp Thr Val Ile Lys Pro Leu Leu Val Glu Pro Glu Gly Leu Glu
65 70 75 80
aag gaa aca aca ttc aac tcc cta ctt tgt cca tca ggt ggt gag gtt 288
Lys Glu Thr Thr Phe Asn Ser Leu Leu Cys Pro Ser Gly Gly Glu Val
85 90 95
tct gaa gaa tta tcc ctg aaa ctg cca cca aat gtg gta gaa gaa tct 336

```