



(11) **EP 2 474 631 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
12.02.2014 Bulletin 2014/07

(51) Int Cl.:
C12Q 1/68 ^(2006.01) **C07K 14/47** ^(2006.01)
C07K 14/705 ^(2006.01) **C07K 16/18** ^(2006.01)
C07K 16/28 ^(2006.01)

(21) Application number: **12163424.0**

(22) Date of filing: **22.12.2003**

(54) **Genetic polymorphisms associated with myocardial infarction, methods of detection and uses thereof**

Mit Myokardinfarkt assoziierte genetische Polymorphismen, Nachweisverfahren und deren Verwendungen

Polymorphismes génétiques associés à l'infarctus du myocarde, procédés de détection et utilisations associées

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**

(30) Priority: **20.12.2002 US 434778 P**
30.04.2003 US 466412 P
10.03.2003 US 453135 P
23.09.2003 US 504955 P

(43) Date of publication of application:
11.07.2012 Bulletin 2012/28

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
03800092.3 / 1 583 771

(73) Proprietor: **Celera Corporation**
Alameda, CA 94502 (US)

(72) Inventors:
• **Lakoubova, Olga**
Alameda, CA 94502 (US)
• **Devlin, James**
Alameda, CA 94502 (US)

• **Cargill, Michele**
Alameda, CA 94502 (US)

(74) Representative: **Jones Day**
Rechtsanwälte, Attorneys-at-Law
Patentanwälte
Prinzregentenstraße 11
80538 München (DE)

(56) References cited:
EP-A2- 0 609 059

• **ZHANG D ET AL: "TAK1 is activated in the
myocardium after pressure overload and is
sufficient to provoke heart failure in transgenic
mice", NATURE MEDICINE, NATURE
PUBLISHING GROUP, NEW YORK, NY, US, vol.
6, no. 5, 1 May 2000 (2000-05-01), pages 556-563,
XP002387985, ISSN: 1078-8956, DOI:
10.1038/75037**

Remarks:

The complete document including Reference Tables
and the Sequence Listing can be downloaded from
the EPO website

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**FIELD OF THE INVENTION**

[0001] The present invention is in the field of myocardial infarction diagnosis. In particular, the present invention relates to a specific single nucleotide polymorphism (SNP) in the human genome, and its association with myocardial infarction (including recurrent myocardial infarction) and related pathologies. Based on differences in allele frequencies in the myocardial infarction patient population relative to normal individuals, the naturally-occurring SNPs disclosed herein can be used as targets for the design of diagnostic reagents and the development of therapeutic agents, as well as for disease association and linkage analysis. In particular, the SNPs disclosed herein are useful for identifying an individual who is at an increased or decreased risk of developing myocardial infarction and for early detection of the disease, for providing clinically important information for the prevention and/or treatment of myocardial infarction, and for screening and selecting therapeutic agents. The SNPs disclosed herein are also useful for human identification applications. Methods, assays, kits, and reagents for detecting the presence of these polymorphisms and their encoded products are disclosed herein.

BACKGROUND OF THE INVENTIONMyocardial Infarction (including Recurrent Myocardial Infarction)

[0002] Myocardial infarction (MI) is the most common cause of mortality in developed countries. It is a multifactorial disease that involves atherogenesis, thrombus formation and propagation. Thrombosis can result in complete or partial occlusion of coronary arteries. The luminal narrowing or blockage of coronary arteries reduces oxygen and nutrient supply to the cardiac muscle (cardiac ischemia), leading to myocardial necrosis and/or stunning. MI, unstable angina, or sudden ischemic death are clinical manifestations of cardiac muscle damage. All three endpoints are part of the Acute Coronary Syndrome since the underlying mechanisms of acute complications of atherosclerosis are considered to be the same.

[0003] Atherogenesis, the first step of pathogenesis of MI, is a complex interaction between blood elements, mechanical forces, disturbed blood flow, and vessel wall abnormality. On the cellular level, these include endothelial dysfunction, monocytes/macrophages activation by modified lipoproteins, monocytes/macrophages migration into the neointima and subsequent migration and proliferation of vascular smooth muscle cells (VSMC) from the media that results in plaque accumulation.

[0004] In recent years, an unstable (vulnerable) plaque was recognized as an underlying cause of arterial thrombotic events and MI. A vulnerable plaque is a plaque, often not stenotic, that has a high likelihood of becoming disrupted or eroded, thus forming a thrombogenic focus. Two vulnerable plaque morphologies have been described. A first type of vulnerable plaque morphology is a rupture of the protective fibrous cap. It can occur in plaques that have distinct morphological features such as large and soft lipid pool with distinct necrotic core and thinning of the fibrous cap in the region of the plaque shoulders. Fibrous caps have considerable metabolic activity. The imbalance between matrix synthesis and matrix degradation thought to be regulated by inflammatory mediators combined with VSMC apoptosis are the key underlying mechanisms of plaque rupture. A second type of vulnerable plaque morphology, known as "plaque erosion", can also lead to a fatal coronary thrombotic event. Plaque erosion is morphologically different from plaque rupture. Eroded plaques do not have fractures in the plaque fibrous cap, only superficial erosion of the intima. The loss of endothelial cells can expose the thrombogenic subendothelial matrix that precipitates thrombus formation. This process could be regulated by inflammatory mediators. The propagation of the acute thrombi for both plaque rupture and plaque erosion events depends on the balance between coagulation and thrombolysis. MI due to a vulnerable plaque is a complex phenomenon that includes: plaque vulnerability, blood vulnerability (hypercoagulation, hypothrombolysis), and heart vulnerability (sensitivity of the heart to ischemia or propensity for arrhythmia).

[0005] Recurrent myocardial infarction can generally be viewed as a severe form of MI progression caused by multiple vulnerable plaques that are able to undergo pre-rupture or a pre-erosive state, coupled with extreme blood coagulability.

[0006] The incidence of MI is still high despite currently available preventive measures and therapeutic intervention. More than 1,500,000 people in the US suffer acute MI each year (many without seeking help due to unrecognized MI), and one third of these people die. The lifetime risk of coronary artery disease events at age 40 years is 42.4% for men (one in two) and 24.9% for women (one in four) (Lloyd-Jones DM; Lancet, 1999 353: 89-92).

[0007] The current diagnosis of MI is based on the levels of troponin I or T that indicate the cardiac muscle progressive necrosis, impaired electrocardiogram (ECG), and detection of abnormal ventricular wall motion or angiographic data (the presence of acute thrombi). However, due to the asymptomatic nature of 25% of acute MIs (absence of atypical chest pain, low ECG sensitivity), a significant portion of MIs are not diagnosed and therefore not treated appropriately (e.g., prevention of recurrent MIs).

[0008] Despite a very high prevalence and lifetime risk of MI, there are no good prognostic markers that can identify an individual with a high risk of vulnerable plaques and justify preventive treatments. MI risk assessment and prognosis is currently done using classic risk factors or the recently introduced Framingham Risk Index. Both of these assessments put a significant weight on LDL levels to justify preventive treatment. However, it is well established that half of all MIs occur in individuals without overt hyperlipidemia. Hence, there is a need for additional risk factors for predicting predisposition to MI.

[0009] Other emerging risk factors are inflammatory biomarkers such as C-reactive protein (CRP), ICAM-1, SAA, TNF α , homocysteine, impaired fasting glucose, new lipid markers (ox LDL, Lp-a, MAD-LDL, etc.) and pro-thrombotic factors (fibrinogen, PAI-1). Despite showing some promise, these markers have significant limitations such as low specificity and low positive predictive value, and the need for multiple reference intervals to be used for different groups of people (e.g., males-females, smokers-non smokers, hormone replacement therapy users, different age groups). These limitations diminish the utility of such markers as independent prognostic markers for MI screening.

[0010] Genetics plays an important role in MI risk. Families with a positive family history of MI account for 14% of the general population, 72% of premature MIs, and 48% of all MIs (Williams R R, Am J Cardiology, 2001; 87:129). In addition, replicated linkage studies have revealed evidence of multiple regions of the genome that are associated with MI and relevant to MI genetic traits, including regions on chromosomes 14, 2, 3 and 7 (Broeckel U, Nature Genetics, 2002; 30: 210; Harrap S, Arterioscler Thromb Vasc Biol, 2002; 22: 874-878, Shearman A, Human Molecular Genetics, 2000, 9; 9,1315-1320), implying that genetic risk factors influence the onset, manifestation, and progression of MI. Recent association studies have identified allelic variants that are associated with acute complications of coronary heart disease, including allelic variants of the ApoE, ApoA5, Lpa, APOCIII, and Klotho genes.

[0011] Genetic markers such as single nucleotide polymorphisms are preferable to other types of biomarkers. Genetic markers that are prognostic for MI can be genotyped early in life and could predict individual response to various risk factors. The combination of serum protein levels and genetic predisposition revealed by genetic analysis of susceptibility genes can provide an integrated assessment of the interaction between genotypes and environmental factors, resulting in synergistically increased prognostic value of diagnostic tests.

[0012] Thus, there is an urgent need for novel genetic markers that are predictive of predisposition to MI, particularly for individuals who are unrecognized as having a predisposition to MI. Such genetic markers may enable prognosis of MI in much larger populations compared with the populations which can currently be evaluated by using existing risk factors and biomarkers. The availability of a genetic test may allow, for example, appropriate preventive treatments for acute coronary events to be provided for susceptible individuals (such preventive treatments may include, for example, statin treatments and statin dose escalation, as well as changes to modifiable risk factors), lowering of the thresholds for ECG and angiography testing, and allow adequate monitoring of informative biomarkers.

[0013] Moreover, the discovery of genetic markers associated with MI will provide novel targets for therapeutic intervention or preventive treatments of MI, and enable the development of new therapeutic agents for treating MI and other cardiovascular disorders.

SNPs

[0014] The genomes of all organisms undergo spontaneous mutation in the course of their continuing evolution, generating variant forms of progenitor genetic sequences (Gusella, Ann. Rev. Biochem. 55, 831-854 (1986)). A variant form may confer an evolutionary advantage or disadvantage relative to a progenitor form or may be neutral. In some instances, a variant form confers an evolutionary advantage to the species and is eventually incorporated into the DNA of many or most members of the species and effectively becomes the progenitor form. Additionally, the effects of a variant form may be both beneficial and detrimental, depending on the circumstances. For example, a heterozygous sickle cell mutation confers resistance to malaria, but a homozygous sickle cell mutation is usually lethal. In many cases, both progenitor and variant forms survive and co-exist in a species population. The coexistence of multiple forms of a genetic sequence gives rise to genetic polymorphisms, including SNPs.

[0015] Approximately 90% of all polymorphisms in the human genome are SNPs. SNPs are single base positions in DNA at which different alleles, or alternative nucleotides, exist in a population. The SNP position (interchangeably referred to herein as SNP, SNP site, or SNP locus) is usually preceded by and followed by highly conserved sequences of the allele (e.g., sequences that vary in less than 1/100 or 1/1000 members of the populations). An individual may be homozygous or heterozygous for an allele at each SNP position. A SNP can, in some instances, be referred to as a "cSNP" to denote that the nucleotide sequence containing the SNP is an amino acid coding sequence.

[0016] A SNP may arise from a substitution of one nucleotide for another at the polymorphic site. Substitutions can be transitions or transversions. A transition is the replacement of one purine nucleotide by another purine nucleotide, or one pyrimidine by another pyrimidine. A transversion is the replacement of a purine by a pyrimidine, or vice versa. A SNP may also be a single base insertion or deletion variant referred to as an "indel" (Weber et al., "Human diallelic insertion/deletion polymorphisms", Am J Hum Genet 2002 Oct;71 (4):854-62).

[0017] A synonymous codon change, or silent mutation/SNP (terms such as "SNP", "polymorphism", "mutation", "mutant", "variation", and "variant" are used herein interchangeably), is one that does not result in a change of amino acid due to the degeneracy of the genetic code. A substitution that changes a codon coding for one amino acid to a codon coding for a different amino acid (i.e., a non-synonymous codon change) is referred to as a missense mutation. A nonsense mutation results in a type of non-synonymous codon change in which a stop codon is formed, thereby leading to premature termination of a polypeptide chain and a truncated protein. A read-through mutation is another type of non-synonymous codon change that causes the destruction of a stop codon, thereby resulting in an extended polypeptide product.

[0018] While SNPs can be bi-, tri-, or tetra- allelic, the vast majority of the SNPs are bi-allelic, and are thus often referred to as "bi-allelic markers", or "di-allelic markers".

[0019] As used herein, references to SNPs and SNP genotypes include individual SNPs and/or haplotypes, which are groups of SNPs that are generally inherited together. Haplotypes can have stronger correlations with diseases or other phenotypic effects compared with individual SNPs, and therefore may provide increased diagnostic accuracy in some cases (Stephens et al. Science 293, 489-493, 20 July 2001).

[0020] Causative SNPs are those SNPs that produce alterations in gene expression or in the expression, structure, and/or function of a gene product, and therefore are most predictive of a possible clinical phenotype. One such class includes SNPs falling within regions of genes encoding a polypeptide product, i.e. cSNPs. These SNPs may result in an alteration of the amino acid sequence of the polypeptide product (i.e., non-synonymous codon changes) and give rise to the expression of a defective or other variant protein. Furthermore, in the case of nonsense mutations, a SNP may lead to premature termination of a polypeptide product. Such variant products can result in a pathological condition, e.g., genetic disease. Examples of genes in which a SNP within a coding sequence causes a genetic disease include sickle cell anemia and cystic fibrosis.

[0021] Causative SNPs do not necessarily have to occur in coding regions; causative SNPs can occur in, for example, any genetic region that can ultimately affect the expression, structure, and/or activity of the protein encoded by a nucleic acid. Such genetic regions include, for example, those involved in transcription, such as SNPs in transcription factor binding domains, SNPs in promoter regions, in areas involved in transcript processing, such as SNPs at intron-exon boundaries that may cause defective splicing, or SNPs in mRNA processing signal sequences such as polyadenylation signal regions. Some SNPs that are not causative SNPs nevertheless are in close association with, and therefore segregate with, a disease-causing sequence. In this situation, the presence of a SNP correlates with the presence of, or predisposition to, or an increased risk in developing the disease. These SNPs, although not causative, are nonetheless also useful for diagnostics, disease predisposition screening, and other uses.

[0022] An association study of a SNP and a specific disorder involves determining the presence or frequency of the SNP allele in biological samples from individuals with the disorder of interest, such as myocardial infarction, and comparing the information to that of controls (i.e., individuals who do not have the disorder; controls may be also referred to as "healthy" or "normal" individuals) who are preferably of similar age and race. The appropriate selection of patients and controls is important to the success of SNP association studies. Therefore, a pool of individuals with well-characterized phenotypes is extremely desirable.

[0023] A SNP may be screened in diseased tissue samples or any biological sample obtained from a diseased individual, and compared to control samples, and selected for its increased (or decreased) occurrence in a specific pathological condition, such as pathologies related to myocardial infarction. Once a statistically significant association is established between one or more SNP(s) and a pathological condition (or other phenotype) of interest, then the region around the SNP can optionally be thoroughly screened to identify the causative genetic locus/sequence(s) (e.g., causative SNP/mutation, gene, regulatory region, etc.) that influences the pathological condition or phenotype. Association studies may be conducted within the general population and are not limited to studies performed on related individuals in affected families (linkage studies).

[0024] Clinical trials have shown that patient response to treatment with pharmaceuticals is often heterogeneous. There is a continuing need to improve pharmaceutical agent design and therapy. In that regard, SNPs can be used to identify patients most suited to therapy with particular pharmaceutical agents (this is often termed "pharmacogenomics"). Similarly, SNPs can be used to exclude patients from certain treatment due to the patient's increased likelihood of developing toxic side effects or their likelihood of not responding to the treatment. Pharmacogenomics can also be used in pharmaceutical research to assist the drug development and selection process. (Linder et al. (1997), Clinical Chemistry, 43, 254; Marshall (1997), Nature Biotechnology, 15, 1249; International Patent Application WO 97/40462, Spectra Biomedical; and Schafer et al. (1998), Nature Biotechnology, 16, 3).

[0025] EP 0 609 059 A2 discloses a method of assessing an individual member from a reference population for the risk of heart disease which comprises determining the leader TaqI and/or kringle TaqI polymorphisms of the Apo(a) gene in a sample of DNA from the individual, and assessing the risk of heart disease by reference to the estimated incidence of heart disease in the reference population of other individuals carrying the same Apo(a) leader TaqI and/or kringle TaqI polymorphisms.

SUMMARY OF THE INVENTION

[0026] The present invention relates to an *in vitro* method for identifying an individual having an altered risk for developing myocardial infarction, comprising detecting the single nucleotide polymorphism (SNP) in the nucleotide sequence of SEQ ID NO: 19038 in said individual's nucleic acids, wherein said individual has an increased risk for developing myocardial infarction due to the presence of C at position 101 of SEQ ID NO: 19038 or the presence of G at position 101 of its complement.

[0027] Further, the present invention relates to the use of an isolated nucleic acid molecule comprising at least 8 contiguous nucleotides of SEQ ID NO: 19038, wherein said at least 8 contiguous nucleotides of SEQ ID NO: 19038 include position 101 of SEQ ID NO: 19038, or the complement thereof, in the method of the present invention.

[0028] The present invention also relates to the use of an isolated nucleic acid molecule that encodes any one of the amino acid sequences in SEQ ID NOS: 845, 846 in the method of the present invention. Also, the present invention relates to the use of an amplified polynucleotide containing the single nucleotide polymorphism (SNP) at position 101 of SEQ ID NO: 19038, or the complement thereof, in the method of the present invention, wherein the amplified polynucleotide is between about 16 and about 1,000 nucleotides in length. Additionally, the present invention relates to the use of an isolated polynucleotide which specifically hybridizes to a nucleic acid molecule containing the single nucleotide polymorphism (SNP) at position 101 of SEQ ID NO: 19038, or the complement thereof, in the method of the present invention.

[0029] Moreover, the present invention relates to the use of a test kit comprising the polynucleotide as used in the method of the present invention, a buffer, and an enzyme in the method of the present invention.

[0030] Finally, the present invention relates to an *in vitro* method of identifying an individual having an altered risk for developing myocardial infarction, comprising detecting the presence of proline (P) or serine (S) at position 157 of the amino acid sequence of SEQ ID NOS:845 or 846, wherein the individual has an increased risk for developing myocardial infarction due to the presence of P at position 157 of the amino acid sequence of SEQ ID NOS:845 or 846.

[0031] The present invention relates to the identification of a SNP that is associated with myocardial infarction (including recurrent myocardial infarction) and related pathologies. The polymorphisms disclosed herein are directly useful as targets for the design of diagnostic reagents and the development of therapeutic agents for use in the diagnosis and treatment of myocardial infarction and related pathologies.

[0032] Based on the identification of a SNP associated with myocardial infarction, the present invention also provides methods of detecting this variant as well as the design and preparation of detection reagents needed to accomplish this task. The invention specifically provides the use of a SNP in genetic sequences involved in myocardial infarction in an *in vitro* method of the present invention. Described herein are variant proteins encoded by nucleic acid molecules containing such SNP, antibodies to the encoded variant proteins, computer-based and data storage systems containing the novel SNP information. The invention provides methods of detecting the SNP in a test sample, methods of identifying individuals who have an altered (i.e., increased or decreased) risk of developing myocardial infarction based on the presence of the SNP disclosed herein or its encoded product. Further, disclosed herein are methods of identifying individuals who are more or less likely to respond to a treatment, methods of screening for compounds useful in the treatment of a disorder associated with a variant gene/protein, compounds identified by these methods, methods of treating disorders mediated by a variant gene/protein, and methods of using the novel SNPs disclosed herein for human identification.

[0033] In Tables 1-2, the present invention provides gene information, transcript sequence (SEQ ID NOS:17-18), encoded amino acid sequences (SEQ ID NOS:845-846), genomic sequence (SEQ ID NO: 17560), transcript-based context sequences (SEQ ID NOS: 2952-2984) and genomic-based context sequences (SEQ ID NOS: 18962-19247) that contain the SNPs disclosed herein, and extensive SNP information that includes observed alleles, allele frequencies, populations/ethnic groups in which alleles have been observed, information about the type of SNP and corresponding functional effect, and; for cSNPs, information about the encoded polypeptide product. The transcript sequences (SEQ ID NOS:1-828), amino acid sequences (SEQ ID NOS:829-1656), genomic sequences (SEQ ID NOS:17,553-18,016), transcript-based SNP context sequences (SEQ ID NOS: 1657-17,552), and genomic-based SNP context sequences (SEQ ID NOS: 18,017-73,085) are also provided in the Sequence Listing.

[0034] In a specific embodiment of the present invention, a naturally-occurring SNP in the human genome is provided. This SNP is associated with myocardial infarction such that it can have a variety of uses in the diagnosis and/or treatment of myocardial infarction. An alternative embodiment of the invention is the use of an amplified polynucleotide, which is produced by amplification of a SNP-containing nucleic acid template in the method of the present invention. In another aspect, described is a variant protein which is encoded by a nucleic acid molecule containing a SNP disclosed herein.

[0035] In yet another embodiment of the invention, the use of a reagent for detecting a SNP in the context of its naturally-occurring flanking nucleotide sequences (which can be, e.g., either DNA or mRNA) in the method of the present invention is provided. In particular, such a reagent may be in the form of, for example, a hybridization probe or an amplification primer that is useful in the specific detection of a SNP of interest. In an alternative aspect disclosed herein,

a protein detection reagent is used to detect a variant protein which is encoded by a nucleic acid molecule containing a SNP disclosed herein. A preferred example of a protein detection reagent is an antibody or an antigen-reactive antibody fragment.

[0036] Also provided in the invention is the use of kits comprising SNP detection reagents, in the method of the present invention and methods for detecting the SNP disclosed herein by employing detection reagents. In a specific embodiment, the present invention provides for a method of identifying an individual having an increased or decreased risk of developing myocardial infarction by detecting the presence or absence of a SNP allele disclosed herein. In another aspect disclosed herein, a method for diagnosis of myocardial infarction by detecting the presence or absence of a SNP allele disclosed herein is provided.

[0037] The nucleic acid molecules for use in the method of the invention can be inserted in an expression vector, such as to produce a variant protein in a host cell. Thus, the disclosed herein is a vector comprising a SNP-containing nucleic acid molecule, genetically-engineered host cells containing the vector, and methods for expressing a recombinant variant protein using such host cells. In another aspect disclosed herein, the host cells, SNP-containing nucleic acid molecules, and/or variant proteins can be used as targets in a method for screening and identifying therapeutic agents or pharmaceutical compounds useful in the treatment of myocardial infarction.

[0038] An aspect disclosed herein is a method for treating myocardial infarction in a human subject wherein said human subject harbors a gene, transcript, and/or encoded protein identified in Tables 1-2, which method comprises administering to said human subject a therapeutically or prophylactically effective amount of one or more agents counteracting the effects of the disease, such as by inhibiting (or stimulating) the activity of the gene, transcript, and/or encoded protein identified in Tables 1-2.

[0039] Another aspect disclosed herein is a method for identifying an agent useful in therapeutically or prophylactically treating myocardial infarction in a human subject wherein said human subject harbors a gene, transcript, and/or encoded protein identified in Tables 1-2, which method comprises contacting the gene, transcript, or encoded protein with a candidate agent under conditions suitable to allow formation of a binding complex between the gene, transcript, or encoded protein and the candidate agent and detecting the formation of the binding complex, wherein the presence of the complex identifies said agent.

[0040] Another aspect disclosed herein is a method for treating myocardial infarction in a human subject, which method comprises:

- (i) determining that said human subject harbors a gene, transcript, and/or encoded protein identified in Tables 1-2, and
- (ii) administering to said subject a therapeutically or prophylactically effective amount of one or more agents counteracting the effects of the disease.

[0041] Many other uses and advantages of the present invention will be apparent to those skilled in the art upon review of the detailed description of the preferred embodiments herein. Solely for clarity of discussion, the invention is described in the sections below by way of non-limiting examples.

DESCRIPTION OF THE SEQUENCE LISTING

[0042]

The Sequence Listing provides the transcript sequences (SEQ ID NOS:1-828) and protein sequences (SEQ ID NOS:829-1656), and genomic sequences (SEQ ID NOS:17,553-18,016), for each myocardial infarction-associated gene that contains one or more SNPs of the present invention. Also provided in the Sequence Listing are context sequences flanking each SNP, including both transcript-based context sequences (SEQ ID NOS: 1657-17,552) and genomic-based context sequences (SEQ ID NOS:18,017-73,085). The context sequences generally provide 100bp upstream (5') and 100bp downstream (3') of each SNP, with the SNP in the middle of the context sequence, for a total of 200bp of context sequence surrounding each SNP.

DESCRIPTION OF TABLE 1 AND TABLE 2

[0043] Table 1 and Table 2 disclose the SNP and associated gene/transcript/protein information used in the method of the present invention. For gene No:8, Table 1 and Table 2 each provide a header containing gene/transcript/protein information, followed by a transcript and protein sequence (in Table 1) or genomic sequence (in Table 2), and then SNP information regarding the SNP used in the method of the present invention found in that gene/transcript.

[0044] NOTE: The SNP is included in both Table 1 and Table 2; Table 1 presents the SNPs relative to their transcript sequences and encoded protein sequences. The SNP can readily be cross-referenced between Tables based on its hCV identification numbers.

[0045] The gene/transcript/protein information includes:

- a gene number (1 through n, where n = the total number of genes in the Table)
- a Celera hCG and UID internal identification numbers for the gene
- 5 - a Celera hCT and UID internal identification numbers for the transcript (Table 1 only)
- a public Genbank accession number (e.g., RefSeq NM number) for the transcript (Table 1 only)
- a Celera hCP and UID internal identification numbers for the protein encoded by the hCT transcript (Table 1 only)
- a public Genbank accession number (e.g., RefSeq NP number) for the protein (Table 1 only)
- an art-known gene symbol
- 10 - an art-known gene/protein name
- Celera genomic axis position (indicating start nucleotide position-stop nucleotide position)
- the chromosome number of the chromosome on which the gene is located
- an OMIM (Online Mendelian Inheritance in Man; Johns Hopkins University/NCBI) public reference number for obtaining further information regarding the medical significance of each gene
- 15 - alternative gene/protein name(s) and/or symbol(s) in the OMIM entry

[0046] NOTE: Due to the presence of alternative splice forms, multiple transcript/protein entries can be provided for a single gene entry in Table 1; i.e., for a single Gene Number, multiple entries may be provided in series that differ in their transcript/protein information and sequences.

20 **[0047]** Following the gene/transcript/protein information is a transcript sequence and protein sequence (in Table 1), or a genomic sequence (in Table 2), for each gene, as follows:

- transcript sequence (Table 1 only) (corresponding to SEQ ID NOS: 17-18 of the Sequence Listing), with SNPs identified by their IUB codes (transcript sequences can include 5' UTR, protein coding, and 3' UTR regions). (NOTE: If there are differences between the nucleotide sequence of the hCT transcript and the corresponding public transcript sequence identified by the Genbank accession number, the hCT transcript sequence (and encoded protein) is provided, unless the public sequence is a RefSeq transcript sequence identified by an NM number, in which case the RefSeq NM transcript sequence (and encoded protein) is provided. However, whether the hCT transcript or RefSeq NM transcript is used as the transcript sequence, the disclosed SNPs are represented by their IUB codes within the transcript.)
- 25 - the encoded protein sequence (Table 1 only) (corresponding to SEQ ID NOS:845-846 of the Sequence Listing)
- the genomic sequence of the gene (Table 2 only), including 6kb on each side of the gene boundaries (i.e., 6kb on the 5' side of the gene plus 6kb on the 3' side of the gene) (corresponding to SEQ ID NO: 17560 of the Sequence Listing).
- 30
- 35

[0048] NOTE: The transcript, protein, and transcript-based SNP context sequences are provided in both Table 1 and in the Sequence Listing. The genomic and genomic-based SNP context sequences are provided in both Table 2 and in the Sequence Listing. SEQ ID NOS are indicated in Table 1 for each transcript sequence (SEQ ID NOS: 17-18), protein sequence (SEQ ID NOS: 845-846), and transcript-based SNP context sequence (SEQ ID NOS: 2952-2984), and SEQ ID NOS are indicated in Table 2 for each genomic sequence (SEQ ID NO:17560), and genomic-based SNP context sequence (SEQ ID NOS:18962-19247).

[0049] The SNP information includes:

- 45 - context sequence (taken from the transcript sequence in Table 1, and taken from the genomic sequence in Table 2) with the SNP represented by its IUB code, including 100 bp upstream (5') of the SNP position plus 100 bp downstream (3') of the SNP position (the transcript-based SNP context sequences in Table 1 are provided in the Sequence Listing as SEQ ID NOS: 2952-2984; the genomic-based SNP context sequences in Table 2 are provided in the Sequence Listing as SEQ ID NOS: 18962-19247).
- 50 - Celera hCV internal identification number for the SNP (in some instances, an "hDV" number is given instead of an "hCV" number)
- SNP position [position of the SNP within the given transcript sequence (Table 1) or within the given genomic sequence (Table 2)]
- SNP source (may include any combination of one or more of the following five codes, depending on which internal sequencing projects and/or public databases the SNP has been observed in: "Applera" = SNP observed during the re-sequencing of genes and regulatory regions of 39 individuals, "Celera" = SNP observed during shotgun sequencing and assembly of the Celera human genome sequence, "Celera Diagnostics" = SNP observed during re-sequencing of nucleic acid samples from individuals who have myocardial infarction or a related pathology, "dbSNP" = SNP
- 55

observed in the dbSNP public database, "HGBASE" = SNP observed in the HGBASE public database, "HGMD" = SNP observed in the Human Gene Mutation Database (HGMD) public database) (NOTE: multiple "Applera" source entries for a single SNP indicate that the same SNP was covered by multiple overlapping amplification products and the re-sequencing results (e.g., observed allele counts) from each of these amplification products is being provided)

- Population/allele/allele count information in the format of [population1(allele1,count|allele2,count) population2(allele1,count|allele2,count) total (allele1,total count|allele2,total count)]. The information in this field includes populations/ethnic groups in which particular SNP alleles have been observed ("cau" = Caucasian, "his" = Hispanic, "chn" = Chinese, and "afr" = African-American, "jpn" = Japanese, "ind" = Indian, "mex" = Mexican, "ain" = "American Indian, "cra" = Celera donor, "no_pop" = no population information available), identified SNP alleles, and observed allele counts (within each population group and total allele counts), where available ["-" in the allele field represents a deletion allele of an insertion/deletion ("indel") polymorphism (in which case the corresponding insertion allele, which may be comprised of one or more nucleotides, is indicated in the allele field on the opposite side of the "|"); "-" in the count field indicates that allele count information is not available].

[0050] NOTE: For SNPs of "Applera" SNP source, genes/regulatory regions of 39 individuals (20 Caucasians and 19 African Americans) were re-sequenced and, since each SNP position is represented by two chromosomes in each individual (with the exception of SNPs on X and Y chromosomes in males, for which each SNP position is represented by a single chromosome), up to 78 chromosomes were genotyped for each SNP position. Thus, the sum of the African-American ("afr") allele counts is up to 38, the sum of the Caucasian allele counts ("cau") is up to 40, and the total sum of all allele counts is up to 78.

[0051] (NOTE: semicolons separate population/allele/count information corresponding to each indicated SNP source; i.e., if four SNP sources are indicated, such as "Celera", "dbSNP", "HGBASE", and "HGMD", then population/allele/count information is provided in four groups which are separated by semicolons and listed in the same order as the listing of SNP sources, with each population/allele/count information group corresponding to the respective SNP source based on order; thus, in this example, the first population/allele/count information group would correspond to the first listed SNP source (Celera) and the third population/allele/count information group separated by semicolons would correspond to the third listed SNP source (HGBASE); if population/allele/count information is not available for any particular SNP source, then a pair of semicolons is still inserted as a place-holder in order to maintain correspondence between the list of SNP sources and the corresponding listing of population/allele/count information)

- SNP type (e.g., location within gene/transcript and/or predicted functional effect) ["MIS-SENSE MUTATION" = SNP causes a change in the encoded amino acid (i.e., a non-synonymous coding SNP); "SILENT MUTATION" = SNP does not cause a change in the encoded amino acid (i.e., a synonymous coding SNP); "STOP CODON MUTATION" = SNP is located in a stop codon; "NONSENSE MUTATION" = SNP creates or destroys a stop codon; "UTR 5" = SNP is located in a 5' UTR of a transcript; "UTR 3" = SNP is located in a 3' UTR of a transcript; "PUTATIVE UTR 5" = SNP is located in a putative 5' UTR; "PUTATIVE UTR 3" = SNP is located in a putative 3' UTR; "DONOR SPLICE SITE" = SNP is located in a donor splice site (5' intron boundary); "ACCEPTOR SPLICE SITE" = SNP is located in an acceptor splice site (3' intron boundary); "CODING REGION" = SNP is located in a protein-coding region of the transcript; "EXON" = SNP is located in an exon; "INTRON" = SNP is located in an intron; "hmCS" = SNP is located in a human-mouse conserved segment; "TFBS" = SNP is located in a transcription factor binding site; "UNKNOWN" = SNP type is not defined; "INTERGENIC" = SNP is intergenic, i.e., outside of any gene boundary]
- Protein coding information (Table 1 only), where relevant, in the format of [protein SEQ ID NO:#, amino acid position, (amino acid-1, codon1) (amino acid-2, codon2)]. The information in this field includes SEQ ID NO of the encoded protein sequence, position of the amino acid residue within the protein identified by the SEQ ID NO that is encoded by the codon containing the SNP, amino acids (represented by one-letter amino acid codes) that are encoded by the alternative SNP alleles (in the case of stop codons, "X" is used for the one-letter amino acid code), and alternative codons containing the alternative SNP nucleotides which encode the amino acid residues (thus, for example, for missense mutation-type SNPs, at least two different amino acids and at least two different codons are generally indicated; for silent mutation-type SNPs, one amino acid and at least two different codons are generally indicated, etc.). In instances where the SNP is located outside of a protein-coding region (e.g., in a UTR region), "None" is indicated following the protein SEQ ID NO.

DESCRIPTION OF TABLE 5

[0052] Table 5 provides sequences (SEQ ID NOS:73557, 73557, 73559) of primers that have been synthesized and used in the laboratory to carry out allele-specific PCR reactions in order to assay the SNPs disclosed in Tables 6-7 during the course of myocardial infarction association studies.

[0053] Table 5 provides the following:

- the column labeled "hCV" provides an hCV identification number for each SNP site
- the column labeled "Alleles" designates the two alternative alleles at the SNP site identified by the hCV identification number that are targeted by the allele-specific primers (the allele-specific primers are shown as "Sequence A" and "Sequence B" in each row)
- the column labeled "Sequence A (allele-specific primer)" provides an allele-specific primer that is specific for the first allele designated in the "Alleles" column
- the column labeled "Sequence B (allele-specific primer)" provides an allele-specific primer that is specific for the second allele designated in the "Alleles" column
- the column labeled "Sequence C (common primer)" provides a common primer that is used in conjunction with each of the allele-specific primers (the "Sequence A" primer and the "Sequence B" primer) and which hybridizes at a site away from the SNP position.

[0054] All primer sequences are given in the 5' to 3' direction.

[0055] Each of the alleles designated in the "Alleles" column matches the 3' nucleotide of the allele-specific primer that is specific for that allele. Thus, the first allele designated in the "Alleles" column matches the 3' nucleotide of the "Sequence A" primer, and the second allele designated in the "Alleles" column matches the 3' nucleotide of the "Sequence B" primer.

DESCRIPTION OF TABLE 6 AND TABLE 7

[0056] Tables 6 and 7 provide results of statistical analyses for SNPs disclosed in Tables 1, 2, and 5 (SNPs can be cross-referenced between Tables based on their hCV identification numbers). Table 6 provides statistical results for association of SNPs with myocardial infarction, and Table 7 provides statistical results for association of SNPs with recurrent myocardial infarction (RMI). The statistical results shown in Tables 6-7 provide support for the association of these SNPs with MI (Table 6) and/or RMI (Table 7). For example, the statistical results provided in Tables 6-7 show that the association of these SNPs with MI and/or RMI is supported by p-values < 0.05 in at least one of three genotypic association tests and/or an allelic association test. Moreover, in general, the SNPs identified in Tables 6-7 are SNPs for which their association with MI and/or RMI has been replicated by virtue of being significant in at least two independently collected sample sets, which further verifies the association of these SNPs with MI and/or RMI. Furthermore, results of stratification-based analyses are also provided; stratified analysis can, for example, enable increased prediction of MI and/or RMI risk via interaction between conventional risk factors (stratum) and SNPs.

[0057] NOTE: SNPs can be cross-referenced between Tables 1, 2, 5, 6, and 7 based on the hCV identification number of each SNP.

TABLE 6 (SNP association with Myocardial Infarction) Description of column headings for Table 6:

Table 6 column heading	Definition
Marker	Internal hCV identification number for the SNP that is tested
Study	Sample set used in the analysis
Stratification	Indicates if the analysis of the dataset was done on a substratum (stratifications are described below)
Strata	Indicates what substratum was used in the analysis (strata are described below)
Status	Identifies the inclusion/exclusion criteria for cases and controls (described below)
Allele1	Nucleotide (allele) of the tested SNP for which statistics are being reported
Case Allele1 frq	Allele frequency of Allele1 in cases
Control Allele1 frq	Allele frequency of Allele1 in controls
Allelic p-value	Result of the Fisher exact test for allelic association
Dom p-value	Result of the asymptotic chi square test for dominant genotypic association
Rec p-value	Result of the asymptotic chi square test for recessive genotypic association
OR	Allelic odds ratio

(continued)

Table 6 column heading	Definition
OR 95%CI L	Lower limit of 95% confidence interval of the allelic odds ratio
OR 95%CI U	Upper limit of 95% confidence interval of the allelic odds ratio

[0058] Definition of entries in the "Stratification" and "Strata" columns (Table 6) for stratification-based analyses:

Stratification	Strata	Definition
no	ALL	All individuals
BMI_GE27	L /H	Individuals with body mass index lower/higher than 27
HTN	Y/N	Individuals with/without (Y/N) history of hypertension
SEX	M/F	Gender male/female
AGE_LT60	Y/O	Individuals younger/older than 60 years
SMOKE	Y/N	Individuals that have/do not have history of smoking

[0059] Definition of entries in the "Status" column (Table 6):

Status	Definition
LT60MI_60TO75noMI	MI cases younger than 60 compared to controls between the ages of 60 to 75
LT60MI_GT75noMI	MI cases younger than 60 compared to controls older than 75
MI-GT75noMI	MI cases compared to controls older than 75
MI_LT75noMI	MI cases compared to controls younger than 75
MI_noASD	Cases with history of MI, controls with no history of atherosclerotic disease
MI-noMI	Cases with history of MI, controls with no history of MI
MI_YOUNGOLD_noASD	Cases with history of MI under the age of 60 controls with no history of atherosclerotic disease over the age of 60
MI_YOUNGOLD_noMI	Cases with history of MI under the age of 60 controls with no history of MI over the age of 60
YoungMI_GT75noASD	Cases with history of MI under the age of 60 controls with no history of atherosclerotic disease over the age of 75

TABLE 7 (SNP association with Recurrent Myocardial Infarction) Description of column headings for Table 7:

Table 7 column heading	Definition
Gene	Locus Link HUGO approved gene symbol
Marker	Internal hCV identification number for the tested SNP
Sample Set	Sample Set used in the analysis (CARE, Pre-CARE or WGS_S0012)
p-value	Result of the asymptotic chi square test for allelic association, dominant genotypic association, recessive genotypic association, or the allelic, dominant, or recessive p-value of the stratified analysis
OR	odds ratio
95%CI	95% confidence interval of the given odds ratio
Case _Freq	Allele frequency of minor allele in cases
Control _Freq	Allele frequency of minor allele in controls

(continued)

Table 7 column heading	Definition
Allele1	Nucleotide (allele) of the tested SNP for which statistics are being reported
Mode	The mode of inheritance
Strata	Indicates if the analysis of the dataset was based on a substratum such as gender, age, BMI, Hypertension, Fasting Glucose levels, etc. (strata are described below)

[0060] Definition of entries in the "Strata" column (Table 7) for stratification-based analyses:

Stratum	Definition
BMI_TERTILE_1	Individuals in the lowest tertile of body mass index
BMI_TERTILE_2	Individuals in the middle tertile of body mass index
BMI_TERTILE_3	Individuals in the highest tertile of body mass index
PLACEBO	Patients who were in placebo arm of the CARE trail
PRAVASTATIN	Patients who were in Pravastatin arm of the CARE trail
MALE	Only males
FEMALE	Only females
HYPERTEN_1	Individuals with history of Hypertension
HYPERTEN_0	Individuals without history of Hypertension
GLUCOSE_TERTILE_1	Individuals in a lowest tertile of Fasting Glucose levels
GLUCOSE_TERTILE_2	Individuals in a middle tertile of Fasting Glucose levels
GLUCOSE_TERTILE_3	Individuals in a highest tertile of Fasting Glucose levels
AGE_TERTILE_1	Individuals in a lowest tertile of age (premature MI)
AGE_TERTILE_2	Individuals in a middle tertile of age
AGE_TERTILE_3	Individuals in a highest tertile age
EVERSMOKED_0	Individuals wh0 never smoked
EVERSMOKED_1	Former smokers
EVERSMOKED_2	Current smokers
FMHX_CHD_1	Individuals with family history of CHD
FMHX_CHD_0	Individuals without family history of CHD

DESCRIPTION OF THE FIGURE

[0061] Figure 1 provides a diagrammatic representation of a computer-based discovery system containing the SNP information of the present invention in computer readable form.

DETAILED DESCRIPTION OF THE INVENTION

[0062] The present invention relates to an *in vitro* method for identifying an individual having an altered risk for developing myocardial infarction, comprising detecting the single nucleotide polymorphism (SNP) in the nucleotide sequence of SEQ ID NO: 19038 in said individual's nucleic acids, wherein said individual has an increased risk for developing myocardial infarction due to the presence of C at position 101 of SEQ ID NO: 19038 or the presence of G at position 101 of its complement.

[0063] Further, the present invention relates to the use of an isolated nucleic acid molecule comprising at least 8 contiguous nucleotides of SEQ ID NO: 19038, wherein said at least 8 contiguous nucleotides of SEQ ID NO: 19038 include position 101 of SEQ ID NO: 19038, or the complement thereof, in the method of the present invention.

[0064] The present invention also relates to the use of an isolated nucleic acid molecule that encodes any one of the amino acid sequences in SEQ ID NOS: 845, 846 in the method of the present invention. Also, the present invention relates to the use of an amplified polynucleotide containing the single nucleotide polymorphism (SNP) at position 101 of SEQ ID NO: 19038, or the complement thereof, in the method of the present invention, wherein the amplified polynucleotide is between about 16 and about 1,000 nucleotides in length. Additionally, the present invention relates to the use of an isolated polynucleotide which specifically hybridizes to a nucleic acid molecule containing the single nucleotide polymorphism (SNP) at position 101 of SEQ ID NO: 19038, or the complement thereof, in the method of the present invention.

[0065] Moreover, the present invention relates to the use of a test kit comprising the polynucleotide as used in the method of the present invention, a buffer, and an enzyme in the method of the present invention.

[0066] Finally, the present invention relates to an *in vitro* method of identifying an individual having an altered risk for developing myocardial infarction, comprising detecting the presence of proline (P) or serine (S) at position 157 of the amino acid sequence of SEQ ID NOS:845 or 846, wherein the individual has an increased risk for developing myocardial infarction due to the presence of P at position 157 of the amino acid sequence of SEQ ID NOS:845 or 846.

[0067] Disclosed herein are SNPs associated with myocardial infarction (including recurrent myocardial infarction), nucleic acid molecules containing SNPs, methods and reagents for the detection of the SNPs disclosed herein, uses of these SNPs for the development of detection reagents, and assays or kits that utilize such reagents. The myocardial infarction-associated SNPs disclosed herein are useful for diagnosing, screening for, and evaluating predisposition to myocardial infarction and related pathologies in humans. Furthermore, such SNPs and their encoded products are useful targets for the development of therapeutic agents.

[0068] A large number of SNPs have been identified from re-sequencing DNA from 39 individuals, and they are indicated as "Applera" SNP source in Tables 1-2. Their allele frequencies observed in each of the Caucasian and African-American ethnic groups are provided. Additional SNPs included herein were previously identified during shotgun sequencing and assembly of the human genome, and they are indicated as "Celera" SNP source in Tables 1-2. Furthermore, the information provided in Table 1-2, particularly the allele frequency information obtained from 39 individuals and the identification of the precise position of each SNP within each gene/transcript, allows haplotypes (i.e., groups of SNPs that are co-inherited) to be readily inferred. The present disclosure encompasses SNP haplotypes, as well as individual SNPs.

[0069] Thus, the present disclosure describes individual SNPs associated with myocardial infarction, as well as combinations of SNPs and haplotypes in genetic regions associated with myocardial infarction, polymorphic/variant transcript sequences (SEQ ID NOS:17-18) and genomic sequences (SEQ ID NO:17560) containing SNPs, encoded amino acid sequences (SEQ ID NOS: 845-846), and both transcript-based SNP context sequences (SEQ ID NOS: 2952-2984) and genomic-based SNP context sequences (SEQ ID NOS:18962-19247) (transcript sequences, protein sequences, and transcript-based SNP context sequences are provided in Table 1 and the Sequence Listing; genomic sequences and genomic-based SNP context sequences are provided in Table 2 and the Sequence Listing), methods of detecting these polymorphisms in a test sample, methods of determining the risk of an individual of having or developing myocardial infarction, methods of screening for compounds useful for treating disorders associated with a variant gene/protein such as myocardial infarction, compounds identified by these screening methods, methods of using the disclosed SNPs to select a treatment strategy, methods of treating a disorder associated with a variant gene/protein (i.e., therapeutic methods), and methods of using the SNPs disclosed herein for human identification.

[0070] Disclosed herein are novel SNPs associated with myocardial infarction, as well as SNPs that were previously known in the art, but were not previously known to be associated with myocardial infarction. Accordingly, disclosed herein are novel compositions and methods based on the novel SNPs disclosed herein, and also describes novel methods of using the known, but previously unassociated, SNPs in methods relating to myocardial infarction (e.g., for diagnosing myocardial infarction, etc.). In Tables 1-2, the SNP is identified based on the public database in which it has been observed, which is indicated as one or more of the following SNP types: "dbSNP" = SNP observed in dbSNP, "HGBASE" = SNP observed in HGBASE, and "HGMD" = SNP observed in the Human Gene Mutation Database (HGMD).

[0071] Particular SNP alleles used in the method of the present invention can be associated with either an increased risk of having or developing myocardial infarction, or a decreased risk of having or developing myocardial infarction. SNP alleles that are associated with a decreased risk of having or developing myocardial infarction may be referred to as "protective" alleles, and SNP alleles that are associated with an increased risk of having or developing myocardial infarction may be referred to as "susceptibility" alleles or "risk factors". Thus, whereas certain SNPs (or their encoded products) can be assayed to determine whether an individual possesses a SNP allele that is indicative of an increased risk of having or developing myocardial infarction (i.e., a susceptibility allele), other SNPs (or their encoded products) can be assayed to determine whether an individual possesses a SNP allele that is indicative of a decreased risk of having or developing myocardial infarction (i.e., a protective allele). Similarly, particular SNP alleles used in the method of the present invention can be associated with either an increased or decreased likelihood of responding to a particular treatment or therapeutic compound, or an increased or decreased likelihood of experiencing toxic effects from a particular

treatment or therapeutic compound. The term "altered" may be used herein to encompass either of these two possibilities (e.g., an increased or a decreased risk/likelihood).

[0072] Those skilled in the art will readily recognize that nucleic acid molecules may be double-stranded molecules and that reference to a particular site on one strand refers, as well, to the corresponding site on a complementary strand. In defining a SNP position, SNP allele, or nucleotide sequence, reference to an adenine, a thymine (uridine), a cytosine, or a guanine at a particular site on one strand of a nucleic acid molecule also defines the thymine (uridine), adenine, guanine, or cytosine (respectively) at the corresponding site on a complementary strand of the nucleic acid molecule. Thus, reference may be made to either strand in order to refer to a particular SNP position, SNP allele, or nucleotide sequence. Probes and primers, may be designed to hybridize to either strand and SNP genotyping methods disclosed herein may generally target either strand. Throughout the specification, in identifying a SNP position, reference is generally made to the protein-encoding strand, only for the purpose of convenience.

[0073] References to variant peptides, polypeptides, or proteins disclosed herein include peptides, polypeptides, proteins, or fragments thereof, that contain at least one amino acid residue that differs from the corresponding amino acid sequence of the art-known peptide/polypeptide/protein (the art-known protein may be interchangeably referred to as the "wild-type", "reference", or "normal" protein). Such variant peptides/polypeptides/proteins can result from a codon change caused by a nonsynonymous nucleotide substitution at a protein-coding SNP position (i.e., a missense mutation) disclosed herein. Variant peptides/polypeptides/proteins described herein can also result from a nonsense mutation, i.e. a SNP that creates a premature stop codon, a SNP that generates a read-through mutation by abolishing a stop codon, or due to any SNP disclosed herein that otherwise alters the structure, function/activity, or expression of a protein, such as a SNP in a regulatory region (e.g. a promoter or enhancer) or a SNP that leads to alternative or defective splicing, such as a SNP in an intron or a SNP at an exon/intron boundary. As used herein, the terms "polypeptide", "peptide", and "protein" are used interchangeably.

ISOLATED NUCLEIC ACID MOLECULES AND SNP DETECTION REAGENTS & KITS

[0074] Tables 1 and 2 provide a variety of information about the SNP used in the method of the present invention that is associated with myocardial infarction, including the transcript sequences (SEQ ID NOS: 17-18), genomic sequences (SEQ ID NO:17560), and protein sequences (SEQ ID NOS:845-846) of the encoded gene products (with the SNPs indicated by IUB codes in the nucleic acid sequences). In addition, Tables 1 and 2 include SNP context sequences, which generally include 100 nucleotide upstream (5') plus 100 nucleotides downstream (3') of each SNP position (SEQ ID NOS:2952-2984 correspond to transcript-based SNP context sequences disclosed in Table 1, and SEQ ID NOS:18962-19247 correspond to genomic-based context sequences disclosed in Table 2), the alternative nucleotides (alleles) at each SNP position, and additional information about the variant where relevant, such as SNP type (coding, missense, splice site, UTR, etc.), human populations in which the SNP was observed, observed allele frequencies, information about the encoded protein, etc.

Isolated Nucleic Acid Molecules

[0075] The present invention provides the use of isolated nucleic acid molecules that contain the SNP disclosed in Table 1 and/or Table 2 in the method of the present invention. Isolated nucleic acid molecules containing the SNP disclosed in Tables 1 and 2 may be interchangeably referred to throughout the present text as "SNP-containing nucleic acid molecules". Isolated nucleic acid molecules may optionally encode a full-length variant protein or fragment thereof. The isolated nucleic acid molecules for use in the method of the present invention also include probes and primers (which are described in greater detail below in the section entitled "SNP Detection Reagents"), which may be used for assaying the disclosed SNPs, and isolated full-length genes, transcripts, cDNA molecules, and fragments thereof, which may be used for such purposes as expressing an encoded protein.

[0076] As used herein, an "isolated nucleic acid molecule" generally is one that contains the SNP used in the method of the present invention or one that hybridizes to such molecule such as a nucleic acid with a complementary sequence, and is separated from most other nucleic acids present in the natural source of the nucleic acid molecule. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule containing the SNP used in the method of the present invention, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. A nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered "isolated". Nucleic acid molecules present in non-human transgenic animals, which do not naturally occur in the animal, are also considered "isolated". For example, recombinant DNA molecules contained in a vector are considered "isolated". Further examples of "isolated" DNA molecules include recombinant DNA molecules maintained in heterologous host cells, and purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include in vivo or in vitro RNA transcripts of the isolated SNP-containing DNA molecules for use in the method of the present invention. Isolated nucleic acid molecules for use in the method of the

present invention further include such molecules produced synthetically.

[0077] Generally, an isolated SNP-containing nucleic acid molecule comprises one or more SNP positions disclosed herein with flanking nucleotide sequences on either side of the SNP positions. A flanking sequence can include nucleotide residues that are naturally associated with the SNP site and/or heterologous nucleotide sequences. Preferably the flanking sequence is up to about 500, 300, 100, 60, 50, 30, 25, 20, 15, 10, 8, or 4 nucleotides (or any other length in-between) on either side of a SNP position, or as long as the full-length gene or entire protein-coding sequence (or any portion thereof such as an exon), especially if the SNP-containing nucleic acid molecule is to be used to produce a protein or protein fragment.

[0078] For full-length genes and entire protein-coding sequences, a SNP flanking sequence can be, for example, up to about 5KB, 4KB, 3KB, 2KB, 1KB on either side of the SNP. Furthermore, in such instances, the isolated nucleic acid molecule comprises exonic sequences (including protein-coding and/or non-coding exonic sequences), but may also include intronic sequences. Thus, any protein coding sequence may be either contiguous or separated by introns. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences and is of appropriate length such that it can be subjected to the specific manipulations or uses described herein such as recombinant protein expression, preparation of probes and primers for assaying the SNP position, and other uses specific to the SNP-containing nucleic acid sequences.

[0079] An isolated SNP-containing nucleic acid molecule can comprise, for example, a full-length gene or transcript, such as a gene isolated from genomic DNA (e.g., by cloning or PCR amplification), a cDNA molecule, or an mRNA transcript molecule. Polymorphic transcript sequences are provided in Table 1 and in the Sequence Listing (SEQ ID NOS: 1-828), and polymorphic genomic sequences are provided in Table 2 and in the Sequence Listing (SEQ ID NOS: 17,553-18,016). Furthermore, fragments of such full-length genes and transcripts that contain one or more SNPs disclosed herein are also described herein, and such fragments may be used, for example, to express any part of a protein, such as a particular functional domain or an antigenic epitope.

[0080] Thus, the present invention also encompasses the use of fragments of the nucleic acid sequences provided in Tables 1-2 (transcript sequences are provided in Table 1 as SEQ ID NOS: 17-18, genomic sequences are provided in Table 2 as SEQ ID NO: 17560, transcript-based SNP context sequences are provided in Table 1 as SEQ ID NO: 2952-2984, and genomic-based SNP context sequences are provided in Table 2 as SEQ ID NO: 18962-19247) and their complements in the method of the present invention. A fragment typically comprises a contiguous nucleotide sequence at least about 8 or more nucleotides, more preferably at least about 12 or more nucleotides, and even more preferably at least about 16 or more nucleotides. Further, a fragment could comprise at least about 18, 20, 22, 25, 30, 40, 50, 60, 100, 250 or 500 (or any other number in-between) nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope-bearing regions of a variant peptide or regions of a variant peptide that differ from the normal/wild-type protein, or can be useful as a polynucleotide probe or primer. Such fragments can be isolated using the nucleotide sequences provided in Table 1 and/or Table 2 for the synthesis of a polynucleotide probe. A labeled probe can then be used, for example, to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in amplification reactions, such as for purposes of assaying one or more SNPs sites or for cloning specific regions of a gene.

[0081] An isolated nucleic acid molecule for use in the method of the present invention further encompasses a SNP-containing polynucleotide that is the product of any one of a variety of nucleic acid amplification methods, which are used to increase the copy numbers of a polynucleotide of interest in a nucleic acid sample. Such amplification methods are well known in the art, and they include but are not limited to, polymerase chain reaction (PCR) (U.S. Patent Nos. 4,683,195; and 4,683,202; PCR Technology: Principles and Applications for DNA Amplification, ed. H.A. Erlich, Freeman Press, NY, NY, 1992), ligase chain reaction (LCR) (Wu and Wallace, Genomics 4:560, 1989; Landegren et al., Science 241:1077, 1988), strand displacement amplification (SDA) (U.S. Patent Nos. 5,270,184; and 5,422,252), transcription-mediated amplification (TMA) (U.S. Patent No. 5,399,491), linked linear amplification (LLA) (U.S. Patent No. 6,027,923), and the like, and isothermal amplification methods such as nucleic acid sequence based amplification (NASBA), and self-sustained sequence replication (Guatelli et al., Proc. Natl. Acad. Sci. USA 87: 1874, 1990). Based on such methodologies, a person skilled in the art can readily design primers in any suitable regions 5' and 3' to a SNP disclosed herein. Such primers may be used to amplify DNA of any length so long that it contains the SNP of interest in its sequence.

[0082] As used herein, an "amplified polynucleotide" for use in the method of the invention is a SNP-containing nucleic acid molecule whose amount has been increased at least two fold by any nucleic acid amplification method performed *in vitro* as compared to its starting amount in a test sample. In other preferred embodiments, an amplified polynucleotide is the result of at least ten fold, fifty fold, one hundred fold, one thousand fold, or even ten thousand fold increase as compared to its starting amount in a test sample. In a typical PCR amplification, a polynucleotide of interest is often amplified at least fifty thousand fold in amount over the unamplified genomic DNA, but the precise amount of amplification needed for an assay depends on the sensitivity of the subsequent detection method used.

[0083] Generally, an amplified polynucleotide is at least about 16 nucleotides in length. More typically, an amplified

polynucleotide is at least about 20 nucleotides in length. In a preferred embodiment, an amplified polynucleotide is at least about 30 nucleotides in length. In a more preferred embodiment, an amplified polynucleotide is at least about 32, 40, 45, 50, or 60 nucleotides in length. In yet another preferred embodiment of the invention, an amplified polynucleotide is at least about 100, 200, or 300 nucleotides in length. While the total length of an amplified polynucleotide of the invention can be as long as an exon, an intron or the entire gene where the SNP of interest resides, an amplified product is typically no greater than about 1,000 nucleotides in length (although certain amplification methods may generate amplified products greater than 1000 nucleotides in length). More preferably, an amplified polynucleotide is not greater than about 600 nucleotides in length. It is understood that irrespective of the length of an amplified polynucleotide, a SNP of interest may be located anywhere along its sequence.

[0084] In a specific embodiment of the invention, the amplified product is at least about 201 nucleotides in length, comprises one of the transcript-based context sequences or the genomic-based context sequences shown in Tables 1-2. Such a product may have additional sequences on its 5' end or 3' end or both. In another embodiment, the amplified product is about 101 nucleotides in length, and it contains a SNP disclosed herein. Preferably, the SNP is located at the middle of the amplified product (e.g., at position 101 in an amplified product that is 201 nucleotides in length, or at position 51 in an amplified product that is 101 nucleotides in length), or within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, or 20 nucleotides from the middle of the amplified product (however, as indicated above, the SNP of interest may be located anywhere along the length of the amplified product).

[0085] The present invention provides the use of isolated nucleic acid molecules that comprise, consist of, or consist essentially of one or more polynucleotide sequences that contain the SNP disclosed herein, complements thereof, and SNP-containing fragments thereof in the method of the present invention.

[0086] Accordingly, the present invention provides the use of nucleic acid molecules that consist of any of the nucleotide sequences shown in Table 1 and/or Table 2 (transcript sequences are provided in Table 1 as SEQ ID NOS: 17-18, genomic sequences are provided in Table 2 as SEQ ID NO: 17560, transcript-based SNP context sequences are provided in Table 1 as SEQ ID NO: 2952-2984, and genomic-based SNP context sequences are provided in Table 2 as SEQ ID NO: 18962-19247), or any nucleic acid molecule that encodes any of the variant proteins provided in Table 1 (SEQ ID NOS: 845-846) in the method of the present invention. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule.

[0087] The present invention further provides the use of nucleic acid molecules that consist essentially of any of the nucleotide sequences shown in Table 1 and/or Table 2 (transcript sequences are provided in Table 1 as SEQ ID NOS: 17-18, genomic sequences are provided in Table 2 as SEQ ID NO: 17560, transcript-based SNP context sequences are provided in Table 1 as SEQ ID NO: 2952-2984, and genomic-based SNP context sequences are provided in Table 2 as SEQ ID NO: 18962-19247), or any nucleic acid molecule that encodes any of the variant proteins provided in Table 1 (SEQ ID NOS: 845-846) in the method of the present invention. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleotide residues in the final nucleic acid molecule.

[0088] The present invention further provides the use of nucleic acid molecules that comprise any of the nucleotide sequences shown in Table 1 and/or Table 2 or a SNP-containing fragment thereof (transcript sequences are provided in Table 1 as SEQ ID NOS: 17-18, genomic sequences are provided in Table 2 as SEQ ID NO: 17560, transcript-based SNP context sequences are provided in Table 1 as SEQ ID NO: 2952-2984, and genomic-based SNP context sequences are provided in Table 2 as SEQ ID NO: 18962-19247), or any nucleic acid molecule that encodes any of the variant proteins provided in Table 1 (SEQ ID NOS: 845-846) in the method of the present invention. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleotide residues, such as residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have one to a few additional nucleotides or can comprise many more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made and isolated is provided below, and such techniques are well known to those of ordinary skill in the art (Sambrook and Russell, 2000, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, NY).

[0089] The isolated nucleic acid molecules can encode mature proteins plus additional amino or carboxyl-terminal amino acids or both, or amino acids interior to the mature peptide (when the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life, or facilitate manipulation of a protein for assay or production. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

[0090] Thus, the isolated nucleic acid molecules include, but are not limited to, nucleic acid molecules having a sequence encoding a peptide alone, a sequence encoding a mature peptide and additional coding sequences such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), a sequence encoding a mature peptide with or without additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5'

and 3' sequences such as transcribed but untranslated sequences that play a role in, for example, transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding, and/or stability of mRNA. In addition, the nucleic acid molecules may be fused to heterologous marker sequences encoding, for example, a peptide that facilitates purification.

[0091] Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form DNA, including cDNA and genomic DNA, which may be obtained, for example, by molecular cloning or produced by chemical synthetic techniques or by a combination thereof (Sambrook and Russell, 2000, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, NY). Furthermore, isolated nucleic acid molecules, particularly SNP detection reagents such as probes and primers, can also be partially or completely in the form of one or more types of nucleic acid analogs, such as peptide nucleic acid (PNA) (U.S. Patent Nos. 5,539,082; 5,527,675; 5,623,049; 5,714,331). The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the complementary non-coding strand (anti-sense strand). DNA, RNA, or PNA segments can be assembled, for example, from fragments of the human genome (in the case of DNA or RNA) or single nucleotides, short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic nucleic acid molecule. Nucleic acid molecules can be readily synthesized using the sequences provided herein as a reference; oligonucleotide and PNA oligomer synthesis techniques are well known in the art (see, e.g., Corey, "Peptide nucleic acids: expanding the scope of nucleic acid recognition", *Trends Biotechnol.* 1997 Jun;15(6):224-9, and Hyrup et al., "Peptide nucleic acids (PNA): synthesis, properties and potential applications", *Bioorg Med Chem.* 1996 Jan;4(1):5-23). Furthermore, large-scale automated oligonucleotide/PNA synthesis (including synthesis on an array or bead surface or other solid support) can readily be accomplished using commercially available nucleic acid synthesizers, such as the Applied Biosystems (Foster City, CA) 3900 High-Throughput DNA Synthesizer or Expedite 8909 Nucleic Acid Synthesis System, and the sequence information provided herein.

[0092] Disclosed herein are nucleic acid analogs that contain modified, synthetic, or non-naturally occurring nucleotides or structural elements or other alternative/modified nucleic acid chemistries known in the art. Such nucleic acid analogs are useful, for example, as detection reagents (e.g., primers/probes) for detecting one or more SNPs identified in Table 1 and/or Table 2. Furthermore, kits/systems (such as beads, arrays, etc.) that include these analogs are also described herein. For example, PNA oligomers that are based on the polymorphic sequences disclosed herein are specifically contemplated. PNA oligomers are analogs of DNA in which the phosphate backbone is replaced with a peptide-like backbone (Lagriffoul et al., *Bioorganic & Medicinal Chemistry Letters*, 4: 1081-1082 (1994), Petersen et al., *Bioorganic & Medicinal Chemistry Letters*, 6: 793-796 (1996), Kumar et al., *Organic Letters* 3(9): 1269-1272 (2001), WO96/04000). PNA hybridizes to complementary RNA or DNA with higher affinity and specificity than conventional oligonucleotides and oligonucleotide analogs. The properties of PNA enable novel molecular biology and biochemistry applications unachievable with traditional oligonucleotides and peptides.

[0093] Additional examples of nucleic acid modifications that improve the binding properties and/or stability of a nucleic acid include the use of base analogs such as inosine, intercalators (U.S. Patent No. 4,835,263) and the minor groove binders (U.S. Patent No. 5,801,115). Thus, references herein to nucleic acid molecules, SNP-containing nucleic acid molecules, SNP detection reagents (e.g., probes and primers), oligonucleotides/polynucleotides include PNA oligomers and other nucleic acid analogs. Other examples of nucleic acid analogs and alternative/modified nucleic acid chemistries known in the art are described in *Current Protocols in Nucleic Acid Chemistry*, John Wiley & Sons, N.Y. (2002).

[0094] The present invention further provides the use of nucleic acid molecules that encode fragments of the variant polypeptides disclosed herein as well as nucleic acid molecules that encode obvious variants of such variant polypeptides in the method of the present invention. Such nucleic acid molecules may be naturally occurring, such as paralogs (different locus) and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, the variants can contain nucleotide substitutions, deletions, inversions and insertions (in addition to the SNPs disclosed in Tables 1-2). Variation can occur in either or both the coding and non-coding regions. The variations can produce conservative and/or non-conservative amino acid substitutions.

[0095] Further variants of the nucleic acid molecules disclosed in Tables 1-2, such as naturally occurring allelic variants (as well as orthologs and paralogs) and synthetic variants produced by mutagenesis techniques, can be identified and/or produced using methods well known in the art. Such further variants can comprise a nucleotide sequence that shares at least 70-80%, 80-85%, 85-90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity with a nucleic acid sequence disclosed in Table 1 and/or Table 2 (or a fragment thereof) and that includes a novel SNP allele disclosed in Table 1 and/or Table 2. Further, variants can comprise a nucleotide sequence that encodes a polypeptide that shares at least 70-80%, 80-85%, 85-90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity with a polypeptide sequence disclosed in Table 1 (or a fragment thereof) and that includes a novel SNP allele disclosed in Table 1 and/or Table 2. Thus, the present invention specifically contemplates the use of isolated nucleic acid molecule that have a certain degree of sequence variation compared with the sequences shown in Tables 1-2, but that contain a novel SNP allele disclosed herein in the method of the present invention. In other words, as long as an isolated nucleic

acid molecule contains a novel SNP allele disclosed herein, other portions of the nucleic acid molecule that flank the novel SNP allele can vary to some degree from the specific transcript, genomic, and context sequences shown in Tables 1-2, and can encode a polypeptide that varies to some degree from the specific polypeptide sequences shown in Table 1.

[0096] To determine the percent identity of two amino acid sequences or two nucleotide sequences of two molecules that share sequence homology, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein, amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0097] The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. (Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch algorithm (J. Mol. Biol. (48):444-453 (1970)) which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

[0098] In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., et al., Nucleic Acids Res. 12(1):387 (1984)), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Myers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4.

[0099] The nucleotide and amino acid sequences for use in the method of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (J. Mol. Biol. 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules used in the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins used in . To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (Nucleic Acids Res. 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. In addition to BLAST, examples of other search and sequence comparison programs used in the art include, but are not limited to, FASTA (Pearson, Methods Mol. Biol. 25, 365-389 (1994)) and KERR (Dufresne et al., Nat Biotechnol 2002 Dec;20(12):1269-71). For further information regarding bioinformatics techniques, see Current Protocols in Bioinformatics, John Wiley & Sons, Inc., N.Y.

[0100] The present invention further provides the use of non-coding fragments of the nucleic acid molecules disclosed in Table 1 and/or Table 2 in the method of the present invention. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, intronic sequences, 5' untranslated regions (UTRs), 3' untranslated regions, gene modulating sequences and gene termination sequences. Such fragments are useful, for example, in controlling heterologous gene expression and in developing screens to identify gene-modulating agents.

SNP Detection Reagents

[0101] In a specific aspect of the present invention, the SNPs disclosed in Table 1 and/or Table 2, and their associated transcript sequences (provided in Table 1 as SEQ ID NOS:17-18), genomic sequences (provided in Table 2 as SEQ ID NO:17560), and context sequences (transcript-based context sequences are provided in Table 1 as SEQ ID NOS:2952-2984 ; genomic-based context sequences are provided in Table 2 as SEQ ID NOS:18962-19247), can be used for the design of SNP detection reagents. As used herein, a "SNP detection reagent" is a reagent that specifically detects a specific target SNP position disclosed herein, and that is preferably specific for a particular nucleotide (allele) of the target SNP position (i.e., the detection reagent preferably can differentiate between different alternative nucleotides at a target SNP position, thereby allowing the identity of the nucleotide present at the target SNP position to be determined).

Typically, such detection reagent hybridizes to a target SNP-containing nucleic acid molecule by complementary base-pairing in a sequence specific manner, and discriminates the target variant sequence from other nucleic acid sequences such as an art-known form in a test sample. An example of a detection reagent is a probe that hybridizes to a target nucleic acid containing one or more of the SNPs provided in Table 1 and/or Table 2. In a preferred embodiment, such a probe can differentiate between nucleic acids having a particular nucleotide (allele) at a target SNP position from other nucleic acids that have a different nucleotide at the same target SNP position. In addition, a detection reagent may hybridize to a specific region 5' and/or 3' to a SNP position, particularly a region corresponding to the context sequences provided in Table 1 and/or Table 2 (transcript-based context sequences are provided in Table 1 as SEQ ID NOS:1657-17,552; genomic-based context sequences are provided in Table 2 as SEQ ID NOS: 18,017-73,085). Another example of a detection reagent is a primer which acts as an initiation point of nucleotide extension along a complementary strand of a target polynucleotide. The SNP sequence information provided herein is also useful for designing primers, e.g. allele-specific primers, to amplify (e.g., using PCR) the SNP used in the method of the present invention.

[0102] In one preferred embodiment of the invention, a SNP detection reagent is an isolated or synthetic DNA or RNA polynucleotide probe or primer or PNA oligomer, or a combination of DNA, RNA and/or PNA, that hybridizes to a segment of a target nucleic acid molecule containing the SNP identified in Table 1 and/or Table 2. A detection reagent in the form of a polynucleotide may optionally contain modified base analogs, intercalators or minor groove binders. Multiple detection reagents such as probes may be, for example, affixed to a solid support (e.g., arrays or beads) or supplied in solution (e.g., probe/primer sets for enzymatic reactions such as PCR, RT-PCR, TaqMan assays, or primer-extension reactions) to form a SNP detection kit.

[0103] A probe or primer typically is a substantially purified oligonucleotide or PNA oligomer. Such oligonucleotide typically comprises a region of complementary nucleotide sequence that hybridizes under stringent conditions to at least about 8, 10, 12, 16, 18, 20, 22, 25, 30, 40, 50, 60, 100 (or any other number in-between) or more consecutive nucleotides in a target nucleic acid molecule. Depending on the particular assay, the consecutive nucleotides can either include the target SNP position, or be a specific region in close enough proximity 5' and/or 3' to the SNP position to carry out the desired assay.

[0104] Other preferred primer and probe sequences can readily be determined using the transcript sequences (SEQ ID NOS:1-828), genomic sequences (SEQ ID NOS:17,553-18,016), and SNP context sequences (transcript-based context sequences are provided in Table 1 as SEQ ID NOS: 1657-17,552; genomic-based context sequences are provided in Table 2 as SEQ ID NOS: 18,017-73,085) disclosed in the Sequence Listing and in Tables 1-2. It will be apparent to one of skill in the art that such primers and probes are directly useful as reagents for genotyping the SNP used in the present invention, and can be incorporated into any kit/system format.

[0105] In order to produce a probe or primer specific for a target SNP-containing sequence, the gene/transcript and/or context sequence surrounding the SNP of interest is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene/SNP context sequence, have a GC content within a range suitable for hybridization, lack predicted secondary structure that may interfere with hybridization, and/or possess other desired characteristics or that lack other undesired characteristics.

[0106] A primer or probe for use in the method of the present invention is typically at least about 8 nucleotides in length. In one embodiment of the invention, a primer or a probe is at least about 10 nucleotides in length. In a preferred embodiment, a primer or a probe is at least about 12 nucleotides in length. In a more preferred embodiment, a primer or probe is at least about 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 nucleotides in length. While the maximal length of a probe can be as long as the target sequence to be detected, depending on the type of assay in which it is employed, it is typically less than about 50, 60, 65, or 70 nucleotides in length. In the case of a primer, it is typically less than about 30 nucleotides in length. In a specific preferred embodiment of the invention, a primer or a probe is within the length of about 18 and about 28 nucleotides. However, in other embodiments, such as nucleic acid arrays and other embodiments in which probes are affixed to a substrate, the probes can be longer, such as on the order of 30-70, 75, 80, 90, 100, or more nucleotides in length (see the section below entitled "SNP Detection Kits and Systems").

[0107] For analyzing SNPs, it may be appropriate to use oligonucleotides specific for alternative SNP alleles. Such oligonucleotides which detect single nucleotide variations in target sequences may be referred to by such terms as "allele-specific oligonucleotides", "allele-specific probes", or "allele-specific primers". The design and use of allele-specific probes for analyzing polymorphisms is described in, e.g., Mutation Detection A Practical Approach, ed. Cotton et al. Oxford University Press, 1998; Saiki et al., Nature 324, 163-166 (1986); Dattagupta, EP235,726; and Saiki, WO 89/11548.

[0108] While the design of each allele-specific primer or probe depends on variables such as the precise composition of the nucleotide sequences flanking a SNP position in a target nucleic acid molecule, and the length of the primer or probe, another factor in the use of primers and probes is the stringency of the condition under which the hybridization between the probe or primer and the target sequence is performed. Higher stringency conditions utilize buffers with lower ionic strength and/or a higher reaction temperature, and tend to require a more perfect match between probe/primer and a target sequence in order to form a stable duplex. If the stringency is too high, however, hybridization may not

occur at all. In contrast, lower stringency conditions utilize buffers with higher ionic strength and/or a lower reaction temperature, and permit the formation of stable duplexes with more mismatched bases between a probe/primer and a target sequence. By way of example and not limitation, exemplary conditions for high stringency hybridization conditions using an allele-specific probe are as follows: Prehybridization with a solution containing 5X standard saline phosphate EDTA (SSPE), 0.5% NaDodSO₄ (SDS) at 55°C, and incubating probe with target nucleic acid molecules in the same solution at the same temperature, followed by washing with a solution containing 2X SSPE, and 0.1% SDS at 55°C or room temperature.

[0109] Moderate stringency hybridization conditions may be used for allele-specific primer extension reactions with a solution containing, e.g., about 50mM KCl at about 46°C. Alternatively, the reaction may be carried out at an elevated temperature such as 60°C. In another embodiment, a moderately stringent hybridization condition suitable for oligonucleotide ligation assay (OLA) reactions wherein two probes are ligated if they are completely complementary to the target sequence may utilize a solution of about 100mM KCl at a temperature of 46°C.

[0110] In a hybridization-based assay, allele-specific probes can be designed that hybridize to a segment of target DNA from one individual but do not hybridize to the corresponding segment from another individual due to the presence of different polymorphic forms (e.g., alternative SNP alleles/nucleotides) in the respective DNA segments from the two individuals. Hybridization conditions should be sufficiently stringent that there is a significant detectable difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles or significantly more strongly to one allele. While a probe may be designed to hybridize to a target sequence that contains a SNP site such that the SNP site aligns anywhere along the sequence of the probe, the probe is preferably designed to hybridize to a segment of the target sequence such that the SNP site aligns with a central position of the probe (e.g., a position within the probe that is at least three nucleotides from either end of the probe). This design of probe generally achieves good discrimination in hybridization between different allelic forms.

[0111] In another embodiment, a probe or primer may be designed to hybridize to a segment of target DNA such that the SNP aligns with either the 5' most end or the 3' most end of the probe or primer. In a specific preferred embodiment which is particularly suitable for use in an oligonucleotide ligation assay (U.S. Patent No. 4,988,617), the 3'most nucleotide of the probe aligns with the SNP position in the target sequence.

[0112] Oligonucleotide probes and primers may be prepared by methods well known in the art. Chemical synthetic methods include, but are limited to, the phosphotriester method described by Narang et al., 1979, Methods in Enzymology 68:90; the phosphodiester method described by Brown et al., 1979, Methods in Enzymology 68:109, the diethylphosphoamidate method described by Beaucage et al., 1981, Tetrahedron Letters 22:1859; and the solid support method described in U.S. Patent No. 4,458,066.

[0113] Allele-specific probes are often used in pairs (or, less commonly, in sets of 3 or 4, such as if a SNP position is known to have 3 or 4 alleles, respectively, or to assay both strands of a nucleic acid molecule for a target SNP allele), and such pairs may be identical except for a one nucleotide mismatch that represents the allelic variants at the SNP position. Commonly, one member of a pair perfectly matches a reference form of a target sequence that has a more common SNP allele (i.e., the allele that is more frequent in the target population) and the other member of the pair perfectly matches a form of the target sequence that has a less common SNP allele (i.e., the allele that is rarer in the target population). In the case of an array, multiple pairs of probes can be immobilized on the same support for simultaneous analysis of multiple different polymorphisms.

[0114] In one type of PCR-based assay, an allele-specific primer hybridizes to a region on a target nucleic acid molecule that overlaps a SNP position and only primes amplification of an allelic form to which the primer exhibits perfect complementarity (Gibbs, 1989, Nucleic Acid Res. 17 2427-2448). Typically, the primer's 3'-most nucleotide is aligned with and complementary to the SNP position of the target nucleic acid molecule. This primer is used in conjunction with a second primer that hybridizes at a distal site. Amplification proceeds from the two primers, producing a detectable product that indicates which allelic form is present in the test sample. A control is usually performed with a second pair of primers, one of which shows a single base mismatch at the polymorphic site and the other of which exhibits perfect complementarity to a distal site. The single-base mismatch prevents amplification or substantially reduces amplification efficiency, so that either no detectable product is formed or it is formed in lower amounts or at a slower pace. The method generally works most effectively when the mismatch is at the 3'-most position of the oligonucleotide (i.e., the 3'-most position of the oligonucleotide aligns with the target SNP position) because this position is most destabilizing to elongation from the primer (see, e.g., WO 93/22456). This PCR-based assay can be utilized as part of the TaqMan assay, described below.

[0115] In a specific embodiment of the invention, a primer for use in the method of the invention contains a sequence substantially complementary to a segment of a target SNP-containing nucleic acid molecule except that the primer has a mismatched nucleotide in one of the three nucleotide positions at the 3'-most end of the primer, such that the mismatched nucleotide does not base pair with a particular allele at the SNP site. In a preferred embodiment, the mismatched nucleotide in the primer is the second from the last nucleotide at the 3'-most position of the primer. In a more preferred embodiment, the mismatched nucleotide in the primer is the last nucleotide at the 3'-most position of the primer.

[0116] In another embodiment of the invention, a SNP detection reagent for use in the method of the invention is

labeled with a fluorogenic reporter dye that emits a detectable signal. While the preferred reporter dye is a fluorescent dye, any reporter dye that can be attached to a detection reagent such as an oligonucleotide probe or primer is suitable for use in the invention. Such dyes include, but are not limited to, Acridine, AMCA, BODIPY, Cascade Blue, Cy2, Cy3, Cy5, Cy7, Dabcyl, Edans, Eosin, Erythrosin, Fluorescein, 6-Fam, Tet, Joe, Hex, Oregon Green, Rhodamine, Rhodol Green, Tamra, Rox, and Texas Red.

[0117] In yet another embodiment of the invention, the detection reagent may be further labeled with a quencher dye such as Tamra, especially when the reagent is used as a self-quenching probe such as a TaqMan (U.S. Patent Nos. 5,210,015 and 5,538,848) or Molecular Beacon probe (U.S. Patent Nos. 5,118,801 and 5,312,728), or other stemless or linear beacon probe (Livak et al., 1995, PCR Method Appl. 4:357-362; Tyagi et al., 1996, Nature Biotechnology 14: 303-308; Nazarenko et al., 1997, Nucl. Acids Res. 25:2516-2521; U.S. Patent Nos. 5,866,336 and 6,117,635).

[0118] The detection reagents used in the method of the invention may also contain other labels, including but not limited to, biotin for streptavidin binding, hapten for antibody binding, and oligonucleotide for binding to another complementary oligonucleotide such as pairs of zipcodes.

[0119] The present invention also contemplates the use of reagents that do not contain (or that are complementary to) a SNP nucleotide identified herein but that are used to assay one or more SNPs disclosed herein. For example, primers that flank, but do not hybridize directly to a target SNP position provided herein are useful in primer extension reactions in which the primers hybridize to a region adjacent to the target SNP position (i.e., within one or more nucleotides from the target SNP site). During the primer extension reaction, a primer is typically not able to extend past a target SNP site if a particular nucleotide (allele) is present at that target SNP site, and the primer extension product can readily be detected in order to determine which SNP allele is present at the target SNP site. For example, particular ddNTPs are typically used in the primer extension reaction to terminate primer extension once a ddNTP is incorporated into the extension product (a primer extension product which includes a ddNTP at the 3'-most end of the primer extension product, and in which the ddNTP corresponds to a SNP disclosed herein, is a composition that is encompassed by the present disclosure). Thus, reagents that bind to a nucleic acid molecule in a region adjacent to a SNP site, even though the bound sequences do not necessarily include the SNP site itself, are also encompassed by the present disclosure.

SNP Detection Kits and Systems

[0120] A person skilled in the art will recognize that, based on the SNP and associated sequence information disclosed herein, detection reagents can be developed and used to assay the SNP as used in the method of the present invention individually or in combination, and such detection reagents can be readily incorporated into one of the established kit or system formats which are well known in the art. The terms "kits" and "systems", as used herein in the context of SNP detection reagents, are intended to refer to such things as combinations of multiple SNP detection reagents, or one or more SNP detection reagents in combination with one or more other types of elements or components (e.g., other types of biochemical reagents, containers, packages such as packaging intended for commercial sale, substrates to which SNP detection reagents are attached, electronic hardware components, etc.). Accordingly, the present invention further provides the use of SNP detection kits and systems in the method of the present invention including but not limited to, packaged probe and primer sets (e.g., TaqMan probe/primer sets), arrays/microarrays of nucleic acid molecules, and beads that contain one or more probes, primers, or other detection reagents for detecting the SNP used in the method of the present invention. The kits/systems can optionally include various electronic hardware components; for example, arrays ("DNA chips") and microfluidic systems ("lab-on-a-chip" systems) provided by various manufacturers typically comprise hardware components. Other kits/systems (e.g., probe/primer sets) may not include electronic hardware components, but may be comprised of, for example, one or more SNP detection reagents (along with, optionally, other biochemical reagents) packaged in one or more containers.

[0121] In some embodiments, a SNP detection kit typically contains one or more detection reagents and other components (e.g., a buffer, enzymes such as DNA polymerases or ligases, chain extension nucleotides such as deoxynucleotide triphosphates, and in the case of Sanger-type DNA sequencing reactions, chain terminating nucleotides, positive control sequences, negative control sequences, and the like) necessary to carry out an assay or reaction, such as amplification and/or detection of a SNP-containing nucleic acid molecule. A kit may further contain means for determining the amount of a target nucleic acid, and means for comparing the amount with a standard, and can comprise instructions for using the kit to detect the SNP-containing nucleic acid molecule of interest. In one embodiment of the present invention, the use of kits is provided which contain the necessary reagents to carry out one or more assays to detect the SNP disclosed herein. In a preferred embodiment of the present invention, SNP detection kits/systems used in the present invention are in the form of nucleic acid arrays, or compartmentalized kits, including microfluidic/lab-on-a-chip systems.

[0122] SNP detection kits/systems may contain, for example, one or more probes, or pairs of probes, that hybridize to a nucleic acid molecule at or near each target SNP position. Multiple pairs of allele-specific probes may be included in the kit/system to simultaneously assay large numbers of SNPs, at least one of which is the SNP used in the method of the present invention. In some kits/systems, the allele-specific probes are immobilized to a substrate such as an array

or bead. For example, the same substrate can comprise allele-specific probes for detecting at least 1; 10; 100; 1000; 10,000; 100,000 (or any other number in-between) or substantially all of the SNPs shown in Table 1 and/or Table 2.

[0123] The terms "arrays", "microarrays", and "DNA chips" are used herein interchangeably to refer to an array of distinct polynucleotides affixed to a substrate, such as glass, plastic, paper, nylon or other type of membrane, filter, chip, or any other suitable solid support. The polynucleotides can be synthesized directly on the substrate, or synthesized separate from the substrate and then affixed to the substrate. In one embodiment, the microarray is prepared and used according to the methods described in U.S. Patent No. 5,837,832, Chee et al., PCT application W095/11995 (Chee et al.), Lockhart, D. J. et al. (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. et al. (1996; Proc. Natl. Acad. Sci. 93: 10614-10619). In other embodiments, such arrays are produced by the methods described by Brown et al., U.S. Patent No. 5,807,522.

[0124] Nucleic acid arrays are reviewed in the following references: Zammattéo et al., "New chips for molecular biology and diagnostics", *Biotechnol Annu Rev.* 2002;8:85-101; Sosnowski et al., "Active microelectronic array system for DNA hybridization, genotyping and pharmacogenomic applications", *Psychiatr Genet.* 2002 Dec;12(4):181-92; Heller, "DNA microarray technology: devices, systems, and applications", *Annu Rev Biomed Eng.* 2002;4:129-53. Epub 2002 Mar 22; Kolchinsky et al., "Analysis of SNPs and other genomic variations using gel-based chips", *Hum Mutat.* 2002 Apr;19(4):343-60; and McGall et al., "High-density genechip oligonucleotide probe arrays", *Adv Biochem Eng Biotechnol.* 2002;77:21-42.

[0125] Any number of probes, such as allele-specific probes, may be implemented in an array, and each probe or pair of probes can hybridize to a different SNP position. In the case of polynucleotide probes, they can be synthesized at designated areas (or synthesized separately and then affixed to designated areas) on a substrate using a light-directed chemical process. Each DNA chip can contain, for example, thousands to millions of individual synthetic polynucleotide probes arranged in a grid-like pattern and miniaturized (e.g., to the size of a dime). Preferably, probes are attached to a solid support in an ordered, addressable array.

[0126] A microarray can be composed of a large number of unique, single-stranded polynucleotides; usually either synthetic antisense polynucleotides or fragments of cDNAs, fixed to a solid support. Typical polynucleotides are preferably about 6-60 nucleotides in length, more preferably about 15-30 nucleotides in length, and most preferably about 18-25 nucleotides in length. For certain types of microarrays or other detection kits/systems, it may be preferable to use oligonucleotides that are only about 7-20 nucleotides in length. In other types of arrays, such as arrays used in conjunction with chemiluminescent detection technology, preferred probe lengths can be, for example, about 15-80 nucleotides in length, preferably about 50-70 nucleotides in length, more preferably about 55-65 nucleotides in length, and most preferably about 60 nucleotides in length. The microarray or detection kit can contain polynucleotides that cover the known 5' or 3' sequence of a gene/transcript or target SNP site, sequential polynucleotides that cover the full-length sequence of a gene/transcript; or unique polynucleotides selected from particular areas along the length of a target gene/transcript sequence, particularly areas corresponding to one or more SNPs disclosed in Table 1 and/or Table 2. Polynucleotides used in the microarray or detection kit can be specific to a SNP or SNPs of interest (e.g., specific to a particular SNP allele at a target SNP site, or specific to particular SNP alleles at multiple different SNP sites), or specific to a polymorphic gene/transcript or genes/transcripts of interest.

[0127] Hybridization assays based on polynucleotide arrays rely on the differences in hybridization stability of the probes to perfectly matched and mismatched target sequence variants. For SNP genotyping, it is generally preferable that stringency conditions used in hybridization assays are high enough such that nucleic acid molecules that differ from one another at as little as a single SNP position can be differentiated (e.g., typical SNP hybridization assays are designed so that hybridization will occur only if one particular nucleotide is present at a SNP position, but will not occur if an alternative nucleotide is present at that SNP position). Such high stringency conditions may be preferable when using, for example, nucleic acid arrays of allele-specific probes for SNP detection. Such high stringency conditions are described in the preceding section, and are well known to those skilled in the art and can be found in, for example, *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6.

[0128] In other embodiments, the arrays are used in conjunction with chemiluminescent detection technology. The following patents and patent applications, provide additional information pertaining to chemiluminescent detection: U.S. patent applications 10/620332 and 10/620333 describe chemiluminescent approaches for microarray detection; U.S. Patent Nos. 6124478, 6107024, 5994073, 5981768, 5871938, 5843681, 5800999, and 5773628 describe methods and compositions of dioxetane for performing chemiluminescent detection; and U.S. published application US2002/0110828 discloses methods and compositions for microarray controls.

[0129] In one embodiment of the invention, a nucleic acid array used in the present invention can comprise an array of probes of about 15-25 nucleotides in length. In further embodiments, a nucleic acid array can comprise any number of probes, in which at least one probe is capable of detecting one or more SNPs disclosed in Table 1 and/or Table 2, and/or at least one probe comprises a fragment of one of the sequences selected from the group consisting of those disclosed in Table 1, Table 2, the Sequence Listing, and sequences complementary thereto, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 12, 15, 16, 18, 20, more preferably 22, 25, 30, 40, 47, 50, 55,

60, 65, 70, 80, 90, 100, or more consecutive nucleotides (or any other number in-between) and containing (or being complementary to) a novel SNP allele disclosed in Table 1 and/or Table 2. In some embodiments, the nucleotide complementary to the SNP site is within 5, 4, 3, 2, or 1 nucleotide from the center of the probe, more preferably at the center of said probe.

[0130] A polynucleotide probe can be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler et al.). In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more polynucleotides, or any other number which lends itself to the efficient use of commercially available instrumentation.

[0131] Using such arrays or other kits/systems, the present invention provides methods of identifying the SNPs disclosed herein in a test sample. Such methods typically involve incubating a test sample of nucleic acids with an array comprising one or more probes corresponding to the SNP position used in the present invention, and assaying for binding of a nucleic acid from the test sample with one or more of the probes. Conditions for incubating a SNP detection reagent (or a kit/system that employs one or more such SNP detection reagents) with a test sample vary. Incubation conditions depend on such factors as the format employed in the assay, the detection methods employed, and the type and nature of the detection reagents used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification and array assay formats can readily be adapted to detect the SNPs disclosed herein.

[0132] A SNP detection kit/system for use in the method of the present invention may include components that are used to prepare nucleic acids from a test sample for the subsequent amplification and/or detection of a SNP-containing nucleic acid molecule. Such sample preparation components can be used to produce nucleic acid extracts (including DNA and/or RNA), proteins or membrane extracts from any bodily fluids (such as blood, serum, plasma, urine, saliva, phlegm, gastric juices, semen, tears, sweat, etc.), skin, hair, cells (especially nucleated cells), biopsies, buccal swabs or tissue specimens. The test samples used in the above-described methods will vary based on such factors as the assay format, nature of the detection method, and the specific tissues, cells or extracts used as the test sample to be assayed. Methods of preparing nucleic acids, proteins, and cell extracts are well known in the art and can be readily adapted to obtain a sample that is compatible with the system utilized. Automated sample preparation systems for extracting nucleic acids from a test sample are commercially available, and examples are Qiagen's BioRobot 9600, Applied Biosystems' PRISM 6700, and Roche Molecular Systems' COBAS AmpliPrep System.

[0133] Another form of kit contemplated for use in the present invention is a compartmentalized kit. A compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include, for example, small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allow one to efficiently transfer reagents from one compartment to another compartment such that the test samples and reagents are not cross-contaminated, or from one container to another vessel not included in the kit, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another or to another vessel. Such containers may include, for example, one or more containers which will accept the test sample, one or more containers which contain at least one probe or other SNP detection reagent for detecting the SNP for use in the method of the present invention, one or more containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and one or more containers which contain the reagents used to reveal the presence of the bound probe or other SNP detection reagents. The kit can optionally further comprise compartments and/or reagents for, for example, nucleic acid amplification or other enzymatic reactions such as primer extension reactions, hybridization, ligation, electrophoresis (preferably capillary electrophoresis), mass spectrometry, and/or laser-induced fluorescent detection. The kit may also include instructions for using the kit. Exemplary compartmentalized kits include microfluidic devices known in the art (see, e.g., Weigl et al., "Lab-on-a-chip for drug development", Adv Drug Deliv Rev. 2003 Feb 24;55(3):349-77). In such microfluidic devices, the containers may be referred to as, for example, microfluidic "compartments", "chambers", or "channels".

[0134] Microfluidic devices, which may also be referred to as "lab-on-a-chip" systems, biomedical micro-electro-mechanical systems (bioMEMs), or multicomponent integrated systems, are exemplary kits/systems for use in the present invention for analyzing SNPs. Such systems miniaturize and compartmentalize processes such as probe/target hybridization, nucleic acid amplification, and capillary electrophoresis reactions in a single functional device. Such microfluidic devices typically utilize detection reagents in at least one aspect of the system, and such detection reagents may be used to detect the SNP used in the method of the present invention. One example of a microfluidic system is disclosed in U.S. Patent No. 5,589,136, which describes the integration of PCR amplification and capillary electrophoresis in chips. Exemplary microfluidic systems comprise a pattern of microchannels designed onto a glass, silicon, quartz, or plastic wafer included on a microchip. The movements of the samples may be controlled by electric, electroosmotic or hydrostatic forces applied across different areas of the microchip to create functional microscopic valves and pumps with no moving

parts. Varying the voltage can be used as a means to control the liquid flow at intersections between the micro-machined channels and to change the liquid flow rate for pumping across different sections of the microchip. See, for example, U.S. Patent Nos. 6,153,073, Dubrow et al., and 6,156,181, Parce et al.

[0135] For genotyping SNPs, an exemplary microfluidic system may integrate, for example, nucleic acid amplification, primer extension, capillary electrophoresis, and a detection method such as laser induced fluorescence detection. In a first step of an exemplary process for using such an exemplary system, nucleic acid samples are amplified, preferably by PCR. Then, the amplification products are subjected to automated primer extension reactions using ddNTPs (specific fluorescence for each ddNTP) and the appropriate oligonucleotide primers to carry out primer extension reactions which hybridize just upstream of the targeted SNP. Once the extension at the 3' end is completed, the primers are separated from the unincorporated fluorescent ddNTPs by capillary electrophoresis. The separation medium used in capillary electrophoresis can be, for example, polyacrylamide, polyethyleneglycol or dextran. The incorporated ddNTPs in the single nucleotide primer extension products are identified by laser-induced fluorescence detection. Such an exemplary microchip can be used to process, for example, at least 96 to 384 samples, or more, in parallel.

USES OF NUCLEIC ACID MOLECULES

[0136] The nucleic acid molecules used in the method of the present invention have a variety of uses, especially in the diagnosis and treatment of myocardial infarction. For example, the nucleic acid molecules are useful as hybridization probes, such as for genotyping SNPs in messenger RNA, transcript, cDNA, genomic DNA, amplified DNA or other nucleic acid molecules, and for isolating full-length cDNA and genomic clones encoding the variant peptides disclosed in Table 1 as well as their orthologs.

[0137] A probe can hybridize to any nucleotide sequence along the entire length of a nucleic acid molecule provided in Table 1 and/or Table 2. Preferably, a probe for use in the method of the present invention hybridizes to a region of a target sequence that encompasses a SNP position indicated in Table 1 and/or Table 2. More preferably, a probe hybridizes to a SNP-containing target sequence in a sequence-specific manner such that it distinguishes the target sequence from other nucleotide sequences which vary from the target sequence only by which nucleotide is present at the SNP site. Such a probe is particularly useful for detecting the presence of a SNP-containing nucleic acid in a test sample, or for determining which nucleotide (allele) is present at a particular SNP site (i.e., genotyping the SNP site).

[0138] A nucleic acid hybridization probe may be used for determining the presence, level, form, and/or distribution of nucleic acid expression. The nucleic acid whose level is determined can be DNA or RNA. Accordingly, probes specific for the SNPs described herein can be used to assess the presence, expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in gene expression relative to normal levels. *In vitro* techniques for detection of mRNA include, for example, Northern blot hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA include Southern blot hybridizations and *in situ* hybridizations (Sambrook and Russell, 2000, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, NY).

[0139] Probes can be used as part of a diagnostic test kit for identifying cells or tissues in which a variant protein is expressed, such as by measuring the level of a variant protein-encoding nucleic acid (e.g., mRNA) in a sample of cells from a subject or determining if a polynucleotide contains a SNP of interest.

[0140] Thus, the nucleic acid molecules used in the method of the invention can be used as hybridization probes to detect the SNPs disclosed herein, thereby determining whether an individual with the polymorphisms is at risk for myocardial infarction or has developed early stage myocardial infarction. Detection of a SNP associated with a disease phenotype provides a diagnostic tool for an active disease and/or genetic predisposition to the disease.

[0141] The nucleic acid molecules used in the method of the invention are also useful as primers to amplify any given region of a nucleic acid molecule, particularly a region containing a SNP identified in Table 1 and/or Table 2.

[0142] The nucleic acid molecules used in the method of the invention are also useful for constructing recombinant vectors (described in greater detail below). Such vectors include expression vectors that express a portion of, or all of, any of the variant peptide sequences provided in Table 1. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced SNPs.

[0143] The nucleic acid molecules used in the method of the invention are also useful for expressing antigenic portions of the variant proteins, particularly antigenic portions that contain a variant amino acid sequence (e.g., an amino acid substitution) caused by a SNP disclosed in Table 1 and/or Table 2.

[0144] The nucleic acid molecules used in the method of the invention are also useful for constructing vectors containing a gene regulatory region of the nucleic acid molecules used in the method of the present invention.

[0145] The nucleic acid molecules used in the method of the invention are also useful for designing ribozymes corresponding to all, or a part, of an mRNA molecule expressed from a SNP-containing nucleic acid molecule described herein.

[0146] The nucleic acid molecules used in the method of the invention are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and variant peptides.

[0147] The nucleic acid molecules used in the method of the invention are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and variant peptides. The production of recombinant cells and transgenic animals having nucleic acid molecules which contain the SNPs disclosed in Table 1 and/or Table 2 allow, for example, effective clinical design of treatment compounds and dosage regimens.

[0148] The nucleic acid molecules used in the method of the invention are also useful in assays for drug screening to identify compounds that, for example, modulate nucleic acid expression.

[0149] The nucleic acid molecules used in the method of the invention are also useful in gene therapy in patients whose cells have aberrant gene expression. Thus, recombinant cells, which include a patient's cells that have been engineered *ex vivo* and returned to the patient, can be introduced into an individual where the recombinant cells produce the desired protein to treat the individual.

SNP Genotyping Methods

[0150] The process of determining which specific nucleotide (i.e., allele) is present at each of one or more SNP positions, such as a SNP position in a nucleic acid molecule disclosed in Table 1 and/or Table 2, is referred to as SNP genotyping. The present disclosure describes methods of SNP genotyping, such as for use in screening for myocardial infarction or related pathologies, or determining predisposition thereto, or determining responsiveness to a form of treatment, or in genome mapping or SNP association analysis, etc.

[0151] Nucleic acid samples can be genotyped to determine which allele(s) is/are present at any given genetic region (e.g., SNP position) of interest by methods well known in the art. The neighboring sequence can be used to design SNP detection reagents such as oligonucleotide probes, which may optionally be implemented in a kit format. Exemplary SNP genotyping methods are described in Chen et al., "Single nucleotide polymorphism genotyping: biochemistry, protocol, cost and throughput", *Pharmacogenomics J.* 2003;3(2):77-96; Kwok et al., "Detection of single nucleotide polymorphisms", *Curr Issues Mol Biol.* 2003 Apr;5(2):43-60; Shi, "Technologies for individual genotyping: detection of genetic polymorphisms in drug targets and disease genes", *Am J Pharmacogenomics.* 2002;2(3):197-205; and Kwok, "Methods for genotyping single nucleotide polymorphisms", *Annu Rev Genomics Hum Genet* 2001;2:235-58. Exemplary techniques for high-throughput SNP genotyping are described in Marnellos, "High-throughput SNP analysis for genetic association studies", *Curr Opin Drug Discov Devel.* 2003 May;6(3):317-21. Common SNP genotyping methods include, but are not limited to, TaqMan assays, molecular beacon assays, nucleic acid arrays, allele-specific primer extension, allele-specific PCR, arrayed primer extension, homogeneous primer extension assays, primer extension with detection by mass spectrometry, pyrosequencing, multiplex primer extension sorted on genetic arrays, ligation with rolling circle amplification, homogeneous ligation, OLA (U.S. Patent No. 4,988,167), multiplex ligation reaction sorted on genetic arrays, restriction-fragment length polymorphism, single base extension-tag assays, and the Invader assay. Such methods may be used in combination with detection mechanisms such as, for example, luminescence or chemiluminescence detection, fluorescence detection, time-resolved fluorescence detection, fluorescence resonance energy transfer, fluorescence polarization, mass spectrometry, and electrical detection.

[0152] Various methods for detecting polymorphisms include, but are not limited to, methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers et al., *Science* 230:1242 (1985); Cotton et al., *PNAS* 85:4397 (1988); and Saleeba et al., *Meth. Enzymol.* 217:286-295 (1992)), comparison of the electrophoretic mobility of variant and wild type nucleic acid molecules (Orita et al., *PNAS* 86:2766 (1989); Cotton et al., *Mutat. Res.* 285:125-144 (1993); and Hayashi et al., *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and assaying the movement of polymorphic or wild-type fragments in polyacrylamide gels containing a gradient of denaturant using denaturing gradient gel electrophoresis (DGGE) (Myers et al., *Nature* 313:495 (1985)). Sequence variations at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or chemical cleavage methods.

[0153] In a preferred embodiment, SNP genotyping is performed using the TaqMan assay, which is also known as the 5' nuclease assay (U.S. Patent Nos. 5,210,015 and 5,538,848). The TaqMan assay detects the accumulation of a specific amplified product during PCR. The TaqMan assay utilizes an oligonucleotide probe labeled with a fluorescent reporter dye and a quencher dye. The reporter dye is excited by irradiation at an appropriate wavelength, it transfers energy to the quencher dye in the same probe via a process called fluorescence resonance energy transfer (FRET). When attached to the probe, the excited reporter dye does not emit a signal. The proximity of the quencher dye to the reporter dye in the intact probe maintains a reduced fluorescence for the reporter. The reporter dye and quencher dye may be at the 5' most and the 3' most ends, respectively, or vice versa. Alternatively, the reporter dye may be at the 5' or 3' most end while the quencher dye is attached to an internal nucleotide, or vice versa. In yet another embodiment, both the reporter and the quencher may be attached to internal nucleotides at a distance from each other such that fluorescence of the reporter is reduced.

[0154] During PCR, the 5' nuclease activity of DNA polymerase cleaves the probe, thereby separating the reporter dye and the quencher dye and resulting in increased fluorescence of the reporter. Accumulation of PCR product is detected directly by monitoring the increase in fluorescence of the reporter dye. The DNA polymerase cleaves the probe between the reporter dye and the quencher dye only if the probe hybridizes to the target SNP-containing template which is amplified during PCR, and the probe is designed to hybridize to the target SNP site only if a particular SNP allele is present.

[0155] Preferred TaqMan primer and probe sequences can readily be determined using the SNP and associated nucleic acid sequence information provided herein. A number of computer programs, such as Primer Express (Applied Biosystems, Foster City, CA), can be used to rapidly obtain optimal primer/probe sets. It will be apparent to one of skill in the art that such primers and probes for detecting the SNP used in the method of the present invention are useful in diagnostic assays for myocardial infarction and related pathologies, and can be readily incorporated into a kit format. The present disclosure also describes modifications of the Taqman assay well known in the art such as the use of Molecular Beacon probes (U.S. Patent Nos. 5,118,801 and 5,312,728) and other variant formats (U.S. Patent Nos. 5,866,336 and 6,117,635).

[0156] Another preferred method for genotyping the SNP used in the method of the present invention is the use of two oligonucleotide probes in an OLA (see, e.g., U.S. Patent No. 4,988,617). In this method, one probe hybridizes to a segment of a target nucleic acid with its 3' most end aligned with the SNP site. A second probe hybridizes to an adjacent segment of the target nucleic acid molecule directly 3' to the first probe. The two juxtaposed probes hybridize to the target nucleic acid molecule, and are ligated in the presence of a linking agent such as a ligase if there is perfect complementarity between the 3' most nucleotide of the first probe with the SNP site. If there is a mismatch, ligation would not occur. After the reaction, the ligated probes are separated from the target nucleic acid molecule, and detected as indicators of the presence of a SNP.

[0157] The following patents, patent applications, and published international patent applications, provide additional information pertaining to techniques for carrying out various types of OLA: U.S. Patent Nos. 6027889, 6268148, 5494810, 5830711, and 6054564 describe OLA strategies for performing SNP detection; WO 97/31256 and WO 00/56927 describe OLA strategies for performing SNP detection using universal arrays, wherein a zipcode sequence can be introduced into one of the hybridization probes, and the resulting product, or amplified product, hybridized to a universal zip code array; U.S. application US01/17329 (and 09/584,905) describes OLA (or LDR) followed by PCR, wherein zipcodes are incorporated into OLA probes, and amplified PCR products are determined by electrophoretic or universal zipcode array readout; U.S. applications 60/427818, 60/445636, and 60/445494 describe SNIPlex methods and software for multiplexed SNP detection using OLA followed by PCR, wherein zipcodes are incorporated into OLA probes, and amplified PCR products are hybridized with a zipchute reagent, and the identity of the SNP determined from electrophoretic readout of the zipchute. In some embodiments, OLA is carried out prior to PCR (or another method of nucleic acid amplification). In other embodiments, PCR (or another method of nucleic acid amplification) is carried out prior to OLA.

[0158] Another method for SNP genotyping is based on mass spectrometry. Mass spectrometry takes advantage of the unique mass of each of the four nucleotides of DNA. SNPs can be unambiguously genotyped by mass spectrometry by measuring the differences in the mass of nucleic acids having alternative SNP alleles. MALDI-TOF (Matrix Assisted Laser Desorption Ionization - Time of Flight) mass spectrometry technology is preferred for extremely precise determinations of molecular mass, such as SNPs. Numerous approaches to SNP analysis have been developed based on mass spectrometry. Preferred mass spectrometry-based methods of SNP genotyping include primer extension assays, which can also be utilized in combination with other approaches, such as traditional gel-based formats and microarrays.

[0159] Typically, the primer extension assay involves designing and annealing a primer to a template PCR amplicon upstream (5') from a target SNP position. A mix of dideoxynucleotide triphosphates (ddNTPs) and/or deoxynucleotide triphosphates (dNTPs) are added to a reaction mixture containing template (e.g., a SNP-containing nucleic acid molecule which has typically been amplified, such as by PCR), primer, and DNA polymerase. Extension of the primer terminates at the first position in the template where a nucleotide complementary to one of the ddNTPs in the mix occurs. The primer can be either immediately adjacent (i.e., the nucleotide at the 3' end of the primer hybridizes to the nucleotide next to the target SNP site) or two or more nucleotides removed from the SNP position. If the primer is several nucleotides removed from the target SNP position, the only limitation is that the template sequence between the 3' end of the primer and the SNP position cannot contain a nucleotide of the same type as the one to be detected, or this will cause premature termination of the extension primer. Alternatively, if all four ddNTPs alone, with no dNTPs, are added to the reaction mixture, the primer will always be extended by only one nucleotide, corresponding to the target SNP position. In this instance, primers are designed to bind one nucleotide upstream from the SNP position (i.e., the nucleotide at the 3' end of the primer hybridizes to the nucleotide that is immediately adjacent to the target SNP site on the 5' side of the target SNP site). Extension by only one nucleotide is preferable, as it minimizes the overall mass of the extended primer, thereby increasing the resolution of mass differences between alternative SNP nucleotides. Furthermore, mass-tagged ddNTPs can be employed in the primer extension reactions in place of unmodified ddNTPs. This increases the mass difference between primers extended with these ddNTPs, thereby providing increased sensitivity and accuracy, and is

particularly useful for typing heterozygous base positions. Mass-tagging also alleviates the need for intensive sample-preparation procedures and decreases the necessary resolving power of the mass spectrometer.

[0160] The extended primers can then be purified and analyzed by MALDI-TOF mass spectrometry to determine the identity of the nucleotide present at the target SNP position. In one method of analysis, the products from the primer extension reaction are combined with light absorbing crystals that form a matrix. The matrix is then hit with an energy source such as a laser to ionize and desorb the nucleic acid molecules into the gas-phase. The ionized molecules are then ejected into a flight tube and accelerated down the tube towards a detector. The time between the ionization event, such as a laser pulse, and collision of the molecule with the detector is the time of flight of that molecule. The time of flight is precisely correlated with the mass-to-charge ratio (m/z) of the ionized molecule. Ions with smaller m/z travel down the tube faster than ions with larger m/z and therefore the lighter ions reach the detector before the heavier ions. The time-of-flight is then converted into a corresponding, and highly precise, m/z . In this manner, SNPs can be identified based on the slight differences in mass, and the corresponding time of flight differences, inherent in nucleic acid molecules having different nucleotides at a single base position. For further information regarding the use of primer extension assays in conjunction with MALDI-TOF mass spectrometry for SNP genotyping, see, e.g., Wise et al., "A standard protocol for single nucleotide primer extension in the human genome using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry", *Rapid Commun Mass Spectrom.* 2003;17(11):1195-202.

[0161] The following references provide further information describing mass spectrometry-based methods for SNP genotyping: Bocker, "SNP and mutation discovery using base-specific cleavage and MALDI-TOF mass spectrometry", *Bioinformatics.* 2003 Jul;19 Suppl 1:144-153; Storm et al., "MALDI-TOF mass spectrometry-based SNP genotyping", *Methods Mol Biol.* 2003;212:241-62; Jurinke et al., "The use of MassARRAY technology for high throughput genotyping", *Adv Biochem Eng Biotechnol.* 2002;77:57-74; and Jurinke et al., "Automated genotyping using the DNA MassArray technology", *Methods Mol Biol.* 2002;187:179-92.

[0162] SNPs can also be scored by direct DNA sequencing. A variety of automated sequencing procedures can be utilized ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO94/16101; Cohen et al., *Adv. Chromatogr.* 36:127-162 (1996); and Griffin et al., *Appl. Biochem. Biotechnol.* 38:147-159 (1993)). The nucleic acid sequences used in the method of the present invention enable one of ordinary skill in the art to readily design sequencing primers for such automated sequencing procedures. Commercial instrumentation, such as the Applied Biosystems 377, 3100, 3700, 3730, and 3730x1 DNA Analyzers (Foster City, CA), is commonly used in the art for automated sequencing.

[0163] Other methods that can be used to genotype the SNP used in the method of the present invention include single-strand conformational polymorphism (SSCP), and denaturing gradient gel electrophoresis (DGGE) (Myers et al., *Nature* 313:495 (1985)). SSCP identifies base differences by alteration in electrophoretic migration of single stranded PCR products, as described in Orita et al., *Proc. Nat. Acad.* Single-stranded PCR products can be generated by heating or otherwise denaturing double stranded PCR products. Single-stranded nucleic acids may refold or form secondary structures that are partially dependent on the base sequence. The different electrophoretic mobilities of single-stranded amplification products are related to base-sequence differences at SNP positions. DGGE differentiates SNP alleles based on the different sequence-dependent stabilities and melting properties inherent in polymorphic DNA and the corresponding differences in electrophoretic migration patterns in a denaturing gradient gel (Erich, ed., *PCR Technology, Principles and Applications for DNA Amplification*, W.H. Freeman and Co, New York, 1992, Chapter 7).

[0164] Sequence-specific ribozymes (U.S. Patent No. 5,498,531) can also be used to score SNPs based on the development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature. If the SNP affects a restriction enzyme cleavage site, the SNP can be identified by alterations in restriction enzyme digestion patterns, and the corresponding changes in nucleic acid fragment lengths determined by gel electrophoresis

[0165] SNP genotyping can include the steps of, for example, collecting a biological sample from a human subject (e.g., sample of tissues, cells, fluids, secretions, etc.), isolating nucleic acids (e.g., genomic DNA, mRNA or both) from the cells of the sample, contacting the nucleic acids with one or more primers which specifically hybridize to a region of the isolated nucleic acid containing a target SNP under conditions such that hybridization and amplification of the target nucleic acid region occurs, and determining the nucleotide present at the SNP position of interest, or, in some assays, detecting the presence or absence of an amplification product (assays can be designed so that hybridization and/or amplification will only occur if a particular SNP allele is present or absent). In some assays, the size of the amplification product is detected and compared to the length of a control sample; for example, deletions and insertions can be detected by a change in size of the amplified product compared to a normal genotype.

[0166] SNP genotyping is useful for numerous practical applications, as described below. Examples of such applications include, but are not limited to, SNP-disease association analysis, disease predisposition screening, disease diagnosis, disease prognosis, disease progression monitoring, determining therapeutic strategies based on an individual's genotype ("pharmacogenomics"), developing therapeutic agents based on SNP genotypes associated with a disease or likelihood of responding to a drug, stratifying a patient population for clinical trial for a treatment regimen, predicting

the likelihood that an individual will experience toxic side effects from a therapeutic agent, and human identification applications such as forensics.

Analysis of Genetic Association Between SNPs and Phenotypic Traits

[0167] SNP genotyping for disease diagnosis, disease predisposition screening, disease prognosis, determining drug responsiveness (pharmacogenomics), drug toxicity screening, and other uses described herein, typically relies on initially establishing a genetic association between one or more specific SNPs and the particular phenotypic traits of interest.

[0168] Different study designs may be used for genetic association studies (Modern Epidemiology, Lippincott Williams & Wilkins (1998), 609-622). Observational studies are most frequently carried out in which the response of the patients is not interfered with. The first type of observational study identifies a sample of persons in whom the suspected cause of the disease is present and another sample of persons in whom the suspected cause is absent, and then the frequency of development of disease in the two samples is compared. These sampled populations are called cohorts, and the study is a prospective study. The other type of observational study is case-control or a retrospective study. In typical case-control studies, samples are collected from individuals with the phenotype of interest (cases) such as certain manifestations of a disease, and from individuals without the phenotype (controls) in a population (target population) that conclusions are to be drawn from. Then the possible causes of the disease are investigated retrospectively. As the time and costs of collecting samples in case-control studies are considerably less than those for prospective studies, case-control studies are the more commonly used study design in genetic association studies, at least during the exploration and discovery stage.

[0169] In both types of observational studies, there may be potential confounding factors that should be taken into consideration. Confounding factors are those that are associated with both the real cause(s) of the disease and the disease itself, and they include demographic information such as age, gender, ethnicity as well as environmental factors. When confounding factors are not matched in cases and controls in a study, and are not controlled properly, spurious association results can arise. If potential confounding factors are identified, they should be controlled for by analysis methods explained below.

[0170] In a genetic association study, the cause of interest to be tested is a certain allele or a SNP or a combination of alleles or a haplotype from several SNPs. Thus, tissue specimens (e.g., whole blood) from the sampled individuals may be collected and genomic DNA genotyped for the SNP(s) of interest. In addition to the phenotypic trait of interest, other information such as demographic (e.g., age, gender, ethnicity, etc.), clinical, and environmental information that may influence the outcome of the trait can be collected to further characterize and define the sample set. In many cases, these factors are known to be associated with diseases and/or SNP allele frequencies. There are likely gene-environment and/or gene-gene interactions as well. Analysis methods to address gene-environment and gene-gene interactions (for example, the effects of the presence of both susceptibility alleles at two different genes can be greater than the effects of the individual alleles at two genes combined) are discussed below.

[0171] After all the relevant phenotypic and genotypic information has been obtained, statistical analyses are carried out to determine if there is any significant correlation between the presence of an allele or a genotype with the phenotypic characteristics of an individual. Preferably, data inspection and cleaning are first performed before carrying out statistical tests for genetic association. Epidemiological and clinical data of the samples can be summarized by descriptive statistics with tables and graphs. Data validation is preferably performed to check for data completion, inconsistent entries, and outliers. Chi-squared tests and t-tests (Wilcoxon rank-sum tests if distributions are not normal) may then be used to check for significant differences between cases and controls for discrete and continuous variables, respectively. To ensure genotyping quality, Hardy-Weinberg disequilibrium tests can be performed on cases and controls separately. Significant deviation from Hardy-Weinberg equilibrium (HWE) in both cases and controls for individual markers can be indicative of genotyping errors. If HWE is violated in a majority of markers, it is indicative of population substructure that should be further investigated. Moreover, Hardy-Weinberg disequilibrium in cases only can indicate genetic association of the markers with the disease (Genetic Data Analysis, Weir B., Sinauer (1990)).

[0172] To test whether an allele of a single SNP is associated with the case or control status of a phenotypic trait, one skilled in the art can compare allele frequencies in cases and controls. Standard chi-squared tests and Fisher exact tests can be carried out on a 2x2 table (2 SNP alleles x 2 outcomes in the categorical trait of interest). To test whether genotypes of a SNP are associated, chi-squared tests can be carried out on a 3x2 table (3 genotypes x 2 outcomes). Score tests are also carried out for genotypic association to contrast the three genotypic frequencies (major homozygotes, heterozygotes and minor homozygotes) in cases and controls, and to look for trends using 3 different modes of inheritance, namely dominant (with contrast coefficients 2, -1, -1), additive (with contrast coefficients 1, 0, -1) and recessive (with contrast coefficients 1, 1, -2). Odds ratios for minor versus major alleles, and odds ratios for heterozygote and homozygote variants versus the wild type genotypes are calculated with the desired confidence limits, usually 95%.

[0173] In order to control for confounders and to test for interaction and effect modifiers, stratified analyses may be performed using stratified factors that are likely to be confounding, including demographic information such as age,

ethnicity, and gender, or an interacting element or effect modifier, such as a known major gene (e.g., APOE for Alzheimer's disease or HLA genes for autoimmune diseases), or environmental factors such as smoking in lung cancer. Stratified association tests may be carried out using Cochran-Mantel-Haenszel tests that take into account the ordinal nature of genotypes with 0, 1, and 2 variant alleles. Exact tests by StatXact may also be performed when computationally possible.

Another way to adjust for confounding effects and test for interactions is to perform stepwise multiple logistic regression analysis using statistical packages such as SAS or R. Logistic regression is a model-building technique in which the best fitting and most parsimonious model is built to describe the relation between the dichotomous outcome (for instance, getting a certain disease or not) and a set of independent variables (for instance, genotypes of different associated genes, and the associated demographic and environmental factors). The most common model is one in which the logit transformation of the odds ratios is expressed as a linear combination of the variables (main effects) and their cross-product terms (interactions) (Applied Logistic Regression, Hosmer and Lemeshow, Wiley (2000)). To test whether a certain variable or interaction is significantly associated with the outcome, coefficients in the model are first estimated and then tested for statistical significance of their departure from zero.

[0174] In addition to performing association tests one marker at a time, haplotype association analysis may also be performed to study a number of markers that are closely linked together. Haplotype association tests can have better power than genotypic or allelic association tests when the tested markers are not the disease-causing mutations themselves but are in linkage disequilibrium with such mutations. The test will even be more powerful if the disease is indeed caused by a combination of alleles on a haplotype (e.g., APOE is a haplotype formed by 2 SNPs that are very close to each other). In order to perform haplotype association effectively, marker-marker linkage disequilibrium measures, both D' and R^2 , are typically calculated for the markers within a gene to elucidate the haplotype structure. Recent studies (Daly et al, Nature Genetics, 29, 232-235, 2001) in linkage disequilibrium indicate that SNPs within a gene are organized in block pattern, and a high degree of linkage disequilibrium exists within blocks and very little linkage disequilibrium exists between blocks. Haplotype association with the disease status can be performed using such blocks once they have been elucidated.

[0175] Haplotype association tests can be carried out in a similar fashion as the allelic and genotypic association tests. Each haplotype in a gene is analogous to an allele in a multi-allelic marker. One skilled in the art can either compare the haplotype frequencies in cases and controls or test genetic association with different pairs of haplotypes. It has been proposed (Schaid et al, Am. J. Hum. Genet., 70, 425-434, 2002) that score tests can be done on haplotypes using the program "haplo.score". In that method, haplotypes are first inferred by EM algorithm and score tests are carried out with a generalized linear model (GLM) framework that allows the adjustment of other factors.

[0176] An important decision in the performance of genetic association tests is the determination of the significance level at which significant association can be declared when the p-value of the tests reaches that level. In an exploratory analysis where positive hits will be followed up in subsequent confirmatory testing, an unadjusted p-value < 0.1 (a significance level on the lenient side) may be used for generating hypotheses for significant association of a SNP with certain phenotypic characteristics of a disease. It is preferred that a p-value < 0.05 (a significance level traditionally used in the art) is achieved in order for a SNP to be considered to have an association with a disease. It is more preferred that a p-value < 0.01 (a significance level on the stringent side) is achieved for an association to be declared. When hits are followed up in confirmatory analyses in more samples of the same source or in different samples from different sources, adjustment for multiple testing will be performed as to avoid excess number of hits while maintaining the experiment-wise error rates at 0.05. While there are different methods to adjust for multiple testing to control for different kinds of error rates, a commonly used but rather conservative method is Bonferroni correction to control the experiment-wise or family-wise error rate (Multiple comparisons and multiple tests, Westfall et al, SAS Institute (1999)). Permutation tests to control for the false discovery rates, FDR, can be more powerful (Benjamini and Hochberg, Journal of the Royal Statistical Society, Series B 57, 1289-1300, 1995, Resampling-based Multiple Testing, Westfall and Young, Wiley (1993)). Such methods to control for multiplicity would be preferred when the tests are dependent and controlling for false discovery rates is sufficient as opposed to controlling for the experiment-wise error rates.

[0177] In replication studies using samples from different populations after statistically significant markers have been identified in the exploratory stage, meta-analyses can then be performed by combining evidence of different studies (Modern Epidemiology, Lippincott Williams & Wilkins, 1998, 643-673). If available, association results known in the art for the same SNPs can be included in the meta-analyses.

[0178] Since both genotyping and disease status classification can involve errors, sensitivity analyses may be performed to see how odds ratios and p-values would change upon various estimates on genotyping and disease classification error rates.

[0179] It has been well known that subpopulation-based sampling bias between cases and controls can lead to spurious results in case-control association studies (Ewens and Spielman, Am. J. Hum. Genet. 62, 450-458, 1995) when prevalence of the disease is associated with different subpopulation groups. Such bias can also lead to a loss of statistical power in genetic association studies. To detect population stratification, Pritchard and Rosenberg (Pritchard et al. Am. J. Hum. Gen. 1999, 65:220-228) suggested typing markers that are unlinked to the disease and using results of association

tests on those markers to determine whether there is any population stratification. When stratification is detected, the genomic control (GC) method as proposed by Devlin and Roeder (Devlin et al. *Biometrics* 1999, 55:997-1004) can be used to adjust for the inflation of test statistics due to population stratification. GC method is robust to changes in population structure levels as well as being applicable to DNA pooling designs (Devlin et al. *Genet. Epidem.* 20001, 21:273-284).

[0180] While Pritchard's method recommended using 15-20 unlinked microsatellite markers, it suggested using more than 30 biallelic markers to get enough power to detect population stratification. For the GC method, it has been shown (Bacanu et al. *Am. J. Hum. Genet.* 2000, 66:1933-1944) that about 60-70 biallelic markers are sufficient to estimate the inflation factor for the test statistics due to population stratification. Hence, 70 intergenic SNPs can be chosen in unlinked regions as indicated in a genome scan (Kehoe et al. *Hum. Mol. Genet.* 1999, 8:237-245).

[0181] Once individual risk factors, genetic or non-genetic, have been found for the predisposition to disease, the next step is to set up a classification/prediction scheme to predict the category (for instance, disease or no-disease) that an individual will be in depending on his genotypes of associated SNPs and other non-genetic risk factors. Logistic regression for discrete trait and linear regression for continuous trait are standard techniques for such tasks (Applied Regression Analysis, Draper and Smith, Wiley (1998)). Moreover, other techniques can also be used for setting up classification. Such techniques include, but are not limited to, MART, CART, neural network, and discriminant analyses that are suitable for use in comparing the performance of different methods (The Elements of Statistical Learning, Hastie, Tibshirani & Friedman, Springer (2002)).

Disease Diagnosis and Predisposition Screening

[0182] Information on association/correlation between genotypes and disease-related phenotypes can be exploited in several ways. For example, in the case of a highly statistically significant association between one or more SNPs with predisposition to a disease for which treatment is available, detection of such a genotype pattern in an individual may justify immediate administration of treatment, or at least the institution of regular monitoring of the individual. Detection of the susceptibility alleles associated with serious disease in a couple contemplating having children may also be valuable to the couple in their reproductive decisions. In the case of a weaker but still statistically significant association between a SNP and a human disease, immediate therapeutic intervention or monitoring may not be justified after detecting the susceptibility allele or SNP. Nevertheless, the subject can be motivated to begin simple life-style changes (e.g., diet, exercise) that can be accomplished at little or no cost to the individual but would confer potential benefits in reducing the risk of developing conditions for which that individual may have an increased risk by virtue of having the susceptibility allele(s).

[0183] The SNP used in the method of the invention may contribute to myocardial infarction in an individual in different ways. Some polymorphisms occur within a protein coding sequence and contribute to disease phenotype by affecting protein structure. Other polymorphisms occur in noncoding regions but may exert phenotypic effects indirectly via influence on, for example, replication, transcription, and/or translation. A single SNP may affect more than one phenotypic trait. Likewise, a single phenotypic trait may be affected by multiple SNPs in different genes.

[0184] As used herein, the terms "diagnose", "diagnosis", and "diagnostics" include, but are not limited to any of the following: detection of myocardial infarction that an individual may presently have or be at risk for, predisposition screening (i.e., determining the increased risk for an individual in developing myocardial infarction in the future, or determining whether an individual has a decreased risk of developing myocardial infarction in the future; in the case of recurrent myocardial infarction (RMI), predisposition screening may typically involve determining the risk that an individual who has previously had a myocardial infarction will develop another myocardial infarction in the future, or determining whether an individual who has not experienced a myocardial infarction will be at risk for developing recurrent myocardial infarctions in the future), determining a particular type or subclass of myocardial infarction in an individual known to have myocardial infarction, confirming or reinforcing a previously made diagnosis of myocardial infarction, pharmacogenomic evaluation of an individual to determine which therapeutic strategy that individual is most likely to positively respond to or to predict whether a patient is likely to respond to a particular treatment, predicting whether a patient is likely to experience toxic effects from a particular treatment or therapeutic compound, and evaluating the future prognosis of an individual having myocardial infarction. Such diagnostic uses are based on the SNPs individually or in a unique combination or SNP haplotypes disclosed herein.

[0185] Haplotypes are particularly useful in that, for example, fewer SNPs can be genotyped to determine if a particular genomic region harbors a locus that influences a particular phenotype, such as in linkage disequilibrium-based SNP association analysis.

[0186] Linkage disequilibrium (LD) refers to the co-inheritance of alleles (e.g., alternative nucleotides) at two or more different SNP sites at frequencies greater than would be expected from the separate frequencies of occurrence of each allele in a given population. The expected frequency of co-occurrence of two alleles that are inherited independently is the frequency of the first allele multiplied by the frequency of the second allele. Alleles that co-occur at expected fre-

quencies are said to be in "linkage equilibrium". In contrast, LD refers to any non-random genetic association between allele(s) at two or more different SNP sites, which is generally due to the physical proximity of the two loci along a chromosome. LD can occur when two or more SNPs sites are in close physical proximity to each other on a given chromosome and therefore alleles at these SNP sites will tend to remain unseparated for multiple generations with the consequence that a particular nucleotide (allele) at one SNP site will show a non-random association with a particular nucleotide (allele) at a different SNP site located nearby. Hence, genotyping one of the SNP sites will give almost the same information as genotyping the other SNP site that is in LD.

[0187] For diagnostic purposes, if a particular SNP site is found to be useful for diagnosing myocardial infarction, then the skilled artisan would recognize that other SNP sites which are in LD with this SNP site would also be useful for diagnosing the condition. Various degrees of LD can be encountered between two or more SNPs with the result being that some SNPs are more closely associated (i.e., in stronger LD) than others. Furthermore, the physical distance over which LD extends along a chromosome differs between different regions of the genome, and therefore the degree of physical separation between two or more SNP sites necessary for LD to occur can differ between different regions of the genome.

[0188] For diagnostic applications, polymorphisms (e.g., SNPs and/or haplotypes) that are not the actual disease-causing (causative) polymorphisms, but are in LD with such causative polymorphisms, are also useful. In such instances, the genotype of the polymorphism(s) that is/are in LD with the causative polymorphism is predictive of the genotype of the causative polymorphism and, consequently, predictive of the phenotype (e.g., myocardial infarction) that is influenced by the causative SNP(s). Thus, polymorphic markers that are in LD with causative polymorphisms are useful as diagnostic markers, and are particularly useful when the actual causative polymorphism(s) is/are unknown.

[0189] Linkage disequilibrium in the human genome is reviewed in: Wall et al., "Haplotype blocks and linkage disequilibrium in the human genome", *Nat Rev Genet.* 2003 Aug;4(8):587-97; Garner et al., "On selecting markers for association studies: patterns of linkage disequilibrium between two and three diallelic loci", *Genet Epidemiol.* 2003 Jan;24(1):57-67; Ardlie et al., "Patterns of linkage disequilibrium in the human genome", *Nat Rev Genet.* 2002 Apr;3(4):299-309 (erratum in *Nat Rev Genet* 2002 Jul;3(7):566); and Remm et al., "High-density genotyping and linkage disequilibrium in the human genome using chromosome 22 as a model", *Curr Opin Chem Biol.* 2002 Feb;6(1):24-30.

[0190] The contribution or association of particular SNPs and/or SNP haplotypes with disease phenotypes, such as myocardial infarction, enables the SNP used in the method of the present invention to be used to develop superior diagnostic tests capable of identifying individuals who express a detectable trait, such as myocardial infarction, as the result of a specific genotype, or individuals whose genotype places them at an increased or decreased risk of developing a detectable trait at a subsequent time as compared to individuals who do not have that genotype. As described herein, diagnostics may be based on a single SNP or a group of SNPs. Combined detection of a plurality of SNPs (for example, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 24, 25, 30, 32, 48, 50, 64, 96, 100, or any other number in-between, or more, of the SNPs provided in Table 1 and/or Table 2) typically increases the probability of an accurate diagnosis. For example, the presence of a single SNP known to correlate with myocardial infarction might indicate a probability of 20% that an individual has or is at risk of developing myocardial infarction, whereas detection of five SNPs, each of which correlates with myocardial infarction, might indicate a probability of 80% that an individual has or is at risk of developing myocardial infarction. To further increase the accuracy of diagnosis or predisposition screening, analysis of the SNP used in the method of the present invention can be combined with that of other polymorphisms or other risk factors of myocardial infarction, such as disease symptoms, pathological characteristics, family history, diet, environmental factors or lifestyle factors.

[0191] It will, of course, be understood by practitioners skilled in the treatment or diagnosis of myocardial infarction that the present invention generally does not intend to provide an absolute identification of individuals who are at risk (or less at risk) of developing myocardial infarction, and/or pathologies related to myocardial infarction, but rather to indicate a certain increased (or decreased) degree or likelihood of developing the disease based on statistically significant association results. However, this information is extremely valuable as it can be used to, for example, initiate preventive treatments or to allow an individual carrying one or more significant SNPs or SNP haplotypes to foresee warning signs such as minor clinical symptoms, or to have regularly scheduled physical exams to monitor for appearance of a condition in order to identify and begin treatment of the condition at an early stage. Particularly with diseases that are extremely debilitating or fatal if not treated on time, the knowledge of a potential predisposition, even if this predisposition is not absolute, would likely contribute in a very significant manner to treatment efficacy.

[0192] The diagnostic techniques of the present invention may employ a variety of methodologies to determine whether a test subject has a SNP or a SNP pattern associated with an increased or decreased risk of developing a detectable trait or whether the individual suffers from a detectable trait as a result of a particular polymorphism/mutation, including, for example, methods which enable the analysis of individual chromosomes for haplotyping, family studies, single sperm DNA analysis, or somatic hybrids. The trait analyzed using the diagnostics of the invention may be any detectable trait that is commonly observed in pathologies and disorders related to myocardial infarction.

[0193] Another aspect of the present invention relates to a method of determining whether an individual is at risk (or

less at risk) of developing one or more traits or whether an individual expresses one or more traits as a consequence of possessing a particular trait-causing or trait-influencing allele. These methods generally involve obtaining a nucleic acid sample from an individual and assaying the nucleic acid sample to determine which nucleotide(s) is/are present at one or more SNP positions, wherein the assayed nucleotide(s) is/are indicative of an increased or decreased risk of developing the trait or indicative that the individual expresses the trait as a result of possessing a particular trait-causing or trait-influencing allele.

[0194] In another embodiment, the SNP detection reagents used in the method of the present invention are used to determine whether an individual has one or more SNP allele(s) affecting the level (e.g., the concentration of mRNA or protein in a sample, etc.) or pattern (e.g., the kinetics of expression, rate of decomposition, stability profile, K_m , V_{max} , etc.) of gene expression (collectively, the "gene response" of a cell or bodily fluid). Such a determination can be accomplished by screening for mRNA or protein expression (e.g., by using nucleic acid arrays, RT-PCR, TaqMan assays, or mass spectrometry), identifying genes having altered expression in an individual, genotyping SNPs disclosed in Table 1 and/or Table 2 that could affect the expression of the genes having altered expression (e.g., SNPs that are in and/or around the gene(s) having altered expression, SNPs in regulatory/control regions, SNPs in and/or around other genes that are involved in pathways that could affect the expression of the gene(s) having altered expression, or all SNPs could be genotyped), and correlating SNP genotypes with altered gene expression. In this manner, specific SNP alleles at particular SNP sites can be identified that affect gene expression.

Pharmacogenomics and Therapeutics/Drug Development

[0195] Disclosed herein are methods for assessing the pharmacogenomics of a subject harboring particular SNP alleles or haplotypes to a particular therapeutic agent or pharmaceutical compound, or to a class of such compounds. Pharmacogenomics deals with the roles which clinically significant hereditary variations (e.g., SNPs) play in the response to drugs due to altered drug disposition and/or abnormal action in affected persons. See, e.g., Roses, *Nature* 405, 857-865 (2000); Gould Rothberg, *Nature Biotechnology* 19, 209-211 (2001); Eichelbaum, *Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996); and Linder, *Clin. Chem.* 43(2):254-266 (1997). The clinical outcomes of these variations can result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the SNP genotype of an individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. For example, SNPs in drug metabolizing enzymes can affect the activity of these enzymes, which in turn can affect both the intensity and duration of drug action, as well as drug metabolism and clearance.

[0196] The discovery of SNPs in drug metabolizing enzymes, drug transporters, proteins for pharmaceutical agents, and other drug targets has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. SNPs can be expressed in the phenotype of the extensive metabolizer and in the phenotype of the poor metabolizer. Accordingly, SNPs may lead to allelic variants of a protein in which one or more of the protein functions in one population are different from those in another population. SNPs and the encoded variant peptides thus provide targets to ascertain a genetic predisposition that can affect treatment modality. For example, in a ligand-based treatment, SNPs may give rise to amino terminal extracellular domains and/or other ligand-binding regions of a receptor that are more or less active in ligand binding, thereby affecting subsequent protein activation. Accordingly, ligand dosage would necessarily be modified to maximize the therapeutic effect within a given population containing particular SNP alleles or haplotypes.

[0197] As an alternative to genotyping, specific variant proteins containing variant amino acid sequences encoded by alternative SNP alleles could be identified. Thus, pharmacogenomic characterization of an individual permits the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic uses based on the individual's SNP genotype, thereby enhancing and optimizing the effectiveness of the therapy. Furthermore, the production of recombinant cells and transgenic animals containing particular SNPs/haplotypes allow effective clinical design and testing of treatment compounds and dosage regimens. For example, transgenic animals can be produced that differ only in specific SNP alleles in a gene that is orthologous to a human disease susceptibility gene.

[0198] Pharmacogenomic uses of the SNP used in the method of the present invention provide several significant advantages for patient care, particularly in treating myocardial infarction. Pharmacogenomic characterization of an individual, based on an individual's SNP genotype, can identify those individuals unlikely to respond to treatment with a particular medication and thereby allows physicians to avoid prescribing the ineffective medication to those individuals. On the other hand, SNP genotyping of an individual may enable physicians to select the appropriate medication and dosage regimen that will be most effective based on an individual's SNP genotype. This information increases a physician's confidence in prescribing medications and motivates patients to comply with their drug regimens. Furthermore, pharmacogenomics may identify patients predisposed to toxicity and adverse reactions to particular drugs or drug dosages. Adverse drug reactions lead to more than 100,000 avoidable deaths per year in the United States alone and therefore represent a significant cause of hospitalization and death, as well as a significant economic burden on the

healthcare system (Pfost et. al., Trends in Biotechnology, Aug. 2000.). Thus, pharmacogenomics based on the SNPs disclosed herein has the potential to both save lives and reduce healthcare costs substantially.

[0199] Pharmacogenomics in general is discussed further in Rose et al., "Pharmacogenetic analysis of clinically relevant genetic polymorphisms", *Methods Mol Med.* 2003;85:225-37. Pharmacogenomics as it relates to Alzheimer's disease and other neurodegenerative disorders is discussed in Cacabelos, "Pharmacogenomics for the treatment of dementia", *Ann Med.* 2002;34(5):357-79, Maimone et al., "Pharmacogenomics of neurodegenerative diseases", *Eur J Pharmacol.* 2001 Feb 9;413(1):11-29, and Poirier, "Apolipoprotein E: a pharmacogenetic target for the treatment of Alzheimer's disease", *Mol Diagn.* 1999 Dec;4(4):335-41. Pharmacogenomics as it relates to cardiovascular disorders is discussed in Siest et al., "Pharmacogenomics of drugs affecting the cardiovascular system", *Clin Chem Lab Med.* 2003 Apr;41(4):590-9, Mukherjee et al., "Pharmacogenomics in cardiovascular diseases", *Prog Cardiovasc Dis.* 2002 May-Jun;44(6):479-98, and Mooser et al., "Cardiovascular pharmacogenetics in the SNP era", *J Thromb Haemost.* 2003 Jul;1(7):1398-402. Pharmacogenomics as it relates to cancer is discussed in McLeod et al., "Cancer pharmacogenomics: SNPs, chips, and the individual patient", *Cancer Invest.* 2003;21(4):630-40 and Watters et al., "Cancer pharmacogenomics: current and future applications", *Biochim Biophys Acta.* 2003 Mar 17;1603(2):99-111.

[0200] The SNP used in the method of the present invention also can be used to identify novel therapeutic targets for myocardial infarction. For example, genes containing the disease-associated variants ("variant genes") or their products, as well as genes or their products that are directly or indirectly regulated by or interacting with these variant genes or their products, can be targeted for the development of therapeutics that, for example, treat the disease or prevent or delay disease onset. The therapeutics may be composed of, for example, small molecules, proteins, protein fragments or peptides, antibodies, nucleic acids, or their derivatives or mimetics which modulate the functions or levels of the target genes or gene products.

[0201] The SNP-containing nucleic acid molecules disclosed herein, and their complementary nucleic acid molecules, may be used as antisense constructs to control gene expression in cells, tissues, and organisms. Antisense technology is well established in the art and extensively reviewed in *Antisense Drug Technology: Principles, Strategies, and Applications*, Crooke (ed.), Marcel Dekker, Inc.: New York (2001). An antisense nucleic acid molecule is generally designed to be complementary to a region of mRNA expressed by a gene so that the antisense molecule hybridizes to the mRNA and thereby blocks translation of mRNA into protein. Various classes of antisense oligonucleotides are used in the art, two of which are cleavers and blockers. Cleavers, by binding to target RNAs, activate intracellular nucleases (e.g., RNaseH or RNase L) that cleave the target RNA. Blockers, which also bind to target RNAs, inhibit protein translation through steric hindrance of ribosomes. Exemplary blockers include peptide nucleic acids, morpholinos, locked nucleic acids, and methylphosphonates (see, e.g., Thompson, *Drug Discovery Today*, 7 (17): 912-917 (2002)). Antisense oligonucleotides are directly useful as therapeutic agents, and are also useful for determining and validating gene function (e.g., in gene knock-out or knock-down experiments).

[0202] Antisense technology is further reviewed in: Lavery et al., "Antisense and RNAi: powerful tools in drug target discovery and validation", *Curr Opin Drug Discov Devel.* 2003 Jul;6(4):561-9; Stephens et al., "Antisense oligonucleotide therapy in cancer", *Curr Opin Mol Ther.* 2003 Apr;5(2):118-22; Kurreck, "Antisense technologies. Improvement through novel chemical modifications", *Eur J Biochem.* 2003 Apr;270(8):1628-44; Dias et al., "Antisense oligonucleotides: basic concepts and mechanisms", *Mol Cancer Ther.* 2002 Mar;1(5):347-55; Chen, "Clinical development of antisense oligonucleotides as anti-cancer therapeutics", *Methods Mol Med.* 2003;75:621-36; Wang et al., "Antisense anticancer oligonucleotide therapeutics", *Curr Cancer Drug Targets.* 2001 Nov;1(3):177-96; and Bennett, "Efficiency of antisense oligonucleotide drug discovery", *Antisense Nucleic Acid Drug Dev.* 2002 Jun;12(3):215-24.

[0203] The SNP used in the method of the present invention are particularly useful for designing antisense reagents that are specific for particular nucleic acid variants. Based on the SNP information disclosed herein, antisense oligonucleotides can be produced that specifically target mRNA molecules that contain one or more particular SNP nucleotides. In this manner, expression of mRNA molecules that contain one or more undesired polymorphisms (e.g., SNP nucleotides that lead to a defective protein such as an amino acid substitution in a catalytic domain) can be inhibited or completely blocked. Thus, antisense oligonucleotides can be used to specifically bind a particular polymorphic form (e.g., a SNP allele that encodes a defective protein), thereby inhibiting translation of this form, but which do not bind an alternative polymorphic form (e.g., an alternative SNP nucleotide that encodes a protein having normal function).

[0204] Antisense molecules can be used to inactivate mRNA in order to inhibit gene expression and production of defective proteins. Accordingly, these molecules can be used to treat a disorder, such as myocardial infarction, characterized by abnormal or undesired gene expression or expression of certain defective proteins. This technique can involve cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible mRNA regions include, for example, protein-coding regions and particularly protein-coding regions corresponding to catalytic activities, substrate/ligand binding, or other functional activities of a protein.

[0205] The SNP used in the method of the present invention are also useful for designing RNA interference reagents that specifically target nucleic acid molecules having particular SNP variants. RNA interference (RNAi), also referred to

as gene silencing, is based on using double-stranded RNA (dsRNA) molecules to turn genes off. When introduced into a cell, dsRNAs are processed by the cell into short fragments (generally about 21-22bp in length) known as small interfering RNAs (siRNAs) which the cell uses in a sequence-specific manner to recognize and destroy complementary RNAs (Thompson, *Drug Discovery Today*, 7 (17): 912-917 (2002)). Thus, because RNAi molecules, including siRNAs, act in a sequence-specific manner, the SNP used in the method of the present invention can be used to design RNAi reagents that recognize and destroy nucleic acid molecules having specific SNP alleles/nucleotides (such as deleterious alleles that lead to the production of defective proteins), while not affecting nucleic acid molecules having alternative SNP alleles (such as alleles that encode proteins having normal function). As with antisense reagents, RNAi reagents may be directly useful as therapeutic agents (e.g., for turning off defective, disease-causing genes), and are also useful for characterizing and validating gene function (e.g., in gene knock-out or knock-down experiments).

[0206] The following references provide a further review of RNAi: Agami, "RNAi and related mechanisms and their potential use for therapy", *Curr Opin Chem Biol.* 2002 Dec;6(6):829-34; Lavery et al., "Antisense and RNAi: powerful tools in drug target discovery and validation", *Curr Opin Drug Discov Devel.* 2003 Jul;6(4):561-9; Shi, "Mammalian RNAi for the masses", *Trends Genet* 2003 Jan;19(1):9-12; Shuey et al., "RNAi: gene-silencing in therapeutic intervention", *Drug Discovery Today* 2002 Oct;7(20):1040-1046; McManus et al., *Nat Rev Genet* 2002 Oct;3(10):737-47; Xia et al., *Nat Biotechnol* 2002 Oct;20(10):1006-10; Plasterk et al., *Curr Opin Genet Dev* 2000 Oct;10(5):562-7; Boshier et al., *Nat Cell Biol* 2000 Feb;2(2):E31-6; and Hunter, *Curr Biol* 1999 Jun 17;9(12):R440-2).

[0207] A subject suffering from a pathological condition, such as myocardial infarction, ascribed to a SNP may be treated so as to correct the genetic defect (see Kren et al., *Proc. Natl. Acad. Sci. USA* 96:10349-10354 (1999)). Such a subject can be identified by any method that can detect the polymorphism in a biological sample drawn from the subject. Such a genetic defect may be permanently corrected by administering to such a subject a nucleic acid fragment incorporating a repair sequence that supplies the normal/wild-type nucleotide at the position of the SNP. This site-specific repair sequence can encompass an RNA/DNA oligonucleotide that operates to promote endogenous repair of a subject's genomic DNA. The site-specific repair sequence is administered in an appropriate vehicle, such as a complex with polyethylenimine, encapsulated in anionic liposomes, a viral vector such as an adenovirus, or other pharmaceutical composition that promotes intracellular uptake of the administered nucleic acid. A genetic defect leading to an inborn pathology may then be overcome, as the chimeric oligonucleotides induce incorporation of the normal sequence into the subject's genome. Upon incorporation, the normal gene product is expressed, and the replacement is propagated, thereby engendering a permanent repair and therapeutic enhancement of the clinical condition of the subject.

[0208] In cases in which a cSNP results in a variant protein that is ascribed to be the cause of, or a contributing factor to, a pathological condition, a method of treating such a condition can include administering to a subject experiencing the pathology the wild-type/normal cognate of the variant protein. Once administered in an effective dosing regimen, the wild-type cognate provides complementation or remediation of the pathological condition.

[0209] Disclosed herein is also a method for identifying a compound or agent that can be used to treat myocardial infarction. The SNPs disclosed herein are useful as targets for the identification and/or development of therapeutic agents. A method for identifying a therapeutic agent or compound typically includes assaying the ability of the agent or compound to modulate the activity and/or expression of a SNP-containing nucleic acid or the encoded product and thus identifying an agent or a compound that can be used to treat a disorder characterized by undesired activity or expression of the SNP-containing nucleic acid or the encoded product. The assays can be performed in cell-based and cell-free systems. Cell-based assays can include cells naturally expressing the nucleic acid molecules of interest or recombinant cells genetically engineered to express certain nucleic acid molecules.

[0210] Variant gene expression in a myocardial infarction patient can include, for example, either expression of a SNP-containing nucleic acid sequence (for instance, a gene that contains a SNP can be transcribed into an mRNA transcript molecule containing the SNP, which can in turn be translated into a variant protein) or altered expression of a normal/wild-type nucleic acid sequence due to one or more SNPs (for instance, a regulatory/control region can contain a SNP that affects the level or pattern of expression of a normal transcript).

[0211] Assays for variant gene expression can involve direct assays of nucleic acid levels (e.g., mRNA levels), expressed protein levels, or of collateral compounds involved in a signal pathway. Further, the expression of genes that are up- or down-regulated in response to the signal pathway can also be assayed. In this embodiment, the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

[0212] Modulators of variant gene expression can be identified in a method wherein, for example, a cell is contacted with a candidate compound/agent and the expression of mRNA determined. The level of expression of mRNA in the presence of the candidate compound is compared to the level of expression of mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of variant gene expression based on this comparison and be used to treat a disorder such as myocardial infarction that is characterized by variant gene expression (e.g., either expression of a SNP-containing nucleic acid or altered expression of a normal/wild-type nucleic acid molecule due to one or more SNPs that affect expression of the nucleic acid molecule) due to one or more SNP used in the method of the present invention. When expression of mRNA is statistically significantly greater in the presence of the candidate

compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

[0213] Further, disclosed herein are methods of treatment, with the SNP or associated nucleic acid domain (e.g., catalytic domain, ligand/substrate-binding domain, regulatory/control region, etc.) or gene, or the encoded mRNA transcript, as a target, using a compound identified through drug screening as a gene modulator to modulate variant nucleic acid expression. Modulation can include either up-regulation (i.e., activation or agonization) or down-regulation (i.e., suppression or antagonization) of nucleic acid expression.

[0214] Expression of mRNA transcripts and encoded proteins, either wild type or variant, may be altered in individuals with a particular SNP allele in a regulatory/control element, such as a promoter or transcription factor binding domain, that regulates expression. In this situation, methods of treatment and compounds can be identified, as discussed herein, that regulate or overcome the variant regulatory/control element, thereby generating normal, or healthy, expression levels of either the wild type or variant protein.

[0215] The SNP-containing nucleic acid molecules for use in the method of the present invention are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of a variant gene, or encoded product, in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as an indicator for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance, as well as an indicator for toxicities. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

[0216] In another aspect disclosed herein is a pharmaceutical pack comprising a therapeutic agent (e.g., a small molecule drug, antibody, peptide, antisense or RNAi nucleic acid molecule, etc.) and a set of instructions for administration of the therapeutic agent to humans diagnostically tested for one or more SNPs or SNP haplotypes disclosed herein.

[0217] The SNPs/haplotypes disclosed herein are also useful for improving many different aspects of the drug development process. For example, individuals can be selected for clinical trials based on their SNP genotype. Individuals with SNP genotypes that indicate that they are most likely to respond to the drug can be included in the trials and those individuals whose SNP genotypes indicate that they are less likely to or would not respond to the drug, or suffer adverse reactions, can be eliminated from the clinical trials. This not only improves the safety of clinical trials, but also will enhance the chances that the trial will demonstrate statistically significant efficacy. Furthermore, the SNPs may explain why certain previously developed drugs performed poorly in clinical trials and may help identify a subset of the population that would benefit from a drug that had previously performed poorly in clinical trials, thereby "rescuing" previously developed drugs, and enabling the drug to be made available to a particular myocardial infarction patient population that can benefit from it.

[0218] SNPs have many important uses in drug discovery, screening, and development. A high probability exists that, for any gene/protein selected as a potential drug target, variants of that gene/protein will exist in a patient population. Thus, determining the impact of gene/protein variants on the selection and delivery of a therapeutic agent should be an integral aspect of the drug discovery and development process. (Jazwinska, A Trends Guide to Genetic Variation and Genomic Medicine, 2002 Mar; S30-S36).

[0219] Knowledge of variants (e.g., SNPs and any corresponding amino acid polymorphisms) of a particular therapeutic target (e.g., a gene, mRNA transcript, or protein) enables parallel screening of the variants in order to identify therapeutic candidates (e.g., small molecule compounds, antibodies, antisense or RNAi nucleic acid compounds, etc.) that demonstrate efficacy across variants (Rothberg, Nat Biotechnol 2001 Mar;19(3):209-11). Such therapeutic candidates would be expected to show equal efficacy across a larger segment of the patient population, thereby leading to a larger potential market for the therapeutic candidate.

[0220] Furthermore, identifying variants of a potential therapeutic target enables the most common form of the target to be used for selection of therapeutic candidates, thereby helping to ensure that the experimental activity that is observed for the selected candidates reflects the real activity expected in the largest proportion of a patient population (Jazwinska, A Trends Guide to Genetic Variation and Genomic Medicine, 2002 Mar; S30-S36).

[0221] Additionally, screening therapeutic candidates against all known variants of a target can enable the early identification of potential toxicities and adverse reactions relating to particular variants. For example, variability in drug absorption, distribution, metabolism and excretion (ADME) caused by, for example, SNPs in therapeutic targets or drug metabolizing genes, can be identified, and this information can be utilized during the drug development process to minimize variability in drug disposition and develop therapeutic agents that are safer across a wider range of a patient population. The SNP used in the method of the present invention, including the variant proteins and encoding polymorphic nucleic acid molecules provided in Tables 1-2, are useful in conjunction with a variety of toxicology methods established in the art, such as those set forth in Current Protocols in Toxicology, John Wiley & Sons, Inc., N.Y.

[0222] Furthermore, therapeutic agents that target any art-known proteins (or nucleic acid molecules, either RNA or DNA) may cross-react with the variant proteins (or polymorphic nucleic acid molecules) disclosed in Table 1, thereby significantly affecting the pharmacokinetic properties of the drug. Consequently, the protein variants and the SNP-containing nucleic acid molecules disclosed in Tables 1-2 are useful in developing, screening, and evaluating therapeutic agents that target corresponding art-known protein forms (or nucleic acid molecules). Additionally, as discussed above, knowledge of all polymorphic forms of a particular drug target enables the design of therapeutic agents that are effective against most or all such polymorphic forms of the drug target.

Pharmaceutical Compositions and Administration Thereof

[0223] Any of the myocardial infarction-associated proteins, and encoding nucleic acid molecules, disclosed herein can be used as therapeutic targets (or directly used themselves as therapeutic compounds) for treating myocardial infarction and related pathologies, and the present disclosure enables therapeutic compounds (e.g., small molecules, antibodies, therapeutic proteins, RNAi and antisense molecules, etc.) to be developed that target (or are comprised of) any of these therapeutic targets.

[0224] In general, a therapeutic compound will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the therapeutic compound i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, and other factors.

[0225] Therapeutically effective amounts of therapeutic compounds may range from, for example, approximately 0.01-50 mg per kilogram body weight of the recipient per day; preferably about 0.1-20 mg/kg/day. Thus, as an example, for administration to a 70 kg person, the dosage range would most preferably be about 7 mg to 1.4 g per day.

[0226] In general, therapeutic compounds will be administered as pharmaceutical compositions by any one of the following routes: oral, systemic (e.g., transdermal, intranasal, or by suppository), or parenteral (e.g., intramuscular, intravenous, or subcutaneous) administration. The preferred manner of administration is oral or parenteral using a convenient daily dosage regimen, which can be adjusted according to the degree of affliction. Oral compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, elixirs, aerosols, or any other appropriate compositions.

[0227] The choice of formulation depends on various factors such as the mode of drug administration (e.g., for oral administration, formulations in the form of tablets, pills, or capsules are preferred) and the bioavailability of the drug substance. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area, i.e., decreasing particle size. For example, U.S. Patent No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a cross-linked matrix of macromolecules. U.S. Patent No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

[0228] Pharmaceutical compositions are comprised of, in general, a therapeutic compound in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the therapeutic compound. Such excipients may be any solid, liquid, semi-solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one skilled in the art.

[0229] Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

[0230] Compressed gases may be used to disperse a compound in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

[0231] Other suitable pharmaceutical excipients and their formulations are described in Remington's Pharmaceutical Sciences, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).

[0232] The amount of the therapeutic compound in a formulation can vary within the full range employed by those skilled in the art. Typically, the formulation will contain, on a weight percent (wt %) basis, from about 0.01-99.99 wt % of the therapeutic compound based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 1-80 wt %.

[0233] Therapeutic compounds can be administered alone or in combination with other therapeutic compounds or in combination with one or more other active ingredient(s). For example, an inhibitor or stimulator of a myocardial infarction-associated protein can be administered in combination with another agent that inhibits or stimulates the activity of

the same or a different myocardial infarction-associated protein to thereby counteract the affects of myocardial infarction.

[0234] For further information regarding pharmacology, see Current Protocols in Pharmacology, John Wiley & Sons, Inc., N.Y.

Human Identification Applications

[0235] In addition to their diagnostic and therapeutic uses in myocardial infarction and related pathologies, the SNPs disclosed herein are also useful as human identification markers for such applications as forensics, paternity testing, and biometrics (see, e.g., Gill, "An assessment of the utility of single nucleotide polymorphisms (SNPs) for forensic purposes", *Int J Legal Med.* 2001;114(4-5):204-10). Genetic variations in the nucleic acid sequences between individuals can be used as genetic markers to identify individuals and to associate a biological sample with an individual. Determination of which nucleotides occupy a set of SNP positions in an individual identifies a set of SNP markers that distinguishes the individual. The more SNP positions that are analyzed, the lower the probability that the set of SNPs in one individual is the same as that in an unrelated individual. Preferably, if multiple sites are analyzed, the sites are unlinked (i.e., inherited independently). Thus, preferred sets of SNPs can be selected from among the SNPs disclosed herein, which may include SNPs on different chromosomes, SNPs on different chromosome arms, and/or SNPs that are dispersed over substantial distances along the same chromosome arm.

[0236] Furthermore, among the SNPs disclosed herein, preferred SNPs for use in certain forensic/human identification applications include SNPs located at degenerate codon positions (i.e., the third position in certain codons which can be one of two or more alternative nucleotides and still encode the same amino acid), since these SNPs do not affect the encoded protein. SNPs that do not affect the encoded protein are expected to be under less selective pressure and are therefore expected to be more polymorphic in a population, which is typically an advantage for forensic/human identification applications. However, for certain forensics/human identification applications, such as predicting phenotypic characteristics (e.g., inferring ancestry or inferring one or more physical characteristics of an individual) from a DNA sample, it may be desirable to utilize SNPs that affect the encoded protein.

[0237] For many of the SNPs disclosed in Tables 1-2 (which are identified as "Applera" SNP source), Tables 1-2 provide SNP allele frequencies obtained by re-sequencing the DNA of chromosomes from 39 individuals (Tables 1-2 also provide allele frequency information for "Celera" source SNPs and, where available, public SNPs from dbEST, HGBASE, and/or HGMD). The allele frequencies provided in Tables 1-2 enable these SNPs to be readily used for human identification applications. Although any SNP disclosed in Table 1 and/or Table 2 could be used for human identification, the closer that the frequency of the minor allele at a particular SNP site is to 50%, the greater the ability of that SNP to discriminate between different individuals in a population since it becomes increasingly likely that two randomly selected individuals would have different alleles at that SNP site. Using the SNP allele frequencies provided in Tables 1-2, one of ordinary skill in the art could readily select a subset of SNPs for which the frequency of the minor allele is, for example, at least 1%, 2%, 5%, 10%, 20%, 25%, 30%, 40%, 45%, or 50%, or any other frequency in-between. Thus, since Tables 1-2 provide allele frequencies based on the re-sequencing of the chromosomes from 39 individuals, a subset of SNPs could readily be selected for human identification in which the total allele count of the minor allele at a particular SNP site is, for example, at least 1, 2, 4, 8, 10, 16, 20, 24, 30, 32, 36, 38, 39, 40, or any other number in-between.

[0238] Furthermore, Tables 1-2 also provide population group (interchangeably referred to herein as ethnic or racial groups) information coupled with the extensive allele frequency information. For example, the group of 39 individuals whose DNA was re-sequenced was made-up of 20 Caucasians and 19 African-Americans. This population group information enables further refinement of SNP selection for human identification. For example, preferred SNPs for human identification can be selected from Tables 1-2 that have similar allele frequencies in both the Caucasian and African-American populations; thus, for example, SNPs can be selected that have equally high discriminatory power in both populations. Alternatively, SNPs can be selected for which there is a statistically significant difference in allele frequencies between the Caucasian and African-American populations (as an extreme example, a particular allele may be observed only in either the Caucasian or the African-American population group but not observed in the other population group); such

[0239] SNPs are useful, for example, for predicting the race/ethnicity of an unknown perpetrator from a biological sample such as a hair or blood stain recovered at a crime scene. For a discussion of using SNPs to predict ancestry from a DNA sample, including statistical methods, see Frudakis et al., "A Classifier for the SNP-Based Inference of Ancestry", *Journal of Forensic Sciences* 2003; 48(4):771-782.

[0240] SNPs have numerous advantages over other types of polymorphic markers, such as short tandem repeats (STRs). For example, SNPs can be easily scored and are amenable to automation, making SNPs the markers of choice for large-scale forensic databases. SNPs are found in much greater abundance throughout the genome than repeat polymorphisms. Population frequencies of two polymorphic forms can usually be determined with greater accuracy than those of multiple polymorphic forms at multi-allelic loci. SNPs are mutationally more stable than repeat polymorphisms. SNPs are not susceptible to artefacts such as stutter bands that can hinder analysis. Stutter bands are frequently encountered when analyzing repeat polymorphisms, and are particularly troublesome when analyzing samples such as

crime scene samples that may contain mixtures of DNA from multiple sources. Another significant advantage of SNP markers over STR markers is the much shorter length of nucleic acid needed to score a SNP. For example, STR markers are generally several hundred base pairs in length. A SNP, on the other hand, comprises a single nucleotide, and generally a short conserved region on either side of the SNP position for primer and/or probe binding. This makes SNPs more amenable to typing in highly degraded or aged biological samples that are frequently encountered in forensic casework in which DNA may be fragmented into short pieces.

[0241] SNPs also are not subject to microvariant and "off-ladder" alleles frequently encountered when analyzing STR loci. Microvariants are deletions or insertions within a repeat unit that change the size of the amplified DNA product so that the amplified product does not migrate at the same rate as reference alleles with normal sized repeat units. When separated by size, such as by electrophoresis on a polyacrylamide gel, microvariants do not align with a reference allelic ladder of standard sized repeat units, but rather migrate between the reference alleles. The reference allelic ladder is used for precise sizing of alleles for allele classification; therefore alleles that do not align with the reference allelic ladder lead to substantial analysis problems. Furthermore, when analyzing multi-allelic repeat polymorphisms, occasionally an allele is found that consists of more or less repeat units than has been previously seen in the population, or more or less repeat alleles than are included in a reference allelic ladder. These alleles will migrate outside the size range of known alleles in a reference allelic ladder, and therefore are referred to as "off-ladder" alleles. In extreme cases, the allele may contain so few or so many repeats that it migrates well out of the range of the reference allelic ladder. In this situation, the allele may not even be observed, or, with multiplex analysis, it may migrate within or close to the size range for another locus, further confounding analysis.

[0242] SNP analysis avoids the problems of microvariants and off-ladder alleles encountered in STR analysis. Importantly, microvariants and off-ladder alleles may provide significant problems, and may be completely missed, when using analysis methods such as oligonucleotide hybridization arrays, which utilize oligonucleotide probes specific for certain known alleles. Furthermore, off-ladder alleles and microvariants encountered with STR analysis, even when correctly typed, may lead to improper statistical analysis, since their frequencies in the population are generally unknown or poorly characterized, and therefore the statistical significance of a matching genotype may be questionable. All these advantages of SNP analysis are considerable in light of the consequences of most DNA identification cases, which may lead to life imprisonment for an individual, or re-association of remains to the family of a deceased individual.

[0243] DNA can be isolated from biological samples such as blood, bone, hair, saliva, or semen, and compared with the DNA from a reference source at particular SNP positions. Multiple SNP markers can be assayed simultaneously in order to increase the power of discrimination and the statistical significance of a matching genotype. For example, oligonucleotide arrays can be used to genotype a large number of SNPs simultaneously. The SNP used in the method of the present invention can be assayed in combination with other polymorphic genetic markers, such as other SNPs known in the art or STRs, in order to identify an individual or to associate an individual with a particular biological sample.

[0244] Furthermore, the SNP used in the method of the present invention can be genotyped for inclusion in a database of DNA genotypes, for example, a criminal DNA databank such as the FBI's Combined DNA Index System (CODIS) database. A genotype obtained from a biological sample of unknown source can then be queried against the database to find a matching genotype, with the SNP used in the method of the present invention providing nucleotide positions at which to compare the known and unknown DNA sequences for identity. Accordingly, disclosed herein is a database comprising novel SNPs or SNP alleles of the present invention (e.g., the database can comprise information indicating which alleles are possessed by individual members of a population at one or more novel SNP sites of the present invention), such as for use in forensics, biometrics, or other human identification applications. Such a database typically comprises a computer-based system in which the SNPs or SNP alleles disclosed herein are recorded on a computer readable medium (see the section of the present specification entitled "Computer-Related Embodiments").

[0245] The SNPs disclosed herein can also be assayed for use in paternity testing. The object of paternity testing is usually to determine whether a male is the father of a child. In most cases, the mother of the child is known and thus, the mother's contribution to the child's genotype can be traced. Paternity testing investigates whether the part of the child's genotype not attributable to the mother is consistent with that of the putative father. Paternity testing can be performed by analyzing sets of polymorphisms in the putative father and the child, with the SNPs of the present disclosure describing nucleotide positions at which to compare the putative father's and child's DNA sequences for identity. If the set of polymorphisms in the child attributable to the father does not match the set of polymorphisms of the putative father, it can be concluded, barring experimental error, that the putative father is not the father of the child. If the set of polymorphisms in the child attributable to the father match the set of polymorphisms of the putative father, a statistical calculation can be performed to determine the probability of coincidental match, and a conclusion drawn as to the likelihood that the putative father is the true biological father of the child.

[0246] In addition to paternity testing, SNPs are also useful for other types of kinship testing, such as for verifying familial relationships for immigration purposes, or for cases in which an individual alleges to be related to a deceased individual in order to claim an inheritance from the deceased individual, etc. For further information regarding the utility of SNPs for paternity testing and other types of kinship testing, including methods for statistical analysis, see Krawczak,

"Informativity assessment for biallelic single nucleotide polymorphisms", Electrophoresis 1999 Jun;20(8):1676-81.

[0247] The use of the SNPs for human identification further extends to various authentication systems, commonly referred to as biometric systems, which typically convert physical characteristics of humans (or other organisms) into digital data. Biometric systems include various technological devices that measure such unique anatomical or physiological characteristics as finger, thumb, or palm prints; hand geometry; vein patterning on the back of the hand; blood vessel patterning of the retina and color and texture of the iris; facial characteristics; voice patterns; signature and typing dynamics; and DNA. Such physiological measurements can be used to verify identity and, for example, restrict or allow access based on the identification. Examples of applications for biometrics include physical area security, computer and network security, aircraft passenger check-in and boarding, financial transactions, medical records access, government benefit distribution, voting, law enforcement, passports, visas and immigration, prisons, various military applications, and for restricting access to expensive or dangerous items, such as automobiles or guns (see, for example, O'Connor, *Stanford Technology Law Review* and U.S. Patent No. 6,119,096).

[0248] Groups of SNPs, particularly the SNPs disclosed herein, can be typed to uniquely identify an individual for biometric applications such as those described above. Such SNP typing can readily be accomplished using, for example, DNA chips/arrays. Preferably, a minimally invasive means for obtaining a DNA sample is utilized. For example, PCR amplification enables sufficient quantities of DNA for analysis to be obtained from buccal swabs or fingerprints, which contain DNA-containing skin cells and oils that are naturally transferred during contact.

[0249] Further information regarding techniques for using SNPs in forensic/human identification applications can be found in, for example, *Current Protocols in Human Genetics*, John Wiley & Sons, N.Y. (2002), 14.1-14.7.

VARIANT PROTEINS, ANTIBODIES,

VECTORS & HOST CELLS, & USES THEREOF

Variant Proteins Encoded by SNP-Containing Nucleic Acid Molecules

[0250] The present invention provides the use of SNP-containing nucleic acid molecules in the present invention many of which encode proteins having variant amino acid sequences as compared to the art-known (i.e., wild-type) proteins. Amino acid sequences encoded by the polymorphic nucleic acid molecules used in the method of the present invention are provided as SEQ ID NOS:845-846 in Table 1 and the Sequence Listing. These variants will generally be referred to herein as variant proteins/peptides/polypeptides, or polymorphic proteins/peptides/polypeptides used in the method of the present invention. The terms "protein", "peptide", and "polypeptide" are used herein interchangeably.

[0251] A variant protein for use in the method of the present invention may be encoded by, for example, a nonsynonymous nucleotide substitution at any one of the cSNP positions disclosed herein. In addition, variant proteins may also include proteins whose expression, structure, and/or function is altered by a SNP disclosed herein, such as a SNP that creates or destroys a stop codon, a SNP that affects splicing, and a SNP in control/regulatory elements, e.g. promoters, enhancers, or transcription factor binding domains.

[0252] As used herein, a protein or peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or chemical precursors or other chemicals. The variant proteins used in the method of the present invention can be purified to homogeneity or other lower degrees of purity. The level of purification will be based on the intended use. The key feature is that the preparation allows for the desired function of the variant protein, even if in the presence of considerable amounts of other components.

[0253] As used herein, "substantially free of cellular material" includes preparations of the variant protein having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the variant protein is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

[0254] The language "substantially free of chemical precursors or other chemicals" includes preparations of the variant protein in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the variant protein having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

[0255] An isolated variant protein may be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant host cells), or synthesized using known protein synthesis methods. For example, a nucleic acid molecule containing SNP(s) encoding the variant protein can be cloned into an expression vector, the expression vector introduced into a host cell, and the variant protein expressed in the host cell. The variant protein can then be isolated from the cells by any appropriate purification scheme using standard protein purification techniques.

Examples of these techniques are described in detail below (Sambrook and Russell, 2000, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

[0256] The present invention provides the use of isolated variant proteins that comprise, consist of or consist essentially of amino acid sequences that contain one or more variant amino acids encoded by one or more codons which contain the SNP used in the method of the present invention.

[0257] Accordingly, the present invention provides the use of variant proteins that consist of amino acid sequences that contain one or more amino acid polymorphisms (or truncations or extensions due to creation or destruction of a stop codon, respectively) encoded by the SNPs provided in Table 1 and/or Table 2 in the method of the present invention. A protein consists of an amino acid sequence when the amino acid sequence is the entire amino acid sequence of the protein.

[0258] The present invention further provides the use of variant proteins that consist essentially of amino acid sequences that contain one or more amino acid polymorphisms (or truncations or extensions due to creation or destruction of a stop codon, respectively) encoded by the SNPs provided in Table 1 and/or Table 2 in the method of the present invention. A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues in the final protein.

[0259] The present invention further provides the use of variant proteins that comprise amino acid sequences that contain one or more amino acid polymorphisms (or truncations or extensions due to creation or destruction of a stop codon, respectively) encoded by the SNPs provided in Table 1 and/or Table 2 in the method of the present invention. A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein may contain only the variant amino acid sequence or have additional amino acid residues, such as a contiguous encoded sequence that is naturally associated with it or heterologous amino acid residues. Such a protein can have a few additional amino acid residues or can comprise many more additional amino acids. A brief description of how various types of these proteins can be made and isolated is provided below.

[0260] The variant proteins used in the method of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a variant protein operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the variant protein. "Operatively linked" indicates that the coding sequences for the variant protein and the heterologous protein are ligated in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the variant protein. In another embodiment, the fusion protein is encoded by a fusion polynucleotide that is synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel et al., Current Protocols in Molecular Biology, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A variant protein-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the variant protein.

[0261] In many uses, the fusion protein does not affect the activity of the variant protein. The fusion protein can include, but is not limited to, enzymatic fusion proteins, for example, beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate their purification following recombinant expression. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence. Fusion proteins are further described in, for example, Terpe, "Overview of tag protein fusions: from molecular and biochemical fundamentals to commercial systems", Appl Microbiol Biotechnol. 2003 Jan;60(5):523-33. Epub 2002 Nov 07; Graddis et al., "Designing proteins that work using recombinant technologies", Curr Pharm Biotechnol. 2002 Dec;3(4):285-97; and Nilsson et al., "Affinity fusion strategies for detection, purification, and immobilization of recombinant proteins", Protein Expr Purif. 1997 Oct;11(1):1-16.

[0262] The present invention also relates to further obvious variants of the variant polypeptides used in the method of the present invention, such as naturally-occurring mature forms (e.g., allelic variants), non-naturally occurring recombinantly-derived variants, and orthologs and paralogs of such proteins that share sequence homology. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude those known in the prior art before the present invention.

[0263] Further variants of the variant polypeptides disclosed in Table 1 can comprise an amino acid sequence that shares at least 70-80%, 80-85%, 85-90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity with an amino acid sequence disclosed in Table 1 (or a fragment thereof) and that includes a novel amino acid residue (allele) disclosed in Table 1 (which is encoded by a novel SNP allele). Thus, disclosed herein are polypeptides that have a certain degree of sequence variation compared with the polypeptide sequences shown in Table 1, but that contain a novel amino acid residue (allele) encoded by a novel SNP allele disclosed herein. In other words, as long as a polypeptide contains a novel amino acid residue disclosed herein, other portions of the polypeptide that flank the novel amino acid residue can vary to some degree from the polypeptide sequences shown in Table 1.

[0264] Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the amino

acid sequences disclosed herein can readily be identified as having complete sequence identity to one of the variant proteins used in the method of the present invention as well as being encoded by the same genetic locus as the variant proteins disclosed herein.

[0265] Orthologs of a variant peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of a variant peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from non-human mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs can be encoded by a nucleic acid sequence that hybridizes to a variant peptide-encoding nucleic acid molecule under moderate to stringent conditions depending on the degree of relatedness of the two organisms yielding the homologous proteins.

[0266] Variant proteins include, but are not limited to, proteins containing deletions, additions and substitutions in the amino acid sequence caused by the SNP used in the method of the present invention. One class of substitutions is conserved amino acid substitutions in which a given amino acid in a polypeptide is substituted for another amino acid of like characteristics. Typical conservative substitutions are replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in, for example, Bowie et al., Science 247:1306-1310 (1990).

[0267] Variant proteins can be fully functional or can lack function in one or more activities, e.g. ability to bind another molecule, ability to catalyze a substrate, ability to mediate signaling, etc. Fully functional variants typically contain only conservative variations or variations in non-critical residues or in non-critical regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree. Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, truncations or extensions, or a substitution, insertion, inversion, or deletion of a critical residue or in a critical region.

[0268] Amino acids that are essential for function of a protein can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham et al., Science 244:1081-1085 (1989)), particularly using the amino acid sequence and polymorphism information provided in Table 1. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as enzyme activity or in assays such as an *in vitro* proliferative activity. Sites that are critical for binding partner/substrate binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith et al., J. Mol. Biol. 224:899-904 (1992); de Vos et al. Science 255:306-312 (1992)).

[0269] Polypeptides can contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Accordingly, the variant proteins for use in the method of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (e.g., polyethylene glycol), or in which additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence for purification of the mature polypeptide or a pro-protein sequence.

[0270] Known protein modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

[0271] Such protein modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as Proteins - Structure and Molecular Properties, 2nd Ed., -T.E. Creighton, W. H. Freeman and Company, New York (1993); Wold, F., Posttranslational Covalent Modification of Proteins, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter et al., Meth. Enzymol. 182: 626-646 (1990); and Rattan et al., Ann. N.Y. Acad. Sci. 663:48-62 (1992).

[0272] Further, disclosed herein are fragments of the variant proteins in which the fragments contain one or more amino acid sequence variations (e.g., substitutions, or truncations or extensions due to creation or destruction of a stop codon) encoded by one or more SNPs disclosed herein. The fragments to which the disclosure pertains, however, are

not to be construed as encompassing fragments that have been disclosed in the prior art before the present disclosure.

[0273] As used herein, a fragment may comprise at least about 4, 8, 10, 12, 14, 16, 18, 20, 25, 30, 50, 100 (or any other number in-between) or more contiguous amino acid residues from a variant protein, wherein at least one amino acid residue is affected by a SNP, e.g., a variant amino acid residue encoded by a nonsynonymous nucleotide substitution at a cSNP position disclosed herein. The variant amino acid encoded by a cSNP may occupy any residue position along the sequence of the fragment. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the variant protein or the ability to perform a function, e.g., act as an immunogen. Particularly important fragments are biologically active fragments. Such fragments will typically comprise a domain or motif of a variant protein disclosed herein, e.g., active site, transmembrane domain, or ligand/substrate binding domain. Other fragments include, but are not limited to, domain or motif-containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known to those of skill in the art (e.g., PROSITE analysis) (Current Protocols in Protein Science, John Wiley & Sons, N.Y. (2002)).

Uses of Variant Proteins

[0274] The variant proteins can be used in a variety of ways, including but not limited to, in assays to determine the biological activity of a variant protein, such as in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another type of immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the variant protein (or its binding partner) in biological fluids; as a marker for cells or tissues in which it is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); as a target for screening for a therapeutic agent; and as a direct therapeutic agent to be administered into a human subject. Any of the variant proteins disclosed herein may be developed into reagent grade or kit format for commercialization as research products. Methods for performing the uses listed above are well known to those skilled in the art (see, e.g., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Sambrook and Russell, 2000, and Methods in Enzymology: Guide to Molecular Cloning Techniques, Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987).

[0275] In a specific embodiment of the invention, the methods of the present invention include detection of one or more variant proteins disclosed herein. Variant proteins are disclosed in Table 1 and in the Sequence Listing as SEQ ID NOS:845-846. Detection of such proteins can be accomplished using, for example, antibodies, small molecule compounds, aptamers, ligands/substrates, other proteins or protein fragments, or other protein-binding agents. Preferably, protein detection agents are specific for a variant protein of the present invention and can therefore discriminate between a variant protein used in the method of the present invention and the wild-type protein or another variant form. This can generally be accomplished by, for example, selecting or designing detection agents that bind to the region of a protein that differs between the variant and wild-type protein, such as a region of a protein that contains one or more amino acid substitutions that is/are encoded by a non-synonymous cSNP, or a region of a protein that follows a nonsense mutation-type SNP that creates a stop codon thereby leading to a shorter polypeptide, or a region of a protein that follows a read-through mutation-type SNP that destroys a stop codon thereby leading to a longer polypeptide in which a portion of the polypeptide is present in one version of the polypeptide but not the other.

[0276] In another specific aspect of the invention, the variant proteins used in the method of the present invention are used as targets for diagnosing myocardial infarction or for determining predisposition to myocardial infarction in a human. Accordingly, the invention provides methods for detecting the presence of, or levels of, one or more variant proteins in a cell, tissue, or organism. Such methods typically involve contacting a test sample with an agent (e.g., an antibody, small molecule compound, or peptide) capable of interacting with the variant protein such that specific binding of the agent to the variant protein can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an array, for example, an antibody or aptamer array (arrays for protein detection may also be referred to as "protein chips"). The variant protein of interest can be isolated from a test sample and assayed for the presence of a variant amino acid sequence encoded by the SNP disclosed by the present invention. The SNPs may cause changes to the protein and the corresponding protein function/activity, such as through non-synonymous substitutions in protein coding regions that can lead to amino acid substitutions, deletions, insertions, and/or rearrangements; formation or destruction of stop codons; or alteration of control elements such as promoters. SNPs may also cause inappropriate post-translational modifications.

[0277] One preferred agent for detecting a variant protein in a sample is an antibody capable of selectively binding to a variant form of the protein (antibodies are described in greater detail in the next section). Such samples include, for example, tissues, cells, and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

[0278] *In vitro* methods for detection of the variant proteins associated with myocardial infarction that are disclosed herein and fragments thereof include, but are not limited to, enzyme linked immunosorbent assays (ELISAs), radioimmunoassays (RIA), Western blots, immunoprecipitations, immunofluorescence, and protein arrays/chips (e.g., arrays

of antibodies or aptamers). For further information regarding immunoassays and related protein detection methods, see Current Protocols in Immunology, John Wiley & Sons, N.Y., and Hage, "Immunoassays", Anal Chem. 1999 Jun 15;71(12):294R-304R.

[0279] Additional analytic methods of detecting amino acid variants include, but are not limited to, altered electrophoretic mobility, altered tryptic peptide digest, altered protein activity in cell-based or cell-free assay, alteration in ligand or antibody-binding pattern, altered isoelectric point, and direct amino acid sequencing.

[0280] Alternatively, variant proteins can be detected *in vivo* in a subject by introducing into the subject a labeled antibody (or other type of detection reagent) specific for a variant protein. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

[0281] Other uses of the variant peptides are based on the class or action of the protein: For example, proteins isolated from humans and their mammalian orthologs serve as targets for identifying agents (e.g., small molecule drugs or antibodies) for use in therapeutic applications, particularly for modulating a biological or pathological response in a cell or tissue that expresses the protein.

[0282] Pharmaceutical agents can be developed that modulate protein activity.

[0283] As an alternative to modulating gene expression, therapeutic compounds can be developed that modulate protein function. For example, many SNPs disclosed herein affect the amino acid sequence of the encoded protein (e.g., non-synonymous cSNPs and nonsense mutation-type SNPs). Such alterations in the encoded amino acid sequence may affect protein function, particularly if such amino acid sequence variations occur in functional protein domains, such as catalytic domains, ATP-binding domains, or ligand/substrate binding domains. It is well established in the art that variant proteins having amino acid sequence variations in functional domains can cause or influence pathological conditions. In such instances, compounds (e.g., small molecule drugs or antibodies) can be developed that target the variant protein and modulate (e.g., up- or down-regulate) protein function/activity.

[0284] The therapeutic methods disclosed herein further include methods that target one or more variant proteins disclosed herein. Variant proteins can be targeted using, for example, small molecule compounds, antibodies, aptamers, ligands/substrates, other proteins, or other protein-binding agents. Additionally, the skilled artisan will recognize that the novel protein variants (and polymorphic nucleic acid molecules) disclosed in Table 1 may themselves be directly used as therapeutic agents by acting as competitive inhibitors of corresponding art-known proteins (or nucleic acid molecules such as mRNA molecules).

[0285] The variant proteins disclosed herein are particularly useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can utilize cells that naturally express the protein, a biopsy specimen, or cell cultures. In one embodiment, cell-based assays involve recombinant host cells expressing the variant protein. Cell-free assays can be used to detect the ability of a compound to directly bind to a variant protein or to the corresponding SNP-containing nucleic acid fragment that encodes the variant protein.

[0286] A variant protein disclosed herein, as well as appropriate fragments thereof, can be used in high-throughput screening assays to test candidate compounds for the ability to bind and/or modulate the activity of the variant protein. These candidate compounds can be further screened against a protein having normal function (e.g., a wild-type/non-variant protein) to further determine the effect of the compound on the protein activity. Furthermore, these compounds can be tested in animal or invertebrate systems to determine *in vivo* activity/effectiveness. Compounds can be identified that activate (agonists) or inactivate (antagonists) the variant protein, and different compounds can be identified that cause various degrees of activation or inactivation of the variant protein.

[0287] Further, the variant proteins can be used to screen a compound for the ability to stimulate or inhibit interaction between the variant protein and a target molecule that normally interacts with the protein. The target can be a ligand, a substrate or a binding partner that the protein normally interacts with (for example, epinephrine or norepinephrine). Such assays typically include the steps of combining the variant protein with a candidate compound under conditions that allow the variant protein, or fragment thereof, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the variant protein and the target, such as any of the associated effects of signal transduction.

[0288] Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam et al., Nature 354:82-84 (1991); Houghten et al., Nature 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang et al., Cell 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

[0289] One candidate compound is a soluble fragment of the variant protein that competes for ligand binding. Other candidate compounds include mutant proteins or appropriate fragments containing mutations that affect variant protein function and thus compete for ligand. Accordingly, a fragment that competes for ligand, for example with a higher affinity,

or a fragment that binds ligand but does not allow release, is described herein.

[0290] Disclosed herein are other end point assays to identify compounds that modulate (stimulate or inhibit) variant protein activity. The assays typically involve an assay of events in the signal transduction pathway that indicate protein activity. Thus, the expression of genes that are up or down-regulated in response to the variant protein dependent signal cascade can be assayed. In one embodiment, the regulatory region of such genes can be operably linked to a marker that is easily detectable, such as luciferase. Alternatively, phosphorylation of the variant protein, or a variant protein target, could also be measured. Any of the biological or biochemical functions mediated by the variant protein can be used as an endpoint assay. These include all of the biochemical or biological events described herein, in the references cited herein, for these endpoint assay targets, and other functions known to those of ordinary skill in the art.

[0291] Binding and/or activating compounds can also be screened by using chimeric variant proteins in which an amino terminal extracellular domain or parts thereof, an entire transmembrane domain or subregions, and/or the carboxyl terminal intracellular domain or parts thereof, can be replaced by heterologous domains or subregions. For example, a substrate-binding region can be used that interacts with a different substrate than that which is normally recognized by a variant protein. Accordingly, a different set of signal transduction components is available as an end-point assay for activation. This allows for assays to be performed in other than the specific host cell from which the variant protein is derived.

[0292] The variant proteins are also useful in competition binding assays in methods designed to discover compounds that interact with the variant protein. Thus, a compound can be exposed to a variant protein under conditions that allow the compound to bind or to otherwise interact with the variant protein. A binding partner, such as ligand, that normally interacts with the variant protein is also added to the mixture. If the test compound interacts with the variant protein or its binding partner, it decreases the amount of complex formed or activity from the variant protein. This type of assay is particularly useful in screening for compounds that interact with specific regions of the variant protein (Hodgson, *Bio/technology*, 1992, Sept 10(9), 973-80).

[0293] To perform cell-free drug screening assays, it is sometimes desirable to immobilize either the variant protein or a fragment thereof, or its target molecule, to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Any method for immobilizing proteins on matrices can be used in drug screening assays. In one embodiment, a fusion protein containing an added domain allows the protein to be bound to a matrix. For example, glutathione-S-transferase/¹²⁵I fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., ³⁵S-labeled) and a candidate compound, such as a drug candidate, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads can be washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of bound material found in the bead fraction quantitated from the gel using standard electrophoretic techniques.

[0294] Either the variant protein or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Alternatively, antibodies reactive with the variant protein but which do not interfere with binding of the variant protein to its target molecule can be derivatized to the wells of the plate, and the variant protein trapped in the wells by antibody conjugation. Preparations of the target molecule and a candidate compound are incubated in the variant protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the protein target molecule, or which are reactive with variant protein and compete with the target molecule, and enzyme-linked assays that rely on detecting an enzymatic activity associated with the target molecule.

[0295] Modulators of variant protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the protein pathway, such as myocardial infarction. These methods of treatment typically include the steps of administering the modulators of protein activity in a pharmaceutical composition to a subject in need of such treatment.

[0296] The variant proteins, or fragments thereof, disclosed herein can themselves be directly used to treat a disorder characterized by an absence of, inappropriate, or unwanted expression or activity of the variant protein. Accordingly, methods for treatment include the use of a variant protein disclosed herein or fragments thereof.

[0297] In yet another aspect disclosed herein, variant proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J. Biol. Chem.* 268:12046-12054; Bartel et al. (1993) *Biotechniques* 14:920-924; Iwabuchi et al. (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300) to identify other proteins that bind to or interact with the variant protein and are involved in variant protein activity. Such variant protein-binding proteins are also likely to be involved in the propagation of signals by the variant proteins or variant protein targets as, for example, elements of a protein-mediated signaling pathway. Alternatively, such variant protein-binding proteins are inhibitors of the variant protein.

[0298] The two-hybrid system is based on the modular nature of most transcription factors, which typically consist of separable DNA-binding and activation domains. Briefly, the assay typically utilizes two different DNA constructs. In one construct, the gene that codes for a variant protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a variant protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene that encodes the protein that interacts with the variant protein.

Antibodies Directed to Variant Proteins

[0299] Disclosed herein are also antibodies that selectively bind to the variant proteins disclosed herein and fragments thereof. Such antibodies may be used to quantitatively or qualitatively detect the variant proteins disclosed herein. As used herein, an antibody selectively binds a target variant protein when it binds the variant protein and does not significantly bind to non-variant proteins, i.e., the antibody does not significantly bind to normal, wild-type, or art-known proteins that do not contain a variant amino acid sequence due to one or more SNPs disclosed herein (variant amino acid sequences may be due to, for example, nonsynonymous cSNPs, nonsense SNPs that create a stop codon, thereby causing a truncation of a polypeptide or SNPs that cause read-through mutations resulting in an extension of a polypeptide).

[0300] As used herein, an antibody is defined in terms consistent with that recognized in the art: they are multi-subunit proteins produced by an organism in response to an antigen challenge. The antibodies disclosed herein include both monoclonal antibodies and polyclonal antibodies, as well as antigen-reactive proteolytic fragments of such antibodies, such as Fab, F(ab)₂, and Fv fragments. In addition, an antibody further includes any of a variety of engineered antigen-binding molecules such as a chimeric antibody (U.S. Patent Nos. 4,816,567 and 4,816,397; Morrison et al., Proc. Natl. Acad. Sci. USA, 81:6851, 1984; Neuberger et al., Nature 312:604, 1984), a humanized antibody (U.S. Patent Nos. 5,693,762; 5,585,089; and 5,565,332), a single-chain Fv (U.S. Patent No. 4,946,778; Ward et al., Nature 334:544, 1989), a bispecific antibody with two binding specificities (Segal et al., J. Immunol. Methods 248:1, 2001; Carter, J. Immunol. Methods 248:7, 2001), a diabody, a triabody, and a tetrabody (Todorovska et al., J. Immunol. Methods, 248:47, 2001), as well as a Fab conjugate (dimer or trimer), and a minibody.

[0301] Many methods are known in the art for generating and/or identifying antibodies to a given target antigen (Harlow, Antibodies, Cold Spring Harbor Press, (1989)). In general, an isolated peptide (e.g., a variant protein disclosed herein) is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit, hamster or mouse. Either a full-length protein, an antigenic peptide fragment (e.g., a peptide fragment containing a region that varies between a variant protein and a corresponding wild-type protein), or a fusion protein can be used. A protein used as an immunogen may be naturally-occurring, synthetic or recombinantly produced, and may be administered in combination with an adjuvant, including but not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substance such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and the like.

[0302] Monoclonal antibodies can be produced by hybridoma technology (Kohler and Milstein, Nature, 256:495, 1975), which immortalizes cells secreting a specific monoclonal antibody. The immortalized cell lines can be created *in vitro* by fusing two different cell types, typically lymphocytes, and tumor cells. The hybridoma cells may be cultivated *in vitro* or *in vivo*. Additionally, fully human antibodies can be generated by transgenic animals (He et al., J. Immunol., 169:595, 2002). Fd phage and Fd phagemid technologies may be used to generate and select recombinant antibodies *in vitro* (Hoogenboom and Chames, Immunol. Today 21:371, 2000; Liu et al., J. Mol. Biol. 315:1063, 2002). The complementarity-determining regions of an antibody can be identified, and synthetic peptides corresponding to such regions may be used to mediate antigen binding (U.S. Patent No. 5,637,677).

[0303] Antibodies are preferably prepared against regions or discrete fragments of a variant protein containing a variant amino acid sequence as compared to the corresponding wild-type protein (e.g., a region of a variant protein that includes an amino acid encoded by a nonsynonymous cSNP, a region affected by truncation caused by a nonsense SNP that creates a stop codon, or a region resulting from the destruction of a stop codon due to read-through mutation caused by a SNP). Furthermore, preferred regions will include those involved in function/activity and/or protein/binding partner interaction. Such fragments can be selected on a physical property, such as fragments corresponding to regions that are located on the surface of the protein, e.g., hydrophilic regions, or can be selected based on sequence uniqueness, or based on the position of the variant amino acid residue(s) encoded by the SNPs disclosed herein. An antigenic fragment will typically comprise at least about 8-10 contiguous amino acid residues in which at least one of the amino

acid residues is an amino acid affected by a SNP disclosed herein. The antigenic peptide can comprise, however, at least 12, 14, 16, 20, 25, 50, 100 (or any other number in-between) or more amino acid residues, provided that at least one amino acid is affected by a SNP disclosed herein.

[0304] Detection of an antibody can be facilitated by coupling (i.e., physically linking) the antibody or an antigen-reactive fragment thereof to a detectable substance. Detectable substances include, but are not limited to, various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

[0305] Antibodies, particularly the use of antibodies as therapeutic agents, are reviewed in: Morgan, "Antibody therapy for Alzheimer's disease", *Expert Rev Vaccines*. 2003 Feb;2(1):53-9; Ross et al., "Anticancer antibodies", *Am J Clin Pathol*. 2003 Apr;119(4):472-85; Goldenberg, "Advancing role of radiolabeled antibodies in the therapy of cancer", *Cancer Immunol Immunother*. 2003 May;52(5):281-96. Epub 2003 Mar 11; Ross et al., "Antibody-based therapeutics in oncology", *Expert Rev Anticancer Ther*. 2003 Feb;3(1):107-21; Cao et al., "Bispecific antibody conjugates in therapeutics", *Adv Drug Deliv Rev*. 2003 Feb 10;55(2):171-97; von Mehren et al., "Monoclonal antibody therapy for cancer", *Annu Rev Med*. 2003;54:343-69. Epub 2001 Dec 03; Hudson et al., "Engineered antibodies", *Nat Med*. 2003 Jan;9(1):129-34; Brekke et al., "Therapeutic antibodies for human diseases at the dawn of the twenty-first century", *Nat Rev Drug Discov*. 2003 Jan;2(1):52-62 (Erratum in: *Nat Rev Drug Discov*. 2003 Mar;2(3):240); Houdebine, "Antibody manufacture in transgenic animals and comparisons with other systems", *Curr Opin Biotechnol*. 2002 Dec;13(6):625-9; Andreaskos et al., "Monoclonal antibodies in immune and inflammatory diseases", *Curr Opin Biotechnol*. 2002 Dec;13(6):615-20; Kellermann et al., "Antibody discovery: the use of transgenic mice to generate human monoclonal antibodies for therapeutics", *Curr Opin Biotechnol*. 2002 Dec;13(6):593-7; Pini et al., "Phage display and colony filter screening for high-throughput selection of antibody libraries", *Comb Chem High Throughput Screen*. 2002 Nov;5(7):503-10; Batra et al., "Pharmacokinetics and biodistribution of genetically engineered antibodies", *Curr Opin Biotechnol*. 2002 Dec;13(6):603-8; and Tangri et al., "Rationally engineered proteins or antibodies with absent or reduced immunogenicity", *Curr Med Chem*. 2002 Dec;9(24):2191-9.

Uses of Antibodies

[0306] Antibodies can be used to isolate the variant proteins from a natural cell source or from recombinant host cells by standard techniques, such as affinity chromatography or immunoprecipitation. In addition, antibodies are useful for detecting the presence of a variant protein in cells or tissues to determine the pattern of expression of the variant protein among various tissues in an organism and over the course of normal development or disease progression. Further, antibodies can be used to detect variant protein *in situ*, *in vitro*, in a bodily fluid, or in a cell lysate or supernatant in order to evaluate the amount and pattern of expression. Also, antibodies can be used to assess abnormal tissue distribution, abnormal expression during development, or expression in an abnormal condition, such as myocardial infarction. Additionally, antibody detection of circulating fragments of the full-length variant protein can be used to identify turnover.

[0307] Antibodies to the variant proteins disclosed herein are also useful in pharmacogenomic analysis. Thus, antibodies against variant proteins encoded by alternative SNP alleles can be used to identify individuals that require modified treatment modalities.

[0308] Further, antibodies can be used to assess expression of the variant protein in disease states such as in active stages of the disease or in an individual with a predisposition to a disease related to the protein's function, particularly myocardial infarction. Antibodies specific for a variant protein encoded by a SNP-containing nucleic acid molecule for use in the method of the present invention can be used to assay for the presence of the variant protein, such as to screen for predisposition to myocardial infarction as indicated by the presence of the variant protein.

[0309] Antibodies are also useful as diagnostic tools for evaluating the variant proteins in conjunction with analysis by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays well known in the art.

[0310] Antibodies are also useful for tissue typing. Thus, where a specific variant protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

[0311] Antibodies can also be used to assess aberrant subcellular localization of a variant protein in cells in various tissues. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting the expression level or the presence of variant protein or aberrant tissue distribution or developmental expression of a variant protein, antibodies directed against the variant protein or relevant fragments can be used to monitor therapeutic efficacy.

[0312] The antibodies are also useful for inhibiting variant protein function, for example, by blocking the binding of a

variant protein to a binding partner. These uses can also be applied in a therapeutic context in which treatment involves inhibiting a variant protein's function. An antibody can be used, for example, to block or competitively inhibit binding, thus modulating (agonizing or antagonizing) the activity of a variant protein. Antibodies can be prepared against specific variant protein fragments containing sites required for function or against an intact variant protein that is associated with a cell or cell membrane. For *in vivo* administration, an antibody may be linked with an additional therapeutic payload such as a radionuclide, an enzyme, an immunogenic epitope, or a cytotoxic agent. Suitable cytotoxic agents include, but are not limited to, bacterial toxin such as diphtheria, and plant toxin such as ricin. The *in vivo* half-life of an antibody or a fragment thereof may be lengthened by pegylation through conjugation to polyethylene glycol (Leong et al., Cytokine 16:106, 2001).

[0313] The disclosure also relates to kits for using antibodies, such as kits for detecting the presence of a variant protein in a test sample. An exemplary kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting variant proteins in a biological sample; means for determining the amount, or presence/absence of variant protein in the sample; means for comparing the amount of variant protein in the sample with a standard; and instructions for use.

Vectors and Host Cells

[0314] Disclosed herein are also vectors containing the SNP-containing nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport a SNP-containing nucleic acid molecule. When the vector is a nucleic acid molecule, the SNP-containing nucleic acid molecule can be covalently linked to the vector nucleic acid. Such vectors include, but are not limited to, a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, or MAC.

[0315] A vector can be maintained in a host cell as an extrachromosomal element where it replicates and produces additional copies of the SNP-containing nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the SNP-containing nucleic acid molecules when the host cell replicates.

[0316] Disclosed herein are vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the SNP-containing nucleic acid molecules. The vectors can function in prokaryotic or eukaryotic cells or in both (shuttle vectors).

[0317] Expression vectors typically contain cis-acting regulatory regions that are operably linked in the vector to the SNP-containing nucleic acid molecules such that transcription of the SNP-containing nucleic acid molecules is allowed in a host cell. The SNP-containing nucleic acid molecules can also be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the SNP-containing nucleic acid molecules from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

[0318] The regulatory sequences to which the SNP-containing nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

[0319] In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

[0320] In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region, a ribosome-binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. A person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors (see, e.g., Sambrook and Russell, 2000, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

[0321] A variety of expression vectors can be used to express a SNP-containing nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors can also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g., cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook and Russell, 2000, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

[0322] The regulatory sequence in a vector may provide constitutive expression in one or more host cells (e.g., tissue

specific expression) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor, e.g., a hormone or other ligand. A variety of vectors that provide constitutive or inducible expression of a nucleic acid sequence in prokaryotic and eukaryotic host cells are well known to those of ordinary skill in the art.

[0323] A SNP-containing nucleic acid molecule can be inserted into the vector by methodology well-known in the art. Generally, the SNP-containing nucleic acid molecule that will ultimately be expressed is joined to an expression vector by cleaving the SNP-containing nucleic acid molecule and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

[0324] The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial host cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic host cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

[0325] As described herein, it may be desirable to express the variant peptide as a fusion protein. Accordingly, disclosed herein are fusion vectors that allow for the production of the variant peptides. Fusion vectors can, for example, increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting, for example, as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired variant peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes suitable for such use include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion expression vectors include pGEX (Smith et al., Gene 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann et al., Gene 69:301-315 (1988)) and pET 11d (Studier et al., Gene Expression Technology: Methods in Enzymology 185:60-89 (1990)).

[0326] Recombinant protein expression can be maximized in a bacterial host by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the SNP-containing nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example, *E. coli* (Wada et al., Nucleic Acids Res. 20:2111-2118 (1992)).

[0327] The SNP-containing nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast (e.g., *S. cerevisiae*) include pYepSec1 (Baldari, et al., EMBO J. 6:229-234 (1987)), pMFa (Kurjan et al., Cell 30:933-943(1982)), pJRY88 (Schultz et al., Gene 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

[0328] The SNP-containing nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith et al., Mol. Cell Biol. 3:2156-2165 (1983)) and the pVL series (Lucklow et al., Virology 170:31-39 (1989)).

[0329] In certain aspects of the disclosure, the SNP-containing nucleic acid, molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. Nature 329:840(1987)) and pMT2PC (Kaufman et al., EMBO J. 6:187-195 (1987)).

[0330] Disclosure herein are also vectors in which the SNP-containing nucleic acid molecules described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to the SNP-containing nucleic acid sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

[0331] The disclosure also relates to recombinant host cells containing the vectors described herein. Host cells therefore include, for example, prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

[0332] The recombinant host cells can be prepared by introducing the vector constructs described herein into the cells by techniques readily available to persons of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those described in Sambrook and Russell, 2000, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

[0333] Host cells can contain more than one vector. Thus, different SNP-containing nucleotide sequences can be introduced in different vectors into the same cell. Similarly, the SNP-containing nucleic acid molecules can be introduced either alone or with other nucleic acid molecules that are not related to the SNP-containing nucleic acid molecules, such

as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced, or joined to the nucleic acid molecule vector.

[0334] In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication can occur in host cells that provide functions that complement the defects.

[0335] Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be inserted in the same vector that contains the SNP-containing nucleic acid molecules described herein or may be in a separate vector. Markers include, for example, tetracycline or ampicillin-resistance genes for prokaryotic host cells, and dihydrofolate reductase or neomycin resistance genes for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait can be effective.

[0336] While the mature variant proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these variant proteins using RNA derived from the DNA constructs described herein.

[0337] Where secretion of the variant protein is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as G-protein-coupled receptors (GPCRs), appropriate secretion signals can be incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

[0338] Where the variant protein is not secreted into the medium, the protein can be isolated from the host cell by standard disruption procedures, including freeze/thaw, sonication, mechanical disruption, use of lysing agents, and the like. The variant protein can then be recovered and purified by well-known purification methods including, for example, ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

[0339] It is also understood that, depending upon the host cell in which recombinant production of the variant proteins described herein occurs, they can have various glycosylation patterns, or may be non-glycosylated, as when produced in bacteria. In addition, the variant proteins may include an initial modified methionine in some cases as a result of a host-mediated process.

[0340] For further information regarding vectors and host cells, see Current Protocols in Molecular Biology, John Wiley & Sons, N.Y.

Uses of Vectors and Host Cells, and Transgenic Animals

[0341] Recombinant host cells that express the variant proteins described herein have a variety of uses. For example, the cells are useful for producing a variant protein that can be further purified into a preparation of desired amounts of the variant protein or fragments thereof. Thus, host cells containing expression vectors are useful for variant protein production.

[0342] Host cells are also useful for conducting cell-based assays involving the variant protein or variant protein fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a variant protein is useful for assaying compounds that stimulate or inhibit variant protein function. Such an ability of a compound to modulate variant protein function may not be apparent from assays of the compound on the native/wild-type protein, or from cell-free assays of the compound. Recombinant host cells are also useful for assaying functional alterations in the variant proteins as compared with a known function.

[0343] Genetically-engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a non-human mammal, for example, a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA containing a SNP disclosed herein which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more of its cell types or tissues. Such animals are useful for studying the function of a variant protein *in vivo*, and identifying and evaluating modulators of variant protein activity. Other examples of transgenic animals include, but are not limited to, non-human primates, sheep, dogs, cows, goats, chickens, and amphibians. Transgenic non-human mammals such as cows and goats can be used to produce variant proteins which can be secreted in the animal's milk and then recovered.

[0344] A transgenic animal can be produced by introducing a SNP-containing nucleic acid molecule into the male pronuclei of a fertilized oocyte, e.g., by microinjection or retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any nucleic acid molecules that contain one or more SNPs disclosed herein can potentially be introduced as a transgene into the genome of a non-human animal.

[0345] Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the variant protein in particular cells or tissues.

[0346] Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described in, for example, U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Patent No. 4,873,191 by Wagner et al., and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes a non-human animal in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

[0347] In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the cre/loxP recombinase system of bacteriophage P1 (Lakso et al. PNAS 89:6232-6236 (1992)). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman et al. Science 251:1351-1355 (1991)). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are generally needed. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected variant protein and the other containing a transgene encoding a recombinase.

[0348] Clones of the non-human transgenic animals described herein can also be produced. according to the methods described in, for example, Wilmut, I. et al. Nature 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell (e.g., a somatic cell) from the transgenic animal can be isolated and induced to exit the growth cycle and enter Go phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell (e.g., a somatic cell) is isolated.

[0349] Transgenic animals containing recombinant cells that express the variant proteins described herein are useful for conducting the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could influence ligand or substrate binding, variant protein activation, signal transduction, or other processes or interactions, may not be evident from *in vitro* cell-free or cell-based assays. Thus, non-human transgenic animals may be used to assay *in vivo* variant protein function as well as the activities of a therapeutic agent or compound that modulates variant protein function/activity or expression. Such animals are also suitable for assessing the effects of null mutations (i.e., mutations that substantially or completely eliminate one or more variant protein functions).

[0350] For further information regarding transgenic animals, see Houdebine, "Antibody manufacture in transgenic animals and comparisons with other systems", Curr Opin Biotechnol. 2002 Dec;13(6):625-9; Petters et al., "Transgenic animals as models for human disease", Transgenic Res. 2000;9(4-5):347-51; discussion 345-6; Wolf et al., "Use of transgenic animals in understanding molecular mechanisms of toxicity", J Pharm Pharmacol. 1998 Jun;50(6):567-74; Echelard, "Recombinant protein production in transgenic animals", Curr Opin Biotechnol. 1996 Oct;7(5):536-40; Houdebine, "Transgenic animal bioreactors", Transgenic Res. 2000;9(4-5):305-20; Pirity et al., "Embryonic stem cells, creating transgenic animals", Methods Cell Biol. 1998;57:279-93; and Robl et al., "Artificial chromosome vectors and expression of complex proteins in transgenic animals", Theriogenology. 2003 Jan 1;59(1):107-13.

COMPUTER-RELATED EMBODIMENTS

[0351] The SNP used in the method of the present invention may be "provided" in a variety of mediums to facilitate use thereof. As used in this section, "provided" refers to a manufacture, other than an isolated nucleic acid molecule, that contains SNP information disclosed herein. Such a manufacture provides the SNP information in a form that allows a skilled artisan to examine the manufacture using means not directly applicable to examining the SNPs or a subset thereof as they exist in nature or in purified form. The SNP information that may be provided in such a form includes any of the SNP information used in the method of the present invention such as, for example, polymorphic nucleic acid and/or amino acid sequence information such as SEQ ID NOS: 17-18, SEQ ID NOS:845-846, SEQ ID NOS:17560, SEQ ID NOS:2952-2984, and SEQ ID NOS: 18962-19247; information about observed SNP alleles, alternative codons, populations, allele frequencies, SNP types, and/or affected proteins; or any other information provided by the present invention in Tables 1-2 and/or the Sequence Listing.

[0352] In one application of this embodiment, the SNP used in the method of the present invention can be recorded on a computer readable medium. As used herein, "computer readable medium" refers to any medium that can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy

discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable media can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence used in the method of the present invention. One such medium is provided with the present application, namely, the present application contains computer readable medium (CD-R) that has nucleic acid sequences (and encoded protein sequences) containing SNPs provided/recorded thereon in ASCII text format in a Sequence Listing along with accompanying Tables that contain detailed SNP and sequence information (transcript sequences are provided as SEQ ID NOS:1-828, protein sequences are provided as SEQ ID NOS:829-1656, genomic sequences are provided as SEQ ID NOS:17,553-18,016, transcript-based context sequences are provided as SEQ ID NOS:1657-17,552, and genomic-based context sequences are provided as SEQ ID NOS:18,017-73,085).

[0353] As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate manufactures comprising the SNP information used in the method of the present invention.

[0354] A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide or amino acid sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide/amino acid sequence information used in the method of the present invention on computer readable medium. For example, the sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, represented in the form of an ASCII file, or stored in a database application, such as OB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (e.g., text file or database) in order to obtain computer readable medium having recorded thereon the SNP information used in the method of the present invention.

[0355] By providing the SNPs disclosed herein in computer readable form, a skilled artisan can routinely access the SNP information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium. Examples of publicly available computer software include BLAST (Altschul et al, J. Mol. Biol. 215:403-410 (1990)) and BLAZE (Brutlag et al, Comp. Chem. 17:203-207 (1993)) search algorithms.

[0356] Further, disclosed herein are systems, particularly computer-based systems, which contain the SNP information described herein. Such systems may be designed to store and/or analyze information on, for example, a large number of SNP positions, or information on SNP genotypes from a large number of individuals. The SNP information used in the method of the present invention represents a valuable information source. The SNP information used in the method of the present invention stored/analyzed in a computer-based system may be used for such computer-intensive applications as determining or analyzing SNP allele frequencies in a population, mapping disease genes, genotype-phenotype association studies, grouping SNPs into haplotypes, correlating SNP haplotypes with response to particular drugs, or for various other bioinformatic, pharmacogenomic, drug development, or human identification/forensic applications.

[0357] As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the SNP information used in the method of the present invention. The minimum hardware means of the computer-based systems of the present invention typically comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are suitable for use in the present invention. Such a system can be changed into a system of the present invention by utilizing the SNP information provided on the CD-R, or a subset thereof, without any experimentation.

[0358] As stated above, the computer-based systems comprise a data storage means having stored therein SNPs and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store SNP information, or a memory access means which can access manufactures having recorded thereon the SNP information disclosed herein.

[0359] As used herein, "search means" refers to one or more programs or algorithms that are implemented on the computer-based system to identify or analyze SNPs in a target sequence based on the SNP information stored within the data storage means. Search means can be used to determine which nucleotide is present at a particular SNP position in the target sequence. As used herein, a "target sequence" can be any DNA sequence containing the SNP position(s) to be searched or queried.

[0360] As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences containing a SNP position in which the sequence(s) is chosen based on a three-dimensional configuration that is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzymatic active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures, and inducible expression elements (protein

binding sequences).

[0361] A variety of structural formats for the input and output means can be used to input and output the information in the computer-based systems disclosed herein. An exemplary format for an output means is a display that depicts the presence or absence of specified nucleotides (alleles) at particular SNP positions of interest. Such presentation can provide a rapid, binary scoring system for many SNPs simultaneously.

[0362] One exemplary embodiment of a computer-based system comprising SNP information disclosed herein is provided in Figure 1. Figure 1 provides a block diagram of a computer system 102 that can be used to implement the present disclosure. The computer system 102 includes a processor 106 connected to a bus 104. Also connected to the bus 104 are a main memory 108 (preferably implemented as random access memory, RAM) and a variety of secondary storage devices 110, such as a hard drive 112 and a removable medium storage device 114. The removable medium storage device 114 may represent, for example, a floppy disk drive, a CD-ROM drive, a magnetic tape drive, etc. A removable storage medium 116 (such as a floppy disk, a compact disk, a magnetic tape, etc.) containing control logic and/or data recorded therein may be inserted into the removable medium storage device 114. The computer system 102 includes appropriate software for reading the control logic and/or the data from the removable storage medium 116 once inserted in the removable medium storage device 114.

[0363] The SNP information may be stored in a well-known manner in the main memory 108, any of the secondary storage devices 110, and/or a removable storage medium 116. Software for accessing and processing the SNP information (such as SNP scoring tools, search tools, comparing tools, etc.) preferably resides in main memory 108 during execution.

EXAMPLES: STATISTICAL ANALYSIS OF SNP ASSOCIATION WITH MYOCARDIAL INFARCTION AND RECURRENT MYOCARDIAL INFARCTION

Myocardial Infarction studies (see Table 6)

[0364] A case-control genetic study to determine the association of SNPs in the human genome with MI was carried out using genomic DNA extracted from 3 independently collected case-control sample sets.

[0365] Study S0012 had 1400 samples, in which patients (cases) had self-reported history of MI, and controls had no history of MI or of acute angina lasting more than 1 hour. Study S0028 had 1500 samples, in which patients (cases) had clinical evidence of history of MI, and controls had no history of MI. Study V0001 had 1288 samples, in which patients (cases) had clinical evidence of history of MI, and controls had no history of MI. All individuals who were included in each study had signed a written informed consent form. The study protocol was IRB approved.

[0366] DNA was extracted from blood samples using conventional DNA extraction methods such as the QIA-amp kit from Qiagen. SNP markers in the extracted DNA were analyzed by genotyping. While some samples were individually genotyped, the same samples were also used for pooling studies, in which DNA from about 50 individuals was pooled, and allele frequencies were determined in pooled DNA. For studies S0012(M) and S0012(F), only male or female cases and controls were used in pooling studies. Genotypes and pool allele frequencies were obtained using a PRISM 7900HT sequence detection PCR system (Applied Biosystems, Foster City, CA) by allele-specific PCR, similar to the method described by Germer et al (Germer S., Holland M.J., Higuchi R. 2000, Genome Res. 10: 258-266). Primers for the allele-specific PCR reactions are provided in Table 5.

[0367] Summary statistics for demographic and environmental traits, history of vascular disease, and allele frequencies for the tested SNPs were obtained and compared between cases and controls. No multiple testing corrections were made.

[0368] Tests of association were calculated for both non-stratified and stratified settings: 1) Fisher's exact test of allelic association, and 2) asymptotic chi-square test of genotypic association, taking two different modes of inheritance into account (dominant, and recessive).

[0369] Effect sizes were estimated through allelic odds ratios, including 95% confidence intervals. The reported Allele 1 may be under-represented in cases (with a lower allele frequency in cases than in controls, indicating that the reported Allele1 is associated with decreased risk and the other allele is a risk factor for disease) or over-represented in cases (indicating that the reported Allele1 is a risk factor in the development of disease).

[0370] A SNP was considered to be a significant genetic marker if it exhibited a p-value < 0.05 in the allelic association test or in any of the 2 genotypic tests (dominant, recessive). The association of a marker with MI was considered replicated if the marker exhibited an allelic or genotypic association test p-value < 0.05 in one of the sample sets and the same test and strata (or substrata) were significant (p < 0.05) in another independent sample set.

[0371] An example of a replicated marker, where the reported Allele1 is associated with decreased risk for MI, is hCV8851074 (Table 6). hCV8851074 shows significant association with all individuals (strata = "ALL") of study S0028 and the non-smoking strata (Stratification = "SMOKE", Strata = "N") of study S0012. The odds ratio in both studies is less than 1 (0.84 and 0.78 respectively), using the same allele (Allele1 = "A") for analysis.

[0372] An example of a replicated marker, where the reported Allele1 is associated with increased risk for MI, is

hCV2716008 (Table 6). hCV2716008 shows significant association with all individuals (strata = "ALL") of study S0028 and all males of study S0012 (strata = "ALL", study="S0012(M)"). The odds ratio in both studies is greater than 1 (1.42 and 1.32 respectively), using the same allele (Allele1 = "C") for analysis.

Recurrent Myocardial Infarction (RMI) studies (see Table 7)

[0373] In order to identify genetic markers associated with recurrent myocardial infarction (RMI), samples from the Cholesterol and Recurrent Events (CARE) study were genotyped utilizing 864 assays for functional SNPs in 500 candidate genes. A well-documented MI was one of the enrollment criteria for the CARE study. Patients were followed up for 5 years and rates of recurrent MI were recorded in Pravastatin treated and placebo groups.

[0374] In the initial analysis (CARE), SNP genotype frequencies were compared in a group of 264 patients who had another MI (second, third, or fourth) during the 5 years of CARE follow-up (cases) versus the frequencies in the group of 1255 CARE patients who had not experienced a second MI (controls).

[0375] To replicate the initial findings, a second group of 394 CARE patients were analyzed who had a history of an MI prior to the MI at CARE enrollment but who had not experienced an MI during trial follow-up (cases), and 1221 CARE MI patients without a second MI were used as controls (Pre-CARE Study). No patients from the CARE Study were used in the Pre-CARE Study.

[0376] The SNPs replicated between CARE and Pre-CARE Studies were also tested for primary MI in study S0012 (UCSF). Study S0012 had 1400 samples. MI patients (cases) in this study had a self-reported history of MI, and controls had no history of MI or of acute angina lasting more than 1 hour. Allele frequencies in this study were detected in pooling experiments, in which DNA from about 50 individuals was pooled, and allele frequencies were determined in pooled DNA. For study S0012, only male cases and controls were used in pooling studies.

[0377] DNA was extracted from blood samples using conventional DNA extraction methods like the QIA-amp kit from Qiagen. Genotypes were obtained on a PRISM 7900HT sequence detection PCR system (Applied Biosystems) by allele-specific PCR, similar to the method described by Germer et al (Germer S., Holland M.J., Higuchi R. 2000, Genome Res. 10: 258-266). Primers for the allele-specific PCR reactions are provided in Table 5.

[0378] Statistical analysis was done using asymptotic chi square test for allelic, dominant, or recessive association, or Armitage trend test for additive genotypic association in the non-stratified as well as in the strata. Replicated SNPs are provided in Table 7. A SNP is considered replicated if the at-risk alleles in both sample sets are identical, the p-values are less than 0.05 in both sample sets, and the significant association is seen in the same stratum in both sets or one stratum is inclusive of the other.

[0379] Effect sizes were estimated through allelic odds ratios and odds ratios for dominant and recessive models, including 95% confidence intervals. Homogeneity of Cochran-Mantel-Haenszel odds ratios was tested across different strata using the Breslow-Day test. A SNP was considered to be a significant genetic marker if it exhibited a p-value < 0.05 in the allelic association test or in any of the 3 genotypic tests (dominant, recessive, additive). Haldane Odds Ratios were used if either case or control count was zero. SNPs with significant HWE violations in both cases and controls ($p < 1 \times 10^{-4}$ in both tests) were not considered for further analysis, since significant deviation from HWE in both cases and controls for individual markers can be indicative of genotyping errors. The association of a marker with RMI was considered replicated if the marker exhibits an allelic or genotypic association test p-value < 0.05 in one of the sample sets and the same test and strata are significant ($p < 0.05$) in either one or two other independent sample sets.

TABLE 1

Gene Number:	8
Celera Gene:	hCG15848 - 146000220474494
Celera Transcript:	hCT6878 - 146000220474495
Public Transcript Accession:	NM_007348
Celera Protein:	hCP35549 - 197000064932342
Public Protein Accession:	NP_031374
Gene Symbol:	ATF6
Protein Name:	activating transcription factor 6
Celera Genomic Axis:	GA_x5YUV32W2NG(94208..292030)
Chromosome:	Chr1
OMIM number:	605537
OMIM Information:	ACTIVATING TRANSCRIPTION FACTOR 6;ATF6

Transcript Sequence (SEQ ID NO:17):

EP 2 474 631 B1

(continued)

Protein Sequence (SEQ ID NO:845):

SNP Information

5	Context (SEQ ID NO:2952):	
	Celera SNP ID:	hCV15860707
	SNP Position Transcript:	224
	SNP Source:	Applera
10	Population(AAllele,Count):	african american(C,30 T,4) caucasian(C,32 T,4) total(C,62 T,8)
	SNP Type:	SILENT MUTATION
	Protein Coding: SEQ ID NO:	845, 35, (L,CTC) (L,CTT)
15	Context (SEQ ID NO:2953):	
	Celera SNP ID:	hCV22275523
	SNP Position Transcript:	552
	SNP Source:	Applera
	Population(AAllele,Count):	african american(C,4 G,28) caucasian(C,5 G,35) total(C,9 G,63)
20	SNP Type:	MISSENSE MUTATION
	Protein Coding:	SEQ ID NO:845, 145, (P,CCG) (A,GCG)
25	Context (SEQ ID NO:2954):	
	Celera SNP ID:	hCV25631983
	SNP Position Transcript:	318
	SNP Source:	Applera
	Population(AAllele,Count):	african american(A,34 G,2) caucasian(A,31 G,3) total(A,65 G,5)
	SNP Type:	MISSENSE MUTATION
30	Protein Coding: SEQ ID NO:	845, 67, (M,ATG) (V,GTG)
	Context (SEQ ID NO:2955):	
	Celera SNP ID:	hCV9079338
	SNP Position Transcript:	5598
	SNP Source:	Celera
35	Population(AAllele,Count):	no_pop(-,2 T,4) total(T,4 -,2)
	SNP Type:	UTR 3
	Protein Coding:	SEQ ID NO:845, None
40	Context (SEQ ID NO:2956):	
	Celera SNP ID:	hCV25631978
	SNP Position Transcript:	178
	SNP Source:	Applera
	Population(AAllele,Count):	african american(T,26) caucasian(C,1 T,37) total(C,1 T,63)
45	SNP Type:	MISSENSE MUTATION
	Protein Coding:	SEQ ID NO:845, 20, (S,TCT) (F,TTT)
50	SNP Source:	Applera
	Population(AAllele,Count):	african american(T,38) caucasian(C,1 T,33) total(C,1 T,71)
	SNP Type:	MISSENSE MUTATION
55		

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:2956):

Protein Coding: SEQ ID NO:845, 20, (S,TCT) (F,TTT)

5

Context (SEQ ID NO:2957):

Celera SNP ID: hCV353390

SNP Position Transcript: 4942

SNP Source: Celera

10

Population(AAllele,Count): no_pop(C,1|T,4) total(C,1|T,4)

SNP Type: UTR 3

Protein Coding: SEQ ID NO:845, None

15

Context (SEQ ID NO:2958):

Celera SNP ID: hCV353391

SNP Position Transcript: 4590

SNP Source: Celera

20

Population(AAllele,Count): no_pop(A,3|G,4) total(G,4|A,3)

SNP Type: UTR 3

Protein Coding: SEQ ID NO:845, None

25

Context (SEQ ID NO:2959):

Celera SNP ID: hCV353392

SNP Position Transcript: 4158

SNP Source: Celera

Population(AAllele,Count): no_pop(T,1|C,5) total(C,5|T,1)

30

SNP Type: UTR 3

Protein Coding: SEQ ID NO:845, None

35

Context (SEQ ID NO:2960):

Celera SNP ID: hCV25631989

SNP Position Transcript: 588

SNP Source: Applera

Population(AAllele,Count): african american(C,28|T,4) caucasian(C,35|T,5) total(C,63|T,9)

SNP Type: MISSENSE MUTATION

40

Protein Coding: SEQ ID NO:845, 157, (P,CCT) (S,TCT)

45

Context (SEQ ID NO:2961):

Celera SNP ID: hCV353393

SNP Position Transcript: 2256

SNP Source: Celera;HGBASE

Population(AAllele,Count): no_pop(A,5|G,5) total(G,5|A,5) no_pop(A,-|G,-)

SNP Type: UTR 3

Protein Coding: SEQ ID NO:845, None

50

55

EP 2 474 631 B1

(continued)

	Context (SEQ ID NO:2962):	
	Celera SNP ID:	hCV353386
5	SNP Position Transcript:	6006
	SNP Source:	Celera
	Population(AAllele,Count):	no_pop(G,1 A,4) total(G,1 A,4)
	SNP Type:	UTR 3
10	Protein Coding:	SEQ ID NO:845, None
	Context (SEQ ID NO:2963):	
	Celera SNP ID:	hCV9620281
	SNP Position Transcript:	1535
15	SNP Source:	Applera
	Population(AAllele,Count):	african american(C,35 T,1) caucasian(C,40) total(C,75 T,1)
	SNP Type:	SILENT MUTATION
	Protein Coding:	SEQ ID NO:845, 472, (N,AAC) (N,AAT)
20	Context (SEQ ID NO:2964):	
	Celera SNP ID:	hCV353388
	SNP Position Transcript:	5484
	SNP Source:	Celera
25	Population(AAllele,Count):	no_pop(G,2 A,4) total(G,2 A,4)
	SNP Type:	UTR 3
	Protein Coding:	SEQ ID NO:845, None
	Context (SEQ ID NO:2965):	
	Celera SNP ID:	hCV26368356
30	SNP Position Transcript:	1149
	SNP Source:	dbSNP
	Population(AAllele,Count):	no_pop(C,- G,-)
	SNP Type:	MISSENSE MUTATION
35	Protein Coding:	SEQ ID NO:84 344, (Q,CAA) (E,GAA)
	Context (SEQ ID NO:2966):	
	Celera SNP ID:	hCV353389
	SNP Position Transcript:	5310
40	SNP Source:	Celera
	Population(AAllele,Count):	no_pop(C,2 G,4) total(G,4 C,2)
	SNP Type:	UTR 3
	Protein Coding:	SEQ ID NO:845, None
45	Context (SEQ ID NO:2967):	
	Celera SNP ID:	hCV27836313
	SNP Position Transcript:	21
	SNP Source:	Applera
50	Population(AAllele,Count):	african american(C,32) caucasian(A,1 C,39) total(A,1 C,71)
55		

EP 2 474 631 B1

(continued)

5 Context (SEQ ID NO:2967):
 SNP Type: UTR 5
 Protein Coding: SEQ ID NO:845, None

10 SNP Source: Applera
 Population(AAllele,Count): african american(C,32) caucasian(A,1|C,31) total(A,1|C,63)
 SNP Type: UTR 5
 Protein Coding: SEQ ID NO:845, None

15 Context (SEQ ID NO:2968):
 Celera SNP ID: hCV9620376
 SNP Position Transcript: 2015
 SNP Source: HGBASE;dbSNP
 Population(AAllele,Count): no_pop(A,-|G,-) ;no_pop(A,-|G,-)
 SNP Type: SILENT MUTATION
 Protein Coding: SEQ ID NO:845, 632, (S,TCA) (S,TCG)

20 Context (SEQ ID NO:2969):
 Celera SNP ID: hCV9620377
 SNP Position Transcript: 2459
 SNP Source: HGBASE
 Population(AAllele,Count): no_pop(A,-|G,-)
 SNP Type: UTR 3
 Protein Coding: SEQ ID NO:845, None

25 Context (SEQ ID NO:2970):
 Celera SNP ID: hCV16188132
 SNP Position Transcript: 4
 SNP Source: Applera
 Population(AAllele,Count): african american(A,2|G,28) caucasian(A,5|G,35) total(A,7|G,63)
 SNP Type: UTR 5
 Protein Coding: SEQ ID NO:845, None

30 Context (SEQ ID NO:2971):
 Celera SNP ID: hCV1498763
 SNP Position Transcript: 389
 SNP Source: Applera
 Population(AAllele,Count): african american(C,3|T,33) caucasian(C,4|T,30) total(C,7|T,63)
 SNP Type: SILENT MUTATION
 Protein Coding: SEQ ID NO:845, 90, (P,CCC) (P,CCT)
 SNP Source: Celera

35
 40
 45
 50
 55

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:2971):
 Population(AAllele,Count): no_pop(C,1|T,6) total(C,1|T,6)
 5 SNP Type: SILENT MUTATION
 Protein Coding: SEQ ID NO:845, 90, (P,CCC) (P,CCT)
 Context (SEQ ID NO:2972):
 Celera SNP ID: hCV1498764
 10 SNP Position Transcript: 428
 SNP Source: Applera
 Population(AAllele,Count): african_american(A,5|G,31) caucasian(A,4|G,30) total(A,9|G,61)
 SNP Type: SILENT MUTATION
 Protein Coding: SEQ ID NO:845, 103, (S,TCA) (S,TCG)
 15
 SNP Source: Celera
 Population(AAllele,Count): no_pop(A,1|G,6) total(G,6|A,1)
 SNP Type: SILENT MUTATION
 Protein Coding: SEQ ID NO:845, 103, (S,TCA) (S,TCG)
 20 Gene Number: 8
 Celera Gene: hCG15848
 Celera Transcript: CRA|335000489562722
 Public Transcript Accession: NM_007348
 25 Celera Protein: CRA|224000006548847
 Public Protein Accession: NP_031374
 Gene Symbol: ATF6
 Protein Name: activating transcription factor 6
 Celera Genomic Axis: GA_x5YUV32W2NG(94208..292030)
 30 Chromosome: Chr1
 OMIM number: 605537
 OMIM Information: ACTIVATING TRANSCRIPTION FACTOR 6;ATF6
 35
 Transcript Sequence (SEQ ID NO:18):
 Protein Sequence (SEQ ID NO:846):
 SNP Information
 40
 Context (SEQ ID NO:2973):
 Celera SNP ID: hCV26368356
 SNP Position Transcript: 1072
 SNP Source: dbSNP
 Population(AAllele,Count): no_pop(C,-|G,-)
 45 SNP Type: MISSENSE MUTATION
 Protein Coding: SEQ ID NO:846, 344, (Q,CAA) (E,GAA)
 Context (SEQ ID NO:2974):
 Celera SNP ID: hCV25631978
 50 SNP Position Transcript: 101
 SNP Source: Applera
 Population(AAllele,Count): african_american(T,26) caucasian(C,1|T,37) total(C,1|T,63)
 SNP Type: MISSENSE MUTATION
 55

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:2974):

Protein Coding: SEQ ID NO:846, 20, (S,TCT) (F,TTT)

SNP Source: Applera

Population(AAllele,Count): african american(T,38) caucasian(C,1|T,33) total(C,1|T,71)

SNP Type: MISSENSE MUTATION

Protein Coding: SEQ ID NO:846, 20, (S,TCT) (F,TTT)

Context (SEQ ID NO:2975):

Celera SNP ID: hCV25631983

SNP Position Transcript: 241

SNP Source: Applera

Population(AAllele,Count): african american(A,34|G,2) caucasian(A,31|G,3) total(A,65|G,5)

SNP Type: MISSENSE MUTATION

Protein Coding: SEQ ID NO:846, 67, (I,ATA) (V,GTA)

Context (SEQ ID NO:2976):

Celera SNP ID: hCV25631989

SNP Position Transcript: 511

SNP Source: Applera

Population(AAllele,Count): african american(C,28|T,4) caucasian(C,35|T,5) total(C,63|T,9)

SNP Type: MISSENSE MUTATION

Protein Coding: SEQ ID NO:846, 157, (P,CCT) (S,TCT)

Context (SEQ ID NO:2977):

Celera SNP ID: hCV1498763

SNP Position Transcript: 312

SNP Source: Applera

Population(AAllele,Count): african american(C,3|T,33) caucasian(C,4|T,30) total(C,7|T,63)

SNP Type: SILENT MUTATION

Protein Coding: SEQ ID NO:846, 90, (P,CCC) (P,CCT)

SNP Source: Celera

Population(AAllele,Count): no_pop(C,1|T,6) total(C,1|T,6)

SNP Type: SILENT MUTATION

Protein Coding: SEQ ID NO:846, 90, (P,CCC) (P,CCT)

Context (SEQ ID NO:2978):

Celera SNP ID: hCV1498764

SNP Position Transcript: 351

SNP Source: Applera

Population(AAllele,Count): african american(A,5|G,31) caucasian(A,4|G,30) total(A,9|G,61)

SNP Type: SILENT MUTATION

Protein Coding: SEQ ID NO:846, 103, (S,TCA) (S,TCG)

SNP Source: Celera

EP 2 474 631 B1

(continued)

5	Context (SEQ ID NO:2978): Population(AAllele,Count): SNP Type: Protein Coding:	no_pop(A,1 G,6) total(G,6 A,1) SILENT MUTATION SEQ ID NO:846, 103, (S,TCA) (S,TCG)
10	Context (SEQ ID NO:2979): Celera SNP ID: SNP Position Transcript: SNP Source: Population(AAllele,Count): SNP Type: Protein Coding:	hCV9620376 1938 HGBASE;dbSNP no_pop(A,- G,-) ;no_pop(A,- G,-) SILENT MUTATION SEQ ID NO:846, 632, (S,TCA) (S,TCG)
15	Context (SEQ ID NO:2980): Celera SNP ID: SNP Position Transcript: SNP Source: Population(AAllele,Count): SNP Type: Protein Coding:	hCV9620377 2375 HGBASE no_pop(A,- G,-) UTR 3 SEQ ID NO:846, None
20	Context (SEQ ID NO:2981): Celera SNP ID: SNP Position Transcript: SNP Source: Population(AAllele,Count): SNP Type: Protein Coding:	hCV15860707 147 Applera african american(C,30 T,4) caucasian(C,32 T,4) total(C,62 T,8) SILENT MUTATION SEQ ID NO:846, 35, (L,CTC) (L,CTT)
25	Context (SEQ ID NO:2982): Celera SNP ID: SNP Position Transcript: SNP Source: Population(AAllele,Count): SNP Type: Protein Coding:	hCV22275523 475 Applera african american(C,4 G,28) caucasian(C,5 G,35) total(C,9 G,63) MISSENSE MUTATION SEQ ID NO:846, 145, (P,CCG) (A,GCG)
30	Context (SEQ ID NO:2983): Celera SNP ID: SNP Position Transcript: SNP Source: Population(AAllele,Count): SNP Type: Protein Coding:	hCV353393 2177 Celera;HGBASE no_pop(A,5 G,5) total(G,5 A,5) no_pop(A,- G,-) UTR 3 SEQ ID NO:846, None
35	Context (SEQ ID NO:2984): Celera SNP ID: SNP Position Transcript: SNP Source: Population(AAllele,Count): SNP Type: Protein Coding:	hCV9620281 1458 Applera african american(C,35 T,1)caucasian(C,40) total(C,75 T,1) SILENT MUTATION SEQ ID NO:846, 472, (N,AAC) (N,AAT)
40		
45		
50		
55		

TABLE 2

Gene Number: 8 / Celera Gene:hCG15848 - 146000220474494 / Gene Symbol: ATF6 / Protein Name:
activating transcription factor 6 / Celera Genomic Axis: GA_x5YUV32W2NG(88209..298031) / Chromosome:
Chr1 / OMIM number: 605537 / OMIM Information: ACTIVATING TRANSCRIPTION FACTOR 6;ATF6 //

Genomic Sequence (SEQ ID NO:17560): / SNP Information //

Context (SEQ ID NO:18962):

Celera SNP ID: hCV9620968 / SNP Position Genomic: 22101 / SNP Source: Celera;HGBASE;dbSNP /

Population(AAllele,Count): no_pop(A,1|G,2) total(G,2|A,1);no_pop(A,-|G,-) ;no_pop(A,-|G,-) SNP Type:
INTRON//

Context (SEQ ID NO:18963):

Celera SNP ID: hCV26368420 / SNP Position Genomic: 187598 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|T,5) total(T,5|-,1) / SNP Type: INTRON//

Context (SEQ ID NO:18964):

Celera SNP ID: hCV9620969 / SNP Position Genomic: 26994 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(T,3|A,6) total(A,6|T,3) ;no_pop(A,-|T,-) / SNP Type: INTRON//

Context (SEQ ID NO:18965):

Celera SNP ID: hCV26368421 / SNP Position Genomic: 192244 / SNP Source: Celera /

Population(AAllele,Count): no_pop(G,1|A,4) total(G,1|A,4) / SNP Type: INTRON//

Context (SEQ ID NO:18966):

Celera SNP ID: hCV26368423 / SNP Position Genomic: 204871 / SNP Source: Celera /

Population(AAllele,Count): SITE (TFBS)// no_pop(-,2|C,4) total(C,4|-,2) / SNP Type:TRANSCRIPTION
FACTOR BINDING

Context (SEQ ID NO:18967):

Celera SNP ID: hCV26368351 / SNP Position Genomic: 39313 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(G,2|C,7) total(G,2|C,7);(C,-|G,-) Caucasian(G,10|C,74)

African-American(G,31|C,53) Asians(G,37|C,47) total(G,78|C,174) / SNP Type: INTRON//

Context (SEQ ID NO:18968):

Celera SNP ID: hCV26368353 / SNP Position Genomic: 41130 / SNP Source: Celera / Population(AAllele,Count):

no_pop(C,1 |T,6) total(C,1|T,6) / SNP Type: INTRON//

Context (SEQ ID NO:18969):

Celera SNP ID: hCV25552 / SNP Position Genomic: 136326 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,3|C,7) total(A,3|C,7) / SNP Type: INTRON/hmCS//

Context (SEQ ID NO:18970):

Celera SNP ID: hCV26368355 / SNP Position Genomic: 45535 / SNP Source: Celera / Population(AAllele,Count):

no_pop(G,1|A,5) total(G,1|A,5) / SNP Type: INTRON//

Context (SEQ ID NO:18971):

Celera SNP ID: hCV26368356 / SNP Position Genomic: 59498 / SNP Source: dbSNP /

Population(AAllele,Count): no_pop(C,-|G,-) / SNP Type: MISSENSE MUTATION//

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:18972):

Celera SNP ID: hCV26368358 / SNP Position Genomic: 60499 / SNP Source: Celera / Population(Allele, Count):
no_pop(C,2|T,8) total(C,2|T,8) / SNP Type: INTRON//

Context (SEQ ID NO:18973):

Celera SNP ID: hCV26368359 / SNP Position Genomic: 61415 / SNP Source: Celera / Population(Allele, Count):
no_pop(T,1|C,8) total(C,8|T,1) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:18974):

Celera SNP ID: hCV15860707 / SNP Position Genomic: 18023 / SNP Source: Applera /
Population(Allele, Count): african american(C,301T,4) caucasian(C,32|T,4) total(C,62|T,8) / SNP Type:
SILENT MUTATION //

Context (SEQ ID NO:18975):

Celera SNP ID: hCV9620976 / SNP Position Genomic: 40604 / SNP Source: Celera;HGBASE;dbSNP /
Population(Allele, Count): no_pop(C,1|T,6) total(C,1|T,6);no_pop(C,-|T,-) ;no_pop(C,-|T,-) / SNP Type:
INTRON//

Context (SEQ ID NO:18976):

Celera SNP ID: hCV9620977 / SNP Position Genomic: 41474 / SNP Source: dbSNP / Population(Allele, Count):
no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:18977):

Celera SNP ID: hCV519954 / SNP Position Genomic: 177032 / SNP Source: Celera / Population(Allele, Count):
no_pop(A,3|G,5) total(G,5|A,3) / SNP Type: INTRON//

Context (SEQ ID NO:18978):

Celera SNP ID: hCV67446 / SNP Position Genomic: 184250 / SNP Source: Celera / Population(Allele, Count):
no_pop(G,3|A,9) total(G,3|A,9) / SNP Type: INTRON//

Context (SEQ ID NO:18979):

Celera SNP ID: hCV67447 / SNP Position Genomic: 185354 / SNP Source: Celera / Population(Allele, Count):
no_pop(A,1|T,3) total(A,1|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:18980):

Celera SNP ID: hCV519956 / SNP Position Genomic: 176303 / SNP Source: Celera / Population(Allele, Count):
no_pop(A,3|T,6) total(A,3|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:18981):

Celera SNP ID: hCV67448 / SNP Position Genomic: 187556 / SNP Source: Celera / Population(Allele, Count):
no_pop(C,1|T,8) total(C,1|T,8) / SNP Type: INTRON //

Context (SEQ ID NO:18982):

Celera SNP ID: hCV519957 / SNP Position Genomic: 179566 / SNP Source: Celera / Population(Allele, Count):
no_pop(C,2|G,4) total(G,4|C,2) / SNP Type: INTRON //

Context (SEQ ID NO:18983):

Celera SNP ID: hCV26368360 / SNP Position Genomic: 64132 / SNP Source: Celera / Population(Allele, Count):
no_pop(A,1|- ,5) total(A,1|- ,5) / SNP Type: INTRON //

Context (SEQ ID NO:18984):

Celera SNP ID: hCV26368362 / SNP Position Genomic: 66677 / SNP Source: Celera / Population(Allele, Count):
no_pop(T,1|- ,5) total(- ,5|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:18985):

Celera SNP ID: hCV26368363 / SNP Position Genomic: 66740 / SNP Source: Celera / Population(Allele, Count):
no_pop(T,1|C,7) total(C,7|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:18986):

Celera SNP ID: hCV427609 / SNP Position Genomic: 143653 / SNP Source: Celera / Population(Allele, Count):
no_pop(C,3|A,7) total(A,7|C,3) / SNP Type: INTRON //

Context (SEQ ID NO:18987):

Celera SNP ID: hCV26368364 / SNP Position Genomic: 69455 / SNP Source: Celera / Population(Allele, Count):
no_pop(G,1|A,4) total(G,1|A,4) / SNP Type: INTRON //

Context (SEQ ID NO:18988):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV26368365 / SNP Position Genomic: 69456 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,1|A,4) total(G,1|A,4) / SNP Type: INTRON //

Context (SEQ ID NO:18989):

Celera SNP ID: hCV26368366 / SNP Position Genomic: 69606 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,2|A,8) total(G,2|A,8) / SNP Type: INTRON //

Context (SEQ ID NO:18990):

Celera SNP ID: hCV26368368 / SNP Position Genomic: 70099 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,5|C,8) total(G,5|C,8) / SNP Type: INTRON //

Context (SEQ ID NO:18991):

Celera SNP ID: hCV26368369 / SNP Position Genomic: 70102 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,3|A,8) total(G,3|A,8) / SNP Type: INTRON //

Context (SEQ ID NO:18992):

Celera SNP ID: hCV16188132 / SNP Position Genomic: 6003 / SNP Source: Applera / Population(Allele,Count):
african american(A,2|G,28) caucasian(A,5|G,35) total(A,7|G,63) / SNP Type: UTR 5 / SNP Source: Applera /
Population(Allele,Count): african american(A,1|G,25) caucasian(A,4|G,30) total(A,5|G,55) / SNP Type: UTR
5 //

Context (SEQ ID NO:18993):

Celera SNP ID: hCV353400 / SNP Position Genomic: 194716 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,4|A,5) total(G,4|A,5) / SNP Type: INTRON //

Context (SEQ ID NO:18994):

Celera SNP ID: hCV353401 / SNP Position Genomic: 194331 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,1|A,6) total(A,6|T,1) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:18995):

Celera SNP ID: hCV353402 / SNP Position Genomic: 193004 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,4|A,5) total(G,4|A,5) / SNP Type: INTRON //

Context (SEQ ID NO:18996):

Celera SNP ID: hCV9078252 / SNP Position Genomic: 187959 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,4|A,5) total(A,5|C,4) / SNP Type: INTRON //

Context (SEQ ID NO:18997):

Celera SNP ID: hCV67450 / SNP Position Genomic: 187716 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,1|G,5) total(G,5|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:18998):

Celera SNP ID: hCV427612 / SNP Position Genomic: 140896 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,2|G,9) total(G,9|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:18999):

Celera SNP ID: hCV26368371 / SNP Position Genomic: 81405 / SNP Source: dbSNP /
Population(Allele,Count): (C,-|T,-) Caucasian(T,84) African-American(T,84) Asians(T,84) total(T,252) / SNP
Type: INTRON //

Context (SEQ ID NO:19000):

Celera SNP ID: hCV26368372 / SNP Position Genomic: 83036 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,3|T,9) total(T,9|-,3) / SNP Type: INTRON //

Context (SEQ ID NO:19001):

Celera SNP ID: hCV515827 / SNP Position Genomic: 133854 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,1|A,5) total(G,1|A,5) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19002):

Celera SNP ID: hCV26368373 / SNP Position Genomic: 85823 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|C,8) total(C,8|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19003):

Celera SNP ID: hCV9621426 / SNP Position Genomic: 58071 / SNP Source: dbSNP / Population(Allele,Count):
(A,-|T,-) Caucasian(A,84) African-American(A,84) Asians(A,84) total(A,252) / SNP Type: INTRON //

Context (SEQ ID NO:19004):

Celera SNP ID: hCV26368374 / SNP Position Genomic: 90980 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|A,5) total(A,5|-,1) / SNP Type: INTRON //

(continued)

Context (SEQ ID NO:19005):

Celera SNP ID: hCV26368375 / SNP Position Genomic: 90981 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|C,5) total(C,5|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19006):

Celera SNP ID: hCV27836313 / SNP Position Genomic: 6020 / SNP Source: Applera / Population(Allele,Count):
african american(C,32) caucasian(A,1|C,39) total(A,1|C,71) / SNP Type: UTR 5/ SNP Source: Applera /
Population(Allele,Count): african american(C,32) caucasian(A,1|C,31) total(A,1|C,63) / SNP Type: UTR 5 //

Context (SEQ ID NO:19007):

Celera SNP ID: hCV26368376 / SNP Position Genomic: 90982 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|A,5) total(A,5|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19008):

Celera SNP ID: hCV26368377 / SNP Position Genomic: 90984 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|C,1|-,5) total(A,1|C,1|-,5) / SNP Type: INTRON //

Context (SEQ ID NO:19009):

Celera SNP ID: hCV26368378 / SNP Position Genomic: 91003 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|C,5) total(C,5|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19010):

Celera SNP ID: hCV315093 / SNP Position Genomic: 180472 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,2|C,7) total(C,7|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19011):

Celera SNP ID: hCV26368379 / SNP Position Genomic: 91005 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|C,5) total(C,5|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19012):

Celera SNP ID: hCV27849856 / SNP Position Genomic: 8031 / SNP Source: Applera / Population(Allele,Count):
african american(A,28|G,8) caucasian(A,34|G,6) total(A,62|G,14) / SNP Type: INTRON //

Context (SEQ ID NO:19013):

Celera SNP ID: hCV26368384 / SNP Position Genomic: 120075 / SNP Source: Celera /
Population(Allele,Count): no_pop(G,1|T,1|-,9) total(G,1|-,9|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19014):

Celera SNP ID: hCV26368385 / SNP Position Genomic: 126994 / SNP Source: Celera /
Population(Allele,Count): no_pop(-,1|T,7) total(T,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19015): Celera SNP ID: hCV26368386 / SNP Position Genomic: 127084 / SNP Source:
Celera / Population(Allele,Count): no_pop(A,1|C,9) total(A,1|C,9) / SNP Type: INTRON //

Context (SEQ ID NO:19016):

Celera SNP ID: hCV25631977 / SNP Position Genomic: 152012 / SNP Source: Applera /
Population(Allele,Count): african american(A,33|G,1) caucasian(A,40) total(A,73|G,1) / SNP Type: INTRON //

Context (SEQ ID NO:19017):

Celera SNP ID: hCV25631978 / SNP Position Genomic: 6177 / SNP Source: Applera / Population(Allele,Count):
african american(T,26) caucasian(C,1|T,37) total(C,1|T,63) / SNP Type: MISSENSE MUTATION/ SNP Source:
Applera / Population(Allele,Count): african american(T,38) caucasian(C,1|T,33) total(C,1|T,71) / SNP Type:
MISSENSE MUTATION //

Context (SEQ ID NO:19018):

Celera SNP ID: hCV25631979 / SNP Position Genomic: 6227 / SNP Source: Applera / Population(Allele,Count):
african american(G,38) caucasian(G,33|T,1) total(G,71|T,1) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19019):

Celera SNP ID: hCV16127908 / SNP Position Genomic: 50937 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|G,6) total(G,6|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19020):

Celera SNP ID: hCV16127909 / SNP Position Genomic: 51158 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,1|A,4) total(A,4|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19021):

Celera SNP ID: hCV26368391 / SNP Position Genomic: 142845 / SNP Source: Celera /
Population(Allele,Count): no_pop(G,2|T,7) total(G,2|T,7) / SNP Type: INTRON

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:19022):

Celera SNP ID: hCV26368392 / SNP Position Genomic: 143571 / SNP Source: Celera / Population(AAllele,Count): no_pop(-,1|T,8) total(T,8|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19023):

Celera SNP ID: hCV26368393 / SNP Position Genomic: 143572 / SNP Source: Celera / Population(AAllele,Count): no_pop(-,1|C,8) total(C,8|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19024):

Celera SNP ID: hCV26368394 / SNP Position Genomic: 143573 / SNP Source: Celera / Population(AAllele,Count): no_pop(-,1|T,8) total(T,8|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19025):

Celera SNP ID: hCV26368395 / SNP Position Genomic: 143574 / SNP Source: Celera / Population(AAllele,Count): no_pop(-,1|G,8) total(G,8|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19026):

Celera SNP ID: hCV150005 / SNP Position Genomic: 180749 / SNP Source: Celera / Population(AAllele,Count): no_pop(C,3|T,8) total(C,3|T,8) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19027):

Celera SNP ID: hCV26368399 / SNP Position Genomic: 155229 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,2|C,10) total(C,10|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19028):

Celera SNP ID: hCV150006 / SNP Position Genomic: 182070 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,3|G,5) total(G,5|A,3) / SNP Type: INTRON //

Context (SEQ ID NO:19029):

Celera SNP ID: Population(AAllele,Count): hCV25631980 / SNP Position Genomic: 6240 / SNP Source: Applera / african american(C,38) caucasian(C,33|T,1) total(C,71|T,1) / SNP Type:INTRON/hmCS //

Context (SEQ ID NO:19030):

Celera SNP ID: hCV25631981 / SNP Position Genomic: 6341 / SNP Source: Applera / Population(AAllele,Count): african american(C,36) caucasian(C,29|T,1) total(C,65|T,1) / SNP Type:INTRON //

Context (SEQ ID NO:19031):

Celera SNP ID: Population(AAllele,Count): hCV25631982 / SNP Position Genomic: 18220 / SNP Source: Applera / african american(C,3|T,33) caucasian(T,38) total(C,3|T,71) / SNP Type:INTRON //

Context (SEQ ID NO:19032):

Celera SNP ID: Population(AAllele,Count): hCV25631983 / SNP Position Genomic: 21708 / SNP Source: Applera / african american(A,34|G,2) caucasian(A,31|G,3) total(A,65|G,5) / SNP Type: //MISSENSE MUTATION

Context (SEQ ID NO:19033):

Celera SNP ID: Population(AAllele,Count): hCV25631984 / SNP Position Genomic: 21933 / SNP Source: Applera / african american(A,31|G,1) caucasian(A,34) total(A,65|G,1) / SNP Type: INTRON //

Context (SEQ ID NO:19034):

Celera SNP ID: Population(AAllele,Count): hCV25631985 / SNP Position Genomic: 22070 / SNP Source: Applera / african american(C,28) caucasian(C,27|G,1) total(C,55|G,1) / SNP Type: INTRON //

Context (SEQ ID NO:19035):

Celera SNP ID: Population(AAllele,Count): hCV25631987 / SNP Position Genomic: 23629 / SNP Source: Applera / african american(A,1|G,35) caucasian(G,36) total(A,1|G,71) / SNP Type: INTRON //

Context (SEQ ID NO:19036):

Celera SNP ID: Population(AAllele,Count): hCV25631988 / SNP Position Genomic: 31020 / SNP Source: Applera / african american(G,4|T,26) caucasian(G,4|T,26) total(G,8|T,52) / SNP Type: INTRON //

Context (SEQ ID NO:19037):

Celera SNP ID: hCV408552 / SNP Position Genomic: 130468 / SNP Source: Celera / Population(AAllele,Count): no_pop(C,1|A,4) total (A,4|C,1) / SNP Type: INTRON //

Context (SEQ ID NO:19038):

Celera SNP ID: Population(AAllele,Count): hCV25631989 / SNP Position Genomic: 31270 / SNP Source: Applera / african american(C,28|T,4) caucasian(C,35|T,5) total(C,63|T,9) / SNP Type: //MISSENSE MUTATION

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:19039):

Celera SNP ID: hCV16127916 // SNP Position Genomic: 98114 / SNP Source: Celera /

Population(AAllele,Count): no_pop(G,1|T,7) total(G,1|T,7) / SNP Type: INTRON //

Context (SEQ ID NO:19040):

Celera SNP ID: hCV9078862 / SNP Position Genomic: 85933 / SNP Source: Celera / Population(AAllele,Count):

no_pop(G,2|A,9) total(G,2|A,9) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19041):

Celera SNP ID: hCV9078867 / SNP Position Genomic: 86064 / SNP Source: Celera / Population(AAllele,Count):

no_pop(T,3|G,10) total(G,10|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19042):

Celera SNP ID: hCV16079344 / SNP Position Genomic: 146402 / SNP Source: Celera /

Population(AAllele,Count): no_pop(A,1|T,6) total(A,1|T,6) / SNP Type: INTRON/hmCS//

Context (SEQ ID NO:19043):

Celera SNP ID: hCV22275513 / SNP Position Genomic: 31057 / SNP Source: Applera /

Population(AAllele,Count): african american(C,25|T,7) caucasian(C,34|T,6) total(C,59|T,13) / SNP Type:INTRON //

Context (SEQ ID NO:19044):

Celera SNP ID: hCV25631990 Population(AAllele,Count): INTRON /// SNP Position Genomic: 59607 / SNP

Source: Applera / african american(C,37|T,1) caucasian(C,40) total(C,77|T,1) / SNP Type:

Context (SEQ ID NO:19045):

Celera SNP ID: hCV25631991 / SNP Position Genomic: 59792 / SNP Source: Applera /

Population(AAllele,Count): african american(G,3|T,35) caucasian(T,40) total(G,3|T,75) / SNP Type:INTRON //

Context (SEQ ID NO:19046):

Celera SNP ID: hCV1498800 / SNP Position Genomic: 75978 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,1|G,5) total(G,5|A,1) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19047):

Celera SNP ID: hCV1498801 / SNP Position Genomic: 76240 / SNP Source: Celera / Population(AAllele,Count):

no_pop(T,2|C,6) total(C,6|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19048):

Celera SNP ID: hCV1498802 / SNP Position Genomic: 76312 / SNP Source: Celera / Population(AAllele,Count):

no_pop(C,2|G,7) total(G,7|C,2) / SNP Type: INTRON //

Context (SEQ ID NO:19049):

Celera SNP ID: hCV1498803 / SNP Position Genomic: 77822 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(C,3|T,6) total(C,3|T,6) ;no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19050):

Celera SNP ID: hCV1498804 / SNP Position Genomic: 78209 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,1|G,8) total(G,8|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19051):

Celera SNP ID: hCV1498805 / SNP Position Genomic: 79662 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,2|T,8) total(A,2|T,8) / SNP Type: INTRON //

Context (SEQ ID NO:19052):

Celera SNP ID: hCV1498806 / SNP Position Genomic: 80107 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,1|G,9) total(G,9|A,1) / SNP Type: INTRON /

Context (SEQ ID NO:19053):

Celera SNP ID: hCV1498807 / SNP Position Genomic: 80140 / SNP Source: Celera / Population(AAllele,Count):

no_pop(T,1|C,8) total(C,8|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19054):

Celera SNP ID: hCV1498808 / SNP Position Genomic: 84582 / SNP Source: Celera / Population(AAllele,Count):

no_pop(G,2|T,6) total(G,2|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19055):

Celera SNP ID: hCV1498809 / SNP Position Genomic: 86007 / SNP Source: Celera / Population(AAllele,Count):

no_pop(G,3|A,9) total(G,3|A,9) / SNP Type: INTRON 11

Context (SEQ ID NO:19056):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV15991581 / SNP Position Genomic: 16645 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,2|T,2) total(C,2|T,2) / SNP Type: INTRON 11

Context (SEQ ID NO:19057):

Celera SNP ID: hCV15991582 / SNP Position Genomic: 16739 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,1|C,2) total(C,2|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19058):

Celera SNP ID: hCV22275523 / SNP Position Genomic: 31234 / SNP Source: Applera /
Population(Allele,Count): african american(C,4|G,28) caucasian(C,5|G,35) total(C,9|G,63) / SNP
Type:MISSENSE MUTATION//

Context (SEQ ID NO:19059):

Celera SNP ID: hCV353386 / SNP Position Genomic: 202285 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,1 |A,4) total(G,1 |A,4) / SNP Type: UTR 3 //

Context (SEQ ID NO:19060):

Celera SNP ID: hCV353388 / SNP Position Genomic: 201763 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,2|A,4) total(G,2|A,4) / SNP Type: UTR 3 //

Context (SEQ ID NO:19061):

Celera SNP ID: hCV11629011 / SNP Position Genomic: 34751 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|G,11) total(G,11|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19062):

Celera SNP ID: hCV1498810 / SNP Position Genomic: 87984 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,1|A,9) total(A,9|C,1) / SNP Type: INTRON //

Context (SEQ ID NO:19063):

Celera SNP ID: hCV353389 / SNP Position Genomic: 201589 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,2|G,4) total(G,4|C,2) / SNP Type: UTR 3 //

Context (SEQ ID NO:19064):

Celera SNP ID: hCV11629012 / SNP Position Genomic: 45559 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,2|A,3) total(A,3|-,2) / SNP Type: INTRON //

Context (SEQ ID NO:19065):

Celera SNP ID: hCV1498811 / SNP Position Genomic: 90983 / SNP Source: Celera / Population(Allele,Count):
no_pop(-,1|C,5) total(C,5|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19066):

Celera SNP ID: hCV1498814 / SNP Position Genomic: 96548 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,2|G,10) total(G,10|A,2) / SNP Type: INTRON //

Context (SEQ ID NO:19067):

Celera SNP ID: hCV1498815 / SNP Position Genomic: 96571 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,5|T,7) total(A,5|T,7) / SNP Type: INTRON //

Context (SEQ ID NO:19068):

Celera SNP ID: hCV1498816 / SNP Position Genomic: 96892 / SNP Source: Celera Population(Allele,Count):
no_pop(A,1|G,13) total(G,13|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19069):

Celera SNP ID: hCV1498817 / SNP Position Genomic: 98646 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,1|T,10) total(G,1|T,10) / SNP Type: INTRON //

Context (SEQ ID NO:19070):

Celera SNP ID: hCV1498818 / SNP Position Genomic: 99126 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,2|A,7) total(G,2|A,7) / SNP Type: INTRON //

Context (SEQ ID NO:19071):

Celera SNP ID: hCV353390 / SNP Position Genomic: 201221 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,1 |T,4) total(C,1 |T,4) / SNP Type: UTR 3 //

Context (SEQ ID NO:19072):

Celera SNP ID: hCV353391 / SNP Position Genomic: 200869 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,3|G,4) total(G,4|A,3) / SNP Type: UTR 3 //

Context (SEQ ID NO:19073):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV353392 / SNP Position Genomic: 200437 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,1|C,5) total(C,5|T,1) / SNP Type: UTR 3 //

Context (SEQ ID NO:19074):

Celera SNP ID: hCV353393 / SNP Position Genomic: 198535 / SNP Source: Celera;HGBASE /
Population(AAllele,Count): no_pop(A,5|G,5) total(G,5|A,5) ;no_pop(A,-|G,-) / SNP Type: UTR 3 //

Context (SEQ ID NO:19075):

Celera SNP ID: hCV353394 / SNP Position Genomic: 197969 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,3|T,5) total(C,3|T,5) / SNP Type: INTRON //

Context (SEQ ID NO:19076):

Celera SNP ID: hCV353395 / SNP Position Genomic: 197852 / SNP Source: Celera / Population(AAllele,Count):
no_pop(G,3|C,5) total(G,3|C,5) / SNP Type: INTRON //

Context (SEQ ID NO:19077):

Celera SNP ID: hCV353396 / SNP Position Genomic: 197146 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,3|G,7) total(G,7|C,3) / SNP Type: INTRON //

Context (SEQ ID NO:19078):

Celera SNP ID: hCV353397 / SNP Position Genomic: 197070 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,3|T,5) total(C,3|T,5) / SNP Type: INTRON //

Context (SEQ ID NO:19079):

Celera SNP ID: hCV353398 / SNP Position Genomic: 196536 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,4|G,4) total(G,4|A,4) / SNP Type: INTRON //

Context (SEQ ID NO:19080):

Celera SNP ID: hCV1498820 / SNP Position Genomic: 100766 / SNP Source: dbSNP /
Population(AAllele,Count): no_pop(A,-|C,-) / SNP Type: INTRON //

Context (SEQ ID NO:19081):

Celera SNP ID: hCV353399 / SNP Position Genomic: 195404 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,5|G,6) total(G,6|C,5) / SNP Type: INTRON//

Context (SEQ ID NO:19082):

Celera SNP ID: hCV1498821 / SNP Position Genomic: 101854 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|G,9) total(G,9|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19083):

Celera SNP ID: hCV1498822 / SNP Position Genomic: 102798 / SNP Source: Applera /
Population(AAllele,Count): african american(G,6|T,14) caucasian(G,10|T,6) total(G,16|T,20) / SNP Type: Celera
/ Population(AAllele,Count): no_pop(G,5|T,6) total(G,5|T,6) / SNP Type:INTRON / SNP Source: INTRON //

Context (SEQ ID NO:19084):

Celera SNP ID: hCV1498823 / SNP Position Genomic: 104402 / SNP Source: Celera / Population(AAllele,Count):
no_pop(G,1|A,12) total(G,1|A,12) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19085):

Celera SNP ID: hCV1498824 / SNP Position Genomic: 104536 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|T,9) total(A,1|T,9) / SNP Type: INTRON //

Context (SEQ ID NO:19086):

Celera SNP ID: hCV1498825 / SNP Position Genomic: 104737 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,2|C,6) total(C,6|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19087):

Celera SNP ID: hCV1498826 / SNP Position Genomic: 104971 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,3|T,8) total(C,3|T,8) / SNP Type: INTRON //

Context (SEQ ID NO:19088):

Celera SNP ID: hCV1498754 / SNP Position Genomic: 6674 / SNP Source: Applera / Population(AAllele,Count):
african american (C,18|T,18) caucasian(C,34|T,6) total(C,52|T,24) / SNP Type: Source: Celera /
Population(AAllele,Count): no_pop(T,1|C,3) total(C,3|T,1) / SNP INTRON/hmCS / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19089):

Celera SNP ID: hCV1498827 / SNP Position Genomic: 106799 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,3|G,7) total(G,7|T,3) / SNP Type: INTRON/TFBS//

Context (SEQ ID NO:19090):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV1498755 / SNP Position Genomic: 9158 / SNP Source: Celera / Population(AAllele,Count):
no_pop(G,1|A,6) total(G,1 |A,6) / SNP Type: INTRON //

Context (SEQ ID NO:19091):

Celera SNP ID: hCV1498828 / SNP Position Genomic: 106946 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,1|C,9) total(C,9|T,1) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19092):

Celera SNP ID: hCV1498756 / SNP Position Genomic: 10278 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,1 |T,6) total(C,1 |T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19093):

Celera SNP ID: hCV1498829 / SNP Position Genomic: 108084 / SNP Source: Celera;HGBASE;dbSNP /
Population(AAllele,Count): no_pop(T,4|C,10) total(C,10|T,4) ;no_pop(C,-|T,-) ;no_pop(C,-|T,-) / SNP Type:
INTRON //

Context (SEQ ID NO:19094):

Celera SNP ID: hCV1498757 / SNP Position Genomic: 10382 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|T,7) total(A,1|T,7) / SNP Type: INTRON //

Context (SEQ ID NO:19095):

Celera SNP ID: hCV1498758 / SNP Position Genomic: 10451 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,3|G,5) total(G,5|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19096):

Celera SNP ID: hCV1498759 / SNP Position Genomic: 11242 / SNP Source: Celera;dbSNP /
Population(AAllele,Count): no_pop(C,3|T,5) total(C,3|T,5) ;no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19097):

Celera SNP ID: hCV16083251 / SNP Position Genomic: 59146 / SNP Source: dbSNP /
Population(AAllele,Count): no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19098):

Celera SNP ID: hCV9079338 / SNP Position Genomic: 201877 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,2|T,4) total(T,4|-,2) / SNP Type: UTR 3 //

Context (SEQ ID NO:19099):

Celera SNP ID: hCV11626121 / SNP Position Genomic: 135717 / SNP Source: Celera /
Population(AAllele,Count): no_pop(A,2|C,6) total(A,2|C,6) / SNP Type: INTRON //

Context (SEQ ID NO:19100):

Celera SNP ID: hCV11454887 / SNP Position Genomic: 19779 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|G,3) total(G,3|A,1) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19101):

Celera SNP ID: hCV11454889 / SNP Position Genomic: 56549 / SNP Source: dbSNP /
Population(AAllele,Count): no_pop(C,-|T,-) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19102):

Celera SNP ID: hCV11626127 / SNP Position Genomic: 142681 / SNP Source: Celera /
Population(AAllele,Count): no_pop(A,3|G,4) total(G,4|A,3) / SNP Type: INTRON //

Context (SEQ ID NO:19103):

Celera SNP ID: hCV1498830 / SNP Position Genomic: 109547 / SNP Source: Celera / Population(AAllele,Count):
no_pop(G,1|A,9) total(G,1 |A,9) SNP Type: INTRON //

Context (SEQ ID NO:19104):

Celera SNP ID: hCV27834348 / SNP Position Genomic: 8206 / SNP Source: Applera / Population(AAllele,Count):
african american(A,26|C,4) caucasian(A,13|C,7) total(A,39|C,11) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19105):

Celera SNP ID: hCV1498760 / SNP Position Genomic: 11586 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,1 |T,7) total(C,1 |T,7) / SNP Type: INTRON //

Context (SEQ ID NO:19106):

Celera SNP ID: hCV1498833 / SNP Position Genomic: 111996 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|G,5) total(G,5|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19107):

(continued)

Celera SNP ID: hCV1498834 / SNP Position Genomic: 112163 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|G,5) total(G,5|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19108):

Celera SNP ID: hCV1498762 / SNP Position Genomic: 19982 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,1|C,3) total(G,1|C,3) / SNP Type: INTRON//

Context (SEQ ID NO:19109):

Celera SNP ID: hCV1498763 / SNP Position Genomic: 23769 / SNP Source: Applera / Population(Allele,Count):
african american(C,3|T,33) caucasian(C,4|,30) total(C,7|T,63) / SNP Type: SILENT MUTATION / SNP Source:
Celera / Population(Allele,Count): no_pop(C,1 |T,6) total(C,1 |T,6) / //SNP Type: SILENT MUTATION

Context (SEQ ID NO:19110):

Celera SNP ID: hCV1498836 / SNP Position Genomic: 120026 / SNP Source: Celera;dbSNP /
Population(Allele,Count): no_pop(C,1|A,9) total(A,9|C,1);no_pop(A,-|C,-) / SNP Type: INTRON //

Context (SEQ ID NO:19111):

Celera SNP ID: hCV1498764 / SNP Position Genomic: 23808 / SNP Source: Applera / Population(Allele,Count):
african SILENT MUTATION / SNP Source: american(A,5|G,31) caucasian(A,4|G,30) total(A,9|G,61) / SNP
Type: Celera / Population(Allele,Count): no_pop(A,1|G,6) total(G,6|A,1) / //SNP Type: SILENT MUTATION

Context (SEQ ID NO:19112):

Celera SNP ID: hCV1498765 / SNP Position Genomic: 23884 / SNP Source: Applera / Population(Allele,Count):
african american (C,5|G,31) caucasian(C,5|G,31) total(C,10|G,62) / SNP Type:

INTRON / SNP Source: Celera;dbSNP / Population(Allele,Count): no_pop(C,1|G,6)
total(G,6|C,1) ;no_pop(C,-|G,-) / SNP Type: 1 NTRON //

Context (SEQ ID NO:19113):

Celera SNP ID: hCV1498766 / SNP Position Genomic: 23971 / SNP Source: Applera / Population(Allele,Count):
african american(A,4|T,26) caucasian(A,2|T,32) total(A,6|T,58) / SNP Type: INTRON / SNP Source:

Celera;HGBASE;dbSNP / Population(Allele,Count): no_pop(A,1|T,7) total(A,1|T,7) ; no_pop(A,-|T,-) ;
no_pop(A,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19114):

Celera SNP ID: hCV1498767 / SNP Position Genomic: 24801 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,1 |T,3) total(C,1 |T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19115):

Celera SNP ID: hCV209451 / SNP Position Genomic: 161493 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,5|G,5) total(G,5|A,5) / SNP Type: INTRON //

Context (SEQ ID NO:19116):

Celera SNP ID: hCV1498769 / SNP Position Genomic: 32582 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|G,3) total(G,3|A,1) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19117):

Celera SNP ID: hCV209453 / SNP Position Genomic: 162444 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,4|C,4) total(A,4|C,4) / SNP Type: INTRON //

Context (SEQ ID NO:19118):

Celera SNP ID: hCV209454 / SNP Position Genomic: 162897 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,3|A,4) total(A,4|C,3) / SNP Type: INTRON //

Context (SEQ ID NO:19119):

Celera SNP ID: hCV209455 / SNP Position Genomic: 162973 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,3|T,3) total(C,3|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19120):

Celera SNP ID: hCV209456 / SNP Position Genomic: 164053 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,4|G,7) total(G,7|C,4) / SNP Type: INTRON //

Context (SEQ ID NO:19121):

Celera SNP ID: hCV209457 / SNP Position Genomic: 165905 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,5|T,8) total(G,5|T,8) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19122):

Celera SNP ID: hCV209458 / SNP Position Genomic: 166971 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,3|G,8) total(G,8|A,3) / SNP Type: INTRON //

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:19123):

Celera SNP ID: hCV209459 / SNP Position Genomic: 167463 / SNP Source: Celera / Population(AAllele,Count):
no_pop(G,3|C,7) total(G,3|C,7) / SNP Type: INTRON //

Context (SEQ ID NO:19124):

Celera SNP ID: hCV9620261 / SNP Position Genomic: 78385 / SNP Source: Celera;dbSNP /
Population(AAllele,Count): no_pop(G,1|T,11) total(G,1|T,11) ;no_pop(G,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19125):

Celera SNP ID: hCV11454893 / SNP Position Genomic: 99304 / SNP Source: Celera;dbSNP /
Population(AAllele,Count): no_pop(G,1|A,7) total(G,1|A,7) ;no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19126):

Celera SNP ID: hCV11626130 / SNP Position Genomic: 155306 / SNP Source: Celera /
Population(AAllele,Count): no_pop(C,4|A,7) total(A,7|C,4) / SNP Type: INTRON //

Context (SEQ ID NO:19127):

Celera SNP ID: hCV9620262 / SNP Position Genomic: 81511 / SNP Source: dbSNP / Population(AAllele,Count):
no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19128):

Celera SNP ID: hCV11626131 / SNP Position Genomic: 156993 / SNP Source: Celera /
Population(AAllele,Count): no_pop(C,1|A,6) total(A,6|C,1) / SNP Type: INTRON //

Context (SEQ ID NO:19129):

Celera SNP ID: hCV9620263 / SNP Position Genomic: 81544 / SNP Source: dbSNP / Population(AAllele,Count):
no_pop(G,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19130):

Celera SNP ID: hCV11626133 / SNP Position Genomic: 157309 / SNP Source: Celera /
Population(AAllele,Count): no_pop(T,1|C,5) total(C,5|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19131):

Celera SNP ID: hCV16206410 / SNP Position Genomic: 131977 / SNP Source: Celera /
Population(AAllele,Count): no_pop(A,1|G,6) total(G,6|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19132):

Celera SNP ID: hCV11626137 / SNP Position Genomic: 166680 / SNP Source: Celera /
Population(AAllele,Count): no_pop(A,3|T,3|-4) total(A,3|-4|T,3) / SNP Type: INTRON/hmCS /

Context (SEQ ID NO:19133):

Celera SNP ID: hCV49962 / SNP Position Genomic: 188275 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,1|A,9) total(A,9|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19134):

Celera SNP ID: hCV49963 / SNP Position Genomic: 188749 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,4|G,5) total(G,5|C,4) / SNP Type: INTRON //

Context (SEQ ID NO:19135):

Celera SNP ID: hCV1498770 / SNP Position Genomic: 33687 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|G,8) total(G,8|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19136):

Celera SNP ID: hCV1498771 / SNP Position Genomic: 33891 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,3|G,8) total(G,8|A,3) / SNP Type: INTRON //

Context (SEQ ID NO:19137):

Celera SNP ID: hCV1498773 / SNP Position Genomic: 35792 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,2|G,5) total(G,5|A,2) / SNP Type: INTRON //

Context (SEQ ID NO:19138):

Celera SNP ID: hCV1498774 / SNP Position Genomic: 36817 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,2|G,4) total(G,4|A,2) / SNP Type: INTRON //

Context (SEQ ID NO:19139):

Celera SNP ID: hCV1498775 / SNP Position Genomic: 38894 / SNP Source: Celera / Population(AAllele,Count):
no_pop(G,1|A,6) total(G,1|A,6) / SNP Type: INTRON //

Context (SEQ ID NO:19140):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV1498776 / SNP Position Genomic: 39094 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,2|T,6) total(G,2|T,6) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19141):

Celera SNP ID: hCV320061 / SNP Position Genomic: 132701 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,3|C,4) total(C,4|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19142):

Celera SNP ID: hCV209461 / SNP Position Genomic: 169426 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|G,4) total(G,4|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19143):

Celera SNP ID: hCV1498778 / SNP Position Genomic: 41243 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,3|A,4) total(G,3|A,4) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19144):

Celera SNP ID: hCV209462 / SNP Position Genomic: 169696 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|G,3) total(G,3|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19145):

Celera SNP ID: hCV9620270 / SNP Position Genomic: 82758 / SNP Source: Celera;dbSNP /
Population(Allele,Count): no_pop(A,1|G,6) total(G,6|A,1) ;no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19146):

Celera SNP ID: hCV9620271 / SNP Position Genomic: 82876 / SNP Source: Celera;dbSNP /
Population(Allele,Count): no_pop(T,1|C,9) total(C,9|T,1) ;no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19147):

Celera SNP ID: hCV9620272 / SNP Position Genomic: 83471 / SNP Source: dbSNP / Population(Allele,Count):
no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19148):

Celera SNP ID: hCV11626142 / SNP Position Genomic: 171336 / SNP Source: Celera /
Population(Allele,Count): no_pop(-,2|A,7) total(A,7|-,2) / SNP Type: INTRON //

Context (SEQ ID NO:19149):

Celera SNP ID: hCV11626143 / SNP Position Genomic: 173001 / SNP Source: Celera /
Population(Allele,Count): no_pop(A,5|G,5) total(G,5|A,5) / SNP Type: INTRON //

Context (SEQ ID NO:19150):

Celera SNP ID: hCV9077759 / SNP Position Genomic: 144100 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,3|G,5) total(G,5|A,3) / SNP Type: INTRON //

Context (SEQ ID NO:19151):

Celera SNP ID: hCV11626146 / SNP Position Genomic: 176636 / SNP Source: Celera /
Population(Allele,Count): no_pop(G,2|A,5) total(G,2|A,5) / SNP Type: INTRON //

Context (SEQ ID NO:19152):

Celera SNP ID: hCV9620279 / SNP Position Genomic: 83976 / SNP Source: Celera;dbSNP /
Population(Allele,Count): no_pop(G,1|T,5) total(G,1|T,5) ;no_pop(G,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19153):

Celera SNP ID: hCV392614 / SNP Position Genomic: 150493 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,3|C,6) total(A,3|C,6) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19154):

Celera SNP ID: hCV11629050 / SNP Position Genomic: 102350 / SNP Source: Celera /
Population(Allele,Count): no_pop(A,2|T,2|-,5) total(A,2|-,5|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19155):

Celera SNP ID: hCV392615 / SNP Position Genomic: 150436 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,3|G,6) total(G,6|A,3) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19156):

Celera SNP ID: hCV392616 / SNP Position Genomic: 148562 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,4|G,7) total(G,7|A,4) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19157):

Celera SNP ID: hCV16206427 / SNP Position Genomic: 142725 / SNP Source: Celera /
Population(Allele,Count): no_pop(A,1|G,6) total(G,6|A,1) / SNP Type: INTRON/TFBS //

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:19158):

Celera SNP ID: hCV16206428 / SNP Position Genomic: 142934 / SNP Source: Celera /

Population(AAllele,Count): no_pop(C,1|G,5) total(G,5|C,1) / SNP Type: INTRON //

Context (SEQ ID NO:19159):

Celera SNP ID: hCV1498780 / SNP Position Genomic: 53262 / SNP Source: Celera / Population(AAllele,Count):

no_pop(T,2|C,6) total(C,6|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19160):

Celera SNP ID: hCV1498781 / SNP Position Genomic: 53868 / SNP Source: Celera / Population(AAllele,Count):

no_pop(G,1|A,6) total(G,1|A,6) / SNP Type: INTRON //

Context (SEQ ID NO:19161):

Celera SNP ID: hCV11629055 / SNP Position Genomic: 109773 / SNP Source: Celera /

Population(AAllele,Count): no_pop(T,2|- ,7) total(- ,7|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19162):

Celera SNP ID: hCV1498783 / SNP Position Genomic: 55578 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,1|G,4) total(G,4|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19163):

Celera SNP ID: hCV1498784 / SNP Position Genomic: 56266 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(A,7|G,7) total(G,7|A,7) ;no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19164):

Celera SNP ID: hCV11629057 / SNP Position Genomic: 109793 / SNP Source: Celera /

Population(AAllele,Count): no_pop(G,2|T,6) total(G,2|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19165):

Celera SNP ID: hCV1498785 / SNP Position Genomic: 58303 / SNP Source: Celera / Population(AAllele,Count):

no_pop(G,4|T,4) total(G,4|T,4) / SNP Type: INTRON //

Context (SEQ ID NO:19166):

Celera SNP ID: hCV11629058 / SNP Position Genomic: 128816 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(T,4|C,7) total(C,7|T,4) ;no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19167):

Celera SNP ID: hCV1498786 / SNP Position Genomic: 59580 / SNP Source: Applera / Population(AAllele,Count):

african american(C,27|G,7) caucasian(C,33|G,5) total(C,60|G,12) / SNP Type: Celera /

Population(AAllele,Count): no_pop(G,1|C,6) total(G,1|C,6) / SNP Type: INTRON / SNP Source: INTRON //

Context (SEQ ID NO:19168):

Celera SNP ID: hCV11629059 / SNP Position Genomic:131103 / SNP Source: Celera /

Population(AAllele,Count): no_pop(G,4|C,5) total(G,4|C,5) / SNP Type: INTRON //

Context (SEQ ID NO:19169):

Celera SNP ID: hCV1498787 / SNP Position Genomic: 60218 / SNP Source: Celera / Population(AAllele,Count):

no_pop(C,2|T,6) total(C,2|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19170):

Celera SNP ID: hCV1498789 / SNP Position Genomic: 61682 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,1|C,8) total(A,1|C,8) / SNP Type: INTRON //

Context (SEQ ID NO:19171):

Celera SNP ID: hCV26368318 / SNP Position Genomic: 12521 / SNP Source: Celera / Population(AAllele,Count):

no_pop(A,1|- ,8) total(A,1|- ,8) / SNP Type: INTRON //

Context (SEQ ID NO:19172):

Celera SNP ID: hCV9620280 / SNP Position Genomic: 85160 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(A,1|G,7) total(G,7|A,1) ;no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19173):

Celera SNP ID: hCV9620281 / SNP Position Genomic: 91564 / SNP Source: Applera / Population(AAllele,Count):

african american(C,35|T,1) caucasian(C,40) total(C,75|T,1) / SNP Type:SILENT MUTATION //

Context (SEQ ID NO:19174):

Celera SNP ID: hCV11626153 / SNP Position Genomic: 177287 / SNP Source: Celera /

Population(AAllele,Count): no_pop(C,1|T,6) total(C,1|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19175):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV11626154 / SNP Position Genomic: 178305 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,4|G,5) total(G,5|A,4) / SNP Type: INTRON //

Context (SEQ ID NO:19176):

Celera SNP ID: hCV11626156 / SNP Position Genomic: 178358 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,4|T,6) total(A,4|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19177):

Celera SNP ID: hCV9620288 / SNP Position Genomic: 93574 / SNP Source: Celera;dbSNP /

Population(AAllele,Count): no_pop(G,2|A,8) total(G,2|A,8) ;(A,-|G,-) Caucasian(A,24|G,56)

African-American(G,19|A,63) Asians(G,37|A,45) total(G,112|A,132) / SNP Type: INTRON //

Context (SEQ ID NO:19178):

Celera SNP ID: hCV26368752 / SNP Position Genomic: 11097 / SNP Source: Celera / Population(AAllele,Count): no_pop(G,1|C,6) total(G,1|C,6) / SNP Type: INTRON //

Context (SEQ ID NO:19179):

Celera SNP ID: hCV26368753 / SNP Position Genomic: 11368 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,1|C,8) total(C,8|T,1) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19180):

Celera SNP ID: hCV11629061 / SNP Position Genomic: 133541 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,1|G,8) total(G,8|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19181):

Celera SNP ID: hCV16206436 / SNP Position Genomic: 149138 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,1|C,8) total(C,8|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19182):

Celera SNP ID: hCV16206437 / SNP Position Genomic: 149913 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,1|G,9) total(G,9|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19183):

Celera SNP ID: hCV1498790 / SNP Position Genomic: 62154 / SNP Source: Celera / Population(AAllele,Count): no_pop(C,1|T,9) total(C,1|T,9) / SNP Type: INTRON //

Context (SEQ ID NO:19184):

Celera SNP ID: hCV11629064 / SNP Position Genomic: 133879 / SNP Source: Celera / Population(AAllele,Count): no_pop(C,1|T,5) total(C,1|T,5) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19185):

Celera SNP ID: hCV69943 / SNP Position Genomic: 155493 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,1|C,5) total(C,5|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19186):

Celera SNP ID: hCV1498791 / SNP Position Genomic: 62156 / SNP Source: Celera / Population(AAllele,Count): no_pop(C,1|T,9) total(C,1|T,9) / SNP Type: INTRON //

Context (SEQ ID NO:19187):

Celera SNP ID: hCV69944 / no_pop(C,1|G,5) total(G,5|C,1) NP Position Genomic: 155499 / SNP Source: Celera / Population(AAllele,Count): / SNP Type: INTRON //

Context (SEQ ID NO:19188):

Celera SNP ID: hCV1498792 / SNP Position Genomic: 63567 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,2|G,7) total(G,7|A,2) / SNP Type: INTRON //

Context (SEQ ID NO:19189):

Celera SNP ID: hCV1498794 / SNP Position Genomic: 64291 / SNP Source: Celera / Population(AAllele,Count): no_pop(C,3|T,4) total(C,3|T,4) / SNP Type: INTRON //

Context (SEQ ID NO:19190):

Celera SNP ID: hCV1498795 / SNP Position Genomic: 65471 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,3|C,4) total(C,4|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19191):

Celera SNP ID: hCV1498796 / SNP Position Genomic: 67304 / SNP Source: Celera / Population(AAllele,Count): no_pop(G,1|A,5) total(G,1|A,5) / SNP Type: INTRON //

Context (SEQ ID NO:19192):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV1498797 / SNP Position Genomic: 67395 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,1|-,4) total(A,1|-,4) / SNP Type: INTRON //

Context (SEQ ID NO:19193):

Celera SNP ID: hCV235862 / SNP Position Genomic: 124706 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,1|T,9) total(C,1|T,9) / SNP Type: INTRON //

Context (SEQ ID NO:19194):

Celera SNP ID: hCV235863 / SNP Position Genomic: 122013 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,4|G,4) total(G,4|A,4) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19195):

Celera SNP ID: hCV9620290 / SNP Position Genomic: 100852 / SNP Source: HGBASE;dbSNP /
Population(Allele,Count): no_pop(C,-|T,-) ;no_pop(C,-|T,-) / SNP Type: INTRON //

Context (SEQ ID NO:19196):

Celera SNP ID: hCV11626161 / SNP Position Genomic: 182699 / SNP Source: Celera /
Population(Allele,Count): no_pop(A,3|G,4) total(G,4|A,3) / SNP Type: INTRON //

Context (SEQ ID NO:19197):

Celera SNP ID: hCV11626162 / SNP Position Genomic: 183219 / SNP Source: Celera /
Population(Allele,Count): no_pop(C,3|A,5) total(A,5|C,3) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19198):

Celera SNP ID: hCV73322 / SNP Position Genomic: 172611 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,4|T,4) total(G,4|T,4) / SNP Type: INTRON //

Context (SEQ ID NO:19199):

Celera SNP ID: hCV11626163 / SNP Position Genomic: 186107 / SNP Source: Celera /
Population(Allele,Count): no_pop(C,1|T,5) total(C,1|T,5) / SNP Type: INTRON //

Context (SEQ ID NO:19200):

Celera SNP ID: hCV73323 / SNP Position Genomic: 172817 / SNP Source: Celera / Population(Allele,Count):
no_pop(G,3|A,6) total(G,3|A,6) / SNP Type: INTRON //

Context (SEQ ID NO:19201):

Celera SNP ID: hCV9080925 / SNP Position Genomic: 29421 / SNP Source: Celera / Population(Allele,Count):
no_pop(C,2|T,3) total(C,2|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19202):

Celera SNP ID: hCV15991281 / SNP Position Genomic: 53476 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,2|C,9) total(C,9|T,2) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19203):

Celera SNP ID: hCV9620297 / SNP Position Genomic: 100884 / SNP Source: HGBASE /
Population(Allele,Count): no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19204):

Celera SNP ID: hCV73325 / SNP Position Genomic: 173500 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,4|G,5) total(G,5|A,4) / SNP Type: INTRON //

Context (SEQ ID NO:19205):

Celera SNP ID: hCV11626167 / SNP Position Genomic: 196931 / SNP Source: Celera /
Population(Allele,Count): no_pop(-,3|A,7) total(A,7|-,3) / SNP Type: INTRON //

Context (SEQ ID NO:19206):

Celera SNP ID: hCV15991282 / SNP Position Genomic: 75844 / SNP Source: Celera / Population(Allele,Count):
no_pop(T,1|G,6) total(G,6|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19207):

Celera SNP ID: hCV9620298 / SNP Position Genomic: 118860 / SNP Source: dbSNP /
Population(Allele,Count): (A,-|G,-) Caucasian(G,7|G,10|A,56|A,77) African-American(G,7|A,73|A,84)
Asians(G,5|A,79|A,84) total(G,29|A,453) / SNP Type: INTRON //

Context (SEQ ID NO:19208):

Celera SNP ID: hCV73326 / SNP Position Genomic: 174464 / SNP Source: Celera / Population(Allele,Count):
no_pop(A,2|G,3) total(G,3|A,2) / SNP Type: INTRON //

Context (SEQ ID NO:19209):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV9620299 / SNP Position Genomic: 119136 / SNP Source: dbSNP /
Population(AAllele,Count): no_pop(A,-|G,-) / SNP Type: INTRON //

Context (SEQ ID NO:19210):

Celera SNP ID: hCV73327 / SNP Position Genomic: 175758 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,1|T,6) total(C,1|T,6) / SNP Type: INTRON //

Context (SEQ ID NO:19211):

Celera SNP ID: hCV73329 / SNP Position Genomic: 176171 / SNP Source: Celera / Population(AAllele,Count):
no_pop(A,1|G,6) total(G,6|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19212):

Celera SNP ID: hCV26368400 / SNP Position Genomic: 155369 / SNP Source: Celera /
Population(AAllele,Count): no_pop(T,1|-|8) total(-,8|T,1) / SNP Type: INTRON //

Context (SEQ ID NO:19213):

Celera SNP ID: hCV26368401 / SNP Position Genomic: 157290 / SNP Source: Celera /
Population(AAllele,Count): no_pop(A,1|G,5) total(G,5|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19214):

Celera SNP ID: hCV26368402 / SNP Position Genomic: 158909 / SNP Source: Celera /
Population(AAllele,Count): no_pop(C,1|T,5) total(C,1|T,5) / SNP Type: INTRON //

Context (SEQ ID NO:19215):

Celera SNP ID: hCV26368403 / SNP Position Genomic: 160761 / SNP Source: Celera /
Population(AAllele,Count): no_pop(G,2|C,3) total(G,2|C,3) / SNP Type: INTRON //

Context (SEQ ID NO:19216):

Celera SNP ID: hCV26368331 / SNP Position Genomic: 16355 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,1|A,4) total(A,4|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19217):

Celera SNP ID: hCV26368404 / SNP Position Genomic: 161570 / SNP Source: Celera /
Population(AAllele,Count): no_pop(G,1|C,9) total(G,1|C,9) / SNP Type: INTRON //

Context (SEQ ID NO:19218):

Celera SNP ID: hCV26368332 / SNP Position Genomic: 16356 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,1|A,4) total(A,4|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19219):

Celera SNP ID: hCV26368406 / SNP Position Genomic: 168383 / SNP Source: Celera /
Population(AAllele,Count): no_pop(T,3|A,6) total(A,6|T,3) / SNP Type: INTRON //

Context (SEQ ID NO:19220):

Celera SNP ID: hCV26368333 / SNP Position Genomic: 16357 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,1|T,4) total(T,4|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19221):

Celera SNP ID: hCV26368407 / SNP Position Genomic: 168384 / SNP Source: Celera /
Population(AAllele,Count): no_pop(-,3|A,6) total(A,6|-,3) / SNP Type: INTRON //

Context (SEQ ID NO:19222):

Celera SNP ID: hCV26368334 / SNP Position Genomic: 16360 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,1|T,4) total(T,4|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19223):

Celera SNP ID: hCV26368408 / SNP Position Genomic: 176079 / SNP Source: Celera /
Population(AAllele,Count): no_pop(A,1|G,6) total(G,6|A,1) / SNP Type: INTRON //

Context (SEQ ID NO:19224):

Celera SNP ID: hCV26368409 / SNP Position Genomic: 177104 / SNP Source: Celera /
Population(AAllele,Count): no_pop(-,3|A,4) total(A,4|-,3) / SNP Type: INTRON //

Context (SEQ ID NO:19225):

Celera SNP ID: hCV73331 / SNP Position Genomic: 176448 / SNP Source: Celera / Population(AAllele,Count):
no_pop(T,2|C,6) total(C,6|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19226):

Celera SNP ID: hCV9620376 / SNP Position Genomic: 198294 / SNP Source: HGBASE;dbSNP /
Population(AAllele,Count): no_pop(A,-|G,-) ;no_pop(A,-|G,-) / SNP Type: SILENT MUTATION //

EP 2 474 631 B1

(continued)

Context (SEQ ID NO:19227):

Celera SNP ID: hCV9620377 / SNP Position Genomic: 198738 / SNP Source: HGBASE /

Population(AAllele,Count): no_pop(A,-|G,-) / SNP Type: UTR 3 //

Context (SEQ ID NO:19228):

Celera SNP ID: hCV11626176 / SNP Position Genomic: 204870 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,2|C,4) total(C,4|-,2) / SNP Type: TRANSCRIPTION FACTOR BINDING SITE (TFBS) //

Context (SEQ ID NO:19229):

Celera SNP ID: hCV15991293 / SNP Position Genomic: 92967 / SNP Source: Applera /

Population(AAllele,Count): african american(A,2|T,26) caucasian(A,13|T,17) total(A,15|T,43) / SNP Type: INTRON/hmCS / SNP Source: Celera / Population(AAllele,Count): no_pop(A,2|T,6) total(A,2|T,6) / SNP Type: INTRON/hmCS //

Context (SEQ ID NO:19230):

Celera SNP ID: hCV16093143 / SNP Position Genomic: 11860 / SNP Source: Celera / Population(AAllele,Count): no_pop(G,1|A,7) total(G,1|A,7) / SNP Type: INTRON //

Context (SEQ ID NO:19231):

Celera SNP ID: hCV26368411 / SNP Position Genomic: 182603 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|T,7) total(T,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19232):

Celera SNP ID: hCV16093144 / SNP Position Genomic: 65348 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,1|G,5) total(G,5|A,1) / SNP Type: INTRON/TFBS //

Context (SEQ ID NO:19233):

Celera SNP ID: hCV26368412 / SNP Position Genomic: 182604 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|T,7) total(T,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19234):

Celera SNP ID: hCV26368413 / SNP Position Genomic: 182605 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|T,7) total(T,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19235):

Celera SNP ID: hCV26368414 / SNP Position Genomic: 182610 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|G,7) total(G,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19236):

Celera SNP ID: hCV26368341 / SNP Position Genomic: 29002 / SNP Source: Celera / Population(AAllele,Count): no_pop(T,2|-,6) total(-,6|T,2) / SNP Type: INTRON //

Context (SEQ ID NO:19237):

Celera SNP ID: hCV26368415 / SNP Position Genomic: 187038 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,2|A,5) total(A,5|-,2) / SNP Type: INTRON //

Context (SEQ ID NO:19238):

Celera SNP ID: hCV26368342 / SNP Position Genomic: 30831 / SNP Source: Celera / Population(AAllele,Count): no_pop(-,1|T,7) total(T,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19239):

Celera SNP ID: hCV16078775 / SNP Position Genomic: 85703 / SNP Source: Celera / Population(AAllele,Count): no_pop(A,1|T,8) total(A,1|T,8) / SNP Type: INTRON //

Context (SEQ ID NO:19240):

Celera SNP ID: hCV26368416 / SNP Position Genomic: 187594 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|T,6) total(T,6|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19241):

Celera SNP ID: hCV26368343 / SNP Position Genomic: 30832 / SNP Source: Celera / Population(AAllele,Count): no_pop(-,1|C,7) total(C,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19242):

Celera SNP ID: hCV26368417 / SNP Position Genomic: 187595 / SNP Source: Celera /

Population(AAllele,Count): no_pop(-,1|T,6) total(T,6|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19243):

EP 2 474 631 B1

(continued)

Celera SNP ID: hCV26368344 / SNP Position Genomic: 30833 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,1|T,7) total(T,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19244):

Celera SNP ID: hCV26368418 / SNP Position Genomic: 187596 / SNP Source: Celera /
Population(AAllele,Count): no_pop(-,1|T,6) total(T,6|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19245):

Celera SNP ID: hCV26368345 / SNP Position Genomic: 30834 / SNP Source: Celera / Population(AAllele,Count):
no_pop(-,1|C,7) total(C,7|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19246):

Celera SNP ID: hCV26368419 / SNP Position Genomic: 187597 / SNP Source: Celera /
Population(AAllele,Count): no_pop(-,1|T,6) total(T,6|-,1) / SNP Type: INTRON //

Context (SEQ ID NO:19247):

Celera SNP ID: hCV26368349 / SNP Position Genomic: 37614 / SNP Source: Celera / Population(AAllele,Count):
no_pop(C,1|G,10) total(G,10|C,1) / SNP Type: INTRON //

TABLE 5, page 1 of 6

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV1022614	G/A	CTGCAGCCTCTCCTACG (SEQ ID NO: 73086)	CCTGCAGCCTCTCCTACA (SEQ ID NO: 73087)	GATCCCCCATCGGTCAATA (SEQ ID NO: 73088)
hCV1026583	A/G	GGAAGTTGAGATCTCTTCAGAA (SEQ ID NO: 73089)	GGAAGTTGAGATCTCTTCAGAG (SEQ ID NO: 73090)	AGTGATTGGAATCCATATTTACTT (SEQ ID NO: 73091)
hCV1065191	G/T	GCCCCACTTTTGCATG (SEQ ID NO: 73092)	AGCCCCACTTTTGCATT (SEQ ID NO: 73093)	ACCCCTGCACAGTTTAGAAC (SEQ ID NO: 73094)
hCV1085595	T/C	GGTGCTCCACCTGGT (SEQ ID NO: 73095)	GTGCTCCACCTGGC (SEQ ID NO: 73096)	GGAGTTCGAACCTAAAGACGTAT (SEQ ID NO: 73097)
hCV1085600	C/G	AGCTGTTCTGTTCTATGATC (SEQ ID NO: 73098)	AGCTGTTCTGTTCTATGATG (SEQ ID NO: 73099)	GAAGTCAACAGTGAACATGTGA (SEQ ID NO: 73100)
hCV11159941	G/A	CCCGTTGGTTCCGAAAG (SEQ ID NO: 73101)	CCCGTTGGTTCCGAAAA (SEQ ID NO: 73102)	TGAATAGCCATTAGAAAAAACTGT (SEQ ID NO: 73103)
hCV1129436	C/A	TTCCAGGGTATATCTCAGAGC (SEQ ID NO: 73104)	GTTCCAGGGTATATCTCAGAGA (SEQ ID NO: 73105)	TTGAAAGAGTGTGAGCAAGATC (SEQ ID NO: 73106)
hCV11359098	G/C	CAAAATGTAGAAGTTTCATATGAG (SEQ ID NO: 73107)	CAAAATGTAGAAGTTTCATATGAC (SEQ ID NO: 73108)	GAGCTGTGTGTTTCTTCTTTGTTCTA (SEQ ID NO: 73109)
hCV11442703	C/T	CCGAGGGCTCCTGAC (SEQ ID NO: 73110)	CCGAGGGCTCCTGAT (SEQ ID NO: 73111)	AAAAAGCCCTTTGGTATTGTATA (SEQ ID NO: 73112)
hCV11482579	T/A	GTTGAAGGGAAGTTTCAGCAT (SEQ ID NO: 73113)	TTGAAGGGAAGTTTCAGCAA (SEQ ID NO: 73114)	TCACGGAGGACAGGTAGAAT (SEQ ID NO: 73115)
hCV11482766	G/A	GCGCACCCAGGTCAG (SEQ ID NO: 73116)	GCGCACCCAGGTCAA (SEQ ID NO: 73117)	CCACGTTCTGGTCGATCTT (SEQ ID NO: 73118)
hCV11482773	A/C	CTGCTGCTGCTCCTGA (SEQ ID NO: 73119)	TGCTGCTGCTCCTGC (SEQ ID NO: 73120)	ACTTGAGCTTCCCTGGAGAAG (SEQ ID NO: 73121)
hCV11484594	C/T	CCACAGCGAGGCTTTTC (SEQ ID NO: 73122)	CCACAGCGAGGCTTTTTT (SEQ ID NO: 73123)	GCACTAATGTGATCGTTGAAAA (SEQ ID NO: 73124)
hCV11486078	C/G	AGCCCCAGAACCTGC (SEQ ID NO: 73125)	AGCCCCAGAACCTGG (SEQ ID NO: 73126)	GGGCTGGGCTTGTAGAATA (SEQ ID NO: 73127)
hCV11506744	T/G	GAAGAAGGAGGATGAAGATGTCT (SEQ ID NO: 73128)	AAAGGAGGATGAAGATGTGCG (SEQ ID NO: 73129)	CACCATGCTCTGCAAAAGAC (SEQ ID NO: 73130)
hCV11592758	T/C	CATCCAACAGCTCTTCTATCAT (SEQ ID NO: 73131)	CATCCAACAGCTCTTCTATCAC (SEQ ID NO: 73132)	CAAAACATCCGAGGACAAG (SEQ ID NO: 73133)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV11628130	A/T	CTGCCCTCTTTTAGCAGA (SEQ ID NO: 73134)	CTGCCCTCTTTTAGCAGT (SEQ ID NO: 73135)	CCCTTTCTCATTCATTCATTTT (SEQ ID NO: 73136)
hCV11655948	A/G	GACGTCTTCCAGTACCA (SEQ ID NO: 73137)	ACGTCTTCCAGTACCG (SEQ ID NO: 73138)	TCTTCCTCGCTCAGAAT (SEQ ID NO: 73139)
hCV11655953	T/C	AGCATTGCCGTCCT (SEQ ID NO: 73140)	GCATTGCCGTCCT (SEQ ID NO: 73141)	CCATGGGTCAAAGAACA (SEQ ID NO: 73142)
hCV11668930	T/C	CCAGATCCAGTTTCTAGCAT (SEQ ID NO: 73143)	CCAGATCCAGTTTCTAGCAC (SEQ ID NO: 73144)	CCCTTCTCCAGCATTATC (SEQ ID NO: 73145)
hCV11689917	T/C	CTGTGAGTGGCCCTTCAT (SEQ ID NO: 73146)	CTGTGAGTGGCCCTTCAC (SEQ ID NO: 73147)	GGAGCCCCGCTTCAT (SEQ ID NO: 73148)
hCV11689926	C/T	GAAGGCCCACTCACAGAC (SEQ ID NO: 73149)	GAAGGCCCACTCACAGAT (SEQ ID NO: 73150)	CGGACCCGGAGACTG (SEQ ID NO: 73151)
hCV11689930	A/G	TCACAGACTGACCGAGTGA (SEQ ID NO: 73152)	CACAGACTGACCGAGTGG (SEQ ID NO: 73153)	CGGACCCGGAGACTGT (SEQ ID NO: 73154)
hCV11696920	A/G	GTCCTTAGAAGCCTCTTCAGAATA (SEQ ID NO: 73155)	CTTTAGAAGCCTCTTCAGAATG (SEQ ID NO: 73156)	CGGCTTTGGCCTACAAAG (SEQ ID NO: 73157)
hCV11758801	C/G	AGTACCTCTTGGTCTCTCTCC (SEQ ID NO: 73158)	AGTACCTCTTGGTCTCTCTCG (SEQ ID NO: 73159)	GCATGTTGTGTTTCTGATTGTAC GATTGTAC (SEQ ID NO: 73160)
hCV11764545	G/T	ACGAAGCTTCCGAGGAAG (SEQ ID NO: 73161)	CGAAGCTTCCGAGGAAT (SEQ ID NO: 73162)	GACACCGGACGAGAGAGAC (SEQ ID NO: 73163)
hCV11789692	G/C	ATCCCCCACAGATCCAG (SEQ ID NO: 73164)	ATCCCCCACAGATCCAC (SEQ ID NO: 73165)	CCAGCTGGACCCAGTAAG (SEQ ID NO: 73166)
hCV11889257	T/C	CTCTCTTTCTAGAACTGAAGAAATT (SEQ ID NO: 73167)	TCTCTTTCTAGAACTGAAGAAATC (SEQ ID NO: 73168)	GGGCAGGGCTAGGAGTAG (SEQ ID NO: 73169)
h6V11937023	C/T	GGATATGAGTTGGACATGAAGAC (SEQ ID NO: 73170)	GGATATGAGTTGGACATGAAGAT (SEQ ID NO: 73171)	CAGGTTTGGTGGGAGAAC (SEQ ID NO: 73172)
hCV11951095	T/C	CGTGACCCCTGCCGT (SEQ ID NO: 73173)	CGTGACCCCTGCCGC (SEQ ID NO: 73174)	GGGCCAGCATGTGGAC (SEQ ID NO: 73175)
hCV11955747	T/G	AAAGGGAAGGAGGTTACTTACT (SEQ ID NO: 73176)	AGGGAAGGAGGTTACTTACG (SEQ ID NO: 73177)	TCCTCTGTGGAGAGGGATAC (SEQ ID NO: 73178)
hCV11972321	G/A	GGTTCTGACATGACTGTGACAG (SEQ ID NO: 73179)	GTTCTGACATGACTGTGACAA (SEQ ID NO: 73180)	CCCCAACCCCAAGGTTTAC (SEQ ID NO: 73181)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV11975283	A/C	GGGCTGGACATTGCAA (SEQ ID NO: 73182)	GGGCTGGACATTGCAC (SEQ ID NO: 73183)	CCAGATTAGCCTTAACCTCTGTTAAC (SEQ ID NO: 73184)
hCV1202883	G/A	GCGTGATGATGAAATCGG (SEQ ID NO: 73185)	GCGTGATGATGAAATCGA (SEQ ID NO: 73186)	AGCCTCTCCTGACTGTCATC (SEQ ID NO: 73187)
hCV1202981	A/G	ACGAGAGCATCATCTGCA (SEQ ID NO: 73188)	CGAGAGCATCATCTGCG (SEQ ID NO: 73189)	CCAGACATTGCAGTTGAAGTC (SEQ ID NO: 73190)
hCV1207994	T/G	GCAGCAGTCGCCCTT (SEQ ID NO: 73191)	GCAGCAGTCGCCCTG (SEQ ID NO: 73192)	CATTTTGTGATGTTTGTCTTAG (SEQ ID NO: 73193)
hCV12083298	A/G	CAGAACGCTGGGAAACA (SEQ ID NO: 73194)	AGAACGCTGGGAAACG (SEQ ID NO: 73195)	CTCCCCACCTGTTGCTC (SEQ ID NO: 73196)
hCV12114319	T/C	GACACTGCCCTCATCGT (SEQ ID NO: 73197)	CACTGCCCTCATCGC (SEQ ID NO: 73198)	CCTGTCCCTTGAGGCTCTGATC (SEQ ID NO: 73199)
hCV1212713	T/C	CCCAGTGGGTCTT (SEQ ID NO: 73200)	CCCAGTGGGTCCC (SEQ ID NO: 73201)	TGCCATCGTTGTTTG (SEQ ID NO: 73202)
hCV1260328	T/C	TCCACGTGGACCAGGT (SEQ ID NO: 73203)	CCACGTGGACCAGGC (SEQ ID NO: 73204)	GCCCAGGTATTTCATCAGC (SEQ ID NO: 73205)
hCV1276216	T/C	TCTTATCAGTCTTGGACAAGAACT (SEQ ID NO: 73206)	TTATCAGTCTTGGACAAGAAC (SEQ ID NO: 73207)	CTCTTGCTTTCTGTCTTCATAGAC (SEQ ID NO: 73208)
hCV1345898	C/T	CAGTTTCCATGGGTCTACTAC (SEQ ID NO: 73209)	CAGTTTCCATGGGTCTACTAT (SEQ ID NO: 73210)	TTATGAAATGGTACAGACAAGTGAT (SEQ ID NO: 73211)
hCV1376137	A/G	CTCCATCATTTGCAGACCA (SEQ ID NO: 73212)	TCCATCATTTGCAGACCG (SEQ ID NO: 73213)	CCAATTCCCCTGATGTTAAA (SEQ ID NO: 73214)
hCV1376266	T/A	CGCCCCCTCCCGCT (SEQ ID NO: 73215)	CGCCCCCTCCCGCA (SEQ ID NO: 73216)	TGATCCCTCCCTTGGATA (SEQ ID NO: 73217)
hCV1376342	C/T	ACTCACCAGTTGTGAAGACTTC (SEQ ID NO: 73218)	TACTCACCAGTTGTGAAGACTTT (SEQ ID NO: 73219)	TCAGGGAGCCTAGATATCTCA (SEQ ID NO: 73220)
hCV1385736	T/C	TGGGTGTCAGGTCCCTT (SEQ ID NO: 73221)	TGGGTGTCAGGTCCCTC (SEQ ID NO: 73222)	GCAGCCACCTTGACACTC (SEQ ID NO: 73223)
hCV1387523	T/C	TGGGGTTGGGTTCT (SEQ ID NO: 73224)	GGGGTTGGGTTCC (SEQ ID NO: 73225)	TGTCCCTGTCTCCTTCAG (SEQ ID NO: 73226)
hCV1403468	T/G	ACTGGCCCCCTTGCAT (SEQ ID NO: 73227)	ACTGGCCCCCTTGCAG (SEQ ID NO: 73228)	AGGAGGGAACCAACCTTA (SEQ ID NO: 73229)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV1419869	G/C	AGTTGTCAGTGCCTCTGTG (SEQ ID NO: 73230)	AGTTGTCAGTGCCTCTGTTC (SEQ ID NO: 73231)	ACAACCCCTTGAATGAGAGAATAC (SEQ ID NO: 73232)
hCV1466546	A/G	TCTGGCTTCCGGAA (SEQ ID NO: 73233)	TCTGGCTTCCGGAG (SEQ ID NO: 73234)	CGTAGCTGTTGACCATCATTTA (SEQ ID NO: 73235)
hCV1486426	A/G	CACAGGCTATGACCTCAACA (SEQ ID NO: 73236)	ACAGGCTATGACCTCAACG (SEQ ID NO: 73237)	AACCATGGAGCTACTCTTCTGTGTA (SEQ ID NO: 73238)
hCV14938	T/C	AGGTGGAGATTCTCAACAGAT (SEQ ID NO: 73239)	AGGTGGAGATTCTCAACAGAC (SEQ ID NO: 73240)	AGGCTGTTCTCATGACATACAT (SEQ ID NO: 73241)
hCV1552900	A/G	GGGATGGAAGAGCTTCA (SEQ ID NO: 73242)	GGGATGGAAGAGCTTCG (SEQ ID NO: 73243)	CTGCAGCCTTCTCTCTGAC (SEQ ID NO: 73244)
hCV15751934	T/C	ACTATGAGAGCATCATGTGT (SEQ ID NO: 73245)	CTATGAGAGCATCATGTGC (SEQ ID NO: 73246)	ACTGAGTCTCTGAAGAAAAATCAG (SEQ ID NO: 73247)
hCV15760070	A/T	TGTCCAGATCCACATAGAACA (SEQ ID NO: 73248)	TTGTCCAGATCCACATAGAACT (SEQ ID NO: 73249)	CTTTATGCAGCGGACCAC (SEQ ID NO: 73250)
hCV15851292	C/T	TCCCCCGAGAAGAAGAC (SEQ ID NO: 73251)	TCCCCCGAGAAGAAGAT (SEQ ID NO: 73252)	TGCTGTCAACCCTCTCTCTT (SEQ ID NO: 73253)
hCV15653800	T/G	ACCTGGTGGCACATCTAATAT (SEQ ID NO: 73254)	CCTGGTGGCACATCTAATAG (SEQ ID NO: 73255)	TGTGGAATGTGGAACAATACT (SEQ ID NO: 73256)
hCV15859395	G/T	GCCTGCATTACTCAAGGG (SEQ ID NO: 73257)	TGCCTGCATTACTCAAGGT (SEQ ID NO: 73258)	CAGAGAGGGTCAGGAGTTGA (SEQ ID NO: 73259)
hCV15859649	G/A	CTGCCATGATGTTACAG (SEQ ID NO: 73260)	CCTGCCATGATGTTTACA (SEQ ID NO: 73261)	GATGGCTGCATAGCAGTAGAT (SEQ ID NO: 73262)
hCV15869253	G/T	CTACTGCCCCGACCTG (SEQ ID NO: 73263)	CTACTGCCCCGACCTT (SEQ ID NO: 73264)	GTCATGGCGCTACTAGATGTAT (SEQ ID NO: 73265)
hCV15878378	T/C	TGCCTGGTGAGGGAGAT (SEQ ID NO: 73266)	TGCCTGGTGAGGGAGAC (SEQ ID NO: 73267)	CACACACAAGAAAGTAAACATGAA (SEQ ID NO: 73268)
hCV15966517	A/G	CACCTTCCAGCCGGTA (SEQ ID NO: 73269)	ACCTTCCAGCCGGTG (SEQ ID NO: 73270)	TGAGTAGCCCCAGTGACTTC (SEQ ID NO: 73271)
hCV15968121	T/C	TCAAAATTCATGGTCACTTTAAT (SEQ ID NO: 73272)	TCAAAATTCATGGTCACTTTAAT (SEQ ID NO: 73273)	CCTCTTCCATGCACTCACTT (SEQ ID NO: 73274)
hCV5976768	C/G	TGGATGTTGGAACAATTGAC (SEQ ID NO: 73275)	TGGATGTTGGAACAATTGAG (SEQ ID NO: 73276)	CCAGCCAGCAAAAGATACATAC (SEQ ID NO: 73277)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV15976971	T/C	GAGCCCTGCGCACT (SEQ ID NO: 73278)	AGCCCTGCGCACC (SEQ ID NO: 73279)	GGCAGGATTGCTCTTAGTGT (SEQ ID NO: 73280)
hCV16033535	G/T	GGTCGTCCTGCACCTTCG (SEQ ID NO: 73281)	TGGTCGTCCTGCACCTTCT (SEQ ID NO: 73282)	CCAGCTCCCTCTCTTTTC (SEQ ID NO: 73283)
hCV16072719	A/G	TCACCTGCTTTTGCCA (SEQ ID NO: 73284)	CACCTGCTTTTGCCG (SEQ ID NO: 73285)	GGCCCAACCCAGTGAC (SEQ ID NO: 73286)
hCV1608777	G/A	CAAGAGGATTTTATGGAATGG (SEQ ID NO: 73287)	ACAAGAGGATTTTATGGAATGA (SEQ ID NO: 73288)	GGTGGTGAGGCTTTGAGTATA (SEQ ID NO: 73289)
hCV16089120	T/C	CCCACATTCTCTCTCCTT (SEQ ID NO: 73290)	CCCACATTCTCTCTCCTC (SEQ ID NO: 73291)	TGAGGAGCGTGGGTGAC (SEQ ID NO: 73292)
hCV16165996	C/T	CTGAGGCCCTATGTCCTC (SEQ ID NO: 73293)	CTGAGGCCCTATGTCCTT (SEQ ID NO: 73294)	AGCTCTCCTTTGTTGCTACTG (SEQ ID NO: 73295)
hCV16166043	T/C	CGGTTGAAGTCCTTGAT (SEQ ID NO: 73296)	CGGTTGAAGTCCTTGAC (SEQ ID NO: 73297)	GGTTGTGCAGAGACATCTGA (SEQ ID NO: 73298)
hCV16110641	G/C	TCTGCTTAAATATGCTTGTG (SEQ ID NO: 73299)	TTCTGCTTAAATATGCTTGTG (SEQ ID NO: 73300)	GCAGTTGTTAAGGACAGAAAATACTT (SEQ ID NO: 73301)
hCV16170651	C/T	AACAAACGTACCATTTGAGGC (SEQ ID NO: 73302)	GAAACAAACGTACCATTTGAGGT (SEQ ID NO: 73303)	TCCACGCTGATCTGAAGATAC (SEQ ID NO: 73304)
hCV16170900	T/C	CGCACACCAAGGTTCTCAT (SEQ ID NO: 73305)	CGCACACCAAGGTTCTCAC (SEQ ID NO: 73306)	GCAACTACCTGGGCCACTATA (SEQ ID NO: 73307)
hCV16170982	C/G	CCCCCACTCTCCAGC (SEQ ID NO: 73308)	CCCCCACTCTCCAGG (SEQ ID NO: 73309)	GGCAAAAAGCACTGTGAAGA (SEQ ID NO: 73310)
hCV16171764	G/A	CCCCAACCAACACG (SEQ ID NO: 73311)	GCCCCAACCAACCACA (SEQ ID NO: 73312)	GCGTTCGGGGAAGACTT (SEQ ID NO: 73313)
hCV16172098	C/T	GACACACAGGGTGGCTC (SEQ ID NO: 73314)	GACACACAGGGTGGCTT (SEQ ID NO: 73315)	TCTGGTCTGCCCTCAGGTAAC (SEQ ID NO: 73316)
hCV16172249	G/C	ACTGTAATTTTTTAAAGGTCCCTG (SEQ ID NO: 73317)	ACTGTAATTTTTTAAAGGTCCCTC (SEQ ID NO: 73318)	GGATGTATATCATCTATCTTTCACAGTATAT (SEQ ID NO: 73319)
hCV16172571	T/C	GGTACCATGGACTGTACTCACT (SEQ ID NO: 73320)	GTACCATGGACTGTACTCACC (SEQ ID NO: 73321)	AGGTTGGTTCTGGAGATGAC (SEQ ID NO: 73322)
hCV16181123	C/A	AATTCAAGACAGTGCAACATC (SEQ ID NO: 73323)	TAAATTCAAGACAGTGCAACATA (SEQ ID NO: 73324)	CACACACCGCACTCTAATTACT (SEQ ID NO: 73325)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV16191372	C/T	AGTGGTGTGGCCGC (SEQ ID NO: 73326)	GAGGTCGTGTGGCCGT (SEQ ID NO: 73327)	GCCTCTCACCACCTTTCTGTAAG (SEQ ID NO: 73328)
hCV16192174	G/A	GAGCACCTTAACATATAGATGGTG (SEQ ID NO: 73329)	TGAGCACCTTAACATATAGATGGTA (SEQ ID NO: 73330)	CTTGTCAGGCACAGAAATAATT (SEQ ID NO: 73331)
hCV16196014	G/C	TGAAGAAGCTAAGGATTGAGG (SEQ ID NO: 73332)	TGAAGAAGCTAAGGATTGAGC (SEQ ID NO: 73333)	CTCTCCCTGGCTGAGTTG (SEQ ID NO: 73334)
hCV16266313	T/C	CACGGCGCACCTTTCTT (SEQ ID NO: 73335)	CACGGCGCACCTTTCTC (SEQ ID NO: 73336)	TGTTTTTTCCTTTGTGTCATCTTATCTA (SEQ ID NO: 73337)
hCV16273460	G/A	GCACTCTTGGACAAGCG (SEQ ID NO: 73338)	TGCACTCTTGGACAAGCA (SEQ ID NO: 73339)	AATGACATCCCCCTATCTTTCTG (SEQ ID NO: 73340)
hCV16276495	C/T	GTACCTTCACCCATGGAAC (SEQ ID NO: 73341)	GTACCTTCACCCATGGAAT (SEQ ID NO: 73342)	TCACTTTCTGTTGATTACATGAGA (SEQ ID NO: 73343)
hCV16399938	T/G	AGTGGAGCTTCAGGGCT (SEQ ID NO: 73344)	TGGAGCTTCAGGGCG (SEQ ID NO: 73345)	CAGTGGAGACAGAGGATGTTTAC (SEQ ID NO: 73346)
hCV1662671	A/G	CAGCCAAGAGCAGGACA (SEQ ID NO: 73347)	AGCCAAGAGCAGGACG (SEQ ID NO: 73348)	CCCAAGACACGTTTCAGAAAT (SEQ ID NO: 73349)
hCV1841898	T/C	TCTCCCTCCTGTTCCCTGT (SEQ ID NO: 73350)	TCCCTCCTGTTCCCTTGC (SEQ ID NO: 73351)	GCAATGTGCTGTGAATGAAG (SEQ ID NO: 73352)
hCV1842400	T/C	TTGGTACCTGGCTCTCT (SEQ ID NO: 73353)	TGGTACCTGGCTCTCC (SEQ ID NO: 73354)	AAACTTCTTAGGACAGAGTGATTAGA (SEQ ID NO: 73355)
hCV1923359	G/C	CCTCAGGCACCGAGAG (SEQ ID NO: 73356)	CCTCAGGCACCGAGAC (SEQ ID NO: 73357)	CTGGACCTCCTGATGATCTC (SEQ ID NO: 73358)
hCV1985481	T/C	CTGGCTGCTCCCTGAT (SEQ ID NO: 73359)	CTGGCTGCTCCCTGAC (SEQ ID NO: 73360)	GCCTGACTGGCTGATCTC (SEQ ID NO: 73361)
hCV1998030	T/C	GGGTGCAGAACCT (SEQ ID NO: 73362)	GGGTGCAGAACCC (SEQ ID NO: 73363)	CAGACCGGCCACTTG (SEQ ID NO: 73364)
hCV2033404	T/C	CCATCTTCTACGCACTGT (SEQ ID NO: 73365)	ATCTTCTACGCACTGC (SEQ ID NO: 73366)	CAGCATAGCCAGGAAGAAGA (SEQ ID NO: 73367)
hCV2033405	G/A	GAGCGGGAGCAACG (SEQ ID NO: 73368)	TGAGCGGGAGCAACA (SEQ ID NO: 73369)	TGGAGCTGCAGGTGATC (SEQ ID NO: 73370)
hCV2038	G/A	CACGGCGGTCATGTG (SEQ ID NO: 73371)	CCACGGCGGTCATGTA (SEQ ID NO: 73372)	GTTTTGTGGAGGAAAGAG (SEQ ID NO: 73373)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV2045908	C/A	CAGTTGTACCCCTACAATAATGTAC (SEQ ID NO: 73374)	CAGTTGTACCCCTACAATAATGTAA (SEQ ID NO: 73375)	TCGGTTTCTCTTCATTCAATTC (SEQ ID NO: 73376)
hCV210485	C/G	GGCGTTCAGTTTTTAGCC (SEQ ID NO: 73377)	GGCGTTCAGTTTTTAGCG (SEQ ID NO: 73378)	CAATACGCCAGCAAAATACC (SEQ ID NO: 73379)
hCV2143205	T/C	TGTCGAATGGGAGTCTTCTT (SEQ ID NO: 73380)	TGTCGAATGGGAGTCTTCTC (SEQ ID NO: 73381)	GGAAGAAACAGCTACCCAGA (SEQ ID NO: 73382)
hCV2146578	C/T	CCTCCTAGAGAAAGATCTGCAC (SEQ ID NO: 73383)	CCTCCTAGAGAAAGATCTGCAT (SEQ ID NO: 73384)	AAGGCCCTCTTCTGTCT (SEQ ID NO: 73385)
hCV2188895	A/G	AGAGAAATGTTACCTCTCCTGA (SEQ ID NO: 73386)	GAGAAATGTTACCTCTCCTGG (SEQ ID NO: 73387)	TTCTCCTGGGTGAGATTCTC (SEQ ID NO: 73388)
hCV2200985	G/C	GCGCACCAGCTTCAG (SEQ ID NO: 73389)	GCGCACCAGCTTCAC (SEQ ID NO: 73390)	TGTAATACATGATTTTCAGACACAC (SEQ ID NO: 73391)
hCV22271841	C/T	CATCACGGAGATCCACC (SEQ ID NO: 73392)	ATCATCACGGAGATCCACT (SEQ ID NO: 73393)	TCAGTCCCAAGGAGATTCTTAG (SEQ ID NO: 73394)
hCV22271999	G/A	GAGCACCTTAACCTATAGATGGTG (SEQ ID NO: 73395)	TGAGCACCTTAACCTATAGATGGTA (SEQ ID NO: 73396)	CTTGTCAAGGCACAGAAATAATT (SEQ ID NO: 73397)
hCV22272267	T/C	CTGGCAGCGGAATGTTAT (SEQ ID NO: 73398)	CTGGCAGCGGAATGTTAC (SEQ ID NO: 73399)	CCTCTAGAAAGAAAATGGACTGTAT (SEQ ID NO: 73400)
hCV22273419	C/T	ACTCCTTTGGACTGGC (SEQ ID NO: 73401)	TGACTCCTTTGGACTGGT (SEQ ID NO: 73402)	TCTTAAATGCTGTGGAATTGTG (SEQ ID NO: 73403)
hCV22274307	C/T	GACCTACGCTCCTTCATCTAAC (SEQ ID NO: 73404)	GACCTACGCTCCTTCATCTAAT (SEQ ID NO: 73405)	CAATGTGATTACCTCAAAATAATATCTAC (SEQ ID NO: 73406)
hCV22274425	C/T	AGAGAGGGTGAAGGTGC (SEQ ID NO: 73407)	GAGAGAGGGTGAAGGTGT (SEQ ID NO: 73408)	CCCTTTAAGGCTGTTATCTGAC (SEQ ID NO: 73409)
hCV22274540	C/T	ACCACTTCTATCCGAGTAGC (SEQ ID NO: 73410)	GACCACTTCTATCCGAGTAGT (SEQ ID NO: 73411)	GGGAAAAATGTGTCTTTTAAATTA (SEQ ID NO: 73412)
hCV22274624	G/A	CCCTACAGAGGATGTCAG (SEQ ID NO: 73413)	CCCTACAGAGGATGTCAA (SEQ ID NO: 73414)	CAGAGCCTCCCTTGTGCAC (SEQ ID NO: 73415)
hCV2253681	T/C	CCTGTGCGGGGAGAT (SEQ ID NO: 73416)	CCTGTGCGGGGAGAC (SEQ ID NO: 73417)	GAAGCGGAGCCTGAGAA (SEQ ID NO: 73418)
hCV2259750	A/T	AGAAACTGGCTCTGAAGACA (SEQ ID NO: 73419)	CAGAAACTGGCTCTGAAGACT (SEQ ID NO: 73420)	GAAGTGGGCATGGGTATAC (SEQ ID NO: 73421)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV2259921	T/C	CTTTCGAAAGTTTCAGTTGAACT (SEQ ID NO: 73422)	TTCGAAAGTTTCAGTTGAAACC (SEQ ID NO: 73423)	GTTTTCCAGCTTGCATGTC (SEQ ID NO: 73424)
hCV2303890	T/G	CACTATCGAAGTCCCCAAAT (SEQ ID NO: 73425)	CACTATCGAAGTCCCCAAAG (SEQ ID NO: 73426)	CATCTGGTTTTTGACAATCATTATA (SEQ ID NO: 73427)
hCV2310409	A/T	CTCAGGGAGGGAGAGAGA (SEQ ID NO: 73428)	CTCAGGGAGGGAGAGAGT (SEQ ID NO: 73429)	ACAGCTCAGGCAGAAACTG (SEQ ID NO: 73430)
hCV2415786	A/G	CTCAGCTGAACCTGGCTA (SEQ ID NO: 73431)	CAGCTGAACCTGGCTG (SEQ ID NO: 73432)	GAAACACCTCCTCCATCTTC (SEQ ID NO: 73433)
hCV2443247	G/T	CCAGAATGCCCTTCATCG (SEQ ID NO: 73434)	GCCAGAATGCCCTTCATCT (SEQ ID NO: 73435)	GCATCACAGCTTGCTTCTATAA (SEQ ID NO: 73436)
hCV2462424	C/T	CTTATGAGATTTTCTGTCCAGTC (SEQ ID NO: 73437)	CTTATGAGATTTTCTGTCCAGTT (SEQ ID NO: 73438)	TGGAATTGCAGATGAACAAG (SEQ ID NO: 73439)
hCV2531086	A/G	GCCCCCTCTCTGAAGA (SEQ ID NO: 73440)	CCCCCTCTCTGAAGG (SEQ ID NO: 73441)	CCAGTTCGTGGTATGTTTCATCT (SEQ ID NO: 73442)
hCV2531732	T/C	CCACTGTTGCCATGAT (SEQ ID NO: 73443)	CCACTGTTGCCATGAC (SEQ ID NO: 73444)	CCTTCGGCTCCATAAG (SEQ ID NO: 73445)
hCV2531795	C/G	CTCTCAGCTCAAGCCTCC (SEQ ID NO: 73446)	TCTCAGCTCAAGCCTCG (SEQ ID NO: 73447)	CCTGCCACCTGTCCTCT (SEQ ID NO: 73448)
hCV2532034	C/T	CTGAAGCGCAACCATAAC (SEQ ID NO: 73449)	CTGAAGCGCAACCATAAT (SEQ ID NO: 73450)	TCCCATAGAAAAATGCACCTAAG (SEQ ID NO: 73451)
hCV25472003	A/C	GGTAAAGCTCCAGCAGCTGTA (SEQ ID NO: 73452)	GGTAAAGCTCCAGCACTGTC (SEQ ID NO: 73453)	TCAGCAGCAGGTTACACATAA (SEQ ID NO: 73454)
hCV25472673	C/T	TGGGCTCCATCCCAC (SEQ ID NO: 73455)	TGGGCTCCATCCCAT (SEQ ID NO: 73456)	CCAATTCCTTTTCTCTTTCAGTT (SEQ ID NO: 73457)
hCV25473150	T/C	CCCACATTCTCTCTCCTT (SEQ ID NO: 73458)	CCCACATTCTCTCTCCTC (SEQ ID NO: 73459)	TGAGGAGCGTGGGTGAC (SEQ ID NO: 73460)
hCV25473653	G/A	CTCTAACATCACCGGTGACG (SEQ ID NO: 73461)	CCTCTAACATCACCGGTGACA (SEQ ID NO: 73462)	GAGCTCTGGGTCAGAACTGT (SEQ ID NO: 73463)
hCV25474627	T/C	GGTACCATGGACTGTACTCACT (SEQ ID NO: 73464)	GTACCATGGACTGTACTCACC (SEQ ID NO: 73465)	AGGTTGGTTCTGGAGATGAC (SEQ ID NO: 73466)
hCV25474661	A/G	AGGAACTACGGCGATATCTAA (SEQ ID NO: 73467)	AGGAACTACGGCGATATCTAG (SEQ ID NO: 73468)	GGGTTTCCCTGGACACAT (SEQ ID NO: 73469)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV25477	A/G	GCACAGCACCTTATGTCCA (SEQ ID NO: 73470)	CACAGCACCTTATGTCCG (SEQ ID NO: 73471)	CCTCTTGAGCATTTCATTGTAATT (SEQ ID NO: 73472)
hCV2548962	C/T	CAAATACATCTCCAGGATC (SEQ ID NO: 73473)	CAAATACATCTCCAGGATT (SEQ ID NO: 73474)	GTTTAAATTGCAGTTTGAATGATAT (SEQ ID NO: 73475)
hCV25591528	T/C	TCCAAAAGGACCTGACAT (SEQ ID NO: 73476)	TCCAAAAGGACCTGACAC (SEQ ID NO: 73477)	GGCTGCAGAAATGGAATTT (SEQ ID NO: 73478)
hCV25594697	C/T	GCGCGTCTTCCCC (SEQ ID NO: 73479)	GCGCGTCTTCCCT (SEQ ID NO: 73480)	TCACGTGCTTGACGATTATC (SEQ ID NO: 73481)
hCV25594815	G/A	GCTGGCCGCTCTG (SEQ ID NO: 73482)	AGCTGGCCGCTCTA (SEQ ID NO: 73483)	CACCGCTCATGGACACA (SEQ ID NO: 73484)
hCV25596020	G/C	AACCCCATTCCTTGAG (SEQ ID NO: 73485)	AACCCCATTCCTTGAC (SEQ ID NO: 73486)	AGGAACTCCCTTTGGAGATAT (SEQ ID NO: 73487)
hCV25598595	A/G	AACAATAGAACATCTTTTCACAA (SEQ ID NO: 73488)	AACAATAGAACATCTTTTCACAG (SEQ ID NO: 73489)	CCAAACACCAACTTGATCATC (SEQ ID NO: 73490)
hCV25602572	G/A	GCAATCCAGGTTGAGAG (SEQ ID NO: 73491)	GCAATCCAGGTTGAGAA (SEQ ID NO: 73492)	CGCACCTGGAGTGATAGAC (SEQ ID NO: 73493)
hCV25603879	C/T	GAAGTCATTCTGCTCTGC (SEQ ID NO: 73494)	ATGAAGTCATTCTGCTCTGT (SEQ ID NO: 73495)	TTTCCATCTCCTAACTCTTTTCTAG (SEQ ID NO: 73496)
hCV25604100	G/A	ACAAAGTGTCTGGCTGAGG (SEQ ID NO: 73497)	CACAAAGTGTCTGGCTGAGA (SEQ ID NO: 73498)	TGCCTCAGACACTGAGAAATTAT (SEQ ID NO: 73499)
hCV25605897	G/T	AAGGCAGGATGGGAGTG (SEQ ID NO: 73500)	AAAGGCAGGATGGGAGTT (SEQ ID NO: 73501)	CGTCAAAGCACTAATGTCATGT (SEQ ID NO: 73502)
hCV25605906	A/G	CTTCAACTTGGAAAGACATCTTA (SEQ ID NO: 73503)	TCAACTTGGAAAGACATCTTG (SEQ ID NO: 73504)	CTTGAAAGGGTAGAAGTAAGACATAT (SEQ ID NO: 73505)
hCV25607736	T/G	GCTGGGAGCTTTCCTGT (SEQ ID NO: 73506)	CTGGGAGCTTTCCTGG (SEQ ID NO: 73507)	CAGGACAGCACTGGTGTCT (SEQ ID NO: 73508)
hCV25607748	A/T	GCTGTGCTGCTGGTACA (SEQ ID NO: 73509)	GGCTGTGCTGCTGGTACT (SEQ ID NO: 73510)	GGCATGACCTCTGACATCTC (SEQ ID NO: 73511)
hCV25608687	C/A	GGGTAGTACCAAAAATATTACTTACTTTC (SEQ ID NO: 73512)	GGGTAGTACCAAAAATATTACTTACTTTA (SEQ ID NO: 73513)	CCAAAGTGGAGAAGTGACTAGACA (SEQ ID NO: 73514)
hCV25608809	G/A	AGGGAACCCCGAGAG (SEQ ID NO: 73515)	AGGGAACCCCGAGAGAA (SEQ ID NO: 73516)	ACAAGTTTCATTGTGAATGTGA (SEQ ID NO: 73517)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV25610227	C/T	GCAGGTGGCGTATCTGTGTC (SEQ ID NO: 73518)	GCAGGTGGCGTATCTGTT (SEQ ID NO: 73519)	CGGCCTGGAAACCTTAGC (SEQ ID NO: 73520)
hCV25610470	T/C	CACAATCACCAACGGTCT (SEQ ID NO: 73521)	ACAATCACCAACGGTCC (SEQ ID NO: 73522)	CCTTCTGCATCAGCATCTTC (SEQ ID NO: 73523)
hCV25610773	C/G	ACCCCCCGAAGAACC (SEQ ID NO: 73524)	CCCCCGAAGAACG ID (SEQ NO: 73525)	GCGTGGAGATCCTGACTC (SEQ ID NO: 73526)
hCV25614016	A/G	AGTGGCCCAAGAACAACCA (SEQ ID NO: 73527)	TGGCCAAAGAACAACCG (SEQ ID NO: 73528)	GGTATGAGGCAAAAGTTCCTG (SEQ ID NO: 73529)
hCV25616048	C/G	GTTGAAAAATGTGAATCAGCAC (SEQ ID NO: 73530)	GTTGAAAAATGTGAATCAGCAG (SEQ ID NO: 73531)	TGCAGAGCTTCCAAGTTTT (SEQ ID NO: 73532)
hCV25617360	C/T	GGGTCTCCCACTCCATAC (SEQ ID NO: 73533)	GGTCTCCCACTCCATAT (SEQ ID NO: 73534)	GGGCCCCATTAACTCACATAC (SEQ ID NO: 73535)
hCV25617557	T/C	GCCTCCCGGAGGACTT (SEQ ID NO: 73536)	GCCTCCCGGAGGACTC (SEQ ID NO: 73537)	GGAACCTAACCATGGCTTCTCTTA (SEQ ID NO: 73538)
hCV25618313	C/T	GGGACCGGATCAGCAC (SEQ ID NO: 73539)	GGGACCGGATCAGCAT (SEQ ID NO: 73540)	GGCAGTAGGATGAATTAGAAAAGTG (SEQ ID NO: 73541)
hCV25620145	T/C	CACACCAGCAATGATGAAACT (SEQ ID NO: 73542)	CACCAGCAATGATGAAACC (SEQ ID NO: 73543)	GGGCTAACTCTTTGCATGTTT (SEQ ID NO: 73544)
hCV25623155	G/A	CGGAATCGAAAACGTGAG (SEQ ID NO: 73545)	CGGAATCGAAAACGTGAA (SEQ ID NO: 73546)	GCACCTCTCGGAGTGATCT (SEQ ID NO: 73547)
hCV25623804	T/C	TTTCAAGCTGTCTCCTACCAT (SEQ ID NO: 73548)	TTTCAAGCTGTCTCCTACCAC (SEQ ID NO: 73549)	GGAGAGAAGGGAAGGACTAAAG (SEQ ID NO: 73550)
hCV25624076	G/A	AGTTTGAGGATGATGCTATTAG (SEQ ID NO: 73551)	AGTTTGAGGATGATGCTATTAA (SEQ ID NO: 73552)	AGTGGGTTATTGTCACCTTTTCTAT (SEQ ID NO: 73553)
hCV25630686	C/T	AGGTTGTACCTGTAGCACTAAGAC (SEQ ID NO: 73554)	TAGGTTGTACCTGTAGCACTAAGAT (SEQ ID NO: 73555)	TGGGCTCCTCAGAGAAAAATAT (SEQ ID NO: 73556)
hCV25631989	C/T	AAGATAAGCCTGTCACTGGTC (SEQ ID NO: 73557)	AAGATAAGCCTGTCACTGGTT (SEQ ID NO: 73558)	CAAGCCAGCCCTAATAAACATAA (SEQ ID NO: 73559)
hCV25632288	A/C	TGAGTGTCTCAGTTTCTAGCA (SEQ ID NO: 73560)	GAGTGTCTCAGTTTCTAGCC (SEQ ID NO: 73561)	CTGAAGAGGACATGTCAAATATTAC (SEQ ID NO: 73562)
hCV25636672	C/T	GAAACATCAAAAATTCTCCAGAC (SEQ ID NO: 73563)	GAAACATCAAAAATTCTCCAGAT (SEQ ID NO: 73564)	TGGGTATTGCATTTTAAAGTTTAG (SEQ ID NO: 73565)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV/25637309	A/T	GGCCACTTTGCCTGAATA (SEQ ID NO: 73566)	GGCCACTTTGCCTGAATT (SEQ ID NO: 73567)	CGAAATGTTTCATTTTAAAGTCAGA (SEQ ID NO: 73568)
hCV/25637537	A/G	ATGGCCAACTCCTTCA (SEQ ID NO: 73569)	TGGCCAACCTCCTTCG (SEQ ID NO: 73570)	GGTCCATGCTCCAGATGA (SEQ ID NO: 73571)
hCV/25638153	G/C	CCTCTCCACAGCGTTTGG (SEQ ID NO: 73572)	CCTCTCCACAGCGTTTTC (SEQ ID NO: 73573)	GCAGCGGCCACAGAG (SEQ ID NO: 73574)
hCV/25638155	C/T	GTGTCTTCCCCCACCAC (SEQ ID NO: 73575)	GTGTCTTCCCCCACCAC (SEQ ID NO: 73576)	TGCGGCAGCAACTGA (SEQ ID NO: 73577)
hCV/25640926	T/C	GGCCAGAGACAGGAAAAT (SEQ ID NO: 73578)	GGCCAGAGACAGGAAAAC (SEQ ID NO: 73579)	GCCTGCCCCCTCTGTTCA (SEQ ID NO: 73580)
hCV/25646248	C/T	ATCTACACCATTGCACAC (SEQ ID NO: 73581)	CATCTACACCATTGCACAT (SEQ ID NO: 73582)	GCCTCCTCCCTTTTCAGT (SEQ ID NO: 73583)
hCV/25651174	T/C	CGCTGCAGGGTCAT (SEQ ID NO: 73584)	CGCTGCAGGGTCAC (SEQ ID NO: 73585)	CCTCCCCGCAGAGAAATTA (SEQ ID NO: 73586)
hCV/25651593	G/T	TGTCCTGGTTCTGCTCAG (SEQ ID NO: 73587)	ATGTCTGGTTCTGCTCAT (SEQ ID NO: 73588)	CTGGTCTCCTTCGTTCCAGA (SEQ ID NO: 73589)
hCV/25652706	C/G	GGTGTTCCTGGCTTCC (SEQ ID NO: 73590)	GGTGTTCCTGGCTTCG (SEQ ID NO: 73591)	GGAGAACCTAACAAAGGATTTTACTA (SEQ ID NO: 73592)
hCV/25653599	C/T	AAGAATTTGAACCTTACTGATGAGAC (SEQ ID NO: 73593)	TAAGAATTTGAACCTTACTGATGAGAT (SEQ ID NO: 73594)	ATGTGGAGTTTGATTTCCCTTATTA (SEQ ID NO: 73595)
hCV/25759173	T/G	AGTCAAAAATGGACACAATCTTCT (SEQ ID NO: 73596)	CAAAAATGGACACAATCTTCG (SEQ ID NO: 73597)	TGAGATAAGGAAACCTCTTGATAA (SEQ ID NO: 73598)
hCV/25771227	G/C	CCTGGATGAGGAATCC TACTG (SEQ ID NO: 73599)	CCTGGATGAGGAATCC TACTC (SEQ ID NO: 73600)	GCTGCCGTACCTGTTGTAG (SEQ ID NO: 73601)
hCV/25772464	C/T	CCCACTGTTGCCATGAC (SEQ ID NO: 73602)	CCCACTGTTGCCATGAT (SEQ ID NO: 73603)	CCTTCCGGCTCCATAAG (SEQ ID NO: 73604)
hCV/25922320	T/C	CTCGCAGCGGTGAGT (SEQ ID NO: 73605)	TCGCAGCGGTGAGC (SEQ ID NO: 73606)	GCTGGCGGGAATTTCT (SEQ ID NO: 73607)
hCV/25924149	G/C	GCAGCGACCATGAG (SEQ ID NO: 73608)	GCAGCGACCATGAC (SEQ ID NO: 73609)	TCCTCTGCTCCACTCTG (SEQ ID NO: 73610)
hCV/25925506	G/T	GTGATAGTGGTGAGAGTGG (SEQ ID NO: 73611)	TGTGATAGTGGTGAGAGTGT (SEQ ID NO: 73612)	ACTGAGCTTGATTTTCTCTTTTAAT (SEQ ID NO: 73613)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV25928236	G/C	GATGGTGAAGTCCTTTATCTCTG (SEQ ID NO: 73614)	GATGGTGAAGTCCTTTATCTCTC (SEQ ID NO: 73615)	AGCACAAATTTGCAATCACAC (SEQ ID NO: 73616)
hCV25928842	A/T	CACGACGATCCCTTCTGA (SEQ ID NO: 73617)	CACGACGATCCCTTCTGT (SEQ ID NO: 73618)	GGCGCGAGCTGTCTG (SEQ ID NO: 73619)
hCV25930271	G/A	GAATCTCATGTTTCAGGAAATG (SEQ ID NO: 73620)	CGAATCTCATGTTTCAGGAAATA (SEQ ID NO: 73621)	GCCATGGCCCATAAAAAC (SEQ ID NO: 73622)
hCV25931060	C/T	CCCAACCCATTGCTCC2 (SEQ ID NO: 73623)	GCCCAACCCATTGCTCT (SEQ ID NO: 73624)	TGTCAGCACAGAATCCACTAC (SEQ ID NO: 73625)
hCV25931248	A/G	CTACTTTATCACCTGGCATCA (SEQ ID NO: 73626)	CTTTATCACCTGGCATCG (SEQ ID NO: 73627)	CACCTCAAGTCTGTGAAGTACTTAC (SEQ ID NO: 73628)
hCV25933600	T/C	GAGGCAGTGACTAAGGT (SEQ ID NO: 73629)	AGGCAGTGACTAAGGC (SEQ ID NO: 73630)	GCATGTCAGAATCCTCAATCTC (SEQ ID NO: 73631)
hCV25941408	G/T	CATGGAGTCAACTCTTGAGG (SEQ ID NO: 73632)	GCATGGAGTCAACTCTTGAGT (SEQ ID NO: 73633)	GGCTGTGCTTTGTCTGATCT (SEQ ID NO: 73634)
hCV25944011	C/A	GTATGACCTGTACTTCTGGAGAC (SEQ ID NO: 73635)	GTATGACCTGTACTTCTGGAGAA (SEQ ID NO: 73636)	AGGCCCCCTCTCAATACT (SEQ ID NO: 73637)
hCV26000635	G/T	TGCTGGAGCAATTGAGAG (SEQ ID NO: 73638)	CTGCTGGAGCAATTGAGAT (SEQ ID NO: 73639)	TCTTCCCCTCGTTTCTTTTC (SEQ ID NO: 73640)
hCV2603660	G/A	GGTCATGGCCTTAGAGACTG (SEQ ID NO: 73641)	GGTCATGGCCTTAGAGACTA (SEQ ID NO: 73642)	GGCCGACCATAGAGATGAG (SEQ ID NO: 73643)
hCV2620926	A/G	CCTAGTCCTGGGCCTGA (SEQ ID NO: 73644)	CCTAGTCCTGGGCCTGG (SEQ ID NO: 73645)	CGCCACCTGCTGAAACA (SEQ ID NO: 73646)
hCV2630153	C/G	AGATGCAACAGAGAAATTTTCTC (SEQ ID NO: 73647)	GATGCAACAGAGAAATTTTCTG (SEQ ID NO: 73648)	CTCTGTTCTTTCAATCTTTAGATG (SEQ ID NO: 73649)
hCV2676035	A/G	AATACACAGTCTTGTTTTAGAATTTAA (SEQ ID NO: 73650)	AATACAGTCTTGTTTTAGAATTTTAG (SEQ ID NO: 73651)	AAATGAGGATGTCTACACAGCTATAT (SEQ ID NO: 73652)
hCV2682687	A/C	AGGAGCTGGAGGAGAA (SEQ ID NO: 73653)	AGGAGCTGGAGGAGAC (SEQ ID NO: 73654)	GAGACCTCAACTTTGTTTGA (SEQ ID NO: 73655)
hCV2716008	G/C	CTGTTGAAGAAAACGCTACCTAG (SEQ ID NO: 73656)	CTGTTGAAGAAAACGCTACCTAC (SEQ ID NO: 73657)	TGCAAGCAGTTGTTTTTATGA (SEQ ID NO: 73658)
hCV2741051	C/T	GCAGCCAGTTTCTCCCC (SEQ ID NO: 73659)	TGCAGCCAGTTTCTCCT (SEQ ID NO: 73660)	CATGAAATGCTTCCAGGTATT (SEQ ID NO: 73661)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV2741083	G/A	GTTCCAAACCAGAGAGAAATG (SEQ ID NO: 73662)	GGTCCAAACCAGAGAGAAATA (SEQ ID NO: 73663)	CTTGCCCCCAACAGATTAG (SEQ ID NO: 73664)
hCV2741104	C/T	CGGTTTACCTTGACATTATTTC (SEQ ID NO: 73665)	CGGTTTACCTTGACATTATTTT (SEQ ID NO: 73666)	GA CTGTTGCCCTTATCTATGT (SEQ ID NO: 73667)
hCV277090	A/G	CTGAAACTTCCGTGGTAGA (SEQ ID NO: 73668)	GAAACTTCCGTGGTAGG (SEQ ID NO: 73669)	GATCAGTCCCTGGTTCTGAA (SEQ ID NO: 73670)
hCV2784540	C/G	ACCTTCAAAGCCTTCAGATC (SEQ ID NO: 73671)	CCTTCAAAGCCTTCAGATG (SEQ ID NO: 73672)	GTGGTTTTGGCTTGATTTTATAT (SEQ ID NO: 73673)
hCV2838900	T/C	CTGATTCTGCCCCCTTTT (SEQ ID NO: 73674)	CTGATTCTGCCCCCTTTT (SEQ ID NO: 73675)	TACATGGAAGCCTGAGACTTAC (SEQ ID NO: 73676)
hCV2838905	C/A	TTCTCCAGAAACTTGACCATC (SEQ ID NO: 73677)	CTTCTCCAGAAACTTGACCATA (SEQ ID NO: 73678)	GAAGATGATGGTAAACCTGAGTTAT (SEQ ID NO: 73679)
hCV2908485	T/C	CAAAGCCATGGTCTTTTCAT (SEQ ID NO: 73680)	CAAAGCCATGGTCTTTTCAC (SEQ ID NO: 73681)	CCGCTCTTCGGTTCTACTC (SEQ ID NO: 73682)
hCV2932115	C/T	TGGACGTGGGCTTTTTC (SEQ ID NO: 73683)	TGGACGTGGGCTTTTTC (SEQ ID NO: 73684)	GCTGCAGCCCCCTTTTCTC (SEQ ID NO: 73685)
hCV2965593	A/G	CAGCAAGTGGAGTTTCCA (SEQ ID NO: 73686)	AGCAAGTGGAGTTTCCG (SEQ ID NO: 73687)	GAGGCAGTGCCAGTG (SEQ ID NO: 73688)
hCV2972952	G/C	TCACACTCTTCCATATTGTCTG (SEQ ID NO: 73689)	GTCACACTCTTCCATATTGTCTC (SEQ ID NO: 73690)	AACAATCCTCACACATCTCTCTT (SEQ ID NO: 73691)
hCV2983036	C/G	GCTGACTTTTTTGCTCTTTC (SEQ ID NO: 73692)	GCTGACTTTTTTGCTCTTTG (SEQ ID NO: 73693)	GCCATTTTCCACAATAAATATTT (SEQ ID NO: 73694)
hCV2983037	C/A	TGAACTTTTCTCAGATTCAGTAAC (SEQ ID NO: 73695)	TGAACTTTTCTCAGATTCAGTAAA (SEQ ID NO: 73696)	GACCACAGCATCAAAGAATGT (SEQ ID NO: 73697)
hCV3011239	G/A	GTTCTGAGCCACTAAGCG (SEQ ID NO: 73698)	GTTCTGAGCCACTAAGCA (SEQ ID NO: 73699)	TCCCCACTTTCTCATTCTTCT (SEQ ID NO: 73700)
hCV3020386	G/T	AGAAATTGTGCCAAAGAAGTTG (SEQ ID NO: 73701)	AAGAATTGTGCCAAAGAAGTTT (SEQ ID NO: 73702)	AACTGGTATAATTTGAATCACATAAAT (SEQ ID NO: 73703)
hCV3035766	T/G	TGCTCTGACCGCAGT (SEQ ID NO: 73704)	GCTCCTGACCGCAGG (SEQ ID NO: 73705)	GGGAGGTCAGCTTTTACAAA (SEQ ID NO: 73706)
hCV3036180	T/G	GGGAATGCGGATGGT (SEQ ID NO: 73707)	GGAATGCGGATGGG (SEQ ID NO: 73708)	CCAGGCGTCGGAACT (SEQ ID NO: 73709)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV3036181	C/G	TCCGAGCCACATCC (SEQ ID NO: 73710)	TCCGAGCCACATCG (SEQ ID NO: 73711)	CATGGAGGAGCTGAGAACAC (SEQ ID NO: 73712)
hCV3046056	A/G	CACCTCCTCATCCACA (SEQ ID NO: 73713)	ACCTCCTCATCCACG (SEQ ID NO: 73714)	CCTCCCCCTCCACAAG (SEQ ID NO: 73715)
hCV305149	T/C	TGAATGGCATCCACAAAAT (SEQ ID NO: 73716)	TGAATGGCATCCACAAAAC (SEQ ID NO: 73717)	CCAAAAGCTAAAAAGCACATCT (SEQ ID NO: 73718)
hCV3068164	C/T	GCATCACCTGCATCCTC (SEQ ID NO: 73719)	GCATCACCTGCATCCTT (SEQ ID NO: 73720)	TGGATTGGTTGCTGTTCA (SEQ ID NO: 73721)
hCV3068176	T/C	TACCACAGCTTGCTCACAT (SEQ ID NO: 73722)	TACCACAGCTTGCTCACAC (SEQ ID NO: 73723)	TTTCCCCCATTTTTCAGTT (SEQ ID NO: 73724)
hCV3084793	C/T	CCCGGCTGGCGCGGACATGGAGGACG TTC (SEQ ID NO: 73725)	CCCGGCTGGCGCGGACATGGAGGACGTTT (SEQ ID NO: 73726)	CAGCTTGGCGCAGGTG (SEQ ID NO: 73727)
hCV3131029	A/G	CAAATAGATGGAACTACTCCA (SEQ ID NO: 73728)	AAATAGATGGAACTACTCCG (SEQ ID NO: 73729)	GCGAGCCCATAGGTTTCT (SEQ ID NO: 73730)
hCV3135085	C/A	CTGGAATGGTTATGGGC (SEQ ID NO: 73731)	TACTGGAATGGTTATGGGA (SEQ ID NO: 73732)	TTTATAGCGTGAAACTAATTCTC (SEQ ID NO: 73733)
hCV3188386	C/T	CGCTGCGCGTGTTC (SEQ ID NO: 73734)	CGCTGCGCGTGTTT (SEQ ID NO: 73735)	GGAGGAAGGGAAGGTACAG (SEQ ID NO: 73736)
hCV3188814	G/A	GCAGATCGAGTTCCG (SEQ ID NO: 73737)	GGCAGATCGAGTTCCA (SEQ ID NO: 73738)	CCTAAACTCCATGAAGAAGACATT (SEQ ID NO: 73739)
hCV3210838	C/T	CTGCATTATTTCTATGACGC (SEQ ID NO: 73740)	TTCTGCATTATTTCTATGACGT (SEQ ID NO: 73741)	CAAAAAATGCCAACACAGTTTAGA (SEQ ID NO: 73742)
hCV3212009	T/C	GTTCTCCCCTTTTCAGTGCT (SEQ ID NO: 73743)	TCTCCCCTTTTCAGTGTC (SEQ ID NO: 73744)	TGTCGGTGACTGTTCTGTAA (SEQ ID NO: 73745)
hCV3215915	G/T	GCATTCGCGAGGGG (SEQ ID NO: 73746)	TGCATTCGCGAGGGT (SEQ ID NO: 73747)	TTTCTTCAGGAGTTGATCTCTA (SEQ ID NO: 73748)
hCV3216553	A/G	GCCATACACCTCTTTTCAGGA (SEQ ID NO: 73749)	CCATACACCTCTTTTCAGGG (SEQ ID NO: 73750)	CCAGGAGGCATGTTGATAAG (SEQ ID NO: 73751)
hCV3216558	C/T	AGTTCCTGACCTTCACATACC (SEQ ID NO: 73752)	AAGTTCCTGACCTTCACATACT (SEQ ID NO: 73753)	CTATGTCAGCATTTGCATCTAA (SEQ ID NO: 73754)
hCV3254661	G/A	GCGTGAGGGTGAGCG (SEQ ID NO: 73755)	GCGTGAGGGTGAGCA (SEQ ID NO: 73756)	GCTAAGGTGACCCAAACTCTTAC (SEQ ID NO: 73757)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV3259537	G/A	ACAGCACAGACTTCACCG (SEQ ID NO: 73758)	GACAGCACAGACTTCACCA (SEQ ID NO: 73759)	GCCATCCCCACAGTG (SEQ ID NO: 73760)
hCV342590	C/T	AGACAAATTCTCTCATGTCCAC (SEQ ID NO: 73761)	AGACAAATTCTCTCATGTCCAT (SEQ ID NO: 73762)	TGTTTTCCAGGAAAAAGATATTC (SEQ ID NO: 73763)
hCV370782	C/T	TTCTACCCAGGTACTTATCATCC (SEQ ID NO: 73764)	CTTTTACCCAGGTACTTATCATCT (SEQ ID NO: 73765)	GGATTCACTGTGAAAGAACAGTAT (SEQ ID NO: 73766)
hCV435368	T/C	GAGGTCTTGAATAACAGGGATT (SEQ ID NO: 73767)	GAGGTCTTGAATAACAGGGATC (SEQ ID NO: 73768)	TCTTGAGAGGTGTGATCACTAACTT (SEQ ID NO: 73769)
hCV517658	T/C	AATGGCCTTGGACTTGAT (SEQ ID NO: 73770)	AATGGCCTTGGACTTGAC (SEQ ID NO: 73771)	CTCTGCCATGCAAAACAC (SEQ ID NO: 73772)
hCV529706	G/C	GCGAGGACGAAGGGG (SEQ ID NO: 73773)	GCGAGGACGAAGGGC (SEQ ID NO: 73774)	GGAGGATGAATGGACAGACAA (SEQ ID NO: 73775)
hCV529710	G/C	CCGACCCGAACATAAGG (SEQ ID NO: 73776)	CCGACCCGAACATAAGC (SEQ ID NO: 73777)	CGCGTTCCCCATGTC (SEQ ID NO: 73778)
hCV5687	T/C	CCCTCAGTGTGACTGAGAT (SEQ ID NO: 73779)	CCCTCAGTGTGACTGAGAC (SEQ ID NO: 73780)	CCAGGCATTTCCCATACAG (SEQ ID NO: 73781)
hCV596331	T/C	TCCACATCAGGAAAAACAGT (SEQ ID NO: 73782)	CCACATCAGGAAAAACAGC (SEQ ID NO: 73783)	CATTGGCCAATGAGAAATATCA (SEQ ID NO: 73784)
hCV598677	G/T	CCAAGCTGAAAGGCAAG (SEQ ID NO: 73785)	CCAAGCTGAAAGGCAAT (SEQ ID NO: 73786)	CAGCCAGGTGGAGAGT (SEQ ID NO: 73787)
hCV7441704	A/G	CCAACCGAGATCAGATTGA (SEQ ID NO: 73788)	CAACCGAGATCAGATTGG (SEQ ID NO: 73789)	TGATGCTGATTGTGGATGATA (SEQ ID NO: 73790)
hCV7449215	T/C	CCCAGGCCCCAGTTTCAT (SEQ ID NO: 73791)	CCCAGGCCCCAGTTTCAC (SEQ ID NO: 73792)	CATGTCTGGCTGGAGAGTA (SEQ ID NO: 73793)
hCV7482175	A/G	TGCCCAGGGCTCTTGATA (SEQ ID NO: 73794)	GCCCAGGGCTCTTGATG (SEQ ID NO: 73795)	CCATCGCATGCTCAATACA (SEQ ID NO: 73796)
hCV7482175	A/G	TGCCCAGGGCTCTTGATA (SEQ ID NO: 73797)	GCCCAGGGCTCTTGATG (SEQ ID NO: 73798)	CCATCGCATGCTCAATACA (SEQ ID NO: 73799)
hCV7489257	A/G	TCCCAAAACCTGGAGACTA (SEQ ID NO: 73800)	CCCAAAACCTGGAGACTG (SEQ ID NO: 73801)	TCATGGCATCTTCCCTCAA (SEQ ID NO: 73802)
hCV7490119	G/C	GCCTTGGGGCACATG (SEQ ID NO: 73803)	GCCTTGGGGCACATC (SEQ ID NO: 73804)	GGAAATTTATGGCAGTTTTAAACAT (SEQ ID NO: 73805)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV7490135	G/A	GCAGTCTGAACAAAGTAGATG (SEQ ID NO: 73806)	CGCAGTCTGAACAAAGTAGATA (SEQ ID NO: 73807)	CGTGCATGTTTTGAAAAATGTA (SEQ ID NO: 73808)
hCV7490146	C/T	GCGCTCTGTTCCTTGTATC (SEQ ID NO: 73809)	GCGCTCTGTTCCTTGTATT (SEQ ID NO: 73810)	CCTTTCTCCTGCAGAAATAAGA (SEQ ID NO: 73811)
hCV7497135	A/G	CACTGGAGAATGCACA (SEQ ID NO: 73812)	CACTGGAGAATGCACG (SEQ ID NO: 73813)	GTGTTCTCGCTCACAGAA (SEQ ID NO: 73814)
hCV7499900	T/C	CACACCAGCAATGATGAAACT (SEQ ID NO: 73815)	CACCAGCAATGATGAAACC (SEQ ID NO: 73816)	GGCGGGTTCCAGACAA (SEQ ID NO: 73817)
hCV7514870	A/C	CATCTTGCCACACAGCAA (SEQ ID NO: 73818)	CATCTTGCCACACAGCAC (SEQ ID NO: 73819)	GAAGTGGGCACTGAACAACT (SEQ ID NO: 73820)
hCV7538986	G/A	CAGGGATGTTATTATGGTCG (SEQ ID NO: 73821)	TCAGGGATGTTATTATGGTCA (SEQ ID NO: 73822)	CTGTCTGGGTGGGAATGTA (SEQ ID NO: 73823)
hCV7559757	T/G	GCATTAAGTCTCTCTGGGT (SEQ ID NO: 73824)	CATTAAGTCTCTCTGGGG (SEQ ID NO: 73825)	CAACACACGAGCTACAAACTACA (SEQ ID NO: 73826)
hCV7565899	C/T	AAGATATCAATGTTCTCTGTCTTC (SEQ ID NO: 73827)	AAGATATCAA TGTTCCTGTCTGTTT (SEQ ID NO: 73828)	ATCACTGGTTCCTTCAACTGT (SEQ ID NO: 73829)
hCV7574719	G/A	CGCTCAGTAACCTGCG (SEQ ID NO: 73830)	GCGCTCAGTAACCTGCA (SEQ ID NO: 73831)	GGTCTGGGCCCTTTCATAAG (SEQ ID NO: 73832)
hCV7591528	G/A	CTCATAGCTTGACCTTCGAG (SEQ ID NO: 73833)	CTCATAGCTTGACCTTCGAA (SEQ ID NO: 73834)	ACTCCTCATCAGTCACAGACAC (SEQ ID NO: 73835)
hCV7615375	C/T	CTTCGTCGCAATGGC (SEQ ID NO: 73836)	TTCTTCGTCGCAATGGT (SEQ ID NO: 73837)	GCAATTTCTGCACAGAAATATT (SEQ ID NO: 73838)
hCV7615376	G/A	GATGAGATCAACACAATCTTCAG (SEQ ID NO: 73839)	GATGAGATCAACACAATCTTCAA (SEQ ID NO: 73840)	CCTGAAGCTCGTTTTTGAATAA (SEQ ID NO: 73841)
hCV761961	G/A	CACAGTCAAAGAAATCAAGCG (SEQ ID NO: 73842)	TCACAGTCAAAGAAATCAAGCA (SEQ ID NO: 73843)	CCGTTTGAATTTTCCAATAAG (SEQ ID NO: 73844)
hCV7798230	G/C	GAGCGAGGGCTCAGG (SEQ ID NO: 73845)	GAGCGAGGGCTCAGC (SEQ ID NO: 73846)	CCTCCCTGGAGAATACTGTG (SEQ ID NO: 73847)
hCV789270	C/G	CCTTTCCCAACACAGC (SEQ ID NO: 73848)	CCTTTCCCAACACAGG (SEQ ID NO: 73849)	CATCTCAGCCTCCTTACCA (SEQ ID NO: 73850)
hCV7900503	C/T	CGTCTCCAGGAAAATCATAAC (SEQ ID NO: 73851)	CGTCTCCAGGAAAATCATAAT (SEQ ID NO: 73852)	TGAGTTATTGCTACTTTCAGAAATCAT (SEQ ID NO: 73853)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV790057	A/G	AGCAGCTCCGAGTCCA (SEQ ID NO: 73854)	AGCAGCTCCGAGTCCG (SEQ ID NO: 73855)	GGCCACACAAGGTGAAAT (SEQ ID NO: 73856)
hCV795441	G/C	TGTGGGCCAGGACG (SEQ ID NO: 73857)	CTGTGGGCCAGGACC (SEQ ID NO: 73858)	ACCCACCAGGACCTAAAAAG (SEQ ID NO: 73859)
hCV795442	G/A	CATTCAATGCAATACGTCG (SEQ ID NO: 73860)	CCATTCAATGCAATACGTCA (SEQ ID NO: 73861)	TGGTCCTGGCCTGAAC (SEQ ID NO: 73862)
hCV8400671	A/G	TTGTTAACATATATACTTACTGGAGA (SEQ ID NO: 73863)	TGTTAACATATATACTTACTGGAGG (SEQ ID NO: 73864)	TGCCTCTTCTTTTATTTATGTC (SEQ ID NO: 73865)
hCV8692704	C/A	CTGAGGACCCCTTGAGAC (SEQ ID NO: 73866)	CTGAGGACCCCTTGAGAA (SEQ ID NO: 73867)	GCTCACCAGCCCCTGAAG (SEQ ID NO: 73868)
hCV8695674	A/C	ACCGGCACAAAGGAGAA (SEQ ID NO: 73869)	ACCGGCACAAAGGAGAC (SEQ ID NO: 73870)	TGCCCTTGTCACCTTCTGTAT (SEQ ID NO: 73871)
hCV8705506	C/G	CCACTTCGGGTTCTCTC (SEQ ID NO: 73872)	CCACTTCGGGTTCTCTG (SEQ ID NO: 73873)	CCCTGGCTTCAACATGA (SEQ ID NO: 73874)
hCV8708464	A/G	CCCTCTCCAGCGGGA (SEQ ID NO: 73875)	CCTCTCCAGCGGGG (SEQ ID NO: 73876)	CCAAAGCAGGGTTCACTACC (SEQ ID NO: 73877)
hCV8709053	G/A	GCCCAGATACCCCAAAG (SEQ ID NO: 73878)	GCCCAGATACCCCAAAA (SEQ ID NO: 73879)	GCGGCTTCAGCAGATC (SEQ ID NO: 73880)
hCV8718197	A/G	CCTCTGAGGCCCTGAGAAA (SEQ ID NO: 73881)	CCTCTGAGGCCCTGAGAA (SEQ ID NO: 73882)	GTCTGATTCTCTCATTCTTTC (SEQ ID NO: 73883)
hCV8722981	G/A	GCGCTGTTTGAGG (SEQ ID NO: 73884)	GCGCTGTTTGAGGA (SEQ ID NO: 73885)	TGGCACAGGCAGTATTAAGTAG (SEQ ID NO: 73886)
hCV8726331	A/G	TGGTCTGTTCCCTGGACA (SEQ ID NO: 73887)	GGTCTGTTCCCTGGACG (SEQ ID NO: 73888)	TGCGGTCACACTGACTGAG (SEQ ID NO: 73889)
hCV8774272	T/C	CCGCTTCCGAGCAGTT (SEQ ID NO: 73890)	CCGCTTCCGAGCAGTC (SEQ ID NO: 73891)	CGTCAACATCCTCTTTGAAGAA (SEQ ID NO: 73892)
hCV8784787	A/C	ACTTCTGGGGCTTAGGAA (SEQ ID NO: 73893)	ACTTCTGGGGCTTAGGAC (SEQ ID NO: 73894)	TTCACCCGGGAACCTCTTGT (SEQ ID NO: 73895)
hCV8827241	G/C	TCAAGAGGACAGTGATGGTG (SEQ ID NO: 73896)	TCAAGAGGACAGTGATGGTC (SEQ ID NO: 73897)	TGGTTAGAATCTGTGAAGGAATA (SEQ ID NO: 73898)
hCV8851065	C/G	CCCCGCAGAGAAATTACC (SEQ ID NO: 73899)	CCCCGCAGAGAAATTACG (SEQ ID NO: 73900)	ACGTCGCTGTGGAAGC (SEQ ID NO: 73901)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV8851074	T/G	TGGCTGTTCCAGTACTCCT (SEQ ID NO: 73902)	GGCTGTTCCAGTACTCCG (SEQ ID NO: 73903)	TTCTGGAGAGATACATCTACAAC (SEQ ID NO: 73904)
hCV8851080	T/C	GGCACTGCCCGCTT (SEQ ID NO: 73905)	GGCACTGCCCGCTC (SEQ ID NO: 73906)	CGCTTCCTGGAGAGATACATC (SEQ ID NO: 73907)
hCV8851084	A/G	CAGTGCCGGACAGGA (SEQ ID NO: 73908)	CAGTGCCGGACAGGG (SEQ ID NO: 73909)	CCGCCCGGCACTAAG (SEQ ID NO: 73910)
hCV8851085	T/C	GCTCGTAGTTGTCTGCAT (SEQ ID NO: 73911)	GCTCGTAGTTGTCTGCAC (SEQ ID NO: 73912)	CGCTTCCTGGAGAGATACAT (SEQ ID NO: 73913)
hCV8884725	T/C	GCATTTTCACTCCGAAGTT (SEQ ID NO: 73914)	GCATTTTCACTCCGAAGTC (SEQ ID NO: 73915)	CATGACTAGTCTCATTTTGATTAG (SEQ ID NO: 73916)
hCV8903097	C/T	GACATGGGGGAGTCAC (SEQ ID NO: 73917)	GACATGGGGGAGTCAT (SEQ ID NO: 73918)	GCATCTCGGGTCTACTT (SEQ ID NO: 73919)
hCV8919441	C/T	AGATCACTAGGAGGGTCCTC (SEQ ID NO: 73920)	AGATCACTAGGAGGGTCCTT (SEQ ID NO: 73921)	GGTTCTCTGGATGAAATTACTA (SEQ ID NO: 73922)
hCV8919442	C/T	TTCTACCTTGGGTCCCTTAC (SEQ ID NO: 73923)	TTCTACCTTGGGTCCCTTAT (SEQ ID NO: 73924)	TTCCAGCCCATATTCTGAA (SEQ ID NO: 73925)
hCV8919444	G/A	TGGTGCTGGAGAAATTCAG (SEQ ID NO: 73926)	TGGTGCTGGAGAAATTCAA (SEQ ID NO: 73927)	GGTGCTCCCACTTTATGTG (SEQ ID NO: 73928)
hCV8919523	T/C	GTGAGGAGGGATTTGAATTAT (SEQ ID NO: 73929)	GTGAGGAGGGATTTGAATTAC (SEQ ID NO: 73930)	AAAAAAGTCTGCACCTCAATTCTAC (SEQ ID NO: 73931)
hCV8921130	C/T	AGGCCGTCACTGTAC (SEQ ID NO: 73932)	AGGCCGTCACTGTAT (SEQ ID NO: 73933)	CCATTTCTCCCAAATACAT (SEQ ID NO: 73934)
hCV8921288	G/T	CCGCAGAGGTGTGGG (SEQ ID NO: 73935)	CCGCAGAGGTGTGGT (SEQ ID NO: 73936)	CATTTTGCGGTGGAAATG (SEQ ID NO: 73937)
hCV8928919	G/A	GGTCCCCGAGGAAAGAAAG (SEQ ID NO: 73938)	GGTCCCCGAGGAAAGAAA (SEQ ID NO: 73939)	CCTCCTCGCTCACCTTAAA (SEQ ID NO: 73940)
hCV8932279	A/G	GAGTGCCCTATCAAAATGTTA (SEQ ID NO: 73941)	GAGTGCCCTATCAAAATGTTG (SEQ ID NO: 73942)	TTTGTTGTGCCTGATGATGTA (SEQ ID NO: 73943)
hCV8933098	A/G	TGCAAACTACTGCCAACCA (SEQ ID NO: 73944)	GCAAACTACTGCCAACCG (SEQ ID NO: 73945)	GAGGCAGTGTGATTTAAGAAGA (SEQ ID NO: 73946)
hCV8934009	C/G	CCTGGTTTCACTGTAGTCACTC (SEQ ID NO: 73947)	CCTGGTTTCACTGTAGTCACTG (SEQ ID NO: 73948)	TGGCCACAGGAATCTGTC (SEQ ID NO: 73949)

5
10
15
20
25
30
35
40
45
50
55

(continued)

hCV	Alleles	Sequence A (allele-specific primer)	Sequence B (allele-specific primer)	Sequence C (common primer)
hCV8952817	G/C	CGCATCCAGAACATTCTATG (SEQ ID NO: 73950)	CGCATCCAGAACATTCTATC (SEQ ID NO: 73951)	GCAGCTTCCCATCATACACT (SEQ ID NO: 73952)
hCV899804	T/G	GTGCATTCTGCTTTTAACTCAT (SEQ ID NO: 73953)	TGCATTCTGCTTTTAACTCAG (SEQ ID NO: 73954)	GCAATTGCCTGCTCATTAGT (SEQ ID NO: 73955)
hCV901792	C/G	ATGACAAAGTCTCTGAATAAGAAGTC (SEQ ID NO: 73956)	TGACAAAGTCTCTGAATAAGAAGTG (SEQ ID NO: 73957)	TGAATGCTTTTGATCACATGAGT (SEQ ID NO: 73958)
hCV9458082	C/T	ACAGTCAGGTGGATCTCC (SEQ ID NO: 73959)	CACAGTCAGGTGGATCTCT (SEQ ID NO: 73960)	CACATTCTTGAGGTGCTAG (SEQ ID NO: 73961)
hCV9458936	A/G	CCCCAACCCCAAGAGAAA (SEQ ID NO: 73962)	CCCCAACCCCAAGAGAAG (SEQ ID NO: 73963)	TCCAGGGCTGCTTACTTC (SEQ ID NO: 73964)
hCV9482394	G/A	CGCGCGCTAACCG (SEQ ID NO: 73965)	CGCGCGCTAACCA (SEQ ID NO: 73966)	CGTCCGTCATGGATCAGA (SEQ ID NO: 73967)
hCV9485713	T/C	GCCCAGAGACAGGAAAAT (SEQ ID NO: 73968)	GCCCAGAGACAGGAAAAC (SEQ ID NO: 73969)	GCCTGCCCTCTGTTCA (SEQ ID NO: 73970)
hCV9494470	C/A	AGGGATCCGCAAAAGC (SEQ ID NO: 73971)	CAGGGATCCGCAAAAGA (SEQ ID NO: 73972)	TCTTTCTGCCAGGTACATCA (SEQ ID NO: 73973)
hCV9506149	T/A	CTGCTGGCCGTCCT (SEQ ID NO: 73974)	TGCTGGCCGTCCA (SEQ ID NO: 73975)	ACTCACGCTTGCTTTGACT (SEQ ID NO: 73976)
hCV9514434	T/C	CCTTGAAAGATCTCCCTCTTT (SEQ ID NO: 73977)	CCTTGAAAGATCTCCCTCTTC (SEQ ID NO: 73978)	CGTAGCCCCCAAAGGATCT (SEQ ID NO: 73979)
hCV9546471	A/C	CTCAGGAAGCTAAAAGGTGA (SEQ ID NO: 73980)	TCAGGAAGCTAAAAGGTGC (SEQ ID NO: 73981)	CCTAATATCCCCCTCCAGAACTAT (SEQ ID NO: 73982)
hCV9596962	G/A	CAGCGGAAGCCCAAGG (SEQ ID NO: 73983)	CAGCGGAAGCCCAAGA (SEQ ID NO: 73984)	CCGGGAGATGAAGAAAGAGA (SEQ ID NO: 73985)
hCV9604851	T/C	CCGCCTTGCAGATGAT (SEQ ID NO: 73986)	CCGCCTTGCAGATGAC (SEQ ID NO: 73987)	GGAGCTGGCCATTAGAATC (SEQ ID NO: 73988)
hCV9615318	C/T	GCTCAGATCTGAACCCCTAACTC (SEQ ID NO: 73989)	GCTCAGATCTGAACCCCTAACTT (SEQ ID NO: 73990)	TCACCCCTCTCTGACC (SEQ ID NO: 73991)
hCV9689262	C/T	GGTTGTGCAGAGCAGTTAAC (SEQ ID NO: 73992)	GGTTGTGCAGAGCAGTTAAT (SEQ ID NO: 73993)	TGGCTGTGTTTTGAAAAACTA (SEQ ID NO: 73994)
hCV216064	A/G	TGCCCAGGGCTCTGATA (SEQ ID NO: 73995)	GCCCAGGGCTCTGATG (SEQ ID NO: 73996)	CCATCGCATGCTCAATACA (SEQ ID NO: 73997)

TABLE 6

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV25632288	S0028	SMOKE	N	MI_YOUNGOLD_noASD	C	0.00	0.06	0.0262			0.07	0	1.68
hCV25632288	S0012(F)	no	ALL	MI_noASD	C	0.02	0.05	0.0056			0.42	0.22	0.77
hCV16196014	S0012	HTN	N	MI_noMI	G	0.07	0.04	0.03	0.02	0.49	1.76	1.09	2.85
hCV16196014	S0012	SMOKE	Y	MI_noASD	G	0.06	0.05	1.00	0.74	0.05	1.01	0.49	2.06
hCV16196014	S0028	no	ALL	MI_noASD	G	0.05	0.03	0.04	0.06	0.19	1.91	1.04	3.51
hCV16196014	S0028	SMOKE	Y	MI_noMI	G	0.05	0.04	0.19	0.32	0.02	1.38	0.88	2.18
hCV8692704	S0028	AGE_LT60	Y	MI_noMI	A	0.57	0.51	0.0379			1.30	1.01	1.66
hCV8692704	S0028	AGE_L T60	Y	MI_noMI	A	0.57	0.51	0.0379			1.30	1.02	1.63
hCV8692704	S0012(F)	no	ALL	YoungMI_GT75noASD	A	0.57	0.49	0.0293			1.39	1.08	1.78
hCV8692704	S0012(M)	no	ALL	LT60MI_GT75noMI	A	0.56	0.47	0.0204			1.41	1.06	1.88
hCV25631989	S0028	no	ALL	MI_noMI	T	0.06	0.09	0.0265			0.68	0.48	0.94
hCV25631989	S0012(M)	AGE_LT60	O	MI_LT75noMI	T	0.05	0.10	0.0335			0.44	0.2	0.93
hCV11506744	S0028	no	ALL	MI_noMI	G	0.40	0.46	0.0033			0.77	0.64	0.91
hCV11506744	S0012(M)	no	ALL	LT60MI_60TO75noMI	G	0.39	0.45	0.0497			0.76	0.57	0.99
hCV8884725	S0028	no	ALL	MI_noASD	C	0.01	0.03	0.0138			0.41	0.21	0.79
hCV8884725	S0028	no	ALL	MI_noMI	C	0.02	0.03	0.0262			0.51	0.28	0.9
hCV8884725	S0012(F)	no	ALL	YoungMI_GT75noASD	C	0.01	0.04	0.0228			0.32	0.12	0.81
hCV25637537	S0012	HTN	Y	MI_noMI	A	0.07	0.05	0.04	0.03		1.53	1.02	2.29
hCV25637537	S0028	no	ALL	MI_noASD	A	0.06	0.03	0.01	0.01	0.44	2.13	1.18	3.82
hCV25637537	S0028	no	ALL	MI_noMI	A	0.06	0.04	0.02	0.02	0.54	1.56	1.08	2.24

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV25637537	S0028	no	ALL	MI_YOUNGOLD_noASD	A	0.05	0.01	0.04	0.03		3.62	1.06	12.32
hCV25637537	S0028	HTN	Y	MI_noASD	A	0.06	0.03	0.06	0.05	0.51	1.95	1.00	3.79
hCV25637537	S0028	HTN	Y	MI_noMI	A	0.06	0.04	0.05	0.04	0.64	1.49	1.01	2.19
hCV8851074	S0012	SMOKE	N	MI_noASD	A	0.28	0.33	0.21	0.03	0.46	0.79	0.56	1.12
hCV8851074	S0012	SMOKE	N	MI_noMI	A	0.28	0.33	0.06	0.03	0.61	0.78	0.60	1.01
hCV8851074	S0028	no	ALL	MI_noASD	A	0.33	0.36	0.27	0.03	0.41	0.88	0.70	1.10
hCV8851074	S0028	no	ALL	MI_noMI	A	0.33	0.37	0.05	0.00	0.73	0.84	0.71	1.00
hCV8851074	S0028	SMOKE	N	MI_noASD	A	0.30	0.39	0.05	0.03	0.45	0.66	0.45	0.98
hCV8851074	S0028	SMOKE	N	MI_noMI	A	0.30	0.39	0.02	0.02	0.21	0.66	0.47	0.93
hCV25931248	V0001	no	ALL	MI_noMI	A	0.16	0.20	0.0202	0.0067	0.9257	0.79	0.65	0.96
hCV25931248	S0012(M)	no	ALL	MI_noMI	G	0.83	0.78	0.0062			1.42	1.1	1.82
hCV1403468	V0001	SEX	M	MI_noMI	G	0.13	0.18	0.0075			0.7	0.4	0.8
hCV1403468	S0012(M)	no	ALL	MI_GT75noMI	G	0.16	0.22	0.0252			0.7	0.4	0.9
hCV11482579	S0012	SEX	M	MI_YOUNGOLD_noASD	T	0.08	0.13	0.26	0.46	0.04	0.59	0.23	1.47
hCV11482579	S0028	SEX	M	MI_YOUNGOLD_noASD	T	0.08	0.09	0.63	0.86	0.05	0.78	0.32	1.90
hCV2620926	S0012	no	ALL	MI_noASD	A	0.10	0.15	0.05	0.02	0.92	0.58	0.35	0.97
hCV2620926	S0012	SMOKE	N	MI_noMI	A	0.08	0.11	0.04	0.04	0.39	0.65	0.43	0.98
hCV2620926	S0028	SMOKE	N	MI_noASD	A	0.09	0.16	0.03	0.03	0.32	0.52	0.29	0.93
hCV2620926	S0028	HTN	Y	MI_noASD	A	0.12	0.17	0.03	0.03	0.21	0.68	0.48	0.95
hCV8903097	S0028	SEX	F	MI_YOUNGOLD_noASD	T	0.82	0.93	0.0380			0.37	0.21	0.63
hCV8903097	S0012(F)	no	ALL	MI_noASD	T	0.86	0.90	0.0178			0.65	0.48	0.87
hCV15869253	S0028	SMOKE	N	MI_noMI	T	0.08	0.13	0.0382	0.0276	0.7584	0.60	0.37	0.97

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV15869253	S0012(M)	no	ALL	MI_noMI	T	0.11	0.15	0.048			0.74	0.55	0.99
hCV15869253	S0012(M)	no	ALL	MI_LT75noMI	T	0.11	0.16	0.009			0.66	0.48	0.9
hCV25931060	S0028	AGE_LT60	Y	MI_noMI	A	0.23	0.16	0.0097			1.51	1.13	1.99
hCV25931060	S0012(F)	AGE_L T60	Y	MI_noASD	A	0.22	0.16	0.0373			1.49	1.1	2.01
hCV7441704	S0028	no	ALL	MI_noASD	G	0.03	0.06	0.0386			0.57	0.35	0.9
hCV7441704	S0012(F)	no	ALL	YoungMI_GT75noASD	G	0.03	0.08	0.0321			0.41	0.19	0.88
hCV25653599	V0001	no	ALL	MI_noMI	C	0.14	0.11	0.0256	0.0312	0.3216	1.31	1.03	1.67
hCV25653599	S0012(M)	no	ALL	MI_noMI	T	0.83	0.87	0.0107			0.70	0.53	0.92
hCV25653599	S0012(M)	no	ALL	MI_LT75noMI	T	0.83	0.88	0.0042			0.64	0.46	0.86
hCV1486426	V0001	no	ALL	MI_noMI	A	0.03	0.05	0.0174	0.0174	0.4091	0.63	0.44	0.91
hCV1486426	S0012(M)	no	ALL	MI_noMI	G	0.97	0.95	0.0418			1.69	1.02	2.8
hCV25928842	S0028	SEX	M	MI_noASD	T	0.96	0.99	0.02180			0.32	0.13	0.76
hCV25928842	S0028	SEX	M	MI_noMI	T	0.96	0.99	0.00935			0.30	0.12	0.7
hCV25928842	S0028	SMOKE	Y	MI_noMI	T	0.97	0.99	0.00710			0.28	0.11	0.68
hCV25928842	S0012(M)	no	ALL	MI_noMI	T	0.95	0.98	0.00185			0.43	0.24	0.74
hCV25928842	S0012(M)	no	ALL	MI_LT75noMI	T	0.95	0.99	0.00000			0.15	0.06	0.38
hCV25928842	S0012(M)	SMOKE	Y	MI_LT75noMI	T	0.97	0.99	0.02132			0.25	0.07	0.86
hCV25928236	V0001	HTN	N	MI_noMI	C	0.01	0.02	0.0399	0.0331		0.46	0.22	0.96
hCV25928236	S0012(M)	no	ALL	MI_noMI	G	1.00	0.98	0.0033			4.51	1.41	14.36
hCV25928236	S0012(M)	no	ALL	MI_LT75noMI	G	1.00	0.98	0.0042			4.81	1.46	15.74
hCV25933600	S0028	no	ALL	MI_noASD	C	0.69	0.76	0.0113			0.73	0.56	0.92
hCV25933600	S0012(M)	no	ALL	MI_noMI	C	0.70	0.74	0.0455			0.80	0.64	0.98

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV3215915	S0028	SMOKE	N	MI_noASD	T	0.71	0.79	0.0289			0.63	0.44	0.88
hCV3215915	S0012(F)	no	ALL	MI_noASD	T	0.66	0.73	0.0108			0.73	0.58	0.9
hCV25930271	S0028	no	ALL	MI_noASD	A	0.95	0.99	0.0000			0.17	0.1	0.25
hCV25930271	S0028	SMOKE	Y	MI_noASD	A	0.95	0.99	0.0045			0.14	0.08	0.22
hCV25930271	S0012(F)	SMOKE	Y	MI_noASD	A	0.96	1.00	0.0052			0.10	0.04	0.21
hCV15853800	S0028	no	ALL	MI_YOUNGOLD_noASD	G	0.01	0.04	0.0165			0.17	0.04	0.64
hCV15853800	S0012(F)	AGE_LT60	Y	MI_noASD	G	0.01	0.03	0.0334			0.28	0.09	0.86
hCV2143205	S0012	no	ALL	MI_YOUNGOLD_noASD	A	0.24	0.33	0.06	0.03	0.65	0.64	0.41	1.01
hCV2143205	S0012	HTN	N	MI_YOUNGOLD_noASD	A	0.21	0.34	0.06	0.02	0.64	0.51	0.26	0.99
hCV2143205	S0028	HTN	N	MI_YOUNGOLD_noASD	A	0.13	0.37	0.03	0.05	0.16	0.24	0.06	0.92
hCV7798230	S0028	AGE_LT60	Y	MI_noASD	C	0.70	0.61	0.0171			1.46	1.12	1.89
hCV7798230	S0012(F)	no	ALL	MI_noASD	C	0.74	0.68	0.0329			1.31	1.04	1.64
hCV7798230	S0012(F)	AGE_LT60	Y	MI_noASD	C	0.74	0.65	0.0113			1.52	1.15	1.99
hCV1276216	S0012	BMI_GE27	H	MI_noMI	G	0.28	0.32	0.06	0.03	0.59	0.80	0.64	1.01
hCV1276216	S0012	SEX	M	MI_noMI	G	0.31	0.35	0.05	0.06	0.20	0.81	0.66	0.99
hCV1276216	S0028	BMI_GE27	H	MI_noMI	G	0.31	0.35	0.08	0.04	0.57	0.83	0.68	1.02
hCV1276216	S0028	SEX	M	MI_noMI	G	0.31	0.35	0.05	0.04	0.36	0.80	0.65	1.00
hCV9506149	S0012	no	ALL	MI_noMI	T	0.28	0.23	0.01	0.02	0.14	1.27	1.05	1.53
hCV9506149	S0028	no	ALL	MI_noMI	T	0.28	0.25	0.09	0.05	0.70	1.17	0.98	1.39
hCV11592758	S0028	no	ALL	MI_noASD	C	0.86	0.81	0.00369			1.42	1.11	1.79

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV11592758	S0028	no	ALL	MI_noMI	C	0.86	0.82	0.01264			1.34	1.06	1.68
hCV11592758	V0001	no	ALL	MI_noMI	C	0.82	0.79	0.0263			1.26	1.02	1.53
hCV11592758	S0012(M)	no	ALL	MI_noMI	C	0.84	0.78	0.00721			1.42	1.1	1.83
hCV11592758	S0012(M)	no	ALL	MI_LT75noMI	C	0.84	0.78	0.00929			1.44	1.1	1.89
hCV2146578	S0028	no	ALL	MI_noMI	T	0.44	0.39	0.0120			1.26	1.07	1.47
hCV2146578	S0012(F)	no	ALL	MI_noASD	T	0.47	0.40	0.0184			1.32	1.07	1.61
hCV7900503	S0028	SMOKE	N	MI_YOUNGOLD_noMI	T	0.15	0.30	0.0301			0.42	0.19	0.86
hCV7900503	S0028	AGE_LT60	O	MI_noASD	T	0.18	0.28	0.0116			0.57	0.37	0.86
hCV7900503	S0012(M)	no	ALL	MI_noMI	T	0.19	0.25	0.0077			0.72	0.56	0.91
hCV8921288	S0012	SEX	M	MI_YOUNGOLD_noASD	G	0.20	0.15	0.20	0.03	0.10	1.40	0.86	2.26
hCV8921288	S0012	SMOKE	N	MI_YOUNGOLD_noASD	G	0.24	0.16	0.02	0.01	0.87	1.68	1.08	2.63
hCV8921288	S0028	no	ALL	MI_noASD	G	0.22	0.17	0.03	0.03	0.46	1.34	1.02	1.77
hCV8921288	S0028	SEX	M	MI_noASD	G	0.22	0.17	0.12	0.02	0.33	1.36	0.93	1.97
hCV16171764	S0012	SEX	M	MI_noASD	T	0.03	0.06	0.06	0.11	0.04	0.47	0.22	1.03
hCV16171764	S0028	SEX	M	MI_YOUNGOLD_noASD	T	0.04	0.07	0.38	0.73	0.05	0.65	0.22	1.88
hCV2259750	S0028	no	ALL	MI_noMI	A	0.38	0.34	0.0324	0.0625	0.0934	1.20	1.02	1.41
hCV2259750	S0012(M)	no	ALL	LT60MI_GT75noMI	A	0.37	0.30	0.040			1.38	1.01	1.86
hCV2259750	S0012(M)	no	ALL	MI_GT75noMI	A	0.37	0.30	0.038			1.36	1.01	1.8
hCV2548962	S0012	no	ALL	MI_noASD	C	0.28	0.36	0.09	0.40	0.02	0.72	0.50	1.05
hCV2548962	S0012	BMI_GE27	L	MI_noMI	C	0.27	0.33	0.07	0.01	0.93	0.78	0.59	1.01

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV2548962	S0028	BMI_GE27	L	MI_noASD	C	0.21	0.33	0.01	0.05	0.01	0.55	0.35	0.87
hCV2548962	S0028	BMI_GE27	L	MI_noMI	C	0.21	0.32	0.01	0.02	0.02	0.58	0.39	0.84
hCV25598595	S0012	no	ALL	MI_YOUNGOLD_noASD	G	0.12	0.09	0.12	0.05	0.37	1.42	0.92	2.18
hCV25598595	S0012	SEX	F	MI_noASD	G	0.11	0.06	0.05	0.02	0.18	1.83	1.00	3.35
hCV25598595	S0012	SEX	F	MI_YOUNGOLD_noASD	G	0.13	0.06	0.02	0.01	0.30	2.30	1.17	4.53
hCV25598595	S0012	HTN	Y	MI_YOUNGOLD_noASD	G	0.13	0.07	0.04	0.04	0.37	1.96	1.02	3.76
hCV25598595	S0028	no	ALL	MI_noASD	G	0.12	0.08	0.0190	0.0286	0.1107	1.55	1.08	2.23
hCV25598595	S0028	no	ALL	MI_noMI	G	0.12	0.09	0.0018	0.0025	0.0943	1.49	1.17	1.91
hCV25598595	S0028	no	ALL	MI_YOUNGOLD_noASD	G	0.12	0.04	0.00	0.00	0.23	3.20	1.49	6.89
hCV25598595	S0028	SEX	F	MI_noMI	G	0.12	0.08	0.0224	0.0462	0.0514	1.61	1.07	2.41
hCV25598595	S0028	SEX	F	MI_YOUNGOLD_noASD	G	0.12	0.05	0.04	0.04	0.32	2.85	1.08	7.53
hCV25598595	S0028	HTN	Y	MI_noASD	G	0.12	0.07	0.01	0.01	0.23	1.86	1.19	2.90
hCV25598595	S0028	HTN	Y	MI_noMI	G	0.12	0.08	0.00	0.00	0.13	1.59	1.22	2.07
hCV25598595	S0028	HTN	Y	MI_YOUNGOLD_noASD	G	0.12	0.04	0.01	0.01	0.28	3.16	1.32	7.56
hCV25598595	S0012(M)	no	ALL	LT60MI_GT75noMI	G	0.18	0.13	0.048			1.50	1	2.24
hCV25598595	S0012(M)	no	ALL	MI_noMI	G	0.17	0.13	0.020			1.39	1.05	1.82
hCV25598595	S0012(M)	no	ALL	MI_LT75noMI	G	0.17	0.13	0.033			1.39	1.03	1.87
hCV2908485	V0001	SEX	M	MI_noMI	G	0.42	0.36	0.0348	0.0783	0.0866	1.27	1.02	1.59
hCV2908485	S0012(M)	no	ALL	MI_LT75noMI	G	0.45	0.38	0.0138			1.31	1.05	1.62
hCV11764545	S0012	no	ALL	MI_noMI	T	0.21	0.18	0.07	0.05	0.61	1.22	0.99	1.50
hCV11764545	S0028	SEX	F	MI_noASD	T	0.22	0.15	0.03	0.03	0.32	1.60	1.05	2.45
hCV25944011	V0001	SEX	M	MI_noMI	A	0.38	0.33	0.0392			1.3	1	1.5

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV25944011	S0012(M)	no	ALL	LT60MI_60TO75noMI	A	0.38	0.27	0.0007			1.7	1.2	2.2
hCV7615375	S0012	no	ALL	MI_noASD	T	0.05	0.02	0.02	0.03	0.40	2.24	1.10	4.58
hCV7615375	S0012	no	ALL	MI_noMI	T	0.05	0.03	0.01	0.01	0.30	1.66	1.13	2.42
hCV7615375	S0028	HTN	N	MI_noASD	T	0.09	0.02	0.04	0.02		5.30	1.14	24.60
hCV7615375	S0028	HTN	N	MI_noMI	T	0.09	0.02	0.03	0.01		4.46	1.32	15.09
hCV3084793	S0012	no	ALL	MI_noASD	C	0.17	0.09	0.00	0.00	0.03	1.97	1.36	2.85
hCV3084793	S0012	no	ALL	MI_YOUNGOLD_noASD	C	0.17	0.09	0.00	0.00	0.05	1.97	1.31	2.96
hCV3084793	S0028	HTN	Y	MI_noASD	C	0.16	0.12	0.16	0.33	0.02	1.32	0.92	1.89
hCV901792	S0012	no	ALL	MI_YOUNGOLD_noASD	G	0.10	0.13	0.07	0.04	0.82	0.69	0.47	1.03
hCV901792	S0012	BMI_GE27	L	MI_YOUNGOLD_noASD	G	0.08	0.14	0.04	0.04	0.39	0.55	0.32	0.97
hCV901792	S0028	BMI_GE27	L	MI_YOUNGOLD_noASD	G	0.03	0.10	0.06	0.04		0.23	0.05	1.11
hCV2188895	S0028	AGE_LT60	O	MI_noMI	G	0.49	0.56	0.0323			0.76	0.59	0.97
hCV2188895	S0028	AGE_LT60	O	MI_noMI	G	0.49	0.56	0.0323			0.76	0.59	0.97
hCV2188895	S0012(M)	no	ALL	LT60MI_GT75noMI	G	0.52	0.60	0.0415			0.74	0.55	0.98
hCV2188895	S0012(M)	no	ALL	MI_GT75noMI	G	0.53	0.60	0.0468			0.75	0.57	0.98
hCV1212713	S0012	no	ALL	MI_noASD	G	0.46	0.35	0.02	0.06	0.05	1.56	1.08	2.26
hCV1212713	S0012	no	ALL	MI_noMI	G	0.46	0.41	0.02	0.10	0.03	1.21	1.03	1.43
hCV1212713	S0012	BMI_GE27	L	MI_noASD	G	0.43	0.28	0.01	0.03	0.05	1.99	1.17	3.40
hCV1212713	S0012	HTN	N	MI_noASD	G	0.47	0.33	0.03	0.22	0.02	1.74	1.05	2.90
hCV1212713	S0028	BMI_GE27	L	MI_noASD	G	0.44	0.34	0.04	0.09	0.08	1.58	1.04	2.38

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV1212713	S0028	HTN	N	MI_noMI	G	0.46	0.40	0.08	0.28	0.04	1.79	0.97	3.31
hCV1212713	V0001	no	ALL	MI_noMI	G	0.45	0.40	0.0078	0.1081	0.0041	1.24	1.06	1.45
hCV1212713	V0001	HTN	N	MI_noMI	G	0.46	0.41	0.0229	0.2744	0.0060	1.25	1.03	1.51
hCV277090	S0028	no	ALL	MI_YOUNGOLD_noASD	G	0.58	0.47	0.0359			1.53	1.17	1.97
hCV277090	S0012(F)	AGE_LT60	O	MI_noASD	G	0.59	0.49	0.0461			1.46	1.02	2.09
hCV2716008	S0028	no	ALL	MI_noMI	C	0.20	0.16	0.0202			1.31	1.04	1.64
hCV2716008	S0028	no	ALL	MI_noASD	C	0.20	0.15	0.0204			1.42	1.05	1.9
hCV2716008	S0012(M)	no	ALL	MI_noMI	C	0.24	0.19	0.0206			1.32	1.04	1.67
hCV2531086	S0028	no	ALL	MI_YOUNGOLD_noMI	G	0.78	0.86	0.0314			0.57	0.33	0.95
hCV2531086	S0012(M)	no	ALL	MI_LT75noMI	G	0.77	0.82	0.0198			0.73	0.55	0.94
hCV9615318	V0001	HTN	Y	MI_noMI	A	0.42	0.33	0.0146	0.0069	0.2786	1.46	1.08	1.96
hCV9615318	S0028	SMOKE	N	MI_YOUNGOLD_noMI	A	0.51	0.30	0.0091			2.41	1.29	4.48
hCV9615318	S0012(M)	no	ALL	MI_LT75noMI	A	0.43	0.36	0.0128			1.32	1.06	1.63
hCV9615318	S0012(M)	no	ALL	MI_LT75noMI	A	0.43	0.36	0.0128			1.32	1.06	1.63
hCV25602572	S0028	no	ALL	MI_noMI	A	0.96	0.98	0.0303			0.54	0.35	0.83
hCV25602572	S0028	SMOKE	Y	MI_noMI	A	0.97	0.99	0.0223			0.39	0.22	0.68
hCV25602572	S0012(F)	SMOKE	Y	MI_noASD	A	0.97	0.99	0.0483			0.17	0.07	0.39
hCV25603879	S0012	BMI_GE27	H	MI_noASD	T	0.02	0.05	0.07	0.04		0.39	0.15	1.00
hCV25603879	S0028	no	ALL	MI_noASD	T	0.01	0.17	0.16	0.03		0.07	0.00	1.38
hCV25603879	S0028	BMI_GE27	H	MI_noASD	T	0.02	0.25	0.14	0.01		0.06	0.00	1.19
hCV1662671	S0028	no	ALL	MI_noASD	G	0.39	0.33	0.0440			1.27	1	1.59
hCV1662671	S0012(M)	no	ALL	LT60MI_GT75noMI	G	0.35	0.26	0.0126			1.50	1.09	2.05

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV517658	S0028	AGE_LT60	Y	MI_noASD	C	0.34	0.26	0.0111			1.51	1.16	1.95
hCV517658	S0012(F)	no	ALL	YoungMI_GT75noASD	C	0.39	0.28	0.0026			1.64	1.26	2.1
hCV25608809	V0001	SMOKE	Y	MI_noMI	A	0.55	0.61	0.0300			0.8	0.6	0.9
hCV25608809	S0012(M)	SMOKE	Y	MI_LT75noMI	A	0.52	0.60	0.0230			0.7	0.5	0.9
hCV12029981	S0028	SMOKE	Y	MI_noMI	G	0.91	0.93	0.03942			0.69	0.48	0.98
hCV12029981	S0012(M)	SMOKE	Y	MI_LT75noMI	G	0.90	0.94	0.02876			0.55	0.32	0.93
hCV3011239	S0028	no	ALL	MI_noASD	A	0.39	0.33	0.0249			1.31	1.03	1.64
hCV3011239	S0012(M)	no	ALL	LT60MI_60TO75noMI	A	0.40	0.33	0.0449			1.35	1.01	1.79
hCV370782	S0012	SMOKE	Y	MI_YOUNGOLD_noASD	C	0.36	0.44	0.07	0.02	0.71	0.71	0.50	1.03
hCV370782	S0028	SMOKE	Y	MI_YOUNGOLD_noASD	C	0.32	0.44	0.04	0.01	0.55	0.61	0.38	0.97
hCV25607748	S0028	SEX	M	MI_noASD	T	0.96	0.91	0.0103			2.25	1.25	4.04
hCV25607748	S0012(M)	no	ALL	MI_noMI	T	0.94	0.92	0.0367			1.53	1.02	2.26
hCV3046056	S0028	no	ALL	MI_noASD	G	0.96	0.93	0.0491			1.65	1.03	2.61
hCV3046056	S0012(M)	no	ALL	MI_GT75noMI	G	0.96	0.93	0.0289			1.95	1.1	3.45
hCV11758801	S0028	no	ALL	MI_noMI	G	0.04	0.02	0.0074			2.03	1.35	3.03
hCV11758801	S0012(F)	no	ALL	YoungMI_GT75noASD	G	0.04	0.01	0.0238			3.77	1.97	7.2
hCV2676035	S0028	no	ALL	MI_noMI	G	0.63	0.59	0.0240			1.23	1.04	1.44
hCV2676035	S0028	no	ALL	MI_noASD	G	0.63	0.58	0.0398			1.27	1.06	1.5
hCV2676035	S0012(F)	no	ALL	MI_noASD	G	0.65	0.60	0.0393			1.28	1.03	1.57
hCV16033535	S0028	SEX	F	MI_noMI	T	0.79	0.73	0.0456			1.41	1.03	1.91

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV16033535	S0028	SEX	F	MI_noASD	T	0.79	0.71	0.0195			1.58	1.14	2.18
hCV16033535	S0012(F)	no	ALL	YoungMI_GT75noASD	T	0.82	0.72	0.0012			1.79	1.31	2.43
hCV25608687	S0028	AGE_LT60	O	MI_noMI	A	0.06	0.03	0.0098			2.21	1.18	4.13
hCV25608687	S0028	AGE_LT60	O	MI_noASD	A	0.06	0.02	0.0110			4.22	1.11	15.96
hCV25608687	S0012(M)	AGE_LT60	O	MI_LT75noMI	A	0.10	0.05	0.0189			2.24	1.14	4.38
hCV9482394	S0028	SMOKE	N	MI_noASD	A	0.12	0.07	0.04386			1.75	1.01	3.02
hCV9482394	S0012(M)	SMOKE	N	MI_LT75noMI	A	0.10	0.06	0.04581			1.91	0.98	3.73
hCV25610470	S0028	no	ALL	MI_noMI	C	0.96	0.94	0.0131			1.66	1.09	2.51
hCV25610470	S0028	no	ALL	MI_noMI	C	0.96	0.94	0.0131			1.66	1.12	2.45
hCV25610470	S0028	AGE_LT60	O	MI_noASD	C	0.97	0.94	0.0495			2.43	1.23	4.78
hCV25610470	S0012(F)	AGE_LT60	O	MI_noASD	C	0.98	0.93	0.0300			3.86	1.15	12.92
hCV25610470	S0012(M)	no	ALL	LT60MI_GT75noMI	C	0.95	0.91	0.0180			2.04	1.15	3.59
hCV25610470	S0012(M)	no	ALL	MI_GT75noMI	C	0.96	0.91	0.0038			2.23	1.31	3.79
hCV25610773	S0028	SEX	M	MI_noASD	G	0.11	0.06	0.0191			1.99	1.11	3.56
hCV25610773	S0012(M)	no	ALL	MI_GT75noMI	G	0.10	0.06	0.0313			1.76	1.04	2.95
hCV16170641	S0028	AGE_LT60	O	MI_noMI	C	0.04	0.07	0.0378			0.56	0.31	0.98
hCV16170641	S0012(M)	AGE_LT60	O	MI_LT75noMI	C	0.03	0.09	0.0126			0.34	0.13	0.82
hCV1387523	S0012	no	ALL	MI_noMI	A	0.17	0.22	0.00	0.01	0.03	0.73	0.59	0.90
hCV1387523	S0028	SMOKE	N	MI_YOUNGOLD_noASD	A	0.09	0.21	0.04	0.02		0.36	0.14	0.94
hCV25610227	S0028	no	ALL	MI_YOUNGOLD_noASD	T	0.03	0.07	0.0220			0.33	0.15	0.71
hCV25610227	S0012(F)	no	ALL	MI_noASD	T	0.02	0.05	0.0490			0.50	0.27	0.93
hCV1385736	S0028	no	ALL	MI_noMI	C	0.51	0.58	0.0005			0.73	0.61	0.87

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV1385736	S0012(M)	AGE_LT60	Y	MI_LT75noMI	C	0.55	0.63	0.0291			0.72	0.54	0.96
hCV8400671	S0028	SMOKE	N	MI_noASD	G	0.81	0.88	0.0352			0.58	0.38	0.87
hCV8400671	S0012(F)	no	ALL	YoungMI_GT75noASD	G	0.78	0.85	0.0278			0.64	0.47	0.86
hCV25614016	S0012	SEX	M	MI_YOUNGOLD_noASD	G	0.19	0.11	0.026	0.021	0.456	1.96	1.08	3.57
hCV25614016	S0012	SMOKE	Y	MI_YOUNGOLD_noASD	G	0.19	0.11	0.029	0.011	0.900	1.84	1.07	3.15
hCV25614016	V0001	no	ALL	MI_noMI	G	0.18	0.15	0.045			1.26	1.01	1.55
hCV25614016	V0001	SEX	M	MI_noMI	G	0.18	0.14	0.019			1.44	1.06	1.95
hCV25614016	V0001	SMOKE	Y	MI_noMI	G	0.18	0.12	0.003			1.63	1.17	2.24
hCV14938	S0028	SMOKE	N	MI_noMI	C	0.33	0.26	0.0452			1.44	1.01	2.04
hCV14938	S0028	SMOKE	N	MI_noMI	C	0.33	0.26	0.0452			1.44	1.04	1.98
hCV14938	S0012(F)	no	ALL	YoungMI_GT75noASD	C	0.35	0.27	0.0236			1.48	1.13	1.91
hCV14938	S0012(M)	no	ALL	MI_noMI	C	0.29	0.24	0.0474			1.25	1	1.56
hCV8932279	S0028	SMOKE	N	MI_noASD	G	0.49	0.39	0.02939			1.47	1.04	2.05
hCV8932279	S0028	SMOKE	N	MI_noMI	G	0.49	0.39	0.02148			1.49	1.07	2.06
hCV8932279	S0012(M)	SMOKE	N	MI_LT75noMI	G	0.51	0.34	0.00011			2.03	1.42	2.89
hCV1022614	S0028	no	ALL	MI_noMI	A	0.16	0.21	0.0129			0.75	0.59	0.93
hCV1022614	S0028	no	ALL	MI_noMI	A	0.16	0.21	0.0129			0.75	0.6	0.91
hCV1022614	S0012(F)	no	ALL	MI_noASD	A	0.17	0.21	0.0500			0.75	0.57	0.98
hCV1022614	S0012(M)	no	ALL	MI_noMI	A	0.17	0.21	0.0382			0.76	0.59	0.98
hCV15851292	S0028	no	ALL	MI_noASD	T	0.10	0.15	0.0074			0.63	0.45	0.87

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV15851292	S0012(M)	no	ALL	MI_GT75noMI	T	0.11	0.16	0.0242			0.63	0.42	0.92
hCV3068176	S0012	BMI_GE27	H	MI_noMI	A	0.43	0.50	0.0444	0.0524	0.1867	0.78	0.62	0.99
hCV3068176	S0028	AGE_LT60	O	MI_noMI	C	0.60	0.52	0.0229			1.34	1.04	1.71
hCV3068176	S0028	AGE_LT60	O	MI_noASD	C	0.60	0.50	0.0316			1.50	1.04	2.16
hCV3068176	S0028	no	ALL	MI_noMI	C	0.61	0.55	0.0042			1.28	1.07	1.5
hCV3068176	S0028	AGE_LT60	O	MI_noASD	C	0.62	0.52	0.00050			1.54	1.2	1.97
hCV3068176	S0028	AGE_LT60	O	MI_noMI	C	0.62	0.54	0.00170			1.44	1.14	1.81
hCV3068176	S0012(M)	AGE_LT60	O	MI_LT75noMI	C	0.70	0.56	0.00149			1.81	1.25	2.61
hCV3068176	S0012(M)	AGE_LT60	O	MI_LT75noMI	C	0.70	0.56	0.0015			1.81	1.25	2.61
hCV25617360	S0028	no	ALL	MI_noMI	T	0.16	0.19	0.0492			0.79	0.63	0.99
hCV25617360	S0012(M)	no	ALL	LT60MI_60T075noMI	T	0.15	0.21	0.0293			0.67	0.47	0.95
hCV795442	S0012	BMI_GE27	L	MI_YOUNGOLD_noASD	G	0.29	0.33	0.30	0.05	0.30	0.82	0.56	1.19
hCV795442	S0028	no	ALL	MI_YOUNGOLD_noASD	G	0.29	0.37	0.07	0.04	0.47	0.72	0.50	1.02
hCV795442	S0028	HTN	N	MI_YOUNGOLD_noASD	G	0.13	0.39	0.0318	0.0084	0.3331	0.22	0.06	0.85
hCV795442	S0012(M)	no	ALL	LT60MI_GT75noMI	G	0.14	0.20	0.023			0.65	0.44	0.94
hCV7490119	S0028	SEX	M	MI_noASD	C	0.29	0.37	0.0151			0.68	0.49	0.93
hCV7490119	S0028	no	ALL	MI_noASD	C	0.27	0.35	0.00009			0.68	0.56	0.82
hCV7490119	S0028	no	ALL	MI_noMI	C	0.27	0.35	0.00007			0.69	0.57	0.82
hCV7490119	S0012(M)	no	ALL	LT60MI_60T075noMI	C	0.28	0.36	0.02106			0.71	0.52	0.94
hCV7490119	S0012(M)	no	ALL	LT60MI_60T075noMI	C	0.28	0.36	0.0211			0.71	0.52	0.94
hCV22274307	S0028	SEX	M	MI_YOUNGOLD_noASD	T	0.73	0.91	0.0025			0.27	0.18	0.4
hCV22274307	S0012(F)	no	ALL	MI_noASD	T	0.72	0.78	0.0240			0.74	0.58	0.92

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV8827241	S0028	AGE_LT60	O	MI_noMI	C	0.66	0.59	0.0237			1.35	1.04	1.74
hCV8827241	V0001	no	ALL	MI_noMI	C	0.64	0.59	0.0244			1.2	1	1.4
hCV8827241	V0001	SEX	M	MI_noMI	C	0.64	0.58	0.0264			1.3	1	1.6
hCV8827241	S0012(M)	no	ALL	LT60MI_60TO75noMI	C	0.67	0.59	0.0120			1.4	1	1.9
hCV8827241	S0012(M)	no	ALL	LT60MI_60TO75noMI	C	0.67	0.59	0.0120			1.44	1.08	1.91
hCV761961	S0012	BMI_GE27	H	MI_noMI	T	0.28	0.21	0.00	0.01	0.02	1.51	1.15	1.98
hCV761961	S0012	HTN	N	MI_noMI	T	0.28	0.23	0.07	0.27	0.01	1.32	0.99	1.77
hCV761961	S0028	no	ALL	MI_noASD	T	0.28	0.20	0.0006	0.0013	0.0416	1.55	1.20	2.00
hCV761961	S0028	no	ALL	MI_noMI	T	0.28	0.24	0.0308	0.0331	0.2243	1.21	1.02	1.45
hCV761961	S0028	BMI_GE27	H	MI_noASD	T	0.30	0.20	0.00	0.00	0.13	1.73	1.25	2.41
hCV761961	S0028	no	ALL	MI_noASD	T	0.27	0.21	0.0047			1.46	1.12	1.89
hCV761961	S0012(M)	no	ALL	MI_LT75noMI	T	0.28	0.23	0.030			1.32	1.03	1.68
hGV761961	S0012(M)	no	ALL	MI_LT75noMI	T	0.28	0.23	0.0298			1.32	1.03	1.68
hCV2972952	S0012	AGE_LT60	Y	MI_noASD	G	0.21	0.36	0.02	0.04	0.02	0.47	0.27	0.84
hCV2972952	S0028	no	ALL	MI_noMI	G	0.17	0.20	0.15	0.44	0.02	0.86	0.70	1.05
hCV2972952	S0028	AGE_LT60	Y	MI_noMI	G	0.16	0.22	0.02	0.05	0.04	0.70	0.52	0.94
hCV7538986	S0012(M)	AGE_LT60	Y	MI_LT75noMI	A	0.06	0.03	0.0484			2.24	1.07	4.67
hCV7538986	S0012(M)	SMOKE	Y	MI_LT75noMI	A	0.07	0.04	0.0461			1.98	1.04	3.73
hCV9514434	S0028	no	ALL	MI_noASD	C	0.94	0.91	0.0495			1.53	1.01	2.31

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV9514434	S0012(M)	SMOKE	Y	MI_LT75noMI	C	0.95	0.91	0.0352			1.80	1.06	3.06
hCV3188386	S0012(M)	no	ALL	MI_noMI	T	0.52	0.58	0.0143			0.78	0.64	0.94
hCV3188386	S0012(M)	no	ALL	LT60MI_GT75noMI	T	0.51	0.61	0.0071			0.67	0.5	0.89
hCV3188386	S0012(M)	AGE_LT60	Y	MI_LT75noMI	T	0.51	0.61	0.0039			0.66	0.49	0.87
hCV3188386	S0012(M)	no	ALL	MI_GT75noMI	T	0.52	0.61	0.0092			0.69	0.52	0.9
hCV1923359	S0028	no	ALL	MI_YOUNGOLD_noMI	C	0.46	0.56	0.0368			0.66	0.45	0.96
hCV1923359	S0028	no	ALL	MI_YOUNGOLD_noASD	C	0.46	0.56	0.0368			0.66	0.51	0.85
hCV1923359	S0012(F)	no	ALL	YoungMI_GT75noASD	C	0.41	0.51	0.0130			0.69	0.53	0.87
hCV1923359	S0012(M)	no	ALL	MI_GT75noMI	C	0.50	0.57	0.0243			0.73	0.56	0.96
hCV25623804	S0028	no	ALL	MI_noASD	C	0.98	0.95	0.0014			2.70	1.55	4.68
hCV25623804	S0012(F)	no	ALL	MI_noASD	C	0.98	0.95	0.0138			2.36	1.24	4.5
hCV25922320	S0028	no	ALL	MI_noMI	C	0.71	0.75	0.0472			0.82	0.68	0.97
hCV25922320	S0012(F)	no	ALL	MI-noASD	C	0.71	0.77	0.0188			0.73	0.58	0.91
hCV2259750	S0028	no	ALL	MI_noMI	A	0.37	0.33	0.0457			1.21	1	1.44
hCV2259750	S0028	no	ALL	MI_noMI	A	0.37	0.33	0.0457			1.21	1.02	1.42
hCV2259750	S0012(F)	AGE_LT60	O	MI_noASD	A	0.48	0.34	0.0033			1.78	1.24	2.53
hCV2259750	S0012(M)	no	ALL	MI_GT75noMI	A	0.37	0.30	0.0381			1.36	1.01	1.8
hCV1842400	S0028	SEX	M	MI_YOUNGOLD_noMI	C	0.18	0.06	0.0296			3.54	1.17	10.66
hCV1842400	S0012(M)	no	ALL	MI_noMI	C	0.17	0.13	0.0169			1.39	1.06	1.82
hCV25477	S0028	SMOKE	Y	MI_noASD	G	0.41	0.50	0.0147			0.69	0.5	0.92
hCV25477	S0012(M)	SMOKE	Y	MI_LT75noMI	G	0.38	0.45	0.0374			0.75	0.57	0.98
hCV216064	S0012(M)	no	ALL	LT60MI_60TO75noMI	C	0.24	0.31	0.0432			0.73	0.53	0.98
hCV216064	S0012(M)	no	ALL	LT60MI_60TO75noMI	C	0.24	0.31	0.0432			0.73	0.53	0.98

(continued)

Marker	Study	Stratification	Strata	Status	Allele1	Case Allele1 frq	Control Allele1 frq	Allelic p-value	Dom p-value	Rec p-value	OR	OR 95%CI L	OR 95%CI U
hCV2682687	S0012(M)	AGE_LT60	Y	MI_LT75noMI	C	0.52	0.43	0.0120			1.43	1.08	1.9
hCV2682687	S0012(M)	AGE_LT60	Y	MI_LT75noMI	C	0.52	0.43	0.0120			1.43	1.08	1.9
hCV3188386	S0012(M)	no	ALL	MI-noMI	T	0.52	0.58	0.0143			0.78	0.64	0.94
hCV3188386	S0012(M)	no	ALL	LT60MI_GT75noMI	T	0.51	0.61	0.0071			0.67	0.5	0.89
hCV3188386	S0012(M)	AGE_L T60	Y	MI_LT75noMI	T	0.51	0.61	0.0039			0.66	0.49	0.87
hCV3188386	S0012(M)	no	ALL	MI_GT75noMI	T	0.52	0.61	0.0092			0.69	0.52	0.9
hCV7538986	S0012(M)	AGE-LT60	Y	MI_LT75noMI	A	0.06	0.03	0.0484			2.24	1.07	4.67
hCV7538986	S0012(M)	SMOKE	Y	MI_LT75noMI	A	0.07	0.04	0.0461			1.98	1.04	3.73

TABLE 7

Gene	Marker	Sample Set	p-value	OR*	95% CI	case_freq	control_freq	Allele 1	mode	Strata
GLIO703	hCV1065191	BMS_CARE	0.016835	0.7090	0.53439-0.941	0.6102	0.6883	T	Dom	ALL
GLIO703	hCV1065191	BMS_PRE-CARE	0.011758	0.5837	0.38306-0.889	0.5873	0.7091	T	Dom	BMI TERTILE 3
GLIO703	hCV1065191	BMS_PRE-CARE	0.036996	0.7356	0.5529-0.979	0.3947	0.4700	T	Allelic	GLUCOSE_TERTILE_2
CSF2RB	hCV11486078	BMS_CARE	0.045129	1.4877	1.00667-2.199	0.1535	0.1087	C	Dom	ALL
CSF2RB	hCV11486078	BMS_PRE-CARE	0.042394	1.8670	1.0146-3.435	0.1776	0.1037	C	Dom	GLUCOSE_TERTILE_1
ITGAL	hCV11789692	BMS_CARE	0.047325	0.5905	0.34955-0.997	0.3662	0.4946	G	Dom	BMI_TERTILE_1
ITGAL	hCV11789692	BMS_PRE-CARE	0.012096	0.6411	0.45179-0.91	0.1984	0.2786	G	Allelic	BMI_TERTILE_1
LRP2	hCV16165996	BMS_CARE	0.036433	0.1535	0.02047-1.151	0.0101	0.0623	T	Rec	BMI_TERTILE_3
LRP2	hCV16165996	BMS_CARE	0.011482	0.2448	0.07533-0.796	0.0197	0.0760	T	Rec	FMHX_CHD-0
LRP2	hCV16165996	BMS_CARE	0.041336	0.2476	0.05822-1.053	0.0168	0.0646	T	Rec	HYPERTEN_1
LRP2	hCV16165996	BMS_CARE	0.049657	0.4346	0.1848-1.022	0.0272	0.0603	T	Rec	MALE
LRP2	hCV16165996	BMS_CARE	0.021819	0.4087	0.18573-0.9	0.0276	0.0648	T	Rec	ALL
LRP2	hCV16165996	BMS_PRE-CARE	0.045253	0.3095	0.0923-1.038	0.0233	0.0714	T	Rec	AGE_TERTILE_2
LRP2	hCV16165996	BMS_PRE-CARE	0.038948	0.3702	0.13918-0.985	0.0289	0.0744	T	Rec	AGE_TERTILE_3
LRP2	hCV16165996	BMS_PRE-CARE	0.010388	0.1835	0.04315-0.781	0.0159	0.0808	T	Rec	BMI_TERTILE_2
LRP2	hCV16165996	BMS_PRE-CARE	0.034392	0.3384	0.11833-0.968	0.0204	0.0580	T	Rec	PREVASTATIN
LRP2	hCV16165996	BMS_PRE-CARE	0.028361	0.4124	0.18255-0.932	0.0385	0.0884	T	Rec	PLACEBO
LRP2	hCV16165996	BMS_PRE-CARE	0.003587	0.2734	0.10784-0.693	0.0236	0.0812	T	Rec	FMHX_CHD-0
LRP2	hCV16165996	BMS_PRE-CARE	0.004987	0.0972	0.01308-0.722	0.0073	0.0703	T	Rec	GLUCOSE TERTILE 3
LRP2	hCV16165996	BMS_PRE-CARE	0.026662	0.3545	0.13642-0.921	0.0303	0.0810	T	Rec	HYPERTEN_1
LRP2	hCV16165996	BMS_PRE-CARE	0.036151	0.4065	0.17053-0.969	0.0282	0.0666	T	Rec	HYPERTEN_0
LRP2	hCV16165996	BMS_PRE-CARE	0.006692	0.4177	0.2182-0.8	0.0318	0.0729	T	Rec	MALE
LRP2	hCV16165996	BMS_PRE-CARE	0.002522	0.3843	0.20212-0.731	0.0291	0.0724	T	Rec	ALL
THBS1	hCV16170900	BMS_CARE	0.018783	6.5926	1.08389-40.098	0.0357	0.0056	G	Rec	BMI_TERTILE_2
THBS1	hCV16170900	BMS_PRE-CARE	0.049388	1.5396	1.00678-2.354	0.1480	0.1014	G	Allelic	BMI_TERTILE_2
MSR1	hCV16172249	BMS_CARE	0.022431	15.7338	-.-	0.0143	0.0000	C	Rec	BMI_TERTILE_1
MSR1	hCV16172249	BMS_PRE-CARE	0.026523	1.5093	1.04709-2.175	0.1303	0.0903	C	Dom	ALL
SELL	hCV16172571	BMS_CARE	0.014986	1.8087	1.11763-2.927	0.3434	0.2243	A	Dom	BMI_TERTILE_3
SELL	hCV16172571	BMS_PRE-CARE	0.019386	2.3618	1.12523-4.957	0.0344	0.0149	A	Rec	ALL
ADAM8	hCV16191372	BMS_CARE	0.033461	2.5166	1.0442-6.065	0.0316	0.0128	T	Rec	ALL
ADAM8	hCV16191372	BMS_PRE-CARE	0.001246	10.0833	1.81263-56.092	0.0526	0.0055	T	Rec	AGE_TERTILE_1

(continued)

Gene	Marker	Sample Set	p-value	OR*	95% CI	case_freq	control_freq	Allele 1	mode	Strata
ADAM8	hCV16191372	BMS_PRE-CARE	0.023749	1.5758	1.06064-2.341	0.3212	0.2310	T	Dom	HYPERTEN_1
BIRC5	hCV16266313	BMS_CARE	0.000147	3.4531	1.76512-6.755	0.1910	0.0640	G	Dom	AGE_TERTILE_3
BIRC5	hCV16266313	BMS_CARE	0.007451	2.2955	1.23349-4.272	0.2169	0.1077	G	Dom	BMI_TERTILE_2
BIRC5	hCV16266313	BMS_CARE	0.029598	1.8335	1.05502-3.186	0.1458	0.0852	G	Dom	PLACEBO
BIRC5	hCV16266313	BMS_CARE	0.033243	1.9616	1.04542-3.681	0.1584	0.0876	G	Dom	FMHX_CHD-1
BIRC5	hCV16266313	BMS_CARE	0.038827	2.1011	1.0259-4.303	0.1494	0.0772	G	Dom	GLUCOSE_TERTILE_1
BIRC5	hCV16266313	BMS_CARE	0.005253	2.6022	1.3547-4.998	0.0792	0.0320	G	Allelic	GLUCOSE TERTILE 3
BIRC5	hCV16266313	BMS_CARE	0.027848	1.6231	1.05066-2.507	0.1448	0.0945	G	Dom	MALE
BIRC5	hCV16266313	BMS_CARE	0.026078	1.6010	1.05444-2.431	0.1344	0.0884	G	Dom	ALL
BIRC5	hCV16266313	BMS_PRE-CARE	0.006131	2.0933	1.26582-3.462	0.1085	0.0550	G	Allelic	AGE TERTILE 2
BIRC5	hCV16266313	BMS_PRE-CARE	0.014989	1.9766	1.15433-3.385	0.0952	0.0506	G	Allelic	BMI_TERTILE_1
BIRC5	hCV16266313	BMS_PRE-CARE	0.00485	2.0006	1.22638-3.263	0.1703	0.0931	G	Dom	PLACEBO
BIRC5	hCV16266313	BMS_PRE-CARE	0.021465	1.6815	1.07589-2.628	0.1604	0.1020	G	Dom	FMHX_CHD-0
BIRC5	hCV16266313	BMS_PRE-CARE	0.010177	2.1607	1.18862-3.928	0.1963	0.1015	G	Dom	GLUCOSE_TERTILE_1
BIRC5	hCV16266313	BMS_PRE-CARE	0.03938	1.8321	1.0367-3.238	0.0790	0.0447	G	Allelic	GLUCOSE_TERTILE_2
BIRC5	hCV16266313	BMS_PRE-CARE	0.00124	4.2532	1.68033-10.766	0.3125	0.0966	G	Dom	FEMALE
BIRC5	hCV16266313	BMS_PRE-CARE	0.014904	1.5129	1.08876-2.102	0.0767	0.0521	G	Allelic	ALL
F13A1	hCV1639938	BMS_CARE	0.034558	0.6743	0.47218-0.963	0.2018	0.2727	A	Allelic	PREVASTATIN
F13A1	hCV1639938	BMS_PRE-CARE	0.012496	0.6988	0.52904-0.923	0.2041	0.2684	A	Allelic	PREVASTATIN
PDGFRA	hCV22271841	BMS_CARE	0.001	9.6964	1.86162-50.505	0.0345	0.0037	C	Rec	PLACEBO
PDGFRA	hCV22271841	BMS_CARE	0.020901	3.2843	1.12941-9.551	0.0236	0.0073	C	Rec	ALL
PDGFRA	hCV22271841	BMS_PRE-CARE	0.026695	8.5140	0.87997-82.375	0.0165	0.0020	C	Rec	PLACEBO
PDGFRA	hCV22271841	BMS_PRE-CARE	0.032784	13.8571	-	0.0313	0.0000	C	Rec	FEMALE
PON1	hCV2259750	BMS_CARE	0.040187	2.7282	1.01487-7.334	0.2121	0.0898	T	Rec	FEMALE
PON1	hCV2259750	BMS_PRE-CARE	0.016576	3.0588	1.18628-7.887	0.8125	0.5862	T	Dom	FEMALE
NPC1	hCV25472673	BMS_CARE	0.008339	1.4341	1.10402-1.863	0.4722	0.3842	C	Allelic	PLACEBO
NPC1	hCV25472673	BMS_PRE-CARE	0.032505	1.6625	1.03965-2.659	0.1758	0.1137	C	Rec	PLACEBO
ICAM3	hCV25473653	BMS_CARE	0.029271	0.5809	0.35844-0.942	0.1447	0.2256	C	Allelic	AGE TERTILE 1
ICAM3	hCV25473653	BMS_CARE	0.027136	0.5830	0.36006-0.944	0.2772	0.3968	C	Dom	GLUCOSE TERTILE 3
ICAM3	hCV25473653	BMS_PRE-CARE	0.033872	0.2336	0.05434-1.004	0.0147	0.0601	C	Rec	GLUCOSE_TERTILE_3
ICAM3	hCV25473653	BMS_PRE-CARE	0.002233	0.3106	0.14087-0.685	0.0186	0.0576	C	Rec	ALL
SELL	hCV25474627	BMS_CARE	0.014986	1.8087	1.11763-2.927	0.3434	0.2243	A	Dom	BMI_TERTILE_3

(continued)

Gene	Marker	Sample Set	p-value	OR*	95% CI	case_freq	control_freq	Allele 1	mode	Strata
SELL	hCV25474627	BMS_PRE-CARE	0.018572	2.3750	1.1315-4.985	0.0345	0.0148	A	Rec	ALL
HADHSC	hCV25594697	BMS_CARE	0.035834	0.6369	0.41669-0.973	0.1102	0.1629	C	Dom	ALL
HADHSC	hCV25594697	BMS_PRE-CARE	0.044643	0.5487	0.3033-0.992	0.0904	0.1533	C	Dom	FMHX_CHD-1
LRP3	hCV25594815	BMS_CARE	0.028825	5.9294	0.97529-36.049	0.0341	0.0059	T	Rec	AGE-TERTILE-2
LRP3	hCV25594815	BMS_CARE	0.004599	3.3092	1.37903-7.941	0.0354	0.0110	T	Rec	ALL
LRP3	hCV25594815	BMS_PRE-CARE	0.021685	9.0480	0.93272-87.771	0.0234	0.0027	T	Rec	AGE_TERTILE_2
LRP3	hCV25594815	BMS_PRE-CARE	0.002526	8.1321	1.62452-40.708	0.0364	0.0046	T	Rec	HYPERTEN_1
LRP3	hCV25594815	BMS_PRE-CARE	0.027417	9.6000	0.84307-109.315	0.0625	0.0069	T	Rec	FEMALE
CR1	hCV25596020	BMS_CARE	0.011753	2.9864	1.22737-7.266	0.0756	0.0267	G	Rec	HYPERTEN_1
CR1	hCV25596020	BMS_PRE-CARE	0.049789	2.4429	0.97421-6.126	0.0552	0.0234	G	Rec	HYPERTEN_1
SLC21A3	hCV25605897	BMS_CARE	0.011574	0.3192	0.12582-0.81	0.0459	0.1309	G	Dom	PREVASTATIN
SLC21A3	hCV25605897	BMS_PRE-CARE	0.002728	0.3812	0.19484-0.746	0.0255	0.0643	G	Allelic	PREVASTATIN
SLC21A3	hCV25605897	BMS_PRE-CARE	0.000185	0.4370	0.27647-0.691	0.0291	0.0642	G	Allelic	ALL
TLR2	hCV25607736	BMS_CARE	0.030239	2.3361	1.06404-5.129	0.1409	0.0656	A	Dom	BMI_TERTILE_1
TLR2	hCV25607736	BMS_PRE-CARE	0.00334	20.1963	-.-	0.0080	0.0000	A	Rec	ALL
PROCR	hCV25620145	BMS_CARE	0.048224	7.8372	0.70242-87.443	0.0227	0.0030	G	Rec	AGE_TERTILE_2
PROCR	hCV25620145	BMS_CARE	0.004479	13.0000	1.33617-126.481	0.0333	0.0027	G	Rec	AGE_TERTILE_3
PROCR	hCV25620145	BMS_CARE	0.005918	7.7474	1.39802-42.933	0.0404	0.0054	G	Rec	BMI_TERTILE_3
PROCR	hCV25620145	BMS_CARE	0.048924	4.4027	0.8799-22.029	0.0197	0.0046	G	Rec	FMHX_CHD-0
PROCR	hCV25620145	BMS_CARE	0.007748	11.6020	1.1938-112.756	0.0297	0.0026	G	Rec	GLUCOSE_TERTILE_3
PROCR	hCV25620145	BMS_CARE	0.008012	1.8020	1.18196-2.747	0.1513	0.0900	G	Allelic	HYPERTEN_1
PROCR	hCV25620145	BMS_CARE	0.001821	7.1451	1.69457-30.127	0.0226	0.0032	G	Rec	MALE
PROCR	hCV25620145	BMS_CARE	0.01132	4.3815	1.25884-15.25	0.0197	0.0046	G	Rec	ALL
PROCR	hCV25620145	BMS_PRE-CARE	0.015417	14.8039	-.-	0.0155	0.0000	G	Rec	AGE_TERTILE_2
PROCR	hCV25620145	BMS_PRE-CARE	0.005072	17.9204	-.-	0.0231	0.0000	G	Rec	AGE_TERTILE_3
PROCR	hCV25620145	BMS_PRE-CARE	0.016453	14.5181	-.-	0.0159	0.0000	G	Rec	BMI_TERTILE_3
PROCR	hCV25620145	BMS_PRE-CARE	0.024768	8.7317	0.89991-84.722	0.0238	0.0028	G	Rec	BMI_TERTILE_2
PROCR	hCV25620145	BMS_PRE-CARE	0.001456	14.3503	1.66518-123.668	0.0275	0.0020	G	Rec	PLACEBO
PROCR	hCV25620145	BMS_PRE-CARE	0.002509	13.1366	1.52306-113.306	0.0301	0.0024	G	Rec	FMHX_CHD-1
PROCR	hCV25620145	BMS_PRE-CARE	0.003635	20.0112	-.-	0.0219	0.0000	G	Rec	GLUCOSE TERTILE 3
PROCR	hCV25620145	BMS_PRE-CARE	0.002342	22.0048	-.-	0.0280	0.0000	G	Rec	GLUCOSE_TERTILE_1
PROCR	hCV25620145	BMS_PRE-CARE	0.001181	24.0464	-.-	0.0242	0.0000	G	Rec	HYPERTEN_1

(continued)

Gene	Marker	Sample Set	p-value	OR*	95% CI	case_freq	control_freq	Allele 1	mode	Strata
PROCR	hCV25620145	BMS_PRE-CARE	0.000459	16.4471	1.97291-137.11	0.0173	0.0011	G	Rec	MALE
PROCR	hCV25620145	BMS_PRE-CARE	0.000376	10.1509	2.09945-49.08	0.0185	0.0019	G	Rec	ALL
TAP1	hCV25630686	BMS_CARE	0.04021	12.6872	..	0.0111	0.0000	T	Rec	AGE_TERTILE_3
TAP1	hCV25630686	BMS_CARE	0.000322	30.4354	..	0.0118	0.0000	T	Rec	ALL
TAP1	hCV25630686	BMS_PRE-CARE	0.018347	2.6736	1.14742-6.23	0.0756	0.0297	T	Dom	AGE_TERTILE_3
LRP2	hCV25646248	BMS_CARE	0.024691	5.3000	1.04828-26.796	0.0411	0.0080	T	Dom	AGE_TERTILE_1
LRP2	hCV25646248	BMS_PRE-CARE	0.022547	3.6575	1.25034-10.699	0.0395	0.0111	T	Allelic	AGE_TERTILE_1
HLA-DPB1	hCV25651174	BMS_CARE	0.029542	1.4094	1.03931-1.911	0.3670	0.2914	G	Allelic	PREVASTATIN
HLA-DPB1	hCV25651174	BMS_PRE-CARE	0.036904	1.4164	1.02066-1.965	0.5714	0.4849	G	Dom	PREVASTATIN
NEUROD1	hCV25651593	BMS_CARE	0.046449	0.1212	..	0.0000	0.0500	T	Dom	AGE_TERTILE_1
NEUROD1	hCV25651593	BMS_PRE-CARE	0.04905	0.5689	0.32222-1.004	0.0397	0.0677	T	Dom	ALL
PON2	hCV2630153	BMS_CARE	0.023524	1.5374	1.05786-2.234	0.4925	0.3870	C	Dom	HYPERTEN_0
PON2	hCV2630153	BMS_PRE-CARE	0.0491	1.2936	1.00802-1.66	0.2747	0.2264	C	Allelic	HYPERTEN_0
ABCA1	hCV2741104	BMS_CARE	0.032615	0.6617	0.45235-0.968	0.3517	0.4506	T	Dom	PLACEBO
ABCA1	hCV2741104	BMS_CARE	0.019416	0.5838	0.37069-0.919	0.3564	0.4868	T	Dom	GLUCOSE_TERTILE_3
ABCA1	hCV2741104	BMS_PRE-CARE	0.000279	0.5035	0.34649-0.732	0.2637	0.4157	T	Dom	PLACEBO
ABCA1	hCV2741104	BMS_PRE-CARE	0.02491	0.7932	0.64814-0.971	0.2050	0.2454	T	Allelic	ALL
PECAM1	hCV435368	BMS_CARE	0.007531	1.6056	1.14496-2.252	0.4464	0.4358	A	Allelic	BMI_TERTILE_2
PECAM1	hCV435368	BMS_CARE	0.014604	1.7845	1.11621-2.853	0.2844	0.1822	A	Rec	PREVASTATIN
PECAM1	hCV435368	BMS_CARE	0.006582	1.5923	1.13818-2.228	0.4318	0.4525	A	Allelic	GLUCOSE_TERTILE_1
PECAM1	hCV435368	BMS_CARE	0.007332	1.4384	1.10501-1.872	0.4482	0.4612	A	Allelic	HYPERTEN_0
PECAM1	hCV435368	BMS_CARE	0.022746	1.2736	1.03468-1.568	0.4774	0.4622	A	Allelic	MALE
PECAM1	hCV435368	BMS_CARE	0.017633	1.2686	1.04533-1.54	0.4822	0.4584	A	Allelic	ALL
PECAM1	hCV435368	BMS_PRE-CARE	0.000689	1.6513	1.23632-2.206	0.4600	0.4155	A	Allelic	BMI_TERTILE_3
PECAM1	hCV435368	BMS_PRE-CARE	0.027008	1.3129	1.03237-1.67	0.5000	0.4324	A	Allelic	PLACEBO
PECAM1	hCV435368	BMS_PRE-CARE	0.0286	1.2842	1.03084-1.6	0.4858	0.4518	A	Allelic	FMHX_CHD-0
PECAM1	hCV435368	BMS_PRE-CARE	0.000544	1.6350	1.23863-2.158	0.4453	0.4325	A	Allelic	GLUCOSE_TERTILE_3
PECAM1	hCV435368	BMS_PRE-CARE	0.011773	1.3314	1.06914-1.658	0.4742	0.4544	A	Allelic	HYPERTEN_0
PECAM1	hCV435368	BMS_PRE-CARE	0.010962	1.2560	1.05449-1.496	0.4913	0.4519	A	Allelic	MALE
PECAM1	hCV435368	BMS_PRE-CARE	0.013871	1.2353	1.04629-1.458	0.4974	0.4500	A	Allelic	ALL
A2M	hCV517658	BMS_CARE	0.028082	1.7449	1.05782-2.878	0.6627	0.5296	G	Dom	BMI_TERTILE_2
A2M	hCV517658	BMS_CARE	0.027305	1.3755	1.03564-1.827	0.6375	0.5611	G	Dom	ALL

(continued)

Gene	Marker	Sample Set	p-value	OR*	95% CI	case_freq	control_freq	Allele 1	mode	Strata
A2M	hCV517658	BMS_PRE-CARE	0.007777	1.7717	1.15969-2.707	0.6532	0.5153	G	Dom	BMI_TERTILE_2
A2M	hCV517658	BMS_PRE-CARE	0.026334	1.7745	1.06464-2.958	0.1718	0.1047	G	Rec	HYPERTEN_1
ADAMTS1	hCV529706	BMS_CARE	0.029249	1.6701	1.05032-2.655	0.5667	0.4392	C	Dom	AGE_TERTILE_3
ADAMTS1	hCV529706	BMS_PRE-CARE	0.02017	1.6853	1.08049-2.629	0.0878	0.0540	C	Rec	ALL
ADAMTS1	hCV529710	BMS_CARE	0.029249	1.6701	1.05032-2.655	0.5667	0.4392	C	Dom	AGE_TERTILE_3
ADAMTS1	hCV529710	BMS_PRE-CARE	0.016984	1.7117	1.09605-2.673	0.0873	0.0529	C	Rec	ALL
HLA-G	hCV7482175	BMS_CARE	0.018628	0.5823	0.3697-0.917	0.2015	0.3023	A	Dom	HYPERTEN_0
HLA-G	hCV7482175	BMS_PRE-CARE	0.037108	0.1101	.-.	0.0000	0.0201	A	Rec	HYPERTEN_0
LYZ	hCV7559757	BMS_CARE	0.024609	15.2416	.-.	0.0133	0.0000	A	Rec	AGE_TERTILE_1
LYZ	hCV7559757	BMS_CARE	0.037787	12.9802	.-.	0.0066	0.0000	A	Rec	FMHX_CHD-0
LYZ	hCV7559757	BMS_CARE	0.003322	21.6600	.-.	0.0079	0.0000	A	Rec	ALL
LYZ	hCV7559757	BMS_PRE-CARE	0.022269	9.9178	0.88758-110.821	0.0267	0.0028	A	Rec	AGE_TERTILE_1
FABP2	hCV761961	BMS_CARE	0.036056	0.5833	0.35113-0.969	0.3684	0.5000	T	Dom	AGE_TERTILE_1
FABP2	hCV761961	BMS_PRE-CARE	0.036725	0.5833	0.35052-0.971	0.3684	0.5000	T	Dom	AGE_TERTILE_1
MMRN	hCV8933098	BMS_CARE	0.04082	1.9109	1.01944-3.582	0.1954	0.1128	G	Dom	AGE_TERTILE_2
MMRN	hCV8933098	BMS_PRE-CARE	0.009857	2.0313	1.17676-3.506	0.1938	0.1058	G	Dom	AGE_TERTILE_2
PON2	hCV8952817	BMS_CARE	0.024507	1.5312	1.05461-2.223	0.4889	0.3845	C	Dom	HYPERTEN_0
PON2	hCV8952817	BMS_PRE-CARE	0.04204	1.2993	1.0124-1.668	0.2747	0.2257	C	Allelic	HYPERTEN_0
MC3R	hCV9485713	BMS_CARE	0.016123	1.9962	1.12843-3.531	0.2529	0.1450	A	Dom	AGE_TERTILE_2
MC3R	hCV9485713	BMS_CARE	0.020971	1.4944	1.06081-2.105	0.2143	0.1543	A	Dom	ALL
MC3R	hCV9485713	BMS_PRE-CARE	0.014872	1.9047	1.12737-3.218	0.2093	0.1220	A	Dom	AGE_TERTILE_2

* Haldane OR was used if either case count or the control count is zero

Certain Embodiments:**[0380]**

- 5 1. A method for identifying an individual who has an altered risk for developing myocardial infarction, comprising detecting a single nucleotide polymorphism (SNP) in any one of the nucleotide sequences of SEQ ID NOS:1-828 and 1657-73,085 in said individual's nucleic acids, wherein the presence of the SNP is correlated with an altered risk for myocardial infarction in said individual.
- 10 2. The method of item 1 in which the altered risk is an increased risk.
3. The method of item 2 in which said individual has previously had a myocardial infarction.
- 15 4. The method of item 1 in which the altered risk is a decreased risk.
5. The method of item 1, wherein the SNP is selected from the group consisting of the SNPs set forth in Tables 6-7.
- 20 6. The method of item 1 in which detection is carried out by a process selected from the group consisting of: allele-specific probe hybridization, allele-specific primer extension, allele-specific amplification, sequencing, 5' nuclease digestion, molecular beacon assay, oligonucleotide ligation assay, size analysis, and single-stranded conformation polymorphism.
- 25 7. An isolated nucleic acid molecule comprising at least 8 contiguous nucleotides wherein one of the nucleotides is a single nucleotide polymorphism (SNP) selected from any one of the nucleotide sequences in SEQ ID NOS:1-828 and 1657-73,085, or a complement thereof.
8. The isolated nucleic acid molecule of item 7, wherein the SNP is selected from the group consisting of the SNPs set forth in Tables 3 and 4.
- 30 9. An isolated nucleic acid molecule that encodes any one of the amino acid sequences in SEQ ID NOS:829-1656.
10. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:829-1656.
- 35 11. An antibody that specifically binds to a polypeptide of item 10, or an antigen-binding fragment thereof.
12. The antibody of item 11 in which the antibody is a monoclonal antibody.
- 40 13. An amplified polynucleotide containing a single nucleotide polymorphism (SNP) selected from any one of the nucleotide sequences of SEQ ID NOS: 1-828 and 1657-73,085, or a complement thereof, wherein the amplified polynucleotide is between about 16 and about 1,000 nucleotides in length.
14. The amplified polynucleotide of item 13 in which the nucleotide sequence comprises any one of the nucleotide sequences of SEQ ID NOS:1-828 and 1657-73,085.
- 45 15. An isolated polynucleotide which specifically hybridizes to a nucleic acid molecule containing a single nucleotide polymorphism (SNP) in any one of the nucleotide sequences in SEQ ID NOS:1-828 and 1657-73,085.
16. The polynucleotide of item 15 which is 8-70 nucleotides in length.
- 50 17. The polynucleotide of item 15 which is an allele-specific probe.
18. The polynucleotide of item 15 which is an allele-specific primer.
- 55 19. The polynucleotide of item 15, wherein the polynucleotide comprises a nucleotide sequence selected from the group consisting of the primer sequences set forth in Table 5 (SEQ ID NOS:73,086-73,997).
20. A kit for detecting a single nucleotide polymorphism (SNP) in a nucleic acid, comprising the polynucleotide of

item 15, a buffer, and an enzyme.

21. A method of detecting a single nucleotide polymorphism (SNP) in a nucleic acid molecule, comprising contacting a test sample with a reagent which specifically hybridizes to a SNP in any one of the nucleotide sequences of SEQ ID NOS: 1-828 and 1657-73,085 under stringent hybridization conditions, and detecting the formation of a hybridized duplex.

22. The method of item 21 in which detection is carried out by a process selected from the group consisting of: allele-specific probe hybridization, allele-specific primer extension, allele-specific amplification, sequencing, 5' nuclease digestion, molecular beacon assay, oligonucleotide ligation assay, size analysis, and single-stranded conformation polymorphism.

23. A method of detecting a variant polypeptide, comprising contacting a reagent with a variant polypeptide encoded by a single nucleotide polymorphism (SNP) in any one of the nucleotide sequences of SEQ ID NOS: 1-828 and 1657-73,085 in a test sample, and detecting the binding of the reagent to the polypeptide.

24. A method for identifying an agent useful in therapeutically or prophylactically treating myocardial infarction, comprising contacting the polypeptide of item 10 with a candidate agent under conditions suitable to allow formation of a binding complex between the polypeptide and the candidate agent, and detecting the formation of the binding complex, wherein the presence of the complex identifies said agent.

Claims

1. An in vitro method for identifying an individual having an altered risk for developing myocardial infarction, comprising detecting the single nucleotide polymorphism (SNP) in the nucleotide sequence of SEQ ID NO: 19038 in said individual's nucleic acids, wherein said individual has an increased risk for developing myocardial infarction due to the presence of C at position 101 of SEQ ID NO: 19038 or the presence of G at position 101 of its complement.
2. The method of claim 1, wherein said nucleic acid has been isolated from cells of a biological sample from said human.
3. The method of claim 2, wherein said biological sample is blood, saliva, or buccal swabs.
4. The method of claim 2, further comprising isolating said nucleic acid from said biological sample prior to said detecting step.
5. The method of claim 1, wherein said detecting step comprises nucleic acid amplification, preferably carried out by polymerase chain reaction.
6. The method of any one of claims 1-5, comprising contacting the nucleic acids of the individual with a detection reagent, and determining which nucleotide is present or absent at position 101 of SEQ ID NO: 19038.
7. The method of claim 6, in which the detection reagent is a polynucleotide probe.
8. The method of claim 7, in which the 3' end of the probe or the center of the probe hybridizes to the SNP in the nucleic acid.
9. The method of claim 7, in which the probe is labeled with a detectable substance, which is preferably a reporter dye.
10. The method of claim 1, wherein said detecting step is performed using allele-specific probe hybridization, allele-specific primer extension, allele-specific amplification, sequencing, 5' nuclease digestion, molecular beacon assay, oligonucleotide ligation assay, size analysis, single-stranded conformation polymorphism analysis, or denaturing gradient gel electrophoresis (DGGE).
11. The method of claim 1, wherein said detection step is carried out using an allele-specific probe or primer comprising SEQ ID NO: 73557, SEQ ID NO: 73558 or SEQ ID NO: 73559.
12. The method of claim 1, wherein said method detects said C or said G.

13. The method of claim 1, wherein said human is homozygous or heterozygous for said C or said G.
14. Use of an isolated nucleic acid molecule comprising at least 8 contiguous nucleotides of SEQ ID NO: 19038, wherein said at least 8 contiguous nucleotides of SEQ ID NO: 19038 include position 101 of SEQ ID NO: 19038, or the complement thereof, in the method according to any one of claims 1-13.
15. Use of an isolated nucleic acid molecule that encodes any one of the amino acid sequences in SEQ ID NOS: 845, 846 in the method according to any one of claims 1-13.
16. Use of an amplified polynucleotide containing the single nucleotide polymorphism (SNP) at position 101 of SEQ ID NO: 19038, or the complement thereof, in the method according to any one of claims 1-13, wherein the amplified polynucleotide is between about 16 and about 1,000 nucleotides in length.
17. Use of claim 16 in which the amplified polynucleotide comprises the nucleotide sequence of SEQ ID NO: 19038.
18. Use of an isolated polynucleotide which specifically hybridizes to a nucleic acid molecule containing the single nucleotide polymorphism (SNP) at position 101 of SEQ ID NO: 19038, or the complement thereof, in the method according to any one of claims 1-13.
19. Use of claim 18, wherein the isolated polynucleotide is 8-70 nucleotides in length or wherein the isolated polynucleotide is an allele-specific probe or an allele-specific primer.
20. Use of claim 18, wherein the isolated polynucleotide comprises a nucleotide sequence selected from the group consisting of the primer sequences set forth in Table 5 (SEQ ID NOS: 73557-73559).
21. Use of a test kit comprising the polynucleotide as defined in any one of claims 14-20, a buffer, and an enzyme in the method according to any one of claims 1-13.
22. An in vitro method of identifying an individual having an altered risk for developing myocardial infarction, comprising detecting the presence of proline (P) or serine (S) at position 157 of the amino acid sequence of SEQ ID NOS:845 or 846, wherein the individual has an increased risk for developing myocardial infarction due to the presence of P at position 157 of the amino acid sequence of SEQ ID NOS:845 or 846.

Patentansprüche

1. In vitro-Verfahren zur Identifizierung eines Individuums, das ein verändertes Risiko besitzt, Myokardinfarkt (MI) zu entwickeln, umfassend das Detektieren des Einzelnukleotidpolymorphismus (Single Nucleotide Polymorphism, SNP) in der Nukleotidsequenz von SEQ ID NO:19038 in den Nukleinsäuren des Individuums, wobei das Individuum aufgrund des Vorliegens von C an Position 101 von SEQ ID NO:19038 oder des Vorliegens von G an Position 101 seiner komplementären Sequenz ein erhöhtes Risiko hat, Myokardinfarkt zu entwickeln.
2. Verfahren nach Anspruch 1, wobei die Nukleinsäure von Zellen einer biologischen Probe des Menschen isoliert wurde.
3. Verfahren nach Anspruch 2, wobei die biologische Probe Blut, Speichel, oder Wangenabstrich ist.
4. Verfahren nach Anspruch 2, ferner umfassend die Isolierung der Nukleinsäure von der biologischen Probe vor dem Detektierungsschritt.
5. Verfahren nach Anspruch 1, bei dem der Detektierungsschritt Nukleinsäureamplifikation umfasst, vorzugsweise durchgeführt mittels Polymerase-Kettenreaktion (polymerase chain reaction).
6. Verfahren nach irgend einem der Ansprüche 1-5, das das Inkontaktbringen der Nukleinsäuren des Individuums mit einem Detektionsreagenz und die Bestimmung, welches Nukleotid an Position 101 von SEQ ID NO:19038 vorliegt oder nicht vorliegt, umfasst.
7. Verfahren nach Anspruch 6, bei dem das Detektionsreagenz eine Polynukleotidsonde ist.

8. Verfahren nach Anspruch 7, bei dem das 3' Ende der Sonde oder die Mitte der Sonde mit dem SNP in der Nukleinsäure hybridisiert.
9. Verfahren nach Anspruch 7, bei dem die Sonde mit einer nachweisbaren Substanz, welche vorzugsweise ein Reporterfarbstoff ist, markiert ist.
10. Verfahren nach Anspruch 1, wobei der Detektierungsschritt unter Verwendung von Allel-spezifischer Sondenhybridisierung, Allel-spezifischer Primerextension, Allel-spezifischer Amplifikation, Sequenzierung, 5'-Nuklease-Verdau, Molecular-Beacon-Assay, Oligonukleotid-Ligationsassay, Größenanalyse, Einzelstrang-Konformationspolymorphismus-Analyse oder denaturierender Gradientengelelektrophorese (DGGE) durchgeführt wird.
11. Verfahren nach Anspruch 1, wobei der Detektierungsschritt unter Verwendung einer Allel-spezifischen Sonde oder eines Primers, umfassend SEQ ID NO:73557, SEQ ID NO:73558 oder SEQ ID NO:73559, durchgeführt wird.
12. Verfahren nach Anspruch 1, wobei das Verfahren das C oder das G detektiert.
13. Verfahren nach Anspruch 1, wobei der Mensch homozygot oder heterozygot für das C oder das G ist.
14. Verwendung eines isolierten Nukleinsäuremoleküls, umfassend mindestens 8 kontinuierliche Nukleotide von SEQ ID NO:19038, wobei die mindestens 8 kontinuierlichen Nukleotide von SEQ ID NO:19038 die Position 101 von SEQ ID NO:1903 einschließen, oder des Komplements davon, in dem Verfahren nach irgend einem der Ansprüche 1-13.
15. Verwendung eines isolierten Nukleinsäuremoleküls, das irgend eine der Aminosäuresequenzen in SEQ ID NOS:845, 846 kodiert, in dem Verfahren nach irgend einem der Ansprüche 1-13.
16. Verwendung eines amplifizierten Polynukleotids, das den Einzelnukleotidpolymorphismus (SNP) an Position 101 von SEQ ID NO:19038 enthält, oder des Komplements davon, in dem Verfahren nach irgend einem der Ansprüche 1-13, wobei das amplifizierte Polynukleotid zwischen etwa 16 und etwa 1000 Nukleotide lang ist.
17. Verwendung nach Anspruch 16, bei der das amplifizierte Polynukleotid die Nukleotidsequenz von SEQ ID NO:1903 umfasst.
18. Verwendung eines isolierten Polynukleotids, das spezifisch mit einem Nukleinsäuremolekül, das einen Einzelnukleotidpolymorphismus (SNP) an Position 101 von SEQ ID NO:19038 enthält, oder dem Komplement davon hybridisiert, in dem Verfahren nach irgend einem der Ansprüche 1-13.
19. Verwendung nach Anspruch 18, bei der das isolierte Polynukleotid 8-70 Nukleotide lang ist oder bei der das isolierte Polynukleotid eine Allel-spezifische Sonde oder ein Allel-spezifischer Primer ist.
20. Verwendung nach Anspruch 18, bei der das isolierte Polynukleotid eine Nukleotidsequenz umfasst, die aus der Gruppe der in Tabelle 5 (SEQ ID NOS:73557-73559) aufgeführten Primer-Sequenzen ausgewählt ist.
21. Verwendung eines Test-Kits, umfassend das Polynukleotid, wie es in irgend einem der Ansprüche 14-20 definiert ist, einen Puffer und ein Enzym, in dem Verfahren nach irgend einem der Ansprüche 1-13.
22. In vitro-Verfahren zur Identifizierung eines Individuums, das ein verändertes Risiko besitzt, Myokardinfarkt zu entwickeln, umfassend das Detektieren des Vorliegens von Prolin (P) oder Serin (S) an Position 157 der Aminosäuresequenz von SEQ ID NOS:845 oder 846, wobei das Individuum aufgrund des Vorliegens von P an Position 157 der Aminosäuresequenz von SEQ ID NOS:845 oder 846 ein erhöhtes Risiko hat, Myokardinfarkt zu entwickeln.

Revendications

1. Procédé *in vitro* d'identification d'un individu qui a un risque modifié d'avoir un infarctus du myocarde, comprenant la détection du polymorphisme d'un seul nucléotide (SNP) dans la séquence de nucléotides de la SEQ ID NO: 19038 dans les acides nucléiques dudit individu, procédé dans lequel ledit individu a un risque accru d'avoir un infarctus du myocarde en raison de la présence de C à la position 101 de la SEQ ID NO: 19038 ou de la présence de G à la position 101 de son complément.

2. Procédé selon la revendication 1, dans lequel ledit acide nucléique a été isolé à partir de cellules d'un échantillon biologique provenant dudit humain.
- 5 3. Procédé selon la revendication 2, dans lequel ledit échantillon biologique est du sang, de la salive ou des prélèvements buccaux.
4. Procédé selon la revendication 2, comprenant en outre l'isolement dudit acide nucléique à partir dudit échantillon biologique avant ladite étape de détection.
- 10 5. Procédé selon la revendication 1, dans lequel ladite étape de détection comprend une amplification de l'acide nucléique, effectuée de préférence au moyen d'une réaction d'amplification en chaîne par polymérase.
6. Procédé selon l'une quelconque des revendications 1 à 5, comprenant la mise en contact des acides nucléiques de l'individu avec un réactif de détection, et la détermination du nucléotide qui est présent ou absent à la position
15 101 de la SEQ ID NO: 19038.
7. Procédé selon la revendication 6, dans lequel l'agent de détection est une sonde de polynucléotide.
8. Procédé selon la revendication 7, dans lequel l'extrémité 3' de la sonde ou le centre de la sonde s'hybride au SNP
20 dans l'acide nucléique.
9. Procédé selon la revendication 7, dans lequel la sonde est marquée avec une substance décelable, qui est de préférence un colorant rapporteur.
- 25 10. Procédé selon la revendication 1, dans lequel ladite étape de détection est effectuée en utilisant une hybridation de sonde spécifique d'un allèle, une extension d'amorce spécifique d'un allèle, une amplification spécifique d'un allèle, un séquençage, une digestion par une 5'-nucléase, une analyse avec balise moléculaire, un test de ligature d'oligonucléotide, une analyse de taille, une analyse de polymorphisme de conformation en simple brin ou une électrophorèse sur gel en gradient dénaturant (DGGE).
- 30 11. Procédé selon la revendication 1, dans lequel ladite étape de détection est effectuée en utilisant une sonde ou une amorce spécifique d'un allèle comprenant la SEQ ID NO: 73557, la SEQ ID NO: 73558 ou la SEQ ID NO: 73559.
12. Procédé selon la revendication 1, ledit procédé détectant ledit C ou ledit G.
- 35 13. Procédé selon la revendication 1, dans lequel ledit humain est homozygote ou hétérozygote pour ledit C ou ledit G.
14. Utilisation d'une molécule d'acide nucléique isolé comprenant au moins 8 nucléotides contigus de la SEQ ID NO: 19038, dans laquelle lesdits au moins 8 nucléotides contigus de la SEQ ID NO: 19038 incluent la position 101 de
40 la SEQ ID NO: 19038, ou du complément de celle-ci, dans le procédé selon l'une quelconque des revendications 1 à 13.
15. Utilisation d'une molécule d'acide nucléique isolé qui code pour l'une quelconque des séquences d'acides aminés dans les SEQ ID NOS: 845, 846 dans le procédé selon l'une quelconque des revendications 1 à 13.
- 45 16. Utilisation d'un polynucléotide amplifié contenant le polymorphisme d'un seul nucléotide (SNP) à la position 101 de la SEQ ID NO: 19038, ou du complément de celle-ci, dans le procédé selon l'une quelconque des revendications 1 à 13, dans laquelle le nucléotide amplifié a une longueur comprise entre environ 16 et environ 1 000 nucléotides.
- 50 17. Utilisation selon la revendication 16 dans laquelle le polynucléotide amplifié comprend la séquence de nucléotides de la SEQ ID NO: 19038.
18. Utilisation d'un polynucléotide isolé qui s'hybride spécifiquement à une molécule d'acide nucléique contenant le polymorphisme d'un seul nucléotide (SNP) à la position 101 de la SEQ ID NO: 19038, ou du complément de celle-ci,
55 dans le procédé selon l'une quelconque des revendications 1 à 13.
19. Utilisation selon la revendication 18, dans laquelle le polynucléotide isolé a une longueur de 8 à 70 nucléotides ou dans laquelle le polynucléotide isolé est une sonde spécifique d'un allèle ou une amorce spécifique d'un allèle.

EP 2 474 631 B1

20. Utilisation selon la revendication 18, dans laquelle le polynucléotide isolé comprend une séquence de nucléotides choisie dans le groupe constitué par les séquences d'amorce présentées dans le Tableau 5 (SEQ ID NOS: 73557 à 73559).

5 **21.** Utilisation d'une trousse d'essai comprenant le polynucléotide tel que défini dans l'une quelconque des revendications 14 à 20, un tampon, et une enzyme dans le procédé selon l'une quelconque des revendications 1 à 13.

10 **22.** Procédé *in vitro* d'identification d'un individu ayant un risque modifié d'avoir un infarctus du myocarde, comprenant la détection de la présence d'une proline (P) ou d'une sérine (S) à la position 157 de la séquence d'acides aminés des SEQ ID NOS: 845 ou 846, dans lequel ledit individu a un risque accru d'avoir un infarctus du myocarde en raison de la présence de P à la position 157 de la séquence d'acides aminés des SEQ ID NOS: 845 ou 846.

15

20

25

30

35

40

45

50

55




FIGURE 1

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9740462 A [0024]
- EP 0609059 A2 [0025]
- US 4683195 A [0081]
- US 4683202 A [0081]
- US 5270184 A [0081]
- US 5422252 A [0081]
- US 5399491 A [0081]
- US 6027923 A [0081]
- US 5539082 A [0091]
- US 5527675 A [0091]
- US 5623049 A [0091]
- US 5714331 A [0091]
- WO 9604000 A [0092]
- US 4835263 A [0093]
- US 5801115 A [0093]
- EP 235726 A, Dattagupta [0107]
- WO 8911548 A, Saiki [0107]
- US 4988617 A [0111] [0156]
- US 4458066 A [0112]
- WO 9322456 A [0114]
- US 5210015 A [0117] [0153]
- US 5538848 A [0117] [0153]
- US 5118801 A [0117] [0155]
- US 5312728 A [0117] [0155]
- US 5866336 A [0117] [0155]
- US 6117635 A [0117] [0155]
- US 5837832 A [0123]
- WO 9511995 A, Chee [0123]
- US 5807522 A, Brown [0123]
- US 620332 A [0128]
- US 10620333 A [0128]
- US 6124478 A [0128]
- US 6107024 A [0128]
- US 5994073 A [0128]
- US 5981768 A [0128]
- US 5871938 A [0128]
- US 5843681 A [0128]
- US 5800999 A [0128]
- US 5773628 A [0128]
- US 20020110828 A [0128]
- WO 95251116 A, Baldeschweiler [0130]
- US 5589136 A [0134]
- US 6153073 A, Dubrow [0134]
- US 6156181 A, Parce [0134]
- US 4988167 A [0151]
- US 6027889 A [0157]
- US 6268148 B [0157]
- US 5494810 A [0157]
- US 5830711 A [0157]
- US 6054564 A [0157]
- WO 9731256 A [0157]
- WO 0056927 A [0157]
- US 0117329 A [0157]
- US 09584905 B [0157]
- US 427818 P [0157]
- US 60445636 B [0157]
- US 60445494 B [0157]
- WO 9416101 A [0162]
- US 5498531 A [0164]
- US 4107288 A [0227]
- US 5145684 A [0227]
- US 6119096 A [0247]
- US 5283317 A [0297]
- WO 9410300 A, Brent [0297]
- US 4816567 A [0300]
- US 4816397 A [0300]
- US 5693762 A [0300]
- US 5585089 A [0300]
- US 5565332 A [0300]
- US 4946778 A [0300]
- US 5637677 A [0302]
- US 4736866 A [0346]
- US 4870009 A, Leder [0346]
- US 4873191 A, Wagner [0346]
- WO 9707668 A [0348]
- WO 9707669 A [0348]

Non-patent literature cited in the description

- LLOYD-JONES DM. *Lancet*, 1999, vol. 353, 89-92 [0006]
- WILLIAMS R R. *Am J Cardiology*, 2001, vol. 87, 129 [0010]
- BROECKEL U. *Nature Genetics*, 2002, vol. 30, 210 [0010]
- HARRAP S. *Arterioscler Thromb Vasc Biol*, 2002, vol. 22, 874-878 [0010]
- SHEARMAN A. *Human Molecular Genetics*, 2000, vol. 9 (9), 1315-1320 [0010]
- GUSELLA. *Ann. Rev. Biochem.*, 1986, vol. 55, 831-854 [0014]

- **WEBER et al.** Human diallelic insertion/deletion polymorphisms. *Am J Hum Genet*, October 2002, vol. 71 (4), 854-62 [0016]
- **STEPHENS et al.** *Science*, 20 July 2001, vol. 293, 489-493 [0019]
- **LINDER et al.** *Clinical Chemistry*, 1997, vol. 43, 254 [0024]
- **MARSHALL.** *Nature Biotechnology*, 1997, vol. 15, 1249 [0024]
- **SCHAFER et al.** *Nature Biotechnology*, 1998, vol. 16, 3 [0024]
- PCR Technology: Principles and Applications for DNA Amplification. Freeman Press [0081]
- **WU ; WALLACE.** *Genomics*, 1989, vol. 4, 560 [0081]
- **LANDEGREN et al.** *Science*, 1988, vol. 241, 1077 [0081]
- **GUATELLI et al.** *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 1874 [0081]
- **SAMBROOK ; RUSSELL.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press, 2000 [0088] [0091] [0138]
- **COREY.** Peptide nucleic acids: expanding the scope of nucleic acid recognition. *Trends Biotechnol.*, June 1997, vol. 15 (6), 224-9 [0091]
- **HYRUP et al.** Peptide nucleic acids (PNA): synthesis, properties and potential applications. *Bioorg Med Chem.*, January 1996, vol. 4 (1), 5-23 [0091]
- **LAGRIFFOUL et al.** *Bioorganic & Medicinal Chemistry Letters*, 1994, vol. 4, 1081-1082 [0092]
- **PETERSEN et al.** *Bioorganic & Medicinal Chemistry Letters*, 1996, vol. 6, 793-796 [0092]
- **KUMAR et al.** *Organic Letters*, 2001, vol. 3 (9), 1269-1272 [0092]
- Current Protocols in Nucleic Acid Chemistry. John Wiley & Sons, 2002 [0093]
- Computational Molecular Biology. Oxford University Press, 1988 [0097]
- Biocomputing: Informatics and Genome Projects. Academic Press, 1993 [0097]
- Computer Analysis of Sequence Data. Humana Press [0097]
- **VON HEINJE, G.** Sequence Analysis in Molecular Biology. Academic Press, 1987 [0097]
- Sequence Analysis Primer. M Stockton Press, 1991 [0097]
- *J. Mol. Biol.*, 1970, 444-453 [0097]
- **DEVEREUX, J. et al.** *Nucleic Acids Res.*, 1984, vol. 12 (1), 387 [0098]
- **E. MYERS ; W. MILLER.** *CABIOS*, 1989, vol. 4, 11-17 [0098]
- **ALTSCHUL et al.** *J. Mol. Biol.*, 1990, vol. 215, 403-10 [0099]
- **ALTSCHUL et al.** *Nucleic Acids Res.*, 1997, vol. 25 (17), 3389-3402 [0099]
- **PEARSON.** *Methods Mol. Biol.*, 1994, vol. 25, 365-389 [0099]
- **DUFRESNE et al.** *Nat Biotechnol*, December 2002, vol. 20 (12), 1269-71 [0099]
- Current Protocols in Bioinformatics. John Wiley & Sons, Inc., [0099]
- Mutation Detection A Practical Approach. Oxford University Press, 1998 [0107]
- **SAIKI et al.** *Nature*, 1986, vol. 324, 163-166 [0107]
- **NARANG et al.** *Methods in Enzymology*, 1979, vol. 68, 90 [0112]
- **BROWN et al.** *Methods in Enzymology*, 1979, vol. 68, 109 [0112]
- **BEAUCAGE et al.** *Tetrahedron Letters*, 1981, vol. 22, 1859 [0112]
- **GIBBS.** *Nucleic Acid Res.*, 1989, vol. 17, 2427-2448 [0114]
- **LIVAK et al.** *PCR Method Appl.*, 1995, vol. 4, 357-362 [0117]
- **TYAGI et al.** *Nature Biotechnology*, 1996, vol. 14, 303-308 [0117]
- **NAZARENKO et al.** *Nucl. Acids Res.*, 1997, vol. 25, 2516-2521 [0117]
- **LOCKHART, D. J. et al.** *Nat. Biotech.*, 1996, vol. 14, 1675-1680 [0123]
- **SCHENA, M. et al.** *Proc. Natl. Acad. Sci.*, 1996, vol. 93, 10614-10619 [0123]
- **ZAMMATTEO et al.** New chips for molecular biology and diagnostics. *Biotechnol Annu Rev.*, 2002, vol. 8, 85-101 [0124]
- **SOSNOWSKI et al.** Active microelectronic array system for DNA hybridization, genotyping and pharmacogenomic applications. *Psychiatr Genet.*, December 2002, vol. 12 (4), 181-92 [0124]
- **HELLER.** DNA microarray technology: devices, systems, and applications. *Annu Rev Biomed Eng.*, 22 March 2002, vol. 4, 129-53 [0124]
- **KOLCHINSKY et al.** Analysis of SNPs and other genomic variations using gel-based chips. *Hum Mutat.*, April 2002, vol. 19 (4), 343-60 [0124]
- **MCGALL et al.** High-density genechip oligonucleotide probe arrays. *Adv Biochem Eng Biotechnol.*, 2002, vol. 77, 21-42 [0124]
- Current Protocols in Molecular Biology. John Wiley & Sons, 1989, 6.3.1-6.3.6 [0127]
- **WEIGL et al.** Lab-on-a-chip for drug development. *Adv Drug Deliv Rev.*, 24 February 2003, vol. 55 (3), 349-77 [0133]
- **CHEN et al.** Single nucleotide polymorphism genotyping: biochemistry, protocol, cost and throughput. *Pharmacogenomics J.*, 2003, vol. 3 (2), 77-96 [0151]
- **KWOK et al.** Detection of single nucleotide polymorphisms. *Curr Issues Mol Biol.*, April 2003, vol. 5 (2), 43-60 [0151]
- **SHI.** Technologies for individual genotyping: detection of genetic polymorphisms in drug targets and disease genes. *Am J Pharmacogenomics*, 2002, vol. 2 (3), 197-205 [0151]
- **KWOK.** Methods for genotyping single nucleotide polymorphisms. *Annu Rev Genomics Hum Genet*, 2001, vol. 2, 235-58 [0151]

- **MARNELLOS.** High-throughput SNP analysis for genetic association studies. *Curr Opin Drug Discov Devel.*, May 2003, vol. 6 (3), 317-21 [0151]
- **MYERS et al.** *Science*, 1985, vol. 230, 1242 [0152]
- **COTTON et al.** *PNAS*, 1988, vol. 85, 4397 [0152]
- **SALEEBA et al.** *Meth. Enzymol.*, 1992, vol. 217, 286-295 [0152]
- **ORITA et al.** *PNAS*, 1989, vol. 86, 2766 [0152]
- **COTTON et al.** *Mutat. Res.*, 1993, vol. 285, 125-144 [0152]
- **HAYASHI et al.** *Genet. Anal. Tech. Appl.*, 1992, vol. 9, 73-79 [0152]
- **MYERS et al.** *Nature*, 1985, vol. 313, 495 [0152] [0163]
- **WISE et al.** A standard protocol for single nucleotide primer extension in the human genome using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun Mass Spectrom.*, 2003, vol. 17 (11), 1195-202 [0160]
- **BOCKER.** SNP and mutation discovery using base-specific cleavage and MALDI-TOF mass spectrometry. *Bioinformatics*, July 2003, vol. 19 (1), 144-153 [0161]
- **STORM et al.** MALDI-TOF mass spectrometry-based SNP genotyping. *Methods Mol Biol.*, 2003, vol. 212, 241-62 [0161]
- **JURINKE et al.** The use of MassARRAY technology for high throughput genotyping. *Adv Biochem Eng Biotechnol.*, 2002, vol. 77, 57-74 [0161]
- **JURINKE et al.** Automated genotyping using the DNA MassArray technology. *Methods Mol Biol.*, 2002, vol. 187, 179-92 [0161]
- *Biotechniques*, 1995, vol. 19, 448 [0162]
- **COHEN et al.** *Adv. Chromatogr.*, 1996, vol. 36, 127-162 [0162]
- **GRIFFIN et al.** *Appl. Biochem. Biotechnol.*, 1993, vol. 38, 147-159 [0162]
- **ORITA et al.** *Proc. Nat. Acad.* [0163]
- PCR Technology, Principles and Applications for DNA Amplification. W.H. Freeman and Co, 1992 [0163]
- Modern Epidemiology. Lippincott Williams & Wilkins, 1998, 609-622 [0168]
- **WEIR B.** Genetic Data Analysis. Sinauer, 1990 [0171]
- **HOSMER ; LEMESHOW.** Applied Logistic Regression. Wiley, 2000 [0173]
- **DALY et al.** *Nature Genetics*, 2001, vol. 29, 232-235 [0174]
- **SCHAIID et al.** *Am. J. Hum. Genet.*, 2002, vol. 70, 425-434 [0175]
- **WESTFALL et al.** Multiple comparisons and multiple tests. SAS Institute, 1999 [0176]
- **BENJAMINI ; HOCHBERG.** *Journal of the Royal Statistical Society, Series B*, 1995, vol. 57, 1289-1300 [0176]
- **WESTFALL ; YOUNG.** Resampling-based Multiple Testing. Wiley, 1993 [0176]
- Modern Epidemiology. Lippincott Williams & Wilkins, 1998, 643-673 [0177]
- **EWENS ; SPIELMAN.** *Am. J. Hum. Genet.*, 1995, vol. 62, 450-458 [0179]
- **PRITCHARD et al.** *Am. J. Hum. Gen.*, 1999, vol. 65, 220-228 [0179]
- **DEVLIN et al.** *Biometrics*, 1999, vol. 55, 997-1004 [0179]
- **DEVLIN et al.** *Genet. Epidemiol.*, 2001, vol. 21, 273-284 [0179]
- **BACANU et al.** *Am. J. Hum. Genet.*, 2000, vol. 66, 1933-1944 [0180]
- **KEHOE et al.** *Hum. Mol. Genet.*, 1999, vol. 8, 237-245 [0180]
- **DRAPER ; SMITH.** Applied Regression Analysis. Wiley, 1998 [0181]
- **HASTIE ; TIBSHIRANI ; FRIEDMAN.** The Elements of Statistical Learning. Springer, 2002 [0181]
- **WALL et al.** Haplotype blocks and linkage disequilibrium in the human genome. *Nat Rev Genet.*, August 2003, vol. 4 (8), 587-97 [0189]
- **GARNER et al.** On selecting markers for association studies: patterns of linkage disequilibrium between two and three diallelic loci. *Genet Epidemiol.*, January 2003, vol. 24 (1), 57-67 [0189]
- **ARDLIE et al.** Patterns of linkage disequilibrium in the human genome. *Nat Rev Genet.*, April 2002, vol. 3 (4), 299-309 [0189]
- *Nat Rev Genet.*, July 2002, vol. 3 (7), 566 [0189]
- **REMM et al.** High-density genotyping and linkage disequilibrium in the human genome using chromosome 22 as a model. *Curr Opin Chem Biol.*, February 2002, vol. 6 (1), 24-30 [0189]
- **ROSES.** *Nature*, 2000, vol. 405, 857-865 [0195]
- **GOULD ROTHBERG.** *Nature Biotechnology*, 2001, vol. 19, 209-211 [0195]
- **EICHELBAUM.** *Clin. Exp. Pharmacol. Physiol.*, 1996, vol. 23 (10-11), 983-985 [0195]
- **LINDER.** *Clin. Chem.*, 1997, vol. 43 (2), 254-266 [0195]
- **PFOST.** *Trends in Biotechnology*, August 2000 [0198]
- **ROSE et al.** Pharmacogenetic analysis of clinically relevant genetic polymorphisms. *Methods Mol Med.*, 2003, vol. 85, 225-37 [0199]
- **CACABELOS.** Pharmacogenomics for the treatment of dementia. *Ann Med.*, 2002, vol. 34 (5), 357-79 [0199]
- **MAIMONE et al.** Pharmacogenomics of neurodegenerative diseases. *Eur J Pharmacol.*, 09 February 2001, vol. 413 (1), 11-29 [0199]
- **POIRIER.** Apolipoprotein E: a pharmacogenetic target for the treatment of Alzheimer's disease. *Mol Diagn.*, December 1999, vol. 4 (4), 335-41 [0199]
- **SIEST et al.** Pharmacogenomics of drugs affecting the cardiovascular system. *Clin Chem Lab Med.*, 2003, vol. 41 (4), 590-9 [0199]

- **MUKHERJEE et al.** Pharmacogenomics in cardiovascular diseases. *Prog Cardiovasc Dis.*, May 2002, vol. 44 (6), 479-98 [0199]
- **MOOSER et al.** Cardiovascular pharmacogenetics in the SNP era. *J Thromb Haemost.*, July 2003, vol. 1 (7), 1398-402 [0199]
- **MCLEOD et al.** Cancer pharmacogenomics: SNPs, chips, and the individual patient. *Cancer Invest.*, 2003, vol. 21 (4), 630-40 [0199]
- **WATTERS et al.** Cancer pharmacogenomics: current and future applications. *Biochim Biophys Acta*, 17 March 2003, vol. 1603 (2), 99-111 [0199]
- Antisense Drug Technology: Principles, Strategies, and Applications. Marcel Dekker, Inc, 2001 [0201]
- **THOMPSON.** *Drug Discovery Today*, 2002, vol. 7 (17), 912-917 [0201] [0205]
- **LAVERY et al.** Antisense and RNAi: powerful tools in drug target discovery and validation. *Curr Opin Drug Discov Devel.*, July 2003, vol. 6 (4), 561-9 [0202] [0206]
- **STEPHENS et al.** Antisense oligonucleotide therapy in cancer. *Curr Opin Mol Ther.*, April 2003, vol. 5 (2), 118-22 [0202]
- **KURRECK.** Antisense technologies. Improvement through novel chemical modifications. *Eur J Biochem.*, April 2003, vol. 270 (8), 1628-44 [0202]
- **DIAS et al.** Antisense oligonucleotides: basic concepts and mechanisms. *Mol Cancer Ther.*, March 2002, vol. 1 (5), 347-55 [0202]
- **CHEN.** Clinical development of antisense oligonucleotides as anti-cancer therapeutics. *Methods Mol Med.*, 2003, vol. 75, 621-36 [0202]
- **WANG et al.** Antisense anticancer oligonucleotide therapeutics. *Curr Cancer Drug Targets*, December 2001, vol. 1 (3), 177-96 [0202]
- **BENNETT.** Efficiency of antisense oligonucleotide drug discovery. *Antisense Nucleic Acid Drug Dev.*, June 2002, vol. 12 (3), 215-24 [0202]
- **AGAMI.** RNAi and related mechanisms and their potential use for therapy. *Curr Opin Chem Biol.*, December 2002, vol. 6 (6), 829-34 [0206]
- **SHI.** Mammalian RNAi for the masses. *Trends Genet.*, January 2003, vol. 19 (1), 9-12 [0206]
- **SHUEY et al.** RNAi: gene-silencing in therapeutic intervention. *Drug Discovery Today*, October 2002, vol. 7 (20), 1040-1046 [0206]
- **MCMANUS et al.** *Nat Rev Genet*, October 2002, vol. 3 (10), 737-47 [0206]
- **XIA et al.** *Nat Biotechnol*, October 2002, vol. 20 (10), 1006-10 [0206]
- **PLASTERK et al.** *Curr Opin Genet Dev*, October 2000, vol. 10 (5), 562-7 [0206]
- **BOSHER et al.** *Nat Cell Biol*, February 2000, vol. 2 (2), E31-6 [0206]
- **HUNTER.** *Curr Biol*, 17 June 1999, vol. 9 (12), R440-2 [0206]
- **KREN et al.** *Proc. Natl. Acad. Sci. USA*, 1999, vol. 96, 10349-10354 [0207]
- **JAZWINSKA.** *A Trends Guide to Genetic Variation and Genomic Medicine*, March 2002, S30-S36 [0218] [0220]
- **ROTHBERG.** *Nat Biotechnol*, March 2001, vol. 19 (3), 209-11 [0219]
- Current Protocols in Toxicology. John Wiley & Sons, Inc, [0221]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1990 [0231]
- Current Protocols in Pharmacology. John Wiley & Sons, Inc, [0234]
- **GILL.** An assessment of the utility of single nucleotide polymorphisms (SNPs) for forensic purposes. *Int J Legal Med.*, 2001, vol. 114 (4-5), 204-10 [0235]
- **FRUDAKIS et al.** A Classifier for the SNP-Based Inference of Ancestry. *Journal of Forensic Sciences*, 2003, vol. 48 (4), 771-782 [0239]
- **KRAWCZAK.** Informativity assessment for biallelic single nucleotide polymorphisms. *Electrophoresis*, June 1999, vol. 20 (8), 1676-81 [0246]
- Current Protocols in Human Genetics. John Wiley & Sons, 2002, 14.1-14.7 [0249]
- **SAMBROOK ; RUSSELL.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 2000 [0255] [0274] [0320] [0321] [0332]
- **AUSUBEL et al.** *Current Protocols in Molecular Biology*, 1992 [0260]
- **TERPE.** Overview of tag protein fusions: from molecular and biochemical fundamentals to commercial systems. *Appl Microbiol Biotechnol.*, 07 November 2002, vol. 60 (5), 523-33 [0261]
- **GRADDIS et al.** Designing proteins that work using recombinant technologies. *Curr Pharm Biotechnol.*, December 2002, vol. 3 (4), 285-97 [0261]
- **NILSSON et al.** Affinity fusion strategies for detection, purification, and immobilization of recombinant proteins. *Protein Expr Purif.*, October 1997, vol. 11 (1), 1-16 [0261]
- **BOWIE et al.** *Science*, 1990, vol. 247, 1306-1310 [0266]
- **CUNNINGHAM et al.** *Science*, 1989, vol. 244, 1081-1085 [0268]
- **SMITH et al.** *J. Mol. Biol.*, 1992, vol. 224, 899-904 [0268]
- **DE VOS et al.** *Science*, 1992, vol. 255, 306-312 [0268]
- **T.E. CREIGHTON.** Proteins - Structure and Molecular Properties. W. H. Freeman and Company, 1993 [0271]
- **WOLD, F.** Posttranslational Covalent Modification of Proteins. Academic Press, 1983, 1-12 [0271]
- **SEIFTER et al.** *Meth. Enzymol.*, 1990, vol. 182, 626-646 [0271]
- **RATTAN et al.** *Ann. N.Y. Acad. Sci.*, 1992, vol. 663, 48-62 [0271]
- Current Protocols in Protein Science. John Wiley & Sons, 2002 [0273]

- **Methods in Enzymology: Guide to Molecular Cloning Techniques.** Academic Press, 1987 [0274]
- **Current Protocols in Immunology.** John Wiley & Sons [0278]
- **HAGE.** Immunoassays. *Anal Chem.*, 15 June 1999, vol. 71 (12), 294R-304R [0278]
- **LAM et al.** *Nature*, 1991, vol. 354, 82-84 [0288]
- **HOUGHTEN et al.** *Nature*, 1991, vol. 354, 84-86 [0288]
- **SONGYANG et al.** *Cell*, 1993, vol. 72, 767-778 [0288]
- **HODGSON.** *Biotechnology*, September 1992, vol. 10 (9), 973-80 [0292]
- **ZERVOS et al.** *Cell*, 1993, vol. 72, 223-232 [0297]
- **MADURA et al.** *J. Biol. Chem.*, 1993, vol. 268, 12046-12054 [0297]
- **BARTEL et al.** *Biotechniques*, 1993, vol. 14, 920-924 [0297]
- **IWABUCHI et al.** *Oncogene*, 1993, vol. 8, 1693-1696 [0297]
- **MORRISON et al.** *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 6851 [0300]
- **NEUBERGER et al.** *Nature*, 1984, vol. 312, 604 [0300]
- **WARD et al.** *Nature*, 1989, vol. 334, 544 [0300]
- **SEGAL et al.** *J. Immunol. Methods*, 2001, vol. 248, 1 [0300]
- **CARTER.** *J. Immunol. Methods*, 2001, vol. 248, 7 [0300]
- **TODOROVSKA et al.** *J. Immunol. Methods*, 2001, vol. 248, 47 [0300]
- **HARLOW.** *Antibodies.* Cold Spring Harbor Press, 1989 [0301]
- **KOHLER ; MILSTEIN.** *Nature*, 1975, vol. 256, 495 [0302]
- **HE et al.** *J. Immunol.*, 2002, vol. 169, 595 [0302]
- **HOOGENBOOM ; CHAMES.** *Immunol. Today*, 2000, vol. 21, 371 [0302]
- **LIU et al.** *J. Mol. Biol.*, 2002, vol. 315, 1063 [0302]
- **MORGAN.** Antibody therapy for Alzheimer's disease. *Expert Rev Vaccines*, February 2003, vol. 2 (1), 53-9 [0305]
- **ROSS et al.** Anticancer antibodies. *Am J Clin Pathol.*, April 2003, vol. 119 (4), 472-85 [0305]
- **GOLDENBERG.** Advancing role of radiolabeled antibodies in the therapy of cancer. *Cancer Immunol Immunother.*, 11 March 2003, vol. 52 (5), 281-96 [0305]
- **ROSS et al.** Antibody-based therapeutics in oncology. *Expert Rev Anticancer Ther.*, February 2003, vol. 3 (1), 107-21 [0305]
- **CAO et al.** Bispecific antibody conjugates in therapeutics. *Adv Drug Deliv Rev.*, 10 February 2003, vol. 55 (2), 171-97 [0305]
- **VON MEHREN et al.** Monoclonal antibody therapy for cancer. *Annu Rev Med.*, 03 December 2001, vol. 54, 343-69 [0305]
- **HUDSON et al.** Engineered antibodies. *Nat Med.*, vol. 9 (1), 129-34 [0305]
- **BREKKE et al.** Therapeutic antibodies for human diseases at the dawn of the twenty-first century. *Nat Rev Drug Discov.*, January 2003, vol. 2 (1), 52-62 [0305]
- *Nat Rev Drug Discov.*, March 2003, vol. 2 (3), 240 [0305]
- **HOUEBINE.** Antibody manufacture in transgenic animals and comparisons with other systems. *Curr Opin Biotechnol.*, December 2002, vol. 13 (6), 625-9 [0305] [0350]
- **ANDREAKOS et al.** Monoclonal antibodies in immune and inflammatory diseases. *Curr Opin Biotechnol.*, December 2002, vol. 13 (6), 615-20 [0305]
- **KELLERMANN et al.** Antibody discovery: the use of transgenic mice to generate human monoclonal antibodies for therapeutics. *Curr Opin Biotechnol.*, December 2002, vol. 13 (6), 593-7 [0305]
- **PINI et al.** Phage display and colony filter screening for high-throughput selection of antibody libraries. *Comb Chem High Throughput Screen*, November 2002, vol. 5 (7), 503-10 [0305]
- **BATRA et al.** Pharmacokinetics and biodistribution of genetically engineered antibodies. *Curr Opin Biotechnol.*, December 2002, vol. 13 (6), 603-8 [0305]
- **TANGRI et al.** Rationally engineered proteins or antibodies with absent or reduced immunogenicity. *Curr Med Chem.*, December 2002, vol. 9 (24), 2191-9 [0305]
- **LEONG et al.** *Cytokine*, 2001, vol. 16, 106 [0312]
- **SMITH et al.** *Gene*, 1988, vol. 67, 31-40 [0325]
- **AMANN et al.** *Gene*, 1988, vol. 69, 301-315 [0325]
- **STUDIER et al.** *Gene Expression Technology: Methods in Enzymology*, 1990, vol. 185, 60-89 [0325]
- **GOTTESMAN, S.** *Gene Expression Technology: Methods in Enzymology.* Academic Press, 1990, vol. 185, 119-128 [0326]
- **WADA et al.** *Nucleic Acids Res.*, 1992, vol. 20, 2111-2118 [0326]
- **BALDARI et al.** *EMBO J.*, 1987, vol. 6, 229-234 [0327]
- **KURJAN et al.** *Cell*, 1982, vol. 30, 933-943 [0327]
- **SCHULTZ et al.** *Gene*, 1987, vol. 54, 113-123 [0327]
- **SMITH et al.** *Mol. Cell Biol.*, 1983, vol. 3, 2156-2165 [0328]
- **LUCKLOW et al.** *Virology*, 1989, vol. 170, 31-39 [0328]
- **SEED, B.** *Nature*, 1987, vol. 329, 840 [0329]
- **KAUFMAN et al.** *EMBO J.*, 1987, vol. 6, 187-195 [0329]
- *Current Protocols in Molecular Biology.* John Wiley & Sons [0340]
- **HOGAN, B.** *Manipulating the Mouse Embryo.* Cold Spring Harbor Laboratory Press, 1986 [0346]
- **LAKSO et al.** *PNAS*, 1992, vol. 89, 6232-6236 [0347]
- **O'GORMAN et al.** *Science*, 1991, vol. 251, 1351-1355 [0347]

- **WILMUT, I. et al.** *Nature*, 1997, vol. 385, 810-813 [0348]
- **PETTERS et al.** Transgenic animals as models for human disease. *Transgenic Res.*, 2000, vol. 9 (4-5), 347-51 [0350]
- **WOLF et al.** Use of transgenic animals in understanding molecular mechanisms of toxicity. *J Pharm Pharmacol.*, June 1998, vol. 50 (6), 567-74 [0350]
- **ECHELARD.** Recombinant protein production in transgenic animals. *Curr Opin Biotechnol.*, October 1996, vol. 7 (5), 536-40 [0350]
- **HOUDEBINE.** Transgenic animal bioreactors. *Transgenic Res.*, 2000, vol. 9 (4-5), 305-20 [0350]
- **PIRITY et al.** Embryonic stem cells, creating transgenic animals. *Methods Cell Biol.*, 1998, vol. 57, 279-93 [0350]
- **ROBL et al.** Artificial chromosome vectors and expression of complex proteins in transgenic animals. *Theriogenology*, 01 January 2003, vol. 59 (1), 107-13 [0350]
- **ALTSCHUL.** *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0355]
- **BRUTLAG.** *Comp. Chem.*, 1993, vol. 17, 203-207 [0355]
- **GERMER S. ; HOLLAND M.J. ; HIGUCHI R.** *Genome Res.*, 2000, vol. 10, 258-266 [0366] [0377]

专利名称(译)	与心肌梗塞相关的遗传多态性，检测方法及其用途		
公开(公告)号	EP2474631B1	公开(公告)日	2014-02-12
申请号	EP2012163424	申请日	2003-12-22
申请(专利权)人(译)	塞雷拉基因组公司		
当前申请(专利权)人(译)	塞雷拉基因组公司		
[标]发明人	LAKOUBOVA OLGA DEVLIN JAMES CARGILL MICHELE		
发明人	LAKOUBOVA, OLGA DEVLIN, JAMES CARGILL, MICHELE		
IPC分类号	C12Q1/68 C07K14/47 C07K14/705 C07K16/18 C07K16/28 A61B C07H21/00 C07H21/04 C12N15/12 C12P19/34 G01N33/53		
CPC分类号	C12Q1/6883 C12Q2600/156		
优先权	60/434778 2002-12-20 US 60/466412 2003-04-30 US 60/453135 2003-03-10 US 60/504955 2003-09-23 US		
其他公开文献	EP2474631A3 EP2474631A2		
外部链接	Espacenet		

摘要(译)

本发明基于与心肌梗塞相关的遗传多态性的发现。特别地，本发明涉及含有多态性的核酸分子，由这种核酸分子编码的变体蛋白，用于检测多态性核酸分子和蛋白质的试剂，以及使用该核酸和蛋白质的方法以及使用方法。用于检测的试剂。




FIGURE 1