



(11)

**EP 3 121 277 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**11.04.2018 Bulletin 2018/15**

(51) Int Cl.:  
**C12N 15/09** <sup>(2006.01)</sup> **A61K 39/395** <sup>(2006.01)</sup>  
**A61P 25/28** <sup>(2006.01)</sup> **C07K 16/18** <sup>(2006.01)</sup>  
**G01N 33/53** <sup>(2006.01)</sup> **C12P 21/08** <sup>(2006.01)</sup>  
**G01N 33/68** <sup>(2006.01)</sup>

(21) Application number: **16183829.7**

(22) Date of filing: **06.02.2009**

(54) **ANTIBODY CAPABLE OF BINDING SPECIFICALLY TO AB-OLIGOMER, AND USE THEREOF**

ZUR SPEZIFISCHEN BINDUNG AN AB-OLIGOMER FÄHIGER ANTIKÖRPER UND VERWENDUNG DAVON

ANTICORPS POUVANT SE LIER SPÉCIFIQUEMENT À AB-OLIGOMÈRE ET SON UTILISATION

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR**

(30) Priority: **08.02.2008 JP 2008028386**  
**01.08.2008 US 85545**  
**04.08.2008 JP 2008201058**

(43) Date of publication of application:  
**25.01.2017 Bulletin 2017/04**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:  
**09707511.3 / 2 246 427**

(73) Proprietors:  
• **Immunas Pharma, Inc.**  
**Kanagawa 213-0012 (JP)**  
• **National Center for Geriatrics and Gerontology**  
**Aichi 4748511 (JP)**

(72) Inventors:  
• **MATSUBARA, Etsuro**  
**Hirosaki-shi Aomori 036-8216 (JP)**  
• **SHIBATA, Masao**  
**Kawasaki-shi Kanagawa 213-0012 (JP)**

• **YOKOSEKI, Tatsuki**  
**Kawasaki-shi Kanagawa 213-0012 (JP)**

(74) Representative: **Bösl, Raphael Konrad**  
**Isenbruck Bösl Hörschler LLP**  
**Patentanwälte**  
**Prinzregentenstraße 68**  
**81675 München (DE)**

(56) References cited:  
**WO-A2-03/104437 WO-A2-2006/055178**

• **MORETTO NADIA ET AL:**  
**"Conformation-sensitive antibodies against Alzheimer amyloid-beta by immunization with a thioredoxin-constrained B-cell epitope peptide", JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, US, vol. 282, no. 15, 1 April 2007 (2007-04-01), pages 11436-11445, XP002451806, ISSN: 0021-9258, DOI: 10.1074/JBC.M609690200**  
• **WILLIAM KLEIN: "Abeta toxicity in Alzheimer's disease: globular oligomers (ADDLs) as new vaccine and drug targets.", NEUROCHEMISTRY INTERNATIONAL, vol. 41, no. 5, 1 November 2002 (2002-11-01), pages 345-352, XP055016511, ISSN: 0197-0186**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 3 121 277 B1**

**Description**

## Technical Field

5 **[0001]** The present invention relates to antibodies that specifically bind to A $\beta$  oligomers and uses thereof.

## Background Art

10 **[0002]** Various evidence has shown that deterioration of memory arises from synaptic dysfunction triggered by soluble A $\beta$  oligomers (see Non-Patent Documents 1 and 2). Excessive accumulation and deposition of A $\beta$  oligomers may be the trigger for a series of pathological cascades that lead to Alzheimer's disease (AD). Therefore, therapeutic intervention targeting A $\beta$  oligomers may be effective for blocking these cascades. However, findings on neurodegeneration mediated by core molecules of this amyloid cascade hypothesis which are responsible for neurodegeneration, particularly by A $\beta$  oligomers, originate from *in vitro* experiments (see Non-Patent Document 3). This neurodegeneration has not been  
 15 proven directly *in vivo*. The greatest defect of previously reported *in vivo* experiments is that they failed to demonstrate synaptic toxicity of endogenous A $\beta$  oligomers due to the lack of conformation-specific molecular tools (see Non-Patent Document 4). There has been known no technique capable of proving the toxicity within the human brain, an aspect which is difficult to demonstrate even in Alzheimer's disease mouse models. Thus, the *in vivo* neurotoxicity of endogenous A $\beta$  has been often disregarded. It has been unknown why NFT formation and loss of nerve cells precede senile plaque  
 20 formation in the human entorhinal cortex, and how A $\beta$  oligomers are involved in this mechanism.

**[0003]** Prior art literature information relating to the present invention is shown below.

[Non-Patent Document 1] Klein WL, Trends Neurosci. 24: 219-224, 2001.

[Non-Patent Document 2] Selkoe DJ, Science 298: 789-791, 2002.

25 [Non-Patent Document 3] Hass C et al.: Nature Review 8: 101-12, 2007.

[Non-Patent Document 4] Lee EB, et al.: J. Biol. Chem. 281: 4292-4299, 2006.

WO 03/104437 A2 discloses polyclonal antibodies which preferentially bind to oligomers of amyloid  $\beta$ . WO 03/104437 A2 further discloses the use of said antibodies for the diagnosis, treatment and prevention of diseases associated with amyloid deposits including Alzheimer's disease.

30 WO 2006/055178 A1 discloses monoclonal antibodies which bind to oligomers of amyloid  $\beta$ . WO 2006/055178 A1 further discloses the use of said antibodies for the detection, treatment and prevention of Alzheimer's disease.

Moretto N. et al. (J. Biol. Chem. (2007); Conformation-sensitive antibodies against Alzheimer amyloid- $\beta$  by immunization with a thioredoxin-constrained B-cell epitope peptide; 282(15): 11436-11445) discloses an antibody which recognizes amyloid- $\beta$  oligomers, and the use thereof for the diagnosis and treatment of Alzheimer's disease in  
 35 human brain tissue and a mouse model.

## Disclosure of the Invention

## [Problems to be Solved by the Invention]

40 **[0004]** The present invention was achieved in view of the above circumstances. An objective of the present invention is to provide antibodies that bind specifically to A $\beta$  oligomers, but not to an A $\beta$  monomer, and uses thereof. More specifically, the present invention provides an antibody that binds specifically to an A $\beta$  oligomer but not to an A $\beta$  monomer, wherein the antibody is a monoclonal antibody and is selected from the group consisting of:

- 45 (1) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 69 as CDR1, the amino acid sequence of SEQ ID NO: 71 as CDR2, and the amino acid sequence of SEQ ID NO: 73 as CDR3 and an L chain having the amino acid sequence of SEQ ID NO: 75 as CDR1, the amino acid sequence of SEQ ID NO: 77 as CDR2, and the amino acid sequence of SEQ ID NO: 79 as CDR3;  
 50 (2) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 65 as VH and an L chain having the amino acid sequence of SEQ ID NO: 67 as VL; and  
 (3) an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 61 and an L chain having the amino acid sequence of SEQ ID NO: 63.

55 The antibody of the present invention can e.g. a chimeric antibody or a humanized antibody. The present invention further provides methods for detecting A $\beta$  oligomers using the antibody, methods for diagnosing Alzheimer's disease using the antibody, and pharmaceutical agents comprising the antibody.  
 In particular, the present invention provides:

- (a) a composition comprising the inventive antibody and a pharmaceutically acceptable carrier;
- (b) an agent for use in treating cognitive impairment, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (c) a therapeutic agent for use in treating Alzheimer's disease, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (d) an agent for use in suppressing the progression of Alzheimer's disease, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (e) an agent for use in suppressing senile plaque formation, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (f) an agent for use in suppressing A $\beta$  accumulation, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (g) an agent for use in neutralizing neurotoxicity, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (h) an agent for use in inhibiting A $\beta$  amyloid fibril formation, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (i) an agent for use in neutralizing synaptic toxicity, which comprises the inventive antibody or the inventive composition as an active ingredient;
- (j) a method for detecting an A $\beta$  oligomer, which comprises the step of detecting an A $\beta$  oligomer contained in a sample collected from a subject using the inventive antibody;
- (k) a method of diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises the step of (a0) below:

(a0) using the inventive antibody of claim 1 or 2 to detect an A $\beta$  oligomer in a sample collected from a subject;

or  
the steps of (a1) to (c1) below:

- (a1) contacting a sample collected from a subject with the inventive antibody;
- (b1) measuring the amount of A $\beta$  oligomer in the sample; and
- (c1) determining that the subject is a possible Alzheimer's disease patient, when the amount measured in step (b1) is higher than that of a healthy individual;

or  
the steps of (a2) to (c2) below:

- (a2) contacting a sample collected from a subject with the inventive antibody and an antibody that binds to an A $\beta$  monomer;
  - (b2) measuring the ratio of A $\beta$  oligomer to A $\beta$  monomer in the sample; and
  - (c2) determining that the subject is a possible Alzheimer's disease patient, when the ratio measured in step (b2) is higher than that of a healthy individual.
- The sample used in the above methods can be blood or cerebrospinal fluid.

(l) A pharmaceutical agent for use in diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises the inventive antibody.

[Means for Solving the Problems]

**[0005]** The present inventors produced monoclonal antibodies that are specific to only soluble amyloid  $\beta$  (A $\beta$ ) oligomers and do not recognize soluble A $\beta$  monomers which are physiological molecules, and confirmed that the antibodies have the following:

- (1) anti-neurotoxic activity;
- (2) activity to suppress A $\beta$  amyloid fibril formation;
- (3) specificity to recognize only A $\beta$  oligomers;
- (4) ability to capture A $\beta$  oligomers in AD brain; and
- (5) ability to prevent the development of Alzheimer's disease-like phenotypes (memory impairment, brain A $\beta$  accumulation) in APPswe transgenic mice (Tg2576).

**[0006]** Using an ultrafiltration/molecular sieve method, among the antibodies produced, monoclonal 1A9 and 2C3 were determined to specifically recognize oligomers of 30 kDa or more, mainly 100 kDa or more, but not monomers of approximately 4.5 kDa. The two antibodies were confirmed to have neurotoxicity-neutralizing activity by evaluating the neutralizing effect against A $\beta$  1-42-induced neurotoxicity in PC12 cells differentiated into nerve cells. Thioflavin T assay and electron microscopy showed that the antibodies have activity to suppress A $\beta$  amyloid fibril formation. The ability of 1A9 and 2C3 to capture A $\beta$  oligomers in AD brain was confirmed by immunoprecipitation using the antibodies in the presence of SDS-stable 4-, 5-, 8-, and 12-mers. Furthermore, to determine the *in vivo* neurotoxicity in the human brain, the amount of polymers recognized by the antibodies was evaluated in the human entorhinal cortex mostly at Braak NFT Stages I to III. By particularly focusing on the 12-mer, which has been reported to have neurotoxicity in animal studies, it was confirmed that the polymer accumulation precedes the occurrence of cognitive impairment, and is increased with the progression of Braak NFT stage. This result shows for the first time that the 12-mer, which is specifically recognized by the antibodies, is a conformational assembly that causes *in vivo* neurotoxicity in the human brain. The present inventors also discovered that the oligomeric conformational structure recognized by the antibodies is present in cerebrospinal fluid (CSF), and is increased in AD patients. The present inventors used 1A9 or 2C3 in passive immunotherapy by intravenous injection as with other neurological disorders. It was confirmed that Tg2576 mice are protected from memory impairment, senile plaque formation, synaptic dysfunction, and A $\beta$  accumulation by subchronic passive immunotherapy, without harmful side-effects. The results obtained by the present inventors demonstrated for the first time that monoclonal 1A9 and 2C3 are promising candidates for therapeutic antibodies for preventing Alzheimer's disease-like phenotypes in Tg2576 mice, which are expected to show their effect by conventional peripheral intravenous administration, and thus there is no need to consider brain transfer.

**[0007]** The present inventors also confirmed that passive immunotherapy using the 1A9 and 2C3 antibodies suppresses senile plaque amyloid formation and swollen dystrophic neurite formation. Furthermore, the present inventors discovered that a fraction of the 1A9 and 2C3 antibodies administered into the blood transfers into the brain.

**[0008]** As described above, the present inventors disclose herein that monoclonal 1A9 and 2C3, which are antibodies that specifically bind to A $\beta$  oligomers, fulfill all of the diagnostic/therapeutic antibody criteria, and are promising candidates for therapeutic antibodies for diagnosing/preventing Alzheimer's disease.

**[0009]** Furthermore, as with the 1A9 and 2C3 antibodies, the present inventors successfully obtained the 5A5, 5A9, 4F7, 4H5, 6E4, and 6H4 antibodies which bind specifically to A $\beta$  oligomers, but do not recognize A $\beta$  monomers. The present inventors discovered that these six types of antibodies have activity to neutralize A $\beta$ -induced neurotoxicity and to suppress A $\beta$  amyloid fibril formation.

**[0010]** The present inventors disclose that the above-mentioned 5A5, 5A9, 4F7, 4H5, 6E4, and 6H4 antibodies are promising candidates for therapeutic antibodies for diagnosing/preventing Alzheimer's disease.

**[0011]** More specifically, the following is disclosed:

- [1] an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 1 and an L chain having the amino acid sequence of SEQ ID NO: 3;
- [2] an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 21 and an L chain having the amino acid sequence of SEQ ID NO: 23;
- [3] an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 41 and an L chain having the amino acid sequence of SEQ ID NO: 43;
- [4] an antibody according to the present invention which binds to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 61 and an L chain having the amino acid sequence of SEQ ID NO: 63;
- [5] an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 81 and an L chain having the amino acid sequence of SEQ ID NO: 83;
- [6] an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 101 and an L chain having the amino acid sequence of SEQ ID NO: 103;
- [7] an antibody of any one of (1) to (38) below:

- (1) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 9 as CDR1, the amino acid sequence of SEQ ID NO: 11 as CDR2, and the amino acid sequence of SEQ ID NO: 13 as CDR3;
- (2) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 15 as CDR1, the amino acid sequence of SEQ ID NO: 17 as CDR2, and the amino acid sequence of SEQ ID NO: 19 as CDR3;
- (3) an antibody that comprises the H chain of (1) and the L chain of (2);
- (4) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 5 as VH;
- (5) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 7 as VL;
- (6) an antibody that comprises the H chain of (4) and the L chain of (5);
- (7) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 29 as CDR1, the

amino acid sequence of SEQ ID NO: 31 as CDR2, and the amino acid sequence of SEQ ID NO: 33 as CDR3;  
 (8) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 35 as CDR1, the  
 amino acid sequence of SEQ ID NO: 37 as CDR2, and the amino acid sequence of SEQ ID NO: 39 as CDR3;  
 (9) an antibody that comprises the H chain of (7) and the L chain of (8);  
 5 (10) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 25 as VH;  
 (11) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 27 as VL;  
 (12) an antibody that comprises the H chain of (10) and the L chain of (11);  
 (13) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 49 as CDR1, the  
 amino acid sequence of SEQ ID NO: 51 as CDR2, and the amino acid sequence of SEQ ID NO: 53 as CDR3;  
 10 (14) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 55 as CDR1, the  
 amino acid sequence of SEQ ID NO: 57 as CDR2, and the amino acid sequence of SEQ ID NO: 59 as CDR3;  
 (15) an antibody that comprises the H chain of (13) and the L chain of (14);  
 (16) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 45 as VH;  
 (17) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 47 as VL;  
 15 (18) an antibody that comprises the H chain of (16) and the L chain of (17);  
 (19) an antibody according to the present invention that comprises an H chain having the amino acid sequence  
 of SEQ ID NO: 69 as CDR1, the amino acid sequence of SEQ ID NO: 71 as CDR2, and the amino acid sequence  
 of SEQ ID NO: 73 as CDR3;  
 20 (20) an antibody according to the present invention that comprises an L chain having the amino acid sequence  
 of SEQ ID NO: 75 as CDR1, the amino acid sequence of SEQ ID NO: 77 as CDR2, and the amino acid sequence  
 of SEQ ID NO: 79 as CDR3;  
 (21) an antibody that comprises the H chain of (19) and the L chain of (20);  
 (22) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 65 as VH;  
 (23) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 67 as VL;  
 25 (24) an antibody that comprises the H chain of (22) and the L chain of (23);  
 (25) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 89 as CDR1, the  
 amino acid sequence of SEQ ID NO: 91 as CDR2, and the amino acid sequence of SEQ ID NO: 93 as CDR3;  
 (26) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 95 as CDR1, the  
 amino acid sequence of SEQ ID NO: 97 as CDR2, and the amino acid sequence of SEQ ID NO: 99 as CDR3;  
 30 (27) an antibody that comprises the H chain of (25) and the L chain of (26);  
 (28) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 85 as VH;  
 (29) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 87 as VL;  
 (30) an antibody that comprises the H chain of (28) and the L chain of (29);  
 (31) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 109 as CDR1, the  
 amino acid sequence of SEQ ID NO: 111 as CDR2, and the amino acid sequence of SEQ ID NO: 113 as CDR3;  
 35 (32) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 115 as CDR1, the  
 amino acid sequence of SEQ ID NO: 117 as CDR2, and the amino acid sequence of SEQ ID NO: 119 as CDR3;  
 (33) an antibody that comprises the H chain of (31) and the L chain of (32);  
 (34) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 105 as VH;  
 40 (35) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 107 as VL;  
 (36) an antibody that comprises the H chain of (34) and the L chain of (35);  
 (37) an antibody that comprises one or more amino acid substitutions, deletions, additions, and/or insertions in  
 the antibody of any one of (1) to (36), which has equivalent activity to the antibody of any one of (1) to (36); and  
 (38) an antibody that binds to the epitope bound by the antibody of any one of (1) to (36);

45 [8] the antibody of [7], wherein the antibody is a chimeric antibody or a humanized antibody;

[9] a composition comprising the antibody of any one of [1] to [8] and a pharmaceutically acceptable carrier;

[10] an agent against cognitive impairment, which comprises the antibody of any one of [1] to [8] or the composition  
 of [9] as an active ingredient;

50 [11] a therapeutic agent for Alzheimer's disease, which comprises the antibody of any one of [1] to [8] or the  
 composition of [9] as an active ingredient;

[12] an agent for suppressing the progression of Alzheimer's disease, which comprises the antibody of any one of  
 [1] to [8] or the composition of [9] as an active ingredient;

[13] an agent for suppressing senile plaque formation, which comprises the antibody of any one of [1] to [8] or the  
 55 composition of [9] as an active ingredient;

[14] an agent for suppressing A $\beta$  accumulation, which comprises the antibody of any one of [1] to [8] or the composition  
 of [9] as an active ingredient;

[15] an anti-neurotoxic agent, which comprises the antibody of any one of [1] to [8] or the composition of [9] as an

active ingredient;

[16] an agent for inhibiting A $\beta$  amyloid fibril formation, which comprises the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[17] an agent against synaptic toxicity, which comprises the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[18] a method for detecting an A $\beta$  oligomer, which comprises the step of detecting an A $\beta$  oligomer contained in a sample collected from a subject using the antibody of any one of [1] to [8];

[19] a method of diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises using the antibody of any one of [1] to [8] to detect an A $\beta$  oligomer in a sample collected from a subject;

[20] a method of diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises the steps of:

(a) contacting a sample collected from a subject with the antibody of any one of [1] to [8]; and

(b) measuring the amount of A $\beta$  oligomer in the sample,

wherein the subject is determined to be a possible Alzheimer's disease patient, when the amount measured in step (b) is higher than that of a healthy individual;

[21] a method of diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises the steps of:

(a) contacting a sample collected from a subject with the antibody of any one of [1] to [8] and an antibody that binds to an A $\beta$  monomer; and

(b) measuring the ratio of A $\beta$  oligomer to A $\beta$  monomer in the sample,

wherein the subject is determined to be a possible Alzheimer's disease patient, when the ratio measured in step (b) is higher than that of a healthy individual;

[22] the method of any one of [18] to [21], wherein the sample is blood or cerebrospinal fluid;

[23] a pharmaceutical agent for use in the method of any one of [18] to [21]; and

[24] a kit for use in the method of any one of [18] to [21].

**[0012]** Furthermore, the following is disclosed:

[25] a method for preventing and/or treating cognitive impairment, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[26] a method for preventing and/or treating Alzheimer's disease, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[27] a method for suppressing the progression of Alzheimer's disease, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[28] a method for suppressing senile plaque formation, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[29] a method for suppressing A $\beta$  accumulation, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[30] a method for neutralizing neurotoxicity, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[31] a method for inhibiting A $\beta$  amyloid fibril formation, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[32] a method for neutralizing synaptic toxicity, which comprises the step of administering the antibody of any one of [1] to [8] or the composition of [9] as an active ingredient;

[33] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent against cognitive impairment;

[34] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of a therapeutic agent for Alzheimer's disease;

[35] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent for suppressing the progression of Alzheimer's disease;

[36] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent for suppressing senile plaque formation;

[37] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent for suppressing A $\beta$  accumulation;

[38] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent for neutralizing (suppressing) neurotoxicity;

[39] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent for inhibiting A $\beta$  amyloid fibril formation;

[40] use of the antibody of any one of [1] to [8] or the composition of [9] in the production of an agent for neutralizing (suppressing) synaptic toxicity;

[41] the antibody of any one of [1] to [8] or the composition of [9] for use in preventing and/or treating cognitive impairment;

[42] the antibody of any one of [1] to [8] or the composition of [9] for use in preventing and/or treating Alzheimer's disease;

[43] the antibody of any one of [1] to [8] or the composition of [9] for use in suppressing the progression of Alzheimer's disease;

[44] the antibody of any one of [1] to [8] or the composition of [9] for use in suppressing senile plaque formation;

[45] the antibody of any one of [1] to [8] or the composition of [9] for use in suppressing A $\beta$  accumulation;

[46] the antibody of any one of [1] to [8] or the composition of [9] for use in neutralizing (suppressing) neurotoxicity;

[47] the antibody of any one of [1] to [8] or the composition of [9] for use in inhibiting A $\beta$  amyloid fibril formation; and

[48] the antibody of any one of [1] to [8] or the composition of [9] for use in neutralizing (suppressing) synaptic toxicity.

#### [Effects of the Invention]

**[0013]** The antibodies provided by the present invention are expected to greatly contribute to the establishment of preventive/therapeutic methods selective to molecules responsible for evoking pathological conditions of Alzheimer's disease, and the establishment of early diagnostic markers for Alzheimer's disease. The present inventors obtained evidence showing that, even in antibody therapy targeting pathological conditions in the brain, peripheral intravenous administration is sufficient and there is no need to consider brain transfer. Thus, the present invention is expected to rapidly accelerate the progress of antibody drugs for Alzheimer's disease.

#### Brief Description of the Drawings

#### **[0014]**

Fig. 1 presents photographs and a graph showing the results of production and characteristic determination of oligomer-specific antibodies. A:

Electrophoresis of immunogens. The A $\beta$  1-42 tetramer (black arrowhead) which is free of contamination of the A $\beta$  1-42 monomer (outlined arrowhead) was isolated using SDS-PAGE. Lane 1: A $\beta$  1-42 dissolved in 10 mM phosphate buffer; and

Lane 2: A $\beta$  1-42 dissolved in distilled deionized water. B: A $\beta$  amyloid, which is insoluble in a buffer but can be extracted using formic acid from the brain of Alzheimer's disease patients, was immunoprecipitated using the supernatant of a positive hybridoma cell culture, and the immune complex was selectively separated using protein-G agarose (Amersham). Nine clones were tested; lane 2 (asterisk) is 1A9 and lane 6 (double asterisk) is 2C3. C: Elution profile of SEC of a conditioned medium. Among the 24 SEC-collected fractions, fractions 8, 13, and 16 were subjected to 1A9 immunoprecipitation. A $\beta$  immunoreactivity was detected using 4G8. The black arrowhead indicates the trimer and the outlined arrowhead indicates the dimer. Asterisk (\*) indicates the anti-mouse IgG light chain.

Fig. 2 presents photographs and a graph showing the antitoxic activity of 1A9 and 2C3. A to F: Representative images of NGF-treated PC12 (PC12N) cells, which were exposed to seed-free A $\beta$  1-42 at 37°C for 48 hours in the presence or absence of the antibodies (left half of each panel). Representative calcein AM/PI staining where live cells were stained green and dead cells were stained red (right half of each panel). G: The viability of cells exposed to seed-free A $\beta$  1-42 (25  $\mu$ M) with the following antibodies: non-specific IgG2b (filled square); 4G8 (open triangle); 1A9 (open square); and 2C3 (filled circle).

Fig. 3 presents photographs and a graph showing the size and morphological characteristics of the toxic A $\beta$  assemblies targeted by 1A9 and 2C3. A: The 540,000 x g supernatant of A $\beta$  1-42 (25  $\mu$ M) was subjected to a continuous molecular sieving process using ultrafiltration membranes having a molecular weight cutoff value of 3, 10, 30, and 100 kDa (Microcon). The four types of filtrates thus fractionated were named as follows: fraction 1 (<3 kDa), fraction 2 (3 to 10 kDa), fraction 3 (10 to 30 kDa), fraction 4 (30 to 100 kDa); and fraction 5 (>100 kDa) which was finally retained. The presence of A $\beta$  1-42 in each of the above-mentioned fractions was detected by 4G8 immunoblotting.

B: Representative images of NGF-treated PC12 (PC12N) cells treated with the five fractions at 37°C for 48 hours. The toxicity of each fraction was evaluated as described above for Fig. 2. C: The viability of cells treated with the 540,000 x g supernatant of A $\beta$  1-42 and the five fractions (fractions 1 to 5). Similar results were obtained from two independent experiments. The values are presented in percentage (mean  $\pm$  SD) with respect to the control. D: Dot blot analysis of the five fractions (fractions 1 to 5). The blots were reacted with A11, 1A9, 2C3, and 4G8. E: AFM images of the five fractions. In fraction 5 (Fr. 5) that had the strongest toxicity, ring-shaped and bead-shaped structures were observed in addition to granular molecules.

Fig. 4 presents photographs and graphs showing the activity of 1A9 and 2C3 to suppress A $\beta$  amyloid fibril formation. A: Amyloid fibril formation of A $\beta$  1-42 at various concentrations (10  $\mu$ M (open square), 25  $\mu$ M (filled diamond), and 50  $\mu$ M (open circle)) was monitored by ThT assay at 37°C for up to 72 hours. B: Coexisting antibody dose-dependent inhibition of amyloid fibril formation of A $\beta$  1-42 was observed for 2C3 (open circle). In contrast, the 1A9 (open square), 4G8 (filled triangle), and non-specific IgG (filled square) antibodies did not inhibit fibril-forming assembly of seed-free A $\beta$  1-42 (ThT-negative 540,000 x g supernatant). C: Coexisting antibody dose-dependent inhibition of fibril-forming assembly of A $\beta$  1-42 was observed for 2C3 (open circle), and nearly complete inhibition was observed also for 1A9 (open square) at 3  $\mu$ M. D: None of the test antibodies added after a 24-hour pre-incubation for A $\beta$  1-42 amyloid fibril formation could dissolve nor disassemble the A $\beta$  1-42 amyloid fibrils. E to G: EM images of A $\beta$  1-42 in the absence (Panel E) and presence of 2C3 (Panel F) and 1A9 (Panel G).

Fig. 5 presents photographs and graphs on toxicity-related A $\beta$  1-42 oligomers. A: Dot blot assay (upper half of Panel A): A $\beta$  1-42 monomers (25  $\mu$ M) were incubated for a specified time (0 to 72 hours) at 37°C, and immobilized onto a nitrocellulose membrane, and subjected to dot blot assay that uses A11, 1A9, 2C3, or 4G8. The emergence of immunoreactivity-positive structures for each antibody was tested. Immunoreactivity intensity analysis (lower half of Panel A): The results of dot blot assay were analyzed semiquantitatively using the Multi Gauge v 3.0 software (Fuji Film, Tokyo). To correlate the oligomer formation and amyloid fibril formation, the ThT fluorescence value (the right Y axis) was overlaid on the same time axis. B: The A $\beta$  1-42 assembly after 0-, 2-, 4-, and 24-hour incubation at 37°C, and the change in A $\beta$  1-42 assembly after further 48-hour incubation. The A $\beta$  1-42 assembly was detected by 4G8 immunoblotting. C: The toxic activity of the above-mentioned various A $\beta$  1-42 assemblies. The viability of nerve cells was determined by the LIVE/DEAD assay as described for Fig. 2. D: The anti-neurotoxic activity of 1A9 and 2C3 was evaluated using various A $\beta$  assemblies (the A $\beta$  1-42 assemblies formed at 37°C for 0 and 2 hours ("0h" and "2h"); and the ThT-positive supernatant collected after ultracentrifugation at 540,000 x g for two hours ("2h sup")). Representative images of PC12N cells exposed to various A $\beta$  1-42 assemblies in the absence or presence of the antibodies are shown in the left half of Panel D (a: "0h"; b: "2h"; c: "2h sup"; d: "2h sup" + IgG2b; e: "2h sup" + 1A9; f: "2h sup" + 2C3). The viability of cells exposed to various A $\beta$  1-42 assemblies in the absence or presence of the antibodies is presented in percentage (mean  $\pm$  SD) with respect to the control, and this is shown in the right half of Panel D. Compared to the "0h" A $\beta$  1-42 assembly, the "2h" A $\beta$  1-42 assembly lowered the neurotoxicity. "2h sup" recovered the neurotoxicity to a degree similar to that of the "0h" A $\beta$  1-42 assembly. Non-specific IgG2b could not block the neurotoxicity induction of the "2h sup" A $\beta$  1-42 assembly. Monoclonal 1A9 completely inhibited the "2h sup"-induced neurotoxicity, while the ability of 2C3 to inhibit the toxicity was slightly inferior. In the experiments using the two monoclonal antibodies (mAbs), the antitoxic activity of the mAbs was observed at a mAb:A $\beta$  mole ratio of 1 : <25 to 50. This suggests that structurally different 1A9- and 2C3-recognized oligomeric assemblies exist at a relatively low concentration.

Fig. 6 presents photographs and graphs showing that soluble 1A9- and 2C3-recognized oligomers exist in the human brain. Antibodies against A $\beta$  oligomers can detect senile plaques and vascular amyloids in AD brain only after pretreatment with Protease K. A: 1A9 staining; B: 2C3 staining; and C: A11 staining. D: 4G8 immunoblotting of 1A9- or 2C3-immunoprecipitated A $\beta$  in buffer-soluble AD brain (lanes 1, 2, 4, and 5) and healthy control brain (lanes 3 and 6). Representative results for 1A9 and 2C3 are shown in the left and right half of the panel, respectively. E and F: Semiquantitative analysis (with actin control) of soluble 1A9-immunoreactive 12-mer (Panel E) and soluble 2C3-immunoreactive 12-mer (Panel F) in the human entorhinal cortex obtained from 50 autopsy cases of a healthy elderly population (Braak NFT Stage I or II: n = 35; Braak NFT Stage III or IV: n = 13; and Braak NFT Stage >IV, AD cases: n = 2).

Fig. 7-1 present graphs showing that soluble 1A9- and 2C3-recognized oligomers exist in human CSF. Pooled whole cerebrospinal fluid (CSF) (AD = 10 and NC = 10) (Panels A and B) and pooled lipoprotein-depleted CSF (AD = 10, and NC = 10) (Panels C and D) were subjected to size exclusion chromatography (SEC). In Panels A and B, the collected fractions were analyzed for the distribution of A $\beta$  40 and A $\beta$  42 monomers by BNT77-BA27 and BNT77-BC05 ELISAs. Panels C and D show the presence of A $\beta$  40 and A $\beta$  42 oligomers captured by 1A9/2C3 mixed antibodies.

Fig. 7-2 is the continuation of Fig. 7-1. The amount of 1A9-recognized oligomeric assembly (1A9-BC05 and 1A9-BA27 ELISAs) and the amount of 2C3-recognized assembly (2C3-BC05 and 2C3-BA27 ELISAs) were measured for 12 AD cases (open circle) and 13 NC cases (filled circle) (Panels E and G). The oligomer/monomer ratio is shown



in Panels F (1A9) and H (2C3).

Fig. 8 presents graphs showing that the onset of memory impairment in Tg2576 mice can be prevented by passive immunization treatment. 13-month-old Tg2576 mice were divided into the following three groups to perform learning/behavior tests: PBS-administered group:  $n = 10$ ; 1A9-administered group:  $n = 13$ ; and 2C3-administered group:  $n = 11$ . All of the measured values were indicated as mean  $\pm$  SE. (A) Y-maze test. Spontaneous alteration behavior was monitored in each group during an eight-minute session of the Y-maze task. The results of one-way ANOVA were as follows:  $F(1, 52) = 3.09$ ,  $p < 0.05$ ; \*  $p < 0.05$  in the comparison with PBS-administered Tg2576 mice. (B) Novel object recognition test. The retention session was performed 24 hours after training. The exploratory preference in a ten-minute session in the novel object recognition test was determined in each group. The results of two-way ANOVA were as follows: training/retention,  $F(1, 64) = 31.53$ ,  $p < 0.01$ ; animal group,  $F(2, 64) \sim 7.49$ ,  $p < 0.01$ ; repeated training/retention by the animal group,  $F(2, 64) = 10.12$ ,  $p < 0.01$ ; \*\*  $p < 0.01$  in the comparison with the corresponding untrained mice, ##  $p < 0.01$  in the comparison with PBS-administered Tg2576 mice. (C) The swimming path length during a 60-second session of water maze test was measured for each group. The results of two-way ANOVA were as follows: trial,  $F(9, 320) = 20.46$ ,  $p < 0.01$ ; animal group,  $F(2, 320) = 12.59$ ,  $p < 0.01$ ; repeated trial by the animal group,  $F(18, 320) = 1.78$ ,  $p < 0.05$ ;  $p < 0.05$ , \*\*  $p < 0.01$  in the comparison with PBS-administered Tg2576 mice. Fear-conditioned learning test: Context-dependent (D) and clue-dependent freezing times (E) were determined. The results of two-way ANOVA were as follows: context-dependent test,  $F(2, 32) = 5.94$ ,  $p < 0.01$ ; clue-dependent test,  $F(2, 32) = 7.33$ ,  $p < 0.01$ ; \*  $p < 0.05$  and \*\*  $p < 0.01$  in the comparison with PBS-administered Tg2576 mice.

Fig. 9 presents graphs and a photograph showing that the brain A $\beta$  accumulation in Tg2576 can be prevented by passive immunotherapy. The hippocampus and cerebral cortex of three groups of 13-month-old Tg2576 mice (PBS-administered group,  $n = 10$ ; 1A9-administered group,  $n = 13$ ; and 2C3-administered group,  $n = 11$ ) were extracted in three continuous steps to prepare the buffer-soluble, SDS-soluble, and formic acid (FA)-extractable fractions. Each of the fractions was subjected to A $\beta$ -specific ELISAs (WAKO kit: BNT77-BA27 for A $\beta$  x-40; BNT77-BC05 for A $\beta$  x-42). The accumulation of A $\beta$  40 (SDS and FA) and A $\beta$  42 (SDS) was found to be significantly suppressed only in the 1A9-treated group. The accumulation-suppressing effect for the A11-positive oligomer (4-mer) was confirmed in the SDS-soluble cerebral cortex fractions from the two antibody-treated groups.

Fig. 10 presents photographs and graphs on A $\beta$  oligomers in the plasma and brain of Tg2576. A and B: As a result of ELISA analysis, no significant difference in the amount of A $\beta$  x-40 and A $\beta$  x-42 in the plasma was observed between the PBS-administered group and the immunotherapy group. C: Similarly, no difference in the A $\beta$  40/A $\beta$  42 ratio was observed among the three groups tested. D: As a result of dot blot analysis using pooled brain homogenates, no difference in the amount of physiological saline-soluble A11-positive oligomer was observed among the three groups tested. Hippocampus (left panel) and cerebral cortex (right panel). PBS-administered group,  $n = 10$ ; 1A9-administered group,  $n = 13$ ; and 2C3-administered group,  $n = 11$ . E: According to immunoblot analysis using the anti-oligomer A11 antibody, the immunoreactivity of the A $\beta$  tetramer in the SDS-extracted cerebral cortex fraction (right panel) was decreased in the 1A9- and 2C3-administered groups compared to the PBS-administered group. On the other hand, this was not observed in the hippocampus (left panel). F: Blood (albumin-depleted plasma, upper part of Panel F; albumin/lipoprotein-depleted plasma, lower part of Panel F) was pooled from each of the groups, and subjected to A11 dot blot analysis. As a result, the A11 immunoreactivity was found to be increased in the 1A9- and 2C3-administered groups compared to the PBS-administered group (Panel F). The proportion of the lipoprotein-bound form of 2C3-recognized oligomers was higher than that of 1A9-recognized oligomers (lower part of Panel F). Furthermore, the A11 immunoblotting also showed positive signals at approximately 200 kDa, and the immunoreactivity was clearly increased in the 1A9- and 2C3-administered groups compared to the PBS-administered group (Panel G). From these results, it is conceivable that the therapeutic effect selective only to target A $\beta$  oligomer molecules was obtained in the antibody-administered groups without affecting physiological molecules.

Fig. 11 presents photographs and graphs showing that senile plaque amyloid formation (A: A $\beta$ -specific antibody staining; and B: thioflavin-S-positive analysis) and swollen dystrophic neurite formation (C: synaptophysin-positive analysis) were suppressed in the Tg2576 mouse brain by passive immunization treatment.

Fig. 12 presents photographs showing the suppression of synaptic degeneration by passive immunization treatment with 1A9 and 2C3. Immunostaining of synaptophysin (left panels) and drebrin (right panels) in presynaptic and postsynaptic dot-like peripheral cells. Top: PBS administration; middle: 1A9 administration; and bottom: 2C3 administration.

Fig. 13 presents photographs showing the brain transfer of the antibodies by passive immunization treatment. The distribution of administered antibodies in the Tg2576 mouse brain is shown. Staining with anti-A $\beta$  antibodies (left panels) and IgG (center panels). 1A9 administration (A), 2C3 administration (B), and PBS administration (C).

Fig. 14 presents photographs showing, by dot blot analysis, that the monoclonal antibodies 5A5, 5A9, 4F7, 4H5, 6E4, and 6H4 are specific to A $\beta$  oligomers (3 to 96 hours), but do not recognize A $\beta$  monomers (0 hour).

Fig. 15 presents graphs showing the A $\beta$  oligomer-selective binding ability of the six types of antibodies (4F7, 4H5,

5A5, 5A9, 6E4, and 6H4). The vertical axis indicates the absorbance at a wavelength of 450 nm, and the horizontal axis indicates the concentration of A $\beta$  oligomer or A $\beta$  monomer used as an inhibitor. In each graph, the dashed line indicates the antibody-binding activity when the A $\beta$  oligomer was used as the inhibitor, and the solid line indicates the antibody-binding activity when the A $\beta$  monomer was used as the inhibitor.

Fig. 16 presents graphs showing the neutralizing activity of the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) against A $\beta$ -induced neurotoxicity. The horizontal axis indicates the amount of antibody added, and the vertical axis shows the cytotoxicity relative to that under the antibody-free condition as the standard (see the equation in the figure). Control IgG (3F1), which is an antibody that does not bind to A $\beta$  42, was used for comparison.

Fig. 17 presents graphs showing the suppressing activity of the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) against A $\beta$  amyloid fibril formation. The antibodies were added at three different concentrations to a A $\beta$  1-42 solution (12.5  $\mu$ M). After incubation at 37°C for 24 hours, the level of A $\beta$  amyloid fibril formation was measured by the ThT fluorescence intensity method. The horizontal axis indicates the amount of antibody added, and the vertical axis shows the level of amyloid fibril formation by the antibody addition that is relative to the level of amyloid fibril formation without antibody addition as the standard.

#### Mode for Carrying Out the Invention

**[0015]** The present invention will be described more specifically below.

**[0016]** As described above, the present inventors succeeded in obtaining antibodies that bind specifically to A $\beta$  oligomers but not to A $\beta$  monomers. That is, the present invention provides antibodies that bind to A $\beta$  oligomers but not to A $\beta$  monomers. The antibodies are preferably isolated or purified.

**[0017]** The terms "isolated" and "purified" used for substances (antibodies and such) of the present invention indicate that the substances do not substantially include at least one substance that may be contained in the natural source. Therefore, "isolated antibodies" and "purified antibodies" refer to antibodies that do not substantially include cell materials such as hydrocarbons, lipids, or other contaminant proteins from the cell or tissue source from which the antibodies (proteins) are derived. When the antibodies are chemically synthesized, the terms refer to antibodies that do not substantially include chemical precursor substances or other chemical substances. In a preferred embodiment, the antibodies of the present invention are isolated or purified.

**[0018]** "Antibodies" refers to glycoproteins that have the same structural characteristics. Antibodies show binding specificity towards specific antigens. Herein, "antigens" refers to proteins that have the ability to bind to the corresponding antibodies, and induce antigen-antibody reactions *in vivo*.

**[0019]** A $\beta$  proteins, which are the major constituents of amyloids, are peptides consisting of 40 to 42 amino acids, and are known to be produced from precursor proteins called amyloid precursor proteins (APPs) by the action of proteases. Besides amyloid fibrils collected in ultracentrifuged sediment fractions, the amyloid molecules produced from APPs include oligomeric non-fibrous assemblies in addition to soluble monomers. "A $\beta$  oligomers" as described herein refer to non-fibrous assemblies. The "A $\beta$  oligomers" as described herein include, for example, A $\beta$ 40 (A $\beta$  1-40) oligomers and A $\beta$ 42 (A $\beta$  1-42) oligomers. For example, "A $\beta$ 42 oligomers" as described herein are molecules showing a molecular weight of 45 to 160 kDa in SDS-PAGE, and 22.5 to 1,035 kDa in Blue Native PAGE. Using molecular sieves, the molecules are collected mainly in the >100 kDa retention solution. When observed under an atomic force microscope, the molecules show mixed morphologies of granular, bead-shaped, and ring-shaped molecules having a height of 1.5 to 3.1 nm. By the gel filtration method, the molecules were eluted in the void volume fraction 8 with a molecular weight of 680 kDa or more, and in fraction 15 with a molecular weight of 17 to 44 kDa.

**[0020]** There is no limitation on the origin and form of the antibodies used in the present invention as long as they bind to A $\beta$  oligomers but not to A $\beta$  monomers.

**[0021]** "Antibodies" of the present invention are monoclonal antibodies. The antibodies of the present invention also include any type of antibodies such as non-human animal antibodies, humanized antibodies, chimeric antibodies, human antibodies, the later-described minibodies, amino acid sequence-modified antibodies, modified antibodies conjugated to other molecules (for example, polymers such as polyethylene glycol), and sugar chain-modified antibodies.

**[0022]** Herein, the term "monoclonal antibodies" refers to antibodies that are obtained from a substantially homogeneous population of antibodies. That is, the individual antibodies constituting the population are identical with the exception of possible natural mutants that may be present in a trace amount. Monoclonal antibodies are highly specific and recognize a single antigenic site. Each of the monoclonal antibodies recognizes a single determinant of the antigen, in contrast to conventional (polyclonal) antibody preparations that typically contain different antibodies against different antigenic determinants (epitopes).

**[0023]** In addition to the above-mentioned specificity, monoclonal antibodies have the advantage that they can be synthesized from a hybridoma culture that is not contaminated with other immunoglobulins. Therefore, "monoclonal" indicates the characteristics of antibodies that can be obtained from a substantially homogeneous antibody population. This term does not indicate the requirement for any specific method for antibody production.

**[0024]** Basically, monoclonal antibodies can be produced by using known techniques. For example, they may be produced by the hybridoma method first described by Kohler and Milstein (Nature 256: 495-7, 1975), or by the recombinant DNA method (Cabilly et al., Proc. Natl. Acad. Sci. USA 81:3273-7, 1984), but the methods are not limited thereto. For example, when using the hybridoma method, an A $\beta$  oligomer (for example, the A $\beta$  tetramer described in the Examples) is used as a sensitizing antigen, and immunization is carried out according to a conventional immunization method. The obtained immune cells are fused with known parent cells by a conventional cell fusion method, and monoclonal antibody-producing cells can be screened and isolated using a conventional screening method.

**[0025]** The monoclonal antibodies of the present invention can be produced as follows. Synthetic A $\beta$  1-42 (Peptide Institute, Inc., Osaka) is dissolved in distilled deionized water or a 10 mM phosphate buffer solution, and this is incubated at 37°C for 18 hours. Then, the peptides are separated by 4-12% SDS-PAGE, and visualized by CBB staining, and the portion of the A $\beta$  1-42 tetramer alone which is not contaminated with the A $\beta$  1-42 monomer is cut out and used as an antigen. On the other hand, a preparation containing a large amount of the A $\beta$  1-40 oligomer is prepared by mixing (i) a modified A $\beta$  1-40 prepared by chemically linking 6-carboxytetramethylrhodamine (6-TAMRA) (SIGMA) to the N terminus of a synthetic A $\beta$  1-40 peptide using a conventional method with (ii) synthetic A $\beta$  1-40 (Peptide Institute, Inc., Osaka) at a ratio of 5:100, 10:100, 20:100, 30:100, 40:100, 50:100, 60:100, 70:100, or 80:100, preferably 90:100, or more preferably 100:100, and carrying out polymerization reaction for three hours, preferably six hours, or more preferably 20 hours. Next, Balb-c mice are immunized with 2.5  $\mu$ g of either the A $\beta$  1-42 tetramer or A $\beta$  1-40 oligomer emulsified using complete Freund's adjuvant by injecting the antigen into their foot pad. Subsequently, booster immunizations are carried out six times. Hybridomas are produced from the inguinal lymph node by fusion with Sp2/O-Ag14 cells using Polyethylene Glycol 1500.

**[0026]** The animals immunized with sensitizing antigens are not particularly limited, but are preferably selected considering the compatibility with parent cells used for cell fusion. Generally, rodents, lagomorphs, or primates are used. Rodents include, for example, mice, rats, and hamsters. Lagomorphs include, for example, rabbits. Primates include, for example, Catarrhini (old-world) monkeys such as *Macaca fascicularis*, *Macaca mulatta*, hamadryas, and chimpanzees.

**[0027]** Animals are immunized with sensitizing antigens according to known methods. For example, as a standard method, immunization is performed by intraperitoneal or subcutaneous injection of a sensitizing antigen into mammals.

**[0028]** An example of the parent cells fused with the aforementioned immunocytes is the Sp2/O-Ag14 cell, which will be described below in the Examples. However, various other known cell lines can be used.

**[0029]** Cell fusion between the aforementioned immunocyte and a myeloma cell can be carried out basically according to known methods including the method by Kohler and Milstein (Kohler G. and Milstein C., Methods Enzymol. (1981) 73, 3-46).

**[0030]** Hybridomas obtained in this manner are selected by culturing them in a conventional selection culture medium such as a HAT culture medium, which contains hypoxanthine, aminopterin, and thymidine. Culturing in the above-mentioned HAT culture medium is generally continued for several days to several weeks for an adequate time for killing cells other than the desired hybridomas (non-fused cells). Next, a conventional limiting dilution method is performed for screening and singly-cloning of a hybridoma that produces the desired antibody.

**[0031]** Thereafter, the obtained hybridoma is transplanted into the abdominal cavity of a mouse, and ascitic fluid containing the desired monoclonal antibodies is extracted. For example, the antibodies can be purified from the ascitic fluid by conventional protein separation and/or purification methods such as a selected combination of column chromatography including, but not limited to, affinity chromatography, filtration, ultrafiltration, salt precipitation, dialysis, SDS polyacrylamide gel electrophoresis, and isoelectric focusing (Antibodies: A Laboratory manual, Harlow and David, Lane (edit.), Cold Spring Harbor Laboratory, 1988).

**[0032]** Protein A columns and Protein G columns can be used for affinity columns. Examples of the Protein A columns used include Hyper D, POROS, and Sepharose F.F. (Pharmacia).

**[0033]** Chromatography (excluding affinity chromatography) includes ion exchange chromatography, hydrophobic chromatography, gel filtration, reverse-phase chromatography, and adsorption chromatography ("Strategies for Protein Purification and Characterization: A Laboratory Course Manual", Daniel R Marshak et al., Cold Spring Harbor Laboratory Press, 1996). When chromatography is carried out, liquid-phase chromatography methods such as HPLC and FPLC can be used.

**[0034]** Monoclonal antibody-producing hybridomas prepared in this manner can be subcultured in a conventional culture medium, and they can be stored for a long time in liquid nitrogen.

**[0035]** Any mammal can be immunized using an immunogen for antibody production. However, when preparing monoclonal antibodies by producing hybridomas, the compatibility with parent cells used in cell fusion for hybridoma production is preferably considered.

**[0036]** Generally, rodents, lagomorphs, or primates are used for the immunization. Rodents include, for example, mice, rats, and hamsters. Lagomorphs include, for example, rabbits. Primates include, for example, Catarrhini (old-world) monkeys such as *Macaca fascicularis*, *Macaca mulatta*, hamadryas, and chimpanzees. The use of transgenic

animals that have a human antibody gene repertoire is known in the art (Ishida I, et al., Cloning and Stem Cells 4: 91-102, 2002). As with other animals, to obtain human monoclonal antibodies, the transgenic animals are immunized, then antibody-producing cells are collected from the animals and fused with myeloma cells to produce hybridomas, and anti-protein human antibodies can be prepared from these hybridomas (see International Publication Nos. WO92/03918, WO94/02602, WO94/25585, WO96/33735, and WO96/34096).

**[0037]** Alternatively, lymphocytes immortalized with oncogenes may be used for monoclonal antibody production. For example, human lymphocytes infected with EB virus or such is immunized *in vitro* with immunogens. Next, the immunized lymphocytes are fused with human-derived myeloma cells (U266, etc) capable of unlimited division, and thus hybridomas that produce the desired human antibodies are obtained (Japanese Patent Application Kokai Publication No. (JP-A) S63-17688 (unexamined, published Japanese patent application)).

**[0038]** Once monoclonal antibodies can be obtained by any of the aforementioned methods, the antibodies may also be prepared using genetic engineering methods (see, for example, Borrebaeck CAK and Larrick JW, Therapeutic Monoclonal Antibodies, MacMillan Publishers, UK, 1990). For example, recombinant antibodies may be prepared by cloning DNAs that encode the desired antibodies from antigen-producing cells such as hybridomas or immunized lymphocytes that produce the antibodies, then inserting the cloned DNAs into appropriate vectors, and transfecting the vectors into suitable host cells. Such recombinant antibodies are also included in the present invention.

**[0039]** Examples of the monoclonal antibodies of the present invention include the 4H5 monoclonal antibody. Preferably, the monoclonal antibodies of the invention include an antibody that binds to an A $\beta$  oligomer but not to an A $\beta$  monomer, wherein the antibody is a monoclonal antibody and is an antibody that comprises an H chain (heavy chain) having the amino acid sequence of SEQ ID NO: 61 and an L chain (light chain) having the amino acid sequence of SEQ ID NO: 63.

In an embodiment, the antibodies of the present invention include minibodies. A minibody contains an antibody fragment lacking a portion of a whole antibody, and is not particularly limited as long as it has the ability to bind to an antigen. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv. Examples of minibodies include Fab, Fab', F(ab')<sub>2</sub>, Fv, scFv (single chain Fv), diabody, and sc(Fv)<sub>2</sub> (single chain (Fv)<sub>2</sub>).

**[0040]** To obtain disclosed polyclonal antibodies against the proteins, blood is removed from a mammal sensitized with an antigen after the serum level of the desired antibody is confirmed to be increased. Serum is separated from blood by a known method. When a polyclonal antibody is used, serum containing the polyclonal antibody may be utilized. Alternatively, if necessary, a fraction containing the polyclonal antibody may be isolated from serum and then used. For example, immunoglobulin G or M can be prepared by obtaining a fraction that specifically recognizes a protein disclosed herein using an affinity column coupled with the protein, and then purifying this fraction using a Protein A or Protein G column.

**[0041]** Disclosed is further an antibody that binds to an A $\beta$  oligomer which is an antibody binding to an A $\beta$  oligomer that binds 1A9, 2C3, 5A5, 5A9, 4F7, 4H5, 6E4, or 6H4. Preferably, the disclosed antibody is any one of the antibodies of (A) to (F) below:

(A) an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 1 and an L chain having the amino acid sequence of SEQ ID NO: 3;

(B) an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 21 and an L chain having the amino acid sequence of SEQ ID NO: 23;

(C) an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 41 and an L chain having the amino acid sequence of SEQ ID NO: 43;

(D) an antibody of the present invention binding to an A $\beta$  oligomers that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 61 and an L chain having the amino acid sequence of SEQ ID NO: 63;

(E) an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 81 and an L chain having the amino acid sequence of SEQ ID NO: 83; and

(F) an antibody binding to an A $\beta$  oligomer that binds to an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 101 and an L chain having the amino acid sequence of SEQ ID NO: 103.

**[0042]** Furthermore, A $\beta$  oligomers are disclosed to which the antibodies bind. Preferably, the antibodies include, for example, the 4H5 monoclonal antibody. Such A $\beta$  oligomers can be used as antigens for preparing antibodies, or vaccines.

**[0043]** In a preferred embodiment, the antibodies disclosed include, for example, the antibody of any one of (1) to (38) below:

(1) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 9 as CDR1, the amino acid sequence of SEQ ID NO: 11 as CDR2, and the amino acid sequence of SEQ ID NO: 13 as CDR3;

(2) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 15 as CDR1, the amino acid sequence of SEQ ID NO: 17 as CDR2, and the amino acid sequence of SEQ ID NO: 19 as CDR3;

- (3) an antibody that comprises the H chain of (1) and the L chain of (2);  
 (4) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 5 as VH;  
 (5) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 7 as VL;  
 (6) an antibody that comprises the H chain of (4) and the L chain of (5);  
 5 (7) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 29 as CDR1, the amino acid sequence of SEQ ID NO: 31 as CDR2, and the amino acid sequence of SEQ ID NO: 33 as CDR3;  
 (8) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 35 as CDR1, the amino acid sequence of SEQ ID NO: 37 as CDR2, and the amino acid sequence of SEQ ID NO: 39 as CDR3;  
 (9) an antibody that comprises the H chain of (7) and the L chain of (8);  
 10 (10) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 25 as VH;  
 (11) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 27 as VL;  
 (12) an antibody that comprises the H chain of (10) and the L chain of (11);  
 (13) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 49 as CDR1, the amino acid sequence of SEQ ID NO: 51 as CDR2, and the amino acid sequence of SEQ ID NO: 53 as CDR3;  
 15 (14) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 55 as CDR1, the amino acid sequence of SEQ ID NO: 57 as CDR2, and the amino acid sequence of SEQ ID NO: 59 as CDR3;  
 (15) an antibody that comprises the H chain of (13) and the L chain of (14);  
 (16) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 45 as VH;  
 (17) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 47 as VL;  
 20 (18) an antibody that comprises the H chain of (16) and the L chain of (17);  
 (19) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 69 as CDR1, the amino acid sequence of SEQ ID NO: 71 as CDR2, and the amino acid sequence of SEQ ID NO: 73 as CDR3;  
 (20) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 75 as CDR1, the amino acid sequence of SEQ ID NO: 77 as CDR2, and the amino acid sequence of SEQ ID NO: 79 as CDR3;  
 25 (21) an antibody that comprises the H chain of (19) and the L chain of (20);  
 (22) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 65 as VH;  
 (23) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 67 as VL;  
 (24) an antibody that comprises the H chain of (22) and the L chain of (23);  
 (25) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 89 as CDR1, the amino acid sequence of SEQ ID NO: 91 as CDR2, and the amino acid sequence of SEQ ID NO: 93 as CDR3;  
 30 (26) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 95 as CDR1, the amino acid sequence of SEQ ID NO: 97 as CDR2, and the amino acid sequence of SEQ ID NO: 99 as CDR3;  
 (27) an antibody that comprises the H chain of (25) and the L chain of (26);  
 (28) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 85 as VH;  
 35 (29) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 87 as VL;  
 (30) an antibody that comprises the H chain of (28) and the L chain of (29);  
 (31) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 109 as CDR1, the amino acid sequence of SEQ ID NO: 111 as CDR2, and the amino acid sequence of SEQ ID NO: 113 as CDR3;  
 (32) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 115 as CDR1, the amino acid sequence of SEQ ID NO: 117 as CDR2, and the amino acid sequence of SEQ ID NO: 119 as CDR3;  
 40 (33) an antibody that comprises the H chain of (31) and the L chain of (32);  
 (34) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 105 as VH;  
 (35) an antibody that comprises an L chain having the amino acid sequence of SEQ ID NO: 107 as VL;  
 (36) an antibody that comprises the H chain of (34) and the L chain of (35);  
 45 (37) an antibody that comprises one or more amino acid substitutions, deletions, additions, and/or insertions in the antibody of any one of (1) to (36), which has equivalent activity as the antibody of any one of (1) to (36); and  
 (38) an antibody that binds to the epitope bound by the antibody of any one of (1) to (36).

**[0044]** An example of the VH in the above-mentioned "H chain having the amino acid sequence of SEQ ID NO: 9 (sequence of the 5A5 antibody H-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 11 (sequence of the 5A5 antibody H-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 13 (sequence of the 5A5 antibody H-chain CDR3) as CDR3" of (1) is a VH having the amino acid sequence of SEQ ID NO: 5 (sequence of the 5A5 antibody VH).

**[0045]** An example of the VL in the above-mentioned "L chain having the amino acid sequence of SEQ ID NO: 15 (sequence of the 5A5 antibody L-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 17 (sequence of the 5A5 antibody L-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 19 (sequence of the 5A5 antibody L-chain CDR3) as CDR3" of (2) is a VL having the amino acid sequence of SEQ ID NO: 7 (sequence of the 5A5 antibody VL).

**[0046]** An example of the VH in the above-mentioned "H chain having the amino acid sequence of SEQ ID NO: 29

(sequence of the 5A9 antibody H-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 31 (sequence of the 5A9 antibody H-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 33 (sequence of the 5A9 antibody H-chain CDR3) as CDR3"of (7) is a VH having the amino acid sequence of SEQ ID NO: 25 (sequence of the 5A9 antibody VH).

**[0047]** An example of the VL in the above-mentioned "L chain having the amino acid sequence of SEQ ID NO: 35 (sequence of the 5A9 antibody L-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 37 (sequence of the 5A9 antibody L-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 39 (sequence of the 5A9 antibody L-chain CDR3) as CDR3"of (8) is a VL having the amino acid sequence of SEQ ID NO: 27 (sequence of the 5A9 antibody VL).

**[0048]** An example of the VH in the above-mentioned "H chain having the amino acid sequence of SEQ ID NO: 49 (sequence of the 4F7 antibody H-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 51 (sequence of the 4F7 antibody H-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 53 (sequence of the 4F7 antibody H-chain CDR3) as CDR3"of (13) is a VH having the amino acid sequence of SEQ ID NO: 45 (sequence of the 4F7 antibody VH).

**[0049]** An example of the VL in the above-mentioned "L chain having the amino acid sequence of SEQ ID NO: 55 (sequence of the 4F7 antibody L-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 57 (sequence of the 4F7 antibody L-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 59 (sequence of the 4F7 antibody L-chain CDR3) as CDR3"of (14) is a VL having the amino acid sequence of SEQ ID NO: 47 (sequence of the 4F7 antibody VL).

**[0050]** An example of the VH in the above-mentioned "H chain having the amino acid sequence of SEQ ID NO: 69 (sequence of the 4H5 antibody H-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 71 (sequence of the 4H5 antibody H-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 73 (sequence of the 4H5 antibody H-chain CDR3) as CDR3"of (19) is a VH having the amino acid sequence of SEQ ID NO: 65 (sequence of the 4H5 antibody VH).

**[0051]** An example of the VL in the above-mentioned "L chain having the amino acid sequence of SEQ ID NO: 75 (sequence of the 4H5 antibody L-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 77 (sequence of the 4H5 antibody L-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 79 (sequence of the 4H5 antibody L-chain CDR3) as CDR3"of (20) is a VL having the amino acid sequence of SEQ ID NO: 67 (sequence of the 4H5 antibody VL).

**[0052]** An example of the VH in the above-mentioned "H chain having the amino acid sequence of SEQ ID NO: 89 (sequence of the 6E4 antibody H-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 91 (sequence of the 6E4 antibody H-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 93 (sequence of the 6E4 antibody H-chain CDR3) as CDR3"of (25) is a VH having the amino acid sequence of SEQ ID NO: 85 (sequence of the 6E4 antibody VH).

**[0053]** An example of the VL in the above-mentioned "L chain having the amino acid sequence of SEQ ID NO: 95 (sequence of the 6E4 antibody L-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 97 (sequence of the 6E4 antibody L-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 99 (sequence of the 6E4 antibody L-chain CDR3) as CDR3"of (26) is a VL having the amino acid sequence of SEQ ID NO: 87 (sequence of the 6E4 antibody VL).

**[0054]** An example of the VH in the above-mentioned "H chain having the amino acid sequence of SEQ ID NO: 109 (sequence of the 6H4 antibody H-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 111 (sequence of the 6H4 antibody H-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 113 (sequence of the 6H4 antibody H-chain CDR3) as CDR3"of (31) is a VH having the amino acid sequence of SEQ ID NO: 105 (sequence of the 6H4 antibody VH).

**[0055]** An example of the VL in the above-mentioned "L chain having the amino acid sequence of SEQ ID NO: 115 (sequence of the 6H4 antibody L-chain CDR1) as CDR1, the amino acid sequence of SEQ ID NO: 117 (sequence of the 6H4 antibody L-chain CDR2) as CDR2, and the amino acid sequence of SEQ ID NO: 119 (sequence of the 6H4 antibody L-chain CDR3) as CDR3"of (32) is a VL having the amino acid sequence of SEQ ID NO: 107 (sequence of the 6H4 antibody VL).

**[0056]** For the disclosed 5A5 antibody, the amino acid sequence and the nucleotide sequence of the full-length H chain are shown in SEQ ID NO: 1 and SEQ ID NO: 2, respectively; the amino acid sequence and the nucleotide sequence of the full-length L chain are shown in SEQ ID NO: 3 and SEQ ID NO: 4, respectively; the amino acid sequence and the nucleotide sequence of the H-chain variable region (VH) are shown in SEQ ID NO: 5 and SEQ ID NO: 6, respectively; the amino acid sequence and the nucleotide sequence of the L-chain variable region (VL) are shown in SEQ ID NO: 7 and SEQ ID NO: 8, respectively; the amino acid sequence and the nucleotide sequence of the H-chain CDR1 are shown in SEQ ID NO: 9 and SEQ ID NO: 10, respectively; the amino acid sequence and the nucleotide sequence of the H-chain CDR2 are shown in SEQ ID NO: 11 and SEQ ID NO: 12, respectively; the amino acid sequence and the nucleotide sequence of the H-chain CDR3 are shown in SEQ ID NO: 13 and SEQ ID NO: 14, respectively; the amino acid sequence



sequence of the full-length L chain are shown in SEQ ID NO: 103 and SEQ ID NO: 104, respectively; the amino acid sequence and the nucleotide sequence of the H-chain variable region (VH) are shown in SEQ ID NO: 105 and SEQ ID NO: 106, respectively; the amino acid sequence and the nucleotide sequence of the L-chain variable region (VL) are shown in SEQ ID NO: 107 and SEQ ID NO: 108, respectively; the amino acid sequence and the nucleotide sequence of the H-chain CDR1 are shown in SEQ ID NO: 109 and SEQ ID NO: 110, respectively; the amino acid sequence and the nucleotide sequence of the H-chain CDR2 are shown in SEQ ID NO: 111 and SEQ ID NO: 112, respectively; the amino acid sequence and the nucleotide sequence of the H-chain CDR3 are shown in SEQ ID NO: 113 and SEQ ID NO: 114, respectively; the amino acid sequence and the nucleotide sequence of the L-chain CDR1 are shown in SEQ ID NO: 115 and SEQ ID NO: 116, respectively; the amino acid sequence and the nucleotide sequence of the L-chain CDR2 are shown in SEQ ID NO: 117 and SEQ ID NO: 118, respectively; and the amino acid sequence and the nucleotide sequence of the L-chain CDR3 are shown in SEQ ID NO: 119 and SEQ ID NO: 120, respectively.

**[0062]** The above-mentioned antibodies of (1) to (38) including the antibodies of the present invention include not only monovalent antibodies but also multivalent antibodies with two or more valencies. The multivalent antibodies of the present invention include multivalent antibodies whose antigen binding sites are all the same and multivalent antibodies whose antigen binding sites are partially or completely different.

**[0063]** Preferably, the above-mentioned antibody of (37) is an antibody with no modified CDRs. For example, the "antibody that comprises one or more amino acid substitutions, deletions, additions, and/or insertions in the antibody of (1), which has equivalent activity as the antibody of (1)" of the above-mentioned antibody of (37) is preferably "an antibody that has equivalent activity as the antibody of (1), and comprises one or more amino acid substitutions, deletions, additions, and/or insertions in the antibody of (1), and comprises an H chain having the amino acid sequence of SEQ ID NO: 9 as CDR1, the amino acid sequence of SEQ ID NO: 11 as CDR2, and the amino acid sequence of SEQ ID NO: 13 as CDR3". Another antibody of the above-mentioned antibody of (37) can be expressed in a similar manner.

**[0064]** Herein, "equivalent activity" means that the antibody of interest has biological or biochemical activity similar to that of an antibody of the present disclosure. Examples of the "activity" as described herein include the activity to bind specifically to A $\beta$  oligomers but not to A $\beta$  monomers, anti-neurotoxic activity, activity to suppress A $\beta$  amyloid fibril formation, anti-synaptic toxicity activity, and anti-memory impairment activity.

**[0065]** Methods for preparing a polypeptide having activity equivalent to that of a certain polypeptide that are well known to those skilled in the art include methods for introducing mutations into a polypeptide. For example, one skilled in the art can prepare an antibody having activity equivalent to that of an antibody of the present invention by introducing appropriate mutations into the antibody using site-directed mutagenesis (Hashimoto-Gotoh, T. et al. (1995) Gene 152, 271-275; Zoller, MJ, and Smith, M. (1983) Methods Enzymol. 100, 468-500; Kramer, W. et al. (1984) Nucleic Acids Res. 12, 9441-9456; Kramer W, and Fritz HJ (1987) Methods. Enzymol. 154, 350-367; Kunkel, TA (1985) Proc. Natl. Acad. Sci. USA. 82, 488-492; Kunkel (1988) Methods Enzymol. 85, 2763-2766) and such. Amino acid mutations may also occur naturally. The antibodies of the present disclosure also include an antibody that comprises an amino acid sequence with one or more amino acid mutations in the amino acid sequence of an antibody of the present invention, and which has activity equivalent to that of the antibody of the present invention. The number of mutated amino acids in such mutants may be generally 50 amino acids or less, preferably 30 amino acids or less, and more preferably ten amino acids or less (for example, five amino acids or less).

**[0066]** Amino acid residues are preferably mutated into other amino acids that conserve the properties of the amino acid side chains. For example, amino acids are categorized as follows depending on the side chain properties: hydrophobic amino acids (A, I, L, M, F, P, W, Y, and V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S, and T), amino acids with aliphatic side chains (G, A, V, L, I, and P), amino acids with hydroxyl-containing side chains (S, T, and Y), amino acids with sulfur atom-containing side chains (C and M), amino acids with carboxylic acid- and amide-containing side chains (D, N, E, and Q), amino acids with basic side chains (R, K, and H), and amino acids with aromatic ring-containing side chains (H, F, Y, and W) (amino acids are represented by one-letter codes in parentheses).

**[0067]** A polypeptide having an amino acid sequence, in which one or more amino acid residues are modified (deleted, added, and/or substituted with other amino acids) in a certain amino acid sequence, is known to retain its original biological activity (function).

**[0068]** In addition to the above-mentioned modifications, the antibodies of the present invention may be conjugated to other substances as long as the activity is maintained. Examples of the substances include peptides, lipids, sugars and sugar chains, acetyl groups, and natural and synthetic polymers. These modifications may be performed to confer additional functions to the antibodies, or to stabilize the antibodies.

**[0069]** Antibodies in which several amino acid residues have been added to the amino acid sequence of an antibody of the present invention include fusion proteins containing the antibody. In the fusion proteins, the antibody is fused with another peptide or protein. Methods for producing a fusion protein can be carried out by ligating a polynucleotide encoding an antibody of the present invention in frame with a polynucleotide encoding another peptide or polypeptide, and inserting this into an expression vector, and expressing the fusion construct in a host. Techniques known to those skilled in the art can be used for this purpose. The peptides or polypeptides fused with an antibody of the present invention include,



for example, known peptides such as FLAG (Hopp, T.P. et al., BioTechnology (1988) 6, 1204-1210), 6x His consisting of six histidine (His) residues, 10x His, Influenza hemagglutinin (HA), human c-myc fragments, VSV-GP fragments, p18HIV fragments, T7-tag, HSV-tag, E-tag, SV40T antigen fragments, Ick tag,  $\alpha$ -tubulin fragments, B-tag, and Protein C fragments; glutathione-S-transferase (GST); immunoglobulin constant regions;  $\beta$ -galactosidase; and maltose-binding protein (MBP), etc. Commercially available polynucleotides encoding these peptides or polypeptides can be fused with polynucleotides encoding the antibodies of the present invention, and the fusion polypeptides can be produced by expressing the fusion polynucleotides thus prepared.

**[0070]** The antibodies of the present invention may differ in the amino acid sequence, molecular weight, presence or absence of sugar chains, structure and such, depending on the cell or host producing the antibodies or the purification method.

**[0071]** Antibodies that bind to an epitope to which an antibody of any one of (1) to (36) above binds can be obtained by methods known to those skilled in the art. For example, the antibodies can be obtained by (i) determining the epitope bound by the antibody of any one of (1) to (36) using a conventional method, and producing the antibodies using a polypeptide comprising an amino acid sequence included in the epitope as an immunogen; or (ii) determining the epitopes of antibodies produced by a conventional method, and selecting antibodies whose epitope is the same as that of the antibody of any one of (1) to (36).

**[0072]** The above-mentioned antibodies of (1) to (38) also include any type of antibodies such as the above-described minibodies, antibodies with modified amino acid sequences such as humanized antibodies and chimeric antibodies, non-human animal antibodies, human antibodies, modified antibodies conjugated to other molecules (for example, polymers such as polyethylene glycol), and sugar chain-modified antibodies.

**[0073]** In a preferred embodiment, the antibodies of the present invention are modified antibodies such as chimeric antibodies and humanized antibodies. Examples of preferred antibodies include (i) a chimeric antibody whose variable region is derived from the 4H5 antibody, and whose constant region is derived from a human immunoglobulin; and (ii) a humanized antibody whose CDR is derived from the 4H5 antibody, and whose FR is derived from a human immunoglobulin, and whose constant region is derived from a human immunoglobulin. These modified antibodies can be produced using known methods.

**[0074]** Since the antigenicity of a chimeric antibody or a humanized antibody in the human body is reduced, such an antibody is useful for administration to humans for therapeutic purposes or such.

**[0075]** Chimeric antibodies are produced by combining sequences derived from different animals. Examples of chimeric antibodies include antibodies comprising the heavy-chain and light-chain variable regions of a mouse antibody and the heavy-chain and light-chain constant regions of a human antibody. The production of chimeric antibodies can be carried out using known methods (see, for example, Jones et al., Nature 321:522-5, 1986; Riechmann et al., Nature 332:323-7, 1988; and Presta, Curr. Opin. Struct. Biol. 2:593-6, 1992). For example, first, genes encoding the variable regions or CDRs of the antibody of interest are prepared from the RNAs of antibody-producing cells by polymerase chain reaction (PCR) or such (see, for example, Larrick et al., "Methods: a Companion to Methods in Enzymology", Vol. 2: 106, 1991; Courtenay-Luck, "Genetic Manipulation of Monoclonal Antibodies" in Monoclonal Antibodies: Production, Engineering and Clinical Application; Ritter et al. (eds.), page 166, Cambridge University Press, 1995, and Ward et al., "Genetic Manipulation and Expression of Antibodies" in Monoclonal Antibodies: Principles and Applications; and Birch et al. (eds.), page 137, Wiley-Liss, Inc., 1995). The prepared genes encoding the variable regions are linked to genes encoding the constant regions or framework regions. The genes encoding the constant regions or framework regions may be determined in a manner similar to that for the CDR-encoding genes, or alternatively, they can be prepared based on the sequence information of known antibodies. DNA sequences encoding chimeric products and CDR-grafted products may be synthesized completely or partially using oligonucleotide synthesis techniques. For example, the oligonucleotide synthesis described by Jones et al. (Nature 321:522-5, 1986) may be performed. Furthermore, in some cases, site-directed mutagenesis and polymerase chain reaction techniques may be appropriately used. Techniques for oligonucleotide-specific mutagenesis of cases, site-directed mutagenesis and polymerase chain reaction techniques may be appropriately used. Techniques for oligonucleotide-specific mutagenesis of known variable regions described by Verhoeven et al. (Science 239: 1534-6, 1988) and Riechmann et al. (Nature 332: 323-7, 1988) may be used for modifying the variable region sequences, for example, to enhance the binding ability of chimeric antibodies. Furthermore, if necessary, enzymatic fill-in of gapped oligonucleotides using T4 DNA polymerase may be performed, for example, as described by Queen et al. (Proc. Natl. Acad. Sci. USA 86: 10029-33, 1989; and WO 90/07861).

**[0076]** For example, CDR-grafting techniques are known in the art ("Immunoglobulin genes", Academic Press (London), pp 260-74, 1989; and Michael A et al., Proc. Natl. Acad. Sci. USA 91: 969-73, 1994). Using the techniques, the CDRs of a certain antibody are replaced with the CDRs of another antibody. Through such replacement, the binding specificity of the former antibody is changed to that of the latter antibody. Among such chimeric antibodies, those in which the framework amino acids are derived from a human antibody are called "humanized antibodies (CDR-grafted antibodies)". When using antibodies to treat humans, human antibodies or humanized antibodies are preferably utilized.

**[0077]** Generally, chimeric antibodies comprise the variable regions of a non-human mammal-derived antibody and

the constant regions derived from a human antibody. On the other hand, humanized antibodies comprise the complementarity-determining regions of a non-human mammal-derived antibody and the framework regions and constant regions derived from a human antibody.

**[0078]** After producing the chimeric antibodies or humanized antibodies, amino acids in the variable regions (for example, FRs) or the constant regions may be substituted with other amino acids.

**[0079]** The origin of the variable regions of the chimeric antibodies or the CDRs of the humanized antibodies is not particularly limited.

**[0080]** Human antibody-derived C-regions are used for the C-regions of the chimeric antibodies and humanized antibodies. For example, C $\gamma$ 1, C $\gamma$ 2, C $\gamma$ 3, C $\gamma$ 4, C $\mu$ , C $\delta$ , C $\alpha$ 1, C $\alpha$ 2, and C $\epsilon$  can be used for the H-chain C-regions, and C $\kappa$  and C $\lambda$  can be used for the L-chain C-regions. Their sequences are known. Furthermore, the human antibody C regions can be modified to improve the stability of the antibodies or their production.

**[0081]** The binding activity of the antibodies of the present invention to the antigens (A $\beta$  oligomers) can be measured using, for example, an absorbance measurement method, an enzyme-linked immunosorbent assay (ELISA) method, an enzyme immunoassay (EIA) method, a radioimmunoassay (RIA) method, and/or a fluoroimmunoassay method. In ELISA, an antibody is immobilized on a plate, and an antigen for the antibody is added to the plate, then a sample containing the desired antibody, such as the culture supernatant of antibody-producing cells or a purified antibody is added. Next, a secondary antibody which recognizes the primary antibody and is tagged with an enzyme such as alkaline phosphatase is added to the plate, and this is preincubated. After washing, an enzyme substrate such as *p*-nitrophenyl phosphate is added to the plate, and the absorbance is measured to evaluate the antigen-binding ability of the sample of interest. The evaluation can be performed using BIAcore (Pharmacia).

**[0082]** Furthermore, the present invention provides compositions comprising the above-mentioned antibody of the present invention and a pharmaceutically acceptable carrier.

**[0083]** As described below, the present invention strongly suggests that monoclonal 1A9 and 2C3 antibody are promising candidates for therapeutic antibodies for preventing Alzheimer-like phenotypes. Memory deterioration has been shown to be related to synaptic dysfunction caused by soluble A $\beta$  oligomers (Klein WL, 2001, Trends Neurosci; and Selkoe DJ, 2002, Science). Excessive accumulation and deposition of A $\beta$  oligomers may trigger the complicated downstream cascades that cause Alzheimer's disease. If this is the case, therapeutic intervention using a composition comprising an antibody of the present invention and a pharmaceutically acceptable carrier could be effective for blocking the pathologic cascades, and thus this could enable the treatment of Alzheimer's disease.

**[0084]** The "treatment" of the present invention does not necessarily have complete therapeutic or preventive effects against organs or tissues exhibiting symptoms of disorders or diseases, but may have partial effects.

**[0085]** "Treatment of Alzheimer's disease" in the present invention means amelioration of at least one symptom that may be caused by Alzheimer's disease, and examples include amelioration or suppression of cognitive impairment, amelioration or suppression of senile plaque formation, amelioration or suppression of synaptic dysfunction, and reduction or suppression of A $\beta$  accumulation in brain tissues, blood, or such. Herein, "cognitive impairment" includes, for example, memory impairment including long term/short term memory impairment, object recognition memory impairment, spatial memory impairment, and associative and emotional memory impairment.

**[0086]** The present invention provides pharmaceutical compositions or pharmaceutical agents which comprise as an active ingredient the above-described composition comprising an antibody of the present invention and a pharmaceutically acceptable carrier.

**[0087]** In the present invention, the phrase "comprising as an active ingredient the above-described composition comprising an antibody of the present invention and a pharmaceutically acceptable carrier" means comprising the above-described composition comprising an antibody of the present invention and a pharmaceutically acceptable carrier as a major ingredient, but does not limit its content rate.

**[0088]** Examples of the above-mentioned pharmaceutical compositions include agents against cognitive impairment, Alzheimer's disease agents, agents for suppressing the progression of Alzheimer's disease, agents for suppressing senile plaque formation, agents for suppressing A $\beta$  accumulation, anti-neurotoxic agents (agents for neutralizing neurotoxicity), agents for inhibiting A $\beta$  amyloid fibril formation, and anti-synaptic toxicity agents (agents for neutralizing synaptic toxicity).

**[0089]** The above-mentioned pharmaceutical composition of the present invention can be used, for example, in methods for suppressing Alzheimer's disease which comprise the step of administering to a subject (individual) the above-described composition comprising an antibody of the present invention and a pharmaceutically acceptable carrier. Other examples of a pharmaceutical composition of the present invention for use include methods for suppressing cognitive impairment, methods for suppressing the progression of Alzheimer's disease, methods for suppressing senile plaque formation, methods for suppressing A $\beta$  accumulation, methods for neutralizing (suppressing) neurotoxic activity, methods for inhibiting A $\beta$  amyloid fibril formation, and methods for neutralizing (suppressing) synaptic toxicity. Further examples of a pharmaceutical composition of the present invention for use include the use in methods for preventing and/or treating cognitive impairment, and methods for preventing and/or treating Alzheimer's disease.

**[0090]** The present invention also provides use of a composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier in the production of the above-mentioned pharmaceutical composition.

**[0091]** Furthermore, the present invention relates to the following compositions.

- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in preventing and/or treating cognitive impairment.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in preventing and/or treating Alzheimer's disease.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in suppressing the progression of Alzheimer's disease.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in suppressing senile plaque formation.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in suppressing A $\beta$  accumulation.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in neutralizing (suppressing) neurotoxic activity.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in inhibiting A $\beta$  amyloid fibril formation.
- A composition comprising the above-described antibody of the present invention and a pharmaceutically acceptable carrier for use in neutralizing (suppressing) synaptic toxicity.

**[0092]** The above-mentioned pharmaceutical agents of the present invention can be administered to humans or other animals. In the present invention, non-human animals to which the pharmaceutical agents are administered include mice, rats, guinea pigs, rabbits, chickens, cats, dogs, sheep, pigs, cattle, monkeys, baboons, and chimpanzees. These animals preferably exhibit at least one symptom selected from, for example, cognitive impairment, senile plaque formation, synaptic dysfunction, A $\beta$  accumulation in brain tissues or blood, etc.

**[0093]** Antibodies contained in the pharmaceutical compositions of the present invention are not particularly limited as long as they are included in the above-mentioned antibodies of the present invention, and examples include the antibodies described herein.

**[0094]** When using the above-mentioned antibodies of the present invention for pharmaceutical compositions, they may be formulated by methods known to those skilled in the art. For example, as necessary, they can be prepared in the form of injectable sterile solutions or suspensions using water or another pharmaceutically acceptable liquid, and can be administered parenterally. For example, the antibodies to be included in the pharmaceutical compositions can be combined with acceptable carriers or media, specifically, sterile water, physiological saline, vegetable oils, emulsifiers, suspensions, surfactants, stabilizers, flavoring agents, excipients, solvents, preservatives, binders, or such, and mixed into a unit dose form required for generally accepted pharmaceutical practice. The phrase "pharmaceutically acceptable" indicates that the substance is inactive, and contains conventional substances used as diluents or vehicles for pharmaceuticals. Suitable excipients and their formulations are described, for example, in Remington's Pharmaceutical Sciences, 16th ed. (1980) Mack Publishing Co., ed. Oslo *et al.*

**[0095]** Physiological saline and other isotonic solutions containing glucose or adjuvants (for example, D-sorbitol, D-mannose, D-mannitol, and sodium chloride) can be used as aqueous solutions for injection. They can be used together with appropriate solubilizers such as alcohols, more specifically, ethanol and polyalcohols (propylene glycol, polyethylene glycol, and such), and non-ionic surfactants (Polysorbate 80<sup>TM</sup>, HCO-50, and such).

**[0096]** Sesame oil or soybean oil can be used as an oleaginous liquid, and benzyl benzoate or benzyl alcohol can be used in combination as a solubilizer. Buffers (for example, phosphate buffer and sodium acetate buffer), soothing agents (for example, procaine hydrochloride), stabilizers (for example, benzyl alcohol and phenol), and antioxidants can be used for the formulations. Prepared injection solutions can be filled into appropriate ampules.

**[0097]** The administration is preferably parenteral administration, and specific examples include administration by injection, transnasal administration, transpulmonary administration, and transdermal administration. Examples of administration by injection include systemic and local administration by intravenous injection, intramuscular injection, intraperitoneal injection, subcutaneous injection, and such.

**[0098]** The pharmaceutical compositions contain a pharmaceutically effective amount of the active component (the above-mentioned antibody of the present invention). "Pharmaceutically effective amount (of a compound)" refers to an amount sufficient for treating and/or preventing disorders in which the antigens for the above-mentioned antibodies of the present invention play an important role. For example, "a pharmaceutically acceptable amount" may be an amount required for reducing A $\beta$  accumulation, neutralizing A $\beta$ -induced toxicity, reducing A $\beta$  fibril formation, or such, thereby treating or preventing conditions caused by Alzheimer's disease, when the compound is administered to individuals

(patients). The reduction or neutralization may be, for example, a reduction or neutralization of at least approximately 5%, 10%, 20%, 30%, 40%, 50%, 75%, 80%, 90%, 95%, 99%, or 100%.

**[0099]** Assessment for determining such a pharmaceutically effective amount of the above-mentioned antibodies of the present invention may be carried out using a standard clinical protocol including histopathological diagnosis.

**[0100]** A suitable administration method may be selected depending on the age and symptoms of the patient. The dosage of an antibody-containing pharmaceutical composition may be selected, for example, within the range of 0.0001 mg to 1000 mg per kilogram body weight for each administration. Alternatively, for example, the dosage for each patient may be selected within the range of 0.001 to 100,000 mg/body; however, the dosage is not necessarily limited to these ranges. Although the dosage and administration methods vary depending on the patient's body weight, age, symptoms, and such, one skilled in the art can appropriately select them. In the later-described animal experiments, the dosage was selected based on the high-dose intravenous immunoglobulin therapy (400 mg/kg) covered by health insurance for humans.

**[0101]** Furthermore, the present invention provides methods for detecting A $\beta$  oligomers (examples include A $\beta$ 40 (A $\beta$  1-40) and A $\beta$ 42 (A $\beta$  1-42) oligomers) in samples. Examples of "samples" as described herein include samples collected from subjects. Specifically, the present methods include the step of detecting A $\beta$  oligomers contained in a sample collected from a subject using an antibody of the present invention. A $\beta$  oligomers in a sample can be detected using, for example, sandwich solid-phase enzyme immunoassay methods that use chemiluminescence (chemiluminescence ELISA), immunoprecipitation methods that use the obtained antibodies, immunoblotting, flow cytometry, mass spectrometry, and immunohistochemical analysis.

**[0102]** When A $\beta$  oligomers are detected in a sample collected from a subject by the above-mentioned measurement methods, the subject may be an Alzheimer's disease patient. For example, when the amount of A $\beta$  oligomers in a sample collected from a subject is compared with that from a healthy individual, and if the amount of A $\beta$  oligomers is greater in the subject than in the healthy individual, the subject is determined to be a possible Alzheimer's disease patient. Whether or not a subject is a possible Alzheimer's disease patient is diagnosed usually by physicians (including individuals under instructions from physicians; same herein below). Data on the amount of A $\beta$  oligomers in samples collected from a subject and a healthy individual, which are obtained by the present methods of diagnosis, will be useful for diagnosis by physicians. Therefore, the present methods of diagnosis can be expressed as methods of collecting and presenting data useful for diagnosis by physicians.

**[0103]** Specifically, the present invention provides methods for diagnosing whether or not a subject is a possible Alzheimer's disease patient, wherein the methods comprise detecting A $\beta$  oligomers in a sample collected from the subject using an antibody of the present invention.

**[0104]** Furthermore, the present invention provides methods of diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprise the steps of:

- (a) contacting a sample collected from a subject with an antibody of the present invention and an antibody that binds to an A $\beta$  monomer; and
- (b) measuring the ratio of A $\beta$  oligomer to A $\beta$  monomer in the sample,

wherein the subject is determined to be a possible Alzheimer's disease patient, if the ratio measured in step (b) is higher than that of a healthy individual.

**[0105]** First, in the present methods, a sample collected from a subject is contacted with an antibody of the present invention and an antibody that binds to an A $\beta$  monomer. Herein, "contact" may be carried out, for example, by adding each of the above-mentioned antibodies to a sample collected from a subject, which is placed in a test tube. In this case, the antibody is added suitably in the form of a solution, a solid obtained by freeze-drying, or such. When adding the antibody as an aqueous solution, the solution may purely contain the antibody alone, or may contain, for example, surfactants, excipients, coloring agents, flavors, preservatives, stabilizers, buffers, suspending agents, tonicity agents, binding agents, disintegrants, lubricants, fluidity promoters, or corrigents. The concentration at which the antibody is added is not particularly limited. For example, as with human immunoglobulin formulations, 500-mg, 1000-mg, and 2500-mg freeze-dried formulations and such may be suitably used.

**[0106]** Next, the ratio of A $\beta$  oligomer to A $\beta$  monomer (herein, this is also referred to as "O/M index") in the aforementioned sample is measured. To measure this ratio, the following method is suitably used. For example, as described below in the Examples, the measurement can be carried out using a method of comparing the oligomer and monomer ELISA values obtained from the same sample.

**[0107]** Then, this ratio is compared with the ratio for a healthy individual. When the ratio is higher in the subject than in the healthy individual, the subject is determined to be a possible Alzheimer's disease patient.

**[0108]** The methods of diagnosis of the present invention can be performed both *in vitro* and *in vivo*, but they are preferably performed *in vitro*.

**[0109]** Preferably, the "sample" described herein is not particularly limited as long as it is a tissue derived from a

subject. Examples include the brain (brain parenchyma, and such), organs, and body fluids (blood, cerebrospinal fluid, and such) of a subject. Preferably, the sample is blood (more preferably, plasma) or cerebrospinal fluid.

**[0110]** Furthermore, the present invention provides pharmaceutical agents for use in the above-mentioned methods of measuring A $\beta$  oligomers in a sample, or methods of diagnosing whether or not a subject is a possible Alzheimer's disease patient.

**[0111]** In the present invention, the pharmaceutical compositions comprising an antibody may be included in products and kits containing materials useful for treating pathological conditions of a subject. The products may comprise any labeled container for a compound. Suitable containers include bottles, vials, and test tubes. The containers may be formed from a variety of materials such as glass and plastic. The label on the container surface should indicate that the composition is used to treat or prevent one or more conditions of the disease. The label may also indicate descriptions for administration, and such.

**[0112]** In addition to the above-mentioned container, a kit containing a pharmaceutical composition comprising an antibody may optionally include a second container that stores a pharmaceutically acceptable diluent. The kit may further include other materials desirable from a commercial and user's standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with descriptions for use.

**[0113]** If necessary, the pharmaceutical compositions may be provided in a pack or dispenser device that may contain one or more unit dosage forms comprising an active ingredient. The pack may comprise metal or plastic foil, and, for example, it is a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

**[0114]** In the above-mentioned pharmaceutical agents and kits, besides the antibody of the present invention that is an active ingredient, sterile water, physiological saline, vegetable oils, surfactants, lipids, solubilizing agents, buffers, protein stabilizers (BSA, gelatin, etc.), preservatives, blocking solutions, reaction solutions, reaction quenching solutions, reagents for treating samples, and such, may be mixed as necessary.

**[0115]** The present inventors showed that the antibodies of the present invention are effective for preventing Alzheimer's disease. That is, the present invention provides methods for suppressing the progression of Alzheimer's disease, wherein the methods comprise the step of administering to an individual affected with Alzheimer's disease, a composition comprising the above-mentioned antibody of the present invention and a pharmaceutically acceptable carrier.

[Examples]

#### Methods

##### Preparation of antigens (1A9 and 2C3)

**[0116]** Synthetic A $\beta$  1-42 (Peptide Institute, Inc., Osaka) was dissolved in distilled water or 10 mM phosphate buffer, and incubated at 37°C for 18 hours. Then, the peptides were separated by SDS-PAGE (4-12% NuPAGE Tris-Glycine gel), and after visualization by CBB staining, just the A $\beta$  1-42 tetramer was excised without contamination of the A $\beta$  1-42 monomer.

##### Preparation of antigens (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4)

**[0117]** A fluorescent dye, 6-carboxytetramethylrhodamine (6-TAMRA) (SIGMA) was chemically linked to the N terminus of a synthetic A $\beta$  1-40 peptide (Peptide Institute, Inc.) to produce a modified A $\beta$ . An oligomer-rich sample (A $\beta$  1-40 oligomer) was prepared by copolymerizing the modified A $\beta$  and synthetic A $\beta$  1-40 peptide. It is preferable to adjust the conditions so that the fluorescence intensity determined by ThT assay, which is described below, is one-fourth or less the fluorescence intensity in the absence of modified A $\beta$ . More specifically, it is preferred that 100  $\mu$ M each of the modified A $\beta$  and synthetic A $\beta$  1-40 peptide are mixed, and polymerized for 20 hours.

##### Preparation of antibody-producing hybridomas

**[0118]** Balb/c mice were immunized by injecting the antigen prepared by the method described above into their foot pads. Then, booster immunization was carried out six times. Hybridomas were prepared from inguinal lymph nodes by fusion with Sp2/O-Ag14 cells using Polyethylene Glycol 1500.

##### Antibody isotyping

**[0119]** Isotyping of purified immunoglobulins was carried out using a Serotec (Oxford, UK) mouse monoclonal antibody isotyping kit.

## Dot blot analysis (primary screening)

**[0120]** The initial screening was carried out by dot blot analysis using a nitrocellulose membrane onto which 2.5  $\mu$ l of A $\beta$  1-42 (2.5  $\mu$ g/dot) pre-incubated for 18 hours was immobilized. Non-specific binding sites on the membrane were blocked with a phosphate buffer containing 5% low-fat milk, 1% BSA, and 0.05% Tween-20, and then the membrane was incubated with a culture supernatant. A $\beta$  oligomer-binding antibodies in the culture supernatant were detected by horseradish peroxidase-labeled goat anti-mouse F(ab')<sub>2</sub> (1:3000; Amersham), and visualized using an enhanced chemiluminescence (ECL) kit and LAS3000 mini (Fujitsu, Tokyo, Japan). Among 400 clones, 16 clones positive in the dot blotting, including 1A9 and 2C3, were subjected to the secondary screening described below.

## Immunoprecipitation and immunoblot analysis (secondary screening)

**[0121]** Immunoprecipitation experiments (Ghisso J, et al., Biochem J, 1993) were conducted using an A $\beta$  oligomer-rich amyloid fraction (Matsubara E, et al., Neurobiol Aging, 2004) for the secondary screening to assess whether the 16 clones selected in the primary screening can capture A $\beta$  oligomers in AD brain. A buffer-insoluble, formic acid-soluble fraction prepared from AD brain was incubated with a culture supernatant and Protein G-Sepharose. The immunoprecipitated A $\beta$  oligomers were separated using a NuPAGE 4-12% Bis-Tris-Glycine gel, and transferred onto a nitrocellulose membrane or Immobilon P (Millipore) using 10 mM 3-cyclohexylamino-1-propane sulfonic acid (pH 11) containing 10% methanol at 400 mA for one hour. Non-specific binding sites on the membrane were blocked with a phosphate buffer containing 5% low-fat milk, 1% BSA, and 0.05% Tween-20 at room temperature for three hours. The immunoprecipitated A $\beta$  oligomers were detected by immunoblotting using the 4G8 (1:1000) or 6E10 (1:1000) monoclonal antibody as described above. Two clones, 1A9 and 2C3, were selected from the 16 clones as candidates for therapeutic antibodies for Alzheimer's disease.

## Antibodies

**[0122]** The 6E10 and 4G8 monoclonal antibodies (Covance Immuno-Technologies, Dedham, MA) recognize the epitopes at amino acid positions 1-16 and 17-24 of the human A $\beta$  sequence, respectively. Polyclonal A11 which specifically recognizes A $\beta$  oligomers was purchased from Biosource (Camarillo, CA). Alex Fluor(AF)488- or 594-conjugated goat anti-mouse IgG and Alex Fluor(AF)488-conjugated goat anti-rat IgG were purchased from Molecular Probes (Eugene, OR). Anti-mouse IgG2b was purchased from Sigma (St. Louis, MO). An anti-synaptophysin antibody was purchased from Santa Cruz (Santa Cruz, CA), and an anti-drebrin antibody was purchased from MBL (Nagoya, Japan).

## Size exclusion chromatography (SEC)

**[0123]** SEC was carried out to assess 1A9 and 2C3 for their size specificity. As previously reported (Matsubara E., et al., Neurobiol Aging, 25: 833-841, 2004), this method can selectively separate A $\beta$  monomers and A $\beta$  oligomers, or lipoprotein-bound A $\beta$  and lipoprotein-free A $\beta$ . The present inventors concentrated the culture supernatant from APP/PS1-overexpressing HEK293 cells about ten-fold using a Microcon 3 kDa molecular weight cut-off filter (Millipore Corp.). Then, this concentrate was fractionated into 28 one-milliliter fractions using a Superose 12 size exclusion column (1 cm x 30 cm; Pharmacia Biotech., Uppsala, Sweden; flow rate of 0.5 ml/min) pre-equilibrated with a phosphate buffer. Half of each fraction was subjected to immunoprecipitation using 1A9 or 2C3. A $\beta$  contained in the resulting immunoprecipitates was detected by immunoblotting using 4G8.

**[0124]** Cerebrospinal fluid (CSF) pooled from ten cases of Alzheimer's disease patients or age-matched healthy individuals, and lipoprotein-depleted CSF from the pools were also fractionated under the same conditions as described above. A $\beta$  in the collected fractions was detected by ELISA analysis. To detect the lipid, the total cholesterol was enzymatically quantified using a standard kit (Wako, Osaka, Japan). Under the experimental conditions of the present inventors, the CSF lipoproteins were eluted at fractions 7 to 14, while fractions 15 to 28 contained cholesterol-free proteins.

Preparation of seed-free A $\beta$  solution

**[0125]** Synthetic A $\beta$  1-42 was dissolved at 250  $\mu$ M in 0.02% ammonia water. Then, in order to prepare a seed-free A $\beta$  solution, insoluble peptides, which may function as a seed, were precipitated by ultracentrifugation using an Optima TL ultracentrifuge (Beckman, USA) at 540,000 x g for three hours. The resulting supernatant was collected, aliquoted, and stored at -80°C until use. Samples were prepared by thawing the A $\beta$  stock solutions immediately before use, and diluting them ten-fold with Tris-buffered saline (TBS; 150 mM NaCl and 10 mM Tris-HCl (pH 7.4)). The resulting 25  $\mu$ M solutions were used in the experiments described below. Synthetic A $\beta$  1-40 (HCL form; Peptide Institute, Inc., Osaka) was prepared at 2x concentration.

A $\beta$  incubation and ThT assay (Yamamoto N, *et al.*, J Biol Chem, 282: 2646-2655, 2007)

**[0126]** An A $\beta$  solution (25  $\mu$ M) was incubated in the presence of a predetermined concentration of an antibody at 37°C for two or 24 hours. The ThT fluorescence intensity of the incubation mixture was determined using a fluorescence spectrophotometer (RF-5300PC; Shimadzu Co., Kyoto, Japan). The optimal fluorescence intensity was determined for A $\beta$  amyloid fibrils at excitation and emission wavelengths of 446 and 490 nm, respectively, using 1.0 ml of a reaction mixture containing 5  $\mu$ M ThT and 50 mM glycine-NaOH (pH 8.5). The fluorescence intensity was determined immediately after preparation of the mixture.

**[0127]** Furthermore, the activity of the 4F7, 4H5, 5A5, 5A9, 6E4, and 6H4 antibodies to suppress A $\beta$  amyloid fibril formation was assessed by the following procedure. An A $\beta$  1-42 solution diluted to 12.5  $\mu$ M with cell culture medium was incubated in the presence or absence of each antibody at 37°C for 24 hours. The amount of formed amyloid fibrils was determined by the above-described ThT fluorescence intensity assay method.

A $\beta$ -induced neurotoxicity assay (Yamamoto N, *et al.*, J Biol Chem, 282: 2646-2655, 2007)

**[0128]** Rat pheochromocytoma PC12 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Invitrogen, Carlsbad, CA) containing 10% heat-inactivated horse serum (Invitrogen) and 5% fetal bovine serum (FBS) (Invitrogen). In order to induce the differentiation into nerve cells, PC12 cells were plated at a density of 20,000 cells/cm<sup>2</sup> in culture dishes coated with poly-L-lysine (10 mg/ml), and cultured for six days in DMEM supplemented with 100 ng/ml nerve growth factor (NGF; Alomone Labs, Jerusalem, Israel) (PC12N). PC12N was exposed to 25  $\mu$ M seed-free A $\beta$  1-42 or pre-incubated A $\beta$  1-42 in the presence or absence of antibody at 4°C for 48 hours. The neurotoxicity induced by A $\beta$  1-42 was assessed by Live/Dead dual-color fluorescence assay according to the supplier's instructions (Molecular Probes, Eugene, Oregon).

**[0129]** Furthermore, the activity of the 4F7, 4H5, 5A5, 5A9, 6E4, and 6H4 antibodies to neutralize A $\beta$ -induced neurotoxicity was assessed by the method described below. First, human neuroblastoma cells (SH-SY5Y) were cultured for 24 hours in DMEM containing 10% FBS, at a density of 150,000 cells/well in 24-well plates. Then, the medium was replaced with serum-free culture medium containing A $\beta$  1-42 (12.5  $\mu$ M) in the presence or absence of antibody, and the cells were cultured for another 24 hours. To determine the cytotoxicity induced by A $\beta$  1-42, the level of dead cell-derived LDH released into the medium was measured by a CytoTox96 kit (Promega).

Ultrafiltration and molecular sieve

**[0130]** In order to determine the size-dependent characteristics of neurotoxic A $\beta$  oligomers, the four types of filtrates (<3 kDa, 3 to 10 kDa, 10 to 30 kDa, 30 to 100 kDa) and the retention solution (>100 kDa) were prepared from a 25  $\mu$ M A $\beta$  oligomer solution by sequential ultrafiltration using Microcon 3 kDa, 10 kDa, 30 kDa, and 100 kDa cut-off membranes. Each of the fractions was subjected to the A $\beta$ -induced neurotoxicity assay described above. PC12N was exposed to each fraction to identify the toxic fraction as described above. The distribution of the three-dimensional structures recognized by A11, 1A9, 2C3, and 4G8 was also identified by the dot blot analysis described above. The morphological characterization of the neurotoxic oligomers was performed by examining each fraction using an atomic force microscope.

Electron microscopy (EM) and atomic force microscopy (AFM)

**[0131]** Samples were diluted with distilled water and sprayed over carbon-coated grids to conduct electron microscopy. The grids were negatively-stained with 1% phosphotungstic acid and observed under a Hitachi H-7000 electron microscope (Tokyo, Japan) with an acceleration voltage of 77 kV. AFM assessment was carried out as recently reported. Drops of the samples were placed onto freshly cleaved mica. The mica was allowed to stand for 30 minutes and then washed with water, and the liquid samples were analyzed using Nanoscope IIIa (Digital Instruments, Santa Barbara, CA, USA) set to the tapping mode (Tero, R, *et al.*, Langmuir 20, 7526-7531, 2004). The cantilever used was OMCL-TR400PSA (Olympus, Japan).

Subject tissues and extraction

**[0132]** The present study was conducted based on autopsy cases (n = 50; 26 male and 24 female cases) from the Tokyo Metropolitan Brain Bank for Aging Research of the Tokyo Metropolitan Institute of Gerontology (Itabashi, Tokyo, Japan). This research project was approved by the institutional ethical committees of the Faculty of Medicine, the University of Tokyo; the Tokyo Metropolitan Geriatric Hospital of the Tokyo Metropolitan Institute of Gerontology; and the National Center of Geriatrics and Gerontology. Details of subjects and sample collection have been reported (Katsuno T, Neurology, 64: 687-692, 2005). However, that study analyzed insoluble brain fractions, whereas in this research

project (Katsuno T, Neurology, 64: 687-692, 2005), the present inventors analyzed soluble brain fractions, which remain uncharacterized in previous studies. Frozen tissue samples (the anterior portion of entorhinal cortex) were homogenized in nine volumes of Tris-buffered saline (TS) containing a protease inhibitor cocktail. The homogenates were ultracentrifuged at 265,000 x g for 20 minutes. One-third aliquots (0.5 ml) of the resulting supernatants were subjected to immunoblot analysis.

#### Immunohistochemistry

**[0133]** The left brain hemispheres of Tg2576 mice were sliced into 30- $\mu$ m-thick sagittal sections using a cryotome (RM 2145; Leica, Wetzlar, Germany), and stained with thioflavin S as previously described (Wyss-Coray *et al.*, 2001). The formation of swollen dystrophic neurites was observed using an anti-synaptophysin antibody (Chemicon, Temecula, CA). The number of thioflavin S-positive plaques and synaptophysin-positive swollen dystrophic neurites was counted by observing four or five sections from the left brain hemisphere of each mouse at 40-fold magnification. To observe A $\beta$  deposition, serial sections briefly pre-treated with formic acid or Protease K were stained using an A $\beta$  immunostaining kit (Sigma, St. Louis, MO), and immuno-positive signals were visualized using an ABC elite kit (Vector Laboratories). Images of the cerebral cortex and hippocampus were recorded using a digital camera connected with a microscope, and analyzed using a simple PCI software (Compix Imaging System, Lake Oswego, OR). The brain translocation of antibodies was observed using a confocal laser microscope (Carl Zeiss LSM510). The number of thioflavin S-positive plaques and synaptophysin-positive swollen dystrophic neurites was determined in a double blind manner.

#### Passive immunotherapy and behavioral analysis

**[0134]** Three-month-old female non-transgenic (non-Tg) mice, and Tg2576 mice having and overexpressing the Swedish-type mutant human APP gene with dual mutations (K670N and M671L) derived from familial AD were purchased from Taconics (Germantown, NY, USA). These mice were reared until 13 months old in the animal facility of the present inventors. To determine whether the Alzheimer-like phenotype is prevented by passive immunotherapy, 1A9 or 2C3 (0.4 mg/kg/week), or PBS was administered into the caudal vein of four-month-old Tg2576, and the administration was continued until 13 months. The memory function was assessed at month 13 as previously described (Mouri A, FASEB J, 21: 2135-2148, 2007), based on the following four behavioral paradigms:

- (1) Y-maze test for short-term memory;
- (2) novel object recognition test;
- (3) Morris water maze test; and
- (4) contextual fear conditioning test.

Three days after the behavioral tests, the mice were sacrificed for biochemical and histological assessments. The experimental results were analyzed by one-way ANOVA and two-way ANOVA. Post-hoc analysis was carried out using Fisher test.

#### Separation and removal of lipoprotein

**[0135]** CSF was collected from 12 AD patients and 13 NC individuals. Then, lipoproteins were removed from 600  $\mu$ l each of the CSF by preparative continuous density gradient ultracentrifugation according to a protocol reported previously (Matsubara E, *et al.*, Ann Neurol, 45: 537-541, 1999). The density of CSF was adjusted to 1.25 g/ml with KBr. The CSF was ultracentrifuged at 100,000 rpm and 16°C for eight hours using a Hitachi RP100AT centrifuge. Lipoproteins floating at a density of 1.25 g/ml and lipoprotein-depleted CSF (LPD-CSF) were subjected to ultrafiltration using a 3 kDa cut-off membrane (Microcon 3; Amicon, Inc), and then frozen and stored, or stored at 4°C, until use.

**[0136]** Lipoproteins were also removed by affinity chromatography using PHML-LIPOSORB (Calbiochem, La Jolla, CA). Each sample (plasma or brain) and PHML-LIPOSORB (Calbiochem, La Jolla, CA) were combined at a ratio of 1.5:1, and mixed for 60 seconds. Then, the mixture was centrifuged at 3,000 rpm for ten minutes. The resulting supernatants (lipoprotein-free samples) were subjected to ELISA using 6E10 for the oligomers. The lipoprotein-bound samples were eluted from PHML-LIPOSORB using 20 mM sodium deoxycholate. The removal of specific lipoproteins was confirmed by agarose electrophoresis using 1% gel (Beckmann), followed by staining with FAST-RED 7B (Wako, Osaka, Japan).

#### Quantification of human A $\beta$

**[0137]** Whole plasma and LPDP A $\beta$  species were specifically quantified by sandwich ELISA as previously described



(Matsubara E, et al., Ann Neurol, 537-541, 1999; Matsubara E, et al., Neurobiol Aging, 25: 833-841, 2004). To analyze brain A $\beta$  species, soluble A $\beta$  species in 100  $\mu$ l of buffer were directly subjected to ELISA, while insoluble A $\beta$  samples extracted with 70% formic acid were neutralized with 1 M Tris-HCl (pH 8.0) and diluted 1,000-fold prior to ELISA. The values obtained by the assay were normalized using the brain wet weight, and ultimately presented in pmol/g. Normalization among plates was done by including the three standard plasma samples in all three plates.

#### A $\beta$ oligomer-specific ELISA

**[0138]** Chemiluminescence-based sandwich solid-phase enzyme immunoassay (chemiluminescent ELISA) was used to specifically detect oligomeric A $\beta$  but not monomeric A $\beta$ . Microplates were coated with monoclonal 1A9 (IgG2b isotype) or 2C3 (IgG2b isotype), or a mixture of 1A9 and 2C3. 100  $\mu$ l of a sample (brain or cerebrospinal fluid) was added and incubated continuously for 24 hours at 4°C. Then, horseradish peroxidase-conjugated BA27 Fab' fragment (anti-A $\beta$  1-40 specific to A $\beta$  40; Wako pure chemical, Osaka, Japan) or horseradish peroxidase-conjugated BCO5 Fab' fragment (anti-A $\beta$  35-43 specific to A $\beta$  42; Wako pure chemical, Osaka, Japan) was added and incubated at 4°C for 24 hours. The chemiluminescence generated using SuperSignal ELISA Pico Chemiluminescent Substrate (Pierce, Rockford, IL, USA) was quantified by a Veritas Microplate Luminometer (Promega).

**[0139]** To assess the *in vivo* efficacy of the peripheral administration of monoclonal 1A9 and 2C3, plasma and organ samples were collected from administered mice and analyzed for A $\beta$  oligomers by ELISA using HRP-labeled 6E10 (Senetek PLC, Napa, CA, USA) specific to the human oligomers. High-sensitivity detection was achieved by using the above-described chemiluminescent system. To avoid interference by lipoprotein-bound A $\beta$  monomers, the present inventors pre-treated plasma and organ samples using PHML-LIPOSORB in the same way as described above. The resulting lipoprotein-depleted samples were used for the assay.

#### Inhibition ELISA

**[0140]** A $\beta$  oligomers used in this assay were prepared by diluting synthetic A $\beta$  1-40 (HCl form) to a concentration of 0.1 mg/ml with PBS and incubating this at 37°C for one hour. Meanwhile, A $\beta$  monomers were prepared by diluting synthetic A $\beta$  1-40 (TFA form) to a concentration of 0.1 mg/ml with PBS. A $\beta$  oligomers were immobilized onto 96-well immunoplates at 400 ng/well, and then the plates were blocked with BSA. Next, the A $\beta$  monomers or oligomers stepwise-diluted in the range of 100 pg/ml to 100  $\mu$ g/ml were reacted with the 4F7, 4H5, 5A5, 5A9, 6E4, or 6H4 antibodies, or the control anti-A $\beta$  antibodies (4G8 and 6E10). After incubation for two hours, the mixtures were added to the above-described 96-well immunoplates, and incubated at room temperature for ten minutes. The binding of immobilized A $\beta$  oligomers to each of the antibodies was detected by measuring the absorbance at 450 nm in the color development reaction using an HRP-labeled anti-mouse IgG antibody and a TMB solution.

[Example 1]

#### Preparation of A $\beta$ oligomer-specific monoclonal antibodies (1A9 and 2C3)

**[0141]** A $\beta$  oligomers and monomers co-exist in a solution. Thus, it is essential to remove A $\beta$  monomers for preparation of antigens to produce A $\beta$  oligomer-specific antibodies. As shown in Fig. 1A, the present inventors succeeded in isolating SDS-stable A $\beta$  tetramers without contamination of A $\beta$  monomers by SDS-PAGE. After *in vivo* immunization with the isolated A $\beta$  tetramers, positive hybridoma clones were selected by two-step screening using dot blot analysis followed by immunoprecipitation. Among 400 clones subjected to dot blot analysis, 16 clones were determined to be positive (positivity rate = 4%). To assess the specificity of the isolated positive clones to the oligomers, a phosphate buffer-insoluble and formic acid (FA)-soluble amyloid fraction derived from AD brain (Matsubara E et al., Neurobiol Aging, 25: 833-841, 2004) was analyzed by immunoprecipitation using the cell culture supernatants of the positive hybridomas (Fig. 1B). The A $\beta$  dimer, a smaller amount of the trimer, and a high-molecular-weight smear characteristic to aggregated A $\beta$  molecular species were detected by immunoblot analysis using anti-A $\beta$  monoclonal 4G8. A very small amount of A $\beta$  monomers dissociated in the presence of SDS was also detected. To further confirm the existence of three-dimensional structures recognized by native 1A9 and 2C3 (*i.e.*, oligomers), the present inventors detected the oligomers in conditioned medium (CM) of human embryonic kidney (HEK) 293 cells transfected with mutant PS1 cDNA (Nakaya Y et al., J Biol Chem, 280: 19070-19077, 2005). The present inventors fractionated HEK293 CM by SEC, and then identified the oligomers. As reported previously (Matsubara E et al., Neurobiol Aging, 25: 833-841, 2004; Yamamoto N, et al., J Biol Chem, 282: 2646-2655, 2007), this method can effectively separate the oligomers (fractions 8 to 13) from monomers (fractions 14 to 20). When immunoprecipitated with monoclonal 1A9, SDS-stable A $\beta$  dimers secreted into CM were precipitated in fraction 8 (>680 kDa); SDS-stable A $\beta$  dimers and trimers were precipitated in fraction 13 (17 to 44 kDa); and a very small amount of the dimers was precipitated in fraction 16 (Fig. 1C). Similar results were obtained when

immunoprecipitation was carried out using 2C3 (data not shown). These data demonstrate that monoclonal 1A9 and 2C3 are exactly specific to A $\beta$  oligomers but do not recognize A $\beta$  monomers.

[Example 2]

#### The anti-neurotoxic activity of monoclonal 1A9 and 2C3

**[0142]** To assess whether monoclonal 1A9 and 2C3 can prevent A $\beta$ -induced neurotoxicity, NGF-differentiated PC12 cells (PC12N) were incubated with 25  $\mu$ M seed-free A $\beta$  1-42 (ThT-negative 540,000 x g supernatant) in the presence or absence of the monoclonal antibodies (mAbs) at 37°C for 48 hours. The viability of nerve cells was determined by LIVE/DEAD assay (Fig. 2). Nerve cell death was detected at a significantly high level (50%) in the presence of A $\beta$  1-42 (Figs. 2B and 2G), as compared to the control assay (Fig. 2A). Non-specific IgG2b (Figs. 2C and 2G) could not inhibit the A $\beta$  1-42-induced neurotoxicity under the same conditions. The commercially available A $\beta$ -specific monoclonal antibody 4G8 (IgG2b isotype; Figs. 2D and 2G) had a tendency to enhance the toxicity. Monoclonal 2C3 (IgG2b isotype; Figs. 2F and 2G) neutralized the neurotoxicity of A $\beta$  1-42 almost completely in a concentration-dependent manner. Thus, the *de novo*-formed neurotoxic three-dimensional structure recognized by 2C3 was speculated to take an oligomer form. Meanwhile, the anti-neurotoxic activity of 1A9 (IgG2b isotype; Figs. 2E and 2G) falls between the anti-neurotoxic activity of 2C3 and non-specific IgG2b. This suggests that the three-dimensional structure recognized by 1A9 is structurally different from the 2C3-recognized oligomers.

[Example 3]

**[0143]** Currently, the determination of the precise size and conformation of neurotoxic A $\beta$  1-42 oligomers is one of the most urgent issues and which is subjected to intense competition. The present inventors succeeded in isolating soluble neurotoxic A $\beta$  1-42 molecular species and fractionating the species into the following five fractions by ultrafiltration and molecular sieving (UC/MS) (Fig. 3A):

- fraction 1, filtrate of <3 kDa (lane 1);
- fraction 2, filtrate of 3 to 10 kDa (lane 2);
- fraction 3, filtrate of 10 to 30 kDa (lane 3);
- fraction 4, filtrate of 30 to 100 kDa (lane 4); and
- fraction 5, retention solution of >100 kDa (lane 5).

**[0144]** The immunoblot analysis using monoclonal 4G8 (Fig. 3A) revealed that:

- fraction 1 does not contain A $\beta$  (lane 1);
- fraction 2 contains A $\beta$  monomers (lane 2);
- lane 3 contains A $\beta$  monomers and a small amount of A $\beta$  dimers;
- fraction 4 contains A $\beta$  monomers to pentamers (lane 4); and
- fraction 5 contains A $\beta$  monomers to pentamers, and molecules of 45 to 160 kDa (lane 5).

**[0145]** These data suggest that 2% SDS depolymerizes high-molecular-weight (HMW) A $\beta$  oligomers into A $\beta$  monomers and low-molecular-weight (LMW) A $\beta$  oligomers. To assess the size distribution of toxic A $\beta$  1-42, the present inventors measured the biological activity of each fraction incubated with PC12N at 37°C for 48 hours. As shown in Figs. 3B and 3C, it was demonstrated that: fraction 1 was non-toxic, and fraction 2 had a very weak toxicity, suggesting that A $\beta$  monomers and dimers are unlikely to be toxic. Fractions 3 to 5 were significantly toxic (one-way ANOVA;  $p < 0.0001$ ), suggesting that the size of neurotoxic oligomers theoretically corresponds to the size of trimers or higher-molecular-weight polymers. The dot blot analysis using the oligomer-specific A11 antibody demonstrated that the above-mentioned three neurotoxic fractions (3 to 5) were positive for A11, supporting the evidence that the neurotoxic molecules are oligomeric (Fig. 3D). The 2C3-recognized oligomers were detected in fractions 4 and 5 (Fig. 3D). Thus, 2C3 was demonstrated to actually react with neurotoxic A $\beta$  oligomers (>30 kDa). Furthermore, the majority of the 2C3-recognized oligomers was detected in fraction 5 (>100 kDa) that was the most toxic, and thus the 2C3-recognized oligomers having a molecular weight over 100 kDa were considered to show a strong neurotoxicity (Fig. 3D). Meanwhile, only an extremely small amount of the 1A9-recognized oligomers was distributed in fraction 5 that was the most toxic. This is consistent with the result that the neutralization of neurotoxicity by 1A9 was insufficient (Figs. 2E and 2G). By contrast, monoclonal 4G8 having no anti-neurotoxic activity detected the A $\beta$  species distributed in all of the fractions (Fig. 3D). This suggests the possibility that non-toxic and toxic oligomers of the same size co-exist.

**[0146]** To further assess the toxicity-structure correlation, each fraction was subjected to atomic force microscopy

(AFM). The presence of globular particle morphology consistent with the fraction size was detected in the three neurotoxic fractions. Fig. 3E shows the atomic force microscopic images of non-toxic fraction 2 (Fr. 2), toxic fractions 3 (Fr. 3) and 4 (Fr. 4), and the most toxic fraction 5 (Fr. 5). The formation of many granular polymer molecules was clearly observed in the toxic fractions. In particular, fraction 5 was revealed to contain heterogeneous toxic molecules including bead-shaped and ring-shaped molecules in addition to various large and small granular molecules.

[Example 4]

#### The activity of 1A9 and 2C3 to suppress A $\beta$ amyloid fibril formation

**[0147]** Next, the present inventors assessed the activity of 1A9 and 2C3 to suppress A $\beta$  amyloid fibril formation. The formation of A $\beta$  1-42 amyloid fibrils (at 0, 10, 25, and 50  $\mu$ M) was assessed by measuring the ThT fluorescence for 72 hours at 37°C. Under the conditions used by the present inventors, seed-free A $\beta$  1-42 (ThT-negative supernatant fraction obtained by ultracentrifugation at 540,000 x g) was polymerized into amyloid fibrils by nucleation-dependent polymerization (Fig. 4A). To assess the activity of 1A9 and 2C3 to suppress A $\beta$  amyloid fibril formation, the present inventors incubated 25  $\mu$ M seed-free A $\beta$  1-42 in the presence or absence of the antibodies at 37°C for 48 hours. As shown in Fig. 4B, the ThT fluorescence intensity was altered in a 2C3 concentration-dependent manner, while none of monoclonal 1A9 and 4G8 and non-specific IgG2b altered the fluorescence intensity. Meanwhile, when A $\beta$  was polymerized by incubation for two hours, 1A9, as well as 2C3, showed the activity to almost completely suppress the fibril formation (Fig. 4C). Since the activity to suppress A $\beta$  amyloid fibril formation was detected even when the molar ratio of 2C3 to A $\beta$  was low, 2C3 was inferred to have the activity to inhibit the polymerization nucleus formed *de novo* or the seed function at an early stage of A $\beta$  1-42 amyloid fibril formation. Similar results were obtained by morphological observation. As shown in Fig. 4E (A $\beta$  42 alone) and Fig. 4F (A $\beta$  42 + 2C3, 25:3), electron microscopy (EM) demonstrated that the formation of A $\beta$  amyloid fibrils was partially inhibited in the presence of monoclonal 2C3, while only a weak inhibitory effect was produced in the presence of 1A9 (Fig. 4G). Meanwhile, none of the test antibodies exhibited the effect of lysing or depolymerizing A $\beta$  1-42 amyloid fibrils that were formed by incubation for 24 hours (Fig. 4D).

[Example 5]

#### Toxicity-related oligomers targeted by 1A9 and 2C3

**[0148]** To elucidate the structural and kinetic connection between the A $\beta$  1-42 oligomerization and amyloid fibril formation, the polymerization time course was analyzed by dot blotting using A11, 1A9, 2C3, and 4G8. As shown in Fig. 5A, the majority of A11 antibody-reactive oligomers was formed during the lag time phase of polymerization (0 to 8 hours), and the ThT fluorescence intensity was relatively weak. During the next fibril extension phase (8 to 24 hours), the level of A11-immunoreactive oligomers reached a plateau, and then was constant (about 20% of the peak level) until 72 hours (plateau phase). It has been demonstrated that, since the anti-oligomer A11 antibody does not recognize amyloid fibrils, the A $\beta$  oligomer formation can be specifically observed using the antibody (Kayed R, et al., Science 300, 486-489, 2003). Hence, the present results suggest that the A $\beta$  oligomer formation precedes amyloid fibril formation, and there is an oligomerization state that does not directly enter the amyloid fibril formation pathway. The 2C3-recognized oligomers were kinetically similar to the A11-recognized oligomers, but not the 1A9-recognized oligomers. The 1A9-recognized oligomers were detected only after four hours, and then the immunoreactivity to 1A9 increased twofold over time. This suggests that the 1A9-recognized oligomers are slowly formed. Meanwhile, it was revealed that the 2C3-recognized oligomers are transiently increased during the lag time phase (0 to 8 hours), and then exist at a very low level (less than 5%) in a oligomerized state from 8 to 72 hours. The above-described data obtained by the present inventors suggest the possibility that the A11-, 1A9-, and 2C3-recognized oligomers have structurally and immunologically different conformations or stability, and the 2C3-recognized oligomers are relatively unstable as compared to the 1A9-recognized oligomers.

**[0149]** To characterize the *de novo* toxic polymerization state, PC12N were exposed at 37°C for 48 hours to seed-free A $\beta$  1-42 (0 hour), or A $\beta$  1-42 pre-incubated for two, four, or 24 hours (Fig. 5B), and the neurotoxic activity was assayed. As shown in Fig. 5B, the immunoblotting analysis using 4G8 revealed that the monomers, dimers, and trimers exist even at the 0 hour time point. The pre-incubation of two or four hours resulted in a high-molecular-weight (HMW) smear pattern of 45 to 160 kDa, in addition to the monomers to pentamers. At the time point of 24 hours, the HMW smear was dramatically reduced, and there were two types of major components: a high-molecular-weight species that could not enter the gel and thus remained in the well, and a small amount of the monomers. The HMW smear disappeared after further incubation at 37°C for 48 hours. As shown in Figs. 3A and 5C, the molecular sieve experiment revealed that seed-free A $\beta$  1-42 is converted into molecular species of 100 kDa or more, and exhibits the strongest toxicity. By SDS-PAGE, it was demonstrated that the toxic molecules include molecular species showing a high-molecular-weight

(HMW) smear pattern of 45 to 160 kDa, in addition to the monomer to pentamer species, and the toxic polymers can be easily depolymerized into low-molecular-weight species in the presence of SDS. However, when seed-free A $\beta$  1-42 was pre-incubated for two, four, and 24 hours, the neurotoxic activity of *de novo*-formed A $\beta$  oligomers was reduced by about 12.5% and 26%, respectively (Fig. 5C). This result suggests that the level of *de novo*-formed A $\beta$  oligomers in the early period of A $\beta$  polymerization is a determining factor for neurotoxicity, and that the formation reaches a peak in the period of zero to two hours, and then the level of formed A $\beta$  oligomers reduces over time. Alternatively, there is a possibility that nuclei for the *de novo* polymerization of A $\beta$  amyloid fibrils, or amyloid fibrils themselves have the neurotoxicity-neutralizing activity. The present inventors incubated A $\beta$  1-42 for two hours, and then removed insoluble A $\beta$  polymerization nuclei and amyloid fibrils by ultracentrifugation for three hours at 540,000 x g. The supernatant and pellet obtained by ultracentrifugation at 540,000 x g exhibited similar levels of thioflavin T signals, suggesting that the 540,000 x g supernatant contains soluble ThT-positive A $\beta$  polymers (The ThT binding indicates structural changes to form a  $\beta$  sheet-rich structure, but not fibril formation). The neurotoxicity was restored and enhanced when PC12N was exposed to the soluble polymers (Fig. 5D). This suggests that insoluble A $\beta$  1-42 itself has the anti-toxic activity. Under the conditions described above, monoclonal 1A9 completely neutralized the neurotoxicity induced by soluble A $\beta$  oligomers enriched in  $\beta$  sheet structures, and this neutralizing activity was greater than that of 2C3. Meanwhile, non-specific IgG2b has no effect on the viability of cultured PC12N. Accordingly, it is speculated that neurotoxic 1A9-recognized polymers are basically soluble toxic oligomers that have been slightly stabilized due to some structural change, while neurotoxic 2C3-recognized polymers are basically short-lived oligomeric intermediates that are very unstable due to drastic structural changes during the early stage of polymerization process.

[Example 6]

#### Monoclonal 1A9 and 2C3 recognize A $\beta$ oligomers in the brain parenchyma

**[0150]** The present inventors demonstrated the specificity and biological activity of 1A9 and 2C3. Furthermore, the inventors detected 1A9 and 2C3 polymers in the brain by immunohistochemistry. The present inventors performed conventional immunohistochemistry methods to enhance immune reaction by formaldehyde fixation, and formic acid, SDS, or microwave treatment of brain sections. The two antibodies exhibited no immunoreactivity to AD brain by any one of the enhancement methods. Thus, the present inventors pre-treated the sections with Protease K, which is known to improve immunostaining (Wrzolek MA, et al., Am J Pathol, 141: 343-355, 1992). As a result, many senile plaques were stained with 1A9 (Fig. 6A), 2C3 (Fig. 6B), and A11 (Fig. 6C). Together with the finding from the *in vitro* experiments by the present inventors that A $\beta$  amyloid fibrils neutralize the A $\beta$  oligomer-induced neurotoxicity, the result described above suggests that a senile plaque serves as a defensive reservoir to isolate and store A $\beta$  oligomers, and thus the interior of the reservoir is hardly accessible for antibodies. Indeed, immunoprecipitation using 1A9 and 2C3 demonstrated that amyloid fractions composed of senile plaques contain A $\beta$  oligomers recognized by the two antibodies. Thus, the hypothesis of the present inventors was proven to be consistent with the *in vivo* finding (see Fig. 1B).

**[0151]** To further assess the existence of "soluble" 1A9- and 2C3-recognized polymers in the brain, the present inventors carried out immunoprecipitation experiments using the two antibodies. Brain homogenates were prepared using Tris-buffered saline (TBS) to avoid chemical modification during the extraction of soluble oligomers. The oligomers having a molecular weight of 4mer, 5mer, 8mer, and 12mer were immunoprecipitated with 1A9 from TBS samples of the cerebral cortex from AD brain (Fig. 6D, lane 2), while the level of the oligomers in the control healthy brain was below the detection limit (lane 3). While the intensity of 4mer, 5mer, and 8mer was comparable between 1A9 (lane 2) and the monoclonal 4G8/6E10 mixture (lane 1), 1A9 appeared to recover a larger amount of 12mer than 4G8/6E10. The immunoprecipitation with 2C3 showed a comparable result (Fig. 6, lanes 4 to 6). Next, the present inventors identified the molecules responsible for the *in vivo* neurotoxicity in the human entorhinal cortex. It is well known that neurofibrillary tangle (NFT) and nerve cell loss precede the formation of senile plaques in lesions in general elderly populations. The present inventors hypothesized that the lack of functional reservoirs such as senile plaques for 1A9- and 2C3-recognized polymers is harmful for entorhinal cortex neurons, and is a possible cause of memory disturbance. The level of 12mer in the buffer-soluble fractions of previously reported 50 autopsy cases was determined by immunoblotting using monoclonal 1A9 and 2C3. The 50 cases include two AD cases, 35 cases at Braak NFT stages I to II, and 13 cases at NFT stages III to IV (Katsuno et al., Neurology, 64: 687-692, 2005). As shown in Figs. 6E (1A9) and 6F (2C3), the immunological activity of 1A9- or 2C3-immunoreactive 12mer relative to actin was significantly higher in AD patients as compared to the healthy control group (Braak NFT stages I to II) and mild cognitive impairment group (Braak stages III to IV). Interestingly, the 12mer was accumulated in the entorhinal cortex of the healthy control group (Braak NFT stages I to II) and mild cognitive impairment group (Braak stages III to IV) at a level of about 40% and 60% (the level of AD cases is 100%), respectively (Figs. 6E and 6F). This result indicates that the accumulation of 12mer precedes the onset of cognitive impairment, and is increased as the Braak NFT stage advances, suggesting that the 1A9- and 2C3-immunoreactive 12mer are polymers responsible for the *in vivo* neurotoxicity.

[Example 7]

Monoclonal 1A9 and 2C3 recognize A $\beta$  oligomers in the cerebrospinal fluid

**[0152]** The A $\beta$  polymers (soluble 1A9- and 2C3-immunoreactive 12mer) responsible for the *in vivo* neurotoxicity were found in the brain parenchyma. Thus, the present inventors speculated that CSF also contains the polymers. To verify this, the present inventors fractionated pooled CSFs from ten AD patients and ten age-matched healthy individuals as a control by SEC, and assayed the fractions by A $\beta$  oligomer-specific sandwich ELISA using monoclonal BC05 or BA27 in the capturing and detection systems. The BC05/BC05 oligomer ELISA detected soluble A $\beta$  1-42 in fraction 13, while the BA27/BA27 ELISA detected soluble A $\beta$  1-40 in fractions 7 to 14 (data not shown). However, in each ELISA, the absorbance (O.D. at 450 nm) was low for sensitive detection of a small amount of A $\beta$  oligomers in CSF. The detection of A $\beta$  monomers in the same fractions by BNT77 ELISA showed that lipoprotein-bound A $\beta$  monomers (fractions 7 to 14) and lipoprotein-free A $\beta$  monomers (fractions 15 to 17) coexist with A $\beta$  oligomers in the fractions (Figs. 7-1A and 7-1B) (Matsubara E, et al., *Neurobiol Aging*, 25: 833-841, 2004). The level of lipoprotein-bound A $\beta$  monomer in AD was comparable to that of the healthy control, while the level of lipoprotein-free A $\beta$ 40 monomer (Fig. 7-1A) and A $\beta$ 42 monomer (Fig. 7-1B) in AD was lower as compared to the age-matched healthy control. The present inventors also found that lipoprotein-bound A $\beta$  monomers, in addition to the oligomers, can be detected when ELISA is designed to use HRP-labeled BC05 or BA27 as a capture antibody. This problem remained unnoticed in the prior art document (Lee EB, et al., *J Biol Chem*, 281: 4292-4299, 2006), which describes assay methods (for example, 6E10/6E10 ELISA) that are similar to the methods described herein. Since the oligomers and lipoprotein-bound A $\beta$  monomers are eluted at a comparable retention time in SEC, it is impossible to distinguish them by oligomer ELISA using the same antibody in capturing and detection. Thus, it was revealed that CSF containing lipoproteins is unsuitable for a test sample when A $\beta$  oligomers are analyzed using A $\beta$  oligomer-nonspecific antibodies.

**[0153]** To overcome the weaknesses of the prior art methods, the present inventors improved the detection antibodies and samples used in ELISA. Lipoproteins were pre-depleted from CSF, and the resulting lipoprotein-depleted CSF (LPD-CSF) was used as an assay sample. A $\beta$  oligomer-specific 1A9 and 2C3 were used as detection antibodies for ELISA. Furthermore, chemiluminescence ELISA was developed to enhance the sensitivity. Pooled LPD-CSF (Figs. 7-1C to D) was fractionated by SEC, and each fraction was analyzed for A $\beta$  oligomer distribution by luminescence ELISA using 1A9 or 2C3 as a detection antibody. As shown in Figs. 7-1C to D, A $\beta$  oligomers were detected in SEC fractions 12 to 15 (relatively large A $\beta$  with a molecular weight ranging within 18 to 108 kDa, which corresponds to the size of 4mer to 24mer). The level of 1A9- and 2C3-recognized oligomers was elevated in all of the AD patient-derived fractions in which the oligomers were detectable. To assess the usefulness of the A $\beta$  oligomers as therapeutic markers, the level of A $\beta$  oligomers in LPD-CSF from AD patients was compared to that from the age-matched healthy control, although a limited number of cases were analyzed. As shown in Fig. 7-2G, 2C3-recognized oligomers composed of A $\beta$  x-42 were significantly increased in the AD patient group as compared to the normal control group (nonparametric analysis;  $p = 0.0103$ ). By contrast, for 2C3-recognized oligomers composed of A $\beta$  x-40, there was no significant difference between the two groups. Meanwhile, the level of 1A9-recognized oligomers composed of A $\beta$  x-42 was higher in AD than in the control, although the difference was not statistically significant. For 1A9-recognized oligomers composed of A $\beta$  x-40, there was no significant difference between the two groups (Fig. 7-2E). The structural change from A $\beta$  monomer to oligomer occurs in the earliest period of the process of A $\beta$  polymerization. The ratio between A $\beta$  oligomer and monomer (O/M index) can be used as a clinical indicator reflecting the pathological conditions of AD. As shown in Figs. 7-2F and 7-2H, the O/M indices for A $\beta$ 42 and A $\beta$ 40 were significantly increased in the AD patient group as compared to the healthy control group (1A9,  $P=0.0137$  for A $\beta$ 42 and  $P=0.0429$  for A $\beta$ 40; 2C3,  $P=0.0012$  for A $\beta$ 42 and  $P=0.0051$  for A $\beta$ 40). The results described above show that the 1A9- and 2C3-positive three-dimensional structures are present as A $\beta$  oligomers in LPD-CSF, and increased in AD patients. In addition, the results obtained by the present inventors demonstrated that the structural conversion of lipoprotein-free soluble A $\beta$  to the oligomeric intermediates occurs in CSF of AD patients, and the oligomers can be detected as useful biological markers for diagnosis of sporadic AD.

[Example 8]

Passive immunotherapy using monoclonal 1A9 and 2C3 prevents the onset of memory disturbance in Tg2576

**[0154]** To assess the *in vivo* preventive/therapeutic effect of passive immunotherapy based on the administration of 1A9 ( $n=13$ ) or 2C3 ( $n=11$ ), the present inventors administered 1A9 or 2C3 (0.4 mg/kg/week), or PBS to Tg2576 mice via the caudal vein during the 4 to 13 month period. The memory function was assessed at 13 months old in terms of the following four types of learning/behavioral paradigms:

- (1) short-term memory in Y-maze test (Fig. 8A);

- (2) object recognition memory in novel object recognition test (Fig. 8B);
- (3) spacial memory in water maze test (Fig. 8C); and
- (4) associative emotional memory in contextual fear conditioning test (Fig. 8D).

**[0155]** As compared to 1A9- and 2C3-administered Tg2576 mice, PBS-administered Tg2576 mice showed significant learning and behavioral impairments (Fig. 8A to 8D). Unlike the memory function of PBS-administered Tg2576 mice (n=10), the memory function of 1A9 and 2C3-administered Tg2576 mice was indistinguishable from that of age-matched non-administered wild type cohort mice, which was previously determined. Therefore, 1A9 and 2C3-administered Tg2576 mice were shown to retain both short- and long-term memory, which were impaired in the PBS administration group. That is, the present inventors obtained evidence supporting the view that the onset of memory disturbance, in particular AD, can be prevented by conducting passive immunotherapy targeting A $\beta$  oligomers before the onset. Furthermore, the result described above presents the first *in vivo* evidence that directly indicates that A $\beta$  oligomers are responsible for the onset of memory disturbance.

[Example 9]

#### Monoclonal 1A9 prevents A $\beta$ accumulation in the brain of Tg2576

**[0156]** Tg2576 mice administered with PBS (n=10) and Tg2576 mice treated with passive immunotherapy during the 4 to 13 month period (1A9 administration group, n=13; 2C3 administration group, n=11) were dissected after the learning/behavioral experiments. The amount of A $\beta$  accumulated in the brain (cerebral cortex vs. hippocampus) was determined in the following three fractions (150 mg/extract) prepared by serial extraction: soluble fraction in Tris buffer containing protease inhibitors; 2% SDS-soluble amyloid fraction; and 2% SDS-insoluble and 70% formic acid-soluble amyloid fraction. It is considered that non-accumulative, physiological A $\beta$  molecules are contained in the Tris buffer fraction, while 2% SDS-soluble A $\beta$  includes A $\beta$  in diffuse senile plaques before amyloid fibril formation, immunocytochemically undetectable A $\beta$ , and conformationally altered, accumulative soluble oligomeric A $\beta$ . A $\beta$  was selectively quantified by A $\beta$ 40 and A $\beta$ 42 end-specific ELISA (BNT77/BA27 specific for A $\beta$ 40, BNT77/BC05 specific for A $\beta$ 42, WAKO kit). There was no marked difference among the three groups in the A $\beta$  concentration in the Tris buffer fraction where the major components were non-accumulative, physiological A $\beta$  molecules (Figs. 9A and 9C, A $\beta$  x-40; Figs. 9B and 9D, A $\beta$  x-42). Regarding soluble A $\beta$  accumulated in the brain (SDS fraction), a significant suppressive effect on the accumulation of A $\beta$  x-40 and A $\beta$  x-42 in the cerebral cortex was observed only in the 1A9 administration group (Fig. 9E, A $\beta$  x-40; Fig. 9F, A $\beta$  x-42). No accumulation-suppressive effect was observed in the hippocampus (Fig. 9G, A $\beta$  x-40; Fig. 9H, A $\beta$  x-42). Meanwhile, regarding insoluble A $\beta$  accumulated in the brain (FA fraction), a significant suppressive effect on the accumulation of A $\beta$  x-40 in the cerebral cortex was observed only in the 1A9 administration group (Fig. 9I, A $\beta$  x-40; Fig. 9J, A $\beta$  x-42). No accumulation-suppressive effect was observed in the hippocampus (Fig. 9K, A $\beta$  x-40; Fig. 9L, A $\beta$  x-42). The A11 immunoblot analysis of the SDS-soluble fractions showed a suppressive effect on the accumulation of A11-positive oligomer (4mer) in the cerebral cortex in the two antibody treatment groups (Fig. 9M).

[Example 10]

#### Plasma A $\beta$ oligomers are increased by passive immunotherapy with 1A9 and 2C3

**[0157]** There was no significant difference in the plasma A $\beta$  concentration among the following three groups: Tg2576 mice administered with PBS (n=10), and Tg2576 mice treated with passive immunotherapy during the 4 to 13 month period (1A9 administration group, n=13; 2C3 administration group, n=11) (Fig. 10A, A $\beta$  x-40; Fig. 10B, A $\beta$  x-42). There was also no significant difference in the A $\beta$ 40/42 ratio (Fig. 10C).

**[0158]** In order to elucidate the mechanism underlying the preventive effect of passive immunotherapy with 1A9 and 2C3 (IVIg) against the AD-like phenotype in Tg2576 mice, the present inventors assessed the level of physiological saline-soluble and -insoluble A $\beta$  oligomers in pooled brain homogenates, and the level of A $\beta$  oligomers in the peripheral blood and plasma. There was no difference in the amount of physiological saline-soluble A $\beta$  oligomers in the pooled brain homogenates among the treatment groups (Fig. 10D). Meanwhile, the amount of insoluble A $\beta$  oligomers was shown to be reduced in the 1A9 and 2C3 treatment groups (Fig. 10E). Furthermore, pooled plasma from each group (albumin-depleted plasma, upper part of Panel F; albumin/lipoprotein-depleted plasma, lower part of Panel F) was assayed for A $\beta$  oligomers by A11 dot blotting. The result shows that the oligomers were present in the plasma from PBS-administered Tg2576 mice (Fig. 10F). A11-positive oligomers in plasma were clearly increased in the passive immunotherapy groups as compared to the PBS administration group (Fig. 10F). The proportion of 2C3-recognized oligomers in a lipoprotein-bound form was greater than that of 1A9-recognized oligomers (lower part of Panel F). Furthermore, plasma A $\beta$  oligomers were detected by A11 immunoprecipitation. The result shows that the oligomers of about

200 kDa were increased in Tg2576 mice treated with passive immunotherapy as compared to the PBS administration group (Fig. 10G). The increase in plasma A $\beta$  oligomers in the passive immunotherapy groups can be considered to directly reflect enhanced cerebral clearance. Thus, the present inventors obtained evidence that direct target molecules for intravenous passive immunotherapy are also present in blood in addition to brain, and that oligomer-selective cerebral clearance can be enhanced through peripheral sites of action. That is, the present inventors showed the clinical usefulness of the intravenous passive immunotherapy.

[Example 11]

Formation of senile amyloid plaques and swollen dystrophic neurites can be suppressed by passive immunotherapy using 1A9 and 2C3

**[0159]** Immunohistochemical A $\beta$  deposition was suppressed in the passive immunotherapy groups (Fig. 11A). The formation of thioflavin S-positive senile amyloid plaques was significantly suppressed in both the cerebral cortex and hippocampus (Fig. 11B, upper part), and the reduction was also clearly demonstrated by histochemistry (Fig. 11B, lower part). The formation of synaptophysin-positive swollen dystrophic neurites was also significantly suppressed in the passive immunotherapy groups (Fig. 11C).

[Example 12]

Immunostaining analysis using anti-synaptophysin and anti-drebrin antibodies

**[0160]** 1A9 and 2C3 suppressed the presynaptic and postsynaptic degeneration in the cerebral neocortex (Fig. 12).

[Example 13]

The antibodies translocate to the brain

**[0161]** The existence/localization of deposited A $\beta$  and cerebral mouse IgG was assessed using a confocal laser microscope. The result shows that mouse IgG is localized almost independently of deposited A $\beta$  within the areas containing diffuse senile plaques. Mouse IgG was observed only in the passive immunotherapy groups (1A9, Fig. 13A; 2C3, Fig. 13B), but not in the PBS administration group (Fig. 13C). Thus, a fraction of the antibodies administered into the blood was considered to translocate to the brain. This result shows that the preventive effect on memory disturbance was produced not only through the direct neutralization of the toxicity of soluble A $\beta$  polymers by the antibodies translocated to the brain, but also through the clearance of soluble A $\beta$  polymers in the form of a complex with the antibodies into the blood. Thus, the therapeutic effect was considered to be based on multiple action mechanisms.

[Example 14]

Preparation of A $\beta$  oligomer-specific monoclonal antibodies (5A5, 5A9, 4F7, 4H5, 6E4, and 6H4) and dot blot analysis

**[0162]** 33 clones prepared by the above-described method that uses the A $\beta$  1-40 oligomer as an antigen were assessed by dot blot analysis. The result showed that the six types of monoclonal antibodies specifically recognize A $\beta$  oligomers. As shown below, the isotype of the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) was determined:

4F7:  $\kappa$  for the L chain, and IgG2a for the H chain;  
 4H5:  $\kappa$  for the L chain, and IgG2a for the H chain;  
 5A5:  $\kappa$  for the L chain, and IgG2b for the H chain;  
 5A9:  $\kappa$  for the L chain, and IgG2b for the H chain;  
 6E4:  $\kappa$  for the L chain, and IgG1 for the H chain; and  
 6H4:  $\kappa$  for the L chain, and IgG2b for the H chain.

Furthermore, the immuno-dot blot analysis showed that, as with 2C3 described above, the 4F7, 4H5, 5A5, 5A9, 6E4, and 6H4 antibodies specifically bind to A $\beta$  oligomers but do not recognize A $\beta$  monomers (see Fig. 14).

[Example 15]

#### Inhibition ELISA

**[0163]** To assess the A $\beta$  oligomer-selective binding activity of the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4), each antibody was mixed with stepwise-diluted A $\beta$  oligomers or monomers ("inhibitors"), and the pre-mixed solutions were added to A $\beta$  oligomer-immobilized 96-well immunoplates, and then incubated (see the "Methods" section). The commercially available 4G8 and 6E10 antibodies were used as control antibodies that nonselectively bind to A $\beta$  oligomers and monomers. When an antibody selectively binds to A $\beta$  oligomers, the antibody pre-mixed with A $\beta$  monomers does not bind to the A $\beta$  monomers in the solution, and therefore can bind to immobilized A $\beta$  oligomers. On the other hand, the antibody pre-mixed with A $\beta$  oligomers binds to the A $\beta$  oligomers in the solution, and therefore the amount of antibody bound to immobilized A $\beta$  oligomers is reduced with the increase in inhibitor concentration. The results for the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) showed concentration-dependent reduction in the amount of bound antibody when A $\beta$  oligomers were used. In contrast, no such strong reduction in binding was detected when A $\beta$  monomers were used (see Fig. 15). Meanwhile, for 4G8 and 6E10, the concentration-dependent reduction in the amount of bound antibody was observed when A $\beta$  monomers and oligomers were used (see Fig. 15). These results suggest that the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) selectively bind to A $\beta$  oligomers.

[Example 16]

#### The activity of the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) to neutralize A $\beta$ -induced neurotoxicity

**[0164]** To assess whether the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) have an activity of neutralizing A $\beta$ -induced neurotoxicity, human neuroblastoma cells (SH-SY5Y) were cultured in a medium containing A $\beta$  1-42 (12.5  $\mu$ M) in the presence or absence of the antibodies for 24 hours, and the change in A $\beta$  1-42-induced cytotoxicity was monitored. As a result, the cytotoxicity was enhanced by addition of control IgG (3F1). Although the cytotoxicity was also increased by addition of the 4F7 and 4H5 antibodies, the increase was smaller than that observed for 3F1 (see Fig. 16). The remaining four types of antibodies (5A5, 5A9, 6E4, and 6H4) were found to markedly reduce the cytotoxicity (see Fig. 16). The results described above demonstrate that the four types of antibodies (5A5, 5A9, 6E4, and 6H4) have a strong activity of neutralizing A $\beta$ -induced neurotoxicity. Since 4F7 and 4H5 lowered the cytotoxicity as compared to control IgG, these antibodies are also inferred to have an activity of neutralizing A $\beta$ -induced neurotoxicity.

[Example 17]

#### The activity of the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) to suppress A $\beta$ amyloid fibril formation

**[0165]** To assess whether the six types of antibodies (4F7, 4H5, 5A5, 5A9, 6E4, and 6H4) have an activity of suppressing A $\beta$  amyloid fibril formation, the formation of A $\beta$  amyloid fibrils was detected by the ThT fluorescence intensity assay method in a solution (medium) whose composition was the same as that used in the experiment for A $\beta$ -induced neurotoxicity (see the "Methods" section). 6E4 and 6H4 were found to suppress the fibril formation in an antibody concentration-dependent manner (see Fig. 17). The other four antibodies (4F7, 4H5, 5A5, and 5A9) were also inferred to have an activity of suppressing the fibril formation, since the antibodies exhibited the tendency of suppressing the fibril formation as compared to control IgG.

#### Discussion

**[0166]** The data obtained by the present inventors show that monoclonal 1A9 and 2C3 specifically recognize the "neurotoxic epitope" and "polymerization epitope" of soluble A $\beta$  polymers that are responsible for the toxic activity and antigen fibril formation activity. Since monoclonal 1A9 and 2C3 do not react with soluble A $\beta$  monomers, which are physiological molecules, it can be concluded that a three-dimensional structure having the epitope that is recognized by 1A9 or 2C3 is specific to soluble oligomeric polymers. The experiments using ultrafiltration and molecular sieve revealed that the size of 1A9- and 2C3-immunoreactive oligomers is greater than 100 kDa (>20mer). The result of morphological observation by AFM demonstrated that the toxic polymers are morphologically heterogeneous (granular, bead-shaped, and ring-shaped).

**[0167]** To demonstrate that the toxic polymers are actually bioactive molecules that exhibit *in vivo* synaptic toxicity, the present inventors commenced treatment of young Tg2576 mice before the onset of memory disturbance with anti-A $\beta$  oligomer passive immunotherapy targeting 1A9- and 2C3-recognized toxic polymers. For the first time, the present inventors presented evidence supporting that age-dependent memory deterioration that naturally develops in Tg2576



mice can be prevented by passive immunotherapy using anti-A $\beta$  oligomer-specific antibodies (1A9 and 2C3). Herein, short-term memory disturbance assessed by the Y-maze test is similar to the A $\beta$  accumulation-associated memory disturbance observed in mild cognitive impairment (MCI) and early AD. The Y-maze test showed excellent and almost normal results in Tg2576 mice administered with 1A9 and 2C3, respectively. When assessed by the novel object recognition task, Morris water maze, and contextual fear conditioning task, the long-term memory was maintained nearly normal by the anti-A $\beta$  oligomer antibodies.

**[0168]** A selective increase in A11-positive oligomers in blood was observed in the antibody-treated mouse groups as compared to the PBS treatment group, which is consistent with the ability of the antibodies to prevent the onset of memory disturbance (the memory maintenance ability). The 1A9 antibody treatment also exhibited the effect of suppressing cerebral A $\beta$  accumulation. The 2C3 antibody treatment demonstrated a higher blood level of A11-positive oligomers as compared to the 1A9 antibody treatment. However, the cerebral A $\beta$  accumulation-suppressing effect of the 2C3 antibody treatment was unclear. Accordingly, 1A9-recognized oligomers were considered to have greater contribution to the cerebral A $\beta$  accumulation than 2C3-recognized oligomers. The involvement of the polymers in cerebral A $\beta$  accumulation can be explained based on the following assumption: neurotoxic 1A9 polymers are soluble toxic oligomers that are somewhat conformationally, while neurotoxic 2C3 polymers are very unstable, short-lived oligomeric intermediates that appear at an early stage of the polymerization process, the conformation of which is easily changed.

**[0169]** The present inventors disclose herein the *in vivo* preventive effect of anti-oligomer antibodies on Alzheimer's disease, and this is the first evidence that directly demonstrates that toxic A $\beta$  oligomers formed *in vivo* can inhibit the functions of nerve cells, thereby inducing the symptoms of Alzheimer's disease.

**[0170]** The data obtained by the present inventors is also the first evidence supporting the view that A $\beta$  exhibits *in vivo* neurotoxicity in the human brain. It is well known that the human entorhinal cortex is an area that is easily affected with AD. In this area, NFT formation and nerve cell loss precede the formation of senile plaques. Thus, the entorhinal cortex is an exceptional area to which the commonly accepted amyloid cascade hypothesis cannot be applied. However, this inconsistency has been neglected and remained unstudied for a long time.

**[0171]** The present inventors proposed and examined the hypothesis that previously unidentifiable, invisible A $\beta$  oligomers are harmful for nerve cells in the entorhinal cortex and cause memory disturbance. To examine this hypothesis, the present inventors performed semi-quantitative analysis of 1A9- and 2C3-immunoreactive 12mer in the entorhinal cortex of elderly individuals who were mostly at Braak NFT stages I to III. The 1A9- and 2C3-immunoreactive 12mer were already present in the entorhinal cortex of healthy individuals at Braak NFT stages I to II, and increased with the advancement of Braak NFT stage. The 12mer was found to be significantly increased in AD. Thus, the appearance of 1A9- and 2C3-immunoreactive 12mer was demonstrated to precede the onset of cognitive impairment in the human brain. On the other hand, by biochemical and immunohistochemical techniques, it was demonstrated that senile plaques contain 1A9- and 2C3-immunoreactive A $\beta$  oligomers. In addition, insolubilized amyloid fibrils themselves were revealed to have an activity of neutralizing the neurotoxicity. These findings suggest that, under conditions where A $\beta$  oligomers are present without senile plaque formation, A $\beta$  oligomers exert *in vivo* toxicity and thus can be a cause of memory disturbance.

**[0172]** As described above, the data of the present inventors show for the first time evidence that directly demonstrates *in vivo* the memory disturbance resulting from synaptic dysfunction caused by endogenous A $\beta$  oligomers. Although active immunotherapy (Janus D, 2000, Nature; Morgan D, 2000, Nature) and passive immunotherapy (Bard F, 2222, Nat med; DeMattos RB, PNAS, 2001) have been used previously, the mechanism by which learning disability and memory disturbance can be prevented has remained a matter of conjecture. One widely proposed possibility is that the antibodies reach the brain through the blood-brain barrier and directly neutralize *in vivo* soluble A $\beta$  oligomers that cause memory impairment. The second possibility, the "sink theory", is that the antibodies act peripherally to deplete the peripheral blood A $\beta$  pool and thus activate A $\beta$  clearance from the brain. DeMattos *et al.* have reported that a peripherally administered anti-A $\beta$  antibody rapidly transports not only cerebral A $\beta$  monomers but also A $\beta$  dimers into plasma, and also cerebral A $\beta$  into CSF (DeMattos RB *et al.*, PNAS, 98; 8850-8855, 2001). The present inventors also revealed that A $\beta$  oligomers are present in human CSF and increased in AD patients. Thus, the present inventors demonstrated that the A $\beta$  oligomers can be used as diagnostic markers for AD. Furthermore, the present inventors presented the first evidence supporting the view that A $\beta$  oligomers are present in the plasma of Tg2576 mice, and, in passive immunotherapy by which A $\beta$  oligomers are specifically captured and neutralized through intravenous injection, intracerebral antibody delivery is not required and the clearance of A $\beta$  oligomers from the brain to blood can be enhanced at the peripheral sites of action, *i.e.*, blood vessels. In addition, the present inventors presented the first evidence that passive immunotherapy can suppress senile amyloid plaque formation, and indirectly suppress nerve cell damage (swollen dystrophic neurite formation) through senile amyloid plaque suppression. These results confirm that the A $\beta$  oligomer is the molecular basis for the onset of Alzheimer's disease, and selective control using oligomer-specific antibodies enables the control of Alzheimer's disease from a prophylactic viewpoint, in addition to a therapeutic viewpoint. Furthermore, a fraction of the administered antibodies was proven to translocate into the brain. This suggests that the effect of suppressing memory disturbance is exerted by a combination of multiple actions such as direct neutralization of soluble A $\beta$  oligomers in the

brain, transport of antibody-A $\beta$  oligomer immune complexes into blood by the neonatal Fc receptor (Deane R, 2005, J Neurosci), and the "sink" action described above.

**[0173]** The establishment of accurate pre-onset diagnosis to identify cases at a high risk of developing AD is essential to design preventive/therapeutic strategies. The significant increase in the CSF O/M ratio in AD, which is reported herein, is expected to be one of the leading candidates for pre-onset diagnostic markers.

#### Industrial Applicability

**[0174]** The antibodies provided by the present invention can be used, for example, in intravenous injection-based preventive passive immunotherapy for Alzheimer's disease, and as biological markers for pre-onset diagnosis, disease monitoring, drug efficacy monitoring/assessment for the disease, and such.

**[0175]** Furthermore, the antibodies of the present invention are expected to greatly contribute to the establishment of preventive/therapeutic methods for Alzheimer's disease that are selective to molecules responsible for evoking the pathological conditions of the disease, and the establishment of early diagnostic markers. The present inventors obtained evidence supporting that antibody therapies, even when they target intracerebral pathological conditions, can be satisfactorily achieved by peripheral intravenous administration, without the need to consider intracerebral transfer of the antibodies. In addition, the present inventors obtained evidence demonstrating that a fraction of administered antibodies translocates to the brain and produces a direct effect even in peripheral intravenous administration therapy, again without the need to consider intracerebral transfer of the antibodies. Thus, the present invention is expected to rapidly accelerate the progress of antibody therapeutics for Alzheimer's disease.

#### SEQUENCE LISTING

##### **[0176]**

<110> IMMUNAS PHARMA, INC.

<120> ANTIBODY CAPABLE OF BINDING SPECIFICALLY TO AB-OLIGOMER, AND USE THEREOF

<130> S67372PCEPT1

<140> PCT/JP2009/052039

<141> 2009-02-06

<150> JP 2008-28386

<151> 2008-02-08

<150> US 61/085,545

<151> 2008-08-01

<150> JP 2008-201058

<151> 2008-08-04

<160> 120

<170> PatentIn version 3.4

<210> 1

<211> 452

<212> PRT

<213> Mus musculus

<400> 1

EP 3 121 277 B1

	Asp	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Arg	Lys	Leu	Ser	Cys	Ala	Thr	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Phe	
				20					25					30			
	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Glu	Lys	Gly	Leu	Glu	Trp	Val	
			35					40					45				
10	Ala	Tyr	Ile	Ser	Ser	Gly	Ser	Ser	Ala	Ile	Tyr	Tyr	Ala	Asp	Thr	Val	
		50					55					60					
	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Pro	Lys	Asn	Thr	Leu	Phe	
	65					70					75					80	
15	Leu	Gln	Met	Thr	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	
					85					90					95		
	Ala	Arg	Ser	Gly	Asp	Thr	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Ser	Val	
				100					105					110			
20	Thr	Val	Ser	Ser	Ala	Lys	Thr	Thr	Pro	Pro	Ser	Val	Tyr	Pro	Leu	Ala	
				115				120					125				
	Pro	Gly	Cys	Gly	Asp	Thr	Thr	Gly	Ser	Ser	Val	Thr	Leu	Gly	Cys	Leu	
25		130					135					140					
	Val	Lys	Gly	Tyr	Phe	Pro	Glu	Ser	Val	Thr	Val	Thr	Trp	Asn	Ser	Gly	
	145					150					155					160	
30	Ser	Leu	Ser	Ser	Ser	Val	His	Thr	Phe	Pro	Ala	Leu	Leu	Gln	Ser	Gly	

EP 3 121 277 B1

	165							170					175			
5	Leu	Tyr	Thr	Met	Ser	Ser	Ser	Val	Thr	Val	Pro	Ser	Ser	Thr	Trp	Pro
				180					185					190		
	Ser	Gln	Thr	Val	Thr	Cys	Ser	Val	Ala	His	Pro	Ala	Ser	Ser	Thr	Thr
			195					200					205			
10	Val	Asp	Lys	Lys	Leu	Glu	Pro	Ser	Gly	Pro	Ile	Ser	Thr	Ile	Asn	Pro
	210						215					220				
	Cys	Pro	Pro	Cys	Lys	Glu	Cys	His	Lys	Cys	Pro	Ala	Pro	Asn	Leu	Glu
	225					230					235					240
15	Gly	Gly	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Asn	Ile	Lys	Asp	Val	Leu
					245					250					255	
	Met	Ile	Ser	Leu	Thr	Pro	Lys	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
				260					265					270		
20	Glu	Asp	Asp	Pro	Asp	Val	Gln	Ile	Ser	Trp	Phe	Val	Asn	Asn	Val	Glu
		275						280					285			
	Val	His	Thr	Ala	Gln	Thr	Gln	Thr	His	Arg	Glu	Asp	Tyr	Asn	Ser	Thr
		290					295					300				
25	Ile	Arg	Val	Val	Ser	Thr	Leu	Pro	Ile	Gln	His	Gln	Asp	Trp	Met	Ser
	305					310					315					320
	Gly	Lys	Glu	Phe	Lys	Cys	Lys	Val	Asn	Asn	Lys	Asp	Leu	Pro	Ser	Pro
30					325					330					335	
	Ile	Glu	Arg	Thr	Ile	Ser	Lys	Ile	Lys	Gly	Leu	Val	Arg	Ala	Pro	Gln
				340					345					350		
	Val	Tyr	Ile	Leu	Pro	Pro	Pro	Ala	Glu	Gln	Leu	Ser	Arg	Lys	Asp	Val
35			355					360					365			
	Ser	Leu	Thr	Cys	Leu	Val	Val	Gly	Phe	Asn	Pro	Gly	Asp	Ile	Ser	Val
		370					375					380				
40	Glu	Trp	Thr	Ser	Asn	Gly	His	Thr	Glu	Glu	Asn	Tyr	Lys	Asp	Thr	Ala
	385					390					395					400
	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Tyr	Phe	Ile	Tyr	Ser	Lys	Leu	Asn
					405					410					415	
45	Met	Lys	Thr	Ser	Lys	Trp	Glu	Lys	Thr	Asp	Ser	Phe	Ser	Cys	Asn	Val
				420					425					430		
	Arg	His	Glu	Gly	Leu	Lys	Asn	Tyr	Tyr	Leu	Lys	Lys	Thr	Ile	Ser	Arg
50			435					440					445			
	Ser	Pro	Gly	Lys												
				450												

<210> 2

55 <211> 1359

<212> DNA

<213> Mus musculus

EP 3 121 277 B1

<400> 2

5	gatgtgcagc tgggtggagtc tgggggaggc ttagtgcagc ctggagggtc ccggaaactc	60
	tcctgtgcaa cctctggatt cactttcagt agctttggaa tgcactgggt tcgtcaggct	120
	ccagagaagg gactggagtg ggtcgcatac attagtagtg gcagtagtgc catctactat	180
10	gcagacacag tgaagggccg attcaccatc tccagagaca atcccaagaa caccctgttc	240
	ctgcaaatga ccagtctaag gtctgaggac acggccatgt attactgtgc aagatctggg	300
15	gatactatgg actactgggg tcaaggaacc tcagtcaccg tctcctcagc caaaacaaca	360
	cccccatcag tctatccact ggcccctggg tgtggagata caactggttc ctccgtgact	420
	ctgggatgcc tgggtcaagg ctacttccct gagtcagtga ctgtgacttg gaactctgga	480
20	tccctgtcca gcagtgtgca caccttccca gctctcctgc agtctggact ctacactatg	540
	agcagctcag tgactgtccc ctccagcacc tggccaagtc agaccgtcac ctgcagcggt	600
	gctcaccag ccagcagcac cacggtggac aaaaaacttg agcccagcgg gccatttca	660
25	acaatcaacc cctgtcctcc atgcaaggag tgtcacaaat gccagctcc taacctcgag	720
	ggtggaccat ccgtcttcat cttccctcca aatatcaagg atgtactcat gatctccctg	780
	acacccaagg tcacgtgtgt ggtggtggat gtgagcgagg atgaccaga cgtccagatc	840
30	agctggtttg tgaacaacgt ggaagtacac acagctcaga cacaaccca tagagaggat	900
	tacaacagta ctatccgggt ggtcagcacc ctccccatcc agcaccagga ctggatgagt	960
35	ggcaaggagt tcaaatgcaa ggtcaacaac aaagacctcc catcacccat cgagagaacc	1020
	atctcaaaaa ttaaagggt agtcagagct ccacaagtat acatcttgcc gccaccagca	1080
	gagcagttgt ccaggaaaga tgtcagtctc acttgccctgg tcgtgggctt caaccctgga	1140
40	gacatcagtg tggagtggac cagcaatggg catacagagg agaactacaa ggacaccgca	1200
	ccagtcctgg actctgacgg ttcttacttc atatatagca agctcaatat gaaaacaagc	1260
	aagtgggaga aaacagattc cttctcatgc aacgtgagac acgagggctt gaaaaattac	1320
45	tacctgaaga agaccatctc ccggtctccg ggtaaatga	1359

<210> 3

<211> 219

50 <212> PRT

<213> Mus musculus

<400> 3

55

# EP 3 121 277 B1

	Asp	Val	Leu	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly
	1				5					10					15	
5	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Asn	Ile	Val	His	Ser
				20					25					30		
	Asn	Gly	Asn	Thr	Tyr	Leu	Glu	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
			35					40					45			
10	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
		50					55					60				
	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
15	65					70					75					80
	Ser	Ser	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Tyr	Cys	Phe	Gln	Val
					85					90					95	
20	Ser	His	Val	Pro	Pro	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys
				100					105					110		
	Arg	Ala	Asp	Ala	Ala	Pro	Thr	Val	Ser	Ile	Phe	Pro	Pro	Ser	Ser	Glu
			115					120					125			
25	Gln	Leu	Thr	Ser	Gly	Gly	Ala	Ser	Val	Val	Cys	Phe	Leu	Asn	Asn	Phe
		130					135					140				
	Tyr	Pro	Lys	Asp	Ile	Asn	Val	Lys	Trp	Lys	Ile	Asp	Gly	Ser	Glu	Arg
	145					150					155					160
30	Gln	Asn	Gly	Val	Leu	Asn	Ser	Trp	Thr	Asp	Gln	Asp	Ser	Lys	Asp	Ser
					165					170					175	
	Thr	Tyr	Ser	Met	Ser	Ser	Thr	Leu	Thr	Leu	Thr	Lys	Asp	Glu	Tyr	Glu
35				180					185					190		
	Arg	His	Asn	Ser	Tyr	Thr	Cys	Glu	Ala	Thr	His	Lys	Thr	Ser	Thr	Ser
			195					200					205			
40	Pro	Ile	Val	Lys	Ser	Phe	Asn	Arg	Asn	Glu	Cys					
		210					215									

<210> 4

<211> 660

<212> DNA

45 <213> Mus musculus

<400> 4

50

55

# EP 3 121 277 B1

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60  
 atctcttgca gatctagtca gaacattgta catagtaatg gaaacaccta tttagaatgg 120  
 5 tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180  
 tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240  
 agcagcgtgg aggctgagga tctgggagtt tattactgct ttcaagtttc acatgttcct 300  
 10 ccgacgttcg gtggaggcac caagctggaa atcaaacggg ctgatgctgc accaactgta 360  
 tccatcttcc caccatccag tgagcagtta acatctggag gtgcctcagt cgtgtgcttc 420  
 ttgaacaact tctaccccaa agacatcaat gtcaagtgga agattgatgg cagtgaacga 480  
 15 caaaatggcg tcctgaacag ttggactgat caggacagca aagacagcac ctacagcatg 540  
 agcagtaccc tcacgttgac caaggacgag tatgaacgac ataacagcta tacctgtgag 600  
 20 gccactcaca agacatcaac ttcacccatt gtcaagagct tcaacaggaa tgagtgttag 660

<210> 5

<211> 116

<212> PRT

25 <213> Mus musculus

<400> 5

30 Asp Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1 5 10 15  
 Ser Arg Lys Leu Ser Cys Ala Thr Ser Gly Phe Thr Phe Ser Ser Phe  
 20 25 30  
 35 Gly Met His Trp Val Arg Gln Ala Pro Glu Lys Gly Leu Glu Trp Val  
 35 40 45  
 Ala Tyr Ile Ser Ser Gly Ser Ser Ala Ile Tyr Tyr Ala Asp Thr Val  
 50 55 60  
 40 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Pro Lys Asn Thr Leu Phe  
 65 70 75 80  
 Leu Gln Met Thr Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys  
 45 85 90 95  
 Ala Arg Ser Gly Asp Thr Met Asp Tyr Trp Gly Gln Gly Thr Ser Val  
 100 105 110  
 Thr Val Ser Ser  
 50 115

<210> 6

<211> 348

<212> DNA

55 <213> Mus musculus

<400> 6

EP 3 121 277 B1

gatgtgcagc tgggtggagtc tggggggaggc ttagtgcagc ctggagggtc ccggaaactc 60  
 tcctgtgcaa cctctggatt cactttcagt agctttggaa tgcactgggt tcgtcaggct 120  
 5 ccagagaagg gactggagtg ggtcgcatac attagtagtg gcagtagtgc catctactat 180  
 gcagacacag tgaagggccg attcaccatc tccagagaca atcccaagaa caccctgttc 240  
 ctgcaaataa ccagtctaag gtctgaggac acggccatgt attactgtgc aagatctggg 300  
 10 gatactatgg actactgggg tcaaggaacc tcagtcaccg tctcctca 348

<210> 7

<211> 112

15 <212> PRT

<213> Mus musculus

<400> 7

20 Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly  
 1 5 10 15  
 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Asn Ile Val His Ser  
 20 25 30  
 25 Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
 35 40 45  
 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
 30 50 55 60  
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80  
 35 Ser Ser Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Val  
 85 90 95  
 Ser His Val Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys  
 100 105 110

40 <210> 8

<211> 336

<212> DNA

<213> Mus musculus

45 <400> 8

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60  
 50 atctcttgca gatctagtca gaacattgta catagtaatg gaaacaccta tttagaatgg 120  
 tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180  
 tctgggggtcc cagacaggtt cagtggcagt ggatcaggga cagatttcac actcaagatc 240  
 55 agcagcgtgg aggctgagga tctgggagtt tattactgct ttcaagtttc acatgttcct 300  
 ccgacgttcg gtggaggcac caagctggaa atcaaa 336



EP 3 121 277 B1

<210> 9  
 <211> 5  
 <212> PRT  
 <213> Mus musculus  
 5  
 <400> 9  
  
 Ser Phe Gly Met His  
 1 5  
 10  
 <210> 10  
 <211> 15  
 <212> DNA  
 <213> Mus musculus  
 15  
 <400> 10  
 agcttggaa tgcac 15  
 20  
 <210> 11  
 <211> 10  
 <212> PRT  
 <213> Mus musculus  
 25  
 <400> 11  
  
 Tyr Ile Ser Ser Gly Ser Ser Ala Ile Tyr  
 1 5 10  
 30  
 <210> 12  
 <211> 30  
 <212> DNA  
 <213> Mus musculus  
 35  
 <400> 12  
 tacattagta gtggcagtag tgccatctac 30  
 40  
 <210> 13  
 <211> 7  
 <212> PRT  
 <213> Mus musculus  
 45  
 <400> 13  
  
 Ser Gly Asp Thr Met Asp Tyr  
 1 5  
 50  
 <210> 14  
 <211> 21  
 <212> DNA  
 <213> Mus musculus  
 55  
 <400> 14  
 tctggggata ctatggacta c 21  
 60  
 <210> 15  
 <211> 16  
 <212> PRT  
 <213> Mus musculus

# EP 3 121 277 B1

<400> 15

	Arg	Ser	Ser	Gln	Asn	Ile	Val	His	Ser	Asn	Gly	Asn	Thr	Tyr	Leu	Glu
	1				5					10					15	

5

<210> 16

<211> 48

<212> DNA

<213> Mus musculus

10

<400> 16

agatctagtc agaacattgt acatagtaat ggaaacacct atttagaa 48

<210> 17

15

<211> 7

<212> PRT

<213> Mus musculus

<400> 17

20

	Lys	Val	Ser	Asn	Arg	Phe	Ser
	1				5		

<210> 18

25

<211> 21

<212> DNA

<213> Mus musculus

<400> 18

30

aaagtttcca accgatttc t 21

<210> 19

<211> 9

<212> PRT

35

<213> Mus musculus

<400> 19

Phe Gln Val Ser His Val Pro Pro Thr

40

1

5

<210> 20

45

<211> 27

<212> DNA

<213> Mus musculus

<400> 20

50

tttcaagttt cacatgttcc tccgacg 27

<210> 21

<211> 455

<212> PRT

55

<213> Mus musculus

<400> 21

EP 3 121 277 B1

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Ile	Leu	Gln	Pro	Ser	Gln
	1				5					10					15	
5	Thr	Leu	Ser	Leu	Thr	Cys	Ser	Phe	Ser	Gly	Phe	Ser	Leu	Thr	Thr	Ser
				20					25					30		
	Ala	Met	Gly	Val	Ser	Trp	Val	Arg	Gln	Pro	Ser	Arg	Lys	Gly	Leu	Glu
			35					40					45			
10	Trp	Leu	Ala	His	Ile	Tyr	Trp	Asp	Asp	Asp	Lys	Arg	Tyr	Asn	Pro	Ser
	50					55						60				
	Leu	Lys	Ser	Arg	Leu	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Ser	Asn	Gln	Val
	65					70					75					80
15	Phe	Leu	Lys	Ile	Thr	Ser	Val	Asp	Thr	Ala	Asp	Thr	Ala	Thr	Tyr	Tyr
					85					90					95	
	Cys	Ala	Arg	Lys	Gly	Leu	Gly	Gly	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly
20				100					105					110		
	Thr	Ser	Val	Thr	Val	Ser	Ser	Ala	Lys	Thr	Thr	Pro	Pro	Ser	Val	Tyr
			115					120					125			
25	Pro	Leu	Ala	Pro	Gly	Cys	Gly	Asp	Thr	Thr	Gly	Ser	Ser	Val	Thr	Leu
	130					135						140				
	Gly	Cys	Leu	Val	Lys	Gly	Tyr	Phe	Pro	Glu	Ser	Val	Thr	Val	Thr	Trp
	145					150					155					160
30	Asn	Ser	Gly	Ser	Leu	Ser	Ser	Ser	Val	His	Thr	Phe	Pro	Ala	Leu	Leu
					165					170					175	
	Gln	Ser	Gly	Leu	Tyr	Thr	Met	Ser	Ser	Ser	Val	Thr	Val	Pro	Ser	Ser
				180					185					190		
35	Thr	Trp	Pro	Ser	Gln	Thr	Val	Thr	Cys	Ser	Val	Ala	His	Pro	Ala	Ser
			195					200					205			
	Ser	Thr	Thr	Val	Asp	Lys	Lys	Leu	Glu	Pro	Ser	Gly	Pro	Ile	Ser	Thr
	210					215						220				
40	Ile	Asn	Pro	Cys	Pro	Pro	Cys	Lys	Glu	Cys	His	Lys	Cys	Pro	Ala	Pro
	225					230					235					240
45	Asn	Leu	Glu	Gly	Gly	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Asn	Ile	Lys

EP 3 121 277 B1

				245					250					255			
		Asp	Val	Leu	Met	Ile	Ser	Leu	Thr	Pro	Lys	Val	Thr	Cys	Val	Val	Val
5					260					265					270		
		Asp	Val	Ser	Glu	Asp	Asp	Pro	Asp	Val	Gln	Ile	Ser	Trp	Phe	Val	Asn
				275					280					285			
		Asn	Val	Glu	Val	His	Thr	Ala	Gln	Thr	Gln	Thr	His	Arg	Glu	Asp	Tyr
10			290					295					300				
		Asn	Ser	Thr	Ile	Arg	Val	Val	Ser	Thr	Leu	Pro	Ile	Gln	His	Gln	Asp
		305				310						315					320
15		Trp	Met	Ser	Gly	Lys	Glu	Phe	Lys	Cys	Lys	Val	Asn	Asn	Lys	Asp	Leu
					325						330					335	
		Pro	Ser	Pro	Ile	Glu	Arg	Thr	Ile	Ser	Lys	Ile	Lys	Gly	Leu	Val	Arg
					340					345					350		
20		Ala	Pro	Gln	Val	Tyr	Ile	Leu	Pro	Pro	Pro	Ala	Glu	Gln	Leu	Ser	Arg
				355					360					365			
		Lys	Asp	Val	Ser	Leu	Thr	Cys	Leu	Val	Val	Gly	Phe	Asn	Pro	Gly	Asp
			370					375					380				
25		Ile	Ser	Val	Glu	Trp	Thr	Ser	Asn	Gly	His	Thr	Glu	Glu	Asn	Tyr	Lys
		385					390					395					400
		Asp	Thr	Ala	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Tyr	Phe	Ile	Tyr	Ser
30					405						410					415	
		Lys	Leu	Asn	Met	Lys	Thr	Ser	Lys	Trp	Glu	Lys	Thr	Asp	Ser	Phe	Ser
				420						425					430		
		Cys	Asn	Val	Arg	His	Glu	Gly	Leu	Lys	Asn	Tyr	Tyr	Leu	Lys	Lys	Thr
35				435					440					445			
		Ile	Ser	Arg	Ser	Pro	Gly	Lys									
			450					455									

40 <210> 22  
 <211> 1368  
 <212> DNA  
 <213> Mus musculus

45 <400> 22

50

55

EP 3 121 277 B1

	caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg	60
	acttgttctt tctctggatt ttactgacc acttctgcta tgggtgtgag ctgggttcgt	120
5	cagccttcaa gaaagggctc ggagtggctg gcacacattt actgggatga tgacaagcgc	180
	tataacccat ccctgaagag ccggctcaca atctccaagg atacctccag caaccaggta	240
10	ttcctcaaga tcaccagtgt ggacactgca gatactgcca catactactg tgctcgaaag	300
	ggactgggag gtgctatgga ctactggggt caaggaacct cagtcaccgt ctctcagcc	360
	aaaacaacac ccccatcagt ctatccactg gccctgggt gtggagatac aactgggtcc	420
15	tccgtgactc tgggatgcct ggtcaagggc tacttccctg agtcagtgac tgtgacttgg	480
	aactctggat ccctgtccag cagtgtgcac accttcccag ctctcctgca gtctggactc	540
20	tacactatga gcagctcagt gactgtcccc tccagcacct ggccaagtca gaccgtcacc	600
	tgcagcgttg ctcacccagc cagcagcacc acggtggaca aaaaacttga gccagcggg	660
	cccatttcaa caatcaaccc ctgtcctcca tgcaaggagt gtcacaaatg cccagctcct	720
25	aacctcgagg gtggaccatc cgtcttcac ttcctccaa atatcaagga tgtactcatg	780
	atctccctga cacccaaggt cacgtgtgtg gtggtggatg tgagcgagga tgaccagac	840
	gtccagatca gctggtttgt gaacaacgtg gaagtacaca cagctcagac acaaaccat	900
30	agagaggatt acaacagtac tatccgggtg gtcagcaccc tccccatcca gcaccaggac	960
	tggatgagtg gcaaggagtt caaatgcaag gtcaacaaca aagacctccc atcacccatc	1020
35	gagagaacca tctcaaaaat taaagggcta gtcagagctc cacaagtata catcttgccg	1080
	ccaccagcag agcagttgtc caggaaagat gtcagtctca cttgcctgggt cgtgggcttc	1140
	aaccctggag acatcagtgt ggagtggacc agcaatgggc atacagagga gaactacaag	1200
40	gacaccgcac cagtcctgga ctctgacggt tcttacttca tatatagcaa gctcaatatg	1260
	aaaacaagca agtgggagaa aacagattcc ttctcatgca acgtgagaca cgagggtctg	1320
45	aaaaattact acctgaagaa gaccatctcc cggctctccg gtaaatga	1368
	<210> 23	
	<211> 219	
	<212> PRT	
	<213> Mus musculus	
50	<400> 23	
55		

# EP 3 121 277 B1

	Asp	Val	Val	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly	
	1				5					10					15		
5	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Leu	His	Ser	
				20					25					30			
	Asn	Gly	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser	
			35					40					45				
10	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
		50					55					60					
	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75					80	
15	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Phe	Cys	Ser	Gln	Ser	
					85					90					95		
	Thr	His	Val	Pro	Leu	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Glu	Leu	Lys	
				100					105					110			
20	Arg	Ala	Asp	Ala	Ala	Pro	Thr	Val	Ser	Ile	Phe	Pro	Pro	Ser	Ser	Glu	
			115					120					125				
	Gln	Leu	Thr	Ser	Gly	Gly	Ala	Ser	Val	Val	Cys	Phe	Leu	Asn	Asn	Phe	
25		130					135					140					
	Tyr	Pro	Lys	Asp	Ile	Asn	Val	Lys	Trp	Lys	Ile	Asp	Gly	Ser	Glu	Arg	
	145					150					155					160	
30	Gln	Asn	Gly	Val	Leu	Asn	Ser	Trp	Thr	Asp	Gln	Asp	Ser	Lys	Asp	Ser	
					165					170					175		
	Thr	Tyr	Ser	Met	Ser	Ser	Thr	Leu	Thr	Leu	Thr	Lys	Asp	Glu	Tyr	Glu	
				180					185					190			
35	Arg	His	Asn	Ser	Tyr	Thr	Cys	Glu	Ala	Thr	His	Lys	Thr	Ser	Thr	Ser	
			195					200					205				
	Pro	Ile	Val	Lys	Ser	Phe	Asn	Arg	Asn	Glu	Cys						
40		210					215										

<210> 24

<211> 660

<212> DNA

45 <213> Mus musculus

<400> 24

50

55

EP 3 121 277 B1

gatgttgtga tgacccaaac tccgctctcc ctgcctgtca gtcttggaga tcaagcctcc 60  
 atctcttgca gatctagtca gagccttcta cacagtaatg gaaacaccta ttacattgg 120  
 5 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180  
 tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240  
 agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttccg 300  
 10 ctacagttcg gtgctgggac caagctggag ctgaaacggg ctgatgctgc accaactgta 360  
 tccatcttcc caccatccag tgagcagtta acatctggag gtgcctcagt cgtgtgcttc 420  
 ttgaacaact tctaccccaa agacatcaat gtcaagtgga agattgatgg cagtgaacga 480  
 15 caaaatggcg tcctgaacag ttggactgat caggacagca aagacagcac ctacagcatg 540  
 agcagtaccc tcacgttgac caaggacgag tatgaacgac ataacagcta tacctgtgag 600  
 20 gccactcaca agacatcaac ttcacccatt gtcaagagct tcaacaggaa tgagtgttag 660

<210> 25

<211> 119

<212> PRT

25 <213> Mus musculus

<400> 25

30 Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln  
 1 5 10 15  
 Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Thr Thr Ser  
 20 25 30  
 35 Ala Met Gly Val Ser Trp Val Arg Gln Pro Ser Arg Lys Gly Leu Glu  
 35 40 45  
 Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser  
 50 55 60  
 40 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn Gln Val  
 65 70 75 80  
 45 Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr  
 85 90 95  
 Cys Ala Arg Lys Gly Leu Gly Gly Ala Met Asp Tyr Trp Gly Gln Gly  
 100 105 110  
 50 Thr Ser Val Thr Val Ser Ser  
 115

<210> 26

<211> 357

55 <212> DNA

<213> Mus musculus

<400> 26

EP 3 121 277 B1

caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60  
 acttggttctt tctctggatt ttcactgacc acttctgcta tgggtgtgag ctgggttcgt 120  
 5 cagccttcaa gaaagggctc ggagtggctg gcacacattt actgggatga tgacaagcgc 180  
 tataacccat ccctgaagag ccggtcaca atctccaagg atacctccag caaccaggta 240  
 ttcctcaaga tcaccagtgt ggacactgca gatactgcca catactactg tgctcgaaag 300  
 10 ggactgggag gtgctatgga ctactggggt caaggaacct cagtcaccgt ctctca 357

<210> 27

<211> 112

15 <212> PRT

<213> Mus musculus

<400> 27

20 Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly  
 1 5 10 15  
 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser  
 20 25 30  
 25 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
 35 40 45  
 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
 50 55 60  
 30 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80  
 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser  
 85 90 95  
 35 Thr His Val Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys  
 100 105 110

40 <210> 28

<211> 336

<212> DNA

<213> Mus musculus

45 <400> 28

gatgttgtga tgacccaaac tccgctctcc ctgcctgtca gtcttgaga tcaagcctcc 60  
 atctcttgca gatctagtca gagccttcta cacagtaatg gaaacaccta ttacattgg 120  
 50 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180  
 tctgggggtcc cagacagggtt cagtggcagt ggatcaggga cagatttcac actcaagatc 240  
 agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttccg 300  
 55 ctacagttcg gtgctgggac caagctggag ctgaaa 336



# EP 3 121 277 B1

<210> 29  
 <211> 7  
 <212> PRT  
 <213> Mus musculus  
 5  
 <400> 29  
  
 Thr Ser Ala Met Gly Val Ser  
 1 5  
 10  
 <210> 30  
 <211> 21  
 <212> DNA  
 <213> Mus musculus  
 15  
 <400> 30  
 acttctgcta tgggtgtgag c 21  
 20  
 <210> 31  
 <211> 9  
 <212> PRT  
 <213> Mus musculus  
 25  
 <400> 31  
  
 His Ile Tyr Trp Asp Asp Asp Lys Arg  
 1 5  
 30  
 <210> 32  
 <211> 27  
 <212> DNA  
 <213> Mus musculus  
 35  
 <400> 32  
 cacatttact gggatgatga caagcgc 27  
 40  
 <210> 33  
 <211> 9  
 <212> PRT  
 <213> Mus musculus  
 45  
 <400> 33  
  
 Lys Gly Leu Gly Gly Ala Met Asp Tyr  
 1 5  
 50  
 <210> 34  
 <211> 27  
 <212> DNA  
 <213> Mus musculus  
 55  
 <400> 34  
 aagggactgg gaggtgctat ggactac 27  
 60  
 <210> 35  
 <211> 16  
 <212> PRT  
 <213> Mus musculus

# EP 3 121 277 B1

<400> 35

5                   Arg Ser Ser Gln Ser Leu Leu His Ser Asn Gly Asn Thr Tyr Leu His  
                  1                   5                   10                   15

<210> 36  
<211> 48  
<212> DNA  
<213> Mus musculus

10  
<400> 36  
agatctagtc agagccttct acacagtaat ggaaacacct attacat           48

15  
<210> 37  
<211> 7  
<212> PRT  
<213> Mus musculus

20  
<400> 37

                  Lys Val Ser Asn Arg Phe Ser  
                  1                   5

25  
<210> 38  
<211> 21  
<212> DNA  
<213> Mus musculus

30  
<400> 38  
aaagttcca accgatttc t           21

35  
<210> 39  
<211> 9  
<212> PRT  
<213> Mus musculus

<400> 39

40                   Ser Gln Ser Thr His Val Pro Leu Thr  
                  1                   5

45  
<210> 40  
<211> 27  
<212> DNA  
<213> Mus musculus

<400> 40  
tctcaaagta cacatgtcc gctcacg           27

50  
<210> 41  
<211> 447  
<212> PRT  
<213> Mus musculus

55  
<400> 41

EP 3 121 277 B1

	Gln	Val	Thr	Leu	Lys	Asp	Ser	Gly	Pro	Gly	Ile	Leu	Gln	Pro	Ser	Gln	
	1				5					10					15		
5	Thr	Leu	Ser	Leu	Thr	Cys	Ser	Phe	Ser	Gly	Phe	Ser	Leu	Ser	Thr	Ser	
				20					25					30			
10	Gly	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Ser	Gly	Lys	Gly	Leu	Glu	
			35					40					45				
15	Trp	Leu	Ala	His	Ile	Tyr	Trp	Asp	Asp	Asp	Lys	Arg	Tyr	Asn	Pro	Ser	
	50						55					60					
20	Leu	Lys	Ser	Arg	Leu	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Ser	Asn	Gln	Val	
	65					70					75					80	
25	Phe	Leu	Lys	Ile	Thr	Ser	Val	Asp	Thr	Ala	Asp	Ser	Ala	Thr	Tyr	Tyr	
					85					90					95		
30	Cys	Ser	Thr	Met	Ile	Thr	Gly	Phe	Val	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	
				100					105					110			
35	Val	Thr	Val	Ser	Ala	Ala	Lys	Thr	Thr	Ala	Pro	Ser	Val	Tyr	Pro	Leu	
			115					120					125				
40	Ala	Pro	Val	Cys	Gly	Asp	Thr	Thr	Gly	Ser	Ser	Val	Ala	Leu	Gly	Cys	
		130					135					140					
45	Leu	Val	Lys	Gly	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Leu	Thr	Trp	Asn	Ser	
	145					150					155					160	
50	Gly	Ser	Leu	Ser	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	
					165					170					175		
55	Asp	Leu	Tyr	Thr	Leu	Ser	Ser	Ser	Val	Thr	Val	Thr	Ser	Ser	Thr	Trp	
				180					185					190			
60	Pro	Ser	Gln	Ser	Ile	Thr	Cys	Asn	Val	Ala	His	Pro	Ala	Ser	Ser	Thr	
			195					200					205				
65	Lys	Val	Asp	Lys	Lys	Ile	Glu	Pro	Arg	Gly	Pro	Thr	Ile	Lys	Pro	Cys	
		210					215					220					
70	Pro	Pro	Cys	Lys	Cys	Pro	Ala	Pro	Asn	Leu	Leu	Gly	Gly	Pro	Ser	Val	
	225					230					235					240	

# EP 3 121 277 B1

	Phe	Ile	Phe	Pro	Pro	Lys	Ile	Lys	Asp	Val	Leu	Met	Ile	Ser	Leu	Ser	
					245					250					255		
5	Pro	Ile	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Glu	Asp	Asp	Pro	Asp	
				260					265					270			
10	Val	Gln	Ile	Ser	Trp	Phe	Val	Asn	Asn	Val	Glu	Val	His	Thr	Ala	Gln	
			275					280					285				
15	Thr	Gln	Thr	His	Arg	Glu	Asp	Tyr	Asn	Ser	Thr	Leu	Arg	Val	Val	Ser	
		290					295					300					
20	Ala	Leu	Pro	Ile	Gln	His	Gln	Asp	Trp	Met	Ser	Gly	Lys	Glu	Phe	Lys	
	305					310					315					320	
25	Cys	Lys	Val	Asn	Asn	Lys	Asp	Leu	Pro	Ala	Pro	Ile	Glu	Arg	Thr	Ile	
				325						330					335		
30	Ser	Lys	Pro	Lys	Gly	Ser	Val	Arg	Ala	Pro	Gln	Val	Tyr	Val	Leu	Pro	
				340					345					350			
35	Pro	Pro	Glu	Glu	Glu	Met	Thr	Lys	Lys	Gln	Val	Thr	Leu	Thr	Cys	Met	
			355					360					365				
40	Val	Thr	Asp	Phe	Met	Pro	Glu	Asp	Ile	Tyr	Val	Glu	Trp	Thr	Asn	Asn	
		370					375					380					
45	Gly	Lys	Thr	Glu	Leu	Asn	Tyr	Lys	Asn	Thr	Glu	Pro	Val	Leu	Asp	Ser	
	385					390					395					400	
50	Asp	Gly	Ser	Tyr	Phe	Met	Tyr	Ser	Lys	Leu	Arg	Val	Glu	Lys	Lys	Asn	
					405					410					415		
55	Trp	Val	Glu	Arg	Asn	Ser	Tyr	Ser	Cys	Ser	Val	Val	His	Glu	Gly	Leu	
				420					425					430			
60	His	Asn	His	His	Thr	Thr	Lys	Ser	Phe	Ser	Arg	Thr	Pro	Gly	Lys		
			435					440					445				
65	<210>	42															
70	<211>	1344															
75	<212>	DNA															
80	<213>	Mus musculus															
85	<400>	42															

EP 3 121 277 B1

	caggttactc tgaaagactc tggccctggg atattgcagc cctcccagac cctcagtctg	60
	acttgttctt tctctgggtt ttcactgagc acttctggta tgggtgtgag ctggattcgt	120
5	cagccttcag gaaagggctc ggagtggctg gcacacattt actgggatga tgacaagcgc	180
	tataacccat ccctgaagag ccggctcaca atctccaagg atacctccag caaccaggta	240
10	ttcctcaaga tcaccagtgt ggacactgca gattctgcca catactactg ttccactatg	300
	attacggggt ttgtttactg gggccaaggg actctgggtca ctgtctctgc agccaaaaca	360
	acagcccat cggtctatcc cctggcccct gtgtgtggag atacaactgg ctccctcgggtg	420
15	gctctaggat gcctgggtcaa gggttatttc cctgagccag tgacctgac ctggaactct	480
	ggatccctgt ccagtgggtg gcacaccttc ccagctgtcc tgcagtctga cctctacacc	540
	ctcagcagct cagtgactgt aacctcgagc acctggccca gccagtccat cacctgcaat	600
20	gtggcccacc cggcaagcag caccaagggtg gacaagaaaa ttgagcccag agggcccaca	660
	atcaagccct gtcctccatg caaatgccc gcacctaacc tcttgggtgg accatccgtc	720
25	ttcatcttcc ctccaaagat caaggatgta ctcatgatct ccctgagccc catagtcaca	780
	tgtgtgggtg tggatgtgag cgaggatgac ccagatgtcc agatcagctg gtttgtgaac	840
	aacgtggaag tacacacagc tcagacacaa acccatagag aggattacaa cagtactctc	900
30	cgggtgggtca gtgccctccc catccagcac caggactgga tgagtggcaa ggagttcaaa	960
	tgcaagggtca acaacaaaga cctcccagcg cccatcgaga gaaccatctc aaaacccaaa	1020
	gggtcagtaa gagctccaca ggtatatgtc ttgcctccac cagaagaaga gatgactaag	1080
35	aaacagggtca ctctgacctg catggtcaca gacttcatgc ctgaagacat ttacgtggag	1140
	tggaccaaca acgggaaaac agagctaaac tacaagaaca ctgaaccagt cctggactct	1200
40	gatggttctt acttcatgta cagcaagctg agagtggaaa agaagaactg ggtggaaaga	1260
	aatagctact cctgttcagt ggtccacgag ggtctgcaca atcaccacac gactaagagc	1320
	ttctcccgga ctccgggtaa atga	1344
45	<210> 43 <211> 219 <212> PRT <213> Mus musculus	
50	<400> 43	
55		

EP 3 121 277 B1

	Asp	Val	Leu	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly
	1				5					10					15	
5	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Ile	Val	His	Ser
				20					25					30		
10	Asn	Gly	Asn	Thr	Tyr	Leu	Glu	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
			35					40					45			
15	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
		50					55					60				
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
	65					70					75					80
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Tyr	Cys	Phe	Gln	Gly
					85					90					95	
30	Ser	His	Val	Pro	Leu	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Glu	Leu	Lys
				100					105					110		
35	Arg	Ala	Asp	Ala	Ala	Pro	Thr	Val	Ser	Ile	Phe	Pro	Pro	Ser	Ser	Glu
			115					120					125			
40	Gln	Leu	Thr	Ser	Gly	Gly	Ala	Ser	Val	Val	Cys	Phe	Leu	Asn	Asn	Phe
		130					135					140				
45	Tyr	Pro	Lys	Asp	Ile	Asn	Val	Lys	Trp	Lys	Ile	Asp	Gly	Ser	Glu	Arg
	145					150					155					160
50	Gln	Asn	Gly	Val	Leu	Asn	Ser	Trp	Thr	Asp	Gln	Asp	Ser	Lys	Asp	Ser
					165					170					175	
55	Thr	Tyr	Ser	Met	Ser	Ser	Thr	Leu	Thr	Leu	Thr	Lys	Asp	Glu	Tyr	Glu
				180					185					190		
60	Arg	His	Asn	Ser	Tyr	Thr	Cys	Glu	Ala	Thr	His	Lys	Thr	Ser	Thr	Ser
			195					200					205			
65	Pro	Ile	Val	Lys	Ser	Phe	Asn	Arg	Asn	Glu	Cys					
		210					215									

<210> 44

<211> 660

<212> DNA

<213> Mus musculus

<400> 44

# EP 3 121 277 B1

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttgaga tcaagcctcc 60  
 atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120  
 5 tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180  
 tctgggggtcc cagacagggtt cagtggcagt ggatcaggga cagatttcac actcaagatc 240  
 10 agtagagtgg aggctgagga tctgggagtt tattactgct ttcaaggttc acatgttccg 300  
 ctcacgttcg gtgctgggac caagctggag ctgaaacggg ctgatgctgc accaactgta 360  
 tccatcttcc caccatccag tgagcagtta acatctggag gtgcctcagt cgtgtgcttc 420  
 15 ttgaacaact tctaccccaa agacatcaat gtcaagtgga agattgatgg cagtgaacga 480  
 caaaatggcg tcctgaacag ttggactgat caggacagca aagacagcac ctacagcatg 540  
 20 agcagtaccc tcacgttgac caaggacgag tatgaacgac ataacagcta tacctgtgag 600  
 gccactcaca agacatcaac ttcacccatt gtcaagagct tcaacaggaa tgagtgttag 660

<210> 45

<211> 117

25 <212> PRT

<213> Mus musculus

<400> 45

30 Gln Val Thr Leu Lys Asp Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln  
 1 5 10 15  
 35 Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser  
 20 25 30  
 40 Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu  
 35 40 45  
 45 Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser  
 50 55 60  
 50 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn Gln Val  
 65 70 75 80  
 55 Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Ser Ala Thr Tyr Tyr  
 85 90 95  
 Cys Ser Thr Met Ile Thr Gly Phe Val Tyr Trp Gly Gln Gly Thr Leu  
 100 105 110  
 Val Thr Val Ser Ala  
 115

# EP 3 121 277 B1

<210> 46  
 <211> 351  
 <212> DNA  
 <213> Mus musculus

5

<400> 46

10

15

20

caggttactc tgaaagactc tggccctggg atattgcagc cctcccagac cctcagtctg 60  
 acttggttctt tctctgggtt ttcactgagc acttctggta tgggtgtgag ctggattcgt 120  
 cagccttcag gaaaggggtct ggagtggctg gcacacattt actgggatga tgacaagcgc 180  
 tataacccat ccctgaagag ccggctcaca atctccaagg atacctccag caaccaggta 240  
 ttccctcaaga tcaccagtgt ggacactgca gattctgcca catactactg ttccactatg 300  
 attacgggggt ttgtttactg gggccaaggg actctgggtca ctgtctctgc a 351

25

<210> 47  
 <211> 112  
 <212> PRT  
 <213> Mus musculus

30

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly  
 1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser  
 20 25 30

35

Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
 35 40 45

40

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
 50 55 60

45

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly  
 85 90 95

50

Ser His Val Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys  
 100 105 110

55

<210> 48  
 <211> 336  
 <212> DNA  
 <213> Mus musculus



EP 3 121 277 B1

<400> 48

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60  
5 atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120  
tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180  
tctgggggtcc cagacaggtt cagtggcagt ggatcagga cagatttcac actcaagatc 240  
10 agtagagtgg aggctgagga tctgggagtt tattactgct ttcaaggttc acatgttccg 300  
ctcacgttcg gtgctgggac caagctggag ctgaaa 336

15 <210> 49  
<211> 7  
<212> PRT  
<213> Mus musculus

20 <400> 49

Thr Ser Gly Met Gly Val Ser  
1 5

25 <210> 50  
<211> 21  
<212> DNA  
<213> Mus musculus

30 <400> 50  
acttctggta tgggtgtgag c 21

<210> 51  
<211> 9  
35 <212> PRT  
<213> Mus musculus

<400> 51

40 His Ile Tyr Trp Asp Asp Asp Lys Arg  
1 5

<210> 52  
<211> 27  
45 <212> DNA  
<213> Mus musculus

<400> 52  
cacatttact gggatgatga caagcgc 27

50 <210> 53  
<211> 7  
<212> PRT  
<213> Mus musculus

55 <400> 53

# EP 3 121 277 B1

Met Ile Thr Gly Phe Val Tyr  
1 5

5 <210> 54  
<211> 21  
<212> DNA  
<213> Mus musculus

10 <400> 54  
atgattacgg ggtttgttta c 21

<210> 55  
<211> 16  
<212> PRT  
15 <213> Mus musculus

<400> 55

20 Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu  
1 5 10 15

<210> 56  
<211> 48  
<212> DNA  
25 <213> Mus musculus

<400> 56  
agatctagtc agagcattgt acatagtaat ggaaacacct atttagaa 48

30 <210> 57  
<211> 7  
<212> PRT  
<213> Mus musculus

35 <400> 57

Lys Val Ser Asn Arg Phe Ser  
1 5

40 <210> 58  
<211> 21  
<212> DNA  
<213> Mus musculus

45 <400> 58  
aaagttcca accgatttc t 21

<210> 59  
<211> 9  
50 <212> PRT  
<213> Mus musculus

<400> 59

55 Phe Gln Gly Ser His Val Pro Leu Thr  
1 5

<210> 60

# EP 3 121 277 B1

<211> 27  
 <212> DNA  
 <213> Mus musculus

5       <400> 60  
 ttccaaggtt cacatgttc gctcacg       27

10       <210> 61  
           <211> 445  
           <212> PRT  
           <213> Mus musculus

15       <400> 61  
           Asp Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
           1                   5                   10                   15

20       Ser Arg Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe

25

30

35

40

45

50

55

EP 3 121 277 B1

	20					25					30					
5	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Glu	Lys	Gly	Leu	Glu	Trp	Val
			35					40					45			
10	Ala	Tyr	Ile	Ser	Ser	Gly	Ser	Ser	Thr	Ile	Tyr	Tyr	Ala	Asp	Thr	Val
		50					55					60				
15	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Pro	Lys	Asn	Thr	Leu	Phe
	65					70					75				80	
20	Leu	Gln	Met	Thr	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys
					85					90					95	
25	Ala	Gly	Thr	Gly	Thr	Arg	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr
				100					105					110		
30	Val	Ser	Ala	Ala	Lys	Thr	Thr	Ala	Pro	Ser	Val	Tyr	Pro	Leu	Ala	Pro
			115					120					125			
35	Val	Cys	Gly	Asp	Thr	Thr	Gly	Ser	Ser	Val	Ala	Leu	Gly	Cys	Leu	Val
		130					135					140				
40	Lys	Gly	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Leu	Thr	Trp	Asn	Ser	Gly	Ser
	145					150					155					160
45	Leu	Ser	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Asp	Leu
					165					170					175	
50	Tyr	Thr	Leu	Ser	Ser	Ser	Val	Thr	Val	Thr	Ser	Ser	Thr	Trp	Pro	Ser
				180					185					190		
55	Gln	Ser	Ile	Thr	Cys	Asn	Val	Ala	His	Pro	Ala	Ser	Ser	Thr	Lys	Val
			195					200					205			
60	Asp	Lys	Lys	Ile	Glu	Pro	Arg	Gly	Pro	Thr	Ile	Lys	Pro	Cys	Pro	Pro
		210					215					220				
65	Cys	Lys	Cys	Pro	Ala	Pro	Asn	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Ile
	225					230					235					240
70	Phe	Pro	Pro	Lys	Ile	Lys	Asp	Val	Leu	Met	Ile	Ser	Leu	Ser	Pro	Ile
					245					250					255	
75	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Glu	Asp	Asp	Pro	Asp	Val	Gln
				260					265					270		

# EP 3 121 277 B1

	Ile	Ser	Trp	Phe	Val	Asn	Asn	Val	Glu	Val	His	Thr	Ala	Gln	Thr	Gln	
			275					280					285				
5	Thr	His	Arg	Glu	Asp	Tyr	Asn	Ser	Thr	Leu	Arg	Val	Val	Ser	Ala	Leu	
		290					295					300					
10	Pro	Ile	Gln	His	Gln	Asp	Trp	Met	Ser	Gly	Lys	Glu	Phe	Lys	Cys	Lys	
	305					310					315					320	
	Val	Asn	Asn	Lys	Asp	Leu	Pro	Ala	Pro	Ile	Glu	Arg	Thr	Ile	Ser	Lys	
				325						330					335		
15	Pro	Lys	Gly	Ser	Val	Arg	Ala	Pro	Gln	Val	Tyr	Val	Leu	Pro	Pro	Pro	
				340					345					350			
20	Glu	Glu	Glu	Met	Thr	Lys	Lys	Gln	Val	Thr	Leu	Thr	Cys	Met	Val	Thr	
			355					360					365				
25	Asp	Phe	Met	Pro	Glu	Asp	Ile	Tyr	Val	Glu	Trp	Thr	Asn	Asn	Gly	Lys	
	370						375					380					
30	Thr	Glu	Leu	Asn	Tyr	Lys	Asn	Thr	Glu	Pro	Val	Leu	Asp	Ser	Asp	Gly	
	385					390					395					400	
	Ser	Tyr	Phe	Met	Tyr	Ser	Lys	Leu	Arg	Val	Glu	Lys	Lys	Asn	Trp	Val	
					405					410					415		
35	Glu	Arg	Asn	Ser	Tyr	Ser	Cys	Ser	Val	Val	His	Glu	Gly	Leu	His	Asn	
			420					425						430			
40	His	His	Thr	Thr	Lys	Ser	Phe	Ser	Arg	Thr	Pro	Gly	Lys				
			435					440					445				

<210> 62

<211> 1338

<212> DNA

45 <213> Mus musculus

<400> 62

50

55

EP 3 121 277 B1

	gatgtgcagc tgggtggagtc tggggggaggc ttagtgcagc ctggaggggc ccggaactc	60
	tcctgtgcag cctctggatt cactttcagt agctttggaa tgcactgggt tcgtcaggct	120
5	ccagagaagg ggctggagtg ggtcgcatac attagtagtg gcagtagtac catctactat	180
	gcagacacag tgaagggccg attcaccatc tccagagaca atcccaagaa caccctgttc	240
10	ctgcaaataa ccagtctaag gtctgaggac acggccatgt attactgtgc cgggactggg	300
	acgagagctt actggggcca agggactctg gtcactgtct ctgcagccaa aacaacagcc	360
	ccatcggtct atcccctggc ccctgtgtgt ggagatacaa ctggctcctc ggtggctcta	420
15	ggatgcctgg tcaagggtta tttccctgag ccagtgcact tgacctggaa ctctggatcc	480
	ctgtccagtg gtgtgcacac cttcccagct gtcctgcagt ctgacctcta caccctcagc	540
20	agctcagtgta ctgtaacctc gagcacctgg cccagccagt ccatcacctg caatgtggcc	600
	cacccggcaa gcagcaccaa ggtggacaag aaaattgagc ccagagggcc cacaatcaag	660
	ccctgtcctc catgcaaata cccagcacct aacctcttgg gtggaccatc cgtcttcac	720
25	ttccctccaa agatcaagga tgtactcatg atctccctga gcccatagt cacatgtgtg	780
	gtgggtggatg tgagcgagga tgaccagat gtccagatca gctggtttgt gaacaacgtg	840
	gaagtacaca cagctcagac acaaaccat agagaggatt acaacagtac tctccgggtg	900
30	gtcagtgccc tccccatcca gcaccaggac tggatgagtg gcaaggagtt caaatgcaag	960
	gtcaacaaca aagacctccc agcgcccatc gagagaacca tctcaaaacc caaaggggtca	1020
35	gtaagagctc cacaggtata tgtcttgcct ccaccagaag aagagatgac taagaaacag	1080
	gtcactctga cctgcatggc cacagacttc atgcctgaag acatttacgt ggagtggacc	1140
	aacaacggga aaacagagct aaactacaag aacactgaac cagtcctgga ctctgatggc	1200
40	tcttacttca tgtacagcaa gctgagagtg gaaaagaaga actgggtgga aagaaatagc	1260
	tactcctggt cagtgggtcca cgagggtctg cacaatcacc acacgactaa gagcttctcc	1320
	cggactccgg gtaaatga	1338
45	<210> 63	
	<211> 219	
	<212> PRT	
	<213> Mus musculus	
50	<400> 63	
55		

# EP 3 121 277 B1

	Asp	Val	Leu	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly	
	1				5					10					15		
5	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Ile	Val	His	Ser	
				20					25					30			
	Asn	Gly	Asn	Thr	Tyr	Leu	Glu	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser	
10			35					40					45				
	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
		50					55					60					
15	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75					80	
	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Tyr	Cys	Phe	Gln	Gly	
20					85					90					95		
	Ser	His	Val	Pro	Pro	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	
25				100					105					110			
	Arg	Ala	Asp	Ala	Ala	Pro	Thr	Val	Ser	Ile	Phe	Pro	Pro	Ser	Ser	Glu	
			115					120					125				
30	Gln	Leu	Thr	Ser	Gly	Gly	Ala	Ser	Val	Val	Cys	Phe	Leu	Asn	Asn	Phe	
		130					135					140					
	Tyr	Pro	Lys	Asp	Ile	Asn	Val	Lys	Trp	Lys	Ile	Asp	Gly	Ser	Glu	Arg	
35		145				150					155					160	
	Gln	Asn	Gly	Val	Leu	Asn	Ser	Trp	Thr	Asp	Gln	Asp	Ser	Lys	Asp	Ser	
40					165					170					175		
	Thr	Tyr	Ser	Met	Ser	Ser	Thr	Leu	Thr	Leu	Thr	Lys	Asp	Glu	Tyr	Glu	
				180					185					190			
45	Arg	His	Asn	Ser	Tyr	Thr	Cys	Glu	Ala	Thr	His	Lys	Thr	Ser	Thr	Ser	
			195					200					205				
	Pro	Ile	Val	Lys	Ser	Phe	Asn	Arg	Asn	Glu	Cys						
50				210			215										

<210> 64

<211> 660

55 <212> DNA

<213> Mus musculus

<400> 64

# EP 3 121 277 B1

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60  
atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120  
5 tacctgcaga aaccaggcca gtctcctaag ctctgatct acaaagtttc caaccgattt 180  
tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240  
aggcagagtgg aggctgagga tctgggagtt tattactgct ttcaaggttc acatgttcct 300  
10 ccgacgttcg gtggaggcac caagctggaa atcaaacggg ctgatgctgc accaactgta 360  
tccatcttcc caccatccag tgagcagtta acatctggag gtgcctcagt cgtgtgcttc 420  
ttgaacaact tctaccccaa agacatcaat gtcaagtgga agattgatgg cagtgaacga 480  
15 caaaatggcg tcctgaacag ttggactgat caggacagca aagacagcac ctacagcatg 540  
agcagtaccc tcacgttgac caaggacgag tatgaacgac ataacagcta tacctgtgag 600  
20 gccactcaca agacatcaac ttcacccatt gtcaagagct tcaacaggaa tgagtgttag 660

<210> 65

<211> 115

<212> PRT

25 <213> Mus musculus

<400> 65

30 Asp Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
1 5 10 15  
Ser Arg Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe  
35 20 25 30  
Gly Met His Trp Val Arg Gln Ala Pro Glu Lys Gly Leu Glu Trp Val  
40 35 40 45  
Ala Tyr Ile Ser Ser Gly Ser Ser Thr Ile Tyr Tyr Ala Asp Thr Val  
50 55 60  
45 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Pro Lys Asn Thr Leu Phe  
65 70 75 80  
Leu Gln Met Thr Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys  
50 85 90 95  
Ala Gly Thr Gly Thr Arg Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr  
100 105 110  
55 Val Ser Ala  
115



# EP 3 121 277 B1

<210> 66  
 <211> 345  
 <212> DNA  
 <213> Mus musculus

5

<400> 66

```

gatgtgcagc tgggtggagtc tggggggaggc ttagtgcagc ctggagggtc cgggaaactc      60
10  tcctgtgcag cctctggatt cactttcagt agctttggaa tgcactgggt tcgtcaggct      120
    ccagagaagg ggctggagtg ggtcgcatac attagtagtg gcagtagtac catctactat      180
    gcagacacag tgaagggccg attcaccatc tccagagaca atcccaagaa caccctgttc      240
15  ctgcaaataa ccagtctaag gtctgaggac acggccatgt attactgtgc cgggactggg      300
    acgagagctt actggggcca agggactctg gtcactgtct ctgca                      345
  
```

<210> 67  
 <211> 112  
 <212> PRT  
 <213> Mus musculus

25 <400> 67

```

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1      5      10      15
30  Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
    20      25      30
35  Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
    35      40      45
40  Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
    50      55      60
45  Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
    65      70      75      80
50  Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly
    85      90      95
55  Ser His Val Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
    100     105     110
  
```

<210> 68  
 <211> 336  
 <212> DNA  
 <213> Mus musculus

55

<400> 68

# EP 3 121 277 B1

	gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc	60
	atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg	120
5	tacctgcaga aaccaggcca gtctcctaag ctctgatct acaaagtttc caaccgattt	180
	tctggggtcc cagacaggtt cagtggcagt ggatcaggga cagatttcac actcaagatc	240
	agcagagtgg aggctgagga tctgggagtt tattactgct ttcaaggttc acatgttcct	300
10	ccgacgttcg gtggaggcac caagctggaa atcaaa	336

	<210> 69
	<211> 5
15	<212> PRT
	<213> Mus musculus

	<400> 69
20	
	Ser Phe Gly Met His
	1 5

	<210> 70
	<211> 15
25	<212> DNA
	<213> Mus musculus

	<400> 70
	agcttggaa tgcac 15

30	
	<210> 71
	<211> 10
	<212> PRT
	<213> Mus musculus

35	<400> 71
----	----------

	Tyr Ile Ser Ser Gly Ser Ser Thr Ile Tyr
	1 5 10

40	
	<210> 72
	<211> 30
	<212> DNA
	<213> Mus musculus

45	
	<400> 72
	tacattagta gtggcagtag taccatctac 30

50	
	<210> 73
	<211> 6
	<212> PRT
	<213> Mus musculus

55	<400> 73
----	----------

	Thr Gly Thr Arg Ala Tyr
	1 5

# EP 3 121 277 B1

<210> 74  
 <211> 18  
 <212> DNA  
 <213> Mus musculus  
 5  
 <400> 74  
 actgggacga gagcttac 18  
 10  
 <210> 75  
 <211> 16  
 <212> PRT  
 <213> Mus musculus  
 15  
 <400> 75  
 Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu  
 1 5 10 15  
 20  
 <210> 76  
 <211> 48  
 <212> DNA  
 <213> Mus musculus  
 25  
 <400> 76  
 agatctagtc agagcattgt acatagtaat ggaaacacct atttagaa 48  
 30  
 <210> 77  
 <211> 7  
 <212> PRT  
 <213> Mus musculus  
 <400> 77  
 35  
 Lys Val Ser Asn Arg Phe Ser  
 1 5  
 40  
 <210> 78  
 <211> 21  
 <212> DNA  
 <213> Mus musculus  
 <400> 78  
 aaagttcca accgatttc t 21  
 45  
 <210> 79  
 <211> 9  
 <212> PRT  
 <213> Mus musculus  
 50  
 <400> 79  
 Phe Gln Gly Ser His Val Pro Pro Thr  
 1 5  
 55  
 <210> 80  
 <211> 27  
 <212> DNA  
 <213> Mus musculus

# EP 3 121 277 B1

<400> 80

tttcaagggtt cacatgttcc tccgacg 27

<210> 81

5 <211> 442

<212> PRT

<213> Mus musculus

<400> 81

10

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala  
1 5 10 15

15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr  
20 25 30

20

Trp Met His Trp Val Lys Leu Arg Pro Gly Gln Gly Phe Glu Trp Ile  
35 40 45

Gly Glu Ile Asn Pro Arg Asn Gly Gly Thr Asn Asn Asn Glu Asn Phe

25

30

35

40

45

50

55

EP 3 121 277 B1

	50		55		60											
5	Lys 65	Arg	Lys	Ala	Thr	Leu 70	Thr	Val	Asp	Lys	Ser 75	Ser	Ser	Thr	Ala	Tyr 80
10	Met	Gln	Leu	Ser	Ser 85	Leu	Thr	Ser	Glu	Asp 90	Ser	Ala	Val	Tyr	Tyr 95	Cys
15	Thr	Arg	Asp	Gly 100	Asn	Tyr	Asp	Pro	Phe 105	Ala	Tyr	Trp	Gly	Gln 110	Gly	Thr
20	Leu	Val	Thr	Val	Ser	Ala	Ala	Lys 120	Thr	Thr	Pro	Pro	Ser 125	Val	Tyr	Pro
25	Leu	Ala	Pro	Gly	Ser	Ala	Ala	Gln	Thr	Asn	Ser	Met 140	Val	Thr	Leu	Gly
30	Cys 145	Leu	Val	Lys	Gly	Tyr 150	Phe	Pro	Glu	Pro	Val 155	Thr	Val	Thr	Trp	Asn 160
35	Ser	Gly	Ser	Leu	Ser 165	Ser	Gly	Val	His	Thr 170	Phe	Pro	Ala	Val	Leu	Gln 175
40	Ser	Asp	Leu	Tyr 180	Thr	Leu	Ser	Ser	Ser 185	Val	Thr	Val	Pro	Ser 190	Ser	Thr
45	Trp	Pro	Ser	Glu	Thr	Val	Thr	Cys 200	Asn	Val	Ala	His	Pro 205	Ala	Ser	Ser
50	Thr	Lys 210	Val	Asp	Lys	Lys	Ile 215	Val	Pro	Arg	Asp	Cys 220	Gly	Cys	Lys	Pro
55	Cys 225	Ile	Cys	Thr	Val	Pro 230	Glu	Val	Ser	Ser	Val 235	Phe	Ile	Phe	Pro	Pro 240
60	Lys	Pro	Lys	Asp	Val 245	Leu	Thr	Ile	Thr	Leu 250	Thr	Pro	Lys	Val	Thr 255	Cys
65	Val	Val	Val	Asp	Ile 260	Ser	Lys	Asp	Asp 265	Pro	Glu	Val	Gln	Phe	Ser	Trp
70	Phe	Val	Asp	Asp	Val 275	Glu	Val	His 280	Thr	Ala	Gln	Thr	Gln 285	Pro	Arg	Glu
75	Glu	Gln	Phe	Asn	Ser	Thr	Phe	Arg	Ser	Val	Ser	Glu	Leu	Pro	Ile	Met
80																

# EP 3 121 277 B1

	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Phe	Lys	Cys	Arg	Val	Asn	Ser	
	305					310					315					320	
5	Ala	Ala	Phe	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	
					325					330					335		
10	Arg	Pro	Lys	Ala	Pro	Gln	Val	Tyr	Thr	Ile	Pro	Pro	Pro	Lys	Glu	Gln	
				340					345					350			
15	Met	Ala	Lys	Asp	Lys	Val	Ser	Leu	Thr	Cys	Met	Ile	Thr	Asp	Phe	Phe	
			355					360					365				
20	Pro	Glu	Asp	Ile	Thr	Val	Glu	Trp	Gln	Trp	Asn	Gly	Gln	Pro	Ala	Glu	
		370					375					380					
25	Asn	Tyr	Lys	Asn	Thr	Gln	Pro	Ile	Met	Asp	Thr	Asp	Gly	Ser	Tyr	Phe	
	385					390					395					400	
30	Val	Tyr	Ser	Lys	Leu	Asn	Val	Gln	Lys	Ser	Asn	Trp	Glu	Ala	Gly	Asn	
					405					410					415		
35	Thr	Phe	Thr	Cys	Ser	Val	Leu	His	Glu	Gly	Leu	Leu	Asn	Leu	His	Thr	
				420					425					430			
40	Glu	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys							
			435					440									

<210> 82  
 <211> 1329  
 <212> DNA  
 <213> Mus musculus

<400> 82

EP 3 121 277 B1

	caggtccaac tccagcagcc tggggctgaa ctggtgaagc ctggggcttc agtgaagttg	60
	tcctgcaagg cttctggcta caccttcacc agctactgga tgcactgggt gaagctgagg	120
5	cctggacaag gctttgagtg gattggagag attaatccta gaaatggtgg tactaacaac	180
	aatgagaact tcaagagaaa ggccacactg actgtagaca aatcctccag cacagcctac	240
	atgcaactca gcagcctgac atctgaggac tctgcggtct attactgtac aagagatggg	300
10	aactacgacc cctttgctta ctggggccaa gggactctgg tcaactgtctc tgcagccaaa	360
	acgacacccc catctgtcta tccactggcc cctggatctg ctgcccacaa taactccatg	420
15	gtgaccctgg gatgcctggg caagggctat ttccctgagc cagtgcaggt gacctggaac	480
	tctggatccc tgtccagcgg tgtgcacacc ttcccagctg tcctgcagtc tgacctctac	540
	actctgagca gctcagtgac tgtcccctcc agcacctggc ccagcgagac cgtcacctgc	600
20	aacgttgccc acccggccag cagcaccaag gtggacaaga aaattgtgcc cagggattgt	660
	ggttgtaagc cttgcatatg tacagtccca gaagtatcat ctgtcttcat cttcccccca	720
25	aagcccaagg atgtgctcac cattactctg actcctaagg tcacgtgtgt tgtggtagac	780
	atcagcaagg atgatcccga ggtccagttc agctggtttg tagatgatgt ggagggtcac	840
	acagctcaga cgcaaccccc ggaggagcag ttcaacagca ctttccgctc agtcagtga	900
30	cttcccatca tgcaccagga ctggctcaat ggcaaggagt tcaaatgcag ggtcaacagt	960
	gcagctttcc ctgcccccat cgagaaaacc atctccaaaa ccaaaggcag accgaaggct	1020
	ccacaggtgt acaccattcc acctcccaag gagcagatgg ccaaggataa agtcagtctg	1080
35	acctgcatga taacagactt cttccctgaa gacattactg tggagtggca gtggaatggg	1140
	cagccagcgg agaactacaa gaacactcag cccatcatgg acacagatgg ctcttacttc	1200
	gtttacagca agctcaatgt gcagaagagc aactgggagg caggaaatac tttcacctgc	1260
40	tctgtgttac atgagggcct gctcaacctc catactgaga agagcctctc cctctctcct	1320
	ggtaaatga	1329
45	<210> 83 <211> 220 <212> PRT <213> Mus musculus	
50	<400> 83	
55		

# EP 3 121 277 B1

	Asp	Val	Val	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly
	1				5					10					15	
5	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser
				20					25					30		
10	Asn	Gly	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
			35					40					45			
15	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
		50					55					60				
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
	65					70					75					80
25	Thr	His	Val	Pro	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile
				100					105					110		
30	Lys	Arg	Ala	Asp	Ala	Ala	Pro	Thr	Val	Ser	Ile	Phe	Pro	Pro	Ser	Ser
			115					120					125			
35	Glu	Gln	Leu	Thr	Ser	Gly	Gly	Ala	Ser	Val	Val	Cys	Phe	Leu	Asn	Asn
		130					135					140				
40	Phe	Tyr	Pro	Lys	Asp	Ile	Asn	Val	Lys	Trp	Lys	Ile	Asp	Gly	Ser	Glu
	145					150					155					160
45	Arg	Gln	Asn	Gly	Val	Leu	Asn	Ser	Trp	Thr	Asp	Gln	Asp	Ser	Lys	Asp
					165					170					175	
50	Ser	Thr	Tyr	Ser	Met	Ser	Ser	Thr	Leu	Thr	Leu	Thr	Lys	Asp	Glu	Tyr
				180					185					190		
55	Glu	Arg	His	Asn	Ser	Tyr	Thr	Cys	Glu	Ala	Thr	His	Lys	Thr	Ser	Thr
			195					200					205			
60	Ser	Pro	Ile	Val	Lys	Ser	Phe	Asn	Arg	Asn	Glu	Cys				
		210					215					220				

<210> 84

<211> 663

<212> DNA

<213> Mus musculus

<400> 84



# EP 3 121 277 B1

	gatgttgtga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc	60
	atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg	120
5	tacctgcaga agccaggcca gtctccaaag ctccctgatct acaaagtttc caaccgattt	180
	tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc	240
	agcagagtgg aggctgagga tctgggagtt tattttctgct ctcaaagtac acatgttcct	300
10	ccgtacacgt tcggaggggg gaccaagctg gaaataaaac gggctgatgc tgcaccaact	360
	gtatccatct tcccaccatc cagtgagcag ttaacatctg gaggtgcctc agtcgtgtgc	420
	ttcttgaaca acttctaccc caaagacatc aatgtcaagt ggaagattga tggcagtga	480
15	cgacaaaatg gcgtcctgaa cagttggact gatcaggaca gcaaagacag cacctacagc	540
	atgagcagta ccctcacgtt gaccaaggac gagtatgaac gacataacag ctatacctgt	600
20	gaggccactc acaagacatc aacttcaccc attgtcaaga gcttcaacag gaatgagtgt	660
	tag	663

<210> 85  
 <211> 118  
 <212> PRT  
 <213> Mus musculus

<400> 85

# EP 3 121 277 B1

	Gln	Val	Gln	Leu	Gln	Gln	Pro	Gly	Ala	Glu	Leu	Val	Lys	Pro	Gly	Ala	
	1				5					10					15		
5	Ser	Val	Lys	Leu	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	
				20					25					30			
10	Trp	Met	His	Trp	Val	Lys	Leu	Arg	Pro	Gly	Gln	Gly	Phe	Glu	Trp	Ile	
			35					40					45				
15	Gly	Glu	Ile	Asn	Pro	Arg	Asn	Gly	Gly	Thr	Asn	Asn	Asn	Glu	Asn	Phe	
		50					55					60					
20	Lys	Arg	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr	
	65					70					75					80	
25	Met	Gln	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		
30	Thr	Arg	Asp	Gly	Asn	Tyr	Asp	Pro	Phe	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
35	Leu	Val	Thr	Val	Ser	Ala											
				115													
40	<210> 86																
	<211> 354																
	<212> DNA																
	<213> Mus musculus																
45	<400> 86																
50	caggtccaac	tccagcagcc	tggggctgaa	ctggtgaagc	ctggggcttc	agtgaagttg											60
55	tcctgcaagg	cttctggcta	caccttcacc	agctactgga	tgactgggt	gaagctgagg											120
	cctggacaag	gctttgagtg	gattggagag	attaatccta	gaaatggtgg	tactaacaac											180
	aatgagaact	tcaagagaaa	ggccacactg	actgtagaca	aatcctccag	cacagcctac											240
	atgcaactca	gcagcctgac	atctgaggac	tctgcggtct	attactgtac	aagagatggt											300
	aactacgacc	cctttgctta	ctggggccaa	gggactctgg	tcactgtctc	tgca											354
60	<210> 87																
	<211> 113																
	<212> PRT																
	<213> Mus musculus																
65	<400> 87																
70	Asp	Val	Val	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly	
	1				5					10					15		

# EP 3 121 277 B1

	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser	
				20				25						30			
5	Asn	Gly	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser	
			35					40					45				
10	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
		50					55					60					
15	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75					80	
20	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Phe	Cys	Ser	Gln	Ser	
					85					90					95		
25	Thr	His	Val	Pro	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	
				100					105					110			
30	Lys																
	<210>	88															
	<211>	339															
	<212>	DNA															
	<213>	Mus musculus															
35	<400>	88															
	gatgtt	gtga	tgaccc	aaac	tccact	tctcc	ctgcct	gtca	gtctt	ggaga	tcaagc	cctcc				60	
40	atctct	ttgca	gatcta	gtca	gagcct	tgtga	cacagta	aatg	gaaacac	ccta	tttacatt	gg				120	
	tacctg	caga	agccagg	ccta	gtctcc	aaag	ctcctga	tct	acaaagt	tttc	caaccg	attt				180	
45	tctggg	gtcc	cagacag	gtt	cagtgg	cagt	ggatcag	gga	cagatt	tcac	actcaag	atc				240	
50	agcaga	gtgg	aggctga	gga	tctggg	agtt	tatttt	ctgct	ctcaaag	tac	acatgtt	cct				300	
	ccgtac	acgt	tcggag	gggg	gaccaag	ctg	gaaata	aaaa								339	
55	<210>	89															
	<211>	5															
	<212>	PRT															
	<213>	Mus musculus															
60	<400>	89															
	Ser	Tyr	Trp	Met	His												
	1				5												
65	<210>	90															
	<211>	15															
	<212>	DNA															
	<213>	Mus musculus															

# EP 3 121 277 B1

<400> 90  
 agctactgga tgcac 15  
  
 <210> 91  
 <211> 10  
 <212> PRT  
 <213> Mus musculus  
  
 <400> 91  
  
 Glu Ile Asn Pro Arg Asn Gly Gly Thr Asn  
 1 5 10  
  
 <210> 92  
 <211> 30  
 <212> DNA  
 <213> Mus musculus  
  
 <400> 92  
 gagattaatc ctagaaatgg tggactaac 30  
  
 <210> 93  
 <211> 9  
 <212> PRT  
 <213> Mus musculus  
  
 <400> 93  
  
 Asp Gly Asn Tyr Asp Pro Phe Ala Tyr  
 1 5  
  
 <210> 94  
 <211> 27  
 <212> DNA  
 <213> Mus musculus  
  
 <400> 94  
 gatgtaact acgaccctt tgcttac 27  
  
 <210> 95  
 <211> 16  
 <212> PRT  
 <213> Mus musculus  
  
 <400> 95  
  
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His  
 1 5 10 15  
  
 <210> 96  
 <211> 48  
 <212> DNA  
 <213> Mus musculus  
  
 <400> 96  
 agatctagtc agagccttgt acacagtaat ggaaacacct attacat 48  
  
 <210> 97

# EP 3 121 277 B1

<211> 7  
<212> PRT  
<213> Mus musculus

5 <400> 97

Lys Val Ser Asn Arg Phe Ser  
1 5

10 <210> 98  
<211> 21  
<212> DNA  
<213> Mus musculus

15 <400> 98  
aaagttcca accgatttc t 21

<210> 99  
<211> 10  
20 <212> PRT  
<213> Mus musculus

<400> 99

25 Ser Gln Ser Thr His Val Pro Pro Tyr Thr  
1 5 10

<210> 100  
<211> 30  
30 <212> DNA  
<213> Mus musculus

<400> 100  
tctcaaagta cacatgttc tccgtacacg 30

35 <210> 101  
<211> 453  
<212> PRT  
<213> Mus musculus

40 <400> 101

45 Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln  
1 5 10 15

Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser  
20 25 30

50 Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly Glu Gly Leu Glu  
35 40 45

55 Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser  
50 55 60

EP 3 121 277 B1

	Leu	Lys	Ser	Arg	Leu	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Arg	Asn	Gln	Val	
	65					70					75					80	
5	Phe	Leu	Lys	Ile	Thr	Ser	Val	Asp	Thr	Ala	Asp	Thr	Ala	Thr	Tyr	Tyr	
					85					90					95		
10	Cys	Gly	Arg	Tyr	Arg	Tyr	Gly	Phe	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	
				100					105					110			
15	Val	Thr	Val	Ser	Ala	Ala	Lys	Thr	Thr	Pro	Pro	Ser	Val	Tyr	Pro	Leu	
			115					120					125				
20	Ala	Pro	Gly	Cys	Gly	Asp	Thr	Thr	Gly	Ser	Ser	Val	Thr	Leu	Gly	Cys	
		130					135					140					
25	Leu	Val	Lys	Gly	Tyr	Phe	Pro	Glu	Ser	Val	Thr	Val	Thr	Trp	Asn	Ser	
	145					150					155					160	
30	Gly	Ser	Leu	Ser	Ser	Ser	Val	His	Thr	Phe	Pro	Ala	Leu	Leu	Gln	Ser	
				165						170					175		
35	Gly	Leu	Tyr	Thr	Met	Ser	Ser	Ser	Val	Thr	Val	Pro	Ser	Ser	Thr	Trp	
				180					185					190			
40	Pro	Ser	Gln	Thr	Val	Thr	Cys	Ser	Val	Ala	His	Pro	Ala	Ser	Ser	Thr	
			195					200					205				
45	Thr	Val	Asp	Lys	Lys	Leu	Glu	Pro	Ser	Gly	Pro	Ile	Ser	Thr	Ile	Asn	
		210					215					220					
50	Pro	Cys	Pro	Pro	Cys	Lys	Glu	Cys	His	Lys	Cys	Pro	Ala	Pro	Asn	Leu	
	225					230					235					240	
55	Glu	Gly	Gly	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Asn	Ile	Lys	Asp	Val	
					245					250					255		
60	Leu	Met	Ile	Ser	Leu	Thr	Pro	Lys	Val	Thr	Cys	Val	Val	Val	Asp	Val	
				260					265					270			
65	Ser	Glu	Asp	Asp	Pro	Asp	Val	Gln	Ile	Ser	Trp	Phe	Val	Asn	Asn	Val	
			275					280					285				
70	Glu	Val	His	Thr	Ala	Gln	Thr	Gln	Thr	His	Arg	Glu	Asp	Tyr	Asn	Ser	
		290					295					300					
75	Thr	Ile	Arg	Val	Val	Ser	Thr	Leu	Pro	Ile	Gln	His	Gln	Asp	Trp	Met	
	305					310					315					320	

# EP 3 121 277 B1

	Ser Gly Lys Glu Phe Lys Cys Lys Val Asn Asn Lys Asp Leu Pro Ser	
	325	330 335
5	Pro Ile Glu Arg Thr Ile Ser Lys Ile Lys Gly Leu Val Arg Ala Pro	
	340	345 350
10	Gln Val Tyr Ile Leu Pro Pro Pro Ala Glu Gln Leu Ser Arg Lys Asp	
	355	360 365
15	Val Ser Leu Thr Cys Leu Val Val Gly Phe Asn Pro Gly Asp Ile Ser	
	370	375 380
20	Val Glu Trp Thr Ser Asn Gly His Thr Glu Glu Asn Tyr Lys Asp Thr	
	385	390 395 400
25	Ala Pro Val Leu Asp Ser Asp Gly Ser Tyr Phe Ile Tyr Ser Lys Leu	
	405	410 415
30	Asn Met Lys Thr Ser Lys Trp Glu Lys Thr Asp Ser Phe Ser Cys Asn	
	420	425 430
35	Val Arg His Glu Gly Leu Lys Asn Tyr Tyr Leu Lys Lys Thr Ile Ser	
	435	440 445
40	Arg Ser Pro Gly Lys	
	450	
45	<210> 102	
	<211> 1362	
	<212> DNA	
	<213> Mus musculus	
50	<400> 102	
55	cagggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg	60
	acttgtttctt tctctgggtt ttactgagc acttctggta tgggtgtgag ctggattcgt	120
	cagccttcag gagaggggtct ggagtggctg gcacacattt actgggatga tgacaagcgc	180
	tataacccat ccctgaagag ccggctcaca atctccaagg atacctccag aaaccaggta	240
	ttcctcaaga tcaccagtgt ggacactgca gatactgcca catactactg tggtcgatat	300
	aggtacggct ttgcttactg gggccaaggg actctgggtca ctgtctctgc agccaaaaca	360
	acacccccat cagtctatcc actggcccct ggggtgtggag atacaactgg ttcctccgtg	420
	actctgggat gcctgggtcaa gggctacttc cctgagtcag tgactgtgac ttggaactct	480
	ggatccctgt ccagcagtgt gcacaccttc ccagctctcc tgcagtctgg actctacact	540
	atgagcagct cagtgactgt cccctccagc acctggccaa gtcagaccgt cacctgcagc	600

EP 3 121 277 B1

gttgctcacc cagccagcag caccacggtg gacaaaaaac ttgagcccag cgggcccatt 660  
tcaacaatca acccctgtcc tccatgcaag gagtgtcaca aatgcccagc tcctaacctc 720  
5 gaggggtggac catccgtctt catcttccct ccaaatatca aggatgtact catgatctcc 780  
ctgacaccca aggtcacgtg tgtggtggtg gatgtgagcg aggatgaccc agacgtccag 840  
atcagctggt ttgtgaacaa cgtggaagta cacacagctc agacacaaac ccatagagag 900  
10 gattacaaca gtactatccg ggtggtcagc accctcccca tccagcacca ggactggatg 960  
agtggcaagg agttcaaag caaggtcaac aacaaagacc tcccatcacc catcgagaga 1020  
accatctcaa aaattaaagg gctagtcaga gctccacaag tatacatctt gccgccacca 1080  
15 gcagagcagt tgtccaggaa agatgtcagt ctcaacttgc ttggtcgtggg cttcaaccct 1140  
ggagacatca gtgtggagtg gaccagcaat gggcatacag aggagaacta caaggacacc 1200  
20 gcaccagtcc tggactctga cggttcttac ttcatatata gcaagctcaa tatgaaaaca 1260  
agcaagtggg agaaaacaga ttccttctca tgcaacgtga gacacgaggg tctgaaaaat 1320  
tactacctga agaagaccat ctcccgtct cggggtaaat ga 1362

<210> 103  
<211> 219  
<212> PRT  
<213> Mus musculus

<400> 103

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly  
1 5 10 15  
35  
Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser  
20 25 30  
40  
Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser  
35 40 45  
45  
Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro  
50 55 60  
50  
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile  
65 70 75 80  
Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly  
85 90 95  
55  
Ser His Val Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys  
100 105 110



**EP 3 121 277 B1**

[illegible]

# EP 3 121 277 B1

<400> 105

5	Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln	1 5 10 15
10	Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser	20 25 30
15	Gly Met Gly Val Ser Trp Ile Arg Gln Pro Ser Gly Glu Gly Leu Glu	35 40 45
20	Trp Leu Ala His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ser	50 55 60
25	Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Arg Asn Gln Val	65 70 75 80
30	Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr	85 90 95
35	Cys Gly Arg Tyr Arg Tyr Gly Phe Ala Tyr Trp Gly Gln Gly Thr Leu	100 105 110
40	Val Thr Val Ser	115

<210> 106

<211> 348

<212> DNA

<213> Mus musculus

<400> 106

40	caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg	60
45	acttggttctt tctctgggtt ttactgagc acttctggta tgggtgtgag ctggattcgt	120
50	cagccttcag gagaggggtct ggagtggctg gcacacattt actgggatga tgacaagcgc	180
55	tataacccat ccctgaagag ccggctcaca atctccaagg atacctccag aaaccaggta	240
60	ttcctcaaga tcaccagtgt ggacactgca gatactgccata cactactactg tggtcgatat	300
65	aggtacggct ttgcttactg gggccaaggg actctgggtca ctgtctct	348

<210> 107

<211> 112

<212> PRT

<213> Mus musculus

<400> 107

# EP 3 121 277 B1

	Asp	Val	Leu	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly	
	1				5					10					15		
5	Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Ile	Val	His	Ser	
				20					25					30			
10	Asn	Gly	Asn	Thr	Tyr	Leu	Glu	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser	
			35					40					45				
15	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
		50					55					60					
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70				75					80		
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Tyr	Cys	Phe	Gln	Gly	
					85					90					95		
30	Ser	His	Val	Pro	Leu	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Glu	Leu	Lys	
				100					105					110			
35	<210> 108																
	<211> 336																
	<212> DNA																
	<213> Mus musculus																
40	<400> 108																
	gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60																
45	atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120																
	tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180																
50	tctgggggtcc cagacaggtt cagtggcagt ggatcagga cagatttcac actcaagatc 240																
	agcagagtgg aggctgagga tctgggagtt tattactgct ttcaaggttc acatgttccg 300																
	ctcacgttcg gtgctgggac caagctggag ctgaaa 336																
55	<210> 109																
	<211> 7																
	<212> PRT																
	<213> Mus musculus																
60	<400> 109																
	Thr Ser Gly Met Gly Val Ser																
	1									5							
65	<210> 110																
	<211> 21																
	<212> DNA																
	<213> Mus musculus																

# EP 3 121 277 B1

<400> 110  
 acttctggta tgggtgtgag c 21  
  
 5 <210> 111  
 <211> 9  
 <212> PRT  
 <213> Mus musculus  
  
 10 <400> 111  
  
 His Ile Tyr Trp Asp Asp Asp Lys Arg  
 1 5  
  
 15 <210> 112  
 <211> 27  
 <212> DNA  
 <213> Mus musculus  
  
 20 <400> 112  
 cacatttact gggatgatga caagcgc 27  
  
 25 <210> 113  
 <211> 7  
 <212> PRT  
 <213> Mus musculus  
  
 30 <400> 113  
  
 Tyr Arg Tyr Gly Phe Ala Tyr  
 1 5  
  
 35 <210> 114  
 <211> 21  
 <212> DNA  
 <213> Mus musculus  
  
 40 <400> 114  
 tataggtacg gcttgctta c 21  
  
 45 <210> 115  
 <211> 16  
 <212> PRT  
 <213> Mus musculus  
  
 50 <400> 115  
  
 Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu  
 1 5 10 15  
  
 55 <210> 116  
 <211> 48  
 <212> DNA  
 <213> Mus musculus  
  
 60 <400> 116  
 agatctagtc agagcattgt acatagtaat ggaaacacct atttagaa 48  
  
 <210> 117

<211> 7  
 <212> PRT  
 <213> Mus musculus

5 <400> 117

Lys Val Ser Asn Arg Phe Ser  
 1 5

10 <210> 118  
 <211> 21  
 <212> DNA  
 <213> Mus musculus

15 <400> 118  
 aaagttcca accgatttc t 21

<210> 119  
 <211> 9  
 20 <212> PRT  
 <213> Mus musculus

<400> 119

25 Phe Gln Gly Ser His Val Pro Leu Thr  
 1 5

<210> 120  
 <211> 27  
 30 <212> DNA  
 <213> Mus musculus

<400> 120  
 35 ttccaaggtt cacatgttcc gctcacg 27

## Claims

1. An antibody that binds to an A $\beta$  oligomer but not to an A $\beta$  monomer,  
 40 wherein the antibody is a monoclonal antibody and selected from the group consisting of:
  - (1) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 69 as CDR1, the amino acid sequence of SEQ ID NO: 71 as CDR2, and the amino acid sequence of SEQ ID NO: 73 as CDR3 and an L chain having the amino acid sequence of SEQ ID NO: 75 as CDR1, the amino acid sequence of SEQ ID NO: 77 as CDR2, and the amino acid sequence of SEQ ID NO: 79 as CDR3;  
 45
  - (2) an antibody that comprises an H chain having the amino acid sequence of SEQ ID NO: 65 as VH and an L chain having the amino acid sequence of SEQ ID NO: 67 as VL; and
  - (3) an antibody comprising an H chain having the amino acid sequence of SEQ ID NO: 61 and an L chain having the amino acid sequence of SEQ ID NO: 63.  
 50
2. The antibody of claim 1, wherein the antibody is a chimeric antibody or a humanized antibody.
3. A composition comprising the antibody of claim 1 or 2 and a pharmaceutically acceptable carrier.
4. An agent for use in treating cognitive impairment, which comprises the antibody of claim 1 or 2 or the composition  
 55 of claim 3 as an active ingredient.
5. A therapeutic agent for use in treating Alzheimer's disease, which comprises the antibody of claim 1 or 2 or the

composition of claim 3 as an active ingredient.

6. An agent for use in suppressing the progression of Alzheimer's disease, which comprises the antibody of claim 1 or 2 or the composition of claim 3 as an active ingredient.

7. An agent for use in suppressing senile plaque formation, which comprises the antibody of claim 1 or 2 or the composition of claim 3 as an active ingredient.

8. An agent for use in suppressing A $\beta$  accumulation, which comprises the antibody of claim 1 or 2 or the composition of claim 3 as an active ingredient.

9. An agent for use in neutralizing neurotoxicity, which comprises the antibody of claim 1 or 2 or the composition of claim 3 as an active ingredient.

10. An agent for use in inhibiting A $\beta$  amyloid fibril formation, which comprises the antibody of claim 1 or 2 or the composition of claim 3 as an active ingredient.

11. An agent for use in neutralizing synaptic toxicity, which comprises the antibody of claim 1 or 2 or the composition of claim 3 as an active ingredient.

12. A method for detecting an A $\beta$  oligomer, which comprises the step of detecting an A $\beta$  oligomer contained in a sample collected from a subject using the antibody of claim 1 or 2.

13. A method of diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises the step of (a0) below:

(a0) using the antibody of claim 1 or 2 to detect an A $\beta$  oligomer in a sample collected from a subject;

or

the steps of (a1) to (c1) below:

(a1) contacting a sample collected from a subject with the antibody of claim 1 or 2;

(b1) measuring the amount of A $\beta$  oligomer in the sample; and

(c1) determining that the subject is a possible Alzheimer's disease patient, when the amount measured in step (b1) is higher than that of a healthy individual;

or

the steps of (a2) to (c2) below:

(a2) contacting a sample collected from a subject with the antibody of claim 1 or 2 and an antibody that binds to an A $\beta$  monomer;

(b2) measuring the ratio of A $\beta$  oligomer to A $\beta$  monomer in the sample; and

(c2) determining that the subject is a possible Alzheimer's disease patient, when the ratio measured in step (b2) is higher than that of a healthy individual.

14. The method of claim 12 or 13, wherein the sample is blood or cerebrospinal fluid.

15. A pharmaceutical agent for use in diagnosing whether or not a subject is a possible Alzheimer's disease patient, which comprises the antibody of claim 1 or 2.

## Patentansprüche

1. Antikörper, der an ein A $\beta$ -Oligomer bindet, aber nicht an einen A $\beta$ -Monomer, wobei der Antikörper ein monoklonaler Antikörper ist und ausgewählt ist aus der Gruppe bestehend aus:

(1) ein Antikörper, der eine H-Kette, die die Aminosäuresequenz SEQ ID Nr: 69 als CDR1, die Aminosäuresequenz von SEQ ID Nr: 71 als CDR2 und die Aminosäuresequenz von SEQ ID Nr: 73 als CDR3 aufweist, und

eine L-Kette, die die Aminosäuresequenz von SEQ ID Nr: 75 als CDR1, die Aminosäuresequenz von SEQ ID Nr: 77 als CDR2 und die Aminosäuresequenz von SEQ ID Nr: 79 als CDR3 aufweist, umfasst;

(2) ein Antikörper, der eine H-Kette, die die Aminosäuresequenz von SEQ ID Nr: 65 als VH aufweist und eine L-Kette, die die Aminosäuresequenz von SEQ ID Nr: 67 als VL aufweist, umfasst; und

(3) ein Antikörper, der eine H-Kette, die die Aminosäuresequenz von SEQ ID Nr: 61 aufweist, und eine L-Kette, die die Aminosäuresequenz von SEQ ID Nr: 63 aufweist, umfasst.

2. Antikörper nach Anspruch 1, wobei der Antikörper ein chimärer Antikörper oder ein humanisierter Antikörper ist.

3. Zusammensetzung umfassend den Antikörper nach Anspruch 1 oder 2, und einen pharmazeutisch annehmbaren Träger.

4. Mittel zur Verwendung bei der Behandlung von kognitiver Beeinträchtigung, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

5. Therapeutisches Mittel zur Verwendung bei der Behandlung der Alzheimer-Erkrankung, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

6. Mittel zur Verwendung beim Unterdrücken des Fortschreitens von Alzheimer-Erkrankung, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

7. Mittel zur Verwendung beim Unterdrücken der Bildung seniler Plaques, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

8. Mittel zur Verwendung beim Unterdrücken der A $\beta$ -Akkumulation, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

9. Mittel zur Verwendung beim Neutralisieren der Neurotoxizität, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

10. Mittel zur Verwendung beim Inhibieren der Bildung von A $\beta$ -Amyloid Fibrillen, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

11. Mittel zur Verwendung beim Neutralisieren der synaptischen Toxizität, das den Antikörper nach Anspruch 1 oder 2 oder die Zusammensetzung nach Anspruch 3 als einen aktiven Wirkstoff umfasst.

12. Verfahren zum Detektieren eines A $\beta$ -Oligomers, das den Schritt des Detektierens eines A $\beta$ -Oligomers umfasst, das in einer Probe enthalten ist, die einem Subjekt entnommen wurde, unter Verwendung des Antikörpers nach Anspruch 1 oder 2.

13. Verfahren zum Diagnostizieren ob oder ob nicht ein Subjekt ein möglicher Alzheimer-Erkrankungs-Patient ist, welches umfasst den Schritt (a0) untenstehend:

(a0) Verwenden des Antikörpers nach Anspruch 1 oder 2, um ein A $\beta$ -Oligomer in einer Probe zu detektieren, die einem Subjekt entnommen wurde; oder

die Schritte von (a1) bis (c1) untenstehend:

(a1) Inkontaktbringen einer Probe, die einem Subjekt entnommen wurde, mit dem Antikörper nach Anspruch 1 oder 2;

(b1) Messen der Menge an A $\beta$ -Oligomer in der Probe; und

(c1) Bestimmen, dass das Subjekt ein möglicher Alzheimer-Erkrankungs-Patient ist, falls die Menge, die in Schritt (b1) gemessen wird, höher ist als diejenige eines gesunden Individuums; oder

die Schritte von (a2) bis (c2) untenstehend:

(a2) Inkontaktbringen der Probe, die einem Subjekt entnommen wurde, mit dem Antikörper nach Anspruch 1

oder 2, und einem Antikörper, der an ein A $\beta$ -Monomer bindet;  
 (b2) Messen des Verhältnisses von A $\beta$ -Oligomer zu A $\beta$ -Monomer in der Probe; und  
 (c2) Bestimmen, dass das Subjekt ein möglicher Alzheimer-Erkrankungs-Patient ist, falls das Verhältnis, das  
 in Schritt (b2) gemessen wird, höher ist als dasjenige eines gesunden Individuums.

14. Verfahren nach Anspruch 12 oder 13, wobei die Probe Blut oder Rückenmarkflüssigkeit ist.

15. Pharmazeutisches Mittel zur Verwendung bei der Diagnose ob oder ob nicht ein Subjekt ein möglicher Alzheimer-Erkrankungs-Patient ist, das den Antikörper nach Anspruch 1 oder 2 umfasst.

## Revendications

1. Un anticorps qui se lie à un oligomère A $\beta$  mais pas à un monomère A $\beta$ , l'anticorps étant un anticorps monoclonal et choisi dans le groupe constitué de :

(1) un anticorps qui comprend une chaîne H ayant la séquence d'acides aminés de SEQ ID NO: 69 comme CDR1, la séquence d'acides aminés de SEQ ID NO: 71 comme CDR2, et la séquence d'acides aminés de SEQ ID NO: 73 comme CDR3 et une chaîne L ayant la séquence d'acides aminés de SEQ ID NO: 75 comme CDR1, la séquence d'acides aminés de SEQ ID NO: 77 comme CDR2, et la séquence d'acides aminés de SEQ ID NO: 79 comme CDR3 ;

(2) un anticorps qui comprend une chaîne H ayant la séquence d'acides aminés de SEQ ID NO: 65 comme VH et une chaîne L ayant la séquence d'acides aminés de SEQ ID NO: 67 comme VL ; et

(3) un anticorps comprenant une chaîne H ayant la séquence d'acides aminés de SEQ ID NO: 61 et une chaîne L ayant la séquence d'acides aminés de SEQ ID NO: 63.

2. Anticorps selon la revendication 1, l'anticorps étant un anticorps chimérique ou un anticorps humanisé.

3. Une composition comprenant l'anticorps de la revendication 1 ou 2 et un véhicule pharmaceutiquement acceptable.

4. Un agent pour son utilisation dans le traitement d'une déficience cognitive, comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

5. Un agent thérapeutique pour son utilisation dans le traitement de la maladie d'Alzheimer, comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

6. Un agent pour son utilisation dans la suppression de la progression de la maladie d'Alzheimer, comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

7. Un agent pour son utilisation dans la suppression de la formation de la plaque sénile, comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

8. Un agent pour son utilisation dans la suppression de l'accumulation de A $\beta$ , comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

9. Un agent pour son utilisation dans la neutralisation de la neurotoxicité, comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

10. Un agent pour son utilisation dans l'inhibition de la formation de fibrilles amyloïdes A $\beta$ , comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

11. Un agent pour son utilisation dans la neutralisation de la toxicité synaptique, comprenant l'anticorps de la revendication 1 ou 2 ou la composition de la revendication 3 comme principe actif.

12. Une méthode de détection d'un oligomère A $\beta$ , comprenant l'étape de détection d'un oligomère A $\beta$  contenu dans un échantillon collecté chez un sujet en utilisant l'anticorps de la revendication 1 ou 2.

13. Une méthode permettant de diagnostiquer si un sujet est possiblement un patient souffrant de la maladie d'Alzheimer,



comprenant  
l'étape (a0) ci-dessous :

5 (a0) utiliser l'anticorps de la revendication 1 ou 2 pour détecter un oligomère A $\beta$  dans un échantillon collecté chez un sujet ;

ou  
les étapes (a1) à (c1) ci-dessous :

10 (a1) mettre en contact un échantillon collecté chez un sujet avec l'anticorps de la revendication 1 ou 2 ;  
(b1) mesurer la quantité d'oligomère A $\beta$  dans l'échantillon ; et  
(c1) déterminer si le sujet est possiblement un patient souffrant de la maladie d'Alzheimer, lorsque la quantité mesurée à l'étape (b1) est supérieure à celle d'un individu sain ;

15 ou  
les étapes (a2) à (c2) ci-dessous :

(a2) mettre en contact un échantillon collecté chez un sujet avec l'anticorps de la revendication 1 ou 2 et un anticorps qui se lie à un monomère A $\beta$  ;  
20 (b2) mesurer le ratio oligomère A $\beta$  sur monomère A $\beta$  dans l'échantillon ; et  
(c2) déterminer si le sujet est possiblement un patient souffrant de la maladie d'Alzheimer, lorsque le ratio mesuré à l'étape (b2) est supérieur à celui d'un individu sain.

25 **14.** La méthode de la revendication 12 ou 13, dans laquelle l'échantillon est du sang ou du liquide céphalorachidien.

**15.** Un agent pharmaceutique pour son utilisation pour diagnostiquer si un sujet est possiblement un patient souffrant de la maladie d'Alzheimer, comprenant l'anticorps de la revendication 1 ou 2.

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7-1






FIG. 7-2




FIG. 8




FIG. 9




FIG. 10




FIG. 11







FIG. 12

**A** 1A9-TREATED Tg2576 MICE

A $\beta$




MOUSE IgG



**B** 2C3-TREATED Tg2576 MICE

A $\beta$

MOUSE IgG



**C** PBS-TREATED Tg2576 MICE

A $\beta$

MOUSE IgG






FIG. 13




FIG. 14






FIG. 15




FIG. 16




FIG. 17

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- WO 03104437 A2 [0003]
- WO 2006055178 A1 [0003]
- WO 9203918 A [0036]
- WO 9402602 A [0036]
- WO 9425585 A [0036]
- WO 9633735 A [0036]
- WO 9634096 A [0036]
- JP S6317688 A [0037]
- WO 9007861 A [0075]
- JP 2009052039 W [0176]
- JP 2008028386 A [0176]
- US 61085545 B [0176]
- JP 2008201058 A [0176]

## Non-patent literature cited in the description

- KLEIN WL. *Trends Neurosci*, 2001, vol. 24, 219-224 [0003]
- SELKOE DJ. *Science*, 2002, vol. 298, 789-791 [0003]
- HASS C et al. *Nature Review*, 2007, vol. 8, 101-12 [0003]
- LEE EB et al. *J. Biol. Chem.*, 2006, vol. 281, 4292-4299 [0003]
- MORETTO N. et al. Conformation-sensitive antibodies against Alzheimer amyloid- $\beta$  by immunization with a thioredoxin-constrained B-cell epitope peptide. *J. Biol. Chem.*, 2007, vol. 282 (15), 11436-11445 [0003]
- KOHLER ; MILSTEIN. *Nature*, 1975, vol. 256, 495-7 [0024]
- CABILLY et al. *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 3273-7 [0024]
- KOHLER G. ; MILSTEIN C. *Methods Enzymol.*, 1981, vol. 73, 3-46 [0029]
- Antibodies: A Laboratory manual. Cold Spring Harbor Laboratory, 1988 [0031]
- DANIEL R MARSHAK et al. Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Cold Spring Harbor Laboratory Press, 1996 [0033]
- ISHIDA I et al. *Cloning and Stem Cells*, 2002, vol. 4, 91-102 [0036]
- BORREBAECK CAK ; LARRICK JW. Therapeutic Monoclonal Antibodies. MacMillan Publishers, 1990 [0038]
- HASHIMOTO-GOTOH, T. et al. *Gene*, 1995, vol. 152, 271-275 [0065]
- ZOLLER, MJ ; SMITH, M. *Methods Enzymol.*, 1983, vol. 100, 468-500 [0065]
- KRAMER, W. et al. *Nucleic Acids Res.*, 1984, vol. 12, 9441-9456 [0065]
- KRAMER W ; FRITZ HJ. *Methods. Enzymol.*, 1987, vol. 154, 350-367 [0065]
- KUNKEL, TA. *Proc. Natl. Acad. Sci. USA.*, 1985, vol. 82, 488-492 [0065]
- KUNKEL. *Methods Enzymol.*, 1988, vol. 85, 2763-2766 [0065]
- HOPP, T.P. et al. *BioTechnology*, 1988, vol. 6, 1204-1210 [0069]
- JONES et al. *Nature*, 1986, vol. 321, 522-5 [0075]
- RIECHMANN et al. *Nature*, 1988, vol. 332, 323-7 [0075]
- PRESTA. *Curr. Opin. Struct. Biol.*, 1992, vol. 2, 593-6 [0075]
- LARRICK et al. *Methods: a Companion to Methods in Enzymology*, 1991, vol. 2, 106 [0075]
- Genetic Manipulation of Monoclonal Antibodies. COURTENAY-LUCK et al. Monoclonal Antibodies: Production, Engineering and Clinical Application. Cambridge University Press, 1995, 166 [0075]
- Genetic Manipulation and Expression of Antibodies. WARD et al. Monoclonal Antibodies: Principles and Applications. Wiley-Liss, Inc, 1995, 137 [0075]
- VERHOEYEN et al. *Science*, 1988, vol. 239, 1534-6 [0075]
- QUEEN et al. *Proc. Natl. Acad. Sci. USA*, 1989, vol. 86, 10029-33 [0075]
- Immunoglobulin genes. Academic Press, 1989, 260-74 [0076]
- MICHAEL A et al. *Proc. Natl. Acad. Sci. USA*, 1994, vol. 91, 969-73 [0076]
- KLEIN WL. *Trends Neurosci*, 2001 [0083]
- SELKOE DJ. *Science*, 2002 [0083]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1980 [0094]
- GHISO J et al. *Biochem J*, 1993 [0121]
- MATSUBARA E et al. *Neurobiol Aging*, 2004 [0121]
- MATSUBARA E. et al. *Neurobiol Aging*, 2004, vol. 25, 833-841 [0123]
- TERO, R et al. *Langmuir*, 2004, vol. 20, 7526-7531 [0131]

- **KATSUNO T.** *Neurology*, 2005, vol. 64, 687-692 [0132]
- **MOURIA.** *FASEB J*, 2007, vol. 21, 2135-2148 [0134]
- **MATSUBARA E et al.** *Ann Neurol*, 1999, vol. 45, 537-541 [0135]
- **MATSUBARA E et al.** *Ann Neurol*, 1999, 537-541 [0137]
- **MATSUBARA E et al.** *Neurobiol Aging*, 2004, vol. 25, 833-841 [0137] [0141] [0152]
- **NAKAYA Y et al.** *J Biol Chem*, 2005, vol. 280, 19070-19077 [0141]
- **YAMAMOTO N et al.** *J Biol Chem*, 2007, vol. 282, 2646-2655 [0141]
- **KAYED R et al.** *Science*, 2003, vol. 300, 486-489 [0148]
- **WRZOLEK MA et al.** *Am J Pathol*, 1992, vol. 141, 343-355 [0150]
- **KATSUNO et al.** *Neurology*, 2005, vol. 64, 687-692 [0151]
- **LEE EB et al.** *J Biol Chem*, 2006, vol. 281, 4292-4299 [0152]
- **JANUS D.** *Nature*, 2000 [0172]
- **MORGAN D.** *Nature*, 2000 [0172]
- **BARD F.** *Nat med*, 2222 [0172]
- **DEMATTOS RB.** *PNAS*, 2001 [0172]
- **DEMATTOS RB et al.** *PNAS*, 2001, vol. 98, 8850-8855 [0172]
- **DEANE R.** *J Neurosci*, 2005 [0172]

专利名称(译)	能够特异性结合ab-寡聚体的抗体及其用途		
公开(公告)号	<a href="#">EP3121277B1</a>	公开(公告)日	2018-04-11
申请号	EP2016183829	申请日	2009-02-06
[标]申请(专利权)人(译)	NAT癌症CENT		
申请(专利权)人(译)	IMMUNAS PHARMA , INC. 国家中心的老年医学和老年病学		
当前申请(专利权)人(译)	IMMUNAS PHARMA , INC. 国家中心的老年医学和老年病学		
[标]发明人	MATSUBARA ETSURO SHIBATA MASAO YOKOSEKI TATSUKI		
发明人	MATSUBARA, ETSURO SHIBATA, MASAO YOKOSEKI, TATSUKI		
IPC分类号	C12N15/09 A61K39/395 A61P25/28 C07K16/18 G01N33/53 C12P21/08 G01N33/68		
CPC分类号	A61K2039/505 A61P17/00 A61P25/00 A61P25/28 C07K16/18 C07K2317/56 C07K2317/565 C07K2317/73 C07K2317/76 G01N33/6896 G01N2333/4709 G01N2800/2821 A61K39/3955		
优先权	2008028386 2008-02-08 JP 12/085545 2008-08-01 US 2008201058 2008-08-04 JP		
其他公开文献	EP3121277A1		
外部链接	<a href="#">Espacenet</a>		

## 摘要(译)

本发明人成功制备了仅对可溶性A 2寡聚体特异的单克隆抗体，但不识别可溶性A 2单体，其为生理分子。已证实该抗体可用作阿尔茨海默氏病的诊断/治疗性单克隆抗体。

```

Asp Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1      5      10
Ser Arg Lys Leu Ser Cys Ala Thr Ser Gly Phe Thr Phe Ser Ser Phe
20      25      30
Gly Met His Trp Val Arg Gln Ala Pro Glu Lys Gly Leu Glu Trp Val
35      40      45
Ala Tyr Ile Ser Ser Gly Ser Ser Ala Ile Tyr Tyr Ala Asp Thr Val
50      55      60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Pro Lys Asn Thr Leu Phe
65      70      75      80
Leu Gln Met Thr Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85      90      95
Ala Arg Ser Gly Asp Thr Met Asp Tyr Trp Gly Gln Gly Thr Ser Val
100     105     110
Thr Val Ser Ser Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala
115     120     125
Pro Gly Cys Gly Asp Thr Thr Gly Ser Ser Val Thr Leu Gly Cys Leu
130     135     140
Val Lys Gly Tyr Phe Pro Glu Ser Val Thr Val Thr Trp Asn Ser Gly
145     150     155     160
Ser Leu Ser Ser Ser Val His Thr Phe Pro Ala Leu Leu Gln Ser Gly

```