




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12Q 1/68</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/52208</b>
			(43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/05143		bridge, MA 02138 (US). YERSHOV, Gennadiy [RU/US]; 105453 Echo Lane #5, Hindsdale, IL 60521 (US). LYSOV, Yuri [RU/US]; 16 West 767 Mockingbird Lane #201, Hindsdale, IL 60521 (US).  (74) Agents: MARTIN, Alice, O.; Barnes & Thornburg, 2600 Chase Plaza, 10 South LaSalle Street, Chicago, IL 60603 (US) et al.  (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 29 February 2000 (29.02.00)			
(30) Priority Data: 09/261,115 3 March 1999 (03.03.99) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/261,115 (CIP) Filed on 3 March 1999 (03.03.99)			
(71) Applicant (for all designated States except US): THE UNIVERSITY OF CHICAGO [-/US]; 5801 South Ellis Avenue, Chicago, IL 60637 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): MIRZABEKOV, Andrei [RU/US]; 1005 Village Court, Darien, IL 60516 (US). GUSCHIN, Dmitry Y. [RU/US]; Apartment T-3, 259 Congressional Lane, Rockville, MD 20852 (US). CHIK, Valentine [RU/US]; Apt. 111, 2540 Crystal Court, Woodridge, IL 60517 (US). DROBYSHEV, Aleksei [RU/RU]; Sportivnaya St., #9A-3, Elektrosol, Moscow Region, 1444009 (RU). FOTIN, Alexander [RU/US]; 24 Everett St. #309, Cam-			

**Published**

Without international search report and to be republished upon receipt of that report.

(54) Title: CUSTOMIZED OLIGONUCLEOTIDE MICROCHIPS THAT CONVERT MULTIPLE GENETIC INFORMATION TO SIMPLER PATTERNS, ARE PORTABLE AND REUSABLE

**(57) Abstract**

This invention relates to using customized oligonucleotide microchips as biosensors for the detection and identification of the nucleic acids specific for different genes, organisms and/or individuals in the environment, in food and in biological samples. The microchips are designed to convert multiple bits of genetic information into simpler patterns of signals that are interpreted as a unit. Because of an improved method of hybridizing oligonucleotides from samples to microchips, microchips are reusable and transportable. For field study, portable laser or bar code scanners are suitable.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**CUSTOMIZED OLIGONUCLEOTIDE MICROCHIPS THAT CONVERT  
MULTIPLE GENETIC INFORMATION TO SIMPLER PATTERNS, ARE  
PORTABLE AND REUSABLE**

5           The U.S. Government has rights to the invention pursuant to Contract  
W-31-109-ENG between the U.S. Department of Energy and the University of  
Chicago representing (Argonne National Laboratory).

          A novel microchip is customized to answer specific questions and has  
oligonucleotides positioned on the microchip so that multiple bits of information are  
10   evidenced to a simpler pattern. A new method of hybridization to a microchip is also  
presented.

**BACKGROUND OF THE INVENTION**

          Differences in nucleotide and amino acid sequences may be exploited to  
analyze environmental, food or biological samples. Detection and identification of  
15   microorganisms is important for clinical purposes and for determination of  
contaminated food, air, water or soil. Studies in environmental microbiology are  
often limited by the inability to unambiguously identify and directly quantify the  
enormous diversity of natural populations. This problem is now changing with  
increasing use of molecular techniques to directly measure different genetic features.  
20   (Mobarry *et al.*, 1996; Stahl, 1995; Wagner *et al.*, 1995) For example, DNA probes  
are now commonly used to detect by hybridization, genes encoding proteins involved  
in specific catabolic functions, and to resolve different genetic populations in the  
environment. In particular, the use of group-specific DNA probes complementary to  
the small subunit (SSU) 16S rRNA has provided a comprehensive framework for  
25   studies of microbial population structure in complex systems. Sequencing of this  
subunit revolutionized microbial classification and led to the discovery of  
archebacteria. (Woese, 1987) A large number of the sequences for different  
organisms has been collected. (Maidak *et al.*, 1996) Every microorganism species is  
characterized by a specific DNA sequence within a variable region of its ribosomal  
30   RNA gene or other genes. A highly efficient procedure for microorganism  
classification and for construction of their evolutionary trees is based on these  
observations. Identification of specific sequences in ribosomal DNA is a reliable  
microbial analysis that can be carried out by direct DNA sequencing. However DNA

sequencing is a rather complicated, expensive and time consuming procedure to use for serial microbial analysis on a commercial scale for environmental or medical applications. Consequently, new methods are needed to make sequence matching commercially feasible.

5           Also, methods are needed that are transportable to the field. A nucleic acid hybridization is a highly specific and sensitive procedure that allows a specific sequence to be detected and identified among other millions of sequences in a genome of higher organisms, or among a mixture of different organisms. The principle of hybridization is that sequences hybridize as a function of the similarity of their linear  
10   nucleotide sequence. The hybridization of DNA or RNA extracted from even a very complicated mixture to a specific oligonucleotide probe has resulted in unambiguous identification of specific microorganisms in an environmental sample, for example. In the course of such an analysis, RNA or DNA is extracted from a sample of microorganisms isolated from water solutions, air or soil, immobilized on a filter and  
15   then hybridized successively with several oligonucleotide probes for different microorganisms. However, for this purpose, the sample needs to be checked for the presence of hundreds or thousands of different oligonucleotides corresponding to various microorganisms which is prohibitively laborious and expensive using present methods and yields results that must be interpreted by a computer in order to decipher  
20   the identification. What is needed is a simplified pattern to provide rapid answers to specific questions, e.g. are any known pathogens in a water sample?

          The scope of applications of nucleotide hybridization is often limited by the nature of the assays, generally involving the independent hybridization and interpretation of multiple environmental samples to multiple DNA probes. In  
25   addition, some detection assays require amplification of the target nucleic acid, for example, via PCR. This may contribute to quantitative biases. Thus, there is need for assays that provide for greater sample through-put capacity and greater sensitivity, rapid read-out of results.

          Another area in which specific DNA or RNA sequences are of interest is  
30   mutation and polymorphism analyses. The number of base changes discovered (mutations) in different genes is growing rapidly. These changes are associated with genetic diseases, with disease predispositions and cancers, with development of drug

resistance in microorganisms, and with genetic polymorphisms. Polymorphisms are useful for determining the source of a sample, e.g. in forensic analyses.

Polymorphisms such as in the HLA system are essential to predict success of tissue transplants. The ability to simultaneously analyze many mutations in a gene in a simple, fast, and inexpensive way is essential in clinical medicine and this need has stimulated the development of different methods for screening mutations, but all have serious limitations. What is needed are kits that are transportable and interpretable, e.g. for use in clinics without high technology microscopes.

Hybridization of filter-immobilized DNA with allele-specific oligonucleotides was suggested as a way to screen for mutations. (Conner *et al.*, 1983) However, the number of alleles that can be assayed at one time is limited, the filters are usable only for a few times, and there is little opportunity for complex analysis or easy interpretation of results.

A possible solution to large scale hybridization is to use microchips for DNA sequence hybridizations (SHOM, sequencing by hybridization with oligonucleotides in a microchip) (e.g. Khrapko, 1996; Yershov, 1996). The development of an array of hundreds or thousands of immobilized oligonucleotides, the so-called "oligonucleotide chips", permits simultaneous analysis of many mutations (for a review, see Mirzabekov, 1994). Such arrays can be manufactured by a parallel synthesis of oligonucleotides (Southern *et al.*, 1992; Fodor *et al.*, 1991; Pease *et al.*, 1994; Matson *et al.*, 1995) or by chemical immobilization of presynthesized oligonucleotides (Khrapko *et al.*, 1991; Lamture *et al.*, 1994; Ghu *et al.*, 1994). Glass surfaces (Southern *et al.*, 1992; Fodor *et al.*, 1991; Ghu *et al.*, 1994), glass pores (Beattie *et al.*, 1995), polypropylene sheets (Matson *et al.*, 1995), and gel pads (Khrapko *et al.*, 1991; Yershov *et al.*, 1996) have been used as solid supports for oligonucleotide immobilization. However "Oligonucleotide array technology has not yet lived up to its promise." Southern, 1996 p. 115.

Some of the deficiencies in the art are unpredictability of the results, lack of knowledge of optimum conditions, and failure to demonstrate accuracy and commercial feasibility. Moreover, analysis of the results of hybridization requires computer programs capable of assimilating and interpreting multiples bits of

information, and high technology microscopes. The microchips are neither portable, reusable, nor easily interpreted.

### **SUMMARY OF THE INVENTION**

This invention embodies applications of oligonucleotide microchip technology wherein the microchip is a biosensor and customized oligonucleotide microchips are designed for specific applications of nucleic acid hybridization.

Hybridization is a process by which, under defined reaction conditions, partially or completely complementary nucleic acids are allowed to join in an antiparallel fashion to form specific and stable hydrogen bonds.

Aspects of the invention include:

1. microchips designed so that multiple bits of genetic information are converted to a pattern, which is interpreted as a unit, wherein the appearance of the pattern provides answers to specific questions; this construction facilitates providing easily interpretable answers provided by hybridization patterns and removes some need for high technology instruments to interpret the results of hybridization; and
2. improved methods of hybridizing oligonucleotides in a sample to oligonucleotides on a customized microchip do not require a washing step but rather measure non-equilibrium melting curves (temperature curves) that do not require washing with a solution that removes immobilized oligonucleotides from microchips; this means that microchips are reusable because the oligonucleotides anchored within the gel elements, do not wash away, and are available for reuse. (Microchips with samples are generally kept in solution, however, microchips can be dried and stored for many months before being reused.)

The patterns exhibited after hybridization to a microchip generally are not directly related to the nature of the hybridizations and are not simply converting a "yes" or a "no" signal, or a "positive" or "negative" signal to a binary outcome, nor are the patterns of the present invention converting a gradation of quantities to another form of gradation, e.g. colorimetric gradations. The deliberate organization of the oligonucleotides on the microchips themselves does not transmit information; only after hybridization with a test nucleic acid will the hybridization signal itself form the pattern. The pattern is then detected by a detection means which can include visual interpretation without the aid of additional detection instrumentation.

By choosing ordered schemes of oligonucleotide positioning on the microchips, visual signals are simplified and enhanced, e.g. the letter "P" is observed if certain pathogenic groups are present; columns of gel elements on the chip that include the same oligonucleotide probes, will be readily detectable as a positive linear  
5 column, if the matching oligonucleotides are in the test sample. The visual appearance may be strong enough to see with the naked eye, may be determined with a UPC (Universal Product Code or "bar code") laser scanner, or with a laser gun. The wavelength of the scanner and the sensor that accepts the signal for a bar code must be concordant with the dye or label used to hybridize the DNA.

10 Of course, aspects one and two do not have to be used together. Designs that result from converting multiple amounts of genetic information obtained by large numbers of hybridizations of oligonucleotides to simpler, readily interpretable patterns, could be done on microchips constructed and analyzed by the methods used prior to the present invention.

15 Similarly, the improved methods of providing hybridization results on microchips could be used on microchips that are not designed to convert multiple pieces of genetic information into a simpler pattern.

Other aspects of the invention include improved predictability, increased accuracy, and standardized factors for detection and identification of nucleotide  
20 sequences. The improvements result from optimizing conditions, methods and compositions for microchip hybridization. Deliberate ordered schemes that are designed to answer specific questions and that convert complex data to simpler patterns, are followed so that much hybridization information can be readily obtained from a single scan of a microchip to detect hybridization of immobilized  
25 oligonucleotides by nucleic acids in a sample to be investigated. Samples include air, water, soil, blood, cells, tissue, tissue culture and a food. An aspect of the invention is that the same microchip can be used for hybridization for more than 20-30 times, without any noticeable deterioration of the hybridization signal because immobilized oligonucleotides are not washed out or stripped. Customized sets of microchips are  
30 obtained for specific applications. Also, parallel hybridization of nucleic acids in a sample to many oligonucleotides on a microchip is possible, allowing replication and standardization. For example, the sequence diversity of SSU rRNAs recovered from

different microbial populations of varying abundances is analyzed by a single hybridization to a microchip. A large number of HLA alleles, are assayed by a single hybridization to a microchip.

The invention relates a method for identifying a nucleotide sequence in a sample using a microchip, said method comprising:

- a) providing a customized matrix of oligonucleotides on the microchip designed to identify genetic sequences in the sample, wherein an ordered scheme positions oligonucleotides to provide a pattern to answer specific questions after hybridization;
- b) hybridizing nucleic acids extracted from the sample as such or after amplification on said microchip; and
- c) identifying the nucleotide sequences represented in said sample by analyzing the pattern of the oligonucleotides which hybridized to the sequences, said pattern provided by signals.

The nucleic acids suitable for the practice of the invention include DNA, mRNA, 16S rRNA sequences and other RNA species.

Customized oligonucleotide microchips are aspects of the invention. The microchip includes a gel-matrix affixed to a support, said matrix is formed by a plurality of gel pad element sites. The number of sites is determined by the number of oligonucleotides in the array. Each gel element contains one chemically immobilized oligonucleotide of a desired sequence, length and concentration; the gel elements being separated from one another by hydrophobic glass spaces and the gel portions having a vertical height above the plane of the interstitial spaces of generally not more than 30  $\mu\text{m}$ . In some applications, the same type of oligonucleotides may be immobilized to different gel pads to form a pattern.

The invention relates screening nucleic acid preparations for genes, RNA transcripts or any other unique nucleotide sequences, for example those that encode microbial 16S ribosomal RNAs. Ratios of DNA/RNA or any other unique nucleotide sequences specific for certain types of organisms are suitable. Multiple labeling allows simultaneous detection and quantitative comparison of different nucleic acid sequences that are hybridized to a microchip.



The methods of the present invention include labeling the oligonucleotide sequence in said sample before bringing it in contact with the array. A suitable label is a fluorescent dye. A plurality of different dyes may be used concurrently.

Oligonucleotides immobilized on a customized microchip include those

- 5 complementary to the beta globin gene, sequences specific for *Salmonella*, or polymorphic HLA allele sequences.

- 10 An oligonucleotide microchip for the detection and classification of nitrifying bacteria has a customized design wherein identifying labels in the cells of the microchip refer to oligonucleotides selected from a class of bacteria, and the selection is designed to answer specific questions regarding classification.

- 15 An embodiment of an application of the present invention is detecting and identifying microorganisms in samples obtained from the environment, e.g. water, air or soil samples to check for pollutants; biological samples obtained for medical diagnosis; or food samples to check for contamination. Other applications include forensic testing to identify DNA in samples obtained for criminal investigations, and detection of chromosomal fragments, or single gene mutations e.g. for diagnosing genetic diseases such as  $\beta$ -thalassemia or types of cancers. Tissue typing for polymorphic HLA alleles for transplantation or studying human diversity is facilitated.

- 20 The nucleic acid preparations are made from samples collected in any type of environment, where detection and identification of the microorganisms in that environment is of interest, or where it is likely that new (previously unidentified) organisms may be discovered.

- 25 DNA and RNA molecules in a sample can be separated from each other during their isolation and labeled with different fluorescent dyes. These RNA and DNA molecules are simultaneously hybridized with oligonucleotides on a microchip that is specific to the sample to be tested. The quantitative monitoring of the simultaneous hybridization of differently labeled DNA and RNA with a microscope that can discriminate multicolors at several wave lengths allows the calculation of DNA/RNA ratios in the sample. For bacterial samples, this ratio determines the state of vitality and physiological activity of the bacterium. In an embodiment, the ratio of RNA/DNA is used to discriminate the dead bacterium cells and spores from the active
- 30

state of microbial growth. In the same way, a DNA or RNA molecule of a bacterial strain stained with one dye can be added in a calculated amount as an internal standard to a sequence or sequences under investigation in which the sequences being investigated stained with a different (second) dye. The fluorescence measurements of hybridization intensities at different wave lengths for the standard and investigated sequences (probes) allow relative quantitative ratios to be determined.

Hybridization on microchips allows unambiguous typing of different groups of chosen bacteria in a sample. Microchip hybridization is a simple, fast, inexpensive and reliable method for bacterial typing.

An aspect of the invention is that there is no limitation on the number of sequences that can be checked or the number of types of microorganisms that can be detected. Instead of multiple sequential hybridizations with different probes of, *e.g.* a 16S rRNA preparation, only one round of hybridization is required to find out what different sequences are in a sample. The volume of hybridizations is dramatically reduced and the assay requires much less RNA or DNA compared with standard techniques. An advantage is that culturing of bacteria and gene amplification can be avoided.

Methods of the invention significantly reduce sample preparation time, avoid the culturing of organisms collected from field situations, and allow the identification of all species of microorganisms contained in a particular sample. Portable microchips are available for field work.

For example, oligonucleotides complementary to small subunit rRNA sequences of selected microbial groups, encompassing key genera of nitrifying bacteria, were shown to selectively retain or hybridize with labeled target nucleic acid derived from either DNA or RNA forms of the target sequences. Methods and compositions of the present invention discriminate among the Genera, *Nitrosomonas*, *Nitrobacter* and *Nitrosovibrio sp.* using fluorescently labeled nucleic acid probes that hybridize to 16S rRNA sequences. Each species has specific DNA sequences within the variable region of its rRNA genes. Since the rRNAs are naturally amplified, often present in thousand of copies per cell, they provide greater sensitivity, eliminating the need for amplification in many applications.

The invention facilitates identification of organisms from environmental samples in a faster, and more economical approach than presently available. In addition, new species may be discovered that would be highly informative regarding taxonomic status of known as well as newly discovered organisms.

5           A diagnostic assay of the present invention for a mutation in a gene, includes the following steps:

                  a.       designing a customized oligonucleotide microchip biosensor comprising oligonucleotides that hybridize to a gene having the mutation, wherein the oligonucleotides are positioned on the microchips so that patterns result depending on  
10   what oligonucleotides are in the sample to answer a specific question(s);

                  b.       contacting a nucleic acid sample to the customized oligonucleotide microchip biosensor under conditions that allow hybridization of the nucleic acid to the microchip; and

                  c.       determining the pattern of hybridization from which  
15   observation the presence of specific nucleic acid sequences is inferred and the specific question is answered.

For diagnostic assays for genetic diseases, sequence analysis of DNA is carried out by hybridization of PCR amplified DNA or its RNA transcripts with oligonucleotide array microchips. Polyacrylamide gel pads containing allele-specific  
20   immobilized oligonucleotides are fixed on a glass slide of the microchip. The RNA transcripts of PCR-amplified genomic DNA are optionally fluorescently labeled by enzymatic or chemical methods and hybridized with the microchip. In the field, the chemical methods are preferred because results are obtained faster, and some chemicals will fragment DNA at the same time which is needed for the sample.

25           When melting curve experiments are performed, both matching and mismatching oligos can be immobilized in the gel pads, and both matching and mismatching nucleic acids can be in the sample. The biochips are reusable in two types of embodiments: 1) the sample or test nucleic acids can be removed or stripped off the chip and a different test sample can be introduced and 2) the same melting  
30   point curve experiments can be run and re-run without any washing.

When experiments are performed with a different test sample, the original sample is removed from the chip by a washing or stripping procedure using distilled

water at 60°C with an hour (or up to overnight) incubation. If the melting curve experiments are repeated (or reused) then the same sample is left in contact with the chip and appearance and disappearance of hybridization signal is observed over a variety of temperatures, usually ranging from 0° - 50° C.

5           When the chips are incubated, in order to remove the sample nucleotides, virtually none of the immobilized oligos are removed in the process. This is because the oligos are covalently linked to the gel matrix of the gel pads that form the microchip.

10           Repeated reuse of the chips in which different samples are applied after sequential removal is usually limited to about 50 uses, because eventually the amount of non-specific or background hybridization signal is greater than one-tenth of a mismatch hybridization signal. The conditions under which a chip would not be reusable (up to 50 times) are very few. Such conditions include allowing the chips to be cooled to -20°C or performing experiments where the chips are heated to above  
15   70°C, conditions that have been shown to cause degradation of the chips, thus rendering them unstable.

          The simultaneous measurement in real time of the hybridization and melting curves on the entire oligonucleotide array is carried out with a fluorescence microscope with a laser light source equipped with CCD camera or a special laser  
20   scanner. Some work only with dried microchips. The monitoring of the hybridization specificity for duplexes with different stabilities and AT content is enhanced by its measurement at optimal discrimination temperatures on melting curves. Microchip diagnostics are optimized by choosing the proper allele-specific oligonucleotides from among the set of overlapping oligomers. The accuracy of mutation detection can be  
25   increased by simultaneous hybridization of the microchip with at least two differently labeled samples of normal and mutated alleles, and by parallel monitoring their hybridization with a multi-wavelength fluorescence microscope. The efficiency and reliability of the sequence analysis was demonstrated by diagnosing  $\beta$ -thalassemia mutations and HLA polymorphisms. Determining levels of gene expression is an  
30   aspect of the invention.

          Because the methods of the present invention require only a simple procedure of hybridization and because only one round of hybridization is necessary, it is fast

and inexpensive. Because the invention allows a lot of information to be obtained from one experiment, in a simple pattern as compared to the analysis of hundreds of data points, it has increased efficiency. The invention is reliable because the microchips are reusable. Immobilized oligonucleotides are not washed out. There is  
5 no waste of hybridization probes, therefore the microchip hybridization is inexpensive and non-isotopic detection simplifies all procedures.

Effective and precise sequence analysis by the hybridization of a probe with rather short microchip-immobilized oligonucleotides depends on many factors. Major factors are the reliability of the discrimination of perfect duplexes from duplexes  
10 containing mismatches, differences in stability of AT- and GC-rich duplexes, the efficiency of the hybridization, and simplicity in the preparation of the labeled samples for hybridization.

Identification of base variations is significantly improved by parallel measuring of the melting curves of the duplexes formed on the entire oligonucleotide  
15 array, as well as by monitoring the simultaneous hybridization of two differently labeled samples at two wavelengths and by choosing proper allele-specific oligonucleotides.

Other factors to be considered for operation of the invention include (1) regulating the flow of the fluid containing a sample to be tested over the microchip  
20 during the hybridization; and (2) control of the temperature of the microchip gel layer and the fluid layer, in a differential manner, by placing a cooling and heating apparatus adjacent to the gel layer and the top fluid layer. The gel layer temperature is controlled in a uniform or gradient manner by a heating/cooling device attached to the glass plate substrate of the gels. For field work, the optimum temperature for a  
25 particular question is determined previously in a laboratory.

A definition of "customized microchip" is a microchip of gel elements on a support, wherein the oligonucleotides are immobilized in gel elements according to an ordered scheme such that multiple bits of information are ordered to a simpler pattern to answer a specific question.

30 Removal of test or sample nucleic acids from microchip is accomplished by an incubation step carried out using distilled water for at least one hour (up to overnight) at 60°C. (This procedure is analogous to the step of "stripping" a filter for re-use in

the standard technique of probing a Southern blot.) The immobilized oligonucleotides in the gel matrix are not removed by this incubation as the oligonucleotides are covalently linked to the gel substrate.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

5           FIG. 1 shows non-equilibrium melting curves of duplexes of RNA with microchip oligonucleotides.

FIG. 2 shows an example of four melting curves for 75-nt-long RNA fragments hybridized with the microchip oligonucleotides. The RNA was derived from a patient having the IVS I/2 T/A mutation in the  $\beta$ -globin gene. The curves were  
10           normalized to the initial hybridization signals. Melting curves 1 and 3 correspond to perfect duplexes; curves 2 and 4 correspond to duplexes containing internal T-T or G-T mismatches, respectively. The curves for the perfect and mismatched duplexes are shifted by about 10°C from each other.

FIG. 3 shows hybridization of fluorescein labelled 16S rRNAs to a microchip.  
15           The microchip with immobilized probes (see Table 1 and Table 2) was hybridized sequentially to *in vitro* transcribed 16S rRNA of *Nitrosovibrio tenuis* (A), *Nitrosomonas europaea* (B), *E. coli* (C), and with *E. coli* rRNA recovered from isolated ribosomes (D). The panels to the right display the number of mismatches between each probe and the RNA.

20           FIGS. 4A and 4B show hybridization of the mixture of differently labelled *E. coli* and *Nitrosovibrio tenuis* rRNAs to the microchip at 10°C and 40°C, measured simultaneously by multicolor detection. A. The microchip was hybridized with a mixture of fluorescein labelled *Nitrosovibrio tenuis* and tetramethylrhodamine labelled *E. coli* 16S rRNA and washed serially at the indicated temperatures, a.u. -  
25           arbitrary units of fluorescence intensities. B. The ratio of the hybridization intensities of *Nitrosovibrio tenuis* ( $I_{Nt}$ ) to *E. coli* ( $I_{E.coli}$ ) 16S RNA measured at 10°C and 40°C.  
 $R = (I_{Nt}/I_{E.coli})$ .

FIG. 5 illustrates the concentration effect of the immobilized oligonucleotides on the hybridization intensities. A microchip with different concentrations of  
30           immobilized oligonucleotides was hybridized with *N. tenuis* 16S rRNA labelled with fluorescein and washed at 20°C. Curve 1 corresponded to Nsv443 (nitrosovibrio-like) probe, curve 2 - Bac338 (Bacteria), curve 3 - Nso1225 (ammonia oxidizers), curve 4 -

Uni1390 (all life), and curve 5 - Nsm-156 (nitrosomonas), a.u. - arbitrary units of fluorescence intensities.

FIG. 6 shows the sequences of  $\beta$ -globin alleles specifying oligonucleotides that were immobilized on a microchip.

5        FIG. 7 shows the experimental design to detect  $\beta$ -globin mutations using oligonucleotide microchips.

FIG. 8 shows results of gene expression studies.

FIG. 9 shows 18 short HLA oligonucleotides.

FIG. 10 shows HLA oligonucleotides hybridized to the microchips.

10       FIG. 11 illustrates a closed microchamber 1 containing a microchip with a gel array 3 on a glass support 4; ports 2 are used merely to provide wetting solution.

FIG. 12 shows an ordered scheme in which a letter "P" will be detected if there is a group of hybridizations of oligonucleotides from a sample that are oligonucleotides from pathogens.

15       FIG. 13 illustrates an ordered scheme on a microchip wherein the presence of *B. anthracis*.

#### **DESCRIPTION OF THE PREFERRED EMBODIMENT**

This invention relates to using customized oligonucleotide microchips as biosensors for the detection and identification of nucleic acids specific for different  
20 genes, organisms and individuals in the environment, in food and in biological samples. "Environment" includes water, air and soil. Biological samples include blood, skin, tumors amniotic fluid, tissues, cells and cell cultures. Detection of sequences in nucleic acids is used to identify microorganisms in a sample, to diagnose genetic defects or polymorphisms, to detect gene expression and for forensic studies.

25       Means for detecting a pattern generated by signals from hybridization within individual gel elements in a microchip of the present invention include a laser scanner, e.g. a laser "gun" such as used to scan bar codes, a CCD camera coupled to a fluorescent microscope. In the field, the naked eye or a scanner is used.

The invention relates to a deliberate and informative arrangement of  
30 oligonucleotides immobilized on a microchip, such that upon hybridization with oligonucleotides in a test sample, a pattern is produced that can be interpreted with a suitable means. Hybridization may be detected by letter (FIG. 12), design or bar code

pattern (FIG. 13) wherein columns "1" and "3" are dark bars signifying the presence of a pathogen, and the specific pathogen in "1" is a different anthrax species from that in "3." By immobilizing all of one type of oligonucleotide in a column, for example, the pattern is readily detected as a linear column, as contrasted to detecting

- 5 hybridization in a single small gel element or elements, which requires a microscope to detect it, and computer programs to analyze it.

A nucleic acid hybridization is a highly specific and sensitive procedure and allows a specific sequence to be detected and identified among other millions of sequences in a genome of an organism. However, nucleic acid hybridization is a  
10 useful but quite a cumbersome procedure. This drawback can be overcome by using oligonucleotide microchips as biosensors for different microorganisms. Within a small area of a few square millimeters or centimeters, hundreds and thousands of synthetic oligonucleotide probes are immobilized that are specific to ribosomal DNA or to other specific nucleic acids. Subsequent hybridization of a DNA or RNA  
15 molecule to the microchip enables a menu of oligonucleotides to be identified in a sample. Instead of having to interpret hundreds or thousands of individual hybridizations, a relative simple pattern produced by hybridizations is analyzed.

For bacterial assays, pure culture microorganisms, purified target nucleic acid or even synthetic oligonucleotides are useful as internal standards, serving to estimate  
20 the efficiency of nucleic acid isolation or the absolute amount of target nucleic acid recovered.

The customized oligonucleotide microchips are produced by chemical immobilization of presynthesized oligonucleotides, or by direct synthesis of oligonucleotides on a microchip. If a microchip contains rather long oligonucleotides,  
25 the former methods are the methods of choice because before immobilization, the oligonucleotides are purified and checked for their quality.

Methods and technologies have been developed for microchip manufacturing, hybridization of fluorescently labeled DNA and RNA with the microchips and monitoring the hybridization with a fluorescence microscope equipped with CCD-  
30 camera, computer and proper software (see U.S. Pat. No. 5,552,270 herein incorporated by reference).



The oligonucleotide microchips consist of many polyacrylamide gel pad elements generally of the size of 40x40x20  $\mu\text{m}$  and larger. The elements are chemically fixed on a glass surface. Each microchip gel element contains a specific presynthesized oligonucleotide that is immobilized through a covalent bond.

5 Hundreds of microchips containing hundreds and thousands of different immobilized oligonucleotides can be manufactured by a specially devised robot. The gel array also offers several advantages over formats using an *in situ* synthesis of the oligonucleotide array. The synthetic oligonucleotides are purified by gel electrophoresis or HPLC prior to immobilization on the microchip. This provides for  
10 stringent quality control of oligonucleotide purity and insures high specificity. The polyacrylamide gel support has a capacity of immobilized oligonucleotides from 0.03 pmol up to 10 pmol per 100x100x20  $\mu\text{m}$  gel pad. This offers improved quantification and better discrimination between perfect and mismatched duplexes. It also provides a way to normalize differences in hybridization signal intensities.

15 Oligonucleotide microchip technology for sequencing by hybridization is available to identify the presence of microorganisms in a sample of any type, or to find new species. As shown in the examples herein, the hybridization of DNA or RNA extracted from even a very complicated mixture to a specific oligonucleotide probe has resulted in unambiguous identification of microorganisms. The nucleotide  
20 sequence of the microorganisms for genes encoding a small subunit of ribosomal 16S rRNA forms the basis for a microchip biosensor. Instead of direct sequencing of the gene, hybridization analysis of DNA or RNA samples with oligonucleotides specific for the microorganisms is performed. This new technology provides efficient microbial analysis and environmental monitoring. Fluorescently labeled DNA and  
25 RNA samples from microorganisms are hybridized with microchips containing oligonucleotides specific for several microorganisms. These microorganisms are reliably identified by microchip hybridization patterns. Microorganism biosensor technology is developed, reusable customized microbial oligonucleotide microchips are produced by methods of the present invention, and methods are developed for  
30 simultaneous quantitative and qualitative microchip analysis of hundreds and thousands of microorganisms in a sample and for discovery of new ones.

## EXAMPLES

The following examples are presented as illustrations of aspects of the invention, rather than limitations on the invention. Other applications include detection of genetic mutations such as are characteristic of hemoglobin disorders; detection of genetic polymorphisms such as HLA; investigation of gene expression; detection of causative agents of diseases; forensic studies; and detection of microbial pollutants.

### **Example 1: Preparation of an Oligonucleotide Microchip Biosensor**

Oligonucleotides are synthesized using a 394 DNA/RNA synthesizer (Applied Biosystems). The synthesis of oligonucleotides for immobilization began with 3-methyluridine at the 3'-terminal position.

In one embodiment, fluorescently labeled RNA was prepared using T7 RNA polymerase. Template DNA (133 and 75 bp long) for *in vitro* transcription was prepared by PCR amplification with the nested primers T7V2L-45, 5'-GGAATTCCTAATACGACTCACTATAGGGACCACC-ATGGTGCACCTGACTCC-3', as well as with the common reverse primer T7-V2L-103 5'-GGAATTCCTAATACGACTCACTATAGGGAGGTGGAACGTGAAGTTGG-3' and 5'-TCTCCTTAAACCTGTCTTGTAACC-3'. Templates were purified using a QIAquick PCR purification kit (QIAGEN) according to the manufacturer's protocol. The RNA polymerase reaction was performed using the MEGAshortscript™ T7 kit (Ambion) with fluorescein 12-UTP (Molecular Probes). Fluorescently labeled ssDNA (single stranded DNA) fragments were prepared by single primer reamplification.

A polyacrylamide gel micromatrix was prepared by photopolymerization of a solution of 4% acrylamide (acrylamide/bisacrylamide 19/1), 40% glycerol, 0.0002% methylene blue, and 0.012% TEMED in 0.1 M sodium-phosphate buffer, pH 7.0. The mixture was applied to an assembled polymerization chamber illuminated with U.V. light.

Two types of microchip matrices (micromatrices) were routinely prepared with gel pad elements of about 60x60x20 µm and 100x100x20 µm that were spaced by 120 and 220 µm, respectively. About 1 nl of activated oligonucleotide solution was transferred to a gel element using either a robot or a simple manual device.

The device includes a Peltier thermostated pin placed under a binocular lens in conjunction with a micromanipulated holder, a power supply, and a refrigerated circulator.

5 The manufacture of microchips of gel-immobilized oligonucleotides basically consists of three steps; shaping the desired topology of oligonucleotides on a gel micromatrix; loading microvolumes of oligonucleotide solutions onto the micromatrix, and immobilizing within the gel oligonucleotides containing the active 3' or 5' terminal aldehyde or amine groups.

10 To avoid the exchange of different oligonucleotide solutions applied on adjacent gel pads, the pads are separated on the micromatrix by a hydrophobic glass surface. Two-dimensional scribing or laser evaporation is used for micromatrix preparation, but these procedures require rather complex equipment and experienced personnel. The photopolymerization method significantly simplifies the procedure and makes it accessible to a biochemical laboratory.

15 Microfabrication by mask-directed photopolymerization (e.g., a photoresist method in microelectronics) is a well developed technique. From several acrylamide photopolymerization techniques tested, modified - methylene - blue induced photopolymerization produced the best results for micromatrix manufacture. The gel matrix consists of gel pads photopolymerized on a glass slide. The gel pads are  
20 formed according to the mask topology due to the lack of photopolymerization in places covered by a nontransparent grid.

The microchip is manufactured by applying the activated oligonucleotide solutions onto the micromatrix of gel elements containing active hydrazide or aldehyde groups. A simple device exists for manual loading of up to 100 different  
25 oligonucleotides on a micromatrix. The transfer is carried out by the hydrophilic upper surface of a pin that is first immersed into, and is wetted with, an oligonucleotide solution, and then is withdrawn from the solution and brought into contact with the gel surface. This transfers about 1 nl of oligonucleotide solution with a reproducibility of  $\pm 10\%$ . The temperature of the pin is maintained near the dew  
30 point of the ambient air to avoid the evaporation of this microvolume solution in the course of transfer.

The oligonucleotides are positioned according to a design wherein hybridization pattern data will be reduced to a readily interpretable pattern.

**Example 2: The Hybridization of Microchips with DNA and RNA using a Hybridization Buffer.**

5

Fluorescently labeled DNA or RNA (5  $\mu$ l, 0.1-1 pmol/ $\mu$ l) were hybridized to a microchip at +5°C in a hybridization buffer containing 1 M NaCl, 1 mM EDTA, 1% Tween-20, and 10 mM sodium phosphate at a pH of 7.0, for between about 2-24 h. The microchip was covered with a cover glass or a Teflon sheet so that a 300- $\mu$ m space is above. Then the hybridization solution containing DNA or RNA fragments was substituted with 10  $\mu$ l of cooled hybridization buffer. The microchip with the cover glass was placed on a thermostabilized table. Hybridization was monitored quantitatively using a specially constructed multicolor epifluorescent microscope with a 4x4 mm observation field equipped with a CCD camera and suitable software.

15 **Example 3: Analysis of Melting Curves; a Hybridization Buffer is Not Required.**

The polyacrylamide gel used on a microchip provides more than 100 times higher capacity for three-dimensional immobilization of oligonucleotides than does a two-dimensional glass surface. The high concentration of immobilized oligonucleotides facilitates the discrimination of mismatched duplexes and enhances the sensitivity of measurements on the microchips. This allows the use of a CCD-camera-equipped fluorescence microscope (Yershov *et al.*, 1996) although it is less sensitive than laser scanning systems (Lipshutz *et al.*, 1995), but offers the advantage of monitoring the hybridization on a microchip at different temperatures in real time for measurement of the melting curves. Melting curves are defined herein as produced by plotting the amount of duplexes [fluorescent intensity] versus temperature. The procedure, the software, and the hybridization microchamber (Yershov *et al.*, 1996) have all been developed for recording melting curves at a wide range of temperatures simultaneously for perfect and mismatched duplexes formed upon hybridization of a probe with all microchip oligonucleotides.

30 A significant amount of time is needed for the microchips hybridized with rather long RNA or DNA probes to achieve equilibrium. Therefore, non-equilibrium dissociation melting curves were measured. However, they are not far away from equilibrium where some difference in heating rate did not significantly affect the

results. The melting curves for hybridization of, for example, synthetic 19-mers with the microchip oligonucleotides reached equilibrium under the same conditions that were used for measuring non-equilibrium RNA and DNA melting curves. The melting curves can also be measured after a few minutes, far away from equilibrium, if an internal standard is added to a tested sample. This standard can be a differently labeled RNA of a normal allele. This significantly speeds up the identification of nucleic acid base changes.

**Example 4: Choice of Optimum Melting Temperatures for Non-Equilibrium Hybridization.**

This invention embodies an improvement in the SHOM technology in which hybridizations between an array of gel-immobilized nucleotides (a microchip) and the unknown nucleotides to be tested are measured at optimal, discriminatory melting temperatures. This improvement is achieved by parallel measuring of the melting curves of the duplexes formed by hybridization on the entire oligonucleotide array, as well as by monitoring the simultaneous hybridization of two samples of nucleotides labeled with different fluorochromes, and judicious choice of proper allele-specific oligonucleotides as the immobilized probes. The fluorochromes chosen for the labeling emit light of sufficiently differing wavelengths, that both types of labels can be measured in the same reaction mixture.

The greatest discrimination between perfect and mismatched duplexes was achieved at a temperature at which the intensity of the hybridization signal from a perfect duplex dropped to one-tenth of its initial value; at such a temperature, the hybridization intensities from mismatched duplexes usually approached the background level. The temperature at which the initial signal of hybridization drops by a factor of 10 is termed the discrimination temperature (Td.).

In the case of beta-thalassemia mutation detections described in Example 6 herein: (1) RNA transcripts of PCR-amplified DNA were hybridized with immobilized oligonucleotides; (2) the Td values for perfect 40% and 70% GC-rich duplexes were 52° and 64°, respectively; (curves 1 and 3 in FIGS. 1A and 1B); (3) the immobilized oligonucleotides were chosen from among a set of overlapping sequences; and (4) the two samples included in the reaction mixture were a mutated

allele RNA labeled with one fluorochrome and a sample of the normal allele RNA labeled with a different fluorochrome.

The Td is determined by hybridization with an RNA sample if an allelic DNA is available. If such DNA is unavailable, the Td can be measured from the  
5 hybridization data resulting from experiments performed with synthetic oligonucleotides corresponding to the mutated allele of interest.

The dissociation curves for perfect and mismatched duplexes are parallel at the range of about 10° (in the middle of the curves) when plotted on a semilogarithmic scale. At this 10°C range, the ratios of the signals for perfect and mismatched  
10 duplexes remain rather constant. This makes the discrimination procedure robust to some inaccuracies in determining Td. The discrimination temperature depends on experimental conditions (rate of heating, ionic strength, probe concentration, extent of fragmentation, and so forth) which can vary from one experiment to another. However, these variations affect Td and the relative intensities of the hybridization  
15 signals to a similar extent for all microchip elements and therefore do not significantly distort the discriminations. Therefore, to provide a reference Td, the oligonucleotides CD26(N) and CD 26 G/A, which form perfect and mismatched duplexes, respectively, with all RNAs tested, were introduced into the microchip.

Since Td is robust to some inaccuracy in measurements, 19-mer  
20 oligodeoxynucleotides were used in these experiments instead of more expensive 19-mer oligoribonucleotides. There are differences in the stability of DNA-DNA homoduplexes relative to DNA-RNA heteroduplexes (Lesnik and Freier, 1995). The pattern of hybridization of the microchip with RNA derived from patients and with 19-mers was rather similar to that from the 10-mers. Hybridization with  
25 corresponding synthetic oligonucleotides is preferred as a control when a mutation is identified in an RNA sample by its hybridization with a diagnostic microchip.

A mixture of fluorescently labeled RNA samples was prepared from two patients; the first sample was TMP-labeled RNA from a patient that is homozygous for the normal CD26 area of the beta-globin allele; the second sample is fluorescein-  
30 labeled RNA from a patient that is heterozygous for the normal CD26 area and a mutation CD26 G/A alleles. This mixture was hybridized with a microchip consisting of two microchip elements that contained the following immobilized oligonucleotides:

SEQUENCE

A sample (*element*)-CD26(N-normal) 5'-GGCCTCACCA-3MeU-3'

B sample (*element*)-CD26(G/A-mutant) 5'-GGCCTTACCA-3MeU-3'

- 5 Usage of different filters during the registration of the signal, allowed the independent, simultaneous registration of the sample, which was marked with the different dyes; TMP (red) and fluorescein (green), on the same element of the microchip. FIG. 2 demonstrates the interaction of sample 1 with the A microchip element; Graph 2 demonstrates the interaction of sample 2 with the A microchip  
10 element; Graph 3 demonstrates the interaction of sample 2 with the B microchip element; and Graph 4 demonstrates the interaction of sample 1 interaction with the B microchip element.

**Example 5: Use of a Customized Microchip Matrix Biosensor to Identify Nitrifying Microorganisms.**

- 15 The results in this example were obtained using methods previously available, not the non-equilibrium melting curves. Microorganisms that degrade nitroaromatic compounds include Pseudonomas, Arthrobacter, Nocarida, Myco-bacterium, and fungi.

- 20 Previously, methods for detection of these bacteria were tedious and inaccurate. For example, to detect Pseudonomas capable of degrading nitroaromatic compounds, 2-nitroluene was tested as a sole carbon, energy and nitrogen source. It was difficult to isolate the bacteria from soil samples to perform the test.

- 25 Nitrifying bacteria have proved particularly difficult to study using cultivation techniques, such as most probable number (MPN) and selective plating because of their long generation times and poor counting efficiencies. Thus, a rapid and non-culture dependent enumeration technique for nitrifiers could greatly facilitate research in their ecology.

- 30 Microchips with 100x100x20 µm gel pads (alternatively 60x60x20 µm), fixed on a glass surface and containing a set of 10 oligonucleotides 15-20 bases-long were manufactured for bacterial typing experiments. The set included oligonucleotides complementary to different regions of 16S ribosomal RNA. Since rRNA's are naturally amplified, and often are present in thousands of copies per cell, they provide

great sensitivity and eliminate the need for amplification in many applications. One oligonucleotide is represented in most living organisms, another is typical for most of bacteria and the rest belong to nitrosos (nitrifying) bacteria only. The group of nitrosos bacteria oligonucleotides consists of two oligonucleotides typical of nitrobacter, two typical of nitrosomonas and one typical of nitrosovibrio. One oligonucleotide was complementary to an antisense strand of rDNA for hybridization with ribosomal dsDNA, that was PCR amplified from genomic or cDNA.

The following scheme for an ordered oligonucleotide loading (placing on a chip) is useful for bacterial (or organism, species) typing. In the micromatrix design shown in Table 1, the first oligonucleotides characterize the highest order (i.e. to distinguish a living organism). [Uni 1390-CIII]. Reducing the order step by step down to the lowest level, i.e. from family, to genus, to species provides further discrimination of oligonucleotides that are present in a sample being investigated. For example, for oligonucleotides used to classify nitrifying bacteria, a bacterial oligonucleotide would be in the next position. [Bac 338 - CI and NonBac338 - CII]. Oligonucleotides specific to nitrobacter [Nb1000 - AI and NIT3 - AII] and ammonia oxidizers [NEU32 or 23? - AIII, Nso190 - AIV and Nso1225 - BI] follow in any order. Finally, oligonucleotides specific to Nitrosomonas [Nsm156 - BII] and Nitrosovibrio [Nsv443 - BIII] complete the micromatrix design.

The microchip was evaluated using three different rRNA preparations (phenol extracts of cellular RNA, RNA isolated from purified ribosomes, and *in vitro* transcripts of cloned ribosomal DNA), and both fragmented double-stranded and single-stranded DNA. Hybridizations were performed in a formamide buffer at low temperature in order to enhance microchip durability and decrease RNA degradation. Although all DNA and RNA preparations could be used, the best discrimination was observed for *in vitro* transcribed rRNAs using the hybridization conditions evaluated in this study.

The hybridization of the microbial microchips was carried out with five different preparations of target nucleic acids. Ribosomal RNA and total RNA were recovered from cells. RNA transcribed *in vitro* as well as single- and double-stranded PCR-amplified 16S rDNA were obtained from plasmids containing the cloned 16S rRNA gene. All of these sample types provided a comparatively reliable



identification of the microorganisms by their hybridization with the microchip-immobilized oligonucleotides and could be used for different purposes. For example, the rRNA provides a naturally amplified target. Also, since cellular ribosome content is well known to vary with growth rate, it is generally thought that direct

5 quantification of rRNA serves to identify the more active environmental populations. In contrast, analysis of PCR amplified rDNA provides a more general measure of all microorganisms present in a sample. Alternatively, these measures could be combined. For example, the RNA and DNA components of an environmental sample could be isolated and labelled with different fluorescent dyes. Following their

10 combined hybridization, the resulting ratio of RNA and DNA hybridizing to an individual gel element could be used to infer the physiological status of the corresponding microbial population.

Table 2 shows the sequences of the oligonucleotides and other characteristics of them.

15

TABLE 1

Micromatrix Design for Nitrifying Microorganisms.

20

	<b>I</b>	<b>II</b>	<b>III</b>	<b>IV</b>
A	Nb1000	NIT3		Nso190
B	Nso1225	Nsm156	Nsv443	
C	Bac338	NonBac338	Uni1390	

TABLE 2

Oligo-nucleotide Name and Position	Sequence (5' to 3')	Specificity	Microchip location Table 1	Td <sup>1</sup> C
Nb1000	5'-tgc gac cgg tca tgg-3'	<i>Nitrobacter</i>	A-I	42°
NIT3	5'-cct gtg ctc cat gct ccg-3'	<i>Nitrobacter</i>	A-II	66° <sup>2</sup>
NEU23	5'-ccc ctc tgc tgc act cta-3'	Ammonia oxidizers	A-III	66° <sup>2</sup>
NSO190	5'-cga tcc cct gct ttt ctc- 3'	Ammonia oxidizers	A-IV	62°
NSO1225	5'-cgc gat tgt att acg tgt ga-3'	Ammonia oxidizers	B-I	51°
NSMO156	5'-tat tag cac atc ttt cga t-3' ?	<i>Nitrosomonas</i>	B-II	46°
NSV443	5'-cgg tga cgg ttt cgt tcc-3'	<i>Nitro-sospira</i> -like	B-III	52°
BAC338	5'-gct gcc tcc cgt agg gat-3'	Bacteria	C-I	54°
NonBAC338	5'-act cct acg gga ggc agc-3'	Eub338 complementary strand	C-II	54°
UNI1390	5'gac ggg cgg tgt gta caa-3'	all life (with a few exceptions)	C-III	44°

<sup>1</sup> Experimentally determined.<sup>2</sup> Estimated from *in situ* hybridization.

A number of hybridization conditions were tested in terms of efficiency and specificity of hybridization. Hybridization in formamide containing buffer at low temperature gave good results. Hybridizations were performed at 5° centigrade in 33% formamide. RNA samples and covalent bonding of oligonucleotides with the support (hence durability of microchips) are more stable at low temperatures. In addition, these conditions were favorable from a point of view of RNA stability and microchip durability similar to other RNA molecules at low temperatures of about 0° - 5°C.

The hybridization on a microbial microchip was carried out with *in vitro* RNA transcripts of 16S rDNA of different nitroso bacteria, total RNA extracts and ribosomal RNA extracted from *E. coli* and *Desulfovibrio vulgurus* as well as PCR amplified double or single stranded DNA of 16S rDNA.

The probes for ammonia oxidizing bacteria show different discrimination specificity under different conditions. FIG. 3 shows the fluorescence of individual gel elements on the microchip following hybridization to the 16S rRNAs of *Nitrosovibrio tenuis* (A), *Nitrosomonas europaea* (B), and *E. coli*, either *in vitro* transcribed (C) or recovered from isolated ribosomes (D). The same microchip was used for each hybridization following washing with distilled water. Each microchip was routinely used for up to 20-30 hybridization experiments. The appropriate pattern of hybridization was observed for all gel elements shown, despite a significant difference in dissociation temperatures ( $T_{ds}$ ) previously determined using membrane support hybridization (Table 1). For hybrids of comparable stability, discrimination is generally achieved by washing at increasing temperatures (described below) or by simultaneously evaluating their melting characteristics, since the fluorescence analyzer can monitor hybridization signals in real time.

FIGURE 4A and 4B shows the results of an experiment evaluating the effect of increasing washing temperature on target RNA retention. A mixture of *Nitrosovibrio tenuis* and *E. coli* 16S rRNA labelled with different fluorescent dyes (fluorescein and tetramethylrhodamine, respectively) was hybridized to the chip at 5°C. The hybridization solution was then replaced with washing buffer and the retention of each RNA species was measured following each 10°C incremental increase in temperature

(up to 60°C) using multicolor detection. Nonspecific hybridization of *E. coli* rRNA to Nso1225 (ammonia oxidizer), Nsm156 (nitrosomonas), and NonBac338 (anti-sense) was observed following the 10°C wash. However, this nonspecific hybridization was significantly reduced following the 40°C wash. In like manner, the 16S rRNA of *Nitrosovibrio tenuis* hybridized to *Nitrosomonas* (Nsm156) at 10°C, but was reduced to near background (compared to NonBac338) following the 40°C wash. Using the methods of the present invention, hybridization buffer is not required. A more complete correction for differences in stabilities of duplexes can be carried out by measuring the equilibrium or non-equilibrium melting curves for all microchip elements. This would provide a basis to compensate for the various factors influencing individual duplex stability, e.g., their length, GC-content, and competition with secondary and tertiary structures in RNA and DNA.

FIG. 4B shows the ratios of hybridization intensities of fluorescein labelled *Nitrosovibrio tenuis* to tetramethylrhodamine labelled *E. coli*. with different microchip oligonucleotides at 10°C and 40°C (the ratios are derived from the data presented on Figure 4A. These ratios were not changed significantly for oligonucleotides specific to bacteria and all living organisms between 10°C and for more stringent conditions at 40°C. However, the ratio is dramatically increased at 40°C (compared to 10°C) for oligonucleotides specific to ammonia oxidizers and nitrosovibrio. This increase reflects the greater duplex stability of *Nitrosovibrio tenuis* RNA with the complementary oligonucleotides compared with *E. coli*. RNA. Although the nitrosomonas ratio increases, the signal originating from each labelled RNA is near background. This experiment demonstrates that the inclusion of second dye-labelled RNA, either isolated from cells or synthesized, could be used as an internal standard for quantitative assessments of hybridization patterns.

Variable hybridization to the different gel elements is the expected consequence of using a single hybridization condition to evaluate an array of probes, each having different kinetics of association and dissociation. To some extent these difference can be normalized by varying the concentration of oligonucleotides in the individual gel elements. For example, the relatively low hybridization signals of Nso1225(b-I) and Uni1390 (c-III) compared to Nsv443 (b-III) could each be elevated by increasing the amount of the corresponding oligonucleotides probes immobilized in the gel. This

approach was evaluated by synthesizing a microchip with selected probes immobilized at several different concentrations, up to 6 times higher than that used in the experiments previously described. This was accomplished by multiple applications of the standard loading solution (100 pmol/ $\mu$ l probe) to each gel element.

- 5 Comparable hybridization of Nso1225 (ammonia oxidizer) and Nsv443 (nitrosovibrio-like) was achieved following three applications of the Nso1225 probe (FIG. 5). Similarly, two applications of Bac338 (bacteria) and five of Uni1390 (all life) resulted in hybridization comparable to Nsv443.

- Strains used.** *Escherichia coli*, *Desulfovibrio vulgaris* strain PT2,  
10 *Nitrosovibrio tenuis* strain NV12, *Nitrosomonas europaea* strain ATCC 19718, and *Nitrosomonas* strain C-56 were used as sources of nucleic acid for these experiments.

- RNA preparation.** Total cellular RNA was isolated by phenol/chloroform extraction. For some of the samples, a ribosome enrichment was performed before RNA extraction. Forty ml of log phase growth *E. coli* or *D. vulgaris* strain PT2 was  
15 centrifuged at 3500 g for 10 min. and resuspended in 4 ml of 4°C ribosome buffer. Ribosome enrichment buffer consisted of 20 mM MgCl<sub>2</sub>, 50 mM KCl, 50 mM Tris at pH 7.5, and 5 mM  $\beta$ -mercaptoethanol in diethyl pyrocarbonate treated double-distilled water. The cell suspension was divided between 4 screwtop microfuge tubes and 0.5 g of 0.1 mm ZrO<sub>2</sub> beads were added. The cell suspensions were disrupted for 2 min.,  
20 put on ice for 5 min., and disrupted again for 2 min. The cell suspensions were centrifuged at 14000 g for 10 minutes. The supernatant, which contained the ribosomes, was recovered and transferred to ultracentrifuge tubes. Ribosomes were pelleted by ultracentrifugation in ribosome buffer at 55,000 rpm (201,000 g average) for 50 minutes in a Beckman Optima Series TL swinging bucket rotor (Beckman,  
25 Fullerton, CA), for a Svedberg sedimentation factor of 70S. After centrifugation, the supernatant was discarded and the RNA was recovered from the pelleted ribosomes by extraction with pH 5.1 phenol/ chloroform. Quality and quantity of extracted RNA was evaluated by polyacrylamide gel electrophoresis and ethidium bromide staining.

- Cloning of 16S rDNA and *in vitro* production of RNA transcripts.** DNA  
30 was extracted from *E. coli*, *Desulfovibrio vulgaris* PT2, *Nitrosovibrio tenuis* NV12, *Nitrosomonas europaea* 19718, and *Nitrosomonas* strain C-56 cell pastes using a guanidine/diatom method. Near-complete 16S rDNA genes (ca. 1500 base pairs)

were recovered from each by PCR amplification using S-D-Bact-0011-a-S-17 (GTTTGATCCTGGCTCAG) and S-D-Bact-1492-a-A-21 (ACGGYTACCTTGTTACGACTT) as primers and a premixed PCR amplification buffer (Pharmacia Biotech Inc. Piscataway, NJ), consisting of 0.2 mM Mg<sup>++</sup>, 2.5 mM each dATP, dCTP, dGTP, dTTP, 0.2 mM of each amplification primer, and 2.5 units of Taq DNA polymerase (Pharmacia). Temperature cycling was done in an Idaho Technology thermocycler (Idaho Falls, ID) using 30 cycles of 15 sec at 94°C, 20 sec at 50°C, and 30 sec at 72°C. The PCR-products were cloned in a pCR plasmid (Invitrogen, San Diego, CA) according to manufacturers instructions. Plasmids were isolated using the Wizard kit (Promega, Madison, WI) and used for *in vitro* transcription of the cloned SSU rRNA genes.

**DNA oligonucleotide probes.** All probes were complementary to the SSU rRNAs and previously characterized using a membrane hybridization format. Five probes hybridize to different groups of ammonia-oxidizing bacteria within the beta-subdivision of the *Proteobacteria*. S-G-Nso-190-b-A-19 (Nso190) and S-G-Nso-1225-a-A-20 (Nso1225) encompass all sequenced ammonia-oxidizers of the beta-subclass of *Proteobacteria*, probe S-G-Nsm-156-a-A-19 (Nsm156) identifies members of the genus *Nitrosomonas* (also including *Nitrosococcus mobilis*), probe S-G-Nsv-443-a-A-20 (Nsv443) is specific for the *Nitrosovibrio/Nitrosolobus/Nitrospira* group, and probe S-G-Nsm-653-a-A-18 (NEU23) is specific for the halotolerant members of *Nitrosomonas*. Probes for members of genus *Nitrobacter* (nitrite oxidizing) were S-G-Nit-1000-b-A-15 (Nb1000) and S-G-Nit-1035-a-A-18 (NIT3). Other probes used were S-D-Bact-0338-a-A-18 (Bac338) which hybridizes to members of the bacterial domain; S-D-NBac-0338-a-S-18 (NonBac338), complementary to the antisense strand of the Bac338, and S-\*-Univ-1390-a-A-18 (Uni1390) complementary to the SSU RNA of nearly all characterized living organisms, with the exception of some protists.

**RNA and DNA labeling and fragmentation.** Single stranded DNA was prepared by asymmetric PCR according to Ausubel *et al.* (1994) using a 100 times excess of the forward primer. Briefly, DNA was partially depurinated in 80% formic acid for 30 min. at 20°C, then incubated in 0.5 M ethylenediamine hydrochloride (pH 7.4) for 3 hr at 37°C, followed by 30 min. at 37°C in the presence of 0.1 M NaBH<sub>4</sub>.

Fluorescein isothiocyanate was incorporated into fragmented DNA by incubation in absolute DMSO at room temperature for 1 hr.

RNA was fragmented by base hydrolysis and dephosphorylated with bovine phosphatase. Fragmented RNA was oxidized by NaIO<sub>4</sub> and labeled either by  
5 ethylenediamine mediated coupling of 6-carboxyfluorescein (FAM) succinamide or by direct incorporation of tetramethylrhodamine-hydrazide (TMR).

**Microchip fabrication.** A matrix of glass-immobilized gel elements measuring 60 x 60 x 20 or 100 x 100 x 20  $\mu$ m each and spaced apart by 120 or 200  $\mu$ m respectively was prepared. The polyacrylamide gel was activated by substitution of  
10 some amide groups with hydrazide groups by hydrazine-hydrate treatment. Oligonucleotides were activated by oxidizing 3'-terminal 3-methyluridine using NaIO<sub>4</sub> to produce dialdehyde groups for coupling with hydrazide groups of the gel and coupled to each micromatrix element by applying 0.5-1 nl of the activated oligonucleotide solution (100 pmol/ $\mu$ l) using a specially devised robot.

15 **Hybridization and image analysis.** Probe binding was quantified by measuring the fluorescence conferred by the binding of fluorescently labeled DNA or RNA (tetramethyl rhodamine or fluorescein) to the individual gel elements. Hybridization and washing was controlled and monitored using a Peltier thermotable (with a working range of -5.0°C to +60.0°C) mounted on the stage of a custom-made  
20 epifluorescent microscope. The microchip was hybridized at 5°C, either overnight or for 6 hr, in 2-5  $\mu$ l of the hybridization buffer [33% formamide, 0.9 M NaCl, 1 mM EDTA, 1% Tween-20, and 50 mM sodium phosphate (pH 7.0)] at a concentration of DNA and RNA between 0.2-2 pmol/ $\mu$ l. The hybridization mixture was replaced with 5-10  $\mu$ l hybridization buffer without formamide immediately prior to microscopic  
25 observation. Exposures were in the range of 0.1-10 sec depending on the signal intensity, but were typically around 1 sec. Fluorescence was monitored either at room temperature or using a range of temperatures between 5 - 60°C.

Conditions for the coupling of micromolecules to the acrylamide gel were devised to rule out the possibility of liquid evaporation during immobilization and to  
30 ensure that covalent bonding of oligonucleotides with the gel matrix proceeds to completion. After the microvolumes of the oligonucleotide solutions have been applied to all cells of the matrix, the micromatrix gel elements were swelled by

condensing water from the ambient air. Then the micromatrix surface was covered with a thin layer of an inert nonluminescent oil, and chemical coupling of the activated oligonucleotides to the activated polyacrylamide was carried out to completion.

#### 5 **Example 6: Use of Microchip Biosensors As Diagnostic Assays**

The microchip technology was successfully tested for identification of single base changes in genomic DNA and RNA for reliable diagnosis of human genetic diseases. A customized microchip contained oligonucleotides specific to  $\beta$ -thalassemia normal and abnormal  $\beta$ -globin genes. The hybridization with PCR-  
10 amplified DNA or RNA samples derived from genomic DNA of subjects allowed unambiguous identification of a mutation in a sample to be tested. Reliability of the identification was enhanced by using simultaneous hybridization with two samples of a normal and mutated RNA stained with different fluorescence dyes and monitoring the hybridization at different wavelengths; by simultaneously measuring the melting  
15 curve for duplexes formed on a microchip, and by using a proper set of several oligonucleotides complementary to the mutated site of the DNA.

A number of the most commonly occurring  $\beta$ -thalassemia mutations with  $\beta$ -globin gene were used in diagnostic assays with oligonucleotide microchip biosensors. These mutations were splice-site mutations for the 1<sup>st</sup>, 2<sup>nd</sup>, 5<sup>th</sup>, and 6<sup>th</sup> nucleotides in  
20 the first intron (IVS I) of the  $\beta$ -globin gene: IVS I/1 G/A (G/A = substitution of G for A), IVS I/2 T/C, IVS I/5 G/T, IVS I/5 G/C, IVS I/6 T/C, and G/A substitution in the 26<sup>th</sup> codon (GAG) of the first exon (FIG. 6), (also known as abnormal hemoglobin E) (see Diaz-Chico *et al.*, 1988 for terminology).

A microchip with 100H100H20  $\mu$ m gel elements (Yershov *et al.*, 1996)  
25 contained immobilized decadeoxyribonucleotides, that is, 10-mers that correspond to normal and mutant  $\beta$ -thalassemia alleles. These 10-mers discriminated mismatches less reliably than 8-mers, but were hybridized more efficiently than 8-mers. 10-mers were, therefore, preferred for this assay. Table 3 shows the sequences of the allele-specific oligonucleotides immobilized on the microchips. It was expected that  
30 mismatches within the duplexes would have a much higher destabilization effect than mismatches at the terminal positions (Khrapko *et al.*, 1991); therefore the mutated bases were placed inside of the immobilized oligonucleotides.



Single- and double-stranded PCR-amplified  $\beta$ -globin DNA fragments of different lengths and collected after a random fragmentation were tested in assays for identification of some of these mutations. However, the hybridization of RNA is preferred over DNA hybridization. RNA fragments were derived from PCR-amplified genomic DNA by transcription with T7 RNA polymerase (Lipshutz *et al.*, 1995). About 100 copies of unlabeled or fluorescently labeled RNA transcripts are synthesized per DNA molecule, providing a convenient way to prepare a sufficient amount of the hybridization probes. RNA is fragmented and one fluorescent dye molecule is introduced per fragment.

Table 3 shows the sequences of the microchip allele specific 10-mers. The oligonucleotides of microchip I are complementary to the coding strand of DNA of the  $\beta$ -globin gene of patients with  $\beta$ -thalassemia single-base mutations (G/A - substitution of A for G) in the 1<sup>st</sup>, 2<sup>nd</sup>, 5<sup>th</sup>, or 6<sup>th</sup> nucleotides of the first intron (IVS I/1, 2, 5, 6) of the  $\beta$ -globin gene and in the codon #-26 (CD-26) of the first exon. Oligonucleotides 1-16 of microchip II correspond to the normal and IVS I/2 G/T allele. The mutated and corresponding normal bases are placed from the 2<sup>nd</sup> to the 9<sup>th</sup> positions of the 10-mers from their 3'-end. The mutated bases are shown in lowercase bold letters and corresponding oligonucleotide bases in the normal allele are underscored. The oligonucleotide synthesis and the microchip manufacturing were described by Yershov *et al.* (1996).

Microchip I was successively hybridized with RNA 75 and 133 nt long without fragmentation or after fragmentation (133~~nt~~, Table 5, probes 3a and 4a) and with 6 synthetic 19-mer oligodeoxyribonucleotides corresponding to  $\beta$ -thalassemia mutations. The RNA and 19-mers were labeled with TMR except for RNA probes 2a, 2b, and 6b, which were labeled with fluorescein (F1). The melting curves (FIG. 1A-B, FIG. 2) were measured simultaneously for all microchip oligonucleotides at each hybridization. These curves provided values of hybridization intensities at the discrimination temperature, T<sub>d</sub>. R is the ratio of the hybridization signal of a mismatched duplex (I<sub>m</sub>) to the signal of the perfect duplex (I<sub>p</sub>) estimated at T<sub>d</sub> in parallel for all microchip oligonucleotides.  $R = I_m/I_p$ . d<sub>19</sub>-synthetic 19-deoxymers were complementary to allele specific 10-mers immobilized on the microchips.

Table 4 shows the effect of the position of the allelic base within 10-mers on mutation detection. Microchip II contains two sets of 10-mers corresponding to the normal and IVS I/2 T/G alleles. The microchip was hybridized with the TMR-labeled normal allele 19-mer and to an RNA 75 nt long.  $T_{0.1}$  is the temperature at which the hybridization signals for a microchip duplex drops to 1/10 of its initial value at 0°C. -  
 $\Delta T_{0.1} = T_{0.1}$  (a perfect duplex) minus  $T_{0.1}$  (the corresponding mismatched duplex.)

Fluorescently labeled RNA probes were prepared from a fragment of the  $\beta$ -globin gene from the first exon (Lawn *et al.*, 1980). PCR amplification of a 1.76-kb fragment of the human  $\beta$ -globin gene mapped from nucleotides -47 to +1714 (Lawn *et al.*, 1980) was carried out with 1  $\mu$ g genomic of DNA (Poncz *et al.*, 1982) and 50 pmol each of the forward primer: 5'-GGAGCCAGGGCTGGGCATAAAAGT-3' (-47->-23) and the reverse primer 5'-ATTTTCCCAAGGTTTGAAGTAGCTC-3' (+1689->+1714). (FIG. 7) The amplification was carried out in a DNA thermal cycler (GeneAmp PCR System 2400, Perkin Elmer Corporation) in 100  $\mu$ l of a buffer containing 200 mM each of dATP, dCTP, dGTP, dTTP, 2.5 mM  $MgCl_2$ , 2 units of Taq DNA polymerase (BioMaster, Russia), 50 mM KCl, 10 mM Tris-HCl, pH 9.0, and 0.1% Triton X-100. The reaction conditions were 30 cycles, with 45 sec at 95°C, 90 sec at 66°C, and 120 sec at 72°C. PCR product was purified from 2% low gel/melting temperature agarose gel (NuSieve agarose, FMC). The 159 bp and 102-bp DNA fragments were amplified with 10 ng of the 1.75 kb DNA with three nested primers, two containing T7 promoter sequence and a common reverse primer. The nested primers were T7-V2L-45. (5'-GGAATTCCTAATACGACTCACTATAGGG ACACCATGGTGCACCTGACTCC- 3'-44->+66); T7-V2L-103 (5'-GGAATTCCTA ATACGACTCACTATAGGGAGGTGAACGTGGATGAAGTTGG-3'; +102->-123); and 5'-TCTCCTTAAACCTGTCTTGTAACC-3' (common reverse; 153->+176). The amplification was carried out in 25 cycles (15 sec at 95°C, 30 sec at 62°C, and 30 sec at 72°C). PCR products were purified by QIAGEN QIAquick PCR Purification Kit. The PCR-amplified 159 or 102 bp DNA (4-5  $\mu$ g) containing T7 promoter was transcribed with 400 units of T7 RNA polymerase (Promega) to produce 133 and 75 nt long RNA in 80  $\mu$ l of buffer containing 300 mM HEPES, pH 7.6, 30 mM  $MgCl_2$ , 16  $\mu$ g of BSA, 40 mM DTT, 30 units of Rnasin (Promega) and 4 mM each of ATP, CTP, GTP, and UTP for 3 h at 38°C. Deproteinization of the reaction mixture was carried

out in 20 mM EGTA, pH 8.0, 2% SDS, and Proteinase K (10 mg/ml) for 15 min at 37°C. The mixture was extracted first with equal volumes of phenol and then with equal volumes of chloroform, precipitated twice by one volume of isopropyl alcohol, from 0.5 M LiClO<sub>4</sub> and dissolved on a Bio-Spin P6 column (BioRad).

5            Fragmentation of 10-100 µg of RNA to an average length of 20- to 40-mers was carried out in 50 µl of 0.1 M KOH for 30 min. at 40°C. Then 5 µl of 1M HEPES, pH 7.6, and 15 µl of 1% HCO<sub>4</sub> were added at 4°C. The pellet of potassium perchlorate was removed by centrifugation and RNA was precipitated by 10 volumes of 2% LiClO<sub>4</sub> in acetone. The RNA was washed twice with acetone and dried for 20-30 min.  
10          at room temperature. The fragmented RNA was dephosphorylated in 50 µl of 20 mM Tris-HCl, pH 8.0, 1 mM MgCl<sub>2</sub>, 1 mM ZnCl<sub>2</sub>, 10 units of Rnasin, 5-7 units of calf intestine phosphatase (CIP) for 1 hour at 37°C. RNA deproteinization and purification was carried out as described herein.

            For chemical fluorescence labeling of RNA the 3'-terminal dephosphorylated  
15          nucleoside was oxidized in 20 µl of 10 mM sodium periodate for 20 min. at room temperature. RNA was precipitated with acetone. A 10 molar excess of 10 mM TMR-hydrazine in 10% acetonitrile was added to oxidized RNA fragments in 20 µl of 20 mM sodium acetate at pH 4.0.

            The reaction mixture was incubated 30-40 min at 37°C, and the hydrazide bond  
20          between the RNA and dye was stabilized by reduction with freshly prepared 1.5 µl of 0.2 M NaCNBH<sub>3</sub> and incubated for 30 min. at room temperature. Then the mixture was extracted four times with water saturated *n*-butanol and precipitated with acetone. Alternatively, RNA was labeled by incorporation of fluorescein-UTP during the transcription with Ambion MEGAscript kit according to the manual.

25            The hybridization of fluorescently labeled RNA (1 pmol/µl) with the microchips was carried out at 0°C for 18 h. In many cases, the intensities of the hybridization signals at 0°C were similar for perfect and mismatched duplexes. The perfect and mismatched duplexes as well as the duplexes having various GC and AT contents displayed different stabilities and therefore were tested at different temperatures.

30            Table 4 summarizes the results of hybridization of the diagnostic microchips with 1) RNA probes derived from a number of homozygous and heterozygous β-thalassemia patients; and 2) with corresponding 19-mers. The table shows the Td

for perfect duplexes formed on each microchip oligonucleotide. The relative intensities,  $R$ , of the hybridization signals for a different microchip oligonucleotides in Table 3 are normalized to the signals for a perfect duplex at the  $T_d$  (estimated as 1.0). In most cases the ratios for mismatched duplexes are less than 0.1 and close to 0.

- 5 These values are low enough to allow unambiguous identification of the homozygous and heterozygous mutations in patients at the  $T_d$  (when the hybridization signals from only perfect duplexes are observed). The hybridization of homozygote RNA (Table 5, probes 1a, 2a, 2b, and 3a) with the microchip shows the distinctive formation of a perfect duplex only with one immobilized oligonucleotide and mismatched duplexes
- 10 with all others. Two perfect duplexes were unambiguously identified upon hybridization with a heterozygote RNA (Table 5, probe 4a).

Table 3. The sequence of the microchip allele specific 10-mers.

	#	Allele	Position of mutated base	Sequence	Location
5					
	1	IVS (N)	-	5'-A TAC CAA CCT-gel	+141
	2	IVS I/1 G/A	8	5'-A TAC CAA tCT-gel	+141
	3	IVS I/1 G/T	8	5'-A TAC CAA aCT-gel	+141
	4	IVS I/2 T/A	7	5'-A TAC Cat CCT-gel	+141
	5	IVS I/2 T/C	7	5'-A TAC Cag CCT-gel	+141
10	6	IVS I/2 T/G	7	5'-A TAC Cac CCT-gel	+141
	7	IVS I/5 G/A	4	5'-A TAt CAA CCT-gel	+141
	8	IVS I/5 G/C	4	5'-A TAg CAA CCT-gel	+141
	9	IVS I/5 G/T	4	5'-A TAa CAA CCT-gel	+141
	10	IVS I/6 T/C	3	5'-A TgC CAA CCT-gel	+141
15	11	CD 26 (N)	-	5'-G GCC TCA CCA-gel	+125
	12	CD 26 G/A	6	5'-G GCC TtA CCA-gel	+125
	1	IVS (N)	9	5'-TGA TAC CAA C-gel	+143
	2	IVS I/2 T/G	9	5'-TGA TAC CAc C-gel	+143
20	3	IVS (N)	8	5'-GA TAC CAA CC-gel	+142
	4	IVS I/2 T/G	8	5'-GA TAC CAc CC-gel	+142
	5	IVS (N)	7	5'-A TAC CAA CCT-gel	+141
	6	IVS I/2 T/G	7	5'-A TAC Cac CCT-gel	+141
	7	IVS (N)	6	5'-TAC CAA CCT G-gel	+140
	8	IVS I/2 T/G	6	5'-TAC CAc CCT G-gel	+140
25	9	IVS (N)	5	5'-AC CAA CCT GC-gel	+139
	10	IVS I/2 T/G	5	5'-AC CAc CCT GC-gel	+139
	11	IVS (N)	4	5'-C CAA CCT GCC-gel	+138
	12	IVS I/2 T/G	4	5'-C CAc CCT GCC-gel	+138
30	13	IVS (N)	3	5'-CAA CCT GCC-gel	+137
	14	IVS I/2 T/G	3	5'-CAc CCT GCC C-gel	+137
	15	IVS (N)	2	5'-AA CCT GCC CA-gel	+136
	16	IVS I/2 T/G	2	5'-Ac CCT GCC CA-gel	+136

TABLE 4  
The effect of the position of the allele base  
within 10-mers on mutation detection.

5

10

15

Position allele	19-mer			RNA		
	$T_{0.1}$ of perfect	$T_{0.1}$ of (G-A)	$\Delta T_{0.1}$	$T_{0.1}$ of perfect	$T_{0.1}$ of (G-A)	$\Delta T_{0.1}$
9	40	32	8	35	37	-2
8	47	32	15	49	38	11
7	42	30	12	44	41	3
6	47	28	19	49	41	8
5	52	38	14	50	42	8
4	54	39	15	54	44	10
3	55	46	9	59	54	5
2	52	46	6	58	53	5

TABLE 5  
Identification of  $\beta$ -thalassaemia  
mutations by hybridization with the microchip.

Hybridized Probe		Immobilized 10-mer oligonucleotide														
		#	Allele	Size (nt)	R at Td=											
					IVS (N)	I/1 G/A	I/1 G/T	I/2 T/A	I/2 T/C	I/2 T/G	I/5 G/A	I/5 G/G	I/5 G/T	I/6 T/C	CD26 (N)	CD26 G/A
1	a		IVS (N)	75	1	0.04	0	0.2	0.05	0.07	0	0	0	0.04	1	-
	b		IVS (N)	19 <sup>a</sup>	1	0.09	0.07	0.02	0.03	0.01	0.03	0.03	0.07	ND	0	0
	a		IVS I/2 T/A	F1 75	0.15	0	0	1	0.12	0.08	0	0	0	0	1	-
	b		IVS I/2 T/A	F1 133	0.03	0	0	1	0	0.3	0	0	0	0	1	0.19
3	c		IVS I/2 T/A	19 <sup>a</sup>	0.01	0	0	1	0.07	0.03	0	0	0	0	0.01	0
	a		IVS I/1 G/A	133 <sub>r</sub>	0.03	1	0	0.01	0	0.02	0	0	0	0	1	-
4	b		IVS I/1 G/A	19 <sup>c</sup>	0.01	1	0.01	0.01	0	0	0.01	0	0	0	0	0
	a		IVS I/1 G/A & IVS I/6 T/C	133 <sub>r</sub>	0.2	0.85	0	0.2	0	0.05	0	0	0	1	1	-
5	b		IVS I/1 G/A	19 <sup>c</sup>	0.01	1	0.01	0.01	0	0	0.01	0	0	0	0	0
	c		IVS I/6 T/C	19 <sup>a</sup>	0.1	0	0	0	0	0	0	0	0	1	0	0
6	a		IVS I/5 G/T	19 <sup>a</sup>	0	0	0	0	0	0	0.03	0.02	1	0	0	0
	b		CD26 (N)	19 <sup>c</sup>	0	0	0	0	0	0	0	0	0	0	1	0.03
7	c		CD26 G/A	19 <sup>a</sup>	0.03	0	0	0	0	0	0	0	0	0	0.04	1
	a		IVS (N)	75	1	0.04	0	0.2	0.05	0.07	0	0	0	0.04	1	-
8	b		IVS I/2 T/A	F1 133	0.03	0	0	1	0	0.3	0	0	0	0	1	-

The noticeable exceptions are oligonucleotides corresponding to IVS ½ T/A and IVS ½ T/G mutations that show strong mismatched signals upon hybridization with non-corresponding samples of IVS (N) and IVS ½ T/A RNA's, respectively (Table 5, 1a, 2b, 4a, 6a and 6b). The relative intensities of these mismatched signals can be significantly decreased by choosing the proper oligonucleotides for immobilization. It appears that the diagnostic assays can be carried out with RNAs 75 nucleotides (nt) long (Table 5, probes 1a, and 6a), and 133 nt long (probes 2a and 6b), as well as with 133 nt long RNA fragmented to pieces 20-40 nt long (probes 3a and 4a). However, the intensities of the hybridization signals after fragmentation are increased by about 5 times and the time of hybridization is decreased from several hours to a tens of minutes.

The longer RNA probes diffuse more slowly into the gel and can form stable secondary structures or aggregates. These factors interfere with their hybridization with rather short immobilized oligonucleotides. Thus, the fragmentation seems to be an essential step in sample preparation, since it enhances and speeds the hybridization.

In addition to the measuring of the melting curves, the reliability of identification of mutations and base changes can be enhanced by the use of a multicolor fluorescence microscope (Yershov *et al.*, 1996). For this purpose, the tested RNA is marked by one fluorescence label and is hybridized with a microchip in the presence of a normal allele sample labeled with a different dye. The pattern and the ratio of hybridization measured with the two dyes will be similar for all microchip oligonucleotides except for those that correspond to different allele bases, i.e., mutations. Table 4 shows the results of such an experiment. The patterns of hybridization detected at two wavelengths are very similar.

As shown in Table 3, the immobilized 10-mers matching the mutations IVS I-2 T/G, IVS I-2 T/C, and IVS I-2 T/A are hybridized rather strongly with some RNA probes that correspond to other alleles. Different structural factors in RNA could cause this hybridization. The effect of these factors can be minimized by placing a variable IVS I-2 base into different positions of the 10-mers. The results of such experiments are shown in Table 4. Microchip II was successively hybridized with fragmented 75-nt-long RNA or with a synthetic DNA 19-mer, both corresponding to the normal allele. Microchip II contained two similar sets of eight overlapped



immobilized 10-mers that are complementary either to a normal allele or to IVS I-2 T/G allele. The allele specific bases A for the first set and C for the second set are located in these 10-mers in all internal positions from the 2<sup>nd</sup> to the 9<sup>th</sup>. These bases form perfect A-T or mismatched A-G base pairs, respectively. The stability of the perfect and mismatched duplexes formed on the microchip is determined as  $T_{0.1}$  the temperature at which the initial hybridization signal of the duplex is decreased to one-tenth of the original intensity.  $\Delta T_{0.1}$  corresponds to the difference in  $T_{0.1}$  between the perfect and similar mismatched duplexes. A better discrimination of the perfect and mismatched duplexes is reflected in higher values of  $\Delta T_{0.1}$ . The discrimination efficiency ( $\Delta T$ ) was lower for hybridized RNAs than for the 19-mers. The discrimination was surprisingly low,  $\Delta T = -2^{\circ}$  and  $3^{\circ}\text{C}$ , when the allelic bases were placed at the 9<sup>th</sup> or 7<sup>th</sup> position, respectively, of the immobilized oligonucleotides. It appears that secondary structures and the presence of similar sequences in other regions of the RNA causes this lowering. These effects can be partly predicted from the sequence of the region that is searched for mutations. However, it is impossible to reach a high discrimination ( $\Delta T = 8-11^{\circ}\text{C}$ ) when allele bases are placed in other positions, for example the 8, 6, 5, or 4 positions.

The hybridization of RNA transcripts of PCR-amplified DNA with oligonucleotide microchips allows the reliable identification of base changes and discrimination of homozygous and heterozygous  $\beta$ -thalassemia mutations in the genomic DNA of patients.

RNA transcribed from PCR-amplified DNA provides an easier method for preparing a sufficient amount of labeled, single-stranded samples than the use of DNA prepared by PCR amplification. RNA can be fragmented and one fluorescent dye molecule can be introduced per fragment.

#### **Example 7: Use of a Customized Microchip Biosensor to Detect Gene Expression**

Gene expression is one of the central themes in modern molecular biology. DNA from well studied genetical sources has already been systematically sequenced. For these sequences hybridization procedures are successfully used to estimate a level of differential gene expression. The results of this estimation are useful for

understanding fundamental mechanisms of development biology, embryology and treatment of genetic and infectious diseases.

To determine whether oligonucleotide microchips are useful to identify gene expression, microchip biosensor hybridization was carried out with ssDNA fragments  
5 isolated from six different genes:

- 205 b fragment from glyceraldehyde 3-phosphate dehydrogenase (G3PDH);
- 281 b fragment from human transferrin receptor (HTR);
- 224 b fragment from human  $\beta_2$ -microglobulin (B2M);
- 545 b fragment from human interleukin-1 receptor (IL1R);
- 10 - 188 b fragment from human NF-kB (p50);
- 224 b fragment from human interferon  $\gamma$  receptor (IGR).

A customized microchip, containing immobilized 60b oligonucleotides, having at the 3'-terminal position 3-methyluridine residues, corresponding to five housekeeping genes (G3PDH, HTR, B2M, IL1R and NF-kb(p50)) (CLONTECH  
15 catalog 94/95 "Tools for the Molecular Biologist", pp. 90-93) were produced for hybridization experiments with complementary ssDNA fragments. Each oligonucleotide was applied at two positions on the microchip in a 1:10 ratio of amount (0.3 pmol:0.03 pmol each). ssDNA fragments complementary to immobilized oligonucleotides were synthesized by asymmetric PCR amplification (using only one  
20 primer) with fluorescently labeled nucleotide triphosphates (*FUORscript T7*, Fluorescein-Labeling In Vitro Transcription Kit). Moreover, the PCR-primer bore a biotin tag that was utilized for following isolation of synthesized ssDNA fragments with avidin carried on a column (Sambrook et al. "Molecular Cloning" 2d edit., p. 12.14). FIG. 8 demonstrates hybridization on the microchip. Intensity of fluorescence  
25 in each spot depends on the amount of immobilized oligonucleotide and on the length of the DNA fragment in the spot. For hybridization, 10  $\mu$ l of Buffer A (50% formamide, 10% dextran sulfate, 1% SDS, 50 mM sodium phosphate at pH 7.4, 750 mM sodium chloride, 5 mM sodium EDTA) containing ssDNA with a concentration of 0.5 pmol/ $\mu$ l (approximately 0.05  $\mu$ g/ $\mu$ l) was incubated for about 6 - 12 h at room  
30 temperature, washed briefly with H<sub>2</sub>O and analyzed with a fluorescent microscope.

Before rehybridization the microchip was treated in Buffer B (50% formamide, 1% Tween 20) for 30 min. at 50°C to completely remove hybridized ssDNA.

These results indicate that concentration of fluorescently labeled ssDNA may be decreased up to 100 fold. Hybridization with individual ssDNA fragments indicates high specificity of studied oligonucleotides. There was no cross-hybridization detected between different tested DNAs and immobilized oligonucleotides. None of the oligonucleotides demonstrated a signal when hybridized with non-specific DNA (e.g. probe IGR). This differentiates "expression" of non-expressed genes from expression of housekeeping genes. Genes that are not expressed in a particular cell or tissue, may actually be picked up in conventional screening procedures as having a low expression, while other genes being expressed in all cells (housekeeping genes) will also be picked up as having low to moderate expression. The housekeeping genes are actually being expressed. In this example a difference in signal is detectable so that low level expression could be unambiguously distinguished from low level background.

The procedure detects expression of genes of high and middle expression level. To determine low level gene expression selective RT-PCR amplification is preferred.

**Example 8: Use of a Customized Microchip Biosensor of the Present Invention to Detect HLA Polymorphisms**

A difficult problem of genotype recognition arises in studying different haplotypes (alleles) of genes encoding Human Leucocyte Antigens (HLA) in regions of histocompatibility genes. The HLA locus (class I and class II genes) is responsible for histocompatibility of tissue transplantation. The need for allele identification is encountered also in various medical and biological tasks involving HLA class II genes. There are many clinical data showing strong association between HLA genotype and susceptibility to some disorders, for example some alleles DQA1/DQB1 are clearly related to IDDM (Insulin - Dependent Diabetes Mellitus), malaria, autoimmune diseases, such as rheumatoid arthritis and pemphigus vulgaris - a skin disease which causes severe blistering. The high level of polymorphism of HLA has been shown to be useful for identification of individuals determining the group of risk for some

diseases. HLA typing is particularly crucial for matching donors for transplants. It is also proposed for infertility work-ups.

In this aspect, the present invention provides a method which allows an array of immobilized 8-12 bp-long oligonucleotides to form an oligonucleotide microchip  
5 thereby facilitating identification of HLA DQA1 alleles.

An algorithm has been designed and special computer programs have been constructed which allow the analysis of the nucleotide sequences of all alleles of various HLA subloci. Forming an optimized set of oligonucleotides provides high reliability of detection of homo- and heterozygotes for the HLA alleles.

10 A customized microchip, containing an array of eighteen PAA - gel immobilized (1pmol of each) short oligonucleotides has been produced for hybridization with fluorescently labelled complementary HLA DQA1 DNA or RNA probe for allele identification. 18 decamers were loaded on the chip in the following order, from left to right:

15

TABLE 6					
first (upper) row	1	2	3	4	g <sup>4</sup> -control oligo
second:	5	6	7	8	g <sup>3</sup> -control oligo
third:	9	10	11	12	13
20 4-th:	14	15	16	17	18

The sequence of the oligos used was as shown in FIG. 9.

The oligonucleotides immobilized on the microchip are complementary to the sense strand of different alleles of DQA1 DNA and some control oligos. A microchip  
25 with 20 oligonucleotides was manufactured for partial identification of 15 different alleles in the HLA DQA1 region. PCR was used to prepare 229 bp (starting from codon 12 to codon 87) DNA fragments of the polymorphic second exon of the DQA1 gene from human genomic DNA. Nested primers were used 2DQAAMP-A:5'-a t ggt gta aac ttg tac cag t; and 2DQAAMP-B:5'tt ggt agc agc ggt aga gtt g. Nested PCR  
30 primers were: T7-2DQAAMP-A and primer B. The first primer containing the promoter for T7 RNA polymerase and PCR product were used for *in vitro* transcription. RNA probes were identical to the coding DNA strand. RNA was

fragmented, labelled with fluorescein and used for hybridization with the microchip. Hybridization conditions were as follows: overnight incubation at 5°C in 1M NaCl, 1mM EDTA, 5 mM Na-phosphate, pH7.0, 1% Tween 20. The temperature was then increased stepwise at 10°C intervals, and fluorescence measurements were taken at  
5 each step. BUFFER WAS NOT CHANGED.

FIG.10 shows the hybridization results and presents schematically the HLA DQA1 - chip for allele identification. In FIG. 10 three diagonally placed oligos (11-0101/0104 allele specific; 6-specific for 0101,01021,01022,0103,00104; 17- correspond to all alleles, except 0502) gave a positive hybridization signal, and are  
10 observed as three diagonally placed bright fluorescence spots. The probes were identified as the 0101 or 0104 allele (both alleles are identical in the second exon). All other oligos yielded much weaker fluorescence signals compared with those described above, because none of them contain sequences complementary to alleles 0101 and 0104. On the other hand any allele different from 0101 or 0104, reveals  
15 another set of hybridization signals.

The brightest fluorescent squares on the chip were: Oligo# 4 which is 03011 or 0302 specific; oligo# 8 is Taq polymerase-specific artifacts; oligo# 18-belong to alleles 0101-05011; oligo 11-0101,0104 allele specific; 17 - corresponds to all alleles, except 0502; g4 is a fluorescent control oligo; #13 - mismatch to #18. All other chip  
20 elements showed significantly less intensive fluorescence. The genotype identified by these probes has a 0101/0104-0302/03011 heterozygote.

**Example 9: Use of a Customized Microchip Biosensor to Detect the Lyme Disease Spirochetes**

25 Bacteria belonging to the species *Barretia burgdorferi* and related species of tick-borne spirochetes are capable of causing human and veterinary disease. Nucleic acid probes are available to detect bacteria causing Lyme disease. These bacteria cannot be identified by standard microbiological methods, although immunological tests are available.

30 Using the methods of the present invention, oligonucleotides are prepared according to Weisburg (1995) and added to a micromatrix designed for use in detecting Lyme disease in a clinical sample.

**Example 10: Use of a Customized Microchip Biosensor to Detect Salmonella In Food Samples**

Salmonella presence is detected most commonly by preparing cultures  
5 according to standard microbiological laboratory procedures, and testing the cultures  
for morphological and biochemical characteristics. After about 48 hours after  
collection of a sample testing begins and takes several days to complete.

However, RNA and DNA probes for Salmonella testing are available. Using  
the methods of the present invention, oligonucleotides are prepared according to Lane  
10 *et al.* (1996) incorporated herein by reference and added to a microchip designed for  
use in detecting Salmonella in food samples by distinguishing rRNA of Salmonella  
from non-Salmonella.

**DOCUMENTS CITED**

- AUSUBEL, F.M., R. BRENT, R.E. KINGSTON, D.D. MORE, J.G. SEIDMAN, J.A. SMITH, and K. STRUHL, Eds., ed. (1994) Current Protocols in Molecular Biology. Greene Wiley Interscience: New York, NY.
- 5 BEATTIE, K.L., BEATTIE, W.G., MENG, L., TURNER, S.L., CORAL-VAZQUEZ, R., SMITH, D.D., McINTYRE, P.M. and DAO, D.D.: Advances in Genosensor Research. Clin. Chem. 41/5 (1995) 700-706.
- CONNER, B.J., REYERS, A.A., MORIN, C., ITAKURA, K.,
- 10 TEPLITZ, R.L. and WALLACE, R.B. (1983) Detection of sickle cell  $\beta$ -globin allele by hybridization with synthetic oligonucleotides. Proc. Natl. Acad. Sci. USA 80: 278-282.
- DIAZ-CHICO, J.C., YANG, K.-G., YANG, K.-Y., EFREMOV, D.G., STOMING, T.A. and HUISMAN, T.H. (1988) J. Biochim. Biophys. Acta 949: 43-48.
- 15 FODOR, S.P.A., READ, J.L., PIRRUNG, M.C., STRYER, L., LU, A.T. and SOLAS, D.: Light-directed, spatially addressable parallel chemical synthesis. Science 251 (1991) 767-773.
- GHU, Z., GUILFOYLE, R.A., THIEL, A.J., WANG, R., and SMITH, L.M.: Direct fluorescence analysis of genetic polymorphisms by hybridization with
- 20 oligonucleotide arrays on glass support. Nucl. Acids Res. 22 (1994) 5456-5465.
- KHRAPKO, K., LYSOV, YU. KHORLIN, A., IVANOV, I., YERSHOV, G., VASILENKO, S., FLORENTIEV, V., and MIRZABEKOV, A. (1991) A method for DNA sequencing by hybridization with oligonucleotide matrix. J. DNA Sequencing 1 375-388.
- 25 KHRAPKO, K. R. *et al.* (1996) U.S. Pat. No. 5,552,270.
- LAMTURE, J.B., BEATTIE, K.L., BURKE, B.E., EGGERS, M.D., EHRlich, D.J., FOWLE, R., HOLLIS, M.A., KOSICKI, B.B., REICH, R.K., SMITH, S.R., VARMA, R.S. and HOGAN, M.E.: Direct detection of nucleic acid hybridization on the surface of a charge-coupled device. Nucl. Acids Res. 22 (1994)
- 30 2121-2125.

- LANE, D.J. *et al.* (1996) Oligonucleotide Probes for Detection of Salmonella, U.S. Pat. No. 5,495,008.
- LAWN, R. EFSTRATIADIS, A., O'CONNEL, C., and MANIATIS, T. (1980) The nucleotide sequence of the human  $\beta$ -globin gene. *Cell* 21: 647-651.
- 5 LESNIK, E.A., and FREIER, S. (1995) Relative thermodynamic stability of DNA, RNA, and DNA:RNA hybrid duplexes: relationship with base composition and structure. *Biochemistry* 34: 10807-10815.
- LIPSHUTZ, R.J., MORRIS, D., CHEM, M., HUBBELL, E., KOZAL, M.J., SHAH, N., SHEN, N., YANG, R. and FODOR, S.P.A. (1995) Using  
10 oligonucleotide probe arrays to access genetic diversity. *BioTechniques* 19: 442-447.
- MAIDAK, B.C. *et al.* (1996) The ribosomal database project (RDP), *Nucleo Acids Res.* 24:82-85.
- MATSON, R.S., RAMPAL, J., PENTONEY, S.L., ANDERSON, P.D. and COASSIN, P.: Biopolymer synthesis on polypropylene support: oligonucleotide  
15 arrays. *Anal. Biochem.* 224 (1995) 110-116.
- MIRZABEKOV, A. (1994) DNA Sequencing by hybridization: a megasequencing method and a diagnostic tool? *Trends in Biotechnology* 12: 27-32.
- MOBARRY, B.K. *et al.* (1996) Phylogenetic probes for analyzing abundance and spatial organization of nitrifying bacteria. *Appl. Environ. Microbiol.*  
20 62:2156-2162.
- PEASE, A.C., SOLAR, D., SULLIVAN, E.J., CRONIN, M.T., HOLMES C.P. and FODOR, S.P.A.: Light-directed oligonucleotide arrays for rapid DNA sequence analysis. *Proc. Natl. Acad. Sci. USA* 91 (1994) 5022-5026.
- PONCZ, M. SOLOWIEJCZYK, D. and HARPEL, B.(1982)  
25 Construction of human gene libraries from small amount of peripheral bloods: analysis of  $\beta$ -like globin genes. *Hemoglobin* 6: 27-40.
- STAHL, D.A. (1995) Application of phylogenetically based hybridization probes to microbial ecology. *Mol. Ecol.* 4:535-542.
- SOUTHERN, E. M. (1996) DNA chips: analyzing sequence by  
30 hybridization to oligonucleotides on a large scale, *TIG* 12:110-115.



SOUTHERN, E.M., MASKOS, U. and ELDER, J.K.: Analyzing and comparing nucleic acid sequences by hybridization to arrays of oligonucleotides: evaluation using experimental models. *Genomics* 13 (1992) 1008-1017.

WAGNER, M. *et al.* (1995) In situ identification of ammoniaoxidizing  
5 bacteria. *Sept. Appl. Microbiol* 18: 251-264.

WEISBURG, W.G. (1995) Nucleic Acid Probes for the Detection of Lyme Disease Spirochetes, U.S. Pat. No. 5,466,577.

WOESE, C.R. (1987) Bacterial evolution, *Microbial. Rev.* 51:221-271.

YERSHOV, G., BARSKY, V., BELGOVSKIY, A., KIRILLOV, E.,  
10 KREINDLIN, E., IVANOV, I., PARINOV, S., GUSHIN, D., DROBYSHEV, A.,  
DUBILEY, S., and MIRZABEKOV, A. (1996) DNA analysis and diagnostics on oligonucleotide microchips. *Proc. Natl. Acad. Sci. USA* 93: 4913-4918.

U.S. Pat. No. 5,552,270

**WE CLAIM:**

1. A method for using a reusable microchip to identify at least one nitrogen base sequence in nucleic acids of a sample, said method comprising:
  - (a) providing a customized matrix of oligonucleotides on the microchip in an arrangement that is an ordered scheme designed to present a detectable pattern as a whole that identifies the nitrogen base sequences of nucleic acids in the sample;
  - (b) hybridizing nucleic acids from the sample on said microchip; and
  - (c) identifying at least one nitrogen base sequence in said sample by the pattern detected as a whole on the microchip, said pattern resulting from the locations of oligonucleotides which hybridized to the sample nucleic acids.
2. The method of claim 1, wherein said nitrogen base sequence is in a DNA molecule.
3. The method of claim 1, wherein said nitrogen base sequence is in a 16S RNA, mRNA or other RNA molecule.
4. The method of claim 1, wherein the customized matrix of oligonucleotides on the reusable microchip is formed by a plurality of gel elements, wherein the number of elements is determined by the number of oligonucleotides in the matrix and wherein each gel element contains one oligonucleotide of a desired nitrogen base sequence length and concentrations, each gel element being separated from another by hydrophobic glass spaces, and wherein the gel elements have a vertical height above the plane of the interstitial spaces of not more than about 30  $\mu\text{m}$ , said oligonucleotides being positioned in specific locations.
5. The method of claim 4, wherein said matrix comprises oligonucleotides having the base sequences:

Oligonucleotide Name and Position	Sequence (5' to 3')
Nb1000	TGC GAC CGG TCA TGG (SEQ ID:6)
NIT3	CCT GTG CTC CAT GCT CCG (SEQ ID:7)
NEU23	CCC CTC TGC TGC ACT CTA (SEQ ID:8)
NSO190	CGA TCC CCT GCT TTT CTC (SEQ ID:9)
NSO1225	CGC GAT TGT ATT ACG TGT GA (SEQ ID:10)
NSMO156	TAT TAG CAC ATC TTT CGA T (SEQ ID:11)
NSV443	CCG TGA CCG TTT CGT TCC (SEQ ID:12)
BAC338	GCT GCC TCC CGT AGG GAT (SEQ ID:13)
NonBAC338	ACT CCT ACG GGA GGC AGC (SEQ ID:14)
UNI1390	GAC GGG CGG TGT GTA CAA (SEQ ID:15)

6. The method of claim 5, wherein the oligonucleotides are arranged in a matrix wherein I, II, III and IV are columns and "ABC" are rows in a microchip:

	I	II	III	IV
A	Nb1000	NIT3	NEU23	Nso190
B	Nso1225	Nsm156	Nsv443	
C	Bac338	NonBac338	Uni1390	

20

7. The method of claim 1, further comprising:

(e) adding a label to the nitrogen base sequences in said sample before hybridizing them to oligonucleotides on the microchip.

8. The method of claim 7, wherein the label is a fluorescent dye.

9. The method of claim 7, wherein the label is a plurality of different dyes.

10. A microchip matrix for the detection and classification of nitrifying bacteria wherein said matrix has the following design:

30

	I	II	III	IV
A	Nb1000	NIT3	NEU23	Nso190
B	Nso1225	Nsm156	Nsv443	
C	Bac338	NonBac338	Uni1390	

5

wherein I, II, III and IV are columns and A, B, C, are row in a microchip, and wherein the base sequences are:

Oligonucleotide Name and Position	Sequence (5' to 3')
Nb1000	5'-TGC GAC CGG TCA TGG-3'
NIT3	5'CCT GTG CTC CAT GCT CCG-3'
NEU23	5'-CCC CTC TGC TGC ACT CTA-3'
NSO190	5'-CGA TCC CCT GCT TTT CTC- 3'
NSO1225	5'-CGC GAT TGT ATT ACG TGT GA-3'
NSMO156	5'-TAT TAG CAC ATC TTT CGA t-3'
NSV443	5'-CCG TGA CCG TTT CGT TCC-3'
BAC338	5'-GCT GCC TCC CGT AGG GAT-3'
NonBAC338	5-'ACT CCT ACG GGA GGC AGC-3'
UNI1390	5'GAC GGG CGG TGT GTA CAA-3'

10

15

20

11. A diagnostic assay for the presence of a mutation in a gene in a sample said assay comprising:

- (a) designing a customized reusable microchip biosensor comprising at least one oligonucleotide that hybridizes to a gene having the mutation, said designing producing an overall pattern that is detectable in the presence of said hybridization of the mutation;
- (b) contacting the sample to the customized microchip biosensor under conditions that allow hybridization of the gene sequence that is a mutation to oligonucleotides on the microchip; and
- (c) determining whether hybridization occurs by analyzing the microchip pattern from which presence of the mutation in the gene is determined.

25

30

12. A method using non-equilibrium melting curves to detect mismatches between -a nitrogen base sequence on a microchip and a nitrogen base sequence to be tested, said method comprising:

- 5 (a) generating the non-equilibrium melting curves by simultaneously monitoring hybridization between the nitrogen base sequence that matches or mismatches with sequences on the microchip at a series of temperatures;
- (b) selecting the temperature at which maximum discrimination occurs between the match and the mismatch; and
- 10 (c) determining the degree of mismatch of the nitrogen base sequence to be tested at the selected temperature.

13. The method of claim 12, wherein the selected temperature is that temperature at which the signal intensity of a mismatched sequence is at least one tenth of the signal intensity of a matched sequence.

- 15 14. A customized oligonucleotide microchip for the detection of a betaglobin mutation, said microchip containing seven oligonucleotides having the following nitrogen base sequences:

	<u>Identifier</u>	<u>Sequences</u>	<u>SEQ ID</u>
	NO:		
20	IVS (N)	CCTGGGCAGGTTGGTATCA;	48
	IVS I/2 T/A	CCTGGGCAGGaTGGTATCA;	49
	IVS I/1 G/A	CCTGGGCAGaTTGGTATCA;	50
	IVS I/6 T/C	CCTGGGCAGGTTGcTATCA;	51
	IVS I/5 G/T	CCTGGGCAGGTTGtTATCA;	52
25	CD26 (N)	GTTGGTGGTGAGGCCCTGG;	53
	CD26 G/A	GTTGGTGGTaAGGCCCTGG	54.

- 15 15. A reusable customized oligonucleotide microchip for quantitation of the expression of a gene, wherein the design of the microchip presents a pattern
- 30 corresponding to the degree of expression.

16. A customized oligonucleotide microchip for the detection of HLA polymorphism, wherein the design of the microchip presents a pattern in which HLA probes are positioned to answer a specific question when hybridization occurs.

17. The customized oligonucleotide microchip of claim 16, comprising the  
5 following oligonucleotides;

10 AGGCAACGTG (1) (SEQ ID:55);  
AGGCGACGTG (2) (SEQ ID:56);  
GGTGAACCTGG (3) (SEQ ID:57);  
TAAATCTGCG (4) (SEQ ID:58);  
AGGCAACATG (5) (SEQ ID:59);  
CAAAACCTCC (6) (SEQ ID:60);  
GCAAACACCA (7) (SEQ ID:61);  
TACACCATAA (8) (SEQ ID:62);  
ACTGCTCATC (9) (SEQ ID:63);  
15 CAATGTCTTC (10) (SEQ ID:64);  
CTCCTCATCT (11) (SEQ ID:65);  
TGCCGGTCAA (12) (SEQ ID:66);  
TTAGGACAGC (13) (SEQ ID:67);  
ACACCACAAG (14) (SEQ ID:68);  
20 CACAATGCCT (15) (SEQ ID:69);  
CAGCAGTAGA (16) (SEQ ID:70);  
TGCGGGTCAA (17) (SEQ ID:71);  
TTAGCACAGC (18) (SEQ ID:72).

25 18. A method for designing a reusable genetic microchip to answer a specific question about a test sample, said answer provided by detection of hybridization patterns of the nitrogen base sequences of nucleic or ribonucleic acids of molecules present in the sample, on the microchip as a whole, said method comprising:



30 (a) determining the specific question; and  
(b) designing the microchip so that hybridization of nucleic acids present in the sample provides a pattern that when detected as a whole provides an answer to the question; and  
(c) detecting the pattern.

19. The method of claim 18, wherein the sample is water, and the question is what pathogens are present.


20. The method of claim 18, wherein the sample is blood, and the question is what genetic mutations are present.

5        21. The method of claim 18, wherein the sample is a food, and the specific question is what containments are present.

22. The method of claim 18, wherein the pattern is a bar code pattern and detecting the pattern is done by a bar code laser scanner.

*Fig. 1A**Fig. 1B*



*Fig. 2*





Fig. 3c








Fig. 3D







*Fig. 5*


- a. IVS (N) 5'-TMR-CCTGGGCAGGTTGGTATCA-3' (SEQ ID NO:48);  
b. IVS I/2 T/A 5'-TMR-CCTGGGCAGGaTGGTATCA-3' (SEQ ID NO:49);  
c. IVS I/1 G/A 5'-TMR-CCTGGGCAGaTTGGTATCA-3' (SEQ ID NO:50);  
d. IVS I/6 T/C 5'-TMR-CCTGGGCAGGTTGtTATCA-3' (SEQ ID NO:51);  
e. IVS I/5 G/T 5'-TMR-CCTGGGCAGGTTGtTATCA-3' (SEQ ID NO:52);  
f. CD26 (N) 5'-TMR-GTTGGTGGTGAGGCCCTGG-3' (SEQ ID NO:53);  
g. CD26 G/A 5'-TMR-GTTGGTGGTaAGGCCCTGG-3' (SEQ ID NO:54);

*Fig. 6*

- 5' -AGGCAACGTG (1) (SEQ ID NO: 55);  
5' -AGGCGACGTG (2) (SEQ ID NO: 56);  
5' -GGTGAACCTG (3) (SEQ ID NO: 57);  
5' -TAAATCTGCG (4) (SEQ ID NO: 58);  
5' -AGGCAACATG (5) (SEQ ID NO: 59);  
5' -CAAAACCTCC (6) (SEQ ID NO: 60);  
5' -GCAAACACCA (7) (SEQ ID NO: 61);  
5' -TACACCATAA (8) (SEQ ID NO: 62);  
5' -ACTGCTCATC (9) (SEQ ID NO: 63);  
5' -CAATGTCTTC (10) (SEQ ID NO: 64);  
5' -CTCCTCATCT (11) (SEQ ID NO: 65);  
5' -TGCCGGTCAA (12) (SEQ ID NO: 66);  
5' -TTAGGACAGC (13) (SEQ ID NO: 67);  
5' -ACACCACAAG (14) (SEQ ID NO: 68);  
5' -CACAATGCCT (15) (SEQ ID NO: 69);  
5' -CAGCAGTAGA (16) (SEQ ID NO: 70);  
5' -TGCGGGTCAA (17) (SEQ ID NO: 71);  
5' -TTAGCACAGC (18) (SEQ ID NO: 72);

*Fig. 7*

Fig. 7









Fig. 8



*Fig. 10*




FIG. 11




FIG. 12




FIG. 13

专利名称(译)	定制的寡核苷酸微芯片，可将多种遗传信息转换为简单模式		
公开(公告)号	<a href="#">EP1208222A2</a>	公开(公告)日	2002-05-29
申请号	EP2000912055	申请日	2000-02-29
[标]申请(专利权)人(译)	芝加哥大学		
申请(专利权)人(译)	芝加哥大学		
当前申请(专利权)人(译)	ENGELHARDT学院分子生物学报		
[标]发明人	MIRZABEKOV ANDREI GUSCHIN DMITRY Y CHIK VALENTINE DROBYSHEV ALEKSEI FOTIN ALEXANDER YERSHOV GENNADIY LYSOV YURI		
发明人	MIRZABEKOV, ANDREI GUSCHIN, DMITRY Y. CHIK, VALENTINE DROBYSHEV, ALEKSEI FOTIN, ALEXANDER YERSHOV, GENNADIY LYSOV, YURI		
IPC分类号	G01N33/53 C12M1/00 C12M1/34 C12N15/09 C12Q1/68 C12Q1/6837 C12Q1/6881 C12Q1/6883 C12Q1/689 G01N37/00		
CPC分类号	C12Q1/6837 B01J2219/00644 C12Q1/6881 C12Q1/6883 C12Q1/689 C12Q2600/156 G16B25/00 C12Q2565/607 C12Q2565/513		
优先权	09/261115 1999-03-03 US		
外部链接	<a href="#">Espacenet</a>		

#### 摘要(译)

本发明涉及使用定制的寡核苷酸微芯片作为生物传感器，用于检测和鉴定对环境，食品和生物样品中的不同基因，生物体和/或个体具有特异性的核酸。微芯片被设计成将多个比特的遗传信息转换成被解释为单元的更简单的信号模式。由于从样品到微芯片的寡核苷酸杂交的改进方法，微芯片是可重复使用的和可运输的。对于现场研究，便携式激光或条形码扫描仪是合适的。