



(11) **EP 1 535 076 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
06.08.2008 Bulletin 2008/32

(21) Application number: **03792291.1**

(22) Date of filing: **11.08.2003**

(51) Int Cl.:
G01N 33/68 (2006.01)

(86) International application number:
PCT/EP2003/008879

(87) International publication number:
WO 2004/019043 (04.03.2004 Gazette 2004/10)

(54) **POLYPEPTIDE BIOMARKERS FOR DIAGNOSING ALZHEIMER'S DISEASE**

POLYPEPTID-BIOMARKER ZUR DIAGNOSE VON MORBUS ALZHEIMER

MARQUEURS BIOLOGIQUES PERMETTANT LE DIAGNOSTIC DE LA MALADIE D'ALZHEIMER

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PT RO SE SI SK TR

(30) Priority: **23.08.2002 EP 02018283**
29.11.2002 EP 02026643

(43) Date of publication of application:
01.06.2005 Bulletin 2005/22

(60) Divisional application:
08002793.1

(73) Proprietor: **Bayer HealthCare AG**
51368 Leverkusen (DE)

(72) Inventors:
• **YALKINOGLU, Özkan**
42115 Wuppertal (DE)
• **KÖNIG, Gerhard**
Arlington, MA 02476 (US)
• **HOCHSTRASSER, Denis, Francois**
CH-1245 Collonge-Bellerive (CH)
• **SANCHEZ, Jean-Charles**
CH-1208 Geneva (CH)
• **CARRETTE, Odile**
F-59100 Roubaix (FR)

(56) References cited:
WO-A-00/25138 **WO-A-01/63294**

- **WEI, LIHONG ET AL: "Cystatin C: Icelandic-like mutation in an animal model of cerebrovascular.beta.-amyloidosis" STROKE (DALLAS) (1996), 27(11), 2080-2085, XP000998699**
- **UTAL A ET AL: "Preliminary expression analysis in Alzheimer's disease using SELDI protein chips." SOCIETY FOR NEUROSCIENCE ABSTRACTS, vol. 26, no. 1-2, 2000, pages Abstract No.-83.7, XP000989844 30th Annual Meeting of the Society of Neuroscience;New Orleans, LA, USA; November 04-09, 2000 ISSN: 0190-5295**
- **DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; March 2003 (2003-03), CARRETTE ODILE ET AL: "A new sensitive and highly specific test for the diagnosis of Alzheimer disease using the ProteinChip(R) technology." XP002265308 Database accession no. PREV200300358956 & FASEB JOURNAL, vol. 17, no. 4-5, March 2003 (2003-03), page Abstract No. 80.3, FASEB Meeting on Experimental Biology: Translating the Genome;San Diego, CA, USA; April 11-15, 2003 ISSN: 0892-6638 (ISSN print)**

EP 1 535 076 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**Field of the invention**

5 **[0001]** The invention is in the field of diagnostics. More specifically, the invention is in the field of assessing the state of Alzheimer's disease in subjects by detection of Alzheimer's disease-specific marker polypeptides.

Background of the invention10 **Alzheimer's disease**

[0002] Alzheimer's disease is an increasingly prevalent form of neurodegeneration that accounts for approximately 50-60% of the overall cases of dementia among people over 65 years of age. Pathologically, Alzheimer's disease neurodegeneration is characterised by prominent atrophy of corticolimbic structures with neuronal death and loss of neuronal synapses, neurofibrillary tangle (NFT) formation, and the formation of senile plaques containing deposits of amyloid β 1-42 (A β 42) aggregates in the brain [Francis PT 1999]. The duration of the progressive cognitive decline is approximately 7 years from the occurrence of first signs until death. It is assumed that the clinical phase is preceded by a 15-30 years preclinical period of continuous deposition of amyloid plaques and neurofibrillary tangles. Age of onset and progression of the disease are largely determined by causative gene mutations and by genetic susceptibility factors. Several environmental risk factors may add to the individual genetic risk factors. Genetic factors known to be involved in the familial form of Alzheimer's disease with early onset of the disease are: mutations in presenilin 1 (PS1), presenilin 2 (PS2), and amyloid precursor protein (APP) genes, and the presence of the apolipoprotein E4 allele. However, the majority (95%) of Alzheimer's disease cases is sporadic and heterogeneous.

[0003] Currently, clinical diagnosis of Alzheimer's disease can only be established at later stages of the disease, when cognitive performance is significantly decreased and paralleled by structural alterations of the brain. The clinical diagnostic work up requires a careful medical history; physical and neurological examination; blood, urine and cerebrospinal fluid (CSF) examinations to exclude metabolic and medical disease states that might masquerade Alzheimer's disease; detailed psychometric examinations to assess mental status and cognitive performance, and imaging techniques such as computed tomographic scan or magnetic resonance imaging of the brain. Diagnostic evaluations at expert centres reach an accuracy of about 80-85%. Due to the fact that these tests are expensive and time consuming, and are particularly inconvenient to patients, there is an increasing need for easy-accessible specific diagnostic biomolecule markers, which can be measured in body fluids, such as CSF, blood or urine, and which have a high positive predictive value for diagnosis of Alzheimer's disease, or would help to distinguish Alzheimer's disease from other forms of dementia. Furthermore, reliable markers sensitive to disease progression may constitute surrogate parameters, a major prerequisite for the evaluation and development of new causal oriented and disease modifying therapeutic strategies in Alzheimer's disease.

[0004] Since CSF directly surrounds the brain, changes in its protein composition may most accurately reflect pathologic conditions that are associated with specific alterations of the protein expression patterns. Over the last decade, a number of biological abnormalities have been reported in the cerebrospinal fluid (CSF) of Alzheimer's disease patients, in particular altered levels of the A β 1-42 fragment of the amyloid precursor protein, and altered levels of the hyperphosphorylated tau protein. The sensitivity and specificity of these markers, however, is low or only modest [The Ronald and Nancy Reagan Research Institute of the Alzheimer's Association and the National Institute on Aging Working Group, 1998, Robles A 1998, Teunissen CE et al., 2002].

[0005] Hence, there is a need for novel biomarkers with sufficient sensitivity and specificity for (i) detecting Alzheimer's disease as early as possible, and (ii) to allow disease differentiation from other types of dementia or neurodegenerative diseases, and (iii) monitoring therapeutic efficacy as surrogate parameter, e.g. in clinical drug development, and to initiate pharmacotherapy as early as possible and postpone loss of memory and disease progression.

Protein Chip Technology

50 **[0006]** A Protein chip technology called Surface Enhanced Laser Desorption/Ionisation time of flight mass spectrometry (SELDI-TOF MS) has recently been developed to facilitate protein profiling of complex biological mixtures [Davies HA 2000, Fung ET 2001, Merchant M 2000].

[0007] Protein chip mass spectrometry has already been used by several groups to detect potentially novel biomarkers of prostate and bladder [Adam BL 2001] or breast cancer [Wulfskuhle JD 2001] in serum, seminal plasma, nipple fluid, urine or cell extracts. For a review on biomarker search using SELDI-TOF MS, see [Issaq HJ 2002].

Cystatin C

[0008] Initially described in 1961 in cerebrospinal fluid (CSF), cystatin C (γ trace or post- γ globulin, Acc. No. P01034) is a small cysteine proteinase inhibitor present in all human body fluids at physiologically relevant concentrations. The physiological role of cystatin C is likely to regulate extracellular cysteine protease activity, which results from microbial invasion or release of lysosomal proteinases from dying or diseased cells. Cystatin C colocalises with β -amyloid ($A\beta$) within the arteriolar walls in Alzheimer's disease brains and cerebral amyloid angiopathy [Levy E 2001]. There are two common haplotypes of the CST3 gene coding for cystatin C (A and B) that differ from each other at three sites: two single base pair changes in the promoter region and one in the signal peptide domain that causes an amino acid substitution (alanine to threonine). Recently, case control studies found associations of CST3 with increased risk for late onset Alzheimer's disease [Crawford FC 2000, Finckh U 2000, Beyer K 2001].

[0009] Hereditary cerebral hemorrhage with amyloidosis, Icelandic type (HCHWA-I), also called hereditary cystatin C amyloid angiopathy (HCCAA), is an autosomal dominant form of cerebral amyloid angiopathy (CAA). The amyloid deposited in the brain vessel's walls is composed mainly of a variant of cystatin C characterised by the presence of the Leu68-Gln substitution [Cohen 1983, Ghiso 1986]. This pathology is also coupled to a decreased concentration of this major cysteine proteinase inhibitor in cerebrospinal fluid and leads to its amyloid deposition in the brain [Grubb AO 1984].

[0010] Leung-Tack *et al* have also purified two N-terminal truncated isoforms of cystatin C in urine from one patient who had received renal transplant. According to their data, (desl-4) cystatin C has an inhibiting effect on two functions of human peripheral mononuclear cells (PMN): O_2^- release and phagocytosis, which may be due to the N-terminal sequence 'KPPR'. Their data support a potentially important role for cystatin C as a possible immunomodulator during inflammation. Accumulating evidence indicates that increased free radical mediated damage to cellular function contributes to the ageing process and age-related neurodegenerative disorders. Oxidative stress may play a role in Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS). Although free-radical damage to neurons may not be the primary event initiating these diseases, it appears that free-radical damage is involved in the pathogenetic cascade of these disorders.

Beta-2-microglobulin

[0011] Beta-2-microglobulin (Acc. No. P01884) constitutes the small constant component of the class I major histocompatibility complex (CMH) and its presence in biological fluids represents the balance between membrane protein turnover and elimination. Since this peptide seems to be increased in some diseases characterised by an elevation of the immune response, its quantification in body fluids has become a useful index of immunological state *in vivo* [Hoekman *et al* 1985]. The function of this protein is unclear, but it seems to be implicated in diseases, which involve glial cell destruction [Ernerudh *et al* 1987].

[0012] The technical problem which is solved by the present invention is the provision of improved methods for diagnosing. Alzheimer's disease and/or monitoring the progression of Alzheimer's disease in a subject.

Neurosecretory Protein (VGF)

[0013] VGF (human VGF, Acc.-No.: O15240) is a secretory peptide precursor that is expressed and processed by neuronal cells [Canu *et al.* 1997]. *In situ* hybridization studies in the adult rat central nervous system have revealed that the VGF mRNA is widely distributed throughout the brain with prominent expression in the hippocampus, entorhinal cortex, and neocortex. Furthermore, it has been shown that VGF transcription and secretion is selectively upregulated by neurotrophins like NGF and BDNF, and by depolarization *in vitro*. Increased BDNF expression can be observed in dentate gyrus and CA3 regions of the hippocampus, which are tissues that appear to die early in Alzheimer Disease pathogenesis.

[0014] WO00/25138 discloses polypeptides as Alzheimer's-disease specific markers being differentially expressed in diseased and healthy tissue.

[0015] WO01/63294 discloses the differential expression of SEQ ID NO:7 as a marker for Bipolar Affective Disorder (BAD).

Description of the invention

[0016] The invention is based on the surprising finding that specific polypeptides are differentially expressed in subjects having Alzheimer's disease when compared to a healthy control group. These differentially expressed polypeptides can be, e.g., detected in samples of cerebrospinal fluid of the subject in which Alzheimer's disease is to be diagnosed. The individual polypeptide used in the invention consisting of the sequence of SEQ ID NO:7 can be detected and/or quantified alone or in combination with other polypeptides.

EP 1 535 076 B1

[0017] The polypeptide markers are defined by their respective molecular weight. Five markers identified by the SAX2 method as described in the Examples show the following molecular masses:

Marker 1 (M1): 4824 ± 20 Da;
 Marker 2 (M2): 7691 ± 20 Da;
 Marker 3 (M3): 11787 ± 20 Da;
 Marker 4 (M4): 11988 ± 20 Da;
 Marker 5 (M5): 13416 ± 20 Da.

[0018] Table one shows the observed molecular weight of polypeptide markers M1 to M5 as determined by SELDI-TOF MS, the amino acid sequences of observed fragments of polypeptide markers M1 to M5, and the protein from which the polypeptide markers M1 to M5 originate.

Table 1:

Marker	SELDI observed MW	Amino Acid Sequence	Protein Name
M1	4823.5 Da ± 1.7	VGEEDEEEAAEAEAEAEER	VGf4.8
M2	7691.4 Da ± 4.9	XXAD (L/I) AGHG (Q/K) EV (L/I) (L/I) R human HGTVV (L/I) TA (L/I) GG (L/I) (L/I) K	myoglobin new variant
M3 M4	11786.9 Da ± 7.6 11988.4 Da ± 5.9	VNHVTLSQLPK VEHSDLFSK IEKVEHSDLFSK SNFLNCYVSGFHPSDIEVDLLK	human beta-2-microglobulin
M5	13416.4 Da ± 9.4	ASNDMYHSR ALDFAVGEYNK RALDFAVGEYNK LVGGPMDASVEEEGVR QIVAGVNYFLDVELGR LVGGPMDASVEEEGVR KQIVAGVNYFLDVELGR TQPNDNCPFHDPHLLK TQPNDNCPFHDPHLLKR SSPGKPPRLVGGPMDASVEEEGVR	human Cystatin C

[0019] The differentially expressed polypeptides can be, e.g., detected in samples of cerebrospinal fluid of the subject in which Alzheimer's disease is to be diagnosed. In addition, depending on the specific embodiment, the source of samples to measure the abundance of the polypeptide markers, can also be blood, serum, or urine, but is not limited to these body compartments.

[0020] As shown in Figure 1, compared to a negative diagnosis (healthy controls), M1 is under-expressed in the CSF of Alzheimer's disease patients ($p < 0.05$), while the markers M2 to M5 are over-expressed in CSF of Alzheimer's disease patients ($p < 0.05$).

[0021] An altered level of one or several polypeptides, compared to the level of polypeptides in healthy control subjects, will allow assessing the state of and/or monitoring the progression of Alzheimer's disease in a subject, will allow monitoring the effectiveness of Alzheimer's disease treatment, and will be useful information for drug development. Furthermore, these biomolecule markers are useful for differentiating Alzheimer's disease from other forms of dementia and neurodegenerative disorders.

[0022] Preferred subjects in which Alzheimer's disease is to be diagnosed or monitored are human subjects. However, diagnosis of Alzheimer's disease according to the invention is also possible with other mammals. If necessary, orthologues of the peptide markers can be used.

[0023] The invention also relates to the use of mass spectrometry (MS) for detecting Alzheimer's disease in human subjects and for assessing the progression of Alzheimer's disease in human subjects by detecting and/or quantifying the amount of specific polypeptides in samples drawn from the subject's body fluids. In a preferred embodiment of the invention, the sample is drawn from the subject's cerebrospinal fluid (CSF).

[0024] Detection and/or quantification of the polypeptides is preferably achieved by quantifying the signal which is detected by MS at specific molecular mass to charge ratios (M/z) which correspond to the M/z ratios of the polypeptides. Preferably, M/z ratios close to the one of the polypeptides are also measured.

[0025] Detection and/or quantification of the polypeptides can also be achieved by using an immunoassay using

specific antibodies raised against the specified marker(s) or polypeptide fragments thereof. Antibodies can be prepared by using the purified marker(s) or fragments thereof, or using synthetic or recombinantly expressed polypeptide(s) consisting of the specific amino acid sequence of the marker(s) using any suitable method known in the art [Coligan 1991]. Such techniques include, but are not limited to, antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or appropriate vectors, as well as preparation of polyclonal and monoclonal antibodies by immunising rabbits or mice [Huse 1989, Ward 1989]. After the antibody is provided, a marker can be detected and/or quantified using any of a number of standard immunological binding assays [US Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168]. Useful assays include, but are not limited to, for example, an enzyme immune assay (EIA) such as enzyme-linked immunosorbent assay (ELISA), a radio immune assay (RIA), a Western blot assay, or a slot or dot blot assay. For a review of the general immunoassays see [Coligan 1991]. Generally, a sample obtained from a subject can be contacted with the antibody that specifically binds the marker. A powerful technique to capture the specified marker (s) from a complex body fluid sample is to use the antibody fixed to solid supports, such as glass or plastic, e.g. microtiter plate, a stick, a bead, or microbead. Alternatively, marker(s) can also be captured from the body fluid sample by the specific antibody immobilised to a probe substrate or a Protein Chip™ array, as described for the SELDI-based immunoassay [Xiao 2001]. After incubating the sample with antibodies, the non-bound material is washed under specified conditions and the antibody-marker complex formed can be detected, using appropriate detection reagents. In an embodiment using the SELDI ProteinChip™ array technique the marker(s) selectively enriched by the immobilised antibody can be detected and quantified by matrix-assisted laser desorption ionisation mass spectrometry.

[0026] Disclosed is

1. a method of assessing the state of Alzheimer's disease in a subject comprising detection of at least one polypeptide comprised in a group of polypeptides consisting of

- i) a polypeptide having a molecular mass of 4824 ± 20 Da,
- ii) a polypeptide having a molecular mass of 7691 ± 20 Da,
- iii) a polypeptide having a molecular mass of 11787 ± 20 Da,
- iv) a polypeptide having a molecular mass of 11988 ± 20 Da, and
- v) a polypeptide having a molecular mass of 13416 ± 20 Da.

Disclosed is further a method of assessing the state of Alzheimer's disease in a subject comprising detection of at least one polypeptide comprised in a group of polypeptides having, respectively, molecular masses of 4824 ± 20 Da, of 7691 ± 20 Da, of 11787 ± 20 Da, of 11988 ± 20 Da, of 13416 ± 20 Da, of 4769 ± 20 Da, of 6958 ± 20 Da, of 6991 ± 20 Da, of 13412 ± 20 Da, of 13787 ± 20 Da, of 17276 ± 20 Da, of 40437 ± 20 Da, of 6895 ± 20 Da, of 6928 ± 20 Da, of 7691 ± 20 Da, of 7769 ± 20 Da, of 7934 ± 20 Da, of 5082 ± 20 Da, of 6267 ± 20 Da, of 6518 ± 20 Da, of 7274 ± 20 Da, and of 8209 ± 20 Da.

Whereas detection of one such polypeptide is in most cases sufficient to reliably diagnose Alzheimer's disease, detection of two or more polypeptides of the invention can increase the sensitivity and robustness of the method. Preferably, 1, 2, 3, 4, 5, 10, and, most preferred, all of said polypeptides will be detected from the same sample. The detection can also be carried out simultaneously with the detection of other polypeptides which are preferably also differentially expressed in subjects having Alzheimer's disease as compared to healthy subjects. "Assessing the state of Alzheimer's disease" shall be understood as diagnosing the presence of Alzheimer's disease in a subject or a patient, as assessing the progression of the disease in a subject or a patient, and/or as assessing the proneness of a subject to develop Alzheimer's disease.

2. Disclosed is further the method of point 1 in which 2, or 3, or 4, or 5 polypeptides of said group of peptides are detected. The invention further relates to a method of point 1 in which 2, or 3, or 4, or 5, or 10 or all polypeptides of said group of peptides are detected.

3. Disclosed is further a method of assessing the state of Alzheimer's disease in a subject comprising detection of at least one polypeptide comprising the sequence of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, and/or SEQ ID NO:16. The invention further relates to a method of assessing the state of Alzheimer's disease in a subject comprising detection of at least one polypeptide comprising the sequence of SEQ ID NO: 1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO: 6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, and/or SEQ ID NO:17. Whereas detection of one such polypeptide is in most cases sufficient to reliably diagnose Alzheimer's disease, detection of two or more polypeptides of the

invention can increase the sensitivity and robustness of the method. Preferably, 1, 2, 3, 4, 5, 10, or all of said polypeptides will be detected from the same sample. The detection can also be carried out simultaneously with the detection of other polypeptides which are preferably also differentially expressed in subjects having Alzheimer's disease as compared to healthy subjects.

5

4. Disclosed is further a method of assessing the state of Alzheimer's disease in a subject comprising detection of at least one polypeptide comprised in a group of polypeptides consisting of

10

- i) human cystatin C,
- ii) human beta-2-microglobulin,
- iii) human myoglobin (new variant),
- iv) a fragment of at least 5, 8, 10, or 20 amino acids of human cystatin C,
- v) a fragment of at least 5, 8, 10, or 20 amino acids of human beta-2-microglobulin, and
- vi) a fragment of at least 5, 8, 10, or 20 amino acids of human myoglobin (new variant).

15

Disclosed is further a method of assessing the state of Alzheimer's disease in a subject comprising detection of at least one polypeptide comprised in a group of polypeptides consisting of

20

- i) human cystatin C,
- ii) human beta-2-microglobulin,
- iii) human myoglobin (new variant),
- iv) human neurosecretory protein VGF,
- v) a fragment of at least 5, 8, 10, or 20 amino acids of human cystatin C,
- vi) a fragment of at least 5, 8, 10, or 20 amino acids of human beta-2-microglobulin,
- vii) a fragment of at least 5, 8, 10, or 20 amino acids of human myoglobin (new variant), and
- viii) a fragment of at least 5, 8, 10, or 20 amino acids of neurosecretory protein VGF.

25

30

Whereas detection of one such polypeptide is in most cases sufficient to reliably diagnose Alzheimer's disease, detection of two or more of said polypeptides can increase the sensitivity and robustness of the method. Preferably, 1, 2, 3, 4, 5, or 6 of said polypeptides will be detected from the same sample. More preferably, 1, 2, 3, 4, 5, 6 or all of said polypeptides will be detected from the same sample. The detection can also be carried out simultaneously with the detection of other polypeptides which are preferably also differentially expressed in subjects having Alzheimer's disease as compared to healthy subjects.

35

5. Disclosed is further a method of investigating the progression of Alzheimer's disease in a subject characterised in that a method of any of points 1 to 4 is performed with at least two distinct samples drawn from the same subject. For this purpose, samples drawn from a subject at different points in time will be analysed. Changes in the amount of the respective polypeptide(s) will allow to draw conclusions on the progression of Alzheimer's disease in the subject

40

6. Disclosed is further a method of any of points 1 to 5, wherein detection of said polypeptide(s) is by SELDI-TOF MS. Other suitable mass spectrometric methods and other methods of detection can alternatively be used. More specifically, the invention relates to a method of any of points 1 to 5, wherein detection of said polypeptide(s) is by SELDI-TOF MS in which the hydrophobic H50, the WCX2, or the IMAC surface is used as a support upon ionisation. Different supports for ionisation yield different sensitivity for specific proteins of interest.

45

7. Disclosed is further a method of any of points 1 to 5, wherein specific antibodies or antibodies recognising said polypeptide(s) are used for detection of said polypeptide(s).

50

8. Disclosed is further a method of any of points 1 to 7, wherein detection is in a sample, comprising CSF of said patient. A sample drawn from a subject can be processed immediately after it has been taken, or it can first be frozen and be analysed later. Samples may also consist of or contain other body fluids such as blood, serum, plasma, urine, seminal plasma, nipple fluid, or cell extracts.

55

9. Disclosed is further a kit comprising a polypeptide having a molecular mass of 4824 ± 20 Da, a polypeptide having a molecular mass of 7691 ± 20 Da, a polypeptide having a molecular mass of 11787 ± 20 Da, a polypeptide having a molecular mass of 11988 ± 20 Da, and/or a polypeptide having a molecular mass of 13416 ± 20 Da. The

invention further relates to a kit comprising a polypeptide having a molecular mass of 4824 ± 20 Da, a polypeptide having a molecular mass of 7691 ± 20 Da, a polypeptide having a molecular mass of 11787 ± 20 Da, a polypeptide having a molecular mass of 11988 ± 20 Da, and a polypeptide having a molecular mass of 13416 ± 20 Da. The invention further relates to a kit comprising polypeptides having a molecular mass of 4824 ± 20 Da, of 7691 ± 20 Da, of 11787 ± 20 Da, of 11988 ± 20 Da, of 13416 ± 20 Da, of 4769 ± 20 Da, of 6958 ± 20 Da, of 6991 ± 20 Da, of 13412 ± 20 Da, of 13787 ± 20 Da, of 17276 ± 20 Da, of 40437 ± 20 Da, of 6895 ± 20 Da, of 6928 ± 20 Da, of 7691 ± 20 Da, of 7769 ± 20 Da, of 7934 ± 20 Da, of 5082 ± 20 Da, of 6267 ± 20 Da, of 6518 ± 20 Da, of 7274 ± 20 Da, and/or of 8209 ± 20 Da. Such a kit can be applied for various purposes, e.g., for use as a standard in one of the above mentioned methods. Said kit can comprise 2, 5, 10, or all of the above polypeptides.

10. Disclosed is further a kit comprising a fragment of at least 5 amino acids of human cystatin C, a fragment of at least 5 amino acids of human beta-2-microglobulin, and a fragment of at least 5 amino acids of human myoglobin. This kit can be applied for various purposes, e.g., for use as a standard in one of the above mentioned methods. Further disclosed is a kit comprising a fragment of at least 5, 10 or 20 amino acids of human cystatin C, a fragment of at least 5, 10 or 20 amino acids of human beta-2-microglobulin, a fragment of at least 5, 10 or 20 amino acids of human myoglobin, and a fragment of at least 5, 10 or 20 amino acids of neurosecretory protein VGF. These kits can be applied for various purposes, e.g., for use as a standard in one of the above mentioned methods.

Brief description of the figures

Figure 1:

[0027] Average intensities of the five marker peptides of Table 1, which are differentially expressed in the diseased group when compared to the control groups.

Examples

[0028] The invention is further described by one or several of the following examples. These examples are not to be understood as restricting the scope of the invention to the examples by any means.

Example 1: Patients evaluation and CSF sampling

[0029] Diagnosis of Alzheimer's disease in human subjects was made according to criteria of the National Institute of Neurologic and Communicative Disorders and Stroke-Alzheimer's disease and Related Disorders Association (NINCDS-ADRDA). The Alzheimer's disease group consisted of 9 patients aged 75 ± 7 years, six men and three women. The group of healthy control subjects consisted of 10 individuals aged 78 ± 14 years, two men and eight women with no history, symptoms or signs of psychiatric or neurological disease.

[0030] Informed consent was given by each patient and the patients' caregivers before the investigation. The study was approved by the local ethics committee. After lumbar puncture, CSF samples were frozen on dry ice immediately upon withdrawal at the bedside in 0.5 ml aliquots and stored at -80°C until analyses.

Example 2: ProteinChip SELDI analysis of CSF on SAX2 chip

[0031] SAX 2 Proteinchip array (CIPHERGEN Biosystems, Fremont, CA, USA) were equilibrated for 5 min with $5\mu\text{l}$ of binding buffer (100mM Na Acetate pH=4.0). The buffer was carefully removed with an handkerchief and $2.5\mu\text{l}$ of binding buffer was added to the wells. Crude CSF samples ($2.5\mu\text{l}$) were added to the wells and incubated for 20 min at room temperature in a humidity chamber on a rocking platform. CSF was removed and the wells were individually washed with $10\mu\text{l}$ of binding buffer for 5 min. The arrays were then placed in a 15ml conical Eppendorf and washed twice with the binding buffer for 5 min. Finally, the chip was rinsed twice with distilled water. Excess of H_2O was removed and while the surface was still moist, two additions per well of $0.5\mu\text{l}$ of sinapinic acid (SPA) (2mg/ml) in 50% (vol/vol) acetonitrile and 0.5% (vol/vol) trifluoroacetic acid was performed and dried. The arrays were then read in a ProteinChip reader system, PBS II series (CIPHERGEN Biosystems). The laser beam was focused on the sample *in vacuo*. This caused the proteins absorbed to the matrix to become ionised and, simultaneously to be desorbed from the Proteinchip array surface. The ionised proteins were detected and their molecular masses were determined according to their time-of-flight (TOF). TOF mass spectra, collected in the positive ion mode were generated using an average of 65 laser shots throughout the spot at a laser power set slightly above threshold (10-15 % higher than the threshold) High mass to acquire was set at 40kDa, optimised from 1 to 15kDa. Spectra were collected and analysed using the CIPHERGEN Proteinchip (version 3.0) software. External calibration of the reader was performed using the "all-in-1" peptide molecular weight standards

EP 1 535 076 B1

(Ciphergen biosystems, Inc.) diluted in the SPA matrix (1:1, vol/vol) and directly applied onto a well. Protein profile comparison was performed after normalisation on total ion current of all the spectra included in the same experiment. The reproducibility was tested by analysing different aliquots of the same CSF sample on 4 different wells of the same proteinchip array (intraassay intrachip reproducibility), on two different chips (intraassay interchip reproducibility) processed in parallel, and reproduced in an other experiment (interassay reproducibility).

[0032] Analysis of CSF samples from 9 patients diagnosed with Alzheimer's disease relative to 10 controls revealed that 5 peaks were significantly differentially expressed between the two groups ($p < 0.05$). The approximate average SELDI mass associated with the five differentially expressed proteins was 4.82kDa, 7.7kDa, 11.8kDa and 12.0kDa and 13.4kDa ($p < 0.05$) (see Table 1, Fig. 1).

Example 3: Strong Anionic exchange chromatography (SAX) purification

[0033] In order to identify the proteins corresponding to these peaks, a fractionation of crude CSF on a SAX spin column was performed. The eluted fractions were analysed by SELDI-TOF MS.

[0034] SAX spin column, lot number SAX2-001116-01, (Ciphergen Biosystems, Fremont, CA, USA) was rehydrated overnight at 4°C in the equilibration buffer (20mM tris(hydroxymethyl)aminomethane hydrochloride (Tris-HCl), 5mM NaCl, pH 9.0). The column was warmed up at room temperature and air bubbles were removed. The equilibration buffer was let flow through column matrix by gravity. Equilibration buffer (0.5ml) was added to the column and passed through the resin twice. Two ml of control CSF was diluted in the equilibration buffer (1:1, vol/vol). Protein sample was loaded to the column by fraction of 0.8ml and allowed to run through the column by gravity until no drops came out of the column. The column was then centrifuged at 150 x g for 1 min. The resin was then washed with an equivalent volume of equilibration buffer. This step was repeated several times in order to load the whole sample onto the resin. Elution of the bound proteins was performed by decreasing the pH. Elution buffer A consisted of 20mM Tris-HCl, 5mM NaCl pH 8.0; elution buffer B = 20mM sodium phosphate pH 7.0; elution buffer C = 20mM sodium phosphate pH 6.0; elution buffer D = 20mM sodium phosphate and citrate pH 5.0; elution buffer E = 20mM sodium phosphate and citrate pH 4.0; elution buffer F = 20mM sodium phosphate and citrate pH 3.4; elution buffer G = 30% acetonitrile in elution buffer F. Elution was performed by applying 2 x 75 µl of the elution buffer and centrifugation at 150 x g for 1 min. Each collected fraction (150 µl) was concentrated on a speed-vac to a volume of 10 µl. Protein profiles were analysed on SELDI-TOF MS using SAX2 Proteinchip arrays. The chip was equilibrated with a binding buffer consisting of 20mM Tris-HCl, 5mM NaCl, pH=9.0. An aliquot of 0.5 µl of each concentrated fraction was applied directly onto 2.5µl of binding buffer per spot and processed as previously described. The rest of the fractions were loaded onto a Tris tricine gel as described below.

[0035] The differentially expressed peak of 13.4kDa was eluted with buffer A (20mM Tris-HCl 5mM NaCl pH 8.0) and B (20mM sodium phosphate pH 7.0). The differentially expressed peaks of 11.8kDa and 12.0kDa were found in the fraction eluted with buffer C (20mM sodium phosphate pH 6.0) and D (20mM sodium phosphate and citrate pH 5.0). The cluster of 7.7kDa was eluted with buffer D (20mM sodium phosphate and citrate pH 5.0) and E (20mM sodium phosphate and citrate pH 4.0).

[0036] Each eluted fraction was loaded on a 16.5% Tris Tricine sodium dodecyl sulfate polyacrylamide gel and electrophoresed (SDS PAGE). After coloration with coomassie blue, the bands seen on the gel confirmed the results obtained by SELDI analysis. The band corresponding to the cluster of 7.7kDa, 11.8kDa and 12.0kDa were cut out. Proteins were extracted as described in Example 6 and identified by Q-TOF. The 7.7kDa peak MS analysis did not match with any known human protein, however, may indicate to a new variant or homologue of myoglobin. Peptide sequences were the following

XXAD(L/I)AGHG(Q/K)EV(L/I)(L/I)R and
HGTVV(L/I)TA(L/I)GG(L/I)(L/I)K.

[0037] The MS analysis of the 11.8kDa and 12.0kDa peaks identified beta-2-microglobulin for both of them.

[0038] Since the cluster of 13.4kDa could not be seen on the Tris Tricine gel, a Tris glycine SDS-PAGE electrophoresis was performed on crude CSF samples. The band corresponding to the beta-2-microglobulin could be easily found on this stained gel. We concluded that the protein migrating just above the beta-2-microglobulin could correspond to the next abundant protein seen on the SELDI profile, namely the 13.4kDa peak. The band was excised from the gel and digested by trypsin before MALDI analysis. The peptide mass fingerprint analysis allowed to identify the Cystatin C. The sequence coverage provided by the analysis was 60%.

Example 4: Monodimensional electrophoresis / Tris Glycine gels

[0039] Twenty µl of CSF were mixed with 10 µl of denaturing Laemmli buffer [Laemmli 1970]. The samples were heated to 95°C for 5 min, and loaded on a 15% T (T = total acrylamide concentration) SDS-polyacrylamide gel according to the method of Laemmli. Gels were stained in a solution containing Coomassie Brilliant Blue R-250 (0.1% w/v) and methanol (50% v/v) for 30 min. Destaining was done in a solution containing methanol (40% v/v) and acetic acid (10% v/v).

Example 5: Monodimensional electrophoresis : Tris Tricine Gels

[0040] Tris tricine SDS-PAGE electrophoresis was performed according to Schagger and von Jagow [1987] using precast 16.5 % T gels (Biorad, Hercules, CA). The anode buffer consisted of 0.2M Tris-HCl, pH 8.9 and the cathode buffer consisted of 0.1M Tris-HCl, 0.1M Tricine, 0.1% SDS, pH 8.25. Samples were diluted in 10 μ l of 50mM Tris-HCl, 4% (w/v) SDS, 12% (w/v) sucrose, 5% (v/v) -mercaptoethanol, and trace of bromophenol blue, pH 6.8. After denaturation at 95°C for 5 min, samples were loaded onto the gel. Gels were run at 80V for 3 hours. After electrophoresis, gels were fixed in 40% methanol, 10% acetic acid for 30min. Gels were then stained with Colloidal blue coomassie G250 overnight and destained in 30% methanol. Bands to be identified were immediately cut, placed in an eppendorf and kept at 4°C until further analysis. The apparent molecular masses were determined by running polypeptide molecular weight (MW) standards: Triosephosphate isomerase MW 26,625; Myoglobin MW 16,950; α -lactalbumin MW 14,437; Aprotinin MW 6,512; Insulin b chain, oxidised MW 3,496 and Bacitracin MW 1,423 (Biorad).

Example 6: Protein digestion and peptide extraction [Bienvenu 1999]

[0041] Fragments of gels containing proteins of interest were cut out for digestion of the proteins with trypsin using previous published procedures [Shevchenko 1996, Hellman 1994, Rosenfeld 1992] and modified as described below. The piece of gel was first destained with 100 μ l of 50 mM ammonium bicarbonate, 30% (v/v) acetonitrile during 15 min at room temperature. Destaining solution was removed and replaced by 25 μ l of 10 mM DL-dithiothreitol (DTT) in 50 mM ammonium bicarbonate and incubated 35 min at 56°C. DTT solution was then replaced by 25 μ l of 55 mM iodoacetamide in 50 mM ammonium bicarbonate and incubated during 45 min at room temperature in the dark. Gel pieces were washed for 10 min with 100 μ l of 50 mM ammonium bicarbonate and for 10 min with 100 μ l of 50 mM ammonium bicarbonate and 30% (v/v) acetonitrile.

[0042] Gel pieces were then dried for 30 min in a Hetovac vacuum centrifuge (HETO, Allerod, Denmark). Dried pieces of gel were rehydrated for 45 min at 4°C in 5-20 μ l of a solution of 50 mM ammonium bicarbonate containing trypsin at 6.25 ng/ μ l. After an over-night incubation at 37°C, gel pieces were dried under high vacuum centrifuge before being rehydrated by the addition of 20 μ l of distilled water and finally dried again in a speed-vac for 30 min. Extraction of the peptides was performed with 20 μ l of 0.1% (v/v) trifluoroacetic acid (TFA) for 20 min at room temperature with occasional shaking. The TFA solution containing the peptides was transferred to a polypropylene tube. A second elution was performed with 20 μ l of 0.1% (v/v) TFA in 50% (v/v) acetonitrile for 20 min at room temperature with occasional shaking. The second TFA solution was pooled with the first one. The volume of the pooled extracts was reduced to 1-2 μ l by evaporation under vacuum. Control extractions (blanks) were performed using pieces of gels devoid of proteins.

Example 7: Protein identification by peptide mass fingerprinting analysis

[0043] 1.5 μ l of sample was placed on a MALDI 100-well target plate. Same volumes of matrix (10 mg/ml α -Cyano-4-hydroxycinnamic acid in 50% (v/v) acetonitrile, 0.1% (v/v) TFA) were added to the previously loaded digest. Samples were dried as quickly as possible using a vacuum container. Mass measurement from liquid solution were conducted with a MALDI-TOF mass spectrometer Voyager™ Elite and Super STR (PerSeptive Biosystems, Framingham MA, USA) equipped with a 337 nm nitrogen laser. The analyser was used in the reflectron mode at an accelerating voltage of 20 kV, a delayed extraction parameter of 100-140 ns and a low mass gate of 850 Da. Laser power was set slightly above threshold (10-15 % higher than the threshold) for molecular ion production. Spectra were obtained by summation of 10 to 256 consecutive laser shots. Masses of the 60 highest peaks were extracted from the spectra and used for protein identification using the SmartIdent peptide mass fingerprint tool [Gras 1999]. The research was conducted against SWISS-PROT and TrEMBL databases. The query was made for the human, the minimum number of matched masses was 4, the maximal tolerance for masses was 50 ppm after an internal calibration using autolysis product of trypsin, at most one missed cleavage for tryptic peptides was allowed, and the modifications accepted were carboxymethylation with iodoacetamide of cysteines and artefactual oxidation of methionines.

Example 8: Protein identification by peptide fragmentation analysis

[0044] Prior to nanoLC (LC = liquid chromatography) separation, the volumes of peptide containing solutions were adjusted to 7 μ l by addition of a 0.1 % (v/v) formic acid solution. Samples were settled in a Triathlon autosampler (Spack, Emmen, Holland). For each experiment, 5 μ l of peptide containing solution were injected on a C18 reverse phase column of 75 μ m inner diameter (YMS-ODS-AQ200, Michrom Bioresource, Auburn, CA). Peptides were eluted with an acetonitrile (ACN) gradient in the presence of 0.1 % (v/v) formic acid, using SunFlow pumps (SunChrom, Friderichsdorf, Germany). A flow splitter was used in order to decrease the flow rate after the pumps from 200 to 0.4 μ l/min. Peptides were analysed with a quadrupole time-of-flight (Q-TOF) mass spectrometer (Micromass, Wythenshawe, England). A 2700 V tension

was applied on the nanoelectrospray capillary (New Objective, Woburn, MA, USA). Argon was used as collision gas. The collision energy was settled as a function of the precursor ion mass. MS/MS spectra were acquired by automatic switching between MS and MS/MS mode. Acquired MS/MS data were converted in a compatible format (DTA files) by ProteinLynx software (Micromass, Wythenshawe, England) and analysed using conventional search engines against SWISS-PROT, TrEMBL, NCBIInr and EST databases. In cases of manual interpretation of MS/MS data, identification was performed by sequence only search.

[0045] It was found that marker M5 was a fragment of Cystatin C, markers M3 and M4 were isoforms of beta-2-microglobulin and M2 was a new variant or homologue of myoglobin. Marker M1 was found to be a fragment of the neurosecretory protein VGF.

Example 9: Statistical Analysis

[0046] P-values were calculated using standard statistical methods known to the person skilled in the art. P-values smaller than 0.05 were considered to be statistically significant.

Example 10: Isolation of the 4.8 kDa Fragment (Marker M1)

[0047] Some CSF samples from control patients were fractionated by Centricon 30 filtration device (Millipore Corp., Bedford, MA) in order to remove the protein with a molecular weight higher than 30 kDa. The salt and polypeptide with a molecular weight lower than 3 kDa were removed using a Centricon 3 (Millipore Corp., Bedford, MA). The Centricon 3 was then washed with ultrapure distilled water. In that wash fraction, the 4.82 kDa was found to be the major component. This liquid fraction was first reduced with a 10 mM solution of 1,4-Dithioerythritol for 1h at 56°C, then alkylated with 54 mM iodoacetamide for 45 min at room temperature. Finally, the polypeptide was digested with 6 mg/l trypsin overnight at 37°C. This liquid fraction was analysed by nanoLC and Q-TOF as previously described.

Example 11: Different Surface Materials for SELDI-TOF Analysis

[0048] Using SELDI-TOF, an analysis of 10 CSF samples from AD patients and 10 controls was performed on three different surfaces: the hydrophobic H50, the WCX2, and the IMAC surface (CIPHERGEN Biosystems, Fremont, California, USA, resp.). Seven differentially expressed peaks were found on the H50, five markers on the WCX2, and five markers on the IMAC surface. A diagnostic test using the markers on the H50 chip revealed a specificity and sensitivity of 100% and 70%, respectively. The combination of the markers found on H50 and WCX2 gave a specificity and sensitivity of 100% and 80%. Finally, the combination of the markers found on H50, WCX2 and IMAC gave a specificity and sensitivity of 100% and 90%.

[0049] The average masses of the differentially expressed polypeptides as determined by SELDI-TOF using different surface materials were as follows:

Surface hydrophobic H50: 7 peaks

Marker 1: 4769 ± s.d. Da

Marker 2: 6958 ± s.d. Da

Marker 3: 6991 ± s.d. Da

Marker 4: 13412 ± s.d. Da

Marker 5: 13787 ± s.d. Da

Marker 6: 17276 ± s.d. Da

Marker 7: 40437 ± s.d. Da

Surface IMAC Cu: 5 peaks

Marker 1: 6895 ± s.d. Da

Marker 2: 6928 ± s.d. Da

Marker 3: 7691 ± s.d. Da

Marker 4: 7769 ± s.d. Da

Marker 5: 7934 ± s.d. Da

Surface WCX2: 5 peaks

Marker 1: 5082 ± s.d. Da

Marker 2: 6267 ± s.d. Da

Marker 3: 6518 ± s.d. Da

Marker 4: 7274 ± s.d. Da

Marker 5: 8209 ± s.d. Da

[0050] The standard deviation (s.d.) is 20 Da for each marker above. However, the standard deviation can also be 40 Da, or 10 Da, or 5 Da for each marker above.

Bibliography

- 5
- [0051]** Adam, B.L. et al. 2001, *Proteomics* 1(10): 1264-1270
- [0052]** Asgeirsson B, Haebel S, Thorsteinsson L, Helgason E, Gudmundsson KO, Gudmundsson G, Roepstorff P. Hereditary cystatin C amyloid angiopathy: monitoring the presence of the Leu68Gln cystatin C variant in cerebrospinal fluid and monocyte cultures by MS. *Biochem J.* 1998, 329, 497-503.
- 10 **[0053]** Beyer K, Lao JI, Gomez M, Riutort N, Latorre P, Mate JL, Ariza A. Alzheimer's disease and the cystatin C gene polymorphism: an association study. *Neuroscience letters.* 2001, 315, 17-20.
- [0054]** Canu N, Possenti R, Ricco AS, Rocchi M, Levi A. Cloning, structural organization analysis, and chromosomal assignment of the human gene for the neurosecretory protein VGF. *Genomics* 1997, 45 (2), 443-446.
- 15 **[0055]** Cohen DH, Feiner H, Jensson O, Frangione B. Amyloid fibril in hereditary cerebral hemorrhage with amyloidosis (HCHWA-I) is related to the gastroentero-pancreatic neuroendocrine protein. *J. Exp. Med.* 1983, 158, 623-628.
- [0056]** *Current Protocols in Immunology*, Eds. Coligan JE, Kruisbeek AM, Margulies DH, Shevach EM, and Strober W., John Wiley & Sons Inc. 1991
- [0057]** Crawford U, Freeman MJ, Schinka JA, Abdullah LI, Gold M, Hartman R, Krivian K, Morris MD, Richards D, Duara R, Anand R, Mullan MJ. A polymorphism in the cystatin C gene is a novel risk factor for late-onset Alzheimer's disease. *Neurology.* 2000, 55, 763-768.
- 20 **[0058]** Davies, H.A. 2000. The ProteinChip(R) System from Ciphergen: A new technique for rapid, micro-scale protein biology. *J. Molecular Medicine*, 78 (7):B29
- Deng A, Irizarry MC, Nitsch RM, Growdon JH, Rebeck GW. Elevation of Cystatin C in susceptible neurons in Alzheimer's disease. *Am. J. Pathol.* 2001, 159, 1061-1068.
- [0059]** Ernerudh J, Olsson T, Berlin G, von Schenck H. Cerebrospinal fluid immunoglobulins and beta 2-microglobulin in lymphoproliferative and other neoplastic diseases of the central nervous system. *Arch Neurol.* 1987, 44(9), 915-20.
- 25 **[0060]** Finckh U, Von der Kammer H, Velden J, Michel T, Andresen B, Deng A, Zhang J, Muller-Thomsen T, Zuchowski K, Menzer G, Mann U, Papassotiropoulos A, Heun R, Zurdel J, Holst F, benussi L, Stoppe G, Reiss J, Miserez AR, Staehelin HB, Rebeck W, Hyman BT, Binetti G, Hock C, Growdon JH, Nitsch RM. Genetic association of a cystatin C gene polymorphism with late-onset Alzheimer's disease. *Arch Neurol.* 2000, 57, 1579-1583.
- 30 **[0061]** Francis, P.T. et al. *J. Neurol. Neurosurg. Psychiatry* 1999, 66(2): 137-147
- Fung, ET et al. Protein biochips for differential profiling. *Current Opinion in Biotechnology*, 2001, 12(1): 65-69
- [0062]** Ghiso J, Jensson O, Frangione B. Amyloid fibrils in hereditary cerebral hemorrhage with amyloidosis of Icelandic type is a variant of gamma-trace basic protein (cystatin C). *Proc Natl Acad Sci USA.* 1986;83(9):2974-8.
- 35 **[0063]** Ghiso J, Pons-Estel B, Frangione B. Hereditary cerebral amyloid angiopathy: the amyloid fibrils contain a protein which is a variant of cystatin C, an inhibitor of lysosomal cysteine proteases. *Biochem Biophys Res Commun.* 1986;136 (2):548-54.
- [0064]** Gras R, Müller M, Gasteiger E, Gay S, Binz PA, Bienvenut WV, Hoogland C, Sanchez JC, Bairoch A, Hochstrasser DF, Appel R. Improving protein identification from peptide mass fingerprinting through a parameterised multi-level scoring algorithm and an optimised peak detection. *Electrophoresis* 1999, 20: 3535-3550
- 40 **[0065]** Grubb AO, Jensson O, Gudmundsson G, Arnason A, Lofberg H, Malm J. Abnormal metabolism of g-trace alkaline microprotein. The basic defect in hereditary cerebral hemorrhage with amyloidosis. *N. Engl. J. Med* 1984, 311, 1547-1549.
- [0066]** Hellman U, Wernstedt C, Góñez, J, Heldin CH. Improvement of an "In-Gel" digestion procedure for the micro-preparation of internal protein fragments for amino acid sequencing. *Anal. Biochem.* 1995, 224, 451-5.
- 45 **[0067]** Hoekman K, Van Nieuwkoop JA, Willemze R. The significance of beta-2 microglobulin in clinical medicine. *Neth J Med.* 1985;28(12):551-7.
- [0068]** Huse WD, Sastry L, Iverson SA, Kang AS, Altling-Mees M, Burton DR, Benkovic SJ, Lemer RA. Generation of a large combinatorial library of the immunoglobulin repertoire in phage lambda. 1989, *Science* 246: 1275-1281.
- 50 **[0069]** Issaq HJ, Veenstra TD, Conrads TP, Felschow D. The SELDI-TOF MS approach to proteomics: protein profiling and biomarker identification. *Biochem. Biophys. Res. Commun.* 2002, 292 (3): 587-592
- [0070]** Kalman J, Marki-Zay J, Juhasz A, Santha A, Dux L, Janka Z. Serum and Cerebrospinal fluid cystatin C levels in vascular and Alzheimer's dementia. *Acta Neurol. Scand.* 2000, 101, 279-282.
- [0071]** Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 1970 Aug 15;227(259):680-5.
- 55 **[0072]** Leung-Tack J, Tavera C, Gensac MC, Martinez J, Colle A. Modulation of phagocytosis-associated respiratory burst by human cystatin C: role of the N-terminal tetrapeptide Lys-Pro-Pro-Arg. *Exp. Cell Research.* 1990, 188, 16-22.
- [0073]** Levy E, Sastre M, Kumar A, Gallo G, Piccardo P, Ghetti B, Tagliavini F. Codeposition of cystatin C with amyloid-beta protein in the brain of Alzheimer's disease patients. *J. Neuropathol Exp Neurol* 2001, 60, 94-104

[0074] Merchant M, Weinberger SR.. Recent advancements in surface-enhanced laser desorption/ionization-time of flight-mass spectrometry. Electrophoresis, 2000, 21: 1164-1167

[0075] Popovic T, Brzin J, Ritonja A, Turk V. Different forms of human Cystatin C. Biol. Chem. Hoppe-Seyler. 1990, 371, 575-580.

5 [0076] Raymackers J, Daniels A, DeBrabandere V, Missiaen C, Dauwe M, Verhaert P, Vanmechelen E, Meheus L. Identification of two dimensionally separated human cerebrospinal fluid proteins by N-terminal sequencing, matrix-assisted laser-desorption/ionization-mass spectrometry, nanoliquid chromatography-electrospray ionization-time of flight-mass spectrometry, and tandem mass spectrometry. Electrophoresis, 2000, 21, 2266-2283.

10 [0077] The Ronald and Nancy Reagan Research Institute of the Alzheimer's Association and the National Institute on Aging Working Group. Consensus Report of the Working Group on: "Molecular and Biochemical Markers of Alzheimer's Disease". Neurobiology of Aging, 1988, 19 (2): 109-116.

[0078] Robles A.. Some Remarks on biological markers of Alzheimer's Disease. Neurobiology of Aging, 1998, 19 (2): 153-157

15 [0079] Rosenfeld J, Capdevielle J, Guillemot JC, Ferrara P. In-gel digestion of proteins for internal sequence analysis after one- or two-dimensional gel electrophoresis. Anal. Biochem. 1992, 203, 173-9.

[0080] Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. Anal. Chem. 1996, 68, 850-8.

[0081] Teunissen CE, de Vente J, Steinbusch HWM, De Bruijn C.. Neurobiology of Aging, 2002, 23: 485-508

20 US Patent 4,366,241

US Patent 4,376,110

US Patent 4,517,288

US Patent 4,837,168

25 [0082] Ward ES et al., Binding activities of a repertoire of single immunoglobulin variable domains secreted from Escherichia coli. Nature 1989, 341, 544-546

[0083] Wei L, Berman Y, Castano EM, Cadene M, Beavis RC, Devi L, Levy E. Instability of the amyloidogenic cystatin C variant of hereditary cerebral hemorrhage with amyloidosis, Icelandic type. J. Biol. Chem. 1998, 273, 11806-11814.

30 [0084] Wulfskuhle JD, McLean KC, Paweletz CP, Sgroi DC, Trock BJ, Steeg PS, Petricoin EF . New approaches to proteomic analysis of breast cancer. Proteomics 2001, 1 (10): 1205-1215

[0085] Xiao Z, Adam BL, Cazares LH, Clements MA, Davis JW, Schellhammer PF, Dalmasso EA, Wright GL. Quantitation of serum prostate-specific membrane antigen by a novel protein biochip immunoassay discriminates benign from malignant prostate disease..2001, Cancer Research 61 (16): 6029-6033

35 SEQUENCE LISTING

[0086]

<110> Bayer AG

40 <120> Biomarkers for the diagnosis of Alzheimer's Disease

<130> LeA 36 293

<160> 17

<170> Patent In version 3.1

45

<210> 1

<211> 15

<212> PRT

<213> Homo sapiens

50

<220>

<221> misc_feature

<222> (1)..(2)

<223> Xaa = any amino acid

<220>

55

<221> misc_feature

<222> (5)..(5)

<223> Xaa = L or I

<220>

EP 1 535 076 B1

<221> misc_feature
 <222> (10)..(10)
 <223> Xaa = Q or K
 <220>
 5 <221> misc_feature
 <222> (13)..(13)
 <223> Xaa = L or I
 <220>
 10 <221> misc_feature
 <222> (14)..(14)
 <223> Xaa = L or I

 <400> 1

 15 Xaa Xaa Ala Asp Xaa Ala Gly His Gly Xaa Glu Val Xaa Xaa Arg
 1 5 10 15

 20 <210> 2
 <211> 14
 <212> PRT
 <213> Homo sapiens
 <220>
 25 <221> misc_feature
 <222> (1)..(14)
 <223> Xaa = L or I
 <220>
 30 <221> misc_feature
 <222> (1)..(14)
 <223> Xaa = Leu or Ile

 <400> 2

 35 His Gly Thr Val Val Xaa Thr Ala Xaa Gly Gly Xaa Xaa Lys
 1 5 10

 40 <210> 3
 <211> 10
 <212> PRT
 <213> Homo sapiens

 45 <400> 3

 Val Asn His Val Thr Leu Ser Gln Pro Lys
 1 5 10
 50

 <210> 4
 <211> 10
 <212> PRT
 55 <213> Homo sapiens

 <400> 4

EP 1 535 076 B1

Val Glu His Ser Asp Leu Ser Phe Ser Lys
1 5 10

5

<210> 5
<211> 13
<212> PRT
<213> Homo sapiens

10

<400> 5

Ile Glu Lys Val Glu His Ser Asp Leu Ser Phe Ser Lys

15

1 5 10

20

<210> 6
<211> 22
<212> PRT
<213> Homo sapiens

<400> 6

25

Ser Asn Phe Leu Asn Cys Tyr Val Ser Gly Phe His Pro Ser Asp Ile
1 5 10 15

30

Glu Val Asp Leu Leu Lys
20

35

<210> 7
<211> 9
<212> PRT
<213> Homo sapiens

<400> 7

40

Ala Ser Asn Asp Met Tyr His Ser Arg
1 5

45

<210> 8
<211> 11
<212> PRT
<213> Homo sapiens

<400> 8

50

Ala Leu Asp Phe Ala Val Gly Glu Tyr Asn Lys
1 5 10

55

<210> 9
<211> 12
<212> PRT

EP 1 535 076 B1

<213> Homo sapiens

<400> 9

5

Arg Ala Leu Asp Phe Ala Val Gly Glu Tyr Asn Lys
1 5 10

10

<210> 10
<211> 16
<212> PRT
<213> Homo sapiens

15

<400> 10

Leu Val Gly Gly Pro Met Asp Ala Ser Val Glu Glu Glu Gly Val Arg
1 5 10 15

20

<210> 11
<211> 16
<212> PRT
<213> Homo sapiens

25

<400> 11

30

Gln Ile Val Ala Gly Val Asn Tyr Phe Leu Asp Val Glu Leu Gly Arg
1 5 10 15

35

<210> 12
<211> 17
<212> PRT
<213> Homo sapiens

40

<400> 12

Leu Val Gly Gly Pro Met Asp Ala Ser Val Glu Glu Glu Gly Val Arg
1 5 10 15

45

Arg

<210> 13
<211> 17
<212> PRT
<213> Homo sapiens

50

<400> 13

55

Lys Gln Ile Val Ala Gly Val Asn Tyr Phe Leu Asp Val Glu Leu Gly
1 5 10 15
Arg

Claims

1. Method of assessing the state of Alzheimer's disease in a subject comprising detection of a polypeptide consisting of the sequence of SEQ ID NO:7 in a sample from the subject.
2. Method of assessing the state of Alzheimer's disease according to claim 1, wherein at least one further polypeptide selected from the group consisting of polypeptides comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17 or selected from the group consisting of human cystatin C, human beta-2-microglobulin, the 7,7 kD homologue of human myoglobin, neurosecretory protein VGF, a fragment of at least 5 amino acids of human cystatin C, a fragment of at least 5 amino acids of human beta-2-microglobulin, a fragment of at least 5 amino acids of the 7,7 kD homologue of human myoglobin, and a fragment of at least 5 amino acids of neurosecretory protein VGF is detected.
3. Method of investigating the progression of Alzheimer's disease in a subject **characterised in that** a method of claim 1 or claim 2 is performed with at least two distinct samples drawn from the same subject.
4. Method of any of claims 1 to 3, wherein detection of said polypeptide(s) is by SELDI-TOF MS.
5. Method of any of claims 1 to 3, wherein specific antibodies or antibodies recognising said polypeptides are used for detection of said polypeptide(s).
6. Method of any of claims 1 to 5, wherein detection is in a sample comprising CSF, blood, serum, plasma, urine, seminal plasma, nipple fluid, and/or cell extracts of said patient.

Patentansprüche

1. Verfahren zur Beurteilung des Standes einer Alzheimer-Erkrankung bei einem Individuum, umfassend das Detektieren eines aus der Sequenz von Seq.-ID Nr. 7 bestehenden Polypeptids in einer Probe von dem Individuum.
2. Verfahren zur Beurteilung des Stands einer Alzheimer-Erkrankung nach Anspruch 1, worin zumindest ein weiteres Polypeptid, das aus der aus den Polypeptiden der Seq.-ID Nr. 1, Seq.-ID Nr. 2, Seq.-ID Nr. 3, Seq.-ID Nr. 4, Seq.-ID Nr. 5, Seq.-ID Nr. 6, Seq.-ID Nr. 8, Seq.-ID Nr. 9, Seq.-ID Nr. 10, Seq.-ID Nr. 11, Seq.-ID Nr. 12, Seq.-ID Nr. 13, Seq.-ID Nr. 14, Seq.-ID Nr. 15, Seq.-ID Nr. 16, Seq.-ID Nr. 17 bestehenden Gruppe ausgewählt ist oder aus der aus menschlichen Cystatin C, menschlichem β -2-Mikroglobulin, dem 7,7-kD-Homolog von menschlichem Myoglobin, dem neurosekretorischen Protein VGF, einem Fragment aus zumindest 5 Aminosäuren von menschlichem Cystatin C, einem Fragment aus zumindest 5 Aminosäuren von menschlichem β -2-Mikroglobulin, einem Fragment aus zumindest 5 Aminosäuren des 7,7-kD-Homologs von menschlichem Myoglobin und einem Fragment aus zumindest 5 Aminosäuren des neurosekretorischen Proteins VGF bestehenden Gruppe ausgewählt ist, detektiert wird.
3. Verfahren zur Untersuchung des Verlaufs einer Alzheimer-Erkrankung bei einem Individuum, **dadurch gekennzeichnet, dass** ein Verfahren nach Anspruch 1 oder Anspruch 2 mit zumindest zwei unterschiedlichen Proben durchgeführt wird, die dem gleichen Individuum abgenommen wurden.
4. Verfahren nach einem der Ansprüche 1 bis 3, worin die Detektion des Polypeptids/der Polypeptide mittels SELDI-TOF-MS erfolgt.
5. Verfahren nach einem der Ansprüche 1 bis 3, worin spezifische Antikörper oder Antikörper, welche die Polypeptide erkennen, zur Detektion des Polypeptids/der Polypeptide eingesetzt werden.
6. Verfahren nach einem der Ansprüche 1 bis 5, worin die Detektion in einer Probe erfolgt, die Zerebrospinalflüssigkeit, Blut, Plasma, Urin, Samenplasma, Brustwarzenflüssigkeit und/oder Zellextrakte vom Patienten umfasst.

Revendications

1. Méthode pour évaluer l'état de la maladie d'Alzheimer chez un sujet, comprenant la détection d'un polypeptide

EP 1 535 076 B1

consistant en la séquence de la SEQ ID N° 7 dans un échantillon provenant du sujet.

- 5
2. Méthode pour évaluer l'état de la maladie d'Alzheimer suivant la revendication 1, dans laquelle au moins un polypeptide supplémentaire choisi dans le groupe consistant en les polypeptides comprenant les SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5, SEQ ID N° 6, SEQ ID N° 8, SEQ ID N° 9, SEQ ID N° 10, SEQ ID N° 11, SEQ ID N° 12, SEQ ID N° 13, SEQ ID N° 14, SEQ ID N° 15, SEQ ID N° 16, SEQ ID N° 17 ou choisi dans le groupe consistant en la cystatine C humaine, la bêta-2-microglobuline humaine, l'homologue de 7,7 kD de la myoglobine humaine, la protéine neurosécrétrice VGF, un fragment d'au moins 5 aminoacides de la cystatine C humaine, un fragment d'au moins 5 aminoacides de la bêta-2-microglobuline humaine, un fragment d'au moins 5 aminoacides de l'homologue de 7,7 kD de la myoglobine humaine et un fragment d'au moins 5 aminoacides de la protéine neurosécrétrice VGF, est détecté.
- 10
3. Méthode d'étude de la progression de la maladie d'Alzheimer chez un sujet, **caractérisée en ce qu'**une méthode de la revendication 1 ou la revendication 2 est mise en oeuvre avec au moins deux échantillons distincts prélevés chez le même sujet.
- 15
4. Méthode suivant l'une quelconque des revendications 1 à 3, dans laquelle la détection dudit ou desdits polypeptides est effectuée par SELDI-TOF MS.
- 20
5. Méthode suivant l'une quelconque des revendications 1 à 3, dans laquelle des anticorps spécifiques ou anticorps reconnaissant lesdits polypeptides sont utilisés pour la détection dudit ou desdits polypeptides.
- 25
6. Méthode suivant l'une quelconque des revendications 1 à 5, dans laquelle la détection est effectuée dans un échantillon comprenant du liquide céphalo-rachidien, du sang, du sérum, du plasma, de l'urine, du plasma séminal, du fluide mammaire et/ou des extraits cellulaires dudit patient.

30

35


40

45

50

55

Fig 1:



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 0025138 A [0014]
- WO 0163294 A [0015]
- US 4366241 A [0025] [0081]
- US 4376110 A [0025] [0081]
- US 4517288 A [0025] [0081]
- US 4837168 A [0025] [0081]

Non-patent literature cited in the description

- **ADAM, B.L. et al.** *Proteomics*, 2001, vol. 1 (10), 1264-1270 [0051]
- **ASGEIRSSON B ; HAEBEL S ; THORSTEINSSON L ; HELGASON E ; GUDMUNDSSON KO ; GUDMUNDSSON G ; ROEPSTORFF P.** Hereditary cystatin C amyloid angiopathy: monitoring the presence of the Leu68Gln cystatin C variant in cerebrospinal fluid and monocyte cultures by MS. *Biochem J.*, 1998, vol. 329, 497-503 [0052]
- **BEYER K ; LAO JI ; GOMEZ M ; RIUTORT N ; LATORRE P ; MATE JL ; ARIZA A.** Alzheimer's disease and the cystatin C gene polymorphism: an association study. *Neuroscience letters*, 2001, vol. 315, 17-20 [0053]
- **CANU N ; POSSENTI R ; RICCO AS ; ROCCHI M ; LEVI A.** Cloning, structural organization analysis, and chromosomal assignment of the human gene for the neurosecretory protein VGF. *Genomics*, 1997, vol. 45 (2), 443-446 [0054]
- **COHEN DH ; FEINER H ; JENSSON O ; FRANGIONE B.** Amyloid fibril in hereditary cerebral hemorrhage with amyloidosis (HCHWA-I) is related to the gastroentero-pancreatic neuroendocrine protein. *J. Exp. Med.*, 1983, vol. 158, 623-628 [0055]
- Current Protocols in Immunology. John Wiley & Sons Inc, 1991 [0056]
- **CRAWFORD U ; FREEMAN MJ ; SCHINKA JA ; ABDULLAHLI ; GOLD M ; HARTMAN R ; KRIVIAN K ; MORRIS MD ; RICHARDS D ; DUARA R.** A polymorphism in the cystatin C gene is a novel risk factor for late-onset Alzheimer's disease. *Neurology*, 2000, vol. 55, 763-768 [0057]
- **DAVIES, H.A.** The ProteinChip(R) System from Ciphergen: A new technique for rapid, micro-scale protein biology. *J. Molecular Medicine*, 2000, vol. 78 (7), B29 [0058]
- **DENG A ; IRIZARRY MC ; NITSCH RM ; GROWDON JH ; REBECK GW.** Elevation of Cystatin C in susceptible neurons in Alzheimer's disease. *Am. J. Pathol.*, 2001, vol. 159, 1061-1068 [0058]
- **ERNERUDH J ; OLSSON T ; BERLIN G ; VON SCHENCK H.** Cerebrospinal fluid immunoglobulins and beta 2-microglobulin in lymphoproliferative and other neoplastic diseases of the central nervous system. *Arch Neurol.*, 1987, vol. 44 (9), 915-20 [0059]
- **FINCKH U ; VON DER KAMMER H ; VELDEN J ; MICHEL T ; ANDRESEN B ; DENG A ; ZHANG J ; MULLER-THOMSEN T ; ZUCHOWSKI K ; MENZER G.** Genetic association of a cystatin C gene polymorphism with late-onset Alzheimer's disease. *Arch Neurol.*, 2000, vol. 57, 1579-1583 [0060]
- **FRANCIS, P.T. et al.** *J. Neurol. Neurosurg. Psychiatry*, 1999, vol. 66 (2), 137-147 [0061]
- **FUNG, ET et al.** Protein biochips for differential profiling. *Current Opinion in Biotechnology*, 2001, vol. 12 (1), 65-69 [0061]
- **GHISO J ; JENSSON O ; FRANGIONE B.** Amyloid fibrils in hereditary cerebral hemorrhage with amyloidosis of Icelandic type is a variant of gamma-trace basic protein (cystatin C). *Proc Natl Acad Sci USA.*, 1986, vol. 83 (9), 2974-8 [0062]
- **GHISO J ; PONS-ESTEL B ; FRANGIONE B.** Hereditary cerebral amyloid angiopathy: the amyloid fibrils contain a protein which is a variant of cystatin C, an inhibitor of lysosomal cysteine proteases. *Biochem Biophys Res Commun*, 1986, vol. 136 (2), 548-54 [0063]
- **GRAS R ; MÜLLER M ; GASTEIGER E ; GAY S ; BINZ PA ; BIENVENUT WV ; HOOGLAND C ; SANCHEZ JC ; BAIROCH A ; HOCHSTRASSER DF.** Improving protein identification from peptide mass fingerprinting through a parameterised multi-level scoring algorithm and an optimised peak detection. *Electrophoresis*, 1999, vol. 20, 3535-3550 [0064]
- **GRUBB AO ; JENSSON O ; GUDMUNDSSON G ; ARNASON A ; LOFBERG H ; MALM J.** Abnormal metabolism of g-trace alkaline microprotein. The basic defect in hereditary cerebral hemorrhage with amyloidosis. *N. Engl. J. Med.*, 1984, vol. 311, 1547-1549 [0065]

- **HELLMAN U ; WERNSTEDT C ; GÓÑEZ, J ; HELDIN CH.** Improvement of an "In-Gel" digestion procedure for the micropreparation of internal protein fragments for amino acid sequencing. *Anal. Biochem.*, 1995, vol. 224, 451-5 [0066]
- **HOEKMAN K ; VAN NIEUWKOOP JA ; WILLEMZE R.** The significance of beta-2 microglobulin in clinical medicine. *Neth J Med.*, 1995, vol. 28 (12), 551-7 [0067]
- **HUSEWD ; SASTRY L ; IVERSON SA ; KANG AS ; ALTING-MEES M ; BURTON DR ; BENKOVIC SJ ; LEMER RA.** Generation of a large combinatorial library of the immunoglobulin repertoire in phage lambda. *Science*, 1989, vol. 246, 1275-1281 [0068]
- **ISSAQ HJ ; VEENSTRA TD ; CONRADS TP ; FELSCHOW D.** The SELDI-TOF MS approach to proteomics: protein profiling and biomarker identification. *Biochem. Biophys. Res. Commun.*, 2002, vol. 292 (3), 587-592 [0069]
- **KALMAN J ; MARKI-ZAY J ; JUHASZ A ; SANTHA A ; DUXL ; JANKAZ.** Serum and Cerebrospinal fluid cystatin C levels in vascular and Alzheimer's dementia. *Acta Neurol. Scand.*, 2000, vol. 101, 279-282 [0070]
- **LAEMMLI UK.** Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 15 August 1970, vol. 227 (259), 680-5 [0071]
- **LEUNG-TACK J ; TAVERA C ; GENSAC MC ; MARTINEZ J ; COLLE A.** Modulation of phagocytosis-associated respiratory burst by human cystatin C: role of the N-terminal tetrapeptide Lys-Pro-Pro-Arg. *Exp. Cell Research*, 1990, vol. 188, 16-22 [0072]
- **LEVY E ; SASTRE M ; KUMAR A ; GALLO G ; PICCARDO P ; GHETTI B ; TAGLIAVINI F.** Codeposition of cystatin C with amyloid-beta protein in the brain of Alzheimer's disease patients. *J. Neuropathol Exp Neurol*, 2001, vol. 60, 94-104 [0073]
- **MERCHANT M ; WEINBERGER SR.** Recent advancements in surface-enhanced laser desorption/ionization-time of flight-mass spectrometry. *Electrophoresis*, 2000, vol. 21, 1164-1167 [0074]
- **POPOVIC T ; BRZIN J ; RITONJA A ; TURK V.** Different forms of human Cystatin. *C. Biol. Chem. Hoppe-Seyler*, 1990, vol. 371, 575-580 [0075]
- **RAYMACKERS J ; DANIELS A ; DEBRABANDERE V ; MISSIAEN C ; DAUWE M ; VERHAERT P ; VANMECHELEN E ; MEHEUS L.** Identification of two dimensionally separated human cerebrospinal fluid proteins by N-terminal sequencing, matrix-assisted laser-desorption/ionization-mass spectrometry, nanoliquid chromatography-electrospray ionization-time of flight-mass spectrometry, and tandem mass spectrometry. *Electrophoresis*, 2000, vol. 21, 2266-2283 [0076]
- **Molecular and Biochemical Markers of Alzheimer's Disease.** *Neurobiology of Aging*, 1988, vol. 19 (2), 109-116 [0077]
- **ROBLES A.** Some Remarks on biological markers of Alzheimer's Disease. *Neurobiology of Aging*, 1998, vol. 19 (2), 153-157 [0078]
- **ROSENFELD J ; CAPDEVIELLE J ; GUILLEMOT JC ; FERRARA P.** In-gel digestion of proteins for internal sequence analysis after one- or two-dimensional gel electrophoresis. *Anal. Biochem.*, 1992, vol. 203, 173-9 [0079]
- **SHEVCHENKO A ; WILM M ; VORM O ; MANN M.** Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.*, 1996, vol. 68, 850-8 [0080]
- **TEUNISSEN CE ; DE VENDE J ; STEINBUSCH HWM ; DE BRUIJN C.** *Neurobiology of Aging*, 2002, vol. 23, 485-508 [0081]
- **WARD ES et al.** Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli*. *Nature*, 1989, vol. 341, 544-546 [0082]
- **WEIL ; BERMAN Y ; CASTANO EM ; CADENE M ; BEAVIS RC ; DEVIL ; LEVY E.** Instability of the amyloidogenic cystatin C variant of hereditary cerebral hemorrhage with amyloidosis, Icelandic type. *J. Biol. Chem.*, 1998, vol. 273, 11806-11814 [0083]
- **WULFKUHLE JD ; MCLEAN KC ; PAWELETZ CP ; SGROI DC ; TROCK BJ ; STEEG PS ; PETRICOIN EF.** New approaches to proteomic analysis of breast cancer. *Proteomics*, 2001, vol. 1 (10), 1205-1215 [0084]
- **XIAO Z ; ADAM BL ; CAZARES LH ; CLEMENTS MA ; DAVIS JW ; SCHELLHAMMER PF ; DALMASSO EA ; WRIGHT GL.** Quantitation of serum prostate-specific membrane antigen by a novel protein biochip immunoassay discriminates benign from malignant prostate disease. *Cancer Research*, 2001, vol. 61 (16), 6029-6033 [0085]

专利名称(译)	用于诊断阿尔茨海默病的多肽生物标志物		
公开(公告)号	EP1535076B1	公开(公告)日	2008-08-06
申请号	EP2003792291	申请日	2003-08-11
[标]申请(专利权)人(译)	拜尔健康护理有限责任公司		
申请(专利权)人(译)	拜耳医药保健公司		
当前申请(专利权)人(译)	拜耳医药保健公司		
[标]发明人	YALKINOGLU OZKAN KONIG GERHARD HOCHSTRASSER DENIS FRANCOIS SANCHEZ JEAN CHARLES CARRETTE ODILE		
发明人	YALKINOGLU, ÖZKAN KÖNIG, GERHARD HOCHSTRASSER, DENIS, FRANCOIS SANCHEZ, JEAN-CHARLES CARRETTE, ODILE		
IPC分类号	G01N33/68 G01N27/62 C07K14/47 C07K14/74 C07K14/805 C07K14/81 C12Q1/37 G01N33/53		
CPC分类号	G01N33/6896 C07K14/70539 C07K14/805 C07K14/8139 G01N2800/2821		
优先权	2002026643 2002-11-29 EP 2002018283 2002-08-23 EP		
其他公开文献	EP1535076A2		
外部链接	Espacenet		

摘要(译)

公开了一种用于评估患者中阿尔茨海默病状态的方法。还公开了一种用于监测患者中阿尔茨海默氏病进展的方法。该方法应用特异性肽标记物的检测，例如，使用质谱分析。

