



(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:
06.12.2006 Bulletin 2006/49

(51) Int Cl.:
G01N 33/74 (2006.01)

(21) Application number: 06008181.7

(22) Date of filing: 13.01.2000

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

• Gao, Ping
San Diego, CA 92129 (US)

(30) Priority: 14.01.1999 US 231422
26.06.1999 US 344639

(74) Representative: Atkinson, Jennifer
Barker Brettell
138 Hagley Road
Edgbaston
Birmingham B16 9PW (GB)

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
00902406.8 / 1 151 307

Remarks:

- This application was filed on 20 - 04 - 2006 as a divisional application to the application mentioned under INID code 62.
- The sequence listing, which is published as annex to the application documents, was filed after the date of filing. The applicant has declared that it does not include matter which goes beyond the content of the application as filed.


(71) Applicant: Scantibodies Laboratory, Inc.
Santee, CA 92071 (US)

(72) Inventors:
• Cantor, Thomas Leslie
El Cajon, CA 92020 (US)

(54) Methods for differentiating and monitoring parathyroid and bone status related diseases

(57) The present invention relates to novel methods and devices for differentiating in a patient parathyroid diseases, such as hyperparathyroidism and related bone diseases, from normal or non-disease states. One detects whole or non-fragmented (1 to 84) parathyroid hormone in a biological sample and also a large non-whole parathyroid hormone peptide fragment that can function as a parathyroid hormone antagonist. By either comparing values or using independently the value of either the large non-whole parathyroid hormone peptide fragment, the whole parathyroid hormone, or the combination of these values one is able to differentiate parathyroid and bone related disease states, as well as differentiate such states from normal states.

FIG. 2



Description**TECHNICAL FIELD**

5 **[0001]** The present invention relates to novel methods and devices for differentiating in a patient parathyroid diseases, such as hyperparathyroidism, from normal or non-disease states. One detects whole or non-fragmented (1 to 84) parathyroid hormone in a biological sample and also a large non-whole parathyroid hormone peptide fragment that can function as a parathyroid hormone antagonist. By either comparing values or using independently the value of either
10 the large non-whole parathyroid hormone peptide fragment, the whole parathyroid hormone, or the combination of these values one can differentiate parathyroid and bone related disease states, as well as differentiate such states from normal states.

RELATED APPLICATIONS

15 **[0002]** The present application is a continuation-in-part of a non-provisional utility patent application filed in the United States Patent and Trademark Office, Serial Number 08/231,422.

BACKGROUND ART

20 **[0003]** Calcium plays an indispensable role in cell permeability, the formation of bones and teeth, blood coagulation, transmission of nerve impulse, and normal muscle contraction. The concentration of calcium ions in the blood is, along with calcitriol and calcitonin, regulated mainly by parathyroid hormone (PTH). Although calcium intake and excretion may vary, PTH serves through a feedback mechanism to maintain a steady concentration of calcium in cells and surrounding fluids. When serum calcium lowers, the parathyroid glands secrete PTH, affecting the release of stored calcium. When
25 serum calcium increases, stored calcium release is retarded through lowered secretions of PTH.

[0004] The complete form of human PTH, sometimes referred to in the art as hPTH but referred to in the present invention either as whole PTH or wPTH, is a unique 84 amino acid peptide (SEQ ID NO. 1), as is shown in FIGURE 1. Researchers have found that this peptide has an anabolic effect on bone that involves a domain for protein kinase C activation (amino acid residues 28 to 34) as well as a domain for adenylate cyclase activation (amino acid residues 1 to 7). However, various catabolic forms of clipped or fragmented PTH peptides also are found in circulation, most likely
30 formed by intraglandular or peripheral metabolism. For example, whole PTH can be cleaved between amino acids 34 and 35 to produce a (1-34) PTH N-terminal fragment and a (35-84) PTH C-terminal fragment. Likewise, clipping can occur between either amino acids 36 and 37 or 37 and 38. Recently, a large PTH fragment referred to as "non-(1-84) PTH" has been disclosed which is clipped closer to the N-terminal end of PTH. (See R. LePage et alia, "A non-(1-84) circulating parathyroid hormone (PTH) fragment interferes significantly with intact PTH commercial assay measurements in uremic samples " Clin Chem (1998); 44: 805-810)

[0005] The clinical need for accurate measurement of PTH is well demonstrated. Serum PTH level is one of the most important indices for patients with the following diseases: familial hypocalciuria; hypercalcemia; multiple endocrine neoplasia types I and II; osteoporosis; Paget's bone disease; primary hyperparathyroidism - caused by primary hyperplasia
40 or adenoma of the parathyroid glands; pseudohypoparathyroidism; and renal failure, which can cause secondary hyperparathyroidism.

[0006] PTH plays a role in the course of disease in a patient with chronic renal failure. Renal osteodystrophy (RO) is a complex skeletal disease comprising osteitis fibrosa cystica (caused by PTH excess), osteomalacia - unmineralized bone matrix (caused by vitamin D deficiency), extraskeletal calcification/ossification (caused by abnormal calcium and phosphorus metabolism), and adynamic bone disease (contributed to by PTH suppression). Chronic renal failure patients can develop. RO Failing kidneys increase serum phosphorus (hyperphosphoremia) and decrease 1,25-dihydroxyvitamin D (1,25-D) production by the kidney. The former results in secondary hyperparathyroidism from decreased gastrointestinal calcium absorption and osteitis fibrosa cystica from increased PTH in response to an increase in serum phosphorus. The later causes hypocalcemia and osteomalacia. With the onset of secondary hyperparathyroidism, the parathyroid
50 gland becomes less responsive to its hormonal regulators because of decreased expression of its calcium and vitamin D receptors. Serum calcium drops. RO can lead to digital gangrene, bone pain, bone fractures, and muscle weakness.

[0007] Determining circulating biologically active PTH levels in humans has been challenging. One major problem is that PTH is found at low levels, normally 10pg/mL to 65 pg/mL. Coupled with extremely low circulating levels is the problem of the heterogeneity of PTH and its many circulating fragments. In many cases, immunoassays have faced
55 substantial and significant interference from circulating PTH fragments. For example, some commercially available PTH kits have almost 100% cross-reactivity with the non-(1-84) PTH fragment, (see the LePage article).

[0008] PTH immunoassays have varied over the years. One early approach is a double antibody precipitation immunoassay found in U. S. 4,369,138 to Arnold W. Lindall *et alia*. A first antibody has a high affinity for a (65-84) PTH

fragment. A radioactive labeled (65-84) PTH peptide is added to the sample with the first antibody to compete for the endogenous unlabeled peptide. A second antibody is added which binds to any first antibody and radioactive labeled PTH fragment complex, thereby forming a precipitate. Both precipitate and supernatant can be measured for radioactive activity, and endogenous PTH levels can be calculated therefrom.

5 [0009] In an effort to overcome PTH fragment interference, immunoradiometric two-site assays for intact PTH (I-PTH) have been introduced, such as Allegro® Intact PTH assay by the Nichol's Institute of San Juan Capistrano, California. In one version, a capture antibody specifically binds to the C-terminal portion of hPTH while a labeled antibody specifically binds to the N-terminal portion of the captured hPTH. In another, two monoclonal antibodies were used, both of which attached to the N-terminal portion of hPTH. Unfortunately, these assays have problems in that they measure but do not discriminate between wPTH and non-whole PTH peptide fragments. This inability comes to the fore in hyperparathyroid patients and renal failure patients who have significant endogenous concentrations of large, non-whole PTH fragments.

10 [0010] Recently, researchers have made a specific binding assay directed to the large N-terminal PTH fragments. (See. Gao, Ping et alia "Immunochemiluminometric assay with two monoclonal antibodies against the N-terminal sequence of human parathyroid hormone", Clinica Chimica Acta 245 (1996) 39-59.) This immunochemiluminometric assay uses two monoclonal antibodies to detect N-terminal (1-34) PTH fragments but not mid-portion PTH fragments or C-terminal PTH fragments. A key factor in the design of these assays is to eliminate any reaction with C-terminal PTH fragments.

20 DISCLOSURE OF THE INVENTION

[0011] The present invention relates to novel methods and devices for differentiating in a patient parathyroid diseases, (such as primary hyperparathyroidism, secondary hyperparathyroidism, and stages thereof), from normal or non-disease states; for monitoring the function of parathyroid glands either during or after treatment, i.e., intra-operation and after operation parathyroid function monitoring as well as therapeutic treatment; and also for monitoring the effects of therapeutic treatments for parathyroid related bone diseases and hyperparathyroidism. One detects the level in the serum or blood of at least one of three different parameters, namely, whole or non-fragmented parathyroid hormone in a biological sample, a large non-whole parathyroid hormone peptide fragment that can function as a parathyroid hormone antagonist, or the combination of the two values. By comparing the two values or by examining independently one of the above three values, one can differentiate parathyroid and bone disease states, as well as differentiate such states from normal states, as the relationship between these values, as well as the values themselves, change significantly between a normal person and a patient with a parathyroid disease.

25 [0012] The present invention incorporates a discovery that a large, non-whole PTH peptide fragment, a peptide having an amino acid sequence from between (SEQ ID No.2 [PTH₃₋₈₄]) and (SEQ ID No. 3 [PTH₃₄₋₈₄]), functions *in vivo* as a wPTH antagonist or inhibitor (PIN), (see FIGURE 12). In other words, the binding of wPTH to PTH receptors and the subsequent biological activity are affected by the presence of this PIN peptide fragment. The PTH receptors can be tied up with respect to PTH or PTH analogs in that the PTH binding site is blocked. The relationship between the concentrations of wPTH and PIN vary with PTH related disease states, and thus, are indicative of such states. Equally useful in view of the discovery of the antagonist nature of PIN, the present invention relates to novel methods and devices for monitoring parathyroid related bone diseases, and resultant bone loss or build-up. Increased amounts of PIN can inhibit the calcium releasing activity of PTH.

30 [0013] In making a measurement of wPTH, one does not want to detect PIN. The method for measuring the amount of wPTH in a sample such as serum, plasma, or blood comprises four general steps which can vary depending upon whether one uses a first antibody or antibody fragment specific for the PTH peptide SER-VAL-SER-GLU-ILE-GLN-LEU-MET (SEQ ID No. 4), wherein at least four amino acids are part of the antibody reactive portion of the peptide either as a signal antibody or a capture antibody in conventional immunoassay formats. (One can also use an analogous peptide present in other species, such as a rat peptide in which the first amino acid serine is substituted with an alanine.) Used either as a signal antibody or as a capture antibody, enough antibody is added to bind all wPTH present. Next, one allows the first antibody to bind to any wPTH present, thereby forming a complex. A specific binding label comprised of a second antibody and a conventional immunoassay label, such as chemiluminescent agents, colorimetric agents, energy transfer agents, enzymes, fluorescent agents, and radioisotopes, is used to label the complex, preferably at the C-terminal end of wPTH, and can be added either substantially simultaneously with the first antibody or subsequent thereto. Finally, one uses conventional techniques to measure the amount of labeled complex, and thereby calculate wPTH levels in the sample. If used as a signal antibody, then the first antibody still attaches at the N-terminal end, but the second antibody would serve as a capture antibody that attaches at the C-terminal end.

35 [0014] In making a measurement of PIN, one can either measure it directly, or indirectly. An indirect measurement can be made by first measuring wPTH and then measuring total PTH. Subtracting the wPTH value from the total PTH value, one derives the PIN value (For the purposes of the present invention, "total PTH" refers to the sum of wPTH, the naturally occurring predominant PTH receptor binding agonist, and PIN, the naturally occurring predominant PTH receptor

binding antagonist.) A total PTH assay detects both PIN and wPTH by detecting the N-terminal end of PTH not at SEQ ID No. 4, the very end of the N-terminal. By detecting between about amino acids 7 to 38 of PTH, the assay can detect both. A commercially available assay for total PTH is available from Scantibodies Laboratory, Inc. of Santee, California. A direct measurement of total PTH can be made by using an antibody or antibody fragment specific for a portion of the PTH peptide LEU-MET-HIS-ASN-LEU-GLY-LYS-HIS-LEU-ALA-SER-VAL-GLU-ARG-MET-GLN-TRP-LEU-ARG-LYS-LYS-LEU-GLN-ASP-VAL-HIS-ASN-PHE-VAL-ALA-LEU-GLY (SEQ ID No. 5), which comprises amino acids 7 to 38 of PTH, (preferably between amino acids 9 to 34), wherein at least four amino acids are part of the antibody reactive portion of the peptide. Such an antibody or antibody fragment can be used in conventional immunoassay formats either as a signal antibody or a capture antibody.

[0015] To differentiate between parathyroid disease states and the normal state or to monitor the effects of therapeutic treatment for parathyroid disease states, one can compare the relationship between the values of wPTH, PIN, or total PTH, (the combination of wPTH and PIN), in other words, the relationship between the values of PIN and total PTH, between PIN and whole PTH, or between whole PTH and total PTH. For example, one can use a proportion between wPTH and total PTH, between PIN and total PTH, or between PIN and wPTH. (Comparisons can even take the form of a neural network of all these factors.) Regardless of the comparative method chosen, these values change significantly between a normal person and a patient with a parathyroid disease and between various stages of parathyroid diseases.

[0016] Alternatively, one can either differentiate between parathyroid disease states and the normal state or monitor the effects of therapeutic treatment for parathyroid disease states by examining independently the value of either wPTH, PIN, or total PTH alone.

[0017] According to one aspect the invention provides a method for differentiating between a person having substantially normal parathyroid function and having hyperparathyroidism comprising determining and comparing at least two of the parameters selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the total parathyroid hormone level in the person. One may measure the whole parathyroid hormone level and the total parathyroid hormone level in the person, determine the parathyroid hormone inhibitory peptide fragment level from these two measurements, and compare the whole parathyroid hormone level to the parathyroid hormone inhibitory peptide fragment level. This comparison may be in the form of a ratio or proportion. The person may be a patient with chronic uremia. According to this aspect of the invention, one may measure and compare the whole parathyroid hormone level and the parathyroid hormone inhibitory peptide fragment level. The comparison may be in the form of a ratio or proportion. The person may be a patient with chronic uremia.

[0018] According to this aspect of the invention one may measure and compare the whole parathyroid hormone level and the total parathyroid hormone level in the person. The comparison may be in the form of a ratio or proportion. The person may be a patient with chronic uremia.

[0019] According to this aspect of the invention one may measure and compare the parathyroid hormone inhibitory peptide fragment level and the total parathyroid hormone level in the person. The comparison may be in the form of a ratio or proportion. The person may be a patient with chronic uremia.

[0020] According to another aspect the invention provides a method for differentiating between a person having substantially normal parathyroid function and having hyperparathyroidism comprising determining one parameter selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and a calculated total parathyroid hormone level. One may determine the parathyroid hormone inhibitory peptide fragment level by measuring the whole parathyroid hormone level and the total parathyroid hormone level. One may determine the total parathyroid hormone level by measuring the whole parathyroid hormone level and the parathyroid hormone inhibitory peptide fragment level. The person may be a patient with chronic uremia.

[0021] According to a further aspect, the invention provides a method for monitoring parathyroid related bone diseases and treatments therefor comprising determining and comparing at least two of the parameters selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the total parathyroid hormone level in the person. The comparison may be in the form of a ratio or proportion.

[0022] In this method one may measure the whole parathyroid hormone level and the total parathyroid hormone level in the person, determine the parathyroid hormone inhibitory peptide fragment level from these two measurements, and compare the whole parathyroid hormone level to the parathyroid hormone inhibitory peptide fragment level. The comparison may be in the form of a ratio or proportion.

[0023] In this method one may measure and compare the whole parathyroid hormone level and the parathyroid hormone inhibitory peptide fragment level. The comparison may be in the form of a ratio or proportion.

[0024] In this method one may measure and compare the whole parathyroid hormone level and the total parathyroid hormone level in the person. The comparison may be in the form of a ratio or proportion.

[0025] In this method one may measure and compare the parathyroid hormone inhibitory peptide fragment level and the total parathyroid hormone level in the person. One may measure the whole parathyroid hormone level in order to calculate the parathyroid hormone inhibitory peptide fragment level from the whole parathyroid hormone level and the total parathyroid hormone level. The comparison may be in the form of a ratio or proportion.

[0026] According to a yet further aspect the invention provides a method for monitoring parathyroid related bone diseases and treatments therefor comprising determining one parameter selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the calculated total parathyroid hormone level. One may determine the parathyroid hormone inhibitory peptide fragment level by measuring the whole parathyroid hormone level and the total parathyroid hormone level. One may determine the total parathyroid hormone level by measuring the whole parathyroid hormone level and the parathyroid hormone inhibitory peptide fragment level.

[0027] According to another aspect, the invention provides a method for monitoring the effects of therapeutic treatment for hyperparathyroidism comprising determining and comparing at least two of the parameters selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the total parathyroid hormone level in the person. The comparison may be in the form of a ratio or proportion. One may measure the whole parathyroid hormone level and the total parathyroid hormone level in the person, determine the parathyroid hormone inhibitory peptide fragment level from these two measurements, and compare the whole parathyroid hormone level to the parathyroid hormone inhibitory peptide fragment level. The comparison may be in the form of a ratio or proportion.

[0028] In this method one may measure and compare the whole parathyroid hormone level and the parathyroid hormone inhibitory peptide fragment level.

[0029] In this method one may measure and compare the whole parathyroid hormone level and the total parathyroid hormone level in the person. The comparison may be in the form of a ratio or proportion.

[0030] In this method one may determine and compare the parathyroid hormone inhibitory peptide fragment level and the total parathyroid hormone level in the person. One may determine the parathyroid hormone inhibitory fragment level by measuring the whole parathyroid hormone and the total parathyroid hormone level. The comparison may be in the form of a ratio or proportion.

[0031] According to a further aspect, the invention provides a method for monitoring the effects of therapeutic treatment for hyperparathyroidism comprising determining one parameter selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the calculated total parathyroid hormone level. One may determine the parathyroid hormone inhibitory peptide fragment level by measuring the whole parathyroid hormone level and the total parathyroid hormone level. One may determine the total parathyroid hormone level by measuring the whole parathyroid hormone level and the parathyroid hormone inhibitory peptide fragment level.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032]

FIGURE 1 is a diagrammatic view of human wPTH.

FIGURE 2 is a diagrammatic view of a wPTH assay using the present antibody as a tracer element.

FIGURE 3 is a diagrammatic view of a wPTH assay using the present antibody as a capture element.

FIGURE 4 is a graph showing a standard curve for a wPTH assay.

FIGURE 5 is a graph comparing a conventional I-PTH assay with the present wPTH assay for healthy normal persons with "normal" PTH values.

FIGURE 6 is a diagrammatic view showing interference from PIN in conventional I-PTH assays.

FIGURE 7 is a graph comparing a conventional I-PTH assay with the present wPTH assay for patients with chronic uremia.

FIGURE 8 is a graph showing the distribution of wPTH values for healthy normal persons, patients with primary hyperparathyroidism, and patients with chronic uremia.

FIGURE 9 is a diagrammatic view showing how PIN blocks the action of wPTH at the receptor level, thereby making the person insensitive to the biological effects of wPTH.

FIGURE 10 is a graph demonstrating complete cross-reactivity of wPTH and PIN in a total PTH assay used in the present invention.

FIGURE 11 is a graph demonstrating how the whole PTH assay used in the present invention does not detect to PIN.

FIGURE 12 is a graph demonstrating how PIN is an *in vivo* inhibitor of wPTH.

BEST MODES FOR CARRYING OUT THE INVENTION

[0033] In disclosing the present invention, one should remember that there are a number of closely analogous, species dependent forms of PTH. The amino acid sequence of hPTH is shown in FIGURE 1. However, for rat PTH, bovine PTH, or porcine PTH, for example, one finds the substitutions at some of the amino acids in the hPTH sequence. For the purposes of the present invention, one can use interchangeably antibodies or antibody fragments to forms of these PTHs, although it is preferred to use an antibody with specificity for PTH having a sequence matching the species in which the PTH measurements are made.

Whole PTH immunoassay

[0034] A preferred embodiment of the present invention is an immunoradiometric assay (IRMA), often referred to as a sandwich assay, as shown FIGURES 2 and 3. Elements employed in such an assay (10) include a capture antibody (12) attached to a solid support (14) and a signal antibody (16) having a label (18), attached thereto (20). Typically, one selects a capture antibody that is specific for C-terminal PTH fragments (22), while the label antibody is specific for the initial wPTH peptide sequence which comprises a domain for adenylate cyclase activation (24), as shown in FIGURE 2. However, one could reverse the specificity of these antibodies, as is shown in FIGURE 3.

[0035] Alternatively, one could create an immunoassay in which wPTH is either precipitated from solution or otherwise differentiated in a solution, as in conventional precipitating assays or turbidometric assays. For example, one can use at least three antibodies to form a precipitating mass. In addition to the initial wPTH sequence antibody and a C-terminal antibody, one can use at least a third antibody which attaches to the mid portion of PTH. The combined mass of wPTH and the at least three antibodies would form a labeled precipitating mass which can be measured by conventional techniques. Another method would be to couple the initial wPTH sequence antibody to colloidal solid supports, such as latex particles.

[0036] More specifically, one can create a signal antibody by iodinating 50 micrograms of affinity purified goat anti-(1-6) PTH antibody (Scantibodies Laboratory, Inc., Santee California, U.S.A.) by oxidation with chloramine T, incubation for 25 seconds at room temperature with 1 millicurie of 125-I radioisotope and reduction with sodium metabisulfate. Unincorporated 125-I radioisotope is separated from the 125-I-Goat anti-(1-6) PTH signal antibody by, passing the iodination mixture over a PD-10 desalting column (Pharmacia, Uppsala, Sweden) and following the manufacturers instructions. The fractions collected from the desalting column are measured in a gamma counter and those fractions representing the 125-I-goat anti-(1-6) PTH antibody are pooled and diluted to approximately 300,000 DPM (disintegrations per minute) per 100 microliters. This solution is the tracer solution to be used in the whole PTH IRMA.

[0037] Capture antibody coated tubes can be created by attaching affinity purified goat anti PTH 39-84 antibody, (Scantibodies Laboratory, Inc., Santee, California, U.S.A.), to 12 x 75 mm polystyrene tubes (Nunc, Denmark) by means of passive absorption techniques which are known to those of skill in the art. The tubes are emptied and dried, creating solid phase antibody coated tubes.

[0038] To conduct a whole PTH assay of a sample, 200 microliter samples of human serum are added to the solid phase antibody coated tubes. To each tube is added 100 microliters of the tracer solution (labeled goat anti-(1-6) PTH signal antibody). The tubes are incubated at room temperature with shaking at 170 rpm for 20-22 hours. During this time the immunochemical reaction of forming the sandwich of (solid phase goat anti-(39-84) PTH antibody) - {whole PTH} -- {125-1-goat anti-(1-6) PTH antibody} takes place. Following this incubation, the test tubes are washed with distilled water. Radioactivity on the solid phase, which amount corresponds to the quantity of wPTH present, is measured using a gamma counter. The radioactivity data for the samples is processed by conventional analysis with use of the results from standards and controls and a computer software in order that the concentration of whole PTH in the samples may be ascertained. FIGURE 4 shows a standard curve for such an assay.

Initial whole PTH sequence peptide

[0039] In order to make the signal antibody in the above assay, first one makes a synthetic PTH peptide corresponding either to hPTH (Ser - Val - Ser - Glu - Ile - Gln - Leu - Met), rat PTH (Ala - Val - Ser - Glu - Ile - Gln - Leu - Met), or at least four amino acids in the common sequence. The selected peptide can play two roles in making an assay, first as a specific source for creating a polyclonal antibody or monoclonal antibody source for signal antibody or capture antibody, and second as part of an affinity purification means for isolating the desired signal antibody or capture antibody.

[0040] Briefly, such a peptide can be synthesized on an Applied Biosystems, Inc. (Foster City, California, U.S.A.)

Model 431 automated peptide synthesizer employing Fmoc (9-fluoronylmethoxycarbonyl) as the alpha-amino protecting group. All amino acids and solvents are from Applied Biosystems and are of synthesis grade. Following synthesis, the peptide is cleaved from the resin, and side chains are de-protected, using a cleavage cocktail containing 6.67% phenol, 4.4% (v/v) thioanisole and 8.8% ethanedithiol in trifluoroacetic acid (TFA). The cleaved peptide is precipitated and washed several times in cold diethyl ether. It is then dissolved in water and lyophilized. The crude peptide is subjected to amino acid analysis (Waters PICO-TAG System, Boston, Massachusetts, U.S.A.) and reversed-phase HPLC using a VYDAC (TM) C8 column with 0.1% TFA in water and 99.9% acetonitrile in 0.1% TFA as the mobile buffers. The presence of a single major peak along with the appropriate amino acid composition is taken as evidence that the peptide is suitable for further use.

[0041] The resulting peptide is then attached to cross linked agarose beads (activated Sepharose 4B from Pharmacia, Uppsala, Sweden) according to instructions from the manufacturer. Armed with the initial peptide sequence on a bead, one can affinity purify a polyclonal antibody serum source to isolate the initial sequence antibody for the wPTH immunoassay.

Initial sequence whole PTH antibody

[0042] To create an affinity-purified anti-(1-6) PTH antibody, one first uses a selected initial PTH sequence peptide as described above as part of an immunogen for injection into a goat. The peptide can be used either by itself as an injectible immunogen, incorporated into a non PTH peptide having a molecular weight, typically, of between about 5,000 and 10,000,000, or as part of the wPTH complete sequence. The immunogen is mixed with an equal volume of Freund's complete adjuvant which is a mixture of light mineral oil, Arlacel detergent, and inactivated mycobacterium tuberculosis bacilli. The resulting mixture is homogenized to produce an aqueous/oil emulsion which is injected into the animal (typically a goat) for the primary immunization. The immunogen dose is approximately 50-400 micrograms. The goats are injected monthly with the same dose of immunogen complex except no mycobacterium tuberculosis bacilli is used in these subsequent injections. The goats are bled monthly, approximately three months after the primary immunization. The serum (or antiserum) is derived from each bleeding by separating the red blood cells from the blood by centrifugation and removing the antiserum which is rich in (1-6) PTH antibodies.

[0043] To purify the antiserum for the desired (1-6) PTH antibody, one packs a separation column with the initial PTH sequence peptide bound beads described above, washes the column and equilibrates it with 0.01 M phosphate buffered saline (PBS). The antiserum is loaded onto the column and washed with 0.01 M PBS in order to remove antibodies without the (1-6) PTH specificity. The bound specific goat anti-(1-6) PTH polyclonal antibody is eluted from the solid phase PTH 1-6 in the column by passing an elution solution of 0.1 M glycine hydrochloride buffer, pH 2.5 through the column. The eluted polyclonal antibody is neutralized after it leaves the column with either the addition of 1.0 M phosphate buffer, pH 7.5 or by a buffer exchange with 0.01 M PBS, as is known to those of skill in the art. The polyclonal antibody is stored at 2-8 degrees centigrade.

Comparison between whole PTH and total PTH assays

[0044] The present w PTH IRMA assay was compared to a conventional intact PTH or I-PTH immunoassay, the Allegro Nichols Intact-PTH assay, (which is commercially available and made by Nichols Institute Diagnostics of San Juan Capistrano, California, U.S.A.), in both PTH normal persons and those suffering from chronic uremia. This I-PTH immunoassay, due to its 100% cross reactivity between PIN and wPTH, is in actuality a total PTH assay, (see FIGURE 10).

[0045] FIGURE 5 shows the results for 34 normal human serum samples from healthy subjects which were assayed both by the present wPTH IRMA and the above I-PTH assay. In every case, the level of wPTH detected by the IRMA is lower than that reported by the I-PTH assay, demonstrating the ability of the present IRMA to avoid detecting the interfering large, non (1-84) PTH fragment detected by the I-PTH assay, (see FIGURE 11). FIGURE 6 illustrates how such interference can occur. An N-terminal PTH specific signal antibody which is not specific to the initial PTH peptide sequence, as in the present invention, can detect not only wPTH (as in the upper part of FIGURE 6), but also can detect PIN, the large, non (1-84) PTH fragment, (as in the lower part of FIGURE 6).

[0046] A comparison of assay results for 157 chronic uremic patients is shown in FIGURE 7. Serum samples from these patients were measured using the wPTH IRMA and the above I-PTH assay. In every case the wPTH levels are lower than I-PTH values.

Clinical Use

[0047] The present wPTH and PIN assays have been used in a clinical setting involving 188 persons. The group included 31 persons having normal healthy parathyroid glands and 157 patients with chronic uremia who are undergoing dialysis on a continuous basis. Each person had a blood sample drawn which was assayed using a wPTH assay from

EP 1 729 135 A2

Scantibodies Laboratory, Inc. as well as an I-PTH assay from Nichols Institute which gave total PTH values.

[0048] Table 1 shows the results individually and comparatively, of the wPTH, PIN, and total PTH assays from chronic uremic patients on dialysis.

TABLE 1

| Patient No. | Total PTH pg/ml | Whole PTH pg/ml | PIN pg/ml | PIN to Total PTH | PIN to Whole PTH | Whole PTH to Total PTH |
|-------------|-----------------|-----------------|-----------|------------------|------------------|------------------------|
| 1 | 1410 | 740 | 670 | 48% | 91% | 52% |
| 2 | 185 | 89 | 96 | 52% | 108% | 48% |
| 3 | 231 | 104 | 127 | 55% | 122% | 45% |
| 4 | 1020 | 590 | 430 | 42% | 73% | 53% |
| 5 | 270 | 159 | 111 | 41% | 70% | 59% |
| 6 | 201 | 100 | 101 | 50% | 101% | 50% |
| 7 | 380 | 100 | 280 | 74% | 280% | 26% |
| 8 | 460 | 277 | 183 | 40% | 66% | 60% |
| 9 | 380 | 197 | 183 | 48% | 93% | 52% |
| 10 | 880 | 522 | 358 | 41% | 69% | 59% |
| 11 | 310 | 154 | 156 | 50% | 101% | 50% |
| 12 | 880 | 451 | 429 | 49% | 95% | 51% |
| 13 | 670 | 418 | 252 | 38% | 60% | 63% |
| 14 | 390 | 221 | 169 | 43% | 76% | 57% |
| 15 | 170 | 108 | 62 | 36% | 57% | 64% |
| 16 | 510 | 381 | 129 | 25% | 34% | 75% |
| 17 | 200 | 67 | 133 | 61% | 199% | 34% |
| 18 | 170 | 109 | 61 | 36% | 56% | 64% |
| 19 | 360 | 199 | 161 | 45% | 81% | 55% |
| 20 | 260 | 164 | 96 | 37% | 59% | 63% |
| 21 | 440 | 372 | 68 | 15% | 18% | 85% |
| 22 | 120 | 51.7 | 68.3 | 57% | 132% | 43% |
| 23 | 600 | 527 | 73 | 12% | 14% | 83% |
| 24 | 220 | 130 | 90 | 41% | 69% | 59% |
| 25 | 190 | 136 | 54 | 28% | 40% | 72% |
| 26 | 220 | 118 | 102 | 46% | 86% | 54% |
| 27 | 630 | 334 | 296 | 47% | 89% | 53% |
| 28 | 150 | 90 | 60 | 40% | 67% | 60% |
| 29 | 170 | 106 | 64 | 38% | 60% | 62% |
| 30 | 810 | 489 | 321 | 40% | 66% | 60% |
| 31 | 570 | 319 | 251 | 44% | 79% | 56% |
| 32 | 570 | 467 | 103 | 18% | 22% | 82% |
| 33 | 400 | 300 | 100 | 25% | 33% | 75% |
| 34 | 560 | 378 | 182 | 33% | 48% | 68% |

EP 1 729 135 A2

(continued)

| | <i>Patient No.</i> | <i>Total PTH pg/ml</i> | <i>Whole PTH pg/ml</i> | <i>PIN pg/ml</i> | <i>PIN to Total PTH</i> | <i>PIN to Whole PTH</i> | <i>Whole PTH to Total PTH</i> |
|----|--------------------|------------------------|------------------------|------------------|-------------------------|-------------------------|-------------------------------|
| 5 | 35 | 310 | 121 | 189 | 61% | 156% | 39% |
| | 36 | 240 | 98 | 142 | 59% | 145% | 41% |
| | 37 | 280 | 133 | 157 | 54% | 118% | 48% |
| 10 | 38 | 230 | 124 | 106 | 46% | 85% | 54% |
| | 39 | 350 | 319 | 31 | 9% | 10% | 91% |
| | 40 | 200 | 133 | 67 | 34% | 50% | 67% |
| | 41 | 920 | 564 | 356 | 39% | 63% | 61% |
| 15 | 42 | 210 | 89 | 121 | 58% | 136% | 42% |
| | 43 | 1990 | 904 | 1086 | 55% | 120% | 45% |
| | 44 | 300 | 212 | 88 | 29% | 42% | 71% |
| 20 | 45 | 260 | 132 | 128 | 49% | 97% | 51% |
| | 46 | 140 | 72 | 68 | 49% | 94% | 51% |
| | 47 | 250 | 129 | 121 | 48% | 94% | 52% |
| | 48 | 130 | 72 | 58 | 45% | 81% | 56% |
| 25 | 49 | 1840 | 1000 | 840 | 46% | 84% | 54% |
| | 50 | 280 | 167 | 113 | 40% | 68% | 60% |
| | 51 | 490 | 268 | 222 | 45% | 83% | 55% |
| 30 | 52 | 150 | 77.1 | 72.9 | 49% | 95% | 51% |
| | 53 | 140 | 58.1 | 81.9 | 59% | 141% | 42% |
| | 54 | 210 | 92.7 | 117.3 | 56% | 127% | 44% |
| | 55 | 160 | 79 | 81 | 51% | 103% | 49% |
| 35 | 56 | 480 | 296 | 184 | 38% | 62% | 62% |
| | 57 | 480 | 281 | 199 | 41% | 71% | 59% |
| | 58 | 270 | 120 | 150 | 56% | 125% | 44% |
| 40 | 59 | 97 | 45 | 52 | 54% | 116% | 46% |
| | 60 | 330 | 154 | 176 | 53% | 114% | 47% |
| | 61 | 110 | 56 | 54 | 49% | 96% | 51% |
| | 62 | 660 | 456 | 204 | 31% | 45% | 69% |
| 45 | 63 | 300 | 137 | 163 | 54% | 119% | 46% |
| | 64 | 240 | 145 | 95 | 40% | 66% | 60% |
| | 65 | 100 | 66.5 | 33.5 | 34% | 50% | 67% |
| 50 | 66 | 410 | 416.3 | -6.3 | -2% | -2% | 102% |
| | 67 | 410 | 235.7 | 174.3 | 43% | 74% | 57% |
| | 68 | 45 | 14.4 | 30.6 | 68% | 213% | 32% |
| | 69 | 200 | 102.3 | 97.7 | 49% | 96% | 51% |
| 55 | 70 | 300 | 134 | 166 | 55% | 124% | 45% |
| | 71 | 320 | 202 | 118 | 37% | 58% | 63% |

EP 1 729 135 A2

(continued)

| <i>Patient No.</i> | <i>Total PTH pg/ml</i> | <i>Whole PTH pg/ml</i> | <i>PIN pg/ml</i> | <i>PIN to Total PTH</i> | <i>PIN to Whole PTH</i> | <i>Whole PTH to Total PTH</i> | |
|--------------------|------------------------|------------------------|------------------|-------------------------|-------------------------|-------------------------------|-----|
| 5 | 72 | 440 | 254 | 186 | 42% | 73% | 58% |
| | 73 | 190 | 99.6 | 90.4 | 48% | 91% | 52% |
| | 74 | 160 | 74.6 | 85.4 | 53% | 114% | 47% |
| 10 | 75 | 600 | 429.8 | 170.2 | 28% | 40% | 72% |
| | 76 | 1140 | 632 | 508 | 45% | 80% | 55% |
| | 77 | 440 | 211 | 229 | 52% | 109% | 48% |
| | 78 | 450 | 276 | 174 | 39% | 63% | 61% |
| 15 | 79 | 510 | 344 | 166 | 33% | 48% | 67% |
| | 80 | 190 | 62.8 | 127.2 | 67% | 203% | 33% |
| | 81 | 170 | 86 | 84 | 49% | 98% | 51% |
| 20 | 82 | 180 | 103.4 | 76.6 | 43% | 74% | 57% |
| | 83 | 78 | 22.7 | 55.3 | 71% | 244% | 29% |
| | 84 | 230 | 117 | 113 | 49% | 97% | 51% |
| | 85 | 160 | 96 | 64 | 40% | 67% | 60% |
| 25 | 86 | 220 | 89 | 131 | 60% | 147% | 40% |
| | 87 | 470 | 321.5 | 148.5 | 32% | 46% | 68% |
| | 88 | 310 | 137 | 173 | 56% | 126% | 44% |
| 30 | 89 | 2050 | 1127 | 923 | 45% | 82% | 55% |
| | 90 | 930 | 414 | 516 | 55% | 125% | 45% |
| | 91 | 180 | 65 | 115 | 64% | 177% | 36% |
| | 92 | 560 | 238 | 322 | 58% | 135% | 43% |
| 35 | 93 | 640 | 597 | 43 | 7% | 7% | 93% |
| | 94 | 590 | 382 | 208 | 35% | 54% | 65% |
| | 95 | 270 | 103 | 167 | 62% | 162% | 38% |
| 40 | 96 | 560 | 349 | 211 | 38% | 60% | 62% |
| | 97 | 180 | 78 | 102 | 57% | 131% | 43% |
| | 98 | 790 | 429 | 361 | 46% | 84% | 54% |
| | 99 | 670 | 372 | 298 | 44% | 80% | 56% |
| 45 | 100 | 140 | 20.4 | 119.6 | 85% | 586% | 15% |
| | 101 | 190 | 117 | 73 | 38% | 62% | 62% |
| | 102 | 190 | 108 | 82 | 43% | 76% | 57% |
| 50 | 103 | 430 | 217 | 213 | 50% | 98% | 50% |
| | 104 | 560 | 439 | 121 | 22% | 28% | 78% |
| | 105 | 500 | 357.7 | 142.3 | 28% | 40% | 72% |
| | 106 | 1560 | 777 | 783 | 50% | 101% | 50% |
| 55 | 107 | 62 | 24.3 | 37.7 | 61% | 155% | 39% |
| | 108 | 430 | 226 | 204 | 47% | 90% | 53% |

EP 1 729 135 A2

(continued)

| <i>Patient No.</i> | <i>Total PTH pg/ml</i> | <i>Whole PTH pg/ml</i> | <i>PIN pg/ml</i> | <i>PIN to Total PTH</i> | <i>PIN to Whole PTH</i> | <i>Whole PTH to Total PTH</i> | |
|--------------------|------------------------|------------------------|------------------|-------------------------|-------------------------|-------------------------------|------|
| 5 | 109 | 160 | 67.2 | 92.8 | 58% | 138% | 42% |
| | 110 | 530 | 346 | 184 | 35% | 53% | 65% |
| | 111 | 260 | 142 | 118 | 45% | 83% | 55% |
| 10 | 112 | 580 | 163 | 417 | 72% | 256% | 28% |
| | 113 | 440 | 579 | -139 | -32% | -24% | 132% |
| | 114 | 500 | 232.3 | 267.7 | 54% | 115% | 46% |
| | 115 | 160 | 60 | 100 | 63% | 167% | 38% |
| 15 | 116 | 340 | 202 | 138 | 41% | 68% | 59% |
| | 117 | 260 | 138 | 122 | 47% | 88% | 53% |
| | 118 | 260 | 119 | 141 | 54% | 118% | 46% |
| 20 | 119 | 160 | 84 | 76 | 48% | 90% | 53% |
| | 120 | 130 | 46 | 84 | 65% | 183% | 35% |
| | 121 | 190 | 104 | 86 | 45% | 83% | 55% |
| | 122 | 420 | 334 | 86 | 20% | 26% | 80% |
| 25 | 123 | 630 | 440 | 190 | 30% | 43% | 70% |
| | 124 | 75 | 26.4 | 48.6 | 65% | 184% | 35% |
| | 125 | 260 | 143 | 117 | 45% | 82% | 55% |
| 30 | 126 | 640 | 409 | 231 | 36% | 56% | 64% |
| | 127 | 130 | 66.7 | 63.3 | 49% | 95% | 51% |
| | 128 | 700 | 381 | 319 | 46% | 84% | 54% |
| | 129 | 560 | 376 | 184 | 33% | 49% | 67% |
| 35 | 130 | 240 | 107 | 133 | 55% | 124% | 45% |
| | 131 | 110 | 63 | 47 | 43% | 75% | 57% |
| | 132 | 420 | 297 | 123 | 29% | 41% | 71% |
| 40 | 133 | 580 | 229 | 351 | 61% | 153% | 39% |
| | 134 | 310 | 201.2 | 108.8 | 35% | 54% | 65% |
| | 135 | 160 | 97.9 | 62.1 | 39% | 63% | 61% |
| | 136 | 290 | 138.7 | 151.3 | 52% | 109% | 48% |
| 45 | 137 | 200 | 96.2 | 103.8 | 52% | 108% | 48% |
| | 138 | 770 | 662.7 | 107.3 | 14% | 16% | 86% |
| | 139 | 290 | 130.7 | 159.3 | 55% | 122% | 45% |
| 50 | 140 | 260 | 219 | 41 | 16% | 19% | 84% |
| | 141 | 350 | 211 | 139 | 40% | 66% | 60% |
| | 142 | 730 | 463.5 | 266.5 | 37% | 57% | 63% |
| | 143 | 490 | 231 | 259 | 53% | 112% | 47% |
| 55 | 144 | 160 | 87 | 73 | 46% | 84% | 54% |
| | 145 | 380 | 222 | 158 | 42% | 71% | 58% |

EP 1 729 135 A2

(continued)

| Patient No. | Total PTH pg/ml | Whole PTH pg/ml | PIN pg/ml | PIN to Total PTH | PIN to Whole PTH | Whole PTH to Total PTH |
|-------------|-----------------|-----------------|-----------|------------------|------------------|------------------------|
| 146 | 210 | 93.5 | 116.5 | 55% | 125% | 45% |
| 147 | 630 | 383.4 | 246.6 | 39% | 64% | 61% |
| 148 | 150 | 83.2 | 66.8 | 45% | 80% | 55% |
| 149 | 320 | 152.5 | 167.5 | 52% | 110% | 48% |
| 150 | 900 | 467.6 | 432.4 | 48% | 92% | 52% |
| 151 | 1180 | 818.6 | 361.4 | 31% | 44% | 69% |
| 152 | 120 | 38.4 | 81.6 | 68% | 213% | 32% |
| 153 | 5230 | 1388 | 3842 | 73% | 277% | 27% |
| 154 | 34 | 10.5 | 23.5 | 69% | 224% | 31% |
| 155 | 1020 | 590.6 | 429.4 | 42% | 73% | 58% |
| 156 | 180 | 76.6 | 103.4 | 57% | 135% | 43% |
| 157 | 120 | 51.1 | 68.9 | 57% | 135% | 43% |
| | | | | | | |
| Median | 300 | 154 | 127 | 46% | 84% | 54% |

[0049] TABLE 2 shows the results, individually and comparatively, of the wPTH, PIN, and total PTH assays from the normals.

TABLE 2

| Patient No. | Total PTH pg/ml | Whole PTH pg/ml | PIN pg/ml | PIN to Total PTH | PIN to Whole PTH | Whole PTH to Total PTH |
|-------------|-----------------|-----------------|-----------|------------------|------------------|------------------------|
| 1 | 17.13 | 3.32 | 13.81 | 81% | 416% | 19% |
| 2 | 32.92 | 10.49 | 22.43 | 68% | 214% | 32% |
| 3 | 31.32 | 10.31 | 21.01 | 67% | 204% | 33% |
| 4 | 41.84 | 12.72 | 29.12 | 70% | 229% | 30% |
| 5 | 33.03 | 10.09 | 22.94 | 69% | 227% | 31% |
| 6 | 44.32 | 14.23 | 30.09 | 68% | 211% | 32% |
| 7 | 31.47 | 6.8 | 24.67 | 78% | 363% | 22% |
| 8 | 20.82 | 10.03 | 10.79 | 52% | 108% | 48% |
| 9 | 34.64 | 15.95 | 18.69 | 54% | 117% | 46% |
| 10 | 23.69 | 5.25 | 18.44 | 78% | 351% | 22% |
| 11 | 53.98 | 17.82 | 36.16 | 67% | 203% | 33% |
| 12 | 52.71 | 18.83 | 33.88 | 64% | 180% | 36% |
| 13 | 26.92 | 5.63 | 21.29 | 79% | 378% | 21% |
| 14 | 39.93 | 11.86 | 28.07 | 70% | 237% | 30% |
| 15 | 48.84 | 20.47 | 28.37 | 58% | 139% | 42% |
| 16 | 29.56 | 13.68 | 15.88 | 54% | 116% | 46% |
| 17 | 36.19 | 14.69 | 21.5 | 59% | 146% | 41% |
| 18 | 20.96 | 6.99 | 13.97 | 67% | 200% | 33% |

EP 1 729 135 A2

(continued)

| <i>Patient No.</i> | <i>Total PTH pg/ml</i> | <i>Whole PTH pg/ml</i> | <i>PIN pg/ml</i> | <i>PIN to Total PTH</i> | <i>PIN to Whole PTH</i> | <i>Whole PTH to Total PTH</i> |
|--------------------|------------------------|------------------------|------------------|-------------------------|-------------------------|-------------------------------|
| 19 | 59.29 | 27.89 | 31.4 | 53% | 113% | 47% |
| 20 | 45.57 | 18.23 | 27.34 | 60% | 150% | 40% |
| 21 | 35.64 | 18.72 | 16.92 | 47% | 90% | 53% |
| 22 | 38.53 | 19.56 | 18.97 | 49% | 97% | 51% |
| 23 | 21.71 | 9.34 | 12.37 | 57% | 132% | 43% |
| 24 | 32.42 | 13.51 | 18.91 | 58% | 140% | 42% |
| 25 | 28.5 | 10.41 | 18.09 | 63% | 174% | 37% |
| 26 | 18.17 | 7.8 | 10.37 | 57% | 133% | 43% |
| 27 | 39.96 | 17.29 | 22.67 | 57% | 131% | 43% |
| 28 | 34.08 | 15.24 | 18.84 | 55% | 124% | 45% |
| 29 | 42.95 | 19.59 | 23.36 | 54% | 119% | 46% |
| 30 | 38.4 | 12.16 | 26.24 | 68% | 216% | 32% |
| 31 | 47.57 | 18.45 | 29.12 | 61% | 158% | 39% |
| Median | 34.64 | 13.51 | 21.5 | 61% | 158% | 39% |

[0050] Clearly, the statistically significant differences in the medians of these two groups demonstrates that one can differentiate between the two by using these assays alone or by comparing their respective values.

TABLE 3

| <i>Sample Type</i> | <i>Total PTH (pg/mL)</i> | <i>Whole PTH (pg/mL)</i> | <i>PIN (pg/mL)</i> | <i>PIN to Total PTH</i> | <i>PIN to Whole PTH</i> | <i>Whole PTH to Total PTH</i> |
|--------------------------------|--------------------------|--------------------------|--------------------|-------------------------|-------------------------|-------------------------------|
| Chronic Uremia (n=157) Medians | 300 | 154 | 127 | 46% | 84% | 55% |
| Normal (n=31) Medians | 34.64 | 13.51 | 21.5 | 61% | 158% | 37% |
| P-Value | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |

[0051] The ordinarily skilled artisan can appreciate that the present invention can incorporate any number of the preferred features described above.

[0052] All publications or unpublished patent applications mentioned herein are hereby incorporated by reference thereto.

[0053] Other embodiments of the present invention are not presented here which are obvious to those of ordinary skill in the art, now or during the term of any patent issuing from this patent specification, and thus, are within the spirit and scope of the present invention.

Annex to the application documents - subsequently filed sequences listing

[0054]

SEQUENCE LISTING

5

<110> Cantor, Thomas L.
Gao, Ping

10

<120> Methods for Differentiating Parathyroid and Bone Status Related Diseases

<160> 5

15

<170> Microsoft Word 7.0

20

<210> 1

<211> 84 [integer length]

25

<212> PRT

30

<400> 1

Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu

35

1 5 10 15

Asn Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp

40

20 25 30

Val His Asn Phe Val Ala Leu Gly Ala Pro Leu Ala Pro Arg Asp

45

35 40 45

Ala Gly Ser Gln Arg Pro Arg Lys Lys Glu Asp Asn Val Leu Val

50

50 55 60

55

5 Glu Ser His Glu Lys Ser Leu Gly Glu Ala Asn Lys Ala Asp Val
 65 70 75

10 Asn Val Leu Thyr Lys Ala Lys Ser Gln
 80

15
<210> 2

20 **<211> 82 [integer length]**

25 **<212> PRT**

30 **<400> 2**

35 Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn Ser
 1 5 10 15

40 Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His
 20 25 30

45 Asn Phe Val Ala Leu Gly Ala Pro Leu Ala Pro Arg Asp Ala Gly
 35 40 45

50 Ser Gln Arg Pro Arg Lys Lys Glu Asp Asn Val Leu Val Glu Ser
 50 55 60

55 His Glu Lys Ser Leu Gly Glu Ala Asn Lys Ala Asp Val Asn Val
 65 70 75

<212> PRT

5

<400> 4

Ser Val Ser Glu Ile Gln Leu Met

10

1 5

15

<210> 5

20

<211> 32 [integer length]

25

<212> PRT

30

<400> 5

Leu Met His Asn Leu Gly Lys His Leu Asn Ser Met Glu Arg Val

35

1 5 10 15

Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His Asn Phe Val Ala

40

20 25 30

Leu Gly

45

50

55

SEQUENCE LISTING

5 <110> SCANTIBODIES LABORATORY, INC.
 CANTOR, Thomas
 GAO, Ping

<120> METHODS FOR DIFFERENTIATING AND MONITORING PARATHYROID AND
 BONE STATUS RELATED DISEASES

10 <130> JL23869P.EPD2

<140> EP 06008181.7
 <141> 2000-01-13

15 <150> EP 00902406.8
 <151> 2000-01-13

<150> PCT/US00/00855
 <151> 2000-01-13

20 <150> 09/344,639
 <151> 1999-06-26

<150> 09/231,422
 <151> 1999-01-14

<160> 5

25 <170> FastSEQ for windows Version 4.0

<210> 1
 <211> 84
 <212> PRT
 <213> Homo sapiens

30 <400> 1
 Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn
 1 5 10 15
 Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His
 20 25 30
 35 Asn Phe Val Ala Leu Gly Ala Pro Leu Ala Pro Arg Asp Ala Gly Ser
 35 40 45
 Gln Arg Pro Arg Lys Lys Glu Asp Asn Val Leu Val Glu Ser His Glu
 50 55 60
 Lys Ser Leu Gly Glu Ala Asn Lys Ala Asp Val Asn Val Leu Thr Lys
 65 70 75 80
 40 Ala Lys Ser Gln

<210> 2
 <211> 82
 <212> PRT
 <213> Homo sapiens

45 <400> 2
 Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu Asn Ser Met
 1 5 10 15
 50 Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val His Asn Phe
 20 25 30
 Val Ala Leu Gly Ala Pro Leu Ala Pro Arg Asp Ala Gly Ser Gln Arg
 35 40 45
 Pro Arg Lys Lys Glu Asp Asn Val Leu Val Glu Ser His Glu Lys Ser
 50 55 60
 55 Leu Gly Glu Ala Asn Lys Ala Asp Val Asn Val Leu Thr Lys Ala Lys
 65 70 75 80
 Ser Gln

<210> 3
 <211> 51
 <212> PRT
 <213> Homo sapiens

5

<400> 3
 Phe Val Ala Leu Gly Ala Pro Leu Ala Pro Arg Asp Ala Gly Ser Gln
 1 5 10 15
 Arg Pro Arg Lys Lys Glu Asp Asn Val Leu Val Glu Ser His Glu Lys
 20 25 30
 Ser Leu Gly Glu Ala Asn Lys Ala Asp Val Asn Val Leu Thr Lys Ala
 35 40 45
 Lys Ser Gln
 50

10

<210> 4
 <211> 8
 <212> PRT
 <213> Homo sapiens

15

<400> 4
 Ser Val Ser Glu Ile Gln Leu Met
 1 5

20

<210> 5
 <211> 32
 <212> PRT
 <213> Homo sapiens

25

<400> 5
 Leu Met His Asn Leu Gly Lys His Leu Asn Ser Met Glu Arg Val Glu
 1 5 10 15
 Trp Leu Arg Lys Lys Leu Gln Asp Val His Asn Phe Val Ala Leu Gly
 20 25 30

30

Claims

35

1. A method for determining whether a person has a parathyroid disease or has substantially normal parathyroid function, which method comprises:

40

comparing at least two parameters selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the total parathyroid hormone level for a sample from the person;
 thereby determining whether the person has a parathyroid disease or has substantially normal parathyroid function.

45

2. A method for monitoring a parathyroid disease or treatment of a parathyroid disease in a person, which method comprises:

50

comparing at least two parameters selected from the group consisting of the whole parathyroid hormone level, the parathyroid hormone inhibitory peptide fragment level, and the total parathyroid hormone level for a sample from the person;
 thereby monitoring a parathyroid disease or the treatment of a parathyroid disease in the person.

3. The method of claim 1 or 2, wherein the sample is a blood or serum sample from a human.

55

4. The method of claim 1 or 2, wherein the level of the parathyroid hormone inhibitory peptide is determined by measuring the level of wPTH and the level of total PTH in the sample, and subtracting the level of wPTH from the level of total PTH.

5. The method of claim 1 or 2, wherein the level of whole parathyroid hormone in the sample is compared with the level of total parathyroid hormone in the sample.
- 5 6. The method of claim 1 or 2, wherein the level of whole parathyroid hormone in the sample is compared with the level of parathyroid hormone inhibitory peptide in the sample.
7. The method of claim 1 or 2, wherein the level of total parathyroid hormone in the sample is compared with the level of parathyroid hormone inhibitory peptide in the sample.
- 10 8. The method of claim 1 or 2, further comprising measuring the level of the whole parathyroid hormone by conventional immunoassay.
9. The method of claim 1 or 2, wherein the parathyroid disease is hyperparathyroidism.
- 15 10. The method of claim 1 or 2, wherein the comparison of parameters is in the form of a ratio or proportion.
11. A method to measure whole parathyroid hormone (wPTH) in a sample, which method comprises:
- 20 obtaining a sample suspected of containing wPTH;
contacting the sample with a first antibody or antibody fragment, wherein the first antibody or antibody fragment is produced using as an immunogen a synthetic peptide corresponding to Ser-Val-Ser-Glu-Ile-Gln-Leu-Met, or Ala-Val-Ser-Glu-Ile-Gln-Leu-Met, or at least four amino acids in the sequence Val-Ser-Glu-Ile-Gln-Leu-Met;
allowing the first antibody or antibody fragment to form a complex with any wPTH present;
contacting the sample with a second antibody or antibody fragment which is specific for a C-terminal PTH
25 fragment, wherein either the first antibody or antibody fragment or the second antibody or antibody fragment is labeled, thereby forming a labeled complex; and
determining the amount of the labeled complex to measure the amount of wPTH in the sample.
12. The method of claim 11, further comprising:
- 30 determining the level of total parathyroid hormone (PTH) in the sample; and
subtracting the level of wPTH in the sample from the level of total PTH in the sample, thereby determining the level of a parathyroid hormone inhibitory peptide (PIN) in the sample.
- 35 13. The method of claim 11, wherein the sample contains a physiological level of whole parathyroid hormone.
14. The method of claim 11 or 12, wherein the sample is a human blood or serum sample.
15. The method of claim 12, wherein the parathyroid inhibitory peptide is human PTH(7-84).
- 40 16. The method of claim 11, wherein the first antibody or antibody fragment is a PTH(1-6) antibody or antibody fragment.
17. A complex comprising:
- 45 whole parathyroid hormone (wPTH) complexed with an antibody or antibody fragment;
wherein the antibody or antibody fragment is specific for the PTH peptide SER-VAL-SER-GLU-ILE-GLN-LEU-MET (SEQ ID NO: 4), and at least four amino acids in this sequence are part of the reactive portion of the peptide sequence with the antibody or antibody fragment; and
wherein the antibody or antibody fragment does not bind to a parathyroid hormone inhibitory peptide.
- 50 18. An isolated antibody or antibody fragment which is produced using an immunogen that comprises a synthetic PTH peptide consisting of at least four amino acids of the sequence VAL-SER-GLU-ILE-GLN-LEU-MET;
wherein the antibody or antibody fragment is affinity purified using a synthetic PTH peptide corresponding to hPTH (1-8), rat PTH(1-8), or at least four amino acids in the common sequence.
- 55 19. A method to produce an antibody or antibody fragment, which method comprises:
- preparing a synthetic peptide corresponding to at least four amino acids of the sequence VAL-SER-GLU-ILE-

GLN-LEU-MET;

using the synthetic peptide to prepare an immunogen;

injecting the immunogen into an animal to elicit antibody formation; and

purifying the antibody or antibody fragment using a synthetic peptide that corresponds to at least four amino acids of the sequence VAL-SER-GLU-ILE-GLN-LEU-MET.

5

20. An isolated antibody or antibody fragment produced by the method of claim 18.

10

21. The method of claim 19 or the antibody or antibody fragment of claim 20, wherein the synthetic peptide used as part of the immunogen is the same as the synthetic peptide used for the affinity purification of the antibody or antibody fragment.

22. The isolated antibody or antibody fragment of claim 18 or claim 20, which is a polyclonal antibody.

15

23. The isolated antibody or antibody fragment of claim 18 or claim 20, which is a monoclonal antibody.

24. A method for measuring the amount of whole parathyroid hormone in a sample, which method is **characterized by**:

20

a) adding to the sample a first antibody or antibody fragment specific for an initial wPTH peptide sequence consisting of amino acids 1 to 6 of human whole parathyroid hormone which comprises a domain for adenylate cyclase activation;

b) allowing the antibody or antibody fragment to bind to any whole parathyroid hormone present, thereby forming a complex;

25

c) adding a second antibody or antibody fragment that is specific for a C-terminal PTH fragment, wherein either the first antibody or antibody fragment or the second antibody or antibody fragment is labeled, thereby forming a second complex; and

d) measuring the amount of labeled complex, thereby determining the amount of whole parathyroid hormone in the sample.

30

25. The method according to claim 24 **characterized in that** the first antibody or antibody fragment acts as the capture antibody in a sandwich assay, and is attached to a solid support; and the second antibody or antibody fragment acts as a signal antibody in a sandwich assay, and is labeled.

35

26. The method according to claim 24 **characterized in that** the second antibody or antibody fragment acts as the capture antibody in a sandwich assay, and is attached to a solid support; and the first antibody or antibody fragment acts as a signal antibody in a sandwich assay, and is labeled.

40

45

50

55

FIG. 1

Whole Human PTH (1-84)




FIG. 2




FIG. 3




FIG. 4

Standard Curve for Whole PTH Assay




FIG. 5

Normal Value Comparison Whole PTH Assay (with PTH 1-8 Antibody as Tracer) versus Nichols' Intact PTH Assay (with PTH 7-84 Interference)





FIG. 6

Whole PTH



Big PTH 7-84 Fragment




FIG. 7




FIG. 8




FIG. 9




FIG. 10

Intact PTH IRMA -Total PTH




FIG. 11





FIG. 12



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 08231422 B [0002]

Non-patent literature cited in the description

- **R. LEPAGE.** A non-(1-84) circulating parathyroid hormone (PTH) fragment interferes significantly with intact PTH commercial assay measurements in uremic samples. *Clin Chem*, 1998, vol. 44, 805-810 [0004]
- **GAO, PING.** Immunochemically luminometric assay with two monoclonal antibodies against the N-terminal sequence of human parathyroid hormone. *Clinica Chimica Acta*, 1996, vol. 245, 39-59 [0010]

| | | | |
|----------------|--|---------|------------|
| 专利名称(译) | 鉴别和监测甲状旁腺和骨骼状态相关疾病的方法 | | |
| 公开(公告)号 | EP1729135A2 | 公开(公告)日 | 2006-12-06 |
| 申请号 | EP2006008181 | 申请日 | 2000-01-13 |
| [标]申请(专利权)人(译) | SCANTIBODIES LAB | | |
| 申请(专利权)人(译) | SCANTIBODIES实验室, INC. | | |
| 当前申请(专利权)人(译) | SCANTIBODIES实验室, INC. | | |
| [标]发明人 | CANTOR THOMAS LESLIE GAO PING | | |
| 发明人 | CANTOR, THOMAS LESLIE GAO, PING | | |
| IPC分类号 | G01N33/74 G01N33/53 C07K14/635 C07K16/26 G01N33/68 G01N33/78 | | |
| 代理机构(译) | 阿特金森JENNIFER | | |
| 优先权 | 09/231422 1999-01-14 US 09/344639 1999-06-26 US | | |
| 其他公开文献 | EP1729135A3 | | |
| 外部链接 | Espacenet | | |

摘要(译)

本发明涉及用于区分患者甲状旁腺疾病(例如甲状旁腺功能亢进和相关骨病)的正常或非疾病状态的新方法和装置。人们可以检测生物样本中的全部或非碎片(1至84)甲状旁腺激素,以及可以作为甲状旁腺激素拮抗剂起作用的大型非整体甲状旁腺激素肽片段。通过比较值或独立使用大的非整体甲状旁腺激素肽片段,整个甲状旁腺激素或这些值的组合的值,能够区分甲状旁腺和骨相关疾病状态,以及区分这些状态来自正常状态。

FIG. 2

