



(11) **EP 2 812 453 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
27.09.2017 Bulletin 2017/39

(51) Int Cl.:
G01N 33/53 (2006.01) G01N 33/569 (2006.01)

(21) Application number: **13747009.2**

(86) International application number:
PCT/CN2013/071467

(22) Date of filing: **06.02.2013**

(87) International publication number:
WO 2013/117162 (15.08.2013 Gazette 2013/33)

(54) **NUCLEIC ACID APTAMERS AGAINST PLASMODIUM LACTATE DEHYDROGENASE AND HISTIDINE-RICH PROTEIN II AND USES THEREOF FOR MALARIA DIAGNOSIS**

NUCLEINSÄUREAPTAMERE GEGEN PLASMODIUM LACTATDEHYDROGENASE UND HISTIDINREICHES PROTEIN II SOWIE VERWENDUNGEN DAVON ZUR DIAGNOSE VON MALARIA

APTAMÈRES D'ACIDE NUCLÉIQUE CONTRE LA LACTICODÉSHYDROGÉNASE DE PLASMODIUM ET LA PROTÉINE II RICHE EN HISTIDINE, ET LEURS UTILISATIONS POUR LE DIAGNOSTIC DU PALUDISME

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **09.02.2012 US 201261596774 P**

(43) Date of publication of application:
17.12.2014 Bulletin 2014/51

(73) Proprietor: **The University of Hong Kong Hong Kong (CN)**

(72) Inventors:
• **TANNER, Julian Alexander Hong Kong (CN)**
• **CHEUNG, Yee Wai Kowloon Hong Kong (CN)**
• **KOTAKA, Masayo Kowloon Hong Kong (CN)**

(74) Representative: **HGF Limited 8th Floor 140 London Wall London EC2Y 5DN (GB)**

(56) References cited:
EP-A1- 2 532 749 WO-A1-2011/147193
US-A1- 2011 117 107

- **Authors Cheung ET AL: "Development of nucleic acid aptamers against Plasmodium falciparum lactate dehydrogenase for improving the diagnosis of malaria", , 11 June 2011 (2011-06-11), XP055198525, Retrieved from the Internet: URL: <http://hdl.handle.net/10722/137696> [retrieved on 2015-06-26]**
- **CHEUNG Y W ET AL: "E2-22: New avenues to malaria diagnosis - nucleic acid aptamers against P-falciparum histidine rich protein 2 and lactate dehydrogenase", FEBS JOURNAL; 35TH CONGRESS OF THE FEDERATION-OF-EUROPEAN-BIOCHEMICAL-SOCIETY ES, WILEY-BLACKWELL PUBLISHING LTD, GB; GOTHENBURG, SWEDEN, vol. 277, no. Suppl. 1, 1 June 2010 (2010-06-01), pages 244-245, XP008153614, ISSN: 1742-464X, DOI: 10.1111/J.1742-4658.2010.07680.X [retrieved on 2010-06-08]**
- **M. MORENO ET AL: "Advances on Aptamers Targeting Plasmodium and Trypanosomatids", CURRENT MEDICINAL CHEMISTRY, vol. 18, no. 32, 1 November 2011 (2011-11-01), pages 5003-5010, XP055033003, ISSN: 0929-8673, DOI: 10.2174/092986711797535218**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 812 453 B1

- **Y.-W. CHEUNG ET AL:** "Structural basis for discriminatory recognition of Plasmodium lactate dehydrogenase by a DNA aptamer", **PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES**, vol. 110, no. 40, 16 September 2013 (2013-09-16), pages 15967-15972, XP055198345, ISSN: 0027-8424, DOI: 10.1073/pnas.1309538110
- **CHEUNG YW ET AL.:** 'E2.22: New avenues to malaria diagnosis-nucleic acid aptamers against P. falciparum histidine rich protein 2 and lactate dehydrogenase.' **THE FEBS JOURNAL** vol. 277, no. SUP.1, 08 June 2010, pages 244 - 245, XP008153614
- **FANG XH ET AL.:** 'Using molecular beacons to probe molecular interactions between lactate dehydrogenase and single-stranded DNA' **ANALYTICAL CHEMISTRY** vol. 72, no. 14, 15 July 2000, pages 3280 - 3285, XP055079487

Description

CROSS-REFERENCE TO RELATED APPLICATION

5 **[0001]** This application claims the benefit of U.S. provisional application Serial No. 61/596,774, filed February 9, 2012.

FIELD OF THE INVENTION

10 **[0002]** The present invention relates to DNA aptamers that specifically bind to *Plasmodium* lactate dehydrogenase (LDH), and uses thereof for diagnosis of malaria.

BACKGROUND

15 **[0003]** Malaria, an infectious disease widespread in tropical and subtropical regions, is caused by infection with the parasitic protist Plasmodium. While there are four species of *Plasmodium* that can infect humans, the species *Plasmodium falciparum* causes the most dangerous form of malaria with the highest mortality rate. According to the World Health Organization (WHO), there are over 200 million confirmed malaria cases annually, resulting in nearly one million deaths.

20 **[0004]** In addition, malaria has a negative impact on the wealth of nations and individuals. According to the World Bank, malaria costs Africa about US\$12 billion per year and slows gross domestic product (GDP) by over 1% each year. The burden of malaria traps families and communities into deeper poverty by lost productivity and the negative impact on domestic and foreign investment and tourism. Furthermore, malaria greatly affects the structure of society as high mortality rates often afflict the age group under five years old.

25 **[0005]** The Centers for Disease Control and Prevention in USA notes that although complete elimination of the malaria parasite would be optimal, this is not realistic for most countries endemic for malaria. Improved management of malaria cases is fundamental to control the spreading of malaria.

30 **[0006]** Rapid diagnostic tests (RDT) are critical for the management of malaria to reduce the morbidity, mortality and transmission of malaria. Commonly-used RDTs for malaria include microscopic, polymerase chain reaction (PCR), and antibody based assays. Microscopic diagnosis of malaria, which is based on the microscopic observation of the malaria parasite in a blood smear, can differentiate various species of Plasmodium. PCR based assays can also distinguish various species of Plasmodium with high sensitivity. However, both the microscopic and PCR-based diagnostic tests require expensive equipment and highly trained healthcare workers that are generally not available to the majority of the at-risk population in developing countries.

35 **[0007]** Antibody based assays utilize antibodies for detection of Plasmodium antigens. The commonly-used Plasmodium antigens for malaria diagnosis include plasmodium lactate dehydrogenase (pLDH), histidine rich protein 2 (PfHRP2), and aldolase. While antibody-based rapid diagnostic assays have greatly benefited malaria management, they suffer from significant shortcomings including thermal instability, batch to batch variations, and high cost of production.

[0008] Therefore, there is a need for improved methods for malaria diagnosis. Cheung and Tanner (2011) disclose the development of aptamers specific for pLDH.

40 BRIEF SUMMARY

[0009] The present invention provides nucleic acid aptamers that bind to Plasmodium lactate dehydrogenase (referred to herein as "malaria aptamers"), and methods for using these malaria aptamers for the diagnosis of malaria. Disclosed are nucleic acid aptamers that target and bind to two malaria proteins, lactate dehydrogenase and histidine rich protein II, with high specificity and affinity. Aptamers against lactate dehydrogenase can be used for pan-species detection of Plasmodium, whilst aptamers against histidine rich protein II can be used for specific detection of *Plasmodium falciparum*. In parallel, these two aptamers can be incorporated into devices both to detect Plasmodium and diagnose malaria generally, and to determine whether the malaria is caused specifically by *Plasmodium falciparum*.

50 **[0010]** The present invention is directed to methods of using malaria aptamers as components of a diagnostic device for the diagnosis of malaria. The method provides specific detection of malaria antigens based on the specificity of binding of the malaria aptamers to the malaria protein targets.

[0011] In certain embodiment, the present invention provides formulations comprising a nucleic acid aptamer of the present invention, or a stabilized derivative of such aptamers, or a pharmaceutically acceptable salt thereof. The formulations may comprise any aptamer that binds to Plasmodium lactate dehydrogenase. Disclosed are methods of using the nucleic acid aptamers of the present invention for the diagnosis or treatment of malaria by specifically binding to and/or inhibiting the function of Plasmodia-encoded proteins.

55 **[0012]** In terms of diagnostics, the present invention may be used in combination with other diagnostic approaches. Malaria aptamers may be administered alone or in combination with other drugs.

[0013] In one embodiment, the present invention also provides diagnostic methods of quantifying Plasmodium protein levels in a sample. The malaria aptamers may be labeled by a detectable substance, including, but not limited to, fluorescent materials, enzymes, luminescent materials and radioactive materials. Such embodiments of the invention can be used to detect Plasmodium protein levels in a biological sample.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014]

FIG 1 is an illustration depicting the *in vitro* selection of aptamers using the "Systematic Evolution of Ligands by Exponential Enrichment" (SELEX™) process.

FIG 2 shows agarose gel analysis of the *in vitro* selection of LDH aptamers. Samples taken at 1st, 3rd, 5th, 7th, 9th, 12th, 15th, 18th, 19th and 20th rounds.

FIG 3 shows the nucleic acid sequences of embodiments of the aptamers of the present invention. The aptamers bind specifically to *Plasmodium falciparum* lactate dehydrogenase (PfLDH) or histidine rich protein 2 (HRP2). The aptamers are cloned and sequenced from the ssDNA pool after 20 rounds of SELEX (including counter SELEX process by using human LDH and magnetic beads) against *Plasmodium falciparum* PfLDH or against *Plasmodium falciparum* Histidine-rich protein II (HRP2), respectively.

FIG 4 is a graph showing the comparison between 2009 (the PfLDH aptamer with two 18-mer sequences on each 5' and 3' end) and 2009s using isothermal titration calorimetry. The left panel is the raw calorimetry data and binding isotherm for 2009, and the right panel is the raw calorimetry data and binding isotherm for 2009s (SEQ ID NO:3). PfLDH is titrated with aptamers by serial injections, and the single-binding site model is used to fit the binding isotherms.

FIG 5 is a graph showing the result of isothermal titration calorimetry analysis of the binding between PfLDH aptamer (SEQ ID NO:1; 2004s) and recombinant *Plasmodium falciparum* lactate dehydrogenase. The upper panel is the raw calorimetry data of the titration of PfLDH with serial injections of PfLDH aptamer (SEQ ID NO:1; 2004s). The lower panel is the binding isotherm resulting from integration of raw calorimetry data after correction for the heat of aptamer dilution.

FIG 6 is a graph showing the result of isothermal titration calorimetry analysis of the binding between PfLDH aptamer (SEQ ID NO:2; 2008s) and recombinant *Plasmodium falciparum* lactate dehydrogenase. The upper panel is the raw calorimetry data of the titration of PfLDH with serial injections of PfLDH aptamer (SEQ ID NO:2; 2008s). The lower panel is the binding isotherm resulting from integration of raw calorimetry data after correction for the heat of aptamer dilution.

FIG 7 is a graph showing the result of isothermal titration calorimetry analysis of the binding between PfLDH aptamer (SEQ ID NO:3; 2009s) and recombinant *Plasmodium falciparum* lactate dehydrogenase. The upper panel is the raw calorimetry data of the titration of PfLDH with serial injections of PfLDH aptamer (SEQ ID NO:3; 2009s). The lower panel is the binding isotherm resulting from integration of raw calorimetry data after correction for the heat of aptamer dilution.

FIG 8 is a graph showing the result of isothermal titration calorimetry analysis of the binding between PfLDH aptamer (SEQ ID NO:4; 2021s) and recombinant *Plasmodium falciparum* lactate dehydrogenase. The upper panel is the raw calorimetry data of the titration of PfLDH with serial injections of PfLDH aptamer (SEQ ID NO:4; 2021s). The lower panel is the binding isotherm resulting from integration of raw calorimetry data after correction for the heat of aptamer dilution.

FIG 9 shows the binding of the tetrameric PfLDH to two DNA aptamers of the present invention. **(A)** Crystal structure of tetrameric LDH in complex with two specific embodiments of the DNA aptamers of the present invention. **(B)** Structure of a specific embodiment of the DNA aptamer of the present invention in the complexed state, as well as the nucleic acid sequence of that DNA aptamer. **(C)** Isothermal titration calorimetry titrations for aptamer binding to PfLDH (left) and human LDHA1 and LDHB (right).

FIG. 10 shows the substrate specificity loop at the protein: the aptamer binding: the aptamer binding interface determines the discrimination of the aptamer for PfLDH over hLDH. **(A)** Direct interactions of DNA aptamer with PfLDH. **(B)** Discrimination of aptamer binding is due to the difference in the substrate specificity loop. Substrate specificity loops of PfLDH and hLDHB are shown in the **(B)**.

FIG. 11 shows aptamers conjugated to gold nanoparticles for use in molecular diagnostics. **(A)** Principle of approach. **(B)** TEM of nanoparticles alone (left), aptamer-conjugated nanoparticles (center) and aptamer-conjugated nanoparticles in the presence of PfLDH (right). **(C)** Qualitative observation of using nanoparticle to detect PfLDH specifically at 25 ng/ μ l. **(D)** Quantitative absorbance of data in **(C)**.

BRIEF DESCRIPTION OF THE SEQUENCES

[0015]

- 5 **SEQ ID NO:1** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* lactate dehydrogenase (LDH).
- SEQ ID NO:2** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* lactate dehydrogenase (LDH).
- 10 **SEQ ID NO:3** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* lactate dehydrogenase (LDH).
- SEQ ID NO:4** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* lactate dehydrogenase (LDH).
- 15 **SEQ ID NO:5** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- SEQ ID NO:6** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- 20 **SEQ ID NO:7** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- SEQ ID NO:8** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- 25 **SEQ ID NO:9** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- SEQ ID NO:10** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- 30 **SEQ ID NO:11** is the nucleic acid sequence of a DNA aptamer binding to *Plasmodium falciparum* histidine-rich protein II (HRP II).
- SEQ ID NO:12** is the nucleic acid sequence of a primer useful according to the present invention.
- 35 **SEQ ID NO:13** is the nucleic acid sequence of a primer useful according to the present invention.
- SEQ ID NO:14** is the nucleic acid sequence useful according to the present invention.
- 40 **SEQ ID NO:15** is the nucleic acid sequence of a primer useful according to the present invention.
- SEQ ID NO:16** is the nucleic acid sequence of a primer useful according to the present invention.
- 45 **SEQ ID NO:17** is the nucleic acid sequence of a primer useful according to the present invention.
- SEQ ID NO:18** is the nucleic acid sequence of a primer useful according to the present invention.
- 50 **SEQ ID NO:19** is the nucleic acid sequence of a primer useful according to the present invention.
- SEQ ID NO:20** is the nucleic acid sequence of a primer useful according to the present invention.
- 55

DETAILED DESCRIPTION

[0016] Provided are nucleic acid aptamers binding to *Plasmodium* lactate dehydrogenase (LDH) or *Plasmodium* his-

tidine-rich protein II (HRPII). The nucleic acid aptamers can comprise DNA or RNA nucleotides. In one embodiment, the present invention provides DNA aptamers binding to Plasmodium lactate dehydrogenase (LDH) Provided are eleven different sequences of DNA aptamers comprising nucleic acid sequences selected from SEQ ID NOs: 1-11. Four of these aptamers (SEQ ID NOs: 1-4) were selected against Plasmodium lactate dehydrogenase and seven of these

5 aptamers (SEQ ID NOs: 5-11) were selected against Plasmodium histidine Rich Protein II.
[0017] Isothermal titration calorimetry results show tight binding of the aptamers of the present invention to their respective targets (Plasmodium lactate dehydrogenase and Plasmodium histidine Rich Protein II) (Figures 4-8). These aptamers also show negligible binding to control proteins, demonstrating specific binding to the respective targets. In particular, the Plasmodium LDH aptamers of the present invention do not bind to human LDH.

10 **[0018]** The present invention also provides uses of the nucleic acid aptamers of the present invention for rapid and sensitive detection of Plasmodium in a sample, as well as for detection of malaria in a subject.

Aptamers Binding to Plasmodium LDH and Plasmodium HRPII

15 **[0019]** The present disclosure provides nucleic acid aptamers binding to Plasmodium lactate dehydrogenase (LDH) or Plasmodium histidine-rich protein II (HRPII). The nucleic acid aptamers can comprise DNA or RNA nucleotides. In one example, the present disclosure provides DNA aptamers binding to *Plasmodium* lactate dehydrogenase (LDH) or *Plasmodium* histidine-rich protein II (HRPII).

20 **[0020]** In one embodiment, the present invention provides nucleic acid aptamers binding to, or binding specifically to, *Plasmodium* lactate dehydrogenase (LDH), wherein the aptamer comprises any one of SEQ ID NOs: 1-4:

5'-ACGCGAGCAGGTGGTAGAATCATAATGGCCTGATC-3' (SEQ ID NO:1) (2004s);
 5'-CTGGGCGGTAGAACCATAGTGACCCAGCCGTCTAC-3'(SEQ ID NO:2) (2008s);
 5'-TAGGTGGCCAGAAGGTAGAACCATAGTGGTCTGGTA-3'(SEQ ID NO:3) (2009s); and
 25 5'-AGAATGGCGGGAGAGCCTTAGCGACCATTTCGTAC-3'(SEQ ID NO:4) (2021s).

Also provided are nucleic acid aptamers binding to, or binding specifically to, *Plasmodium* histidine-rich protein II (HRPII), wherein the aptamer comprises any one of SEQ ID NOs: 5-7:

30 5'-AGCGCATTTCATGCGCTCCCGCTTATGCGGGGCGGCCACGTGGAAACCCGGT TTCGCTTGTCT-
 GCTAGCC-3' (SEQ ID NO:5) (2101s);
 5'-TGCCCACTTATGTTCCGCCCCCCCCCTTGTCTC-3'(SEQ ID NO:6) (2105s);
 5'-GCTTATCCGATGCAGACCCCTTCGGTCCTGCCCTC-3'(SEQ ID NO:7) (2106s);
 5'-TGGTCATGCCGTTGGGAGTATCATTCCCCGTACGC-3'(SEQ ID NO:8) (2112s);
 35 5'-CACTCCACTGAGAACTTGCGAGTGGTCCCATTTTACCTAGCCGTCCCGCACT GCTGCTTTCTGTGCG-
 GACCGTATC-3'(SEQ ID NO:9) (2115s);
 5'-CTGGGGGGGTTCTAGGGGGGGGGCACTTATCTGCA-3'(SEQ ID NO:10) (2126s); and
 5'-TTATTGGGGGGGTTAGGGGGGGGCTTTTATCACT-3'(SEQ ID NO:11) (2144s).

40 **[0021]** As shown in the results of isothermal titration calorimetry (Figure 4), the aptamer with the addition of two 18-mer sequences on both 5' and 3' ends of SEQ ID NO:3 has similar stoichiometry, dissociation constant, and binding affinity to Plasmodium LDH (similar binding affinity in nanomolar range), when compared to the aptamer having SEQ ID NO:3.

45 **[0022]** In certain embodiments, the present invention provides nucleic acid aptamers binding to Plasmodium lactate dehydrogenase (LDH), wherein the aptamer comprises a nucleic acid sequence having at least about 80% identity to any of SEQ ID NOs: 1-4, or any percentages higher than 80%, such as, at least 85%, 90%, 92%, 93%, 94%, 95%, or 98%.

[0023] In certain examples, the present disclosure provides nucleic acid aptamers binding to Plasmodium histidine-rich protein II (HRPII), wherein the aptamer comprises a nucleic acid sequence having at least about 80% identity to any of SEQ ID NOs: 5-11, or any percentages higher than 80%, such as, at least 85%, 90%, 92%, 93%, 94%, 95%, or 98%.

50 **[0024]** In certain embodiments, the present invention provides nucleic acid aptamers binding to Plasmodium lactate dehydrogenase (LDH), wherein the aptamer comprises a nucleic acid sequence with no more than 8, 7, 6, 5, 4, 3, 2, or 1 modifications (such as addition, deletion, substitution) of nucleic acids to any of SEQ ID NOs: 1-4. Provided are also nucleic acid aptamers binding to *Plasmodium* histidine-rich protein II (HRPII), wherein the aptamer comprises a nucleic acid sequence with no more than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 modifications (such as insertion,
 55 deletion, substitution) of nucleic acids to SEQ ID NOs: 5-11. Disclosed are nucleic acid aptamers binding specifically to *Plasmodium* lactate dehydrogenase (LDH) or *Plasmodium* histidine-rich protein II (HRPII). Further disclosed are nucleic acid aptamers binding specifically to *Plasmodium falciparum* lactate dehydrogenase (LDH) or *Plasmodium falciparum* histidine-rich protein II (HRPII).

[0025] "Specific binding" or "specificity" refers to the ability of an aptamer to detectably bind an epitope presented on an antigen, while having relatively little detectable reactivity with other proteins or structures. Specificity can be exhibited by, e.g., an about 10:1, about 20:1, about 50:1, about 100:1, 10,000:1 or greater ratio of affinity/avidity in binding to the specific antigen versus nonspecific binding to other irrelevant molecules.


[0026] Unless otherwise specified, as used herein percent sequence identity and/or similarity of two sequences can be determined using the algorithm of Karlin and Altschul (1990), modified as in Karlin and Altschul (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul *et al.* (1990). BLAST searches can be performed with the NBLAST program, score = 100, wordlength = 12, to obtain sequences with the desired percent sequence identity. To obtain gapped alignments for comparison purposes, Gapped BLAST can be used as described in Altschul *et al.* (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (NBLAST and XBLAST) can be used. See NCBI/NIH website.

Modification of Aptamer Sequences

[0027] Various modifications can be made to the aptamers to reduce exonuclease degradation and increase their lifetimes for diagnostics or for therapy. Modification of the 3' end of the aptamer with inverted thymidine, deoxythymidine nucleotide, and polyethylene glycol (PEG) can reduce degradation of the oligonucleotide aptamer and increases stability of the aptamer. In one embodiment, PEG has an average molecular weight from about 20 to 80 kDa.

[0028] Further, the phosphodiester linkages of the deoxyribose-phosphate backbone of the aptamer can also be modified to improve stability.


[0029] In one embodiment, the nucleic acid aptamer of the present invention is an oligonucleic acid molecule comprising repeating units of the structure shown in Formula 1. Wavy lines demarcate one nucleotide and/or repeat unit from a neighboring nucleotide and/or repeat unit.



[0030] In one embodiment, each repeat unit of Formula 1 has a deoxyribose moiety linked to one of the common nucleotide bases (B), such as, adenosine, cytidine, guanosine, thymidine, and uridine. The base (B) for each repeating unit is independent from the other repeat units. The nucleotide sequences disclosed herein describe the order of appearance of bases (B) in an aptamer from the repeat unit on the 5' end of the aptamer to the 3' end of the aptamer.

[0031] In one embodiment, "L" is a linker group that links the deoxyribose moiety of adjacent repeat units. In the well-known structure of DNA, the L group is a phosphate group PO_4H , which can exist as a salt or in a neutral protonated form. The deoxyribose moiety together with the linker group forms the backbone of the aptamer, where the nucleotide base "B" varies independently between repeat units. The majority of the linker groups (L) forming the repeat units of Formula 1 in the aptamer are phosphate groups. As such, a majority of the backbone of the aptamer can be referred to as a deoxyribose-phosphate backbone. Many nuclease enzymes exist that can degrade oligonucleotide molecules without specificity for the specific nucleotide base sequence of the oligonucleotide molecule. Without wishing to be bound by any one particular theory, linker groups "L" other than phosphate can be incorporated into an oligonucleotide or aptamer to prevent degradation by nucleases.


[0032] In one embodiment, L can be replaced with a group as shown in Formula 2, where X_{1-4} are independently O or S. X_2 and X_3 can be bonded to either the 3' carbon or the 5' carbon of a deoxyribose moiety. In one embodiment, X_1 is O and X_4 is O that can be either protonated or unprotonated. In another embodiment, one or more of X_2 and/or X_3 is S and X_1 and X_4 are O, where O can be either protonated or unprotonated. Where one of X_2 and/or X_3 are S, the aptamer can be referred to as having a thioester linkage in the deoxyribose-phosphate backbone.



5

[0033] In another embodiment, the linker group "L" is an amide-containing group as shown in Formula 3, where R can be selected from hydrogen substituted or unsubstituted C₁-C₁₀ hydrocarbyl group. A "hydrocarbon" or "hydrocarbyl" refers to organic compounds or radicals consisting exclusively of the elements carbon and hydrogen. Hydrocarbyl includes alkyl, alkenyl, alkynyl, and aryl moieties. Hydrocarbyl also includes alkyl, alkenyl, alkynyl, and aryl moieties substituted with other aliphatic, cyclic or aryl hydrocarbon groups, such as alkaryl, alkenaryl and alkynaryl. In one embodiment, the linker group "L" is a group having Formula 3, and the aptamer comprises amide linkage(s) in the deoxyribose-phosphate backbone. The "NR" group of Formula 3 can be bonded to either the 3' carbon or the 5' carbon of a deoxyribose moiety. In one embodiment, R is methoxymethyl or methoxyethyl.

20



[0034] In some embodiments, the nucleic acid aptamer has from about 14 to about 100 nucleotide bases and/or repeat units, or any number of nucleotide bases or repeating units between 14 to 100, such as between 20 to 50 nucleotide bases and/or repeating units. In some embodiments, the aptamer has from about 14 to about 50 nucleotide bases and/or repeat units. In some embodiments, the aptamer has from about 30 to about 35 nucleotide bases and/or repeat units.

[0035] In some embodiments, the aptamer comprises one or more repeating units having the linker "L" selected from Formulae 2-3. In some embodiments of the aptamer of the present invention, the number of repeating units having the linker "L" selected from Formulae 2-3 is between 1 to 15, or any number therebetween such as for example, 1 to 10, 2 to 8, and 3 to 5. Linker groups in repeat units not selected from Formulae 2-3 are phosphate.

[0036] In some embodiments of the aptamer of the present invention, about 10% to about 100% (or any percentages therebetween such as about 20% to about 90%, about 30% to about 50%) of the repeat units have the linker "L" selected from Formulae 2-3. In some embodiments of the aptamer of the present invention, about 10% to about 70% of the repeat units have the linker "L" selected from Formulae 2-3. In some embodiments of the aptamer of the present invention, about 10% to about 50% of the repeat units have the linker "L" selected from Formulae 2-3. In some embodiments of the aptamer of the present invention, about 10% to about 30% of the repeat units have the linker "L" selected from Formulae 2-3. In some embodiments of the aptamer of the present invention, about 10% to about 20% of the repeat units have the linker "L" selected from Formulae 2-3. Linker groups in repeat units not selected from Formulae 2-3 are phosphate.

[0037] Various nucleases are exonucleases that degrade oligonucleotides from the 5' or 3' end. As such, in one embodiment a linker group L selected from Formulae 2-3 is located within about 5 repeat units from the 5' or the 3' end of the aptamer. In another embodiment, a linker group L selected from Formulae 2-3 is located within about 3 repeat units from the 5' or the 3' end of the aptamer. In yet another embodiment, a linker group L selected from Formulae 2-3 is part of the repeat unit on the 5' or the 3' end of the aptamer.

[0038] Degradation of the aptamers can also be reduced by the inclusion of modified nucleotide bases (B). The pyrimidine nucleotide bases, cytosine, thymine and uracil can be replaced with alkylated pyrimidines. Examples of alkylated pyrimidines include pseudoisocytosine; N4,N4-ethanocytosine; 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil; 5-fluorouracil; 5-bromouracil; 5-carboxymethylaminomethyl-2-thiouracil; 5-carboxymethylaminomethyl uracil; dihydrouracil; 1-methylpseudouracil; 3-methylcytosine; 5-methylcytosine; 5-methylaminomethyl uracil; 5-methoxy amino methyl-2-thiouracil; 5-methoxycarbonylmethyluracil; 5-methoxyuracil; uracil-5-oxyacetic acid methyl ester; pseudouracil; 2-thiocytosine; 5-methyl-2-thiouracil, 2-thiouracil; 4-thiouracil; 5-methyluracil; N-uracil-5-oxyacetic acid methylester; uracil 5-oxyacetic acid; 2-thiocytosine; 5-propyluracil; 5-propylcytosine; 5-ethyluracil; 5-ethylcytosine; 5-butyluracil; 5-pentyluracil; 5-pentylcytosine; methylpseudouracil; and 1-methylcytosine. The purine nucleotide bases, adenine and guanine, can be replaced by alkylated purines. Examples alkylated purines include 8-hydroxy-N6-methyladenine; inosine; N6-isopentyl-adenine; 1-methyladenine; 1-methylguanine; 2,2-dimethylguanine; 2-methyladenine; 2-methylguanine; N6-methyladenine; 7-methylguanine; 2-methylthio-N6-isopentenyladenine; and 1-methylguanine.

55

[0039] In one embodiment, at least one deoxyribose or ribose of the nucleic acid aptamer is replaced with a morpholine ring. In one embodiment, at least one phosphorothioate or phosphodiester linkage of the nucleic acid aptamer is replaced with phosphorodiamidate.

[0040] The nucleotides that can be substituted for natural nucleotides of DNA have a base moiety that can include, but is not limited to, inosine, 5-fluorouracil, 5-bromouracil, hypoxanthine, 1-methylguanine, 5-methylcytosine, and tritylated bases. The sugar moiety of the nucleotide in a sequence can also be modified and includes, but is not limited to, arabinose, xylulose, and hexose. In addition, the adenine, cytosine, guanine, thymine, and uracil bases of the nucleotides can be modified with acetyl, methyl, and/or thio groups. Sequences containing nucleotide substitutions, deletions, and/or insertions can be prepared and tested using standard techniques known in the art.

Detection of Plasmodium Infections

[0041] The present invention also provides methods for diagnosis of malaria and for detection of Plasmodium (such as *Plasmodium falciparum*) in a sample.

[0042] In one example, the method for diagnosing Plasmodium infection or malaria in a subject comprises:

obtaining a biological sample from a subject;
 contacting the biological sample with a nucleic acid aptamer comprising any one of SEQ ID NOs: 1-11, or a sequence having at least 85% identity to any one of SEQ ID NOs: 1-11;
 determining the formation of the binding complex between the nucleic acid aptamer and Plasmodium lactate dehydrogenase (LDH) or Plasmodium histidine-rich protein II (HRP II),
 wherein the presence of the binding complex indicates that the subject has Plasmodium infection.

[0043] In one example, the method comprises contacting a sample with a nucleic acid aptamer comprising any one of SEQ ID NOs: 1-4, or a sequence having at least 85% identity to any one of SEQ ID NOs: 1-4, and determining the formation of the binding complex between the nucleic acid aptamer and Plasmodium lactate dehydrogenase (LDH). Disclosed is a method for detecting the presence of Plasmodium in a sample, wherein the method comprises:

contacting a sample with a nucleic acid aptamer comprising any one of SEQ ID NOs: 1-11, or a sequence having at least 85% identity to any one of SEQ ID NOs: 1-11;
 determining the formation of the binding complex between the nucleic acid aptamer and Plasmodium lactate dehydrogenase (LDH) or Plasmodium histidine-rich protein II (HRP II), wherein the formation of the binding complex indicates the presence of Plasmodium in the sample.

[0044] In a further example, the method further determines the concentration or amount of Plasmodium in the sample comprising determining the concentration or amount of the binding complex between the aptamer of the present invention and Plasmodium lactate dehydrogenase (LDH) or Plasmodium histidine-rich protein II (HRP II).

[0045] In another example, the method comprises contacting a sample with a nucleic acid aptamer comprising any one of SEQ ID NOs: 5-11, or a sequence having at least 85% identity to any one of SEQ ID NOs: 5-11, and determining the formation of the binding complex between the nucleic acid aptamer and Plasmodium histidine-rich protein II (HRP II).

[0046] Aptamers against lactate dehydrogenase can be used for pan-species detection of Plasmodium, whilst aptamers against histidine rich protein II can be used for specific detection of *Plasmodium falciparum*.

[0047] The term "subject," as used herein, describes an organism, including mammals such as primates. Mammalian species that can benefit from the subject methods include, but are not limited to, apes, chimpanzees, orangutans, humans, monkeys; and domesticated and/or laboratory animals such as dogs, cats, horses, cattle, pigs, sheep, goats, chickens, mice, rats, guinea pigs, and hamsters. Typically, the subject is a human.

[0048] In certain embodiments, the present invention can be used to detect the presence of Plasmodium species including, but not limited to, *Plasmodium falciparum*, *Plasmodium yoelii*, *Plasmodium knowlesi*, *Plasmodium brasilanum*, *Plasmodium ovale*, *Plasmodium chabaudi*, *Plasmodium vivax*, *Plasmodium malaria*, *Plasmodium berghei*, *Plasmodium reichenowi*, and *Plasmodium gallinaceum*.

[0049] In one specific embodiment, the present invention can be used to detect the presence of *Plasmodium falciparum* in a sample, as well as to diagnose whether a subject has malaria caused by *Plasmodium falciparum* infection.

[0050] The aptamers of the present invention can be labeled with a detectable substance and localized in biological samples based upon the presence of the detectable substance. Examples of detectable substances include, but are not limited to, the following radioisotopes (e.g., ³H, ¹⁴C, ³⁵S, ¹²⁵I, ¹³¹I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), luminescent labels such as luminol; enzymatic labels (e.g., horseradish peroxidase, beta-galactosidase, luciferase, alkaline phosphatase, acetylcholinesterase), biotinyl groups (which can be detected by marked avidin, e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods),

predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags).

[0051] The terms "detecting" or "detect" include assaying or otherwise establishing the presence or absence of the target Plasmodium LDH or HRP II, or combinations of agent bound targets.

[0052] A "sample" (biological sample) can be any composition of matter of interest from a human or non-human subject, in any physical state (e.g., solid, liquid, semi-solid, vapor) and of any complexity. Preferably, the sample is a fluid (biological fluid). Samples preferably include human samples. The sample may be contained within a test tube, culture vessel, multi-well plate, or any other container or supporting substrate. The sample can be, for example, a cell culture or human tissue.

[0053] The term "biological sample," as used herein, includes but is not limited to a sample containing tissues, cells, and/or biological fluids isolated from a subject. Examples of biological samples include but, are not limited to, tissues, cells, biopsies, blood, lymph, serum, plasma, urine, saliva, and tears. In various embodiments, biological samples are obtained from, or derived from, blood, including plasma, serum, and blood cells.

[0054] In certain embodiment, the detection and diagnostic assays of the present invention can be performed using assay devices, such as a lateral flow device, and a device for colorimetric assays.

Kits

[0055] In one aspect, the present invention includes kits comprising the required elements for detection of Plasmodium in a sample. In one embodiment, the kit comprises one or more nucleic acid aptamers of the present invention.

[0056] Preferably, the kits comprise a container for collecting samples and an agent for detecting the presence of Plasmodium lactate dehydrogenase (LDH) or Plasmodium histidine-rich protein II (HRP II). Other components of the kit may include but are not limited to, means for collecting biological samples, means for labeling the detecting agent (binding agent), solid support, buffers, labels, tags, and preservatives. The kit can also contain a solid support such as microtiter multi-well plates, standards, assay diluent, wash buffer, adhesive plate covers, and/or instructions for carrying out a method of the invention using the kit. In one embodiment, the kit includes one or protease inhibitors (e.g., a protease inhibitor cocktail) to be applied to the biological sample to be assayed (such as blood or urine).

[0057] As used herein, the terms "label" and "tag" refer to substances that may confer a detectable signal, and include, but are not limited to, enzymes such as alkaline phosphatase, glucose-6-phosphate dehydrogenase, and horseradish peroxidase, ribozyme, a substrate for a replicase such as QB replicase, promoters, dyes, fluorescers, such as fluorescein, rhodamine compounds, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde, and fluorescamine, chemiluminescers such as isoluminol, sensitizers, coenzymes, enzyme substrates, radiolabels, particles such as latex or carbon particles, liposomes, cells, etc., which may be further labeled with a dye, catalyst or other detectable group.

[0058] As used herein, the term "conjugate" refers to a compound comprising two or more molecules bound together, optionally through a linking group, to form a single structure. The binding can be made by a direct connection (e.g., a chemical bond) between the molecules or by use of a linking group.

[0059] As used herein, the terms solid "support", "substrate", and "surface" refer to a solid phase which is a porous or non-porous water insoluble material that can have any of a number of shapes, such as strip, rod, particle, beads, or multi-welled plate. In some embodiments, the support has a fixed organizational support matrix that preferably functions as an organization matrix, such as a microtiter tray. Solid support materials include, but are not limited to, cellulose, polysaccharide such as Sephadex, glass, polyacryloylmorpholide, silica, controlled pore glass (CPG), polystyrene, polystyrene/latex, polyethylene such as ultrahigh molecular weight polyethylene (UPE), polyamide, polyvinylidene fluoride (PVDF), polytetrafluoroethylene (PTFE; TEFLON), carboxyl modified teflon, nylon, nitrocellulose, and metals and alloys such as gold, platinum and palladium. The solid support can be biological, non-biological, organic, inorganic, or a combination of any of these, existing as particles, strands, precipitates, gels, sheets, pads, cards, strips, dipsticks, test strips, tubing, spheres, containers, capillaries, pads, slices, films, plates, slides, etc., depending upon the particular application. Preferably, the solid support is planar in shape, to facilitate contact with a biological sample such as urine, whole blood, plasma, serum, peritoneal fluid, or ascites fluid. Other suitable solid support materials will be readily apparent to those of skill in the art. The solid support can be a membrane, with or without a backing (e.g., polystyrene or polyester card backing), such as those available from Millipore Corp. (Bedford, MA), e.g., Hi-Flow™ Plus membrane cards. The surface of the solid support may contain reactive groups, such as carboxyl, amino, hydroxyl, thiol, or the like for the attachment of nucleic acids, proteins, etc. Surfaces on the solid support will sometimes, though not always, be composed of the same material as the support. Thus, the surface can be composed of any of a wide variety of materials, such as polymers, plastics, resins, polysaccharides, silica or silica-based materials, carbon, metals, inorganic glasses, membranes, or any of the aforementioned support materials (e.g., as a layer or coating).

EXAMPLES

[0060] Following are examples that illustrate procedures and embodiments for practicing the invention. The examples should not be construed as limiting.

EXAMPLE 1 - PROPERTIES OF APTAMERS THAT BIND TO PLASMODIUM LDH AND HRPII

[0061] The thermodynamics of the interaction between the claimed aptamers and their targets were studied by using isothermal titration calorimetry. As shown in Figures 4-8, the aptamers claimed were observed to specifically bind with nanomolar affinity (K_d) to their targets without binding to controls.

[0062] In the upper panel of Figure 5, an exothermic heat pulse was observed after each injection of 2004s to PflDH solution. Using a single site binding model to fit the data then the result shows the stoichiometry of the 2004s-PflDH complex is 0.152 ± 0.003 sites, which indicates more than one 2004s may bind per PflDH molecule in solution at 25 °C.

[0063] In the upper panel of Figure 6, an exothermic heat pulse was observed after each injection of 2008s to PflDH solution. Using a single site binding model to fit the data then the result shows the stoichiometry of 2008s-PflDH complex is 0.166 ± 0.003 sites, which indicates more than one 2008s may bind per PflDH molecule in solution at 25 °C.

[0064] In the upper panel of Figure 7, an exothermic heat pulse was observed after each injection of 2009s to PflDH solution. Using a single site binding model to fit the data then the result shows the stoichiometry of 2009s-PflDH complex is 0.451 ± 0.007 sites, which indicates two 2009s molecules may bind per PflDH molecule in solution at 25 °C.

[0065] In the upper panel of Figure 8, an exothermic heat pulse was observed after each injection of 2021s to PflDH solution. Using a single site binding model fit the data then the result shows the stoichiometry of 2021s-PflDH complex is 0.177 ± 0.003 sites, which indicates more than one 2021s may bind per PflDH molecule in solution at 25 °C.

Materials and Methods

[0066] The present disclosure describes aptamers that bind to PflDH and HRP. PflDH was obtained by molecular cloning of cDNA of PflDH from *Plasmodium falciparum* 3D7 cDNA library. The coding region of PflDH was amplified by PCR with the forward primer 5'-ATTATTGCTAGCATGGCACCAAAGCAAAAATCGTTTTAGTTG-3' (SEQ ID NO:12) and the reverse primer 5'-ATTATTCTCGAGTTAAGCTAATGCCTTCATTCTCTTAGTTTCA-3' (SEQ ID NO:13). The PCR product was digested with *NheI/XhoI*, purified by gel purification and ligated to the *NheI/XhoI* digested pET-28a vector to become the expression plasmid pET28a-PflDH.

[0067] PflDH was heterologously expressed in *Escherichia coli* BL21(DE3) pLysS. pET28a-PflDH was transformed into *Escherichia coli* BL21(DE3) pLysS cells for recombinant protein expression. An overnight culture was prepared by inoculating a single colony into LB medium with 50 µg/mL of kanamycin, and followed by incubating in 37°C overnight. For PflDH expression, the overnight culture was diluted in LB medium in 1:50 ratio and incubated at 37°C until OD₆₀₀ reached 0.5, then isopropyl-D-thiogalactopyranoside (IPTG) was added to the culture to a final concentration of 0.5 mM to induce the PflDH expression. The induced culture was further incubated in room temperature for 4 hours and cells were harvested by centrifugation.

[0068] For the extraction of recombinant PflDH from *Escherichia coli* BL21 (DE3) pLysS cells, the harvested cells were suspended into lysis buffer (50 mM Tris-HCl, pH 7.5, 0.3 M NaCl, 20 mM imidazole with protease inhibitor and Benzonase) in 1:50 of lysis buffer to cell culture ratio. The cell suspension was incubated on ice for 30 minutes and followed by sonicating for 10 minutes in 9 seconds on and 5 seconds off cycle. The bacterial cell lysate was centrifuged at 18000xg for 20 minutes and the supernatant, soluble protein extract, was collected for PflDH purification.

[0069] Recombinant PflDH was purified by affinity chromatography. The soluble protein extract was loaded on nickel affinity column and PflDH was eluted by different 150 mM of imidazole in 50 mM Tris-HCl, pH7.5, 0.3 M NaCl.

[0070] PflDH aptamers were selected by SELEX performed on PflDH immobilised magnetic beads and counter SELEX performed on human lactate dehydrogenase immobilised magnetic beads and the naked magnetic beads. The selection of PflDH aptamers was started by incubating a single-stranded DNA (ssDNA) library containing 35mer random region flanked by two 18mer priming regions (5'-CGTACGGTCGACGCTAGC-[N35]-CACGTGGAGCTCGGATCC-3') (SE ID NO:14) with the PflDH immobilised magnetic beads. After incubation, the unbound species were removed and the ssDNA-PflDH-magnetic beads complexes were suspended in 10 µL of water.

[0071] PCR was carried out for the amplification of PflDH bound species. The reaction mixtures for PCR contained 10 µL Pwo SuperYield PCR buffer, 10 mM dNTPs, 20 mM forward primer (5'-CGTACGGTCGACGCTAGC-3') (SEQ ID NO:15), 20 mM reverse primer (5'-biotin-GGATCCGAGCTCCACGTG-3') (SEQ ID NO:16), 10 µL of magnetic beads suspension, 2.5 U Pwo SuperYield DNA polymerase and water to a final volume of 100 µL. PCR conditions were denaturation at 95°C for 15 seconds, annealing at 50°C for 30 seconds and elongation at 72°C for 15 seconds for 10 cycles.

[0072] PCR products were collected and incubated with streptavidin magnetic beads. Unbiotinylated ssDNAs were eluted by adding 100 mM NaOH. The first to third round of selections, eluted ssDNAs were concentrated by ethanol

precipitation whereas QIAquick gel extraction kit was used for the following rounds. The yielded ssDNA pool was used for the next round of selection. Counter selections by using naked magnetic beads were included in the fourth and seventh rounds, hLDHA1 immobilised magnetic beads were included in sixth and tenth rounds and hLDHB immobilised magnetic beads were included in fifth and ninth rounds. Ultimately, the ssDNA pool from the twentieth round was cloned and the colonies were picked for sequencing.

[0073] A similar approach was taken for the expression, purification and selection of aptamers against HRP2.

[0074] For isothermal titration calorimetry, PfLDH aptamers and PfLDH were dialysed in 25 mM Tris-HCl, pH 7.5 containing 0.1 M NaCl and 20 mM imidazole. All the buffers, protein and aptamers were degassed for 10 minutes at 25°C prior to ITC experiments. All the experiments were performed in 25 mM Tris-HCl, pH 7.5 containing 0.1 M NaCl and 20 mM imidazole at 25°C using iTC₂₀₀ microcalorimeter (MicroCal Inc.). ITC experiment was carried out by injecting PfLDH aptamer into PfLDH. The ITC data were analysed by using Origin v7.0 software (MicroCal Inc.) to integrate the titration curves for extracting the thermodynamic parameters, stoichiometry (N), equilibrium association constant [K_d ($=K_d^{-1}$)] and the binding enthalpy (ΔH).

[0075] A similar approach was taken for the investigation of the binding of aptamers to HRPII.

EXAMPLE 2 - STRUCTURAL BASIS OF DNA APTAMER RECOGNITION OF PLASMODIUM LACTATE DEHYDROGENASE

[0076] This Example shows the crystal structure and application of one embodiment of a DNA aptamer against a malaria panspecies diagnostic target, *Plasmodium falciparum* lactate dehydrogenase (PfLDH). This example also shows a new mechanism of molecular recognition of *Plasmodium falciparum* lactate dehydrogenase (PfLDH) by the hairpin DNA aptamer of the present invention.

[0077] PfLDH and two related human homologues hLDHA1 and hLDHB were cloned, expressed in *E. coli* and purified. DNA aptamers were selected from a pool with a 35-base random region against purified PfLDH and counterselected to remove aptamers bound to the closely related human homologues hLDHA1 and hLDHB. After 20 rounds of selection, 51 sequences were obtained and aligned, revealing the presence of conserved sequence signatures and motifs. A number of these aptamers were observed to specifically bind to PfLDH (the most promising with K_d in the range 20-50 nM) without binding to hLDHA1 and hLDHB, as determined by isothermal titration calorimetry (**Fig. 9c**), electrophoretic mobility shift assay, and surface plasmon resonance (SPR) spectroscopy.

[0078] A single DNA aptamer which showed significant promise (2008s) (SEQ ID NO:2) in the three binding assays (with sequence shown in **Fig. 9b**) was taken forward for structural analysis.

[0079] The crystal structure of the PfLDH: aptamer complex was determined by X-ray crystallography (**Fig. 9a**). The four molecules of PfLDH in the asymmetric unit are organized into a tetramer with characteristic 222 symmetry, similar to previously determined PfLDH X-ray crystallographic structures.

[0080] Each of the two ssDNA aptamers were seen bound to the Q-axis dimers formed by molecules A and B (or molecules C and D), spanning across the large cofactor domains of the two juxtaposed monomers.

[0081] The aptamer did not form a G-quadruplex structure, but showed a hairpin structure (**Fig. 9b**). Clear electron density was observed for the first 27 bases of both aptamers in the asymmetric unit; no electron density was observed beyond the hairpin structure for the last 8 bases at the 3' end of the aptamer, probably because the aptamer tail is disordered. The global fold of the aptamer adopted a distorted hairpin structure with a C¹⁵A¹⁶T¹⁷A¹⁸ apical tetraloop, an asymmetric internal loop with six bases in one strand (C⁶G⁷G⁸T⁹A¹⁰G¹¹) and one base in the other (A²²), and a terminal B-form helix. The aptamer contains seven Watson-Crick type base-pairs, five of which form the terminal B-form helical stem DNA duplex. The remaining two Watson-Crick type base pairs, together with non-canonical base pairing between A¹² and G²¹, form the DNA duplex flanking between the internal loop and apical loop. Similar to the terminal B-helical stem, this short duplex assumes B-helical geometry. The two B-helices are oriented in an ~140° angle, resulting in an overall bent shape for the DNA aptamer. C⁶ of the internal loop is involved in stacking interaction with its neighbors, G⁵ of the stem B-helix. As a result, C⁶, G⁷ and G⁸ of the internal loop remain within the stem B-helix as they continuously stack. G⁷ and A¹⁰ form a non-canonical base pair in the internal loop. T⁹ is flipped out of the internal loop and is engaged in aptamer: PfLDH interactions as described below. In the apical tetraloop, both T¹⁷ and A¹⁸ are involved in stacking interactions and remain in the helical axis of the short duplex flanking the apical loop and internal loop. C¹⁵ and A¹⁶ are flipped out of the apical loop, the position of C¹⁵ is stabilised by interaction of N4 with the backbone phosphate of C¹⁴, while A¹⁶ is involved in indirect interactions with PfLDH via waters.

[0082] Each aptamer interacts through extensive salt bridges with the cofactor binding sites of two PfLDH molecules of the Q-axis dimer (**Fig. 10a**). The interface between the aptamer and PfLDH buries a solvent-accessible surface of 1276Å² on the aptamer and 895Å² and 394Å² for molecules A/C and B/D of PfLDH respectively. The aptamer interacts with the substrate specificity loop at the tip of the cofactor binding sites of the two adjacent PfLDH molecules via distinct regions. Molecule A/C interacts with the internal loop whilst molecule B/D interacts with the apical tetraloop.

[0083] The aptamer interacts with Molecule A of the PfLDH dimer via the internal loop. The bases C⁶, G⁷, G⁸, G¹¹

and A²² of the internal loop of the DNA aptamer form extensive base interactions with residues 102A-108A of the extended substrate specificity loop, backbone contacts are also seen with Lys102A and Lys106A forming contacts with the phosphates of A¹⁰ and A²² respectively. The flipped base T⁹ extends away from the bent hairpin structure into the adenine end of the NADH binding cleft and is stabilized by hydrogen bonds between N3 of the T⁹ base and the sidechain of Asp53A, and O4 of the T⁹ base with the amino nitrogen of Ile54A.

[0084] Compared to the interaction of the internal loop and Molecule A of PflDH, the interaction of the apical tetraloop with Molecule B of PflDH is less extensive and involves indirect contacts via water molecules. Direct backbone contacts are observed with phosphate groups of A18 and G19 of the tetraloop interacting with the sidechains of Lys62B and His250B of the NADH binding cleft. Similar to T9 in the internal loop, the flipped base A16 extends into the adenine end of the NADH binding cleft of molecule B. However, instead of direct interactions with the protein, the position of A16 of the apical loop is stabilised via indirect interactions with the protein via water molecules (**Fig. 10a**).

[0085] The extensive interaction of this aptamer to the unique extended substrate specificity loop of PflDH (amino acids 102-108) contributes to the high specificity of aptamer binding to PflDH and not human LDH. Indeed, superposition of human LDHB (PDB accession code: 1T2F) with our PflDH: aptamer structure revealed that human LDH lacks the extension at the substrate specificity loop, resulting in an inability to reach the aptamer (**Fig. 10b**). Thus, binding of the aptamer to human LDH would not occur, which both proves that the counterselection strategy was successful and provides a structural explanation for the specificities determined in our binding assays.

[0086] The PflDH aptamer was conjugated to gold nanoparticles (AuNPs) to develop a colorimetric assay for PflDH (**Fig. 11a**). Transmission electron microscopy revealed that the AuNP aggregated specifically in the presence of PflDH (**Fig. 11b**). The characteristic ruby red color (absorbance peak of 520 nm) was specifically lost in the presence of 25 ng/ μ l PflDH but not in the presence of hLDHA1, hLDHB or other controls (**Fig. 11c and 11d**). The limit of detection was determined to be 0.73 ng/ μ l PflDH.

[0087] The results show that the DNA aptamer identified can specifically recognize PflDH due to its unique loop that is not present in human LDH. The results also show that the DNA aptamers of the invention can be used for diagnostic of malaria.

[0088] In conclusion, this Example shows the identification, structural characterization, and application of a specific embodiment of a hairpin DNA aptamer against PflDH. This Example shows that the specificity of the DNA aptamer is achieved by discriminatory binding to a loop that is present only in the *Plasmodium*, but not present in the human lactate dehydrogenase, proving at the molecular level the utility of counter-selection in aptamer selection. The results show that the DNA aptamers of the present invention can be used for malaria diagnosis.

Materials and Methods

Cloning, expression and purification of PflDH and human lactate dehydrogenases (hLDHA1 & hLDHB)

[0089] Open reading frame (ORF) of *PflDH* was amplified by PflDH-S(ATTATTGCTAGCATGGCACAAAAG-CAAAAATCGTTTTAGTTG) (SEQ ID NO:12), and PflDH-AS (ATTATTCTCGAGTTAAGCTAATGCCTTCATTCTCT-TAGTTTCA) (SEQ ID NO:13), followed by ligation into the *NcoI/XhoI* digested pET-28a (Novagen) vector.

[0090] ORF of *hLDHA1* was amplified by hLDHA1-S (ATTATTGAATTCATGGCAACTCTAAAGGATCAGCTGA) (SEQ ID NO:17), and hLDHA1-AS (ATTATTAAGCTTTTTAAAATTGCAGCTCCTTTTGGATC) (SEQ ID NO:18), followed by ligation into the *NheI/HindIII* digested pET-28a (Novagen) vector.

[0091] ORF of *hLDHB* was amplified by hLDHB-S (ATTATTGCTAGCATGGCAACTCTTAAGGAAAACTCATT-GCACC) (SEQ ID NO:19) and hLDHB-AS (ATTATTGCGGCCGCTCACAGGTCTTTTAGGTCCTTCTGG) (SEQ ID NO:20), followed by ligation into the *NheI/NotI* digested pET-28a (Novagen) vector.

[0092] The plasmids were transformed into *E. coli* BL21 (DE3) pLysS for IPTG induced expression. The expressed proteins were further purified by using HisTrap column (GE Lifesciences).

In vitro selection of DNA aptamers

[0093] A single-stranded DNA (ssDNA) library containing a 35-mer random region flanked by two 18-mer priming regions (5'-CGTACGGTCGACGCTAGC-[N35]-CACGTGGAGCTCGGATCC-3' (SEQ ID NO:14)) was used as starting material for *in vitro* selection. 2 nmol of ssDNA library was incubated with 1nmol of target protein that was conjugated with Ni-NTA magnetic beads. The unbound species were removed and the ssDNA-protein-magnetic beads complexes were suspended in 10 μ L of water for PCR amplification of PflDH bound species by using forward primer (5'-CGTACG-GTCGACGCTAGC-3') (SEQ ID NO:15) and reverse primer with biotinylated 5' end (5'-biotin-GGATCCGAGCTC-CACGTG-3') (SEQ ID NO:16).

[0094] PCR conditions were denaturation at 95°C for 15 seconds, annealing at 50°C for 30 seconds and elongation at 72°C for 15 seconds for 10 cycles. Enriched DNA aptamer pool was purified by streptavidin magnetic beads followed

by alkynylation to remove the unbiotinylated complementary strand. The resultant ssDNA pool was used for the next round of selection. Counter-selections by using Ni-NTA magnetic agarose beads, hLDHA1 immobilized Ni-NTA magnetic agarose beads and hLDHB immobilized Ni-NTA magnetic agarose beads were incorporated in between the SELEX cycles.

5

Crystallization and structure solution

[0095] The complex between PflDH and the DNA aptamer was formed by mixing protein and DNA in a 1:1 molar ratio for crystallization. Screening of crystallization conditions was performed using MOSQUITO® liquid handler (TTP Labtech) at 291 K using sitting-drop vapor diffusion method by mixing 100 nl protein:DNA complex with an equal volume of crystallization solution. Of the 576 conditions screened, several hits for protein:DNA complex were observed. The best crystals were obtained under condition 26 of the Hampton Research Matrix Screen (0.2 M KC1, 0.1 M magnesium acetate, 0.05 M sodium cacodylate pH 6.5, 10% w/v PEG 8000).

10

[0096] The crystal of PflDH:aptamer complex was cryoprotected in a solution of 30% glycerol mixed with the reservoir solution. Data were collected at beamline 13B1, NSRRC, Taiwan. Data were indexed and integrated with HKL2000. The structure was solved by molecular replacement using the program PHASER with the structure of LDH (PDB accession number 1LDG). The structure of the DNA aptamer was built manually with COOT and the complex structure was refined in Refmac.

15

Isothermal titration calorimetry (ITC)

[0097] For isothermal titration calorimetry, PflDH aptamers and PflDH were dialyzed in 25 mM Tris-HCl, pH 7.5 containing 0.1 M NaCl and 20 mM imidazole. All the buffers, protein and aptamers were degassed for 10 minutes at 25°C prior to ITC experiments. All the experiments were performed in 25 mM Tris-HCl, pH 7.5 containing 0.1 M NaCl and 20 mM imidazole at 25°C using iTC200 microcalorimeter (MicroCal Inc.). ITC experiment was carried out by injecting PflDH aptamer into PflDH. The ITC data were analyzed by using Origin v7.0 software (MicroCal Inc.) to integrate the titration curves for extracting the thermodynamic parameters, stoichiometry (N), equilibrium association constant [$K_a (=Kd^{-1})$] and the binding enthalpy (ΔH).

25

Electrophoretic mobility shift assay (EMSA)

[0098] Electrophoresis mobility shift assay was carried out by incubating 25 nM PflDH aptamer 2008s with different proteins at concentrations ranging from 0 - 22.8 μM in 25 mM Tris-HCl, pH 7.5 containing 0.1 M NaCl and 20 mM imidazole in room temperature for 1 hour. Reactions were loaded on 12% native polyacrylamide gels and visualised by SYBR® gold nucleic acid gel stain (Invitrogen).

35

Surface plasmon resonance

[0099] Surface plasmon resonance (SPR) measurement was performed using a Biacore X100 instrument (GE Healthcare). PflDH was captured on the surface of NTA sensor chip (GE Healthcare). A running buffer containing 25 mM Tris-HCl, pH7.5, 100 mM NaCl, 20 mM imidazole and 0.005% (v/v) TWEEN 20 was used for ligand capturing. The surface of the NTA chip was primed with running buffer and PflDH aptamer, 2008s, was injected in triplicate with concentrations ranging from 0.625 - 10 μM by single-cycle analysis.

40

[0100] All experiments were performed at a flow rate of 30 $\mu\text{L min}^{-1}$ and at 25 °C. All data were referenced for surface without captured PflDH and blank injections of buffer. Sensorgrams were analyzed with Biacore X100 Plus Package evaluation software (GE Lifesciences).

45

Reparation of aptamer-immobilized gold nanoparticles

[0101] The PflDH aptamer, 2008s, was conjugated to 10 nm gold colloid (Sigma) according to the protocol described by Taton⁶ with modifications. All the incubations were carried out at room temperature in the dark. Disulfide-functionalized aptamer was reduced by 0.1 M dithiothreitol (DTT) for 30 minutes at room temperature prior to use. The reduced aptamers were purified using Sephadex G-25 (Pharmacia Biotech.). 2.5 nmol of reduced aptamers were then added to 500 μL of AuNP.

50

[0102] After 24 hours of incubation, 1 M NaCl and 0.1 M sodium phosphate buffer were added to reach final concentrations of 0.1 M NaCl and 10 mM phosphate buffer respectively. The aptamer/gold nanoparticle mixture was aged at room temperature for another 24 hours. 2008s-AuNP was separated from the solution by centrifugation followed by washing in 0.1 M NaCl/10 mM sodium phosphate buffer, pH 7. A 37-mer poly thymine (polyT) was conjugated to AuNP

55

by the above procedures as the control experiment. The aptamer-AuNP conjugates were stored in 0.3 M NaCl/0.01% sodium azide/10 mM sodium phosphate buffer, pH 7.0.

Characterization of 2008s-AuNP: UV-Vis spectrometry

[0103] The spectrometry analyses were carried out by Varioskan Flash Multimode Reader (Thermo Scientific). The spectrum was derived from at least three experiments.

Characterization of 2008s-AuNP: Transmission electron microscopy (TEM)

[0104] Oligonucleotides immobilised AuNPs were characterised by Philips CM100 TEM. Approximately 10 μ L of AuNP conjugates were coated on a 200 mesh copper grid by adding sample to the grid and dried at room temperature.

[0105] The terms "a" and "an" and "the" and similar referents as used in the context of describing the invention are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context.

[0106] Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. Unless otherwise stated, all exact values provided herein are representative of corresponding approximate values (*e.g.*, all exact exemplary values provided with respect to a particular factor or measurement can be considered to also provide a corresponding approximate measurement, modified by "about," where appropriate).

[0107] The use of any and all examples, or exemplary language (*e.g.*, "such as") provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise indicated. No language in the specification should be construed as indicating any element is essential to the practice of the invention unless as much is explicitly stated.

[0108] The description herein of any aspect or embodiment of the invention using terms such as "comprising", "having", "including" or "containing" with reference to an element or elements is intended to provide support for a similar aspect or embodiment of the invention that "consists of", "consists essentially of", or "substantially comprises" that particular element or elements, unless otherwise stated or clearly contradicted by context (*e.g.*, a composition described herein as comprising a particular element should be understood as also describing a composition consisting of that element, unless otherwise stated or clearly contradicted by context).

[0109] It should be understood that the examples and embodiments described herein are for illustrative purposes only.

SEQUENCE LISTING

[0110]

<110> The University of Hong Kong
Tanner, Julian
Cheung, Yee Wai

<120> NUCLEIC ACID APTAMERS AGAINST PLASMODIUM LACTATE DEHYDROGENASE AND HISTIDINE-RICH PROTEIN II AND USES THEREOF FOR MALARIA DIAGNOSIS

<130> UHK.167X

<150> 61/596,774

<151> 2012-02-09

<160> 20

<170> PatentIn version 3.5

<210> 1

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

EP 2 812 453 B1

<223> nucleic acid

<400> 1
 acgcgagcag gtggtagaat cataatggcc tgatc 35

5

<210> 2
 <211> 35
 <212> DNA
 <213> Artificial Sequence

10

<220>
 <223> nucleic acid

<400> 2
 ctgggcgcta gaaccatagt gaccagccg tctac 35

15

<210> 3
 <211> 36
 <212> DNA
 <213> Artificial Sequence

20

<220>
 <223> nucleic acid

<400> 3
 taggtggcca gaaggtagaa ccatagtggc ctggta 36

25

<210> 4
 <211> 34
 <212> DNA
 <213> Artificial Sequence

30

<220>
 <223> nucleic acid

35

<400> 4
 agaatggcgg gagagcctta gcgaccattc gtac 34

40

<210> 5
 <211> 70
 <212> DNA
 <213> Artificial Sequence

45

<220>
 <223> nucelic acid

<400> 5

50 **agcgcattca tgcgctcccg cttatgcggg gcggccacgt ggaaaccg tttcgcttgt 60**

tctgctagcc 70

<210> 6
 <211> 35
 <212> DNA
 <213> Artificial Sequence

55

EP 2 812 453 B1

<220>
<223> nucleic acid

<400> 11
5 ttattggggg ggtaggggg gggctttat tcaact 35

<210> 12
<211> 43
<212> DNA
10 <213> Artificial Sequence

<220>
<223> nucleic acid

<400> 12
15 attattgcta gcatggcacc aaaagcaaaa atcgtttag ttg 43

<210> 13
<211> 43
20 <212> DNA
<213> Artificial Sequence

<220>
<223> artificial sequence

<400> 13
25 attattctcg agttaagcta atgcctcat tctcttagtt tca 43

<210> 14
30 <211> 37
<212> DNA
<213> Artificial Sequence

<220>
35 <223> nucleic acid

<220>
<221> misc_feature
<222> (19) .. (19)
40 <223> n is a, c, g, or t

<400> 14
cgtagcgtcg acgctagcnc acgtggagct cggatcc 37

<210> 15
45 <211> 18
<212> DNA
<213> Artificial Sequence

<220>
50 <223> nucleic acid

<400> 15
55 cgtacggtcg acgctagc 18

<210> 16
<211> 18
<212> DNA

<213> Artificial Sequence
 <220>
 <223> nucleic acid
 5
 <400> 16
 ggatccgagc tccacgtg 18
 <210> 17
 10 <211> 37
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> nucleic acid
 <400> 17
 attattgaat tcatggcaac tctaaaggat cagctga 37
 20 <210> 18
 <211> 37
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> nucleic acid
 <400> 18
 30 attattaagc tttaaaatt gcagctcct ttggatc 37
 <210> 19
 <211> 44
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> nucleic acid
 <400> 19
 40 attattgcta gcatggcaac tcttaaggaa aaactcattg cacc 44
 <210> 20
 <211> 39
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> nucleic acid
 50 <400> 20
 attattgctg cgcgtcacag gtctttagg tcctctgg 39

Claims

- 55
1. An aptamer capable of binding to Plasmodium lactate dehydrogenase (LDH) comprising a nucleic acid sequence having at least 85% identity with any portion of SEQ ID NOs: 1, 2, 3 or 4, wherein the aptamer is from 30 nucleotide bases to 100 nucleotide bases in length.

2. An aptamer as claimed in claim 1, comprising a nucleic acid sequence selected from any of SEQ ID NOs: 1-4.
3. An aptamer as claimed in any preceding claim, wherein the aptamer is conjugated with one or more moieties selected from deoxythymidine nucleotide, inverted thymidine, and polyethylene glycol.
- 5
4. An aptamer as claimed in any of claims 1 to 3, wherein the aptamer is
- (a) an oligonucleotide having a backbone formed of deoxyribose-phosphate linkages, or
- (b) an oligonucleotide whose backbone comprises one or more deoxyribose-phosphate linkages stabilized by one or more thioester linkages and/or one or more amide linkages.
- 10
5. A kit comprising an aptamer of any preceding claim, and a carrier.
6. A method for detecting the presence of Plasmodium in a sample, comprising:
- 15
- contacting a sample with a nucleic acid aptamer as claimed in any of claims 1 to 4 determining the formation of the binding complex between the nucleic acid aptamer and LDH, wherein the formation of the binding complex indicates the presence of Plasmodium in the sample.
7. A method as claimed in claim 6 for use in the diagnosis of a Plasmodium infection in a subject, comprising:
- 20
- contacting a biological sample from the subject with a nucleic acid aptamer as claimed in any of claims 1 to 4; determining the formation of the binding complex between the nucleic acid aptamer and LDH, wherein the presence of the binding complex indicates that the subject has Plasmodium infection.
- 25
8. A method as claimed in claim 7, wherein the biological sample is a blood or urine sample.
9. A method as claimed in claim 6, further comprising determining the concentration or amount of the binding complex formed between the nucleic acid aptamer and LDH.
- 30
10. A method as claimed in any of claims 6 to 9 wherein the nucleic acid aptamer comprises a sequence selected from any of SEQ ID NOs: 1-4.
11. A method as claimed in claim 6 wherein the presence of the binding complex indicates the presence of *Plasmodium falciparum*.
- 35
12. A method as claimed in claims 7 or claim 8 wherein the presence of the binding complex indicates that the subject has *Plasmodium falciparum* infection.

40

Patentansprüche

1. Aptamer, das in der Lage ist, an Plasmodium-Lactatdehydrogenase (Plasmodium-LDH) zu binden, umfassend eine Nucleinsäuresequenz mit mindesten 85 % Identität mit einem Abschnitt der SEQ ID NO: 1, 2, 3 oder 4, wobei das Aptamer eine Länge von 30 Nucleotidbasen bis 100 Nucleotidbasen hat.
- 45
2. Aptamer nach Anspruch 1, umfassend eine Nucleinsäuresequenz, ausgewählt aus einer der SEQ ID NO: 1-4.
3. Aptamer nach einem vorherigen Anspruch, wobei das Aptamer mit einem oder mehreren Resten ausgewählt aus Deoxythymidinnucleotid, invertiertes Thymidin und Polyethylenglycol konjugiert ist.
- 50
4. Aptamer nach einem der Ansprüche 1 bis 3, wobei das Aptamer wie folgt ist
- (a) ein Oligonucleotid mit einem Rückgrat, das aus Desoxyribose-Phosphat-Verbindungen gebildet ist, oder
- (b) ein Oligonucleotid, dessen Rückgrat eine oder mehrere Desoxyribose-Phosphat-Verbindungen umfasst, die durch eine oder mehrere Thioester-Verbindungen und/oder eine oder mehrere Amid-Verbindungen sind.
- 55
5. Kit, umfassend ein Aptamer nach einem vorherigen Anspruch und einen Träger.

EP 2 812 453 B1

6. Verfahren zum Detektieren der Anwesenheit von Plasmodium in einer Probe, umfassend:

Inkontaktbringen einer Probe mit einem Nukleinsäure-Aptamer nach einem der Ansprüche 1 bis 4,
Bestimmen der Bildung des Bindungskomplexes zwischen dem Nukleinsäure-Aptamer und LDH, wobei die
Bildung des Bindungskomplexes die Anwesenheit von Plasmodium in der Probe anzeigt.

7. Verfahren nach Anspruch 6 zur Verwendung in der Diagnose einer Plasmodium-Infektion in einem Subjekt, umfassend:

Inkontaktbringen einer biologischen Probe von dem Subjekt mit einem Nukleinsäure-Aptamer nach einem der
Ansprüche 1 bis 4;
Bestimmen der Bildung des Bindungskomplexes zwischen dem Nukleinsäure-Aptamer und LDH,
wobei die Anwesenheit des Bindungskomplexes anzeigt, dass das Subjekt eine Plasmodium-Infektion hat.

8. Verfahren nach Anspruch 7, wobei die biologische Probe eine Blut- oder Urinprobe ist.

9. Verfahren nach Anspruch 6, ferner umfassend ein Bestimmen der Konzentration oder Menge des Bindungskomplex, der zwischen dem Nukleinsäure-Aptamer und LDH gebildet ist.

10. Verfahren nach einem der Ansprüche 6 bis 9, wobei das Nukleinsäure-Aptamer eine Sequenz ausgewählt aus einer der SEQ ID NO: 1-4 umfasst.

11. Verfahren nach Anspruch 6, wobei die Anwesenheit des Bindungskomplexes die Anwesenheit von *Plasmodium falciparum* anzeigt.

12. Verfahren nach Anspruch 7 oder Anspruch 8 wobei die Anwesenheit des Bindungskomplexes anzeigt, dass das Subjekt eine Infektion mit *Plasmodium falciparum* hat.

Revendications

1. Aptamère capable de se lier à la lactico-déshydrogénase de Plasmodium (LDH) comprenant une séquence d'acides nucléiques présentant une identité d'au moins 85 % avec une quelconque portion des SEQ ID n° : 1, 2, 3 ou 4, ledit aptamère faisant 30 bases nucléotidiques à 100 bases nucléotidiques de long.

2. Aptamère selon la revendication 1, comprenant une séquence d'acides nucléiques choisie parmi l'une quelconque des SEQ ID n° : 1 à 4.

3. Aptamère selon l'une quelconque des revendications précédentes, ledit aptamère étant conjugué à un ou plusieurs groupements choisis parmi un nucléotide de désoxythymidine, une thymidine inversée et un polyéthylène glycol.

4. Aptamère selon l'une quelconque des revendications 1 à 3, ledit aptamère étant

(a) un oligonucléotide possédant un squelette formé de liaisons désoxyribose-phosphate, ou
(b) un oligonucléotide dont le squelette comprend une ou plusieurs liaisons désoxyribose-phosphate stabilisées par une ou plusieurs liaisons thioester et/ou une ou plusieurs liaisons amide.

5. Kit comprenant un aptamère selon l'une quelconque des revendications précédentes, et un véhicule.

6. Procédé de détection de la présence de Plasmodium dans un échantillon, comprenant :

la mise en contact d'un échantillon avec un aptamère d'acides nucléiques selon l'une quelconque des revendications 1 à 4,
la détermination de la formation du complexe de liaison entre l'aptamère d'acides nucléiques et la LDH, ladite formation du complexe de liaison indiquant la présence de Plasmodium dans l'échantillon.

7. Procédé selon la revendication 6 destiné à être utilisé dans le diagnostic d'une infection par Plasmodium chez un sujet, comprenant :

EP 2 812 453 B1

la mise en contact d'un échantillon biologique provenant du sujet avec un aptamère d'acides nucléiques selon l'une quelconque des revendications 1 à 4 ;
la détermination de la formation du complexe de liaison entre l'aptamère d'acides nucléiques et la LDH, ladite présence du complexe de liaison indiquant que le sujet présente une infection par Plasmodium.

5

8. Procédé selon la revendication 7, ledit échantillon biologique étant un échantillon de sang ou d'urine.

9. Procédé selon la revendication 6, comprenant en outre la détermination de la concentration ou de la quantité du complexe de liaison formé entre l'aptamère d'acides nucléiques et la LDH.

10

10. Procédé selon l'une quelconque des revendications 6 à 9, ledit aptamère d'acides nucléiques comprenant une séquence choisie parmi l'une quelconque des SEQ ID n° : 1 à 4.

11. Procédé selon la revendication 6, la présence du complexe de liaison indiquant la présence de *Plasmodium falciparum*.

15

12. Procédé selon l'une quelconque des revendications 7 ou 8, la présence du complexe de liaison indiquant que le sujet présente une infection par *Plasmodium falciparum*.

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2

Aptamer Target	Aptamer Name	Sequence (5' to 3')	Length (bases)	K _D by ITC (nM)
LDH	2004s SEQ ID NO:1	ACGCGAGCAGGTGGTAGAATCATAAT GGCCTGATC	35	26
LDH	2008s SEQ ID NO:2	CTGGGCGGTAGAACCATAGTGACCCA GCCGTCTAC	35	43
LDH	2009s SEQ ID NO:3	TAGGTGGCCAGAAGGTAGAACCATAG TGGTCTGGTA	36	42
LDH	2021s SEQ ID NO:4	AGAATGGCGGGAGAGCCTTAGCGAC CATTCGTAC	34	29
HRP2	2101s SEQ ID NO:5	AGCGCATTCATGCGCTCCCGCTTATGC GGGGCGGCCACGTGGAAACCCGGTT TCGCTTGTCTGCTAGCC	70	85
HRP2	2105s SEQ ID NO:6	TGCCCACTTATGTTGCCCCCCCCCTC TTGTTCTC	35	ND*
HRP2	2106s SEQ ID NO:7	GCTTATCCGATGCAGACCCCTTCGGT CCTGCCCTC	35	ND*
HRP2	2112s SEQ ID NO:8	TGGTCATGCCGTTGGGAGTATCATTCC CCGTACGC	35	ND*
HRP2	2115s SEQ ID NO:9	CACTCCACTGAGAACTTGCGAGTGGT CCCATTTTACCTAGCCGTCCCGCACTG CTGCTTTCTGTGCGGACCGTATC	76	35
HRP2	2126s SEQ ID NO:10	CTGGGGGGGTTCTAGGGGGGGGGCA CTTATCTGCA	35	48
HRP2	2144s SEQ ID NO:11	TTATTGGGGGGGTTAGGGGGGGGCTT TTATTCATC	35	ND*

*K_D could not be determined by ITC for these aptamers.

FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8





FIG. 9A



SEQ ID NO:2

5'-ctgggcggtagaaccatagtgaccagccgtctac-3'

FIG. 9B




FIG. 9C




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61596774 B [0001]

专利名称(译)	针对乳酸疟原虫脱氢酶和富含组氨酸的蛋白质ii的核酸适体及其用于疟疾诊断的用途		
公开(公告)号	EP2812453A4	公开(公告)日	2015-12-16
申请号	EP2013747009	申请日	2013-02-06
[标]申请(专利权)人(译)	香港大学		
申请(专利权)人(译)	香港大学		
当前申请(专利权)人(译)	香港大学		
[标]发明人	TANNER JULIAN ALEXANDER CHEUNG YEE WAI KOTAKA MASAYO		
发明人	TANNER, JULIAN ALEXANDER CHEUNG, YEE WAI KOTAKA, MASAYO		
IPC分类号	C12Q1/68 C07H21/04 G01N33/53 G01N33/569		
CPC分类号	C12Q1/6893 G01N33/5308 G01N33/56905 G01N2333/445 Y02A50/58		
优先权	61/596774 2012-02-09 US		
其他公开文献	EP2812453A1 EP2812453B1		
外部链接	Espacenet		

摘要(译)

本发明提供了与疟原虫蛋白乳酸脱氢酶和富含组氨酸的蛋白II结合的核酸适体，及其用于诊断疟疾的用途。针对富含组氨酸的蛋白质II的适体通常可用于检测疟原虫物种的存在，而针对乳酸脱氢酶的适体可用于特异性地检测恶性疟原虫 (Plasmodium falciparum)。