



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : G01N 33/573, C12Q 1/37, G01N 33/569, 33/68, A61K 38/00, 38/55</p>	A1	<p>(11) International Publication Number: <b>WO 00/45175</b></p> <p>(43) International Publication Date: 3 August 2000 (03.08.00)</p>
<p>(21) International Application Number: PCT/EP00/00522</p> <p>(22) International Filing Date: 24 January 2000 (24.01.00)</p> <p>(30) Priority Data: 199 03 034.0 26 January 1999 (26.01.99) DE 199 20 704.6 5 May 1999 (05.05.99) DE</p> <p>(71) Applicant (for all designated States except US): TECHNOLOGIE INTEGRALE LTD. [GB/GB]; Water Street, Ramsey, Isle of Man IM8 IRD (GB).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): PAPUASHVILI, Marina, N. [RU/RU]; Millyonshikova Strasse 18/50, Moscow, 115446 (RU).</p> <p>(74) Agent: BECKER-KURIG-STRAUS; Bavariastrasse 7, D-80336 München (DE).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: URINARY TRYPSIN INHIBITOR TO DIAGNOSE AIDS</p>		
<p>(57) Abstract</p> <p>The present invention pertains to the use of urinary trypsin inhibitor for the diagnosis of the onset of AIDS and to a kit for performing such an assay. The present invention also relates to the use of urinary trypsin inhibitor for testing the efficacy of anti-HIV drugs.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav Republic of Macedonia	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece			<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>ML</b>	Mali	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MN</b>	Mongolia	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MR</b>	Mauritania	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MW</b>	Malawi	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>MX</b>	Mexico	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NE</b>	Niger	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NL</b>	Netherlands	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NO</b>	Norway	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's Republic of Korea	<b>NZ</b>	New Zealand		
<b>CM</b>	Cameroon	<b>KR</b>	Republic of Korea	<b>PL</b>	Poland		
<b>CN</b>	China	<b>KZ</b>	Kazakhstan	<b>PT</b>	Portugal		
<b>CU</b>	Cuba	<b>LC</b>	Saint Lucia	<b>RO</b>	Romania		
<b>CZ</b>	Czech Republic	<b>LI</b>	Liechtenstein	<b>RU</b>	Russian Federation		
<b>DE</b>	Germany	<b>LK</b>	Sri Lanka	<b>SD</b>	Sudan		
<b>DK</b>	Denmark	<b>LR</b>	Liberia	<b>SE</b>	Sweden		
<b>EE</b>	Estonia			<b>SG</b>	Singapore		

**URINARY TRYPSIN INHIBITOR TO DIAGNOSE AIDS**

5

The present invention pertains to the use of the urinary trypsin inhibitor for the diagnosis of the onset of AIDS. In particular the present invention relates to the use of antibodies directed against said urinary trypsin inhibitor for the diagnosis of the onset of AIDS. In another embodiment of the invention the antibodies are provided in a kit for determining the onset of AIDS.

In the early 80's patients were observed in hospitals that had developed severe immunodeficiencies. Some of them developed unusual opportunistic infections while others have been found to suffer from Kaposi's sarcoma, a hitherto rare skin tumor. An immunologic evaluation of these patients showed a marked deficiency of the cellular immune function and a selective decrease of the number of T4 (helper) cells, a subpopulation of T cells that mediate the cellular immune response. This condition with all of its accompanied symptoms was termed "Acquired Immune Deficiency Syndrome" (AIDS). The disease itself was subsequently found to be caused by a pathogenic retrovirus called human immunodeficiency virus (HIV - virus).

Soon after HIV was found to be the origin of AIDS it was observed that this virus was in particular cytopathic for T4 cells. It was therefore hypothesized that the immunological abnormalities in AIDS resulted at least in part from a progressive depletion of T4 cells due to their infection and destruction by HIV.

From the clinical point of view the "disease" AIDS is defined as the presence of a reliably diagnosed disease at least moderately indicative of cellular immune deficiency with the absence of known underlying immune deficiency and of any other reduced resistance reported to be associated with the disease, such as e.g. immunosuppressive  
5 therapy or lymphoreticular malignancy.

The disease AIDS, that is the last stage of a slowly but constantly progressing decline of the body's normal constitution, is preceded by a variety of diseases that are generically designated as "predromal AIDS". Such disorders include ARC (Aids  
10 Related Complex) which state of disease is considered to be highly predictive of the development of subsequent AIDS. ARC includes the symptoms of e.g. having fever for more than 3 months, exhibiting a loss of weight by more than 10 %, suffering from diarrhoea and showing a reduced number of T4 cells. This state of disease is known to eventually develop into the so called LAIDS (Lesser AIDS) which state is  
15 characterized by the development of oral candidiasis, herpes zoster, idiopathic thrombocytopenia and other diseases that are not lethal, do, however, indicate immune suppression.

Another predictive symptom for the prospective development of AIDS is the so called  
20 LAS (lymphadenopathy syndrome), which is identified by the presence of at least three or more lymph nodes outside of the regio inguinalis having a diameter of more than 1 cm.

Yet, another indication for the coming development of AIDS is the ADC symptoms  
25 (AIDS dementia complex) which is seen in the increasing number of detrimental influences on brain activity.

With the progress of molecular biology and cell biology techniques the determination of various antigenic determinants of the HIV virus itself has become possible. To this  
30 end, antibodies have been prepared with which the presence of the virus in the blood of

a patient can be detected. With these tests it is, however, merely possible to determine the presence or absence of an HIV infection while the imminent manifestation of the disease as such can not reliably be predicted thereby. Hence, due to the complex picture of the prodromal disease states mentioned above it is often difficult for the physician to  
5 determine whether the patient is going to develop AIDS or not within the next time.

Consequently, there exists a need in the art for a reliable and quick diagnostic tool to reliably predict the onset of the disease.

10 The development of an HIV infection in the human body is in general determined by two interacting factors. The unique property of HIV to weaken the infected individual's immune system and the immune response of the host against the invader, which develops along with the infection's progress.

15 Prior to the development of the profound immunodeficiency status the immune system still performs its task to a certain extent producing particular molecules that may be detected in the infected individual long before the various conditions described above are perceived. It was therefore considered to use such molecules for the determination of the disease.

20

Molecules, which are presently used for this purpose, are e.g.  $\beta$ -microglobulin or neopterin. The use of these compounds for the determination of the disease suffers, however, from the fact that they are not particularly specific for an HIV infection and that they have to be prepared from blood samples in a complicated manner, which  
25 impedes the procedure of diagnosis. Further, the increase in the amount of those molecules in body fluids can only be determined at a stage at which other immune system parameters and virological assays already clearly indicate the development of AIDS.

30 Therefore, the problem underlying the present invention is to provide a novel

diagnostic tool for predicting the onset of AIDS already at an early developmental stage of the disease, preferably already during predromal AIDS conditions.

The above problem has been solved by using a particular compound to be found in  
5 body fluids, the urinary trypsin inhibitor, as a diagnostic tool, and identifying its amount in said body fluids.

During the extensive experiments leading to the present invention it has been found that the amount of urinary trypsin inhibitor (UTI), which may ordinarily be present in a  
10 healthy person in the urine in an amount of up to about 3  $\mu\text{g/ml}$  is markedly increased in patients suffering from an HIV infection in whom the disease is just about to develop.

Thus, according to the present invention there is provided the use of UTI in the  
15 diagnosis of the onset of AIDS, wherein the amount of UTI in a body fluid sample of an AIDS patient is determined, which body fluid may be blood or preferably urine. If the amount exceeds a certain level, in the urine 3  $\mu\text{g/ml}$ , this may be taken as a reliable signal that this patient will develop the disease.

20 The amount of UTI in a body fluid may be determined according to methods known in the art, such as by means of antibodies, that may be polyclonal or monoclonal antibodies or an antibody containing serum. Other methods of UTI determination include measurement of the antitryptic activity or antichymotryptic activity in the sample.

25

The present invention also provides a kit for carrying out the determination of UTI-level in a body fluid sample, providing all the necessary ingredients and buffers for performing the assay.

30 In the figures,

Fig. 1 is a standard curve showing the dependency of the extinction at 492 nm on the UTI concentration

5 During the studies leading to the present invention the manifestation of many proteinase inhibitors and the dependency of their levels on tumoral processes in a body suffering from an immunodeficiency condition was examined. One of the blood plasma inhibitors is the inter- $\alpha$ -trypsin inhibitor (ITI), a protein the function of which has not yet been fully elucidated. ITI is a glycoprotein with a molecular weight of about 240  
10 kDa that is synthesized in the liver. Its concentration in blood is about 450  $\mu$ g/ml.

Immunologically affixed to ITI there has been found an inhibitor having a lower molecular weight. This specific inhibitor represents a fragment of the ITI light chain containing a large number of bisulfide links which determine its acid-stability. Due to  
15 the fact that in certain pathologic conditions this inhibitor is secreted from blood into the urine said molecule is termed urinary trypsin inhibitor (UTI).

UTI is an acidic glycoprotein with a molecular weight of about 44 kDa. About 30% of its mass is comprised by two major carbohydrate chains. The molecular weight of UTI,  
20 without carbohydrates is about 30 kDa. In terms of its tertiary structure, UTI is a two-domain molecule, even though it is comprised of only one polypeptide chain. A structural homology has been noticed between both domains and also to the main pancreatic proteinase inhibitor (MPPI), present in organs of cattle. This essentially also applies to the UTI trypsin-binding domain and to its chymotrypsin-binding domain, in  
25 the region of reactive centers.

The structural homology to MPPI allows to correlate UTI to the "Kunin" inhibitors. Due to the fact that both domains of UTI are homologous to the MPPI, this polypeptide is sometimes called "Bi-Kunin". The UTI's property of inactivating elastase and  
30 neutrophilic granulocytes G cathepsin gave rise to the hypothesis that it might exhibit

an anti-inflammatory effect.

On the other hand, already in 1974 a particular protein termed EDC-1 (mol. weight = 27 kDa) was isolated from urine of a patient suffering from acute myeloid leucosis. In most of the oncological patients examined so far the level of EDC-1 in urine was found to correspond to the clinical progress of the disease. It decreases significantly before the appearance of clinical signs of remission and increases prior to a clinical relapse.

The analysis of the EDC-1 amino acid sequence has shown that its N-terminal sequence generally corresponds to that of UTI while the C-terminal sequences differ. It was sometimes believed that EDC-1 and UTI may emerge by lysis of various peptide links in ITI.

Meanwhile, a potential mechanism for the increase of EDC-1's content in oncological patients has been proposed. At first stages of tumoral growth the tumor-associated proteolytic activity induces an enhanced formation of  $\alpha$ -1-proteinase inhibitor in the host, which is substantially inactivated by the tumor's proteinases. It is only in the second turn that the tumor-associated proteinases decompose the plasma ITI until EDC-1 is formed. As a result, the concentration of EDC-1 in the serum increases while the concentration of ITI decreases.

In urine and blood serum of patients suffering from different carcinomas, such as e.g. carcinoma of the stomach, the esophagus or of the large intestine, an increased level of UTI has been noticed, which level is in correlation with the progress of the disease. After a successful application of a chemotherapy the level of UTI was found to be decreased.

Moreover, in a healthy kidney UTI has been discovered in the cytoplasm of proximal canal epithelial cells, which suggests its secretion into the urine by those cells. UTI has not been discovered in glomerules, distal canals or vascular walls. Moreover, also

interstices did not contain UTI, while the tissue surrounding the tumor showed a level of UTI just like in a healthy kidney. It was, therefore, suggested that cancer cells obviously do not produce UTI.

5 According to the present invention the increase of the UTI level in body fluids, such like blood or, more preferably urine, may now be utilized for determining the onset of the development of diseases associated with AIDS or, more preferably predromal AIDS. In an HIV-infected patient, in whom the HIV virus is presently dormant and who does not suffer from the syndromes associated with AIDS the level of UTI in the  
10 urine is in the normal range of up to 3  $\mu\text{g/ml}$  like in normal healthy people. However, when the diseases making up the syndrome are about to emerge due to the patient's immune system being weakened to an extent such that it may no longer defeat "normal" bacteriological, viral or oncological occurrences in the body so that the syndrome is starting to emerge the level of UTI in urine markedly exceeds the normal  
15 level of up to 3  $\mu\text{g/ml}$ .

This increase of UTI-level, measurable long before the symptoms of the diseases associated with the syndrome are noticeable, may be taken as an indication of the onset of the development of AIDS. This knowledge will allow the physician to start with an  
20 appropriate treatment of the patient before the respective person will effectively fall ill.

In addition by measuring the amount of UTI during a treatment of a patient with drugs, such as anti-HIV drugs or drugs specifically directed against the particular disease the patient has acquired, e.g. candidiasis, the efficacy of said drug in an AIDS patient may  
25 be easily monitored. Consequently, when e.g. applying drugs, such as AZT or the so called "triple therapy", to a patient suffering from diseases associated with AIDS the decline of the disease may be monitored by determining the UTI level in the patient.

Moreover, the present invention also provides for the advantage that due to the  
30 possibility to monitor and HIV-infected person for the onset of the disease the amount

of drugs administered to said patient before AIDS breaks out anyway may be reduced, so that the immense costs involved in such a therapy may be saved and in addition the possibility of creating resistance in the virus will be reduced as well.

- 5 Since the assay may also be carried out on a biological material easily available, i.e. urine, the assay is easy to perform. No skilled person, such as a physician or a medical technician taking the blood is necessary.

The present invention also provides for a kit to perform the assay. The kit may have  
10 e.g. simply the form of a strip on which polyclonal or monoclonal antibodies or an antibody containing serum directed against UTI have been immobilized. Thus, a sample of a biological material is simply brought in contact with the strip and a solution containing an antibody against UTI is subsequently contacted with the strip to form a sandwich with the UTI molecule. The sandwich may then be visualized by  
15 ordinary means such as e.g. applying an antibody directed to the constant part of the antibody and being linked to a marker, such as e.g. a dye, horse raddish peroxidase etc..

It will be appreciated that the skilled person will, based on his own knowledge, design the kit according to the particular needs, such as an ELISA and others. Hence in order  
20 to provide for an accurate reading of the amount of UTI present in a sample the person skilled in the art will consider the use of a corresponding apparatus, such as e.g. a photometer. In addition it will also be appreciated that techniques for raising polyclonal or monoclonal antibodies against UTI are well within the ordinary skill.

- 25 Depending on the sort of kit provided the affected person may well perform the assay at home for a general indication with e.g. dyes indicating a particular UTI content. In addition the UTI level may also be accurately determined in a laboratory with e.g. using a photometer.

## Examples

### Example 1

#### Determination of UTI via antibodies

5

To detect the UTI, the following procedure was used: Three monoclonal antibodies, M2, B6, P1 were raised against UTI in a manner known per se and examined in the presence of the trypsin binding domain for their binding capacity. The M2 antibody was shown to have the highest affinity for this domain. On the basis of the mab M2, a competitive ELISA of UTI concentration in urine has been set up.

10

The methodology that was used to detect UTI with a view of determining its value in predicting the progress of HIV infection at different stages of the disease, has revealed the following:

15

## Results

### Groups of Patients

Seventeen HIV-1-infected patients and 30 healthy volunteers were examined; 4 patients had no clinical symptoms; 8 patients had different opportunistic infections, and 5 patients had already developed AIDS.

20

### Materials and Methods

UTI ELISA (mab to UTI: secreting hybridoma were derived from a fusion of non-secreting mouse myeloma cells (clone P3 O1) with spleen cells of BALB/c mice that have been immunized with human UTI) level in urine, residual nitrogen and creatinin concentrations and a number of immunological parameter's (CD4, CD8, CD16, B-lymphocytes, HLA-DR), p24 concentration were tested.

25

The preparation of urinary protein concentrates:

1. Collection of urine (preferably, the morning portion); then treatment with acetone (5 ml of urine + 10 ml of acetone cooled down to +4°C);
2. The acetone precipitation of urinary proteins may be kept at +4°C for one month.
- 5 3. Spinning-down of protein precipitates (1.000 g, 10 min.).
4. Preparation of suspension of the precipitates obtained (precipitate +2,5 ml 0,05 M buffer, pH 7,8 the precipitate is triturated with a glass rod) the concentration and depigmentation of the urine.
5. The undissolved precipitate is centrifuged.
- 10 6. The supernatant thus obtained is used for the determination of the UTI.

ELISA-Test

1. The UTI solution is adsorbed on a plate having 96-wells for one night at a temperature of 4°C;
- 15 2. The plate is rinsed with water and knocked out;
3. The filling of vacant binding-sites with albumin (in each well +200µl of 1% albumin solution in 0,05 M phosphate buffer, pH 7,4 for one-hour incubation;
4. The plate is rinsed with water and knocked out.
5. The titration into the plate's well of: (a) urine specimen (see point 6 above; (b) the  
20 standard UTI solution for the calibration curve; and (c) the control sample with the known UTI content; thereto the solution of the UTI monoclonal antibodies conjugate with horse-radish peroxidase is added and one-hour incubation; the binding of UTI from the sample and that adsorbed on the plate with the antibodies.
6. The plate is rinsed with the buffer (with 0,1% albumin and Tween-20), and with  
25 water, and knocked out.
7. Development: addition of the substrate (ortho-phenylene-diamine in the citrate-phosphate buffer); incubating for 10 minutes.
8. The stopping of the reaction: +10% sulphuric acid.
9. The measurement of optical density with an ELISA-reader at 492 nm.

10. Drawing a calibration curve with known amounts of UTI; the standard curve was constructed using 2-fold dilution of UTI in the wells of the microtiter plate; the linear segment of the standard curve is to cover the working concentration range (see Fig. 1);
- 5 11. Determination of the UTI concentration of the specimen therefrom.

In order to determine the UTI concentration in blood, the following equipment may be used:

- 10 1. A 80-channel pipette, 40 x 200  $\mu$ l.  
2. An Eppendorf centrifuge for test-tubes (microfuge).  
3. An ELISA-reader with a filter for 492 nm for plates.  
4. Plates for the agents.  
5. The measuring of the UTI activity, in addition to concentration, would require a  
15 photometer with a 405-nm filter.  
6. A washer.

### Example 2

#### Quantitative ATA determination

20

The urine sample was obtained as described in example 1.

#### Reagents used:

- 1) Trypsin concentrate: 1 mg trypsin was dissolved in 1 ml of 0.0025 M HCl with 0.1  
25 M  $\text{CaCl}_2$ ; the solution may be stored at 4 °C up to 10 days;  
2) working trypsin solution; dilute trypsin concentrate 50 times by 0.0025 M HCl with  
0.1 M  $\text{CaCl}_2$ ;  
3) 0.2 M TRIS-HCl pH 7.8 - 8.0;  
4) BAPNA concentrate (p-nitroanilide N-benzoyl-DL-arginine); dissolve 11.4 mg  
30 BAPNA in 6 ml dehydrated DMSO; store in darkness;

- 5) BAPNA working solution; dilute BAPNA concentrate with distilled water in a ratio of 1 : 2;

ATA measurements in the urine sample

- 5 1) add 40  $\mu$ l of trypsin working solution and 2 - 80  $\mu$ l of the sample to be tested into a spectrophotometer cuvette;
- 2) add 0.2 M TRIS-HCl, pH 7,8 - 8,0 for a total volume up to 160  $\mu$ l;
- 3) mix solution and incubate for 5 min at 37 °C;
- 4) add 40  $\mu$ l of BAPNA working solution;
- 10 5) mix solution and measure the initial optical density ( $A_1$ ) at 405 or 410 nm in the spectrophotometer;
- 6) incubate the solution for 30 min at 37 °C;
- 7) mix solution and measure the final optical density ( $A_2$ ) at 405 or 410 nm in the spectrophotometer.

15

Processing of the results obtained

The quantity of ATA is calculated with the following formulae:

20

$$ATA = \varepsilon (B_2 - B_1 - A_2 + A_1) / (t V),$$

25

wherein  $A_1$  und  $A_2$  are the initial and final optical density in the presence of a tested sample,  $B_1$  and  $B_2$  are the initial and final optical density in the absence of the sample. V is the volume of the urine sample added, t is the reaction time and  $\varepsilon$  is a tabulated extinction coefficient of p-nitroaniline in dependence of the wavelength used and the total solution volume.

The most accurate results are obtained if

30

$$0.4 (B_2 - B_1) < A_2 - A_1 < 0.6 (B_2 - B_1)$$

Claims

1. Use of urinary trypsin inhibitor for the diagnosis of the onset of AIDS.
- 5 2. The use according to claim 1, wherein the amount of UTI is determined in urine or blood samples.
3. The use according to any of the claims 1 or 2, wherein the amount of UTI is  
10 determined by means of polyclonal or monoclonal antibodies or antibody containing sera.
4. The use according to any of the claims 1 or 2, wherein the amount of UTI is determined via the measurement of the antitryptic activity.
- 15 5. The use according to any of the claims 1 or 2, wherein the amount of UTI is determined via the measurement of the antichymotryptic measurement.
6. The use according to claim 3, wherein the antibodies are labeled with a marker.
- 20 7. The use according to claim 6, wherein the marker is a dye, a fluorescent or a radioactive marker.
8. The use according to claim 7, wherein the antibodies are applied in an ELISA  
25 assay.
9. A kit for the diagnosis of urinary trypsin inhibitor as an indicator for the onset of AIDS comprising at least one antibody against UTI.
- 30 10. The kit according to claim 7, which includes a carrier onto which a polyclonal antibody has been immobilized.
11. The kit according to claim 10, which additionally includes a monoclonal antibody and an anti-antibody to which a marker has been fixed.

12. The use for UTI for determining the efficacy of an anti-HIV drug.




FIG. 1

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 00/00522

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC 7 G01N33/573 C12Q1/37 G01N33/569 G01N33/68 A61K38/00  
 A61K38/55

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**  
 Minimum documentation searched (classification system followed by classification symbols)  
 IPC 7 G01N C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 371 706 A (NIHON CHEMICAL RESEARCH KK) 6 June 1990 (1990-06-06) examples 1-3	1-12
X	PATENT ABSTRACTS OF JAPAN vol. 18, no. 111 (C-1170), 23 February 1994 (1994-02-23) & JP 05 304983 A (NISSIN FOOD PRODUCTION COMPANY LTD) abstract	9-11

Further documents are listed in the continuation of box C.       Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family
---	---

Date of the actual completion of the international search  <b>15 May 2000</b>	Date of mailing of the international search report  <b>24/05/2000</b>
---	---

Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer   <b>Van Bohemen, C</b>
--	---

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/00522

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 371706      A	06-06-1990	JP 2145527 A JP 2804979 B DE 68907569 D DE 68907569 T US 5504065 A	05-06-1990 30-09-1998 19-08-1993 28-10-1993 02-04-1996
JP 05304983    A	19-11-1993	NONE	

专利名称(译)	尿胰蛋白酶抑制剂诊断艾滋病		
公开(公告)号	<a href="#">EP1147420A1</a>	公开(公告)日	2001-10-24
申请号	EP2000902627	申请日	2000-01-24
申请(专利权)人(译)	TECHNOLOGIE INTEGRALE LTD.		
当前申请(专利权)人(译)	TECHNOLOGIE INTEGRALE LTD.		
[标]发明人	PAPUASHVILI MARINA N		
发明人	PAPUASHVILI, MARINA, N.		
IPC分类号	G01N33/53 A61K38/00 A61K38/55 C12Q1/00 C12Q1/37 C12Q1/70 G01N33/543 G01N33/569 G01N33/573 G01N33/577 G01N33/68		
CPC分类号	G01N33/56988 G01N2333/16 G01N2333/811 Y10S435/975		
优先权	19903034 1999-01-26 DE 19920704 1999-05-05 DE		
其他公开文献	EP1147420B1		
外部链接	<a href="#">Espacenet</a>		

#### 摘要(译)

本发明涉及尿胰蛋白酶抑制剂用于诊断AIDS发作的用途和用于进行这种测定的试剂盒。本发明还涉及尿胰蛋白酶抑制剂在测试抗HIV药物功效中的用途。