



US 20190324041A1

(19) **United States**

(12) **Patent Application Publication**  
**Wang et al.**

(10) **Pub. No.: US 2019/0324041 A1**

(43) **Pub. Date: Oct. 24, 2019**

(54) **IMMUNO-BASED RETARGETED  
ENDOPEPTIDASE ACTIVITY ASSAYS**

(71) Applicant: **Allergan, Inc.**, Irvine, CA (US)

(72) Inventors: **Joanne Wang**, Irvine, CA (US); **Hong  
Zhu**, San Diego, CA (US); **Dianne D.  
Hodges**, Tustin, CA (US); **Ester  
Fernandez-Salas**, San Mateo, CA (US)

(21) Appl. No.: **16/504,870**

(22) Filed: **Jul. 8, 2019**

**Related U.S. Application Data**

(63) Continuation of application No. 15/593,727, filed on May 12, 2017, now Pat. No. 10,345,306, which is a continuation of application No. 14/695,938, filed on Apr. 24, 2015, now Pat. No. 9,651,553, which is a continuation of application No. 13/801,711, filed on Mar. 13, 2013, now abandoned, which is a continuation of application No. 12/723,595, filed on Mar. 12, 2010, now Pat. No. 8,455,203.

(60) Provisional application No. 61/160,217, filed on Mar. 13, 2009.

**Publication Classification**

(51) **Int. Cl.**

**G01N 33/573** (2006.01)

**G01N 33/50** (2006.01)

**C12N 9/50** (2006.01)

**C07K 16/18** (2006.01)

**G01N 33/53** (2006.01)

**C12Q 1/37** (2006.01)

(52) **U.S. Cl.**

CPC . **G01N 33/573** (2013.01); **G01N 2333/96425**

(2013.01); **G01N 33/5035** (2013.01); **C12N**

**9/50** (2013.01); **C07K 16/18** (2013.01); **C12Y**

**304/24069** (2013.01); **G01N 33/53** (2013.01);

**C12Q 1/37** (2013.01); **G01N 2333/33**

(2013.01); **C07K 2317/34** (2013.01); **C07K**

**2317/92** (2013.01); **C07K 2317/56** (2013.01);

**G01N 2333/47** (2013.01); **G01N 2333/952**

(2013.01); **G01N 33/5014** (2013.01)

(57)

**ABSTRACT**

The present specification discloses a retargeted endopeptidase pharmaceutical wherein the activity has been determined by the methods disclosed.

**Specification includes a Sequence Listing.**

FIG. 1A.





FIG. 1B.



**FIG. 2.**




FIG. 3.




FIG. 4.




FIG. 5.




FIG. 6.




FIG. 7A.





FIG. 7B.



**IMMUNO-BASED RETARGETED  
ENDOPEPTIDASE ACTIVITY ASSAYS****CROSS REFERENCE TO RELATED  
APPLICATIONS**

[0001] This application is a continuation of U.S. patent application Ser. No. 15/593,727, filed May 12, 2017, now U.S. Pat. No. 10,345,306, which is a continuation of U.S. patent application Ser. No. 14/695,938, filed Apr. 24, 2015, now U.S. Pat. No. 9,651,553, which is a continuation to U.S. patent application Ser. No. 13/801,711, filed Mar. 13, 2013, which is a continuation of U.S. patent application Ser. No. 12/723,595, filed Mar. 12, 2010, now U.S. Pat. No. 8,455,203, which claims priority to U.S. Provisional Patent Application Ser. No. 61/160,217 filed Mar. 13, 2009, each of which is incorporated entirely by reference.

[0002] The sequences disclosed in the present specification are contained in the Sequence Listing submitted with the present specification which is hereby incorporated by reference in its entirety.

[0003] The ability of Clostridial toxins, such as, e.g., Botulinum neurotoxins (BoNTs), BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F and BoNT/G, and Tetanus neurotoxin (TeNT), to inhibit neuronal transmission are being exploited in a wide variety of therapeutic and cosmetic applications, see e.g., William J. Lipham, *Cosmetic and Clinical Applications of Botulinum Toxin* (Slack, Inc., 2004). Clostridial toxins commercially available as pharmaceutical compositions include, BoNT/A preparations, such as, e.g., BOTOX® (Allergan, Inc., Irvine, Calif.), DYS-PORT®/RELOXIN®, (Ipsen Ltd., Slough, England), PURTOX® (Mentor Corp., Santa Barbara, Calif.), XEOMIN® (Merz Pharmaceuticals, GmbH, Frankfurt, Germany), NEURONOX® (Medy-Tox, Inc., Ochang-myeon, South Korea), BTX-A (Biogen-tech Ltd., University, Yantai, Shandong, China); and BoNT/B preparations, such as, e.g., MYOBLOMNEUROBLOC® (Solstice Neurosciences, Inc., South San Francisco, Calif.). As an example, BOTOX® is currently approved in one or more countries for the following indications: achalasia, adult spasticity, anal fissure, back pain, blepharospasm, bruxism, cervical dystonia, essential tremor, glabellar lines or hyperkinetic facial lines, headache, hemifacial spasm, hyperactivity of bladder, hyperhidrosis, juvenile cerebral palsy, multiple sclerosis, myoclonic disorders, nasal labial lines, spasmodic dysphonia, strabismus and VII nerve disorder.

[0004] A Clostridial toxin treatment inhibits neurotransmitters and neuropeptide release by disrupting the exocytotic process used to secrete the neurotransmitters and neuropeptides into the synaptic cleft. There is a great desire by the pharmaceutical industry to expand the use of Clostridial toxin therapies beyond its current myo-relaxant applications to treat sensory nerve-based ailment, such as, e.g., various kinds of chronic pain, neurogenic inflammation and urogenital disorders, as well as other disorders, such as, e.g., pancreatitis. One approach that is currently being exploited to expand Clostridial toxin-based therapies involves modifying a Clostridial toxin so that the modified toxin has an altered cell targeting capability for a neuronal or non-neuronal cell of interest. Called either re-targeted endopeptidase or Targeted Vesicular Exocytosis Modulator Proteins (TVEMPs), these molecules achieve their exocytosis inhibitory effects by using a target receptor present on the neuronal or non-neuronal target cell of interest. This

re-targeted capability is achieved by replacing a naturally-occurring binding domain of a Clostridial toxin with a targeting domain showing a selective binding activity for a non-Clostridial toxin receptor present in a neuronal or non-neuronal target cell of interest. Such modifications to a binding domain result in a molecule that is able to selectively bind to a non-Clostridial toxin receptor present on the target cell. A re-targeted endopeptidase can bind to a target receptor, translocate into the cytoplasm, and exert its proteolytic effect on the SNARE complex of the neuronal or non-neuronal target cell of interest.

[0005] One group of re-targeted endopeptidase comprises molecules having an opioid targeting domain. These opioid re-targeted endopeptidases comprise an opioid targeting domain, a Clostridial toxin translocation domain, and a Clostridial toxin enzymatic domain. Non-limiting examples of opioid re-targeted endopeptidase, or opioid-TVEMPs, are described in, e.g., Keith A. Foster et al., *Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions*, U.S. Pat. No. 5,989,545; J. Oliver Dolly et al., *Activatable Recombinant Neurotoxins*, U.S. Pat. No. 7,132,259; Stephan Donovan, *Clostridial Toxin Derivatives and Methods For Treating Pain*, U.S. Pat. No. 7,244,437; Stephan Donovan, *Clostridial Toxin Derivatives and Methods For Treating Pain*, U.S. Pat. No. 7,413,742; Stephan Donovan, *Clostridial Toxin Derivatives and Methods For Treating Pain*, U.S. Pat. No. 7,415,338; Lance E. Steward et al., *Multivalent Clostridial Toxin Derivatives and Methods of Their Use*, U.S. Pat. No. 7,514,088; Keith A. Foster, *Fusion Proteins*, U.S. Patent Publication 2008/0064092; Keith A. Foster, *Fusion Proteins*, U.S. Patent Publication 2009/0035822; Lance E. Steward et al., *Multivalent Clostridial Toxin Derivatives and Methods of Their Use*, U.S. Patent Publication 2009/0048431; Keith A. Foster, *Non-Cytotoxic Protein Conjugates*, U.S. Patent Publication 2009/0162341; Keith A. Foster et al., *Re-targeted Toxin Conjugates*, International Patent Publication WO 2005/023309; and Lance E. Steward, *Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Capabilities for Non-Clostridial Toxin Target Cells*, International Patent Application WO 2008/008805; each of which is hereby incorporated by reference in its entirety.

[0006] One general difference between re-targeted endopeptidases and Clostridial toxins is that because re-targeted endopeptidases typically do not target motor neurons, the lethality associated with over-dosing a mammal with a re-targeted endopeptidase is greatly minimized, if not avoided altogether. For example, opioid re-targeted endopeptidases can be administered at 10,000 times the therapeutically effective dose before evidence of lethality is observed, and this lethality is due to the passive diffusion of the molecule and not via the intoxication process. Thus, for all practical purposes re-targeted endopeptidases are non-lethal molecules. Although this non-lethal property is of great therapeutic benefit, a manufacturing problem arises because the standard activity assay used to manufacture Clostridial toxin-based biologics is a mouse LD<sub>50</sub> bioassay, a lethality test. S. S. Aron et al., *JAMA* 285: 1059-1070 (2001). Currently a mouse LD<sub>50</sub> bioassay is used by all pharmaceutical manufacturers to express the potency of their Clostridial toxin preparations. In fact, the activity units for Clostridial toxins are mouse LD<sub>50</sub> units. However, because re-targeted endopeptidases are essentially non-lethal, a mouse LD<sub>50</sub> bioassay cannot be used to assess the

potency of these molecules. Thus, a simple, reliable, validated, and governmental agency acceptable activity assay that can evaluate the integrity of all the steps necessary in re-targeted endopeptidase uptake would be of significant value. The present specification provides novel compositions, cells, and methods for assaying the activity of re-targeted endopeptidases useful for various industries, such as, e.g., the pharmaceutical and food industries, and provides related advantages as well. Such compositions, cells, and methods do not use live animals or tissues taken from live animals, but can evaluate all the steps necessary for re-targeted endopeptidase action.

#### DETAILED DESCRIPTION OF THE DRAWINGS

**[0007]** FIGS. 1A and 1B show a schematic of the current paradigm of neurotransmitter release and Clostridial toxin intoxication in a central and peripheral neuron. FIG. 1A shows a schematic for the neurotransmitter release mechanism of a central and peripheral neuron. The release process can be described as comprising two steps: 1) vesicle docking, where the vesicle-bound SNARE protein of a vesicle containing neurotransmitter molecules associates with the membrane-bound SNARE proteins located at the plasma membrane; and 2) neurotransmitter release, where the vesicle fuses with the plasma membrane and the neurotransmitter molecules are exocytosed. FIG. 1B shows a schematic of the intoxication mechanism for tetanus and botulinum toxin activity in a central and peripheral neuron. This intoxication process can be described as comprising four steps: 1) receptor binding, where Clostridial toxin binds to a Clostridial receptor complex and initiates the intoxication process; 2) complex internalization, where after toxin binding, a vesicle containing a toxin/receptor system complex is endocytosed into the cell; 3) light chain translocation, where multiple events are thought to occur, including changes in the internal pH of the vesicle, formation of a channel pore comprising the HN domain of Clostridial toxin heavy chain, separation of the Clostridial toxin light chain from the heavy chain, and release of the light chain and 4) enzymatic target modification, where the light chain of Clostridial toxin proteolytically cleaves its target SNARE substrates, such as, e.g., SNAP-25, VAMP or Syntaxin, thereby preventing vesicle docking and neurotransmitter release.

**[0008]** FIG. 2 shows a full dose response to the re-targeted endopeptidase Noc/A in the ORL-1Clone #6 clonal cell line overexpressing ORL-1. Specific uptake of Noc/A can be observed in the ORL-1Clone #6 clonal cell line overexpressing ORL-1. The treatment with Noc/A (LHN/A plus binding ligand nociceptin variant) and LHN/A (LC/A and HN without any binding domain) performed on ORL-1 stable cell line clone #6 in the ECL ELISA for cleaved SNAP-25197 demonstrated that uptake of Noc/A is specific in this clonal cell line. The clonal cell line also show great sensitivity for Noc/A with an  $EC_{50}$  of 1.2 nM.

**[0009]** FIG. 3 shows a full dose response to Noc/A in the SK-N-DZ single-cell derived clones #3 and #22. Specific uptake of Noc/A on SK-N-DZ clones #3 and #22 when compared to LHN/A (n=4 independent experiments run). Cells were plated on poly-D-lysine 96-well plates in RPMI SFM+N2+B27+NGF. Treatment with compounds was for 22 hours. ECL ELISA for cleaved SNAP-25197 demonstrated that uptake of Noc/A is specific in this clonal cell

lines. The clonal cell lines also show great sensitivity for Noc/A with an  $EC_{50}$  of 0.3 nM for clone #3 and an  $EC_{50}$  of 0.9 nM for clone #22.

**[0010]** FIG. 4 shows an ECL sandwich ELISA assay results from ORL1 ND7 clones 1C11, 4B7, and 4C9 treated with re-targeted endopeptidase Noc/A. Parental ND7 and ORL1 ND7 clones were treated for 24 hours with Noc/A followed by two days of incubation. Parental ND7  $EC_{50}$  could not be calculated since it only reached approximately 50% SNAP-25197 cleavage. Clones 4B7 and 1C11 reach more than 80% SNAP-25197 cleavage.  $EC_{50}$  values were calculated to be  $5.7\pm 0.5$ ,  $6.7\pm 1$ , and  $8.6\pm 2$  nM respectively.

**[0011]** FIG. 5 shows an anti-nociceptin polyclonal antibodies can block re-targeted endopeptidase Noc/A uptake in SK-N-DZ clone #3, clone #22, and AGN P33 ORL-1 clone #6 cell lines. Cells were plated on poly-D-lysine 96-well plates in RPMI SFM+N2+B27+NGF and treated for 22 hours in serum-free media containing with the Anti-nociceptin polyclonal antibodies at different dilutions (0-3  $\mu$ g/mL) in 1 nM Noc/A.

**[0012]** FIG. 6 shows cells from SiMa clone AF4 and the established cell line PC-12 were treated with the re-targeted endopeptidase Dyn/A at concentrations from 0.017 nM to 1  $\mu$ M as depicted in the Western blot image. Dose-dependent uptake could be observed for both cell lines.

**[0013]** FIGS. 7A and 7B show normalized BIAcore SPR curves of 7.8 nM of the antibodies 2E2A6, 1D3B8, 3C1A5 and 2C9B10 and commercial MC-6050 and MC-6053. FIG. 7A shows the normalized data for the on-rate of each antibody. FIG. 7B shows the normalized data for the off-rate of each antibody.

#### DETAILED DESCRIPTION

**[0014]** The present specification provides novel assays for determining the presence or absence of an active re-targeted endopeptidase in a sample and for determining the activity/potency of a re-targeted endopeptidase. The novel cell-based assays disclosed in the present specification rely on cells, reagents and detection methods that enable the assay to detect nanomolar quantities of a re-targeted endopeptidase in a sample. The cell-based assays disclosed in the present specification serve to analyze multiple functions a re-targeted endopeptidase, namely, re-targeted endopeptidase binding to a cell surface receptor, internalization of the endopeptidase-receptor complex, enzymatic domain translocation into the cytoplasm, enzymatic domain cleavage of substrate. As discussed further below, the novel methods and compositions can be used to analyze crude and bulk samples as well as highly purified di-chain re-targeted endopeptidases and formulated re-targeted endopeptidase products and further are amenable to automated high throughput assay formats.

**[0015]** Thus, one aspect disclosed in the present specification provides immune response inducing compositions for producing  $\alpha$ -SNAP-25 antibodies that can selectively bind to an epitope comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. Immune response inducing compositions can comprise an adjuvant and an immune response inducing composition including a SNAP-25 antigen, a carrier linked to a SNAP-25 antigen, or a carrier linked to a flexible spacer linked to a SNAP-25 antigen, where the flexible linker intervenes between the SNAP-25 antigen and the carrier. It is envisioned that any and all SNAP-25

antigens that triggers an immune response that produce a  $\alpha$ -SNAP-25 antibody that can selectively bind to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be useful as a SNAP-25 antigen, including, without limitation, a SNAP-25 antigen derived from a naturally occurring SNAP-25, a SNAP-25 antigen derived from a non-naturally occurring SNAP-25, and a SNAP-25 antigen comprising an immunoreactive fragment of the SNAP-25, the SNAP-25 from a naturally occurring SNAP-25 or a non-naturally occurring SNAP-25. SNAP-25 antigens useful for producing  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond include, without limitation, SNAP-25 antigens comprising a SNAP-25 peptide having a carboxylated C-terminal glutamine linked to a carrier peptide, including, without limitation SEQ ID NO:38. Other immune response inducing compositions useful for making  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond include, without limitation, an immune response inducing composition comprising a carrier linked to a flexible linker linked to a SNAP-25 antigen a carboxylated C-terminal glutamine, wherein the flexible linker intervenes between the SNAP-25 antigen and the carrier. It is envisioned that any and all adjuvants can be useful in such an immune response inducing composition, including, without limitation, polyethylene glycol (PEG), monomethoxypolyethylene glycol (mPEG), polyvinyl alcohol (PVA), complete and incomplete Freund's adjuvant.

**[0016]** Another aspect disclosed in the present specification provides methods of producing an  $\alpha$ -SNAP-25 antibody that can selectively bind to an epitope comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. Aspects of this method comprise the steps of (a) administering to an animal a SNAP-25 immune response inducing composition disclosed in the present specification; (b) collecting from the animal a sample containing an  $\alpha$ -SNAP-25 antibody or  $\alpha$ -SNAP-25 antibody-producing cell; and (c) isolating the  $\alpha$ -SNAP-25 antibody from the sample. The methods disclosed are useful for making either  $\alpha$ -SNAP-25 monoclonal antibodies that can selectively bind to an epitope comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond or  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to an epitope comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

**[0017]** Still another aspect disclosed in the present specification provides  $\alpha$ -SNAP-25 antibodies that selectively bind to an epitope comprising a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. Such  $\alpha$ -SNAP-25 antibodies include both naturally-occurring and non-naturally-occurring antibodies, as well as, monoclonal  $\alpha$ -SNAP-25 antibodies or polyclonal  $\alpha$ -SNAP-25 antibodies. Monoclonal  $\alpha$ -SNAP-25 antibodies useful as  $\alpha$ -SNAP-25 antibodies that selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, include, without limitation, the monoclonal  $\alpha$ -SNAP-25 antibodies produced from hybridoma cell lines 103B8, 2C9B10, 2E2A6, 3C1A5 and 3C3E2.

**[0018]** Yet another aspect disclosed in the present specification provides immuno-based methods of detecting re-targeted endopeptidase activity. Aspects of this method comprise the steps of (a) treating a cell from an established cell line with a sample comprising a re-targeted endopeptidase, wherein the cell from an established cell line is susceptible to re-targeted endopeptidase activity by the re-targeted endopeptidase; (b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; (c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody disclosed in the present specification; and (d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; wherein detection by the antibody-antigen complex is indicative of re-targeted endopeptidase activity. The  $\alpha$ -SNAP-25 antibody of step (c) can optionally be linked to a solid phase support.

**[0019]** Yet another aspect disclosed in the present specification provides immuno-based methods of detecting opioid-TVEMP activity. Aspects of this method comprise the steps of (a) treating a cell from an established cell line with a sample comprising a re-targeted endopeptidase, wherein the cell from an established cell line can uptake a re-targeted endopeptidase; (b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; (c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody disclosed in the present specification; and (d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; wherein detection by the antibody-antigen complex is indicative of re-targeted endopeptidase activity. The  $\alpha$ -SNAP-25 antibody of step (c) can optionally be linked to a solid phase support.

**[0020]** A further aspect disclosed in the present specification provides methods of determining re-targeted endopeptidase immunoresistance in a mammal. Aspects of this method comprise the steps of (a) adding a re-targeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; (b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line is susceptible to re-targeted endopeptidase activity; (c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; (d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody disclosed in the present specification; (e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; (f) repeating steps a-e with a negative control sample instead of a test sample; and (g) comparing the amount of antibody-antigen complex detected in step (e) to the amount of antibody-antigen complex detected in step (f), wherein detection of a lower amount of antibody-antigen complex detected in step (e) relative to the amount of antibody-antigen complex detected in step (f) is indicative of the presence of  $\alpha$ -re-targeted

endopeptidase neutralizing antibodies. The  $\alpha$ -SNAP-25 antibody of step (d) can optionally be linked to a solid phase support. The control sample in step (f) can also include a positive control sample, in addition to the negative control sample.

**[0021]** Clostridia toxins produced by *Clostridium botulinum*, *Clostridium tetani*, *Clostridium baratii* and *Clostridium butyricum* are the most widely used in therapeutic and cosmetic treatments of humans and other mammals. Strains of *C. botulinum* produce seven antigenically-distinct serotypes of botulinum toxins (BoNTs), which have been identified by investigating botulism outbreaks in man (BoNT/A, BoNT/B, BoNT/E and BoNT/F), animals (BoNT/C1 and BoNT/D), or isolated from soil (BoNT/G). While all seven botulinum toxin serotypes have similar structure and biological properties, each also displays heterogeneous characteristics, such as, e.g., different pharmacological properties. In contrast, tetanus toxin (TeNT) is produced by a uniform group of *C. tetani*. Two other species of Clostridia, *C. baratii* and *C. butyricum*, also produce toxins similar to BoNT/F and BoNT/E, respectively.

**[0022]** Clostridial toxins are each translated as a single chain polypeptide of approximately 150 kDa that is subsequently cleaved by proteolytic scission within a disulfide loop by a naturally-occurring protease, such as, e.g., an endogenous Clostridial toxin protease or a naturally-occurring protease produced in the environment. This posttranslational processing yields a di-chain molecule comprising an approximately 50 kDa light chain (LC) and an approximately 100 kDa heavy chain (HC) held together by a single disulfide bond and noncovalent interactions. Each mature di-chain molecule comprises three functionally distinct domains: 1) an enzymatic domain located in the LC that includes a metalloprotease region containing a zinc-dependent endopeptidase activity which specifically targets core components of the neurotransmitter release apparatus; 2) a translocation domain contained within the amino-terminal half of the HC ( $H_N$ ) that facilitates release of the LC from intracellular vesicles into the cytoplasm of the target cell; and 3) a binding domain found within the carboxyl-terminal half of the HC ( $H_C$ ) that determines the binding activity and binding specificity of the toxin to the receptor complex located at the surface of the target cell.

**[0023]** The binding, translocation and enzymatic activity of these three functional domains are all necessary for toxicity. While all details of this process are not yet precisely known, the overall cellular intoxication mechanism whereby Clostridial toxins enter a neuron and inhibit neurotransmitter release is similar, regardless of serotype or subtype. Although the applicants have no wish to be limited by the following description, the intoxication mechanism can be described as comprising at least four steps: 1) receptor binding, 2) complex internalization, 3) light chain translocation, and 4) enzymatic target modification (FIG. 1). The process is initiated when the HC domain of a Clostridial toxin binds to a toxin-specific receptor system located on the plasma membrane surface of a target cell. The binding specificity of a receptor complex is thought to be achieved, in part, by specific combinations of gangliosides and protein receptors that appear to distinctly comprise each Clostridial toxin receptor complex. Once bound, the toxin/receptor complexes are internalized by endocytosis and the internalized vesicles are sorted to specific intracellular routes. The translocation step appears to be triggered by the acidification

of the vesicle compartment. This process seems to initiate important pH-dependent structural rearrangements that increase hydrophobicity, promote pore formation, and facilitate separation of the heavy and light chains of the toxin. Once separated, the light chain endopeptidase of the toxin is released from the intracellular vesicle into the cytosol where it appears to specifically target core components of the neurotransmitter release apparatus. These core proteins, vesicle-associated membrane protein (VAMP)/synaptobrevin, synaptosomal-associated protein of 25 kDa (SNAP-25) and Syntaxin, are necessary for synaptic vesicle docking and fusion at the nerve terminal and constitute members of the soluble N-ethylmaleimide-sensitive factor-attachment protein-receptor (SNARE) family. BoNT/A and BoNT/E cleave SNAP-25 in the carboxyl terminal region, releasing a nine or twenty six amino acid fragment, respectively, and BoNT/C1 also cleaves SNAP-25 near the carboxyl terminus releasing an eight amino acid fragment. The botulinum serotypes BoNT/B, BoNT/D, BoNT/F and BoNT/G, and tetanus toxin, act on the conserved central portion of VAMP, and release the amino terminal portion of VAMP into the cytosol. BoNT/C1 cleaves syntaxin at a single site near the cytosolic membrane surface. The selective proteolysis of synaptic SNAREs accounts for the block of neurotransmitter release caused by Clostridial toxins in vivo. The SNARE protein targets of Clostridial toxins are common to exocytosis in a variety of non-neuronal types; in these cells, as in neurons, light chain peptidase activity inhibits exocytosis, see, e.g., Yann Humeau et al., *How Botulinum and Tetanus Neurotoxins Block Neurotransmitter Release*, 82(5) *Biochimie*. 427-446 (2000); Kathryn Turton et al., *Botulinum and Tetanus Neurotoxins: Structure, Function and Therapeutic Utility*, 27(11) *Trends Biochem. Sci.* 552-558. (2002); Giovanna Lalli et al., *The Journey of Tetanus and Botulinum Neurotoxins in Neurons*, 11(9) *Trends Microbiol.* 431-437, (2003).

**[0024]** Re-targeted endopeptidases generally substitute the naturally-occurring di-chain loop protease cleavage site with an exogenous protease cleavage site. See e.g., Dolly, J.O. et al., *Activatable Clostridial Toxins*, U.S. Pat. No. 7,419,676, which is hereby incorporated by reference. Although re-targeted endopeptidases vary in their overall molecular weight because of the size of the targeting moiety, the activation process and its reliance on cleavage at the exogenous cleavage site to produce a di-chain molecule is essentially the same as that for Clostridial toxins. See e.g., Steward, L. E. et al., *Activatable Clostridial Toxins*, U.S. Patent Publication 2009/0005313; Steward, L. E. et al., *Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Non-Clostridial Toxin Target Cells*, U.S. patent application Ser. No. 11/776,075; Steward, L. E. et al., *Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity for Clostridial Toxin Target Cells*, U.S. Patent Publication 2008/0241881, each of which is hereby incorporated by reference.

**[0025]** Aspects of the present disclosure comprise, in part, an immune response inducing composition for producing  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. As used herein, the term "immune response inducing composition" refers to a composition comprising a SNAP-25 antigen which, when administered to an animal, stimulates an immune response

against the SNAP-25 antigen, thereby producing  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. The term “immune response” refers to any response by the immune system of an animal to an immune response inducing composition. Exemplary immune responses include, but are not limited to, cellular as well as local and systemic humoral immunity, such as, e.g., CTL responses, including antigen-specific induction of CD8+ CTLs, helper T-cell responses, including T-cell proliferative responses and cytokine release, and B-cell responses including, e.g., an antibody producing response. The term “inducing an immune response” refers to administration of an immune response inducing composition or a polynucleotide encoding the immune response inducing composition, where an immune response is affected, i.e., stimulated, initiated or induced.

**[0026]** An SNAP-25 immune response inducing composition comprises a SNAP-25 antigen. As used herein, the term “antigen” refers to a molecule that elicits an immune response and includes, without limitation, peptides, polysaccharides and conjugates of lipids, such as, e.g., lipoproteins and glycolipids. As used herein, the term “SNAP-25 antigen” refers to any antigen which has a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond that can elicit an immune response. A SNAP-25 antigen used in an immune response inducing composition must be large enough to be substantially unique in sequence, thus reducing the possibility of producing antibodies that are cross reactive against antigens other than SNAP-25. In addition, a SNAP-25 antigen used in an immune response inducing composition must be small enough to only trigger an immune response substantially against a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond, thus increasing the possibility of producing  $\alpha$ -SNAP-25 antibodies that can distinguish a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond from a SNAP-25 lacking a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. Furthermore, it is also very desirable to generate  $\alpha$ -SNAP-25 antibodies of a single amino acid sequence in a good yield that are reproducibly selective and which bind with acceptable avidity in order to permit the design of a highly sensitive assay.

**[0027]** The sequence surrounding a BoNT/A cleavage site present in SNAP-25 is denoted as  $P_5-P_4-P_3-P_2-P_1-P_1'-P_2'-P_3'-P_4'-P_5'$ , with  $P_1-P_1'$  representing the scissile bond. Upon cleavage by retargeted endopeptidase, the resulting cleavage products produced comprise a fragment including the  $P_5-P_4-P_3-P_2-P_1$  sequence and a fragment including the  $P_1'-P_2'-P_3'-P_4'-P_5'$ . Thus, as used herein, the term “SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond” refers to any SNAP-25 having the  $P_1$  residue as its carboxyl-terminal amino acid. For example,  $Q_{197}-R_{198}$  of human SNAP-25 (SEQ ID NO:5) represents the  $P_1-P_1'$  scissile bond for the BoNT/A cleavage site. As such, “SNAP-25 having a carboxyl-terminus glutamine of the BoNT/A cleavage site scissile bond” would be any SNAP-25 cleavage product having a glutamine at its carboxyl-terminal amino acid where the glutamine represents  $Q_{197}$  of the scissile bond. As another example,  $K_{204}-H_{205}$  of *Torpedo marmorata* SNAP-25 (SEQ ID NO:16) represents the  $P_1-P_1'$  scissile bond for the BoNT/A cleavage site. As such,

“SNAP-25 having a carboxyl-terminus lysine of the BoNT/A cleavage site scissile bond” would be any SNAP-25 cleavage product having a lysine at its carboxyl-terminal amino acid where the lysine represents  $K_{204}$  of the scissile bond.

**[0028]** The SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond from the BoNT/A cleavage site can be modified to enhance the immunogenicity of a SNAP-25 antigen, a hapten, or any other antigenic compound that is immunogenic, non-immunogenic, or weakly immunogenic when not associated with the modification. In an aspect of this embodiment, the carboxyl-terminal  $P_1$  residue from the scissile bond of a SNAP-25 antigen can be carboxylated. Carboxylation increases the desired immunogenic properties of a SNAP-25 antigen in two respects. First, because charged amino acids enhance immunogenicity, adding a COO— group to the carboxyl-terminal residue will increase the overall immunogenicity of a SNAP-25 antigen. Second, because the  $P_1$  residue of the BoNT/A cleavage site scissile bond is in a charged state upon cleavage, adding a COO— group to the carboxyl-terminal residue will better mimic the actual antigen that the  $\alpha$ -SNAP-25 antibodies disclosed in the present specification are designed to selectively bind.

**[0029]** In an aspect of this embodiment, the amino-terminal residue from a SNAP-25 antigen can be modified by the addition of an amino acid adapted to attach the SNAP-25 antigen to a carrier protein, such as, e.g., a keyhole limpet hemocyanin (KLH), an ovalbumin (OVA), a thyroglobulin (THY), a bovine serum albumin (BSA), a soybean trypsin inhibitor (STI), or a multiple attachment peptide (MAP). For example, a cysteine residue can be placed at the amino-terminus in order to conjugate the carrier protein KLH.

**[0030]** Thus, an embodiment, a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can be, e.g., at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 25, or at least 30 amino acids in length. In another embodiment, a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can be, e.g., at most 5, at most 6, at most 7, at most 8, at most 9, at most 10, at most 11, at most 12, at most 13, at most 14, at most 15, at most 16, at most 17, at most 18, at most 19, at most 20, at most 25, or at most 30 amino acids in length. In still another embodiment, a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can be, e.g., between 7-12 amino acids, between 10-15 amino acids, or between 13-18 amino acids.

**[0031]** In another embodiment, the SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond comprises SEQ ID NO:33. In aspects of this embodiment, the SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond comprises SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, or SEQ ID NO:39. In a further embodiment, the SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond comprises SEQ ID NO:40.

**[0032]** In yet another embodiment, the SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond comprises SEQ ID NO:41. In

aspects of this embodiment, the SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond comprises SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46. In a further embodiment, the SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond comprises SEQ ID NO:47.

**[0033]** It is envisioned that any and all SNAP-25 antigens that triggers an immune response that produces  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be useful as a SNAP-25 antigen. Thus, amino acid sequence variants comprising SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:46 can be useful as a SNAP-25 antigen to trigger an immune response that produces  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. Thus, in an embodiment, a SNAP-25 antigen can substitute at least 1, at least 2, at least 3, at least 4, or at least 5 amino acid substitutions, deletions or additions to the SNAP-25 antigens comprising SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:46. In still another embodiment, a SNAP-25 antigen can have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity to the SNAP-25 antigens comprising SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:46.

**[0034]** It is envisioned that one or more carriers may be linked to a SNAP-25 antigen in order to enhance the immunogenicity of a SNAP-25 antigen that is immunogenic, non-immunogenic, or weakly immunogenic when not associated with the carrier. Non-limiting examples, include, e.g., a keyhole limpet hemocyanin (KLH), an ovalbumin (OVA), a thyroglobulin (THY), a bovine serum albumin (BSA), a soybean trypsin inhibitor (STI), or a multiple attachment peptide (MAP). As is well known in the art, a non-antigenic or weakly antigenic antigen can be made antigenic by coupling the antigen to a carrier. Various other carrier and methods for coupling an antigen to a carrier are well known in the art. See, e.g., Harlow and Lane, *supra*, 1998a; Harlow and Lane, *supra*, 1998b; and David W. Waggoner, Jr. et al., *Immunogenicity-enhancing carriers and compositions thereof and methods of using the same*, U.S. Patent Publication No. 20040057958 (Mar. 25, 2004). An epitope can also be generated by expressing the epitope as a fusion protein. Methods for expressing polypeptide fusions are well known to those skilled in the art as described, for example, in Ausubel et al., *Current Protocols in Molecular Biology* (Supplement 47), John Wiley & Sons, New York (1999). As the carboxyl-terminal end of the SNAP-25 antigen must be the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, a carrier must be linked to the amino end of the SNAP-25 antigen.

**[0035]** It is envisioned that one or more flexible spacers may be linked to a SNAP-25 antigen in order to enhance the

immunogenicity of a SNAP-25 antigen that is immunogenic, non-immunogenic, or weakly immunogenic when not associated with the flexible linkers. A flexible spacer increases the overall peptide length of the SNAP-25 antigen and provides flexibility, thereby facilitating the proper presentation of the SNAP-25 antigen to the immune cells. As a non-limiting example, a SNAP-25 immune response inducing composition can comprise a SNAP-25 antigen linked to one or more flexible spacers in tandem to better present SNAP-25 antigen to immune cells, thereby facilitating the immune response.

**[0036]** A flexible space comprising a peptide is at least one amino acid in length and comprises non-charged amino acids with small side-chain R groups, such as, e.g., glycine, alanine, valine, leucine or serine. Thus, in an embodiment a flexible spacer can be, e.g., at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 amino acids in length. In another embodiment, a flexible spacer can be, e.g., at least 1, at most 2, at most 3, at most 4, at most 5, at most 6, at most 7, at most 8, at most 9, or at most 10 amino acids in length. In still another embodiment, a flexible spacer can be, e.g., between 1-3 amino acids, between 2-4 amino acids, between 3-5 amino acids, between 4-6 amino acids, or between 5-7 amino acids. Non-limiting examples of a flexible spacer include, e.g., a G-spacers such as GGG, GGGG (SEQ ID NO:57), and GGGGS (SEQ ID NO:58) or an A-spacers such as AAA, AAAA (SEQ ID NO:59) and AAAAV (SEQ ID NO:60). A flexible spacer is linked in-frame to the SNAP-25 antigen as a fusion protein.

**[0037]** As discussed above, a flexible spacer is used, in part, to increase the overall peptide length of the SNAP-25 antigen. For example, a 5-10 amino acid SNAP-25 antigen can have its overall length increased by linking a 3-5 amino acid flexible space to the amino-end of the SNAP-25 antigen. As another example, a 5-10 amino acid SNAP-25 antigen can have its overall length increased by linking a 4-6 amino acid flexible space to the amino-end of the SNAP-25 antigen. As another example, a 5-10 amino acid SNAP-25 antigen can have its overall length increased by linking a 7-10 amino acid flexible space to the amino-end of the SNAP-25 antigen. As another example, a 7-12 amino acid SNAP-25 antigen can have its overall length increased by linking a 1-3 amino acid flexible space to the amino-end of the SNAP-25 antigen. As another example, a 7-12 amino acid SNAP-25 antigen can have its overall length increased by linking a 4-6 amino acid flexible space to the amino-end of the SNAP-25 antigen. The increased length provided by the flexible spacer allows for the selection of a small sized SNAP-25 antigen, thereby increasing the likelihood that the SNAP-25 antigen will only trigger an immune response substantially against a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, thus increasing the possibility of producing  $\alpha$ -SNAP-25 antibodies that can distinguish a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 lacking a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

**[0038]** It is envisioned that a SNAP-25 immune response inducing composition disclosed in the present specification can optionally comprise a SNAP-25 antigen disclosed in the present specification and one or more adjuvants. As used herein, the term "adjuvant" when used in reference to a

SNAP-25 immune response inducing composition refers to any substance or mixture of substances that increases or diversifies the immune response to a SNAP-25 antigen. An immune response inducing adjuvant can, for example, serve to reduce the number of immunizations or the amount of antigen required for protective immunization. The use of immune response inducing adjuvants in an immune response inducing composition is well known. The main objective of these adjuvants is to allow an increase in the immune response. Non-limiting adjuvants include, e.g., liposomes, oily phases, including, without limitation, the Freund type of adjuvants, such as, e.g., Freund's complete adjuvant (FCA); Freund's incomplete adjuvant (FIA); sapogenin glycosides, such as, e.g., saponins; carbopol; N-acetylmuramyl-L-alanyl-D-isoglutamine (commonly known as muramyl dipeptide or "MDP"); and lipopolysaccharide (LPS). Such adjuvants are generally used in the form of an emulsion with an aqueous phase, or, more commonly, may consist of water-insoluble inorganic salts. These inorganic salts may consist, for example, of aluminum hydroxide, zinc sulfate, colloidal iron hydroxide, calcium phosphate or calcium chloride. Aluminum hydroxide (Al(OH)<sub>3</sub>) is a commonly used adjuvant. Currently, the only FDA-approved adjuvant for use in humans is aluminum salts (Alum) which are used to "depot" antigens by precipitation of the antigens. Adjuvants provided above are merely exemplary. In fact, any immune response inducing adjuvant may be used in an immune response inducing composition disclosed in the present specification as long as the adjuvant satisfies the requisite characteristics for inducing an immune response.

**[0039]** A carrier disclosed in the present specification may also act as an adjuvant. Specific adjuvants and methods of making and using are described in, e.g., Gupta et al. *Vaccine*, 11: 993-306, 1993; Arnon, R. (Ed.) *Synthetic Vaccines 1:83-92*, CRC Press, Inc., Boca Raton, Fla., 1987; and David W. Waggoner, Jr. et al., *Immunogenicity-Enhancing Carriers and Compositions Thereof and Methods of Using the Same*, U.S. Patent Publication No. 20040057958 (Mar. 25, 2004). Additional adjuvants include any compound described in Chapter 7 (pp 141-227) of "Vaccine Design, The Subunit and Adjuvant Approach" (eds. Powell, M. F. and Newman, M. J.) *Pharmaceutical Biotechnology*, Volume 6, Plenum Press (New York). Examples from this compendium include Muramyl Dipeptide (MDP) and Montanide 720. Molecules such as Poly Inosine:Cytosine (Poly I:C) or plasmid DNA containing CpG motifs can also be administered as adjuvants in combination with antigens encapsulated in microparticles. In another example, the adjuvant is an agent that facilitates entry of the antigenic compound into the cytoplasm of a cell such as listeriolysin, streptolysin or a mixture thereof.

**[0040]** Thus, in an embodiment, a SNAP-25 immune response inducing composition comprises a SNAP-25 antigen having a carboxylated carboxyl-terminal glutamine linked to a carrier peptide. In aspects of this embodiment, a SNAP-25 antigen having a carboxylated carboxyl-terminal glutamine comprises SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, or SEQ ID NO:39. In another aspect of this embodiment, a SNAP-25 antigen comprises SEQ ID NO:40. In aspects of this embodiment, the carrier peptide is a keyhole limpet hemocyanin (KLH), an ovalbumin (OVA), a

thyroglobulin (THY), a bovine serum albumin (BSA), a soybean trypsin inhibitor (STI) or a multiple attachment peptide (MAP).

**[0041]** In another embodiment, a SNAP-25 immune response inducing composition comprises a SNAP-25 antigen having a carboxylated carboxyl-terminal lysine linked to a carrier peptide. In aspects of this embodiment, SNAP-25 antigen having a carboxylated carboxyl-terminal lysine comprises SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:46. In another aspect of this embodiment, a SNAP-25 antigen comprises SEQ ID NO:47. In aspects of this embodiment, the carrier peptide is a keyhole limpet hemocyanin (KLH), an ovalbumin (OVA), a thyroglobulin (THY), a bovine serum albumin (BSA), a soybean trypsin inhibitor (STI) or a multiple attachment peptide (MAP).

**[0042]** In yet another embodiment, a SNAP-25 immune response inducing composition comprises a SNAP-25 antigen having a carboxylated C-terminal glutamine linked to one or more flexible linkers and a carrier peptide wherein the flexible linkers intervene between the SNAP-25 antigen and the carrier peptide. In aspects of this embodiment, SNAP-25 antigen having a carboxylated carboxyl-terminal glutamine comprises SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, or SEQ ID NO:39. In another embodiment, a SNAP-25 antigen comprises SEQ ID NO:46. In aspects of this embodiment, the carrier peptide is a keyhole limpet hemocyanin (KLH), an ovalbumin (OVA), a thyroglobulin (THY), a bovine serum albumin (BSA), a soybean trypsin inhibitor (STI) or a multiple attachment peptide (MAP). In aspects of this embodiment, the flexible linker is a G-spacer or an A-spacer.

**[0043]** In still another embodiment, a SNAP-25 immune response inducing composition comprises a SNAP-25 antigen having a carboxylated C-terminal lysine linked to a flexible linker and a carrier peptide wherein the flexible linker intervenes between the SNAP-25 antigen and the carrier peptide. In aspects of this embodiment, SNAP-25 antigen having a carboxylated carboxyl-terminal lysine comprises SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:46. In another aspect of this embodiment, a SNAP-25 antigen comprises SEQ ID NO:47. In aspects of this embodiment, the carrier peptide is a keyhole limpet hemocyanin (KLH), an ovalbumin (OVA), a thyroglobulin (THY), a bovine serum albumin (BSA), a soybean trypsin inhibitor (STI) or a multiple attachment peptide (MAP). In aspects of this embodiment, the flexible linker is a G-spacer or an A-spacer.

**[0044]** Aspects of the present disclosure comprise, in part, a method for producing  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. An  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be produced by a wide variety of methods that are well known in the art. Specific protocols for making and using antibodies as well as detecting, and measuring antibody binding specificity, binding affinity and binding avidity are known in the art. See, e.g., *ANTIBODIES: A LABORATORY MANUAL* (Edward Harlow & David Lane, eds., Cold Spring Harbor Laboratory Press, 2<sup>nd</sup> ed. 1998a); and *USING ANTIBODIES: A LABORATORY MANUAL: PORTABLE PROTO-*

COL No. I (Edward Harlow & David Lane, Cold Spring Harbor Laboratory Press, 1998b); Molecular Cloning, A Laboratory Manual, 2001; and Current Protocols in Molecular Biology, 2004; David Anderson et al., *Therapeutic Polypeptides, Nucleic Acids Encoding Same, and Methods of Use*, U.S. Pat. No. 7,034,132 (Apr. 25, 2005); and Beatriz M. Carreno et al., *Antibodies Against CTLA4*, U.S. Pat. No. 7,034,121 (Apr. 25, 2006).

**[0045]** As a non-limiting example,  $\alpha$ -SNAP-25 polyclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be produced by injecting an animal, such as, e.g., a rabbit, a goat, a mouse or another mammal, with one or more injections of an immune response inducing composition disclosed in the present specification. As another non-limiting example,  $\alpha$ -SNAP-25 polyclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be produced by injecting an egg, such as, e.g., a chicken egg, with one or more injections of an immune response inducing composition disclosed in the present specification. The antibody titer in the immunized animal can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized antigen or a cell-based activity assay. If desired, polyclonal antibodies for an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as protein A affinity chromatography to obtain the IgG fraction, or by affinity purification against the peptide used for producing the antibodies.

**[0046]** As another non-limiting example,  $\alpha$ -SNAP-25 monoclonal antibody that selectively binds to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be produced using a hybridoma method. See e.g., Chapter 6 *Monoclonal Antibodies*, pp. 196-244, Harlow & Lane, supra, 1998a; and Chapter 7 *Growing Hybridomas*, pp. 245-282, Harlow & Lane, supra, 1998a; and Goding, pp. 59-103, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986). In this method, a host animal, such as, e.g., a mouse, a hamster, or another appropriate host animal, is typically exposed to one or more injections of a SNAP-25 antigen disclosed in the present specification to elicit lymphocytes that produce or are capable of producing  $\alpha$ -SNAP-25 antibodies that will specifically bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. The antibody titer in the immunized animal can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized antigen or a cell-based activity assay. Alternatively, the lymphocytes can be immunized in vitro using a suitable cell culture line. At an appropriate time after immunization, e.g., when the antibody titers are highest, antibody-producing cells are isolated from the animal. Generally, either peripheral blood lymphocytes are used, if cells of human origin are desired, or spleen cells or lymph node cells are used, if non-human mammalian sources are desired. The isolated antibody-producing cells are fused with an immortal cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell. Immortalized cell lines are usually transformed mammalian cells,

particularly myeloma cells of rodent, bovine and human origin. Typically, a murine myeloma cell line is fused with splenocytes harvested from an appropriately immunized mouse to produce the hybridoma. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine (HAT). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/0-Ag14 myeloma lines. Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days in culture because they are not transformed). The culture medium in which the hybridoma cells are grown can then be assayed for the presence of  $\alpha$ -SNAP-25 monoclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. For example, hybridoma supernatants can be screened using  $\alpha$ -SNAP-25 positive media in an immunoprecipitation assay, in vitro binding assay, such as, e.g., a radioimmunoassay (RIA) or an enzyme-linked immunosorbent assay (ELISA), or in a cell-based activity assay. Such techniques and assays are known in the art. See e.g., Chapter 11 *Immunoprecipitation*, pp. 421-470, Harlow & Lane, supra, 1998a; Chapter 12 *Immunoblotting*, pp. 471-510, Harlow & Lane, supra, 1998a; Chapter 14 *Immunoassays*, pp. 553-612, Harlow & Lane, supra, 1998a. Additional studies can then be done to determine whether the antibody is also unreactive to a SNAP-25 lacking a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. The binding affinity of an  $\alpha$ -SNAP-25 monoclonal antibody can also be determined, e.g., by Scatchard analysis. See, e.g., Peter J. Munson and David Rodbard, *Ligand: A Versatile Computerized Approach For Characterization of Ligand-Binding Systems*, 107(1) *Anal. Biochem.* 220-239 (1980). After the desired hybridoma cells are identified, limiting dilution procedures are used to isolate clones originating from a single cell until a clonal cell line expressing the desired monoclonal antibody is obtained. Those antibodies sufficiently selective for a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond and bind with sufficiently high avidity are chosen for further characterization and study.

**[0047]** Another alternative for preparing an  $\alpha$ -SNAP-25 monoclonal antibody that selectively binds to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond is by screening a recombinant combinatorial immunoglobulin library, such as, e.g., an antibody phage display library, with a SNAP-25 peptide and isolate immunoglobulin library members that bind a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. Kits for generating and screening phage display libraries are commercially available, such as, e.g., the Recombinant Phage Antibody System (Amersham GE Healthcare, Piscataway, N.J.); and the SurfZAP™ Phage Display Kit (Stratagene, La Jolla, Calif.). Additionally, examples of methods and reagents useful in generating and screening antibody display library can be found in, for example, Ladner et al. U.S. Pat. No. 5,223,409; Borrebaeck et al. U.S. Pat. No. 5,712,089; Griffiths et al. U.S. Pat. No. 5,885,793; Griffiths et al. U.S. Pat. No. 5,962,255; McCafferty et al. U.S. Pat. No. 5,969,108; Griffiths et al. U.S. Pat. No. 6,010,884; Jespers et al. U.S. Pat.

No. 6,017,732; Borrebaeck et al. U.S. Pat. No. 6,027,930; Johnson et al. U.S. Pat. No. 6,140,471; McCafferty et al. U.S. Pat. No. 6,172,197, each of which is hereby incorporated by reference in its entirety.

**[0048]** Aspects of the present disclosure comprise, in part, collecting a sample containing an  $\alpha$ -SNAP-25 antibody or  $\alpha$ -SNAP-25 antibody-producing cells. As used herein, the term “sample containing an  $\alpha$ -SNAP-25 antibody or  $\alpha$ -SNAP-25 antibody-producing cell” refers to any biological matter that contains or potentially contains at least one an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. It is envisioned that any and all samples that can contain an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be used in this method, including, without limitation, blood, plasma, serum and lymph fluid. It is also envisioned that any cell capable of producing an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be used in this method, including, without limitation, a CD8 cells, a CTL cell, a helper T-cell and a B-cell. A variety of well known methods can be used for collecting from an individual a sample containing the  $\alpha$ -SNAP-25 antibody or  $\alpha$ -SNAP-25 antibody-producing cell, see, e.g., Harlow & Lane, supra, 1998a; and Harlow & Lane, supra, 1998b. Similarly, a variety of well known methods can be used for processing a sample to isolate an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. A procedure for collecting a sample can be selected based on the type of antibody to be isolated. As a non-limiting example, when isolating an  $\alpha$ -SNAP-25 polyclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, an appropriate sample can be a blood sample containing such  $\alpha$ -SNAP-25 antibodies, whereas when isolating an  $\alpha$ -SNAP-25 monoclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, an appropriate sample can be an  $\alpha$ -SNAP-25 antibody-producing cell such as a spleen cell or hybridoma.

**[0049]** Aspects of the present disclosure comprise, in part, isolating an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from the sample. Methods of isolating an such  $\alpha$ -SNAP-25 antibodies, such as, e.g.,  $\alpha$ -SNAP-25 polyclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond or  $\alpha$ -SNAP-25 monoclonal antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond are well known to those skilled in the art. See, e.g., Harlow and Lane, supra, 1998a; and Harlow and Lane, supra, 1998b. For example, such  $\alpha$ -SNAP-25 polyclonal antibodies can be isolated from the sample by well known techniques, such as, e.g., affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of immune serum. Subsequently, or alternatively, a specific SNAP-25 antigen can be immobilized on a column or magnetic beads to purify the  $\alpha$ -SNAP-25 polyclonal antibodies that selectively binds to a

SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond by immunoaffinity chromatography. An  $\alpha$ -SNAP-25 monoclonal antibody that selectively binds to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be isolated from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, e.g., protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

**[0050]** Thus, in an embodiment, a method of producing a  $\alpha$ -SNAP-25 antibody that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond comprises the steps (a) administering to an animal a SNAP-25 immune response inducing composition comprising a SNAP-25 antigen having a carboxylated C-terminal glutamine linked to a carrier peptide; (b) collecting from the animal a sample containing an  $\alpha$ -SNAP-25 antibody or  $\alpha$ -SNAP-25 antibody-producing cell; and (c) isolating the  $\alpha$ -SNAP-25 antibody component from the sample. In an aspect of this embodiment, the  $\alpha$ -SNAP-25 antibody that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond is a polyclonal antibody. In another aspect of this embodiment,  $\alpha$ -SNAP-25 antibody that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond is a monoclonal antibody. In a further aspect of this embodiment, an  $\alpha$ -SNAP-25 monoclonal antibody that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond produced is an IgG subtype. In other aspects of this embodiment, SNAP-25 immune response inducing composition further comprises an adjuvant, such as, e.g., polyethylene glycol (PEG), monomethoxypolyethylene glycol (mPEG), or polyvinyl alcohol (PVA).

**[0051]** In another embodiment, a method of producing  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond comprises the steps (a) administering to an animal a SNAP-25 immune response inducing composition comprising a SNAP-25 peptide having a carboxylated C-terminal glutamine linked to a flexible linker and a carrier peptide wherein the flexible linker intervenes between the SNAP-25 peptide and the carrier peptide; (b) collecting from the animal a sample containing an  $\alpha$ -SNAP-25 antibody or  $\alpha$ -SNAP-25 antibody-producing cell; and (c) isolating the  $\alpha$ -SNAP-25 antibody from the sample. In an aspect of this embodiment, the  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond is a polyclonal antibody. In another aspect of this embodiment,  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond is a monoclonal antibody. In a further aspect of this embodiment,  $\alpha$ -SNAP-25 monoclonal antibody that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond produced is an IgG subtype. In other aspects of this embodiment, SNAP-25 immune response inducing composition further comprises an adjuvant, such as, e.g., polyethylene glycol (PEG), monomethoxypolyethylene glycol (mPEG), or polyvinyl alcohol (PVA).

**[0052]** Aspects of the present disclosure comprise, in part, an isolated  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. As used herein, the term “antibody” refers to a molecule generated by an immune system that was made in response to a particular antigen that specifically binds to that antigen, and includes both naturally occurring antibodies and non-naturally occurring antibodies. As used herein, the term “isolated” refers to separating a molecule from its natural environment by the use of human intervention. For example, an antibody can be a polyclonal antibody, a monoclonal antibody, a dimer, a multimer, a multispecific antibody, a humanized antibody, a chimeric antibody, bi-functional antibody, a cell-associated antibody like an Ig receptor, a linear antibody, a diabody, or a minibody, so long as the fragment exhibits the desired biological activity, and single chain derivatives of the same. An antibody can be a full-length immunoglobulin molecule comprising the V<sub>H</sub> and V<sub>L</sub> domains, as well as a light chain constant domain (C<sub>L</sub>) and heavy chain constant domains, C<sub>H1</sub>, C<sub>H2</sub> and C<sub>H3</sub>, or an immunologically active fragment of a full-length immunoglobulin molecule, such as, e.g., a Fab fragment, a F(ab')<sub>2</sub> fragment, a Fc fragment, a Fd fragment, a Fv fragment. An antibody can be derived from any vertebrate species (e.g., human, goat, horse, donkey, murine, rat, rabbit, or chicken), and can be of any type (e.g., IgG, IgE, IgM, IgD, and IgA), class (e.g., IgA, IgD, IgE, IgG, and IgM) or subclass (IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2). For general disclosure on the structure of naturally occurring antibodies, non-naturally occurring antibodies, and antigenic compound-binding fragments thereof, see, e.g., Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994); Borrabeck, *Antibody Engineering*, 2nd ed. (Oxford University Press 1995), each of which is hereby incorporated by reference in its entirety.

**[0053]** Naturally-occurring antibodies are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V<sub>H</sub>) followed by a number of constant domains. Each light chain has a variable domain at one end (V<sub>L</sub>) and a constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains.

**[0054]** The complete antigen-recognition and antigen-binding site is contained within the variable domains of the antibody, i.e., the Fv fragment. This fragment includes a dimer of one heavy chain variable domain (V<sub>H</sub>) and one light chain variable domain (V<sub>L</sub>) in tight, non-covalent association. Each domain comprises four framework regions (FR), which largely adopting a  $\beta$ -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases form part of, the  $\beta$ -sheet structure. Each hypervariable region comprises an amino

acid sequence corresponding to a complementarity determining region (CDRs). Collectively, it the three-dimensional configuration of the six CDR regions that define an antigen-binding site on the surface of the V<sub>H</sub>-V<sub>L</sub> dimer that confers antigen-binding specificity. See e.g., Cyrus Chothia, et al., *Conformations of Immunoglobulin Hypervariable Regions*, *Nature* 342(6252): 877-883 (1989); Elvin A. Kabat, et al *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991), each of which is incorporated by reference in its entirety. The constant domains of the antibody are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity.

**[0055]** A target antigen generally has one or more binding sites, also called epitopes, which are recognized by the CDR-formed antigen-binding site. As used herein, an “epitope” is synonymous with “antigenic determinant” and refers to the site on a target antigen, such as, e.g., a peptide, polysaccharide or lipid-containing molecule, capable of specific binding to an immunoglobulin or T-cell receptor or otherwise interacting with a molecule. Each antibody that specifically binds to a different epitope has a different structure. Thus, one antigen may have more than one corresponding antibody.

**[0056]** Polyclonal antibodies refer to a heterogeneous population of antibody molecules that contain at least two species of antibody capable of binding to a particular antigen. By definition, a polyclonal antibody includes two different antibodies that bind to at least two different epitopes. As used herein, the term “monoclonal antibody” or “monoclonal antibodies” refer to a substantially homogeneous population of antibody molecules that contain only one species of antibody capable of binding a particular antigen i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. By definition, a monoclonal antibody binds to a single epitope. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibodies, each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present disclosure may be made by the hybridoma method first described by Kohler et al (1975) *Nature* 256:495, or may be made by recombinant DNA methods (see for example: U.S. Pat. No. 4,816,567; U.S. Pat. No. 5,807,715). The monoclonal antibodies may also be isolated from phage antibody libraries using the techniques described in Clackson et al (1991) *Nature*, 352: 624-628; Marks et al (1991) *J. Mol. Biol.*, 222:581-597; for example.

**[0057]** Thus, in an embodiment, an  $\alpha$ -SNAP-25 antibody comprises a heavy chain variable domain (V<sub>H</sub>) and a light chain variable domain (V<sub>L</sub>) that selectively binds to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. In an aspect of this

embodiment, the heavy chain variable domain ( $V_H$ ) is SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:80, SEQ ID NO:82, or SEQ ID NO:133. In another aspect of this embodiment, the light chain variable domain ( $V_L$ ) is SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, or SEQ ID NO:92.

**[0058]** In another embodiment, a nucleic acid sequence encodes an  $\alpha$ -SNAP-25 antibody comprising a heavy chain variable domain ( $V_H$ ) and a light chain variable domain ( $V_L$ ) that selectively binds to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. In an aspect of this embodiment, the heavy chain variable domain ( $V_H$ ) is encoded by the nucleic acid sequence SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, or SEQ ID NO:132. In another aspect of this embodiment, the heavy chain variable domain ( $V_H$ ) is encoded by a nucleic acid sequence that is at least 70% identical to, at least 75% identical to, at least 80% identical to, at least 85% identical to, at least 90% identical to, at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical, or at least 99% identical to SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, or SEQ ID NO:132. In yet another aspect of this embodiment, the light chain variable domain ( $V_L$ ) is encoded by SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, or SEQ ID NO:91. In still another aspect of this embodiment, the light chain variable domain ( $V_L$ ) is encoded by a nucleic acid sequence that is at least 70% identical to, at least 75% identical to, at least 80% identical to, at least 85% identical to, at least 90% identical to, at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical, or at least 99% identical to SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, or SEQ ID NO:91.

**[0059]** In another embodiment, an  $\alpha$ -SNAP-25 antibody comprises a heavy chain variable domain ( $V_H$ ) CDR1 region, a CDR2 region, a CDR3 region, or any combination thereof that selectively binds to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. In an aspect of this embodiment, the heavy chain variable domain ( $V_H$ ) CDR1 region is SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:118, SEQ ID NO:119, or SEQ ID NO:120. In another aspect of this embodiment, the heavy chain variable domain ( $V_H$ ) CDR2 region is SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:121, SEQ ID NO:122, or SEQ ID NO:123. In yet another aspect of this embodiment, the heavy chain variable domain ( $V_H$ ) CDR3 region is SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:124, SEQ ID NO:134, or SEQ ID NO:135.

**[0060]** In another embodiment, an  $\alpha$ -SNAP-25 antibody comprises a light chain variable domain ( $V_L$ ) CDR1 region, a CDR2 region, a CDR3 region, or any combination thereof that selectively binds to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. In an aspect of this embodiment, the light chain variable domain ( $V_L$ ) CDR1 region is SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:125, SEQ ID NO:126, SEQ ID NO:127, SEQ ID NO:128, or SEQ ID NO:129. In another aspect of this embodiment, the light chain variable domain ( $V_L$ ) CDR2 region is SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID

NO:111, or SEQ ID NO:112. In yet another aspect of this embodiment, the light chain variable domain ( $V_L$ ) CDR3 region is SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, or SEQ ID NO:117.

**[0061]** In yet another embodiment, an  $\alpha$ -SNAP-25 antibody specifically binds an epitope comprising a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. In an aspect of this embodiment, the epitope comprises SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, or SEQ ID NO:37. In an aspect of this embodiment, the epitope comprises SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, or SEQ ID NO:44.

**[0062]** As discussed above, the sequence surrounding a BoNT/A cleavage site present in SNAP-25 is denoted  $P_5-P_4-P_3-P_2-P_1-P_1'-P_2'-P_3'-P_4'-P_5'$ , with  $P_1-P_1'$  representing the scissile bond. Upon cleavage by BoNT/A, the resulting cleavage products produced comprise a fragment including the  $P_5-P_4-P_3-P_2-P_1$  sequence and a fragment including the  $P_1'-P_2'-P_3'-P_4'-P_5'$ . As used herein, the term “ $\alpha$ -SNAP-25 antibodies that selectively bind to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond” refers to  $\alpha$ -SNAP-25 antibodies that selectively bind to any SNAP-25 cleavage product fragment comprising the  $P_5-P_4-P_3-P_2-P_1$  sequence, but not to any SNAP-25 cleavage product fragment comprising the  $P_1'-P_2'-P_3'-P_4'-P_5'$  sequence or to any SNAP-25 having an intact  $P_1-P_1'$  scissile bond of a BoNT/A cleavage site. As used herein, the term “ $\alpha$ -SNAP-25<sub>197</sub> antibody” refers to an antibody that selectively binds to a SNAP-25 having a carboxyl-terminus  $P_1$  residue that corresponds to glutamine 197 of SEQ ID NO:5. As used herein, the term “ $\alpha$ -SNAP-25<sub>204</sub> antibody” refers to an antibody that selectively binds to a SNAP-25 having a carboxyl-terminus  $P_1$  residue that corresponds to lysine 204 of SEQ ID NO:16.

**[0063]** As used herein, the term “selectively” refers to having a unique effect or influence or reacting in only one way or with only one thing. As used herein, the term “selectively binds”, or “selective binding” when made in reference to an antibody, refers to the discriminatory binding of the antibody to the indicated target epitope such that the antibody does not substantially cross react with non-target epitopes. The minimal size of a peptide epitope, as defined herein, is about five amino acids, and a peptide epitope typically comprises at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, or at least 20 amino acids. A peptide epitope may be discontinuous, i.e., it comprises amino acid residues that are not adjacent in the primary structure of the peptide but are brought together into an epitope by way of the secondary, tertiary, or quaternary structure of the peptide. Furthermore, it is also noted that an epitope might comprise a portion of a molecule other than an amino acid sequence, such as, e.g., a carbohydrate moiety, a lipid moiety like lipoproteins or glycolipids, or a chemically-modified amino acid moiety like a phosphorylated amino acid. In aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can selectively bind a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond comprising at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, or at least 20 amino acids. In other aspects of this embodi-

ment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can selectively bind a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond comprising at most 5, at most 6, at most 7, at most 8, at most 9, at most 10, at most 15, or at most 20 amino acids.

**[0064]** Selective binding includes binding properties such as, e.g., binding affinity, binding specificity, and binding avidity. See David J. King, *Applications and Engineering of Monoclonal Antibodies*, pp. 240 (1998). Binding affinity refers to the length of time the antibody resides at its epitope binding site, and can be viewed as the strength with which an antibody binds its epitope. Binding affinity can be described an antibody's equilibrium dissociation constant (KD), which is defined as the ratio  $K_d/K_a$  at equilibrium. Where  $K_a$  is the antibody's association rate constant and  $k_d$  is the antibody's dissociation rate constant. Binding affinity is determined by both the association and the dissociation and alone neither high association or low dissociation can ensure high affinity. The association rate constant ( $K_a$ ), or on-rate constant ( $K_{on}$ ), measures the number of binding events per unit time, or the propensity of the antibody and the antigen to associate reversibly into its antibody-antigen complex. The association rate constant is expressed in  $M^{-1} s^{-1}$ , and is symbolized as follows:  $[Ab] \times [Ag] \times K_{on}$ . The larger the association rate constant, the more rapidly the antibody binds to its antigen, or the higher the binding affinity between antibody and antigen. The dissociation rate constant ( $K_d$ ), or off-rate constant ( $K_{off}$ ), measures the number of dissociation events per unit time propensity of an antibody-antigen complex to separate (dissociate) reversibly into its component molecules, namely the antibody and the antigen. The dissociation rate constant is expressed in  $s^{-1}$ , and is symbolized as follows:  $[Ab+Ag] \times K_{off}$ . The smaller the dissociation rate constant, the more tightly bound the antibody is to its antigen, or the higher the binding affinity between antibody and antigen. The equilibrium dissociation constant (KD) measures the rate at which new antibody-antigen complexes formed equals the rate at which antibody-antigen complexes dissociate at equilibrium. The equilibrium dissociation constant is expressed in M, and is defined as  $K_{off}/K_{on} = [Ab] \times [Ag] / [Ab+Ag]$ , where  $[Ab]$  is the molar concentration of the antibody,  $[Ag]$  is the molar concentration of the antigen, and  $[Ab+Ag]$  is the molar concentration of the antibody-antigen complex, where all concentrations are of such components when the system is at equilibrium. The smaller the equilibrium dissociation constant, the more tightly bound the antibody is to its antigen, or the higher the binding affinity between antibody and antigen.

**[0065]** Thus, in an embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an association rate constant of, e.g., less than  $1 \times 10^5 M^{-1} s^{-1}$ , less than  $1 \times 10^6 M^{-1} s^{-1}$ , less than  $1 \times 10^7 M^{-1} s^{-1}$ , or less than  $1 \times 10^8 M^{-1} s^{-1}$ . In another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an association rate constant of, e.g., more than  $1 \times 10^5 M^{-1} s^{-1}$ , more than  $1 \times 10^6 M^{-1} s^{-1}$ , more than  $1 \times 10^7 M^{-1} s^{-1}$ , or more than  $1 \times 10^8 M^{-1} s^{-1}$ . In other aspects, the binding affinity of an

$\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an association rate constant between  $1 \times 10^5 M^{-1} s^{-1}$  to  $1 \times 10^8 M^{-1} s^{-1}$ ,  $1 \times 10^6 M^{-1} s^{-1}$  to  $1 \times 10^8 M^{-1} s^{-1}$ ,  $1 \times 10^5 M^{-1} s^{-1}$  to  $1 \times 10^7 M^{-1} s^{-1}$ , or  $1 \times 10^6 M^{-1} s^{-1}$  to  $1 \times 10^7 M^{-1} s^{-1}$ .

**[0066]** In another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have a disassociation rate constant of less than  $1 \times 10^{-3} s^{-1}$ , less than  $1 \times 10^{-4} s^{-1}$ , or less than  $1 \times 10^{-5} s^{-1}$ . In other aspects of this embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have a disassociation rate constant of, e.g., less than  $1.0 \times 10^{-4} s^{-1}$ , less than  $2.0 \times 10^{-4} s^{-1}$ , less than  $3.0 \times 10^{-4} s^{-1}$ , less than  $4.0 \times 10^{-4} s^{-1}$ , less than  $5.0 \times 10^{-4} s^{-1}$ , less than  $6.0 \times 10^{-4} s^{-1}$ , less than  $7.0 \times 10^{-4} s^{-1}$ , less than  $8.0 \times 10^{-4} s^{-1}$ , or less than  $9.0 \times 10^{-4} s^{-1}$ . In another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have a disassociation rate constant of, e.g., more than  $1 \times 10^{-3} s^{-1}$ , more than  $1 \times 10^{-4} s^{-1}$ , or more than  $1 \times 10^{-5} s^{-1}$ . In other aspects of this embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have a disassociation rate constant of, e.g., more than  $1.0 \times 10^{-4} s^{-1}$ , more than  $2.0 \times 10^{-4} s^{-1}$ , more than  $3.0 \times 10^{-4} s^{-1}$ , more than  $4.0 \times 10^{-4} s^{-1}$ , more than  $5.0 \times 10^{-4} s^{-1}$ , more than  $6.0 \times 10^{-4} s^{-1}$ , more than  $7.0 \times 10^{-4} s^{-1}$ , more than  $8.0 \times 10^{-4} s^{-1}$ , or more than  $9.0 \times 10^{-4} s^{-1}$ .

**[0067]** In another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an equilibrium dissociation constant of less than 0.500 nM. In aspects of this embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an equilibrium dissociation constant of, e.g., less than 0.500 nM, less than 0.450 nM, less than 0.400 nM, less than 0.350 nM, less than 0.300 nM, less than 0.250 nM, less than 0.200 nM, less than 0.150 nM, less than 0.100 nM, or less than 0.050 nM. In another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an equilibrium dissociation constant of more than 0.500 nM. In aspects of this embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an equilibrium dissociation constant of, e.g., more than 0.500 nM, more than 0.450 nM, more than 0.400 nM, more than 0.350 nM, more than 0.300 nM, more than 0.250 nM, more than 0.200 nM, more than 0.150 nM, more than 0.100 nM, or more than 0.050 nM.

**[0068]** In yet another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can have an association

rate constant of for the intact SNAP-25 of, e.g., less than  $1 \times 10^0 \text{ M}^{-1} \text{ s}^{-1}$ , less than  $1 \times 10^1 \text{ M}^{-1} \text{ s}^{-1}$ , less than  $1 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$ , less than  $1 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ , or less than  $1 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ . In another embodiment, the binding affinity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can have an association rate constant of for the intact SNAP-25 of, e.g., at most  $1 \times 10^0 \text{ M}^{-1} \text{ s}^{-1}$ , at most  $1 \times 10^1 \text{ M}^{-1} \text{ s}^{-1}$ , at most  $1 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$ , at most  $1 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ , or at most  $1 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ .

**[0069]** Binding specificity is the ability of an antibody to discriminate between a molecule containing its epitope and a molecule that does not contain that epitope. One way to measure binding specificity is to compare the Kon association rate of the antibody for a molecule containing its epitope relative to the Kon association rate of the antibody for a molecule that does not contain that epitope. For example, comparing the association rate constant (Ka) of an  $\alpha$ -SNAP-25 antibody for a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond relative to a SNAP-25 not comprising that epitope, such as, e.g., a SNAP-25 epitope lacking a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond or a SNAP-25 epitope having an intact  $P_1$ — $P_1'$  scissile bond of a BoNT/A cleavage site. In aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has an association rate constant (Ka) for a SNAP-25 not comprising its epitope(s) of, e.g., less than  $1 \times 10^0 \text{ M}^{-1} \text{ s}^{-1}$ , less than  $1 \times 10^1 \text{ M}^{-1} \text{ s}^{-1}$ , less than  $1 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$ , less than  $1 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$  or less than  $1 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ . In other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has an association rate constant (Ka) for a SNAP-25 not comprising its epitope(s) of, e.g., at most  $1 \times 10^0 \text{ M}^{-1} \text{ s}^{-1}$ , at most  $1 \times 10^1 \text{ M}^{-1} \text{ s}^{-1}$ , at most  $1 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$ , at most  $1 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$  or at most  $1 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ .

**[0070]** In yet aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has an association rate constant (Ka) for its epitope relative to a SNAP-25 not comprising that epitope of, e.g., at least 2-fold more, at least 3-fold more, at least 4-fold more, at least 5-fold more, at least 6-fold more, at least 7-fold more, at least 8-fold more, or at least 9-fold more. In further aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_i$  residue of the BoNT/A cleavage site scissile bond has an association rate constant (Ka) for its epitope relative to a SNAP-25 not comprising that epitope of, e.g., at least 10-fold more, at least 100-fold more, at least 1,000-fold more or at least 10,000-fold more. In yet other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_i$  residue of the BoNT/A cleavage site scissile bond has an association rate constant (Ka) for its epitope relative to a SNAP-25 not comprising that epitope of, e.g., at most 1-fold more, at most 2-fold more, at most 3-fold more, at most 4-fold more, at most 5-fold more, at most 6-fold more, at most 7-fold more, at most 8-fold more, or at most 9-fold more. In yet other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a

SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has an association rate constant (Ka) for its epitope relative to a SNAP-25 not comprising that epitope of, e.g., at most 10-fold more, at most 100-fold more, at most 1,000-fold more or at most 10,000-fold more.

**[0071]** The binding specificity of an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can also be characterized as a ratio that such an  $\alpha$ -SNAP-25 antibody can discriminate its SNAP-25 epitope relative to a SNAP-25 not comprising that epitope, such as, e.g., a SNAP-25 epitope lacking a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond or a SNAP-25 epitope having an intact  $P_1$ — $P_1'$  scissile bond of a BoNT/A cleavage site. In aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has a binding specificity ratio for its SNAP-25 epitope relative to a SNAP-25 not comprising that epitope of, e.g., at least 2:1, at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 35:1, or at least 40:1. In yet other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has a binding specificity ratio for its SNAP-25 epitope relative to a SNAP-25 lacking a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond of, e.g., at least 2:1, at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 35:1, or at least 40:1. In still other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond has a binding specificity ratio for its SNAP-25 epitope relative to a SNAP-25 having an intake  $P_i$ — $P_i'$  scissile bond of a BoNT/A cleavage site of, e.g., at least 2:1, at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 35:1, or at least 40:1.

**[0072]** Binding avidity, also known as functional affinity, refers to the sum total of the functional binding strength between a multivalent antibody and its antigen. Antibody molecules can have more than one binding site (e.g., 2 for IgG, 10 for IgM), and many antigens contain more than one antigenic site. While binding avidity of an antibody depends on the binding affinities of the individual antibody binding sites, binding avidity is greater than the binding affinity as all the antibody-antigen interactions must be broken simultaneously for the antibody to dissociate completely. It is envisioned that an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond can selectively bind to any and all epitopes for that antibody.

**[0073]** Thus, in an embodiment, an  $\alpha$ -SNAP-25 antibody is an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. In aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody is an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus glutamine or an  $\alpha$ -SNAP-25 antibody

that selectively binds to a SNAP-25 epitope having a carboxyl-terminus lysine. In other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody is an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus P<sub>1</sub> residue that corresponds to glutamine 197 of SEQ ID NO:5 or an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus P<sub>1</sub> residue that corresponds to lysine 204 of SEQ ID NO:16. In still other aspects of this embodiment, an  $\alpha$ -SNAP-25 antibody is an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminal amino acid sequence of SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:46.

**[0074]** Aspects of the present disclosure comprise, in part, an immuno-based method of detecting retargeted endopeptidase activity. The immuno-based methods disclosed in the present specification can be evaluated by several parameters including, e.g., accuracy, precision, limit of detection (LOD), limits of quantitation (LOQ), range, specificity, selectivity, linearity, ruggedness, and system suitability. The accuracy of a method is the measure of exactness of an analytical method, or the closeness of agreement between the measured value and the value that is accepted as a conventional true value or an accepted reference value. The precision of a method is the degree of agreement among individual test results, when the procedure is applied repeatedly to multiple samplings of a homogeneous sample. As such, precision evaluates 1) within assay variability; 2) within-day variability (repeatability); and 3) between-day variability (intermediate precision); and 4) between-lab variability (reproducibility). Coefficient of variation (CV %) is a quantitative measure of precision expressed relative to the observed or theoretical mean value.

**[0075]** An immuno-based method disclosed in the present specification must be able to detect, over background, the presence of an  $\alpha$ -SNAP-25 antibody-antigen complex comprising a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. The limit of detection (LOD) of a method refers to the concentration of analyte which gives rise to a signal that is significantly different from the negative control or blank and represents the lowest concentration of analyte that can be distinguished from background.

**[0076]** Thus, in an embodiment, the immuno-based method disclosed in the present specification can detect the LOD of retargeted endopeptidase at an amount that is significantly different from a negative control or blank. In aspect of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 10 ng or less, 9 ng or less, 8 ng or less, 7 ng or less, 6 ng or less, 5 ng or less, 4 ng or less, 3 ng or less, 2 ng or less, 1 ng or less of a retargeted endopeptidase. In still other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 900 pg or less, 800 pg or less, 700 pg or less, 600 pg or less, 500 pg or less, 400 pg or less, 300 pg or less, 200 pg or less, 100 pg or less of a retargeted endopeptidase. In further aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 90 pg or less, 80 pg or less, 70 pg or less, 60 pg or less, 50 pg or less, 40 pg or less, 30 pg or less, 20 pg or less, 10 pg or less of a

retargeted endopeptidase. In other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 9 pg or less, 8 pg or less, 7 pg or less, 6 pg or less, 5 pg or less, 4 pg or less, 3 pg or less, 2 pg or less, 1 pg or less of a retargeted endopeptidase. In yet other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 0.9 pg or less, 0.8 pg or less, 0.7 pg or less, 0.6 pg or less, 0.5 pg or less, 0.4 pg or less, 0.3 pg or less, 0.2 pg or less, 0.1 pg or less of a retargeted endopeptidase.

**[0077]** In another aspect of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 100 nM or less or less, 90 nM or less or less, 80 nM or less or less, 70 nM or less or less, 60 nM or less or less, 50 nM or less or less, 40 nM or less or less, 30 nM or less or less, 20 nM or less or less, 10 nM or less or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, or 1 nM or less of a retargeted endopeptidase. In other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 900 pM or less, 800 pM or less, 700 pM or less, 600 pM or less, 500 pM or less, 400 pM or less, 300 pM or less, 200 pM or less, or 100 pM or less of a retargeted endopeptidase. In other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 100 pM or less, 90 pM or less, 80 pM or less, 70 pM or less, 60 pM or less, 50 pM or less, 40 pM or less, 30 pM or less, 20 pM or less, or 10 pM or less of a retargeted endopeptidase. In yet other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOD of, e.g., 10 pM or less of a retargeted endopeptidase, 9 pM or less, 8 pM or less, 7 pM or less, 6 pM or less, 5 pM or less, 4 pM or less, 3 pM or less, 2 pM or less, or 1 pM or less of a retargeted endopeptidase.

**[0078]** The limits of quantitation (LOQ) are the lowest and the highest concentrations of analyte in a sample or specimen that can be measured with an acceptable level of accuracy and precision. The lower limit of quantitation refers to the lowest dose that a detection method can measure consistently from the background. The upper limit of quantitation is the highest dose that a detection method can measure consistently before saturation of the signal occurs. The linear range of the method is the area between the lower and the upper limits of quantitation. The linear range is calculated by subtracting lower limit of quantitation from the upper limit of quantitation. As used herein, the term "signal to noise ratio for the lower asymptote" refers to the signal detected in the method at the lower limit of detection divided by the background signal. As used herein, the term "signal to noise ratio for the upper asymptote" refers to the signal detected in the method at the upper limit of detection divided by the background signal.

**[0079]** Thus, in an embodiment, the immuno-based method disclosed in the present specification can detect the LOQ of retargeted endopeptidase at an amount that is significantly different from a negative control or blank. In aspect of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 10 ng or less, 9 ng or less, 8 ng or less, 7 ng or less, 6 ng or less, 5 ng or less, 4 ng or less, 3 ng or less, 2 ng or less, 1 ng or less of a retargeted endopeptidase. In still other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 900 pg or less, 800

pg or less, 700 pg or less, 600 pg or less, 500 pg or less, 400 pg or less, 300 pg or less, 200 pg or less, 100 pg or less of a retargeted endopeptidase. In further aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 90 pg or less, 80 pg or less, 70 pg or less, 60 pg or less, 50 pg or less, 40 pg or less, 30 pg or less, 20 pg or less, 10 pg or less of a retargeted endopeptidase. In other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 9 pg or less, 8 pg or less, 7 pg or less, 6 pg or less, 5 pg or less, 4 pg or less, 3 pg or less, 2 pg or less, 1 pg or less of a retargeted endopeptidase. In yet other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 0.9 pg or less, 0.8 pg or less, 0.7 pg or less, 0.6 pg or less, 0.5 pg or less, 0.4 pg or less, 0.3 pg or less, 0.2 pg or less, 0.1 pg or less of a retargeted endopeptidase.

**[0080]** In another aspect of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 100 nM or less, 90 nM or less, 80 nM or less, 70 nM or less, 60 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, 10 nM or less, 9 nM or less, 8 nM or less, 7 nM or less, 6 nM or less, 5 nM or less, 4 nM or less, 3 nM or less, 2 nM or less, or 1 nM or less of a retargeted endopeptidase. In other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 900 pM or less, 800 pM or less, 700 pM or less, 600 pM or less, 500 pM or less, 400 pM or less, 300 pM or less, 200 pM or less, or 100 pM or less of a retargeted endopeptidase. In other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 100 pM or less, 90 pM or less, 80 pM or less, 70 pM or less, 60 pM or less, 50 pM or less, 40 pM or less, 30 pM or less, 20 pM or less, or 10 pM or less of a retargeted endopeptidase. In yet other aspects of this embodiment, the immuno-based method disclosed in the present specification has an LOQ of, e.g., 10 pM or less of a retargeted endopeptidase, 9 pM or less, 8 pM or less, 7 pM or less, 6 pM or less, 5 pM or less, 4 pM or less, 3 pM or less, 2 pM or less, or 1 pM or less of a retargeted endopeptidase.

**[0081]** An immuno-based assay useful to practice aspect of the disclosed methods must have a precision of no more than 50%. In aspects of this embodiment, an immuno-based assay has a precision of no more than 50%, no more than 40%, no more than 30%, or no more than 20%. In other aspects of this embodiment, an immuno-based assay has a precision of no more than 15%, no more than 10%, or no more than 5%. In other aspects of this embodiment, an immuno-based assay has a precision of no more than 4%, no more than 3%, no more than 2%, or no more than 1%.

**[0082]** An immuno-based assay useful to practice aspect of the disclosed methods must have an accuracy of at least 50%. In aspects of this embodiment, an immuno-based assay has an accuracy of at least 50%, at least 60%, at least 70%, or at least 80%. In other aspects of this embodiment, an immuno-based assay has an accuracy of at least 85%, at least 90%, or at least 95%. In other aspects of this embodiment, an immuno-based assay has an accuracy of at least 96%, at least 97%, at least 98%, or at least 99%.

**[0083]** An immuno-based method disclosed in the present specification must have a signal to noise ratio for the lower asymptote that is statistically significant and a signal to noise

ratio for the upper asymptote that is statistically significant. In aspects of this embodiment, an immuno-based method disclosed in the present specification has a signal to noise ratio for the lower asymptote of, e.g., at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1 or at least 20:1. In other aspects of this embodiment, an immuno-based method has a signal to noise ratio for the upper asymptote of, e.g., at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 35:1, at least 40:1, at least 45:1, at least 50:1, at least 60:1, at least 70:1, at least 80:1, at least 90:1, or at least 100:1, at least 150:1, at least 200:1, at least 250:1, at least 300:1, at least 350:1, at least 400:1, at least 450:1, at least 500:1, at least 550:1, or at least 600:1.

**[0084]** The specificity of a method defines the ability of the method to measure the analyte of interest to the exclusion of other relevant components, such as, e.g., partially-active or inactive analyte. The selectivity of a method describes the ability of an analytical method to differentiate various substances in a sample. The linearity of a method is its ability to elicit results that are directly, or by a well defined mathematical transformation, proportional to the concentration of analyte in the sample. Thus in an embodiment, an immuno-based method disclosed in the present specification can distinguish a fully-active retargeted endopeptidase from a partially-active retargeted endopeptidase having, e.g., 70% or less, 60% or less, 50% or less, 40% or less, 30% or less, 20% or less, or 10% or less the activity of a fully-active retargeted endopeptidase.

**[0085]** The ruggedness of the method is the reproducibility of the test results obtained for identical samples under normal (but variable) test conditions. Robustness of a procedure is a measure of its capacity to remain unaffected by small but deliberate variations in the method parameters and provides an indication of its reliability in normal usage. Thus, whereas ruggedness evaluates unavoidable changes, robustness evaluates deliberate changes. Typical parameters evaluated by ruggedness and robustness include the effects of freeze/thaw, incubation times, incubation temperature, longevity of reagent, sample preparation, sample storage, cell passage number, lots of re-targeted endopeptidase, variability between purifications, and variability between nicking reactions. Robustness parameters for cell-based assays include the cell bank (beginning, middle and end of freeze), cell passage level, cell seeding density, cell stock density (how many days in culture), cell age in flask (waiting time to seeding), incubation time, different plates, excessive amounts of serum, and source of reagents. The system suitability of the method is the determination of assay performance, including the performance of reagents and instruments, overtime by analysis of a reference standard or reference molecule. System suitability is stressed in FDA guidance referring to the fact that equipment, electronics, assay performance, and samples to be analyzed, constitute an integrated system. System suitability can be evaluated by testing for parallelism, which is when plotting the log dose versus the response, serial dilutions of the reference and serial dilutions of the samples should give rise to parallel curves.

**[0086]** Aspects of the present disclosure comprise, in part, a cell from an established cell line. As used herein, the term "cell" refers to any eukaryotic cell susceptible to retargeted endopeptidase activity by a retargeted endopeptidase or any eukaryotic cell that can uptake a retargeted endopeptidase.

The term cell encompasses cells from a variety of organisms, such as, e.g., murine, rat, porcine, bovine, equine, primate and human cells; from a variety of cell types such as, e.g., neuronal and non-neuronal; and can be isolated from or part of a heterogeneous cell population, tissue or organism. As used herein, the term “established cell line” is synonymous with “immortal cell line,” or “transformed cell line” and refers to a cell culture of cells selected for indefinite propagation from a cell population derived from an organism, tissue, or organ source. By definition, an established cell line excludes a cell culture of primary cells. As used herein, the term “primary cells” are cells harvested directly from fresh tissues or organs and do not have the potential to propagate indefinitely. An established cell line can comprise a heterogeneous population of cells or a uniform population of cells. An established cell line derived from a single cell is referred to as a clonal cell line. An established cell line can be one whose cells endogenously express all component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate and encompasses the binding of a retargeted endopeptidase to its receptor, the internalization of the endopeptidase/receptor complex, the translocation of the retargeted endopeptidase light chain from an intracellular vesicle into the cytoplasm and the proteolytic cleavage of a SNAP-25. Alternatively, an established cell line can be one whose cells have had introduced from an exogenous source at least one component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate and encompasses the binding of a retargeted endopeptidase to its receptor, the internalization of the endopeptidase/receptor complex, the translocation of the retargeted endopeptidase light chain from an intracellular vesicle into the cytoplasm and the proteolytic cleavage of a SNAP-25. Also referred to as a genetically-engineered cell line, cells from such an established cell line may, e.g., express an exogenous retargeted endopeptidase, such as, e.g., an exogenous ORL1, an exogenous DOR, an exogenous KOR, an exogenous MOR, an exogenous Galanin receptor 1, an exogenous Galanin receptor 2, an exogenous Galanin receptor 3, or any combination thereof.

**[0087]** Aspects of the present disclosure comprise, in part, a cell from an established cell line susceptible to retargeted endopeptidase activity. As used herein, the terms “cell(s) susceptible to retargeted endopeptidase activity,” “cell(s) susceptible to retargeted endopeptidase activity by a retargeted endopeptidase,” or “cell(s) from an established cell line susceptible to retargeted endopeptidase activity by a retargeted endopeptidase” refer to cell(s) that can undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate thereby inhibiting exocytosis and encompasses the binding of a retargeted endopeptidase to its receptor, the internalization of the endopeptidase/receptor complex, the translocation of the retargeted endopeptidase activity chain from an intracellular vesicle into the cytoplasm and the proteolytic cleavage of a SNAP-25. By definition, cell(s) susceptible to retargeted endopeptidase activity must express, or be engineered to express, at least one retargeted endopeptidase receptor and at least one SNAP-25 substrate. As used herein, the terms “cell(s) that can uptake retargeted endopeptidase” or “cell(s) comprising an established cell line that can uptake

retargeted endopeptidase” refer to cells that can undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate thereby inhibiting exocytosis and encompasses the binding of a retargeted endopeptidase to its receptor, the internalization of the endopeptidase/receptor complex, the translocation of the retargeted endopeptidase light chain from an intracellular vesicle into the cytoplasm and the proteolytic cleavage of a SNAP-25. By definition, cell(s) that can uptake retargeted endopeptidase must express, or be engineered to express, at least one retargeted endopeptidase receptor and at least one SNAP-25 substrate.

**[0088]** Thus in an embodiment, cells from an established cell line are susceptible to retargeted endopeptidase activity. In aspects of this embodiment, cells from an established cell line are susceptible to retargeted endopeptidase activity by, e.g., about 100 nM or less or less, about 90 nM or less or less, about 80 nM or less or less, about 70 nM or less or less, about 60 nM or less or less, about 50 nM or less or less, about 40 nM or less or less, about 30 nM or less or less, about 20 nM or less or less, about 10 nM or less or less of a retargeted endopeptidase. In other aspects, cells from an established cell line are susceptible to retargeted endopeptidase activity by, e.g., about 9 nM or less, about 8 nM or less, about 7 nM or less, about 6 nM or less, about 5 nM or less, about 4 nM or less, about 3 nM or less, about 2 nM or less, or about 1 nM or less of a retargeted endopeptidase. In yet other aspects, cells from an established cell line are susceptible to retargeted endopeptidase activity by, e.g., about 0.9 nM or less, about 0.8 nM or less, about 0.7 nM or less, about 0.6 nM or less, about 0.5 nM or less, about 0.4 nM or less, about 0.3 nM or less, about 0.2 nM, or about 0.1 nM or less of a retargeted endopeptidase. As used herein, the term “about” when qualifying a value of a stated item, number, percentage, or term refers to a range of plus or minus ten percent of the value of the stated item, percentage, parameter, or term.

**[0089]** In another embodiment, cells comprising an established cell line can uptake a retargeted endopeptidase. In aspects of this embodiment, cells comprising an established cell line can uptake, e.g., about 100 nM or less or less, about 90 nM or less or less, about 80 nM or less or less, about 70 nM or less or less, about 60 nM or less or less, about 50 nM or less or less, about 40 nM or less or less, about 30 nM or less or less, about 20 nM or less or less, about 10 nM or less or less of a retargeted endopeptidase. In other aspects, cells comprising an established cell line possess the ability to uptake about 9 nM or less, about 8 nM or less, about 7 nM or less, about 6 nM or less, about 5 nM or less, about 4 nM or less, about 3 nM or less, about 2 nM or less, or about 1 nM or less of a retargeted endopeptidase. In yet other aspects, cells comprising an established cell line possess the ability to uptake about 0.9 nM or less, about 0.8 nM or less, about 0.7 nM or less, about 0.6 nM or less, about 0.5 nM or less, about 0.4 nM or less, about 0.3 nM or less, about 0.2 nM or less, or about 0.1 nM or less of a retargeted endopeptidase.

**[0090]** Aspects of the present disclosure comprise, in part, cells from an established cell line that exhibit a selective binding for a retargeted endopeptidase disclosed in the present specification. As used herein, the term “selectively binds”, or “selective binding” when made in reference to a retargeted endopeptidase, refers to the discriminatory binding of a retargeted endopeptidase to the indicated target

receptor such that the retargeted endopeptidase does not substantially bind to a non-target receptor. The degree to which cells from an established cell line exhibit selective binding for a retargeted endopeptidase can be measured by the extent these cells exhibit non-selective uptake for a molecule lacking the targeting domain of the retargeted endopeptidase. One way to assess non-selective uptake for a molecule lacking the targeting domain of the retargeted endopeptidase is to measure the non-selective uptake of a  $LH_N$  fragment. An  $LH_N$  fragment is one that comprises a Clostridial toxin translocation domain and a Clostridial toxin enzymatic domain, but lacks any targeting domain altogether. Non-limiting examples of a  $LH_N$  fragments include a  $LH_N/A$  fragment, a  $LH_N/B$  fragment, a  $LH_N/C$  fragment, a  $LH_N/D$  fragment, a  $LH_N/E$  fragment, a  $LH_N/F$  fragment, and a  $LH_N/G$  fragment. An exemplary  $LH_N/A$  fragment is SEQ ID NO:146 which is encoded by the polynucleotide molecule SEQ ID NO:147.

**[0091]** Thus, in an embodiment, cells from an established cell line exhibit selective binding for a retargeted endopeptidase. In aspects of this embodiment, cells from an established cell line exhibit selective binding for a retargeted endopeptidase that represents, e.g., at least 75% of the total activity assayed, at least 80% of the total activity assayed, at least 85% of the total activity assayed, at least 90% of the total activity assayed, or at least 95% of the total activity assayed. In other aspects of this embodiment, cells from an established cell line exhibit selective binding for a retargeted endopeptidase that represents, e.g., about 75% to about 100% of the total activity assayed, about 80% to about 100% of the total activity assayed, about 85% to about 100% of the total activity assayed, about 90% to about 100% of the total activity assayed.

**[0092]** In another embodiment, cells from an established cell line exhibit minimal non-selective uptake of a  $LH_N$  fragment. In aspects of this embodiment, cells from an established cell line exhibit non-selective uptake of a  $LH_N$  fragment that is, e.g., at most 25% of the total uptake measured, at most 20% of the total uptake measured, at most 15% of the total uptake measured, at most 10% of the total uptake measured, or at most 5% of the total uptake measured. In other aspects of this embodiment, cells from an established cell line exhibit non-selective uptake of a  $LH_N$  fragment that is, e.g., about 0% to about 25% of the total uptake measured, about 0% to about 20% of the total uptake measured, about 0% to about 15% of the total uptake measured, about 0% to about 10% of the total uptake measured, or about 0% to about 5% of the total uptake measured.

**[0093]** In yet another embodiment, cells from an established cell line exhibit minimal non-selective uptake of a  $LH_N/A$  fragment. In aspects of this embodiment, cells from an established cell line exhibit non-selective uptake of a  $LH_N/A$  fragment that is, e.g., at most 25% of the total uptake measured, at most 20% of the total uptake measured, at most 15% of the total uptake measured, at most 10% of the total uptake measured, or at most 5% of the total uptake measured. In other aspects of this embodiment, cells from an established cell line exhibit non-selective uptake of a  $LH_N/A$  fragment that is, e.g., about 0% to about 25% of the total uptake measured, about 0% to about 20% of the total uptake measured, about 0% to about 15% of the total uptake

measured, about 0% to about 10% of the total uptake measured, or about 0% to about 5% of the total uptake measured.

**[0094]** Aspects of the present disclosure comprise, in part, cells from an established cell line that exhibit a sufficient number of receptor binding sites on the plasma membrane to confer sensitive and selective binding for a retargeted endopeptidase. An equilibrium saturation binding assay measures the total and non-specific binding of a ligand at various concentrations. The equilibrium dissociation constant ( $K_d$ ) for the ligand and the maximal number of receptor binding sites,  $B_{max}$ , can be calculated from the specific binding using non-linear regression analysis. Specific binding is calculated by subtracting the non-specific binding of a ligand from the total binding observed.  $K_d$  is the concentration of ligand required to reach half-maximal binding and is measured in terms of molarity.  $B_{max}$  is the maximal number of binding sites present on the plasma membrane and is measured in terms of pmol/mg, pmol/cell, fmol/cell, or sites/cell.

**[0095]** Thus, in an embodiment, cells from an established cell line exhibit a sufficient number of receptor binding sites on the plasma membrane to confer sensitive and selective binding for a retargeted endopeptidase. In aspects of this embodiment, cells from an established cell line exhibit a  $B_{max}$  value of, e.g., at least 0.1 fmol/cell, at least 0.2 fmol/cell, at least 0.3 fmol/cell, at least 0.4 fmol/cell, at least 0.5 fmol/cell, at least 0.6 fmol/cell, at least 0.7 fmol/cell, at least 0.8 fmol/cell, at least 0.9 fmol/cell, or at least 1.0 fmol/cell, for the targeting ligand of a retargeted endopeptidase. In other aspects of this embodiment, cells from an established cell line exhibit a  $B_{max}$  value of, e.g., at least 1 fmol/cell, at least 2 fmol/cell, at least 3 fmol/cell, at least 4 fmol/cell, at least 5 fmol/cell, at least 6 fmol/cell, at least 7 fmol/cell, at least 8 fmol/cell, at least 9 fmol/cell, or at least 10 fmol/cell, for the targeting ligand of a retargeted endopeptidase.

**[0096]** Aspects of the present disclosure comprise, in part, cells from an established clonal cell line susceptible to re-targeted endopeptidase activity that are more stable than cells from the parental cell line from which the clonal cell line was derived. As used herein, the term "stable" refers to cells from an established clonal cell line for a particular passage number that exhibit a relative  $EC_{50}$ , sensitivity, efficacy, well-defined upper asymptote, and/or a well-defined dose-response curve for re-targeted endopeptidase activity that is similar to the values for relative  $EC_{50}$ , sensitivity, efficacy, well-defined upper asymptote, and/or a well-defined dose-response curve exhibited by cells from the parental cell line from which the clonal cell line was derived, at the same or similar passage number, where the same assay conditions and the same re-targeted endopeptidase are used in both assays.

**[0097]** Thus in an embodiment, cells from an established clonal cell line are more stable as compared to from the parental cell line from which the clonal cell line was derived. In an aspect of this embodiment, cells from an established clonal cell line are more stable as compared to the parental SK-N-DZ cell line. In another aspect of this embodiment, cells from an established clonal cell line are more stable as compared to the parental SK-N-DZ cell line ATCC CRL-2149. In other aspects of this embodiment, cells from an established clonal cell line are more stable for, e.g., at least 5 more passages, at least 10 more passages, at least 15 more

passages, at least 20 more passages, at least 25 more passages, or at least 30 more passages, as compared to from the parental cell line from which the clonal cell line was derived. In yet other aspects of this embodiment, cells from an established clonal cell line are more stable for, e.g., at least 5 more passages, at least 10 more passages, at least 15 more passages, at least 20 more passages, at least 25 more passages, or at least 30 more passages, as compared to from the parental cell line from which the clonal cell line was derived.

**[0098]** Aspects of the present disclosure comprise, in part, cells from an established clonal cell line susceptible to re-targeted endopeptidase activity that are stable over a plurality of cell passages. As used herein, the term “stable” refers to cells from an established clonal cell line for a particular passage number that exhibit a relative  $EC_{50}$ , sensitivity, efficacy, well-defined upper asymptote, and/or a well-defined dose-response curve for re-targeted endopeptidase activity that is similar to the values for relative  $EC_{50}$ , sensitivity, efficacy, well-defined upper asymptote, and/or a well-defined dose-response curve exhibited by cells from the same established clonal cell line, but from a prior passage or passages, where the same assay conditions and the same re-targeted endopeptidase are used in both assays.

**[0099]** Cells from an established cell line disclosed in the present specification can exhibit a consistent sensitivity for re-targeted endopeptidase activity over a plurality of cell passages. As used herein, the term “sensitivity for re-targeted endopeptidase activity” refers to the lowest dose that an assay can measure consistently above the signal detected by a non-treatment control or background signal.

**[0100]** Thus, in an embodiment, cells from the established clonal cell line exhibit a sensitivity for re-targeted endopeptidase activity for any given passages that is e.g., 100 nM or less, about 80 nM or less, about 70 nM or less, about 60 nM or less, about 50 nM or less, about 40 nM or less, about 30 nM or less, about 20 nM or less, about 10 nM or less, about 1 nM or less, about 0.9 nM or less, about 0.8 nM or less, about 0.7 nM or less, about 0.6 nM or less, about 0.5 nM or less, about 0.4 nM or less, about 0.3 nM or less, about 0.2 nM or less, or about 0.1 nM or less of a re-targeted endopeptidase. In aspects of this embodiment, cells from the established clonal cell line exhibit a sensitivity for re-targeted endopeptidase activity for any given passages that is, e.g., about 0.01 nM to about 100 nM, about 0.01 nM to about 75 nM, about 0.01 nM to about 50 nM, about 0.01 nM to about 25 nM, about 0.01 nM to about 20 nM, about 0.01 nM to about 15 nM, about 0.01 nM to about 10 nM, about 0.01 nM to about 5 nM, about 0.001 nM to about 100 nM, about 0.001 nM to about 75 nM, about 0.001 nM to about 50 nM, about 0.001 nM to about 25 nM, about 0.001 nM to about 20 nM, about 0.001 nM to about 15 nM, about 0.001 nM to about 10 nM, or about 0.001 nM to about 5 nM of a re-targeted endopeptidase.

**[0101]** In another embodiment, cells from the established clonal cell line exhibit a sensitivity for re-targeted endopeptidase activity that is about 100 nM or less, about 75 nM or less, about 50 nM or less, about 25 nM or less, less about 20 nM or less, about 15 nM or less, about 10 nM or less, or about 1 nM or less for, e.g., 5 or more cell passages, 10 or more cell passages, 15 or more cell passages, 20 or more cell passages, 25 or more cell passages, 30 or more cell passages, 35 or more cell passages, 40 or more cell passages, 45 or more cell passages, or 50 or more cell passages. In other

aspects of this embodiment, cells from the established clonal cell line exhibit a sensitivity for re-targeted endopeptidase activity that is about 100 nM or less, about 75 nM or less, about 50 nM or less, about 25 nM or less, less about 20 nM or less, about 15 nM or less, about 10 nM or less, or about 1 nM or less for, e.g., about 15 to about 60 passages, about 20 to about 60 passages, about 25 to about 60 passages, about 30 to about 60 passages, about 35 to about 60 passages, about 40 to about 60 passages, about 45 to about 60 passages, about 50 to about 60 passages, about 15 to about 50 passages, about 20 to about 50 passages, about 25 to about 50 passages, about 30 to about 50 passages, about 35 to about 50 passages, about 40 to about 50 passages, about 15 to about 40 passages, about 20 to about 40 passages, about 25 to about 40 passages, or about 30 to about 40 passages.

**[0102]** Cells from an established cell line disclosed in the present specification can exhibit a consistent relative efficacy of re-targeted endopeptidase uptake or re-targeted endopeptidase activity over a plurality of cell passages. As used herein, the term “relative efficacy” refers to how well the upper asymptote for the re-targeted endopeptidase activity detected in the current assay run compares to the upper asymptote for the re-targeted endopeptidase activity detected in a reference standard, a reference molecule, or a reference passage number used on that assay. As used herein, the term “signal to noise ratio for the upper asymptote” refers to the signal detected in an assay at the upper limit of detection divided by the signal detected by a non-treatment control or background signal. The upper limit of detection is the highest dose that an assay can measure consistently before saturation of the signal occurs.

**[0103]** Thus, in an embodiment, cells from an established cell line disclosed in the present specification can exhibit a well defined upper asymptote over a plurality of cell passages and maintain a signal to noise ratio that is consistent and adequate for the assay. In aspects of this embodiment, cells from an established cell line disclosed in the present specification must have a well defined signal to noise ratio for the upper asymptote for re-targeted endopeptidase activity of, e.g., at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 35:1, at least 40:1, at least 45:1, at least 50:1, at least 60:1, at least 70:1, at least 80:1, at least 90:1, or at least 100:1, at least 150:1, at least 200:1, at least 250:1, at least 300:1, at least 350:1, at least 400:1, at least 450:1, at least 500:1, at least 550:1, or at least 600:1, over, e.g., 5 or more cell passages, 10 or more cell passages, 15 or more cell passages, 20 or more cell passages, 25 or more cell passages, 30 or more cell passages, 35 or more cell passages, 40 or more cell passages, 45 or more cell passages, or 50 or more cell passages. In other aspects of this embodiment, cells from an established cell line disclosed in the present specification must have a well defined signal to noise ratio for the upper asymptote for re-targeted endopeptidase activity of, e.g., at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 35:1, at least 40:1, at least 45:1, at least 50:1, at least 60:1, at least 70:1, at least 80:1, at least 90:1, or at least 100:1, at least 150:1, at least 200:1, at least 250:1, at least 300:1, at least 350:1, at least 400:1, at least 450:1, at least 500:1, at least 550:1, or at least 600:1, over, e.g., about 15 to about 60 passages, about 20 to about 60

passages, about 25 to about 60 passages, about 30 to about 60 passages, about 35 to about 60 passages, about 40 to about 60 passages, about 45 to about 60 passages, about 50 to about 60 passages, about 15 to about 50 passages, about 20 to about 50 passages, about 25 to about 50 passages, about 30 to about 50 passages, about 35 to about 50 passages, about 40 to about 50 passages, about 15 to about 40 passages, about 20 to about 40 passages, about 25 to about 40 passages, or about 30 to about 40 passages.

**[0104]** Cells from an established cell line disclosed in the present specification can exhibit a well defined dose-response curve for re-targeted endopeptidase activity over a plurality of cell passages. As used herein, the term “dose-response curve” refers to the how well the raw data fits the statistical model of choice for that assay. As a non-limiting example, a sigmoidal curve with a four parameter logistics fit is a dose-response curve for an enzymatic activity assay, such as, e.g. a potency assay. As another non-limiting example, a ligand binding with one site saturation fit is a dose-response curve for a ligand/antibody binding assay.

**[0105]** Thus, in an embodiment, cells from an established cell line disclosed in the present specification exhibit a well defined dose-response curve for re-targeted endopeptidase activity over a plurality of cell passages. In aspects of this embodiment, cells from an established cell line disclosed in the present specification exhibit a well defined dose-response curve for re-targeted endopeptidase activity over, e.g., 5 or more cell passages, 10 or more cell passages, 15 or more cell passages, 20 or more cell passages, 25 or more cell passages, 30 or more cell passages, 35 or more cell passages, 40 or more cell passages, 45 or more cell passages, or 50 or more cell passages. In other aspects of this embodiment, cells from an established cell line disclosed in the present specification exhibit a well defined dose-response curve for re-targeted endopeptidase activity over, e.g., about 15 to about 60 passages, about 20 to about 60 passages, about 25 to about 60 passages, about 30 to about 60 passages, about 35 to about 60 passages, about 40 to about 60 passages, about 45 to about 60 passages, about 50 to about 60 passages, about 15 to about 50 passages, about 20 to about 50 passages, about 25 to about 50 passages, about 30 to about 50 passages, about 35 to about 50 passages, about 40 to about 50 passages, about 15 to about 40 passages, about 20 to about 40 passages, about 25 to about 40 passages, or about 30 to about 40 passages.

**[0106]** Cells from an established cell line disclosed in the present specification can exhibit a consistent relative  $EC_{50}$  value for re-targeted endopeptidase activity over a plurality of cell passages. As used herein, the term “relative  $EC_{50}$ ” or “relative  $EC_{50}$  value” refers to an  $EC_{50}$  value for re-targeted endopeptidase activity that is normalized against the  $EC_{50}$  calculated for a reference standard, a reference molecule, or a reference passage number used on that assay.

**[0107]** Thus, in an embodiment, cells from an established clonal cell line exhibit a consistent relative  $EC_{50}$  for re-targeted endopeptidase activity over a plurality of cell passages. In aspects of this embodiment, cells from an established clonal cell line exhibit a consistent relative  $EC_{50}$  for re-targeted endopeptidase activity that is, e.g., about  $\pm 10\%$ , about  $\pm 20\%$ , about  $\pm 30\%$ , about  $\pm 40\%$ , about  $\pm 50\%$ , about  $\pm 60\%$ , about  $\pm 70\%$ , or about  $\pm 75\%$  the relative  $EC_{50}$  for re-targeted endopeptidase activity over, e.g., 5 or more cell passages, 10 or more cell passages, 15 or more cell passages, 20 or more cell passages, 25 or more cell passages, 30 or

more cell passages, 35 or more cell passages, 40 or more cell passages, 45 or more cell passages, or 50 or more cell passages. In other aspects of this embodiment, cells from an established clonal cell line exhibit a relative  $EC_{50}$  for re-targeted endopeptidase activity that is, e.g., about  $\pm 10\%$  to about 75%, about  $\pm 10\%$  to about 70%, about  $\pm 10\%$  to about 60%, about  $\pm 10\%$  to about 50%, about  $\pm 10\%$  to about 40%, about  $\pm 10\%$  to about 30%, or about  $\pm 10\%$  to about 20% the relative  $EC_{50}$  for re-targeted endopeptidase activity over, e.g., 5 or more cell passages, 10 or more cell passages, 15 or more cell passages, 20 or more cell passages, 25 or more cell passages, 30 or more cell passages, 35 or more cell passages, 40 or more cell passages, 45 or more cell passages, or 50 or more cell passages.

**[0108]** Aspects of the present disclosure comprise, in part, a re-targeted endopeptidase. As used herein, the term “re-targeted endopeptidase” is synonymous with “Targeted Vesicular Exocytosis Modulator Protein” or “TVEMP.” Non-limiting examples of re-targeted endopeptidase are disclosed in, e.g., Keith A. Foster et al., Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions, U.S. Pat. No. 5,989,545; Clifford C. Shone et al., Recombinant Toxin Fragments, U.S. Pat. No. 6,461,617; Conrad P. Quinn et al., Methods and Compounds for the Treatment of Mucus Hypersecretion, U.S. Pat. No. 6,632,440; Lance E. Steward et al., Methods And Compositions For The Treatment Of Pancreatitis, U.S. Pat. No. 6,843,998; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Patent Publication 2002/0037833; Keith A. Foster et al., Inhibition of Secretion from Non-neural Cells, U.S. Patent Publication 2003/0180289; J. Oliver Dolly et al., Activatable Recombinant Neurotoxins, WO 2001/014570; Keith A. Foster et al., Re-targeted Toxin Conjugates, International Patent Publication WO 2005/023309; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. patent application Ser. No. 11/376,696; Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Non-Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,075; Dolly, J.O. et al., Activatable Clostridial Toxins, U.S. patent application Ser. No. 11/829,475; Foster, K. A. et al., Fusion Proteins, International Patent Publication WO 2006/059093; and Foster, K. A. et al., Non-Cytotoxic Protein Conjugates, International Patent Publication WO 2006/059105, each of which is incorporated by reference in its entirety. Non-limiting examples of re-targeted endopeptidases include SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:130, and SEQ ID NO:131.

**[0109]** Thus in an embodiment, the re-targeted endopeptidase activity being detected is from a re-targeted endopeptidase. In aspects of this embodiment, the re-targeted endopeptidase activity being detected is from a re-targeted endopeptidase disclosed in Keith A. Foster et al., Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions, U.S. Pat. No. 5,989,545; Clifford C. Shone et al., Recombinant Toxin Fragments, U.S. Pat. No. 6,461,617; Conrad P. Quinn et al., Methods and Compounds for the Treatment of Mucus Hypersecretion, U.S. Pat. No. 6,632,440; Lance E. Steward et al., Methods And Compositions For The Treatment Of Pancreatitis, U.S. Pat. No. 6,843,998; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Patent Publication 2002/0037833; Keith A. Foster et al., Inhibition of Secretion from

Non-neural Cells, U.S. Patent Publication 2003/0180289; J. Oliver Dolly et al., Activatable Recombinant Neurotoxins, WO 2001/014570; Keith A. Foster et al., Re-targeted Toxin Conjugates, International Patent Publication WO 2005/023309; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. patent application Ser. No. 11/376,696; Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Non-Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,075; Dolly, J. O. et al., Activatable Clostridial Toxins, U.S. patent application Ser. No. 11/829,475; Foster, K. A. et al., Fusion Proteins, International Patent Publication WO 2006/059093; and Foster, K. A. et al., Non-Cytotoxic Protein Conjugates, International Patent Publication WO 2006/059105, each of which is incorporated by reference in its entirety. In aspects of this embodiment, a retargeted endopeptidase is SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:130, or SEQ ID NO:131.

**[0110]** In another embodiment, the retargeted endopeptidase activity being detected is from a retargeted endopeptidase having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with a retargeted endopeptidase disclosed in Keith A. Foster et al., Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions, U.S. Pat. No. 5,989,545; Clifford C. Shone et al., Recombinant Toxin Fragments, U.S. Pat. No. 6,461,617; Conrad P. Quinn et al., Methods and Compounds for the Treatment of Mucus Hypersecretion, U.S. Pat. No. 6,632,440; Lance E. Steward et al., Methods And Compositions For The Treatment Of Pancreatitis, U.S. Pat. No. 6,843,998; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Patent Publication 2002/0037833; Keith A. Foster et al., Inhibition of Secretion from Non-neural Cells, U.S. Patent Publication 2003/0180289; J. Oliver Dolly et al., Activatable Recombinant Neurotoxins, WO 2001/014570; Keith A. Foster et al., Re-targeted Toxin Conjugates, International Patent Publication WO 2005/023309; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. patent application Ser. No. 11/376,696; Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Non-Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,075; Dolly, J. O. et al., Activatable Clostridial Toxins, U.S. patent application Ser. No. 11/829,475; Foster, K. A. et al., Fusion Proteins, International Patent Publication WO 2006/059093; and Foster, K. A. et al., Non-Cytotoxic Protein Conjugates, International Patent Publication WO 2006/059105, each of which is incorporated by reference in its entirety. In another embodiment, the retargeted endopeptidase activity being detected is from a retargeted endopeptidase having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with a retargeted endopeptidase of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:130, or SEQ ID NO:131.

**[0111]** In other aspects of this embodiment, the retargeted endopeptidase activity being detected is from a re-targeted endopeptidase having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino acid substitutions, deletions, or additions relative to a retargeted endo-

peptidase disclosed in Keith A. Foster et al., Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions, U.S. Pat. No. 5,989,545; Clifford C. Shone et al., Recombinant Toxin Fragments, U.S. Pat. No. 6,461,617; Conrad P. Quinn et al., Methods and Compounds for the Treatment of Mucus Hypersecretion, U.S. Pat. No. 6,632,440; Lance E. Steward et al., Methods And Compositions For The Treatment Of Pancreatitis, U.S. Pat. No. 6,843,998; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Patent Publication 2002/0037833; Keith A. Foster et al., Inhibition of Secretion from Non-neural Cells, U.S. Patent Publication 2003/0180289; J. Oliver Dolly et al., Activatable Recombinant Neurotoxins, WO 2001/014570; Keith A. Foster et al., Re-targeted Toxin Conjugates, International Patent Publication WO 2005/023309; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. patent application Ser. No. 11/376,696; Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Non-Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,075; Dolly, J.O. et al., Activatable Clostridial Toxins, U.S. patent application Ser. No. 11/829,475; Foster, K. A. et al., Fusion Proteins, International Patent Publication WO 2006/059093; and Foster, K. A. et al., Non-Cytotoxic Protein Conjugates, International Patent Publication WO 2006/059105, each of which is incorporated by reference in its entirety. In other aspects of this embodiment, the retargeted endopeptidase activity being detected is from a retargeted endopeptidase having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino acid substitutions, deletions, or additions relative to a retargeted endopeptidase of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:130, or SEQ ID NO:131.

**[0112]** In yet other aspects of this embodiment, the retargeted endopeptidase activity being detected is from a non-naturally occurring retargeted endopeptidase variant having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to a retargeted endopeptidase disclosed in Keith A. Foster et al., Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions, U.S. Pat. No. 5,989,545; Clifford C. Shone et al., Recombinant Toxin Fragments, U.S. Pat. No. 6,461,617; Conrad P. Quinn et al., Methods and Compounds for the Treatment of Mucus Hypersecretion, U.S. Pat. No. 6,632,440; Lance E. Steward et al., Methods And Compositions For The Treatment Of Pancreatitis, U.S. Pat. No. 6,843,998; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Patent Publication 2002/0037833; Keith A. Foster et al., Inhibition of Secretion from Non-neural Cells, U.S. Patent Publication 2003/0180289; J. Oliver Dolly et al., Activatable Recombinant Neurotoxins, WO 2001/014570; Keith A. Foster et al., Re-targeted Toxin Conjugates, International Patent Publication WO 2005/023309; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. patent application Ser. No. 11/376,696; Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting

Activity For Non-Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,075; Dolly, J. O. et al., Activatable Clostridial Toxins, U.S. patent application Ser. No. 11/829,475; Foster, K. A. et al., Fusion Proteins, International Patent Publication WO 2006/059093; and Foster, K. A. et al., Non-Cytotoxic Protein Conjugates, International Patent Publication WO 2006/059105, each of which is incorporated by reference in its entirety. In yet other aspects of this embodiment, the retargeted endopeptidase activity being detected is from a non-naturally occurring retargeted endopeptidase variant having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to a retargeted endopeptidase of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:130, or SEQ ID NO:131.

**[0113]** In yet another embodiment, the retargeted endopeptidase activity being detected is from an opioid retargeted endopeptidase. Non-limiting examples of opioid retargeted endopeptidase, or opioid-TVEMPs, are described in, e.g., Keith A. Foster et al., Clostridial Toxin Derivatives Able To Modify Peripheral Sensory Afferent Functions, U.S. Pat. No. 5,989,545; J. Oliver Dolly et al., Activatable Recombinant Neurotoxins, U.S. Pat. No. 7,132,259; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Pat. No. 7,244,437; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Pat. No. 7,413,742; Stephan Donovan, Clostridial Toxin Derivatives and Methods For Treating Pain, U.S. Pat. No. 7,415,338; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. Pat. No. 7,514,088; Keith A. Foster, Fusion Proteins, U.S. Patent Publication 2008/0064092; Keith A. Foster, Fusion Proteins, U.S. Patent Publication 2009/0035822; Lance E. Steward et al., Multivalent Clostridial Toxin Derivatives and Methods of Their Use, U.S. Patent Publication 2009/0048431; Keith A. Foster, Non-Cytotoxic Protein Conjugates, U.S. Patent Publication 2009/0162341; Keith A. Foster et al., Re-targeted Toxin Conjugates, International Patent Publication WO 2005/023309; and Lance E. Steward, Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Capabilities for Non-Clostridial Toxin Target Cells, International Patent Application WO 2008/008805; each of which is hereby incorporated by reference in its entirety.

**[0114]** In yet another embodiment, the retargeted endopeptidase activity being detected is from a galanin retargeted endopeptidase. Non-limiting examples of galanin re-targeted endopeptidase, or galanin-TVEMPs, are described in, e.g., Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capability and Enhanced Targeting Activity, U.S. patent application Ser. No. 11/776,043 (Jul. 11, 2007); Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,052 (Jul. 11, 2007); and Steward, L. E. et al., Modified Clostridial Toxins with Enhanced Translocation Capabilities and Altered Targeting Activity For Non-Clostridial Toxin Target Cells, U.S. patent application Ser. No. 11/776,075 (Jul. 11, 2007), each of which is incorporated by reference in its entirety.

**[0115]** Aspects of the present disclosure comprise, in part, a SNAP-25. As used herein, the term “SNAP-25” refers to

a naturally-occurring SNAP-25 or a non-naturally occurring SNAP-25 which is preferentially cleaved by a retargeted endopeptidase. As used herein, the term “preferentially cleaved” refers to that the cleavage rate of SNAP-25 by a retargeted endopeptidase is at least one order of magnitude higher than the cleavage rate of any other substrate by a retargeted endopeptidase. In aspects of this embodiment, the cleavage rate of SNAP-25 by a retargeted endopeptidase is at least two orders of magnitude higher, at least three orders of magnitude higher, at least four orders of magnitude higher, or at least five orders of magnitude higher than that the cleavage rate of any other substrate by retargeted endopeptidase.

**[0116]** As used herein, the term “naturally occurring SNAP-25” refers to any SNAP-25 produced by a naturally-occurring process, including, without limitation, SNAP-25 isoforms produced from a post-translational modification, an alternatively-spliced transcript, or a spontaneous mutation, and SNAP-25 subtypes. A naturally occurring SNAP-25 includes, without limitation, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0117]** As used herein, the term “non-naturally occurring SNAP-25” refers to any SNAP-25 whose structure was modified with the aid of human manipulation, including, without limitation, a SNAP-25 produced by genetic engineering using random mutagenesis or rational design and a SNAP-25 produced by in vitro chemical synthesis. Non-limiting examples of non-naturally occurring SNAP-25s are described in, e.g., Steward, L. E. et al., FRET Protease Assays for Clostridial Toxins, U.S. Pat. No. 7,332,567; Fernandez-Salas et al., Lipophilic Dye-based FRET Assays for Clostridial Toxin Activity, U.S. Patent Publication 2008/0160561, each of which is hereby incorporated by reference in its entirety. A non-naturally occurring SNAP-25 may substitute, delete or add, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0118]** Thus in an embodiment, a SNAP-25 is a naturally occurring SNAP-25. In aspects of this embodiment, the SNAP-25 is a SNAP-25 isoform or a SNAP-25 subtype. In aspects of this embodiment, the naturally occurring SNAP-25 is the naturally occurring SNAP-25 of SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9,

SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24. In other aspects of this embodiment, the SNAP-25 is a naturally occurring SNAP-25 having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0119]** In another embodiment, a SNAP-25 is a non-naturally occurring SNAP-25. In other aspects of this embodiment, the SNAP-25 is a non-naturally occurring SNAP-25 having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, or SEQ ID NO:4. In other aspects of this embodiment, the SNAP-25 is a non-naturally occurring SNAP-25 having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24. In yet other aspects of this embodiment, the SNAP-25 is a non-naturally occurring SNAP-25 having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0120]** A SNAP-25 can be an endogenous SNAP-25 or an exogenous SNAP-25. As used herein, the term “endogenous SNAP-25” refers to a SNAP-25 naturally present in the cell because it is naturally encoded within the cell’s genome, such that the cell inherently expresses the SNAP-25 without the need an external source of SNAP-25 or an external source of genetic material encoding a SNAP-25. The expression of an endogenous SNAP-25 may be with or without environmental stimulation such as, e.g., cell differentiation. By definition, an endogenous SNAP-25 can only be a naturally-occurring SNAP-25 or variants thereof. For example, the following established cell lines express an endogenous SNAP-25: BE(2)-M17, Kelly, LA1-55n, N1E-115, N4TG3, N18, Neuro-2a, NG108-15, PC12, SH-SY5Y, SiMa, SK-N-DZ, and SK-N-BE(2)-C.

**[0121]** As used herein, the term “exogenous SNAP-25” refers to a SNAP-25 expressed in a cell through the introduction of an external source of SNAP-25 or an external source of genetic material encoding a SNAP-25 by human manipulation. The expression of an exogenous SNAP-25

may be with or without environmental stimulation such as, e.g., cell differentiation. As a non-limiting example, cells from an established cell line can express an exogenous SNAP-25 by transient or stably transfection of a SNAP-25. As another non-limiting example, cells from an established cell line can express an exogenous SNAP-25 by protein transfection of a SNAP-25. An exogenous SNAP-25 can be a naturally-occurring SNAP-25 or variants thereof, or a non-naturally occurring SNAP-25 or variants thereof.

**[0122]** Thus in an embodiment, cells from an established cell line express an endogenous SNAP-25. In aspects of this embodiment, the endogenous SNAP-25 expressed by cells from an established cell line is a naturally-occurring SNAP-25. In other aspects of this embodiment, the endogenous SNAP-25 expressed by cells from an established cell line is SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24. In yet aspects of this embodiment, the endogenous SNAP-25 expressed by cells from an established cell line is a naturally occurring SNAP-25, such as, e.g., a SNAP-25 isoform or a SNAP-25 subtype. In other aspects of this embodiment, the endogenous SNAP-25 expressed by cells from an established cell line is a naturally occurring SNAP-25 having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0123]** In another embodiment, cells from an established cell line are transiently or stably engineered to express an exogenous SNAP-25. In an aspect of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally-occurring SNAP-25. In other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express the naturally-occurring SNAP-25 of SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24. In yet other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally occurring SNAP-25, such as, e.g., a SNAP-25 isoform or a SNAP-25 subtype. In still other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally occurring SNAP-25 having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0124]** In another aspect of the embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring SNAP-25. In other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring SNAP-25 having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24. In other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring SNAP-25 having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24. In yet other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring SNAP-25 having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:24.

**[0125]** Assays that detect the cleavage of a SNAP-25 after exposure to a retargeted endopeptidase can be used to assess whether a cell is expressing an endogenous or exogenous SNAP-25. In these assays, generation of a SNAP-25 cleavage-product would be detected in cells expressing a SNAP-25 after retargeted endopeptidase treatment. Non-limiting examples of specific Western blot analysis, as well as well-characterized reagents, conditions and protocols are readily available from commercial vendors that include, without limitation, Amersham Biosciences, Piscataway, N.J.; Bio-Rad Laboratories, Hercules, Calif.; Pierce Biotechnology, Inc., Rockford, Ill; Promega Corporation, Madison, Wis., and Stratagene, Inc., La Jolla, Calif. It is understood that these and similar assays for SNAP-25 cleavage can be useful in identifying cells expressing an endogenous or an exogenous SNAP-25.

**[0126]** As non-limiting examples, Western blot analysis using an antibody that recognize a SNAP-25 cleavage product or both the cleaved and uncleaved forms of SNAP-25 can be used to assay for uptake of retargeted endopeptidase. Examples of  $\alpha$ -SNAP-25 antibodies useful for these assays include, without limitation,  $\alpha$ -SNAP-25 mouse monoclonal antibody SMI-81 (Sternberger Monoclonals Inc., Lutherville, Md.), mouse  $\alpha$ -SNAP-25 monoclonal antibody CI 71.1 (Synaptic Systems, Goettingen, Germany),

$\alpha$ -SNAP-25 mouse monoclonal antibody CI 71.2 (Synaptic Systems, Goettingen, Germany),  $\alpha$ -SNAP-25 mouse monoclonal antibody SP12 (Abcam, Cambridge, Mass.),  $\alpha$ -SNAP-25 rabbit polyclonal antiserum (Synaptic Systems, Goettingen, Germany),  $\alpha$ -SNAP-25 rabbit polyclonal antiserum (Abcam, Cambridge, Mass.), and  $\alpha$ -SNAP-25 rabbit polyclonal antiserum S9684 (Sigma, St Louis, Mo.).

**[0127]** Aspects of the present disclosure comprise, in part, a retargeted endopeptidase receptor. As used herein, the term “retargeted endopeptidase receptor” refers to either a naturally-occurring retargeted endopeptidase receptor or a non-naturally occurring retargeted endopeptidase receptor which preferentially interacts with a retargeted endopeptidase in a manner that elicits a retargeted endopeptidase activity response. As used herein, the term “preferentially interacts” refers to that the equilibrium dissociation constant (KD) of retargeted endopeptidase for a retargeted endopeptidase receptor is at least one order of magnitude less than that of retargeted endopeptidase for any other receptor at the cell surface. The equilibrium dissociation constant, a specific type of equilibrium constant that measures the propensity of an retargeted endopeptidase-retargeted endopeptidase receptor complex to separate (dissociate) reversibly into its component molecules, namely the retargeted endopeptidase and the retargeted endopeptidase receptor, is defined as  $KD=K_a/K_d$  at equilibrium. The association constant ( $K_a$ ) is defined as  $K_a=[C]/[L][R]$  and the disassociation constant ( $K_d$ ) is defined as  $K_d=[L][R]/[C]$ , where [L] equals the molar concentration of retargeted endopeptidase, [R] is the molar concentration of a retargeted endopeptidase receptor, and [C] is the molar concentration of the endopeptidase-receptor complex, and where all concentrations are of such components when the system is at equilibrium. The smaller the dissociation constant, the more tightly bound the retargeted endopeptidase is to its receptor, or the higher the binding affinity between retargeted endopeptidase and retargeted endopeptidase receptor. In aspects of this embodiment, the disassociation constant of retargeted endopeptidase for its receptor is at least two orders of magnitude less, at least three orders of magnitude less, at least four orders of magnitude less, or at least five orders of magnitude less than that of retargeted endopeptidase for any other receptor. In other aspects of this embodiment, the binding affinity of a retargeted endopeptidase that preferentially interacts with its receptor can have an equilibrium disassociation constant (KD) of, e.g., of 500 nM or less, 400 nM or less, 300 nM or less, 200 nM, or less 100 nM or less. In other aspects of this embodiment, the binding affinity of a retargeted endopeptidase that preferentially interacts with its receptor can have an equilibrium disassociation constant (KD) of, e.g., of 90 nM or less, 80 nM or less, 70 nM or less, 60 nM, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM, or less 10 nM or less. As used herein, the term “elicits a retargeted endopeptidase activity response” refers to the ability of a retargeted endopeptidase receptor to interact with a retargeted endopeptidase to form a endopeptidase/receptor complex and the subsequent internalization of that complex into the cell cytoplasm.

**[0128]** As used herein, the term “naturally occurring retargeted endopeptidase receptor” refers to any retargeted endopeptidase receptor produced by a naturally-occurring process, including, without limitation, retargeted endopeptidase receptor isoforms produced from a post-translational modification, an alternatively-spliced transcript, or a spontaneous

mutation, and retargeted endopeptidase receptor subtypes. A naturally occurring retargeted endopeptidase receptor includes, without limitation, naturally occurring opioid receptors like a opiate-like receptor 1 (ORL1), a  $\delta$ -opioid receptor (DOR), a  $\kappa$ -opioid receptor (KOR), and a  $\mu$ -opioid receptor (MOR), such as those described in Christopher Evans et al., Opioid Receptor Genes, U.S. Pat. No. 6,265,563; Christopher Evans et al., Methods of Screening Modulators of Opioid Receptor Activity, U.S. Pat. No. 6,432,652; Christopher Evans et al., Opioid Receptors and Methods of Use, U.S. Pat. No. 7,282,563; and Christopher Evans et al., Delta Opioid Receptor Proteins, U.S. Patent Publication 2008/0219925, each of which is hereby incorporated by reference in its entirety. Other examples of a naturally occurring retargeted endopeptidase receptor includes, without limitation, the galanin receptor 1, the galanin receptor 2, and the galanin receptor 3. Naturally occurring opioid receptors from other vertebrate species are known in the art, such as, e.g., primate, cow, dog, mouse, rat, chicken, and fish, and can be used in aspects of the present specification.

**[0129]** A naturally occurring ORL1 includes, without limitation, SEQ ID NO:25 and SEQ ID NO:26, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:25 or SEQ ID NO:26. A naturally occurring DOR includes, without limitation, SEQ ID NO:27 and SEQ ID NO:28, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:27 or SEQ ID NO:28. A naturally occurring KOR includes, without limitation, SEQ ID NO:29 and SEQ ID NO:30, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:29 or SEQ ID NO:30. A naturally occurring MOR includes, without limitation, SEQ ID NO:31, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:31.

**[0130]** A naturally occurring galanin receptor 1 includes, without limitation, SEQ ID NO:136, SEQ ID NO:137, and SEQ ID NO:138, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:136, SEQ ID NO:137, or SEQ ID NO:138. A naturally occurring galanin receptor 2 includes, without limitation, SEQ ID NO:139, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:139. A naturally occurring galanin receptor 3 includes, without limitation, SEQ ID NO:140, or one that substitutes, deletes or adds, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:140.

**[0131]** As used herein, the term “non-naturally occurring retargeted endopeptidase receptor variant” refers to any retargeted endopeptidase receptor produced with the aid of human manipulation or design, including, without limitation, a retargeted endopeptidase receptor produced by genetic engineering using random mutagenesis or rational design and a retargeted endopeptidase receptor produced by chemical synthesis. Non-limiting examples of non-naturally occurring retargeted endopeptidase receptor variants include, e.g., conservative retargeted endopeptidase receptor variants, non-conservative retargeted endopeptidase receptor variants, retargeted endopeptidase receptor chimeric variants and active retargeted endopeptidase receptor fragments.

**[0132]** As used herein, the term “non-naturally occurring retargeted endopeptidase receptor” refers to any retargeted endopeptidase receptor whose structure was modified with the aid of human manipulation, including, without limitation, a retargeted endopeptidase receptor produced by genetic engineering using random mutagenesis or rational design and a retargeted endopeptidase receptor produced by in vitro chemical synthesis. A non-naturally occurring retargeted endopeptidase receptor may substitute, delete or add, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more amino acids from SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140.

**[0133]** Thus in an embodiment, a retargeted endopeptidase receptor is a naturally occurring retargeted endopeptidase receptor such as, e.g., ORL1, DOR, KOR, or MOR. In aspects of this embodiment, the retargeted endopeptidase receptor is a retargeted endopeptidase receptor isoform or a retargeted endopeptidase receptor subtype. In aspects of this embodiment, the naturally occurring retargeted endopeptidase receptor is the naturally occurring retargeted endopeptidase receptor of SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, or SEQ ID NO:31. In other aspects of this embodiment, the retargeted endopeptidase receptor is a naturally occurring retargeted endopeptidase receptor having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, or SEQ ID NO:31.

**[0134]** In another embodiment, a retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor, such as, e.g., a genetically-engineered ORL1, a genetically-engineered DOR, a genetically-engineered KOR, or a genetically-engineered MOR. In other aspects of this embodiment, the retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, or SEQ ID NO:31. In other aspects of this embodiment, the retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino

acid substitutions, deletions, or additions relative to SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, or SEQ ID NO:31. In yet other aspects of this embodiment, the retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, or SEQ ID NO:31.

**[0135]** In another embodiment, a retargeted endopeptidase receptor is a naturally occurring retargeted endopeptidase receptor such as, e.g., galanin receptor 1, galanin receptor 2, or galanin receptor 3. In aspects of this embodiment, the retargeted endopeptidase receptor is a retargeted endopeptidase receptor isoform or a retargeted endopeptidase receptor subtype. In aspects of this embodiment, the naturally occurring retargeted endopeptidase receptor is the naturally occurring retargeted endopeptidase receptor of SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In other aspects of this embodiment, the retargeted endopeptidase receptor is a naturally occurring retargeted endopeptidase receptor having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140.

**[0136]** In another embodiment, a retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor, such as, e.g., a genetically-engineered galanin receptor 1, a genetically-engineered galanin receptor 2, or a genetically-engineered galanin receptor 3. In other aspects of this embodiment, the retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In other aspects of this embodiment, the retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In yet other aspects of this embodiment, the retargeted endopeptidase receptor is a non-naturally occurring retargeted endopeptidase receptor having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140.

**[0137]** A retargeted endopeptidase receptor can be an endogenous retargeted endopeptidase receptor or an exogenous retargeted endopeptidase receptor. As used herein, the term “endogenous retargeted endopeptidase receptor” refers to a retargeted endopeptidase receptor naturally present in the cell because it is naturally encoded within the cell’s genome, such that the cell inherently expresses the retar-

geted endopeptidase receptor without the need an external source of retargeted endopeptidase receptor or an external source of genetic material encoding a retargeted endopeptidase receptor. Expression of an endogenous retargeted endopeptidase receptor may be with or without environmental stimulation such as e.g., cell differentiation or promoter activation. For example, the following established cell lines express at least one endogenous retargeted endopeptidase receptor: AGN P33, Neuro-2a, SiMa, and SK-N-DZ. An endogenous retargeted endopeptidase receptor can only be a naturally-occurring retargeted endopeptidase receptor or naturally-occurring variants thereof.

**[0138]** As used herein, the term “exogenous retargeted endopeptidase receptor” refers to a retargeted endopeptidase receptor expressed in a cell through the introduction of an external source of retargeted endopeptidase receptor or an external source of genetic material encoding a retargeted endopeptidase receptor by human manipulation. The expression of an exogenous retargeted endopeptidase receptor may be with or without environmental stimulation such as, e.g., cell differentiation or promoter activation. As a non-limiting example, cells from an established cell line can express one or more exogenous retargeted endopeptidase receptors by transient or stably transfection of a polynucleotide molecule encoding a retargeted endopeptidase receptor, such as, e.g., an ORL1, a DOR, a KOR, a MOR, a galanin receptor 1, a galanin receptor 2, or a galanin receptor 3. As another non-limiting example, cells from an established cell line can express one or more exogenous retargeted endopeptidase receptors by protein transfection of the retargeted endopeptidase receptors, such as, e.g., an ORL1, a DOR, a KOR, a MOR, a galanin receptor 1, a galanin receptor 2, or a galanin receptor 3. An exogenous retargeted endopeptidase receptor can be a naturally-occurring retargeted endopeptidase receptor or naturally occurring variants thereof, or non-naturally occurring retargeted endopeptidase receptor or non-naturally occurring variants thereof.

**[0139]** Thus in an embodiment, cells from an established cell line express an endogenous retargeted endopeptidase receptor. In aspects of this embodiment, the endogenous retargeted endopeptidase receptor expressed by cells from an established cell line is a naturally-occurring retargeted endopeptidase receptor. In other aspects of this embodiment, the endogenous retargeted endopeptidase receptor expressed by cells from an established cell line is SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In yet aspects of this embodiment, the endogenous retargeted endopeptidase receptor expressed by cells from an established cell line is a naturally occurring retargeted endopeptidase receptor, such as, e.g., a retargeted endopeptidase receptor isoform or a retargeted endopeptidase receptor subtype. In other aspects of this embodiment, the endogenous retargeted endopeptidase receptor expressed by cells from an established cell line is a naturally occurring retargeted endopeptidase receptor having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140.

**[0140]** In another embodiment, cells from an established cell line are transiently or stably engineered to express an exogenous retargeted endopeptidase receptor. In an aspect of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally-occurring retargeted endopeptidase receptor. In other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express the naturally-occurring retargeted endopeptidase receptor of SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In yet other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally occurring retargeted endopeptidase receptor, such as, e.g., a retargeted endopeptidase receptor isoform or a retargeted endopeptidase receptor subtype. In still other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally occurring retargeted endopeptidase receptor having, e.g., at least 70% amino acid identity, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140.

**[0141]** In another aspect of the embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring retargeted endopeptidase receptor. In other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring retargeted endopeptidase receptor having, e.g., at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid identity with SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring retargeted endopeptidase receptor having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more non-contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140. In yet other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally occurring retargeted endopeptidase receptor having, e.g., 1 or more, 2 or more, 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, 20 or more, 30 or more, 40 or more, 50 or more, or 100 or more contiguous amino acid substitutions, deletions, or additions relative to SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:136, SEQ ID NO:137, SEQ ID NO:138, SEQ ID NO:139, or SEQ ID NO:140.

**[0142]** In another embodiment, cells from an established cell line are transiently or stably engineered to express an exogenous ORL1, an exogenous DOR, an exogenous KOR,

an exogenous MOR, or any combination thereof. In aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally-occurring ORL1, a naturally-occurring DOR, a naturally-occurring KOR, a naturally-occurring MOR, or any combination thereof. In yet other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally-occurring ORL1, a non-naturally-occurring DOR, a non-naturally-occurring KOR, a non-naturally-occurring MOR, or any combination thereof. In still other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express either a naturally-occurring ORL1 or a non-naturally-occurring ORL1, a naturally-occurring DOR or a non-naturally-occurring DOR, a naturally-occurring KOR or a non-naturally-occurring KOR, a naturally-occurring MOR or a non-naturally-occurring MOR, or any combination thereof.

**[0143]** In yet another embodiment, cells from an established cell line are transiently or stably engineered to express an exogenous galanin receptor 1, an exogenous galanin receptor 2, an exogenous galanin receptor 3, or any combination thereof. In aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a naturally-occurring galanin receptor 1, a naturally-occurring galanin receptor 2, a naturally-occurring galanin receptor 3, or any combination thereof. In yet other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express a non-naturally-occurring galanin receptor 1, a non-naturally-occurring galanin receptor 2, a non-naturally-occurring galanin receptor 3, or any combination thereof. In still other aspects of this embodiment, cells from an established cell line are transiently or stably engineered to express either a naturally-occurring galanin receptor 1 or a non-naturally-occurring galanin receptor 1, a naturally-occurring galanin receptor 2 or a non-naturally-occurring galanin receptor 2, a naturally-occurring galanin receptor 3 or a non-naturally-occurring galanin receptor 3, or any combination thereof.

**[0144]** Cells that express one or more endogenous or exogenous re-targeted endopeptidase receptors can be identified by routine methods including direct and indirect assays for re-targeted endopeptidase uptake. Assays that determine retargeted endopeptidase binding or uptake properties can be used to assess whether a cell is expressing a retargeted endopeptidase receptor. Such assays include, without limitation, cross-linking assays using labeled retargeted endopeptidase, such as, e.g., [1251] retargeted endopeptidase, see, e.g., Noriko Yokosawa et al., *Binding of Clostridium botulinum type C neurotoxin to different neuroblastoma cell lines*, 57(1) *Infect. Immun.* 272-277 (1989); Noriko Yokosawa et al., *Binding of botulinum type C1, D and E neurotoxins to neuronal cell lines and synaptosomes*, 29(2) *Toxicol.* 261-264 (1991); and Tei-ichi Nishiki et al., *Identification of protein receptor for Clostridium botulinum type B neurotoxin in rat brain synaptosomes*, 269(14) *J. Biol. Chem.* 10498-10503 (1994). Other non-limiting assays include immunocytochemical assays that detect retargeted endopeptidase binding using labeled or unlabeled antibodies, see, e.g., Atsushi Nishikawa et al., *The receptor and transporter for internalization of Clostridium botulinum type C progenitor toxin into HT-29 cells*, 319(2) *Biochem. Biophys. Res. Commun.* 327-333 (2004) and immunoprecipitation assays, see, e.g., Yukako Fujinaga et al., *Molecu-*

lar characterization of binding subcomponents of *Clostridium botulinum* type C progenitor toxin for intestinal epithelial cells and erythrocytes, 150(Pt 5) Microbiology 1529-1538 (2004), that detect bound re-targeted endopeptidase using labeled or unlabeled antibodies. Antibodies useful for these assays include, without limitation, antibodies selected against re-targeted endopeptidase and/or antibodies selected against a re-targeted endopeptidase receptor, such as, e.g., ORL1, DOR, KOR, MOR, galanin receptor 1, galanin receptor 2, or galanin receptor 3. If the antibody is labeled, the binding of the molecule can be detected by various means, including Western blot analysis, direct microscopic observation of the cellular location of the antibody, measurement of cell or substrate-bound antibody following a wash step, flow cytometry, electrophoresis or capillary electrophoresis, employing techniques well-known to those of skill in the art. If the antibody is unlabeled, one may employ a labeled secondary antibody for indirect detection of the bound molecule, and detection can proceed as for a labeled antibody. It is understood that these and similar assays that determine re-targeted endopeptidase uptake properties or characteristics can be useful in identifying cells expressing endogenous or exogenous or re-targeted endopeptidase receptors.

**[0145]** Assays that monitor the release of a molecule after exposure to a re-targeted endopeptidase can also be used to assess whether a cell is expressing one or more endogenous or exogenous re-targeted endopeptidase receptors. In these assays, inhibition of the molecule's release would occur in cells expressing a re-targeted endopeptidase receptor after re-targeted endopeptidase treatment. Well known assays include methods that measure inhibition of radio-labeled catecholamine release from neurons, such as, e.g., [3H] noradrenaline or [3H] dopamine release, see e.g., A Fassio et al., *Evidence for calcium-dependent vesicular transmitter release insensitive to tetanus toxin and botulinum toxin type F*, 90(3) Neuroscience 893-902 (1999); and Sara Stigliani et al., *The sensitivity of catecholamine release to botulinum toxin C1 and E suggests selective targeting of vesicles set into the readily releasable pool*, 85(2) J. Neurochem. 409-421 (2003), or measures catecholamine release using a fluorometric procedure, see, e.g., Anton de Paiva et al., *A role for the interchain disulfide or its participating thiols in the internalization of botulinum neurotoxin A revealed by a toxin derivative that binds to ecto-acceptors and inhibits transmitter release intracellularly*, 268(28) J. Biol. Chem. 20838-20844 (1993); Gary W. Lawrence et al., *Distinct exocytotic responses of intact and permeabilized chromaffin cells after cleavage of the 25-kDa synaptosomal-associated protein (SNAP-25) or synaptobrevin by botulinum toxin A or B*, 236(3) Eur. J. Biochem. 877-886 (1996); and Patrick Foran et al., *Botulinum neurotoxin C1 cleaves both syntaxin and SNAP-25 in intact and permeabilized chromaffin cells: correlation with its blockage of catecholamine release*, 35(8) Biochemistry 2630-2636 (1996). Other non-limiting examples include assays that measure inhibition of hormone release from endocrine cells, such as, e.g., anterior pituitary cells or ovarian cells. It is understood that these and similar assays for molecule release can be useful in identifying cells expressing endogenous or exogenous or re-targeted endopeptidase receptors.

**[0146]** Assays that detect the cleavage of a SNAP-25 substrate after exposure to a re-targeted endopeptidase can also be used to assess whether a cell is expressing one or

more endogenous or exogenous re-targeted endopeptidase receptors. In these assays, generation of a SNAP-25 cleavage-product, or disappearance of the intact SNAP-25, would be detected in cells expressing a re-targeted endopeptidase receptor after a re-targeted endopeptidase treatment. Non-limiting examples of specific Western blot analysis, as well as well-characterized reagents, conditions and protocols are readily available from commercial vendors that include, without limitation, Amersham Biosciences, Piscataway, N.J.; Bio-Rad Laboratories, Hercules, Calif.; Pierce Biotechnology, Inc., Rockford, Ill.; Promega Corporation, Madison, Wis., and Stratagene, Inc., La Jolla, Calif. It is understood that these and similar assays for SNAP-25 cleavage can be useful in identifying cells expressing endogenous or exogenous re-targeted endopeptidase receptors.

**[0147]** As non-limiting examples, Western blot analysis using an antibody that recognizes SNAP-25-cleaved product or both the cleaved and uncleaved forms of SNAP-25 can be used to assay for uptake of a re-targeted endopeptidase. Examples of  $\alpha$ -SNAP-25 antibodies useful for these assays include, without limitation, SMI-81  $\alpha$ -SNAP-25 mouse monoclonal antibody (Sternberger Monoclonals Inc., Lutherville, Md.), CI 71.1 mouse  $\alpha$ -SNAP-25 monoclonal antibody (Synaptic Systems, Goettingen, Germany), CI 71.2  $\alpha$ -SNAP-25 mouse monoclonal antibody (Synaptic Systems, Goettingen, Germany), SP12  $\alpha$ -SNAP-25 mouse monoclonal antibody (Abcam, Cambridge, Mass.),  $\alpha$ -SNAP-25 rabbit polyclonal antiserum (Synaptic Systems, Goettingen, Germany),  $\alpha$ -SNAP-25 rabbit polyclonal antiserum S9684 (Sigma, St. Louis, Mo.), and  $\alpha$ -SNAP-25 rabbit polyclonal antiserum (Abcam, Cambridge, Mass.).

**[0148]** Aspects of the present disclosure provide cells that through genetic manipulation or recombinant engineering are made to express an exogenous SNAP-25 and/or one or more exogenous re-targeted endopeptidase receptors. Cells useful to express an exogenous SNAP-25 and/or one or more exogenous re-targeted endopeptidase receptors through genetic manipulation or recombinant engineering include neuronal cells and non-neuronal cells that may or may not express an endogenous SNAP-25 and/or one or more endogenous re-targeted endopeptidase receptors. It is further understood that such genetically manipulated or recombinantly engineered cells may express an exogenous SNAP-25 and one or more exogenous re-targeted endopeptidase receptors under control of a constitutive, tissue-specific, cell-specific or inducible promoter element, enhancer element or both. It is understood that any cell is useful as long as the cell can be genetically manipulated or recombinantly engineered to express an exogenous SNAP-25 and/or one or more exogenous re-targeted endopeptidase receptors and is capable of undergoing re-targeted endopeptidase activity.

**[0149]** Methods useful for introducing into a cell an exogenous polynucleotide molecule encoding a component necessary for the cells to undergo the overall cellular mechanism whereby a re-targeted endopeptidase proteolytically cleaves a SNAP-25 substrate, such as, e.g., a SNAP-25, an ORL1, a DOR, a KOR, or a MOR, include, without limitation, chemical-mediated delivery methods, such as, e.g., calcium phosphate-mediated, diethyl-aminoethyl (DEAE) dextran-mediated, lipid-mediated, polyethyleneimine (PEI)-mediated, polylysine-mediated and polybrene-mediated; physical-mediated delivery methods, such as, e.g., biolistic particle delivery, microinjection, protoplast fusion and electroporation; and viral-mediated delivery methods, such as,

e.g., retroviral-mediated transfection, see e.g., *Introducing Cloned Genes into Cultured Mammalian Cells*, pp. 16.1-16.62 (Sambrook & Russell, eds., *Molecular Cloning A Laboratory Manual*, Vol. 3, 3<sup>rd</sup> ed. 2001); Alessia Colosimo et al., *Transfer and Expression of Foreign Genes in Mammalian Cells*, 29(2) *Biotechniques* 314-318, 320-322, 324 (2000); Philip Washbourne & A. Kimberley McAllister, *Techniques for Gene Transfer into Neurons*, 12(5) *Curr. Opin. Neurobiol.* 566-573 (2002); and *Current Protocols in Molecular Biology*, John Wiley and Sons, pp. 9.16.4-9.16.11 (2000), each of which is incorporated by reference in its entirety. One skilled in the art understands that selection of a specific method to introduce a polynucleotide molecule into a cell will depend, in part, on whether the cell will transiently or stably contain a component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate. Non-limiting examples of polynucleotide molecule encoding a component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate as follows: ORL1 polynucleotide molecule of SEQ ID NO:61 or SEQ ID NO:62; DOR polynucleotide molecule of SEQ ID NO:63 or SEQ ID NO:64; KOR polynucleotide molecule of SEQ ID NO:65 or SEQ ID NO:66; MOR polynucleotide molecule of SEQ ID NO:67; galanin receptor 1 polynucleotide molecule of SEQ ID NO:141, SEQ ID NO:142, or SEQ ID NO:143, galanin receptor 2 polynucleotide molecule of SEQ ID NO:144, or galanin receptor 3 polynucleotide molecule of SEQ ID NO:145, and SNAP-25 polynucleotide molecule of SEQ ID NO:68, or SEQ ID NO:69.

**[0150]** Chemical-mediated delivery methods are well-known to a person of ordinary skill in the art and are described in, e.g., Martin Jordan & Florian Worm, *Transfection of Adherent and Suspended Cells by Calcium Phosphate*, 33(2) *Methods* 136-143 (2004); Chun Zhang et al., *Polyethylenimine Strategies for Plasmid Delivery to Brain-Derived Cells*, 33(2) *Methods* 144-150 (2004), each of which is hereby incorporated by reference in its entirety. Such chemical-mediated delivery methods can be prepared by standard procedures and are commercially available, see, e.g., CellPfect Transfection Kit (Amersham Biosciences, Piscataway, N.J.); Mammalian Transfection Kit, Calcium phosphate and DEAE Dextran, (Stratagene, Inc., La Jolla, Calif.); Lipofectamine<sup>TM</sup> Transfection Reagent (Invitrogen, Inc., Carlsbad, Calif.); ExGen 500 Transfection kit (Fermentas, Inc., Hanover, Md.), and SuperFect and Effectene Transfection Kits (Qiagen, Inc., Valencia, Calif.).

**[0151]** Physical-mediated delivery methods are well-known to a person of ordinary skill in the art and are described in, e.g., Jeike E. Biewenga et al., *Plasmid-Mediated Gene Transfer in Neurons using the Biolistics Technique*, 71(1) *J. Neurosci. Methods.* 67-75 (1997); John O'Brien & Sarah C. R. Lummis, *Biolistic and Diolistic Transfection: Using the Gene Gun to Deliver DNA and Lipophilic Dyes into Mammalian Cells*, 33(2) *Methods* 121-125 (2004); M. Golzio et al., *In Vitro and In Vivo Electric Field-Mediated Permeabilization, Gene Transfer, and Expression*, 33(2) *Methods* 126-135 (2004); and Oliver Greschet et al., *New Non-Viral Method for Gene Transfer into Primary Cells*, 33(2) *Methods* 151-163 (2004), each of which is hereby incorporated by reference in its entirety.

**[0152]** Viral-mediated delivery methods are well-known to a person of ordinary skill in the art and are described in,

e.g., Chooi M. Lai et al., *Adenovirus and Adeno-Associated Virus Vectors*, 21(12) *DNA Cell Biol.* 895-913 (2002); Ilya Frolov et al., *Alphavirus-Based Expression Vectors: Strategies and Applications*, 93(21) *Proc. Natl. Acad. Sci. U.S.A.* 11371-11377 (1996); Roland Wolkowicz et al., *Lentiviral Vectors for the Delivery of DNA into Mammalian Cells*, 246 *Methods Mol. Biol.* 391-411 (2004); A. Huser & C. Hofmann, *Baculovirus Vectors: Novel Mammalian Cell Gene-Delivery Vehicles and Their Applications*, 3(1) *Am. J. Pharmacogenomics* 53-63 (2003); Tiziana Tonini et al., *Transient Production of Retroviral-and Lentiviral-Based Vectors for the Transduction of Mammalian Cells*, 285 *Methods Mol. Biol.* 141-148 (2004); Manfred Gossen & Hermann Bujard, *Tight Control of Gene Expression in Eukaryotic Cells by Tetracycline-Responsive Promoters*, U.S. Pat. No. 5,464,758; Hermann Bujard & Manfred Gossen, *Methods for Regulating Gene Expression*, U.S. Pat. No. 5,814,618; David S. Hogness, *Polynucleotides Encoding Insect Steroid Hormone Receptor Polypeptides and Cells Transformed With Same*, U.S. Pat. No. 5,514,578; David S. Hogness, *Polynucleotide Encoding Insect Ecdysone Receptor*, U.S. Pat. No. 6,245,531; Elisabetta Vegeto et al., *Progesterone Receptor Having C. Terminal Hormone Binding Domain Truncations*, U.S. Pat. No. 5,364,791; Elisabetta Vegeto et al., *Mutated Steroid Hormone Receptors, Methods for Their Use and Molecular Switch for Gene Therapy*, U.S. Pat. No. 5,874,534, each of which is hereby incorporated by reference in its entirety. Such viral-mediated delivery methods can be prepared by standard procedures and are commercially available, see, e.g., ViraPower<sup>TM</sup> Adenoviral Expression System (Invitrogen, Inc., Carlsbad, Calif.) and ViraPower<sup>TM</sup> Adenoviral Expression System Instruction Manual 25-0543 version A, Invitrogen, Inc., (Jul. 15, 2002); and AdEasy<sup>TM</sup> Adenoviral Vector System (Stratagene, Inc., La Jolla, Calif.) and AdEasy<sup>TM</sup> Adenoviral Vector System Instruction Manual 064004f, Stratagene, Inc. Furthermore, such viral delivery systems can be prepared by standard methods and are commercially available, see, e.g., BD<sup>TM</sup> Tet-Off and Tet-On Gene Expression Systems (BD Biosciences-Clontech, Palo Alto, Calif.) and BD<sup>TM</sup> Tet-Off and Tet-On Gene Expression Systems User Manual, PT3001-1, BD Biosciences Clontech, (Mar. 14, 2003), GeneSwitch<sup>TM</sup> System (Invitrogen, Inc., Carlsbad, Calif.) and GeneSwitch<sup>TM</sup> System A Mifepristone-Regulated Expression System for Mammalian Cells version D, 25-0313, Invitrogen, Inc., (Nov. 4, 2002); ViraPower<sup>TM</sup> Lentiviral Expression System (Invitrogen, Inc., Carlsbad, Calif.) and ViraPower<sup>TM</sup> Lentiviral Expression System Instruction Manual 25-0501 version E, Invitrogen, Inc., (Dec. 8, 2003); and Complete Control<sup>®</sup> Retroviral Inducible Mammalian Expression System (Stratagene, La Jolla, Calif.) and Complete Control<sup>®</sup> Retroviral Inducible Mammalian Expression System Instruction Manual, 064005e.

**[0153]** Thus, in an embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain a polynucleotide molecule encoding a component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate. In another embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain a polynucleotide molecule encoding a plurality of components necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically

cleaves a SNAP-25 substrate. In aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain a polynucleotide molecule encoding ORL1, DOR, KOR, MOR, or SNAP-25. In aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:61, or SEQ ID NO:62 encoding ORL1. In other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:63, or SEQ ID NO:64 encoding DOR. In yet other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:65, or SEQ ID NO:66 encoding KOR. In still other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:67 encoding MOR.

**[0154]** In other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:141, SEQ ID NO:142, or SEQ ID NO:143 encoding Galanin receptor 1. In yet other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:144 encoding Galanin receptor 2. In still other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:145 encoding Galanin receptor 3. In further aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity transiently contain the polynucleotide molecule of SEQ ID NO:68 or SEQ ID NO:69 encoding SNAP-25.

**[0155]** In another embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain a polynucleotide molecule encoding a component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate. In another embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain a polynucleotide molecule encoding a plurality of components necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate. In aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:61, or SEQ ID NO:62 encoding ORL1. In other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:63, or SEQ ID NO:64 encoding DOR. In yet other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:65, or SEQ ID NO:66 encoding KOR. In still other aspects of this embodiment, cells from an established cell

line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:67 encoding MOR.

**[0156]** In other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:141, SEQ ID NO:142, or SEQ ID NO:143 encoding Galanin receptor 1. In yet other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:144 encoding Galanin receptor 2. In yet other aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:145 encoding Galanin receptor 3. In further aspects of this embodiment, cells from an established cell line susceptible to retargeted endopeptidase activity stably contain the polynucleotide molecule of SEQ ID NO:68 or SEQ ID NO:69 encoding SNAP-25.

**[0157]** As mentioned above, an exogenous component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate, such as, e.g., a SNAP-25, an ORL1, a DOR, a KOR, a MOR, a Galanin receptor 1, a Galanin receptor 2, or a Galanin receptor 3, disclosed in the present specification can be introduced into a cell. Any and all methods useful for introducing such an exogenous component with a delivery agent into a cell population can be useful with the proviso that this method transiently introduces the exogenous component disclosed in the present specification in at least 50% of the cells within a given cell population. Thus, aspects of this embodiment can include a cell population in which, e.g., at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% of the given cell population transiently contains an exogenous component necessary for the cells to undergo the overall cellular mechanism whereby a retargeted endopeptidase proteolytically cleaves a SNAP-25 substrate, such as, e.g., a SNAP-25, an ORL1, a DOR, a KOR, a MOR, a Galanin receptor 1, a Galanin receptor 2, or a Galanin receptor 3, disclosed in the present specification. As used herein, the term "delivery agent" refers to any molecule that enables or enhances internalization of a covalently-linked, non-covalently-linked or in any other manner associated with a polypeptide into a cell. Thus, the term "delivery agent" encompasses, without limitation, proteins, peptides, peptidomimetics, small molecules, polynucleotide molecules, liposomes, lipids, viruses, retroviruses and cells that, without limitation, transport a covalently or non-covalently linked molecule to the cell membrane, cell cytoplasm or nucleus. It further is understood that the term "delivery agent" encompasses molecules that are internalized by any mechanism, including delivery agents which function via receptor mediated endocytosis and those which are independent of receptor mediated endocytosis.

**[0158]** A delivery agent can also be an agent that enables or enhances cellular uptake of a covalently linked component, like SNAP-25, ORL1, DOR, KOR, MOR, Galanin receptor 1, Galanin receptor 2, or Galanin receptor 3, such as, e.g., by chemical conjugation or by genetically produced fusion proteins. Methods that covalently link delivery agents and methods of using such agents are described in, e.g., Steven F. Dowdy, Protein Transduction System and Methods of Use Thereof, International Publication No WO 00/34308;

Gérard Chassaing & Alain Prochiantz, Peptides which can be Used as Vectors for the Intracellular Addressing of Active Molecules, U.S. Pat. No. 6,080,724; Alan Frankel et al., Fusion Protein Comprising TAT-derived Transport Moiety, U.S. Pat. No. 5,674,980; Alan Frankel et al., TAT-derived Transport Polypeptide Conjugates, U.S. Pat. No. 5,747,641; Alan Frankel et al., TAT-derived Transport Polypeptides and Fusion Proteins, U.S. Pat. No. 5,804,604; Peter F. J. O'Hare et al., Use of Transport Proteins, U.S. Pat. No. 6,734,167; Yao-Zhong Lin & Jack J. Hawiger, Method for Importing Biologically Active Molecules into Cells, U.S. Pat. No. 5,807,746; Yao-Zhong Lin & Jack J. Hawiger, Method for Importing Biologically Active Molecules into Cells, U.S. Pat. No. 6,043,339; Yao-Zhong Lin et al., Sequence and Method for Genetic Engineering of Proteins with Cell Membrane Translocating Activity, U.S. Pat. No. 6,248,558; Yao-Zhong Lin et al., Sequence and Method for Genetic Engineering of Proteins with Cell Membrane Translocating Activity, U.S. Pat. No. 6,432,680; Jack J. Hawiger et al., Method for Importing Biologically Active Molecules into Cells, U.S. Pat. No. 6,495,518; Yao-Zhong Lin et al., Sequence and Method for Genetic Engineering of Proteins with Cell Membrane Translocating Activity, U.S. Pat. No. 6,780,843; Jonathan B. Rothbard & Paul A Wender, Method and Composition for Enhancing Transport Across Biological Membranes, U.S. Pat. No. 6,306,993; Jonathan B. Rothbard & Paul A Wender, Method and Composition for Enhancing Transport Across Biological Membranes, U.S. Pat. No. 6,495,663; and Pamela B. Davis et al., Fusion Proteins for Protein Delivery, U.S. Pat. No. 6,287,817, each of which is incorporated by reference in its entirety.

**[0159]** A delivery agent can also be an agent that enables or enhances cellular uptake of a non-covalently associated component, like SNAP-25, ORL1, DOR, KOR, MOR, Galanin receptor 1, Galanin receptor 2, or Galanin receptor 3. Methods that function in the absence of covalent linkage and methods of using such agents are described in, e.g., Gilles Divita et al, Peptide-Mediated Transfection Agents and Methods of Use, U.S. Pat. No. 6,841,535; Philip L Feigner and Olivier Zelphati, Intracellular Protein Delivery Compositions and Methods of Use, U.S. Patent Publication No. 2003/0008813; and Michael Karas, Intracellular Delivery of Small Molecules, Proteins and Nucleic Acids, U.S. Patent Publication 2004/0209797, each of which is incorporated by reference in its entirety. Such peptide delivery agents can be prepared and used by standard methods and are commercially available, see, e.g. the CHARIOT™ Reagent (Active Motif, Carlsbad, Calif.); BIO-PORTER® Reagent (Gene Therapy Systems, Inc., San Diego, Calif.), BIO TREK™ Protein Delivery Reagent (Stratagene, La Jolla, Calif.), and PRO-JECT™ Protein Transfection Reagent (Pierce Biotechnology Inc., Rockford, Ill.).

**[0160]** Aspects of the present disclosure comprise, in part, a sample comprising a retargeted endopeptidase. As used herein, the term “sample comprising a retargeted endopeptidase” refers to any biological matter that contains or potentially contains an active retargeted endopeptidase. A variety of samples can be assayed according to a method disclosed in the present specification including, without limitation, purified, partially purified, or unpurified retargeted endopeptidase; recombinant single chain or di-chain retargeted endopeptidase with a naturally or non-naturally occurring sequence; recombinant retargeted endopeptidase with a modified protease specificity; recombinant retargeted

endopeptidase with an altered cell specificity; bulk retargeted endopeptidase; a formulated retargeted endopeptidase product; and cells or crude, fractionated or partially purified cell lysates from, e.g., bacteria, yeast, insect, or mammalian sources; blood, plasma or serum; raw, partially cooked, cooked, or processed foods; beverages; animal feed; soil samples; water samples; pond sediments; lotions; cosmetics; and clinical formulations. It is understood that the term sample encompasses tissue samples, including, without limitation, mammalian tissue samples, livestock tissue samples such as sheep, cow and pig tissue samples; primate tissue samples; and human tissue samples. Such samples encompass, without limitation, intestinal samples such as infant intestinal samples, and tissue samples obtained from a wound. As non-limiting examples, a method of detecting picomolar amounts of retargeted endopeptidase activity can be useful for determining the presence or activity of a retargeted endopeptidase in a food or beverage sample; to assay a sample from a human or animal, for example, exposed to a retargeted endopeptidase or having one or more symptoms of botulism; to follow activity during production and purification of bulk retargeted endopeptidase; to assay a formulated retargeted endopeptidase product used in pharmaceutical or cosmetics applications; or to assay a subject's blood serum for the presence or absence of neutralizing  $\alpha$ -retargeted endopeptidase antibodies.

**[0161]** Thus, in an embodiment, a sample comprising a retargeted endopeptidase is a sample comprising any amount of a retargeted endopeptidase. In aspects of this embodiment, a sample comprising a retargeted endopeptidase comprises about 100 ng or less, about 10 ng or less, about 1 ng or less, about 100 pg or less, about 10 pg or less, or about 1 pg or less of a retargeted endopeptidase. In other aspects of this embodiment, a sample comprising a retargeted endopeptidase comprises about 1  $\mu$ M or less, about 100 nM or less, about 10 nM or less, about 1 nM or less, about 100 nM or less, about 10 nM or less, about 1 nM or less of a retargeted endopeptidase.

**[0162]** Aspects of the present disclosure comprise, in part, isolating from the treated cell a SNAP-25 component comprising a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. As used herein, the term “SNAP-25 component comprising a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond” refers to a cellular component containing the SNAP-25 cleavage product. It is envisioned that any method suitable for enriching or isolating a SNAP-25 component can be useful, including, without limitation, cell lysing protocols, spin-column purification protocols, immunoprecipitation, affinity purification, and protein chromatography.

**[0163]** Aspects of the present disclosure comprise, in part, an  $\alpha$ -SNAP-25 antibody linked to a solid phase support. As used herein, the term “solid-phase support” is synonymous with “solid phase” and refers to any matrix that can be used for immobilizing an  $\alpha$ -SNAP-25 antibody disclosed in the present specification. Non-limiting examples of solid phase supports include, e.g., a tube; a plate; a column; pins or “dipsticks”; a magnetic particle, a bead or other spherical or fibrous chromatographic media, such as, e.g., agarose, sepharose, silica and plastic; and sheets or membranes, such as, e.g., nitrocellulose and polyvinylidene fluoride (PVDF). The solid phase support can be constructed using a wide variety of materials such as, e.g., glass, carbon, polystyrene, poly-

vinylchloride, polypropylene, polyethylene, dextran, nylon, diazocellulose, or starch. The solid phase support selected can have a physical property that renders it readily separable from soluble or unbound material and generally allows unbound materials, such as, e.g., excess reagents, reaction by-products, or solvents, to be separated or otherwise removed (by, e.g., washing, filtration, centrifugation, etc.) from solid phase support-bound assay component. Non-limiting examples of how to make and use a solid phase supports are described in, e.g., *Molecular Cloning, A Laboratory Manual*, supra, (2001); and *Current Protocols in Molecular Biology*, supra, (2004), each of which is hereby incorporated by reference in its entirety.

**[0164]** Aspects of the present disclosure comprise, in part, detecting the presence of an antibody-antigen complex comprising an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond and a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. It is envisioned that any detection system can be used to practice aspects of this disclosed immuno-based method, with the provision that the signal to noise ratio can distinguish to a statistically significant degree the signal from the antibody-antigen complex from the background signal. Non-limiting examples of immuno-based detection systems include immunoblot analysis, like Western blotting and dot-blotting, immunoprecipitation analysis, enzyme-linked immunosorbent analysis (ELISA), and sandwich ELISA. The detection of the signal can be achieved using autoradiography with imaging or phosphorimaging (AU), chemiluminescence (CL), electrochemiluminescence (ECL), bioluminescence (BL), fluorescence, resonance energy transfer, plane polarization, colorimetric, or flow cytometry (FC). Descriptions of immuno-based detection systems are disclosed in, e.g., Michael M. Rauhut, *Chemiluminescence*, In *Kirk-Othmer Concise Encyclopedia of Chemical Technology* (Ed. Grayson, 3rd ed, John Wiley and Sons, 1985); A. W. Knight, *A Review of Recent Trends in Analytical Applications of Electrogenenerated Chemiluminescence*, *Trends Anal. Chem.* 18(1): 47-62 (1999); K. A. Fahrnich, et al., *Recent Applications of Electrogenenerated Chemiluminescence in Chemical Analysis*, *Talanta* 54(4): 531-559 (2001); *Commonly Used Techniques in Molecular Cloning*, pp. A8.1-A8-55 (Sambrook & Russell, eds., *Molecular Cloning A Laboratory Manual*, Vol. 3, 3<sup>rd</sup> ed. 2001); *Detection Systems*, pp. A9.1-A9-49 (Sambrook & Russell, eds., *Molecular Cloning A Laboratory Manual*, Vol. 3, 3<sup>rd</sup> ed. 2001); *Electrogenenerated Chemiluminescence*, (Ed. Allen J. Bard, Marcel Dekker, Inc., 2004), each of which is hereby incorporated by reference in its entirety.

**[0165]** A sandwich ELISA (or sandwich immunoassay) is a method based on two antibodies, which bind to different epitopes on the antigen. A capture antibody having a high binding specificity for the antigen of interest, is bound to a solid surface. The antigen is then added followed by addition of a second antibody referred to as the detection antibody. The detection antibody binds the antigen to a different epitope than the capture antibody. The antigen is therefore 'sandwiched' between the two antibodies. The antibody binding affinity for the antigen is usually the main determinant of immunoassay sensitivity. As the antigen concentration increases the amount of detection antibody increases leading to a higher measured response. To quantify the

extent of binding different reporter systems can be used, such as, e.g., an enzyme attached to the secondary antibody and a reporter substrate where the enzymatic reaction forms a readout as the detection signal. The signal generated is proportional to the amount of target antigen present in the sample. The reporter substrate used to measure the binding event determines the detection mode. A spectrophotometric plate reader is used for colorimetric detection. Chemiluminescent and electrochemiluminescence substrates have been developed which further amplify the signal and can be read on a luminescent reader. The reporter can also be a fluorescent readout where the enzyme step of the assay is replaced with a fluorophore and the readout is then measured using a fluorescent reader. Reagents and protocols necessary to perform an ECL sandwich ELISA are commercially available, including, without exception, MSD sandwich ELISA-ECL detection platform (Meso Scale Discovery, Gaithersburg, Md.).

**[0166]** Thus, in an embodiment, detecting the presence of an antibody-antigen complex comprising an  $\alpha$ -SNAP-25 antibody that selectively binds to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond and a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond can be performed using an immuno-blot analysis, an immunoprecipitation analysis, an ELISA, or a sandwich ELISA. In aspects of this embodiment, the detection is performed using a AU, CL, ECL, or BL immuno-blot analysis, a AU, CL, ECL, BL, or FC immunoprecipitation analysis, a AU, CL, ECL, BL, or FC ELISA, or a AU, CL, ECL, BL, or FC sandwich ELISA.

**[0167]** Aspects of the present disclosure can be practiced in a singleplex or multiplex fashion. An immuno-based method of detecting retargeted endopeptidase activity practiced in a single-plex fashion is one that only detects the presence of an antibody-antigen complex comprising an  $\alpha$ -SNAP-25 antibody and a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. An immuno-based method of detecting retargeted endopeptidase activity practiced in a multiplex fashion is one that concurrently detects the presence of two or more antibody-antigen complexes; one of which is the antibody-antigen complex comprising an  $\alpha$ -SNAP-25 antibody and a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; and the other(s) of which is antibody-antigen complex to a second, third, fourth, etc. different protein. A second protein can be used, e.g., as an internal control to minimize sample to sample variability by normalizing the amount of  $\alpha$ -SNAP-25/SNAP-25 antibody-antigen complex detected to the amount of antibody-antigen complex detected for the second protein. As such, the second protein is usually one that is consistently expressed by the cell, such as a house-keeping protein. Non-limiting examples of a useful second protein, include, e.g., a Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH), Syntaxin, cytokines. Methods of performing an immuno-based assay in a multiplex fashion are described in, e.g., U. B. Nielsen and B. H. Geierstanger, *Multiplexed Sandwich Assays in Microarray Format*, *J. Immunol. Methods*. 290 (1-2): 107-120 (2004); R. Barry and M. Soloviev, *Quantitative Protein Profiling using Antibody Arrays*, *Proteomics*, 4(12): 3717-3726 (2004); M. M. Ling et al., *Multiplexing Molecular Diagnostics and Immunoassays using Emerging*

*Microarray Technologies*, Expert Rev Mol Diagn. 7(1): 87-98 (2007); S. X. Leng et al., *ELISA and Multiplex Technologies for Cytokine Measurement in Inflammation and Aging Research*, J Gerontol A Biol Sci Med Sci. 63(8): 879-884 (2008), each of which is hereby incorporated by reference in its entirety.

[0168] Thus, in one embodiment, an immuno-based method of detecting re-targeted endopeptidase activity practiced in a single-plex fashion by only detecting the presence of an antibody-antigen complex comprising an  $\alpha$ -SNAP-25 antibody and a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. In another embodiment, immuno-based method of detecting re-targeted endopeptidase activity practiced in a multiplex fashion by concurrently detecting the presence of an antibody-antigen complex comprising an  $\alpha$ -SNAP-25 antibody and a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond and at least one other antibody-antigen complex to a protein other than SNAP-25, such as, e.g., GAPDH or Syntaxin.

[0169] Aspects of the present disclosure provide, in part, a method of determining re-targeted endopeptidase immunoresistance. As used herein, the term "re-targeted endopeptidase immunoresistance" means a mammal that does not fully respond to a re-targeted endopeptidase therapy, or shows a reduced beneficial effect of a re-targeted endopeptidase therapy because the immune response of that mammal, either directly or indirectly, reduces the efficacy of the therapy. A non-limiting example of reduced efficacy would be the presence in a mammal of at least one neutralizing a-re-targeted endopeptidase antibody that binds to a re-targeted endopeptidase in a manner that reduces or prevents the specificity or activity of the re-targeted endopeptidase. As used herein, the term "re-targeted endopeptidase therapy" means a treatment, remedy, cure, healing, rehabilitation or any other means of counteracting something undesirable in a mammal requiring neuromodulation using a re-targeted endopeptidase or administering to a mammal one or more controlled doses of a medication, preparation or mixture of a re-targeted endopeptidase that has medicinal, therapeutic, curative, cosmetic, remedial or any other beneficial effect. Re-targeted endopeptidase therapy encompasses, without limitation, the use of any naturally occurring or modified fragment thereof, in any formulation, combined with any carrier or active ingredient and administered by any route of administration.

[0170] Aspects of the present disclosure provide, in part, a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies. As used herein, the term "test sample" refers to any biological matter that contains or potentially contains at least one  $\alpha$ -re-targeted endopeptidase antibody. An  $\alpha$ -re-targeted endopeptidase antibody can be a neutralizing  $\alpha$ -re-targeted endopeptidase antibody or a non-neutralizing  $\alpha$ -re-targeted endopeptidase antibody. As used herein, the term "neutralizing  $\alpha$ -re-targeted endopeptidase antibodies" means any a-re-targeted endopeptidase antibody that will, under physiological conditions, bind to a region of a re-targeted endopeptidase in such a manner as to reduce or prevent the re-targeted endopeptidase from exerting its effect in a re-targeted endopeptidase therapy. As used herein, the term "non-neutralizing a-re-targeted endopeptidase antibodies" means any  $\alpha$ -re-targeted endopeptidase antibody

that will, under physiological conditions, bind to a region of a re-targeted endopeptidase, but not prevent the re-targeted endopeptidase from exerting its effect in a re-targeted endopeptidase therapy. It is envisioned that any and all samples that can contain  $\alpha$ -re-targeted endopeptidase antibodies can be used in this method, including, without limitation, blood, plasma, serum and lymph fluid. In addition, any and all organisms capable of raising  $\alpha$ -re-targeted endopeptidase antibodies against a re-targeted endopeptidase can serve as a source for a sample including, but not limited to, birds and mammals, including mice, rats, goats, sheep, horses, donkeys, cows, primates and humans. Non-limiting examples of specific protocols for blood collection and serum preparation are described in, e.g., Marjorie Schaub Di Lorenzo & Susan King Strasinger, BLOOD COLLECTION IN HEALTH-CARE (F. A. Davis Company, 2001); and Diana Garza & Kathleen Becan-McBride, PHLEBOTOMY HANDBOOK: BLOOD COLLECTION ESSENTIALS (Prentice Hall, 6<sup>th</sup> ed., 2002). These protocols are routine procedures well within the scope of one skilled in the art and from the teaching herein. A test sample can be obtained from an organism prior to exposure to a re-targeted endopeptidase, after a single re-targeted endopeptidase treatment, after multiple re-targeted endopeptidase treatments, before onset of resistance to a re-targeted endopeptidase therapy, or after onset of resistance to a re-targeted endopeptidase therapy.

[0171] Aspects of the present disclosure provide, in part, a control sample. As used herein, the term "control sample" means any sample in which the presence or absence of the test sample is known and includes both negative and positive control samples. With respect to neutralizing  $\alpha$ -re-targeted endopeptidase antibodies, a negative control sample can be obtained from an individual who had never been exposed to re-targeted endopeptidase and may include, without limitation, a sample from the same individual supplying the test sample, but taken before undergoing a re-targeted endopeptidase therapy; a sample taken from a different individual never been exposed to re-targeted endopeptidase; a pooled sample taken from a plurality of different individuals never been exposed to BoNT/A. With respect to neutralizing  $\alpha$ -re-targeted endopeptidase antibodies, a positive control sample can be obtained from an individual manifesting re-targeted endopeptidase immunoresistance and includes, without limitation, individual testing positive in a patient-based testing assays; individual testing positive in an in vivo bioassay; and individual showing hyperimmunity, e.g., a re-targeted endopeptidase vaccinated individual.

[0172] It is further foreseen that  $\alpha$ -re-targeted endopeptidase antibodies can be purified from a sample.  $\alpha$ -Re-targeted endopeptidase antibodies can be purified from a sample, using a variety of procedures including, without limitation, Protein A/G chromatography and affinity chromatography. Non-limiting examples of specific protocols for purifying antibodies from a sample are described in, e.g., ANTIBODIES: A LABORATORY MANUAL (Edward Harlow & David Lane, eds., Cold Spring Harbor Laboratory Press, 2<sup>nd</sup> ed. 1998); USING ANTIBODIES: A LABORATORY MANUAL: PORTABLE PROTOCOL No. I (Edward Harlow & David Lane, Cold Spring Harbor Laboratory Press, 1998); and MOLECULAR CLONING, A LABORATORY MANUAL, supra, (2001), which are hereby incorporated by reference. In addition, non-limiting examples of antibody purification methods as well as well-characterized reagents, conditions and protocols are readily available from

commercial vendors that include, without limitation, Pierce Biotechnology, Inc., Rockford, Ill.; and Zymed Laboratories, Inc., South San Francisco, Calif. These protocols are routine procedures well within the scope of one skilled in the art.

**[0173]** Thus, in an embodiment, a sample comprises blood. In aspect of this embodiment, the sample comprises mouse blood, rat blood, goat blood, sheep blood, horse blood, donkey blood, cow blood, primate blood or human blood. In another embodiment, a sample comprises plasma. In an aspect of this embodiment, a test sample comprises mouse plasma, rat plasma, goat plasma, sheep plasma, horse plasma, donkey plasma, cow plasma, primate plasma or human plasma. In another embodiment, a sample comprises serum. In an aspect of this embodiment, the sample comprises mouse serum, rat serum, goat serum, sheep serum, horse serum, donkey serum, cow serum, primate serum and human serum. In another embodiment, a sample comprises lymph fluid. In aspect of this embodiment, a sample comprises mouse lymph fluid, rat lymph fluid, goat lymph fluid, sheep lymph fluid, horse lymph fluid, donkey lymph fluid, cow lymph fluid, primate lymph fluid or human lymph fluid. In yet another embodiment, a sample is a test sample. In yet another embodiment, a sample is a control sample. In aspects of this embodiment, a control sample is a negative control sample or a positive control sample.

**[0174]** Aspects of the present disclosure provide, in part, comparing the amount of SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond detected in step (d) to the amount of SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond detected in step (e). In an embodiment, the amount of SNAP-25 cleavage product in the test sample is higher as compared to the amount of SNAP-25 cleavage product in the control sample. In an aspect of this embodiment, a higher amount of SNAP-25 cleavage product in the test sample as compared to a positive control sample indicates a reduction in or lack of re-targeted endopeptidase immunoresistance in the mammal. In another aspect of this embodiment, an equivalent amount of SNAP-25 cleavage product in the test sample as compared to a negative control sample indicates a reduction in or lack of re-targeted endopeptidase immunoresistance in the mammal. In another embodiment, the amount of SNAP-25 cleavage product in the test sample is lower as compared to the amount of SNAP-25 cleavage product in the control sample. In an aspect of this embodiment, a lower or equivalent amount of SNAP-25 cleavage product in the test sample as compared to a positive control sample indicates an increase in or presence of re-targeted endopeptidase immunoresistance in the mammal. In another aspect of this embodiment, a lower amount of SNAP-25 cleavage product in the test sample as compared to a negative control sample indicates an increase in or presence of re-targeted endopeptidase immunoresistance in the mammal.

**[0175]** It is envisioned that any and all assay conditions suitable for detecting the present of a neutralizing  $\alpha$ -re-targeted endopeptidase antibody in a sample are useful in the methods disclosed in the present specification, such as, e.g., linear assay conditions and non-linear assay conditions. In an embodiment, the assay conditions are linear. In an aspect of this embodiment, the assay amount of a re-targeted endopeptidase is in excess. In another aspect of this embodiment, the assay amount of a re-targeted endopeptidase is

rate-limiting. In another aspect of this embodiment, the assay amount of a test sample is rate-limiting.

**[0176]** Aspects of the present disclosure can also be described as follows:

**[0177]** 1. A method of detecting re-targeted endopeptidase activity, the method comprising the steps of: a) treating a cell from an established cell line with a sample comprising a re-targeted endopeptidase, wherein the cell from an established cell line is susceptible to re-targeted endopeptidase activity by a re-targeted endopeptidase; b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond; c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of re-targeted endopeptidase activity.

**[0178]** 2. A method of detecting re-targeted endopeptidase activity, the method comprising the steps of: a) treating a cell from an established cell line with a sample comprising a re-targeted endopeptidase, wherein the cell from an established cell line is susceptible to re-targeted endopeptidase activity by a re-targeted endopeptidase; b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond; c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody linked to a solid phase support, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of re-targeted endopeptidase activity.

**[0179]** 3. A method of detecting re-targeted endopeptidase activity, the method comprising the steps of: a) treating a cell from an established cell line with a sample comprising a re-targeted endopeptidase, wherein the cell from an established cell line is susceptible to re-targeted endopeptidase activity by a re-targeted endopeptidase; b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond; c) fixing the SNAP-25 component to a solid phase support; d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of re-targeted endopeptidase activity.

- [0180] 4. A method of detecting retargeted endopeptidase activity, the method comprising the steps of: a) treating a cell from an established cell line with a sample comprising a retargeted endopeptidase, wherein the cell from an established cell line can uptake retargeted endopeptidase; b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of retargeted endopeptidase activity.
- [0181] 5. A method of detecting retargeted endopeptidase activity, the method comprising the steps of: a) treating a cell from an established cell line with a sample comprising a retargeted endopeptidase, wherein the cell from an established cell line can uptake retargeted endopeptidase; b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody linked to a solid phase support, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of retargeted endopeptidase activity.
- [0182] 6. A method of detecting retargeted endopeptidase activity, the method comprising the steps of: a) treating a cell from an established cell line with a sample comprising a retargeted endopeptidase, wherein the cell from an established cell line can uptake retargeted endopeptidase; b) isolating from the treated cell a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; c) fixing the SNAP-25 component to a solid phase support; d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of retargeted endopeptidase activity.
- [0183] 7. A method of determining retargeted endopeptidase immunoresistance in a mammal comprising the steps of: a) adding a retargeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -retargeted endopeptidase neutralizing antibodies; b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line is susceptible to retargeted endopeptidase activity; c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; f) repeating steps b-e with a negative control sample instead of a test sample, the negative control sample comprising a retargeted endopeptidase and a serum known not to contain  $\alpha$ -retargeted endopeptidase neutralizing antibodies; and g) comparing the amount of antibody-antigen complex detected in step e to the amount of antibody-antigen complex detected in step f, wherein detection of a lower amount of antibody-antigen complex detected in step e relative to the amount of antibody-antigen complex detected in step f is indicative of the presence of  $\alpha$ -retargeted endopeptidase neutralizing antibodies.
- [0184] 8. A method of determining retargeted endopeptidase immunoresistance in a mammal comprising the steps of: a) adding a retargeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -retargeted endopeptidase neutralizing antibodies; b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line is susceptible to retargeted endopeptidase activity; c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody linked to a solid phase support, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -retargeted endopeptidase antibody and the SNAP-25 cleavage product; f) repeating steps b-e with a negative control sample instead of a test sample, the negative control sample comprising a re-targeted endopeptidase and a serum known not to contain  $\alpha$ -retargeted endopeptidase neutralizing antibodies; and g) comparing the amount of antibody-antigen complex detected in step e to the amount of antibody-antigen complex detected in step f, wherein detection of a lower amount of antibody-antigen complex detected in step e relative to the amount of antibody-antigen complex detected in step f is indicative of the presence of  $\alpha$ -retargeted endopeptidase neutralizing antibodies.
- [0185] 9. A method of determining retargeted endopeptidase immunoresistance in a mammal comprising the steps of: a) adding a retargeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -retargeted endopeptidase neutralizing antibodies; b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line is susceptible to retargeted

- endopeptidase activity; c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; d) fixing the SNAP-25 component to a solid phase support; e) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; f) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; g) repeating steps b-f with a negative control sample instead of a test sample, the negative control sample comprising a re-targeted endopeptidase and a serum known not to contain  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; and h) comparing the amount of antibody-antigen complex detected in step f to the amount of antibody-antigen complex detected in step g, wherein detection of a lower amount of antibody-antigen complex detected in step f relative to the amount of antibody-antigen complex detected in step g is indicative of the presence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies.
- [0186] 10. A method of determining re-targeted endopeptidase immunoresistance in a mammal comprising the steps of: a) adding a re-targeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line can uptake re-targeted endopeptidase; c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; f) repeating steps b-e with a negative control sample instead of a test sample, the negative control sample comprising a re-targeted endopeptidase and a serum known not to contain  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; and g) comparing the amount of antibody-antigen complex detected in step e to the amount of antibody-antigen complex detected in step f, wherein detection of a lower amount of antibody-antigen complex detected in step e relative to the amount of antibody-antigen complex detected in step f is indicative of the presence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies.
- [0187] 11. A method of determining re-targeted endopeptidase immunoresistance in a mammal comprising the steps of: a) adding a re-targeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line can uptake re-targeted endopeptidase; c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; d) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; e) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; f) repeating steps b-f with a negative control sample instead of a test sample, the negative control sample comprising a re-targeted endopeptidase and a serum known not to contain  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; and g) comparing the amount of antibody-antigen complex detected in step f to the amount of antibody-antigen complex detected in step g, wherein detection of a lower amount of antibody-antigen complex detected in step f relative to the amount of antibody-antigen complex detected in step g is indicative of the presence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies.
- [0188] 12. A method of determining re-targeted endopeptidase immunoresistance in a mammal comprising the steps of: a) adding a re-targeted endopeptidase to a test sample obtained from a mammal being tested for the presence or absence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; b) treating a cell from an established cell line with the test sample, wherein the cell from an established cell line can uptake re-targeted endopeptidase; c) isolating from the treated cells a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond; d) fixing the SNAP-25 component to a solid phase support; e) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; f) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; g) repeating steps b-f with a negative control sample instead of a test sample, the negative control sample comprising a re-targeted endopeptidase and a serum known not to contain  $\alpha$ -re-targeted endopeptidase neutralizing antibodies; and h) comparing the amount of antibody-antigen complex detected in step f to the amount of antibody-antigen complex detected in step g, wherein detection of a lower amount of antibody-antigen complex detected in step f relative to the amount of antibody-antigen complex detected in step g is indicative of the presence of  $\alpha$ -re-targeted endopeptidase neutralizing antibodies.
- [0189] 13. The method of 1-3 and 7-9, wherein the cell is susceptible to re-targeted endopeptidase activity by about 500 nM or less, by about 400 nM or less, by about 300 nM or less, by about 200 nM or less, by about 100 nM or less of a re-targeted endopeptidase.
- [0190] 14. The method of 4-6 and 10-12, wherein the cell can uptake about 500 nM or less, by about 400 nM or less, by about 300 nM or less, by about 200 nM or less, by about 100 nM or less of re-targeted endopeptidase.

- [0191] 15. The method of 1-6, wherein the sample comprises about 100 ng or less, about 10 ng or less, about 1 ng or less, 100 fg or less, 10fg or less, or 1 fg or less of a retargeted endopeptidase.
- [0192] 16. The method of 1-6, wherein the sample comprises about 100 nM or less, about 10 nM or less, about 1 nM or less, about 100 nM or less, about 10 nM or less, about 1 nM or less, about 0.5 nM or less, or about 0.1 nM or less, of a retargeted endopeptidase.
- [0193] 17. The method of 1-12, wherein the presence of an antibody-antigen complex is detected by an immuno-blot analysis, an immunoprecipitation analysis, an ELISA, or a sandwich ELISA.
- [0194] 18. The method of 1-12, wherein the method has a signal-to-noise ratio for the lower asymptote of at least 3:1, at least 5:1, at least 10:1, at least 20:1, at least 50:1, or at least 100:1.
- [0195] 19. The method of 1-12, wherein the method has a signal-to-noise ratio for the higher asymptote of at least 10:1, at least 20:1, at least 50:1, at least 100:1, at least 200:1, at least 300:1, at least 400:1, at least 500:1, or at least 600:1.
- [0196] 20. The method of 1-12, wherein the method can detect the  $EC_{50}$  activity of, e.g., at least 100 ng, at least 50 ng, at least 10 ng, at least 5 ng, at least 100 pg, at least 50 pg, at least 10 pg, at least 5 pg, at least 100 fg, at least 50 fg, at least 10 fg, or at least 5 fg of a retargeted endopeptidase.
- [0197] 21. The method of 1-12, wherein the method can detect the  $EC_{50}$  activity of, e.g., at least 10 nM, at least 5 nM, at least 100 nM, at least 50 nM, at least 10 nM, at least 5 nM, at least 1 nM, at least 0.5 nM, or at least 0.1 nM of a retargeted endopeptidase.
- [0198] 22. The method of 1-12, wherein the method has an LOD of, e.g., 10 pg or less, 9 pg or less, 8 pg or less, 7 pg or less, 6 pg or less, 5 pg or less, 4 pg or less, 3 pg or less, 2 pg or less, 1 pg or less of a retargeted endopeptidase.
- [0199] 23. The method of 1-12, wherein the method has an LOD of, e.g., 100 nM or less, 90 nM or less, 80 nM or less, 70 nM or less, 60 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, or 10 nM or less of a retargeted endopeptidase.
- [0200] 24. The method of 1-12, wherein the method has an LOQ of, e.g., 10 pg or less, 9 pg or less, 8 pg or less, 7 pg or less, 6 pg or less, 5 pg or less, 4 pg or less, 3 pg or less, 2 pg or less, 1 pg or less of a retargeted endopeptidase.
- [0201] 25. The method of 1-12, wherein the method has an LOQ of, e.g., 100 nM or less, 90 nM or less, 80 nM or less, 70 nM or less, 60 nM or less, 50 nM or less, 40 nM or less, 30 nM or less, 20 nM or less, or 10 nM or less of a retargeted endopeptidase.
- [0202] 26. The method of 1-12, wherein the method can distinguish a fully-active retargeted endopeptidase from a partially-active retargeted endopeptidase having 70% or less, 60% or less, 50% or less, 40% or less, 30% or less, 20% or less, or 10% or less the activity of a fully-active retargeted endopeptidase A.
- [0203] 27. The method of 1-12, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the  $P_1$  residue from the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product.
- [0204] 28. The method of 27, wherein the  $\alpha$ -SNAP-25 antibody has an association rate constant for an epitope not comprising a carboxyl-terminus glutamine of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product of less than  $1 \times 10^1 \text{ M}^{-1} \text{ s}^{-1}$  and wherein the  $\alpha$ -SNAP-25 antibody has an equilibrium disassociation constant for the epitope of less than 0.450 nM.
- [0205] 29. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody has a heavy chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:80, and SEQ ID NO:82; and a light chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, and SEQ ID NO:92.
- [0206] 30. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises at least the  $V_H$  CDR1 of SEQ ID NO:93, the  $V_H$  CDR1 of SEQ ID NO:94, the  $V_H$  CDR1 of SEQ ID NO:95, the  $V_H$  CDR1 of SEQ ID NO:118, the  $V_H$  CDR1 of SEQ ID NO:119, or the  $V_H$  CDR1 of SEQ ID NO:120.
- [0207] 32. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises at least the  $V_H$  CDR2 of SEQ ID NO:96, the  $V_H$  CDR2 of SEQ ID NO:97, the  $V_H$  CDR2 of SEQ ID NO:98, the  $V_H$  CDR2 of SEQ ID NO:99, the  $V_H$  CDR2 of SEQ ID NO:121, the  $V_H$  CDR2 of SEQ ID NO:122, or the  $V_H$  CDR2 of SEQ ID NO:123.
- [0208] 33. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises at least the  $V_H$  CDR3 of SEQ ID NO:100, the  $V_H$  CDR3 of SEQ ID NO:101, the  $V_H$  CDR3 of SEQ ID NO:102, or the  $V_H$  CDR3 of SEQ ID NO:124.
- [0209] 34. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises at least the  $V_L$  CDR1 of SEQ ID NO:103, the  $V_L$  CDR1 of SEQ ID NO:104, the  $V_L$  CDR1 of SEQ ID NO:105, the  $V_L$  CDR1 of SEQ ID NO:106, the  $V_L$  CDR1 of SEQ ID NO:107, the  $V_L$  CDR1 of SEQ ID NO:125, the  $V_L$  CDR1 of SEQ ID NO:126, the  $V_L$  CDR1 of SEQ ID NO:127, the  $V_L$  CDR1 of SEQ ID NO:128, or the  $V_L$  CDR1 of SEQ ID NO:129.
- [0210] 35. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises at least the  $V_L$  CDR2 of SEQ ID NO:108, the  $V_L$  CDR2 of SEQ ID NO:109, the  $V_L$  CDR2 of SEQ ID NO:110, the  $V_L$  CDR2 of SEQ ID NO:111, or the  $V_L$  CDR2 of SEQ ID NO:112.
- [0211] 36. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises at least the  $V_L$  CDR3 of SEQ ID NO:113, the  $V_L$  CDR3 of SEQ ID NO:114, the  $V_L$  CDR3 of SEQ ID NO:115, the  $V_L$  CDR3 of SEQ ID NO:116, or the  $V_L$  CDR3 of SEQ ID NO:117.
- [0212] 37. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody comprises a heavy chain variable region comprising SEQ ID NO:93, SEQ ID NO:121 and SEQ ID NO:100; and a light chain variable region comprising SEQ ID NO:105, SEQ ID NO:110 and SEQ ID NO:115.
- [0213] 38. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody selectively binds the SNAP-25 epitope of SEQ ID NO:32, SEQ ID NO:33, SEQ ID

NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:147 or SEQ ID NO:148.

[0214] 39. The method of 27, wherein the isolated  $\alpha$ -SNAP-25 antibody selectively binds the SNAP-25 epitope of SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, or SEQ ID NO:44.

## EXAMPLES

### Example I

#### Screening of Candidate Cell Lines for Endogenous Re-Targeted Endopeptidase Receptor Expression

[0215] The following example illustrates how to identify established cell lines possessing the re-targeted endopeptidase uptake capacity required to develop a cell-based potency assay.

[0216] 1. Growth of Stock Culture of Candidate Cell Lines.

[0217] To grow the cell lines, a suitable density of cells from the cell line being tested were plated in a 162 cm<sup>2</sup> tissue culture flask containing 30 mL of a suitable growth medium (see Table 1), and grown in a 37° C. incubator under 5% or 10% carbon dioxide until cells reached the desired density.

TABLE 1

Media Used in Cell Line Screening.	
Cell Line	Serum Growth Media Composition
SiMa and SiMa clones	RPMI 1640, 10% fetal bovine serum, 1% Penicillin-Streptomycin, 2 mM L-Glutamine
PC12	RPMI 1640, 5% heat-inactivated fetal bovine serum, 10% equine serum, 2 mM GlutaMAX™, 10 mM HEPES, 1 mM sodium pyruvate, 1% Penicillin-Streptomycin
N18	90% DMEM, 10% heat-inactivated fetal bovine serum, 2 mM Glutamine, 2 mM glucose
ND8/34	
NG108-15	
SK-N-DZ	90% DMEM, 10% heat-inactivated fetal bovine serum, 4 mM Glutamine, 4 mM glucose,
SK-N-F1	0.1 mM non-essential amino acids, 1.5 g/L NaHCO <sub>3</sub>
SK-N-SH	
BE(2)-C	EMEM(11090-081, Gibco), Ham's F12 (11765-054, Gibco), 10% heat-inactivated fetal
SK-N-BE(2)	bovine serum, 2 mM Glutamine, 0.1 mM non-essential amino acids,
SH-SY5Y	
ND3, ND7, ND15	DMEM media with 2 mM Glutamine (Invitrogen, Cat #. 11885), 10% Fetal Bovine Serum (Invitrogen, Cat #. 16140), and 1x antibiotic/antimycotic
Neuro-2a	EMEM, 10% heat-inactivated fetal bovine serum, 2 mM Glutamine, 0.1 mM non-essential amino acids, 1.5 g/L NaHCO <sub>3</sub> , 1 mM Sodium pyruvate

[0218] 2. Screening of Cells Expressing Target Receptor on Cell Surface.

[0219] The cell lines were screened for the presence of the desired target receptor using flow cytometry and/or ligand binding assays. Although the examples below used reagents to identify opioid or opioid-like receptor in the plasma membrane, the approaches disclosed below can be used to identify the cognate receptor for any of re-targeted endopeptidase.

[0220] a. Identification of Cell Lines Using Flow Cytometry.

[0221] To identify cells comprising established cell lines that express target receptors for a re-targeted endopeptidase on the cell surface, flow cytometry analysis was conducted. Cells for each candidate cell line were grown as described in Section 1, trypsin-treated, washed in staining buffer com-

prising 1× PBS, 0.5% BSA, and centrifuged at 1200 rpm for 3 minutes. Pelleted cells were resuspended in staining buffer and about 2.0×10<sup>6</sup> cells were transferred to new tubes, two for each receptor tested. To screen for the presence of an opioid or opioid-like receptors, about 2.0-5.0  $\mu$ L of  $\alpha$ -ORL-1 RA14133 (Neuromics, Edina, Minn.),  $\alpha$ -DOR rabbit polyclonal antibody RA10101 (Neuromics, Edina, Minn.),  $\alpha$ -KOR rabbit polyclonal antibody RA10103 (Neuromics, Edina, Minn.), or  $\alpha$ -MOR rabbit polyclonal antibody RA10104 (Neuromics, Edina, Minn.), was added to one tube and the mixture was incubated at 4° C. for 1 hour. The second tube was incubated at 4° C. for 1 hour without any antibodies and served as a negative control. After the antibody incubation, 1.0 mL of staining buffer was added to each tube and centrifuge at 1200 rpm for 3 minutes. The cell pellet was washed once more with 1.0 mL of staining buffer. The cell pellet was resuspend in 200  $\mu$ L of staining buffer, and 2.0  $\mu$ L of goat anti-rabbit IgG FITC antibody was added to each tube and incubated at 4° C. for 1 hour in the dark. After incubation with the secondary antibody, 1.0 mL of staining buffer was added to each tube and centrifuge at 1200 rpm for 3 minutes. The cell pellet was washed once more with 1.0 mL of staining buffer and the pellet resuspend in 500  $\mu$ L of staining buffer. The sample was analyzed using a flow cytometer and the data was displayed as an overlay the anti-receptor antibody staining over the rabbit IgG FITC staining.

[0222] The results indicate that. Of the cell lines tested, ORL-1 was expressed on the cell surface of about 50% of the cells comprising SiMa, SiMa P>33, clone H10, ND7, and SK-N-DZ established cell lines; was expressed on the cell surface of between about 25% to about 50% of the cells comprising SH-SY5Y and ND15 established cell lines; and was expressed on the cell surface of less than about 25% of the cells comprising ND3, ND8, N18, and Neuro-2a established cell lines (Table 2). The results also indicate that KOR was expressed on the cell surface of about 50% of the cells comprising SH-SY5Y and ND7 established cell lines; was expressed on the cell surface of between about 25% to about 50% of the cells comprising SiMa clone H10, SiMa P>33, ND15, and Neuro-2a established cell lines; and was expressed on the cell surface of less than about 25% of the cells comprising ND3, ND8, and N18 established cell lines (Table 2). The results also revealed that MOR was expressed

on the cell surface of about 50% of the cells comprising ND7, ND15, and SiMa P>33 established cell lines; was expressed on the cell surface of between about 25% to about 50% of the cells comprising SH-SY5Y, SiMa clone H10, ND8, and Neuro-2a established cell lines; and was expressed on the cell surface of less than about 25% of the cells comprising ND3 and N18 established cell lines (Table 2). The  $\alpha$ -DOR rabbit polyclonal antibody RA10101 failed to work properly and no useable data was generated.

**[0223]** b. Identification of Cell Lines Using Ligand Binding.

**[0224]** To identify cells comprising established cell lines that express target receptors for a re-targeted endopeptidase on the cell surface, ligand binding analysis was conducted. Cells from the candidate cell lines to be tested were seeded on a black-clear bottom 96-well plate for about 4 hours to promote attachment. To screen for the presence of an opioid or opioid-like receptors, media was then aspirated from each well and replaced with 50  $\mu$ L of ligand solution containing either 0 (untreated control), 0.001 nM, 0.01 nM, 0.1 nM, or 1 nM of FAM-nociceptin (Phoenix Pharmaceuticals, Inc, Burlingame, Calif.); or either 0 (untreated control), 0.001 nM, 0.01 nM, 0.1 nM, or 1 nM of FAM-dynorphin A (Phoenix Pharmaceuticals, Inc, Burlingame, Calif.). Cells were incubated with the ligand solution for 1 hour in the 37° C. incubator under 5% carbon dioxide. The cells were washed to remove unbound ligand by washing the cells three times with 100  $\mu$ L of 1 $\times$  PBS. The plate was scanned on the Typhoon (Ex 488 and Em 520 nm), and then read on the M5 Plate Reader (Ex 495 and Em 520 nm) for RFU signals. The results indicate that cells comprising the SiMa clone Hi 0, SH-SY5Y, and SK-N-DZ established cell lines bound nociceptin, whereas cells comprising the SiMa clone H10 also bound Dynorphin (Table 2).

able density of cells from a stock culture of the cell line being tested was plated into the wells of 24-well tissue culture plates containing 1 mL of an appropriate serum growth medium (Table 1). The cells were grown in a 37° C. incubator under 5% carbon dioxide until cells reached the desired density (approximately 18 to 24 hours). To assess the uptake of an opioid re-targeted endopeptidase, the growth media was aspirated from each well and replaced with either 1) fresh growth media containing no opioid re-targeted endopeptidase (untreated cell line) or 2) fresh growth media containing 30 nM for the nociceptin re-targeted endopeptidase (Noc/A) or 100 nM for the dynorphin re-targeted endopeptidase (Dyn/A) (treated cell line). After an overnight incubation, the cells were washed by aspirating the growth media and rinsing each well with 200  $\mu$ L of 1 $\times$  PBS. To harvest the cells, the 1 $\times$  PBS was aspirated, the cells were lysed by adding 50  $\mu$ L of 2 $\times$  SDS Loading Buffer, the lysate was transferred to a clean test tube and the sample was heated to 95° C. for 5 minutes.

**[0228]** To detect for the presence of both uncleaved SNAP-25 substrate and cleaved SNAP-25 products, an aliquot from each harvested sample was analyzed by Western blot. In this analysis, a 12  $\mu$ L aliquot of the harvested sample was separated by MOPS polyacrylamide gel electrophoresis using NuPAGE® Novex 12% Bis-Tris precast polyacrylamide gels (Invitrogen Inc., Carlsbad, Calif.) under denaturing, reducing conditions. Separated peptides were transferred from the gel onto polyvinylidene fluoride (PVDF) membranes (Invitrogen Inc., Carlsbad, Calif.) by Western blotting using a TRANS-BLOT® SD semi-dry electrophoretic transfer cell apparatus (Bio-Rad Laboratories, Hercules, Calif.). PVDF membranes were blocked by incubating at room temperature for 2 hours in a solution containing Tris-Buffered Saline (TBS) (25 mM 2-amino-2-

TABLE 2

Cell lines Expressing Target Receptors on Cell Surface					
Cell Lines Identified					
Target	Flow Cytometry			Ligand Binding	
	More than 50% Expression	25% to 50% Expression	Less than 50% Expression	Nociceptin	Dynorphin A
ORL-1	AGN P33, SiMa, SiMa clone H10, ND7, SK-N-DZ	SH-SY5Y, ND15	ND3, ND8, N18, Neuro-2a	SiMa clone H10, SH-SY5Y, SK-N-DZ	—
DOR	ND	ND	ND	ND	ND
KOR	SH-SY5Y, ND7	SiMa clone H10, AGN P33, ND15, Neuro-2a	ND3, ND8, N18	—	SiMa clone H10
MOR	ND7, ND15, AGN P33	SH-SY5Y, SiMa clone H10, ND8, Neuro-2a	ND3, N18	ND	ND

**[0225]** Using a similar approach, cell lines comprising cells having cognate receptors for other re-targeted endopeptidases can be identified by FAM-labelling the targeting domain for these endopeptidases and screening cell lines as described above.

**[0226]** 3. Single Dose Screening of Candidate Cell Lines Using Re-Targeted Endopeptidase Molecule.

**[0227]** To determine whether a cell line was able to uptake the appropriate re-targeted endopeptidase molecule, a suit-

hydroxymethyl-1,3-propanediol hydrochloric acid (Tris-HCl) (pH 7.4), 137 mM sodium chloride, 2.7 mM potassium chloride), 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate), 2% Bovine Serum Albumin (BSA), 5% nonfat dry milk. Blocked membranes were incubated at 4° C. for overnight in TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate), 2% BSA, and 5% nonfat dry milk containing either 1) a 1:5,000 dilution of an  $\alpha$ -SNAP-25 mouse monoclonal antibody as the primary antibody (SMI-81; Sternberger Monoclonals Inc., Luther-

ville, Md.); or 2) a 1:5,000 dilution of S9684  $\alpha$ -SNAP-25 rabbit polyclonal antiserum as the primary antibody (Sigma, St. Louis, Mo.). Both  $\alpha$ -SNAP-25 mouse monoclonal and rabbit polyclonal antibodies can detect both the uncleaved SNAP-25 substrate and the SNAP-25 cleavage product,

C, and SK-N-DZ (Table 3); whereas the following cell lines exhibited uptake of 100 nM Dyn/A: N18TG2, Neuro-2a, PC12, and SiMa. Some of these sensitive cell lines were tested with lower doses of compounds and/or with full dose responses.

TABLE 3

Single-Dose Screening of Candidate Cell Lines Using Retargeted Noc/A and Dyn/A				
Cell Line	Description	Source	30 nM Noc/A Uptake	100 nM Dyn/A Uptake
BE(2)-C	Human neuroblastoma	ATCC CRL-2268	Yes	NT
N18TG2	Mouse neuroblastoma	DSMZ ACC 103	Yes	Yes
ND3	Mouse neuroblastoma/primary neonatal rat DRG hybrid	ECACC 92090901	NDA	NDA
ND7/23	Mouse neuroblastoma/primary rat DRG hybrid	ECACC 92090903	No	No
ND8	Mouse neuroblastoma/primary neonatal rat DRG hybrid	ECACC 92090904	NDA	NDA
ND15	Mouse neuroblastoma/primary neonatal rat DRG hybrid	ECACC 92090907	No	No
Neuro-2a	Mouse neuroblastoma	ATCC CCL-131	Yes	Yes
NG108-15	Mouse neuroblastoma/rat glioma hybrid	ECACC 88112302	No	NT
PC12	Rat pheochromocytoma	ATCC CRL-1721	NT	Yes
SH-SY5Y	Human neuroblastoma	ATCC CRL-2266	No	NT
SiMa	Human neuroblastoma	DSMZ ACC 164	Yes	Yes
SK-N-BE(2)-C	Human neuroblastoma	ATCC CRL-2271	Yes	NT
SK-N-DZ	Human neuroblastoma	ATCC CRL-2149	Yes	NT
SK-N-F1	Human neuroblastoma	ATCC CRL-2142	No	NT
SK-N-SH	Human neuroblastoma	ECACC 86012802	No	NT

NT: Not tested.

NDA: No detectable amount of SNAP-25 was detected in this cell line.

allowing for the assessment of overall SNAP-25 expression in each cell line and the percent of SNAP-25 cleaved after re-targeted endopeptidase treatment as a parameter to assess the amount of re-targeted endopeptidase uptake. Primary antibody probed blots were washed three times for 15 minutes each time in TBS, TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Washed membranes were incubated at room temperature for 2 hours in TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate), 2% BSA, and 5% nonfat dry milk containing either 1) a 1:10,000 dilution of goat polyclonal anti-mouse immunoglobulin G, heavy and light chains (IgG, H+L) antibody conjugated to horseradish peroxidase (Zymed, South San Francisco, Calif.) as a secondary antibody; or 2) a 1:10,000 dilution of goat polyclonal anti-rabbit immunoglobulin G, heavy and light chains (IgG, H+L) antibody conjugated to horseradish peroxidase (Zymed, South San Francisco, Calif.) as a secondary antibody. Secondary antibody-probed blots were washed three times for 15 minutes each time in TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Signal detection of the labeled SNAP-25 products were visualized using the ECL Plus™ Western Blot Detection System (GE Healthcare, Amersham Biosciences, Piscataway, N.J.) and the membrane was imaged and the percent of cleaved quantified with a Typhoon 9410 Variable Mode Imager and Imager Analysis software (GE Healthcare, Amersham Biosciences, Piscataway, N.J.). The choice of pixel size (100 to 200 pixels) and PMT voltage settings (350 to 600, normally 400) depended on the individual blot.

**[0229]** Based on the detection of SNAP-25 cleavage product, the following cell lines exhibited uptake of 30 nM Noc/A: BE(2)-C, N18TG2, Neuro-2a, SiMa, SK-N-BE(2)-

**[0230]** Using a similar approach, cell lines comprising cells having cognate receptors for other re-targeted endopeptidases can be assessed for re-targeted endopeptidase uptake.

#### Example II

##### Screening of Candidate Clonal Cell Lines Endogenous Re-Targeted Endopeptidase Receptor Expression

**[0231]** 1. Single Dose Re-Targeted Endopeptidase Screening of Candidate Clonal Cell Lines from a Parental SiMa Cell Line.

**[0232]** Companion patent application Zhu Hong et al., Cell Lines Useful in Immuno-Based Botulinum Toxin Serotype A Activity Assays, U.S. Patent Application Ser. No. 61/160,199 discloses clonal cell lines derived from a parental SiMa cell line that were useful in a BoNT/A potency assay, as described in Ester Fernandez-Salas, et al., Immuno-Based Botulinum Toxin Serotype A Activity Assays, U.S. patent application Ser. No. 12/403,531, each of which is hereby incorporated by reference in its entirety. To determine whether these clonal cell lines were able to uptake the appropriate re-targeted endopeptidase, each was screened using an ECL sandwich ELISA assay.

**[0233]** To prepare a lysate treated with a retargeted endopeptidase, a suitable density of cells from a stock culture of the cell line being tested was plated into the wells of 96-well tissue culture plates containing 100  $\mu$ L of an appropriate serum growth medium (Table 1) overnight. The media from the seeded cells was aspirated from each well and replaced with fresh media containing either 30 nM of a Noc/A re-targeted endopeptidase or 80 nM of a Dyn/A re-targeted

endopeptidase. After a 24 hour incubation, the cells were washed by aspirating the growth media and rinsing each well with 200  $\mu$ L of 1 $\times$  PBS. To harvest the cells, 1 $\times$  PBS was aspirated, the cells lysed by adding 30  $\mu$ L of Lysis Buffer comprising 20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100 to each well, and the plate incubated on a shaker rotating at 500 rpm for 30 minutes at 4° C. The plate was centrifuged at 4000 rpm for 20 minutes at 4° C. to pellet cellular debris and the supernatant was transferred to a capture antibody coated 96-well plate to perform the detection step.

**[0234]** To prepare an  $\alpha$ -SNAP-25197 capture antibody solution, the  $\alpha$ -SNAP-25197 mouse monoclonal antibody contained in the ascites from hybridoma cell line 2E2A6 (Example XI) was purified using a standard Protein A purification protocol.

**[0235]** To prepare an  $\alpha$ -SNAP-25 detection antibody solution,  $\alpha$ -SNAP-25 rabbit polyclonal antibody S9684 (Sigma, St. Louis, MO) was conjugated to Ruthenium(II)-tris-bipyridine-(4-methylsulfonate) NHS ester labeling reagent (Meso Scale Discovery, Gaithersburg, Md.) according to the manufacturer's instructions (Meso Scale Discovery, Gaithersburg, Md.). The conjugation reaction was performed by adding to 30  $\mu$ L of distilled water reconstituted MSD SULFO-TAG™ stock solution to 200  $\mu$ L of 2 mg/mL  $\alpha$ -SNAP-25 polyclonal antibodies and incubating the reaction at room temperature for 2 hours in the dark. The labeled antibodies were purified using a standard spin column protocol and the protein concentration determined using a standard colorimetric protein assay. The absorbance of the  $\alpha$ -SNAP-25 antibody/MSD SULFO-TAG™ conjugate was measured at 455 nm using a spectrophotometer to determine the concentration in moles per liter. The detection antibody solution was stored at 4° C. until needed. Long term storage of unused aliquots was at -20° C.

**[0236]** To prepare an  $\alpha$ -SNAP-25 solid phase support comprising an  $\alpha$ -SNAP-25197 capture antibody, approximately 5  $\mu$ L of the appropriate  $\alpha$ -SNAP-25197 monoclonal antibody solution (20  $\mu$ g/mL in 1 $\times$  PBS) is added to each well of a 96-well MSD High Bind plate and the solution is allowed to air dry in a biological safety cabinet for 2-3 hours in order to liquid evaporate the solution. Blocked plates were sealed and stored at 4° C. until needed.

**[0237]** To detect the presence of a cleaved SNAP-25 product by ECL sandwich ELISA, the capture antibody-bound wells were then blocked by adding 150  $\mu$ L of Blocking Buffer comprising 2% Amersham Blocking Reagent (GE Life Sciences, Piscataway, N.J.) and 10% goat serum (VWR, West Chester, Pa.) at room temperature for 2 hours. The Blocking Buffer was aspirated, 25  $\mu$ L of a lysate from cells treated with re-targeted endopeptidase was added to each well and the plates were incubated at 4° C. for overnight. Plate wells were washed three times by aspirating the cell lysate and rinsing each well three times with 200  $\mu$ L 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, 25  $\mu$ L of 5  $\mu$ g/mL  $\alpha$ -SNAP-25 detection antibody solution comprising 2% Amersham Blocking Reagent in 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate) was added to each well, sealed, and incubated at room temperature for 1 hour with shaking. After  $\alpha$ -SNAP-25 detection antibody incubation, the wells were washed three times with 200  $\mu$ L 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing 150  $\mu$ L of 1 $\times$  Read Buffer (Meso

Scale Discovery, Gaithersburg, MD) was added to each well and the plates were read using a SECTOR™ Imager 6000 Image Reader (Meso Scale Discovery, Gaithersburg, Md.). The raw data was collected using the ECL imager.

**[0238]** The results indicate that the parental SiMa cell line as well as clonal cell line H10 showed good uptake of the Noc/A retargeted endopeptidase (Table 4). In addition, these results reveal that many cell lines exhibited uptake of the Dyn/A retargeted endopeptidase (Table 4). Three clonal cell lines (1E11, AF4, and DC4) showed good uptake of the Dyn/A retargeted endopeptidase; eleven clonal cell lines (1E3, 2D2, 2D6, 3D8, 5010, 5F3, BB10, BFB, CG8, CG10, and DE7) exhibited moderate uptake of the Dyn/A retargeted endopeptidase; and (3B8, 2B9, CE6, YB8, 408, 2F5, AC9, CD6, DD10, YF5) showed minimal uptake of the Dyn/A retargeted endopeptidase. Some of these candidate cell lines were tested in a full dose-response assay with the corresponding retargeted endopeptidase.

TABLE 4

Single-Dose Screening of Candidate Clonal Cell Lines Using Retargeted Noc/A and Dyn/A		
Cell Line	30 nM Noc/A Uptake	80 nM Dyn/A Uptake
AGN P33	+++	NT
A10	-	NT
D11	-	NT
H1	-	-
H10	+++	-
1D4	NT	-
2E4	NT	-
3D5	NT	-
3G10	NT	-
4D3	NT	-
BB3	NT	-
CC11	NT	-
DF5	NT	-
YB7	NT	-
BE3	NT	-
4B5	NT	-
2B9	NT	+
2F5	NT	+
3B8	NT	+
4C8	NT	+
AC9	NT	+
CD6	NT	+
CE6	NT	+
DD10	NT	+
YB8	NT	+
YF5	NT	+
1E3	NT	++
2D2	NT	++
2D6	NT	++
3D8	NT	++
5C10	NT	++
5F3	NT	++
BFB	NT	++
BB10	NT	++
CG8	NT	++
CG10	NT	++
DE7	NT	++
1E11	NT	+++
AF4	NT	+++
DC4	NT	+++

NT: Not Tested  
 -: no uptake;  
 +: minimal uptake;  
 ++: moderate uptake;  
 +++: good uptake

**[0239]** 2. Full Dose Response Screening of Candidate Cell Lines.

**[0240]** Established cell lines identified above, were subsequently evaluated using a full dose response of the appropriate re-targeted endopeptidase. Cells from the different cell lines were plated in 96-well plates and exposed to various concentrations of Noc/A (0, 0.14 nM, 0.4 nM, 1.23 nM, 3.7 nM, 11.1 nM, 33.3 nM, and 100 nM) or of Dyn/A (0.017 nM,

sensitivity of each one of the cell lines tested, and EC<sub>50</sub> value was calculated for each cell line. The values for the Noc/A retargeted endopeptidase are summarized in Table 5. Full dose response of retargeted endopeptidase Dyn/A were only performed in PC12 and clone AF4. In both cases the assay did not reach an upper asymptote and an EC<sub>50</sub> could not be calculated. The lower dose that produced a signal for the AF4 clone was 12 nM for both cell lines.

TABLE 5

Full-Dose Screening of Candidate Cell Lines Using Retargeted Noc/A and Dyn/A				
Cell Line	Description	Source	EC <sub>50</sub> Noc/A Uptake	EC <sub>50</sub> Dyn/A Uptake
AGN P33	Human neuroblastoma	—	5-10 nM	NT
BE(2)-C	Human neuroblastoma	ATCC CRL-2268	NT	NT
N18TG2	Mouse neuroblastoma	DSMZ ACC 103	NT	NT
N18	Mouse Neuroblastoma	ECACC 88112301	>100 nM	NT
ND3	Mouse neuroblastoma/primary neonatal rat DRG hybrid	ECACC 92090901	NDA	NT
ND7/23	Mouse neuroblastoma/primary rat DRG hybrid	ECACC 92090903	>100 nM	NT
ND8	Mouse neuroblastoma/primary neonatal rat DRG hybrid	ECACC 92090904	NDA	NT
ND15	Mouse neuroblastoma/primary neonatal rat DRG hybrid	ECACC 92090907	>100 nM	NT
Neuro-2a	Mouse neuroblastoma	ATCC CCL-131	30 nM	NT
NG108-15	Mouse neuroblastoma/rat glioma hybrid	ECACC 88112302	NT	NT
PC12	Rat pheochromocytoma	ATCC CRL-1721	NT	>1000 nM
SH-SY5Y	Human neuroblastoma	ATCC CRL-2266	NT	NT
SiMa	Human neuroblastoma	DSMZ ACC 164	30 nM	NT
SiMa clone AF4	Human neuroblastoma	—	NT	>300 nM
SiMa clone H1	Human neuroblastoma	—	>100 nM	NT
SiMa clone H10	Human neuroblastoma	—	20 nM	NT
SK-N-BE(2)-C	Human neuroblastoma	ATCC CRL-2271	NT	NT
SK-N-DZ	Human neuroblastoma	ATCC CRL-2149	0.5-2 nM	NT
SK-N-F1	Human neuroblastoma	ATCC CRL-2142	>100 nM	NT
SK-N-SH	Human neuroblastoma	ECACC 86012802	>100 nM	NT

NT: Not tested.

NDA: No detectable amount of SNAP-25 was detected in this cell line.

0.05 nM, 0.15 nM, 0.45 nM, 1.4 nM, 4.1 nM, 12 nM, 37 nM, 111 nM, 333 nM, and 1000 nM) for 24 hours. Retargeted endopeptidase-containing medium was then removed and replaced with fresh complete medium. Plates were incubated another 24 hours under 5% CO<sub>2</sub> at 37° C. to allow for the cleavage of SNAP-25. Cells were lysed in the lysis buffer (Table 5) and plates centrifuged to eliminate debris. The lysates were used either in a Western blot assay or in a sandwich ELISA.

**[0241]** For the Western blot analysis, samples were assayed for the presence of both the intact SNAP-25 and the SNAP-25 cleavage product as described in Example I.

**[0242]** For the sandwich ELISA, ELISA plates coated with 2E2A6 monoclonal antibody were blocked with 150  $\mu$ L Blocking Buffer at room temperature for 2 hours. After blocking buffer was removed, 25  $\mu$ L of cell lysate was added to each well and the plates were incubated at 4° C. for 2 hours. Plates were washed three times with PBS-T and 25  $\mu$ L of SULFO-TAG NHS-Ester labeled detection anti-SNAP25 pAb antibody at 5  $\mu$ g/mL in 2% blocking reagent in PBS-T was added to the bottom corner of wells. The plates were sealed and shaken at room temperature for 1 hour, followed by three washes with PBS-T. After washes were completed, 150  $\mu$ L of 1 $\times$  Read Buffer per well was added and the plate was read in the SI6000 Image reader. To determine the

**[0243]** Using a similar approach, clonal cell lines comprising cells having cognate receptors for other re-targeted endopeptidases can be screened and assessed for retargeted endopeptidase uptake.

### Example III

#### Evaluation of Growth Conditions on Retargeted Endopeptidase Uptake in Candidate Cell Lines

**[0244]** The following example illustrates how to determine culture conditions, growth and differentiation, for established cell lines that maximize re-targeted endopeptidase uptake.

**[0245]** 1. Effects of Cell Differentiation and Trophic Factors on Re-Targeted Endopeptidase Uptake of Candidate Cell Lines.

**[0246]** To determine whether cell differentiation or the present of trophic factors in the growth media improved re-targeted endopeptidase uptake, cell lines exhibiting good uptake of Noc/A were tested with different media compositions. A suitable density of cells from a stock culture of the SiMa P>30 cell line being tested was plated into the wells of 96-well tissue culture plates containing 100  $\mu$ L of a serum-free medium containing RPM11640, 1% Penicillin-Streptomycin, 2 mM L-Glutamine, supplemented with B27, and N2, or 100  $\mu$ L of a serum-free medium containing

RPM11640, 1% Penicillin-Streptomycin, 2 mM L-Glutamine, supplemented with B27, N2, and NGF (Nerve Growth Factor, 100 ng/mL). These cells were incubated in a 37° C. incubator under 5% carbon dioxide until the cells differentiated, as assessed by standard and routine morphological criteria, such as growth arrest and neurite extension (approximately 1 to 2 days). As a control, a suitable density of cells from a stock culture of the cell line being tested was plated into the wells of 96-well tissue culture plates containing 100 µl of an appropriate growth medium (Table 1) without or with NGF (100 ng/mL). These undifferentiated control cells were grown in a 37° C. incubator under 5% carbon dioxide until cells reach the desired density (approximately 18 to 24 hours). The media from both differentiated and undifferentiated control cultures was aspirated from each well and replaced with fresh media containing either 0 (untreated sample) or to various concentrations of Noc/A (0.14, 0.4, 1.23, 3.7, 11.1, 33.3, and 100 nM). After a 24 hours treatment, the cells were washed and incubated for 24 hours in retargeted endopeptidase-free media in order to increase the amount of SNAP25197 produced. Cells were then washed and harvested for the ECL sandwich ELISA assay as described in Example II.

[0247] Effects of trophic factors were also tested on the SK-N-DZ cell line. SK-N-DZ cells were plated on a poly-D-lysine coated 96-well plate at 25,000 cells per well in eight different SM media (Table 6) for 72 hours. Cells were treated in the same eight media with Noc/A at doses of 0, 0.3 nM, 3 nM, and 30 nM. After a 24 hour treatment, the cells were washed and incubated for 24 hours in retargeted endopeptidase-free media in order to increase the amount of SNAP-25197 cleavage product produced. Cells were then washed and harvested for the Western blot assay as described in Example 1.

[0248] Differentiation did not have an effect on Noc/A uptake in the SiMa>P30 cell line while it seemed to improve uptake in the SK-N-DZ cell line. The basal media had a significant effect on Noc/A uptake in the SK-N-DZ cell line with RPM11640 comprising trophic factors N2 and B27 being the best combination. The presence of NGF in the media did not seem to improve uptake in the two cell lines tested.

[0249] Using a similar approach, the growth and differentiation conditions for clonal cell lines comprising cells having cognate receptors for other re-targeted endopeptidases can be assessed.

#### Example IV

##### Development of Established Cell Lines Expressing Exogenous Re-Targeted Endopeptidase Receptors

[0250] The following example illustrates how to make an established cell line expressing an exogenous receptor for a re-targeted endopeptidase.

[0251] 1. Transfection of Target Receptor Into Cells Comprising a Candidate Cell Line.

[0252] The re-targeted endopeptidase Noc/A comprises the nociceptin targeting domain which is the natural ligand of Opioid Receptor Like-1 (ORL-1). To obtain an expression construct comprising an open reading frame for an ORL-1, the expression construct pReceiver-M02/ORL-1 was obtained from GeneCopoeia (GeneCopoeia, Germantown, Md.).

[0253] Alternatively, a polynucleotide molecule based on an ORL-1 amino acid sequence (e.g., the amino acid sequences SEQ ID NO:25 or SEQ ID NO:26) can be synthesized using standard procedures (BlueHeron® Biotechnology, Bothell, Wash.). Oligonucleotides of 20 to 50 bases in length are synthesized using standard phosphoramidite synthesis. These oligonucleotides will be hybridized into double stranded duplexes that are ligated together to assemble the full-length polynucleotide molecule. This polynucleotide molecule will be cloned using standard molecular biology methods into a pUCBHB1 vector at the SmaI site to generate pUCBHB1/ORL-1. The synthesized polynucleotide molecule is verified by sequencing using Big Dye Terminator™ Chemistry 3.1 (Applied Biosystems, Foster City, Calif.) and an ABI 3100 sequencer (Applied Biosystems, Foster City, Calif.). If desired, an expression optimized polynucleotide molecule based on an ORL-1 amino acid sequence (e.g., the amino acid sequences SEQ ID NO:25 or SEQ ID NO:26) can be synthesized in order to improve expression in an Escherichia coli strain. The poly-

TABLE 6

Effects of Trophic Factors and Cell Differentiation on Noc/A Uptake of Candidate Cell Lines.

		EC <sub>50</sub> Noc/A Uptake	
		AGN P33	SK-N-DZ
Undifferentiated	Differentiated		
DMEM, 10% FBS	—	NT	>30 nM
DMEM, 10% FBS, N2, B27	—	NT	3 nM
DMEM, 10% FBS, N2, B27, NGF	—	NT	3 nM
DMEM, 10% FBS, N2, B27, RA	—	NT	>30 nM
RPM11640, 10% FBS	—	NT	10 nM
RPM11640, 10% FBS, N2, B27	—	7.2 nM	1 nM
RPM11640, 10% FBS, N2, B27, NGF	—	9.1 nM	1 nM
RPM11640, 10% FBS, N2, B27, RA	—	NT	10 nM
—	RPM11640, N2, B27	10.2 nM	1 nM
—	RPM11640, N2, B27, NGF	9.8 nM	0.6 nM

NGF: Nerve Growth Factor;

RA: Retinoic Acid

NT: Not tested

nucleotide molecule encoding the ORL-1 can be modified to 1) contain synonymous codons typically present in native polynucleotide molecules of an *Escherichia coli* strain; 2) contain a G+C content that more closely matches the average G+C content of native polynucleotide molecules found in an *Escherichia coli* strain; 3) reduce polymononucleotide regions found within the polynucleotide molecule; and/or 4) eliminate internal regulatory or structural sites found within the polynucleotide molecule, see, e.g., Lance E. Steward et al., Optimizing Expression of Active Botulinum Toxin Type A, U.S. Patent Publication 2008/0057575 (Mar. 6, 2008); and Lance E. Steward et al., Optimizing Expression of Active Botulinum Toxin Type E, U.S. Patent Publication 2008/0138893 (Jun. 12, 2008). Once sequence optimization is complete, oligonucleotides of 20 to 50 bases in length are synthesized using standard phosphoramidite synthesis. These oligonucleotides are hybridized into double stranded duplexes that are ligated together to assemble the full-length polynucleotide molecule. This polynucleotide molecule is cloned using standard molecular biology methods into a pUCBHB1 vector at the SmaI site to generate pUCBHB1/ORL-1. The synthesized polynucleotide molecule is verified by DNA sequencing. If so desired, expression optimization to a different organism, such as, e.g., a yeast strain, an insect cell-line or a mammalian cell line, can be done, see, e.g., Steward, U.S. Patent Publication 2008/0057575, supra, (2008); and Steward, U.S. Patent Publication 2008/0138893, supra, (2008). Exemplary polynucleotide molecules encoding an ORL-1 include SEQ ID NO:61 and SEQ ID NO:62.

**[0254]** To construct an expression contract encoding an ORL-1, a pUCBHB 1/ORL-1 construct will be digested with restriction endonucleases that 1) excised the polynucleotide molecule encoding the open reading frame of ORL-1; and 2) enabled this polynucleotide molecule to be operably-linked to a pcDNA3 vector (Invitrogen, Inc., Carlsbad, Calif.). This insert will be subcloned using a T4 DNA ligase procedure into a pcDNA3 vector that will be digested with appropriate restriction endonucleases to yield pcDNA3/ORL-1. The ligation mixture will be transformed into electro-competent *E. coli* BL21(DE3) cells (Edge Biosystems, Gaithersburg, Md.) using an electroporation method, and the cells will be plated on 1.5% Luria-Bertani agar plates (pH 7.0) containing 50 µg/mL of ampicillin, and will be placed in a 37° C. incubator for overnight growth. Bacteria containing expression constructs will be identified as ampicillin resistant colonies. Candidate constructs will be isolated using an alkaline lysis plasmid mini-preparation procedure and will be analyzed by restriction endonuclease digest mapping to determine the presence and orientation of the insert. This cloning strategy will yielded a pcDNA3 expression construct comprising the polynucleotide molecule encoding an ORL-1.

**[0255]** The re-targeted endopeptidase Dyn/A comprises the dynorphin targeting domain which is the natural ligand of κ-opioid receptor (KOR). To obtain an expression construct comprising an open reading frame for an ORL-1, the expression construct pReceiver-M02/KOR-1 was obtained from GeneCopoeia (GeneCopoeia, Germantown, Md.). Alternatively, expression constructs encoding the KOR can be synthesized and subcloned using an approach similar to the one described above to produce the expression construct pcDNA3.1/KOR. Exemplary KOR amino acid sequences

include SEQ ID NO:29 and SEQ ID NO:30; exemplary polynucleotide molecules encoding a KOR include SEQ ID NO:65 and SEQ ID NO:66.

**[0256]** Similar cloning strategies can be used to make expression constructs encoding other re-target endopeptidase receptors, such as, e.g., pcDNA3.1/DOR or pcDNA3.1/MOR, pcDNA3.1/Galanin receptor 1, pcDNA3.1/Galanin receptor 2, or pcDNA3.1/Galanin receptor 3. Exemplary DOR amino acid sequences include SEQ ID NO:27 and SEQ ID NO:28; exemplary MOR amino acid sequences include SEQ ID NO:31 exemplary Galanin receptor 1 amino acid sequences include SEQ ID NO:136, SEQ ID NO:137, and SEQ ID NO:138; exemplary Galanin receptor 2 amino acid sequences include SEQ ID NO:139; and exemplary Galanin receptor 3 amino acid sequences include SEQ ID NO:140. Exemplary polynucleotide molecules encoding a DOR include SEQ ID NO:63 and SEQ ID NO:64; exemplary polynucleotide molecules encoding a MOR include SEQ ID NO:67; exemplary polynucleotide molecules encoding a Galanin receptor 1 include SEQ ID NO:141, SEQ ID NO:142, and SEQ ID NO:143; exemplary polynucleotide molecules encoding a Galanin receptor 2 include SEQ ID NO:144; and exemplary polynucleotide molecules encoding a Galanin receptor 3 include SEQ ID NO:145.

**[0257]** To introduce an expression construct encoding a re-targeted endopeptidase receptor, cell lines were transfected with an expression construct encoding a re-targeted endopeptidase receptor. To transfect a cell line with an opioid or opioid-like receptor, cells from a candidate cell line were seeded at a density of  $1 \times 10^7$  cells in a 1175 Collagen IV coated flask and grown in a 37° C. incubator under 5% carbon dioxide until cells reach the desired density. A 4.2 mL transfection solution is prepared by adding 4 mL of OPTI-MEM Reduced Serum Medium containing 200 µL of LipofectAmine 2000 (Invitrogen, Carlsbad, Calif.) incubated at room temperature for 5 minutes to 4 mL of OPTI-MEM Reduced Serum Medium containing 20 µg of pReceiver-M02/ORL-1 or 20 µg of pReceiver-M02/KOR-1. This transfection was incubated at room temperature for approximately 20 minutes. The media was replaced with 8 mL of fresh serum-free and antibiotic-free media and the transfection solution was added to the cells. The cells were then incubated in a 37° C. incubator under 5% carbon dioxide for approximately 16-18 hours. Transfection media was replaced with fresh growth media and incubate cells in a 37° C. incubator under 5% carbon dioxide. After 24 hours, the growth media were replaced with fresh growth media containing the antibiotic G418 at 1 mg/mL in the culture medium (selection medium) and cells were incubated for 7 days. The selection medium was changed every week for a total of 4 weeks (about 90% cells died and were removed during the weekly media changes).

**[0258]** Candidate cell lines transfected with the ORL-1 receptor included SiMa>P30, ND15, ND7, NG108-T15 and SK-N-DZ cell lines. Candidate cell lines transfected with the KOR-1 receptor included SiMa, SiMa>P30, ND15, ND7, NG108-115 and SK-N-DZ cell lines. Transfected NG108-115 cells did not survive selection in G418.

**[0259]** 2. Single Dose and Dose Response Screening of Stable Transfected Cell Lines Using Re-Targeted Endopeptidase Molecules.

**[0260]** Cells from the transfected and selected candidate cell lines from the previous section were plated on 96-well poly-D-lysine or Collagen IV coated plates at  $1 \times 10^5$  cells/

well in medium RPM11640 containing N2 and B27 supplements, and NGF (50-100 ng/ml) for 20±4 hours before compound treatment. Then the cells stably transfected with the ORL-1 receptor were treated with re-targeted endopeptidase Noc/A at 30 nM in the same medium for 24±2 hours, except for the SK-N-DZ cell line that was treated at 10 nM. Cells were lysed in 120 µL lysis buffer, and 20 µL of the lysate was mixed with the 2× SDS buffer for the Western Blot assay that was performed as detailed in Example I. All of the cells lines displayed increase uptake of the re-targeted Noc/A compound when transfected with the ORL-1 receptor (Table 7).

TABLE 7

Cell lines stably transfected with ORL-1 tested with re-targeted endopeptidase Noc/A			
% Cleaved SNAP25 at 30 nM			
Cell line	Description	Non-transfected	Transfected
AGN P33	Human Neuroblastoma	20%	40%
SK-N-DZ	Human Neuroblastoma	25% @ 10 nM	40% @ 10 nM
ND7	Mouse Neuroblastoma and rat DRG cells hybrid	10%	42%
ND15	Mouse Neuroblastoma and rat DRG cells hybrid	8%	20%
NG108-T15	Mouse neuroblastoma/rat glioma hybrid	No cells survive	No cells survive

**[0261]** Cells from the transfected and selected candidate cell lines from the previous section were plated on 96-well poly-D-lysine or Collagen IV coated plates at 1×10<sup>5</sup> cells/well in medium RPM11640 containing 10% FBS and N2 and B27 supplements for 20±4 hours before compound treatment. The cells stably transfected with the KOR-1 receptor were treated with re-targeted endopeptidase Dyn/A at 100 nM in the same medium for 24±2 hours. Cells were lysed in 120 µL lysis buffer, and 20 µL of the lysate was mixed with the 2× SDS buffer for the Western Blot assay that was performed as detailed in Example I. All of the cells lines displayed increase uptake of the re-targeted Dyn/A compound when transfected with the human KOR-1 receptor.

**[0262]** 3. Selection of Stably Transfected Clonal Cell Lines Exhibiting High Sensitivity by Serial Dilution

**[0263]** The following example illustrates how to identify clonal cells from a stably transfected established cell line that are susceptible to re-targeted endopeptidase action or have re-targeted endopeptidase uptake capacity.

**[0264]** For single cell cloning of the selected cells described above, the limited-dilution cell line cloning method was employed. Cells were trypsinized, counted, diluted to achieve 0.5-1 cell per 100 µL, and plated on selection media in five poly-D-lysine coated 96-well plates at 100 µL per well. Cells were incubated for more than 2 weeks until colonies formed on the bottom of the well. The positive colonies originating from single cells were marked. Pictures of single-cell derived clones were taken using a microscope camera. Cells from wells with single clones were grown for an additional week and transferred into 24 well plates about 4 weeks after cloning started.

**[0265]** For clone selection, the main parameter used to screen for positive clones was the highest amount of SNAP-25 cleavage obtained after Noc/A or Dyn/A treatment measured using the Western Blot analysis with the antibody that

recognizes both intact and cleaved SNAP-25. Clones over-expressing ORL-1 were tested with 10 nM and 30 nM re-targeted endopeptidase Noc/A overnight as soon as enough cells became available (Table 8). Clones over-expressing KOR-1 were tested with 100 nM re-targeted endopeptidase Dyn/A overnight (Table 9). In addition, clones over-expressing KOR-1 were tested on the Dynorphin binding assay as described in Example I.

TABLE 8

Single-Dose Screening of Candidate Clonal Cell Lines Stably transfected with ORL-1 Using re-targeted endopeptidase Noc/A				
Cell Line	Clone number	10 nM Noc/A Uptake	30 nM Noc/A Uptake	Second screening @ 1 nM (% cleaved)
AGN P33	1	+	+	28%
AGN P33	2	++	+++	50%
AGN P33	3	-	+	NT
AGN P33	4	ND	ND	NT
AGN P33	5	-	+	31%
AGN P33	6	++	+++	60%
AGN P33	7	+	+	14%
AGN P33	8	+	+	NT
AGN P33	9	+	+	38%
AGN P33	10	+	++	29%
AGN P33	11	+	+	NT
AGN P33	12	+	+	27%
ND7	1C11	NT	++	NT
ND7	2F3	NT	-	NT
ND7	1D10	NT	-	NT
ND7	1F9	NT	-	NT
ND7	1G10	NT	-	NT
ND7	2D8	NT	-	NT
ND7	2E2	NT	-	NT
ND7	4B7	NT	+++	NT
ND7	3C11	NT	-	NT
ND7	3C3	NT	+	NT
ND7	3E8	NT	-	NT
ND7	3E11	NT	-	NT
ND7	2G3	NT	-	NT
ND7	4D5	NT	+	NT
ND7	4D8	NT	+	NT
ND7	4C8	NT	-	NT
ND7	4C9	NT	+++	NT
ND7	4E8	NT	+	NT
ND7	2E6	NT	++	NT
ND7	4F4	NT	+++	NT
ND7	5D6	NT	-	NT
ND7	5G3	NT	-	NT
ND7	4D5	NT	++	NT
ND15	1C10	NT	+	NT
ND15	1F10	NT	++	NT
ND15	2D8	NT	++	NT
ND15	2E11	NT	-	NT
ND15	2F4	NT	++	NT
ND15	2F10	NT	++	NT
ND15	2F11	NT	-	NT
ND15	3C4	NT	+	NT
ND15	3C7	NT	++	NT
ND15	3E8	NT	+++	NT
ND15	4C8	NT	+	NT
ND15	4D8	NT	+	NT
SK-N-DZ	#2	-	-	NT
SK-N-DZ	#4	-	-	NT
SK-N-DZ	#5	+++	++	NT
SK-N-DZ	#6	NT	++	NT
SK-N-DZ	#7	+	NT	NT
SK-N-DZ	#8	-	NT	NT
SK-N-DZ	#9	+	NT	NT
SK-N-DZ	#10	-	NT	NT
SK-N-DZ	#11	+	+++	NT
SK-N-DZ	#12	-	NT	NT
SK-N-DZ	#14	++	NT	NT
SK-N-DZ	#16	-	NT	NT

TABLE 8-continued

Single-Dose Screening of Candidate Clonal Cell Lines Stably transfected with ORL-1 Using re-targeted endopeptidase Noc/A				
Cell Line	Clone number	10 nM Noc/A Uptake	30 nM Noc/A Uptake	Second screening @ 1 nM (% cleaved)
SK-N-DZ	#17	+	+++	NT
SK-N-DZ	#19	+	+++	NT
SK-N-DZ	#20	-	NT	NT
SK-N-DZ	#23	NT	++	NT
SK-N-DZ	#25	-	NT	NT
SK-N-DZ	#26	-	++	NT
SK-N-DZ	#27	+	NT	NT
SK-N-DZ	#28	++	+	NT
SK-N-DZ	#30	++	NT	NT
SK-N-DZ	#31	-	NT	NT
SK-N-DZ	#32	++	++	NT
SK-N-DZ	#33	+	NT	NT
SK-N-DZ	#34	+++	ND	NT
SK-N-DZ	#35	+	++	NT
SK-N-DZ	#36	-	NT	NT
SK-N-DZ	#37	+++	++	NT
SK-N-DZ	#42	-	NT	NT
SK-N-DZ	#43	+	++	NT

ND: Not Determined;

NT: Not Tested.

-: no uptake;

+: minimal uptake;

++: moderate uptake;

+++: good uptake

TABLE 9

Single-Dose Screening of Candidate Clonal Cell Lines Stably transfected with KOR-1 Using re-targeted endopeptidase Dyn/A				
Cell Line	Clone number	100 nM Dyn/A Uptake	100 nM Dyn binding	Selected future testing
SiMa	2	-	-	No
SiMa	6	+	+	No
SiMa	8	+	+	No
SiMa	12	+++	++	Yes
SiMa	14	++	++	No
SiMa	20	+	++	No
SiMa	25	++	++	No
AGN P33	1	+++	+	Yes
AGN P33	3	++	+	No
AGN P33	5	++	+	Yes
AGN P33	6	++	+	No
AGN P33	7	+++	+	Yes
AGN P33	8	++	+	Yes
AGN P33	9	+++	+	Yes
AGN P33	10	+++	+	Yes
AGN P33	11	++	+	No
AGN P33	12	+++	+	Yes
AGN P33	14	+	+	No
AGN P33	16	++	+	No
AGN P33	17	+++	+	Yes
AGN P33	21	+	++	No
ND7	A1	+	+	No
ND7	A2	-	-	No
ND7	A3	-	-	No
ND7	A4	-	-	No
ND7	A5	-	-	No
ND7	A6	-	-	No
ND7	A7	-	-	No
ND7	A8	-	-	No
ND7	A9	-	-	No
ND7	A10	-	-	No
ND7	A11	-	-	No
ND7	A12	+++	+++	Yes
ND7	B1	-	-	No

TABLE 9-continued

Single-Dose Screening of Candidate Clonal Cell Lines Stably transfected with KOR-1 Using re-targeted endopeptidase Dyn/A				
Cell Line	Clone number	100 nM Dyn/A Uptake	100 nM Dyn binding	Selected future testing
ND7	B2	-	-	No
ND7	B3	-	-	No
ND7	B4	-	-	No
ND7	B5	+	+	Yes
ND7	B6	-	-	No
ND7	B7	-	-	No
ND7	B8	-	-	No
ND7	B9	-	-	No
ND7	B10	-	-	No
ND7	B11	-	-	No
ND7	B12	-	-	No
ND7	C1	-	-	No
ND7	C2	-	-	No
ND7	C3	-	-	No
ND7	C4	-	-	No
ND7	C5	-	-	No
ND7	C6	+	+	No
ND7	C7	-	-	No
ND7	C8	-	-	No
ND7	C9	-	-	No
ND7	C10	-	-	No
ND7	C11	-	-	No
ND7	C12	-	-	No
ND7	D1	-	-	No
ND7	D2	-	-	No
ND7	D3	-	-	No
ND7	D4	-	-	No
ND7	D5	-	-	No
ND7	D6	++	++	Yes
ND7	D7	++	++	Yes
ND7	D8	-	-	No
ND7	D9	-	-	No
ND7	D10	-	-	No
ND7	D11	-	-	No
ND7	D12	-	-	No
ND7	E1	-	-	No
ND7	E2	-	-	No
ND7	E3	-	-	No
ND7	E4	-	-	No
ND7	E5	-	-	No
ND7	E6	-	-	No
ND7	E7	-	-	No
ND7	E8	-	-	No
ND7	E9	-	-	No
ND7	E10	-	-	No
ND7	E11	-	-	No
ND7	E12	++	++	Yes
ND7	F1	-	-	No
ND7	F2	-	-	No
ND7	F3	-	-	No
ND7	F4	-	-	No
ND15	A1	-	-	No
ND15	A2	-	-	No
ND15	A3	+	-	No
ND15	A4	+	-	No
ND15	A5	-	-	No
ND15	A6	++	-	No
ND15	A7	++	-	No
ND15	A8	++	-	No
ND15	A9	+	-	No
ND15	A10	+	-	No
ND15	A11	-	-	No
ND15	A12	-	-	No
ND15	B1	-	-	No
ND15	B2	++	-	No
ND15	B3	-	-	No
ND15	B4	-	-	No
ND15	B5	+++	-	Yes
ND15	B6	+	-	No
ND15	B7	-	-	No
ND15	B8	-	-	No

TABLE 9-continued

Single-Dose Screening of Candidate Clonal Cell Lines Stably transfected with KOR-1 Using re-targeted endopeptidase Dyn/A				
Cell Line	Clone number	100 nM Dyn/A Uptake	100 nM Dyn binding	Selected future testing
ND15	B9	-	-	No
ND15	B10	-	-	No
ND15	B11	-	-	No
ND15	B12	-	-	No
ND15	C1	-	-	No
ND15	C2	+++	+	Yes
ND15	C3	-	-	No
ND15	C4	-	-	No
ND15	C5	+	NT	No
ND15	C6	+++	NT	Yes
SK-N-DZ	#11	NT	NT	ND

ND: Not Determined;  
 NT: Not Tested.  
 -: no uptake;  
 +: minimal uptake;  
 ++: moderate uptake;  
 +++: good uptake

**[0266]** 4. Dose Response Screening of Stably Transfected Clonal Cell Lines Using Re-Targeted Endopeptidase.

**[0267]** Candidate stably transfected clonal cell lines from section 3 showing good uptake of the re-targeted endopeptidase Noc/A were tested in full dose response experiments to determine sensitivity and efficacy towards the re-targeted endopeptidase Noc/A. Cells were plated on 96-well poly-D-lysine or Collagen IV coated plates at  $1 \times 10^5$  cells/well in medium RPM11640 containing N2 and B27 supplements, and NGF (50-100 ng/mL) for 20 ± 4 hours before compound treatment. Cells from the parental AGN P33 cell line and the ND7 clonal cell lines were treated with 0, 0.14 nM, 0.4 nM, 1.23 nM, 3.7 nM, 11.1 nM, 33.3 nM, and 100 nM of Noc/A in the same medium for 24 hours plus 24 hours incubation in re-targeted endopeptidase free media to allow for cleavage of SNAP-25. Cells from the parental AGN P33 cell line were also treated with 0, 0.03 nM, 0.08 nM, 0.24 nM, 0.74 nM, 2.22 nM, 6.67 nM, and 20 nM of Noc/A in the same medium for 24 hours plus 24 hours incubation in re-targeted endopeptidase free media to allow for cleavage of SNAP-25. Media was removed and cells washed and lysed for the ECL sandwich ELISA assay as detailed in Example II. The data from the AGN P33 parental and the clonal cell lines stably transfected with the ORL-1 receptor is summarized in Table 10. Clones #2 and #6 demonstrated better sensitivity and efficacy towards the re-targeted endopeptidase Noc/A than the parental cell line. Moreover, the increased sensitivity of the new clonal cell lines allowed the use of lower concentrations for the dose-response confirming the new clonal cell lines are more sensitive.

TABLE 10

Summary table of signal-to-noise ratios (S/N) and EC50 values of the three most sensitive clones overexpressing ORL-1 in the AGN P33 background cell line.				
	Parental	Clone 2	Clone 6	Clone 8
S/N Ratio 0.03 nM/BK		41	26	1.8
S/N Ratio 20 nM/BK		259	522	33.1
S/N Ratio 0.14 nM/BK				
S/N Ratio 100 nM/BK				
EC <sub>50</sub> (nM)	6.8 ± 1.1	0.6 ± 0.1	0.7 ± 0.07	0.3 ± 0.2

**[0268]** The data from the ND7 parental and the clonal cell lines stably transfected with the ORL-1 receptor is summarized in Table 11. All the clones tested demonstrated improved sensitivity and efficacy towards the re-targeted endopeptidase Noc/A than the parental cell line ND7. Clones 4B7, 1 E6, and 1C11 were the most sensitive with EC<sub>50</sub> values lower than 10 pM.

TABLE 11

Summary table of signal-to-noise ratios (S/N) and EC <sub>50</sub> values of the six most sensitive clones overexpressing ORL-1 in the ND7 background cell line.							
	Parental	1C11	4B7	4C9	4F4	1E6	3E9
S/N Ratio 0.14 nM/BK	1.7	9.3	11.1	5.3	3.6	5.8	5.1
S/N Ratio 100 nM/BK	53	217	243	126	169	123	121
EC <sub>50</sub> (nM)	>50	8.6 ± 2	5.7 ± 0.5	33 ± 11	24 ± 5	6.7 ± 1	>30 nM

**[0269]** Table 12 summarizes the results obtained with the generation and testing of clonal cell lines overexpressing the ORL-1 receptor in different cell backgrounds.

TABLE 12

Summary of clonal cell lines overexpressing the Human ORL-1 receptor tested with Noc/A			
Background Cell Line	Species	Stable cell lines tested with full dose of Noc/A	EC <sub>50</sub> (nM)
AGN P33	Human neuroblastoma	Three	0.6-2.5
ND7	Rat neuroblastoma and DRG hybrid	Six	3.7-8
SK-N-DZ	Human neuroblastoma	None (seven stable clones selected for further studies)	N/A

Example V

Development of Clonal Cell Lines from a Parental SK-N-DZ Cell Line

**[0270]** The following example illustrates how to identify clonal cells from a parental established cell line that are susceptible to re-targeted endopeptidase inhibition of exocytosis or have re-targeted endopeptidase uptake capacity.

**[0271]** 1. Isolation of Clonal Cell Lines.

**[0272]** During characterization of the SK-N-DZ cell line, it was discovered that the cells comprising this established cell line comprised at least five different cellular phenotypes. To determine whether any one of these phenotypically-distinct cell types was responsible for the susceptibility of this cell line to re-targeted endopeptidase inhibition of exocytosis, two different limited-dilution screens were conducted to obtain single colony isolates for each phenotypically-distinct cell type.

**[0273]** A suitable density of cells from a SK-N-DZ stock were grown in DMEM, 10% Fetal Bovine Serum (heat-inactivated), 0.1 mM Non-Essential Amino-Acids, 10 mM HEPES, 1 mM Sodium Pyruvate, 100 U/ml Penicillin, 100 µg/ml Streptomycin, contained in a T175 Collagen IV coated flask. After the second passage, the cells were

trypsin-treated to produce a cell suspension and the cell concentration was determined. About  $4.0 \times 10^6$  cells from this cell suspension was transferred into a 50 mL tube and the cells were dissociated into single cells by repeated vigorous expulsion through an 18.5 gauge needle using a 10 mL syringe. Cells from this disassociated single-cell suspension were then diluted to a concentration of  $0.2 \times 10^6$  cells/mL by adding 15 mL of fresh growth medium, and 2.5  $\mu$ L of this dilution was added to 50 mL of fresh growth medium to obtain a concentration of 10 cells/mL. From this final dilution stock, 100  $\mu$ L of growth medium was added to each well of a 96-well Collagen IV coated plates and the cells were grown undisturbed in a 37° C. incubator under 5% carbon dioxide for four weeks. Four 96-well plates were setup for analysis. After four weeks, each well was microscopically examined to identify growing single colonies, and for each colony identified 100  $\mu$ L of fresh growth medium was added to each well and the cells were grown undisturbed in a 37° C. incubator under 5% carbon dioxide for two weeks. After two additional weeks of growth, the growing single colonies were trypsin-treated and transferred to a new 96-well plate for continued growth. Once colonies grew to about 1,000 cells, based on visual inspection, the cells were trypsin-treated and each cell-suspension was transferred into a new well from a 24-well Collagen IV-coated plate. The cells were grown in a 37° C. incubator under 5% carbon dioxide with fresh growth medium being replenished every 2-3 days, if needed. The cells were grown until the culture reached approximately 60% confluence or greater, at which point the cells were trypsin-treated and each cell-suspension was transferred into a 25 cm<sup>2</sup> Collagen IV-coated flask, based on the confluence of the cells in the 24-well plate. The cells were grown in a 37° C. incubator under 5% carbon dioxide with fresh growth medium being replenished every 2-3 days, if needed. Once the cells in the flask reached 70-80% confluence, they were frozen and stored in liquid nitrogen until the clonal cell lines were tested to determine their susceptibility to Noc/A inhibition of exocytosis. Of the 384 colony isolates initially setup from both screens, 24 clonal cell lines were selected based on viability and growth criteria and expanded for subsequent screening procedures. Of those, 12 fast growing cell lines were identified.

**[0274]** 2. Primary Screen for Re-targeted Endopeptidase Activity Susceptibility of Cells from a Clonal Cell Line Using a Re-Targeted Endopeptidase.

**[0275]** To determine whether cells from a clonal cell line were susceptible to re-targeted endopeptidase Noc/A activity, a primary screen was conducted using an immuno-based method for determining endopeptidase activity.

**[0276]** Thirteen SK-N-DZ clones (#3, #4, #5, #8, #9, #10, #13, #15, #16, #17, #18, #22, and #23) plus SK-N-DZ parental cells were plated on a 96-well plate (unknown cell number per well) in EMEM, 10% FBS, 1x B27, and 1xN2 and incubated overnight. Cells were treated with 1 nM Noc/A for 24 hours. Cells were lysed with 100  $\mu$ L of lysis buffer for 20 minutes and centrifuged at 4000 rpm for 20 minutes. Fifty microliters of 2x SDS Sample buffer were added to 50  $\mu$ L of cell lysate and heated at 95° C. for 5 minutes. Ten microliters of protein sample were loaded per lane on 12% NuPage gels and a Western Blot assay was performed as described in Example I. Evaluation of the total SNAP-25 and the cleaved SNAP-25 demonstrated that clones #3, #8, #15, and #22 were at least as good as parental cells for Noc/A uptake. Full dose response treatment and

analysis with the ECL sandwich ELISA assay was conducted after the cells were scale up.

**[0277]** 3. Secondary Response Screening of Clonal Cell Lines Using Re-targeted Endopeptidase Molecule.

**[0278]** To determine whether cells from a clonal cell line were susceptible to re-targeted endopeptidase Noc/A activity, a secondary screen was conducted using an immuno-based method for determining endopeptidase activity.

**[0279]** To further compare these SK-N-DZ cloned cell lines, the ECL sandwich ELISA assay was carried out. Five clones (#3, #9, #15, #16, #22) plus SK-N-DZ parental cells were plated on one 96-well Poly-D-lysine coated plate per cell line at 25,000 cells per well in RPMI 1640, 10% FBS, 1x B27, and 1x N2 media (no NGF) over the weekend. Cells were treated with Noc/A at doses from 0 to 20 nM (0, 0.03, 0.08, 0.24, 0.74, 2.22, 6.67, 20 nM) for 24 hours. The cleaved SNAP-25197 was quantified with the ECL ELISA assay as detailed in Example I.

**[0280]** Table 13 shows the EC<sub>50</sub> values and signal to noise for the five clones and their parental cell line. Three clones named #3, #9, and #15 generated lower EC<sub>50</sub> values (<1 nM) and clone #16 and #22 generated similar EC<sub>50</sub> values when compared to the parental cell line (~2 nM). However, the total signals from cleaved SNAP25 were higher in clones #3, #22, and the parental cells. Clones #9, #16, and #15 had lower total signals when compared to the rest of the cell lines.

TABLE 13

Summary table of signal-to-noise ratios (S/N) and EC <sub>50</sub> values of the five clones obtained from SK-N-DZ cells by limited-dilution cloning.						
	Parental	3	9	19	16	22
S/N Ratio	2	3	2	2	2	3
0.03 nM/BK						
S/N Ratio	19	27	12	8	14	20
20 nM/BK						
EC <sub>50</sub> (nM)	2.6 ± 1.5	0.8 ± 0.07	0.7 ± 0.04	0.6 ± 0.1	2.2 ± 0.8	1.9 ± 0.6

**[0281]** Conditions for Noc/A treatment on SK-N-DZ clones were optimized and an assay was run comparing clones #3, #15, and #22, and the parental heterogeneous SK-N-DZ cell line. Table 14 shows the result of the comparison and demonstrated that assay optimization has greatly improved the signal to noise for the assay. Clones #3 and #22 were selected for further assay development as they possess excellent sensitivity and efficacy.

TABLE 14

Summary table of signal-to-noise ratios (S/N) and EC <sub>50</sub> values of three clones obtained from SK-N-DZ cells using optimized conditions.				
	Parental	3	15	22
S/N Ratio 0.03 nM/BK	15	8	5	10
S/N Ratio 20 nM/BK	107	89	33	60
EC <sub>50</sub> (nM)	0.6 ± 0.2	0.9 ± 0.2	0.6 ± 0.1	0.4 ± 0.09

#### Example VI

##### Characterization and Comparison of Clonal Cell Lines for Re-Targeted Endopeptidase Uptake

**[0282]** The following example illustrates how to characterize and compare clonal cell lines originated either from an

established cell line comprising a heterogeneous population or by transfection of the target receptor and subsequent cloning of the cell line.

**[0283]** To evaluate the specificity or selectivity of re-targeted endopeptidase uptake, non-specific uptake assays were performed using of a re-targeted endopeptidase lacking the targeting domain. For opioid re-targeted endopeptidase, cells from the AGN P33 clone #6 cell line (comprising cells stably transformed with an expression construct encoding an ORL-1 receptor) and the SK-N-DZ clonal cell lines #3 and #22 (comprising cells that express endogenous ORL-1 receptor) were seeded at 150,000 cells per well on a 96-well Poly-D-lysine coated plate in RPMI 1640 serum free media containing N2 and B27 supplements, and NGF (50 ng/mL) and incubated for 20 ±4 hours at 37° C. in a 5% CO<sub>2</sub> incubator before compound treatment. Cells were treated with 8-doses of Noc/A ranging from 0-20 nM or 0-40 nM and/or eight doses of LH<sub>N</sub>/A ranging from 0 to 400 nM or 0 to 40 nM in the same medium for 22 hours. Media was removed and cells washed, lysed, and centrifuged to eliminate debris in preparation for a sandwich ELISA assay. An ELISA plate coated with 2E2A6 monoclonal antibody was blocked with 150 µL Blocking Buffer at room temperature for 1 hour. After blocking buffer was removed, 30 µL of cell lysate was added to each well and the plate was incubated at 4° C. for 2 hours. Plates were washed three times with PBS-T and 30 µL of SULFO-TAG NHS-Ester labeled detection α-SNAP25 polyclonal antibodies at 5 µg/mL in 2% blocking reagent in PBS-T was added to the bottom corner of wells. The plate was sealed and shaken at room temperature for 1 hour, followed by three washes with PBS-T. After washes were completed, 150 µL of 1× Read Buffer per well was added and the plate was read in the S16000 Image reader. The results comparing Noc/A uptake relative to the negative control LH<sub>N</sub>/A are shown in Table 15 and Table 16. These results indicate that there was good separation between Noc/A and LH<sub>N</sub>/A uptake in both cell lines demonstrating specific uptake of Noc/A.

TABLE 15

Non-specific uptake for SK-N-DZ clone #3. Summary of four independent experiments		
nM	% non-specific uptake	SEM (standard error of the mean)
0	2	0.5
1	6	0.5
2	8	0.5
5	10	1
15	19	0.9
44	33	1.5
133	65	2.4
400	93	2.3

TABLE 16

Non-specific uptake for hORL-1 #6 cells. Summary of three independent experiments		
nM	% non-specific uptake	SEM (standard error of the mean)
0	1	0.2
1	2	0.2
2	3	0.6
5	3	0.3
15	8	1.3
44	12	1.9
133	22	3.0
400	32	3.0

**[0284]** Table 17 summarizes the results for the characterization and comparison of the three cell lines. SK-N-DZ clonal cell lines #3 and #22 possess a sensitivity identical to the primary eDRG and an excellent signal-to-noise to develop a robust assay for re-targeted endopeptidase Noc/A. The AGN P33 clonal cell line #6 is also an excellent candidate with low non-specific uptake and adequate sensitivity.

TABLE 17

Parameter	SK-N-DZ clone 3	SK-N-DZ clone 22	AGN P33 clone 6	eDRG
Cell line species	Human Clonal	Human Clonal	Human Clonal	Rat primary
Cell Receptor	Human ORL1	Human ORL1	Human ORL1	Rat ORL1
Expression	endogenous	endogenous	transfected	endogenous
Dynamic Range	0.03 to 20 nM dose response	0.03 to 20 nM dose response	0.04 to 40 nM dose response	0.17 to 20 nM dose response
Sensitivity (EC <sub>50</sub> )	EC <sub>50</sub> = 0.75 ± 0.1 (N = 10)	EC <sub>50</sub> = 0.8 ± 0.2 (N = 9)	EC <sub>50</sub> = 2.4 ± 0.2 (N = 21)	EC <sub>50</sub> = 0.8 ± 0.15 (N = 6)
ULOQ	20 nM	20 nM	20 nM	10-20 nM
S/N ULOQ/background	98 ± 15 (N = 10)	86 ± 17 (N = 9)	385 ± 32 (N = 19)	~300
S/N LLOQ/background	12 ± 2 (N = 11)	10 ± 2 (N = 9)	29 ± 7 (N = 18)	N/A
Specificity vs LH <sub>N</sub> /A	≥2 logs (N = 4)	≥2 logs (N = 4)	≥2 logs (N = 3)	N/A
SNAP-25 Expression	Endogenous	Endogenous	Endogenous	Endogenous
Competition with Nociceptin Var.	Full competition (n = 4)	Full competition (n = 4)	Partial competition (n = 4)	N/A
Inhibition by Ab Anti-nociceptin	Full competition (n = 4)	Full competition (n = 4)	Full competition (n = 4)	N/A
Inhibition by Anti-868 Ab	Partial competition (N = 3)	Partial competition (N = 3)	Partial competition (N = 3)	N/A

**[0285]** To evaluate the sensitivity of re-targeted endopeptidase uptake, ligand saturation binding assays were performed. The interaction of most ligands with their binding sites can be characterized in terms of binding affinity (NIH Assay Guidance). In general, high affinity binding involves a longer residence time for the ligand at its receptor binding site than in the case of low affinity binding. The dissociation constant is commonly used to describe the affinity between a ligand (L) (such as a drug) and a protein (P) i.e. how tightly a ligand binds to a particular protein. An equilibrium saturation binding experiment measures total and non-specific binding (NSB) at various radioligand concentrations. The equilibrium dissociation constant or affinity for the radioligand,  $K_d$ , and the maximal number of receptor binding sites,  $B_{max}$ , can be calculated from specific binding (total—NSB) using non-linear regression analysis. The  $K_d$  for the specific binding can be calculated using a one-site binding hyperbola nonlinear regression analysis (i.e. GraphPad Prism) as shown in the equation below, where  $B_{max}$  is the maximal number of binding sites (pmol/mg, or pmol/cells, or sites/cells), and  $K_d$  (nM, pM, etc.) is the concentration of radioligand required to reach half-maximal binding:

$$Bound = \frac{B_{max} \times [L]}{[L] + K_d}$$

**[0286]** For opioid re-targeted endopeptidase, cells from the AGN P33 clone #6 cell line (comprising cells stably transformed with an expression construct encoding an ORL-1 receptor), the SK-N-DZ parental cell line, and the SK-N-DZ clonal cell lines #3, #15, and #22 (comprising cells that express endogenous ORL-1 receptor) were plated at 200,000 cells per well on a 48-well Poly-D-lysine coated plate in RPMI 1640 serum free media containing  $1 \times N2$  and  $1 \times B27$  supplements and incubated overnight at  $37^\circ C$ . in a  $5\% CO_2$  incubator. Media was removed and cells and  $150 \mu L$  of Tris binding buffer was added to wells used to assess total binding and  $100 \mu L$  of Tris binding buffer was added to well used to assess non-specific binding. About  $50 \mu L$  of  $4 \times$  final concentration cold nociceptin ( $2.5 \mu M$  to SK-N-DZ cell lines and  $1 \mu M$  to AGN P33 clonal cell line #6) was added to the non-specific binding wells, and  $50 \mu L$  of  $4 \times$  final concentrations of  $^3H$ -nociceptin ( $0 nM$ ,  $0.05 nM$ ,  $0.1 nM$ ,  $0.2 nM$ ,  $0.4 nM$ ,  $0.8 nM$ ,  $1.6 nM$ ,  $3.1 nM$ ,  $6.3 nM$ ,  $12.5 nM$ ,  $25 nM$ , and  $50 nM$  to the SK-N-DZ cell lines and  $0$ ,  $0.01 nM$ ,  $0.02 nM$ ,  $0.039 nM$ ,  $0.078 nM$ ,  $0.156 nM$ ,  $0.313 nM$ ,  $0.625 nM$ ,  $1.25 nM$ ,  $2.5 nM$ ,  $5.0 nM$ , and  $10 nM$  to the AGN P33 clonal cell line #6) was added to both the total binding wells and the non-specific binding wells to a final volume of  $200 \mu L$ . After incubation at  $37^\circ C$ . for 30 minutes, the wells were washed twice in  $0.5 mL$  of cold Washing Buffer. Cells were then denatured in  $200 \mu L$   $2 N NaOH$  and transferred to  $20 mL$  scintillation vials containing  $5 mL$  of scintillation fluid. Raw data were used to plot the dose-response graphs and calculate the  $K_d$  for each sample. The raw data obtained were transferred to SigmaPlot v10.0 and One Site Saturation fit was used to define the dose-response curves under the equation category of Ligand Binding. Graphical reports were generated and contained the following parameters:  $R^2$  (correlation coefficient),  $B_{max}$ , and  $K_d \pm SE$  (Coefficient  $\pm$  standard error). Graphs of total binding, specific binding, and non-specific binding were obtained on the assay performed on the SK-N-DZ clonal cell lines #3, #15,

and #22, and AGN P33 clonal cell line #6 cells. SK-N-DZ clonal cell lines #3 and #22 produced a concentration-dependent and saturable binding of  $^3H$ -nociceptin. Under the same experimental conditions, SK-N-DZ clonal cell line #15 produced a dose-dependent response of  $^3H$ -nociceptin, but not saturated at the highest dose of  $50 nM$ . Compared to the SK-N-DZ cell lines expressing endogenous ORL-1, cells from the AGN P33 clonal cell line #6 had significantly higher affinity binding to  $^3H$ -nociceptin (highest dose was  $10 nM$  versus  $50 nM$  in SK-N-DZ) with low non-specificity binding.

**[0287]** The saturation binding curves of SK-N-DZ clonal cell lines #3, #22, #15 and AGN P33 clonal cell line #6 were used to estimate  $K_d$  and  $B_{max}$  values from three independent binding experiments per cell line performed in three different days. The rank order of these four cell lines is: AGN P33 clonal cell line #6 ( $K_d=1.86 nM$  and  $B_{max}=2.9 fmol/cell$ )  $>$ SK-N-DZ clonal cell line #3 ( $K_d=14 nM$  and  $B_{max}=0.6 fmol/cell$ ) SK-N-DZ clonal cell line #22 ( $K_d=17 nM$  and  $B_{max}=0.6 fmol/cell$ )  $\gg$ SK-N-DZ clonal cell line #15 ( $K_d > 50 nM$ ). To get a saturated dose-response for SK-N-DZ clonal cell line #15, a higher dose range of  $^3H$ -nociceptin needs to be used. Table 16 summarizes the data regarding the characterization of the specific plasma membrane nociceptin-binding sites in three SK-N-DZ clonal cell lines, #3, #15, and #22, and AGN P33 clonal cell line #6 stable cell lines. The data showed the following: 1) a high affinity site with very low non-specific binding ( $K_d$ ,  $1.8 nM$ , and  $B_{max}$   $2.9 fmol$  per cell) in the AGN P33 clonal cell line #6; 2) nociceptin-binding can be performed on SK-N-DZ native cells expressing endogenous receptor; 3) AGN P33 clonal cell line #6 had about 10-fold higher affinity to nociceptin than the SK-N-DZ cell lines; 4) as seen in the cell-based potency assay, SK-N-DZ clonal cell lines #3 and #22 ( $K_d$   $14-17 nM$ ,  $B_{max}$   $0.6 fmol$  per cell) had more receptor sites per cell than SK-N-DZ clonal cell line #15 (not saturable under the same dose range).

TABLE 18

Summary of $^3H$ -nociceptin saturation binding assay for four leading cell lines (n = 3 independent experiments)		
Cell Lines	$K_d$ (nM $\pm$ SD)	$B_{max}$ (fmol/cell)
SK-N-DZ #3	$14 \pm 1.6$	0.59
SK-N-DZ #15	$>50$	ND
SK-N-DZ #22	$16.7 \pm 1.1$	0.58
AGN P33 clonal cell line #6	$1.86 \pm 0.1$	2.89

**[0288]** To evaluate the sensitivity of re-targeted endopeptidase uptake, the amount of re-targeted endopeptidase receptor expressed at the mRNA level was assessed using RT-PCR. The amount of receptor expressed in the cells is an important aspect of the characterization of the cell line being used for testing and it is related to the sensitivity to re-targeted endopeptidases. The amount of expressed re-targeted endopeptidase receptor can also be a tool for screening other potential cell lines and to eliminate cell lines that do not express the target receptor. One method of measuring receptor expression is to quantify the amount of re-targeted endopeptidase receptor mRNA using real time PCR (RT-PCR).

**[0289]** For opioid re-targeted endopeptidase, RNA was isolated from cells of a non-transfected parental SiMa cell

line, cells from the AGN P33 clonal cell line #6, cells from the parental SK-N-DZ cell line, and cells from the SK-N-DZ clonal cell lines #3 and #22 grown in either serum free media or media with serum. The mRNA was converted to cDNA and the ORL-1 was amplified and measured real time to determine the relative amount present in each cell line using the following oligonucleotide primers for ORL-1: forward 5'-CACTCGGCTGGTGGTGG-3' (SEQ ID NO:148) and reverse 5'-AATGGCCACGGCAGTCTCGC-3' (SEQ ID NO:149). The DNA is quantified by using SYBR® green which fluoresces relative to the amount of double stranded DNA (PCR product) present in the reaction. Plotting the amount of fluorescence vs. number of cycles gives a logistic curve for each reaction. The faster a reaction reaches the linear phase of the curve the more ORL-1 receptor cDNA there is in the reaction. A control RT reaction where no enzyme is added was used to determine if there is contamination. Since there is no RT enzyme present in this reaction, no cDNA will be produced. A PCR product cannot be produced using a RNA template, so if a PCR curve appears in the -RT reaction, the only possibility is genomic DNA contamination. In the -RT reactions, no PCR plots appears, confirming there was minimal genomic DNA contamination (data not shown). Table 18 lists the cell lines with their CT value. The CT is the number of PCR cycles it took for that corresponding PCR reaction to produce a signal above a set threshold. The amount of ORL-1 receptor mRNA in a cell line can be compared to another by looking at their corresponding CT values. According to the CT values, cells from the AGN P33 clonal cell line #6 had much more ORL-1 mRNA than cells from the parental SiMa cell line in serum free media (Ave CT: 28.6 vs. 17.3) and in media with serum (Ave CT: 26.1 vs. 16.5). Also, there appears to be minimal difference in mRNA obtained from cells at passage 6 vs. passage 16 in the AGN P33 clonal cell line #6. Also, there are minimal differences in CT values and plots in the parental SK-N-DZ cell line vs. clonal cell line #3 and #22. This conclusion is true in cells grown in media with serum and serum free media and reflects the similarity of these cell lines observed in the cell-based potency assay for Noc/A.

TABLE 19

Average CT Values for ORL-1 Expression in Cell Lines		
Media	Cell line	CT average
Serum Free Media	SiMa Parental p26	28.6
	SiMa hORL-1 clone #6 p6	17.3
	SiMa hORL-1 clone #6 p16	17.3
Complete Media	SiMa Parental p26	26.1
	SiMa hORL-1 clone #6 p6	16.4
	SiMa hORL-1 clone #6 p16	16.6
Serum Free Media	SK-N-DZ	26.3
	SK-N-DZ clone #3	25.9
	SK-N-DZ clone #22	26.6
Complete Media	SK-N-DZ	26.2
	SK-N-DZ clone #3	25.8
	SK-N-DZ clone #22	26.4

## Example VII

Development of  $\alpha$ -SNAP-25 Monoclonal  
Antibodies that Selectively Bind a SNAP-25  
Epitope Having a Free Carboxyl-terminus at the P<sub>1</sub>  
Residue of the BoNT/A Cleavage Site Scissile  
Bond

**[0290]** The following example illustrates how to make  $\alpha$ -SNAP-25 monoclonal antibodies that can selectively bind to a SNAP-25 epitope having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

**[0291]** 1. Generation of  $\alpha$ -SNAP-25 Monoclonal Antibodies.

**[0292]** To develop monoclonal  $\alpha$ -SNAP-25 antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, the 13-residue peptide CDSNKTRIDEANQ-cooH (SEQ ID NO:38) was designed as a SNAP-25 cleavage product antigen. This peptide comprises a flexible linker region and a N-terminal Cysteine residue for conjugation to KLH and amino acids 186-197 of human SNAP-25 (SEQ ID NO:5) with a carboxylated C-terminal glutamine (SEQ ID NO:38). The generation of monoclonal antibodies to well-chosen, unique peptide sequences provides control over epitope specificity, allowing the identification of a particular subpopulation of protein among a pool of closely related isoforms. Blast searches revealed that this peptide has high homology only to SNAP-25 and almost no possible cross-reactivity with other proteins in neuronal cells. The sequence was also carefully scrutinized by utilizing computer algorithms to determine hydrophathy index, protein surface probability, regions of flexibility, and favorable secondary structure, followed by proper orientation and presentation of the chosen peptide sequence. The peptide was synthesized and conjugated to Keyhole Limpet Hemocyanin (KLH) to increase immunogenicity. Six Balb/c mice were immunized with this peptide, and after three immunizations in about eight weeks, the mice were bled for testing. The blood was allowed to clot by incubating at 4° C. for 60 minutes. The clotted blood was centrifuged at 10,000×g at 4° C. for 10 minutes to pellet the cellular debris. The resulting serum sample was dispensed into 50  $\mu$ l aliquots and stored at -20° C. until needed.

**[0293]** A similar strategy based on other SNAP-25 antigens disclosed in the present specification is used to develop  $\alpha$ -SNAP-25 monoclonal antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond. For example, the SNAP-25 antigen of SEQ ID NO:45 can be conjugated to KLH instead of the SNAP-25 antigen of SEQ ID NO:38. As another example, the amino acids 186-197 of human SNAP-25 from the SNAP-25 antigen of SEQ ID NO:38 can be replaced with SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, or SEQ ID NO:44.

**[0294]** 2. Screening for the Presence of  $\alpha$ -SNAP-25 Monoclonal Antibodies.

**[0295]** To determine the presence of an  $\alpha$ -SNAP-25 monoclonal antibody that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, comparative ELISA and cell-based cleavage assay were performed using the extracted mouse serum. For comparative ELISA, two fusion

proteins were constructed: BirA-HisTag®-SNAP-25<sub>134-197</sub> of SEQ ID NO:48 and the BirA-HisTag®-SNAP-25<sub>134-206</sub> of SEQ ID NO:49. BirA-HisTag®-SNAP-25<sub>134-197</sub> comprised a naturally-biotinylated 16 amino acid BirA peptide of SEQ ID NO:50 amino-terminally linked to a SNAP-25 peptide comprising amino acids 134-197 of SEQ ID NO:5. BirA-HisTag®-SNAP-25<sub>134-206</sub> comprised a naturally-biotinylated 16 amino acid BirA peptide of SEQ ID NO:50 amino-terminally linked to a SNAP-25 peptide comprising amino acids 134-206 of SEQ ID NO:5. These two substrates were suspended in 1× PBS at a concentration of 10 µg/mL BirA-HisTag®-SNAP-25<sub>134-197</sub> and the BirA-HisTag®-SNAP-25<sub>134-206</sub>. The BirA-HisTag®-SNAP-25<sub>134-197</sub> and the BirA-HisTag®-SNAP-25<sub>134-206</sub> were coated onto separate plates by adding approximately 100 µl of the appropriate Substrate Solution and incubating the plates at room temperature for one hour. Washed plates were incubated at 37° C. for one hour in 0.5% BSA in 1× TBS containing a 1:10 to 1:100 dilution of an antibody-containing serum derived from one of the six immunized mice (Mouse 1, Mouse 2, Mouse 3, Mouse 4, Mouse 5, and Mouse 6). Primary antibody probed plates were washed four times for 5 minutes each time in 200 µl TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Washed plates were incubated at 37° C. for 1 hour in 1× TBS containing a 1:10,000 dilution of goat polyclonal anti-mouse IgG antibody conjugated to Horseradish peroxidase (Pierce Biotechnology, Rockford, Ill.) as a secondary antibody. Secondary antibody-probed plates were washed four times in 200 µl TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Chromogenic detection of the labeled SNAP-25 products were visualized by chromogenic detection using ImmunoPure TMB substrate kit (Pierce Biotechnology, Rockford, Ill.). The development of a yellow color in the BirA-HisTag®-SNAP-25<sub>134-197</sub> coated plates, but not the BirA-HisTag®-SNAP-25<sub>134-206</sub> coated plates, indicated that the α-SNAP-25 antibody preferentially recognized the SNAP-25<sub>197</sub> cleavage product. The result indicated that of the six mice used for immunization, three mice (Mouse 2, Mouse 3, and Mouse 4) had higher titers and more specificity towards a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

[0296] These results were confirmed using an ELISA light chain activity assay. A 96-well Reacti-Bind Streptavidin coated plates (Pierce Biotechnology, Rockford, Ill.) were prepared by adding approximately 100 µl of the following Substrate Solution: Rows A-C were coated with 100 µl of BirA-HisTag®-SNAP-25<sub>134-197</sub> at twelve different concentrations; Rows D-H were coated with 100 µl of BirA-HisTag®-SNAP-25<sub>134-206</sub> at 10 µg/mL. The plates were washed by aspirating the Substrate Solution and rinsing each well three times with 200 µl TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Dilutions of BoNT/A were pre-reduced at 37° C. for 20 minutes in BoNT/A Incubation Buffer (50 mM HEPES, pH 7.4, 1% fetal bovine serum, 10 µM ZnCl<sub>2</sub>, 10 mM dithiothreitol) and 100 µl of the pre-reduced BoNT/A was added to the substrate-coated plates and incubated at 37° C. for 90 minutes. BoNT/A treated plates were washed by aspirating the BoNT/A Incubation Buffer and rinsing each plate three times with 200 µl TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Washed plates were incubated at 37° C. for one hour in 0.5% BSA in 1× TBS

containing a 1:10 to 1:100 dilution of the antibody-containing serum being tested. Primary antibody probed plates were washed four times for 5 minutes each time in 200 µl TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Washed plates were incubated at 37° C. for 1 hour in 1× TBS containing a 1:10,000 dilution of goat polyclonal anti-mouse IgG antibody conjugated to Horseradish peroxidase (Pierce Biotechnology, Rockford, Ill.) as a secondary antibody. Secondary antibody-probed plates were washed four times in 200 µl TBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). Chromogenic detection of the labeled SNAP-25 products were visualized by chromogenic detection using ImmunoPure TMB substrate kit (Pierce Biotechnology, Rockford, Ill.). The development of a yellow color, which correlated with the presence of the SNAP-25197 cleavage product was detected in BoNT/A treated samples, but not untreated controls, using antibody-containing serum derived from all six immunized mice (Mouse 1, Mouse 2, Mouse 3, Mouse 4, Mouse 5, and Mouse 6). Thus, the comparative ELISA analysis indicated that of the mice used for immunization, three mice (Mouse 2, Mouse 3, and Mouse 4) had higher titers and more specificity towards a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

[0297] For cell-based cleavage assay, a suitable density of PC12 cells were plated into 60 mm<sup>2</sup> tissue culture plates containing 3 mL of an appropriate serum medium (Table 1). The cells were grown in a 37° C. incubator under 5% carbon dioxide until cells reached the appropriate density. A 500 µL transfection solution was prepared by adding 250 µL of OPTI-MEM Reduced Serum Medium containing 15 µL of LipofectAmine 2000 (Invitrogen Inc., Carlsbad, Calif.) incubated at room temperature for 5 minutes to 250 µL of OPTI-MEM Reduced Serum Medium containing 10 µg of a pQBI-25/GFP-BoNT/A-LC expression construct (SEQ ID NO:51). The pQBI-25/GFP-BoNT/A-LC expression construct comprises a pQBI-25 expression vector (Qbiogene Inc., Carlsbad, Calif.) whose promoter elements are functionally linked to a polynucleotide encoding the GFP-BoNT/A light chain of SEQ ID NO:52. This transfection mixture was incubated at room temperature for approximately 20 minutes. The media was replaced with fresh unsupplemented media and the 500 µL transfection solution was added to the cells. The cells were then incubated in a 37° C. incubator under 5% carbon dioxide for approximately 6 to 18 hours. The cells were washed and harvested as described in Example II. To detect for the presence of the cleaved SNAP-25197 product, an aliquot from each harvested sample was analyzed by Western blot as described in Example II, except that the primary antibody used was a 1:1,000 dilution of the antibody-containing serum and the secondary antibody used was a 1:20,000 of mouse α-IgG Horseradish Peroxidase (Pierce Biotechnology, Rockford, Ill.). A single band corresponding to the SNAP-25197 cleavage product was detected in BoNT/A treated samples, but not untreated controls, using antibody-containing serum derived from three mice (Mouse 2, Mouse 3, and Mouse 4). Thus, the cell-based cleavage assay indicated that of the mice used for immunization, three mice (Mouse 2, Mouse 3, and Mouse 4) had higher titers and more specificity towards a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

**[0298]** 3. Production of Hybridomas.

**[0299]** To make hybridomas producing  $\alpha$ -SNAP-25 monoclonal antibodies that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, the spleen from Mouse 2 was harvested three days subsequent to a final “booster” immunization and the spleen cells were fused with myeloma cells P3-X63 Ag8.653 using standard hybridoma protocols. These cells were plated into five 96-well plates and hybrids were selected using HAT medium. Within 8-14 days after fusion, the first screening of the approximately 480 parent clones was carried out using comparative ELISA with the BirA-HisTag®-SNAP-25<sub>134-197</sub> and the BirA-HisTag®-SNAP-25<sub>134-206</sub> peptides coated in two separate plates. The comparative ELISA provided a quick screen method to identify hybridomas producing antibodies specific for the cleaved SNAP-25197. The top 18 clones were subjected to further screening using the cell-based cleavage assay described above and immunostaining of LC/A transfected cells. (Table 20).

TABLE 20

Analysis of Supernatants Containing $\alpha$ -SNAP-25 Monoclonal Antibody						
Clone	Comparative ELISA				Cell-Based Assay	
	OD		Ratio		SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>
	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>	Ratio <sub>197/206</sub>	Ratio <sub>296/197</sub>		
1D3	1.805	0.225	8.02	0.13	+++	-
1F12	0.365	0.093	3.92	0.25	-	-
1G10	0.590	0.137	4.31	0.23	++	-
1H1	0.335	0.121	2.77	0.36	-	-
1H8	0.310	0.302	1.03	0.97	+	-
2C9	0.139	0.274	0.51	1.97	-	-
2E2	0.892	0.036	24.78	0.04	++	-
2E4	0.228	0.069	3.30	0.30	+	-
2F11	1.095	1.781	0.61	1.63	-	-
3C1	1.268	0.053	23.92	0.04	++	-
3C3	0.809	0.052	15.56	0.06	++	-
3E1	0.086	0.155	0.55	1.80	0	-
3E8	2.048	0.053	38.64	0.03	+++	-
3G2	0.053	0.158	0.34	2.98	-	-
4D1	0.106	0.218	0.49	2.06	-	-
4G6	0.061	0.159	0.38	2.61	-	-
5A5	0.251	0.106	2.37	0.42	+	-
5F11	0.243	0.061	3.98	0.25	-	-

**[0300]** Clones 103, 1G10, 2E2, 3C1, 3C3, and 3E8 were further cloned by limiting dilution because the conditioned media produced by these clones comprised  $\alpha$ -SNAP-25 antibodies with a preferential binding specificity having a ratio<sub>197/206</sub> of at least 4:1 for the SNAP-25<sub>197</sub> cleavage product relative to the SNAP-25<sub>206</sub> uncleaved substrate and detected the SNAP-25<sub>197</sub>-cleavage product using the cell-based cleavage assay and the immunostaining of PC12 cells transfected with GFP-LC/A. Similarly clones 2C9, 2F11, 3G2, 4D1 and 4G6 were further cloned by limiting dilution because the conditioned media produced by these clones comprised  $\alpha$ -SNAP-25 antibodies with a preferential binding specificity having a ratio<sub>206/197</sub> of at least 1.5:1 for the SNAP-25<sub>206</sub> uncleaved substrate relative to the SNAP-25<sub>197</sub> cleavage product and detected the SNAP-25<sub>206</sub>-uncleaved substrate using the cell-based cleavage assay. These single-cell derived clones were screened again using comparative ELISA, cell-based cleavage, and immunostaining to confirm

their affinity and specificity, and the antibodies were isotyped using standard procedures. Ascites were produced from clones 103B8 (IgM.k), 1G10A12 (IgG3.k), 2C9B10 (IgG3.k), 2E2A6 (IgG3.k), 2F11B6 (IgM.k), 3C1A5 (IgG2a.k), and 3C3E2 (IgG2a.k). Clone 3E8 stopped producing antibodies during the cloning process and could not be further evaluated.

**[0301]** 4. Evaluation of Binding Specificity of  $\alpha$ -SNAP-25 Monoclonal Antibodies.

**[0302]** To evaluate binding specificity of an  $\alpha$ -SNAP-25 monoclonal antibody that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, ascites from clones 103B8, 1G10A12, 2C9B10, 2E2A6, 2F11B6, 3C1A5, and 3C3E2 were used to detect SNAP-25 cleavage product using the cell-based activity assay, immunocytochemistry and immunoprecipitation.

**[0303]** For the cell-based activity assay, binding specificity was determined by analyzing the ability of  $\alpha$ -SNAP-25 antibody-containing ascites to detect the uncleaved SNAP-25206 substrate and the cleaved SNAP-25197 product by Western blot analysis. A suitable density of PC12 cells were plated into 60 mm<sup>2</sup> tissue culture plates containing 3 mL of an appropriate serum medium, grown in a 37° C. incubator under 5% carbon dioxide until an appropriate cell density was reached, and transfected with either a transfection solution lacking the pQBI-25/GFP-BoNT/A-LC expression construct (untransfected cells) or a transfection solution containing the pQBI-25/GFP-BoNT/A-LC expression construct (transfected cells) as described above. The cells were washed and harvested as described in Example I. To detect for the presence of both the uncleaved SNAP-25<sub>206</sub> substrate and the cleaved SNAP-25<sub>197</sub> product, an aliquot from each harvested sample was analyzed by Western blot as described in Example I, except that the primary antibody used was a 1:100 dilution of the  $\alpha$ -SNAP-25 monoclonal antibody-containing ascites and the secondary antibody used was a 1:20,000 of  $\alpha$ -mouse IgG conjugated to Horseradish Peroxidase (Pierce Biotechnology, Rockford, Ill.). In addition, three commercially available mouse  $\alpha$ -SNAP-25 monoclonal antibodies were tested. SMI-81 (Sternberger Monoclonals Inc., Lutherville, Md.), an  $\alpha$ -SNAP-25 antibody the manufacturer indicates detects both the uncleaved SNAP-25<sub>206</sub> substrate and the cleaved SNAP-25<sub>197</sub> product, was used at a 15,000 dilution according to the manufacturer's recommendations. MC-6050 (Research & Diagnostic Antibodies, Las Vegas, Nev.), an  $\alpha$ -SNAP-25 antibody the manufacturer indicates detects both the uncleaved SNAP-25<sub>206</sub> substrate and the cleaved SNAP-25<sub>197</sub> product, was used at a 1:100 dilution according to the manufacturer's recommendations. MC-6053 (Research & Diagnostic Antibodies, Las Vegas, Nev.), an  $\alpha$ -SNAP-25 antibody the manufacturer indicates detects only the cleaved SNAP-25<sub>197</sub> product, was used at a 1:100 dilution according to the manufacturer's recommendations.

**[0304]** Table 21 indicates the  $\alpha$ -SNAP-25 antibody-containing ascites that detected only the SNAP-25<sub>197</sub> cleavage product. The cell-based cleavage assay indicated that ascites produced from clones 1D3B8, 2C9B10, 2E2A6, 3C1A5, and 3C3E2 synthesize an  $\alpha$ -SNAP-25 monoclonal antibody having high binding specificity for the SNAP-25<sub>197</sub> cleavage product that allows for the selective recognition of this cleavage product relative to the SNAP-25<sub>206</sub> uncleaved substrate. Commercial antibody SMI-81 detected the SNAP-

25<sub>206</sub> uncleaved substrate, but only poorly recognized the SNAP-25<sub>197</sub> cleavage product (Table 21). Surprisingly, commercial antibody MC-6050 only detected the SNAP-25<sub>206</sub> uncleaved substrate, and failed to recognize the SNAP-25<sub>197</sub> cleavage product (Table 21). Even more surprisingly, commercial antibody MC-6050 only detected the SNAP-25<sub>206</sub> uncleaved substrate, and failed to recognize the SNAP-25<sub>197</sub> cleavage product, even though the manufacturer advertises that this antibody selectively detects the SNAP-25<sub>197</sub> cleavage product (Table 21). Thus, this analysis indicates that while 1D3B8, 2C9B10, 2E2A6, 3C1A5, and 3C3E2 exhibit suitable selectivity for the SNAP-25<sub>197</sub> cleavage product, 1G10A12 and 2F11B6 do not. In addition, commercial antibodies SMI-81, MC-6050 and MC-6053 all are unsuitable for the immuno-based methods disclosed in the present application because all failed to selectively detect the SNAP-25<sub>197</sub> cleavage product.

**[0305]** For immunocytochemistry analysis, binding specificity was determined by analyzing the ability of  $\alpha$ -SNAP-25 antibody-containing ascites to detect the uncleaved SNAP-25<sub>206</sub> substrate and the cleaved SNAP-25<sub>197</sub> product by immunostaining. See e.g., Ester Fernandez-Salas et al., *Plasma Membrane Localization Signals in the Light Chain of Botulinum Neurotoxin*, *Proc. Natl. Acad. Sci., U.S.A.* 101(9): 3208-3213 (2004). A suitable density of PC12 cells were plated, grown, and transfected with either a transfection solution lacking the pQBI-25/GFP-BoNT/A-LC expression construct (untransfected cells) or a transfection solution containing the pQBI-25/GFP-BoNT/A-LC expression construct (transfected cells) as described above. The cells were washed in 1 $\times$  PBS and fixed in 5 mL of PAF at room temperature for 30 minutes. Fixed cells were washed in phosphate buffered saline, incubated in 5 mL of 0.5% Triton<sup>®</sup> X-100 (polyethylene glycol octylphenol ether) in 1 $\times$  PBS, washed in 1 $\times$  PBS, and permeabilized in 5 mL of methanol at -20 $^{\circ}$  C. for six minutes. Permeabilized cells were blocked in 5 mL of 100 mM glycine at room temperature for 30 minutes, washed in 1 $\times$  PBS, and blocked in 5 mL of 0.5% BSA in 1 $\times$  PBS at room temperature for 30 minutes. Blocked cells were washed in 1 $\times$  PBS and incubated at room temperature for two hours in 0.5% BSA in 1 $\times$  PBS containing a 1:10 dilution of an ascites from a clonal hybridoma cell line being tested. Primary antibody probed cells were washed three times for 5 minutes each time in 1 $\times$  PBS. Washed cells were incubated at room temperature for 2 hours in 1 $\times$  PBS containing a 1:200 dilution of goat polyclonal anti-mouse immunoglobulin G, heavy and light chains (IgG, H+L) antibody conjugated to ALEXA<sup>®</sup> FLUOR 568 (Invitrogen Inc., Carlsbad, Calif.) as a secondary antibody. Secondary antibody-probed cells were washed three times for 5 minutes each time in 1 $\times$  PBS. Washed cells were prepared for microscopic examination by mounting in VECTASHIELD<sup>®</sup> Mounting Media (Vector Laboratories, Burlingame, Calif.) and coverslipped. Images of signal detection were obtained with a Leica confocal microscope using appropriate laser settings. Table 21 indicates that the  $\alpha$ -SNAP-25 antibody-containing ascites that specifically detected the SNAP-25<sub>197</sub>-cleavage product. The immuno-

cytochemistry analysis indicated that ascites produced from clones 1D3B8, 2C9B10, 2E2A6, 3C1A5, and 3C3E2 synthesize an  $\alpha$ -SNAP-25 monoclonal antibody having high binding specificity for the SNAP-25<sub>197</sub> cleavage product that allows for the preferential recognition of this cleavage product relative to the SNAP-25<sub>206</sub> uncleaved substrate.

**[0306]** For immunoprecipitation analysis, binding specificity was determined by analyzing the ability of Protein A (Hilrap<sup>™</sup> Protein A HP Columns, GE Healthcare, Amersham, Piscataway, N.J.), purified  $\alpha$ -SNAP-25 monoclonal antibodies to precipitate the uncleaved SNAP-25<sub>206</sub> substrate and the cleaved SNAP-25<sub>197</sub> product. See e.g., Chapter 8 Storing and Purifying Antibodies, pp. 309-311, Harlow & Lane, supra, 1998a. A suitable density of PC12 cells were plated, grown, and transfected with either a transfection solution containing a pQBI-25/GFP expression construct (control cells; SEQ ID NO:53) or a transfection solution containing the pQBI-25/GFP-BoNT/A-LC expression construct (experimental cells) as described above. The pQBI-25/GFP expression construct comprises an expression vector whose promoter elements are functionally linked to a polynucleotide encoding GFP of SEQ ID NO:54. After an overnight incubation, the cells were washed by aspirating the growth media and rinsing each well with 200  $\mu$ L 1 $\times$  PBS. To harvest the cells, the PBS was aspirated, the cells were lysed by adding an Immunoprecipitation Lysis Buffer comprising 50 mM HEPES, 150 mM NaCl, 1.5 mM MgCl<sub>2</sub>, 1 mM EGDT, 10% glycerol, 1% Triton<sup>®</sup> X-100 (polyethylene glycol octylphenol ether) and a 1 $\times$  COMPLETE<sup>™</sup> Protease inhibitor cocktail (Roche Applied Biosciences, Indianapolis, Ind.) and incubating at 4 $^{\circ}$  C. for one hour. The lysed cells were centrifuged at 3,000 $\times$  g at 4 $^{\circ}$  C. for 10 minutes to remove cellular debris and the supernatant transferred to a clean tube and diluted to a protein concentration of approximately 1 mg/mL. Approximately 5  $\mu$ g of purified monoclonal antibody was added to 0.5 mL of diluted supernatant and incubated at 4 $^{\circ}$  C. for two hours. After primary antibody incubation, approximately 50  $\mu$ L of immobilized Protein G (Pierce Biotechnology, Rockford, Ill.) was added to the diluted supernatant and incubated at 4 $^{\circ}$  C. for one hour. The incubated supernatant was washed three times for 30 minutes each time by adding 0.5 mL of Immunoprecipitation Lysis Buffer, centrifuging at 300 $\times$  g at 4 $^{\circ}$  C. for one minute to pellet the immobilized Protein G, and decanting the supernatant. After washing, the pellet was resuspended in 30  $\mu$ L of 1 $\times$  SDS Loading Buffer and the sample was heated to 95 $^{\circ}$  C. for 5 minutes. To detect for the presence of both the uncleaved SNAP-25<sub>206</sub> substrate and the cleaved SNAP-25<sub>197</sub> product, an aliquot from each harvested sample was analyzed by Western blot as described in Example I, except that the primary antibody used was a 1:1,000 dilution of the  $\alpha$ -SNAP-25 polyclonal antibody serum (see Example V) and the secondary antibody used was a 1:20,000 of rabbit  $\alpha$ -IgG Horseradish Peroxidase (Pierce Biotechnology, Rockford, Ill.). Table 21 indicates the  $\alpha$ -SNAP-25 antibody-containing ascites that specifically pulled down the SNAP-25<sub>197</sub>-cleavage product by immunoprecipitation analysis. The immunoprecipitation analysis indicated that ascites produced from clones 2E2A6 and 3C1A5 synthesize an  $\alpha$ -SNAP-25 monoclonal antibody having high binding specificity for the SNAP-25<sub>197</sub> cleavage product that allows for the preferential recognition of this cleavage product relative to the SNAP-25<sub>206</sub> uncleaved substrate.

TABLE 21

Analysis of Clone Ascites Containing $\alpha$ -SNAP-25 Monoclonal Antibody						
Clone	Cell-Based Assay		Immunocytochemistry		Immunoprecipitation	
	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>
1D3B8	++	-	++	-	Not Tested	Not Tested
1G10A12	++	++	Not Tested	Not Tested	Not Tested	Not Tested
2C9B10	++	-	++	-	Not Tested	Not Tested
2E2A6	++	-	++	-	++	-
2F11B6	+	+	+	+	Not Tested	Not Tested
3C1A5	++	-	++	-	++	-
3C3E2	+	-	Not Tested	Not Tested	Not Tested	Not Tested
MC-6050	-	+	Not Tested	Not Tested	Not Tested	Not Tested
MC-6053	-	+	Not Tested	Not Tested	Not Tested	Not Tested
SMI-81	-/+	++	Not Tested	Not Tested	Not Tested	Not Tested

### [0307] 5. Evaluation of Binding Affinity of $\alpha$ -SNAP-25 Monoclonal Antibodies

**[0308]** To determine the binding affinity of an  $\alpha$ -SNAP-25 monoclonal antibody showing high binding specificity for either the SNAP-25<sub>197</sub> cleavage product or the SNAP-25<sub>206</sub> uncleaved substrate, binding affinity assays were performed on a BIAcore® 3000 instrument using carboxymethyl dextran (CM5) sensor chips (BIAcore, Inc., Piscataway, N.J.). Runs were conducted at 25° C. with HBS-EP buffer comprising 10 mM HEPES (pH 7.4), 150 mM sodium chloride, 3 mM EDTA, 0.005% (v/v) surfactant P20 at a flow rate of 10  $\mu$ L/min. SNAP-25 peptides comprising amino acids 134-197 of SEQ ID NO:5 (SNAP-25<sub>134-197</sub>) or amino acids 134-206 of SEQ ID NO:5 (SNAP-25<sub>134-206</sub>) were covalently attached to the surface of the CM5 sensor chips using standard amine coupling. Briefly, the CM5 chips were activated by a 7 minute injection of a mixture of 0.2 M 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide and 0.05 M N-hydroxysuccinimide; the SNAP-25 peptides were then injected in 10 mM sodium acetate (pH 4.0) for 20 min at a flow rate of 10  $\mu$ L/min; and unreacted succinimide esters were blocked by a 7-min injection of 1 M ethanolamine hydrochloride, pH 8.5. The immobilized amount of SNAP-25<sub>134-197</sub> or SNAP-25<sub>134-206</sub> on the chip was reflected by a 100-150 increase in response units (about 0.10-0.15 ng/mm<sup>2</sup>). Antibody samples comprising either ascites or purified monoclonal antibodies produced from clones 1D3B8, 2C9B10, 2E2A6, 3C1A5, and 3C3E2, as well as, commercially available  $\alpha$ -SNAP-25 antibodies were passed over the surface of the CM5 chips allowing an association time of 10 min and a dissociation time of 20 min. The surfaces were regenerated between runs by a 1 minute injection of 10 mM glycine-HCl (pH 2.5) at a flow rate of 15  $\mu$ L/min. Sensorgram curves were fitted to a 1:1 kinetic binding model with the BIAevaluation 3.0 software.

**[0309]** The results indicate that both 2E2A6 and 3C1A5 were highly specific for cleaved SNAP-25<sub>197</sub> product over SNAP-25 uncleaved substrate (Table 22). When compared to the binding affinities of MC-6050 and MC-6053, 1D3B8 had an approximately 10-fold higher equilibrium dissociation constant for the SNAP-25 cleavage product relative to these commercial antibodies (Table 22). Interestingly, 2E2A6 had only a slightly lower equilibrium dissociation constant for the SNAP-25 cleavage product relative to these commercial antibodies (0.405 nM versus 0.497 and 0.508) (Table 22). As neither of these commercial  $\alpha$ -SNAP-25 antibodies selectively recognized the SNAP-25 cleavage

product (Table 21), an equilibrium dissociation constant lower than about 0.5 nM appears, in part, critical to achieve such selectivity. Similarly, when compared to the binding affinities of MC-6050 and MC-6053, 2E2A6 had an about at least one-fold slower off rate/dissociation constant ( $6.74 \times 10^{-5}$  versus  $8.82 \times 10^{-4} \text{ s}^{-1}$  and  $1.18 \times 10^{-3} \text{ s}^{-1}$ ) (Table 22). This further suggests that an off rate/dissociation constant lower than about  $8.82 \times 10^{-4}$  appears, in part, critical to achieve selective binding for the SNAP-25 cleavage product. This result is consistent with 1D3B8, which had an off rate/dissociation constant of  $5.78 \times 10^{-5} \text{ s}^{-1}$  (Table 22).

TABLE 22

Analysis of Binding Affinity $\alpha$ -SNAP-25 Monoclonal Antibodies				
SPR	1D3B8		2E2A6	
Parameter	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub> <sup>a</sup>	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub> <sup>b</sup>
Ka ( $\text{M}^{-1} \text{ s}^{-1}$ )	$1.06 \times 10^6$	—	$1.70 \times 10^6$ ( $1.66 \times 10^5$ )	— (—)
Kd ( $\text{s}^{-1}$ )	$5.78 \times 10^{-5}$	—	$1.53 \times 10^{-4}$ ( $6.74 \times 10^{-5}$ )	— (—)
KD (nM)	0.050	—	0.090 (0.405)	— (—)
SPR	3C1A5		2C9B10	
Parameter	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub> <sup>c</sup>	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub> <sup>d</sup>
Ka ( $\text{M}^{-1} \text{ s}^{-1}$ )	$2.17 \times 10^5$	—	$1.15 \times 10^4$	—
Kd ( $\text{s}^{-1}$ )	$2.88 \times 10^{-4}$	—	$3.11 \times 10^{-4}$	—
KD (nM)	1.33	—	27.1	—
SPR	MC-6050		MC-6053	
Parameter	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>	SNAP-25 <sub>197</sub>	SNAP-25 <sub>206</sub>
Ka ( $\text{M}^{-1} \text{ s}^{-1}$ )	$1.78 \times 10^6$	$3.06 \times 10^2$	$2.32 \times 10^6$	$1.06 \times 10^2$
Kd ( $\text{s}^{-1}$ )	$8.82 \times 10^{-4}$	$6.07 \times 10^{-3}$	$1.18 \times 10^{-3}$	$2.56 \times 10^{-5}$
KD (nM)	0.497	19,800	0.508	240

<sup>a</sup>No binding was observed when up to 125 nM of  $\alpha$ -SNAP-25 monoclonal antibody 1D3B8 was passed over the surface of the CM5 sensor chip after a 10 minute association time.

<sup>b</sup>No binding was observed when up to 10  $\mu$ M of  $\alpha$ -SNAP-25 monoclonal antibody 2E2A6 was passed over the surface of the CM5 sensor chip after a 10 minute association time.

<sup>c</sup>No binding was observed when up to 100 nM of  $\alpha$ -SNAP-25 monoclonal antibody 3C1A5 was passed over the surface of the CM5 sensor chip after a 10 minute association time.

<sup>d</sup>No binding was observed when up to 100 nM of  $\alpha$ -SNAP-25 monoclonal antibody 2C9B10 was passed over the surface of the CM5 sensor chip after a 10 minute association time.

**[0310]** To compare the six different antibodies, the on-rate (ka) and off-rate (kd) for each was normalized using a program from the BIA evaluation 4.1 software. For comparison of the on-rates, the data were first individually

trimmed by deleting the re-generation portion and the injection spikes, and then normalized to a 0 to 100 scale. For comparison of the off-rate, the data were normalized to the injection stop/top point. This analysis showed that 209B10 had a much slower on-rate than the other antibodies (FIG. 7A), and that MC-6053 has a much faster off-rate (dissociation) than the other antibodies (FIG. 7B). The fast off-rate of MC-6053 indicates that this antibody will not perform well in the methods disclosed in the present specification because this antibody will have difficulty staying bound to the substrate antigen during the washing steps.

**[0311]** 6. Sequencing of the Epitope from Isolated  $\alpha$ -SNAP-25 Monoclonal Antibodies.

**[0312]** To determine the epitope of an isolated  $\alpha$ -SNAP-25 monoclonal antibody that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond, the polynucleotide molecule encoding the variable heavy (V<sub>H</sub>) and variable light (V<sub>L</sub>) chains of the  $\alpha$ -SNAP-25 monoclonal antibody produced by hybridomas 1D3B8, 2C9B10, 2E2A6, 3C1A5 and 3C3E2 were sequenced. mRNA was extracted and purified from each hybridoma using standard protocols and reversed transcribed into cDNA using either an oligo dT anti-sense primer or a gene-specific (murine IgG1 CH and kappa CL) anti-sense primer. Specific murine and human constant domain primers were used to amplify the cDNA by PCR after cDNA production to determine the isotype of the antibody. Degenerate V<sub>H</sub> and V<sub>L</sub> primers were used to amplify the variable domains from the cDNA. For 5'RACE, a homopolymeric dCTP tail was added to the 3' end of the cDNA. The heavy and light chains were then amplified with an oligo dG sense primer and a gene specific (CH/KC) anti-sense primer. PCR products included the sequence of the signal peptide, variable domains and constant domains up to the anti-sense primer. The PCR products were gel purified to remove small fragments, and cloned into a blunt or TA vector for sequencing. Five independent clones for each chain were sequenced and alignments of V<sub>H</sub> and V<sub>L</sub> chains and consensus sequences were determined. Methods used to determine the V<sub>H</sub> and V<sub>L</sub> amino acid sequences are described in, e.g., Roger A. Sabbadini, et al., Novel Bioactive Lipid Derivatives and Methods of Making and Using Same, U.S. Patent Publication 2007/0281320; and Peter Amersdorfer, et al., Molecular Characterization of Murine Humoral Immune Response to Botulinum Neurotoxin Type A Binding Domain as Assessed by Using Phage Antibody Libraries, 65(9) Infect. Immun. 3743-3752, each of which is hereby incorporated by reference in its entirety. In addition, commercial services are available to sequence the variable heavy (V<sub>H</sub>) and variable light (V<sub>L</sub>) chains of an antibody and identify the CDR regions, see, e.g., Fusion Antibodies Ltd., Northern Ireland. In one case, for the 3C1A5 V<sub>L</sub> region, the amino acid sequence was also determined by separating the affinity purified antibody by high resolution 2DE electrophoresis and then subjecting the protein to peptide fragmentation fingerprinting analysis using high resolution nanoLC-MSMS after proteolytic digestion.

**[0313]** The polynucleotide sequence comprising the V<sub>H</sub> and V<sub>L</sub> chains of the  $\alpha$ -SNAP-25 monoclonal antibody produced by the hybridomas disclosed in the present specification is as follows: 1D3B8 V<sub>H</sub> (SEQ ID NO:71), 2C9B10 V<sub>H</sub> (SEQ ID NO:73), 2E2A6 V<sub>H</sub> (SEQ ID NO:75), 3C1A5 V<sub>H</sub> (SEQ ID NO:77), 3C3E2 V<sub>H</sub> variant 1 (SEQ ID NO:79), 3C3E2 V<sub>H</sub> variant 2 (SEQ ID NO:81), 3C3E2 V<sub>H</sub> variant 3

(SEQ ID NO:132), 1D3B8 V<sub>L</sub> (SEQ ID NO:83), 2C9B10 V<sub>L</sub> (SEQ ID NO:85), 2E2A6 V<sub>L</sub> (SEQ ID NO:87), 3C1A5 V<sub>L</sub> (SEQ ID NO:89), and 3C3E2 V<sub>L</sub> (SEQ ID NO:91). The amino acid sequence comprising the V<sub>H</sub> and V<sub>L</sub> chains of the  $\alpha$ -SNAP-25 monoclonal antibody produced by the hybridomas disclosed in the present specification is as follows: 1D3B8 V<sub>H</sub> (SEQ ID NO:72), 2C9B10 V<sub>H</sub> (SEQ ID NO:74), 2E2A6 V<sub>H</sub> (SEQ ID NO:76), 3C1A5 V<sub>H</sub> (SEQ ID NO:78), 3C3E2 V<sub>H</sub> variant 1 (SEQ ID NO:80), 3C3E2 V<sub>H</sub> variant 2 (SEQ ID NO:82); 3C3E2 V<sub>H</sub> variant 2 (SEQ ID NO:133), 1D3B8 V<sub>L</sub> (SEQ ID NO:84), 2C9B10 V<sub>L</sub> (SEQ ID NO:86), 2E2A6 V<sub>L</sub> (SEQ ID NO:88), 3C1A5 V<sub>L</sub> (SEQ ID NO:90), and 3C3E2 V<sub>L</sub> (SEQ ID NO:92). The amino acid sequences comprising the V<sub>H</sub> and V<sub>L</sub> CDR domains of the  $\alpha$ -SNAP-25 monoclonal antibody produced by the hybridomas 1D3B8, 2C9B10, 2E2A6, 3C1A5, and 3C3E2 are given in Table 23.

TABLE 23

CDR Sequences of V <sub>H</sub> and V <sub>L</sub> domains from $\alpha$ -SNAP-25 Monoclonal Antibodies			
CDR	Sequence	Identified In	SEQ ID NO:
V <sub>H</sub> CDR 1	TFTDHSIH	2E2A6 2C9B10 3C1A5	93
V <sub>H</sub> CDR 1	TFTNYVIH	3C3E2	94
V <sub>H</sub> CDR 1	IFTDHALH	1D3B8	95
V <sub>H</sub> CDR 2	YIFPGNGNIEYNDKFKG	2E2A6	96
V <sub>H</sub> CDR 2	YIFPGNGNIEYNDKFKG	2C9B10 3C1A5	97
V <sub>H</sub> CDR 2	YINPYNDGSKYNEKFKG	3C3E2	98
V <sub>H</sub> CDR 2	YIFPGNGNIEYNEKFKG	1D3B8	99
V <sub>H</sub> CDR 3	KRMGY	2E2A6 3C1A5	100
V <sub>H</sub> CDR 3	KKMDY	2C9B10 1D3B8	101
V <sub>H</sub> CDR 3	ARMYD	3C3E2var1	102
V <sub>H</sub> CDR 3	ARMGY	3C3E2var2	134
V <sub>H</sub> CDR 3	ARHLANTYFFDY	3C3E2var3	135
V <sub>L</sub> CDR 1	RSSQSIVHSNGNTYLE	1D3B8	103
V <sub>L</sub> CDR 1	RTTENIYSYFV	2C9B10	104
V <sub>L</sub> CDR 1	KSSQSLLYTNKTYLT	2E2A6	105
V <sub>L</sub> CDR 1	KSSQSLLNTNGKTYLT	3C1A5	106
V <sub>L</sub> CDR 1	RASQNIGNYLH	3C3E2	107
V <sub>L</sub> CDR 2	KVSNRFS	1D3B8	108
V <sub>L</sub> CDR 2	NAKSLAE	2C9B10	109
V <sub>L</sub> CDR 2	LVSELDLS	2E2A6	110
V <sub>L</sub> CDR 2	LVSKLDS	3C1A5	111
V <sub>L</sub> CDR 2	YASQSIG	3C3E2	112
V <sub>L</sub> CDR 3	FQGSHPPT	1D3B8	113

TABLE 23-continued

CDR Sequences of $V_H$ and $V_L$ domains from $\alpha$ -SNAP-25 Monoclonal Antibodies			
CDR	Sequence	Identified In	SEQ ID NO:
$V_L$ CDR 3	QHHTGTPYT	2C9B10	114
$V_L$ CDR 3	LQSAHFPPFT	2E2A6	115
$V_L$ CDR 3	LQSSHPPFT	3C1A5	116
$V_L$ CDR 3	QQSDTWPLT	3C3E2	117

**[0314]** Non-limiting examples of amino acid sequences comprising  $V_H$  CDR domain variants of the  $\alpha$ -SNAP-25 monoclonal antibody produced by the hybridomas disclosed in the present specification include  $V_H$  CDR1 variant SEQ ID NO:118 for 1D3B8;  $V_H$  CDR1 variant SEQ ID NO:119 for 2C9B10, 2E2A6 and 3C1A5  $V_H$ ;  $V_H$  CDR1 variant SEQ ID NO:120 for 3C1A5  $V_H$  and 3C3E2 variant 3;  $V_H$  CDR2 variant SEQ ID NO:121 for 1C3B8 and 2E2A6;  $V_H$  CDR2 variant SEQ ID NO:122 for 2C9B10 and 3C1A5  $V_H$ ;  $V_H$  CDR2 variant SEQ ID NO:123 for 3C1A5  $V_H$  and 3C3E2 variant 3;  $V_H$  CDR3 variant MDY for 1D3B8 and 2C9B10;  $V_H$  CDR3 variant MGY for 2E2A6 and 3C1A5  $V_H$ ; and  $V_H$  CDR3 variant SEQ ID NO:124 for 3C1A5  $V_H$  and 3C3E2 variant 3. Non-limiting examples of amino acid sequences comprising  $V_L$  CDR domain variants of the  $\alpha$ -SNAP-25 monoclonal antibody produced by the hybridomas disclosed in the present specification include  $V_L$  CDR1 variant SEQ ID NO:125 for 1D3B8;  $V_L$  CDR1 variant SEQ ID NO:126 for 2C9B10;  $V_L$  CDR1 variant SEQ ID NO:127 for 2E2A6;  $V_L$  CDR1 variant SEQ ID NO:128 for 3C1A5;  $V_L$  CDR1 variant SEQ ID NO:129 for 3C3E2;  $V_L$  CDR2 variant KVS for 103B8;  $V_L$  CDR2 variant NAK for 2C9B10;  $V_L$  CDR2 variant LVS for 2E2A6;  $V_L$  CDR2 variant YAT for 3C1A5; and  $V_L$  CDR2 variant YAS for 3C3E2.

#### Example VIII

##### Development of $\alpha$ -SNAP-25 Polyclonal Antibodies that Selectively Bind a SNAP-25 Epitope Having a Free Carboxyl-terminus at the $P_1$ Residue of the BoNT/A Cleavage Site Scissile Bond

**[0315]** The following example illustrates how to make  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to a SNAP-25 epitope having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond.

**[0316]** To develop  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond, the 10-residue peptide CGGGRIDEANQ (SEQ ID NO:46) was designed as a SNAP-25 cleavage product antigen. This peptide comprising a N-terminal Cysteine residue for conjugation to KLH, a G-spacer flexible spacer (GGG) linked to amino acids 191-197 of human SNAP-25 (SEQ ID NO:5) and has a carboxylated C-terminal glutamine. Blast searches revealed that this peptide has high homology only to SNAP-25 and almost no possible cross-reactivity with other proteins in neuronal cells. The sequence was also carefully scrutinized by utilizing computer algorithms to determine hydropathy index, protein surface probability, regions of flexibility, and favorable secondary structure, followed by proper orientation and

presentation of the chosen peptide sequence. The peptide was synthesized and conjugated to Keyhole Limpet Hemocyanin (KLH) to increase immunogenicity. Before the animals were immunized, naive rabbits were first screened against cell lysates from candidate cell lines in a Western blot in order to identify animals that had no immunoreactivity to the proteins present in the cell lysates. Two pre-screened rabbits were immunized with this peptide, and after three immunizations in about eight weeks, the rabbits were bled for testing. The blood was allowed to clot by incubating at 4° C. for 60 minutes. The clotted blood was centrifuged at 10,000x g at 4° C. for 10 minutes to pellet the cellular debris. The resulting serum sample was dispensed into 50  $\mu$ L aliquots and stored at -20° C. until needed.

**[0317]** A similar strategy based on other SNAP-25 antigens disclosed in the present specification is used to develop  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to a SNAP-25 having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. For example, the SNAP-25 antigen of SEQ ID NO:47 can be conjugated to KLH instead of the SNAP-25 antigen of SEQ ID NO:46. As another example, the amino acids 191-197 of human SNAP-25 from the SNAP-25 antigen of SEQ ID NO:38 can be replaced with SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, or SEQ ID NO:44.

#### 2. Screening for the Presence of $\alpha$ -SNAP-25 Polyclonal Antibodies.

**[0318]** To determine the presence of  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond, comparative ELISA and cell-based cleavage assays were performed using the extracted rabbit serum as described in Example III. The serum from both rabbits contained  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond. The  $\alpha$ -SNAP-25 rabbit polyclonal antibodies were designated as NTP 22 and NTP 23.

#### 3. Purification of $\alpha$ -SNAP-25 Polyclonal Antibodies.

**[0319]** To purify  $\alpha$ -SNAP-25 polyclonal antibodies that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond, NTP 22 and NTP 23 antibodies from rabbit serum were purified using affinity columns containing the SNAP-25 antigen of SEQ ID NO:46.

#### 4. Evaluation of Binding Specificity of $\alpha$ -SNAP-25 Polyclonal Antibodies.

**[0320]** To evaluate binding specificity of an  $\alpha$ -SNAP-25 polyclonal antibody that can selectively bind to a SNAP-25 antigen having a carboxyl-terminus at the  $P_1$  residue of the BoNT/A cleavage site scissile bond, purified NTP 22 and NTP 23  $\alpha$ -SNAP-25 polyclonal antibodies were used to detect cleavage product using the cell-based activity assay, immunocytochemistry and immunoprecipitation as described in Example III. The cell-based cleavage assay, immunocytochemistry analysis and immunoprecipitation analysis all indicated that NTP 22 and NTP 23  $\alpha$ -SNAP-25 polyclonal antibodies did not cross-react with uncleaved

SNAP-25. Thus both NTP 22 and NTP 23 have high binding specificity for the SNAP-25<sub>197</sub> cleavage product that allows for the preferential recognition of this cleavage product relative to the SNAP-25<sub>206</sub> uncleaved substrate. Affinity for the antigens can be determined using SPR in the BiAcore as described in Example III.

#### Example IX

##### Component and Condition Preparation for a Sandwich ELISA

**[0321]** The following example illustrates how to identify and prepare the components and conditions necessary to perform a sandwich ELISA useful for conducting immunobased methods of detecting retargeted endopeptidase activity by detecting a SNAP-25 cleavage product using an  $\alpha$ -SNAP-25 monoclonal antibody specific for a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond.

##### 1. Preparation of Cell Lysates from Cells Treated with Re-Targeted Endopeptidase.

**[0322]** To obtain a re-targeted endopeptidase treated cell lysate for analysis, a suitable density of cells from a stock culture of Neuro-2a was seeded into a T175 flask containing 50 mL of a serum-free medium containing Minimum Essential Medium, 2 mM GlutaMAX™ 1 with Earle's salts, 1× B27 supplement, 1×N2 supplement, 0.1 mM Non-Essential Amino Acids, 10 mM HEPES. These cells were incubated in a 37° C. incubator under 5% carbon dioxide until the cells differentiated, as assessed by standard and routine morphological criteria, such as growth arrest and neurite extension (approximately 2 to 3 days).

**[0323]** As a control, a suitable density of cells from a stock culture of Neuro-2a was seeded into a T175 flask containing 50 mL of an appropriate growth medium (Table 1). These undifferentiated control cells were grown in a 37° C. incubator under 5% carbon dioxide until 50% confluence was reached (approximately 18 hours). The media from both differentiated and undifferentiated control cultures was aspirated from each well and replaced with fresh media containing either 0 (untreated sample) or 10 nM of a re-targeted endopeptidase. After an overnight incubation, the cells were washed and the cells harvested by lysing in freshly prepared Triton X-100 Lysis Buffer (50 mM HEPES, 150 mM NaCl, 1.5 mM MgCl<sub>2</sub>, 1 mM EGTA, 1% Triton X-100) at 4° C. for 30 minutes with constant agitation. Lysed cells were centrifuged at 4000 rpm for 20 min at 4° C. to eliminate debris using a bench-top centrifuge. The protein concentrations of cell lysates were measured by Bradford assay.

##### 2. Preparation and Identification of Sandwich ELISA Components.

**[0324]** To identify an appropriate capture antibody-detection antibody pair an ECL sandwich ELISA analysis was conducted on twenty-six different combinations of capture and detection antibody pairs comprising eleven different  $\alpha$ -SNAP-25 capture antibodies and seven different  $\alpha$ -SNAP-25 detection antibodies (Table 12). The  $\alpha$ -SNAP-25 antibodies used were 2E2A6 and 3C1A5  $\alpha$ -SNAP-25 mouse monoclonal antibodies disclosed in the present specification, SMI-81, MC-6050, and MC-6053  $\alpha$ -SNAP-25 mouse monoclonal antibodies disclosed in the present specification, NTP 23  $\alpha$ -SNAP-25 rabbit polyclonal antibodies

disclosed in the present specification, S9684  $\alpha$ -SNAP-25 rabbit polyclonal antibodies (Sigma, St. Louis, Mo.), H-50  $\alpha$ -SNAP-25 rabbit polyclonal antibodies (Santa Cruz Biotechnology, Inc., Santa Cruz, Calif.), C-18  $\alpha$ -SNAP-25 goat polyclonal antibodies (Santa Cruz Biotechnology, Inc., Santa Cruz, Calif.), N-19  $\alpha$ -SNAP-25 goat polyclonal antibodies (Santa Cruz Biotechnology, Inc., Santa Cruz, Calif.), and SP12  $\alpha$ -SNAP-25 mouse polyclonal antibodies (Santa Cruz Biotechnology, Inc., Santa Cruz, Calif.).

**[0325]** To prepare the capture antibody solution, the  $\alpha$ -SNAP-25 monoclonal antibodies contained in the ascites from hybridoma cell lines 2E2A6 and 3C1A5 as well as the  $\alpha$ -SNAP-25 MC-6050 and MC-6053 monoclonal antibodies were purified using a standard Protein A purification protocol. All other  $\alpha$ -SNAP-25 antibodies were purchased as purified antibodies.

**[0326]** To prepare the detection antibody solution, the appropriate  $\alpha$ -SNAP-25 antibody was conjugated to Ruthenium(II)-tris-bipyridine-(4-methylsulfonate) NHS ester labeling reagent (Meso Scale Discovery, Gaithersburg, Md.) according to the manufacturer's instructions (Meso Scale Discovery, Gaithersburg, Md.). The conjugation reaction was performed by adding 30  $\mu$ L of distilled water reconstituted MSD SULFO-TAG™ stock solution to 200  $\mu$ L of 2 mg/mL  $\alpha$ -SNAP-25 polyclonal antibodies and incubating the reaction at room temperature for 2 hours in the dark. The labeled antibodies were purified using a standard spin column protocol and the protein concentration determined using a standard colorimetric protein assay. The absorbance of the  $\alpha$ -SNAP-25 antibody/MSD SULFO-TAG™ conjugate was measured at 455 nm using a spectrophotometer to determine the concentration in moles per liter. The detection antibody solution was stored at 4° C. until needed.

**[0327]** To prepare the solid phase support comprising the capture antibody that is specific for a SNAP-25 cleavage product, approximately 5  $\mu$ L of the appropriate  $\alpha$ -SNAP-25 monoclonal antibody solution (20  $\mu$ g/mL in 1× PBS) is added to each well of a 96-well MSD High Bind plate and the solution is allowed to air dry in a biological safety cabinet for 2-3 hours in order to liquid evaporate the solution. The capture antibody-bound wells were then blocked by adding 150  $\mu$ L of Blocking Buffer comprising 2% Amersham Blocking Reagent (GE Life Sciences, Piscataway, N.J.) and 10% goat serum (VWR, West Chester, Pa.) at room temperature for 2 hours. Blocked plates were sealed and stored at 4° C. until needed.

**[0328]** To detect the presence of a cleaved SNAP-25 cleavage product by ECL sandwich ELISA analysis, the Blocking Buffer from stored plates was aspirated from the wells, 25  $\mu$ L of a lysate from cells treated with re-targeted endopeptidase, as described above, was added to each well and the plates were incubated at 4° C. for overnight. Plate wells were washed three times by aspirating the cell lysate and rinsing each well three times with 200  $\mu$ L 1× PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, 25  $\mu$ L of 5  $\mu$ g/mL detection antibody solution comprising 2% Amersham Blocking Reagent in 1× PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate) was added to each well, the plate was sealed, and the sealed plate was incubated at room temperature at room temperature for 1 hour with shaking. After detection antibody incubation, the wells were washed three times with 200  $\mu$ L 1× PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing 150  $\mu$ L of 1× Read

Buffer (Meso Scale Discovery, Gaithersburg, MD) was added to each well and the plates were read using a SECTOR™ Imager 6000 Image Reader (Meso Scale Discovery, Gaithersburg, Md.). A ratio was calculated by dividing the signal obtained at the 10 nM dose for each antibody-pair by the signal obtained at the 0 nM dose for each antibody-pair (Table 24). These results indicated that among the twenty-six different combinations of antibody pairs tested, only three antibody pairs had signal-to-noise ratios above 10:1 for the higher dose tested: Pair No. 1 (2E2A6 mouse mAb and S9684 rabbit pAb), Pair No. 4 (3C1A5 mouse mAb and S9684 rabbit pAb), and Pair No. 18 (S9684 rabbit pAb and 2E2A6 mouse mAb). Antibody Pair 1 was chosen for further assay development.

aspirated from each well and replaced with fresh media containing either 0 (untreated sample) or one of the doses determined from a dose-response experiment for that re-targeted endopeptidase. After 24 hours incubation, the cells were washed and harvested.

**[0331]** To prepare the  $\alpha$ -SNAP-25 capture antibody solution, the  $\alpha$ -SNAP-25 monoclonal antibody contained in the ascites from hybridoma cell line 2E2A6 was purified using a standard Protein A purification protocol. To prepare the  $\alpha$ -SNAP-25 detection antibody solution,  $\alpha$ -SNAP-25 rabbit polyclonal antibody S9684 (Sigma, St. Louis, Mo.) was conjugated to Ruthenium(II)-tris-bipyridine-(4-methylsulfonate) NHS ester labeling reagent (Meso Scale Discovery, Gaithersburg, Md.) according to the manufacturer's instruc-

TABLE 24

Screening of $\alpha$ -SNAP-25 Antibody Combinations					
Antibody Pair No.	Capture Antibody	Detection Antibody	Detection SNAP-25 cleavage product	Detection SNAP-25 uncleaved substrate	Signal/Noise Ratio (10 nM/0 nM)
1	2E2A6 mouse mAb	S9684 rabbit pAb	Yes	No	26.6:1
2	2E2A6 mouse mAb	N-19 goat pAb	Yes	No	7.3:1
3	2E2A6 mouse mAb	H-50 rabbit pAb	Yes	No	0.9:1
4	3C1A5 mouse mAb	S9684 rabbit pAb	Yes	No	12.1:1
5	3C1A5 mouse mAb	N-19 goat pAb	Yes	No	1.9:1
6	3C1A5 mouse mAb	H-50 rabbit pAb	Yes	No	0.9:1
7	C-18 goat pAb	S9684 rabbit pAb	No	No	0.8:1
8	C-18 goat pAb	N-19 goat pAb	No	No	0.9:1
9	C-18 goat pAb	H-50 rabbit pAb	No	No	0.9:1
10	H-50 rabbit pAb	2E2A6 mouse mAb	Yes	No	0.9:1
11	H-50 rabbit pAb	C-18 goat pAb	No	No	1.0:1
12	N-19 goat pAb	2E2A6 mouse mAb	Yes	No	0.9:1
13	N-19 goat pAb	C-18 goat pAb	No	No	1.1:1
14	NTP 23 rabbit pAb	N-19 goat pAb	Yes	No	1.2:1
15	NTP 23 rabbit pAb	C-18 goat pAb	No	No	1.1:1
16	NTP 23 rabbit pAb	SP12 mouse pAb	Yes	No	1.3:1
17	NTP 23 rabbit pAb	H-50 rabbit pAb	Yes	No	1.1:1
18	S9684 rabbit pAb	2E2A6 mouse mAb	Yes	No	21.3:1
19	S9684 rabbit pAb	C-18 goat pAb	No	No	0.7:1
20	S9684 rabbit pAb	SMI-81 mouse mAb	Yes	Yes	1.2:1
21	SMI-81 mouse mAb	S9684 rabbit pAb	Yes	Yes	1.1:1
22	SMI-81 mouse mAb	N-19 goat pAb	Yes	Yes	1.0:1
23	SMI-81 mouse mAb	C-18 goat pAb	No	No	0.8:1
24	SP12 mouse pAb	C-18 goat pAb	No	No	1.0:1
25	MC-6050 mouse mAb	S9684 rabbit pAb	Yes	Yes	5.0:1
26	MC-6053 mouse mAb	S9684 rabbit pAb	Yes	Yes	7.1:1

### Example X

#### Immuno-Based method of Detecting Re-targeted Endopeptidase Having a BoNT/A Light Chain Enzymatic Activity Using ECL Sandwich ELISA

**[0329]** The following example illustrates immuno-based methods of detecting re-targeted endopeptidase activity by detecting a SNAP-25 cleavage product using an  $\alpha$ -SNAP-25 monoclonal antibody specific for a SNAP-25 cleavage product having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond by ECL sandwich ELISA.

**[0330]** To prepare a lysate from cells treated with a re-targeted endopeptidase having BoNT/A light chain enzymatic activity, a suitable density of cells from an established cell line was plated into the wells of 96-well tissue culture plates containing 100  $\mu$ L of the appropriate media. These cells were incubated in a 37° C. incubator under 5% carbon dioxide for about 24 hours. The media from the cells was

(Meso Scale Discovery, Gaithersburg, Md.). The conjugation reaction, purification of labeled  $\alpha$ -SNAP-25 antibody, concentration determination and storage were as described in Example VI.

**[0332]** To prepare the solid phase support comprising the capture antibody that is specific for a SNAP-25 cleaved product, approximately 5  $\mu$ L of  $\alpha$ -SNAP-25 monoclonal antibody 2E2A6 solution (20  $\mu$ g/mL in 1 $\times$  PBS) was added to each well of a 96-well MSD High Bind plate and the solution is allowed to air dry in a biological safety cabinet for 2-3 hours in order to liquid evaporate the solution. The capture antibody-bound wells were then blocked and used directly to detect re-targeted endopeptidase activity.

**[0333]** To detect the presence of a cleaved SNAP-25 product by ECL sandwich ELISA analysis, the Blocking Buffer from stored plates was aspirated from the wells, 25  $\mu$ L of a lysate from cells treated with re-targeted endopeptidase was added to each well and the plates were incubated at 4° C. for overnight. Plate wells were washed three times

by aspirating the cell lysate and rinsing each well three times with 200  $\mu$ L 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, 25  $\mu$ L of 5 pg/mL detection antibody solution comprising 2% Amersham Blocking Reagent in 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate) was added to each well, the plate was sealed, and the sealed plate was incubated at room temperature at room temperature for 1 hour with shaking. After detection antibody incubation, the wells were washed three times with 200  $\mu$ L 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing 150  $\mu$ L of 1 $\times$  Read Buffer (Meso Scale Discovery, Gaithersburg, Md.) was added to each well and the plates were read using a SECTOR™ Imager 6000 Image Reader (Meso Scale Discovery, Gaithersburg, Md.). The collected data was analyzed and the EC<sub>50</sub> calculated as described in Example VI. For opioid, re-targeted endopeptidases, these results indicated that on average 1.0 nM of Noc/A at the EC<sub>50</sub> was detected (a range of about 0.3 nM to about 2.0 nM) with a signal-to-noise ratio for the lower asymptote of about 15:1 to about 20:1 and a signal-to-noise ratio for the upper asymptote of about 180:1 to about 300:1.

#### Example XI

##### Immuno-Based Method of Detecting Re-Targeted Endopeptidase Activity Using CL Sandwich ELISA

**[0334]** The following example illustrates immuno-based methods of detecting re-targeted endopeptidase activity by detecting a SNAP-25 cleavage product using an  $\alpha$ -SNAP-25 monoclonal antibody specific for a SNAP-25 having a carboxyl-terminus at the P<sub>1</sub> residue of the BoNT/A cleavage site scissile bond by CL sandwich ELISA.

**[0335]** Lysate from cells treated with a re-targeted endopeptidase and the  $\alpha$ -SNAP-25 capture antibody solution will be prepared as described in Example VII.

**[0336]** To prepare the  $\alpha$ -SNAP-25 detection antibody solution,  $\alpha$ -SNAP-25 polyclonal antibody S9684 (Sigma, St. Louis, Mo.) will be conjugated to Horseradish peroxidase (HRP) according to the manufacturer's instructions (Pierce Biotechnology, Inc., Rockford, Ill.). The conjugation reaction will be performed by adding to 500  $\mu$ L of 1 mg/mL  $\alpha$ -SNAP-25 polyclonal antibodies to a vial containing lyophilized activated peroxidase, mixing the components, and then adding 10  $\mu$ L of sodium cyanoborohydride. This reaction mixture will be incubated at room temperature for 1 hour in a fume hood. After quenching the reaction, the labeled antibodies will be purified using a standard spin column protocol and the protein concentration will be determined using a standard colorimetric protein assay. The absorbance of the  $\alpha$ -SNAP-25 polyclonal antibody/HRP conjugate will be measured at 455 nm using a spectrophotometer to determine the concentration in moles per liter. The  $\alpha$ -SNAP-25 detection antibody solution will be stored at 4° C. until needed.

**[0337]** To prepare the solid phase support comprising the  $\alpha$ -SNAP-25 capture antibody that is specific for the SNAP-25 cleaved product, approximately 100  $\mu$ L of  $\alpha$ -SNAP-25 monoclonal antibody 2E2A6 solution (1 mg/mL in 1 $\times$  PBS) will be added to each well of a 96-well Greiner white plate and the plates will be incubated at 4° C. overnight, and then any excess antibody solution will be discarded. The capture antibody-bound wells will then be blocked by adding 150  $\mu$ L of Blocking Buffer comprising 2% Amersham Blocking

Reagent (GE Life Sciences, Piscataway, N.J.) and 10% goat serum (VWR, West Chester, Pa.) at room temperature for 1 hour. The blocking buffer will be discarded and the plates will be blotted dry on paper towels by inverting and tapping. The capture antibody-bound wells will then be blocked and will be used directly to detect re-targeted endopeptidase activity.

**[0338]** To detect the presence of a cleaved SNAP-25 product by CL sandwich ELISA analysis, 50  $\mu$ L of a lysate from cells treated with re-targeted-endopeptidase will be added to each well, the plate will be sealed, and the sealed plate will be incubated on a shaker rotating at 500 rpm at 4° C. for 2-4 hours to overnight. Plate wells will be washed three times by aspirating the cell lysate and rinsing each well three times with 200  $\mu$ L 1 $\times$  PBS, 0.05% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, 100  $\mu$ L of 1 mg/mL  $\alpha$ -SNAP-25 polyclonal antibody/HRP detection antibody solution comprising 2% Amersham Blocking Reagent in 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate) will be added to each well, the plate will be sealed, and the sealed plate will be incubated on a shaker rotating at 650 rpm at room temperature for 1 hour. After detection antibody incubation, the wells will be washed three times with 200  $\mu$ L 1 $\times$  PBS, 0.05% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing 100  $\mu$ L of SuperSignal ELISA Pico 1:1 mixture (Pierce Biotechnology, Inc., Rockford, Ill.) will be added to each well and the plates will be read using a luminometer (Molecular Devices, Sunnyvale, Calif.) at 395 nm. The collected data will be analyzed and the EC<sub>50</sub> will be calculated as described in Example VI.

#### Example XII

##### Immuno-Based Method of Detecting Re-Targeted Endopeptidase Activity Using Multiplex ECL Sandwich ELISA

**[0339]** The following example illustrates multiplex immuno-based methods of detecting re-targeted endopeptidase activity by detecting a SNAP-25 cleavage product using an  $\alpha$ -SNAP-25 monoclonal antibody specific for a SNAP-25 cleavage product and a second antibody pair for a different protein.

**[0340]** A re-targeted endopeptidase potency assay can be performed using a multiplex ECL sandwich ELISA. Such an assay is described in companion patent application Ester Fernandez-Salas, et al., Immuno-Based Botulinum Toxin Serotype A Activity Assays, U.S. patent application Ser. No. 12/403,531, which is hereby incorporated by reference in its entirety, and can be used using the cell lines and re-targeted endopeptidases and the corresponding cell lines disclosed in the present specification.

#### Example XIII

##### Immuno-Based Method of Detecting Re-Targeted Endopeptidase Activity Using Multiplex EC Sandwich ELISA

**[0341]** The following example illustrates multiplex immuno-based methods of detecting re-targeted endopeptidase activity by detecting a SNAP-25 cleavage product using an  $\alpha$ -SNAP-25 monoclonal antibody specific for a SNAP-25 cleavage product and a second antibody pair for a different protein.

**[0342]** A re-targeted endopeptidase potency assay can be performed using a multiplex EC sandwich ELISA. Such an assay is described in companion patent application Ester Fernandez-Salas, et al., Immuno-Based Botulinum Toxin Serotype A Activity Assays, U.S. patent application Ser. No. 12/403,531, which is hereby incorporated by reference in its entirety, and can be used using the cell lines and re-targeted endopeptidases and the corresponding cell lines disclosed in the present specification.

#### Example XIV

##### Immuno-Based Method to Detect Nanomolar Amounts of Retargeted Endopeptidases

**[0343]** The following example illustrates how to perform immuno-based methods of detecting Nanomolar amounts of retargeted endopeptidase activity.

##### 1. Immuno-Based Method of Detecting Retargeted Endopeptidases Using ECL Sandwich ELISA.

**[0344]** To prepare a lysate from cells treated with a retargeted endopeptidase, approximately 50,000 to 150,000 cells from an established cell line suitable for the assay were plated into the wells of 96-well tissue culture poly-D-lysine plates containing 100  $\mu$ L of the appropriate media (see Examples I and II). These cells were incubated in a 37° C. incubator under 5% carbon dioxide for 24 hours. The media from the cells was aspirated from each well and replaced with fresh media containing either 0 (untreated sample) and the appropriate dose response as described for each retargeted endopeptidase in this application. After a 24 hour incubation, the cells were washed and harvested or incubated for an additional two days without retargeted endopeptidase before harvesting. To harvest the cells, the medium was aspirated, washed with 1 $\times$  PBS, and lysed by adding 30  $\mu$ L of Lysis Buffer comprising 50 mM HEPES, 150 mM NaCl, 1.5 mM MgCl<sub>2</sub>, 1 mM EGTA, 1% Triton X-100 to each well, and the plate incubated on a shaker rotating at 500 rpm for 30 minutes at 4° C. The plate was centrifuged at 4000 rpm for 20 minutes at 4° C. to pellet cellular debris and the supernatant was transferred to a capture antibody coated 96-well plate to perform the detection step.

**[0345]** The  $\alpha$ -SNAP-25 capture antibody solution, the  $\alpha$ -SNAP-25 detection antibody solution, and the solid phase support comprising the capture antibody that is specific for a SNAP-25 cleaved product were prepared as described in Example VII.

**[0346]** To detect the presence of a cleaved SNAP-25 product by ECL sandwich ELISA analysis, the Blocking Buffer from stored plates was aspirated, 25-30  $\mu$ L of a lysate from cells treated with retargeted endopeptidase was added to each well and the plates were incubated at 4° C. for either 2 hours or 24 hours. Plate wells were washed three times by aspirating the cell lysate and rinsing each well three times with 200  $\mu$ L 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, 25  $\mu$ L of 5 pg/mL  $\alpha$ -SNAP-25 detection antibody solution comprising 2% Amersham Blocking Reagent in 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate) was added to each well, the plate was sealed, and the sealed plate was incubated at room temperature for 1 hour with shaking. After  $\alpha$ -SNAP-25 detection antibody incubation, the wells

were washed three times with 200  $\mu$ L 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, the plates were processed, collected data was analyzed, and the EC<sub>50</sub> calculated as described in Example VI. These results indicated that on average 1.0 nM of Noc/A at the EC<sub>50</sub> was detected when using SK-N-DZ clonal cell line #3 cells (a range of about 0.3 nM to about 2.0 nM) with a signal-to-noise ratio for the upper asymptote of about 20:1 to about 300:1. Moreover, on average 3.7 nM of Noc/A at the EC<sub>50</sub> was detected when using AGN P33 clonal cell line #6 cells (a range of about 2.0 nM to about 5.5 nM) with a signal-to-noise ratio for the upper asymptote of about 20:1 to about 500:1. For the SK12 cells that are specific for the retargeted endopeptidase containing a dynorphin A ligand, on average 8.4 nM of Dyn/A at the EC<sub>50</sub> was detected when using SK12 cells (a range of about 4.5 nM to about 10.0 nM) with a signal-to-noise ratio for the upper asymptote of about 10:1 to about 20:1. Additionally, on average 8.8 nM of TVEMP-galanin at the EC<sub>50</sub> was detected when using Neuro-2a clonal cell line #7 cells (a range of about 5.0 nM to about 15.5 nM) with a signal-to-noise ratio for the upper asymptote of about 20:1 to about 200:1. This method can also be performed in a multiplex fashion as described in Example IX.

**[0347]** Lysate from cells treated with a retargeted endopeptidase and the  $\alpha$ -SNAP-25 capture antibody solution will be prepared as described in Example VII. The  $\alpha$ -SNAP-25 detection antibody solution and solid phase support comprising the capture antibody that is specific for a SNAP-25 cleaved product will be prepared as described in Example VIII.

**[0348]** To detect the presence of a cleaved SNAP-25 product by CL sandwich ELISA analysis, 100  $\mu$ L of a lysate from cells treated with a retargeted endopeptidase will be added to each well, the plate will seal, and the sealed plate will be incubated on a shaker rotating at 500 rpm at 4° C. for either 2 hours or 24 hours. Plate wells will be washed three times by aspirating the cell lysate and rinsing each well three times with 200  $\mu$ L 1 $\times$  PBS, 0.05% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing, 100  $\mu$ L of 1 mg/mL  $\alpha$ -SNAP-25 polyclonal antibody/HRP detection antibody solution comprising 2% Amersham Blocking Reagent in 1 $\times$  PBS, 0.1% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate) will be added to each well, the plate will be sealed, and the sealed plate will be incubated on a shaker rotating at 650 rpm at room temperature for 1 hour. After detection antibody incubation, the wells will be washed three times with 200  $\mu$ L 1 $\times$  PBS, 0.05% TWEEN-20® (polyoxyethylene (20) sorbitan monolaurate). After washing 100  $\mu$ L of SuperSignal ELISA Pico 1:1 mixture (Pierce Biotechnology, Inc., Rockford, Ill.) will be added to each well and the plates will be read using a luminometer (Molecular Devices, Sunnyvale, Calif.) at 395 nm. The collected data will be analyzed and the EC<sub>50</sub> will be calculated as described in Example VI. This method can also be performed in a multiplex fashion as described in Example IX.

#### Example XV

##### Immuno-Based Method to Detect Neutralizing $\alpha$ -Retargeted-Endopeptidase Antibodies

**[0349]** The following example illustrates how to perform an immuno-based method that can detect the presence of neutralizing  $\alpha$ -Noc/A antibodies.

**[0350]** Noc/A, is currently being evaluated for treating painful conditions, some of them chronic. With repeated long-term treatment of Noc/A, a patient may develop neutralizing  $\alpha$ -Noc/A antibodies to the retargeted endopeptidase leading to immunoresistance. Neutralizing  $\alpha$ -Noc/A antibodies will inhibit retargeted endopeptidase activity by stopping the retargeted endopeptidase's uptake into neuronal and other target cells by binding to the targeting ligand and/or the translocation domain (HN) of the retargeted endopeptidase. There is not established assay to determine the presence of the neutralizing  $\alpha$ -Noc/A antibodies in patient's blood. It would be more cost and time efficient if a cell-based assay could be developed to detect neutralizing antibodies in patients treated with retargeted endopeptidases.

**[0351]** To detect the presence or absence of neutralizing  $\alpha$ -Noc/A antibodies, the immuno-based methods of determining retargeted endopeptidase activity disclosed in the present specification can be used. One way is to determine the amount of SNAP-25 cleavage product present after treatment with various concentrations of Noc/A using a Western blot detection method, the other way was to use an ECL sandwich ELISA detection method.

**[0352]** To prepare a sample comprising neutralizing  $\alpha$ -Noc/A antibodies, serum was isolated from blood of a monkey immunized with Noc/A and the antibodies were affinity purified. Rabbits were also immunized with the nociceptin variant peptide, the targeting ligand present in the Noc/A molecule, their serum collected, and the antibodies affinity purified (anti-nociceptin polyclonal antibodies).

**[0353]** To prepare a lysate from cells treated with a sample comprising Noc/A, SK-N-DZ clonal cell line #3 cells and AGN P33 clonal cell line #6 cells were seeded in poly-D-lysine 96-well plates for 16-18 hours. Anti-nociceptin pAb at 0-3  $\mu\text{g}/\text{mL}$  was diluted in RPMI SFM (with N2, B27, and NGF supplements) containing 1 nM of Noc/A and the mix was pre-incubated at room temperature for 1 hour. Then the solutions were added to the cells and incubated for 24 h before performing the ECL ELISA assay. This anti-nociceptin variant antibody totally blocked 1 nM Noc/A uptake at 1  $\mu\text{g}/\text{mL}$  (>90% inhibition) on both cell lines. Anti-Noc/A monkey polyclonal antibody was also assayed these cell lines. Cells were plated in a 96-well poly-D-lysine plate at 100,000 cells per well for 24 hours in RPMI growth media supplemented with N2, B27, and NGF. Anti-Noc/A polyclonal antibodies at 0-20  $\mu\text{g}/\text{mL}$  was diluted in media containing 1 nM Noc/A and the mix was pre-incubated at room temperature for 1 h. Then the mix was added to the cells and incubated for 24 h before performing the ECL ELISA assay. Up to 60% inhibition was seen at the higher concentrations of 6-20  $\mu\text{g}/\text{mL}$  of anti-Noc/A pAb on the SK-N-DZ cell line and about 30% on the AGN P33 clonal cell line #6 cell line. This may due to the fact that the anti-Noc/A polyclonal antibodies is not specific to the binding site and contains other antibodies that bind other parts of the molecule producing only partial blocking at the concentrations tested. Higher concentrations maybe needed to achieve complete blocking.

**[0354]** To detect the presence of a cleaved SNAP-25 product by Western blot analysis, the media will aspirated from each well, the cells suspended in 50  $\mu\text{L}$  of SDS-PAGE loading buffer, and then heated to 95° C. for 5 minutes. An aliquot from each harvested sample will be analyzed by Western blot as described in Example I, except that harvested samples will be separated by SDS-PAGE using 12% 26-well Criterion gels (Bio-Rad Laboratories, Hercules, Calif.), and the rabbit polyclonal  $\alpha$ -SNAP-25<sub>1,97</sub> antibody serum will be used as the primary antibody (see Example V). The results will reveal the lowest concentration of retargeted endopeptidase that will produce a detectable band of SNAP-25 cleavage product in the Western blot.

**[0355]** To detect the presence of a cleaved SNAP-25 product by ECL Sandwich ELISA, the media was removed from each well and the cells were lysed as described in Example VI. The  $\alpha$ -SNAP-25 capture antibody solution, the  $\alpha$ -SNAP-25 detection antibody solution, and the  $\alpha$ -SNAP-25 solid phase support were prepared as described in Example VIII. Supernatants were transferred to the  $\alpha$ -SNAP-25 solid phase support and an ECL sandwich ELISA assay was performed as detailed in Example VI. The collected data was analyzed and the  $\text{EC}_{50}$  calculated as described in Example VI, except that the  $\text{EC}_{50}$  is the serum dilution needed to inhibit the activity of the retargeted endopeptidase to  $\frac{1}{2}$  its maximum and the ratio of maximal signal ( $\text{Signal}_{\text{Max}}$ ) to minimum signal ( $\text{Signal}_{\text{Min}}$ ) was obtained by dividing the SNAP-25 cleavage product signal obtained with the highest dilution of antibody by the signal obtained with the lowest antibody dilution.

**[0356]** The results indicate that the presence of neutralizing  $\alpha$ -Noc/A antibodies in monkey serum and the presence of  $\alpha$ -nociceptin variant antibodies from rabbit could be detected. The activity of the Noc/A molecule incubated in affinity purified antibodies from the immunized animal decreased as the antibody dilution decreased. The same assay will be performed with the Dyn/A and the TVEMP-galanin compounds utilizing the cell lines specific for each compound to be tested.

#### Example XV

##### Development of a Cell-Based Assay for a Galanin Re-Targeted Endopeptidase

**[0357]** The following example illustrates how to identify established cell lines possessing the re-targeted endopeptidase uptake capacity required to develop a cell-based potency assay.

##### 1. Growth of Stock Culture of Candidate Cell Lines.

**[0358]** To grow the cell lines, a suitable density of cells from the cell line being tested were plated in a 162  $\text{cm}^2$  tissue culture flask containing 30 mL of a suitable growth medium (see Table 25), and grown in a 37° C. incubator under 5% or 10% carbon dioxide until cells reached the desired density.

TABLE 25

Summary table of all cell lines and their respective media.		
Cell type; description; source	Complete Media (CM) All from Invitrogen (unless otherwise specified)	Serum Free media (SFM) All from Invitrogen, (unless otherwise specified)
SiMa (Human neuroblastoma cell line, DSMZ# ACC 164, Braunschweig, Germany)	RPMI 1640 (90%) Fetal Bovine Serum (FBS, 10%)	RPMI 1640 (90%) NEAA (0.1 mM), HEPES (10 mM), Sodium Pyruvate (1 mM)
SiMa H1 (cloned cell line from SiMa cells)	NEAA (0.1 mM), HEPES (10 mM), Sodium Pyruvate (1 mM) Penicillin (100 U per ml) Streptomycin (100 µg per ml),	Penicillin (100 U per ml) Streptomycin (100 µg per ml) N2 supplement (1x) B27 supplement (1x)
Neuro-2a (Mouse neuroblastoma: (ATCC#CC1131, Manassas, VA,))	Earle's MEM (90%) Fetal Bovine Serum 10%) NEAA (0.1 mM), HEPES (10 mM), Sodium Pyruvate (1 mM), Penicillin (100 U per ml), Streptomycin (100 µg per ml)	EMEM (90%) NEAA (0.1 mM), HEPES (10 mM), Sodium Pyruvate (1 mM), Penicillin (100 U per ml), Streptomycin (100 µg per ml)
PC-12 Rat Pheochromocytoma (ATCC # CRL-1721)	RPMI 1640 (90%) Dialyzed FBS (5%) Horse serum (10%) HEPES (10 mM) Sodium Pyruvate (1 mM) D-glucose (0.5%, Sigma) Penicillin (100 U per ml); Streptomycin (100 µg per ml) N2 supplement (1x)	Differentiation media: RPMI 1640 (90%) HEPES (10 mM) Sodium Pyruvate (1 mM) D-glucose (0.5%, Sigma) Penicillin (100 U per ml); Streptomycin (100 µg per ml) N2 supplement (1x) Bovine serum albumin (0.2% w/v) NGF (50 ng per ml, Promega)
P19 Mouse embryonic carcinoma (ATCC #CRL-1825)	Alpha MEM (90%) Bovine Calf Serum (7.5%) FBS (2.5%) Penicillin (100 U per ml); Streptomycin (100 µg per ml)	Alpha MEM (90%) FBS (2.5%) Penicillin (100 U per ml); Streptomycin (100 µg per ml)

NEAA: Non-Essential Amino Acids, MEM: Minimum Essential Media. DMEM: Dulbecco's MEM. EMEM—Earle's MEM. Please note PC-12 cells were differentiated in differentiation media and not SFM.

## 2. Screening of Commercial Cell Lines for Sensitivity to Galanin TVEMP-Galanin Compounds

**[0359]** Commercial cell lines were screened for their sensitivity to TVEMP-galanin compounds as measured by the cleavage of SNAP25 after treatment with the corresponding compounds. Various TVEMP-galanin compounds were used for screening and testing. PC-12, Neuro-2a, SiMa, and P19 cells were plated in serum free media for three days or in CM for one day. These differentiated and naive cells were treated for 18 hours with TVEMP-galanin Batch A at concentrations of 0 and 75 nM. TVEMP-galanin Batch A showed activity in both PC-12 and Neuro-2a cells as seen by the increased presence of cleaved SNAP25, and Neuro-2a cells in the differentiated condition are more sensitive to TVEMP compounds with galanin ligand, than the naive cells. The rank order in activity of the cells shows PC-12 have the most

activity, followed by Neuro-2a and finally SiMa cells. It was necessary to determine if the uptake was specific for these galanin-retargeted compounds and therefore it was important to test the cells with other compounds that do not contain the galanin ligand. Noc/A is a retargeted compound which contains a nociceptin variant ligand, and LH<sub>N</sub>/A (a negative control) a compound that lacks the binding domain. The uptake of LH<sub>N</sub>/A is non-specific and should have significantly lower activity than the TVEMP-galanin compound if the cell line possess specific uptake for the retargeted compound. The Noc/A compound has been shown previously to have specific uptake in the SiMa cells and will be used as a baseline to test the cell lines. A favorable cell line should have low uptake of the LH<sub>N</sub>/A and the Noc/A compound and high uptake of the TVEMP-galanin compound. Table 26 displays the results from this experiment.

TABLE 26

Screening of PC-12, Neuro-2a, and SiMa cells at different conditions using TVEMP-galanin.					
		TVEMP-galanin Batch A	TVEMP-galanin Batch B	LH <sub>N</sub> /A	Noc/A
Conc. (mg/mL)		0.168	0.175	1.63	1.00
EC <sub>50</sub>	PC-12, naïve	73.4 ± 10.7	105.6 ± 16.0	>200	72.9 ± 26.9
Values	SiMa, naïve	138.6 ± 43.9	133.8 ± 24.2	>300	48.3 ± 18.1
(nM)	Neuro-2a, naïve	122.4 ± 15.7	116 ± 17.5	>200	>150
	SiMa, Dif O/N	>400	>150	>400	16.1 ± 11.9

TABLE 26-continued

Screening of PC-12, Neuro-2a, and SiMa cells at different conditions using TVEMP-galanin.				
	TVEMP-galanin Batch A	TVEMP-galanin Batch B	LH <sub>N</sub> /A	Noc/A
Neuro-2a Dif 4 d		34.5 ± 7.5	39.7 ± 5.6	105.9 ± 44.3
SiMa, Dif 4 d	101.8 ± 20.5	65.3 ± 7.8	>150	88.7 ± 23.3

TVEMP-galanin Batches A and B, and LH<sub>N</sub>/A and Noc/A controls testing on various cell lines and growth/differentiation conditions. Summary chart showing details of each compound tested plus EC<sub>50</sub> values.

**[0360]** The results show that TVEMP-galanin Batch A and TVEMP-galanin Batch B had plots or EC<sub>50</sub> values that were either similar to, or only 1-2 fold more active than the negative controls in the cell lines tested. This data implies that the native cells are not sensitive enough and that these cells will have to be transfected with the plasmids encoding galanin receptor proteins GalR1 or GalR2 receptors.

### 3. Stable Transfection of PC-12, Neuro-2a, and SiMa Cells with GalR.

**[0361]** One day before transfection, cells were seeded at densities of  $0.5 \times 10^6$  cells/well in either a 6-well Collagen IV coated plate (Cat #354554: BD Biosciences) (SiMa, PC-12) or a 6 well Costar plate (Cat #3516: Corning) (Neuro-2a). Transfections were performed by diluting 12  $\mu$ l of Lipofectamine™ 2000 (Cat #52758, Invitrogen) in 250  $\mu$ l Opti-MEM® I Reduced Serum Medium (Cat #3195, Invitrogen) followed by incubation at room temperature for 5 min. Four micrograms of GalR plasmid DNA was mixed with 0.4  $\mu$ g pAdVantage™ vector (1 mg per ml, Cat #E1711, Promega) in 250  $\mu$ l Opti-MEM® I Reduced Serum Media for 5 minutes. After 5 minutes of incubation, the diluted Lipofectamine™ 2000 and the diluted plasmids DNA were mixed and incubated for an additional 20 min at room temperature, for complex formation. In the meantime, the cells were washed with OPTI-MEM® and 0.5 ml OPTI-MEM® was added to each well. After the 20 minute incubation, 0.5 ml containing the complexes of diluted Lipofectamine™ 2000 and diluted plasmids DNA was carefully added to the wells containing cells in 0.5 ml OPTI-MEM®. The plate was incubated at 37° C. for 5 hours, after which 1 ml of complete media was added. The next day, the medium was replaced with growth media for 48 hours. On day 4, after cells were recovered from the transfection, the growth media was replaced with fresh growth media containing Geneticin® (Cat #10131: Invitrogen) at 0.5 mg per ml (1:100 dilution) and incubated for an additional 3 days. On day 7 post transfection, the cells were transferred to a 75 cm Collagen IV flask (Cat #35423: BD Biosciences) containing growth medium and geneticin (0.5 mg per ml, 1:100 dilution). On this transfer, approximately 90% of the cells were dead and were removed during the media change. Growth media containing geneticin (0.5 mg per ml, 1:100 dilution) was changed every two days till day 21.

**[0362]** For the selection of stable cells able to take up galanin TVEMP compounds, the parameters were to screen for clones that produced the highest percentage of SNAP25 cleavage with TVEMP-galanin treatment in the ECL Sandwich ELISA using monoclonal 2E2A6 coated plates for capture and polyclonal SNAP25 (Sigma Cat #S9684) sulfotagged antibody for detection. The EC<sub>50</sub> values in Table 27 show that TVEMP-galanin Batch D exhibits at least 10-fold greater uptake than the negative control in the SiMa and

Neuro-2a cells transfected with GalR1 and GalR2, and only 2-4 fold greater uptake in transfected PC-12 cells. Since the PC-12 transfected cells seem to have lower sensitivity and specificity than the SiMa and Neuro-2a cells they will not be cloned. Also, since the galanin 1-16 mer ligand in the TVEMP-Galanin compounds binds to GALR1 receptor with more affinity than the GALR2, only cells transfected with GALR1 will be cloned. The figure also shows that TVEMP-galanin Batches C and D exhibit 9-10 fold greater uptake than both LH<sub>N</sub>/A and the retargeted nociceptin compound TVEMP-nociceptin in the Neuro-2a GalR1.

TABLE 27

Test of stably transfected but non-clonal populations of SiMa, Neuro-2a, and PC12 transfected with GalR1 or GalR2 receptors				
	TVEMP- galanin C	TVEMP- galanin D	LH <sub>N</sub> /A	TVEMP- nociceptin
Conc. (mg/mL)	1.260	0.303	1.46	1.00
EC <sub>50</sub> Values (nM)				
SiMa GalR1		36.2 ± 8.6	>300	
SiMa GalR2		26.6 ± 6.7	>300	
PC-12 GalR1		64.1 ± 19.5	202.7	
PC-12 GalR2		>150	>300	
Neuro-2a GalR1	32.2 ± 3.3	40.8 ± 6.0	>300	>300
Neuro-2a GalR2	35.2 ± 3.1	46.0 ± 6.1	>300	>300

**[0363]** The non-clonal selected populations are not a good population of cells to use on a regular basis because they contain a mix of cells expressing different levels of receptor and these populations can change overtime. To obtain stable cell lines derived from single cells, a dilutional cloning approach was initiated. On day 21, the transfected cells were trypsinized, needle-dissociated and counted. The remaining transfected cell lines were frozen for future use. The cells were serially diluted to 10 cells per ml in growth media containing geneticin (0.5 mg per ml, 1:100 dilution). Either 2x96-well Collagen IV coated plates (SiMa, PC-12) or 2x96-well Costar plates (Neuro-2a) were plated at 100  $\mu$ l per well to achieve 1 cell per well density. The plates were returned to the incubator and left untouched for two weeks for colony formation. After two weeks (day 35), the wells were carefully checked for the presence of single colonies formed on the bottom of the well (the entire well was carefully checked for multiple colonies). When a well was identified with a single cluster of cells, that entire well was carefully scrutinized to make sure that one and only one cluster of cells was present. A picture of that single cluster was taken. If there were any queries about additional clusters, the well was not selected. On day 36, the clones that were selected were detached with TrypLE and 0.5 ml of complete media containing geneticin (0.5 mg per ml, 1:100

dilution) was added to stop the trypsin reaction. This entire volume was transferred to 6-well plates and further diluted with 3.0 ml additional complete media containing geneticin (0.5 mg per ml, 1:100 dilution). The clones were allowed to grow to 90% confluence, then trypsinized again and transferred to 75 cm Collagen IV or Costar flasks with 10.0 ml complete media containing geneticin (0.5 mg per ml, 1:100 dilution). Once the cells were 90% confluent again, the cells were used to either fill three cryovials for frozen storage or used for screening in the ELISA assay for galanin retargeted compounds.

**[0364]** The reference compound TVEMP-galanin Batch C was used to test these clones using two operators performing independent tests. The SiMa GalR1 clones grew slowly and were not available for testing at this time. Fortunately, the Neuro-2a clones grew faster, and soon sufficient quantities of 8 of the 12 clones were available for testing. These Neuro-2a GalR1 clonal cells were tested with a full dose range of TVEMP-galanin compounds (0-300 nM) and the results of nine of these clones is shown below. The remaining four clones grew very slowly and were not tested. The selected but non-clonal parental cells were plated along with the clones to use as a benchmark. Table 28 shows the activity of each of the eight clones together with the selected non-clonal Neuro-2a GalR1 cells, when tested with TVEMP-galanin compound. Out of the eight clones tested, only clones #4, 7 and 12 showed good uptake of the TVEMP-galanin compound with acceptable EC<sub>50</sub> values. Neuro-2a GalR1 clones #1, 3 and 10 did not take up the TVEMP-galanin compound, while clones #5, 11 and 13 together with the non-clonal population generated very high EC<sub>50</sub> values and no further testing was done with these cells.

TABLE 28

Results of screening Neuro-2a GalR1 single-cell derived clones with the TVEMP-galanin Batch C.				
		EC50 ± Std. error (nM)		
Plate	Cell type	Operator 1	Operator 2	
1	N2A Non-clonal	82.1 ± 9.6	92.0 ± 10.8	
1	N2A GALR1 Clone #1	>300	>300	
1	N2A GALR1 Clone #3	>300	>300	
1	N2A GALR1 Clone #4	39.7 ± 3.4	39.4 ± 6.6	
2	N2A Non-clonal	211.2 ± 167.7	116.0 ± 26.8	
2	N2A GALR1 Clone #5	202.6 ± 82.9	113.0 ± 18.1	
2	N2A GALR1 Clone #7	23.1 ± 3.3	15.5 ± 1.8	
2	N2A GALR1 Clone #10	>300	>300	
3	N2A GALR1 Clone #7	20.3 ± 1.6	38.0 ± 6.3	
3	N2A GALR1 Clone #11	270.0 ± 243	247.0 ± 101	
3	N2A GALR1 Clone #12	43.2 ± 5.2	57.5 ± 14.3	
3	N2A GALR1 Clone #13	144.1 ± 143	184.7 ± 15.6	

#### 4. Characterization of GalR1 Expression in the Clonal Cell Lines

**[0365]** The screening of the clones showed that only clones #4, 7, and 12 are more sensitive than the non-clonal cells. Messenger RNA (mRNA) was extracted from these 3 clones as well as the non-transfected parental and stably transfected non-clonal Neuro-2a cells for characterization by RT-PCR using the RT-PCR conditions described in Example V and the primers described in Table 29.

TABLE 29

Specific GALR1 and GALR2 primers		
Name	Sequence	SEQ ID NO:
GALR1 fwd	5' 'CCCCATCATGTCCATCCACCT 3'	150
GALR1 rev	5' ATGGGGTTCCACCGAGGAGTT 3'	151
GALR2 fwd	5' CATCGTGGCGGTGCTTTT 3'	152
GALR2 rev	5' AGCGGGAAGCGACCAAAC 3'	153

**[0366]** The results in Table 30 show that the transfected non-clonal cells and clones have much greater amounts of GALR1 mRNA than the parental cells. In the TVEMP-Galanin cell screening, Clone #7 was shown to be the most sensitive to TVEMP-galanin. Clone #7 also is shown to have the highest amount of GALR1 mRNA according to Table 30. The CT values for Neuro-2a GalR1 clone 7 (Neuro-2a #7) was the lowest, followed by clone 4 and then clone 12. The non-clonals tested at this time provided a CT close to clone 12, however, these cells contain a constantly changing population of cells containing varying concentrations of GalR1 receptor, and therefore were not considered a good population for future work. Of the three clones with low EC<sub>50</sub>'s, Neuro-2a clone GalR1 clone #12 (Neuro-2a #12) grew the fastest, followed by Neuro-2a clone #7 and lastly Neuro-2a clone #4. In addition to its slow growth rate, Neuro-2a clone #4 was not tested further because the sensitivity of Neuro-2a clone #7 was much better than for clone #4.

TABLE 30

Large differences in GALR1 mRNA in Neuro-2a transfected clonal cells vs. transfected non-clonal and parental cells.					
Cell line	Parental	Non-Clonal	Clone 4	Clone 7	Clone 12
Ave CT	32.0	21.7	20.8	19.3	21.6
fold mRNA change	1.0	1269.5	2418.7	6793.8	1332.6

#### 5. Comparison of the Sensitivity and Specificity of Neuro-2a clones #7 and #12 with TVEMP-Galanin Compounds

**[0367]** The two clones were tested side-by-side in an attempt to identify the most sensitive and selective of the two, so that data could be confidently collected from the best performing clone. Table 31 shows the results of these two clones when treated with TVEMP-galanin Batch C and LH<sub>α</sub>/A for sensitivity and selectivity respectively. Both clones exhibit high Signal-to-Noise ratios. Neuro-2a Clone #7 has an EC<sub>50</sub> of 5.5 nM while the EC<sub>50</sub> for Neuro-2a clone #12 is 68.4 nM. The Neuro-2a clone #12 has to be tested with a dose range of 0-300 nM, while the Neuro-2a Clone #7 can be tested with a dose range of 0-30 nM to elicit a plateau at the highest concentration used. Both clones show good separation between the LH<sub>α</sub>/A and TVEMP-galanin Batch C, Neuro-2a Clone #12 shows some non-specific uptake at the high concentrations, while Neuro-2a Clone #7 does not. As seen in the tabulated results, the range for testing with Neuro-2a #7 cells is 10-fold lower than that for Neuro-2a #12 cells resulting in a 10-fold less compound being used for Neuro-2a #7 than Neuro-2a #12. Neuro-2a #7 is 8-fold more selective than Neuro-2a clone #12 when

LH<sub>N</sub>/A was used as a comparison. The Signal-to-Noise ratio is over 100 for both clones, however a ratio of 10 would be sufficient to develop a cell based potency assay. The EC<sub>50</sub> for the Neuro-2a #7 clone is 5.5 nM about 12-fold lower than that for Neuro-2a #12, whose EC<sub>50</sub> is 68.4 nM. The lower dose-range for testing, the 24-fold selectivity over LH<sub>N</sub>/A, the high signal-to-noise ratio, the excellent sensitivity resulting in low EC<sub>50</sub>, and the low amount of protein required for each test, all imply that Neuro-2a Clone #7 would be the clone to go forward with the cell-based potency assay for use in determining potency ratios for TVEMP-galanin compounds.

TABLE 31

Comparison of characteristics of Neuro-2a clone #7 and #12.		
	Neuro-2a #7	Neuro-2a #12
Range	0-30 nM	0-300 nM
Selectivity	24-fold	3-fold
Signal-to-Noise ratio	190	547
Percent of max LH <sub>N</sub> /A signal over max TVEMP-Gal signal	4.3%	37.6%
EC <sub>50</sub>	5.5 nM	68.4 nM
Protein required	~1 µg	~10 µg

Neuro-2a # 7 and #12 were treated with TVEMP-galanin Batch C and LH<sub>N</sub>/A for 16 hours in CM. Activity was detected using ECL-ELISA.

## Example XVI

#### Generation of Clonal Cell Lines Overexpressing the KOR-1 Receptor for Dynorphin A Retargeted Endopeptidase Uptake

[0368] The following example illustrates how to characterize and compare several clonal cell lines originated from an established cell line transfected with the target receptor and subsequent cloning of the cell line. This specific example refers to the identification and characterization of clonal cell lines transfected with hKOR-1 that were first described in Example III, Table 9.

[0369] Four of the AGN P33-KOR clones (clones number 8, 9, 10, and 12 Table 9 in Example III) were selected and tested with Dyn/A with a full dose response of 0-150 nM. At the same time, two SiMa-KOR clones (clones number 12, and 16 from Table 9 in Example III) selected and tested with Dyn/A with a full dose response of 0-150 nM. In this experiment, AGN P33-KOR clones 8, 9, and 12 produced very low uptake and were therefore discarded; AGN P33-KOR clone 10 displayed good uptake and an EC<sub>50</sub> of 30.3 nM was obtained. The two SiMa-KOR clones tested displayed good uptake and an EC<sub>50</sub> of 26.6 nM was obtained for clone 16 and an EC<sub>50</sub> of 11.8 nM was obtained for clone 12. These three clones were then tested for sensitivity and selectivity by comparing the uptake of the target Dyn/A compound against the negative control LH<sub>N</sub>/A that lacks a targeting ligand and the Noc/A control. The comparison of the three clones and the parental SiMa cells utilizing a full dose response of 0-150 nM is summarized in Table 32.

TABLE 32

Cell Line	EC <sub>50</sub> Dyn/A (nM)	EC <sub>50</sub> LHN/A (nM)	EC <sub>50</sub> Noc/A (nM)
SiMa Parental	>100	>100	5.4
AGN P33-KOR clone 10	9.7	>150	9.4

TABLE 32-continued

Cell Line	EC <sub>50</sub> Dyn/A (nM)	EC <sub>50</sub> LHN/A (nM)	EC <sub>50</sub> Noc/A (nM)
SiMa-KOR clone 16	10.6	>100	1.6
SiMa-KOR clone 12	4.65	>150	19.7

[0370] There was a marked increase in Dyn/A uptake in the KOR-1 transfected clones treated with Dyn/A while the parental SiMa cells showed minimal uptake of the compound (uptake was similar to the negative control LH<sub>N</sub>/A). There is some Noc/A in all the cell lines including parental SiMa cells. This is not surprising as uptake of Noc/A in SiMa cells was observed during the assay development for this retargeted compound. Moreover, Noc/A uptake is best in the AGN P33 cell line that was specifically derived for this retargeted endopeptidase. The difference between Noc/A uptake and Dyn/A compound uptake is greater in the clonal SiMa-KOR clone 12 (SK12) cells. In all the graphs, activity of the negative control, LH<sub>N</sub>/A, is minimal, showing that in the absence of the binding domain there is no specific uptake in these cell lines and the lowest was in the SK12 cells showing that the uptake of the Dyn/A compound is highly specific. From these results, the SK12 clone was selected for future optimization and characterization.

[0371] Optimization studies were performed with the SK12 cells in order to develop a robust, specific and sensitive assay. Several parameters were assayed including plating media and plating densities, treatment media, and treatment time. A summary of the data obtained during the optimization is provided in Table 33.

TABLE 33

Medium used		Treat-ment	Cells/well				
plating	treating	time	25000	50000	75000	100000	150000
complete	complete	6 hr + o/n	51.3	76	13.4	9.2	n/a
complete	complete	16 hr	21.3	19.0	4.96	4.64	n/a
complete	complete	16 hr	n/a	n/a	n/a	2.1	15.3
serum free	serum free	16 hr	n/a	n/a	n/a	9.0	12.1
complete	serum free	16 hr	n/a	10.3	5.4	8.97	8.38
complete	complete	16 hr	n/a	7.7	4.86	13.72	11.26
serum free	serum free	16 hr	n/a	11.2	8.5	8.4	9.2

[0372] Table B shows that cells plated at 100,000 cells per well in CM and treated with compounds in CM showed more variability in EC<sub>50</sub> values from one experiment to the next (4.6; 1.2 and 13.72 nM) while cells plated at 100,000 cells per well in SFM and treated with compounds diluted in SFM provided the best curves and consistent EC<sub>50</sub> values (9.0 and 8.4 nM). In future, cells would be plated at 100,000 cells per well in SFM and treated with compounds in SFM too.

[0373] SK12 plated on PDL plates at 100,000 cells per well in SFM for 24 hours, followed by treatment in SFM for 16 hours yielded the lowest EC<sub>50</sub> value of 8.4+/-1.1 nM and a Signal-to Noise ratio of 12. Both these values would be acceptable for future use of this cell in CBPA.

#### Characterization of SK12 Cells with the Saturation Binding Assay

**[0374]** The saturation Binding assay utilized here was described in detail in Example V. Saturation binding studies were performed using the KOR-1 antagonist <sup>3</sup>H-diprenorphine to evaluate binding. The total, specific, and non-specific binding were measured in several experiments. A saturation binding curve of <sup>3</sup>H-diprenorphine with the receptor was generated from two independent experiments. It appears that about 25% binding is non-specific and 75% specific binding of the molecule to the receptor. The affinity of the molecule to the receptor is adequate at 6.5 nM. The Bmax indicates that there are 23 fmol KOR-1 receptors per cell on the SK12 cells.

**[0375]** Unless otherwise indicated, all numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

**[0376]** The terms “a,” “an,” “the” and similar referents used in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. Recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of the invention.

**[0377]** Groupings of alternative elements or embodiments of the invention disclosed herein are not to be construed as limitations. Each group member may be referred to and claimed individually or in any combination with other members of the group or other elements found herein. It is anticipated that one or more members of a group may be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

**[0378]** Certain embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Of course, variations on these described embodiments will become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventor expects skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

**[0379]** Specific embodiments disclosed herein may be further limited in the claims using consisting of or consisting essentially of language. When used in the claims, whether as filed or added per amendment, the transition term “consisting of” excludes any element, step, or ingredient not specified in the claims. The transition term “consisting essentially of” limits the scope of a claim to the specified materials or steps and those that do not materially affect the basic and novel characteristic(s). Embodiments of the invention so claimed are inherently or expressly described and enabled herein.

**[0380]** Furthermore, numerous references have been made to patents and printed publications throughout this specification. Each of the above-cited references and printed publications are individually incorporated herein by reference in their entirety.

**[0381]** In closing, it is to be understood that the embodiments of the invention disclosed herein are illustrative of the principles of the present invention. Other modifications that may be employed are within the scope of the invention. Thus, by way of example, but not of limitation, alternative configurations of the present invention may be utilized in accordance with the teachings herein. Accordingly, the present invention is not limited to that precisely as shown and described.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 153  
 <210> SEQ ID NO 1  
 <211> LENGTH: 903  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Retargeted endopeptidase

-continued

&lt;400&gt; SEQUENCE: 1

Met Gly Ser Met Glu Phe Val Asn Lys Gln Phe Asn Tyr Lys Asp Pro  
 1 5 10 15  
 Val Asn Gly Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn Ala Gly Gln  
 20 25 30  
 Met Gln Pro Val Lys Ala Phe Lys Ile His Asn Lys Ile Trp Val Ile  
 35 40 45  
 Pro Glu Arg Asp Thr Phe Thr Asn Pro Glu Glu Gly Asp Leu Asn Pro  
 50 55 60  
 Pro Pro Glu Ala Lys Gln Val Pro Val Ser Tyr Tyr Asp Ser Thr Tyr  
 65 70 75 80  
 Leu Ser Thr Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly Val Thr Lys  
 85 90 95  
 Leu Phe Glu Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met Leu Leu Thr  
 100 105 110  
 Ser Ile Val Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr Ile Asp Thr  
 115 120 125  
 Glu Leu Lys Val Ile Asp Thr Asn Cys Ile Asn Val Ile Gln Pro Asp  
 130 135 140  
 Gly Ser Tyr Arg Ser Glu Glu Leu Asn Leu Val Ile Ile Gly Pro Ser  
 145 150 155 160  
 Ala Asp Ile Ile Gln Phe Glu Cys Lys Ser Phe Gly His Glu Val Leu  
 165 170 175  
 Asn Leu Thr Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile Arg Phe Ser  
 180 185 190  
 Pro Asp Phe Thr Phe Gly Phe Glu Glu Ser Leu Glu Val Asp Thr Asn  
 195 200 205  
 Pro Leu Leu Gly Ala Gly Lys Phe Ala Thr Asp Pro Ala Val Thr Leu  
 210 215 220  
 Ala His Glu Leu Ile His Ala Gly His Arg Leu Tyr Gly Ile Ala Ile  
 225 230 235 240  
 Asn Pro Asn Arg Val Phe Lys Val Asn Thr Asn Ala Tyr Tyr Glu Met  
 245 250 255  
 Ser Gly Leu Glu Val Ser Phe Glu Glu Leu Arg Thr Phe Gly Gly His  
 260 265 270  
 Asp Ala Lys Phe Ile Asp Ser Leu Gln Glu Asn Glu Phe Arg Leu Tyr  
 275 280 285  
 Tyr Tyr Asn Lys Phe Lys Asp Ile Ala Ser Thr Leu Asn Lys Ala Lys  
 290 295 300  
 Ser Ile Val Gly Thr Thr Ala Ser Leu Gln Tyr Met Lys Asn Val Phe  
 305 310 315 320  
 Lys Glu Lys Tyr Leu Leu Ser Glu Asp Thr Ser Gly Lys Phe Ser Val  
 325 330 335  
 Asp Lys Leu Lys Phe Asp Lys Leu Tyr Lys Met Leu Thr Glu Ile Tyr  
 340 345 350  
 Thr Glu Asp Asn Phe Val Lys Phe Phe Lys Val Leu Asn Arg Lys Thr  
 355 360 365  
 Tyr Leu Asn Phe Asp Lys Ala Val Phe Lys Ile Asn Ile Val Pro Lys  
 370 375 380  
 Val Asn Tyr Thr Ile Tyr Asp Gly Phe Asn Leu Arg Asn Thr Asn Leu

-continued

---

385	390	395	400
Ala Ala Asn Phe	Asn Gly Gln Asn Thr Glu Ile Asn Asn Met Asn Phe		
	405	410	415
Thr Lys Leu Lys	Asn Phe Thr Gly Leu Phe Glu Phe Tyr Lys Leu Leu		
	420	425	430
Cys Val Asp Gly	Ile Ile Thr Ser Lys Thr Lys Ser Asp Asp Asp Asp		
	435	440	445
Lys Phe Gly Gly	Phe Thr Gly Ala Arg Lys Ser Ala Arg Lys Arg Lys		
	450	455	460
Asn Gln Ala Leu	Ala Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly		
	465	470	475
Gly Gly Gly Ser	Ala Leu Val Leu Gln Cys Ile Lys Val Asn Asn Trp		
	485	490	495
Asp Leu Phe Phe	Ser Pro Ser Glu Asp Asn Phe Thr Asn Asp Leu Asn		
	500	505	510
Lys Gly Glu Glu	Ile Thr Ser Asp Thr Asn Ile Glu Ala Ala Glu Glu		
	515	520	525
Asn Ile Ser Leu	Asp Leu Ile Gln Gln Tyr Tyr Leu Thr Phe Asn Phe		
	530	535	540
Asp Asn Glu Pro	Glu Asn Ile Ser Ile Glu Asn Leu Ser Ser Asp Ile		
	545	550	555
Ile Gly Gln Leu	Glu Leu Met Pro Asn Ile Glu Arg Phe Pro Asn Gly		
	565	570	575
Lys Lys Tyr Glu	Leu Asp Lys Tyr Thr Met Phe His Tyr Leu Arg Ala		
	580	585	590
Gln Glu Phe Glu	His Gly Lys Ser Arg Ile Ala Leu Thr Asn Ser Val		
	595	600	605
Asn Glu Ala Leu	Leu Asn Pro Ser Arg Val Tyr Thr Phe Phe Ser Ser		
	610	615	620
Asp Tyr Val Lys	Lys Val Asn Lys Ala Thr Glu Ala Ala Met Phe Leu		
	625	630	640
Gly Trp Val Glu	Gln Leu Val Tyr Asp Phe Thr Asp Glu Thr Ser Glu		
	645	650	655
Val Ser Thr Thr	Asp Lys Ile Ala Asp Ile Thr Ile Ile Ile Pro Tyr		
	660	665	670
Ile Gly Pro Ala	Leu Asn Ile Gly Asn Met Leu Tyr Lys Asp Asp Phe		
	675	680	685
Val Gly Ala Leu	Ile Phe Ser Gly Ala Val Ile Leu Leu Glu Phe Ile		
	690	695	700
Pro Glu Ile Ala	Ile Pro Val Leu Gly Thr Phe Ala Leu Val Ser Tyr		
	705	710	715
Ile Ala Asn Lys	Val Leu Thr Val Gln Thr Ile Asp Asn Ala Leu Ser		
	725	730	735
Lys Arg Asn Glu	Lys Trp Asp Glu Val Tyr Lys Tyr Ile Val Thr Asn		
	740	745	750
Trp Leu Ala Lys	Val Asn Thr Gln Ile Asp Leu Ile Arg Lys Lys Met		
	755	760	765
Lys Glu Ala Leu	Glu Asn Gln Ala Glu Ala Thr Lys Ala Ile Ile Asn		
	770	775	780
Tyr Gln Tyr Asn	Gln Tyr Thr Glu Glu Glu Lys Asn Asn Ile Asn Phe		
	785	790	795
			800

-continued

---

```

Asn Ile Asp Asp Leu Ser Ser Lys Leu Asn Glu Ser Ile Asn Lys Ala
      805                      810                      815
Met Ile Asn Ile Asn Lys Phe Leu Asn Gln Cys Ser Val Ser Tyr Leu
      820                      825                      830
Met Asn Ser Met Ile Pro Tyr Gly Val Lys Arg Leu Glu Asp Phe Asp
      835                      840                      845
Ala Ser Leu Lys Asp Ala Leu Leu Lys Tyr Ile Tyr Asp Asn Arg Gly
      850                      855                      860
Thr Leu Ile Gly Gln Val Asp Arg Leu Lys Asp Lys Val Asn Asn Thr
      865                      870                      875                      880
Leu Ser Thr Asp Ile Pro Phe Gln Leu Ser Lys Tyr Val Asp Asn Gln
      885                      890                      895
Arg Leu Leu Ser Thr Leu Asp
      900

```

```

<210> SEQ ID NO 2
<211> LENGTH: 908
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Retargeted endopeptidase

```

```

<400> SEQUENCE: 2

```

```

Met Gly Ser Met Glu Phe Val Asn Lys Gln Phe Asn Tyr Lys Asp Pro
1      5      10      15
Val Asn Gly Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn Ala Gly Gln
      20      25      30
Met Gln Pro Val Lys Ala Phe Lys Ile His Asn Lys Ile Trp Val Ile
      35      40      45
Pro Glu Arg Asp Thr Phe Thr Asn Pro Glu Glu Gly Asp Leu Asn Pro
      50      55      60
Pro Pro Glu Ala Lys Gln Val Pro Val Ser Tyr Tyr Asp Ser Thr Tyr
      65      70      75      80
Leu Ser Thr Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly Val Thr Lys
      85      90      95
Leu Phe Glu Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met Leu Leu Thr
      100      105      110
Ser Ile Val Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr Ile Asp Thr
      115      120      125
Glu Leu Lys Val Ile Asp Thr Asn Cys Ile Asn Val Ile Gln Pro Asp
      130      135      140
Gly Ser Tyr Arg Ser Glu Glu Leu Asn Leu Val Ile Ile Gly Pro Ser
      145      150      155      160
Ala Asp Ile Ile Gln Phe Glu Cys Lys Ser Phe Gly His Glu Val Leu
      165      170      175
Asn Leu Thr Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile Arg Phe Ser
      180      185      190
Pro Asp Phe Thr Phe Gly Phe Glu Glu Ser Leu Glu Val Asp Thr Asn
      195      200      205
Pro Leu Leu Gly Ala Gly Lys Phe Ala Thr Asp Pro Ala Val Thr Leu
      210      215      220
Ala His Glu Leu Ile His Ala Gly His Arg Leu Tyr Gly Ile Ala Ile
      225      230      235      240

```

-continued

---

Asn Pro Asn Arg Val Phe Lys Val Asn Thr Asn Ala Tyr Tyr Glu Met  
                   245                                  250                                  255

Ser Gly Leu Glu Val Ser Phe Glu Glu Leu Arg Thr Phe Gly Gly His  
                   260                                  265                                  270

Asp Ala Lys Phe Ile Asp Ser Leu Gln Glu Asn Glu Phe Arg Leu Tyr  
                   275                                  280                                  285

Tyr Tyr Asn Lys Phe Lys Asp Ile Ala Ser Thr Leu Asn Lys Ala Lys  
                   290                                  295                                  300

Ser Ile Val Gly Thr Thr Ala Ser Leu Gln Tyr Met Lys Asn Val Phe  
 305                                  310                                  315                                  320

Lys Glu Lys Tyr Leu Leu Ser Glu Asp Thr Ser Gly Lys Phe Ser Val  
                   325                                  330                                  335

Asp Lys Leu Lys Phe Asp Lys Leu Tyr Lys Met Leu Thr Glu Ile Tyr  
                   340                                  345                                  350

Thr Glu Asp Asn Phe Val Lys Phe Phe Lys Val Leu Asn Arg Lys Thr  
                   355                                  360                                  365

Tyr Leu Asn Phe Asp Lys Ala Val Phe Lys Ile Asn Ile Val Pro Lys  
                   370                                  375                                  380

Val Asn Tyr Thr Ile Tyr Asp Gly Phe Asn Leu Arg Asn Thr Asn Leu  
 385                                  390                                  395                                  400

Ala Ala Asn Phe Asn Gly Gln Asn Thr Glu Ile Asn Asn Met Asn Phe  
                   405                                  410                                  415

Thr Lys Leu Lys Asn Phe Thr Gly Leu Phe Glu Phe Tyr Lys Leu Leu  
                   420                                  425                                  430

Cys Val Asp Gly Ile Ile Thr Ser Lys Thr Lys Ser Asp Asp Asp Asp  
                   435                                  440                                  445

Lys Tyr Gly Gly Phe Leu Arg Arg Ile Arg Pro Lys Leu Lys Trp Asp  
                   450                                  455                                  460

Asn Gln Ala Leu Ala Gly Gly Gly Ser Gly Gly Gly Ser Gly  
 465                                  470                                  475                                  480

Gly Gly Gly Ser Ala Leu Val Leu Gln Cys Ile Lys Val Asn Asn Trp  
                   485                                  490                                  495

Asp Leu Phe Phe Ser Pro Ser Glu Asp Asn Phe Thr Asn Asp Leu Asn  
                   500                                  505                                  510

Lys Gly Glu Glu Ile Thr Ser Asp Thr Asn Ile Glu Ala Ala Glu Glu  
                   515                                  520                                  525

Asn Ile Ser Leu Asp Leu Ile Gln Gln Tyr Tyr Leu Thr Phe Asn Phe  
                   530                                  535                                  540

Asp Asn Glu Pro Glu Asn Ile Ser Ile Glu Asn Leu Ser Ser Asp Ile  
 545                                  550                                  555                                  560

Ile Gly Gln Leu Glu Leu Met Pro Asn Ile Glu Arg Phe Pro Asn Gly  
                   565                                  570                                  575

Lys Lys Tyr Glu Leu Asp Lys Tyr Thr Met Phe His Tyr Leu Arg Ala  
                   580                                  585                                  590

Gln Glu Phe Glu His Gly Lys Ser Arg Ile Ala Leu Thr Asn Ser Val  
                   595                                  600                                  605

Asn Glu Ala Leu Leu Asn Pro Ser Arg Val Tyr Thr Phe Phe Ser Ser  
                   610                                  615                                  620

Asp Tyr Val Lys Lys Val Asn Lys Ala Thr Glu Ala Ala Met Phe Leu  
 625                                  630                                  635                                  640

-continued

---

Gly	Trp	Val	Glu	Gln	Leu	Val	Tyr	Asp	Phe	Thr	Asp	Glu	Thr	Ser	Glu		
				645					650					655			
Val	Ser	Thr	Thr	Asp	Lys	Ile	Ala	Asp	Ile	Thr	Ile	Ile	Ile	Pro	Tyr		
			660					665					670				
Ile	Gly	Pro	Ala	Leu	Asn	Ile	Gly	Asn	Met	Leu	Tyr	Lys	Asp	Asp	Phe		
		675					680					685					
Val	Gly	Ala	Leu	Ile	Phe	Ser	Gly	Ala	Val	Ile	Leu	Leu	Glu	Phe	Ile		
	690					695					700						
Pro	Glu	Ile	Ala	Ile	Pro	Val	Leu	Gly	Thr	Phe	Ala	Leu	Val	Ser	Tyr		
705					710					715					720		
Ile	Ala	Asn	Lys	Val	Leu	Thr	Val	Gln	Thr	Ile	Asp	Asn	Ala	Leu	Ser		
			725						730					735			
Lys	Arg	Asn	Glu	Lys	Trp	Asp	Glu	Val	Tyr	Lys	Tyr	Ile	Val	Thr	Asn		
			740					745					750				
Trp	Leu	Ala	Lys	Val	Asn	Thr	Gln	Ile	Asp	Leu	Ile	Arg	Lys	Lys	Met		
	755						760					765					
Lys	Glu	Ala	Leu	Glu	Asn	Gln	Ala	Glu	Ala	Thr	Lys	Ala	Ile	Ile	Asn		
	770					775						780					
Tyr	Gln	Tyr	Asn	Gln	Tyr	Thr	Glu	Glu	Glu	Lys	Asn	Asn	Ile	Asn	Phe		
785					790					795					800		
Asn	Ile	Asp	Asp	Leu	Ser	Ser	Lys	Leu	Asn	Glu	Ser	Ile	Asn	Lys	Ala		
				805					810					815			
Met	Ile	Asn	Ile	Asn	Lys	Phe	Leu	Asn	Gln	Cys	Ser	Val	Ser	Tyr	Leu		
		820						825					830				
Met	Asn	Ser	Met	Ile	Pro	Tyr	Gly	Val	Lys	Arg	Leu	Glu	Asp	Phe	Asp		
		835					840					845					
Ala	Ser	Leu	Lys	Asp	Ala	Leu	Leu	Lys	Tyr	Ile	Tyr	Asp	Asn	Arg	Gly		
	850				855							860					
Thr	Leu	Ile	Gly	Gln	Val	Asp	Arg	Leu	Lys	Asp	Lys	Val	Asn	Asn	Thr		
865					870					875					880		
Leu	Ser	Thr	Asp	Ile	Pro	Phe	Gln	Leu	Ser	Lys	Tyr	Val	Asp	Asn	Gln		
				885					890					895			
Arg	Leu	Leu	Ser	Thr	Leu	Glu	Ala	Leu	Ala	Ser	Gly						
		900						905									

<210> SEQ ID NO 3  
 <211> LENGTH: 906  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Retargeted endopeptidase

<400> SEQUENCE: 3

Ile	Ser	Glu	Phe	Gly	Ser	Met	Glu	Phe	Val	Asn	Lys	Gln	Phe	Asn	Tyr		
1				5					10					15			
Lys	Asp	Pro	Val	Asn	Gly	Val	Asp	Ile	Ala	Tyr	Ile	Lys	Ile	Pro	Asn		
			20					25					30				
Ala	Gly	Gln	Met	Gln	Pro	Val	Lys	Ala	Phe	Lys	Ile	His	Asn	Lys	Ile		
		35					40					45					
Trp	Val	Ile	Pro	Glu	Arg	Asp	Thr	Phe	Thr	Asn	Pro	Glu	Glu	Gly	Asp		
	50					55					60						
Leu	Asn	Pro	Pro	Pro	Glu	Ala	Lys	Gln	Val	Pro	Val	Ser	Tyr	Tyr	Asp		
65					70					75					80		

-continued

---

Ser Thr Tyr Leu Ser Thr Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly  
 85 90 95  
 Val Thr Lys Leu Phe Glu Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met  
 100 105 110  
 Leu Leu Thr Ser Ile Val Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr  
 115 120 125  
 Ile Asp Thr Glu Leu Lys Val Ile Asp Thr Asn Cys Ile Asn Val Ile  
 130 135 140  
 Gln Pro Asp Gly Ser Tyr Arg Ser Glu Glu Leu Asn Leu Val Ile Ile  
 145 150 155 160  
 Gly Pro Ser Ala Asp Ile Ile Gln Phe Glu Cys Lys Ser Phe Gly His  
 165 170 175  
 Glu Val Leu Asn Leu Thr Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile  
 180 185 190  
 Arg Phe Ser Pro Asp Phe Thr Phe Gly Phe Glu Glu Ser Leu Glu Val  
 195 200 205  
 Asp Thr Asn Pro Leu Leu Gly Ala Gly Lys Phe Ala Thr Asp Pro Ala  
 210 215 220  
 Val Thr Leu Ala His Glu Leu Ile His Ala Gly His Arg Leu Tyr Gly  
 225 230 235 240  
 Ile Ala Ile Asn Pro Asn Arg Val Phe Lys Val Asn Thr Asn Ala Tyr  
 245 250 255  
 Tyr Glu Met Ser Gly Leu Glu Val Ser Phe Glu Glu Leu Arg Thr Phe  
 260 265 270  
 Gly Gly His Asp Ala Lys Phe Ile Asp Ser Leu Gln Glu Asn Glu Phe  
 275 280 285  
 Arg Leu Tyr Tyr Tyr Asn Lys Phe Lys Asp Ile Ala Ser Thr Leu Asn  
 290 295 300  
 Lys Ala Lys Ser Ile Val Gly Thr Thr Ala Ser Leu Gln Tyr Met Lys  
 305 310 315 320  
 Asn Val Phe Lys Glu Lys Tyr Leu Leu Ser Glu Asp Thr Ser Gly Lys  
 325 330 335  
 Phe Ser Val Asp Lys Leu Lys Phe Asp Lys Leu Tyr Lys Met Leu Thr  
 340 345 350  
 Glu Ile Tyr Thr Glu Asp Asn Phe Val Lys Phe Phe Lys Val Leu Asn  
 355 360 365  
 Arg Lys Thr Tyr Leu Asn Phe Asp Lys Ala Val Phe Lys Ile Asn Ile  
 370 375 380  
 Val Pro Lys Val Asn Tyr Thr Ile Tyr Asp Gly Phe Asn Leu Arg Asn  
 385 390 395 400  
 Thr Asn Leu Ala Ala Asn Phe Asn Gly Gln Asn Thr Glu Ile Asn Asn  
 405 410 415  
 Met Asn Phe Thr Lys Leu Lys Asn Phe Thr Gly Leu Phe Glu Phe Tyr  
 420 425 430  
 Lys Leu Leu Cys Val Asp Gly Ile Ile Thr Ser Lys Thr Lys Ser Leu  
 435 440 445  
 Ile Glu Gly Arg Asn Lys Ala Leu Asn Asp Leu Cys Ile Lys Val Asn  
 450 455 460  
 Asn Trp Asp Leu Phe Phe Ser Pro Ser Glu Asp Asn Phe Thr Asn Asp  
 465 470 475 480  
 Leu Asn Lys Gly Glu Glu Ile Thr Ser Asp Thr Asn Ile Glu Ala Ala

-continued

485				490				495							
Glu	Glu	Asn	Ile	Ser	Leu	Asp	Leu	Ile	Gln	Gln	Tyr	Tyr	Leu	Thr	Phe
		500						505					510		
Asn	Phe	Asp	Asn	Glu	Pro	Glu	Asn	Ile	Ser	Ile	Glu	Asn	Leu	Ser	Ser
		515						520				525			
Asp	Ile	Ile	Gly	Gln	Leu	Glu	Leu	Met	Pro	Asn	Ile	Glu	Arg	Phe	Pro
	530					535						540			
Asn	Gly	Lys	Lys	Tyr	Glu	Leu	Asp	Lys	Tyr	Thr	Met	Phe	His	Tyr	Leu
	545				550					555					560
Arg	Ala	Gln	Glu	Phe	Glu	His	Gly	Lys	Ser	Arg	Ile	Ala	Leu	Thr	Asn
			565						570					575	
Ser	Val	Asn	Glu	Ala	Leu	Leu	Asn	Pro	Ser	Arg	Val	Tyr	Thr	Phe	Phe
			580					585					590		
Ser	Ser	Asp	Tyr	Val	Lys	Lys	Val	Asn	Lys	Ala	Thr	Glu	Ala	Ala	Met
		595						600				605			
Phe	Leu	Gly	Trp	Val	Glu	Gln	Leu	Val	Tyr	Asp	Phe	Thr	Asp	Glu	Thr
	610					615						620			
Ser	Glu	Val	Ser	Thr	Thr	Asp	Lys	Ile	Ala	Asp	Ile	Thr	Ile	Ile	Ile
	625				630					635					640
Pro	Tyr	Ile	Gly	Pro	Ala	Leu	Asn	Ile	Gly	Asn	Met	Leu	Tyr	Lys	Asp
			645						650					655	
Asp	Phe	Val	Gly	Ala	Leu	Ile	Phe	Ser	Gly	Ala	Val	Ile	Leu	Leu	Glu
		660						665				670			
Phe	Ile	Pro	Glu	Ile	Ala	Ile	Pro	Val	Leu	Gly	Thr	Phe	Ala	Leu	Val
	675						680					685			
Ser	Tyr	Ile	Ala	Asn	Lys	Val	Leu	Thr	Val	Gln	Thr	Ile	Asp	Asn	Ala
	690					695					700				
Leu	Ser	Lys	Arg	Asn	Glu	Lys	Trp	Asp	Glu	Val	Tyr	Lys	Tyr	Ile	Val
	705				710					715					720
Thr	Asn	Trp	Leu	Ala	Lys	Val	Asn	Thr	Gln	Ile	Asp	Leu	Ile	Arg	Lys
			725						730					735	
Lys	Met	Lys	Glu	Ala	Leu	Glu	Asn	Gln	Ala	Glu	Ala	Thr	Lys	Ala	Ile
			740					745					750		
Ile	Asn	Tyr	Gln	Tyr	Asn	Gln	Tyr	Thr	Glu	Glu	Glu	Lys	Asn	Asn	Ile
		755					760					765			
Asn	Phe	Asn	Ile	Asp	Asp	Leu	Ser	Ser	Lys	Leu	Asn	Glu	Ser	Ile	Asn
	770					775					780				
Lys	Ala	Met	Ile	Asn	Ile	Asn	Lys	Phe	Leu	Asn	Gln	Cys	Ser	Val	Ser
	785				790					795					800
Tyr	Leu	Met	Asn	Ser	Met	Ile	Pro	Tyr	Gly	Val	Lys	Arg	Leu	Glu	Asp
			805						810					815	
Phe	Asp	Ala	Ser	Leu	Lys	Asp	Ala	Leu	Leu	Lys	Tyr	Ile	Tyr	Asp	Asn
		820						825					830		
Arg	Gly	Thr	Leu	Ile	Gly	Gln	Val	Asp	Arg	Leu	Lys	Asp	Lys	Val	Asn
		835					840					845			
Asn	Thr	Leu	Ser	Thr	Asp	Ile	Pro	Phe	Gln	Leu	Ser	Lys	Tyr	Val	Asp
	850					855					860				
Asn	Gln	Arg	Leu	Ser	Thr	Leu	Glu	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly
	865				870				875						880
Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Ala	Leu	Val	Val	Gly	Arg	Pro	Glu
			885						890					895	

-continued

---

Trp Trp Met Asp Tyr Gln Lys Arg Tyr Gly  
 900 905

<210> SEQ ID NO 4  
 <211> LENGTH: 919  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Retargeted endopeptidase

<400> SEQUENCE: 4

Met Glu Phe Val Asn Lys Gln Phe Asn Tyr Lys Asp Pro Val Asn Gly  
 1 5 10 15

Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn Ala Gly Gln Met Gln Pro  
 20 25 30

Val Lys Ala Phe Lys Ile His Asn Lys Ile Trp Val Ile Pro Glu Arg  
 35 40 45

Asp Thr Phe Thr Asn Pro Glu Glu Gly Asp Leu Asn Pro Pro Pro Glu  
 50 55 60

Ala Lys Gln Val Pro Val Ser Tyr Tyr Asp Ser Thr Tyr Leu Ser Thr  
 65 70 75 80

Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly Val Thr Lys Leu Phe Glu  
 85 90 95

Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met Leu Leu Thr Ser Ile Val  
 100 105 110

Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr Ile Asp Thr Glu Leu Lys  
 115 120 125

Val Ile Asp Thr Asn Cys Ile Asn Val Ile Gln Pro Asp Gly Ser Tyr  
 130 135 140

Arg Ser Glu Glu Leu Asn Leu Val Ile Ile Gly Pro Ser Ala Asp Ile  
 145 150 155 160

Ile Gln Phe Glu Cys Lys Ser Phe Gly His Glu Val Leu Asn Leu Thr  
 165 170 175

Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile Arg Phe Ser Pro Asp Phe  
 180 185 190

Thr Phe Gly Phe Glu Glu Ser Leu Glu Val Asp Thr Asn Pro Leu Leu  
 195 200 205

Gly Ala Gly Lys Phe Ala Thr Asp Pro Ala Val Thr Leu Ala His Glu  
 210 215 220

Leu Ile His Ala Gly His Arg Leu Tyr Gly Ile Ala Ile Asn Pro Asn  
 225 230 235 240

Arg Val Phe Lys Val Asn Thr Asn Ala Tyr Tyr Glu Met Ser Gly Leu  
 245 250 255

Glu Val Ser Phe Glu Glu Leu Arg Thr Phe Gly Gly His Asp Ala Lys  
 260 265 270

Phe Ile Asp Ser Leu Gln Glu Asn Glu Phe Arg Leu Tyr Tyr Tyr Asn  
 275 280 285

Lys Phe Lys Asp Ile Ala Ser Thr Leu Asn Lys Ala Lys Ser Ile Val  
 290 295 300

Gly Thr Thr Ala Ser Leu Gln Tyr Met Lys Asn Val Phe Lys Glu Lys  
 305 310 315 320

Tyr Leu Leu Ser Glu Asp Thr Ser Gly Lys Phe Ser Val Asp Lys Leu  
 325 330 335

-continued

---

Lys Phe Asp Lys Leu Tyr Lys Met Leu Thr Glu Ile Tyr Thr Glu Asp  
 340 345 350

Asn Phe Val Lys Phe Phe Lys Val Leu Asn Arg Lys Thr Tyr Leu Asn  
 355 360 365

Phe Asp Lys Ala Val Phe Lys Ile Asn Ile Val Pro Lys Val Asn Tyr  
 370 375 380

Thr Ile Tyr Asp Gly Phe Asn Leu Arg Asn Thr Asn Leu Ala Ala Asn  
 385 390 395 400

Phe Asn Gly Gln Asn Thr Glu Ile Asn Asn Met Asn Phe Thr Lys Leu  
 405 410 415

Lys Asn Phe Thr Gly Leu Phe Glu Phe Tyr Lys Leu Leu Cys Val Asp  
 420 425 430

Gly Ile Ile Thr Ser Lys Thr Lys Ser Asp Asp Asp Asp Lys Tyr Gly  
 435 440 445

Gly Phe Met Thr Ser Glu Lys Ser Gln Thr Pro Leu Val Thr Leu Phe  
 450 455 460

Lys Asn Ala Ile Ile Lys Asn Ala Tyr Lys Lys Gly Glu Ala Leu Ala  
 465 470 475 480

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ala  
 485 490 495

Leu Val Leu Gln Cys Ile Lys Val Asn Asn Trp Asp Leu Phe Phe Ser  
 500 505 510

Pro Ser Glu Asp Asn Phe Thr Asn Asp Leu Asn Lys Gly Glu Glu Ile  
 515 520 525

Thr Ser Asp Thr Asn Ile Glu Ala Ala Glu Glu Asn Ile Ser Leu Asp  
 530 535 540

Leu Ile Gln Gln Tyr Tyr Leu Thr Phe Asn Phe Asp Asn Glu Pro Glu  
 545 550 555 560

Asn Ile Ser Ile Glu Asn Leu Ser Ser Asp Ile Ile Gly Gln Leu Glu  
 565 570 575

Leu Met Pro Asn Ile Glu Arg Phe Pro Asn Gly Lys Lys Tyr Glu Leu  
 580 585 590

Asp Lys Tyr Thr Met Phe His Tyr Leu Arg Ala Gln Glu Phe Glu His  
 595 600 605

Gly Lys Ser Arg Ile Ala Leu Thr Asn Ser Val Asn Glu Ala Leu Leu  
 610 615 620

Asn Pro Ser Arg Val Tyr Thr Phe Phe Ser Ser Asp Tyr Val Lys Lys  
 625 630 635 640

Val Asn Lys Ala Thr Glu Ala Ala Met Phe Leu Gly Trp Val Glu Gln  
 645 650 655

Leu Val Tyr Asp Phe Thr Asp Glu Thr Ser Glu Val Ser Thr Thr Asp  
 660 665 670

Lys Ile Ala Asp Ile Thr Ile Ile Ile Pro Tyr Ile Gly Pro Ala Leu  
 675 680 685

Asn Ile Gly Asn Met Leu Tyr Lys Asp Asp Phe Val Gly Ala Leu Ile  
 690 695 700

Phe Ser Gly Ala Val Ile Leu Leu Glu Phe Ile Pro Glu Ile Ala Ile  
 705 710 715 720

Pro Val Leu Gly Thr Phe Ala Leu Val Ser Tyr Ile Ala Asn Lys Val  
 725 730 735

-continued

---

```

Leu Thr Val Gln Thr Ile Asp Asn Ala Leu Ser Lys Arg Asn Glu Lys
    740                                745                                750

Trp Asp Glu Val Tyr Lys Tyr Ile Val Thr Asn Trp Leu Ala Lys Val
    755                                760                                765

Asn Thr Gln Ile Asp Leu Ile Arg Lys Lys Met Lys Glu Ala Leu Glu
    770                                775                                780

Asn Gln Ala Glu Ala Thr Lys Ala Ile Ile Asn Tyr Gln Tyr Asn Gln
    785                                790                                795                                800

Tyr Thr Glu Glu Glu Lys Asn Asn Ile Asn Phe Asn Ile Asp Asp Leu
    805                                810                                815

Ser Ser Lys Leu Asn Glu Ser Ile Asn Lys Ala Met Ile Asn Ile Asn
    820                                825                                830

Lys Phe Leu Asn Gln Cys Ser Val Ser Tyr Leu Met Asn Ser Met Ile
    835                                840                                845

Pro Tyr Gly Val Lys Arg Leu Glu Asp Phe Asp Ala Ser Leu Lys Asp
    850                                855                                860

Ala Leu Leu Lys Tyr Ile Tyr Asp Asn Arg Gly Thr Leu Ile Gly Gln
    865                                870                                875                                880

Val Asp Arg Leu Lys Asp Lys Val Asn Asn Thr Leu Ser Thr Asp Ile
    885                                890                                895

Pro Phe Gln Leu Ser Lys Tyr Val Asp Asn Gln Arg Leu Leu Ser Thr
    900                                905                                910

Leu Glu Ala Leu Ala Ser Gly
    915

<210> SEQ ID NO 5
<211> LENGTH: 206
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg
1      5      10      15

Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
20     25     30

Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
35     40     45

Met Leu Asp Glu Gln Gly Glu Gln Leu Asp Arg Val Glu Glu Gly Met
50     55     60

Asn His Ile Asn Gln Asp Met Lys Glu Ala Glu Lys Asn Leu Lys Asp
65     70     75     80

Leu Gly Lys Cys Cys Gly Leu Phe Ile Cys Pro Cys Asn Lys Leu Lys
85     90     95

Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
100    105    110

Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
115    120    125

Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn
130    135    140

Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile Gly Asn Leu
145    150    155    160

Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg
165    170    175
    
```

-continued

Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Thr Arg Ile  
 180 185 190

Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly  
 195 200 205

<210> SEQ ID NO 6  
 <211> LENGTH: 206  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg  
 1 5 10 15

Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met  
 20 25 30

Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val  
 35 40 45

Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met  
 50 55 60

Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp  
 65 70 75 80

Leu Gly Lys Phe Cys Gly Leu Cys Val Cys Pro Cys Asn Lys Leu Lys  
 85 90 95

Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val  
 100 105 110

Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala  
 115 120 125

Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn  
 130 135 140

Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile Gly Asn Leu  
 145 150 155 160

Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg  
 165 170 175

Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Thr Arg Ile  
 180 185 190

Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly  
 195 200 205

<210> SEQ ID NO 7  
 <211> LENGTH: 206  
 <212> TYPE: PRT  
 <213> ORGANISM: Macaca mulatta

<400> SEQUENCE: 7

Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg  
 1 5 10 15

Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met  
 20 25 30

Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val  
 35 40 45

Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met  
 50 55 60

Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp  
 65 70 75 80



-continued

&lt;213&gt; ORGANISM: Rattus norvegicus

&lt;400&gt; SEQUENCE: 9

```

Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg
1           5           10           15
Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
           20           25           30
Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
           35           40           45
Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met
           50           55           60
Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp
           65           70           75           80
Leu Gly Lys Phe Cys Gly Leu Cys Val Cys Pro Cys Asn Lys Leu Lys
           85           90           95
Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
           100          105          110
Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
           115          120          125
Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn
           130          135          140
Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile Gly Asn Leu
           145          150          155          160
Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg
           165          170          175
Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Thr Arg Ile
           180          185          190
Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly
           195          200          205

```

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 206

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 10

```

Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg
1           5           10           15
Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
           20           25           30
Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
           35           40           45
Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met
           50           55           60
Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp
           65           70           75           80
Leu Gly Lys Phe Cys Gly Leu Cys Val Cys Pro Cys Asn Lys Leu Lys
           85           90           95
Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
           100          105          110
Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
           115          120          125
Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn

```



-continued

35					40					45					
Met	Leu	Asp	Glu	Gln	Gly	Glu	Gln	Leu	Glu	Arg	Ile	Glu	Glu	Gly	Met
50						55					60				
Asp	Gln	Ile	Asn	Lys	Asp	Met	Lys	Asp	Ala	Glu	Lys	Asn	Leu	Asn	Asp
65				70					75					80	
Leu	Gly	Lys	Phe	Cys	Gly	Leu	Cys	Ser	Cys	Pro	Cys	Asn	Lys	Met	Lys
			85						90					95	
Ser	Gly	Gly	Ser	Lys	Ala	Trp	Gly	Asn	Asn	Gln	Asp	Gly	Val	Val	Ala
			100					105					110		
Ser	Gln	Pro	Ala	Arg	Val	Val	Asp	Glu	Arg	Glu	Gln	Met	Ala	Ile	Ser
		115					120					125			
Gly	Gly	Phe	Ile	Arg	Arg	Val	Thr	Asp	Asp	Ala	Arg	Glu	Asn	Glu	Met
	130					135					140				
Asp	Glu	Asn	Leu	Glu	Gln	Val	Gly	Gly	Ile	Ile	Gly	Asn	Leu	Arg	His
145					150					155					160
Met	Ala	Leu	Asp	Met	Gly	Asn	Glu	Ile	Asp	Thr	Gln	Asn	Arg	Gln	Ile
			165						170					175	
Asp	Arg	Ile	Met	Glu	Lys	Ala	Asp	Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu
		180						185					190		
Ala	Asn	Gln	Arg	Ala	Thr	Lys	Met	Leu	Gly	Ser	Gly				
	195						200								

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 203

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Carassius auratus

&lt;400&gt; SEQUENCE: 13

Met	Ala	Asp	Glu	Ala	Asp	Met	Arg	Asn	Glu	Leu	Thr	Asp	Met	Gln	Ala
1				5					10					15	
Arg	Ala	Asp	Gln	Leu	Gly	Asp	Glu	Ser	Leu	Glu	Ser	Thr	Arg	Arg	Met
			20					25					30		
Leu	Gln	Leu	Val	Glu	Glu	Ser	Lys	Asp	Ala	Gly	Ile	Arg	Thr	Leu	Val
		35				40						45			
Met	Leu	Asp	Glu	Gln	Gly	Glu	Gln	Leu	Glu	Arg	Ile	Glu	Glu	Gly	Met
	50				55						60				
Asp	Gln	Ile	Asn	Lys	Asp	Met	Lys	Glu	Ala	Glu	Lys	Asn	Leu	Thr	Asp
65				70					75					80	
Leu	Gly	Asn	Leu	Cys	Gly	Leu	Cys	Pro	Cys	Pro	Cys	Asn	Lys	Leu	Lys
			85						90					95	
Gly	Gly	Gly	Gln	Ser	Trp	Gly	Asn	Asn	Gln	Asp	Gly	Val	Val	Ser	Ser
			100					105					110		
Gln	Pro	Ala	Arg	Val	Val	Asp	Glu	Arg	Glu	Gln	Met	Ala	Ile	Ser	Gly
		115					120					125			
Gly	Phe	Ile	Arg	Arg	Val	Thr	Asn	Asp	Ala	Arg	Glu	Asn	Glu	Met	Asp
	130					135					140				
Glu	Asn	Leu	Glu	Gln	Val	Gly	Ser	Ile	Ile	Gly	Asn	Leu	Arg	His	Met
145				150						155					160
Ala	Leu	Asp	Met	Gly	Asn	Glu	Ile	Asp	Thr	Gln	Asn	Arg	Gln	Ile	Asp
			165						170					175	
Arg	Ile	Met	Asp	Met	Ala	Asp	Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu	Ala
		180							185					190	

-continued

---

```

Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly
      195                200

<210> SEQ ID NO 14
<211> LENGTH: 204
<212> TYPE: PRT
<213> ORGANISM: Danio rerio

<400> SEQUENCE: 14

Met Ala Glu Asp Ser Asp Met Arg Asn Glu Leu Ala Asp Met Gln Gln
1          5          10          15

Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
      20          25          30

Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
      35          40          45

Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met
      50          55          60

Asp Gln Ile Asn Lys Asp Met Lys Asp Ala Glu Lys Asn Leu Asn Asp
      65          70          75          80

Leu Gly Lys Phe Cys Gly Leu Cys Ser Cys Pro Cys Asn Lys Met Lys
      85          90          95

Ser Gly Ala Ser Lys Ala Trp Gly Asn Asn Gln Asp Gly Val Val Ala
      100         105         110

Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala Ile Ser
      115         120         125

Gly Gly Phe Ile Arg Arg Val Thr Asp Asp Ala Arg Glu Asn Glu Met
      130         135         140

Asp Glu Asn Leu Glu Gln Val Gly Gly Ile Ile Gly Asn Leu Arg His
      145         150         155         160

Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg Gln Ile
      165         170         175

Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Thr Arg Ile Asp Glu
      180         185         190

Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly
      195                200

```

```

<210> SEQ ID NO 15
<211> LENGTH: 203
<212> TYPE: PRT
<213> ORGANISM: Danio rerio

<400> SEQUENCE: 15

Met Ala Asp Glu Ser Asp Met Arg Asn Glu Leu Asn Asp Met Gln Ala
1          5          10          15

Arg Ala Asp Gln Leu Gly Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
      20          25          30

Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
      35          40          45

Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met
      50          55          60

Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp
      65          70          75          80

Leu Gly Asn Leu Cys Gly Leu Cys Pro Cys Pro Cys Asn Lys Leu Lys
      85          90          95

```

-continued

---

Gly Gly Gly Gln Ser Trp Gly Asn Asn Gln Asp Gly Val Val Ser Ser  
100 105 110

Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala Ile Ser Gly  
115 120 125

Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn Glu Met Asp  
130 135 140

Glu Asn Leu Glu Gln Val Gly Ser Ile Ile Gly Asn Leu Arg His Met  
145 150 155 160

Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg Gln Ile Asp  
165 170 175

Arg Ile Met Asp Met Ala Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala  
180 185 190

Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly  
195 200

<210> SEQ ID NO 16  
<211> LENGTH: 210  
<212> TYPE: PRT  
<213> ORGANISM: *Torpedo marmorata*

<400> SEQUENCE: 16

Met Glu Asn Ser Val Glu Asn Ser Met Asp Pro Arg Ser Glu Gln Glu  
1 5 10 15

Glu Met Gln Arg Cys Ala Asp Gln Ile Thr Asp Glu Ser Leu Glu Ser  
20 25 30

Thr Arg Arg Met Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile  
35 40 45

Arg Thr Leu Val Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile  
50 55 60

Glu Glu Gly Met Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys  
65 70 75 80

Asn Leu Ser Asp Leu Gly Lys Cys Cys Gly Leu Cys Ser Cys Pro Cys  
85 90 95

Asn Lys Leu Lys Asn Phe Glu Ala Gly Gly Ala Tyr Lys Lys Val Trp  
100 105 110

Gly Asn Asn Gln Asp Gly Val Val Ala Ser Gln Pro Ala Arg Val Met  
115 120 125

Asp Asp Arg Glu Gln Met Ala Met Ser Gly Gly Tyr Ile Arg Arg Ile  
130 135 140

Thr Asp Asp Ala Arg Glu Asn Glu Met Glu Glu Asn Leu Asp Gln Val  
145 150 155 160

Gly Ser Ile Ile Gly Asn Leu Arg His Met Ala Leu Asp Met Ser Asn  
165 170 175

Glu Ile Gly Ser Gln Asn Ala Gln Ile Asp Arg Ile Val Val Lys Gly  
180 185 190

Asp Met Asn Lys Ala Arg Ile Asp Glu Ala Asn Lys His Ala Thr Lys  
195 200 205

Met Leu  
210

<210> SEQ ID NO 17  
<211> LENGTH: 206  
<212> TYPE: PRT  
<213> ORGANISM: *Xenopus laevis*

-continued

&lt;400&gt; SEQUENCE: 17

```

Met Ala Asp Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg
1           5           10           15
Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
20           25           30
Leu Gln Tyr Val Glu Gly Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
35           40           45
Met Leu Asp Glu Gln Gly Glu Gln Leu Asp Arg Val Glu Glu Gly Met
50           55           60
Asn His Ile Asn Gln Asp Met Lys Glu Ala Glu Lys Asn Leu Lys Asp
65           70           75           80
Leu Gly Lys Cys Cys Gly Leu Phe Ile Cys Pro Cys Asn Lys Leu Lys
85           90           95
Ser Ser Gly Ala Tyr Asn Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
100          105          110
Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
115          120          125
Ile Ser Gly Gly Phe Val Arg Arg Val Thr Asn Asp Ala Arg Glu Thr
130          135          140
Glu Met Asp Glu Asn Leu Glu Gln Val Gly Gly Ile Ile Gly Asn Leu
145          150          155          160
Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg
165          170          175
Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Ala Arg Ile
180          185          190
Asp Glu Ala Asn Lys His Ala Thr Lys Met Leu Gly Ser Gly
195          200          205

```

&lt;210&gt; SEQ ID NO 18

&lt;211&gt; LENGTH: 206

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Xenopus laevis*

&lt;400&gt; SEQUENCE: 18

```

Met Ala Asp Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg
1           5           10           15
Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
20           25           30
Leu Gln Tyr Val Glu Gly Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
35           40           45
Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met
50           55           60
Glu Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp
65           70           75           80
Leu Gly Lys Phe Cys Gly Leu Cys Val Cys Pro Cys Asn Lys Leu Lys
85           90           95
Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
100          105          110
Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
115          120          125
Ile Ser Gly Gly Phe Val Arg Arg Val Thr Asn Asp Ala Arg Glu Thr
130          135          140

```

-continued

Glu Met Asp Glu Asn Leu Glu Gln Val Gly Gly Ile Ile Gly Asn Leu  
145 150 155 160

Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg  
165 170 175

Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Ala Arg Ile  
180 185 190

Asp Glu Ala Asn Lys His Ala Thr Lys Met Leu Gly Ser Gly  
195 200 205

<210> SEQ ID NO 19

<211> LENGTH: 212

<212> TYPE: PRT

<213> ORGANISM: Strongylocentrotus purpuratus

<400> SEQUENCE: 19

Met Glu Asp Gln Asn Asp Met Asn Met Arg Ser Glu Leu Glu Glu Ile  
1 5 10 15

Gln Met Gln Ser Asn Met Gln Thr Asp Glu Ser Leu Glu Ser Thr Arg  
20 25 30

Arg Met Leu Gln Met Ala Glu Glu Ser Gln Asp Met Gly Ile Lys Thr  
35 40 45

Leu Val Met Leu Asp Glu Gln Gly Glu Gln Leu Asp Arg Ile Glu Glu  
50 55 60

Gly Met Asp Gln Ile Asn Thr Asp Met Arg Glu Ala Glu Lys Asn Leu  
65 70 75 80

Thr Gly Leu Glu Lys Cys Cys Gly Ile Cys Val Cys Pro Trp Lys Lys  
85 90 95

Leu Gly Asn Phe Glu Lys Gly Asp Asp Tyr Lys Lys Thr Trp Lys Gly  
100 105 110

Asn Asp Asp Gly Lys Val Asn Ser His Gln Pro Met Arg Met Glu Asp  
115 120 125

Asp Arg Asp Gly Cys Gly Gly Asn Ala Ser Met Ile Thr Arg Ile Thr  
130 135 140

Asn Asp Ala Arg Glu Asp Glu Met Asp Glu Asn Leu Thr Gln Val Ser  
145 150 155 160

Ser Ile Val Gly Asn Leu Arg His Met Ala Ile Asp Met Gln Ser Glu  
165 170 175

Ile Gly Ala Gln Asn Ser Gln Val Gly Arg Ile Thr Ser Lys Ala Glu  
180 185 190

Ser Asn Glu Gly Arg Ile Asn Ser Ala Asp Lys Arg Ala Lys Asn Ile  
195 200 205

Leu Arg Asn Lys  
210

<210> SEQ ID NO 20

<211> LENGTH: 212

<212> TYPE: PRT

<213> ORGANISM: Drosophila melanogaster

<400> SEQUENCE: 20

Met Pro Ala Asp Pro Ser Glu Glu Val Ala Pro Gln Val Pro Lys Thr  
1 5 10 15

Glu Leu Glu Glu Leu Gln Ile Asn Ala Gln Gly Val Ala Asp Glu Ser  
20 25 30

-continued

---

Leu Glu Ser Thr Arg Arg Met Leu Ala Leu Cys Glu Glu Ser Lys Glu  
           35                                  40                                  45  
 Ala Gly Ile Arg Thr Leu Val Ala Leu Asp Asp Gln Gly Glu Gln Leu  
           50                                  55                                  60  
 Asp Arg Ile Glu Glu Gly Met Asp Gln Ile Asn Ala Asp Met Arg Glu  
           65                                  70                                  75                                  80  
 Ala Glu Lys Asn Leu Ser Gly Met Glu Lys Cys Cys Gly Ile Cys Val  
                                   85                                  90                                  95  
 Leu Pro Cys Asn Lys Ser Gln Ser Phe Lys Glu Asp Asp Gly Thr Trp  
                                   100                                  105                                  110  
 Lys Gly Asn Asp Asp Gly Lys Val Val Asn Asn Gln Pro Gln Arg Val  
                                   115                                  120                                  125  
 Met Asp Asp Arg Asn Gly Met Met Ala Gln Ala Gly Tyr Ile Gly Arg  
           130                                  135                                  140  
 Ile Thr Asn Asp Ala Arg Glu Asp Glu Met Glu Glu Asn Met Gly Gln  
           145                                  150                                  155                                  160  
 Val Asn Thr Met Ile Gly Asn Leu Arg Asn Met Ala Leu Asp Met Gly  
                                   165                                  170                                  175  
 Ser Glu Leu Glu Asn Gln Asn Arg Gln Ile Asp Arg Ile Asn Arg Lys  
                                   180                                  185                                  190  
 Gly Glu Ser Asn Glu Ala Arg Ile Ala Val Ala Asn Gln Arg Ala His  
                                   195                                  200                                  205  
 Gln Leu Leu Lys  
           210

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 212

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Hirudo medicinalis

&lt;400&gt; SEQUENCE: 21

Met Ala Lys Asp Ile Lys Pro Lys Pro Ala Asn Gly Arg Asp Ser Pro  
 1                                  5                                  10                                  15  
 Thr Asp Leu Gln Glu Ile Gln Leu Gln Met Asn Ala Ile Thr Asp Asp  
                                   20                                  25                                  30  
 Ser Leu Glu Ser Thr Arg Arg Met Leu Ala Met Cys Glu Glu Ser Lys  
           35                                  40                                  45  
 Asp Ala Gly Ile Arg Thr Leu Val Met Leu Asp Glu Gln Gly Glu Gln  
           50                                  55                                  60  
 Leu Asp Arg Ile Glu Glu Gly Met Asp Gln Ile Asn Gln Asp Met Arg  
           65                                  70                                  75                                  80  
 Asp Ala Glu Lys Asn Leu Glu Gly Met Glu Lys Cys Cys Gly Leu Cys  
                                   85                                  90                                  95  
 Ile Leu Pro Trp Lys Arg Thr Lys Asn Phe Asp Lys Gly Ala Glu Trp  
                                   100                                  105                                  110  
 Asn Lys Gly Asp Glu Gly Lys Val Asn Thr Asp Gly Pro Arg Leu Val  
           115                                  120                                  125  
 Val Gly Asp Gly Asn Met Gly Pro Ser Gly Gly Phe Ile Thr Lys Ile  
           130                                  135                                  140  
 Thr Asn Asp Ala Arg Glu Glu Glu Met Glu Gln Asn Met Gly Glu Val  
           145                                  150                                  155                                  160  
 Ser Asn Met Ile Ser Asn Leu Arg Asn Met Ala Val Asp Met Gly Ser

-continued

---

```

                165                170                175
Glu Ile Asp Ser Gln Asn Arg Gln Val Asp Arg Ile Asn Asn Lys Met
                180                185                190
Thr Ser Asn Gln Leu Arg Ile Ser Asp Ala Asn Lys Arg Ala Ser Lys
                195                200                205
Leu Leu Lys Glu
                210

```

```

<210> SEQ ID NO 22
<211> LENGTH: 212
<212> TYPE: PRT
<213> ORGANISM: Loligo pealei

```

```

<400> SEQUENCE: 22

```

```

Met Ser Ala Asn Gly Glu Val Glu Val Pro Lys Thr Glu Leu Glu Glu
1                5                10                15
Ile Gln Gln Gln Cys Asn Gln Val Thr Asp Asp Ser Leu Glu Ser Thr
                20                25                30
Arg Arg Met Leu Asn Met Cys Glu Glu Ser Lys Glu Ala Gly Ile Arg
                35                40                45
Thr Leu Val Met Leu Asp Glu Gln Gly Glu Gln Leu Asp Arg Ile Glu
                50                55                60
Glu Gly Leu Asp Gln Ile Asn Gln Asp Met Lys Asp Ala Glu Lys Asn
65                70                75                80
Leu Glu Gly Met Glu Lys Cys Cys Gly Leu Cys Val Leu Pro Trp Lys
                85                90                95
Arg Gly Lys Ser Phe Glu Lys Ser Gly Asp Tyr Ala Asn Thr Trp Lys
                100                105                110
Lys Asp Asp Asp Gly Pro Thr Asn Thr Asn Gly Pro Arg Val Thr Val
                115                120                125
Gly Asp Gln Asn Gly Met Gly Pro Ser Ser Gly Tyr Val Thr Arg Ile
                130                135                140
Thr Asn Asp Ala Arg Glu Asp Asp Met Glu Asn Asn Met Lys Glu Val
145                150                155                160
Ser Ser Met Ile Gly Asn Leu Arg Asn Met Ala Ile Asp Met Gly Asn
                165                170                175
Glu Ile Gly Ser Gln Asn Arg Gln Val Asp Arg Ile Gln Gln Lys Ala
                180                185                190
Glu Ser Asn Glu Ser Arg Ile Asp Glu Ala Asn Lys Lys Ala Thr Lys
                195                200                205
Leu Leu Lys Asn
                210

```

```

<210> SEQ ID NO 23
<211> LENGTH: 220
<212> TYPE: PRT
<213> ORGANISM: Lymnaea stagnalis

```

```

<400> SEQUENCE: 23

```

```

Met Thr Thr Asn Gly Glu Ile Leu Pro Val Gly Glu Glu Glu Glu
1                5                10                15
Glu Leu Gly Glu Asp Ala Leu Leu Arg Lys Gln Ile Asp Cys Asn Thr
                20                25                30
Asn Glu Ser Leu Glu Ser Thr Arg Arg Met Leu Ser Leu Cys Glu Glu

```

-continued

---

	35					40						45			
Ser	Lys	Glu	Ala	Gly	Ile	Lys	Thr	Leu	Val	Met	Leu	Asp	Glu	Gln	Gly
	50					55					60				
Glu	Gln	Leu	Asp	Arg	Ile	Glu	Glu	Gly	Met	Gly	Gln	Ile	Asn	Gln	Asp
65					70					75				80	
Met	Arg	Asp	Ala	Glu	Lys	Asn	Leu	Glu	Gly	Leu	Glu	Lys	Cys	Cys	Gly
			85						90					95	
Leu	Cys	Val	Leu	Pro	Trp	Lys	Arg	Ser	Lys	Asn	Phe	Glu	Lys	Gly	Ser
			100					105						110	
Asp	Tyr	Asn	Lys	Thr	Trp	Lys	Ala	Ser	Glu	Asp	Gly	Lys	Ile	Asn	Thr
	115						120					125			
Asn	Gly	Pro	Arg	Leu	Val	Val	Asp	Gln	Gly	Asn	Gly	Ser	Gly	Pro	Thr
	130					135					140				
Gly	Gly	Tyr	Ile	Thr	Arg	Ile	Thr	Asn	Asp	Ala	Arg	Glu	Asp	Glu	Met
145					150					155					160
Glu	Gln	Asn	Ile	Gly	Glu	Val	Ala	Gly	Met	Val	Ser	Asn	Leu	Arg	Asn
			165						170					175	
Met	Ala	Val	Asp	Met	Gly	Asn	Glu	Ile	Glu	Ser	Gln	Asn	Lys	Gln	Leu
			180					185						190	
Asp	Arg	Ile	Asn	Gln	Lys	Gly	Gly	Ser	Leu	Asn	Val	Arg	Val	Asp	Glu
	195					200						205			
Ala	Asn	Lys	Arg	Ala	Asn	Arg	Ile	Leu	Arg	Lys	Gln				
	210					215					220				

<210> SEQ ID NO 24  
 <211> LENGTH: 207  
 <212> TYPE: PRT  
 <213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 24

Met	Ser	Gly	Asp	Asp	Asp	Ile	Pro	Glu	Gly	Leu	Glu	Ala	Ile	Asn	Leu
1			5					10						15	
Lys	Met	Asn	Ala	Thr	Thr	Asp	Asp	Ser	Leu	Glu	Ser	Thr	Arg	Arg	Met
		20					25						30		
Leu	Ala	Leu	Cys	Glu	Glu	Ser	Lys	Glu	Ala	Gly	Ile	Lys	Thr	Leu	Val
		35					40						45		
Met	Leu	Asp	Asp	Gln	Gly	Glu	Gln	Leu	Glu	Arg	Cys	Glu	Gly	Ala	Leu
	50					55					60				
Asp	Thr	Ile	Asn	Gln	Asp	Met	Lys	Glu	Ala	Glu	Asp	His	Leu	Lys	Gly
65					70					75					80
Met	Glu	Lys	Cys	Cys	Gly	Leu	Cys	Val	Leu	Pro	Trp	Asn	Lys	Thr	Asp
			85						90					95	
Asp	Phe	Glu	Lys	Thr	Glu	Phe	Ala	Lys	Ala	Trp	Lys	Lys	Asp	Asp	Asp
			100					105						110	
Gly	Gly	Val	Ile	Ser	Asp	Gln	Pro	Arg	Ile	Thr	Val	Gly	Asp	Ser	Ser
		115						120					125		
Met	Gly	Pro	Gln	Gly	Gly	Tyr	Ile	Thr	Lys	Ile	Thr	Asn	Asp	Ala	Arg
	130					135						140			
Glu	Asp	Glu	Met	Asp	Glu	Asn	Val	Gln	Gln	Val	Ser	Thr	Met	Val	Gly
145					150						155				160
Asn	Leu	Arg	Asn	Met	Ala	Ile	Asp	Met	Ser	Thr	Glu	Val	Ser	Asn	Gln
			165							170					175

-continued

---

```

Asn Arg Gln Leu Asp Arg Ile His Asp Lys Ala Gln Ser Asn Glu Val
      180                      185                      190

Arg Val Glu Ser Ala Asn Lys Arg Ala Lys Asn Leu Ile Thr Lys
      195                      200                      205

<210> SEQ ID NO 25
<211> LENGTH: 370
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Met Glu Pro Leu Phe Pro Ala Pro Phe Trp Glu Val Ile Tyr Gly Ser
1      5      10      15

His Leu Gln Gly Asn Leu Ser Leu Leu Ser Pro Asn His Ser Leu Leu
      20      25      30

Pro Pro His Leu Leu Leu Asn Ala Ser His Gly Ala Phe Leu Pro Leu
      35      40      45

Gly Leu Lys Val Thr Ile Val Gly Leu Tyr Leu Ala Val Cys Val Gly
      50      55      60

Gly Leu Leu Gly Asn Cys Leu Val Met Tyr Val Ile Leu Arg His Thr
      65      70      75      80

Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala
      85      90      95

Asp Thr Leu Val Leu Leu Thr Leu Pro Phe Gln Gly Thr Asp Ile Leu
      100     105     110

Leu Gly Phe Trp Pro Phe Gly Asn Ala Leu Cys Lys Thr Val Ile Ala
      115     120     125

Ile Asp Tyr Tyr Asn Met Phe Thr Ser Thr Phe Thr Leu Thr Ala Met
      130     135     140

Ser Val Asp Arg Tyr Val Ala Ile Cys His Pro Ile Arg Ala Leu Asp
      145     150     155     160

Val Arg Thr Ser Ser Lys Ala Gln Ala Val Asn Val Ala Ile Trp Ala
      165     170     175

Leu Ala Ser Val Val Gly Val Pro Val Ala Ile Met Gly Ser Ala Gln
      180     185     190

Val Glu Asp Glu Glu Ile Glu Cys Leu Val Glu Ile Pro Thr Pro Gln
      195     200     205

Asp Tyr Trp Gly Pro Val Phe Ala Ile Cys Ile Phe Leu Phe Ser Phe
      210     215     220

Ile Val Pro Val Leu Val Ile Ser Val Cys Tyr Ser Leu Met Ile Arg
      225     230     235     240

Arg Leu Arg Gly Val Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg
      245     250     255

Asn Leu Arg Arg Ile Thr Arg Leu Val Leu Val Val Val Ala Val Phe
      260     265     270

Val Gly Cys Trp Thr Pro Val Gln Val Phe Val Leu Ala Gln Gly Leu
      275     280     285

Gly Val Gln Pro Ser Ser Glu Thr Ala Val Ala Ile Leu Arg Phe Cys
      290     295     300

Thr Ala Leu Gly Tyr Val Asn Ser Cys Leu Asn Pro Ile Leu Tyr Ala
      305     310     315     320

Phe Leu Asp Glu Asn Phe Lys Ala Cys Phe Arg Lys Phe Cys Cys Ala
      325     330     335

```

-continued

---

Ser Ala Leu Arg Arg Asp Val Gln Val Ser Asp Arg Val Arg Ser Ile  
 340 345 350

Ala Lys Asp Val Ala Leu Ala Cys Lys Thr Ser Glu Thr Val Pro Arg  
 355 360 365

Pro Ala  
 370

<210> SEQ ID NO 26  
 <211> LENGTH: 365  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Met Glu Pro Leu Phe Pro Ala Pro Phe Trp Glu Val Ile Tyr Gly Ser  
 1 5 10 15

His Leu Gln Gly Asn Leu Ser Leu Leu Ser Pro Asn His Ser Leu Leu  
 20 25 30

Pro Pro His Leu Leu Leu Asn Ala Ser His Gly Ala Phe Leu Pro Leu  
 35 40 45

Gly Leu Lys Val Thr Ile Val Gly Leu Tyr Leu Ala Val Cys Val Gly  
 50 55 60

Gly Leu Leu Gly Asn Cys Leu Val Met His Thr Lys Met Lys Thr Ala  
 65 70 75 80

Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala Asp Thr Leu Val Leu  
 85 90 95

Leu Thr Leu Pro Phe Gln Gly Thr Asp Ile Leu Leu Gly Phe Trp Pro  
 100 105 110

Phe Gly Asn Ala Leu Cys Lys Thr Val Ile Ala Ile Asp Tyr Tyr Asn  
 115 120 125

Met Phe Thr Ser Thr Phe Thr Leu Thr Ala Met Ser Val Asp Arg Tyr  
 130 135 140

Val Ala Ile Cys His Pro Ile Arg Ala Leu Asp Val Arg Thr Ser Ser  
 145 150 155 160

Lys Ala Gln Ala Val Asn Val Ala Ile Trp Ala Leu Ala Ser Val Val  
 165 170 175

Gly Val Pro Val Ala Ile Met Gly Ser Ala Gln Val Glu Asp Glu Glu  
 180 185 190

Ile Glu Cys Leu Val Glu Ile Pro Thr Pro Gln Asp Tyr Trp Gly Pro  
 195 200 205

Val Phe Ala Ile Cys Ile Phe Leu Phe Ser Phe Ile Val Pro Val Leu  
 210 215 220

Val Ile Ser Val Cys Tyr Ser Leu Met Ile Arg Arg Leu Arg Gly Val  
 225 230 235 240

Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg Arg Ile  
 245 250 255

Thr Arg Leu Val Leu Val Val Val Ala Val Phe Val Gly Cys Trp Thr  
 260 265 270

Pro Val Gln Val Phe Val Leu Ala Gln Gly Leu Gly Val Gln Pro Ser  
 275 280 285

Ser Glu Thr Ala Val Ala Ile Leu Arg Phe Cys Thr Ala Leu Gly Tyr  
 290 295 300

Val Asn Ser Cys Leu Asn Pro Ile Leu Tyr Ala Phe Leu Asp Glu Asn



-continued

---

Ala Leu Gly Tyr Ala Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe  
305 310 315 320

Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Gln Leu Cys Arg Lys Pro  
325 330 335

Cys Gly Arg Pro Asp Pro Ser Ser Phe Ser Arg Ala Arg Glu Ala Thr  
340 345 350

Ala Arg Glu Arg Val Thr Ala Cys Thr Pro Ser Asp Gly Pro Gly Gly  
355 360 365

Gly Ala Ala Ala  
370

<210> SEQ ID NO 28  
<211> LENGTH: 372  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

Met Glu Pro Ala Pro Ser Ala Gly Ala Glu Leu Gln Pro Pro Leu Phe  
1 5 10 15

Ala Asn Ala Ser Asp Ala Tyr Pro Ser Ala Phe Pro Ser Ala Gly Ala  
20 25 30

Asn Ala Ser Gly Pro Pro Gly Ala Arg Ser Ala Ser Ser Leu Ala Leu  
35 40 45

Ala Ile Ala Ile Thr Ala Leu Tyr Ser Ala Val Cys Ala Val Gly Leu  
50 55 60

Leu Gly Asn Val Leu Val Met Phe Gly Ile Val Arg Tyr Thr Lys Met  
65 70 75 80

Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala  
85 90 95

Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Ala Lys Tyr Leu Met Glu  
100 105 110

Thr Trp Pro Phe Gly Glu Leu Leu Cys Lys Ala Val Leu Ser Ile Asp  
115 120 125

Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Thr Met Met Ser Val  
130 135 140

Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu Asp Phe Arg  
145 150 155 160

Thr Pro Ala Lys Ala Lys Leu Ile Asn Ile Cys Ile Trp Val Leu Ala  
165 170 175

Ser Gly Val Gly Val Pro Ile Met Val Met Ala Val Thr Arg Pro Arg  
180 185 190

Asp Gly Ala Val Val Cys Met Leu Gln Phe Pro Ser Pro Ser Trp Tyr  
195 200 205

Trp Asp Thr Val Thr Lys Ile Cys Val Phe Leu Phe Ala Phe Val Val  
210 215 220

Pro Ile Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Leu Leu Arg Leu  
225 230 235 240

Arg Ser Val Arg Leu Leu Ser Gly Ser Lys Glu Lys Asp Arg Ser Leu  
245 250 255

Arg Arg Ile Thr Arg Met Val Leu Val Val Val Gly Ala Phe Val Val  
260 265 270

Cys Trp Ala Pro Ile His Ile Phe Val Ile Val Trp Thr Leu Val Asp  
275 280 285

-continued

---

Ile Asp Arg Arg Asp Pro Leu Val Val Ala Ala Leu His Leu Cys Ile  
 290 295 300

Ala Leu Gly Tyr Ala Asn Ser Ser Leu Asn Pro Val Leu Tyr Ala Phe  
 305 310 315 320

Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Gln Leu Cys Arg Lys Pro  
 325 330 335

Cys Gly Arg Pro Asp Pro Ser Ser Phe Ser Arg Ala Arg Glu Ala Thr  
 340 345 350

Ala Arg Glu Arg Val Thr Ala Cys Thr Pro Ser Asp Gly Pro Gly Gly  
 355 360 365

Gly Ala Ala Ala  
 370

<210> SEQ ID NO 29  
 <211> LENGTH: 380  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

Met Asp Ser Pro Ile Gln Ile Phe Arg Gly Glu Pro Gly Pro Thr Cys  
 1 5 10 15

Ala Pro Ser Ala Cys Leu Pro Pro Asn Ser Ser Ala Trp Phe Pro Gly  
 20 25 30

Trp Ala Glu Pro Asp Ser Asn Gly Ser Ala Gly Ser Glu Asp Ala Gln  
 35 40 45

Leu Glu Pro Ala His Ile Ser Pro Ala Ile Pro Val Ile Ile Thr Ala  
 50 55 60

Val Tyr Ser Val Val Phe Val Val Gly Leu Val Gly Asn Ser Leu Val  
 65 70 75 80

Met Phe Val Ile Ile Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile  
 85 90 95

Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala Leu Val Thr Thr Thr Met  
 100 105 110

Pro Phe Gln Ser Thr Val Tyr Leu Met Asn Ser Trp Pro Phe Gly Asp  
 115 120 125

Val Leu Cys Lys Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr  
 130 135 140

Ser Ile Phe Thr Leu Thr Met Met Ser Val Asp Arg Tyr Ile Ala Val  
 145 150 155 160

Cys His Pro Val Lys Ala Leu Asp Phe Arg Thr Pro Leu Lys Ala Lys  
 165 170 175

Ile Ile Asn Ile Cys Ile Trp Leu Leu Ser Ser Ser Val Gly Ile Ser  
 180 185 190

Ala Ile Val Leu Gly Gly Thr Lys Val Arg Glu Asp Val Asp Val Ile  
 195 200 205

Glu Cys Ser Leu Gln Phe Pro Asp Asp Asp Tyr Ser Trp Trp Asp Leu  
 210 215 220

Phe Met Lys Ile Cys Val Phe Ile Phe Ala Phe Val Ile Pro Val Leu  
 225 230 235 240

Ile Ile Ile Val Cys Tyr Thr Leu Met Ile Leu Arg Leu Lys Ser Val  
 245 250 255

Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg Arg Ile

-continued

---

```

                260                265                270
Thr Arg Leu Val Leu Val Val Val Ala Val Phe Val Val Cys Trp Thr
   275                280                285

Pro Ile His Ile Phe Ile Leu Val Glu Ala Leu Gly Ser Thr Ser His
   290                295                300

Ser Thr Ala Ala Leu Ser Ser Tyr Tyr Phe Cys Ile Ala Leu Gly Tyr
  305                310                315                320

Thr Asn Ser Ser Leu Asn Pro Ile Leu Tyr Ala Phe Leu Asp Glu Asn
   325                330                335

Phe Lys Arg Cys Phe Arg Asp Phe Cys Phe Pro Leu Lys Met Arg Met
   340                345                350

Glu Arg Gln Ser Thr Ser Arg Val Arg Asn Thr Val Gln Asp Pro Ala
   355                360                365

Tyr Leu Arg Asp Ile Asp Gly Met Asn Lys Pro Val
   370                375                380

<210> SEQ ID NO 30
<211> LENGTH: 380
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

Met Glu Ser Pro Ile Gln Ile Phe Arg Gly Glu Pro Gly Pro Thr Cys
  1                5                10                15

Ala Pro Ser Ala Cys Leu Pro Pro Asn Ser Ser Ala Trp Phe Pro Gly
   20                25                30

Trp Ala Glu Pro Asp Ser Asn Gly Ser Ala Gly Ser Glu Asp Ala Gln
   35                40                45

Leu Glu Pro Ala His Ile Ser Pro Ala Ile Pro Val Ile Ile Thr Ala
   50                55                60

Val Tyr Ser Val Val Phe Val Val Gly Leu Val Gly Asn Ser Leu Val
   65                70                75                80

Met Phe Val Ile Ile Arg Tyr Thr Lys Met Lys Thr Ala Thr Asn Ile
   85                90                95

Tyr Ile Phe Asn Leu Ala Leu Ala Asp Ala Leu Val Thr Thr Thr Met
  100                105                110

Pro Phe Gln Ser Thr Val Tyr Leu Met Asn Ser Trp Pro Phe Gly Asp
  115                120                125

Val Leu Cys Lys Ile Val Ile Ser Ile Asp Tyr Tyr Asn Met Phe Thr
  130                135                140

Ser Ile Phe Thr Leu Thr Met Met Ser Val Asp Arg Tyr Ile Ala Val
  145                150                155                160

Cys His Pro Val Lys Ala Leu Asp Phe Arg Thr Pro Leu Lys Ala Lys
  165                170                175

Ile Ile Asn Ile Cys Ile Trp Leu Leu Ser Ser Ser Val Gly Ile Ser
  180                185                190

Ala Ile Val Leu Gly Gly Thr Lys Val Arg Glu Asp Val Asp Val Ile
  195                200                205

Glu Cys Ser Leu Gln Phe Pro Asp Asp Asp Tyr Ser Trp Trp Asp Leu
  210                215                220

Phe Met Lys Ile Cys Val Phe Ile Phe Ala Phe Val Ile Pro Val Leu
  225                230                235                240

```

-continued

---

```

Ile Ile Ile Val Cys Tyr Thr Leu Met Ile Leu Arg Leu Lys Ser Val
      245                250                255

Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg Arg Ile
      260                265                270

Thr Arg Leu Val Leu Val Val Val Ala Val Phe Val Val Cys Trp Thr
      275                280                285

Pro Ile His Ile Phe Ile Leu Val Glu Ala Leu Gly Ser Thr Ser His
      290                295                300

Ser Thr Ala Ala Leu Ser Ser Tyr Tyr Phe Cys Ile Ala Leu Gly Tyr
      305                310                315                320

Thr Asn Ser Ser Leu Asn Pro Ile Leu Tyr Ala Phe Leu Asp Glu Asn
      325                330                335

Phe Lys Arg Cys Phe Arg Asp Phe Cys Phe Pro Leu Lys Met Arg Met
      340                345                350

Glu Arg Gln Ser Thr Ser Arg Val Arg Asn Thr Val Gln Asp Pro Ala
      355                360                365

Tyr Leu Arg Asp Ile Asp Gly Met Asn Lys Pro Val
      370                375                380
    
```

```

<210> SEQ ID NO 31
<211> LENGTH: 493
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

```

<400> SEQUENCE: 31
    
```

```

Met Cys Leu His Arg Arg Val Pro Ser Glu Glu Thr Tyr Ser Leu Asp
1      5      10      15

Arg Phe Ala Gln Asn Pro Pro Leu Phe Pro Pro Pro Ser Leu Pro Ala
      20      25      30

Ser Glu Ser Arg Met Ala His Ala Pro Leu Leu Gln Arg Cys Gly Ala
      35      40      45

Ala Arg Thr Gly Phe Cys Lys Lys Gln Gln Glu Leu Trp Gln Arg Arg
      50      55      60

Lys Glu Ala Ala Glu Ala Leu Gly Thr Arg Lys Val Ser Val Leu Leu
      65      70      75      80

Ala Thr Ser His Ser Gly Ala Arg Pro Ala Val Ser Thr Met Asp Ser
      85      90      95

Ser Ala Ala Pro Thr Asn Ala Ser Asn Cys Thr Asp Ala Leu Ala Tyr
      100     105     110

Ser Ser Cys Ser Pro Ala Pro Ser Pro Gly Ser Trp Val Asn Leu Ser
      115     120     125

His Leu Asp Gly Asn Leu Ser Asp Pro Cys Gly Pro Asn Arg Thr Asp
      130     135     140

Leu Gly Gly Arg Asp Ser Leu Cys Pro Pro Thr Gly Ser Pro Ser Met
      145     150     155     160

Ile Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser Ile Val Cys Val Val
      165     170     175

Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val Ile Val Arg Tyr Thr
      180     185     190

Lys Met Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala
      195     200     205

Asp Ala Leu Ala Thr Ser Thr Leu Pro Phe Gln Ser Val Asn Tyr Leu
      210     215     220
    
```

-continued

---

```

Met Gly Thr Trp Pro Phe Gly Thr Ile Leu Cys Lys Ile Val Ile Ser
225                230                235                240

Ile Asp Tyr Tyr Asn Met Phe Thr Ser Ile Phe Thr Leu Cys Thr Met
                245                250                255

Ser Val Asp Arg Tyr Ile Ala Val Cys His Pro Val Lys Ala Leu Asp
                260                265                270

Phe Arg Thr Pro Arg Asn Ala Lys Ile Ile Asn Val Cys Asn Trp Ile
                275                280                285

Leu Ser Ser Ala Ile Gly Leu Pro Val Met Phe Met Ala Thr Thr Lys
290                295                300

Tyr Arg Gln Gly Ser Ile Asp Cys Thr Leu Thr Phe Ser His Pro Thr
305                310                315

Trp Tyr Trp Glu Asn Leu Leu Lys Ile Cys Val Phe Ile Phe Ala Phe
                325                330                335

Ile Met Pro Val Leu Ile Ile Thr Val Cys Tyr Gly Leu Met Ile Leu
340                345                350

Arg Leu Lys Ser Val Arg Met Leu Ser Gly Ser Lys Glu Lys Asp Arg
355                360                365

Asn Leu Arg Arg Ile Thr Arg Met Val Leu Val Val Val Ala Val Phe
370                375                380

Ile Val Cys Trp Thr Pro Ile His Ile Tyr Val Ile Ile Lys Ala Leu
385                390                395

Val Thr Ile Pro Glu Thr Thr Phe Gln Thr Val Ser Trp His Phe Cys
405                410                415

Ile Ala Leu Gly Tyr Thr Asn Ser Cys Leu Asn Pro Val Leu Tyr Ala
420                425                430

Phe Leu Asp Glu Asn Phe Lys Arg Cys Phe Arg Glu Phe Cys Ile Pro
435                440                445

Thr Ser Ser Asn Ile Glu Gln Gln Asn Ser Thr Arg Ile Arg Gln Asn
450                455                460

Thr Arg Asp His Pro Ser Thr Ala Asn Thr Val Asp Arg Thr Asn His
465                470                475                480

Gln Leu Glu Asn Leu Glu Ala Glu Thr Ala Pro Leu Pro
485                490

```

```

<210> SEQ ID NO 32
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-
terminus at the P1

```

```

<400> SEQUENCE: 32

```

```

Asp Glu Ala Asn Gln
1          5

```

```

<210> SEQ ID NO 33
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus
at the P1 residue of the scissile bond of the BoNT/A cleavage site

```

```

<400> SEQUENCE: 33

```

-continued

---

Ile Asp Glu Ala Asn Gln  
1 5

<210> SEQ ID NO 34  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 34

Arg Ile Asp Glu Ala Asn Gln  
1 5

<210> SEQ ID NO 35  
<211> LENGTH: 8  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 35

Thr Arg Ile Asp Glu Ala Asn Gln  
1 5

<210> SEQ ID NO 36  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 36

Lys Thr Arg Ile Asp Glu Ala Asn Gln  
1 5

<210> SEQ ID NO 37  
<211> LENGTH: 10  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1

<400> SEQUENCE: 37

Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln  
1 5 10

<210> SEQ ID NO 38  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 38

Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln  
1 5 10

<210> SEQ ID NO 39

-continued

---

<211> LENGTH: 12  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 39

Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln  
1 5 10

<210> SEQ ID NO 40  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxylated  
carboxyl-terminus at the P1 residue of the scissile bond of the  
BoNT/A cleavage site  
<220> FEATURE:  
<221> NAME/KEY: SITE  
<222> LOCATION: (13)...(13)  
<223> OTHER INFORMATION: carboxylated glutamine

<400> SEQUENCE: 40

Cys Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln  
1 5 10

<210> SEQ ID NO 41  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1

<400> SEQUENCE: 41

Arg Ile Asp Glu Ala Asn Lys  
1 5

<210> SEQ ID NO 42  
<211> LENGTH: 8  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 42

Ala Arg Ile Asp Glu Ala Asn Lys  
1 5

<210> SEQ ID NO 43  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 43

Lys Ala Arg Ile Asp Glu Ala Asn Lys  
1 5

<210> SEQ ID NO 44  
<211> LENGTH: 10

-continued

---

<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 44

Asn Lys Ala Arg Ile Asp Glu Ala Asn Lys  
1 5 10

<210> SEQ ID NO 45  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 45

Met Asn Lys Ala Arg Ile Asp Glu Ala Asn Lys  
1 5 10

<210> SEQ ID NO 46  
<211> LENGTH: 12  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxyl-terminus  
at the P1 residue of the scissile bond of the BoNT/A cleavage site

<400> SEQUENCE: 46

Asp Met Asn Lys Ala Arg Ile Asp Glu Ala Asn Lys  
1 5 10

<210> SEQ ID NO 47  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen having a free carboxylated  
carboxyl-terminus at the P1 residue of the scissile bond of the  
BoNT/A cleavage site  
<220> FEATURE:  
<221> NAME/KEY: SITE  
<222> LOCATION: (13)...(13)  
<223> OTHER INFORMATION: Carboxylated lysine

<400> SEQUENCE: 47

Cys Asp Met Asn Lys Ala Arg Ile Asp Glu Ala Asn Lys  
1 5 10

<210> SEQ ID NO 48  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: SNAP-25 antigen

<400> SEQUENCE: 48

Cys Gly Gly Gly Arg Ile Asp Glu Ala Asn Gln  
1 5 10

<210> SEQ ID NO 49  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence

-continued

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: SNAP-25 antigen

&lt;400&gt; SEQUENCE: 49

Cys Gly Gly Gly Arg Ile Asp Glu Ala Asn Lys  
 1                    5                    10

&lt;210&gt; SEQ ID NO 50

&lt;211&gt; LENGTH: 88

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: BirA-HisTag?-SNAP-25-134-197

&lt;400&gt; SEQUENCE: 50

Met Gly Gly Gly Leu Asn Asp Ile Phe Glu Ala Gln Lys Ile Glu Trp  
 1                    5                    10                    15

His His His His His His His His Ile Arg Arg Val Thr Asn Asp Ala  
                   20                    25                    30

Arg Glu Asn Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile  
                   35                    40                    45

Gly Asn Leu Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr  
                   50                    55                    60

Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys  
 65                    70                    75                    80

Thr Arg Ile Asp Glu Ala Asn Gln  
                   85

&lt;210&gt; SEQ ID NO 51

&lt;211&gt; LENGTH: 97

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: BirA-HisTag?-SNAP-25-134-206

&lt;400&gt; SEQUENCE: 51

Met Gly Gly Gly Leu Asn Asp Ile Phe Glu Ala Gln Lys Ile Glu Trp  
 1                    5                    10                    15

His His His His His His His His Ile Arg Arg Val Thr Asn Asp Ala  
                   20                    25                    30

Arg Glu Asn Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile  
                   35                    40                    45

Gly Asn Leu Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr  
                   50                    55                    60

Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys  
 65                    70                    75                    80

Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser  
                   85                    90                    95

Gly

&lt;210&gt; SEQ ID NO 52

&lt;211&gt; LENGTH: 16

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: BirA peptide

&lt;400&gt; SEQUENCE: 52

-continued

---

Gly Gly Gly Leu Asn Asp Ile Phe Glu Ala Gln Lys Ile Glu Trp His  
1 5 10 15

<210> SEQ ID NO 53  
 <211> LENGTH: 7570  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: pQBI-25/GFP-BoNT/A-LC expression construct.

<400> SEQUENCE: 53

gacggatcgg gagatctccc gatcccctat ggtcgactct cagtacaatc tgctctgatg 60  
 ccgcatagtt aagccagtat ctgtccctg cttgtgtgtt ggaggtcgct gagtagtgcg 120  
 cgagcaaaat ttaagtacata acaaggcaag gcttgaccga caattgcatg aagaatctgc 180  
 ttagggtagt gcgttttgcg ctgcttcgcc tcgaggcctg gccattgcat acgttgatc 240  
 catatcataa tatgtacatt tatattggct catgtccaac attaccgcca tgttgacatt 300  
 gattattgac tagttattaa tagtaatcaa ttacggggtc attagtcat agcccatata 360  
 tggagtccg cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc 420  
 cccgccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc 480  
 attgacgtca atgggtggag tatttacggt aaactgcca cttggcagta catcaagtgt 540  
 atcatatgcc aagtaagccc cctattgacg tcaatgacgg taaatggccc gcttggcatt 600  
 atgcccagta catgacctta tgggactttc ctacttgga gtacatctac gtattagtca 660  
 tcgctattac catggtgatg cggttttggc agtacatcaa tgggctgga tagcggtttg 720  
 actcacgggg atttccaagt ctccaccoca ttgacgtcaa tgggagtttg ttttggcacc 780  
 aaaatcaacg ggactttcca aaatgtcgta acaactccgc cccattgacg caaatggcg 840  
 gtaggcgtgt acggtgggag gtctatataa gcagagctcg tttagtgaac cgtcagatcg 900  
 cctggagacg ccatccacgc tgttttgacc tccatagaag acaccgggac cgatccagcc 960  
 tccgcgggcc accatggagg gcccggttac cgttacgga tccagatc tggcgggccg 1020  
 ctcagcaagc ttccgaatt cgggaggcgg aggtggagct agcaaaggag aagaactctt 1080  
 cactggagtt gtcccattc ttgttgaatt agatggtgat gttaacggcc acaagttctc 1140  
 tgtcagtga gaggtgaag gtgatgcaac atacggaaaa cttaccctga agttcatctg 1200  
 cactactggc aaactgcctg ttccatggcc aacactagtc actactctgt gctatgggtg 1260  
 tcaatgcttt tcaagatacc cggatcatat gaaacggcat gactttttca agagtgccat 1320  
 gcccgaaggt tatgtacagg aaaggaccat cttcttcaaa gatgacggca actacaagac 1380  
 acgtgctgaa gtcaagtttg aaggtgatac ccttgtaaat agaatcgagt taaaaggat 1440  
 tgacttcaag gaagatggca acattctggg acacaaattg gaatacaact ataactcaca 1500  
 caatgtatac atcatggcag acaaacaaaa gaatggaatc aaagtgaact tcaagaccgc 1560  
 ccacaacatt gaagatggaa cgtttcaact agcagaccat tatcaacaaa atactccaat 1620  
 tggcagatggc cctgtccttt taccagacaa ccattacctg tccacacaat ctgccctttc 1680  
 gaaagatccc aacgaaaaga gagaccacat ggtccttctt gagttttaa cagctgctgg 1740  
 gattacacat ggcctggatg aactgtacaa catcgatgga ggcggagggtg gaccttttgt 1800  
 taataacaaa ttttaattata aagatcctgt aaatggtgtt gatattgctt atataaaaa 1860  
 tccaaatgca ggacaaatgc aaccagtaaa agcttttaaa attcataata aaatatgggt 1920

-continued

---

tattccagaa agagatacat ttacaaatcc tgaagaagga gatttaaate caccaccaga	1980
agcaaaaaca gttccagttt catattatga ttcaacatat ttaagtacag ataataaaaa	2040
agataattat ttaaagggag ttacaaaatt atttgagaga atttattcaa ctgatcttgg	2100
aagaatggtg ttaacatcaa tagtaagggg aataccattt tgggggtggaa gtacaataga	2160
tacagaatta aaagtatttg atactaattg tattaatgtg atacaaccag atggtagtta	2220
tagatcagaa gaacttaate tagtaataat aggacctca gctgatatta tacagtttga	2280
atgtaaaagc tttggacatg aagttttgaa tcttacgcga aatggttatg gctctactca	2340
atacattaga ttagcccag attttacatt tggttttgag gagtcaactg aagttgatac	2400
aatcctctt ttaggtgagc gcaaatgtg tacagatcca gcagtaacat tagcacatga	2460
actatacat gctggacata gattatatgg aatagcaatt aatccaaata gggtttttaa	2520
agtaaatact aatgcctatt atgaaatgag tgggttagaa gtaagctttg aggaacttag	2580
aacatttggg ggacatgatg caaagtttat agatagtta caggaaaacg aatttcgtct	2640
atattattat aataagttta aagatatagc aagtacactt aataaagcta aatcaatagt	2700
aggtaactt gcttcattac agtatatgaa aaatgttttt aaagagaaat atctcctatc	2760
tgaagataca tctggaaaa tttcggtaga taaattaaaa tttgataagt tatacaaaat	2820
gttaacagag atttacacag aggataattt tgttaagttt tttaaagtac ttaacagaaa	2880
aacatatttg aattttgata aagccgtatt taagataaat atagtaccta aggtaaatta	2940
cacaatata gatggattta atttaagaaa tacaattta gcagcaaac ttaatggta	3000
aaatacagaa attaataata tgaattttac taaactaaaa aattttactg gattgtttga	3060
attttataag ttgctatgtg taagagggat aatcacttgc aaatgaacgc gttggcccta	3120
ttctatagtg tcacctaaat gctagagctc gctgatcagc ctgcactgtg ccttctagtt	3180
gccagccatc tgttgtttgc cctcccccg tgccttcctt gacctggaa ggtgccactc	3240
ccactgtcct ttctaataa aatgagaaa ttgcatcgca ttgtctgagt aggtgtcatt	3300
ctattctggg ggggtgggtg gggcaggaca gcaaggggga ggattgggaa gacaatagca	3360
ggcatgctgg ggatgcggtg ggctctatgg cttctgaggc ggaagaacc agctggggct	3420
ctaggggta tccccacgcg ccctgtagcg gcgcattaag cgcggcgggt gtgggtgta	3480
cgcgcagcgt gaccgtaca cttgccagcg ccctagcgcg cgctccttc gctttcttc	3540
cttctttct cgcacgttc gccggcttc cccgtcaagc tctaatacgg ggcacccct	3600
tagggttccg atttagtct ttacggcacc tcgacccca aaaacttgat tagggtgatg	3660
gttcacgtag tgggcatcg ccctgataga cggtttttcg cctttgacg ttggagtcca	3720
cgttctttaa tagtggactc ttgttccaaa ctggaacaac actcaaccct atctcggct	3780
attctttga tttataaggg attttgggga tttcggccta ttggttaaaa aatgagctga	3840
tttaacaaaa atttaacgcg aattaattct gtggaatgtg tgtcagttag ggtgtggaaa	3900
gtccccaggc tccccaggca ggcagaagta tgcaaagcat gcactcfaat tagtcagcaa	3960
ccaggtgtgg aaagtccca ggcctccag caggcagaag tatgcaaagc atgcatctca	4020
attagtcagc aaccatagtc ccgcccctaa ctccgcccct cccgccccta actccgccc	4080
gttccgccc tctccgccc catggctgac taattttttt tatttatgca gaggcgagg	4140
ccgctctgc ctctgagcta ttccagaagt agtgaggagg cttttttgga ggctaggt	4200

-continued

---

tttgcaaaaa gctccccgga gcttgatat ccattttcgg atctgatcaa gagacaggat	4260
gaggatcggt tcgcatgatt gaacaagatg gattgcacgc aggttctccg gccgcttggg	4320
tggagaggct attcggctat gactgggcac aacagacaat cggctgctct gatgccgccg	4380
tgttccggct gtcagcgcag gggcgcccgg ttctttttgt caagaccgac ctgtccggtg	4440
ccctgaatga actgcaggac gaggcagcgc ggctatcgtg gctggccacg acgggcggtc	4500
cttgccgacg tgtgctcgac gttgtcactg aagcgggaag ggactggctg ctattgggcg	4560
aagtgccggg gcaggatctc ctgtcatctc accttgctcc tgccgagaaa gtatccatca	4620
tggctgatgc aatgcggcgg ctgcatacgc ttgatccggc tacctgccc ttcgaccacc	4680
aagcgaaca tcgcatcgag cgagcacgta ctcgatgga agccggtctt gtcgatcagg	4740
atgatctgga cgaagagcat caggggctcg cgccagccga actgttcgcc aggctcaagg	4800
cgcgcatgcc cgacggcgag gatctcgtcg tgaccatgg cgatgcctgc ttgccgaata	4860
tcattggtgga aaatggccc tttctggat tcactgactg tggccggctg ggtgtggcgg	4920
accgctatca ggacatagcg ttggctaccc gtgatattgc tgaagagctt ggcggcgaat	4980
gggctgaccg ctccctcgtg ctttacggta tcgccgctcc cgattcgcag cgcatcgct	5040
tctatcgctt tcttgacgag ttcttctgag cgggactctg gggttcgaaa tgaccgacca	5100
agcagcgcgc aacctgccat cagcagatct cgattccacc gccgcttct atgaaagggt	5160
gggcttcgga atcgttttcc gggacgccgg ctggatgatc ctccagcgcg gggatctcat	5220
gctggagttc ttgcccccacc ccaacttgtt tattgcagct tataatggtt acaataaaag	5280
caatagcadc acaaatttca caaataaagc atttttttca ctgcattcta gttgtggttt	5340
gtccaaaact atcaatgtat cttatcatgt ctgtataccg tcgacctcta gctagagctt	5400
ggcgtaatca tggatcatag tgtttcctgt gtgaaattgt tatccgctca caattccaca	5460
caacatacga gccggaagca taaagtgtaa agcctggggt gcctaagtag tgagctaaact	5520
cacattaatt gcggttcgct cactgccgcg tttccagtcg ggaaacctgt cgtgccagct	5580
gcattaatga atcgccaac gcgcggggag aggcggtttg cgtattgggc gctcttcgc	5640
ttcctcgtc actgactcgc tgccctcggc cgttcggctg cggcgagcgg tatcagctca	5700
ctcaaaggcg gtaatacggc tatccacaga atcaggggat aacgcaggaa agaacaatgtg	5760
agcaaaaagg cagcaaaaag ccaggaaccg taaaaaggcc gcggttctgg cgtttttcca	5820
taggctccgc ccccctgacg agcatcaca aaatcgacgc tcaagtcaga ggtggcgaaa	5880
cccgacagga ctataaagat accagcgtt tccccctgga agctccctcg tgcgctctcc	5940
tgttccgacc ctgcccctta ccggatacct gtccgccttt ctcccttcgg gaagcgtggc	6000
gctttctcaa tgctcagct gtaggtatct cagttcggtg taggtcgttc gctccaagct	6060
gggctgtgtg cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg	6120
tcttgagtcc aacctggtaa gacacgactt atcgccactg gcagcagcca ctggtaacag	6180
gattagcaga gcgaggtatg taggcggtgc tacagagttc ttgaagtggg gcctaaacta	6240
cggtacact agaaggacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg	6300
aaaaagagtt ggtagctctt gatccggcaa acaaaccacc gctggtagcg gtggtttttt	6360
tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt	6420
ttctacgggg tctgacgctc agtggaaaca aaactcagct taagggattt tggatcatgag	6480

-continued

---

```

attatcaaaa aggatcttca cctagatcct tttaaattaa aatgaagtt ttaatcaat 6540
ctaaagtata tatgagtaaa cttgggtctga cagttaccaa tgcttaatca gtgaggcacc 6600
tatctcagcg atctgtctat ttcgttcac ccatagttgcc tgactccccg tcgtgtagat 6660
aactacgata cgggagggct taccatctgg ccccagtgtc gcaatgatac cgcgagaccc 6720
acgtccaccg gctccagatt tatcagcaat aaaccagcca gccggaaggg ccgagcgcag 6780
aagtggctct gcaactttat cgcctccat ccagtctatt aattggtgcc gggagactag 6840
agtaagtagt tcgccagtta atagtttgcg caacgttggt gccattgcta caggcatcgt 6900
gggtgcacgc tcgtcgtttg gtatggcttc atcagctcc ggttcccaac gatcaaggcg 6960
agttacatga tccccatgt tgtgcaaaaa agcggttagc tccttcggtc ctccgatcgt 7020
tgtcagaagt aagtggccg cagtgttacc actcatgggt atggcagcac tgcataattc 7080
tcttactgtc atgccatccg taagatgctt ttctgtgact ggtgagtact caaccaagtc 7140
attctgagaa tagtgatgc ggcgaccgag ttgctcttgc cggcgtcaa tacgggataa 7200
taccgcgcca catagcagaa ctttaaaagt gctcatcatt ggaaaacgtt ctcgggggcg 7260
aaaactctca aggatcttac cgctgttgag atccagttcg atgtaacca ctcgtgcacc 7320
caactgatct tcagcatctt ttaactttcac cagcgtttct gggtgagcaa aaacaggaag 7380
gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa tgttaatac tcatactctt 7440
cctttttcaa tattattgaa gcatttatca gggttattgt ctcatgagcg gatacatatt 7500
tgaatgtatt tagaaaaata acaaaatagg ggttccgcgc acatttcccc gaaaagtgcc 7560
acctgacgtc 7570

```

```

<210> SEQ ID NO 54
<211> LENGTH: 682
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: GFP-BoNT/A light chain amino acid sequence.

```

<400> SEQUENCE: 54

```

Ala Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val
1           5           10          15
Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu
20          25          30
Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys
35          40          45
Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu
50          55          60
Cys Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Arg
65          70          75          80
His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg
85          90          95
Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val
100         105         110
Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile
115         120         125
Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn
130         135         140

```

-continued

---

Tyr	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	145	150	155	160
Ile	Lys	Val	Asn	Phe	Lys	Thr	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	165	170	175	
Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	Pro	180	185	190	
Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln	Ser	Ala	Leu	Ser	195	200	205	
Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe	Val	210	215	220	
Thr	Ala	Ala	Gly	Ile	Thr	His	Gly	Met	Asp	Glu	Leu	Tyr	Asn	Ile	Asp	225	230	235	240
Gly	Gly	Gly	Gly	Gly	Pro	Phe	Val	Asn	Lys	Gln	Phe	Asn	Tyr	Lys	Asp	245	250	255	
Pro	Val	Asn	Gly	Val	Asp	Ile	Ala	Tyr	Ile	Lys	Ile	Pro	Asn	Ala	Gly	260	265	270	
Gln	Met	Gln	Pro	Val	Lys	Ala	Phe	Lys	Ile	His	Asn	Lys	Ile	Trp	Val	275	280	285	
Ile	Pro	Glu	Arg	Asp	Thr	Phe	Thr	Asn	Pro	Glu	Glu	Gly	Asp	Leu	Asn	290	295	300	
Pro	Pro	Pro	Glu	Ala	Lys	Gln	Val	Pro	Val	Ser	Tyr	Tyr	Asp	Ser	Thr	305	310	315	320
Tyr	Leu	Ser	Thr	Asp	Asn	Glu	Lys	Asp	Asn	Tyr	Leu	Lys	Gly	Val	Thr	325	330	335	
Lys	Leu	Phe	Glu	Arg	Ile	Tyr	Ser	Thr	Asp	Leu	Gly	Arg	Met	Leu	Leu	340	345	350	
Thr	Ser	Ile	Val	Arg	Gly	Ile	Pro	Phe	Trp	Gly	Gly	Ser	Thr	Ile	Asp	355	360	365	
Thr	Glu	Leu	Lys	Val	Ile	Asp	Thr	Asn	Cys	Ile	Asn	Val	Ile	Gln	Pro	370	375	380	
Asp	Gly	Ser	Tyr	Arg	Ser	Glu	Glu	Leu	Asn	Leu	Val	Ile	Ile	Gly	Pro	385	390	395	400
Ser	Ala	Asp	Ile	Ile	Gln	Phe	Glu	Cys	Lys	Ser	Phe	Gly	His	Glu	Val	405	410	415	
Leu	Asn	Leu	Thr	Arg	Asn	Gly	Tyr	Gly	Ser	Thr	Gln	Tyr	Ile	Arg	Phe	420	425	430	
Ser	Pro	Asp	Phe	Thr	Phe	Gly	Phe	Glu	Glu	Ser	Leu	Glu	Val	Asp	Thr	435	440	445	
Asn	Pro	Leu	Leu	Gly	Ala	Gly	Lys	Phe	Ala	Thr	Asp	Pro	Ala	Val	Thr	450	455	460	
Leu	Ala	His	Glu	Leu	Ile	His	Ala	Gly	His	Arg	Leu	Tyr	Gly	Ile	Ala	465	470	475	480
Ile	Asn	Pro	Asn	Arg	Val	Phe	Lys	Val	Asn	Thr	Asn	Ala	Tyr	Tyr	Glu	485	490	495	
Met	Ser	Gly	Leu	Glu	Val	Ser	Phe	Glu	Glu	Leu	Arg	Thr	Phe	Gly	Gly	500	505	510	
His	Asp	Ala	Lys	Phe	Ile	Asp	Ser	Leu	Gln	Glu	Asn	Glu	Phe	Arg	Leu	515	520	525	
Tyr	Tyr	Tyr	Asn	Lys	Phe	Lys	Asp	Ile	Ala	Ser	Thr	Leu	Asn	Lys	Ala	530	535	540	
Lys	Ser	Ile	Val	Gly	Thr	Thr	Ala	Ser	Leu	Gln	Tyr	Met	Lys	Asn	Val				

-continued

---

545	550	555	560
Phe Lys Glu Lys Tyr Leu Leu Ser Glu Asp Thr Ser Gly Lys Phe Ser	565	570	575
Val Asp Lys Leu Lys Phe Asp Lys Leu Tyr Lys Met Leu Thr Glu Ile	580	585	590
Tyr Thr Glu Asp Asn Phe Val Lys Phe Phe Lys Val Leu Asn Arg Lys	595	600	605
Thr Tyr Leu Asn Phe Asp Lys Ala Val Phe Lys Ile Asn Ile Val Pro	610	615	620
Lys Val Asn Tyr Thr Ile Tyr Asp Gly Phe Asn Leu Arg Asn Thr Asn	625	630	640
Leu Ala Ala Asn Phe Asn Gly Gln Asn Thr Glu Ile Asn Asn Met Asn	645	650	655
Phe Thr Lys Leu Lys Asn Phe Thr Gly Leu Phe Glu Phe Tyr Lys Leu	660	665	670
Leu Cys Val Arg Gly Ile Ile Thr Ser Lys	675	680	

<210> SEQ ID NO 55  
 <211> LENGTH: 6259  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: pQBI-25/GFP expression construct.

<400> SEQUENCE: 55

```

gacggatcgg gagatctccc gatcccctat ggtcgactct cagtacaatc tgetctgatg      60
ccgcatagtt aagccagtat ctgtccctcg cttgtgtggt ggaggtcgct gtagtagtgcg      120
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc      180
ttagggttag cggttttgcg ctgcttcgcc tcgaggcctg gccattgcat acgttgtatc      240
catatcataa tatgtacatt tatattggct catgtccaac attaccgcca tgttgacatt      300
gattattgac tagttattaa tagtaatcaa ttacggggtc attagttcat agcccatata      360
tggagttccg cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc      420
cccgccatt gagtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc      480
attgacgtca atgggtggag tatttacggt aaactgccca cttggcagta catcaagtgt      540
atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt      600
atgccagta catgacctta tgggactttc ctacttggca gtacatctac gtattagtca      660
tcgtattac catggtgatg cggttttggc agtacatcaa tggcgtgga tagcggtttg      720
actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc      780
aaaatcaacg ggactttcca aaatgtcgta acaactccgc ccattgacg caaatgggcg      840
gtaggcgtgt acggtgggag gtctatataa gcagagctcg tttagtgaac cgtcagatcg      900
cctggagacg ccatccacgc tgttttgacc tccatagaag acaccgggac cgatccagcc      960
tccgcgggcc accatggagg gcccggttac cggtaccgga tccagatata tgggcggccg     1020
ctcagcaagc ttocgaatt cgggaggcgg aggtggagct agcaaaggag aagaactctt     1080
cactggagtt gtoccaattc ttgtgaatt agatggtgat gttaacggcc acaagttctc     1140
tgtcagtga gaggtgaag gtgatgcaac atacggaaaa cttaccctga agttcatctg     1200
    
```

-continued

---

cactactggc aaactgcctg ttccatggcc aacactagtc actactctgt gctatgggtg	1260
tcaatgcttt tcaagatacc eggatcatat gaaacggcat gactttttca agagtgccat	1320
gcccgaaggt tatgtacagg aaaggaccat cttcttcaaa gatgacggca actacaagac	1380
acgtgctgaa gtcaagtttg aaggtgatac ccttgtaaat agaatcgagt taaaaggat	1440
tgacttcaag gaagatggca acattctggg acacaattg gaatacaact ataactcaca	1500
caatgtatac atcatggcag acaaacaaaa gaatggaatc aaagtgaact tcaagaccg	1560
ccacaacatt gaagatggaa gcgttcaact agcagaccat tatcaacaaa atactccaat	1620
tggcgatggc cctgtccttt taccagacaa ccattacctg tccacacaat ctgccctttc	1680
gaaagatccc aacgaaaaa gagaccacat ggtccttctt gagttttaa cagctgctgg	1740
gattacacat ggcctggatg aactgtacaa catcgatgga ggcggagggtg gatgaacgag	1800
ttggccctat tctatagtgt cacctaaatg ctagagctcg ctgatcagcc tcgactgtgc	1860
cttctagtgt ccagccatct gttgtttgcc cctccccctg gccttccttg accctggaag	1920
gtgccactcc cactgtcctt tcctaataaa atgaggaaat tgcacgcacat tgtctgagta	1980
gggtgcattc tattctgggg ggtgggggtgg ggcaggacag caagggggag gattgggaag	2040
acaatagcag gcctgctggg gatgcgggtg gctctatggc ttctgaggcg gaaagaacca	2100
gctggggctc tagggggat cccacgcgc cctgtagcgg cgcattaagc gcggcgggtg	2160
tggtgggtac gcgcagcgtg accgctacac ttgccagcgc cctagcgcgc gctcctttcg	2220
ctttcttccc ttctttctc gccacgttcg ccggctttcc cgcgcaagct ctaaatcggg	2280
gcatcccttt agggttccga tttagtgtt tacggcacct cgaccccaaa aaacttgatt	2340
agggatgagg ttcacgtagt gggccatcgc cctgatagac ggtttttcgc cctttgacgt	2400
tggagtccac gttctttaat agtggactct tgttccaaac tggacaaca ctaacccta	2460
tctcggctca tttctttgat ttataaggga ttttggggat ttcggcctat tggtaaaaa	2520
atgagctgat ttaacaaaaa ttaaacgca attaatctg tggaatgtg gtcagttagg	2580
gtgtgaaag tccccaggct cccaggcag gcagaagat gcaaaagcat catctcaatt	2640
agtcagcaac caggtgtgga aagtccccag gctccccagc aggcagaagt atgcaagca	2700
tgcactcaa ttagtcagca accatagtcc cgcctaac tccgccatc ccgccctaa	2760
ctccgccag tccgccat tctccgccc atggctgact aatTTTTTTT atttatgcag	2820
aggccgaggc cgcctctgcc tctgagctat tccagaagta gtgaggaggc ttttttgag	2880
gcctaggctt ttgcaaaaaa ctccccggag cttgtatata cttttcgga tctgatcaag	2940
agacaggatg aggatcgtt cgcctgattg aacaagatg attgcacga ggttctccg	3000
ccgcttgggt ggagaggcta ttcggctatg actgggcaca acagacaatc ggtgctctg	3060
atgccgcgt gttccggctg tcagcgcagg ggcgcccgtt tctttttgtc aagaccgac	3120
tgtccggtgc cctgaatgaa ctgcaggacg aggcagcgcg gctatcgtgg ctggccacga	3180
cggcggttcc ttgcgcagct gtgctcgacg ttgtcactga agcgggaagg gactggtgc	3240
tattgggoga agtccgggg caggatctcc tgtcatctca ccttctcct gccgagaaag	3300
tatccatcat ggtgatgca atgcggcggc tgcatacgt tgatccggt acctgccat	3360
tcgaccacca agcgaacat cgcctcagc gagcacgtac tcggatggaa gccggtcttg	3420
tcgatcagga tgactggag gaagagcacc aggggctcgc gccagccgaa ctgttcgca	3480

-continued

---

ggctcaaggc gcgcatgccc gacggcgagg atctcgtcgt gacccatggc gatgcctgct	3540
tgcggaatat catggtgaa aatggccgct tttctggatt catcgactgt ggcggctgg	3600
gtgtggcgga ccgctatcag gacatagcgt tggctaccgg tgatattgct gaagagcttg	3660
gcgcggaatg ggctgaccgc ttctcgtgc tttacggat cgccgctccc gattcgcagc	3720
gcatacgcctt ctatgcctt cttgacgagt tcttctgagc gggactctgg ggttcgaaat	3780
gaccgaccaa gcgacgccc acctgccatc acgagatttc gattccaccg ccgccttcta	3840
tgaagggttg ggcttcggaa tcgttttccg ggacgcccgc tggatgatcc tccagcggg	3900
ggatctcatg ctggagttct tcgcccacc caactgtttt attgcagctt ataatggta	3960
caaataaagc aatagcatca caaatttcc acataaagca ttttttccac tgcattctag	4020
ttgtggttg tccaaactca tcaatgtatc ttatcatgct tgtataccgt cgacctag	4080
ctagagcttg gcgtaatcat ggtcatagct gtttctgtg tgaattggt atccgctcac	4140
aattccacac aacatacagc ccggaagcat aaagtgtaaa gcctggggtg cctaagtagt	4200
gagctaacct acattaattg cgttgccctc actgcccgtt tccagtcgg gaaacctgtc	4260
gtgccagctg cattaatgaa tcggccaacg cgcggggaga ggcggttgc gtattggcg	4320
ctcttccgct tcctcgtca ctgactcgtc gcgctcggc gttcggctgc ggcgagcgt	4380
atcagctcac tcaaaggcgg taatacgggt atccacagaa tcaggggata acgcaggaaa	4440
gaacatgtga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggcgg cgttgctggc	4500
gtttttccat aggtccgccc cccctgacga gcatcacaaa aatcgacgct caagtccagag	4560
gtggcgaaac ccgacaggac tataaagata ccaggcgttt cccctggaa gctccctcgt	4620
gcgctctcct gttccgacc tcgcccctac cggatacctg tccgcttcc tccctcggg	4680
aagcgtggcg ctttctcaat gctcacgctg taggtatctc agttcgggtg aggtcgttcg	4740
ctccaagctg ggtgtgtgc acgaaccccc cgttcagccc gaccgctgcg ccttatccgg	4800
taactatcgt cttgagtcca acccggttaag acacgactta tcgcccactgg cagcagccac	4860
tggtaacagg attagcagag cgaggtatgt aggcgggtgct acagagttct tgaagtgtg	4920
gcctaactac ggctacacta gaaggcagct atttggtatc tgcgctctgc tgaagccagt	4980
taccttcgga aaaagagttg gtagctcttg atccggcaaa caaaccaccg ctggtagcgg	5040
tggttttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc aagaagatcc	5100
tttgatcttt tctacggggt ctgacgctca gtggaacgaa aactcacgct aagggtttt	5160
ggctcatgaga ttatcaaaaa ggatcttcac ctagatcctt ttaaatataa aatgaagttt	5220
taaatcaatc taaagtatat atgagtaaac ttggtctgac agttaccaat gcttaatcag	5280
tgaggcacct atctcagcga tctgtctatt tcgttcatcc atagttgcct gactccccgt	5340
cgtgtagata actacgatac gggagggctt accatctggc cccagtgctg caatgatacc	5400
gcgagacca cgctcaccgg ctccagattt atcagcaata aaccagccag ccggaagggc	5460
cgagcgcaga agtggctcctg caactttatc cgctccatc cagtctatta attggtgccc	5520
ggaagctaga gtaagttagt cgccagtaa tagtttgccc aacgttggtg ccattgctac	5580
aggcatcgtg gtgtcacgct cgtcgtttgg tatggcttca ttcagctccg gttcccaacg	5640
atcaaggcga gttacatgat cccccatggt gtgcaaaaaa gcggttagct ccttcggctc	5700
tccgatcgtt gtcagaagta agttggccgc agtggtatca ctcatggtta tggcagcact	5760

-continued

---

```

gcataattct cttactgtca tgccatccgt aagatgcttt tctgtgactg gtgagtactc 5820
aaccaagtca ttctgagaat agtgtatgcg gcgaccgagt tgctcttgcc cggcgtcaat 5880
acgggataat accgcgccac atagcagaac tttaaaagtg ctcacatttg gaaaacgttc 5940
ttcggggcga aaactctcaa ggatcttacc gctggtgaga tccagttcga tgtaaccac 6000
tcgtgcaccc aactgatctt cagcatcttt tactttcacc agcgtttctg ggtgagcaaa 6060
aacaggaagg caaaatgccg caaaaaaggg aataaggcgc acacggaaat gttgaatact 6120
catactcttc ctttttcaat attattgaag catttatcag ggttattgtc tcatgagcgg 6180
atacatatth gaatgtatth agaaaaataa acaaataggg gttccgcgca catttccccg 6240
aaaagtgccca cctgacgtc 6259

```

&lt;210&gt; SEQ ID NO 56

&lt;211&gt; LENGTH: 245

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: GFP amino acid sequence.

&lt;400&gt; SEQUENCE: 56

```

Ala Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val
1           5           10           15
Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu
20          25          30
Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys
35          40          45
Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu
50          55          60
Cys Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Arg
65          70          75          80
His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg
85          90          95
Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val
100         105         110
Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile
115         120         125
Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn
130         135         140
Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly
145         150         155         160
Ile Lys Val Asn Phe Lys Thr Arg His Asn Ile Glu Asp Gly Ser Val
165         170         175
Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro
180         185         190
Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser
195         200         205
Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val
210         215         220
Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Asn Ile Asp
225         230         235         240
Gly Gly Gly Gly Gly
245

```

-continued

---

<210> SEQ ID NO 57  
<211> LENGTH: 4  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: G-spacer flexible spacer

<400> SEQUENCE: 57

Gly Gly Gly Gly  
1

<210> SEQ ID NO 58  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: G-spacer flexible spacer

<400> SEQUENCE: 58

Gly Gly Gly Gly Ser  
1 5

<210> SEQ ID NO 59  
<211> LENGTH: 4  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: A-spacer flexible spacer

<400> SEQUENCE: 59

Ala Ala Ala Ala  
1

<210> SEQ ID NO 60  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: A-spacer flexible spacer

<400> SEQUENCE: 60

Ala Ala Ala Ala Val  
1 5

<210> SEQ ID NO 61  
<211> LENGTH: 3359  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

gtcgaggtgc tcatagtgga gccctggctc ccgggaggac ggagccgcac ggtagtagat 60  
gggggtgtgg ccgtggcccc cgactctgct cggcggggcc gttctgtctt tgccatccgt 120  
gtgggacttc caccacagtg gaggaacgag agctggggccc catatgctgc ttgccagct 180  
tgggaaagag gaggtctgtg caaaggaccg atcggcggct tggggtgcc ggctcactcg 240  
gctgctgctg ctggtctggc gtctgtgag aagatcctct tctaccctgc tctgcacctg 300  
tgctcgaactg ccagccggct gagggcgggg gtctccacgg tggtoaccag tcccaaggag 360  
gttgacagaag taocgtacag agtggatttg cagggcagtg goatggagcc cctcttcccc 420  
cgcctgttct gggaggttat ctacggcagc caccttcagg gcaacctgtc cctcctgagc 480

-continued

---

cccaaccaca gctgctgccc cccgcactctg ctgctcaatg ccagccacgg cgccttctctg	540
cccctcgggc tcaaggteac catcggtggg ctctacctgg ccgtgtgtgt cggagggtctc	600
ctggggaact gccttgatc gtacgtcacc ctcaggcaca ccaaatgaa gacagccacc	660
aatatttaca tctttaacct ggccttggcc gacactctgg tctgctgac gctgcccttc	720
cagggcacgg acatcctctc gggcttctgg ccggttggga atgcgctgtg caagacagtc	780
attgccattg actactacaa catgttcacc agcaccttca cctaactgc catgagtgtg	840
gatcgctatg tagccatctg ccaccccatc cgtgccctcg acgtccgcac gtccagcaaa	900
gcccaggctg tcaatgtggc catctggggc ctggcctctg ttgtcgggtg tcccgttggc	960
atcatgggct cggcacaggt cgaggatgaa gagatcgagt gcctggtgga gatccctacc	1020
cctcaggatt actggggccc ggtgtttgac atctgcatct tcccttctc cttcatcgtc	1080
cccgtgctcg tcaatctctg ctgctacagc ctcatgatcc ggcggctccg tggagtccgc	1140
ctgctctcgg gctcccagaa gaaggaccgg aaacctcggc gcatcactcg gctggtgctg	1200
gtgtagtggt ctgtgttctg gggctgctgg acgcctgtcc aggtcttctg gctggcccaa	1260
gggctggggg ttcagccgag cagcagagct gccgtggcca ttctgcgctt ctgcacggcc	1320
ctgggctacg tcaacagctg cctcaacccc atcctctacg ccttctctga tgagaacttc	1380
aaggcctgct tccgcaagtt ctgctgtgca tctgccctgc gccgggacgt gcagggtgtct	1440
gaccgcgctg gcagcattgc caaggacgtg gccctggcct gcaagacctc tgagacggta	1500
ccgcggcccg catgactagg cgtggacctg cccatggtgc ctgtcagccc gcagagccca	1560
tctacgccc aacacagagct cacacaggtc actgctctct aggcggacac acctggggc	1620
ctgagcatcc agagcctggg atgggctttt cctgtggggc cagggatgct cgggtcccaga	1680
ggaggaccta gtgacatcat gggacaggtc aaagcattag ggccacctcc atggccccag	1740
acagactaaa gctgccctcc tgggtgcaggg ccgaggggac acaaggacct acctggaagc	1800
agctgacatg ctgggtggac gccgtgactg gagcccgctc cctccctcc cctgcttca	1860
tgtgactctt ggcctctctg ctgctgctgt ggcagaaacc tgggtgggca ggcaccggga	1920
ggaggagcag cagctgtgtc atcctgtgcc ccccatgtgc tgtgtgctgt ttgcatggca	1980
gggctccagc tgccctcagc cctgtgacgt ctctcaggg cagctggaca ggcttggcac	2040
tgcccgggaa gtgcagcagg cagcttttct ttggggggg acttgccctg agcttggagc	2100
tgccacctgg aggacttgcc tggctcagct ccacctgtgc agccggggcc acccaggag	2160
aaagtgtcca ggtgggggct ggcagtcctc ggctgcagac cccgagctgg cctggggcca	2220
gccgcacctc tgaaggtttt ctgtgtgctg cacgggtgcag gctcctccc tgactgcagc	2280
ttgactctgg gcccacccc catttccctt caggagacca gcgagaggcc ctggcccatt	2340
ccctccagcg gtgcaatgaa ctatcatgct gtggaccgtc aaccagccc tgcttctcag	2400
tgtggggcag gtgtctcagg acgaaggcgc ccgctgacca catgggcagc tctgttcaca	2460
aagtggaggc ctgcttttcc tggctctgac tgctctgttt ggggtgggaga agattctctg	2520
ggggtcccca catcctccca aggtccctc cacagcctct ccttctgttg aagccagagg	2580
tcagtggcgg tctgtgtgtg cgggggaagc tgtgtggaag gagaagctgg tggccacagc	2640
agagtcctgc tctggggacg cctgcttcat ttacaagcct caagatggct ctgtgtaggg	2700
cctgagcttg ctgcccacag ggaggatggc ttcacagcag agccagcatg aggggtgggg	2760

-continued

---

```

cctggcaggg cttgcttgag ccaaactgca aaggctgtgg tggctgtgag gacactgcgg 2820
gggttggggg gggggcgtct gtacctcagg ggatgccccg ctgtggtcac ccagagaatc 2880
acccttctct gtctacagat ggaagctgca ggttggtgac tttgcaaatg cacttctctac 2940
agatgaacta ttaaaagacc tgcaacattg aaaaaactca ttttttccac caaaccttg 3000
gccaggtaac ctaccttagg cacctgcaaa gaacaggaag tgatggctgt ctgcacaacag 3060
agcctgggct gctcctcctg ctctggggag tctagccctg ggggactggt ctggggaggc 3120
tcatgtctgc tccatgacgt ctgtggcagg agtccctgag gacgggagct gcctagctac 3180
agttttcttg ccaaggcagag gtgttttgag aatctgtgct gatgtaatgt gcaccttcac 3240
gtatttatgc atgtggcaag cgttacttcc tgtgcacgta gccagccctg ggtctgtctc 3300
tggggtaatg aaaaaggacc ctaataaaca cctgctcact ggctgggtat tcttcgtaa 3359

```

```

<210> SEQ ID NO 62
<211> LENGTH: 1134
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 62

```

```

ttgcagggca gtggcatgga gccctcttc cccgcgccgt tctgggaggt tatctacggc 60
agccaccttc agggcaacct gtccctcctg agccccaacc acagtctgct gcccccgcat 120
ctgctgtctc atgccagcca cggcgccttc ctgcccctcg ggctcaaggt caccatcgtg 180
gggctctacc tggcctgtgt gtgtggaggg ctctggggga actgcctgt catgcacacc 240
aaaatgaaga cagccaccaa tattacatc ttaacctgg cctggcoga cactctggtc 300
ctgctgacgc tgcccttcca gggcacggac atcctcctgg gcttctggcc gtttgggaat 360
gcgctgtgca agacagtcac tgccattgac tactacaaca tgttaccag caccttacc 420
ctaactgcca tgagtgtgga tcgctatgta gccatctgcc accccatccg tgccctcgac 480
gtccgcacgt ccagcaaacg ccaggctgtc aatgtggcca tctgggccct ggctctgtt 540
gtcgggttcc cgttgccat catgggctcg gcacaggctg aggatgaaga gatcgagtgc 600
ctggtggaga tccctacccc tcaggattac tggggcccgg tgtttgcat ctgcatcttc 660
ctcttctcct tcatcgtccc cgtgctcgtc atctctgtct gctacagcct catgatccgg 720
cggctccgtg gactccgctt gctctcgggc tcccagaga aggaccgaa cctgcggcgc 780
atcactcggc tgggtctggt ggtagtggct gtgttcgtgg gctgctggac gcctgtccag 840
gtcttctgtc tggcccaagg gctggggggt cagccgagca gcgagactgc cgtggccatt 900
ctgcgcttct gcacggcctt gggctacgtc aacagctgcc tcaacccat cctctacgcc 960
ttctggatg agaacttcaa ggctgcttc cgcaagttct gctgtgcatc tgccctgcgc 1020
cgggacgtgc aggtgtctga ccgcgtgcgc agcattgcca aggacgtggc cctggcctgc 1080
aagacctctg agacggtacc gcggcccgca tgactaggcg tggacctgcc catg 1134

```

```

<210> SEQ ID NO 63
<211> LENGTH: 1774
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 63

```

```

ccgaggagcc tgcgctgct cctggctcac agcgtcccg gcgaggagag cgggcggacg 60

```

-continued

---

```

ccgsgggctg ggcgggtgcg ggcggcgagg caggcggagc aggcgcagag acagcggggc 120
ggcggggggcg cggcagccgg cggcgctcggg gccggggcct ctgccttgcc gctcccctcg 180
cgtcggatcc ccgcgcccag ggcgcacggg ggagagggac gcggcggagc cggccggcag 240
ccatggaacc gggccccctc gccggcgccg agctgcagcc cccgctcttc gccaacgcct 300
cggacgccta ccctagcgcg tgccccagcg ctggcgccaa tgcgtcgggg ccgccaggcg 360
cgcggagcgc ctgcctccct gccctggcaa tcgccatcac cgcgctctac tcggccgtgt 420
gcgccgtggg gctgctgggc aacgtgcttg tcatgttcgg catcgtccgg tacactaaga 480
tgaagacggc caccaacatc tacatcttca acctggcctt agccgatgcg ctggccacca 540
gcacgctgcc tttccagagt gccaaagtacc tgatggagac gtggcccttc ggcgagctgc 600
tctgcaaggc tgtgctctcc atcgactact acaatatgtt caccagcadc ttcacgctca 660
ccatgatgag tgttgaccgc tacatcgcgt tctgccacco tgtaaggcc ctggacttcc 720
gcacgcctgc caaggccaag ctgatcaaca tctgtatctg ggtcctggcc tcaggcgttg 780
gctgcccctc catggtcatg gctgtgaccc gtcccgggga cggggcagtg gttgcatgc 840
tccagttccc cagccccagc tggtaactggg acacggtgac caagatctgc gtgttctct 900
tcgcttctgt ggtgcccctc ctcatcatca ccgtgtgcta tggcctcatg ctgctgcgcc 960
tgccagtggt gcgcctgctg tcgggctcca aggagaagga ccgcagcctg cggcgcatca 1020
cgcgcatggt gctggtggtt ttggggcctc tcgtggtgtg ttggggcgcc atccacatct 1080
tcgtcatcgt ctggacgctg ttggacatcg accggcgcgca cccgctggtg gttgctgcgc 1140
tgcacctggt catcgcgctg ggctacgcca atagcagcct caaccccgtg ctctacgctt 1200
tcctcgacga gaacttcaag cgctgcttcc gccagctctg ccgcaagccc tgcggccgcc 1260
cagaccccag cagcttcagc cgcgcccggg aagccacggc ccgcgagcgt gtcaccgct 1320
gcaccccgtc cgtggttccc ggcgggtggcg ctgccgctcg accaggccat ccggccccca 1380
gagcggccct ccctagtgc cggaggcca catgagtccc agtgggaggc gcgagccatg 1440
atgtggagtg gggcagtaga aggtcggagg cttgggaccg ccagatgggg cctctgtttc 1500
ggagacggga ccgggcccgt agatgggcat ggggtgggccc tctggtttgg ggcgaggcag 1560
aggacagatc aatggcgcag tgcctctggt ctgggtgccc cgtccacggc tctagtgagg 1620
gcgggaaagc cagtactcc aggagaggag cgggacctgt ggctataca ctgagtcctt 1680
aaacagggca tctccaggaa ggcggggcct caaccttgag acagcttcgg tttctaactt 1740
ggagccggac tttcggagtt gggggctcgg gcc 1774

```

&lt;210&gt; SEQ ID NO 64

&lt;211&gt; LENGTH: 1773

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 64

```

ccgaggagcc tgcgctgctc ctggctcaca gcgctccggg cgaggagagc gggcggaccg 60
gggggctggg ccggtgcggg cggcgaggca ggcggacgag gcgcagagac agcggggcg 120
ccggggcgcg gcacggcgcg ggtcggggcc gccctctgcc ttgcccctcc cctcgcgctg 180
gatccccgcg cccaggcagc cggtgagagag ggacgcggcg gacgcccgca gccatggaac 240
cggccccctc cgcggcgccg gagctgcagc ccccgtctt cggcaacgcc tcggacgct 300

```

-continued

---

```

accctagcgc cttccccagc gctggcgcca atgctcggg gccgccaggc gcgcgagcg 360
cctcgteect cgccctggca atcgccatca ccgctctta ctggcgctg tgcgctgg 420
ggctgctggg caacgtgctt gtcattgctg gcatcgctcg gtacactaag atgaagacgg 480
ccaccaacat ctacatcttc aacctggcct tagccgatgc gctggccacc agcacgctgc 540
ctttccagag tgccaagtac ctgatggaga cgtggccctt cggcgagctg ctctgcaagg 600
ctgtgctctc catcgactac tacaatatgt tcaccagcat cttcacgctc accatgatga 660
gtgttgaccg ctacatcgct gtctgccacc ctgtcaaggc cctggacttc cgcacgcctg 720
ccaaggccaa gctgatcaac atctgtatct gggctctggc ctgagcgctt ggcgtgccca 780
tcatggctcat ggtgtgacc cgtccccggg acggggcagt ggtgtgcatg ctccagtcc 840
ccagccccag ctggactgg gacacgggta ccaagatctg cgtgttctc ttcgccttcg 900
tgggtcccat cctcatcctc accgtgtgct atggcctcat gctgctgccc ctgcccagtg 960
tgccctgct gtcgggctcc aaggagaagg accgcagcct gcggcgctc acgcgcatgg 1020
tgctgggtgt tgtggggcgc ttcgtggtgt gttggggccc catccacatc ttcgtcatcg 1080
tctggacgct ggtggacatc gaccggcgcg acccgctggt ggtggctgcg ctgcacctgt 1140
gcatcgctcg gggctacgcc aatagcagcc tcaacccctg gctctacgct ttcctcgacg 1200
agaactcaa gcgctgcttc cgccagctct gccgcaagcc ctgcccggc ccagacccca 1260
gcagcttcag ccgccccgc gaagccaagg ccccgagcgc tgtaaccgcc tgcacccct 1320
ccgatggtcc cggcgggtgc gctgcccct gaccaggcca tccggcccc agaccccct 1380
ccctagtgtt acccgaggc cacatgagtc ccagtgggag gcgagccca tgatgtggag 1440
tggggccagt agataggtcg gaggctttg ggaccgccag atggggcctc tgtttcggag 1500
acgggaccgg gccgctagat gggcatgggg tggcctctg gtttggggcg aggcagagga 1560
cagatcaatg gcgagtgcc tctggtctgg gtgccccct ccaaggctct aggtggggcg 1620
ggaaaaccag tgactccagg agaggagcgg gacctgtggc tctacaactg agtccttaaa 1680
cagggcatct ccaggaaggc ggggcttcaa ccttgagaca gcttcggttt ctaacttga 1740
gccggacttt cggagtggg gggccgggg ccc 1773

```

&lt;210&gt; SEQ ID NO 65

&lt;211&gt; LENGTH: 1154

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 65

```

atggactccc cgatccagat cttccgggg gagccgggcc ctacctgccc ccgagcggc 60
tgctgcccc ccaacagcag cgctggttt cccggctggg ccgagcccga cagcaacggc 120
agcgcgggct cggaggagcc gcagctggag cccgcgcaca tctccccgc catcccgtc 180
atcatcacgg cgtctactc cgtagtgttc gtcgtgggct tgggggcaa ctgctggtc 240
atgttcgtga tcattcgata cacaagatg aagacagcaa ccaacattta catattaac 300
ctggctttgg cagatgcttt agttactaca accatgcctt ttcagagtac ggtctactg 360
atgaattcct ggcttttgg ggatgtgctg tgcaagatag taatttccat tgattactac 420
aacatgttca ccagcatctt caccttgacc atgatgagcg tggaccgcta cattgcccgtg 480
tgccaccccg tgaaggcttt ggaactccgc acacccttga aggcacaagat catcaatctc 540

```

-continued

---

```

tgcactctggc tgcctgctgc atctgttggc atctctgcaa tagtccttgg aggcacccaaa 600
gtcaggggaag acgtcgatgt cattgagtgc tccttgcaagt tcccagatga tgactactcc 660
tgggtgggacc tcttcatgaa gatctgctgc ttcactcttg ccttcgtgat cctgtctctc 720
atcatcatcg tctgctacac cctgatgac ctgcgtctca agagcgtccg gctcctttct 780
ggctcccag agaaagatcg caacctgcgt aggatcacca gactggctct ggtggtggtg 840
gcagtctctcg tcgtctgctg gactccatt cacatattca tcctggtgga ggcctctggg 900
agcacctccc acagcacagc tgcctctctc agctattact tctgcatcgc cttaggctat 960
accaacagta gctgaatcc cattctctac gcctttcttg atgaaaactt caagcggtgt 1020
ttccgggact tctgctttcc actgaagatg aggatggagc ggcagagcac tagcagagtc 1080
cgaatacag ttcaggatcc tgcctacctg agggacatcg atgggatgaa taaaccagta 1140
tgactagtcg tgga 1154

```

```

<210> SEQ ID NO 66
<211> LENGTH: 1143
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 66

```

```

atggaatccc cgattcagat cttccgctgg gagccgggccc ctacctgcgc cccgagcgc 60
tgctgcccc ccaacagcag cgctctggtt cccgctggg cagagcccga cagcaacggc 120
agcgcctggc cggaggagc gcagctggag cccgcgcaca tctcccggc catcccggc 180
atcatcagg cggtctactc cgtagtgttc gtcgtgggct tgggtgggcaa ctgctggtc 240
atgtctgtga tcatccgata cacaaagatg aagacagcaa ccaacattta catatttaac 300
ctggctttgg cagatgcttt agttactaca accatgcctt ttcagagtac ggtctacttg 360
atgaattcct ggccttttgg ggatgtgctg tgcaagatag taatttccat tgattactac 420
aacatgttca ccagcatctt caccttgacc atgatgagcg tggaccgcta cattgccgtg 480
tgccacccc tgaaggcttt ggacttcgc acaccttga agcacaagat catcaatctc 540
tgcactctggc tgcctgctgc atctgttggc atctctgcaa tagtccttgg aggcacccaaa 600
gtcaggggaag acgtcgatgt cattgagtgc tccttgcaagt tcccagatga tgactactcc 660
tgggtgggacc tcttcatgaa gatctgctgc ttcactcttg ccttcgtgat cctgtctctc 720
atcatcatcg tctgctacac cctgatgac ctgcgtctca agagcgtccg gctcctttct 780
ggctcccag agaaagatcg caacctgcgt aggatcacca gactggctct ggtggtggtg 840
gcagtctctcg tcgtctgctg gactccatt cacatattca tcctggtgga ggcctctggg 900
agcacctccc acagcacagc tgcctctctc agctattact tctgcatcgc cttaggctat 960
accaacagta gctgaatcc cattctctac gcctttcttg atgaaaactt caagcggtgt 1020
ttccgggact tctgctttcc actgaagatg aggatggagc ggcagagcac tagcagagtc 1080
cgaatacag ttcaggatcc tgcctacctg agggacatcg atgggatgaa taaaccagta 1140
tga 1143

```

```

<210> SEQ ID NO 67
<211> LENGTH: 2738
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

-continued

&lt;400&gt; SEQUENCE: 67

---

cctgtaaaga aatgatgag ggctaaatcc atcagcacca aagetgggaa gcctccagg	60
ttcatttga agaaaaact cctctgagct caaaggaagt gtgatctgtc acaatattgt	120
atgcctgcac taagtttgca tcctgaaaac tcaactggaag ataggaaagc aagcatgaaa	180
aagcagccgg gtcagacagg cttctggatt cagtgtgtgg acatgacttt gcctgcatga	240
attgcccctt tctctacaa acaagagaat tcggtcaagt ggatgtggca gaactgggct	300
gctctgagat gatagaaaa ggctcctgct tttcctgtaa ttgcagcccc ttgttcttgt	360
ggttgctaca tgcaataaat gtaattctat gagaaggacc agcccttaca tcccatcaaa	420
atgtttctg gaaacctgga gcacagaact ctgatatcct ctcacactgt ggcaggagaa	480
gcagcacaag gcacaatgct gaaatagcat ggtccaggat gtgtttgcac agaagagtgc	540
ccagtgaaga gacctactcc ttggatcgct ttgcgcaaaa tccacccctt ttcctcctc	600
cctcccttcc agcctccgaa tcccgcattg cccacgctcc cctcctgcag cggtgccggg	660
cagccaggac tggtttctgt aagaacacagc aggagctgtg gcagcggcga aaggaagcgg	720
ctgaggcget tggaaacctg aaagtctcgg tgcctctggc tacctcgcac agcggtgccc	780
gcccggccgt cagtaccatg gacagcagcg ctgccccac gaacgccagc aattgcaactg	840
atgccttggc gtactcaagt tgcctcccag caccagccc cggttcctgg gtcaacttgt	900
cccacttaga tggcaacctg tccgacctat gcggtccgaa ccgcaccgac ctgggcccga	960
gagacagcct gtgcctccg accggcagtc cctccatgat cacggccatc acgatcatgg	1020
ccctctactc catcgtgtgc gtggtggggc tcttcggaaa cttcctggtc atgtatgtga	1080
ttgtcagata caccaagatg aagactgcca ccaacatcta cattttcaac cttgctctgg	1140
cagatgcctt agccaccagt accctgcctt tccagagtgt gaattaccta atgggaacat	1200
ggccatttgg aaccatcctt tgcaagatag tgatctccat agattactat aacatgttca	1260
ccagcatatt caccctctgc accatgagtg ttgatcgata cattgcagtc tgccaccctg	1320
tcaaggcctt agatttccgt actccccgaa atgccaaaat tatcaatgtc tgcaactgga	1380
tcctctcttc agccatttgt cttcctgtaa tgttcatggc tacaacaaaa tacaggcaag	1440
gttccataga ttgtacacta acattctctc atccaacctg gtactgggaa aacctgctga	1500
agatctgtgt tttcatcttc gccttcatta tgccagtgct catcattacc gtgtgctatg	1560
gactgatgat cttgcgcctc aagagtgctc gcatgctctc tggctccaaa gaaaaggaca	1620
ggaatcttctg aaggatcacc aggatggtgc tgggtggtgt ggctgtgttc atcgtctgct	1680
ggactcccat tcacatttac gtcacatta aagccttggc tacaatccca gaaactacgt	1740
tccagactgt ttcttggcac ttctgcattg ctctaggtta cacaaacagc tgcctcaacc	1800
cagtcttcta tgcatttctg gatgaaaact tcaaacgatg cttcagagag ttctgtatcc	1860
caacctcttc caacattgag caacaaaact ccaactcgaat tgcgcagaac actagagacc	1920
accctccac ggccaataca gtggatagaa ctaatcatca gctagaaaat ctggaagcag	1980
aaactgctcc gttgcctaa cagggtctca tgccattccg acctcacca agcttagaag	2040
ccaccatgta tgtggaagca ggttgcctca agaattgtga ggaggctcta attctctagg	2100
aaagtgcctg cttttaggtc atccaacctc tttcctctct ggccactctg ctctgcacat	2160
tagagggaca gccaaaagta agtggagcat ttggaaggaa aggaatatac cacaccgagg	2220

-continued

---

agtccagttt gtgcaagaca cccagtgga ccaaaaccca tcgtggatg tgaattgaag	2280
tcatcataaa aggtgacct tctgtctgta agattttatt ttcaagcaaa tatttatgac	2340
ctcaacaaag aagaaccatc ttttgtaag ttcaccgtag taacacataa agtaaatgct	2400
acctctgac aaagcacctt gaatggaagg tccgagtctt tttagtgtt tgcaagggaa	2460
tgaatccatt attctatttt agacttttaa cttcacctta aaattagcat ctggctaagg	2520
catcattttc acctccattt cttggtttg tattgtttaa aaaaataaca tctctttcat	2580
ctagctccat aattgcaagg gaagagatta gcatgaaagg taatctgaaa cacagtcag	2640
tgtagctgt agaaaggtg attctcatgc actgcaata cttccaaaga gtcacatgg	2700
gggatttttc attcttaggc tttcagtggt ttgttct	2738

&lt;210&gt; SEQ ID NO 68

&lt;211&gt; LENGTH: 2053

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 68

catctttgat gagggcagag ctcacgttgc attgaagacg aaacctcggg gaggtcaggc	60
gctgtctttc cttccctccc tgctcggcgg ctcaccaca gttgcaacct gcagaggccc	120
ggagaacaca accctcccga gaagcccagg tccagagcca aacctcgcac tgacccccca	180
gccaggcgc ccagccactc cccaccgcta ccatggccga agacgcagac atgacgaatg	240
agctggagga gatgcagca agggctgacc agttggctga tgagtcgctg gaaagcaccc	300
gtcgtatgct gcaactggtt gaagagagta aagatgctgg taccaggact ttggttatgt	360
tggatgaaca aggagaacaa ctcgatcgtg tcgaagaagg catgaacct atcaaccaag	420
acatgaagga ggctgagaaa aatttaaaag atttagggaa atgctgtggc cttttcatat	480
gtcctttaa caagcttaa tcaagtgat cttacaaaa agcctggggc aataatcagg	540
acggagtggg gccagccag cctgctcgtg tagtgagca acgggagcag atggccatca	600
gtggcggctt catccgcagg gtaacaaatg atgccgaga aaatgaaatg gatgaaaacc	660
tagagcaggt gagcggcatc atcgggaacc tccgtcacat ggcctggat atgggcaatg	720
agatcgatac acagaatcgc cagatcgaca ggatcatgga gaagctgat tccaacaaaa	780
ccagaattga tgaggccaac caactgcaa caaagatgct gggaaagtgt taagtgtgcc	840
caccctggtt ctcctccaaa tgctgctcgg caagatagct cttcatgct tttctcatgg	900
tattatctag taggtctgca cacataacac acatcagtc acccccattg tgaatgttgt	960
cctgtgtcat ctgtcagctt cccaacaata ctttgtgtct tttgttctct cttggtctct	1020
ttctttccaa aggtgtgaca tagtggctcat ttggtggctc taactccttg atgtcttgag	1080
tttcattttt cattttctct cctcgtggc atttgctgaa taacaacaat ttaggaatgc	1140
tcaatgtgct gttgattctt tcaatccaca gtattgttct tgtaaaactg tgacattcca	1200
cagagttact gccacggtcc tttgagtgc aggctctgaa tctctcaaaa tgtgccgtct	1260
ttggttcctc atggtgttta tctgtcttta tgatttcag attagacaat gtggaattac	1320
ataacaggca ttgcactaaa agtgatgtga tttatgcatt tatgatgag aactaaatag	1380
atTTTTAGAT tctacttaa acaaaaactt tccatgacag tagcatactg atgagacaac	1440
acacacacac acaaaaacac agcaacaaca acagaacaac aacaagcat gctcagtatt	1500

-continued

---

gagacactgt caagattaag ttataccagc aaaagtgcag tagtgtcact tttttcctgt	1560
caatatatag agacttctaa atcataatca tcctttttta aaaaaaagaa ttttaaaaa	1620
gatggatttg acacactcac catttaatca tttccagcaa aatatatggt tggctgaaat	1680
tatgtcaaat ggatgtaata tagggtttgt ttgctgcttt tgatggctac gttttggaga	1740
gagcaatcct gctgtgaaac agtgtggatg taaatttat aaggctgact cttactaac	1800
accatttccc ctgtggtttg ttatcagtac aattccttgt tgcttaatct agagctatgc	1860
acaccaaatt gctgagatgt ttagtagctg ataaagaaac cttttaaaaa aataatataa	1920
atgaatgaaa tataaactgt gagataata tcattatagc atgtaatatt aaattcctcc	1980
tgtctcctct gtcagtttgt gaagtgattg acattttgta gctagttaa aattattaaa	2040
aattatagac tcc	2053

&lt;210&gt; SEQ ID NO 69

&lt;211&gt; LENGTH: 2053

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 69

catctttgat gagggcagag ctcacgttgc attgaagacg aaacctcggg gaggtcagc	60
gctgtctttc cttccctccc tgctcggcgg ctcaccaca gttgcaacct gcagaggccc	120
ggagaacaca accctcccga gaagcccagg tccagagcca aaccgctcac tgacccccca	180
gcccaggcgc ccagccactc cccaccgcta ccatggccga agacgcagac atgcgcaatg	240
agctggagga gatgcagcga agggctgacc agttggctga tgagtcgctg gaaagcacc	300
gtcgtatgct gcaactggtt gaagagagta aagatgctgg taccaggact ttggttatgt	360
tggatgaaca aggagaacaa ctggaacgca ttgaggaagg gatggaccaa atcaataagg	420
acatgaaaga agcagaaaag aatttgacgg acctaggaag attctgcggg ctttgtgtgt	480
gtccctgtaa caagcttaaa tcaagtgatg cttacaaaa agcctggggc aataatcagg	540
acggagtggg ggcagcccag cctgctcgtg tagtggacga acgggagcag atggccatca	600
gtggcggcct catccgcagg gtaacaaatg atgcccgaga aatgaaatg gatgaaaacc	660
tagagcaggt gagcggcatc atcgggaacc tccgtcacat ggcctggat atgggcaatg	720
agatcgatc acagaatcgc cagatcgaca ggatcatgga gaaggctgat tccaacaaaa	780
ccagaattga tgaggccaac caacgtgcaa caaagatgct gggaaagtgt taagtgtgcc	840
caccctgtgt ctccctcaaa tgctgtcggg caagatagct ccttcattgt tttctcatgg	900
tattatctag taggtctgca cacataaac acatcagtc accccattg tgaatgttgt	960
cctgtgtcat ctgtcagctt cccaacaata ctttgtgtct tttgttctct cttggtctct	1020
ttctttccaa aggtgtgaca tagtggctcat ttggtggctc taactccttg atgtcttgag	1080
tttcattttt cattttctct cctcgggtggc atttgctgaa taacaacaat ttaggaatgc	1140
tcaatgtgct gttgattctt tcaatccaca gtattgttct tgtaaaactg tgacattcca	1200
cagagttact gccacggtcc tttgagtgtc aggctctgaa tctctcaaaa tgtgccgtct	1260
ttggttcctc atggetgtta tctgtcttta tgatttcatt attagacaat gtggaattac	1320
ataacaggca ttgcactaaa agtgatgtga tttatgcatt tatgcatgag aactaaatag	1380
atttttagat tctacttaa acaaaaactt tccatgacag tagcactatg atgagacaa	1440

-continued

---

```

acacacacac acaaaacaac agcaacaaca acagaacaac aacaaagcat gctcagatatt 1500
gagacactgt caagattaag ttataccagc aaaagtgcag tagtgctcact tttttcctgt 1560
caatatatag agacttctaa atcataatca tcctttttta aaaaaaagaa ttttaaaaaa 1620
gatggatttg acacactcac catttaatca tttccagcaa aatatatggt tggctgaaat 1680
tatgtcaaat ggatgtaata tagggtttgt ttgctgcttt tgatggctac gttttggaga 1740
gagcaatcct gctgtgaaac agtgtggatg taaattttat aaggctgact cttactaacc 1800
accatttccc ctgtggtttg ttatcagtac aattccttgt tgcttaactc agagctatgc 1860
acaccaaatt gctgagatgt ttagtagctg ataaagaaac cttttaaaaa aataatataa 1920
atgaatgaaa tataaactgt gagataaata tcattatagc atgtaatatt aaattcctcc 1980
tgtctcctct gtcagtttgt gaagtgattg acattttgta gctagtttaa aattattaaa 2040
aattatagac tcc 2053

```

```

<210> SEQ ID NO 70
<211> LENGTH: 266
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 70

```

```

Met Val Ser Trp Gly Arg Phe Ile Cys Leu Val Val Val Thr Met Ala
1 5 10 15
Thr Leu Ser Leu Ala Arg Pro Ser Phe Ser Leu Val Glu Asp Thr Thr
20 25 30
Leu Glu Pro Glu Glu Pro Pro Thr Lys Tyr Gln Ile Ser Gln Pro Glu
35 40 45
Val Tyr Val Ala Ala Pro Gly Glu Ser Leu Glu Val Arg Cys Leu Leu
50 55 60
Lys Asp Ala Ala Val Ile Ser Trp Thr Lys Asp Gly Val His Leu Gly
65 70 75 80
Pro Asn Asn Arg Thr Val Leu Ile Gly Glu Tyr Leu Gln Ile Lys Gly
85 90 95
Ala Thr Pro Arg Asp Ser Gly Leu Tyr Ala Cys Thr Ala Ser Arg Thr
100 105 110
Val Asp Ser Glu Thr Trp Tyr Phe Met Val Asn Val Thr Asp Ala Ile
115 120 125
Ser Ser Gly Asp Asp Glu Asp Asp Thr Asp Gly Ala Glu Asp Phe Val
130 135 140
Ser Glu Asn Ser Asn Asn Lys Arg Ala Pro Tyr Trp Thr Asn Thr Glu
145 150 155 160
Lys Met Glu Lys Arg Leu His Ala Val Pro Ala Ala Asn Thr Val Lys
165 170 175
Phe Arg Cys Pro Ala Gly Gly Asn Pro Met Pro Thr Met Arg Trp Leu
180 185 190
Lys Asn Gly Lys Glu Phe Lys Gln Glu His Arg Ile Gly Gly Tyr Lys
195 200 205
Val Arg Asn Gln His Trp Ser Leu Ile Met Glu Ser Val Val Pro Ser
210 215 220
Asp Lys Gly Asn Tyr Thr Cys Val Val Glu Asn Glu Tyr Gly Ser Ile
225 230 235 240
Asn His Thr Tyr His Leu Asp Val Val Gly Glu Ser Ala Ser Pro Arg

```

-continued

---

245	250	255	
Val Ala Ala Ala Tyr Gln Pro Ile Leu Ala			
260	265		

<210> SEQ ID NO 71  
 <211> LENGTH: 336  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 71

caggtgaagc tgcaggagtc tggcgctgag ttggtgaaac ctggggcttc agtgaagata	60
tcctgcaagg cttctggcta catcttcaact gaccatgctc ttcaactgggt gaggcagaag	120
cctgaacagg gcctggaatg gattgggtat atttttcccg gaaatggtaa tattgagtac	180
aatgagaagt tcaagggcaa ggccacactg actgcagaca aatcctccag tactgcctac	240
atgcagctca acagcctgac atctggagat tctgcaatgt atttctgtaa aaagatggac	300
tactggggcc aagggaccac ggtcaccgtc tcctca	336

<210> SEQ ID NO 72  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 72

Gln Val Lys Leu	Gln Glu Ser Gly	Ala Glu Leu Val	Lys Pro Gly Ala		
1	5	10	15		
Ser Val Lys Ile	Ser Cys Lys Ala	Ser Gly Tyr Ile	Phe Thr Asp His		
20	25	30			
Ala Leu His Trp	Val Arg Gln Lys	Pro Glu Gln Gly	Leu Glu Trp Ile		
35	40	45			
Gly Tyr Ile Phe	Pro Gly Asn Gly	Asn Ile Glu Tyr	Asn Glu Lys Phe		
50	55	60			
Lys Gly Lys Ala	Thr Leu Thr Ala	Asp Lys Ser Ser	Ser Thr Ala Tyr		
65	70	75	80		
Met Gln Leu Asn	Ser Leu Thr Ser	Gly Asp Ser Ala	Met Tyr Phe Cys		
85	90	95			
Lys Lys Met Asp	Tyr Trp Gly Gln	Gly Thr Thr Val	Thr Val Ser Ser		
100	105	110			

<210> SEQ ID NO 73  
 <211> LENGTH: 336  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 73

caggtgaagc tgcaggagtc tggcgctgag ttggtgaaac ctggggcttc agtgaagatc	60
tcctgcaagg cttctggcta caccttcaact gaccattcta ttcaactgggt gaagcagaag	120
cctggacagg gcctagaatg gattggatat ctttttcccg gaaatggtaa ttttgaatat	180
aatgagaaat tcaagggcaa ggccacactg actgcagaca aatcctccag cactgcctac	240
atgcacctca acagcctgac atctggagat tctgcagtgt atttctgtaa aaagatggac	300
tactggggcc aagggaccac ggtcaccgtc tcctca	336

<210> SEQ ID NO 74

-continued

---

```

<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 74
Gln Val Lys Leu Gln Glu Ser Gly Ala Glu Leu Val Lys Pro Gly Ala
1          5          10          15
Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp His
20        25        30
Ser Ile His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
35        40        45
Gly Tyr Leu Phe Pro Gly Asn Gly Asn Phe Glu Tyr Asn Glu Lys Phe
50        55        60
Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
65        70        75        80
Met His Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85        90        95
Lys Lys Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
100       105       110

```

```

<210> SEQ ID NO 75
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 75
caggttcagc tgcagcagtc cgacgctgag ttggtgaaac ctggggcttc agtgaagata    60
tctctgcaggg cttctgggcta caccttcaact gaccattcta ttcactgggt gaagcagcag    120
cctggccagg gcttgggaatg gatcgatgat atttttcccg gaaatggaaa tattgaatac    180
aatgacaaat tcaagggcaa ggccacactg actgcagaca aatcctccgg cactgcctac    240
atgcagctca acagcctgac atctgaggat tctgcagtgt atttctgtaa aaggatgggg    300
tactggggtc aaggaacctc agtcaccgct tcctca                                336

```

```

<210> SEQ ID NO 76
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 76
Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Ala
1          5          10          15
Ser Val Lys Ile Ser Cys Arg Ala Ser Gly Tyr Thr Phe Thr Asp His
20        25        30
Ser Ile His Trp Val Lys Gln Gln Pro Gly Gln Gly Leu Glu Trp Ile
35        40        45
Gly Tyr Ile Phe Pro Gly Asn Gly Asn Ile Glu Tyr Asn Asp Lys Phe
50        55        60
Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Gly Thr Ala Tyr
65        70        75        80
Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85        90        95
Lys Arg Met Gly Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser
100       105       110

```

-continued

---

```

<210> SEQ ID NO 77
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 77
cagggtcaagc tgcaggagtc tggcgctgag ttggtgaaac ctggggcttc agtgaagatc   60
tcctgcaagg cttctggcta caccttcaact gaccattcta ttcactgggt gaagcagaag   120
cctggacagg gcttagaatg gattggatat ctttttcccg gaaatggtaa ttttgagtac   180
aatgaaaaat tcaagggcaa ggccacactg actgcagaca aatcctccag cactgtctac   240
atgtacctca acagcctgac atctgaggat tctgcagtgt atttctgtaa aaggatgggg   300
tactggggcc aagggaccac ggtcaccgtc tctca                                     336

```

```

<210> SEQ ID NO 78
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 78
Gln Val Lys Leu Gln Glu Ser Gly Ala Glu Leu Val Lys Pro Gly Ala
1          5          10          15
Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp His
20        25        30
Ser Ile His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
35        40        45
Gly Tyr Leu Phe Pro Gly Asn Gly Asn Phe Glu Tyr Asn Glu Lys Phe
50        55        60
Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Val Tyr
65        70        75        80
Met Tyr Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85        90        95
Lys Arg Met Gly Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
100       105       110

```

```

<210> SEQ ID NO 79
<211> LENGTH: 336
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 79
cagggtcaagc tgcaggagtc tggacctgaa ctggtaaagc ctggggcttc agtgaagatg   60
tcctgcaagg cttctggata cacattcaact aactatgta tacactgggt gaagcaaaag   120
cctgggcagg gcttgtagtg gattggatat attaatcctt acaatgatgg ctctaagtac   180
aatgagaagt tcaaaggcaa ggcctcactg acttcagaca aatcctccag cacagcctac   240
atggagctca gcagcctgac ctctgaggac tctgcggtct attactgtgc aagaatggac   300
tactggggcc aagggaccac ggtcaccgtc tctca                                     336

```

```

<210> SEQ ID NO 80
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 80

```

-continued

---

Gln Val Gln Leu Gln Glu Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
 1 5 10 15  
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr  
 20 25 30  
 Val Ile His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile  
 35 40 45  
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Ser Lys Tyr Asn Glu Lys Phe  
 50 55 60  
 Lys Gly Lys Ala Ser Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr  
 65 70 75 80  
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
 85 90 95  
 Ala Arg Met Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
 100 105 110

<210> SEQ ID NO 81  
 <211> LENGTH: 336  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 81

cagggtcaagc tgcaggagtc tggacctgaa ctggtaaagc ctggggcttc agtgaagatg 60  
 tcctgcaagg cttctggata cacattcact aactatgta taaactgggt gaagcaaaag 120  
 cctgggcagg gccttgagtg gattggatat attaatcctt acaatgatgg ctctaagtac 180  
 aatgagaagt tcaaaaggcaa ggctcactg acttcagaca aatcctccag cacagcctac 240  
 atggagctca gcagcctgac ctctgaggac tctgcggtct attactgtgc aagaatgggg 300  
 tactggggcc aagggaccac ggctaccgtc tcctca 336

<210> SEQ ID NO 82  
 <211> LENGTH: 112  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 82

Gln Val Gln Leu Gln Glu Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
 1 5 10 15  
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr  
 20 25 30  
 Val Ile His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile  
 35 40 45  
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Ser Lys Tyr Asn Glu Lys Phe  
 50 55 60  
 Lys Gly Lys Ala Ser Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr  
 65 70 75 80  
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
 85 90 95  
 Ala Arg Met Gly Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
 100 105 110

<210> SEQ ID NO 83  
 <211> LENGTH: 342  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

-continued

&lt;400&gt; SEQUENCE: 83

```

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttgaga tcaagcctcc    60
atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg    120
tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt    180
tctgggggcc cagacaggtt cagtggcagt ggatcagggc cagatttcac actcaagatc    240
agcagagtgg aggtcgagga tctggggagt tattactgct ttcaaggttc acatgttctc    300
cctacgttcg gtgctgggac caagctggag ctgaaacggg ct                          342

```

&lt;210&gt; SEQ ID NO 84

&lt;211&gt; LENGTH: 113

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 84

```

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1           5           10          15
Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
20          25          30
Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35          40          45
Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50          55          60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65          70          75          80
Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly
85          90          95
Ser His Val Pro Pro Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
100         105         110

```

Arg

&lt;210&gt; SEQ ID NO 85

&lt;211&gt; LENGTH: 324

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 85

```

gacatccaga tgactcagtc tccagcctcc ctatctgcat ctgtgggaga aactgtcacc    60
atcacatgtc gaacaactga aaatatttac agttattttg tatggtctca gcagagacag    120
ggaaaatctc ctacagctccg ggtctataat gcaaaatcct tagcagaagg tgtgccatca    180
agtttcaatg tcagtgtatc aggcacacag ttttctctga agatcaatag cctgcagcct    240
gaagattttg ggacttatca ctgtcaaac cattatggta ctccgtacac gttcggaggg    300
gggaccaggc tggaaataag acgg                          324

```

&lt;210&gt; SEQ ID NO 86

&lt;211&gt; LENGTH: 108

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 86

```

Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
1           5           10          15

```

-continued

Glu Thr Val Thr Ile Thr Cys Arg Thr Thr Glu Asn Ile Tyr Ser Tyr  
 20 25 30  
 Phe Val Trp Ser Gln Gln Arg Gln Gly Lys Ser Pro Gln Leu Arg Val  
 35 40 45  
 Tyr Asn Ala Lys Ser Leu Ala Glu Gly Val Pro Ser Ser Phe Asn Val  
 50 55 60  
 Ser Val Ser Gly Thr Gln Phe Ser Leu Lys Ile Asn Ser Leu Gln Pro  
 65 70 75 80  
 Glu Asp Phe Gly Thr Tyr His Cys Gln His His Tyr Gly Thr Pro Tyr  
 85 90 95  
 Thr Phe Gly Gly Gly Thr Arg Leu Glu Ile Arg Arg  
 100 105

<210> SEQ ID NO 87  
 <211> LENGTH: 339  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 87

gatgttttg atgacccaaac tccactcact ttgtcgggta ccattggaca accagcttcc 60  
 atctcttgc aagtccagtca gagcctctta tatactaata gaaaaaccta tttgacttgg 120  
 ttattccag aggccaggcca gtctocaaaa cgctaatct atctggtgtc tgaattggac 180  
 tctggagtc cctgacaggtt cagtggcagt ggttcagga cagatttcac actggaatc 240  
 accagagtg gaggctgagga tttgggagtt tattactgct tgcagagtgc acattttcca 300  
 ttcacgttc ggctcgggcac caagctggaa atcaaacgg 339

<210> SEQ ID NO 88  
 <211> LENGTH: 113  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 88

Asp Val Leu Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly  
 1 5 10 15  
 Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Thr  
 20 25 30  
 Asn Gly Lys Thr Tyr Leu Thr Trp Leu Phe Gln Arg Pro Gly Gln Ser  
 35 40 45  
 Pro Lys Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Gly Val Pro  
 50 55 60  
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Glu Ile  
 65 70 75 80  
 Thr Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Leu Gln Ser  
 85 90 95  
 Ala His Phe Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys  
 100 105 110

Arg

<210> SEQ ID NO 89  
 <211> LENGTH: 339  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 89

-continued

---

```

gatgttgga tgacccaac tccactcact ctgtcgggga ccattggaca accagcgttc    60
atctcttgca agtccagtca gagcctcttt aacactaatg gcaaaccta tttgacttgg    120
ttaattcaga ggcacaggcca gtctccacag cgectgatct atctgggtgc caaattggac    180
tctggcgctcc cggacaggtt cagtggcagt ggctcagga cagatttcac actgaaaatc    240
agcagagtgg aggtgagga tctgggagtt tattactgcc tgcagagtag ccattttccg    300
tttacgttcg gctcgggcac caagctggaa atcaaacgg    339

```

```

<210> SEQ ID NO 90
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 90

```

```

Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly
1          5          10          15
Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Thr
          20          25          30
Asn Gly Lys Thr Tyr Leu Thr Trp Leu Ile Gln Arg Pro Gly Gln Ser
          35          40          45
Pro Gln Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
          50          55          60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65          70          75          80
Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Leu Gln Ser
          85          90          95
Ser His Phe Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
          100          105          110

```

```

Arg

```

```

<210> SEQ ID NO 91
<211> LENGTH: 327
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 91

```

```

gatgttgatgc taactcagtc tcctgccacc ctgtctgtga ctccaggaga tagagtcagt    60
ctttcctgca gggccagcca aaatattggc aactacctac actggtatca acagaaatca    120
catgagtctc caaggtctct catcaagtat gcttcccagt ccatctctgg gatcccctec    180
aggttcagtg gcagtgatc agtcacagat ttcactctca atatcaacag tgtggagact    240
gaagattttg gaatgtattt ctgtcaacag agtgacacct ggcctctcac gttcgggtct    300
gggaccaagc tggagctgaa acgggct    327

```

```

<210> SEQ ID NO 92
<211> LENGTH: 108
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 92

```

```

Asp Val Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
1          5          10          15
Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Asn Ile Gly Asn Tyr

```

-continued

---

20	25	30
Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile		
35	40	45
Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly		
50	55	60
Ser Gly Ser Val Thr Asp Phe Thr Leu Asn Ile Asn Ser Val Glu Thr		
65	70	75
Glu Asp Phe Gly Met Tyr Phe Cys Gln Gln Ser Asp Thr Trp Pro Leu		
85	90	95
Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg		
100	105	

<210> SEQ ID NO 93  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 93

Thr Phe Thr Asp His Ser Ile His  
 1 5

<210> SEQ ID NO 94  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 94

Thr Phe Thr Asn Tyr Val Ile His  
 1 5

<210> SEQ ID NO 95  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 95

Ile Phe Thr Asp His Ala Leu His  
 1 5

<210> SEQ ID NO 96  
 <211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 96

Tyr Ile Phe Pro Gly Asn Gly Asn Ile Glu Tyr Asn Asp Lys Phe Lys  
 1 5 10 15

Gly

<210> SEQ ID NO 97  
 <211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 97

Tyr Leu Phe Pro Gly Asn Gly Asn Phe Glu Tyr Asn Glu Lys Phe Lys  
 1 5 10 15

Gly

-continued

---

<210> SEQ ID NO 98  
<211> LENGTH: 17  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 98

Tyr Ile Asn Pro Tyr Asn Asp Gly Ser Lys Tyr Asn Glu Lys Phe Lys  
1           5                   10                   15

Gly

<210> SEQ ID NO 99  
<211> LENGTH: 17  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 99

Tyr Ile Phe Pro Gly Asn Gly Asn Ile Glu Tyr Asn Glu Lys Phe Lys  
1           5                   10                   15

Gly

<210> SEQ ID NO 100  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 100

Lys Arg Met Gly Tyr  
1           5

<210> SEQ ID NO 101  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 101

Lys Lys Met Asp Tyr  
1           5

<210> SEQ ID NO 102  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 102

Ala Arg Met Asp Tyr  
1           5

<210> SEQ ID NO 103  
<211> LENGTH: 16  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 103

Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu  
1           5                   10                   15

<210> SEQ ID NO 104  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

-continued

---

<400> SEQUENCE: 104

Arg Thr Thr Glu Asn Ile Tyr Ser Tyr Phe Val  
1           5                   10

<210> SEQ ID NO 105

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 105

Lys Ser Ser Gln Ser Leu Leu Tyr Thr Asn Gly Lys Thr Tyr Leu Thr  
1           5                   10                   15

<210> SEQ ID NO 106

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 106

Lys Ser Ser Gln Ser Leu Leu Asn Thr Asn Gly Lys Thr Tyr Leu Thr  
1           5                   10                   15

<210> SEQ ID NO 107

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 107

Arg Ala Ser Gln Asn Ile Gly Asn Tyr Leu His  
1           5                   10

<210> SEQ ID NO 108

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 108

Lys Val Ser Asn Arg Phe Ser  
1           5

<210> SEQ ID NO 109

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 109

Asn Ala Lys Ser Leu Ala Glu  
1           5

<210> SEQ ID NO 110

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 110

Leu Val Ser Glu Leu Asp Ser  
1           5

<210> SEQ ID NO 111

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

-continued

---

<400> SEQUENCE: 111

Leu Val Ser Lys Leu Asp Ser  
1 5

<210> SEQ ID NO 112

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 112

Tyr Ala Ser Gln Ser Ile Ser  
1 5

<210> SEQ ID NO 113

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 113

Phe Gln Gly Ser His Val Pro Pro Thr  
1 5

<210> SEQ ID NO 114

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 114

Gln His His Tyr Gly Thr Pro Tyr Thr  
1 5

<210> SEQ ID NO 115

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 115

Leu Gln Ser Ala His Phe Pro Phe Thr  
1 5

<210> SEQ ID NO 116

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 116

Leu Gln Ser Ser His Phe Pro Phe Thr  
1 5

<210> SEQ ID NO 117

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 117

Gln Gln Ser Asp Thr Trp Pro Leu Thr  
1 5

<210> SEQ ID NO 118

<211> LENGTH: 5

<212> TYPE: PRT

---

-continued

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 118

Asp His Ser Ile His  
1 5

<210> SEQ ID NO 119

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 119

Asn Tyr Val Ile His  
1 5

<210> SEQ ID NO 120

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 120

Asp His Ala Leu His  
1 5

<210> SEQ ID NO 121

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 121

Ile Phe Pro Gly Asn Gly Asn Ile Glu  
1 5

<210> SEQ ID NO 122

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 122

Leu Phe Pro Gly Asn Gly Asn Phe Glu  
1 5

<210> SEQ ID NO 123

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 123

Ile Asn Pro Tyr Asn Asp Gly Ser Lys  
1 5

<210> SEQ ID NO 124

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 124

His Leu Ala Asn Thr Tyr Tyr Tyr Phe Asp Tyr  
1 5 10

<210> SEQ ID NO 125

<211> LENGTH: 5

-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 125

Ser Asn Gly Asn Thr
1           5

<210> SEQ ID NO 126
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 126

Glu Asn Ile Tyr Ser
1           5

<210> SEQ ID NO 127
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 127

Thr Ser Gly Tyr Ser
1           5

<210> SEQ ID NO 128
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 128

Gln Asp Ile Lys Ser
1           5

<210> SEQ ID NO 129
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 129

Gln Asn Ile Gly Asn
1           5

<210> SEQ ID NO 130
<211> LENGTH: 911
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Retargeted endopeptidase

<400> SEQUENCE: 130

Ile Ser Glu Phe Gly Ser Met Glu Phe Val Asn Lys Gln Phe Asn Tyr
1           5           10           15

Lys Asp Pro Val Asn Gly Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn
           20           25           30

Ala Gly Gln Met Gln Pro Val Lys Ala Phe Lys Ile His Asn Lys Ile
           35           40           45

Trp Val Ile Pro Glu Arg Asp Thr Phe Thr Asn Pro Glu Glu Gly Asp
           50           55           60

Leu Asn Pro Pro Pro Glu Ala Lys Gln Val Pro Val Ser Tyr Tyr Asp
65           70           75           80

```

-continued

---

Ser Thr Tyr Leu Ser Thr Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly  
                   85  90  95

Val Thr Lys Leu Phe Glu Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met  
                   100  105  110

Leu Leu Thr Ser Ile Val Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr  
                   115  120  125

Ile Asp Thr Glu Leu Lys Val Ile Asp Thr Asn Cys Ile Asn Val Ile  
                   130  135  140

Gln Pro Asp Gly Ser Tyr Arg Ser Glu Glu Leu Asn Leu Val Ile Ile  
                   145  150  155  160

Gly Pro Ser Ala Asp Ile Ile Gln Phe Glu Cys Lys Ser Phe Gly His  
                   165  170  175

Glu Val Leu Asn Leu Thr Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile  
                   180  185  190

Arg Phe Ser Pro Asp Phe Thr Phe Gly Phe Glu Glu Ser Leu Glu Val  
                   195  200  205

Asp Thr Asn Pro Leu Leu Gly Ala Gly Lys Phe Ala Thr Asp Pro Ala  
                   210  215  220

Val Thr Leu Ala His Glu Leu Ile His Ala Gly His Arg Leu Tyr Gly  
                   225  230  235  240

Ile Ala Ile Asn Pro Asn Arg Val Phe Lys Val Asn Thr Asn Ala Tyr  
                   245  250  255

Tyr Glu Met Ser Gly Leu Glu Val Ser Phe Glu Glu Leu Arg Thr Phe  
                   260  265  270

Gly Gly His Asp Ala Lys Phe Ile Asp Ser Leu Gln Glu Asn Glu Phe  
                   275  280  285

Arg Leu Tyr Tyr Tyr Asn Lys Phe Lys Asp Ile Ala Ser Thr Leu Asn  
                   290  295  300

Lys Ala Lys Ser Ile Val Gly Thr Thr Ala Ser Leu Gln Tyr Met Lys  
                   305  310  315  320

Asn Val Phe Lys Glu Lys Tyr Leu Leu Ser Glu Asp Thr Ser Gly Lys  
                   325  330  335

Phe Ser Val Asp Lys Leu Lys Phe Asp Lys Leu Tyr Lys Met Leu Thr  
                   340  345  350

Glu Ile Tyr Thr Glu Asp Asn Phe Val Lys Phe Phe Lys Val Leu Asn  
                   355  360  365

Arg Lys Thr Tyr Leu Asn Phe Asp Lys Ala Val Phe Lys Ile Asn Ile  
                   370  375  380

Val Pro Lys Val Asn Tyr Thr Ile Tyr Asp Gly Phe Asn Leu Arg Asn  
                   385  390  395  400

Thr Asn Leu Ala Ala Asn Phe Asn Gly Gln Asn Thr Glu Ile Asn Asn  
                   405  410  415

Met Asn Phe Thr Lys Leu Lys Asn Phe Thr Gly Leu Phe Glu Phe Tyr  
                   420  425  430

Lys Leu Leu Cys Val Asp Gly Ile Ile Thr Ser Lys Thr Lys Ser Leu  
                   435  440  445

Ile Glu Gly Arg Phe Gly Gly Phe Thr Gly Ala Arg Lys Ser Ala Arg  
                   450  455  460

Lys Arg Lys Asn Gln Ala Leu Ala Gly Gly Gly Gly Ser Gly Gly Gly  
                   465  470  475  480

-continued

---

Gly	Ser	Gly	Gly	Gly	Gly	Ser	Ala	Leu	Val	Leu	Gln	Cys	Ile	Lys	Val
				485					490					495	
Asn	Asn	Trp	Asp	Leu	Phe	Phe	Ser	Pro	Ser	Glu	Asp	Asn	Phe	Thr	Asn
			500					505					510		
Asp	Leu	Asn	Lys	Gly	Glu	Glu	Ile	Thr	Ser	Asp	Thr	Asn	Ile	Glu	Ala
		515					520					525			
Ala	Glu	Glu	Asn	Ile	Ser	Leu	Asp	Leu	Ile	Gln	Gln	Tyr	Tyr	Leu	Thr
	530				535						540				
Phe	Asn	Phe	Asp	Asn	Glu	Pro	Glu	Asn	Ile	Ser	Ile	Glu	Asn	Leu	Ser
545				550						555					560
Ser	Asp	Ile	Ile	Gly	Gln	Leu	Glu	Leu	Met	Pro	Asn	Ile	Glu	Arg	Phe
				565					570					575	
Pro	Asn	Gly	Lys	Lys	Tyr	Glu	Leu	Asp	Lys	Tyr	Thr	Met	Phe	His	Tyr
			580					585					590		
Leu	Arg	Ala	Gln	Glu	Phe	Glu	His	Gly	Lys	Ser	Arg	Ile	Ala	Leu	Thr
		595					600					605			
Asn	Ser	Val	Asn	Glu	Ala	Leu	Leu	Asn	Pro	Ser	Arg	Val	Tyr	Thr	Phe
	610					615					620				
Phe	Ser	Ser	Asp	Tyr	Val	Lys	Lys	Val	Asn	Lys	Ala	Thr	Glu	Ala	Ala
625					630					635					640
Met	Phe	Leu	Gly	Trp	Val	Glu	Gln	Leu	Val	Tyr	Asp	Phe	Thr	Asp	Glu
				645					650					655	
Thr	Ser	Glu	Val	Ser	Thr	Thr	Asp	Lys	Ile	Ala	Asp	Ile	Thr	Ile	Ile
			660					665					670		
Ile	Pro	Tyr	Ile	Gly	Pro	Ala	Leu	Asn	Ile	Gly	Asn	Met	Leu	Tyr	Lys
		675					680					685			
Asp	Asp	Phe	Val	Gly	Ala	Leu	Ile	Phe	Ser	Gly	Ala	Val	Ile	Leu	Leu
	690					695					700				
Glu	Phe	Ile	Pro	Glu	Ile	Ala	Ile	Pro	Val	Leu	Gly	Thr	Phe	Ala	Leu
705					710					715					720
Val	Ser	Tyr	Ile	Ala	Asn	Lys	Val	Leu	Thr	Val	Gln	Thr	Ile	Asp	Asn
				725					730					735	
Ala	Leu	Ser	Lys	Arg	Asn	Glu	Lys	Trp	Asp	Glu	Val	Tyr	Lys	Tyr	Ile
			740					745					750		
Val	Thr	Asn	Trp	Leu	Ala	Lys	Val	Asn	Thr	Gln	Ile	Asp	Leu	Ile	Arg
		755						760				765			
Lys	Lys	Met	Lys	Glu	Ala	Leu	Glu	Asn	Gln	Ala	Glu	Ala	Thr	Lys	Ala
	770					775						780			
Ile	Ile	Asn	Tyr	Gln	Tyr	Asn	Gln	Tyr	Thr	Glu	Glu	Glu	Lys	Asn	Asn
785					790						795				800
Ile	Asn	Phe	Asn	Ile	Asp	Asp	Leu	Ser	Ser	Lys	Leu	Asn	Glu	Ser	Ile
			805						810					815	
Asn	Lys	Ala	Met	Ile	Asn	Ile	Asn	Lys	Phe	Leu	Asn	Gln	Cys	Ser	Val
			820					825					830		
Ser	Tyr	Leu	Met	Asn	Ser	Met	Ile	Pro	Tyr	Gly	Val	Lys	Arg	Leu	Glu
		835					840					845			
Asp	Phe	Asp	Ala	Ser	Leu	Lys	Asp	Ala	Leu	Leu	Lys	Tyr	Ile	Tyr	Asp
	850					855						860			
Asn	Arg	Gly	Thr	Leu	Ile	Gly	Gln	Val	Asp	Arg	Leu	Lys	Asp	Lys	Val
865					870					875					880
Asn	Asn	Thr	Leu	Ser	Thr	Asp	Ile	Pro	Phe	Gln	Leu	Ser	Lys	Tyr	Val

-continued

---

	885							890							895
Asp	Asn	Gln	Arg	Leu	Leu	Ser	Thr	Leu	Glu	Ala	Leu	Ala	Ser	Gly	
	900							905						910	

<210> SEQ ID NO 131  
 <211> LENGTH: 904  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Retargeted endopeptidase

<400> SEQUENCE: 131

Met	Gly	Ser	Met	Glu	Phe	Val	Asn	Lys	Gln	Phe	Asn	Tyr	Lys	Asp	Pro
1			5						10					15	
Val	Asn	Gly	Val	Asp	Ile	Ala	Tyr	Ile	Lys	Ile	Pro	Asn	Ala	Gly	Gln
			20					25						30	
Met	Gln	Pro	Val	Lys	Ala	Phe	Lys	Ile	His	Asn	Lys	Ile	Trp	Val	Ile
		35					40					45			
Pro	Glu	Arg	Asp	Thr	Phe	Thr	Asn	Pro	Glu	Glu	Gly	Asp	Leu	Asn	Pro
	50					55					60				
Pro	Pro	Glu	Ala	Lys	Gln	Val	Pro	Val	Ser	Tyr	Tyr	Asp	Ser	Thr	Tyr
65					70					75					80
Leu	Ser	Thr	Asp	Asn	Glu	Lys	Asp	Asn	Tyr	Leu	Lys	Gly	Val	Thr	Lys
				85					90					95	
Leu	Phe	Glu	Arg	Ile	Tyr	Ser	Thr	Asp	Leu	Gly	Arg	Met	Leu	Leu	Thr
			100					105					110		
Ser	Ile	Val	Arg	Gly	Ile	Pro	Phe	Trp	Gly	Gly	Ser	Thr	Ile	Asp	Thr
		115					120					125			
Glu	Leu	Lys	Val	Ile	Asp	Thr	Asn	Cys	Ile	Asn	Val	Ile	Gln	Pro	Asp
	130					135					140				
Gly	Ser	Tyr	Arg	Ser	Glu	Glu	Leu	Asn	Leu	Val	Ile	Ile	Gly	Pro	Ser
145				150						155				160	
Ala	Asp	Ile	Ile	Gln	Phe	Glu	Cys	Lys	Ser	Phe	Gly	His	Glu	Val	Leu
			165						170					175	
Asn	Leu	Thr	Arg	Asn	Gly	Tyr	Gly	Ser	Thr	Gln	Tyr	Ile	Arg	Phe	Ser
			180					185					190		
Pro	Asp	Phe	Thr	Phe	Gly	Phe	Glu	Glu	Ser	Leu	Glu	Val	Asp	Thr	Asn
		195					200					205			
Pro	Leu	Leu	Gly	Ala	Gly	Lys	Phe	Ala	Thr	Asp	Pro	Ala	Val	Thr	Leu
	210					215					220				
Ala	His	Glu	Leu	Ile	His	Ala	Gly	His	Arg	Leu	Tyr	Gly	Ile	Ala	Ile
225					230					235					240
Asn	Pro	Asn	Arg	Val	Phe	Lys	Val	Asn	Thr	Asn	Ala	Tyr	Tyr	Glu	Met
			245						250					255	
Ser	Gly	Leu	Glu	Val	Ser	Phe	Glu	Glu	Leu	Arg	Thr	Phe	Gly	Gly	His
		260						265					270		
Asp	Ala	Lys	Phe	Ile	Asp	Ser	Leu	Gln	Glu	Asn	Glu	Phe	Arg	Leu	Tyr
		275					280					285			
Tyr	Tyr	Asn	Lys	Phe	Lys	Asp	Ile	Ala	Ser	Thr	Leu	Asn	Lys	Ala	Lys
	290					295					300				
Ser	Ile	Val	Gly	Thr	Thr	Ala	Ser	Leu	Gln	Tyr	Met	Lys	Asn	Val	Phe
305					310					315					320
Lys	Glu	Lys	Tyr	Leu	Leu	Ser	Glu	Asp	Thr	Ser	Gly	Lys	Phe	Ser	Val



-continued

---

Lys Trp Asp Glu Val Tyr Lys Tyr Ile Val Thr Asn Trp Leu Ala Lys  
 740 745 750

Val Asn Thr Gln Ile Asp Leu Ile Arg Lys Lys Met Lys Glu Ala Leu  
 755 760 765

Glu Asn Gln Ala Glu Ala Thr Lys Ala Ile Ile Asn Tyr Gln Tyr Asn  
 770 775 780

Gln Tyr Thr Glu Glu Glu Lys Asn Asn Ile Asn Phe Asn Ile Asp Asp  
 785 790 795 800

Leu Ser Ser Lys Leu Asn Glu Ser Ile Asn Lys Ala Met Ile Asn Ile  
 805 810 815

Asn Lys Phe Leu Asn Gln Cys Ser Val Ser Tyr Leu Met Asn Ser Met  
 820 825 830

Ile Pro Tyr Gly Val Lys Arg Leu Glu Asp Phe Asp Ala Ser Leu Lys  
 835 840 845

Asp Ala Leu Leu Lys Tyr Ile Tyr Asp Asn Arg Gly Thr Leu Ile Gly  
 850 855 860

Gln Val Asp Arg Leu Lys Asp Lys Val Asn Asn Thr Leu Ser Thr Asp  
 865 870 875 880

Ile Pro Phe Gln Leu Ser Lys Tyr Val Asp Asn Gln Arg Leu Leu Ser  
 885 890 895

Thr Leu Glu Ala Leu Ala Ser Gly  
 900

<210> SEQ ID NO 132  
 <211> LENGTH: 360  
 <212> TYPE: DNA  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 132

cagggtgaagc tgcaggagtc tggacctgaa ctggtaaagc ctggggcttc agtgaagatg 60

tcttgcaagg cttctggata cacattcact aactatgta tacactgggt gaagcaaaag 120

cctgggcagg gccttgagtg gattggatat attaacctt acaatgatgg ctctaagtac 180

aatgagaagt tcaaaggcaa ggcctcactg acttcagaca aatcctccag cacagcctac 240

atggagctca gcagcctgac ctctgaggac tctgcggtct attactgtgc aagacatctc 300

gctaatacct actactactt tgactactgg ggccaaggca ccaactctcac agtctcctca 360

<210> SEQ ID NO 133  
 <211> LENGTH: 120  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 133

Gln Val Gln Leu Gln Glu Ser Gly Pro Glu Leu Val Lys Pro Gly Ala  
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr  
 20 25 30

Val Ile His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile  
 35 40 45

Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Ser Lys Tyr Asn Glu Lys Phe  
 50 55 60

Lys Gly Lys Ala Ser Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr  
 65 70 75 80

-continued

---

Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

Ala Arg His Leu Ala Asn Thr Tyr Tyr Tyr Phe Asp Tyr Trp Gly Gln  
100 105 110

Gly Thr Thr Leu Thr Val Ser Ser  
115 120

<210> SEQ ID NO 134  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 134

Ala Arg Met Gly Tyr  
1 5

<210> SEQ ID NO 135  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 135

Ala Arg His Leu Ala Asn Thr Tyr Tyr Tyr Phe Asp Tyr  
1 5 10

<210> SEQ ID NO 136  
<211> LENGTH: 349  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 136

Met Glu Leu Ala Val Gly Asn Leu Ser Glu Gly Asn Ala Ser Trp Pro  
1 5 10 15

Glu Pro Pro Ala Pro Glu Pro Gly Pro Leu Phe Gly Ile Gly Val Glu  
20 25 30

Asn Phe Val Thr Leu Val Val Phe Gly Leu Ile Phe Ala Leu Gly Val  
35 40 45

Leu Gly Asn Ser Leu Val Ile Thr Val Leu Ala Arg Ser Lys Pro Gly  
50 55 60

Lys Pro Arg Ser Thr Thr Asn Leu Phe Ile Leu Asn Leu Ser Ile Ala  
65 70 75 80

Asp Leu Ala Tyr Leu Leu Phe Cys Ile Pro Phe Gln Ala Thr Val Tyr  
85 90 95

Ala Leu Pro Thr Trp Val Leu Gly Ala Phe Ile Cys Lys Phe Ile His  
100 105 110

Tyr Phe Phe Thr Val Ser Met Leu Val Ser Ile Phe Thr Leu Ala Ala  
115 120 125

Met Ser Val Asp Arg Tyr Val Ala Ile Val His Ser Arg Arg Ser Ser  
130 135 140

Ser Leu Arg Val Ser Arg Asn Ala Leu Leu Gly Val Gly Cys Ile Trp  
145 150 155 160

Ala Leu Ser Ile Ala Met Ala Ser Pro Val Ala Tyr His Gln Gly Leu  
165 170 175

Phe His Pro Arg Ala Ser Asn Gln Thr Phe Cys Trp Glu Gln Trp Pro  
180 185 190

-continued

---

Asp Pro Arg His Lys Lys Ala Tyr Val Val Cys Thr Phe Val Phe Gly  
 195 200 205

Tyr Leu Leu Pro Leu Leu Leu Ile Cys Phe Cys Tyr Ala Lys Val Leu  
 210 215 220

Asn His Leu His Lys Lys Leu Lys Asn Met Ser Lys Lys Ser Glu Ala  
 225 230 235 240

Ser Lys Lys Lys Thr Ala Gln Thr Val Leu Val Val Val Val Phe  
 245 250 255

Gly Ile Ser Trp Leu Pro His His Ile Ile His Leu Trp Ala Glu Phe  
 260 265 270

Gly Val Phe Pro Leu Thr Pro Ala Ser Phe Leu Phe Arg Ile Thr Ala  
 275 280 285

His Cys Leu Ala Tyr Ser Asn Ser Ser Val Asn Pro Ile Ile Tyr Ala  
 290 295 300

Phe Leu Ser Glu Asn Phe Arg Lys Ala Tyr Lys Gln Val Phe Lys Cys  
 305 310 315 320

His Ile Arg Lys Asp Ser His Leu Ser Asp Thr Lys Glu Asn Lys Ser  
 325 330 335

Arg Ile Asp Thr Pro Pro Ser Thr Asn Cys Thr His Val  
 340 345

<210> SEQ ID NO 137  
 <211> LENGTH: 349  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 137

Met Glu Leu Ala Val Gly Asn Leu Ser Glu Gly Asn Ala Ser Trp Pro  
 1 5 10 15

Glu Pro Pro Ala Pro Glu Pro Gly Pro Leu Phe Gly Ile Gly Val Glu  
 20 25 30

Asn Phe Val Thr Leu Val Val Phe Gly Leu Ile Phe Ala Leu Gly Val  
 35 40 45

Leu Gly Asn Ser Leu Val Ile Thr Val Leu Ala Arg Ser Lys Pro Gly  
 50 55 60

Lys Pro Arg Ser Thr Thr Asn Leu Phe Ile Leu Asn Leu Ser Ile Ala  
 65 70 75 80

Asp Leu Ala Tyr Leu Leu Phe Cys Ile Pro Phe Gln Ala Thr Val Tyr  
 85 90 95

Ala Leu Pro Thr Trp Val Leu Gly Ala Phe Ile Cys Lys Phe Ile His  
 100 105 110

Tyr Phe Phe Thr Val Ser Met Leu Val Ser Ile Phe Thr Leu Ala Ala  
 115 120 125

Met Ser Val Asp Arg Tyr Val Ala Ile Val His Ser Arg Arg Ser Ser  
 130 135 140

Ser Leu Arg Val Ser Arg Asn Ala Leu Leu Gly Val Gly Cys Ile Trp  
 145 150 155 160

Ala Leu Ser Ile Ala Met Ala Ser Pro Val Ala Tyr His Gln Gly Leu  
 165 170 175

Phe His Pro Arg Ala Ser Asn Gln Thr Phe Cys Trp Glu Gln Trp Pro  
 180 185 190

Asp Pro Arg His Lys Lys Ala Tyr Val Val Cys Thr Phe Val Phe Gly  
 195 200 205

-continued

---

Tyr Leu Leu Pro Leu Leu Leu Ile Cys Phe Cys Tyr Ala Lys Val Leu  
 210 215 220  
 Asn His Leu His Lys Lys Leu Lys Asn Met Ser Lys Lys Ser Glu Ala  
 225 230 235 240  
 Ser Lys Lys Lys Thr Ala Gln Thr Val Leu Val Val Val Val Phe  
 245 250 255  
 Gly Ile Ser Trp Leu Pro His His Ile Ile His Leu Trp Ala Glu Phe  
 260 265 270  
 Gly Val Phe Pro Leu Thr Pro Ala Ser Phe Leu Phe Arg Ile Thr Ala  
 275 280 285  
 His Cys Leu Ala Tyr Ser Asn Ser Ser Val Asn Pro Ile Ile Tyr Ala  
 290 295 300  
 Phe Leu Ser Glu Asn Phe Arg Lys Ala Tyr Lys Gln Val Phe Lys Cys  
 305 310 315 320  
 His Ile Arg Lys Asp Ser His Leu Ser Asp Thr Lys Glu Ser Lys Ser  
 325 330 335  
 Arg Ile Asp Thr Pro Pro Ser Thr Asn Cys Thr His Val  
 340 345

&lt;210&gt; SEQ ID NO 138

&lt;211&gt; LENGTH: 349

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 138

Met Glu Leu Ala Val Gly Asn Leu Ser Glu Gly Asn Ala Ser Cys Pro  
 1 5 10 15  
 Glu Pro Pro Ala Pro Glu Pro Gly Pro Leu Phe Gly Ile Gly Val Glu  
 20 25 30  
 Asn Phe Val Thr Leu Val Val Phe Gly Leu Ile Phe Ala Leu Gly Val  
 35 40 45  
 Leu Gly Asn Ser Leu Val Ile Thr Val Leu Ala Arg Ser Lys Pro Gly  
 50 55 60  
 Lys Pro Arg Ser Thr Thr Asn Leu Phe Ile Leu Asn Leu Ser Ile Ala  
 65 70 75 80  
 Asp Leu Ala Tyr Leu Leu Phe Cys Ile Pro Phe Gln Ala Thr Val Tyr  
 85 90 95  
 Ala Leu Pro Thr Trp Val Leu Gly Ala Phe Ile Cys Lys Phe Ile His  
 100 105 110  
 Tyr Phe Phe Thr Val Ser Met Leu Val Ser Ile Phe Thr Leu Ala Ala  
 115 120 125  
 Met Ser Val Asp Arg Tyr Val Ala Ile Val His Ser Arg Arg Ser Ser  
 130 135 140  
 Ser Leu Arg Val Ser Arg Asn Ala Leu Leu Gly Val Gly Cys Ile Trp  
 145 150 155 160  
 Ala Leu Ser Ile Ala Met Ala Ser Pro Val Ala Tyr His Gln Gly Leu  
 165 170 175  
 Phe His Pro Arg Ala Ser Asn Gln Thr Phe Cys Trp Glu Gln Trp Pro  
 180 185 190  
 Asp Pro Arg His Lys Lys Ala Tyr Val Val Cys Thr Phe Val Phe Gly  
 195 200 205  
 Tyr Leu Leu Pro Leu Leu Leu Ile Cys Phe Cys Tyr Ala Lys Val Leu



-continued

---

```

Ser Gly Ala Arg Arg Ala Lys Arg Lys Val Thr Arg Met Ile Leu Ile
225                230                235                240

Val Ala Ala Leu Phe Cys Leu Cys Trp Met Pro His His Ala Leu Ile
                245                250                255

Leu Cys Val Trp Phe Gly Gln Phe Pro Leu Thr Arg Ala Thr Tyr Ala
                260                265                270

Leu Arg Ile Leu Ser His Leu Val Ser Tyr Ala Asn Ser Cys Val Asn
                275                280                285

Pro Ile Val Tyr Ala Leu Val Ser Lys His Phe Arg Lys Gly Phe Arg
                290                295                300

Thr Ile Cys Ala Gly Leu Leu Gly Arg Ala Pro Gly Arg Ala Ser Gly
305                310                315                320

Arg Val Cys Ala Ala Ala Arg Gly Thr His Ser Gly Ser Val Leu Glu
                325                330                335

Arg Glu Ser Ser Asp Leu Leu His Met Ser Glu Ala Ala Gly Ala Leu
                340                345                350

Arg Pro Cys Pro Gly Ala Ser Gln Pro Cys Ile Leu Glu Pro Cys Pro
                355                360                365

Gly Pro Ser Trp Gln Gly Pro Lys Ala Gly Asp Ser Ile Leu Thr Val
370                375                380

Asp Val Ala
385

```

```

<210> SEQ ID NO 140
<211> LENGTH: 368
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 140

```

```

Met Ala Asp Ala Gln Asn Ile Ser Leu Asp Ser Pro Gly Ser Val Gly
1                5                10                15

Ala Val Ala Val Pro Val Val Phe Ala Leu Ile Phe Leu Leu Gly Thr
                20                25                30

Val Gly Asn Gly Leu Val Leu Ala Val Leu Leu Gln Pro Gly Pro Ser
35                40                45

Ala Trp Gln Glu Pro Gly Ser Thr Thr Asp Leu Phe Ile Leu Asn Leu
50                55                60

Ala Val Ala Asp Leu Cys Phe Ile Leu Cys Cys Val Pro Phe Gln Ala
65                70                75                80

Thr Ile Tyr Thr Leu Asp Ala Trp Leu Phe Gly Ala Leu Val Cys Lys
85                90                95

Ala Val His Leu Leu Ile Tyr Leu Thr Met Tyr Ala Ser Ser Phe Thr
100               105               110

Leu Ala Ala Val Ser Val Asp Arg Tyr Leu Ala Val Arg His Pro Leu
115               120               125

Arg Ser Arg Ala Leu Arg Thr Pro Arg Asn Ala Arg Ala Ala Val Gly
130               135               140

Leu Val Trp Leu Leu Ala Ala Leu Phe Ser Ala Pro Tyr Leu Ser Tyr
145               150               155               160

Tyr Gly Thr Val Arg Tyr Gly Ala Leu Glu Leu Cys Val Pro Ala Trp
165               170               175

Glu Asp Ala Arg Arg Arg Ala Leu Asp Val Ala Thr Phe Ala Ala Gly
180               185               190

```

-continued

Tyr Leu Leu Pro Val Ala Val Val Ser Leu Ala Tyr Gly Arg Thr Leu  
 195 200 205  
 Arg Phe Leu Trp Ala Ala Val Gly Pro Ala Gly Ala Ala Ala Ala Glu  
 210 215 220  
 Ala Arg Arg Arg Ala Thr Gly Arg Ala Gly Arg Ala Met Leu Ala Val  
 225 230 235 240  
 Ala Ala Leu Tyr Ala Leu Cys Trp Gly Pro His His Ala Leu Ile Leu  
 245 250 255  
 Cys Phe Trp Tyr Gly Arg Phe Ala Phe Ser Pro Ala Thr Tyr Ala Cys  
 260 265 270  
 Arg Leu Ala Ser His Cys Leu Ala Tyr Ala Asn Ser Cys Leu Asn Pro  
 275 280 285  
 Leu Val Tyr Ala Leu Ala Ser Arg His Phe Arg Ala Arg Phe Arg Arg  
 290 295 300  
 Leu Trp Pro Cys Gly Arg Arg Arg Arg His Arg Ala Arg Arg Ala Leu  
 305 310 315 320  
 Arg Arg Val Arg Pro Ala Ser Ser Gly Pro Pro Gly Cys Pro Gly Asp  
 325 330 335  
 Ala Arg Pro Ser Gly Arg Leu Leu Ala Gly Gly Gly Gln Gly Pro Glu  
 340 345 350  
 Pro Arg Glu Gly Pro Val His Gly Gly Glu Ala Ala Arg Gly Pro Glu  
 355 360 365

&lt;210&gt; SEQ ID NO 141

&lt;211&gt; LENGTH: 1050

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 141

```

atggagctgg cggtcgggaa cctcagcgag ggcaacgcga gctggccgga gcccccgcc 60
ccggagcccg ggccgctgtt cggcatcggc gtggagaact tcgtcacgct ggtggtgttc 120
ggcctgatct tcgctgtggg tgtgctgggc aacagcctag tgatcacgct gctggcgcgc 180
agcaagccgg gcaagccgcg gacgaccacc aacctgttca tcctcaacct gagcatcgcc 240
gacctggcct acctgctctt ctgcatcccc ttccaggcca ccgtgtacgc gctgcccacc 300
tgggtgctgg ggcgcttcat ctgcaagtcc atccactact tcttaccgct gtccatgctg 360
gtgagcatct tcaccctggc cgcgatgtcc gtggaccgct acgtggccat cgtgcactcg 420
cggcgctcct cctcctcag ggtgtcccgc aacgcgctgc tgggcgtggg ctgcatctgg 480
gcgctgtcca ttgccatggc ctcgcccgtg gcttaccacc agggcctctt ccaccgcgcc 540
gccagcaacc agacctctg ctgggagcag tggcccgcacc ctgcaccaca gaaggcctac 600
gtggtgtgca ccttctgctt cggctacctg ctgccgctcc tgetcatctg cttctgetat 660
gccaaggtcc ttaatcactt gcataaaaag ttgaagaaca tgtcaaagaa gtctgaagca 720
tccaagaaaa agactgcaca gacagttctg gtggtggttg tgggttttg aatctcctgg 780
ctgccgcacc acatcatcca tctctgggct gagtttgag ttttcccgt gacgcccgtc 840
tccttctctc tcagaatcac gcgccactgc ctggcgtaga gcaattctc cgtgaatcct 900
atcatttatg catttctctc tgaaaatttc aggaagcctc ataaacaagt gttcaagtgt 960
cacattcgca aagattcaca cctgagtgat actaaagaaa ataaaagtcg aatagacacc 1020
  
```

-continued

---

ccaccatcaa ccaattgtac tcatgtgtga 1050

<210> SEQ ID NO 142

<211> LENGTH: 1050

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 142

atggagctgg cggtcgggaa cctcagcgag ggcaacgcga gctggccgga gccccccgcc 60  
ccggagcccg ggccgctgtt cggcatcggc gtggagaact tcgtcacgct ggtggtgttc 120  
ggcctgatct tcgcgtggg tgtgctgggc aacagcctag tgatcacgct getggcgcgc 180  
agcaagcccg gcaagcccg gagcaccacc aacctgttca tctcaacct gagcatcgcc 240  
gacctggcct acctgctctt ctgcatcccc ttccaggcca cgtgtacgc gctgcccacc 300  
tgggtgctgg gcgccttcat ctgcaagtcc atccactact tcttcaccgt gtccatgctg 360  
gtgagcatct tcacctggc cgcgatgtcc gtggaccgct acgtggccat cgtgcaactc 420  
cggcgctcct cctccctcag ggtgtcccgc aacgcgctgc tgggcgtggg ctgcatctgg 480  
gcgctgtcca ttgccatggc ctgcgccgtg gcctaccacc agggcctctt ccaccgcgc 540  
gccagcaacc agacctctg ctgggagcag tggcccgacc ctgcacaaa gaaggcctac 600  
gtggtgtgca ccttcgtctt cggctacctg ctgccgctcc tgcctcatctg cttctgctat 660  
gccaaggctc ttaatcactt gcataaaaag ttgaagaaca tgtcaaagaa gtctgaagca 720  
tccaagaaaa agactgcaca gacagtctct gtggtggttg tgggtgttgg aatctcctgg 780  
ctgccgcacc acatcatcca tctctgggct gagtttgag ttttcccgt gaegccggct 840  
tccttctctc tcagaatcac gcgccactgc ctggcgta gcaattctc cgtgaatcct 900  
atcatttatg catttctctc tgaaaattc aggaaggcct ataaacaagt gttcaagtgt 960  
cacattcgca aagattcaca cctgagtgat actaaagaaa gtaaaagtgc aatagacacc 1020  
ccaccatcaa ccaattgtac tcatgtgtga 1050

<210> SEQ ID NO 143

<211> LENGTH: 1050

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 143

atggagctgg cggtcgggaa cctcagcgag ggcaacgcga gctgtccgga gccccccgcc 60  
ccggagcccg ggccgctgtt cggcatcggc gtggagaact tcgtcacgct ggtggtgttc 120  
ggcctgatct tcgcgtggg cgtgctgggc aacagcctag tgatcacgct getggcgcgc 180  
agcaagcccg gcaagcccg gagcaccacc aacctgttca tctcaacct gagcatcgcc 240  
gacctggcct acctgctctt ctgcatcccc ttccaggcca cgtgtacgc gctgcccacc 300  
tgggtgctgg gcgccttcat ctgcaagtcc atccactact tcttcaccgt gtccatgctg 360  
gtgagcatct tcacctggc cgcgatgtcc gtggaccgct acgtggccat cgtgcaactc 420  
cggcgctcct cctccctcag ggtgtcccgc aacgcgctgc tgggcgtggg ctgcatctgg 480  
gcgctgtcca ttgccatggc ctgcgccgtg gcctaccacc agggcctctt ccaccgcgc 540  
gccagcaacc agacctctg ctgggagcag tggcccgacc ctgcacaaa gaaggcctac 600  
gtggtgtgca ccttcgtctt cggctacctg ctgccgctcc tgcctcatctg cttctgctat 660

-continued

---

```

gccaaggtcc ttaatcactt gcataaaaag ttgaagaaca tgtcaaagaa gtctgaagca 720
tccaagaaaa agactgcaca gacagttctg gtggtggttg tgggtgttgg aatctcctgg 780
ctgccgcacc acatcatcca tctctgggct gagtttgag ttttccgct gaecgceggt 840
tccttctctc tcagaatcac cgcccactgc ctggcgta ca gcaattctc cgtgaatcct 900
atcatttatg catttctctc tgaaaatttc aggaaggcct ataaacaagt gttcaagtgt 960
cacattcgca aagattcaca cctgagtgat actaaagaaa ataaaagtcg aatagacacc 1020
ccaccatcaa ccaattgtac tcatgtgtga 1050

```

```

<210> SEQ ID NO 144
<211> LENGTH: 1164
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 144

```

```

atgaacgtct cgggctgccc aggggcccgg aacgcgagcc aggcgggccc cgggggaggc 60
tggaaccccc aggcgggtcat cgtgccctg ctcttccgct tcactcttct cgtgggcacc 120
gtgggcaaca cgtcgtgctt ggcggtgctg ctgcgcccgg gccaggcggc cagcaactacc 180
aacctgttca tccttaacct gggcgtggcc gacctgtgtt tcactctgtg ctgcgtgccc 240
ttccaggcca ccactacac cctggacggc tgggtgttcg gctcgtgct gtgcaaggcg 300
gtgcacttcc tcacttctc caccatgcac gccagcagct tcacgctggc cgcgctctcc 360
ctggacaggt atctggccat ccgctacccc ctgcactccc gcgagctgag caegcctcga 420
aacgcgctgg cagccatcgg gctcactcgg gggctgtcgc tgcctctctc cgggcccctac 480
ctgagctact accgccagtc gcagctggcc aacctgaccg tgtgccatcc cgcgtggagc 540
gcccctcgcc gccgcgccat ggacatctgc acctctgtct tcagctacct gcttctctgtg 600
ctggttctcg gctgaccta cgcgcgcacc ttgcgctacc tctggcgcgc cgtcgacccc 660
gtggccgccc gctcgggtgc ccggcgcgcc aagcgcgaagg tgacacgcat gatcctcacc 720
gtggccgccc tcttctgctt ctgtggatg ccccaccacg cgctcactct ctgcgtgtgg 780
ttcggccagt tcccgtcac gcgcgcacct tatgccttc gcactctctc gcacctggtc 840
tcctacgcca actcctgctt caacccatc gtttacgcgc tggctcctca gcaactccgc 900
aaaggcttcc gaacgatctg cgcgggcccgt ctgggcccgt ccccaggccc agcctcgggc 960
cgtgtgtgag ctgcccgcgc gggcaccac agtggcagcg tgttgagagc cgagtccagc 1020
gacctgttgc acatgagcga ggcggcgggg gccctctgct cctgcccggc cgttcccagc 1080
ccatgcatcc tcgagccctg tcctggcccg tcctggcagg gcccaaggc aggcgacagc 1140
atcctgacgg ttgatgtggc ctga 1164

```

```

<210> SEQ ID NO 145
<211> LENGTH: 1107
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 145

```

```

atggctgatg cccagaacat ttcactggac agcccaggga gtgtgggggc cgtggcagtg 60
cctgtggtct ttgccctaat ctctctgctg ggcacagtgg gcaatgggct ggtgctggca 120
gtgctcctgc agctggccc gagtgctggc caggagcctg gcagcaccac ggacctgttc 180

```

-continued

---

```

atcctcaacc tggcggtagc tgacctctgc ttcactctgt gctgcgtgcc cttccaggcc 240
accatctaca cgctggatgc ctggctcttt ggggcccctcg tctgcaaggc cgtgcacctg 300
ctcatctacc tcaccatgta cggcagcagc tttacgctgg ctgctgtctc cgtggacagg 360
tacctggccg tgcggcaccg gctgcgctcg cgcgccctgc gcacgcccg taacgcccgc 420
gccgcagtgg ggctgggtgtg gctgctggcg gcgctcttct cggcgccta cctcagctac 480
tacggcaccg tgcgtacgg cgcgctggag ctctgctgc cgcctggga ggacgcgcgc 540
cgccgcgcc tggactggc caccttgcgt gccggctacc tgtgcccgt ggctgtggtg 600
agcctggcct acgggcgcac gctgcgcttc ctgtgggccc cgtgggtcc cgcgggcgcg 660
gcgggggccc aggcgcggcg gagggcgacg ggccgcgcgg ggccgcctat gctggcggtg 720
gccgcgctct acgcgctctg ctggggtccg caccacgcgc tcatctgtg cttctggtac 780
ggccgcttcg ccttcagccc gggcacctac gcctgccgcc tggcctcaca ctgctggcc 840
tacgccaact cctgcctcaa cccgctcgtc tacgcgctcg cctcgcgcca cttccgcgcg 900
cgcttcgcc gcctgtggcc gtgcccgcgc cgacgccgc accgtgcccg cgcgccttg 960
cgtcgcgtcc gccccgcgtc ctcgggcccc cccggtgcc ccggagacgc ccggcctagc 1020
gggaggctgc tggtggtgg cggccagggc ccggagccca gggagggacc cgtccacggc 1080
ggagaggctg cccgaggacc ggaataa 1107
    
```

```

<210> SEQ ID NO 146
<211> LENGTH: 886
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: LHn/A fragment
    
```

<400> SEQUENCE: 146

```

Met Pro Phe Val Asn Lys Gln Phe Asn Tyr Lys Asp Pro Val Asn Gly
1           5           10           15

Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn Ala Gly Gln Met Gln Pro
          20           25           30

Val Lys Ala Phe Lys Ile His Asn Lys Ile Trp Val Ile Pro Glu Arg
          35           40           45

Asp Thr Phe Thr Asn Pro Glu Glu Gly Asp Leu Asn Pro Pro Pro Glu
          50           55           60

Ala Lys Gln Val Pro Val Ser Tyr Tyr Asp Ser Thr Tyr Leu Ser Thr
          65           70           75           80

Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly Val Thr Lys Leu Phe Glu
          85           90           95

Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met Leu Leu Thr Ser Ile Val
          100          105          110

Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr Ile Asp Thr Glu Leu Lys
          115          120          125

Val Ile Asp Thr Asn Cys Ile Asn Val Ile Gln Pro Asp Gly Ser Tyr
          130          135          140

Arg Ser Glu Glu Leu Asn Leu Val Ile Ile Gly Pro Ser Ala Asp Ile
          145          150          155          160

Ile Gln Phe Glu Cys Lys Ser Phe Gly His Glu Val Leu Asn Leu Thr
          165          170          175

Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile Arg Phe Ser Pro Asp Phe
    
```

-continued

180				185				190							
Thr	Phe	Gly	Phe	Glu	Glu	Ser	Leu	Glu	Val	Asp	Thr	Asn	Pro	Leu	Leu
		195					200					205			
Gly	Ala	Gly	Lys	Phe	Ala	Thr	Asp	Pro	Ala	Val	Thr	Leu	Ala	His	Glu
	210					215						220			
Leu	Ile	His	Ala	Gly	His	Arg	Leu	Tyr	Gly	Ile	Ala	Ile	Asn	Pro	Asn
225					230					235					240
Arg	Val	Phe	Lys	Val	Asn	Thr	Asn	Ala	Tyr	Tyr	Glu	Met	Ser	Gly	Leu
			245						250					255	
Glu	Val	Ser	Phe	Glu	Glu	Leu	Arg	Thr	Phe	Gly	Gly	His	Asp	Ala	Lys
			260						265				270		
Phe	Ile	Asp	Ser	Leu	Gln	Glu	Asn	Glu	Phe	Arg	Leu	Tyr	Tyr	Tyr	Asn
		275					280						285		
Lys	Phe	Lys	Asp	Ile	Ala	Ser	Thr	Leu	Asn	Lys	Ala	Lys	Ser	Ile	Val
	290					295					300				
Gly	Thr	Thr	Ala	Ser	Leu	Gln	Tyr	Met	Lys	Asn	Val	Phe	Lys	Glu	Lys
305					310					315					320
Tyr	Leu	Leu	Ser	Glu	Asp	Thr	Ser	Gly	Lys	Phe	Ser	Val	Asp	Lys	Leu
			325						330					335	
Lys	Phe	Asp	Lys	Leu	Tyr	Lys	Met	Leu	Thr	Glu	Ile	Tyr	Thr	Glu	Asp
		340						345					350		
Asn	Phe	Val	Lys	Phe	Phe	Lys	Val	Leu	Asn	Arg	Lys	Thr	Tyr	Leu	Asn
		355					360						365		
Phe	Asp	Lys	Ala	Val	Phe	Lys	Ile	Asn	Ile	Val	Pro	Lys	Val	Asn	Tyr
	370					375					380				
Thr	Ile	Tyr	Asp	Gly	Phe	Asn	Leu	Arg	Asn	Thr	Asn	Leu	Ala	Ala	Asn
385					390					395					400
Phe	Asn	Gly	Gln	Asn	Thr	Glu	Ile	Asn	Asn	Met	Asn	Phe	Thr	Lys	Leu
			405						410					415	
Lys	Asn	Phe	Thr	Gly	Leu	Phe	Glu	Phe	Tyr	Lys	Leu	Leu	Cys	Val	Arg
			420						425				430		
Gly	Ile	Ile	Thr	Ser	Lys	Thr	Lys	Ser	Leu	Gly	Gly	Gly	Gly	Glu	Asn
		435					440						445		
Leu	Tyr	Phe	Gln	Gly	Gly	Gly	Gly	Gly	Asp	Lys	Gly	Tyr	Asn	Lys	Ala
	450					455					460				
Phe	Asn	Asp	Leu	Cys	Ile	Lys	Val	Asn	Asn	Trp	Asp	Leu	Phe	Phe	Ser
465					470					475					480
Pro	Ser	Glu	Asp	Asn	Phe	Thr	Asn	Asp	Leu	Asn	Lys	Gly	Glu	Glu	Ile
			485						490					495	
Thr	Ser	Asp	Thr	Asn	Ile	Glu	Ala	Ala	Glu	Glu	Asn	Ile	Ser	Leu	Asp
			500						505				510		
Leu	Ile	Gln	Gln	Tyr	Tyr	Leu	Thr	Phe	Asn	Phe	Asp	Asn	Glu	Pro	Glu
		515					520						525		
Asn	Ile	Ser	Ile	Glu	Asn	Leu	Ser	Ser	Asp	Ile	Ile	Gly	Gln	Leu	Glu
	530					535					540				
Leu	Met	Pro	Asn	Ile	Glu	Arg	Phe	Pro	Asn	Gly	Lys	Lys	Tyr	Glu	Leu
545					550					555					560
Asp	Lys	Tyr	Thr	Met	Phe	His	Tyr	Leu	Arg	Ala	Gln	Glu	Phe	Glu	His
			565						570					575	
Gly	Lys	Ser	Arg	Ile	Ala	Leu	Thr	Asn	Ser	Val	Asn	Glu	Ala	Leu	Leu
			580						585					590	

-continued

---

Asn Pro Ser Arg Val Tyr Thr Phe Phe Ser Ser Asp Tyr Val Lys Lys  
 595 600 605

Val Asn Lys Ala Thr Glu Ala Ala Met Phe Leu Gly Trp Val Glu Gln  
 610 615 620

Leu Val Tyr Asp Phe Thr Asp Glu Thr Ser Glu Val Ser Thr Thr Asp  
 625 630 635 640

Lys Ile Ala Asp Ile Thr Ile Ile Ile Pro Tyr Ile Gly Pro Ala Leu  
 645 650 655

Asn Ile Gly Asn Met Leu Tyr Lys Asp Asp Phe Val Gly Ala Leu Ile  
 660 665 670

Phe Ser Gly Ala Val Ile Leu Leu Glu Phe Ile Pro Glu Ile Ala Ile  
 675 680 685

Pro Val Leu Gly Thr Phe Ala Leu Val Ser Tyr Ile Ala Asn Lys Val  
 690 695 700

Leu Thr Val Gln Thr Ile Asp Asn Ala Leu Ser Lys Arg Asn Glu Lys  
 705 710 715 720

Trp Asp Glu Val Tyr Lys Tyr Ile Val Thr Asn Trp Leu Ala Lys Val  
 725 730 735

Asn Thr Gln Ile Asp Leu Ile Arg Lys Lys Met Lys Glu Ala Leu Glu  
 740 745 750

Asn Gln Ala Glu Ala Thr Lys Ala Ile Ile Asn Tyr Gln Tyr Asn Gln  
 755 760 765

Tyr Thr Glu Glu Glu Lys Asn Asn Ile Asn Phe Asn Ile Asp Asp Leu  
 770 775 780

Ser Ser Lys Leu Asn Glu Ser Ile Asn Lys Ala Met Ile Asn Ile Asn  
 785 790 795 800

Lys Phe Leu Asn Gln Cys Ser Val Ser Tyr Leu Met Asn Ser Met Ile  
 805 810 815

Pro Tyr Gly Val Lys Arg Leu Glu Asp Phe Asp Ala Ser Leu Lys Asp  
 820 825 830

Ala Leu Leu Lys Tyr Ile Tyr Asp Asn Arg Gly Thr Leu Ile Gly Gln  
 835 840 845

Val Asp Arg Leu Lys Asp Lys Val Asn Asn Thr Leu Ser Thr Asp Ile  
 850 855 860

Pro Phe Gln Leu Ser Lys Tyr Val Asp Asn Gln Arg Leu Leu Ser Thr  
 865 870 875 880

His His His His His His  
 885

<210> SEQ ID NO 147  
 <211> LENGTH: 2661  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: LHn/A fragment

<400> SEQUENCE: 147

atgccgttcg taaacaaaca gttcaactat aaagaccag tcaacggcgt ggacattgcc 60  
 tatatcaaaa tccgaatgc gggcctcaatg cagcccgatg aagcatttaa aatccataac 120  
 aaaatttggg tgatcccga ggcgatatc ttcacgaacc cggaagaagg agatttaaac 180  
 ccaccgctg aggtctaaaca ggtcccgtg tcttactatg atagcacata cctgagtacc 240

-continued

---

gacaatgaaa aggacaacta cctgaaagggt gttaccaaac tggtcgagcg cattttattcg	300
acagatctcg gtcgcatggt gctgacttct attgtgctcg gcattccggt ttgggggtggt	360
agcaccatcg atacagaact caaagtgatt gacaccaact gcataaatgt gattcagcct	420
gatgggagct accggtccga agagcttaac ctcgtaatca ttggcccagag cgcggatatt	480
atccaattcg aatgtaaatc ttttgggcat gaagtcctga atctgacgcg gaatggctat	540
ggatcgacgc agtatattcg tttttctoca gatttcacat ttggatttga agaaagcctc	600
gaagttgata cgaaccctct tttaggcgcg ggaaaattcg cgacggacc cgcggtgacc	660
ttggcacatg aacttattca tgccgggcat cgcttgatg gaatgcgat taaccggaac	720
cgtgttttca aggtgaatac gaacgcgtat tacgagatgt cgggcttaga agtgccttt	780
gaagaactgc gcacgttttg cggctatgat gcaaaattta ttgatagtct gcaagaaaa	840
gaatttcggc tgtactatta caataaatc aaagacattg catcaacctt aaacaaggcg	900
aaaagcattg tgggtaccac ggctagctta caatatatga aaaacgtttt caaagaaaa	960
tacctctta gcgaagacac ttccggcaaa ttctctgtcg ataaactgaa atttgataaa	1020
ctgtataaaa tgcctaccga gatctacaca gaggataact ttgtcaaat cttcaaggtc	1080
ttgaatcggg aaacctatct gaacttcgat aaagccgtct ttaagatcaa catcgtaccg	1140
aaagttaact acaccatcta tgatggcttt aatctgctca atacgaatct ggcggcgacc	1200
tttaacggcc agaaccacca aatcaacaac atgaacttta ctaaactgaa aaattttacc	1260
ggcttgtttg aattctataa gctcctgtgt gtcccggtta ttatcaccag caaaacaaa	1320
tccttgggcg gtgggtggca aaacctgtac ttccaggcg gtggcggtgg tgataagggc	1380
tataacaagg ccttcaatga tttatgcac aaggtgaaca actgggactt gttttctct	1440
ccatctgaag ataattttac taacgacttg aacaaaggag aggaaattac ttccgatacc	1500
aacatcgaag cagcgggaag gaatattagt ctagatctta ttcaacaata ttacctgacc	1560
tttaattttg ataacgagcg tgagaacatt tccattgaga atctcagctc tgacatcatc	1620
ggccagctgg aactgatgcc gaatatacga cgctttccta atggaaagaa atatgaattg	1680
gacaaataca ccatgtcca ctatctccg gcgcaggagt ttgagcacgg caagtctcgt	1740
attgctctga ccaattcggg aaacgaagcc cttttaaact cttcgcgtgt gtacacctt	1800
ttctcaagcg attatgttaa aaaagtgaac aaggcgaccg aagcggcgat gtttttggga	1860
tgggtggaac aactggtata tgactttacg gatgaaactt ctgaagtctc gaccaccgac	1920
aaaattgccc atattaccat tatcattccc tatattggcc ctgcaactgaa catttgtaac	1980
atgctgtata aagatgattt tgtgggccc ctgatctttt caggcgtgt tatcctgctg	2040
gaatttatcc cggaaatcgc cattccagta ctcggtacct ttgcgctggt gtcctatac	2100
gcaacaaaag ttttgactgt ccagacgatc gacaacgcgc tcagtaaacg taacgaaaa	2160
tgggatgagg tgtataagta tattgttacc aactggctcg ctaaagtaaa caccagatt	2220
gacctgattc gcaagaagat gaaagaagcg ctggaaaacc aagcagaagc gaccaaagct	2280
attatcaact atcaatataa ccagtacaca gaggaagaaa agaataacat caacttcaac	2340
atcgacgact tatcttcaaa gctgaatgaa tctattaaca aagcgtatg taatattaac	2400
aagttcttga accaatgtag tgtcagctat ctgatgaact cgatgatccc ttacggtgtg	2460
aaacgtctgg aagacttcca tgcaagcctt aaagatgccc ttctgaagta tatttacgat	2520

---

-continued

---

aatcgcgga cttattgg ccaagtgat cgcttaaag ataaagcaa caacacgctg 2580  
agtacagaca tccctttca gctgtctaaa tatgtggaca atcagcgct gctgtccacg 2640  
caccatcacc atcaccacta a 2661

<210> SEQ ID NO 148  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: ORL-1 forward oligonucleotide primer  
  
<400> SEQUENCE: 148

cactcggctg gtgctggtgg 20

<210> SEQ ID NO 149  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: ORL-1 reverse oligonucleotide primer  
  
<400> SEQUENCE: 149

aatggccacg gcagtctcgc 20

<210> SEQ ID NO 150  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Galanin receptor 1 forward oligonucleotide  
primer  
  
<400> SEQUENCE: 150

ccccatcatg tcaccacct 20

<210> SEQ ID NO 151  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Galanin receptor 1 reverse oligonucleotide  
primer  
  
<400> SEQUENCE: 151

atggggttca cegaggagtt 20

<210> SEQ ID NO 152  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Galanin receptor 2 forward oligonucleotide  
primer  
  
<400> SEQUENCE: 152

catcgtggcg gtgctttt 18

<210> SEQ ID NO 153  
<211> LENGTH: 18

- continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Galanin receptor 2 reverse oligonucleotide
        primer

<400> SEQUENCE: 153
agcgggaagc gaccaaac

```

---

18

We claim:

1. A method of detecting retargeted endopeptidase activity, the method comprising the steps of:

- a) treating a cell from an established cell line with a sample comprising a retargeted endopeptidase, wherein the cell from an established cell line is susceptible to retargeted endopeptidase activity by a retargeted endopeptidase;
- b) isolating from the treated cell line a SNAP-25 component comprising a SNAP-25 cleavage product having a carboxyl-terminus at the P1 residue of the BoNT/A cleavage site scissile bond;
- c) contacting the SNAP-25 component with an  $\alpha$ -SNAP-25 antibody, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P1 residue of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product; and
- d) detecting the presence of an antibody-antigen complex comprising the  $\alpha$ -SNAP-25 antibody and the SNAP-25 cleavage product; wherein detection by the antibody-antigen complex is indicative of retargeted endopeptidase activity, wherein the  $\alpha$ -SNAP-25 antibody binds an epitope comprising a carboxyl-terminus at the P1 residue from the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product and wherein the isolated  $\alpha$ -SNAP-25 antibody has a  $V_H$  CDR1 of SEQ ID NO:93, a  $V_H$  CDR2 of SEQ ID NO:96, a  $V_H$  CDR3 of SEQ ID NO:100, a  $V_L$  CDR1 of SEQ ID NO:105, a  $V_L$  CDR2 of SEQ ID NO:110, and a  $V_L$  CDR3 of SEQ ID NO:115.

2. The method of claim 1, wherein the  $\alpha$ -SNAP-25 antibody is linked to a solid phase support.

3. The method of claim 2, comprising a step of fixing the SNAP-25 component to said solid phase support.

4. The method of claim 1, wherein the cell is susceptible to retargeted endopeptidase activity by about 500 nM or less, by about 400 nM or less, by about 300 nM or less, by about 200 nM or less, by about 100 nM or less of a retargeted endopeptidase.

5. The method of claim 1, wherein the cell can uptake about 500 nM or less, by about 400 nM or less, by about 300 nM or less, by about 200 nM or less, by about 100 nM or less of retargeted endopeptidase.

6. The method of claim 1, wherein the sample comprises about 100 ng or less, about 10 ng or less, about 1 ng or less, 100 fg or less, 10 fg or less, or 1 fg or less of a retargeted endopeptidase.

7. The method of claim 1, wherein the sample comprises about 100 nM or less, about 10 nM or less, about 1 nM or less, about 0.5 nM or less, or about 0.1 nM or less, of a retargeted endopeptidase.

8. The method of claim 1, wherein the presence of an antibody-antigen complex is detected by an immuno-blot analysis, an immunoprecipitation analysis, an ELISA, or a sandwich ELISA.

9. The method of claim 1, wherein the  $\alpha$ -SNAP-25 antibody has an association rate constant for an epitope not comprising a carboxyl-terminus glutamine of the BoNT/A cleavage site scissile bond from a SNAP-25 cleavage product of less than  $1 \times 10^1 \text{ M}^{-1} \text{ s}^{-1}$ ; and wherein the  $\alpha$ -SNAP-25 antibody has an equilibrium disassociation constant for the epitope of less than 0.450 nM.

10. An isolated monoclonal  $\alpha$ -SNAP-25 antibody which has a  $V_H$  CDR1 of SEQ ID NO:93, a  $V_H$  CDR2 of SEQ ID NO:96, a  $V_H$  CDR3 of SEQ ID NO:100, a  $V_L$  CDR1 of SEQ ID NO:105, a  $V_L$  CDR2 of SEQ ID NO:110, and a  $V_L$  CDR3 of SEQ ID NO:115.

\* \* \* \* \*

专利名称(译)	基于免疫的重定位内肽酶活性测定		
公开(公告)号	<a href="#">US20190324041A1</a>	公开(公告)日	2019-10-24
申请号	US16/504870	申请日	2019-07-08
[标]申请(专利权)人(译)	阿勒根公司		
申请(专利权)人(译)	ALLERGAN , INC.		
当前申请(专利权)人(译)	ALLERGAN , INC.		
[标]发明人	WANG JOANNE ZHU HONG HODGES DIANNE D FERNANDEZ SALAS ESTER		
发明人	WANG, JOANNE ZHU, HONG HODGES, DIANNE D. FERNANDEZ-SALAS, ESTER		
IPC分类号	G01N33/573 G01N33/50 C12N9/50 C07K16/18 G01N33/53 C12Q1/37		
CPC分类号	C12Q1/37 G01N2333/33 G01N33/53 C12N9/50 G01N33/573 C07K16/18 C07K2317/92 G01N2333/47 C07K2317/34 G01N2333/96425 G01N2333/952 G01N33/5035 C12Y304/24069 G01N33/5014 C07K2317/56 C12N9/52		
优先权	61/160217 2009-03-13 US		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

摘要(译)

本说明书公开了重靶向的内肽酶药物，其中活性已经通过所公开的方法确定。

