

**(12) United States Patent**
Styrkarsdottir et al.**(10) Patent No.: US 6,630,304 B1**
(45) Date of Patent: Oct. 7, 2003**(54) HUMAN OSTEOPOROSIS GENE****(75) Inventors: Unnur Styrkarsdottir, Reykjavik (IS);
Vala Drofn Johannsdottir, Reykjavik (IS)****(73) Assignee: deCODE genetics ehf., Reykjavik (IS)****(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.****(21) Appl. No.: 09/661,887****(22) Filed: Sep. 14, 2000****(51) Int. Cl.⁷ C12Q 1/68; C07H 19/00****(52) U.S. Cl. 435/6; 536/23.5****(58) Field of Search 435/6; 536/23.5****(56) References Cited**

U.S. PATENT DOCUMENTS

5,266,683 A 11/1993 Oppermann et al.
5,863,758 A 1/1999 Oppermann et al.

FOREIGN PATENT DOCUMENTS

WO WO 99/57261 11/1999
WO WO 00/04183 1/2000

OTHER PUBLICATIONS

Sambrook et al in *Molecular Cloning*, 1989, Cold Spring Harbor Laboratory, CSH, NY, Ch. 11 & 17.*Wallace et al in *Methods Enzymol* 152: 432-439, 1987.*Dale, L. and C.M. Jones, "BMP Signalling in Early Xenopus Development", *Bioessays*, 21(9):751-760 (1999).Zimmerman, L.B., et al., "The Spemann Organizer Signal Noggin Binds and Inactivates Bone Morphogenetic Protein 4", *Cell*, 86(4):599-606 (1996).Krisch, T., et al., "Crystal Structure of the BMP-2-BRIA Ectodomain Complex", *Nature Structural Biology*, 7(6):492-496 (2000).Massague, J., "TGF- β Signal Transduction", *Annu. Rev. Biochem.*, 67:753-791 (1998).Aspenberg, P., et al., "The Bone Morphogenetic Proteins Antagonist Noggin Inhibits Membranous Ossification", *J. Bone Miner Res.*, 16(3):497-500 (2001).Chen, D., et al., "Bone morphogenetic protein 2 (BMP-2) enhances BMP-3, BMP-4, and bone cell differentiation marker gene expression during the induction of mineralized bone matrix formation in cultures of fetal rat calvarial osteoblasts.", *Calcif. Tissue Int.*, 60:283-90 (1997).Chen, D., et al., "Differential roles for bone morphogenetic protein (BMP) receptor type IB and IA in differentiation and specification of mesenchymal precursor cells to osteoblast and adipocyte lineages.", *J. Cell Biol.*, 142:295-305 (1997).Fujii, M., et al., "Roles of bone morphogenetic protein type I receptors and Smad proteins in osteoblast and chondroblast differentiation." *Mol. Biol. Cell*, 10:3801-13 (1999).Miyazono, K., "Signal transduction by bone morphogenetic protein receptors: functional roles of Smad proteins." *Bone*, 25:91-3 (1999).Takazawa, Y., et al., "An osteogenesis-related transcription factor, core-binding factor A1, is constitutively expressed in the chondrocytic cell line TC6, and its expression is upregulated by bone morphogenetic protein-2." *J. Endocrinol.*, 165:579-586 (2000).Thirunavukkarasu, K., et al., "The osteoblast-specific transcription factor Cbfa1 regulates the expression of osteoprotegerin (OPG), a potent inhibitor of osteoclast differentiation and function.", *J. Biol. Chem.*, 275(33):25163-25172 (2000).Thompson, D.B., et al., "Linkage between stature and a region on chromosome 20 and analysis of a candidate gene, bone morphogenetic protein 2.", *Am. J. Med. Genet.*, 59:495-500 (1995).Sugiura, T., "Cloning and functional characterization of the 5'-flanking region of the human bone morphogenetic protein-2 gene.", *Biochem. J.*, 338:433-40 (1999).Wang, E.A., "Bone morphogenetic proteins (BMPs): therapeutic potential in healing bony defects." *Trends Biotechnol.*, 11:379-83 (1993).Wozney, J.M., et al., "Novel regulators of bone formation: molecular clones and activities." *Science*, 242:1528-34 (1988).Christiansen, M., et al., "CBFA1 and topoisomerase I mRNA levels decline during cellular aging of human trabecular osteoblasts," *J Gerontol A. Biol. Sci. Med. Sci.*, 55 (4): B194-200 (2000).

GenBank Seq. AL035668, Mar. 2001.

Helvering, L.M., et al., "Regulation of the Promoters for the Human Bone Morphogenetic Protein 2 and 4 Genes", *Gene*, 256:123-138 (2000).

Birren, B. et al., "Homo sapiens chromosome 14, clone RP11-19605," [online] Accession No. ac026511, Mar. 2000. Retrieved from EMBL Database.

Thompson, D.B. et al., "Linkage Between Stature and a Region on Chromosome 20 and Analysis of a Candidate Gene, Bone Morphogenetic Protein 2," *Am. J. of Med. Genet.* 59:595-500 (1995).

* cited by examiner

Primary Examiner—Christina Chan

Assistant Examiner—Phuong N. Huynh

(74) Attorney, Agent, or Firm—Hamilton, Brook, Smith & Reynolds, P.C.

(57) ABSTRACT

A role of the human BMP2 gene in osteoporosis is disclosed. Methods for diagnosis, prediction of clinical course and treatment for osteoporosis using polymorphisms in the BMP2 gene are also disclosed.

9 Claims, No Drawings

HUMAN OSTEOPOROSIS GENE**BACKGROUND OF THE INVENTION**

Osteoporosis is a debilitating disease characterized by low bone mass and deterioration of bone tissue, as defined by decreased bone mineral density (BMD). A direct result of the experienced microarchitectural deterioration is susceptibility to fractures and skeletal fragility, ultimately causing high mortality, morbidity and medical expenses worldwide. Postmenopausal women are at greater risk than others because the estrogen deficiency and corresponding decrease in bone mass experienced during menopause increase both the probability of osteoporotic fracture and the number of potential fracture sites. Yet aging women are not the only demographic group at risk. Young women who are malnourished, amenorrheic, or insufficiently active are at risk of inhibiting bone mass development at an early age. Furthermore, androgens play a role in the gain of bone mass during puberty, so elderly or hypogonadal men face the risk of osteoporosis if their bones were insufficiently developed.

The need to find a cure for this disease is complicated by the fact that there are many contributing factors that cause osteoporosis. Nutrition (particularly calcium, vitamin D and vitamin K intake), hormone levels, age, sex, race, body weight, activity level, and genetic factors all account for the variance seen in bone mineral density among individuals. Currently, the drugs approved to treat osteoporosis act as inhibitors of bone reabsorption, and include methods such as hormone replacement therapy (HRT), selective estrogen receptor modulators, calcitonin, and bisphosphonates. However, these treatments may not individually reduce risk with consistent results and while some therapies improve BMD when coadministered, others show no improvement or even lose their efficacy when used in combination. Clearly, as life expectancy increases and health and economic concerns of osteoporosis grow, a solution for the risks associated with this late-onset disease is in great demand. Early diagnosis of the disease or predisposition to the disease would be desirable.

SUMMARY OF THE INVENTION

As described herein, it has been discovered that polymorphisms in the gene for human bone morphogenetic protein 2 (BMP2) have been correlated through human linkage studies to a number of osteoporosis phenotypes. In particular, it has been discovered that one or more single nucleotide polymorphisms within the nucleotide sequence encoding the BMP2 gene product is correlated to osteoporosis. Accordingly, this invention pertains to an isolated nucleic acid molecule containing the BMP2 gene of SEQ ID NO:1 (Table 1) having at least one altered nucleotide and to gene products encoded thereby (referred to herein as a "variant BMP2 gene" or "variant BMP2 gene product").

A number of polymorphisms have been observed in the BMP2 gene, as follows:

Promoter Region:

- A to G at nucleotide position 420;
- A to G at nucleotide position 472;
- G to C at nucleotide position 1464;
- G to A at nucleotide position 1722;
- C to A at nucleotide position 1914;

Coding Region:

- T to G at nucleotide position 3747; resulting in an amino acid change from serine to alanine at amino acid position 37;

A to G at nucleotide position 3899; no amino acid change; G to T at nucleotide position 3918; resulting in an amino acid change from alanine to serine at amino acid position 94;

A to T at nucleotide position 11980; resulting in an amino acid change from arginine to serine at amino acid position 189;

3' UTR and Downstream Region:

C to T at nucleotide position 12571;

T to C at nucleotide position 13066;

A to G at nucleotide position 13209;

C to A at nucleotide position 13296; and

at least one deletion in nucleotides at positions 13533–13536.

All numbering is relative to SEQ ID NO. 1. Thus, in preferred embodiments, the isolated nucleic acid molecule of the invention can have one or a combination of these single nucleotide polymorphisms. These polymorphisms may be part of a group of other polymorphisms in the BMP2 gene which contributes to the presence, absence or severity of osteoporosis.

The invention also relates to DNA constructs comprising the nucleic acid molecules described herein operatively linked to a regulatory sequence, and to recombinant host cells, such as bacterial cells, fungal cells, plant cells, insect cells and mammalian cells, comprising the nucleic acid molecules described herein operatively linked to a regulatory sequence.

The invention also pertains to methods of diagnosing osteoporosis in an individual. The methods include detecting the presence of a mutation in the BMP2 gene. The invention additionally pertains to pharmaceutical compositions comprising the BMP2 nucleic acids of the invention and to kits for carrying out the methods described herein. The methods of the invention allow the accurate diagnosis of osteoporosis at or before disease onset, thus reducing or minimizing the debilitating effects of osteoporosis.

DETAILED DESCRIPTION OF THE INVENTION

As described herein, Applicants have completed a genome wide scan on patients with various forms of osteoporosis and identified a region on chromosome 20 linked to osteoporosis. Until now there have been no known linkage studies of osteoporosis in humans showing any connection to this region of the chromosome. Based on the linkage studies conducted, Applicants have discovered a direct relationship between BMP2 and osteoporosis. Although the BMP2 gene from normal individuals is known, there have been no studies directly investigating BMP2 and osteoporosis. Moreover, there have been no variant forms reported that have been associated with osteoporosis. The linkage studies are based on four genome wide scans encompassing affected persons having different osteoporosis phenotypes; i.e., hip, spine, combined and combined severe (e.g., patients having vertebral compression fracture, hip fracture, other osteoporosis related low impact fracture). From the data obtained in the linkage study, a region on chromosome 20, specifically the BMP2 gene, was identified. The variant BMP2 gene has previously unreported nucleotide changes that were observed in the patient population, as follows:

Promoter Region:

A to G at nucleotide position 420;

A to G at nucleotide position 472;

G to C at nucleotide position 1464;

G to A at nucleotide position 1722;

C to G at nucleotide position 1914;

Coding Region:

T to G at nucleotide position 3747; resulting in an amino acid change from serine to alanine at amino acid position 37;

A to G at nucleotide position 3899; no amino acid change;

G to T at nucleotide position 3918; resulting in an amino acid change from alanine to serine at amino acid position 94;

A to T at nucleotide position 11980; resulting in an amino acid change from arginine to serine at amino acid position 189;

3' UTR and Downstream Region:

C to T at nucleotide position 12571;

T to C at nucleotide position 13066;

A to G at nucleotide position 13209;

C to A at nucleotide position 13296; and

at least one deletion in nucleotides at positions 13533–13536.

All nucleotide positions are relative to SEQ ID NO: 1. The polymorphism at nucleotide position 3747 appears statistically more frequent in the osteoporosis test population than in the control population.

NUCLEIC ACIDS OF THE INVENTION

Accordingly, the invention pertains to an isolated nucleic acid molecule comprising the human BMP2 gene having at least one nucleotide alteration and correlated with incidence of osteoporosis. The term, “variant BMP2”, as used herein, refers to an isolated nucleic acid molecule in chromosome 20 having at least one altered nucleotide that is associated with a susceptibility to a number of osteoporosis phenotypes, and also to a portion or fragment of the isolated nucleic acid molecule containing the alteration (e.g., cDNA or the gene) and encoding a variant BMP2 polypeptide (e.g., the polypeptide having SEQ ID NO: 2, as shown in Table 1). In a preferred embodiment, the isolated nucleic acid molecules comprises a polymorphism selected from the group consisting of: A to G at nucleotide position 420; A to G at nucleotide position 472; G to C at nucleotide position 1464; G to A at nucleotide position 1722; C to G at nucleotide position 1914; T to G at nucleotide position 3747, resulting in an amino acid change from serine to alanine at amino acid position 37; A to G at nucleotide position 3899, no amino acid change; G to T at nucleotide position 3918, resulting in an amino acid change from alanine to serine at amino acid position 94; A to T at nucleotide position 11980, resulting in an amino acid change from arginine to serine at amino acid position 189; C to T at nucleotide position 12571; T to C at nucleotide position 13066; A to G at nucleotide position 13209; C to A at nucleotide position 13296; at least one deletion in nucleotides at positions 13533–13536 and combinations thereof, of the BMP2 gene.

The isolated nucleic acid molecules of the present invention can be RNA, for example, mRNA, or DNA, such as cDNA and genomic DNA. DNA molecules can be double-stranded or single-stranded; single stranded RNA or DNA can be either the coding, or sense, strand or the non-coding, or antisense, strand. The nucleic acid molecule can include all or a portion of the coding sequence of the gene and can further comprise additional non-coding sequences such as introns and non-coding 3' and 5' sequences (including regulatory sequences, for example). Additionally, the nucleic

acid molecule can be fused to a marker sequence, for example, a sequence that encodes a polypeptide to assist in isolation or purification of the polypeptide. Such sequences include, but are not limited to, those which encode a glutathione-S-transferase (GST) fusion protein and those which encode a hemagglutinin A (HA) polypeptide marker from *influenza*.

An “isolated” nucleic acid molecule, as used herein, is one that is separated from nucleic acids which normally flank the gene or nucleotide sequence (as in genomic sequences) and/or has been completely or partially purified from other transcribed sequences (e.g., as in an RNA library). For example, an isolated nucleic acid of the invention may be substantially isolated with respect to the complex cellular milieu in which it naturally occurs, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. In some instances, the isolated material will form part of a composition (for example, a crude extract containing other substances), buffer system or reagent mix. In other circumstances, the material may be purified to essential homogeneity, for example as determined by PAGE or column chromatography such as HPLC. Preferably, an isolated nucleic acid molecule comprises at least about 50, 80 or 90% (on a molar basis) of all macromolecular species present. With regard to genomic DNA, the term “isolated” also can refer to nucleic acid molecules which are separated from the chromosome with which the genomic DNA is naturally associated. For example, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotides which flank the nucleic acid molecule in the genomic DNA of the cell from which the nucleic acid molecule is derived.

The nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated. Thus, recombinant DNA contained in a vector is included in the definition of “isolated” as used herein. Also, isolated nucleic acid molecules include recombinant DNA molecules in heterologous host cells, as well as partially or substantially purified DNA molecules in solution. “Isolated” nucleic acid molecules also encompass *in vivo* and *in vitro* RNA transcripts of the DNA molecules of the present invention. An isolated nucleic acid molecule or nucleotide sequence can include a nucleic acid molecule or nucleotide sequence which is synthesized chemically or by recombinant means. Therefore, recombinant DNA contained in a vector are included in the definition of “isolated” as used herein. Also, isolated nucleotide sequences include recombinant DNA molecules in heterologous organisms, as well as partially or substantially purified DNA molecules in solution. *In vivo* and *in vitro* RNA transcripts of the DNA molecules of the present invention are also encompassed by “isolated” nucleotide sequences. Such isolated nucleotide sequences are useful in the manufacture of the encoded polypeptide, as probes for isolating homologous sequences (e.g., from other mammalian species), for gene mapping (e.g., by *in situ* hybridization with chromosomes), or for detecting expression of the gene in tissue (e.g., human tissue), such as by Northern blot analysis.

The present invention also pertains to nucleic acid molecules which are not necessarily found in nature but which encode a BMP2 polypeptide (e.g., a polypeptide having the amino acid sequence of SEQ ID NO: 2). Thus, for example, DNA molecules which comprise a sequence that is different from the naturally-occurring nucleotide sequence but which, due to the degeneracy of the genetic code, encode an BMP2 polypeptide of the present invention are also the subject of

this invention. The invention also encompasses variants of the nucleotide sequences of the invention, such as those encoding portions, analogues or derivatives of the BMP2 polypeptide. Such variants can be naturally-occurring, such as in the case of allelic variation, or non-naturally-occurring, such as those induced by various mutagens and mutagenic processes. Intended variations include, but are not limited to, addition, deletion and substitution of one or more nucleotides which can result in conservative or non-conservative amino acid changes, including additions and deletions. Preferably the nucleotide (and/or resultant amino acid) changes are silent or conserved; that is, they do not alter the characteristics or activity of the BMP2 polypeptide.

Other alterations of the nucleic acid molecules of the invention can include, for example, labeling, methylation, internucleotide modifications such as uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates), charged linkages (e.g., phosphorothioates, phosphorodithioates), pendent moieties (e.g., polypeptides), intercalators (e.g., acridine, psoralen), chelators, alkylators, and modified linkages (e.g., alpha anomeric nucleic acids). Also included are synthetic molecules that mimic nucleic acid molecules in the ability to bind to a designated sequences via hydrogen bonding and other chemical interactions. Such molecules include, for example, those in which peptide linkages substitute for phosphate linkages in the backbone of the molecule.

The invention also pertains to nucleic acid molecules which hybridize under high stringency hybridization conditions, such as for selective hybridization, to a nucleotide sequence described herein (e.g., nucleic acid molecules which specifically hybridize to a nucleotide sequence encoding polypeptides described herein, and, optionally, have an activity of the polypeptide). In one embodiment, the invention includes variants described herein which hybridize under high stringency hybridization conditions (e.g., for selective hybridization) to a nucleotide sequence comprising a nucleotide sequence selected from SEQ ID NO: 1 or the complement of SEQ ID NO: 1.

Such nucleic acid molecules can be detected and/or isolated by specific hybridization (e.g., under high stringency conditions). "Stringency conditions" for hybridization is a term of art which refers to the incubation and wash conditions, e.g., conditions of temperature and buffer concentration, which permit hybridization of a particular nucleic acid to a second nucleic acid; the first nucleic acid may be perfectly (i.e., 100%) complementary to the second, or the first and second may share some degree of complementarity which is less than perfect (e.g., 70%, 75%, 85%, 95%). For example, certain high stringency conditions can be used which distinguish perfectly complementary nucleic acids from those of less complementarity. "High stringency conditions", "moderate stringency conditions" and "low stringency conditions" for nucleic acid hybridizations are explained on pages 2.10.1-2.10.16 and pages 6.3.1-6.3.6 in *Current Protocols in Molecular Biology* (Ausubel, F. M. et al., "Current Protocols in Molecular Biology", John Wiley & Sons, (1998), the entire teachings of which are incorporated by reference herein). The exact conditions which determine the stringency of hybridization depend not only on ionic strength (e.g., 0.2xSSC, 0.1xSSC), temperature (e.g., room temperature, 42° C., 68° C.) and the concentration of destabilizing agents such as formamide or denaturing agents such as SDS, but also on factors such as the length of the nucleic acid sequence, base composition, percent mismatch between hybridizing sequences and the frequency of occurrence of subsets of that sequence within other non-

identical sequences. Thus, equivalent conditions can be determined by varying one or more of these parameters while maintaining a similar degree of identity or similarity between the two nucleic acid molecules. Typically, conditions are used such that sequences at least about 60%, at least about 70%, at least about 80%, at least about 90% or at least about 95% or more identical to each other remain hybridized to one another. By varying hybridization conditions from a level of stringency at which no hybridization occurs to a level at which hybridization is first observed, conditions which will allow a given sequence to hybridize (e.g., selectively) with the most similar sequences in the sample can be determined.

Exemplary conditions are described in Krause, M. H. and S. A. Aaronson, *Methods in Enzymology*, 200:546-556 (1991). Also, in Ausubel, et al., "Current Protocols in Molecular Biology", John Wiley & Sons, (1998), which describes the determination of washing conditions for moderate or low stringency conditions. Washing is the step in which conditions are usually set so as to determine a minimum level of complementarity of the hybrids. Generally, starting from the lowest temperature at which only homologous hybridization occurs, each ° C. by which the final wash temperature is reduced (holding SSC concentration constant) allows an increase by 1% in the maximum extent of mismatching among the sequences that hybridize. Generally, doubling the concentration of SSC results in an increase in T_m of ~17° C. Using these guidelines, the washing temperature can be determined empirically for high, moderate or low stringency, depending on the level of mismatch sought.

For example, a low stringency wash can comprise washing in a solution containing 0.2xSSC/0.1% SDS for 10 min at room temperature; a moderate stringency wash can comprise washing in a prewarmed solution (42° C.) solution containing 0.2xSSC/0.1% SDS for 15 min at 42° C.; and a high stringency wash can comprise washing in prewarmed (68° C.) solution containing 0.1xSSC/0.1% SDS for 15 min at 68° C. Furthermore, washes can be performed repeatedly or sequentially to obtain a desired result as known in the art. Equivalent conditions can be determined by varying one or more of the parameters given as an example, as known in the art, while maintaining a similar degree of identity or similarity between the target nucleic acid molecule and the primer or probe used.

The percent identity of two nucleotide or amino acid sequences can be determined by aligning the sequences for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first sequence). The nucleotides or amino acids at corresponding positions are then compared, and the percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity=# of identical positions/total # of positionsx100). In certain embodiments, the length of a sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 60%, and even more preferably at least 70%, 80% or 90% of the length of the reference sequence. The actual comparison of the two sequences can be accomplished by well-known methods, for example, using a mathematical algorithm. A preferred, non-limiting example of such a mathematical algorithm is described in Karlin et al., *Proc. Natl. Acad. Sci. USA*, 90:5873-5877 (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) as described in Altschul et al., *Nucleic Acids Res.*, 25:389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs

(e.g., NBLAST) can be used. In one embodiment, parameters for sequence comparison can be set at score=100, wordlength=12, or can be varied (e.g., W=5 or W=20).

Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package (Accelrys, Cambridge, UK). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art and include ADVANCE and ADAM as described in Torellis and Robotti (1994) *Comput. Appl. Biosci.*, 10:3-5; and FASTA described in Pearson and Lipman (1988) *PNAS*, 85:2444-8.

In another embodiment, the percent identity between two amino acid sequences can be accomplished using the GAP program in the CGC software package using either a Blossom 63 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. In yet another embodiment, the percent identity between two nucleic acid sequences can be accomplished using the GAP program in the CGC software package, using a gap weight of 50 and a length weight of 3.

The present invention also provides isolated nucleic acid molecules that contain a fragment or portion that hybridizes under highly stringent conditions to a nucleotide sequence comprising a nucleotide sequence selected from SEQ ID NO: 1 and the complement of SEQ ID NO: 1. The nucleic acid fragments of the invention are at least about 15, preferably at least about 18, 20, 23 or 25 nucleotides, and can be 30, 40, 50, 100, 200 or more nucleotides in length. Longer fragments, for example, 30 or more nucleotides in length, which encode antigenic polypeptides described herein are particularly useful, such as for the generation of antibodies as described below.

In a related aspect, the nucleic acid fragments of the invention are used as probes or primers in assays such as those described herein. "Probes" are oligonucleotides that hybridize in a base-specific manner to a complementary strand of nucleic acid molecules. Such probes include polypeptide nucleic acids, as described in Nielsen et al., *Science*, 254, 1497-1500 (1991). Typically, a probe comprises a region of nucleotide sequence that hybridizes under highly stringent conditions to at least about 15, typically about 20-25, and more typically about 40, 50 or 75, consecutive nucleotides of a nucleic acid molecule comprising a nucleotide sequence selected from SEQ ID NO: 1 and the complement of SEQ ID NO: 1. More typically, the probe further comprises a label, e.g., radioisotope, fluorescent compound, enzyme, or enzyme co-factor.

As used herein, the term "primer" refers to a single-stranded oligonucleotide which acts as a point of initiation of template-directed DNA synthesis using well-known methods (e.g., PCR, LCR) including, but not limited to those described herein. The appropriate length of the primer depends on the particular use, but typically ranges from about 15 to 30 nucleotides.

The nucleic acid molecules of the invention such as those described above can be identified and isolated using standard molecular biology techniques and the sequence information provided in SEQ ID NO: 1. For example, nucleic acid molecules can be amplified and isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based on one or more of the sequences provided

in SEQ ID NO: 1 and/or the complement of SEQ ID NO: 1. See generally *PCR Technology: Principles and Applications for DNA Amplification* (ed. H. A. Erlich, Freeman Press, NY, N.Y., 1992); *PCR Protocols: A Guide to Methods and Applications* (Eds. Innis, et al., Academic Press, San Diego, Calif., 1990); Mattila et al., *Nucleic Acids Res.*, 19:4967 (1991); Eckert et al., *PCR Methods and Applications*, 1:17 (1991); PCR (eds. McPherson et al., IRL Press, Oxford); and U.S. Pat. No. 4,683,202. The nucleic acid molecules can be amplified using cDNA, mRNA or genomic DNA as a template, cloned into an appropriate vector and characterized by DNA sequence analysis.

Other suitable amplification methods include the ligase chain reaction (LCR) (see Wu and Wallace, *Genomics*, 4:560 (1989), Landegren et al., *Science*, 241:1077 (1988), transcription amplification (Kwoh et al., *Proc. Natl. Acad. Sci. USA*, 86:1173 (1989)), and self-sustained sequence replication (Guatelli et al., *Proc. Nat. Acad. Sci. USA*, 87:1874 (1990)) and nucleic acid based sequence amplification (NASBA). The latter two amplification methods involve isothermal reactions based on isothermal transcription, which produce both single stranded RNA (ssRNA) and double stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

The amplified DNA can be radiolabelled and used as a probe for screening a cDNA library derived from human cells, mRNA in zap express, ZIPLOX or other suitable vector. Corresponding clones can be isolated, DNA can be obtained following in vivo excision, and the cloned insert can be sequenced in either or both orientations by art recognized methods to identify the correct reading frame encoding a polypeptide of the appropriate molecular weight. For example, the direct analysis of the nucleotide sequence of nucleic acid molecules of the present invention can be accomplished using well-known methods that are commercially available. See, for example, Sambrook et al., *Molecular Cloning, A Laboratory Manual* (2nd Ed., CSHP, New York 1989); Zyskind et al., *Recombinant DNA Laboratory Manual*, (Acad. Press, 1988)). Using these or similar methods, the polypeptide and the DNA encoding the polypeptide can be isolated, sequenced and further characterized.

Antisense nucleic acid molecules of the invention can be designed using the nucleotide sequences of SEQ ID NO: 1 and/or the complement of SEQ ID NO: 1, and/or a portion of SEQ ID NO: 1 or the complement of SEQ ID NO: 1, and constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid molecule (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Alternatively, the antisense nucleic acid molecule can be produced biologically using an expression vector into which a nucleic acid molecule has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid molecule will be of an antisense orientation to a target nucleic acid of interest).

In general, the isolated nucleic acid sequences of the invention can be used as molecular weight markers on Southern gels, and as chromosome markers which are labeled to map related gene positions. The nucleic acid sequences can also be used to compare with endogenous DNA sequences in patients to identify genetic disorders

(e.g., a predisposition for or susceptibility to osteoporosis), and as probes, such as to hybridize and discover related DNA sequences or to subtract out known sequences from a sample. The nucleic acid sequences can further be used to derive primers for genetic fingerprinting, to raise anti-polypeptide antibodies using DNA immunization techniques, and as an antigen to raise anti-DNA antibodies or elicit immune responses. Portions or fragments of the nucleotide sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. Additionally, the nucleotide sequences of the invention can be used to identify and express recombinant polypeptides for analysis, characterization or therapeutic use, or as markers for tissues in which the corresponding polypeptide is expressed, either constitutively, during tissue differentiation, or in diseased states.

Another aspect of the invention pertains to nucleic acid constructs containing a nucleic acid molecule selected from the group consisting of SEQ ID NO: 1 and the complement of SEQ ID NO: 1 (or a portion thereof). The constructs comprise a vector (e.g., an expression vector) into which a sequence of the invention has been inserted in a sense or antisense orientation. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors, expression vectors, are capable of directing the expression of genes to which they are operably linked. In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses) that serve equivalent functions.

Preferred recombinant expression vectors of the invention comprise a nucleic acid molecule of the invention in a form suitable for expression of the nucleic acid molecule in a host cell. This means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably or operatively linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (e.g., in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, *Gene Expression*

Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed and the level of expression of polypeptide desired. The expression vectors of the invention can be introduced into host cells to thereby produce polypeptides, including fusion polypeptides, encoded by nucleic acid molecules as described herein.

The recombinant expression vectors of the invention can be designed for expression of a polypeptide of the invention in prokaryotic or eukaryotic cells, e.g., bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors), yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, supra. Alternatively, the recombinant expression vector can be transcribed and translated in vitro, for example using T7 promoter regulatory sequences and T7 polymerase.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, a nucleic acid molecule of the invention can be expressed in bacterial cells (e.g., *E. coli*), insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing a foreign nucleic acid molecule (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, et al. (supra), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., for resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid molecules encoding a selectable marker can be introduced into a host cell on the same vector as the nucleic acid molecule of the invention or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid molecule can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (i.e., express) a polypeptide of the invention. Accordingly, the invention further provides methods for producing a polypeptide using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding a polypeptide of the invention has been introduced) in a suitable medium such that the polypeptide is produced. In another embodiment, the method further comprises isolating the polypeptide from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which a nucleic acid molecule of the invention has been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous nucleotide sequences have been introduced into the genome or homologous recombinant animals in which endogenous nucleotide sequences have been altered. Such animals are useful for studying the function and/or activity of the nucleotide sequence and polypeptide encoded by the sequence and for identifying and/or evaluating modulators of their activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens and amphibians. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, an "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009, U.S. Pat. No. 4,873,191 and in Hogan, *Manipulating the Mouse Embryo* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley (1991) *Current Opinion in Bio/Technology*, 2:823-829 and in PCT Publication Nos. WO 90/11354, WO 91/01140, WO 92/0968, and WO 93/04169. Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut et al. (1997) *Nature*, 385:810-813 and PCT Publication Nos. WO 97/07668 and WO 97/07669.

POLYPEPTIDES OF THE INVENTION

The present invention also pertains to isolated BMP2 polypeptides, e.g., proteins, and variants thereof, as well as polypeptides encoded by nucleotide sequences described herein (e.g., other splicing variants). The term "polypeptide" refers to a polymer of amino acids, and not to a specific length; thus, peptides, oligopeptides and proteins are included within the definition of a polypeptide. As used

herein, a polypeptide is said to be "isolated" or "purified" when it is substantially free of cellular material when it is isolated from recombinant and non-recombinant cells, or free of chemical precursors or other chemicals when it is chemically synthesized. A polypeptide, however, can be joined to another polypeptide with which it is not normally associated in a cell and still be "isolated" or "purified."

The polypeptides of the invention can be purified to homogeneity. It is understood, however, that preparations in which the polypeptide is not purified to homogeneity are useful. The critical feature is that the preparation allows for the desired function of the polypeptide, even in the presence of considerable amounts of other components. Thus, the invention encompasses various degrees of purity. In one embodiment, the language "substantially free of cellular material" includes preparations of the polypeptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins.

When a polypeptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20%, less than about 10%, or less than about 5% of the volume of the polypeptide preparation. The language "substantially free of chemical precursors or other chemicals" includes preparations of the polypeptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the polypeptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

In one embodiment, a polypeptide comprises an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 1 and complements and portions thereof, e.g., SEQ ID NO: 2, or a portion of SEQ ID NO: 2. However, the invention also encompasses sequence variants. Variants include a substantially homologous polypeptide encoded by the same genetic locus in an organism, i.e., an allelic variant, as well as other splicing variants. Variants also encompass polypeptides derived from other genetic loci in an organism, but having substantial homology to a polypeptide encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 1 and complements and portions thereof. Variants also include polypeptides substantially homologous or identical to these polypeptides but derived from another organism, i.e., an ortholog. Variants also include polypeptides that are substantially homologous or identical to these polypeptides that are produced by chemical synthesis. Variants also include polypeptides that are substantially homologous or identical to these polypeptides that are produced by recombinant methods.

As used herein, two polypeptides (or a region of the polypeptides) are substantially homologous or identical when the amino acid sequences are at least about 45-55%, typically at least about 70-75%, more typically at least about 80-85%, and most typically greater than about 90% or more homologous or identical. A substantially homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid molecule hybridizing to SEQ ID NO: 1, or portion thereof, under stringent conditions as more particularly described above.

To determine the percent homology or identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of one polypeptide or nucleic acid molecule for optimal alignment with the other polypeptide or nucleic acid molecule). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in one sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the other sequence, then the molecules are homologous at that position. As used herein, amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity". The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (i.e., percent homology equals the number of identical positions/total number of positions times 100).

The invention also encompasses polypeptides having a lower degree of identity but having sufficient similarity so as to perform one or more of the same functions performed by a polypeptide encoded by a nucleic acid molecule of the invention. Similarity is determined by conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Conservative substitutions are likely to be phenotypically silent. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie et al., *Science* 247:1306-1310 (1990).

A variant polypeptide can differ in amino acid sequence by one or more substitutions, deletions, insertions, inversions, fusions, and truncations or a combination of any of these. Further, variant polypeptides can be fully functional or can lack function in one or more activities. Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree. Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham et al., *Science*, 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity in vitro, or in vitro proliferative activity. Sites that are critical for polypeptide activity can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith et al., *J. Mol. Biol.*, 224:899-904 (1992); de Vos et al. *Science*, 255:306-312 (1992)).

The invention also includes polypeptide fragments of the polypeptides of the invention. Fragments can be derived from a polypeptide encoded by a nucleic acid molecule comprising SEQ ID NO: 1 or a portion thereof and the

complements thereof. However, the invention also encompasses fragments of the variants of the polypeptides described herein. As used herein, a fragment comprises at least 6 contiguous amino acids. Useful fragments include those that retain one or more of the biological activities of the polypeptide as well as fragments that can be used as an immunogen to generate polypeptide-specific antibodies.

Biologically active fragments (peptides which are, for example, 6, 9, 12, 15, 16, 20, 30, 35, 36, 37, 38, 39, 40, 50, 100 or more amino acids in length) can comprise a domain, segment, or motif that has been identified by analysis of the polypeptide sequence using well-known methods, e.g., signal peptides, extracellular domains, one or more transmembrane segments or loops, ligand binding regions, zinc finger domains, DNA binding domains, acylation sites, glycosylation sites, or phosphorylation sites.

Fragments can be discrete (not fused to other amino acids or polypeptides) or can be within a larger polypeptide. Further, several fragments can be comprised within a single larger polypeptide. In one embodiment a fragment designed for expression in a host can have heterologous pre- and pro-polypeptide regions fused to the amino terminus of the polypeptide fragment and an additional region fused to the carboxyl terminus of the fragment.

The invention thus provides chimeric or fusion polypeptides. These comprise a polypeptide of the invention operatively linked to a heterologous protein or polypeptide having an amino acid sequence not substantially homologous to the polypeptide. "Operatively linked" indicates that the polypeptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the polypeptide. In one embodiment the fusion polypeptide does not affect function of the polypeptide per se. For example, the fusion polypeptide can be a GST-fusion polypeptide in which the polypeptide sequences are fused to the C-terminus of the GST sequences. Other types of fusion polypeptides include, but are not limited to, enzymatic fusion polypeptides, for example β -galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions and Ig fusions. Such fusion polypeptides, particularly poly-His fusions, can facilitate the purification of recombinant polypeptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a polypeptide can be increased by using a heterologous signal sequence. Therefore, in another embodiment, the fusion polypeptide contains a heterologous signal sequence at its N-terminus.

EP-A-O 464 533 discloses fusion proteins comprising various portions of immunoglobulin constant regions. The Fc is useful in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). In drug discovery, for example, human proteins have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists. Bennett et al., *Journal of Molecular Recognition*, 8:52-58 (1995) and Johanson et al., *The Journal of Biological Chemistry*, 270,16:9459-9471 (1995). Thus, this invention also encompasses soluble fusion polypeptides containing a polypeptide of the invention and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclass (IgG, IgM, IgA, IgE).

A chimeric or fusion polypeptide can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including auto-

mated DNA synthesizers. Alternatively, PCR amplification of nucleic acid fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive nucleic acid fragments which can subsequently be annealed and re-amplified to generate a chimeric nucleic acid sequence (see Ausubel et al., *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A nucleic acid molecule encoding a polypeptide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the polypeptide.

The isolated polypeptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. In one embodiment, the polypeptide is produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the polypeptide is cloned into an expression vector, the expression vector introduced into a host cell and the polypeptide expressed in the host cell. The polypeptide can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques.

In general, polypeptides of the present invention can be used as a molecular weight marker on SDS-PAGE gels or on molecular sieve gel filtration columns using art-recognized methods. The polypeptides of the present invention can be used to raise antibodies or to elicit an immune response. The polypeptides can also be used as a reagent, e.g., a labeled reagent, in assays to quantitatively determine levels of the polypeptide or a molecule to which it binds (e.g., a receptor or a ligand) in biological fluids. The polypeptides can also be used as markers for cells or tissues in which the corresponding polypeptide is preferentially expressed, either constitutively, during tissue differentiation, or in a diseased state. The polypeptides can be used to isolate a corresponding binding partner, e.g., receptor or ligand, such as, for example, in an interaction trap assay, and to screen for peptide or small molecule antagonists or agonists of the binding interaction.

ANTIBODIES OF THE INVENTION

Polyclonal and/or monoclonal antibodies that specifically bind one form of the gene product but not to the other form of the gene product are also provided. Antibodies are also provided that bind a portion of either the variant or the reference gene product that contains the polymorphic site or sites. The invention provides antibodies to the polypeptides and polypeptide fragments of the invention, e.g., having an amino acid sequence encoded by SEQ ID NO: 2, or a portion thereof, or having an amino acid sequence encoded by a nucleic acid molecule comprising all or a portion of SEQ ID NO: 1. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that specifically binds an antigen. A molecule that specifically binds to a polypeptide of the invention is a molecule that binds to that polypeptide or a fragment thereof, but does not substantially bind other molecules in a sample, e.g., a biological sample, which naturally contains the polypeptide. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab)₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind to a polypeptide of the invention. The term "monoclonal antibody" or "monoclonal antibody

composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of a polypeptide of the invention. A monoclonal antibody composition thus typically displays a single binding affinity for a particular polypeptide of the invention with which it immunoreacts.

Polyclonal antibodies can be prepared as described above by immunizing a suitable subject with a desired immunogen, e.g., polypeptide of the invention or fragment thereof. The antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized polypeptide. If desired, the antibody molecules directed against the polypeptide can be isolated from the mammal (e.g., from the blood) and further purified by well-known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature*, 256:495-497, the human B cell hybridoma technique (Kozbor et al. (1983) *Immunol. Today*, 4:72), the EBV-hybridoma technique (Cole et al. (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing hybridomas is well known (see generally *Current Protocols in Immunology* (1994) Coligan et al. (eds.) John Wiley & Sons, Inc., New York, N.Y.). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with an immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds a polypeptide of the invention.

Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating a monoclonal antibody to a polypeptide of the invention (see, e.g., *Current Protocols in Immunology*, supra; Galfre et al. (1977) *Nature*, 266:55052; R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, N.Y. (1980); and Lerner (1981) *Yale J. Biol. Med.*, 54:387-402. Moreover, the ordinarily skilled worker will appreciate that there are many variations of such methods that also would be useful.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal antibody to a polypeptide of the invention can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with the polypeptide to thereby isolate immunoglobulin library members that bind the polypeptide. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene SurfZAP™ Phage Display Kit, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, U.S. Pat. No. 5,223,409; PCT Publication No. WO 92/18619; PCT Publication No. WO 91/17271; PCT Publication No. WO 92/20791; PCT Publication No. WO 92/15679; PCT Publication No. WO 93/01288; PCT Publication No. WO 92/01047; PCT Publication No. WO 92/09690; PCT Publication No. WO 90/02809; Fuchs et al. (1991) *BioTechnology*, 9:1370-1372; Hay et al. (1992)

Hum. Antibod. Hybridomas, 3:81-85; Huse et al. (1989) *Science*, 246:1275-1281; Griffiths et al. (1993) *EMBO J.*, 12:725-734.

Additionally, recombinant antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art.

In general, antibodies of the invention (e.g., a monoclonal antibody) can be used to isolate a polypeptide of the invention by standard techniques, such as affinity chromatography or immunoprecipitation. A polypeptide-specific antibody can facilitate the purification of natural polypeptide from cells and of recombinantly produced polypeptide expressed in host cells. Moreover, an antibody specific for a polypeptide of the invention can be used to detect the polypeptide (e.g., in a cellular lysate, cell supernatant, or tissue sample) in order to evaluate the abundance and pattern of expression of the polypeptide. Antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ¹²⁵I, ¹³¹I, ¹³¹S or ³H.

DIAGNOSTIC AND SCREENING ASSAYS OF THE INVENTION

The present invention also pertains to a method of diagnosing or aiding in the diagnosis of osteoporosis associated with the presence of the BMP2 gene or gene product in an individual. Diagnostic assays can be designed for determining BMP2 polypeptide and/or nucleic acid expression as well as activity of polypeptides of the invention, in the context of a biological sample (e.g., blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with osteoporosis, or is at risk for (has a predisposition for or a susceptibility to) developing osteoporosis. The invention also provides for prognostic (or predictive) assays for determining whether an individual is susceptible to developing osteoporosis. For example, mutations in the gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of symptoms associated with osteoporosis. Another aspect of the invention pertains to monitoring the influence of agents (e.g., drugs, compounds or other agents) on the expression or activity of polypeptides of the invention. These and other agents are described in further detail in the following sections.

DIAGNOSTIC ASSAYS

The nucleic acids, polypeptides and antibodies described herein can be used in methods of diagnosis of a suscepti-

bility to osteoporosis, as well as in kits useful for diagnosis of a susceptibility to osteoporosis.

In one embodiment of the invention, diagnosis of a susceptibility to osteoporosis is made by detecting a polymorphism in BMP2 as described herein. The polymorphism can be a mutation in BMP2, such as the insertion or deletion of a single nucleotide, or of more than one nucleotide, resulting in a frame shift mutation; the change of at least one nucleotide, resulting in a change in the encoded amino acid; the change of at least one nucleotide, resulting in the generation of a premature stop codon; the deletion of several nucleotides, resulting in a deletion of one or more amino acids encoded by the nucleotides; the insertion of one or several nucleotides, such as by unequal recombination or gene conversion, resulting in an interruption of the coding sequence of the gene; duplication of all or a part of the gene; transposition of all or a part of the gene; or rearrangement of all or a part of the gene. More than one such mutation may be present in a single gene. Such sequence changes cause a mutation in the polypeptide encoded by an BMP2 gene. For example, if the mutation is a frame shift mutation, the frame shift can result in a change in the encoded amino acids, and/or can result in the generation of a premature stop codon, causing generation of a truncated polypeptide. Alternatively, a polymorphism associated with a susceptibility to osteoporosis can be a synonymous mutation in one or more nucleotides (i.e., a mutation that does not result in a change in the polypeptide encoded by an BMP2 gene). Such a polymorphism may alter splicing sites, affect the stability or transport of mRNA, or otherwise affect the transcription or translation of the gene. An BMP2 gene that has any of the mutations described above is referred to herein as a "mutant gene."

In a first method of diagnosing a susceptibility to osteoporosis, hybridization methods, such as Southern analysis, Northern analysis, or in situ hybridizations, can be used (see *Current Protocols in Molecular Biology*, Ausubel, F. et al., eds., John Wiley & Sons, including all supplements through 1999). For example, a biological sample from a test subject (a "test sample") of genomic DNA, RNA, or cDNA, is obtained from an individual suspected of having, being susceptible to or predisposed for, or carrying a defect for, osteoporosis (the "test individual"). The individual can be an adult, child, or fetus. The test sample can be from any source which contains genomic DNA, such as a blood sample, sample of amniotic fluid, sample of cerebrospinal fluid, or tissue sample from skin, muscle, buccal or conjunctival mucosa, placenta, gastrointestinal tract or other organs. A test sample of DNA from fetal cells or tissue can be obtained by appropriate methods, such as by amniocentesis or chorionic villus sampling. The DNA, RNA, or cDNA sample is then examined to determine whether a polymorphism in *bmp2* is present. The presence of the polymorphism can be indicated by hybridization of the gene in the genomic DNA, RNA, or cDNA to a nucleic acid probe. A "nucleic acid probe", as used herein, can be a DNA probe or an RNA probe; the nucleic acid probe contains at least one polymorphism in BMP2. The probe can be any of the nucleic acid molecules described above (e.g., the gene, a fragment, a vector comprising the gene, etc.).

To diagnose a susceptibility to osteoporosis, a hybridization sample is formed by contacting the test sample containing BMP2, with at least-one nucleic acid probe. A preferred probe for detecting mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to mRNA or genomic DNA sequences described herein. The nucleic acid probe can be, for example, a full-length nucleic acid

molecule, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to appropriate mRNA or genomic DNA. For example, the nucleic acid probe can be all or a portion of SEQ ID NO: 1, or the complement of SEQ ID NO: 1, or a portion thereof. Other suitable probes for use in the diagnostic assays of the invention are described herein.

The hybridization sample is maintained under conditions which are sufficient to allow specific hybridization of the nucleic acid probe to BMP2. "Specific hybridization", as used herein, indicates exact hybridization (e.g., with no mismatches). Specific hybridization can be performed under high stringency conditions or moderate stringency conditions, for example, as described above. In a particularly preferred embodiment, the hybridization conditions for specific hybridization are high stringency.

Specific hybridization, if present, is then detected using standard methods. If specific hybridization occurs between the nucleic acid probe and BMP2 in the test sample, then BMP2 has the polymorphism that is present in the nucleic acid probe. More than one nucleic acid probe can also be used concurrently in this method. Specific hybridization of any one of the nucleic acid probes is indicative of a polymorphism in BMP2, and is therefore diagnostic for a susceptibility to osteoporosis.

In another hybridization method, Northern analysis (see Current Protocols in Molecular Biology, Ausubel, F. et al., eds., John Wiley & Sons, supra) is used to identify the presence of a polymorphism associated with a susceptibility to osteoporosis. For Northern analysis, a test sample of RNA is obtained from the individual by appropriate means. Specific hybridization of a nucleic acid probe, as described above, to RNA from the individual is indicative of a polymorphism in BMP2, and is therefore diagnostic for a susceptibility to osteoporosis.

For representative examples of use of nucleic acid probes, see, for example, U.S. Pat. Nos. 5,288,611 and 4,851,330.

Alternatively, a peptide nucleic acid (PNA) probe can be used instead of a nucleic acid probe in the hybridization methods described above. PNA is a DNA mimic having a peptide-like, inorganic backbone, such as N-(2-aminoethyl) glycine units, with an organic base (A, G, C, T or U) attached to the glycine nitrogen via a methylene carbonyl linker (see, for example, Nielsen, P. E. et al., *Bioconjugate Chemistry*, 1994, 5, American Chemical Society, p. 1 (1994)). The PNA probe can be designed to specifically hybridize to a gene having a polymorphism associated with a susceptibility to osteoporosis. Hybridization of the PNA probe to BMP2 is diagnostic for a susceptibility to osteoporosis.

In another method of the invention, mutation analysis by restriction digestion can be used to detect a mutant gene, or genes containing a polymorphism(s), if the mutation or polymorphism in the gene results in the creation or elimination of a restriction site. A test sample containing genomic DNA is obtained from the individual. Polymerase chain reaction (PCR) can be used to amplify BMP2 (and, if necessary, the flanking sequences) in the test sample of genomic DNA from the test individual. RFLP analysis is conducted as described (see Current Protocols in Molecular Biology, supra). The digestion pattern of the relevant DNA fragment indicates the presence or absence of the mutation or polymorphism in BMP2, and therefore indicates the presence or absence of this susceptibility to osteoporosis.

Sequence analysis can also be used to detect specific polymorphisms in BMP2. A test sample of DNA or RNA is

obtained from the test individual. PCR or other appropriate methods can be used to amplify the gene, and/or its flanking sequences, if desired. The sequence of BMP2, or a fragment of the gene, or cDNA, or fragment of the cDNA, or mRNA, or fragment of the mRNA, is determined, using standard methods. The sequence of the gene, gene fragment, cDNA, cDNA fragment, mRNA, or mRNA fragment is compared with the known nucleic acid sequence of the gene, cDNA (e.g., SEQ ID NO: 1) or mRNA, as appropriate. The presence of a polymorphism in BMP2 indicates that the individual has a susceptibility to osteoporosis.

Allele-specific oligonucleotides can also be used to detect the presence of a polymorphism in BMP2, through the use of dot-blot hybridization of amplified oligonucleotides with allele-specific oligonucleotide (ASO) probes (see, for example, Saiki, R. et al., (1986), *Nature (London)* 324:163-166). An "allele-specific oligonucleotide" (also referred to herein as an "allele-specific oligonucleotide probe") is an oligonucleotide of approximately 10-50 base pairs, preferably approximately 15-30 base pairs, that specifically hybridizes to BMP2, and that contains a polymorphism associated with a susceptibility to osteoporosis. An allele-specific oligonucleotide probe that is specific for particular polymorphisms in BMP2 can be prepared, using standard methods (see Current Protocols in Molecular Biology, supra). To identify polymorphisms in the gene that are associated with a susceptibility to osteoporosis, a test sample of DNA is obtained from the individual. PCR can be used to amplify all or a fragment of BMP2, and its flanking sequences. The DNA containing the amplified BMP2 (or fragment of the gene) is dot-blotted, using standard methods (see Current Protocols in Molecular Biology, supra), and the blot is contacted with the oligonucleotide probe. The presence of specific hybridization of the probe to the amplified BMP2 is then detected. Specific hybridization of an allele-specific oligonucleotide probe to DNA from the individual is indicative of a polymorphism in BMP2, and is therefore indicative of a susceptibility to osteoporosis.

In another embodiment, arrays of oligonucleotide probes that are complementary to target nucleic acid sequence segments from an individual, can be used to identify polymorphisms in BMP2. For example, in one embodiment, an oligonucleotide array can be used. Oligonucleotide arrays typically comprise a plurality of different oligonucleotide probes that are coupled to a surface of a substrate in different known locations. These oligonucleotide arrays, also described as "Genechips," have been generally described in the art, for example, U.S. Pat. No. 5,143,854 and PCT patent publication Nos. WO 90/15070 and 92/10092. These arrays can generally be produced using mechanical synthesis methods or light directed synthesis methods which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis methods. See Fodor et al., *Science*, 251:767-777 (1991), Pirrung et al., U.S. Pat. No. 5,143,854 (see also PCT Application No. WO 90/15070) and Fodor et al., PCT Publication No. WO 92/10092 and U.S. Pat. No. 5,424,186, the entire teachings of each of which are incorporated by reference herein. Techniques for the synthesis of these arrays using mechanical synthesis methods are described in, e.g., U.S. Pat. No. 5,384,261, the entire teachings of which are incorporated by reference herein.

Once an oligonucleotide array is prepared, a nucleic acid of interest is hybridized with the array and scanned for polymorphisms. Hybridization and scanning are generally carried out by methods described herein and also in, e.g., Published PCT Application Nos. WO 92/10092 and WO 95/11995, and U.S. Pat. No. 5,424,186, the entire teachings

of which are incorporated by reference herein. In brief, a target nucleic acid sequence which includes one or more previously identified polymorphic markers is amplified by well known amplification techniques, e.g., PCR. Typically, this involves the use of primer sequences that are complementary to the two strands of the target sequence both upstream and downstream from the polymorphism. Asymmetric PCR techniques may also be used. Amplified target, generally incorporating a label, is then hybridized with the array under appropriate conditions. Upon completion of hybridization and washing of the array, the array is scanned to determine the position on the array to which the target sequence hybridizes. The hybridization data obtained from the scan is typically in the form of fluorescence intensities as a function of location on the array.

Although primarily described in terms of a single detection block, e.g., for detection of a single polymorphism, arrays can include multiple detection blocks, and thus be capable of analyzing multiple, specific polymorphisms. In alternate arrangements, it will generally be understood that detection blocks may be grouped within a single array or in multiple, separate arrays so that varying, optimal conditions may be used during the hybridization of the target to the array. For example, it may often be desirable to provide for the detection of those polymorphisms that fall within G-C rich stretches of a genomic sequence, separately from those falling in A-T rich segments. This allows for the separate optimization of hybridization conditions for each situation.

Additional description of use of oligonucleotide arrays for detection of polymorphisms can be found, for example, in U.S. Pat. Nos. 5,858,659 and 5,837,832, the entire teachings of which are incorporated by reference herein.

Other methods of nucleic acid analysis can be used to detect polymorphisms in BMP2. Representative methods include direct manual sequencing (Church and Gilbert, (1988) *Proc. Natl. Acad. Sci. USA* 81:1991-1995; Sanger, F. et al. (1977) *Proc. Natl. Acad. Sci.* 74:5463-5467; Beavis et al. U.S. Pat. No. 5,288,644); automated fluorescent sequencing; single-stranded conformation polymorphism assays (SSCP); clamped denaturing gel electrophoresis (CDGE); denaturing gradient gel electrophoresis (DGGE) (Sheffield, V. C. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:232-236), mobility shift analysis (Orita, M. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:2766-2770), restriction enzyme analysis (Flavell et al. (1978) *Cell* 15:25; Geever, et al. (1981) *Proc. Natl. Acad. Sci. USA* 78:5081); heteroduplex analysis; chemical mismatch cleavage (CMC) (Cotton et al. (1985) *Proc. Natl. Acad. Sci. USA* 85:4397-4401); RNase protection assays (Myers, R. M. et al. (1985) *Science* 230:1242); use of polypeptides which recognize nucleotide mismatches, such as *E. coli* mutS protein; allele-specific PCR, for example.

In another embodiment of the invention, diagnosis of a susceptibility to osteoporosis can also be made by examining expression and/or composition of an BMP2 polypeptide, by a variety of methods, including enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. A test sample from an individual is assessed for the presence of an alteration in the expression and/or an alteration in composition of the polypeptide encoded by BMP2. An alteration in expression of a polypeptide encoded by BMP2 can be, for example, an alteration in the quantitative polypeptide expression (i.e., the amount of polypeptide produced); an alteration in the composition of a polypeptide encoded by BMP2 is an alteration in the qualitative polypeptide expression (e.g., expression of a mutant BMP2 polypeptide or of a different splicing

variant). Both such alterations (quantitative and qualitative) can also be present. An "alteration" in the polypeptide expression or composition, as used herein, refers to an alteration in expression or composition in a test sample, as compared with the expression or composition of polypeptide by BMP2 in a control sample. A control sample is a sample that corresponds to the test sample (e.g., is from the same type of cells), and is from an individual who is not affected by osteoporosis. An alteration in the expression or composition of the polypeptide in the test sample, as compared with the control sample, is indicative of a susceptibility to osteoporosis. Various means of examining expression or composition of the polypeptide encoded by BMP2 can be used, including spectroscopy, colorimetry, electrophoresis, isoelectric focusing, and immunoassays (e.g., David et al., U.S. Pat. No. 4,376,110) such as immunoblotting (see also Current Protocols in Molecular Biology, particularly chapter 10). For example, in one embodiment, an antibody capable of binding to the polypeptide (e.g., as described above), preferably an antibody with a detectable label, can be used. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin.

Western blotting analysis, using an antibody as described above that specifically binds to a polypeptide encoded by a mutant BMP2, or an antibody that specifically binds to a polypeptide encoded by a non-mutant gene, can be used to identify the presence in a test sample of a polypeptide encoded by a polymorphic or mutant BMP2, or the absence in a test sample of a polypeptide encoded by a non-polymorphic or non-mutant gene. The presence of a polypeptide encoded by a polymorphic or mutant gene, or the absence of a polypeptide encoded by a non-polymorphic or non-mutant gene, is diagnostic for a susceptibility to osteoporosis.

In one embodiment of this method, the level or amount of polypeptide encoded by BMP2 in a test sample is compared with the level or amount of the polypeptide encoded by BMP2 in a control sample. A level or amount of the polypeptide in the test sample that is higher or lower than the level or amount of the polypeptide in the control sample, such that the difference is statistically significant, is indicative of an alteration in the expression of the polypeptide encoded by BMP2, and is diagnostic for a susceptibility to osteoporosis. Alternatively, the composition of the polypeptide encoded by BMP2 in a test sample is compared with the composition of the polypeptide encoded by BMP2 in a control sample. A difference in the composition of the polypeptide in the test sample, as compared with the composition of the polypeptide in the control sample, is diagnostic for a susceptibility to osteoporosis. In another embodiment, both the level or amount and the composition of the polypeptide can be assessed in the test sample and in the control sample. A difference in the amount or level of the polypeptide in the test sample, compared to the control sample; a difference in composition in the test sample, compared to the control sample; or both a difference in the

amount or level, and a difference in the composition, is indicative of a susceptibility to osteoporosis.

Kits useful in the methods of diagnosis comprise components useful in any of the methods described herein, including for example, hybridization probes, restriction enzymes (e.g., for RFLP analysis), allele-specific oligonucleotides, antibodies which bind to mutant or to non-mutant (native) BMP2 polypeptide (e.g., to SEQ ID NO:2), means for amplification of nucleic acids comprising BMP2, or means for analyzing the nucleic acid sequence of BMP2 or for analyzing the amino acid sequence of an BMP2 polypeptide, etc.

SCREENING ASSAYS AND AGENTS IDENTIFIED THEREBY

The invention provides methods (also referred to herein as "screening assays") for identifying agents (e.g., antisense, polypeptides, peptidomimetics, small molecules or other drugs) which alter (e.g., increase or decrease) the activity of the polypeptides described herein. For example, such agents can be agents which bind to nucleic acid molecules or polypeptides described herein; which have a stimulatory or inhibitory effect on, for example, expression or activity of the nucleic acid molecules or polypeptides of the invention; or which change (e.g., enhance or inhibit) the ability of the polypeptides of the invention to interact with BMP2 binding agents (e.g., receptors).

In one embodiment, the invention provides assays for screening candidate or test agents that bind to or modulate the activity of polypeptides described herein (or biologically active portion(s) thereof). The test agents can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K. S. (1997) *Anticancer Drug Des.*, 12:145).

In one embodiment, to identify agents which alter the activity of an BMP2 polypeptide, a cell, cell lysate, or solution containing or expressing an BMP2 polypeptide (e.g., SEQ ID NO:2), or an active fragment or derivative thereof (as described above), can be contacted with an agent to be tested; alternatively, the polypeptide can be contacted directly with the agent to be tested. The level (amount) of BMP2 activity is assessed, and is compared with the level of activity in a control (i.e., the level of activity of the BMP2 polypeptide or active fragment or derivative thereof in the absence of the agent to be tested). If the level of the activity in the presence of the agent differs, by an amount that is statistically significant, from the level of the activity in the absence of the agent, then the agent is an agent that alters the activity of BMP2 polypeptide. An increase in the level of BMP2 activity relative to a control, indicates that the agent is an agent that enhances (is an agonist of) BMP2 activity. Similarly, a decrease in the level of BMP2 activity relative to a control, indicates that the agent is an agent that inhibits (is an antagonist of) BMP2 activity. In another embodiment, the level of activity of an BMP2 polypeptide or derivative or fragment thereof in the presence of the agent to be tested, is compared with a control level that has previously been established. A level of the activity in the presence of the

agent that differs from the control level by an amount that is statistically significant indicates that the agent alters BMP2 activity.

The present invention also relates to an assay for identifying agents which alter the expression of BMP2. For example, a cell, cell lysate, or solution containing a nucleic acid encoding BMP2 polypeptide (e.g., BMP2) can be contacted with an agent to be tested. The level and/or pattern of BMP2 expression (e.g., the level and/or pattern of mRNA or of protein expressed) is assessed, and is compared with the level and/or pattern of expression in a control (i.e., the level and/or pattern of the BMP2 expression in the absence of the agent to be tested). If the level and/or pattern in the presence of the agent differs, by an amount or in a manner that is statistically significant, from the level and/or pattern in the absence of the agent, then the agent is an agent that alters the expression of BMP2. Enhancement of BMP2 expression indicates that the agent is an agonist of BMP2 activity. Similarly, inhibition of BMP2 expression indicates that the agent is an antagonist of BMP2 activity. In another embodiment, the level and/or pattern of an BMP2 polypeptide in the presence of the agent to be tested, is compared with a control level and/or pattern that has previously been established. A level and/or pattern in the presence of the agent that differs from the control level and/or pattern by an amount or in a manner that is statistically significant indicates that the agent alters BMP2 expression.

In other embodiments of the invention, assays can be used to assess the impact of a test agent on the activity of a polypeptide in relation to a BMP2 binding partner. For example, a cell that expresses a compound that interacts with BMP2 (herein referred to as a "BMP2 binding partner", which can be a polypeptide or other molecule that interacts with BMP2, such as a receptor) is contacted with BMP2 in the presence of a test agent, and the ability of the test agent to alter the interaction between BMP2 and the BMP2 binding partner is determined. Alternatively, a cell lysate or a solution containing the BMP2 binding partner, can be used. An agent which binds to BMP2 or the BMP2 binding partner can alter the interaction by interfering with, or enhancing the ability of BMP2 to bind to, associate with, or otherwise interact with the BMP2 binding partner. Determining the ability of the test agent to bind to BMP2 or an BMP2 binding partner can be accomplished, for example, by coupling the test agent with a radioisotope or enzymatic label such that binding of the test agent to the polypeptide can be determined by detecting the labeled with ¹²⁵I, ³⁵S, ¹⁴C, or ³H, either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, test agents can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. It is also within the scope of this invention to determine the ability of a test agent to interact with the polypeptide without the labeling of any of the interactants. For example, a microphysiometer can be used to detect the interaction of a test agent with BMP2 or an BMP2 binding partner without the labeling of either the test agent, BMP2, or the BMP2 binding partner. McConnell, H. M. et al. (1992) *Science*, 257:1906-1912. As used herein, a "microphysiometer" (e.g., Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between ligand and polypeptide. See the Examples Section for a discussion of known BMP2

binding partners. Thus, these receptors can be used to screen for compounds that are BMP2 receptor agonists for use in treating osteoporosis or BMP2 receptor antagonists for studying osteoporosis. The linkage data provided herein, for the first time, provides such correction to osteoporosis. Drugs could be designed to regulate BMP2 receptor activation which in turn can be used to regulate signaling pathways and transcription events of genes downstream, such as Cbfa1.

In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either BMP2, the BMP2 binding partner, or other components of the assay on a solid support, in order to facilitate separation of complexed from uncomplexed forms of one or both of the polypeptides, as well as to accommodate automation of the assay. Binding of a test agent to the polypeptide, or interaction of the polypeptide with a binding partner in the presence and absence of a test agent, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein (e.g., a glutathione-S-transferase fusion protein) can be provided which adds a domain that allows BMP2 or an BMP2 binding agent to be bound to a matrix or other solid support.

In another embodiment, modulators of expression of nucleic acid molecules of the invention are identified in a method wherein a cell, cell lysate, or solution containing a nucleic acid encoding BMP2 is contacted with a test agent and the expression of appropriate mRNA or polypeptide in the cell, cell lysate, or solution, is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test agent is compared to the level of expression of mRNA or polypeptide in the absence of the test agent. The test agent can then be identified as a modulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater (statistically significantly greater) in the presence of the test agent than in its absence, the test agent is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less (statistically significantly less) in the presence of the test agent than in its absence, the test agent is identified as an inhibitor of the mRNA or polypeptide expression. The level of mRNA or polypeptide expression in the cells can be determined by methods described herein for detecting mRNA or polypeptide.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a test agent that is a modulating agent, an antisense nucleic acid molecule, a specific antibody, or a polypeptide-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

PHARMACEUTICAL COMPOSITIONS

The present invention also pertains to pharmaceutical compositions comprising nucleic acids described herein,

particularly nucleotides encoding the polypeptides described herein; comprising polypeptides described herein (e.g., one or more of SEQ ID NO:2); and/or comprising the agent that alters (e.g., enhances or inhibits) BMP2 polypeptide activity described herein. For instance, a polypeptide, protein, fragment, fusion protein or prodrug thereof, or a nucleotide or nucleic acid construct (vector) comprising a nucleotide of the present invention, or an agent that alters BMP2 polypeptide activity, can be formulated with a physiologically acceptable carrier or excipient to prepare a pharmaceutical composition. The carrier and composition can be sterile. The formulation should suit the mode of administration.

Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions (e.g., NaCl), saline, buffered saline, alcohols, glycerol, ethanol, gum arabic, vegetable oils, benzyl alcohols, polyethylene glycols, gelatin, carbohydrates such as lactose, amylose or starch, dextrose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, etc., as well as combinations thereof. The pharmaceutical preparations can, if desired, be mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with the active agents.

The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, polyvinyl pyrrolidone, sodium saccharine, cellulose, magnesium carbonate, etc.

Methods of introduction of these compositions include, but are not limited to, intradermal, intramuscular, intraperitoneal, intraocular, intravenous, subcutaneous, topical, oral and intranasal. Other suitable methods of introduction can also include gene therapy (as described below), rechargeable or biodegradable devices, particle acceleration devices ("gene guns") and slow release polymeric devices. The pharmaceutical compositions of this invention can also be administered as part of a combinatorial therapy with other agents.

The composition can be formulated in accordance with the routine procedures as a pharmaceutical composition adapted for administration to human beings. For example, compositions for intravenous administration typically are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water, saline or dextrose/water. Where the composition is administered by injection, an ampule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

For topical application, nonsprayable forms, viscous to semi-solid or solid forms comprising a carrier compatible

with topical application and having a dynamic viscosity preferably greater than water, can be employed. Suitable formulations include but are not limited to solutions, suspensions, emulsions, creams, ointments, powders, enemas, lotions, sols, liniments, salves, aerosols, etc., which are, if desired, sterilized or mixed with auxiliary agents, e.g., preservatives, stabilizers, wetting agents, buffers or salts for influencing osmotic pressure, etc. The agent may be incorporated into a cosmetic formulation. For topical application, also suitable are sprayable aerosol preparations wherein the active ingredient, preferably in combination with a solid or liquid inert carrier material, is packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant, e.g., pressurized air.

Agents described herein can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The agents are administered in a therapeutically effective amount. The amount of agents which will be therapeutically effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. In addition, in vitro or in vivo assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the symptoms of osteoporosis, and should be decided according to the judgment of a practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use of sale for human administration. The pack or kit can be labeled with information regarding mode of administration, sequence of drug administration (e.g., separately, sequentially or concurrently), or the like. The pack or kit may also include means for reminding the patient to take the therapy. The pack or kit can be a single unit dosage of the combination therapy or it can be a plurality of unit dosages. In particular, the agents can be separated, mixed together in any combination, present in a single vial or tablet. Agents assembled in a blister pack or other dispensing means is preferred. For the purpose of this invention, unit dosage is intended to mean a dosage that is dependent on the individual pharmacodynamics of each agent and administered in FDA approved dosages in standard time courses.

METHODS OF THERAPY

The present invention also pertains to methods of treatment (prophylactic and/or therapeutic) for osteoporosis, using an BMP2 therapeutic agent. An "BMP2 therapeutic agent" is an agent that alters (e.g., enhances or inhibits) BMP2 polypeptide activity and/or BMP2 expression, as described herein (e.g., an BMP2 agonist or antagonist).

The therapy is designed to inhibit, replace or supplement activity of an BMP2 polypeptide in an individual (for example, by administering a nucleic acid encoding an BMP2 polypeptide or a derivative or active fragment thereof including BMP2 from normal individuals not having osteoporosis; by administering an BMP2 polypeptide(s) or a derivative or active fragment thereof including BMP2 from normal individuals not having osteoporosis, or by administering a different splicing variant of the BMP2 polypeptide (s) or a derivative or active fragment thereof; and/or by administering an agent that alters the activity of the BMP2 polypeptide). The BMP2 therapeutic agent can be a nucleic acid (e.g., a gene, cDNA, mRNA, a nucleic acid encoding an BMP2 polypeptide or active fragment or derivative thereof, or an oligonucleotide); a protein, polypeptide, peptide, or peptidomimetic (e.g., an BMP2 polypeptide, such as SEQ ID NO:2, or an active fragment or derivative thereof), an antibody (e.g., an antibody to a mutant BMP2 polypeptide, or an antibody to a non-mutant BMP2 polypeptide, as described above); a ribozyme; a small molecule or other agent that alters BMP2 polypeptide activity and/or gene expression (e.g., which upregulate or downregulate expression of BMP2). More than one BMP2 therapeutic agents can be used concurrently, if desired.

The BMP2 therapeutic agent(s) are administered in a therapeutically effective amount (i.e., an amount that is sufficient to treat the disease, such as by ameliorating symptoms associated with the disease, preventing or delaying the onset of the disease, and/or also lessening the severity or frequency of symptoms of the disease). The amount which will be therapeutically effective in the treatment of a particular individual's disorder or condition will depend on the symptoms and severity of the disease, and can be determined by standard clinical techniques. In addition, in vitro or in vivo assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of a practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

In one embodiment of the invention, a nucleic acid is used in the treatment of osteoporosis. The term, "treatment" as used herein, refers not only to ameliorating symptoms associated with the disease, but also preventing or delaying the onset of the disease, and also lessening the severity or frequency of symptoms of the disease. In one embodiment, a nucleic acid of the invention (e.g., a nucleic acid encoding an BMP2 polypeptide, such as SEQ ID NO: 1; or another nucleic acid that encodes an BMP2 polypeptide or a splicing variant, derivative or fragment thereof) can be used, either alone or in a pharmaceutical composition as described above. For example, BMP2 or a cDNA encoding the BMP2 polypeptide, either by itself or included within a vector, can be introduced into cells (either in vitro or in vivo) such that the cells produce native BMP2 polypeptide. If necessary, cells that have been transformed with the gene or cDNA or a vector comprising the gene or cDNA can be introduced (or re-introduced) into an individual affected with the disease. Thus, cells which, in nature, lack native BMP2 expression and activity, or have mutant BMP2 expression and activity, or have expression of a disease-associated BMP2 polypeptide variant, can be engineered to express BMP2 polypeptide or an active fragment of the BMP2 polypeptide (or a different variant of BMP2 polypeptide). In a preferred embodiment, nucleic acid encoding the BMP2 polypeptide,

or an active fragment or derivative thereof, can be introduced into an expression vector, such as a viral vector, and the vector can be introduced into appropriate cells which lack native BMP2 expression in an animal. In such methods, a cell population can be engineered to inducibly or constitutively express active BMP2 polypeptide. Other gene transfer systems, including viral and nonviral transfer systems, can be used. Alternatively, nonviral gene transfer methods, such as calcium phosphate coprecipitation, mechanical techniques (e.g., microinjection); membrane fusion-mediated transfer via liposomes; or direct DNA uptake, can also be used.

Alternatively, in another embodiment of the invention, a nucleic acid of the invention; a nucleic acid complementary to a nucleic acid of the invention; or a portion of such a nucleic acid (e.g., an oligonucleotide as described below), can be used in "antisense" therapy, in which a nucleic acid (e.g., an oligonucleotide) which specifically hybridizes to the mRNA and/or genomic DNA of BMP2 is administered or generated in situ. The antisense nucleic acid that specifically hybridizes to the mRNA and/or DNA inhibits expression of the BMP2 polypeptide, e.g., by inhibiting translation and/or transcription. Binding of the antisense nucleic acid can be by conventional base pair complementarity, or, for example, in the case of binding to DNA duplexes, through specific interaction in the major groove of the double helix.

An antisense construct of the present invention can be delivered, for example, as an expression plasmid as described above. When the plasmid is transcribed in the cell, it produces RNA which is complementary to a portion of the mRNA and/or DNA which encodes BMP2 polypeptide. Alternatively, the antisense construct can be an oligonucleotide probe which is generated *ex vivo* and introduced into cells; it then inhibits expression by hybridizing with the mRNA and/or genomic DNA of BMP2. In one embodiment, the oligonucleotide probes are modified oligonucleotides which are resistant to endogenous nucleases, e.g. exonucleases and/or endonucleases, thereby rendering them stable in vivo. Exemplary nucleic acid molecules for use as antisense oligonucleotides are phosphoramidate, phosphothioate and methylphosphonate analogs of DNA (see also U.S. Pat. Nos. 5,176,996; 5,264,564; and 5,256,775). Additionally, general approaches to constructing oligomers useful in antisense therapy are also described, for example, by Van der Krol et al. ((1988) *Biotechniques* 6:958-976); and Stein et al. ((1988) *Cancer Res* 48:2659-2668). With respect to antisense DNA, oligodeoxyribonucleotides derived from the translation initiation site, e.g. between the -10 and +10 regions of BMP2 sequence, are preferred.

To perform antisense therapy, oligonucleotides (mRNA, cDNA or DNA) are designed that are complementary to mRNA encoding BMP2. The antisense oligonucleotides bind to BMP2 mRNA transcripts and prevent translation. Absolute complementarity, although preferred, is not required. A sequence "complementary" to a portion of an RNA, as referred to herein, indicates that a sequence has sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid, as described in detail above. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures.

The oligonucleotides used in antisense therapy can be DNA, RNA, or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotides can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotides can include other appended groups such as peptides (e.g. for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre et al., (1987), *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT International Publication No. W088/09810) or the blood-brain barrier (see, e.g., PCT International Publication No. W089/10134), or hybridization-triggered cleavage agents (see, e.g., Krol et al. (1988) *Biotechniques* 6:958-976) or intercalating agents. (See, e.g., Zon, (1988), *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule (e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent).

The antisense molecules are delivered to cells which express BMP2 *in vivo*. A number of methods can be used for delivering antisense DNA or RNA to cells; e.g., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the desired cells (e.g., antisense linked to peptides or antibodies that specifically bind receptors or antigens expressed on the target cell surface) can be administered systematically. Alternatively, in a preferred embodiment, a recombinant DNA construct is utilized in which the antisense oligonucleotide is placed under the control of a strong promoter (e.g., pol III or pol II). The use of such a construct to transfect target cells in the patient results in the transcription of sufficient amounts of single stranded RNAs that will form complementary base pairs with the endogenous BMP2 transcripts and thereby prevent translation of the BMP2 mRNA. For example, a vector can be introduced *in vivo* such that it is taken up by a cell and directs the transcription of an antisense RNA. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art and described above. For example, a plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site. Alternatively, viral vectors can be used which selectively infect the desired tissue, in which case administration may be accomplished by another route (e.g., systematically).

Endogenous BMP2 expression can also be reduced by inactivating or "knocking out" BMP2 or its promoter using targeted homologous recombination (e.g., see Smithies et al. (1985) *Nature* 317:230-234; Thomas & Capecchi (1987) *Cell* 51:503-512; Thompson et al. (1989) *Cell* 5:313-321). For example, a mutant, non-functional BMP2 (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous BMP2 (either the coding regions or regulatory regions of BMP2) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express BMP2 *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of BMP2. The recombinant DNA constructs can be directly administered or targeted to the required site *in vivo* using appropriate vectors, as described above. Alternatively, expression of non-mutant BMP2 can be increased using a similar method: targeted homologous recombination can be used to insert a DNA construct com-

prising a non-mutant, functional BMP2 (e.g., a gene having SEQ ID NO: 1), or a portion thereof, in place of a mutant BMP2 in the cell, as described above. In another embodiment, targeted homologous recombination can be used to insert a DNA construct comprising a nucleic acid that encodes an BMP2 polypeptide variant that differs from that present in the cell.

Alternatively, endogenous BMP2 expression can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of BMP2 (i.e., the BMP2 promoter and/or enhancers) to form triple helical structures that prevent transcription of BMP2 in target cells in the body. (See generally, Helene, C. (1991) *Anticancer Drug Des.*, 6(6):569-84; Helene, C., et al. (1992) *Ann. N.Y. Acad. Sci.*, 660:27-36; and Maher, L. J. (1992) *Bioassays* 14(12):807-15). Likewise, the antisense constructs described herein, by antagonizing the normal biological activity of one of the BMP2 proteins, can be used in the manipulation of tissue, e.g. tissue differentiation, both in vivo and for ex vivo tissue cultures. Furthermore, the anti-sense techniques (e.g. microinjection of antisense molecules, or transfection with plasmids whose transcripts are anti-sense with regard to an BMP2 mRNA or gene sequence) can be used to investigate role of BMP2 in developmental events, as well as the normal cellular function of BMP2 in adult tissue. Such techniques can be utilized in cell culture, but can also be used in the creation of transgenic animals.

In yet another embodiment of the invention, polypeptides and/or agents that alter (e.g., enhance or inhibit) BMP2 polypeptide activity, as described herein, can be used in the treatment or prevention of osteoporosis. The polypeptides or agents can be delivered in a composition, as described above, or by themselves. They can be administered systemically, or can be targeted to a particular tissue. The polypeptides and/or agents can be produced by a variety of means, including chemical synthesis; recombinant production; in vivo production (e.g., a transgenic animal, such as U.S. Pat. No. 4,873,316 to Meade et al.), for example, and can be isolated using standard means such as those described herein.

A combination of any of the above methods of treatment (e.g., administration of non-mutant BMP2 polypeptide in conjunction with antisense therapy targeting mutant BMP2 mRNA; or administration of a first variant of BMP2 polypeptide in conjunction with antisense therapy targeting nucleic acid encoding a second variant of BMP2 polypeptide), can also be used.

The invention will be further described by the following non-limiting examples. The teachings of all publications cited herein are incorporated herein by reference in their entirety.

EXEMPLIFICATION

Identification of the BMP2 Gene With Linkage to Osteoporosis Phenotype and Family Construction

Patients who have low impact fractures and/or take bisphosphonates for treating osteoporosis are automatically treated as affecteds. People with low bone mass density (BMD) measurements are considered to be osteoporotic, and have been shown to have substantially increased risk of fractures. BMD measurements are taken for both the hip and the spine. For each person with BMD measurements, a standardized BMD score is computed (mean 0, standard deviation 1 for the population), which is adjusted for sex,

age, body weight and hormone replacement therapy (HRT). For the combined analysis, the two measurements are summed. Population BMD data from Iceland and the United States are used for standardization and adjustment. For example, a person with a positive BMD score is above average and one with a negative score is below average for his/her age, body weight and possibly HRT. Assuming approximate normality, a score of -1 corresponds approximately to the lower 16th percentile, etc.

For analysis, we start with a current list of primary people, people who have BMD measurements and/or are severely affected, and for whom we have genotypes. We then use the genealogy database to create family clusters linking these primary people using a threshold distance of 5 meiotic events. This procedure produced 190 potentially informative clusters with a total of 1215 primary people.

Linkage data

Four genome wide scans (GWS) were performed using osteoporotic phenotypes at different skeletal sites; the hip, the spine, and combined phenotypes. All GWS analysis located at 20 cM region on Chr20, between 10 cM and 30 cM based on the Marshfield map.

All of the analyses were performed using the Allegro linkage program developed at deCODE (Gudbjartsson et al., *Nature Genetics*, 25: 12-13, May 2000). The allele sharing analysis uses the Spairs scoring function of GENEHUNTER (Kruglyak et al., *Am. J. Hum. Genet.*, 46: 1347-1363, 1996), but families were weighted using a scheme which is a compromise between weighting families equally and weighting affected pairs equally. The allele-sharing LOD scores were computed using the 'exponential model' described in Kong and Cox, *Am. J. Hum. Genet.*, 61: 1179-1188 (1997).

Hip

The phenotype used was age, sex, weight and HRT corrected BMD < -1 SD at the hip (total hip). Hip fracture cases and bisphosphonate users are also considered affected even if values are above -1 SD. A total of 346 affected were used in this analysis. The GWS resulted in a LOD score of 3.1 using our standard set of markers. Adding 10 extra markers at the region on interest, between 11 cM and 39 cM, resulted in a LOD score of 3.3.

Spine The phenotype was age, sex, weight and HRT corrected BMD < -1 SD at lumbar spine (L2-L4). Vertebral compression fracture cases and bisphosphonate users are also considered affected even if values are above -1SD. A total of 402 affected people were used in this analysis. The GWS resulted in a LOD score of 2.4 at the same location as in the hip analysis using the standard set of markers, but a LOD score of 2.9 with the extra marker set.

Combined

The phenotype used was the sum of corrected BMD < -1.5 SD. Vertebral compression fracture, hip fracture, other osteoporosis related low impact fracture (at least two fractures) and bisphosphonate users (BMD measurements before treatment start are used if available) are all considered affected. A total of 522 affected were used in this analysis. The GWS resulted in a LOD score of 2.5 with the standard marker set, but a LOD score of 3.9 using the extra markers in the region.

Combined Severe

The phenotype used was the sum of the age, sex, weight and HRT corrected BMD < -2.3 SD. Vertebral compression fracture, hip fracture, other osteoporosis related low impact fracture (at least two fractures) and bisphosphonate users affected. The number of affected in this analysis was 290. The GWS resulted in a LOD score of 3.8 with the standard

set but a LOD score of 4.7 was reached using the extra 10 markers in addition.

Corticosteroid users and women with early menopause were excluded as affected in all analysis.

The BMP2 Gene

The BMP2 gene is located in this region. Only 5 kb are between the marker D20S846, which gives the highest LOD score, and the 3' end of the gene. The gene has been sequenced and characterized in terms of exon/intron structures, promoter region and transcriptional start sites. This information are publicly available.

A number of nucleotide changes are observed in the Icelandic population. These changes have not to our knowledge been described before, as follows:

Promoter Region:

- A to G at nucleotide position 420;
- A to G at nucleotide position 472;
- G to C at nucleotide position 1464;
- G to A at nucleotide position 1722;
- C to G at nucleotide position 1914;

Coding Region:

- T to G at nucleotide position 3747; resulting in an amino acid change from serine to alanine at amino acid position 37;
- A to G at nucleotide position 3899; no amino acid change;
- G to T at nucleotide position 3918; resulting in an amino acid change from alanine to serine at amino acid position 94;
- A to T at nucleotide position 11980; resulting in an amino acid change from arginine to serine at amino acid position 189;

3' UTR and Downstream Region:

- C to T at nucleotide position 12571;
- T to C at nucleotide position 13066;
- A to G at nucleotide position 13209;
- C to A at nucleotide position 13296; and

at least one deletion in nucleotides at positions 13533–13536 (all numbering is relative to SEQ ID NO: 1).

BMP2 binds to the receptors BMPR-IA or BMPR-IB, and BMPR-II, leading to formation of receptor complex heterodimer and phosphorylation of the BMPR-IA or BMPR-IB receptors. Once activated, these receptors subsequently phosphorylate SMAD1, SMAD5 or SMAD8, which in turn form complexes with SMAD4 and translocate to the nucleus where the transcription of specific genes is affected (Massague, *J. Annu. Rev. Biochem.*, 67:753–791 (1998); Chen, D. et al., *J. Cell Biol.*, 142(1):295–305 (1998)). SMADs 6 and 7 block signals by preventing the activation of SMAD1, SMAD5 or SMAD8 by the BMP2 receptors and have been shown to inhibit osteoblast differentiation (Miyazono, K., *Bone*, 25(1):91–93 (1999); Fujii, M., et al., *Mol. Biol. Cell*, 1(11):3801–3813 (1999)). BMP2 stimulates Cbfa1, alkaline phosphatase and Collagen type I (osteoblast specific proteins) expression through BMPR-IB (Chen, D. et al., *J. Cell Biol.*, 142(1):295–305 (1998)). Cbfa1 regulates the expression of osteoprotegerin (OPG), which is an osteoblast-secreted glycoprotein that functions as a potent inhibitor of osteoblast differentiation and thus of bone resorption (Thirunavukkarasu, K., et al., *J. Biol. Chem.*, (2000)). Cbfa1 controls osteoblast differentiation and bone formation. During cellular aging of human osteoblasts, there is a significant reduction (up to 50%) of Cbfa1 mRNA

(Christiansen, M., et al., *J. Gerontol. A Biol. Sci. Med. Sci.*, 55(4):B194–200 (2000)).

Frequency of the nucleotide changes at positions 3747, 3899 and 11980 relative to SEQ ID NO 1 found in the coding region of the gene BMP2 in the Icelandic population.

Affected-unrelated: are all unrelated patients with the combined phenotype. Most also have the hip and spine phenotype.

Controls-unrelated: are all unrelated controls, affected status unknown. Fam_aff: are affected in our familial material, divided into different phenotypes as appropriate.

No difference in frequency of the nucleotide changes is observed between the affected-unrelated and fam_affected.

A→G transition at nucleotide position 3899 in SEQ ID NO: 1 does not affect the amino acid composition.

Wt/wtwt/GG/Gtotalaffected-unrelated186551134Controls-unrelated166345124fam_aff_combined258167173fam_aff_hip176456137fam_aff_spine227255149

A→T transversion at nucleotide position 11980 in SEQ ID NO: 1, changing the amino acid arginine to serine.

wt/wtwt/TT/totalaffected-unrelated216750138controls-unrelated167150137fam_aff_hip145139104fam_aff_spine18513910

T→G transversion at nucleotide position 3747 in SEQ ID NO: 1, changing the amino acid serine to alanine.

Wt/wtwt/GG/Gtotalaffected-unrelated128100138controls-unrelated25470267fam_aff_combined275190294fam_aff_spine238180256

Of the 10 affected-unrelated that are wt/G, 9 also have the spine phenotype and 8 have the hip phenotype.

Results and Discussion

As a result of the linkage studies, the analysis shows that this locus is involved in multiple osteoporosis phenotypes. Furthermore, mutation within the human BMP2 gene is likely to explain the phenotypes in these families. Sporadic occurrence of osteoporosis, i.e., occurrence without familial connection, can also be determined using the information contained herein.

Osteoporosis could be caused by a defect in the BMP2 gene as follows: A mutation in the BMP2 gene (transcription, splice, protein variant etc.) could lead to a reduction of its action on Cbfa1 through BMPR-IB and the subsequent signaling pathway, that would lead to less bone formation because of fewer and less active osteoblasts and more bone resorption because of less OPG and more osteoclasts. This would lead to bone loss. Since a significant reduction of Cbfa1 levels is associated with aging osteoblasts, this effect could become more important with older age.

While this invention has been particularly shown and described with reference to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended claims.

TABLE 1

LOCUS — 14759 bp DNA

DEFINITION Human bone morphogenetic protein 2 (SEQ ID NO:2) (BMP2) gene, complete cds (SEQ ID NO:1), complete sequence.

KEYWORDS .

SOURCE human.

ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1-14759)

AUTHORS Blakey, S.

TITLE Direct Submission

JOURNAL Submitted (04-APR-2000) Sanger Centre, Hinxton, Cambridgeshire, CB10 1SA, UK. E-mail enquiries: humquery@sanger.ac.uk Clone requests: clonerequest@sanger.ac.uk

COMMENT This sequence was taken from GenBank sequence AL035668 (VERSION AL035668.15, GI:4995292), bp 118501 . . . 133259.

FEATURES Location/Qualifiers

source 1 . . . 14759
/organism="Homo sapiens"
/db_xref="taxon: 9606"
/chromosome="20"
/map="20p12"
/clone="RP5-859D4"
/clone_lib="RPCI-5"

gene 2072 . . . 12634
/gene="BMP2"
/note="BMP2A"
/db_xref="LocusID:650"
/db_xref="MIM:112261"

exon 2072 . . . 2387
/gene="BMP2"
/number=1

exon 3632 . . . 3984
/gene="BMP2"
/number=2

CDS /join(3639 . . . 3984, 11757 . . . 12601)
/gene="BMP2"
/note="BMP2 exons defined by comparison to mRNA sequence (NM_001200)
/codon_start=1
/product="bone morphogenetic protein 2 precursor"
/protein_id="NP_001191.1"
/db_xref="GI:4557369"

/translation="MVAGTRCLLALLLPQVLLGGAAGLVPELGRRKFAAASSGRPSSQ
PSDEVLSEFELRLLSMFGLKQRPTPSRDAVVPYMLDLYRRHSGQPGSPAPDHLRLERA
ASRANTVRSFHHESLEELPETSQKTRRRFFNLSSIPTEEFITSAELQVFRQMQDA
LGNNSFFHHRINIYEIIKPATANSKFPVTRLLDTRLVNQNASRWESFDVTPAVMRWTA
QGHANHGFFVVEVAHLEEKQGVSKRHVRSRSLHQDEHSWSQIRPLLVTFGHDGKGHPL
HKREKRQAKHKQRKRLKSSCKRHPLYVDFSDVGNWDWIVAPPGYHAFYCHGECFFPLA
DHLNSTNHAIIVQTLVNSVNSKIPKACCVPTLSAISMLYLDENEKVVVKNYQDMVVEG

CGCR"
exon 11757 . . . 12634
/gene="BMP2"
/number=3

BASE COUNT 3913 a 3147 c 3351 g 4348 t

CCTTGGTTTTTGGGGATCATTGGGCAAGCCCGAGGTGCTGTGCATGGGGGCTCCTGGAAT
CCTGGGAAGGGCAGAAAGCCTTGGCCCCAGACTCATCGTCAGCAGCTCTGAGCAGTATT
TCGGCTGAGGAGTGAAGTTCAGTGAATATTCAGCTGAGGAGTGAAGTGGCCACGTGTCACA
GCCCTACTTCTTGGGGCCTGGTGAAGAGGGTGGCGTAGAAGGTCCAAGGTCCCAAAC
TGAATGTCTCCTGTATGCTTGGTTCACACAGTGCCTTATTTTACCTTCTCTGAGCTGCT
AATCGCCTGCCTCTGAGCTGGGTGAGATAAATATCACAAAGGCACAAAGTATTGTACAAT

TABLE 1-continued

AAAAAAATCAAATCCCTCCCATCCATCCTTCAGTCTGCCACACACGCAGTCTACGTTACA
CACATGTCACGTAAAGCAGGATGACATCCATGTCACATACATAGACATATTAACCGAAAT
GTGGCCCTTCGGTTGCATATATTTCTCATACATGAATATATTTATAGAAATATATGCACAT
ATTTTTGTATATGGATATATTTATGTAACATAAAATTTACATGCGTATGGATATGAAAA
TAAATGCATACACATTTATGTAATAAAAAATTTGTACACATGCATTTACATATGTAATACA
TACATCTCTATGTATTAATGTTTAAAAACACTCAATTTCCAGCCTGCTGTTTTCTTTTAA
TTTTCTCTATTCGGGAAACAGAAGCGTGGATCCCACGCTATGCTATGCCAAAATA
CGCTGTAATTGAGGTGTTTTGTTTTGTTTTGTTTTGAAATCGTATATTACCAGAAAAAC
TTCAAAGTAAAGTTGAATAACGGGCCAGCGGGGAAATAAGAGGCCAGACCCCTGACCCCT
GCATTTGTCTCGGATTTCCGCTCCAGAGTCCCGCGAGGGTCCGGCGCGCCAGCTGATCT
CTCCTTTGAGAGCAGGGAGTGGAGGCGGAGCGCCCCCTTGGCGGGCCGCGCCCCCGC
CCTCCGCCCCACCCCGCGGCTGCCGGGGCGCGCTCCACACCCCTGCGCGCAGCTC
CCGCCCCGTCGGGGATCCCGCGGAGCCGCGCGCGAAGGGGGAGGTGTTCCGGCCGCGC
CGGGAGGGAGCCGGCAGGCGGCTCCCTTTAAAAGCCGCGAGCGCCGCGCCACGGCGCC
GCCGCCCGCTCGCCGCCCGGAGTCCCTCGCCCCGCGCGCTGCGCCCCGCTCGCGCTG
CGTAGTCGCTCCGCTTCCACACCCCGCGGGACTGGCAGCCGCCCGCCACATCTGC
CGCCACAGCCTCCGCGGCTACCCGAACGTTCTCGGGGCCAGCGCGAGTGGATCACCGG
GGACCCGAGGACCCCGCGCGCCGAGACCCCGCGGGCTGGAGCACCCGGCAGAGCGC
GCCACAGCCCGTGGCTCTGCTGCCGGGCTGCCAGAGCCCGGACGGGCGCGCAGA
GCGCCGGGACTCCGGAGCCGATCCCTAGCGCCGCGATGCGGAGCACCTACTGCAGGAGA
TCGGGGGCTGGGACCGCTGGCCGAGGTGTGATCGGACCCAGGCTAGCCACAAGGGC
ACTTGGCCCCAGGGCTAGGAGAGCGAGGGAGAGCACAGCCACCCGCTCGCGGCCCGG
GACTCGGCTCGACTCGCCGAGAATGCCCGGAGGACGACGGGGCCAGAGCCCGGCTG
CTTTCAACTGGCGAGCGCAATGGGGTGCCTGGAGTAAGGCAGAGTGTGCGGGGGG
CAACTCGCTGGCACCGAGATCGCCCGCTGCCCTTCCCTGGACCCGCGCTCGCCCAGGA
TGGCTGCCCGAGCCATGGGCGCGCGGAGTAGCGCGGAGCGCCGACCCCTCGACCCC
CGAGTCCCGAGCCGCCCGCGGGGCCACGCTCCCTCGGGCGCTGGTTCTTAAGGA
GGACGACAGCACCAGCTTCTCTTTCTCCCTTCCCTTCCCTGCCCGCACTCCTCCCT
GCTCGCTGTGTGTGTGTGTCAGCACTTGGTGGGACTTCTTGAATTCAGGGAGAATA
ACTTGGCACCCCACTTTGCGCCGCTGCTTTGCCCGAGCGGAGCCTGCTTCGCCATCTC
CGAGCCCCACCGCCCTCCACTCCTCGGCCCTGCGCGCACTGAGACGCTGTTCCAGCG
TGAAAAGAGAGACTGCGCGGCCGCAACCCGGGAGAAGGAGGAGCAAGAAAAGGAACGG
ACATTCGGTCCCTGCGCCAGGTCCTTTGACCAGAGTTTTTCCATGTGGACGCTCTTTCAA
TGGACGTGTCGCCCGCTTCTTAGACGACTGCGGTCTCCTAAAGGTAGAGGACGCGG
GCCAGGGCCCGGGTGGGTGGTGGGAGGGGATTTGGGCAGCCACTGCGGTAGAGC
CCTTCTTACGTCCAGGCCAGAAGTAACAGACCCCTCTCCAGT&CACGTGCAACGGAGC
CCTGCAGGGGCTCCCACTTCCAGCTGCCCGGGCGACCGTAAGCCTCACCCCTCCCGGCC
GCACCTTCCACCCCTCTTTCTTCCCTCTCCCTGGAATACTTTTGGAGCTGTTAAACT
TAGATGAGGTGTTTTATTTATTTATTTATTTATTTTAAATTTTTTAAAAACTTTTTTGG
GTCAAAAGAAATCCCTTTGAGAGGGTAGCCCTGGGTTTACCCGTTAGCTGAGAACCTGT

TABLE 1-continued

GTTTTCTCCCAGTGTGAAGGGTGACTTATACTTGTGAGAGTATTTGTTGGTAATGGGAAT
 AAGTCCCAAAGGCAAGCCACATAGCAGAAGATACGTTCTCATTGAGGCAGCTACACATTA
 CGACGGGGACACTGAATTGATCATCAGTTCATTTACAAGCACATTTCTAAGTGAGGTGCT
 CTCTGTAGCAGAAATCAGATTTGAAAGGCAGTAAGATCTCACTCCACTCTTTGAGAATT
 CATCCAATGAAAGCAGAAATCACCTGTTGTATATGTAATAATTTGTTGTATGTGTACAT
 TCTGCCATCTTAACCTGAAATGATTATAGATCCAGCTAATCATTCCCAGGTAATGCTGA
 TTAGAATACTTTTTTTTTGTATAGGAATGTAATAAGAACAACGTTTTAGACACCTCTT
 CTGGAATTTAGCATGGAAGCTCTCAACTTTATTTTTAAGGCCGGAAGATGCTGTGTCT
 CTGTTACAACCTTAAAGGAAGATCATTTAAGTTAGTTAACACCTAAAACATTCATTGTG
 TGAGGATTTTATCAGTGATGTCTGCATATTTTCATCATTCATCTAGAAGTGGTTGATCA
 GAACTAAACAGGCTACACGTTATTTCAACTGTGTTATTTAACTTAAAAGCATGCTTGAG
 TTTATAAAATCAGAAATTTATATCTTTGTGAGTGTAAATGTTACCTGAGAAACAGTACAGA
 AGTGACCAACTTGATTTAAATCAACTTGAATAACTTCAGGTCTTAATGCAGTTAGATAA
 TGGAGAAAAGCTATGTAATTTTGGCCCAAATTTCAACTAATCCATTTCTGTCTCATTAT
 GACTAATATATCATCTTAATCTGGATGGATATAGCACTTTTTTCAAGACTAATCATTGT
 TGTATACACCCAGGATTTGCTTTTGATAAACATCTCTGTGCCATGCATGCCACGAAAAA
 GTTTTTGGTAAACCATGTGATGAAGTTGCTGGCTCAAGAACAGAATTTAGTTTCTACAG
 CATTAAATGAGCATTATTTGAAAAAGACCATAAAGACCCAATCATAAGAATTACCTGTT
 GGGTTTTCTTTGTAGGTGTGATCGAATGGTTGGTGGAAATACTCGACGAGATATCATGA
 TAGCATTCTTTCAACCAATATGAGTATAATGCGACCATATCATAGGGATCTGAGACAGA
 ATTATCAGTTGTATTTTTCCTATGAAATTTGTCTAGTCCCTTCTCCAGTGGCTTTTATT
 TGGGAGAATATCAGCTTTGCTAAAATGTTATTTGTTTCAAGATCATTA AAAAGTCTTCA
 GCTACATAGACCTTTGGAAACGCCATTGAACATAGAAAAGTCAGTTCTGCAAGTGGAAA
 GAGTGTTTTGTGTATTGCTGTAGTTGGAACACATTGAAACTGGTTGACTTCACTGGCCC
 TCCAAAAGTCTTTATGCTTTTTTGTGATGGGAGAGAGAAAGACCAGGTGCTTCTTGT
 TCTCCTCACTCTGAAGGCACAGTCTTCTTCTACATGAAATAACTGGATTATTTGCTC
 TGTGACTGAAGCTTTCAAATAGAGATTAACCTCTTTCCACAAATATAATTTATATGAAA
 ATATCCATATAATAGAAAAGTTCAAGAAATAACTATTGCCCTGCATTAGAGACTTTGTGG
 CACAAATCCCCGTCAAACAACAGATTTGGACACATAGATCCACCAAAACCAACTTT
 ACCTGGTATGGTTCCTAGTGGCCCAGGTATTTCAATTGTCATTACAGAGGCCACATTAA
 GTAGGAAAATTACTCTATTTTGAAATGGTTGTTGAGATTGAGGCTTTGGTGTCCAGTGAT
 ACTTCCTGGCACTGACATTTTCCGTTCCACCTGTTTTTTAGTGGTTCCCTAAATTTCT
 CTTAATCCCTTTGCACTGAACTATTTTGGCTTCTTAGACTTGCTCTTTGTGTATTTTAC
 TGAGACAATAAGAGAATATTTTCATCATTCGAAGGTGTTGGTGTAAAGGGTGGCAGAGG
 CCAAATCAGGGTTGTTGATGACAACCATGCTCTCTATTCTTTATTTGCCATTTCCCTGT
 TGTATTTTTTTAAAATGGAATGTTTTTAACCTTTTGTATTTGATATTTTTTTCTCCTT
 GATCAGTTGCTGTATTTTATTATCTGGAAAATCTTATATTACTCAGCCTCTTTCAT
 TTTGTGTTAGGGCAGTGACTTCCAGCCTTACTGATTGCCAGCATATCCCCAGGTTTTGTT
 GTTGTGTTGTTGTTTTACTGGAGATTTTTTAGCCCAAAGTGTGTTTTAAAATCCTCGAA
 GCATAACGGTAACTTACTTTTTTGATAAAACTTACCATACTTTATTTAGAACAAAAGGGC

TABLE 1-continued

AGCCACAAAATAGCAGTGGCTCCTTATAAAAATAGACACATTCCAGTGGGCCCCGTCACCTT
TTCTGCTCATTCTGTCTGTTCTGTCCATCATACCTAAGTCATATATTTCTGTTCATTTA
GTTGGGACAGAACTCACCAATGTTATCATGTACTAAATATAAATGTGCCCTAATGGT
TTTGACTTTTGTCTAAGTTTTTGTAGTCCCTCATGTATGTTAGGTAGTGCCTCTAGTAGCC
AGAAATTTGGAACTGGCTGGCATGATGGCTAATACCTGTAATCCCAGCACTTTGGGAG
GCTTAGGTGGTGGATCACTTGAGGTCAGGAGTTCAGACCAGCCTGGCCAACATGGTGA
AACCACATCTCTACTAAAATATAAAAAATAGCCAGGTATGATGGCCATGCTGTAAATC
CGAGCTAATTTGGGAGGCTGAGATGGGAGGACTGCTTGAACCTGGGAGGTGGAGGCTGCTG
TGAGCCAAGATTGTGCCACTGCACTCCAGCCTGGGCAACAGAGTGAGACCTGTGTGACA
AAAACAAGAAACAAAACAAAACAAAAGACAAGAAACCTGAGAAGCGCAGTAGATTCATTT
ATATATATCTACTTTTAAATTTGCTAGCTCTGTGACCTTAGGAAAGTTACATAACCTCTCT
GAACTGCAACTGTTTCATTTACAAAATGGAGATAATGATAGTTTTTCTCTAATTTGGTTT
TTGTGAGATAATTCATATAAAGCTGATGGTCCAGATTACACTCAAAAAAGCATTCAGC
TGTCATTATCATTATGACTTCTTTTGTAAATGTTATAGCCTTTCCTTCTCTAGGAAAAG
GAGGCCAGAGTGGACCTAGGCTGACTGAGAGAATTCAGCTCAGTCTTTTGAATTTTGT
AGGTAGAGGAATGATGATATAGTATAGATTATTAATTAGGACTTCACTTTTGGAGAAA
AGTTCAGATATCATTGTTGTCTTATTTTTTCTTCACTTTCCACATTTTTGCAGCCATAGC
TCCATCCATTTGGTTAAGAACTTAGAAGCTCACAACTCGGGTCAAAGACAGGTCGAAAT
CCTCAAATCCCTTAAGAACTTCAGCTTATTCAGGAAGGGATATTTACAGAAAACCTAGCAA
TTGTATAAGTCTCAAAAAAGCATACTACTTGAGGATCCATATATTTTTGGCATCCTC
AGGGTTGCTGTGATGATTTATAGAAGTTTGTATTTAATTTACTTTATTTCAAATAGG
TTTTAATTTTTGTACCCCTTAAGAAAAGATTCGTACTCTTCCCTGGCAGATTAAGAAAAT
GAGCGTATATTCCTAACCTTGCCAGTTACTTTCTGGGTTTGAGGGTTCTGTGAACG
TCTAACTTACCTCTGTGACCTGTTTCTGCAACCAGGGGTGTGCAATGGATGCTTTTGTG
TTGAGGATGGGACCTTCAAGAAACAGATTCAGTGGTGCAGTGGGAAGGTCAGAGAAA
GATCTTCGTATCGCCTATTATTTATTTGCTCGTCTATTTTTTCTCCTTTCTTAAGCCACT
AACTGATTCCTCTTGTCTAAGGCTGCTTACTTCCACTGAGACCTTGAACCACATGAAAT
GTTGTTGTCTGTGTTCTGTTCAAATAGTGCAATTTTGTATGATTCATCTTGTTCATTT
AATTTTTTGGGAGGTTATTTCTATTTTCTATACCTTTTTTATACCCATCTTCTTACTTC
ATTTACCTGTCCCTCATACTTGACTTGTAGCTTGTCCCTTCACTGTCTATGCTGGCCAT
GTGGGTGTGTACGTGTCTGCGACAGAGAGAATGTGTGAGAATGTATCTTTCTTTATGCAT
TGGGATTTAGGTTTTTCTGCAATGTGATTTCTCTGGGCACTTTTGTTAATATAGCTA
GTCAGCGAGTGTCTAGATAATTTTCTTGCCTCCCCCTTTTGAAGAAAAGAGGGTGT
TCTTAGATGATTTCTTATCAGATAAGCCAGTAGCTCAGGTGCTGGTCTGGCTTTGGTGT
ATTGGGGTCTGAGGTTGCTGACTTTTACCTTCTCTGCTGAAAAATTACCTTCAGCAGAAA
CGTCTGAATTCGAAGGAGAAGGAGAAAAAACAGGCCAACACAGTCTTGGTACTCCTT
GGGAGCCACTGAGAAGAGTCCAGGTTCAAATGGTCAGAAGGTTATTTAATGATTTGTGTC
TGGCCTAAAGTACCATTAGCTTCCAGTGGAGTTTGAATGTGGATGGATCCTGAAAGGTA
TTCCCCAGAGGTTTGGATTAATAGGCACAAGGGAACCTAAAGGACTCTATTGGCCTGAT

TABLE 1-continued

ACTCCCATATCCACGTAGAGAGCTTTAGAAGAACCTTCTGTTCTGAGACCCCTGGCTGG
GCCACCCAGAGCTGGCCCATCAACTCTTACTCCTTTGCCACCCTAATGGTTCTTCTA
CTAGTTTTTATATTATTTAACAAAAAGGCACCTTAAAAATGCACTCCTGGCAATCTATAC
TGGAATATGAAAAACATGCTGCAAAACCTTGACACTCCAAGTGTGGTCTTACAGTTCCCA
GAATCCCCTCCTTGAGGAGCTGCTAGAAATGCTGAATCTCAAGCATCTCCCCAGACCTAC
TGAATCAGAGCCTGCATCTGAAGCTTTACGGTGTACAAGCTGTTTTATGTGAAGGCTGAA
GTTTAAAAAGCACTGCATTAAGCGTTAGTTTGGTATAAACTGCCCTGACTGAACTGGT
GTGTCCACTTAGCTTGCATGATGACTGTTGCTTTGATGATGAAGCTTACACGGGTAGAT
CCTTTGAGTGAGTGATCTGACATGATTCTCCTTTGCTAAGGCATCTAGATTTCAGTGCACA
ACTTACAGCTGTTGCTTTAGGGGAAATACAACGTAAAATTAATAAAAAACATAGTCTC
TTCTTATGATAACATGGAACGATGGCAAAATAGATTTTGTAGCACTTGGGTAGGAATTC
TGAATGAAGCAGGCAAAATCTGTTGGCAGTGAAATGATAGGATGTGGTAAAGTTAGAATA
AAATAAACTTAAATGTCTCAAACCTCATGGTATATACTACCAGTTTAATAAATGTTG
TACCTTTGATGATTTGCAGACTACAAGCATCAAGGTGCTGTGTATATATTAATCTTGT
GGAGAATAATACCTTAAAAATGAAATTCAGAAATTTTAAATCAGACAAAGCTTTTGT
GCATGGCCCACTTAAATGGCTATTTGAAATAATGATAGTGGATATAGAAGGATTATTCT
GTAATAGGATGAGACTGTTCCCTTTGTCATGGAGATCATAATCATATTTTGTAAATTTT
TATTATTTTTTGGTTTTGTGCCATCTGCACACTATTACTGGGTAGGTACATGGTTTTT
TTAACATGTTTTATCTTCAAACCTATAAAGGCATGCAAAACAGAAGACAGGTCATTTAT
TTTTCTTCAAAGCATCTAAAATGAGATTTTGATATTTGAGGTCATAAAGAGGTGAGAG
AACAGACAACAGTTGGGAAAGCTATTTCTCTTGAATTTGTTGGCCTTAATTACTACAGT
GTCCTAGTACCACCCATACGTTTCAAAGAAGTAGATCCCTGTAATGCCTTTGTCCTCG
GACTTTTGAGTAAAATAGTAGGGTGTGCTTTGCAAAATGTCATCGTTGATGTTGAGTTTC
AGAGTCTTAAATAGGAAGCTGAAATCTGTATATCGAGATTTGTAATCATCTAAATTC
AGAGTAATGTTTTAGAACTACTGCTTAAGGGATTGGCATTAAAGCCTTTTTTAAAAAGAA
ATGCAATAATTTCTCAAATCCTCACTCATTAGACCTCTACTAATACTAGTGCAGACTTT
TTTTTTTTTTTACCCTAAAGTCTGGAATTCCAAAGAAATGCTTACCATTTCCTCCATTA
TTATAGCCACCTGGAAGCAGTATTCATGTATTAGATCAAAAACACAACAAGAAATATGA
AAGGTTGTTTCTGTTATGCAATGCATGATGACATGAACTTACAGAACAGAGAGAAGGGA
GGCTCCATGTTTATTTAAAGAGGAAATTTTTATTTCTGGTTACCTACTTTTACATGGGT
TACATCAAATCCACGATGAGGTTTAAAAATCTCATAGATAATCAAACGTCATTACTTG
GCTTACTGAAATTCAGACTTTTCTTTTTTCTTCCCTGTTTTTCTCTATCAAATAGAAATC
TTTGAAGAACTACCAGAAACGAGTGGGAAAAACCCGGAGATTCTTCTTAAATTAAG
TTCTATCCCACGGAGGAGTTTATCACCTCAGCAGAGCTTCAGGTTTTCCGAGAACAGAT
GCAAGATGCTTTAGGAAACAATAGCAGTTTCCATCACCGAATTAATATTTATGAAATCAT
AAAACCTGCAACAGCCAACCTCGAAATTCCTCGTGACCAGACTTTTGGACACCAGGTTGGT
GAATCAGAATGCAAGCAGGTGGGAAAGTTTTGATGTCACCCCGCTGTGATGCGGTGGAC
TGCACAGGGACACGCCAACCATGGATTTCGTGGTGAAGTGGCCCACTTGGAGGAGAAACA
AGGTGTCTCCAAGAGACATGTTAGGATAAGCAGGCTTTGACCAAGATGAACACAGCTG
GTCACAGATAAGGCCATTGCTAGTAACTTTTGGCCATGATGGAAAAGGCATCCTCTCCA

TABLE 1-continued

CCTTTTATCTCTAGTTCTATTGTTCTTGATTAGGTGAGAACCATTAGATCATACCCAAC
 TGAGGGGATGGGGTCTTGTGTTGTTCTCCAGCTGTTCTTACCCTCTATTGCCATGGACA
 TGAAGGACAGACTGCACGGTCTTAACATGTTAAAACGAATGACCCATGTTTTCATAT

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 2

<210> SEQ ID NO 1
 <211> LENGTH: 14759
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapien
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (3639)...(3984)
 <221> NAME/KEY: CDS
 <222> LOCATION: (11757)...(12601)

<400> SEQUENCE: 1

ccttggtttt ggggatcatt tgggcaagcc cgaggtgctg tgcattggggg ctcctggaat	60
cctgggaagg gcagaaaacc ttggcccccag actcatcgtg cagcagctct gagcagtatt	120
tgggctgagg agtgacttca gtgaatattc agctgaggag tgacttggcc acgtgtcaca	180
gccctacttc ttggggggcct ggtggaagag ggtggcgtag aaggttccaa ggtcccaaac	240
tggaattgtc ctgtatgctt ggttcacaca gtgcgttatt ttaccttctc ctgagctgct	300
aatcgctgco ctctgagctg ggtgagataa ataccacaag gcacaaagtg attgtacaat	360
aaaaaaaaatca aatccctccc atccatcctt cagtctgcca cacacgcagt ctacgttaca	420
cacatgtcac gtaaagcagg atgacatcca tgtcacatac atagacatat taaccgaaat	480
gtggcccttc ggttgcatat attctcatac atgaatatac ttatagaaat atatgcacat	540
atttttgtat attggatata tttatgtaac tataaattta catgctgatg gatatgaaaa	600
taaatgcata cacatttatg taaaaaaatt tgtacacatg catttacata tgtaataaca	660
tacatctcta tgtattaatg tttaaaaaca ctcaatttcc agcctgctgt tttcttttaa	720
ttttcctcct attccgggga aacagaagcg tggatccccc gtctatgcta tgccaaaata	780
cgctgtaatt gagggtgttt gttttgtttt gttttttgaa atcgtatatt accgaaaaac	840
ttcaaaactga aagtgaata acggggcccag cggggaaata agaggccaga ccctgaccct	900
gcatttgtcc tggatttcgc ctccagagtc cccgcgaggg tccggcgcgc cagctgatct	960
ctcctttgag agcaggaggt ggaggcgcga gcgccccctc tggcggccgc gcgccccgc	1020
cctccgcccc accccgcgcg ggctgcccgc gcgcgccgtc cacaccctcgc cgcgcagctc	1080
ccgcccgcgc ggggatcccc ggcgagcccgc gccgcgaagg gggaggtgtt cggccgcggc	1140
cgggagggag ccggcagggc gcgtcccctt taaaagccgc gagcgcgcgc ccacggcgc	1200
gccgcccgcg tcgcccgcgc cggagtcttc gccccgcgcg gctgcccgcg gctcgcgctg	1260
cgctagtcgc tccgcttccc acaccgccgc ggggactggc agccgcgcgc gcacatctgc	1320
cgccacagcc tccgcccgcg acccgaacgt tctcggggcc agcgcgcgagt ggatcaccgg	1380
ggaccgcgag gcaccccgcg gccgcagacc ccgcgcgggc tggagcaccg gccagagcgc	1440
gccacagcgc cgtggcctct gctgcccggg ctgcgccaga gccgcggagc ggcgcgcaga	1500
gcgcccggga ctccggagcc gatccctagc gccgcgatgc ggagcaccta ctgcaggaga	1560

-continued

tcgggggcct gggacgcgct ggccgaggtg tgatcggacc ccaggctagc cacaaagggc	1620
acttgccccc agggctagga gagcagggg agagcacagc cacccgcctc ggcggcccgg	1680
gactcggcto gactcgcgg agaatgcgcc cgaggacgac ggggcgccag agccgcggtg	1740
ctttcaactg gcgagcgcga atgggggtgc actggagtaa ggacagagtga tgcggggggg	1800
caactcgcct ggcaccgaga tcgccgccgt gcccttcctt gaaccggcg tcgccagga	1860
tggtgcccc gagccatggg ccgcggcgga gctagcgcgg agcggccgac cctcgacccc	1920
cgagtcccgg agccggcccc gcgcggggcc acgcgtccct cgggcgctgg ttctaagga	1980
ggacgacagc accagctctt cctttctccc tcccttccc tgcccgcac tctcccctt	2040
gctcgtggtt gttgtgtgc agcaattggc tggggacttc ttgaactgc agggagaata	2100
acttgcgcac cccactttgc gccggtgct ttgcccagc ggagcctgct tcgccatctc	2160
cgagccccc acgcccctca ctctcggcc ttgcccgaca ctgagacgct gttcccagcg	2220
tgaaaagaga gactgcgcgg ccggcaccgg ggagaaggag gaggcaaaga aaaggaacgg	2280
acattcggto cttgcgccag gtcctttgac cagagttttt ccatgtggac gctctttcaa	2340
tgagcgtgto cccgcgtgct tcttagacgg actgcggtct cctaaaggta gaggacgcgg	2400
gccagggccc ggggtgggtg gtgggtggga gggggatttg ggcagccact gcggtagagc	2460
ccttccttac gtccaggcca gaagtaaca gaccctctc cagtccacgt gcaacggagc	2520
cctgcagggg ctcccacttc cagctgcccc gggcgaccgt aagcctcacc ctcccggccc	2580
gcactcttcc acccctcttt ctccctctt ccctggaata cttttggagc tgtaaacact	2640
tagatgaggt gttttattta tttatttatt ttttttaat ttttttaaaa acttttttgg	2700
gtcaaagaaa tccctttgag agggtagccc ctgggtttca cccgttagct gagaacctgt	2760
ccgctctgcc atggtgatct ccattcttca agtgtttccg ggagacttgg tttctttgct	2820
cagagccgtg tcccatttag gaaagtacta ggagtttggg gttctccta cttgtttcca	2880
gaaatgcgag gggtcagtac tgaaggatca cttggtactg tgttttaac agctgacacg	2940
tgcatataa gatattcacc atttacgtaa tcccgggaag atacatgtgt atcttgactg	3000
cactgtgggg atgcgggatg gagctgcctt tcgagacacc cctgagggta gggcctggg	3060
acacaagtca taagtggctt cagaagttgt ggccttgagc ttacagggtc tggaagctat	3120
aagggtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt caggaagttc tatacagtgc	3180
ctctaaggaa gtcacatgca ccatttatgt gtgtttatat gccagacagc gctcagcact	3240
ccgcatthtg gtttgatag gggacgcagg gtgtcagatc aagcgtggt tttcccaggt	3300
tcccggcatt ggctgtcagc gctgtgtcac acacaaaaa gtgacagtca ttggcgtgg	3360
tttggttggg ggggagggca aatcccaaat ctgatgtcag acgagctaag cgttggtatg	3420
gagcgataaa tcatctggtt caggaacttg ggacccttca ttatcccaaa cgtttgagct	3480
tcggtcggto ttacctagac tcgtgagtgt gccaaaccag gaggcatcc tggaggaggc	3540
acgccagcca aatgggagac cgggcccggg gggcgcgagg ggggaggact gggcggggaa	3600
ctcgggtgac tcacgtcggg cctgtccgca ggtcgacc atg gtg gcc ggg acc cgc	3656
	Met Val Ala Gly Thr Arg
	1 5
tgt ctt cta gcg ttg ctg ctt ccc cag gtc ctc ctg ggc ggc gcg gct	3704
Cys Leu Leu Ala Leu Leu Leu Pro Gln Val Leu Leu Gly Gly Ala Ala	
10 15 20	
ggc ctc gtt ccg gag ctg ggc cgc agg aag ttc gcg gcg gcg tcg tcg	3752
Gly Leu Val Pro Glu Leu Gly Arg Arg Lys Phe Ala Ala Ala Ser Ser	

-continued

25	30	35	
ggc cgc ccc tca tcc cag ccc tct gac gag gtc ctg agc gag ttc gag			3800
Gly Arg Pro Ser Ser Gln Pro Ser Asp Glu Val Leu Ser Glu Phe Glu			
40	45	50	
ttg cgg ctg ctc agc atg ttc ggc ctg aaa cag aga ccc acc ccc agc			3848
Leu Arg Leu Leu Ser Met Phe Gly Leu Lys Gln Arg Pro Thr Pro Ser			
55	60	65	70
agg gac gcc gtg gtg ccc ccc tac atg cta gac ctg tat cgc agg cac			3896
Arg Asp Ala Val Val Pro Pro Tyr Met Leu Asp Leu Tyr Arg Arg His			
	75	80	85
tca ggt cag cgg ggc tca ccc gcc cca gac cac cgg ttg gag agg gca			3944
Ser Gly Gln Pro Gly Ser Pro Ala Pro Asp His Arg Leu Glu Arg Ala			
	90	95	100
gcc agc cga gcc aac act gtg cgc agc ttc cac cat gaa g gtgagcatg			3994
Ala Ser Arg Ala Asn Thr Val Arg Ser Phe His His Glu			
105	110	115	
gagcagggcg tgggggocggg gagtcaccct gcaaagccct ccaccgtggg cagactgcag			4054
ccgtccctgt agaggcagct tggccggggc accagcggac gttccactc ttgcttctgt			4114
actatcgttt ctgaatctga ttttaactca ctgcttctgt ggtgggggag ccagggattc			4174
cccttagta actccgcacc ctcttctctg cttgcagcca gaagagctac tctcctgga			4234
agaattggag agaaatcaag tgatggggaa gatgagggca aaaggcatgc ctctagtcat			4294
ctaaactgac aagaattcca cagagggaaa aggagaaaaa gggaggcaga ttgagatttc			4354
tttaagtctg tttggaagct tttgctctat aaatctgccc cttagccag ggttttaggg			4414
tagacagagc caagggcaga gttttcagag atagtattga aaaatcaaag ccagggccc			4474
caaagtcttt ctaatttata gttgatctgg gcctgggttg gaagattttg aatcccaatc			4534
taatccccgt gggagatcaa tactacaatc aatcttattg tttccacaat gactttcttg			4594
tctgtgctt aaatctgaga taggctctga gtagagacaa ggcaagcctt cagataaaag			4654
cgtttgtagc agctgcctgt ttttttttca tgtgcaccga aatgtggatt tttttttctt			4714
ttatgatact acatgtgggt tttctaaggt gggatatttc tgcttgtttc atcagaaggg			4774
catttagtgg actgaaatg tcttacagca gctattgagg tctgctgtac ctaagtcttt			4834
agagcaatta gtcaaaaata tgttccactt caattctttt tctacacttt taaatgcttc			4894
tttgcttaa tacattttaa atagagcatg gtttcttca attcctagaa aagagtacaa			4954
aagtgtatat cacagagcaa ccaactggca gatatttggg gagttgggag tgaagtctc			5014
tttctgctt ttcctgctt aggtggtaaa tttcaagtgg gaaatttaca ctgataatag			5074
actaatggga aatggcactt ccagatggtt tctcccagtg tgaagggtga cttatacttg			5134
tgagagtatt tgttggtaat ggaataaagt cccaaaggca agccacatag cagaagatac			5194
gttctcattg aggcagctac acattacgac ggggacactg aattgatcat cagttcattt			5254
acaagcacat ttctaagtga ggtgctctct gctagcagaa atcagatttg aaaggcagta			5314
agatctcact ccaactcttc agaattcacc caatgaaagc agaaatcacc tgttgtcata			5374
tgtaaaattt ggtgtatgt gtacattctg ccatctaac cctgaaatga ttatagatcc			5434
agctaactat tcccagtaa tgctgattag aatacttttt tttttgtata ggaatgtaat			5494
aagaacaact gtttagaca cctctctctg aaatttagca tggaagctct caactttatt			5554
tttaaggcct ggaagatgct gtgtctctgt tacaacttaa aaggaagatc atttaagtta			5614
gttaacacct aaaacattcc attgtgtgag gattttatca gtgatgtctg catattctca			5674
tcattcatct agaagtggtt tgatcagaac taaacaggct acacgttatt caactgtggt			5734

-continued

atTTTaaCtT aaaaagcatg cttgagttta taaaatcaga atttatactt ttgtgagtgt	5794
aaatgttacc tgagaaacag tacagaagtg accaacttga ttaaaatcaa cttgtaataa	5854
cttcaggtct taatgcagtt agataatgga gaaaagctat gtaatTTTgc cccaaatttc	5914
aactaatcca tttcttTgtct cattatgact aatatatcat ccttaactctg gatggatata	5974
gcactTTTTt caagactaat cattgttTgta tacaccCagg atttgctTTt gataaacatc	6034
cttTgtccat gcatgccacg aaaaagTtt ttggtaaacc atgtgatgaa ggtTgtggc	6094
tcaagaacag aatttagTtt ctacagcatt aatgagcatt tatttGaaa aagaccataa	6154
agaccaatc ataagaatta cctgttgggt tttctTTgta ggtgtgatcg aatggTTtg	6214
tggaattact cgacgagata tcatgatagc attctTTcaa ccaatatgag tataatgcga	6274
ccatatcata ggggatctga gacagaatta tcagtTgtat ttttctatt gaattTTgtc	6334
tagtcttTt tccagtggct tttattTggg agaatatcag cttTgctaaa atgttattgt	6394
tttcaagatc attaaaaagt gcttcagcta catagacctt tggaaactgc cattgaacat	6454
agaaaagtca gttctgcaag tggaaagagt gtttTgtgta ttgctgtagt tggaaacaca	6514
ttgaaactgg ttgacttCac tggccctcca aaaagtctt atgctTTTTt gtcagatggg	6574
agagagaaag accaggtgct tcttTtctc ctactctga aggacacagt cttctTTcta	6634
catgaaataa ctggattatt tgcctctTg actgaagctt tcaaatagag attaacctc	6694
ttccacaaa tataattatt atgaaaatat ccatataata gaaaagtTca agaaataact	6754
attgcctgc attagagact ttgtggcaca aattccccg tgcaaacaac agattTggac	6814
acatagatcc accaaaacca atacttacct ggtatggTtC cctagtggc ccaggtattt	6874
cattgtcatt acagaggcca cattaagtag gaaaattact ctattTgga atggtTgtg	6934
agattgaggc tttggTgtcc agtgatactt ccttggcact gacatttTc gttccacctg	6994
tttttagtg gttccctaa atttctotta atcccttTg agtgaactat tttgcgtTt	7054
tagacttTgct ctttTgtat ttctactgag acaataagag aatatttcat cattccgaag	7114
gtgtTgtgt taagggTgg cagaggcaa atcagggtTg ttgatgacaa ccatgctctc	7174
tattcttTta tttgocattc cctgtTgta tttttTtaa aatggaatgt ttttaacctt	7234
ttgtattTga tattttttt ctcttTgac agttgtctgt tattttatta tctggaaaat	7294
cttatattat actcagctc tttcattTtTg tTtagggca gtgacttcca gccttactga	7354
ttgccagcat atccccaggt tttgtTgtTg ttgtTgtTgt tttactggag attttttagc	7414
ccaaagtTg ttttaaaatc ctCGaagcat aacggtaact tactttttTg ataaaactta	7474
ccatactTta tttagaacaa agggcagcc acaaaatagc agtggctcct tataaaatag	7534
acacattcca gtgggcccc tcactttTct gctcattTct gtctgttctg tccatcatac	7594
ctaagtcata tatttctgtt catttagTtG ggacagaact caccaatgt tatcattTga	7654
ctaaatataa atgtgcccct aatggTttTg actttTgctt aagttttTga gtcctcatgt	7714
atgttagTga gtgccatcta gtagccagaa atttgggaac tggctgggca tgatggctaa	7774
tacctgtaat cccagcactt tgggaggctt aggtgggtTg atcactTgag gtcaggagtT	7834
ccagaccagc ctggccaaca tggTgaaacc acatctctac taaaatataa aaaaatagcc	7894
aggatgatg gcccatgct gtaatccgag ctaattggga ggctgagatg ggaggactgc	7954
ttgaacctgg gaggtggagg ctgctgtgag ccaagattgt gccactgcac tccagcctgg	8014
gcaacagagt gagaccctgt gtcacaaaa caagaacaa aacaaaacaa aagacaagaa	8074

-continued

acctgagaag	cgcagtagat	tcaattatat	atatctactt	ttaatttgct	agctctgtga	8134
ccttaggaaa	gttacataac	ctctctgaac	tgcaactggt	tcatttacia	aatggagata	8194
atgatagttt	ttctctaatt	ggtttgtgt	gagataattc	atataaagct	gatggtgcca	8254
gattacacto	aaaaaaaa	ttcagctgtc	attatcatta	tgacttcttt	tgtaaatggt	8314
atagccttto	cttctctag	gaaaaggagg	ccagagtgga	cctaggctga	ctgagagaa	8374
tcagctcagt	cttttgaatt	atthttgagg	agaggaatga	ttgatatagt	atagattatt	8434
aaattaggac	ttcacttttg	gagaaaagtt	cagatatcat	tggtgtotta	tttttcttca	8494
ctttcccaca	tttttgagc	catagctcca	tccatttggt	taagaactta	gaagctcaca	8554
aactcgggtc	aaagacaggt	cgaatcctc	aaatccctta	agaacttcag	cttattcagg	8614
aagggatatt	tacagaaaac	tagcaattgt	ataagtctcc	aaaaaagcat	acattacttg	8674
aggatccata	tatttttggc	atcctcagg	ttgctgtgat	gatttataga	aggtttgttt	8734
athtaattta	ctttatttca	aataggtttt	aatttttga	ccottaagaa	aagattcgta	8794
ctcttccctg	gcagattaaa	gaaaatgagc	gtatattccc	taacctggc	cagttacttt	8854
cctgggtttg	agggtttctg	tgaacgtcta	acttacctct	gtgacctgtt	tctgcaacca	8914
gggtgtttgc	aatggatgct	tttgtcttga	ggatgggacc	ttcaagaaa	cagattcact	8974
gaggtgcagt	gggaaggtca	gagaaagatc	ttcgtatcgc	ctattattat	ttgctcgtct	9034
atthtttctc	ctttcttaag	gccactaact	gattctcctt	tgtaaggct	gcctacttcc	9094
actgagacct	tgaaccacat	gaaattggtg	ttgtctgtgt	ttctgggtcaa	atagtggoaa	9154
ttttgtatga	ttcaactctg	tcatttaatt	ttttgggagg	ttattattct	atthcatacc	9214
ttttttatac	ccatctctct	tacttcattt	acctgtccct	catacttgac	ttgtagcttg	9274
tcocctcoact	gtcatcgtct	ggccatgtgg	gtgtgtactg	gtgtgcgaga	gagagaatgt	9334
gtgagaatgt	atgtttcttt	atgcattggg	atthtagggt	ttcttgcaa	ttgtgatthc	9394
tctgggcact	tttgtaata	tagctagtca	gcgagtgtc	tagataattt	tccttgctc	9454
cccctctttg	aaagaaaaga	gggtgttctt	agatgtattc	ttatcagata	agccagtagc	9514
tcaggtgctg	gtctggcttt	ggtgtcattg	gggtctgagg	ttgotgactt	ttaccttctc	9574
tgctgaaaa	ttaccttcag	cagaaacgtc	tgaattgcaa	ggagaaggag	aaaaaacag	9634
gccaacacaca	gtccttggtta	ctccttgga	gccactgaga	agagtcocag	ttcaaatggt	9694
cagaaggtta	ttttaatgat	tgtgtctggc	ctaaagtacc	attagcttcc	agtgaggttt	9754
agaatgtgga	tggatcctga	aaggatttcc	ccagagggtt	ggattaatag	gcacaagggga	9814
accctaaagg	actctattgg	cctgatactc	cccatatcca	cgtagaagag	ctthtagaaga	9874
accttctgtt	ctgagacctt	ggctggggcc	accagagct	ggccattca	actcttactc	9934
ctttgccacc	actaatggtt	cttctactag	tttttatatt	atthaacaaa	aaggcacttt	9994
aaaaatgcac	tcctggcaat	ctatactgga	atatgaaaa	catgctgcaa	aaccttgaca	10054
ctccaagtgt	ggtcttacag	ttcccagaat	cccctccttg	aggagctgct	agaaatgctg	10114
aatctcaagc	atctcccag	acctactgaa	tcagagcctg	catctgaagc	tttacggtgt	10174
acaagctggt	ttatgtgaa	gctgaagttt	gaaaagcact	gcattaaagc	gttagtttgg	10234
tataaactgc	cctgactgaa	cttggtgtgt	ccacttagct	tgcatgatga	ctggtgcttt	10294
gatgatgaag	gcttacacgg	gtagatcctt	tgagtgagtg	atctgacatg	attctccttt	10354
gctaaggcat	ctagattcag	tgcaaacctt	acagctgttt	gtctthaggg	gaaatacaac	10414
tgtaaaatta	ataaaaaacat	agtctcttct	tatgataaca	tggaacgatg	gcaaaataga	10474

-continued

ttttgttagc acttggttag gaattctgaa tgaagcaggc aaattctggt ggcagtgaaa	10534
tgataggatg tggtaaaagt agaataaaat aaacttaaat gtctcaaact ctcatggtat	10594
atactaccag ttttaataata atgtgtgacc tttgatgatt tgcagactac aagcattcaa	10654
ggtgctgtgt tataatattac ttgcttgag aataataact cttaaaaatt gaaattcaga	10714
aattttaaat cagacaaagc ttttgtgcat ggcccactta aatggctatt ttgaaataat	10774
gatagtggat atagaaggat tattctgtaa taggatgaga ctgttccttt tgtcatggag	10834
atcataatca tatttttgta aatttttatt atttttttgg ttttgtgtcc atcctgcaca	10894
ctattactgg gttagtacct ggttttttaa catggtttat ctttcaaac tataaaggca	10954
ttgcaaacag aagacaggtc atttattttt cttccaaaag catctaaaat gagattttga	11014
tatttgaggt cataaagagg tgagagaaca gacaacagtt gggaaagcta tttctcttga	11074
aattgtttgg cottaattac tacagtgtcc tagtaccacc catacgtttc caaagaagta	11134
gatccctgta aatgcctttg tctctggact tttgagtaaa atagtagggt gtgctttgca	11194
aaatgtcatc gttgatgttg agtttcagag tctttaatta ggaagctgaa atctgtatat	11254
cgagatttgt aaatcatcta aattgcagag taatgtttta gaatactgct taagggattg	11314
gcattaaagc ctttttttaa aaagaaatgc aataatttc tcaaatctc actcattaga	11374
cctctactaa ctatagtgtc gacttttttt tttttttacc ctaaagtctg gaattccaaa	11434
gaaatgcttc accatttccc ccattattat agccacctgg aagcagtatt catgtattag	11494
atcaaaaaca caacaaagaa ttatgaaagg ttgtttcctg gtatgcaatg catgatgaca	11554
tgaacttaca gaacagagag aaggaggct ccatgtttat ttaaagagga aatttttatt	11614
ttctggttac ctacttttac atgggttaca tcaaatccca cgatgaggtt taaaaattct	11674
catagataat caaacgtcat tacttggtt actgaaattc agacttttct tttttcttcc	11734
ctgtttttct ctatcaaatt ag aa tct ttg gaa gaa cta cca gaa acg agt	11785
Glu Ser Leu Glu Glu Leu Pro Glu Thr Ser	
120 125	
ggg aaa aca acc cgg aga ttc ttc ttt aat tta agt tct atc ccc acg	11833
Gly Lys Thr Thr Arg Arg Phe Phe Asn Leu Ser Ser Ile Pro Thr	
130 135 140	
gag gag ttt atc acc tca gca gag ctt cag gtt ttc cga gaa cag atg	11881
Glu Glu Phe Ile Thr Ser Ala Glu Leu Gln Val Phe Arg Glu Gln Met	
145 150 155	
caa gat gct tta gga aac aat agc agt ttc cat cac cga att aat att	11929
Gln Asp Ala Leu Gly Asn Asn Ser Ser Phe His His Arg Ile Asn Ile	
160 165 170	
tat gaa atc ata aaa cct gca aca gcc aac tcg aaa ttc ccc gtg acc	11977
Tyr Glu Ile Ile Lys Pro Ala Thr Ala Asn Ser Lys Phe Pro Val Thr	
175 180 185	
aga ctt ttg gac acc agg ttg gtg aat cag aat gca agc agg tgg gaa	12025
Arg Leu Leu Asp Thr Arg Leu Val Asn Gln Asn Ala Ser Arg Trp Glu	
190 195 200 205	
agt ttt gat gtc acc ccc gct gtg atg cgg tgg act gca cag gga cac	12073
Ser Phe Asp Val Thr Pro Ala Val Met Arg Trp Thr Ala Gln Gly His	
210 215 220	
gcc aac cat gga ttc gtg gtg gaa gtg gcc cac ttg gag gag aaa caa	12121
Ala Asn His Gly Phe Val Val Glu Val Ala His Leu Glu Glu Lys Gln	
225 230 235	
ggt gtc tcc aag aga cat gtt agg ata agc agg tct ttg cac caa gat	12169
Gly Val Ser Lys Arg His Val Arg Ile Ser Arg Ser Leu His Gln Asp	
240 245 250	

-continued

gaa cac agc tgg tca cag ata agg cca ttg cta gta act ttt ggc cat	12217
Glu His Ser Trp Ser Gln Ile Arg Pro Leu Leu Val Thr Phe Gly His	
255 260 265	
gat gga aaa ggg cat cct ctc cac aaa aga gaa aaa cgt caa gcc aaa	12265
Asp Gly Lys Gly His Pro Leu His Lys Arg Glu Lys Arg Gln Ala Lys	
270 275 280 285	
cac aaa cag cgg aaa cgc ctt aag tcc agc tgt aag aga cac cct ttg	12313
His Lys Gln Arg Lys Arg Leu Lys Ser Ser Cys Lys Arg His Pro Leu	
290 295 300	
tac gtg gac ttc agt gac gtg ggg tgg aat gac tgg att gtg gct ccc	12361
Tyr Val Asp Phe Ser Asp Val Gly Trp Asn Asp Trp Ile Val Ala Pro	
305 310 315	
ccg ggg tat cac gcc ttt tac tgc cac gga gaa tgc cct ttt cct ctg	12409
Pro Gly Tyr His Ala Phe Tyr Cys His Gly Glu Cys Pro Phe Pro Leu	
320 325 330	
gct gat cat ctg aac tcc act aat cat gcc att gtt cag acg ttg gtc	12457
Ala Asp His Leu Asn Ser Thr Asn His Ala Ile Val Gln Thr Leu Val	
335 340 345	
aac tct gtt aac tct aag att cct aag gca tgc tgt gtc ccg aca gaa	12505
Asn Ser Val Asn Ser Lys Ile Pro Lys Ala Cys Cys Val Pro Thr Glu	
350 355 360 365	
ctc agt gct atc tcg atg ctg tac ctt gac gag aat gaa aag gtt gta	12553
Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp Glu Asn Glu Lys Val Val	
370 375 380	
tta aag aac tat cag gac atg gtt gtg gag ggt tgt ggg tgt cgc tag	12601
Leu Lys Asn Tyr Gln Asp Met Val Val Glu Gly Cys Gly Cys Arg *	
385 390 395	
tacagcaaaa ttaaatacat aaatatatat atatatatat attttagaaa aaagaaaaa	12661
acaaacaac aaaaaaaccc caccagtt gacactttaa ttttcccaa tgaagacttt	12721
atztatggaa tggaatggaa aaaaaacag ctattttgaa aatataatta tatctacgaa	12781
aagaagtgg gaaaacaaat attttaatca gagaattatt ccttaaagat ttaaaatgta	12841
tttagttgta cattttatat gggttcaacc ccagcacatg aagtataatg gtcagattta	12901
ttttgtattt atttactatt ataaccactt tttaggaaaa aaatagctaa tttgtattta	12961
tatgtaatca aaagaagtat cgggtttgta cataattttc caaaaattgt agttgttttc	13021
agttgtgtgt atttaagatg aaaagtctac atggaaggtt actctggcaa agtgcttagc	13081
acgtttgctt ttttgcatg ctactgttga gttcacaagt tcaagtccag aaaaaaaag	13141
tgataatcc actctgtgta ctttcaagat tattatatta ttcaattctc aggaatgttg	13201
cagagtgatt gtccaatcca tgagaattta catccttatt aggtggaata tttggataag	13261
aaccagacat tgctgatcta ttatagaaac tctcctctg cccottaatt tacagaaaga	13321
ataaagcagg atccatagaa ataattagga aaacgatgaa cctgcaggaa agtgaatgat	13381
ggttgtgtgt tcttctttcc taaattagtg atcccttcaa aggggctgat ctggccaaag	13441
tattcaataa aacgtaagat ttcttcatta ttgatattgt ggtcatatat atttaaaatt	13501
gatatctcgt ggcctcatc aagggttga aatttatttg tgttttacct ttacctcatc	13561
tgagagctct ttattctcca aagaaccag ttttctaact ttttgcccaa cacgcagcaa	13621
aattatgca acgtgttttt ctgcccacc tctgttctct gacotatcag cttgcttttc	13681
ttccaaggt tgtgtgtttg aacacatttc tccaaatggt aaacctattt cagataataa	13741
atatcaaatc tctggcattt cattctataa agtccaacct gtaagagaaa atggtgcatt	13801
tgtatagcgc ttacaatgat gacctgtgt ttgcattttt gtttctgaag ttatatattt	13861
tagaggggt ggggaaagg taatgaatgg ctggaaaatt gcaggcaagt tatttgataa	13921

-continued

gtcatatttg cactaaaggt gttaccagtg atttagtatt tttcaaatga acttctttgg 13981
 ggcaagaaga ttaagggaa aactaaagcc tacaaaacaa gcaaaacctg gataaccgga 14041
 gataaagttt cagagataat agcccatgca acagaggcaa cggtgccaga aaattagaaa 14101
 gggaaagtgt cggagatcag cttctataag aacatctgcc agttggactg acgcccacac 14161
 agaatgaagt caaattagcg tgctcagatt gaacacttac cagagtgtca gggcttctgt 14221
 accctggggt agaatcagac caaggaaggg ttcagcagat gttcataaga gcagggcacc 14281
 cacaactacc cactatttta ctggcagtat tttaggtcag tttccaggac tttgcatccc 14341
 ctctgatcct gccatgcatg attggtgaaa cctacctcta atctccttgg aattggctaa 14401
 aaaacagtgt gtttataatg gaacagactg ttataatcaa attcttctta ggaattaact 14461
 tttgatgact atgagcttag ttacagttcg gaggttatga ggttatgtaa accttatctt 14521
 taaatgtgca tgacagttat cttttactaa tgctgggttaa cttttaaact cttgcagctc 14581
 ctttttatct ctagttctat tgttcttgat taggtgagaa ccattagatc ataccacact 14641
 gaggggattg gggctctgtt tgttctocag ctggtcttca ccctctattg ccatggacat 14701
 gaaggacaga ctgcacggtc ttaacatggt aaaacgaatg acctatggtt tctcatat 14759

<210> SEQ ID NO 2
 <211> LENGTH: 396
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 2

Met Val Ala Gly Thr Arg Cys Leu Leu Ala Leu Leu Leu Pro Gln Val
 1 5 10 15
 Leu Leu Gly Gly Ala Ala Gly Leu Val Pro Glu Leu Gly Arg Arg Lys
 20 25 30
 Phe Ala Ala Ala Ser Ser Gly Arg Pro Ser Ser Gln Pro Ser Asp Glu
 35 40 45
 Val Leu Ser Glu Phe Glu Leu Arg Leu Leu Ser Met Phe Gly Leu Lys
 50 55 60
 Gln Arg Pro Thr Pro Ser Arg Asp Ala Val Val Pro Pro Tyr Met Leu
 65 70 75 80
 Asp Leu Tyr Arg Arg His Ser Gly Gln Pro Gly Ser Pro Ala Pro Asp
 85 90 95
 His Arg Leu Glu Arg Ala Ala Ser Arg Ala Asn Thr Val Arg Ser Phe
 100 105 110
 His His Glu Glu Ser Leu Glu Glu Leu Pro Glu Thr Ser Gly Lys Thr
 115 120 125
 Thr Arg Arg Phe Phe Phe Asn Leu Ser Ser Ile Pro Thr Glu Glu Phe
 130 135 140
 Ile Thr Ser Ala Glu Leu Gln Val Phe Arg Glu Gln Met Gln Asp Ala
 145 150 155 160
 Leu Gly Asn Asn Ser Ser Phe His His Arg Ile Asn Ile Tyr Glu Ile
 165 170 175
 Ile Lys Pro Ala Thr Ala Asn Ser Lys Phe Pro Val Thr Arg Leu Leu
 180 185 190
 Asp Thr Arg Leu Val Asn Gln Asn Ala Ser Arg Trp Glu Ser Phe Asp
 195 200 205
 Val Thr Pro Ala Val Met Arg Trp Thr Ala Gln Gly His Ala Asn His
 210 215 220

-continued

Gly	Phe	Val	Val	Glu	Val	Ala	His	Leu	Glu	Glu	Lys	Gln	Gly	Val	Ser
225					230					235					240
Lys	Arg	His	Val	Arg	Ile	Ser	Arg	Ser	Leu	His	Gln	Asp	Glu	His	Ser
			245						250					255	
Trp	Ser	Gln	Ile	Arg	Pro	Leu	Leu	Val	Thr	Phe	Gly	His	Asp	Gly	Lys
			260					265					270		
Gly	His	Pro	Leu	His	Lys	Arg	Glu	Lys	Arg	Gln	Ala	Lys	His	Lys	Gln
		275					280					285			
Arg	Lys	Arg	Leu	Lys	Ser	Ser	Cys	Lys	Arg	His	Pro	Leu	Tyr	Val	Asp
	290					295					300				
Phe	Ser	Asp	Val	Gly	Trp	Asn	Asp	Trp	Ile	Val	Ala	Pro	Pro	Gly	Tyr
305					310					315					320
His	Ala	Phe	Tyr	Cys	His	Gly	Glu	Cys	Pro	Phe	Pro	Leu	Ala	Asp	His
				325					330					335	
Leu	Asn	Ser	Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu	Val	Asn	Ser	Val
			340					345					350		
Asn	Ser	Lys	Ile	Pro	Lys	Ala	Cys	Cys	Val	Pro	Thr	Glu	Leu	Ser	Ala
		355					360					365			
Ile	Ser	Met	Leu	Tyr	Leu	Asp	Glu	Asn	Glu	Lys	Val	Val	Leu	Lys	Asn
	370					375					380				
Tyr	Gln	Asp	Met	Val	Val	Glu	Gly	Cys	Gly	Cys	Arg				
385				390						395					

What is claimed is:

1. A method of diagnosing a susceptibility to osteoporosis in an individual, comprising detecting a polymorphism in a human BMP2 gene of SEQ ID NO: 1, wherein the presence of a "T" at nucleotide position 11980 is indicative of a susceptibility to osteoporosis, compared with an individual having an "A" at nucleotide position 11980.
2. A method of diagnosing a susceptibility to osteoporosis in an individual, comprising detecting a polymorphism in a human BMP2 gene of SEQ ID NO: 1, wherein the polymorphism is selected from the group consisting of A to G at nucleotide position 420; A to G at nucleotide position 472; G to C at nucleotide position 1464; G to A at nucleotide position 1722; C to G at nucleotide position 1914; T to G at nucleotide position 3747; A to G at nucleotide position 3899; G to T at nucleotide position 3918; A to T at nucleotide position 11980; C to T at nucleotide position 12571; T to C at nucleotide position 13066; A to G at nucleotide position 13209; C to A at nucleotide position 13296; at least one deletion in nucleotides at positions 13533-13536 and combinations thereof.
3. A method of diagnosing a susceptibility to osteoporosis in an individual, comprising detecting a polymorphism in a human BMP2 gene of SEQ ID NO: 1, wherein the presence of a "G" at nucleotide position 3747 is indicative of susceptibility to osteoporosis compared to an individual having a "T" at nucleotide position 3747.
4. The method of claim 1, wherein the polymorphism is detected in a sample from a source selected from the group consisting of: blood, serum, cells and tissue.
5. The method of claim 1, wherein detecting the polymorphism is performed by a method selected from the group consisting of: allele-specific hybridization, using oligonucleotide arrays, direct manual sequencing, automated fluorescent sequencing, single-stranded conformation polymorphism assays, clamped denaturing gel electrophoresis,

- denaturing gradient gel electrophoresis, mobility shift analysis, restriction enzyme analysis, heteroduplex analysis, chemical mismatch cleavage, RNase protection assays, use of polypeptides that recognize nucleotide mismatches and allele-specific PCR.
6. The method of claim 2, wherein the polymorphism is detected in a sample from a source selected from the group consisting of: blood, serum, cells and tissue.
7. The method of claim 2, wherein detecting the polymorphism is performed by a method selected from the group consisting of: allele-specific hybridization, using oligonucleotide arrays, direct manual sequencing, automated fluorescent sequencing, single-stranded conformation polymorphism assays, clamped denaturing gel electrophoresis, denaturing gradient gel electrophoresis, mobility shift analysis, restriction enzyme analysis, heteroduplex analysis, chemical mismatch cleavage, RNase protection assays, use of polypeptides that recognize nucleotide mismatches and allele-specific PCR.
8. The method of claim 3, wherein the polymorphism is detected in a sample from a source selected from the group consisting of: blood, serum, cells and tissue.
9. The method of claim 3, wherein detecting the polymorphism is performed by a method selected from the group consisting of: allele-specific hybridization, using oligonucleotide arrays, direct manual sequencing, automated fluorescent sequencing, single-stranded conformation polymorphism assays, clamped denaturing gel electrophoresis, denaturing gradient gel electrophoresis, mobility shift analysis, restriction enzyme analysis, heteroduplex analysis, chemical mismatch cleavage, RNase protection assays, use of polypeptides that recognize nucleotide mismatches and allele-specific PCR.

* * * * *

专利名称(译)	人骨质疏松症基因		
公开(公告)号	US6630304	公开(公告)日	2003-10-07
申请号	US09/661887	申请日	2000-09-14
[标]申请(专利权)人(译)	解码遗传学私营有限责任公司		
申请(专利权)人(译)	deCODE Genetics公司EHF.		
当前申请(专利权)人(译)	deCODE Genetics公司EHF.		
[标]发明人	STYRKARSDOTTIR UNNUR JOHANNSDOTTIR VALA DROFN		
发明人	STYRKARSDOTTIR, UNNUR JOHANNSDOTTIR, VALA DROFN		
IPC分类号	C07K14/435 C07K14/51 C12Q1/68 G01N33/53 A61K48/00 A61P19/10 A61P43/00 C07K14/705 C07K16/18 C07K19/00 C12N1/15 C12N1/19 C12N1/21 C12N5/10 C12N15/09 C12P21/02 C07H19/00		
CPC分类号	C07K14/51 C12Q1/6883 C12Q2600/158 C12Q2600/156 A61P19/10		
审查员(译)	陈, CHRISTINA		
外部链接	Espacenet USPTO		

摘要(译)

公开了人BMP2基因在骨质疏松症中的作用。还公开了使用BMP2基因中的多态性诊断, 预测临床过程和治疗骨质疏松症的方法。