



US007271251B2

(12) **United States Patent**  
**Boren et al.**

(10) **Patent No.:** **US 7,271,251 B2**  
(45) **Date of Patent:** **Sep. 18, 2007**

(54) **HELICOBACTER PYLORI ADHESIN BINDING GROUP ANTIGEN**

FOREIGN PATENT DOCUMENTS

FR 2724936 A1 9/1994

(76) Inventors: **Thomas Boren**, Torelvagen 68 S-906 28, Umea (SE); **Anna Arnqvist**, O. Brinkvagen 56 S-903 21, Umea (SE); **Lennart Hammarstrom**, Immunologi Huddinge Sjukhua S-141 86, Huddinge (SE); **Staffan Normark**, Vallhallavagen 126 S-1141 41, Stockholm (SE); **Dag Ilver**, Bankgatan 18 S-902 35, Umea (SE)

(Continued)

OTHER PUBLICATIONS

Amara, A et al (1995), Feb. 13, Voo. 185(3) Neuroscience letters (Ireland), pp. 147-150, Molecular detection of methionine in rat brain using specific antibodies, abstract only.\*

(Continued)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 35 days.

*Primary Examiner*—Mark Navarro  
*Assistant Examiner*—Ginny Allen Portner  
(74) *Attorney, Agent, or Firm*—Birch, Stewart, Kolasch & Birch LLP

(21) Appl. No.: **10/761,201**

(22) Filed: **Jan. 22, 2004**

(57) **ABSTRACT**

(65) **Prior Publication Data**

US 2004/0234529 A1 Nov. 25, 2004

**Related U.S. Application Data**

(62) Division of application No. 09/202,178, filed on Feb. 10, 1999, now Pat. No. 6,709,656.

(51) **Int. Cl.**  
*A61K 39/40* (2006.01)  
*C07K 16/12* (2006.01)

(52) **U.S. Cl.** ..... **530/388.4**; 424/164.1; 530/388.7

(58) **Field of Classification Search** ..... 530/350, 530/388.1, 388.4; 424/184.1, 164.1, 190.1, 424/234.1, 235.1; 435/7.32; 514/25  
See application file for complete search history.

A novel *Helicobacter pylori* blood group antigen binding (BAB) adhesin protein was isolated and purified, whereby said protein or fractions thereof bind specifically to fucosylated blood group antigens. The protein sequence of said adhesin is disclosed in this application. Simultaneously the DNA sequences for two genes, babA and babB, producing highly similar proteins, are disclosed. Said adhesin and/or DNA is useful for diagnose and therapy and/or prophylaxis directed against *H. pylori* induced infections, e.g. gastritis and acid peptic disease, i.e. active vaccination.

A new immunoglobulin composition, which exhibits specific activity to a Lewis<sup>b</sup> antigen binding *Helicobacter pylori* adhesin, or preferably, monoclonal and/or polyclonal antibodies to said adhesin offer a new and more efficient method of treatment and/or prevention of gastrointestinal diseases, caused by *Helicobacter pylori* or other *Helicobacter* species, i.e. passive vaccination.

(56) **References Cited**


U.S. PATENT DOCUMENTS

4,444,879 A \* 4/1984 Foster et al. .... 435/7.95  
5,258,177 A \* 11/1993 Uemura et al. .... 424/176.1

(Continued)

**7 Claims, 10 Drawing Sheets**

Antibody preparations used for Western blot detection of the BabA adhesin and *H. pylori* protein antigens;



## U.S. PATENT DOCUMENTS

5,262,156	A *	11/1993	Alemohammad	435/7.32
5,538,729	A	7/1996	Czinn et al.	
5,625,124	A	4/1997	Falk et al.	
5,837,502	A	11/1998	Smith	
5,846,751	A	12/1998	Pronovost et al.	
5,883,079	A	3/1999	Zopf et al.	
5,897,475	A	4/1999	Pace et al.	
6,051,416	A *	4/2000	Pace et al.	435/252.1
6,410,719	B1 *	6/2002	Boren et al.	536/23.7
6,576,244	B1 *	6/2003	Weltzin et al.	424/234.1
6,709,656	B1 *	3/2004	Boren et al.	424/190.1
2004/0265427	A1 *	12/2004	Boren	426/52
2005/0009037	A1 *	1/2005	Chang et al.	435/6

## FOREIGN PATENT DOCUMENTS

WO	WO92/01718	2/1992
WO	WO97/12908	4/1995

## OTHER PUBLICATIONS

- Aucher, Pe tal, J. Clinical Microbiology, vol. 36(6), pp. 931-936, Apr. 1998.\*
- Bai, Y et al., World J. Gastroenterology, vol. 10(17), pp. 2560-2562, 2004, Cloning and expression and immunogenicity of *Helicobacter pylori* BabA2 gene.\*
- Bina, J et al, Journal of Bacteriology, May 2000, vol. 182(9), pp. 2370-2375, Functional expression in *Escherichia coli* and membrane topology of porin HopE, a member of a large family of conserved proteins in *Helicobacter pylori*.\*
- Boren and Falk, 1995, *Helicobacter pylori* binds to blood group antigens. Scientific American Science and Medicine, Sep./Oct. pp. 28-37.\*
- Cover, Timothy L, Crisp abstract, identification No: 5R01DK053623-07, *H. pylori* factors associated with peptic ulcer disease (abstract only).\*
- Doig, P et al, Journal of Bacteriology, Oct. 1995, vol. 177(19), pp. 5447-5452, Isolation and Characterization of a conserved porin protein from *Helicobacter pylori*.\*
- Durrant, LG et al, Hybridoma, vol. 12(6), 1993, Development of Second generation Monoclonal antibodies recognizing LewisY/B antigen by Anti-idiotypic immunization.\*
- Exner, M.M et al, Infection and Immunity, vol. 63(4), pp. 1567-1572, Apr. 1995, Isolation and Characterization of a Family of Porin Proteins from *Helicobacter pylori*.\*
- Hennig, E.E. et al, Infection and Immunity, Jun. 2004, vol. 72(6), pp. 3429-3435, Heterogeneity among *Helicobacter pylori* strains in expression of the outer membrane protein BabA.\*
- Pride, David T. et al, Infection and Immunity, vol. 69(3), pp. 1160-1171, Feb. 2001, Allelic Variation within *Helicobacter pylori* babA and babB.\*
- Durrant et al (1993), reference of record.\*
- Essery et al (1994), FEMS Immunology Medical Microbiology, Jun. , vol. 9(1), pp. 15-21, Detection of microbial surface antigens that bind Lewis(a) antigen.\*
- Bina et al , reference of record.\*
- Cover Crisp Document reference of record.\*
- Pride et al (2001) reference of record.\*
- Boren et al (1995) reference of record.\*
- Boren et al (1995, reference of record).\*
- Durrant et al (1993, reference of record).\*
- Huang, J et al (1992), Journal of General Microbiologi, vol. 138, pp. 1503-1513, Identification of erythrocyte-binding antigens in *Helicobacter pylori*.\*
- Aspholm-Hurtig et al, Science, vol. 305, Jul. 23, 2004, pp. 5619-522.\*
- Hennig, Ewa E. et al, Infection and Immunity, Jun. 2004, pp. 3429-3435, vol. 72(6), Heterogeneity among *Helicobacter pylori* strain in Expression of the outer membrane protein BabA.\*
- Pride, David T et al, Infection and Immunity, Feb. 2001, pp. 1160-1171, vol. 69(2), Allelic Variation within *Helicobacter pylori* babA and babB.\*
- Henning, Ewa E et al, Infection and Immunity, vol. 74(5), pp. 3046-3051, May 2006, Multiple Chromosomal Loci for the babA Gene in *Helicobacter pylori*.\*
- Solnick, Jay V e tal, PNAS, Feb. 17, 2004, vol. 101(7), pp. 2106-2111, Modification of *Helicobacter pylori* outer membrane protein expression during Experimental infection of rehesus macaques.□□\*
- Boren et al., Science, vol. 262, p. 1892, (Dec. 1993).
- Alkout et al., Dialog Information Services, File 34, Scisearch, Dialog accession No. 15596361, Gastroenterology, vol. 112, No. 4, pp. 1179-1187, (1997).
- Clyne et al., Dialog Information Services, File 34, Scisearch, Dialog accession No. 15313697, FEMS Immunology and Medical Microbiology, vol. 16, No. 2, pp. 141-155, (1996).
- Chan et al., Gycobiology, vol. 5, No. 7, pp. 683-688, (1995).
- Alkout et al., Gut, vol. 37, (Suppl. 7), p. A21, Abstract 84, (Jul. 1995).
- Ho et al., European Journal of Gastroenterology and Hepatology, vol. 7, pp. 121-124, (1995).
- Huang et al., Journal of General Microbiology, vol. 138, pp. 1503-1513, (1992).
- McKibbin et al., The Journal of Biological Chemistry, vol. 257, No. 2, pp. 755-760, (Jan. 1982).
- Oriol, Transplantation proceedings, vol. 19, No. 6, pp. 4416-4420, (Dec. 1987).
- Smith, Australian Vet. Pract., vol. 26, No. 3, pp. 140-145, (Sep. 1996).
- Chmiela et al., FEMS Immunology and Medical Microbiology, vol. 10, Nos. 3-4, pp. 307-326, (Feb. 1995).
- Lelwala-Guruge et al., FEMS Immunology and Medical Microbiology, vol. 11, No. 1, pp. 73-77, (Mar. 1995).
- Jiang et al., Molecular Microbiology, vol. 20, No. 4, pp. 833-842.
- Exner et al., Infection and Immunity, vol. 63, No. 4, pp. 1567-1572, (Apr. 1995).
- Doig et al., Journal of Bacteriology, vol. 177, No. 19, pp. 5447-5452, (Oct. 1995).
- Hirmo et al., Glycoconj . J., vol. 13, No. 6, pp. 1005-1011, (Dec. 1996).
- Huang et al., Journal of General Microbiology, vol. 138, pp. 1503-1513, (1992).
- Cameron et al., Dialog Information Services, File 34, Scisearch, Dialog accession No. 14645778, Infection and Immunity, vol. 64, No. 3, pp. 891-896, Mar. 1996.
- Farokhnia, Antibody specific for a plasmid encoded protein of *Helicobacter pylori*, A Thesis for the degree of master of Science in the graduate School of the Texas Woman's University, (Dec. 1995).
- Amano, Akita J. Med., vol. 24, pp. 100-108, English translation, (1997).
- Sjostedt et al., Ann. Surg., vol. 207, No. 3, pp. 341-346, abstract, (Mar. 1998).
- Blakely et al., J. Med. Microbiol., vol. 15, No. 4, pp. 519-529, abstract, (Nov. 1982).
- Huang et al., FEMS Microbiology Letters, vol. 56, No. 1, pp. 109-112, (1988).
- Newell, Journal of General Microbiology, vol. 133, pp. 163-170.
- Essery et al., FEMS Immunology Med. Microiol., vol. 91, No. 1, pp. 15-21, (Jun. 1994).
- Lelwala-Guruge et al., Zbl. Bakt, vol. 280, pp. 93-106, (1993).
- Falk et al., Proc. Natl. Acad. Sci., vol. 90, pp. 2035-2039, (Mar. 1993).
- Boren et al., Trends in Microbiology, vol. 2, No. 7, pp. 221-228, (Jul. 1994).
- Smith et al., The Lancet, vol. 343, No. 8896, pp. 543, abstract, (Feb. 1994).
- Fauchere et al., Microbial Pathogenesis, vol. 9, pp. 427-439, (1990).
- Enders et al., Infection Immunity, vol. 63, No. 7, pp. 2473-2477, (Jul. 1995).

Moran, FEMS Immunology and Medical Microbiology, vol. 10, pp. 271-280, (1995).

Lingwood et al., Lancet, vol. 2, No. 8657, pp. 238-241, (Jul. 1989).

Thomas Boren et al., "Attachment of *Helicobacter pylori* . . .," Science, Dec. 17, 1993, vol. 262, pp. 1892-1895.

Mark A. Agius et al., "Monoclonal anti-idiotypic antibodies against . . .," The Journal of Immunology, Jan. 1, 1988, vol. 140, No. 1, pp. 62-68.

\* cited by examiner




Fig. 1A




Fig. 1B




Fig. 1C



Fig.2A




Fig.2B



Fig.2C




Fig.2D







Fig. 3



\* Biotin tagged adhesin protein

Fig.4A

Fig.4C



A = Adhesin

Fig.4B






Fig. 6

Antibody preparations used for Western blot detection of the BabA  
adhesin and *H. pylori* protein antigens;

Human Sera      Chicken Antibody      Bovine Antibodies  
I      II      III      I      II      III      II/III



Lanes: 1 2 3 4 5 6 7 8 9 10 11

Fig.7




Fig.8

**HELICOBACTER PYLORI ADHESIN BINDING  
GROUP ANTIGEN**

This application is a Divisional of application Ser. No. 09/202,178, filed on Feb. 10, 1999, now U.S. Pat. No. 6,709,656 the entire contents of which are hereby incorporated by reference and for which priority is claimed under 35 U.S.C. § 120; and this application claims priority of Application No. 9602287-6 filed in Sweden on Jun. 10, 1996 and Application No. 9701014-4 filed in Sweden on Mar. 19, 1997 under 35 U.S.C. § 119.

## FIELD OF THE INVENTION

The present invention relates to materials and methods for prevention, treatment and diagnosing of infections caused by *Helicobacter pylori*. More specifically the present invention relates to polypeptides and antibodies useful in vaccines for the treatment and prevention of pathologic infections caused by *Helicobacter pylori* strains. The present invention specifically relates to a bacterial blood group antigen binding adhesin (BAB-adhesin). The present invention further relates to polynucleotides useful for the recombinant production of said polypeptides and for use in immunisation therapies. In addition, it relates to polypeptides, antibodies, and polynucleotides used for the detection of said bacteria.

The present invention further relates to new immunoglobulins, which exhibit specific activity to a blood group binding adhesin, expressed by *Helicobacter pylori*, methods for the production of said immunoglobulins, their isolation and use. The present invention further relates to the treatment and prevention of *H. pylori* induced infections in the gastrointestinal tract.

## BACKGROUND OF THE INVENTION

*Helicobacter pylori* is a causative agent for acid peptic disease and the presence of this organism is highly correlated to the development of gastric adenocarcinoma. Bacterial adherence to the human gastric epithelial lining was recently shown to be mediated by fucosylated blood group antigens.

Recent research has focused on the role of *Helicobacter pylori* in the development of ulcers in the gastric mucosa. Recent findings show a strong connection between *H. pylori* and chronic, active gastritis and gastric ulcers. Furthermore, there appears to be a strong correlation between ventricular cancer and gastric ulcers. Traditional treatment of gastric ulcers has involved gastric resection, the administration of bismuth compositions, the administration of H<sub>2</sub>-blockers and the administration of pH-buffering agents, to mention a few examples.

More recently, various forms of treatment have been supplemented with the administration of antibiotics. One problem with presently known treatments is the risk for treatment failure. Furthermore, not only do microbes develop antibiotic resistance, the antibiotics administered often upset the natural balance of benign microbes, colonising the intestinal tract. This leads to diarrhoea and other signs of intestinal discomfort, in addition to destabilising the benign flora in the intestines. Other treatments, e. g. H<sub>2</sub>-blockers often require life-long medication to prevent the recurrence of disease.

The foregoing, together with the fact that *H. pylori* is very widely spread among humans—according to a conservative estimate approximately 60% of all adult humans in the

industrialised countries are infected—makes the diagnosing, prevention and treatment of *H. pylori* infections an urgent task.

Further, the fact that developing countries frequently lack the resources for conventional treatment of gastric ulcers further underlines the importance of finding new ways of treatment and prevention of *H. pylori* induced infections. It is obvious, for many reasons, that disease prevention with vaccines is a preferable mode. A vaccine would provide an easily administered and economical prophylactic regimen against *H. pylori* infections. An effective vaccine against *H. pylori* is nevertheless presently lacking.

## STATE OF THE ART

*H. pylori* colonises the human gastric mucosa, in an equilibrium between adherence to the epithelial surface mucous cells and the mucous layer lining the gastric epithelium. Once infected, bacteria seems to colonise for a lifetime. Attachment to the epithelial lining protects the bacteria from the anti-microbial effects of the acidic gastric juice of the stomach lumen, as well as from physical forces such as peristalsis. For survival in this hostile ecological niche, *H. pylori* has developed a battery of virulence factors; such as production of the enzyme urease that buffers the micro-environment around the bacteria and the polar flagellae to ensure high motility, a prerequisite in an ecological niche where the turnover of the mucous layer is in the range of hours. A subset of *H. pylori* strains produces the vacuolating cytotoxin, VacA, and the cytotoxin associated antigen CagA.

Attachment is essential for colonisation of the epithelial lining and bacteria express surface associated adhesion molecules that recognise specific carbohydrate or protein receptors on the cell surfaces or mucous lining. The specificity in this interaction in combination with the genetically regulated receptor distribution results in a restricted range of cell lineages and tissues available for colonisation. Several putative receptor structures have been described for *H. pylori*, such as the hemagglutinin-sialic acid, sulphated glycoconjugates and sulphatides. Recently, the fucosylated blood group antigens H-1 and Lewis<sup>b</sup> were described (Borén et al., Science, 262, 18921993), mediating specific adherence of *H. pylori* to human and rhesus monkey gastric surface mucous cells in situ. The H-1 and Lewis<sup>b</sup> antigens are part of the blood group antigens that define blood group O in the ABO system.

Surface-exposed proteins are often constituents of the outer membrane. The outer membrane has a structural role and acts as a selective barrier, determining what enters the cell and what molecules are secreted. One class of outer membrane proteins are called porins, and create hydrophilic pores through the outer membrane where specific metabolites, such as sugar molecules, can cross. Recently the finding of a number of outer membrane proteins in *H. pylori*, was reported, which proteins were suggested to constitute a family of porin proteins.

The BAB adhesin has previously been identified and shown to be localised on the bacterial surface of *H. pylori* (SE 9602287-6). The blood group binding activity was shown to be pH dependent and the present inventors present evidence that the binding affinity to the Lewis<sup>b</sup> receptor reveals a high equilibrium constant. For the purification of the BAB adhesin, a crosslinker-labelled receptor conjugate was used in order to mediate specific transfer of biotin to the adhesins on the bacterial surface. Thereafter the biotin-labelled adhesin could be extracted by streptavidin coated

magnetic beads. Determination of the amino terminal amino acid sequence of the purified BAB adhesin exhibit homologies to outer membrane proteins of *H. pylori* porins.

Intensive research has been directed to the immunological treatment and prevention of *H. pylori* induced infections. EP 0 484 148 (Ando & Nakamura) describes a method for treating and/or preventing upper gastrointestinal disease in mammals, said method comprising orally administering to a patient in need thereof an effective amount of a pharmaceutical composition comprising anti-*Helicobacter pylori* polyclonal immunoglobulins and a pharmaceutically acceptable carrier. Said description further dwells on the combination of said treatment in combination with the administration of antibiotics. As the method of producing said polyclonal antibodies, EP 0 484 148 describes the isolation and purification of anti-*H. pylori* immunoglobulins from the sera and milk of mammals. *H. pylori* itself was not found in the stomachs of cows, goats, sheep, swine or horses, according to EP 0 484 148, but it was assumed that these animal species have colonizing microorganisms with antigenic determinants similar to those of *H. pylori* because they have immunoglobulins which cross-react to strains of *H. pylori* found in humans. Preferably, according to EP 0 484 148, large mammals, e.g. pregnant cows, are immunized with whole cells of *H. pylori* and the immunoglobulins subsequently extracted from the milk or colostrum. In the immunization experiments, NCTC Strain 11362 and clinical isolate *H. pylori* No. 153 were used to trigger the production of immunoglobulins. On the other hand, NCTC Strain 11637 was used for analysing purposes. Immunization is claimed to yield an anti-*H. pylori* titer in the milk of such magnitude, that daily doses of 0.01–0.1 g/day immunoglobulin composition, are sufficient for successful therapy. The claimed interval of 0.01–0.1 g/day is however not supported by the experiments presented by Ando & Nakamura and so low doses have hitherto not proven efficient in clinical tests. The doses actually used in example 5 and 7 are in the order of magnitude of 1 g/day, i.e. 10-fold the upper limit of the given interval. Furthermore, it is very unlikely, that unspecific immunoglobulin mixtures as those manufactured by Ando & Nakamura, would be effective in claimed doses as similar doses are ineffective against other gastrointestinal pathogens. The simultaneous administration of antibiotics, extensively discussed in the description, underlines the insufficiency of the disclosed immunoglobulins.

EP 0 469 359 (Cordle & Schaller) likewise describes the immunization of mammals, preferably pregnant cows, with formalin killed *H. pylori* bacteria (ATCC Strain 26695). Anti-*H. pylori* polyclonal antibodies were isolated and purified from the milk and finally fed to piglets, in amounts of about 0.5 g immunoglobulins, three times daily. The results were assessed by determination of the number of biopsy specimens, which were positive for Gram-negative bacteria after the trial. Gram-negative bacteria was found in 78% of the piglets fed a non-immune nutrient but only (sic!) in 35% of the piglets fed a nutrient containing so called specific anti-*H. pylori* antibodies.

Anti-*H. pylori* polyclonal antibodies, effective to cause aggregation of *H. pylori*, have thus been administered orally as a regimen in the treatment and prevention of *H. pylori* induced infections in the gastrointestinal tract. Nevertheless, as also noted in EP 0 484 148 A1, it is still not clear, how many antigenic determinants are present on the surface of *H. pylori*. The occurrence of a wide variety of *H. pylori* strains, makes questionable the practical efficiency of any polyclonal immunological therapy according to the state of the art. Immunization using whole bacteria will always trigger

a highly polyclonal immuneresponse with a low level of antibodies against a given antigenic determinant. This is underlined e.g. by the results presented by Cordle & Schaller, where, although the number of *Helicobacter* positive biopsies were reduced, complete cure was not obtained through the treatment according to their invention.

It is notable, that the dose of immunoglobulin needed for oral prophylaxis or therapy has not yet been clearly defined. In a normal human adult, approximately 5 g IgA is produced and secreted at mucosal surfaces each day. Obviously, doses of this magnitude are economically and practically unfeasible for large-scale therapy or prophylaxis. In studies on the effect of oral immunoglobulin on rotavirus infection, daily doses in the interval of 600 to 9000 mg have been tried in clinical tests. Successful intervention has also been reported when treating *H. pylori* and cryptosporidial infections with daily administrations of 3 to 15 g immunoglobulin from immunized cows (Hammarström et al., Immunol Rev, 139 (1994) 43–70). Generally speaking, all studies hitherto point to the necessity of using high doses of immunoglobulins when trying to combat an ongoing infection. The need for more specific immunoglobuline preparations, allowing the use of smaller doses, is thus an urgent one.

To maximize the potency of an immunological regimen for the treatment and prevention of *H. pylori*, it is of great importance to find a specific conserved antigenic determinant, which plays a central role for the pathogenicity of *H. pylori*. Using such an antigenic determinant would make it possible to produce highly specific and therapeutically efficient novel polyclonal and/or monoclonal immunoglobulin preparations.

#### SUMMARY OF THE INVENTION

The above problem of providing specific, cost-efficient and therapeutically superior immunoglobulin preparations for the treatment and prevention of *H. pylori* has now been solved through the composition and methods according to the attached patent claims. The present inventors have now surprisingly shown, that highly specific and therapeutically efficient polyclonal and/or monoclonal immunoglobulin preparations can be provided through the immunization of an animal with an adhesin protein, specific for *H. pylori*. The invention will now be described in closer detail with reference to the attached, non-limiting figures and examples.

One objective of the present invention was to further purify and characterize the *H. pylori* blood group antigen binding (BAB) adhesin to make possible the development of methods and materials for specific and selective diagnosing and treatment of *H. pylori* induced infections and related diseases and the development of said methods and materials. A further and equally important objective was to determine the DNA sequences of the genes involved in the expression of this protein. These objectives were fulfilled through the protein, the DNA and the methods and materials specified herein The DNA sequences are SEQ ID NOS: 1 and 2, disclosing the babA (SEQ ID NO:1) and babB (SEQ ID NO:2) sequences, respectively. The full protein sequences are disclosed in SEQ ID NOS: 3 and 4.

#### DESCRIPTION OF THE FIGURES

FIG. 1A) illustrates the bacterial binding to soluble blood group antigens. *H. pylori* strains were incubated with <sup>125</sup>I-labeled blood group antigen glycoconjugates and bound <sup>125</sup>I-activity was measured (Note the absence of blood group antigen binding shown for strains MO19 and 26695.).

FIG. 1B) illustrates a receptor displacement assay. Strain CCUG 17875 was first incubated with 10 ng <sup>125</sup>I-labeled Le<sup>b</sup> antigen glycoconjugate and the complex was then challenged (1 h) with an excess of unlabeled Le<sup>b</sup> or Le<sup>a</sup> glycoconjugate, before the <sup>125</sup>I-activity in the bacterial pellet was measured. Concentrations of the unlabeled glycoconjugate ranged from 50 ng to 8 μg and C) shows the results of a Scatchard analysis of the *H. pylori*-Le<sup>b</sup> antigen interaction. Bacterial binding to the Le<sup>b</sup> glycoconjugate was titrated to an affinity constant (K<sub>a</sub>) value of 8×10<sup>-10</sup> M<sup>-1</sup> (13).

FIG. 2: Upper panel: Prevalence of the BabA adhesin in the bacterial population. Cells of strain CCUG 17875 were incubated with biotinylated Le<sup>b</sup> (A) or Le<sup>b</sup> (B) glycoconjugate. Bound biotinylated Lewis-conjugate was detected with FITC-labeled streptavidin (green fluorescence) and bacteria were counterstained with propidium iodine (red fluorescence). Lower panel: Localisation of the BabA adhesin. For electron microscopy (15) cells of strain CCUG 17875 were incubated with biotinylated Le<sup>b</sup> (C) or Le<sup>a</sup> (D).

FIG. 3 shows the characterization of the molecular weight of the BabA adhesin by the use of receptor overlay analysis (A, B) and receptor activity directed affinity tagging of BabA (C).

FIG. 4 shows receptor activity directed affinity tagging and protein purification of the BabA (SEQ ID NOS: 7 and 8) adhesin.

FIG. 5 shows the translated amino acid sequences for the babA (SEQ ID NO: 7) and babB (SEQ ID NO: 9) genes, corresponding to the N-terminal domain of the BabA adhesin.

FIG. 6 shows the procentual inhibition of *H. pylori* binding to <sup>125</sup>I-labeled Lewis b antigen for different preparations as a function of the antibody titre.

FIG. 7 shows a Western blot detection of the BabA adhesin by the different antibody preparations.

FIG. 8 shows four Western blot analyses of *H. pylori* proteins by the different antibody preparations.

#### DESCRIPTION OF THE INVENTION

The blood group antigen binding adhesin, BabA, has now been biochemically characterized and purified by a novel technique, receptor Activity Directed Affinity Tagging (Retagging). Two genes, babA and babB were found to code for two different but very similar proteins. The present invention thus comprises a novel blood group antigen binding adhesin. The DNA sequences are disclosed in SEQ ID NO:1 (babA) and SEQ ID NO:2 (babB). The protein sequences are disclosed in SEQ ID NOS: 3 and 4. The invention also includes any pharmaceutical composition comprising said adhesin protein and/or fractions thereof. Examples of such pharmaceutical compositions are for example medicaments for the prevention or treatment of *Helicobacter pylori* induced gastritis, gastric and duodenal ulcers and gastric adenocarcinoma. Optionally said pharmaceutical composition additionally encompasses pharmaceutically acceptable excipients.

Further, the present invention comprises the BAB-adhesin gene or genes for expression of an adhesin protein according to the invention. Said invention also comprises a novel method for the isolation and purification of said adhesin. The disclosed genes are contemplated to function as a cassette system, the organism alternating between these to avoid immunity in the host. It is very likely, that homologies of the disclosed sequences exist and additionally supplement said cassette function in other strains of *H. pylori*. Also, genes corresponding to a homology of the first 40 amino acids or

genes, corresponding to a homology of the last, about 300 amino acids, can function to this effect. It is further highly likely, that *Helicobacter pylori* is able to switch between several genes, similar to the disclosed genes, in a so-called cassette system.

The invention additionally comprises monospecific antisera produced using the novel adhesin protein and/or fractions thereof. Said monospecific antisera is preferably produced according to any suitable, conventional method for producing monospecific antisera in vitro or in vivo, e.g. by inoculating a suitable animal. Such methods are familiar to a person skilled in the art. Antibodies raised in a suitable animal or in the patient to be treated, can subsequently be administered locally, e.g. orally to the patient.

The invention further comprises the use of said monospecific antisera for the manufacturing of a test kit for quantitative or qualitative determinations of adhesin protein or fractions thereof in cells, tissues or body fluids.

The invention further comprises the use of said adhesin protein or corresponding DNA for use in therapy or immunisation and/or in the manufacture of compositions for said uses. The invention specifically encompasses the use of said DNA for immunisation therapy and for the manufacture for compositions for such therapy. Preferably, in an immunisation therapy where said composition is administered orally to a patient, the adhesin protein, fractions thereof or said DNA is administered in combination with a pharmaceutically suitable immunostimulating agent. Examples of such agents include, but are not limited to the following: cholera toxin and/or derivatives thereof, heat labile toxins, such as *E. coli* toxin and similar agents. The composition according to the present invention can further include conventional and pharmaceutically acceptable adjuvants, familiar to a person skilled in the art of immunisation therapy. Preferably, in an immunisation therapy using the inventive DNA or fractions thereof, said DNA is preferably administered intramuscularly, whereby said DNA is incorporated in suitable plasmid carriers. An additional gene or genes encoding a suitable immunostimulating agent can preferably be incorporated in the same plasmide.

Said immunisation therapies are not restricted to the above-described routes of administration, but can naturally be adapted to any one of the following routes of administration: oral, nasal, subcutaneous and intramuscular. Especially the oral and nasal methods of administration are promising, in particular for large-scale immunisations.

The present inventors have surprisingly shown, that highly specific and therapeutically efficient polyclonal and/or monoclonal immunoglobulin preparations can be provided through the immunisation of an animal with an adhesin protein or fractions thereof, specific for *H. pylori*. When considering immunisation against *H. pylori*, it is worth noting that the infection is known to be lifelong despite a vigorous immune response in the gastric mucosa. An increased local production of IgA in the mucosa is not necessarily enough and the administration of monospecific antibodies directed against a central virulence factor, such as the adhesin according to the present invention, may constitute a more effective approach.

The term "immunisation" refers here to a method for inducing a continuous high level of antibody and/or cellular immunresponse. The term "animal" here preferentially denotes any member of the subphylum Vertebrata, a division that includes all animals, including mammals, which are characterized by a segmented bony or cartilaginous spinal column. All vertebrates have a functional immune system and respond to antigens by producing antibodies. The term

“protein” is used here to denote a naturally occurring polypeptide and the term “polypeptide” is used here in its widest meaning, i.e. any amino acid polymer (di-peptide or longer) linked through peptide bonds. Accordingly the term “polypeptide” comprises proteins, oligopeptides, protein fragments, analogues, muteins, fusion proteins and the like. The term “antibody” as used in this context includes an antibody belonging to any of the immunological classes, such as immunoglobulins A, D, E, G or M. Of particular interest are nevertheless immunoglobulin A (IgA) since this is the principle immunoglobulin produced by the secretory system of warm-blooded animals. However, in cow colostrum, the main antibody class is IgG 1.

Borén et al. have recently isolated and characterized a Lewis<sup>b</sup> binding protein with a molecular weight of about 73500 Da (See the priority applications SE 9602287-6 and SE 9701014-4, which are referred to in their entirety). This adhesin protein is thought to be a conserved structure and specific for pathogenic strains of *H. pylori*. Said protein is specific for at least one of the *H. pylori* strains included in the following group: CCUG 17875, NCTC 11637, A5, P466, G109, G56, Ba 185, Ba 99, 931 and 932.

This adhesin protein or immunologically effective fractions thereof are characterized in that the following amino acid sequence (SEQ ID NO:5) is included:

EDDGFYTSVGYQIGEEAAQMV or homologues thereof.

The following DNA sequence (SEQ ID NO:6) or homologues thereof is included in DNA for expression of said adhesin protein or fractions thereof:

5'-GAAGACGACGGCTTTTACACAAGCGTAG-GCTATCAAATCGGT

GAAGCCGCTCAAATGGTA-3'

According to one embodiment of the invention, a pregnant mammal, preferably a cow or another suitable domestic animal, is immunised with said Lewis<sup>b</sup> binding adhesin protein or fractions thereof. The adhesin protein or fractions thereof is/are preferably injected intramuscularly or subcutaneously in the chosen animal, optionally together with suitable adjuvants. Examples of such adjuvants include, but are not limited to immunostimulating agents such as cholera toxin and/or derivatives thereof, heat labile toxins, such as *E. coli* toxin and similar, conventional agents, such as classical adjuvants including mineral and vegetable oils. Subsequent to the regimen of immunization, comprising a necessary amount of doses, including so called booster-doses, over a time span allowing for optimal immunoglobulin expression, milk or sera is collected from said animal. Preferably the cow colostrum, which is specially high in immunoglobulins, is collected. The specific immunoglobulin fraction according to the present invention is then separated and purified in a conventional manner, e.g. including separation of fats, protein precipitation and concentration by ultrafiltration.

According to another embodiment of the invention, a bird, preferably a chicken or another suitable domestic bird, is immunized with said Lewis<sup>b</sup> binding adhesin protein or fractions thereof. The adhesin protein or fractions thereof is preferably injected intramuscularly or subcutaneously in the chosen bird, optionally together with suitable adjuvants. Examples of such adjuvants include, but are not limited to immunostimulating agents such as cholera toxin and/or derivatives thereof, heat labile toxins, such as *E. coli* toxin and similar, conventional agents, such as classical adjuvants including mineral and vegetable oils. Subsequent to the regimen of immunization, comprising a necessary amount of doses, including so called booster-doses, over a time span allowing for optimal immunoglobulin expression, sera or eggs is/are collected from said animal. Preferably the egg

yolk, which is specially high in immunoglobulins, is collected. The specific immunoglobulin fraction according to the present invention is then separated and purified in a conventional manner, e.g. including protein precipitation and ultrafiltration. Alternatively, the egg yolk being of high nutritional value in addition to containing a high titer of specific antibodies according to the present invention, can be administered as such.

According to a preferred embodiment of the present invention, monoclonal immunoglobulin is produced by establishing transgenic animals. Said transgenic animals can be chosen from the following group of species: mammals, e.g. cow, goat and rabbit, and birds: e.g. chicken, duck, turkey. The mammal most preferably used is cow and the most preferable bird is chicken. Further developments of transgenic animals such as mice and rats could also offer new possibilities. The choice of animal is naturally governed by availability and local adaptation.

According to one embodiment, a stock of transgenic animals according to the present invention, adapted to the local conditions, are kept locally, e.g. in villages in developing countries to function as local units for the production of immunoglobulins for oral administration. For example transgenic cows, goats or chicken are suitable for this purpose and preferably chicken are used. Consumption of the milk or preferably the eggs, produced by the transgenic animals, can help to eradicate presently very difficult infectious diseases, e.g. diseases caused by *H. pylori*.

According to yet another embodiment of the present invention, monoclonal antibodies can be produced using the hybridoma method. The hybridoma method is well known to a skilled worker in the field of biochemistry and it is described e.g. in Galfre, G. And Milstein, C., Preparation of monoclonal antibodies: strategies and procedures (Methods in Enzymology, 73:3-46, 1981). A suitable host animal is immunized with the Lewis<sup>b</sup> binding adhesin protein or fractions thereof. When the immunization is accomplished, the animal is sacrificed, spleen cells collected and fused with cells from a neoplastic cell line, preferably myeloma cells. By choosing the growth conditions, the successfully fused hybridoma cells can be selected. The monoclonal antibodies produced by the hybridoma cell line can then be administered orally in a regimen for treatment and/or prevention of *H. pylori* infections.

Preferably the polyclonal and/or monoclonal antibodies are purified prior to administration and, more preferably, admixed with pharmaceutically suitable carriers and/or adjuvants. Examples of suitable carriers are saline, pharmaceutically acceptable fats, oils, carbohydrates and proteins. The carrier or carriers is/are preferably chosen so that the solubility and absorption of the immunoglobulin in the mucus layer lining the stomach is enhanced. Using suitable adjuvants the stability, therapeutic efficiency and nutritional value of the composition can be improved. To improve stability under storage, the immunoglobulin composition can be lyophilized. Regardless of the exact preparation and formulation, it is of central importance to avoid denaturing the immunoglobulins.

The higher specificity, exhibited by the immunoglobulin preparation of polyclonal and/or monoclonal antibodies according to the invention, makes it possible use substantially lower doses compared to those presently used, thus lowering the cost and improving the availability of the treatment. The use of specific, monoclonal antibodies can make it possible to further lower the doses. The doses are in all cases a function of the antibody titer of the preparation. A high titer naturally allows the use of lower doses.

According to one embodiment of the invention, an immunoglobulin preparation is manufactured as follows: an animal is immunized with a Lewis<sup>b</sup> binding adhesin protein or fractions thereof, expressed by *Helicobacter pylori*, the immunoglobulin fraction is isolated from a excretion of said animal and subsequently purified. The purified immunoglobulin composition is admixed with suitable carriers and adjuvants to form an immunoglobulin preparation for the prevention or treatment of *H. pylori* infections. In cases where the antibody titer is sufficiently high and the other constituents of the immunoglobulin composition isolated from the animal are harmless, for example in the case of colostrum from immunized cows or egg yolk from immunized chicken, there is always the option of administering the colostrum or egg yolk to the patient without any further treatment of the colostrum or egg yolk.

The immunoglobulin composition according to the invention is preferably administered orally to the patient, in the smallest therapeutically or prophylactically effective dose. Presently conceived are doses in the interval of 0.1 to 1000 mg/day, preferably in the interval of 0.1 to 100 mg/day. The chosen doses naturally depend on the antibody titer of the preparation in question. The exact doses and the regimen of administration can be chosen by the physician responsible for the patient, infected by *Helicobacter pylori*. Routine experimentation and later, with increasing experience of this method, empirical information will suffice to establish the required amount. Multiple dosages may be used, as needed, to provide the desired level of therapeutic or prophylactic effect. The immunoglobulin preparations according to the present invention can also, being free from adverse side effects and imposing practically no danger of overdosing, be taken prophylactically or therapeutically by a person without medical supervision.

A therapeutical effect can be attained, except with the specific antibody according to the present invention, also with at least two Fab-fragments of said antibody. Said embodiment is also encompassed by the scope of the present invention.

According to yet another embodiment, avirulent microorganisms, preferably bacteria, are used as expression systems for the specific antibody according to the present invention. An "avirulent microorganism" in this context is a microorganism which has the ability to colonize and replicate in an infected individual, but which does not cause disease symptoms associated with virulent strains of the same species of microorganism. The definition inherent in the GRAS (Generally Regarded As Safe) concept can be applied here. A GRAS-organism is suitable for use according to the present invention, provided that the organism externalises the antibody or can be modified to this effect. The term "microorganism" as used herein includes bacteria, protozoa and unicellular fungi. Preferably, bacteria are used as expression systems, e.g. bacteria of the genus *Lactobacillus*, *Streptococcus* or *Enterobacteriae*. The above mentioned expression system can be utilised in vitro for the production of the specific antibody according to the present invention or, according to a further embodiment of the invention, the micro-organism constituting the expression system can be administered directly to the patient. The micro-organisms can be harvested and administered as such, but they are preferably mixed with a suitable carrier, mixed in a suitable foodstuff, lyophilised, encapsulated or treated in any other conventional way, used for the delivery of viable micro-organisms to the gastrointestinal tract.

According to yet another embodiment, avirulent microorganisms, preferably bacteria, are used as expression sys-

tems for the specific adhesin protein according to the present invention. An "avirulent microorganism" in this context is a microorganism which has the ability to colonize and replicate in an infected individual, but which does not cause disease symptoms associated with virulent strains of the same species of microorganism. The definition inherent in the GRAS (Generally Regarded As Safe) concept can be applied here. A GRAS-organism is suitable for use according to the present invention, provided that the organism externalises the adhesin protein or can be modified to this effect. The term "microorganism" as used herein includes bacteria, protozoa and unicellular fungi. Preferably, bacteria are used as expression systems, e.g. bacteria of the genus *Lactobacillus*, *Streptococcus* or *Enterobacteriae*. The above mentioned expression system can be utilised in vitro for the production of the specific adhesin according to the present invention or, according to a further embodiment of the invention, the micro-organism constituting the expression system can be administered directly to the patient. The micro-organisms can be harvested and administered as such, but they are preferably mixed with a suitable carrier, mixed in a suitable foodstuff, lyophilised, encapsulated or treated in any other conventional way, used for the delivery of viable micro-organisms to the gastrointestinal tract.

The exact doses and the regimen of administration of said micro-organisms can be chosen by the physician responsible for the patient, infected by *Helicobacter pylori*. Routine experimentation and later, with increasing experience of this method, empirical information will suffice to establish the required amount. Multiple dosages may be used, as needed, to provide the desired level of therapeutic or prophylactic effect. The avirulent micro-organism expressing the antibody or adhesin protein according to the present invention can also, being free from adverse side effects and imposing practically no danger of overdosing, be taken prophylactically or therapeutically by a person without medical supervision. A preferred carrier in this specific application is a foodstuff, e.g. a fermented product such as fermented cereal or dairy product.

The creation of previously mentioned expression systems and still earlier mentioned methods of creating hybridomas and transgenic animals can include steps involving recombinant DNA techniques. Recombinant DNA techniques are now sufficiently well known and widespread so as to be considered routine. In very general and broad terms, recombinant DNA techniques consist of transferring part of the genetic material of one organism into a second organism, so that the transferred genetic material becomes a permanent part of the genetic material of the organism to which it is transferred. Methods for achieving this are well known and the mere choice of specific methods for achieving the objectives, set out in the present description and claims, fall under the scope of the invention.

It is possible, that *H. pylori* alone or together with related slow-acting bacteria are involved in the genesis and aggravation of other chronic inflammatory diseases in the gastrointestinal tract. It is obvious for a skilled practitioner how to modify the present invention, within the scope of the claims, to gain utility in the treatment and/or prevention of such diseases. Examples of such diseases are ulcerative colitis, Crohn's disease, sarcoidosis, Wegener's granulomatosis and other vasculitic disorders, as well as various neoplasms, including carcinomas of the colon, pancreas and prostate.

*H. pylori* strain CCUG 17875 was obtained from CCUG, Göteborg, Sweden. Strain A5, a gastric ulcer isolate, from Astra Arcus, Södertälje, Sweden. Strains P466 and MO19 were described previously (Borén et. al, Science, 262, 1892(1993)). Strain 26695 came from Dr. K. A. Eaton, The Ohio State University and its genome was recently sequenced by TIGR, Rockville, Md., USA. The panel of 45 *H. pylori* clinical isolates came from the University Hospital in Uppsala, Sweden. Bacteria were grown at 37° C. in 10% CO<sub>2</sub> and 5% O<sub>2</sub> for 48 h.

All blood group antigen glycoconjugates used, i.e. semi-synthetic glycoproteins constructed by the conjugation of purified fucosylated oligosaccharides to serum albumin were from IsoSep AB, Tullinge, Sweden. The RIA was performed according to Falk et al. (Meth. Enzymol., 236, 353, 1994) with some modifications; the H-1, Le<sup>b</sup>, Le<sup>a</sup>, H-2, Le<sup>x</sup> and Le<sup>y</sup> glycoconjugates were 125I-labeled by the Chloramine T method. 1 ml of bacteria (A600=OD 0.10) was incubated with 300 ng of 125I-labelled conjugate (i.e. an excess of receptors) for 30 min. in phosphate buffered saline (PBS), 0.5% albumin, 0.05% Tween-20 (BB-buffer). After centrifugation, 125I-activity in the bacterial pellet was measured by gamma scintillation counting.

In this study the present inventors' first biochemically characterized and identified the *H. pylori* blood group antigen binding adhesin, BabA. *H. pylori* strains were analysed for binding to soluble <sup>125</sup>I-labeled fucosylated blood group antigens (FIG. 1A). Binding of these strains to the soluble blood group antigens correlate with adherence in situ. The prevalence of blood group antigen binding (BAB)-activity was assessed among 45 clinical *H. pylori* isolates and the majority of the isolates, 71%, express Le<sup>b</sup> antigen binding properties (data not shown). In contrast, none of the reference strains (FIG. 1A), or strains from the panel of 45 clinical isolates, bind to the Le<sup>a</sup>, H-2, Le<sup>x</sup>, or Le<sup>y</sup> antigens. These results support our previous findings of high receptor specificity for the Le<sup>b</sup> and H-1 blood group antigens and demonstrate the high prevalence of BAB activity among clinical isolates.

Based on the presence or absence of virulence factors such as the Cytotoxin associated gene A (CagA) and the Vacuolating cytotoxin A (VacA), *H. pylori* strains are classified as type I or type II strains. *H. pylori* isolates from patients with duodenal ulcers most often express the VacA and the CagA-proteins, i.e. type-I strains. By definition, type II strains express neither markers. Twenty-one clinical isolates previously defined for expression of CagA and VacA were analysed for Le<sup>b</sup> antigen binding properties. Expression of CagA was found to correlate with bacterial binding to the Le<sup>b</sup> antigen (Table 1). The cagA gene belongs to a 40 kb pathogenicity island that encodes components of secretion and transport systems. These findings could indicate functional crosstalk between the cag pathogenicity island and the BabA adhesin gene, for the correct presentation of the BabA adhesin protein in the bacterial outer membrane.

To further characterize BabA, the present inventors determined the affinity constant (K<sub>a</sub>) between BabA and the Le<sup>b</sup> antigen. Since K<sub>a</sub>-values are based on equilibrium conditions (13), the present inventors first analysed the interaction by performing receptor displacement analysis. *H. pylori* CCUG 17875 (positive for Le<sup>b</sup> binding, FIG. 1A) was first incubated with <sup>125</sup>I-labeled Le<sup>b</sup> glycoconjugate. Then unlabeled Le<sup>b</sup> glycoconjugate was added in a dilution series. The unlabeled Le<sup>b</sup> conjugate displaced the bound <sup>125</sup>I-labeled Le<sup>b</sup> glycoconjugate efficiently (FIG. 1B). The results dem-

onstrate that the receptor-adhesin complex formed is in a true state of equilibrium. An equivalent excess of Le<sup>a</sup> glycoconjugate did not dissociate the Le<sup>b</sup>-BabA complex, verifying the high receptor specificity (FIG. 1B). The K<sub>a</sub>-value for the Le<sup>b</sup>-BabA complex of strain CCUG 17875 was titrated with Le<sup>b</sup> glycoconjugate in concentrations from 10 ng to 260 ng/ml and determined to be of a high affinity close to 1×10<sup>10</sup>M<sup>-1</sup>(FIG. 1C). The number of Le<sup>b</sup> glycoconjugate molecules bound to BabA on the bacterial cell surface was calculated to be around 500 per cell. This number is similar to the number of fimbriae organelles on the surface of *E. coli* (14). However, for the BabA adhesin, the calculations are based on the assumption that the majority of bacterial cells in the experiment exhibit an equal number of adhesin molecules with Le<sup>b</sup> antigen binding properties.

TABLE 1

BAB activity among <i>H. pylori</i> Type I and Type II strains			
Type	Strain	BAB activity	
Type I CagA <sup>+</sup> , VacA <sup>+</sup>	CCUG 17874	-	
	G39	-	
	G11	-	
	G20	-	
	G27	+	
	G56	+	
	G106	-	
	G109	+	
	932	+	
	Ba185	+	
	87A300	+	
	931	+	
	Ba99	+	
	Ba179	+	
Ba194	+		
Type Ib CagA <sup>-</sup> , VacA <sup>+</sup>	G12	-	
	Type Id ΔcagA, VacA <sup>+</sup>	G104	-
		Tx30	-
	Type II CagA <sup>-</sup> , VacA <sup>-</sup>	G21	-
		G50	-
		G198	-

To determine the prevalence of BabA in the bacterial population, strain CCUG 17875 was incubated with Le<sup>b</sup> or Le<sup>a</sup> antigens, and bacterial binding activity was visualised by confocal fluorescence microscopy (FIG. 2, upper panel). The analyses demonstrate the high prevalence of BabA binding activity in the bacterial population to the Le<sup>b</sup> antigen (FIG. 2A, green staining) and the complete lack of binding to the Le<sup>a</sup> antigen (FIG. 2B, red counter staining).

Next, the localisation and density of BabA on the bacterial cell surfaces was investigated by immunogold electron microscopy. The Le<sup>b</sup> antigen binding activity of the adhesin localised gold particles to the bacterial outer membrane (FIG. 2C). Individual bacterial cells exhibit an equal number of gold particles (data not shown). When the Le<sup>b</sup> antigen was substituted with the Le<sup>a</sup> antigen (lacking receptor activity), no gold particles were detected (FIG. 2D).

The molecular weight of BabA was characterized by receptor overlay analysis. A protein extract of strain CCUG 17875 was separated on SDS-PAGE and blotted to a membrane. The membrane was incubated with biotinylated Le<sup>b</sup> glycoconjugate, followed by detection with streptavidin and enhanced chemiluminescence. The BabA adhesin activity corresponds to a single 74 kDa band (FIG. 3A). The 40 kDa band is presumably endogenous peroxidase activity since it stains independently of the Le<sup>b</sup> conjugate overlay (lane 3). BabA was very heat stable and could regain some activity

after heating to 97° C. (FIG. 3A, lane 2). The panel of strains exhibited the same molecular weight of BabA (FIG. 3B).

To purify BabA, a novel technique was developed, Receptor Activity Directed Affinity Tagging (ReTagging). Multi-functional crosslinking agents with radiolabeled donating tags have been previously used for receptor-ligand characterization studies. However, the use of affinity donating tags, such as biotin residues presented on flexible spacer structures, adds a new dimension to the applicability of crosslinker technology. An affinity tag, biotin, is transferred to the adhesin protein by the receptor activity and is used for further identification and for affinity purification of the adhesin part of the interaction, by streptavidin (FIG. 4A, B).

A multi-functional crosslinking agent with a biotin donating handle was attached to the Le<sup>b</sup> glycoconjugate. The receptor activity of the Le<sup>b</sup> glycoconjugate subsequently directed the targeted biotin tagging of the BabA adhesin protein (FIG. 4A, B). After crosslinking, the bacterial protein from strains A5, P466, and CCUG 17875 were separated on SDS-PAGE. Immunodetection with streptavidin demonstrated a biotin tagged protein, with the molecular weight of 74 kDa (FIG. 3C) (28). These results support the estimates of the molecular weight from the previous overlay analyses (FIG. 3B). Strain MO19 devoid of Le<sup>b</sup> antigen binding properties (FIG. 3B) (FIG. 1A), was negative for binding also in this set of analyses (FIG. 3C).

The high specificity in the ReTagging technique provided a method for purification of the adhesin protein. Strains CCUG 17875 and A5, that both express the BabA adhesin (FIG. 1A) were processed by the ReTagging technique using crosslinker labelled Le<sup>b</sup> receptor conjugate as the biotin donor. After crosslinking, bacteria were suspended in SDS sample buffer. Streptavidin coated magnetic beads were subsequently added to the solubilised proteins, and biotin tagged BabA was extracted (FIG. 4C). The N-terminal 20 amino acid sequences of the BabA adhesins from strains CCUG 17875 (Australia) and A5 (Sweden) were found to be identical, indicating a biologically conserved protein (FIG. 5). Recently, a series of outer membrane proteins from *H. pylori* were characterized. These proteins, HopA-E, are homologous in their N-terminal sequences to BabA (17), possible indicating a motif for a common secretion mechanism. The biotin tagged BabA adhesin was purified more than 3000-fold from the cell extract, and the yield was calculated to 20%. However, based on data from the Scatchard plots, the level of available BabA adhesin would be about 5-times higher, i.e. approximately 1 mg adhesin/750 mg bacterial protein, which nevertheless could be the reason for the high signal to noise ratio (FIG. 3B). The purification of BabA via the ReTagging technique indicates the potential of this technique for the purification of lectins in complex receptor-ligand interactions, such as the selectin family of cell adhesion molecules.

To clone the gene encoding BabA, the N-terminal 20 aa sequence was utilised for the construction of degenerate primers (18). Two sets of clones were identified which both encode two different but very similar proteins. Both genes code for proteins having almost identical N-terminal domains and identical C-terminal domains, complicating the identification of the functional BabA gene. (FIG. 5). To identify the corresponding gene, the BabA adhesin was purified in large scale by ReTagging. This provided enough protein for an extended amino terminal sequence. 41 amino acids were identified and these residues unambiguously discriminated between the two genes by the differences in aa-positions 28, 35, 37, 38 and 41 (FIG. 5). The gene encoding BabA was named babA and correspond to a basic

protein with a pI of 9.4 and a molecular weight of 78 kDa, i.e. of slightly higher molecular weight than that predicted from the SDS PAGE analyses (FIG. 3). The other gene, babB, corresponds to a protein of a calculated molecular weight of 75.5 kDa. In contrast to babA, the babB gene contains a predicted translational initiation codon (FIG. 5). This could indicate the existence of a third bab gene in the genome or mechanisms for recombination activities. Interestingly, the bab-genes were also detected in strains lacking Lewis b binding properties (data not shown). Gene cassette systems have been shown to promote antigenic variation in *Neisseria gonorrhoeae* (19). Another possibility would be the presence of similar genes coding for adhesins with differences in receptor specificity/host tissue tropism (20). Gene inactivation experiments targeting the bab-genes could aid in understanding this complex gene organisation.

Immunisation experiments with adhesins from *Bordetella pertussis* (21) indicate the potential for outer membrane proteins to act as vaccine candidates (discussed in ref. 22). In a mouse model for persistent *H. pylori* infection, oral immunisation with *H. pylori* antigens proved protective against *H. pylori* infection (10). However, results from animal models are difficult to evaluate for human specific pathogens, such as *H. pylori* and Polio virus. For Polio, an animal model has been achieved by expressing the virus receptor in transgenic mice (23). A similar strategy was taken for *H. pylori*. A transgenic mouse was constructed by the use of an  $\alpha$ 1,3/4-fucosyltransferase, driving the synthesis of the human specific Le<sup>b</sup> antigen in the gastrointestinal tract (24). The Lewis b mouse can be useful for the evaluation of the role of the BabA adhesin as a colonisation/virulence factor and in addition for the evaluation of BabA as a vaccine candidate against acid peptic disease and gastric adenocarcinoma.

In the present study the ReTagging technique was used for the purification of the adhesin part of the microbial receptor-ligand interaction. By the use of purified adhesin/lectin-protein, the ReTagging technique could, in addition, be used to further study the receptor part of the interaction. Identification of the biologically active receptor structure, carrying Le<sup>b</sup> oligosaccharides, would aid in the understanding of the mechanisms supporting the chronic *H. pylori* infection.

Inhibition of *H. pylori* binding to <sup>125</sup>I-labeled Lewis b antigen by preparations is presented graphically, as a function of antibody concentration (mg/ml) in FIG. 6: 1 ml aliquots of *H. pylori* bacteria (A<sub>600</sub>=OD 0.10) were pre-incubated with dilution series of antibody preparations, in 0.01–10 mg/ml for 2 hours in phosphate buffered saline (PBS), 0.5% albumin, 0.05% Tween-20™. Then 500 ng of <sup>125</sup>I-labeled conjugate (i.e. an excess of receptor structure) was added and incubated for 30 minutes. After centrifugation, <sup>125</sup>I-activity in the bacterial pellet was measured by gamma scintillation counting. The Lewis b blood group antigen glycoconjugates used, i.e. semi-synthetic glycoproteins constructed by the conjugation of purified fucosylated oligosaccharides to serum albumin were from IsoSep NB, Tullinge, Sweden.

Western blot detection of the BabA adhesin by the different antibody preparations is presented in FIG. 7: Molecular weight rainbow marker (2 μL) from Amersham, Buckinghamshire, England, was dissolved in SDS sample buffer (lane 1). Approx. 100 ng of purified BabA adhesin (approx. 74 kDa with degradation product of approx. 55 kDa) was dissolved in SDS sample buffer (lane 2). SDS solubilized protein extracts of strain CCUG 17875 were prepared by dissolving the bacterial pellet corresponding to 0.15 ml of bacteria (A<sub>600</sub>=OD 0.10) by SDS sample buffer (lane 3). The

3 protein samples were then boiled at 100° C. for 5 minutes. The proteins were separated on SDS-PAGE, and transferred to a PVDF-membrane for Western blot immuno analysis. Five sets of PVDF-membranes were prepared. The PVDF membranes were blocked/incubated overnight with 4% human sera/plasma, in phosphate buffered saline, from a patient with no *H. pylori* infection, i.e. with no serum antibodies against *H. pylori*. The membrane was then washed in phosphate buffered saline (PBS), 0.5% albumin, 0.05% Tween-20, followed by the addition of the antibody preparations. The sets of membranes were incubated with the following 5 antibody preparations; 1) pooled human sera from *H. pylori* infected patients, diluted 1:500. 2) Chicken antibodies (positive) 1 mg/ml diluted 1:100x, 3) Bovine I preparation of antibodies, 1 mg/ml diluted 1:100x. 4) Bovine II preparation of antibodies, 1 mg/ml diluted 1:100x. 5) Bovine III preparation of antibodies, 1 mg/ml diluted 1:100x (indicated in the figure). These antibodies were incubated with the membrane for 2 hours followed by extensive washings in phosphate buffered saline (PBS), 0.05% Tween-20, followed by the addition of secondary anti-human, anti-chicken, and anti-bovine antibodies labeled with HRP-peroxidase (from DAKO, Denmark), all diluted 1:2000x. Membranes were incubated for 1 hour, followed by extensive washings in phosphate buffered saline (PBS), 0.05% Tween-20. The membranes were developed with enhanced chemoluminescens (ECL) from Amersham. The results show, that the antigenic response against the adhesin is strongly enhanced in the bovine preparations. This finding is also supported by the inhibition data in FIG. 6.

Western blot analyses of *H. pylori* proteins by the different antibody preparations are shown in FIG. 8. 2 clinical isolates (1-2) from Dr. Lars Engstrand, Department of Clinical Microbiology and Cancer epidemiology, University Hospital, Uppsala, Sweden and strain CCUG 17875 (3), from Culture Collection, University of Göteborg, Department of Clinical Bacteriology, Göteborg, Sweden, and strain 52 (4) from Prof. Torkel Wadström, Dept. Medical Microbiology, Lunds University, were prepared for SDS-PAGE electrophoresis. Bacterial pellets corresponding to 0.15 ml of bacteria ( $A_{600} = OD 0.10$ ) were dissolved in SDS sample buffer and heated to 100° C. for 5 minutes. The proteins were separated on SDS-PAGE, and transferred to PVDF-membranes for Western blot immuno analysis. The western blot analyses were as described above, i.e. the sets of membranes were incubated with the following 4 antibody preparations; 1) pooled human sera from *H. pylori* infected patients, diluted 1:500. 2) Chicken antibodies (positive) 1 mg/ml diluted 1:100x, 3) Bovine I preparation of antibodies, 1 mg/ml diluted 1:100x. 4) Bovine III preparation of antibodies, 1 mg/ml diluted 1:100x (indicated in the figure). These antibodies were incubated with the membrane for 2 hours followed by extensive washings in phosphate buffered saline (PBS), 0.05% Tween-20, followed by the addition of secondary anti-human, anti-chicken, and anti-bovine antibodies labeled with HRP-peroxidase (from DAKO, Denmark), all diluted 1:2000x. Membranes were incubated for 1 hour, followed by extensive washings in phosphate buffered saline (PBS), 0.05% Tween-20. The membranes were developed with enhanced chemoluminescens (ECL) from Amersham. The results show, that the chicken antibodies and the bovine preparations reacts nearly identically against all four strains, indicating conserved properties in strains of different geographical origin.

Although the invention has been described with regard to its preferred embodiments, which constitute the best mode presently known to the inventors, it should be understood

that various changes and modifications as would be obvious to one having the ordinary skill in this art may be made without departing from the scope of the invention which is set forth in the claims appended hereto.

## REFERENCES AND NOTES

1. J. R. Warren, *Lancet*, i, 1273, 1983, B. Marshall, *Lancet*, i, 1273, 1983.
2. A. Dubois, *Emerging Infectious Diseases* 1, 79 (1995).
3. M. J. Blaser, *Sci. Amer.* 2, 92 (1996).
4. M. J. Blaser, *Trends Microbiol.* 7, 255 (1993), D. E. Kirschner and M. J. Blaser, *J. Theor. Biol.* 176, 281 (1995).
5. P. Falk, T. Borén, and S. Normark, *Meth. Enzymol.* 236, 353 (1994).
6. D. G. Evans, D. J. Evans Jr., J. J. Moulds and D. Y. Graham, *Infect. Immun.* 56, 2896 (1988), S. Hirno, M. Utt, M. Ringner and T. Wadström, *FEMS Immunol. and Med. Microbiol.* 10, 301 (1995), T. Saitoh, et al *FEBS Lett.* 282, 385 (1991).
7. T. Borén, P. Falk, K. A. Roth, G. Larson, and S. Normark, *Science.* 262, 1892 (1993), P. Falk, et al *Proc. Natl. Acad. Sci. U.S.A.* 90, 2035 (1993).
8. *H. pylori* strain CCUG 17875 was obtained from CCUG, Göteborg, Sweden. Strain A5, a gastric ulcer isolate, came from Astra Arcus, Södertälje, Sweden. Strains P466 and MO19 were described previously (7). Strain 26695 came from Dr. K. A. Eaton, The Ohio State University, and its genome was recently sequenced by The Institute for Genomic Research (TIGR), Rockville, Md. (J.-F. Tomb, et al, abstract 3B: 059, IX International Workshop on Gastrointestinal Pathology and *Helicobacter pylori*, Copenhagen, Denmark, 1996). The panel of 45 *H. pylori* clinical isolates came from the University Hospital in Uppsala, Sweden. Bacteria were grown at 37°C in 10% CO<sub>2</sub> and 5% O<sub>2</sub> for 48 h.
9. All blood group antigen glycoconjugates used, i.e. semi-synthetic glycoproteins constructed by the conjugation of purified fucosylated oligosaccharides to serum albumin (7, 25), were from IsoSep AB, Tullinge, Sweden. The RIA was performed according to ref. 26 with some modifications; The H-1, Le<sup>b</sup>, Le<sup>a</sup>, H-2, Le<sup>x</sup>, and Le<sup>y</sup> glycoconjugates were <sup>125</sup>I-labeled by the Chloramine T method. 1 ml of bacteria ( $A_{600} = OD 0.10$ ) was incubated with 300 ng of <sup>125</sup>I-labeled conjugate (i.e. an excess of receptors) for 30 min. in phosphate buffered saline (PBS), 0.5% albumin, 0.05% Tween-20 (BB-buffer). After centrifugation, <sup>125</sup>I-activity in the bacterial pellet was measured by gamma scintillation counting.
10. A. Covacci, et al, *Proc. Natl. Acad. Sci. U.S.A.* 90, 5791 (1993), M. Marchetti, et al, *Science* 267, 1655 (1995).
11. S. Censini et al, *Proc. Natl. Acad. Sci. U.S.A.* 93, 14648, (1996).
12. Z. Xiang, et al, *Infect. Immun.* 63, 94 (1995).
13. A. G. Scatchard, *Ann. N. Y. Acad. Sci.* 51, 600 (1949).
14. O. Mol, and B. Oudega, *FEMS. Microbiol. Reviews*, 19, 25 (1996).
15. Confocal microscopy was performed on a Nikon/Multiprobe 2001 instrument (Molecular Dynamics, Sunnyvale, Calif.). Electron microscopy was performed on a JEOL 100 CX instrument.
16. J. Brunner, *Trends in Cell Biol.* 6, 154 (1996), J. D. Bleil and P. M. Wassarman, *Proc. Natl. Acad. Sci. U.S.A.* 87, 5563, (1990).

17. M. M. Exner, P. Doig, T. J. Trust, and R. E. W. Hancock, *Infect. Immun.* 63, 1567 (1995), P. Doig, M. M. Exner, R. E. W. Hancock and T. J. Trust, *J. Bacteriol.* 177, 5447 (1995).
18. The BabA N-terminal sequence analysis was used to make degenerate oligonucleotides, which were used in PCR to obtain an amplified fragment from the chromosome of the babA gene. A 59 bp fragment was identified and used as probe for the screening of a low-copy plasmide (pACYC184) library of Sau3A partially digested chromosomal DNA from strain CCUG 17875.
19. P. Hagblom, E. Segal, E. Billyard, and M. So, *Nature*, 315, 156 (1985), R. Haas and T. F. Meyer, *Cell*, 44, 107 (1986).
20. A.-B. Jonsson, D. Ilver, P. Falk, J. Pepose, and S. Normark, *Mol. Microbiol.* 13, 403 (1994), N. Strömberg, P. G. Nyholm, I. Pascher, and S. Normark, *Proc. Natl. Acad. Sci USA* 88, 9340 (1991).
21. A. Kimura, K. T. Mountzouros, D. A. Relman, S. Falkow, J. L. Cowell, *Infect. Immun.* 58, 7 (1990).
22. T. Borén, and P. Falk, *Sci. Amer., Sci. & Med.* 4 (1994), L. S. Tompkins and S. Falkow, *Science* 267, 1621 (1995).
23. R. B. Ren, et al, *Cell* 63, 353 (1990).
24. P. G. Falk, L. Bry, J. Holgersson, and J. I. Gordon, *Proc. Natl. Acad. Sci. U.S.A.* 92, 1515 (1995).
25. P. D. Rye, *Nature Biotechnology* 2, 155 (1996).
26. P. Falk, T. Borén, D. Haslam, M. G. Caparon, *Meth. Cell Biol.* 45, 161 (1994)
27. Cell extracts were prepared in SDS sample buffer without mercapto ethanol and heated at 37° C. or 97° C. for 10 min. before separation on SDS-PAGE. Proteins were blotted onto a PVDF membrane. The membrane was incubated with 1 µg/ml biotinylated Le<sup>b</sup> glycoconjugate or biotinylated albumin (negative control) overnight,

- labelled as described in ref 7. After washing in PBS/0.05% Tween-20, the biotinylated structures bound by the BabA band were probed by HRP-streptavidin and detected using ECL reagents (Amersham, Buckinghamshire, England).
28. The bacterial suspension was incubated with Le<sup>b</sup> glycoconjugate, to which the Sulfo-SBED crosslinker (Pierce, Rockville, Ill.) had been conjugated by the N-hydroxysuccinimide ester (NHS), according to the manufacturers specifications. The aryl azide crosslinker group was activated by UV irradiation (360 nm). Bacteria were washed with PBS pH 7.6, 0.05% Tween-20 and protease inhibitors (EDTA and benzamidine) under reducing conditions with 50 mM dithiothreitol (DTT). Bacterial proteins were separated on SDS-PAGE, and the biotin tagged BabA protein was detected by immunodetection (PVDF membrane/HRP-streptavidin and ECL) (FIG. 3C).
29. Strains CCUG 17875 and A5 were first processed by crosslinking and DTT treatment, as above (28), followed by solubilisation in SDS sample buffer. The biotin tagged BabA protein was then extracted with streptavidin coated magnetic beads (Advanced Magnetics Inc., Cambridge, Mass.). The beads were boiled in SDS sample buffer, and bound proteins were eluted and alkylated. The protein preparation was further fractionated by preparative SDS-PAGE (Prep-Cell 491, BioRad, Hercules, Calif.). Fractions with the biotin tagged protein, i.e. the BabA fractions, were identified by immunodetection using streptavidin/ECL. The pooled BabA preparation was then separated on SDS-PAGE and transferred to PVDF membrane. The BabA band was excised and the BabA protein was N-terminally sequenced using a Procise™ 494 instrument (Applied Biosystems, Foster City, Calif.).

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1

<211> LENGTH: 3340

<212> TYPE: DNA

<213> ORGANISM: Helicobacter pylori

<400> SEQUENCE: 1

```

tttcagtcaa gcccaaaagt atgcgcacaaa cgcttatgct aaagagaatt tacaagcaca    60
gccgtccaag tatcaaaaca gcgctgcctga aatcaatatt gatgaagaag aaatcccctt    120
taagggttaa aattaaggag acattatgga aagaaaacgc tattcaaac gctattgcaa    180
atacactgaa gctaaaaatca gctttattga ctataaagat ttggacatgc tcaagcacac    240
gctatcagag cgctataaaa tcatgccaaag gaggttgaca ggcaatagca aaaagtggca    300
agagaggggtg gaagttagcg atcaaaagag cccgccacat ggctttaatc ccctacattg    360
tggataggaa aaaagtcgtg gatagccctt ttaaacagca ctgaattttt gattagggct    420
aatagggggc atgcctttta atcttgttta atcttggtc tttttttgtt aaacatcggc    480
tataaaagcg ttaaaagcac ttttaaaatc caattaaaag cgttcaaaag taacgcaaaa    540
aatcaaaaaa atgacaaaaa ttttaagaaa atgacaaaaa aaaaaaaaac gctttatgct    600
ataatattcc aaatacattc taatgcaaat gcattctaag gcaaatgtat aatgaatgta    660
tgaaatccct aatattcaat ccaatttaat ccaaaaagga gaaaaaacac atcctttcat    720

```

-continued

---

taactttagc ctcgctttta gtttccactt tgagcgctga agacgacggc ttttacacaa	780
gcgtaggcta tcaaatcggg gaagccgctc aaatggtaac aaacaccaa ggcacccaag	840
atctttcaga caactatgaa aacttgagca aacttttgac cggatacagc accctaaaca	900
cccttatcaa attgtccgct gatccgagcg cgattaacgc ggacagtgaa aatctgggcg	960
cgagcgcgaa gaacttgatc ggcgatacca aaaattcccc cgcctatcaa gccgtgcttt	1020
tggcgatcaa tgcggcggta gggttttgga atgtcttagg ctatgctacg caatgcgggg	1080
gtaacgctaa tggtaagaa agcactctt caaccacat cttcaacaac gagccagggt	1140
atcgatccac ttccatcact tgcagtttga acaggtataa gcctggatac tacggcccta	1200
tgagcattga aaatttcaaa aagcctaacg aagcctatca aatcctcaa acggctttaa	1260
ataaaggcct acccgcgctc aaagaaaaca acggaacggt cagtgtaac tacacctaca	1320
catgctcagg ggaaggaat gataactgct cgaaaaaagc cacaggtgta agtgaccaa	1380
atggcggaac caaaactaaa acccaaacca tagacggcaa aaccgtaacc accacgatca	1440
gttcaaaagt cgttgatagt caggcaaaag gtaatacaac aagggtgtcc tacaccgaaa	1500
tcactaacia attagacggt gtgcctgata gcgctcaagc gctcttgcg caagcgagca	1560
cgctcatcaa caccatcaac acggcatgcc cgtattttag tgtaactaat aaaagtggtg	1620
gtccacagat ggaaccgact agagggaaat tgtgcggttt tacagaagaa atcagcgcga	1680
tccaaaagat gatcacagac gcgcaagagc tggtaaatca aacgagcgtc attaacgagc	1740
atgaacaatc aaccccggtg ggcggtaata atggcaagcc tttcaaccct tcacggagc	1800
caagcttcgc tcaaggcatg ctcgctaacg ctagtgcgca agcctaaatg ctcaatctag	1860
cccatcaagt ggggcaaac attaacctg acaatcttac cgggactttt aaaaattttg	1920
ttacaggctt tttagccaca tgcaacaaca aatcaacagc tggcactagt ggcacacaag	1980
gttcacctcc tggcacagta accactcaaa ctttcgcttc cggttgcgcc tatgtggagc	2040
aaaccataac gaatctaac aacagcatcg ctcattttgg cactcaagag cagcagatac	2100
agcaagctga aaacatcgct gacactctag tgaatttcaa atctagatac agcgaattag	2160
ggaatactta taacagcatc actactgcgc tctccaaagt ccctaacgcg caaagcttgc	2220
aaaacgtggt gggaaaaaag aataaccct atagcccga aggcatagaa accaattact	2280
acttgaatca aaactcttac aaccaaacc aaaccatcaa ccaagaatta ggcgtaacc	2340
cotttaggaa agtgggcato gtcagttctc aaaccaaca tggtgccatg aatgggatcg	2400
gtatccaggt gggctacaag caattctttg ggcaaaaaag gaaatggggt gcaagatac	2460
acggcttttt tgattacaac catgcttca ttaaatccag cttcttcaac tcggcttctg	2520
acgtgtggac ttatggtttt ggagcggacg ctctttataa cttcatcaac gataaagcca	2580
ccaatttctt aggcaaaaac aacaagcttt ctgtggggct ttttgccggg attgcgttag	2640
cgggcacttc atggcttaat tctgaatagc tgaatttagc caccatgaat aacgtctata	2700
acgctaaaat gaacgtggcg aactccaat tcttattcaa catgggagtg aggatgaatt	2760
tagccagatc caagaaaaa ggcagcagatc atgcggctca gcatggcatt gagttagggc	2820
ttaaaatccc caccattaac acgaactact attcctttat gggggctgaa ctcaaatacc	2880
gcaggctcta tagcgtgat ttgaattatg tgttcgctta ctaaaaacta aaaatccttt	2940
gtggaactcc ctttttaagg ggtttctttt aaagccttta ttttttttg gaggggtta	3000
atttttttga aacctttgtt tttgaattct ctttttaatg ggtttctttt ttgaactctt	3060

-continued

---

tgttttgaac tccttttttt gaactccott ttttaaaccc tttctttttt aaaattctct	3120
tttttggggg gtttgatgaa aaatcctttt ttagcgtttt ggtattgggt agtggaaaac	3180
ttgatactaa ttttaagcga agttttttaa aagtgcctct ttaatatagg gggtttaagt	3240
tggtgattaa aaggggggaa tggtttcaaa gcgcttccta tccctttaag aaaataaaat	3300
aaaactttaa taaaatgagt tttacaacaa aatgagatcc	3340

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 2781

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Helicobacter pylori

&lt;400&gt; SEQUENCE: 2

catttgatcg cattggattht caaagaaggc cgttttgtga aaggctttgg tcaagcttat	60
gatatttttag gcgacaaaaat cgcttatggt gggggtaaag gcaaccacac caatttcgct	120
cacaagaaat aaactttctc acccataagg ggcaaacgcc cccaaaagag tgctttttta	180
agaggttaag gcaaaatcaa gctctttagt atttaactct aaaaaactact aaaagccttt	240
ttatgggcta acaccacaca aaaagcgtca aaatcaaaaa aatgacaaaa ttttcccaa	300
atgacaaaaa aaaaaaaaaa cgattttatg ctatattaac gaaatcttgt gataagatct	360
tattctttta aaagatttac ctaaccattht taatttcaag gagaaaacat gaaaaaaaaac	420
ccttttactc tctctctctc tctctctgtht ttgctccacg ctgaagacga cggcttttac	480
acaagcgtag gctatcaaat cgggtgaagcc gctcaaatgg taaccaacac caaaggcctc	540
caacagcttht cagacaatta tgaaaagctg aacaactcttht tgaataatta cagcacccta	600
aacaccctta tcaaattatc cgctgatccg agtgcgatta acgacgcaag ggataatcta	660
ggctcaagtg ctaagaattht gcttgatgtht aaaaccaact ccccgcccta tcaagccgtg	720
cttttagcgt tgaatgcggc ggtggggtht tggcaagtha caagctacgc ttttactgct	780
tgtggtcctg gcagtaacga gagcgcaaat ggaggtatcc aaacttttaa taatgtgcca	840
ggacaaaaga cgacaacat cacttgcaat tcgtattatc aaccaggaca tggggggcct	900
atatccactg caaactatgc aaaaatcaat caagcctatc aaatcattca aaaggcttht	960
acagccaatg aagctaagtg agatggggtht cccgthttta gcgacaccac tacaaaactt	1020
gatttcaacta ttcaaggaga caaagaacg ggtggccgac caaatacacc taaaaagtht	1080
ccatggagtg atgggaaata tattcacacc caatggattht acacaacacc acaatcaaca	1140
gaaacaaaaga tcaacacaga aaataacgct caagagcttht taaaacaagc gagcatcatt	1200
atcactacc taaatgagtg atgcccacac ttccaaaatg gtggtagcgg ttattggcaa	1260
gggataagcg gcaatgggac aatgtgtggg atgtttaaga atgaaatcag cgctatccaa	1320
ggcatgatcg ctaacgcgca agaagctgtht gcgcaagtha aaatcgtht tgaaaatgcg	1380
caaaatcaaa acaacttgga tactggaaaa ccattcaacc cttcacgga cgctagcttc	1440
gctcaaaagc tgctcaaaaa cgctcaagcc caagcagaga ttttaaacca agccgaacaa	1500
gtggtgaaaa actttgaaaa aatccctaaa aatthttgat cagactcttht aggggtgtht	1560
tatgaagagc aaggggtht gcgtagggg accaatccag gtcaggttac ttctaact	1620
ttcgcttccg gttgcgcta tgtggagcaa accataacga atctaaacaa cagcatcgct	1680
cattttggca ctcaagagca gcagatacag caagctgaaa acatcgctga cactctagtht	1740
aatttcaaat ctagatacag cgaattaggg aatacttata acagcatcac tactgcgctc	1800
tccaaagtht ctaacgcgca aagcttgcaa aacgtggtht gaaaaaagaa taaccctat	1860

-continued

```

agcccgcaag gcatagaaac caattactac ttgaatcaaa actcttacia ccaaatccaa 1920
accatcaacc aagaattagg gcgtaacccc tttaggaaag tgggcatcgt cagttctcaa 1980
accaacaatg gtgccatgaa tgggatcggg atccaggtgg gctacaagca attccttggg 2040
caaaaaagga aatgggggtgc aagatactac ggcttttttg attacaacca tgcgttcatt 2100
aaatccagct tcttcaactc ggcttctgac gtgtggactt atggtttttg agcggagcgt 2160
ctttataact tcataacga taaagccacc aatttcttag gcaaaaacia caagctttct 2220
gtggggcttt ttggcgggat tgcgtagcg ggcacttcat ggctaattc tgaatacgtg 2280
aatttagcca ccatgaataa cgtctataac gctaaaatga acgtggcgaa cttccaattc 2340
ttattcaaca tgggagtgg gatgaattta gccagatcca agaaaaagg cagcgatcat 2400
gcggtcagc atggcattga gttagggtt aaaatcccca ccattaacac gaactactat 2460
tcctttatgg gggctgaact caaataccgc aggctctata gogtgtattt gaattatgtg 2520
ttcgttact agaaactaaa aatcctttgt ggaactccct ttttaagggg tttcttttaa 2580
agcctttatt tttttttgga ggggtttaat tttttgaaa cctttgtttt tgaattctct 2640
ttttaatggg tttctttttt gaactctttg ttttgaactc ctttttttga actccctttt 2700
ttaaacctt tcttttttaa aattctcttt tttggggggg ttgatgaaaa atcctttttt 2760
agcgttttgg tattggtag t 2781

```

```

<210> SEQ ID NO 3
<211> LENGTH: 744
<212> TYPE: PRT
<213> ORGANISM: Helicobacter pylori
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (24)..(64)
<223> OTHER INFORMATION: N-terminal domain of the BabA adhesin
<400> SEQUENCE: 3

```

```

Ser Lys Lys Glu Lys Lys His Ile Leu Ser Leu Thr Leu Gly Ser Leu
 1           5           10          15
Leu Val Ser Thr Leu Ser Ala Glu Asp Asp Gly Phe Tyr Thr Ser Val
 20          25
Gly Tyr Gln Ile Gly Glu Ala Ala Gln Met Val Thr Asn Thr Lys Gly
 35          40          45
Ile Gln Asp Leu Ser Asp Asn Tyr Glu Asn Leu Ser Lys Leu Leu Thr
 50          55          60
Arg Tyr Ser Thr Leu Asn Thr Leu Ile Lys Leu Ser Ala Asp Pro Ser
 65          70          75          80
Ala Ile Asn Ala Ala Arg Glu Asn Leu Gly Ala Ser Ala Lys Asn Leu
 85          90          95
Ile Gly Asp Thr Lys Asn Ser Pro Ala Tyr Gln Ala Val Leu Leu Ala
100         105         110
Ile Asn Ala Ala Val Gly Phe Trp Asn Val Leu Gly Tyr Ala Thr Gln
115         120         125
Cys Gly Gly Asn Ala Asn Gly Gln Glu Ser Thr Ser Ser Thr Thr Ile
130         135         140
Phe Asn Asn Glu Pro Gly Tyr Arg Ser Thr Ser Ile Thr Cys Ser Leu
145         150         155         160
Asn Arg Tyr Lys Pro Gly Tyr Tyr Gly Pro Met Ser Ile Glu Asn Phe
165         170         175
Lys Lys Leu Asn Glu Ala Tyr Gln Ile Leu Gln Thr Ala Leu Asn Lys

```

-continued

180					185					190					
Gly	Leu	Pro	Ala	Leu	Lys	Glu	Asn	Asn	Gly	Thr	Val	Ser	Val	Thr	Tyr
	195						200					205			
Thr	Tyr	Thr	Cys	Ser	Gly	Glu	Gly	Asn	Asp	Asn	Cys	Ser	Lys	Lys	Ala
	210					215					220				
Thr	Gly	Val	Ser	Asp	Gln	Asn	Gly	Gly	Thr	Lys	Thr	Lys	Thr	Gln	Thr
	225					230					235				240
Ile	Asp	Gly	Lys	Thr	Val	Thr	Thr	Thr	Ile	Ser	Ser	Lys	Val	Val	Asp
				245					250					255	
Ser	Gln	Ala	Lys	Gly	Asn	Thr	Thr	Arg	Val	Ser	Tyr	Thr	Glu	Ile	Thr
			260					265					270		
Asn	Lys	Leu	Asp	Gly	Val	Pro	Asp	Ser	Ala	Gln	Ala	Leu	Leu	Ala	Gln
			275				280					285			
Ala	Ser	Thr	Leu	Ile	Asn	Thr	Ile	Asn	Thr	Ala	Cys	Pro	Tyr	Phe	Ser
	290					295					300				
Val	Thr	Asn	Lys	Ser	Gly	Gly	Pro	Gln	Met	Glu	Pro	Thr	Arg	Gly	Lys
	305					310					315				320
Leu	Cys	Gly	Phe	Thr	Glu	Glu	Ile	Ser	Ala	Ile	Gln	Lys	Met	Ile	Thr
				325					330					335	
Asp	Ala	Gln	Glu	Leu	Val	Asn	Gln	Thr	Ser	Val	Ile	Asn	Glu	His	Glu
			340					345					350		
Gln	Ser	Thr	Pro	Val	Gly	Gly	Asn	Asn	Gly	Lys	Pro	Phe	Asn	Pro	Phe
		355					360					365			
Thr	Asp	Ala	Ser	Phe	Ala	Gln	Gly	Met	Leu	Ala	Asn	Ala	Ser	Ala	Gln
	370					375					380				
Ala	Lys	Met	Leu	Asn	Leu	Ala	His	Gln	Val	Gly	Gln	Thr	Ile	Asn	Pro
	385					390					395				400
Asp	Asn	Leu	Thr	Gly	Thr	Phe	Lys	Asn	Phe	Val	Thr	Gly	Phe	Leu	Ala
			405					410						415	
Thr	Cys	Asn	Asn	Lys	Ser	Thr	Ala	Gly	Thr	Ser	Gly	Thr	Gln	Gly	Ser
		420						425					430		
Pro	Pro	Gly	Thr	Val	Thr	Thr	Gln	Thr	Phe	Ala	Ser	Gly	Cys	Ala	Tyr
		435					440					445			
Val	Glu	Gln	Thr	Ile	Thr	Asn	Leu	Asn	Asn	Ser	Ile	Ala	His	Phe	Gly
	450					455					460				
Thr	Gln	Glu	Gln	Gln	Ile	Gln	Gln	Ala	Glu	Asn	Ile	Ala	Asp	Thr	Leu
	465					470					475				480
Val	Asn	Phe	Lys	Ser	Arg	Tyr	Ser	Glu	Leu	Gly	Asn	Thr	Tyr	Asn	Ser
			485					490						495	
Ile	Thr	Thr	Ala	Leu	Ser	Lys	Val	Pro	Asn	Ala	Gln	Ser	Leu	Gln	Asn
			500					505					510		
Val	Val	Gly	Lys	Lys	Asn	Asn	Pro	Tyr	Ser	Pro	Gln	Gly	Ile	Glu	Thr
		515					520					525			
Asn	Tyr	Tyr	Leu	Asn	Gln	Asn	Ser	Tyr	Asn	Gln	Ile	Gln	Thr	Ile	Asn
	530					535					540				
Gln	Glu	Leu	Gly	Arg	Asn	Pro	Phe	Arg	Lys	Val	Gly	Ile	Val	Ser	Ser
				545		550					555				560
Gln	Thr	Asn	Asn	Gly	Ala	Met	Asn	Gly	Ile	Gly	Ile	Gln	Val	Gly	Tyr
				565				570						575	
Lys	Gln	Phe	Phe	Gly	Gln	Lys	Arg	Lys	Trp	Gly	Ala	Arg	Tyr	Tyr	Gly
			580					585					590		
Phe	Phe	Asp	Tyr	Asn	His	Ala	Phe	Ile	Lys	Ser	Ser	Phe	Phe	Asn	Ser
		595					600					605			

-continued

Ala Ser Asp Val Trp Thr Tyr Gly Phe Gly Ala Asp Ala Leu Tyr Asn  
610 615 620

Phe Ile Asn Asp Lys Ala Thr Asn Phe Leu Gly Lys Asn Asn Lys Leu  
625 630 635 640

Ser Val Gly Leu Phe Gly Gly Ile Ala Leu Ala Gly Thr Ser Trp Leu  
645 650 655

Asn Ser Glu Tyr Val Asn Leu Ala Thr Met Asn Asn Val Tyr Asn Ala  
660 665 670

Lys Met Asn Val Ala Asn Phe Gln Phe Leu Phe Asn Met Gly Val Arg  
675 680 685

Met Asn Leu Ala Arg Ser Lys Lys Lys Gly Ser Asp His Ala Ala Gln  
690 695 700

His Gly Ile Glu Leu Gly Leu Lys Ile Pro Thr Ile Asn Thr Asn Tyr  
705 710 715 720

Tyr Ser Phe Met Gly Ala Glu Leu Lys Tyr Arg Arg Leu Tyr Ser Val  
725 730 735

Tyr Leu Asn Tyr Val Phe Ala Tyr  
740

<210> SEQ ID NO 4  
<211> LENGTH: 707  
<212> TYPE: PRT  
<213> ORGANISM: Helicobacter pylori  
<220> FEATURE:  
<221> NAME/KEY: PEPTIDE  
<222> LOCATION: (46)..(59)  
<223> OTHER INFORMATION: Corresponding to the N-terminal domain of the  
BabA  
adhesin

<400> SEQUENCE: 4

Met Lys Lys Asn Pro Phe Thr Leu Ser Leu Ser Leu Ser Phe Leu Leu  
1 5 10 15

His Ala Glu Asp Asp Gly Phe Tyr Thr Ser Val Gly Tyr Gln Ile Gly  
20 25 30

Glu Ala Ala Gln Met Val Thr Asn Thr Lys Gly Ile Gln Gln Leu Ser  
35 40 45

Asp Asn Tyr Glu Lys Leu Asn Asn Leu Leu Asn Asn Tyr Ser Thr Leu  
50 55 60

Asn Thr Leu Ile Lys Leu Ser Ala Asp Pro Ser Ala Ile Asn Asp Ala  
65 70 75 80

Arg Asp Asn Leu Gly Ser Ser Ala Lys Asn Leu Leu Asp Val Lys Thr  
85 90 95

Asn Ser Pro Ala Tyr Gln Ala Val Leu Leu Ala Leu Asn Ala Ala Val  
100 105 110

Gly Leu Trp Gln Val Thr Ser Tyr Ala Phe Thr Ala Cys Gly Pro Gly  
115 120 125

Ser Asn Glu Ser Ala Asn Gly Gly Ile Gln Thr Phe Asn Asn Val Pro  
130 135 140

Gly Gln Lys Thr Thr Thr Ile Thr Cys Asn Ser Tyr Tyr Gln Pro Gly  
145 150 155 160

His Gly Gly Pro Ile Ser Thr Ala Asn Tyr Ala Lys Ile Asn Gln Ala  
165 170 175

Tyr Gln Ile Ile Gln Lys Ala Leu Thr Ala Asn Glu Ala Asn Gly Asp  
180 185 190

Gly Val Pro Val Leu Ser Asp Thr Thr Thr Lys Leu Asp Phe Thr Ile

-continued

195			200			205									
Gln	Gly	Asp	Lys	Arg	Thr	Gly	Gly	Arg	Pro	Asn	Thr	Pro	Lys	Lys	Phe
210						215					220				
Pro	Trp	Ser	Asp	Gly	Lys	Tyr	Ile	His	Thr	Gln	Trp	Ile	Asp	Thr	Thr
225					230					235				240	
Pro	Gln	Ser	Thr	Glu	Thr	Lys	Ile	Asn	Thr	Glu	Asn	Asn	Ala	Gln	Glu
				245					250					255	
Leu	Leu	Lys	Gln	Ala	Ser	Ile	Ile	Ile	Thr	Thr	Leu	Asn	Glu	Ala	Cys
			260					265					270		
Pro	Asn	Phe	Gln	Asn	Gly	Gly	Ser	Gly	Tyr	Trp	Gln	Gly	Ile	Ser	Gly
		275					280					285			
Asn	Gly	Thr	Met	Cys	Gly	Met	Phe	Lys	Asn	Glu	Ile	Ser	Ala	Ile	Gln
290					295						300				
Gly	Met	Ile	Ala	Asn	Ala	Gln	Glu	Ala	Val	Ala	Gln	Ser	Lys	Ile	Val
305				310						315					320
Ser	Glu	Asn	Ala	Gln	Asn	Gln	Asn	Asn	Leu	Asp	Thr	Gly	Lys	Pro	Phe
			325					330						335	
Asn	Pro	Phe	Thr	Asp	Ala	Ser	Phe	Ala	Gln	Ser	Met	Leu	Lys	Asn	Ala
			340					345					350		
Gln	Ala	Gln	Ala	Glu	Ile	Leu	Asn	Gln	Ala	Glu	Gln	Val	Val	Lys	Asn
		355					360						365		
Phe	Glu	Lys	Ile	Pro	Lys	Asn	Phe	Val	Ser	Asp	Ser	Leu	Gly	Val	Cys
	370					375					380				
Tyr	Glu	Glu	Gln	Gly	Gly	Glu	Arg	Arg	Gly	Thr	Asn	Pro	Gly	Gln	Val
385					390					395					400
Thr	Ser	Asn	Thr	Phe	Ala	Ser	Gly	Cys	Ala	Tyr	Val	Glu	Gln	Thr	Ile
			405						410					415	
Thr	Asn	Leu	Asn	Asn	Ser	Ile	Ala	His	Phe	Gly	Thr	Gln	Glu	Gln	Gln
		420						425					430		
Ile	Gln	Gln	Ala	Glu	Asn	Ile	Ala	Asp	Thr	Leu	Val	Asn	Phe	Lys	Ser
		435					440					445			
Arg	Tyr	Ser	Glu	Leu	Gly	Asn	Thr	Tyr	Asn	Ser	Ile	Thr	Thr	Ala	Leu
	450				455						460				
Ser	Lys	Val	Pro	Asn	Ala	Gln	Ser	Leu	Gln	Asn	Val	Val	Gly	Lys	Lys
465				470						475					480
Asn	Asn	Pro	Tyr	Ser	Pro	Gln	Gly	Ile	Glu	Thr	Asn	Tyr	Tyr	Leu	Asn
			485						490					495	
Gln	Asn	Ser	Tyr	Asn	Gln	Ile	Gln	Thr	Ile	Asn	Gln	Glu	Leu	Gly	Arg
		500						505					510		
Asn	Pro	Phe	Arg	Lys	Val	Gly	Ile	Val	Ser	Ser	Gln	Thr	Asn	Asn	Gly
		515					520						525		
Ala	Met	Asn	Gly	Ile	Gly	Ile	Gln	Val	Gly	Tyr	Lys	Gln	Phe	Phe	Gly
	530				535						540				
Gln	Lys	Arg	Lys	Trp	Gly	Ala	Arg	Tyr	Tyr	Gly	Phe	Phe	Asp	Tyr	Asn
545					550					555					560
His	Ala	Phe	Ile	Lys	Ser	Ser	Phe	Phe	Asn	Ser	Ala	Ser	Asp	Val	Trp
			565					570						575	
Thr	Tyr	Gly	Phe	Gly	Ala	Asp	Ala	Leu	Tyr	Asn	Phe	Ile	Asn	Asp	Lys
			580					585					590		
Ala	Thr	Asn	Phe	Leu	Gly	Lys	Asn	Asn	Lys	Leu	Ser	Val	Gly	Leu	Phe
		595					600						605		
Gly	Gly	Ile	Ala	Leu	Ala	Gly	Thr	Ser	Trp	Leu	Asn	Ser	Glu	Tyr	Val
	610						615						620		

-continued

Asn Leu Ala Thr Met Asn Asn Val Tyr Asn Ala Lys Met Asn Val Ala  
 625 630 635 640

Asn Phe Gln Phe Leu Phe Asn Met Gly Val Arg Met Asn Leu Ala Arg  
 645 650 655

Ser Lys Lys Lys Gly Ser Asp His Ala Ala Gln His Gly Ile Glu Leu  
 660 665 670

Gly Leu Lys Ile Pro Thr Ile Asn Thr Asn Tyr Tyr Ser Phe Met Gly  
 675 680 685

Ala Glu Leu Lys Tyr Arg Arg Leu Tyr Ser Val Tyr Leu Asn Tyr Val  
 690 695 700

Phe Ala Tyr  
 705

<210> SEQ ID NO 5  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Helicobacter pylori

<400> SEQUENCE: 5

Glu Asp Asp Gly Phe Tyr Thr Ser Val Gly Tyr Gln Ile Gly Glu Ala  
 1 5 10 15

Ala Gln Met Val  
 20

<210> SEQ ID NO 6  
 <211> LENGTH: 60  
 <212> TYPE: DNA  
 <213> ORGANISM: Helicobacter pylori

<400> SEQUENCE: 6

gaagacgacg gcttttacac aagcgtaggc tatcaaatcg gtgaagccgc tcaaatggta 60

<210> SEQ ID NO 7  
 <211> LENGTH: 41  
 <212> TYPE: PRT  
 <213> ORGANISM: Helicobacter pylori

<400> SEQUENCE: 7

Glu Asp Asp Gly Phe Tyr Thr Ser Val Gly Tyr Gln Ile Gly Glu Ala  
 1 5 10 15

Ala Gln Met Val Thr Asn Thr Lys Gly Ile Gln Asp Leu Ser Asp Asn  
 20 25 30

Tyr Glu Asn Leu Ser Lys Leu Leu Thr  
 35 40

<210> SEQ ID NO 8  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Helicobacter pylori

<400> SEQUENCE: 8

Glu Asp Asp Gly Phe Tyr Thr Ser Val Gly Tyr Gln Ile Gly Glu Ala  
 1 5 10 15

Ala Gln Met Val  
 20

<210> SEQ ID NO 9  
 <211> LENGTH: 41  
 <212> TYPE: PRT  
 <213> ORGANISM: Helicobacter pylori

-continued

&lt;400&gt; SEQUENCE: 9

Glu Asp Asp Gly Phe Tyr Thr Ser Val Gly Tyr Gln Ile Gly Glu Ala  
 1                    5                    10                    15  
 Ala Gln Met Val Thr Asn Thr Lys Gly Ile Gln Gln Leu Ser Asp Asn  
                   20                    25                    30  
 Tyr Glu Lys Leu Asn Asn Leu Leu Asn  
                   35                    40

The invention claimed is:

1. A test kit comprising a monospecific antiserum that recognizes a BabA antigen and comprises an immunoglobulin that binds said BabA antigen via a variable region, wherein said Bab A adhesin protein comprises SEQ ID NO:5 and is produced using an isolated and purified bacterial blood group antigen binding protein (BabA) from *Helicobacter pylori* species, wherein said BabA protein binds specifically to fucosylated Lewis<sup>b</sup> type I and H-1 blood group antigen-glycoconjugates and,

wherein said BabA protein contains less than 20% bacterial protein impurities, has a molecular weight in the interval of 70 to 77 kDa as determined by sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE), and is not a HopA, HopB, HopC, HopD, or HopE protein.

2. The test kit according to claim 1, wherein said BabA adhesin protein has a molecular weight in the interval of about 73 to 75 kDa as determined by SDS-PAGE.

3. The test kit according to claim 1, wherein the molecular weight of said BabA adhesin protein is about 73.5 kDa.

4. A test kit comprising an isolated monospecific immunoglobulin composition that recognizes a BabA antigen and

15 comprises an immunoglobulin that binds said BabA antigen via a variable region, wherein said Bab A adhesin protein comprises SEQ ID NO:5 and is produced using an isolated and purified bacterial blood group antigen binding protein (BabA) from *Helicobacter pylori* species, wherein said BabA protein binds specifically to fucosylated Lewis<sup>b</sup> type I and H-1 blood group antigen-glycoconjugates and,

wherein said BabA protein contains less than 20% bacterial protein impurities, has a molecular weight in the interval of 70 to 77 kDa as determined by sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE), and is not a HopA, HopB, HopC, HopD, or HopE protein.

5. The immunoglobulin according to claim 4, wherein the molecular weight of said BabA adhesin protein is about 73.5 kDa.

6. The test kit according to claim 4, wherein said BabA adhesin protein has a molecular weight in the interval of about 73 to 75 kDa as determined by SDS-PAGE.

7. The test kit according to claim 4, wherein said immunoglobulin is a monoclonal antibody.

\* \* \* \* \*

专利名称(译)	幽门螺杆菌粘附素结合基因抗原		
公开(公告)号	<a href="#">US7271251</a>	公开(公告)日	2007-09-18
申请号	US10/761201	申请日	2004-01-22
[标]申请(专利权)人(译)	BOREN THOMAS ARNQVIST ANNA HAMMARSTROM伦纳特 NORMARK斯塔 ILVER DAG		
申请(专利权)人(译)	BOREN THOMAS ARNQVIST ANNA HAMMARSTROM伦纳特 NORMARK斯塔 ILVER DAG		
当前申请(专利权)人(译)	BOREN THOMAS ARNQVIST ANNA HAMMARSTROM伦纳特 NORMARK斯塔 ILVER DAG		
[标]发明人	BOREN THOMAS ARNQVIST ANNA HAMMARSTROM LENNART NORMARK STAFFAN ILVER DAG		
发明人	BOREN, THOMAS ARNQVIST, ANNA HAMMARSTROM, LENNART NORMARK, STAFFAN ILVER, DAG		
IPC分类号	A61K39/40 C07K16/12 G01N33/53 A61K39/00 A61K39/02 A61K39/116 A61K39/395 A61K48/00 A61P1/04 A61P31/04 C07K1/14 C07K14/195 C07K14/205 C12N15/09 C12P21/08 G01N33/569		
CPC分类号	C07K14/205 A61K39/00 A61K2039/51 A61P1/04		
优先权	09/202178 1999-02-10 US		
其他公开文献	US20040234529A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

#### 摘要(译)

分离并纯化新型幽门螺杆菌血型抗原结合 (BAB) 粘附素蛋白, 其中所述蛋白质或其部分特异性结合岩藻糖基化血型抗原。在本申请中公开了所述粘附素的蛋白质序列。同时公开了产生高度相似蛋白质的两种基因babA和babB的DNA序列。所述粘附素和/或DNA可用于诊断和治疗和/或预防针对幽门螺杆菌诱导的感染, 例如, 胃炎和酸性消化性疾病, 即主动接种疫苗。一种新的免疫球蛋白组合物, 其对结合幽门螺杆菌粘附素的Lewisb抗原表现出特异性活性, 或优选地, 对所述粘附素的单克隆和/或多克隆抗体提供了一种新的和更有效的治疗和/或预防由幽门螺杆菌引起的胃肠疾病的方法。幽门螺杆菌或其他螺杆菌属物种, 即被动疫苗接种。

Antibody preparations used for Western blot detection of the BabA adhesin and *H. pylori* protein antigens:

