



US 20060024749A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2006/0024749 A1**
Dambinova (43) **Pub. Date: Feb. 2, 2006**

(54) **RAPID MULTIPLE PANEL OF BIOMARKERS
IN LABORATORY BLOOD TESTS FOR
TIA/STROKE**

(60) Provisional application No. 60/301,297, filed on Jun. 27, 2001.

(76) Inventor: **Svetlana A. Dambinova**, Atlanta, GA (US)

Correspondence Address:
KING & SPALDING LLP
191 PEACHTREE STREET, N.E.
45TH FLOOR
ATLANTA, GA 30303-1763 (US)

Publication Classification

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G01N 33/53 (2006.01)
(52) **U.S. Cl.** **435/7.1; 435/6**

(21) Appl. No.: **11/076,074**

(22) Filed: **Mar. 9, 2005**

Related U.S. Application Data

(63) Continuation of application No. 09/922,011, filed on Aug. 2, 2001, now Pat. No. 6,896,872.

(57) **ABSTRACT**

A methods, kits and compositions for diagnosing a central nervous system disorder, particularly transient ischemic attack or stroke, comprising measuring the level of NR2A and/or NR2B NMDA receptor or fragment thereof, in a biological sample from a human subject, and optionally measuring other biomarkers such as homocysteine and glutamate. The method is particularly useful for identifying individuals that are at risk for stroke, and for diagnosing stroke in an emergency room setting.

RAPID MULTIPLE PANEL OF BIOMARKERS IN LABORATORY BLOOD TESTS FOR TIA/STROKE

RELATION TO PRIOR APPLICATIONS

[0001] This application is a continuation of U.S. Utility Application No. 09/922,011, filed Aug. 2, 2001 (allowed), and claims priority under 35 U.S.C. § 119(e) to U.S. Provisional Application No. 60/301,297, filed Jun. 27, 2001, and under 35 U.S.C. § 120 to U.S. Utility Application No. 09/632,749, filed Aug. 4, 2000 (currently pending), of which this application is a continuation-in-part.

FIELD OF THE INVENTION

[0002] The present invention relates generally to the diagnosis, management and therapy of central nervous system disorders such as stroke, transient ischemic attack, and traumatic brain injury. In particular, the invention relates to methods and kits for evaluating these central nervous system disorders, in order to better respond to episodes of focal cerebral ischemia, and to best manage the risk associated with future acute incidences.

BACKGROUND INFORMATION

[0003] Stroke or "brain attack" is clinically defined as a rapidly developing syndrome of vascular origin that manifests itself in focal loss of cerebral function. In more severe situations, the loss of cerebral function is global. A stroke occurs when the blood supply to the part of the brain is suddenly interrupted (ischemic) or when a blood vessel in the brain bursts, spilling blood into the spaces surrounding the brain cells (hemorrhagic). The symptoms of stroke are easy to spot: sudden numbness or weakness, especially on one side of the body; sudden confusion or trouble speaking or understanding speech; sudden trouble seeing in one or both eyes; sudden trouble walking; dizziness; or loss of balance or coordination. (National Institute of Neurological Disorders and Stroke, 2001). Stroke is the most common devastating neurologic disease in the world, and the third leading cause of death in world after heart disease and cancer. Despite recent progress understanding stroke mechanisms, stroke management is still not optimal for a number of reasons.

[0004] The importance of promptly diagnosing a stroke after symptoms appear cannot be overstated. Delays in diagnosis and medical intervention beyond 3 hours after stroke onset may contribute to clinical deterioration and disability. An early diagnosis enables doctors to more effectively choose the emergency intervention such as antiplatelet or/and neuroprotective therapy, and also to make better prognoses of disease outcome. Successful treatment of stroke requires rapid state diagnosis. The delay in achieving an accurate and certain diagnosis wastes the limited amount of time available in which the brain can respond to reperfusion, and significantly increases the risk of hemorrhage after most of the permanent injury has occurred (Marler J. R. *Annl. Emergency Med.* 1999, 33: 450-451).

[0005] Unfortunately, however, many people who have a stroke either do not seek immediate medical care or suffer from delays in medical care even in countries where stroke care is advanced, such as the United States and Europe (Alberts M J, Hademenos G, Latchaw R E, et al. *JAMA* 2000; 23:3102-3109). Several clinical criteria can be

employed to diagnose whether a patient is having a stroke, but even all these criteria do not always allow one to differentiate the episode from other disorders, such as epilepsy, syncope, and migraine (Toole J F. *Cerebrovascular Disorders*. 1999. Lippincott, Williams & Wilkins, New York, 5th Ed., 542 p). Moreover, progressing stroke is only partially predictable based on clinical and neuroimaging data that is currently available to neurologists.

[0006] Transient ischemic attack (TIA) is a short-lived episode of focal neurologic deficit which often precedes the cerebral infarction of a stroke. It occurs when the blood supply to part of the brain is briefly interrupted, and is typically accompanied by permanent brain damage (albeit less severe damage than normally results from a stroke). TIA symptoms, which usually occur suddenly, are similar to those of stroke but do not last as long. Most symptoms of a TIA disappear within an hour, although they may persist for up to 24 hours. Symptoms can include: numbness or weakness in the face, arm, or leg, especially on one side of the body; confusion or difficulty in talking or understanding speech; trouble seeing in one or both eyes; and difficulty with walking, dizziness, or loss of balance and coordination. (National Institute of Neurological Disorders and Stroke, 2001). Patients who have suffered a TIA have 9.5 times greater risk of having a future stroke than those who have not had a TIA, and about one third of patients who suffer a TIA will have an acute stroke in the future. (American Stroke Association, 2001). However, because the symptoms of TIA are short term, many patients do not recognize the event as a TIA or perceive the event as a warning of a potentially impending stroke.

[0007] Standard treatments to reduce the risk of future stroke include the use of antiplatelet agents, particularly aspirin. People with atrial fibrillation (irregular beating of the heart) may be prescribed anticoagulants. The most important treatable factors linked to TIAs and stroke are high blood pressure, cigarette smoking, heart disease, carotid artery disease, diabetes, and heavy use of alcohol. Lifestyle changes can often be implemented to reduce these factors. However, it is necessary to diagnose the TIA as a warning sign of impending stroke before such treatments can be administered. Therefore, a laboratory blood test to detect TIA or stroke, or the risk of suffering a TIA or stroke in the future, would be of tremendous benefit.

[0008] During the past 5 years a number of molecular and immunochemical assays have been evaluated for clinical use in neurology. (Schenone A. et al. *Current Opinion in Neurology*. 1999, 12: 603-604; Honnorat J. *J. Neurosurg. Psychiatry*. 1996, 61:270-278). At present, the Thrombogene V and two Thrombx tests are available for diagnosing stroke/thrombosis from Athena Diagnostic. These tests evaluate the frequent deep vein thrombosis and hypercoagulation states of patients to evaluate the need for intravenous anticoagulant therapy. The Thrombogene V test detects the Factor V Leiden mutation by Polymerase Chain Reaction (PCR) in the blood of patients. The other two tests monitor changes of different blood coagulation markers: antithrombin Im protein C, factor IX, and anticardiolipin antibodies (IgG, IgM, IgA) by use of ELISA technique. These tests thus reveal the hypercoagulation state as a result of a thrombotic events, such as stroke stroke.

[0009] Stroke can be related to different types of venous thromboembolisms, which are common disorders with con-

siderable morbidity and potential for mortality (Anderson, D.; Wells, P. Cur. Opin. in Hemat. 2000, 7: 296-301). The biochemical marker: D-dimer, a breakdown product of a cross-linked fibrin blood clot that indicates the occurrence of plasmin mediated lysis of cross-linked fibrin, has been extensively evaluated for use in diagnostic tests for indicating acute venous thromboembolism. Indeed, a fully automated, semi-quantitative latex agglutination assays that uses turbidimetric or agglutination endpoints has been developed that provides results within 20 minutes with sensitivity between 89% and 95% (Roussi J.; Bentolila L.; Contribution of D-dimer determination in the exclusion of deep venous thrombosis in spinal cord injury patients. Spinal Cord, 1999; v. 37: p. 548-552). Unfortunately, however, the presence of D-dimer may also be increased in other settings that result in fibrin generation, including recent surgery, hemorrhage, trauma, cancer, and pregnancy (Anderson D R., Wells P S.; Thromb. Haemost.; 1999; 82:878-886).

[0010] However, the foregoing tests do not elucidate upon the TIA/stroke mechanisms that are actually responsible for the damage associated with neurotoxic molecular events. It is necessary to find out specific and sensitive biomarkers which could be helpful to recognize initial brain damage and which could help to choose not only the appropriate anticoagulant treatment, but also emergency or regular neuroprotective therapy.

[0011] It is well known that two of the three leading causes of death, namely cardiovascular diseases and stroke, are the end result of atherosclerosis. Thus, it is not surprising that several biochemical markers implicated in thromboembolic processes are also reported to be associated with stroke and stroke risk. Among these are homocysteine, cholesterol and LDL (Cerebrovascular Disorders ed. by J. E. Toole. Lippincott Williams & Wilkins. 1999, pp. 34-35), which are also classified as risk factors to cardiovascular and cerebrovascular diseases. (Hankey G J., and Eikelboom J W. Lancet. 354: 407-413 (1999). Approximately one fourth of patients with symptomatic atherosclerosis have elevated plasma homocysteine levels caused by various factors. High levels of homocysteine may run in families with increased susceptibility to heart attack and stroke (Graham I. J. Ir. Call. Phys. Surg. 1995; 24: 25-30). Elevated plasma homocysteine may be a causal and modifiable risk-factor for ischemic stroke, but the results of previous studies have been conflicting (Deulofeu V N R, Chamorro A, Piera C. Med Clin (Barc). 1998; 110: 605-608; Yamamoto T, Rossi S, Stiefel M F, Doppenberg E, Zauner A, Bullock R, Marmarou A. Acta Neurochir. Suppl. (Wien) 1999; 75:17-19).

[0012] The neurotoxic effect of excitatory amino acids (glutamate, aspartate) in the brain has also been well documented. The results of this work show a correlation between glutamate content in the blood and the severity of acute ischaemia (Castillo J, Dávalos A, Naveiro J, Noya M. Stroke 1996, 27:1060-1065; Castillo J, Dávalos A, Noya M. Lancet. 1997; 349:79-83). Cerebral damage and its association with progressing stroke has been attributed to increased glutamate release, or low glutamate reuptake, both in animals and in humans (Dávalos A, Castillo J. Serena J. Noya M. Stroke 1997; 28:708-710).

[0013] However, only 56% of patients with progressing stroke are reported to have high glutamate content in their blood serum (Dávalos A., Toni D., Iweins F., et al., 1999, 30:

2631-2636). Moreover, even though glutamate is considered the strongest biochemical predictor of progressing stroke (Davalos A, and Castillo J. In Book: *Cerebrovascular Disease*. Current Med. Inc.: Philadelphia. 2000 Chapter 16, pp. 169-181), this marker remains non-specific for TIA. Glutamate changes have also been observed in the blood of patients with epilepsy and other nervous system disorders (Meldrum B S. J. Nutrition. 2000, 130:1007S-1015S).

[0014] Over the last three decades substantial progress has been made in elucidating the mechanisms by which cerebral ischemia leads to brain damage. The cellular and molecular mechanisms of cerebral ischemia abnormalities have been better defined through the role of glutamate and glutamate receptors, one of the most distributed excitatory neuroreceptors in brain, in regulating of initial stages of brain damage. Indeed, numerous molecular investigators consider glutamate receptors to be one of the key biological receptors involved in the molecular mechanisms of TIA and stroke (Meldrum B S. J. Nutrition. 2000, 130:1007S-1015S). According to a leading hypothesis, ischemia-induced glutamate release activates these glutamate receptors. It has been shown that glutamate and homocysteine (the sulfonic analog of aspartate) activate the glutamate binding site of NMDA receptors and participate in neurotoxic processes (Lipton S. A., Kim W. K., Choi Y. B., Kumar S., et al. PNAS. 1997, 94: 5923-5928).

[0015] Glutamate receptors are divided into two main groups: ionotropic and metabotropic. The ionotropic neuroreceptors are ligand-gated ion channels that are subdivided into NMDA, AMPA and kainate receptor subtypes. There are four NR2 subunits: NR2A, NR2B, NR2C and NR2D, which is responsible for Ca²⁺-permeability regulation. NMDA receptors can be modified by ischemia, resulting in changes of ion permeability and/or ion selectivity.

[0016] Recent research findings indicate that the blood of patients with CNS disorders other than TIA or stroke exhibit properties of autoimmunization to products of nerve cell degradation (Vincent A., Oliver L., Pallace J. J Neuroimmun. 1999; 100: 169-180). For example, a correlation between AMPA GluR1 autoantibodies and common epilepsy has been shown (Dambinova et al. J. Neurol. Sci. 1997; 152: 93-97; Dambinova et al. J. Neurochem. 1998;71: 2088-2093), as has a correlation between AMPA GluR 3 autoantibodies and Rasmussen's encephalitis (Rogers S W, Andrews PI, Gahring L C, et al. Science. 1995;265:648-651; Twyman R E, Gahring L C, Spiess J, Rogers S W. Neuron. 1995; 14:755-762; Gahring L C, Twyman R I, Greenlee J E, Rogers S W. Mol. Med. 1995; 1:245-253).

[0017] In a similar vein, several researchers have reported an increase in NMDA receptor synthesis, the appearance of high levels of receptor antigen, and the generation of autoantibodies to the receptors during the initial stages of cerebral ischemia (Gusev et al., J. Neurol. & Psych. 1996, 5:68-72; Dambinova et al. J. Neurol. Sci. 1997, 152:93-97; Dambinova et al. J. Neurochem. 1998, 71: 2088-2093). Acting on this research, one company developed a laboratory kit (cerebral ischaemia (CIS)-test) that detects autoantibodies to the N-terminus domain of the NR2A subunit in the blood of patients with TIA or stroke (Gusev E. I., Skvortsova V I, Alekseev A A, Izykenova G A, Dambinova S A. S. S Korsakov's J. Neurol. & Psych. 1996; 5:68-72). The N-terminus domain of the NR2A subunit of NMDA

receptors was selected as the immunoreactive epitope on the basis of molecular biological and experimental studies showing that this epitope is the most immunoreactive region of the receptor (Dambinova S A, Izykenova G A. J. High Nervous Activity. 1997; 47:439-446).

[0018] More recently, researchers have reported a correlation between the effectiveness of a stroke treatment regimen and the levels of autoantibodies to the NR2A and NR2B subunits of NMDA. In particular, these researchers have reported increased titers of autoantibodies to the NR2A and NR2B subunits of NMDA in the blood of patients severely affected by stroke, and a reduction of the autoantibodies, accompanied by an improvement in neurological function, during therapy by glycine—a non-specific agonist of NMDA receptors (Gusev et.al. Cerebrovascular Diseases. 2000, 10: 49-60). Patients that responded positively to glycine had lower autoantibody titers than patients who were not treated, and had levels of autoantibodies that were close to the levels of autoantibodies in control subjects.

[0019] Unfortunately, the use of NR2A and NR2B autoantibodies in the diagnosis of stroke or TIA does not provide a real-time assessment of the damage being done by a stroke or TIA. Rather, because of the time the immune system requires to mount an immune response, and to generate NR2A and NR2B autoantibodies, methods that test for these antibodies at best provide a delayed assessment of the extent and severity of stroke or TIA.

[0020] Investigators from Canada (Hill M. D., Jackowski G., Bayer N., Lawrence M., Jaeschke R. Can. Med. Assoc. J. 2000, 163: 1139-1140) have proposed a new diagnostic laboratory assay for differentiating stroke subtype. They designed a preliminary prospective cohort study to test a panel of biochemical markers (neuron-specific enolase [NSE], myelin basic protein [MBP], S-100 [beta] protein and thrombomodulin [Tm]) in blood samples from patients with acute ischemic stroke. These markers were chosen because they cover important cellular components of the brain that might be damaged in acute stroke. The 4 biochemical markers were assayed using a standard ELISA technique.

[0021] The results of this investigation demonstrated elevated levels of NSE in 89% of the patients admitted in hospitals, Tm in 43%, MBP in 39% and S-100 [beta] in 32%. At least one of the markers was elevated on admission in 93% of the acute stroke patients. By stroke type, 100% of the patients with lacunar stroke, 100% of those with posterior circulation stroke and 90% of those with partial anterior circulation stroke had elevated NSE levels on admission. Conversely, none of the patients with lacunar stroke had an elevated S-100 [beta] level initially or subsequently. Peak levels of NSE, S-100 [beta] and MBP, but not of Tm, were significantly correlated with admission NIHSS scores ($p < 0.05$).

[0022] For stroke, 3 hours is an outside limit for administering appropriate therapies. The focus must change from extensive evaluation before any action to a well-planned acute emergency therapy developed using an appropriate diagnostic strategy. Every future advance to improve the outcome after TIA/stroke will depend on a fast initial response—within minutes and not hours (Marler J. R. Annl. Emergency Med. 1999, 33: 450-451). Therefore, it is especially important to develop a fast and simple method (within

one hour) of detecting brain and blood biomarkers capable of recognizing the initial processes of TIA/stroke before irreparable ischemic damage ensues.

OBJECTS OF INVENTION

[0023] Therefore, it is an object of the invention to provide biochemical methods and kits for diagnosing central nervous system disorders such as TIA and stroke.

[0024] It is another object of the present invention to improve upon the accuracy of currently available methods for diagnosing TIA and stroke, and to more accurately diagnose TIA and stroke to the exclusion of other nervous system disorders or traumatic brain injury.

[0025] It is still another object of the present invention to provide methods of diagnosing stroke using biochemical markers that distinguish between hemorrhagic and ischemic stroke.

[0026] Still another object of the invention is to provide biochemical analyses of the extent and progression of TIA or stroke, or the infarction resulting from the TIA or stroke.

[0027] It is another object of the present invention to provide rapid biochemical methods and kits for diagnosing TIA and stroke, to provide real-time assessments of TIA or stroke, within a window of time that permits effective therapeutic intervention.

[0028] It is another object of the present invention to provide rapid and inexpensive biochemical methods and kits for diagnosing TIA and stroke, which can be performed at frequent intervals to monitor the progression of a TIA or stroke, or the effectiveness of therapy administered against TIA or stroke.

[0029] Still another object of the present invention is to provide diagnostic methods and kits for assessing the risk of incurring a TIA or stroke, and for monitoring the remission of risk factors for TIA or stroke.

[0030] Still another object of the invention is to provide a panel of rapid multiple panel of biomarkers for assessing the nature, severity and progression of TIA or stroke, and thereby to enable a more effective selection of intervention therapy.

SUMMARY OF THE INVENTION

[0031] It has unexpectedly been discovered that levels of circulating NMDA receptor proteins or fragments thereof can be assessed using diagnostic kits and processes, and that levels of these proteins or fragments can be used to clinically evaluate patients suffering from ischemic central nervous system disorders such as stroke or TIA. When analyzed in combination with other biomarkers for stroke and TIA, such as the thromboembolic marker homocysteine, or the excitatory amino acid glutamate, these proteins can diagnose the existence of a stroke with remarkable accuracy (generally greater than 89%). In contrast, the efficacy of single parameters for early diagnosis of stroke is 58% for glutamate, 66% for homocysteine, and 79% for NMDA receptors. The rapid evaluation of these neural ischemic biomarkers in an emergency room setting will greatly enhance the confidence of physicians when diagnosing stroke or TIA, and significantly improve the speed at which therapy against the stroke or TIA can be administered.

[0032] The biomarkers also yield extensive evidence about the nature of the stroke or TIA and the type therapy which should be administered. For example, the respective levels of biomarkers can be evaluated to determine whether the patient is suffering an ischemic or hemorrhagic stroke, or whether the patient is suffering from a traumatic brain injury. The data from the biomarkers can also be used to monitor or evaluate the progression of the ischemic episode, as well as the damage that has resulted as a consequence of the ischemia. High levels of all parameters reflect the neurological deficit and may be also used for prognosis of disease outcome. Moreover, a relationship has been observed between the respective levels of the biomarkers and the degree of thromboembolic and neurotoxicity in brain processes under the stroke. Once again, these relationships can be put to extensive use when evaluating the choice of emergency therapy in short time frames, such as anti-platelet and neuroprotective therapy. The data can be used independently of other diagnostic strategies, but preferably forms an integral part of a comprehensive diagnostic strategy employing conventional diagnostic techniques.

[0033] The data obtained from the NMDA biomarkers, especially when combined with data from other biomarkers such as glutamate and homocysteine, can also be used to monitor the efficacy of a treatment regime. It has surprisingly been found that the NMDA biomarkers provide real time evidence of neurotoxicity, and that reductions in levels of circulating NR1A or NR2A NMDA receptors or fragments thereof correspond well with reductions in neurotoxic mechanisms. By obtaining data at appropriate intervals using rapid laboratory techniques such as latex agglutination, one is able to monitor the progression of the episode in response to the therapeutic regime.

[0034] A latex agglutination technique has also been developed which dramatically increases the speed of diagnosis obtained by the methods of this invention, and thereby improves the effectiveness of the methods in emergency-room settings. The technique can be adapted for use in the detection of NMDA receptors, homocysteine, glutamate, or any other suitable biomarker against central nervous system disorders. Using the latex agglutination technique, one is able to provide real-time biochemical diagnosis and monitoring of TIA/stroke patients (within about 30 minutes), and thereby dramatically improve the effectiveness of response to TIA/stroke. This is surprising because these biomarkers are naturally occurring and, in contrast to viruses for which latex agglutination methods were originally developed, show much lower strengths of association with their corresponding antibodies.

[0035] This semi-quantitative method gives reliable data quickly in a format that is simple for interpretation. Surprisingly, the technique shows greater accuracy than even well established methods based upon HPLC and ELISA. The application of the latex agglutination technique to the analysis of brain biomarkers for stroke will decrease the cost of analysis, provide the opportunity to monitor real-time progress of a treatment procedure, and allow physicians to determine the efficacy of medication administered in the treatment of TIA or stroke.

[0036] The methods of the present invention also can be employed in a non-emergency setting, when evaluating the risk that an individual will suffer a stroke or TIA. In addition,

based upon results showing an increased risk of suffering TIA or stroke, prevention therapy can be administered, and the effectiveness of the therapy monitored using the methods of the present invention.

[0037] The invention also relates to indirect methods for measuring levels of NR2A and NR2B NMDA receptor proteins or fragments thereof. Thus, analytical techniques can be used to evaluate indirect measures of NR2A and NR2B NMDA receptor proteins or fragments thereof, such as autoantibodies specific for the proteins, or cDNA that encodes for the proteins.

[0038] Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION OF THE INVENTION

[0039] The present invention may be understood more readily by reference to the following detailed description of preferred embodiments of the invention and the Examples included therein. Before the present methods and techniques are disclosed and described, it is to be understood that this invention is not limited to specific analytical or synthetic methods as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

DEFINITIONS AND USE OF TERMS

[0040] As used in this specification and in the claims which follow, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a fragment" includes mixtures of fragments, reference to "an cDNA oligonucleotide" includes more than one oligonucleotide, and the like.

[0041] An analogue of a protein, peptide, or polypeptide means a protein, peptide, or polypeptide that contains one or more amino acid substitutions, deletions, additions, or rearrangements. For example, it is well known in the art of protein biochemistry that an amino acid belonging to a grouping of amino acids having a particular size or characteristic (such as charge, hydrophobicity, and hydrophilicity) can often be substituted for another amino acid without altering the activity of the protein, particularly in regions of the protein that are not directly associated with biological activity. Thus, an analogue of an NMDA receptor or fragment thereof is useful in the present invention if it includes amino acid substitutions, deletions, additions or rearrangements at sites such that antibodies raised against the analogue are still specific against the NMDA receptor or fragment.

[0042] Preferably, an NMDA analogue has at least 80%, 85%, 90%, or 95% amino acid identity with naturally occurring NMDA. Amino acid identity is defined by a

analogy comparison between the analogue and naturally occurring NMDA. The two amino acid sequences are aligned in such a way that maximizes the number of amino acids in common along the length of their sequences; gaps in either or both sequences are permitted in making the alignment in order to maximize the number of common amino acids. The percentage amino acid identity is the higher of the following two numbers: (1) the number of amino acids that the two polypeptides have in common with the alignment, divided by the number of amino acids in the NMDA analogue or fragment thereof, multiplied by 100, or (2) the number of amino acids that the two polypeptides have in common with the alignment, divided by the number of amino acids in naturally occurring NMDA or fragment thereof, multiplied by 100.

[0043] NMDA derivatives, and derivatives of NMDA fragments, include naturally occurring NMDA and NMDA analogues and fragments thereof that are chemically or enzymatically derivatized at one or more constituent amino acids, including side chain modifications, backbone modifications, and N- and C-terminal modifications, by for example acetylation, hydroxylation, methylation, amidation, phosphorylation or glycosylation. The term also includes NMDA salts such as zinc NMDA and ammonium NMDA.

[0044] A protein or peptide is measured "directly" in the sense that the protein or peptide is itself measured in the biological sample, as opposed to some other indirect measure of the protein or peptide such as autoantibodies to the protein or peptide, or cDNA associated with the expression of the protein or peptide.

[0045] The term "antibody" is intended to be synonymous with "immunoglobulin." As used herein, the term "antibody" is meant to include both the native antibody, and biologically active derivatives of antibodies, such as, for example, Fab', F(ab')₂ or Fv as well as single-domain and single-chain antibodies. A biologically active derivative of an antibody retains the ability to bind antigen.

General Discussion

[0046] The present disclosure describes diagnostic and therapeutic applications that result from the realization that genetic or accidental increase of NMDA receptors synthesis in the brain reflects a neurological ischemic deficit, and may be used for early diagnoses of stroke or TIA. NMDA receptors that are abnormally expressed in the brain are quickly metabolized and, following penetration of the blood brain barrier, these metabolic destruction products enter the circulatory system. The immune system recognizes these peptides and protein fragments as foreign antigens and responds by generating autoantibodies to them.

[0047] In one aspect of the present invention, the correlation between increased NMDA receptor synthesis, and the appearance of high levels of the receptors in blood sera of individuals during the initial stages of cerebral ischemia, is used for diagnostic and therapeutic applications. Experiments in rats with focal ischemia have demonstrated that NR2A mRNA expression in the cortex and hippocampus can be measured within two hours of the onset of the ischemic episode, and thus provide an opportunity for real time measurement of ischemic processes and damage resulting therefrom. At the same time, meaningful expression of NR2C and NR2D mRNA is not observed in brain structures

that showed no changes in NR1 mRNA expression in rat ischemic brain. These changes in NR2-receptor mRNA expression in the early stages of ischemia are observed prior to morphological evidence of neuronal damage or appearance of autoantibodies to them in blood serum specimens.

[0048] Thus, in one aspect the present invention provides a method for diagnosing a central nervous system disorder comprising measuring the level of NR2A and/or NR2B NMDA receptor or fragment thereof in a biological sample. Elevated levels of NR2A and NR2B NMDA receptors are specific to brain injury, and are expressed in ischemic brain tissue at higher rates than other NMDA receptors, and thus are uniquely suited for assessing ischemic brain episodes such as TIA or stroke. Baseline levels for determining whether the measured levels are elevated, and hence indicative of a central nervous system disorder, can be obtained from population norms or, preferably, from a patient's own test history.

[0049] The biological sample tested for the receptor or fragment can be derived from blood, urine, blood plasma, blood serum, cerebrospinal fluid, saliva, perspiration, or brain tissue. In a preferred embodiment, the biological sample is a blood sample. In an even more preferred embodiment the biological sample is a blood sample diluted to a ratio of from about 1:2 to about 1:32 (v:v).

[0050] Immunoassay techniques are generally preferred for measuring the proteins or peptides of the present invention, as discussed in greater detail herein, although other analytical techniques are also available as known to those skilled in the art, such as HPLC. The amino acid sequences of the NR2A and NR2B subunits, and antigenic fragments thereof, are recited in SEQ ID NOS. 1, 2, 3, 10, 11, and 12, and any fragment of these sequences can be employed in methods for directly detecting the receptors as long as sufficient antigenicity is maintained. However, when using immunoassays it has been found that the antigenic determinants are concentrated in the N-terminal domain of the NR2A and/or NR2B NMDA receptor, and that antibodies raised against the N-terminal domains and fragments thereof should be employed for optimal test results. The inventors have sequenced the amino acid chain of the N-terminal domains for these receptors, and set forth the sequences as SEQ ID NOS. 2 and 11, respectively, at the end of this document.

[0051] In a preferred embodiment, other biomarkers of central nervous system disorders are also measured to improve the accuracy of the diagnosis, and to provide further information about the nature, severity, or progression of the disorder. Particularly useful markers are directly implicated in the NMDA receptor pathway, and include naturally occurring agonists and antagonists of the NMDA receptors. An exemplary antagonist is glycine. Exemplary agonists include glutamate, polyglutamate, aspartate, polyaspartate, homocysteine, and polyhomocysteine. A particularly preferred agonist for measuring the activity of the receptors is glutamate or polyglutamate.

[0052] In another embodiment, thromboembolic biomarkers are measured to obtain a simultaneous reading of the likelihood for clotting in the brain. Exemplary thromboembolic biomarkers include homocysteine or polyhomocysteine.

[0053] Titers of higher than 2.63 for combined levels of NR2A and NR2B, especially when combined with titers

higher than 3.34 for glutamate and/or 2.23 for homocysteine, are remarkably predictive of the occurrence of stroke and typically justify immediate therapeutic intervention for the TIA or stroke or risk of stroke. These titers can be translated into absolute concentrations by reference to the examples hereof.

[0054] The methods of the present invention are preferably performed by directly measuring the levels of NR2A and/or NR2B biomarkers in a selected biological sample, using immunoassay techniques employing antibodies raised against the biomarkers, or through quantitative techniques such as HPLC. However, it is also possible to measure the presence of the NR2A and/or NR2B biomarkers indirectly. This can be done by directly measuring autoantibodies of the biomarkers, or by directly measuring the cDNA nucleic acid intermediates involved in expression of these biomarkers. If autoantibodies are measured, they are preferably measured using one or more antigenic fragments of the NR2A and/or NR2B receptors as the target of the antibody, as opposed to a whole NR2A and/or NR2B protein. Healthy persons generally have NR2A autoantibodies in an amount of about 1.0-2.0 ng/ml. Healthy persons generally have NR2A cDNA levels of about 1.0-1.5 pg/ml.

Latex Agglutination and Other Diagnostic Techniques

[0055] A number of immunoassays can be employed in accordance with the principles of the present invention. Examples include radioimmunoassays, enzyme immunoassays (e.g. ELISA), immunofluorescence, immunoprecipitation, latex agglutination, hemagglutination, and histochemical tests. A particularly preferred method, however, because of its speed and ease of use, is latex agglutination.

[0056] Latex agglutination assays have been described in Beltz, G. A. et al., in *Molecular Probes: Techniques and Medical Applications*, A. Albertini et al., eds., Raven Press, New York, 1989, incorporated herein by reference. In the latex agglutination assay, antibody raised against a particular biomarker is immobilized on latex particles. A drop of the latex particles is added to an appropriate dilution of the serum to be tested and mixed by gentle rocking of the card. With samples lacking sufficient levels of the biomarkers, the latex particles remain in suspension and retain a smooth, milky appearance. However, if biomarkers reactive with the antibody are present, the latex particles clump into visibly detectable aggregates.

[0057] An agglutination assay can also be used to detect biomarkers wherein the corresponding antibody is immobilized on a suitable particle other than latex beads, for example, on gelatin, red blood cells, nylon, liposomes, gold particles, etc. The presence of antibodies in the assay causes agglutination, similar to that of a precipitation reaction, which can then be detected by such techniques as nephelometry, turbidity, infrared spectrometry, visual inspection, colorimetry, and the like.

[0058] The term latex agglutination is employed generically herein to refer to any method based upon the formation of detectable agglutination, and is not limited to the use of latex as the immunosorbent substrate. While preferred substrates for the agglutination are latex based, such as polystyrene and polypropylene, particularly polystyrene, other well-known substrates include beads formed from glass, paper, dextran, and nylon. The immobilized antibodies may

be covalently, ionically, or physically bound to the solid-phase immunoabsorbent, by techniques such as covalent bonding via an amide or ester linkage, ionic attraction, or by adsorption. Those skilled in the art will know many other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

[0059] Thus, in one embodiment, the method of measuring the NR2A and/or NR2B NMDA receptor, fragment thereof, or other biomarker is by latex agglutination comprising:

[0060] (i) contacting the biological sample with poly- or monoclonal antibodies bound on an agglutinating carrier to target biomarkers for a sufficient time period and under conditions to promote agglutination; and

[0061] (ii) reading a signal generated from the agglutination; wherein the amount of signal detected correlates to the titer of biomarkers present in the sample.

[0062] The reaction is preferably read macroscopically against a dark background for a sufficient time period. The method preferably yields a clinically useful reading within about 30 minutes or less.

[0063] It has been experimentally found that latex beads having a mean diameter of from about 0.25 to about 0.4 μm are particularly preferred in the practice of this invention. The poly- or monoclonal antibodies are preferably present in a ratio with the latex beads of about 1:1.

[0064] Latex beads having the foregoing characteristics can be prepared generally by adding antibodies to the target biomarker to a carrier solution that contains a 1% concentration (by weight) of latex beads, until the concentration of the antibodies in the carrier solution reaches about 2 mg/ml, and allowing the ingredients a sufficient time to covalently link, typically about 1 hour, in the presence of a linking agent such as glutaraldehyde.

[0065] Conventional methods can be used to prepare the antibodies. For example, by using a peptide of a NMDA protein, polyclonal antisera or monoclonal antibodies can be made using standard methods. A mammal, (e.g., a mouse, hamster, or rabbit) can be immunized with an immunogenic form of the peptide (preferably the NR2A and/or NR2B receptor, an antigenic determinant of the NR2A and/or NR2B receptor, or an analogue or derivative thereof) which elicits an antibody response in the mammal. Techniques for conferring immunogenicity on a peptide include conjugation to carriers or other techniques well known in the art. For example, the peptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassay procedures can be used with the immunogen as antigen to assess the levels of antibodies. Following immunization, antisera can be administered and, if desired, polyclonal antibodies isolated from the sera.

[0066] To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused with myeloma cells by standard somatic cell fusion procedures thus immortalizing these cells and yielding hybridoma cells. Such techniques are well known in the art, (e.g., the hybridoma technique originally developed by Kohler and Milstein (*Nature* 256, 495-497 (1975))) as well as other techniques such as the human B-cell

hybridoma technique (Kozbor et al., *Immunol. Today* 4, 72 (1983)), the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al. *Monoclonal Antibodies in Cancer Therapy* (1985) Allen R. Bliss, Inc., pages 77-96), and screening of combinatorial antibody libraries (Huse et al., *Science* 246, 1275 (1989)). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with the peptide and the monoclonal antibodies can be isolated. Therefore, the invention also contemplates hybridoma cells secreting monoclonal antibodies with specificity for NR2A or NR2B NMDA proteins or fragments thereof as described herein.

[0067] When the NR2A and/or NR2B receptors are detected indirectly, by measuring the cDNA expression of the NR2A and/or NR2B receptors, the measuring step in the present invention may be carried out by traditional PCR assays such as cDNA hybridization, Northern blots, or Southern blots. These methods can be carried out using oligonucleotides encoding the polypeptide antigens of the invention. Therefore, in one embodiment the methods are performed employing oligonucleotides that encode the amino acid sequence of SEQ ID NO: 2, which is preferably represented by nucleotides 371-1978 of SEQ ID NO: 6. More preferably, the nucleic acid construct comprises a oligonucleotide consisting of a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 3, which is preferably represented by oligonucleotides 1790-1852 of SEQ ID NO: 7.

[0068] Thus, in one embodiment the methods of this invention include measuring an increase of NR2A and/or NR2B cDNA expression by contacting the total DNA isolated from a biological sample with oligonucleotide primers attached to a solid phase, for a sufficient time period. In another preferred embodiment, NR2A and/or NR2B cDNA expression is measured by contacting an array of total DNA bound to a solid matrix with a ready-to-use reagent mixture containing oligonucleotide primers for a sufficient time period. Expressed NR2A cDNA is revealed by the complexation of the cDNA with an indicator reagent that comprises a counterpart oligonucleotide to the cDNA attached to a signal-generating compound. The signal-generating compound is preferably selected from the group consisting of horseradish peroxidase, alkaline phosphatase, urinase and non-enzyme reagents. The signal-generating compound is most preferably a non-enzyme reagent.

[0069] In a preferred embodiment, the solid phase is a polymer matrix. More preferably, the polymer matrix is polyacrylate, polystyrene, or polypropylene. In one preferred embodiment the solid phase is a microplate. In another preferred embodiment, the solid phase is a nitrocellulose membrane or a charged nylon membrane.

[0070] As mentioned above, the methods of performing the present invention also may be performed by measuring the levels of autoantibodies specific for the NR2A and/or NR2B subunits. These autoantibodies may be measured by any suitable immunoassay such as, for example, a radioimmunoassay, an immunofluorescence assay, an enzyme-linked immunosorbent assay (ELISA), an immunocytochemical assay, and immunoblotting. In a preferred embodiment, the antigen to which the anti-NR2A and/or NR2B autoantibodies bind is a polypeptide or protein fragment of the N-terminal domain of the NR2A and/or NR2B

receptor. More preferably, the antigen comprises a polypeptide or protein fragment of amino acid SEQ ID NO:2, 3, 4, 11, 12, or 13, or a suitable analogue or derivative thereof.

[0071] Thus, in yet another embodiment the methods of the present invention are performed by measuring the levels of anti-NR2A and/or anti-NR2B autoantibodies, by contacting a biological sample with a polypeptide or protein fragment of the NR2A and/or NR2B receptor (preferably the N-terminal domain) (or an analogue or derivative thereof) attached to a solid phase, for a sufficient time period and under conditions to allow a complex to form between any NR2A and/or NR2B autoantibodies which may be present in the sample and the polypeptide or protein fragment, contacting the complex with an indicator reagent comprising a secondary antibody specific for the species of the mammal attached to a signal-generating compound (or for the polypeptide or protein fragment); and measuring the signal generated. The peptide can be obtained directly from biological samples, by recombinant DNA techniques, or by direct chemical synthesis. The signal-generating compound is preferably selected from horseradish peroxidase, alkaline phosphatase, and urinase. More preferably, the signal-generating compound is horseradish peroxidase. Most preferably, the indicator reagent is rabbit anti-human IgG attached to horseradish peroxidase. The amount of signal detected is correlated to the amount of anti-NR2A and/or NR2B autoantibodies present in the biological sample.

[0072] In this method it is preferred that the solid phase be a polymer matrix. More preferably, the polymer matrix is selected from the group consisting of polyacrylate, polystyrene, and polypropylene. In one preferred embodiment the solid phase is a microplate. In another preferred embodiment, the solid phase is a nitrocellulose membrane or a charged nylon membrane.

[0073] The immunosorbent of the present invention for measuring levels of autoantibody can be produced as follows. A fragment of the receptor protein is fixed, preferably by covalent bond or an ionic bond, on a suitable carrier such as polystyrene or nitrocellulose. If the standard polystyrene plate for immunological examinations is employed, it is first subjected to the nitration procedure, whereby free nitrogroups are formed on the plate surface, which are reduced to amino groups and activated with glutaric dialdehyde serving as a linker. Next the thus-activated plate is incubated with about 2 to 50 nM of the target peptide for the purpose of chemically fixing the respective immunogenic fragment of the receptor protein for a time and at a temperature sufficient to assure fixation (i.e. for about 16 hours at 4° C.).

[0074] The amount of protein below 2 nM affects adversely the reliability of the findings, whereas its amount exceeding 50 nM is inexpedient due to an increase in the nonspecific binding of autoantibodies with the immunosorbents. The plate is then washed with an aqueous solution of sodium boron hydride and an aqueous solution of sodium chloride, vacuum-dried, enclosed in a hermetically sealed package, and put under storage at 4° C.

[0075] It is also practicable to produce the immunosorbent by fixing the respective fragment of the receptor protein on nitrocellulose strips by virtue of ionic interaction. The respective fragment of the receptor protein isolated from the mammals' brain is applied to nitrocellulose and incubated for 15 min at 37° C. Then nitrocellulose is washed with a

0.5% solution of Tween-20, and the resultant immunosorbent is dried at room temperature and stored in dry place for one year period.

Emergency Room Diagnosis and Prognosis

[0076] As mentioned above, the methods of the present invention are especially well suited for use in emergency room settings. There are two reasons for this. First, the method is extremely useful in an emergency room setting because NR2A and NR2B NMDA receptor levels are elevated at a very early stage of ischemic insult, and thus provide a real time indication of neurotoxic events. This is in contrast to autoantibodies which require that an immune response first be mounted by the insulted organism.

[0077] The second reason the method is useful in an emergency room setting is the speed and ease with which the latex agglutination procedure can be employed. Using the latex agglutination processes described herein, one is able to turn laboratory results around often in less than 30 or even 20 minutes. Thus, using the methods of the present invention real-time data can be obtained that permits a therapeutic response within the window for an effective response to stroke.

[0078] Therefore, in one embodiment the invention provides a method for diagnosing the existence of a central nervous system disorder such as TIA or stroke, further comprising withdrawing the biological sample from a human subject, wherein the biological sample is withdrawn within three hours of the onset of symptoms of the central nervous system disorder. In still another embodiment of the invention, the amount of time elapsed between withdrawing the biological sample from the subject, and detecting or measuring the presence or quantity of the NR2A and/or NR2B NMDA receptor, is less than about one hour, 45 minutes, or 30 minutes.

[0079] One of the principal advantages of the present invention is the ability to distinguish ischemic episodes such as stroke from other brain injuries such as traumatic brain injury. Thus, in another embodiment, the invention provides a method for diagnosing the existence of TIA or stroke further comprising evaluating from the level of NR2A and/or NR2B NMDA receptor whether the brain injury is a traumatic brain injury or stroke/TIA, and administering traumatic brain injury or stroke/TIA therapy as appropriate.

[0080] Another advantage of the methods of the present invention which is extremely useful in an emergency room setting, is the ability to determine from the test data the type of stroke involved. In particular, if a stroke is suspected, the method will help diagnose whether the stroke is an ischemic or hemorrhagic insult. Thus, in another embodiment the invention provides a method for diagnosing the existence of TIA or stroke further comprising, when the diagnosis confirms a stroke, evaluating from the level of NR2A and/or NR2B NMDA receptor whether the stroke is ischemic or hemorrhagic and administering ischemic or hemorrhagic stroke therapy as appropriate.

[0081] Another advantage of the present invention is the ability to evaluate infarction volume and extent of neurotoxicity from NMDA expression data. NMDA receptor expression research in an animal model of middle carotid artery occlusion has been employed to demonstrate such correlation. Thus, in still another embodiment the invention

provides a method for diagnosing the existence of TIA or stroke further comprising, if TIA and/or stroke is confirmed, evaluating from the level of NR2A and/or NR2B NMDA receptor cranial infarct volume, and administering therapy appropriate to the infarct volume.

[0082] Moreover, one can periodically repeat the procedure, to provide continuous monitoring of a patient's state as interventional therapy is administered, to monitor the effectiveness of a particular therapeutic regime. In this embodiment, it is preferable for the mammal to be concurrently undergoing treatment for the disorder. More preferably, the samples are collected at intervals from about 20 min to about 1 month. Even more preferably, the interval is from about 20 min. to about 2 hours. Most preferably the samples are collected at an interval of about 30 minutes. Thus, in still another embodiment the invention provides a method for diagnosing the progression of TIA or stroke further comprising detecting or measuring the presence or quantity of a NR2A and/or NR2B NMDA receptor in a biological sample one or more additional times, at a frequency of less than about 6 hours.

Primary Care Physician Setting

[0083] In another application the method is used in a clinical setting to determine an individual's risk of stroke, or to monitor the effectiveness of risk reduction therapies. As mentioned above, a number of therapies can be employed to reduce the risk of stroke in an individual. The use of antiplatelet agents, particularly aspirin, is a standard treatment for patients at risk for stroke. People with atrial fibrillation (irregular beating of the heart) may be prescribed anticoagulants. The most important treatable factors linked to TIAs and stroke are high blood pressure, cigarette smoking, heart disease, carotid artery disease, diabetes, and heavy use of alcohol. Medical help is available to reduce and eliminate these factors. Lifestyle changes such as eating a balanced diet, maintaining healthy weight, exercising, and enrolling in smoking and alcohol cessation programs can also reduce these factors. When these therapies are administered it is desirable to determine the effectiveness of the therapy.

[0084] Therefore, in one embodiment the invention provides a method for evaluating an individual's risk for TIA or stroke comprising measuring levels of NR2A and/or NR2B NMDA receptors or fragments thereof in a biological sample from the individual, and comparing the levels to a baseline level. In one embodiment the baseline levels are derived from population averages. In another embodiment the baseline levels are derived from the individual's own medical history.

[0085] In another embodiment the method is performed more than once to monitor the reduction or increase in risk for stroke or TIA, optionally in conjunction with the administration of risk reduction therapy. In one embodiment the method is performed at a frequency of from about one week to about six months. In another embodiment the method is performed at a frequency of from about one month to about three months.

[0086] In a particularly preferred embodiment other biomarkers are also measured to assess the risk for stroke or TIA. Particularly preferred biomarkers for risk of stroke or TIA are glutamate and homocysteine.

Novel Kits of the Present Invention

[0087] In another embodiment the invention provides kits for diagnosing central nervous system disorders such as TIA, stroke, and traumatic brain injury. NR2A and/or NR2B antibodies or antigens may be incorporated into immunoassay diagnostic kits depending upon whether autoantibodies or NMDA receptors are being measured. A first container may include a composition comprising an antigen or antibody preparation. Both antibody and antigen preparations should preferably be provided in a suitable titrated form, with antigen concentrations and/or antibody titers given for easy reference in quantitative applications.

[0088] The kits may also include an immunodetection reagent or label for the detection of specific immunoreaction between the provided antigen and/or antibody, as the case may be, and the diagnostic sample. Suitable detection reagents are well known in the art as exemplified by radioactive, enzymatic or otherwise chromogenic ligands, which are typically employed in association with the antigen and/or antibody, or in association with a second antibody having specificity for first antibody. Thus, the reaction is detected or quantified by means of detecting or quantifying the label. Immunodetection reagents and processes suitable for application in connection with the novel methods of the present invention are generally well known in the art.

[0089] The reagents may also include ancillary agents such as buffering agents and protein stabilizing agents, e.g., polysaccharides and the like. The diagnostic kit may further include where necessary agents for reducing background interference in a test, agents for increasing signal, apparatus for conducting a test, calibration curves and charts, standardization curves and charts, and the like.

[0090] In a more particular aspect the invention relates to a rapid multiple panel containing antibodies to the thromboembolic and neurotoxicity biomarkers glutamate, homocysteine and NMDA receptors that employs latex agglutination. Thus, in one embodiment the invention provides a kit for diagnosing central nervous system disorders comprising: (1) an agglutinating immunosorbent for NR2A and/or NR2B NMDA receptors or fragments thereof, and (2) a control such as saline or a known concentration of NR2A and/or NR2B receptor or fragment thereof. In a more preferred embodiment the kit further comprises an agglutinating immunosorbent for another biomarker for TIA/stroke, such as an agonist or antagonist of NR2A and/or NR2B, a thromboembolic marker, or more particularly glutamate or polyglutamate, and/or an agglutinating immunosorbent for homocysteine or polyhomocysteine. The agglutinating immunosorbent is preferably of the type discussed in greater detail above.

[0091] In another embodiment the invention relates to a kit for detecting NR2A and/or NR2B receptors or fragments thereof that does not employ latex agglutination. Thus, in another embodiment the invention provides a kit for diagnosing central nervous system disorders comprising: (1) an immunosorbent for NR2A and/or NR2B NMDA receptors or fragments thereof, and (2) an indicator reagent comprising secondary antibodies attached to a signal generating compound. The secondary antibodies can be specific for the receptor or fragment, or for the primary antibodies in the immunosorbent. In a preferred embodiment the kits further comprise an immunosorbent for glutamate or polyglutamate,

and/or an immunosorbent for homocysteine or polyhomocysteine, and secondary antibodies against the glutamate and/or homocysteine, or to the primary antibodies on the immunosorbents against the glutamate or homocysteine. The immunosorbent preferably comprises anti-antibodies for the biomarkers bound to a solid support.

[0092] In another aspect the present invention relates to a test-kit that relies upon PCR amplification for measuring NR2A and/or NR2B levels. Thus, in another embodiment the invention provides a kit comprising: (a) one or more oligonucleotide primers (preferably of SEQ ID NO: 8) attached to a solid phase, (b) indicator reagent attached to a signal-generating compound capable of generating a detectable signal from oligonucleotides, and (c) a control sample (i.e. template cDNA). The reagents may also include ancillary agents such as buffering agents, polymerase agents, and the like. The diagnostic kit may further include, where necessary, other members of the signal-producing system of which system the detectable group is a member (e.g., enzyme and non-enzyme substrates), agents for reducing background interference in a test, agents for increasing the signal, apparatus for conducting a test, and the like.

[0093] In another embodiment of test-kit comprises (a) a solid phase to which biological fluids for receiving total DNA including NR2A cDNA could be attached, (b) oligonucleotide primers, preferably in a ready-to-use PCR buffer, and (c) a control sample (i.e. template cDNA). Ancillary agents as described above may similarly be included.

[0094] In another embodiment the invention provides a diagnostic kit for detecting NR2A and/or NR2B autoantibodies comprising (a) a polypeptide of the N-terminal domain of the NR2A and/or NR2B receptor, fragment thereof, or analog or derivative thereof, (b) an indicator reagent comprising a secondary antibody specific for the autoantibody or the polypeptide attached to a signal-generating compound; and (c) a control sample, such as a known concentration of NR2A and/or NR2B polyclonal antibodies. The reagents may also include ancillary agents such as buffering agents and protein stabilizing agents, e.g., polysaccharides and the like. The diagnostic kit may further include, where necessary, other members of the signal-producing system of which system the detectable group is a member (e.g., enzyme and non-enzyme substrates), agents for reducing background interference in a test, agents to increase the signal, apparatus for conducting a test, calibration and standardization information or instructions, and the like.

Novel Compositions of the Invention

[0095] The methods of the present invention rely upon a series of novel compositions which themselves form a part of the invention. Thus, in one series of embodiments the invention provides an isolated polypeptide fragment of the NR2A and/or NR2B NMDA receptor, comprising:

[0096] 1. An antigenic determinant of the NR2A NMDA receptor,

[0097] 2. An antigenic determinant of the NR2B NMDA receptor,

[0098] 3. The N-terminal domain of the NR2A NMDA receptor,

[0099] 4. The N-terminal domain of the NR2B NMDA receptor,

- [0100] 5. SEQ ID NO. 2,
- [0101] 6. SEQ ID NO. 3,
- [0102] 7. SEQ ID NO. 4,
- [0103] 8. SEQ ID NO. 11,
- [0104] 9. SEQ ID NO. 12, and
- [0105] 10. SEQ ID NO. 13,

[0106] or an antigenic fragment, analog, or derivative thereof. In another series of embodiments the invention provides any of the foregoing polypeptides linked covalently to a distinct antigenic determinant, such as human serum albumin. In still another series of embodiments the invention provides any of the foregoing polypeptides linked to any of the immunosorbent materials discussed above. The immunosorbent can be in the form of a bead for latex agglutination, in the size ranges discussed above, or in the form of a synthetic plate for conventional immunoassay analysis. The polypeptide can be linked to the immunosorbent using any conventional means of linkage, including covalent linkage, ionic linkage, and adsorption.

[0107] In another series of embodiments the present invention relates to the novel monoclonal and polyclonal antibodies specific for and/or raised against the foregoing polypeptides, including the foregoing polypeptides linked to distinct antigenic determinants. Thus, in one embodiment the invention provides non-human antibodies against any of the foregoing peptides or polypeptides or antigenic fragment, analog, or derivative thereof. In another embodiment the invention provides immunosorbents to which such antibodies are linked.

[0108] In another series of embodiments the present invention provides oligonucleotides that encode the foregoing peptides and polypeptides and fragments, analogs, and derivatives thereof, and to recombinant expression vectors that include such oligonucleotides. Such oligonucleotides include, without limitation, the oligonucleotides defined by SEQ ID NO:6, 7, 14, and 15, and fragments thereof which encode antigenic determinants.

[0109] In still another embodiment the present invention relates to isolated oligonucleotide sequences that are useful in the cDNA PCR analytical techniques of the present invention. Thus, the invention further provides oligonucleotides comprising the nucleotide sequences of SEQ ID NOS:7, 8, 9, 15, 16, and 17.

BRIEF DESCRIPTION OF THE SEQUENCE LISTINGS

[0110] The features, aspects, and advantages of the present invention will become better understood with regard to the following sequence listings where, in the sequence the recited amino acid position numbering reflects that used throughout this document.

[0111] SEQ ID NO:1. shows the full-length amino acid sequence of the mature NR2A receptor subunit, as follows:

Sequence Listing

Peptide

[0112] *Homo sapiens* glutamate receptor. ionotropic, N-methyl D-aspartate 2A

[0113] Science 256:1217-1221(1992)

```

NCBI/NM 000833.2
      1      11      21      31      41      51
1  MGRLGYWTL  VLPALLVWRD PAQNAAAEKG PPALNIAVLL GHSHDVTERE LRNLWGPEQA 60
61  TGLPLDVNVV ALLMNRTPDK SLITHVCDLM SGARIHGLVF GDDTDQEAVA QMIDFISSQT 120
121 FIPILGIHGG ASMIMADKDP TSTFFQFGAS IQQQATVMLK IMQDYDWHVE SLVTTIFPGY 180
181 RDFISFJKTT VDNSFVGWDM QNVITLDTSF EDAKTQVQLK KIHSSVILLY CSKDEAVLIL 240
241 SEARSLGLTG YDFFW1VPSL VSGNTELIPIK EFPSGLISVS YDDWDYSLEA RVRDGLGILT 300
301 TAASSMLEKF SYIPEAKASC YGQAEKPETP LHTLHQFMVN VTWDGKDLSE TEEGYQVHPR 360
361 LVVJVLNKDR EWEKVGKVEN QTLSLRHAVW PRYKSFSDCE PDDNHLISIVT LEEAPFVIVE 420
421 DIDPLTETCV RNTVPCRKFV KINNSTNEGM NVKCKCGFC IDILKKLSRT VKFTYDLYLV 480
481 TNGKHGKKVN NVWNGMIGEV VYQRAVMAVG SLTLNEERSE VVDFSVPFVE TGISVMVSRS 540
541 NGTVSPSAFL EPFSASVWVM MFVMLLIVSA LAVFVFEYFS PVGYNRNLAK GKAPHGPSFT 600
601 IGKAIWLLWG LVFNNSVPVQ NPKGTTSKIM VSVWAPFAVI FLASYTANLA AFMIQEEFVD 660
661 QVTGLSDKKF QRPDYSPPF RFGTVPNGST ERNIRNNYPY MHQYMTFRNQ RGVEDALVSL 720
721 KTGKLDAFIY DAAVLNYKAG RDEGCKLVTI GSGYIFASTG YGIALQKQSP WKRQIDLALL 780
781 QFVGDGEMEE LETLWLTGIC HNEKNEVMSS QLDIDNMAGV FYMLAAMAL SLITFIWEHL 840
841 FYWKLRFCTF GVCSDRPGLL FISISRGIYSC IHGVHIEEK KSPDFNLTGS QSNMLKLLRS 900
901 AKNISNMSNM NSSRMDSPKR ATDFIQRQSL IVDMVSCKGN LIYSDNRSFQ GKDSIFGDNM 960
    
```

-continued

961 NELQTFVANR HKDNLSNYVF QGQHPLTLNE SNPNTVEVAV STESKGNRP RQLWKKSME 1020
 1021 LRQDSLQNP VSQRDEKTAE NRTHSLKSPR YLPEEVAHSD ISETSSRATC HREPDNNKNH 1080
 1081 KTKDNFKRSM ASKYPKDCSD VDRTYMKTKA SSPRKIYTI DGEKEPSFHL DPPQFVENIT 1140
 1141 LPENGVFPDT YQDHNEFRK GDSTLPMNRN PLHNEDGLPN NDQYKLYAKH FTLKDKGSPH 1200
 1201 SEGSDRYRQN STHCRSCLSN LPTYSGHFTM RSPFKCACL RMGNLYDIDE DQMLQETGNP 1260
 1261 ATREEVYQQD WSQNNALQFQ KNKLRINRQH SYDNILDKPR EIDLRSRPS ISLKDRERLL 1320
 1321 EGNLYGSLFS VPSSKLLGNK SSLFPQGLE DSKRSKLLPD HASDNPFLHT YGDDQRLVIG 1380
 1381 RCPSPDPYKHS LPSQAVNSY LRSSLRSTAS YCSRDSRGS DVYISEHVMP YAANKNTMYS 1440
 1441 TPRVLNSCSN RRVYKMP SI ESDV

[0114] SEQ ID NO:2. shows the amino acid sequence of the auto-antigenic region of the N-terminal domain of the NR2A subunit, as follows:

[0115] SEQ ID NO:2

[0118] SEQ ID NO:4. shows a 22 amino acid antigenic peptide, corresponding to a fragment of the NR2A N-terminal domain another such peptide, modified by an N-terminal Cys for attachment to a carrier protein):

HOMO SAPIENS

PAQNAAAEKG PPALNIAVLL GHSEVTERE LRNLWGPEQA 60
 61 TGLPLDVNVV ALLMNRDTPK SLITHVCDLM SGARIHGLVF GDDTDQEAVA QMLDFISSQT 120
 121 FIPILGIHGG ASMIMADKDP TSTFFQFGAS IQQQATVMLK IMQDYDWHVF SLVTTIFPGY 180
 181 RDFISFIKTT VDNSFVGWDM QNVITLDTSF EDAKTQVQLK KIHSSVILLY CSKDEAVLIL 240
 241 SEARSLGLTG YDFFWIVPSL VSGNTELIK EPFSGSLISVS YDDWDYSLEA RVRDGLGLLT 300
 301 TAASSMLEKF SYIPEAKASC YGQAEKPEP LHTLHQFMVN VTWDGKDLSE TEEGYQVHPR 360
 361 LVVIVLNKDR EWEKVGKVEN QTLNLRHAWV PRYKSFSDCE PDDNHSIIVT LEEAPFVIVE 420
 421 DIDPLTETCV RNTVPCRKFV KINNSTNEGM NVKCKCKGFC IDILKLSRT VKFTYDLYLV 480
 481 TNGKHGKQVN NVWNGMIGEV VYQRAVMAVG SLT1NEERSE VVDFSVPFVE TGISVMVSR 540
 541 NGTVSPSAFL EPFSAS

[0116] SEQ ID NO:3; shows a 21 amino acid antigenic peptide, corresponding to a fragment of the NR2A N-terminal domain another such peptide (21 amino acids derived from the NR2A sequence and an N-terminal Cys for attachment to a carrier protein), as follows:

SEQ ID NO:3

[0117] *Homo sapiens*

NGMIGEVVYQRAVMAVGLTI

[0119] Artificial Sequence

CNGMIGEVVYQRAVMAVGLTI

Full

Base Count Origin

[0120] *Homo sapiens* glutamate receptor, ionotropic, N-methyl D-aspartate 2A (GRIN 2A) mRNA

[0121] SEQ ID NO:5. shows the Oligonucleotide position numbering used throughout in reference to NR2A oligonucleotide sequences, as follows:

SEQ ID NO:5

[0122] Science 256:1217-1221(1992)

[0123] May 22, 1992

NIGB/NM_000833

1 atcatgggac cgggtgagcg ctgagaatcg cggccgcagc catcagccct ggagatgacc
61 aggagcggcc actgctgaga actatgtgga gagaggctgc gagccctgct gcagagcctc
121 cggctgggat agccgcccc cgtggggcg atgcggacag cgcgggacag ccaggggagc
181 gcgctggggc cgcagcatgc gggaaaccgc taaaccggg ggctgctgag gcggccgaga
241 tgctcgtgcg cgcagcgcgc cccactgcat cctcgacctt ctcgggctac agggaccgtc
301 agtggcgact atgggcagag tgggctattg gacctgctg gtgctgccgg cccttctggt
361 ctggcgcggt ccggcgcga gcgcggcggc ggagaagggt cccccgcgc taaatattgc
421 ggtgatgctg ggtcacagcc acgacgtgac agagcgcgaa ctctgaacac tgtggggccc
481 cgagcagcgg cgcgggctgc ccctggacgt gaacgtggtg gctctgctga tgaaccgcac
541 cgaccccaag agcctcatca cgcacgtgtg cgacctcatg tccggggcac gcatccacgg
601 cctcgtgttt ggggacgaca cggaccagga ggcgtagcc cagatgctgg attttatctc
661 ctcccacacc ttctgcccc tcttgggcat tcatgggggc gcatctatga tcatggctga
721 caaggatccg acgtctacct tcttccagtt tggagcgtcc atccagcagc aagccacggt
781 catgctgaag atcatgcagg attatgactg gcatgtcttc tcctgggtga ccaactatctt
841 ccctggctac agggaattca tcagcttctg caagaccaca gtggacaaca gctttgtggg
901 ctgggacatg cagaatgtga tcacactgga cacttccttt gaggatgcaa agacacaagt
961 ccagctgaag aagatccact cttctgtcat cttgctctac tgttccaaag acgaggctgt
1021 tctcattctg agttagggcc gctcccttgg cctcaccggg tatgatttct tctggattgt
1081 ccccagcttg gtctctggga acacggagct catcccaaaa gagtttccat cgggactcat
1141 ttctgtctcc tacgatgact gggactacag cctggaggcg agagtgaggg acggcattgg
1201 catcctaacc accgctgcat cttctatgct ggagaagttc tcctacatcc ccgaggccaa
1261 ggccagctgc tacgggcaga tggagaggcc agaggtccc atgcacacct tgcaccatt
1321 tatggtcaat gttacatggg atggcaaaga cttatccttc actgaggaag gctaccaggt
1381 gacccccagg ctggtggtga ttgtgctgaa caaagaccgg gaatgggaaa aggtgggcaa
1441 gtgggagaac catcagctga gcctgaggca cgcctgtggt cccagggtaca agtccttctc
1501 cgactgtgag ccgatgaca accatctcag catcgtcacc ctggaggagg ccccatctgt
1561 catcgtgga gacatagacc ccctgaccga gacgtgtgtg aggaacaccg tgccatgtcg
1621 gaagtctcgc aaaatcaaca attcaaccaa tgaggggatg aatgtgaaga aatgctgcaa
1681 ggggttctgc attgatattc tgaagaagct ttccagaact gtgaagtta cttacgacct
1741 ctatctggtg accaatggga agcatggcaa gaaagttaac aatgtgtgga atggaatgat
1801 cggatgaagt gtctatcaac gggcagtcac ggcagttggc tcgctcacca tcaatgagga
1861 acgttctgaa gtggtgact tctctgtgcc ctttgtggaa acgggaatca gtgtcatggt
1921 ttcaagaagt aatggcaccg tctcaccttc tgcttttota gaaccattca gcgcctctgt
1981 ctgggtgatg atgtttgtga tgotgctcat tgtttctgcc atagctgttt ttgtctttga
2041 atacttcagc cctgttggat acaacagaaa cttagccaaa gggaaagcac cccatgggccc

-continued

2101 ttcttttaca attgaaaaag ctatatggct tctttggggc ctgggtttca ataactccgt
2161 gctgtccag aatcctaaag ggaccaccag caagatcatg gtatctgtat gggccttctt
2221 cgctgtcata ttcttgcta gctacacagc caatctggct gccttcatga tccaagagga
2281 atttgtggac caagtgaccg gcctcagtga caaaaagttt cagagaccctc atgactattc
2341 cccacctttt cgatttggga cagtgcctaa tggaaacacg gagagaaca ttcggaataa
2401 ctatccctac atgcatcagt acatgaccaa atttaatcag aaaggagtag aggacgcctt
2461 ggtcagcctg aaaacgggga agctggacgc tttcatctac gatgccgcag tcttgaatta
2521 caaggctggg agggatgaag gctgcaagct ggtgaccatc gggagtgggt acatctttgc
2581 caccaccggt tatggaattg cccttcagaa aggctctcct tggaaagagc agatcgacct
2641 ggccttgctt cagtttggg gtgatggtga gatggaggag ctggagacct tgtggctcac
2701 tgggatctgc cacaacgaga agaacgaggt gatgagcagc cagctggaca ttgacaacat
2761 ggcgggctga ttctacatgc tggctgccgc catggccctt agcctcatca ccttcatctg
2821 ggagcacctc ttctactgga agctgcgctt ctgtttcacg ggcgtgtgct ccgaccggcc
2881 tgggttgcct ttctccatca gcaggggcat ctacagctgc attcatggag tgcacattga
2941 agaaaagaag aagtctccag acttcaatct gacgggatcc cagagcaaca tgttaaaact
3001 cctccggtea gccaaaaaca tttccagcat gtccaacatg aactcctcaa gaatggactc
3061 acccaaaaga gctgctgact tcatccaaag aggttccctc atcatggaca tggtttcaga
3121 taaggggaat ttgatgtact cagacaacag gtcctttcag gggaaagaga gcatttttgg
3181 agacaacatg aacgaactcc aacatttgt ggccaaccgg cagaaggata acctcaataa
3241 ctatgtattc cagggacaac atctcttac tctcaatgag tccaacccta acacggtgga
3301 ggtggccgtg agcacagaat ccaaagcgaa ctctagacct cggcagctgt ggaagaaatc
3361 cgtggattcc atacgccag attcactatc ccagaatcca gtctcccaga gggatgaggc
3421 aacagcagag aataggacct actccctaaa gagccctagg tatcttccag aagagatggc
3481 ccaactctgac atttcagaaa cgtcaaatcg ggccacgtgc cacagggaac ctgacaacag
3541 taagaaccac aaaaccaag acaactttaa aaggtcagtg gcctccaaat accccaagga
3601 ctgtagtgag gtcgagcgca cctacctgaa aaccaaatca agctccccta gagacaagat
3661 ctacactata gatgggtgaga aggagcctgg tttccactta gatccacctc agtttgttga
3721 aaatgtgacc ctgcccgaga acgtggactt cccggacctc taccaggatc ccagtgaaaa
3781 cttccgcaag ggggactcca cgctgccaat gaaccggaac cccttgcata atgaagaggg
3841 gctttccaac aacgaccagt ataaactcta ctccaagcac ttcaccttga aagacaaggg
3901 ttccccgcac agtgagacca gcgagcgata ccggcagaac tccacgcact gcagaagctg
3961 cctttccaac atgcccacct attcaggcca cttcaccatg aggtccccct tcaagtgcga
4021 tgctgcctg cggtggggga acctctatga catcgatgaa gaccagatgc ttcaggagac
4081 aggtaaccga gccaccgggg agcaggtcta ccagcaggac tgggcacaga acaatgccct
4141 tcaattacaa aagaacaagc taaggattag ccgtcagcat tcctacgata acattgtcga
4201 caaacctagg gagctagacc ttagcaggcc ctcccggagc ataagcctca aggacagggg
4261 acggcttctg gaggaaatt tttacggcag cctgtttagt gtccccctca gcaactctc
4321 ggggaaaaaa agctcccttt tcccccaagg tctggaggac agcaagagga gcaagtctct
4381 cttgccagac cacacctccg ataacccttt cctccactcc cacagggatg accaacgctt

-continued

4441 ggttattggg agatgccctt cggaccctta caaacactcg ttgccatccc aggcggtgaa
 4501 tgacagctat cttcggctcg ccttgaggtc aacggcatcg tactgttcca gggacagtcg
 4561 gggccacaat gatgtgtata tttcggagca tgttatgcct tatgctgcaa ataagaataa
 4621 tatgtactct acccccaggg ttttaaattc ctgcagcaat agacgcgtgt acaagaaaat
 4681 gcctagtatc gaatctgatg tttaaaaatc ttccattaat gttttatcta tagggaaata
 4741 cacgtaatgg ccaatgttct ggagggtaaa tgttgatgt ccaatagtgc cctgctaaga
 4801 ggaagaagat gtagggaggt attttgttgt tgttggtgtt ggctcttttg cacacggctt
 4861 catgccataa tcttccactc aaggaatctt gtgaggtgtg tgctgagcat ggcagacacc
 4921 agataggtag gtccttaacc aaaaataact aactacataa gggcaagtct ccgggacatg
 4981 cctactgggt atgttggcaa taatgatgca ttggatgcca atggtgatgt tatgatttcc
 5041 tatattccaa attccattaa ggtcagccca ccatgtaatt ttctcatcag aaatgcctaa
 5101 tggtttctct aatacagaat aagcaatag gtgtgcatgt aaacctgaca cagacaaaat
 5161 aaaaacagtt aagaatgcat ctgcactgta gtcggatttg aacatgtgca agagattagg
 5221 aagtttgctt cgtaaacagtt tcagctttct tgttatgcct tccatcacag ccagggctca
 5281 ccccaagaac tccaggctcc cctaaagaat agcaaatcag tgtgttcgtg atgactgtgc
 5341 taccttcatt atagttcatt tccaagacac atctggagcc aaaggcccga gggaccctca
 5401 ggtggggaga gctacaggaa tctctttgga tgttgatgtg tgtttctctc taccctcggc
 5461 ttogatggct ttgttcagag ctgcataaac taacacattt atgtctccga gatctaagtg
 5521 tggatcttct gtctgtgaca cagtggccat tgtagtattt ccgaagacg cctatgtacg
 5581 taagtttgca tttcctccct tctggtgatg actcagggtt gtatagtatc tgttaccctt
 5641 tccctccag agtaaccata actcgttccg tttccaaaca gccatggtgg tgtccaatta
 5701 gctgtgatc gctcttccca gagttgttaa tgtggtgaca tgcaccaaca gccgtatgtg
 5761 tactgtgatc tgtaagaagt acaatgccat ctgtctgccg aaggctagca tggttttagg
 5821 tttatcttcc ttcacatcca gaaattctgt tggacactca ctccacccc aaactcctca
 5881 aatcaaaagc cttcaaaaca cgaggcactc ttggatctac cctgagtatc ctccaaactg
 5941 tggatacagt ttagtgagac aagcaatttc tccctctga gttattctct ctgttggtgg
 6001 caaaccactt catagcacca acagagatgt aggaaaaatt cctcaaagta tttgtcattt
 6061 ctgagtcgac tgcattatcc cattcttatt ctctcaaac ctgtgcatat atgacatgaa
 6121 atgatatcca tttttttttt aagttagaaa cagagagggg aatacttatg catggggagc
 6181 ctgtagcacc agtgcctgcc acaaaaacaa gtgccccga caagatagtt gctatgttat
 6241 gacactttct cagateagga ttttctagtt taaaaattaa atatcataaa acg

[0124] SEQ ID NO:6. shows the oligonucleotide sequence of the auto-antigenic region of the N-terminal domain of the NR2A subunit, as follows:

[0125] SEQ ID NO:6

N-terminal nucleotide sequence

371 ccggcgcgca gcgcggcggc ggagaagggt cccccgcgc taaatattgc

421 ggtgatgctg ggtcacagcc acgacgtgac agagcgcgaa cttcgaacac tgtggggccc

-continued

481 cgagcaggcg gcggggctgc ccctggacgt gaacgtggta gctctgctga tgaaccgcac
 541 cgaccccaag agcctcatca cgcacgtgtg cgacctcatg tccggggcac gcatccacgg
 601 cctcgtgttt ggggacgaca eggaccagga ggccgtagcc cagatgctgg attttatctc
 661 ctcccacacc ttcgtcccca tcttgggcat tcatgggggc gcatctatga tcatggctga
 721 caaggatccg acgtctacct tcttccagtt tggagcgtcc atccagcagc aagccacggt
 781 catgctgaag atcatgcagg attatgactg gcatgtcttc tcctggtga ccaactatct
 841 ccctggctac agggaattca tcagcttcgt caagaccaca gtggacaaca gctttgtggg
 901 ctgggacatg cagaatgtga tcacactgga cacttccttt gaggatgcaa agacacaagt
 961 ccagctgaag aagatccact cttctgtcat cttgctctac tgttccaaag acgaggctgt
 1021 tctcattctg agtgaggccc gctcccttgg cctcaccggg tatgattttct tctggattgt
 1081 ccccagcttg gtctctggga acacggagct catcccaaaa gagtttccat cgggactcat
 1141 ttctgtctcc tacgatgact gggactacag cctggaggcg agagtgaggg acggcatttg
 1201 catcctaacc accgctgcat cttctatgct ggagaagttc tcctacatcc ccgaggccaa
 1261 gccagctgc tacgggcaga tggagaggcc agagggtccc atgcacacct tgcaccatt
 1321 tatggtcaat gttacatggg atggcaaaga cttatccttc actgaggaag gctaccaggt
 1381 gcaccccaag ctggtggtga ttgtgctgaa caaagaccgg gaatgggaaa aggtgggcaa
 1441 gtgggagaac catacgtga gcctgaggca gcctgtgtgg cccaggtaca agtccttctc
 1501 cgactgtgag ccgatgaca accatctcag catcgtcacc ctggaggagg cccattctgt
 1561 catcgtgaa gacatagacc ccctgaccga gacgtgtgtg aggaacaccg tgccatgtcg
 1621 gaagttctgc aaaatcaaca attcaaccaa tgaggggatg aatgtgaaga aatgctgcaa
 1681 ggggttctgc attgatattc tgaagaagct ttccagaact gtgaagtta cttacgacct
 1741 ctatctggtg accaatggga agcatggcaa gaaagttaac aatgtgtgga atggaatgat
 1801 cggtgaagtg gtctatcaac gggcagtcac ggcagttggc tcgctacca tcaatgagga
 1861 acgttctgaa gtggtggact tctctgtgcc ctttgtgaa acgggaatca gtgcatggt
 1921 ttcaagaagt aatggcaccg tctcaccttc tgettttcta gaaccattca gcgcctct

[0126] SEQ ID NO:7 shows a 62 oligonucleotide fragment target, as follows:

[0127] SEQ ID NO:7

atggaatgatcgggtgaagtggctctatcaacgggcagtcatggcagttggc

tcgctcaccatc

[0128] SEQ ID NO:8 shows one oligonucleotide primer, as follows:

[0129] SEQ ID NO:8

agcatggcaagaaagttaaca

[0130] SEQ ID NO:9 shows a second oligonucleotide primer, as follows:

[0131] SEQ ID NO:9

acgttctgaagtggtagactt

[0132] SEQ ID NO:10. shows the full-length amino acid sequence of the mature NR2B receptor subunit, as follows:

Peptide

[0133] *Homo sapiens* glutamate receptor, ionotropic, N-methyl D-aspartate 2B

[0134] Biochim. Biophys. Acta 1260:105-108(1995).

sequence NME2_HUMAN (Q13224)

1 11 21 31 41 51

1 MKPRAECCSP KFWLVLAFLA VSGSRARSOKSPPSIGIAVILVGTSDVEVAIKDAHEKDDDFH60

61HLSVVPVELVAMNETDPKSIITRICDLMSDRKIQGVVFAADTDQEAIAOILDFISAOTL120

121 TPILGIHGGS SMIMADKDES SMFFQFGPSI EQQASVMLNI MEEYDWYIFS IVTTYFPGYQ 180

181 DFNKIRSTI ENSFVGWELE EVLLLDMSLD DGDSKIQNL KKLQSPIILL YCTKEEATYI 240

241 FEVANSVGLT GYGYTWIVPS LVAGDQDTVP AEFPTGLISV SYDEWDYGLP ARVRDGI AII 300

301 TTAASDMLSE HSFPEPKSS CYNTEKRIY QSNMLNRYLI NVTFEGRNLS FSEDGYQMHP 360

361 KLVIIILLNKE RKWERVGVKWK DKSLOMKYYV WPRMCPETEE QEDDHLISIVT LEEAPFVIVE 420

421 SVDPLSGTCM RNTVPCOKRI VTENKTDEEP GYIKKCKGF CIDILKISK SVKFTYDLYL 480

481 VTNGKHGKKI NGTWNMIGE VVMKRAYMAV GSLTINEERS EVVDFSVVPI ETGISVMVSR 540

541 SNGTVSPSAF LEFVSADVVW MMFVMLLIVS AVAVFVFEYF SPVGYNRCLA DGREPGGPF 600

601 TIGKAIWLLW GLVFNNVSPV QNPKGTTSKI MVSVWAFFAV TFLASYTANL AAFMIQEYV 660

661 DQVSGLSDKK FQRPNDFSP FRFGTVPNGS TERNIRNNYA EMUAYMGKFN QRGVDDALLS 720

721 LKTGKLDLAFI YDAAVLNMA GRDEGCKLVT IGSGKVPFAST GYGIAIQKDS GWKRQVDLAI 780

781 LQLFGDGEME ELEALWLTGI CHNEKNEVMS SQLDIDNMAG VFYMLGAAMA LSLITFICEH 840

841 LFYWQFRHCF MGVCSGKPGM VFSISRGIYS CIHGAIEER QSMNSPTAT MNNTHSNLR 900

901 LLRTAKNMAN LSGVNGSPQS ALDFIRRESS VYDISEHRRS FTHSDCKSYN NPPCEENLFS 960

961 DYISEVERTF GNLQLKDSNV YQDHYHHHR PHSIGSASSI DGLYDCDNP FTTQSRISK 1020

1021 KPLDIGLPSS KHSQSLDLYG KFSFKSDRYG GHDDLIRSDV SDISTHTVTY GNTGNAAKR 1080

1081 RKQYKDSLK KRPASAKSRR EFDEIELAYR RPPRSPDHK RYFRDKEGLR DFYLDQFRTK 1140

1141 ENSPHWEHVD LTDIYKERSD DFKRDSVSGG GPCTNRSHIK HGTGDKHGVV SGVPAPWEKN 1200

1201 LTNVEWEDRS GGNFCRSCPS KLHNYSTTVT QNSGRQACI RCEACKKAGN LYDISEDNSL 1260

1261 QELDQPAAPV AVTASNATTK YPQSPNTSKA QKKNRNKLR QHSYDTFVDL QKEEAALAPR 1320

1321 SVSLKDKGRF MDGSPYAHMF EMSAGESTFA NNKSSVPTAG HHHHNNPGGG YMLSKSLYPD 1380

1381 RVTQNPFIPT FGDDQCLLHG SKSYFFRQPT VAGASKARPD FRALVTNKPV VSALHGAVPA 1440

1441 RFQKDCIGN QSNPCVPNNK NPROFNGSSN GHVYEKLSI

[0135] SEQ ID NO:11. shows the amino acid sequence of the auto-antigenic region of the N-terminal domain of the NR2B subunit, as follows:

[0136] SEQ ID NO:11

Homo sapiens

RSQK SPPSIGIAVI LVGTSDVEVAI KDAHEKDDDFH 60

61 HLSVVPVEL VAMNETDPKS IITRICDLMS DRKIQGVVFA DTDQEAIAQ ILDFISAQTL 120

121 TPILGIHGGS SMIMADKDES SMFFQFGPSI EQQASVMLNI MEEYDWYIFS IVTTYFPGYQ 180

181 DFNKIRSTI ENSFVGWELE EVLLLDMSLD DGDSKIQNL KKLQSPIILL YCTKEEATYI 240

241 FEVANSVGLT GYGYTWIVPS LVAGDQDTVP AEFPTGLISV SYDEWDYGLP ARVRDGI AII 300

301 TTAASDMLSE HSFPEPKSS CYNTEKRIY QSNMLNRYLI NVTFEGRNLS FSEDGYQMHP 360

361 KLVIIILLNKE RKWERVGVKWK DKSLOMKYYV WPRMCPETEE QEDDHLISIVT LEEAPFVIVE 420

-continued

421 SVDPLSGTCM RNTVPCQKRI VTENKTDEEP GYIKKCKGF CIDILKKISK SVKFTYDLYL 480

481 VTNGKHGKKI NGTWNGMIGE VVMKRAYMAV GSLTINEERS EVVDFSVVPI ETGISVMVSR 540

541 SNGTVSPSAF LEPFSAD

[0137] SEQ ID NO:12; shows a 20 amino acid antigenic peptide fragment of the NR2B subunit, as follows:

[0140] SEQ ID NO:13.

[0138] SEQ ID NO:12

Artificial Sequence (21 amino acids)
CGYIKKCKGF CIDILKKISK

Homo sapiens
GYIKKCKGF CIDILKKISK

Full
Base Count Origin

[0139] SEQ ID NO:13 shows a 21 amino acid sequence of an antigenic fragment of the NR2B subunit modified by an N-terminal Cys for attachment to a carrier protein, as follows:

[0141] SEQ ID NO:14 shows the oligonucleotide position numbering used throughout in reference to NR2B oligonucleotide sequences, as follows:

[0142] SEQ. NO. 14

Homo sapiens glutamate receptor, ionotropic, N-methyl D-aspartate 2B mRNA

1 ttgaatttgc atctcttcaa gacacaagat taaaacaaaa tttacgctaa attggatttt
61 aaattatctt ccgttcattt atccttcgtc tttcttatgt ggatattgcaa gcgagaagaa
121 gggactggac attccaaca tgctcactcc cttaactctgt ccgtctagag gtttggtctc
181 tacaaccaa gggagtcgac gagttgaaga tgaagcccag agcggagtgc tgttctccca
241 agttctgggt ggtgttgccc gtccctggccg tgtcaggeag cagagctcgt tctcagaaga
301 gccccccag cattggcatt gctgtcatcc tcgtgggcaac ttcgacgag gtggccatca
361 aggatgocca cgagaaagat gatttcacc atctctccgt ggtaccccg gtggaactgg
421 tagccatgaa tgagaccgac ccaaagaca tcatcaccg catctgtgat ctcatgtctg
481 accggaagat ccagggggtg gtgtttgctg atgacacaga ccaggaagcc atcgcccaga
541 tcctcgattt catttcagca cagactctca ccccgatcct gggcatccac gggggtcct
601 ctatgataat ggcagataag gatgaatcct ccatgttctt ccagtttggc ccatcaattg
661 aacagcaagc ttccgtaatg ctcaacatca tggagaata tgactggtac atcttttcta
721 tcgtcaccac ctatttcctt ggctaccagg actttgtaaa caagatccgc agcaccattg
781 agaatagctt tgtgggctgg gagctagagg aggtcctcct actggacatg tcctgggacg
841 atggagattc taagatccag aatcagctca agaaacttca aagccccatc attcttcttt
901 actgtaccaa ggaagaagcc acctacatct ttgaagtggc caactcagta gggctgactg
961 gctatggcta cacgtggatc gtgccagtc tgggtggcagg ggatacagac acagtgcctg
1021 cggagttccc cactgggctc atctctgtat catatgatga atgggactat ggcctccccg
1081 ccagagttag agatggaatt gccataatca ccaactgctc ttctgacatg ctgtctgagc
1141 acagcttcat cctgagccc aaaagcagtt gttacaacac ccacgagaag agaatctacc
1201 agtccaatat gctaaatagg tatctgatca atgtcacttt tgaggggagg aatttgcct
1261 tcagtgaaga tggctaccag atgcaccga aactgggtgat aattcttctg aacaaggaga
1321 ggaagtggga aagggtgggg aagtggaaa acaagtccct gcagatgaag tactatgtgt
1381 ggccccgaat gtgtccagag actgaagagc aggaggatga ccatctgagc attgtgacct

-continued

1441 tggaggaggc accatttgtc attgtgaaa gtgtggacc tctgagtga acctgcatga
1501 ggaacacagt cccctgcca aaacgcatag tcaactgaga taaaacagac gaggagccgg
1561 gttacatcaa aaaatgctgc aaggggttct gtattgacat ccttaagaaa atttctaatt
1621 ctgtgaagtt cacctatgac ctttacctg ttaccaatgg caagcatggg aagaaaatca
1681 atggaacctg gaatggtatg attggagagg tggcatgaa gagggcctac atggcagtgg
1741 gctcactcac catcaatgag gaacgatcgg aggtggtoga cttctctgtg cccttcatag
1801 agacaggcat cagtgtcatg gtgtcacgca gcaatgggac tgtctcacct tctgccttct
1861 tagagccatt cagcgtgac gtatgggtga tgatgttgt gatgctgctc atcgtctcag
1921 ccgtggctgt ctttgtcttt gagtacttca gccctgtggg ttataacagg tgcctcgtg
1981 atggcagaga gcctgggtga ccctctttca ccactcggca agctatttgg ttgctctggg
2041 gtctggtgtt taacaactcc gtacctgtgc agaaccctaaa ggggaccacc tccaagatca
2101 tgggtcagc gtggcccttc tttgctgca tcttctctggc cagctacact gccaaactag
2161 ctgccttcat gatccaagag gaatatgtgg accaggtttc tggcctgagc gacaaaaagt
2221 tccagagacc taatgacttc tcaccccctt tccgcttgg gaccgtgccc aacggcagca
2281 cagagagaaa tattcgcaat aactatgagc aatgcatgc ctacatggga aagttcaacc
2341 agaggggtgt agatgatgca ttgctctccc tgaacacagg gaaactggat gccttcatct
2401 atgatgcagc agtgctgaac tatatggcag gcagagatga aggctgcaag ctggtgacca
2461 ttggcagtgg gaaggtcttt gcttccactg gctatggcat tgccatccaa aaagattctg
2521 ggtggaagcg ccaggtggac cttgctatcc tgcagctctt tggagatggg gagatggaag
2581 aactggaagc tctctggctc actggcattt gtcacaatga gaagaatgag gtcatgagca
2641 gccagctgga cattgacaac atggcagggg tcttctacat gttggggcg gccatggctc
2701 tcagcctcat caccttcatc tgogaacacc ttttctattg gcagttccga cattgcttta
2761 tgggtgtctg ttctggcaag cctggcatgg tcttctccat cagcagaggt atctacagct
2821 gcatccatgg ggtggcgatc gaggagcggc agtctgtaat gaactcccc accgcaacca
2881 tgaacaacac aactccaac atcctcggcc tgetgcgcac ggccaagaac atggctaacc
2941 tgtctggtgt gaatggtca ccgcagagcg ccctggactt catccgacgg gagtcatccg
3001 tctatgacat ctcaagcagc cgcgcagct tcacgcattc tgactgcaa tcctacaaca
3061 acccgcctg tgaggagaac ctotctcagt actacatcag tgaggtagag agaacttctg
3121 ggaacctgca gctgaaggac agcaactgtt accaagatca ctaccaccat caccaccggc
3181 cccatagtat tggcagtgcc agctccatcg atgggctcta cgactgtgac aaccaccct
3241 tcaccacca gtccaggctc atcagcaaga agcccctgga catcggcctc ccctcctcca
3301 agcacagcca gctcagtgac ctgtaccgca aattctcctt caagagcagc cgctacagtg
3361 gccacgacga cttgatccgc tccgatgtct ctgacatctc aaccacacc gtcacctatg
3421 ggaacatcga gggcaatgcc gccaaaggc gtaagcagca atataaggac agcctgaaga
3481 agcggcctgc ctccgccaag tcccgcaggg agtttgacga gatcagagctg gcctaccgtc
3541 gccgaccgcc ccgtccctc gaccacaagc gctacttcag ggacaaggaa gggctacggg
3601 acttctacct ggaccagtgc cgaacaaaag agaactcacc cactgggag cacgtagacc
3661 tgaccgacat ctacaaggag cggagtgtat actttaagcg cgactccatc agcggaggag

-continued

3721 ggccctgtac caacaggtct cacatcaagc acgggacggg cgacaaacac ggcgtggtca
3781 gcgggggtacc tgcaccttg gagaagaacc tgaccaacgt ggagtgggag gaccgggtccg
3841 ggggcaactt ctgccgcagc tgtccctcca agctgcacaa ctactccacg acggtgacgg
3901 gtcagaactc gggcaggcag gcgtgcatcc ggtgtgaggc ttgcaagaaa gcaggcaacc
3961 tgtatgacat cagtgaggac aactccctgc aggaactgga ccagccggct gccccagtgg
4021 cggtgacgtc aaacgcctcc accactaagt accctcagag cccgactaat tccaaggccc
4081 agaagaagaa ccggaacaaa ctgcccggc agcactccta cgacaccttc gtggacctgc
4141 agaaggaaga agccgcctg gccccgcga gcgtaagcct gaaagacaag ggccgattca
4201 tggatgggag cccctacgcc cacatgtttg agatgtcagc tggcgagagc acctttgcca
4261 acaacaagtc ctcagtgcc actgcccggc atcaccacca caacaacccc ggcggcgggt
4321 acatgctcag caagtcgctc taccctgacc gggtcacgca aaacccttcc atccccactt
4381 ttggggacga ccagtgttg ctccatggca gcaaatccta cttcttcagg cagcccacgg
4441 ttggcggggc gtcgaaagcc aggccggact tccgggcccct tgcaccaac aagccgggtg
4501 tctcggccct tcatggggcc gtgccagccc gtttccagaa ggacatctgt ataggaacc
4561 agtccaaccc ctgtgtgctt aacaacaaa accccagggc tttcaatggc tccagcaatg
4621 ggcatgttta tgagaaactt tctagtattg agtctgatgt ctgagtgagg gaacagagag
4681 gttaagggtg gtacgggagg gtaaggctgt gggtcgcgtg atgcgcatgt cacggagggt
4741 gacgggggtg aacttggttc ccatttgctc ctttctgtt ttaatttatt tatgggatcc
4801 tggagtcttg gttcctactg ggggcaaccc tggtgaccag caccatctct cctccttttc
4861 acagttctct ctttcttccc ccogctgtca gccattctct tcccatgag atgatgccat
4921 gggccctctc agcaggggag ggtagagcgg agaaaggaag ggctgcatgc gggcttcctc
4981 ctggtgtgga agagctcctt gatatcctct ttgagtgaag ctgggagaac caaaaagagg
5041 ctatgtgagc acaaagtag cttttcccaa actgatcttt tcathtaggt gaggaagcaa
5101 aagcatctat gtgagaccat ttagcacact gcttgtgaaa gaaaagaggc tctggctaaa
5161 ttcagtctgc ttagatgaca tctgtctagg aatcatgtgc caagcagagg ttgggaggcc
5221 atttgtgttt atatataagc ccaaaaatgc ttgcttcaac cccatgagac tcgatagtgg
5281 ttggtaacag aaccaaggt cattggtggc agagtggatt cttgaacaaa ctggaaagta
5341 cgttatgata gtgtccccc gtgccttggg gacaagagca ggtggattgt gcgtgcatgt
5401 gtgttcatgc acacttgac ccatgtgtag tcagggtcct caagagaagg caaccttgac
5461 tctttctatt gtttcttca atatcccaa gcagtgtgat tgtttggctt atatacagac
5521 agagatggcc atgtattacc tgaattttg ctgtgtctcc cttcatcctt ctggaataag
5581 gagaatgaaa attcttgata aagaagattc tgtggtctaa acaaaaaaag gcggtgagca
5641 atcctgcaag aacaaggtac ataaacaagt cctcagtggt tggcaattgt ttcaaccagt
5701 ttgaaccaag aactttccag gaaggctaaa gggaaaccga attttcacag ccatgattct
5761 tttgccaca cttgggagca aaagattcta caaagctctt ttgagcattt agactctcga
5821 ctggccaagg tttggggaag aocgaagcca ctttgaaga agtaaggagt cgtgtatggt
5881 agggtaagtg agagaggggg atgtttccaa tgctttgatc ctttcttact taacctgaag
5941 ctagacgagc aggtcttctc cccccaaaac tgattacaac tgctacagag cagacagtta
6001 agagaaatga gcttgacctt taagagaaat gagctgcact ccatgagtgc agctctggag

-continued

6061 gtacgaaaag aggggaagag acttggaat gggagacggg gccagagagg gaccctccac

6121 caccctctttg gccctggctc cctgggaatg tgacttgagc ccagagtga cactcttggg

6181 agaagccctt ctaccttctt gcaacacctt gttccctct cagattgtac cattgag

[0143] SEQ ID NO:15 shows a 60 oligonucleotide fragment target, as follows:

[0144] SEQ ID NO:15

g gttacatcaa aaaatgctgc aaggggttct gtattgacat
ccttaagaaa atttctaaa

[0145] SEQ ID NO:16 shows one oligonucleotide primers (21 nucleotides), as follows:

[0146] SEQ ID NO:16

tcactgagaa taaaacagac g

[0147] SEQ ID NO:17 shows one oligonucleotide primers (21 nucleotides), as follows:

[0148] SEQ ID NO:17

t cacctatgac ctttacctgg

EXAMPLES

[0149] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in ° C. or is at room temperature, and pressure is at or near atmospheric.

Example 1

Preparation of Polyclonal Antibodies (IGG) to Glutamate and Homocysteine

[0150] Glutamate (polyglutamate, 10 aminoacids) or homocysteine (polyhomocysteine, 10 aminoacids) alone will not generate antibodies when injected into an animal. Therefore, polyglutamate and polyhomocysteine were conjugated with human serum albumin for the immunization to obtain polyclonal antibodies. For glutaraldehyde conjugation, polyglutamate or homocysteine (10 mg) and 40 mg bovine serum albumin (BSA, Sigma, St. Louis, Mo.) were incubated for 2 hr at room temperature in 4 ml of PBS containing 5% glutaraldehyde. The reaction was stopped by adding glycine to a final concentration of 0.2 M, and the conjugate was dialyzed against PBS.

[0151] Rabbits were given initial injections of 1 mg of conjugated glutamate (polyglutamate) or homocysteine

(polyhomocysteine) in complete Freund's and subsequent increased doses of injections (2 mg) in incomplete Freund's adjuvant at successive 2 week intervals. All injections were given subcutaneously. The immunization period lasted for 110 days. Antibodies (IgG) were affinity purified according to standard procedures (Warr, G. W., Purification of antibodies, In: Antibody as a Tool, Eds., Marchalonis, J. J., and G. W. Warr, J. Wiley, UK, pp. 59-96 (1982)) and were shown to be selective for glutamate or homocysteine by ELISA assay.

Example 2

Preparation of Polyclonal Antibodies (IGG) to NR2A Receptor Peptide

[0152] Using computer analysis of the hydrophobicity profile of human NR2A and NR2B NMDA receptors to predict the antigenic determinants in the protein structure, we selected fragments corresponding to the N-terminal sequence of human NR2A and NR2B receptor peptides for synthesis. The fragments corresponded to the N-terminal sequence of the NR2A and NR2B receptors, represented by SEQ ID NO: 1 and SEQ ID NO: 2 for the NR2A and NR2B receptors, respectively. The peptide fragments were reproduced using solid-phase synthesis, and had a purity ranging from 90% to 98%. The peptide sequences were verified by amino acid analysis after acid hydrolysis. A mixture of NR2A and NR2B peptides (1:1) was conjugated with human serum albumin for the immunization to obtain polyclonal antibodies. For glutaraldehyde conjugation, 10 mg of the mixture of peptides and 40 mg human serum albumin (Sigma, St. Louis, Mo.) were incubated for 1.45 hr at room temperature in 4 ml of PBS containing 5% glutaraldehyde. The reaction was stopped by adding glycine to a final concentration of 0.2 M, and the conjugate was dialyzed against PBS.

[0153] Rabbit polyclonal antibodies were raised against the NR2A-B peptides. Rabbits were given initial injections of 1 mg of conjugated peptides in complete Freund's adjuvant and subsequent injections (0.5 mg) in incomplete Freund's adjuvant at successive 2 week intervals. Antibodies were affinity purified according to standard procedures (Warr, G. W., Purification of antibodies, In: Antibody as a Tool, Eds., Marchalonis, J. J., and G. W. Warr, J. Wiley, UK, pp. 59-96 (1982)) and were shown to be selective for NR2A and NR2B NMDA receptors using an ELISA assay.

Example 3

Preparation of Latex Beads Containing Biomarker Antibodies

[0154] Three different sensitized latex beads containing IgG against glutamate, homocysteine and NR2A-B receptor peptides were prepared using two types of blue polystyrene latex beads (diameter, 0.25 and 0.4 μm; Sigma, St. Louis,

Mo.) as follows. A 1% suspension of latex beads in 50 mM PBS (1 ml, pH 7.0) was mixed with an equal volume of corresponding IgG (2 mg/ml) and incubated on a shaker at room temperature for 2 hours. The mixture was then washed twice with PBS by centrifugation at 9,500 g for 5 min. The pellet was suspended in PBS containing 1% BSA overnight at 4° C. After being washed twice with PBS, the sensitized latex beads were resuspended in latex diluent (50 mM PBS with 1% BSA) at a concentration of 0.4% and stored at 4° C. until used.

[0155] Preliminary experiments with latex agglutination (LA) alone were performed to identify problems and to select the most desirable latex particle size. Two types of commercial latex beads were coated with antibodies at various concentrations. Tests were initially performed with the corresponding aminoacid or NR2A and NR2B receptor peptides as controls. Particle size and IgG concentration were found to be the primary factors affect the sensitivity of the test. The most desirable particle size was found to be 0.25 μm (blue latex) because particles of this size agglutinated each aminoacid and peptide specifically. Higher IgG concentrations showed higher sensitivities. Using blue latex bead coated with 2 mg of IgG per ml, agglutination could be observed within 30 min.

Example 4

Latex Agglutination Analysis of Blood Serum Specimens

[0156] Blood samples (5 ml) were collected using standard venipuncture clinical protocol, from patients with TIA, stroke and brain injury (n=30) and examined at the laboratory of CIS Biotech, Inc. in Atlanta (Ga., USA). None of the patients had been treated with anticoagulants, and serum samples were obtained from the clotted blood. All specimens were free of visible lipids, white blood cells, platelets, fibrin, mucus or other contaminants that could cause "false positive" reactions. Platelets, white blood cells, mucus and fibrin were removed by centrifugation. Lipids were removed by filtration.

[0157] Specimens to be tested within 72 hours after collection were stored at 2-8° C. For longer storage periods, -20° C. or colder is recommended.

[0158] The semi-quantitative analysis of glutamate, homocysteine and NR2A-B receptor peptides in the serum samples is basically a three step process: serum sample dilution, reaction of latex beads with serum samples, and product analysis.

[0159] In previous experiments serial dilutions of the serum samples from 1:4 to 1:64 in saline containing 4% glycerol for better agglutination were performed. The highest dilution in which agglutination was observed corresponded to the sample titer.

[0160] Two 25 μl aliquots of coated latex beads containing the corresponding IgG were layered on a double-concave slide (Fisher Sci., Norcross, Ga.), one with 25 μl of the serum sample in serial dilution to be tested and one with 25 μl of PBS as a negative control. After gentle mixing with vortex, agglutination was judged macroscopically against a dark background. A negative reaction corresponded to a homogeneous lactescent background with no agglutination;

a positive reaction corresponded to a clearly visible agglutination against the black background and weakly visible agglutination on a slightly lactescent background.

[0161] The highest dilution at which agglutination occurs gives the titer of the sample. To obtain the approximate titer in $\mu\text{g/ml}$ we used the following calculation:

$$\text{Titer } \mu\text{g/ml} = A \times D$$

where A is the test sensitivity, and D is the highest dilution at which agglutination occurs.

Example 5

Description of Patients

[0162] Patients observed in trials (n=68) included 9 with pre-stroke, 9 with TIA (mean age 52.0 \pm 3.0), 31 with acute ischemic stroke (mean age 54.7 \pm 1.4) and 11 with mild brain injury (mean age 53.0 \pm 4.4). Clinical evaluation of patients by neuroimaging (CT, MRI, arteriography, Doppler ultrasonography, EEG), detailed physical and neurologic examination and laboratory tests was performed. Patients with TIA were characterized by contra lateral weakness, dysphasia, transient blurring of vision or blindness, abnormal pulsation of the common carotid arteries, microemboli confined to the ipsilateral retina. Untreated patients with pre-stroke demonstrated altered state of consciousness, severe headache, nausea and vomiting, visual disturbances, and focal neurological deficit, with some patients experiencing seizures.

[0163] The N-Score rating scale reported in "MCA Infarction" (Orgogozo, 1986) was used for evaluating the neurologic deficit in patients with acute cerebral stroke. The total score of acute cerebral stroke clinical manifestation differentiated severe patients (n=9, 11-35 scores) from patients with mild (n=12, 36-55 scores) and moderate patients (n=10, 60-90 scores). Most patients with acute cerebral ischaemia (61.3%) suffered ischemia in the carotid artery of left hemisphere. Arterial hypertension and cerebral atherosclerosis etiologically corresponded in all patients.

[0164] The patients with ischemia were divided into groups based on the differences between TIA, pre-stroke and acute ischemic pathogenic mechanisms. The clinical diagnosis was established on the basis of routine observations which included detailed neurological examination and neuroimaging. Groups of TIA (n=9) and pre-stroke patients with chronic cerebral blood insufficiency (n=9) were identified by neurophysiological investigations.

Example 6

Detection of Glutamate and Homocysteine in the Blood of Patients

[0165] Glutamate and homocysteine content were measured by standard high performance liquid chromatography (HPLC) according to methods described (Perry I. J., Refsum H., Morris R. W., Ebrahim S. B., Ueland P. M., Shaper A. G. Lancet. 1995, 346:1395-1398; Yamamoto T., Rossi S., Stiefel M., Doppenberg E., Zauner A., Bullock R., Marmarou A. Acta Neurochir. Suppl. 1999, 75: 17-19). The limits of the normal range were 165.0 $\mu\text{mol/L}$ for glutamate (Table 1) and 8.0 $\mu\text{mol/L}$ for homocysteine (Table 2). Elevated glutamate and homocysteine amounts were detected in the blood of patients with acute stroke. However, approximately

66% of these patients had additional risk factors indicative of atherosclerotic processes such as high cholesterol and LDL levels (Denisenko T. V., Skuliabin D., Gromov I., Cherkas Yi., Iluchina A., Dambinova S. A., 1998. *Vopr. Med. Khimii.* 44, 584-590, in Russian).

[0166] Abnormal glutamate and homocysteine plasma concentrations were observed more frequently in patients with TIA than in patients with acute stroke. The positive predictive efficiency of plasma glutamate for TIA patients was 56%. The positive predictive efficiency of plasma homocysteine for TIA patients was 66%. Baseline concentrations for glutamate and homocysteine are 160 $\mu\text{mol/L}$ and 10 $\mu\text{mol/L}$, respectively. Routine treatment for TIA was found to consistently decrease the glutamate and homocysteine levels in the blood of patients (data not shown).

[0167] In patients with pre-stroke, slightly elevated levels homocysteine were observed; levels of glutamate were unchanged (Tables 1, 2). In patients with traumatic brain injury (TBI), glutamate levels were observed that were nearly twice the glutamate levels in healthy individuals; levels of homocysteine were up to 57% higher.

TABLE 1

Glutamate concentration in the blood of patients detected by HPLC						
Group	Total N	Glutamate $\mu\text{mol/L}$	HPLC predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	165.0 \pm 28.2	19	67.8	9	32.2
TIA	9	200.0 \pm 11.7	4	44.4	5	55.6
Pre-stroke	9	163.7 \pm 10.4	5	55.6	4	44.4
Acute stroke	31	172.1 \pm 20.6	13	41.9	18	58.1
TBI	11	305.0 \pm 28.8	4	36.4	7	63.6

[0168] We also compared homocysteine concentrations in the blood of patients with TIA and pre-stroke to homocysteine concentrations in the blood of patients who have had stroke onsets. We observed that homocysteine content in the blood of patients depended on stage of the stroke, but that homocysteine concentration did not correlate with the severity of the cerebral ischemia. A significant decrease in homocysteine levels in patients with acute stroke was observed after emergency therapy (data not shown).

[0169] Latex agglutination was also employed to detect TIA/stroke biomarkers in the blood serum of patients. The titer of plasma glutamate determined by latex agglutination was 3.34 ± 0.25 in the group of healthy volunteers. Homocysteine and glutamate trends observed using HPLC were similarly observed for different groups of patients observed by using the LA technique (Tables 3, 4). Thus, increased levels of glutamate and homocysteine were similarly observed in the blood of patients with TIA and acute stroke using LA.

[0170] With respect to predictive efficiency, however, LA showed a surprising improvement over HPLC. For example, the LA method improved the positive predictive efficiency of patients with TIA and acute stroke on the basis of glutamate content to more than 63% (Tables 1, 3). The negative predictive value for healthy patients was similarly

improved when using the LA technique (Tables 3, 4). The predictive value of the LA technique in the group of patients with TBI was identical to the predictive value using HPLC.

TABLE 2

Homocysteine concentration in the blood of patients detected by HPLC						
Group	Total N	Homocysteine $\mu\text{mol/L}$	HPLC predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	8.0 \pm 1.7	20	71.4	8	28.6
TIA	9	10.8 \pm 1.3	3	33.3	6	66.4
Pre-stroke	9	9.0 \pm 1.2	4	44.4	5	55.6
Acute stroke	31	11.5 \pm 1.1	11	35.5	20	64.5
TBI	11	12.6 \pm 2.1	4	36.4	7	63.6

[0171]

TABLE 3

Detection of glutamate in the blood of patients by latex agglutination						
Group	Total N	Glutamate Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	3.34 \pm 0.25	22	78.6	6	21.4
TIA	9	4.52 \pm 0.38	3	33.3	6	66.4
Pre-stroke	9	3.57 \pm 0.32	4	44.4	5	55.6
Acute stroke	31	4.34 \pm 0.47	11	35.5	20	64.5
TBI	11	5.12 \pm 0.62	4	36.4	7	63.6

[0172]

TABLE 4

Detection of homocysteine the blood of patients by latex agglutination						
Group	Total N	Homocysteine Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	2.23 \pm 0.21	21	75.0	7	25.0
TIA	9	3.95 \pm 0.37	3	33.3	6	66.4
Pre-stroke	9	2.89 \pm 0.12	4	44.4	5	55.6
Acute stroke	31	4.01 \pm 0.41	10	32.3	21	67.7
TBI	11	4.74 \pm 0.38	4	36.4	7	63.6

Example 7

The Detection of NR2A-B in the Blood of Patients

[0173] The excessive activation and damage of NMDA receptors is the result of glutamate, aspartate and homocysteine neurotoxicity. Autoantibodies to have been detected in previous work in the blood of patients with TIA and pre-stroke, supporting our hypothesis that cerebral ischemia causes neuronal damage and the appearance of autoantibodies to NMDA receptor subunits (Gusev E. I., Skvortsova V. I., Alekseev A. A., Izykenova G. A., Dambinova S. A. S. S

Korsakov's J. Neurol. & Psych. 1996, 5:68-72; Dambinova S. A., Izykenova G. A. J. High Nervous Activity. 1997, 47: 439-446).

[0174] The titer of NR2A-B receptor peptides in the blood of healthy volunteers determined by LA was 2.63 ± 0.92 . Using the LA technique, we observed an increase in the test efficiency in the group of healthy persons up to 89% (Table 5). We also observed an improvement in the sensitivity of the LA test over ELISA. For example, patients with pre-stroke had slightly increased levels of NR2A-B receptor peptides over healthy volunteers when tested by ELISA, but had nearly double the level of NR2A-B receptor peptides when measured by LA (Table 5, 6). We detected high levels of NR2A-B receptor peptides using both ELISA and LA in the blood of patients with TIA and acute stroke, and observed comparable levels of predictive efficiency for each test.

[0175] Patients with TIA received routine treatment to improve brain circulation. Upon receiving treatment, NR2A-B levels decreased to levels corresponding to those observed for the healthy individuals as the patient's state normalized. As mentioned earlier, glutamate and homocysteine contents also decreased during treatment, but it never reached the levels observed in healthy individuals.

TABLE 5

Detection of NR2A-B receptor peptides in the blood of patients by latex agglutination						
Group	Total N	NR2A-B Titer	LA predictive value			
			Negative		Positive	
			N	%	N	%
Healthy individuals	28	2.63 ± 0.92	25	89.3	3	10.7
TIA	9	7.34 ± 0.43	2	22.2	7	77.8
Pre-stroke	9	4.21 ± 0.26	2	22.2	7	77.8
Acute stroke	31	5.20 ± 1.71	4	9.7	27	87.1
TBI	11	3.99 ± 0.44	2	18.8	9	81.8

[0176] Completely different profiles of NR2A-B were revealed in the blood of patients with acute ischemic stroke. In the blood of patients (n=8) with severe cerebral ischaemia (30.4 ± 3.2 Orgogozo scores) NR2A-B receptor peptides titer was 4 times higher than that for control group of healthy individuals. The peptides titer for patients with mild to moderate ischemic stroke (n=22, 49-62 Orgogozo scores) was slightly elevated in comparison with those with TIA. The tendency of slight decreases in NR2A-B receptor peptide levels was observed to the end of 30 days of patients' routine treatment, correlating with improvement in the neurological state.

TABLE 6

Group	Total N	Detection of NR2A-B receptor peptides in the blood of patients by ELISA				
		NR2A-B Ng/ml	ELISA assay results			
			Negative	Positive		
		N	%	N	%	
Healthy individuals	28	18.2 ± 2.1	20	71.4	8	28.6
TIA	9	66.6 ± 4.1	2	22.2	7	77.8
Pre-stroke	9	23.7 ± 1.9	3	33.3	6	66.7
Acute stroke	31	73.4 ± 6.5	5	16.1	26	83.9
TBI	11	54.3 ± 4.9	3	27.3	8	72.7

[0177] It is necessary notice that efficiency of both laboratory assays to detect the NR2A-B receptor peptides in the blood of patients with TIA/stroke and traumatic brain injury have been determined as 78 and 82% correspondingly.

[0178] The simultaneous detection of all brain damage biomarkers: glutamate, homocysteine and NR2A-B receptor peptides in the blood patients by latex agglutination allowed to diagnose rapidly TIA/stroke with efficiency up to 85-89%. The simultaneously increased levels of all biomarkers in the blood reflect the neurological deficit and may be used also for prognosis of diseases outcome. The relation between these biomarkers is showing the degree of thromboembolic and neurotoxicity involvement in brain processes underlying the ischemia. That fact is very important for choosing the strategy of emergency therapy in short time.

[0179] Using the latex agglutination technique allowed us significantly cut off the time of blood analysis from 3-8 hours in ELISA or HPLC to 30 min in LA. This RMP semi-quantitative test demonstrated the fast, simple for interpretation and reliable data.

Example 8

Identification of cDNA Sequence Encoding Antigenic Determinants of NMDA Receptors

[0180] It was necessary to first determine the cDNA sequence coding the immunological fragment of NMDA receptors responsible for the appearance of autoantibodies appearance. To find the most active peptide fragment of NMDA receptors a standard molecular biology procedure was used. Immunopositive phage GT11 containing cDNA coding NMDA receptors was isolated from a human cDNA library using autoantibodies to NMDA receptors isolated from blood samples of patients with severe cerebral ischemia or polyclonal antibodies to the NR2A receptor. An *E. Coli* bacterial system was employed to express the phage GT11 cDNA (600 bp). The expression product was transferred to a MBmp11 vector and a restriction map was constructed by use of a standard restrictases' kit. Three unique sites of the cDNA fragment (PstI, BamHI, and PsaI) were revealed, and the 5'-3' oligonucleotide sequence orientation using KpnII, BamHI and EcoR1 was deduced. The oligonucleotide (target cDNA) obtained was sequenced and compared to the sequence of the NR2A glutamate receptor (SEQ ID NO: 5) from the NCBI library. The target cDNA corresponded to the N-terminal domain of the NR2A recep-

tor (620 bp) of SEQ ID NO: 6, namely SEQ ID NO: 7. Primers for this target nucleotide were designed. All the oligonucleotides were prepared by the phosphoramidite method on an Applied Biosystem 394 synthesizer and were purified by reverse-phase high-pressure liquid chromatography (HPLC). The oligonucleotides used for detection and capture were synthesized with an amine arm at the 5' end.

Example 9

PCR Analysis of Blood Serum Specimens

[0181] Blood samples (5 ml collected by venipuncture) from patients with TIA and pre-stroke (n=30) were collected according to standard clinical protocol and examined at the Department of Neurology of Human Brain Institute, St. Petersburg Russia. The blood specimens were used for total DNA isolation or applied on FTA paper circles.

[0182] The quantitative analysis of NR2A cDNA expression in the serum samples is basically a three step process: Total DNA isolation and purification from sera of individuals; specific cDNA coding NR2A receptor amplification; and product analysis.

[0183] The total DNA isolated by DNAzol (Mol. Res. Center, Inc., Cincinnati, Ohio) or bound on FTA blood staining collection cards (Life Technologies, Inc., Gaithersburg, Md.) serves as a template for the polymerase chain reaction (PCR). In the first variant, the PCR assay uses a set of specially designed primers (50 pmol), immobilized on solid matrix of microplates and amplifies a specific cDNA sequence (620 bp) coding the NR2A glutamate receptor. In a second variant, the PCR assay uses a master ready-to-use buffer and amplifies cDNA bound on FTA paper. Following amplification, the quantity of a product is determined by enzyme or non-enzyme color reaction with a substrate.

[0184] Using the DNAzol reagent for DNA isolation, the whole blood of each individual (0.5 ml) was combined with 1 ml DNAzol (Mol. Res. Center, Inc., Cincinnati, Ohio) for 5 min at room temperature and lysed (Mackey K. et al. Mol. Biotechnol. 9: 1-5 (1997)). The organic phase (0.4 ml) of each sample was transferred to a clean tube and 0.4 ml isopropanol was added. The mixture was incubated for 5 min at room temperature and centrifugated at 6,000 g for 6 minutes. The pellet was washed in 0.5 ml DNAzol and centrifugated at the same conditions. The total DNA pellet was mixed with 1 ml of 75% ethanol and centrifuged at 6,000 g for 5 minutes. Then the DNA pellet was diluted in 200 μ l of 8 mM NaOH and incubated at room temperature for 5 min followed by vortexing. Alkaline DNA solution was then neutralized with 160 μ l of 0.1 M HEPES, pH 7.4.

[0185] Immobilization of oligonucleotide probes (primers, SEQ ID NO:8) was performed as follows. A total of 100 μ l of 3 \times PBS buffer containing the primers (150 nM) was dropped into each well of a 96-well microtiter plate (Fisher Sci., Suwanee, Ga.). After incubation for 2 h at 37 $^{\circ}$ C. or overnight at room temperature, the plate was washed three times with 1 \times PBS buffer containing 0.05% (w/vol) Tween 20. The oligonucleotide-coated plates were stable for 2 months at 4 $^{\circ}$ C.

[0186] Direct PCR reactions were performed in a final volume of 50 μ l (Sisk R B. in book: Molecular diagnostics: for the clinical laboratorian. Ed. by Coleman W B., and

Tsongalis G J. Humana Press Inc., Totowa, N.J. 1997, pp. 103-121). The total DNA (5 μ l), isolated from blood samples of individuals, to oligonucleotide-coated plate in duplicates and 45 μ l of master ready-to-use buffer containing 1 μ l TaKaRa Z-Taq DNA polymerase (TaKaRa Biomedicals, Otsu, Shiga, Japan) 10 μ l AMV/Tfl 5 \times reaction buffer, 1 μ l dNTP mix (Promega, Madison, Wis.) 2 μ l of 25 mM MgSO₄ were added and sealed. The 30-thermal cycles (98 $^{\circ}$ C.-5 s, 66 $^{\circ}$ C.-2 sec) amplification using programmable Gene Cyclyer thermocycler (Bio-Rad Lab., Hempstead, UK) for 20 minutes was performed. Then 50 μ l of PicoGreen reagent (Mol. Probes, Inc., Eugene, Oreg.) were added to each PCR products and mixed on a shaker (BioTechniques 20:676 (1996)). Samples were incubated 5 min at room temperature, protected from light. After incubation the fluorescence of the samples was measured using a fluorescence microplate reader (Mol. Device, Sunnyvale, Calif.) and standard fluorescein wavelengths (excitation 480 nm, emission 520 nm). The fluorescence value of the reagent blank was subtracted from that of each of the samples, and the data was employed to generate five-point standardization curves of fluorescence versus DNA concentration, from 25 pg/ml to 25 ng/ml reaction of control target cDNA (50 ng/ml stock) with the same Pico Green reagent.

[0187] The other method of total DNA isolation is follows. Whole blood was spotted onto FTA paper and lysed, and samples of DNA immobilized within the matrix of the stain card were punched into a 3 mm (1/8") diameter paper (1 mm or 2 mm Harris Micro-PunchTM) and amplified directly by the amplification mix (Mackey K. et al. Mol. Biotechnol. 9: 1-5 (1997)).

[0188] The FTA Bloodstain Card is divided into 4 circles for at least 4 different 120 μ l samples of EDTA collected whole blood. Samples of blood were dried at room temperature for at least 1 hour. A circle was drawn with a #2 pencil around each blood to visualize where the blood had been spotted after the FTA paper processing. The FTA Bloodstain Card was then placed in a small plastic tray and 50 μ l of FTA Purification Reagent was added and incubated on a shaker for 5 minutes. FTA Purification Reagent was replace 3 times with 25-50 μ l of the fresh solution and shaken for an additional 5 minutes. Then 25-50 μ l of TE-4 (10 mM Tris-HCl pH 8.0; 0.1 mM EDTA pH 8.0) was added and the mixture incubated twice on a shaker for 5 minutes. The FTA Bloodstain Card was allowed to air dry completely during 2 hours at room temperature. The samples were then punched from the cards using a 3 mm diameter punch or the Harris Micro-Punch (1.2 mm or 2.0 mm), and transferred into corresponding microplate wells. PCR was then performed using the above-described procedure using regular PCR microplates and a ready-to-use buffer containing primers.

[0189] Patients (n=30, the age of 44-77) were divided into two groups. The first group of patients (n=12) were diagnosed with TIA in the carotid circulatory system, according to the following neurological criteria. Neural dysfunction was localized to a specific vascular distribution; the duration of the attack was usually less than 15 minutes and never exceeded 24 hours; and the patients did not have abnormal neurologic signs between attacks. The second pre-stroke group (n=18) were diagnosed with TIA in the vertebral-basilar. circulatory system. The second group of patients was subdivided on the basis of compensation or non-compensa-

tion of neurological deficit. The third group (n=12) included patients with migraine and epilepsy.

[0190] The control group of healthy individuals (n=20) showed a level of NR2A cDNA expression of 1.2 0.11 pg/ml. The first group demonstrated slightly elevated levels of NR2A cDNA expression of 1.7 0.13 pg/ml. The patients with compensation of neurological deficit from the second group showed a level of NR2A cDNA expression of 1.8 1.4 pg/ml. At the same time, the patients without compensation of neurological deficit that possessed more severe symptoms of TIA showed levels of NR2A cDNA expression of 3 times the levels seen in healthy individuals. Patients suffering migraine and epilepsy did not show any increase of NR2A cDNA expression when compared with the control group.

Example 10

Immunological Analysis of Blood Serum Specimens

[0191] Blood samples (10 ml, collected by venipuncture) from patients with cerebral ischemia (n=70), and healthy individuals (n=200), collected according to standard clinical protocols, were examined at the Neurology Hospital of Russian Medical Academy (Moscow, Russia). The blood specimens were centrifugated (4000 g, 5 min, +4° C.) and the collected serum stored at -70° C. for further analysis.

[0192] Computer analysis was employed to predict the antigenic determinants in the NR2A receptor protein structure based on hydrophobicity profile (Hopp, T. P. and K. R. Woods, *Proc. Natl. Acad. Sci. USA* 6:3824-3828 (1981)) and antigenicity (Welling, G. W., et al., *FEBS Lett.* 188:215-218 (1985)). Based upon this analysis, the N-terminal sequence of the NR2A NMDA receptor was synthesized. This synthetic peptide, which corresponded to amino acid sequence (494-514) (Grandy, D K., et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:9762-9766(1989) (SEQ ID NO:3) of human NR2A NMDA, was produced by solid-phase synthesis in a NPS-400 semi-automated synthesizer (Neosystem Lab, France) on MBHA resin using the BOC/Bzl strategy for the first two amino acids. The peptides were purified by preparative HPLC on a DELTAPAC™ C18 column (Waters Chromatography, Milford, Mass.) in a H₂O/acetonitrile/0.015 TFA system. The purity of the peptides was determined by analytical BPLC and ranged from 90% to 98%. The peptide sequence was verified by amino acid analysis after acid hydrolysis. This peptide was used in immunoassays of blood serum from patients and healthy individuals.

[0193] A quantitative analysis of the level of NR2A autoantibodies in serum samples was performed by enzyme-linked immunosorbent assay (ELISA) (Ngo, T. T. and H. M. Lenhoff, *FEBS Lett.* 116:285-288 (1980)). The diluted blood sera (1:50) and polyclonal antibodies to the NR2A peptide as a standard (0.01 ng/ml-400 ng/ml) were applied to the immunosorbent. The plate was incubated for 1 h at 25° C. and then washed by 0.05 M phosphate buffer, pH 7.4, containing 0.05% of TWEEN-20™. Rabbit antibodies to the human immunoglobulin labeled with horseradish peroxidase were added (Sigma, St. Louis, Mo.; 1:1000), and the plate was incubated for 1 h at 25° C. After incubation the wells were washed twice in the same buffer. The reaction was revealed by o-phenylenediamine in 0.05 M citrate buffer, pH 4.3 monitored at 490 nm on a microplate reader (BioRad,

UK). The titer of NR2A autoantibodies in blood serum was determined by ELISA using a standard curve of the absorbance units of NR2A autoantibodies versus their concentration in a microtiter well plate.

[0194] The synthetic peptide corresponding to the NR2A NMDA glutamate receptors (3 μg) were immobilized on a nitrocellulose membrane (0.45 μm, Schleicher-Shuell, Germany) in phosphate-buffered saline (PBS), pH 7.4, then washed 2-3 times in the same buffer. Membranes with immobilized peptide were incubated with the diluted serum (1:50) of cerebral ischemia patients and other subjects for 1 h at 25° C., and then rinsed 4 times with the PBS buffer. Secondary rabbit anti-human immunoglobulins conjugated with horseradish peroxidase (Sigma, St. Louis, Mo.; 1:1000) were incubated with the membrane for 1 h at 25° C., then washed 4 times with PBS. The development of brown color was registered and then quantitated by densitometry.

[0195] To provide a positive control or standard, rabbit polyclonal antibodies were raised against NR2A synthetic peptide corresponding to amino acid sequence predicted from the cloned human NR2A protein (Science 256:1217-1221 (1992); SEQ ID NO:1). For glutaraldehyde conjugation, 10 mg of peptide and 40 mg of human serum albumin (Sigma, St. Louis, Mo.) were incubated for 1.45 h at room temperature in 4 ml of PBS containing 5% glutaraldehyde. The reaction was stopped by adding glycine to a final concentration of 0.2 M, and the conjugate was dialyzed against PBS. Rabbits were given initial injections of 1 mg of conjugated peptide in complete Freund's and subsequent injections of 0.5 mg of peptide in incomplete Freund's adjuvant at successive 2 week intervals. Antibodies were affinity purified according standard procedure (Warr, G. W., Purification of antibodies, In: *Antibody as a Tool*, Eds., Marchalonis, J. J., and G. W. Warr, J. Wiley, UK, pp. 59-96 (1982)) and were shown to be selective for the NR2A NMDA glutamate receptor using Western blot analysis.

[0196] The patients (men, n=30; women, n=40; age of 40-75) were admitted in the hospital within no more than six hours after the onset of an ischemic episode. All patients were divided into three groups according to the severity of the stroke: The first group had moderate ischemic stroke (n=25), manifested by moderate focal deficit (>60-Orgogozo scale). The second group had severe stroke (n=30), manifested by mild disorders of consciousness, severe headache, meningeal sings, and pronounced focal deficit (30-60-Orgogozo scale). The third group had extremely severe stroke (n=15), accompanied by stupor-coma, signs of brain edema, autonomic dysfunction, and severe focal deficit (<30-Orgogozo scale).

[0197] The level of NR2A autoantibodies was measured in the blood serum of healthy persons (n=200, age 35-75) as a control, and ranged from 0.3-1.5 ng/ml. The NR2A antibody level in the 55 patients of the first and second groups was significantly greater than that in the control group (p<0.01). Levels of NR2A autoantibodies were monitored every three hours during the first day, and then up to 5th day after stroke. The level of NR2A autoantibodies in the blood serum of patients with severe stroke was significantly higher than that in the blood serum of patients with moderate stroke, especially in the 9-12 hours after the onset of a stroke (p<0.05). The tendency for NR2A autoantibodies level to decrease to the control level on the first day of stroke was

registered in group of patients with good neurological recovery (90,50,5 units on Orgogozo scale). It can be concluded that the dynamic changes in NR2A autoantibodies level may predict a recovery period of patients after ischemic stroke.

Example 11

SPRIA Assay of Autoantibodies

[0198] The solid-phase radioimmunoassay (SPRIA) of autoantibodies is performed as follows: a 10% acetic acid solution is added for one minute to the Cooker microtiter microplates (available from Dynatech Co., USA) for activation, whereupon 0.1 ml of the blood serum under analysis (diluted 1:40) is applied to the microplates and subjected to incubation for four hours at 25° C. Then the microplate are washed with a 0.14 M sodium chloride solution and 0.1 ml of a mixture of the respective fragment of the mammal's brain protein labeled by 125I in the presence of nonlabelled one. The plates are incubated for 20 hours at 4° C. On completion of incubation, the microplates are washed with a 0.14M sodium chloride solution, after which each of the wells of the microplates is cut off and placed in gamma-counting vials.

Example 12

Elisa Assay of Autoantibodies

[0199] The enzyme-linked immunosorbent assay (ELISA) of autoantibodies is carried out as follows: the samples of the blood serum diluted 1:40 or 1:50 are applied to the respective immunosorbent. Then the plate carrying the immunosorbent is incubated for 30 min at 37° C., whereupon the wells of the plate are washed with a 0.05 M phosphate buffer, containing 0.05% of Tween-20. Rabbit antibodies to human immunoglobulin labeled with horseradish peroxidase (conjugate) are added thereto, and the plate is reincubated for 35 min at 37° C., then washed by the aforementioned buffer and distilled water. The reaction with conjugate is determined by adding chromogen, i.e., orthophenylenediamine in the presence of 30% hydrogen peroxide. The intensity of color development is evaluated by using the rider (available Multiskan microplate rider) at the 492 nm wavelength.

[0200] It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

What is claimed is:

1) A method for diagnosis of a stroke in a patient comprising directly or indirectly measuring, within three hours after stroke onset the level of NR2A and/or NR2B NMDA receptor or fragment thereof in a subject.

2) The method of claim 1 further comprising comparing the level of NR2A and/or NR2B NMDA receptor or fragment thereof in the biological sample to a baseline level selected from population norms and prior levels measured for the subject.

3) The method of claim 1 wherein the the level of NR2A and/or NR2B NMDA receptor or fragment thereof is measured directly from the level of the N-terminal domain of a NR2A and/or NR2B NMDA receptor or fragment thereof in the biological sample.

4) The method of claim 1 wherein the level of NR2A and/or NR2B NMDA receptor or fragment thereof is measured indirectly from the level of antibodies against the N-terminal domain of a NR2A and/or NR2B NMDA receptor, analogue thereof, or fragment thereof in the biological sample.

5) The method of claim 1 wherein the level of NR2A and/or NR2B NMDA receptor or fragment thereof is measured indirectly from the mRNA encoding the NR2A and/or NR2B NMDA receptor or fragment thereof in the biological sample.

6) The method of claim 1 wherein the the level of NR2A and/or NR2B NMDA receptor or fragment thereof is measured directly from the level of the peptide represented by SEQ ID NO:2, 3, 11, and/or 12.

7) The method of claim 1 wherein the the level of NR2A and/or NR2B NMDA receptor or fragment thereof is measured indirectly from the level of antibodies against a peptide of SEQ ID NO:2, 3, 11, and/or 12, analogue thereof, or fragment thereof, in the biological sample.

8) The method of claim 1 wherein the the level of NR2A and/or NR2B NMDA receptor or fragment thereof is measured indirectly from the level of cDNA corresponding to SEQ ID NOS: 5, 6, 7, 14, and/or 15, analogue thereof, or fragment thereof, in the biological sample.

9) The method of claim 1 for diagnosing the existence of TIA or stroke, further comprising withdrawing the biological sample from a human subject, wherein the biological sample is withdrawn within three hours of the onset of symptoms of TIA or stroke.

10) The method of claim 9 for diagnosing the existence of TIA or stroke, wherein the amount of time elapsed between withdrawing the biological sample from the subject, and detecting or measuring the presence or quantity of the NR2A and/or NR2B NMDA receptor or fragment thereof, is less than about one hour.

11) The method of claim 1 for diagnosing the existence of TIA or stroke further comprising, when the diagnosis confirms a stroke, evaluating from the level of NR2A and/or NR2B NMDA receptor or fragment thereof whether the stroke is ischemic or hemorrhagic and administering ischemic or hemorrhagic stroke therapy as appropriate.

12) The method of claim 1 for diagnosing the existence of TIA or stroke further comprising, if TIA and/or stroke is confirmed, evaluating from the level of NR2A and/or NR2B NMDA receptor or fragment thereof whether the episode is TIA or stroke and administering TIA or stroke therapy as appropriate.

13) The method of claim 1 for diagnosing the existence of TIA or stroke further comprising, if TIA and/or stroke is confirmed, evaluating from the level of NR2A and/or NR2B NMDA receptor or fragment thereof cranial infarct volume, and administering therapy consistent with the extent of cranial infarct.

14) The method of claim 1 wherein the NR2A and/or NR2B NMDA receptor or fragment thereof is measured by immunoassay.

15) The method of claim 1 wherein the NR2A and/or NR2B NMDA receptor or fragment thereof is detected or measured by agglutination comprising:

- a) contacting the biological sample with poly- or monoclonal antibodies bound on an agglutinating carrier for sufficient time and under conditions to promote agglutination, wherein the antibodies are specific for the NR2A and/or NR2B NMDA receptor, analogue thereof, or fragment thereof; and
- b) reading a signal generated from the agglutination; wherein the signal correlates to the titer of NR2A and/or NR2B NMDA receptor or fragment thereof present in the sample.

16) The method of claim 15, wherein the sufficient time period is 30 minutes or less.

17) The method of claim 15 wherein the carrier comprises polystyrene latex beads having a mean diameter of from about 0.25 to about 0.4 μm .

18) The method of claim 1, wherein the biological sample is blood, urine, blood plasma, blood serum, cerebrospinal fluid, saliva, perspiration or brain tissue, or a derivative thereof.

19) The method of claim 1, wherein the biological sample is blood diluted to a ratio of from about 1:2 to about 1:32.

20) The method of claim 1, wherein levels of cDNA are measured by:

- a) complexing the biological sample with one or more oligonucleotide primers to the cDNA for a sufficient time period and under conditions to promote amplification;
- b) contacting the complex with an indicator reagent comprising a secondary oligonucleotide complementary to the cDNA attached to a signal-generating compound; and
- c) measuring the signal.

21) The method of claim 1 wherein levels of NR2A and/or NR2B autoantibody are measured comprising:

- a) contacting the biological sample with a protein fragment of the N-terminal domain of the NR2A and/or NR2B receptor for a time sufficient and under conditions to form a complex between autoantibodies in the sample and the protein fragment;
- b) contacting the complex with an indicator reagent comprising a secondary antibody attached to a signal generating compound; and
- c) measuring the signal.

* * * * *

专利名称(译)	在TIA /中风的实验室血液测试中快速多组生物标记物		
公开(公告)号	US20060024749A1	公开(公告)日	2006-02-02
申请号	US11/076074	申请日	2005-03-09
[标]申请(专利权)人(译)	DAMBINOVA SVETLANA一个		
申请(专利权)人(译)	DAMBINOVA SVETLANA一个		
当前申请(专利权)人(译)	CIS BIOTECH , INC.		
[标]发明人	DAMBINOVA SVETLANA A		
发明人	DAMBINOVA, SVETLANA A.		
IPC分类号	C12Q1/68 G01N33/53 G01N33/68 G01N33/94		
CPC分类号	G01N33/6893 G01N33/9406 G01N2800/52 G01N2800/2871 G01N2333/70571		
优先权	60/301297 2001-06-27 US		
其他公开文献	US7622100		
外部链接	Espacenet USPTO		

摘要(译)

一种用于诊断中枢神经系统疾病，特别是短暂性脑缺血发作或中风的方法，试剂盒和组合物，包括测量来自人受试者的生物样品中的NR2A和/或NR2B NMDA受体或其片段的水平，并任选地测量其他生物标志物如同型半胱氨酸和谷氨酸。该方法对于识别有中风风险的个体以及在急诊室环境中诊断中风特别有用。

TABLE 6

Detection of NR2A-B receptor peptides in the blood of patients by ELISA

Group	Total N	Ng/ml	ELISA assay results			
			NR2A-B			
			Negative	Positive	N	%
Healthy individuals	28	18.2 ± 2.1	20	71.4	8	28.6
TIA	9	66.6 ± 4.1	2	22.2	7	77.8
Pre-stroke	9	23.7 ± 1.9	3	33.3	6	66.7
Acute stroke	31	73.4 ± 6.5	5	16.1	26	83.9
TBI	11	54.3 ± 4.9	3	27.3	8	72.7