



US 20130089877A1


(19) **United States**(12) **Patent Application Publication**
Kim et al.(10) **Pub. No.: US 2013/0089877 A1**(43) **Pub. Date: Apr. 11, 2013**(54) **THE NOVEL ABETA22(PE)-42 PEPTIDE THAT BINDS TO AMYLOID-BETA ANTIBODY SPECIFICALLY IN BLOOD AND PHARMACEUTICAL COMPOSITION FOR THE DIAGNOSIS OF DEMENTIA DISEASE CONTAINING THE SAME AS AN ACTIVE INGREDIENT**(71) Applicant: **Korea Center For Disease Control And Prevention, Cheongwon-gun (KR)**(72) Inventors: **Young-Youl Kim, Chungcheongbuk-do (KR); Seungwoo Kim, Chungcheongbuk-do (KR); Hyo-Soon Cheon, Chungcheongnam-do (KR); Sang Ick Park, Daejeon (KR)**(73) Assignee: **KOREA CENTER FOR DISEASE CONTROL AND PREVENTION, Cheongwon-gun (KR)**(21) Appl. No.: **13/645,741**(22) Filed: **Oct. 5, 2012**(30) **Foreign Application Priority Data**

Oct. 7, 2011 (KR) KR10-2011-0102611

Publication Classification(51) **Int. Cl.**
G01N 33/53 (2006.01)
G01N 21/76 (2006.01)(52) **U.S. Cl.**
USPC **435/7.92; 436/501; 435/7.9; 435/7.1**(57) **ABSTRACT**


The present inventors screened peptides having a specific sequence specifically binding to amyloid-beta antibody and accordingly confirmed that A β 22(pE)-42 peptide showed higher reactivity to amyloid-beta antibody in serum of Alzheimer's disease patients. Therefore, the said A β 22(pE)-42 peptide can be used as an active ingredient for the kit for diagnosing dementia and thus it can be said that the peptide can be effectively used for the diagnosis of dementia whose early diagnosis is hardly possible.

[Fig. 1]



[Fig. 2]

A β 22(pE)-42



**THE NOVEL ABETA22(PE)-42 PEPTIDE THAT
BINDS TO AMYLOID-BETA ANTIBODY
SPECIFICALLY IN BLOOD AND
PHARMACEUTICAL COMPOSITION FOR
THE DIAGNOSIS OF DEMENTIA DISEASE
CONTAINING THE SAME AS AN ACTIVE
INGREDIENT**

[0001] The novel A β 22(pE)-42 peptide that binds to amyloid-beta antibody specifically in blood and pharmaceutical composition for the diagnosis of dementia disease containing the same as an active ingredient

RELATED APPLICATIONS

[0002] This application claims priority to Korean Patent Application No. 1020110102611, filed on Oct. 7, 2011, the entire content of which is hereby incorporated herein in its entirety.

BACKGROUND OF THE INVENTION

[0003] 1. Field of the Invention

[0004] The present invention relates to a kit for diagnosis of dementia comprising the novel A β 22(pE)-42 peptide binding specifically to amyloid-beta antibody in blood as an active ingredient

[0005] 2. Description of the Related Art

[0006] Population aging in Korea and in most of other countries increases interest in senile diseases and accordingly studies have been actively undergoing about senile diseases. Alzheimer's disease is a kind of degenerative diseases and one of the most representative dementia related diseases, which puts heavy mental and physical burden on the shoulder of not only patient himself but also his family. Alzheimer's disease is found in 10% of people at the age of 65~74, 19% of people at the age of 75~84, and 47% of people at the age of 85 and up. Incidence rate of the disease rises every year, which becomes a serious social problem.

[0007] None of the prevention method for Alzheimer's disease or the method for early diagnosis has been known so far. Diagnosis of the disease has depended on doctor's clinical opinion and neuropsychological test, indicating only after the disease progresses far, diagnosis can be made, which makes the treatment difficult. Any effective therapeutic agent has not been developed, yet. Compositions for alleviating symptoms are just used.

[0008] The most acceptable theory to understand the mechanism of Alzheimer's disease now is "amyloid hypothesis". That is, amyloid-beta existed in patients or plaque generated by amyloid-beta accumulation induces continuous apoptosis of neurons, resulting in the suppression of neuronal transmission and damages in recognition, leading to the outbreak of Alzheimer's disease. Senile plaque and neurofibrillary tangles are observed in the brains of expired patients due to Alzheimer's disease, which are major pathological characteristics of Alzheimer's disease. Particularly, senile plaque is formed by the accumulation of protein and dead cells in outer cell whose major component is a peptide called amyloid-beta (A β) (Hardy, J. et al, Nat Neurosci. 1:355-358, 1998). Gradual damage in recognition, which is the major characteristics of Alzheimer's disease, is believed to be caused by abnormal accumulation of amyloid-beta that is produced by proteolysis of amyloid precursor protein (referred as "APP" hereinafter). The precursor APP is decomposed by β -secretase (BSCE) and γ -secretase to generate amyloid-beta (Cra-

ven, R., Nat Rev. Neurosci. 2: 533, 2001; David, H. S. et al., Nat Rev. Neurosci. 2: 595-598, 2001; Yankner, B. A., Neuron 16: 921-932, 1996; Selkoe, D. J., Nature 399: A23-A31, 1999). Development of diagnosis indexes for Alzheimer's disease from blood is actively undergoing. Recently, amyloid protein (amyloid-beta, A β 42, and A β 40) known as a key player in Alzheimer's disease development is used as standard for the diagnosis of Alzheimer's disease.

[0009] In animal models having Alzheimer's disease, the levels of A β 42 and A β 40 in serum and cerebrospinal fluid are increased in proportion to aging. Once cognitive impairment starts, the levels of those proteins are decreased. Thus, taking the amyloid protein level in serum as a standard index for the diagnosis of Alzheimer's disease is under controversy. It has been known that amyloid antibody is observed in the brain, blood, and cerebrospinal fluid of both normal healthy people and Alzheimer's disease patients, but the amount or characteristics of amyloid antibody have not been explained, yet.

[0010] The method for early diagnosis of Alzheimer's disease and the prevention method thereof have recently been focused on measuring amyloid-beta protein in body by using amyloid-beta antibody. In this method, it is very important to identify amyloid-beta antigen having a specific arrangement binding specifically to the antibody growing specifically in the blood of Alzheimer's disease patients, unlike in normal people.

[0011] Therefore, the present inventors screened peptides having specific patterns that bind specifically to amyloid-beta antibody and then confirmed that A β 22(pE)-42 peptide had comparatively high reactivity against amyloid-beta in serum of Alzheimer's disease patients, compared with in normal people. Accordingly, the present inventors completed this invention by confirming that the said A β 22(pE)-42 peptide can be used as an active ingredient for the diagnostic kit of dementia and for the diagnosis of dementia whose early diagnosis is very difficult so far.

SUMMARY OF THE INVENTION

[0012] It is an object of the present invention to provide a method for diagnosis of dementia using the novel A β 22(pE)-42 peptide binding specifically to amyloid-beta antibody in blood.

[0013] To achieve the above object, the present invention provides a kit for diagnosis of dementia comprising the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 as an active ingredient.

[0014] The present invention also provides a method for diagnosis of dementia comprising the following steps:

[0015] 1) treating blood taken from a test subject on the solid board attached with the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1, followed by washing thereof; and

[0016] 2) diagnosing the subject with dementia or high risk of dementia when the binding intensity of amyloid-beta antibody on the solid board is significantly higher than that of normal people.

[0017] The present invention further provides a method for diagnosis of dementia comprising the following steps:

[0018] 1) treating blood taken from a test subject on the solid board, followed by washing thereof;

[0019] 2) treating the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board, followed by washing thereof; and

[0020] 3) diagnosing the subject with dementia or high risk of dementia when the binding intensity of the A β 22(pE)-42

peptide represented by SEQ. ID. NO: 1 on the solid board is significantly higher than that of normal people.

ADVANTAGEOUS EFFECT

[0021] As explained hereinbefore, the present inventors screened peptides having a specific sequence specifically binding to amyloid-beta antibody and accordingly confirmed that A β 22(pE)-42 peptide showed higher reactivity to amyloid-beta antibody in serum of mild cognitive impairment patient. Therefore, the said A β 22(pE)-42 peptide can be effectively used as an active ingredient for the kit for diagnosis of dementia whose early diagnosis has been hardly made.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] The application of the preferred embodiments of the present invention is best understood with reference to the accompanying drawings, wherein:

[0023] FIG. 1 is a diagram illustrating the immune reactivity of A β 22(pE)-42 peptide to amyloid-beta antibody:

[0024] Normal: normal people;

[0025] MCI: mild cognitive impairment;

[0026] sevAD: severe Alzheimer's disease;

[0027] proAD: probable Alzheimer's disease; and

[0028] Negative con: negative control.

[0029] FIG. 2 is a graph illustrating the result of quantitative analysis with the said FIG. 1.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0030] Hereinafter, the present invention is described in detail.

[0031] The present invention provides a kit for diagnosis of dementia comprising the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 as an active ingredient.

[0032] The said peptide can be the one binding to amyloid-beta antibody, but not always limited thereto.

[0033] The said dementia can be selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease, and cerebral amyloid angiopathy, but not always limited thereto.

[0034] Dementia is defined by declined intellectual functions as severe as causing problems in social life or daily life resulted from decrease of cognitive functions in a variety of fields including memory, language, judgment, etc., which is acquired by structural lesion in the brain. For example, mild cognitive impairment, Alzheimer's disease, and cerebral amyloid angiopathy are included in the criteria of dementia. The generation and accumulation of amyloid might be a reason of the said disease, suggesting that the peptide of the present invention can be applied for diagnosis of such disease.

[0035] The peptide herein can be conjugated with chromogenic enzyme, chromogenic material or fluorescent molecule. The chromogenic enzyme herein is HRP (horseradish peroxidase) or alkaline phosphatase. The chromogenic material can be colloid gold, and the fluorescent molecule in this invention can be selected from the group consisting of poly L-lysine-fluorescein isothiocyanate (FITC), rhodamine-B-isothiocyanate (RITC), Cy2, Cy3, and Cy5, but not always limited thereto.

[0036] In particular, when secondary antibody-signal molecule complex is used as a detecting tool, the signal molecule can be selected from the group consisting of fluorescent molecule (ex; Cy3, CY5, FITC, GFP (green fluorescent protein), RFP (red fluorescent protein), and Texas Red), luminescent

material, radio isotope, and such enzymes as HRP (horse radish peroxidase), alkaline phosphatase, β -galactosidase, and luciferase, etc, but not always limited thereto.

[0037] The kit for diagnosis of dementia of the present invention makes the diagnosis of dementia possible by analyzing quantitatively or qualitatively the binding reactions such as antigen/antibody binding or protein/ligand binding. The said binding reaction herein can be measured by ELISA, radioimmunoassay (RIA), sandwich assay, western blotting, immunoprecipitation, immunohistochemical staining, immunofluorescence method, enzyme-substrate colorimetry, and antigen/antibody agglutination assay.

[0038] The kit is characteristically composed of:

[0039] 1) reactor coated with the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1;

[0040] 2) secondary antibody conjugate conjugated with chromogenic marker whose color is developed by the reaction with substrate;

[0041] 3) chromogenic substrate solution for the color development reaction with the marker;

[0042] 4) washing solution for each reaction step; and

[0043] 5) stop solution to terminate the enzyme reaction.

[0044] The secondary antibody conjugate marker can be selected from the group consisting of HRP (horseradish peroxidase), alkaline phosphatase, colloid gold, fluorescent molecule, and dye, but not always limited thereto.

[0045] The chromogenic substrate can be selected from the group consisting of TMB (3,3',5,5'-tetramethyl bezidine), ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)], and OPD (o-phenylenediamine), but not always limited thereto.

[0046] To investigate immune reactivity of A β 22(pE)-42 peptide to amyloid-beta antibody, the present inventors separated antibodies from plasma of normal people (48 people), mild cognitive impairment (MCI) patients (49), probable Alzheimer's disease patients (11), and severe Alzheimer's disease patients (62).

[0047] Immuno reactivity was measured by using protein chip. As a result, reactivity of A β 22(pE)-42 peptide to amyloid-beta antibody was increased in blood obtained from mild cognitive impairment and Alzheimer's disease patients, compared with that in normal people (see FIG. 1). Particularly, reactivity of A β 22(pE)-42 peptide was specifically increased in mild cognitive impairment patients (143%, $p < 0.01$), probable Alzheimer's disease patients (146%, $p < 0.01$), and severe Alzheimer's disease patients (148%, $p < 0.001$), compared with in normal people (see FIG. 2).

[0048] As explained hereinbefore, it was confirmed that immune reactivity of A β 22(pE)-42 peptide to amyloid-beta antibody was higher in serum of mild cognitive impairment patients. Therefore, the said A β 22(pE)-42 peptide can be effectively used as an active ingredient for the kit for diagnosis of dementia, in other words it can be effectively used for the diagnosis of dementia whose early diagnosis has been hardly made.

[0049] The present invention also provides a method for diagnosis of dementia using A β 22(pE)-42 peptide, which comprises the following steps:

[0050] 1) treating blood taken from a test subject on the solid board attached with the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1, followed by washing thereof; and

[0051] 2) diagnosing the subject with dementia or high risk of dementia when the binding intensity of amyloid-beta antibody on the solid board is significantly higher than that of normal people.

[0052] The present invention further provides a method for diagnosis of dementia using A β 22(pE)-42 peptide, which comprises the following steps:

[0053] 1) treating blood taken from a test subject on the solid board, followed by washing thereof;

[0054] 2) treating the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board, followed by washing thereof; and

[0055] 3) diagnosing the subject with dementia or high risk of dementia when the binding intensity of the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board is significantly higher than that of normal people.

[0056] In the above method, the subject of step 1) is a vertebrate including human, and preferably a mammal particularly exemplified by human, apes, cow, pig, rat, rabbit, guinea pig, hamster, dog, and cat, but not always limited thereto.

[0057] In the above method, the binding intensity between A β 22(pE)-42 peptide and amyloid-beta antibody can be measured by a method selected from the group consisting of Western blotting, enzyme-linked immunosorbent assay (ELISA), immunohistochemical staining, immunoprecipitation, and immunofluorescence assay, but not always limited thereto.

[0058] Since it was confirmed that A β 22(pE)-42 peptide demonstrated higher immune reactivity against amyloid-beta antibody in serum of Alzheimer's disease patients than normal people, the A β 22(pE)-42 peptide can be effectively used for the diagnosis of dementia.

[0059] The present invention also provides a method for the detection of amyloid-beta antibody to provide information useful for the diagnosis of dementia using A β 22(pE)-42. The method herein is composed of the following steps:

[0060] 1) treating blood taken from a test subject on the solid board attached with the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1, followed by washing thereof; and

[0061] 2) diagnosing the subject with dementia or high risk of dementia when the binding intensity of amyloid-beta antibody on the solid board is significantly higher than that of normal people.

[0062] In addition, the present invention provides a method for detecting amyloid-beta antibody to provide information useful for the diagnosis of dementia using A β 22(pE)-42 peptide. This method is composed of the following steps:

[0063] 1) treating blood taken from a test subject on the solid board, followed by washing thereof;

[0064] 2) treating the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board, followed by washing thereof; and

[0065] 3) diagnosing the subject with dementia or high risk of dementia when the binding intensity of the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board is significantly higher than that of normal people.

[0066] Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples, Experimental Examples and Manufacturing Examples.

[0067] However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

EXAMPLE 1

Antigen-Antibody Separation

[0068] Plasma was obtained from each of normal people (48 people), mild cognitive impairment patients (MCI,

probable Alzheimer's disease patients (11 patients), and severe Alzheimer's disease patients (62 people), which was diluted in dissociation buffer (TBS buffer, 1.5% BSA, 0.2M glycine-acetate, pH 2.5) at the ratio of 1:500, followed by standing at room temperature for 20 minutes. The diluted Plasma was loaded in Amicon Ultra-0.5 ml 10K Ultracel (Millipore, Ireland) filter, followed by centrifugation at 8000 g at room temperature for 20 minutes. The filter was made inside-out and loaded in a new Amicon tube, followed by centrifugation at 1000 g for 2 minutes. At that time, pH was regulated to 7.0 by using 1 M Tris buffer (pH 9.0).

EXAMPLE 2

Protein Chip Assay

[0069] 6 $\times$ His-antibody (Abcam, UK) was diluted in TBS-glycerol buffer (30% glycerol, pH 7.4) at the concentration of 1 μ g/ml, which was loaded on protein chip (Proteogen, South Korea) by 1 μ l at a time, which stood at 4 $^{\circ}$ C. for overnight or at 37 $^{\circ}$ C. for 3 hours. 6 $\times$ His-antibody on the chip was eliminated by using 3M paper, followed by washing softly with washing buffer (TBS buffer, 0.1% tween 20) three times at 10 minutes intervals. The remaining washing buffer on the chip was washed out with distilled water by shaking lightly for 10 minutes, followed by centrifugation at 1500 rpm for 3 minutes to eliminate the remaining solution. Blocking buffer (TBS buffer, 0.1% tween20, 3% BSA) was added thereto, followed by reaction with shaking lightly for 1 hour at room temperature. Washing was performed by the same manner as described above. Supernatant was eliminated by centrifugation. A β 22(pE)-42 peptide (pyrEDVGSNKGAIIGLMVGVGVIA, SEQ. ID. NO: 1) (Peptide 2.0 Inc, www.peptide2.com) (TBS-glycerol buffer, 3% BSA) was loaded on the chip by 1 μ l at a time, which stood at 37 $^{\circ}$ C. for 1 hour. Then, liquid on the chip was eliminated. The antibody separated in Example 1 was diluted with antibody dilution buffer (TBS-glycerol buffer, 3% BSA) at the ratio of 1:4, which was loaded on the chip by 1 μ l. Then the chip was stood at 37 $^{\circ}$ C. for 1 hour. One hour later, liquid on the chip was eliminated and the secondary antibody (Anti-human IgG, cy5) diluted in antibody dilution buffer at the ratio of 1:5000 was loaded on the chip by 1 μ l. Then the chip was stood at 37 $^{\circ}$ C. for 30 minutes. 30 minutes later, liquid on the chip was eliminated. The result was confirmed by using chip scanner.

[0070] Immuno reactivity was measured by using protein chip. As a result, immune reactivity of A β 22(pE)-42 peptide to amyloid-beta antibody was increased in blood obtained from mild cognitive impairment patients and Alzheimer's disease patients, compared with that in blood of normal people (FIG. 1). Particularly, A β 22(pE)-42 peptide demonstrated higher immune reactivity to amyloid-beta antibody in blood of mild cognitive impairment patients (143%, $p < 0.01$), probable Alzheimer's disease patients (146%, $p < 0.01$), and severe Alzheimer's disease patients (148%, $p < 0.001$) than that in blood of normal people (FIG. 2).

[0071] Those skilled in the art will appreciate that the conceptions and specific embodiments disclosed in the foregoing description may be readily utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present invention. Those skilled in the art will also appreciate that such equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended Claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 1

<210> SEQ ID NO 1

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile Gly Leu Met Val Gly
1          5          10          15
Gly Val Val Ile Ala
                20

```

What is claimed is:

1. A method for diagnosis of dementia comprising the following steps:

- 1) treating blood taken from a test subject on the solid board attached with the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1, followed by washing thereof; and
- 2) diagnosing the subject with dementia or high risk of dementia when the binding intensity of amyloid-beta antibody on the solid board is significantly higher than that of normal people.

2. The method for diagnosis of dementia according to claim 1, wherein the peptide is characteristically conjugated to amyloid-beta antibody.

3. The method for diagnosis of dementia according to claim 1, wherein the dementia is selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease, and cerebral amyloid angiopathy.

4. The method for diagnosis of dementia according to claim 1, wherein the peptide is characteristically conjugated with chromogenic enzyme, chromogenic material, or fluorescent molecule.

5. The method for diagnosis of dementia according to claim 4, wherein the chromogenic enzyme is HRP (horseradish peroxidase) or alkaline phosphatase.

6. The method for diagnosis of dementia according to claim 4, wherein the chromogenic material is colloid gold.

7. The method for diagnosis of dementia according to claim 4, wherein the fluorescent molecule is characteristically selected from the group consisting of poly L-lysine-flourescein isothiocynate (FITC), rhodamine-B-isothiocyanate (RITC), CY2, CY3, and CY5.

8. The method for diagnosis of dementia according to claim 1, wherein the binding intensity between A β 22(pE)-42 peptide and amyloid-beta antibody is measured by a method selected from the group consisting of Western blotting, enzyme-linked immunosorbent assay (ELISA), immunohistochemical staining, immunoprecipitation, and immunofluorescence assay.

9. A method for diagnosis of dementia comprising the following steps:

- 1) treating blood taken from a test subject on the solid board, followed by washing thereof;
- 2) treating the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board, followed by washing thereof; and
- 3) diagnosing the subject with dementia or high risk of dementia when the binding intensity of the A β 22(pE)-42 peptide represented by SEQ. ID. NO: 1 on the solid board is significantly higher than that of normal people.

* * * * *

专利名称(译)	新的ABETA22 (pE) -42与血液和药物组合物中的淀粉样抗体特异性结合的肽，用于诊断含有相同成分的痴呆症作为活性成分		
公开(公告)号	US20130089877A1	公开(公告)日	2013-04-11
申请号	US13/645741	申请日	2012-10-05
[标]申请(专利权)人(译)	韩国CENT的疾病控制与预防		
申请(专利权)人(译)	韩国疾病控制中心和预防		
当前申请(专利权)人(译)	韩国疾病控制中心和预防		
[标]发明人	KIM YOUNG YOUL KIM SEUNGWOO CHEON HYU SOON PARK SANG ICK		
发明人	KIM, YOUNG-YOUL KIM, SEUNGWOO CHEON, HYU-SOON PARK, SANG ICK		
IPC分类号	G01N33/53 G01N21/76		
CPC分类号	C07K14/4711 G01N33/6896 G01N2800/2821 G01N2800/2814 G01N2333/4709		
优先权	1020110102611 2011-10-07 KR		
其他公开文献	US8679768		
外部链接	Espacenet USPTO		

摘要(译)

本发明人筛选了具有特异性结合淀粉样蛋白-β抗体的特异性序列的肽，因此证实Aβ22 (pE) -42肽对阿尔茨海默病患者血清中的淀粉样蛋白-β抗体显示出更高的反应性。因此，所述Aβ22 (pE) -42肽可以用作诊断痴呆的试剂盒的有效成分，因此可以说该肽可以有效地用于诊断难以早期诊断的痴呆。

[Fig. 1]

