



US 20180235195A1

(19) **United States**(12) **Patent Application Publication****Dennis et al.**(10) **Pub. No.: US 2018/0235195 A1**(43) **Pub. Date: Aug. 23, 2018**(54) **TRANSFERRIN RECEPTOR TRANSGENIC MODELS**

filed on Aug. 10, 2017, provisional application No. 62/543,559, filed on Aug. 10, 2017, provisional application No. 62/583,314, filed on Nov. 8, 2017.

(71) Applicant: **Denali Therapeutics Inc.**, South San Francisco, CA (US)(72) Inventors: **Mark S. Dennis**, South San Francisco, CA (US); **Adam P. Silverman**, South San Francisco, CA (US); **Joy Yu Zuchero**, South San Francisco, CA (US)(73) Assignee: **Denali Therapeutics Inc.**, South San Francisco, CA (US)(21) Appl. No.: **15/923,928**(22) Filed: **Mar. 16, 2018****Related U.S. Application Data**

(63) Continuation of application No. PCT/US2018/018302, filed on Feb. 15, 2018.

(60) Provisional application No. 62/460,692, filed on Feb. 17, 2017, provisional application No. 62/543,658,

Publication Classification(51) **Int. Cl.****A01K 67/027** (2006.01)**C07K 14/705** (2006.01)**G01N 33/53** (2006.01)(52) **U.S. Cl.**CPC **A01K 67/0278** (2013.01); **C07K 14/705** (2013.01); **G01N 33/5308** (2013.01); **G01N 2500/10** (2013.01); **A01K 2217/075** (2013.01); **A01K 2227/105** (2013.01); **A01K 2267/03** (2013.01); **A01K 2217/072** (2013.01)

(57)

ABSTRACT

In some aspects, the present invention provides chimeric transferrin receptor (TfR) polynucleotides and polypeptides. In other aspects, this invention provides chimeric TfR transgenic animal models and methods of using the animal models to identify therapeutics that can cross the blood-brain barrier.

Specification includes a Sequence Listing.




FIG. 1A




FIG. 1B




FIG. 1C

IV 5 mg/kg anti-TfR

IV 5 mg/kg anti-TfR




FIG. 2B

FIG. 2A

IV 50 mg/kg anti-TfR/BACE1




FIG. 3B

IV 50 mg/kg anti-BACE1




FIG. 3A




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D

TRANSFERRIN RECEPTOR TRANSGENIC MODELS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application is a continuation of International Patent Application Serial No. PCT/US2018/018302, filed Feb. 15, 2018, which application claims the benefit of U.S. Patent Application Ser. No. 62/460,692, filed Feb. 17, 2017, U.S. Patent Application Ser. No. 62/543,658, filed Aug. 10, 2017, U.S. Patent Application Ser. No. 62/543,559, filed Aug. 10, 2017 and U.S. Patent Application Ser. No. 62/583,314, filed Nov. 8, 2017, the contents of which are incorporated herein by reference for all purposes.

BACKGROUND OF THE INVENTION

[0002] The blood-brain barrier (BBB) blocks the passage of most macromolecules from the periphery into the brain and thus limits the uses of large molecule therapeutics where brain exposure is required. Transferrin receptor (TfR) is highly expressed at the BBB and can be used to transport such therapeutics across the BBB via a receptor-mediated transcytosis. Mouse models previously have been developed, in which the mouse TfR was replaced with a full-length human TfR cDNA, with the objective of evaluating the ability of potential therapeutics to cross the BBB. However, these transgenic mice were unhealthy and showed abnormally high TfR expression, low red blood cell count, and high serum iron concentration. Yu et al., *Science Trans. Med.*, 6(261):261ra154 (2014). As a result, these existing mouse models are not suited for use as tools to evaluate therapeutics that are capable of crossing the BBB to treat brain diseases; models that are more representative of endogenous TfR expression and phenotype are required.

BRIEF SUMMARY OF THE INVENTION

[0003] In one aspect, this disclosure provides a polynucleotide comprising a nucleic acid sequence encoding a chimeric transferrin receptor (TfR) polypeptide that comprises a non-human mammalian transferrin binding site and a heterologous apical domain having an amino acid sequence at least 80% identical to SEQ ID NO:1. In some embodiments, the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:1. In some embodiments, the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:7, SEQ ID NO:8, or SEQ ID NO:9.

[0004] In some embodiments, the non-human mammalian transferrin binding site is a native (e.g., from the same species as the transmembrane and/or intracellular domain of the TfR) transferrin binding site, e.g., a native mouse transferrin binding site. In some embodiments, the chimeric TfR polypeptide has at least 80% amino acid sequence identity, or at least 85%, 90%, or 95% identity, to SEQ ID NO:3. In some embodiments, the chimeric TfR polypeptide comprises the amino acid sequence of SEQ ID NO:3. In some embodiments, the region of the nucleic acid sequence encoding the heterologous apical domain of the chimeric TfR polypeptide has at least 70% nucleotide sequence identity to SEQ ID NO:2. In some embodiments, the region of the nucleic acid sequence encoding the heterologous apical domain of the chimeric TfR polypeptide comprises the nucleotide sequence of SEQ ID NO:2. In some embodiments, the polynucleotide encoding the chimeric TfR com-

prises exons and introns of a mouse transferrin receptor gene and the nucleic acid sequence encoding the heterologous apical domain is positioned after the fourth exon of a mouse transferrin receptor gene to replace the apical binding domain of the mouse transferrin receptor gene.

[0005] In another aspect, provided herein is a chimeric TfR polypeptide that comprises a non-human mammalian transferrin binding site and a heterologous apical domain having an amino acid sequence at least 80% identical to SEQ ID NO:1. In some embodiments, the chimeric TfR polypeptide comprises a native TfR polypeptide in which only the native apical domain is replaced by a heterologous apical domain. In some embodiments, a chimeric TfR polypeptide comprises a native TfR binding site and an apical binding domain that is heterologous to the native TfR binding site, e.g., wherein at least one domain, or region thereof, in addition to the apical domain comprises a non-native amino acid sequence. In some embodiments, the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:1. In some embodiments, the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:7, SEQ ID NO:8, or SEQ ID NO:9. In some embodiments, the chimeric TfR has at least 80%, 90%, 95%, or 98% amino acid sequence identity to SEQ ID NO: 3. In some embodiments, the chimeric TfR polypeptide comprises the amino acid sequence of SEQ ID NO:3.

[0006] In a further aspect, provided herein are host cells that express a chimeric transferrin receptor as described above. In some embodiments, a host cell comprises a polynucleotide that encodes the chimeric transferrin receptor polypeptide. In some embodiments, the host cell is a mouse cell. In some embodiments, the chimeric TfR polypeptide expressed by the host cell comprises (a) a heterologous apical domain in place of the endogenous apical domain of the TfR polypeptide and (b) the endogenous transferrin binding site. In some embodiments, the heterologous apical domain has an amino acid sequence at least 80% identical to SEQ ID NO: 1. In some embodiments, the host cell expresses a chimeric TfR in which only the apical domain of the endogenous TfR is replaced by a heterologous apical domain. In some embodiments, a host cell expresses a chimeric TfR comprising an endogenous TfR binding site and a heterologous apical domain, e.g., wherein at least one domain, or region thereof, in addition to the apical domain comprises a non-native amino acid sequence. In some embodiments, the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:1. In some embodiments, the nucleic acid sequence that encodes the heterologous apical domain in the cell comprises the nucleotide sequence of SEQ ID NO:2. In some embodiments, the host cell is a mouse cell. In some embodiments, the nucleic acid sequence encoding the apical domain in the cell is positioned after the fourth exon of a mouse transferrin receptor gene. In some embodiments, the host cell is ex vivo. In some embodiments, the host cell is an embryonic stem cell. In some embodiments, the genome of the host cell comprises a deletion of the apical domain of the native TfR.

[0007] In an additional aspect, the disclosure provides a non-human transgenic animal that expresses a chimeric TfR polypeptide, wherein the chimeric TfR polypeptide comprises a heterologous apical domain that replaces the apical domain of the TfR polypeptide endogenous to the non-human transgenic animal. In some embodiments, the genome of the non-human transgenic animal comprises a

transferrin receptor gene that encodes a heterologous apical domain in place of the apical domain of the endogenous TfR of the non-human transgenic animal. In some embodiments, the non-human transgenic animal expresses a chimeric TfR comprising a heterologous apical domain in place of the native domain of the TfR of the non-human transgenic animal and a native transferrin binding site. In some embodiments, a non-human transgenic animal expresses a chimeric transferrin receptor in which only the apical domain of the endogenous transferrin receptor is replaced by a heterologous apical domain. In some embodiments, a non-human transgenic animal expresses a chimeric TfR polypeptide comprising an endogenous TfR binding site and an apical binding domain that is heterologous to the endogenous TfR binding site, e.g., wherein at least one domain, or region thereof, in addition to the apical domain comprises a non-native amino acid sequence. In some embodiments, the non-human transgenic animal comprises the host cells as described above. In some embodiments, the transgenic animal is a rodent. In some embodiments, the transgenic animal is a mouse or a rat. In some embodiments, the transgenic animal is homozygous for the chimeric TfR. In some embodiments, the transgenic animal is heterozygous for the chimeric TfR.

[0008] In another aspect, provided herein is a method of screening for an apical domain binding polypeptide (ADBP) that binds to a chimeric TfR, the method comprising contacting a candidate ADBP with a chimeric TfR polypeptide as described above; and determining the amount of the candidate ADBP that binds to the chimeric TfR polypeptide. In some embodiments, the step of contacting the candidate ADBP with the chimeric TfR polypeptide comprises contacting the ADBP with a host cell that expresses the chimeric TfR polypeptide. In some embodiments, the step of contacting the candidate ADBP with the chimeric TfR polypeptide comprises contacting the ADBP with an endothelium that expresses the chimeric TfR polypeptide. In some embodiments, the endothelium is a blood-brain barrier endothelium. In some embodiments, the amount of the candidate ADBP that binds the chimeric TfR polypeptide is determined by immunoassay. In some embodiments, the amount of the candidate ADBP that binds the chimeric TfR polypeptide is determined by surface plasmon resonance. In some embodiments, wherein contacting step is performed in vivo. In some embodiments the candidate ADBP is coupled to an effector molecule. In some embodiments, the effector molecule is a small molecule, RNA, DNA, or polypeptide. In some embodiments, the effector molecule is a polypeptide. In some embodiments, the polypeptide is an antibody or an antigen-binding fragment thereof.

[0009] In yet another aspect, provided herein is a method of measuring the amount of an ADBP that binds to a chimeric TfR polypeptide, the method comprising contacting the ADBP with a chimeric TfR polypeptide disclosed above; and determining the amount of ADBP bound to the chimeric TfR polypeptide by immunoassay or surface plasmon resonance.

[0010] In yet another aspect, provided herein is a method of screening for an ADBP that crosses the blood-brain barrier, the method comprising: (a) administering an ADBP that binds an apical domain having at least 80% amino acid sequence identity to SEQ ID NO:1 to a non-human transgenic animal as disclosed herein; and (b) measuring the presence or an activity of the ADBP in the brain of the

non-human transgenic animal. In some embodiments, the ADBP is coupled to an effector molecule. In some embodiments, the effector molecule is a small molecule, RNA, DNA, or polypeptide. In some embodiments, the polypeptide is an antibody or an antigen-binding fragment thereof. In some embodiments, the determining step comprises performing a quantitative immunoassay. In some embodiments, the measuring step comprises contacting the brain or brain tissue of the animal with an agent that binds to the effector molecule to determine the level of the effector molecule in the brain. In some embodiments, the measuring step comprises measuring a pharmacodynamic (PD) effect of the effector molecule. In some embodiments, the effector molecule is an anti-BACE1 antibody or an antigen-binding fragment thereof and the measuring step comprises measuring the level of soluble ABeta40 in the brain. In some embodiments, the effector molecule is an antibody or an antigen-binding fragment thereof that binds to a target in the brain.

[0011] In another aspect, provided herein is a method of monitoring an ADBP that crosses the blood-brain barrier, the method comprising: (a) administering an ADBP that binds an apical domain having at least 80% amino acid sequence identity to SEQ ID NO:1 to a non-human transgenic animal as disclosed herein; and (b) measuring the presence or an activity of the ADBP in the brain of the non-human transgenic animal. In some embodiments, the ADBP is coupled to an effector molecule. In some embodiments, the effector molecule is a small molecule, RNA, DNA, or polypeptide. In some embodiments, the polypeptide is an antibody or an antigen-binding fragment thereof. In some embodiments, the determining step comprises performing a quantitative immunoassay. In some embodiments, the determining step comprises contacting the effector molecule with an agent that binds to the effector molecule and determining the level of level of the effector molecule present in the brain. In some embodiments, the effector molecule is an antibody or an antigen-binding fragment thereof that binds to a target in the brain. In some embodiments, the measuring step comprises measuring a PD effect of the effector molecule binding to the target. In some embodiments, the effector molecule is an anti-BACE1 antibody or an antigen-binding fragment thereof and the measuring step comprises measuring the level of soluble ABeta40 in the brain.

[0012] In yet another aspect, provided herein is a method of generating a transgenic non-human single cell embryo that expresses a chimeric transferrin receptor (TfR) polypeptide, the method comprising replacing the apical domain of the endogenous TfR in the non-human single cell embryo with a heterologous apical domain having at least 80% identity to SEQ ID NO:1. In some embodiments, replacing the apical domain is performed by homologous recombination. In some embodiments, the method comprises contacting a Cas9 protein, at least one single guide RNA (sgRNA), and a donor DNA comprising a nucleic acid sequence encoding the heterologous apical domain, wherein the heterologous apical domain is flanked by a left homology arm and a right homology arm, such that the heterologous apical domain coding sequence replaces the apical domain of the endogenous TfR in the genome of the non-human single cell embryo. In some embodiments, the heterologous apical domain is codon-optimized for expression in the non-human single cell embryo. In some embodiments the non-human single cell embryo is a mouse embryo. In some embodi-

ments, the donor DNA is positioned after the fourth exon of a mouse transferrin receptor gene.

[0013] In yet another aspect, provided herein is a method of generating a non-human transgenic animal comprising (a) transferring the transgenic non-human single cell embryo disclosed above to a pseudo pregnant female of the same animal species as the non-human single cell embryo, and (b) selecting a non-human transgenic animal from the progeny produced by the female, wherein the non-human transgenic animal comprises a chimeric transferrin receptor (TfR) polypeptide in which the apical domain of an endogenous TfR has been replaced with a heterologous apical domain having an amino acid sequence of at least 80% identity to SEQ ID NO:1.

[0014] In yet another aspect, provided herein is a method of generating a non-human transgenic animal that expresses a chimeric transferrin receptor (TfR) polypeptide, the method comprising (a) introducing into an embryonic cell of the animal a polynucleotide encoding an apical domain having at least 80% identity to SEQ ID NO:1, wherein the polynucleotide is targeted to a region of an endogenous TfR gene that encodes an endogenous TfR apical domain and wherein the polynucleotide encoding the apical domain having at least 80% identity to SEQ ID NO:1 replaces the region of the endogenous TfR gene that encodes the endogenous apical domain, and (b) developing the cell or progeny thereof into a non-human transgenic animal.

[0015] The foregoing general description and the following detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed. Other objects, advantages and novel features will be readily apparent to those skilled in the art from the following detailed description of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] FIGS. 1A-1C show the results of a complete blood count analysis of wild-type, huTfR^{apical+/-}, and huTfR^{apical+/+} mice. No genotype-specific differences were observed in total red blood cells, hemoglobin, or hematocrit. Graphs in the figure represent mean±SD, n=3 per group.

[0017] FIGS. 2A-2B show the brain vascular localization of systemically administered anti-TfR in huTfR^{apical+/-} mice. 5 mg/kg of a human-apical-specific anti-TfR was intravenously administered into either C57Bl6 wild-type or chimeric huTfR^{apical+/-} heterozygous mice. After one hour, mice were perfused with PBS, and brains were stained for antibody distribution. Representative images show prominent vascular localization of systemically injected anti-TfR (apical domain-specific) in the huTfR^{apical+/-} heterozygous, but not in the wild-type mice. This indicates that the chimeric huTfR protein is expressed at the BBB.

[0018] FIGS. 3A-3B show the brain parenchymal distribution of systemically administered anti-TfR/BACE1 in huTfR^{apical+/+} mice. Representative images of cortical brain sections from huTfR mice show broad parenchymal distribution of intravenously-injected anti-TfR/BACE1 (50 mg/kg, 24 hrs post-dose). In contrast, no appreciable staining was observed in brain sections from huTfR^{apical+/-} huTfR mice injected with anti-BACE1.

[0019] FIGS. 4A-4D show the brain uptake of an anti-TfR/BACE1 bispecific antibody and A-beta reduction followed by the administration of the bispecific antibody in huTfR^{apical+/+} mice. FIG. 4A shows plasma huIgG1 concentration in huTfR^{apical+/+} mice 24 hours after the mice were

administered with 50 mg/kg of anti-TfR/BACE1 or anti-BACE1. The results showed that TfR-mediated clearance of anti-TfR/BACE1 was enhanced relative to that of anti-BACE1. FIG. 4B shows mean brain uptake of anti-TfR/BACE1 following a systemic dosing of the antibody. The results show an increase of about 28-fold of the accumulation of anti-TfR/BACE1 as compared to that of anti-BACE1 in mouse brains. FIG. 4C shows that A-beta in the brain was reduced by 49% in mice treated with anti-TfR/BACE1 as compared to mice treated with anti-BACE1. FIG. 4D shows a reduction in the plasma A-beta level in huTfR^{apical+/+} mice that had been treated with anti-BACE1 or anti-TfR/BACE1 as compared to untreated wild-type mice. All graphs represent mean±SD, n=8 per group (n=2 for untreated wild-type mice).

[0020] FIGS. 5A-5D show the expression of TfR in various tissues in huTfR^{apical+/+} mice as compared to wild-type mice; no marked differences in total TfR expression in brain (FIG. 5A), liver (FIG. 5B), kidney (FIG. 5C), and lung (FIG. 5D) were observed. All graphs represent mean±SD, n=4-8 per group.

DETAILED DESCRIPTION OF THE INVENTION

[0021] We have developed chimeric forms of the transferrin receptor that include a non-human (e.g., mouse) mammalian transferrin binding site and an apical domain that is heterologous to the domain containing the transferrin binding site. These chimeric receptors can be expressed in transgenic animals, particularly where the transferrin binding site is derived from the transgenic animal species and where the apical domain is derived from a primate (e.g., human or monkey). The present invention therefore provides a polynucleotide encoding a chimeric transferrin receptor that comprises a non-human mammalian transferrin binding site and an apical domain having an amino acid sequence at least 80% identical to SEQ ID NO: 1. The invention also provides a non-human, for example, non-primate, transgenic animal expressing such chimeric TfRs and the use of the non-human transgenic animal to screen for polypeptides that can cross the BBB by binding to human transferrin receptor (huTfR) in vivo. In some embodiments, the non-human transgenic animal contains a native transferrin receptor (such as a mouse transferrin receptor (mTfR)), in which the apical domain is replaced with an orthologous apical domain having an amino acid sequence at least 80% identical to SEQ ID NO:1, thereby leaving the native transferrin binding site and the majority, e.g., at least 70%, or at least 75%, of the sequence encoding the transferrin receptor intact. This non-human transgenic animal thus maximally retains the transferrin-binding functionality of the endogenous transferrin receptor of the non-human animal, including the ability to maintain proper iron homeostasis as well as bind and transport transferrin. As a result, the transgenic animal is healthy and suitable for use in discovery and development of therapeutics for treating brain diseases.

Terminology

[0022] As used herein, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “an antibody” optionally includes a combination of two or more such molecules, and the like.

[0023] As used herein, the terms “about” and “approximately,” when used to modify an amount specified in a numeric value or range, indicate that the numeric value as well as reasonable deviations from the value known to the skilled person in the art, for example, $\pm 20\%$, $\pm 10\%$, or $\pm 5\%$, are within the intended meaning of the recited value.

[0024] A “transferrin receptor” as used herein refers to transferrin receptor protein 1. The human transferrin receptor 1 polypeptide sequence is set forth in SEQ ID NO:6. Transferrin receptor protein 1 sequences from other species are also known (e.g., chimpanzee, accession number XP_003310238.1; rhesus monkey, NP_001244232.1; dog, NP_001003111.1; cattle, NP_001193506.1; mouse, NP_035768.1; rat, NP_073203.1; and chicken, NP_990587.1). The term “transferrin receptor” also encompasses allelic variants of exemplary reference sequences, e.g., human sequences, that are encoded by a gene at a transferrin receptor protein 1 chromosomal locus. Full length transferrin receptor protein includes a short N-terminal intracellular region, a transmembrane region, and a large extracellular domain. The extracellular domain is characterized by three domains: a protease-like domain, a helical domain, and an apical domain.

[0025] The term “chimeric TfR” as used herein refers to a transferrin receptor protein that has all or a subregion of the apical domain replaced with a corresponding apical domain region from a heterologous transferrin receptor.

[0026] A “transferrin binding site” as used herein refers to regions in the helical and protease-like domain of a TfR protein that mediate binding of transferrin, e.g., iron-bound transferrin, to the receptor. The transferrin binding site is distal to the apical domain.

[0027] A “non-human mammalian transferrin binding site” as used herein refers to a sequence from the transferrin binding site of the native transferrin receptor of a non-human mammal, or a functional derivative thereof that is capable of binding to the native non-human mammalian transferrin. In some embodiments, the non-human mammalian transferrin binding site comprises an amino acid sequence that is at least 80%, at least 90%, at least 95%, or at least 98% identical to the transferrin binding site of the native transferrin receptor of the non-human mammal. Examples of the non-human mammals include mouse, rat, rabbit, bovine, ovine, canine, feline, equine, porcine, non-human primates, and the like.

[0028] As used herein, an “huTfR^{apical+/+} mouse” refers to a transgenic mouse in which the apical domain of the mouse transferrin receptor has been replaced with the apical domain of a human transferrin receptor, and the transgenic mouse being homozygous for the transgene.

[0029] As used herein, an “huTfR^{apical+/-} mouse” refers to a transgenic mouse in which the apical domain of the mouse transferrin receptor has been replaced with the apical domain of a human transferrin receptor; and the transgenic mouse being heterozygous for the transgene.

[0030] As used herein, the terms “wild-type,” “native,” and “naturally occurring” with respect to a transferrin receptor or a domain thereof, refer to a transferrin receptor or a domain thereof that has a sequence that occurs in nature.

[0031] An “endogenous” transferrin receptor or domain thereof as used herein refers to a transferrin receptor that naturally occurs in a cell or non-human animal, i.e., in the absence of genetic modification to the cell or animal.

[0032] As used herein, the term “heterologous” with respect to a domain of a transferrin receptor, e.g., the apical domain, refers to a domain of the transferrin receptor that is expressed outside its native context, e.g., separated from transferrin receptor sequences with which it typically is in proximity in nature, or adjacent (or contiguous with) transferrin receptor sequences with which it typically is not in proximity.

[0033] The term “amino acid” refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids.

[0034] Naturally-occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate and O-phosphoserine. Naturally-occurring α -amino acids include, without limitation, alanine (Ala), cysteine (Cys), aspartic acid (Asp), glutamic acid (Glu), phenylalanine (Phe), glycine (Gly), histidine (His), isoleucine (Ile), arginine (Arg), lysine (Lys), leucine (Leu), methionine (Met), asparagine (Asn), proline (Pro), glutamine (Gln), serine (Ser), threonine (Thr), valine (Val), tryptophan (Trp), tyrosine (Tyr), and combinations thereof. Stereoisomers of a naturally-occurring α -amino acids include, without limitation, D-alanine (D-Ala), D-cysteine (D-Cys), D-aspartic acid (D-Asp), D-glutamic acid (D-Glu), D-phenylalanine (D-Phe), D-histidine (D-His), D-isoleucine (D-Ile), D-arginine (D-Arg), D-lysine (D-Lys), D-leucine (D-Leu), D-methionine (D-Met), D-asparagine (D-Asn), D-proline (D-Pro), D-glutamine (D-Gln), D-serine (D-Ser), D-threonine (D-Thr), D-valine (D-Val), D-tryptophan (D-Trp), D-tyrosine (D-Tyr), and combinations thereof.

[0035] Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission.

[0036] The terms “polypeptide,” “peptide,” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymers. Amino acid polymers may comprise entirely L-amino acids, entirely D-amino acids, or a mixture of L and D amino acids.

[0037] “Conservatively modified variant” refers to an alteration that results in the substitution of an amino acid with another amino acid that can be categorized as having a similar feature. Examples of categories of conservative amino acid groups defined in this manner can include: a “charged/polar group” including Glu (Glutamic acid or E), Asp (Aspartic acid or D), Asn (Asparagine or N), Gln (Glutamine or Q), Lys (Lysine or K), Arg (Arginine or R), and His (Histidine or H); an “aromatic group” including Phe (Phenylalanine or F), Tyr (Tyrosine or Y), Trp (Tryptophan or W), and (Histidine or H); and an “aliphatic group” including Gly (Glycine or G), Ala (Alanine or A), Val (Valine or V), Leu (Leucine or L), Ile (Isoleucine or I), Met (Methionine or M), Ser (Serine or S), Thr (Threonine or T), and Cys (Cysteine or C). Within each group, subgroups can also be identified. For example, the group of charged or polar amino acids can be sub-divided into sub-groups including: a “positively-charged sub-group” comprising Lys, Arg and His; a “negatively-charged sub-group” com-

prising Glu and Asp; and a “polar sub-group” comprising Asn and Gln. In another example, the aromatic or cyclic group can be sub-divided into sub-groups including: a “nitrogen ring sub-group” comprising Pro, His and Trp; and a “phenyl sub-group” comprising Phe and Tyr. In another further example, the aliphatic group can be sub-divided into sub-groups, e.g., an “aliphatic non-polar sub-group” comprising Val, Leu, Gly, and Ala; and an “aliphatic slightly-polar sub-group” comprising Met, Ser, Thr, and Cys. Examples of categories of conservative mutations include amino acid substitutions of amino acids within the sub-groups above, such as, but not limited to: Lys for Arg or vice versa, such that a positive charge can be maintained; Glu for Asp or vice versa, such that a negative charge can be maintained; Ser for Thr or vice versa, such that a free —OH can be maintained; and Gln for Asn or vice versa, such that a free —NH₂ can be maintained. In some embodiments, hydrophobic amino acids are substituted for naturally occurring hydrophobic amino acid, e.g., in the active site, to preserve hydrophobicity.

[0038] The terms “identical” or percent “identity,” in the context of two or more polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues, e.g., at least 60% identity, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% or greater, that are identical over a specified region when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one a sequence comparison algorithm or by manual alignment and visual inspection.

[0039] For sequence comparison of polypeptides, typically one amino acid sequence acts as a reference sequence, to which a candidate sequence is compared. Alignment can be performed using various methods available to one of skill in the art, e.g., visual alignment or using publicly available software using known algorithms to achieve maximal alignment. Such programs include the BLAST programs, ALIGN, ALIGN-2 (Genentech, South San Francisco, Calif.) or Megalign (DNASTAR). The parameters employed for an alignment to achieve maximal alignment can be determined by one of skill in the art. For sequence comparison of polypeptide sequences for purposes of this application, the BLASTP algorithm standard protein BLAST for aligning two proteins sequence with the default parameters is used.

[0040] The term “comprising” is intended to mean that the compositions and methods include the recited elements, but do not exclude others. “Consisting essentially of” when used to define compositions and methods, refers to the specified materials or steps and those that do not materially affect the basic and novel characteristic(s) of the claimed invention. “Consisting of” shall mean excluding more than trace amounts of other ingredients and substantial method steps recited. Embodiments defined by each of these transition terms are within the scope of this invention.

[0041] The terms “polynucleotide,” “nucleic acid,” and “oligonucleotide” are used interchangeably and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides or analogs thereof. Polynucleotides can have any three-dimensional structure and may perform any function, known or unknown. The following are non-limiting examples of polynucleotides: a gene or gene fragment (for example, a probe, primer, EST or SAGE tag), exons, introns, messenger RNA (mRNA), transfer

RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes and primers. A polynucleotide can comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure can be imparted before or after assembly of the polynucleotide. The sequence of nucleotides can be interrupted by non-nucleotide components. A polynucleotide can be further modified after polymerization, such as by conjugation with a labeling component. The term also refers to both double and single stranded molecules. Unless otherwise specified or required, any embodiment of this invention that is a polynucleotide encompasses both the double stranded form and each of two complementary single stranded forms known or predicted to make up the double stranded form.

[0042] A polynucleotide is composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); thymine (T); and uracil (U) for thymine when the polynucleotide is RNA. Thus, the term “polynucleotide sequence” is the alphabetical representation of a polynucleotide molecule.

[0043] The term “knock-in” refers to a one-for-one substitution of DNA sequence information in a predetermined genetic locus or the insertion of sequence information not found within the locus. Those skilled in the art will readily appreciate how to use various genetic approaches, e.g., CRISPR/Cas9 systems, ZFN, TALEN, transposon-mediated insertion, to knock in a target polynucleotide sequence in a specific locus of the genome.

[0044] The term “blood-brain barrier” or “BBB” refers to a highly selective semipermeable membrane barrier that separates the circulating blood from the brain extracellular fluid in the central nervous system (CNS). The blood-brain barrier is formed by brain endothelial cells, which are connected by tight junctions.

Transferrin Receptor

[0045] A transferrin receptor mediates cellular uptake of iron via receptor-mediated endocytosis of ligand-occupied transferrin receptor. TfR is present both in human and non-human species, such as non-human primates and rodents. The native human TfR (huTfR), Uniprot P02786, SEQ ID NO:6, is a homodimeric type II transmembrane protein; it has a cytoplasmic domain, a transmembrane region, and an extracellular domain, which comprises an apical domain and a transferrin-binding domain. Each monomer of the huTfR has three structurally distinct domains: a protease-like domain proximal to the membrane, a helical domain accounting for all the dimer contacts, and a membrane-distal apical domain (Lawrence et al., *Science*, 286 (1999), pp. 779-782). HuTfR dimer has a molecular weight of about 190,000 Daltons. The apical domain of the huTfR, which has a sequence of SEQ ID NO:1 (encoded by SEQ ID NO:2), does not participate in the interaction between transferrin and TfR. It has been speculated that this domain may provide contact surface for other proteins to bind the TfR. The native cynomolgous monkey, native rhesus monkey, and native chimpanzee TfRs are also known, as represented, for example, by accession numbers XP_005545315, NP_001244232.1, and XP_003310238.1, respectively. The apical domain of the native cynomolgous monkey, native rhesus monkey, and native chimpanzee TfRs

share about 96%, 95%, and 98% sequence identity, respectively, with the apical domain of the native human TfR of SEQ ID NO:1.

[0046] The native mouse TfR (mTfR), Uniprot Q62351, SEQ ID NO:5 has about 77% amino acid sequence identity with huTfR. The apical domain of the native mTfR is about 74% identical to that of the native huTfR. The mTfR contains the three structurally distinct domains that are similar to the human counterparts. The complete gene sequence for mouse TfR, with annotated exons and introns, can be found from the NCBI database (Gene ID: 22042). Mouse TfR is found on chromosome 16 (NCBI reference sequence NC 000082.6).

[0047] One aspect includes a chimeric TfR polypeptide. In some embodiments, the chimeric TfR comprises a non-human mammalian transferrin binding site and a heterologous apical domain that shares an amino acid sequence identity, e.g., at least 75%, at least 77%, at least 80%, at least 85%, at least 90%, or at least 95%, with the apical domain of huTfR, SEQ ID NO:1. In some embodiments, the heterologous apical domain has a sequence of SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:8, or SEQ ID NO:9.

[0048] The non-human mammalian transferrin binding site of the chimeric TfR allows for the specific binding of the non-mammalian transferrin to the chimeric TfR. In some embodiments, the non-human mammalian transferrin binding site is a native transferrin binding site, e.g., a mouse transferrin receptor binding site.

[0049] In some embodiments, a chimeric TfR polypeptide comprises a native TfR polypeptide in which only the native apical domain is replaced by a heterologous apical domain. In some embodiments, a chimeric TfR polypeptide comprises a native TfR binding site and an apical binding domain that is heterologous to the native TfR binding site, e.g., wherein at least one domain, or region thereof, in addition to the apical domain has a non-native amino acid sequence.

[0050] In some embodiments, the chimeric TfR polypeptide has at least 80%, at least 85%, at least 85%, at least 92%, at least 95%, or at least 98% amino acid sequence identity to SEQ ID NO:3. In one embodiment, the polynucleotide encoding the chimeric TfR polypeptide comprises exons and introns of a mouse transferrin receptor gene and a nucleic acid sequence encoding the huTfR apical domain. In one embodiment, a non-human mammalian TfR apical domain is replaced by a huTfR apical domain coding sequence, for example by replacing the corresponding exons in the non-human mammalian TfR gene with a huTfR apical domain sequence. In an exemplary embodiment, the non-human mammalian TfR gene is a mouse TfR gene. In one embodiment, a mTfR apical domain is replaced by a huTfR apical domain coding sequence, which is positioned, for example, after the fourth exon of the mouse transferrin receptor gene in order to produce the chimeric TfR.

[0051] In some aspects, the invention provides isolated nucleic acids comprising a nucleic acid sequence encoding any of the polypeptides comprising a chimeric TfR polypeptide described herein. In some embodiments, the region of the nucleic acid sequence encoding the heterologous apical domain in the chimeric TfR polypeptide shares a nucleic acid sequence identity of at least 75%, at least 77%, at least 80%, at least 85%, at least 90%, or at least 95% with the coding sequence of the apical domain of the native huTfR, SEQ ID NO:2.

[0052] In another aspect, polynucleotides are provided that comprise a nucleotide sequence that encodes the chimeric transferrin receptor described herein. The polynucleotides may be single-stranded or double-stranded. In some embodiments, the polynucleotide is DNA. In particular embodiments, the polynucleotide is cDNA. In some embodiments, the polynucleotide is RNA.

Codon Optimization

[0053] In some embodiments, the coding sequence for the chimeric TfR, especially the sequence that encodes the huTfR apical domain, is codon optimized in order to improve the expression of the chimeric TfR in mouse. Methods for codon optimization are readily available, for example, optimizer, which is accessible at <http://genomes.urv.es/OPTIMIZER>, and GeneGPS® Expression Optimization Technology from DNA 2.0 (Newark, Calif.). In a preferred embodiment, the coding sequence is codon-optimized for expression in mouse using the OptimumGene™ algorithm from GenScript (Piscataway, N.J.).

Methods for Replacing the Apical Domain of the Non-Human Mammalian Transferrin Receptor with a Desired Apical Domain

[0054] A non-human transgenic animal comprising a knock-in of a heterologous apical domain as disclosed herein can be generated using a variety of methods, for example, a zinc finger nuclease (ZFN), a Tale-effector domain nuclease (TALEN), a transposon-mediated system, and the CRISPR/Cas9 system. These methods typically comprise administering to the cell one or more polynucleotides encoding one or more nucleases such that the nuclease mediates modification of the endogenous gene by cleaving the DNA to create 5' and 3' cut ends in the DNA strand. In the presence of a donor sequence that is flanked by a left and a right homology arms that are substantially homologous to a sequence extending 5' from the 5' end and a sequence extending 3' from the 3' end, the donor is integrated into the endogenous gene targeted by the nuclease via homology-directed repair (HDR). In some embodiments, the knock-in is conducted using the CRISPR/Cas9 system. For example, a nucleic acid sequence encoding a heterologous apical domain is introduced into an endogenous TfR gene to generate a chimeric TfR, which results in that the naturally occurring sequences that encodes the apical domain is replaced but the overall structure of the gene is maintained.

CRISPR

[0055] In some embodiments, the knock-in of the apical domain that is at least 80% identical to SEQ ID NO:1 is performed using the CRISPR/Cas9 system. The CRISPR/Cas9 system includes a Cas9 protein and at least one to two ribonucleic acids that are capable of directing the Cas9 protein to and hybridizing to a target motif in the apical domain of the transferrin receptor that is to be replaced. These ribonucleic acids are commonly referred to as the "single guide RNA" or "sgRNA." The Cas9 protein then cleaves the target motif, which results in a double-strand break or a single-strand break. In the presence of a donor DNA that comprises huTfR apical domain coding sequence flanked by two homology arms, the donor DNA is inserted into the target transferrin receptor DNA, replacing the apical domain.

[0056] The Cas9 protein used in the invention can be a naturally occurring Cas9 protein or a functional derivative thereof. A “functional derivative” of a native sequence polypeptide is a compound having a qualitative biological property in common with a native sequence polypeptide. “Functional derivatives” include, but are not limited to, fragments of a native sequence and derivatives of a native sequence polypeptide and its fragments, provided that they have a biological activity in common with a corresponding native sequence polypeptide. A biological activity contemplated herein is the ability of the functional derivative of Cas9 to hydrolyze a DNA substrate into fragments. Suitable functional derivatives of a Cas9 polypeptide or a fragment thereof include but are not limited to mutants, fusions, covalent modifications of Cas9 protein or a fragment thereof.

[0057] In some embodiments, the Cas9 protein is from *Streptococcus pyogenes*. Cas9 contains 2 endonuclease domains, including a RuvC-like domain which cleaves target DNA that is noncomplementary to the sgRNA, and an HNH nuclease domain which cleave target DNA complementary to sgRNA. The double-stranded endonuclease activity of Cas9 also requires that a short conserved sequence (2-5 nucleotides), known as a protospacer-associated motif (PAM), follows immediately 3' of a target motif in the target sequence. In some embodiments, the PAM motif is an NGG motif. In one illustrative embodiment, the apical domain in a mouse is replaced by using the Cas9 protein, which is directed by sgRNAs to the region between exons 4 and 9 of the mouse gene. A donor DNA is introduced to the reaction. The donor DNA comprises a human apical domain coding sequence that is between a left homology arm that is homologous to the mouse TfR sequence starting upstream of exon 4 and a right homology arm that is homologous to the mouse TfR sequence starting within exon 9. In certain embodiments, the left homology arm overlaps the mouse TfR sequence by 817 nucleotides, starting upstream of exon 4, and the right homology arm overlaps the mouse TfR sequence by 807 nucleotides, starting within exon 9. As a result, the nucleotide sequence encoding a desired apical domain, such as the one that has an amino acid sequence at least 80% identical to SEQ ID NO:1, can be inserted after the fourth mouse exon, and the inserted nucleotide sequence is flanked at the 3' end by the appropriately following mouse exon. In some embodiments, the human apical domain coding sequence that is inserted into the mouse TfR gene is codon-optimized for mouse expression.

[0058] The sgRNAs can be selected depending on the particular CRISPR/Cas9 system employed and the sequence of the target polynucleotide. In some embodiments, the one to two ribonucleic acids are designed to hybridize to a target motif immediately adjacent to a deoxyribonucleic acid motif recognized by the Cas9 protein. In some embodiments, each of the one to two ribonucleic acids are designed to hybridize to target motifs immediately adjacent to deoxyribonucleic acid motifs recognized by the Cas9 protein, wherein the target motifs flank the genomic sequence to be replaced. Guide RNAs can be designed using software that is readily available, for example, at <http://crispr.mit.edu>. Illustrative sgRNAs that can be used to generate a chimeric TfR transgenic mouse include SEQ ID NOs:10-11.

[0059] The donor DNA as disclosed herein comprises a nucleotide sequence that encodes an amino acid sequence at

least 75% identical to SEQ ID NO:1. In some embodiments, the donor DNA comprises a sequence encoding SEQ ID NO:1 or encoding a sequence that shares at least 75%, at least 77%, at least 80%, at least 85%, at least 90%, or at least 95% amino acid sequence identity with SEQ ID NO:1. In some embodiments, the donor DNA comprises the nucleotide sequence of SEQ ID NO:2 or a sequence that shares at least 60%, at least 70%, at least 77%, at least 80%, at least 85%, at least 90%, or at least 95% sequence identity with SEQ ID NO:2. The donor DNA as disclosed herein further comprises a left homology arm and a right homology arm that flank the apical domain coding sequence and are designed to overlap the 5' and 3' exon sequences relative to the cleave site by the Cas9 protein. The homology arms may extend beyond the 5' and 3' exon sequences, and each of the homology arms may be at least 20, 30, 40, 50, 100, or 150 nucleotides in length. One of skilled in the art can readily determine the optimal length of the homology arm required for the experiment. In one illustrative embodiment, the left homology arm of the donor DNA spans nucleotides 1-817 of SEQ ID NO:4 and the right homology arm spans nucleotides 1523-2329 of SEQ ID NO:4. In some embodiments, the left homology arm shares at least 85%, at least 90%, at least 95%, or at least 98% sequence identity to nucleotides 1-817 of SEQ ID NO:4. In some embodiments, the right homology arm shares at least 85%, at least 90%, at least 95%, or at least 98% sequence identity to nucleotides 1523-2329 of SEQ ID NO:4.

[0060] In some embodiments, the sgRNAs can also be selected to minimize hybridization with nucleic acid sequences other than the target polynucleotide sequence. In some embodiments, the one to two ribonucleic acids are designed to hybridize to a target motif that contains at least two mismatches when compared with all other genomic nucleotide sequences in the cell to minimize off-target effects of the CRISPR/Cas9 system. Those skilled in the art will appreciate that a variety of techniques can be used to select suitable target motifs for minimizing off-target effects (e.g., bioinformatics analyses). Methods of using the CRISPR/Cas9 system to reduce gene expression are described in various publications, e.g., US Pat. Pub. Nos. 2014/0170753 and 2016/0257974, the disclosures of which hereby are incorporated by reference in their entirety.

Zinc Finger Nuclease (ZFN)

[0061] In some embodiments, the chimeric TfR is produced by knocking-in the huTfR apical domain using a ZFN. ZFNs are fusion proteins that comprise a non-specific cleavage domain (N) of FokI endonuclease and a zinc finger protein (ZFP). A pair of ZNFs are involved to recognize a specific locus in a target gene: one that recognizes the sequence upstream and the other that recognizes the sequence downstream of the site to be modified. The nuclease portion of the ZFN cuts at the specific locus. The donor DNA as described above can then be inserted into the specific locus. Methods of using the ZFNs to reduce gene expression are well known, for example, as disclosed in U.S. Pat. No. 9,045,763 and also in Durai et al., “Zinc Finger Nucleases: Custom-Designed Molecular Scissors for Genome Engineering of Plant and Mammalian cells,” *Nucleic Acid Research*, 33 (18):5978-5990 (2005), the disclosures of which are incorporated by reference in their entirety.

Transcription Activator-Like Effector Nucleases (TALENs)

[0062] In some embodiments, the chimeric TfR is produced by knocking-in the huTfR apical domain with TALENs. TALENs are similar to ZFNs in that they bind as a pair around a genomic site and direct the same non-specific nuclease, FokI, to cleave the genome at a specific site, but instead of recognizing DNA triplets, each domain recognizes a single nucleotide. Methods of using the ZFNs to reduce gene expression are also well known, for example, as disclosed in U.S. Pat. No. 9,005,973 and also Christian et al., "Targeting DNA Double-Strand Breaks with TAL Effector Nucleases," *Genetics*, 186(2): 757-761 (2010), the disclosures of which are incorporated by reference in their entirety.

Host Cells/Transgenic Animals Expressing the Chimeric TfR

[0063] In some embodiments, the invention provides a host cell that expresses the chimeric TfR, e.g., comprising a nucleic acid sequence encoding the chimeric transferrin receptor described above. In some embodiments, the host cell is a non-human mammalian cell. Any of knock-in methods described above, i.e., CRISPR, TALEN, Zinc finger nuclease, can be used to replace the apical domain of the native transferrin receptor in the host cells with a heterologous apical domain that has an amino acid sequence at least 80% identical to SEQ ID NO:1. In some embodiments, the host cell is eukaryotic, e.g., a mouse cell. at least 80% identical to SEQ ID NO:1. In some cases, the host cell is contacted with the sgRNA and Cas9, a donor DNA that comprises the nucleic acid sequence encoding the heterologous apical domain, the nucleic acid sequence being flanked with a left and a right homology arms. The sgRNA and homology arms having sequences such that the heterologous apical domain coding sequence is inserted into the location in the genome to replace the coding sequence of the apical domain of the native transferrin receptor in the host cell. In some embodiments, the host cell is a cell from a non-primate mammal, such as a mouse, rat, rabbit, bovine, ovine, canine, feline, equine, porcine, and the like.

[0064] In some embodiments, the method of knock-in is performed in an embryonic stem (ES) cell to produce an ES cell that expresses the chimeric transferrin receptor polypeptide. The embryonic stem cell may then be developed into a progeny cell or a non-human transgenic animal whose genome comprises the nucleic acid encoding the chimeric transferrin receptor polypeptide. In some embodiments, the ES cell is introduced into blastocysts and transferred into pseudo pregnant females. In some cases, a founder male harboring the transgene can be selected and bred to wild-type females to generate F1 heterozygous mice. Homozygous non-human animals can be subsequently generated from breeding of F1 generation heterozygous non-human animals. Methods for culturing ES cells and introducing nucleotide sequences to target the genome of an ES cell to produce a transgenic animal are well known, for example, as described in Ramirez-Solis et al., "Gene targeting in mouse embryonic stem cells," *Methods Enzymol.*, 225:855-878 (1993); and US Pat. Pub. No. 2013/0318643, the disclosures of which are incorporated by reference in their entirety. In some embodiments, embryonic stem cells from a transgenic animal that has a chimeric TfR of the present invention can be used as a source to provide progeny of the transgenic animal.

[0065] In some embodiments, the method of knocking-in is carried out in single-cell non-human animal. In one illustrative embodiment, sgRNAs, Cas9, and a donor polynucleotide comprising the apical domain coding sequence that is at least 80% identical to SEQ ID NO:1, where the coding sequence being flanked by a left homology arm and a right homology arm, are introduced into single cell embryos via pronuclear microinjection. The recipient embryos are then transferred to pseudo pregnant females. The sgRNAs form a complex with the Cas9 protein, which then targets the coding sequence of the apical domain of the transferrin receptor in the non-human animal embryos. As a result, the non-human animal transferrin receptor apical domain is cleaved and replaced with the transferrin receptor apical domain coding sequence from the donor polynucleotide. In some cases, a founder male harboring the transgene can be selected and bred to wild-type females to generate F1 heterozygous mice. Homozygous non-human animals can be subsequently generated from breeding of F1 generation heterozygous non-human animals. The transgenic animals disclosed herein can be a rodent, for example, a mouse or a rat.

[0066] In one illustrative embodiment, in part due to the fact that the non-human transgenic animal, e.g., a non-primate mammal, retains introns and the transferrin binding domain of the native TfR, the transgenic animals generated by knocking-in the apical domain that has an amino acid sequence at least 80% identical to SEQ ID NO:1 are generally healthy and demonstrate physiological conditions that are similar to those of wild-type mice of the same species. In one embodiment, all introns outside the apical domain of the TfR are retained. For example, the TfR expression levels are similar to a wild-type animal of the same species; the expression level in the transgenic mouse is no more than 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% less than that of the wild-type mouse or is no more than 10%, 20%, 30%, 40%, 50%, 75%, 100%, 150%, 200%, 300%, or 500% greater than that of the wild-type mouse. The red blood cell count, the level of hemoglobin, and/or hematocrit level are also similar to those in wild-type animals of the same species; the difference is no greater than 50%, e.g., no greater than 40%, no greater than 30%, no greater than 20%, or no greater than 10%. In typical embodiments, transgenic animals in accordance with the invention retain selective BBB transport that enables import of nutrients and proteins and retain the ability to protect CNS from toxins; the presence of transgene doesn't interfere with transferrin binding or FcRn binding to antibodies that bind to the apical domain, described below. Typically, TfR-mediated cellular trafficking in the transgenic animal is also similar to those wild-type animals. The transgenic animals in accordance with the invention are more relevant as a model for pharmacokinetic or pharmacodynamic studies of human BBB-penetrating drugs than the wild-type mice that lack the human TfR entirely or than transgenic animal models that express the entire huTfR extracellular domain (e.g., express the entire huTfR protein).

[0067] Although the present invention is illustrated in mouse as shown in the examples, one of ordinary skill in the art would understand that other non-human mammals, for example, rodent, rabbit, bovine, ovine, canine, feline, equine, porcine, camelid, non-human primate, and other mammals, can also be engineered to express the chimeric

TfR in a similar fashion, and these transgenic animals can also be used for applications as disclosed herein.

The Apical Domain Binding Polypeptide

[0068] An “apical domain binding polypeptide” or “ADBP” as used herein refers to a polypeptide that binds to the apical domain having an amino acid sequence at least 80% identical to SEQ ID NO:1. The ADBP can be an antibody or any polypeptide that is capable of binding to the apical domain of the huTfR of the chimeric TfR. In some embodiments, the ADBP is an agent that is to be delivered across the blood-brain barrier. In some embodiments, the ADBP further comprises an effector molecule coupled to it, e.g., by covalent linkage. The effector molecule may be a therapeutic agent, a labelling agent, or diagnostic agent. In certain embodiments, the effector molecule is a polypeptide, such as a therapeutic or diagnostic antibody, or a polypeptide that has an enzymatic activity or inhibitory activity on an enzyme or a signaling molecule. In certain embodiments, the effector molecule comprises a small molecule, RNA, DNA, or protein.

[0069] In some embodiments, the ADBP is a bispecific antibody, with the apical domain binding region being an antibody that recognizes the apical domain and the effector molecule being an antibody that recognizes a different antigen, e.g., an enzyme or a signaling molecule, and the binding of the effector moiety either activates or inhibits the enzyme or the signaling molecule.

Screen for ADBPs that Bind the Chimeric TfR

[0070] The chimeric TfRs disclosed herein can be used to screen for ADBPs that are capable of binding to the TfR. The screening method comprises contacting a candidate ADBP with a chimeric TfR disclosed above and determining the amount of candidate ADBP that binds to the chimeric TfR. In some embodiments, the step of contacting the candidate ADBP with the chimeric TfR comprises contacting the ADBP with a host cell that expresses the chimeric TfR.

[0071] In some cases, the step of contacting the candidate ADBP with the chimeric TfR comprises contacting the ADBP with an endothelium that expresses the chimeric TfR. In some embodiments, the endothelium is a BBB endothelium.

[0072] Interactions between the candidate ADBP and TfR can be measured using methods well known in the art, for example, immunoassays or SPR. In some embodiments, the binding of the candidate ADBP to the chimeric TfR is measured by ELISA, a Biacore™ system, or coimmunoprecipitation.

Screen for ADBPs that can Cross the BBB

[0073] The non-human transgenic animals expressing the chimeric TfR as described above can be used to characterize the ability of ADBP to bind the apical domain of the chimeric TfR and ultimately the ability to cross the BBB.

[0074] Typically, to evaluate the ability of an ADBP to cross BBB, the ADBP is administered to the transgenic animal carrying the chimeric TfR disclosed herein, preferably through intravenous injection. After a period of time, e.g., at least 10 min, at least 20 min, at least 30 min, at least 60 min, at least 90 min, at least 120 min, at least 180 min, or at least 240 min, the transgenic animal is sacrificed and brain tissues are analyzed to determine the presence of the ADBP. The presence of the ADBP can be determined by assaying for the presence of the ADBP and/or an effector molecule joined thereto. In some embodiments, the brain

tissues are perfused with saline, e.g., PBS, and fixed before detection. The presence of the effector molecule in the sections can be detected using standard imaging methods, for example, immunohistochemical or immunofluorescent methods. A positive detection of the effector molecule in the brain tissue indicates that the effector molecule can cross the BBB. In some cases, determining the presence of the ADBP in the brain comprises performing a quantitative immunoassay. The assay for measuring transport across the BBB using the chimeric TfR transgenic mouse is robust and can measure greater than a 10-, 20-, 30-, 40-, or 50-fold improvement in uptake of an ADBP.

[0075] In some embodiments, in addition to using imaging methods or immunoassays to detect the presence of the ADBP in the brain, methods of detecting changes of a substrate of the effector molecule can also be used to evaluate the brain uptake of the effector molecule. In one illustrative embodiment, the ADBP comprises an effector molecule that can inhibit the enzymatic activity of an enzyme in the brain. In some embodiments, the brain uptake of an ADBP, i.e., which reflects its ability of BBB transport, can be measured by assessing enzymatic activity of an enzyme that is modulated by either the ADBP or an effector molecule joined thereto.

[0076] In some embodiments, the brain uptake of candidate ADBPs is measured in the brain. Plasma can also be monitored and the pharmacokinetic profiles evaluated. Following the administration of a candidate effector molecule, an increase in the brain-to-plasma ratio as compared to a non-BBB penetrating molecule indicates that the candidate ADBP can cross the BBB.

[0077] In some cases, a non-human transgenic animal comprising a polynucleotide encoding the chimeric TfR can be crossed with a non-human transgenic animal that has been engineered to show a certain disease phenotype. In some cases, the non-human transgenic animal is a transgenic mouse that can be crossed with various mouse models, for example, an ALS mouse model, such as described in U.S. Pat. No. 8,476,485; an AD mouse model, such as described in U.S. Pat. No. 5,898,094 and U.S. Pat. No. 6,175,057; a TSPO mouse model, such as described in US Pat. Pub. No. 2016/0050895, an autism spectrum disorder (ASD) mouse model, such as described in US Pat. Pub. No. 2014/0041062. The entire content of these aforementioned patents and patent applications are hereby incorporated by reference. In some cases, the hybrid mice produced by such crosses can be used to evaluate both the distribution of an ADBP comprising an effector molecule in the brain as well as the efficacy of the ADBP or effector molecule in treating brain diseases.

KITS

[0078] In some embodiments, kits comprising a chimeric transferrin receptor polynucleotide or polypeptide, or cells that express such polypeptides, as described herein are provided. In some embodiments, the kits are for use in screening for ADBPs as described above.

[0079] In some embodiments, the kit further comprises buffers and vessels that can be used in the assay to detect the binding between the chimeric TfR polypeptide and a candidate ADBP. In some embodiments, the kit further comprises instructional materials containing directions (i.e., protocols) for the practice of the methods described herein (e.g., instructions for using the kit for administering a composition

across the blood-brain barrier). While the instructional materials typically comprise written or printed materials, they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to, electronic storage media (e.g., magnetic discs, tapes, cartridges, chips), optical media (e.g., CD-ROM), and the like. Such media may include addresses to internet sites that provide such instructional materials.

EXAMPLES

[0080] The following examples are for illustrative purposes only and should not be interpreted as limitations of the claimed invention. There are a variety of alternative techniques and procedures available to those of skill in the art which would similarly permit one to successfully perform the intended invention.

Example 1: HUTFR Mouse Generation and Characterization

[0081] Methods for generating knock-in/knock-out mice have been published in the literature and are well known to those with skill in the art. In brief, C57Bl6 mice were used to generate a knock-in of the human apical TfR mouse line via pronuclear microinjection into single cell embryos, followed by embryo transfer to pseudo pregnant females. Specifically, Cas9, sgRNAs SEQ ID NOs:10-11, and a donor DNA, SEQ ID NO:4, were introduced into the embryos. The donor DNA comprised the human apical domain coding sequence that has been codon optimized for expression in mouse, SEQ ID NO:2. The apical domain coding sequence was flanked with a left (nucleotides 1-817 of SEQ ID NO:4) and right homology arm (nucleotides 1523-2329 of SEQ ID NO:4). The donor sequence was designed in this manner such that the apical domain was to be inserted after the fourth mouse exon, and was immediately flanked at the 3' end by the ninth mouse exon. A founder male from the progeny of the female that received the embryos was bred to wild-type females to generate F1 heterozygous mice. Homozygous mice were subsequently generated from breeding of F1 generation heterozygous mice.

Example 2: Generation of Tool Antibodies for Monitoring Antibody Brain Uptake

[0082] Tool antibodies targeting human TfR or human/mouse BACE1 were generated by transforming Expi293 or ExpiCHO cells with expression plasmids containing DNA encoding the heavy and light chains and using protocols familiar to those with skill in the art. Bispecific antibodies were generated using the “knobs-into-holes” technology; knob and hole half antibodies were separately expressed and then joined using published methods. Antibodies were purified first with Protein A and then by size-exclusion chromatography. The antibodies generated for these studies were as follows:

anti-TfR: human IgG1 antibody that binds to human TfR apical domain.

anti-BACE1: human IgG1 antibody that binds to human BACE1 and cross-reacts with mouse BACE1. This antibody inhibits the enzymatic activity of BACE1.

anti-TfR/BACE1: human IgG1 knobs-into-holes bispecific antibody that binds to human TfR apical domain, as well as human and mouse BACE1. The knob half-antibody has the

variable domain from the anti-BACE1 antibody; the hole half-antibody has the variable domain from the anti-TfR antibody.

Example 3: Blood Analysis of HUTFR^{apical+/-} and HUTFR^{apical+/+} Mice

[0083] Blood was collected from wild-type C57Bl6, huTfR^{apical+/-}, and huTfR^{apical+/+} mice (n=3/group) and a standard complete blood count (CBC) analysis was performed. No genotype-specific differences were observed in any red blood cell parameters, including total red blood cells, hemoglobin, and hematocrit levels (FIG. 1).

Example 4: Brain Localization of TfR-Targeted Antibodies in HUTFR^{apical+/-} and HUTFR^{apical+/+} Mice

[0084] In this example, the anti-TfR antibody was generated to evaluate brain uptake of TfR-targeted therapeutics in huTfR^{apical+/-} mice. huTfR^{apical+/-} mice or wild-type C57Bl6 were intravenously injected with 5 mg/kg of the anti-TfR antibody. After 1 hour, the mice were sacrificed and perfused with PBS. Hemi-brains were drop fixed in 4% PFA overnight followed by 30% sucrose preservation. Sagittal brain sections (35 μ m) were cut using a microtome, blocked in 5% BSA+0.3% Triton X-100, followed by fluorescent secondary staining with Alexa488 anti-huIgG1 (1:500). Brain images were taken using a Zeiss widefield microscope with a 20 $\times$ objective. Significant vascular staining was observed in the huTfR^{apical+/-}, indicating robust binding of human apical-specific anti-TfR on brain endothelial cells at the BBB where TfR is highly expressed (FIG. 2). In contrast, very little staining was observed in the wild-type mice.

[0085] To confirm the TfR-specific BBB transport, the anti-BACE1 antibody and the anti-TfR/BACE1 bispecific antibody were tested using a similar approach as described above. huTfR^{apical+/+} mice were intravenously injected with 50 mg/kg of either antibody. After 24 hours, mice were perfused with PBS, and hemi-brains were processed and stained as described above for the huTfR^{apical+/-} mice. Broad brain parenchymal staining was observed for anti-TfR/BACE1, while no staining was observed for anti-BACE1, indicating that a TfR-apical domain-binding polypeptide is required for BBB transcytosis in these mice (FIG. 3).

Example 5: Brain and Plasma PK/PD of Antibodies in HUTFR^{apical+/+} Mice

[0086] In this example, huTfR^{apical+/+} mice were intravenously injected with 50 mg/kg of either the anti-BACE1 antibody or the anti-TfR/BACE1 bispecific antibody. After 24 hours, blood was collected via cardiac puncture, and the mice were perfused with PBS. Brain tissue was homogenized in 10 $\times$ tissue weight of lysis buffer containing 1% NP-40 in PBS. Blood was collected in EDTA tubes to prevent clotting and spun at 14000 rpm for 7 minutes to isolate plasma. Antibody concentrations in mouse plasma and brain lysates were quantified using a generic human IgG assay (MSD human IgG kit #K150JLD) following the manufacturer's instructions. Briefly, pre-coated plates were blocked for 30 min with MSD Blocker A. Plasma samples were diluted 1:10,000 using a Hamilton Nimbus liquid handler and added in duplicate to the blocked plates. Brain

samples were homogenized in 1% NP40 lysis buffer and lysates diluted 1:10 for PK analysis. Dosing solutions were also analyzed on the same plate to confirm the correct dosage. The standard curve, 0.78-200 ng/mL IgG, was fit using a four-parameter logistic regression.

[0087] After 24 hours, the plasma levels of anti-TfR/BACE1 were lower than the levels for anti-BACE1, likely due to clearance of this antibody via binding to peripherally-expressed huTfR^{apical} (FIG. 4A). In brain, a ~28-fold increase in the concentration of anti-TfR/BACE1 compared to anti-BACE1 was observed (FIG. 4B). The significant accumulation of the anti-TfR/BACE1 is due to TfR-mediated transcytosis at the BBB, this result validates the huTfR^{apical/+} mice as a tool for measuring BBB uptake for human TfR-apical domain-binding polypeptides.

[0088] BACE1 inhibition of amyloid precursor protein (APP) cleavage was used as a pharmacodynamic readout of antibody activity in plasma and brain. Brain tissue was homogenized in 10× tissue weight of 5M guanidine-HCl and then diluted 1:10 in 0.25% casein buffer in PBS. Mouse A1340 levels in plasma and brain lysate were measured using a sandwich ELISA. A 384-well MaxiSorp plate was coated overnight with a polyclonal capture antibody specific for the C-terminus of the A1340 peptide (Millipore #ABN240). Casein-diluted guanidine brain lysates were further diluted 1:2 on the ELISA plate and added concurrently with the detection antibody, biotinylated M3.2. Plasma was analyzed at a 1:5 dilution. Samples were incubated overnight at 4° C. prior to addition of streptavidin-HRP followed by TMB substrate. The standard curve, 0.78-50 pg/mL msAβ40, was fit using a four-parameter logistic regression.

[0089] Compared to anti-BACE1, anti-TfR/BACE1 treatment resulted in an increased reduction of A-beta in huTfR^{apical/+} mice, indicating BACE1 target engagement in the

brain is achieved with anti-TfR/BACE1 (FIG. 4C). Plasma A-beta was reduced to a similar extent for both anti-TfR/BACE1 and anti-BACE1, as compared to untreated wild-type mice (FIG. 4D). These data support the use of the huTfR^{apical/+} mice in target engagement studies that require human TfR-mediated brain uptake, particularly for evaluation of human TfR-apical domain-binding polypeptides.

Example 6: Expression of TfR in HUTFR^{apical/+} Mice

[0090] Brain and various peripheral tissues were isolated from wild-type and huTfR^{apical/+} mice in order to determine whether TfR expression levels are altered in the huTfR^{apical/+} mice. Brain, liver, lung, and kidney were taken from mice following perfusion with PBS. Tissues were homogenized in 10× tissue weight of lysis buffer containing 1% NP-40 in PBS. Samples were run on western blots and TfR expression levels were determined using a TfR antibody recognizing the intracellular portion of TfR and thus cross-reactive to both wild-type and huTfR^{apical/+} (1:2000; ThermoFisher #13-6800). Quantification of TfR expression was expressed as a ratio to actin (1:5000; Abcam 8227). FIGS. 5A-5D show that TfR expression in the huTfR^{apical/+} mice is very similar to wild-type mice in brain (FIG. 5A), liver (FIG. 5B), kidney (FIG. 5C), and lung (FIG. 5D).

[0091] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, sequence accession numbers, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

TABLE OF ILUSTRATIVE SEQUENCES

SEQ ID NO: 1: Protein sequence of human apical domain insert
AQN~~SV~~I~~VD~~K~~NG~~R~~LV~~Y~~LV~~EN~~PG~~GY~~V~~AYS~~KA~~AT~~VT~~G~~KL~~V~~H~~AN~~FG~~T~~KK~~DF~~ED~~LY~~TP~~V~~NG~~SI~~IV~~RA
G~~K~~IT~~FA~~E~~K~~V~~AN~~ES~~LN~~AIG~~VL~~IY~~MD~~Q~~TK~~FP~~IV~~NAEL~~S~~FF~~GH~~AHL~~GT~~GD~~P~~Y~~TP~~GF~~PS~~FN~~HT~~Q~~PP~~
SR~~SS~~SL~~PN~~IP~~VQ~~TI~~S~~RAAAE~~KL~~FG~~N~~ME~~GD~~CP~~SD~~WK~~TD~~ST~~CR~~M~~VT~~SE~~SK~~N~~V~~KL~~TV~~SN

SEQ ID NO: 2: DNA sequence of human apical domain insert
G~~CT~~C~~AG~~A~~ACT~~C~~CG~~T~~G~~A~~T~~C~~AT~~C~~GT~~G~~G~~A~~T~~A~~GA~~A~~CG~~C~~CG~~C~~T~~G~~GT~~G~~T~~A~~CT~~G~~GT~~G~~G~~A~~ACC~~T~~G~~
G~~CG~~G~~A~~T~~AC~~G~~T~~G~~GC~~T~~T~~A~~CT~~T~~A~~G~~CC~~G~~C~~T~~AC~~C~~GT~~G~~AC~~G~~CA~~A~~GC~~T~~G~~G~~T~~G~~C~~A~~CG~~C~~CA~~A~~CT~~T~~C~~G~~G~~
A~~ACC~~A~~AG~~A~~AG~~C~~ATT~~T~~G~~A~~GG~~A~~CT~~G~~T~~A~~C~~A~~C~~A~~CG~~A~~GT~~G~~A~~A~~CG~~G~~CT~~C~~T~~A~~TC~~G~~T~~A~~TC~~G~~T~~G~~C~~G~~CT~~
G~~GA~~A~~AG~~A~~T~~C~~AC~~CT~~T~~C~~G~~C~~G~~A~~AG~~G~~T~~G~~CT~~A~~AC~~G~~CC~~G~~AG~~C~~CT~~G~~A~~A~~CG~~C~~CA~~T~~C~~G~~G~~C~~T~~G~~CT~~G~~A~~
T~~CT~~A~~C~~A~~T~~G~~G~~A~~T~~C~~AG~~A~~CA~~A~~AG~~T~~T~~T~~CC~~A~~T~~C~~GT~~G~~A~~A~~CG~~C~~T~~G~~AG~~C~~T~~G~~T~~C~~T~~T~~T~~C~~TT~~G~~G~~A~~C~~A~~CG~~C~~T~~A
C~~CT~~G~~G~~C~~AC~~C~~CG~~A~~CC~~A~~T~~A~~C~~A~~C~~C~~CG~~A~~T~~T~~CC~~C~~T~~A~~G~~C~~T~~T~~T~~A~~CC~~A~~C~~A~~CC~~A~~GT~~T~~CC~~C~~CC~~C~~T~~
T~~CC~~A~~GG~~T~~C~~T~~A~~G~~CG~~A~~CT~~G~~CC~~A~~AA~~C~~AT~~C~~CC~~C~~GT~~G~~C~~A~~CA~~A~~T~~C~~AG~~C~~AG~~A~~CC~~G~~CT~~G~~CC~~G~~A~~A~~GC~~
T~~GT~~T~~T~~G~~CA~~A~~C~~A~~T~~G~~G~~A~~GG~~A~~GA~~C~~T~~G~~CC~~C~~T~~C~~CG~~A~~T~~T~~G~~A~~AG~~A~~CC~~G~~ACT~~T~~A~~C~~AT~~G~~T~~A~~G~~A~~T~~G~~GT~~
G~~AC~~C~~T~~C~~CG~~A~~G~~T~~CA~~A~~AA~~A~~T~~G~~T~~C~~AA~~A~~CT~~A~~CC~~G~~CT~~G~~T~~C~~CA~~T

SEQ ID NO: 3: Chimeric TfR sequence expressed in transgenic mouse (The italicized portion represents the cytoplasmic domain, the bolded portion represents the transmembrane domain, the portion in grey represents the extracellular domain, and the bold and underlined portion represents the apical domain)
MMDQARS~~AF~~SNLFGGEPLSYTRFSLARQVDGNSH~~VE~~MKLA~~AD~~EENADNNMKASVRKPKRFNG

RLCFAAIALVIFFLIGFM~~SG~~LYG~~Y~~K~~R~~VEQ~~KE~~ECV~~KL~~AETET~~DK~~SEIMETEDVPTSSRLI~~W~~AD

LKTL~~SE~~KLNSIEFADTIKQLSQNTYTPREAGSQKDESLAYIENQPH~~EF~~K~~PS~~KYWRDEHYVKI

QVKSSAQN~~SV~~I~~VD~~K~~NG~~R~~LV~~Y~~LV~~EN~~PG~~GY~~V~~AYS~~KA~~AT~~VT~~G~~KL~~V~~H~~AN~~FG~~T~~KK~~DF~~ED~~LY~~TP~~V~~NG~~SI

-continued

TABLE OF ILUSTRATIVE SEQUENCES

VIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFFIVNAELSF~~FGHAHLGTGDPYTPGFPSFNH~~

TQFPSPRSSGLPNIPVQ~~TISR~~AAAEKLF~~GNMEGDCPSDKIDSTCRMVTSESKNVKLTVSNVLK~~

ERRILNIPGVIKGYEEDRYVVG~~AORDALGAGVAAKSSVGTGLLLKLAQVPSDMISKDGFRPS~~

RSIIFASWTAGDFGAVGATEWLEGYLSSHLK~~AFYINLDKVVLGTSNFKVSASPLLYTLMGKI~~

MODVKHPVDGKSLYRDSNWISKVEKLSFDN~~AAYPPLAYSGIPAVSPCFCEADADYPYLGTRLDY~~

EALTKQV~~PQLNQMVRTAAEVAGOLIIKLTHDVELNLDYEMYNKLLSFMKDLNQPKTDIEDMGL~~

SLOWLYSARGDYFRATSR~~LTTDPHNAEKTNRFFVMREINDRIMKVEYHFLSPYVSPRESPRHF~~

WGS~~GSHTLSALVENLKLQKNITAFNETLFRNQLALATWTIOGVANALSGDIWNIDNEF~~

SEQ ID NO: 4: Sequence of full donor DNA (Left homology arm: 1-917; Right homology arm: 1523-2329; Human Apical Domain: 941-1492; Codon Optimized Sequence: 821-1522)

CTATACAGATATATAAGGATGGGGCTTTTTTTTTTTAATTTTAAAAAAGATTTGTTTATTATT
ATATGTAAGTACACTGTAGCTGTCTTCAGACACTCCAGAAGAGGGCATCAGATCTCATTACAGA
TGGTTGTGAGCTACCATGTGGTCACTGGGATTTGAACTCAGGACCTTCAGAAGAGCAGTCAGTG
CTCTTAAGTATAAGTTAATAAAGTTAACTGATAAGGTAATAAAGGTCCTTATGAAAAGGG
TTCAGACCCAAAGAGTCAGAGATCCACAGGTTGAGAACCTCCTGCCCTAAATCTTGTTGCTCTC
CTTATTCAAGACCACTCCTGTTGCAGTTGCTCTTAAGCATGAGTATGCTCCCTCTGAAAGTCT
CCATAGCAGCCATCTCTCCAGCCCCAGAGTGAGGCTTTTAAAGGAATCTTCATGATAAATAGAA
TTTTTAAAAAGTAAGTAACTGAAGTAACTTAAGGTGTTAAGGTACATTTTATCCCTCAGTAACTGG
TTAATCTAGCAGTTTGGAGTCATACTTCATTTATCTTGACTTTGAAGAGTAAGATATTTAAACA
ATTTGCTTGATCCTTGAAGTAAGTATTAAATAGACATTTAATGCAGACTTTTTTGTGTTGAC
TGGTGGTGTGACGCTGCTCAATCCAAGTACTCATGGGAGGCAGAGGCAGGAGGATCTCTCTCT
AGACCAGCCTGGTCTATAGAGCAAGTCCAGGACAGCCAGGCTACACAGAAACCTTGTTTCAA
ACAAGACTTTTATCCTTCCAGGCAGCTGAGCCAGAATACATACACTCCTAGGGAAGCTGGTTCA
CAGAAGGACGAATCCCTGGCATACTACATCGAGAATCAGTTTCACGAGTTCAAGTTTAGCAAAG
TCTGGAGAGATGAGCACTACGTGAAGATCCAGGTGAAGAGCTCCGCTCAGAAGTCCGTGATCAT
CGTGGATAAGAACGGCCGGCTGGTGTACCTGGTGGAGAACCCTGGCGGATACGTGGCTTACTCT
AAGGCCGCTACCGTGACAGGCAAGCTGGTGCACGCCAAGTTCGGAACCAAGAAGGACTTTGAGG
ATCTGTACACACCAAGTGAACGGCTCTATCGTGATCGTGCAGCTGGAAAGATCACCTTCGCCGA
GAAGGTGGCTAACGCCGAGAGCTGAACGCCATCGGCGTGCTGATCTACATGGATCAGACAAAG
TTTCCCATCGTGAACGCTGAGCTGTCTTTCTTTGGACAGCTCACCTGGGCACCGGAGACCCAT
ACACACCCGGATTCCCTAGCTTTAACCACACCCAGTTCCCTCCCTTCAGGCTAGCGGACTGCC
AAACATCCCGTGACAGCAATCAGCAGAGCCGCTGCCGAGAAGCTGTTTGGCAACATGGAGGGA
GACTGCCCTCCGATTGGAAGACCGACTCTACATGTAGGATGGTGACCTCCGAGTCAAAAAATG
TCAAAGTCAACCGTGCTCAATGTGCTGAAAGAACGACGCATCCTGAATATCTTTGGAGTTATTAA
AGGTTATGAGGAACCAAGTAAAGACCTGCTTTGTACTTTTCACTTTACTGTTTGTGCTTACTGT
AGATAGGTCTAGTGCAGGAAGGAGAAGGATGCTAGCTTGGCATGAAGTGTATATCTTGTGTTGT
CCTAATGTGAACCTTTGTAAATATATGTGTATATAACATAATATGGCCATGTAAGTGTATGGAG
AGGCCAGAGTTAAGTATTAAATATCTTTCTGTAATCATTTAAATTTTACATATGAAGGTGAGT
GAACAGATTGAAGGAGTTTGTCCAGGTGGGACTTGGATCTAAATTTTACAAATGCCTGGCAG
CAAACACCTTTTAAATCACTGAGCTGTCTCCCAATAAAGTGAATGTGATATCAGCTTGTGG
ATAATTTTTTTTGTGCTTTGATAAGTGGTTTTCTTACAGGATCACATACAGTTCTGTCCAT
AGCATTAAACAACATAACTGTGCATGAGTAGATTAAATGTGAGGACACATCCAACAGTCACAT
TTATTAATAGGACAAAAAGTTGGACCTTATATGTAGCACACCTATAATCCAGTGTAGGAAGA
TCCGGGTAGGAGATCCTTAGTTCGGTGTACTTAGTGAGGTTTGTTCAAAAACAAAAGCTA
TGATGGTGTGTTGCCTTTTTCTTTTAGACCGTTATGTGTAGTAGGAGCCAGAGAGACGCTT
TGGGTGCTGTGTTGCGGCGAAGTCCAGTGTGGGAACAGGTCTTCTGTTGAACTTGCCCAAGT
ATTCTCAGATATGATTTCAAAAGGT

SEQ ID NO: 5 Mouse TfR protein sequence (Uniprot Q62351) (The italicized portion represents the cytoplasmic domain, the bolded portion represents the transmembrane domain, the portion in grey represents the extracellular domain, and the bold and underlined portion represents the apical domain)

MMDQARS~~AFSNLPGGEPLSYTRFSLARQVDGDN~~SHVEMKLA~~AD~~EEENADNNMKASVRKPKRFNG

RLCFAAIALVIFFLIGFMSGYLGYCKRVEQKEECVKLAETEETDKSETMETEDVPTSSRLYWAD

LKTL~~LSEK~~LSIEFADTIKQLSONTYTFREAGSQ~~DR~~SLAYYIENQFHEPKFSKVRDEHYVKI

-continued

TABLE OF ILUSTRATIVE SEQUENCES

QVKSSIGQNMVTIVQSNENLDPVESPEGYVAFSKPTEVSGKLVHANFGTKKDFEELSYSVNGSL

VIVRAGEITFAEKVANAQSFNAIGVLIYMDKNKFPVVEADLALFGHAHLGTGDPYTPGFPSPFNH

TQFPSPQSSGLPNIPVQTISRAAAELFGKMEGSCPARWNIDSSCKLELSONQNVKLIVKNVLK

ERRILNIPGVIKGYEEDRYVVVGAQRDALGAGVAAKSSVGTGLLLKLAQVFSNMISKDGFRRPS

RSIIFASWTAGDFGAVGATEWLEGYLSSLHLKAPTYYINLDKVVLTGSNFKVSASPLLYTLMGKI

MODVKHPVDGKSLYRDSNWISKVEKLSFDNAAYPFLAYSGIPAVSFCFCEDADYPYLGTRLDTY

EALTQKVPOLNQMVRTAEVAGQLIIKLTHDVELNLDYEMYNKLLSFMKDLNQFKTDIRDML

SLQWLYSARGDYFRATSRLLTTDFHNAEKTNRFFVMREINDRIMKVEYHFLSPYVSPRESPPRHIF

WGSGSHTLSALVENLKLKQKNITAFNETLFFNQALATWTIQGVANALSGDIWNIDNEF

SEQ ID NO: 6: Human TfR protein sequence (Uniprot P02786) (The italicized portion represents the cytoplasmic domain, the bolded portion represents the transmembrane domain, the portion in grey represents the extracellular domain, and the bold and underlined portion represents the apical domain)

MMDQARSAPSNLFGGEPLSYTRPSLARQVDGGNSHVEMKLAVIDEENADNNTKANVTKPKRCSG

SICYGTIAVIVFFLIGFMIGYLGYCKQVEPKTECERLAGTESPVREEPGEDFPAARRLYWDLLK

RKLSEKLDSTDFGTIKLNNENSYVPREAGSOKDENLALYVENQPREFKLSKVMRDQHFVKIQV

KDSAQNSVIVDKNGRLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLYTPVNGSIV

VRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSFPGHAHLGTGDPYTPGFPSPFNHTQ

FPSPRSSGLPNIPVQTISRAAAELFGNMEGDCPSDWKTDSTCRMVTSSEKNVKLTVSNVLKEI

KILNIPGVIRGEVEPDHYVVVGAQRDANGPGAAGSGVGTALLLKLQMFSDMVLKDCGQPSRSI

IFASWSAGDFGAVGATEWLEGYLSSLHLKAPTYYINLDKAVLTGSNFKVSASPLLYTLIEKTMQN

VKHPVTGQFLYQDSNMAKVEKLTLDNAAPPFLAYSGIPAVSFCFCEDTDYPYLGTMTDITYKEL

IERIPELNKVARAAAEVAGQFVIKLTHDVELNLDYERYNSQLLSFVRDLNQYRADIKEMGLSLQ

WLYSARGDFFRATSRLLTTDFGNAEKTDRFVMKKLNDRVMRVEYHFLSPYVSPKESPPRHVFWGS

GSHTLPALLENLKLKQKNNGAFNETLFFNQALATWTIQGAANALSGDVWDIDNEF

SEQ ID NO: 7: The apical domain of macaca mulatta (rhesus monkey) TfR (NCBI Reference Sequence NP_001244232.1); it has 95% identity to the apical domain of the native human TfR
AQN^SVIIIVDKNGGLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLDSPVNGSIVIVRA
GKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSFPGHAHLGTGDPYTPGFPSPFNHTQFPP
SQSSGLPNIPVQTISRAAAELFGNMEGDCPSDWKTDSTCRMVTSSENKSVKLTVSN

SEQ ID NO: 8: The apical domain of chimpanzee TfR (NCBI Reference Sequence XP_003310238.1); it is 98% identical to the apical domain of the native human TfR
AQN^SVIIIVDKNGSLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLHTPVNGSIVIVRA
GKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSFPGHAHLGTGDPYTPGFPSPFNHTQFPP
SRSSGLPNIPVQTISRAAAELFGNMEGDCPSDWKTDSTCRMVTSSENKSVKLTVSN

-continued

TABLE OF ILLUSTRATIVE SEQUENCES

SEQ ID NO: 9: The apical domain of *Mascaca fascicularis* (cynomolgous monkey) Tfr (NCBI Reference Sequence XP_005545315); it is 96% identical to the apical domain of the native human Tfr
 AQNSVVIIVDKNGGLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLDSPVNGSIVIVRA
 GKITFAEKVANAESLNAIGVLIYMDQTKFPPIVKADLSFFGHAHLGTGDPYTPGPPSFNHTQFPP
 SQSSGLPNIPVQTISRAAAELFGNMEGDCPSDWKTDSTCKMVTSENKSVKLTVSN

SEQ ID NO: 10
 GAATACATACACTCCTCGTGAGG

SEQ ID NO: 11
 AGAAGAATACTTAACATCTTTGG

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1
 <211> LENGTH: 184
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Ala Gln Asn Ser Val Ile Ile Val Asp Lys Asn Gly Arg Leu Val Tyr
 1 5 10 15

Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys Ala Ala Thr
 20 25 30

Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys Lys Asp Phe
 35 40 45

Glu Asp Leu Tyr Thr Pro Val Asn Gly Ser Ile Val Ile Val Arg Ala
 50 55 60

Gly Lys Ile Thr Phe Ala Glu Lys Val Ala Asn Ala Glu Ser Leu Asn
 65 70 75 80

Ala Ile Gly Val Leu Ile Tyr Met Asp Gln Thr Lys Phe Pro Ile Val
 85 90 95

Asn Ala Glu Leu Ser Phe Phe Gly His Ala His Leu Gly Thr Gly Asp
 100 105 110

Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His Thr Gln Phe Pro Pro
 115 120 125

Ser Arg Ser Ser Gly Leu Pro Asn Ile Pro Val Gln Thr Ile Ser Arg
 130 135 140

Ala Ala Ala Glu Lys Leu Phe Gly Asn Met Glu Gly Asp Cys Pro Ser
 145 150 155 160

Asp Trp Lys Thr Asp Ser Thr Cys Arg Met Val Thr Ser Glu Ser Lys
 165 170 175

Asn Val Lys Leu Thr Val Ser Asn
 180

<210> SEQ ID NO 2
 <211> LENGTH: 552
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

gctcagaact ccgtgatcat cgtggataag aacggccgce tgggtgtacct ggtggagaac 60

-continued

```

cctggcgcat acgtggctta ctctaaggcc gctaccgtga caggcaagct ggtgcacgcc 120
aacttcggaa ccaagaagga ctttgaggat ctgtacacac cagtgaacgg ctctatcgtg 180
atcgtgcgcg ctggaagat caccttcgcc gagaaggtgg ctaacgccga gagcctgaac 240
gccatcggcg tgctgatcta catggatcag acaaagtctt ccatcgtgaa cgctgagctg 300
tctttctttg gacacgctca cctgggcacc ggagacccat acacacccgg attccctage 360
tttaaccaca ccagttccc cccttcagg tctagcggac tgccaaacat ccccgctgcag 420
acaatcagca gagccgctgc cgagaagctg ttgggaaca tggagggaga ctgccctcc 480
gattggaaga ccgactctac atgtaggatg gtgacctcg agtcaaaaaa tgtcaaaactc 540
accgtgtcca at 552

```

```

<210> SEQ ID NO 3
<211> LENGTH: 763
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic chimeric transferrin receptor (TfR)
      polypeptide sequence
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(67)
<223> OTHER INFORMATION: Cytoplasmic Domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (68)..(88)
<223> OTHER INFORMATION: Transmembrane Domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (89)..(763)
<223> OTHER INFORMATION: Extracellular Domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (198)..(381)
<223> OTHER INFORMATION: Apical Domain

```

```

<400> SEQUENCE: 3

```

```

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu
1           5           10          15
Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp
20          25          30
Asn Ser His Val Glu Met Lys Leu Ala Ala Asp Glu Glu Glu Asn Ala
35          40          45
Asp Asn Asn Met Lys Ala Ser Val Arg Lys Pro Lys Arg Phe Asn Gly
50          55          60
Arg Leu Cys Phe Ala Ala Ile Ala Leu Val Ile Phe Phe Leu Ile Gly
65          70          75          80
Phe Met Ser Gly Tyr Leu Gly Tyr Cys Lys Arg Val Glu Gln Lys Glu
85          90          95
Glu Cys Val Lys Leu Ala Glu Thr Glu Glu Thr Asp Lys Ser Glu Thr
100         105         110
Met Glu Thr Glu Asp Val Pro Thr Ser Ser Arg Leu Tyr Trp Ala Asp
115        120        125
Leu Lys Thr Leu Leu Ser Glu Lys Leu Asn Ser Ile Glu Phe Ala Asp
130        135        140
Thr Ile Lys Gln Leu Ser Gln Asn Thr Tyr Thr Pro Arg Glu Ala Gly
145        150        155        160
Ser Gln Lys Asp Glu Ser Leu Ala Tyr Tyr Ile Glu Asn Gln Phe His

```

-continued

165							170							175						
Glu	Phe	Lys	Phe	Ser	Lys	Val	Trp	Arg	Asp	Glu	His	Tyr	Val	Lys	Ile					
			180				185						190							
Gln	Val	Lys	Ser	Ser	Ala	Gln	Asn	Ser	Val	Ile	Ile	Val	Asp	Lys	Asn					
			195				200						205							
Gly	Arg	Leu	Val	Tyr	Leu	Val	Glu	Asn	Pro	Gly	Gly	Tyr	Val	Ala	Tyr					
			210				215						220							
Ser	Lys	Ala	Ala	Thr	Val	Thr	Gly	Lys	Leu	Val	His	Ala	Asn	Phe	Gly					
			230						235			240								
Thr	Lys	Lys	Asp	Phe	Glu	Asp	Leu	Tyr	Thr	Pro	Val	Asn	Gly	Ser	Ile					
			245						250			255								
Val	Ile	Val	Arg	Ala	Gly	Lys	Ile	Thr	Phe	Ala	Glu	Lys	Val	Ala	Asn					
			260						265			270								
Ala	Glu	Ser	Leu	Asn	Ala	Ile	Gly	Val	Leu	Ile	Tyr	Met	Asp	Gln	Thr					
			275						280			285								
Lys	Phe	Pro	Ile	Val	Asn	Ala	Glu	Leu	Ser	Phe	Phe	Gly	His	Ala	His					
			290			295			300											
Leu	Gly	Thr	Gly	Asp	Pro	Tyr	Thr	Pro	Gly	Phe	Pro	Ser	Phe	Asn	His					
305			310						315			320								
Thr	Gln	Phe	Pro	Pro	Ser	Arg	Ser	Ser	Gly	Leu	Pro	Asn	Ile	Pro	Val					
			325						330			335								
Gln	Thr	Ile	Ser	Arg	Ala	Ala	Ala	Glu	Lys	Leu	Phe	Gly	Asn	Met	Glu					
			340			345						350								
Gly	Asp	Cys	Pro	Ser	Asp	Trp	Lys	Thr	Asp	Ser	Thr	Cys	Arg	Met	Val					
			355			360						365								
Thr	Ser	Glu	Ser	Lys	Asn	Val	Lys	Leu	Thr	Val	Ser	Asn	Val	Leu	Lys					
			370			375						380								
Glu	Arg	Arg	Ile	Leu	Asn	Ile	Phe	Gly	Val	Ile	Lys	Gly	Tyr	Glu	Glu					
385			390						395			400								
Pro	Asp	Arg	Tyr	Val	Val	Val	Gly	Ala	Gln	Arg	Asp	Ala	Leu	Gly	Ala					
			405						410			415								
Gly	Val	Ala	Ala	Lys	Ser	Ser	Val	Gly	Thr	Gly	Leu	Leu	Leu	Lys	Leu					
			420			425						430								
Ala	Gln	Val	Phe	Ser	Asp	Met	Ile	Ser	Lys	Asp	Gly	Phe	Arg	Pro	Ser					
			435			440						445								
Arg	Ser	Ile	Ile	Phe	Ala	Ser	Trp	Thr	Ala	Gly	Asp	Phe	Gly	Ala	Val					
			450			455						460								
Gly	Ala	Thr	Glu	Trp	Leu	Glu	Gly	Tyr	Leu	Ser	Ser	Leu	His	Leu	Lys					
465			470						475			480								
Ala	Phe	Thr	Tyr	Ile	Asn	Leu	Asp	Lys	Val	Val	Leu	Gly	Thr	Ser	Asn					
			485						490			495								
Phe	Lys	Val	Ser	Ala	Ser	Pro	Leu	Leu	Tyr	Thr	Leu	Met	Gly	Lys	Ile					
			500			505						510								
Met	Gln	Asp	Val	Lys	His	Pro	Val	Asp	Gly	Lys	Ser	Leu	Tyr	Arg	Asp					
			515			520						525								
Ser	Asn	Trp	Ile	Ser	Lys	Val	Glu	Lys	Leu	Ser	Phe	Asp	Asn	Ala	Ala					
			530			535						540								
Tyr	Pro	Phe	Leu	Ala	Tyr	Ser	Gly	Ile	Pro	Ala	Val	Ser	Phe	Cys	Phe					
545			550						555			560								
Cys	Glu	Asp	Ala	Asp	Tyr	Pro	Tyr	Leu	Gly	Thr	Arg	Leu	Asp	Thr	Tyr					
			565						570			575								

-continued

Glu Ala Leu Thr Gln Lys Val Pro Gln Leu Asn Gln Met Val Arg Thr
 580 585 590
 Ala Ala Glu Val Ala Gly Gln Leu Ile Ile Lys Leu Thr His Asp Val
 595 600 605
 Glu Leu Asn Leu Asp Tyr Glu Met Tyr Asn Ser Lys Leu Leu Ser Phe
 610 615 620
 Met Lys Asp Leu Asn Gln Phe Lys Thr Asp Ile Arg Asp Met Gly Leu
 625 630 635 640
 Ser Leu Gln Trp Leu Tyr Ser Ala Arg Gly Asp Tyr Phe Arg Ala Thr
 645 650 655
 Ser Arg Leu Thr Thr Asp Phe His Asn Ala Glu Lys Thr Asn Arg Phe
 660 665 670
 Val Met Arg Glu Ile Asn Asp Arg Ile Met Lys Val Glu Tyr His Phe
 675 680 685
 Leu Ser Pro Tyr Val Ser Pro Arg Glu Ser Pro Phe Arg His Ile Phe
 690 695 700
 Trp Gly Ser Gly Ser His Thr Leu Ser Ala Leu Val Glu Asn Leu Lys
 705 710 715 720
 Leu Arg Gln Lys Asn Ile Thr Ala Phe Asn Glu Thr Leu Phe Arg Asn
 725 730 735
 Gln Leu Ala Leu Ala Thr Trp Thr Ile Gln Gly Val Ala Asn Ala Leu
 740 745 750
 Ser Gly Asp Ile Trp Asn Ile Asp Asn Glu Phe
 755 760

<210> SEQ ID NO 4
 <211> LENGTH: 2329
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic donor DNA sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(817)
 <223> OTHER INFORMATION: left homology arm
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (821)..(1522)
 <223> OTHER INFORMATION: codon-optimized sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (941)..(1492)
 <223> OTHER INFORMATION: human apical domain
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1523)..(2329)
 <223> OTHER INFORMATION: right homology arm

<400> SEQUENCE: 4

ctatacagat atataaggat ggggcttttt ttttttaatt tttaaaaaag atttgtttat	60
tattatatgt aagtacactg tagctgtctt cagacactcc agaagagggc atcagatctc	120
attacagatg gttgtgagct accatgtggt cactgggatt tgaactcagg accttcagaa	180
gagcagtcag tgctcttaac tgataagtta ataataagtt aactgataag gtaataaagg	240
tcccctatga aaaggggttc gacccaaaga gtcagagatc cacagggtga gaacctcctg	300
ccctaaatct tgttgccttc cttattcaag accactcctg ttgcagttgc tcttaagcat	360
gagtatgctc ccttctgaaa gtctccatag cagccatctc tccagcccca gagtgaggct	420

-continued

```

tttaaaggaa tcttcatgat aaatagaatt tttaaaaaag taactgaagt tacttaaggt 480
gttaaggtag atttttattcc ctcaagtaact ggtaaatcta gcagttttga gtcatacttc 540
atttatcttg actttgaaga gtaagatatt aaaacaattt gcttgatcct tgaagtaagt 600
atttaaatag acattttaat gcagactttt tttagttgac tgggtggtgtt gcacgtgggc 660
aatccaagta ctcatgggag gcagaggcag gaggatctct ctctagacca gctggtcta 720
tagagcaagt tccaggacag ccagggtctac acagaaacct tgtttcaaac aagactttta 780
tccttccagg cagctgagcc agaatacata cactcctagg gaagctggtt cacagaagga 840
cgaatccctg gcatactaca tcgagaatca gtttcacgag ttcaagtta gcaaagtctg 900
gagagatgag cactacgtga agatccaggt gaagagctcc gctcagaact ccgtgatcat 960
cgtggataag aacggccggc tgggtgacct ggtggagaac cctggcggat acgtggctta 1020
ctctaaggcc gctaccgtga caggcaagct ggtgcacgcc aacttcggaa ccaagaagga 1080
ctttgaggat ctgtacacac cagtgaacgg ctctatcgtg atcgtgcgcg ctggaaagat 1140
caccttcgcc gagaagggtg ctaacgccga gagcctgaac gccatcggcg tgctgatcta 1200
catggatcag acaaagtttc ccatcgtgaa cgtgagctg tctttctttg gacacgctca 1260
cctgggcacc ggagacccat acacaccgg attccctagc tttaaccaca cccagttccc 1320
cccttcagg tctagcggac tgccaaacat ccccgtcag acaatcagca gagccgctgc 1380
cgagaagctg tttggcaaca tggagggaga ctgcccctcc gattggaaga ccgactctac 1440
atgtaggatg gtgacctccg agtcaaaaaa tgtcaaaact accgtgtcca atgtgtgaa 1500
agaacgacgc atcctgaata tctttggagt tattaaggt tatgaggaac caggtaaaga 1560
cctgctttgt actttttcac tttactgttt tgcttactgt agataggtct agtgcaggaa 1620
ggagaaggat gctagcttgg catgaactgc tatatcttgt ttgtccta atgtgaactttg 1680
taatatatgt gtatataaca cataatatgg ccatgtaagt gtatggagag gccagagtta 1740
agtattaaat atctttctgt aatcatttaa aattttacat atgaaggta gtgaacagat 1800
tgaaggagtt ttgtccaggt gggacttga tctaaattt ttacaatgcc tggcagcaaa 1860
caccttttta atcaactgag ctgtctcccc aaataaagt aatgtgatat cagcttgtgg 1920
ataatttttt tttgttctt tgataagtgg ttttcttaca ggatcacata ccagttctgt 1980
ccatagcatt aaacaacat aactgtcatg cagtagatta atgtgcaggg cacatccaac 2040
agtcacattt attaatagga caaaaagttg gaccttataat gtagcacacc tataattcca 2100
gtgctaggaa gatccgggta ggagatcctt agttcgggtgc tacttagtga gggtttgttt 2160
caaaaaacaa aagctatgat ggtgtgttgc cttttttctt ttagaccgtt atgtttagt 2220
aggagccag agagacgctt tgggtgctgg tgttcggcg aagtcagtg tgggaacagg 2280
tcttctgttg aaacttgccc aagtattctc agatatgatt tcaaaaggt 2329

```

```

<210> SEQ ID NO 5
<211> LENGTH: 763
<212> TYPE: PRT
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(67)
<223> OTHER INFORMATION: cytoplasmic domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN

```

-continued

```

<222> LOCATION: (68)..(88)
<223> OTHER INFORMATION: transmembrane domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (89)..(763)
<223> OTHER INFORMATION: extracellular domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (199)..(381)
<223> OTHER INFORMATION: apical domain

<400> SEQUENCE: 5

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu
1          5          10          15

Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp
          20          25          30

Asn Ser His Val Glu Met Lys Leu Ala Ala Asp Glu Glu Asn Ala
          35          40          45

Asp Asn Asn Met Lys Ala Ser Val Arg Lys Pro Lys Arg Phe Asn Gly
          50          55          60

Arg Leu Cys Phe Ala Ala Ile Ala Leu Val Ile Phe Phe Leu Ile Gly
65          70          75          80

Phe Met Ser Gly Tyr Leu Gly Tyr Cys Lys Arg Val Glu Gln Lys Glu
          85          90          95

Glu Cys Val Lys Leu Ala Glu Thr Glu Glu Thr Asp Lys Ser Glu Thr
          100          105          110

Met Glu Thr Glu Asp Val Pro Thr Ser Ser Arg Leu Tyr Trp Ala Asp
          115          120          125

Leu Lys Thr Leu Leu Ser Glu Lys Leu Asn Ser Ile Glu Phe Ala Asp
          130          135          140

Thr Ile Lys Gln Leu Ser Gln Asn Thr Tyr Thr Pro Arg Glu Ala Gly
145          150          155          160

Ser Gln Lys Asp Glu Ser Leu Ala Tyr Tyr Ile Glu Asn Gln Phe His
          165          170          175

Glu Phe Lys Phe Ser Lys Val Trp Arg Asp Glu His Tyr Val Lys Ile
          180          185          190

Gln Val Lys Ser Ser Ile Gly Gln Asn Met Val Thr Ile Val Gln Ser
          195          200          205

Asn Gly Asn Leu Asp Pro Val Glu Ser Pro Glu Gly Tyr Val Ala Phe
          210          215          220

Ser Lys Pro Thr Glu Val Ser Gly Lys Leu Val His Ala Asn Phe Gly
225          230          235          240

Thr Lys Lys Asp Phe Glu Glu Leu Ser Tyr Ser Val Asn Gly Ser Leu
          245          250          255

Val Ile Val Arg Ala Gly Glu Ile Thr Phe Ala Glu Lys Val Ala Asn
          260          265          270

Ala Gln Ser Phe Asn Ala Ile Gly Val Leu Ile Tyr Met Asp Lys Asn
          275          280          285

Lys Phe Pro Val Val Glu Ala Asp Leu Ala Leu Phe Gly His Ala His
          290          295          300

Leu Gly Thr Gly Asp Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His
305          310          315          320

Thr Gln Phe Pro Pro Ser Gln Ser Ser Gly Leu Pro Asn Ile Pro Val
          325          330          335

```

-continued

Gln	Thr	Ile	Ser	Arg	Ala	Ala	Ala	Glu	Lys	Leu	Phe	Gly	Lys	Met	Glu
			340					345					350		
Gly	Ser	Cys	Pro	Ala	Arg	Trp	Asn	Ile	Asp	Ser	Ser	Cys	Lys	Leu	Glu
		355					360					365			
Leu	Ser	Gln	Asn	Gln	Asn	Val	Lys	Leu	Ile	Val	Lys	Asn	Val	Leu	Lys
		370				375					380				
Glu	Arg	Arg	Ile	Leu	Asn	Ile	Phe	Gly	Val	Ile	Lys	Gly	Tyr	Glu	Glu
385					390					395					400
Pro	Asp	Arg	Tyr	Val	Val	Val	Gly	Ala	Gln	Arg	Asp	Ala	Leu	Gly	Ala
			405						410					415	
Gly	Val	Ala	Ala	Lys	Ser	Ser	Val	Gly	Thr	Gly	Leu	Leu	Leu	Lys	Leu
		420						425					430		
Ala	Gln	Val	Phe	Ser	Asp	Met	Ile	Ser	Lys	Asp	Gly	Phe	Arg	Pro	Ser
		435					440					445			
Arg	Ser	Ile	Ile	Phe	Ala	Ser	Trp	Thr	Ala	Gly	Asp	Phe	Gly	Ala	Val
		450				455					460				
Gly	Ala	Thr	Glu	Trp	Leu	Glu	Gly	Tyr	Leu	Ser	Ser	Leu	His	Leu	Lys
465					470				475						480
Ala	Phe	Thr	Tyr	Ile	Asn	Leu	Asp	Lys	Val	Val	Leu	Gly	Thr	Ser	Asn
			485						490					495	
Phe	Lys	Val	Ser	Ala	Ser	Pro	Leu	Leu	Tyr	Thr	Leu	Met	Gly	Lys	Ile
			500					505					510		
Met	Gln	Asp	Val	Lys	His	Pro	Val	Asp	Gly	Lys	Ser	Leu	Tyr	Arg	Asp
		515					520					525			
Ser	Asn	Trp	Ile	Ser	Lys	Val	Glu	Lys	Leu	Ser	Phe	Asp	Asn	Ala	Ala
		530				535					540				
Tyr	Pro	Phe	Leu	Ala	Tyr	Ser	Gly	Ile	Pro	Ala	Val	Ser	Phe	Cys	Phe
545					550				555						560
Cys	Glu	Asp	Ala	Asp	Tyr	Pro	Tyr	Leu	Gly	Thr	Arg	Leu	Asp	Thr	Tyr
			565						570					575	
Glu	Ala	Leu	Thr	Gln	Lys	Val	Pro	Gln	Leu	Asn	Gln	Met	Val	Arg	Thr
			580					585					590		
Ala	Ala	Glu	Val	Ala	Gly	Gln	Leu	Ile	Ile	Lys	Leu	Thr	His	Asp	Val
		595					600					605			
Glu	Leu	Asn	Leu	Asp	Tyr	Glu	Met	Tyr	Asn	Ser	Lys	Leu	Leu	Ser	Phe
		610				615					620				
Met	Lys	Asp	Leu	Asn	Gln	Phe	Lys	Thr	Asp	Ile	Arg	Asp	Met	Gly	Leu
625					630					635					640
Ser	Leu	Gln	Trp	Leu	Tyr	Ser	Ala	Arg	Gly	Asp	Tyr	Phe	Arg	Ala	Thr
			645						650					655	
Ser	Arg	Leu	Thr	Thr	Asp	Phe	His	Asn	Ala	Glu	Lys	Thr	Asn	Arg	Phe
			660					665					670		
Val	Met	Arg	Glu	Ile	Asn	Asp	Arg	Ile	Met	Lys	Val	Glu	Tyr	His	Phe
		675				680						685			
Leu	Ser	Pro	Tyr	Val	Ser	Pro	Arg	Glu	Ser	Pro	Phe	Arg	His	Ile	Phe
		690				695					700				
Trp	Gly	Ser	Gly	Ser	His	Thr	Leu	Ser	Ala	Leu	Val	Glu	Asn	Leu	Lys
705					710					715					720
Leu	Arg	Gln	Lys	Asn	Ile	Thr	Ala	Phe	Asn	Glu	Thr	Leu	Phe	Arg	Asn
			725						730					735	
Gln	Leu	Ala	Leu	Ala	Thr	Trp	Thr	Ile	Gln	Gly	Val	Ala	Asn	Ala	Leu

-continued

740	745	750
Ser Gly Asp Ile Trp Asn Ile Asp Asn Glu Phe		
755	760	

<210> SEQ ID NO 6
 <211> LENGTH: 760
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(67)
 <223> OTHER INFORMATION: cytoplasmic domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (68)..(89)
 <223> OTHER INFORMATION: transmembrane domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (90)..(760)
 <223> OTHER INFORMATION: extracellular domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (196)..(379)
 <223> OTHER INFORMATION: apical domain

<400> SEQUENCE: 6

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu		
1	5	10 15
Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp		
	20	25 30
Asn Ser His Val Glu Met Lys Leu Ala Val Asp Glu Glu Asn Ala		
	35	40 45
Asp Asn Asn Thr Lys Ala Asn Val Thr Lys Pro Lys Arg Cys Ser Gly		
	50	55 60
Ser Ile Cys Tyr Gly Thr Ile Ala Val Ile Val Phe Phe Leu Ile Gly		
65	70	75 80
Phe Met Ile Gly Tyr Leu Gly Tyr Cys Lys Gly Val Glu Pro Lys Thr		
	85	90 95
Glu Cys Glu Arg Leu Ala Gly Thr Glu Ser Pro Val Arg Glu Glu Pro		
	100	105 110
Gly Glu Asp Phe Pro Ala Ala Arg Arg Leu Tyr Trp Asp Asp Leu Lys		
	115	120 125
Arg Lys Leu Ser Glu Lys Leu Asp Ser Thr Asp Phe Thr Gly Thr Ile		
	130	135 140
Lys Leu Leu Asn Glu Asn Ser Tyr Val Pro Arg Glu Ala Gly Ser Gln		
145	150	155 160
Lys Asp Glu Asn Leu Ala Leu Tyr Val Glu Asn Gln Phe Arg Glu Phe		
	165	170 175
Lys Leu Ser Lys Val Trp Arg Asp Gln His Phe Val Lys Ile Gln Val		
	180	185 190
Lys Asp Ser Ala Gln Asn Ser Val Ile Ile Val Asp Lys Asn Gly Arg		
	195	200 205
Leu Val Tyr Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys		
	210	215 220
Ala Ala Thr Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys		
225	230	235 240
Lys Asp Phe Glu Asp Leu Tyr Thr Pro Val Asn Gly Ser Ile Val Ile		
	245	250 255

-continued

Val	Arg	Ala	Gly	Lys	Ile	Thr	Phe	Ala	Glu	Lys	Val	Ala	Asn	Ala	Glu	
			260					265					270			
Ser	Leu	Asn	Ala	Ile	Gly	Val	Leu	Ile	Tyr	Met	Asp	Gln	Thr	Lys	Phe	
		275					280					285				
Pro	Ile	Val	Asn	Ala	Glu	Leu	Ser	Phe	Phe	Gly	His	Ala	His	Leu	Gly	
		290				295					300					
Thr	Gly	Asp	Pro	Tyr	Thr	Pro	Gly	Phe	Pro	Ser	Phe	Asn	His	Thr	Gln	
305					310					315					320	
Phe	Pro	Pro	Ser	Arg	Ser	Ser	Gly	Leu	Pro	Asn	Ile	Pro	Val	Gln	Thr	
				325					330					335		
Ile	Ser	Arg	Ala	Ala	Ala	Glu	Lys	Leu	Phe	Gly	Asn	Met	Glu	Gly	Asp	
			340					345					350			
Cys	Pro	Ser	Asp	Trp	Lys	Thr	Asp	Ser	Thr	Cys	Arg	Met	Val	Thr	Ser	
		355					360					365				
Glu	Ser	Lys	Asn	Val	Lys	Leu	Thr	Val	Ser	Asn	Val	Leu	Lys	Glu	Ile	
		370				375					380					
Lys	Ile	Leu	Asn	Ile	Phe	Gly	Val	Ile	Lys	Gly	Phe	Val	Glu	Pro	Asp	
385				390						395					400	
His	Tyr	Val	Val	Val	Gly	Ala	Gln	Arg	Asp	Ala	Trp	Gly	Pro	Gly	Ala	
			405					410						415		
Ala	Lys	Ser	Gly	Val	Gly	Thr	Ala	Leu	Leu	Leu	Lys	Leu	Ala	Gln	Met	
			420					425					430			
Phe	Ser	Asp	Met	Val	Leu	Lys	Asp	Gly	Phe	Gln	Pro	Ser	Arg	Ser	Ile	
		435					440					445				
Ile	Phe	Ala	Ser	Trp	Ser	Ala	Gly	Asp	Phe	Gly	Ser	Val	Gly	Ala	Thr	
	450					455					460					
Glu	Trp	Leu	Glu	Gly	Tyr	Leu	Ser	Ser	Leu	His	Leu	Lys	Ala	Phe	Thr	
465					470					475					480	
Tyr	Ile	Asn	Leu	Asp	Lys	Ala	Val	Leu	Gly	Thr	Ser	Asn	Phe	Lys	Val	
			485						490					495		
Ser	Ala	Ser	Pro	Leu	Leu	Tyr	Thr	Leu	Ile	Glu	Lys	Thr	Met	Gln	Asn	
			500					505					510			
Val	Lys	His	Pro	Val	Thr	Gly	Gln	Phe	Leu	Tyr	Gln	Asp	Ser	Asn	Trp	
		515					520					525				
Ala	Ser	Lys	Val	Glu	Lys	Leu	Thr	Leu	Asp	Asn	Ala	Ala	Phe	Pro	Phe	
	530					535					540					
Leu	Ala	Tyr	Ser	Gly	Ile	Pro	Ala	Val	Ser	Phe	Cys	Phe	Cys	Glu	Asp	
545					550					555					560	
Thr	Asp	Tyr	Pro	Tyr	Leu	Gly	Thr	Thr	Met	Asp	Thr	Tyr	Lys	Glu	Leu	
			565						570					575		
Ile	Glu	Arg	Ile	Pro	Glu	Leu	Asn	Lys	Val	Ala	Arg	Ala	Ala	Ala	Glu	
			580					585					590			
Val	Ala	Gly	Gln	Phe	Val	Ile	Lys	Leu	Thr	His	Asp	Val	Glu	Leu	Asn	
		595					600					605				
Leu	Asp	Tyr	Glu	Arg	Tyr	Asn	Ser	Gln	Leu	Leu	Ser	Phe	Val	Arg	Asp	
	610					615					620					
Leu	Asn	Gln	Tyr	Arg	Ala	Asp	Ile	Lys	Glu	Met	Gly	Leu	Ser	Leu	Gln	
625				630						635					640	
Trp	Leu	Tyr	Ser	Ala	Arg	Gly	Asp	Phe	Phe	Arg	Ala	Thr	Ser	Arg	Leu	
				645					650						655	

-continued

Thr	Thr	Asp	Phe	Gly	Asn	Ala	Glu	Lys	Thr	Asp	Arg	Phe	Val	Met	Lys
			660					665					670		
Lys	Leu	Asn	Asp	Arg	Val	Met	Arg	Val	Glu	Tyr	His	Phe	Leu	Ser	Pro
		675						680				685			
Tyr	Val	Ser	Pro	Lys	Glu	Ser	Pro	Phe	Arg	His	Val	Phe	Trp	Gly	Ser
	690					695					700				
Gly	Ser	His	Thr	Leu	Pro	Ala	Leu	Leu	Glu	Asn	Leu	Lys	Leu	Arg	Lys
705					710					715					720
Gln	Asn	Asn	Gly	Ala	Phe	Asn	Glu	Thr	Leu	Phe	Arg	Asn	Gln	Leu	Ala
			725					730						735	
Leu	Ala	Thr	Trp	Thr	Ile	Gln	Gly	Ala	Ala	Asn	Ala	Leu	Ser	Gly	Asp
		740					745						750		
Val	Trp	Asp	Ile	Asp	Asn	Glu	Phe								
		755				760									

<210> SEQ ID NO 7

<211> LENGTH: 184

<212> TYPE: PRT

<213> ORGANISM: Macaca mulatta

<400> SEQUENCE: 7

Ala	Gln	Asn	Ser	Val	Ile	Ile	Val	Asp	Lys	Asn	Gly	Gly	Leu	Val	Tyr
1			5					10					15		
Leu	Val	Glu	Asn	Pro	Gly	Gly	Tyr	Val	Ala	Tyr	Ser	Lys	Ala	Ala	Thr
		20					25					30			
Val	Thr	Gly	Lys	Leu	Val	His	Ala	Asn	Phe	Gly	Thr	Lys	Lys	Asp	Phe
		35				40					45				
Glu	Asp	Leu	Asp	Ser	Pro	Val	Asn	Gly	Ser	Ile	Val	Ile	Val	Arg	Ala
	50					55				60					
Gly	Lys	Ile	Thr	Phe	Ala	Glu	Lys	Val	Ala	Asn	Ala	Glu	Ser	Leu	Asn
65				70				75						80	
Ala	Ile	Gly	Val	Leu	Ile	Tyr	Met	Asp	Gln	Thr	Lys	Phe	Pro	Ile	Val
			85					90						95	
Lys	Ala	Asp	Leu	Ser	Phe	Phe	Gly	His	Ala	His	Leu	Gly	Thr	Gly	Asp
		100					105					110			
Pro	Tyr	Thr	Pro	Gly	Phe	Pro	Ser	Phe	Asn	His	Thr	Gln	Phe	Pro	Pro
	115					120						125			
Ser	Gln	Ser	Ser	Gly	Leu	Pro	Asn	Ile	Pro	Val	Gln	Thr	Ile	Ser	Arg
	130				135						140				
Ala	Ala	Ala	Glu	Lys	Leu	Phe	Gly	Asn	Met	Glu	Gly	Asp	Cys	Pro	Ser
145				150				155						160	
Asp	Trp	Lys	Thr	Asp	Ser	Thr	Cys	Lys	Met	Val	Thr	Ser	Glu	Asn	Lys
		165					170						175		
Ser	Val	Lys	Leu	Thr	Val	Ser	Asn								
		180													

<210> SEQ ID NO 8

<211> LENGTH: 184

<212> TYPE: PRT

<213> ORGANISM: Pan troglodytes

<400> SEQUENCE: 8

Ala	Gln	Asn	Ser	Val	Ile	Ile	Val	Asp	Lys	Asn	Gly	Ser	Leu	Val	Tyr
1			5					10					15		

-continued

Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys Ala Ala Thr
 20 25 30
 Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys Lys Asp Phe
 35 40 45
 Glu Asp Leu His Thr Pro Val Asn Gly Ser Ile Val Ile Val Arg Ala
 50 55 60
 Gly Lys Ile Thr Phe Ala Glu Lys Val Ala Asn Ala Glu Ser Leu Asn
 65 70 75 80
 Ala Ile Gly Val Leu Ile Tyr Met Asp Gln Thr Lys Phe Pro Ile Val
 85 90 95
 Asn Ala Glu Leu Ser Phe Phe Gly His Ala His Leu Gly Thr Gly Asp
 100 105 110
 Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His Thr Gln Phe Pro Pro
 115 120 125
 Ser Arg Ser Ser Gly Leu Pro Asn Ile Pro Val Gln Thr Val Ser Arg
 130 135 140
 Ala Ala Ala Glu Lys Leu Phe Gly Asn Met Glu Gly Asp Cys Pro Ser
 145 150 155 160
 Asp Trp Lys Thr Asp Ser Thr Cys Arg Met Val Thr Ser Glu Ser Lys
 165 170 175
 Asn Val Lys Leu Thr Val Ser Asn
 180

<210> SEQ ID NO 9

<211> LENGTH: 184

<212> TYPE: PRT

<213> ORGANISM: Macaca fascicularis

<400> SEQUENCE: 9

Ala Gln Asn Ser Val Ile Ile Val Asp Lys Asn Gly Gly Leu Val Tyr
 1 5 10 15
 Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys Ala Ala Thr
 20 25 30
 Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys Lys Asp Phe
 35 40 45
 Glu Asp Leu Asp Ser Pro Val Asn Gly Ser Ile Val Ile Val Arg Ala
 50 55 60
 Gly Lys Ile Thr Phe Ala Glu Lys Val Ala Asn Ala Glu Ser Leu Asn
 65 70 75 80
 Ala Ile Gly Val Leu Ile Tyr Met Asp Gln Thr Lys Phe Pro Ile Val
 85 90 95
 Lys Ala Asp Leu Ser Phe Phe Gly His Ala His Leu Gly Thr Gly Asp
 100 105 110
 Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His Thr Gln Phe Pro Pro
 115 120 125
 Ser Gln Ser Ser Gly Leu Pro Asn Ile Pro Val Gln Thr Ile Ser Arg
 130 135 140
 Ala Ala Ala Glu Lys Leu Phe Gly Asn Met Glu Gly Asp Cys Pro Ser
 145 150 155 160
 Asp Trp Lys Thr Asp Ser Thr Cys Lys Met Val Thr Ser Glu Asn Lys
 165 170 175
 Ser Val Lys Leu Thr Val Ser Asn
 180

-continued

<210> SEQ ID NO 10
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic sgRNA sequence

<400> SEQUENCE: 10

gaatacacatc actcctcgtg agg

23

<210> SEQ ID NO 11
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic sgRNA sequence

<400> SEQUENCE: 11

agaagaatac ttaacatctt tgg

23

What is claimed is:

1. A non-human transgenic animal that expresses a chimeric transferrin receptor (TfR) polypeptide, wherein the chimeric TfR polypeptide comprises (a) a heterologous apical domain that replaces the apical domain of the TfR polypeptide endogenous to the non-human transgenic animal and (b) an endogenous transferrin binding site.

2. The non-human transgenic animal of claim 1, wherein the heterologous apical domain comprises an amino acid sequence having at least 80% identity to SEQ ID NO:1.

3. The non-human transgenic animal of claim 2, wherein the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:1.

4. The non-human transgenic animal of claim 1, wherein the heterologous apical domain comprises an amino acid sequence having at least 80% identity to SEQ ID NO:7, SEQ ID NO:8, or SEQ ID NO:9.

5. The non-human transgenic animal of claim 1, wherein the chimeric TfR polypeptide comprises an amino acid sequence having at least 80% identity to SEQ ID NO:3.

6. The non-human transgenic animal of claim 1, wherein the non-human transgenic animal is a mouse or rat.

7. The non-human transgenic animal of claim 6, wherein the non-human transgenic animal comprises a level of expression of the chimeric TfR polypeptide in brain, liver, kidney, or lung tissue similar to the level of expression of TfR in a wild-type animal of the same species.

8. The non-human transgenic animal of claim 6, wherein the non-human transgenic animal comprises a red blood cell count, level of hemoglobin, or level of hematocrit similar to a wild-type animal of the same species.

9. The non-human transgenic animal of claim 1, wherein the non-human transgenic animal is a mouse, the heterologous apical domain comprises an amino acid sequence having at least 80% identity to SEQ ID NO:1, and the mouse comprises a level of expression of the chimeric TfR polypeptide in brain, liver, kidney, or lung tissue similar to the level of expression of TfR in a wild-type mouse.

10. The non-human transgenic animal of claim 1, wherein the genome of the non-human transgenic animal comprises a chimeric TfR nucleic acid sequence in which the apical

domain of an endogenous TfR gene of the non-human transgenic animal has been replaced with a heterologous apical domain.

11. The non-human transgenic animal of claim 10, wherein all introns outside the heterologous apical domain are retained.

12. The non-human transgenic animal of claim 10, wherein the heterologous apical domain is positioned after the fourth exon of the endogenous TfR gene.

13. The non-human transgenic animal of claim 10, wherein the heterologous apical domain comprises a nucleic acid sequence having at least 70% identity to SEQ ID NO:2.

14. The non-human transgenic animal of claim 13, wherein the heterologous apical domain comprises the nucleic acid sequence of SEQ ID NO:2.

15. The non-human transgenic animal of claim 10, wherein the transgenic animal is homozygous for the chimeric TfR nucleic acid sequence.

16. The non-human transgenic animal of claim 10, wherein the transgenic animal is heterozygous for the chimeric TfR nucleic acid sequence.

17. The non-human transgenic animal of claim 10, wherein the non-human transgenic animal is a mouse, the heterologous apical domain comprises a nucleic acid sequence having at least 70% identity to SEQ ID NO:2, and the mouse is homozygous for the chimeric TfR nucleic acid sequence.

18. An embryonic stem (ES) cell that expresses a chimeric transferrin receptor (TfR) polypeptide, wherein the chimeric TfR polypeptide comprises (a) a heterologous apical domain in place of the endogenous apical domain of the TfR polypeptide and (b) an endogenous transferrin binding site.

19. The ES cell of claim 18, wherein the ES cell is a mouse ES cell.

20. The ES cell of claim 18, wherein the heterologous apical domain comprises an amino acid sequence having at least 80% identity to SEQ ID NO:1.

21. The ES cell of claim 20, wherein the heterologous apical domain comprises the amino acid sequence of SEQ ID NO:1.

22. The ES cell of claim **18**, wherein the heterologous apical domain comprises an amino acid sequence having at least 80% identity to SEQ ID NO:7, SEQ ID NO:8, or SEQ ID NO:9.

23. The ES cell of claim **18**, wherein the chimeric TfR polypeptide comprises an amino acid sequence having at least 80% identity to SEQ ID NO:3.

24. A method of screening for an apical domain binding polypeptide (ADBP) that crosses the blood-brain barrier, the method comprising administering an ADBP to the non-human transgenic animal of claim **1**.

25. The method of claim **24**, wherein the method further comprises measuring the presence or an activity of the ADBP in the brain of the non-human transgenic animal.

26. The method of claim **25**, wherein the ADBP is coupled to an effector molecule.

27. The method of claim **26**, wherein the effector molecule is a small molecule, RNA, DNA, or polypeptide.

28. The method of claim **27**, wherein the polypeptide is an antibody or an antigen-binding fragment thereof.

29. The method of claim **26**, wherein the measuring step comprises contacting the brain or brain tissue of the non-human transgenic animal with an agent that binds to the effector molecule and determining the level of the effector molecule present in the brain.

30. The method of claim **26**, wherein the measuring step comprises measuring a pharmacodynamic (PD) effect of the effector molecule.

* * * * *

专利名称(译)	转铁蛋白受体转基因模型		
公开(公告)号	US20180235195A1	公开(公告)日	2018-08-23
申请号	US15/923928	申请日	2018-03-16
申请(专利权)人(译)	Denali Therapeutics Inc..		
当前申请(专利权)人(译)	Denali Therapeutics Inc..		
[标]发明人	DENNIS MARK S SILVERMAN ADAM P ZUCHERO JOY YU		
发明人	DENNIS, MARK S. SILVERMAN, ADAM P. ZUCHERO, JOY YU		
IPC分类号	A01K67/027 C07K14/705 G01N33/53		
CPC分类号	A01K67/0278 C07K14/705 G01N33/5308 A01K2217/072 A01K2217/075 A01K2227/105 A01K2267/03 G01N2500/10 A01K67/0275 A01K2217/00 C07K14/70582 C07K14/79 C07K2319/00 G01N33/5088 G01N33/68		
优先权	62/460692 2017-02-17 US 62/543658 2017-08-10 US 62/543559 2017-08-10 US 62/583314 2017-11-08 US		
其他公开文献	US10143187		
外部链接	Espacenet USPTO		

摘要(译)

在一些方面，本发明提供嵌合转铁蛋白受体（TfR）多核苷酸和多肽。在其他方面，本发明提供了嵌合TfR转基因动物模型和使用动物模型鉴定可以穿过血脑屏障的治疗剂的方法。

