



US007217535B2

(12) **United States Patent**  
**MacPhee et al.**(10) **Patent No.:** **US 7,217,535 B2**  
(45) **Date of Patent:** **\*May 15, 2007**(54) **LIPOPROTEIN ASSOCIATED  
PHOSPHOLIPASE A2, INHIBITORS  
THEREOF AND USE OF THE SAME IN  
DIAGNOSIS AND THERAPY**(75) Inventors: **Colin H. MacPhee**, Berwyn, PA (US);  
**David Graham Tew**, Leicester (GB)(73) Assignee: **SmithKline Beecham P.L.C.**,  
Brentford, Middlesex (GB)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 149 days.This patent is subject to a terminal dis-  
claimer.(21) Appl. No.: **10/973,467**(22) Filed: **Oct. 26, 2004**(65) **Prior Publication Data**

US 2005/0106611 A1 May 19, 2005

**Related U.S. Application Data**(63) Continuation of application No. 10/173,233, filed on  
Jun. 14, 2002, now Pat. No. 7,052,862, which is a  
continuation of application No. 09/569,899, filed on  
May 12, 2000, which is a continuation of application  
No. 09/294,384, filed on Apr. 20, 1999, now Pat. No.  
6,177,257, which is a continuation of application No.  
08/387,858, filed as application No. PCT/GB94/  
01374 on Jun. 24, 1994, now Pat. No. 5,981,252.(30) **Foreign Application Priority Data**Jun. 25, 1993 (GB) ..... 9313144.9  
Jan. 11, 1994 (GB) ..... 9400413.2(51) **Int. Cl.**  
**C12Q 1/34** (2006.01)(52) **U.S. Cl.** ..... **435/18**; 435/4; 435/198;  
435/252.3; 435/320.1(58) **Field of Classification Search** ..... 435/18,  
435/198, 252.3, 320.1, 4  
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**5,019,508 A 5/1991 Johnson et al. .... 435/198  
5,108,899 A 4/1992 Allen ..... 435/7.21  
5,210,017 A 5/1993 Carlsson et al. .... 435/7.8  
5,279,957 A 1/1994 Gross ..... 435/240.2  
5,527,698 A 6/1996 Knopf et al.  
5,530,101 A 6/1996 Queen et al. .... 530/387.3  
5,532,152 A 7/1996 Cousens et al.  
5,605,801 A 2/1997 Cousens et al.  
5,641,669 A 6/1997 Cousens et al.  
5,656,431 A 8/1997 Cousens et al.  
5,698,403 A 12/1997 Cousens et al.  
5,847,088 A 12/1998 Cousens et al.  
5,880,273 A 3/1999 Adachi et al.  
5,968,818 A 10/1999 Gloger et al.5,977,308 A 11/1999 Cousens et al.  
5,981,252 A 11/1999 MacPhee et al.  
6,045,794 A 4/2000 Cousens et al.  
6,099,836 A 8/2000 Cousens et al.  
6,146,625 A 11/2000 Cousens et al.  
6,177,257 B1 \* 1/2001 Macphee et al. .... 435/7.4  
6,203,790 B1 3/2001 Cousens et al.  
7,045,329 B2 5/2006 MacPhee et al. .... 435/198  
7,052,862 B2 \* 5/2006 MacPhee et al. .... 435/18**FOREIGN PATENT DOCUMENTS**EP 359425 8/1989  
EP 509719 10/1992  
EP 658205 3/2000  
EP 673426 6/2001  
JP 04346797 8/1988  
JP 07059597 10/1989  
JP 04346797 12/1992  
JP 07059597 3/1995  
JP 10017600 1/1998  
WO WO 89/09818 10/1989  
WO WO 94/20069 9/1994  
WO WO 95/00649 1/1995  
WO WO 95/09921 4/1995  
WO WO 97/12984 4/1997  
WO WO 98/44158 10/1998  
WO WO 99/09147 2/1999  
WO WO 99/64627 12/1999

(Continued)

**OTHER PUBLICATIONS**Yamamoto et al. (1990) Formation of lipid hydroperoxides in the  
cupric ion-induced oxidation of plasma and low density lipoprotein.  
Oxid. Damage Repair, Int. Soc. Free Radical Res. Bienn. Meeting.  
Meeting Date 1990, pp. 287-291. Editor: K. J. A. Da, Jan. 1990.  
Stafforini et al., "The Platelet-activating Factor Acetylhydrolase  
from human erythrocytes", Journal of Biological Chemistry 1993  
vol. 268; 6; 3856-3865.  
Stafforini et al., "Platelet activating factor acetylhydrolase from  
human plasma", Methods in Enzymology 1990 187: 344-57. Kohler  
G, et al., Nature 256:495-497 (1975).  
Sevier, Ed, et al., Clinical Chem. 27:1797-1806 (1981).  
Chandrakumar NS, et al.; :Stereospecific Synthesis of Ehter  
Phospholipids. Preparation of a-Alkyl-2-(acylamino)-2  
deoxyglycerophosphoryl cholines, J. Org. Chem., 1197-1202  
(1983).  
Ponpipom MM, et al., Synthesis of Azide and Amide Analogs of  
Platelet-Activating Factors and Related Derivatives, Chem. Phys.  
Lipids;35:29-37 (1984).

(Continued)

*Primary Examiner*—Tekchand Saidha(74) *Attorney, Agent, or Firm*—Licata & Tyrrell P.C.;  
Nathan P. Letts(57) **ABSTRACT**The enzyme Lp-PLA<sub>2</sub> in purified form, an isolated nucleic  
acid molecule encoding Lp-PLA<sub>2</sub>, the use of an inhibitor of  
the enzyme Lp-PLA<sub>2</sub> in therapy and a method of screening  
compounds to identify those compounds.

## FOREIGN PATENT DOCUMENTS

WO	WO 00/24910	5/2000
WO	WO 00/32808	6/2000

## OTHER PUBLICATIONS

Stafforini D, et al., "Human plasma platelet-activating factor acetylhydrolase," *J. Biol. Chem.* 262(9): 4224-4230 (1987).

Stafforini D, et al., "Human macrophages secrete plate-activating factor acetylhydrolase," *J. Biol. Chem.* 265(17):9682-9687 (1990).

Langlais J, et al., "Platelet-activating factor acetylhydrolase in human semen: a novel decapacitation factor? *Biol. of Reprod.* 44 (Suppl. 1):94 (1991).

San H, et al., "Safety and Short Term Toxicity of a Novel and Cationic Formulation for Human Gene Therapy," *Human Gene Therapy*: 781-798 (1993).

Kim UT, et al., "Synthesis of Phospholipid Head groups via Nucleophilic Ring Opening of 1,3,2-Dioxaphospholanes," *J. Chem. Commun.*: 70-71 (1993).

Christoffersen R, et al., "Ribozymes as Human Therapeutic Agents," *J. of Medicinal Chem.*: 2023-2037 (1994).

Stull R, et al., "Antigene, Ribozyme, and Aptamer Nucleic Acid Drugs: Progress and Prospects," *Pharm. Res.* 1995.

Kohler G, et al., 256:495-497 (1975).

U.S. Appl. No. 08/886,152, filed Jun. 18, 1996, related to ref. AW of Oct. 26, 2004 IDS.

Hirashima et al., "Platelet-activating factor (PAF) concentration and PAF acetylhydrolase activity in cerebrospinal fluid of patients with subarachnoid hemorrhage," *J. Neurosurg* 1994 80:31-36.

Kuby, Janis, "Immunology" 1991 W.H. Freeman 80-83.

Satoh et al., "Platelet-Activating Factor Acetylhydrolase in Plasma Lipoproteins From Patients With Ischemic Stroke", *Stroke* 1992 23:1090-1092.

Satoh et al., "Activity of Platelet-Activating Factor (PAF) Acetylhydrolase in Plasma from Patients with Ischemic Cerebrovascular Disease", *Prostaglandins* 1988 35 (5) :685-698.

Stafforini et al., "Platelet-activating Factor Acetylhydrolases", *J. Biol. Chem.* 1997 272(29) :17895-17898.

Washburn, W.N. and Dennis, E.A. "Novel General Approach for the Assay and Inhibition of Hydrolytic Enzymes Utilizing Suicide-Inhibitory Bifunctional Linked substrates (SIBLINKS) : Exemplified in a Phospholipase Assay" *J. Am. Chem. Soc.* 1990 112:2040-2041.

Washburn, W.N. and Dennis, E.A. "Suicide-inhibitory Bifunctionally Linked Substrates (SIBLINKS) as Phospholipase A2 Inhibitors" *J. Biol. Chem.* 1991 266(8) :5042-5048.

Yoshida et al., "Platelet-activating factor acetylhydrolase activity in red blood cell-stroma from patients with cerebral thrombosis", *Acta Neurol Scand* 1992 86:199-203.

Adjei et al., "Pulmonary Delivery of Peptide Drugs: Effect of Particle Size on Bioavailability of Leuprolide Acetate in Healthy Male Volunteers", *Pharm. Res.* 7(6):565-569 (1990).

Basran et al., "Properties of Platelet Activating Factor (PAF-ACETHER) Which Suggest Involvement in Chronic Inflammation and Persistent Asthma", *Br. J. Pharmacol.* 77:437 (1982).

Braquet et al., "Effect of Endothelin-1 on Blood Pressure and Bronchopulmonary System of the Guinea Pig", *J. Cardio. Pharm.* 13 (Supp.5):S143-S146 (1989).

Brenner, "The Molecular Evolution of Genes and Proteins: A Tale of Two Serines", *Nature* 334:528-530 (1988).

Campbell, *Biology* 3rd Ed., p. 104-106 (1993).

Capecchi, "Altering the Genome by Homologous Recombination", *Science* 244:1288-1292 (1989).

Caplan et al., "Role of Platelet Activating Factor and Tumor Necrosis Factor- $\alpha$  in Neonatal Necrotizing Enterocolitis", *J. Pediatr.* 116(6):960-964 (1990).

Chapus et al., "Minireview on Pancreatic Lipase and Colipase", *Biochimie* 70:1223-1234 (1988).

deBoer et al., "The tac Promoter: A Functional Hybrid Derived from the trp and lac Promoters", *Proc. Natl. Acad. Sci. USA* 80:21-25 (1983).

Debs et al., "Lung-Specific Delivery of Cytokines Induces Sustained Pulmonary and Systemic Immunomodulation in Rats", *J. Immunol.* 140(10):3482-3488 (1988).

Denizot et al., "PAF-Acether and Acetylhydrolase in Stool of Patients with Crohn's Disease", *Digestive Diseases and Sciences* 37(3):432-437 (1992).

Furukawa et al., "Platelet-activating Factor-induced Ischemic Bowel Necrosis: The Effect of Platelet-Activating Factor Acetylhydrolase", *Ped. Res.* 34(2):237-241 (1993).

Grino J.M., "BN 52021: A Platelet Activating Factor Antagonist for Preventing Post-Transplant Renal Failure", *Ann. Int. Med.* 121(5):345-347 (1994).

Hahn et al., "The Complete Sequences of Plasmids pFNeo and pMH-Neo: Convenient Expression Vectors for High-level Expression of Eukaryotic Genes in Hematopoietic Cell Lines", *Gene* 127:267-268 (1993).

Harlow et al., "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, p. 423 (1988).

Handley et al., "Platelet Activating Factor and Inflammation in Atherogenesis: Targets for Drug Development", *Drug Dev. Res.* 7:361-375 (1986).

Hattori et al., "The Catalytic Subunit of Bovine Brain Platelet-activating Factor Acetylhydrolase is a Novel Type of Serine Esterase", *J. Biol. Chem.* 269(37):23150-23155 (1994).

Hattori et al., "Purification and Characterization of Bovine Brain Platelet-activating Factor Acetylhydrolase", *J. Biol. Chem.* 268(25):18748-18753 (1993).

Henriques et al., "Endothelin-1 Inhibits PAF-induced Paw Oedema and Pleurisy in the Mouse", *Br. J. Pharmacol.* 106:579-582 (1992).

Heuer et al., "Current Status of PAF Antagonist", *Clin. Exp. Allergy* 22:980-983 (1992).

Hoffman et al., "Detection of Platelet-activating Factor in Amniotic Fluid of Complicated Pregnancies", *Am. J. Obstet Gynecol.* 162(2):525-528 (1990).

Horwitz et al., "DNA Sequence of the araBAD-araC Controlling Region in *Salmonella typhimurium* LT2", *Gene* 14:309-319 (1981).

Hsieh et al., "Increased Plasma Platelet-activating Factor in Children with Acute Asthmatic Attacks and Decreased in vivo and in vitro Production of Platelet-activating Factor After Immunotherapy", *J. Allergy Clin. Immunol.* 91:650-657 (1993).

Hsueh et al., "Platelet-activating Factor, Tumor Necrosis Factor, Hypoxia and Necrotizing Enterocolitis", *Acta Paediatr. Suppl.* 396:11-17 (1994).

Hubbard et al., "Anti-Neutrophil-Elastase Defenses of the Lower Respiratory Tract in  $\alpha$ 1-Antitrypsin Deficiency Directly Augmented with an Aerosol of A-1-Antitrypsin", *Ann. Int. Med.* 111(3):206-212 (1989).

Kald et al., "Release of Platelet-Activating Factor in Acute Experimental Pancreatitis", *Pancreas* 8(4):440-442 (1993).

Kirsch et al., "Mechanism of Platelet Activating Factor-Induced Vascular Leakage in the Rat Trachea", *Exp. Lung Res.* 18:447-459 (1992).

Kurosawa et al., "Increased Levels of Blood Platelet-activating Factor in Bronchial Asthmatic Patients with Active Symptoms", *Allergy* 49:60-63 (1994).

LaVallie et al., "A Thioredoxin Gene Fusion Expression System That Circumvents Inclusion Body Formation in the *E. coli* Cytoplasm", *Bio/Technology* 11:187-193 (1993).

Lellouch-Tubiana et al., "Eosinophil Recruitment into Guinea Pig Lungs after PAF-acether and Allergen Administration", *Am. Rev. Respir. Dis.* 137:948-954 (1988).

Lellouch-Tubiana et al., "Ultrastructural Evidence for Extravascular Platelet Recruitment in the Lung upon Intravenous Injection of Platelet-activating Factor (PAF-acether) to Guinea-pigs", *Br. J. Exp. Path.* 66:345-355 (1985).

Lewin B., Genes V, Oxford University Press, New York, New York, Chapter 6, pp. 136-141 (1994).

Lindsberg et al., "Evidence for Platelet-Activating Factor as a Novel Mediator in Experimental Stroke in Rabbits", *Stroke* 21:1452-1457 (1990).

Lindsberg et al., "Platelet-activating Factor in Stroke And Brain Injury", *Ann. Neurol.* 30(2):117-129 (1991).

- Maki et al., "Platelet-activating Factor Acetylhydrolase Activity in Maternal, Fetal, and Newborn Rabbit Plasma During Pregnancy and Lactation", *Proc. Natl. Acad. Sci. USA* 85:728-732 (1988).
- Matsumoto et al., "Platelet-Activating Factor in Bronchoalveolar Lavage Fluid of Patients with Adult Respiratory Distress Syndrome", *Clin. Exp. Pharmacol. Physiol.* 19:509-515 (1992).
- Matsuzaki et al., "PAF Acetylhydrolase Activities in Human Systemic Lupus Erythematosus and Lupus-Prone Mice", *Clinica Chimica Acta* 210:139-144 (1992).
- Mezzano et al., "Detection of Platelet-Activating Factor in Plasma of Patients with *Streptococcal nephritis*", *J. Am. Soc. Nephrol.* 4:235-242 (1993).
- Miwa et al., "Characterization of Serum Platelet-activating Factor (PAF) Acetylhydrolase", *J. Clin. Invest.* 82:1983-1991 (1988).
- Rabinovici et al., "Platelet Activating Factor Mediates Interleukin-2-induced Lung Injury in the Rat", *J. Clin. Invest.* 89:1669-1673 (1992).
- Rabinovici et al., "ARDS-like Lung Injury Produced by Endotoxin in Platelet-activating Factor-primed Rats", *J. Appl. Physiol.* 74(4):1791-1802 (1993).
- Rodríguez-Roisin et al., "Platelet-activating Factor Causes Ventilation-Perfusion Mismatch in Humans", *J. Clin. Invest.* 93:188-194 (1994).
- Roitt et al., *Immunology*, Gower Medical Publishing Co., p. 25.7 (1985).
- Sandhu et al., "Amplification of Reproducible Allele Markets for Amplified Fragment Length Polymorphism Analysis", *Biotechniques* 12(1):16-22 (1992).
- Satoh et al., "Platelet-activating Factor (PAF) Stimulates the Production of PAF Acetylhydrolase by the Human Hepatoma Cell Line, HepG2", *J. Clin. Invest.* 87:476-481 (1991).
- Smith et al., "Pulmonary Deposition and Clearance of Aerosolized Alpha-1-Proteinase Inhibitor Administered to Dogs and to Sheep", *J. Clin. Invest.* 84:1145-1154 (1989).
- Stafforini et al., "Human Plasma Platelet-activating Factor Acetylhydrolase: Association with Lipoprotein Particles and Role in the Degradation of Platelet-activating Factor", *J. Biol. Chem.* 262(9):4215-4222 (1987).
- Stafforini et al., "Platelet-activating Factor Acetylhydrolase Activity in Human Tissues and Blood Cells", *Lipids*, 26(12):979-985 (1991).
- Stafforini et al., "Lipoproteins Alter the Catalytic Behavior of the Platelet-activating Factor Acetylhydrolase in Human Plasma", *Proc. Natl. Acad. Sci. USA* 86:2393-2397 (1989).
- Stremmler et al., "An Oxidized Derivative of Phosphatidylcholine Is a Substrate for the Platelet-activating Factor Acetylhydrolase from Human Plasma", *J. Biol. Chem.* 264(10): 5331-5334 (1989).
- Stremmler et al., "Human Plasma Platelet-activating Factor Acetylhydrolase", *J. Biol. Chem.* 266(17):11095-11103 (1991).
- Tarbet et al., "Liver cells Secrete the Plasma Form of Platelet-activating Factor Acetylhydrolase", *J. Biol. Chem.* 266(25):16667-16673 (1991).
- Tjoelker et al., "Anti-inflammatory Properties of a Platelet-activating Factor Acetylhydrolase", *Nature* 374:549-553 (1995).
- Venable et al., "Platelet-activating Factor: a Phospholipid Autacoid with Diverse Actions", *J. Lipid Res.* 34:691-702 (1993).
- Von Heijne, "A New Method for Predicting Signal Sequence Cleavage Sites", *Nuc. Acids. Res.* 14(11):4683-4690 (1986).
- Wada et al., "Codon Usage Tabulated from the GenBank Genetic Sequence Data", *Nuc. Acids Res.* 19S:1981-1986 (1991).
- Watanabe et al., "Pharmacological Analysis of Neutrophil Chemotactic Factor Production by Leucocytes and Roles of PAF in Allergic Inflammation in Rats", *Br. J. Pharmacol.* 111:123-130 (1994).
- Watson et al., "The Platelet-Activating Factor Antagonist Web 2170; Its Beneficial Effect on Dog Renal Allograft Survival", *Transplantation* 56(4):1047-1049 (1993).
- Yasuda et al., "The Hormonal Regulation of Platelet-Activating Factor-Acetylhydrolase in the Rat", *Endocrinology* 130(2):708-716 (1992).
- Zarco et al., "Involvement of Platelet-activating Factor and Tumour Necrosis Factor in the Pathogenesis of Joint Inflammation in Rabbits", *Clin. Exp. Immunol.* 88:318-323 (1992).
- Elstad et al., "Platelet-activating Factor Acetylhydrolase Increases during Macrophage Differentiation" *J. Biol. Chem.* 264:8467-8470 (1989).
- Smiley et al., "Oxidatively Fragmented Phosphatidylcholines Activate Human Neutrophils through the Receptor for Platelet-activating Factor," *J. Biol. Chem.* 266:11104-11110 (1991).
- Tjoelker et al., "Plasma Platelet-activating Factor Acetylhydrolase is a Secreted Phospholipase A2 with a Catalytic Triad," *J. Biol. Chem.* 270:25481-25487 (1995).
- Yasuda et al., "Influence of a Cigarette Smoke Extract on the Hormonal Regulation of Platelet-activating Factor Acetylhydrolase in Rats," *Biol. Repro.* 53:244-252 (1995).
- Sigma-Aldrich Catalog Product Detail, PAF (2005).

\* cited by examiner




Fig.1 Compound 4 inhibits copper ( $5\mu\text{M}$ ) - stimulated LDL ( $150\mu\text{g/ml}$ ) oxidation.

## FIGURE 1



Lanes 2, 4 and 6 contain adjacent fractions of purified native Lp-PLA2. Lanes 1, 3 and 5 are fractions 2, 4 and 6 respectively after N-deglycosylation.

**FIGURE 2**



which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low-melting vegetable or synthetic waxes or fats.

Preferably the composition is in unit dose form such as a tablet or capsule.

Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base.

The daily dosage regimen for an adult patient may be, for example, an oral dose of between 1 mg and 500 mg, preferably between 1 mg and 250 mg, or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg, of the compound of the formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy.

The enzyme, lipoprotein associated Lp-PLA<sub>2</sub> has not hitherto been available in isolated purified form. The present invention therefore provides in a further aspect, the enzyme lipoprotein associated Lp-PLA<sub>2</sub> in purified form. By purified form is meant at least 80%, more preferably 90%, still more preferably 95% and most preferably 99% pure with respect to other protein contaminants.

The enzyme Lp-PLA<sub>2</sub> may be characterised by one or more partial peptide sequences selected from SEQ ID NOS: 1, 2, 3, 4, 10 and 11 or by the partial peptide sequence comprising residues 271 to 441 or consisting of residues 1 to 441 of SEQ ID NO: 9. The enzyme Lp-PLA<sub>2</sub> may further or alternatively characterised by its molecular weight found to be 45 kDa, at least 45 kDa, 45–47 kDa, 46–47 kDa or 45–50 kDa.

The invention also provides fragments of the enzyme having Lp-PLA<sub>2</sub> activity.

The enzyme can be isolated and purified using the methods hereafter described. Once isolated, the protein sequence of the enzyme can be obtained using standard techniques. In identifying said sequence, a number of protein fragments have been identified, each of which comprises part of the whole sequence of the enzyme. These sequences are themselves novel and form a further aspect of the invention.

This invention also provides isolated nucleic acid molecules encoding the enzyme, including mRNAs, DNAs, cDNAs as well as antisense analogs thereof and biologically active and diagnostically or therapeutically useful fragments thereof.

In particular, the invention provides an isolated nucleic acid molecule consisting of bases 1 to 1361 or 38 to 1361 or comprising the sequence corresponding to bases 848 to 1361 of SEQ ID NO: 9.

This invention also provides recombinant vectors, such as cloning and expression plasmids useful as reagents in the recombinant production of the enzyme, as well as recombinant prokaryotic and/or eukaryotic host cells comprising the novel nucleic acid sequence.

This invention also provides nucleic acid probes comprising nucleic acid molecules of sufficient length to specifically hybridize to the novel nucleic acid sequences.

This invention also provides an antisense oligonucleotide having a sequence capable of binding with mRNAs encoding the enzyme so as to prevent the translation of said mRNA.

This invention also provides transgenic non-human animals comprising a nucleic acid molecule encoding the enzyme. Also provided are methods for use of said transgenic animals as models for mutation and SAR (structure/activity relationship) evaluation as well as in drug screens.

This invention further provides a method of screening compounds to identify those compounds which inhibit the enzyme comprising contacting isolated enzyme with a test compound and measuring the rate of turnover of an enzyme substrate as compared with the rate of turnover in the absence of test compound.

“Recombinant” polypeptides refer to polypeptides produced by recombinant DNA techniques, i.e., produced from cells transformed by an exogenous DNA construct encoding the desired polypeptide. “Synthetic” polypeptides are those prepared by chemical synthesis.

A “replicon” is any genetic element (e.g., plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication in vivo: i.e., capable of replication under its own control.

A “vector” is a replicon, such as a plasmid, phage, or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A “double-stranded DNA molecule” refers to the polymeric form of deoxyribonucleotides (bases adenine, guanine, thymine, or cytosine) in a double-stranded helix, both relaxed and supercoiled. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the sense strand of DNA.

A DNA “coding sequence of” or a “nucleotide sequence encoding” a particular protein, is a DNA sequence which is transcribed and translated into a polypeptide when placed under the control of appropriate regulatory sequences.

A “promoter sequence” is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. Within the promoter sequence will be found a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain “TATA” boxes and “CAT” boxes.

DNA “control sequences” refers collectively to promoter sequences, ribosome binding sites, polyadenylation signals, transcription termination sequences, upstream regulatory domains, enhancers, and the like, which collectively provide for the expression (i.e., the transcription and translation) of a coding sequence in a host cell.

A control sequence “directs the expression” of a coding sequence in a cell when RNA polymerase will bind the promoter sequence and transcribe the coding sequence into mRNA, which is then translated into the polypeptide encoded by the coding sequence.

A “host cell” is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous DNA sequence.

A cell has been “transformed” by exogenous DNA when such exogenous DNA has been introduced inside the cell membrane. Exogenous DNA may or may not be integrated (covalently linked) into chromosomal DNA making up the

genome of the cell. In prokaryotes and yeasts, for example, the exogenous DNA may be maintained on an episomal element, such as a plasmid. With respect to eukaryotic cells, a stably transformed or transfected cell is one in which the exogenous DNA has become integrated into the chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cell containing the exogenous DNA.

A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth in vitro for many generations.

Two DNA or polypeptide sequences are "substantially homologous" or "substantially the same" when at least about 85% (preferably at least about 90%, and most preferably at least about 95%) of the nucleotides or amino acids match over a defined length of the molecule and includes allelic variations. As used herein, substantially homologous also refers to sequences showing identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See e.g., A "Current Protocols in Mol. Biol." Vol. I & II, Wiley Interscience, Ausbel et al. (ed.) (1992). Protein sequences that are substantially the same can be identified by proteolytic digestion, gel electrophoresis and microsequencing.

The term "functionally equivalent" intends that the amino acid sequence of the subject protein is one that will exhibit enzymatic activity of the same kind as that of Lp-PLA<sub>2</sub>.

A "heterologous" region of a DNA construct is an identifiable segment of DNA within or attached to another DNA molecule that is not found in association with the other molecule in nature.

This invention provides an isolated nucleic acid molecule encoding the enzyme Lp-PLA<sub>2</sub>. One means for isolating the coding nucleic acid is. to probe a human genomic or cDNA library with a natural or artificially designed probe using art recognized procedures (See for example: "Current Protocols in Molecular Biology", Ausubel, F. M., et al. (eds.) Greene Publishing Assoc and John Wiley Interscience, N.Y., 1989, 1992); for example using the protein fragment information disclosed herein. The enzyme of this invention may be made by recombinant genetic engineering techniques. The isolated nucleic acids particularly the DNAs can be introduced into expression vectors by operatively linking the DNA to the necessary expression control regions (e.g. regulatory regions) required for gene expression. The vectors can be introduced into the appropriate host cells such as prokaryotic (e.g., bacterial), or eukaryotic (e.g. yeast, insect or mammalian) cells by methods well known in the art (Ausubel et al., supra). The coding sequences for the desired proteins having been prepared or isolated, can be cloned into a suitable vector or replicon. Numerous cloning vectors are known to those of skill in the art, and the selection of an appropriate cloning vector is a matter of choice. Examples of recombinant DNA vectors for cloning and host cells which they can transform include the bacteriophage  $\lambda$  (*E. coli*), pBR322 (*E. coli*), pACYC177 (*E. coli*), pKT230 (gram-negative bacteria), pGV1106 (gram-negative bacteria), pLAFR1 (gram-negative bacteria), pME290 (non-*E. coli* gram-negative bacteria), pHV14 (*E. coli* and *Bacillus subtilis*), pBD9

(*Bacillus*), pIJ61 (*Streptomyces*), pUC6 (*Streptomyces*), YIp5 (*Saccharomyces*), a baculovirus insect cell system, , YCp19 (*Saccharomyces*). See generally "DNA Cloning": Vols. I & II, Glover et al. ed IRL Press Oxford (1985) (1987) and; T. Maniatis et. al. ("Molecular Cloning" Cold Spring Harbor Laboratory (1982).

The gene can be placed under the control of a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator (collectively referred to herein as "control" elements), so that the DNA sequence encoding the desire protein is transcribed into RNA in the host cell transformed by a vector containing this expression construction. The coding sequence may or may not contain a signal peptide or leader sequence. The protein sequences of the present invention can be expressed using, for example, the *E. coli* tac promoter or the protein A gene (spa) promoter and signal sequence. Leader sequences can be removed by the bacterial host in post-translational processing. See. e.g. U.S. Pat. Nos. 4,431,739; 4,425,437; 4,338,397.

In addition to control sequences, it may be desirable to add regulatory sequences which allow for regulation of the expression of the protein sequences relative to the growth of the host cell. Regulatory sequences are known to those of skill in the art, and examples include those which cause the expression of a gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound Other types of regulatory elements may also be present in the vector, for example, enhancer sequences.

An expression vector is constructed so that the particular coding sequence is located in the vector with the appropriate regulatory sequences, the positioning and orientation of the coding sequence with respect to the control sequences being such that the coding sequence is transcribed under the "control" of the control sequences (i.e., RNA polymerase which binds to the DNA molecule at the control sequences transcribes the coding sequence). Modification of the coding sequences may be desirable to achieve this end. For example, in some cases it may be necessary to modify the sequence so that it may be attached to the control sequences with the appropriate orientation; i.e., to maintain the reading frame. The control sequences and other regulatory sequences may be ligated to the coding sequence prior to insertion into a vector, such as the cloning vectors described above. Alternatively, the coding sequence can be cloned directly into an expression vector which already contains the control sequences and an appropriate restriction site. Modification of the coding sequences may also be performed to alter codon usage to suit the chosen host cell, for enhanced expression

In some cases, it may be desirable to add sequences which cause the secretion of the polypeptide from the host organism, with subsequent cleavage of the secretory signal. It may also be desirable to produce mutants or analogs of the enzyme of interest. Mutants or analogs may be prepared by the deletion of a portion of the sequence encoding the protein, by insertion of a sequence, and/or by substitution of one or more nucleotides within the sequence. Techniques for modifying nucleotide sequences, such as site-directed mutagenesis, are well known to those skilled in the art. See e.g., T. Maniatis et al., supra; *DNA Cloning*. Vols. I and II, supra; *Nucleic Acid Hybridization*, supra.

A number of prokaryotic expression vectors are known in the art. See. e.g., U.S. Pat. Nos. 4,578,355; 4,440,859; 4,436,815; 4,431,740; 4,431,739; 4,428,941; 4,425,437;



7

4,418,149; 4,411,994; 4,366,246; 4,342,832; see also U.K. Patent Applications GB 2,121,054; GB 2,008,123; GB 2,007,675; and European Patent Application 103,395. Yeast expression vectors are also known in the art See. e.g. U.S. Pat. Nos. 4,446,235; 4,443,539; 4,430,428; see also European Patent Applications 103,409; 100,561; 96,491. pSV2neo (as described in *J. Mol. Appl. Genet.* 1:327-341) which uses the SV40 late promoter to drive expression in mammalian cells or pCDNA1 neo, a vector derived from pCDNA1 (*Mol. Cell Biol.* 7:4125-29) which uses the CMV promoter to drive expression. Both these latter two vectors can be employed for transient or stable(using G418 resistance) expression in mammalian cells. Insect cell expression systems, e.g., *Drosophila*, are also useful, see for example, PCT applications U.S. Ser. No. 89/05155 and U.S. Ser. No. 91/06838 as well as EP application 88/3040933.

Depending on the expression system and host selected, the enzyme of the present invention may be produced by growing host cells transformed by an expression vector described above under conditions whereby the protein of interest is expressed. The protein is then isolated from the host cells and purified. If the expression system secretes the protein into growth media, the protein can be purified directly from the media. If the protein is not secreted, it is isolated from cell lysates or recovered from the cell membrane fraction. Where the protein is localized to the cell surface, whole cells or isolated membranes can be used as an assayable source of the desired gene product Protein expressed bacterial hosts such as *E. coli* may require isolation from inclusion bodies and refolding. The selection of the appropriate growth conditions and recovery methods are within the skill of the art.

The identification of this novel target for the treatment of atherosclerosis, also leads to a novel diagnostic method to diagnose a patient's susceptibility to developing atherosclerotic disease.

The present invention therefore provides in a still further aspect a diagnostic method comprising isolating a sample of blood from the patient, and assaying said sample for Lp-PLA<sub>2</sub> activity. Patients that are susceptible to atherosclerotic disease are expected to have elevated levels of the Lp-PLA<sub>2</sub> enzyme, and hence the levels of Lp-PLA<sub>2</sub> provides an indication of the patients susceptibility to atherosclerotic disease. Moreover, Lp-PLA<sub>2</sub> is found located predominantly on dense subfraction(s) of LDL which are known to be very atherogenic. Plasma Lp-PLA<sub>2</sub> levels could therefore provide a ready measure of these very atherogenic small dense LDL particles.

It is expected that the presence of the enzyme in the blood sample of the patient can be assayed by analysis of enzyme activity (i.e. by an assay set up against the purified enzyme as standard); or alternatively by assaying protein content of the sample by using polyclonal or monoclonal antibodies prepared against the purified enzyme. Monoclonal (and polyclonal) antibodies raised against the purified enzyme or fragments thereof are themselves novel and form a further aspect of the invention.


## DATA AND EXAMPLES

### 1. Screen for Lp-PLA<sub>2</sub> Inhibition.

Enzyme activity was determined by measuring the rate of turnover of the artificial substrate (A) at 37° C. in 50 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid) buffer containing 150 mM NaCl, pH 7.4.

8



(A)



Assays were performed in 96 well titre plates.

Lp-PLA<sub>2</sub> was pre-incubated at 37° C. with vehicle or test compound for 10 min in a total volume of 180 μl. The reaction was then initiated by the addition of 20 μl 10× substrate (A) to give a final substrate concentration of 20 μM. The reaction was followed at 405 nm for 20 minutes using a plate reader with automatic mixing. The rate of reaction was measured as the rate of change of absorbance.

Results:

Compound No	XR <sup>1</sup>	R	Z	IC <sub>50</sub> (μM)
1	O(CH <sub>2</sub> ) <sub>15</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH	N <sup>+</sup> (CH <sub>3</sub> ) <sub>3</sub>	0.8
2	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH	N <sup>+</sup> (CH <sub>3</sub> ) <sub>3</sub>	3.5
3	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH	S <sup>+</sup> (CH <sub>3</sub> ) <sub>2</sub>	1.0
4	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH	SCH <sub>3</sub>	0.08
5	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH	OH	0.45
6	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH	OAc	0.2
7	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH		0.5
8	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CH <sub>3</sub> CONH		0.55
9	O(CH <sub>2</sub> ) <sub>17</sub> CH <sub>3</sub>	CF <sub>3</sub> CONH	N <sup>+</sup> (CH <sub>3</sub> ) <sub>3</sub>	2.5

### 2. Copper Stimulated LDL Oxidation:

Copper stimulated oxidation of LDL is routinely measured by following the increase in conjugated diene formation by monitoring the change in absorption at 234 nm. This assay can be used to study inhibitors of oxidative modification of LDL. FIG. 1 demonstrates that Lp-PLA<sub>2</sub> inhibitors are effective inhibitors of LDL oxidation through a prolongation of the lag phase, using compound 4 as an example.

### 3. Inhibition of Cu<sup>2+</sup> Stimulated Lyso-Phosphatidylcholine (Lyso-PtdCho) Formation.

A 1 ml aliquot of human LDL (0.25 mg protein/ml) was incubated for 15 min at 37° C. with compound or vehicle. 5 μM Cu<sup>2+</sup> was then added to allow oxidation/lyso-PtdCho formation to occur. The incubation was terminated by the addition of 3.75 ml chloroform/methanol/c HCl (200:400:5, v/v/v). Following the addition of 125 ml chloroform and 1.25 ml 0.1M HCl, the mixture was vortexed and centrifuged. The lower phase was carefully removed and the upper phase re-extracted with an equal volume of synthetic lower phase. The extracts were pooled and dried under nitrogen.

Phospholipids were reconstituted into 50  $\mu$ l chloroform-methanol (2:1 v/v). 10  $\mu$ l aliquots were spotted on to pre-run silica gel HPTLC plates and then developed in chloroform/methanol 25–30% methylamine (60:20:5 v/v/v). Plates were subsequently sprayed with the fluorescent indicator, 2-p-toluidinylnaphthalene-6-sulphonic acid (1 mM in 50 mM Tris/HCl, pH 7.4) to identify phospholipid components. Fluorescence was measured at 222 nm using a CAMAG TLC scanner. Lipoprotein lyso-PtdCho content was quantified using a standard curve (0.05–0.5  $\mu$ g) prepared in parallel.

Compound 4 dose dependently inhibits LDL lyso-PtdCho accumulation stimulated by copper ions with an IC<sub>50</sub> value of ~30  $\mu$ M

#### 4. Purification of Lipoprotein Associated Lp-PLA<sub>2</sub>

Low density lipoprotein (LDL) was obtained by plasma apheresis. The LDL was dialysed against 0.5 M NaCl, 50 mM MES (4-morpholine ethane sulphonic acid), pH=6.0 overnight at 4° C. Solid CHAPS (3-[(3-cholamidopropyl)dimethylamino]-1-propane sulphonate) was added to 10 mM and the LDL stirred for 30 minutes to effect solubilisation. The solubilised LDL was pumped onto a pre-equilibrated Blue Sepharose 6FF (Pharmacia) column (2.6x20 cm). The column was then washed with 50 mM MES, 10 mM CHAPS, 0.5 M NaCl, pH=6.0 followed by 50 mM Tris, 10 mM CHAPS, 0.5 M NaCl, pH=8.0 until the absorbance (280 nm) of the eluate reached zero. Lp-PLA<sub>2</sub> was eluted using 50 mM Tris, 10 mM CHAPS, 1.5 M NaCl, pH=8.0. The Lp-PLA<sub>2</sub> fraction was then concentrated and dialysed overnight against 50 mM Tris, 10 mM CHAPS, pH=8.0.

The dialysed Lp-PLA<sub>2</sub> was submitted to anion exchange chromatography on a mono Q column (Pharmacia) using 50 mM Tris, 10 mM CHAPS, pH=8.0 with a NaCl gradient from zero to 0.3 M. The Lp-PLA<sub>2</sub> fractions obtained from the mono Q column were applied directly to a Hi Trap Blue cartridge (Pharmacia). The cartridge was washed with 50 mM Tris, 10 mM CHAPS, 0.5 M NaCl, pH=8.0 until the absorbance of the eluate (280 nm) was zero. Lp-PLA<sub>2</sub> was then eluted using 50 mM Tris, 10 mM CHAPS, 1.5 M NaCl, pH=8.0. This gave Lp-PLA<sub>2</sub> which is greater than 95% pure as shown in FIG. 2. This also demonstrates that the native enzyme is extensively glycosylated.

#### 5. Enzyme Sequence

The purity of the final enzyme preparation was verified by five criteria 1) SDS-polyacrylamide gel electrophoresis gave one band for both native and de-glycosylated forms. 2) Reverse phase high pressure liquid chromatography (RP-HPLC) gave a single peak, 3) The intact preparation gave no results by protein sequencing, implying that the protein was N-terminally blocked and free of any contaminants with open N-terminals, 4) By laser desorption mass spectrometry only one broad peak was observed with de-glycosylated protein, and 5) none of the sections of extended peptide data from sequencing gave any database matches indicative of contaminating proteins. Three cleavage strategies were used to obtain internal sequence information; trypsin (after de-glycosylation), cyanogen bromide (methionine cleavage) and BNPS-Skatol (tryptophan cleavage). The resulting peptides were separated by RP-HPLC, collected and sequenced. The accumulated sequence data allowed several extended stretches of primary structure of the Lp-PLA<sub>2</sub> enzyme to be verified. These are shown below as Peptides 1, 2, 3 and 4 (SEQ ID Nos 1 to 4). When searched against the National Centre for Biotechnological information (NCBI) non-redundant peptide sequence databases no high similarity matches were obtained. Estimation of the molecular weight of pure,

de-glycosylated protein by laser desorption mass spectrometry gives values in the region of 45–47 kDa (separately 45 kDa and 46–47 kDa), indicating that the sequences constitute approximately 15 to 20% of the protein.

#### 6. Gene Sequence

Three expressed sequence tags (ESTs) from human cDNA libraries have been found to have extensive alignments with the Peptide Sequences 1 to 3. These EST's are shown below as Nucleotide Sequences 1 to 3 (SEQ ID NOS: 5 to 7) Nucleotide Sequence 1 is a 420 base sequence derived from a human foetal spleen library. Nucleotide Sequence 2 is a 379 base sequence derived from a 12-week human embryo library. Nucleotide Sequence 3 is a 279 base sequence derived from a T-cell lymphoma library. The identities at both the nucleic acid and amino acid level justified an overlapping alignment of the cDNA of all three ESTs, Nucleotide Sequences 3 (bases 1–278), 1 (bases 1–389) (in reverse orientation) and 2 (bases 1–304) with the Peptide Sequences 1, 2 and 3 (partially). Beyond these limits, the poor resolution of the raw sequence data precludes accurate base calling.

There are two remaining unassigned peptide sections from Peptides 3 and 4, both of which are expected to be present in the complete protein, -Q-Y-I-N-P-A-V-, and W-L-M-G-N-I-L-R-L-L-F-G-S-M-T-T-P-A-N-.

#### 7. Isolation of Full-Length Lp-PLA<sub>2</sub> cDNA

The full DNA sequence was determined for the clone (HLTA145) from which the Lymphoma EST (SEQ ID NO: 7) was derived, giving a total of 572 bases; SEQ ID NO: 8. There is one base difference between this sequence and the EST (between bases 1 to 256 of the EST); at position 27 of HLTA145 there is an A compared with a T in the EST. This would cause a coding change; L in HLTA145 compared with F in the EST. Clone HLTA145 was used as a radiolabelled probe to screen the Lymphoma cDNA library in order to isolate the full-length Lp-PLA<sub>2</sub> clone. The library was prepared in the bacteriophage  $\lambda$  vector, Unizap XR (Stratagene).

#### Preparation of the Filters for Screening

The library was plated out at a density of 20,000 plaques per 150 mm petri dish onto *E.coli* XL-1 Blue host cells (ie. 200,000 plaques on 10 dishes). An overnight of XL-1 Blue was prepared in 100 mls LB/0.2% w/v Maltose/10 mM MgSO<sub>4</sub>. The cells were pelleted, resuspended in 50 mls 10 mM MgSO<sub>4</sub> and stored on ice. 180  $\mu$ l of the library bacteriophage stock (23,400 pfu's) were added to 7 mls XL-1 Blue cells, mixed and divided into 10 aliquots of 615  $\mu$ l. The 10 tubes were incubated at 37° C. for 30 minutes. 7 mls of molten (@45° C.) top agarose (0.7% wv agarose in LB) were added, mixed well and poured onto 150 mm LB agar plates (1.5% w/v agar in LB). The plates were inverted and incubated at 37° C. for approximately 7.5 hour. The plates were held at 4° C. until needed.

The plaques were transferred to 132 mm Hybond-N nylon filters (Amersham International) by laying the filters on the plates for 2 minutes (4 minutes for the duplicate). The DNA's on the filters were denatured for 2 minutes (0.5M NaCl, 1.5M NaOH), neutralised for 4 minutes (15M NaCl, 1.0M Tris pH7.4) and the filters placed on 2xSSC for 1 minute. The filters were then dried and the DNA cross-linked to the filter using a Stratalinker UV 2400 (Stratagene) at 120,000  $\mu$ Joules/cm<sup>2</sup>.

The filters were pre-hybridised in 1 mM EDTA, 0.5M NaHPO<sub>4</sub>, 7% SDS (Church, G M. and Gilbert, W. (1984) PNAS USA 81 p1991–1995) in a Techne HB2 hybridisation

oven at 55° C. for 3 hours. Each bottle contained 2 filters and 25 mls prehybridization solution.

#### Preparation of the Radiolabelled Probe

The probe cDNA (from HLTA 145) was excised from pBluescript II SK+/- as an approximately 600 bp EcoRI-XhoI fragment and approximately 100 ng of gel purified fragment were labelled using 1.2 MBq <sup>32</sup>P dATP and 1.2 MBq <sup>32</sup>P dCTP by PCR labelling using Taq DNA polymerase (Boehringer Mannheim) and primers designed to prime at the 5' and 3' ends of the EST sequence. The labelling reaction was carried out in a total volume of 200 µl and included unlabelled dNTP's at the following concentrations:

dATP 20 µM  
dCTP 20 µM  
dGTP 200 µM  
dTTP 200 µM

The PCR reaction was carried out over 35 cycles of:

94° C. for 30 s  
60° C. for 30 s  
72° C. for 30 s

#### Screening

The radiolabelled probe was denatured at 98° C. for 5 minutes and divided into 10 aliquots of 20 µl. One aliquot was added per hybridisation bottle. Hybridisation was carried out over 16 hours at 55° C. The filters were washed at 60° C. (2x10 minutes) with 0.1% w/y SDS, 0.1xSSC (50 mls per wash per bottle). The filters were autoradiographed and the films (Fuji Medical X-Ray Film) developed after 5 days exposure.

Duplicate positives were progressed to a secondary screen. The plaques were cored out into 1 ml SM (100 mM NaCl, 10 mM MgSO<sub>4</sub>, 1M Tris, pH7.4), titrated and plated onto 90 mm petri dishes at between 20 and 200 pfu's per dish. The secondary screen was carried out as described for the primary screen except the filters were washed at 65° C. The autoradiographs were developed after 16 hours exposure.

#### DNA Sequencing

The duplicated positive clones from the secondary screen were excised from the λ Unizap XR bacteriophage vector into the Bluescript phagemid (according to the Stratagene manual) for characterisation. One of the clones, carrying an insert of approximately 1.5 kb, was sequenced on both strands (using the USB Sequenase 2.0 DNA sequencing kit) by primer walking (SEQ ID NO: 9). The cDNA has an open reading frame with the potential to code for a polypeptide of 441 amino acids.

The 3' region of the full-length cDNA aligns with the HLTA145 sequence with the exception of 3 mismatches (see below). The predicted polypeptide sequence of the lymphoma Lp-PLA<sub>2</sub> is shown as SEQ ID NO: 9.

Inspection of the full length cDNA (SEQ ID NO: 9) reveals probable errors in Peptide 3. One of these errors is in the assignment of continuity between V-M which is incompatible with the perfect sequence match with the cDNA after this position. It seems likely that a short peptide, containing the sequence -Q-Y-I-N-P-, co-purified with the longer cyanogen bromide partial cleavage peptide and, by being present in greater quantity, was assigned as the major sequence and contiguous with the subsequent amino acids. The remaining section of Peptide 3 and the whole of Peptide

4 can be identified in the predicted full length enzyme sequence (SEQ ID No: 9). It thus seems likely that Peptide 3 is in fact two separate Peptides 5 (SEQ ID No:10) and 6 (SEQ ID NO: 11). The second probable error has occurred in the transcription from the raw data for Peptide 3 which on checking was consistent with Peptide 5 having the sequence -Q-Y-I-N-P-V-A, rather than Q-Y-I-N-P-A-V-.

The 3 base differences are as follows:

- 1) T at 859 is A in HLTA145; aminoacid change F in full-length, L in HLTA145. (Note that the original EST is identical with the full-length cDNA at position 859).
- 2) C at 1173 is T in HLTA145; aminoacid change A in full-length, V in HLTA145.
- 3) T at 1203 is C in HLTA145; aminoacid change L in full-length, S in HLTA145.

The peptide data and the Foetal Spleen EST sequence (SEQ ID NO: 5) support the full-length cDNA sequence for differences (2) and (3) although the Human Embryo EST (SEQ ID NO: 6) is identical to the Lymphoma EST (SEQ ID NO: 7) at position 1173. The Human Embryo EST (SEQ ID NO: 6) has a further difference (4) corresponding to position 1245 in the full-length Lymphoma sequence (SEQ ID NO: 9)(comparison between bases 2 to 304 of the Human Embryo EST and the full-length Lymphoma cDNA).

- 4) A at 1245 is T in the Embryo EST (SEQ ID NO: 6)(amino acid change D to V in the Embryo EST). Peptide data covering this region supports the Lymphoma DNA sequence (SEQ ID NO: 9).

The Lp-PLA<sub>2</sub> DNA sequence from 848 to 1361 of SEQ ID NO: 9 (amino acid residues 271 to 441 of SEQ ID NO: 9) is the region for which all major data sets agree substantially, ie. the peptide data, the Foetal spleen, full-length Lymphoma and it includes the known active site and is therefore believed to be a significant characterising region for the Lp-PLA<sub>2</sub> enzyme.

The predicted MW for the full reading frame is 50090. This is in excess of that determined for the de-glycosylated, purified protein but post-translational events could account for this discrepancy. The most likely of these are the removal of an N-terminal signal peptide and/or limited proteolytic degradation of the protein C-terminal. The latter could occur in-vivo, during purification, or under the conditions of de-glycosylation.

#### Diagnostic Method

A sample of blood is taken from a patient, the plasma/serum sample prepared and passed through a dextran sulphate column pre-equilibrated with 0.9% (w/v) NaCl solution. Following washes with the same salt solution Lp-PLA<sub>2</sub> is eluted with a 4.1% (w/v) NaCl solution. Heparin agarose columns can also be used with the wash and elution solutions containing 0.7% and 2.9% NaCl, respectively. Enzyme present in the sample is determined by assaying for either (a) enzyme activity:

The substrate (A) (see structure in 1) is used to assay Lp-PLA<sub>2</sub> activity by monitoring the absorbance change at 400 nm. Purified enzyme is pre-incubated at 37° C. and substrate (50 µM) is added after 5 minutes. The absorbance change at 400 nm is monitored for 20-minutes. This substrate has previously been reported as a substrate for classical calcium-dependent PLA<sub>2</sub>s. (Washburn, W. N. and Dennis, E. A., J. Amer Chem. Soc., 1990, 112, 2040-2041); or

## (b) protein content

Total protein content (i.e. enzyme content) can be determined using polyclonal antiserum raised against purified human Lp-PLA<sub>2</sub>. The antisera recognises both native and glycosylated enzyme as measured by immunoprecipitation of activity and Western Blot analysis.

Polyclonal antiserum was prepared as follows. Immunisation of rabbits involved mixing 0.5 ml of purified human Lp-PLA<sub>2</sub> (=100 µg) with an equal volume of Freund's complete adjuvant. The final emulsion was given subcutaneously in 4x0.25 ml injections. Boosting using a Freund's incomplete adjuvant antigen mixture (4x0.25 ml subcut; dosage=50 µg) took place 4 weeks later. Adequate titre was evident at between 6–8 weeks from initial injection.

## IN THE FIGURES

FIG. 1 is a graph of absorbance at 234 nm against time (min) in a study of inhibition of copper (5 µM)-stimulated LDL (150 µg/ml) oxidation by compound 4 vs control vehicle.

FIG. 2 is an analysis the purified Lp-PLA<sub>2</sub> material of Example 4 following separation by polyacrylamide gel electrophoresis. Lanes 2, 4 and 6 contain adjacent fractions of purified native Lp-PLA<sub>2</sub>. Lanes 1, 3 and 5 are fractions 2, 4 and 6 respectively after N-deglycosylation.

## SEQUENCE DATA

SEQ. ID. No: 1 - Peptide 1

-M-L-K-L-K-G-D-I-D-S-N-A-A-I-D-L-S-N-K-A-S-L-A-F-L-Q-K-H-L-G-L-H-K-D-F-D-Q-

SEQ. ID. No: 2 - Peptide 2

-W-M-F-P-L-G-D-E-V-Y-S-R-I-P-Q-P-L-F-F-I-N-S-E-Y-F-Q-Y-P-A-N-

SEQ. ID. No: 3 - Peptide 3

-Q-Y-I-N-P-A-V-M-I-T-I-R-G-S-V-H-Q-N-F-A-D-F-T-F-A-T-G-

SEQ. ID. No: 4 - Peptide 4

-W-L-M-G-N-I-L-R-L-L-F-G-S-M-T-T-P-A-N

SEQ. ID. No: 5 - Nucleotide Sequence 1

```

1  AAAAAACCTA TTTTAATCCT AATTGTATTT CTCTATTCCT GAAGAGTTCT
51  GTAACATGAT GTGTTGATTG GTTGTGTTAA TGTGTTGCC TGGAAATAAGA
101 TTCTCATCAT CTCCTTCAAT CAAGCAGTCC CACTGATCAA AATCTTTATG
151 AAGTCCTAAA TGCTTTTGTA AGAATGCTAA TGAAGCTTTG TTGCTAAGAT
201 CAATAGCTGC ATTTGAATCT ATGTCTCCCT TTAATTTGAG CATGTGTCCA
251 ATTATTTTGC CAGTNGCAA AGTGAAGTCA GCAAAATTCT GGTGGACTGA
301 ACCCCTGATT GTAATCATCT TTCTTTCTTT ATCAGGTGAG TAGCATTTTT
351 TCATTTTAT GATATTAGCA GGATATTGGA AATATTGAGN GTTGNTAAAA
401 AGNGGNGGCT GAGGGATTCT

```

SEQ. ID. No: 6 - Nucleotide Sequence 2

```

1  TCGTAATATC ATAAAAATGA AAAAATGCTA CTCACCTGAT AAAGAAAGAA
51  AGATGATTAC AATCAGGGGT TCAGTCCACC AGANTTTTGC TGACTTCACT
101 TTTGCAACTG GCAAAATAAT TGGACACATG CTCAAATTAA AGGGAGACAT
151 AGATTCAAAT GTAGCTATTG ATCTTAGCAA CAAAGCTTCA TTAGCATTCT
201 TACAAAAGCA TTTAGGACTT CATAAAGATT TTGTTCAAGT GGAAGTCTTG
251 ATTGAAGGAG ATGATGAGAA TCTTATTCCA GGGACCAACA TTAACACAAC
301 CAATTCACAA CATCATGTTT ACAGAACTTC TTCCAGGGAA TAGGAGGAAA
351 TACAATTGGG GTTTAAATA GGTTTTTTTT

```

SEQ. ID. No: 7 - Nucleotide Sequence 3

```

1  GAAGAATGCA TTAGATTTAA AGTTTGATAT GGAACAACCTG AAGGACTCTA
51  TTGATAGGGA AAAAATAGCA GTAATTGGAC ATTCTTTTGG TGGAGCAACG
101 GTTATTCAGA CTCTTAGTGA AGATCAGAGA TTCAGATGTG GTATTGCCCT
151 GGATGCATGG ATGTTTCCAC TGGGTGATGA AGTATATTCC AGAATTCCTC

```

-continued

201 AGCCCCCTCTT TTTTATCAAC TCTGAATATT TCCAATATCC TGCTAATATC

251 ATAAAAANTGG AAAAAATGCTA CTCACCTGG

Seq. ID. No:8 - DNA sequence of HLTA145

10	20	30	40	50
AAAATAGCAG	TAATTGGACA	TTCTTTAGGT	GGAGCAACGG	TTATTCAGAC

60	70	80	90	100
TCTTAGTGAA	GATCAGAGAT	TCAGATGTGG	TATTGCCCTG	GATGCATGGA

110	120	130	140	150
TGTTTCCACT	GGGTGATGAA	GTATATTCCA	GAATTCCTCA	GCCCCCTTTT

160	170	180	190	200
TTTATCAACT	CTGAATATTT	CCAATATCCT	GCTAATATCA	TAAAAATGAA

210	220	230	240	250
AAAATGCTAC	TCACCTGATA	AAGAAAGAAA	GATGATTACA	ATCAGGGGTT

260	270	280	290	300
CAGTCCACCA	GAATTTTGCT	GACTTCACTT	TTGCAACTGG	CAAAATAATT

310	320	330	340	350
GGACACATGC	TCAAATTAAT	GGGAGACATA	GATTCAAATG	TAGCTATTGA

360	370	380	390	400
TCTTAGCAAC	AAAGCTTCAT	CAGCATTCCT	ACAAAAGCAT	TTAGGACTTC

410	420	430	440	450
ATAAAGATTT	TGATCAGTGG	GACTGCTTGA	TTGAAGGAGA	TGATGAGAAT

460	470	480	490	500
CTTATTCCAG	GGACCAACAT	TAACACAACC	AATCAACACA	TCATGTTACA

510	520	530	540	550
GAACTCTTCA	GGAATAGAGA	AATACAATTA	GGATTAAAAT	AGGTTTTTTA

560	570
AAAAAAAAAA	AAAAAAACT CG

SEQ. ID. No:9 - cDNA Sequence of Lymphoma Lp-PLA<sub>2</sub>

10	20	30	40	50
TGAGAGACTAAGCTGAACTGCTGCTCAGCTCCCAAGATGGTGCCACCCA				
		M	V	P P K

60	70	80	90	100
AATTGCATGTGCTTTTCTGCCTCTGCGGCTGCCTGGCTGTGGTTTATCCT				
L	H V L F C L C G C L A V V Y P			

110	120	130	140	150
TTTGACTGGCAATACATAAATCCTGTTGCCATATGAAATCATCAGCATG				
F D W Q Y I N P V A H M K S S A W				

160	170	180	190	200
GGTCAACAAAATACAAGTACTGATGGCTGCTGCAAGCTTTGGCCAAACTA				
V N K I Q V L M A A A S F G Q T K				

210	220	230	240	250
AAATCCCCGGGAAATGGGCCTTATCCGTTGGTTGTACAGACTTAATG				
I P R G N G P Y S V G C T D L M				

260	270	280	290	300
TTTGATCACACTAATAAGGCACCTTCTTGCGTTTATATTATCCATCCCA				
F D H T N K G T F L R L Y Y P S Q				

310	320	330	340	350
AGATAATGATCGCCTTGACACCCCTTTGGATCCCAAATAAGAAATATTTT				
D N D R L D T L W I P N K E Y F W				

360	370	380	390	400
GGGGTCTTAGCAAATTTCTTGGAACACACTGGCTTATGGGCAACATTTTG				
G L S K F L G T H W L M G N I L				

410	420	430	440	450
AGGTTACTCTTTGGTTCAATGACAACTCCTGCAACTGGAATTCCTCT				
R L L F G S M T T P A N W N S P L				

-continued

460 470 480 490 500  
GAGGCTGGTGAAAAATATCCACTTGTGTTTTTCTCATGGTCTTGGGG  
R P G E K Y P L V V F S H G L G A

510 520 530 540 550  
CATTCAGGACACTTTATTCGCTATTGGCATTGACCTGGCATCTCATGGG  
F R T L Y S A I G I D L A S H G

560 570 580 590 600  
TTTATAGTTGCTGCTGTAGAACACAGAGATAGATCTGCATCTGCAACTTA  
F I V A A V E H R D R S A S A T Y

610 620 630 640 650  
CTATTTCAAGGACCAATCTGCTGCAGAAATAGGGGACAAGTCTTGGCTCT  
Y F K D Q S A A E I G D K S W L Y

660 670 680 690 700  
ACCTTAGAACCTGAAACAAGAGGAGGAGACATATACGAAATGAGCAG  
L R T L K Q E E E T H I R N E Q

710 720 730 740 750  
GTACGGCAAAGAGCAAAGAATGTTCCAAGCTCTCAGTCTGATTCTTGA  
V R Q R A K E C S Q A L S L I L D

760 770 780 790 800  
CATTGATCATGGAAGCCAGTGAAGAATGCATTAGATTTAAAGTTTGATA  
I D H G K P V K N A L D L K F D M

810 820 830 840 850  
TGGAACAACTGAAGGACTCTATTGATAGGGAAAAATAGCAGTAATTGGA  
E Q L K D S I D R E K I A V I G

860 870 880 890 900  
CATTCTTTTGGTGGAGCAACGGTTATTCAGACTCTTAGTGAAGATCAGAG  
H S F G G A T V I Q T L S E D Q R

910 920 930 940 950  
ATTGAGATGTGGTATTGCCCTGGATGCATGGATGTTTCCACTGGGTGATG  
F R C G I A L D A W M F P L G D E

960 970 980 990 1000  
AAGTATATTCAGAATTCTCAGCCCTCTTTTTTATCAACTCTGAATAT  
V Y S R I P Q P L F F I N S E Y

1010 1020 1030 1040 1050  
TTCCAATATCCTGCTAATATCATAAAATGAAAAATGCTACTCACCTGA  
F Q Y P A N I I K M K K C Y S P D

1060 1070 1080 1090 1100  
TAAAGAAAGAAAGATGATTACAATCAGGGGTTGAGTCCACCAGAAATTTG  
K E R K M I T I R G S V H Q N F A

1110 1120 1130 1140 1150  
CTGACTTCACTTTTGCAACTGGCAAAATAATTGGACACATGCTCAAATTA  
D F T F A T G K I I G H M L K L

1160 1170 1180 1190 1200  
AAGGGAGACATAGATTCAAATGCAGCTATTGATCTTAGCAACAAAGCTTC  
K G D I D S N A A I D L S N K A S

1210 1220 1230 1240 1250  
ATTAGCATTCTTACAAAAGCATTAGGACTTCATAAGATTTTGATCAGT  
L A F L Q K H L G L H K D F D Q W

1260 1270 1280 1290 1300  
GGGACTGCTTGATTGAAGGAGATGATGAGAATCTTATCCAGGGACCAAC  
D C L I E G D D E N L I P G T N

1310 1320 1330 1340 1350  
ATTAACACAACCAATCAACACATCATGTTACAGAACTCTTCAGGAATAGA  
I N T T N Q H I M L Q N S S G I E

1360  
GAAATACAATT  
K Y N

-continued

SEQ. ID. No: 10 - Peptide 5  
 -Q-Y-I-N-P-V-A-

SEQ. ID. No: 11 - Peptide 6  
 -M-I-T-I-R-G-S-V-H-Q-N-F-A-D-F-T-F-A-T-G-

---

 SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 11

## (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 37 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: <Unknown>  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met Leu Lys Leu Lys Gly Asp Ile Asp Ser Asn Ala Ala Ile Asp Le  
 1 5 10 15

Ser Asn Lys Ala Ser Leu Ala Phe Leu Gln Lys His Leu Gly Leu Hi  
 20 25 30

Lys Asp Phe Asp Gln  
 35

## (2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 30 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Trp Met Phe Pro Leu Gly Asp Glu Val Tyr Ser Arg Ile Pro Gln Pr  
 1 5 10 15

Leu Phe Phe Ile Asn Ser Glu Tyr Phe Gln Tyr Pro Ala Asn  
 20 25 30

## (2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 27 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

-continued

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Gln Tyr Ile Asn Pro Ala Val Met Ile Thr Ile Arg Gly Ser Val Hi  
 1                      5                      10                      15

Gln Asn Phe Ala Asp Phe Thr Phe Ala Thr Gly  
                     20                      25

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 19 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Trp Leu Met Gly Asn Ile Leu Arg Leu Leu Phe Gly Ser Met Thr Th  
 1                      5                      10                      15

Pro Ala Asn

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 420 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

AAAAAACCTA TTTTAATCCT AATTGTATTT CTCTATTCCT GAAGAGTTCT GTAACATGAT      60

GTGTTGATTG GTTGTGTTAA TGTTGGTCCC TGAATAAGA TTCTCATCAT CTCCTTCAA      120

CAAGCAGTCC CACTGATCAA AATCTTTATG AAGTCCTAAA TGCTTTTGTA AGAATGCTA      180

TGAAGCTTTG TTGCTAAGAT CAATAGCTGC ATTTGAATCT ATGTCTCCCT TTAATTGA      240

CATGTGTCCA ATTATTTTGC CAGTNGCAAA AGTGAAGTCA GCAAAATTCT GGTGGACTG      300

ACCCCTGATT GTAATCATCT TTCCTTCTTT ATCAGGTGAG TAGCATTTTT TCATTTTAA      360

GATATTAGCA GGATATTGGA AATATTAGN GTTGNTAAAA AGNGGNGGCT GAGGGATTC      420

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 379 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO



-continued

---

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

TGCTAATATC ATAAAAATGA AAAAATGCTA CTCACCTGAT AAAGAAAGAA AGATGATTAC	60
AATCAGGGGT TCAGTCCACC AGANTTTTGC TGACTTCACT TTTGCAACTG GCAAAATAA	120
TGGACACATG CTCAAATTAA AGGGAGACAT AGATTCAAAT GTAGCTATTG ATCTTAGCA	180
CAAAGCTTCA TTAGCATTCT TACAAAAGCA TTTAGGACTT CATAAGATT TTGTTCACT	240
GGACTGCTTG ATTGAAGGAG ATGATGAGAA TCTTATTCCA GGGACCAACA TTAACACAA	300
CAATTCAACA CATCATGTTT ACAGAACTTC TTCCAGGGAA TAGGAGGAAA TACAATTGG	360
GTTTAAAATA GGTTTTTTT	379

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 279 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GAAGAATGCA TTAGATTTAA AGTTTGATAT GGAACAACG AAGGACTCTA TTGATAGGGA	60
AAAAATAGCA GTAATTGGAC ATTCTTTTGG TGGAGCAACG GTTATTCAGA CTCTTAGTG	120
AGATCAGAGA TTCAGATGTG GTATTGCCCT GGATGCATGG ATGTTTCCAC TGGGTGATG	180
AGTATATTCC AGAATTCCTC AGCCCCCTCT TTTTATCAAC TCTGAATATT TCCAATATC	240
TGCTAATATC ATAAAAANTGG AAAAATGCTA CTCACCTGG	279

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 572 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

AAAATAGCAG TAATTGGACA TTCTTTAGGT GGAGCAACGG TTATTCAGAC TCTTAGTGAA	60
GATCAGAGAT TCAGATGTGG TATTGCCCTG GATGCATGGA TGTTCCTACT GGGTGATGA	120
GTATATTCCA GAATTCCTCA GCCCTCTTT TTTATCAACT CTGAATATTT CCAATATCC	180
GCTAATATCA TAAAAATGAA AAAATGCTAC TCACCTGATA AAGAAAGAAA GATGATTAC	240
ATCAGGGGTT CAGTCCACCA GAATTTTGCT GACTTCACTT TTGCAACTGG CAAAATAAT	300
GGACACATGC TCAAAATAAA GGGAGACATA GATTCAAATG TAGCTATTGA TCTTAGCAA	360
AAAGCTTCAT CAGCATCTT ACAAAAGCAT TTAGGACTTC ATAAAGATTT TGATCAGTG	420
GACTGCTTGA TTGAAGGAGA TGATGAGAAT CTTATTCCAG GGACCAACAT TAACACAAC	480
AATCAACACA TCATGTTACA GAACTCTTCA GGAATAGAGA AATACAATTA GGATTAATA	540
AGGTTTTTTT AAAAAAAAAA AAAAAAACT CG	572

-continued

## (2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1361 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 38..1360

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TGAGAGACTA AGCTGAACT GCTGCTCAGC TCCCAAG ATG GTG CCA CCC AAA TTG	55
Met Val Pro Pro Lys Leu	
1 5	
CAT GTG CTT TTC TGC CTC TGC GGC TGC CTG GCT GTG GTT TAT CCT TTT	103
His Val Leu Phe Cys Leu Cys Gly Cys Leu Ala Val Val Tyr Pro Phe	
10 15 20	
GAC TGG CAA TAC ATA AAT CCT GTT GCC CAT ATG AAA TCA TCA GCA TGG	151
Asp Trp Gln Tyr Ile Asn Pro Val Ala His Met Lys Ser Ser Ala Trp	
25 30 35	
GTC AAC AAA ATA CAA GTA CTG ATG GCT GCT GCA AGC TTT GGC CAA ACT	199
Val Asn Lys Ile Gln Val Leu Met Ala Ala Ala Ser Phe Gly Gln Thr	
40 45 50	
AAA ATC CCC CGG GGA AAT GGG CCT TAT TCC GTT GGT TGT ACA GAC TTA	247
Lys Ile Pro Arg Gly Asn Gly Pro Tyr Ser Val Gly Cys Thr Asp Leu	
55 60 65 70	
ATG TTT GAT CAC ACT AAT AAG GGC ACC TTC TTG CGT TTA TAT TAT CCA	295
Met Phe Asp His Thr Asn Lys Gly Thr Phe Leu Arg Leu Tyr Tyr Pro	
75 80 85	
TCC CAA GAT AAT GAT CGC CTT GAC ACC CTT TGG ATC CCA AAT AAA GAA	343
Ser Gln Asp Asn Asp Arg Leu Asp Thr Leu Trp Ile Pro Asn Lys Glu	
90 95 100	
TAT TTT TGG GGT CTT AGC AAA TTT CTT GGA ACA CAC TGG CTT ATG GGC	391
Tyr Phe Trp Gly Leu Ser Lys Phe Leu Gly Thr His Trp Leu Met Gly	
105 110 115	
AAC ATT TTG AGG TTA CTC TTT GGT TCA ATG ACA ACT CCT GCA AAC TGG	439
Asn Ile Leu Arg Leu Leu Phe Gly Ser Met Thr Thr Pro Ala Asn Trp	
120 125 130	
AAT TCC CCT CTG AGG CCT GGT GAA AAA TAT CCA CTT GTT GTT TTT TCT	487
Asn Ser Pro Leu Arg Pro Gly Glu Lys Tyr Pro Leu Val Val Phe Ser	
135 140 145 150	
CAT GGT CTT GGG GCA TTC AGG ACA CTT TAT TCT GCT ATT GGC ATT GAC	535
His Gly Leu Gly Ala Phe Arg Thr Leu Tyr Ser Ala Ile Gly Ile Asp	
155 160 165	
CTG GCA TCT CAT GGG TTT ATA GTT GCT GCT GTA GAA CAC AGA GAT AGA	583
Leu Ala Ser His Gly Phe Ile Val Ala Ala Val Glu His Arg Asp Arg	
170 175 180	
TCT GCA TCT GCA ACT TAC TAT TTC AAG GAC CAA TCT GCT GCA GAA ATA	631
Ser Ala Ser Ala Thr Tyr Tyr Phe Lys Asp Gln Ser Ala Ala Glu Ile	
185 190 195	
GGG GAC AAG TCT TGG CTC TAC CTT AGA ACC CTG AAA CAA GAG GAG GAG	679
Gly Asp Lys Ser Trp Leu Tyr Leu Arg Thr Leu Lys Gln Glu Glu Glu	
200 205 210	
ACA CAT ATA CGA AAT GAG CAG GTA CGG CAA AGA GCA AAA GAA TGT TCC	727



-continued

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

```

Met Ile Thr Ile Arg Gly Ser Val His Gln Asn Phe Ala Asp Phe Th
1           5           10          15

Phe Ala Thr Gly
                20
  
```

The invention claimed is:

1. A method of diagnosing a patient's susceptibility to atherosclerotic disease which comprises assaying a sample from a patient for Lp-PLA<sub>2</sub> enzyme activity wherein elevated Lp-PLA<sub>2</sub> enzyme activity in the sample of the patient as compared to a standard is indicative of the patient's susceptibility to atherosclerotic disease.

2. The method according to claim 1 wherein the atherosclerotic disease is selected from the group comprising atherosclerosis, stroke and myocardial infarction.

3. The method according to claim 1 wherein the atherosclerotic disease is atherosclerosis.

4. The method according to claim 1 wherein the atherosclerotic disease is stroke.

5. The method according to claim 1 wherein the atherosclerotic disease is myocardial infarction.

6. The method according to claim 1 wherein the sample is selected from the group comprising blood, serum and plasma.

7. The method according to claim 1 wherein the sample is blood.

8. The method according to claim 1 wherein the sample is serum.

9. The method according to claim 1 wherein the sample is plasma.

10. The method according to claim 1 wherein the sample is assayed in a 96 well titre plate.

11. The method of according to claim 1 wherein the sample is assayed for enzymatic activity via an absorbance assay.

12. The method according to claim 11 wherein absorbance at 400 nm or 405 nm is monitored.

13. The method of according to claim 1 wherein the sample is assayed for enzymatic activity via an absorbance assay measuring rate of turnover of an artificial substrate.

14. The method according to claim 13 wherein absorbance at 400 nm or 405 nm is monitored.

15. The method according to claim 13 wherein the artificial substrate comprises a nitrophenyl group.

16. A method of diagnosis of a patient's susceptibility to atherosclerotic disease which comprises analyzing a sample from a patient for the presence of the enzyme Lp-PLA<sub>2</sub> by assaying the sample for enzyme activity via an absorbance assay measuring rate of turnover of an artificial substrate wherein elevated Lp-PLA<sub>2</sub> enzyme activity in the sample of the patient as compared to a standard are indicative of the patient's susceptibility to atherosclerotic disease.

17. The method of claim 16 wherein the artificial substrate comprises a nitrophenyl group.

18. A method according to claim 16 wherein the disease is atherosclerosis, stroke or myocardial infarction.

19. The method according to claim 16 wherein the sample is selected from the group consisting of blood, serum, and plasma.

20. The method according to claim 16 wherein the sample is blood.

21. The method according to claim 16 wherein the sample is plasma.

22. The method according to claim 16 wherein the sample is serum.

23. The method according to claim 16 wherein the disease is atherosclerosis.

24. The method according to claim 16 wherein the disease is stroke.

25. The method according to claim 16 wherein the disease is myocardial infarction.

26. The method according to claim 16 wherein the absorbance assay is performed in a 96 well titre plate.


27. The method according to claim 16 wherein absorbance at 400 nm or 405 nm is monitored.

\* \* \* \* \*

专利名称(译)	脂蛋白相关磷脂酶A2，其抑制剂及其在诊断和治疗中的用途		
公开(公告)号	<a href="#">US7217535</a>	公开(公告)日	2007-05-15
申请号	US10/973467	申请日	2004-10-26
[标]申请(专利权)人(译)	MACPHEE COLIN ^ h TEW DAVID 摹		
申请(专利权)人(译)	MACPHEE COLIN H. TEW DAVID G.		
当前申请(专利权)人(译)	史克必成P.L.C.		
[标]发明人	MACPHEE COLIN H TEW DAVID GRAHAM		
发明人	MACPHEE, COLIN H. TEW, DAVID GRAHAM		
IPC分类号	C12Q1/34 G01N33/53 A61K31/739 A61K38/00 A61K38/55 A61K39/395 A61P9/10 C07H21/04 C07K14/47 C07K16/40 C12N1/21 C12N9/16 C12N9/18 C12N9/20 C12N15/00 C12N15/09 C12N15/55 C12N15/74 C12P21/08 C12Q1/44 G01N33/573		
CPC分类号	A61K38/005 C12N9/18 C12N9/20 C12Q1/44 G01N33/6893 G01N2800/323 G01N2800/324 Y10S435/81		
优先权	PCT/GB1994/001374 1994-06-24 WO 1993013144 1993-06-25 GB 1994000413 1994-01-11 GB		
其他公开文献	US20050106611A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

#### 摘要(译)

纯化形式的酶Lp-PLA2，编码Lp-PLA2的分离的核酸分子，酶Lp-PLA2的抑制剂在治疗中的用途以及筛选化合物以鉴定那些化合物的方法。



(I)