



US 20050221348A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2005/0221348 A1**
Ray et al. (43) **Pub. Date: Oct. 6, 2005**

(54) **METHODS AND COMPOSITIONS FOR
DIAGNOSIS, STRATIFICATION, AND
MONITORING OF ALZHEIMER'S DISEASE
AND OTHER NEUROLOGICAL DISORDERS
IN BODY FLUIDS**

filed on Apr. 30, 2004. Provisional application No.
60/566,782, filed on Apr. 30, 2004.

Publication Classification

(76) Inventors: **Sandip Ray**, San Francisco, CA (US);
Anton Wyss-Coray, Belmont, CA (US)

(51) **Int. Cl.⁷** **C12Q 1/68**; G01N 33/53;
G01N 33/567
(52) **U.S. Cl.** **435/6**; 435/7.2

Correspondence Address:
MORRISON & FOERSTER LLP
755 PAGE MILL RD
PALO ALTO, CA 94304-1018 (US)

(57) **ABSTRACT**

(21) Appl. No.: **10/993,813**

(22) Filed: **Nov. 19, 2004**

Related U.S. Application Data

(60) Provisional application No. 60/523,796, filed on Nov.
19, 2003. Provisional application No. 60/566,783,

The inventors have discovered a collection of proteinaceous biomarkers ("AD biomarkers") which can be measured in peripheral biological fluid samples to aid in the diagnosis of neurodegenerative disorders, particularly Alzheimer's disease and mild cognitive impairment (MCI). The invention further provides methods of identifying candidate agents for the treatment of Alzheimer's disease by testing prospective agents for activity in modulating AD biomarker levels.




Figure 1A




Figure 1B




Figure 1C




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7

METHODS AND COMPOSITIONS FOR DIAGNOSIS, STRATIFICATION, AND MONITORING OF ALZHEIMER'S DISEASE AND OTHER NEUROLOGICAL DISORDERS IN BODY FLUIDS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit of U.S. Provisional Patent Application No. 60/523,796, filed Nov. 19, 2003, U.S. Provisional Patent Application Ser. No. 60/566,783, filed Apr. 30, 2004, and U.S. Provisional Patent Application No. 60/566,782, filed Apr. 30, 2004, all of which are incorporated by reference herein in their entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] Not applicable.

REFERENCE TO A COMPACT DISK APPENDIX

[0003] Not applicable

BACKGROUND OF THE INVENTION

[0004] An estimated 4.5 million Americans have Alzheimer's Disease ("AD"). By 2050, the estimated range of AD prevalence will be 11.3 million to 16 million. Currently, the societal cost of AD to the U.S. is \$100 billion per year, including \$61 billion born by U.S. businesses. Neither Medicare nor most private health insurance covers the long-term care most patients need.

[0005] Alzheimer's Disease is a neurodegenerative disease of the central nervous system associated with progressive memory loss resulting in dementia. Two pathological characteristics are observed in AD patients at autopsy: extracellular plaques and intracellular tangles in the hippocampus, cerebral cortex, and other areas of the brain essential for cognitive function. Plaques are formed mostly from the deposition of amyloid beta ("A β "), a peptide derived from amyloid precursor protein ("APP"). Filamentous tangles are formed from paired helical filaments composed of neurofilament and hyperphosphorylated tau protein, a microtubule-associated protein. It is not clear, however, whether these two pathological changes are only associated with the disease or truly involved in the degenerative process. Late-onset/sporadic AD has virtually identical pathology to inherited early-onset/familial AD (FAD), thus suggesting common pathogenic pathways for both forms of AD. To date, genetic studies have identified three genes that cause autosomal dominant, early-onset AD, amyloid precursor protein ("APP"), presenilin 1 ("PS1"), and presenilin 2 ("PS2"). A fourth gene, apolipoprotein E ("ApoE"), is the strongest and most common genetic risk factor for AD, but does not necessarily cause it. All mutations associated with APP and PS proteins can lead to an increase in the production of A β peptides, specifically the more amyloidogenic form, A β ₄₂. In addition to genetic influences on amyloid plaque and intracellular tangle formation, environmental factors (e.g., cytokines, neurotoxins, etc.) may also play important roles in the development and progression of AD.

[0006] The main clinical feature of AD is a progressive cognitive decline leading to memory loss. The memory dysfunction involves impairment of learning new informa-

tion which is often characterized as short-term memory loss. In the early (mild) and moderate stages of the illness, recall of remote well-learned material may appear to be preserved, but new information cannot be adequately incorporated into memory. Disorientation to time is closely related to memory disturbance.

[0007] Language impairments are also a prominent part of AD. These are often manifest first as word finding difficulty in spontaneous speech. The language of the AD patient is often vague, lacking in specifics and may have increased automatic phrases and cliches. Difficulty in naming everyday objects is often prominent. Complex deficits in visual function are present in many AD patients, as are other focal cognitive deficits such as apraxia, acalculia and left-right disorientation. Impairments of judgment and problems solving are frequently seen.

[0008] Non-cognitive or behavioral symptoms are also common in AD and may account for an event larger proportion of caregiver burden or stress than the cognitive dysfunction. Personality changes are commonly reported and range from progressive passivity to marked agitation. Patients may exhibit changes such as decreased expressions of affection. Depressive symptoms are present in up to 40%. A similar rate for anxiety has also been recognized. Psychosis occurs in 25%. In some cases, personality changes may predate cognitive abnormality.

[0009] Currently, the primary method of diagnosing AD in living patients involves taking detailed patient histories, administering memory and psychological tests, and ruling out other explanations for memory loss, including temporary (e.g., depression or vitamin B₁₂ deficiency) or permanent (e.g., stroke) conditions. These clinical diagnostic methods, however, are not foolproof.

[0010] One obstacle to diagnosis is pinpointing the type of dementia; AD is only one of seventy conditions that produce dementia. Because of this, AD cannot be diagnosed with complete accuracy until after death, when autopsy reveals the disease's characteristic amyloid plaques and neurofibrillary tangles in a patient's brain. In addition, clinical diagnostic procedures are only helpful after patients have begun displaying significant, abnormal memory loss or personality changes. By then, a patient has likely had AD for years.

[0011] Given the magnitude of the public health problem posed by AD, considerable research efforts have been undertaken to elucidate the etiology of AD as well as to identify biomarkers (secreted proteins or metabolites) that can be used to diagnose and/or predict whether a person is likely to develop AD. Because AD the CNS is relatively isolated from the other organs and systems of the body, most research (in regards to both disease etiology and biomarkers) has focused on events, gene expression, biomarkers, etc. within the central nervous system. With regards to biomarkers, the proteins amyloid beta and tau are probably the most well characterized. Research has shown that cerebrospinal fluid ("CSF") samples from AD patients contain higher than normal amounts of tau, which is released as neurons degenerate, and lower than normal amounts of beta amyloid, presumably because it is trapped in the brain in the form of amyloid plaques. Because these biomarkers are released into CSF, a lumbar puncture (or "spinal tap") is required to obtain a sample for testing.

[0012] A number of U.S. patents have been issued relating to methods for diagnosing AD, including U.S. Pat. Nos.

4,728,605, 5,874,312, 6,027,896, 6,114,133, 6,130,048, 6,210,895, 6,358,681, 6,451,547, 6,461,831, 6,465,195, 6,475,161, and 6,495,335. Additionally, a number of reports in the scientific literature relate to certain biochemical markers and their correlation/association with AD, including Fahnestock et al., 2002, *J. Neural. Transm. Suppl.* 2002(62):241-52; Masliah et al., 1195, *Neurobiol. Aging* 16(4):549-56; Power et al., 2001, *Dement. Geriatr. Cogn. Disord.* 12(2):167-70; and Burbach et al., 2004, *J. Neurosci.* 24(10):2421-30. Additionally, Li et al. (2002, *Neuroscience* 113(3):607-15) and Sanna et al. (2003, *J. Clin. Invest.* 111(2):241-50) have investigated Leptin in relation to memory and multiple sclerosis, respectively.

[0013] All patents and publications cited herein are incorporated by reference in their entirety.

BRIEF SUMMARY OF THE INVENTION

[0014] The inventors have discovered a collection of biochemical markers, present in the serum of individuals, which are altered in individuals with Alzheimer's Disease ("AD"). Accordingly, these biomarkers ("AD diagnosis biomarkers") may be used to assess cognitive function, to diagnose or aid in the diagnosis of AD and/or to measure progression of AD in AD patients. AD diagnosis markers may be used individually or in combination for diagnosing or aiding in the diagnosis of AD. The invention provides methods for the diagnosis of AD or aiding the diagnosis of AD in an individual by measuring the amount of one or more AD diagnosis biomarkers in a biological fluid sample, such as a peripheral biological fluid sample from the individual and comparing the measured amount with a reference value for each AD diagnosis biomarker measured. The information thus obtained may be used to aid in the diagnosis or to diagnose AD in the individual. Accordingly, the present invention provides a method of aiding diagnosis of Alzheimer's disease ("AD"), comprising comparing a measured level of at least one AD diagnosis biomarker in a biological fluid sample from an individual to a reference level for the biomarker, wherein the AD diagnosis biomarker is selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In some examples, the AD diagnosis biomarker is selected from the group consisting of basic fibroblast growth factor (bFGF); BB homodimeric platelet derived growth factor (PDGF-BB); brain derived neurotrophic factor (BDNF); epidermal growth factor (EGF); fibroblast growth factor 6 (FGF-6); interleukin-3 (IL-3); soluble interleukin-6 receptor (sIL-6R); leptin (also known as ob); macrophage inflammatory protein-1 delta (MIP-1δ); macrophage stimulating protein alpha chain (MSP-α); neurotrophin-3 (NT-3); neutrophil activating peptide-2 (NAP-2); RANTES; soluble tumor necrosis factor receptor-2 (sTNF RII); stem cell factor (SCF); thrombopoietin (TPO); tissue inhibitor of metalloproteases-1 (TIMP-1); tissue inhibitor of metalloproteases-2 (TIMP-2); transforming growth factor-beta 3 (TGF-β3); and tumor necrosis factor beta (TNF-β). In other examples, the AD diagnosis marker is selected from the group consisting of BDNF, sIL-6R, IL-8, leptin, MIP-1δ, PDGF-BB, and TIMP-1. In

yet other examples, the AD diagnosis marker is selected from the group consisting of sIL-6R, IL-8, and TIMP-1. In further examples, the AD diagnosis marker is selected from the group consisting of BDNF, MIP-1δ, and TIMP-1. In additional examples, the AD diagnosis marker is selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES. In additional examples, the AD diagnosis marker comprises BDNF, PDGF-BB, leptin and RANTES.

[0015] Provided herein are methods of aiding diagnosis of Alzheimer's disease ("AD"), comprising comparing a measured level of at least four AD diagnosis biomarkers, wherein said biomarkers comprise BDNF, PDGF-BB, leptin and RANTES, in a biological fluid sample from an individual to a reference level for each AD diagnosis biomarker. In some examples, AD is diagnosed when BDNF is decreased at least about 20% as compared to a reference level of BDNF. In other examples, AD is diagnosed when Leptin is decreased at least about 25% as compared to a reference level of Leptin. In additional examples, AD is diagnosed when RANTES is decreased at least about 16% as compared to a reference level of RANTES. In further examples, severe AD is diagnosed when PDGF-BB is decreased at least about 85% as compared to a reference level of PDGF-BB. In yet further examples, the biological fluid sample is a peripheral biological fluid sample.

[0016] Provided herein are methods for monitoring progression of Alzheimer's disease (AD) in an AD patient, comprising: comparing a measured level of at least one AD diagnosis biomarker in a biological fluid sample from an individual to a reference level for the biomarker, wherein the AD diagnosis biomarker is selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In some examples, the AD diagnosis biomarker is selected from the group consisting of basic fibroblast growth factor (bFGF); BB homodimeric platelet derived growth factor (PDGF-BB); brain derived neurotrophic factor (BDNF); epidermal growth factor (EGF); fibroblast growth factor 6 (FGF-6); interleukin-3 (IL-3); soluble interleukin-6 receptor (sIL-6R); leptin (also known as ob); macrophage inflammatory protein-1 delta (MIP-1δ); macrophage stimulating protein alpha chain (MSP-α); neurotrophin-3 (NT-3); neutrophil activating peptide-2 (NAP-2); RANTES; soluble tumor necrosis factor receptor-2 (sTNF RII); stem cell factor (SCF); thrombopoietin (TPO); tissue inhibitor of metalloproteases-1 (TIMP-1); tissue inhibitor of metalloproteases-2 (TIMP-2); transforming growth factor-beta 3 (TGF-β3); and tumor necrosis factor beta (TNF-β). In other examples, the AD diagnosis marker is selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES.

[0017] The inventors have also discovered methods of identifying individuals with mild cognitive deficit (MCI), a clinically recognized disorder considered distinct from AD in which cognition and memory are mildly deficient. The inventors have found that the biomarker RANTES is decreased in individuals with MCI. Individuals with MCI can be distinguished from those with AD by measuring biomarkers which are reduced in AD patients, but not those

individuals with MCI (e.g., Leptin). Accordingly, the invention provides methods for diagnosing or aiding in the diagnosis of MCI by obtaining a measured value for the level of RANTES in a peripheral biological fluid sample and comparing that measured value against a reference value. In certain embodiments, such methods include obtaining a measuring value for Leptin levels in the peripheral biological fluid sample and comparing that measured level against a reference value. The information thus obtained may be used to aid in the diagnosis or to diagnose MCI in the individual.

[0018] Further, the inventors have discovered methods of stratifying AD patients (i.e., sorting individuals with a probable diagnosis of AD or diagnosed with AD into different classes of AD) by obtaining measured values for brain derived neurotrophic factor (BDNF) and BB-homodimer platelet derived growth factor (PDGF-BB) levels in a peripheral biological fluid sample from an AD patient. The measured levels of these two biomarkers are compared with reference values. The information thus obtained may be used to aid in stratification of the AD diagnosis (or probable AD diagnosis) of the individual. Accordingly, the present invention provides methods for stratifying Alzheimer's disease (AD) in an individual, comprising comparing measured values for brain derived neurotrophic factor (BDNF) and BB homodimeric platelet derived growth factor (PDGF-BB) levels in a biological fluid sample from said patient with reference values for BDNF and PDGF-BB. In some examples, the biological fluid sample is a peripheral fluid sample, including blood, serum or plasma. In other examples, the method further comprises comparing measured values for leptin and Rantes levels with reference values for leptin and Rantes, wherein reference values for BDNF, PDGF-BB, leptin and Rantes are for samples from individuals with MMSE scores from 25 to 28, wherein an increase in leptin and PDGF-BB levels and wherein levels of BDNF and RANTES stay substantially the same indicate mild AD as indicated by an MMSE score of 20-25. In additional examples, the method further comprises comparing measured values for leptin and Rantes levels with reference values for leptin and Rantes, wherein reference values for BDNF, PDGF-BB, leptin and Rantes are for samples from individuals with MMSE scores from 20-25, wherein a decrease in Rantes, BDNF, and PDGF levels and wherein levels of Leptin stays substantially the same indicate moderate AD as indicated by an MMSE score of 10-20.

[0019] In one aspect, the invention provides methods of aiding in the diagnosis of Alzheimer's disease ("AD") by obtaining a measured level of at least one AD diagnosis biomarker in a peripheral biological fluid sample from an individual, where the AD diagnosis biomarker is from the group consisting of basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), Leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1 δ), macrophage stimulating protein alpha chain (MSP- α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF- β 13), and tumor

necrosis factor beta (TNF- β), and comparing the measured level to the reference level. In some embodiments, measured levels are obtained for at least two, three, four, or five AD diagnosis biomarkers. In some embodiments, the comparison of the measured value and the reference value includes calculating a fold difference between the measured value and the reference value. In some embodiments the measured value is obtained by measuring the level of the AD diagnosis biomarker(s) in the sample, while in other embodiments the measured value is obtained from a third party. Also provided are methods of aiding in the diagnosis of Alzheimer's disease ("AD") by comparing a measured level of at least one AD diagnosis biomarker in a peripheral biological fluid sample from an individual with a reference level. Further provided are methods of aiding in the diagnosis of Alzheimer's disease ("AD") by measuring a level of at least one AD diagnosis biomarker in a peripheral biological fluid sample from an individual, wherein a decrease as compared to a reference level suggests a diagnosis of AD.

[0020] In another aspect, the invention provides methods for aiding in the diagnosis of mild cognitive impairment (MCI) by obtaining a measured level for RANTES in a peripheral biological fluid sample from an individual, and comparing the measured level to a reference level. In some embodiments, the method for aiding in the diagnosis of MCI also includes obtaining a measured value for Leptin in the peripheral biological fluid sample and comparing measured value for Leptin to a reference level. In certain embodiments, the measured value is obtained by measuring the level of RANTES (and/or Leptin) in the sample, while in other embodiments, the measured value(s) is obtained from a third party. Also provided are methods of aiding in the diagnosis of mild cognitive impairment (MCI) by comparing a measured level for RANTES, and optionally Leptin, in a peripheral biological fluid sample from an individual with a reference level. Further provided are methods for aiding in the diagnosis of MCI by measuring a level for RANTES, and optionally Leptin, in a peripheral biological fluid sample from an individual, wherein a reduction in the RANTES level as compared to a reference level suggests a diagnosis of MCI (in embodiments in which Leptin is measured, a Leptin level that is equal to or greater than the reference level also suggests MCI).

[0021] In a further aspect, the invention provides methods for monitoring progression of Alzheimer's disease (AD) in an AD patient by obtaining a measured value for Leptin in a peripheral biological fluid sample; and comparing said measured value for Leptin with a reference value. In certain embodiments, the measured value is obtained by measuring the level of Leptin in the sample to produce, while in other embodiments, the measured value is obtained from a third party. Also provided are methods for monitoring progression of AD in an AD patient by comparing a measured value for Leptin in a peripheral biological fluid sample with a reference value. Further provided are methods for monitoring progression of AD in an AD patient by measuring a level for Leptin in a peripheral biological fluid sample, wherein a decrease in Leptin as compared with a reference value suggests progression (increased severity) of the AD.

[0022] In another aspect, the invention provides methods for stratifying AD in an AD patient. In some embodiments, stratification between mild and more advanced AD is carried out by obtaining a measured value for brain derived neu-

rotrophic factor (BDNF) levels in a peripheral biological fluid sample from an AD patient, and comparing the measured value with reference values for BDNF. In other embodiments, stratification between mild, moderate, and severe AD is carried out by obtaining levels for BDNF and BB homodimeric platelet derived growth factor (PDGF-BB), and comparing the measured levels with reference levels for BDNF and PDGF-BB. In certain embodiments, the measured value is obtained by measuring the level(s) of BDNF (and PDGF-BB) in the sample to produce the measured value(s), while in other embodiments, the measured value(s) is obtained from a third party. Also provided are methods for stratifying AD in an AD patient by comparing a BDNF (and, optionally, PDGF-BB) level in a peripheral biological fluid sample from an AD patient with a reference value for BDNF (and PDGF-BB when appropriate). Further provided are methods for stratifying AD in an AD patient by measuring a BDNF level (and, optionally, a PDGF-BB level) in a peripheral biological fluid sample, wherein a low level of BDNF (as compared to a reference value) suggests mild AD, a high level of BDNF (as compared to a reference value) suggests more advanced AD, a high level of BDNF and a low level of PDGF-BB (as compared to reference values) suggests moderate AD, and a high level of BDNF and a high level of PDGF-BB (as compared to reference values) suggests severe AD.

[0023] In some embodiments, the peripheral biological fluid sample is a blood sample. In certain embodiments the peripheral biological fluid sample is a plasma sample. In other embodiments, the peripheral biological fluid sample is a serum sample.

[0024] In yet another aspect, the invention provides methods of identifying candidate agents for treatment of Alzheimer's Disease by assaying a prospective candidate agent for activity in modulating an AD biomarker, where the AD biomarker is from the group consisting of basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), Leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), tumor necrosis factor beta (TNF-β). Provided herein are methods of identifying a candidate agent for treatment of Alzheimer's Disease, comprising: assaying a prospective candidate agent for activity in modulating an AD biomarker, said AD biomarker selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In some examples, the AD biomarkers are selected from the group consisting of BDNF, PDGF-BB, Leptin and RANTES.

[0025] In a further aspect, the invention provides kits for diagnosing Alzheimer's disease (AD) including at least one reagent specific for an AD diagnosis marker, where the AD diagnosis biomarker is from the group consisting of basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), Leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), tumor necrosis factor beta (TNF-β), and instructions for carrying out a method of aiding in the diagnosis of AD described herein. Provided herein are kits comprising at least one reagent specific for at least one AD diagnosis marker, said at least one AD diagnosis biomarker selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R and instructions for carrying out methods provided herein. Additionally, provided herein are sets of reference values for AD diagnosis biomarkers comprising BDNF, PDGF-BB, Leptin and RANTES and set of reagents specific for AD diagnosis biomarkers, wherein said biomarkers comprise BDNF, PDGF-BB, Leptin and RANTES.

[0026] In another aspect, the invention provides kits for identifying individuals with mild cognitive impairment (MCI) including at least one reagent specific for RANTES; and instructions for carrying out method of aiding in the diagnosis of MCI described herein. In certain embodiments, kits for identifying individuals with MCI may also include a reagent specific for Leptin.

[0027] In yet another aspect, the invention provides kits for monitoring progression of Alzheimer's disease (AD) in AD patients including at least one reagent specific for Leptin; and instructions for carrying out a method of monitoring AD progression described herein.

[0028] In a further aspect, the invention provides kits for stratifying an Alzheimer's disease (AD) patients including at least one reagent specific for brain derived neurotrophic factor (BDNF), at least one reagent specific for BB homodimeric platelet derived growth factor (PDGF-BB), and instructions for carrying out a method of stratifying an AD patient described herein. In yet further examples, kits comprise AD diagnosis markers are selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES. In further examples of kits, the reagent specific for the AD diagnosis biomarker is an antibody, or fragment thereof, that is specific for said AD diagnosis biomarker. In further examples kits further comprise at least one reagent specific for a biomarker that measures sample characteristics.

[0029] Provided herein are surfaces comprising attached thereto, at least one reagent specific for each AD diagnosis

biomarker in a set of AD diagnosis biomarkers, wherein said set of AD diagnosis biomarkers comprises BDNF, PDGF-BB, leptin and RANTES. Provided herein are surfaces comprising attached thereto, at least one reagent specific for each AD diagnosis biomarker in a set of AD diagnosis biomarkers, wherein said set of AD diagnosis biomarkers consists of BDNF, PDGF-BB, leptin and RANTES; and at least one reagent specific for a biomarker that measures sample characteristics. In further examples, provided herein are surfaces wherein said reagent specific for said AD diagnosis biomarker is an antibody, or fragment thereof, that is specific for said AD diagnosis biomarker.

[0030] Provided herein are combinations comprising the surfaces as described herein having attached thereto at least one reagent specific for each AD diagnosis biomarker and a peripheral biological fluid sample from an individual. In some examples, the individual is at least 60, 65, 70, 75, 80, or 85 years of age.

[0031] Provided herein are methods for obtaining values for the comparison of the measured level to the reference level of biological fluid samples. The present invention provides computer readable formats comprising the values obtained by the methods described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032] **FIGS. 1A-1C** show ELISA results for 3 proteins, **FIG. 1A** BDNF; **FIG. 1B** Leptin; and **FIG. 1C** RANTES, selected from the list from Table 3 shown herein in the Examples. 95 plasma samples from individuals having AD and having mean MMSE scores of 20, and mean age of 74, was compared to plasma sample from 88 age-matched controls having mean MMSE score of 30. Non-parametric, unpaired t tests comparing the mean concentration of each protein was used to determine statistical significance (p-value).

[0033] **FIG. 2** shows a Cell Bar Chart for concentration of BDNF in plasma. (Cell Bar Chart Grouping Variable(s): stage Error Bars: ± 1 Standard error(s) Inclusion criteria: Sparks from Center All)

[0034] **FIG. 3** shows BDNF in control vs AD for male and female. (Cell Bar Chart Grouping Variable(s): Disease Split By: sex Error Bars: ± 1 Standard Error(s) Row exclusion: Center All)

[0035] **FIG. 4** shows RANTES concentration in plasma. (Cell Bar Chart Grouping Variable(s): stage Error Bars: ± 1 Standard Error(s) Row exclusion: Center All)

[0036] **FIG. 5** shows concentration of Leptin in plasma. (Cell Bar Chart Grouping Variable(s): stage Error Bars: ± 1 Standard Error(s) Row exclusion: Center All)

[0037] **FIG. 6** shows PDGF-BB concentration in plasma. (Cell Bar Chart Grouping Variable(s): stage Error Bars: ± 1 Standard Error(s) Row exclusion: Center All)

[0038] **FIG. 7** shows BDNF concentration in plasma. (Cell Bar Chart Grouping Variable(s): stage Error Bars: ± 1 Standard Error(s) Row exclusion: Center All)

DETAILED DESCRIPTION OF THE INVENTION

[0039] Inflammation and injury responses are invariably associated with neuron degeneration in AD, PD, frontotem-

poral dementia, cerebrovascular disease, multiple sclerosis, and neuropathies. The brain and CNS are not only immunologically active in there own accord, but also have complex peripheral immunologic interactions. Fiala et al. (1998 Mol Med. July; 4(7):480-9) has shown that in Alzheimer's disease, alterations in the permeability of the blood-brain barrier and chemotaxis, in part mediated by chemokines and cytokines, may permit the recruitment and transendothelial passage of peripheral cells into the brain parenchyma. A paradigm of the blood-brain barrier was constructed utilizing human brain endothelial and astroglial cells with the anatomical and physiological characteristics observed in vivo. This model was used to test the ability of monocytes/macrophages to transmigrate when challenged by A beta 1-42 on the brain side of the blood-brain barrier model. In that model A beta 1-42 and monocytes on the brain side potentiated monocyte transmigration from the blood side to the brain side. In some individuals, circulating monocytes/macrophages, when recruited by chemokines produced by activated microglia and macrophages, could add to the inflammatory destruction of the brain in Alzheimer's disease.

[0040] The inventors assert that the monitoring for relative concentrations of many secreted markers measured simultaneously in the serum is a more sensitive method for monitoring the progression of disease than the absolute concentration of any single biochemical markers have been able to achieve. A composite or array embodying the use of 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200 markers in Table 7 simultaneously, consisting of antibodies bound to a solid support or protein bound to a solid support, for the detection of inflammation and injury response markers associated with neuron degeneration in AD, PD, frontotemporal dementia, cerebrovascular disease, multiple sclerosis, and neuropathies.

[0041] The inventors have discovered a collection of biochemical markers (collectively termed "AD biomarkers") useful for diagnosis of AD, aiding in diagnosis of AD, monitoring AD in AD patients (e.g., tracking disease progression in AD patients, which may be useful for tracking the effect of medical or surgical therapy in AD patients), stratifying AD patients, and diagnosing or aiding in the diagnosis of mild cognitive impairment (MCI) as well as diagnosing or aiding in the diagnosis of cognitive impairment. The AD biomarkers are present in biological fluids of individuals. In some examples, the AD biomarkers are present in peripheral biological fluids (e.g., blood) of individuals, allowing collection of samples by procedures that are relatively non-invasive, particularly as compared to the lumbar puncture procedure commonly used to collect cerebrospinal fluid samples.

[0042] Definitions

[0043] As used herein, the terms "Alzheimer's patient", "AD patient", and "individual diagnosed with AD" all refer to an individual who has been diagnosed with AD or has been given a probable diagnosis of Alzheimer's Disease (AD).

[0044] As used herein, the phrase "AD biomarker" refers to a biomarker that is an AD diagnosis biomarker.

[0045] The term "AD biomarker polynucleotide", as used herein, refers to any of: a polynucleotide sequence encoding

a AD biomarker, the associated trans-acting control elements (e.g., promoter, enhancer, and other gene regulatory sequences), and/or mRNA encoding the AD biomarker.

[0046] As used herein, methods for “aiding diagnosis” refer to methods that assist in making a clinical determination regarding the presence, or nature, of the AD or MCI, and may or may not be conclusive with respect to the definitive diagnosis. Accordingly, for example, a method of aiding diagnosis of AD can comprise measuring the amount of one or more AD biomarkers in a biological sample from an individual.

[0047] As used herein, the term “stratifying” refers to sorting individuals into different classes or strata based on the features of a neurological disease. For example, stratifying a population of individuals with Alzheimer’s disease involves assigning the individuals on the basis of the severity of the disease (e.g., mild, moderate, advanced, etc.).

[0048] As used herein, the term “predicting” refers to making a finding that an individual has a significantly enhanced probability of developing a certain neurological disease.

[0049] As used herein, the phrase “neurological disease” refers to a disease or disorder of the central nervous system. Neurological diseases include multiple sclerosis, neuropathies, and neurodegenerative disorders such as AD, Parkinson’s disease, amyotrophic lateral sclerosis (ALS), mild cognitive impairment (MCI) and frontotemporal dementia.

[0050] As used herein, “biological fluid sample” encompasses a variety of fluid sample types obtained from an individual and can be used in a diagnostic or monitoring assay. The definition encompasses blood, cerebral spinal fluid (CSF), urine and other liquid samples of biological origin. The definition also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as proteins or polynucleotides.

[0051] As used herein, the term “peripheral biological fluid sample” refers to a biological fluid sample that is not derived from the central nervous system (i.e., is not a CSF sample) and includes blood samples and other biological fluids not derived from the CNS.

[0052] A “blood sample” is a biological sample which is derived from blood, preferably peripheral (or circulating) blood. A blood sample may be, for example, whole blood, plasma or serum.

[0053] An “individual” is a mammal, more preferably a human. Mammals include, but are not limited to, humans, primates, farm animals, sport animals, rodents and pets.

[0054] A “Normal” individual or sample from a “Normal” individual as used herein for quantitative and qualitative data refers to an individual who has or would be assessed by a physician as not having AD or MCI, and has an Mini-Mental State Examination (MMSE) (referenced in Folstein et al., *J. Psychiatr. Res* 1975; 12:1289-198) score or would achieve a MMSE score in the range of 25-30. A “Normal” individual is generally age-matched within a range of 5 to 10 years, including but not limited to an individual that is age-matched, with the individual to be assessed.

[0055] An “individual with mild AD” is an individual who (a) has been diagnosed with AD or has been given a

diagnosis of probable AD, and (b) has either been assessed with the Mini-Mental State Examination (MMSE) (referenced in Folstein et al., *J. Psychiatr. Res* 1975; 12:1289-198) and scored 22-27 or would achieve a score of 22-27 upon MMSE testing. Accordingly, “mild AD” refers to AD in a individual who has either been assessed with the MMSE and scored 22-27 or would achieve a score of 22-27 upon MMSE testing.

[0056] An “individual with moderate AD” is an individual who (a) has been diagnosed with AD or has been given a diagnosis of probable AD, and (b) has either been assessed with the MMSE and scored 16-21 or would achieve a score of 16-21 upon MMSE testing. Accordingly, “moderate AD” refers to AD in a individual who has either been assessed with the MMSE and scored 16-21 or would achieve a score of 16-21 upon MMSE testing.

[0057] An “individual with severe AD” is an individual who (a) has been diagnosed with AD or has been given a diagnosis of probable AD, and (b) has either been assessed with the MMSE and scored 12-15 or would achieve a score of 12-15 upon MMSE testing. Accordingly, “severe AD” refers to AD in a individual who has either been assessed with the MMSE and scored 12-15 or would achieve a score of 12-15 upon MMSE testing.

[0058] As used herein, the term “treatment” refers to the alleviation, amelioration, and/or stabilization of symptoms, as well as delay in progression of symptoms of a particular disorder. For example, “treatment” of AD includes any one or more of: elimination of one or more symptoms of AD, reduction of one or more symptoms of AD, stabilization of the symptoms of AD (e.g., failure to progress to more advanced stages of AD), and delay in progression (i.e., worsening) of one or more symptoms of AD.

[0059] As used herein, the phrase “fold difference” refers to a numerical representation of the magnitude difference between a measured value and a reference value for an AD biomarker. Fold difference is calculated mathematically by division of the numeric measured value with the numeric reference value. For example, if a measured value for an AD biomarker is 20 nanograms/milliliter (ng/ml), and the reference value is 10 ng/ml, the fold difference is 2 (20/10=2). Alternatively, if a measured value for an AD biomarker is 10 nanograms/milliliter (ng/ml), and the reference value is 20 ng/ml, the fold difference is 10/20 or -0.50 or -50%.

[0060] As used herein, a “reference value” can be an absolute value; a relative value; a value that has an upper and/or lower limit; a range of values; an average value; a median value, a mean value, or a value as compared to a particular control or baseline value. A reference value can be based on an individual sample value, such as for example, a value obtained from a sample from the individual with AD, MCI or cognitive impairment, but at an earlier point in time, or a value obtained from a sample from an AD patient other than the individual being tested, or a “normal” individual, that is an individual not diagnosed with AD. The reference value can be based on a large number of samples, such as from AD patients or normal individuals or based on a pool of samples including or excluding the sample to be tested.

[0061] As used herein, “a”, “an”, and “the” can mean singular or plural (i.e., can mean one or more) unless indicated otherwise.

[0062] Methods of the Invention

[0063] Methods for Identifying Biomarkers

[0064] The invention provides methods for identifying one or more biomarkers useful for diagnosis, aiding in diagnosis, stratifying, assessing risk, monitoring, and/or predicting a neurological disease. In certain aspects of the invention, levels of a group of biomarkers are obtained for a set of peripheral biological fluid samples from one or more individuals. The samples are selected such that they can be segregated into one or more subsets on the basis of a neurological disease (e.g., samples from normal individuals and those diagnosed with amyotrophic lateral sclerosis or samples from individuals with mild Alzheimer's disease and those with severe Alzheimer's disease). The measured values from the samples are compared to each other to identify those biomarkers which differ significantly amongst the subsets. Those biomarkers that vary significantly amongst the subsets may then be used in methods for aiding in the diagnosis, diagnosis, stratification, monitoring, and/or prediction of neurological disease. In other aspects of the invention, measured values for a set of peripheral biological fluid samples from one or more individuals (where the samples can be segregated into one or more subsets on the basis of a neurological disease) are compared, wherein biomarkers that vary significantly are useful for aiding in the diagnosis, diagnosis, stratification, monitoring, and/or prediction of neurological disease. In further aspects of the invention, levels of a set of peripheral biological fluid samples from one or more individuals (where the samples can be segregated into one or more subsets on the basis of a neurological disease) are measured to produced measured values, wherein biomarkers that vary significantly are useful for aiding in the diagnosis, diagnosis, stratification, monitoring, and/or prediction of neurological disease.

[0065] The instant invention utilizes a set of peripheral biological fluid samples, such as blood samples, that are derived from one or more individuals. The set of samples is selected such that it can be divided into one or more subsets on the basis of a neurological disease. The division into subsets can be on the basis of presence/absence of disease, stratification of disease (e.g., mild vs. moderate), or subclassification of disease (e.g., relapsing/remitting vs. progressive relapsing).

[0066] Biomarkers measured in the practice of the invention may be any proteinaceous biological marker found in a peripheral biological fluid sample. Table 7 contains a collection of exemplary biomarkers. Additional biomarkers are described herein.

[0067] Accordingly, the invention provides methods identifying one or more biomarkers which can be used to aid in the diagnosis, diagnose, detect, stratify, and/or predict neurological diseases such as neurodegenerative disorders. The methods of the invention are carried out by obtaining a set of measured values for a plurality of biomarkers from a set of peripheral biological fluid samples, where the set of peripheral biological fluid samples is divisible into at least two subsets in relation to a neurological disease, comparing said measured values between the subsets for each biomarker, and identifying biomarkers which are significantly different between the subsets.

[0068] The process of comparing the measured values may be carried out by any method known in the art,

including Significance Analysis of Microarrays, Tree Harvesting, CART, MARS, Self Organizing Maps, Frequent Item Set, or Bayesian networks.

[0069] In one aspect, the invention provides methods for identifying one or more biomarkers useful for the diagnosis of a neurological disease by obtaining measured values from a set of peripheral biological fluid samples for a plurality of biomarkers, wherein the set of peripheral biological fluid samples is divisible into subsets on the basis of a neurological disease, comparing the measured values from each subset for at least one biomarker; and identifying at least one biomarker for which the measured values are significantly different between the subsets. In some embodiments, the comparing process is carried out using Significance Analysis of Microarrays. In certain embodiments, the neurodegenerative disease is from the group consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease, and amyotrophic lateral sclerosis (ALS).

[0070] In another aspect, the invention provides methods for identifying at least one biomarker useful for aiding in the diagnosis of a neurological disease by obtaining measured values from a set of peripheral biological fluid samples for a plurality of biomarkers, wherein the set of peripheral biological fluid samples is divisible into subsets on the basis of a neurological disease, comparing the measured values from each subset for at least one biomarker; and identifying biomarkers for which the measured values are significantly different between the subsets.

[0071] In a further aspect, the invention provides methods for identifying at least one biomarker useful for the stratification of a neurological disease by obtaining measured values from a set of peripheral biological fluid samples for a plurality of biomarkers, wherein the set of peripheral biological fluid samples is divisible into subsets on the basis of strata of a neurological disease, comparing the measured values from each subset for at least one biomarker; and identifying biomarkers for which the measured values are significantly different between the subsets.

[0072] In another aspect, the invention provides methods for identifying at least one biomarker useful for the monitoring of a neurological disease by obtaining measured values from a set of peripheral biological fluid samples for a plurality of biomarkers, wherein the set of peripheral biological fluid samples is divisible into subsets on the basis of strata of a neurological disease, comparing the measured values from each subset for at least one biomarker; and identifying biomarkers for which the measured values are significantly different between the subsets.

[0073] In yet another aspect, the invention provides methods for identifying at least one biomarker useful for the prediction of a neurological disease by obtaining measured values from a set of peripheral biological fluid samples for a plurality of biomarkers, wherein the set of peripheral biological fluid samples is divisible into subsets on the basis of a neurological disease, comparing the measured values from each subset for at least one biomarker; and identifying biomarkers for which the measured values are significantly different between the subsets.

[0074] Methods of Assessing Cognitive Function

[0075] Provided herein are methods for assessing cognitive function, assessing cognitive impairment, diagnosing or

aiding diagnosis of cognitive impairment by obtaining measured levels of one or more AD diagnosis biomarkers in a biological fluid sample from an individual, such as for example, a peripheral biological fluid sample from an individual, and comparing those measured levels to reference levels. Reference to "AD diagnosis markers" herein is a term of convenience to refer to the markers described herein and their use, and is not intended to indicate the markers are only used to diagnose AD. As this disclosure makes clear, these biomarkers are useful for, for example, assessing cognitive function, assessing MCI, assessing risk of developing AD, stratifying AD, etc. AD biomarkers include but are not limited to secreted proteins or metabolites present in a person's biological fluids (that is, a biological fluid sample), such as for example, blood, including whole blood, plasma or serum; urine; cerebrospinal fluid; tears; and saliva. Biological fluid samples encompass clinical samples, and also includes serum, plasma, and other biological fluids. As described herein, assessment of results can depend on whether the data were obtained by the qualitative or quantitative methods described herein and/or type of reference point used. For example, as described in Example 4, qualitative measurement of AD biomarker levels relative to another reference level, which may be relative to the level of another AD biomarker, may be obtained. In other methods described herein, such as in Example 7, quantitative or absolute values, that is protein concentration levels, in a biological fluid sample may be obtained. "Quantitative" result or data refers to an absolute value (see Example 7), which can include a concentration of a biomarker in pg/ml or ng/ml of molecule to sample. An example of a quantitative value is the measurement of concentration of protein levels directly for example by ELISA. "Qualitative" result or data provides a relative value which is as compared to a reference value. In some examples herein (Example 4), qualitative measurements are assessed by signal intensity on a filter. In some examples herein, multiple antibodies specific for AD biomarkers are attached to a suitable surface, e.g. as slide or filter.

[0076] In one aspect, the present invention provides methods of aiding diagnosis of Alzheimer's disease ("AD") and diagnosing AD, by obtaining measured levels of one or more AD diagnosis biomarkers in a biological fluid sample from an individual, such as for example, a peripheral biological fluid sample from an individual, and comparing those measured levels to reference levels. In some examples, the AD diagnosis biomarkers are selected from the group shown in Table 7. In other examples, the AD diagnosis biomarkers are selected from the group GCSF; IFN- γ ; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1 α ; MSP-a; SCF; TGF- β 3; TNF- β MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In yet other examples, the AD diagnosis biomarker are selected from the group shown in Table 3. In further examples, the AD diagnosis biomarkers are selected from the group consisting of BDNF, PDGF-BB, Leptin and RANTES. As shown herein in the examples, quantitative Leptin and BDNF levels have a statistically significant positive correlation with MMSE scores; quantitative PDGF-BB levels have a statistically significant negative correlation with MMSE scores in men; and quantitative RANTES levels

have a statistically significant positive correlation with PDGF-BB and BDNF. In some examples, the AD diagnosis biomarkers for use in methods of aiding diagnosis of Alzheimer's disease ("AD") and diagnosing AD include two or more of the following 4 biomarkers: BDNF, PDGF-BB, Leptin and RANTES. In further examples, the AD diagnosis biomarkers for use in methods of aiding diagnosis of Alzheimer's disease ("AD") and diagnosing AD comprise Leptin and RANTES; Leptin and BDNF; Leptin and PDGF-BB; Leptin, RANTES and BDNF; Leptin, RANTES and PDGF-BB; Leptin, BDNF and PDGF-BB; RANTES and BDNF; RANTES and PDGF-BB; BDNF and PDGF-BB; or Leptin, RANTES, BDNF and PDGF-BB. In some examples, the AD diagnosis markers for use in methods of aiding diagnosis of AD or diagnosing AD comprise Leptin, RANTES, BDNF and PDGF-BB. In other examples, the AD diagnosis markers for use in methods of aiding diagnosis of AD or diagnosing AD consist essentially of or consist of Leptin, RANTES, BDNF and PDGF-BB.

[0077] Methods of assessing cognitive function, aiding diagnosis of AD and diagnosing AD as described herein may comprise any of the following steps of obtaining a biological fluid sample from an individual, measuring the level of at least one AD diagnosis biomarker in the sample and comparing the measured level to an appropriate reference; obtaining measured levels of at least one AD diagnosis biomarker in a sample and comparing the measured level to an appropriate reference; comparing measured levels of at least one AD diagnosis biomarker obtained from a sample to an appropriate reference; measuring the level of at least one AD diagnosis biomarker in a sample; measuring the level of at least one AD diagnosis biomarker in a sample and comparing the measured level to an appropriate reference; diagnosing AD based on comparison of measured levels to an appropriate reference; or obtaining a measured value for at least one AD diagnosis biomarker in a sample. Comparing a measured level of an AD diagnosis biomarker to a reference level or obtaining a measured value for an AD diagnosis biomarker in a sample may be performed for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more AD diagnosis biomarker(s). The present invention also provides methods of evaluating results of the analytical methods described herein. Such evaluation generally entails reviewing such results and can assist, for example, in advising regarding clinical and/or diagnostic follow-up and/or treatment options. The present invention also provides methods for assessing a biological fluid sample for an indicator of any one or more of the following: cognitive function and/or impairment; MCI; AD; extent of AD, such as, for example, mild, moderate, severe; progression of AD; by measuring the level of or obtaining the measured level of or comparing a measured level of an AD diagnosis biomarker as described herein. Methods of assessing cognitive impairment includes the ADAS-COG, which is generally accepted to be equivalent to MMSE scoring.

[0078] For methods of diagnosing AD as described herein, the reference level is generally a predetermined level considered 'normal' for the particular AD diagnosis biomarker (e.g., an average level for age-matched individuals not diagnosed with AD), although reference levels which are determined contemporaneously (e.g., a reference value that is derived from a pool of samples including the sample being tested) are also contemplated. Also provided are methods of aiding in the diagnosis of Alzheimer's disease ("AD") by

comparing a measured level of at least one AD diagnosis biomarker in a biological fluid sample, such as, for example, a peripheral biological fluid sample from an individual with a reference level. Further provided are methods of aiding in the diagnosis of Alzheimer's disease ("AD") by measuring a level of at least one AD diagnosis biomarker in a biological fluid sample, such as, for example, a peripheral biological fluid sample from an individual. For the AD diagnosis biomarkers disclosed herein, a measurement for a marker which is below the reference level suggests (i.e., aids in the diagnosis of) or indicates a diagnosis of AD.

[0079] In another aspect, the invention provides methods of identifying individuals with mild cognitive impairment (MCI), by obtaining a quantitative measured level for RANTES in a biological fluid sample, such as, for example, a peripheral biological fluid sample from an individual, and comparing that level to a reference level. Generally, the reference level for RANTES is a predetermined level considered 'normal' for RANTES, and may be an age-matched normal level for RANTES, although reference levels which are determined contemporaneously (e.g., a reference value that is derived from a pool of samples including the sample being tested) are also contemplated. Also provided are methods of aiding in the diagnosis of MCI by comparing a quantitative measured level for RANTES in a biological fluid sample, such as, for example, a peripheral biological fluid sample from an individual with a reference level. Further provided are methods for aiding in the diagnosis of MCI by measuring a level for RANTES in a biological fluid sample, such as, for example, a peripheral biological fluid sample from an individual. A finding that the quantitative level of RANTES is low (below the reference level) in the biological fluid sample, such as, for example, the peripheral biological fluid sample from the individual suggests (i.e., aids in the diagnosis of) or indicates a diagnosis of MCI. In certain embodiments, such methods further include measuring, obtaining, and/or comparing the quantitative level of Leptin in the biological fluid sample, such as, for example, a peripheral biological sample. When both RANTES and Leptin levels are utilized, a finding that the quantitative RANTES level is low while the quantitative Leptin level is not (i.e., is substantially the same as or higher than the Leptin reference value) suggests (i.e., aids in the diagnosis of) or indicates a diagnosis of MCI. Accordingly the present invention provides methods for aiding in the diagnosis of mild cognitive impairment (MCI), comprising comparing a measured level for RANTES in a biological fluid sample obtained from an individual to a reference level. In some examples, the methods further comprise comparing a measured value for leptin in the biological fluid sample obtained from the individual to a reference level. In yet other examples, the methods further comprises measuring a level for leptin in said biological fluid sample, thereby producing said measured value for leptin. In yet other examples, the methods comprise measuring a level for RANTES in said biological fluid sample, thereby producing said measured value for RANTES. In yet other examples, the biological fluid sample is a peripheral fluid sample.

[0080] In a further aspect, the invention provides methods of monitoring progression of AD in an AD patient. As shown in Example 7, the inventors have found that quantitative levels of RANTES are decreased in AD patients with Questionable AD (MMSE=25-28); and that quantitative levels of RANTES are decreased in AD patients with mild

AD (MMSE=20-25), and RANTES levels decrease further as the severity of the AD intensifies. An individual with "Questionable AD" as used herein for quantitative data (also called absolute measurement) is an individual who (a) has been diagnosed with AD or has been given a diagnosis of probable AD, and (b) has either been assessed with the Mini-Mental State Examination (MMSE) (referenced in Folstein et al., *J. Psychiatr. Res* 1975; 12:1289-198) and scored 25-28 or would achieve a score of 25-28 upon MMSE testing. Accordingly, "Questionable AD" refers to AD in an individual having scored 25-28 on the MMSE and or would achieve a score of 25-28 upon MMSE testing. The reference level may be a predetermined level considered 'normal' for the particular RANTES (e.g., an average level for age-matched individuals not diagnosed with AD or MCI), or may be a historical reference level for the particular patient (e.g., a RANTES level that was obtained from a sample derived from the same individual, but at an earlier point in time). Reference levels which are determined contemporaneously (e.g., a reference value that is derived from a pool of samples including the sample being tested) are also contemplated. Accordingly, the invention provides methods for monitoring progression of AD in an AD patient by obtaining a quantitative value for RANTES from a biological fluid sample, such as for example, a peripheral biological fluid sample and comparing measured value to a reference value. Also provided are methods for monitoring progression of AD in an AD patient by comparing a measured value for leptin in a biological fluid sample, such as for example, a peripheral biological fluid sample with a reference value. Further provided are methods for monitoring progression of AD in an AD patient by measuring a level for leptin in a biological fluid sample, such as for example, a peripheral biological fluid sample. A decrease in the measured value indicates or suggests (diagnoses or suggests a diagnosis) progression (e.g., an increase in the severity) of AD in the AD patient.

[0081] In a further aspect, the inventors have found that quantitative Leptin levels are decreased in AD patients with Questionable AD; and that the quantitative levels of Leptin are decreased in AD patients with mild AD, and quantitative Leptin levels decrease further as the severity of the AD intensifies; and the quantitative levels of Leptin are positively correlated with MMSE scores (as described in Example 7). The reference level may be a predetermined level considered 'normal' for the particular Leptin (e.g., an average level for age-matched individuals not diagnosed with AD or MCI), or may be a historical reference level for the particular patient (e.g., a Leptin level that was obtained from a sample derived from the same individual, but at an earlier point in time). Quantitative reference levels which are determined contemporaneously (e.g., a reference value that is derived from a pool of samples including the sample being tested) are also contemplated. Accordingly, the invention provides methods for monitoring progression of AD in an AD patient by obtaining a quantitative measured value for Leptin from a biological fluid sample, such as for example, a peripheral biological fluid sample and comparing measured value to a reference value. Also provided are methods for monitoring progression of AD in an AD patient by comparing a measured value for Leptin in a biological fluid sample, such as for example, a peripheral biological fluid sample with a reference value. Further provided are methods for monitoring progression of AD in an AD patient by measuring a level for Leptin in a biological fluid sample,

such as for example, a peripheral biological fluid sample. A decrease in the quantitative measured value indicates or suggests (diagnoses or suggests a diagnosis) progression (e.g., an increase in the severity) of AD in the AD patient.

[0082] The inventors have found that quantitative BDNF levels are decreased in AD patients with mild AD, and that the quantitative BDNF levels in women are correlated with MMSE scores and BDNF levels decrease further as the severity of the AD intensifies (as described in Example 7). The reference level may be a predetermined level considered 'normal' for the particular BDNF (e.g., an average level for age-matched individuals not diagnosed with AD or MCI), or may be a historical reference level for the particular patient (e.g., a BDNF level that was obtained from a sample derived from the same individual, but at an earlier point in time). Reference levels which are determined contemporaneously (e.g., a reference value that is derived from a pool of samples including the sample being tested) are also contemplated. Accordingly, the invention provides methods for monitoring progression of AD in an AD patient by obtaining a quantitative measured value for BDNF from a biological fluid sample, such as for example, a peripheral biological fluid sample and comparing measured value to a reference value. Also provided are methods for monitoring progression of AD in an AD patient by comparing a quantitative measured value for BDNF in a biological fluid sample, such as for example, a peripheral biological fluid sample with a reference value. Further provided are methods for monitoring progression of AD in an AD patient by measuring a level for BDNF in a biological fluid sample, such as for example, a peripheral biological fluid sample. Generally speaking, a decrease in the measured value indicates or suggests (diagnoses or suggests a diagnosis) progression (e.g., an increase in the severity) of AD in the AD patient.

[0083] The inventors have found that quantitative PDGF-BB levels are decreased in AD patients with Questionable AD; that PDGF-BB levels are decreased in Questionable AD compared to Mild AD; and that the MMSE scores for male AD patients are negatively correlated with PDGF-BB levels (as described in Example 7). The reference level may be a predetermined level considered 'normal' for the PDGF-BB (e.g., an average level for age-matched male individuals not diagnosed with AD or MCI), or may be a historical reference level for the particular patient (e.g., a PDGF-BB level that was obtained from a sample derived from the same male individual, but at an earlier point in time). Reference levels which are determined contemporaneously (e.g., a reference value that is derived from a pool of samples including the sample being tested) are also contemplated. Accordingly, the invention provides methods for monitoring progression of AD in an AD patient by obtaining a measured value for PDGF-BB from a biological fluid sample from a male, such as for example, a peripheral biological fluid sample and comparing measured value to a reference value. Also provided are methods for monitoring progression of AD in an AD patient by comparing a measured value for PDGF-BB in a biological fluid sample, such as for example, a peripheral biological fluid sample with a reference value. Further provided are methods for monitoring progression of AD in an AD patient by measuring a level for PDGF-BB in a biological fluid sample such as for example, a peripheral biological fluid sample. A decrease in the measured value

indicates or suggests (diagnoses or suggests a diagnosis) progression (e.g., an increase in the severity) of AD in the AD patient.

[0084] Additionally, the invention provides methods of stratifying individuals diagnosed with (or having a probable diagnosis of) AD. The inventors have found that analysis of the levels of BDNF, or BDNF and PDGF-BB in biological fluid samples, such as, peripheral biological fluid samples provides information as to the severity of the AD in the AD patient from whom the peripheral biological fluid sample is derived. The reference values for BDNF and PDGF-BB used in these aspects of the invention are most commonly obtained from a population of AD patients other than the AD patient who is the source of the sample being tested (e.g., a mean or median value derived from a large number of AD patients), although reference levels for BDNF and PDGF-BB which are determined contemporaneously (e.g., a reference values that is derived from a pool of samples including the sample being tested) are also contemplated. Accordingly, the invention provides methods of stratifying AD patients into mild, and more advanced (e.g., moderate and severe) stages of AD ("staging") by obtaining a measured level for BDNF, and comparing the measured value with a reference value for BDNF. Accordingly, the invention provides methods of stratifying AD in an AD patient by obtaining a measured value for BDNF, and, optionally, PDGF-BB, in a biological fluid sample, such as a peripheral biological fluid sample, and comparing the measured level to a reference level. The invention also provides methods of stratifying AD in an AD patient by comparing a measured value for BDNF, and, optionally, PDGF-BB, in a biological fluid sample, such as a peripheral biological fluid sample with a reference value. The invention further provides methods of stratifying AD in an AD patient by measuring BDNF and, optionally, PDGF-BB, in a biological fluid sample, such as a peripheral biological fluid sample. As described in Example 4, and under the experimental conditions disclosed in Example 4 which provide qualitative results, samples which have BDNF levels lower than the reference level suggest or indicate mild AD, while samples with BDNF levels higher than the reference level suggest more advanced AD (i.e., moderate or severe AD). Amongst those samples with BDNF levels higher than the reference level, those also having PDGF-BB levels below the reference level suggest or indicate moderate AD, while those samples also having PDGF-BB levels above the reference level suggest or indicate severe AD. It has been found that for Questionable AD (MMSE score in the range of 25-28) the levels of Leptin and PDGF-BB increase significantly whereas BDNF and RANTES do not change significantly. It has been found that from Mild AD (MMSE score in the range of 20-25) to Moderate AD (MMSE score in the range of 10-20) the level of LEPTIN does not decline whereas the levels for RANTES, BDNF and PDGF-BB declines. Accordingly, in some embodiments (as defined by the above MMSE scores from Example 7), Mild AD is indicated in quantitative assays when the levels of Leptin and/or PDGF-BB increase significantly whereas BDNF and RANTES do not change significantly as compared to Questionable AD as a reference. Accordingly, in some embodiments, (as defined by the above MMSE scores from Example 7), Moderate AD is indicated when Leptin does not decline whereas the levels for RANTES, BDNF and PDGF declines as compared to Mild AD as a reference. Accordingly, provided herein are

methods comprising comparing measured values for RANTES and Leptin levels in a biological fluid sample from said patient with reference values for RANTES and Leptin; comparing measured values for brain derived neurotrophic factor (BDNF), Leptin, and RANTES, levels in a biological fluid sample from said patient with reference values for BDNF, Leptin, and RANTES; comparing measured values for Leptin and BB homodimeric platelet derived growth factor (PDGF-BB) levels in a biological fluid sample from said patient with reference values for Leptin and PDGF-BB. Accordingly, the present invention provides methods for stratifying Alzheimer's disease (AD) in an individual, comprising comparing measured values for brain derived neurotrophic factor (BDNF) and BB homodimeric platelet derived growth factor (PDGF-BB) levels in a biological fluid sample from said patient with reference values for BDNF and PDGF-BB. In some examples, the methods further comprise comparing measured values for leptin and Rantes levels with reference values for leptin and Rantes, wherein reference values for BDNF, PDGF-BB, leptin and Rantes are for samples from individuals with MMSE scores from 25 to 28, wherein an increase in leptin and PDGF-BB levels and wherein levels of BDNF and RANTES stay substantially the same indicate mild AD as indicated by an MMSE score of 20-25. The present invention also provides methods of further comprising comparing measured values for leptin and Rantes levels with reference values for leptin and Rantes, wherein reference values for BDNF, PDGF-BB, leptin and Rantes are for samples from individuals with MMSE scores from 20-25, wherein a decrease in Rantes, BDNF, and PDGF levels and wherein levels of Leptin stays substantially the same indicate moderate AD as indicated by an MMSE score of 10-20. An AD biomarker that stays "substantially the same" means that there is not a significant change, and that the values stay about the same. In some embodiments, substantially the same is a change less than any of about 12%, 10%, 5%, 2%, 1%. In some embodiments, a significant change means not statistically significant using standard methods in the art. The methods described above are also applicable to methods for assessing progression of AD. It is understood that the cognitive function indicated by the markers herein can be by other measurements with results or indicia that corresponds to approximately the same level of cognitive function as the MMSE scores provided herein.

[0085] The present invention also provides methods of aiding diagnosis of Alzheimer's disease ("AD"), comprising comparing a measured level of at least one AD diagnosis biomarker in a biological fluid sample from an individual to a reference level for the biomarker for each biomarker measured, wherein the at least one AD diagnosis biomarker is selected from Table 7 and has a statistically significant positive correlation with MMSE scores that is comparable to BDNF and/or Leptin correlation with MMSE scores, and wherein the at least one AD diagnosis biomarker is not statistically correlated with age. An AD diagnosis biomarker that has a statistically significant positive correlation with MMSE scores that is comparable to BDNF and/or leptin correlation with MMSE scores means that the biomarker is an AD diagnosis marker. In some examples, the AD diagnosis biomarker is selected from the group of biomarkers consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL

R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R and in other examples is selected from the group of biomarkers consisting of basic fibroblast growth factor (bFGF); BB homodimeric platelet derived growth factor (PDGF-BB); brain derived neurotrophic factor (BDNF); epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), and tumor necrosis factor beta (TNF-β).

[0086] The results of the comparison between the measured value(s) and the reference value(s) are used to diagnose or aid in the diagnosis of AD or MCI, to stratify AD patients according to the severity of their disease, or to monitor progression of AD in an AD patient. Accordingly, if the comparison indicates a difference between the measured value(s) and the reference value(s) that is suggestive/indicative of AD or MCI, then the appropriate diagnosis is aided in or made. Conversely, if the comparison of the measured level(s) to the reference level(s) does not indicate differences that suggest or indicate a diagnosis of AD or MCI, then the appropriate diagnosis is not aided in or made. Likewise, when comparison of a measured level for Leptin in a sample derived from an AD patient is decreased in comparison to the reference value, diagnosis of progression of the patient's AD is made or aided in. Similarly, when the comparison of levels of BDNF and PDGF-BB levels in a sample obtained from an AD patient indicates or suggests a particular stage of AD, the diagnosis of the particular stage of AD (mild, moderate or severe) is aided in or made.

[0087] As will be understood by those of skill in the art, when, in the practice of the AD diagnosis methods of the invention (i.e., methods of diagnosing or aiding in the diagnosis of AD), more than one AD diagnosis biomarker is used but the markers do not unanimously suggest or indicate a diagnosis of AD, the 'majority' suggestion or indication (e.g., when the method utilizes five AD diagnosis biomarkers, 3 of which suggest/indicate AD, the result would be considered as suggesting or indicating a diagnosis of AD for the individual) is considered the result of the assay. However, in some embodiments in which measured values for at least two AD diagnosis biomarkers are obtained and one of the measured values is for Leptin, the measured value for Leptin must be less than the reference value to indicate or suggest a diagnosis of AD. As will be appreciated by one of skill in the art, methods disclosed herein may include the use of any of a variety of biological markers (which may or may not be AD markers) to determine the integrity and/or characteristics of the biological sample(s). For example, Leptin levels, which are generally higher in females, may be measured as a marker of gender.

[0088] In certain embodiments of the invention, levels for AD biomarkers are obtained from an individual at more than one time point. Such "serial" sampling is well suited for the

aspects of the invention related to monitoring progression of AD in an AD patient. Serial sampling can be performed on any desired timeline, such as monthly, quarterly (i.e., every three months), semi-annually, annually, biennially, or less frequently. The comparison between the measured levels and the reference level may be carried out each time a new sample is measured, or the data relating to levels may be held for less frequent analysis.

[0089] As will be understood by those of skill in the art, biological fluid samples including peripheral biological fluid samples are usually collected from individuals who are suspected of having AD, or developing AD or MCI. The invention also contemplates samples from individuals for whom cognitive assessment is desired. Alternatively, individuals (or others involved in for example research and/or clinicians may desire such assessments without any indication of AD, suspected AD, at risk for AD. For example, a normal individual may desire such information. Such individuals are most commonly 65 years or older, although individuals from whom biological fluid samples, such as peripheral biological fluid samples are taken for use in the methods of the invention may be as young as 35 to 40 years old, when early onset AD or familial AD is suspected.

[0090] The invention also provides methods of screening for candidate agents for the treatment of AD and/or MCI by assaying prospective candidate agents for activity in modulating AD biomarkers. The screening assay may be performed either in vitro and/or in vivo. Candidate agents identified in the screening methods described herein may be useful as therapeutic agents for the treatment of AD and/or MCI.

[0091] The probability P that the composite is more predictive than any subset of markers present in the composite can be expressed mathematically as:

$$P=1-(1-P_1)(1-P_2)(1-P_3) \dots (1-P_n)$$

[0092] Where the probability P_1 , P_2 , P_n represent the probability of individual marker being able to predict clinical phenotypes, and where $1-P_n$ represents the complement of that probability. Any subset of the composite, will always therefore have a smaller value for P.

[0093] In accordance with a further embodiment of the present invention, the relative concentrations in serum, CSF, or other fluids of the biomarkers cited in Table 7 as a composite, or collective, or any subset of such a composite, composed of 5 (five) or more elements is more predictive than the absolute concentration of any individual marker in predicting clinical phenotypes, disease detection, stratification, monitoring, and treatment of AD, PD, frontotemporal dementia, cerebrovascular disease, multiple sclerosis, and neuropathies.

[0094] AD Diagnosis Biomarkers

[0095] Immune mechanisms are an essential part of the host defense system and typically feature prominently in the inflammatory response. A growing number of studies are discovering intriguing links between the immune system and the CNS. For example, it has become clear that the CNS is not entirely sheltered from immune surveillance and that various immune cells can traverse the blood-brain barrier. Invading leukocytes can attack target antigens in the CNS or produce growth factors that might protect neurons against

degeneration (Hohlfeld et al., 2000, *J. Neuroimmunol.* 107, 161-166). These responses are elicited through a variety of protein mediators, including but not limited to cytokines, chemokines, neurotrophic factors, collectins, kinins, and acute phase proteins in the immune and inflammatory systems, in intercellular communication across neurons, glial cells, endothelial cells and leukocytes. Without being bound by theory, it is hypothesized that the cytokines, chemokines, neurotrophic factors, collectins, kinins, and acute phase proteins listed in Table 7 are differentially expressed in serum associated with neurodegenerative and inflammatory diseases such as Alzheimer's, Parkinson's disease, Multiple Sclerosis, and neuropathies. Cytokines are a heterogeneous group of polypeptide mediators that have been associated with activation of numerous functions, including the immune system and inflammatory responses. Peripheral cytokines also penetrate the blood-brain barrier directly via active transport mechanisms or indirectly via vagal nerve stimulation. Cytokines can act in an autocrine manner, affecting the behavior of the cell that releases the cytokine, or in a paracrine manner, affecting the behavior of adjacent cells. Some cytokines can act in an endocrine manner, affecting the behavior of distant cells, although this depends on their ability to enter the circulation and on their half-life. The cytokine families include, but are not limited to, interleukins (IL-1 alpha, IL-1 beta, IL1ra and IL-2 to IL-18), tumor necrosis factors (TNF-alpha and TNF-beta), interferons (INF-alpha, beta and gamma), colony stimulating factors (G-CSF, M-CSF, GM-CSF, IL-3 and some of the other ILs), and growth factors (EGF, FGF, PDGF, TGF alpha, TGF betas, BMPs, GDFs, CTGF, and ECGF).

[0096] The inventors have discovered a collection of biochemical markers present in peripheral bodily fluids that may be used to assess cognitive function, including diagnose or aid in the diagnosis of AD. These "AD diagnosis markers" include, but are not limited to GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In other examples, these "AD diagnosis biomarkers" are: basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), Leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neurophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), tumor necrosis factor beta (TNF-β). In other examples, the AD diagnosis markers include one or more of Leptin, RANTES, PDFG-BB and BDNF.

[0097] The AD diagnosis biomarkers discovered by the inventors are all known molecules. Brain derived neurotrophic factor (BDNF) is described in, for example Rosenthal et al., 1991, *Endocrinology* 129(3):1289-94. Basic fibroblast growth factor (bFGF) is described in, for

example Abraham et al., 1986, *EMBO J.* 5(10):2523-28. Epidermal growth factor (EGF) is described in, for example Gray et al., 1983, *Nature* 303(5919):722-25. Fibroblast growth factor 6 (FGF-6) is described in, for example Marics et al., 1989, *Oncogene* 4(3):335-40. Interleukin-3 (IL-3) is described in, for example Yang et al., 1986, *Cell* 47(1):3-10. Soluble interleukin-6 receptor (sIL-6R) is described in, for example, Taga et al., 1989, *Cell* 58(3):573-81. Leptin (also known as "ob") is described in, for example Masuzaki et al. 1995, *Diabetes* 44(7):855-58. Macrophage inflammatory protein-1 delta (MIP-1δ) is described in, for example Wang et al., 1998, *J. Clin. Immunol.* 18(3):214-22. Macrophage stimulating protein alpha chain (MSP-α) is described in, for example, Yoshimura et al., 1993, *J. Biol. Chem.* 268 (21), 15461-68, and Yoshikawa et al., 1999, *Arch. Biochem. Biophys.* 363(2):356-60. Neutrophil activating peptide-2 (NAP-2) is described in, for example Walz et al., 1991, *Adv. Exp. Med. Biol.* 305:39-46. Neurotrophin-3 (NT-3) is described in, for example Hohn et al., 1990, *Nature* 344(6264):339-41. BB homodimeric platelet derived growth factor (PDGF-BB) is described in, for example Collins et al., 1985, *Nature* 316(6030):748-50. RANTES is described in, for example Schall et al., 1988, *J. Immunol.* 141(3):1018-25. Stem cell factor (SCF) is described in, for example Zsebo et al., 1990, *Cell* 63(1):213-24. Soluble tumor necrosis factor receptor-2 (sTNF RII) is described in, for example Schall et al., 1990, *Cell* 61(2):361-70. Transforming growth factor-beta 3 (TGF-β3) is described in, for example ten Dijke et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85 (13):4715-19. Tissue inhibitor of metalloproteases-1 (TIMP-1) is described in, for example, Docherty et al., 1985, *Nature* 318(6041):66-69 and Gasson et al., 1985, *Nature* 315(6022):768-71. Tissue inhibitor of metalloproteases-2 (TIMP-2) is described in, for example, Stetler-Stevenson et al., 1990, *J. Biol. Chem.* 265(23):13933-38. Tumor necrosis factor beta (TNF-β) is described in, for example Gray et al., 1984, *Nature* 312(5996):721-24. Thrombopoietin (TPO) is described in, for example, Foster et al., 1994, *Proc. Natl. Acad. Sci. U.S.A.* 91(26):13023-27.

[0098] Although the inventors have found acceptable levels of sensitivity and specificity with single AD diagnosis biomarkers for practice of the AD diagnosis methods, the effectiveness (e.g., sensitivity and/or specificity) of the methods of the AD diagnosis methods of the instant invention are generally enhanced when at least two AD diagnosis biomarkers are utilized. In some examples, the methods of the AD diagnosis methods of the instant invention are generally enhanced when at least four AD diagnosis biomarkers are utilized. Multiple AD diagnosis biomarkers may be selected from the AD diagnosis biomarkers disclosed herein by a variety of methods, including "q value" and/or by selecting for cluster diversity. AD diagnosis biomarkers may be selected on the basis of "q value", a statistical value that the inventors derived when identifying the AD diagnosis biomarkers (see Table 3 in Example 1). "q values" for selection of AD diagnosis biomarkers range from less than about 0.0001 to about 0.05 and in some examples, range from about 0.01 to about 0.05. Alternately (or additionally), AD diagnosis biomarkers may be selected to preserve cluster diversity. The inventors have separated the AD diagnosis biomarkers into a number of clusters (see Table 1). Here the clusters are formed by qualitative measurements for each biomarker which are most closely correlated. As used herein, "correlate" or "correlation" is a simultaneous change

in value of two numerically valued random variables such as MMSE scores and quantitative protein concentrations or qualitative protein concentrations. As used herein "discriminate" or "discriminatory" is refers to the quantitative or qualitative difference between two or more samples for a given variable. The cluster next to such a cluster is a cluster that is most closely correlated with the cluster. The correlations between biomarkers and between clusters can be represented by a hierarchical tree generated by unsupervised clustering using a public web based software called wCLUTO available at: cluto.cccb.umn.edu/cgi-bin/wCluto/wCluto.cgi. If more than one AD diagnosis biomarker is selected for testing, in some examples, the AD diagnosis biomarkers selected are at least partially diverse (i.e., the AD diagnosis biomarkers represent at least two different clusters, for example, a set of AD diagnosis biomarkers comprising Leptin, BDNF and/or PDGF-BB from cluster 4 in Table 1 and RANTES from cluster 3 of Table 1), and in some instances the AD diagnosis biomarkers are completely diverse (i.e. no two of the selected AD diagnosis biomarkers are from the same cluster). Accordingly, the invention provides a number of different embodiments for diagnosing or aiding in the diagnosis of AD.

TABLE 1

Cluster	Biomarker
0	bFGF
1	TPO
2	FGF-6
	IL-3
	sIL-6 R
	MIP-1δ
	sTNF RII
	TNF-β
3	RANTES
	TIMP-1
	TIMP-2
4	BDNF
	EGF
	LEPTIN(OB)
	MSP-α
	NAP-2
	NT-3
	PDGF-BB
	SCF
	TGF-β3

[0099] In some embodiments, the level of a single AD diagnosis biomarker in a peripheral biological fluid sample is obtained and the measured level is compared to a reference level to diagnose or aid in diagnosing AD. In certain embodiments where measured level for a single AD diagnosis biomarker is obtained for the practice of the invention, the measured level is for RANTES in the peripheral biological fluid sample.

[0100] In other embodiments, the levels of at least two AD diagnosis biomarkers in a peripheral biological fluid sample are obtained and compared to reference levels for each of the markers. Accordingly, the invention provides methods for diagnosing and/or aiding in the diagnosis of AD by measuring the levels of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15, or 20 AD diagnosis biomarkers and comparing the measured levels with reference levels. Exemplary embodiments utilize 2, 3, 4, or 5 AD diagnosis biomarkers. In some embodiments, provided herein are methods for diagnosing

and/or aiding in the diagnosis of AD by measuring the levels of at least Leptin, RANTES, BDGF, and PDGF-BB.

[0101] For those embodiments which utilize more than one AD diagnosis biomarker (i.e., those embodiments in which measured values are obtained for more than one AD diagnosis biomarker), exemplary combinations of AD diagnosis biomarkers shown in Table 3 include (1) Leptin in combination with any of the other AD diagnosis biomarkers (i.e., Leptin and BDNF, Leptin and bFGF, Leptin and EGF, Leptin and FGF-6, Leptin and IL-3, Leptin and sIL-6R, Leptin and MIP-1 δ , Leptin and MSP- α , Leptin and NAP-2, Leptin and NT-3, Leptin and PDGF-BB, Leptin and RANTES, Leptin and SCF, Leptin and sTNF RII, Leptin and TGF- β 3, Leptin and TIMP-1, Leptin and TIMP-2, Leptin and TNF- β , and Leptin and TPO); (2) RANTES in combination with any of the other AD diagnosis biomarkers (i.e., RANTES and BDNF, RANTES and bFGF, RANTES and EGF, RANTES and FGF-6, RANTES and IL-3, RANTES and sIL-6R, RANTES and Leptin, RANTES and MIP-1 δ , RANTES and MSP- α , RANTES and NAP-2, RANTES and NT-3, RANTES and PDGF-BB, RANTES and SCF, RANTES and sTNF RII, RANTES and TGF- β 3, RANTES and TIMP-1, RANTES and TIMP-2, RANTES and TNF- β , and RANTES and TPO); (3) PDGF-BB and any of the other AD diagnosis biomarkers (i.e., PDGF-BB and BDNF, PDGF-BB and bFGF, PDGF-BB and EGF, PDGF-BB and FGF-6, PDGF-BB and IL-3, PDGF-BB and sIL-6R, PDGF-BB and Leptin, PDGF-BB and MIP-1 δ , PDGF-BB and MSP- α , PDGF-BB and NAP-2, PDGF-BB and NT-3, PDGF-BB and RANTES, PDGF-BB and SCF, PDGF-BB and sTNF RII, PDGF-BB and TGF- β 3, PDGF-BB and TIMP-1, PDGF-BB and TIMP-2, PDGF-BB and TNF- β , and PDGF-BB and TPO); (4) BDNF in combination with any of the other AD diagnosis biomarkers (i.e., BDNF and bFGF, BDNF and EGF, BDNF and FGF-6, BDNF and IL-3, BDNF and sIL-6R, BDNF and Leptin, BDNF and MIP-1 δ , BDNF and MSP- α , BDNF and NAP-2, BDNF and NT-3, BDNF and PDGF-BB, BDNF and RANTES, BDNF and SCF, BDNF and sTNF RII, BDNF and TGF- β 3, BDNF and TIMP-1, BDNF and TIMP-2, BDNF and TNF- β , and BDNF and TPO); (5) RANTES, PDGF-BB, and NT-3; (6) Leptin, PDGF-BB, and RANTES; (7) BDNF, PDGF-BB, and RANTES; (8) BDNF, Leptin, and RANTES; (9) BDNF, Leptin, and PDGF-BB; (10) PDGF-BB, EGF, and NT-3; (11) PDGF-BB, NT 3, and Leptin; (12) BDNF, Leptin, PDGF-BB, RANTES; and (13) RANTES, PDGF-BB, NT-3, EGF, NAP-2, and Leptin. Additional exemplary combinations of AD diagnosis biomarkers include (14) Leptin in combination with any of the other AD diagnosis biomarkers disclosed herein (i.e., Leptin and GCSF, Leptin and IFN- γ , Leptin and IGFBP-1, Leptin and BMP-6, Leptin and BMP-4, Leptin and Eotaxin-2, Leptin and IGFBP-2, Leptin and TARC, Leptin and ANG, Leptin and PARC, Leptin and Acrp30, Leptin and AgRP(ART), Leptin and ICAM-1, Leptin and TRAIL R3, Leptin and uPAR, Leptin and IGFBP-4, Leptin and IL-1Ra, Leptin and AXL, Leptin and FGF-4, Leptin and CNTF, Leptin and MCP-1, Leptin and MIP 1b, Leptin and VEGF-B, Leptin and IL-8, Leptin and FAS and Leptin and EGF-R), (15) RANTES in combination with any of the other AD diagnosis biomarkers disclosed herein (i.e., RANTES and GCSF, RANTES and IFN- γ , RANTES and IGFBP-1, RANTES and BMP-6, RANTES and BMP-4, RANTES and Eotaxin-2, RANTES and IGFBP-2, RANTES and TARC, RANTES and ANG, RANTES and PARC,

RANTES and Acrp30, RANTES and AgRP(ART), RANTES and ICAM-1, RANTES and TRAIL R3, RANTES and uPAR, RANTES and IGFBP-4, RANTES and IL-1Ra, RANTES and AXL, RANTES and FGF-4, RANTES and CNTF, RANTES and MCP-1, RANTES and MIP1b, RANTES and VEGF-B, RANTES and IL-8, RANTES and FAS and RANTES and EGF-R), (16) PDGF-BB in combination with any of the other AD diagnosis biomarkers disclosed herein (i.e., PDGF-BB and GCSF, PDGF-BB and IFN- γ , PDGF-BB and IGFBP-1, PDGF-BB and BMP-6, PDGF-BB and BMP-4, PDGF-BB and Eotaxin-2, PDGF-BB and IGFBP-2, PDGF-BB and TARC, PDGF-BB and ANG, PDGF-BB and PARC, PDGF-BB and Acrp30, PDGF-BB and AgRP(ART), PDGF-BB and ICAM-1, PDGF-BB and TRAIL R3, PDGF-BB and uPAR, PDGF-BB and IGFBP-4, PDGF-BB and IL-1Ra, PDGF-BB and AXL, PDGF-BB and FGF-4, PDGF-BB and CNTF, PDGF-BB and MCP-1, PDGF-BB and MIP1b, PDGF-BB and VEGF-B, PDGF-BB and IL-8, PDGF-BB and FAS and PDGF-BB and EGF-R), (17) BDNF in combination with any of the other AD diagnosis biomarkers disclosed herein (i.e., BDNF and GCSF, BDNF and IFN- γ , BDNF and IGFBP-1, BDNF and BMP-6, BDNF and BMP-4, BDNF and Eotaxin-2, BDNF and IGFBP-2, BDNF and TARC, BDNF and ANG, BDNF and PARC, BDNF and Acrp30, BDNF and AgRP(ART), BDNF and ICAM-1, BDNF and TRAIL R3, BDNF and uPAR, BDNF and IGFBP-4, BDNF and IL-1Ra, BDNF and AXL, BDNF and FGF-4, BDNF and CNTF, BDNF and MCP-1, BDNF and MIP1b, BDNF and VEGF-B, BDNF and IL-8, BDNF and FAS and BDNF and EGF-R).

[0102] Measuring Levels of AD Biomarkers

[0103] There are a number of statistical tests for identifying biomarkers which vary significantly between the subsets, including the conventional t test. However, as the number of biomarkers measured increases, it is generally advantageous to use a more sophisticated technique, such as SAM (see Tusher et al., 2001, *Proc. Natl. Acad. Sci. U.S.A.* 98(9):5116-21). Other useful techniques include Tree Harvesting (Hastie et al., *Genome Biology* 2001, 2:research0003.1-0003.12), Self Organizing Maps (Kohonen, 1982b, *Biological Cybernetics* 43(1):59-69), Frequent Item Set (Agrawal et al., 1993 "Mining association rules between sets of items in large databases." In Proc. of the ACM SIGMOD Conference on Management of Data, pages 207-216, Washington, D.C., May 1993), Bayesian networks (Gottardo, Statistical analysis of microarray data, A Bayesian approach. Biostatistics (2001), 1,1, pp 1-37), and the commercially available software packages CART and MARS.

[0104] The SAM technique assigns a score to each biomarker on the basis of change in expression relative to the standard deviation of repeated measurements. For biomarkers with scores greater than an adjustable threshold, the algorithm uses permutations of the repeated measurements to estimate the probability that a particular biomarker has been identified by chance (calculated as a "q-value"), or a false positive rate which is used to measure accuracy. The SAM technique can be carried out using publicly available software called Significance Analysis of Microarrays (see www-stat.class.stanford.edu/~tibs/clickwrap/sam.html).

[0105] A biomarkers is considered "identified" as being useful for aiding in the diagnosis, diagnosis, stratification,

monitoring, and/or prediction of neurological disease when it is significantly different between the subsets of peripheral biological samples tested. Levels of a biomarker are “significantly different” when the probability that the particular biomarker has been identified by chance is less than a predetermined value. The method of calculating such probability will depend on the exact method utilized to compare the levels between the subsets (e.g., if SAM is used, the q-value will give the probability of misidentification, and the p value will give the probability if the t test (or similar statistical analysis) is used). As will be understood by those in the art, the predetermined value will vary depending on the number of biomarkers measured per sample and the number of samples utilized. Accordingly, predetermined value may range from as high as 50% to as low as 20, 10, 5, 3, 2, or 1%.

[0106] As described herein, the level of at least one AD diagnosis biomarker is measured in a biological sample from an individual. The AD biomarker level(s) may be measured using any available measurement technology that is capable of specifically determining the level of the AD biomarker in a biological sample. The measurement may be either quantitative or qualitative, so long as the measurement is capable of indicating whether the level of the AD biomarker in the peripheral biological fluid sample is above or below the reference value.

[0107] The measured level may be a primary measurement of the level of a particular biomarker or a measurement of the quantity of biomarker itself (quantitative data, such as in Example 7), such as by detecting the number of biomarker molecules in the sample) or it may be a secondary measurement of the biomarker (a measurement from which the quantity of the biomarker can be but not necessarily deduced (qualitative data, such as Example 4), such as a measure of enzymatic activity (when the biomarker is an enzyme) or a measure of mRNA coding for the biomarker). Qualitative data may also be derived or obtained from primary measurements.

[0108] Although some assay formats will allow testing of peripheral biological fluid samples without prior processing of the sample, it is expected that most peripheral biological fluid samples will be processed prior to testing. Processing generally takes the form of elimination of cells (nucleated and non-nucleated), such as erythrocytes, leukocytes, and platelets in blood samples, and may also include the elimination of certain proteins, such as certain clotting cascade proteins from blood. In some examples, the peripheral biological fluid sample is collected in a container comprising EDTA.

[0109] Commonly, AD biomarker levels will be measured using an affinity-based measurement technology. “Affinity” as relates to an antibody is a term well understood in the art and means the extent, or strength, of binding of antibody to the binding partner, such as an AD diagnosis biomarker as described herein (or epitope thereof). Affinity may be measured and/or expressed in a number of ways known in the art, including, but not limited to, equilibrium dissociation constant (K_D or K_d), apparent equilibrium dissociation constant (K_D' or K_d'), and IC_{50} (amount needed to effect 50% inhibition in a competition assay; used interchangeably herein with “150”). It is understood that, for purposes of this invention, an affinity is an average affinity for a given

population of antibodies which bind to an epitope. Values of K_D reported herein in terms of mg IgG per ml or mg/ml indicate mg Ig per ml of serum, although plasma can be used.

[0110] Affinity-based measurement technology utilizes a molecule that specifically binds to the AD biomarker being measured (an “affinity reagent,” such as an antibody or aptamer), although other technologies, such as spectroscopy-based technologies (e.g., matrix-assisted laser desorption ionization-time of flight, or MALDI-TOF, spectroscopy) or assays measuring bioactivity (e.g., assays measuring mitogenicity of growth factors) may be used.

[0111] Affinity-based technologies include antibody-based assays (immunoassays) and assays utilizing aptamers (nucleic acid molecules which specifically bind to other molecules), such as ELONA. Additionally, assays utilizing both antibodies and aptamers are also contemplated (e.g., a sandwich format assay utilizing an antibody for capture and an aptamer for detection).

[0112] If immunoassay technology is employed, any immunoassay technology which can quantitatively or qualitatively measure the level of a AD biomarker in a biological sample may be used. Suitable immunoassay technology includes radioimmunoassay, immunofluorescent assay, enzyme immunoassay, chemiluminescent assay, ELISA, immuno-PCR, and western blot assay.

[0113] Likewise, aptamer-based assays which can quantitatively or qualitatively measure the level of a AD biomarker in a biological sample may be used in the methods of the invention. Generally, aptamers may be substituted for antibodies in nearly all formats of immunoassay, although aptamers allow additional assay formats (such as amplification of bound aptamers using nucleic acid amplification technology such as PCR (U.S. Pat. No. 4,683,202) or isothermal amplification with composite primers (U.S. Pat. Nos. 6,251,639 and 6,692,918).

[0114] A wide variety of affinity-based assays are known in the art. Affinity-based assays will utilize at least one epitope derived from the AD biomarker of interest, and many affinity-based assay formats utilize more than one epitope (e.g., two or more epitopes are involved in “sandwich” format assays; at least one epitope is used to capture the marker, and at least one different epitope is used to detect the marker).

[0115] Affinity-based assays may be in competition or direct reaction formats, utilize sandwich-type formats, and may further be heterogeneous (e.g., utilize solid supports) or homogenous (e.g., take place in a single phase) and/or utilize or immunoprecipitation. Most assays involve the use of labeled affinity reagent (e.g., antibody, polypeptide, or aptamer); the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the probe are also known; examples of which are assays which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA and ELONA assays. Herein, the examples referred to as “quantitative data” the biomarker concentrations were obtained using ELISA. Either of the biomarker or reagent specific for the biomarker can be attached to a surface and levels can be measured directly or indirectly.

[0116] In a heterogeneous format, the assay utilizes two phases (typically aqueous liquid and solid). Typically an AD biomarker-specific affinity reagent is bound to a solid support to facilitate separation of the AD biomarker from the bulk of the biological sample. After reaction for a time sufficient to allow for formation of affinity reagent/AD biomarker complexes, the solid support or surface containing the antibody is typically washed prior to detection of bound polypeptides. The affinity reagent in the assay for measurement of AD biomarkers may be provided on a support (e.g., solid or semi-solid); alternatively, the polypeptides in the sample can be immobilized on a support or surface. Examples of supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates), polyvinylidene fluoride, diazotized paper, nylon membranes, activated beads, glass and Protein A beads. Both standard and competitive formats for these assays are known in the art. Accordingly, the provided herein are complexes comprising at least one AD diagnosis biomarker bound to a reagent specific for the biomarker, wherein said reagent is attached to a surface. Also provided herein are complexes comprising at least one AD diagnosis biomarker bound to a reagent specific for the biomarker, wherein said biomarker is attached to a surface.

[0117] Array-type heterogeneous assays are suitable for measuring levels of AD biomarkers when the methods of the invention are practiced utilizing multiple AD biomarkers. Array-type assays used in the practice of the methods of the invention will commonly utilize a solid substrate with two or more capture reagents specific for different AD biomarkers bound to the substrate a predetermined pattern (e.g., a grid). The peripheral biological fluid sample is applied to the substrate and AD biomarkers in the sample are bound by the capture reagents. After removal of the sample (and appropriate washing), the bound AD biomarkers are detected using a mixture of appropriate detection reagents that specifically bind the various AD biomarkers. Binding of the detection reagent is commonly accomplished using a visual system, such as a fluorescent dye-based system. Because the capture reagents are arranged on the substrate in a predetermined pattern, array-type assays provide the advantage of detection of multiple AD biomarkers without the need for a multiplexed detection system.

[0118] In a homogeneous format the assay takes place in single phase (e.g., aqueous liquid phase). Typically, the biological sample is incubated with an affinity reagent specific for the AD biomarker in solution. For example, it may be under conditions that will precipitate any affinity reagent/antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

[0119] In a standard (direct reaction) format, the level of AD biomarker/affinity reagent complex is directly monitored. This may be accomplished by, for example, determining the amount of a labeled detection reagent that forms is bound to AD biomarker/affinity reagent complexes. In a competitive format, the amount of AD biomarker in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled AD biomarker (or

other competing ligand) in the complex. Amounts of binding or complex formation can be determined either qualitatively or quantitatively.

[0120] The methods described in this patent may be implemented using any device capable of implementing the methods. Examples of devices that may be used include but are not limited to electronic computational devices, including computers of all types. When the methods described in this patent are implemented in a computer, the computer program that may be used to configure the computer to carry out the steps of the methods may be contained in any computer readable medium capable of containing the computer program. Examples of computer readable medium that may be used include but are not limited to diskettes, CD-ROMs, DVDs, ROM, RAM, and other memory and computer storage devices. The computer program that may be used to configure the computer to carry out the steps of the methods may also be provided over an electronic network, for example, over the internet, world wide web, an intranet, or other network.

[0121] In one example, the methods described in this patent may be implemented in a system comprising a processor and a computer readable medium that includes program code means for causing the system to carry out the steps of the methods described in this patent. The processor may be any processor capable of carrying out the operations needed for implementation of the methods. The program code means may be any code that when implemented in the system can cause the system to carry out the steps of the methods described in this patent. Examples of program code means include but are not limited to instructions to carry out the methods described in this patent written in a high level computer language such as C++, Java, or Fortran; instructions to carry out the methods described in this patent written in a low level computer language such as assembly language; or instructions to carry out the methods described in this patent in a computer executable form such as compiled and linked machine language.

[0122] Complexes formed comprising AD biomarker and an affinity reagent are detected by any of a number of known techniques known in the art, depending on the format of the assay and the preference of the user. For example, unlabelled affinity reagents may be detected with DNA amplification technology (e.g., for aptamers and DNA-labeled antibodies) or labeled "secondary" antibodies which bind the affinity reagent. Alternately, the affinity reagent may be labeled, and the amount of complex may be determined directly (as for dye-(fluorescent or visible), bead-, or enzyme-labeled affinity reagent) or indirectly (as for affinity reagents "tagged" with biotin, expression tags, and the like). Herein the examples provided referred to as "qualitative data" filter based antibody arrays using chemiluminescence were used to obtain measurements for biomarkers.

[0123] As will be understood by those of skill in the art, the mode of detection of the signal will depend on the exact detection system utilized in the assay. For example, if a radiolabeled detection reagent is utilized, the signal will be measured using a technology capable of quantitating the signal from the biological sample or of comparing the signal from the biological sample with the signal from a reference sample, such as scintillation counting, autoradiography (typically combined with scanning densitometry), and the

like. If a chemiluminescent detection system is used, then the signal will typically be detected using a luminometer. Methods for detecting signal from detection systems are well known in the art and need not be further described here.

[0124] When more than one AD biomarker is measured, the biological sample may be divided into a number of aliquots, with separate aliquots used to measure different AD biomarkers (although division of the biological sample into multiple aliquots to allow multiple determinations of the levels of the AD biomarker in a particular sample are also contemplated). Alternately the biological sample (or an aliquot therefrom) may be tested to determine the levels of multiple AD biomarkers in a single reaction using an assay capable of measuring the individual levels of different AD biomarkers in a single assay, such as an array-type assay or assay utilizing multiplexed detection technology (e.g., an assay utilizing detection reagents labeled with different fluorescent dye markers).

[0125] It is common in the art to perform 'replicate' measurements when measuring biomarkers. Replicate measurements are ordinarily obtained by splitting a sample into multiple aliquots, and separately measuring the biomarker(s) in separate reactions of the same assay system. Replicate measurements are not necessary to the methods of the invention, but many embodiments of the invention will utilize replicate testing, particularly duplicate and triplicate testing.

[0126] Reference Levels

[0127] The reference level used for comparison with the measured level for a AD biomarker may vary, depending on aspect of the invention being practiced, as will be understood from the foregoing discussion. For AD diagnosis methods, the "reference level" is typically a predetermined reference level, such as an average of levels obtained from a population that is not afflicted with AD or MCI, but in some instances, the reference level can be a mean or median level from a group of individuals including AD patients. In some instances, the predetermined reference level is derived from (e.g., is the mean or median of) levels obtained from an age-matched population.

[0128] For MCI diagnosis methods (i.e., methods of diagnosing or aiding in the diagnosis of MCI), the reference level is typically a predetermined reference level, such as an average of levels obtained from a population that is not afflicted with AD or MCI, but in some instances, the reference level can be a mean or median level from a group of individuals including MCI and/or AD patients. In some instances, the predetermined reference level is derived from (e.g., is the mean or median of) levels obtained from an age-matched population.

[0129] For AD monitoring methods (e.g., methods of diagnosing or aiding in the diagnosis of AD progression in an AD patient), the reference level may be a predetermined level, such as an average of levels obtained from a population that is not afflicted with AD or MCI, a population that has been diagnosed with MCI or AD, and, in some instances, the reference level can be a mean or median level from a group of individuals including MCI and/or AD patients. Alternately, the reference level may be a historical reference level for the particular patient (e.g., a Leptin level that was obtained from a sample derived from the same

individual, but at an earlier point in time). In some instances, the predetermined reference level is derived from (e.g., is the mean or median of) levels obtained from an age-matched population.

[0130] For AD stratification methods (i.e., methods of stratifying AD patients into mild, moderate and severe stages of AD), the reference level is normally a predetermined reference level that is the mean or median of levels from a population which has been diagnosed with AD or MCI (preferably a population diagnosed with AD). In some instances, the predetermined reference level is derived from (e.g., is the mean or median of) levels obtained from an age-matched population.

[0131] Age-matched populations (from which reference values may be obtained) are ideally the same age as the individual being tested, but approximately age-matched populations are also acceptable. Approximately age-matched populations may be within 1, 2, 3, 4, or 5 years of the age of the individual tested, or may be groups of different ages which encompass the age of the individual being tested. Approximately age-matched populations may be in 2, 3, 4, 5, 6, 7, 8, 9, or 10 year increments (e.g. a "5 year increment" group which serves as the source for reference values for a 62 year old individual might include 58-62 year old individuals, 59-63 year old individuals, 60-64 year old individuals, 61-65 year old individuals, or 62-66 year old individuals).

[0132] Comparing Levels of AD Biomarkers

[0133] The process of comparing a measured value and a reference value can be carried out in any convenient manner appropriate to the type of measured value and reference value for the AD biomarker at issue. As discussed above, 'measuring' can be performed using quantitative or qualitative measurement techniques, and the mode of comparing a measured value and a reference value can vary depending on the measurement technology employed. For example, when a qualitative calorimetric assay is used to measure AD biomarker levels, the levels may be compared by visually comparing the intensity of the colored reaction product, or by comparing data from densitometric or spectrometric measurements of the colored reaction product (e.g., comparing numerical data or graphical data, such as bar charts, derived from the measuring device). However, it is expected that the measured values used in the methods of the invention will most commonly be quantitative values (e.g., quantitative measurements of concentration, such as nanograms of AD biomarker per milliliter of sample, or absolute amount). As with qualitative measurements, the comparison can be made by inspecting the numerical data, by inspecting representations of the data (e.g., inspecting graphical representations such as bar or line graphs).

[0134] A measured value is generally considered to be substantially equal to or greater than a reference value if it is at least 95% of the value of the reference value (e.g., a measured value of 1.71 would be considered substantially equal to a reference value of 1.80). A measured value is considered less than a reference value if the measured value is less than 95% of the reference value (e.g., a measured value of 1.7 would be considered less than a reference value of 1.80).

[0135] The process of comparing may be manual (such as visual inspection by the practitioner of the method) or it may

be automated. For example, an assay device (such as a luminometer for measuring chemiluminescent signals) may include circuitry and software enabling it to compare a measured value with a reference value for an AD biomarker. Alternately, a separate device (e.g., a digital computer) may be used to compare the measured value(s) and the reference value(s). Automated devices for comparison may include stored reference values for the AD biomarker(s) being measured, or they may compare the measured value(s) with reference values that are derived from contemporaneously measured reference samples.

[0136] In some embodiments, the methods of the invention utilize 'simple' or 'binary' comparison between the measured level(s) and the reference level(s) (e.g., the comparison between a measured level and a reference level determines whether the measured level is higher or lower than the reference level). For AD diagnosis biomarkers, a comparison showing that the measured value for the biomarker is lower than the reference value indicates or suggests a diagnosis of AD. For methods relating to the diagnosis of MCI, a comparison showing that measured value for RANTES is lower than the reference value indicates or suggests a diagnosis of AD. In those embodiments relating to diagnosis of MCI which additionally utilize a measured value for Leptin, a comparison showing that RANTES is less than the reference value while Leptin is substantially equal to or greater than the reference level suggests or indicates a diagnosis of MCI.

[0137] As described herein, biological fluid samples may be measured quantitatively (absolute values) or qualitatively (relative values). The respective AD biomarker levels for a given assessment may or may not overlap. As described herein, for some embodiments, qualitative data indicate a given level of cognitive impairment (mild, moderate or severe AD) (which can be measured by MMSE scores) and in other embodiments, quantitative data indicate a given level of cognitive impairment. As shown in Example 4 and under the conditions provided in Example 4 (qualitative data), in those embodiments relating to stratification of AD, a comparison which shows BDNF levels lower than the reference level suggests or indicates mild AD, while a comparison which shows BDNF levels higher than the reference level suggests more advanced AD (i.e., moderate or severe AD), and amongst those samples with BDNF levels higher than the reference level, those also having PDGF-BB levels below the reference level suggest or indicate moderate AD, while those samples also having PDGF-BB levels above the reference level suggest or indicate severe AD. In those embodiments relating to stratification of AD shown in Example 7 (quantitative data), a comparison which shows BDNF levels lower than the reference level where the reference level is Normal suggests or indicates mild AD, while a comparison which shows BDNF levels lower than the reference level where the reference level is Mild AD suggests more advanced AD (i.e., moderate, severe AD), while those samples with leptin levels equal to the reference level where the reference level is Mild AD, those having RANTES levels below the reference level suggest or indicate moderate AD, while those samples with leptin levels equal to the reference level where the reference level is Moderate AD those having PDGF-BB, RANTES, or BDNF levels lower than the reference level suggest or indicate severe AD.

[0138] However, in certain aspects of the invention, the comparison is performed to determine the magnitude of the difference between the measured and reference values (e.g., comparing the 'fold' or percentage difference between the measured value and the reference value). A fold difference that is about equal to or greater than the minimum fold difference disclosed herein suggests or indicates a diagnosis of AD, MCI, progression from MCI to AD, or progression from mild AD to moderate AD, as appropriate to the particular method being practiced. A fold difference can be determined by measuring the absolute concentration of a protein and comparing that to the absolute value of a reference, or a fold difference can be measured by the relative difference between a reference value and a sample value, where neither value is a measure of absolute concentration, and/or where both values are measured simultaneously. A fold difference can be in the range of 10% to 95%. An ELISA measures the absolute content or concentration of a protein from which a fold change is determined in comparison to the absolute concentration of the same protein in the reference. An antibody array measures the relative concentration from which a fold change is determined. Accordingly, the magnitude of the difference between the measured value and the reference value that suggests or indicates a particular diagnosis will depend on the particular AD biomarker being measured to produce the measured value and the reference value used (which in turn depends on the method being practiced). Tables 2A-2B list minimum fold difference values for AD biomarkers for use in methods of the invention which utilize a fold difference in making the comparison between the measured value and the reference value. In those embodiments utilizing fold difference values, a fold difference of about the fold difference indicated in Table 2A suggests a diagnosis of AD, wherein the fold change is a negative value. For example, as described herein, BDNF levels (as measured by ELISA) are decreased in AD patients with mild AD, and BDNF levels decrease further as the severity of the AD intensifies. As shown in Table 6, a BDNF fold change of -46% means a reduction of BDNF levels by 46%. As shown in Table 2A, for qualitative measurements using antibodies, a BDNF fold change of 0.60 means a reduction in BDNF levels by about 60%. Table 2B provides additional information regarding fold changes.

TABLE 2A

Biomarker	Fold Change (as negative value or decrease)
BDNF	0.60
bFGF	0.75
EGF	0.60
FGF-6	0.70
IL-3	0.80
sIL-6 R	0.75
Leptin	0.55
MIP-1 β	0.60
MSP- α	0.80
NAP-2	0.75
NT-3	0.75
PDGF-BB	0.60
RANTES	0.75
SCF	0.80
sTNF RII	0.75
TGF- β 3	0.80
TIMP-1	0.75

TABLE 2A-continued

Biomarker	Fold Change (as negative value or decrease)
TIMP-2	0.80
TNF- β	0.70
TPO	0.75

[0139]

TABLE 2B

Protein	Relative Fold Change (n = 51)	q-value	Absolute Fold Change (n = 187)	p-value
MIP-1d	-0.54291	0.0165		
PDGF-BB	-0.53687	0.0165	-0.135	0.891
LEPTIN(OB)	-0.47625	0.0165	-0.357	0.0018
IL-6 R	-0.6763	0.0165		
BDNF	-0.53628	0.0165	-0.355	0.0006
TIMP-1	-0.71622	0.0165		
RANTES	-0.68299	0.0165	-0.184	0.0144
EGF	-0.56182	0.0165		
TIMP-2	-0.75011	0.0165		
NAP-2	-0.67257	0.0165		
sTNF RII	-0.70029	0.0165		
TNF-b	-0.64998	0.0165		
TPO	-0.71405	0.0165		
FGF-6	-0.66467	0.0165		
NT-3	-0.69805	0.0165		
bFGF	-0.67351	0.0165		
IL-3	-0.75802	0.0165		
SCF	-0.73041	0.0165		
TGF-b3	-0.76912	0.0165		
MSP-a	-0.76466	0.0165		

[0140] As will be apparent to those of skill in the art, when replicate measurements are taken for the biomarker(s) tested, the measured value that is compared with the reference value is a value that takes into account the replicate measurements. The replicate measurements may be taken into account by using either the mean or median of the measured values as the “measured value.”

[0141] Screening Prospective Agents for AD Biomarker Modulation Activity

[0142] The invention also provides methods of screening for candidate agents for the treatment of AD and/or MCI by assaying prospective candidate agents for activity in modulating AD biomarkers. The screening assay may be performed either in vitro and/or in vivo. Candidate agents identified in the screening methods described herein may be useful as therapeutic agents for the treatment of AD and/or MCI.

[0143] The screening methods of the invention utilize the AD biomarkers described herein and AD biomarker polynucleotides as “drug targets.” Prospective agents are tested for activity in modulating a drug target in an assay system. As will be understood by those of skill in the art, the mode of testing for modulation activity will depend on the AD biomarker and the form of the drug target used (e.g., protein or gene). A wide variety of suitable assays are known in the art.

[0144] When the AD biomarker protein itself is the drug target, prospective agents are tested for activity in modulat-

ing levels or activity of the protein itself. Modulation of levels of an AD biomarker can be accomplished by, for example, increasing or reducing half-life of the biomarker protein. Modulation of activity of an AD biomarker can be accomplished by increasing or reducing the availability of the AD biomarker to bind to its cognate receptor(s) or ligand(s).

[0145] When an AD biomarker polynucleotide is the drug target, the prospective agent is tested for activity in modulating synthesis of the AD biomarker. The exact mode of testing for modulatory activity of a prospective agent will depend, of course, on the form of the AD biomarker polynucleotide selected for testing. For example, if the drug target is an AD biomarker polynucleotide, modulatory activity is typically tested by measuring either mRNA transcribed from the gene (transcriptional modulation) or by measuring protein produced as a consequence of such transcription (translational modulation). As will be understood by those in the art, many assay formats will utilize a modified form of the AD biomarker gene where a heterologous sequence (e.g., encoding an expression marker such as an enzyme or an expression tag such as oligo-histidine or a sequence derived from another protein, such as myc) is fused to (or even replaces) the sequence encoding the AD biomarker protein. Such heterologous sequence(s) allow for convenient detection of levels of protein transcribed from the drug target.

[0146] Prospective agents for use in the screening methods of the invention may be chemical compounds and/or complexes of any sort, including both organic and inorganic molecules (and complexes thereof). As will be understood in the art, organic molecules are most commonly screened for AD biomarker modulatory activity. In some situations, the prospective agents for testing will exclude the target AD biomarker protein.

[0147] Screening assays may be in any format known in the art, including cell-free in vitro assays, cell culture assays, organ culture assays, and in vivo assays (i.e., assays utilizing animal models of AD and MCI). Accordingly, the invention provides a variety of embodiments for screening prospective agents to identify candidate agents for the treatment of AD and/or MCI.

[0148] In some embodiments, prospective agents are screened to identify candidate agents for the treatment of AD and/or MCI in a cell-free assay. Each prospective agent is incubated with the drug target in a cell-free environment, and modulation of the AD biomarker is measured. Cell-free environments useful in the screening methods of the invention include cell lysates (particularly useful when the drug target is an AD biomarker gene) and biological fluids such as whole blood or fractionated fluids derived therefrom such as plasma and serum (particularly useful when the AD biomarker protein is the drug target). When the drug target is an AD biomarker gene, the modulation measured may be modulation of transcription or translation. When the drug target is the AD biomarker protein, the modulation may be of the half-life of the protein or of the availability of the AD biomarker protein to bind to its cognate receptor or ligand.

[0149] In other embodiments, prospective agents are screened to identify candidate agents for the treatment of AD and/or MCI in a cell-based assay. Each prospective agent is incubated with cultured cells, and modulation of target AD biomarker is measured. In certain embodiments, the cultured

cells are astrocytes, neuronal cells (such as hippocampal neurons), fibroblasts, or glial cells. When the drug target is an AD biomarker gene, transcriptional or translational modulation may be measured. When the drug target is the AD biomarker protein, the AD biomarker protein is also added to the assay mixture, and modulation of the half-life of the protein or of the availability of the AD biomarker protein to bind to its cognate receptor or ligand is measured.

[0150] Further embodiments relate to screening prospective agents to identify candidate agents for the treatment of AD and/or MCI in organ culture-based assays. In this format, each prospective agent is incubated with either a whole organ or a portion of an organ (such as a portion of brain tissue, such as a brain slice) derived from a non-human animal and modulation of the target AD biomarker is measured. When the drug target is an AD biomarker gene, transcriptional or translational modulation may be measured. When the drug target is the AD biomarker protein, the AD biomarker protein is also added to the assay mixture, and modulation of the half-life of the protein or of the availability of the AD biomarker protein to bind to its cognate receptor is measured.

[0151] Additional embodiments relate to screening prospective agents to identify candidate agents for the treatment of AD and/or MCI utilizing in vivo assays. In this format, each prospective agent is administered to a non-human animal and modulation of the target AD biomarker is measured. Depending on the particular drug target and the aspect of AD and/or MCI treatment that is sought to be addressed, the animal used in such assays may either be a "normal" animal (e.g., C57 mouse) or an animal which is a model of AD or MCI. A number of animal models of AD are known in the art, including the 3xTg-AD mouse (Caccamo et al., 2003, *Neuron* 39(3):409-21), mice over expressing human amyloid beta precursor protein (APP) and presenilin genes (Westaway et al., 1997, *Nat. Med.* 3(1):67-72), and others (see Higgins et al., 2003, *Behav. Pharmacol.* 14(5-6):419-38). When the drug target is an AD biomarker gene, transcriptional or translational modulation may be measured. When the drug target is the AD biomarker protein, modulation of the half-life of the target AD biomarker or of the availability of the AD biomarker protein to bind to its cognate receptor or ligand is measured. The exact mode of measuring modulation of the target AD biomarker will, of course, depend on the identity of the AD biomarker, the format of the assay, and the preference of the practitioner. A wide variety of methods are known in the art for measuring modulation of transcription, translation, protein half-life, protein availability, and other aspects which can be measured. In view of the common knowledge of these techniques, they need not be further described here.

[0152] Kits

[0153] The invention provides kits for carrying out any of the methods described herein. Kits of the invention may comprise at least one reagent specific for an AD biomarker, and may further include instructions for carrying out a method described herein. Kits may also comprise AD biomarker reference samples, that is, useful as reference values. "AD diagnosis markers" for use in kits provided herein include, but are not limited to GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2;

ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In other examples, "AD diagnosis biomarkers" for use in kits provided herein include but are not limited to basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), Leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), tumor necrosis factor beta (TNF-β). In other examples, kits comprise any one, two, three or four of the AD diagnosis markers Leptin, RANTES, PDGF-BB and BDNF.

[0154] More commonly, kits of the invention comprise at least two different AD biomarker-specific affinity reagents, where each reagent is specific for a different AD biomarker. In some embodiments, kits comprise at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 reagents specific for an AD biomarker. In some embodiments, the reagent(s) specific for an AD biomarker is an affinity reagent.

[0155] Kits comprising a single reagent specific for an AD biomarker will generally have the reagent enclosed in a container (e.g., a vial, ampoule, or other suitable storage container), although kits including the reagent bound to a substrate (e.g., an inner surface of an assay reaction vessel) are also contemplated. Likewise, kits including more than one reagent may also have the reagents in containers (separately or in a mixture) or may have the reagents bound to a substrate.

[0156] In some embodiments, the AD biomarker-specific reagent(s) will be labeled with a detectable marker (such as a fluorescent dye or a detectable enzyme), or be modified to facilitate detection (e.g., biotinylated to allow for detection with an avidin- or streptavidin-based detection system). In other embodiments, the AD biomarker-specific reagent will not be directly labeled or modified.

[0157] Certain kits of the invention will also include one or more agents for detection of bound AD biomarker-specific reagent. As will be apparent to those of skill in the art, the identity of the detection agents will depend on the type of AD biomarker-specific reagent(s) included in the kit, and the intended detection system. Detection agents include antibodies specific for the AD biomarker-specific reagent (e.g., secondary antibodies), primers for amplification of an AD biomarker-specific reagent that is nucleotide based (e.g., aptamer) or of a nucleotide 'tag' attached to the AD biomarker-specific reagent, avidin- or streptavidin-conjugates for detection of biotin-modified AD biomarker-specific reagent(s), and the like. Detection systems are well known in the art, and need not be further described here. Accordingly, provided herein are kits for identifying an individual with mild cognitive impairment (MCI), comprising at least

one reagent specific for RANTES; and instructions for carrying out the method. In some examples, the kits further comprise a reagent specific for leptin. In other examples, provided herein are kits for monitoring progression of Alzheimer's disease (AD) in an AD patient, comprising at least one reagent specific for leptin; and instructions for carrying out the method. Also provided herein are kits for stratifying an Alzheimer's disease (AD) patient, comprising at least one reagent specific for brain derived neurotrophic factor (BDNF); at least one reagent specific for BB homodimeric platelet derived growth factor (PDGF-BB); and instructions for carrying out the method.

[0158] A modified substrate or other system for capture of AD biomarkers may also be included in the kits of the invention, particularly when the kit is designed for use in a sandwich-format assay. The capture system may be any capture system useful in an AD biomarker assay system, such as a multi-well plate coated with an AD biomarker-specific reagent, beads coated with an AD biomarker-specific reagent, and the like. Capture systems are well known in the art and need not be further described here.

[0159] In certain embodiments, kits according to the invention include the reagents in the form of an array. The array includes at least two different reagents specific for AD biomarkers (each reagent specific for a different AD biomarker) bound to a substrate in a predetermined pattern (e.g., a grid). Accordingly, the present invention provides arrays comprising "AD diagnosis markers" including, but not limited to GCSF; IFN- γ ; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; AcRP30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1 α ; MSP-a; SCF; TGF- β 3; TNF- β MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. In other examples, "AD diagnosis biomarkers" include but are not limited to basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), Leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1 δ), macrophage stimulating protein alpha chain (MSP- α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF- β 3), tumor necrosis factor beta (TNF- β). In other examples, arrays comprise any one, two, three or four of the AD diagnosis markers Leptin, RANTES, PDGF-BB and BDNF. The localization of the different AD biomarker-specific reagents (the "capture reagents") allows measurement of levels of a number of different AD biomarkers in the same reaction. Kits including the reagents in array form are commonly in a sandwich format, so such kits may also comprise detection reagents. Normally, the kit will include different detection reagents, each detection reagent specific to a different AD biomarker. The detection reagents in such embodiments are normally reagents specific for the same AD biomarkers as the reagents bound to the substrate (although the detection reagents typically bind to a different portion or site on the AD biomarker target than the substrate-

bound reagents), and are generally affinity-type detection reagents. As with detection reagents for any other format assay, the detection reagents may be modified with a detectable moiety, modified to allow binding of a separate detectable moiety, or be unmodified. Array-type kits including detection reagents that are either unmodified or modified to allow binding of a separate detectable moiety may also contain additional detectable moieties (e.g., detectable moieties which bind to the detection reagent, such as labeled antibodies which bind unmodified detection reagents or streptavidin modified with a detectable moiety for detecting biotin-modified detection reagents).

[0160] The instructions relating to the use of the kit for carrying out the invention generally describe how the contents of the kit are used to carry out the methods of the invention. Instructions may include information as sample requirements (e.g., form, pre-assay processing, and size), steps necessary to measure the AD biomarker(s), and interpretation of results.

[0161] Instructions supplied in the kits of the invention are typically written instructions on a label or package insert (e.g., a paper sheet included in the kit), but machine-readable instructions (e.g., instructions carried on a magnetic or optical storage disk) are also acceptable. In certain embodiments, machine-readable instructions comprise software for a programmable digital computer for comparing the measured values obtained using the reagents included in the kit.

[0162] The following Examples are provided to illustrate the invention, but are not intended to limit the scope of the invention in any way.

EXAMPLES

Example 1

AD Diagnosis Biomarkers

[0163] We compared plasma protein expression levels for 120 proteins in 32 cases of serum collected from patients with Alzheimer's Disease (with a mean age of 74) to 19 cases of serum collected from control subjects (also with mean age of 74). Alzheimer's Disease subjects were clinically diagnosed with AD by a neurologist, and had Mini Mental State Exam (MMSE) scores ranging from 26-14.

[0164] Plasma samples were assayed using a sandwich-format ELISA on a nitrocellulose filter substrate. Plasma samples were diluted 1:10 in phosphate buffer and incubated with the capture substrate (a nitrocellulose membrane spotted with capture antibodies). The samples were incubated with the capture substrate for two hours at room temperature, then decanted from the capture substrate. The substrate was washed twice with 2 ml of washing buffer (1 $\times$ PBS; 0.05% Tween-20) at room temp, then incubated with biotinylated detection antibodies for two hours at room temperature. The capture antibody solution was decanted and the substrate was washed twice for 5 min with washing buffer. The washed substrate was then incubated with horseradish peroxidase/streptavidin conjugate for 45 minutes, at which time the conjugate solution was decanted and the membranes were washed with washing buffer twice for 5 minutes. The substrate was transferred onto a piece of filter paper, incubated in enhanced chemiluminescence (ECL)

Detection Buffer solution purchased from Raybiotech, Inc. Chemiluminescence was detected and quantified with a chemiluminescence imaging camera. Signal intensities were normalized to standard proteins blotted on the substrate and used to calculate relative levels of biomarkers. In other examples, signal intensities were normalized to the median and used to calculate relative levels of biomarkers.

[0165] Relative biomarker levels in plasma are compared between control and AD groups revealing 46 discriminatory biomarkers: GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R. An unsupervised clustering (that is, the clustering algorithm does not know which cases are AD and which are normal) of the 46 discriminatory markers results in the clustering of the samples into 2 groups or clusters, a cluster of control samples, and a cluster of AD samples. Sensitivity was calculated as the number of correctly classed AD samples in the AD cluster/total number of AD samples, which is 29/32 or 90.6%. Specificity was calculated as total number of correctly classed control samples in the control cluster/total number of controls, which is (14/19=73.6%).

[0166] Biomarker levels were compared between control and AD groups, revealing 20 biomarkers (shown in Table 3) that are differentially regulated (each is decreased in AD as compared to control) between the two groups. Statistical analysis was performed to find the probability that the finding of differential levels was in error (the "q" value) for any one biomarker. Biomarkers with differential levels and associated q values (shown as percentage values) are shown in Table 3 (fold change indicates the fold change between levels in control vs. AD samples). Sensitivity was calculated as number of AD samples in AD cluster/total number of AD samples, which is 29/32 or 90.6%. Specificity was calculated as total correctly predicted AD/total predicted AD (29/34=85%).

TABLE 3

Qualitative Biomarker	Fold Change (as negative value or decrease)	q-value (%)
Brain derived neurotrophic factor (BDNF)	0.536	1.656
Basic fibroblast growth factor (bFGF)	0.673	1.656
Epidermal growth factor (EGF)	0.561	1.656
Fibroblast growth factor-6 (FGF-6)	0.664	1.656
Interleukin-3 (IL-3)	0.758	1.656
Soluble interleukin-6 receptor (sIL-6 R)	0.676	1.656
Leptin (also known as OB)	0.476	1.656
Macrophage inflammatory protein 1-delta (MIP-1δ)	0.542	1.656
MSP-a	0.764	1.656
NAP-2	0.672	1.656
Neurotrophin-3 (NT-3)	0.698	1.656
Platelet derived growth factor, BB dimer (PDGF-BB)	0.536	1.656
RANTES	0.682	1.656
Stem cell factor (SCF)	0.730	1.656
sTNF RII	0.700	1.656
Transforming growth factor beta-3 (TGF-β3)	0.769	1.656

TABLE 3-continued

Qualitative Biomarker	Fold Change (as negative value or decrease)	q-value (%)
Tissue inhibitor of metalloproteases-1 (TIMP-1)	0.716	1.656
Tissue inhibitor of metalloproteases-2 (TIMP-2)	0.750	1.656
Tumor necrosis factor beta (TNF-β)	0.649	1.656
TPO	0.714	1.656

Example 2

Decision Trees from AD Diagnosis Marker Data

[0167] Upon further analysis of the data from example 1, two different decision trees were formulated for diagnosis of AD using AD diagnosis biomarkers.

[0168] The first decision tree utilizes sIL-6R, IL-8, and TIMP-1 levels. The rules which make up the decision tree are: (1) If sIL-6R ≤ 5.18 and IL-8 is ≤ 0.957 , the indication is normal; (2) if sIL-6R ≤ 5.18 and IL-8 > 0.957 , the indication is AD; (3) if sIL-6R > 5.18 and TIMP-1 ≤ 7.978 , the indication is AD; and (4) if sIL-6R > 5.18 and TIMP-1 is > 7.978 , the indication is normal, wherein the values expressed are relative concentrations.

[0169] Accuracy of this decision tree was measured using 10-fold cross-validation testing feature in CART to generate misclassification rates for learning samples and testing samples. Sensitivity was calculated from the testing scores as number of AD samples correctly predicted as AD/total number of AD samples (29/32=0.906). Specificity was calculated from the testing scores as total correctly predicted cases of AD/total number of cases predicted AD (29/33=0.878).

[0170] A second decision tree was formulating using BDNF, TIMP-1 and MIP-1δ levels. The rules which make up the decision tree are: (1) if BDNF > 4.476 , the indication is normal; (2) if BDNF ≤ 4.476 and TIMP-1 ≤ 8.942 , the indication is AD; (3) if BDNF ≤ 4.476 , TIMP-1 > 8.942 , and MIP-1δ ≤ 1.89 , the indication is AD; and (4) if BDNF < 4.476 , TIMP-1 > 8.942 , and MIP-1δ > 1.89 , the indication is normal. Accuracy of this decision tree was measured using 10-fold cross-validation testing feature in CART to generate misclassification rates for learning samples and testing samples. Sensitivity was calculated from the testing scores as number of AD samples correctly predicted as AD/total number of AD samples (0.875). Specificity was calculated from the testing scores as total correctly predicted cases of AD/total number of cases predicted AD (0.82).

Example 3

Diagnosis of MCI

[0171] Levels of RANTES and Leptin were measured in 18 samples from control subjects (mean age=74) and 6 samples from patients diagnosed with mild cognitive impairment (MCI). MCI patients had been clinically diagnosed by a neurologist, and had an AULT-A7 score of less than 5 and Mini Mental State Exam (MMSE) scores ranging from

30-28. Control subjects had an AULT-A7 score greater than or equal to 5 and MMSE score ranging from 30-28.

[0172] RANTES and Leptin levels were measured using an ELISA kit from R&D systems according to the manufacturer's instructions. The raw ELISA expressions values were normalized by dividing each value by the median of all the samples. Analysis of the data showed (a) Leptin is not decreased in MCI patients as compared to control subjects (in the six MCI samples, Leptin was actually 11% higher than the control subjects), and (b) a bimodal distribution of RANTES, where MCI patients had RANTES levels of between 1.043 and 1.183 (levels from control subjects were either ≤ 1.043 or > 1.183). However, closer inspection of the data led us to believe that those control subjects with RANTES ≤ 1.043 had been incorrectly classified as normal (and should have been diagnosed as MCI).

[0173] Reclassification of control subjects with RANTES ≤ 1.043 as MCI patients allows the creation of a simple rule: if RANTES ≤ 1.183 and Leptin ≥ 0.676 , the indication is MCI. Sensitivity and specificity, calculated as described in Example 2, were 83.3% and 88.8%, respectively.

Example 4

Monitoring and Stratification of AD Patients

[0174] Levels of RANTES, Leptin, PDGF-BB, and BDNF were measured in serum samples collected from 36 patients diagnosed with Alzheimer's Disease. (mean age of 74) using ELISA kits from R&D systems according to the manufacturer's instructions. The raw ELISA expressions values were normalized by dividing each value by the median of all the samples. The samples were grouped into three classes on the basis of MMSE score: Class 1 (mild AD), MMSE 27-22; Class 2 (moderate AD), MMSE 21-16; and Class 3 (severe AD), MMSE 15-12.

[0175] Upon analysis of the ELISA data, we formulated a decision tree using BDNF and PDGF-BB. The rules which make up the decision tree are: (1) if BDNF ≤ 0.626 , the indication is mild AD; (2) if BDNF > 0.626 and PDGF-BB ≤ 0.919 , the indication is moderate AD; and (3) if BDNF > 0.626 and PDGF-BB > 0.919 , the indication is severe AD. The values expressed are relative concentrations that have been normalized to the median. Average normalized levels for Leptin were: Class I=0.886; class II=0.757; class III=0.589. Average normalized levels for BDNF were: Class I=0.595; class II=0.956; class III=1.23. When applied to a set of "test" data, the decision tree produced 58%, 47%, and 57% percent correct stratification of the test samples into mild, moderate, and severe categories.

Example 5

Four Discriminatory Markers

[0176] The absolute concentrations in plasma of only 4 discriminatory markers, BDNF, PDGF-BB, LEPTIN, and RANTES measured by ELISA was used to classify samples. ELISA kits were purchased from R&D Systems, and measurements were obtained according to manufacturer recommendations. For example for RANTES, the following protocol was followed.

[0177] 1. Add 50 μ L standards, specimens or controls to appropriate wells.

[0178] 2. Add 50 μ L anti-RANTES Biotin Conjugate to each well.

[0179] 3. Incubate wells at 37° C. for 1 hour.

[0180] 4. Aspirate and wash wells 4x with Working Wash Buffer.

[0181] 5. Add 100 μ L Streptavidin-HRP Working Conjugate to each well.

[0182] 6. Incubate for 30 minutes at room temperature.

[0183] 7. Aspirate and wash wells 4x with Working Wash Buffer.

[0184] 8. Add 100 μ L of Stabilized Chromogen to each well.

[0185] 9. Incubate at room temperature for 30 minutes in the dark.

[0186] 10. Add 100 μ L of Stop Solution to each well.

[0187] 11. Read absorbance at 450 nm.

[0188] Following the above protocol, an unsupervised clustering of BDNF, PDGF-BB, LEPTIN, and RANTES was performed using the publicly available web based clustering software wCLUTO at cluto.ccg.umn.edu/cgi-bin/wCluto/wCluto.cgi. Here the clustering of the 4 proteins resulted in the clustering of the samples into 2 groups or clusters, a cluster of control samples and a cluster of AD samples. Sensitivity was calculated as the number of correctly classed AD samples in the AD cluster/total number of AD samples, which is 21/24 or 87.5%. Specificity was calculated as total number of correctly classed control samples in the control cluster/total number of controls, which is 20/24=83.3%.

[0189] Additionally, absolute biomarker levels in plasma (as measured by ELISA) for BDNF, PDGF-BB, and LEPTIN, were correlated with MMSE scores (range 12-30). AD could be identified in MMSE scores in a range of 12-28 and control samples were identified in MMSE scores in the range of 25-30. Table 4 shows the correlations and their statistical significance (p-value). The upper and lower correlations show whether the upper end of the range of MMSE scores and biomarker concentrations or the lower end of the range of MMSE scores and biomarker concentrations are more correlated. Therefore, the correlations show that higher levels of BDNF and Leptin are significantly correlated with better MMSE scores, and that increase in the concentration of BDNF and Leptin from a reference point or an earlier collection is an indication of improvement in cognition as measured by MMSE. Simultaneously, or by itself, the lower the levels of PDGF-BB in men is significantly correlated with better MMSE scores, and a decrease in the concentration of PDGF-BB in male sample compared to an earlier collection in that male, is an indication of improvement in cognition as measured by MMSE.

[0190] The results show (Table 4) the correlation between the plasma concentration of 3 discriminatory proteins for AD to the MMSE score of the subjects and the correlation between concentrations of proteins that are discriminatory for AD. There was no correlation between MMSE score and Age among AD subjects and there was no correlation between Age and the concentration of BDNF, PDGF-BB, or

LEPTIN in plasma among AD subjects. The p-values show that the correlations are statistically significant. The count shows the number of cases. BDNF has a statistically significant positive correlation with MMSE scores. PDGF-BB has a statistically significant negative correlation with MMSE scores in men. LEPTIN has a statistically significant positive correlation with MMSE scores. This experiment demonstrates that plasma concentrations for PDGF-BB, LEPTIN, and BDNF can be used to monitor the progression of cognitive decline.

TABLE 4

	Correlation	Count	Z-value	P-value	95% Lower	95% Upper
BDNF to MMSE	0.184	165	2.373	0.0176	0.032	0.328
BDNF to MMSE (Females)	0.229	91	2.18	0.0289	0.024	0.415
PDGF-BB to MMSE (Males)	-0.207	74	-1.769	0.0768	-0.416	0.023
LEPTIN to MMSE	0.193	164	2.478	0.0132	0.041	0.336
BDNF to PDGF-BB	0.700	181	11.575	0.0001	0.617	0.768
PDGF-BB to RANTES	0.563	181	8.5	0.0001	0.454	0.655
BDNF to RANTES	0.714	181	11.9	0.0001	0.634	0.779

[0191] Controls and AD cases were age matched, and had a mean age of 74. The mean MMSE score for AD cases (n=24) was 20, while the mean MMSE score for Control cases (n=24) was 30. Classification of the samples was performed with unsupervised clustering of protein concentration. The total accuracy of classification was 85.4%. This results demonstrated that plasma protein concentrations for BDNF, PDGF-BB, LEPTIN, and RANTES, as measured by ELISA can be used to accurately discriminate between AD and controls.

Example 6

Validation of Mean Protein Concentrations in AD and Controls by ELISA

[0192] Protein concentrations for proteins, LEPTIN, BDNF and RANTES, in plasma samples of AD (n=95) to age matched Controls (n=88) are shown in FIGS. 1A-1C. One of the four proteins we measured was Brain Derived Neurotrophic Factor (BDNF). The mean concentration of BDNF in AD plasma was 8.1 ng/ml (SE+/-0.4) compared to the mean of control plasma 10.8ng/ml (SE+/-0.68) and the difference was found to be extremely statistically significant (p-value=0.0006). We also found that the concentrations of BDNF were lower in other forms of dementia (5.74ng/ml, n=20) than AD. The mean concentration of a second protein Leptin in AD plasma was found to be 10.9 ng/ml (SE+/-1.06) compared to the mean of control plasma 17.4 ng/ml (SE+/-1.8) and the difference was found to be statistically very significant (p-value=0.0018). The mean concentration of a third protein Rantes in AD plasma was found to be 66.3 ng/ml (SE+/-2.4) compared to control samples 74.5 ng/ml (SE+/-3.2) and the difference was found to be statistically significant (p-value=0.0403). No difference in the means of concentrations for RANTES, PDGF-BB, and BDNF were observed among AD subjects with MMSE scores >=20 (n=54) and those <20 (n=41).

Example 7

Absolute Biomarker Concentrations in Plasma

[0193] Additionally, absolute biomarker concentrations in plasma were measured for BDNF, and mean concentrations for Controls was compared to MCI (Mild Cognitive Impairment), MMSE 25-28, MMSE 20-25, and MMSE 10-20. For the purposes of this experiment, the index used in the following example is: questionable AD is=MMSE score in the range of 25-28; mild AD=MMSE score in the range of

20-25; and moderate AD=MMSE score in the range of 10-20 and severe AD=MMSE score in the range of 10-20. For the purpose of Example 7, all individuals assessed as having Questionable AD were diagnosed by a physician as having AD. The FIG. 2 shows that mean concentrations of BDNF in plasma for MMSE 25-28; MMSE 20-25; MMSE 10-20 are significantly lower than the mean concentration in Controls (Normal, mean age 74) and the mean concentration of BDNF in MCI is significantly higher than in Controls and all cases of AD. FIG. 2.

Unpaired t-test for BDNF plasma

Grouping Variable: stage

Hypothesized Difference = 0

Inclusion criteria: Sparks from Center All

	Mean Diff.	DF	t-Value	P-Value
MCI, mild	6349.252	47	3.050	.0038
MCI, moderate	6828.574	31	2.651	.0125
MCI, normal	3961.358	86	1.442	.1529
MCI, questionable	7547.218	17	2.550	.0207
mild, moderate	479.322	68	.460	.6467
mild, normal	-2387.894	123	-2.270	.0250
mild, questionable	1197.966	54	.969	.3369
moderate, normal	-2867.216	107	-2.175	.0319
moderate, questionable	718.644	38	.475	.6372
normal, questionable	3585.860	93	1.993	.0492

[0194]

Group Info for BDNF plasma

Grouping Variable: stage

Inclusion criteria: Sparks from Center All

	Count	Mean	Variance	Std. Dev.	Std. Err
MCI	6	14879.833	85932530.967	9269.980	3784.454
mild	43	8530.581	15299257.963	3911.427	596.487
moderate	27	8051.259	22317487.815	4724.139	909.161

-continued

Group Info for BDNF plasma					
Grouping Variable: stage					
Inclusion criteria: Sparks from Center All					
	Count	Mean	Variance	Std. Dev.	Std. Err
normal	82	10918.476	39478328.993	6283.178	693.861
question- able	13	7332.615	15122872.923	3888.814	1078.563

[0195] Additionally, absolute concentrations of BDNF, in plasma samples collected from four separate Alzheimer's Centers was compared for gender differences in mean concentrations between AD (Females) and Control (Females) and AD (Males) and Control (Males). FIG. 3 shows that there is 40% difference in the concentration of BDNF in AD Females compared to Control Females and the difference is highly statistically significant (p-value=0.004). The difference in the mean concentration of BDNF for all AD cases compared to all Control case was found to be extremely statistically significant (p-value=0.0006).

Unpaired t-test for BDNF plasma				
Grouping Variable: Disease				
Split By: sex				
Hypothesized Difference = 0				
Row exclusion: Center All				
	Mean Diff.	DF	t-Value	P-Value
AD, Control: Total	-2974.140	187	-3.482	.0006
AD, Control: F	-3939.353	87	-2.924	.0044
AD, Control: M	-1348.601	92	-1.165	.2469

[0196] Results for totals may not agree with results for individual cells because of missing values for split variables.

Group Info for BDNF plasma					
Grouping Variable: Disease					
Split By: sex					
Row exclusion: Center All					
	Count	Mean	Variance	Std. Dev.	Std. Err
AD: Total	106	5596.113	24323422.844	4931.878	479.026
AD: F	38	5775.921	25121499.318	5012.135	813.076
AD: M	62	5396.774	24336564.079	4933.210	626.518
Control: Total	83	8570.253	46322420.606	6806.058	747.062
Control: F	51	9715.275	50173107.603	7083.298	991.860
Control: M	32	6745.375	36011373.274	6000.948	1060.828

[0197] Results for totals may not agree with results for individual cells because of missing values for split variables.

[0198] Additionally, absolute biomarker concentrations in plasma were measured for RANTES in plasma samples collected from four different Alzheimer's Centers, and mean concentrations for Controls were compared to MCI (Mild Cognitive Impairment), MMSE 25-28; (MMSE 20-25; MMSE 10-20; and MMSE 10-20. The index is described above. The mean differences between Mild AD compared to

Moderate AD, Mild AD compared to Normal, Mild AD compared to Severe AD, Moderate AD compared to Normal, Questionable AD compared to Normal, Normal to Severe AD were all found to be statistically significant. FIG. 4.

Unpaired t-test for RANTES ELISA				
Grouping Variable: stage				
Hypothesized Difference = 0				
Row exclusion: Center All				
	Mean Diff.	DF	t-Value	P-Value
MCI, mild	84.789	64	.007	.9945
MCI, moderate	12454.688	51	1.042	.3022
MCI, normal	-10422.892	106	-.866	.3884
MCI, questionable	9682.438	29	.682	.5007
MCI, severe	50349.200	10	1.647	.1305
mild, moderate	12369.899	97	1.814	.0728
mild, normal	-10507.681	152	-1.775	.0780
mild, questionable	9597.649	75	1.081	.2830
mild, severe	50264.411	56	2.031	.0470
moderate, normal	-22877.580	139	-3.606	.0004
moderate, questionable	-2772.250	62	-.315	.7535
moderate, severe	37894.512	43	1.647	.1069
normal, questionable	20105.330	117	2.353	.0203
normal, severe	60772.092	98	2.395	.0185
questionable, severe	40666.762	21	1.624	.1192

[0199]

Group Info for RANTES ELISA					
Grouping Variable: stage					
Row exclusion: Center All					
	Count	Mean	Variance	Std. Dev.	Std. Err
MCI	10	54919.200	1729660285.733	41589.185	13151.655
mild	56	54834.411	1203622609.701	34693.265	4636.082
moderate	43	42464.512	1036226732.256	32190.476	4909.002
normal	98	65342.092	1275358885.672	35712.167	3607.474
question- able	21	45236.762	1201710117.890	34665.691	7564.674
severe	2	4570.000	2976800.000	1725.341	1220.000

[0200] Additionally, absolute biomarker concentrations in plasma were measured for Leptin in plasma samples collected from four different Alzheimer's Centers, and mean concentrations for Controls were compared to MCI (Mild Cognitive Impairment); MMSE 25-28; MMSE 20-25; MMSE 10-20; and MMSE 10-20. The mean differences between Questionable AD compared to MCI, Mild AD compared to Normal, Mild AD compared to Questionable AD, Questionable AD compared to Normal, and Moderate AD compared to Normal were all found to be statistically significant. FIG. 5.

-continued

Unpaired t-test for Leptin ELISA Grouping Variable: stage Hypothesized Difference = 0 Row exclusion: Center All				
	Mean Diff.	DF	t-Value	P-Value
MCI, mild	4164.889	64	1.338	.1856
MCI, moderate	4707.044	51	1.061	.2939
MCI, normal	-650.092	105	-.123	.9022
MCI, questionable	7793.348	29	2.000	.0550
MCI, severe	8187.800	10	.739	.4767
mild, moderate	542.155	97	.272	.7860
mild, normal	-4814.981	151	-2.117	.0359
mild, questionable	3628.458	75	1.897	.0617
mild, severe	4022.911	56	.734	.4661
moderate, normal	-5357.136	138	-1.963	.0516
moderate, questionable	3086.303	62	1.085	.2822
moderate, severe	3480.756	43	.403	.6892
normal, questionable	8443.439	116	2.368	.0195
normal, severe	8837.892	97	.778	.4383
questionable, severe	394.452	21	.078	.9383

[0201]

Group Info for Leptin ELISA Grouping Variable: stage Row exclusion: Center All					
	Count	Mean	Variance	Std. Dev.	Std. Err
MCI	10	15727.300	225300738.678	15010.021	4746.585
mild	56	11562.411	58790550.756	7667.500	1024.613
moderate	43	11020.256	145797834.909	12074.677	1841.371
normal	97	16377.392	255125297.032	15972.642	1621.776
questionable	21	7933.952	47833192.348	6916.154	1509.229
severe	2	7539.500	16125520.500	4015.659	2839.500

[0202] Additionally, absolute biomarker concentrations in plasma were measured for PDGF-BB in plasma samples collected from four different Alzheimer's Centers, and mean concentrations for Controls were compared to MCI (Mild Cognitive Impairment); MMSE 25-28; MMSE 20-25; MMSE 10-20; and MMSE 10-20. The mean differences between Questionable AD compared to Mild AD, Mild AD compared to Severe AD, Moderate AD compared to Severe AD, Normal compared to Questionable AD, and Normal to Severe AD were all found to be statistically significant. FIG. 6.

Unpaired t-test for PDGF-BB ELISA Grouping Variable: stage Hypothesized Difference = 0 Row exclusion: Center All				
	Mean Diff.	DF	t-Value	P-Value
MCI, mild	-62.275	58	-.286	.7756
MCI, moderate	81.595	44	.411	.6831
MCI, normal	-42.865	103	-.210	.8343
MCI, questionable	191.571	28	.810	.4246
MCI, severe	637.000	9	1.072	.3117
mild, moderate	143.869	86	1.285	.2023
mild, normal	19.410	145	.199	.8426

Unpaired t-test for PDGF-BB ELISA Grouping Variable: stage Hypothesized Difference = 0 Row exclusion: Center All				
	Mean Diff.	DF	t-Value	P-Value
mild, questionable	253.846	70	1.812	.0742
mild, severe	699.275	51	1.745	.0871
moderate, normal	-124.459	131	-1.201	.2320
moderate, questionable	109.977	56	.869	.3885
moderate, severe	555.405	37	1.716	.0945
normal, questionable	234.436	115	1.767	.0799
normal, severe	679.865	96	1.696	.0931
questionable, severe	445.429	21	1.278	.2153

[0203]

Group Info for PDGF-BB ELISA Grouping Variable: stage Row exclusion: Center All					
	Count	Mean	Variance	Std. Dev.	Std. Err
MCI	9	731.000	650139.000	806.312	268.771
mild	51	793.275	315391.883	561.598	78.639
moderate	37	649.405	204231.470	451.920	74.295
normal	96	773.865	318171.171	564.067	57.570
questionable	21	539.429	233024.657	482.726	105.340
severe	2	94.000	648.000	25.456	18.000

[0204] Additionally, absolute biomarker concentrations in plasma were measured for BDNF in plasma samples collected from four different Alzheimer's centers, and means concentrations for Controls were compared to MCI (Mild Cognitive Impairment), Questionable AD (MMSE 25-28), Mild differences between MCI compared to Moderate AD, MCI compared to Questionable AD, Mild AD to Normal, Mild AD to severe AD, Moderate to Normal, Normal to Questionable AD, and Normal to Severe were all found to be statistically significant. FIG. 7.

Unpaired t-test for BDNF plasma Grouping Variable: stage Hypothesized Difference = 0 Row exclusion: Center All				
	Mean Diff.	DF	t-Value	P-Value
MCI, mild	2819.186	64	1.433	.1568
MCI, moderate	4071.016	51	1.877	.0663
MCI, normal	124.278	106	.053	.9578
MCI, questionable	4535.757	29	1.806	.0813
MCI, severe	8660.400	10	1.202	.2570
mild, moderate	1251.831	97	1.262	.2098
mild, normal	-2694.908	152	-2.638	.0092
mild, questionable	1716.571	75	1.447	.1520
mild, severe	5841.214	56	1.726	.0898
moderate, normal	-3946.739	139	-3.431	.0008
moderate, questionable	464.741	62	.360	.7199
moderate, severe	4589.384	43	1.265	.2128
normal, questionable	4411.480	117	2.868	.0049
normal, severe	8536.122	98	1.781	.0781
questionable, severe	4124.643	21	1.321	.2006

[0205]

Example 9

Additional Biomarkers

Group Info for BDNF plasma Grouping Variable: stage Row exclusion: Center All					
	Count	Mean	Variance	Std. Dev.	Std. Err
MCI	10	9511.900	96113654.322	9803.757	3100.220
mild	56	6692.714	22509096.208	4744.375	633.994
moderate	43	5440.884	25765123.534	5075.936	774.073
normal	98	9387.622	45504479.969	6745.701	681.419
question- able	21	4976.143	18681976.129	4322.265	943.196
severe	2	851.500	63724.500	252.437	178.500

[0206] It has been found that for Questionable AD (MMSE score in the range of 25-28) the levels of Leptin and

[0208] Additionally, qualitative biomarker levels for GDNF, SDF-1, IGFBP3, FGF-6, TGF-b3, BMP-4, NT-3, EGF, BDNF, IGFBP-2 were correlated with MMSE scores (range 12-30) for AD (MMSE range 12-28) and control samples (MMSE range 25-30). Table 5 shows the correlations and their statistical significance (p-value). The upper and lower correlations show whether the upper end of the range of MMSE Scores and biomarker concentrations or the lower end of the range of MMSE scores and biomarker concentrations are more correlated. A negative correlation means that MMSE scores increase with decreasing levels of biomarker and vice versa. A positive correlation mean that MMSE scores increase with increasing levels of biomarker.

TABLE 5

	Correlation	Count	Z-value	P-value	95% Lower	95% Upper
GDNF to MMSE	-0.258	42	-1.646	0.0997	-0.521	0.05
SDF-1 to MMSE	-0.363	42	-2.375	0.0175	-0.601	-0.066
IGFBP-3 to MMSE	0.293	42	1.886	0.0593	-0.012	0.548
FGF-6 to MMSE	0.471	42	3.192	0.0014	0.195	0.687
TGF-b3 to MMSE	0.317	42	2.049	0.0405	0.014	0.566
BMP-4 to MMSE	0.294	42	1.845	0.0583	-0.011	0.545
NT-3 to MMSE	0.327	42	2.118	0.0342	0.025	0.574
EGF to MMSE	0.409	42	2.711	0.0067	0.12	0.634
BDNF to MMSE	0.464	42	3.139	0.0017	0.187	0.673
IGFBP-2 to MMSE (Females)	0.498	24	2.5	0.0123	0.118	0.75

PDGF-BB increase significantly whereas BDNF and RANTES do not change significantly. It has been found that from Mild AD (MMSE score in the range of 20-25) to Moderate AD (MMSE score in the range of 10-20) the level of LEPTIN does not decline whereas the levels for RANTES, BDNF and PDGF-BB declines.

Example 8

[0207] In an attempt to identify proteins that are altered in the peripheral immune system in AD, expression levels of 120 cytokines, chemokines, and growth factors in plasma from 32 AD patients and 19 nondemented age-matched controls were measured using spotted antibody microarrays on filters. Statistical analysis identified 20 proteins as significantly different between AD and controls. Six of them including brain derived neurotrophic factor (BDNF) and NT-3, and PDGF-BB, EGF, FGF-6, bFGF, TGF-b3 have known neurotrophic activity and were significantly reduced in AD plasma. BDNF levels correlated with better cognitive function in the mini mental state exam (MMSE). BDNF measurements in plasma from two hundred AD cases and controls using commercial sandwich ELISA showed a highly significant 25% reduction in AD cases. Consistent with the array data, reduced plasma BDNF levels were associated with impaired memory function. BDNF is critical for neuronal maintenance, survival, and function. Without being bound by theory decreased blood levels of neurotrophins and BDNF may be linked with neurodegeneration and cognitive dysfunction in AD.

Example 10

[0209] This example shows Table 6, a Summary of Quantitative Markers for Identification and Stratification of AD.

TABLE 6

References	Samples	Plasma BioMarker	% Difference in Samples	p-value
Normal	Questionable AD	BDNF	-46%	0.0049
Normal	Questionable AD	Leptin	-52%	0.0195
Normal	Questionable AD	RANTES	-31%	0.0203
Normal	Questionable AD	PDGF-BB	-30%	0.0799
Normal	Mild AD	BDNF	-29%	0.0092
Normal	Mild AD	Leptin	-29%	0.0359
Normal	Mild AD	RANTES	-16%	0.0780
Normal	Moderate AD	BDNF	-42%	0.0008
Normal	Moderate AD	Leptin	-33%	0.0359
Normal	Moderate AD	RANTES	-35%	0.0004
Normal	Severe AD	BDNF	-90%	0.0781
Normal	Severe AD	RANTES	-93%	0.0185
Normal	Severe AD	PDGF-BB	-89%	0.0931
Questionable AD	Mild AD	Leptin	45%	0.0617
Questionable AD	Mild AD	PDGF-BB	46%	0.0742
Mild AD	Moderate AD	RANTES	-23%	0.0780
Mild AD	Severe AD	BDNF	-87%	0.0898
Mild AD	Severe AD	RANTES	-92%	0.0470
Mild AD	Severe AD	PDGF-BB	-88%	0.0871
Questionable AD	MCI	BDNF	91%	0.0813
Questionable AD	MCI	Leptin	98%	0.0550
MCI	Mild AD	BDNF	-42%	0.0038

[0210] Accordingly, the present invention provides methods of aiding diagnosis of Alzheimer's disease ("AD"), comprising comparing a measured level of at least 4 AD diagnosis biomarkers, wherein said biomarkers comprise BDNF, PDGF-BB, Leptin and RANTES, in a biological fluid sample from an individual to a reference level for each AD diagnosis biomarker. Accordingly, methods are provided in which BDNF decreased at least about 10%, about 15%, about 20%, about 25% or about 30% as compared to a reference level of BDNF, indicates cognitive impairment, such as for example, an indication of AD. Accordingly, methods are provided in which Leptin decreased at least

about 10%, about 15%, about 20%, about 25% or about 30% as compared to a reference level of Leptin, indicates cognitive impairment, such as for example, an indication of AD. Accordingly, methods are provided in which RANTES decreased at least about 5%, about 10%, or about 15% as compared to a reference level of RANTES, indicates cognitive impairment, such as for example, an indication of AD. Accordingly, methods are provided in which PDGF-BB decreased at least about 80%, about 85% or about 90% as compared to a reference level of PDGF-BB, indicates cognitive impairment, such as for example, an indication of severe AD.

TABLE 7

Protein	Alternate names	Class	Protein ID
alpha-1 acid glycoprotein		acute phase	
alpha-1 antitrypsin		acute phase	
Ceruloplasmin		acute phase	
Haptoglobin		acute phase	
Hemopexin		acute phase	
Hemoxygenase		acute phase	
plasminogen activator inhibitor-1	PAI-1	acute phase	
serum amyloid A	SAA	acute phase	
serum amyloid P	SAP	acute phase	
4-11313 ligand	4-1BBL/CD137L	apoptosis	P41273
BAFF	TALL-1	apoptosis	Q9Y275
soluble TRAIL receptor 3	TRAIL sR3/TNFR S10C	apoptosis	014755
soluble TRAIL receptor 4	TRAIL sR4/TNFR S10D	apoptosis	Q9UBN6
TNF-related death ligand 1a	TRDL-1a/APRIL	apoptosis	AF046888
TNFSF-14	LIGHT	apoptosis	043557
TRAIL	Apo2L	apoptosis	P50591
BCA-1	BLC	chemokine	043927
CCL-28	CCK-1	chemokine	
cutaneous T cell attracting chemokine	CTACK, CCL27	chemokine	Qgz1X0
ENA-78		chemokine	P42830
Eotaxin-1		chemokine	P51671
Eotaxin-2	MPIF-2	chemokine	000175
Eotaxin-3	CCL26	chemokine	Q9Y258
Fractalkine	neurotactin	chemokine	P78423
Granulocyte chemotactic protein 2	GCP-2	chemokine	P80162
GRO alpha	MGSA	chemokine	P09341
GRO beta	MIP-2alpha	chemokine	P19875
GRO gamma	MIP-2beta	chemokine	P19876
haemoinfiltrate CC chemokine 1	HCC-1	chemokine	Q16627
haemoinfiltrate CC chemokine 4	HCC-4/CCL16	chemokine	015476
I-309	TCA-3/CCL-1	chemokine	P22362
IFNgamma inducible protein-10	IP-10	chemokine	P02778
IFN-inducible T cell alpha chemokine	I-TAC/CXCL11	chemokine	AF030514
interleukin-8	IL-8/NAP-1	chemokine	P10145
leucocyte cell-derived chemotaxin-2	LECT2	chemokine	
Lungkine	CXCL-15/WECHE	chemokine	
Lymphotactin	Lptn/ATAC	chemokine	P47992
macrophage inflammatory protein 1alpha	CCL3	chemokine	MIP-1alpha/
macrophage inflammatory protein 1beta	MIP-1beta/ACT-2/CCL4	chemokine	pLD78/
macrophage inflammatory protein 1d	MIP-1d/CCL15/LKN-1	chemokine	P10147
macrophage inflammatory protein 1gamma	MIP-1gamma/CCL9/MIP-3alpha/CCL20/	chemokine	P13236
macrophage inflammatory protein 3alpha	LARC	chemokine	P78556
macrophage inflammatory protein 3beta	MIP-3beta/ELC/CCL19	chemokine	Q99731
macrophage-derived chemokine	MDC/STCP-1	chemokine	000626
monocyte chemoattractant protein-1	MCP-1/CCL2	chemokine	P13500
monocyte chemoattractant protein-2	MCP-2/CCL8	chemokine	P78388
monocyte chemoattractant protein-3	MCP-3/CCL7	chemokine	P80098
monocyte chemoattractant protein-4	MCP-4/CCL13	chemokine	Q99616
monocyte chemoattractant protein-5	MCP-5/CCL12	chemokine	
monokine induced by IFN gamma	MIG	chemokine	Q07325
mucosa-associated chemokine	MEC	chemokine	AF266504
Myeloid progenitor inhibitory factor	MPIF/CKbeta8/CCL23	chemokine	
platelet basic protein	PBP/CTAP-III/NAP-2	chemokine	P02775

TABLE 7-continued

Protein	Alternate names	Class	Protein ID
platelet factor 4	PF-4/CXCL4	chemokine	P02776
pulmonary activation regulated chemokine	PARC/CCL18/MIP-4	chemokine	
RANTES	CCL5	chemokine	P13501
secondary lymphoid tissue chemokine	SLC/6Ckine	chemokine	000585
stromal cell derived factor 1	SDF-1/CXCL12	chemokine	P48061
thymus activation regulated chemokine	TARC/CCL17	chemokine	Q92583
thymus expressed chemokine	TECK/CCL25	chemokine	015444
Clq		collectin	
mannose binding lectin	MBL	collectin	
surfactant protein A	SP-A	collectin	
surfactant protein D	SP-D	collectin	
C1 inhibitor		complement	
C3a		complement	
Cob binding protein	C4BP	complement	
C5a		complement	
complement C3	C3	complement	
complement C5	C5	complement	
complement C8	C8	complement	
complement C9	C9	complement	
decay accelerating factor	DAF	complement	
Factor H		complement	
membrane inhibitor of reactive lysis	MIRL/CD59	complement	
Properdin		complement	
soluble complement receptor 1	sCR1	complement	
soluble complement receptor 2	sCR2	complement	
cardiotrophin-1	CT-1	cytokine	Q16619
CD27		cytokine	P26842
CD27L	CD70	cytokine	P32970
CD30	Ki-1	cytokine	P28908
CD30L	TNFSF8	cytokine	P32971
CD40L	TRAP/CD154	cytokine	P29965
interferon alpha	IFNalpha	cytokine	P01562
interferon beta	IFNbeta	cytokine	P01574
interferon gamma	IFNgamma	cytokine	P01579
interferon omega	IFNomega	cytokine	P05000
interferon-sensitive gene 15	ISG-15	cytokine	P05161
Leptin	OB	cytokine	P41159
leukemia inhibitory factor	LIF/CNDF	cytokine	P15018
Lymphotoxin	LT/TNF beta	cytokine	P01374
macrophage colony stimulating factor	M-CSF/CSF-1	cytokine	P09603
macrophage stimulating protein-alpha	MSPalpha/HGF1	cytokine	P26927
macrophage stimulating protein-beta	MSPbeta/HGF1	cytokine	P26927
migration inhibition factor	MIF/GIF	cytokine	P14174
oncostatin M	OSM	cytokine	P13725
RANKL	TRANCE/TNFSF-11	cytokine	014788
soluble IL6 R complex	sIL6RC (gp130 + sIL6R)	cytokine	
soluble Fas ligand	sCD95L	cytokine	P48023
TNF type I receptor	TNF-RI p55	cytokine	P19438
TNF type II receptor	TNF-R p75	cytokine	P20333
TNFSF-18	GITRL/AITRL	cytokine	095852
tumor necrosis factor alpha	TNF-alpha/Apo3L/DR3-L/ TNFSF-12	cytokine	P01375
TWEAK		cytokine	043508
acidic fibroblast growth factor	aFGF	growth factor	P05230
activin beta A		growth factor	P08476
agouti related protein	AGRP	growth factor	AAB52240
Amphiregulin	AR/SDGF	growth factor	P15514
angiopoietin-like factor	ALF	growth factor	
basic fibroblast growth factor	bFGF	growth factor	P09038
Betacellulin		growth factor	P35070
bone morphogenic protein 2	BMP2	growth factor	P12643
bone morphogenic protein 4	BMP4	growth factor	
bone morphogenic protein 5	BMP5	growth factor	
bone morphogenic protein 6	BMP6	growth factor	
bone morphogenic protein 7	BMP7	growth factor	
cripto-1	CRGF	growth factor	
epidermal growth factor	EGF	growth factor	P01133
Erythropoietin	Epo	growth factor	
fibroblast growth factor 17	FGF-17	growth factor	
fibroblast growth factor 18	FGF-18	growth factor	
fibroblast growth factor 19	FGF-19	growth factor	
fibroblast growth factor 2	FGF-2	growth factor	
fibroblast growth factor 4	FGF-4	growth factor	

TABLE 7-continued

Protein	Alternate names	Class	Protein ID
fibroblast growth factor 6	FGF-6	growth factor	
fibroblast growth factor 7	FGF-7/KGF	growth factor	
fibroblast growth factor 8	FGF-8	growth factor	
fibroblast growth factor 9	FGF-9	growth factor	
Flt3 ligand	Flt L	growth factor	P49771
Follistatin	FSP	growth factor	
Granulocyte colony stimulating factor	G-CSF	growth factor	P09919
granulocyte/macrophage CSF	GM-CSF	growth factor	P04141
growth and differentiation factor 11	GDF-11	growth factor	
growth and differentiation factor 15	GDF-15	growth factor	
growth arrest specific gene 6	Gas-6	growth factor	
heparin-binding epidermal growth factor	HB-EGF	growth factor	Q99075.
hepatocyte growth factor	HGF/SF	growth factor	P14210
hepatopoietin A	HPTA/HRG alpha/ neuregulin	growth factor	
heregulin alpha	NDF/HRG beta/neuregulin/	growth factor	
heregulin beta	NDF	growth factor	
IGF binding protein-1	IGFBP-1	growth factor	
IGF binding protein-2	IGFBP-2	growth factor	
IGF binding protein-3	IGFBP-3	growth factor	
IGF binding protein-4	IGFBP-4	growth factor	
inhibin A		growth factor	
inhibin B		growth factor	
insulin-like growth factor IA	IGF-IA	growth factor	P01343
insulin-like growth factor IB	IGF-IB	growth factor	P05019
insulin-like growth factor II	IGF-II	growth factor	P01344
macrophage galatose-specific lectin 1	MAC-1	growth factor	
Neuritin		growth factor	
Neurturin		growth factor	
orexin A		growth factor	
Osteonectin	SPARC	growth factor	
Osteoprotegrin	TNFRSF11B	growth factor	
placenta growth factor	PGIF	growth factor	
platelet derived growth factor alpha	PDGF-A	growth factor	P04085
platelet derived growth factor beta	PDGF-B	growth factor	P01127
pregnancy zone protein		growth factor	
Prolactin	PRL	growth factor	P01236
sensory and motor neuron-derived factor	SMDF	growth factor	
soluble GM-CSF receptor	sGM-CSF R	growth factor	P15509
stem cell factor	SLF/SCF/kit ligand/MGF	growth factor	P21583
Thrombopoietin	TPO/c-MPL ligand	growth factor	P40225
thymic stromal lymphoprotein	TSLP	growth factor	
Thymopoietin	Tpo	growth factor	
transforming growth factor alpha	TGF-alpha	growth factor	P01135
transforming growth factor beta 1	TGF-beta1	growth factor	P01137
transforming growth factor beta 2	TGF-beta2	growth factor	P08112
transforming growth factor beta 3	TGF-beta3	growth factor	P10600
vascular endothelial growth factor	VEGF	growth factor	P15692
interleukin-1 receptor antagonist	IL1Ra	interleukin	P18510
interleukin-10	IL-10	interleukin	P22301
interleukin-11	IL-11	interleukin	P20809
interleukin-12p35	IL-12p35	interleukin	P29459
interleukin-12p40	IL-12p40	interleukin	P29460
interleukin-13	IL-13	interleukin	P35225
interleukin-14	IL-14	interleukin	L15344
interleukin-15	IL-15	interleukin	P40933
interleukin-16	IL-16	interleukin	Q14005
interleukin-17	IL-17	interleukin	Q16552
interleukin-18	IL-18	interleukin	Q14116
interleukin-1alpha	IL-1a1.pha	interleukin	P01583
interleukin-1beta	IL-1beta	interleukin	P01584
interleukin-2	IL-2	interleukin	P01585
interleukin-3	IL-3	interleukin	P08700
interleukin-4	IL-4	interleukin	P05112
interleukin-5	IL-5	interleukin	P05113
interleukin-6	IL-6	interleukin	P05231
interleukin-7	IL-7	interleukin	P13232
interleukin-9	IL-9	interleukin	P15248
soluble interleukin-1 receptor I	sIL1R/CD121a	interleukin	P14778
soluble interleukin-1 receptor II	sIL1R/CD121b	interleukin	P27930
soluble interleukin-2 receptor	IL-2R/CD25	interleukin	P01589
soluble interleukin-5 receptor	sIL-5R/CD125	interleukin	Q01344
soluble interleukin-6 receptor	sIL-6R/CD126	interleukin	P08887

TABLE 7-continued

Protein	Alternate names	Class	Protein ID
soluble interleukin-7 receptor	sIL-7R/CD127	interleukin	P16871
soluble interleukin-9 receptor	sIL-9R	interleukin	PQ01113
AD7C	NTP	neuronal	AF010144
alpha synuclein		neuronal	AAH13293
GAP-43		neuronal	
Neurofilament		neuronal	
Synaptogamin		neuronal	
Synaptophysin		neuronal	
tau P199		neuronal	
brain derived neurotrophic factor	BDNF	neurotrophin	P23560
ciliary neurotrophic factor	CNTF	neurotrophin	P26441
glial derived neurotrophic factor	GDNF	neurotrophin	P39905
nerve growth factor	NGF	neurotrophin	P01138
neurotrophin 3	NT-3	neurotrophin	P20783
neurotrophin 4	NT-4	neurotrophin	P34130
soluble CNTF receptor	sCNTFR	neurotrophin	P26992
alpha2-macroglobulin	alpha 2M	others	
Alzheimer associated protein	ALZAS	others	
amyloid beta protein	Abeta 1-x	others	
apolipoprotein A	apoA	others	
apolipoprotein B	apoB	others	
apolipoprotein D	apoD	others	
apolipoprotein E	apoE	others	
apolipoprotein J	apoD/clusterin	others	
C reactive protein	CRP	others	
clara cell protein	CC16	others	
glial fibrillary acidic protein	GFAP	others	
Melanotransferrin		others	
soluble transferrin receptor	TfR	others	
Thrombomodulin		others	
Thrombospondin	Tsp	others	
tissue transglutaminase		others	
Transferrin		others	
alpha 1-antichymotrypsin	ACT	protease	NP001076
Clr		protease	
Cls		protease	
complement C2	C2	protease	
Factor B		protease	
Factor D	adipsin	protease	
FactorI		protease	
Kallikrein		protease	
MBL-associated serine protease 1	MASP-1	protease	
MBL-associated serine protease 2	MASP-2	protease	
Neuroserpin		protease	AAH18043
secretory leukocyte protease inhibitor	SLPI	protease	
Angiogenin		vascular	P00747
Angiostatin		vascular	
Endostatin		vascular	
Endothelin		vascular	
soluble E selectin	s E selectin	vascular	
vascular endothelial growth inhibitor	VEGI	vascular	

[0211] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be apparent to those skilled in the art that certain changes and modifications may be practiced. Therefore, the descriptions and examples should not be construed as limiting the scope of the invention.

We claim:

1. A method of aiding diagnosis of Alzheimer's disease ("AD"), comprising comparing a measured level of at least one AD diagnosis biomarker in a biological fluid sample from an individual to a reference level for the biomarker, wherein the AD diagnosis biomarker is selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC;

Acrcp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RI; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R.

2. The method of claim 1 wherein said AD diagnosis biomarker is selected from the group consisting of basic fibroblast growth factor (bFGF); BB homodimeric platelet derived growth factor (PDGF-BB); brain derived neurotrophic factor (BDNF); epidermal growth factor (EGF); fibroblast growth factor 6 (FGF-6); interleukin-3 (IL-3); soluble interleukin-6 receptor (sIL-6R); leptin (also known as ob); macrophage inflammatory protein-1 delta (MIP-1δ); macrophage stimulating protein alpha chain (MSP-α); neurotrophin-3 (NT-3); neutrophil activating peptide-2 (NAP-

2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF- β 3), and tumor necrosis factor beta (TNF- β).

3. The method of claim 1, wherein said AD diagnosis marker is selected from the group consisting of BDNF, sIL-6R, IL-8, leptin, MIP-1 δ , PDGF-BB, and TIMP-1.

4. The method of claim 3, wherein said AD diagnosis marker is selected from the group consisting of sIL-6R, IL-8, and TIMP-1.

5. The method of claim 3, wherein said AD diagnosis marker is selected from the group consisting of BDNF, MIP-1 δ , and TIMP-1.

6. The method of claim 1 wherein said AD diagnosis marker is selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES.

7. The method of claim 1, comprising comparing the measuring level of at least two AD diagnosis biomarkers to a reference level for the biomarkers.

8. The method of claim 7, wherein one of said at least two AD diagnosis biomarkers is leptin.

9. The method of claim 7, wherein one of said at least two AD diagnosis biomarkers is BDNF.

10. The method of claim 7 wherein the AD diagnosis biomarkers are selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES.

11. The method of claim 1, comprising comparing the measured level of at least three AD diagnosis biomarkers to a reference level for the biomarkers.

12. The method of claim 1, comprising comparing the measured level of at least four AD diagnosis biomarkers to a reference level for the biomarkers.

13. The method of claim 12, comprising comparing the measured level of at least BDNF, PDGF-BB, leptin and RANTES to a reference level for the biomarkers.

14. The method of claim 1, wherein comparing the measured value to a reference value for each AD diagnosis biomarker measured comprises calculating the fold difference between the measured value and the reference value.

15. The method of claim 14, further comprising comparing the fold difference for each AD diagnosis biomarker measured with a minimum fold difference value.

16. A method of aiding diagnosis of Alzheimer's disease ("AD"), comprising:

comparing a measured level of at least 4 AD diagnosis biomarkers, wherein said biomarkers comprise BDNF, PDGF-BB, leptin and RANTES, in a biological fluid sample from an individual to a reference level for each AD diagnosis biomarker.

17. The method of claim 16 wherein BDNF is decreased at least about 20% as compared to a reference level of BDNF.

18. The method of claim 16 wherein leptin is decreased at least about 25% as compared to a reference level of Leptin.

19. The method of claim 16 wherein RANTES is decreased at least about 16% as compared to a reference level of RANTES.

20. The method of claim 16 wherein PDGF-BB is decreased at least about 85% as compared to a reference level of PDGF-BB in a normal individual is associated with severe AD.

21. The method of claim 1, wherein said biological fluid sample is a peripheral biological fluid sample.

22. The method of claim 16 or 21, wherein said peripheral biological fluid sample is blood, serum or plasma.

23. The method of claim 1 or 16, further comprising obtaining a measured level of said AD biomarker in said biological fluid sample.

24. A method for monitoring progression of Alzheimer's disease (AD) in an AD patient, comprising: comparing a measured level of at least one AD diagnosis biomarker in a biological fluid sample from an individual to a reference level for the biomarker, wherein the AD diagnosis biomarker is selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(Ob); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R.

25. The method of claim 24 wherein said AD diagnosis biomarker is selected from the group consisting of basic fibroblast growth factor (bFGF); BB homodimeric platelet derived growth factor (PDGF-BB); brain derived neurotrophic factor (BDNF); epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1 δ), macrophage stimulating protein alpha chain (MSP- α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF- β 3), and tumor necrosis factor beta (TNF- β).

26. The method of claim 24 wherein the AD diagnosis marker is selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES.

27. The method of claim 24 wherein the AD diagnosis marker is leptin.

28. The method of claim 27, further comprising measuring a level for leptin in said biological fluid sample, thereby producing said measured value for leptin.

29. The method of claim 24, wherein said reference value is a value obtained from a biological fluid sample from the same AD patient at an earlier point in time.

30. The method of claim 24 wherein the biological fluid sample is a peripheral biological fluid sample.

31. A method for stratifying Alzheimer's disease (AD) in an individual, comprising:

comparing measured values for brain derived neurotrophic factor (BDNF) and BB homodimeric platelet derived growth factor (PDGF-BB) levels in a biological fluid sample from said patient with reference values for BDNF and PDGF-BB.

32. The method of claim 31 wherein said biological fluid sample is a peripheral fluid sample, including blood, serum or plasma.

33. The method of claim 31, further comprising comparing measured values for leptin and Rantes levels with reference values for leptin and Rantes, wherein reference values for BDNF, PDGF-BB, leptin and Rantes are for samples from individuals with MMSE scores from 25 to 28,

wherein an increase in leptin and PDGF-BB levels and wherein levels of BDNF and RANTES stay substantially the same indicate mild AD as indicated by an MMSE score of 20-25.

34. The method of claim 31, further comprising comparing measured values for leptin and Rantes levels with reference values for leptin and Rantes, wherein reference values for BDNF, PDGF-BB, leptin and Rantes are for samples from individuals with MMSE scores from 20-25, wherein a decrease in Rantes, BDNF, and PDGF levels and wherein levels of Leptin stays substantially the same indicate moderate AD as indicated by an MMSE score of 10-20.

35. A method of identifying a candidate agent for treatment of Alzheimer's Disease, comprising: assaying a prospective candidate agent for activity in modulating an AD biomarker, said AD biomarker selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R.

36. The method of claim 35 wherein the AD biomarker is selected from the group consisting of basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF), fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), tumor necrosis factor beta (TNF-β).

37. The method of claim 35 comprising assaying a prospective candidate agent for activity in modulating a set of AD biomarkers, wherein said set comprises BDNF, PDGF-BB, leptin and RANTES.

38. The method of claim 35 wherein said assaying is performed in vivo.

39. A kit comprising:

at least one reagent specific for at least one AD diagnosis marker, said at least one AD diagnosis biomarker selected from the group consisting of GCSF; IFN-g; IGFBP-1; BMP-6; BMP-4; Eotaxin-2; IGFBP-2; TARC; RANTES; ANG; PARC; Acrp30; AgRP(ART); TIMP-1; TIMP-2; ICAM-1; TRAIL R3; uPAR; IGFBP-4; LEPTIN(OB); PDGF-BB; EGF; BDNF; NT-3; NAP-2; IL-1ra; MSP-a; SCF; TGF-b3; TNF-b; MIP-1d; IL-3; FGF-6; IL-6 R; sTNF RII; AXL; bFGF; FGF-4; CNTF; MCP-1; MIP-1b; TPO; VEGF-B; IL-8; FAS; EGF-R and instructions for carrying out the method of claim 1.

40. The kit of claim 39 wherein said AD diagnosis biomarker is selected from the group consisting of basic fibroblast growth factor (bFGF), BB homodimeric platelet derived growth factor (PDGF-BB), brain derived neurotrophic factor (BDNF), epidermal growth factor (EGF),

fibroblast growth factor 6 (FGF-6), interleukin-3 (IL-3), soluble interleukin-6 receptor (sIL-6R), leptin (also known as ob), macrophage inflammatory protein-1 delta (MIP-1δ), macrophage stimulating protein alpha chain (MSP-α), neurotrophin-3 (NT-3), neutrophil activating peptide-2 (NAP-2), RANTES, soluble tumor necrosis factor receptor-2 (sTNF RII), stem cell factor (SCF), thrombopoietin (TPO), tissue inhibitor of metalloproteases-1 (TIMP-1), tissue inhibitor of metalloproteases-2 (TIMP-2), transforming growth factor-beta 3 (TGF-β3), tumor necrosis factor beta (TNF-β).

41. The kit of claim 39 comprising at least one reagent specific for each of at least two AD diagnosis markers.

42. The kit of claim 39 comprising at least one reagent specific for each of at least three AD diagnosis markers.

43. The kit of claim 39 comprising at least one reagent specific for each of at least four AD diagnosis markers.

44. The kit of any one of claim 41, 42 or 43 wherein the AD diagnosis markers are selected from the group consisting of BDNF, PDGF-BB, leptin and RANTES.

45. The kit of claim 44 wherein the reagent specific for the AD diagnosis biomarker is an antibody, or fragment thereof, that is specific for said AD diagnosis biomarker.

46. The kit of claim 39 further comprising at least one reagent specific for a biomarker that measures sample characteristics.

47. A surface comprising attached thereto, at least one reagent specific for each AD diagnosis biomarker in a set of AD diagnosis biomarkers, wherein said set of AD diagnosis biomarkers comprises BDNF, PDGF-BB, leptin and RANTES.

48. A surface comprising attached thereto,

a. at least one reagent specific for each AD diagnosis biomarker in a set of AD diagnosis biomarkers, wherein said set of AD diagnosis biomarkers consists of BDNF, PDGF-BB, leptin and RANTES; and

b. at least one reagent specific for a biomarker that measures sample characteristics.

49. The surface of claim 47 or 48 wherein said reagent specific for said AD diagnosis biomarker is an antibody, or fragment thereof, that is specific for said AD diagnosis biomarker.

50. A combination comprising the surface of claim 48 or 49 and a peripheral biological fluid sample from an individual.

51. The combination of claim 50 wherein said individual is at least 60, 65, 70, 75, 80, or 85.

52. The method of claim 1 further comprising the step of obtaining a value for the comparison of the measured level to the reference level.

53. The method of claim 13 further comprising the step of obtaining a value for the comparison of the measured level to the reference level.

54. A computer readable format comprising the values obtained by the method of claim 52 or 53.

55. A set of reference values for AD diagnosis biomarkers comprising BDNF, PDGF-BB, Leptin and RANTES.

56. A set of reagents specific for AD diagnosis biomarkers, wherein said biomarkers comprise BDNF, PDGF-BB, Leptin and RANTES.

专利名称(译)	用于诊断，分层和监测体内阿尔茨海默病和其他神经障碍的方法和组合物		
公开(公告)号	US20050221348A1	公开(公告)日	2005-10-06
申请号	US10/993813	申请日	2004-11-19
[标]申请(专利权)人(译)	RAY的Sandip WYSS科赖安东		
申请(专利权)人(译)	RAY的Sandip WYSS - 科赖安东		
当前申请(专利权)人(译)	规则为基础的医学，INC. 董事会利兰·斯坦福，齐齐哈尔大学，信托		
[标]发明人	RAY SANDIP WYSS CORAY ANTON		
发明人	RAY, SANDIP WYSS-CORAY, ANTON		
IPC分类号	C12Q1/68 G01N33/53 G01N33/567 G01N33/68		
CPC分类号	G01N33/6896 G01N2333/4745 G01N2800/2821		
优先权	60/523796 2003-11-19 US 60/566783 2004-04-30 US 60/566782 2004-04-30 US		
其他公开文献	US7598049		
外部链接	Espacenet USPTO		

摘要(译)

发明人已经发现了一组蛋白质生物标志物（“AD生物标志物”），其可以在外周生物液体样品中测量，以帮助诊断神经变性疾病，特别是阿尔茨海默病和轻度认知障碍（MCI）。本发明还提供了通过测试预期试剂调节AD生物标志物水平的活性来鉴定用于治疗阿尔茨海默病的候选药剂的方法。






Figure 1C