



US007691384B2

(12) **United States Patent**  
**Frank et al.**

(10) **Patent No.:** **US 7,691,384 B2**  
(45) **Date of Patent:** **\*Apr. 6, 2010**

(54) **METHOD TO DETECT IGE**

(75) Inventors: **Glenn R. Frank**, Wellington, CO (US);  
**James P. Porter**, Fort Collins, CO (US);  
**Keith E. Rushlow**, Fort Collins, CO  
(US); **Donald L. Wassom**, Fort Collins,  
CO (US)

(73) Assignee: **Heska Corporation**, Loveland, CO (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **11/683,854**

(22) Filed: **Mar. 8, 2007**

(65) **Prior Publication Data**

US 2007/0218565 A1 Sep. 20, 2007

#### **Related U.S. Application Data**

(62) Division of application No. 11/305,447, filed on Dec.  
16, 2005, now Pat. No. 7,195,767, which is a division  
of application No. 10/763,400, filed on Jan. 23, 2004,  
now abandoned, which is a division of application No.  
09/944,277, filed on Aug. 30, 2001, now Pat. No.  
6,682,894, which is a division of application No.  
09/285,873, filed on Mar. 31, 1999, now Pat. No.  
6,309,832, which is a division of application No.  
08/756,387, filed on Nov. 26, 1996, now Pat. No.  
5,945,294.

(51) **Int. Cl.**

**A61K 39/00** (2006.01)

**A61K 39/385** (2006.01)

(52) **U.S. Cl.** ..... **424/185.1**; 424/193.1; 530/300;  
530/350; 530/402; 536/23.1; 536/23.5

(58) **Field of Classification Search** ..... 424/185.1,  
424/193.1; 530/300, 350, 402; 536/23.1,  
536/23.5

See application file for complete search history.

(56) **References Cited**

#### **U.S. PATENT DOCUMENTS**

4,208,479 A	6/1980	Zuk et al.
4,962,035 A	10/1990	Leder et al.
5,091,318 A	2/1992	Anawis et al.
5,770,396 A	6/1998	Kinet
5,945,294 A	8/1999	Frank et al.
6,060,326 A *	5/2000	Frank et al. .... 436/513

#### **FOREIGN PATENT DOCUMENTS**

JP	5113443	5/1993
JP	07-072150	3/1995
WO	WO 90/04640	5/1990
WO	WO 91/06570	5/1991
WO	WO 97/24617	7/1997

#### **OTHER PUBLICATIONS**

Kochan, et al., (1988) *Nucleic Acids Res.* 16(8), p. 3584.  
Küster, et al., (1990) *J. Biol. Chem.* 265(11), pp. 6448-6452.  
Küster, et al., (1992) *J. Biol. Chem.* 267(18), pp. 12782-12787.  
Lowenthal, et al., (1993) *Annals of Allergy* 71, pp. 481-484.  
Pang, et al., "Characterization of the Gene for the Human High  
Affinity IgE Receptor (FcεRI) α-Chain," (1993) *J. Immunol.*  
151(11), pp. 6166-6174.  
Shimizu, et al., (1988), *Proc. Natl. Acad. Sci. USA* (85), pp. 1907-  
1911.  
Patent Abstract of Japan, vol. 095, No. 007, Aug. 31, 1995 and JP 07  
092167 A (Kinki Univ.; Others: 01), Apr. 7, 1995.  
Patent Abstract of Japan, vol. 095, No. 006, Jul. 31, 1995 and JP 07  
072150 A (Tonen Corp.; Others: 01), Mar. 7, 1995.

\* cited by examiner

*Primary Examiner*—Rodney P. Swartz

(74) *Attorney, Agent, or Firm*—Sheridan Ross P.C.

(57) **ABSTRACT**

The present invention includes a method to detect IgE using a  
human Fc epsilon receptor (Fc<sub>ε</sub>R) to detect IgE antibodies in  
a biological sample from a cat, a dog, or a horse. The present  
invention also relates to kits to perform such methods.

**12 Claims, 11 Drawing Sheets**






Fig. 1

**SAMPLES****Fig. 2**




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7






Fig. 8




Fig. 9




Fig. 10




Fig. 11

## METHOD TO DETECT IGE

## CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional of U.S. patent application Ser. No. 11/305,447, filed Dec. 16, 2005, now issued as U.S. Pat. No. 7,195,767 B2; which is a divisional of U.S. patent application Ser. No. 10/763,400, filed Jan. 23, 2004, now abandoned; which is a divisional of U.S. patent application Ser. No. 09/944,277, filed Aug. 30, 2001, now issued as U.S. Pat. No. 6,682,894 B2; which is a divisional of U.S. application Ser. No. 09/285,873, filed Mar. 31, 1999, now U.S. Pat. No. 6,309,832 B1; which is a divisional of U.S. application Ser. No. 08/756,387, filed Nov. 26, 1996, now issued as U.S. Pat. No. 5,945,294; each of the foregoing entitled "METHOD TO DETECT IgE" and incorporated by reference herein.

## FIELD OF THE INVENTION

The present invention relates to a novel method to detect epsilon immunoglobulin (IgE). The present invention also includes novel kits to detect IgE as well as methods to produce the detection reagent.

## BACKGROUND OF THE INVENTION

Diagnosis of disease and determination of treatment efficacy are important tools in medicine. In particular, detection of IgE production in an animal can be indicative of disease. Such diseases include, for example, allergy, atopic disease, hyper IgE syndrome, internal parasite infections and B cell neoplasia. In addition, detection of IgE production in an animal following a treatment is indicative of the efficacy of the treatment, such as when using treatments intended to disrupt IgE production.

Until the discovery of the present invention, detection of IgE in samples obtained from non-human animals has been hindered by the absence of suitable reagents for detection of IgE. Various compounds have been used to detect IgE in IgE-containing compositions. In particular, antibodies that bind selectively to epsilon idiotype antibodies (i.e., anti-IgE antibodies) have been used to detect IgE. These anti-IgE antibodies, however, can cross-react with other antibody idiotypes, such as gamma isotype antibodies. The discovery of the present invention includes the use of a Fe epsilon receptor (Fc<sub>ε</sub>R) molecule to detect the presence of IgE in a putative IgE-containing composition. A Fc<sub>ε</sub>R molecule provides an advantage over, for example anti-IgE antibodies, to detect IgE because a Fc<sub>ε</sub>R molecule can bind to an IgE with more specificity (i.e., less idiotype cross-reactivity) and more sensitivity (i.e., affinity) than anti-IgE binding antibodies.

Lowenthal et al., 1993, *Annals of Allergy* 71:481-484, teach that dog serum can transfer cutaneous reactivity to a human. While it is possible that Lowenthal et al. properly teach the binding of human Fc<sub>ε</sub>R to canine IgE. Lowenthal et al., however, do not provide data defining the particular cellular proteins responsible for the transfer of cutaneous reactivity. As such, a skilled artisan would conclude that the transfer of cutaneous reactivity taught by Lowenthal et al. could be due to a variety of different molecular interactions and that the conclusion drawn by Lowenthal et al. is merely an interpretation. In addition, Lowenthal et al. do not teach the use of purified human Fc<sub>ε</sub>R to detect canine IgE. The subunits of human Fc<sub>ε</sub>R have been known as early as 1988 and have never been used to detect canine, feline or equine IgE. Indeed, U.S. Pat. No. 4,962,035, to Leder et al., issued Oct. 9, 1990,

discloses human Fc<sub>ε</sub>R but does not disclose the use of such a human Fc<sub>ε</sub>R to detect human or non-human IgE. The use of purified human Fc<sub>ε</sub>R avoids complications presented by use of Fc<sub>ε</sub>R bound to a cell, such as non-specific binding of the Fc<sub>ε</sub>R-bearing cell due to additional molecules present on the cell membrane. That purified human Fc<sub>ε</sub>R detects non-human IgE is unexpected because inter-species binding between a Fc<sub>ε</sub>R and an IgE is not predictable. For example, human Fc<sub>ε</sub>R binds to rat IgE but rat Fc<sub>ε</sub>R does not bind to human IgE.

The high affinity Fc<sub>ε</sub>R consists of three protein chains, alpha, beta and gamma. Prior investigators have disclosed the nucleic acid sequence for: the alpha chain (Kochan et al., *Nucleic Acids Res.* 16:3584, 1988; Shimizu et al., *Proc. Natl. Acad. Sci. USA* 85:1907-1911, 1988; and Pang et al., *J. Immunol.* 151:6166-6174, 1993); the beta chain (Kuster et al., *J. Biol. Chem.* 267: 12782-12787, 1992); and the gamma chain (Kuster et al., *J. Biol. Chem.* 265-6448-6452, 1990).

Thus, methods and kits are needed in the art that will provide specific detection of non-human IgE.

## SUMMARY OF THE INVENTION

The present invention includes detection methods and kits that detect IgE. One embodiment of the present invention is a method to detect IgE comprising: (a) contacting an isolated human Fc<sub>ε</sub>receptor (Fc<sub>ε</sub>R) molecule with a putative IgE-containing composition under conditions suitable for formation of a Fc<sub>ε</sub>R molecule:IgE complex, wherein the IgE is selected from the group consisting of canine IgE, feline IgE and equine IgE; and (b) determining the presence of IgE by detecting the Fc<sub>ε</sub>R molecule:IgE complex, the presence of the Fc<sub>ε</sub>R molecule:IgE complex indicating the presence of IgE. A preferred Fc<sub>ε</sub>R molecule is one in which a carbohydrate group of the Fc<sub>ε</sub>R molecule is conjugated to biotin.

Another embodiment of the present invention is a method to detect IgE comprising: (a) contacting a recombinant cell with a putative IgE-containing composition under conditions suitable for formation of a recombinant cell:IgE complex, in which the recombinant cell includes: a recombinant cell expressing a human Fc<sub>ε</sub>R molecule; and a recombinant cell expressing an antibody that binds selectively to an IgE including canine IgE, feline IgE and equine IgE; and (b) determining the presence of IgE by detecting the recombinant cell:IgE complex, the presence of the recombinant cell:IgE complex indicating the presence of IgE. A preferred recombinant cell includes a RBL-hFc<sub>ε</sub>R cell.

Another embodiment of the present invention is a method to detect flea allergy dermatitis comprising, (a) immobilizing a flea allergen on a substrate; (b) contacting the flea allergen with a putative IgE-containing composition under conditions suitable for formation of an antigen:IgE complex bound to said substrate; (c) removing non-bound material from the substrate under conditions that retain antigen:IgE complex binding to the substrate; and (c) detecting the presence of the antigen:IgE complex by contacting the antigen:IgE complex with a Fc<sub>ε</sub>R molecule. Preferably, the flea allergen is a flea saliva antigen and more preferably flea saliva products and/or flea saliva proteins.

The present invention also includes a kit for performing methods of the present invention. One embodiment is a kit for detecting IgE comprising a human Fc<sub>ε</sub> receptor (Fc<sub>ε</sub>R) molecule and a means for detecting an IgE including canine IgE, feline IgE and equine IgE. Another embodiment is a general allergen kit comprising an allergen common to all regions of the United States and a human Fc<sub>ε</sub> receptor (Fc<sub>ε</sub>R) molecule.

Another embodiment is a kit for detecting flea allergy dermatitis comprising a human Fc<sub>ε</sub> receptor (Fc<sub>ε</sub>R) molecule and a flea allergen.

Another embodiment of the present invention is an isolated human Fc<sub>ε</sub> receptor (Fc<sub>ε</sub>R) alpha chain protein, in which a carbohydrate group of the Fc<sub>ε</sub>R alpha chain protein is conjugated to biotin. A preferred Fc<sub>ε</sub>R alpha chain protein comprises PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect canine IgE antibodies.

FIG. 2 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect plant allergen-specific canine IgE antibodies.

FIG. 3 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect human or canine IgE antibodies.

FIG. 4 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect flea allergen-specific canine IgE antibodies.

FIG. 5 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect flea allergen-specific and heartworm antigen-specific canine IgE antibodies.

FIG. 6 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect flea saliva-specific canine IgE antibodies.

FIG. 7 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect heartworm antigen-specific feline IgE antibodies.

FIG. 8 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect heartworm antigen-specific feline IgE antibodies.

FIG. 9 depicts ELISA results using biotinylated alpha chain of human Fc<sub>ε</sub>R to detect antigen-specific equine IgE antibodies.

FIG. 10 depicts ELISA results using basophilic leukemia cells expressing alpha chain of human Fc<sub>ε</sub>R to detect canine IgE antibodies in sera from heartworm-infected dogs (diamonds) and heat inactivated sera from heartworm-infected dogs (squares).

FIG. 11 depicts ELISA results using basophilic leukemia cells expressing alpha chain of human Fc<sub>ε</sub>R to detect canine IgE antibodies in sera from flea saliva sensitized dogs (diamonds) and heat inactivated sera from flea saliva sensitized dogs (squares).

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the discovery that purified high affinity human Fc epsilon receptor (referred to herein as Fc<sub>ε</sub>R) can be used in certain non-human (i.e., canine, feline or equine) epsilon immunoglobulin (referred to herein as IgE or IgE antibody)-based detection (e.g., diagnostic, screening) methods and kits. The use of human Fc<sub>ε</sub>R to detect non-human IgE is unexpected because canine, feline and equine immune systems are different from the human immune system, as well as from each other (i.e., molecules important to the immune system usually are species specific).

One embodiment of the present invention is a method to detect a non-human IgE using an isolated human Fc<sub>ε</sub>R molecule. It is to be noted that the term "a" entity or "an" entity refers to one or more of that entity; for example, a protein refers to one or more proteins or at least one protein. As such, the terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the

terms "comprising", "including", and "having" can be used interchangeably. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. Furthermore, a compound "selected from the group consisting of" refers to one or more of the compounds in the list that follows, including mixtures (i.e., combinations) of two or more of the compounds.

According to the present invention, an isolated, or biologically pure, Fc<sub>ε</sub>R molecule, is a molecule that has been removed from its natural milieu. As such, "isolated" and "biologically pure" do not necessarily reflect the extent to which the molecule has been purified. An isolated human Fc<sub>ε</sub>R molecule of the present invention can be obtained from its natural source (e.g., from a human mast cell), can be produced using recombinant DNA technology or can be produced by chemical synthesis.

A Fc<sub>ε</sub>R molecule (also referred to herein as Fc<sub>ε</sub>R or Fc<sub>ε</sub>R protein) of the present invention can be a full-length protein, a portion of a full-length protein or any homolog of such a protein. As used herein, a protein can be a polypeptide or a peptide. A Fc<sub>ε</sub>R molecule of the present invention can comprise a complete Fc<sub>ε</sub>R (i.e., alpha, beta and gamma Fc<sub>ε</sub>R chains), an alpha Fc<sub>ε</sub>R chain (also referred to herein as Fc<sub>ε</sub>R α chain) or portions thereof. Preferably, a Fc<sub>ε</sub>R molecule comprises at least a portion of a Fc<sub>ε</sub>R α chain that binds to IgE, i.e., that is capable of forming an immunocomplex with an IgE constant region. Preferably, a Fc<sub>ε</sub>R molecule of the present invention binds to IgE with an affinity of about K<sub>d</sub>≈10<sup>8</sup>, more preferably with an affinity of about K<sub>d</sub>≈10<sup>9</sup> and even more preferably with an affinity of about K<sub>d</sub>≈10<sup>10</sup>.

An isolated Fc<sub>ε</sub>R molecule of the present invention, including a homolog, can be identified in a straight-forward manner by the Fc<sub>ε</sub>R molecule's ability to form an immunocomplex with an IgE. Examples of Fc<sub>ε</sub>R homologs include Fc<sub>ε</sub>R proteins in which amino acids have been deleted (e.g., a truncated version of the protein, such as a peptide), inserted, inverted, substituted and/or derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation and/or addition of glycerophosphatidyl inositol) such that the homolog includes at least one epitope capable of forming an immunocomplex with an IgE.

Fc<sub>ε</sub>R homologs can be the result of natural allelic variation or natural mutation. Fc<sub>ε</sub>R homologs of the present invention can also be produced using techniques known in the art including, but not limited to, direct modifications to the protein or modifications to the gene encoding the protein using, for example, classic or recombinant DNA techniques to effect random or targeted mutagenesis.

According to the present invention, a human Fc<sub>ε</sub>R α chain of the present invention is encoded by at least a portion of the nucleic acid sequence of the coding strand of a cDNA encoding a full-length Fc<sub>ε</sub>R α chain protein represented herein as SEQ ID NO:1, the portion at least encoding the IgE binding site of the Fc<sub>ε</sub>R α chain protein. The double-stranded nucleic acid molecule including both the coding strand having SEQ ID NO:1 and the complementary non-coding strand (the nucleic acid sequence of which can be readily determined by one skilled in the art and is shown herein as SEQ ID NO:3) is referred to herein as Fc<sub>ε</sub>R nucleic acid molecule nhFc<sub>ε</sub>Rα<sub>1198</sub>. Translation of SEQ ID NO:1 suggests that nucleic acid molecule nhFc<sub>ε</sub>Rα<sub>1198</sub> encodes a full-length Fc<sub>ε</sub>R α chain protein of about 257 amino acids, referred to herein as PhFc<sub>ε</sub>Rα<sub>257</sub>, represented by SEQ ID NO:2, assuming an open reading frame having an initiation (start) codon spanning from about nucleotide 107 through about nucleotide 109 of SEQ ID NO:1 and a termination (stop) codon spanning

from about nucleotide 878 through about nucleotide 880 of SEQ ID NO:1. The coding region encoding PhFc<sub>ε</sub>Rα<sub>257</sub>, including the stop codon, is represented by nucleic acid molecule nhFc<sub>ε</sub>Rα<sub>774</sub>, having a coding strand with the nucleic acid sequence represented herein as SEQ ID NO:4. SEQ ID NO:1 encodes a signal peptide of about 25 amino acids as well as a mature protein of about 232 amino acids, denoted herein as PhFc<sub>ε</sub>Rα<sub>232</sub>, the amino acid sequence of which is represented herein as SEQ ID NO:6. The nucleic acid molecule encoding the apparent mature protein is referred to as nhFc<sub>ε</sub>Rα<sub>699</sub>, the nucleic acid sequence of the coding strand of which is denoted herein as SEQ ID NO:7. SEQ ID NO:1 also encodes a hydrophobic transmembrane domain and a cytoplasmic tail which as a group extend from about amino acid 205 to about amino acid 257 of SEQ ID NO:2. Knowledge of these nucleic acid and amino acid sequences allows one skilled in the art to make modifications to the respective nucleic acid molecules and proteins to, for example, develop a Fc<sub>ε</sub>R α chain protein with increased solubility and/or a truncated protein (e.g., a peptide) capable of detecting IgE, e.g., PhFc<sub>ε</sub>Rα<sub>197</sub> and PhFc<sub>ε</sub>Rα<sub>172</sub>. Preferred Fc<sub>ε</sub>R molecules include PhFc<sub>ε</sub>Rα<sub>257</sub>, PhFc<sub>ε</sub>Rα<sub>197</sub>, PhFc<sub>ε</sub>Rα<sub>232</sub> and PhFc<sub>ε</sub>Rα<sub>172</sub>. Preferred nucleic acid molecules to encode a Fc<sub>ε</sub>R molecules include nhFc<sub>ε</sub>Rα<sub>774</sub>, nhFc<sub>ε</sub>Rα<sub>1198</sub>, nhFc<sub>ε</sub>Rα<sub>612</sub>, nhFc<sub>ε</sub>Rα<sub>591</sub>, nhFc<sub>ε</sub>Rα<sub>699</sub> and/or nhFc<sub>ε</sub>Rα<sub>516</sub>.

Isolated Fc<sub>ε</sub>R molecule protein of the present invention can be produced by culturing a cell capable of expressing the protein under conditions effective to produce the protein, and recovering the protein. A preferred cell to culture is a recombinant cell that is capable of expressing the protein, the recombinant cell being produced by transforming a host cell with one or more nucleic acid molecules of the present invention. Transformation of a nucleic acid molecule into a cell can be accomplished by any method by which a nucleic acid molecule can be inserted into the cell. Transformation techniques include, but are not limited to, transfection, electroporation, microinjection, lipofection, adsorption, and protoplast fusion. A recombinant cell may remain unicellular or may grow into a tissue, organ or a multicellular organism. Transformed nucleic acid molecules of the present invention can remain extrachromosomal or can integrate into one or more sites within a chromosome of the transformed (i.e., recombinant) cell in such a manner that their ability to be expressed is retained. Suitable and preferred nucleic acid molecules with which to transform a cell are as disclosed herein for suitable and preferred Fc<sub>ε</sub>R nucleic acid molecules per se. Particularly preferred nucleic acid molecules to include in recombinant cells of the present invention include nhFc<sub>ε</sub>Rα<sub>774</sub>, nhFc<sub>ε</sub>Rα<sub>1198</sub>, nhFc<sub>ε</sub>Rα<sub>612</sub>, nhFc<sub>ε</sub>Rα<sub>591</sub>, nhFc<sub>ε</sub>Rα<sub>699</sub> and/or nhFc<sub>ε</sub>Rα<sub>516</sub>.

Suitable host cells to transform include any cell that can be transformed with a nucleic acid molecule of the present invention. Host cells can be either untransformed cells or cells that are already transformed with at least one nucleic acid molecule. Host cells of the present invention either can be endogenously (i.e., naturally) capable of producing a Fc<sub>ε</sub>R molecule protein of the present invention or can be capable of producing such proteins after being transformed with at least one nucleic acid molecule of the present invention. Host cells of the present invention can be any cell capable of producing at least one protein of the present invention, and include bacterial, fungal (including yeast), parasite (including protozoa and ectoparasite), insect, other animal and plant cells.

Preferably, a recombinant cell is transfected with a recombinant molecule of the present invention that can include at least one of any nucleic acid molecule heretofore described operatively linked to at least one of any transcription control

sequence capable of effectively regulating expression of the nucleic acid molecule(s) in the cell to be transformed, examples of which are disclosed herein. A particularly preferred recombinant molecule includes pVL-nhFc<sub>ε</sub>Rα<sub>612</sub>. Details regarding the production of Fc<sub>ε</sub>R molecule nucleic acid molecule-containing recombinant molecules are disclosed herein. Particularly preferred recombinant cell of the present invention includes *Trichoplusia ni*-pVL-nhFc<sub>ε</sub>Rα<sub>612</sub>.

A Fc<sub>ε</sub>R molecule of the present invention can include chimeric molecules comprising a portion of a Fc<sub>ε</sub>R molecule that binds to an IgE and a second molecule that enables the chimeric molecule to be bound to a substrate in such a manner that the Fc<sub>ε</sub>R portion binds to IgE in essentially the same manner as a Fc<sub>ε</sub>R molecule that is not bound to a substrate. An example of a suitable second molecule includes a portion of an immunoglobulin molecule.

A Fc<sub>ε</sub>R molecule of the present invention can be contained in a formulation, herein referred to as a Fc<sub>ε</sub>R formulation. For example, a Fc<sub>ε</sub>R can be combined with a buffer in which the Fc<sub>ε</sub>R is solubilized, and/or a carrier. Suitable buffers and carriers are known to those skilled in the art. Examples of suitable buffers include any buffer in which a Fc<sub>ε</sub>R can function to selectively bind to IgE, such as, but not limited to, phosphate buffered saline, water, saline, phosphate buffer, bicarbonate buffer, HEPES buffer (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid buffered saline), TES buffer (Tris-EDTA buffered saline), Tris buffer and TAE buffer (Tris-acetate-EDTA). Examples of carriers include, but are not limited to, polymeric matrices, toxoids, and serum albumins, such as bovine serum albumin. Carriers can be in mixed with Fc<sub>ε</sub>R or conjugated (i.e., attached) to Fc<sub>ε</sub>R in such a manner as to not substantially interfere with the ability of the Fc<sub>ε</sub>R to selectively bind to IgE.

A Fc<sub>ε</sub>R of the present invention can be bound to the surface of a cell expressing the Fc<sub>ε</sub>R. A preferred Fc<sub>ε</sub>R-bearing cell includes a recombinant cell expressing a nucleic acid molecule encoding a human Fc<sub>ε</sub>R alpha chain of the present invention. A more preferred recombinant cell of the present invention expresses a nucleic acid molecule that encodes at least one of the following proteins: PhFc<sub>ε</sub>Rα<sub>257</sub> and PhFc<sub>ε</sub>Rα<sub>232</sub>. An even more preferred recombinant cell expresses a nucleic acid molecule including nhFc<sub>ε</sub>Rα<sub>612</sub>, nhFc<sub>ε</sub>Rα<sub>591</sub>, nhFc<sub>ε</sub>Rα<sub>699</sub> and/or nhFc<sub>ε</sub>Rα<sub>516</sub> with a recombinant cell expressing a nucleic acid molecule comprising a nucleic acid sequence including SEQ ID NO:1 or SEQ ID NO:4, or a nucleic acid molecule comprising an allelic variant of a nucleic acid molecule comprising SEQ ID NO:1 or SEQ ID NO:4, being even more preferred. An even more preferred recombinant cell is a RBL-hFc<sub>ε</sub>R cell.

In addition, a Fc<sub>ε</sub>R formulation of the present invention can include not only a Fc<sub>ε</sub>R but also one or more additional antigens or antibodies useful in detecting IgE. As used herein, an antigen refers to any molecule capable of being selectively bound by an antibody. As used herein, specific binding of a first molecule to a second molecule refers to the ability of the first molecule to preferentially bind (e.g., having higher affinity higher avidity) to the second molecule when compared to the ability of a first molecule to bind to a third molecule. The first molecule need not necessarily be the natural ligand of the second molecule. Examples of such antibodies include, but are not limited to, antibodies that bind selectively to the constant region of an IgE heavy (i.e., anti-IgE isotype antibody) or antibodies that bind selectively to an IgE having a specific antigen specificity (i.e., anti-IgE idiotype antibody). Examples of such antigens include any antigen known to induce the production of IgE. Preferred antigens include allergens and parasite antigens. Allergens of the

present invention are preferably derived from fungi, trees, weeds, shrubs, grasses, wheat, corn, soybeans, rice, eggs, milk, cheese, bovines (or cattle), poultry, swine, sheep, yeast, fleas, flies, mosquitos, mites, midges, biting gnats, lice, bees, wasps, ants, true bugs or ticks. A suitable flea allergen includes an allergen derived from a flea, in particular flea saliva antigen. A preferred flea allergen includes a flea saliva antigen. Preferred flea saliva antigens include antigens such as those disclosed in PCT Patent Publication No. WO 96/11271, published Apr. 18, 1996, by Frank et al. (this publication is incorporated by reference herein in its entirety), with flea saliva products and flea saliva proteins being particularly preferred. According to the present invention, a flea saliva protein includes a protein produced by recombinant DNA methods, as well as proteins isolated by other methods disclosed in PCT Patent Publication No. WO 96/11271.

Preferred general allergens include those derived from grass, Meadow Fescue, Curly Dock, plantain, Mexican Firebush, Lamb's Quarters, pigweed, ragweed, sage, elm, cocklebur, Box Elder, walnut, cottonwood, ash, birch, cedar, oak, mulberry, cockroach, *Dermatophagoides*, *Alternaria*, *Aspergillus*, *Cladosporium*, *Fusarium*, *Helminthosporium*, *Mucor*, *Penicillium*, *Pulularia*, *Rhizopus* and/or *Tricophyton*. More preferred general allergens include those derived from Johnson Grass, Kentucky Blue Grass, Meadow Fescue, Orchard Grass, Perennial Rye Grass, Redtop Grass, Timothy Grass, Bermuda Grass, Brome Grass, Curly Dock, English Plantain, Mexican Firebush, Lamb's Quarters, Rough Pigweed Short Ragweed, Wormwood Sage, American Elm, Common Cocklebur, Box Elder, Black Walnut, Eastern Cottonwood, Green Ash, River Birch, Red Cedar, Red Oak, Red Mulberry, Cockroach, *Dermatophagoides farinae*, *Alternaria alternata*, *Aspergillus ummigatus*, *Cladosporium herbarum*, *Fusarium vasinfectum*, *Helminthosporium sativum*, *Mucor recemosus*, *Penicillium notatum*, *Pullularia pullulans*, *Rhizopus nigricans* and/or *Tricophyton* spp. Preferred parasite antigens include, but are not limited to, helminth antigens, in particular heartworm antigens, such as Di33 (described in U.S. patent application Ser. No. 08/715,628, filed Sep. 18, 1996, to Grieve et al.). The term "derived from" refers to a natural allergen of such plants or organisms (i.e., an allergen directly isolated from such plants or organisms), as well as, non-natural allergens of such plants or organisms that possess at least one epitope capable of eliciting an immune response against an allergen (e.g., produced using recombinant DNA technology or by chemical synthesis).

The present invention also includes human Fc<sub>ε</sub>R mimetopes and use thereof to detect IgE. In accordance with the present invention, a "mimotope" refers to any compound that is able to mimic the ability of a Fc<sub>ε</sub>R molecule to bind to IgE. A mimotope can be a peptide that has been modified to decrease its susceptibility to degradation but that still retains IgE-binding activity. Other examples of mimetopes include, but are not limited to, carbohydrate-based compounds, lipid-based compounds, nucleic acid-based compounds, natural organic compounds, synthetically derived organic compounds, anti-idiotypic antibodies and/or catalytic antibodies, or fragments thereof. A mimotope can be obtained by, for example, screening libraries of synthetic compounds for compounds capable of binding to IgE. A mimotope can also be obtained by, for example, rational drug design. In a rational drug design procedure, the three-dimensional structure of a compound of the present invention can be analyzed by, for example, nuclear magnetic resonance (NMR) or x-ray crystallography. The three-dimensional structure can then be used to predict structures of potential mimetopes by, for example, computer modeling. The predicted mimotope structures can

then be produced by, for example, chemical synthesis, recombinant DNA technology, or by isolating a mimotope from a natural source. Specific examples of Fc<sub>ε</sub>R mimetopes include anti-idiotypic antibodies, oligonucleotides produced using Selex technology, peptides identified by random screening of peptide libraries and proteins identified by phage display technology.

One embodiment of the present invention is a method to detect non-human IgE which includes the steps of: (a) contacting an isolated human Fc<sub>ε</sub>R receptor (Fc<sub>ε</sub>R) molecule with a putative IgE-containing composition under conditions suitable for formation of an Fc<sub>ε</sub>R molecule:IgE complex; and (b) detecting levels of IgE by detecting said Fc<sub>ε</sub>R molecule:IgE complex. Presence of such a Fc<sub>ε</sub>R molecule:IgE complex indicates that the animal is producing IgE. Preferred non-human IgE to detect using a human Fc<sub>ε</sub>R molecule include canine IgE, feline IgE and equine IgE. The present method can further include the step of determining whether an IgE complexed with a Fc<sub>ε</sub>R molecule is heat labile. Methods to determine heat lability of IgE are disclosed in the Examples section. Preferably, an IgE is heat labile when incubated at about 56° C. for about 4 hours. Without being bound by theory, Applicants believe that heat labile forms of IgE bind to certain allergens and non-heat labile forms of IgE bind to other types of allergens. As such, detection of heat labile IgE compared with non-heat labile IgE can be used to discriminate between allergen sensitivities. For example, Applicants believe that IgE antibodies that bind to certain flea allergens and heartworm allergens are heat labile while IgE antibodies that bind to certain plant allergens are not heat labile. Thus, the presence of non-heat labile IgE can indicate that an animal is sensitive to certain plant allergens but not to certain flea or heartworm allergens. Moreover, Applicants believe that identification of heat labile IgE and non-heat labile IgE using a human Fc<sub>ε</sub>R suggests the presence of different sub-populations of IgE that may or may not have substantially similar structures to known IgE. As such, a Fc<sub>ε</sub>R molecule of the present invention may be useful for detecting molecules bound by the Fc<sub>ε</sub>R molecule but not identical to a known IgE.

As used herein, canine refers to any member of the dog family, including domestic dogs, wild dogs and zoo dogs. Examples of dogs include, but are not limited to, domestic dogs, wild dogs, foxes, wolves, jackals and coyotes. As used herein, a feline refers to any member of the cat family, including domestic cats, wild cats and zoo cats. Examples of cats include, but are not limited to, domestic cats, lions, tigers, leopards, panthers, cougars, bobcats, lynx, jaguars, cheetahs, and servals. As used herein, equine refers to any member of the horse family, including horses, donkeys, mules and zebras.

As used herein, the term "contacting" refers to combining or mixing, in this case a putative IgE-containing composition with a human Fc<sub>ε</sub>R molecule. Formation of a complex between a Fc<sub>ε</sub>R and an IgE refers to the ability of the Fc<sub>ε</sub>R to selectively bind to the IgE in order to form a stable complex that can be measured (i.e., detected). As used herein, the term selectively binds to an IgE refers to the ability of a Fc<sub>ε</sub>R of the present invention to preferentially bind to IgE, without being able to substantially bind to other antibody isotypes. Binding between a Fc<sub>ε</sub>R and an IgE is effected under conditions suitable to form a complex; such conditions (e.g., appropriate concentrations, buffers, temperatures, reaction times) as well as methods to optimize such conditions are known to those skilled in the art, and examples are disclosed herein. Examples of complex formation conditions are also disclosed in, for example, in Sambrook et al., *Molecular Cloning: A*



*Laboratory Manual*, Cold Spring Harbor Labs Press, 1989, the reference Sambrook et al., *ibid.*, is incorporated by reference herein in its entirety.

As used herein, the term "detecting complex formation" refers to determining if any complex is formed, i.e., assaying for the presence (i.e., existence) of a complex. If complexes are formed, the amount of complexes formed can, but need not be, determined. Complex formation, or selective binding, between Fc<sub>ε</sub>R and any IgE in the composition can be measured (i.e., detected, determined) using a variety of methods standard in the art (see, for example, Sambrook et al. *ibid.*), examples of which are disclosed herein.

In one embodiment, a putative IgE-containing composition of the present method includes a biological sample from an animal. A suitable biological sample includes, but is not limited to, a bodily fluid composition or a cellular composition. A bodily fluid refers to any fluid that can be collected (i.e., obtained) from an animal, examples of which include, but are not limited to, blood, serum, plasma, urine, tears, aqueous humor, central nervous system fluid (CNF), saliva, lymph, nasal secretions, milk and feces. Such a composition of the present method can, but need not be, pretreated to remove at least some of the non-IgE isotypes of immunoglobulin and/or other proteins, such as albumin, present in the fluid. Such removal can include, but is not limited to, contacting the bodily fluid with a material, such as Protein G, to remove IgG antibodies and/or affinity purifying IgE antibodies from other components of the body fluid by exposing the fluid to, for example, Concanavalin A. In another embodiment, a composition includes collected bodily fluid that is pretreated to concentrate immunoglobulin contained in the fluid. For example, immunoglobulin contained in a bodily fluid can be precipitated from other proteins using ammonium sulfate. A preferred composition of the present method is serum.

In another embodiment, a composition of the present method includes an IgE-producing cell. Such a cell can have IgE bound to the surface of the cell and/or can secrete IgE. Examples of such cells include basophil cells and myeloma cells. IgE can be bound to the surface of a cell either directly to the membrane of a cell or bound to a molecule (e.g., an antigen) bound to the surface of the cell.

A complex can be detected in a variety of ways including, but not limited to use of one or more of the following assays: an enzyme-linked immunoassay, a radioimmunoassay, a fluorescence immunoassay, a chemiluminescent assay, a lateral flow assay, an agglutination assay, a particulate-based assay (e.g., using particulates such as, but not limited to, magnetic particles or plastic polymers, such as latex or polystyrene beads), an immunoprecipitation assay, a BioCore™ assay (e.g., using colloidal gold) and an immunoblotting assay (e.g., a western blot). Such assays are well known to those skilled in the art. Assays can be used to give qualitative or quantitative results depending on how they are used. Some assays, such as agglutination, particulate separation, and immunoprecipitation, can be observed visually (e.g., either by eye or by a machine, such as a densitometer or spectrophotometer) without the need for a detectable marker. In other assays, conjugation (i.e., attachment) of a detectable marker to the Fc<sub>ε</sub>R or to a reagent that selectively binds to the Fc<sub>ε</sub>R or to the IgE being detected (described in more detail below) aids in detecting complex formation. Examples of detectable markers include, but are not limited to, a radioactive label, a fluorescent label, a chemiluminescent label, a chromophoric label or a ligand. A ligand refers to a molecule that binds selectively to another molecule. Preferred detectable markers include, but are not limited to, fluorescein, a radioisotope, a phosphatase (e.g., alkaline phosphatase), biotin, avidin, a

peroxidase (e.g., horseradish peroxidase) and biotin-related compounds or avidin-related compounds (e.g., streptavidin or ImmunoPure® NeutrAvidin). Preferably, biotin is conjugated to an alpha chain of a Fc<sub>ε</sub>R. Preferably a carbohydrate group of the Fc<sub>ε</sub>R alpha chain is conjugated to biotin. A preferred Fc<sub>ε</sub>R molecule conjugated to biotin comprises PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (the production of which is described in the Examples section).

In one embodiment, a complex is detected by contacting a putative IgE-containing composition with a Fc<sub>ε</sub>R molecule that is conjugated to a detectable marker. A suitable detectable marker to conjugate to a Fc<sub>ε</sub>R molecule includes, but is not limited to, a radioactive label, a fluorescent label, a chemiluminescent label or a chromophoric label. A detectable marker is conjugated to a Fc<sub>ε</sub>R molecule or a reagent in such a manner as not to block the ability of the Fc<sub>ε</sub>R or reagent to bind to the IgE being detected. Preferably, a carbohydrate group of a Fc<sub>ε</sub>R is conjugated to biotin.

In another embodiment, a Fc<sub>ε</sub>R molecule:IgE complex is detected by contacting a putative IgE-containing composition with a Fc<sub>ε</sub>R molecule and then contacting the complex with an indicator molecule. Suitable indicator molecules of the present invention include molecules that can bind to either the Fc<sub>ε</sub>R molecule or to the IgE antibody. As such, an indicator molecule can comprise, for example, a Fc<sub>ε</sub>R molecule, an antigen, an antibody and a lectin, depending upon which portion of the Fc<sub>ε</sub>R molecule:IgE complex being detected. Preferred identifying labeled compounds that are antibodies include, for example, anti-IgE antibodies and anti-Fc<sub>ε</sub>R antibodies. Preferred lectins include those lectins that bind to high-mannose groups. More preferred lectins bind to high-mannose groups present on a Fc<sub>ε</sub>R molecule of the present invention produced in insect cells. An indicator molecule itself can be attached to a detectable marker of the present invention. For example, an antibody can be conjugated to biotin, horseradish peroxidase, alkaline phosphatase or fluorescein.

In one preferred embodiment, a Fc<sub>ε</sub>R molecule:IgE complex is detected by contacting the complex with a reagent that selectively binds to a Fc<sub>ε</sub>R molecule of the present invention. Examples of such a reagent includes, but are not limited to, an antibody that selectively binds to a Fc<sub>ε</sub>R molecule (referred to herein as an anti-Fc<sub>ε</sub>R antibody) or a compound that selectively binds to a detectable marker conjugated to a Fc<sub>ε</sub>R molecule. Fc<sub>ε</sub>R molecules conjugated to biotin are preferably detected using streptavidin, more preferably using ImmunoPure® NeutrAvidin (available from Pierce, Rockford, Ill.).

In another preferred embodiment, a Fc<sub>ε</sub>R molecule:IgE complex is detected by contacting the complex with a reagent that selectively binds to an IgE antibody (referred to herein as an anti-IgE reagent). Examples of such an anti-IgE reagent include, but are not limited to, a secondary antibody that is an anti-isotype antibody (e.g., an antibody that selectively binds to the constant region of an IgE), an antibody-binding bacterial surface protein (e.g., Protein A or Protein G), an antibody-binding cell (e.g., a B cell, a T cell, a natural killer cell, a polymorphonuclear leukocyte cell, a monocyte cell or a macrophage cell), an antibody-binding eukaryotic cell surface protein (e.g., an Fc receptor), and an antibody-binding complement protein. Preferred anti-IgE reagents include, but are not limited to, D9, and CMI antibody #9, CMI antibody #19, CMI antibody #59 and CMI antibody #71 (available from Custom Monoclonal International, West Sacramento, Calif.). In particular, as used herein, an anti-IgE antibody includes not only a complete antibody but also any subunit or portion thereof that is capable of selectively binding to an IgE heavy chain constant region. For example, a portion of an

anti-IgE reagent can include an Fab fragment or a F(ab')<sub>2</sub> fragment, which are described in detail in Janeway et al., in *Immunobiology, the Immune System in Health and Disease*, Garland Publishing, Inc., NY, 1996 (which is incorporated herein by this reference in its entirety).

In one embodiment a complex can be formed and detected in solution. In another embodiment, a complex can be formed in which one or more members of the complex are immobilized on (e.g., coated onto) a substrate. Immobilization techniques are known to those skilled in the art. Suitable substrate materials include, but are not limited to, plastic, glass, gel, celluloid, paper, PVDF (poly-vinylidene-fluoride), nylon, nitrocellulose, and particulate materials such as latex, polystyrene, nylon, nitrocellulose, agarose and magnetic resin. Suitable shapes for substrate material include, but are not limited to, a well (e.g., microtiter dish well), a plate, a dipstick, a bead, a lateral flow apparatus, a membrane, a filter, a tube, a dish, a celluloid-type matrix, a magnetic particle, and other particulates. A particularly preferred substrate comprises an ELISA plate, a dipstick, a radioimmunoassay plate, agarose beads, plastic beads, latex beads, immunoblot membranes and immunoblot papers. In one embodiment, a substrate, such as a particulate, can include a detectable marker.

A preferred method to detect IgE is an immunosorbent assay. An immunoabsorbent assay of the present invention comprises a capture molecule and an indicator molecule. A capture molecule of the present invention binds to an IgE in such a manner that the IgE is immobilized to a substrate. As such, a capture molecule is preferably immobilized to a substrate of the present invention prior to exposure of the capture molecule to a putative IgE-containing composition. An indicator molecule of the present invention detects the presence of an IgE bound to a capture molecule. As such, an indicator molecule preferably is not immobilized to the same substrate as a capture molecule prior to exposure of the capture molecule to a putative IgE-containing composition.

A preferred immunoabsorbent assay method includes a step of either: (a) binding an Fc<sub>ε</sub>R molecule to a substrate prior to contacting a Fc<sub>ε</sub>R molecule with a putative IgE-containing composition to form a Fc<sub>ε</sub>R molecule-coated substrate; or (b) binding a putative IgE-containing composition to a substrate prior to contacting a Fc<sub>ε</sub>R molecule with a putative IgE-containing composition to form a putative IgE-containing composition-coated substrate. Preferably, the substrate includes of a non-coated substrate, a Fc<sub>ε</sub>R molecule-coated substrate, an antigen-coated substrate or an anti-IgE antibody-coated substrate.

Both a capture molecule and an indicator molecule of the present invention are capable of binding to an IgE. Preferably, a capture molecule binds to a different region of an IgE than an indicator molecule, thereby allowing a capture molecule to be bound to an IgE at the same time as an indicator molecule. The use of a reagent as a capture molecule or an indicator molecule depends upon whether the molecule is immobilized to a substrate when the molecule is exposed to an IgE. For example, a Fc<sub>ε</sub>R molecule of the present invention is used as a capture molecule when the Fc<sub>ε</sub>R molecule is bound to a substrate. Alternatively, a Fc<sub>ε</sub>R molecule is used as an indicator molecule when the Fc<sub>ε</sub>R molecule is not bound to a substrate. Suitable molecule for use as capture molecules or indicator molecules include, but are not limited to, a Fc<sub>ε</sub>R molecule of the present invention, an antigen reagent or an anti-IgE antibody reagent of the present invention.

An immunoabsorbent assay of the present invention can further comprise one or more layers and/or types of secondary molecules or other binding molecules capable of detecting the presence of an indicator molecule. For example, an

untagged (i.e., not conjugated to a detectable marker) secondary antibody that selectively binds to an indicator molecule can be bound to a tagged (i.e., conjugated to a detectable marker) tertiary antibody that selectively binds to the secondary antibody. Suitable secondary antibodies, tertiary antibodies and other secondary or tertiary molecules can be selected by those of skill in the art. Preferred secondary molecules of the present invention include, an antigen, an anti-IgE idiotype antibody and an anti-IgE isotypic. Preferred tertiary molecules can be selected by a skilled artisan based upon the characteristics of the secondary molecule. The same strategy is applied for subsequent layers.

In one embodiment, a desired antigen is used as a capture molecule by being immobilized on a substrate, such as a microtiter dish well or a dipstick. Preferred antigens include those disclosed herein. A biological sample collected from an animal is applied to the substrate and incubated under conditions suitable (i.e., sufficient) to allow for antigen:IgE complex formation bound to the substrate (i.e., IgE in a sample binds to an antigen immobilized on a substrate). Excess non-bound material (i.e., material from the biological sample that has not bound to the antigen), if any, is removed from the substrate under conditions that retain antigen:IgE complex binding to the substrate. Preferred conditions are disclosed herein in the Examples section and generally in Sambrook et al., *ibid*. An indicator molecule that can selectively bind to an IgE bound to the antigen, the indicator molecule can be conjugated to a detectable marker (preferably to an enzyme label, to a colorimetric label, to a fluorescent label, to a radioisotope, or to a ligand such as of the biotin or avidin family), is added to the substrate and incubated to allow formation of a complex between the indicator molecule and the antigen:IgE complex. Excess indicator molecule is removed, a developing agent is added if required, and the substrate is submitted to a detection device for analysis. A preferred indicator molecule for this embodiment is a Fc<sub>ε</sub>R molecule, preferably conjugated to biotin, to a fluorescent label or to an enzyme label.

In one embodiment, a Fc<sub>ε</sub>R molecule is used as a capture molecule by being immobilized on a substrate, such as a microtiter dish well or a dipstick. A biological sample collected from an animal is applied to the substrate and incubated under conditions suitable to allow for Fc<sub>ε</sub>R molecule:IgE complex formation bound to the substrate. Excess non-bound material, if any, is removed from the substrate under conditions that retain Fc<sub>ε</sub>R molecule:IgE complex binding to the substrate. An indicator molecule that can selectively bind to an IgE bound to the Fc<sub>ε</sub>R is added to the substrate and incubated to allow formation of a complex between the indicator molecule and the Fc<sub>ε</sub>R molecule:IgE complex. Preferably, the indicator molecule is conjugated to a detectable marker (preferably to an enzyme label, to a colorimetric label, to a fluorescent label, to a radioisotope, or to a ligand such as of the biotin or avidin family). Excess indicator molecule is removed, a developing agent is added if required, and the substrate is submitted to a detection device for analysis. A preferred indicator molecule for this embodiment is an antigen that will bind to IgE in the biological sample or an anti-IgE isotype or idiotype antibody, either preferably being conjugated to fluorescein or biotin.

In one embodiment, an anti-IgE antibody (e.g., isotype or idiotype specific antibody) is used as a capture molecule by being immobilized on a substrate, such as a microtiter dish well or a dipstick. A biological sample collected from an animal is applied to the substrate and incubated under conditions suitable to allow for anti-IgE antibody:IgE complex formation bound to the substrate. Excess non-bound material, if any, is removed from the substrate under conditions that

retain anti-IgE antibody:IgE complex binding to the substrate. A Fc<sub>ε</sub>R molecule is added to the substrate and incubated to allow formation of a complex between the Fc<sub>ε</sub>R molecule and the anti-IgE antibody:IgE complex. Preferably, the Fc<sub>ε</sub>R molecule is conjugated to a detectable marker (preferably to biotin, an enzyme label or a fluorescent label). Excess Fc<sub>ε</sub>R molecule is removed, a developing agent is added if required, and the substrate is submitted to a detection device for analysis.

In one embodiment, an immunosorbent assay of the present invention does not utilize a capture molecule. In this embodiment, a biological sample collected from an animal is applied to a substrate, such as a microtiter dish well or a dipstick, and incubated under conditions suitable to allow for IgE binding to the substrate. Any IgE present in the bodily fluid is immobilized on the substrate. Excess non-bound material, if any, is removed from the substrate under conditions that retain IgE binding to the substrate. A Fc<sub>ε</sub>R molecule is added to the substrate and incubated to allow formation of a complex between the Fc<sub>ε</sub>R molecule and the IgE. Preferably, the Fc<sub>ε</sub>R molecule is conjugated to a detectable marker (preferably to biotin, an enzyme label or a fluorescent label). Excess Fc<sub>ε</sub>R molecule is removed, a developing agent is added if required, and the substrate is submitted to a detection device for analysis.

Another preferred method to detect IgE is a lateral flow assay, examples of which are disclosed in U.S. Pat. No. 5,424, 193, issued Jun. 13, 1995, by Pronovost et al.; U.S. Pat. No. 5,415,994, issued May 16, 1995, by Imrich et al; WO 94/29696, published Dec. 22, 1994, by Miller et al.; and WO 94/01775, published Jan. 20, 1994, by Pawlak et al.; each of these patent publications is incorporated by reference herein in its entirety. In one embodiment, a biological sample is placed in a lateral flow apparatus that includes the following components: (a) a support structure defining a flow path; (b) a labeling reagent comprising a bead conjugated to an antigen, the labeling reagent being impregnated within the support structure in a labeling zone; and (c) a capture reagent comprising an IgE-binding composition. Preferred antigens include those disclosed herein. The capture reagent is located downstream of the labeling reagent within a capture zone fluidly connected to the labeling zone in such a manner that the labeling reagent can flow from the labeling zone into the capture zone. The support structure comprises a material that does not impede the flow of the beads from the labeling zone to the capture zone. Suitable materials for use as a support structure include ionic (i.e., anionic or cationic) material. Examples of such a material include, but are not limited to, nitrocellulose (NC), PVDF, carboxymethylcellulose (CM). The support structure defines a flow path that is lateral and is divided into zones, namely a labeling zone and a capture zone. The apparatus can further comprise a sample receiving zone located along the flow path, more preferably upstream of the labeling reagent. The flow path in the support structure is created by contacting a portion of the support structure downstream of the capture zone, preferably at the end of the flow path, to an absorbent capable of absorbing excess liquid from the labeling and capture zones.

In this embodiment, the biological sample is applied to the sample receiving zone which includes a portion of the support structure. The labeling zone receives the sample from the sample receiving zone which is directed downstream by the flow path. The labeling zone comprises the labeling reagent that binds to IgE. A preferred labeling reagent is an antigen conjugated, either directly or through a linker, to a plastic bead substrate, such as to a latex bead. The substrate also includes a detectable marker, preferably a colorimetric

marker. Typically, the labeling reagent is impregnated to the support structure by drying or lyophilization. The sample structure also comprises a capture zone downstream of the labeling zone. The capture zone receives labeling reagent from the labeling zone which is directed downstream by the flow path. The capture zone contains the capture reagent, in this case a Fc<sub>ε</sub>R molecule, as disclosed above, that immobilizes the IgE complexed to the antigen in the capture zone. The capture reagent is preferably fixed to the support structure by drying or lyophilizing. The labeling reagent accumulates in the capture zone and the accumulation is assessed visually or by an optical detection device.

In another embodiment, a lateral flow apparatus used to detect IgE includes: (a) a support structure defining a flow path; (b) a labeling reagent comprising a Fc<sub>ε</sub>R molecule as described above, the labeling reagent impregnated within the support structure in a labeling zone; and (c) a capture reagent comprising an antigen, the capture reagent being located downstream of the labeling reagent within a capture zone fluidly connected to the labeling zone in such a manner that the labeling reagent can flow from the labeling zone into the capture zone. The apparatus preferably also includes a sample receiving zone located along the flow path, preferably upstream of the labeling reagent. The apparatus preferably also includes an absorbent located at the end of the flow path.

One embodiment of the present invention is an inhibition assay in which the presence of IgE in a putative IgE-containing composition is determined by adding such composition to a Fc<sub>ε</sub>R molecule of the present invention and an isolated IgE known to bind to the Fc<sub>ε</sub>R molecule. The absence of binding of the Fc<sub>ε</sub>R molecule to the known IgE indicating the presence of IgE in the putative IgE-containing composition.

The present invention also includes kits to detect IgE based on each of the disclosed detection methods. One embodiment is a kit to detect IgE comprising a human Fc<sub>ε</sub> receptor (Fc<sub>ε</sub>R) molecule and a means for detecting an IgE including canine IgE, feline IgE and/or equine IgE. Suitable and preferred Fc<sub>ε</sub>R molecules are disclosed herein. Suitable means of detection include compounds disclosed herein that bind to either the Fc<sub>ε</sub>R molecule or to an IgE. A preferred kit of the present invention further comprises a detection means including one or more antigens disclosed herein, an antibody capable of selectively binding to an IgE disclosed herein and/or a compound capable of binding to a detectable marker conjugated to a Fc<sub>ε</sub>R molecule (e.g., avidin, streptavidin and ImmunoPure® NeutrAvidin when the detectable marker is biotin). Such antigens preferably induce IgE antibody production in animals including canines, felines and/or equines.

A preferred embodiment of a kit of the present invention is a flea allergen kit comprising a flea allergen such as those disclosed herein. A particularly preferred flea allergen for use with a flea allergen kit includes a flea saliva product or a flea saliva protein.

Another preferred kit of the present invention is a general allergen kit comprising an allergen common to all regions of the United States and a human Fc<sub>ε</sub>R molecule of the present invention. As used herein, a "general allergen" kit refers to a kit comprising allergens that are found substantially throughout the United States (i.e., essentially not limited to certain regions of the United States). A general allergen kit provides an advantage over regional allergen kits because a single kit can be used to test an animal located in most geographical locations on the United States. Suitable and preferred general allergens for use with a general allergen kit of the present invention include those general allergens disclosed herein.

Another preferred kit of the present invention is a food allergen kit comprising a food allergen including beef,

15

chicken, pork, a mixture of fish, such as cod, halibut or and tuna, egg, milk, Brewer's yeast, whole wheat, corn, soybean, cheese and rice, and a human  $Fc_\epsilon R$  molecule of the present invention. Preferably, the beef, chicken, pork, fish, corn and rice, are cooked.

A preferred kit of the present invention includes those in which the allergen is immobilized to a substrate. If a kit comprises two or more antigens, the kit can comprise one or more compositions, each composition comprising one antigen. As such, each antigen can be tested separately. A kit can also contain two or more diagnostic reagents for IgE, additional isolated IgE antigens and/or antibodies as disclosed herein. Particularly preferred are kits used in a lateral flow assay format. It is within the scope of the present invention that a lateral flow assay kit can include one or more lateral flow assay apparatuses. Multiple lateral flow apparatuses can be attached to each other at one end of each apparatus, thereby creating a fan-like structure.

In particular, a method and kit of the present invention are useful for diagnosing abnormal conditions in animals that are associated with changing levels of IgE. Particularly preferred conditions to diagnose include allergies, parasitic infections and neoplasia. For example, a method and kit of the present invention are particularly useful for detecting flea allergy dermatitis (FAD), when such method or kit includes the use of a flea saliva antigen. FAD is defined as a hypersensitive response to fleabites. Preferably, a putative IgE-containing composition is obtained from an animal suspected of having FAD. Preferred animals include those disclosed herein, with dogs and cats being more preferred. In addition, methods and kits of the present invention are particularly useful for detecting helminth infection, in particular heartworm infection, when such methods or kits include the use of a helminth antigen, such as Di33. Preferably, a putative IgE-containing composition is obtained from an animal suspected of having a helminth infection. Preferred animals include those disclosed herein, with dogs and cats being more preferred.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention.

## EXAMPLES

### Example 1

This example describes the construction of a recombinant baculovirus expressing a truncated portion of the  $\alpha$ -chain of the human  $Fc_\epsilon$  receptor.

Recombinant molecule pVL-nhFc $\epsilon$ R $\alpha_{612}$ , containing a nucleic acid molecule encoding the extracellular domain of the  $Fc_\epsilon R \alpha$  chain, operatively linked to baculovirus polyhedron transcription control sequences was produced in the following manner. A cDNA clone encoding the full-length alpha chain ( $\alpha$  chain) of the human  $Fc_\epsilon$  receptor was obtained from Dr. Jean-Pierre Kinet (Harvard University, Cambridge, Mass.). The cDNA clone included an about 1198 nucleotide insert, referred to herein as nhFc $\epsilon$ R $\alpha_{1198}$ . The nucleic acid sequence of the coding strand of nhFc $\epsilon$ R $\alpha_{1198}$  is denoted herein as SEQ ID NO:1. Translation of SEQ ID NO:1 indicates that nucleic acid molecule nhFc $\epsilon$ R $\alpha_{1198}$  encodes a full-length human  $Fc_\epsilon$  receptor  $\alpha$  chain protein of about 257 amino acids, referred to herein as PhFc $\epsilon$ R $\alpha_{257}$ , having amino acid sequence SEQ ID NO:2, assuming an open reading frame in which the initiation codon spans from about nucleotide 107 through about nucleotide 109 of SEQ ID NO:1 and the termination codon spans from about nucleotide 878 through about nucleotide 880 of SEQ ID NO:1. The complement of

16

SEQ ID NO:1 is represented herein by SEQ ID NO:3. The proposed mature protein In (i.e., Fc $\epsilon$ R $\alpha$  chain from which the signal sequence has been cleaved), denoted herein as PhFc $\epsilon$ R $\alpha_{232}$ , contains about 232 amino acids which is represented herein as SEQ ID NO:6. The nucleic acid molecule encoding PhFc $\epsilon$ R $\alpha_{232}$  is denoted herein as nhFc $\epsilon$ R $\alpha_{696}$ , the coding strand of which is represented by SEQ ID NO:7.

To produce a secreted form of the extracellular domain of the  $Fc_\epsilon R \alpha$  chain, the hydrophobic transmembrane domain and the cytoplasmic tail of the  $Fc_\epsilon R \alpha$  chain encoded by nhFc $\epsilon$ R $\alpha_{1198}$  were removed as follows. A  $Fc_\epsilon R \alpha$  chain extracellular domain nucleic acid molecule-containing fragment of about 612 nucleotides was PCR amplified from nhFc $\epsilon$ R $\alpha_{1198}$  using a forward primer EJH 040 containing a BamHI site, having the nucleic acid sequence 5' CGC GOA TCC TAT AAT ATG GCT CCT GCC ATG G 3' (denoted SEQ ID NO:8) and a reverse primer IgE ANTI-SENSE containing an EcoRI site, having the nucleic acid sequence 5' GGC GAA TTC TTA AGC TTT TAT TAC AG 3' (denoted herein as SEQ ID NO:9). The resulting PCR product was digested with BamHI and EcoRI to produce nhFc $\epsilon$ R $\alpha_{612}$ . Nucleic acid molecule nhFc $\epsilon$ R $\alpha_{612}$  contained an about 591 nucleotide fragment encoding the extracellular domain of the human  $Fc_\epsilon R \alpha$  chain, extending from about nucleotide 107 to about nucleotide 697 of SEQ ID NO 1, denoted herein as nucleic acid molecule nhFc $\epsilon$ R $\alpha_{591}$ , the coding strand of which has a nucleic acid sequence denoted SEQ ID NO:10. Translation of SEQ ID NO:10 indicates that nucleic acid molecule nhFc $\epsilon$ R $\alpha_{612}$  encodes a  $Fc_\epsilon R$  protein of about 197 amino acids, referred to herein as PhFc $\epsilon$ R $\alpha_{197}$ , having amino acid sequence SEQ ID NO:11. Nucleic acid molecule nhFc $\epsilon$ R $\alpha_{612}$  encodes a secretable form of the human  $Fc_\epsilon R \alpha$  chain which does not possess a leader sequence, which is denoted herein as PhFc $\epsilon$ R $\alpha_{172}$  having amino acid sequence SEQ ID NO:13. The coding region for PhFc $\epsilon$ R $\alpha_{172}$  is denoted nhFc $\epsilon$ R $\alpha_{516}$ , the coding strand of which has a nucleic acid sequence denoted SEQ ID NO:12. The complement of SEQ ID NO:12 is represented herein by SEQ ID NO:14.

In order to produce a baculovirus recombinant molecule capable of directing the production of PhFc $\epsilon$ R $\alpha_{197}$ , the nucleic acid molecule nhFc $\epsilon$ R $\alpha_{612}$  was subcloned into unique BamHI and EcoRI sites of pVL1392 baculovirus shuttle plasmid (available from Pharmingen, San Diego, Calif.) to produce a recombinant molecule referred to herein as pVL-nhFc $\epsilon$ R $\alpha_{612}$ . The resultant recombinant molecule pVL-nhFc $\epsilon$ R $\alpha_{612}$  was verified for proper insert orientation by restriction mapping.

### Example 2

This example describes the production of PhFc $\epsilon$ R $\alpha_{172}$  protein.

The recombinant molecule pVL-nhFc $\epsilon$ R $\alpha_{612}$  was co-transfected with a linear Baculogold baculovirus DNA (available from Pharmingen) into *Trichoplusia ni* cells (available from Invitrogen Corp., San Diego, Calif.; High Five™ cells) using the following method. About 1.5 liter cultures of serum-free ex-Cell Medium (available from Invitrogen) were seeded with about  $1 \times 10^6$  cells per ml of medium. The *Trichoplusia ni* cells were infected with recombinant molecule pVL-nhFc $\epsilon$ R $\alpha_{612}$  at a multiplicity of infection (MOI) of about 2 to about 5 particle forming units (pfu) per cell to produce recombinant cell *Trichoplusia ni*-pVL-nhFc $\epsilon$ R $\alpha_{612}$ . The infection was allowed to proceed at a controlled temperature of 27° C. for 48 hours, to produce recombinant protein

## 17

PhFc<sub>ε</sub>Rα<sub>172</sub>. Following infection, cells were separated from the medium by centrifugation, and the medium was frozen at -70° C.

PhFc<sub>ε</sub>Rα<sub>172</sub> was purified from the culture medium described immediately above by affinity chromatography using an IgE antibody produced by the myeloma cell line U266DI (American Tissue Type Catalogue No. TIB196) linked to sepharose 4B. The amino acid composition and N-terminal amino acid sequence of the affinity purified PhFc<sub>ε</sub>Rα<sub>172</sub> were determined using methods standard in the art. The results indicated that PhFc<sub>ε</sub>Rα<sub>172</sub> was properly processed by the *Trichoplusia ni* cells.

## Example 3

This example describes the biotinylation of a recombinant human Fc<sub>ε</sub>R alpha chain protein.

Affinity purified recombinant protein PhFc<sub>ε</sub>Rα<sub>172</sub>, prepared as described above in Example 2, was biotinylated as follows. About 440 micrograms (μg) of PhFc<sub>ε</sub>Rα<sub>172</sub> were diluted in about 1.5 milliliter (ml) of acetate buffer (0.1 M NaAc, pH 5.5) containing about 200 microliter (μl) of 0.1 M NaIO<sub>4</sub>. The mixture was incubated for about 20 minutes, on ice, and about 2 μl of glycerol was added following the incubation. The mixture was then dialyzed against about 2 liters of acetate buffer in a 3 ml Slide-A-Lyzer cassette (available from Pierce, Rockford, Ill.), 2 times for about 2 hours each time. About 3.72 μg of biotin-LC-hydrazide (available from Pierce) was dissolved in about 200 μl of dimethylsulfoxide (DMSO) and injected into the cassette. The cassette was then rocked at room temperature for about 2 hours. Following the incubation, the mixture containing recombinant protein and biotin dialyzed 3 times, a first time for about 18 hours and two times for about 2 hours, each time at 5° C. against phosphate buffered saline. The biotinylated protein was recovered from the dialysis, and is referred to herein as PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

## Example 4

This example describes detection of canine IgE in a solid-phase ELISA using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Wells of two Immulon II microtiter plates (available from Dynatech, Alexandria, Va.) were coated with duplicate samples of about 100 μl/well of various concentrations of purified canine IgE as denoted in FIG. 1. The canine IgE was obtained from a canine IgE producing hybridoma, such as heterohybridoma 2.39 (described in Gebhard et al., *Immunology* 85:429-434, 1995) and was diluted in a CBC buffer (15 mM Na<sub>2</sub>CO<sub>3</sub> and 34.8 mM NaHCO<sub>3</sub>, pH 9.6. The coated plates were incubated overnight at 4° C. Following incubation, the canine IgE-containing solution was removed from each plate, and the plates were blotted dry. The plates were then blocked using about 200 μl/well of 0.25% bovine serum albumin (BSA) contained in phosphate buffered saline (PBSB) for about 1 hour at room temperature. The plates were then washed four times with 0.05% Tween-20 in PBS (PBST) using an automatic washer (available from Dynatech). Experimental samples consisting of about 100 μl/well of a 1:4000 dilution of 40 μg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (about 145 μg/ml; described in Example 3), contained in PBSB with 0.05% Tween-20 (PESBT) were added to each well of one plate coated with canine IgE.

Control samples consisting of about 100 μl of biotinylated anti-canine IgE monoclonal antibody D9 (supplied by Dr. DeBoer, U. of Wisconsin, Madison, Wis.) were added to each well of the other plate coated with canine IgE. The plates were incubated for 1 hour at room temperature and then washed

## 18

four times with PBST. About 100 μl of about 0.25 μg/ml streptavidin conjugated to horseradish peroxidase (available from Kirkegaard and Perry Laboratories (KPL), Gaithersburg, Md.; diluted in PBST) was added to each well that received experimental or control samples. The plates were then incubated for 1 hour at room temperature and washed four times with PBST. About 100 μl of TMB substrate (available from available from KPL), that had been pre-warmed to room temperature, was added. Plates were then incubated for 10 minutes at room temperature and then about 100 μl/well of Stop Solution (available from KPL) was added. Optical densities of wells were read on a Spectramax Microtiter Plate (available from Molecular Devices Inc.) reader at 450 nm within 10 minutes of adding the stop solution.

The results shown in FIG. 1 indicate that the alpha chain of human Fc<sub>ε</sub>R detects the presence of canine IgE (closed squares) in a solid-phase assay in a similar manner as the control antibody that binds specifically to canine IgE (D9; closed circles).

## Example 5

This example describes detection of plant allergen-specific canine IgE using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Immulon II microtiter plate (available from Dynatech) were coated with either about 100 μl/well of 1 μg/ml of Kentucky Blue Grass allergen or about 100 μl/well of about 1 μg/ml of Green Ash allergen (both available from Greer Inc., Lenoir, N.C.) both diluted in CBC buffer. The plate was incubated overnight at 4° C. The plate was blocked and washed as described in Example 4. Two different pools of canine sera were then added to the antigen-coated wells. The first pool consisted of sera isolated from 8 dogs reported to be allergen reactive. The second pool consisted of sera isolated from 8 dogs reported to be allergen non-reactive. Each pool of sera was diluted 1:10 or 1:100 in PBST. About 100 μl of each concentration of each diluted sera sample was added to the wells and incubated for 1 hour at room temperature. The plate was then washed four times with PBST. About 100 μl/well of a 1:4000 dilution of 40 g/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3), contained in PBSBT was added to the antigen-coated wells. The plate was incubated for 1 hour at room temperature. The plate was then washed four times with PBST. About 100 μl/well of about 0.25 μg/ml of neutravidin conjugated to horseradish peroxidase (available from Pierce) contained in PBSBT, was added. The plate was incubated for 1 hour at room temperature. The plate was then washed and the presence of neutravidin bound to the plate detected using the method described in Example 4.

The results shown in FIG. 2 indicate that the alpha chain of human Fc<sub>ε</sub>R detects the presence of canine IgE antibodies that bind specifically to a common grass allergen or to a common tree allergen. In addition, detection of canine IgE antibodies is dose dependent.

## Example 6

This example describes detection of total canine IgE using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Immulon II microtiter plate (available from Dynatech) were coated with about 100 μl/well of about 1 μg/ml CMI anti-canine IgE antibody #6 (available from Custom Monoclonals International, West Sacramento, Calif.) diluted in CBC buffer. The plate was incubated overnight at 4° C. The plate was blocked and washed as described in Example 4. About 100 μl/well of a 1:60 dilution in PBSBT of sera samples from a variety of sources were then added to

multiple wells coated with anti-IgE antibody. The samples included: (1) serum from a dog known to be allergic to flea saliva; (2) serum from dogs infected with *D. immitis*; (3) and (4) a pool of dog sera from defined as canine allergy calibrators (available from BioProducts DVM, Tempe, Ariz.); (5) pools of dog sera containing antibodies that have low binding to Kentucky Blue Grass allergen; (6) pools of dog sera that have high binding to Kentucky Blue Grass allergen; (7) a pool of dog sera from dogs known to be allergic to flea saliva, the sample was heat inactivated (at 56° C. for 4 hours); (8) a pool of dog sera from dogs known to be allergic to flea saliva; or (9) a pool of dog sera from dogs raised in a barrier facility (i.e., negative control). A set of positive control samples consisting of IgE derived from the canine heterohybridoma described in Example 4 were also added to the plate to generate a standard curve. The plate was incubated for 1 hour at room temperature and then washed four times with PBST. The presence of canine IgE was detected using either about 100 µl/well of a 1:4000 dilution of 40 µg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3) or about 100 µl/well of about 1 µg/ml CMI anti-canine IgE antibody #19 (available from Custom Monoclonals International), both contained in PBSBT. The plate was incubated for 1 hour at room temperature. The plate was then washed, contacted with about 0.25 µg/ml streptavidin conjugated to horseradish peroxidase, washed again, and the presence of streptavidin bound to the plate was detected using the method described in Example 4. The optical density readings obtained for the control samples were used to generate a standard curve that was used to determine the total IgE bound to wells that had received test samples.

The results shown in FIG. 3 indicate that canine IgE from a variety of dog sera are detected using the alpha chain of human Fc<sub>ε</sub>R in a manner similar to using an antibody that binds specifically to canine IgE. The absence of detectable amounts of IgE in the heat treated sample (Sample 7) indicates that the antibody detected by PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT is IgE. In addition, the results indicate that PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT is an effective reagent for detecting IgE that binds to allergen Kentucky Blue Grass, Samples 5 and 6), as well as a parasite antigen (*D. immitis*, Sample 2).

#### Example 7

This example describes detection of canine IgE in dog sera isolated from dogs known to be allergic to flea saliva, using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Immulon IT microtiter plate were coated with about 100 µl/well of varying concentrations of flea saliva recombinant protein fspN (described in PCT Patent Publication No. WO 96/11271, *ibid.*; concentrations shown in FIG. 4) diluted in CBC buffer. The plate was incubated overnight at 4° C. The plate was then blocked and washed as described in Example 4. About 100 µl/well of a 1:10 dilution in PBSBT of a pool of sera isolated from dogs known to produce IgE that binds specifically to flea saliva. Some wells did not receive dog sera so that background binding levels could be determined. The plate was incubated for 1 hour at room temperature and then washed four times with PBST. About 100 µl/well of a 1:4000 dilution of 40 µg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3) contained in PBSBT was added. The plate was incubated for 1 hour at room temperature. The plate was then washed, contacted with about 0.25 µg/ml streptavidin-conjugated to horseradish peroxidase, washed again, and the presence of streptavidin bound to the plate was detected using the method described in Example 4.

The results shown in FIG. 4 indicate that canine IgE that binds specifically to a flea saliva antigen is detected using the alpha chain of human Fc<sub>ε</sub>R.

#### Example 8

This example describes detection of total canine IgE in dog sera isolated from dogs known to be allergic to flea saliva, heartworm-infected dogs and specific pathogen free (SPF) dogs, using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Immulon II microtiter plate were coated with about 100 µl/well of about 1 µg/ml CMI anti-canine IgE antibody #6 (available from Custom Monoclonals International) in CBC buffer. The plate was incubated overnight at 4° C. The plate was blocked and washed as described in Example 4. About 100 µl/well of different samples of IgE-containing fluids in PBSBT were added to multiple wells coated with the anti-canine IgE antibody. The samples included: (1) 100 µg/ml of canine IgE purified from the heterohybridoma described in Example 4; (2) a 1:10 dilution of a pool of sera from dogs known to be allergic to flea saliva, (3) a 1:10 dilution of the same sera pool as in (2) but heat inactivated; (4) a 1:10 dilution of serum from a dog known to have clinical flea allergy dermatitis (dog CPO2); (5) a 1:10 dilution of heat inactivated CPO2 serum; (6) a 1:10 dilution of serum from a heartworm-infected dog (dog 417); (7) a 1:10 dilution of heat inactivated 417 serum; (8) a 1:10 dilution of a pool of sera from heartworm-infected dogs; (9) a 1:10 dilution of the same sera pool as in (8) but heat inactivated; and (10) a pool of sera from dogs raised in a barrier facility. Each sample was diluted in PBSBT. The plate was incubated for 1 hour at room temperature and then washed four times with PBST. About 100 µl/well of a 1:4000 dilution of 40 µg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3) in PBSBT was added. The plate was incubated for 1 hour at room temperature. The plate was then washed, contacted with about 0.25 µg/ml streptavidin-conjugated to horseradish peroxidase, washed again, and the presence of streptavidin bound to the plate was detected using the method described in Example 4.

The results shown in FIG. 5 indicate that canine IgE from dogs allergic to flea saliva and from dogs infected with heartworm are detected using the alpha chain of human Fc<sub>ε</sub>R. In addition, the absence of colorimetric signal in samples of heat inactivated sera indicates that antibody bound to the anti-IgE antibody and detected by Fc<sub>ε</sub>R alpha chain is an epsilon isotype antibody and not another isotype.

#### Example 9

This example describes detection of IgE that specifically binds to flea saliva, using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Immulon II microtiter plate were coated with about 100 µl/well of about 0.1 µg/ml of flea saliva collected using the method described in PCT Patent Publication No. WO 96/11271, *ibid.*, in CBC buffer. The plate was incubated, blocked and washed as described in Example 4. The IgE-containing samples described in Example 8 were then applied to the flea saliva coated plate. The plate was then treated using the method described in Example 8.

The results shown in FIG. 6 indicate that canine IgE that binds specifically to flea saliva, contained in serum, is detected using the alpha chain of human Fc<sub>ε</sub>R. In addition, the absence of colorimetric signal in samples of heat inactivated serum indicates that antibody bound to the flea saliva protein and detected by Fc<sub>ε</sub>R alpha chain is an epsilon isotype antibody.

## 21

## Example 10

This example describes the detection of feline IgE using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Trnmulon II microtiter plate were coated with about 100 μl/well of about 10 μg/ml Di33 protein (described in U.S. Pat. No. 6,391,569 B1, *ibid.*) or 10 μg/ml crude homogenate of heartworm, both in CBC buffer. Crude homogenate of heartworm is the clarified supernatant of adult heartworms homogenized in PBS. The plate was incubated overnight at 4° C. The plate was blocked and washed as described in Example 4. Serum samples from 2 heartworm infected cats were then added to Di33-coated wells and to heartworm antigen-coated wells. About 100 μl/well of a 1:10 dilution in PBSBT of sera from heartworm-infected cat #AXH3 or from cat #MGC2 were added to the plate. Negative control samples consisting of serum from pre-infection bleeds of cat #AXH3 and cat #MGC2 were also added to the plate at a dilution of 1:10 in PBSBT. A positive control sample consisting of a pool of sera from heartworm-infected dogs was also added to the plate at a dilution of 1:10 in PBSBT. The plate was incubated for 1 hour at room temperature and then washed four times with PBST. About 100 μl/well of a 1:4000 dilution of 40 μg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3) in PBSBT was added. The plate was incubated for 1 hour at room temperature. The plate was then washed, contacted with 1:4000 dilution of a 0.5 mg/ml solution of streptavidin-conjugated to horseradish peroxidase, washed again, and the presence of streptavidin bound to the plate was detected using the method described in Example 4.

The results shown in FIG. 7 indicate that feline IgE that binds specifically to crude homogenate of heartworm or Di33 protein is detected using the alpha chain of human Fc<sub>ε</sub>R.

## Example 11

This example describes detection of feline IgE using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Multiple wells of an Immulon II microtiter plate were coated with Di33 as described in Example 10, in CBC buffer. The plate was incubated overnight at 4° C. The plate was blocked and washed as described in Example 4. Serum samples from 2 heartworm infected cats were then added to Di33-coated wells. About 100 μl/well of a 1:10 dilution in PBSBT of serum from heartworm-infected cat # MGC2 and a pool of sera from heartworm-infected cats, as well as heat inactivated samples of each of these sera, were added to the plate. A positive control sample consisting of a pool of sera from heartworm-infected dogs was also added to the plate at a dilution of 1:10 in PBSBT. The plate was incubated for 1 hour at room temperature and then washed four times with PBST. About 100 μl/well of a 1:4000 dilution of 40 μg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3) in PBSBT was added. The plate was incubated for 1 hour at room temperature. The plate was then washed, contacted with streptavidin-conjugated to horseradish peroxidase, washed again, and the presence of streptavidin bound to the plate was detected using the method described in Example 4.

The results shown in FIG. 8 indicate that feline IgE from heartworm-infected cats that specifically binds to the heartworm antigen Di33 is detected using the alpha chain of human Fc<sub>ε</sub>R. In addition, the absence of calorimetric signal in

## 22

samples of heat inactivated sera indicates that antibody bound to the Di33 protein and detected by Fc<sub>ε</sub>R alpha chain is an epsilon isotype antibody.

## Example 12

This example describes detection of equine IgE in a solid-phase ELISA using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT.

Horse sera from a horse known to be allergic to certain allergens and horse sera from a horse known not to be allergic the same allergens, were assayed for the presence of IgE using PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT as follows. A North Atlantic/Ohio Valley Regional Panel plate of a Canitec™ Allergen-Specific IgE Kit (available from BioProducts DVM) was blocked and washed as described in Example 4. Two samples of about 1:10 dilutions of the two horse sera were prepared using PBSBT. The two samples were added to the blocked plate and the plate was incubated for 1 hour at room temperature. The plate was washed as described in Example 4. About 100 μl/well of a 1:4000 dilution of 40 μg/ml PhFc<sub>ε</sub>Rα<sub>172</sub>-BIOT (described in Example 3), contained in PBSBT was added to each well. The plate was then washed, contacted with 1:4000 dilution of a 0.5 mg/ml solution of streptavidin-conjugated to horseradish peroxidase, washed again, and the presence of streptavidin bound to the plate was detected using the method described in Example 4.

The results shown in FIG. 9 indicate that equine IgE from a horse known to be allergic to certain allergens specifically binds to certain plant and mite allergens is detected using the alpha chain of human Fc<sub>ε</sub>R.

## Example 13

This example describes detection of canine IgE in a solid-phase ELISA using basophilic cells transfected with human Fc<sub>ε</sub>R alpha chain.

Rat basophilic leukemia (RBL) cells transfected with a nucleic acid molecule encoding a human Fc<sub>ε</sub>R alpha chain (referred to herein as RBL-hFc<sub>ε</sub>R cells; described in Miller et al., *Science* 244:334-337, 1989) were used to detect canine IgE as follows. About 4×10<sup>4</sup> RBL-hFc<sub>ε</sub>R cells contained in Earles Modified Eagles Medium containing 10% fetal bovine serum (EMEM-FBS) were added to each well of 96-well flat bottom tissue culture plates. The RBL-hFc<sub>ε</sub>R cells were incubated overnight at 37° C. Following the incubation the plates were washed 4 times with PBST. The cells were then fixed for about 2 minutes using about 200 μl per well of absolute alcohol at room temperature. The plates were then washed 8 times with PBST to remove residual alcohol.

Serial dilutions in EMEM-FBS (concentrations shown in FIG. 10) were prepared using a pool of sera from dogs infected with heartworm. Serial dilutions in EMEM-FBS (concentrations shown in FIG. 11) were prepared using a pool of sera from dogs sensitized to flea saliva. Additional samples were prepared in which both pools of sera were heat inactivated for about 4 hours at 56° C. The heat treated samples were diluted as described above.

About 100 μl of each dilution of each serum sample was added to separate wells containing fixed RBL-hFc<sub>ε</sub>R cells and the plates were incubated at 37° C. for about 1 hour. Following the incubation, the plates were washed 4 times with PBST. About 5 μg of a murine IgG monoclonal antibody anti-canine IgE antibody (i.e., Custom Monoclonal Antibody #71; available from Custom Monoclonal International) in 100 μl of EMEM-FBS was added to each well. The plates were incubated for about 30 minutes at 37° C. Following the incubation, the plates were washed 4 times with PBST. About 100

ng of horseradish peroxidase labelled donkey anti-murine IgG (available from Jackson Laboratories, Westgrove, Pa.) in 100  $\mu$ l EMEM-FBS was added to each well, and the plates were incubated for about 30 minutes at room temperature. Following the incubation, the plates were washed 4 times with PBST. The presence of anti-murine IgG bound to the plates thereby indicating the ability of RBL-hFc $\epsilon$ R cells to bind to canine IgE was detected using the method described in Example 4.

The results shown in FIG. 10 indicate that canine IgE from heartworm-infected dogs ( $\blacklozenge$ ) is detected using RBL-h Fc $\epsilon$ R cells expressing the alpha chain of human Fc $\epsilon$ R. In addition, the absence of colorimetric signal in samples of heat inactivated samples of such sera ( $\blacksquare$ ) indicates that antibody detected by the Fc $\epsilon$ R alpha chain on the RBL-h Fc $\epsilon$ R cells is

an epsilon isotype antibody. Similarly, the results shown in FIG. 11 indicate that canine IgE from dogs sensitized with flea saliva ( $\blacklozenge$ ) is detected using RBL-h Fc $\epsilon$ R cells expressing the alpha chain of human Fc $\epsilon$ R. In addition, the absence of colorimetric signal in samples of heat inactivated samples of such sera ( $\blacksquare$ ) indicates that antibody detected by the Fc $\epsilon$ R alpha chain on the RBL-h Fc $\epsilon$ R cells is an epsilon isotype antibody.

While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1

<211> LENGTH: 1198

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (107)..(877)

<400> SEQUENCE: 1

tactaagagt	ctccagcatc	ctccacctgt	ctaccaccga	gcatgggcct	atatttgaag	60
ccttagatct ctccagcaca gtaagcacca ggagtccatg aagaag atg gct cct						115
Met Ala Pro						
1						
gcc atg gaa tcc cct act cta ctg tgt gta gcc tta ctg ttc ttc gct	163					
Ala Met Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu Phe Phe Ala						
5 10 15						
cca gat ggc gtg tta gca gtc cct cag aaa cct aag gtc tcc ttg aac	211					
Pro Asp Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val Ser Leu Asn						
20 25 30 35						
cct cca tgg aat aga ata ttt aaa gga gag aat gtg act ctt aca tgt	259					
Pro Pro Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr Leu Thr Cys						
40 45 50						
aat ggg aac aat ttc ttt gaa gtc agt tcc acc aaa tgg ttc cac aat	307					
Asn Gly Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp Phe His Asn						
55 60 65						
ggc agc ctt tca gaa gag aca aat tca agt ttg aat att gtg aat gcc	355					
Gly Ser Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile Val Asn Ala						
70 75 80						
aaa ttt gaa gac agt gga gaa tac aaa tgt cag cac caa caa gtt aat	403					
Lys Phe Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln Gln Val Asn						
85 90 95						
gag agt gaa cct gtg tac ctg gaa gtc ttc agt gac tgg ctg ctc ctt	451					
Glu Ser Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp Leu Leu Leu						
100 105 110 115						
cag gcc tct gct gag gtg gtg atg gag ggc cag ccc ctc ttc ctc agg	499					
Gln Ala Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu Phe Leu Arg						
120 125 130						
tgc cat ggt tgg agg aac tgg gat gtg tac aag gtg atc tat tat aag	547					
Cys His Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys						
135 140 145						
gat ggt gaa gct ctc aag tac tgg tat gag aac cac aac atc tcc att	595					
Asp Gly Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile						



-continued

150	155	160	
aca aat gcc aca gtt gaa gac agt gga acc tac tac tgt acg ggc aaa			643
Thr Asn Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys			
165	170	175	
gtg tgg cag ctg gac tat gag tct gag ccc ctc aac att act gta ata			691
Val Trp Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile Thr Val Ile			
180	185	190	195
aaa gct ccg cgt gag aag tac tgg cta caa ttt ttt atc cca ttg ttg			739
Lys Ala Pro Arg Glu Lys Tyr Trp Leu Gln Phe Phe Ile Pro Leu Leu			
200	205	210	
gtg gtg att ctg ttt gct gtg gac aca gga tta ttt atc tca act cag			787
Val Val Ile Leu Phe Ala Val Asp Thr Gly Leu Phe Ile Ser Thr Gln			
215	220	225	
cag cag gtc aca ttt ctc ttg aag att aag aga acc agg aaa ggc ttc			835
Gln Gln Val Thr Phe Leu Leu Lys Ile Lys Arg Thr Arg Lys Gly Phe			
230	235	240	
aga ctt ctg aac cca cat cct aag cca aac ccc aaa aac aac			877
Arg Leu Leu Asn Pro His Pro Lys Pro Asn Pro Lys Asn Asn			
245	250	255	
tgatataatt actcaagaaa tatttgcaac attagttttt ttccagcatc agcaattgct			937
actcaattgt caaacacagc ttgcaatata catagaaacg tctgtgctca aggatttata			997
gaaatgcttc attaaactga gtgaaactgg ttaagtggca tgtaaatagta agtgctcaat			1057
taacattggt tgaataaatg agagaatgaa tagattcatt tattagcatt tgtaaaagag			1117
atgttcaatt tcaataaaaa aaatataaaa ccatgtaaca gaatgcttct gagtaaaaaa			1177
aaaaaaaaa aaaaaaaaaa a			1198
<210> SEQ ID NO 2			
<211> LENGTH: 257			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 2			
Met Ala Pro Ala Met Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu			
1	5	10	15
Phe Phe Ala Pro Asp Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val			
20	25	30	
Ser Leu Asn Pro Pro Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr			
35	40	45	
Leu Thr Cys Asn Gly Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp			
50	55	60	
Phe His Asn Gly Ser Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile			
65	70	75	80
Val Asn Ala Lys Phe Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln			
85	90	95	
Gln Val Asn Glu Ser Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp			
100	105	110	
Leu Leu Leu Gln Ala Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu			
115	120	125	
Phe Leu Arg Cys His Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile			
130	135	140	
Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn			
145	150	155	160
Ile Ser Ile Thr Asn Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys			
165	170	175	

-continued

Thr	Gly	Lys	Val	Trp	Gln	Leu	Asp	Tyr	Glu	Ser	Glu	Pro	Leu	Asn	Ile
			180					185						190	
Thr	Val	Ile	Lys	Ala	Pro	Arg	Glu	Lys	Tyr	Trp	Leu	Gln	Phe	Phe	Ile
		195					200					205			
Pro	Leu	Leu	Val	Val	Ile	Leu	Phe	Ala	Val	Asp	Thr	Gly	Leu	Phe	Ile
	210					215					220				
Ser	Thr	Gln	Gln	Gln	Val	Thr	Phe	Leu	Leu	Lys	Ile	Lys	Arg	Thr	Arg
225					230					235				240	
Lys	Gly	Phe	Arg	Leu	Leu	Asn	Pro	His	Pro	Lys	Pro	Asn	Pro	Lys	Asn
			245					250						255	

Asn

<210> SEQ ID NO 3  
 <211> LENGTH: 1198  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 3

```

tttttttttt tttttttttt tttttttact cagaagcatt ctgttacatg gttttatatt      60
tattttattg aaattgaaca tctcttttac aaatgctaataaatgaatct attcattctc      120
tcattttattc aaccaatggt aattgagcac ttactattac atgccactta accagtttca      180
ctcagtttaa tgaagcattt ctataaatcc ttgagcacag acgtttctat gtatattgca      240
agctgtgttt gacaattgag tagcaattgc tgatgctgga aaaaaactaa tgttgcaaat      300
atttcttgag taattatatt agttgttttt ggggtttggc ttaggatgtg ggttcagaag      360
tctgaagcct ttcctgtgtc tcttaattct caagagaaat gtgacctgct gctgagttga      420
gataaataat cctgtgtcca cagcaaacag aatcaccacc aacaatggga taaaaaattg      480
tagccagtac ttctcacgag gagcttttat tacagtaatg ttgaggggct cagactcata      540
gtccagctgc cacactttgc ccgtacagta gtaggttcca ctgtcttcaa ctgtggcatt      600
tgtaatggag atgttgttgt tctcatacca gtacttgaga gcttcacat ccttataata      660
gatcaccttg tacacatccc agttctccca accatggcac ctgaggaaga ggggctggcc      720
ctccatcacc acctcagcag aggcctgaag gagcagccag tcaactgaaga cttccaggta      780
cacaggttca ctctcattaa cttgttggtg ctgacatttg tattctccac tgtcttcaaa      840
tttggcattc acaatattca aactgaatt tgtctcttct gaaaggctgc cattgtggaa      900
ccatttggtg gaactgactt caaagaaatt gttccatta catgtaagag tcacattctc      960
tcctttaaat attctattcc atggaggggt caaggagacc ttaggtttct gagggactgc     1020
taacacgcca tctggagcga agaacagtaa ggctacacac agtagagtag gggattccat     1080
ggcaggagcc atctcttcca tggactcctg gtgcttactg tgctggagag atctaaggct     1140
tcaaatatag gccatgctc ggtggttagc aggtggagga tgctggagac tcttagta      1198

```

<210> SEQ ID NO 4  
 <211> LENGTH: 774  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: exon  
 <222> LOCATION: (1)..(774)

&lt;400&gt; SEQUENCE: 4

```

atg gct cct gcc atg gaa tcc cct act cta ctg tgt gta gcc tta ctg      48
Met Ala Pro Ala Met Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu
1          5          10          15

```

-continued

ttc ttc gct cca gat ggc gtg tta gca gtc cct cag aaa cct aag gtc Phe Phe Ala Pro Asp Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val 20 25 30	96
tcc ttg aac cct cca tgg aat aga ata ttt aaa gga gag aat gtg act Ser Leu Asn Pro Pro Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr 35 40 45	144
ctt aca tgt aat ggg aac aat ttc ttt gaa gtc agt tcc acc aaa tgg Leu Thr Cys Asn Gly Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp 50 55 60	192
ttc cac aat ggc agc ctt tca gaa gag aca aat tca agt ttg aat att Phe His Asn Gly Ser Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile 65 70 75 80	240
gtg aat gcc aaa ttt gaa gac agt gga gaa tac aaa tgt cag cac caa Val Asn Ala Lys Phe Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln 85 90 95	288
caa gtt aat gag agt gaa cct gtg tac ctg gaa gtc ttc agt gac tgg Gln Val Asn Glu Ser Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp 100 105 110	336
ctg ctc ctt cag gcc tct gct gag gtg gtg atg gag ggc cag ccc ctc Leu Leu Leu Gln Ala Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu 115 120 125	384
ttc ctc agg tgc cat ggt tgg agg aac tgg gat gtg tac aag gtg atc Phe Leu Arg Cys His Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile 130 135 140	432
tat tat aag gat ggt gaa gct ctc aag tac tgg tat gag aac cac aac Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn 145 150 155 160	480
atc tcc att aca aat gcc aca gtt gaa gac agt gga acc tac tac tgt Ile Ser Ile Thr Asn Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys 165 170 175	528
acg ggc aaa gtg tgg cag ctg gac tat gag tct gag ccc ctc aac att Thr Gly Lys Val Trp Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile 180 185 190	576
act gta ata aaa gct ccg cgt gag aag tac tgg cta caa ttt ttt atc Thr Val Ile Lys Ala Pro Arg Glu Lys Tyr Trp Leu Gln Phe Phe Ile 195 200 205	624
cca ttg ttg gtg gtg att ctg ttt gct gtg gac aca gga tta ttt atc Pro Leu Leu Val Val Ile Leu Phe Ala Val Asp Thr Gly Leu Phe Ile 210 215 220	672
tca act cag cag cag gtc aca ttt ctc ttg aag att aag aga acc agg Ser Thr Gln Gln Gln Val Thr Phe Leu Leu Lys Ile Lys Arg Thr Arg 225 230 235 240	720
aaa ggc ttc aga ctt ctg aac cca cat cct aag cca aac ccc aaa aac Lys Gly Phe Arg Leu Leu Asn Pro His Pro Lys Pro Asn Pro Lys Asn 245 250 255	768
aac tga Asn	774

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 774

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 5

tcagttgttt ttgggggttg gcttaggatg tgggttcaga agtctgaagc ctttctggt	60
tctcttaatc ttcaagagaa atgtgacctg ctgctgagtt gagataaata atcctgtgtc	120
cacagcaaac agaatcacca ccaacaatgg gataaaaaat tgtagccagt acttctcacg	180
cggagctttt attacagtaa tgttgagggg ctcagactca tagtcagct gccacacttt	240

-continued

```

ccccgtacag tagtaggttc cactgtcttc aactgtggca tttgtaatgg agatgttgtg 300
gtttctcatc cagtacttga gagcttcacc atccttataa tagatcacct tgtacacatc 360
ccagttcctc caaccatggc acctgaggaa gaggggctgg cctccatca ccacctcagc 420
agaggcctga aggagcagcc agtcactgaa gacttccagg tacacagggt cactctcatt 480
aactgtttgg tgotgacatt tgtattctcc actgtcttca aatttggcat tcacaatatt 540
caaaacttgaa tttgtctctt ctgaaagggt gccattgtgg aaccatttgg tggaactgac 600
ttcaaagaaa ttgttcccat tacatgtaag agtcacattc tctcctttaa atattctatt 660
ccatggaggg ttcaaggaga ccttaggttt ctgagggact gctaacacgc catctggagc 720
gaagaacagt aaggctacac acagtagagt aggggattcc atggcaggag ccat 774

```

```

<210> SEQ ID NO 6
<211> LENGTH: 232
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 6

```

```

Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro Trp Asn Arg Ile
1           5           10          15
Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly Asn Asn Phe Phe
                20          25          30
Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser Leu Ser Glu Glu
        35          40          45
Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe Glu Asp Ser Gly
        50          55          60
Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser Glu Pro Val Tyr
65          70          75          80
Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala Ser Ala Glu Val
        85          90          95
Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His Gly Trp Arg Asn
        100         105         110
Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys
        115         120         125
Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn Ala Thr Val Glu
        130         135         140
Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu Asp Tyr
145         150         155         160
Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala Pro Arg Glu Lys
        165         170         175
Tyr Trp Leu Gln Phe Phe Ile Pro Leu Leu Val Val Ile Leu Phe Ala
        180         185         190
Val Asp Thr Gly Leu Phe Ile Ser Thr Gln Gln Gln Val Thr Phe Leu
        195         200         205
Leu Lys Ile Lys Arg Thr Arg Lys Gly Phe Arg Leu Leu Asn Pro His
210         215         220
Pro Lys Pro Asn Pro Lys Asn Asn
225         230

```

```

<210> SEQ ID NO 7
<211> LENGTH: 699
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: exon

```

-continued

&lt;222&gt; LOCATION: (1) .. (699)

&lt;400&gt; SEQUENCE: 7

gtc cct cag aaa cct aag gtc tcc ttg aac cct cca tgg aat aga ata 48  
 Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro Trp Asn Arg Ile  
 1 5 10 15

ttt aaa gga gag aat gtg act ctt aca tgt aat ggg aac aat ttc ttt 96  
 Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly Asn Asn Phe Phe  
 20 25 30

gaa gtc agt tcc acc aaa tgg ttc cac aat ggc agc ctt tca gaa gag 144  
 Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser Leu Ser Glu Glu  
 35 40 45

aca aat tca agt ttg aat att gtg aat gcc aaa ttt gaa gac agt gga 192  
 Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe Glu Asp Ser Gly  
 50 55 60

gaa tac aaa tgt cag cac caa caa gtt aat gag agt gaa cct gtg tac 240  
 Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser Glu Pro Val Tyr  
 65 70 75 80

ctg gaa gtc ttc agt gac tgg ctg ctc ctt cag gcc tct gct gag gtg 288  
 Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala Ser Ala Glu Val  
 85 90 95

gtg atg gag ggc cag ccc ctc ttc ctc agg tgc cat ggt tgg agg aac 336  
 Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His Gly Trp Arg Asn  
 100 105 110

tgg gat gtg tac aag gtg atc tat tat aag gat ggt gaa gct ctc aag 384  
 Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys  
 115 120 125

tac tgg tat gag aac cac aac atc tcc att aca aat gcc aca gtt gaa 432  
 Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn Ala Thr Val Glu  
 130 135 140

gac agt gga acc tac tac tgt acg ggc aaa gtg tgg cag ctg gac tat 480  
 Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu Asp Tyr  
 145 150 155 160

gag tct gag ccc ctc aac att act gta ata aaa gct ccg cgt gag aag 528  
 Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala Pro Arg Glu Lys  
 165 170 175

tac tgg cta caa ttt ttt atc cca ttg ttg gtg gtg att ctg ttt gct 576  
 Tyr Trp Leu Gln Phe Phe Ile Pro Leu Leu Val Val Ile Leu Phe Ala  
 180 185 190

gtg gac aca gga tta ttt atc tca act cag cag cag gtc aca ttt ctc 624  
 Val Asp Thr Gly Leu Phe Ile Ser Thr Gln Gln Gln Val Thr Phe Leu  
 195 200 205

ttg aag att aag aga acc agg aaa ggc ttc aga ctt ctg aac cca cat 672  
 Leu Lys Ile Lys Arg Thr Arg Lys Gly Phe Arg Leu Leu Asn Pro His  
 210 215 220

cct aag cca aac ccc aaa aac aac tga 699  
 Pro Lys Pro Asn Pro Lys Asn Asn  
 225 230

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 32

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Primer

&lt;400&gt; SEQUENCE: 8

cgcggtatcct ataatatgg ctctgccaat gg 32

&lt;210&gt; SEQ ID NO 9

&lt;211&gt; LENGTH: 26

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 9

ggcgaattct taagctttta ttacag                26

<210> SEQ ID NO 10
<211> LENGTH: 591
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(591)

<400> SEQUENCE: 10

atg gct cct gcc atg gaa tcc cct act cta ctg tgt gta gcc tta ctg      48
Met Ala Pro Ala Met Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu
1          5          10          15

ttc ttc gct cca gat ggc gtg tta gca gtc cct cag aaa cct aag gtc      96
Phe Phe Ala Pro Asp Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val
20          25          30

tcc ttg aac cct cca tgg aat aga ata ttt aaa gga gag aat gtg act     144
Ser Leu Asn Pro Pro Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr
35          40          45

ctt aca tgt aat ggg aac aat ttc ttt gaa gtc agt tcc acc aaa tgg     192
Leu Thr Cys Asn Gly Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp
50          55          60

ttc cac aat ggc agc ctt tca gaa gag aca aat tca agt ttg aat att     240
Phe His Asn Gly Ser Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile
65          70          75          80

gtg aat gcc aaa ttt gaa gac agt gga gaa tac aaa tgt cag cac caa     288
Val Asn Ala Lys Phe Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln
85          90          95

caa gtt aat gag agt gaa cct gtg tac ctg gaa gtc ttc agt gac tgg     336
Gln Val Asn Glu Ser Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp
100         105         110

ctg ctc ctt cag gcc tct gct gag gtg gtg atg gag ggc cag ccc ctc     384
Leu Leu Leu Gln Ala Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu
115        120        125

ttc ctc agg tgc cat ggt tgg agg aac tgg gat gtg tac aag gtg atc     432
Phe Leu Arg Cys His Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile
130        135        140

tat tat aag gat ggt gaa gct ctc aag tac tgg tat gag aac cac aac     480
Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn
145        150        155        160

atc tcc att aca aat gcc aca gtt gaa gac agt gga acc tac tac tgt     528
Ile Ser Ile Thr Asn Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys
165        170        175

acg ggc aaa gtg tgg cag ctg gac tat gag tct gag ccc ctc aac att     576
Thr Gly Lys Val Trp Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile
180        185        190

act gta ata aaa gct
Thr Val Ile Lys Ala
195

<210> SEQ ID NO 11
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

-continued

---

Met Ala Pro Ala Met Glu Ser Pro Thr Leu Leu Cys Val Ala Leu Leu  
1 5 10 15

Phe Phe Ala Pro Asp Gly Val Leu Ala Val Pro Gln Lys Pro Lys Val  
20 25 30

Ser Leu Asn Pro Pro Trp Asn Arg Ile Phe Lys Gly Glu Asn Val Thr  
35 40 45

Leu Thr Cys Asn Gly Asn Asn Phe Phe Glu Val Ser Ser Thr Lys Trp  
50 55 60

Phe His Asn Gly Ser Leu Ser Glu Glu Thr Asn Ser Ser Leu Asn Ile  
65 70 75 80

Val Asn Ala Lys Phe Glu Asp Ser Gly Glu Tyr Lys Cys Gln His Gln  
85 90 95

Gln Val Asn Glu Ser Glu Pro Val Tyr Leu Glu Val Phe Ser Asp Trp  
100 105 110

Leu Leu Leu Gln Ala Ser Ala Glu Val Val Met Glu Gly Gln Pro Leu  
115 120 125

Phe Leu Arg Cys His Gly Trp Arg Asn Trp Asp Val Tyr Lys Val Ile  
130 135 140

Tyr Tyr Lys Asp Gly Glu Ala Leu Lys Tyr Trp Tyr Glu Asn His Asn  
145 150 155 160

Ile Ser Ile Thr Asn Ala Thr Val Glu Asp Ser Gly Thr Tyr Tyr Cys  
165 170 175

Thr Gly Lys Val Trp Gln Leu Asp Tyr Glu Ser Glu Pro Leu Asn Ile  
180 185 190

Thr Val Ile Lys Ala  
195

<210> SEQ ID NO 12  
 <211> LENGTH: 516  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(516)

<400> SEQUENCE: 12

gtc cct cag aaa cct aag gtc tcc ttg aac cct cca tgg aat aga ata Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro Trp Asn Arg Ile 1 5 10 15	48
ttt aaa gga gag aat gtg act ctt aca tgt aat ggg aac aat ttc ttt Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly Asn Asn Phe Phe 20 25 30	96
gaa gtc agt tcc acc aaa tgg ttc cac aat ggc agc ctt tca gaa gag Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser Leu Ser Glu Glu 35 40 45	144
aca aat tca agt ttg aat att gtg aat gcc aaa ttt gaa gac agt gga Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe Glu Asp Ser Gly 50 55 60	192
gaa tac aaa tgt cag cac caa caa gtt aat gag agt gaa cct gtg tac Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser Glu Pro Val Tyr 65 70 75 80	240
ctg gaa gtc ttc agt gac tgg ctg ctc ctt cag gcc tct gct gag gtg Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala Ser Ala Glu Val 85 90 95	288
gtg atg gag ggc cag ccc ctc ttc ctc agg tgc cat ggt tgg agg aac Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His Gly Trp Arg Asn 100 105 110	336

-continued

tgg gat gtg tac aag gtg atc tat tat aag gat ggt gaa gct ctc aag	384
Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys	
115 120 125	
tac tgg tat gag aac cac aac atc tcc att aca aat gcc aca gtt gaa	432
Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn Ala Thr Val Glu	
130 135 140	
gac agt gga acc tac tac tgt acg ggc aaa gtg tgg cag ctg gac tat	480
Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu Asp Tyr	
145 150 155 160	
gag tct gag ccc ctc aac att act gta ata aaa gct	516
Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala	
165 170	

<210> SEQ ID NO 13  
 <211> LENGTH: 172  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

Val Pro Gln Lys Pro Lys Val Ser Leu Asn Pro Pro Trp Asn Arg Ile	
1 5 10 15	
Phe Lys Gly Glu Asn Val Thr Leu Thr Cys Asn Gly Asn Asn Phe Phe	
20 25 30	
Glu Val Ser Ser Thr Lys Trp Phe His Asn Gly Ser Leu Ser Glu Glu	
35 40 45	
Thr Asn Ser Ser Leu Asn Ile Val Asn Ala Lys Phe Glu Asp Ser Gly	
50 55 60	
Glu Tyr Lys Cys Gln His Gln Gln Val Asn Glu Ser Glu Pro Val Tyr	
65 70 75 80	
Leu Glu Val Phe Ser Asp Trp Leu Leu Leu Gln Ala Ser Ala Glu Val	
85 90 95	
Val Met Glu Gly Gln Pro Leu Phe Leu Arg Cys His Gly Trp Arg Asn	
100 105 110	
Trp Asp Val Tyr Lys Val Ile Tyr Tyr Lys Asp Gly Glu Ala Leu Lys	
115 120 125	
Tyr Trp Tyr Glu Asn His Asn Ile Ser Ile Thr Asn Ala Thr Val Glu	
130 135 140	
Asp Ser Gly Thr Tyr Tyr Cys Thr Gly Lys Val Trp Gln Leu Asp Tyr	
145 150 155 160	
Glu Ser Glu Pro Leu Asn Ile Thr Val Ile Lys Ala	
165 170	

What is claimed is:

1. A method to detect a plant allergy in a canid, felid or equid comprising:

- (a) contacting an allergen from a plant with a putative IgE containing composition from said animal, under conditions suitable for formation of an allergen:IgE complex;
- (b) contacting the putative allergen:IgE complex with a human FcεR molecule under conditions suitable for formation of an allergen:IgE:FcεR molecule complex, wherein either the allergen or the human FcεR molecule is labeled by conjugation to a detectable marker;
- (c) washing the allergen:IgE:FcεR mixture to remove non-specifically bound labeled- molecule; and
- (d) detecting the presence of an allergen:IgE:FcεR complex, if present, by detecting the presence of detectable marker, wherein the presence of an allergen:IgE:FcεR complex indicates the animal is allergic to said plant.

50

2. The method of claim 1, wherein said putative IgE containing composition comprises an animal fluid is selected from the group consisting of serum, plasma, blood, urine, tears, saliva, nasal secretions and milk.

55 3. The method of claim 1, wherein said allergen is selected from the group consisting of a grass allergen, a plantain allergen, a pigweed allergen, a ragweed allergen, a sage allergen, an elm allergen, a cocklebur allergen, an elder allergen, a walnut allergen, a cottonwood allergen, an ash allergen, a birch allergen, a cedar allergen, an oak allergen, and a mulberry allergen.

60 4. The method of claim 1, wherein said allergen is selected from the group consisting of a Johnson Grass allergen, a Kentucky Blue Grass allergen, a Meadow Fescue allergen, an Orchard Grass allergen, a Perennial Rye Grass allergen, a Redtop Grass allergen, a Timothy Grass allergen, a Bermuda Grass allergen, a Brome Grass allergen, a Curly Dock aller-



41

gen, an English Plantain allergen, a Mexican Firebush allergen, a Lamb's Quarters allergen, a Rough Pigweed allergen, a Short Ragweed allergen, a Wormwood Sage allergen, an American Elm allergen, a Common Cocklebur allergen, a Box Elder allergen, a Black Walnut allergen, an Eastern Cottonwood allergen, a Green Ash allergen, a River Birch allergen, a Red Cedar allergen, a Red Oak allergen, and a Red Mulberry allergen.

5 The method of claim 1, wherein said FcεR molecule is encoded by a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10 and SEQ ID NO:12.

10 The method of claim 1, wherein said allergen is bound to a substrate.

7. A method to detect plant-allergen-specific IgE in a canid, felid or equid, said method comprising:

- (a) contacting a plant allergen with a putative IgE containing composition from said animal, under conditions suitable for formation of an allergen:IgE complex;
- (b) contacting the putative allergen:IgE complex with a human FcεR molecule under conditions suitable for formation of an allergen:IgE:FcεR molecule complex, wherein either the allergen or the human FcεR molecule is labeled by conjugation to a detectable marker;
- (c) washing the allergen:IgE:FcεR mixture to remove non-specifically bound labeled-molecule; and
- (d) detecting the presence of an allergen:IgE:FcεR complex, if present, by detecting the presence of detectable marker, wherein the presence of an allergen:IgE:FcεR molecule complex indicates the presence of plant-allergen-specific IgE.

42

8. The method of claim 7, wherein said putative IgE containing composition comprises an animal fluid is selected from the group consisting of serum, plasma, blood, urine, tears, saliva, nasal secretions and milk.

5 9. The method of claim 7, wherein said allergen is selected from the group consisting of a grass allergen, a plantain allergen, a pigweed allergen, a ragweed allergen, a sage allergen, an elm allergen, a cocklebur allergen, an elder allergen, a walnut allergen, a cottonwood allergen, an ash allergen, a birch allergen, a cedar allergen, an oak allergen, and a mulberry allergen.

10 10. The method of claim 7, wherein said allergen is selected from the group consisting of a Johnson Grass allergen, a Kentucky Blue Grass allergen, a Meadow Fescue allergen, an Orchard Grass allergen, a Perennial Rye Grass allergen, a Redtop Grass allergen, a Timothy Grass allergen, a Bermuda Grass allergen, a Brome Grass allergen, a Curly Dock allergen, an English Plantain allergen, a Mexican Firebush allergen, a Lamb's Quarters allergen, a Rough Pigweed allergen, a Short Ragweed allergen, a Wormwood Sage allergen, an American Elm allergen, a Common Cocklebur allergen, a Box Elder allergen, a Black Walnut allergen, an Eastern Cottonwood allergen, a Green Ash allergen, a River Birch allergen, a Red Cedar allergen, a Red Oak allergen, and a Red Mulberry allergen.

15 11. The method of claim 7, wherein said FcεR molecule is encoded by a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10 and SEQ ID NO:12.

20 12. The method of claim 7, wherein said allergen is bound to a substrate.

\* \* \* \* \*

专利名称(译)	检测IgE的方法		
公开(公告)号	<a href="#">US7691384</a>	公开(公告)日	2010-04-06
申请号	US11/683854	申请日	2007-03-08
[标]申请(专利权)人(译)	FR GLENN - [R PORTER JAMES P RUSHLOW KEITHê WASSOM DONALD大号		
申请(专利权)人(译)	FRANK GLENN - [R PORTER JAMES P RUSHLOW KEITHê WASSOM DONALD大号		
当前申请(专利权)人(译)	HESKA CORPORATION		
[标]发明人	FRANK GLENN R PORTER JAMES P RUSHLOW KEITH E WASSOM DONALD L		
发明人	FRANK, GLENN R. PORTER, JAMES P. RUSHLOW, KEITH E. WASSOM, DONALD L.		
IPC分类号	A61K39/00 A61K39/385 G01N33/566 C07K14/735 C12N5/10 C12N15/09 G01N33/53 G01N33/68		
CPC分类号	G01N33/6854 G01N2800/24 G01N2333/70535		
代理机构(译)	SHERIDAN ROSS P.C.		
其他公开文献	US20070218565A1		
外部链接	<a href="#">Espacenet</a> <a href="#">USPTO</a>		

#### 摘要(译)

本发明包括使用人Fcε受体 ( Fc68R ) 检测IgE以检测来自猫，狗或马的生物样品中的IgE抗体的方法。本发明还涉及进行这些方法的试剂盒。




Fig. 1